



US 20050170392A1

(19) **United States**
 (12) **Patent Application Publication** (10) **Pub. No.: US 2005/0170392 A1**
Zerangue (43) **Pub. Date: Aug. 4, 2005**

(54) **OAT3 TRANSPORTERS EXPRESSED IN BLOOD BRAIN BARRIER CELLS**

Related U.S. Application Data

(60) Provisional application No. 60/540,772, filed on Jan. 30, 2004.

(75) Inventor: **Noa Zerangue**, Sunnyvale, CA (US)

Publication Classification

Correspondence Address:
**TOWNSEND AND TOWNSEND AND CREW,
 LLP
 TWO EMBARCADERO CENTER
 EIGHTH FLOOR
 SAN FRANCISCO, CA 94111-3834 (US)**

(51) **Int. Cl.⁷** **C12Q 1/68**; G01N 33/53;
 G01N 33/567; A61K 31/60;
 A61K 31/225; A61K 31/19
 (52) **U.S. Cl.** **435/6**; 514/2; 514/159; 514/356;
 514/548; 514/557; 435/7.2

(57) **ABSTRACT**

OAT3 is consistently expressed at high levels in brain microvessel endothelial cells. Disclosed herein are assays for determining whether a test material/molecule is a substrate for, and/or is actively transported by, the OAT3 transporter, and therefore a candidate substrate for crossing the blood brain barrier. The assays are useful in screening for therapeutic, cytotoxic or imaging compounds used in the treatment or diagnosis of neurological diseases.

(73) Assignee: **Xenoport, Inc.**, Santa Clara, CA


(21) Appl. No.: **11/027,581**

(22) Filed: **Dec. 30, 2004**


Figure 1

OAT3 SUBSTRATES


Para-amino-hippurate




Ochratoxin A



Estrone-3-sulphate



Benzylpenicillin



Probenecid




Figure 2

**Uptake of PAH into Xenopus
Oocytes Expressing OAT3**




Figure 3




Figure 4




Figure 5





Figure 6



OAT3 TRANSPORTERS EXPRESSED IN BLOOD BRAIN BARRIER CELLS

CONTINUITY

[0001] This application claims the benefit of U.S. Provisional Application No. 60/540,772, filed Jan. 30, 2004, the disclosure of which is incorporated by reference herein in its entirety.

TECHNICAL FIELD

[0002] The disclosures herein relate to assays and methods of using the same for screening compounds and/or chemical moieties for their ability to be actively transported across the blood brain barrier.

BACKGROUND

[0003] The capillaries that supply blood to the tissues of the brain constitute the blood brain barrier (Goldstein et al. (1986) *Scientific American* 255:74-83; Pardridge, W. M. (1986) *Endocrin. Rev.* 7:314-330). The endothelial cells which form the brain capillaries are different from those found in other tissues in the body. Brain capillary endothelial cells are joined together by tight intercellular junctions which form a continuous wall against the passive diffusion of molecules from the blood to the brain and other parts of the central nervous system (CNS). These cells are also different in that they have few pinocytic vesicles which in other tissues allow somewhat unselective transport across the capillary wall. Also lacking are continuous gaps or channels running between the cells which would allow unrestricted passage.

[0004] The blood-brain barrier functions to ensure that the environment of the brain is constantly controlled. The levels of various substances in the blood, such as hormones, amino acids and ions, undergo frequent small fluctuations which can be brought about by activities such as eating and exercise (Goldstein et al., cited supra). If the brain was not protected by the blood brain barrier from these variations in serum composition, the result could be uncontrolled neural activity.

[0005] The isolation of the brain from the bloodstream is not complete. If this were the case, the brain would be unable to function properly due to a lack of nutrients and because of the need to exchange chemicals with the rest of the body. The presence of specific transport systems within the capillary endothelial cells assures that the brain receives, in a controlled manner, all of the compounds required for normal growth and function. In many instances, these transport systems consist of membrane-associated proteins which selectively bind and certain molecules across the barrier membranes. These transporter proteins are known as solute carrier transporters.

[0006] The problem posed by the blood-brain barrier is that, in the process of protecting the brain, it excludes many potentially useful therapeutic agents. Presently, only substances which are sufficiently lipophilic can penetrate the blood-brain barrier (Goldstein et al., cited supra; Pardridge, W. M., cited supra). Some drugs can be modified to make them more lipophilic and thereby increase their ability to cross the blood brain barrier. However, each modification must be tested individually on each drug and the modification can alter the activity of the drug.

[0007] Because the blood brain barrier is composed of brain microvessel endothelial cells, these cells have been isolated and cultured for use in in vitro model systems for studying the blood brain barrier (Bowman et. al, Brain microvessel endothelial cells in tissue culture: A model for study of blood-brain barrier permeability, *Ann. Neurol.* 14, 396-402 (1983); Audus and Borchardt, Characterization of an in vitro blood-brain barrier model system for studying drug transport and metabolism, *Pharm. Res.* 3, 81-87 (1986)). In vitro model systems of the blood brain barrier have been successfully derived from bovine, canine, human, murine, porcine, and rat cells, and have similar permeability properties due to similarity of the physiological characteristics of the blood brain barrier among mammals (Cserr et al., Blood-brain interfaces in vertebrates: a comparative approach, *Am. J. Physiol.* 246, R277-R288 (1984); Audus et al., The use of cultured epithelial and endothelial cells for drug transport and metabolism studies, *Pharm. Res.* 7, 435-451 (1990)). In these models, the cultured endothelial cells retain the characteristics of brain endothelial cells in vivo, such as morphology, specific blood brain barrier enzyme markers, and tight intercellular junctions. The cells can also be used for the study of passive diffusion, carrier mediated transport, and metabolism to specific factors affecting the blood brain barrier permeability. However, passaging of brain microvessel endothelial cells results in loss of specific endothelial and blood brain barrier markers as well as tight intercellular junctions (Brightman and Neuwelt (ed.), *Implications of the blood-brain barrier and its manipulation*, Vol. 1, Plenum Medical, New York, pp. 53-83 (1989)).

[0008] Currently, primary cultures of brain microvessel endothelial cells are the principal tool for in vitro prediction of blood brain barrier permeability. Isolated and cultured primary brain cells developed previously have exhibited different properties primarily due to considerable variability in the starting material. For example, with respect to transcellular transport, rigorous comparison of data between different laboratories has been very difficult (Pardridge et al., Comparison of in vitro and in vivo models of drug transcytosis through the blood-brain barrier, *J. Pharmacol. Exp. Thera.* 253, 884-891 (1990); Masereeuw et al., In vitro and in vivo transport of zidovudine (AZT) across the blood-brain barrier and the effect of transport inhibitors. *Pharm. Res.*, 11, 324-330 (1994)). Passaging primary cells can affect the differentiation of cells and lead to the selection of the most rapidly proliferating clones. Furthermore, the expression of some marker enzymes such as gamma-glutamyl transpeptidase as well as tight junctional complexity has been shown to decrease with time in culture and passage number (Meresse et. al., Bovine brain endothelial cells express tight junctions and monoamine oxidase activity in long-term culture, *J. Neurochem.* 53, 1363-1371 (1989)). Some transporter substrates have been demonstrated to accumulate in the brain (see U.S. Pat. No. 6,489,302).

[0009] Thus, it is apparent that the presently available clones of immortalized brain microvessel endothelial cell cultures suffer from individual drawbacks in terms of phenotype expression and homogeneous maintenance of that expression. This leads to difficulties with respect to accuracy and reproducibility in studies utilizing brain microvessel endothelial cells to model passage of chemical compounds and moieties, e.g., potential therapeutic compounds and/or drug moieties, across the blood brain barrier.

SUMMARY

[0010] Disclosed herein are methods of screening agents, conjugates or conjugate moieties for the ability to enter the CNS by crossing the blood brain barrier in order to treat or diagnose conditions within the CNS. These methods entail providing a cell expressing an OAT3 transporter, the transporter being situated in the plasma membrane of the cell. The cell is contacted with an agent, conjugate or conjugate moiety. Whether the agent, conjugate or conjugate moiety passes through the plasma membrane via the OAT3 transporter is determined. If the method comprises contacting the cell with an agent, the agent is a neuropharmaceutical agent or an imaging component. If the method comprises contacting the cell with a conjugate, the conjugate comprises an agent that is a neuropharmaceutical agent or an imaging component. If the method comprises contacting the cells with a conjugate moiety, the method further comprises linking the conjugate moiety to an agent that is a neuropharmaceutical agent or an imaging component.

[0011] In some methods, the cell endogenously expresses an OAT3 transporter. In other methods a nucleic acid molecule encoding an OAT3 transporter has been transfected or injected into the cell. In some methods the cell is a brain microvessel endothelial cell. In other methods the cell is a choroid plexus epithelial cell. In other methods the cell is an oocyte. In other methods the cell is a human embryonic kidney (HEK) cell. In other methods the cell is a Madin Darby canine kidney cell (MDCK). In still other methods, the cell is constructed to conditionally express the transporter.

[0012] In some methods the agent, conjugate or conjugate moiety comprises a xenobiotic molecule containing one or more ring structures and one or more anionic groups such as carboxylates, sulphates, and phosphates. In some methods the agent, conjugate or conjugate moiety is administered to an undiseased animal and any toxic effects are determined. In some methods the neuropharmaceutical agent is a cytotoxic neuropharmaceutical agent selected from the group consisting of platinum, nitrosourea, a phosphoramidate group that is selectively cytotoxic to brain tumor cells, nitroimidazole, and nitrogen mustard.

[0013] Disclosed herein are methods of screening agents, conjugates or conjugate moieties for the ability to enter the CNS by crossing the blood brain barrier wherein a cell used for testing is a brain microvessel endothelial cell that is one of a plurality of brain microvessel endothelial cells forming a polarized monolayer. An agent, conjugate or conjugate moiety is contacted to one side of the polarized monolayer and whether the agent, conjugate or conjugate moiety is transported into the brain microvessel endothelial cell(s) or to the opposite side of the polarized monolayer is determined. Some methods further comprise administering the agent, conjugate, or conjugate moiety to a peripheral tissue of an animal and measuring the amount of agent, conjugate, or conjugate moiety that passes through the blood brain barrier into the brain of the animal. In some methods, the animal is a non-human animal.

[0014] Disclosed herein are methods of screening an agent, conjugate or conjugate moiety for neuropharmacological activity useful for treating neurological disorders. In these methods, one determines whether the agent, conjugate or conjugate moiety is transported through an OAT3 trans-

porter. One then administers the agent, conjugate or conjugate moiety to a test animal and determines whether the agent, conjugate or conjugate moiety is actively transported across the blood brain barrier by measuring agent, conjugate or conjugate moiety concentrations found in the CNS of the animal. For those agents, conjugates or conjugate moieties that are transported in sufficient quantities, the agents, conjugates or conjugate moieties can be further tested in animals suffering from a particular neurological disorder to determine whether the agents, conjugates or conjugate moieties have the requisite therapeutic neuropharmacological activity for treating such neurological disorder.

[0015] Also disclosed herein are methods for in vitro screening of agents, conjugates or conjugate moieties for improved retention in the CNS. In these methods, one determines the substrate properties of a compound on both uptake transporters and efflux transporters. An agent, conjugate or conjugate moiety is first tested for activity on the OAT3 transporter. The agent, conjugate or conjugate moiety is then tested for substrate activity on an efflux transporter, such as P Glycoprotein (PgP). Those agents, conjugates or conjugate moieties active on both the efflux transporter and OAT3 are then modified and tested for a reduction of efflux substrate activity and retested for retention of activity on the OAT3 transporter. This iterative process produces an agent, conjugate or conjugate moiety with an increased ratio of substrate activities in the uptake and efflux systems, and improved retention of pharmacological levels of the modified agent, conjugate or conjugate moiety in the CNS.

[0016] Disclosed herein are methods of screening an agent, conjugate or conjugate moiety for capacity to be transported into the brain, comprising determining whether the agent, conjugate or conjugate moiety specifically binds to an OAT3 transporter, contacting the agent to one side of a polarized monolayer of cells, and determining whether the agent is actively transported across the polarized monolayer. In some methods the specific binding is determined by contacting a cell expressing the OAT3 transporter, the transporter being situated in the plasma membrane of the cell, with a substrate of the OAT3 transporter, and determining whether the agent inhibits transport of the substrate across the polarized monolayer.

[0017] Disclosed herein are pharmaceutical compositions comprising a therapeutic neuropharmaceutical agent, a cytotoxic neuropharmaceutical agent or an imaging component linked to a conjugate moiety to form a conjugate in which the conjugate moiety has a higher V_{max} for the OAT3 transporter than the therapeutic neuropharmaceutical agent, cytotoxic neuropharmaceutical agent or imaging component alone. Some pharmaceutical compositions have at least 5 times the V_{max} for OAT3 than the neuropharmaceutical agent or the imaging component alone. In some pharmaceutical compositions the conjugate has a V_{max} for OAT3 that is at least 5% of the V_{max} for OAT3 of a compound selected from the group comprising para-amino-hippurate (PAH), ochratoxin A, estrone-3-sulphate, benzylpenicillin, and probenecid. In some pharmaceutical compositions the conjugate has a lower V_{max} for an efflux transporter than the neuropharmaceutical agent or the imaging component alone.

[0018] Disclosed herein are methods of formulating a therapeutic neuropharmaceutical agent, a cytotoxic neuropharmaceutical agent or an imaging component. These meth-

ods entail linking the therapeutic neuropharmaceutical agent, the cytotoxic neuropharmaceutical agent or the imaging component to a conjugate moiety to form a conjugate, wherein the conjugate moiety has a greater V_{max} for an OAT3 transporter than the component alone. The conjugate is formulated with a pharmaceutical carrier as a pharmaceutical composition.

[0019] Disclosed herein are methods of delivering a therapeutic neuropharmaceutical agent, a cytotoxic neuropharmaceutical agent or an imaging component. Some methods involve administering to a patient a pharmaceutical composition comprising a therapeutic neuropharmaceutical agent, a cytotoxic neuropharmaceutical agent or an imaging component linked to a conjugate moiety to form a conjugate, wherein the conjugate has a higher V_{max} for an OAT3 transporter than the therapeutic neuropharmaceutical agent, cytotoxic neuropharmaceutical agent or imaging component alone, whereby the conjugate passes through brain microvessel endothelial cells which make up the blood brain barrier, via the OAT3 transporter, into the CNS of the patient. Also disclosed herein are methods of delivering a conjugate, comprising administering to a patient a pharmaceutical composition comprising a neuropharmaceutical agent or imaging component linked to a conjugate moiety to form the conjugate, wherein the conjugate has a higher V_{max} for an OAT3 transporter than the neuropharmaceutical agent or imaging component alone. In some methods the V_{max} of the conjugate is at least two-fold higher than that of the neuropharmaceutical agent or imaging component alone. In some methods the neuropharmaceutical agent is a cytotoxic neuropharmaceutical selected from the group consisting of platinum, nitrosourea, a phosphoramidate group selectively cytotoxic to brain tumor cells, nitroimidazole, and nitrogen mustard.

[0020] Disclosed herein are methods of treating neurological disorders. These methods entail administering to a patient an effective amount of an agent that is transported by OAT3, wherein the agent is a conjugate comprising a therapeutic neuropharmaceutical agent, a cytotoxic neuropharmaceutical agent or an imaging component linked to a conjugate moiety.

[0021] Disclosed herein are methods of screening an agent for decreased side effects in the central nervous system (CNS), comprising providing an agent having a pharmacological activity, wherein the pharmacological activity is useful for treating a disease present in a tissue other than the CNS, and the pharmacological activity results in undesired side effects in the CNS if the agent enters the CNS, modifying the agent, providing a cell expressing at least one efflux transporter protein that transports substrates out of the CNS, contacting the cell with the modified agent, and determining whether the modified agent is transported by the at least one efflux transporter protein with a higher V_{max} than the agent, a higher V_{max} indicating that the modification increases the capacity of the modified agent relative to the agent to be transported out of the CNS, thereby decreasing undesired side effects in the CNS.

BRIEF DESCRIPTION OF THE FIGURES

[0022] FIG. 1 shows the structures of known substrates of the OAT3 transporter.

[0023] FIG. 2 shows uptake of para-amino-hippurate (PAH) into oocytes injected with OAT3 cRNA.

[0024] FIG. 3 shows uptake of ^3H -Ochratoxin A and competition with PAH using HEK cells transfected with a tetracycline-inducible OAT3 expression vector.

[0025] FIG. 4 shows an efflux transporter ATPase activity assay using membrane preparations containing the PgP efflux transporter and the PgP substrate verapamil.

[0026] FIG. 5 shows an efflux transporter competition assay using the reporter molecule calcein-AM and the PgP substrate verapamil.

[0027] FIG. 6 shows a direct efflux transport assay using a polarized monolayer of MDCK cells transfected with a tetracycline-inducible PgP expression construct.

DEFINITIONS

[0028] "Transport by passive diffusion" refers to transport of an agent that is not mediated by a specific transporter protein. An agent that is substantially incapable of passive diffusion has a permeability across a standard cell monolayer (e.g., Caco-2 or MDCK cells or an artificial bilayer (PAMPA)) of less than 5×10^{-6} cm/sec, and usually less than 1×10^{-6} cm/sec in the absence of an efflux mechanism.

[0029] A "substrate" of a transporter protein is a compound whose uptake into or passage through the plasma membrane of a cell is facilitated at least in part by a transporter protein.

[0030] The term "ligand" of a transporter protein includes compounds that bind to the transporter protein. Some ligands are transported and are thereby also substrates. Some ligands inhibit or antagonize transport of a substrate by the transporter protein. Some ligands bind in a manner non-competitive with substrates and modulate the transport of substrates by the transporter protein.

[0031] The term "neuropharmaceutical agent" is used to describe a compound that has or may have a pharmacological activity in the treatment or prophylaxis of a neurological disorder. Neuropharmaceutical agents include compounds that are known drugs, compounds for which pharmacological activity has been identified but which are undergoing further therapeutic evaluation, and compounds that are members of collections and libraries that are to be screened for a pharmacological activity. The neuropharmaceutical agent can be a compound having a therapeutic, prophylactic or cytotoxic effect on a neurological disease including any condition which affects biological functioning of the central nervous system. Examples of neurological diseases include cancer (e.g., brain tumors), Acquired Immune Deficiency Syndrome (AIDS), stroke, epilepsy, Parkinson's disease, multiple sclerosis, neurodegenerative disease, trauma, depression, Alzheimer's disease, migraine, pain, or a seizure disorder. Classes of neuropharmaceutical agents include proteins, antibiotics, adrenergic agents, anticonvulsants, small molecules, nucleotide analogs, chemotherapeutic agents, anti-trauma agents, peptides and other classes of agents used in treatment or prophylaxis of a neurological disorder. Examples of proteins include CD4 (including soluble portions thereof), growth factors (e.g., nerve growth factor and interferon), dopamine decarboxylase and tricosanthin. Examples of antibiotics include amphotericin B, gentamycin sulfate, and pyrimethamine. Examples of adrenergic agents (including blockers) include dopamine and atenolol. Examples of chemotherapeutic agents include

adriamycin, methotrexate, cyclophosphamide, etoposide, and carboplatin. An example of an anticonvulsant which can be used is valproate and an anti-trauma agent which can be used is superoxide dismutase. Examples of peptides are somatostatin analogues and enkephalinase inhibitors. Nucleotide analogs which can be used include azido thymidine (hereinafter AZT), dideoxy Inosine (ddI) and dideoxy Cytidine (ddC).

[0032] The term "agent" is used to describe a compound that has or may have a pharmacological activity. Agents include compounds that are known drugs, compounds for which pharmacological activity has been identified but which are undergoing further therapeutic evaluation, and compounds that are members of collections and libraries that are to be screened for a pharmacological activity.

[0033] A "pharmacological" activity means that an agent exhibits an activity in a screening system that indicates that the agent is or may be useful in the prophylaxis or treatment of a disease. The screening system can be in vitro, cellular, animal or human. Agents can be described as having pharmacological activity notwithstanding that further testing may be required to establish actual prophylactic or therapeutic utility in treatment of a disease.

[0034] An agent is "orally active" if it can exert a pharmacological activity when administered via an oral route.

[0035] A "peripheral tissue" means a tissue other than the CNS.

[0036] A "conjugate" refers to a compound comprising a neuropharmaceutical agent or imaging component and a chemical moiety bound thereto, which moiety by itself or in combination with the neuropharmaceutical agent or imaging component renders the conjugate a substrate for active transport, for example rendering the conjugate to be a substrate for a transporter protein. The chemical moiety may or may not be subject to cleavage from the neuropharmaceutical agent or imaging component upon uptake and metabolism of the conjugate in the patient's body. In other words, the moiety may be cleavably bound to the neuropharmaceutical agent or imaging component or non-cleavably bound to the neuropharmaceutical agent or imaging component. The bond can be a direct (i.e., covalent) bond or the bond can be through a linker. In cases where the bond/linker is cleavable by metabolic processes, the neuropharmaceutical agent or imaging component, or a further metabolite of the neuropharmaceutical agent or imaging component, is the therapeutic or imaging entity. In cases where the bond/linker is not cleavable by metabolic processes, the conjugate itself is the therapeutic or imaging entity. Most typically, the conjugate comprises a prodrug having a metabolically cleavable moiety, where the conjugate itself does not have pharmacological activity but the component to which the moiety is cleavably bound does have pharmacological activity. Typically, the moiety facilitates therapeutic use of the neuropharmaceutical agent or imaging component by promoting uptake of the conjugate via a transporter. Thus, for example, a conjugate comprising a neuropharmaceutical agent and a conjugate moiety may have a V_{max} for a transporter that is at least 2, 5, 10, 20, 50 or 100-fold higher than that of the neuropharmaceutical agent or imaging component alone. A conjugate moiety can itself be a substrate for a transporter or can become a substrate when linked to the neuropharmaceutical agent or

imaging component. Examples of preferred conjugate moieties are para-amino-hippurate, ochratoxin A, estrone-3-sulphate, benzylpenicillin, and probenecid. Thus, a conjugate formed from a neuropharmaceutical agent or imaging component and a conjugate moiety can have higher CNS uptake activity than either the neuropharmaceutical agent, the imaging component, or the conjugate moiety alone.

[0037] A "xenobiotic" is a molecule that is not naturally produced by mammalian organisms.

[0038] A "neuropharmacological" activity means that a neuropharmaceutical agent exhibits an activity in a screening system that indicates that the neuropharmaceutical agent is or may be useful in the prophylaxis or treatment of a neurological disease. The screening system can be in vitro, cellular, animal or human. Neuropharmaceutical agents can be described as having neuropharmacological activity notwithstanding that further testing may be required to establish actual prophylactic or therapeutic utility in treatment of a disease.

[0039] V_{max} and K_m of a compound for a transporter are defined in accordance with convention. V_{max} is the number of molecules of compound transported per second at saturating concentration of the compound. K_m is the concentration of the compound at which the compound is transported at half of V_{max} . When the goal is to transport an agent, conjugate or conjugate moiety into the CNS, a high V_{max} for an influx transporter such as OAT3 is generally desirable. Likewise for the same goal, a low value of K_m is typically desirable for transport of a compound present at low blood concentrations. In some cases a high value of K_m is acceptable for the transport of compounds present at high concentrations in the blood. For these reasons, the intrinsic capacity of a compound to be transported by a particular transporter is usually expressed as the ratio V_{max} of the compound/ V_{max} of a reference compound known to be a substrate for the transporter. V_{max} is affected both by the intrinsic turnover rate of a transporter (molecules/transporter protein) and transporter density in the plasma membrane, which depends on expression level. In certain instances, the goal is to avoid transport into the CNS. In these instances, low V_{max} for all influx transporters and a high V_{max} for all efflux transporters expressed in the blood brain barrier is desirable.

[0040] "EC50", or "effective concentration 50", is a measurement of the substrate concentration that results in a turnover rate 50% of the maximal turnover rate for the substrate ($0.5 V_{max}$).

[0041] A plasma membrane containing a monolayer of cells in physical contact with each other and having different sets of proteins embedded in the plasma membranes facing either side of the monolayer is described as being "polarized". For example, brain microvessel endothelial cells in the blood brain barrier have a luminal side facing capillaries and exposed to blood, and an abluminal side facing cells of the central nervous system and exposed to cerebrospinal fluid. The luminal plasma membrane contains a different set of transmembrane and membrane-associated components than the abluminal plasma membrane of the same cell. Brain microvessel endothelial cells in culture can also be polarized, where the cells form a monolayer in culture that has a luminal and abluminal side. MDCK cells, when grown on filter membranes in transwell dishes, form a polarized monolayer in which one side of the monolayer is the apical

side and the other is the basolateral side. Similarly, choroid plexus epithelial cells in the blood brain barrier have a luminal side facing capillaries and exposed to blood, and an abluminal side facing cells of the central nervous system and exposed to cerebrospinal fluid. The luminal plasma membrane contains a different set of transmembrane and membrane-associated components than the abluminal plasma membrane of the same cell. Choroid plexus epithelial cells in culture can also be polarized, where the cells form a monolayer in culture that has a luminal and abluminal side.

[0042] "Sustained release" refers to release of a therapeutic or prophylactic amount of a drug or an active metabolite thereof over a period of time that is longer than a conventional formulation of the drug. For oral formulations, the term "sustained release" typically means release of the drug within the GI tract lumen over a period of from about 2 to about 30 hours, more typically over a period of about 4 to about 24 hours. Sustained release formulations achieve therapeutically effective concentrations of the drug in the systemic blood circulation over a prolonged period of time relative to that achieved by oral administration of a conventional formulation of the drug. "Delayed release" refers to release of the drug or an active metabolite thereof into the gastrointestinal lumen after a delay time period, typically a delay of about 1 to about 12 hours, relative to that achieved by oral administration of a conventional formulation of the drug.

[0043] The phrase "specifically binds" when referring to a substrate or ligand of an OAT3 transporter refers to a specific interaction between a substrate or ligand and the OAT3 transporter in which the substrate or ligand binds preferentially with an OAT3 transporter and does not bind in a significant amount to most or any other proteins present in a biological sample. A substrate or ligand that specifically binds to an OAT3 transporter often has an association constant of $10 \cdot 10^3 \text{ M}^{-1}$, 10^5 M^{-1} , 10^6 M^{-1} or 10^7 M^{-1} , preferably 10^8 M^{-1} to 10^9 M^{-1} or higher. However, some substrates or ligands of OAT3 transporters have much lower affinities and yet the binding can still be shown to be specific. Substrates of OAT3 can specifically bind to OAT3 and other proteins such as efflux transporters without specifically binding to other proteins.

[0044] " P_{app} ", or "apparent permeability", is a value that reflects the permeability of a test compound through a cell layer such as a polarized monolayer. The equation for determining P_{app} is as follows:

$$P_{app} = \frac{V \cdot dC}{A \cdot C_0 \cdot dt} \text{ (cm/sec)}$$

[0045] where,

[0046] V=volume of receiving chamber (in cm^3 , i.e., ml);

[0047] dC/dt =steady state rate of appearance of applied compound in receiving chamber after primary lag time (in $\mu\text{M}/\text{sec}$);

[0048] C_0 =concentration of compound in the donor chamber (in μM)

[0049] A=area of the cell layer (in cm^2)

[0050] "Allelic variants" at the DNA level are the result of genetic variation between individuals of the same species. Some allelic variants at the DNA level that cause substitution, deletion or insertion of amino acids in proteins encoded by the DNA result in corresponding allelic variation at the protein level.

[0051] "Cognate forms" of a gene refers to variation between structurally and functionally related genes between species. For example, the human gene showing the greatest sequence identity and closest functional relationship to a mouse gene is the human cognate form of the mouse gene.

[0052] For sequence comparison, typically one sequence acts as a reference sequence, to which test sequences are compared. When using a sequence comparison algorithm, test and reference sequences are input into a computer, subsequence coordinates are designated, if necessary, and sequence algorithm program parameters are designated. The sequence comparison algorithm then calculates the percent sequence identity for the test sequence(s) relative to the reference sequence, based on the designated program parameters.

[0053] Optimal alignment of sequences for comparison can be conducted, e.g., by the local homology algorithm of Smith & Waterman, *Adv. Appl. Math.* 2:482 (1981), by the homology alignment algorithm of Needleman & Wunsch, *J. Mol. Biol.* 48:443 (1970), by the search for similarity method of Pearson & Lipman, *Proc. Nat'l. Acad. Sci. USA* 85:2444 (1988), by computerized implementations of these algorithms (GAP, BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group, 575 Science Dr., Madison, Wis.), or by visual inspection (see generally Ausubel et al., *supra*).

[0054] Another example of an algorithm that is suitable for determining percent sequence identity and sequence similarity is the BLAST algorithm, which is described in Altschul et al., *J. Mol. Biol.* 215:403-410 (1990). Software for performing BLAST analyses is publicly available through the National Center for Biotechnology Information web site. This algorithm involves first identifying high scoring sequence pairs (HSPs) by identifying short words of length W in the query sequence, which either match or satisfy some positive-valued threshold score T when aligned with a word of the same length in a database sequence. T is referred to as the neighborhood word score threshold (Altschul et al., *supra*). These initial neighborhood word hits act as seeds for initiating searches to find longer HSPs containing them. The word hits are then extended in both directions along each sequence for as far as the cumulative alignment score can be increased. Cumulative scores are calculated using, for nucleotide sequences, the parameters M (reward score for a pair of matching residues; always >0) and N (penalty score for mismatching residues; always <0). For amino acid sequences, a scoring matrix is used to calculate the cumulative score. Extension of the word hits in each direction are halted when: the cumulative alignment score falls off by the quantity X from its maximum achieved value; the cumulative score goes to zero or below, due to the accumulation of one or more negative-scoring residue alignments; or the end of either sequence is reached. For identifying whether a nucleic acid or polypeptide is within the scope hereof, the default parameters of the BLAST programs are suitable. The BLASTN program (for nucleotide

sequences) uses as defaults a word length (W) of 11, an expectation (E) of 10, M=5, N=-4, and a comparison of both strands. For amino acid sequences, the BLASTP program uses as defaults a word length (W) of 3, an expectation (E) of 10, and the BLOSUM62 scoring matrix. The TBLASTN program (using protein sequence for nucleotide sequence) uses as defaults a word length (W) of 3, an expectation (E) of 10, and a BLOSUM 62 scoring matrix. (see Henikoff & Henikoff, Proc. Natl. Acad. Sci. USA 89:10915 (1989)).

[0055] In addition to calculating percent sequence identity, the BLAST algorithm also performs a statistical analysis of the similarity between two sequences (see, e.g., Karlin & Altschul, Proc. Nat'l. Acad. Sci. USA 90:5873-5787 (1993)). One measure of similarity provided by the BLAST algorithm is the smallest sum probability (P(N)), which provides an indication of the probability by which a match between two nucleotide or amino acid sequences would occur by chance. For example, a nucleic acid is considered similar to a reference sequence if the smallest sum probability in a comparison of the test nucleic acid to the reference nucleic acid is less than about 0.1, more preferably less than about 0.01, and most preferably less than about 0.001.

DETAILED DESCRIPTION

[0056] I. General

[0057] OAT3 is shown herein to be expressed in brain microvessel endothelial cells and in choroid plexus epithelial cells. This finding can be used to generate or isolate conjugates and agents having neuropharmacological or imaging activity useful for treatment, prophylaxis or diagnosis of neurological diseases. The invention provides methods of identifying agents, conjugates or conjugate moieties that are substrates for OAT3. For therapeutic purposes, agents or conjugates having inherent neuropharmacologic activity can be screened to determine whether they are substrates for OAT3. Alternatively, a conjugate moiety lacking such activity can be screened, and linked to a neuropharmacologic agent after screening. Agents or conjugates that both have neuropharmacologic activity and are substrates for OAT3 are preferentially transported into the CNS via OAT3 transporters after administration to a patient. Such an agent or conjugate by itself or in combination with another agent is effective in treatment or prophylaxis of a neurological disease. An analogous approach is used for imaging features of the brain. Agents and conjugates that have an imaging component and are substrates for OAT3 are preferentially transported into the CNS via OAT3 transporters. The imaging component is then detected by various methods such as detecting radioactive decay of the imaging component. The agents and conjugates can be used to image brain tumors overexpressing the OAT3 transporter. Optionally, the agents or conjugates have inherent affinity for, or are provided with a conjugate moiety that confers affinity for, a particular antigen or cell type within the brain. For example, the agents or conjugates can be provided with a targeting moiety to A β to allow imaging of plaques in Alzheimer's patients.

[0058] II. OAT3 Transporter

[0059] The family of organic anion transporters (OATs) and organic cation transporters contains at least 25 members in humans (SLC21A1-12; SLC22A1-13). Organic anion and

cation transporters have 12 putative transmembrane domains, with both the amino and carboxy termini located on the cytoplasmic side. A sub-family of organic anion transporters (OAT1-4) are expressed in the kidney, liver, and barrier tissues such as intestine, testis, and blood brain barrier. Members of the OAT family recognize a diverse group of organic anion compounds such as ochratoxin A, para-amino-hippurate, and estrone-3-sulphate. OAT transport is coupled to the countertransport of metabolic anions such as ketoglutarate.

[0060] It is now shown that OAT3 is expressed in non-human brain microvessel endothelial cells. OAT3 is expressed at a level more than 15-fold higher than other OAT family transporters with similar substrate specificity. OAT3 is also expressed in choroid plexus epithelial cells. It is desirable to generate agents, conjugates, and conjugate moieties for transport into the CNS that have activity for OAT3 due to this high expression level. The GenBank accession number for human OAT3 is AF097491 (incorporated by reference). Unless otherwise apparent from the context, reference to a transporter includes the amino acid sequence described in or encoded by the GenBank reference number AF097491 (SEQ ID NO:1), and, allelic, cognate and induced variants and fragments thereof retaining essentially the same transporter activity. Usually such variants show at least 90% sequence identity to the exemplary Genbank nucleic acid or amino acid sequence.

[0061] III. Methods of Screening to Identify OAT3 Substrates

[0062] Agents known or suspected to have a neuropharmaceutical activity or to comprise an imaging component can be screened directly for their capacity to act as substrates of OAT3. Alternatively, conjugate moieties can be screened as substrates, and the conjugate moieties are then linked to a neuropharmaceutical agent or imaging component. In such methods, the conjugate moieties can optionally be linked to a neuropharmaceutical agent or imaging component, or other molecule during the screening process. If another molecule is used in place of a neuropharmaceutical agent or imaging component, the molecule can be chosen to resemble the structure of a neuropharmaceutical agent or imaging component ultimately intended to be linked to the conjugate moiety for neuropharmaceutical use. Alternatively, a conjugate moiety can be screened for a substrate activity alone and linked to a neuropharmaceutical agent or imaging component after screening.

[0063] Preferred substrates for OAT3 are xenobiotic molecules containing one or more ring structures and one or more anionic groups such as carboxylates, sulphates, and phosphates. Table 1 lists examples of substrates of OAT3. The structure of each compound listed in Table 1 is depicted in FIG. 1.

TABLE 1

SUBSTRATES	Reported Affinity for OAT3
Para-amino-hippurate	0.2 mM
Ochratoxin A	0.05 mM
Estrone-3-sulphate	0.04 mM

TABLE 1-continued

SUBSTRATES	Reported Affinity for OAT3
Benzylpenicillin	0.3 mM
Probenecid	0.05 mM

[0064] Para-amino-hippurate, ochratoxin A, estrone-3-sulphate, benzyl-penicillin, and probenecid are examples of OAT3 substrates that are candidates for conjugation to therapeutic neuropharmaceutical agents, cytotoxic neuropharmaceutical agents and imaging components.

[0065] In some screening methods, the cells are transfected with DNA encoding the OAT3 transporter. HEK (human embryonic kidney) and CHO (Chinese hamster ovary) cells, for example, are suitable for transfection. Oocytes can be injected with OAT3 cRNA to express OAT3 transporter. In some methods, the only transporter expressed by the cells is the OAT3 transporter. In other methods, cells express OAT3 in combination with other transporters. In still other methods, agents, conjugate moieties or conjugates are screened on different cells expressing different transporters. Agents, conjugate moieties or conjugates can be screened either for specificity for the OAT3 transporter or for transport into cells endogenously expressing a plurality of transporters. In some methods, the results of a screening method (e.g., a competition uptake, exchange or direct uptake assay) using a cell expressing the OAT3 transporter can be compared with the results of a control cell(s) lacking the OAT3 transporter or in the presence of a specific inhibitor of the OAT3 transporter.

[0066] In some methods, cells endogenously expressing the OAT3 transporter are used. Brain microvessel endothelial cells, for example, endogenously express the OAT3 transporter, as demonstrated in Example 1. Agents, conjugate moieties or conjugates can be screened for transport into cultured brain microvessel endothelial cells. Passaging cultures of brain microvessel endothelial cells typically causes the cells to lose differentiation characteristics such as the ability to form tight junctions. The propensity of passaged cells to lose differentiation characteristics can be avoided through the use of brain microvessel endothelial cells that are transformed with an SV40 large T antigen. See Terasaki et al., *Drug Discovery Today* 8:944-954 (2003). Inducible expression of the SV40 large T antigen allows cells to divide when the antigen is expressed and differentiate when the antigen is not expressed. Brain microvessel endothelial cells can be isolated from animals transgenic for the SV40 large T antigen, which can be expressed in a temperature-sensitive fashion. The cells are stimulated to divide by being cultured at the temperature at which the antigen is expressed. Once the cells have formed a monolayer, they are placed at a temperature at which the antigen is not expressed, causing the cells to stop dividing and differentiate. Differentiation results in the formation of tight junctions and the polarization of the plasma membranes. Monolayers of polarized cells are tested for the ability to transport agents, conjugates or conjugate moieties.

[0067] In other methods, choroid plexus epithelial cells endogenously expressing the OAT3 transporter are used. Agents, conjugate moieties or conjugates can be screened

for transport into cultured choroid plexus epithelial cells. Passaging cultures of choroid plexus epithelial cells typically causes the cells to lose differentiation characteristics such as the ability to form tight junctions. The propensity of passaged cells to lose differentiation characteristics can be avoided through the use of choroid plexus epithelial cells that are transformed with an SV40 large T antigen. See Zheng & Zhao, Establishment and characterization of an immortalized Z310 choroidal epithelial cell line from murine choroid plexus, *Brain Res.* 958(2):371-80 (2002). Inducible expression of the SV40 large T antigen allows cells to divide when the antigen is expressed and differentiate when the antigen is not expressed. Choroid plexus epithelial cells can be isolated from animals transgenic for the SV40 large T antigen, which can be expressed in a temperature-sensitive fashion. The cells are stimulated to divide by being cultured at the temperature at which the antigen is expressed. Once the cells have formed a monolayer, they are placed at a temperature at which the antigen is not expressed, causing the cells to stop dividing and differentiate. Differentiation results in the formation of tight junctions and the polarization of the plasma membranes. Monolayers of polarized cells are tested for the ability to transport agents, conjugates or conjugate moieties.

[0068] In some methods, the ability of an agent, conjugate or conjugate moiety to specifically bind to an OAT3 transporter is tested. A known substrate of the OAT3 transporter and the agent, conjugate or conjugate moiety are added to cells expressing the OAT3 transporter. The amount or rate of transport of the substrate in the presence of the agent, conjugate or conjugate moiety is compared to the amount or rate of transport of the agent, conjugate or conjugate moiety in the absence of the test compound. If the amount or rate of transport of the substrate is decreased by the presence of the agent, conjugate or conjugate moiety, the agent, conjugate or conjugate moiety binds the OAT3 transporter. Agents, conjugates or conjugate moieties that bind the OAT3 transporter can be further analyzed to determine if they are transported by the OAT3 transporter or only adhere to the exterior of the transporter. Agents, conjugates or conjugate moieties that are transported by the OAT3 transporter can be further tested to determine if they are transported from one side of a monolayer of polarized cells to the other side, such as a monolayer of brain microvessel endothelial cells or a monolayer of choroid plexus epithelial cells. Agents and conjugates having neuropharmaceutical activity and that are transported by the OAT3 transporter can be used to form pharmaceutical compositions. Conjugate moieties that are transported by the OAT3 transporter can be linked to a therapeutic or cytotoxic neuropharmaceutical agent or an imaging component.

[0069] Transport of a compound into a cell can be detected by detecting a signal from within a cell from any of a variety of reporters. The reporter can be as simple as a label such as a fluorophore, a chromophore, or a radioisotope. Confocal imaging can also be used to detect internalization of a label as it provides sufficient spatial resolution to distinguish between fluorescence on a cell surface and fluorescence within a cell; alternatively, confocal imaging can be used to track the movement of compounds over time. In another approach, transport of a compound is detected using a reporter that is a substrate for an enzyme expressed within a cell. Once the compound is transported into the cell, the substrate is metabolized by the enzyme and generates an

optical signal that can be detected. Light emission can be monitored by commercial PMT-based instruments or by CCD-based imaging systems. In addition, assay methods utilizing liquid chromatography-mass spectroscopy (LC-MS-MS) detection of the transported compounds or electrophysiological signals indicative of transport activity are also employed. Mass spectroscopy is a powerful tool because it allows detection of very low concentrations of almost any compound, especially molecules for which a radiolabeled version is not available. It can also be used to distinguish substrates from non-transported ligands. These same detection methods can be used to determine if a compound is transported from one side of a monolayer of polarized cells to the other side by administering the compound to one side of the monolayer and sampling the media on the other side of the monolayer after a predetermined period of time.

[0070] In some methods, multiple agents, conjugates or conjugate moieties are screened simultaneously and the identity of each agent, conjugate or conjugate moiety is tracked using tags linked to the agents, conjugates or conjugate moieties. In some methods, a preliminary step is performed to determine binding of an agent, conjugate or conjugate moiety to a transporter. Although not all agents, conjugates or conjugate moieties that bind to a transporter are substrates of the transporter, observation of binding is an indication that allows one to reduce the number of candidates from an initial repertoire. In some methods, the transport rate of an agent, conjugate or conjugate moiety is tested in comparison with the transport rate of a reference substrate for that transporter. For example, para-amino-hippurate, a natural substrate of OAT3, can be used as a reference. The comparison can be performed in separate parallel assays in which an agent, conjugate or conjugate moiety under test and the reference substrate are compared for uptake on separate samples of the same cells. Alternatively, the comparison can be performed in a competition format in which an agent, conjugate or conjugate moiety under test and the reference substrate are applied to the same cells. Typically, the agent, conjugate or conjugate moiety and the reference substrate are differentially labeled in such assays.

[0071] In comparative assays, the V_{\max} of an agent, conjugate or conjugate moiety tested can be compared with that of a reference substrate. If an agent, conjugate moiety or conjugate has a V_{\max} of at least 1%, 5%, 10%, 20%, and most preferably at least 50% of the reference substrate for the OAT3 transporter, then the agent, conjugate moiety or conjugate is also a substrate for the OAT3 transporter. If transport of the agent, conjugate moiety or conjugate into the CNS is desired, a higher V_{\max} of the agent, conjugate moiety or conjugate relative to that of the reference substrate is preferred. Therefore, agents, conjugate moieties or conjugates having V_{\max} 's of at least 1%, 5%, 10%, 20%, 50%, 100%, 150% or 200% (i.e., two-fold) of the V_{\max} of a reference substrate (e.g., para-amino-hippurate) for the transporter are screened in some methods. The components to which conjugate moieties are linked can by themselves show little or no detectable substrate activity for the transporter (e.g., V_{\max} relative to that of a reference substrate of less than 0.1% or 1%). Preferred agents, conjugates or conjugate moieties have a V_{\max} for OAT3 that is at least 5% of the V_{\max} for OAT3 of para-amino-hippurate. Preferred conjugates comprising a neuropharmaceutical agent or

imaging component linked to a conjugate moiety preferably have a greater V_{\max} for OAT3 than the neuropharmaceutical agent or imaging component alone.

[0072] Having determined that an agent, conjugate or conjugate moiety is a substrate for OAT3, a further screen can be performed to determine its therapeutic activity in treatment or prophylaxis of a disease, or its cytotoxic activity against brain tumor cells. Usually the disease is neurological (i.e., the pathology occurs in the CNS). Alternatively, the diseased tissue is non-CNS tissue but is responsive to treatment by an agent that exerts a pharmacological effect on the CNS that in turn causes an effect on the diseased non-CNS tissue, such as an effect caused by the release of hormones from the CNS. Diseases of this type are also considered to be diseases of the CNS unless otherwise apparent from context. If the agent, conjugate or conjugate moiety does not have inherent therapeutic or cytotoxic activity, it is first linked to another chemical component having such therapeutic or cytotoxic properties. The agent, conjugate or conjugate moiety is then contacted with cells expressing OAT3. The contacting can be performed either on a population of cells in vitro, or the brain microvessel endothelial cells of a test animal via administration of the agent, conjugate or conjugate moiety to a test animal. Similarly, the contacting can be performed on choroid plexus epithelial cells of a test animal via administration of the agent, conjugate or conjugate moiety to a test animal. The therapeutic or cytotoxic activity of the agent, conjugate or conjugate moiety is then determined from established protocols for that particular disease. Optionally, the effect of the agent, conjugate or conjugate moiety can be compared with a placebo.

[0073] A further screen can be performed to determine toxicity of the agent, conjugate, or conjugate moiety to normal cells. The agent, conjugate or conjugate moiety is administered to a laboratory animal that is preferably in an undiseased state. Various tissues of the animal, such as liver, kidney, heart and brain are then examined for signs of pathology. Cells in the animal can also be analyzed for uptake of the agent, conjugate, or conjugate moiety.

[0074] IV. Iterative Modification and Testing of OAT3 Substrates

[0075] Having determined that an agent, conjugate or conjugate moiety is a substrate for OAT3, the agent, conjugate or conjugate moiety can be modified to improve its properties as a substrate. The modified agent, conjugate or conjugate moiety is then tested for transport by OAT3. Modified agents, conjugates or conjugate moieties that are transported by OAT3 at a higher V_{\max} compared to the unmodified agent, conjugate or conjugate moiety are preferred. The process of modifying agents, conjugates or conjugate moieties and testing for transport by OAT3 can be repeated until a desired level of transport is reached.

[0076] Agents, conjugates or conjugate moieties that are substrates of OAT3 can also be modified for decreased capacity to be transported out of cells by efflux transporters. An agent, conjugate or conjugate moiety transported by OAT3 is assayed to determine whether it is also a substrate for one or more efflux transporters. If the agent, conjugate or conjugate moiety is transported by an efflux transporter, the agent, conjugate or conjugate moiety is modified and tested for both reduced transport by an efflux transporter and retention of OAT3 substrate activity.

[0077] In some instances, the specific efflux transporter responsible for transporting an agent, conjugate or conjugate moiety is known. The agent, conjugate or conjugate moiety is modified, preferably by addition of a chemical group that differs in chemical characteristics from other known substrates of the efflux transporter. The modified agent, conjugate or conjugate moiety is then tested for retained capacity to be transported by OAT3 and a diminished capacity to be transported by an efflux transporter. It is not necessary that the modified agent, conjugate or conjugate moiety retain the same kinetic properties of OAT3 transporter substrate as the unmodified agent, conjugate or conjugate moiety as long as some OAT3 substrate activity is retained. Examples of efflux transporters are the P-glycoprotein (PgP), multidrug resistance protein (MRP1), and breast cancer resistance protein (BCRP). Preferred agents, conjugates or conjugate moieties have an OAT3 transport:efflux transport ratio of at least 1.1:1.0, more preferably, 2.0:1.0, and more preferably 5.0:1.0 and more preferably 10.0:1.0 or higher at a given concentration of agent, conjugate or conjugate moiety.

[0078] Efflux transporter activity can be measured in several ways. First, functional assays can be performed in which interaction of compounds with efflux transporters is measured by stimulation of efflux transporter ATPase activity in cellular membrane fragments or vesicles. Second, competition assays can be performed in which test compounds compete with known efflux substrates in whole cells. Third, direct transport assays can be performed in which the transport of compounds is measured across a polarized monolayer of cells. Other assays besides these three can also be used to directly or indirectly measure the efflux substrate characteristics of a test compound.

[0079] The efflux transporter ATPase assay is based on the fact that most efflux substrates increase the ATPase activity of efflux transporters upon binding. In one type of assay, Baculovirus membrane fragments or vesicles containing an efflux transporter such as PgP, as well as control membrane fragments or vesicles not containing the efflux transporter, are either prepared or obtained from commercial suppliers. The ATPase activity of the membrane fragments or vesicles is measured in the presence of various concentrations of the test compound. An agent, conjugate, or conjugate moiety that is transported by OAT3 is added to the ATPase assay reaction and the amount of ATPase activity is measured at various concentrations of agent, conjugate, or conjugate moiety. Parallel experiments are performed in which ATPase activity is measured under addition of the same concentrations of modified agent, conjugate, or conjugate moiety that retain OAT3 substrate activity. Reduced ATPase activity caused by the modified agent, conjugate, or conjugate moiety compared to the unmodified agent, conjugate, or conjugate moiety indicates that the modified agent, conjugate, or conjugate moiety is a better candidate for retention in the CNS.

[0080] In the competition assay, the test compound is assayed for competition with a known efflux substrate. For example, calcein-AM is a non-fluorescent compound that is a substrate of PgP and MRP1. Calcein-AM is initially loaded into the cells, for example, by transport by passive diffusion. Cells expressing these efflux transporters actively efflux nearly all of the calcein-AM that is present in the cells. However, when other efflux transporter substrates are present, these other substrates compete with calcein-AM for

efflux, resulting in more calcein-AM accumulating inside the cells. Intracellular esterases convert the non-fluorescent calcein-AM to fluorescent calcein which can be measured spectrophotometrically. An agent, conjugate, or conjugate moiety that is transported by OAT3 is loaded into efflux transporter-containing cells by either OAT3 transport or passive diffusion. Calcein-AM is also loaded into the cells by active transport or transport by passive diffusion. Accumulation of calcein-AM is measured and compared to the amount of accumulation in the absence of the agent, conjugate, or conjugate moiety. Parallel experiments are performed in which a modified agent, conjugate, or conjugate moiety that is transported by OAT3 is loaded into the cells. Accumulation of calcein-AM is measured and compared to the amount of accumulation in the absence of the modified agent, conjugate, or conjugate moiety. Decreased calcein-AM accumulation inside the cells caused by the presence of a modified agent, conjugate, or conjugate moiety compared to calcein-AM accumulation in the presence of unmodified agent, conjugate, or conjugate moiety indicates that the modified agent, conjugate, or conjugate moiety is a better candidate for retention inside the CNS.

[0081] The cells used for competition assays can be cells that either express a high endogenous level of the efflux transporter of interest or are transformed with an expression vector containing the efflux transporter gene. Suitable cell lines for efflux assays are, for example, HEK and MDCK cell lines into which the PgP gene has been transfected, or MES-SA/Dx5 uterine sarcoma cells grown in the presence of 500 nM doxorubicin, which express a high endogenous level of PgP. These cells can optionally be transfected with the OAT3 transporter gene. Preferred cells express both one or more efflux transporter genes such as PgP and the OAT3 gene, either endogenously or through transfection of expression vectors.

[0082] A third type of efflux transporter assay is the cellular transwell monolayer efflux assay. In this assay, cells expressing efflux transporters, such as MDCK cells containing the TReX-PgP expression vector, are seeded and grown in transwell dishes on filter membranes made of substances such as polycarbonate. The cells form a polarized monolayer. The transwell dishes have apical and basolateral chambers that are separated by the filter membrane on which the polarized monolayer is situated. Assays are performed by placing a test compound in either the apical or basolateral chamber, followed by sampling the opposite chamber after a predetermined period of time such as 60-120 minutes and measuring the amount of the test compound. The test compound can be measured by methods such as radiolabel detection or LC-MS-MS analysis. Assays are performed in the presence and absence of an efflux transporter inhibitor or competitor. Efflux transporter inhibitors or competitors increase apical to basolateral transport and decrease basolateral to apical transport of compounds that are efflux transporter substrates. Apparent permeability (P_{app}) of test compounds is measured. Test compounds that are substrates of efflux transporters generate a P_{app} (basolateral to apical)/ P_{app} (apical to basolateral) ratio of greater than 2.0, while test compounds that are not substrates generate a ratio of 1.5 or less. Test compounds that generate ratios between 1.5 and 2.0 require additional testing to determine if they are efflux transporter substrates. An agent, conjugate, or conjugate moiety that is an OAT3 substrate and also generates a ratio of greater than 2.0 can be modified. A modified agent,

conjugate, or conjugate moiety that retains OAT3 substrate activity and generates a lower ratio compared to the unmodified agent, conjugate, or conjugate moiety indicates that the modified agent, conjugate, or conjugate moiety is a better candidate for retention inside the CNS.

[0083] An additional screen can be performed to determine whether agents, conjugates or conjugate moieties have substantial capacity for passive diffusion across the brain microvessel endothelial cells making up the blood brain barrier. Such assays can be performed using cells lacking OAT3 transporters. That is, the agents, conjugates or conjugate moieties are exposed to cells that lack OAT3 transporters, and the amount of agents, conjugates or conjugate moieties that are present inside the cell is measured.

[0084] V. Modification of Compounds Having Non-Neuropharmacological Activity

[0085] In some instances it is desirable to modify an agent to reduce its capacity to be transported from the blood into the brain. Reduced capacity to enter the brain is desirable for agents having a pharmacological activity that is useful in a tissue outside the CNS, but which causes undesired side effects when the agent enters the CNS. Most typically, such agents are drugs administered to treat a non-neurological disease, and which exert a useful therapeutic pharmacological effect on cells, tissues, or molecules located outside of the CNS. When such drugs are transported from the blood into the brain, serious side effects can occur. Many known drugs exhibit undesirable side effects from penetrating the CNS. Examples include drowsiness experienced by patients taking antihistamines, nonsteroidal anti-inflammatory drugs (NSAIDS), anti-asthmatics and antihypertensives.

[0086] The methods are performed on an agent having an intended site of pharmacological activity that is located outside of the CNS. The agent is known or suspected to enter the CNS. In some instances, the agent is known to be transported by OAT3. The agent is covalently attached to a conjugate moiety and the resulting conjugate is tested for transport into the brain. The assay can be performed on brain microvessel endothelial cells, such cells transformed with an OAT3 expression vector, a polarized monolayer of such cells, or an actual blood brain barrier via administration to a test animal. The assay also can be performed on choroid plexus epithelial cells, such cells transformed with an OAT3 expression vector, a polarized monolayer of such cells, or an actual blood brain barrier via administration to a test animal. Transport of the conjugate is then compared with transport of the agent alone (i.e., without the conjugate moiety). Conjugates having a lower V_{max} for transport than the agent alone are less likely to exhibit undesirable CNS side effects caused by unwanted transport from the blood into the brain. For example, preferred conjugates include those having a lower V_{max} for transport by OAT3 than the agent alone.

[0087] Some methods comprise providing an agent having a pharmacological activity, wherein the pharmacological activity is useful for treating a disease present in a tissue other than the CNS, and the pharmacological activity results in undesired side effects in the CNS if the agent enters the CNS, modifying the agent, providing a cell expressing at least one transporter protein that transports substrates across the blood brain barrier, contacting the cell with the modified agent, and determining whether the modified agent passes through the plasma membrane via the transporter protein

with a lower V_{max} than the agent, a lower V_{max} indicating that the modification decreases the capacity of the modified agent relative to the agent to cross the blood brain barrier, thereby decreasing undesired side effects in the CNS. In some methods the at least one transporter protein is OAT3. In some methods the cell is transformed or injected with a nucleic acid encoding a transporter or the cell is a brain microvessel endothelial cell. In some methods the cell is transformed or injected with a nucleic acid encoding a transporter or the cell is a choroid plexus epithelial cell. In some methods the modifying step comprises linking the agent to a conjugate moiety to form a conjugate, preferably wherein the conjugate moiety is an inhibitor of the OAT3 transporter.

[0088] Other methods comprise providing an agent having a pharmacological activity, wherein the pharmacological activity is useful for treating a disease present in a tissue other than the CNS, and the pharmacological activity results in undesired side effects in the CNS if the agent enters the CNS, modifying the agent, providing a cell expressing at least one efflux transporter protein that transports substrates out of the CNS, contacting the cell with the modified agent, and determining whether the modified agent is transported by the at least one efflux transporter protein with a higher V_{max} than the agent, a higher V_{max} indicating that the modification increases the capacity of the modified agent relative to the agent to be transported out of the CNS, thereby decreasing undesired side effects in the CNS. In some methods the at least one efflux transporter protein is P-glycoprotein (PgP), multidrug resistance protein (MRP1), or breast cancer resistance protein (BCRP). In some methods the cell is transformed or injected with a nucleic acid encoding an efflux transporter or the cell is a brain microvessel endothelial cell, a choroid plexus epithelial cell, a kidney-derived cell, or a uterine sarcoma cell. In some methods the modifying step comprises linking the agent to a conjugate moiety to form a conjugate, preferably wherein the conjugate moiety is a substrate of the efflux transporter.

[0089] VI. Sources of Neuropharmaceutical Agents, Imaging Components, and Conjugate Moieties

[0090] Therapeutic neuropharmaceutical agents, cytotoxic neuropharmaceutical agents, imaging components and conjugate moieties can be obtained from natural sources such as, e.g., marine microorganisms, algae, plants, and fungi. Alternatively, these compounds can be from combinatorial libraries, including peptides or small molecules, or from existing repertoires of chemical compounds synthesized in industry, e.g., by the chemical, pharmaceutical, environmental, agricultural, marine, cosmeceutical, drug, and biotechnological industries. Neuropharmaceutical compounds can include proteins, antibiotics, adrenergic agents, anticonvulsants, small molecules, nucleotide analogs, chemotherapeutic agents, anti-trauma agents, peptides and other classes of agents used in treatment or prophylaxis of a neurological disease. Examples of such proteins include CD4 (including soluble portions thereof), growth factors (e.g., nerve growth factor and interferon), dopamine decarboxylase and tricosanthin. Examples of such antibiotics include amphotericin B, gentamycin sulfate, and pyrimethamine. Examples of such adrenergic agents (including blockers) include dopamine and atenolol. Examples of such chemotherapeutic agents include adriamycin, methotrexate, cyclophosphamide, etoposide, and carboplatin. An example of an anticonvulsant

which can be used is valproate and an anti-trauma agent which can be used is superoxide dismutase. Examples of such peptides are somatostatin analogues and enkephalinase inhibitors. Nucleotide analogs which can be used include azido thymidine (hereinafter AZT), dideoxy Inosine (ddI) and dideoxy cytosine (ddc).

[0091] Typically if an agent is being screened as a substrate, the agent is known or suspected to have an inherent therapeutic neuropharmaceutical, cytotoxic neuropharmaceutical or imaging activity. If a conjugate is being screened, the conjugate usually comprises such an agent or component. If a conjugate moiety is being screened, the conjugate moiety typically lacks a therapeutic, cytotoxic or imaging activity and an agent or component that has this activity is added after screening.

[0092] Suitable cytotoxic agents for incorporation into conjugates or linkage to conjugate moieties after screening include platinum, nitrosourea, nitrogen mustard, nitroimidazole, and a phosphoramidate group that is only cytotoxic to brain tumor cells. The choice of imaging component depends on the means of detection. For example, a fluorescent imaging component is suitable for optical detection. A paramagnetic imaging component is suitable for topographic detection without surgical intervention. Radioactive labels can also be detected using positron emission tomography or single photon emission computed tomography.

[0093] The agents, conjugates or conjugate moieties to be screened, optionally linked to a neuropharmaceutical agent or an imaging component if not inherently present, are preferably small molecules having molecular weights of less than 1000 Da and preferably less than 500 Da.

[0094] VII. Linkage of Neuropharmaceutical Agents or Imaging Components to Substrates

[0095] Conjugates can be prepared by either by direct conjugation of a neuropharmaceutical agent or an imaging component to a substrate of OAT3 with a covalent bond (optionally cleavable in vivo), or by covalently coupling a difunctionalized linker precursor with the neuropharmaceutical agent or imaging component and substrate. The linker precursor is selected to contain at least one reactive functionality that is complementary to at least one reactive functionality on the neuropharmaceutical agent or imaging component and at least one reactive functionality on the substrate. Optionally, the linker is cleavable. Suitable complementary reactive groups are well known in the art as illustrated below:

COMPLEMENTARY BINDING CHEMISTRIES		
First Reactive Group	Second Reactive Group	Linkage
hydroxyl	carboxylic acid	ester
hydroxyl	haloformate	carbonate
thiol	carboxylic acid	thioester
thiol	haloformate	thiocarbonate
amine	carboxylic acid	amide
hydroxyl	isocyanate	carbamate
amine	haloformate	carbamate
amine	isocyanate	urea
carboxylic acid	carboxylic acid	anhydride

-continued

COMPLEMENTARY BINDING CHEMISTRIES		
First Reactive Group	Second Reactive Group	Linkage
hydroxyl	phosphorus acid	phosphonate or phosphate ester

[0096] The same methods of chemical modification can be used to form conjugates for the purpose of inhibiting transport into the CNS, for inhibiting efflux from the CNS, or for enhancing efflux from the CNS.

[0097] VIII. Pharmaceutical Compositions

[0098] The above screening processes can identify one or more types of compounds that can be incorporated into pharmaceutical compositions. These compounds include agents that are both substrates for OAT3 and have an inherent neuropharmaceutical activity or imaging activity. The compounds also include conjugates in which a neuropharmaceutical agent or imaging component is linked to a substrate for OAT3. Conjugates comprising an agent with a pharmacological activity and a conjugate moiety having decreased substrate capacity for OAT3 relative to the agent alone are also provided for the purpose of reducing transport of the agent into the CNS, where the agent would confer undesired side effects.

[0099] One or more of the above entities can be combined with pharmaceutically-acceptable, non-toxic carriers or diluents, which are defined as vehicles commonly used to formulate pharmaceutical compositions for animal or human administration. The diluent is selected so as not to affect the biological activity of the combination. Examples of such diluents are distilled water, buffered water, physiological saline, phosphate buffered saline (PBS), Ringer's solution, dextrose solution, and Hank's solution. In addition, the pharmaceutical composition or formulation can also include other carriers, adjuvants, or non-toxic, nontherapeutic, non-immunogenic stabilizers, excipients and the like. The compositions can also include additional substances to approximate physiological conditions, such as pH adjusting and buffering agents, toxicity adjusting agents, wetting agents, detergents and the like (see, e.g., Remington's Pharmaceutical Sciences, Mace Publishing Company, Philadelphia, Pa., 17th ed. (1985); for a brief review of methods for drug delivery, see, Langer, Science 249:1527-1533 (1990); each of these references is incorporated by reference in its entirety).

[0100] Pharmaceutical compositions can be administered orally, intranasally, intradermally, subcutaneously, intrathecally, intramuscularly, topically, intravenously, or injected directly to a site of cancerous tissue. For parenteral administration, the compounds disclosed herein can be administered as injectable dosages of a solution or suspension of the compound in a physiologically acceptable diluent with a pharmaceutical carrier which can be a sterile liquid such as water, oils, saline, glycerol, or ethanol. Additionally, auxiliary substances, such as wetting or emulsifying agents, surfactants, pH buffering substances and the like can be present in the pharmaceutical compositions. Other components of pharmaceutical compositions are those of petroleum, animal, vegetable, or synthetic origin, for example,

peanut oil, soybean oil, and mineral oil. In general, glycols such as propylene glycol or polyethylene glycol are preferred liquid carriers, particularly for injectable solutions.

[0101] Typically, compositions are prepared as injectables, either as liquid solutions or suspensions; solid forms suitable for solution in, or suspension in, liquid vehicles prior to injection can also be prepared. The preparation also can be emulsified or encapsulated in liposomes or micro particles such as polylactide, polyglycolide, or a copolymer thereof for enhanced adjuvant effect, as discussed above (see Langer, *Science* 249, 1527 (1990) and Hanes, *Advanced Drug Delivery Reviews* 28, 97-119 (1997). The pharmaceutical compositions disclosed herein can be administered in the form of a depot injection or implant preparation which can be formulated in such a manner as to permit a sustained or pulsatile release of the active ingredient.

[0102] Pharmaceutical compositions for oral administration can be in the form of e.g., tablets, pills, powders, lozenges, sachets, cachets, elixirs, suspensions, emulsions, solutions, or syrups. Some examples of suitable excipients include lactose, dextrose, sucrose, sorbitol, mannitol, starches, gum acacia, calcium phosphate, alginates, tragacanth, gelatin, calcium silicate, microcrystalline cellulose, polyvinylpyrrolidone, cellulose, sterile water, syrup, and methylcellulose. Preserving agents such as methyl- and propylhydroxy-benzoates; sweetening agents; and flavoring agents can also be included. Depending on the formulation, compositions can provide quick, sustained or delayed release of the active ingredient after administration to the patient. Polymeric materials can be used for oral sustained release delivery (see "Medical Applications of Controlled Release," Langer and Wise (eds.), CRC Pres., Boca Raton, Fla. (1974); "Controlled Drug Bioavailability," Drug Product Design and Performance, Smolen and Ball (eds.), Wiley, New York (1984); Ranger and Peppas, 1983, *J Macromol. Sci. Rev. Macromol Chem.* 23:61; see also Levy et al., 1985, *Science* 228: 190; During et al., 1989, *Ann. Neurol.* 25:351; Howard et al., 1989, *J. Neurosurg.* 71:105). Sustained release can be achieved by encapsulating conjugates within a capsule, or within slow-dissolving polymers. Preferred polymers include sodium carboxymethylcellulose, hydroxypropylcellulose, hydroxypropylmethylcellulose and hydroxyethylcellulose (most preferred, hydroxypropyl methylcellulose). Other preferred cellulose ethers have been described (Alderman, *Int. J. Pharm. Tech. & Prod. Mfr.*, 1984, 5(3) 1-9). Factors affecting drug release have been described in the art (Bamba et al., *Int. J. Pharm.*, 1979, 2, 307). For administration by inhalation, the compounds for use according to the disclosures herein are conveniently delivered in the form of an aerosol spray preparation from pressurized packs or a nebulizer, with the use of a suitable propellant, e.g., dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, carbon dioxide or other suitable gas, or from propellant-free, dry-powder inhalers. In the case of a pressurized aerosol the dosage unit can be determined by providing a valve to deliver a metered amount. Capsules and cartridges of, e.g., gelatin for use in an inhaler or insufflator can be formulated containing a powder mix of the compound and a suitable powder base such as lactose or starch.

[0103] Effective dosage amounts and regimes (amount and frequency of administration) of the pharmaceutical

compositions are readily determined according to any one of several well-established protocols. For example, animal studies (e.g., mice, rats) are commonly used to determine the maximal tolerable dose of the bioactive agent per kilogram of weight. In general, at least one of the animal species tested is mammalian. The results from the animal studies can be extrapolated to determine doses for use in other species, such as humans for example.

[0104] The components of pharmaceutical compositions are preferably of high purity and are substantially free of potentially harmful contaminants (e.g., at least National Food (NF) grade, generally at least analytical grade, and more typically at least pharmaceutical grade).

[0105] To the extent that a given compound must be synthesized prior to use, the resulting product is typically substantially free of any potentially toxic agents, particularly any endotoxins, which may be present during the synthesis or purification process. Compositions are usually made under GMP conditions. Compositions for parenteral administration are usually sterile and substantially isotonic.

[0106] IX. Methods of Treatments

[0107] Pharmaceutical compositions disclosed herein are used in methods of treatment of prophylaxis of neurological diseases. Examples of such diseases amenable to treatment are cancer (e.g., brain tumors), Acquired Immune Deficiency Syndrome (AIDS), stroke, epilepsy, Parkinson's disease, multiple sclerosis, neurodegenerative disease, trauma, depression, Alzheimer's disease, migraine, pain, seizure disorders, inflammation, and allergic diseases.

[0108] Other pharmaceutical compositions disclosed herein are used in methods of treatment and prophylaxis of non-neurological diseases. Examples of such diseases amenable to treatment are cancer (e.g., tumors of non-CNS tissue), inflammation, and allergic diseases.

[0109] In prophylactic applications, pharmaceutical compositions are administered to a patient susceptible to, or otherwise at risk of, a disease in an amount and frequency sufficient to eliminate or reduce the risk, lessen the severity, or delay the outset of the disease, including biochemical, histologic and/or behavioral symptoms of the disease, its complications and intermediate pathological phenotypes presenting during development of the disease. In therapeutic applications, pharmaceutical compositions are administered to a patient suspected of, or already suffering from such a disease in an amount and frequency sufficient to cure, or at least partially arrest, the symptoms of the disease (biochemical, histologic and/or behavioral), including its complications and intermediate pathological phenotypes in development of the disease. An amount of pharmaceutical composition sufficient to achieve at least one of the above objects is referred to as an effective amount, and a combination of amount and frequency sufficient to achieve at least one of the above objects is referred to as an effective regime.

[0110] X. Methods of Imaging

[0111] As discussed above, the invention provides conjugates comprising a conjugate moiety, which is a substrate of OAT3, linked to an imaging component, as well as agents that are substrates for OAT3 and have an inherent imaging activity. Optionally, the agents also have inherent affinity for a particular antigen or cell type found in the CNS, or the

conjugate is provided with an additional conjugate moiety having such affinity. The additional moiety is referred to as a targeting moiety. The targeting moiety can be an antibody or fragment thereof, or any other molecule that specifically binds to a desired antigen or cell type within the brain. The invention further provides pharmaceutical compositions comprising all of these entities. These pharmaceutical compositions can be used for in vivo imaging. The compositions are administered to a patient and preferentially taken up by central nervous system cells after being actively transported from the blood into the brain by brain microvessel endothelial cells expressing OAT3 in the patient. The compositions also can be administered to a patient and preferentially taken up by central nervous system cells after being actively transported from the cerebrospinal fluid into the brain by choroid plexus epithelial cells expressing OAT3 in the patient. The imaging activity is then detected. In some methods, the imaging component is also a cytotoxic agent. For example many radioisotopes are suitable for both imaging and tumor cytotoxic activity. In such cases, methods of imaging and methods of treatment can be combined. Currently used diagnostic imaging techniques include positron emission tomography (PET), magnetic resonance imaging (MRI), and computed tomography (CT). Actively transported imaging components provide information about, for example, the presence and/or size of a brain tumor. The cell assay methods provided herein can also be used to identify imaging compounds for use outside the CNS, wherein such imaging agents exert undesirable side effect on the CNS. As can be appreciated from the disclosure above, the present invention has a wide variety of applications. For example, the OAT3 transporter can be used to identify an agent or conjugate that is a substrate for the transporter and can cross the blood brain barrier, and can therefore treat the CNS. The OAT3 transporter also can be used to increase the capacity of an agent to cross the blood brain barrier by identifying a conjugate moiety that is a substrate for the OAT3 transporter and linking the conjugate moiety to the agent. Accordingly, the following examples are offered by way of illustration, not by way of limitation.

EXAMPLES

Example 1

Quantitative PCR Detection of OAT3 Expression in Brain Endothelial Cells

[0112] Quantitative PCR was performed to analyze OAT3 expression in human, mouse and rat brain endothelial cells. Microvessel endothelial cells from mouse and rat brains were isolated as follows: To isolate an adequate number of brain endothelial cells, brains were removed from 10 adult rats or 20 adult mice. The brains were washed in 70% ethanol, and placed in sterile phosphate buffered saline. Meninges and surface vessels were removed. Cortical gray

matter was minced, placed in preparation medium (1 g/L glucose, 25 mM HEPES, 100 U/ml penicillin, 100 μ g/ml streptomycin, 10 μ g/ml DNase I, 1 mg/ml collagenase/dispase, in DMEM, adjusted to a pH of 7.4) and incubated for 1 hour at 37° C. Samples were centrifuged for 10 minutes at 1000xg. Fat, cell debris, and myelin were discarded. The pellet was resuspended in fresh preparation medium and incubated for an additional 3 hours at 37° C. in a shaking bath. Medium was filtered through a 230 μ M nylon sieve followed by a 150 μ M nylon sieve. Microvessels were collected by retention on a 60 μ M nylon sieve. Capillaries were washed with preparation medium, and then pelleted for RNA isolation.

[0113] Human brain tissue was obtained from epileptic foci surgically removed from human patients. Human brain microvessel endothelial cells were isolated essentially as described above.

[0114] Total RNA was isolated from the brain endothelial cells using the standard protocol for the RNEasy RNA Isolation Kit (Qiagen). Cells were resuspended in RLT lysis buffer at 10 mls per 0.4 grams of cells. Lysates were vortexed and run through a QiaShredder Column (Qiagen) prior to RNA isolation. Once isolated, the RNA was quantified, run on a 1% agarose gel to ensure integrity, and then stored at -80° C.

[0115] Prior to cDNA synthesis, total RNA was DNase I treated to destroy genomic DNA contamination (Invitrogen DNaseI Kit). Twenty microliters of oligo dT primed single-stranded cDNA was then synthesized from 1 μ g total RNA (Invitrogen ThermoScript cDNA Synthesis Kit). The cDNA was treated with RNase H and stored at -20° C.

[0116] Quantitative PCR was performed in a 96-well format using the MJ Research DNA Engine Opticon. For each transporter, a pair of 26 base oligonucleotide primers were used to amplify the specific transporter. Primers were designed to recognize the non-conserved 3' ends of OAT transporter mRNA. The single stranded cDNA was used as a template for a PCR reaction containing human, mouse or rat primers and SYBR Green master mix (Applied Biosystems). Fluorescent signal was read and graphed each cycle. A CT value, or cycle threshold value, was determined for each reaction. This value was defined as the point at which the fluorescent signal of the reaction exceeds background fluorescence. Background fluorescence was calculated as 20 standard deviations above the average signal from cycles 3 through 10. Transcript abundance was normalized to GAPDH transcript levels. Averaged results from several experiments in which rat OAT family transporters with similar substrate specificity were amplified are shown below in Table 2. The units of measurement are mRNA transcripts detected per PCR reaction. Results from two mouse OAT3 amplification experiments are shown below in Table 3. Average results from 2 human OAT3 amplification experiments are shown below in Table 4.

TABLE 2

OAT family mRNA Expression in Rat Capillary Endothelial Cells			
Gene	transcripts	forward primer	reverse primer
OAT1	758	gccgtctacatgaggaggggagtga	ggcggagtgcaccatgagcaggaagat
		(SEQ ID NO:2)	(SEQ ID NO:3)

TABLE 2-continued

<u>OAT family mRNA Expression in Rat Capillary Endothelial Cells</u>			
Gene	transcripts	forward primer	reverse primer
OAT2	0	tcctgatgccctctgcctgtgctctc (SEQ ID NO:4)	cgccatgagaggggagtccagaatc (SEQ ID NO:5)
OAT3	11,763	agccctgccattcttctgtcaaccct (SEQ ID NO:6)	aaggagagagtttgggcaagtaaga (SEQ ID NO:7)
OAT4	0	tgcaaatcaaaagaataagagaacc (SEQ ID NO:8)	agcttctggcatgattagtagcattc (SEQ ID NO:9)

[0117]

TABLE 3

<u>OAT3 mRNA Expression in Mouse Capillary Endothelial Cells</u>			
Mouse	transcripts	forward primer	reverse primer
Exp. #1	8,639	ccctagccatactcactcctgcactc (SEQ ID NO:10)	actgccaggaatctcaaagggaaat (SEQ ID NO:11)
Exp. #2	14,886	ccctagccatactcactcctgcactc (SEQ ID NO:10)	actgccaggaatctcaaagggaaat (SEQ ID NO:11)
Exp. #3	17,977	ccctagccatactcactcctgcactc (SEQ ID NO:10)	actgccaggaatctcaaagggaaat (SEQ ID NO:11)

[0118]

TABLE 4

<u>OAT3 mRNA Expression in Human Capillary Endothelial Cells</u>			
Human	transcripts	forward primer	reverse primer
Exp.	548	gcatggtgtcccgcctggtgaaaatc (SEQ ID NO:12)	cctctggctcctgctttggcttctt (SEQ ID NO:13)

[0119] The enrichment of OAT3 transcripts in brain capillary endothelial cells (BMECs) relative to total brain transcripts was also determined by quantitative PCR as described above. Total RNA was isolated from whole brain samples. OAT3 transcript levels were normalized to GLUT1 transcript levels. GLUT1 transcript levels were determined

using the human, mouse and rat GLUT1 primers described in Table 7 below. Table 5 below shows the average OAT3 transcript levels, normalized transcript levels, and ratio of OAT3 transcripts in BMEC versus brain in human, mouse and rat brain.

TABLE 5

<u>OAT3 mRNA Expression in Human, Mouse and Rat Brain Microvessel Endothelial Cells</u>									
	Average BMEC			BMEC % GLUT1			BMEC:Brain Ratio		
	Human	Mouse	Rat	Human	Mouse	Rat	Human	Mouse	Rat
OAT3	548	17,977	11,763	0.1	10.6	8.7	0	3.0	1.6
GLUT1	802,859	169,616	135,976	100	100	100	43.1	3.1	9.2

[0120] To confirm the purity of the brain endothelial cell RNA preparations, samples of RNA from each preparation were tested by quantitative PCR for mRNA transcript levels of capillary (GLUT1), neuronal (BNPI) and glial (GFAP) cell markers. The quantitative PCR analysis was conducted as described above. The primers used are described in Table 6 below. The results of the transcript quantitation are described in Table 7 below.

sequenced, and subcloned into plasmids that can be used for expression in mammalian cells or *Xenopus* oocytes. For expression in *Xenopus* oocytes, in vitro OAT3 cRNA was prepared and injected into defolliculated oocytes.

[0123] Oocytes expressing OAT3 exhibited higher levels of ³H-PAH uptake than noninjected controls, as shown in FIG. 2. To measure directly the uptake of possible sub-

TABLE 6

<u>Primers for Quantitative Analysis of Control Genes</u>		
Gene	forward primer	reverse primer
Human GLUT1	ggggcatgattggctccttctctgtg (SEQ ID NO:14)	aggccgcagtacacaccgatgatgaa (SEQ ID NO:15)
Mouse GLUT1	cgcccattcctgtctcttctaccaca (SEQ ID NO:16)	tcattggtgtttgtgtggccctcagtg (SEQ ID NO:17)
Rat GLUT1	gaaccacagggaaagcaactctaate (SEQ ID NO:18)	tcgggtcattatctttcacgtttcca (SEQ ID NO:19)
Human BNPI	cacccccgccttctcttctctccag (SEQ ID NO:20)	ctgctggtagggagatgtgaagtgg (SEQ ID NO:21)
Mouse BNPI	acgggggacatcactcagaattacat (SEQ ID NO:22)	ttcttctcttttctcccagccgtagg (SEQ ID NO:23)
Rat BNPI	gccacacacagcacagttcagcctcc (SEQ ID NO:24)	ggacagcactgggcacaaggggaagac (SEQ ID NO:25)
Mouse GFAP	aggaaattgctggaggggcgaagaaaa (SEQ ID NO:26)	caccatcccgcattctccacagtcttt (SEQ ID NO:27)
Rat GFAP	ggtgggcaggtgaggaagaaatggag (SEQ ID NO:28)	tagcagaggtgacaaggggggagtg (SEQ ID NO:29)

[0121]

TABLE 7

<u>Control Gene mRNA Transcript Levels</u>				
Control Gene	Source	Control Gene mRNA Transcript Abundance		
		Human	Rat	Mouse
GLUT1 (Capillary marker)	Capillaries	802859	169616	135976
	Whole	11120	13546	5278
	Brain			
BNPI (Neuronal marker)	Capillaries	2614	5	343
	Whole	222285	67705	122509
	Brain			
GFAP (Glial marker)	Capillaries	Not determined	561	670
	Whole	Not determined	68789	24032
	Brain	determined		

Example 2

Studies of Cloned OAT3 Transporters: Oocyte Expression

[0122] To assess transport function of a specific transporter protein, it can be desirable to clone the cDNA and express the protein in cells that have low endogenous transport activity. Human OAT3 was cloned by PCR, fully

strates, an oocyte uptake assay can be performed in which uptake of compounds is measured by mass spectroscopy. For example, uptake of PAH can be measured. Oocytes injected with OAT3 cRNA can be incubated at 16-18° C. until maximal transporter expression is reached. Oocytes from the same batch, not injected with cRNA, can be used as a control. A 0.5 mM solution of PAH can be prepared in oocyte ringers (ND96) buffer (90 mM NaCl, 10 mM HemiNa HEPES, 2 mM KCl, 1 mM MgCl₂, 1.8 mM CaCl₂, pH adjusted to 7.4) containing 0.5% bovine serum albumin. The PAH is, for example, administered to pools of 8 oocytes for a 20 minute duration. Following the incubation, the pools of oocytes are washed with 0.5% BSA ND96 buffer and separated into subpools containing, for example, 4 oocytes each. Subpools are homogenized in 150 μ l of ice cold 80% MeOH/H₂O and lysed manually with a P200 pipettor. Lysates can be vortexed before being centrifuged at, for example, 13.2 krpm for 15 minutes. Approximately 110 μ l of lysate is removed from the tubes and placed in a 96-well plate and analyzed for PAH concentration by LC-MS-MS.

[0124] Samples are analyzed by LC-MS-MS as follows. A specific method can be developed for each test compound, and calibrated against a series of dilutions of known compound concentrations spiked into cellular extract. Measurements are performed using, for example, an API 2000 LC-MS-MS spectrometer equipped with Agilent 1100 binary transporters and a CTC HTS-PAL autosampler. Analyte fragmentation peaks are integrated, for example, using

Analyst 1.2 quantitation software, and concentrations are calculated using a calibration curve of signals produced by known concentrations of the compound.

Example 3

Studies of Cloned OAT3 Transporters: Uptake into Mammalian Cells

[0125] To increase the throughput of OAT3 compound screening, a mammalian cell assay was developed using an inducible expression system. OAT3 was subcloned into a plasmid that allows for inducible expression by tetracycline (TREx plasmid, Invitrogen Inc., Carlsbad, Calif.). The OAT3-HEK plasmid was transiently transfected into a human embryonic kidney (HEK) cell line engineered for tetracycline-inducible expression (HEK-TREx, Invitrogen) and stable clones were isolated by G418 selection and flow activated cell sorting (FACS).

[0126] To determine whether a compound interacts with the OAT3 transporter, a competition binding assay was developed. This assay measures how different concentrations of a test compound block the uptake of a radiolabeled substrate such as ^3H -Ochratoxin A. The half-maximal inhibitory concentration (IC_{50}) for inhibition of transport of a substrate by a test compound is an indication of the affinity of the test compound for the OAT3 transporter. Competition binding studies were performed as follows. OAT3-HEK/TREx cells were plated in 96-well plates at 100,000 cells/well and incubated at 37° C. for 24 hours and tetracycline (1 $\mu\text{g}/\text{mL}$) was added to each well for an additional 24 hours to induce OAT3 expression. Radiolabeled ^3H -Ochratoxin A (~50,000 cpm/well) was added to each well in the presence and absence of various concentrations of PAH in duplicate or triplicate. Plates were incubated at room temperature for 20 minutes. Excess ^3H -Ochratoxin A was removed and cells were washed three times with a 96-well plate washer with cold assay buffer. Scintillation fluid was added to each well, and the plates were sealed and counted in a 96-well plate-based scintillation counter. Data was graphed and analyzed using non-linear regression analysis with Prism Software (GraphPad, Inc., San Diego, Calif.). An example of results obtained using this assay is shown in **FIG. 3**.

[0127] Competition binding studies only demonstrate that a molecule interacts with the OAT3 protein, but do not demonstrate whether the molecule is a substrate and is translocated across the plasma membrane or is non-transported inhibitor or a non-transported ligand. In order to measure whether test compounds are actively translocated across the membrane, and to determine the maximal transport rate, a direct uptake method can be used in which transport of a test compound is measured by mass spectrometry. For direct uptake measurements using mass spectrometry, OAT3-HEK/TREx cells can be prepared similarly to those used for competition studies (described above). OAT3-expressing cells are washed and incubated with test compounds such as PAH. Excess substrate is removed by washing with cold assay buffer. Cells are lysed with 50% ethanol/water and the cell debris is pelleted by centrifugation. The supernatant is analyzed by LC-MS-MS. As a negative control, uptake is measured in cells not treated with tetracycline, or by competition with another compound such as PAH or ochratoxin.

Example 4

Efflux Assays

[0128] **FIG. 4** depicts the results of an efflux experiment in which the PgP substrate verapamil was added to commercial Baculovirus membranes (purchased from BD Biosciences) at various concentrations depicted on the X axis followed by ATPase activity measurement. The ATPase activity measurement was performed using the lactate dehydrogenase/pyruvate kinase coupled enzyme system described by Tietz & Ochoa, Arch. Biochim. Biophys. Acta 78:477 (1958) to follow the decrease in absorbance at 340 nm resulting from the oxidation of NADH, which is proportional to ATPase activity. 5 mM sodium azide (NaN_3), 1 mM EGTA, and 0.5 mM Ouabain, each of which inhibit non-specific ATPases in the membranes, were added to the reactions to further enhance the specificity of the PgP ATPase signal. The other components in the assay mixture were 25 mM Tris, pH 7.8, 100 mM NaCl, 10 mM KCl, 5 mM MgCl_2 , 1 mM DTT, 2 mM phosphoenolpyruvate, 1 mM NADH, 0.1 mg/ml lactate dehydrogenase, 0.1 mg/ml pyruvate kinase, 5 mM ATP, and 6 μg PgP or control membranes. **FIG. 4** demonstrates that as the concentration of verapamil was increased, the ATPase activity in PgP-containing membranes but not in control membranes also increased.

[0129] **FIG. 5** depicts the results of an efflux competition assay. A tetracycline-inducible PgP expression construct (TREx-PgP) was transfected into HEK cells. The cells were incubated with PgP substrate 5 μM calcein-AM, which passively diffuses into the cells, as well as with various concentrations of the PgP substrate verapamil as shown in **FIG. 5**. As the concentration of PgP substrate verapamil was increased, more calcein-AM accumulated in the cells and was converted to the fluorescent product calcein.

[0130] **FIG. 6** depicts the results of a cellular transwell monolayer efflux assay. MDCK cells transfected with the tetracycline-inducible TREx-PgP expression vector were seeded on polycarbonate filter membranes in transwell dishes and grown for 3-5 days, yielding a polarized monolayer with tight junctions between cells. In this example, apical to basolateral and basolateral to apical transport of 2.5 nM (approximately 100,000 cpm) radiolabeled PgP substrate ^3H -vinblastine was measured in the absence and presence of 250 μM of the inhibitor/competitor verapamil. The left set of bars depicts apical to basolateral transport, while the right set of bars depicts basolateral to apical transport. Apical to basolateral transport of ^3H -vinblastine was strongly increased and basolateral to apical transport of ^3H -vinblastine was strongly decreased in the presence of verapamil, indicating that ^3H -vinblastine is a substrate of PgP.

Example 5

Localization of OAT3 in the Human Brain

[0131] A commercial rabbit anti-OAT3 antibody (4ADI) was used to stain sections of human brain using tissue slides from Lifespan Technologies. Paraffin-embedded human brain sections were deblocked, heat treated for antigen retrieval, and stained with the OAT3 antibody and an alkaline-phosphatase conjugated goat anti-rabbit antibody (Jackson Labs). Staining was detected by the DAB colorimetric

method. The brain sections showed no vessel staining in the cerebral cortex and cerebellum, but weak, diffuse staining in the choroid plexus epithelium.

[0132] Although the foregoing compounds, conjugates and methods have been described in detail for purposes of clarity of understanding, it will be obvious that certain

modifications may be practiced within the scope of the claim(s) granted herefrom. Unless otherwise apparent from the context, any element, embodiment, or step can be used in combination with any other. All publications and patent documents cited herein are hereby incorporated by reference in their entirety for all purposes to the same extent as if each were so individually denoted.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 29

<210> SEQ ID NO 1

<211> LENGTH: 1707

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 1

```

atgaccttct cggagatcct ggaccgtgtg ggaagcatgg gccatttcca gttcctgcat    60
gtagccatac tgggcctccc gatcctcaac atggccaacc acaacctgct gcagatcttc    120
acagccgcca cccctgtcca ccaactgtcg cgcgccaca atgcctccac agggccttgg    180
gtgtccccc tgggcccata tgggaagcct gagaggtgcc tccgttttgt acatccgccc    240
aatgccagcc tgcccataga caccagagg gccatggagc catgcctgga tggctgggtc    300
tacaacagca ccaaggactc cattgtgaca gagtgggact tgggtgtgca ctccaacaaa    360
ctgaaggaga tggcccagtc tatcttcctg gcaggtatac tgattggagg gctcgtgctt    420
ggagacctgt ctgacaggtt tggccgcacc atcctgacct gcagctacct gctgctggca    480
gccagcggtt ccggtgcagc cttcagcccc accttcccca tctacatggt ctccgcttc    540
ctgtgtggct ttggcatctc aggcattacc ctgagcaccg tcatctttaa tggggaatgg    600
gtgcctaccc ggatgcgggc catcatgtcg acagcactcg ggtactgcta cacctttggc    660
cagttcattc tgcccggcct ggctacgcc atccccagc ggcgttggct gcagttaact    720
gtgtccattc cttctctcgt ctcttccta tcatcctggt ggacaccaga gtccatacgc    780
tgggtgtctt gtctggaagt cctcgaagc cctgaagata ctccggcggg tggctgtctt    840
caatggcaag aagagggaga aaggctcagc ttggaggagc tcaaaactca cctgcagaag    900
gagatctcct tggccaaggc caagtacacc gcaagtgacc tgttccggat acccatcggt    960
gcgcccagtc acctctgct tccctggcc tggtttgcta ccggttttgc ctactatagt   1020
ttggctatgg gtgtggaaga atttgagtc aacctctaca tctccagat catctttggt   1080
ggggtcagat tcccagccaa gttcatcacc atcctctcct taagctacct gggccggcat   1140
accactcagg ggcgctgccc tgcctctggc agaggggcca tcttgctct cacccttgtg   1200
cccttgagct tgcagaccgt ggagacagta ttggctgtgt ttgggaaggg atgcctatcc   1260
agctccttca gctgcctctt cctctacaca agtgaattat accccacagt catcaggcaa   1320
acaggtatgg gcgtaagtaa cctgtggacc cgcgtgggaa gcatggtgtc ccgctgggtg   1380
aaaatcacgg gtgaggtaca gccttctac cccaatatca tctttacggg atctaccgcc   1440
ctcctcgggg gcagtgtctc cctcttctg cctgagacc tgaacagccc ttgccagaga   1500
cgatcgaaga cctggaaaac tggctagtc ctgcctctgg ccccatcagt gctcctcct   1560
ggggaagcag gctctggccc agggcttttc cttagctctc tgtccctagg totgcccga   1620

```

-continued

aagaagccaa agcaggagcc agaggtggaa aaggcctccc agaggatccc tctacagcct 1680

cacggaccag gcctgggctc cagctga 1707

<210> SEQ ID NO 2
<211> LENGTH: 26
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: OAT1 forward primer

<400> SEQUENCE: 2

gccgtctaca tgaggagggg gagtga 26

<210> SEQ ID NO 3
<211> LENGTH: 26
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: OAT1 reverse primer

<400> SEQUENCE: 3

ggcggagtga ccatgagcag gaagat 26

<210> SEQ ID NO 4
<211> LENGTH: 26
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: OAT2 forward primer

<400> SEQUENCE: 4

tcctgatgcc ctctgctgt gctctc 26

<210> SEQ ID NO 5
<211> LENGTH: 26
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: OAT2 reverse primer

<400> SEQUENCE: 5

cggccatgag aggggagtcc agaatc 26

<210> SEQ ID NO 6
<211> LENGTH: 26
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: OAT3 forward primer

<400> SEQUENCE: 6

agccctgcca ttcttctgtc aaccct 26

<210> SEQ ID NO 7
<211> LENGTH: 26
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: OAT3 reverse primer

<400> SEQUENCE: 7

aaggagagag tttgggcaag gtaaga 26

-continued

<210> SEQ ID NO 8
<211> LENGTH: 26
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: OAT4 forward primer

<400> SEQUENCE: 8

tgcaaatcaa aagaaataag agaacc 26

<210> SEQ ID NO 9
<211> LENGTH: 26
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: OAT4 reverse primer

<400> SEQUENCE: 9

agcttctggc atgattagta cgattc 26

<210> SEQ ID NO 10
<211> LENGTH: 26
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: Mouse OAT3 forward primer

<400> SEQUENCE: 10

ccctagccat actcactcct gcactc 26

<210> SEQ ID NO 11
<211> LENGTH: 26
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: Mouse OAT3 reverse primer

<400> SEQUENCE: 11

actgccaggg aatctcaaag ggaatc 26

<210> SEQ ID NO 12
<211> LENGTH: 26
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: Human OAT3 forward primer

<400> SEQUENCE: 12

gcatggtgtc cccgctggtg aaaatc 26

<210> SEQ ID NO 13
<211> LENGTH: 26
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: Human OAT3 reverse primer

<400> SEQUENCE: 13

cctctggctc ctgcttggc ttcttt 26

<210> SEQ ID NO 14
<211> LENGTH: 26
<212> TYPE: DNA

-continued

<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: Human GLUT1 forward primer

<400> SEQUENCE: 14
ggggcatgat tggctccttc tctgtg 26

<210> SEQ ID NO 15
<211> LENGTH: 26
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: Human GLUT1 reverse primer

<400> SEQUENCE: 15
aggccgcagt acacaccgat gatgaa 26

<210> SEQ ID NO 16
<211> LENGTH: 26
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: Mouse GLUT1 forward primer

<400> SEQUENCE: 16
cgccccattcc tgtctcttcc taccca 26

<210> SEQ ID NO 17
<211> LENGTH: 26
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: Mouse GLUT1 reverse primer

<400> SEQUENCE: 17
tcatgtgtgtt tgtgtggccc tcaagt 26

<210> SEQ ID NO 18
<211> LENGTH: 26
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: Rat GLUT1 forward primer

<400> SEQUENCE: 18
gaaccacag gaaagcaact ctaatc 26

<210> SEQ ID NO 19
<211> LENGTH: 26
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: Rat GLUT1 reverse primer

<400> SEQUENCE: 19
tcgggtcatt attttccag tttcca 26

<210> SEQ ID NO 20
<211> LENGTH: 26
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: Human BNP1 forward primer

-continued

<400> SEQUENCE: 20

cacccccgc tttcctttat ctccag 26

<210> SEQ ID NO 21

<211> LENGTH: 26

<212> TYPE: DNA

<213> ORGANISM: Artificial

<220> FEATURE:

<223> OTHER INFORMATION: Human BNP1 reverse primer

<400> SEQUENCE: 21

ctgctggtag gggagatgtg aagtgg 26

<210> SEQ ID NO 22

<211> LENGTH: 26

<212> TYPE: DNA

<213> ORGANISM: Artificial

<220> FEATURE:

<223> OTHER INFORMATION: Mouse BNP1 forward primer

<400> SEQUENCE: 22

acgggggaca tcactcagaa ttacat 26

<210> SEQ ID NO 23

<211> LENGTH: 26

<212> TYPE: DNA

<213> ORGANISM: Artificial

<220> FEATURE:

<223> OTHER INFORMATION: Mouse BNP1 reverse primer

<400> SEQUENCE: 23

ttcttccttt ttctcccagc cgttag 26

<210> SEQ ID NO 24

<211> LENGTH: 26

<212> TYPE: DNA

<213> ORGANISM: Artificial

<220> FEATURE:

<223> OTHER INFORMATION: Rat BNP1 forward primer

<400> SEQUENCE: 24

gccacacaca gcacagttca gcctcc 26

<210> SEQ ID NO 25

<211> LENGTH: 26

<212> TYPE: DNA

<213> ORGANISM: Artificial

<220> FEATURE:

<223> OTHER INFORMATION: Rat BNP1 reverse primer

<400> SEQUENCE: 25

ggacagcact gggcacaagg gaagac 26

<210> SEQ ID NO 26

<211> LENGTH: 26

<212> TYPE: DNA

<213> ORGANISM: Artificial

<220> FEATURE:

<223> OTHER INFORMATION: Mouse GFAP forward primer

<400> SEQUENCE: 26

aggaaattgc tggagggcga agaaaa 26

-continued

```

<210> SEQ ID NO 27
<211> LENGTH: 26
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: Mouse GFAP reverse primer

<400> SEQUENCE: 27

caccatcccg catctccaca gtcttt                26

<210> SEQ ID NO 28
<211> LENGTH: 26
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: Rat GFAP forward primer

<400> SEQUENCE: 28

ggtgggcagg tgaggaagaa atggag                26

<210> SEQ ID NO 29
<211> LENGTH: 26
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: Rat GFAP reverse primer

<400> SEQUENCE: 29

tagcagaggt gacaaggggg ggagtg                26

```

1. A method of screening an agent, conjugate or conjugate moiety for activity useful for treating or diagnosing a disease, comprising:

- (a) providing a cell expressing an OAT3 transporter, the transporter being situated in the plasma membrane of the cell;
- (b) contacting the cell with an agent, conjugate or conjugate moiety; and
- (c) determining whether the agent, conjugate or conjugate moiety passes through the plasma membrane via the OAT3 transporter, passage through the OAT3 transporter being useful for treatment or diagnosis of the disease;

wherein:

if step (b) comprises contacting the cell with the agent, the agent is a neuropharmaceutical agent or an imaging component;

if step (b) comprises contacting the cell with the conjugate, the conjugate comprises an agent that is a neuropharmaceutical agent or an imaging component; or

if step (b) comprises contacting the cells with the conjugate moiety, the method further comprises linking the conjugate moiety to an agent that is a neuropharmaceutical agent or an imaging component.

2. The method of claim 1, wherein:

(i) the cell endogenously expresses the OAT3 transporter; or

(ii) a nucleic acid molecule encoding the OAT3 transporter has been transfected or injected into the cell.

3. The method of claim 2, wherein the cell is a brain microvessel endothelial cell or a choroid plexus epithelial cell.

4. The method of claim 2, wherein the cell is an oocyte.

5. The method of claim 2, wherein the cell is a human embryonic kidney (HEK) cell.

6. The method of claim 2, wherein the cell is transformed with an SV40 large T antigen that can be expressed in a temperature sensitive fashion.

7. The method of claim 2, wherein the determining is performed by a competition assay.

8. The method of claim 2, wherein the determining is performed by a direct uptake assay.

9. The method of claim 3, wherein the cell is a brain microvessel endothelial cell, and the brain microvessel endothelial cell is one of a plurality of brain microvessel endothelial cells forming a polarized monolayer, the agent, conjugate or conjugate moiety is contacted to one side of the polarized monolayer, and the determining comprises determining whether the agent, conjugate or conjugate moiety is transported into the brain microvessel endothelial cells or to the opposite side of the polarized monolayer.

10. The method of claim 3, wherein the cell is a choroid plexus epithelial cell, and the choroid plexus epithelial cell is one of a plurality of choroid plexus epithelial cells

forming a polarized monolayer, the agent, conjugate or conjugate moiety is contacted to one side of the polarized monolayer, and the determining comprises determining whether the agent, conjugate or conjugate moiety is transported into the choroid plexus epithelial cells or to the opposite side of the polarized monolayer.

11. The method of claim 1, further comprising administering the agent, conjugate, or conjugate moiety to a peripheral tissue of an animal and measuring the amount of agent, conjugate, or conjugate moiety that passes through the blood brain barrier into the brain of the animal.

12. The method of claim 1, wherein the determining step determines that the agent, conjugate or conjugate moiety passes through the plasma membrane via the OAT3 transporter; and the method further comprises:

(d) modifying the agent, conjugate or conjugate moiety; and

(e) determining if the modified agent, conjugate or conjugate moiety is transported with a higher V_{max} by the OAT3 transporter than the agent, conjugate or conjugate moiety.

13. The method of claim 1, wherein the neuropharmaceutical agent is a cytotoxic neuropharmaceutical agent selected from the group consisting of platinum, nitrosourea, a phosphoramidate group that is selectively cytotoxic to brain tumor cells, nitroimidazole, and nitrogen mustard.

14. The method of claim 1, wherein the agent, conjugate or conjugate moiety comprises a xenobiotic molecule containing one or more ring structures and at least one anionic groups selected from the list comprising carboxylates, sulphates, and phosphates.

15. The method of claim 14, wherein the xenobiotic molecule is selected from the list comprising para-amino-hippurate, ochratoxin A, estrone-3-sulphate, benzylpenicillin, and probenecid.

16. The method of claim 1, further comprising administering the agent, conjugate or conjugate moiety to an undiseased animal and determining any toxic effects.

17. The method of claim 1, further comprising determining that the agent, conjugate or conjugate moiety is transported by at least one efflux transporter.

18. The method of claim 17, further comprising:

(d) modifying the agent, conjugate or conjugate moiety;

(e) establishing that the modified agent, conjugate or conjugate moiety retains OAT3 substrate activity; and

(f) comparing the ratio of OAT3 substrate activity to the ratio of efflux substrate activity for the agent, conjugate or conjugate moiety and the modified agent, conjugate or conjugate moiety wherein an increased ratio of OAT3 substrate activity to efflux substrate activity demonstrates that the modification improves the usefulness of the agent, conjugate or conjugate moiety for treatment or diagnosis of the disease.

19. The method of claim 18, wherein the efflux substrate activity is determined by conducting an assay selected from the group consisting of:

(a) an efflux transporter ATPase activity assay;

(b) an efflux transporter competition assay; and

(c) a direct efflux transport assay across a polarized monolayer of cells.

20. A conjugate comprising a neuropharmaceutical agent or imaging component which is actively transported across the blood brain barrier, identified by the method of claim 11.

21. Use of an OAT3 transporter to identify an agent or conjugate that is a substrate for the transporter and can cross the blood brain barrier and can therefore treat a CNS disease.

22. A pharmaceutical composition comprising a neuropharmaceutical agent or an imaging component linked to a conjugate moiety to form a conjugate, wherein the conjugate has a higher V_{max} for OAT3 than the neuropharmaceutical agent or the imaging component alone.

23. The pharmaceutical composition of claim 22, wherein the conjugate has at least 5 times the V_{max} for OAT3 than the neuropharmaceutical agent or the imaging component alone.

24. The pharmaceutical composition of claim 22, wherein the conjugate has a V_{max} for OAT3 that is at least 5% of the V_{max} for OAT3 of para-amino-hippurate.

25. The pharmaceutical composition of claim 22, wherein the conjugate has a lower V_{max} for an efflux transporter than the neuropharmaceutical agent or the imaging component alone.

26. A method of screening an agent, conjugate or conjugate moiety for capacity to be transported into the brain, comprising:

(a) determining whether the agent, conjugate or conjugate moiety specifically binds to an OAT3 transporter; and

(b) contacting the agent to one side of a polarized monolayer of brain microvessel endothelial or choroid plexus epithelial cells; and

(c) determining whether the agent is transported across the polarized monolayer.

27. The method of claim 26, wherein step (a) is performed by contacting a cell expressing the OAT3 transporter, the transporter being situated in the plasma membrane of the cell, with a substrate of the OAT3 transporter, and determining whether the agent inhibits transport of the substrate across the polarized monolayer.

28. The method of claim 26, further comprising administering the agent to an undiseased animal and determining any toxic effects.

29. A method of formulating a conjugate, comprising:

(a) linking a neuropharmaceutical agent or imaging component to a conjugate moiety to form the conjugate, wherein the conjugate has a greater V_{max} for an OAT3 transporter than the neuropharmaceutical agent or imaging component alone; and

(b) formulating the conjugate with a pharmaceutical carrier as a pharmaceutical composition.

30. A method of delivering a conjugate, comprising administering to a patient a pharmaceutical composition comprising a neuropharmaceutical agent or imaging component linked to a conjugate moiety to form the conjugate, wherein the conjugate has a higher V_{max} for an OAT3 transporter than the neuropharmaceutical agent or imaging component alone.

31. The method of claim 30, wherein the V_{max} of the conjugate is at least two-fold higher than that of the neuropharmaceutical agent or imaging component alone.

32. The method of claim 30, wherein the neuropharmaceutical agent is a cytotoxic neuropharmaceutical selected from the group consisting of platinum, nitrosourea, a phos-

phoramide group selectively cytotoxic to brain tumor cells, nitroimidazole, and nitrogen mustard.

33. A method of screening an agent or imaging component for decreased side effects in the central nervous system (CNS), comprising:

- (a) providing
 - (i) an agent having a pharmacological activity, wherein the pharmacological activity is useful for treating a disease present in a tissue other than the CNS, and the agent causes undesired side effects in the CNS if the agent enters the CNS; or
 - (ii) an imaging component useful for imaging a tissue other than the CNS, and the imaging component causes undesired side effects in the CNS if the imaging component enters the CNS;
- (b) modifying the agent or imaging component;
- (c) providing a cell expressing at least one OAT3 transporter protein that transports substrates across the blood brain barrier,

- (d) contacting the cell with the modified agent or modified imaging component; and

- (e) determining whether the modified agent or modified imaging component passes through the plasma membrane via the transporter protein with a lower V_{\max} than the agent, a lower V_{\max} indicating that the modification decreases the capacity of the modified agent or modified imaging component relative to the agent or imaging component to cross the blood brain barrier, thereby decreasing undesired side effects in the CNS.

34. The method of claim 33, wherein the cell is transformed or injected with a nucleic acid encoding a transporter or the cell is a brain microvessel endothelial cell or a choroid plexus epithelial cell.

35. The method of claim 33, wherein the modifying comprises linking the agent or imaging component to a conjugate moiety to form a conjugate.

* * * * *

专利名称(译)	OAT3转运蛋白在血脑屏障细胞中表达		
公开(公告)号	US20050170392A1	公开(公告)日	2005-08-04
申请号	US11/027581	申请日	2004-12-30
[标]申请(专利权)人(译)	什诺波特有限公司		
申请(专利权)人(译)	XENOPORT INC.		
当前申请(专利权)人(译)	XENOPORT INC.		
[标]发明人	ZERANGUE NOA		
发明人	ZERANGUE, NOA		
IPC分类号	A61K31/19 A61K31/225 A61K31/60 A61K47/48 C12Q1/68 G01N33/53 G01N33/567 G01N33/68		
CPC分类号	G01N33/6872 G01N2500/10 G01N33/6896		
优先权	60/540772 2004-01-30 US		
外部链接	Espacenet USPTO		

摘要(译)

OAT3在脑微血管内皮细胞中始终以高水平表达。本文公开了用于确定测试材料/分子是否是OAT3转运蛋白的底物和/或由OAT3转运蛋白活性转运的测定法，并因此确定用于穿过血脑屏障的候选底物。该测定法可用于筛选用于治疗或诊断神经疾病的治疗性，细胞毒性或成像化合物。

