



US007883858B2

(12) **United States Patent**
Hood et al.

(10) **Patent No.:** **US 7,883,858 B2**
(45) **Date of Patent:** **Feb. 8, 2011**

(54) **METHODS FOR IDENTIFYING AND MONITORING DRUG SIDE EFFECTS**

(75) Inventors: **Leroy Hood**, Seattle, WA (US);
Biaoyang Lin, Bothell, WA (US)

(73) Assignee: **Institute for Systems Biology**, Seattle, WA (US)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

(21) Appl. No.: **12/468,834**

(22) Filed: **May 19, 2009**

(65) **Prior Publication Data**

US 2010/0075866 A1 Mar. 25, 2010

Related U.S. Application Data

(63) Continuation of application No. 11/342,367, filed on Jan. 27, 2006, now abandoned.

(60) Provisional application No. 60/683,004, filed on May 20, 2005, provisional application No. 60/647,792, filed on Jan. 27, 2005.

(51) **Int. Cl.**
G01N 33/53 (2006.01)

(52) **U.S. Cl.** **435/7.1; 435/7.2; 436/518**

(58) **Field of Classification Search** None
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

4,764,456 A * 8/1988 Watanabe et al. 430/550
5,824,467 A * 10/1998 Mascarenhas 435/4
6,210,971 B1 * 4/2001 Messenger et al. 436/98
2004/0137440 A1 * 7/2004 Lin 435/6

OTHER PUBLICATIONS

Merrick et al. (Environ Health Perspectives 2003 vol. 111, p. A578-579).*

* cited by examiner

Primary Examiner—Jacob Cheu

(74) *Attorney, Agent, or Firm*—Arnall Golden Gregory LLP

(57) **ABSTRACT**

The present invention relates generally to methods for identifying drug side effects by detecting perturbations in organ-specific molecular blood fingerprints. The invention further relates to methods for identifying drug-specific organ-specific molecular blood fingerprints. As such, the present invention provides compositions comprising organ-specific proteins, detection reagents for detecting such proteins, and panels and arrays for determining organ-specific molecular blood fingerprints.

22 Claims, No Drawings

METHODS FOR IDENTIFYING AND MONITORING DRUG SIDE EFFECTS

STATEMENT OF GOVERNMENT INTEREST

This invention was made with government support under Grant Nos. P50 CA097186 and P01 CA085857 awarded by the National Cancer Institute. The government may have certain rights in this invention.

STATEMENT REGARDING SEQUENCE LISTING

The Sequence Listing associated with this application submitted Nov. 13, 2009 as a text file named "25409_6_8404_2009_11_13_AMD_AFD_Sequence_Listing.txt," created on Oct. 19, 2009, and having a size of 3,396 kilobytes is hereby incorporated by reference pursuant to 37 C.F.R. §1.52 (e)(5).

BACKGROUND OF THE INVENTION

1. Field of the Invention

The present invention relates to organ-specific molecular blood fingerprints and methods for using the same in identifying and/or monitoring drug side effects.

2. Description of the Related Art

Side effects, particularly adverse side effects, are monitored and tested throughout the developmental path of all drugs. A variety of in vitro and in vivo toxicity assays are available in the art for testing on and off-target effects of drugs during development including metabolic effects on liver P450 enzymes. Clearly, however, as is evidenced by the recent controversy surrounding COX-2 inhibitors, off-target side effects can be subtle and difficult to detect. Therefore, monitoring and identifying drug side effects remains an important issue with regard to drug safety of developing drugs and approved drugs already on the market. The COX-2 story highlights the need in the art for improved methods to detect sometimes subtle adverse off-target effects of drugs.

The present invention provides methods that satisfy this and other needs.

BRIEF SUMMARY OF THE INVENTION

One aspect of the present invention provides a method for detecting a drug side effect comprising measuring in the blood of a subject taking the drug the level of a plurality of organ-specific proteins secreted from an organ wherein the levels of the plurality of organ-specific proteins together provide an organ-specific molecular blood fingerprint that indicates a drug side effect on the organ in the subject. In one embodiment, the level of the plurality of organ-specific proteins is measured with any of a variety of methods, including but not limited to mass spectrometry, such as tandem mass spectrometry, an immunoassay, such as an ELISA, Western blot, microfluidics/nanotechnology sensors, and aptamer capture assay. In this regard, an aptamer may be used in a similar manner to an antibody in a variety of appropriate binding assays known to the skilled artisan and described herein. In a further embodiment, the plurality of organ-specific proteins is measured using tandem mass spectrometry. In one embodiment, the level of one or more organ-specific proteins is measured. In yet an additional embodiment, the plurality of organ-specific proteins comprises from at least about 2 organ-specific proteins to about 100, 150, 160, 170, 180, 190, 200 or more organ-specific proteins. In this regard,

the plurality of organ-specific proteins may comprise at least 2, 3, 4, 5, 6, 7, 8, 9, 10, or more organ-specific proteins. In one embodiment, the plurality of organ-specific proteins comprises about 5, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 or 20 organ-specific proteins. In another embodiment, the organ-specific proteins comprise proteins from any of a number of organs, such as, but not limited to the liver or kidney. In a further embodiment the organ-specific proteins comprise cardiac-specific proteins. In yet another embodiment, the organ-specific proteins are from an organ other than the expected therapeutic target of the drug.

Another aspect of the present invention provides a method for detecting a drug side effect comprising measuring in the blood of a subject taking the drug the level of one or more organ-specific proteins secreted from an organ wherein the level of the one or more organ-specific proteins together provide an organ-specific molecular blood fingerprint that indicates a drug side effect on the organ in the subject.

Another aspect of the invention provides a method for determining the presence or absence of a drug side effect in a subject taking the drug comprising, detecting a level of each of a plurality of organ-specific proteins in a blood sample from the subject, wherein the plurality of organ-specific proteins are secreted from the same organ; comparing said level of each of the plurality of organ-specific proteins in the blood sample from the subject to a level of each of the plurality of organ-specific proteins in a control sample of drug-free blood; wherein a statistically significant altered level of one or more of the plurality of organ-specific proteins in the blood is indicative of the presence or absence of a drug side effect.

As would be readily appreciated by the skilled artisan, an altered level can mean an increase in the level or a decrease in the level. In this regard, the skilled artisan would readily appreciate that a variety of statistical tests can be used to determine if an altered level is significant. The Z-test (Man, M. Z., et al., *Bioinformatics*, 16: 953-959, 2000) or other appropriate statistical tests can be used to calculate P values for comparison of protein expression levels. In certain embodiments, the level of each of the plurality of organ-specific proteins in the blood sample from the subject is compared to a previously determined normal control level of each of the plurality of organ-specific proteins taking into account standard deviation (see e.g., U.S. Patent Application No. 20020095259). In one embodiment, the level of each of the plurality of organ-specific proteins is detected using any one or more methods, such as, but not limited to mass spectrometry (e.g., tandem mass spectrometry or other spectrometry-based techniques), and immunoassays (e.g., ELISA, Western blot, or other immunoaffinity-based assays). In an additional embodiment, the method level of each of the plurality of organ-specific proteins is measured using an antibody array. In yet an additional embodiment, the method provides for determining the presence or absence of a drug side effect wherein the organ-specific proteins comprise liver-specific proteins or kidney-specific proteins. In certain embodiments, the organ-specific proteins are from an organ other than the expected therapeutic target of the drug.

A further aspect of the present invention provides a method for detecting perturbation of a normal biological state induced by a drug, contacting a blood sample from a subject taking the drug with a plurality of detection reagents each specific for an organ-specific protein secreted into blood, wherein each organ-specific protein is secreted from the same organ; measuring the amount of the organ-specific protein detected in the blood sample by each detection reagent, comparing the amount of the organ-specific protein detected in the blood sample by each detection reagent to a predetermined

control amount for each organ-specific protein; wherein an altered level in one or more of the organ-specific proteins indicates a perturbation in the normal biological state induced by the drug. In this regard, the plurality of detection reagents may comprise from at least about 2 detection reagents to about 100, 150, 160, 170, 180, 190, 200 or more detection reagents. In one embodiment, the plurality of detection reagents comprises about 5, 10, or 20 detection reagents. In yet another embodiment, the organ-specific proteins comprise kidney-specific proteins, liver-specific proteins, or cardiac-specific proteins. As would be recognized by the skilled artisan upon reading the present disclosure, the organ-specific proteins can be derived from any organ in the body as described further herein.

In yet a further aspect, the invention provides a diagnostic panel for determining a drug side effect in a subject taking the drug comprising, a plurality of detection reagents each specific for detecting one of a plurality of organ-specific proteins present in a blood/serum/plasma sample; wherein the organ-specific proteins are secreted from the same organ and wherein detection of the plurality of organ-specific proteins with the plurality of detection reagents results in an organ-specific molecular blood fingerprint indicative of the drug side effect in the subject. In certain embodiments, the fingerprint (e.g., the pattern of interaction of the detection reagents with each of the plurality of organ-specific proteins) is the combination of, a snapshot of sorts, of the different quantitative levels of the organ-specific proteins detected. Thus, in other words, the fingerprint is a set of numbers, each number corresponding to a level of a particular organ-specific protein. This set of numbers and the specific organ-specific proteins that they correspond to together make up the unique fingerprint that defines a biological condition. In this regard, the detection reagents may comprise antibodies or antigen-binding fragments thereof or monoclonal antibodies, or antigen-binding fragments thereof. The panels of the present invention may comprise from at least about 2 detection reagents to about 100, 150, 160, 170, 180, 190, 200 or more detection reagents. In one embodiment, the panel comprises about 5, 10, or 20 detection reagents. In a further embodiment, the plurality of detection reagents are specific for kidney-specific proteins, liver-specific proteins, cardiac-specific proteins, or indeed specific for proteins derived from any organ as described herein.

BRIEF DESCRIPTION OF THE SEQUENCE IDENTIFIERS

SEQ ID NO:1 is the cDNA sequence that encodes the WDR19 prostate specific secreted protein.

SEQ ID NO:2. is the amino acid sequence of the WDR19 prostate specific secreted protein.

SEQ ID NOs:3-72 are MPSS signature sequences that correspond to differentially expressed genes in LNCaP cells (early prostate cancer phenotype) to androgen-independent CL1 cells (late prostate cancer phenotype) (see Table 1).

SEQ ID NOs:73-593 are MPSS signature sequences that correspond to differentially expressed genes in prostate cancer cell lines LNCaP and CL1 that encode secreted proteins (see Table 3).

SEQ ID NOs:594-1511 are the GENBANK sequences of differentially expressed genes that encode predicted secreted proteins as referred to in Table 3. Both polynucleotide and amino acid sequences are provided for each GENBANK accession number.

SEQ ID NOs:1512-1573 are the amino acid sequences from GENBANK of prostate-specific proteins potentially secreted into blood as described in Table 4.

SEQ ID NOs:1574-1687 are the GENBANK sequences of examples of differentially expressed genes as described in Table 1. Both polynucleotide and amino acid sequences are provided where available for each GENBANK accession number.

SEQ ID NOs:1688-1796 are MPSS signature sequences that correspond to prostate-specific/enriched genes as described in Table 5.

SEQ ID NOs:1797-1947 are the GENBANK sequences of prostate-specific genes as described in Table 5. Both polynucleotide and amino acid sequences are provided where available for each GENBANK accession number.

DETAILED DESCRIPTION OF THE INVENTION

A powerful new systems approach to disease is revealing powerful new blood diagnostics/monitoring approaches. Particularly, in specific cells there are protein and gene regulatory networks that mediate the normal functions of the cell. The disease process causes one or more of these networks to be perturbed, either genetically or environmentally (e.g. infections). The disease-altered networks result in altered patterns of protein expression—and some of the transcripts with altered expression levels are organ (cell)-specific and some of these organ-specific transcripts encode secreted proteins. Hence disease leads to altered expression patterns of organ-specific, secreted proteins in the blood. Drugs also cause altered expression of organ-specific secreted proteins in the blood. In particular, the liver and kidney are organs that often reflect the side effects of drugs.

Hence the blood may be viewed as a window into the health and disease of an individual. The levels of organ-specific secreted proteins present in the blood taken together represent molecular fingerprints in the blood that reflect the operation of normal organs. Each organ has a specific quantitative molecular fingerprint. When a drug has a side effect on a particular organ, that blood fingerprint changes, for example, in the levels of these proteins expressed in the blood and the change in the fingerprint correlates with the specific effect the drug has on the organ. The changes in the fingerprints occur as a consequence of virtually any disease or organ perturbation (e.g., drug effect) with each disease or drug effect resulting in a unique fingerprint. The changes in the fingerprints are sufficiently informative to visualize side effects of drugs, be they adverse or positive. Thus, as used herein, side effect refers to any unintended effect of a drug, either positive (e.g., a previously unrecognized positive indication for a drug) or negative (e.g., toxicity or other adverse effect). The drug-altered fingerprints are determined by comparing the blood from normal individuals against that from patients on a particular drug regimen. Not only will the absolute levels of the changes in the proteins constituting individual fingerprints be determined, but all the protein changes (e.g. N changed proteins) will be compared against one another to generate an N-dimensional shape space that will correlate even more powerfully with the stratifications of drug-induced alterations as described herein (see e.g., U.S. Patent Application No. 20020095259).

The studies described herein use prostate cancer as a model for studying perturbation of organ-specific molecular blood fingerprints. The same principles apply in the setting of determining perturbations that result from drugs. In the studies described herein, the transcriptomes of two prostate cancer cell lines were analyzed: LNCaP, an androgen sensitive cell

line, and hence a model for early stage of prostate cancer; and a variant of this cell, CL1, an androgen unresponsive cell line, thus, a model for late stage of prostate cancer. Analyses of the transcriptomes of these two cell lines revealed changes in cellular states that occur with the progression of prostate cancer. These transcriptomes were also compared to normal prostate tissue, prostate cancer tissues and prostate cancer metastases. These prostate transcriptomes were compared against their counterparts from 29 other tissues to identify those transcripts that are primarily expressed in the prostate. Computational approaches were used to predict which of these transcripts encode secreted proteins. Further, a prostate protein, referred to as WDR19, that was previously shown by microarray and northern analysis to be prostate-specific, was used in a multiparameter analysis of prostate cancer samples.

Thus, the present invention is generally directed to methods for identifying organ-specific secreted proteins present in the blood. The present invention is also directed to methods for defining organ-specific molecular blood fingerprints and further provides defined examples of predicted organ-specific molecular blood fingerprints. Additionally, the present invention is directed to panels of reagents or proteomic techniques employing mass spectrometry that detect organ-specific secreted proteins in the blood for use in identifying side effects of drugs, evaluating drug toxicity, and other related applications.

By predefining the components of a given molecular blood fingerprint using the methods described herein, the present invention alleviates the need to blindly search for protein patterns using blood proteomics. Thus, the present invention enables the skilled artisan to 1) identify blood proteins which collectively constitute unique organ-specific molecular blood fingerprints for healthy, diseased individuals and individuals affected (either adversely or positively) by one or more drugs; 2) identify unique organ-specific molecular blood fingerprints associated with the direct (e.g., intended) effects of drugs or the side effects of different drugs; 3) identify fingerprints that can uniquely distinguish the different types of side effects. Importantly, the organ-specific, secreted blood fingerprints can be predicted from a combination of quantitative comparative transcriptome studies and computational methods to predict which transcripts encode secreted proteins. The methods for determining the organ-specific, blood fingerprints for all organs described herein allow drug effects (either adverse or positive) on any organ to be easily identified. Further, the present invention can be used to determine distinct, normal organ-specific molecular blood fingerprints, such as in different populations of people. In this regard, there may be differences in normal organ-specific molecular blood fingerprints between populations of individuals that permit the stratification of patients into classes of individuals who would respond positively to a particular drug and those who would not. Thus, the present invention provides the ability to determine those individuals who may have adverse reactions to drugs.

A drug, as used herein, refers to any substance (synthetic or natural) which when administered to or otherwise absorbed into a living organism or system derived therefrom, may modify one or more of its functions.

Methods for Identifying Organ-Specific Proteins Secreted Into the Blood.

The invention provides methods for identifying organ-specific secreted proteins. In this regard, as used herein, the term "organ" is defined as would be understood in the art. Thus, the term, "organ-specific" as used herein refers to proteins (or transcripts) that are primarily expressed in a single organ. It

should be noted that the skilled artisan would readily appreciate upon reading the instant specification that cell-specific transcripts and proteins and tissue-specific transcripts and proteins are also contemplated in the present invention. As such, and as discussed further herein, in certain embodiments, organ-specific protein is defined as a protein encoded by a transcript that is expressed at a level of at least 3 copies/million (as measured, for example, by massively parallel signature sequencing (MPSS) in the cell/tissue/organ of interest but is expressed at less than 3 copies/million in other cells/tissues/organs. In a further embodiment, an organ-specific protein is one that is encoded by a transcript that is expressed 95% in one organ and the remaining 5% in one or more other organs. (In this context, total expression across all organs examined is taken as 100%).

In certain embodiments, an organ-specific protein is one that is encoded by a transcript that is expressed at about 50%, 55%, 60%, 65%, 70%, 75%, 80% to about 90% in one organ and wherein the remaining 10%-50% is expressed in one or more other organs. As would be readily recognized by the skilled artisan upon reading the present disclosure, in certain embodiments, an organ-specific molecular blood fingerprint can readily be discerned even if some expression of an "organ-specific" protein from a particular organ is detected at some level in another organ, or even more than one organ. For example, the organ-specific molecular blood fingerprint from prostate can conclusively identify a particular prostate disease (and stage of disease) despite expression of one or more protein members of the fingerprint in one or more other organs. Thus, an organ-specific protein as described herein may be predominantly or differentially expressed in an organ of interest rather than uniquely or specifically expressed in the organ. In this regard, in certain embodiments, differentially expressed means at least 1.5 fold expression in the organ of interest as compared to other organs. In another embodiment, differentially expressed means at least 2 fold expression in the organ of interest as compared to expression in other organs. In yet a further embodiment, differentially expressed means at least 2.5, 3, 3.5, 4, 4.5, 5 fold or higher expression in the organ of interest as compared to expression of the protein in other organs. As described elsewhere herein, "protein" expression can be determined by analysis of transcript expression using a variety of methods.

In one embodiment, the organ-specific proteins are identified by preparing a cDNA library from an organ of interest. Any organ of a mammalian body is contemplated herein. Illustrative organs include, but are not limited to, heart, kidney, ureter, bladder, urethra, liver, prostate, heart, blood vessels, bone marrow, skeletal muscle, smooth muscle, brain (amygdala, caudate nucleus, cerebellum, corpus callosum, fetal, hypothalamus, thalamus), spinal cord, peripheral nerves, retina, nose, trachea, lungs, mouth, salivary gland, esophagus, stomach, small intestines, large intestines, hypothalamus, pituitary, thyroid, pancreas, adrenal glands, ovaries, oviducts, uterus, placenta, vagina, mammary glands, testes, seminal vesicles, penis, lymph nodes, PBMC, thymus, and spleen. As noted above, upon reading the present disclosure, the skilled artisan would recognize that cell-specific and tissue-specific proteins are contemplated herein and thus, proteins specifically expressed in cells or tissues that make up such organs are also contemplated herein. In certain embodiments, in each of these organs, transcriptomes are obtained for the cell types in which the disease of interest arises. For example, in the prostate there are two dominant types of cells—epithelial cells and stromal cells. About 98% of prostate cancers arise in epithelial cells. As such, in certain embodiments, "organ-specific" means the transcripts that are

expressed in particular cell types of the organ of interest (e.g., prostate epithelial cells). In this regard, any cell type that makes up any of the organs described herein is contemplated herein. Illustrative cell types include, but are not limited to, epithelial cells, stromal cells, endothelial cells, endodermal cells, ectodermal cells, mesodermal cells, lymphocytes (e.g., B cells and T cells including CD4+ T helper 1 or T helper 2 type cells, CD8+ cytotoxic T cells), erythrocytes, keratinocytes, and fibroblasts. Particular cell types within organs or tissues may be obtained by histological dissection, by the use of specific cell lines (e.g., prostate epithelial cell lines), by cell sorting or by a variety of other techniques known in the art.

It should be noted that in certain embodiments, fingerprints can be determined from "organ-specific" proteins from multiple organs, such as from organs that share a common function or make up a system (e.g., digestive system, circulatory system, respiratory system, the immune system (including the different cells of the immune system, such as, but not limited to, B cells, T cells including CD4+ T helper 1 or T helper 2 type cells, regulatory T cells, CD8+ cytotoxic T cells, NK cells, dendritic cells, macrophages, monocytes, neutrophils, granulocytes, mast cells, etc.), cardiovascular system, the sensory system, the skin, brain and the nervous system, and the like).

Complementary DNA (cDNA) libraries can be generated using techniques known in the art, such as those described in Ausubel et al. (2001 *Current Protocols in Molecular Biology*, Greene Publ. Assoc. Inc. & John Wiley & Sons, Inc., NY, N.Y.); Sambrook et al. (1989 *Molecular Cloning*, Second Ed., Cold Spring Harbor Laboratory, Plainview, N.Y.); Maniatis et al. (1982 *Molecular Cloning*, Cold Spring Harbor Laboratory, Plainview, N.Y.) and elsewhere. Further, a variety of commercially available kits for constructing cDNA libraries are useful for making the cDNA libraries of the present invention. Libraries are constructed from organs/tissues/cells procured from normal subjects.

All or substantially all of the transcripts of the cDNA library, e.g., representing virtually or substantially all genes functioning in the organ of interest, are cloned and sequenced using any of a variety of techniques known in the art. In this regard, in certain embodiments, substantially all refers to a sample representing at least 80% of all genes functioning in the organ of interest. In a further embodiment, substantially all refers to a sample representing at least 85%, 90%, 95%, 96%, 97%, 98% 99% or higher of all genes functioning in the organ of interest. In one embodiment, substantially all the transcripts from a cDNA library are amplified, sorted and signature sequences generated therefrom according to the methods described in U.S. Pat. Nos. 6,013,445; 6,172,218; 6,172,214; 6,140,489 and Brenner, P., et al., *Nat Biotechnol*, 18:630-634 2000. Briefly, polynucleotide templates from a cDNA library of interest are cloned into a vector system that contains a vast set of minimally cross-hybridizing oligonucleotide tags (see U.S. Pat. No. 5,863,722). The number of tags is usually at least 100 times greater than the number of cDNA templates (see e.g., U.S. Pat. No. 6,013,445 and Brenner, P., et al., supra). Thus, the set of tags is such that a 1% sample taken of template-tag conjugates ensures that essentially every template in the sample is conjugated to a unique tag and that at least one of each of the different template cDNAs is represented in the sample with >99% probability (U.S. Pat. No. 6,013,445 and Brenner, P., et al., supra). The conjugates are then amplified and hybridized under stringent conditions to microbeads each of which has attached thereto a unique complementary, minimally cross-hybridizing oligonucleotide tag. The transcripts are then directly sequenced simultaneously in a flow cell using a ligation-based sequenc-

ing method (see e.g., U.S. Pat. No. 6,013,445). A short signature sequence of about 17-20 base pairs is generated simultaneously from each of the hundreds of thousands of beads (or more) in the flow cell, each having attached thereto copies of a unique transcript from the sample. This technique is termed massively parallel signature sequencing (MPSS).

In certain embodiments, other techniques may be used to evaluate the transcripts from a particular cDNA library, including microarray analysis (Han, M., et al., *Nat Biotechnol*, 19: 631-635, 2001; Bao, P., et al., *Anal Chem*, 74: 1792-1797, 2002; Schena et al., *Proc. Natl. Acad. Sci. USA* 93:10614-19, 1996; and Heller et al., *Proc. Natl. Acad. Sci. USA* 94:2150-55, 1997) and SAGE (serial analysis of gene expression). Like MPSS, SAGE is digital and can generate a large number of signature sequences. (see e.g., Velculescu, V. E., et al., *Trends Genet*, 16: 423-425., 2000; Tuteja R. and Tuteja N. *Bioessays*. 2004 August; 26(8):916-22) although the coverage is not nearly as deep as with MPSS.

The resulting sequences, (e.g., MPSS signature sequences), are generally about 20 bases in length. However, in certain embodiments, the sequences can be about 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, or 100 or more bases in length. The sequences are annotated using annotated human genome sequence (such as human genome release hg16, released in November, 2003, or other public databases) and the human Unigene (Unigene build #184) using methods known in the art, such as the method described by Meyers, B. C., et al., *Genome Res*, 14: 1641-1653, 2004. Other databases useful in this regard include Genbank, EMBL, or other publicly available databases. In certain embodiments, transcripts are considered only for those with 100% matches between an MPSS or other type of signature and a genome signature. As would be readily appreciated by the skilled artisan upon reading the present disclosure, this is a stringent match criterion and in certain embodiments, it may be desirable to use less stringent match criteria. Indeed, polymorphisms could lead to variations in transcripts that would be missed if only exact matches were used. For example, it may be desirable to consider signature sequences that match a genome signature with 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identity. In one embodiment, signatures that are expressed at less than 3 transcripts per million in libraries of interest are disregarded, as they might not be reliably detected since this, in effect, represents less than one transcript per cell (see for example, Jongeneel, C. V., et al., *Proc Natl Acad Sci U S A*, 2003). cDNA signatures are classified by their positions relative to polyadenylation signals and poly (A) tails and by their orientation relative to the 5'→3' orientation of source mRNA. Full-length sequences corresponding to the signature sequences can be thus identified.

In order to identify organ-specific transcripts, the resulting annotated transcripts are compared against public and/or private sequence databases, such as a variety of annotated human genome sequence databases (e.g., Genebank, the EMBL and Japanese databases and databases generated and compiled from other normal tissues, to identify those transcripts that are expressed primarily in the organ of interest but are not expressed in other organs. As noted elsewhere herein, some expression in organs other than the organ of interest does not necessarily preclude the use of a particular transcript in a blood molecular signature panel of the present invention.

Comparisons of the transcripts between databases can be made using a variety of computer analysis algorithms known in the art. As such, alignment of sequences for comparison may be conducted by the local identity algorithm of Smith and Waterman (1981) *Add. APL Math* 2:482, by the identity alignment algorithm of Needleman and Wunsch (1970) *J.*

Mol. Biol. 48:443, by the search for similarity methods of Pearson and Lipman (1988) *Proc. Natl. Acad. Sci. USA* 85: 2444, by computerized implementations of these algorithms (GAP, BESTFIT, BLAST, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group (GCG), 575 Science Dr., Madison, Wis.), or by inspection. As would be understood by the skilled artisan, many algorithms are available and are continually being developed. Appropriate algorithms can be chosen based on the specific needs for the comparisons being made (See also, e.g., J. A. Cuff, et al., *Bioinformatics*, 16(2):111-116, 2000; S. F. Altschul and B. W. Erickson. *Bulletin of Mathematical Biology*, 48(5/6):603-616, 1986; S. F. Altschul and B. W. Erickson. *Bulletin of Mathematical Biology*, 48(5/6):633-660, 1986; S. F. Altschul, et al., *J. Mol. Bio.*, 215:403-410, 1990; K. Buckalassen, et al., *BIOINFORMATICS*, 15(2):122-130, 1999; K.-M. Chao, et al., *Bulletin of Mathematical Biology*, 55(3): 503-524, 1993; W. M. Fitch and T. F. Smith. *Proceedings of the National Academy of Sciences*, 80:1382-1386, 1983; A. D. Gordon. *Biometrika*, 60:197-200, 1973; O. Gotoh. *J Mol Biol*, 162:705-708, 1982; O. Gotoh. *Bulletin of Mathematical Biology*, 52(3):359-373, 1990; X. Huang, et al., *CABIOS*, 6:373-381, 1990; X. Huang and W. Miller. *Advances in Applied Mathematics*, 12:337-357, 1991; J. D. Thompson, et al., *Nucleic Acids Research*, 27(13):2682-2690, 1999).

In certain embodiments, a particular transcript is considered to be organ-specific when the number of transcripts/million as determined by MPSS is 3 or greater in the organ of interest but is less than 3 in all other organs. In another embodiment, a transcript is considered organ-specific if it is expressed in the organ of interest at a detectable level using a standard measurement (e.g., microarray analysis, quantitative real-time RT-PCR, MPSS, etc.) in the organ of interest but is not detectably expressed in other organs, using appropriate negative and positive controls as would be familiar to the skilled artisan. In a further embodiment, an organ-specific transcript is one that is expressed 95% in one organ and the remaining 5% in one or more other organs. (In this context, total expression across all organs examined is taken as 100%). In certain embodiments, an organ-specific transcript is one that is expressed at about 50%, 55%, 60%, 65%, 70%, 75%, 80% to about 90% in one organ and wherein the remaining 10%-50% is expressed in one or more other organs.

In another embodiment, organ-specific transcripts are identified by determining the ratio of expression of a transcript in the organ of interest as compared to other organs. In this regard, expression levels in the organ of interest of at least 2.0, 2.5, 3.0, 3.5, 4.0, 4.5, 5.0, 5.5, 6.0, 6.5, 7.0 fold or higher as compared to expression in all other organs is considered to be organ-specific expression.

As would be readily recognized by the skilled artisan upon reading the present disclosure, in certain embodiments, an organ-specific molecular blood fingerprint can readily be discerned even if some expression of an "organ-specific" protein from a particular organ is detected at some level in another organ, or even more than one organ. This is because the fingerprint (e.g., the combination of the levels of multiple proteins; the pattern of the expression levels of multiple markers) itself is unique despite that the expression levels of one or more individual members of the fingerprint may not be unique to a particular organ. For example, the organ-specific molecular blood fingerprint from prostate can conclusively identify a particular prostate disease (and stage of disease) despite some expression of one or more members of the fingerprint in one or more other organs. Thus the present invention relates to determining the presence or absence of a disease or condition or stage of disease based on a pattern

(e.g., fingerprint) of markers measured concurrently using any one or more of a variety of methods described herein (e.g., antibody binding, mass spectrometry, and the like), rather than the measure of individual markers.

In further embodiments, specificity can be confirmed at the protein level using immunohistochemistry (IHC) and/or other protein measurement techniques known in the art (e.g., isotope-coded affinity tags and mass spectrometry, such as described by Han, D. K., et al., *Nat Biotechnol.* 19: 946-951, 2001). The Z-test (Man, M. Z., et al., *Bioinformatics*, 16: 953-959, 2000) or other appropriate statistical tests can be used to calculate P values for comparison of gene and protein expression levels between libraries from organs of interest.

Organ-specific sequences identified as described herein are further analyzed to determine which of the sequences encode secreted proteins. Proteins with signal peptides (classical secretory proteins) can be predicted using computation analysis known in the art. Illustrative methods include, but are not limited to the criteria described by Chen et al., *Mamm Genome*, 14: 859-865, 2003. In certain embodiments, such analyses are carried out using prediction servers, for example SignalP 3.0 server developed by The Center for Biological Sequence Analysis, Lyngby, Denmark (<http://www.cbs.dtu.dk/services/SignalP-3.0/>); see also, J. D. Bendtsen, et al., *J. Mol. Biol.*, 340:783-795, 2004.) and the TMHMM2.0 server (see for example A. Krogh, et al., *Journal of Molecular Biology*, 305(3):567-580, January 2001; E. L. L. Sonnhammer, et al., In J. Glasgow, T. Littlejohn, F. Major, R. Lathrop, D. Sankoff, and C. Sensen, editors, *Proceedings of the Sixth International Conference on Intelligent Systems for Molecular Biology*, pages 175-182, Menlo Park, Calif., 1998. AAAI Press). Other prediction methods that can be used in the context of the present invention include those described for example, in S. Moller, M.D.R. et al., *Bioinformatics*, 17(7):646-653, July 2001. Nonclassical secretory secreted proteins (without signal peptides) can be predicted using, for example, the SecretomeP 1.0 server, (<http://www.cbs.dtu.dk/services/SecretomeP-1.0/>) with an odds ratio score > 3.0. Updated versions of these analysis programs are also contemplated for use in the present methods as are other methods known in the art (e.g., PSORT (<http://nibb.ac.jp/>) and Sigfind (<http://colobal.com/sigfind/>)).

Confirmation that the identified secreted proteins are present in blood can be carried out using a variety of methods known in the art. For example, the proteins can be expressed, purified, and specific antibodies can be made against them. The specific antibodies can then be used to test the presence of the protein in blood/serum/plasma by a variety of immunoaffinity based techniques (e.g., immunoblot, Western analysis, immunoprecipitation, ELISA, etc.). Antibodies specific for the organ-specific protein identified herein can also be used to study expression patterns of the identified proteins. It should be noted that in certain circumstances, the secreted protein may not be detectable in normal blood samples but will be detected in the blood as a result of perturbation due to disease or other environmental factors. Accordingly, both normal and disease samples are tested for the presence of the secreted protein and particularly for changes in levels of expression in the two states. As an alternative, aptamers (short DNA or RNA fragments with binding complementarity to the proteins of interest) may be used in assays similar to those described for antibodies (see for example, *Biotechniques*. 2001 February; 30(2):290-2, 294-5; *Clinical Chemistry*. 1999;45:1628-1650). In addition, antibodies or aptamers may be used in connection with nanowires to create highly sensitive detec-

tions systems (see e.g., J. Heath et al., *Science*. 2004 Dec. 17; 306(5704):2055-6). In further embodiments, mass spectrometry-based methods can be used to confirm the presence of a particular protein in the blood.

As would be recognized by the skilled artisan, while the organ-specific secreted proteins, the levels of which make up a given fingerprint, need not be isolated, in certain embodiments, it may be desirable to isolate such proteins (e.g., for antibody production). As such, the present invention provides for isolated organ-specific secreted proteins or fragments or portions thereof and polynucleotides that encode such proteins. As used herein, the terms protein and polypeptide are used interchangeably. The terms "polypeptide" and "protein" encompass amino acid chains of any length, including full-length endogenous (i.e., native) proteins and variants of endogenous polypeptides described herein. Illustrative polypeptides of the present invention are described in Table 1 and Tables 3-5, the section entitled "Brief Description of the Sequence Identifiers" and are set forth in the sequence listing. "Variants" are polypeptides that differ in sequence from the polypeptides of the present invention only in substitutions, deletions and/or other modifications, such that either the variants' disease-specific expression patterns are not significantly altered or the polypeptides remain useful for diagnostics/detection of organ-specific blood fingerprints as described herein. For example, modifications to the polypeptides of the present invention may be made in the laboratory to facilitate expression and/or purification and/or to improve immunogenicity for the generation of appropriate antibodies and other binding agents, etc. Modified variants (e.g., chemically modified) of the polypeptides of organ-specific, secreted proteins may be useful herein, (e.g., as standards in mass spectrometry analyses of the corresponding proteins in the blood, and the like). As such, in certain embodiments, the biological function of a variant protein is not relevant for utility in the methods for detection and/or diagnostics described herein. Polypeptide variants generally encompassed by the present invention will typically exhibit at least about 70%, 75%, 80%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% or more identity along its length, to a polypeptide sequence set forth herein. Within a polypeptide variant, amino acid substitutions are usually made at no more than 50% of the amino acid residues in the native polypeptide, and in certain embodiments, at no more than 25% of the amino acid residues. In certain embodiments, such substitutions are conservative. A conservative substitution is one in which an amino acid is substituted for another amino acid that has similar properties, such that one skilled in the art of peptide chemistry would expect the secondary structure and hydrophobic nature of the polypeptide to be substantially unchanged. In general, the following amino acids represent conservative changes: (1) ala, pro, gly, glu, asp, gln, asn, ser, thr; (2) cys, ser, tyr, thr; (3) val, ile, leu, met, ala, phe; (4) lys, arg, his; and (5) phe, tyr, trp, his. Thus, a variant may comprise only a portion of a native polypeptide sequence as provided herein. In addition, or alternatively, variants may contain additional amino acid sequences (such as, for example, linkers, tags and/or ligands), usually at the amino and/or carboxy termini. Such sequences may be used, for example, to facilitate purification, detection or cellular uptake of the polypeptide.

When comparing polypeptide sequences, two sequences are said to be "identical" if the sequence of amino acids in the two sequences is the same when aligned for maximum correspondence, as described below. Comparisons between two sequences are typically performed by comparing the sequences over a comparison window to identify and com-

pare local regions of sequence similarity. A "comparison window" as used herein, refers to a segment of at least about 20 contiguous positions, usually 30 to about 75, 40 to about 50, in which a sequence may be compared to a reference sequence of the same number of contiguous positions after the two sequences are optimally aligned.

Optimal alignment of sequences for comparison may be conducted using the Megalign program in the Lasergene suite of bioinformatics software (DNASTAR, Inc., Madison, Wis.), using default parameters. This program embodies several alignment schemes described in the following references: Dayhoff, M. O. (1978) A model of evolutionary change in proteins—Matrices for detecting distant relationships. In Dayhoff, M. O. (ed.) *Atlas of Protein Sequence and Structure*, National Biomedical Research Foundation, Washington D.C. Vol. 5, Suppl. 3, pp. 345-358; Hein J. (1990) Unified Approach to Alignment and Phylogenesis pp. 626-645 *Methods in Enzymology* vol. 183, Academic Press, Inc., San Diego, Calif.; Higgins, D. G. and Sharp, P. M. (1989) *CABIOS* 5:151-153; Myers, E. W. and Muller W. (1988) *CABIOS* 4:11-17; Robinson, E. D. (1971) *Comb. Theor* 11:105; Saitou, N. Nei, M. (1987) *Mol. Biol. Evol.* 4:406-425; Sneath, P. H. A. and Sokal, R. R. (1973) *Numerical Taxonomy—the Principles and Practice of Numerical Taxonomy*, Freeman Press, San Francisco, Calif.; Wilbur, W. J. and Lipman, D. J. (1983) *Proc. Natl. Acad., Sci. USA* 80:726-730.

Alternatively, optimal alignment of sequences for comparison may be conducted by the local identity algorithm of Smith and Waterman (1981) *Add. APL. Math* 2:482, by the identity alignment algorithm of Needleman and Wunsch (1970) *J. Mol. Biol.* 48:443, by the search for similarity methods of Pearson and Lipman (1988) *Proc. Natl. Acad. Sci. USA* 85: 2444, by computerized implementations of these algorithms (GAP, BESTFIT, BLAST, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group (GCG), 575 Science Dr., Madison, Wis.), or by inspection.

Illustrative examples of algorithms that are suitable for determining percent sequence identity and sequence similarity include the BLAST and BLAST 2.0 algorithms, which are described in Altschul et al. (1977) *Nucl. Acids Res.* 25:3389-3402 and Altschul et al. (1990) *J. Mol. Biol.* 215:403-410, respectively. BLAST and BLAST 2.0 can be used, for example, to determine percent sequence identity for the polynucleotides and polypeptides of the invention. Software for performing BLAST analyses is publicly available through the National Center for Biotechnology Information.

An isolated polypeptide is one that is removed from its original environment. For example, a naturally occurring protein or polypeptide is isolated if it is separated from some or all of the coexisting materials in the natural system. In certain embodiments, such polypeptides are also purified, e.g., are at least about 90% pure, in some embodiments, at least about 95% pure and in further embodiments, at least about 99% pure.

In one embodiment of the present invention, a polypeptide comprises a fusion protein comprising an organ-specific secreted polypeptide. The present invention further provides, in other aspects, fusion proteins that comprise at least one polypeptide as described herein, as well as polynucleotides encoding such fusion proteins. The fusion proteins may comprise multiple polypeptides or portions/variants thereof, as described herein, and may further comprise one or more polypeptide segments for facilitating the expression, purification, detection, and/or activity of the polypeptide(s).

In certain embodiments, the proteins and/or polynucleotides, and/or fusion proteins are provided in the form of

compositions, e.g., pharmaceutical compositions, vaccine compositions, compositions comprising a physiologically acceptable carrier or excipient. Such compositions may comprise buffers such as neutral buffered saline, phosphate buffered saline and the like; carbohydrates such as glucose, mannose, sucrose or dextrans, mannitol; proteins; polypeptides or amino acids such as glycine; antioxidants; chelating agents such as EDTA or glutathione; adjuvants (e.g., aluminum hydroxide); and preservatives.

In general, organ-specific secreted polypeptides and polynucleotides encoding such polypeptides as described herein, may be prepared using any of a variety of techniques that are well known in the art. For example, a DNA sequence encoding an organ-specific secreted protein may be prepared by amplification from a suitable cDNA or genomic library using, for example, polymerase chain reaction (PCR) or hybridization techniques. Libraries may generally be prepared and screened using methods well known to those of ordinary skill in the art, such as those described in Sambrook et al., *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratories, Cold Spring Harbor, N.Y., 1989. cDNA libraries may be prepared from any of a variety of organs, tissues, cells, as described herein. Other libraries that may be employed will be apparent to those of ordinary skill in the art upon reading the present disclosure. Primers for use in amplification may be readily designed based on the polynucleotide sequences encoding organ-specific polypeptides as provided herein, for example, using programs such as the PRIMER3 program (http://double slash www-genome dot wi dot mit dot edu/cgi-bin/primer/primer3_www dot cgi).

Polynucleotides encoding the organ-specific secreted polypeptides as described herein are also provided by the present invention. A polynucleotide as used herein may be single-stranded (coding or antisense) or double-stranded, and may be DNA (genomic, cDNA or synthetic) or RNA molecules. Thus, within the context of the present invention, a polynucleotide encoding a polypeptide may also be a gene. A gene is a segment of DNA involved in producing a polypeptide chain; it includes regions preceding and following the coding region (leader and trailer) as well as intervening sequences (introns) between individual coding segments (exons). Additional coding or non-coding sequences may, but need not, be present within a polynucleotide of the present invention, and a polynucleotide may, but need not, be linked to other molecules and/or support materials. An isolated polynucleotide, as used herein, means that a polynucleotide is substantially away from other coding sequences, and that the DNA molecule does not contain large portions of unrelated coding DNA, such as large chromosomal fragments or other functional genes or polypeptide coding regions. Of course, this refers to the DNA molecule as originally isolated, and does not exclude genes or coding regions later added to the segment by the hand of man.

Polynucleotides of the present invention may comprise a native sequence (i.e., an endogenous polynucleotide, for instance, a native or non-artificially engineered or naturally occurring gene as provided herein) encoding an organ-specific secreted protein, an alternate form of such a sequence, or a portion or splice variant thereof or may comprise a variant of such a sequence. Polynucleotide variants may contain one or more substitutions, additions, deletions and/or insertions such that the polynucleotide encodes a polypeptide useful in the methods described herein, such as for the detection of organ-specific proteins (e.g., wherein said polynucleotide variants encode polypeptides that can be used to generate detection reagents as described herein that are specific for an organ-specific secreted protein). In certain embodiments,

variants exhibit at least about 70% identity, and in other embodiments, exhibit at least about 80%, 85%, 86%, 87%, 88%, 89%, identity and in yet further embodiments, at least about 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% identity to a polynucleotide sequence that encodes a native organ-specific secreted polypeptide or an alternate form or a portion thereof. Illustrative polynucleotides of the present invention are described in Table 1 and Tables 3-5, the section entitled "Brief Description of the Sequence Identifiers" and are set forth in the sequence listing. The percent identity may be readily determined by comparing sequences using computer algorithms well known to those having ordinary skill in the art and described herein.

Polynucleotides that are complementary to the polynucleotides described herein, or that have substantial identity to a sequence complementary to a polynucleotide as described herein are also within the scope of the present invention. "Substantial identity", as used herein refers to polynucleotides that exhibit at least about 70% identity, and in certain embodiments, at least about 80%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% identity to a polynucleotide sequence that encodes a native organ-specific secreted polypeptide as described herein. Substantial identity can also refer to polynucleotides that are capable of hybridizing under stringent conditions to a polynucleotide complementary to a polynucleotide encoding an organ-specific secreted protein. Suitable hybridization conditions include prewashing in a solution of 5xSSC, 0.5% SDS, 1.0 mM EDTA (pH 8.0); hybridizing at 50-65° C., 5xSSC, overnight; followed by washing twice at 65° C. for 20 minutes with each of 2x, 0.5x and 0.2xSSC containing 0.1% SDS. Nucleotide sequences that, because of code degeneracy, encode a polypeptide encoded by any of the above sequences are also encompassed by the present invention.

Oligonucleotide primers for amplification of the polynucleotides encoding organ-specific secreted proteins are also within the scope of the present invention. Many amplification methods are known in the art such as PCR, RT-PCR, quantitative real-time PCR, and the like. The PCR conditions used can be optimized in terms of temperature, annealing times, extension times and number of cycles depending on the oligonucleotide and the polynucleotide to be amplified. Such techniques are well known in the art and are described in, for example, Mullis et al., *Cold Spring Harbor Symp. Quant. Biol.*, 51:263, 1987; Erlich ed., *PCR Technology*, Stockton Press, NY, 1989. Oligonucleotide primers can be anywhere from 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, or 30 nucleotides in length. In certain embodiments, the oligonucleotide primers of the present invention are typically 35, 40, 45, 50, 55, 60, or more nucleotides in length.

Organ-Specific Molecular Blood Fingerprints

The present invention also provides methods for defining organ-specific molecular blood fingerprints. Additionally, the present invention provides defined examples of organ-specific molecular blood fingerprints as described further herein.

Each normal organ controls the expression of a variety of genes, some of which are expressed at major levels at other organs or tissues in the body and some of which are expressed only in the organ of interest or at significantly increased levels in the organ of interest as compared to expression in other organs/tissues (e.g., at least 2 fold, at least 2.5 fold, at least 3.0 fold, at least 3.5 fold, at least 4.0 fold, at least 4.5 fold, or higher fold expression in the organ of interest as compared to other tissues. Some of the organ-specific transcripts encode proteins which can be secreted into the blood. Hence these

secreted proteins constitute an organ-specific molecular fingerprint for that organ in the blood. Analysis of levels of these proteins in the blood provides organ-specific molecular blood fingerprints that are indicative of biological states. A biological state may be a normal, healthy state or a disease state (e.g., perturbation from normal). Thus, there are molecular fingerprints in the blood that reflect the operation of normal organs and each organ has a specific molecular fingerprint. These organ-specific blood fingerprints are perturbed when disease, or other agents such as drugs, affects the organ. Different diseases will alter the organ-specific blood fingerprints in different ways (e.g. alter the expression levels of the corresponding secreted proteins). Likewise, different drugs will alter the organ-specific blood fingerprints in different ways. Thus, a unique perturbed blood molecular fingerprint is associated with each type of distinct disease and with each drug or combination of drugs. In effect, each drug or combination of drugs will create a unique organ-specific molecular blood fingerprint for each organ that it affects. As would be readily appreciated by the skilled artisan, each disease or stage of a disease or drug or combination of drugs can affect multiple organs. For example, in kidney cancer, a primary perturbation in the kidney-specific molecular blood fingerprint would occur. However, a secondary or indirect effect may also be observed in the bladder-specific molecular blood fingerprint. As another example, in liver cancer, perturbation of a liver-specific blood fingerprint as a primary indicator of disease would occur. However, secondary or indirect effects at other sites, for example in a lymphocyte-specific blood fingerprint, would also be observed. As described elsewhere herein, each disease type and stage results in a unique, identifiable fingerprint for each organ that it affects, for primary and secondary organs affected. Likewise, each drug or combination of drugs results in a unique, identifiable fingerprint for each organ that it affects, both primary and secondary organs. Thus, multiple organ-specific molecular blood fingerprints can be used in combination to determine a particular drug side effect and the fingerprints may include those for the primary organ affected and/or for a secondary or indirect organ that is affected by a particular drug or combination of drugs.

Most common diseases such as prostate cancer actually represent multiple distinct diseases that initially appear similar (e.g., benign and very slowly growing prostate cancer, slowly invasive prostate cancer and rapidly metastatic prostate cancer represent three different types of prostate cancer—the process of dividing individual prostate cancers into one of these three types is called stratification). The blood molecular fingerprints will be distinct for each of these disease types, thus allowing for the stratification of similar diseases and rapid intervention where necessary. The blood fingerprints will also be perturbed in unique ways as each type of disease progresses—hence the blood fingerprints will also permit the progression of disease to be followed. The blood fingerprints also change with therapy, and hence will permit the effectiveness of therapy to be followed, thereby allowing a physician to alter treatment accordingly. Importantly, the blood fingerprints change with exposure to a variety of environmental factors, such as drugs, and can be used to assess toxic or off target damage by the drug and will even permit following the subsequent recovery from such adverse drug exposure.

One of the advantages of the organ-specific, secreted blood fingerprints is the possibility that very subtle side effects of drugs can be detected, either adverse effects, or previously unrecognized positive effects.

Thus, an organ-specific molecular blood fingerprint for a given setting (e.g., one or more particular drug side effects) is

defined by the levels in the blood of the organ-specific proteins that make up the fingerprint. As such, an organ-specific molecular blood fingerprint for a given organ at any given time and in any given setting (e.g., drug-induced perturbation) is determined by measuring the levels of each of a plurality of organ-specific proteins in the blood. It is the combination of the different levels in the blood of the organ-specific proteins that reveals a unique pattern that defines the fingerprint. Equally important, each of the levels of the proteins can be compared against one another to create an N-dimensional measure of the fingerprint space, a very powerful correlate to health, disease, and drug-induced changes (see e.g., U.S. Patent Application No 20020095259). It should be noted that, in certain embodiments, an organ-specific molecular blood fingerprint may be comprised of the determined level in the blood of one or more organ-specific secreted proteins. In one embodiment, an organ-specific molecular blood fingerprint may comprise the determined level in the blood of anywhere from about 2 to more than about 100, 200 or more organ-specific secreted proteins from a particular organ of interest. In one embodiment, the organ-specific molecular blood fingerprint comprises the quantitatively measured level in the blood of at least 2, 3, 4, 5, 6, 7, 8, 9, or 10 organ-specific secreted proteins. In another embodiment, the organ-specific molecular blood fingerprint comprises the determined level in the blood of at least, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 28, 29, or 30 organ-specific secreted proteins. In a further embodiment, the organ-specific molecular blood fingerprint comprises the determined level in the blood of at least, 31, 32, 33, 34, 35, 36, 37, 38, 39, or 40 organ-specific secreted proteins. In yet a further embodiment, the organ-specific molecular blood fingerprint comprises the determined level in the blood of at least, 41, 42, 43, 44, 45, 46, 47, 48, 49, or 50 organ-specific secreted proteins. In an additional embodiment, the organ-specific molecular blood fingerprint comprises the determined level in the blood of 51, 52, 53, 54, 55, 56, 57, 58, 59, or 60 organ-specific secreted proteins. In another embodiment, the organ-specific molecular blood fingerprint comprises the determined level in the blood of 61, 62, 63, 64, 65, 66, 67, 68, 69, or 70 organ-specific secreted proteins. In further embodiments, the organ-specific molecular blood fingerprint comprises the determined level in the blood of 75, 80, 85, 90, 100, or more organ-specific secreted proteins.

It should be noted that in certain circumstances, an organ-specific molecular blood fingerprint can be defined (in part or entirely) merely by the presence or absence of one or a plurality of organ-specific proteins, and determining the exact level of each of a plurality of organ-specific proteins in the blood may not be necessary.

In a further embodiment, the fingerprints associated with a particular drug and side effects thereof are determined by comparing the blood from normal individuals against that from subjects taking a particular drug of interest. As such, a statistically significant change in the levels (e.g., an increase or a decrease) of one or more of the organ-specific proteins that comprise the fingerprint as compared to normal is indicative of a perturbation of the fingerprint and is useful in identifying direct effects or side effects of the drug of interest. The skilled artisan would readily appreciate that a variety of statistical tests can be used to determine if an altered level of a given protein is significant. The Z-test (Man, M. Z., et al., *Bioinformatics*, 16: 953-959, 2000) or other appropriate statistical tests can be used to calculate P values for comparison of protein expression levels. In certain embodiments, the level of each of the plurality of organ-specific proteins in the blood of each of the plurality of organ-specific proteins in the blood sample from the subject is compared to a previously deter-

mined normal control level of each of the plurality of organ-specific proteins taking into account standard deviation. Thus, the present invention provides determined normal control levels of each of a plurality of organ-specific proteins that make up a particular molecular blood fingerprint.

In an additional embodiment, the present invention can be used to determine distinct, normal organ-specific molecular blood fingerprints, such as in different populations of people. In this regard, differences in normal organ-specific molecular blood fingerprints between populations of individuals can be defined and these differences permit the stratification of patients into classes of individuals who would respond positively to a particular drug and those who would not. Thus, the present invention provides the ability to determine those individuals who may have adverse reactions to drugs.

Organ-specific molecular blood fingerprints can be determined using any of a variety of detection reagents in the context of a variety of methods for measuring protein levels. Any detection reagent that can specifically bind to or otherwise detect an organ-specific secreted protein as described herein is contemplated as a suitable detection reagent. Illustrative detection reagents include, but are not limited to antibodies, or antigen-binding fragments thereof, yeast ScFv, DNA or RNA aptamers, isotope labeled peptides, microfluidic/nanotechnology measurement devices and the like.

In one illustrative embodiment, a detection reagent is an antibody or an antigen-binding fragment thereof. Antibodies may be prepared by any of a variety of techniques known to those of ordinary skill in the art. See, e.g., Harlow and Lane, *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory, 1988. In general, antibodies can be produced by cell culture techniques, including the generation of monoclonal antibodies as described herein, or via transfection of antibody genes into suitable bacterial or mammalian cell hosts, in order to allow for the production of recombinant antibodies. In one technique, an immunogen comprising the polypeptide is initially injected into any of a wide variety of mammals (e.g., mice, rats, rabbits, sheep or goats). In this step, the polypeptides of this invention may serve as the immunogen without modification. Alternatively, particularly for relatively short polypeptides, a superior immune response may be elicited if the polypeptide is joined to a carrier protein, such as bovine serum albumin or keyhole limpet hemocyanin. The immunogen is injected into the animal host, usually according to a predetermined schedule incorporating one or more booster immunizations, and the animals are bled periodically. Polyclonal antibodies specific for the polypeptide may then be purified from such antisera by, for example, affinity chromatography using the polypeptide coupled to a suitable solid support.

In one embodiment, multiple target proteins or peptides are used in a single immune response to generate multiple useful detection reagents simultaneously. In one embodiment, the individual specificities are later separated out.

In certain embodiments, antibody can be generated by phage display methods (such as described by Vaughan, T. J., et al., *Nat Biotechnol*, 14: 309-314, 1996; and Knappik, A., et al., *Mol Biol*, 296: 57-86, 2000); ribosomal display (such as described in Hanes, J., et al., *Nat Biotechnol*, 18: 1287-1292, 2000), or periplasmic expression in *E. coli* (see e.g., Chen, G., et al., *Nat Biotechnol*, 19: 537-542, 2001.). In further embodiments, antibodies can be isolated using a yeast surface display library. See e.g., nonimmune library of 10^9 human antibody scFv fragments as constructed by Feldhaus, M. J., et al., *Nat Biotechnol*, 21: 163-170, 2003. There are several advantages of this yeast surface display compared to more traditional large nonimmune human antibody repertoires such as phage

display, ribosomal display, and periplasmic expression in *E. coli* 1). The yeast library can be amplified 10^{10} -fold without measurable loss of clonal diversity and repertoire bias as the expression is under control of the tightly GAL1/10 promoter and expansion can be done under non induction conditions; 2) nanomolar-affinity scFvs can be routinely obtained by magnetic bead screening and flow-cytometric sorting, thus greatly simplified the protocol and capacity of antibody screening; 3) with equilibrium screening, a minimal affinity threshold of the antibodies desired can be set; 4) the binding properties of the antibodies can be quantified directly on the yeast surface; 5) multiplex library screening against multiple antigens simultaneously is possible; and 6) for applications demanding picomolar affinity (e.g. in early diagnosis), subsequent rapid affinity maturation (Kieckhefer, M. C., et al., *J Mol Biol*, 307: 1305-1315, 2001.) can be carried out directly on yeast clones without further re-cloning and manipulations.

Monoclonal antibodies specific for an organ-specific secreted polypeptide of interest may be prepared, for example, using the technique of Kohler and Milstein, *Eur. J. Immunol.* 6:511-519, 1976, and improvements thereto. Briefly, these methods involve the preparation of immortal cell lines capable of producing antibodies having the desired specificity (i.e., reactivity with the polypeptide of interest). Such cell lines may be produced, for example, from spleen cells obtained from an animal immunized as described above. The spleen cells are then immortalized by, for example, fusion with a myeloma cell fusion partner, in certain embodiments, one that is syngeneic with the immunized animal. A variety of fusion techniques may be employed. For example, the spleen cells and myeloma cells may be combined with a nonionic detergent for a few minutes and then plated at low density on a selective medium that supports the growth of hybrid cells, but not myeloma cells. An illustrative selection technique uses HAT (hypoxanthine, aminopterin, thymidine) selection. After a sufficient time, usually about 1 to 2 weeks, colonies of hybrids are observed. Single colonies are selected and their culture supernatants tested for binding activity against the polypeptide. Hybridomas having high reactivity and specificity are preferred.

Monoclonal antibodies may be isolated from the supernatants of growing hybridoma colonies. In addition, various techniques may be employed to enhance the yield, such as injection of the hybridoma cell line into the peritoneal cavity of a suitable vertebrate host, such as a mouse. Monoclonal antibodies may then be harvested from the ascites fluid or the blood. Contaminants may be removed from the antibodies by conventional techniques, such as chromatography, gel filtration, precipitation, and extraction. The polypeptides of this invention may be used in the purification process in, for example, an affinity chromatography step.

A number of therapeutically useful molecules are known in the art which comprise antigen-binding sites that are capable of exhibiting immunological binding properties of an antibody molecule. The proteolytic enzyme papain preferentially cleaves IgG molecules to yield several fragments, two of which (the "F(ab)" fragments) each comprise a covalent heterodimer that includes an intact antigen-binding site. The enzyme pepsin is able to cleave IgG molecules to provide several fragments, including the "F(ab)₂" fragment which comprises both antigen-binding sites. An "Fv" fragment can be produced by preferential proteolytic cleavage of an IgM, and on rare occasions IgG or IgA immunoglobulin molecule. Fv fragments are, however, more commonly derived using recombinant techniques known in the art. The Fv fragment includes a non-covalent V_H::V_L heterodimer including an antigen-binding site which retains much of the antigen rec-

ognition and binding capabilities of the native antibody molecule. Inbar et al. (1972) *Proc. Nat. Acad. Sci. USA* 69:2659-2662; Hochman et al. (1976) *Biochem* 15:2706-2710; and Ehrlich et al. (1980) *Biochem* 19:4091-4096.

A single chain Fv ("sFv") polypeptide is a covalently linked $V_H::V_L$ heterodimer which is expressed from a gene fusion including V_H and V_L -encoding genes linked by a peptide-encoding linker. Huston et al. (1988) *Proc. Nat. Acad. Sci. USA* 85(16):5879-5883. A number of methods have been described to discern chemical structures for converting the naturally aggregated—but chemically separated—light and heavy polypeptide chains from an antibody V region into an sFv molecule which will fold into a three dimensional structure substantially similar to the structure of an antigen-binding site. See, e.g., U.S. Pat. Nos. 5,091,513 and 5,132,405, to Huston et al.; and U.S. Pat. No. 4,946,778, to Ladner et al.

Each of the above-described molecules includes a heavy chain and a light chain CDR set, respectively interposed between a heavy chain and a light chain FR set which provide support to the CDRs and define the spatial relationship of the CDRs relative to each other. As used herein, the term "CDR set" refers to the three hypervariable regions of a heavy or light chain V region. Proceeding from the N-terminus of a heavy or light chain, these regions are denoted as "CDR1," "CDR2," and "CDR3" respectively. An antigen-binding site, therefore, includes six CDRs, comprising the CDR set from each of a heavy and a light chain V region. A polypeptide comprising a single CDR, (e.g., a CDR1, CDR2 or CDR3) is referred to herein as a "molecular recognition unit." Crystallographic analysis of a number of antigen-antibody complexes has demonstrated that the amino acid residues of CDRs form extensive contact with bound antigen, wherein the most extensive antigen contact is with the heavy chain CDR3. Thus, the molecular recognition units are primarily responsible for the specificity of an antigen-binding site.

As used herein, the term "FR set" refers to the four flanking amino acid sequences which frame the CDRs of a CDR set of a heavy or light chain V region. Some FR residues may contact bound antigen; however, FRs are primarily responsible for folding the V region into the antigen-binding site, particularly the FR residues directly adjacent to the CDRs. Within FRs, certain amino residues and certain structural features are very highly conserved. In this regard, all V region sequences contain an internal disulfide loop of around 90 amino acid residues. When the V regions fold into a binding-site, the CDRs are displayed as projecting loop motifs which form an antigen-binding surface. It is generally recognized that there are conserved structural regions of FRs which influence the folded shape of the CDR loops into certain "canonical" structures—regardless of the precise CDR amino acid sequence. Further, certain FR residues are known to participate in non-covalent interdomain contacts which stabilize the interaction of the antibody heavy and light chains.

The detection reagents of the present invention may comprise any of a variety of detectable labels. The invention contemplates the use of any type of detectable label, including, e.g., visually detectable labels, fluorophores, and radioactive labels. The detectable label may be incorporated within or attached, either covalently or non-covalently, to the detection reagent.

Methods for measuring organ-specific protein levels from blood/serum/plasma include, but are not limited to, immunoaffinity based assays such as ELISAs, Western blots, and radioimmunoassays, and mass spectrometry based methods (matrix-assisted laser desorption ionization (MALDI), MALDI-Time-of-Flight (TOF), Tandem MS (MS/MS), elec-

troscopy ionization (ESI), Surface Enhanced Laser Desorption Ionization (SELDI)-TOF MS, liquid chromatography (LC)-MS/MS, etc). Other methods useful in this context include isotope-coded affinity tag (ICAT) followed by multi-dimensional chromatography and MS/MS. The procedures described herein for analysis of blood organ-specific protein fingerprints can be modified and adapted to make use of microfluidics and nanotechnology in order to miniaturize, parallelize, integrate and automate diagnostic procedures (see e.g., L. Hood, et al., *Science* 306:640-643; R. H. Carlson, et al., *Phys. Rev. Lett.* 79:2149 (1997); A. Y. Fu, et al., *Anal. Chem.* 74:2451 (2002); J. W. Hong, et al., *Nature Biotechnol.* 22:435 (2004); A. G. Hadd, et al., *Anal. Chem.* 69:3407 (1997); I. Karube, et al., *Ann. N.Y. Acad. Sci.* 750:101 (1995); L. C. Waters et al., *Anal. Chem.* 70:158 (1998); J. Fritz et al., *Science* 288, 316 (2000)).

It should be noted that when the term "blood" is used herein, any part of the blood is intended. Accordingly, for determining molecular blood fingerprints, whole blood may be used directly where appropriate, or plasma or serum may be used.

Panels/Arrays for Detecting Organ-Specific Molecular Blood Fingerprints

The present invention also provides panels for detecting the organ-specific blood fingerprints at any given time in a subject. The term "subject" is intended to include any mammal or indeed any vertebrate that may be used as a model system for human disease. Examples of subjects include humans, monkeys, apes, dogs, cats, mice, rats, fish, zebra fish, birds, horses, pigs, cows, sheep, goats, chickens, ducks, donkeys, turkeys, peacocks, chinchillas, ferrets, gerbils, rabbits, guinea pigs, hamsters and transgenic species thereof. Further subjects contemplated herein include, but are not limited to, reptiles and amphibians, e.g., lizards, snakes, turtles, frogs, toads, salamanders, and newts. In one embodiment, the panel/array of the present invention comprises one detection reagent that specifically detects an organ-specific secreted protein. In another embodiment, the panel/arrays are comprised of a plurality of detection reagents that each specifically detects an organ-specific secreted protein, wherein the levels of organ-specific secreted proteins taken together form a unique pattern that defines the fingerprint. In certain embodiments, detection reagents can be bispecific such that the panel/array is comprised of a plurality of bispecific detection reagents that may specifically detect more than one organ-specific secreted protein. The term "specifically" is a term of art that would be readily understood by the skilled artisan to mean, in this context, that the protein of interest is detected by the particular detection reagent but other proteins are not detected in a statistically significant manner under the same conditions. Specificity can be determined using appropriate positive and negative controls and by routinely optimizing conditions.

The panels/arrays may be comprised of a solid phase surface having attached thereto a plurality of detection reagents each attached at a distinct location. As would be recognized by the skilled artisan, the number of detection reagents on a given panel would be determined from the number of organ-specific secreted proteins in the fingerprint to be measured. In one embodiment, the panel/array comprises one or more detection reagents. In a further embodiment, the panel/array comprises a plurality of detection reagents, wherein the plurality of detection reagents may be anywhere from about 2 to about 100, 150, 160, 170, 180, 190, 200, or more detection reagents each specific for an organ-specific secreted protein. In one embodiment, the panel/array comprises at least 2, 3, 4,

5, 6, 7, 8, 9, or 10 detection reagents each specific for one of the plurality of organ-specific secreted proteins that make up a given fingerprint. In another embodiment, the panel/array comprises at least 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 detection reagents each specific for one of the plurality of organ-specific secreted proteins that make up a given fingerprint. In a further embodiment, the panel/array comprises at least 21, 22, 23, 24, 25, 26, 27, 28, 29, or 30 detection reagents each specific for one of the plurality of organ-specific secreted proteins that make up a given fingerprint. In an additional embodiment, the panel/array comprises at least 31, 32, 33, 34, 35, 36, 37, 38, 39, or 40 detection reagents each specific for one of the plurality of organ-specific secreted proteins that make up a given fingerprint. In yet a further embodiment, the panel/array comprises at least 41, 42, 43, 44, 45, 46, 47, 48, 49, or 50 detection reagents each specific for one of the plurality of organ-specific secreted proteins that make up a given fingerprint. In an additional embodiment, the panel/array comprises at least 51, 52, 53, 54, 55, 56, 57, 58, 59, or 60 detection reagents each specific for one of the plurality of organ-specific secreted proteins that make up a given fingerprint. In one embodiment, the panel/array comprises at least 61, 62, 63, 64, 65, 66, 67, 68, 69, or 70 detection reagents each specific for one of the plurality of organ-specific secreted proteins that make up a given fingerprint. In one embodiment, the panel/array comprises at least 75, 80, 85, 90, 100, 150, 160, 170, 180, 190, 200, or more, detection reagents each specific for one of the plurality of organ-specific secreted proteins that make up a given fingerprint.

Further in this regard, the solid phase surface may be of any material, including, but not limited to, plastic, polycarbonate, polystyrene, polypropylene, polyethylene, glass, nitrocellulose, dextran, nylon, metal, silicon and carbon nanowires, nanoparticles that can be made of a variety of materials and photolithographic materials. In certain embodiments, the solid phase surface is a chip. In another embodiment, the solid phase surface may comprise microtiter plates, beads, membranes, microparticles, the interior surface of a reaction vessel such as a test tube or other reaction vessel. In other embodiments the peptides will be fractionated by one or more one-dimensional columns using size separations, ion exchange or hydrophobicity properties and, for example, deposited in a MALDI 96 or 384 well plate and then injected into an appropriate mass spectrometer.

In one embodiment, the panel/array is an addressable array. As such, the addressable array may comprise a plurality of distinct detection reagents, such as antibodies or aptamers, attached to precise locations on a solid phase surface, such as a plastic chip. The position of each distinct detection reagent on the surface is known and therefore "addressable". In one embodiment, the detection reagents are distinct antibodies that each have specific affinity for one of a plurality of organ-specific polypeptides.

In one embodiment, the detection reagents, such as antibodies, are covalently linked to the solid surface, such as a plastic chip, for example, through the Fc domains of antibodies. In another embodiment, antibodies are adsorbed onto the solid surface. In a further embodiment, the detection reagent, such as an antibody, is chemically conjugated to the solid surface. In a further embodiment, the detection reagents are attached to the solid surface via a linker. In certain embodiments, detection with multiple specific detection reagents is carried out in solution.

Methods of constructing protein arrays, including antibody arrays, are known in the art (see, e.g., U.S. Pat. No. 5,489,678; U.S. Pat. No. 5,252,743; Blawas and Reichert, 1998, *Biomaterials* 19:595-609; Firestone et al., 1996, *J. Amer. Chem. Soc.*

18, 9033-9041; Mooney et al., 1996, *Proc. Natl. Acad. Sci.* 93,12287-12291; Pirrung et al, 1996, *Bioconjugate Chem.* 7, 317-321; Gao et al, 1995, *Biosensors Bioelectron* 10, 317-328; Schena et al, 1995, *Science* 270, 467-470; Lom et al., 1993, *J. Neurosci. Methods*, 385-397; Pope et al., 1993, *Bioconjugate Chem.* 4, 116-171; Schramm et al., 1992, *Anal. Biochem.* 205, 47-56; Gombotz et al., 1991, *J Biomed. Mater. Res.* 25, 1547-1562; Alarie et al., 1990, *Analy. Chim. Acta* 229, 169-176; Owaku et al, 1993, *Sensors Actuators B*, 13-14, 723-724; Bhatia et al., 1989, *Analy. Biochem.* 178, 408-413; Lin et al., 1988, *IEEE Trans. Biomed. Engng.*, 35(6), 466-471).

In one embodiment, the detection reagents, such as antibodies, are arrayed on a chip comprised of electronically activated copolymers of a conductive polymer and the detection reagent. Such arrays are known in the art (see e.g., U.S. Pat. No. 5,837,859 issued Nov. 17, 1998; PCT publication WO 94/22889 dated Oct. 13, 1994). The arrayed pattern may be computer generated and stored. The chips may be prepared in advance and stored appropriately. The antibody array chips can be regenerated and used repeatedly.

Using the methods described herein, a vast array of organ-specific molecular blood fingerprints can be defined for any of a variety of drugs as described further herein. As such, the present invention further provides information databases comprising data that make up molecular blood fingerprints as described herein. As such, the databases may comprise the defined differential expression levels as determined using any of a variety of methods such as those described herein, of each of the plurality of organ-specific secreted proteins that make up a given fingerprint in any of a variety of settings (e.g., normal or drug-associated fingerprints).

Methods of Use

The present invention provides methods for identifying organ-specific secreted proteins and methods for identifying organ-specific molecular blood fingerprints. The present invention further provides panels/arrays of detection reagents for detecting such fingerprints. The present invention also provides defined organ-specific molecular blood fingerprints for normal and disease settings and for fingerprints associated with/resulting from a particular drug or combination of drugs. As such, the present invention provides for methods for identifying and monitoring drug effects in any of a variety of settings. Further, the present invention provides methods for following responses to therapy in a variety of disease settings such that any adverse or other drug side effects can be monitored. The present invention also provides methods of detecting disease, stratifying disease, monitoring the progression of disease, and monitoring responses to therapy such as described in U.S. Provisional Application Nos. 60/647685 and 60/683071 filed Jan. 27, 2005 and May 20, 2005, respectively, and Copending U.S. application Ser. No. 11,342,366 entitled *Methods for Identifying and Using Organ-Specific Proteins in Blood*, filed concurrently on Jan. 27, 2006.

The present invention can be used as a standard screening test. In this regard, one or more of the detection panels described herein can be run on an individual taking a particular drug and any statistically significant deviation from a normal organ-specific molecular blood fingerprint would indicate that drug-related perturbation was present. Thus, the present invention provides a standard or "normal" blood fingerprint for any given organ. In certain embodiments, a normal blood fingerprint is determined by measuring the normal range of levels of the individual protein members of a fingerprint. Any deviation therefrom or perturbation of the normal fingerprint that is outside the standard deviation (normal

range) has utility in determining drug side effects (see also U.S. Patent Application No. 0020095259). As would be recognized by the skilled artisan, the significance of any deviation in the levels of (e.g., a significantly altered level of one or more of) the individual protein members of a fingerprint can be determined using statistical methods known in the art and described herein. In certain embodiments, a normal standard can be generated from a blood sample taken from the individual prior to administration of the drug such that comparisons can be made thereto.

Further, the present invention provides methods for determining and evaluating not only the absolute levels of the changes in the proteins constituting individual fingerprints, but also for evaluating all the protein changes (e.g. N changed proteins) and comparing them against one another to generate an N-dimensional shape space that provides more powerful correlation with the stratifications of drug-induced alterations described above (see e.g., U.S. Patent Application No. 20020095259).

In a further embodiment, the present invention can be used to determine the risk of having one or more side effects from a drug or combination thereof. A statistically significant alteration (e.g., increase or decrease) in the levels of one or more members of a particular molecular blood fingerprint may signify a risk of developing a one or more side effects from a drug or combination thereof.

The organ-specific molecular blood fingerprints of the present invention can be used to detect side effects, either positive or negative (or lack thereof) from any of a variety of drugs. As would be recognized by the skilled artisan, the present invention can be used to detect the side effects (or lack thereof) from virtually any drug. In this regard, any drug either currently under development or already approved and on the market is contemplated in the context of this invention. In particular, the present invention provides methods for monitoring the organ-specific molecular blood fingerprint of organs/tissues/cells that fall outside the expected therapeutic targets (e.g., monitoring off-target effects of drugs). For example, the liver and kidney are organs that often reflect the side effects of drugs. Further, as demonstrated by the side-effects of COX-2 inhibitors, drugs can have off-target effects on the cardiovascular system, or any other cell/organ/tissue/system as described herein. Thus, the present invention also provides methods for monitoring non-target organs for any drug, including drugs under development and drugs currently on the market for which subtle side-effects may not have been detected.

In a further embodiment, the present invention can be used to determine side effects of combinations of drugs on any organ. In a further embodiment, the organ-specific molecular blood fingerprints can be monitored in subjects taking very low (non-toxic) doses of drugs to determine whether subtle side effects are occurring.

Thus, the organ-specific molecular blood fingerprints of the present invention can be used to detect direct effects or side effects of any drug on the heart, kidney, ureter, bladder, urethra, liver, prostate, heart, blood vessels, bone marrow, skeletal muscle, smooth muscle, various specific regions of the brain (including, but not limited to the amygdala, caudate nucleus, cerebellum, corpus callosum, fetal, hypothalamus, thalamus), spinal cord, peripheral nerves, retina, nose, trachea, lungs, mouth, salivary gland, esophagus, stomach, small intestines, large intestines, hypothalamus, pituitary, thyroid, pancreas, adrenal glands, ovaries, oviducts, uterus, placenta, vagina, mammary glands, testes, seminal vesicles, penis, lymph nodes, thymus, and spleen. The present invention can be used to detect drug side effects on the cardiovas-

cular system, neurological system, metabolic system, respiratory system, the immune system, etc. As would be recognized by the skilled artisan, the present invention can be used to detect any side effects wherein the side effects cause perturbation in organ-specific secreted proteins. In this regard, a side effect may be an adverse effect or may be a positive effect. Accordingly, the present invention can be used to identify potential new indications for a particular drug.

In an additional embodiment, the present invention can be used to determine distinct, normal organ-specific molecular blood fingerprints, such as in different populations of people. In this regard, there may be differences in normal organ-specific molecular blood fingerprints between populations of individuals that permit the stratification of patients into classes of individuals who would respond positively to a particular drug and those who would not. Thus, the present invention provides the ability to determine those individuals who may have adverse reactions to drugs. Additionally, in certain embodiments, the nature of the drug-induced changes in one or more organ-specific molecular blood fingerprints can be used to predict which patients might effectively respond to the drug. Thus, the organ-specific molecular blood fingerprints of the present invention provide the ability to stratify patients with regard to drug response and the ability to assess the toxicity of drugs.

Once a side effect is detected by identifying a perturbation in an organ-specific molecular blood fingerprint, the effect can be further mapped using systems approaches by mapping the gene and protein networks perturbed (see Example 1). In certain embodiments, a single organ-specific secreted protein may be perturbed (as indicated by detection of an increase or decrease in the level of the protein in the blood). In further embodiments, more than one organ-specific secreted protein may be perturbed.

To monitor the monitor responses to therapy or responses to any drug, one or more organ-specific molecular blood fingerprints are detected/measured as described herein using any of the methods as described herein at one time point and detected/measured again at subsequent time points, thereby monitoring responses to therapy or to any drug.

Organ-specific molecular blood fingerprints can also be defined and/or perturbations thereof tested in any of a variety of animal models. Animals that can be used in this context, include, for example, mice, rats, rabbits, pigs, monkeys, apes, zebra fish, etc, and transgenic species thereof.

Business Methods

A further embodiment of the present invention comprises a business method of identifying a particular drug side effect in a subject taking a drug that comprises detecting an organ-specific molecular blood fingerprint as described herein.

Thus, the present invention contemplates methods for (a) manufacturing one or more of the detection reagents, panels, arrays, (b) providing diagnostic services for determining organ-specific blood fingerprints, and identifying particular drug side effects (c) providing manufacturers of genomics devices the use of the detection reagents, panels, arrays, blood fingerprints or transcriptomes described herein to develop diagnostic devices, where the genomics device includes any device that may be used to define differences in a blood sample between the normal and disturbed state resulting from one or more drug side effects (d) providing manufacturers of proteomics devices the use of the detection reagents, panels, arrays, blood fingerprints or transcriptomes described herein to develop diagnostic devices, where the proteomics device includes any device that may be used to define differences in a blood sample between the normal and disturbed state result-

ing from a drug side effect and (e) providing manufacturers of imaging devices the use of the detection reagents, panels, arrays, blood fingerprints or transcriptomes described herein to develop diagnostic devices, where the proteomics device includes any device that may be used to define differences in a blood sample between the normal and disturbed state resulting from one or more drug side effects (f) providing manufacturers of molecular imaging devices the use of the detection reagents, panels, arrays, blood fingerprints or transcriptomes described herein to develop diagnostic devices, where the proteomics device includes any device that may be used to define differences in a blood sample between the normal and disturbed state resulting from one or more drug side effects and g) marketing to healthcare providers the benefits of using the detection reagents, panels, arrays, and diagnostic services of the present invention to enhance diagnostic capabilities and thus, to better treat patients.

Another aspect of the invention relates to a method for conducting a business, which includes: (a) manufacturing one or more of the detection reagents, panels, arrays, (b) providing services for determining organ-specific molecular blood fingerprints and (c) marketing to healthcare providers the benefits of using the detection reagents, panels, arrays, and services of the present invention to enhance capabilities to identify drug side effects and thus, to better treat patients.

Another aspect of the invention relates to a method for conducting a business, comprising: (a) providing a distribution network for selling the detection reagents, panels, arrays, diagnostic services, and access to organ-specific molecular blood fingerprint databases (b) providing instruction material to physicians or other skilled artisans for using the detection reagents, panels, arrays, and organ-specific molecular blood fingerprint databases to improve the ability to identify drug side effects for patients.

Yet another aspect of the invention relates to a method for conducting a business, comprising: (a) identifying organ-specific secreted proteins in the blood sera, etc. (b) determining the organ-specific molecular fingerprints as described herein and (c) providing a distribution network for selling access to the database of organ-specific molecular fingerprints identified in step (b).

For instance, the subject business method can include an additional step of providing a sales group for marketing the database, or panels, or arrays, to healthcare providers.

Another aspect of the invention relates to a method for conducting a business, comprising: (a) determining one or more organ-specific molecular blood fingerprints and (b) licensing, to a third party, the rights for further development and sale of panels, arrays, and information databases related to the organ-specific molecular blood fingerprints of (a).

The business methods of the present application relate to the commercial and other uses, of the methodologies, panels, arrays, organ-specific secreted proteins, organ-specific molecular blood fingerprints, and databases comprising identified fingerprints of the present invention. In one aspect, the business method includes the marketing, sale, or licensing of the present invention in the context of providing consumers, i.e., patients, medical practitioners, medical service providers, and pharmaceutical distributors and manufacturers, with all aspects of the invention described herein, (e.g., the methods for identifying organ-specific secreted proteins, detection reagents for such proteins, molecular blood fingerprints, etc., as provided by the present invention).

In a particular embodiment of the present invention, a business method relating to providing information related to molecular blood fingerprints (e.g., levels of the plurality of organ-specific secreted proteins that make up a given finger-

print), method for determining fingerprints and sale of panels for determining such molecular blood fingerprints. In a specific embodiment, that method may be implemented through the computer systems of the present invention. For example, a user (e.g. a health practitioner such as a physician or a diagnostic laboratory technician) may access the computer systems of the present invention via a computer terminal and through the Internet or other means. The connection between the user and the computer system is preferably secure.

In practice, the user may input, for example, information relating to a patient such as the patient's disease state and/or drugs that the patient is taking, e.g., levels determined for the proteins that make up a given molecular blood fingerprint using a panel or array of the present invention. The computer system may then, through the use of the resident computer programs, provide a diagnosis or determination of drug side effects that fits with the input information by matching the fingerprint parameters (e.g., levels of the proteins present in the blood as detected using a particular panel or array of the present invention) with a database of fingerprints.

A computer system in accordance with a preferred embodiment of the present invention may be, for example, an enhanced IBM AS/400 mid-range computer system. However, those skilled in the art will appreciate that the methods and apparatus of the present invention apply equally to any computer system, regardless of whether the computer system is a complicated multi-user computing apparatus or a single user device such as a personal computer or workstation. Computer systems suitably comprise a processor, main memory, a memory controller, an auxiliary storage interface, and a terminal interface, all of which are interconnected via a system bus. Note that various modifications, additions, or deletions may be made to the computer system within the scope of the present invention such as the addition of cache memory or other peripheral devices.

The processor performs computation and control functions of the computer system, and comprises a suitable central processing unit (CPU). The processor may comprise a single integrated circuit, such as a microprocessor, or may comprise any suitable number of integrated circuit devices and/or circuit boards working in cooperation to accomplish the functions of a processor.

In a preferred embodiment, the auxiliary storage interface allows the computer system to store and retrieve information from auxiliary storage devices, such as magnetic disk (e.g., hard disks or floppy diskettes) or optical storage devices (e.g., CD-ROM). One suitable storage device is a direct access storage device (DASD). A DASD may be a floppy disk drive that may read programs and data from a floppy disk. It is important to note that while the present invention has been (and will continue to be) described in the context of a fully functional computer system, those skilled in the art will appreciate that the mechanisms of the present invention are capable of being distributed as a program product in a variety of forms, and that the present invention applies equally regardless of the particular type of signal bearing media to actually carry out the distribution. Examples of signal bearing media include: recordable type media such as floppy disks and CD ROMS, and transmission type media such as digital and analog communication links, including wireless communication links.

The computer systems of the present invention may also comprise a memory controller, through use of a separate processor, which is responsible for moving requested information from the main memory and/or through the auxiliary storage interface to the main processor. While for the purposes of explanation, the memory controller is described as a

separate entity, those skilled in the art understand that, in practice, portions of the function provided by the memory controller may actually reside in the circuitry associated with the main processor, main memory, and/or the auxiliary storage interface.

Furthermore, the computer systems of the present invention may comprise a terminal interface that allows system administrators and computer programmers to communicate with the computer system, normally through programmable workstations. It should be understood that the present invention applies equally to computer systems having multiple processors and multiple system buses. Similarly, although the system bus of the preferred embodiment is a typical hard-wired, multidrop bus, any connection means that supports bidirectional communication in a computer-related environment could be used.

The main memory of the computer systems of the present invention suitably contains one or more computer programs relating to the organ-specific molecular blood fingerprints and an operating system. Computer program is used in its broadest sense, and includes any and all forms of computer programs, including source code, intermediate code, machine code, and any other representation of a computer program. The term "memory" as used herein refers to any storage location in the virtual memory space of the system. It should be understood that portions of the computer program and operating system may be loaded into an instruction cache for the main processor to execute, while other files may well be stored on magnetic or optical disk storage devices. In addition, it is to be understood that the main memory may comprise disparate memory locations.

All of the U.S. patents, U.S. patent application publications, U.S. patent applications, foreign patents, foreign patent applications and non-patent publications referred to in this specification and/or listed in the Application Data Sheet, are incorporated herein by reference, in their entirety. Moreover, all numerical ranges utilized herein explicitly include all integer values within the range and selection of specific numerical values within the range is contemplated depending on the particular use. Further, the following examples are offered by way of illustration, and not by way of limitation.

EXAMPLES

Example 1

Evidence For the Presence of Disease-Perturbed Networks in Prostate Cancer Cells By Genomic and Proteomic Analysis: A Systems Approach to Disease

The following example demonstrates the presence of disease-perturbed networks in prostate. This provides a model for studying perturbation of organ-specific molecular blood fingerprints. The same principles apply in the setting of determining perturbations that result from drugs.

Prostate cancer is the most common nondermatological cancer in the United States (Greenlee, R. T., et al., *CA Cancer J Clin*, 50: 7-33., 2000). Initially, its growth is androgen-dependent (AD); early-stage therapies, including chemical and surgical castration, kill cancerous cells by androgen deprivation. Although such therapies produce tumor regression, they eventually fail because most prostate carcinomas become androgen-independent (AI) (Isaacs, J. T. *Urol Clin North Am*, 26: 263-273., 1999). To improve the efficacy of prostate cancer therapy, it is necessary to understand the molecular mechanisms underlying the transition from androgen dependence to androgen independence.

The transition from AD to AI status likely results from multiple processes, including activation of oncogenes, inactivation of tumor suppressor genes, and changes in key components of signal transduction pathways and gene regulatory networks. Systems approaches to biology and disease are predicated on the identification of the elements of the systems, the delineation of their interactions and their changes in distinct disease states. Biological information is of two types: the digital information of the genome (e.g. genes and cis-control elements) and environmental cues. Proteins rarely act in isolation; rather, they form parts of molecular machines or participate in network interactions mediating cellular functions such as signal transduction and developmental or physiological response patterns. Gene regulatory networks, whose architecture and linkages are established by cis-control elements, integrate information from signal transduction networks and output it to developmental or physiological batteries or networks of effector proteins. Normal protein and gene regulatory networks may be perturbed by disease—through genetic and/or environmental perturbations and understanding these differences lies at the heart of systems approaches to disease. Disease-perturbed networks initiate altered responses that bring about pathologic phenotypes such as the invasiveness of cancer cells.

To map network perturbations in cancer initiation and progression, changes in expression levels of virtually all transcripts must be measured. Certain low-abundance transcripts, such as those encoding transcription factors and signal transducers, wield significant regulatory influences in spite of the fact they may be present in the cell at very low copy numbers. Differential display (Bussemakers, M. J., et al., *Cancer Res*, 59: 5975-5979, 1999) or cDNA microarrays (Vaarala, M. H., et al., *Lab Invest*, 80: 1259-1268, 2000; Chang, G. T., et al., *Cancer Res*, 57: 4075-4081, 1997) have been used to profile changes in gene expression during the AD to AI transition; however, those technologies can identify only a limited number of more abundant mRNAs, and they miss many low-abundance mRNAs due to their low detection sensitivities. Massively parallel signature sequencing (MPSS), allows 20-nucleotide signature sequences to be determined in parallel for more than 1,000,000 DNA sequences (Brenner, et al., 2000, supra). MPSS technology allows identification and cataloging of almost all mRNAs that are changed between two cell states, even those with one or a few transcripts per cell, or between different organs or tissues. Differentially expressed genes thus identified can be mapped onto cellular networks to provide a systemic understanding of changes in cellular state.

Although transcriptome (mRNA levels) differences are easier to study than proteome (protein levels) differences and provide extremely valuable information, cellular functions are usually performed by proteins. RNA expression profiling studies do not address how the encoded proteins function biologically, and transcript abundance levels do not always correlate with protein abundance levels (Chen, G., et al., *Mol Cell Proteomics*, 1: 304-313, 2002). Therefore, the mRNA expression profiling described herein was complemented with a more limited protein profiling by using isotope-coded affinity tags (ICAT) coupled with tandem mass spectrometry (MS/MS) (Gygi, S. P., et al., *Nat Biotechnol*, 17: 994-999., 1999).

The LNCaP cell line is a widely used androgen-sensitive model for early-stage prostate cancer from which androgen-independent sublines have been generated (Vaarala, M. H., et al., 2000, supra; Chang, G. T., et al., 1997, supra; Patel, B. J., et al., *J Urol*, 164: 1420-1425., 2000). The cells of one such variant, CL-1, in contrast to their LNCaP progenitors, are

highly tumorigenic, and exhibit invasive and metastatic characteristics in intact and castrated mice (Patel, G. J., et al., 2000, supra; Tso, C. L., et al., *Cancer J Sci Am*, 6: 220-233., 2000). Thus CL-1 cells model late-stage prostate cancer. MPSS and ICAT data extracted from these model cell lines can be validated by real-time RT-PCR or western blot analysis in more relevant biological models (tumor xenografts) and in tumor biopsies.

An MPSS analysis of about 5 million signatures was conducted for the androgen-dependent LNCaP cell line and its androgen-independent derivative CL1. The resulting database offers the first comprehensive view of the digital transcriptomes of prostate cancer cells and allows exploration of the cellular pathways perturbed during the transition from AD to AI growth. Additionally, protein expression profiles between LNCaP and CL1 cells were compared using ICAT/MS/MS technology. Further, computational analysis was used to identify those proteins that are secreted. Once such protein was further investigated and shown to be a diagnostic marker for prostate cancer used either alone, or in combination with the known PSA prostate cancer marker.

MPSS analysis: LNCaP and CL1 cells were grown using methods known in the art, for example, as described by Tso et al. 2000, supra). RNAs were isolated using Trizol (Life Technologies) according to the manufacturer's protocols (see, e.g., as described by Nelson et al. *Proc Natl Acad Sci USA*, 99: 11890-11895, 2002). MPSS cDNA libraries were constructed, individual cDNA sequences were amplified and attached to individual beads and sequenced as described by Brenner, et al., 2000, supra. The resulting signatures, generally 20 bases in length, were annotated using the then most recently annotated human genome sequence (human genome release hg16, released in November, 2003) and the human Unigene (Unigene build #184) according to a previously published method (Meyers, B. C., et al., *Genome Res*, 14: 1641-1653, 2004). Only 100% matches between an MPSS signature and a genome signature were considered. Those signatures that expressed at less than 3 tpm in both LNCaP and CL1 libraries were also excluded, as they might not be reliably detected (this represents less than one transcript per cell) (Jongeneel, C. V., et al., *Proc Natl Acad Sci USA*, 2003). Additionally, cDNA signatures were classified by their positions relative to polyadenylation signals and poly (A) tails and by their orientation relative to the 5'→3' orientation of source mRNA. The Z-test (Man, M. Z., et al., *Bioinformatics*, 16: 953-959, 2000) was used to calculate P values for comparison of gene expression levels between the cell lines.

Isotope-Coded Affinity Tag (ICAT) analysis: ICAT reagents were purchased from Applied Biosystems Inc. Fractionation of cells into cytosolic, microsomal and nuclear fractions, as well as ICAT labeling, MS/MS, and data analyses were performed as described by Han et al. *Nat Biotechnol*, 19: 946-951, 2001. In addition, probability score analysis (Keller, A., et al., *Anal Chem*, 74: 5383-5392, 2002) and ASAPRatio (Automated Statistical Analysis on Protein Ratio) (Li, X. J., et al., *Anal Chem*, 75: 6648-6657, 2003) were used to assess the quality of MS spectra and to calculate protein ratios from multiple peptide ratios. (Briefly, and as described at <http://regis.systemsbio.org/software>, Automated Statistical Analysis on Protein Ratio (ASAPRatio) accurately calculates the relative abundances of proteins and the corresponding confidence intervals from ICAT-type ESI-LC/MS data. The software first uses a Savitzky-Golay smoothing filter to reconstruct LC spectra of a peptide and its partner in a single charge state, subtracts background noise from each spectrum, and calculates light:heavy ratio of the peptide in that charge state. The ratios of the same peptide in different charge states

are averaged and weighted by the corresponding spectrum intensity to obtain the peptide light:heavy ratio and its error. Subsequently, all unique peptides identified for a given protein are collected, their ratios and errors calculated, outliers are checked for using Dixon's tests, and the relative abundance and confidence interval for the protein are calculated by applying statistics for weighed samples. The software quickly generates a list of interesting proteins based on their relative abundance. A byproduct of the software is to identify outlier peptides which may be misidentified or, more interestingly, post-translationally modified.) To compare protein and mRNA expression levels, the Unigene numbers of the differentially expressed proteins were used to find MPSS signatures and their expression levels in transcripts per million (tpm). If one Unigene had more than one MPSS signature, likely due to alternative terminations, the average tpm of all signatures was taken.

Real-time RT-PCR: All primers were designed with the PRIMER3 program (<http://www.genome.gov/cgi-bin/primer/primer3.cgi>) and BLAST-searched against the human cDNA and EST database for uniqueness. Real-time PCR was performed on an ABI 7700 machine (PE Biosystems) and the SYBR Green dye (Molecular Probe Inc.) was used as a reporter. PCR conditions were designed to give bands of the expected size with minimal primer dimer bands.

Identification of perturbed networks: Genes in the 328 Biocarta and Kyoto Encyclopedia of Genes and Genomes (KEGG) pathways or networks (<http://www.ncbi.nlm.nih.gov/Pathways/>) were downloaded and compared with the MPSS data, using Unigene IDs as identifiers. If a Unigene ID or an E.C. number corresponded to multiple signatures, potentially due to multiple alternatively terminated isoforms, the tpm counts of the isoforms were combined and then subjected to the Z-test (Man, M. Z., et al., 2000, supra). Genes with P values of 0.001 or less were considered to be significantly differentially expressed. The following criteria were used to identify perturbed networks: a perturbed network must have more than 3 genes represented our differentially expressed gene list ($p < 0.001$) and at least 50% of those genes must be up regulated, it was considered an up-regulated pathway (vice versa for the down-regulated pathways).

Display of KEGG networks by Cytoscape: Cytoscape software was used (www.cytoscape.org) (Shannon, P., et al., *Genome Res*, 13: 2498-2504, 2003), to map the data onto the web of intracellular molecular interactions. We imported metabolic network maps and related information such as enzymes, substrates, and reactions from the recently developed KEGG (<http://www.genome.ad.jp/>) API 2.0 web server into the Cytoscape program. Expression data were thus automatically mapped to the KEGG and Biocarta pathways/networks and visualized by Cytoscape.

MPSS analyses of the androgen-dependent LNCaP cell line and its androgen-independent variant CL1: Using MPSS technology, 2.22 million signature sequences were sequenced for LNCaP cells and 2.96 million for CL1 cells.

A total of 19,595 unique transcript signatures expressed at levels >3 tpm in at least one of the samples were identified. The signatures were classified into three major categories: 1093 signatures matched repeat sequences; 15,541 signatures matched unique cDNAs or ESTs, and 2961 signatures had no matches to any cDNA or EST sequences (but did match genomic sequences). The last category included sequences falling into one of three different categories: signatures representing new transcripts yet to be defined, signatures repre-

senting polymorphisms in cDNA sequences (a match of an MPSS sequence to cDNA or EST sequences requires 100% sequence identity), or errors in the MPSS reads. Transcript tags with matches to a cDNA or EST sequence were further classified based on the signatures' relative orientation to transcription direction and their position relative to a polyadenylation site and/or poly(A) tail. A searchable MySQL database (www dot mysql dot corn) was also built containing the expression levels (tpm), the genomic locations of the MPSS sequences, the cDNAs or EST matches, and the classification of each signature.

The first analysis was restricted to those MPSS signatures corresponding to cDNAs with poly(A) tails and/or polyadenylation sites, so that corresponding genes could be conclusively identified. The Z-test was used to compare differential gene expression between LNCaP cells and CL1 cells (Mann, et al., 2000, supra). Using very stringent P values (less than 0.001), 2088 MPSS signatures were identified (corresponding to 1987 unique genes, as some genes have two or more MPSS signatures, due to alternative usages of polyadenylation sites) with significant differential expression. Of these, 1011 signatures (965 genes) were overexpressed in CL1 cells, and 1077 signatures (1022 genes) were overexpressed in LNCaP cells. The significance score of Z-test was dependent on the expression level. If a cut off P value of less than 0.001 was taken in the dataset, the expression level in tpm changed from 0 to 26 tpm for the most lowly expressed transcript (>26 fold); and changed from 7591 and 11206 tpm for the most highly expressed transcript (1.48 fold).

The expression levels of nine randomly chosen genes were identified using the MPSS and quantitative real-time RT-PCR techniques and showed that both RNA data sets were concordant. The MPSS expression profiling data were consistent with the available published data. For example, using RT-PCR, Patel et al. (Patel, B. J., et al., *J Urol*, 164: 1420-1425, 2000) showed that CL1 tumors express barely detectable prostate-specific antigen (PSA) and androgen receptor (AR) mRNAs as compared with LNCaP cells. The present MPSS results indicated that LNCaP cells expressed 584 tpm of androgen receptor (AR) and 841 tpm of PSA; CL1 cells did not express either AR or PSA (0 tpm in both cases). Freedland et al. found that CD10 expression was lost in CL1 cells compared with LNCaP cells (Freedland, S. J., et al., *Prostate*, 55: 71-80, 2003); the present study found that CD10 was expressed at 0 tpm in CL1 cells but at 56 tpm in LNCaP cells. Using cDNA microarrays, Vaarala et al. (Vaarala, M. H., et al., *Lab Invest*, 80: 1259-1268, 2000) compared LNCaP cells and another androgen-independent variant, non-PSA-producing LNCaP line, which is similar to CL1, and identified a total of 56 differentially expressed genes. We found completely concordant expression changes in these 56 genes between LNCaP and CL1 (in contrast to 1987 found by MPSS), and between LNCaP and non-PSA-producing LNCaP cells. This underscores the striking differences in sensitivity between the MPSS and cDNA microarray techniques.

CL1 cells do not express AR and thus lack the AR-mediated response program. To distinguish androgen response from other programs contributing to prostate cancer progression, the list of genes differentially expressed between LNCaP and CL1 cells were compared with a complementary list derived from MPSS analysis of LNCaP cells grown in the presence or absence of androgens (LNCaP R+/R-). From the 1987 differentially expressed gene between LNCaP and CL1, 525 genes were identified that were also differentially expressed in the LNCaP R+/R- dataset. Differential expression of these genes between LNCaP and CL1 cells probably

reflects the fact that LNCaP cells express AR but CL1 does not, and the fact that normal medium contains some androgen. The remaining 1462 differentially expressed genes were not directly related to cellular AR status.

To compare the sensitivity of the MPSS and cDNA microarray procedures, cDNA microarrays containing 40,000 human cDNAs were hybridized to the same LNCaP and CL1 RNAs that were used for MPSS. Three replicate array hybridizations were performed. MPSS signatures and array clone IDs were mapped to Unigene IDs for data extraction and comparisons. The results showed that only those genes expressed at >40 tpm by MPSS could be reliably detected as changing levels by cDNA microarray hybridizations [judged by an expression level twice the standard deviation of the background, a standard cutoff value for microarray data analysis]. This observation is consistent with the 33-60 tpm sensitivity of microarrays estimated from the experiment performed by Hill et al. *Science*, 290: 809-812, 2000, in which known concentrations of synthetic transcripts were added. In LNCaP and CL1 cells, about 68.75% (13,471 of 19,595) of MPSS signatures (>3 tpm) were expressed at a level below 40 tpm; changes in the levels of these genes will be missed by microarray methods. Many attempts have been made to increase the sensitivity of DNA array technology (Han, M., et al., *Nat Biotechnol*, 19: 631-635, 2001; Bao, P., et al., *Anal Chem*, 74: 1792-1797, 2002.), however, the present study has not compared these new improvements against MPSS but it is clear that there will still be significant differences in the levels of change that can be detected.

SAGE (serial analysis of gene expression) (Velculescu, V. E., et al., *Trends Genet*, 16: 423-425., 2000) is another technology for gene expression profiling; like MPSS, it is digital and can generate a large number of signature sequences. However, MPSS, which can sequence ~1 million signatures per sample, can achieve a much deeper coverage than SAGE (typical ~10,000-100,000 signatures sequenced/sample) at reasonable cost. The MPSS data on LNCaP cells was compared against publicly available SAGE data on LNCaP cells (NCBI SAGE database) through common Unigene IDs. The SAGE library GSM724 (total SAGE tags sequenced: 22,721) (Lal, A., et al., *Cancer Res*, 59: 5403-5407, 1999) was derived from LNCaP cells with an inactivated PTEN gene; it is the SAGE library most similar to the LNCaP cells. Only 400 (about 20%) of the 1987 significantly differentially expressed genes (P<0.001) had any SAGE tag entry in GSM724. These data illustrate the importance of deep sequence coverage in identifying state changes in transcripts expressed at low abundance levels.

Functional classifications of genes differentially expressed between LNCaP and CL1 cells: Examination of the GO (Gene Ontology) classification of the 1987 genes revealed that multiple cellular processes change during the transition from LNCaP cells to CL1 cells. The most interesting groups, categorized by function, are shown in Table 1.

Nineteen differentially expressed proteins are related to apoptosis. Twelve of these are up regulated in CL1 cells, including the apoptosis inhibitors Tax1 (human T-cell leukemia virus type I) binding protein 1 (TAX1BP1) and CASP8 and FADD-like apoptosis regulator. Seven are down regulated in CL1, including programmed cell death 8 and 5 (apoptosis-inducing factors), and BCL2-like 13 (an apoptosis facilitator). Since CL1 cells have increased expression of apoptosis inhibitors and decreased expression of apoptosis inducers, net inhibition of apoptosis may contribute to their greater tumorigenicity.

TABLE 1

EXAMPLES OF DIFFERENTIALLY EXPRESSED GENES AND THEIR FUNCTIONAL CLASSIFICATIONS					
Signatures (tpm) (tpm)	LNCaP Description	CL1 Gen- Bank IDNOS:		SEQ ID	
<u>Apoptosis related</u>					
GATCAATGTGTGGCCT (SEQ ID NO: 3)	0	3609	lectin, galactoside- BC001693 binding, soluble, 1 (galectin 1),	1574-1575	
GATCATAATGTTAACTA (SEQ ID NO: 4)	0	14	pleiomorphic adenoma gene-like 1 (PLAGL1)	NM_002656	1576-1577
GATCATCCAGAGGAGCT (SEQ ID NO: 5)	0	16	caspase 7, apoptosis-related cysteine protease	U40281	1578-1579
GATCGCGGTATTAATC (SEQ ID NO: 6)	0	15	tumor necrosis factor receptor superfamily, member 12	U75380	1580-1581
GATCTCCTGCCATCAG (SEQ ID NO: 7)	0	24	interleukin 1, beta	M15330	1582-1583
GATCCCCTTCAAGGACA (SEQ ID NO: 8)	1	19	nudix (nucleoside diphosphate linked moiety X)-type motif 1	NM_006024	1584-1585
GATCATGCCATCACCA (SEQ ID NO: 9)	51	278	EST, Highly similar to AL832733 CUL2_HUMAN CULLIN HOMOLOG 2	1586	
GATCTGAAAATTCTTGG (SEQ ID NO: 10)	16	56	CASP8 and FADD-like apoptosis regulator	U97075	1587-1588
GATCCACCTTGGCCTCC (SEQ ID NO: 11)	49	149	tumor necrosis factor receptor superfamily, member 10b	NM_003842	1589-1590
GATCATGAATGACTGAC (SEQ ID NO: 12)	118	257	cytochrome c	BC009582	1591-1592
GATCAAGTCCTTTGTGA (SEQ ID NO: 13)	299	102	programmed cell death 8 (apoptosis- inducing factor)	H20713	1593
GATCACAAAACCTGAT (SEQ ID NO: 14)	72	24	BCL2-like 13 (apoptosis facilitator)	BM904887	1594
GATCAATCTGAACTATC (SEQ ID NO: 15)	563	146	apoptosis related protein APR-3 (APR-3)	NM_016085	1595-1596
GATCCCTCTGTACAGGC (SEQ ID NO: 16)	83	13	unc-13-like (C. elegans) (UNC13), mRNA.	NM_006377	1597-1598
GATCTGGTTGAAAATTG (SEQ ID NO: 17)	1006	49	CED-6 protein (CED-6), mRNA.	NM_016315	1599-1600
GATCTCCCATGTTGGCT (SEQ ID NO: 18)	86	4	CASP2 and RIPK1 domain containing adaptor with death domain	BC017042	1601-1602

TABLE 1-continued

EXAMPLES OF DIFFERENTIALLY EXPRESSED GENES AND THEIR FUNCTIONAL CLASSIFICATIONS					
Signatures (tpm) (tpm)	LNCaP Description	CL1 Gen- Bank IDNOS:		SEQ ID	
GATCAGAAAATCCCTCT (SEQ ID NO: 19)	27	1	DEAD/H (Asp- Glu-Ala-Asp/His) box polypeptide 20, 103kDa	BC011556	1603-1604
GATCAAGGATGAAAGCT (SEQ ID NO: 20)	50	3	programmed cell death 2	D20426	1605
GATCTGATTATTTACTT (SEQ ID NO: 21)	1227	321	programmed cell death 5	NM_004708	1606-1607
GATCAAGTCCTTTGTGA (SEQ ID NO: 22)	299	102	programmed cell death 8 (apoptosis- inducing factor)	NM_004208	1608-1609
<u>Cyclins</u>					
GATCCTGTCAAATAGT (SEQ ID NO: 23)	2	47	MCT-1 protein (MCT-1), mRNA.	NM_014060	1610-1611
GATCATTATATCATTGG (SEQ ID NO: 24)	3	39	cyclin-dependent kinase inhibitor 2B (CDKN2B)	NM_078487	1612-1613
GATCATCAGTCACCGAA (SEQ ID NO: 25)	38	396	cyclin-dependent kinase inhibitor 2A (p16)	BM054921	1614
GATCGGGGGCGTAGCAT (SEQ ID NO: 26)	5	43	cyclin D1	NM_053056	1615-1616
GATCTACTCTGTATGGG (SEQ ID NO: 27)	40	144	cyclin fold protein 1	BG119256	1617
GATCAGCACTCTACCAC (SEQ ID NO: 28)	530	258	cyclin B1	BM973693	1618
GATCTGGTGTAGTATAT (SEQ ID NO: 29)	210	77	cyclin G2	BM984551	1619
GATCAGTACACAATGAA (SEQ ID NO: 30)	642	224	cyclin G 1,	BC000196	1620-1621
GATCTCAGTTCTGCGTT (SEQ ID NO: 31)	918	308	CDK2-associated protein 1 (CDK2AP1), mRNA.	NM_004642	1622-1623
GATCCTGAGCTCCCTTT (SEQ ID NO: 32)	2490	650	cyclin I,	BC000420	1624-1625
GATCATGCAGTGACATA (SEQ ID NO: 33)	15	1	KIAA1028 protein	AL122055	1626-1627
GATCTGTATGTGATTGG (SEQ ID NO: 34)	28	1	cyclin M3	AA489077	1628
<u>Kallikreins</u>					
GATCCACACTGAGAGAG (SEQ ID NO: 35)	841	0	KLK3	AA523902	1629
GATCCAGAAATAAAGTC (SEQ ID NO: 36)	385	0	KLK4	AA595489	1630
GATCCTCCTATGTTGTT (SEQ ID NO: 37)	314	0	KLK2	S39329	1631-1633

TABLE 1-continued

EXAMPLES OF DIFFERENTIALLY EXPRESSED GENES AND THEIR FUNCTIONAL CLASSIFICATIONS					
Signatures (tpm) (tpm)	LNCaP Description	CL1 Gen- Bank IDNOS:			SEQ ID
CD markers					
GATCAGAGAAGATGATA (SEQ ID NO: 38)	0	810	CD213a2, interleukin 13 receptor, alpha 2	U70981	1634-1635
GATCCCTAGGTCTTGGG (SEQ ID NO: 39)	23	161	CD213a1, interleukin 13 receptor, alpha 1	AW874023	1636
GATCCACATCTCTACA (SEQ ID NO: 40)	0	63	CD33, CD33 antigen (gp67)	BC028152	1637-1638
GATCAATAAATAATGAGG (SEQ ID NO: 41)	0	151	CD44, CD44 antigen	AL832642	1639-1640
GATCCTTCAGCCTTCAG (SEQ ID NO: 42)	0	35	CD73, 5'- nucleotidase, ecto (CD73)	A1831695	1641
GATCTGGAACCTCAGCC (SEQ ID NO: 43)	1	50	CD49e, integrin, alpha 5	BC008786	1642-1643
GATCAGAGATGCACCAC (SEQ ID NO: 44)	8	122	CD138, syndecan 1	BM974052	1644
GATCAAAGGTTTAAAGT (SEQ ID NO: 45)	38	189	CD 166, activated leukocyte cell adhesion molecule	AL833702	1645
GATCAGCTGTTTGTCAT (SEQ ID NO: 46)	53	295	CD71, transferrin receptor (p90, CD71)	BC001188 1646-1647	1646-1647
GATCGGTGCGTTCTCCT (SEQ ID NO: 47)	287	509	CD107a, A1521424 lysosomal- associated membrane protein 1	1648	
GATCTACAAGGCCATG (SEQ ID NO: 48)	161	681	CD29, integrin, beta 1	NM_002211	1649-1650
GATCATTTATTTAAGC (SEQ ID NO: 49)	56	0	CD10 (neutral endopeptidase, enkephalinase)	BQ013520	1651
GATCAGTCTTTATTAAT (SEQ ID NO: 50)	150	50	CD107b, lysosomal- associated membrane protein 2	A1459107	16652
GATCTTGCTGTATTTA (SEQ ID NO: 51)	84	1014	CD59 antigen p18- 20	NM_000611	1653-1654
GATCTTGCTGTGCTA (SEQ ID NO: 52)	408	234	CD9 antigen (p24)	NM_001769	1655-1656
Transcription factors					
GATCAAATAACAAGTCT (SEQ ID NO: 53)	0	62	transcription factor BMAL2	BM854818	1657
GATCTCTATGTTTACTT (SEQ ID NO: 54)	0	27	transcription factor BMAL2	BG163364	1658
GATCCTGACACATAAGA (SEQ ID NO: 55)	12	74	transcription factor BMAL2	BF055294	1659

TABLE 1-continued

EXAMPLES OF DIFFERENTIALLY EXPRESSED GENES AND THEIR FUNCTIONAL CLASSIFICATIONS					
Signatures (tpm) (tpm)	LNcaP Description	CL1 Gen- Bank IDNOS:			SEQ ID
GATCATTTTGTATTAAT (SEQ ID NO: 56)	10	61 transcription factor NRF	BC047878		1660-1661
GATCGTCTCATATTGC (SEQ ID NO: 57)	52	0 transcriptional coactivator tubedown-100	NM_025085		1662-1663
GATCCCCCTCTTCAATG (SEQ ID NO: 58)	0	31 transcriptional co- activator with PDZ-binding motif	AJ299431		1664-1665
GATCAAATGCTATTGCA (SEQ ID NO: 59)	1	55 transcriptional regulator interacting with the PHS-bromodomain 2	A1126500	1666	
GATCTGTGACAGCAGCA (SEQ ID NO: 60)	140	35 transducer of ERBB2, 1	BC031406		1667-1668
GATCAAATCTGTACAGT (SEQ ID NO: 61)	239	23 transducer of ERBB2, 2	A694240		1669
Annexins and their ligands					
GATCCTGTGCAACAAGA (SEQ ID NO: 62)	0	69 annexin A10	BC007320		1670-1671
GATCTGTGGTGGCAATG (SEQ ID NO: 63)	41	630 annexin A11	AL576782		1672
GATCAGAATCATGGTCT (SEQ ID NO: 64)	0	1079 annexin A2	BC001388		1673-1674
GATCTCTTTGACTGCTG (SEQ ID NO: 65)	210	860 annexin A5	BC001429		1675-1676
GATCCAAAACATCCTG (SEQ ID NO: 66)	83	241 annexin A6	AI566871		1677
GATCAGAAGACTTTAAT (SEQ ID NO: 67)	0	695 annexin A1	B001597		1678-1679
GATCAGGACACTTAGCA (SEQ ID NO: 68)	0	2949 S100 calcium binding protein A10 (annexin II ligand)	BC015973		1680-1681
Matrix metalloproteinase					
GATCATCACAGTTTGAG (SEQ ID NO: 69)	0	38 matrix metalloproteinase 10 (stromelysin 2)	BC002591		1682-1683
GATCCCAGAGAGCAGCT (SEQ ID NO: 70)	0	108 matrix metalloproteinase 1 (interstitial collagenase)	BC013118		1684-1685
GATCGGCCATCAAGGGA (SEQ ID NO: 71)	0	25 matrix metalloproteinase 13 (collagenase 3)	AI370581		16686
GATCTGGACCAGAGACA (SEQ ID NO: 72)	0	10 matrix metalloproteinase 2 (gelatinase A)	BG332150		1687

Matrix metalloproteinases (MMPs), which degrade extracellular matrix components that physically impede cell migration, are implicated in tumor cell growth, invasion, and metastasis. MMP1, 2, 10 and 13 were found to be significantly overexpressed in CL1 cells (Table 1), which may partially explain these cells' aggressive and metastatic behavior.

CD (cluster designation of monoclonal antibodies) markers are generally localized at the cell surface; some may be associated with prostate cancer (Liu, A. Y., et al., *Prostate*, 40: 192-199, 1999). All currently identified CD markers (CD1 to CD247) from the PROW CD index database (<http://colondoubleslashwww.ncbi.nlm.nih.gov/prow/guide/45277084.htm>) were converted to UniGene numbers and the UniGene numbers used to identify their signatures and their expression levels. Fifteen CD markers were identified that were differentially expressed between LNCaP and CL1 cells (Z score < 0.001) (Table 1). Eleven CD markers, including CD213a2 and CD213a1, which encode IL-13 receptors alpha 1 and 2, are up regulated in CL1 cells; three CD markers, CD9, CD10, and CD107, were downregulated in these cells (Table 1). Six CD markers went from 0 or 1 tpm to >35 tpm (Table 1), making them good digital or absolute markers or therapeutic targets. These data suggest that carefully selected CD markers may be useful in following the progression of prostate cancer, and indeed could serve as potential targets for antibody-mediated therapies (Liu, A. Y., et al., *Prostate*, 40: 192-199, 1999).

Delineation of disease-perturbed networks in prostate cancer cells. Genes and proteins rarely act alone but rather gen-

erally operate in networks of interactions. Identifying key nodes (proteins) in the disease-perturbed networks may provide insights into effective drug targets. Comparing the genes (proteins) currently available in the 314 BioCarta and 155 KEGG pathway or network (<http://colondoubleslashcgapdotnci.nih.gov/Pathways/>) databases with the MPSS data through UniGene IDs, we identified 37 BioCarta and 14 KEGG pathways that are up regulated and 23 BioCarta and 22 KEGG pathways down regulated in LNCaP cells versus CL1 cells (Table 2). The number of genes whose expression patterns changed in each pathway is listed in Table 2. Each gene along with its expression level in LNCaP and CL1 cells is listed pathway by pathway in our database (<ftp://colondoubleslashftp.systemsbiology.net/blin/mpss>). Changes in these pathways reveal the underlying phenotypic differences between LNCaP and CL1 cells. For example, multiple networks involved in modulating cell mobility, adhesion and spreading are up regulated in CL1 cells, which are more metastatic and invasive than LNCaP cells (Table 2). In the uCalpain and Friends in Cell Spread pathway, calpains are calcium-dependent thiol proteases implicated in cytoskeletal rearrangements and cell migration. During cell migration, calpain cleaves target proteins such as talin, ezrin, and paxillin at the leading edge of the membrane, while at the same time cleaving the cytoplasmic tails of the integrins β 1(a) and β 3(b) to release adhesion attachments at the trailing membrane edge. Increased activity of calpains increases migration rates and facilitates cell invasiveness (Liu, A. et al., *Prostate*, 40: 192-199, 1999).

TABLE 2

PATHWAYS THAT ARE UP OR DOWN REGULATED COMPARING LNCAP TO CL1 CELLS.				
Pathways	# Genes hits in a pathway	# p < 0.001 & LNCA > CL1	# p < 0.001 & LNCA < CL1	# no change
Up-regulated Pathways in LNCAP cells				
BioCarta Pathways				
Mechanism of Gene Regulation by Peroxisome Proliferators via PPAR α	35	9	2	24
T Cell Receptor Signaling Pathway	21	6	2	13
ATM Signaling Pathway	15	5	2	8
CARM1 and Regulation of the Estrogen Receptor	18	5	2	11
HIV-1 Nef negative effector of Fas and TNF	33	5	2	26
EGF Signaling Pathway	17	5	1	11
Role of BRCA1 BRCA2 and ATR in Cancer Susceptibility	16	5	1	10
TNFR1 Signaling Pathway	17	5	1	11
Toll-Like Receptor Pathway	17	5	1	11
FAS signaling pathway CD95	17	4	1	12
VEGF Hypoxia and Angiogenesis	16	4	1	11
Bone Remodelling	9	3	1	5
ER associated degradation	11	3	1	7
ERAD Pathway				
Estrogen-responsive protein Efp controls cell cycle and breast tumors growth	11	3	1	7
Influence of Ras and Rho proteins on G1 to S Transition	16	3	1	12
Inhibition of Cellular Proliferation by Gleevec	13	3	1	9
Map Kinase Inactivation of SMRT Corepressor	9	3	1	5

TABLE 2-continued

PATHWAYS THAT ARE UP OR DOWN REGULATED COMPARING LNCAp TO CL1 CELLS.				
Pathways	# Genes hits in a pathway	# p < 0.001 & LNCA > CL1	# p < 0.001 & LNCA < CL1	# no change
NFkB activation by Nontypeable Hemophilus influenzae	16	3	1	12
RB Tumor Suppressor Checkpoint Signaling in response to DNA damage	10	3	1	6
Transcription Regulation by Methyltransferase of CARM1	10	3	1	6
Ceramide Signaling Pathway	13	4	0	9
Cystic fibrosis transmembrane conductance regulator and beta 2 adrenergic receptor pathway	7	4	0	3
Nerve growth factor pathway NGF	11	4	0	7
PDGF Signaling Pathway	16	4	0	12
TNF Stress Related Signaling	14	4	0	10
Activation of Csk by cAMP- dependent Protein Kinase Inhibits Signaling through the T Cell Receptor	9	3	0	6
AKAP95 role in mitosis and chromosome dynamics	11	3	0	8
Attenuation of GPCR Signaling	7	3	0	4
Chaperones modulate interferon Signaling Pathway	11	3	0	8
ChREBP regulation by carbohydrates and cAMP	12	3	0	9
IGF-1 Signaling Pathway	11	3	0	8
Insulin Signaling Pathway	11	3	0	8
NF-kB Signaling Pathway	11	3	0	8
Protein Kinase A at the Centrosome	12	3	0	9
Regulation of ck1 cdk5 by type 1 glutamate receptors	10	3	0	7
Role of Mitochondria in Apoptotic Signaling	10	3	0	7
Signal transduction through IL1R	14	3	0	11
KEGG Pathways				
Aminosugars metabolism	24	9	4	11
Androgen and estrogen metabolism	37	13	5	19
Benzoate degradation via hydroxylation	5	3	1	1
C21-Steroid hormone metabolism	4	1	0	3
C5-Branched dibasic acid metabolism	2	2	0	0
Carbazole degradation	1	1	0	0
Terpenoid biosynthesis	6	4	1	1
Chondroitin_heparan sulfate biosynthesis	14	8	3	3
Fatty acid biosynthesis (path 1)	3	2	0	1
Fluorene degradation	3	2	0	1
Pentose and glucuronate interconversions	19	9	1	9
Phenylalanine, tyrosine and tryptophan biosynthesis	10	5	2	3
Porphyrin and chlorophyll metabolism	28	13	3	12
Streptomycin biosynthesis	6	4	1	1
Up-regulated Pathways in CL1 cells				
BioCarta Pathways				
Rho cell motility signaling pathway	18	2	6	10
Trefoil Factors Initiate Mucosal Healing	14	1	6	7
Integrin Signaling Pathway	14	1	5	8

TABLE 2-continued

PATHWAYS THAT ARE UP OR DOWN REGULATED COMPARING LNCAp TO CL1 CELLS.				
Pathways	# Genes hits in a pathway	# p < 0.001 & LNCA > CL1	# p < 0.001 & LNCA < CL1	# no change
Ca Calmodulin-dependent Protein Kinase Activation	7	1	4	2
Effects of calcineurin in Keratinocyte Differentiation	9	1	4	4
Angiotensin II mediated activation of JNK Pathway via Pyk2 dependent signaling	12	1	3	8
Bioactive Peptide Induced Signaling Pathway	16	1	3	12
CBL mediated ligand-induced downregulation of EGF receptors	6	1	3	2
Control of skeletal myogenesis by HDAC calcium calmodulin-dependent kinase CaMK	12	1	3	8
How does salmonella hijack a cell	8	1	3	4
Melanocyte Development and Pigmentation Pathway	4	1	3	0
Overview of telomerase protein component gene hTert	7	1	3	3
Transcriptional Regulation				
Regulation of PGC-1a	9	0	4	5
ADP-Ribosylation Factor	9	0	3	6
Downregulated of MTA-3 in ER-negative Breast Tumors	7	0	3	4
Endocytotic role of NDK	7	0	3	4
Phosphins and Dynamin				
Mechanism of Protein Import into the Nucleus	7	0	3	4
Nuclear Receptors in Lipid Metabolism and Toxicity	7	0	3	4
Pertussis toxin-insensitive CCR5 Signaling in Macrophage	9	0	3	6
Platelet Amyloid Precursor Protein Pathway	5	0	3	2
Role of Ran in mitotic spindle regulation	8	0	3	5
Sumoylation by RanBP2	8	0	3	5
Regulates Transcriptional Repression				
uCalpain and friends in Cell spread	5	0	3	2
KEGG Pathways				
Arginine and proline metabolism	45	7	16	22
ATP synthesis	31	7	15	9
Biotin metabolism	5	1	3	1
Blood group glycolipid biosynthesis-lactoseries	12	1	6	5
Cyanoamino acid metabolism	5	0	3	2
Ethylbenzene degradation	9	1	3	5
Ganglioside biosynthesis	16	2	6	8
Globoside metabolism	17	3	8	6
Glutathione metabolism	26	4	10	12
Glycine, serine and threonine metabolism	32	6	14	12
Glycosphingolipid metabolism	35	6	18	11
Glycosylphosphatidylinositol(GPI)- anchor biosynthesis	26	5	12	9
Glyoxylate and dicarboxylate metabolism	9	1	6	2
Huntington's disease	25	4	10	11
Methane metabolism	9	1	3	5
O-Glycans biosynthesis	19	3	8	8
One carbon pool by folate	12	2	8	2
Oxidative phosphorylation	93	21	45	27
Parkinson's disease	30	5	14	11
Phospholipid degradation	21	4	12	5

TABLE 2-continued

PATHWAYS THAT ARE UP OR DOWN REGULATED COMPARING LNCAP TO CL1 CELLS.				
Pathways	# Genes hits in a pathway	# p < 0.001 & LNCA > CL1	# p < 0.001 & LNCA < CL1	# no change
Synthesis and degradation of ketone bodies	7	1	3	3
Urea cycle and metabolism of amino groups	18	2	8	8

Many pathways we identified as perturbed in the LNCaP and CL1 comparison are interconnected to form networks (in fact there are probably no discrete pathways, only networks). For example, the insulin signaling pathway, the signal transduction through IL1R pathway, NF-kB signaling pathway are interconnected through c-Jun, IL1R and NF-kB. The mapping of genes onto networks/pathways will be an ongoing objective as more networks/pathways become available. Our transcriptome data will be an invaluable resource in delineating these relationships.

As gene regulatory networks controlled by transcription factors form the top layer of the hierarchy that controls the physiological network, we sought to identify differentially expressed transcription factors. Of 554 transcription factors expressed in LNCaP and CL1 cells, 112 showed significantly different levels between the cell lines ($P < 0.001$) This clearly demonstrated significant difference in the functioning of the corresponding gene regulatory networks during the progression of prostate cancer from the early to late stages.

Quantitative Proteomics Analysis of Prostate Cancer Cells. We quantitatively profiled the protein expression changes between LNCaP and CL1 cells using the ICAT-MS/MS protocol described by Han et al. *Nat Biotechnol*, 19: 946-951, 2001. To increase proteome coverage, cells were separated into nuclear, cytosolic and microsomal fractions prior to ICAT analysis as described in Han et al., 2001, supra. We generated a total of 142,849 tandem mass spectra, 7282 of which corresponded to peptides with a mass spectrum quality score P value (Keller, A., et al., *Anal Chem.* 2002 Oct 15;74 (20):5383-92) greater than 0.9 (allowing unambiguous identification of peptides). These 7282 peptides represented 971 proteins (Keller, A., et al., 2002, supra). We obtained quantitative peptide ratios for 4583 peptides corresponding to 941 proteins. The number of peptides is greater than the number of proteins because 1) mass spectrometry identified multiple peptides from the same protein and 2) the ionization step of mass spectrometry created different charge states for the same peptide. The protein ratios were calculated from multiple peptide ratios using an algorithm for the automated statistical analysis of protein abundance ratios (ASAPRatio) (Li, X. J., et al., *Anal Chem*, 75: 6648-6657, 2003). In the end, we identified 82 proteins that are down regulated and 108 proteins that are up regulated by at least 1.8-fold in LNCaP cells compared with CL1 cells. For example, five proteins belong to annexins that were markers for prostate and other cancers (Hayes, M. J. and Moss, S. E. *Biochem Biophys Res Commun*, 322: 1166-1170, 2004), seven are involved in fatty acids and lipid metabolism that are involved in the carcinogenesis and progression of prostate cancer (Pandian, S. S., et al., *J R Coll Surg Edinb*, 44: 352-361, 1999), five are related to apoptosis, 11 are cancer related, and five proteins are putative transcription factors. As we only identified a limited number of proteins that are significantly differentially

expressed due to low sensitivity of ICAT technology, we were only able to identify a few pathways that are perturbed based on ICAT data alone (using the stringent criteria discussed above). This also illustrated the importance of MPSS analysis described earlier.

103 of 190 (54%) differentially expressed proteins identified have enzymatic activity and hence many are involved in metabolism. Notably, many of the proteins identified are involved in fatty acid and lipid metabolism, including fatty acid synthase, carnitine palmitoyltransferase II and propionyl Coenzyme A carboxylase alpha polypeptide. Fatty acid and lipid metabolism is known to be perturbed in prostate cancer (Fleshner, N., et al., *J Urol*, 171: S19-24, 2004). Additionally, many genes involved in lipid transport were altered, including the annexins, prosaposin, and fatty acid binding protein 5. Annexin A1 has previously been shown to be overexpressed in non-PSA-producing LNCaP cells as compared with PSA-producing LNCaP cells (Vaarala, M. H., et al., 2000, supra). Annexin A7 is postulated to be a prostate tumor suppressor gene (Cardo-Vila, M., et al., *Pharmacogenomics J*, 1: 92-94, 2001). Annexin A2 expression is reduced or lost in prostate cancer cells, and its re-expression inhibits prostate cancer cell migration (Liu, J. W., et al., *Oncogene*, 22: 1475-1485, 2003).

Other genes identified here have been implicated in carcinogenesis, including tumor suppressor p16 and insulin-like growth factor 2 receptor (Chi, S. G., et al., *Clin Cancer Res*, 3: 1889-1897, 1997; Kiess, W., et al., *Horm Res*, 41 Suppl 2: 66-73, 1994). Some genes have previously been implicated in prostate cancer, such as prostate cancer over expressed gene 1 POV1, which is over expressed in prostate cancer (Cole, K. A., et al., *Genomics*, 51: 282-287, 1998), and delta 1 and alpha 1 catenin (cadherin-associated protein) and junction plakoglobin, which are down regulated in prostate cancer cells (Kallakury, B. V., et al., *Cancer*, 92: 2786-2795, 2001). However, the potential relationships of most of the proteins identified here to prostate cancer require further elucidation. For example, transmembrane protein 4 (TMEM4), a gene predicted to encode a 182-amino acid type II transmembrane protein, is downregulated about twofold in CL1 cells compared with LNCaP cells. MPSS data also indicated that TMEM4 is down regulated about twofold in CL1 cells. Many type II transmembrane proteins, such as TMPRSS2, are overexpressed in prostate cancer patients (Vaarala, M. H., et al., *Int J Cancer*, 94: 705-710, 2001). It will be interesting to see whether TMEM4 overexpression plays a primary role in prostate carcinogenesis. We also identified 12 proteins that have not been annotated or functionally characterized.

The mRNA expression level of eight proteins change from 0 tpm in LNCaP cells to greater than 50 tpm (we called them 'digital changes' because they go from zero to some expression) in CL1 cells, and that of one protein changed from 0 tpm in CL1 cells to greater than 50 in LNCaP cells. These genes can be used as digital diagnostic signals. Twenty-two of the

differentially expressed proteins were predicted to be secreted proteins (See Table 3) and can be further evaluated as serum marker (see also Example 2 below).

Additionally, we sought to compare the expression at the protein level with that at the mRNA level. We converted the protein IDs and MPSS signatures to Unigene IDs to compare the MPSS data with the ICAT-MS/MS data. We limited this comparison to those with common Unigene IDs and with reliable ICAT ratios (standard deviation less than 0.5) and ended up with a subset of 79 proteins. Of these, 66 genes (83.5%) were concordant in their changes in mRNA and protein levels of expression and 13 genes (16.5%) were discordant, i.e. having higher protein expression but lower

mRNA expression or vice versa. There are no functional similarities among the discordant genes. As these mRNAs and proteins are expressed at relatively high levels, discordance due to measurement errors is unlikely. Clearly post-transcriptional mechanism(s) of protein expression are functioning, although the elucidation of the specific mechanism(s) awaits further studies.

This system provides a model for studying perturbation of organ-specific molecular blood fingerprints. These results, and those described in the Examples below, indicate a systems approach will offer powerful tools for disease diagnostics, drug side effects diagnostics, and therapeutics.

TABLE 3

DIFFERENTIALLY EXPRESSED GENES THAT ENCODE PREDICTED SECRETED PROTEINS.

Signature	SEQ ID NO: Number	Accession Number	SEQ ID NOS:	Description
GATCAGCATGGGCCACG	73	NM_001928	594-595	D component of complement (adipsin)
GATCTACTACTGGCCT	74	NM_006280	596-597	signal sequence receptor, delta (translocon associated protein delta)
GATCCTGTTGGGAAAGA	75	NM_203329	598-599	CD59 antigen p18-20 (antigen identified by monoclonal antibodies 16.3A5, EJ16, EJ30, EL32 and G344)
GATCCTGTTGGGAAAGA	76	NM_203331	600-601	CD59 antigen p18-20 (antigen identified by monoclonal antibodies 16.3A5, EJ16, EJ30, EL32 and G344)
GATCCCTGAAGTTGCC	77	NM_203331	600-601	CD59 antigen p18-20 (antigen identified by monoclonal antibodies 16.3A5, EJ16, EJ30, EL32 and G344)
GATCTGGCTGTATTTA	78	NM_203331	600-601	CD59 antigen p18-20 (antigen identified by monoclonal antibodies 16.3A5, EJ16, EJ30, EL32 and G344)
GATCCCTGAAGTTGCC	79	NM_203330	602-603	CD59 antigen p18-20 (antigen identified by monoclonal antibodies 16.3A5, EJ16, EJ30, EL32 and G344)
GATCCTGTTGGGAAAGA	80	NM_203330	602-603	CD59 antigen p 18-20 (antigen identified by monoclonal antibodies 16.3A5, EJ16, EJ30, EL32 and G344)
GATCTGGCTGTATTTA	81	NM_203330	602-603	CD59 antigen p18-20 (antigen identified by monoclonal antibodies 16.3A5, EJ16, EJ30, EL32 and G344)

TABLE 3-continued

DIFFERENTIALLY EXPRESSED GENES THAT ENCODE PREDICTED SECRETED PROTEINS.				
Signature	SEQ ID Accession		SEQ ID	
	NO:	Number	NOS:	Description
GATCCCTGAAGTTGCC	82	NM_203329	598-599	CD59 antigen p18-20 (antigen identified by monoclonal antibodies 16.3A5, EJ16, EJ30, EL32 and G344)
GATCTTGCTGTATTTA	83	NM_000611	604-605	CD59 antigen p18-20 (antigen identified by monoclonal antibodies 16.3A5, EJ16, EJ30, EL32 and G344)
GATCCCTGAAGTTGCC	84	NM_000611	604-605	CD59 antigen p18-20 (antigen identified by monoclonal antibodies 16.3A5, EJ16, EJ30, EL32 and G344)
GATCCTGTGGGAAAGA	85	NM_000611	604-605	CD59 antigen p18-20 (antigen identified by monoclonal antibodies 16.3A5, EJ16, EJ30, EL32 and G344)
GATCTTGCTGTATTTA	86	NM_203329	598-599	CD59 antigen p18-20 (antigen identified by monoclonal antibodies 16.3A5, EJ16, EJ30, EL32 and G344)
GATCTGTGCTGACCCCA	87	NM_002982	606-607	chemokine (C-C motif) ligand 2
GATCTCTTGAATGACA	88	NM_012242	608-609	dickkopf homolog 1 (<i>Xenopus laevis</i>)
GATCACCATCAAGCCAG	89	NM_012242	608-609	dickkopf homolog 1 (<i>Xenopus laevis</i>)
GATCAACAGCTCTAGT	90	NM_016308	610-611	UMP-CMP kinase
GATCCCTGTTACGACA	91	NM_014155	612-613	HSPC063 protein
GATCTCTGATTACCAGC	92	NM_025205	614-615	mediator of RNA polymerase II transcription, subunit 28 homolog (yeast)
GATCATTGAACGAGACA	93	NM_031903	616-617	mitochondrial ribosomal protein L32
GATCACAGACCACGAGT	94	NM_178507	618-619	NS5ATP13TP2 protein
GATCTGCATCAGTTGTA	95	NM_148170	620-621	cathepsin C
GATCTCTTGCTAGATTT	96	NM_005059	622-623	relaxin 2
GATCACAAGGCTGCCTG	97	NM_000405	624-625	GM2 ganglioside activator
GATCGTTTCTCATCTCT	98	NM_006432	626-627	Niemann-Pick disease, type C2
GATCCCGCGATACTTC	99	NM_015921	628-629	chromosome 6 open reading frame 82

TABLE 3-continued

DIFFERENTIALLY EXPRESSED GENES THAT ENCODE PREDICTED SECRETED PROTEINS.				
Signature	SEQ ID NO:	Accession Number	SEQ ID NOS:	Description
GATCTTTTTTGGATAT	100	NM_181777	630-631	ubiquitin-conjugating enzyme E2A (RAD6 homolog)
GATCCGAGAGTAAGGAA	101	NM_032488	632-633	cornifelin
GATCATGTGTTTCCATG	102	NM_014435	634-635	N-acylsphingosine amidohydrolase (acid ceramidase)-like
GATCTCAGAACAACCTT	103	NM_016029	636-637	dehydrogenase/reductase (SDR family) member 7
GATCTTACCTCCTGATA	104	NM_020467	638-639	hypothetical protein from clone 643
GATCCCAGACTGGTTCT	105	NM_003782	640-641	UDP-Gal:betaGlcNAc beta 1,3-galactosyltransferase, polypeptide 4
GATCAAGTGCATTTGAC	106	NM_173631	642-643	zinc finger protein 547
GATCAGTGCATGATGGA	107	NM_005423	644-645	trefoil factor 2 (spasmolytic protein 1)
GATCCAAGAGGAAGAAT	108	NM_014402	646-647	low molecular mass ubiquinone-binding protein (9.5kD)
GATCCAGCAAACAGGTT	109	NM_003851	648-649	cellular repressor of E1A-stimulated genes 1
GATCATAGAAGGCTATT	110	NM_181834	650-651	neurofibromin 2 (bilateral acoustic neuroma)
GATCCCCCTTCATTGA	111	NM_004862	652-653	lipopolysaccharide induced TNF factor
GATCCCAAATTGAAGT	112	NM_001685	654-655	ATP synthase, H ⁺ transporting, mitochondrial F0 complex, subunit F6
GATCTGCTTTCTGTAAT	113	NM_002406	656-657	mannosyl (alpha-1,3-)-glycoprotein beta-1,2-N-acetylglucosaminyl-transferase
GATCACTCCTTATTTGC	114	NM_019021	658-659	hypothetical protein FLJ20010
GATCACCTTCGACGACT	115	NM_003130	660-661	sorcin
GATCTCTATTGTAATCT	116	NM_002489	662-663	NADH dehydrogenase (ubiquinone) 1 alpha subcomplex, 4, 9kDa
GATCTCCTGGTGCAAA	117	NM_138429	664-665	claudin 15
GATCCCAGTCTCTGCCA	118	NM_201397	666-667	glutathione peroxidase 1
GATCTTCTTTATAATTC	119	NM_004048	668-669	beta-2-microglobulin

TABLE 3-continued

DIFFERENTIALLY EXPRESSED GENES THAT ENCODE PREDICTED SECRETED PROTEINS.				
Signature	SEQ ID NO:	Accession Number	SEQ ID NOS:	Description
GATCTGTTCAAACAGCA	120	NM_024060	670-671	hypothetical protein MGC5395
GATCGTGCTCACAGGCA	121	NM_033280	672-673	SEC11-like 3 (<i>S. cerevisiae</i>)
GATCAATATGTAAATAT	122	NM_020199	674-675	chromosome 5 open reading frame 15
GATCAGCTTTGCTCCTG	123	NM_207495	676-677	hypothetical protein DKFZp686115217
GATCTCTATGGCTGTAA	124	NM_033211	678-679	hypothetical gene supported by AF038182; BC009203
GATCTCAGAACCCTGT	125	NM_001001436	680-681	similar to RIKEN cDNA 4921524J17
GATCCAGCCATTACTAA	126	NM_016205	682-683	platelet derived growth factor C
GATCTTCCCAAGATTG	127	NM_001001434	684-685	syntaxin 16
GATCGATTCTGTGACAC	128	NM_181726	686-687	low density lipoprotein receptor- related protein binding protein
GATCTATTTTTTCTAAA	129	NM_004125	688-689	guanine nucleotide binding protein (G protein), gamma 10
GATCAAGAATCCTGCTC	130	NM_006332	690-691	interferon, gamma- inducible protein 30
GATCGGTGGAGAACCTC	131	NM_175742	692-693	melanoma antigen, family A, 2
GATCGGTGGAGAACCTC	132	NM_175743	694-695	melanoma antigen, family A, 2
GATCGGTGGAGAACCTC	133	NM_153488	696-697	melanoma antigen, family A, 2B
GATCATGGGTGAGGGGT	134	NM_001483	698-699	glioblastoma amplified sequence
GATCCCCCTCACCATGA	135	NM_032621	700-701	brain expressed X- linked 2
GATCAACTAATAGCTCT	136	NM_181892	702-703	ubiquitin-conjugating enzyme E2D 3 (UBC4/5 homolog, yeast)
GATCAAAATAAGTTATA	137	NM_181892	702-703	ubiquitin-conjugating enzyme E2D 3 (UBC4/5 homolog, yeast)
GATCAAGGAGACCCGGA	138	NM_024540	704-705	mitochondrial ribosomal protein L24
GATCAAGGAGACCCGGA	139	NM_145729	706-707	mitochondrial ribosomal protein L24
GATCCTAAGCCATAGAC	140	NM_025075	708-709	Ngg1 interacting factor 3 like 1 binding protein 1

TABLE 3-continued

DIFFERENTIALLY EXPRESSED GENES THAT ENCODE PREDICTED SECRETED PROTEINS.				
Signature	SEQ ID NO:	Accession Number	SEQ ID NOS:	Description
GATCCATTGAGCCCAGC	141	M181725	710-711	hypothetical protein FLJ12760
GATCTGAGGGCGTCTTC	142	NM_012153	712-713	ets homologous factor
GATCTCGGTAGTTACGT	143	NM_012153	712-713	ets homologous factor
GATCCCAAGATGATTAA	144	NM_014177	714-715	chromosome 18 open reading frame 55
GATCTCAAACCTGTCTT	145	NM_003350	716-717	ubiquitin conjugating enzyme E2 variant 2
GATCATAGTTATTATAC	146	NM_032466	718-719	aspartate beta- hydroxylase
GATCCCAACTGCTCCTG	147	NM_005947	720-721	metallothionein 1B (functional)
GATCAAATGCTAAAAC	148	NM_016311	722-723	ATPase inhibitory factor 1
GATCTGTTTGTCCCTG	149	NM_013411	724-725	adenylate kinase 2
GATCAACAGTGGCAATG	150	NM_001001392	726-727	CD44 antigen (homing function and Indian blood group system)
GATCAATAAATGAGG	151	NM_001001392	726-727	CD44 antigen (homing function and Indian blood group system)
GATCAACTAATAGCTCT	152	NM_181890	728-729	ubiquitin-conjugating enzyme E2D 3 (UBC4/5 homolog, yeast)
GATCAAATAAAGTTATA	153	NM_181891	730-731	ubiquitin-conjugating enzyme E2D 3 (UBC4/5 homolog, yeast)
GATCAAATAAAGTTATA	154	NM_181890	728-729	ubiquitin-conjugating enzyme E2D 3 (UBC4/5 homolog, yeast)
GATCAAATAAAGTTATA	155	NM_181889	732-733	ubiquitin-conjugating enzyme E2D 3 (UBC4I5 homolog, yeast)
GATCAACTAATAGCTCT	156	NM_003340	734-735	ubiquitin-conjugating enzyme E2D 3 (UBC4/5 homolog, yeast)
GATCAACTAATAGCTCT	157	NM_181888	736-737	ubiquitin-conjugating enzyme E2D 3 (UBC4/5 homolog, yeast)
GATCAAATAAAGTTATA	158	NM_181888	736-737	ubiquitin-conjugating enzyme E2D 3 (UBC4/5 homolog, yeast)

TABLE 3-continued

DIFFERENTIALLY EXPRESSED GENES THAT ENCODE PREDICTED SECRETED PROTEINS.				
Signature	SEQ ID NO:	Accession Number	SEQ ID NOS:	Description
GATCAACTAATAGCTCT	159	NM_181891	730-731	ubiquitin-conjugating enzyme E2D 3 (UBC4/5 homolog, yeast)
GATCAACTAATAGCTCT	160	NM_181887	738-739	ubiquitin-conjugating enzyme E2D 3 (UBC4/5 homo log, yeast)
GATCAAATAAAGTTATA	161	NM_181887	738-739	ubiquitin-conjugating enzyme E2D 3 (UBC4/5 homolog, yeast)
GATCAACTAATAGCTCT	162	NM_181886	740-741	ubiquitin-conjugating enzyme E2D 3 (UBC4/5 homolog, yeast)
GATCAAATAAAGTTATA	163	NM_181886	740-741	ubiquitin-conjugating enzyme E2D 3 (UBC4/5 homolog, yeast)
GATCAAATAAAGTTATA	164	NM_003340	734-735	ubiquitin-conjugating enzyme E2D 3 (UBC4/5 homolog, yeast)
GATCAACTAATAGCTCT	165	NM_181889	732-733	ubiquitin-conjugating enzyme E2D 3 (UBC4/5 homolog, yeast)
GATCTGATTTTTTCCCC	166	NM_145751	742-743	TNF receptor-associated factor 4
GATCAGAAATGACTGTG	167	NM_018509	744-745	hypothetical protein PR01855
GATCACTGAGAAAAAAT	168	NM_152407	746-747	GrpE-like 2, mitochondrial (<i>E. coli</i>)
GATCCAAGAGTTTAGTG	169	NM_006807	748-749	chromobox homolog 1 (HP1 beta homolog <i>Drosophila</i>)
GATCTTTGCTGGCAAGC	170	NM_002954	750-751	ribosomal protein S2 7a
GATCCACTGAGAGAG	171	NM_145864	752-753	kallikrein 3, (prostate specific antigen)
GATCTGTATTATTAAT	172	NM_032549	754-755	IMP2 inner mitochondrial membrane protease-like (<i>S. cerevisiae</i>)
GATCTGTTTGTCCCTG	173	NM_172199	756-757	adenylate kinase 2
GATCCCCTGCCTGGTGC	174	NM_001312	758-759	cysteine rich protein 2
GATCAACTAATAGCTCT	175	NM_181893	760-761	ubiquitin-conjugating enzyme E2D 3 (UBC4/5 homolog, yeast)

TABLE 3-continued

DIFFERENTIALLY EXPRESSED GENES THAT ENCODE PREDICTED SECRETED PROTEINS.				
Signature	SEQ ID Accession		SEQ ID	
	NO:	Number	NOS:	Description
GATCAATAAAGTTATA	176	NM_181893	760-761	ubiquitin-conjugating enzyme E2D 3 (UBC4/5 homolog, yeast)
GATCTTTTCAAGTCTT	177	NM_012071	762-763	COMM domain containing 3
GATCATGTATGAGATAG	178	NM_012460	764-765	translocase of inner mitochondrial membrane 9 homolog (yeast)
GATCCTTCAGGCAGTAA	179	NM_176805	766-767	mitochondrial ribosomal protein S11
GATCTTTTTTGGATAT	180	NM_003336	768-769	ubiquitin-conjugating enzyme E2A (RAD6 homolog)
GATCCCAGTCTCTGCCA	181	NM_000581	770-771	glutathione peroxidase 1
GATCAAGACGAGCCTGC	182	NM_004864	772-773	growth differentiation factor 15
GATCCCAGCTGATGTAG	183	NM_001885	774-775	crystallin, alpha B
GATCATGAAGACCTGCT	184	NM_003754	776-777	eukaryotic translation initiation factor 3, subunit 5 epsilon, 47kDa
GATCTCAAGTTGATAG	185	NM_003864	778-779	sin3-associated polypeptide, 30kDa
GATCACCAGGCTGCCCA	186	NM_148571	780-781	mitochondrial ribosomal protein L27
GATCAAAATGCTAAAC	187	NM_178190	782-783	ATPase inhibitory factor 1
GATCAAGATGACTGA	188	NM_004483	784-785	glycine cleavage system protein H (aminomethyl carrier)
GATCGGGAACCTCTGCT	189	NM_005952	786-787	metallothionein 1X
GATCTTGCTTTAAAC	190	NM_015646	788-789	RAP1B, member of RAS oncogene family
GATCCACACGTTGGT	191	NM_003255	790-791	tissue inhibitor of metalloproteinase 2
GATCATCAGTCACCGAA	192	NM_000077	792-793	cyclin-dependent kinase inhibitor 2A (melanoma, p16, inhibits CDK4)
GATCCAGTATTCACTCA	193	NM_002166	794-795	inhibitor of DNA binding 2, dominant negative helix-loop-helixprotein
GATCCTGCAGGGAGCT	194	NM_015343	796-797	dullard homolog (<i>Xenopus laevis</i>)
GATCTCCTTGCCCCAGC	195	NM_015343	796-797	dullard homolog (<i>Xenopus laevis</i>)

TABLE 3-continued

DIFFERENTIALLY EXPRESSED GENES THAT ENCODE PREDICTED SECRETED PROTEINS.				
Signature	SEQ ID Accession		SEQ ID	
	NO:	Number	NOS:	Description
GATCGCCTAGTATGTTC	196	NM_003897	798-799	immediate early response 3
GATCAGACTGTATTAAA	197	NM_032052	800-801	zinc finger protein 278
GATCGGCCCTACTAGAT	198	NM_032052	800-801	zinc finger protein 278
GATCTCCCACTGCGGGG	199	NM_032052	800-801	zinc finger protein 278
GATCTGTGATGGTCAGC	200	NM_000232	802-803	sarcoglycan, beta (43kDa dystrophin-associated glycoprotein)
GATCACTGTGGTATCTA	201	NM_052822	804-805	secretory carrier membrane protein 1
GATCATCAGTCACCGAA	202	NM_058197	806-807	cyclin-dependent kinase inhibitor 2A (melanoma, p16, inhibits CDK4)
GATCATTTGTTTATTAA	203	NM_022334	808-809	integrin beta 1 binding protein 1
GATCAAATATGTAAAAAT	204	NM_004842	810-811	A kinase (PRKA) anchor protein 7
GATCTCTTGCTAGATTT	205	NM_134441	812-813	relaxin 2
GATCACCTTCGACGACT	206	NM_198901	814-815	sorcin
GATCGGATTGATTAATA	207	NM_020353	816-817	phospholipid scramblase 4
GATCTAGTTGGGAGATA	208	NM_153367	818-819	chromosome 10 open reading frame 56
GATCTTTTTGGCTACT	209	NM_018424	820-821	erythrocyte membrane protein band 4.1 like 4B
GATCACATTTTCTGTTG	210	NM_201436	822-823	H2A histone family, member V
GATCACCTGGGTTTCTT	211	NM_021999	824-825	integral membrane protein 2B
GATCTATTAGATCAAAA	212	NM_021105	826-827	phospholipid scramblase 1
GATCTCTTATTTTACAA	213	NM_000546	828-829	tumor protein p53 (Li-Fraumeni syndrome)
GATCATAGAAGGCTATT	214	NM_181835	830-831	neurofibromin 2 (bilateral acoustic neuroma)
GATCTTCTGGACAGGA	215	NM_152992	832-833	POM (POM121 homolog, rat) and ZP3 fusion
GATCAAGGACCGGCCCA	216	NM_032391	834-835	small nuclear protein PRAC
GATCGCATTTTGTAAA	217	NM_058171	836-837	inhibitor of growth family, member 2
GATCCATCCTCATCTCC	218	NM_020188	838-839	DC13 protein

TABLE 3-continued

DIFFERENTIALLY EXPRESSED GENES THAT ENCODE PREDICTED SECRETED PROTEINS.			
Signature	SEQ ID NO:	Accession Number	SEQ ID NOS: Description
GATCGATGGTGGCGCTT	219	NM_138992	beta-site APP- cleaving enzyme 2
GATCTTATAAAAAGAAA	220	NM_017998	840-841 chromosome 9 open reading frame 40
GATCTGAACGATGCCGT	221	NM_024579	842-843 hypothetical protein FLJ23221
GATCTCCCCGCCGAGC	222	NM_015973	844-845 galanin
GATCGTCGTCCAGGCCA	223	NM_032920	846-847 chromosome 21 open reading frame 124
GATCGTTGGGGAACCCC	224	NM_199483	848-849 chromosome 20 open reading frame 24
GATCCTATATGTCCTGT	225	NM_152344	850-851 hypothetical protein FLJ30656
GATCGATGGTTGACAAT	226	NM_004552	852-853 NADH dehydrogenase (ubiquinone) Fe-S proteins, 15kDa (NADH-coenzyme Q reductase)
GATCTTGTACTAACTTA	227	NM_019059	854-855 translocase of outer mitochondrial membrane 7 homolog (yeast)
GATCCCGATGTTCTTAA	228	NM_001806	856-857 CCAAT/enhancer binding protein (C/EBP), gamma
GATCCTGTTTAACAAAG	229	NM_015469	858-859 nipsnap homolog 3A (<i>C. elegans</i>)
GATCACGCACACACAAT	230	NM_198337	860-861 insulin induced gene 1
GATCCAGCCAGACTTGC	231	NM_144772	862-863 apolipoprotein A-I binding protein
GATCCACACTGGAGAGA	232	NM_003450	864-865 zinc finger protein 174
GATCTCAGTTCTGCGTT	233	NM_004642	866-867 CDK2-associated protein 1
GATCTACACCTCTTGCC	234	NM_052845	868-869 methylmalonic aciduria (cobalamin deficiency) type B
GATCCAGCTGGAAAGCT	235	NM_006406	870-871 peroxiredoxin 4
GATCCTTCAGGCAGTAA	236	NM_022839	872-873 mitochondrial ribosomal protein S11
GATCCACACTGAGAGAG	237	NM_001648	874-875 kallikrein 3, (prostate specific antigen)
GATCACCTTATGGATGT	238	NM_003932	876-877 suppression of tumorigenicity 13 (colon carcinoma) (Hsp70 interacting protein)
GATCTAGTTATTTTAAT	239	NM_172178	878-879 mitochondrial ribosomal protein L42

TABLE 3-continued

DIFFERENTIALLY EXPRESSED GENES THAT ENCODE PREDICTED SECRETED PROTEINS.				
Signature	SEQ ID Accession		SEQ ID	
	NO:	Number	NOS:	Description
GATCATTGAGAATGCAG	240	NM_206966	880-881	similar to AVLV472 deoxyribonuclease II
GATCATGCCAAGTGGTG	241	NM_058248	882-883	beta H2A histone family,
GATCACATTTTCTGTTG	242	NM_201516	884-885	member V
GATCAGAAAGAAACCTT	243	NM_006744	886-887	retinol binding protein 4, plasma
GATCCGTGGCAGGGCTG	244	NM_031901	888-889	mitochondrial ribosomal protein S21
GATCCGTGGCAGGGCTG	245	NM_018997	890-891	mitochondrial ribosomal protein S21
GATCTATCACCCAAACA	246	NM_198157	892-893	ubiquitin conjugating enzyme E2L3
GATCAAGCGTGCTTTCC	247	NM_000995	894-895	ribosomal protein L34
GATCAAGCGTGCTTTCC	248	NM_033625	896-897	ribosomal protein L34
GATCCCTCATCCCTGAA	249	NM_014098	898-899	peroxiredoxin 3
GATCCACCTTGGCCTCC	250	NM_147187	900-901	tumor necrosis factor receptor superfamily, member 10b
GATCTTAGGGAGACAAA	251	NM_182529	902-903	THAP domain containing 5
GATCAAGATACGGAAGA	252	NM_177924	904-905	N-acylsphingosine amidohydrolase (acid ceramidase) 1
GATCTGTTTGTTCCTG	253	NM_001625	906-907	adenylate kinase 2
GATCAGCAAAAGCCAAA	254	NM_201263	908-909	tryptophanyl tRNA synthetase 2 (mitochondrial)
GATCGGGGAGGGTAAA	255	NM_004544	910-911	NADH dehydrogenase (ubiquinone) 1 alpha subcomplex, 10, 42kDa
GATCGTGGAGGAGGGAC	256	NM_016310	912-913	polymerase (RNA) III (DNA directed) polypeptide K, 12.3 kDa
GATCACTTTTGAAAGCA	257	NM_018465	914-915	chromosome 9 open reading frame 46
GATCTGATTTGCTAGTT	258	NM_015147	916-917	KIAA0582
GATCCTAGGGGTTTTG	259	NM_015147	916-917	KIAA0582
GATCTAAGTTGCCTACC	260	NM_014176	918-919	HSPC 150 protein similar to ubiquitin conjugating enzyme
GATCTTTGTTCTTGACC	261	NM_020531	920-921	chromosome 20 open reading frame 3

TABLE 3-continued

DIFFERENTIALLY EXPRESSED GENES THAT ENCODE PREDICTED SECRETED PROTEINS.				
Signature	SEQ ID Accession NO: Number	SEQ ID NOS:	Description	
GATCTCTTAGCCAGAGG	262 NM_153333	922-923	transcription elongation factor A (SII)-like 8	
GATCTCTCTCACCTACA	263 NM_003287	924-925	tumor protein D52-like 1	
GATCAGAGGTGAAGGGA	264 NM_007021	926-927	chromosome 10 open reading frame 10	
GATCTCATTGATGTACA	265 NM_032947	928-929	putative small membrane protein NID67	
GATCTGTGCCGGCTTCC	266 NM_005656	930-931	transmembrane protease, serine 2	
GATCCGTCTGTGCACAT	267 NM_005656	930-931	transmembrane protease, serine 2	
GATCGCTCTGGGAGAC	268 NM_006315	932-933	ring finger protein 3	
GATCGATTAATGAAGTG	269 NM_016326	934-935	chemokine-like factor	
GATCCTGGACTGGGTAC	270 NM_006830	936-937	ubiquinol-cytochrome c reductase (6.4kD) subunit	
GATCTTGAGAATGTGA	271 NM_001216	938-939	carbonic anhydrase IX	
GATGTTTTTTTGGATAT	272 NM_181762	940-941	ubiquitin-conjugating enzyme E2A (RAD6 homolog)	
GATCTAGTTATTTTAAT	273 NM_014050	942-943	mitochondrial ribosomal protein L42	
GATCTAGTTATTTTAAT	274 NM_172177	944-945	mitochondrial ribosomal protein L42	
GATCAAGGGACGGCTGA	275 NM_000978	946-947	ribosomal protein L23	
GATCAGAAGGCTCTGGT	276 NM_018442	948-949	IQ motif and WD repeats 1	
GATCAATGTTGAAGAAT	277 NM_018442	948-949	IQ motif and WD repeats 1	
GATCCTGCACTCTAACA	278 NM_203339	950-951	clusterin (complement lysis inhibitor, SP-40,40, sulfated glycoprotein 2, testosterone-repressed prostate message 2, apolipoprotein J)	
GATCTGATTATTTACTT	279 NM_004708	952-953	programmed cell death 5	
GATCCTTGAAGGCAGCT	280 NM_197958	954-955	acheron	
GATCCCTTTTCTTACTA	281 NM_153713	956-957	hypothetical protein MGC46719	
GATCTGTCCACTTCTGG	282 NM_153713	956-957	hypothetical protein MGC46719	

TABLE 3-continued

DIFFERENTIALLY EXPRESSED GENES THAT ENCODE PREDICTED SECRETED PROTEINS.				
Signature	SEQ ID NO:	Accession Number	SEQ ID NOS:	Description
GATCAGATACCACCAAG	283	NM_001001503	958-959	NADH dehydrogenase (ubiquinone) flavoprotein 3, 10kDa
GATCCTTTGGATTAATC	284	NM_016138	960-961	coenzyme Q7 homolog, ubiquinone (yeast)
GATCATTATTTCTGTCT	285	NM_018184	962-963	ADP-ribosylation factor-like 10C
GATCAGCCCTCAAAGAA	286	NM_018184	962-963	ADP-ribosylation factor-like 10C
GATCAGCAAAAATAAAG	287	NM_016096	964-965	HSPC038 protein
GATCTCAGCGGCATTAA	288	NM_052951	966-967	deoxynucleotidy ltransferase, terminal, interacting protein 1
GATCCCTGGAGTGCCTT	289	NM_003226	968-969	trefoil factor 3 (intestinal)
GATCTGTTTCTACCAAT	290	NM_183045	970-971	ring finger protein (C3H2C3 type) 6
GATCCTGGTGTGAAAGG	291	NM_153750	972-973	chromosome 21 open reading frame 81
GATCTTGAAAGTGCCTG	292	NM_022130	974-975	golgi phosphoprotein 3 (coat-protein)
GATCAATAGAATAACAA	293	NM_003479	976-977	protein tyrosine phosphatase type IVA, member 2
GATCTCCTATGAGAACA	294	NM_003479	976-977	protein tyrosine phosphatase type IVA, member 2
GATCAATACAATAACAA	295	NM_080391	978-979	protein tyrosine phosphatase type IVA, member 2
GATCTCCTATGAGAACA	296	NM_080391	978-979	protein tyrosine phosphatase type IVA, member 2
GATCCAACCCTGTACTG	297	NM_177969	980-981	protein phosphatase 18 (formerly 2C), magnesium- dependent, beta isoform
GATCTCTACCATTTAAT	298	NM_001017	982-983	ribosomal protein S13
GATCCAGAAATACTTAA	299	NM_005410	984-985	selenoprotein P, plasma, 1
GATCCAATGCCTAAACTC	300	NM_005410	984-985	selenoprotein P, plasma, 1
GATCAAATGAGAATAAAA	301	NM_182620	986-987	family with sequence similarity 33, member A
GATCCTTGCCACAAGAA	302	NM_004034	988-989	annexin A7
GATCAGACTGTATTAAA	303	NM_032051	990-991	zinc finger protein 278

TABLE 3-continued

DIFFERENTIALLY EXPRESSED GENES THAT ENCODE PREDICTED SECRETED PROTEINS.			
Signature	SEQ ID Accession NO: Number	SEQ ID NOS: Description	
GATCTCCCACTGCGGGG	304 NM_032051	990-991 zinc finger protein 278	
GATCGGCCCTACTAGAT	305 NM_032051	990-991 zinc finger protein 278	
GATCAAAAAGCAAGCAG	306 NM_015972	992-993 polymerase (RNA) 1 polypeptide D, 16kDa	
GATCACTTCAGCTGCCT	307 NM_019007	994-995 armadillo repeat containing, X-linked 6	
GATCACCGACTGAAAAT	308 NM_002165	996-997 inhibitor of DNA binding 1, dominant negative helix-loop helix rotein	
GATCAATGAAGTGAGAA	309 NM_003094	998-999 small nuclear ribonucleoprotein polypeptide E	
GATCATCTCAGAAGTCT	310 NM_018683	1000-1001 zinc finger protein 313	
GATCAGGAAGGACTTGT	311 NM_018683	1000-1001 zinc finger protein 313	
GATCATTCCATTTTCAT	312 NM_002583	1002-1003 PRKC, apoptosis, WT1, regulator	
GATCGCTTTCTACTG	313 NM_006926	1004-1005 surfactant, pulmonary-associated protein A2	
GATCAGTTAGCTTTTAT	314 NM_014335	1006-1007 CREBBP/EP300 inhibitor 1	
GATCAGTAGTTCAACAG	315 NM_175061	1008-1009 juxtaposed with another zinc finger gene 1	
GATCCGATAAGTTATTG	316 NM_004707	1010-1011 APG12 autophagy 12-like (<i>S. cerevisiae</i>)	
GATCAGTGGGCACAGTT	317 NM_006818	1012-1013 ALL1-fused gene from chromosome 1q	
GATCAGTGCCAGAAGTC	318 NM_016303	1014-1015 WW domain binding protein 5	
GATCAGAGAAGTAAGTT	319 NM_004871	1016-1017 golgi SNAP receptor complex member 1	
GATCTCACTTTCCCTT	320 NM_015373	1018-1019 PKD2 interactor, golgi and endoplasmic reticulum associated 1	
GATCAGGCAGTTCCTGG	321 NM_213720	1020-1021 chromosome 22 open reading frame 16	
GATCCTTGCCACAAGAA	322 NM_001156	1022-1023 annexin A7	
GATCAAGAAAATAAGG	323 NM_000999	1024-1025 ribosomal protein L38	
GATCGATTCTTTCTC	324 NM_021102	1026-1027 serine protease inhibitor, Kunitz type, 2	

TABLE 3-continued

DIFFERENTIALLY EXPRESSED GENES THAT ENCODE PREDICTED SECRETED PROTEINS.			
Signature	SEQ ID Accession NO: Number	SEQ ID NOS: Description	
GATCATAGAAGGCTATT	325 NM_181826	1028-1029 neurofibromin 2 (bilateral acoustic neuroma)	
GATCCGGTGCGCCATGT	326 NM_002638	1030-1031 protease inhibitor 3, skin-derived (SKALP)	
GATCGCAGTTTGGAAAC	327 NM_005461	1032-1033 v-maf musculoaponeurotic fibrosarcoma oncogene homolog B (avian)	
GATCAATTTCAAACCTT	328 NM_005461	1032-1033 v-maf musculoaponeurotic fibrosarcoma oncogene homolog B (avian)	
GATCTCCTATGAGAACA	329 NM_080392	1034-1035 protein tyrosine phosphatase type IVA, member 2	
GATCAATACAATAACAA	330 NM_080392	1034-1035 protein tyrosine phosphatase type IVA, member 2	
GATGCTACCACCTACTG	331 NM_018281	1036-1037 hypothetical protein FLJ10948	
GATCATTGTTTATTAA	332 NM_004763	1038-1039 integrin beta 1 binding protein 1	
GATCAAAATGCTAAAC	333 NM_178191	1040-1041 ATPase inhibitory factor 1	
GATCTGGGGTGGGAGTA	334 NM_002773	1042-1043 protease, serine, 8 (prostasin)	
GATCATGCTTGTGTGAG	335 NM_018648	1044-1045 nucleolar protein family A, member 3 (H/ACA small nucleolar RNPs)	
GATCAAATATGTAAAAT	336 NM_138633	1046-1047 A kinase (PRKA) anchor protein 7	
GATCAGACTTCTCAGCT	337 NM_006856	1048-1049 activating transcription factor 7	
GATCATAGAAGGCTATT	338 NM_181827	1050-1051 neurofibromin 2 (bilateral acoustic neuroma)	
GATCCACCTTGGCCTCC	339 NM_003842	1052-1053 tumor necrosis factor receptor superfamily, member 10b	
GATCTCTGGCCCTCAG	340 NM_198527	1054-1055 Similar to RIKEN cDNA 1110033009 gene	
GATCCTCATTGAGCCAC	341 NM_024866	1056-1057 adrenomedullin 2	
GATCCAGTGGGGTCGGG	342 NM_002475	1058-1059 myosin light chain 1 slow a	
GATCATTTGTATTAAT	343 NM_017544	1060-1061 NF-kappa B repressing factor	
GATCAGAAAAAGAAAGA	344 NM_000982	1062-1063 ribosomal protein L21	

TABLE 3-continued

DIFFERENTIALLY EXPRESSED GENES THAT ENCODE PREDICTED SECRETED PROTEINS.			
Signature	SEQ ID Accession NO: Number	SEQ ID NOS: Description	
GATCCTGTTCTCTGTCAC	345 NM_203413	1064-1065 S-phase 2 protein	
GATCATGGTTCTCTTTG	346 NM_000202	1066-1067 iduronate 2-sulfatase (Hunter syndrome)	
GATCCTCTGACCGCTGG	347 NM_022365	1068-1069 DnaJ (Hsp40) homolog, subfamily C, member 1	
GATCTGCTATATGCCAGC	348 NM_016399	1070-1071 hypothetical protein HSPC132	
GATCCTGGAAATGTCAG	349 NM_001233	1072-1073 caveolin 2	
GATCAGTCTCAAGTGTC	350 NM_003702	1074-1075 regulator of G- protein signalling 20	
GATCAGTTAGCAAATG	351 NM_004331	1076-1077 BCL2/adenovirus E1B 19kDa interacting protein 3- like	
GATCAGTATGCTGTTTT	352 NM_004968	1078-1079 islet cell autoantigen 1,69kDa	
GATCTGGTTTCTAGCAA	353 NM_024096	1080-1081 XTP3-transactivated protein A	
GATCTAATTAATAAAT	354 NM_000903	1082-1083 NAD(P)H dehydrogenase, quinone 1	
GATCCTGGGTTTTTGTG	355 NM_017830	1084-1085 OCIA domain containing 1	
GATCACCGACTGAAAT	356 NM_181353	1086-1087 inhibitor of DNA binding 1, dominant negative helix-loop helixprotein	
GATCAGGTAACCGAGC	357 NM_002488	1088-1089 NADH dehydrogenase (ubiquinone) 1 alpha subcomplex, 2, 8kDa	
GATCAGTGAACACTAAC	358 NM_016645	1090-1091 mesenchymal stem cell protein DSC92	
GATCTCAGATGCTAGAA	359 NM_016567	1092-1093 BRCA2 and CDKN1A interacting protein	
GATCGCTCTGCCATGT	360 NM_016567	1092-1093 BRCA2 and CDKN1A interacting protein	
GATCAGTCCGTGGGGC	361 NM_152398	1094-1095 OCIA domain containing 2	
GATCATGCCCCAAAGTT	362 NM_152398	1094-1095 OCIA domain containin 2	
GATCTGGCACTGTGGTT	363 NM_000998	1096-1097 ribosomal protein L37a	
GATCTGGCACTGTGGGT	364 NM_000998	1096-1097 ribosomal protein L37a	
GATCTCAGATGCTAGAA	365 NM_078468	1098-1099 BRCA2 and CDKN1A interacting	

TABLE 3-continued

DIFFERENTIALLY EXPRESSED GENES THAT ENCODE PREDICTED SECRETED PROTEINS.			
Signature	SEQ ID Accession NO: Number	SEQ ID NOS: Description	
GATCGCTCTGCCCATGT	366 NM_078468	1098-1099 protein BRCA2 and CDKN1A interacting protein	
GATCTGCTGTGGAATTG	367 NM_172316	1100-1101 Meis1, myeloid ecotropic viral integration site 1 homolog 2 (mouse)	
GATCGTTCTTGATTTTG	368 NM_032476	1102-1103 mitochondrial ribosomal protein S6	
GATCTTGGTTTCATGTG	369 NM_032476	1102-1103 mitochondrial ribosomal protein S6	
GATCATTCTTGATTTTG	370 NM_032476	1102-1103 mitochondrial ribosomal protein S6	
GATCCATATGGAAGAA	371 NM_014171	1104-1105 postsynaptic protein CRIPT	
GATCTGCCCCACTGTC	372 NM_138929	1106-1107 diablo homolog (<i>Drosophila</i>)	
GATCGCCTAGTATGTTT	373 NM_052815	1108-1109 immediate early response 3	
GATCAATGCTAATATGA	374 NM_005805	1110-1111 proteasome (prosome, macropain) 26S subunit, non-ATPase, 14	
GATCAGCATCAGGCTGT	375 NM_012459	1112-1113 translocase of inner mitochondrial membrane 8 homolog B (yeast)	
GATCTGGAAGTGAAACA	376 NM_134265	1114-1115 WD repeat and SOCS box-containing 1	
GATCCACGTGTGAGGGA	377 NM_182640	1116-1117 mitochondrial ribosomal protein S9	
GATCACAGAAAAATTAA	378 NM_182640	1116-1117 mitochondrial ribosomal protein S9	
GATCTCTCTGCGTTTGA	379 NM_012445	1118-1119 spondin 2, extracellular matrix protein	
GATCTCAGAAGTTTGA	380 NM_138459	1120-1121 chromosome 6 open reading frame 68	
GATCCGGACTTTTAAA	381 NM_006339	1122-1123 high-mobility group 20B	
GATCATAGTTATTATAC	382 NM_032467	1124-1125 aspartate beta- hydroxylase	
GATCCTGCCCTGCTCTC	383 NM_003145	1126-1127 signal sequence receptor, beta (translocon- associated protein beta)	
GATCGATTGAGAAGTTA	384 NM_012110	1128-1129 cysteine-rich hydrophobic domain 2	

TABLE 3-continued

DIFFERENTIALLY EXPRESSED GENES THAT ENCODE PREDICTED SECRETED PROTEINS.			
Signature	SEQ ID Accession NO: Number	SEQ ID NOS: Description	
GATCCAAGTACTCTCTC	385 NM_175081	1130-1131 purinergic receptor P2X, ligand-gated ion channel, 5	
GATCATAACCTGCTCA	386 NM_001009	1132-1133 ribosomal protein S5	
GATCCTGGATGCCACGA	387 NM_174889	1134-1135 hypothetical protein LOC91942	
GATCCCTGCCACAAGTT	388 NM_006923	1136-1137 stromal cell-derived factor 2	
GATCAGACGAGGCCATG	389 NM_006107	1138-1139 cisplatin resistance- associated overexpressed protein	
GATCTTTCAGGAAAGAC	390 NM_033011	1140-1141 plasminogen activator, tissue	
GATCTTTTAAAAATATA	391 NM_001914	1142-1143 cytochrome b-5	
GATCGTTTTGTTTTGTT	392 NM_021149	1144-1145 coactosin-like 1 (<i>Dictyostelium</i>)	
GATCTATGGCCTCTGGT	393 NM_021643	1146-1147 tribbles homolog 2 (<i>Drosophila</i>)	
GATCCTAAATCATTTTG	394 NM_022783	1148-1149 DEP domain containing 6	
GATCTAAGAAGAACTA	395 NM_005765	1150-1151 ATPase, H+ transporting, lysosomal accessory protein 2	
GATCTTGGTGTCAAAA	396 NM_001497	1152-1153 UDP- Gal:betaGlcNAc beta 1,4- galactosyltransferase, polypeptide 1	
GATCCCTCATCCCTGAA	397 NM_006793	1154-1155 peroxiredoxin 3	
GATCTGCAGTGCTTCAC	398 NM_178181	1156-1157 CUB domain- containing protein 1	
GATCTATGCCCTTGTTA	399 NM_033167	1158-1159 UDP- Gal:betaGlcNAc beta 1,3- galactosyltransferase, polypeptide 3	
GATCTATGCCCTTGTTA	400 NM_033169	1160-1161 UDP- Gal:betaGlcNAc beta 1,3- galactosyltransferase, polypeptide 3	
GATCAGTTTATTATTGA	401 NM_033169	1160-1161 UDP Gal:betaGlcNAc beta 1,3- galactosyltransferase, polypeptide 3	
GATCTATGCCCTTGTTA	402 NM_033168	1162-1163 UDP- Gal:betaGlcNAc beta 1,3- galactosyltransferase, polypeptide 3	

TABLE 3-continued

DIFFERENTIALLY EXPRESSED GENES THAT ENCODE PREDICTED SECRETED PROTEINS.				
Signature	SEQ ID Accession		SEQ ID	
	NO:	Number	NOS:	Description
GATCAGTTTATTATTGA	403	NM_033167	1158-1159	UDP-Gal:betaGlcNAc beta 1,3-galactosyltransferase, polypeptide 3
GATCTATGCCCTTGTTA	404	NM_003781	1164-1165	UDP-Gal:betaGlcNAc beta 1,3-galactosyltransferase, polypeptide 3
GATCAGTTTATTATTGA	405	NM_003781	1164-1165	UDP-Gal:betaGlcNAc beta 1,3-galactosyltransferase, polypeptide 3
GATCAGTTTATTATTGA	406	NM_033168	1162-1163	UDP-Gal:betaGlcNAc beta 1,3-galactosyltransferase, polypeptide 3
GATCGAGTCAAGATGAG	407	NM_013442	1166-1167	stomatin (EPB72)-like 2
GATCACCATGATGCAGA	408	NM_031905	1168-1169	SVH protein
GATCCCGTGTGTGTGTG	409	NM_031905	1168-1169	SVH protein
GATCATGGTTCTCTTTG	410	NM_006123	1170-1171	iduronate 2-sulfatase (Hunter syndrome)
GATCCGCAGGCAGAAGC	411	NM_002775	1172-1173	protease, serine, 11 (IGF binding)
GATCGATGGTGGCGCTT	412	NM_138991	1174-1175	beta-site APP-cleaving enzyme 2
GATCTGCATCAGTTGTA	413	NM_001814	1176-1177	cathepsin C
GATCTCTACTACCACAA	414	NM_001908	1178-1179	cathepsin B
GATCTCTACTACCACAA	415	NM_147780	1180-1181	cathepsin B
GATCTCTACTACCACAA	416	NM_147781	1182-1183	cathepsin B
GATCTCTACTACCACAA	417	NM_147782	1184-1185	cathepsin B
GATCTCTACTACCACAA	418	NM_147783	1186-1187	cathepsin B
GATCGATGGTGGCGCTT	419	NM_012105	1188-1189	beta-site APP-cleaving enzyme 2
GATCTTTCAGGAAAGAC	420	NM_000931	1190-1191	plasminogen activator, tissue
GATCAAATGCAAATA	421	NM_153705	1192-1193	KDEL (Lys-Asp Glu-Leu) containing 2
GATCTTATTTTCTGAGA	422	NM_014584	1194-1195	ERO 1-like (<i>S. cerevisiae</i>)
GATCCACAAGGCCTGAG	423	NM_001185	1196-1197	alpha-2-glycoprotein 1, zinc
GATCTAGGCCTCATCTT	424	NM_016352	1198-1199	carboxypeptidase A4
GATCCCTTTGAAATTTT	425	NM_001219	1200-1201	calumenin

TABLE 3-continued

DIFFERENTIALLY EXPRESSED GENES THAT ENCODE PREDICTED SECRETED PROTEINS.			
Signature	SEQ ID NO: Number	Accession	SEQ ID NOS: Description
GATCTACAACATATAAA	426	NM_020648	1202-1203 twisted gastrulation homolog 1 (<i>Drosophila</i>)
GATCAGTTTTTTCACCT	427	NM_001901	1204-1205 connective tissue growth factor
GATCAGAGTGTGAGAGA	428	NM_007224	1206-1207 neurexophilin 4
GATCGTTACTATGTGTC	429	NM_004541	1208-1209 NADH dehydrogenase (ubiquinone) 1 alpha subcomplex, 1, 7.5kDa
GATCATTGACCTCTGTG	430	NM_006459	1210-1211 SPFH domain family, member 1
GATCTGAAGCCCAGGTT	431	NM_024514	1212-1213 cytochrome P450, family 2, subfamily R, polypeptide 1
GATCTGTTAAAAA	432	NM_147159	1214-1215 opioid receptor, sigma 1
GATCTTTCAGGAAAGAC	433	NM_000930	1216-1217 plasminogen activator, tissue
GATCATAAGACAATGGA	434	NM_001657	1218-1219 amphiregulin (schwannoma- derived growth factor)
GATCAGTCTTTATTAAT	435	NM_013995	1220-1221 lysosomal-associated membrane protein 2
GATCCAGGGTCACTGTG	436	NM_005250	1222-1223 forkhead box L1
GATCAAATAATGCGACG	437	NM_018064	1224-1225 chromosome 6 open reading frame 166
GATCTTGGTTTTCCATG	438	NM_003000	1226-1227 succinate dehydrogenase complex, subunit B, iron sulfur (Ip)
GATCTGTTAGTCAAGTG	439	NM_005313	1228-1229 glucose regulated protein, 58kDa
GATCATTTCTGGTAAAT	440	NM_005313	1228-1229 glucose regulated protein, 58kDa
GATCAAAGCACTCTTCC	441	NM_005313	1228-1229 glucose regulated protein, 58kDa
GATCATGCCAAGTGGTG	442	NM_021233	1230-1231 deoxyribonuclease II beta
GATCATGCCTCCCTGG	443	NM_006216	1232-1233 serine (or cysteine) proteinase inhibitor, clade E (nexin, plasminogen activator inhibitor type 1), member 2
GATCACCAGGCTGCCCA	444	NM_016504	1234-1235 mitochondrial ribosomal protein L27
GATCGGATGGGCAAGTC	445	NM_002178	1236-1237 insulin-like growth factor binding protein 6

TABLE 3-continued

DIFFERENTIALLY EXPRESSED GENES THAT ENCODE PREDICTED SECRETED PROTEINS.			
Signature	SEQ ID Accession NO: Number	SEQ ID NOS: Description	
GATCTCAAGACCAAAGA	446 NM_030810	1238-1239 thioredoxin domain containing 5	
GATCTCACATTGTGCC	447 NM_014254	1240-1241 transmembrane protein 5	
GATCAGTCTTTATTAAT	448 NM_002294	1242-1243 lysosomal-associated membrane protein 2	
GATCAGAGAAGATGATA	449 NM_000640	1244-1245 interleukin 13 receptor, alpha 2	
GATCAGGTAACCAGAGC	450 NM_000591	1246-1247 CD14 antigen	
GATCATCAGTAAATTTG	451 NM_031284	1248-1249 ADP-dependent glucokinase	
GATCAATAAATGTGAT	452 NM_002658	1250-1251 plasminogen activator, urokinase	
GATCCCTCGGGTTTTGT	453 NM_006350	1252-1253 follistatin	
GATCTTGCAACTCCATT	454 NM_006350	1252-1253 follistatin	
GATCCAGCATGGAGGCC	455 NM_018664	1254-1255 Jun dimerization protein p21SNFT	
GATCATTGTGAAGGCAG	456 NM_001511	1256-1257 chemokine (C-X-C motif) ligand 1 (melanoma growth stimulating activity, alpha)	
GATCTGCCAGCAGTGTT	457 NM_002004	1258-1259 farnesyl diphosphatesynthase (farnesylpyrophosphate synthetase, dimethylallyltranstransferase, geranyltranstransferase)	
GATCAGAGGTTACTAGG	458 NM_006408	1260-1261 anterior gradient 2 homolog (<i>Xenopus laevis</i>)	
GATCCACAGGGTGGTG	459 NM_000602	1262-1263 serine (or cysteine) proteinase inhibitor, clade (nexin, plasminogen activator inhibitor type 1), member 1	
GATCACAGGGGGGAT	460 NM_016588	1264-1265 neuritin 1	
GATCTGTGTTTGGACTA	461 NM_004109	1266-1267 ferredoxin 1	
GATCTAACCTGGCTTGT	462 NM_004109	1266-1267 ferredoxin 1	
GATCAGCAAGTGTCCTT	463 NM_000935	1268-1269 procollagen-lysine, 2- oxoglutarate 5- dioxygenase 2	
GATCTAGTGGTTCACAC	464 NM_003236	1270-1271 transforming growth factor, alpha	
GATCAAACAGTTTCTGG	465 NM_016139	1272-1273 coiled-coil-helix- coiled-coil-helix domain containing 2	
GATCATCAAGAAAAAAG	466 NM_018464	1274-1275 chromosome 10 open reading frame 70	
GATCCCAGAGAGCAGCT	467 NM_002421	1276-1277 matrix metalloproteinase 1 (interstitial collagenase)	

TABLE 3-continued

DIFFERENTIALLY EXPRESSED GENES THAT ENCODE PREDICTED SECRETED PROTEINS.				
Signature	SEQ ID NO:	Accession Number	SEQ ID NOS:	Description
GATCTTGTGTATTTTTG	468	NM_020440	1278-1279	prostaglandin F2 receptor negative regulator
GATCTATGTTCTCTCAG	469	NM_013363	1280-1281	procollagen C- endopeptidase enhancer 2
GATCAGCAAGTGCCCTT	470	NM_182943	1282-1283	procollagen-lysine, 2- oxoglutarate 5- dioxygenase 2
GATCATGTGCTACTGGT	471	NM_003172	1284-1285	surfeit
GATCTGTAATAAAATC	472	NM_130781	1286-1287	RAB24, member RAS oncogene family
GATCAGGGCTGAGGGTA	473	NM_000157	1288-1289	glucosidase, beta; acid (includes glucosylceramidase)
GATCCTCCTATGTTGTT	474	NM_005551	1290-1291	kallikrein 2, prostatic
GATCAGAGATGCACCAC	475	NM_002997	1292-1293	syndecan 1
GATCTGTCTGTTGCTTG	476	NM_005570	1294-1295	lectin, mannose- binding, 1
GATCACCATGAAAGAAG	477	NM_003873	1296-1297	neuropilin 1
GATCTGTATAAAAAA	478	NM_005866	1298-1299	opioid receptor, sigma 1
GATCAATCCCTTGAAT	479	NM_138322	1300-1301	proprotein convertase subtilisin/kexin type 6
GATCCCAGACCAACCCT	480	NM_024642	1302-1303	UDP-N-acetyl-alpha galactosamine:polypeptide N-acetylgalactosaminyltransferase 12 (GalNAc-T 12)
GATCATCACAGTTTGAG	481	NM_002425	1304-1305	matrix metalloproteinase 10 (stromelysin 2)
GATCGAACAGCTCCTT	482	NM_178154	1306-1307	fucosyltransferase 8 (alpha (1,6) fucosyltransferase)
GATCGAACAGCTCCTT	483	NM_178155	1308-1309	fucosyltransferase 8 (alpha (1,6) fucosyltransferase)
GATCGAACAGCTCCTT	484	NM_178156	1310-1311	fucosyltransferase 8 (alpha (1,6) fucosyltransferase)
GATCTGTGGGCCAGTC	485	NM_004077	1312-1313	citrate synthase
GATCAACCTTAAAGGAA	486	NM_000143	1314-1315	fumarate hydratase
GATCTTCTACTTGCCCTG	487	NM_000302	1316-1317	procollagen-lysine 1, 2-oxoglutarate 5- dioxygenase 1
GATCACCAGCCATGTGC	488	NM_004390	1318-1319	cathepsin H
GATCACCGAGGTCAGT	489	NM_016026	1320-1321	retinol dehydrogenase 11 (all-trans and 9-cis)

TABLE 3-continued

DIFFERENTIALLY EXPRESSED GENES THAT ENCODE PREDICTED SECRETED PROTEINS.			
Signature	SEQ ID Accession NO: Number	SEQ ID NOS: Description	
GATCTATTTTATGCATG	490 NM_020792	1322-1323 KIAA1363 protein	
GATCTGTTAAAAA	491 NM_147157	1324-1325 opioid receptor, sigma 1	
GATCATTTTGGTTCGTG	492 NM_016417	1326-1327 chromosome 14 open reading frame 87	
GATCACTTGTGTACGAA	493 NM_024641	1328-1329 mannosidase, endo- alpha	
GATCCCTCCACCCCAT	494 NM_001441	1330-1331 fatty acid amide hydrolase	
GATCCAAAGTCATGTGT	495 NM_058172	1332-1333 anthrax toxin receptor 2	
GATCCATAAATATTTAT	496 NM_058172	1332-1333 anthrax toxin receptor2	
GATCTGCCTGCATCCTG	497 NM_003225	1334-1335 trefoil factor 1 (breast cancer, estrogen- inducible sequence expressed in)	
GATCCAGTGTCCATGGA	498 NM_007085	1336-1337 follistatin-like 1	
GATCAATTCCTTGAAT	499 NM_138324	1338-1339 proprotein convertase subtilisin/kexin type 6	
GATCCGTGTGCTTGGGC	500 NM_018143	1340-1341 keich-like 11 (<i>Drosophila</i>)	
GATCCAGGTCCCCCAG	501 NM_004911	1342-1343 protein disulfide isomerase related protein (calcium- binding protein, intestinal-related)	
GATCATGGGACCTCTC	502 NM_003032	1344-1345 sialyltransferase 1 (beta-galactoside alpha-2,6- sialyltransferase)	
GATCATGGGACCTCTC	503 NM_173216	1346-1347 sialyltransferase 1 (beta-galactoside alpha-2,6- sialyltransferase)	
GATCTCACTGTTATTAT	504 NM_007115	1348-1349 tumor necrosis factor, alpha-induced protein 6	
GATCCTGTATCCAAATC	505 NM_007115	1348-1349 tumor necrosis factor, alpha-induced protein 6	
GATCAGTTTCTCTTAA	506 NM_024769	1350-1351 adipocyte-specific adhesion molecule	
GATCTACCAGATAACCT	507 NM_000522	1352-1353 homeo box A13	
GATCCTAGTAATTGCCT	508 NM_054034	1354-1355 fibronectin 1	
GATCAATGCAACGACGT	509 NM_006833	1355-1356 COP9 constitutive photomorphogenic homolog subunit 6 (<i>Arabidopsis</i>)	
GATCAATTCCTTGAAT	510 NM_138325	1358-1359 proprotein convertase subtilisin/kexin type 6	

TABLE 3-continued

DIFFERENTIALLY EXPRESSED GENES THAT ENCODE PREDICTED SECRETED PROTEINS.			
Signature	SEQ ID NO: Number	Accession Number	SEQ ID NOS: Description
GATCAATCCCTTGAAT	511	NM_138323	1360-1361 proprotein convertase subtilisin/kexin type 6
GATCCAGAGGGATGCA	512	NM_024040	1362-1363 CUE domain containing 2
GATCATCAAAAATGCTA	513	NM_017898	1364-1365 hypothetical protein FLJ20605
GATCCCTCGGTTTTGT	514	NM_013409	1366-1367 follistatin
GATCTTGCAACTCCATT	515	NM_013409	1366-1367 follistatin
GATCTTGTTAATGCATT	516	NM_001873	1368-1369 carboxypeptidase E
GATCAAAGGTTTAAAGT	517	NM_001627	1370-1371 activated leukocyte cell adhesion molecule
GATCACCAAGATGCTTC	518	NM_018371	1372-1373 chondroitin beta 1,4 N-acetylgalactosaminyltransferase
GATCAAATGTGCCTTAA	519	NM_014918	1374-1375 carbohydrate (chondroitin) synthase 1
GATCTTCGGCCTCATT	520	NM_017860	1376-1377 hypothetical protein FLJ20519
GATCCCTTCTGCCCTGG	521	NM_022367	1378-1379 sema domain, immunoglobulin domain (Ig), transmembrane domain (TM) and short cytoplasmic domain, (semaphorin) 4A
GATCCAACCGACTGAAT	522	NM_006670	1380-1381 trophoblast glycoprotein
GATCTCTGCAGATCCA	523	NM_004750	1382-1383 cytokine receptor-like factor 1
GATCACAAAATGTTGCC	524	NM_001077	1384-1385 UDP- glycosyltransferase 2 family, polypeptide B17
GATCTCTCTTTCTCTCT	525	NM_031882	1386-1387 protocadherin alpha subfamily C, 1
GATCTCTCTTTCTCTCT	526	NM_031860	1388-1389 protocadherin alpha 10
GATCTCTCTTTCTCTCT	527	NM_018906	1390-1391 protocadherin alpha 3
GATCTCTCTTTCTCTCT	528	NM_031411	1392-1393 protocadherin alpha 1
GATCACAGGCGTGAGCT	529	NM_032620	1394-1395 GTP binding protein 3 (mitochondrial)
GATCAACATCTTTCTTT	530	NM_004343	1396-1397 calreticulin
GATCTCTGATTTAACCG	531	NM_002185	1398-1399 interleukin 7 receptor
GATCTCTCTTTCTCTCT	532	NM_031497	1400-1401 protocadherin alpha 3
GATCCATTTTAAATGGT	533	NM_198278	1402-1403 hypothetical protein LOC255743
GATCTTTTCTAAATGTT	534	NM_005699	1404-1405 interleukin 18 binding protein

TABLE 3-continued

DIFFERENTIALLY EXPRESSED GENES THAT ENCODE PREDICTED SECRETED PROTEINS.			
Signature	SEQ ID Accession NO: Number	SEQ ID NOS: Description	
GATCTCTCTTTCTCTCT	535 NM_031410	1406-1407	protocadherin alpha I
GATCGGTGCGTTCTCCT	536 NM_005561	1408-1409	lysosomal-associated membrane protein 1
GATCTTTTCTAAATGTT	537 NM_173042	1410-1411	interleukin 18 binding protein
GATCTTTTCTAAATGTT	538 NM_173043	1412-1413	interleukin 18 binding protein
GATCTCTCTTTCTCTCT	539 NM_031496	1414-1415	protocadherin alpha 2
GATCCTGTTGGATGTGA	540 NM_080927	1416-1417	discoidin, CUB and LCCL domain containing 2
GATCTCTCTTTCTCTCT	541 NM_031864	1418-1419	protocadherin alpha 12
GATCTCTCTTTCTCTCT	542 NM_031849	1420-1421	protocadherin alpha 6
GATCCTGTGCTTCTGCA	543 NM_006464	1422-1423	trans-golgi network protein 2
GATCTCTCTTTCTCTCT	544 NM_031865	1424-1425	protocadherin alpha 13
GATCTGATGAAGTATAT	545 NM_022746	1426-1427	hypothetical protein FLJ22390
GATCACTTGCTTGTGG	546 NM_006988	1428-1429	a disintegrin-like and metalloprotease (reprolysin type) with thrombospondin type 1 motif, 1
GATCTTTTCTAAATGTT	547 NM_173044	1430-1431	interleukin 18 binding protein
GATCTCTCTTTCTCTCT	548 NM_031856	1432-1433	protocadherin alpha 8
GATCTCTCTTTCTCTCT	549 NM_031500	1434-1435	protocadherin alpha 4
GATCAGCACTGCCAGTG	550 NM_016592	1436-1437	GNAS complex locus
GATCCGAAAGATGAAT	551 NM_144640	1438-1439	interleukin 17 receptor E
GATCTCTCTTTCTCTCT	552 NM_031501	1440-1441	protocadherin alpha 5
GATCTCTCTTTCTCTCT	553 NM_031495	1442-1443	protocadherin alpha 2
GATCTAATGTAAAATCC	554 NM_002354	1444-1445	tumor-associated calcium signal transducer 1
GATCTTCTTTTGTAAATG	555 NM_032780	1446-1447	transmembrane protein 25
GATCAATAAATGAGG	556 NM_001001390	1448-1449	CD44 antigen (homing function and Indian blood group system)
GATCAACAGTGGCAATG	557 NM_001001390	1448-1449	CD44 antigen (homing function and Indian blood group system)

TABLE 3-continued

DIFFERENTIALLY EXPRESSED GENES THAT ENCODE PREDICTED SECRETED PROTEINS.			
Signature	SEQ ID NO: Number	Accession	SEQ ID NOS: Description
GATCAACAGTGGCAATG	558	NM_001001391	1450-14551CD44 antigen (homing function and Indian blood group system)
GATCAATAATAATGAGG	559	NM_001001391	1450-1451CD44 antigen (homing function and Indian blood group system)
GATCATTGCTCCTTCTC	560	NM_004872	1452-1453chromosome 1 open reading frame 8
GATCTCTGCATTTTATA	561	NM_020198	1454-1455GK001 protein
GATCTATGAAATCTGTG	562	NM_020198	1454-1455GK001 protein
GATCTCTCTTCTCTCT	563	NM_018901	1456-1457protocadherin alpha 10
GATCACTGGAGCTGTGG	564	NM_002116	1458-1459major histocompatibility complex, class I, A
GATCATCCAGTTTGCTT	565	NM_004540	1460-1461neural cell adhesion molecule 2
GATCAAAATGTTACCC	566	NM_004540	1460-1461neural cell adhesion molecule 2
GATCAACAGTGGCAATG	567	NM_001001389	1462-1463CD44 antigen (homing function and Indian blood group system)
GATCAATAATAATGAGG	568	NM_001001389	1462-1463CD44 antigen (homing function and Indian blood group system)
GATCAACAGTGGCAATG	569	NM_000610	1464-1465CD44 antigen (homing function and Indian blood group system)
GATCAATAATAATGAGG	570	NM_000610	1464-1465CD44 antigen (homing function and Indian blood group system)
GATCCATACTGTTTGGA	571	NM_001792	1466-1467cadherin 2, type 1, N- cadherin (neuronal)
GATCTGCATTTTCAGAA	572	NM_015544	1468-1469DKFZP564K1964 protein
GATCCATTTTTTGTA	573	NM_000574	1470-1471decay accelerating factor for complement (CD55, Cromer blood group system)
GATCTGCAGTGCTTAC	574	NM_022842	1472-1473CUB domain- containing protein 1
GATCTGTTAAAAAAA	575	NM_147160	1474-1475opioid receptor, sigma 1
GATCATAGGCTTGACA	576	NM_014045	1476-1477low density lipoprotein receptor- related protein 10

TABLE 3-continued

DIFFERENTIALLY EXPRESSED GENES THAT ENCODE PREDICTED SECRETED PROTEINS.			
Signature	SEQ ID Accession NO: Number	SEQ ID NOS: Description	
GATCTAATACTACTGTC	577 NM_001110	1478-1479 a disintegrin and metalloproteinase domain 10	
GATCTCTTGAGGCTGGG	578 NM_016371	1480-1481 hydroxysteroid (17-beta) dehydrogenase 7	
GATCGTTCATTGCCTTT	579 NM_001746	1482-1483 calnexin	
GATCTCTCTTTCTCTCT	580 NM_018900	1484-1485 protocadherin alpha 1	
GATCTGACCTGGTGAGA	581 NM_004393	1486-1487 dystroglycan 1 (dystrophin-associated glycoprotein 1)	
GATCATCTTTCCTGTTT	582 NM_002117	1488-1489 major histocompatibility complex, class I, C	
GATCGTAAAATTTTAAG	583 NM_003816	1490-1491 a disintegrin and metalloproteinase domain 9 (meltrin gamma)	
GATCTCTCTTTCTCTCT	584 NM_018904	1492-1493 protocadherin alpha 13	
GATCTCTCTTTCTCTCT	585 NM_018911	1494-1495 protocadherin alpha 8	
GATCTCTCITTCTCTCT	586 NM_018905	1496-1497 protocadherin alpha 2	
GATCTCTCTTTCTCTCT	587 NM_018903	1498-1499 protocadherin alpha 12	
GATCTCTCTTTCTCTCT	588 NM_018907	1500-1501 protocadherin alpha 4	
GATCTCTCTTTCTCTCT	589 NM_018908	1502-1503 protocadherin alpha 5	
GATCCGAAAGATGAAT	590 NM_153480	1504-1505 interleukin 17 receptor E	
GATCCGAAAGATGAAT	591 NM_153483	1506-1507 interleukin 17 receptor E	
GATCTCTGTAATTTTAT	592 NM_021923	1508-1509 fibroblast growth factor receptor-like 1	
GATCTAAGAGATTAATA	593 NM_004362	1510-1511 calmegin	

Example 2

Identification of Secreted Proteins By Computational Analysis of MPSS Signature Sequences

Secreted proteins can readily be exploited for blood cancer diagnosis and prognosis. As such, the differentially expressed genes identified in Example 1 were further analyzed to determine how many of the differentially expressed genes encode secreted proteins. Proteins with signal peptides (classical secretory proteins) were predicted using the same criteria described by Chen et al., *Mamm Genome*, 14: 859-865, 2003, with the SignalP 3.0 server developed by The Center for Biological Sequence Analysis, Lyngby, Denmark (<http://www.cbs.dtu.dk/services/SignalP-3.0/>; see also, J. D. Bendtsen, et al., *J. Mol. Biol.*, 340:783-795, 2004.) and the TMHMM2.0 server (see for example A.

50

Krogh, et al., *Journal of Molecular Biology*, 305(3):567-580, January 2001; E. L. L. Sonnhammer, et al., In J. Glasgow, T. Littlejohn, F. Major, R. Lathrop, D. Sankoff, and C. Sensen, editors, *Proceedings of the Sixth International Conference on Intelligent Systems for Molecular Biology*, pages 175-182, Menlo Park, Calif., 1998. AAAI Press). Putatively nonclassical secretory secreted proteins (without signal peptides) were predicted based on the SecretomeP 1.0 server, (<http://www.cbs.dtu.dk/services/SecretomeP-1.0/>) and required an odds ratio score >3.

55

60

65

Five hundred and twenty one signatures belonging to 460 genes potentially encoding secreted proteins (Table 3) were identified. Among these, 287 (259 genes) and 234 (201 genes) signatures were overexpressed or underexpressed in CL1 cells compared with LNCaP cells. Thus these proteins can be used in blood diagnostics to follow prostate cancer progres-

sion. Additionally, these proteins can be used in other settings, such as for identifying drug side effects.

Example 3

Prostate Cancer Diagnostics Using Multiparameter Analysis

This example describes a multiparameter diagnostic fingerprint using the WDR19 prostate-specific secreted protein in combination with PSA. The WDR19 prostate-specific protein is diagnostically superior to PSA when used alone and further improved prostate cancer detection when used in combination with PSA.

WDR19 was previously identified as relatively tissue-specific by cDNA array studies and Northern blot analysis (see e.g., U.S. Patent Application Publication No. 20020150893). This protein was selected, expressed as protein, purified and antibodies were made against it, all using standard techniques known in the art (the cDNA encoding the WDR19 protein is provided in SEQ ID NO:1, the amino acid sequence is provided in SEQ ID NO:2). The WDR19-specific antibody was shown to be an excellent tissue-specific marker of prostate cancer with staining of the specific epithelial cells being directly proportional to the progression of the cancer. In this regard it is very different from the well-established PSA marker which is not a good prostate tissue cancer marker.

The WDR19 antibodies and those for the well-established PSA prostate cancer blood marker were used to analyze 10 blood samples from normal individuals, 10 blood samples from early prostate cancer patients and 10 blood samples from late prostate cancer patients. The results showed that WDR19 reacted against no normals, against 5/10 early cancers, and against 5/10 late cancers, whereas PSA reacted against no normals, no early cancers and 7/10 late cancers. The two markers together detected all the late cancers. Thus the multiparameter analysis of blood markers (e.g. the analyses of multiple markers) for prostate cancer was far more powerful than using each marker alone.

Accordingly, the results show a molecular blood fingerprint that comprises the WDR19 and PSA proteins. This fingerprint allows superior diagnostic power to PSA alone and further improves prostate cancer detection.

WDR19 was also shown to be an effective histochemical marker for prostate cancer. Two hundred and seventy-five tissue cores that contain both stromal and epithelial cells from cancer patients, 17 from benign prostatic hyperplasia (BPH) and 12 from normal individuals were examined. The mean WDR19 protein staining intensities were 2.52 [standard error (S.E.), 0.05; 95% confidence interval (CI), 2.41-2.61] for prostate cancer; 1.03 BPH (S.E. 0.03; 95% CI, 0.96-1.09); and 1.0 (S.E., 0, 95% CI 1.0-1.0) for normal individuals. Pair-wise comparisons (using independent t-test) demonstrated that WDR19 staining intensity is significantly different between prostate cancer and BPH (mean difference 1.49; $P < 0.0001$) and between prostate cancers and normal (mean difference 1.52; $P < 0.0001$). These data suggested that WDR19, in addition to being a prostate-specific blood biomarker, is a quantitative cancer-specific marker for prostate tissues.

Example 4

Identification of Organ-Specific Secreted Proteins Using MPSS and Computational Analysis

MPSS as described in Example 1 and in the detailed description, was used to identify more than 2 million transcripts from each of the prostate cell lines (see Example 1) and in normal prostate tissue. The MPSS signature sequences from normal prostate were compared against 29 other tissues each with about 1 million or more mRNA transcripts. This comparison revealed that about 300 of these transcripts are organ-specific and about 60 of these organ-specific transcripts are potentially secreted into the blood. (See Table 4).

TABLE 4

PROSTATE-SPECIFIC PROTEINS POTENTIALLY SECRETED INTO BLOOD		
Accession No.	SEQ ID NO:	Annotations/Description
NP_001176	1512	alpha-2-glycoprotein 1, zinc; Alpha-2-glycoprotein, zinc [<i>Homo sapiens</i>]
NP_001719	1513	basigin isoform 1; OK blood group; collagenase stimulatory factor; M6 antigen; extracellular matrix metalloproteinase inducer [<i>Homo sapiens</i>]
NP_940991	1514	basigin isoform 2; OK blood group; collagenase stimulatory factor; M6 antigen; extracellular matrix metalloproteinase inducer [<i>Homo sapiens</i>]
NP_004039	1515	beta-2-microglobulin precursor [<i>Homo sapiens</i>]
NP_002434	1516	beta-microseminoprotein isoform a precursor; seminal plasma beta-inhibin; prostate secreted seminal plasma protein; immunoglobulin binding factor; prostatic secretory protein 94 [<i>Homo sapiens</i>]
NP_619540	1517	beta-microseminoprotein isoform b precursor; seminal plasma beta-inhibin; prostate secreted seminal plasma protein; immunoglobulin binding factor; prostatic secretory protein 94 [<i>Homo sapiens</i>]
NP_817089	1518	cadherin-like 26 isoform a; cadherin-like protein VR20 [<i>Homo sapiens</i>]
NP_068582	1519	cadherin-like 26 isoform b; cadherin-like protein VR20 [<i>Homo sapiens</i>]
NP_001864	1520	carboxypeptidase E precursor [<i>Homo sapiens</i>]
NP_004807	1521	chromosome 9 open reading frame 61; Friedreich ataxia region gene X123 [<i>Homo sapiens</i>]

TABLE 4-continued

PROSTATE-SPECIFIC PROTEINS POTENTIALLY SECRETED INTO BLOOD		
Accession No.	SEQ ID NO:	Annotations/Description
NP_001271	1522	cold inducible RNA binding protein; Cold-inducible RNA-binding protein; cold inducible RNA-binding protein; glycine-rich RNA binding protein [<i>Homo sapiens</i>]
NP_008977	1523	elastin microfibril interfacier 1; TNF? elastin microfibril interface located protein; elastin microfibril interface located protein [<i>Homo sapiens</i>]
NP_004104	1524	fibroblast growth factor 12 isoform 2; fibroblast growth factor 12B; fibroblast growth factor homologous factor 1; myocyte-activating factor; fibroblast growth factor FGF-12b [<i>Homo sapiens</i>]
NP_005962	1525	FXYP domain containing ion transport regulator 3 isoform 1 precursor; phospholemman-like protein; FXYP domain-containing ion transport regulator 3 [<i>Homo sapiens</i>]
NP_068710	1526	FXYP domain containing ion transport regulator 3 isoform 2 precursor; phospholemman-like protein; FXYP domain-containing ion transport regulator 3 [<i>Homo sapiens</i>]
NP_006352	1527	homeo box B13; homeobox protein HOX-B13 [<i>Homo sapiens</i>]
NP_002139	1528	homeo box D10; homeobox protein Hox-D10; homeo box 4D; Hox-4
NP_000513	1529	homeobox protein A13; homeobox protein HOXA13; homeo box 1J; transcription factor HOXA13 [<i>Homo sapiens</i>]
NP_060819	1530	hypothetical protein FLJ11175 [<i>Homo sapiens</i>]
NP_078985	1531	hypothetical protein FLJ14146 [<i>Homo sapiens</i>]
NP_061894	1532	hypothetical protein FLJ20010 [<i>Homo sapiens</i>]
NP_115617	1533	hypothetical protein FLJ23544; QM gene; DNA segment on chromosome X (unique) 648 expressed sequence; 60S ribosomal protein L10; tumor suppressor QM; Wilms tumor-related protein; laminin receptor homolog [<i>Homo sapiens</i>]
NP_057582	1534	hypothetical protein HSPC242 [<i>Homo sapiens</i>]
NP_116285	1535	hypothetical protein MGC14388 [<i>Homo sapiens</i>]
NP_116293	1536	hypothetical protein MGC14433 [<i>Homo sapiens</i>]
NP_077020	1537	hypothetical protein MGC4309 [<i>Homo sapiens</i>]
NP_061074	1538	hypothetical protein PRO1741 [<i>Homo sapiens</i>]
NP_563614	1539	hypothetical protein similar to KIAA0187 gene product [<i>Homo sapiens</i>]
NP_951038	1540	I-mfa domain-containing protein isoform p40 [<i>Homo sapiens</i>]
NP_005542	1541	kallikrein 2, prostatic isoform 1; glandular kallikrein 2 [<i>Homo sapiens</i>]
NP_004908	1542	kallikrein 4 preproprotein; protease, serine, 17; enamel matrix serine protease 1; kallikrein-like protein 1; protase; androgen-regulated message 1 [<i>Homo sapiens</i>]
NP_002328	1543	low density lipoprotein receptor-related protein associated protein 1; lipoprotein receptor associated protein; alpha-2-MRAP; alpha-2-macroglobulin receptor-associated protein 1; low density lipoprotein-related protein-associated protein 1; low density li
NP_859077	1544	low density lipoprotein receptor-related protein binding protein [<i>Homo sapiens</i>]
NP_000897	1545	natriuretic peptide receptor A/guanylate cyclase A (atriuretic peptide receptor A); Natriuretic peptide receptor A/guanylate cyclase A [<i>Homo sapiens</i>]
NP_085048	1546	Nedd4 family interacting protein 1; Nedd4 WW domain-binding protein 5 [<i>Homo sapiens</i>]
NP_000896	1547	neuropeptide Y [<i>Homo sapiens</i>]
NP_039227	1548	olfactory receptor, family 10, subfamily H, member 2 [<i>Homo sapiens</i>]
NP_000599	1549	orosomucoid 2; alpha-1-acid glycoprotein, type 2 [<i>Homo sapiens</i>]
NP_002643	1550	prolactin-induced protein; prolactin-inducible protein [<i>Homo sapiens</i>]
NP_057674	1551	prostate androgen-regulated transcript 1 protein; prostate-specific and androgen-regulated cDNA 14D7 protein [<i>Homo sapiens</i>]
NP_001639	1552	prostate specific antigen isoform 1 preproprotein; gamma-seminoprotein; semenogelase; seminin; P-30 antigen [<i>Homo sapiens</i>]

TABLE 4-continued

PROSTATE-SPECIFIC PROTEINS POTENTIALLY SECRETED INTO BLOOD		
Accession No.	SEQ ID NO:	Annotations/Description
NP_665863	1553	prostate specific antigen isoform 2; gamma-seminoprotein; semenogelase; seminolisin; P-30 antigen [<i>Homo sapiens</i>]
NP_001090	1554	prostatic acid phosphatase precursor [<i>Homo sapiens</i>]
NP_001000	1555	ribosomal protein S5; 40S ribosomal protein S5 [<i>Homo sapiens</i>]
NP_005658	1556	ring finger protein 103; Zinc finger protein expressed in cerebellum; zinc finger protein 103 homolog (mouse) [<i>Homo sapiens</i>]
NP_937761	1557	ring finger protein 138 isoform 2 [<i>Homo sapiens</i>]
NP_002998	1558	semenogelin I isoform a preproprotein [<i>Homo sapiens</i>]
NP_937782	1559	semenogelin I isoform b preproprotein [<i>Homo sapiens</i>]
XP_353669	1560	similar to HIC protein isoform p32 [<i>Homo sapiens</i>]
NP_003855	1561	sin3 associated polypeptide p30 [<i>Homo sapiens</i>]
NP_036581	1562	six transmembrane epithelial antigen of the prostate; six transmembrane epithelial antigen of the prostate (NOTE: non-standard symbol and name) [<i>Homo sapiens</i>]
NP_008868	1563	SMT3 suppressor of mif two 3 homolog 2; SMT3 (suppressor of mif two 3, yeast) homolog 2 [<i>Homo sapiens</i>]
NP_066568	1564	solute carrier family 15 (H+/peptide transporter), member 2 [<i>Homo sapiens</i>]
NP_055394	1565	solute carrier family 39 (zinc transporter), member 2 [<i>Homo sapiens</i>]
NP_003209	1566	telomeric repeat binding factor 1 isoform 2; Telomeric repeat binding factor 1; telomeric repeat binding protein 1 [<i>Homo sapiens</i>]
NP_110437	1567	thioredoxin domain containing 5 isoform 1; thioredoxin related protein; endothelial protein disulphide isomerase [<i>Homo sapiens</i>]
NP_004863	1568	thymic dendritic cell-derived factor 1; liver membrane-bound protein [<i>Homo sapiens</i>]
NP_665694	1569	TNF receptor-associated factor 4 isoform 2; tumor necrosis receptor-associated factor 4A; malignant 62; cysteine-rich domain associated with ring and TRAF domain [<i>Homo sapiens</i>]
NP_005647	1570	transmembrane protease, serine 2; epitheliasin [<i>Homo sapiens</i>]
NP_008931	1571	uropod 1A [<i>Homo sapiens</i>]
NP_036609	1572	WW domain binding protein 1 [<i>Homo sapiens</i>]
NP_009062	1573	zinc finger protein 75 [<i>Homo sapiens</i>]

Example 5

Comparison of Localized Prostate Cancer and Prostate Cancer Metastases in the Liver

In an additional experiment, the transcriptome from normal prostate tissue was compared to the transcriptome of each of the LNCaP and CL-1 prostate cancer cell lines. The comparison showed that the transcriptomes were distinct for the normal tissue, the early prostate cancer and the late prostate cancer. An additional comparison was carried out between localized prostate cancer and metastases in the liver. About 6,000 genes were identified that were significantly changed between the localized prostate cancer and the metastasized cancer and again, many of the changed genes encoded secreted proteins that can be part of the blood fingerprints indicative of the more advanced disease status of metastases. The metastases-altered blood fingerprints may indicate the site of metastases.

These experiments demonstrate that there are continuous changes in the two types of networks as prostate cancer progresses—from localized to androgen independence to metastases. These graded network transitions suggest that one will be able to detect the very earliest stages of prostate cancer and, accordingly, that the organ-specific, molecular

blood fingerprints approach described herein will permit a very early diagnosis of prostate and other types of cancers. These experiments further support the notion that the drug-related side effects that impact the prostate can be identified and monitored using the organ-specific, molecular blood fingerprints.

Example 6

MPSS Analysis in a Yeast Model System

This experiment demonstrates perturbation-specific fingerprints of patterns of gene expression for nuclear, cytoplasmic, membrane-bound and secreted proteins in the yeast metabolic system that converts the sugar galactose into glucose-6-phosphate (the gal system).

The gal systems includes 9 proteins. In the course of studying how this systems works, 9 new strains of yeast were created, each with a different one of the 9 relevant genes destroyed (gene knockouts). Yeast is a single celled eukaryote organism with about 6,000 genes. The expression patterns of each of the 6,000 genes was studied in the wild type yeast and each of the 9 knockout strains. The data from these experiments showed: 1) the wild type and each of knock out strains exhibited statistically significant changes in patterns of gene

expression from the wild type strain ranging from 89 to 465 altered patterns of gene expression; 2) each of these patterns of changed gene expression were unique; and 3) on average about 15% of the genes with changed expression patterns encoded proteins that were potentially secreted (as determined by computational analysis from the sequence of the gene). These genes are as follows: (listed by gene name as available through the public yeast genome database at <http://colon.double slash www dot yeastgenome dot org/>. The genomic DNA, cDNA and amino acid sequences corresponding to each of the listed genes are publicly available, for example, through the yeast genome database.) YGL102C, YGL069C, YLL044W, YMR321C, YKL153W, YMR195W, YHL015W, YNL096C, YGR030C, YDR123C, YKL186C, YOR234C, YKL001C, YJL188C, YDL023C, YPL143W, YEL039C, YKL006W, YGR280C, YBR285W, YKR091W, YDR064W, YBR047W, YGR243W, YOR309C, YDR461W, YHR053C, YHR055C, YGR148C, YGL187C, YIL018W, YFR003C, YPL107W, YBR185C, YNR014W, YJL067W, YDR451C, YGL031C, YHR141C, YNL162W, YBR046C, YNL036W, YDL136W, YDL191W, YLR257W, YNL057W, YGL068W, YKR057W, YLR201C, YHL001W, YDR010C, YPL138C, YOR312C, YPL276W, YML114C, YLR327C, YBR191W, YOR257W, YOR096W, YPL223C, YJL136C, YAL044C, YER079W, YMR107W, YPL079W, YDR175C, YGR035C, YDR153C, YDR337W, YOR167C, YMR194W, YOR194C, YHR090C, YGR110W, YMR242C, YHR198C, YPL177C, YLR164W, YMR143W, YDL083C, YLR325C, YOR203W, YMR193W, YLR062C, YOR383C, YLR300W, YJL079C, YJL158C, YHR139C, YGL032C, YER150W, YNL160W, YDR382W, YMR305C, YKL096W, YKR013W, YCL043C, YLR042C, YDR055W, YPL163C, YEL040W, YJL171C, YLR121C, YOR382W, YLR250W, YGR189C, YJL159W, YMR215W, YDR519W, YIL162W, YKL163W, YDR518W, YDR534C, YPR157W, YML130C, YML128C, YBR092C, YDR032C, YLR120C, YBR093C, YHR215W, YAR071W, YDL130W, YDR144C, YPR123C, YGR174C, YOR327C, YNL058C, YGR265W, YGR160W, YIL117C, YOL053W, YGR236C, YGR060W, YKL120W, YDL046W, YHR132C, YMR058W, YLR332W, YKR061W, YEL001C,

YKL154W, YKL073W, YMR238W, YJR020W, YIL136W, YHL028W, YDL010W, YLR339C, YNL217W, YHR063C.

The different knockout strains can be thought of as analogous to genetic disease mutants. Accordingly, these data further support the notion that each disease has a unique expression fingerprint and that each disease generates unique collections of secreted proteins that constitute molecular fingerprints capable of identifying the corresponding disease.

Example 7

Identification of Prostate-Specific/Enriched Genes Using a 2.5 Fold Over-Expression Cut-Off

Organ specific/enriched expression can be determined by the ratio of the expression (e.g., measured in transcripts per million (tpm)) in a particular organ as compared to other organs. In this example, prostate enriched/specific expression was analyzed by comparing the expression level (tpm counts) of MPSS signature sequences identified from normal prostate tissue to their corresponding expression levels in 33 normal tissues. A particular gene that demonstrated at least a 2.5-fold increase in expression in prostate as compared to all tissues examined (each tissue evaluated individually) was considered to be prostate-specific/enriched. The tissues examined were adrenal gland, bladder, bone marrow, brain (amygdala, caudate nucleus, cerebellum, corpus callosum, hypothalamus, and thalamus), whole fetal brain, heart, kidney, liver (new cloning), lung, mammary gland, monocytes, peripheral blood lymphocytes, pituitary gland, placenta, pancreas, prostate, retina, spinal cord, salivary gland, small intestine, stomach, spleen, testis, thymus, trachea, thyroid, and uterus. This analysis identified 109 unique genes (with mpss signature sequence belonging to class 1-4, i.e. with confirmed match to cDNAs) whose expression was at least 2.5 fold that observed in other normal tissues. The list of prostate-specific/enriched genes is provided in Tables 5A-5D with the expression level in tpm in prostate shown. This list includes KLK2, KLK3, KLK4, TMPRSS2, which are genes previously shown to be prostate-specific.

TABLE 5A

PROSTATE ENRICHED GENES IDENTIFIED BY RATIO SCHEMA (RATIO > 2.5)*					
MPSS Signature	MPSS Sig. SEQ ID NO:	Genbank Name	Genbank SEQ Accession No.	Tissue Names	
				ID NOS:	Description
GATCTCAGAACACCTT	1688	DHRS7	BC000637	1797-1798	Dehydrogenase/reductase (SDR family) member 7
GATCCAGCCCAGAGACA	1689	NPY	BC029497	1799-1800	Neuropeptide Y
GATCACTCCTTATTTGC	1690	FLJ20010	AW172826	1801	Hypothetical protein FLJ20010
GATCCCTCTCCTCTCTG	1691	C9orf61	B1771919	1802	Chromosome 9 open reading frame 61
GATCTGACTTTTTACTT	1691	Lrp2bp	BU853306	1803	Ankyrin repeat domain 37
GATCGTTAGCCTCATAT	1693	HOXB13	BC007092	1804-1805	Homeo box B13
GATCACAAGGAATCCTG	169	CREB3L4	BC038962	1806-1807	CAMP responsive element binding protein 3-like 4
GATCTCATGGATGATTA	1695	LEPREL1	BC005029	1808-1809	Leprecan-like 1
GATCCAGAAATAAAGTC	1696	KLK4	CB051271	1810	Kallikrein 4 (prostase, enamel matrix, prostate)

TABLE 5A-continued

PROSTATE ENRICHED GENES IDENTIFIED BY RATIO SCHEMA (RATIO > 2.5)*					
MPSS Signature	MPSS Sig.	Genbank Name	Genbank SEQ Accession No.	Tissue Names	Description
	SEQ ID NO:			ID NOS:	
GATCTCACAGAAGATGT	1697	MGC35558	NM_145013	1811-1812	Chromosome 11 open reading frame 45
GATCCAAAATCACCAAG	1698	HAX1	BU157155	1813	HCLS1 associated protein X-1
GATCCTGGGCTGGAAGG	1699	0	AW207206	1814	Hypothetical gene supported by AY338954
GATCCAGATGCAGGACT	1700	0	BC013389	1815	LOC440156
GATCTGTGCTCATCTGT	1701	TMEM16G	BC028162	1816-1817	Transmembrane protein 16G
GATCATTTTATATCAAT	1702	MGC31963	X099160	1818	Chromosome 1 open reading frame 85
GATCCACACTGAGAGAG	1703	KLK3	BC005307	1819-1820	Kallikrein 3, (prostate specific antigen)
GATCCGCTGTGCACAT	1704	TMPRSS2	NM_005656	1821-1822	Transmembrane protease, serine 2
GATCATGTAGGGTAAC	1705	LOC221442	BC026923	1823	Hypothetical protein LOC221442
GATCAGCCCTCAAAAAA	1706	ARL10C	BU159800	1824	ADP-ribosylation factor-like 8B
GATCTGGATTGAGGACC	170	MGC13102	NM_032323	1825-1826	Hypothetical protein MGC13102
GATCAAAAATAAAATGT	1708	0	A1954252	1827	Hypothetical gene supported by AK022914; AK095211; BC016035; BC041856; BX248778
GATCCGCTCTGGTCAAC	1709	SEPX1	BQ941313	1828	Selenoprotein X, 1
GATCCCTCAAGACTGGT	1710	ACPP	BC007460	1829-1830	Acid phosphatase, prostate
GATCCACAAGACGAGG	1711	BIN3	B1911790	1831	Bridging integrator 3
GATCTCTCTGCGTTTGA	171	SPON2	BC002707	1832-1833	Spondin 2, extracellular matrix protein
GATCTCAACCTCGCTTG	1713	0	AK026938	1834	Hypothetical gene supported by AL713796
GATCAAGTTCCCGCTGG	1714	RPL18A	BG818587	1835	Ribosomal protein L18a
GATCATAATGAGGThITG	1715	ABCC4	NM_005845	1836-1837	ATP-binding cassette, sub-family C (CFTRIMRP), member 4
GATCGGTGACATCGTAA	1716	RPS11	A888242	1838	Ribosomal protein S11
GATCCACCAGCTGATAA	1717	SNEP1	CN353139	1839	Y box binding protein 1
GATCAACACTTTTATT	1718	FLJ22955	AA256381	1840	Hypothetical protein FLJ22955
GATCCCTTCCTITCCTCT	1719	HOXD11	AA513505	1841	Homeo box D11
GATCAGGACACAGACTT	1720	ORM1	BG564253	1842	Orosomuroid 1
GATCCTGCAATCTTGTA	1721	HTPAP	A1572087	1843	Phosphatidic acid phosphatase type 2 domain containin 1B
GATCCTCCTATGTTGTT	1722	KLK2	AA259243	1844	Kallikrein 2, prostatic
GATCTGTACCTTGCTA	1723	SLC2A12	A1675682	1845	Solute carrier family 2 (facilitated glucose transporter), member 12

TABLE 5A-continued

PROSTATE ENRICHED GENES IDENTIFIED BY RATIO SCHEMA (RATIO > 2.5)*					
MPSS Signature	MPSS Sig. SEQ ID NO:	Genbank Name	Genbank SEQ Accession No.	Tissue Names	
				ID NOs:	Description
GATCGGGGCAAGAGAGG	1724	NDRG1	NM_006096	1846-1847	N-myc downstream regulated gene 1
GATCCCCTCCCCTCCCC	1725	NPRI	NM_000906	1848-1849	Natriuretic peptide receptor A/guanylate cyclase A (atriuretic peptide receptor A)
GATCCTACAAAGAAG- GApotein FLJ21511	1726	FLJ21511	NM_025087	1850-1851	Hypothetica
GATCATTGCGAGTAAAG	1727	FOXA1	NM_004496	1852-1853	Forkhead box A1
GATCTGTCTCCTGCTCT	1728	ENPP3	A1535878	1854	Ectonucleotide pyrophosphatase/phosphodiesterase 3
GATCCTTCCCAAGGTAC	1729	GATA2	NM_032638	1855-1856	GATA binding protein 2
GATCTTGTGAAGTCAA	1730	ARG2	BX331427	1857	Arginase, type II
GATCGCACCACTGTACA	1731	XPO1	A1569484	1858	Exportin 1 (CRM1 homolog, yeast)
GATCATTTTCTGCTTTA	1732	ASB3	BC009569	1859-1860	Ankyrin repeat and SOCS box-containing 3
GATCCCACACTTGTC	1733	0	AK000028	1861	Hypothetical LOC90024
GATCTGGAATGTGATA	1734	KLF3	BX100634	1862	Kruppel-like factor 3 (basic)
GATCAATAAGCTTTAAA	1735	TGM4	BC007003	1863-1864	Transglutaminase 4 (prostate)
GATCAATGTTGTAGAT	1736	FLJ16231	NM_001008401	1865-1866	FLJ16231 protein
GATCTACATGTCTATCA	1737	BLNK	BX113323	1867	B-cell linker
GATCTGTTTTAAATGAG	1738	SLC14A1	NM_015865	1868-1869	Solute carrier family 14 (urea transporter), member 1 (Kidd blood group)
GATCAAAAAATGCTGCA	1739	PTPLB	A1017286	1870	Protein tyrosine phosphatase-like (proline instead of catalytic arginine), member b
GATCATGTCTTCATTTT	1740	OR51E2	NM_030774	1871-1872	Olfactory receptor, family 51, subfamily E, member 2
GATCCCTCCACCCCAT	1741	AAH	NM_001441	1873-1874	Fatty acid amide hydrolase
GATCCTAAGCCATAAAT	1742	STAT6	AL044554	1875	Signal transducer and activator of transcription 6, interleukin-4 induced
GATCATCGTCCTCATCG	1743	ANKH	CB049466	1876	Ankylosis, progressive homolog (mouse)
GATCATCATTGTGTCATT	1744	DSCR1L2	AW575747	1877	Down syndrome critical region gene 1-like 2
GATCTAATTTGAAAAAC	1745	TRPM8	NM_024080	1878-1879	Transient receptor potential cation channel, subfamily M, member 8
GATCTTCCTTGATCAT	1746	TMC4	AV724505	1880	Transmembrane channel-like 4
GATCTCCCCATGCCTG	1747	ZNF589	BC005859	1881-1882	Zinc finger protein 589
GATCAAATTTAGTATTT	1748	LRRK1	BC005408	1883-1884	Leucine-rich repeat kinase 1

TABLE 5A-continued

PROSTATE ENRICHED GENES IDENTIFIED BY RATIO SCHEMA (RATIO > 2.5)*					
MPSS Signature	MPSS Sig.	Genbank Name	Genbank SEQ Accession No.	Tissue Names	
	SEQ ID NO:			ID NOs:	Description
GATCTGCCTTATAACA	1749	STEAP2	AA177004	1885	Six transmembrane epithelial antigen of the prostate 2
GATCAGAAAATGAGCTC	1750	SAFB2	BC001216	1886	Scaffold attachment factor B2
GATCACCGTGGAGGTTA	1751	CPE	BG707154	1887	Carboxypeptidase E
GATCCCTCTGTGCTTCT	1752	GNB2L1	AA024878	1888	Guanine nucleotide binding protein (G protein), beta polypeptide 2-like 1
GATCTCATTTTTAGAGC	1753	LOC92689	BU688574	1889	Hypothetical protein BC001096
GATCATCACATTTTCGTG	1754	DLG1	BC042118	1890	Discs, large homolog 1 (<i>Drosophila</i>)
GATCATTTTCTGCTTCA	1755	SEMG1	NM) 003007	1891-1892	Semenogelin I
GATCAATGAAGGAGAGA	1756	SPATA 13	BM875598	1893	Spermatogenesis associated 13
GATCCCAACTACTCGGG	1757	LOC157657	NM_177965	1894-1895	Chromosome 8 open reading frame 37
GATCAGTTTTTCTGTAA	1758	KIAA1411	CA433208	1896	KIAA1411
GATCAAAATTTTAAAAA	1759	MGC20781	BM984931	1897	5'-nucleotidase, cytosolic III-like
GATCACCTTCTCTTCC	1760	LOC255189	BC035335	1898-1899	Phospholipase A2, group IVF
GATCCTGGGTACTGAAA	1761	ERBB2	BC080193	1900	V-erb-b2 erythroblastic leukemia viral oncogene homolog 2, neuro/glioblastoma derived oncogene homolog (avian)
GATCGTTCTAAGAGTGT	1762	ZFP64	NM_199427	1901-1902	Zinc finger protein 64 homolog (mouse)
GATCATCATCAAGGGCT	1763	SUHW2	BC042370	1903	Suppressor of hairy wing homolog 2 (<i>Drosophila</i>)
GATCAAAATGATTTTCA	1764	ELOVL7	AL137506	1904-1905	ELOVL family member 7, elongation of long chain fatty acids (yeast)
GATCTGATTTTTTTCCC	1765	TRAF4	AI888175	1906	TNF receptor-associated factor 4
GATCCCATTTCTCACCC	1766	SLC39A2	AI669751	1907	Solute carrier family 39 (zinc transporter, member 2)
GATCCTCCCGCCTTGCC	1767	HNF4G	AI088739	1908	Hepatocyte nuclear factor 4, gamma
GATCTTTCTTTTTTTGT	1768	SLC22A3	BC070300	1909	Solute carrier family 22 (extraneuronal monoamine transporter), member 3
GATCTTAACTGTCTCCT	1769	HIST2H2BE	BC005827	1910	Histone 2, H2be
GATCAGTTTGATTTCTGT	1770	AMD1	BC041345	1911-1912	Adenosylmethionine decarboxylase 1
GATCATGATGTAGAGGG	1771	TYMS	BX390036	1913	Thymidylate synthetase
GATCGCACCACTACAGT	1772	PHC3	AK022455	1914	Polyhomeotic like 3 (<i>Drosophila</i>)

TABLE 5A-continued

PROSTATE ENRICHED GENES IDENTIFIED BY RATIO SCHEMA (RATIO > 2.5)*					
MPSS Signature	MPSS Sig. SEQ ID NO:	Genbank Name	Genbank SEQ Accession No.	Tissue Names ID NOS:	Description
GATCTCAAAGTGCCTTC	1773	SARG	AL832940	1915-1916	Chromosome 1 open reading frame 116
GATCAATGTCAAACCTC	1774	MTERF	BC000965	1917-1918	Mitochondrial transcription termination factor
GATCTCCAGAGTCTAA	1775	CYP4F8	NM_007253	1919-1920	Cytochrome P450, family 4, subfamily F, polypeptide 8
GATCCTGATGGCTGTGT	1776	PPAP2A	AK124401	1921	Phosphatidic acid phosphatase type 2A
GATCACTTCCGCAGTC	1777	KIAA0056	BC011408	1922-1923	KIAA0056 protein
GATCTCAAAGGAACCAA	1778	MSMB	AA469293	1924	Microseminoprotein, beta-
GATCTGTGCCAGGGTTA	1779	VEGF	AK056914	1925	Vascular endothelial growth factor
GATCTCTTTTTATTAA	1780	CDH1	NM_004360	1926-1927	Cadherin 1, type 1, E-cadherin (epithelial)
GATCTCCAGCACCAATC	1781	TARP	BC062761	1928-1929	TCR gamma alternate reading frame protein
GATCTGGCGCTTGGGGG	1782	RFP2	NM_001007278	1930-1931	Ret finger protein 2
GATCCCGACGGGGGCAT	1783	MESP1	NM_018670	1932-1933	Mesoderm posterior 1 homolog (mouse)
GATCCCGGCGTATATC	1784	TRPM4	AA026974	1934	Transient receptor potential cation channel, subfamily M, member 4
GATCTTTCTCAAATAAT	1785	PAK11P1	AI468032	1935	PAKI interacting protein 1
GATCGTGACGCTTAATA	1786	HRPA1	CF122297	1936	Heterogeneous nuclear ribonucleoprotein A1
GATCGCATAATTTTTAA	1787	HNF207	CB053869	1937	Zinc finger protein 207
GATCCCAACTGAAGG	1788	WNK4	NM_032387	1938-1939	WNK lysine deficient protein kinase 4
GATCTTAAAACTGCAG	1789	APXL2	BQ448015	1940	Apical protein 2
GATCATTTTTTCTATCA	1790	MED28	AI554477	1941	Mediator of RNA polymerase II transcription, subunit 28 homolog (yeast)
GATCCCATTGTGTATAT	1791	LOC285300	AK095655	1942	Hypothetical protein LOC285300
GATCTCAAAGGAAAAAA	1792	0	AW291753	1943	Transcribed locus
GATCTTCTGTATATTTT	1793	0	BM023121	1944	Full length insert cDNA clone ZD79H10
GATCCACAACATACAGC	1794	0	AY338953	1945	Prostate-specific P712P mRNA sequence
GATCTGTGCAGTTGTAA	1795	0	AY533562	1946	KLK16 mRNA, partial sequence
GATCTACTATGCCAAAT	1796	0	BC030554	1947	(clone HGT25) T cell receptor gamma-chain mRNA, V region

*ratio of prostate expression in tpm to other organs greater than 2.5

TABLE 5B

PROSTATE ENRICHED GENES IDENTIFIED BY RATIO SCHEMA (RATIO >2.5)*				
Genbank Accession No.	Genbank SEQ ID NOs:	Name	SignalP3.0 prediction Prediction	SignalP3.0 prediction Signal peptide probability
BC000637	1797-1798	DHRS7	Signal peptide	0.999
BC029497	1799-1800	NPY	Signal peptide	0.998
AW172826	1801	FLJ20010	Non-secretory protein	0.001
BI771919	1802	C9orf61	Signal peptide	0.994
BU853306	1803	Lrp2bp	Non-secretory protein	0
BC007092	1804-1805	HOXB13	Non-secretory protein	0
BC038962	1806-1807	CREB3L4	Non-secretory protein	0
BC005029	1808-1809	LEPREL1	Signal peptide	0.995
CB051271	1810	KLK4	Signal peptide	0.988
NM_145013	1811-1812	MGC35558	Signal peptide	0.935
BU157155	1813	HAX1	Non-secretory protein	0.001
AW207206	1814	0	Non-secretory protein	0.001
BC013389	1815	0	Non-secretory protein	0
BC028162	1816-1817	TMEM16G	Non-secretory protein	0.001
BX099160	1818	MGC31963	Signal peptide	0.994
BC005307	1819-1820	KLK3	Signal peptide	0.992
NM_005656	1821-1822	TMPRSS2	Non-secretory protein	0
BC026923	1823	LOC221442	Signal anchor	0.01
BU159800	1824	ARL10C	Non-secretory protein	0
NM_032323	1825-1826	MGC13102	Non-secretory protein	0
AI954252	1827	0	Non-secretory protein	0.128
BQ941313	1828	SEPX1	Non-secretory protein	0
BC007460	1829-1830	ACPP	Signal peptide	1
BI911790	1831	BIN3	Non-secretory protein	0
BC002707	1832-1833	SPON2	Signal peptide	0.998
AK026938	1834	0	Signal peptide	0.587
BG818587	1835	RPL18A	Non-secretory protein	0
NM_005845	1836-1837	ABCC4	Non-secretory protein	0
AA888242	1838	RPS11	Non-secretory protein	0
CN353139	1839	NSEP1	Non-secretory protein	0.001
AA256381	1840	FLJ22955	Non-secretory protein	0.06
AA513505	1841	HOXD11	Non-secretory protein	0
BG564253	1842	ORM1	Signal peptide	1
AI572087	1843	HTPAP	Non-secretory protein	0.021
AA259243	1844	KLK2	Signal peptide	0.985
AI675682	1845	SLC2A12	Non-secretory protein	0
NM_006096	1846-1847	NDRG1	Non-secretory protein	0
NM_000906	1848-1849	NPR1	Signal peptide	0.997
NM_025087	1850-1851	FLJ21511	Non-secretory protein	0.005
NM_004496	1852-1853	FOXA1	Non-secretory protein	0
AI535878	1854	ENPP3	Non-secretory protein	0.069
NM_032638	1855-1856	GATA2	Non-secretory protein	0
BX331427	1857	ARG2	Non-secretory protein	0.014
AI569484	1858	XPO1	Non-secretory protein	0
BC009569	1859-1860	ASB3	Non-secretory protein	0
AK000028	1861	0	Non-secretory protein	0.001
BX100634	1862	KLF3	Non-secretory protein	0
BC007003	1863-1864	TGM4	Non-secretory protein	0
NM_001008401	1865-1866	FLJ16231	Non-secretory protein	0
BX113323	1867	BLNK	Non-secretory protein	0
NM_015865	1868-1869	SLC14A1	Non-secretory protein	0
AI017286	1870	PTPLB	Non-secretory protein	0.06
NM_030774	1871-1872	OR51E2	Non-secretory protein	0.008
NM_001441	1873-1874	FAAH	Signal peptide	0.805
AL044554	1875	STAT6	Non-secretory protein	0
CB049466	1876	ANKH	Non-secretory protein	0.001
AW575747	1877	DSCR1L2	Non-secretory protein	0
NM_024080	1878-1879	TRPM8	Non-secretory protein	0
AV724505	1880	TMC4	Non-secretory protein	0
BC005859	1881-1882	ZNF589	Non-secretory protein	0
BC005408	1883-1884	LRRK1	Non-secretory protein	0
AA177004	1885	STEAP2	Non-secretory protein	0
BC001216	1886	SAFB2	Non-secretory protein	0
BG707154	1887	CPE	Signal peptide	1
AA024878	1888	GNB2L1	Non-secretory protein	0
BU688574	1889	LOC92689	Non-secretory protein	0
BC042118	1890	DLG1	Non-secretory protein	0
NM_003007	1891-1892	SEMG1	Signal peptide	0.922
BM875598	1893	SPATA13	Non-secretory protein	0
NM_177965	1894-1895	LOC157657	Non-secretory protein	0
CA433208	1896	KIAA1411	Non-secretory protein	0

TABLE 5B-continued

PROSTATE ENRICHED GENES IDENTIFIED BY RATIO SCHEMA (RATIO >2.5)*				
Genbank Accession No.	Genbank SEQ ID NOs:	Name	SignalP3.0 prediction Prediction	SignalP3.0 prediction Signal peptide probability
BM984931	1897	MGC20781	Non-secretory protein	0
BC035335	1898-1899	LOC255189	Non-secretory protein	0
BC080193	1900	ERBB2	Non-secretory protein	0
NM_199427	1901-1902	ZFP64	Non-secretory protein	0
BC042370	1903	SUHW2	Non-secretory protein	0
AL137506	1904-1905	ELOVL7	Non-secretory protein	0
AI888175	1906	TRAF4	Non-secretory protein	0
AI669751	1907	SLC39A2	Signal peptide	0.982
AI088739	1908	HNF4G	Non-secretory protein	0.001
BC070300	1909	SLC22A3	Signal anchor	0.097
BC005827	1910	HIST2H2BE	Non-secretory protein	0
BC041345	1911-1912	AMD1	Non-secretory protein	0
BX390036	1913	TYMS	Non-secretory protein	0
AK022455	1914	PHC3	Non-secretory protein	0
AL832940	1915-1916	SARG	Non-secretory protein	0
BC000965	1917-1918	MTERF	Non-secretory protein	0
NM_007253	1919-1920	CYP4F8	Signal peptide	1
AK124401	1921	PPAP2A	Non-secretory protein	0.348
BC011408	1922-1923	KIAA0056	Non-secretory protein	0
AA469293	1924	MSMB	Signal peptide	0.997
AK056914	1925	VEGF	Non-secretory protein	0
NM_004360	1926-1927	CDH1	Signal peptide	0.896
BC062761	1928-1929	TARP	Non-secretory protein	0
NM_001007278	1930-1931	RFP2	Non-secretory protein	0
NM_018670	1932-1933	MESP1	Signal anchor	0.004
AA026974	1934	TRPM4	Non-secretory protein	0
AI468032	1935	PAK1IP1	Non-secretory protein	0.001
CF122297	1936	HNRPA1	Non-secretory protein	0
CB053869	1937	ZNF207	Non-secretory protein	0
NM_032387	1938-1939	WNK4	Non-secretory protein	0
BQ448015	1940	APXL2	Non-secretory protein	0
AI554477	1941	MED28		
AK095655	1942	LOC285300		
AW291753	1943	0		
BM023121	1944	0		
AY338953	1945	0		
AY533562	1946	0		
BC030554	1947	0		

*ratio of prostate expression in tpm to other organs greater than 2.5

TABLE 5C

PROSTATE ENRICHED GENES IDENTIFIED BY RATIO SCHEMA (RATIO > 2.5)*					
Genbank Accession No.	Genbank SEQ ID NOs:	name	SignalP3.0 prediction Max cleavage site probability	SecretomeP2.0 prediction Secreted potential (Odds)	TMHMM 2.0 prediction Pred trans- membrane domains
BC000637	1797-1798	DHRS7	0.599 between pos. 28 and 29	6.3	1
BC029497	1799-1800	NPY	0.520 between pos. 28 and 29	6.09	1
AW172826	1801	FLJ20010	0.000 between pos. 46 and 47	6.06	0
BI771919	1802	C9orf61	0.534 between pos. 29 and 30	5.9	2
BU853306	1803	Lrp2bp	0.000 between pos. 55 and 56	5.62	0
BC007092	1804-1805	HOXB13	0.000 between pos. -1 and 0	5.14	0
BC038962	1806-1807	CREB3L4	0.000 between pos. -1 and 0	4.72	0
BC005029	1808-1809	LEPREL1	0.991 between pos. 24 and 25	4.59	0
CB051271	1810	KLK4	0.401 between pos. 29 and 30	4.57	1

TABLE 5C-continued

PROSTATE ENRICHED GENES IDENTIFIED BY RATIO SCHEMA (RATIO > 2.5)*					
Genbank Accession No.	Genbank SEQ ID NOs:	name	SignalP3.0 prediction Max cleavage site probability	SecretomeP2.0 prediction Secreted potential (Odds)	TMHMM 2.0 prediction Pred trans- membrane domains
NM_145013	1811-1812	MGC35558	0.901 between pos. 22 and 23	4.47	0
BU157155	1813	HAX1	0.001 between pos. 18 and 19	4.41	0
AW207206	1814	0	0.001 between pos. 20 and 21	4.39	0
BC013389	1815	0	0.000 between pos. 27 and 28	4.3	0
BC028162	1816-1817	TMEM16G	0.001 between pos. 22 and 23	4.29	7
BX099160	1818	MGC31963	0.855 between pos. 35 and 36	4.22	2
BC005307	1819-1820	KLK3	0.525 between pos. 23 and 24	3.938	0
NM_005656	1821-1822	TMPRSS2	0.000 between pos. -1 and 0	3.86	1
BC026923	1823	LOC221442	0.004 between pos. 50 and 51	3.81	0
BU159800	1824	ARL10C	0.000 between pos. 35 and 36	3.76	0
NM_032323	1825-1826	MGC13102	0.000 between pos. -1 and 0	3.69	5
AI954252	1827	0	0.121 between pos. 42 and 43	3.58	0
BQ941313	1828	SEPX1	0.000 between pos. 13 and 14	3.49	0
BC007460	1829-1830	ACPP	0.975 between pos. 32 and 33	3.49	1
BI911790	1831	BIN3	0.000 between pos. -1 and 0	3.41	0
BC002707	1832-1833	SPON2	0.829 between pos. 26 and 27	3.06	0
AK026938	1834	0	0.568 between pos. 27 and 28	3.02	0
BG818587	1835	RPL18A	0.000 between pos. 24 and 25	2.8	0
NM_005845	1836-1837	ABCC4	0.000 between pos. -1 and 0	2.67	11
AA888242	1838	RPS11	0.000 between pos. -1 and 0	2.64	0
CN353139	1839	NSEP1	0.000 between pos. 25 and 26	2.35	0
AA256381	1840	FLJ22955	0.038 between pos. 15 and 16	2.19	1
AA513505	1841	HOXD11	0.000 between pos. 20 and 21	2.14	0
BG564253	1842	ORM1	0.923 between pos. 18 and 19	2.03	0
AI572087	1843	HTPAP	0.009 between pos. 63 and 64	2.01	4
AA259243	1844	KLK2	0.455 between pos. 17 and 18	1.81	0
AI675682	1845	SLC2A12	0.000 between pos. 51 and 52	1.79	12
NM_006096	1846-1847	NDRG1	0.000 between pos. -1 and 0	1.76	0
NM_000906	1848-1849	NPR1	0.960 between pos. 32 and 33	1.75	0
NM_025087	1850-1851	FLJ21511	0.005 between pos. 20 and 21	1.75	10
NM_004496	1852-1853	FOXA1	0.000 between pos. -1 and 0	1.71	0
AI535878	1854	ENPP3	0.036 between pos. 42 and 43	1.69	1
NM_032638	1855-1856	GATA2	0.000 between pos. 22 and 23	1.65	0
BX331427	1857	ARG2	0.013 between pos. 36 and 37	1.56	0
AI569484	1858	XPO1	0.000 between pos. -1 and 0	1.54	0

TABLE 5C-continued

PROSTATE ENRICHED GENES IDENTIFIED BY RATIO SCHEMA (RATIO > 2.5)*					
Genbank Accession No.	Genbank SEQ ID NOs:	name	SignalP3.0 prediction Max cleavage site probability	SecretomeP2.0 prediction Secreted potential (Odds)	TMHMM 2.0 prediction Pred trans- membrane domains
BC009569	1859-1860	ASB3	0.000 between pos. -1 and 0	1.53	0
AK000028	1861	0	0.000 between pos. 22 and 23	1.46	0
BX100634	1862	KLF3	0.000 between pos. -1 and 0	1.4	0
BC007003	1863-1864	TGM4	0.000 between pos. -1 and 0	1.36	0
NM_001008401	1865-1866	FLJ16231	0.000 between pos. -1 and 0	1.21	0
BX113323	1867	BLNK	0.000 between pos. -1 and 0	1.21	0
NM_015865	1868-1869	SLC14A1	0.000 between pos. -1 and 0	1.2	8
AI017286	1870	PTPLB	0.028 between pos. 63 and 64	1.2	4
NM_030774	1871-1872	OR51E2	0.003 between pos. 22 and 23	1.2	7
NM_001441	1873-1874	FAAH	0.549 between pos. 28 and 29	1.2	1
AL044554	1875	STAT6	0.000 between pos. -1 and 0	1.17	0
CB049466	1876	ANKH	0.000 between pos. 26 and 27	1.15	8
AW575747	1877	DSCR1L2	0.000 between pos. -1 and 0	1.12	0
NM_024080	1878-1879	TRPM8	0.000 between pos. -1 and 0	1.07	8
AV724505	1880	TMC4	0.000 between pos. -1 and 0	1.06	8
BC005859	1881-1882	ZNF589	0.000 between pos. -1 and 0	0.99	1
BC005408	1883-1884	LRRK1	0.000 between pos. -1 and 0	0.99	0
AA177004	1885	STEAP2	0.000 between pos. -1 and 0	0.95	6
BC001216	1886	SAFB2	0.000 between pos. -1 and 0	0.95	0
BG707154	1887	CPE	0.859 between pos. 27 and 28	0.93	0
AA024878	1888	GNB2L1	0.000 between pos. 33 and 34	0.92	0
BU688574	1889	LOC92689	0.000 between pos. -1 and 0	0.91	0
BC042118	1890	DLG1	0.000 between pos. -1 and 0	0.87	0
NM_003007	1891-1892	SEMG1	0.515 between pos. 23 and 24	0.85	0
BM875598	1893	SPATA13	0.000 between pos. -1 and 0	0.81	0
NM_177965	1894-1895	LOC157657	0.000 between pos. -1 and 0	0.81	0
CA433208	1896	KIAA1411	0.000 between pos. -1 and 0	0.8	0
BM984931	1897	MGC20781	0.000 between pos. 25 and 26	0.79	0
BC035335	1898-1899	LOC255189	0.000 between pos. 23 and 24	0.78	0
BC080193	1900	ERBB2	0.000 between pos. -1 and 0	0.74	2
NM_199427	1901-1902	ZFP64	0.000 between pos. -1 and 0	0.68	0
BC042370	1903	SUHW2	0.000 between pos. -1 and 0	0.67	0
AL137506	1904-1905	ELOVL7	0.000 between pos. -1 and 0	0.67	7
AI888175	1906	TRAF4	0.000 between pos. -1 and 0	0.63	0
AI669751	1907	SLC39A2	0.297 between pos. 23 and 24	0.62	8

TABLE 5C-continued

PROSTATE ENRICHED GENES IDENTIFIED BY RATIO SCHEMA (RATIO > 2.5)*					
Genbank Accession No.	Genbank SEQ ID NOs:	name	SignalP3.0 prediction Max cleavage site probability	SecretomeP2.0 prediction Secreted potential (Odds)	TMHMM 2.0 prediction Pred trans-membrane domains
AI088739	1908	HNFB4G	0.001 between pos. 21 and 22	0.59	0
BC070300	1909	SLC22A3	0.048 between pos. 33 and 34	0.58	12
BC005827	1910	HIST2H2BE	0.000 between pos. -1 and 0	0.58	0
BC041345	1911-1912	AMD1	0.000 between pos. -1 and 0	0.58	0
BX390036	1913	TYMS	0.000 between pos. -1 and 0	0.57	0
AK022455	1914	PHC3	0.000 between pos. -1 and 0	0.57	0
AL832940	1915-1916	SARG	0.000 between pos. 21 and 22	0.56	0
BC000965	1917-1918	MTERF	0.000 between pos. 14 and 15	0.56	0
NM_007253	1919-1920	CYP4F8	0.781 between pos. 36 and 37	0.56	1
AK124401	1921	PPAP2A	0.226 between pos. 30 and 31	0.53	5
BC011408	1922-1923	KIAA0056	0.000 between pos. -1 and 0	0.52	0
AA469293	1924	MSMB	0.928 between pos. 20 and 21	0.51	1
AK056914	1925	VEGF	0.000 between pos. -1 and 0	0.485	0
NM_004360	1926-1927	CDH1	0.487 between pos. 22 and 23	0.36	1
BC062761	1928-1929	TARP	0.000 between pos. 20 and 21	0.35	1
NM_001007278	1930-1931	RFP2	0.000 between pos. 24 and 25	0.32	1
NM_018670	1932-1933	MESP1	0.002 between pos. 20 and 21	0.31	0
AA026974	1934	TRPM4	0.000 between pos. -1 and 0	0.3	5
AI468032	1935	PAK1IP1	0.000 between pos. 25 and 26	0.27	0
CF122297	1936	HNRPA1	0.000 between pos. 32 and 33	0.22	0
CB053869	1937	ZNF207	0.000 between pos. -1 and 0	0.21	0
NM_032387	1938-1939	WNK4	0.000 between pos. -1 and 0	0.2	0
BQ448015	1940	APXL2	0.000 between pos. 41 and 42	0.19	0
AI554477	1941	MED28		#N/A	#N/A
AK095655	1942	LOC285300		#N/A	#N/A
AW291753	1943	0		#N/A	#N/A
BM023121	1944	0		#N/A	#N/A
AY338953	1945	0		#N/A	#N/A
AY533562	1946	0		#N/A	#N/A
BC030554	1947	0		#N/A	#N/A

*ratio of prostate expression in tpm to other organs greater than 2.5

TABLE 5D

PROSTATE ENRICHED GENES IDENTIFIED BY RATIO SCHEMA (RATIO > 2.5)*					
Genbank Accession No.	Genbank SEQ ID NOs:	name	NN-score	Odds	Prostate Expression (tmp)
BC000637	1797-1798	DHRS7	0.92	6.302	754
BC029497	1799-1800	NPY	0.911	6.099	642
AW172826	1801	FLJ20010	0.911	6.061	92

TABLE 5D-continued

PROSTATE ENRICHED GENES IDENTIFIED BY RATIO SCHEMA (RATIO > 2.5)*					
Genbank Accession No.	Genbank SEQ ID NOs:	name	NN-score	Odds	Prostate Expression (tmp)
BI771919	1802	C9orf61	0.906	5.902	91
BU853306	1803	Lrp2bp	0.895	5.626	95
BC007092	1804-1805	HOXB13	0.875	5.145	344

TABLE 5D-continued

PROSTATE ENRICHED GENES IDENTIFIED BY RATIO SCHEMA (RATIO > 2.5)*					
Genbank Accession No.	Genbank SEQ ID NOs:	name	NN-score	Odds	Prostate Expression (tmp)
BC038962	1806-1807	CREB3L4	0.866	4.721	334
BC005029	1808-1809	LEPREL1	0.857	4.594	118
CB051271	1810	KLK4	0.856	4.575	360
NM_145013	1811-1812	MGC35558	0.86	4.477	53
BU157155	1813	HAX1	0.854	4.412	67
AW207206	1814	0	0.854	4.391	279
BC013389	1815	0	0.85	4.304	64
BC028162	1816-1817	TMEM16G	0.843	4.293	281
BX099160	1818	MGC31963	0.846	4.222	53
BC005307	1819-1820	KLK3	0.838	3.938	24771
NM_005656	1821-1822	TMPRSS2	0.816	3.861	1425
BC026923	1823	LOC221442	0.8	3.812	104
BU159800	1824	ARL10C	0.822	3.76	167
NM_032323	1825-1826	MGC13102	0.788	3.699	238
AI954252	1827	0	0.814	3.589	159
BQ941313	1828	SEPX1	0.798	3.492	56
BC007460	1829-1830	ACPP	0.815	3.495	55
BI911790	1831	BIN3	0.806	3.41	54
BC002707	1832-1833	SPON2	0.766	3.063	873
AK026938	1834	0	0.769	3.025	304
BG818587	1835	RPL18A	0.768	2.806	58
NM_005845	1836-1837	ABCC4	0.747	2.671	454
AA888242	1838	RPS11	0.754	2.645	50
CN353139	1839	NSEP1	0.733	2.358	179
AA256381	1840	FLJ22955	0.688	2.196	57
AA513505	1841	HOXD11	0.715	2.142	99
BG564253	1842	ORM1	0.691	2.034	180
AI572087	1843	HTPAP	0.677	2.013	332
AA259243	1844	KLK2	0.676	1.816	7988
AI675682	1845	SLC2A12	0.499	1.792	127
NM_006096	1846-1847	NDRG1	0.667	1.765	2688
NM_000906	1848-1849	NPR1	0.658	1.755	150
NM_025087	1850-1851	FLJ21511	0.605	1.756	230
NM_004496	1852-1853	FOXA1	0.627	1.711	793
AI535878	1854	ENPP3	0.635	1.693	54
NM_032638	1855-1856	GATA2	0.598	1.659	238
BX331427	1857	ARG2	0.621	1.56	150
AI569484	1858	XPO1	0.604	1.54	68
BC009569	1859-1860	ASB3	0.607	1.538	2781
AK000028	1861	0	0.595	1.466	55
BX100634	1862	KLF3	0.581	1.401	136
BC007003	1863-1864	TGM4	0.59	1.368	5602
NM_001008401	1865-1866	FLJ16231	0.55	1.21	254
BX113323	1867	BLNK	0.559	1.211	183
NM_015865	1868-1869	SLC14A1	0.335	1.208	255
AI017286	1870	PTPLB	0.457	1.201	102
NM_030774	1871-1872	OR51E2	0.522	1.208	420
NM_001441	1873-1874	FAAH	0.535	1.206	476
AL044554	1875	STAT6	0.547	1.174	71
CB049466	1876	ANKH	0.335	1.153	58
AW575747	1877	DSCR1L2	0.471	1.123	225
NM_024080	1878-1879	TRPM8	0.519	1.077	267
AV724505	1880	TMC4	0.402	1.064	120
BC005859	1881-1882	ZNF589	0.491	0.992	156
BC005408	1883-1884	LRRK1	0.499	0.999	202
AA177004	1885	STEAP2	0.482	0.954	2156
BC001216	1886	SAFB2	0.427	0.954	76
BG707154	1887	CPE	0.464	0.933	148
AA024878	1888	GNB2L1	0.465	0.921	59
BU688574	1889	LOC92689	0.461	0.918	82
BC042118	1890	DLG1	0.457	0.872	50
NM_003007	1891-1892	SEMG1	0.447	0.853	4660
BM875598	1893	SPATA13	0.422	0.812	79
NM_177965	1894-1895	LOC157657	0.434	0.819	92
CA433208	1896	KIAA1411	0.427	0.809	69
BM984931	1897	MGC20781	0.417	0.795	117
BC035335	1898-1899	LOC255189	0.49	1.04	56
BC080193	1900	ERBB2	0.377	0.743	1770
NM_199427	1901-1902	ZFP64	0.374	0.688	80
BC042370	1903	SUHW2	0.364	0.678	587
AL137506	1904-1905	ELOVL7	0.322	0.673	256
AI888175	1906	TRAF4	0.343	0.631	50

5

TABLE 5D-continued

PROSTATE ENRICHED GENES IDENTIFIED BY RATIO SCHEMA (RATIO > 2.5)*					
Genbank Accession No.	Genbank SEQ ID NOs:	name	NN-score	Odds	Prostate Expression (tmp)
AI669751	1907	SLC39A2	0.34	0.629	60
AI088739	1908	HNF4G	0.32	0.593	225
BC070300	1909	SLC22A3	0.294	0.581	77
BC005827	1910	HIST2H2BE	0.306	0.587	912
BC041345	1911-1912	AMD1	0.317	0.588	438
BX390036	1913	TYMS	0.306	0.571	67
AK022455	1914	PHC3	0.287	0.57	105
AL832940	1915-1916	SARG	0.302	0.563	158
BC000965	1917-1918	MTERF	0.3	0.56	190
NM_007253	1919-1920	CYP4F8	0.28	0.566	54
AK124401	1921	PPAP2A	0.211	0.533	75
BC011408	1922-1923	KIAA0056	0.281	0.527	287
AA469293	1924	MSMB	0.27	0.517	275
AK056914	1925	VEGF	0.256	0.485	202
NM_004360	1926-1927	CDH1	0.179	0.362	192
BC062761	1928-1929	TARP	0.174	0.353	564
NM_001007278	1930-1931	RFP2	0.162	0.322	192
NM_018670	1932-1933	MESP1	0.154	0.315	133
AA026974	1934	TRPM4	0.147	0.305	290
AI468032	1935	PAK1IP1	0.13	0.271	74
CF122297	1936	HNRPA1	0.106	0.228	104
CB053869	1937	ZNF207	0.099	0.212	72
NM_032387	1938-1939	WNK4	0.089	0.201	100
BQ448015	1940	APXL2	0.083	0.19	244
AI554477	1941	MED28			700
AK095655	1942	LOC285300			84
AW291753	1943	0			310
BM023121	1944	0			178
AY338953	1945	0			166
AY533562	1946	0			67
BC030554	1947	0			66

35 *ratio of prostate expression in tmp to other organs greater than 2.5

Additional analysis was carried out to determine the secretion potential of the prostate-specific genes identified. The analysis programs used included SignalP 3.0, Secretome 2.0 and TMHMM 2.0 (see <http://colon.double.slash.www.cbs.dtu.dk/services/>). The SignalP analysis identifies classical secreted proteins and was conducted using the classical secretion pathway prediction as described at <http://colon.double.slash.www.cbs.dtu.dk/services/SignalP/> (see Jannick Dyrlov Bendtsen, et al. *J. Mol. Biol.*, 340:783-795, 2004; Henrik Nielsen et al., *Protein Engineering*, 10:1-6, 1997; Henrik Nielsen and Anders Krogh. *Proceedings of the Sixth International Conference on Intelligent Systems for Molecular Biology (ISMB 6)*, AAAI Press, Menlo Park, Calif., pp. 122-130, 1998). The Secretome2.0 analysis identifies nonclassical secreted proteins (see J. Dyrlov Bendtsen, et al., *Protein Eng. Des. Sel.*, 17(4):349-356, 2004). TMHMM uses hidden Markov model for three-state (TM-helix, inside, outside) topology prediction of transmembrane proteins (see Erik L. L. Sonnhammer, et al., *Proc. of Sixth Int. Conf. on Intelligent Systems for Molecular Biology*, p. 175-182 Ed. J. Glasgow, T. Littlejohn, F. Major, R. Lathrop, D. Sankoff, and C. Sensen, Menlo Park, Calif.: AAAI Press, 1998). According to the SignalP analysis method, proteins with an odds scoring 3 or higher have a high confidence of being secreted. However, it should be noted that several proteins scoring well below 3 by this method are known to be secreted proteins detected in the blood (see e.g., Table 5, KLK2). Further, these analyses do not take into account proteins that may be shed.

In summary, this example identifies prostate-specific and potentially secreted prostate-specific proteins that can be used in diagnostic panels for the detection of diseases of the prostate.

Example 8

Prostate Cancer Diagnostics Using Multiparameter Analysis

This example describes a multiparameter diagnostic fingerprint using the NDRG1 prostate-specific protein in combination with PSA. The NDRG1 prostate-specific protein further improved prostate cancer detection when used in combination with PSA.

Commercially available antibodies specific for numerous proteins encoded by prostate-specific genes as described in Table 5 were used to determine which proteins would be useful in a multiparameter diagnostic assay for prostate cancer. Most of the commercially available antibodies were not suitable (e.g., were not sensitive enough or showed non-specific binding). However, the antibody available for NDRG1 (anti-NDRG1(C terminal) poly IgY ; Cat #A22272B; GenWay Inc) was shown to specifically bind to NDRG 1 from serum. NDRG 1 is a member of the N-myc downregulated gene (NDRG) family that belongs to the alpha/beta hydrolase superfamily. It is classified as a tumor suppressor and heavy metal-response protein. Its expression is modulated by diverse physiological and pathological con-

ditions including hypoxia, cellular differentiation, heavy metal, N-myc and neoplasia (Lachat P, et al.; Histochem Cell Biol. 2002 Nov;118(5):399-408).

NDRG1 protein expression was analyzed in serum samples from 18 advanced prostate cancer patients, 21 prostate cancer patients with localized cancer, and 22 normal controls. Western blot analysis was used to measure serum protein expression as follows: Serum was diluted (1:10) with lysis buffer (50 mm Hepes, pH 7.4, 4 mm EDTA, 2 mm EGTA, 2 μM PMSF, 20 μg/ml, leupeptine (or 1× protease inhibitor cocktail), 1 mm Na₃VO₄, 10 mm NaF, 2 mm Na pyrophosphate, 1% Triton X-100). Protein concentration was determined using the Bio-Rad protein assay kit. Serum proteins (50 μg) were subjected to SDS-PAGE electrophoresis and transferred to a PVDF membrane (Hybond-P, Amersham Pharmacia Biotech, Piscataway, N.J.). The membrane was blocked with 4% non-fat milk in TBS (25 mm Tris, pH 7.4, 125 mm NaCl) for 1 h at room temperature, followed by incubation with primary antibodies against NDRG1 IgY (1:500) overnight at 4° C. The membranes were washed 3 times with TBS, and then incubated with horseradish peroxidase conjugated anti-rabbit IgY (1:16,000) for 1 h. The immunoblot was then washed five times with TBS and developed using an ECL (Amersham). The intensities of the single band corresponding to the NDRG 1 protein were then scored. The results are summarized in Table 6 together with serum PSA measurements performed using a commercial ELISA kit.

TABLE 6

COMBINED ANALYSIS OF NDRG1 AND PSA SERUM EXPRESSION INCREASES PROSTATE CANCER DIAGNOSIS CONFIDENCE.				
cancer status	NDRG-1 intensity (scores*)	PSA values (ng/ml)	serum diagnosis by PSA	serum diagnosis by NDRG1
Advanced	3	70.48	identified as cancer by PSA assay	identified as cancer by NDRG1 assay
Advanced	4	127.3	identified as cancer by PSA assay	identified as cancer by NDRG1 assay
Advanced	4	422.1	identified as cancer by PSA assay	identified as cancer by NDRG1 assay
Advanced	4	1223	identified as cancer by PSA assay	identified as cancer by NDRG1 assay
Advanced	4	71.28	identified as cancer by PSA assay	identified as cancer by NDRG1 assay
Advanced	2	133.2	identified as cancer by PSA assay	missed by NDRG1 assay
Advanced	4	353.7	identified as cancer by PSA assay	identified as cancer by NDRG1 assay
Advanced	1	73.95	identified as cancer by PSA assay	missed by NDRG1 assay
Advanced	3	454.8	identified as cancer by PSA assay	identified as cancer by NDRG1 assay
Advanced	4	474	identified as cancer by PSA assay	identified as cancer by NDRG1 assay
Advanced	6	150.1	identified as cancer by PSA assay	identified as cancer by NDRG1 assay

TABLE 6-continued

COMBINED ANALYSIS OF NDRG1 AND PSA SERUM EXPRESSION INCREASES PROSTATE CANCER DIAGNOSIS CONFIDENCE.				
cancer status	NDRG-1 intensity (scores*)	PSA values (ng/ml)	serum diagnosis by PSA	serum diagnosis by NDRG1
Advanced	0	1375	identified as cancer by PSA assay	missed by NDRG1 assay
Advanced	6	71.28	identified as cancer by PSA assay	identified as cancer by NDRG1 assay
Advanced	6	4066	identified as cancer by PSA assay	identified as cancer by NDRG1 assay
Advanced	4	1199	identified as cancer by PSA assay	identified as cancer by NDRG1 assay
Advanced	1	38.14	identified as cancer by PSA assay	missed by NDRG1 assay
Advanced	6	552.6	identified as cancer by PSA assay	identified as cancer by NDRG1 assay
Advanced	5	321	identified as cancer by PSA assay	identified as cancer by NDRG1 assay
Primary	-1	14.2	possibly cancer	
Primary	2	6.27	Grey Zone of diagnosis by Psa	
Primary	2	9.2	Grey Zone of diagnosis by Psa	
Primary	1	8.57	Grey Zone of diagnosis by Psa	
Primary	0	5.67	Grey Zone of diagnosis by Psa	
Primary	2	11.3	possibly cancer	
Primary	0	4.58	Grey Zone of diagnosis by Psa	
Primary	0	5.67	Grey Zone of diagnosis by Psa	
Primary	-1	6.48	Grey Zone of diagnosis by Psa	
Primary	3	12.71	possibly cancer	strong NDRG-1 expression reinforces the diagnosis of this patient as cancer
Primary	3	4.93	Grey Zone of diagnosis by Psa	strong NDRG-1 expression reinforces the diagnosis of this patient as cancer
Primary	1	3.16	Grey Zone of diagnosis by Psa	
Primary	1	4.87	Grey Zone of diagnosis by Psa	
Primary	1	4.66	Grey Zone of diagnosis by Psa	
Primary	1	6.87	Grey Zone of diagnosis by Psa	
Primary	0	3.91	Grey Zone of diagnosis by Psa	
Primary	0	6.48	Grey Zone of diagnosis by Psa	
Primary	2	13.1	possibly cancer	
Primary	0	4.58	Grey Zone of diagnosis by Psa	
Primary	1	4.72	Grey Zone of diagnosis by Psa	
Primary	4	12.71	possibly cancer	strong NDRG-1 expression reinforces the diagnosis of this patient as cancer
Normal	-1	0.8	Normal	normal
Normal	-1	0.8	Normal	normal
Normal	0	0.6	Normal	normal
Normal	1	1	Normal	normal
Normal	-1	1.2	Normal	normal
Normal	-1	1.91	Normal	normal
Normal	2	0.6	Normal	normal
Normal	-1	0.3	Normal	normal
Normal	0	1	Normal	normal

TABLE 6-continued

COMBINED ANALYSIS OF NDRG1 AND PSA SERUM EXPRESSION INCREASES PROSTATE CANCER DIAGNOSIS CONFIDENCE.				
cancer status	NDRG-1 intensity (scores*)	PSA values (ng/ml)	serum diagnosis by PSA	serum diagnosis by NDRG1
Normal	-1	0.4	Normal	normal
Normal	-1	0.8	Normal	normal
Normal	0	1	Normal	normal
Normal	1	0.8	Normal	normal
Normal	2	0.6	Normal	normal
Normal	1	0.5	Normal	normal
Normal	1	1	Normal	normal
Normal	-1	0.7	Normal	normal
Normal	-1	1.2	Normal	normal
Normal	-1	1.1	Normal	normal
Normal	0	0.8	Normal	normal
Normal	0	0.7	Normal	normal
Normal	0	0.6	Normal	normal

*scores: no expression, -1; no expression to very faint, 0; expression levels then scored from 1 to 6 by intensities

PSA was detected in 100% of the advanced prostate cancers. NDRG1 was detected in 14 out of 18 advanced cancers. (78%) (see Table 6, scores greater than 3). Serum PSA levels below 15 ng/ml, particularly, levels between 4-10 ng/ml (often referred to as the 'grey zone' in the PSA assay) cannot reliably detect prostate cancer as PSA levels in this range may be the result of other factors such as infection (prostatitis) or benign prostatic hyperplasia (BPH), a common condition in older men. Additionally, the normal range of PSA values increases with patient age. NDRG1 detection in serum reinforced the diagnosis of three prostate cancer patients with PSA levels between 4.9 ng/ml and 15 ng/ml. In these three

patients, the NDRG1 scores were 3 or 4, significantly higher than the NDRG1 scores in a cohort of 22 normal individuals (average 0.09, range -1 to 2).

Thus, this example illustrates that the use of two or more prostate specific/enriched cancer markers such as NDRG1 and PSA can improve prostate cancer diagnosis to reduce false positive and false negative rates.

From the foregoing it will be appreciated that, although specific embodiments of the invention have been described herein for purposes of illustration, various modifications may be made without deviating from the spirit and scope of the invention. Accordingly, the invention is not limited except as by the appended claims.

SEQUENCE LISTING

The patent contains a lengthy "Sequence Listing" section. A copy of the "Sequence Listing" is available in electronic form from the USPTO web site (<http://seqdata.uspto.gov/?pageRequest=docDetail&DocID=US07883858B2>). An electronic copy of the "Sequence Listing" will also be available from the USPTO upon request and payment of the fee set forth in 37 CFR 1.19(b)(3).

What is claimed is:

1. A method for detecting a drug side effect comprising measuring in the blood of a subject taking the drug the level of a plurality of organ-specific proteins secreted from an organ wherein the levels of the plurality of organ-specific proteins together provide an organ-specific molecular blood fingerprint that indicates a drug side effect on the organ in the subject.

2. The method of claim 1 wherein the level of the plurality of organ-specific proteins is measured using a method selected from the group consisting of mass spectrometry, immunoassay, tandem mass spectrometry, ELISA, Western blot, microfluidics/nanotechnology sensors, and aptamer capture assay.

3. The method of claim 2 wherein the level of the plurality of organ-specific proteins is measured using tandem mass spectrometry.

4. The method of claim 2 wherein the level of the plurality of organ-specific proteins is measured using ELISA.

5. The method of claim 1 wherein the plurality of organ-specific proteins comprises 5 organ-specific proteins.

6. The method of claim 1 wherein the plurality of organ-specific proteins comprises 20 organ-specific proteins.

7. The method of claim 1 wherein the organ-specific proteins comprise liver-specific proteins.

8. The method of claim 1 wherein the organ-specific proteins comprise kidney-specific proteins.

9. The method of claim 1 wherein the organ-specific proteins comprise prostate-specific proteins.

10. The method of claim 2 wherein the level of each of the plurality of organ-specific proteins is measured using an antibody array.

11. The method of claim 1 wherein the plurality of organ-specific proteins comprises 10 organ-specific proteins.

135

12. The method of claim 1 wherein the plurality of organ-specific proteins comprises 50 organ-specific proteins.

13. The method of claim 1 wherein the organ-specific proteins are from an organ other than the expected therapeutic target of the drug.

14. The method of claim 1 wherein measuring the level of the plurality of organ-specific proteins is accomplished by detecting the level of each of the plurality of organ-specific proteins in a blood sample from the subject.

15. The method of claim 1 wherein a statistically significant altered level of one or more of the plurality of organ-specific proteins in the blood is indicative of the presence or absence of a drug side effect.

16. The method of claim 15 where the altered level of one or more of the plurality of organ-specific proteins is determined by comparing the level of each of the plurality of organ-specific proteins in the blood sample from the subject to a level of each of the plurality of organ-specific proteins in a control sample of blood.

136

17. The method of claim 1 wherein the plurality of organ-specific proteins are secreted from the same organ.

18. The method of claim 1 wherein the plurality of organ-specific proteins comprises 100 organ-specific proteins.

19. The method of claim 1 wherein the plurality of organ-specific proteins comprises 5 organ-specific proteins from each particular organ.

20. The method of claim 1 wherein the plurality of organ-specific proteins comprises 10 organ-specific proteins from each particular organ.

21. The method of claim 1 wherein the plurality of organ-specific proteins comprises 20 organ-specific proteins from each particular organ.

22. The method of claim 1 wherein the plurality of organ-specific proteins comprises 50 organ-specific proteins from each particular organ.

* * * * *

专利名称(译)	鉴定和监测药物副作用的方法		
公开(公告)号	US7883858	公开(公告)日	2011-02-08
申请号	US12/468834	申请日	2009-05-19
[标]申请(专利权)人(译)	系统生物学研究所		
申请(专利权)人(译)	系统生物学研究所		
当前申请(专利权)人(译)	系统生物学研究所		
[标]发明人	HOOD LEROY LIN BIAOYANG		
发明人	HOOD, LEROY LIN, BIAOYANG		
IPC分类号	G01N33/53		
CPC分类号	G01N33/6842 G01N33/6893 G01N33/57434		
优先权	60/683004 2005-05-20 US 60/647792 2005-01-27 US		
其他公开文献	US20100075866A1		
外部链接	Espacenet USPTO		

摘要(译)

本发明一般涉及通过检测器官特异性分子血液指纹中的扰动来鉴定药物副作用的方法。本发明还涉及鉴定药物特异性器官特异性分子血液指纹的方法。因此，本发明提供了包含器官特异性蛋白质的组合物，用于检测此类蛋白质的检测试剂，以及用于确定器官特异性分子血液指纹的面板和阵列。

TABLE 3
DIFFERENTIALLY EXPRESSED GENES THAT ENCODE PREDICTED SECRETED PROTEINS

Signature	SEQ ID NO.	Accession Number	SEQ ID NO.	Description
GATCAGCATGGCCACG	73	NM_001928	594-595	D component of complement (cd5p1n)
GATCTACTACTGGCCT	74	NM_006280	596-597	signal sequence receptor, delta (extralocus associated protein delta)
GATCCTGTTGGAAAGA	75	NM_203329	598-599	CD59 antigen p18-20 (antigen identified by monoclonal antibodies 16.3A5, E31c, E33c, EL32 and G344)
GATCCTGTTGGAAAGA	76	NM_203331	600-601	CD59 antigen p18-20 (antigen identified by monoclonal antibodies 16.3A5, E31c, E33c, EL32 and G344)
GATCCCTGAAGTGCC	77	NM_203331	600-601	CD59 antigen p18-20 (antigen identified by monoclonal antibodies 16.3A5, E31c, E33c, EL32 and G344)
GATCTGGCTGTATTA	78	NM_203331	600-601	CD59 antigen p18-20 (antigen identified by monoclonal antibodies 16.3A5, E31c, E33c, EL32 and G344)
GATCCCTGAAGTGCC	79	NM_203330	602-603	CD59 antigen p18-20 (antigen identified by monoclonal antibodies 16.3A5, E31c, E33c, EL32 and G344)
GATCCTGTTGGAAAGA	80	NM_203330	602-603	CD59 antigen p18-20 (antigen identified by monoclonal antibodies 16.3A5, E31c, E33c, EL32 and G344)
GATCTGGCTGTATTA	81	NM_203330	602-603	CD59 antigen p18-20 (antigen identified by monoclonal antibodies 16.3A5, E31c, E33c, EL32 and G344)