



US009476868B2

(12) **United States Patent**  
**Al-Murrani et al.**

(10) **Patent No.:** **US 9,476,868 B2**  
(45) **Date of Patent:** **Oct. 25, 2016**

(54) **COMPOSITIONS AND METHODS FOR  
DIAGNOSING AND TREATING KIDNEY  
DISORDERS IN A CANINE**

(75) Inventors: **Samer Al-Murrani**, Topeka, KS (US);  
**Sukhaswami Malladi**, Lawrence, KS  
(US); **Xiangming Gao**, Topeka, KS  
(US)

(73) Assignee: **Hill's Pet Nutrition, Inc.**, Topeka, KS  
(US)

(\* ) Notice: Subject to any disclaimer, the term of this  
patent is extended or adjusted under 35  
U.S.C. 154(b) by 0 days.

(21) Appl. No.: **13/519,048**

(22) PCT Filed: **Dec. 23, 2010**

(86) PCT No.: **PCT/US2010/062031**

§ 371 (c)(1),

(2), (4) Date: **Jun. 25, 2012**

(87) PCT Pub. No.: **WO2011/079280**

PCT Pub. Date: **Jun. 30, 2011**

(65) **Prior Publication Data**

US 2012/0283129 A1 Nov. 8, 2012

**Related U.S. Application Data**

(60) Provisional application No. 61/289,773, filed on Dec.  
23, 2009.

(51) **Int. Cl.**

**C12Q 1/68** (2006.01)

**G01N 33/53** (2006.01)

**G01N 33/50** (2006.01)

(52) **U.S. Cl.**

CPC ..... **G01N 33/50** (2013.01); **C12Q 1/6883**  
(2013.01); **C12Q 2600/158** (2013.01); **G01N**  
**2800/347** (2013.01)

(58) **Field of Classification Search**

None

See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

4,683,195 A 7/1987 Mullis et al.  
4,683,202 A 7/1987 Mullis et al.  
4,800,159 A 1/1989 Mullis et al.  
4,965,188 A 10/1990 Mullis et al.  
5,333,675 A 8/1994 Mullis et al.  
5,445,934 A 8/1995 Fodor et al.  
5,541,061 A 7/1996 Fodor et al.  
5,547,839 A 8/1996 Dower et al.  
5,677,195 A 10/1997 Winkler et al.  
5,744,305 A 4/1998 Fodor et al.  
5,770,722 A 6/1998 Lockhart et al.  
5,800,992 A 9/1998 Fodor et al.  
5,856,092 A 1/1999 Dale et al.  
5,858,659 A 1/1999 Sapolsky et al.  
5,902,723 A 5/1999 Dower et al.

6,013,449 A 1/2000 Hacia et al.  
6,020,135 A 2/2000 Levine et al.  
6,033,860 A 3/2000 Lockhart et al.  
6,040,138 A 3/2000 Lockhart et al.  
6,040,193 A 3/2000 Winkler et al.  
6,156,501 A 12/2000 McGall et al.  
6,177,248 B1 1/2001 Oliner et al.  
6,197,506 B1 3/2001 Fodor et al.  
6,203,825 B1 3/2001 Hodgkins  
6,284,460 B1 9/2001 Fodor et al.  
6,300,063 B1 10/2001 Lipshutz et al.  
6,300,070 B1 10/2001 Boles et al.  
6,309,822 B1 10/2001 Fodor et al.  
6,333,179 B1 12/2001 Matsuzaki et al.  
6,368,799 B1 4/2002 Chee  
7,300,652 B2 11/2007 Bailly et al.  
7,482,135 B2 1/2009 Ver Heijen et al.  
2002/0142284 A1 10/2002 Raha et al.  
2003/0073100 A1 4/2003 Peyman et al.  
2003/0087259 A1\* 5/2003 Clancy et al. .... 435/6  
2004/0038856 A1 2/2004 Chakravarty et al.  
2004/0101924 A1 5/2004 Ver Heijen et al.  
2005/0032063 A1 2/2005 Kopreski  
2005/0124534 A1 6/2005 Noble et al.  
2006/0008804 A1 1/2006 Chibout et al.  
2006/0240437 A1 10/2006 Krolewski et al.  
2006/0253262 A1\* 11/2006 Ching et al. .... 702/20  
2007/0015227 A1 1/2007 Moses et al.  
2007/0243214 A1 10/2007 Schiemann et al.  
2007/0254290 A1 11/2007 Yen et al.  
2009/0117604 A1 5/2009 Ver Heijen et al.  
2009/0203547 A1\* 8/2009 Banes et al. .... 506/16

FOREIGN PATENT DOCUMENTS

EP 0985932 3/2000  
EP 1808694 7/2007  
JP S60-153762 A 8/1985  
JP H08-191668 A 7/1996  
WO WO 01/88544 11/2001  
WO WO 2004/005544 1/2004  
WO WO 2004/005934 1/2004

OTHER PUBLICATIONS

Greer et al in "Gene expression analysis in a canine model of  
X-linked Alport syndrome" (Mammalian Genome: 2006, vol. 17,  
pp. 976-990).\*

(Continued)

Primary Examiner — Catherine S Hibbert

(57) **ABSTRACT**

The present invention provides methods for: diagnosing of,  
devising and monitoring a treatment plan for, and monitor-  
ing the status of a kidney disorder characterized by an  
abnormal loss of renal function, renal failure, reduced glom-  
erular filtration rate or glomerulonephritis, in a canine,  
wherein the kidney disorder is detectable by utilizing at least  
one relevant biomarker isolated and measured from a bio-  
logical test sample taken from such canine. The invention  
additionally relates to compositions, reagents and kits for  
carrying out the specified methods.

**5 Claims, No Drawings**

(56)

## References Cited

## OTHER PUBLICATIONS

- Quan Dissertation entitled "Expression Profile of Drug Absorption Related Genes in MDCK In Vitro Model" (Sep. 2008).\*
- Mody et al entitled "Decreased clearance of serum retinol-binding protein and elevated levels of transthyretin in insulin-resistant ob/ob mice" (*Am J Physiol Endocrinol Metab* 294: E785-E793, published Feb. 19, 2008).\*
- Neame et al (CMLS, *Cell Mol Life Sci* vol. 57 2000, pp. 859-863).\*
- Altschul et al., 1990, "Basic Local Alignment Search Tool", *Journal of Molecular Biology* 215(3):403-410.
- Altschul et al., 1997 "Gapped BLAST and PSI-Blast: a New Generation of Protein Database Search Programs", *Nucleic Acids Research* 25(17):3389-3402.
- Braun et al., 2002, "Plasma Cystatin C in the Dog: Reference Values and Variations with Renal Failure", *Comparative Clinical Pathology* 11(1):44-49.
- Eckert et al., 1991, "DNA Polymerase Fidelity and the Polymerase Chain Reaction", *PCR Methods & Applications* 1:17-24.
- Fan et al., 2005, "A Class of Models for Analyzing GeneChip®, Gene Expression Analysis Array Data", *BMC Genomics* 16:6-16.
- Folli et al., 2001, "Identification, Retinoid Binding, and X-ray Analysis of a Human Retinol-binding Protein", *Proceedings of the National Academy of Sciences* 98(7):3710-3715.
- Forterre et al., 2004, "Protein Profiling of Urine from Dogs with Renal Disease using ProteinChip Analysis", *Journal of Veterinary Diagnostic Investigation* 16(4):271-277.
- Gerhard et al., 2004, "The Status, Quality, and Expansion of the NIH Full-Length cDNA Project: The Mammalian Gene Collection (MGC)", *Genome Research* 14:2121-2127.
- Hein et al., 2005, "BGX: a Fully Bayesian Integrated Approach to the Analysis of Affymetrix GeneChip Data", *Biostatistics* 6:349-373.
- Ho et al., 2007, "Down-regulation of Retinol Binding Protein 5 is Associated with Aggressive Tumor Features in Hepatocellular Carcinoma", *Journal of Cancer Research & Clinical Oncology* 133(12):929-936.
- Irizarry et al. 2003, "Exploration, Normalization, and Summaries of High Density Oligonucleotide Array Probe Level Data", *Biostatistics* 4:249-264.
- Irizarry et al., 2003, "Summaries of Affymetrix GeneChip Probe Level Data", *Nucleic Acid Research* 31(4):e15.
- Lee et al, 2006, "Secreted Frizzled Related Protein 2 (sFRP2) Decreases Susceptibility to UV-Induced Apoptosis in Primary Culture of Canine Mammary Gland Tumors by NF- $\kappa$ B Activation or JNK Suppression", *Breast Cancer Research and Treatment* 100(1):49-58.
- Karlin et al., 1990, "Methods for Assessing the Statistical Significance of Molecular Sequence Features by Using General Scoring Schemes", *Proceedings of the National Academy of Sciences* 87:2264-2268.
- Laterza et al., 2002, "Cystatin C: An Improved Estimator of Glomerular Filtration Rate?", *Clinical Chemistry* 48:699-707.
- Lenz et al., 1998, "Glomerular Endothelial Cell; Synthesize Collagens but Little Gelatinase A and B.", *Journal of American Society of Nephrology* 9:2040-2047.
- Lenz et al., 2000, "Matrix Metalloproteinases in Renal Development and Disease", *Journal of American Society of Nephrology* 11:574-581.
- Li et al., 2001, "Model-based Analysis of Oligonucleotide Arrays: Expression Index Computation and Outlier Detection", *Proceedings of the National Academy of Sciences* 98(1):31-36.
- Mattila et al., 1991, "Fidelity of DNA Synthesis by the *Thermococcus Litoralis* DNA Polymerase—an Extremely Heat Stable Enzyme with Proofreading Activity", *Nucleic Acids Research* 19(18):4967-4973.
- Miyagawa et al., 2009, "Evaluation of the Measurement of Serum Cystatin C by an Enzyme-linked Immunosorbent Assay for Humans as a Marker of the Glomerular Filtration Rate in Dogs", *Journal of Veterinary Medical Science* 71(9):1169-1176.
- Nagase et al., 1999, "Matrix Metalloproteinases", *Journal of Biological Chemistry* 274(31):21491-21494.
- Newman, 2002, "Cystatin C", *Annals of Clinical Biochemistry* 39:89-104.
- Nielsen et al., 1991, "Sequence-selective recognition of DNA by strand displacement with a thymine-substituted polyamide", *Science* 254:1497-1500.
- Nielsen, 1999, "Applications of Peptide Nucleic Acids", *Current Opinion in Biotechnology* 10(1):71-75.
- Onetti Muda et al., 1998, "Expression of TGF- $\beta$  Receptors Type I and II in Human Glomerulonephritis", *Nephrology, dialysis, transplantation* 13(2):279-284.
- Paciello et al., 2008, "G.P.5.12 Decorin and lumican are differentially expressed in canine masticatory muscle myositis", *Neuromuscular Disorders* 18(9-10):772-773, abstract No. G.P.5.12. PCT/US2010/062031—ISR and Written Opinion mailed Sep. 22, 2011.
- PCT/US2010/062031—Written Opinion mailed Feb. 23, 2011.
- Piantedosi et al., 2005, "Cellular Retinol-binding Protein Type III Is Needed for Retinoid Incorporation into Milk", *Journal of Biological Chemistry* 280(25):24286-24292.
- Raila et al., 2000, "The Distribution of Vitamin A and Retinol-binding Protein in the Blood Plasma, Urine, Liver and Kidneys of Carnivores", *Veterinary Research* 31(6):541-551.
- Sanders et al., 2004, "Renal Expression of Matrix Metalloproteinases in Human ANCA-associated Glomerulonephritis", *Nephrology Dialysis Transplantation* 19:1412-1419.
- Sato et al., 2005, "Roles of Membrane-type Matrix Metalloproteinase-1 in Tumor Invasion and Metastasis", *Cancer Science* 96(4):212-217.
- Schaefer et al., 2001, "Small Proteoglycans in Human Diabetic Nephropathy: Discrepancy Between Glomerular Expression and Protein Accumulation of Decorin, Biglycan, Lumican, and Fibromodulin", *FASEB Journal*, Published online.
- Schaeffer et al., 2000, "Small Proteoglycans of Normal Adult Human Kidney: Distinct Expression Patterns of Decorin, Biglycan, Fibromodulin, and Lumican", *Kidney International* 58:1557-1568.
- Schmidt et al., 2009, "Effect of Azotemia on Serum N-terminal proBNP Concentration in Dogs with Normal Cardiac Function: A Pilot Study", *Journal of Veterinary Cardiology* 11(suppl 1):S81-S86.
- Schwab et al., 2004, "Microarray Analysis of Focal Segmental Glomerulosclerosis", *American Journal of Nephrology* 24:438-447.
- van Hoek et al., 2008, "Immunoassay of Urinary Retinol Binding Protein as Putative Renal Marker in Cats", *Journal of Immunization Methods* 329:208-213.
- Werten et al., 2000, "Gecko t-crystallin: How Cellular Retinol-binding Protein Became an Eye Lens Ultraviolet Filter", *Proceedings of the National Academy of Sciences* 97(7):3282-3287.
- Zheng, 1999, "Global Gene Expression Profiling on a Canine Model of X-linked Alport Syndrome: Cross-species cDNA Microarray Analysis Provides Insight into the Early Molecular Alterations Associated with the Onset of this Progressive Renal Disease", *Poster Presentation*, abstract No. 480.
- Zhou et al., 2005, "An Expression Index for Affymetrix GeneChips Based on the Generalized Logarithm", *Bioinformatics* 21(21):3983-3989.
- U.S. Appl. No. 60/319,253, filed May 17, 2002 by Shoulian et al.
- Almy et al., 2002, "Evaluation of cystatin C as an endogenous marker of glomerular filtration rate in dogs," *J. Vet. Intern. Med.* 16(1):45-51.
- Clements et al., 2009, "Cartilage gene expression correlates with radiographic severity of canine elbow osteoarthritis," *Vet. J.* 179(2):211-218.
- Hoffer et al., 1993, "The mouse apolipoprotein C1 gene: structure and expression," *Genomics* 18(1):37-42.
- Kawakami et al., 2001, "Relationship between testicular transferrin and plasma estradiol-17 $\beta$  concentrations of dogs with azoospermia and dogs with sertoli cell tumors," *J. Vet. Med. Sci.* 63(5):579-581.
- Lee et al., 2004, "Secreted frizzled-related protein 2 (SFRP2) is highly expressed in canine mammary gland tumors but not in normal mammary glands," *Breast Cancer Res. Treat.* 84(2):139-149.

(56)

**References Cited**

## OTHER PUBLICATIONS

Maddens et al., 2010, "Validation of immunoassays for the candidate renal markers C-reactive protein, immunoglobulin G, thromboxane B2 and retinol binding protein in canine urine," *Vet. Immunol. Immunopathol.* 134(3-4):259-264.

Yokota et al., 2001, "High expression of 92 kDa type IV collagenase (matrix metalloproteinase-9) in canine mammary adenocarcinoma," *Biochim. Biophys. Acta* 1568(1):7-12.

Zheng et al., 2009, "Microarray identifies extensive downregulation of noncollagen extracellular matrix and profibrotic growth factor genes in chronic isolated mitral regurgitation in the dog," *Circulation* 119(15):2086-2095.

Tsuchiya et al, 2004, "Urinary protein analysis and renal mRNA expression in dogs with renal papillary necrosis induced by

nefiroacetam, a neurotransmission enhancer," *J. Toxicol. Sci.*, 29(4):447 (P7-18).

Hellman, N., et al., 'Matrix metalloproteinase 13 (MMP13) and tissue inhibitor of matrix metalloproteinase 1 (TIMP1), regulated by the MAPK pathway, are both necessary for Madin-Darby canine kidney tubulogenesis', *Journal of Biological Chemistry*, 2008, vol. 283, No. 7, pp. 4272-4282.

Ojaimi, C., et al., 'Altered expression of a limited number of genes contributes to cardiac decompensation during chronic ventricular tachypacing in dogs', *Physiological Genomics*, 2007, vol. 29, No. 1, pp. 76-83.

Suematsu, N., et al., 'Hyperhomocysteinemia alters cardiac substrate metabolism by impairing nitric oxide bioavailability through oxidative stress', *Circulation*, 2007, vol. 115, No. 2, pp. 255-262.

\* cited by examiner

## COMPOSITIONS AND METHODS FOR DIAGNOSING AND TREATING KIDNEY DISORDERS IN A CANINE

This application is a national stage entry under 35 U.S.C. §371 of International Patent Application No. PCT/US2010/062031, filed 23 Dec. 2010, which claims priority to U.S. Provisional Patent Application Ser. No. 61/289,773, filed on 23 Dec. 2009, which are incorporated herein by reference.

### FIELD OF THE INVENTION

The present invention relates to compositions and methods for: diagnosing of, devising and monitoring a treatment plan for, and monitoring the status of a kidney disorder characterized by an abnormal loss of renal function, renal failure, reduced glomerular filtration rate or glomerulonephritis, in a canine, wherein the kidney disorder is detectable by utilizing at least one relevant biomarker isolated and measured from a biological test sample taken from such canine. A relevant biomarker for practice of the compositions and methods of the present invention comprises an RNA transcript or its translation product taken from such biological test sample of such canine. A biological test sample for the practice of the method of the invention comprises a tissue sample of a kidney of such canine or a specimen of a biological fluid taken from such canine.

The present invention also relates to the identification of combinations of novel biomarkers for use in diagnosing of, devising and monitoring a treatment plan for, and monitoring the status of, a kidney disorder characterized by an abnormal loss of renal function, renal failure, reduced glomerular filtration rate or glomerulonephritis, in a canine.

### BACKGROUND OF THE INVENTION

Glomerulonephritis or glomerular nephritis ("GN") is a renal disease which is characterized by inflammation of the glomeruli or capillary loops of the kidney. It is a pathologic process associated with a number of diverse underlying diseases. The condition occurs in acute, sub-acute and chronic forms and also secondary to an infection. The former conditions, where a concurrent illness cannot be found, are generally referred to as idiopathic glomerulonephritis. The latter conditions are generally referred to as secondary GN. Whatever the underlying cause, immune complexes form and result in a series of events leading to glomerular injury and loss of renal function, proteinuria and ultimately, in some cases, renal failure.

Nephritis is an inflammation of the kidney, which may be a focal or diffuse proliferative or destructive disease involving the glomerulus, renal tubule or the kidney interstitial (or connective) tissue. The most common form of nephritis is glomerulonephritis. Nephritis may progress through a number of stages ending in end-stage kidney disease or end-stage renal failure.

Renal failure results from the inability of the kidney to maintain its normal functions. As a result, metabolic waste products and metabolites accumulate in the blood. These waste products and metabolites may adversely affect most bodily systems. Disturbances in the maintenance of fluid and electrolyte balances are characteristics of renal failure.

Acute renal failure may occur suddenly due to trauma, infection, inflammation or exposure to nephrotoxic substances. This condition may result in dehydration, hypotension and circulatory collapse. Acute renal failure is frequently segregated into three categories: (1) pre-renal

failure, which is associated with decreased renal blood flow; (2) intra-renal failure, which is associated with ischemia and toxins; and (3) post-renal failure, which results from obstruction of urine flow.

Chronic renal failure involves a progressive loss of kidney function that may eventually progress to end-stage renal disease or failure. At inception, chronic renal failure begins as a diminishing kidney function, without appreciable accumulation of metabolic waste products in the blood. As the glomerular filtration rate slows due to inflammation, waste products begin to accumulate. The disease progresses to uremia due to low kidney function, and high levels of protein end products start to accumulate and impair bodily functions. Common causes of chronic renal failure include: inflammation, infection, urinary tract obstruction and certain systemic diseases and toxicities, including hypercalcemia, lupus erythematosus, diabetes mellitus and hypertension.

End-stage renal disease is marked by irreversible chronic renal failure. Serum creatinine and blood urea nitrogen levels continue to rise and the resulting uremia impairs all bodily systems. The kidney can suffer permanent and almost complete loss of function on the order of 10% or less of normal kidney function. One cause of end-stage kidney disease is glomerulonephritis. Other causes include those mentioned for chronic renal failure.

Glomerulonephritis may arise as a result of a biological insult to the immune system. Foreign substances may adhere to the basement membrane and cause an immune response resulting in the production of antibodies. These antibodies may combine with the foreign substances to cause immune complexes that become deposited on the walls of the tiny glomerular capillaries, resulting in damage to the nephron. Alternatively, in some individuals the immune system can create autoantibodies which are immunoglobulins that may attack kidney cells resulting in a so-called autoimmune response. If proteins in the body are altered, an autoantibody response may ensue because the autoantibodies recognize the altered proteins as non-self. These autoantibody-protein complexes may likewise be deposited on the basement membrane of the glomerulus causing a disruption of the functioning of the nephron.

Glomerulonephritis is a common cause of proteinuria in dogs and may be either the idiopathic or secondary form of the condition. In the latter situation, the condition may develop secondary to neoplasia, inflammatory diseases, endocrine malfunctions, infections or familial nephropathies. As in humans, GN in dogs is mediated immunologically, involving immunoglobulins and complement factors in the body of the animal. Injury occurs within the glomeruli of the kidney resulting in morphological changes to the glomeruli. Eventually the injury is irreversible and leads to malfunction of the nephrons.

It is generally accepted in the scientific community that the regulation of gene expression plays a key role in the development of some diseases or conditions that affect an animal's health and well being. Similarly, a differential expression of genes is one factor in the development of such diseases and conditions and the evaluation of gene expression patterns has become widely recognized as important to understanding the development and control of such diseases and conditions at the molecular level. To advance the understanding of genes and their relationship to disease, a number of methods have been developed for studying differential gene expression, e.g., DNA microarrays, expressed tag sequencing (EST), serial analysis of gene expression (SAGE), subtractive hybridization, subtractive cloning and differential display (DD) for mRNA, RNA-arbitrarily

primed PCR (RAP-PCR), real-time PCR (RT-PCR), representational difference analysis (RDA), two-dimensional gel electrophoresis, mass spectrometry, and protein microarray based antibody-binding for proteins.

Due to the complexity of the biological pathways implicated in renal disease and the inherent molecular interactions and intercellular signaling processes, it is highly desirable to understand at a genetic level the interactions that are taking place. Detection of dysregulated genes in the early stages of loss of renal function in canines is helpful in understanding the biology of renal disease, especially glomerulonephritis on a genome-wide basis. The fact that gene dysregulation may be detected at an early stage of disease development in animals subjected to repeated ischemic injury is helpful in designing methods for diagnosing of, and devising and monitoring a treatment plan for, an abnormal loss of kidney function, renal failure, reduced glomerular filtration rate or glomerulonephritis, in a canine.

A more detailed understanding of the biological pathways involved through gene expression profiling will aid in the development of diagnostic procedures, reagents and test kits as well as salutary pharmaceutical, nutraceutical and nutritional (dietary) interventions in the disease pathways. These approaches may enable early detection and potentially prevention or treatment of the underlying kidney disorder, particularly glomerulonephritis, as well as in monitoring the prognosis of early stage renal failure and glomerulonephritis, especially in canines. Dysregulated genes involved in the pathology of such disorders may serve as important biomarkers for diagnosis and potentially prevention or treatment of the disorder and to optimize selection of appropriate pharmaceutical, nutraceutical and nutritional (dietary) interventions.

The level of gene expression and/or the determination of the level of functioning of an expressed gene product in a canine may be used to select an appropriate agent for therapeutic or prophylactic use. This data may be employed by the skilled worker in selecting appropriate drugs as agents for the prevention or treatment of renal diseases in canines through gene expression profiling. Gene expression data and analysis may also be used to select nutritional compositions, dietary supplements, and nutraceuticals having a salutary effect on kidney performance by utilizing biomarkers indicative of a healthy state of kidney functioning.

Only very limited work has been done to date in screening the canine genome for gene expression profiles in connection with the diagnosis of diseases in dogs. Certain work employing animal models has utilized cDNA array technologies to screen for gene expression in renal tissues associated with renal disease.

Studies in healthy populations of canines versus populations having a disease such as a kidney disease and loss of kidney function as described in this specification have not been extensively conducted. Little data is available with respect to the expression profile of the canine genome, especially with respect to the development of renal diseases in canines over time. Gene expression data contained in this specification identifies the genes associated with renal function in dogs. Such gene expression data enables development of compositions and methods for diagnosing of, devising and monitoring a treatment plan for, and monitoring the status of, a kidney disorder characterized by an abnormal loss of renal function, renal failure, reduced glomerular filtration rate or glomerulonephritis, in a canine, wherein the kidney disorder is detectable by utilizing at least one rel-

evant biomarker isolated and measured from a biological test sample taken from such canine.

Gene expression data contained in the specification and examples enables a variety of desirable inventions based on the gene expression profiles described in the specification and examples of this patent application. The data permits identification and quantification of gene expression products as biomarkers of disease prevention, identification and treatment of the underlying renal disease. Gene expression data elicited as a result of the practice of the methods of the invention also permits monitoring the progression of such renal disease. These inventions further include genetic testing to identify susceptible subpopulations of animals likely to be afflicted with such renal disease, to identify optimal diets for the prevention or treatment of such renal diseases, to identify pharmaceutical, nutraceutical and nutritional (dietary) interventions based on the findings set forth on this specification in order to treat the underlying renal diseases. The inventions also include biomarkers for early disease detection, targeted therapeutics, diagnostic reagents and kits for the analysis of tissue and blood samples from canines susceptible to or having such a renal disease.

#### SUMMARY OF THE INVENTION

The present invention relates to compositions and methods for: diagnosing of, devising and monitoring a treatment plan for, and monitoring the status of a kidney disorder characterized by an abnormal loss of renal function, renal failure, reduced glomerular filtration rate or glomerulonephritis, in a canine, wherein the kidney disorder is detectable by utilizing at least one relevant biomarker isolated and measured from a biological test sample taken from such canine.

A relevant biomarker for practice of the compositions and methods of the present invention comprises an RNA transcript or its translation product taken from such biological test sample of such canine. A biological test sample for the practice of the method of the invention comprises a tissue sample of a kidney of such canine or a specimen of a biological fluid taken from such canine.

In particular, this invention relates to methods for: diagnosing of, devising and monitoring a treatment plan for, and monitoring the status of, a kidney disorder characterized by an abnormal loss of renal function, renal failure, reduced glomerular filtration rate or glomerulonephritis, in a canine, through use of gene expression analysis.

The present invention also relates to the identification of combinations of novel biomarkers for use in determining diagnosing of, devising and monitoring a treatment plan for, and monitoring the status of, a kidney disorder characterized by an abnormal loss of renal function, renal failure, reduced glomerular filtration rate or glomerulonephritis, in a canine.

The invention additionally relates to compositions, reagents and kits for carrying out the specified methods.

The invention is based, in part, on the discovery that particular gene expression profiles in canines correlate with a change in such animal from a normal to an abnormal biological process in the kidney that may lead to a decline in renal function over time. A correlation of a particular gene expression profile with the risk of experiencing a decline in renal function can be predicted, detected and diagnosed in a canine without rendering a conventional clinical diagnosis based on art-recognized clinical signs and symptoms of renal disease. An altered gene expression profile in a canine is, therefore, predictive of a decline in renal function, as might otherwise be diagnosed at a later time by art-recog-

nized measurements of renal function. Such art-recognized measurements of renal function typically may include, for example, one of the following measurements: glomerular filtration rate, creatinine clearance rate, urinary protein levels, serum creatinine levels, urinary creatinine levels, blood urea nitrogen (BUN) levels, radioisotope metabolic labeling, soft tissue imaging, including sonography, magnetic resonance imaging and/or computed tomography. Non-intrusive assays such as serum creatinine and BUN levels typically show poor correlation with kidney histopathology and generally would not be predictive of future changes in the kidney.

The methods of evaluating a canine to diagnose a kidney disorder involve evaluating the gene expression level or activity of one or more of the following genes or the expression (translation) product of such genes in a canine, selected from the group consisting of: secreted frizzled-related protein 2 (sFRP2); matrilin-2 (Matn2); lumican (LUM); collagen alpha 1 (III) chain, variant 12 (COL3A1); decorin (DCN); retinol binding protein 4 (rbp4); MMP-9; clusterin (CLU); transferrin (TF); Apo-C-1 (ApoC1); and inhibin beta A (INHBA).

The methods of evaluating a canine to diagnose a kidney disorder involve evaluating the gene expression level or activity of one or more of the following genes or the expression product of such gene in a canine selected from the group consisting of: secreted frizzled-related protein 2 (sFRP2) or matrilin-2 (Matn2); and optionally, a second group consisting of: lumican (LUM); decorin (DCN); collagen alpha 1 (III) chain, variant 12 (COL3A1); retinol binding protein 4 (rbp4); MMP-9; clusterin (CLU); transferrin (TF); Apo-C-1 (ApoC1); and inhibin beta A (INHBA).

The biomarkers useful in the practice of the present invention are: lumican (LUM); collagen alpha 1 (III) chain, variant 12 (COL3A1); decorin (DCN); secreted frizzled-related protein 2 (sFRP2); matrilin-2 (Matn2); retinol binding protein 4 (rbp4); MMP-9; clusterin (CLU); transferrin (TF); Apo-C-1 (ApoC1); and inhibin beta A (INHBA), as more fully described below and in the sequence listings appended to this specification.

Canine lumican. Affymetrix probe CfaAFFx.10198.1.S1\_s\_at corresponds to NCBI Reference Sequence: XP\_539716.1, accession XP\_539716, predicted to be similar to Lumican precursor (keratin sulfate proteoglycan lumican [KSPG lumican]) [*Canis familiaris*]. Canine lumican is a 338 amino acid leucine-rich repeat protein coded by XM\_539716.2 at GeneID:482599 [SEQ. ID. NO: 1].

The terms "Lumican" and "LUM" and "Lum" denote a protein having substantially the amino acid sequence as shown in SEQ. ID. NO: 1. Preferably, lumican is a protein substantially consisting of the amino acid sequence as shown in SEQ. ID. NO: 1. Lumican also includes protein variants of the sequence as shown in SEQ. ID. NO: 1, such as allelic variants and other mutations such as substitutions, additions and/or deletions. The term lumican also refers to the nucleic acid sequence encoding the protein. The sequence corresponds to XM\_539716.2 and appended SEQ. ID. NO: 12. Lumican also refers to nucleic acid sequences that hybridize to SEQ. ID. NO: 12 and to fragments of SEQ. ID. NO: 12.

Canine collagen alpha 1 (III), variant 12 (COL3A1). Affymetrix probe CfaAffx.22804.1S1\_s\_at corresponds to NCBI Reference Sequence: XP\_863148.1, accession XP\_863148, predicted to be similar to Collagen alpha (III)

chain precursor isoform 12. Canine COL3A1 is a 1446 amino acid protein coded by XM\_863148.1 at GeneID: 478835 [SEQ. ID. NO: 2].

The terms "Collagen alpha (III) chain precursor isoform 12," "Col3a1" and "COL3A1" denote a protein having substantially the amino acid sequence as shown in SEQ. ID. NO: 2. Preferably, COL3A1 is a protein substantially consisting of the amino acid sequence as shown in SEQ. ID. NO: 2. COL3A1 also includes protein variants of the sequence as shown in SEQ. ID. NO: 2, such as allelic variants and other mutations such as substitutions, additions and/or deletions. The term COL3A1 also refers to the nucleic acid sequence encoding the protein. The sequence corresponds to XM\_863148.1 and appended SEQ. ID. NO: 13. COL3A1 also refers to nucleic acid sequences that hybridize to SEQ. ID. NO: 13 and to fragments of SEQ. ID. NO: 13.

Canine decorin (DCN). Affymetrix probe Cfa.6065.1.A1\_s\_at corresponds to NCBI Reference Sequence: NP\_001003228.1, accession NP\_001003228, decorin [*Canis lupus familiaris*]. Canine decorin is a 360 amino acid leucine-rich repeat protein coded by NM\_001003228.1 at GeneID: 403904 [SEQ. ID. NO: 3].

The terms "Decorin," "Dcn" and "DCN" denote a protein having substantially the amino acid sequence as shown in SEQ. ID. NO: 3. Preferably, decorin is a protein substantially consisting of the amino acid sequence as shown in SEQ. ID. NO: 3. Decorin also includes protein variants of the sequence as shown in SEQ. ID. NO: 3, such as allelic variants and other mutations such as substitutions, additions and/or deletions. The term decorin also refers to the nucleic acid sequence encoding the protein. The sequence corresponds to NM\_001003228.1 and appended SEQ. ID. NO: 14. Decorin also refers to nucleic acid sequences that hybridize to SEQ. ID. NO: 14 and to fragments of SEQ. ID. NO: 14.

Canine secreted frizzled-related protein 2 (sFRP2). Affymetrix probe Cfa.1200.1.S1\_s\_at corresponds to NCBI Reference Sequence: NP\_001002987.1, accession NP\_001002987 XP\_532695, secreted frizzled-related protein 2 (sFRP2) [*Canis lupus familiaris*]. Canine secreted frizzled-related protein 2 (sFRP2) is a 294 amino acid protein coded by NM\_001002987.1 at GeneID: 475471 [SEQ. ID. NO: 4].

The terms "secreted frizzle-related protein 2," "sFRP2," and "sFRP2" denote a protein having substantially the amino acid sequence as shown in SEQ. ID. NO: 4. Preferably, sFRP2 is a protein substantially consisting of the amino acid sequence as shown in SEQ. ID. NO: 4. sFRP2 also includes protein variants of the sequence as shown in SEQ. ID. NO: 4, such as allelic variants and other mutations such as substitutions, additions and/or deletions. The term sFRP2 also refers to the nucleic acid sequence encoding the protein. The sequence corresponds to NM\_001002987.1 and appended SEQ. ID. NO: 15. sFRP2 also refers to nucleic acid sequences that hybridize to SEQ. ID. NO: 15 and to fragments of SEQ. ID. NO: 15.

Canine matrilin-2 (Matn2). Affymetrix probe Cfa.9487.1A1\_at corresponds to NCBI Reference Sequence: XP\_5485552.2, accession NP\_5485552, similar to matrilin 2 isoform a precursor [*Canis lupus familiaris*]. Canine matrilin-2 is a 978 amino acid protein coded by XM\_5485552.2 at GeneID: 491431 [SEQ. ID. NO: 5].

The terms "matrilin 2 isoform a precursor," "matrilin 2" and "Matn2" denote a protein having substantially the amino acid sequence as shown in SEQ. ID. NO: 5. Preferably, Matn2 is a protein substantially consisting of the amino acid

sequence as shown in SEQ. ID. NO: 5. Matn2 also includes protein variants of the sequence as shown in SEQ. ID. NO: 5, such as allelic variants and other mutations such as substitutions, additions and/or deletions. The term Matn2 also refers to the nucleic acid sequence encoding the protein. The sequence corresponds to XM\_548552.2 and appended SEQ. ID. NO: 16. Matn2 also refers to nucleic acid sequences that hybridize to SEQ. ID. NO: 16 and to fragments of SEQ. ID. NO: 16.

Canine retinol binding protein 4 (rbp4). Affymetrix probe Cfa.15489.1.S1\_at corresponds to NCBI Reference Sequence: XP\_534969.2 accession NP\_534969, similar to retinol binding protein 4, plasma precursor (rbp4) [*Canis lupus familiaris*]. Canine retinol binding protein 4 (rbp4) is a 267 amino acid protein coded by XM\_534969.2 at GeneID: 477775 [SEQ. ID. NO: 6].

The terms “retinol binding protein 4,” “plasma precursor,” “retinol” and “Rbp4” denote a protein having substantially the amino acid sequence as shown in SEQ. ID. NO: 6. Preferably, Rbp4 is a protein substantially consisting of the amino acid sequence as shown in SEQ. ID. NO: 6. Rbp4 also includes protein variants of the sequence as shown in SEQ. ID. NO: 6, such as allelic variants and other mutations such as substitutions, additions and/or deletions. The term Rbp4 also refers to the nucleic acid sequence encoding the protein. The sequence corresponds to XM\_534969.2 and appended SEQ. ID. NO: 17. Rbp4 also refers to nucleic acid sequences that hybridize to SEQ. ID. NO: 17 and to fragments of SEQ. ID. NO: 17.

Canine matrix metalloproteinase 9 (MMP-9). Affymetrix probe Cfa.3470.1.S1\_at corresponds to NCBI Reference Sequence: NP\_001003219.1 accession NP\_001003219, matrix metalloproteinase 9 [*Canis lupus familiaris*]. Canine matrix metalloproteinase 9 (MMP9) is a 704 amino acid protein coded by NM\_001003219.1 at GeneID: 403885 [SEQ. ID. NO: 7].

The terms “matrix metalloproteinase 9” and “MMP9” denote a protein having substantially the amino acid sequence as shown in SEQ. ID. NO: 7. Preferably, MMP9 is a protein substantially consisting of the amino acid sequence as shown in SEQ. ID. NO: 7. MMP9 also includes protein variants of the sequence as shown in SEQ. ID. NO: 7, such as allelic variants and other mutations such as substitutions, additions and/or deletions. The term MMP9 also refers to the nucleic acid sequence encoding the protein. The sequence corresponds to NM\_001003219.1 and appended SEQ. ID. NO: 18. MMP9 also refers to nucleic acid sequences that hybridize to SEQ. ID. NO: 18 and to fragments of SEQ. ID. NO: 18.

Canine clusterin (CLU). Affymetrix probe Cfa1254.S1\_s\_at corresponds to NCBI Reference Sequence: NM\_001003370.1 accession NM\_001003370 canine clusterin. Canine clusterin is a 445 amino acid protein coded by NM\_001003370.1 at GeneID: 442971 [SEQ. ID. NO: 8].

The terms “clusterin” and “Clu” denote a protein having substantially the amino acid sequence as shown in SEQ. ID. NO: 8. Preferably, clusterin is a protein substantially consisting of the amino acid sequence as shown in SEQ. ID. NO: 8. Clusterin also includes protein variants of the sequence as shown in SEQ. ID. NO: 8, such as allelic variants and other mutations such as substitutions, additions and/or deletions. The term clusterin also refers to the nucleic acid sequence encoding the protein. The sequence corresponds to NM\_001003370.1 and appended SEQ. ID. NO: 19. Clusterin also refers to nucleic acid sequences that hybridize to SEQ. ID. NO: 19 and to fragments of SEQ. ID. NO: 19

Canine transferrin (TF). Affymetrix probe Cfa2217.1.A1\_at corresponds to NCBI Reference Sequence: XP\_534268.2 accession XP\_534268 similar to canine serotransferrin precursor (Transferrin) isoform 1. Canine transferrin is a 705 amino acid protein coded by XM\_534268.2 at GeneID: 477072 [SEQ. ID. NO: 9].

The terms “canine serotransferrin precursor (Transferrin) isoform 1,” “transferrin” and “TF” denote a protein having substantially the amino acid sequence as shown in SEQ. ID. NO: 9. Preferably, transferrin is a protein substantially consisting of the amino acid sequence as shown in SEQ. ID. NO: 9. Transferrin also includes protein variants of the sequence as shown in SEQ. ID. NO: 9, such as allelic variants and other mutations such as substitutions, additions and/or deletions. The term transferrin also refers to the nucleic acid sequence encoding the protein. The sequence corresponds to XM\_534268.2 and appended SEQ. ID. NO: 20. Transferrin also refers to nucleic acid sequences that hybridize to SEQ. ID. NO: 20 and to fragments of SEQ. ID. NO: 20.

Canine apolipoprotein C-1 (ApoC1). Affymetrix probe Cfa1254.S1\_s\_at corresponds to NCBI Reference Sequence: XP\_533643.2 accession XP\_533643 similar to canine apolipoprotein C-I precursor (ApoCI). Canine apolipoprotein C-I (ApoCI) is an 88 amino acid protein coded by NM\_533643.2 at GeneID: 113459 [SEQ. ID. NO: 10].

The terms “apolipoprotein C-1” and “ApoC1” denote a protein having substantially the amino acid sequence as shown in SEQ. ID. NO: 10. Preferably, transferrin is a protein substantially consisting of the amino acid sequence as shown in SEQ. ID. NO: 10. ApoC1 also includes protein variants of the sequence as shown in SEQ. ID. NO: 10, such as allelic variants and other mutations such as substitutions, additions and/or deletions. The term ApoC1 also refers to the nucleic acid sequence encoding the protein. The sequence corresponds to NM\_533643.2 and appended SEQ. ID. NO: 21. ApoC1 also refers to nucleic acid sequences that hybridize to SEQ. ID. NO: 21 and to fragments of SEQ. ID. NO: 21.

Inhibin beta A (INHBA). Affymetrix probe Cfa596.1A1\_at corresponds to NCBI Reference Sequence: XP\_540364 accession XP\_540364 similar to inhibin beta A chain precursor (INHBA) (Activin beta-A-chain) (erythroid differentiation protein; EDF). Canine inhibin beta A chain precursor (INHBA) is a 424 amino acid protein coded by XM\_540364.2 at GeneID: 483245 [SEQ. ID. NO: 11].

The terms “Inhibin beta A chain precursor” and “INHBA” denote a protein having substantially the amino acid sequence as shown in SEQ. ID. NO: 11. Preferably, inhibin is a protein substantially consisting of the amino acid sequence as shown in SEQ. ID. NO: 11. Inhibin also includes protein variants of the sequence as shown in SEQ. ID. NO: 11, such as allelic variants and other mutations such as substitutions, additions and/or deletions. The term inhibin also refers to the nucleic acid sequence encoding the protein. The sequence corresponds to XM\_540364.2 and appended SEQ. ID. NO: 22. Transferrin also refers to nucleic acid sequences that hybridize to SEQ. ID. NO: 22 and to fragments of SEQ. ID. NO: 22.

In a preferred embodiment of the present invention, a method for diagnosing a kidney disorder in a canine comprises the steps of: (a) measuring the level of expression of at least one biomarker in a biological sample from the canine, wherein the at least one biomarker is selected from secreted frizzled-related protein-2 (SFRP2); matrilin-2 (Matn2); lumican (LUM); decorin (DCN); collagen alpha 1 (III) chain, variant 12 (COL3A1); retinol binding protein 4

(rbp4); MMP-9; clusterin (CLU); transferrin (TF); Apo-C-1 (ApoC1); and inhibin beta A (INHBA); and (b) identifying the canine as having a kidney disorder wherein differences in expression of the one or more biomarkers in the sample relative to a control value for expression in a sample from a normal animal indicates the existence of a kidney disorder.

In another preferred embodiment of the present invention, the canine has normal kidney function as defined by art-recognized clinical measurements, e.g., glomerular filtration rate, creatinine clearance, urinary protein levels, blood creatinine levels, urinary creatinine levels and/or blood urea nitrogen levels and the methods of the invention may be used to detect and diagnose in such canine a change from a normal state to an abnormal state leading to a kidney disorder characterized by reduced renal function, renal failure, reduced glomerular filtration rate and glomerulonephritis.

In another preferred embodiment, an elevated level of activity or expression of one or more genes in a canine selected from the group consisting of: lumican (LUM); collagen alpha 1 (III) chain, variant 12 (COL3A1); decorin (DCN); secreted frizzled-related protein 2 (sFRP2); and matrilin-2 (Matn2) correlates with a kidney disorder characterized by an abnormal loss of renal function, renal failure, reduced glomerular filtration rate or glomerulonephritis. The level of activity or expression of one or more genes may be determined by measuring the expression product of such genes which may be a polynucleotide or a polypeptide or protein, typically an RNA transcript or its translation product.

In another preferred embodiment, differential expression of one or more genes in a canine selected from at least one RNA transcript or its translation product selected from the group consisting of: secreted frizzled-related protein 2 (sFRP2); matrilin-2 (Matn2); lumican (LUM); decorin (DCN); collagen alpha 1 (III) chain, variant 12 (COL3A1); retinol binding protein 4 (rbp4); MMP-9; clusterin (CLU); transferrin (TF); Apo-C-1 (ApoC1); and inhibin beta A (INHBA) correlates with an abnormal loss of renal function, renal failure, reduced glomerular filtration rate or glomerulonephritis.

In another preferred embodiment, differential expression of one or more genes in a canine selected from a first group of at least one RNA transcript or its translation product selected from the group consisting of: secreted frizzled-related protein 2 (sFRP2) or matrilin-2 (Matn2); and optionally, a second group of at least one RNA transcript or its translation product selected from the group consisting of: lumican (LUM); decorin (DCN); collagen alpha 1 (III) chain, variant 12 (COL3A1); retinol binding protein 4 (rbp4); MMP-9; clusterin (CLU); transferrin (TF); Apo-C-1 (ApoC1); and inhibin beta A (INHBA) correlates with an abnormal loss of renal function, renal failure, reduced glomerular filtration rate or glomerulonephritis. It is to be understood that the present invention contemplates combination of biomarkers comprising genes or their expression products that are both (i) selected from a first group of at least one RNA transcript or its translation product selected from the group consisting of: secreted frizzled-related protein 2 (sFRP2) or matrilin-2 (Matn2); and optionally, a second group of at least one RNA transcript or its translation product selected from the group consisting of: lumican (LUM); decorin (DCN); collagen alpha 1 (III) chain, variant 12 (COL3A1); retinol binding protein 4 (rbp4); MMP-9; clusterin (CLU); transferrin (TF); Apo-C-1 (ApoC1); and inhibin beta A (INHBA); as well as (ii) selected from a first group of at least one RNA transcript or its translation

product selected from the group consisting of: secreted frizzled-related protein 2 (sFRP2) or matrilin-2 (Matn2); and a second group of at least one RNA transcript or its translation product selected from the group consisting of: lumican (LUM); decorin (DCN); collagen alpha 1 (III) chain, variant 12 (COL3A1); retinol binding protein 4 (rbp4); MMP-9; clusterin (CLU); transferrin (TF); Apo-C-1 (ApoC1); and inhibin beta A (INHBA). Embodiments of the invention contemplate constructing panels of biomarkers from various combinations of the two groupings of genes and their expression products.

In another preferred embodiment, the invention encompasses measurement and correlation of differential expression in a canine of one or more genes selected from at least one RNA transcript or its translation product selected from the group consisting of: secreted frizzled-related protein 2 (sFRP2); matrilin-2 (Matn2); lumican (LUM); decorin (DCN); collagen alpha 1 (III) chain, variant 12 (COL3A1); retinol binding protein 4 (rbp4); MMP-9; clusterin (CLU); transferrin (TF); Apo-C-1 (ApoC1); and inhibin beta A (INHBA), with accelerated or early loss of renal function in canines as evidenced by an abnormal process leading to a kidney disorder characterized by reduced renal function, reduced glomerular filtration rate, glomerulonephritis or renal failure.

In one aspect the invention includes contacting a tissue sample or bodily fluid specimen with an agent that detects in a canine one or more genes or the expression product of such one or more genes selected from at least one RNA transcript or its translation product selected from the group consisting of: secreted frizzled-related protein 2 (sFRP2); matrilin-2 (Matn2); lumican (LUM); decorin (DCN); collagen alpha 1 (III) chain, variant 12 (COL3A1); retinol binding protein 4 (rbp4); MMP-9; clusterin (CLU); transferrin (TF); Apo-C-1 (ApoC1); and inhibin beta A (INHBA). The agent can be an antibody or a nucleic acid probe used in conjunction with conventional assay means such as immobilization on a solid phase, microtiter wells, tubes, dipsticks or other conventional means.

Another embodiment of the method of the invention encompasses use of conventional assay means to determine gene expression in a canine either alone or in conjunction with gene expression array displays employing polypeptides and/or polynucleotides, such conventional assay means comprising one or more of ELISA, RIA, immunoblot assays, in situ hybridization, Northern blot analysis, Western blot analysis and Luminex X-Map® analysis.

Another embodiment of the invention encompasses diagnosing a kidney disorder characterized by an abnormal loss of renal function, renal failure, reduced glomerular filtration rate or glomerulonephritis over time by determining gene expression profiles for one or more genes or their expression products selected from at least one RNA transcript or its translation product selected from the group consisting of: secreted frizzled-related protein 2 (sFRP2); matrilin-2 (Matn2); lumican (LUM); decorin (DCN); collagen alpha 1 (III) chain, variant 12 (COL3A1); retinol binding protein 4 (rbp4); MMP-9; clusterin (CLU); transferrin (TF); Apo-C-1 (ApoC1); and inhibin beta A (INHBA) in conjunction with one or more conventional diagnostic measurements selected from the group consisting of: determining levels of albumin, serum cystatin C, blood creatinine, urinary creatinine, creatinine clearance, urinary protein, blood urea nitrogen and glomerular filtration rate, to choose a course of treatment for such canine.

In another embodiment, the invention features a kit for diagnosing and/or monitoring a kidney disorder in a canine

over time, the kit comprising one or more agents capable of detecting expression profiles for one or more genes or their expression products selected from at least one RNA transcript or its translation product selected from the group consisting of: secreted frizzled-related protein 2 (sFRP2); matrilin-2 (Matn2); lumican (LUM); decorin (DCN); collagen alpha 1 (III) chain, variant 12 (COL3A1); retinol binding protein 4 (rbp4); MMP-9; clusterin (CLU); transferrin (TF); Apo-C-1 (ApoC1); and inhibin beta A (INHBA), and instructions for using said one or more agents to evaluate the risk of such canine developing a kidney disorder characterized by an abnormal loss of renal function, renal failure, reduced glomerular filtration rate or glomerulonephritis over time.

In another aspect, the invention includes contacting a tissue sample or bodily fluid specimen with an agent that detects in one or more genes or their expression products selected from at least one RNA transcript or its translation product selected from the group consisting of: secreted frizzled-related protein 2 (sFRP2); matrilin-2 (Matn2); lumican (LUM); decorin (DCN); collagen alpha 1 (III) chain, variant 12 (COL3A1); retinol binding protein 4 (rbp4); MMP-9; clusterin (CLU); transferrin (TF); Apo-C-1 (ApoC1); and inhibin beta A (INHBA). The agent can be an antibody or a nucleic acid probe used in conjunction with conventional assay means such as immobilization on a solid phase, microtiter wells, tubes, dipsticks or other conventional means.

Particularly preferred embodiments of the present invention follow below. In each of the following embodiments, the present invention contemplates a test sample comprising either solid kidney tissue samples, or samples of biological fluids from a canine. Differential expression of the recited genes contemplates a significant difference in absolute terms. In preferred embodiments the differential expression may be greater than about 1.1, 1.2, 1.3, 1.4, 1.5, 1.6, 1.7, 1.8, 1.9, 2.0, 2.1, 2.2, 2.3, 2.4, 2.5, or 3.0 fold or greater. Especially preferred differential expression values are greater than about 2 fold or one standard deviation around the mean. Also, the embodiments of the invention may employ a variety of art-recognized detection means, including an array of one or more hybridization probes, panels of one or more antibodies and combinations of these technologies. When immunoassays are desired, the skilled worker can select among many art-recognized techniques, including immunoassays selected from the group consisting of a competitive binding assay, a non-competitive binding assay, a radioimmunoassay, an enzyme linked immunosorbent assay (ELISA), a sandwich assay, a precipitin reaction, a gel diffusion immunodiffusion assay, an agglutination assay, a fluorescent immunoassay, chemiluminescence immunoassay, immunoPCR immunoassay, a protein A or protein G immunoassay and an immunoelectrophoresis assay. When conventional diagnostic measurements are specified, these may be selected from the group consisting of: determining levels of albumin, serum cystatin C, blood creatinine, urinary creatinine, creatinine clearance, urinary protein, blood urea nitrogen and glomerular filtration rate

In addition, the methods and compositions of the invention contemplate utilizing at least one RNA transcript or its translation product which may be an altered form of the gene or its translation product.

A still further embodiment of the invention comprises a method for diagnosing a kidney disorder in a canine, comprising the steps of: (a) measuring the level of expression of at least one biomarker in a biological sample from the canine, wherein the at least one biomarker is selected from

at least one gene or the translation product of such gene selected from the group consisting of secreted frizzled-related protein-2 (SFRP2); matrilin-2 (Matn2); lumican (LUM); decorin (DCN); collagen alpha 1 (III) chain, variant 12 (COL3A1); retinol binding protein 4 (rbp4); MMP-9; clusterin (CLU); transferrin (TF); Apo-C-1 (ApoC1); and inhibin beta A (INHBA); and (b) identifying the canine as having a kidney disorder wherein difference in expression of the one or more biomarkers in the sample relative to a control value for expression in a sample from a normal animal indicates the existence of a kidney disorder.

Another embodiment comprises a kit for the diagnosis, prognosis or monitoring a kidney disorder in a canine, comprising at least one or more agent capable of detecting at least one or more RNA transcript or its translation product selected from a first group of at least one gene or the translation product of such gene selected from the group consisting of: secreted frizzled-related protein-2 (SFRP2); matrilin-2 (Matn2); lumican (LUM); decorin (DCN); collagen alpha 1 (III) chain, variant 12 (COL3A1); retinol binding protein 4 (rbp4); MMP-9; clusterin (CLU); transferrin (TF); Apo-C-1 (ApoC1); and inhibin beta A (INHBA); and instructions for using such at least one or more agent to measure the expression of the one or more biomarkers and to diagnose a kidney disease in such canine.

In a further aspect, the invention relates to compositions comprising one or more nucleic acid probes that specifically hybridize to a nucleic acid, or fragment thereof, encoding a biomarker of the present invention selected from the group consisting of: secreted frizzled-related protein-2 (SFRP2); matrilin-2 (Matn2); lumican (LUM); decorin (DCN); collagen alpha 1 (III) chain, variant 12 (COL3A1); retinol binding protein 4 (rbp4); matrix metalloproteinase 9 (MMP9); clusterin (CLU); transferrin (TF); Apo-C-1 (ApoC1); and inhibin beta A (INHBA).

In an additional aspect, the invention relates to compositions comprising antibodies that specifically bind to a polypeptide encoded by a gene expressing a biomarker of the present invention selected from the group consisting of: secreted frizzled-related protein-2 (SFRP2); matrilin-2 (Matn2); lumican (LUM); decorin (DCN); collagen alpha 1 (III) chain, variant 12 (COL3A1); retinol binding protein 4 (rbp4); matrix metalloproteinase 9 (MMP9); clusterin (CLU); transferrin (TF); Apo-C-1 (ApoC1); and inhibin beta A (INHBA).

It is further contemplated herein that the methods of the present invention may be used in combination with traditional diagnostic techniques that are able to detect the physical and morphological characteristics of kidney disorders. Thus, for example, the characterization of differential expression in genes for kidney in cells obtained from a tissue samples or bodily fluid specimens of a canine may be combined with conventional diagnostic (e.g., radiological) techniques in order to corroborate a diagnosis of a kidney disorder in a canine, including, for example, glomerulonephritis.

The invention also relates to kits to diagnose an abnormal kidney disorder in a canine comprising components that can be used to detect expression of the biomarkers of the present invention, including, but not limited to, the compositions and microarrays described herein.

A further aspect of the invention is a method for diagnosis and/or prognosis of kidney disorder in a canine, wherein the method comprises the steps of: obtaining at least one tissue sample or bodily fluid specimen from the animal; determining the amount of one or more biomarkers selected from Table 3 in said at least one sample or specimen obtained

from the animal, wherein said biomarker is a polypeptide, protein, RNA, DNA, polynucleotide or metabolite thereof.

Yet another embodiment of the invention is a kit for diagnosis and/or prognosis of a kidney disorder in a canine, particularly for carrying out the method for diagnosis and/or prognosis of glomerulonephritis in a canine wherein the method comprises the steps of: obtaining at least one tissue sample or bodily fluid specimen from the animal; determining the amount of one or more biomarkers selected from Table 3 in said at least one sample or specimen obtained from the animal, wherein said biomarker is a polypeptide, protein, RNA, DNA, polynucleotide or metabolite thereof, and optionally, further comprising a detectable agent linked to said biomarker.

A still further embodiment of the invention is a reagent for diagnosis and/or prognosis of glomerulonephritis in a canine particularly for carrying out the method for diagnosis and/or prognosis of glomerulonephritis in a canine, wherein the method comprises the steps of: obtaining at least one tissue sample or bodily fluid specimen from the animal; determining the amount of one or more biomarkers selected from Table 3 in said at least one sample or specimen obtained from a canine, wherein said biomarker is a polypeptide, protein, RNA, DNA, polynucleotide or metabolite thereof, and optionally further comprising a detectable agent linked to said biomarker.

A further aspect of the invention is a method for diagnosis and/or prognosis of kidney disorder in a canine, wherein the method comprises the steps of: obtaining at least one tissue sample or bodily fluid specimen from the animal; determining the amount of one or more biomarkers selected from Tables 3 and 4 in said at least one sample or specimen obtained from the animal, wherein said biomarker is a polypeptide, protein, RNA, DNA, polynucleotide or metabolite thereof.

Another embodiment of the invention is the use of one or more polypeptides, proteins, RNAs, DNAs, polynucleotides or metabolites thereof selected from the group consisting of secreted frizzle-related protein-2 (SFRP2); matrilin-2 (Matn2); lumican (LUM); decorin (DCN); collagen alpha 1 (III) chain, variant 12 (COL3A1); retinol binding protein 4 (rbp4); matrix metalloproteinase 9 (MMP9); clusterin (CLU); transferrin (TF); Apo-C-1 (ApoC1); and inhibin beta A (INHBA) as a biomarker for diagnosis and/or prognosis of a kidney disorder, particularly for forming a kit for diagnosis or prognosis of a kidney disorder in a canine.

Another embodiment of the invention is the use of one or more polypeptides, proteins, RNAs, DNAs, polynucleotides or metabolites thereof selected from the group consisting of: secreted frizzle-related protein-2 (SFRP2); matrilin-2 (Matn2); lumican (LUM); decorin (DCN); collagen alpha 1 (III) chain, variant 12 (COL3A1); retinol binding protein 4 (rbp4); matrix metalloproteinase 9 (MMP9); clusterin (CLU); transferrin (TF); Apo-C-1 (ApoC1); and inhibin beta A (INHBA) as a biomarker for diagnosis and/or prognosis of a kidney disorder, particularly for forming a kit for diagnosis or prognosis of a kidney disorder in a canine.

Yet another embodiment is such kit, wherein the reagents and equipment comprise DNA microarray analysis materials including oligonucleotide microarray, c-DNA microarray, and focused gene chip, or a combination thereof.

Another embodiment of the invention is a method of assessing the progression of a course of nutritional management for a canine suffering from a kidney disorder, the method comprising (a) measuring a first level of one or more biomarker polypeptides, proteins, RNAs, DNAs, polynucleotides or metabolites thereof selected from the group con-

sisting of: secreted frizzle-related protein-2 (SFRP2); matrilin-2 (Matn2); lumican (LUM); decorin (DCN); collagen alpha 1 (III) chain, variant 12 (COL3A1); retinol binding protein 4 (rbp4); matrix metalloproteinase 9 (MMP9); clusterin (CLU); transferrin (TF); Apo-C-1 (ApoC1); and inhibin beta A (INHBA) in a tissue sample or a specimen of bodily fluid from said canine at a first point in time during the course of treatment, (b) measuring a second level of said biomarker in said sample or specimen from said canine at a second point in time during the course of treatment, and (c) comparing the measurements of the biomarker at said first point and said second point; wherein expression of the biomarker has at least about a 1.0, 1.1, 1.2, 1.3, 1.4, 1.5, 1.6, 1.7, 1.8, 1.9, 2.0, 2.5, or 3.0 fold differential or greater difference in gene expression compared with expression in a cell of a control animal.

Still another embodiment of the invention is a method for identifying a plurality of genes that are differentially expressed between tissue samples or specimens of a bodily fluid for use in an informative array, comprising: providing a first set of heterogeneous nucleic acid probes derived from a first tissue sample; providing a second set of heterogeneous nucleic acid probes derived from a second tissue sample; hybridizing a nucleic acid array comprising a plurality of sequences derived from genes of a biological process with the first set of probes and determining a first level of expression for sequences of the array; hybridizing the array with said second set of probes and determining a second level of expression for sequences of the array; identifying a plurality of genes that are differentially expressed in said biological process by comparing the first level of expression with said second level of expression for hybridized sequences; and establishing a ranking of the identified genes by a step selected from the group of steps consisting of: determining an absolute value of the difference between the first level of expression and the second level of expression, and ranking genes having a higher difference over genes having a lower difference; and determining a standard deviation of the difference between the first level of expression and the second level of expression, and ranking genes having a higher standard deviation over genes having a lower standard deviation, wherein the genes are selected from the group consisting of: secreted frizzle-related protein-2 (SFRP2); matrilin-2 (Matn2); lumican (LUM); decorin (DCN); collagen alpha 1 (III) chain, variant 12 (COL3A1); retinol binding protein 4 (rbp4); matrix metalloproteinase 9 (MMP9); clusterin (CLU); transferrin (TF); Apo-C-1 (ApoC1); and inhibin beta A (INHBA).

Another embodiment of the invention is a method for converting a nucleic acid array into an informative array, comprising: providing a first set of heterogeneous nucleic acid probes derived from a first tissue sample; providing a different, second set of heterogeneous nucleic acid probes derived from a second tissue sample; hybridizing a nucleic acid array comprising a plurality of sequences with the first set of probes and determining a first level of expression for sequences of the array; hybridizing the array with said second set of probes and determining a second level of expression for sequences of the array; identifying a plurality of genes that are differentially expressed in said biological process based on a difference between the first level of expression and the second level of expression for identified genes, by a step selected from the group of steps consisting of: determining an absolute value for the difference between the first level of expression and the second level of expression, and ranking genes having a higher difference over genes having a lower difference; and determining a standard

deviation of the difference between the first level of expression and the second level of expression, and ranking genes having a higher standard deviation over genes having a lower standard deviation; and selecting genes from the plurality of identified differentially expressed genes for inclusion on the informative array, wherein said genes are selected from the group consisting of: secreted frizzle-related protein-2 (SFRP2) and matrilin-2 (Matn2); and, optionally, one or more genes selected from the group consisting of: lumican (LUM); decorin (DCN); collagen alpha 1 (III) chain, variant 12 (COL3A1); retinol binding protein 4 (rbp4); matrix metalloproteinase 9 (MMP9); clusterin (CLU); transferrin (TF); Apo-C-1 (ApoC1); and inhibin beta A (INHBA).

Another embodiment of the invention is a method of determining the efficacy of a treatment for a kidney disorder, comprising the steps of: (a) providing a biological sample from a canine affected by a kidney disorder, who has been subjected to said treatment, (b) determining the level in said sample of one or more biomarkers for glomerulonephritis, to create an expression profile for said canine, and (c) comparing said expression profile with: i) a comparable expression profile obtained from said test canine before initiation of said treatment, and/or ii) a comparable expression profile obtained from said test canine at an earlier stage of said treatment, and/or iii) a comparable expression profile characteristic of a subject who is unaffected by a kidney disorder, wherein the one or more biomarkers for a kidney disorder, comprise expression products of one or more gene selected from the group consisting of secreted frizzle-related protein-2 (SFRP2) and matrilin-2 (Matn2); and, optionally, one or more genes selected from the group consisting of: lumican (LUM); decorin (DCN); collagen alpha 1 (III) chain, variant 12 (COL3A1); retinol binding protein 4 (rbp4); matrix metalloproteinase 9 (MMP9); clusterin (CLU); transferrin (TF); Apo-C-1 (ApoC1); and inhibin beta A (INHBA).

Other and further objects, features, and advantages of the present invention will be readily apparent to those skilled in the art.

#### DETAILED DESCRIPTION OF THE INVENTION

As used herein, an abnormal canine is a canine that has a risk of developing, predisposition for, or an abnormal loss of kidney function, renal failure, reduced glomerular filtration rate or glomerulonephritis, wherein the abnormality is detectable by utilizing at least one relevant biomarker isolated and measured from a biological specimen taken from such canine.

Throughout this disclosure, various aspects of this invention can be presented in a range format. It should be understood that the description in range format is merely for convenience and brevity and should not be construed as an inflexible limitation on the scope of the invention. Accordingly, the description of a range should be considered to have specifically disclosed all the possible subranges as well as individual numerical values within that range. For example, description of a range such as from 1 to 5 should be considered to have specifically disclosed subranges such as from 1 to 3, from 1 to 4, from 2 to 4, from 2 to 5, from 3 to 5 etc., as well as individual numbers within that range, for example, 1, 2, 3, 4, and 5. This applies regardless of the breadth of the range.

The practice of the present invention may employ, unless otherwise indicated, conventional techniques and descriptions of organic chemistry, polymer technology, molecular

biology (including recombinant techniques), cell biology, biochemistry, and immunology, which are within the skill of the art. Such conventional techniques include polymer array synthesis, hybridization, ligation, and detection of hybridization using a label.

The present invention also contemplates many uses for polymers attached to solid substrates. These uses include gene expression monitoring, profiling, library screening, genotyping and diagnostics.

Those skilled in the art will recognize that the products and methods embodied in the present invention may be applied to a variety of systems, including commercially available gene expression monitoring systems involving nucleic acid probe arrays, membrane blots, microwells, beads and sample tubes, constructed with various materials using various methods known in the art. Accordingly, the present invention is not limited to any particular environment, and the following description of specific embodiments of the present invention are for illustrative purposes only.

The gene expression monitoring system, in a preferred embodiment, may comprise a nucleic acid probe array (including an oligonucleotide array, a cDNA array, a spotted array, and the like), membrane blot (such as used in hybridization analysis such as Northern, Southern, dot, and the like), or microwells, sample tubes, beads or fibers (or any solid support comprising bound nucleic acids). The gene expression monitoring system may also comprise nucleic acid probes in solution.

The present invention also contemplates sample preparation involving amplification. A genomic sample may be amplified by a variety of mechanisms, some of which may employ PCR. The sample may be amplified on the array.

Other suitable amplification methods include the ligase chain reaction (LCR) (e.g., Wu and Wallace, *Genomics* 4, 560 (1989), Landegren et al., *Science* 241, 1077 (1988) and Barringer et al. *Gene* 89:117 (1990)), transcription amplification (Kwoh et al., *Proc. Natl. Acad. Sci. USA* 86, 1173 (1989) and WO88/10315), self-sustained sequence replication (Guatelli et al., *Proc. Nat. Acad. Sci. USA*, 87, 1874 (1990) and WO90/06995), selective amplification of target polynucleotide sequences (U.S. Pat. No. 6,410,276), consensus sequence primed polymerase chain reaction (CP-PCR) (U.S. Pat. No. 4,437,975), arbitrarily primed polymerase chain reaction (AP-PCR) (U.S. Pat. Nos. 5,413,909, 5,861,245) and nucleic acid based sequence amplification (NABSA).

The gene expression monitoring system according to the present invention may be used to facilitate a comparative analysis of expression in different cells or tissues, different subpopulations of the same cells or tissues, different physiological states of the same cells or tissue, different developmental stages of the same cells or tissue, or different cell populations of the same tissue. In a preferred embodiment, the proportional amplification methods of the present invention can provide reproducible results (i.e., within statistically significant margins of error or degrees of confidence) sufficient to facilitate the measurement of quantitative as well as qualitative differences in the tested samples.

The term "antibody" means any immunoglobulin that binds to a specific antigen, including IgG, IgM, IgA, IgD, and IgE antibodies. The term includes polyclonal, monoclonal, monovalent, humanized, heteroconjugate, antibody compositions with polyepitopic specificity, chimeric, bispecific antibodies, diabodies, single-chain antibodies and antibody fragments such as Fab, Fab', F(ab')<sub>2</sub>, and Fv, or other antigen-binding fragments.

The term "array" means an ordered arrangement of at least two probes on a substrate. At least one of the probes is a control or standard and at least one of the probes is a diagnostic probe. The arrangement of from about two to about 40,000 probes on a substrate assures that the size and signal intensity of each labeled complex formed between a probe and a sample polynucleotide or polypeptide is individually distinguishable. The collection of molecules deposited on the array may be prepared either synthetically or biosynthetically. The array may take a variety of forms including libraries of soluble molecules, libraries of compounds tethered to resin beads, silica chips or other solid supports. The nucleic acid array may include libraries of nucleic acids which can be prepared by spotting nucleic acids in essentially any length (for example, from 1 to about 1,000 nucleotides in length) onto a substrate. A nucleic acid probe array preferably comprises nucleic acids bound to a substrate in known locations. In other embodiments, the system may include a solid support or substrate, such as a membrane, filter, microscope slide, microwell, sample tube, bead, bead array, or the like. The solid support may be made of various materials, including paper, cellulose, nylon, polystyrene, polycarbonate, plastics, glass, ceramic, stainless steel, or the like. The solid support may preferably have a rigid or semi-rigid surface, and may preferably be spherical (e.g., bead) or substantially planar (e.g., flat surface) with appropriate wells, raised regions, etched trenches, or the like. The solid support may also include a gel or matrix in which nucleic acids may be embedded.

The term "biomarker" refers to genes and gene products encoded by the gene of the invention, i.e., where the gene has been determined to have been differentially regulated as a result of a kidney disorder. In addition, the term can be generally used to refer to any portion of such gene or protein that can identify or correlate with the full-length gene or protein, for example, in an assay or other method of the invention.

Biomarker expression can also be identified by detection of biomarker translation (i.e., detection of biomarker protein in a sample). Methods suitable for the detection of biomarker protein include any suitable method for detecting and/or measuring proteins from a cell or cell extract. Such methods include, but are not limited to, immunoblot (e.g., Western blot), enzyme-linked immunosorbent assay (ELISA), radioimmunoassay (RIA), immunoprecipitation, immunohistochemistry and immunofluorescence. Particularly preferred methods for detection of proteins include any single-cell assay, including immunohistochemistry and immunofluorescence assays. Such methods are well known in the art. Furthermore, antibodies against certain of the biomarkers described herein are known in the art and are described in the public literature, and methods for their preparation are well known to the skilled worker.

The term "differential expression" or "differentially expressed" means increased or unregulated gene expression or means decreased or downregulated gene expression as detected by the absence, presence, or at least two-fold, or at least a 2.0, 1.9, 1.8, 1.7, 1.6, 1.5, 1.4, 1.3, 1.2, 1.1 or 1.0-fold change in the amount of transcribed messenger RNA or translated protein in a sample.

The term "fold" when used as a measure of differential gene expression means an amount of gene expression in a canine that is a multiple or a fraction of gene expression compared to the amount of gene expression against a control value. For example, a gene that is expressed 2 times as much in the animal as compared to the control value has a 2-fold differential gene expression and a gene that is expressed

one-third as much in the animal as compared to the control value also has a 2-fold differential gene expression.

The term "fragment" means (1) an oligonucleotide or polynucleotide sequence that is a portion of a complete sequence and that has the same or similar activity for a particular use as the complete polynucleotide sequence or (2) a peptide or polypeptide sequence that is a portion of a complete sequence and that has the same or similar activity for a particular use as the complete polypeptide sequence. Such fragments can comprise any number of nucleotides or amino acids deemed suitable for a particular use. Generally, oligonucleotide or polynucleotide fragments contain at least about 10, 50, 100, or 1000 nucleotides and polypeptide fragments contain at least about 4, 10, 20, or 50 consecutive amino acids from the complete sequence. The term encompasses polynucleotides and polypeptides variants of the fragments. A polynucleotide, for example, can be broken up, or fragmented into, a plurality of segments.

Various methods of fragmenting nucleic acid are well known in the art. These methods may be, for example, either chemical or physical in nature. Chemical fragmentation may include partial degradation with a DNase; partial depurination with acid; the use of restriction enzymes; intron-encoded endonucleases; DNA-based cleavage methods, such as triplex and hybrid formation methods, that rely on the specific hybridization of a nucleic acid segment to localize a cleavage agent to a specific location in the nucleic acid molecule; or other enzymes or compounds which cleave DNA at known or unknown locations. Physical fragmentation methods may involve subjecting the DNA to a high shear rate. High shear rates may be produced, for example, by moving DNA through a chamber or channel with pits or spikes, or forcing the DNA sample through a restricted size flow passage, e.g., an aperture having a cross sectional dimension in the micron or submicron scale. Other physical methods include sonication and nebulization. Combinations of physical and chemical fragmentation methods may likewise be employed such as fragmentation by heat and ion-mediated hydrolysis. These methods can be optimized to digest a nucleic acid into fragments of a selected size range. Useful size ranges may be from 100, 200, 400, 700 or 1000 to 500, 800, 1500, 2000, 4000 or 10,000 base pairs. However, larger size ranges such as 4000, 10,000 or 20,000 to 10,000, 20,000 or 500,000 base pairs may also be useful.

The term "gene" or "genes" means a complete or partial segment of DNA involved in producing a polypeptide, including regions preceding and following the coding region (leader and trailer) and intervening sequences (introns) between individual coding segments (exons). The term encompasses any DNA sequence that hybridizes to the complement of gene coding sequences.

The term "homolog" means (1) a polynucleotide, including polynucleotides from the same or different animal species, having greater than 30%, 50%, 70%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% sequence similarity to a polynucleotide and having the same or substantially the same properties and performing the same or substantially the same function as the complete polynucleotide, or having the capability of specifically hybridizing to a polynucleotide under stringent conditions or (2) a polypeptide, including polypeptides from the same or different animal species, having greater than 30%, 50%, 70%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% sequence similarity to a polypeptide identified by the expression of polynucleotides and having the same or substantially the same properties and performing the same or substantially

the same function as the complete polypeptide, or having the capability of specifically binding to a polypeptide identified by the expression of polynucleotides. Sequence similarity of two polypeptide sequences or of two polynucleotide sequences is determined using methods known to skilled artisans, e.g., the algorithm of Karlin and Altschul (Proc. Natl. Acad. Sci. USA 87:2264-2268 (1990)). Such an algorithm is incorporated into the NBLAST and XBLAST programs of Altschul et al. (J. Mol. Biol. 215:403-410 (1990)). To obtain gapped alignments for comparison purposes, Gapped Blast can be utilized as described in Altschul et al. (Nucl. Acids Res. 25: 3389-3402 (1997)). When utilizing BLAST and Gapped BLAST programs, the default parameters of the respective programs (e.g., XBLAST and NBLAST) are used. See <http://www.ncbi.nlm.nih.gov>.

The term "hybridization" refers to the process in which two single-stranded polynucleotides bind non-covalently to form a stable double-stranded polynucleotide. The term "hybridization" may also refer to triple-stranded hybridization. The resulting (usually) double-stranded polynucleotide is a "hybrid." The proportion of the population of polynucleotides that forms stable hybrids is referred to herein as the "degree of hybridization".

Hybridization reactions can be performed in absolute or differential hybridization formats. In the absolute hybridization format, polynucleotides derived from one sample are hybridized to the probes in a nucleic acid array. Signals detected after the formation of hybridization complexes correlate to the polynucleotide levels in the sample. In the differential hybridization format, polynucleotides derived from two samples are labeled with different labeling moieties. A mixture of these differently labeled polynucleotides is added to a nucleic acid array. The nucleic acid array is then examined under conditions in which the emissions from the two different labels are individually detectable. In one embodiment, the fluorophores Cy3 and Cy5 (Amersham Pharmacia Biotech, Piscataway, N.J.) are used as the labeling moieties for the differential hybridization format.

Signals gathered from nucleic acid arrays can be analyzed using commercially available software, such as those provided by Affymetrix or Agilent Technologies. Controls, such as for scan sensitivity, probe labeling and cDNA or cRNA quantization, are preferably included in the hybridization experiments. Hybridization signals can be scaled or normalized before being subject to further analysis. For instance, hybridization signals for each individual probe can be normalized to take into account variations in hybridization intensities when more than one array is used under similar test conditions. Hybridization signals can also be normalized using the intensities derived from internal normalization controls contained on each array. In addition, genes with relatively consistent expression levels across the samples can be used to normalize the expression levels of other genes. In one embodiment, probes for certain maintenance genes are included in a nucleic acid array of the present invention. These genes are chosen because they show stable levels of expression across a diverse set of tissues. Hybridization signals can be normalized and/or scaled based on the expression levels of these maintenance genes.

The term "hybridization complex" means a complex that is formed between sample polynucleotides when the purines of one polynucleotide hydrogen bond with the pyrimidines of the complementary polynucleotide, e.g., 5'-A-G-T-C-3' base pairs with 3'-T-C-A-G-5'. The degree of complementarity and the use of nucleotide analogs affect the efficiency and stringency of hybridization reactions.

The term "hybridization probes" includes nucleic acids (such as oligonucleotides) capable of binding in a base-specific manner to a complementary strand of nucleic acid.

The term "kidney disease" or "kidney disorder" or analogously "renal disease" or "renal disorder" is intended to cover an acute or chronic abnormal loss of kidney function, such as renal failure, reduced glomerular filtration rate and glomerulonephritis. Glomerulonephritis may take the form of membranous glomerulonephritis which involves thickening of the glomerular basement membrane. Alternatively, glomerulonephritis may take the form of proliferative or mesangioproliferative glomerulonephritis, which is characterized by proliferation of cells in the mesangial matrix. In addition, glomerulonephritis may take the form of membranoproliferative glomerulonephritis which involves a combination of the foregoing changes. Glomerulosclerosis is a severe form of glomerulonephritis. Kidney disease or kidney disorders also include nephritis, nephropathy, hyperfiltration, mild microalbuminuria, clinical albuminuria, advanced clinical nephropathy, chronic renal insufficiency, injuries to renal papilla, tubular necrosis and diabetic nephropathy, all as differentially diagnosed by veterinarians of ordinary skill in the art. The term is not intended to encompass polycystic kidney disease of genetic origin.

The term "polynucleotide" or "oligonucleotide" means a polymer of nucleotides. The term encompasses DNA and RNA (including cDNA and mRNA) molecules, either single or double stranded and, if single stranded, its complementary sequence in either linear or circular form. The term also encompasses fragments, variants, homologs, and alleles, as appropriate for the sequences that have the same or substantially the same properties and perform the same or substantially the same function as the original sequence. The sequences may be fully complementary (no mismatches) when aligned or may have up to about a 30% sequence mismatch. Preferably, for polynucleotides, the chain contains from about 50 to 10,000 nucleotides, more preferably from about 150 to 3,500 nucleotides. Preferably, for oligonucleotides, the chain contains from about 2 to 100 nucleotides, more preferably from about 6 to 30 nucleotides. The exact size of a polynucleotide or oligonucleotide will depend on various factors and on the particular application and use of the polynucleotide or oligonucleotide. The term includes nucleotide polymers that are synthesized and that are isolated and purified from natural sources. The term "polynucleotide" is inclusive of "oligonucleotide."

The term "polypeptide," "peptide," or "protein" means a polymer of amino acids. The term encompasses naturally occurring and non-naturally occurring (synthetic) polymers and polymers in which artificial chemical mimetics are substituted for one or more amino acids. The term also encompasses fragments, variants, and homologs that have the same or substantially the same properties and perform the same or substantially the same function as the original sequence. The term encompasses polymers of any length, preferably polymers containing from about 2 to 1000 amino acids, more preferably from about 5 to 500 amino acids. The term includes amino acid polymers that are synthesized and that are isolated and purified from natural sources.

The term "probe" means (1) an oligonucleotide or polynucleotide, either RNA or DNA, whether occurring naturally as in a purified restriction enzyme digest or produced synthetically, that is capable of annealing with or specifically hybridizing to a polynucleotide with sequences complementary to the probe or (2) a peptide or polypeptide capable of specifically binding a particular protein or protein fragment to the substantial exclusion of other proteins or protein

fragments. An oligonucleotide or polynucleotide probe may be either single or double stranded. The exact length of the probe will depend upon many factors, including temperature, source, and use. For example, for diagnostic applications, depending on the complexity of the target sequence, an oligonucleotide probe typically contains about 10 to 100, 15 to 50, or 15 to 25 nucleotides. In certain diagnostic applications, a polynucleotide probe contains about 100-1000, 300-600, nucleotides, preferably about 300 nucleotides. The probes herein are selected to be "substantially" complementary to different strands of a particular target sequence. This means that the probes must be sufficiently complementary to specifically hybridize or anneal with their respective target sequences under a set of predetermined conditions. Therefore, the probe sequence need not reflect the exact complementary sequence of the target. For example, a noncomplementary nucleotide fragment may be attached to the 5' or 3' end of the probe, with the remainder of the probe sequence being complementary to the target sequence. Alternatively, noncomplementary bases or longer sequences can be interspersed into the probe provided that the probe sequence has sufficient complementarity with the sequence of the target polynucleotide to specifically anneal to the target polynucleotide. A peptide or polypeptide probe may be any molecule to which the protein or peptide specifically binds, including DNA (for DNA binding proteins), antibodies, cell membrane receptors, peptides, cofactors, lectins, sugars, polysaccharides, cells, cell membranes, organelles and organellar membranes.

The terms "sample" and "specimen" mean any animal tissue or fluid containing, e.g., polynucleotides, polypeptides, antibodies, metabolites, and the like, including cells and other tissue containing DNA and RNA. Examples include: blood, cartilage, connective, epithelial, lymphoid, muscle, nervous, sputum, and the like. A sample may be solid or liquid and may be DNA, RNA, cDNA, bodily fluids such as blood or urine, cells, cell preparations or soluble fractions or media aliquots thereof, chromosomes, organelles, and the like.

The term "single package" means that the components of a kit are physically associated in or with one or more containers and considered a unit for manufacture, distribution, sale, or use. Containers include, but are not limited to, bags, boxes, bottles, shrink wrap packages, stapled or otherwise affixed components, or combinations thereof. A single package may be containers of individual food compositions physically associated such that they are considered a unit for manufacture, distribution, sale, or use.

The "specifically bind" means a special and precise interaction between two molecules which is dependent upon their structure, particularly their molecular side groups. For example, the intercalation of a regulatory protein into the major groove of a DNA molecule, the hydrogen bonding along the backbone between two single stranded nucleic acids, or the binding between an epitope of a protein and an agonist, antagonist, or antibody.

The term "specifically hybridize" means an association between two single stranded polynucleotides of sufficiently complementary sequence to permit such hybridization under predetermined conditions generally used in the art (sometimes termed "substantially complementary"). For example, the term may refer to hybridization of a polynucleotide probe with a substantially complementary sequence contained within a single stranded DNA or RNA molecule according to an aspect of the invention, to the substantial

exclusion of hybridization of the polynucleotide probe with single stranded polynucleotides of non-complementary sequence.

The term "stringent conditions" means (1) hybridization in 50% (vol/vol) formamide with 0.1% bovine serum albumin, 0.1% Ficoll, 0.1% polyvinylpyrrolidone, 50 mM sodium phosphate buffer at pH 6.5 with 750 mM NaCl, 75 mM sodium citrate at 42° C., (2) hybridization in 50% formamide, 5xSSC (0.75 M NaCl, 0.075 M sodium citrate), 50 mM sodium phosphate (pH 6.8), 0.1% sodium pyrophosphate, 5xDenhardt's solution, sonicated salmon sperm DNA (50 µg/ml), 0.1% SDS, and 10% dextran sulfate at 42° C.; with washes at 42° C. in 0.2xSSC and 0.1% SDS or washes with 0.015 M NaCl, 0.0015 M sodium citrate, 0.1% Na<sub>2</sub>SO<sub>4</sub> at 50° C. or similar procedures employing similar low ionic strength and high temperature washing agents and similar denaturing agents.

The term "substance" means an element, compound, molecule, or a mixture thereof or any other material that could potentially be useful for diagnosing, prognosing, or modulating the onset or severity of a kidney disorder in an animal, including any drug, chemical entity, or biologic entity.

The term "useful variations" means (1) for a polynucleotide, the complements of the polynucleotide; the homologs of the polynucleotide and its complements; the variants of the polynucleotide, its complements, and its homologs; and the fragments of the polynucleotide, its complements, its homologs, and its variants and (2) for a polypeptide, the homologs of the polypeptide; the variants of the polypeptide and its homologs; and the fragments of the polynucleotide, its homologs, and its variants.

The term "variant" means (1) a polynucleotide sequence containing any substitution, variation, modification, replacement, deletion, or addition of one or more nucleotides from or to a polynucleotide sequence and that has the same or substantially the same properties and performs the same or substantially the same function as the original sequence and (2) a polypeptide sequence containing any substitution, variation, modification, replacement, deletion, or addition of one or more amino acids from or to a polypeptide sequence and that has the same or substantially the same properties and performs the same or substantially the same function as the original sequence. The term therefore includes single nucleotide polymorphisms (SNPs) and allelic variants and includes conservative and non-conservative amino acid substitutions in polypeptides. The term also encompasses chemical derivatization of a polynucleotide or polypeptide and substitution of nucleotides or amino acids with nucleotides or amino acids that do not occur naturally, as appropriate.

The term "virtual package" means that the components of a kit are associated by directions on one or more physical or virtual kit components instructing the user how to obtain the other components, e.g., in a bag containing one component and directions instructing the user to go to a website, contact a recorded message, view a visual message, or contact a caregiver or instructor to obtain instructions on how to use the kit.

The invention is not limited to the particular methodology, protocols, and reagents described herein because they may vary. Further, the terminology used herein is for the purpose of describing particular embodiments only and is not intended to limit the scope of the present invention. As used herein and in the appended claims, the singular forms "a," "an," and "the" include plural references unless the context clearly dictates otherwise, e.g., reference to "a variant"

includes a plurality of variants. Further, defined terms include variations of the terms used in the proper grammatical context, e.g., the term “specifically binds” includes “specific binding” and other forms of the term. Similarly, the words “comprise,” “comprises,” and “comprising” are to be interpreted inclusively rather than exclusively.

Unless defined otherwise, all technical and scientific terms and any acronyms used herein have the same meanings as commonly understood by one of ordinary skill in the art in the field of the invention. Although any compositions, methods, articles of manufacture, or other means or materials similar or equivalent to those described herein can be used in the practice of the present invention, the preferred compositions, methods, articles of manufacture, or other means or materials are described herein.

All patents, patent applications, publications, and other references cited or referred to herein are incorporated herein by reference to the extent allowed by law. The discussion of those references is intended merely to summarize the assertions made therein. No admission is made that any such patents, patent applications, publications or references, or any portion thereof, is relevant prior art for the present invention and the right to challenge the accuracy and pertinence of such patents, patent applications, publications, and other references is specifically reserved.

In one embodiment, the present invention encompasses one or more genes or gene segments (“genes” as defined herein) that are differentially expressed in abnormal animals compared to a control value obtained from one or more normal animals. The invention is based upon the discovery of polynucleotides that are differentially expressed in abnormal animals compared to normal animals. The genes were identified by comparing the expression of genes in tissue samples taken from animals diagnosed as abnormal with genes in tissue samples from animals diagnosed as normal using Affymetrix GeneChip® technology.

The polynucleotides and genes are identified by measuring differences in gene expression from tissue samples taken from canines diagnosed as abnormal and having a kidney disorder against a control value, wherein the control value is obtained from a sample from one or more normal animals. Changes in gene expression can be determined by any method known to skilled artisans. Generally, changes in gene expression are determined by measuring transcription (determining the amount of mRNA produced by a gene) or measuring translation (determining the amount of protein produced by a gene). The amount of RNA or protein produced by a gene can be determined using any method known to skilled artisans for quantifying polynucleotides and proteins.

Generally, mRNA expression is determined using polymerase chain reaction (PCR) (including, without limitation, reverse transcription-PCR (RT-PCR) and quantitative real-time PCR (qPCR)), short or long oligonucleotide arrays, cDNA arrays, EST sequencing, Northern blotting, SAGE, MPSS, MS, bead arrays and other hybridization methods. The RNA measured is typically in the form of mRNA or reverse transcribed mRNA.

Protein or polypeptide expression is determined using various colorimetric and spectroscopic assays and methods such as quantitative Western blots, ELISA, 2D-gels, gas or liquid chromatography, mass-spec, the lowry assay, the biuret assay, fluorescence assays, turbidimetric methods, the bicinchoninic assay, protein chip technology, infrared absorbance, ninhydrin, the Bradford assay, and ultraviolet absorbance.

In a preferred method, changes in gene expression are determined using Affymetrix Canine-1 and Canine-2 gene chips available from Affymetrix, Inc. and the instructions for using such chips to determine gene expression. Gene chips allow a large-scale study of biological processes and the measurement of activity within a cell at a certain point in time. Microarray analysis permits one to account for differences in phenotypes on a large-scale genetic basis. Actual measurement of gene expression products is a more accurate indicator of gene function than determining sequences per se. Microarray analysis is based upon quantifying the concentration of a gene’s mRNA transcript in a cell at a given time. DNA is immobilized on a medium and labeled target mRNA is hybridized with probes on the array. Binding of the labeled mRNA to the probes is measured by laser analysis. The measurement is a count of photons emitted. The entire chip is scanned and digitally imaged. The image is processed to locate probes and to assign intensity measurements to each probe. In this manner up- and down-regulated genes may be determined. The analysis enables the skilled person to find groups of genes with similar expression profiles and to determine tissues with similar expression profiles. In this manner, genes that explain the observed differences in tissue samples can be identified.

Affymetrix Gene Chips typically employ probes of 25 bp and probe sets of 11 to 20 probes corresponding to a particular gene or EST. The chip is constructed with a perfect match and mismatch probe of 25 bp each, the former being perfectly complementary to a specific region of a gene and the latter having the 13<sup>th</sup> by substituted to make a mismatch. A probe summarization algorithm is used to determine background correction, normalization and probe summarization, which is the conversion of probe values to probe expression values. RMA is one of the algorithms that may be used for this purpose. The algorithm performs the last two steps of analysis, normalization and summarization of probe-level intensity measurements. The perfect match values are, therefore, background corrected, normalized and summarized into a set of expression measurements.

The raw data is analyzed using GeneSpring version 7.0 (GS) software (Agilent Corporation) and validated using the R-Bioconductor (RB) freeware. Both software packages are used to compute probe intensities from the CEL files generated by the Affymetrix Instrument. The Present/Absent/Marginal calls per probe and P-values are computed using the R-Bioconductor and GeneSpring software separately.

Generally, differential gene expression is determined by measuring the expression of at least one gene. Preferably, the expression of two or more differentially expressed genes is measured to provide a gene expression pattern or gene expression profile. More preferably, the expression of a plurality of differentially expressed genes is measured to provide additional information for a more significant gene expression pattern or profile.

The present invention provides a plurality of markers that together or alone are or can be used as markers of renal disease. In especially useful embodiments of the invention, a plurality of these markers can be selected and their mRNA expression may be measured simultaneously to provide expression profiles for use in various aspects of the inventions described in this application. In a preferred embodiment of the present methods and compositions, at least 2, 3, 4, 5, 6, 7, 8, 9, 10, or 11 markers are selected among lumican (LUM); collagen alpha 1 (III) chain, variant 12 (COL3A1); decorin (DCN); secreted frizzled-related protein 2 (sFRP2); matrilin-2 (Matn2); retinol binding protein 4 (rbp4); MMP-9; clusterin (CLU); transferrin (TF), Apo-C-1 (ApoC1); and

inhibin beta A (INHBA) and can be used for determination of gene expression profiles employed in the practice of the methods of the invention. Each marker can be particularly linked to certain aspects of kidney disease.

In another embodiment of the invention, the level of gene expression can be alternatively determined by detecting the protein corresponding to the gene expression product. Secretable proteins of the denominated genes are preferred.

In another aspect, the invention provides a device suitable for detecting the expression of a plurality of genes differentially expressed in abnormal animals compared to a control values. The device comprises a substrate having a plurality of the oligonucleotide or polynucleotide probes of the present invention affixed to the substrate at known locations. The device is essentially an immobilized version of the oligonucleotide or polynucleotide probes described herein. The device is useful for rapid and specific detection of genes and polynucleotides and their expression patterns and profiles. Typically, such probes are linked to a substrate or similar solid support and a sample containing one or more polynucleotides (e.g., a gene, a PCR product, a ligase chain reaction (LCR) product, a DNA sequence that has been synthesized using amplification techniques, or a mixture thereof) is exposed to the probes such that the sample polynucleotide(s) can hybridize to the probes. Either the probes, the sample polynucleotide(s), or both, are labeled, typically with a fluorophore or other tag such as streptavidin, and detected using methods known to skilled artisans. If the sample polynucleotide(s) is labeled, hybridization may be detected by detecting bound fluorescence. If the probes are labeled, hybridization is typically detected by label quenching. If both the probe and the sample polynucleotide(s) are labeled, hybridization is typically detected by monitoring a color shift resulting from proximity of the two bound labels. A variety of labeling strategies and labels are known to skilled artisans, particularly for fluorescent labels. Preferably, the probes are immobilized on substrates suitable for forming an array (known by several names including DNA microarray, gene chip, biochip, DNA chip, and gene array) comparable to those known in the art.

The polypeptide probes may be made according to conventional methods, e.g., using the nucleotide sequence data provided for polynucleotides of the present invention and methods known in the art. Such methods include, but are not limited to, isolating polypeptide directly from cells, isolating or synthesizing DNA or RNA encoding the polypeptides and using the DNA or RNA to produce recombinant products, synthesizing the polypeptides chemically from individual amino acids, and producing polypeptide fragments by chemical cleavage of existing polypeptides.

In another aspect, the invention provides a device suitable for detecting the expression of a plurality of genes differentially expressed in abnormal animals compared to a control value. The device comprises a substrate having a plurality of the peptide or polypeptide probes of the present invention affixed to the substrate at known locations. The device is essentially an immobilized version of the peptide or polypeptide probes described herein. The device is useful for the rapid and specific detection of proteins and their expression patterns. Typically, such probes are linked to a substrate and a sample containing one or more proteins is exposed to the probes such that the sample proteins can hybridize to the probes. In certain embodiments, the probes, the sample proteins, or both, are labeled and detected, typically with a fluorophore or other agent known to skilled artisans. Generally, the same methods and instrumentation used for reading polynucleotide microarrays is applicable to

protein arrays. Preferably, the probes are immobilized on a substrate suitable for forming an array.

Methods for determining the amount or concentration of protein in a sample are known to skilled artisans. Such methods include radioimmunoassays, competitive-binding assays, Western blot analysis, and ELISA assays. For methods that use antibodies, polyclonal and monoclonal antibodies are suitable. Such antibodies may be immunologically specific for a protein, protein epitope, or protein fragment.

Some embodiments of the invention utilize antibodies for the detection and quantification of proteins produced by expression of the polynucleotides of the present invention. Although proteins may be detected by immunoprecipitation, affinity separation, Western blot analysis, protein arrays, and the like, a preferred method utilizes ELISA technology wherein the antibody is immobilized on a solid support and a target protein or peptide is exposed to the immobilized antibody. Either the probe, or the target, or both, can be labeled using known methods.

In some embodiments, expression patterns or profiles of a plurality of genes differentially expressed in abnormal animals are observed utilizing an array of probes for detecting polynucleotides or polypeptides. In one embodiment, arrays of oligonucleotide or polynucleotide probes may be utilized, whereas another embodiment may utilize arrays of antibodies or other proteins that specifically bind to the differentially expressed gene products of the present invention. Such arrays may be commercially available or they may be custom made using methods known to skilled artisans, e.g., in-situ synthesis on a solid support or attachment of pre-synthesized probes to a solid support via micro-printing techniques. In various embodiments, arrays of polynucleotides or polypeptides probes are custom made to specifically detect transcripts or proteins produced by the differentially expressed genes of the present invention.

In one embodiment, arrays of polynucleotide or polypeptide probes are custom made to specifically detect transcripts or proteins produced by two or more polynucleotides or genes identified in Table 2. These probes are designed to detect genes associated with lipid and glucose metabolism pathways in animals. In another embodiment, arrays of polynucleotide or polypeptide probes are custom made to specifically detect transcripts or proteins produced by two or more polynucleotides or genes identified in Tables 3-5. These probes are designed to detect genes that are particularly relevant to abnormal canines compared to normal canines.

These probes are exposed to a sample to form hybridization complexes that are detected and compared with those of a standard. The differences between the hybridization complexes from the sample and standard indicate differential expression of polynucleotides and therefore genes differentially expressed in abnormal canines compared to a control value in the sample. In a preferred embodiment, probes are made to specifically detect polynucleotides or fragments thereof produced by one or more of the genes or gene fragments identified by the present invention. Methods for detecting hybridization complexes are known to skilled artisans.

In another aspect, the invention provides a method for detecting the differential expression of genes differentially expressed in abnormal canines compared to normal canines in a sample. The method comprises (a) reacting a combination comprising a plurality of polypeptide probes with proteins in the sample under conditions that allow specific binding between the probes and the proteins to occur, wherein the proteins bound by the probes are differentially

expressed in an abnormal canine compared to a normal canine; (b) optionally, reacting a combination comprising a plurality of polypeptide probes with proteins in a standard under conditions that allow specific binding between the probes and the proteins to occur, wherein the proteins bound by the probes are differentially expressed in an abnormal canine compared to a normal canine; (c) detecting specific binding in the sample and, optionally, the standard from step (b); and (d) comparing the specific binding in the sample with that of a standard, wherein differences between the specific binding in the standard and the sample indicate differential expression of genes differentially expressed in abnormal canines compared to normal canines in the sample.

These probes are exposed to a sample to form specific binding that is detected and compared with those of a standard. The differences between the specific binding from the sample and standard indicate differential expression of proteins and therefore genes differentially expressed in abnormal canines compared to normal canines, particularly abnormal-associated genes, in the sample. In a preferred embodiment, probes are made to specifically detect proteins or fragments thereof produced by one or more of the genes or gene fragments identified by the present invention.

In one embodiment, the method further comprises exposing the canine or sample to a test substance before reacting the polypeptides with the proteins. Then, the comparison is indicative of whether the test substance altered the expression of genes differentially expressed in abnormal canines compared to normal canines, particularly abnormal-associated genes, in the sample.

Example 1

Classification of Canines with Chronic Kidney Disease According to the Guidelines of the International Renal Interest Society

In the Examples that follow, canines exhibiting clinical signs of chronic kidney disease were tested versus animals not exhibiting signs or symptoms of chronic kidney disease. Pathological diagnoses of chronic kidney disease were made based on the criteria set forth in Tables 1 and 2 below and in accordance with the guidelines of the International Renal Interest Society (IRIS).

Staging of chronic kidney disease (CKD) is undertaken following the diagnosis of CKD in order to facilitate appropriate treatment and monitoring of the subject animal. Staging is based initially on fasting plasma creatinine, assessed on at least two occasions in the stable animal. Canines demonstrating normal renal function and no clinical signs or symptoms of CKD were grouped as non-disease canines. Stage 1 in canines corresponds to prior classifications of early renal disease with no biochemical evidence of CKD to renal insufficiency, where no azotemia is detected, but where glomerular filtration rate (GFR) may be reduced and there may be a poor concentrating ability of the kidneys. Stage 2 corresponds to the prior classification of early renal failure. In Stage 2, mild azotemia is noted. Stage 3 corresponds to prior classification of uremic renal failure, where moderate azotemia is detected. Systemic signs of uremic renal failure may be present such as bone pain, uremic gastritis, anemia and metabolic acidosis. Stage 4 corresponds to end-stage renal failure, which is characterized by severe azotemia and increasing systemic clinical signs of uremic crisis.

Table 1 identifies the five categories of canines studied, respectively. A total of 42 dogs diagnosed as not having

CKD were studied. A total of 14 Stage 1 canines exhibited minimal glomerulonephritis (GN). The number of dogs studied exhibiting advanced stages of CKD were: Stage 2 mild GN=24; Stage 3 moderate GN=8 and Stage 4 marked GN=13. Plasma creatinine levels for each of the groups of dogs are shown in Table 2 as mean and median plasma creatinine levels for each group of dogs.

TABLE 1

Staging for Canines	
IRIS CKD Staging	Plasma Creatinine range mg/dl
Non-Disease	<1.4 with no firm evidence of disease
Stage 1	<1.4 (<125 μmol/l) with evidence of disease. Non-azotemic. Some other renal abnormality present (e.g. inadequate concentrating ability without identifiable non-renal cause; abnormal renal palpation and/or renal imaging findings; proteinuria of renal origin; abnormal renal biopsy)
Stage 2	1.4 to 2.0 (125-179 μmol/l) Mild renal azotemia. Clinical signs usual mild or absent.
Stage 3	2.1 to 5.0 (180-439 μmol/l) Moderate renal azotemia. Many clinical signs may be present.
Stage 4	>5.0 (>440 μmol/l) Sever renal azotemia. Many extra-renal clinical signs present.

TABLE 2

Creatinine Levels		
Pathological Diagnosis	Mean Plasma Creatinine mg/dl	Median Plasma Creatinine mg/dl
Non-Disease (n = 42)	0.8	0.7
Minimal GN (n = 14)	0.7	0.7
Mild GN (n = 24)	0.9	0.7
Moderate GN (n = 8)	1.5	1.5
Marked GN (n = 13)	6.7	6.2

Example 2

Candidate Selection Criteria

The raw data in the following examples was analyzed using GeneSpring version 7.0 (GS) software (Agilent Corporation) and validated using the R-Bioconductor (RB) freeware. Both software packages are used to compute probe intensities from the CEL files generated by the Affymetrix Instrument. The Present/Absent/Marginal calls per probe and P-values are computed using the R-Bioconductor and GeneSpring software separately.

The gene expression data is determined to be either “up” or “down”-regulated for any given analysis. The decision on whether a gene is “up” or “down” is based on the fold change, which is calculated as treatment intensity/control intensity for each individual probe. The fold change is considered down-regulated if its value is <1/2 and is up-regulated if it is >2.0. Also, a probe is considered significant for further scrutiny if it is called as present in only one of the conditions being compared (treatment or control) and is “absent” or “marginal” in the other and the fold change is significant according to the software used.

## RNA Isolation Procedures

Materials and Methods. The following general procedures may be used to isolate RNA from tissue samples of dogs and cats for gene expression profiling utilizing gene chips as further described in the Examples of this specification. It will be apparent to a person of ordinary skill in the art that these procedures or modifications thereof, as recognized within the art, can be applied to isolate RNA from tissue or body fluid samples for further gene expression analysis using a variety of analytical procedures available to a person of ordinary skill in the art, in particular microarray technologies.

## Isolation of Ribonucleic Acid (RNA) from Tissue.

Tissue samples may be collected, frozen in liquid nitrogen, thawed and then, ground in a mortar and pestle, homogenized and transferred to a 50 ml conical flask. The homogenized tissue sample is then processed using a TRIzol® RNA extraction method according to the manufacturer's instructions (Invitrogen) to produce good quality RNA which is then subjected to further genomic analysis.

Materials: Ice, liquid nitrogen, frozen canine tissue, TRIzol® lysis reagent, chloroform minimum 99%, isopropyl alcohol, 70% ethanol (prepared with ethanol, absolute and deionized, RNase-free water), RNase Zap®, deionized water, RNA Storage Solution®, from Ambion.

Equipment: Ultra-Turrax T25 Power Homogenizer, Beckman Coulter Allegra 25R Centrifuge, Eppendorf Centrifuge, forceps, scalpel, hard cutting surface, i.e. cutting board, 1.5 mL DNase and RNase free/sterile microcentrifuge tubes, 50 mL DNase and RNase free/sterile disposable polypropylene tubes, P1000, P200, P20, P10 and P2 Rainin Pipetman pipettes, filter pipette tips for P1000, P200, P20, P10 and P2 pipettes, DNase and RNase free/sterile, and lint free wipes.

Preparations: Prepare 50 mL polypropylene tubes with 4 mL TRIzol® (one tube for each tissue selected for RNA isolation).

Tissue Homogenization: Fill a container capable of holding liquid nitrogen with 3-4 scoops of liquid nitrogen. Place a piece of frozen tissue immediately into the aforementioned container (the tissue should be about the size of a pea) and place the tissue into the appropriate labeled 50 mL polypropylene tube (that already contains 4 mL TRIzol®). Immediately begin homogenization using the Ultra-Turrax T25 Power Homogenizer. Homogenize on the highest setting (6) for 10-15 seconds. Cool the sample on ice for another 10-15 seconds and then repeat. Continue until the tissue is fully homogenized and the solution is cloudy. Upon complete homogenization, cap the 50 mL tube and return to the ice. Incubate the homogenized tissues at room temperature for 5 minutes before proceeding with the isolation procedure.

## RNA Preparation Procedures

RNA Isolation: The procedures given in the Invitrogen instructions provided with the TRIzol® reagent are generally followed. Separate the homogenized sample into four 1 mL aliquots in four 1.5 mL microcentrifuge tubes. Add 200 uL of chloroform to each 1 mL aliquot. Cap the tubes, vortex for 15 seconds and then shake up and down. The result should be a pink milky liquid. Incubate the tubes at room temperature for 2-3 minutes. Centrifuge the tubes for 15 minutes at 14,000 rpm and 4° C. Transfer the aqueous phase

(top layer) to a sterile 1.5 mL microcentrifuge tube. The typical volume of the aqueous phase which should be transferred to the new tube is about 500 uL. Be sure not to transfer any of the intermediate or lower phase. Precipitate the RNA from solution by adding 500 uL of Isopropyl Alcohol to each microcentrifuge tube containing the aqueous layer. Shake the tubes up and down for at least 20 seconds. Incubate the samples at room temperature for 10 minutes. Centrifuge the samples for 10 minutes, 14,000 rpm at 4° C. Remove the supernatant carefully by aspirating off the liquid being sure not to lose the pellet. Add 1 mL of 70% ethanol to wash the pellet. Dislodge the pellet by flicking the tube (or tapping the tube on the bench top) and shake to mix. Centrifuge for 5 minutes, 8,200 rpm at 4° C. Remove the supernatant carefully by aspirating off the liquid being sure not to lose the pellet. Use a lint free wipe to carefully soak up excess ethanol to make sure the pellet is dry. Resuspend each pellet into 30 uL of RNA Storage Solution. Mix gently by pipetting until the RNA goes back into solution and then store at -80° C. It may be necessary to vortex the sample for a few seconds at a low speed to facilitate the resuspension of the RNA. If this is necessary, spin down the samples, using the microcentrifuge, prior to freezing.

RNA Cleaning: The procedures given in the RNeasy® Mini Handbook are followed.

RNA Isolation from Cells Cultured in OptiCell Chambers Using the RNeasy Mini Kit. Cells cultured from mammalian cell lines are used to isolate good quality RNA which is then used for future downstream genomic analysis. All work related to the culturing of the cells is to be done under strict aseptic conditions.

Reagents: 10xPBS, deionized H<sub>2</sub>O, absolute ethanol, RNA Storage Solution, β-Mercaptoethanol, RNase Zap®, Buffer RLT, and Buffer RW1 and Buffer RPE (provided in the RNeasy Mini Kit).

Equipment/Materials: RNeasy Mini Kit, QIAshredder spin columns, OptiCell knife, 20 mL sterile syringe, OptiCell tips, Cell scraper, P1000 Pipetman pipette, Rainin, P200 Pipetman pipette, Rainin, 100-100 uL filtered pipette tips, 1-200 uL filtered pipette tips, sterile transfer pipettes, 55 mL sterile solution basin, 1.5 mL sterile microcentrifuge tubes, and Eppendorf Microcentrifuge.

Solutions: Buffer RLT (stock provided in RNeasy Mini Kit); —Add 100 uL of β-Mercaptoethanol per 10 mL of Buffer RLT prior to beginning protocol. 70% Ethanol: Make 50 mL of 70% ethanol by adding 35 mL absolute ethanol to 15 mL deionized, RNase-free water. 1xPBS: RNase-free water. Filter the solution using a 0.22 um filter.

Procedure: Removing Cells from the OptiCell Chamber (proceed one OptiCell at a time). Check the cells under a microscope to ensure that the cells are alive before isolating RNA. Remove and discard the cell culture medium. Using the OptiCell knife, cut away the top membrane exposing the cells on the lower membrane. Wash the membrane to which the cells are attached three times with 1xPBS. Pipette 600 uL of the Buffer RLT solution (containing β-Mercaptoethanol) onto the center of the membrane to which the cells are attached. Using the cell scraper, gently spread the Buffer RLT over the entire surface of the membrane, and then collect the liquid in one corner. Pipette off the entire volume of Buffer RLT and place into a QIAshredder spin column.

RNA Isolation: Centrifuge the QIAshredder spin columns at 14,000 rpm for 2 minutes. Discard the spin column but keep the collection tube and its contents. Add 600 uL of 70% ethanol to the collection tube and mix well by pipetting (the total volume now=1.2 mL). Transfer 600 uL of the cell lysate to an RNeasy mini column and centrifuge for 15

seconds at 14,000 rpm. Discard the flow through but keep the collection tube and the spin column. Transfer the remaining volume of cell lysate (~600 uL) to the spin column and repeat the centrifugation. Discard the flow through but keep the collection tube and the spin column. Add 700 uL Buffer RW1 to the spin column. Centrifuge for 15 seconds at 14,000 rpm to wash the column. Discard the flow through and the collection tube. Transfer the spin column to a new 2 mL collection tube and add 500 uL Buffer RPE to the column. Centrifuge for 15 seconds at 14,000 rpm. Discard the flow through, keep the collection tube/column. Add another 500 uL Buffer RPE to the column. Centrifuge for 2 minutes at 14,000 rpm. Transfer the spin column to a 1.5 mL collection tube. Add 30 uL of RNA Storage Solution directly to the silica gel membrane and centrifuge for 1 minute at 14,000 rpm to elute the RNA. Store the final RNA at -70° C.

#### RNA 6000 Nano Assay.

Using the Agilent 2100 Bioanalyzer and the RNA 6000 Nano Assay, analyze RNA isolated from cultured mammalian cells, lymphocytes or tissues for quality.

Reagents: RNA 6000 Nano gel matrix, RNA 6000 Nano dye concentrate, RNA 6000 Nano Marker, (all of the above reagents are contained in the RNA 6000 Nano Assay kit, Agilent), RNA 6000 ladder, RNase Zap, and RNase-free water, from Ambion.

Equipment/Other Materials: Agilent Chip Priming Station, Agilent, RNA 6000 chip, Agilent, electrode cleaners, P2, P10, P200, and P1000 Rainin Pipetman pipettes, sterile, DNase/RNase free filtered pipette tips, 1.5 mL microcentrifuge tubes, sterile, vortex, IKA vortex mixer, microcentrifuge, and heating block.

Procedure: The procedure is given in the Reagent Kit Guide, RNA 6000 Nano Assay, Edition November 2003, by Agilent Technologies. The procedures are followed as given in the Guide, with the following modifications: Preparing the Gel, pg. 17—rather than separating the filtered gel into aliquots of 65 uL each, keep the stock filtered gel in the original microcentrifuge tube and aliquot the 65 uL as needed. Loading the RNA 6000 Nano Marker, pg. 22—add 1 uL of RNase-free water (instead of RNA 6000 Nano Marker) to each sample well that will not contain sample. Not only will this conserve the amount of Marker used but also serves as a negative control to see that none of the reagents are contaminated, including the RNase-free water. Loading the Ladder and Samples, pg. 23—heat denature the samples and RNA 6000 Ladder for an additional 30 seconds (total of 2.5 minutes) at 71° C. Starting the Chip Run, pg. 26—choose the “Eukaryote Total RNA Nano” option from the assay menu.

#### Example 5

##### Affymetrix GeneChip Expression Analysis

Gene expression is analyzed using Affymetrix Canine 1 and Canine 2 GeneChip® Arrays which are commercially available from Affymetrix, Inc., Santa Clara, Calif. 95051.

Total RNA is reverse transcribed into cDNA. The cDNA is used to generate cRNA which is fragmented and used as probes for GeneChip hybridization. The gene chip is washed and the hybridization signal is measured with an Affymetrix laser scanner. The hybridization data is then validated and normalized for further analysis in accordance with instructions from the manufacturer.

Materials: Affymetrix provides most of the reagents and kit. Other reagents listed in the Affymetrix Manual but not supplied in the kit may be obtained separately (refer to GeneChip Expression Analysis Technical Manual (701021 Rev.4) for details), RNase Zap® and deionized water.

Equipment: Eppendorf microcentrifuge, 1.5 mL DNase and RNase free/sterile microcentrifuge tubes, 50 mL DNase and RNase free/sterile disposable polypropylene tubes, P1000, P200, P20, P10 and P2 Rainin Pipetman pipettes, Filter pipette tips for P1000, P200, P20, P10 and P2 pipettes, DNase and RNase free/sterile, and Peltier Thermal Cycler PTC-200.

Procedure: follow all procedures exactly as described in GeneChip Expression Analysis Technical Manual (Affymetrix Copyright 1999-2003). Use 5 microgram of total RNA for the first strand cDNA synthesis. Use either Peltier Thermal Cycler PTC-200 or heat block for temperature control on reactions and probe denaturing. The quality control is performed using RNA NanoDrop chips with BioAnalyser 2100. Use 100 Format (Midi Array) for the canine genechip.

#### Example 6

##### Gene Expression in Dogs with Chronic Kidney Disease

Studies were conducted in accordance with the previous Examples 1-5 using dogs having various stages of chronic kidney disease to determine the underlying gene expression differences between dogs with normal renal function and dogs having minimal, mild, moderate and marked glomerulonephritis corresponding to Stages 1 through 4 as presented on Table 1. Procedures as described in the Examples of this specification were used to prepare tissue and bodily fluid samples from 42 dogs having normal renal function, 14 dogs having minimal glomerulonephritis, 24 dogs having mild glomerulonephritis, 8 dogs having moderate glomerulonephritis and 13 dogs having marked glomerulonephritis, as determined by plasma creatinine levels presented in Table 2 and by clinical observation.

Based on gene expression data comparing the dogs with normal renal function versus dogs having glomerulonephritis, as defined in the preceding examples, the five genes listed on Table 3 were identified as meeting the selection criteria of Example 2 as potential biomarkers of chronic kidney disorders in dogs. The genes include lumican (LUM); collagen alpha 1 (III) chain, variant 12 (COL3A1); Decorin (DCN); secreted frizzled-related protein 2 (SFRP2); and Matrilin-2 (Matn2). Analogous human synonyms and mRNA and protein accession numbers are listed on Table 3 for each gene. Each of the proteins is a secreted protein.

TABLE 3

List of Potential Biomarkers						
Gene	Symbol in Canine/Human	Human Synonyms	Canine Synonym	Gene Description	mRNA	Protein
Lumican	LUM	LDC; SLRR2D			XM_539716.2	XP_539716.1
Collagen alpha 1 (III) chain, variant 12	COL3A1	EDS4A; FLJ34534			XM_858055.1	XP_863148

TABLE 3-continued

List of Potential Biomarkers						
Gene	Symbol in Canine/Human	Human Synonyms	Canine Synonym	Gene Description	mRNA	Protein
Decorin	DCN	CSCD; DSPG2; PG40; PGII; PGS2; SLRR1B			NM_001003228.1	NP_001003228.1
SFRP2	SFRP2	FRP-2; SARP1; SDF-5		secreted frizzled-related protein 2	NM_001002987.1	NP_001002987.1
Matrilin-2	Matn2	Crtm2; MGC102477; matrilin-2		matrilin 2	XM_548552.2	XP_548552.2

TABLE 4

List of Potential Biomarkers						
Gene	Symbol in Canine/Human	Human Synonyms	Canine Synonym	Gene Description	mRNA	Protein
Retinol Binding Protein 4	Rbp4	Rbp-4		retinol binding protein 4, plasma matrix	XM_534969.2	XP_534969.2
MMP-9	MMP9	CLG4B; GELB; MMP-9		metallopeptidase 9 (gelatinase B, 92 kDa gelatinase, 92 kDa type IV collagenase)	NM_001003219.1	NP_001003219.1
Clusterin	CLU	AAG4; APOJ; CLI; KUB1; MGC24903; SGP-2; SGP2; SP-40; TRPM-2; TRPM2	GP80		NM_001003370.1	NP_001003370.1

TABLE 5

List of Potential Biomarkers						
Gene	Symbol in Canine/Human	Human Synonyms	Canine Synonym	Gene Description	mRNA	Protein
Transferrin	TF	DKFZp781D0156; PRO1557; PRO2086			XM_534268	XP_534268
Apo-C-I	APOC1			apolipoprotein C-I	XM_533643	XP_533643
Inhibin beta A	INHBA	EDF; FRP			XM_540364	XP_540364

## SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 22

<210> SEQ ID NO 1

<211> LENGTH: 338

<212> TYPE: PRT

<213> ORGANISM: *Canis familiaris*

<400> SEQUENCE: 1

Met Asn Leu Gly Val Phe Thr Leu Leu Ala Leu Ile Gly Gly Ala  
1 5 10 15

Ser Gly Gln Tyr Tyr Asp Tyr Asp Phe Ser Leu Pro Ile Tyr Gly Leu  
20 25 30

Ser Ser Pro Asn Cys Ala Pro Glu Cys Asn Cys Pro Glu Ser Tyr Pro  
35 40 45

Ser Ala Met Tyr Cys Asp Glu Leu Lys Leu Lys Ser Val Pro Met Val  
50 55 60

-continued

---

```

Pro Pro Gly Ile Lys Tyr Leu Tyr Leu Arg Asn Asn Gln Ile Asp His
65          70          75          80

Ile Asp Glu Lys Ala Phe Glu Asn Val Thr Asp Leu Gln Trp Leu Ile
85          90          95

Leu Asp His Asn Leu Leu Glu Asn Ser Lys Ile Lys Gly Arg Ile Phe
100         105         110

Ser Lys Leu Lys Gln Leu Lys Lys Leu His Ile Asn His Asn Asn Leu
115         120         125

Thr Glu Ser Val Gly Pro Leu Pro Lys Ser Leu Val Asp Leu Gln Leu
130         135         140

Thr His Asn Lys Ile Gln Lys Leu Gly Ser Phe Asp Gly Leu Val Asn
145         150         155         160

Leu Thr Phe Val His Leu Gln His Asn Gln Leu Lys Glu Asp Thr Val
165         170         175

Ser Ala Ala Phe Lys Gly Leu Lys Ser Leu Glu Tyr Leu Asp Leu Ser
180         185         190

Tyr Asn Gln Met Ala Lys Leu Pro Ser Gly Leu Pro Ala Ser Leu Leu
195         200         205

Thr Leu Tyr Leu Asp Asn Asn Lys Ile Ser Asn Ile Pro Asp Glu Tyr
210         215         220

Phe Lys Arg Phe Asn Gly Leu Gln Tyr Leu Arg Leu Ser His Asn Glu
225         230         235         240

Leu Ala Asp Ser Gly Val Pro Gly Asn Ser Phe Asn Val Ser Ser Leu
245         250         255

Leu Glu Leu Asp Leu Ser Tyr Asn Lys Leu Lys Asn Ile Pro Thr Val
260         265         270

Asn Glu Asn Leu Glu Asn Tyr Tyr Leu Glu Val Asn Glu Leu Glu Lys
275         280         285

Phe Glu Val Lys Ser Phe Cys Lys Ile Leu Gly Pro Leu Ser Tyr Ser
290         295         300

Lys Ile Lys His Leu Arg Leu Asp Gly Asn Arg Leu Thr His Thr Ser
305         310         315         320

Leu Pro Pro Asp Met Tyr Glu Cys Leu Arg Val Ala Asn Glu Ile Thr
325         330         335

```

Val Asn

```

<210> SEQ ID NO 2
<211> LENGTH: 1466
<212> TYPE: PRT
<213> ORGANISM: Canis familiaris

```

&lt;400&gt; SEQUENCE: 2

```

Met Thr Ser Phe Val Gln Lys Gly Thr Trp Leu Leu Leu Ala Leu Leu
1          5          10          15

Gln Pro Ala Val Ile Ser Ala Gln Gln Gln Ala Ile Asp Gly Gly Cys
20         25         30

Ser His Leu Gly Gln Ser Tyr Ala Asp Arg Asp Val Trp Lys Pro Glu
35         40         45

Pro Cys Gln Ile Cys Val Cys Asp Ser Gly Ser Val Leu Cys Asp Asp
50         55         60

Ile Ile Cys Asp Glu Gln Glu Leu Asp Cys Pro Asn Pro Glu Ile Pro
65         70         75         80

Phe Gly Glu Cys Cys Ala Val Cys Pro Gln Pro Pro Thr Ser Pro Pro
85         90         95

```

-continued

---

Arg Pro Pro Asn Gly His Gly Pro Gln Gly Pro Lys Gly Asp Pro Gly  
 100 105 110

Pro Pro Gly Ile Pro Gly Arg Asn Gly Asp Pro Gly Ile Pro Gly Gln  
 115 120 125

Pro Gly Ser Pro Gly Ser Pro Gly Pro Pro Gly Ile Cys Glu Ser Cys  
 130 135 140

Pro Thr Gly Pro Gln Pro Asn Tyr Ser Pro Gln Phe Glu Ser Tyr Asp  
 145 150 155 160

Val Lys Ala Gly Val Ala Gly Gly Gly Ile Gly Gly Tyr Pro Gly Pro  
 165 170 175

Ala Gly Pro Pro Gly Pro Pro Gly Pro Pro Gly Thr Ser Gly His Pro  
 180 185 190

Gly Ser Pro Gly Ser Pro Gly Tyr Gln Gly Pro Pro Gly Glu Pro Gly  
 195 200 205

Gln Ala Gly Pro Ala Gly Pro Pro Gly Pro Pro Gly Ala Met Gly Pro  
 210 215 220

Ser Gly Pro Ala Gly Lys Asp Gly Glu Ser Gly Arg Pro Gly Arg Pro  
 225 230 235 240

Gly Glu Arg Gly Leu Pro Gly Pro Pro Gly Met Lys Gly Pro Ala Gly  
 245 250 255

Met Pro Gly Phe Pro Gly Met Lys Gly His Arg Gly Phe Asp Gly Arg  
 260 265 270

Asn Gly Glu Lys Gly Asp Thr Gly Ala Pro Gly Leu Lys Gly Glu Asn  
 275 280 285

Gly Leu Pro Gly Glu Asn Gly Ala Pro Gly Pro Met Gly Pro Arg Gly  
 290 295 300

Ala Pro Gly Glu Arg Gly Arg Pro Gly Leu Pro Gly Ala Ala Gly Ala  
 305 310 315 320

Arg Gly Asn Asp Gly Ala Arg Gly Ser Asp Gly Gln Pro Gly Pro Pro  
 325 330 335

Gly Pro Pro Gly Thr Ala Gly Phe Pro Gly Ser Pro Gly Ala Lys Gly  
 340 345 350

Glu Val Gly Pro Ala Gly Ser Pro Gly Ser Asn Gly Ser Pro Gly Gln  
 355 360 365

Arg Gly Glu Pro Gly Pro Gln Gly His Ala Gly Ala Pro Gly Pro Pro  
 370 375 380

Gly Pro Pro Gly Ser Asn Gly Ser Pro Gly Gly Lys Gly Glu Met Gly  
 385 390 395 400

Pro Ala Gly Ile Pro Gly Ala Pro Gly Leu Ile Gly Ala Arg Gly Pro  
 405 410 415

Pro Gly Pro Pro Gly Thr Asn Gly Ala Pro Gly Gln Arg Gly Gly Ala  
 420 425 430

Gly Glu Pro Gly Lys Asn Gly Ala Lys Gly Glu Pro Gly Pro Arg Gly  
 435 440 445

Glu Arg Gly Glu Ala Gly Ser Pro Gly Ile Pro Gly Pro Lys Gly Glu  
 450 455 460

Asp Gly Lys Asp Gly Ser Pro Gly Glu Pro Gly Ala Asn Gly Leu Pro  
 465 470 475 480

Gly Ala Ala Gly Glu Arg Gly Ala Pro Gly Phe Arg Gly Pro Ala Gly  
 485 490 495

Ala Asn Gly Leu Pro Gly Glu Lys Gly Pro Ala Gly Glu Arg Gly Gly  
 500 505 510

Pro Gly Pro Ala Gly Pro Arg Gly Ala Pro Gly Glu Pro Gly Arg Asp

-continued

515				520				525							
Gly	Val	Pro	Gly	Gly	Pro	Gly	Met	Arg	Gly	Met	Pro	Gly	Ser	Pro	Gly
530						535					540				
Gly	Pro	Gly	Ser	Asp	Gly	Lys	Pro	Gly	Pro	Pro	Gly	Ser	Gln	Gly	Glu
545				550						555					560
Ser	Gly	Arg	Pro	Gly	Pro	Pro	Gly	Pro	Ser	Gly	Pro	Arg	Gly	Gln	Pro
			565						570					575	
Gly	Val	Met	Gly	Phe	Pro	Gly	Pro	Lys	Gly	Asn	Asp	Gly	Ala	Pro	Gly
		580						585				590			
Lys	Asn	Gly	Glu	Arg	Gly	Gly	Pro	Gly	Gly	Pro	Gly	Pro	Gln	Gly	Pro
	595						600					605			
Ala	Gly	Lys	Asn	Gly	Glu	Thr	Gly	Pro	Gln	Gly	Pro	Pro	Gly	Pro	Thr
610						615					620				
Gly	Pro	Ala	Gly	Asp	Lys	Gly	Asp	Ala	Gly	Pro	Pro	Gly	Pro	Gln	Gly
625				630						635					640
Leu	Gln	Gly	Leu	Pro	Gly	Thr	Ser	Gly	Pro	Pro	Gly	Glu	Asn	Gly	Lys
			645						650					655	
Pro	Gly	Glu	Pro	Gly	Pro	Lys	Gly	Glu	Ser	Gly	Ser	Pro	Gly	Val	Pro
			660						665					670	
Gly	Gly	Lys	Gly	Asp	Ser	Gly	Ala	Pro	Gly	Glu	Arg	Gly	Pro	Pro	Gly
		675					680					685			
Ala	Ala	Gly	Pro	Met	Gly	Pro	Arg	Gly	Gly	Ala	Gly	Pro	Pro	Gly	Pro
690						695					700				
Glu	Gly	Gly	Lys	Gly	Ala	Ala	Gly	Pro	Pro	Gly	Pro	Pro	Gly	Ser	Ala
705				710						715					720
Gly	Thr	Pro	Gly	Leu	Gln	Gly	Met	Pro	Gly	Glu	Arg	Gly	Gly	Pro	Gly
			725						730					735	
Gly	Pro	Gly	Pro	Lys	Gly	Asp	Lys	Gly	Glu	Pro	Gly	Ser	Ala	Gly	Ala
			740						745				750		
Asp	Gly	Ala	Pro	Gly	Lys	Asp	Gly	Pro	Arg	Gly	Pro	Thr	Gly	Pro	Ile
	755						760					765			
Gly	Pro	Pro	Gly	Pro	Ala	Gly	Gln	Pro	Gly	Asp	Lys	Gly	Glu	Gly	Gly
770						775					780				
Ala	Pro	Gly	Leu	Pro	Gly	Ile	Ala	Gly	Pro	Arg	Gly	Gly	Pro	Gly	Glu
785					790					795					800
Arg	Gly	Glu	His	Gly	Pro	Pro	Gly	Pro	Ala	Gly	Phe	Pro	Gly	Ala	Pro
			805						810					815	
Gly	Gln	Asn	Gly	Glu	Pro	Gly	Ala	Lys	Gly	Glu	Arg	Gly	Ala	Pro	Gly
		820						825					830		
Glu	Lys	Gly	Glu	Gly	Gly	Pro	Pro	Gly	Val	Ala	Gly	Pro	Pro	Gly	Gly
	835					840					845				
Ala	Gly	Pro	Ala	Gly	Pro	Pro	Gly	Pro	Gln	Gly	Val	Lys	Gly	Glu	Arg
850						855					860				
Gly	Ser	Pro	Gly	Gly	Pro	Gly	Ala	Ala	Gly	Phe	Pro	Gly	Gly	Arg	Gly
865					870					875				880	
Leu	Pro	Gly	Pro	Pro	Gly	Asn	Asn	Gly	Asn	Pro	Gly	Pro	Pro	Gly	Ser
			885						890					895	
Ser	Gly	Ala	Pro	Gly	Lys	Asp	Gly	Pro	Pro	Gly	Pro	Pro	Gly	Asn	Asn
		900							905					910	
Gly	Ala	Pro	Gly	Ser	Pro	Gly	Val	Ser	Gly	Pro	Lys	Gly	Asp	Ala	Gly
	915					920						925			
Gln	Pro	Gly	Glu	Lys	Gly	Ser	Pro	Gly	Pro	Gln	Gly	Pro	Pro	Gly	Ala
930						935					940				

-continued

---

Pro Gly Pro Leu Gly Ile Ala Gly Ile Thr Gly Ala Arg Gly Leu Ala  
 945                                    950                                    955                                    960

Gly Pro Pro Gly Ile Pro Gly Pro Arg Gly Ser Pro Gly Pro Gln Gly  
    965                                    970                                    975

Val Lys Gly Glu Asn Gly Lys Pro Gly Pro Ser Gly His Asn Gly Glu  
    980                                    985                                    990

Arg Gly Pro Pro Gly Pro Gln Gly Leu Pro Gly Leu Ala Gly Thr Ala  
    995                                    1000                                    1005

Gly Glu Pro Gly Arg Asp Gly Asn Pro Gly Ser Asp Gly Leu Pro  
    1010                                    1015                                    1020

Gly Arg Asp Gly Ala Pro Gly Gly Lys Gly Asp Arg Gly Glu Asn  
    1025                                    1030                                    1035

Gly Ser Pro Gly Ala Pro Gly Ala Pro Gly His Pro Gly Pro Pro  
    1040                                    1045                                    1050

Gly Pro Val Gly Pro Ala Gly Lys Ser Gly Asp Arg Gly Glu Thr  
    1055                                    1060                                    1065

Gly Pro Ala Gly Pro Ser Gly Ala Pro Gly Pro Ala Gly Ser Arg  
    1070                                    1075                                    1080

Gly Pro Pro Gly Pro Gln Gly Pro Arg Gly Asp Lys Gly Glu Thr  
    1085                                    1090                                    1095

Gly Glu Arg Gly Leu Asn Gly Ile Lys Gly His Arg Gly Phe Pro  
    1100                                    1105                                    1110

Gly Asn Pro Gly Ala Pro Gly Ser Pro Gly Pro Ala Gly His Gln  
    1115                                    1120                                    1125

Gly Ala Ile Gly Ser Pro Gly Pro Ala Gly Pro Arg Gly Pro Val  
    1130                                    1135                                    1140

Gly Pro Ser Gly Pro Pro Gly Lys Asp Gly Thr Ser Gly His Pro  
    1145                                    1150                                    1155

Gly Pro Ile Gly Pro Pro Gly Pro Arg Gly Asn Arg Gly Glu Arg  
    1160                                    1165                                    1170

Gly Ser Glu Gly Ser Pro Gly His Pro Gly Gln Pro Gly Pro Pro  
    1175                                    1180                                    1185

Gly Pro Pro Gly Ala Pro Gly Pro Cys Cys Gly Gly Gly Ala Ala  
    1190                                    1195                                    1200

Ala Leu Ala Ala Ala Gly Gly Glu Lys Ala Gly Gly Phe Ala Pro  
    1205                                    1210                                    1215

Tyr Tyr Gly Asp Glu Pro Met Asp Phe Lys Ile Asn Thr Glu Glu  
    1220                                    1225                                    1230

Ile Met Thr Ser Leu Lys Ser Val Ser Gly Gln Ile Glu Ser Leu  
    1235                                    1240                                    1245

Val Ser Pro Asp Gly Ser Arg Lys His Pro Ala Arg Asn Cys Arg  
    1250                                    1255                                    1260

Asp Leu Lys Phe Cys His Pro Glu Leu Lys Ser Gly Glu Tyr Trp  
    1265                                    1270                                    1275

Val Asp Pro Asn Gln Gly Cys Lys Leu Asp Ala Ile Lys Val Phe  
    1280                                    1285                                    1290

Cys Asn Met Glu Thr Gly Glu Thr Cys Leu Asn Ala Ser Pro Gly  
    1295                                    1300                                    1305

Ser Val Pro Arg Lys Asn Trp Trp Thr Asp Ser Gly Ala Glu Lys  
    1310                                    1315                                    1320

Lys His Val Trp Phe Gly Glu Ser Met Asp Gly Gly Phe Gln Phe  
    1325                                    1330                                    1335

-continued

---

Gly Tyr Gly Asn Pro Glu Leu Pro Glu Asp Val Leu Asp Val Gln  
 1340 1345 1350  
 Leu Ala Phe Leu Arg Leu Leu Ser Ser Arg Ala Ser Gln Asn Ile  
 1355 1360 1365  
 Thr Tyr His Cys Lys Asn Ser Ile Ala Tyr Met Asp His Ala Ser  
 1370 1375 1380  
 Gly Asn Val Lys Lys Ala Leu Arg Leu Met Gly Ser Asn Glu Gly  
 1385 1390 1395  
 Glu Phe Lys Ala Glu Gly Asn Ser Lys Phe Thr Tyr Thr Val Leu  
 1400 1405 1410  
 Glu Asp Gly Cys Thr Lys His Thr Gly Glu Trp Gly Lys Thr Val  
 1415 1420 1425  
 Phe Glu Tyr Arg Thr Arg Lys Ala Val Arg Leu Pro Ile Ile Asp  
 1430 1435 1440  
 Ile Ala Pro Tyr Asp Val Gly Gly Pro Asp Gln Glu Phe Gly Val  
 1445 1450 1455  
 Asp Val Gly Pro Val Cys Phe Leu  
 1460 1465

<210> SEQ ID NO 3  
 <211> LENGTH: 360  
 <212> TYPE: PRT  
 <213> ORGANISM: Canis familiaris

<400> SEQUENCE: 3

Met Lys Ala Thr Ile Ile Phe Leu Leu Leu Ala Gln Val Ser Trp Ala  
 1 5 10 15  
 Gly Pro Phe Gln Gln Arg Gly Leu Phe Asp Phe Met Leu Glu Asp Glu  
 20 25 30  
 Ala Ser Gly Ile Gly Pro Glu Asp Arg Ala Pro Asp Met Pro Asp Leu  
 35 40 45  
 Glu Leu Leu Gly Pro Val Cys Pro Phe Arg Cys Gln Cys His Leu Arg  
 50 55 60  
 Val Val Gln Cys Ser Asp Leu Gly Leu Asp Lys Val Pro Lys Asp Leu  
 65 70 75 80  
 Pro Pro Asp Thr Thr Leu Leu Asp Leu Gln Asn Asn Lys Ile Thr Glu  
 85 90 95  
 Ile Lys Asp Gly Asp Phe Lys Asn Leu Lys Asn Leu His Thr Leu Ile  
 100 105 110  
 Leu Val Asn Asn Lys Ile Ser Lys Ile Ser Pro Gly Ala Phe Thr Pro  
 115 120 125  
 Leu Leu Lys Leu Glu Arg Leu Tyr Leu Ser Lys Asn His Leu Lys Glu  
 130 135 140  
 Leu Pro Glu Lys Met Pro Lys Thr Leu Gln Glu Leu Arg Ala His Glu  
 145 150 155 160  
 Asn Glu Ile Thr Lys Val Arg Lys Ala Val Phe Asn Gly Leu Asn Gln  
 165 170 175  
 Met Ile Val Val Glu Leu Gly Thr Asn Pro Leu Lys Ser Ser Gly Ile  
 180 185 190  
 Glu Asn Gly Ala Phe Gln Gly Met Lys Lys Leu Ser Tyr Ile Arg Ile  
 195 200 205  
 Ala Asp Thr Asn Ile Thr Thr Ile Pro Gln Gly Leu Pro Pro Ser Leu  
 210 215 220  
 Thr Glu Leu His Leu Glu Gly Asn Lys Ile Thr Lys Val Asp Ala Ser  
 225 230 235 240

-continued

Ser Leu Lys Gly Leu Asn Asn Leu Ala Lys Leu Gly Leu Ser Phe Asn  
 245 250 255

Ser Ile Ser Ala Val Asp Asn Gly Thr Leu Ala Asn Thr Pro His Leu  
 260 265 270

Arg Glu Leu His Leu Asp Asn Asn Lys Leu Ile Arg Val Pro Gly Gly  
 275 280 285

Leu Ala Glu His Lys Tyr Ile Gln Val Val Tyr Leu His Asn Asn Asn  
 290 295 300

Ile Ser Ala Val Gly Ser Asn Asp Phe Cys Pro Pro Gly Tyr Asn Thr  
 305 310 315 320

Lys Lys Ala Ser Tyr Ser Gly Val Ser Leu Phe Ser Asn Pro Val Gln  
 325 330 335

Tyr Trp Glu Ile Gln Pro Ser Thr Phe Arg Cys Val Tyr Val Arg Ser  
 340 345 350

Ala Ile Gln Leu Gly Asn Tyr Lys  
 355 360

&lt;210&gt; SEQ ID NO 4

&lt;211&gt; LENGTH: 294

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Canis familiaris

&lt;400&gt; SEQUENCE: 4

Met Pro Arg Gly Pro Gly Ser Leu Leu Leu Leu Val Leu Ala Ser His  
 1 5 10 15

Cys Cys Leu Gly Ser Ala Arg Gly Leu Phe Phe Gly Gln Pro Asp Phe  
 20 25 30

Ser Tyr Lys Arg Ser Asn Cys Lys Pro Ile Pro Ala Asn Leu Gln Leu  
 35 40 45

Cys His Gly Ile Glu Tyr Gln Asn Met Arg Leu Pro Asn Leu Leu Gly  
 50 55 60

His Glu Thr Met Lys Glu Val Leu Glu Gln Ala Gly Ala Trp Ile Pro  
 65 70 75 80

Leu Val Met Lys Gln Cys His Pro Asp Thr Lys Lys Phe Leu Cys Ser  
 85 90 95

Leu Phe Ala Pro Val Cys Leu Asp Asp Leu Asp Glu Thr Ile Gln Pro  
 100 105 110

Cys His Ser Leu Cys Val Gln Val Lys Asp Arg Cys Ala Pro Val Met  
 115 120 125

Ser Ala Phe Gly Phe Pro Trp Pro Asp Met Leu Glu Cys Asp Arg Phe  
 130 135 140

Pro Gln Asp Asn Asp Leu Cys Ile Pro Leu Ala Ser Ser Asp His Leu  
 145 150 155 160

Leu Pro Ala Thr Glu Glu Ala Pro Lys Val Cys Glu Ala Cys Lys Asn  
 165 170 175

Lys Asn Glu Asp Asp Asn Asp Ile Met Glu Thr Leu Cys Lys Asn Asp  
 180 185 190

Phe Ala Leu Lys Ile Lys Val Lys Glu Ile Thr Tyr Ile Asn Arg Asp  
 195 200 205

Thr Lys Ile Ile Leu Glu Thr Lys Ser Lys Thr Ile Tyr Lys Leu Asn  
 210 215 220

Gly Val Ser Glu Arg Asp Leu Lys Lys Ser Val Leu Trp Leu Lys Asp  
 225 230 235 240

Ser Leu Gln Cys Thr Cys Glu Glu Met Asn Asp Ile Asn Ala Pro Tyr



-continued

---

Ala Lys Ala His Gly Cys Glu Gln Leu Cys Val Asn Val Leu Gly Ser  
325 330 335

Phe Val Cys Gln Cys Tyr Ser Gly Phe Thr Leu Ala Glu Asp Gly Lys  
340 345 350

Thr Cys Ala Ala Val Asp Tyr Cys Ala Ser Glu Asn His Gly Cys Glu  
355 360 365

His Glu Cys Val Asn Ala Asp Ser Ser Tyr Phe Cys Arg Cys Arg Lys  
370 375 380

Gly Phe Thr Leu Asn Pro Asp Lys Lys Thr Cys Ala Lys Ile Asp Tyr  
385 390 395 400

Cys Ala Ser Pro Asn Gln Gly Cys Gln His Glu Cys Val Asn Thr Asp  
405 410 415

Asp Ser Tyr Ser Cys Arg Cys Leu Lys Gly Phe Thr Leu Asn Pro Asp  
420 425 430

Lys Lys Thr Cys Arg Arg Ile Asn Tyr Cys Ala Leu Asn Lys Pro Gly  
435 440 445

Cys Glu His Glu Cys Ile Asn Ile Glu Asp Gly Tyr Tyr Cys Arg Cys  
450 455 460

Arg Gln Gly Tyr Thr Leu Asp Pro Asn Gly Lys Thr Cys Ser Arg Val  
465 470 475 480

Asp His Cys Ala Glu Gln Asp His Gly Cys Glu Gln Leu Cys Leu Asn  
485 490 495

Thr Glu Asp Ser Tyr Val Cys Gln Cys Ser Glu Gly Phe Leu Ile Asn  
500 505 510

Asp Asp Leu Lys Thr Cys Ser Arg Ala Asp Tyr Cys Leu Leu Ser Asn  
515 520 525

His Gly Cys Glu Tyr Ser Cys Val Asn Thr Asp Arg Ser Phe Val Cys  
530 535 540

Arg Cys Pro Glu Gly Tyr Val Leu Arg Ser Asp Gly Lys Thr Cys Ala  
545 550 555 560

Lys Leu Asp Ser Cys Ala Val Gly Asp His Gly Cys Glu His Ser Cys  
565 570 575

Val Ser Ser Gly Asp Ser Phe Val Cys Gln Cys Phe Glu Gly Tyr Ile  
580 585 590

Leu Arg Glu Asp Gly Lys Thr Cys Arg Arg Lys Gly Val Cys Gln Ser  
595 600 605

Val Asn His Gly Cys Glu His Ile Cys Val Asn Ser Asp Glu Ser Tyr  
610 615 620

Ile Cys Lys Cys Arg Glu Gly Phe Arg Leu Ala Glu Asp Gly Lys Arg  
625 630 635 640

Cys Arg Arg Lys Asp Val Cys Lys Ser Thr Tyr His Gly Cys Glu His  
645 650 655

Ile Cys Val Asn His Gly Asn Ser Tyr Ile Cys Lys Cys Ser Glu Gly  
660 665 670

Phe Val Leu Ala Glu Asp Gly Lys Arg Cys Lys Arg Cys Thr Glu Gly  
675 680 685

Pro Leu Asp Leu Val Phe Val Ile Asp Gly Ser Lys Ser Leu Gly Glu  
690 695 700

Glu Asn Phe Glu Ile Val Lys Gln Phe Val Ala Gly Ile Ile Asp Ser  
705 710 715 720

Leu Ala Val Ser Pro Lys Ala Ala Arg Val Gly Leu Leu Gln Tyr Ser  
725 730 735

Thr Gln Val Arg Thr Glu Phe Thr Leu Gly Asp Phe Ser Ser Ala Arg

-continued

740				745				750							
Asp	Met	Lys	Lys	Ala	Val	Ala	His	Met	Lys	Tyr	Met	Gly	Lys	Gly	Ser
		755					760					765			
Met	Thr	Gly	Leu	Ala	Leu	Lys	His	Met	Phe	Glu	Arg	Ser	Phe	Asn	Pro
	770					775					780				
Val	Glu	Gly	Ala	Arg	Pro	Val	Ser	Pro	Gly	Val	Ser	Arg	Val	Ala	Ile
785					790					795					800
Val	Phe	Thr	Asp	Gly	Arg	Ala	Gln	Asp	Asp	Val	Ser	Ala	Trp	Ala	Arg
			805						810						815
Arg	Ala	Arg	Ala	Gly	Gly	Ile	Thr	Met	Tyr	Ala	Val	Gly	Val	Gly	Lys
			820						825						830
Ala	Ile	Glu	Glu	Glu	Leu	Gln	Glu	Ile	Ala	Ser	Glu	Pro	Thr	Asp	Lys
		835					840								845
His	Leu	Phe	Tyr	Ala	Glu	Asp	Phe	Ser	Thr	Met	Gly	Glu	Ile	Ser	Glu
	850					855					860				
Lys	Leu	Lys	Lys	Gly	Ile	Cys	Glu	Ala	Leu	Glu	Asp	Ser	Asp	Gly	Ser
865					870					875					880
Gln	Asp	Ser	Pro	Ala	Gly	Glu	Leu	Pro	Lys	Arg	Val	His	Gln	Pro	Thr
			885						890						895
Glu	Ser	Glu	Pro	Val	Thr	Ile	Asn	Ile	Arg	Asp	Leu	Leu	Ser	Cys	Ser
			900						905						910
Asn	Phe	Ala	Val	Gln	His	Arg	Tyr	Leu	Phe	Glu	Glu	Asp	Ser	Val	Ser
		915					920					925			
Arg	Ser	Thr	Gln	Lys	Leu	Phe	His	Ser	Thr	Lys	Ser	Ser	Gly	Ser	Pro
	930					935					940				
Leu	Glu	Glu	Lys	His	Asp	Gln	Cys	Lys	Cys	Glu	Asn	Leu	Ile	Met	Phe
945					950					955					960
Gln	Asn	Leu	Ala	Asn	Glu	Glu	Val	Arg	Lys	Leu	Thr	Gln	Arg	Tyr	Ile
			965						970						975

Leu Phe

&lt;210&gt; SEQ ID NO 6

&lt;211&gt; LENGTH: 267

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Canis familiaris

&lt;400&gt; SEQUENCE: 6

Met	Gln	Val	Leu	Ala	Ala	Gly	Arg	Arg	Leu	Pro	Ser	Val	Phe	His	Pro
1				5					10					15	
Gly	Arg	Tyr	Glu	Ser	Glu	Thr	Pro	Ser	Arg	Pro	Gly	Tyr	Lys	Ala	Gly
			20						25				30		
Gly	Pro	Ala	Ala	Pro	Ala	Asp	Arg	Arg	Leu	Pro	Trp	Leu	His	Ala	Arg
		35					40					45			
Pro	Ala	Pro	Arg	Pro	Gly	Leu	Arg	Ala	Ala	Pro	Val	Gly	Gly	Leu	Pro
		50				55					60				
Gly	Thr	Met	Glu	Trp	Val	Trp	Ala	Leu	Val	Leu	Leu	Ala	Ala	Leu	Gly
65					70					75					80
Ser	Ala	Arg	Ala	Glu	Ser	Asp	Cys	Arg	Val	Ser	Asn	Phe	Gln	Val	Lys
				85					90					95	
Lys	Asn	Phe	Asp	Lys	Ala	Arg	Phe	Ala	Gly	Thr	Trp	Tyr	Ala	Met	Ala
				100					105					110	
Lys	Lys	Asp	Pro	Glu	Gly	Leu	Phe	Leu	Gln	Asp	Asn	Ile	Val	Ala	Glu
			115						120					125	
Phe	Ser	Val	Asp	Glu	Asn	Gly	Arg	Met	Ser	Ala	Thr	Ala	Lys	Gly	Arg



-continued

---

Tyr Ser Ala Cys Thr Thr Asp Gly Arg Ser Asp Asp Thr Pro Trp Cys  
 245 250 255  
 Ser Thr Thr Ala Asp Tyr Asp Thr Asp Arg Arg Phe Gly Phe Cys Pro  
 260 265 270  
 Ser Glu Lys Leu Tyr Thr Gln Asp Gly Asn Gly Asp Gly Lys Pro Cys  
 275 280 285  
 Val Phe Pro Phe Thr Phe Glu Gly Arg Ser Tyr Ser Thr Cys Thr Thr  
 290 295 300  
 Asp Gly Arg Ser Asp Gly Tyr Arg Trp Cys Ser Thr Thr Gly Asp Tyr  
 305 310 315 320  
 Asp Gln Asp Lys Leu Tyr Gly Phe Cys Pro Thr Arg Val Asp Ser Ala  
 325 330 335  
 Val Thr Gly Gly Asn Ser Ala Gly Glu Pro Cys Val Phe Pro Phe Ile  
 340 345 350  
 Phe Leu Gly Lys Gln Tyr Ser Thr Cys Thr Arg Glu Gly Arg Gly Asp  
 355 360 365  
 Gly His Leu Trp Cys Ala Thr Thr Ser Asn Phe Asp Arg Asp Lys Lys  
 370 375 380  
 Trp Gly Phe Cys Pro Asp Gln Gly Tyr Ser Leu Phe Leu Val Ala Ala  
 385 390 395 400  
 His Glu Phe Gly His Ala Leu Gly Leu Asp His Ser Ser Val Pro Glu  
 405 410 415  
 Ala Leu Met Tyr Pro Met Tyr Ser Phe Thr Glu Gly Pro Pro Leu His  
 420 425 430  
 Glu Asp Asp Val Arg Gly Ile Gln His Leu Tyr Gly Pro Arg Pro Glu  
 435 440 445  
 Pro Glu Pro Gln Pro Pro Thr Ala Pro Pro Thr Ala Pro Pro Thr Val  
 450 455 460  
 Cys Ala Thr Gly Pro Pro Thr Thr Arg Pro Ser Glu Arg Pro Thr Ala  
 465 470 475 480  
 Gly Pro Thr Gly Pro Pro Ala Ala Gly Pro Thr Gly Pro Pro Thr Ala  
 485 490 495  
 Gly Pro Ser Glu Ala Pro Thr Val Pro Val Asp Pro Ala Glu Asp Ile  
 500 505 510  
 Cys Lys Val Asn Ile Phe Asp Ala Ile Ala Glu Ile Arg Asn Tyr Leu  
 515 520 525  
 His Phe Phe Lys Glu Gly Lys Tyr Trp Arg Phe Ser Lys Gly Lys Gly  
 530 535 540  
 Arg Arg Val Gln Gly Pro Phe Leu Ile Thr Asp Thr Trp Pro Ala Leu  
 545 550 555 560  
 Pro Arg Lys Leu Asp Ser Ala Phe Glu Asp Gly Leu Thr Lys Lys Thr  
 565 570 575  
 Phe Phe Phe Ser Gly Arg Gln Val Trp Val Tyr Thr Gly Thr Ser Val  
 580 585 590  
 Val Gly Pro Arg Arg Leu Asp Lys Leu Gly Leu Gly Pro Glu Val Thr  
 595 600 605  
 Gln Val Thr Gly Ala Leu Pro Gln Gly Gly Gly Lys Val Leu Leu Phe  
 610 615 620  
 Ser Arg Gln Arg Phe Trp Ser Phe Asp Val Lys Thr Gln Thr Val Asp  
 625 630 635 640  
 Pro Arg Ser Ala Gly Ser Val Glu Gln Met Tyr Pro Gly Val Pro Leu  
 645 650 655  
 Asn Thr His Asp Ile Phe Gln Tyr Gln Glu Lys Ala Tyr Phe Cys Gln

-continued

---

Asp	Arg	Phe	Tyr	Trp	Arg	Val	Asn	Ser	Arg	Asn	Glu	Val	Asn	Gln	Val	660	665	670	
		675					680					685							
Asp	Glu	Val	Gly	Tyr	Val	Thr	Phe	Asp	Ile	Leu	Gln	Cys	Pro	Glu	Asp	690	695	700	
<p>&lt;210&gt; SEQ ID NO 8          &lt;211&gt; LENGTH: 445          &lt;212&gt; TYPE: PRT          &lt;213&gt; ORGANISM: Canis familiaris</p> <p>&lt;400&gt; SEQUENCE: 8</p>																			
Met	Met	Lys	Thr	Leu	Leu	Leu	Leu	Val	Gly	Leu	Leu	Leu	Thr	Trp	Asp	1		10	15
Asn	Gly	Arg	Val	Leu	Gly	Asp	Gln	Ala	Val	Ser	Asp	Thr	Glu	Leu	Gln	20	25	30	
Glu	Met	Ser	Thr	Glu	Gly	Ser	Lys	Tyr	Ile	Asn	Lys	Glu	Ile	Lys	Asn	35	40	45	
Ala	Leu	Lys	Gly	Val	Lys	Gln	Ile	Lys	Thr	Leu	Ile	Glu	Gln	Thr	Asn	50	55	60	
Glu	Glu	Arg	Lys	Ser	Leu	Leu	Ser	Asn	Leu	Glu	Glu	Ala	Lys	Lys	Lys	65	70	75	80
Lys	Glu	Asp	Ala	Leu	Asn	Asp	Thr	Lys	Asp	Ser	Glu	Thr	Lys	Leu	Lys	85	90	95	
Ala	Ser	Gln	Gly	Val	Cys	Asn	Asp	Thr	Met	Met	Ala	Leu	Trp	Glu	Glu	100	105	110	
Cys	Lys	Pro	Cys	Leu	Lys	Gln	Thr	Cys	Met	Lys	Phe	Tyr	Ala	Arg	Val	115	120	125	
Cys	Arg	Ser	Gly	Ser	Gly	Leu	Val	Gly	His	Gln	Leu	Glu	Glu	Phe	Leu	130	135	140	
Asn	Gln	Ser	Ser	Pro	Phe	Tyr	Phe	Trp	Met	Asn	Gly	Asp	Arg	Ile	Asp	145	150	155	160
Ser	Leu	Leu	Glu	Asn	Asp	Arg	Gln	Gln	Thr	His	Ala	Leu	Asp	Val	Met	165	170	175	
Gln	Asp	Ser	Phe	Asn	Arg	Ala	Ser	Ser	Ile	Met	Asp	Glu	Leu	Phe	Gln	180	185	190	
Asp	Arg	Phe	Phe	Thr	Arg	Glu	Pro	Gln	Asp	Thr	Tyr	His	Tyr	Ser	Pro	195	200	205	
Phe	Ser	Leu	Phe	Gln	Arg	Arg	Pro	Phe	Phe	Asn	Pro	Lys	Phe	Arg	Ile	210	215	220	
Ala	Arg	Asn	Ile	Ile	Pro	Phe	Pro	Arg	Phe	Gln	Pro	Leu	Asn	Phe	His	225	230	235	240
Asp	Met	Phe	Gln	Pro	Phe	Phe	Asp	Met	Ile	His	Gln	Ala	Gln	Gln	Ala	245	250	255	
Met	Asp	Val	Asn	Leu	His	Arg	Ile	Pro	Tyr	His	Phe	Pro	Ile	Glu	Phe	260	265	270	
Pro	Glu	Glu	Asp	Asn	Arg	Thr	Val	Cys	Lys	Glu	Ile	Arg	His	Asn	Ser	275	280	285	
Thr	Gly	Cys	Leu	Lys	Met	Lys	Asp	Gln	Cys	Glu	Lys	Cys	Gln	Glu	Ile	290	295	300	
Leu	Ser	Val	Asp	Cys	Ser	Ser	Asn	Asn	Pro	Ala	Gln	Val	Gln	Leu	Arg	305	310	315	320
Gln	Glu	Leu	Ser	Asn	Ser	Leu	Gln	Ile	Ala	Glu	Lys	Phe	Thr	Lys	Leu	325	330	335	

-continued

---

Tyr Asp Glu Leu Leu Gln Ser Tyr Gln Glu Lys Met Phe Asn Thr Ser  
                   340                  345                  350  
 Ser Leu Leu Lys Gln Leu Asn Glu Gln Phe Ser Trp Val Ser Gln Leu  
                   355                  360                  365  
 Ala Asn Leu Thr Gln Ser Glu Asp Pro Phe Tyr Leu Gln Val Thr Thr  
                   370                  375                  380  
 Val Gly Ser Gln Thr Ser Asp Ser Asn Val Pro Val Gly Phe Thr Lys  
 385                  390                  395                  400  
 Val Val Val Lys Leu Phe Asp Ser Asp Pro Ile Thr Val Met Ile Pro  
                   405                  410                  415  
 Glu Ala Val Ser Arg Asn Asn Pro Lys Phe Met Glu Thr Val Ala Glu  
                   420                  425                  430  
 Lys Ala Leu Gln Glu Tyr Arg Gln Lys His Arg Glu Glu  
                   435                  440                  445

&lt;210&gt; SEQ ID NO 9

&lt;211&gt; LENGTH: 705

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Canis familiaris

&lt;400&gt; SEQUENCE: 9

Met Arg Leu Ala Val Arg Ala Leu Leu Ala Cys Ala Val Leu Gly Leu  
 1                  5                  10                  15  
 Cys Leu Ala Val Ser Pro Glu Lys Thr Val Arg Trp Cys Thr Val Ser  
                   20                  25                  30  
 Asn His Glu Ala Ser Lys Cys Ser Ser Phe Met Glu Asn Met Lys Thr  
                   35                  40                  45  
 Val Leu Glu Asn Gly Pro Phe Val Ser Cys Val Lys Arg Thr Ser Tyr  
                   50                  55                  60  
 Leu Glu Cys Ile Lys Ala Ile Trp Ala Asn Glu Ala Asp Ala Val Thr  
 65                  70                  75                  80  
 Leu Asp Ala Gly Leu Val Phe Glu Ala Gly Leu Asn Pro Tyr Asn Leu  
                   85                  90                  95  
 Lys Pro Val Val Ala Glu Phe Tyr Gly Lys Asp Lys Gln Thr Arg Tyr  
                   100                  105                  110  
 Tyr Ala Val Ala Val Val Lys Lys Ser Ser Thr Phe Asn Leu Asn Gln  
                   115                  120                  125  
 Leu Gln Gly Lys Lys Ser Cys His Thr Gly Leu Gly Arg Ser Ala Gly  
                   130                  135                  140  
 Trp Asn Ile Pro Met Gly Leu Leu Tyr Trp Lys Leu Pro Glu Pro Arg  
 145                  150                  155                  160  
 Glu Ser Leu Gln Lys Ala Ala Ser Ser Phe Phe Thr Ala Ser Cys Val  
                   165                  170                  175  
 Pro Cys Val Asp Arg Thr Ala Phe Pro Lys Leu Cys Gln Leu Cys Val  
                   180                  185                  190  
 Gly Lys Gly Thr Asp Lys Cys Ala Cys Ser Asn His Glu Pro Tyr Phe  
                   195                  200                  205  
 Gly Tyr Ser Gly Ala Phe Lys Cys Leu Met Glu Asp Ala Gly Glu Val  
                   210                  215                  220  
 Ala Phe Val Lys His Ser Thr Val Leu Glu Asn Leu Pro Asn Lys Ala  
 225                  230                  235                  240  
 Asp Leu Asp Gln Tyr Glu Leu Leu Cys Pro Asp Asn Lys Arg Lys Pro  
                   245                  250                  255  
 Val Gly Glu Tyr Lys Gln Cys Tyr Leu Ala Lys Val Pro Ser His Ala  
                   260                  265                  270

-continued

---

Val Val Ala Arg Ser Val Gly Gly Lys Glu Asp Leu Ile Trp Glu Leu  
 275 280 285  
 Leu Asn Gln Ala Gln Glu His Tyr Gly Lys Asp Lys Ser Lys Val Phe  
 290 295 300  
 Gln Leu Phe Ser Ser Thr Leu Gly Lys Asp Leu Leu Phe Lys Asp Ser  
 305 310 315 320  
 Ala Glu Gly Phe Phe Arg Ile Pro Pro Lys Met Asp Thr Trp Leu Tyr  
 325 330 335  
 Leu Gly Tyr Glu Tyr Val Thr Ala Leu Arg Asn Leu Arg Glu Asp Val  
 340 345 350  
 Arg Pro Asp Thr Pro Arg Asp Glu Cys Lys Lys Val Lys Trp Cys Ala  
 355 360 365  
 Val Gly His His Glu Ile Ala Lys Cys Asp Glu Trp Ser Val Asn Ser  
 370 375 380  
 Glu Gly Lys Ile Glu Cys Glu Ser Ala Glu Ser Thr Glu Asp Cys Ile  
 385 390 395 400  
 Ala Lys Ile Ala Lys Gly Glu Ala Asp Ala Met Ser Leu Asp Gly Gly  
 405 410 415  
 Tyr Ile Tyr Ile Ala Gly Gln Cys Gly Leu Val Pro Val Leu Ala Glu  
 420 425 430  
 Asn Tyr Lys Thr Gln Gly Ser Thr Cys Ser Asn Thr Ala Glu Glu Gly  
 435 440 445  
 Tyr Leu Ala Val Ala Val Val Lys Arg Leu Asp Lys Thr Ile Ser Trp  
 450 455 460  
 Asn Asn Leu Gln Gly Arg Lys Ser Cys His Thr Ala Val Asp Arg Thr  
 465 470 475 480  
 Ala Gly Trp Asn Ile Pro Met Gly Leu Leu Tyr Asn Arg Ile Asn His  
 485 490 495  
 Cys Glu Phe Asp Lys Phe Phe Ser Gln Gly Cys Ala Pro Gly Ser Met  
 500 505 510  
 Arg Asn Ser Ser Leu Cys Ala Leu Cys Ile Gly Ser Ala Asn Val Pro  
 515 520 525  
 Gly Lys Glu Cys Val Pro Asn Asn His Glu Arg Tyr Tyr Gly Tyr Thr  
 530 535 540  
 Gly Ala Phe Arg Cys Leu Val Glu Lys Gly Asp Val Ala Phe Val Lys  
 545 550 555 560  
 Asp Gln Thr Val Leu Gln Asn Thr Gly Gly Lys Asn Thr Glu Asp Trp  
 565 570 575  
 Ala Lys Asp Leu Lys Glu Glu Asp Phe Glu Leu Leu Cys Pro Asp Gly  
 580 585 590  
 Gln Arg Lys Ser Val Asp Lys Ala Pro Glu Cys Phe Leu Ala Lys Ala  
 595 600 605  
 Pro Asn His Ala Val Val Ser Arg Lys Asp Lys Ala Ser Cys Val Ser  
 610 615 620  
 Lys Met Leu Leu Asp Gln Gln Leu Leu Phe Gly Arg Asn Gly Asn Asp  
 625 630 635 640  
 Cys Ser Gly Lys Phe Cys Leu Phe His Ser Ala Thr Lys Asp Leu Leu  
 645 650 655  
 Phe Arg Asp Asp Thr Gln Cys Leu Ala Lys Leu Pro Glu Asp Thr Thr  
 660 665 670  
 Tyr Lys Ser Tyr Leu Gly Ala Glu Tyr Ile Thr Ala Val Ala Asn Leu  
 675 680 685

-continued

---

Arg Gln Cys Ser Thr Ser Lys Leu Leu Glu Ala Cys Thr Phe His Thr  
690 695 700

Ser  
705

<210> SEQ ID NO 10  
<211> LENGTH: 88  
<212> TYPE: PRT  
<213> ORGANISM: Canis familiaris

<400> SEQUENCE: 10

Met Arg Leu Ile Leu Ser Leu Pro Val Leu Val Val Val Leu Ser Met  
1 5 10 15

Val Leu Glu Gly Pro Ala Pro Ala Gln Ala Ala Gly Glu Ile Ser Ser  
20 25 30

Thr Phe Glu Arg Ile Pro Asp Lys Leu Lys Glu Phe Gly Asn Thr Leu  
35 40 45

Glu Asp Lys Ala Arg Ala Ala Ile Glu Ser Ile Lys Lys Ser Asp Ile  
50 55 60

Pro Ala Lys Thr Arg Asn Trp Phe Ser Glu Ala Phe Lys Lys Val Lys  
65 70 75 80

Glu His Leu Lys Thr Ala Phe Ser  
85

<210> SEQ ID NO 11  
<211> LENGTH: 424  
<212> TYPE: PRT  
<213> ORGANISM: Canis familiaris

<400> SEQUENCE: 11

Met Pro Leu Leu Trp Leu Arg Gly Phe Leu Val Ala Ser Cys Trp Ile  
1 5 10 15

Ile Val Arg Ser Ser Pro Thr Pro Gly Pro Glu Gly Pro Gly Ala Ala  
20 25 30

Pro Ala Cys Pro Ala Cys Ala Leu Thr Ala Leu Pro Arg Asp Ala Pro  
35 40 45

Asn Ser Gln Pro Glu Met Val Glu Ala Val Lys Lys His Ile Leu Asn  
50 55 60

Met Leu His Leu Lys Lys Arg Pro Glu Val Thr Gln Pro Val Pro Lys  
65 70 75 80

Ala Ala Leu Leu Asn Ala Ile Arg Lys Leu His Val Gly Lys Val Gly  
85 90 95

Glu Asn Gly Phe Val Glu Ile Glu Asp Asp Ile Gly Arg Arg Ala Glu  
100 105 110

Met Asn Glu Leu Met Glu Gln Thr Ser Glu Ile Ile Thr Phe Ala Glu  
115 120 125

Ser Gly Thr Ala Arg Lys Thr Leu His Phe Glu Ile Ser Lys Glu Gly  
130 135 140

Ser Asp Leu Ser Val Val Glu Arg Ala Glu Val Trp Leu Phe Leu Lys  
145 150 155 160

Val Pro Lys Ala Asn Arg Thr Arg Thr Lys Val Thr Ile Arg Leu Leu  
165 170 175

Gln Lys His Pro Gln Gly Ser Leu Asp Ala Gly Glu Glu Ala Glu Asp  
180 185 190

Met Gly Phe Pro Glu Glu Arg Asn Glu Val Leu Ile Ser Glu Lys Val  
195 200 205

-continued

---

Val Asp Ala Arg Lys Ser Thr Trp His Ile Phe Pro Val Ser Ser Ser  
 210 215 220

Ile Gln Arg Leu Leu Asp Gln Gly Arg Ser Ser Leu Asp Val Arg Ile  
 225 230 235 240

Ala Cys Glu Gln Cys His Glu Thr Gly Ala Ser Leu Val Leu Leu Gly  
 245 250 255

Lys Lys Lys Lys Lys Glu Glu Glu Gly Glu Gly Lys Lys Lys Asp Gly  
 260 265 270

Gly Asp Ala Gly Ala Gly Gly Asp Glu Asp Lys Glu Gln Ser His Arg  
 275 280 285

Pro Phe Leu Met Leu Gln Ala Arg Gln Ser Glu Asp His Pro His Arg  
 290 295 300

Arg Arg Arg Arg Gly Leu Glu Cys Asp Gly Lys Val Asn Ile Cys Cys  
 305 310 315 320

Lys Lys Gln Phe Phe Val Ser Phe Lys Asp Ile Gly Trp Asn Asp Trp  
 325 330 335

Ile Ile Ala Pro Ser Gly Tyr His Ala Asn Tyr Cys Glu Gly Gly Cys  
 340 345 350

Pro Ser His Ile Ala Gly Thr Ser Gly Ser Ser Leu Ser Phe His Ser  
 355 360 365

Thr Val Ile Asn His Tyr Arg Leu Arg Gly His Ser Pro Phe Thr Asn  
 370 375 380

Leu Lys Ser Cys Cys Val Pro Thr Lys Leu Arg Pro Met Ser Met Leu  
 385 390 395 400

Tyr Tyr Asp Asp Gly Gln Asn Ile Ile Lys Lys Asp Ile Gln Asn Met  
 405 410 415

Ile Val Glu Glu Cys Gly Cys Ser  
 420

<210> SEQ ID NO 12  
 <211> LENGTH: 2028  
 <212> TYPE: DNA  
 <213> ORGANISM: Canis familiaris

<400> SEQUENCE: 12

```

agctagagta ttggactag ttaagcatga atctctctcc cattocatag ggaatgagct    60
gggcagtcct tctcctccc gtccacctgc acttcgtag agagcagaat gctcacatgc    120
caccaccaca gatcccccaga gtgacataac tccatgcaga gactggcgtg actgggctgg    180
gtttcccccc cttcagctct tgtatcacta agaatctggc agccagttcc gtctgacag    240
agttcacagc atatattggt ggattcttgt ccatagtgca tctgctttaa gaattaacga    300
aagcagtgtc aagacagaag gattcaaagc atttgccaaa aatgaatcta ggtgtgttta    360
ctctactctt ggcattaatt ggtggtgcca gcgccagta ctatgattac gattttctcc    420
taccgattta tgggctgtca tcaccaaact gtgcaccaga atgtaactgc cctgaaagct    480
accctcggc catgtactgc gatgagctga aattgaaaag tgtaccaatg gtgcctcctg    540
gaatcaagta tctttacctt aggaataacc agatcgacca tattgatgaa aaggcctttg    600
aaaacgtaac tgacctgcag tggetcattc tggatcataa ccttctagaa aattccaaga    660
ttaaaggaag aattttctct aaactgaaac aactgaagaa gctgcatata aatcacaaca    720
acctgacaga gtctgtgggc ccgcttccca aatctctggt ggacctgcag cttacacaca    780
acaagatcca gaagctgggc tcttcgatg gactggtaaa cctgaccttt gtcacacctc    840
aacacaatca gctgaaagag gatactgtgt cagctgcttt taaaggtctt aagtcctctg    900
    
```

-continued

---

```

agtacctga cttgagctac aatcagatgg ccaaactgcc ttctggcttc ccagcatctc   960
ttctgactct ctacttgac aacaataaga tcagcaacat ccctgatgag tatttcaagc  1020
gttttaatgg actgcagtat ctgctgttat ctcataatga actggcggat agtggagtac  1080
ctgaaaatc ttttaatgta tcatocctgc tggagctgga totctcctac aataagctta  1140
aaaacatacc gactgtcaat gagaacctg aaaactatta cctggaggtc aatgaactcg  1200
aaaagtttga ggtaaagagc ttctgtaaga tccttgacc attatcctac tccaagatta  1260
aacatttgcg tctggatggc aaccgtctta cccacaccag tctgccacct gatatgtacg  1320
aatgtctacg tgttgcaaat gaaatcactg ttaattaaca tctactaatt ccaattctat  1380
tgagggttac cctggagcaa cattttatgg tttatgtttt tgtgtgtgtc agttttcata  1440
gtattcaatt ttttgggtca ctttattagt ttcataaatt tcagattctg agggaaatgt  1500
tttgtaaaca ttttactact tttttttta agataagatg aaaggcaggc ctatttcatc  1560
acaagcacag acatagacac acagagacaa caaacttatg ctttattgtt aaatttagtg  1620
ttttctccat ctctactgct aatgatgtg caaagtcttc tcttgggtga atggaagtca  1680
gccaaggatt ataactcctg aatcttaact taatatgcca aacagcatgg gctacacaca  1740
tatgaatatt taatatcttg aaccaaaatga tctcaaatc caatgttcag atttgettta  1800
ctgaaaaata ggtggttaaat ttaagactg atgattttgc aaaatactag accaaattca  1860
tgaagccaca tacacttaga gcaatatttt tagtattaag aattatatag ttacctagtg  1920
aaacttctct tttctagaat tatttttccac tctaagtcac gtgtatgttc ctttttaatc  1980
atttgcattg tatgtttaac aagctcatag caaaaataaaa catagcaa           2028

```

&lt;210&gt; SEQ ID NO 13

&lt;211&gt; LENGTH: 5511

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Canis familiaris

&lt;400&gt; SEQUENCE: 13

```

gaagccaaac ttttctcat ttaaggcga agcgaaggaa tctcagtggc tgagttttat   60
gacgggcccg gtgctgcagg gcagggaaaca tctgatggtg ctactttgag ctgctgctct  120
ccccctctcg ctgcacaaaag agtctcatgt ctacattta gacatgacga gctttgtgca  180
aaaggggacc tgggtgcttc tcgctctgct tcagcccgtc gtcactctcg cacagcagca  240
agctattgat ggaggatgct cccactctgg tcagtcctat gcggatagag atgtctggaa  300
gccagaaccg tgccaaatat gcgtctgtga ctcaggatct gttctctgcg atgacataat  360
atgtgacgaa caagaattag actgtcccaa cccagagatc ccatttggag aatggtgtgc  420
agtttgccca cagcctccaa catctctctc tcgcccctct aatggtcatt gacctcaagg  480
ccctaagggg gatccaggcc ctctctggtat tctctggaga aatggcgacc cgggtattcc  540
cggacagcca ggctcccctg gttctctggt cccccctgga atctgtgaat cctgcctctc  600
tggctcctcag cctaactatt ctcccagtt cgagtccat gacgtcaagg ctggtgtcgc  660
aggaggcggc atcggaggct accccgggac agctggtccc cctggctctc ctggtcccc  720
tggtagatct ggtcatcctg gctccccggg ttcccagga taccaggcc cccctggtga  780
gcctgggcaa getggtcctg cgggcccctc aggaccccct ggtgctatgg gtccatctgg  840
tcctgctgga aaagatgggg aatcaggaag acctggacga cctggagagc gagggctgcc  900
tggacctcca ggtatgaaaag gtccagctgg catgcctgga ttcccggta tgaaggaca  960

```

-continued

---

tagaggcttt gatggacgaa atggagaaaa gggtgataca ggtgctctg gattaaaggg	1020
tgaaaatggc ctcccagggtg aaaatggagc ccccggaccc atgggtcccc gaggtgctcc	1080
tggtagcgga ggacggccag gcctccccgg ggccgagggt gctcgaggta acgatgggtc	1140
tcgtggaagt gacggacaac caggccctcc cggccccct ggaactgcag gattccccgg	1200
ttcccctggt gctaaggggtg aagttggacc cgcgggatct cctggttcaa acggctcccc	1260
tggacaaaaga ggagaacctg gacctcaggg acatgctggt gctccaggcc ctctggccc	1320
ccctgggagc aatggtagtc ctggcggtaa agtgaaatg ggtcccctg gcattcctgg	1380
agcccctgga ctgataggag cccggggccc tcttgacca cccgtacca atgggtgctcc	1440
tggacagcga ggtggtgcag gtgaaccogg taaaatggt gccaaaggag agccaggccc	1500
acgcggtgaa cgtggtgaa ctggttctcc aggtatccca ggacctaaag gtgaagatgg	1560
caaagatggc tcgcctggag aaccaggtgc aaacggactt cggggagctg ccggagaaaag	1620
gggtgcacct ggattccgag gacctgctgg agcaaatggc ctccaggag aaaagggtcc	1680
cgtgggggag cgtggtggtc caggccctgc agggcccaga ggagctcctg gagaaccggg	1740
ccgagatggt gtcccaggag gtccaggaat gagggcatg cccggaagcc caggagggcc	1800
aggcagcgt gggaaaccag ggcctccogg aagtcaagga gaaagtggtc gaccaggctc	1860
tccaggcccg tctggtcccc gaggtcagcc tggcgtcatg ggtttccctg gtccctaaag	1920
aatgatggt gacctggca agaatggaga gcgagtggt cctggaggtc cggccctca	1980
gggtcctgct gaaagaatg gtgaaactgg acctcagggt cccccgggc ctactgggcc	2040
ggctggtgac aaaggagacg cgggaccccc tggccccaa gggctacagg gtttgcccg	2100
aaccagtgga ccccaggag aaaaaggaaa acccggtgaa ccaggcccga agggatgaatc	2160
tggttcacct ggagttccag gaggcaaggc cgactcaggt gcaccgggcg agcgtggtcc	2220
tccccgggct gcaggccca tggcccccg aggtggagct ggaaccccag gcccgaagg	2280
agggaagggc gctgctggcc cccctgggc acccgctcg gcaggtaacc ctggtctgca	2340
gggatgccc ggagagagag gaggccccg cggccccgc cccaagggtg acaagggcga	2400
accaggcagt gcaggtgcag atggagcccc ggggaaggat ggtcccagg gtcctaccg	2460
tcccatcggc cccctggcc cagccggtca gcctggagat aagggtgaag gtggtgcccc	2520
cggcctcccg gcatagctg gtctcgcgg tggccccgc gagagagggt agcacggccc	2580
cccggaacc gccggtctcc ccggagctcc cggacagaac ggcgagccc gegctaagg	2640
cgaaggggc gccccggcg agaagggtga aggagcccc cccggagtgg caggcccccc	2700
cggaggagcc gggcctgcg gtccccccg tcccagggt gtcaaagggt aacgtggcag	2760
tcccgcggt cccggtgccc ctggctccc tgggtgctgt ggtcttctg gccctcctgg	2820
caataatggt aaccaggcc cccaggtc cagtgggtct ccaggcaagg atgggcccc	2880
cgtccaccc ggtaacaacg gcgtcctgg cagccctgg gtgtccggac ctaaagggtga	2940
tgctggccaa cccggtgaga agggatcgc tggccccag ggcctcccg gagccccagg	3000
ccctctcgga attgctggga tcaccggggc cagggtctc gcgggaccac caggcattcc	3060
aggctcctagg ggaagcccag gccacaggc cgtcaagggt gaaaatggga agccaggacc	3120
cagtggctac aacggagagc gcggtcctcc cgggccccag gggcttccc gcctggcccg	3180
caaggctggt gaacctgaa gagatgaaa cccaggtccc gacgggctgc caggccgcga	3240
cggcgtccc ggccgcaagg gtgatcgtgg ggagaacggg tctcctggcg ccccgggcgc	3300
ccccggccac cccggcccc ccggccccgt gggccccggc gggaaagagc gegaccgagg	3360

-continued

---

```

cgagacgggc cccgctggtc cctccggcgc tcttggctct gcaggctccc gagggccccc 3420
tggcccccaa ggtccacgcg gtgacaaaagg tgaaccgggt gaacgtggcc ttaacggcat 3480
caaaggccat cgaggattcc ccgtaaccc aggtgcccc ggctctccgg gtectgcagg 3540
ccaccaggga gcaatcggtg gtccaggccc cgcaggcccc agaggaccag ttggaccag 3600
tggacccccct ggtaaagacg gaacaagtgg acatccaggt cccattggac caccaggccc 3660
tcgaggtaac agagagaaa gaggatctga gggtccccc ggccaccag gacagccagg 3720
tccccccggg cctcctggtg cccctggacc ctgctgtggt gggggcgtg ccgccctggc 3780
tgccgccgga ggggagaaa ctggcgggtt tgccccctat tacggagatg agccaatgga 3840
tttcaaaatc aacaccgagg agattatgac ttcgctcaaa tccgtcagtg gacaaataga 3900
aagcctcgtt agccccgacg ggtcccgcaa acaccccgct cggaaactgcc gggacctcaa 3960
attctgccat ccagagctca agagcggaga gtactgggtt gacccaacc aaggctgcaa 4020
gttgatgcg atcaaagtat tctgcaacat gaaaaccggg gagacgtgcc tgaatgccag 4080
tccccgggagc gttccccgga agaactgggt gacagattct ggtgctgaga agaaacatgt 4140
ttggtttgga gaatccatgg atggtggctt ccagtttggc tacggcaatc ccgagctgcc 4200
tgaagacgtc ctcgatgtcc agctggcatt cctccgcctt ctctccagcc gggcctccca 4260
gaacatcacg tatcactgca agaatagcat cgcgtacatg gatcacgcca gtgggaatgt 4320
aaagaaagcc ctgaggctga tggggtcaaa tgaagtgaa ttcaaggctg aaggaaacag 4380
caaattcaca tacacagctc tggaggatgg ttgcactaaa cacactgggg aatggggcaa 4440
aacagtcttc gaataccgaa cacgcaaggc tgtgagacta cctattatag atattgcacc 4500
ctatgatggt ggtggtcctg atcaagaatt tgggtggac gttggcctg tttgcttttt 4560
ataaaccaaa ctctatctga aaccccagca aaaaatttca cactccatat gtgttcctct 4620
cgttctaacc ttgtcaacca gtacaagtga ccaactcaat tccagttatt tatttccaaa 4680
agttttggaa aaaagtataa tttggcaaaa aaaaagatac cttttttttt tttttttttg 4740
ctgttacacc aaatacagtt caaatgcttt ttgttctatt tttttaccaa tttcaatttc 4800
aaaatgtctc aatggtgcta taataataa acttcaacac tttacaata aactgtgtt 4860
atattctttg aatcctagcc catttgcaga gcaatgacta ggcttaccat taaaagataa 4920
cctttcttcc tgaatatgtc aaacatgaaa ttagagaaga cctcctggt tcaactacct 4980
caacctggtc agaatacag atgaattcta gaagcctgg aagatgaaaa aaatcataa 5040
ttagcaataa tttcctaaaa tattgataaa aaagtatagt acgagaattt aaagaaaaa 5100
tttttttaaa tctacgattt taatcctgga tatcaactgc ttttaaagggt gcttttctcc 5160
tttctgtgca ttgctgaaat ggtcaagatt acccacattt ggggaaggctt taaagacaca 5220
tggtatgggt ctaatgtact ttcactttaa aaactccagg tcagaattgt tgccttgcac 5280
tcaaacataa aatgcacacc atctttcaaa aaaatttatt gccgtgtcac ttcaggggac 5340
tctctctccc ttgatcctgt aaaagtcaac aacaacaaca acaacaaca aaaattatgg 5400
ggctgctttt gtcacagtaa cacagagaat gtgctgaaat ttaactttgt aagcttgcac 5460
gtggtgtgtg gtcttttttt tttttctctc cagacacca taataaata t 5511

```

&lt;210&gt; SEQ ID NO 14

&lt;211&gt; LENGTH: 1470

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Canis familiaris

-continued

&lt;400&gt; SEQUENCE: 14

```

taaaaccagg aggttctcaa cctattaaca ttcacaaaac agcacctacc cctcctcct    60
ttccacacct gcaaactctt ttgcttgggc tgaatattta gtttaattac atctcacct    120
taagggtccc tgtggcaaat ccccgatta aaaggttccct tggctgtgaa aatacataac    180
ctaaacctat aaggcaacta tcatcttctc cctgcttgca caagtttctc gggctgggccc    240
gttccaacag agaggcttat ttgactttat gctagaagat gaggcttccg ggataggccc    300
ggaggaccgt gcacctgaca tgectgacct cgagcttctg ggacctgtgt gtccttccg    360
ctgtcagtgc catctccgag tggctccagt ttccgacctg ggtctggaca aagtacccaa    420
agatcttccc cctgacacta cgctgctoga cttgcaaac aacaaaatca ccgaaatcaa    480
agatggagac tcaagaacct tcaagaacct gcataccttg attctttaa acaacaaaat    540
tagcaaaatc agccctggag catttacacc tttgtgaaa ttggaacgac tttatctgtc    600
caagaatcat ctgaaggaat tgccagaaaa aatgcccaaa actcttcagg agctgcgtgc    660
ccatgagaat gagatcacca aagttcgaaa agctgtgttc aatggactga accagatgat    720
cgtcgtagag ctgggcacca atcccctgaa gagttcaggg attgaaaatg gagccttcca    780
gggaatgaag aagctctcct atatccgeat tgctgatacc aatataacta ccatccctca    840
aggtcttctc ccttccctta ctgaattaca tcttgaaggc aacaaaatca ccaaggttga    900
tgcactagc ctgaaaggac tgaataattt ggctaagttg ggactgagtt ttaacagcat    960
ctccgctggt gacaatggca ctctagccaa cactcctcat ctgagggagc ttcacttgga   1020
caacaataag ctcatcagag taccgggtgg gctggcggag cataagtaca tccaggttgt   1080
ctaccttcat aacaacaata tatctgcagt cggatctaata gacttctgcc cactggata   1140
caacacccaa aaggcttctt attcaggtgt gagccttttc agcaaccag tgcagtactg   1200
ggagatccag ccatccacct tccgggtgtg ctacgtgcgc tctgccatcc agcttgga   1260
ttataataa ttoccaagaa agccctcgtt ttataacct ggcaaaatcc cattaatc   1320
attgctcaaa aaaaaaaaaa aaaaaaacta aatacgggaa acttagctgc aatgtgaatg   1380
tttccccct aatttataat actaaaccaa attgcattta aaatgccaaa aaaaaaaaaa   1440
ttggccgccc aaaaaaaaaa aaaaaaaaaa                                     1470

```

&lt;210&gt; SEQ ID NO 15

&lt;211&gt; LENGTH: 1760

&lt;212&gt; TYPE: DNA

<213> ORGANISM: *Canis familiaris*

&lt;400&gt; SEQUENCE: 15

```

attcggcacg aggatgccgc ggggccccgg ctccgtgctg ctgctcgtcc tagcgtcga    60
ctgctgcttg gctcggcgc gcggtctctt cttcggccag cccgacttct cctacaagcg    120
cagcaactgc aagcccatcc cggccaacct gcagctgtgc cacggcatcg agtaccagaa    180
catcgggctg cccaacctgc tgggccacga gacctgaag gaggtgctgg agcaggcggg    240
cgcttgatc ccgctggtea tgaagcagtg ccaccggac accaagaagt tcctctgctc    300
gctcttggcc ccgctgtgcc tcgacgacct ggacgagacc atccagcgt gccactctct    360
ctgctgcag gtgaaggacc gctgcgcccc ggtcatgtcg gccttcggct tcccctggcc    420
ggacatgctc gagtgcgacc gattccccca ggacaacgac ctctgcatac cctcgtctag    480
cagcgaccat ctctgcggc ccaccgagga agctccaaag gtatgcgaag cctgcaaaaa    540
taaaaatgag gatgacaacg acataatgga aactctttgt aaaaatgatt ttgcaactgaa    600

```

-continued

---

```

aataaaagtg aaggagataa cctacatcaa cagagatacc aaaatcatcc tggagaccaa 660
gagcaagacc atttacaagc tgaacgggtg gtctgaaagg gacctgaaga aatccgtgct 720
gtggctgaaa gacagcctgc agtgcacctg cgaggagatg aatgacatca acgcgcctta 780
tctggtcacg ggacagaaac taggtgggga gctggtgatc acctcagtga agcgggtggca 840
aaaggggcag agagagtcca agcgcacctc ccgaagcadc cgcaagttgc agtgctagt 900
tgcccccccc tccccaggcc gcctccagag ccaggctgac cacttccgct ctgggtcccc 960
agctctcadc cccccaagca caggcccttg cagctctggc cccagcctgg agcagcttcc 1020
cttgcccttt gcacgtttgc acctggcat ctctgaatt ataaggcctt aggaagcttc 1080
aggagtggat actgttttaa acatacggga ggaaccacc cagatcttgt agaaatgttc 1140
aaactaataa aatcatggaa tatttttacg gaagttttta aatagctcaa ctttagtttt 1200
gaataggtac agctgtgact tgggtctgac tttccttctt tctgtttggt ttgggccaac 1260
tgattttcaa tttcgtggta aggttgccgt aacgtgcaaa tggcttcatt tttcagtg 1320
gccccaaaatg tgggtgggta caatccttcg ttgagataaa gctggctggt attcgacgtt 1380
tctcagctcc agcctgagac ttcagagcct aagtcttata ataattcacc tgcattttta 1440
tgccccggtt gggaaactac aacagtagca gcatgtactg atttccgggt agagtacttc 1500
catttatcaa aagcacatta accaccatag catgattctt cgaataaagg gcaaacaga 1560
ttttataatt gacctgagta ctttaagctt ttgtttaaaa cttttttac ttaattttgc 1620
aatttaaacc attgtagctt acctgtaata tacatagtag ttgacctaa aaagttgtaa 1680
aatattgct ttaaccaaca ctgtaaatat ttcagataag cattatattc ttgtatataa 1740
actttgcttc ctgttttaaa 1760

```

&lt;210&gt; SEQ ID NO 16

&lt;211&gt; LENGTH: 2937

&lt;212&gt; TYPE: DNA

<213> ORGANISM: *Canis familiaris*

&lt;400&gt; SEQUENCE: 16

```

atggaggctg aagtgccatg gaaagtggag catgtcgtca ctggtgctag taaacagcgc 60
cagaggcact tccaagggac cagttgccac ttggagctcc agactatgga aaagatgctg 120
gtgggctgct tctgtttggt cctcggacag atcctccttg tcctccctgc tgaggccaaa 180
gagcggactc cctggaggtc catctccaga gggagacatg cttggacca tccccagacg 240
gccctcctgg agagctcctg tgagaataaa caggcagacc tggcttcat catcgacagc 300
tcccgcagtg tcaaacacca tgactatgca aaagttaagg agttcatcct ggacatcttg 360
cagttcttgg acatcagctc tgacctcacc cgagtgggccc tgtcgcagta tggcagcacc 420
atcaagaacg agttctccct caagacctc aagaagaagt ctgaagtgga gcgtgccctc 480
aagaggatga ggcacctgtc cacaggcacc atgacggggc tggccatcca gtacgcctg 540
aacattgcat tctcagaagc agagggggca cggcccctga gggagaacgt gctgcgggtc 600
ataatgattg tgaccgacgg gaggccacag gactccgtgg ccgaggtggc tgccaaggca 660
cgtgacacag gcacctgat ctttgccatc ggtgtgggccc aggtggacct caacacactg 720
aaggccattg ggagtgagcc ccatgaggac catgtcttcc tgggtggccaa cttcagccag 780
atggagtccc tgacctcggg gttccagaag aagttgtgca tggctccatc gtgcagcgtc 840
ctggagcata actgtgcccc cttctgcadc aatacccccg gctcatacgt ctgcaggtgc 900

```

-continued

---

```

aaacaagggg acatcctcaa ctcagatcag acgacttgca gaatccagga tctgtgtgcg 960
gccaaagccc atgggtgtgt gcagctctgt gtcaatgtgc tcggctcctt tgtctgccag 1020
tgttacagcg gcttcacact ggctgaggac ggaagacgt gtgcggctgt ggactactgt 1080
gcctcagaga accacggatg tgaacatgag tgtgtgaatg ctgacagctc ctacttttgc 1140
cggtgccgta aaggatttac tcttaacca gataaaaaa catgtgcaa gatagactac 1200
tgtgcctcgc ctaatcaagg atgtcagcac gagtgtgtta acacagatga ctctattcc 1260
tgtcgtgcc taaaaggttt taccttgaat ccagataaga aaacctgcag aaggatcaac 1320
tactgtgcat tgaacaaacc gggctgtgag catgaatgca tcaacataga ggatggctac 1380
tactgccgct gccccaggg ctacacctg gacccaatg gcaagacttg cagcagggtg 1440
gatcactgcg cggagcagga ccatggctgt gacagctgt gcctgaacac tgaggattct 1500
tacgtctgcc agtgcctaga aggttcctc atcaacgacg acctcaagac ctgctccagg 1560
gcggtattat gcttgtgag caaccatggt tgcgaatac cctgcgtcaa cactgaccga 1620
tcctttgttt gtccgtgtcc tgagggatac gtgctccgca gtgacgggaa gacctgtgcc 1680
aaattggact cctgtgctgt gggagaccat ggctgtgaac attcatgtgt aagcagtgga 1740
gactcttttg tgtgccagtg ctttgaaggg tatatactcc gtgaagatgg gaaaacctgc 1800
agaaggaaag gtgtctgcca atcagtaaac catggctgtg agcatatttg tgtgaacagt 1860
gatgagtcgt acatctgcaa gtgccgggaa ggattcaggc ttgctgaaga tgggaagcgc 1920
tgcagaagga aggatgtctg caaatcgacc taccacggct gtgaacacat ttgtgttaat 1980
catggcaact cctacatctg caaatgctca gagggatttg ttctagctga ggatggaaaa 2040
cggtgcaaga gatgactgag agggccgctt gacctggtct ttgtgatcga tgggtccaag 2100
agcctcgggg aagagaatth cgagattgtg aagcagttcg tcgcccgcac tatagattcc 2160
ttggcggttt ccccaaaagc gcgccgagtg gggctgctgc agtactccac gcaggtccgc 2220
acggagttca cctcgggaga cttcagctcg gccagagaca tgaaaaaagc cgtggccac 2280
atgaagtaca tgggcaaggg ctctatgacg gggctggccc tgaaacacat gtttgagaga 2340
agttttaacc cagtggaaag ggcagaccc gtgtccccg ggggtccag agtggccatc 2400
gtgttcacgg acggccgggc ccaggacgac gtctccgct gggcccgcag agcccggcc 2460
ggcggcataa ctatgtatgc tgtcgggta gaaaagcta ttgaggaaga gctacaagag 2520
attgcctctg agcccacaga caagcatctc ttctatgccc aagacttcag cacaatggga 2580
gagataagtg aaaaactaaa gaagggata tgtgaagctc tggaagactc tgatggaagt 2640
caggactccc cagcagggga actgccaag agggccacc agccaacaga atctgagcca 2700
gtcaccataa atatccgaga cctactttcc tgttctaatt ttgcagtgca acaccggtat 2760
ctgtttgaag aagacagtgt ttcacggctc acacaaaaa tttttcattc acaaaaatct 2820
tcaggaagtc ctttgaaga aaaacacgat cagtgcaaat gtgaaaatct tataatgttc 2880
cagaaccttg cgaatgaaga agtgagaaag ttaacacaac gctatatcct tttctaa 2937

```

```

<210> SEQ ID NO 17
<211> LENGTH: 1054
<212> TYPE: DNA
<213> ORGANISM: Canis familiaris

```

```

<400> SEQUENCE: 17

```

```

ttatgcaagt gctggccgcc ggccggcgcc tcccctcggc etttcacccc ggccgttacg 60
aaagcgagac cccctcccgc cccggctata aagcggggcg cccggcgccc cggcgggatc 120

```

-continued

---

```

cccgccttcc ctggetccac gcgcgccccg caccgcggcc aggcttgcgc gcagctccgg 180
tggggcgact cccgggcacg atggagtggg tgtgggcgct cgtgctgctg gcggcgctgg 240
gcagcgcccc ggccggagagc gactgcccag tgagcaactt ccaagtcaag aagaacttcg 300
acaaggetcg ctctgcgggg acctggtacg ccatggccaa gaaggacccc gagggcctct 360
ttctgcagga caacatcgtc gctgagttct cggtgatga gaatggccgg atgagcgcca 420
cggccaaggg ccgagtcggg cttttgaata actgggacgt gtgtgcagac atggttggca 480
ccttcacaga caccgaggac cctgctaaat tcaagatgaa gtattggggc gtagcgtcct 540
tcctccagaa aggaaatgat gaccactgga tcatcgacac ggactatgac acctatgccg 600
tgcagtattc ctgcccctc ctgaacctcg atggcacctg tgctgacagc tactccttcg 660
tgttttcccg tgacccaat ggcttcccc tggaggcaca aaagatcgtg aggagagggc 720
aggaggagct gtgcctggcc aggcagtaca ggctgatcgt tcacaatggg tattgtgacg 780
gcagatcaga accaaacct ttgtagcaac gtggagtttc atttgaag ttccatata 840
ttcatgcagc cttcaattgt ctctatctt aggggtttag tttgccctgc tctgcacccc 900
tgcctccgc aaacatggaa tctcagcaca cctaaaaata tcggtgggga tcagtgaatc 960
tgcactcaga tgactgtctt tcctggagtt ttctaaggaa ttgtttcaag cttaggattc 1020
cagactttga tttattaaaa tgtagtcacc tggt 1054

```

```

<210> SEQ ID NO 18
<211> LENGTH: 2138
<212> TYPE: DNA
<213> ORGANISM: Canis familiaris

```

```

<400> SEQUENCE: 18

```

```

gagacacacc actaccctca ccatgagccc caggcagccc ctggtcctgg tgttctcgtt 60
gctgggctgc tgctctgcag ctcccagacc acacaagccc accgttctgg tctttccagg 120
agacctgaga actaatctca ctgacaagca gctggcagag gaatatctgt ttcgctatgg 180
ctacactcaa gtggccgagc tgagcgacga caagcagtc ctgagtcgcg ggctgcggct 240
tctccagagg cgctggctc tgctgagac tggagagctg gacaaaacca cctggaggc 300
catcggggcc ccgctgctg gcgtcccga cctgggcaaa ttccagacct ttgagggcga 360
cctcaagtgg caccacaacg acatcactta ctggatacaa aactactcgg aagacttgc 420
ccgcgacgtg atcgacgacg cctttgcccg agccttcgcg gtctggagcg cggtgacacc 480
gctcaccttc actcgcgtgt acggcccga agccgacatc atcattcagt ttggtgtag 540
ggagcaegga gatgggtatc ccttcgatgg gaagaacggg cttctggctc acgcctttcc 600
tcccggcccc ggcaattcag gagacgccc cttcgacgac gaggagtat ggactctggg 660
caagggcgtc gtggttccga cccacttcgg aaacgcagat gggccccct gccacttccc 720
cttcacctc gagggccgct cctactcggc ctgcaccacg gacggccgct ccgatgacac 780
gccctggtgc agcaccacgg ccgactatga caccgaccgt cggttcggct tctgccccag 840
cgagaaactc tacaccacag acggcaatgg ggacggcaag cctcgcgtgt ttccgttca 900
cttcgagggc cgctcctact ccacgtgac caccgacggc cgtcggacg gctaccgctg 960
gtgctccacc accggcgact acgaccagga caaactctac ggcttctgcc caaccgagt 1020
cgattccgcg gtgaccgggg gcaactccgc cggggagccc tgtgtcttcc cttcatctt 1080
cctgggcaag cagtactcga cgtgaccag ggaggcccgc ggagatgggc acctctggtg 1140

```

-continued

---

```

cgccaccact tcgaactttg acagagacaa gaagtggggc ttctgcccgg accaaggata 1200
cagcctgttc cttgtggccg cccatgagtt cggccacgcg ctgggttttag atcattcacc 1260
ggtgccagaa gcgctcatgt accccatgta cagcttcacc gagggccccc cctgcatga 1320
agacgacgtg aggggcatcc agcatctgta cggctcctcg cctgaacctg agccacagcc 1380
tccaaaccgc ccgcccaccg ccccgccac cgtctgcgct actggtcctc ccaccaccg 1440
cccctcagag cgcccactg ctggcccac aggccccct gcagctggcc ccacgggtcc 1500
ccccactgct ggcccctctg agggccctac agtgccctgt gatccggcag aggatatatg 1560
caaagtgaac atcttcgacg ccatcgcgga gatcaggaac tacttgcat tcttcaagga 1620
agggaaagtac tggcgattct ccaagggcaa gggacgccg gtgcagggcc ccttccttat 1680
caccgacacg tggcctgcgc tgccccgcaa gctggactcc gcctttgagg acgggctcac 1740
caagaagact ttcttcttct ctggggcgca agtgtgggtg tacacaggca cgtcggtggt 1800
aggcccagg cgtctggaca agctgggctt gggcccggag gttaccaag tcaccgggc 1860
cctcccgcaa ggggggggta aggtgctgct gttcagcagg cagcgttctt ggagtttga 1920
cgtgaagacg cagaccgtgg atcccaggag cgcggctcg gtggaacaga tgtaccggg 1980
ggtgcccttg aacacgcatg acatcttcca gtaccaagag aaagcctact tctgccagga 2040
ccgcttctac tggcgtgtga attctcgaa tgaggtgaac caggtggacg aagtgggcta 2100
cgtgaccttt gacattttgc agtgccctga ggattaga 2138

```

&lt;210&gt; SEQ ID NO 19

&lt;211&gt; LENGTH: 1636

&lt;212&gt; TYPE: DNA

<213> ORGANISM: *Canis familiaris*

&lt;400&gt; SEQUENCE: 19

```

cgccggcgcc gctgaccgag gcctgctggg attccagaat tggagaagga ggcatgatga 60
agactctcct gctgctgggt gggctgctgc tgacctggga caatggaagg gttctgggag 120
accaggctgt ctcagacacc gagctccagg aaatgtccac cgaaggtagt aagtacatta 180
ataaggaat aaaaaatgct ctcaaggggg tgaacagat aaagacccta atagaacaga 240
caaacgaaga gcgcaaatca ctgctcagca acctggagga agccaagaag aagaaagg 300
atgcaactaa tgataccaag gattctgaaa cgaagctgaa ggcgtcccag ggggtgtgca 360
atgacacccat gatggccctc tgggaggagt gcaagccctg cctgaaacag acctgcatga 420
agttctacgc acgtgtctgc agaagtggct cggggctagt tggccaccag ctcgaggagt 480
tctgaaacca gagttctccc ttctacttct ggatgaatgg cgatcgtatc gactccctgc 540
tggagaacga ccggcagcag acccacgcc tggacgtcat gcaggacagc ttcaaccggg 600
catccagcat catggatgag cttttccagg acagattctt caccgggag ccccaggata 660
cttaccacta ctcacccttc agcttattcc agaggaggcc tttcttcaat cccaaatttc 720
gcatcgcccg gaacataata cctttcccta gattccagcc cttaacttc catgacatgt 780
ttcagccctt ctttgacatg atacaccagg ctcaacaggc catggatggt aacctgcata 840
gaattcctta ccacttcca attgaattcc cagaagaaga taaccgcacc gtgtgcaagg 900
agatccgtca caactccaca ggttgccctga aaatgaagga ccagtgtgaa aagtgccagg 960
agatcttctc agtggattgt tcttccaaca accctgcgca ggtccagctg cgacaggaa 1020
ttagtaattc cctccagatt cgggagaagt tcaccaagct ctacgacgag ctgctgcagt 1080
cctaccagga gaagatgttc aacacgtcct ccctgctgaa gcagctgaa gagcagttta 1140

```

-continued

---

```

gctgggtgtc ccaactggct aacctcactc aaagtgaaga cccattctat ctccagggtca 1200
cgacgggtggg ttctcagact tctgactcca atgttcccgt gggcttcaact aagggtggtg 1260
tgaagctcctt tgattctgac cccatcacgg tgatgatccc agaagcagtc tccaggaaca 1320
atcctaaatt tatggagact gtggcagaga aggctcttca ggagtatcgc caaaagcacc 1380
gagaggaatg agatgtgaac attgctcttc caaatatggg agggctgagt tcctttgacc 1440
ccaagatgag tgctaggccc ctagagagag ctctgcatgt caccaagtga ccaggccttg 1500
cctcgaggcg ctctgtcttc ctaaccacgc tttccttctc ctggactccg cactgtaacg 1560
cctacgtttg ctgatcatgg gaagaactcc tgtgtgccac taactcaata aaaccgcca 1620
ggaatctgaa aaaaaat 1636

```

&lt;210&gt; SEQ ID NO 20

&lt;211&gt; LENGTH: 2321

&lt;212&gt; TYPE: DNA

<213> ORGANISM: *Canis familiaris*

&lt;400&gt; SEQUENCE: 20

```

caggctgtgt ctccagggtc ctagccacag gcccggaag atgaggctcg cegtcgcgc 60
tctgctggcc tgcgcggtcc tggggctgtg tctggtgtc tcccctgaga aaactgtgag 120
atggtgcact gtctcaaatc atgaggccag taagtgttcc agtttcatgg aaaatatgaa 180
aaccgtcctt gaaaatggtc cttttgtcag ctgtgtgaag agaacctcct accttgagtg 240
catcaaggct atttgggcaa atgaagcaga tgctgtgaca cttgatgcag gtttggtgtt 300
tgaggcaggc ctgaacccct acaacctaaa gcctgtagtg gcagagtctc atgggaaaga 360
taagcaaac cgctattatg ctgtggctgt ggtgaagaag agcagtaagt tcaatctgaa 420
tcagctccaa ggcaagaaat cctgccacac aggccttggc aggtctgctg ggtggaacat 480
ccccatgggc ttactttatt ggaagttgac agagccactg gaatctcttc agaaggcagc 540
gtccagtttc ttcacggcca gctgtgttcc ctgtgtggat cggacagctt tcccctaaact 600
gtgtcaactg tgtgtgggga aaggacgga caagtgtgcc tgctccaacc atgagccata 660
cttcggctac tcgggtgctt tcaagtgtct gatggaggac gctggggaag tggcctttgt 720
caagcattca acagtgttgg agaacctgcc aaacaaagct gacctggacc agtatgagct 780
gctctgccc aacaacaaac gaaagccagt gggggaatat aagcagtgtc acctggccaa 840
ggtcccttcc catgctgttg tggcccgaag tgtggggggc aaggaagact tgatctggga 900
gcttctaaac caggcccagc aacattatgg caaagacaaa tctaaagtct tccagctcct 960
cagctcaact ttggggaagg acctgctgtt taaagactct gccgaagggt tttttaggat 1020
tcctcctaaa atggacacct ggctgtacct gggatatgag tatgtcactg ctcttcggaa 1080
cctaagggaa gatgtgccc cagatacccc aagggatgaa tgcaagaagg tgaatggtg 1140
tgcaagtagt caccatgaga ttgccaagtg tgatgagtgg agtgtaaca gcgaaggga 1200
aatagagtgt gaatcagcag agtctactga agactgtatt gccaaagatt cgaaaggaga 1260
ggctgatgcc atgagcttgg atggaggcta tatctacata gggggcagc gcggtctggt 1320
gcctgtcctg gcagagaact acaaaactca gggctctaca tgtagcaaca cagcagagga 1380
agggtatcct gctgtggccg tagttaagag attagataaa accatctctt ggaataatct 1440
gcaaggcagg aagtcctgcc aactgcagc agatagaact gcaggctgga acattcccat 1500
gggctgtctc tacaacagga tcaaccactg tgaatttgat aaatttttct ctcaaggctg 1560

```

-continued

---

```

tgcccccgga tctatgcgga attccagtct ctgtgctctg tgtattggct cagcaaatgt 1620
tccaggaaag gagtgtgttc ccaataacca tgagagatac tatggctaca caggggcttt 1680
caggtgtctg gttgagaagg gagatgtggc ctttgtgaaa gaccagactg tcttgcagaa 1740
cactggggga aagaacactg aagattgggc taagatctg aaggaagagg actttgagct 1800
gctgtgcct gatggccaaa ggaagtctgt ggacaaggct cctgagtgtc tcctagccaa 1860
agccccaaat catgctgtgg tctcaaggaa agacaaggca tcttgtgtta gcaaaatgtt 1920
acttgaccag cagcttttgt ttggaagaaa tggaaatgac tgctgggca agttttgctt 1980
gttccactca gcaaccaagg accttctgtt cagggacgac acacaatgtt tggccaaact 2040
tccagaggac acaacatata aatcatactt aggagcagag tatatcacgg ctgttgctaa 2100
cctgagacia tgctccactc cgaaactcct ggaggcctgc actttccata catcttaaaa 2160
tccaagaggc ggagacacac cagatgtgta tgggagctca cgtgacccat aagtccctc 2220
ggggcctcac tggcctgggt ggtttctctg gtttcacagt tcagcgggtg aacctctgta 2280
gaacattaaa taacattat tattggtgtt atcttttaaa a 2321

```

```

<210> SEQ ID NO 21
<211> LENGTH: 576
<212> TYPE: DNA
<213> ORGANISM: Canis familiaris

```

```

<400> SEQUENCE: 21

```

```

ctttccgaga ccgctccagg ttggtgctcg atacctgatg cccctgatgc cccgggaggg 60
ctgcccctcc aagaacacta gaaaaggggt ggagggaact cgggggaggg aggaggggtg 120
tccctgatcg ctcccacccc tgccttctgt tccagagagc cctccggcc ccgcatgag 180
gctcattctg tcgctcccgg ttttgggtgt ggttctgtcg atggttttgg aaggtccagc 240
ccccgccagc ggggcccagg aaatctccag cacttttgag cgcaccccgg ataagctgaa 300
ggagtttggt aacaccctgg aagacaaggc cggggcagcc attgagagca tcaagaagag 360
cgacattcct gcaaaagacc gaaactgggt ttctgaggct ttcaagaaag tgaaggagca 420
tctcaaaact gccttctcct gaacaccagg agagccgccc ctctactctg gcctgtgtgc 480
cccaggaggg ggcctgaaa tttccatccc cctggctcct tgccaaggac ttcattgatg 540
tcatgtctac ccccaacctc caataaaaaa cctata 576

```

```

<210> SEQ ID NO 22
<211> LENGTH: 1275
<212> TYPE: DNA
<213> ORGANISM: Canis familiaris

```

```

<400> SEQUENCE: 22

```

```

atgcccttgc tctggtgag aggattcttg gtggcgagtt gctggattat agtgccggagc 60
tccccgacct cggggcccga ggggcccggc gcggcccccg cctgccgggc ctgcgcgctc 120
accgccctgc ccagggatgc ccccaactcc cagcccgaga tgggtggaggc cgtcaagaag 180
cacatctca acatgtgca cttgaagaag agaccggaag tcaccagacc ggtgcccaag 240
gcggcgcttc tgaacgcgat ccgcaagctg cacgtaggca aagtcgggga gaacgggttc 300
gtggagatag aggatgacat cggcaggagg gcagaaatga atgaactcat ggagcagacc 360
tcggagatca tcacgtctgc ggaatcaggc acagccagga aaacgctgca ctttgagatt 420
tccaaagaag gcagtgacct gtcgggtggg gagcgtgcag aagtctggct ctctctcaaa 480
gtccccaagg ccaacaggac caggacaaa gtcaccatcc ggctcttgca gaagcacc 540

```

-continued

---

cagggcagct tggatgcggg ggaggaggcc gaggacatgg gcttcccga ggagaggaac	600
gaggtgttga tttctgaaaa ggtggtggac gcacggaaga gcacctggca catcttccct	660
gtctccagca gcatccagcg cttgctggac cagggcagga gctccctgga cgttcggatt	720
gcctgcgagc agtcccacga gacgggcgcc agcctggtgc tcctgggcaa gaagaagaag	780
aaggaggagg agggggaagg gaagaagaag gacggaggag acgcaggggc cgggggggac	840
gaggacaagg agcagtccca cagaccttcc ctcatgctgc aggcccgcca gtctgaagac	900
cacctcacc ggccggcgcg gcggggcctg gactgtgacg gcaaggtcaa catctgctgt	960
aagaaacagt tctttgtgag cttcaaggac attggctgga acgactggat cattgcccc	1020
tccggctatc acgccaacta ctgcgagggg gggtgccca gccacatagc aggcacgtcg	1080
gggtcctcgc tctccttca ctcgaccgtc atcaaccact accgctcgc gggtcacagc	1140
cccttcacca acctcaagtc gtgctgtgtg cccaccaagc tgagaccaat gtccatgctg	1200
tactacgatg atgggcaaaa catcatcaaa aaggacattc agaacatgat cgtggaggag	1260
tgccgggtgct catag	1275

---

What is claimed is:

1. A method of treating, ameliorating, or delaying the progression of a kidney disorder in a canine, comprising the steps of:

(a) diagnosing the kidney disorder by measuring the level of expression of at least one biomarker in a biological sample from the canine, wherein the at least one biomarker is selected from secreted frizzle-related protein-2 (SFRP2); matrilin-2 (Matn2); lumican (LUM); retinol binding protein 4 (rbp4); transferrin (TF); Apo-C-1 (ApoC1); and inhibin beta A (INHBA), and wherein the test sample is a sample of solid kidney tissue or a bodily fluid;

(b) identifying the canine as having a kidney disorder wherein differences in expression of one or more biomarkers in the sample relative to a control value for expression in a sample from a normal animal indicates the existence of a kidney disorder, and wherein the differences in expression are determined with a custom array consisting of two or more probes specific to the biomarkers selected from matrilin-2 (Matn2); lumican (LUM); retinol binding protein 4 (rbp4); transferrin (TF); Apo-C-1 (ApoC1); and inhibin beta A (INHBA); and

(c) treating, ameliorating, or delaying the progression of the kidney disorder in the canine identified in step (b); wherein the kidney disorder is indicated by a significant difference in expression of one or more biomarkers relative to control expression values wherein a significant difference in the case of increased expression is an increase of at least twofold and in the case of decreased expression is a decrease of at least 50% and wherein one or more biomarkers is selected from: matrilin-2 (Matn2); lumican (LUM); retinol

25

30

35

40

45

50

55

binding protein 4 (rbp4); transferrin (TF); Apo-C-1 (ApoC1); and inhibin beta A (INHBA), and wherein the canine has normal renal function, wherein normal renal function is determined by one or more of: glomerular filtration rate, urine protein level, blood creatinine level, urine creatinine level, creatinine clearance and blood urea nitrogen.

2. The method of claim 1, wherein the differential expression of step (b) is determined with a panel of one or more antibodies that specifically bind to the at least one biomarker.

3. The method of claim 1, wherein the differential expression of step (b) is determined by immunoassay to detect the at least one biomarker.

4. The method of claim 3, wherein the immunoassay is selected from the group consisting of a competitive binding assay, a non-competitive binding assay, a radioimmunoassay, an enzyme linked immunosorbent assay (ELISA), a sandwich assay, a precipitin reaction, a gel diffusion immunodiffusion assay, an agglutination assay, a fluorescent immunoassay, chemiluminescence immunoassay, immunoPCR immunoassay, a protein A or protein G immunoassay, Northern blot analysis, Western blot analysis, bead-based multiplexed assay and an immunoelectrophoresis assay.

5. The method of claim 1, further comprising use of one or more conventional diagnostic measurement selected from the group consisting of: determining levels of albumin, determining levels of serum cystatin C, determining levels of blood creatinine, determining levels of urinary creatinine, determining levels of creatinine clearance, determining levels of urinary protein, determining levels of blood urea nitrogen and determining glomerular filtration rate.

\* \* \* \* \*

专利名称(译)	用于诊断和治疗犬肾脏疾病的组合物和方法		
公开(公告)号	<a href="#">US9476868</a>	公开(公告)日	2016-10-25
申请号	US13/519048	申请日	2010-12-23
[标]申请(专利权)人(译)	希尔氏宠物营养品公司		
申请(专利权)人(译)	希尔思宠物营养, INC		
当前申请(专利权)人(译)	希尔思宠物营养, INC.		
[标]发明人	AL MURRANI SAMER MALLADI SUKHASWAMI GAO XIANGMING		
发明人	AL-MURRANI, SAMER MALLADI, SUKHASWAMI GAO, XIANGMING		
IPC分类号	C12Q1/68 G01N33/53 G01N33/50		
CPC分类号	G01N33/50 C12Q1/6883 C12Q2600/158 G01N2800/347 A61P13/12		
优先权	61/289773 2009-12-23 US		
其他公开文献	US20120283129A1		
外部链接	<a href="#">Espacenet</a> <a href="#">USPTO</a>		

摘要(译)

本发明提供了以下方法：诊断，设计和监测治疗计划，并监测肾脏疾病的状态，所述肾脏疾病的特征在于犬的肾功能，肾衰竭，肾小球滤过率降低或肾小球肾炎的异常丧失，其中通过利用从取自这种犬科动物的生物学测试样品中分离和测量的至少一种相关生物标志物，可以检测到肾脏疾病。本发明还涉及用于实施特定方法的组合物，试剂和试剂盒。

(continued on next page)

TABLE 3

List of Potential Biomarkers

Gene	Symbol in Canine/Human	Human Synonyms	Canine Synonym	Gene Description	mRNA	Protein
Lumican	LUM	LDC; SLRR2D			XM_539716.2	XP_539716.1
Collagen alpha 1 (III) chain, variant 12	COL3A1	EDS4A; FLJ34534			XM_858055.1	XP_863148