



US 20060024719A1

(19) **United States**(12) **Patent Application Publication**
Colgin et al.(10) **Pub. No.: US 2006/0024719 A1**(43) **Pub. Date: Feb. 2, 2006**(54) **METHODS AND DEVICES FOR DIAGNOSIS
OF APPENDICITIS****Publication Classification**(76) Inventors: **Mark A. Colgin**, Castle Rock, CO
(US); **John F. Bealer**, Englewood, CO
(US); **Richard Donnelly**, Fort Collins,
CO (US); **Diane Newman**, Castle Rock,
CO (US)(51) **Int. Cl.****C12Q 1/68** (2006.01)**G01N 33/53** (2006.01)**G01N 33/558** (2006.01)(52) **U.S. Cl. 435/6; 435/7.1; 436/514**


Correspondence Address:

GREENLEE WINNER AND SULLIVAN P C
4875 PEARL EAST CIRCLE
SUITE 200
BOULDER, CO 80301 (US)(57) **ABSTRACT**

A method is provided for diagnosing appendicitis in a patient that includes identifying at least one symptom of appendicitis in the patient and identifying the presence of at least one molecule differentially associated with appendicitis in a fluid or tissue sample of said patient. MRP-8/14 and haptoglobin are examples of molecules differentially associated with appendicitis. Devices and kits for performing the appendicitis assays of this invention are also provided. In one embodiment, the device is in a flow-through immunoassay format for testing blood samples. Further, methods for screening for molecules differentially associated with appendicitis are provided that include the use of samples from patients being operated on for suspected appendicitis.

(21) Appl. No.: **11/189,120**(22) Filed: **Jul. 25, 2005****Related U.S. Application Data**

(60) Provisional application No. 60/590,631, filed on Jul. 23, 2004.

Relative MRP 8/14 serum values




FIGURE 1A




FIGURE 1B




FIGURE 2




FIGURE 3

Relative MRP 8/14 serum values




FIGURE 4




FIGURE 5A




FIGURE 5B




FIGURE 6A




FIGURE 6B




FIGURE 7A




FIGURE 7B

METHODS AND DEVICES FOR DIAGNOSIS OF APPENDICITIS

CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] This application claims priority to U.S. provisional application Ser. No. 60/590,631 filed Jul. 23, 2004.

BACKGROUND OF THE INVENTION

[0002] Appendicitis is a common acute surgical problem affecting human beings of a wide age range. There are approximately 700,000 cases annually in the United States. A large proportion of cases occur in the 10 to 30 age group. An accurate diagnosis at a sufficiently early stage is a significant factor in achieving a successful outcome.

[0003] Many people present to their physician with symptoms suggestive of appendicitis but caused by other ailments such as viral infections. Differentiating the appendicitis patients from those affected with other ailments is a daunting clinical task that physicians face daily. While medical science has an excellent understanding of appendicitis and its treatment, it is very limited in its ability to accurately recognize or diagnose the disease.

[0004] Complicating the goal of an accurate and early diagnosis is the considerable overlap of genuine appendicitis with other clinical conditions. There appears to be no individual sign, symptom, test, or procedure capable of providing a reliable indication of appendicitis. Imaging technology is inadequate in identifying and characterizing the appendix, especially in the early stages of the disease when treatment is likely to be most effective. Imaging technology is further handicapped by its expense and its dependence upon the availability of highly trained and experienced people to interpret the studies. This limitation affects thousands of people every year by inaccurately diagnosing their problem or by delaying the accurate diagnosis. In cases of appendicitis, delays in diagnosis are the single most important factor leading to worsening of the condition and more complications related to the disease. The misdiagnosis of appendicitis can lead not only to unnecessary surgery but also to delay of proper therapy for the actual underlying condition.

[0005] A dilemma for surgeons is how to minimize the negative appendectomy rate without increasing the incidence of perforation among patients referred for suspected appendicitis. What is desperately needed to more effectively treat this very common ailment is a simple, reliable diagnostic test that is capable of recognizing the earliest stages of the disease process.

[0006] The typical pathogenesis in appendicitis begins with obstruction of the lumen, although an initial inflammation of the organ can precede and even contribute to the obstruction. The secreted mucus of the appendix fills the closed lumen, causing an increase in intraluminal pressure and distension. The increased intraluminal pressure can exceed the level of capillary perfusion pressure, resulting in perturbation of normal lymphatic and circulatory drainage. Ultimately the appendix can become ischemic. The appendix mucosa is compromised, which can allow invasion of intraluminal bacteria. In advanced cases, perforation of the appendix may also occur with spillage of pus into the peritoneal cavity.

[0007] Currently, the diagnosis of appendicitis is difficult, and the difficulty persists during various stages in the progression of the condition. The following represents a hypothetical portrayal of stages and associated clinical presentations. Artisans of ordinary skill will recognize that a considerable degree of variation will occur in a given patient population.

[0008] At the earliest stages of inflammation, a patient can present with a variety of non-specific signs and symptoms. Upon obstruction, presentation can involve periumbilical pain, mild cramping, and loss of appetite. The progress toward increased luminal pressure and distension can be associated with presentation involving the localization of pain to the right lower quadrant of the abdomen, nausea, vomiting, diarrhea, and low grade fever. If perforation occurs, a patient can present with severe pain and high fever. At this very advanced stage, sepsis can be a serious risk with a potentially fatal outcome.

[0009] Practitioners currently use several tools to aid in appendicitis diagnosis. These tools include physical examination, laboratory tests, and other procedures. Routine laboratory tests include complete blood count (CBC) with or without differential and urinalysis (UA). Other tests include a computed tomography (CT) scan of the abdomen and abdominal ultrasonography. Procedures can include, for example, laparoscopic examination and exploratory surgery.

[0010] Flum et al. attempted to determine whether the frequency of misdiagnosis preceding appendectomy has decreased with increased availability of certain techniques (Flum D R et al., 2001). These techniques included computed tomography (CT), ultrasonography, and laparoscopy, which have been suggested for patients presenting with equivocal signs of appendicitis. Flum et al. concluded as follows: "Contrary to expectation, the frequency of misdiagnosis leading to unnecessary appendectomy has not changed with the introduction of computed tomography, ultrasonography, and laparoscopy, nor has the frequency of perforation decreased. These data suggest that on a population level, diagnosis of appendicitis has not improved with the availability of advanced diagnostic testing." The rate of misdiagnosis of appendicitis is about 9 percent in men and about 23.2 percent in women (Neary, W., 2001).

[0011] Myeloid-related Protein Complex 8/14 (MRP-8/14) is a heterodimeric complex associated with acute inflammatory conditions (for review see Striz and Trebichavsky, 2004). The complex belongs to the S100 superfamily of proteins and is also referred to S100A8/9, L1, macrophage inhibitory related protein and calprotectin. The heterodimer consists of an 8 kilodalton (MRP-8) and 14 kilodalton (MRP-14) subunit. MRP-8 and MRP-14 are alternatively named S100A8/calgranulin and S100A9/calgranulin b, respectively. MRP-8/14 is a calcium binding protein originally discovered in macrophages. Neutrophils expressing high concentrations of MRP-8/14 are found in a variety of inflammatory conditions, including rheumatoid arthritis, inflammatory bowel disease and allograft rejections (Frosch et al., 2000; Limburg et al., 2000; Burkhardt et al., 2001).

[0012] MRP-8/14 is not always diagnostic of inflammation. For example, it does not reliably indicate the presence of inflammatory diverticuli (Gasché, C. 2005). Lymphocytes do not generally contain MRP-8/14 (Hycult Biotechnology, Monoclonal Antibody to Human S100A8/A9), and therefore

MRP-8/14 is not diagnostic of inflammation characterized by the presence of lymphocytes but not neutrophils. Also, this protein is not always associated with opportunistic infections (Froland, M. F., et al., 1994).

[0013] Haptoglobin is an acute phase protein that binds free hemoglobin following hemolysis. The haptoglobin-hemoglobin complex is removed by the liver. Haptoglobin is a heterotetramer composed of two alpha and two beta subunits. The alpha and beta units are derived from a single polypeptide chain precursor that is enzymatically cleaved to produce the subunits. The molecular weights of the subunits are approximately 9 kd-18 kd and 38 kd for alpha and beta, respectively.

[0014] In addition to being a hemoglobin scavenger, haptoglobin has a wide range of biological functions (Dobryszka, 1997). Haptoglobin has been shown to be upregulated and modulate the immune response in certain infection and inflammatory conditions perhaps by regulating monocyte function (Arredouani et al., 2005). The alpha subunit has been demonstrated to be a potentially useful serum marker for ovarian cancer (Ye et al., 2003).

[0015] The ability to accurately diagnose appendicitis would be greatly augmented by the identification of molecules differentially associated with appendicitis.

SUMMARY OF THE INVENTION

[0016] This invention provides a method for diagnosing appendicitis in a patient comprising identifying at least one classical symptom of appendicitis in said patient and identifying the presence of at least one molecule differentially associated with appendicitis in a fluid or tissue sample of said patient. It is recognized in the art that the diagnosis of appendicitis is difficult, and that it is often misdiagnosed. Thus the term "diagnosing appendicitis" as used herein does not necessarily mean diagnosing appendicitis with more than usual accuracy. However, in fact, the methods of the present invention have been shown to provide improvements in correct diagnosis, with almost no false positives and few false negatives.

[0017] The term "differentially associated" with respect to a molecule "differentially associated with appendicitis" refers: (1) to a molecule present in a patient with appendicitis and not present in a patient not having appendicitis; (2) to a molecule whose relative level (amount) is distinguishing between appendicitis and non-appendicitis; (3) to a molecule present, or present at a level, in conjunction with the presence of other symptoms of appendicitis, that is diagnostic of appendicitis; and/or (4) to a molecule present, or present at a level, in conjunction with the lack of symptoms associated with conditions other than appendicitis in which the presence of the molecule occurs, that is diagnostic of appendicitis.

[0018] The diagnostic level of such a molecule is also referred to herein as the "threshold amount" or "threshold level." The molecules differentially associated with appendicitis are preferably protein antigens.

[0019] Classical symptoms of appendicitis include: pain in the abdomen; pain that starts near the navel, then moves to the lower right quadrant of the abdomen; anorexia (loss of appetite); trouble eating accompanied by sleepiness; nausea starting after onset of pain; vomiting starting after onset of

pain; vomiting accompanied by fatigue; constipation; small stools with mucus; diarrhea; inability to pass gas; low-grade fever; abdominal swelling; pain in the abdomen worsening; tenesmus (feeling of needing to move the bowels); high fever; and leukocytosis. Increased plasma viscosity is also associated with appendicitis. In one embodiment of the invention at least two or more symptoms of appendicitis are identified.

[0020] In one embodiment of this invention patients are screened to determine whether or not they have an "interfering condition," i.e., another condition in which the molecule is present in the type of sample being tested. Patients are tested for the presence of the molecule if they do not have such an interfering condition; or are tested for the presence of appendicitis-diagnostic levels of the molecule if they do have such an interfering condition. Appendix-diagnostic levels when the patient has an interfering condition are levels higher than those present in patients who have the interfering condition but do not have appendicitis. Interfering conditions include recent allograft; septicemia; meningitis; pneumonia; tuberculosis; rheumatoid arthritis; gastrointestinal cancer; inflammatory bowel disease; skin cancer, periodontitis, preeclampsia, and AIDS.

[0021] A sample can be a fluid or tissue, and can contain whole blood, plasma, serum, milk, urine, saliva and/or cells. Fecal samples may also be used. Preferably tissue and fecal samples are liquefied before testing.

[0022] In one embodiment of this invention two or more molecules differentially associated with appendicitis are tested for. Identification of additional molecules provides greater accuracy to the method.

[0023] One molecule differentially associated with appendicitis is MRP-8/14. Another is haptoglobin. Both these molecules can be tested for in diagnosing appendicitis.

[0024] MRP-8/14 levels in the range of about 1 to about 11 $\mu\text{g/ml}$ are present in patients without appendicitis. Levels higher than this provide increased accuracy in diagnosing appendicitis. Levels higher than about 10, 11, 13, 15 or 20 $\mu\text{g/ml}$ of MRP-8/14 can be used to diagnose appendicitis. Haptoglobin levels in the range of about 27-139 mg/dL are found in patients without appendicitis. Levels higher than this, e.g., higher than 125, 130, 135, 139 and 150 provide increased accuracy in diagnosing appendicitis.

[0025] Other molecules that can be tested for in the methods of this invention, or that can be tested for in addition to the foregoing molecules, include unique structural proteins of the gastrointestinal tract, stress-related inflammatory mediators, immunologic factors, indicators of intestinal bacterial flora, Plasminogen Activator Inhibitor-1, fatty acid binding proteins, nuclear factor kappa beta (NF κ B), specific appendix antigens (HLA-DR), inflammation associated antigens; and nucleic acids coding for any of the foregoing, including nucleic acids coding for MRP-8/14 and haptoglobin. Methods for testing for the presence of nucleic acids are known to the art.

[0026] The methods of this invention involving obtaining a first sample from a patient suspected of having appendicitis can also comprise identifying at least one molecule differentially associated with appendicitis by a process including obtaining a second fluid or tissue sample from a second patient, wherein the second patient has appendicitis;

obtaining a third fluid or tissue sample from a third patient wherein the third patient has a non-appendicitis condition characterized by at least one symptom of appendicitis; and analyzing the second and third samples so as to detect a molecule differentially associated with the appendicitis in the second patient, and then identifying the presence of that molecule, or presence of an increased level of that molecule, in the first sample, thereby diagnosing appendicitis. Candidate molecules for this process of identifying molecules differentially associated with appendicitis include unique structural proteins of the gastrointestinal tract, stress-related inflammatory mediators, immunologic factors, indicators of intestinal bacterial flora, Plasminogen Activator Inhibitor-1, fatty acid binding proteins, nuclear factor kappa beta (NF κ B), specific appendix antigens (HLA-DR), inflammation associated antigens, and nucleic acids coding for any of the foregoing.

[0027] This invention also provides a method for identifying a molecule differentially associated with appendicitis, the method comprising obtaining a sample from each of a plurality of patients who are undergoing surgery for suspected appendicitis; determining during surgery whether each said patient has appendicitis or not; and analyzing said samples for the presence of a molecule differentially associated with appendicitis. The samples can be blood samples or samples of appendix tissue. This method can also include determining the amount of each molecule found to be differentially associated with appendicitis in the sample. In one embodiment of the invention, following identification of the molecule in tissue, it is also identified in plasma. This requires that samples of blood be taken from patients suspected of having appendicitis. The amount of the molecule differentially associated with appendicitis in patients who have appendicitis compared with those who do not is also determined.

[0028] The methods for diagnosing appendicitis of this invention can include using test devices, e.g., cartridge test devices and dipstick test devices, and/or other means for determining the presence or absence of a molecule differentially associated with appendicitis, e.g., performing western blots, northern blots, ELISA tests, protein function tests, PCR and other assays known to the art. In testing molecules differentially associated with appendicitis that are present in patients without appendicitis, but upregulated in patients with appendicitis, assays that test for the relative amount of the molecule present in patient fluids or tissues as well as the mere presence of the molecule are required. Cartridge immunoassays can be designed to provide information on relative amounts of such molecules as described herein. Other assays known to the art including ELISAs and hospital assay devices such as the Synchron LX system of Beckman Coulter can be used to provide the amount of such molecules present in the patient, which can then be compared with amounts present in patients without appendicitis to determine whether or not the patient has appendicitis.

[0029] The methods for diagnosing appendicitis can include performing an immunological assay using a monoclonal or polyclonal antibody to the molecule differentially associated with appendicitis. Such antibodies are known to the art or can be generated by means known to the art without undue experimentation.

[0030] This invention also provides an immunoassay test device for detecting the presence of a molecule differentially

associated with appendicitis in a sample. The device comprises a first monoclonal or polyclonal antibody specific to the molecule, a support for the first monoclonal or polyclonal antibody, means for contacting the first monoclonal or polyclonal antibody with the sample, and an indicator capable of detecting binding of the first monoclonal or polyclonal antibody with the molecule.

[0031] Detecting binding of the antibody with the molecule can include binding the antibody/molecule complex to a second, labeled antibody which binds to the molecule or to the antibody of the complex.

[0032] Test devices can be in the form of cartridges, dipsticks, or other conformations known to the art. The test device can also be part of a kit which can contain instructions for use, instructions for comparison of test results with results of the same test done on non-appendicitis patient, additional reagents, such as cells or fluids from non-appendicitis patients, and other reagents known to the art. These types of assay devices are known to the art and described, e.g., in U.S. patent publication No. 2003/0224452.

[0033] The methods for diagnosing appendicitis can include comparing the level of the molecule in the sample with a background level of the same molecule in persons not having appendicitis. This comparison can be made by any means known to the art. It can include comparing sample results with results from a second sample taken from a person known not to have appendicitis, or comparing sample results with a photograph or other representation of results from a person not having appendicitis. Test devices having means for masking non-appendicitis levels, e.g. a support having the same color or tone as indicators showing non-appendicitis levels, or a filter having the same color or tone as a non-appendicitis level, so that only higher, appendicitis-indicating levels of the molecule are detectable, e.g., by eye, can also be used. The methods of this invention can include use of control fluids having background levels of the molecule typical of non-appendicitis samples, as well as colored supports and/or light filters as discussed above.

[0034] When the sample is blood, the method can also include processing the blood by a means known to the art, such as filtration or centrifugation, for separating plasma or serum which is to be assayed.

[0035] Antibody supports are known to the art. In an embodiment of this invention, antibody supports are absorbent pads to which the antibodies are removably or fixedly attached. In the devices of this invention, any indicator means known to the art to detect antibody binding with the molecule can be used. The indicator means can include second, labeled, monoclonal or polyclonal antibodies which bind to the selected protein, which preferably bind to a substantially different epitope on the selected protein from that to which the first monoclonal or polyclonal antibodies bind, such that binding of the first monoclonal or polyclonal antibody will not block binding of the second antibody, or vice versa. The indicator means can also include a test window through which labeled antibodies can be viewed. Any label known to the art can be used for labeling the second antibody. In an embodiment of this invention, the label is colloidal gold. The second antibody can be monoclonal or polyclonal. In an embodiment of this invention, the first antibody is a polyclonal or a monoclonal antibody made using a specific polypeptide sequence of the molecule dif-

ferentially associated with appendicitis, and the second antibody is a different monoclonal or polyclonal antibody which binds to a different site of the molecule or binds to the first antibody. Antibodies for MRP-8 and MRP-14 are commercially available through Cell Sciences, Canton, M A. Monoclonal antibodies to haptoglobin useful in the methods of this invention are also known to the art, e.g., as described in U.S. Pat. No. 5,552,295.

[0036] In one embodiment of this invention, the sample to be assayed is a liquid, and the immunoassay test device is a lateral flow device comprising inlet means for flowing a liquid sample into contact with the antibodies. The test device can also include a flow control means for assuring that the test is properly operating. Such flow control means can include control antigens bound to a support that capture detection antibodies as a means of confirming proper flow of sample fluid through the test device. Alternatively, the flow control means can include capture antibodies in the control region which capture the detection antibodies, again indicating that proper flow is taking place within the device.

[0037] Methods for detecting the presence of a molecule differentially associated with appendicitis using the foregoing devices are also provided, the methods comprising: providing an immunoassay test device of this invention; contacting a first antibody with a sample; and reading an indicator which is capable of detecting binding of the first antibody. Preferably, binding indicates appendicitis in the patient being tested. Methods of using these devices can be performed in the doctor's office, emergency room, or surgery, rather than requiring sending the patient or the sample to a separate laboratory.

[0038] The devices of this invention are useful for testing the above-mentioned samples. When cells are tested, e.g., when the molecule differentially associated with appendicitis is suspected to be in blood or tissue cells rather than serum, the method and/or device can include a cell-lysing step or means using detergent, puncture or other physical or chemical process known to the art.

BRIEF DESCRIPTION OF THE FIGURE

[0039] **FIG. 1:** Two-dimensional electrophoresis image of proteins from (A) normal and (B) diseased appendix tissue. Proteins were separated by isoelectric focusing on the x axis and by molecular weight on the y axis. The molecular weight in kilodaltons is shown on the left. The arrow indicates the upregulated protein, AP-93.

[0040] **FIG. 2:** MRP-14 western blot analysis of normal (N) and diseased (A) tissue. The numbers are sample ID numbers. Molecular weights are shown in kilodaltons.

[0041] **FIG. 3:** MRP-8 western blot analysis of normal (N) and diseased (A) tissue. The numbers are sample ID numbers. Molecular weights are shown in kilodaltons.

[0042] **FIG. 4:** Relative levels of MRP-8/14 in normal and appendicitis serum as determined by ELISA. The levels are given as a fraction of the mean for the patients not having appendicitis, said fraction also being referred to herein as a "fold increase." Dark bars represent samples from patients having appendicitis. White bars represent samples from patients not having appendicitis.

[0043] **FIG. 5:** Two-dimensional electrophoresis image of proteins in depleted serum samples from (A) normal and (B)

appendicitis patients. Proteins were separated by isoelectric focusing on the x-axis and by molecular weight on the y-axis. The molecular weight in kilodaltons is shown in the right. The tailed arrow indicates the upregulated protein, AP-77 (haptoglobin alpha subunit). The untailed arrow indicates a control protein that is equally abundant in diseased vs. normal.

[0044] **FIG. 6:** Two-dimensional electrophoresis image of proteins from (A) normal and (B) diseased (perforated) appendix tissue. Proteins were separated by isoelectric focusing on the x axis and by molecular weight on the y axis. The molecular weight in kilodaltons is shown on the left. The arrow indicates the upregulated protein, AP-91 (haptoglobin alpha subunit).

[0045] **FIG. 7:** Haptoglobin distribution. Haptoglobin western blot analysis of normal (N) and diseased (A) tissue. The numbers are sample ID numbers. Molecular weights are shown in kilodaltons. The alpha and beta subunits are >20 kd and 38 kd, respectively.

DETAILED DESCRIPTION

[0046] The vermiform appendix is recognized as a separate organ from the large and small intestines. It extends as a finger-like pouch from the base of the ascending colon, which is also called the cecum. The appendix, like the large intestine, is hollow and composed of the same three tissue layers. These three layers are a mucosa, muscularis and a serosa. The appendiceal lumen communicates with the lumen of the cecum via a round opening (os) through which the appendix adds its secretions to the fecal stream. These secretions are excess mucus produced from the appendiceal mucosa. In addition to containing mucus, the appendix also contains numerous bacteria common to the right colon. Obstruction of the appendiceal lumen is the dominant factor causing acute appendicitis. While fecaliths are the usual cause of appendiceal obstruction, hypertrophied lymphoid tissue, inspissated barium from previous x-ray studies, vegetable and fruit seeds, and intestinal worms like ascarids can also block the appendiceal lumen.

[0047] Following luminal obstruction an escalating cycle of events ensues. The proximal obstruction of the appendix produces a closed-loop obstruction that blocks the normal flow of appendiceal mucus into the cecum. The continuing normal secretion of the appendiceal mucus very rapidly fills the luminal capacity of the appendix (approximately 0.1 cc). Once the luminal capacity of the appendix is reached additional mucus production from the obstructed appendix rapidly elevates the intraluminal pressure within the organ. This elevated intraluminal pressure is exerted outward against the appendiceal wall and causes the appendix to distend. Distention stimulates nerve endings of the visceral afferent pain fibers, producing vague, dull, diffuse pain in the midabdomen or lower epigastrium. Peristalsis is also stimulated by the rather sudden distention, so that some cramping may be superimposed on the visceral pain early in the course of appendicitis.

[0048] Distention of the appendix continues, not only from continued mucosal secretion, but also from rapid multiplication of the resident bacteria of the appendix. As pressure in the organ increases, venous pressure within the appendiceal wall is exceeded. This rising intraluminal pressure then occludes capillaries and venules, but arteriolar

inflow continues, resulting in engorgement and vascular congestion. Distention of this magnitude usually causes reflex nausea and vomiting, and the diffuse visceral pain becomes more severe. The inflammatory process soon involves the serosa of the appendix and in turn parietal peritoneum in the region, producing the characteristic shift in pain to the right lower quadrant (RLQ). The disease process is fairly advanced when pain is localized to the RLQ.

[0049] The mucosa of the gastrointestinal tract, including the appendix, is very susceptible to impaired blood supply. Thus mucosal integrity is compromised early in the process, allowing bacterial invasion of the deeper tissue layers. This bacterial invasion leads to appendiceal destruction and systemic liberation of various bacterial toxins. Fever, tachycardia, and leukocytosis develop as a consequence of this systemic release of dead tissue products and bacterial toxins. As progressive appendiceal distention rises, encroaching on the arteriolar pressure, ellipsoidal infarcts develop in the antimesenteric border of the appendiceal serosa. As distention, bacterial invasion, compromise of vascular supply and infarction progress, perforation occurs through one of the infarcted areas on the antimesenteric border. This perforation then releases the bacteria and its toxins into the abdominal cavity.

[0050] Appendicitis has been called the "great imitator," as its symptoms are frequently confused with those of other conditions. This confusion stems from the nonspecific nature of the pain early in its course and the variability in how appendicitis progresses. Pain in the right lower quadrant of the abdomen is the hallmark of appendicitis but this is not typically what the patient first perceives. When the appendiceal lumen first obstructs, the patient will have few if any symptoms because the appendiceal lumen has not yet had the chance to fill with mucus. The time required to fill the appendiceal lumen is proportional to the lumen volume available behind the obstruction. This is variable and unpredictable, as that volume is dependent upon the individual's appendix size and precisely where the fecalith or other obstruction is located along that length. Should the fecalith or other obstruction be close to the tip of the appendix the available volume is relatively small and the time to symptoms or perforation short. In contrast, the opposite will be true should the fecalith or other obstruction be near the base of the appendix and provide for the largest possible appendiceal volume.

[0051] Once the appendix begins to distend, the appendicitis patient will begin to experience a nonspecific discomfort usually in the mid portion of the abdomen. This discomfort can be easily confused with common ailments such as indigestion, constipation or a viral illness. Continued appendiceal distention is also accompanied by some nausea and frequently vomiting. Rarely is the vomiting severe or unrelenting, which reinforces the confusion with common ailments.

[0052] Later in the progression of appendicitis, inflammation will have progressed to the outermost layer of the appendix. This outermost layer is called the serosa and it touches the inner lining of the abdominal cavity called the peritoneum. This contact irritates the peritoneum, producing peritonitis that is perceived by the appendicitis patient as focal pain wherever the appendix is touching the perito-

neum. This too can vary between different individuals. The appendix is most usually located in the right lower quadrant under an area known as McBurney's point. McBurney's point is a position on the abdomen that is approximately two-thirds of the distance from the anterior superior iliac spine in a straight line toward the umbilicus. The appendix can, however, reside in other locations in which case the peritonitis produced by the appendix will be in an atypical location. This again is a common factor producing an erroneous diagnosis and delays surgical treatment in cases of appendicitis.

[0053] Regardless of its location, if appendicitis is allowed to progress the organ will eventually perforate. This contaminates the abdominal cavity around the perforated appendix with bacteria producing a severe infection. This infection will usually lead to a localized intra-abdominal abscess or phlegmon and can produce generalized sepsis.

[0054] To identify molecules differentially associated with appendicitis, a proteomic approach was used. A protein complex, MRP-8/14, that is present in appendix tissue in patients with acute appendicitis was identified. The highly correlative nature of this complex with appendicitis led us to examine MRP-8/14 serum levels in patients with apparent appendicitis. MRP-8/14 is significantly elevated ($p < 0.02$) in patients with appendicitis as compared to levels in patients with apparent appendicitis yet having no appendiceal inflammation. The source of MRP-8/14 in the serum is the inflamed appendix tissue. This is consistent with the known functions of MRP-8/14.

[0055] The role of MRP-8/14 in inflammation is not fully understood but it does seem to play a vital role in retaining leukocytes in microcapillaries. Extracellular MRP-8/14 interacts with endothelial cells by binding to heparin sulfate and specifically carboxylated glycans (Robinson et al., 2002). The intracellular signal pathways and effector mechanisms induced by binding of MRP-8/14 to endothelial cells are not well defined. However, interaction of MRP-8/14 with phagocytes increases binding activity of the integrin receptor CD11 b-CD18. This is one of the major adhesion pathways of leukocytes to vascular endothelium (Ryckman et al., 2003). It is believed that the MRP-8/14 utilizes the receptor for advanced glycation end products (RAGE). A relative of MRP-8/14, S100A12, is a specific ligand of RAGE expressed by endothelial cells and their interaction activates NF-kappa B binding in these cells (Hsieh et al., 2004). The NF-kappa B binding subsequently induces expression of many proinflammatory molecules, such as various cytokines or adhesion molecules. Thus, release and extracellular functions of S100 proteins represent a positive feedback mechanism by which phagocytes promote further recruitment of leukocytes to sites of inflammation. Taken together, these proteins appear to play a role in a fundamental inflammatory response in certain inflammatory conditions, and are excellent markers of appendix tissue inflammation.

[0056] Neutrophils are white blood cells that are the first to migrate from the circulation into sites of inflammation. Within neutrophils, constituting approximately 40% of total cytosolic proteins is the MRP-8/14 complex. This protein is specifically expressed only in cells of macrophage lineage, making blood monocytes and acutely activated macrophages other potential white blood cell sources of these pro-

teins. MRP-8/14 is not usually expressed in lymphocytes nor resident macrophages or those macrophages involved in chronic inflammation. These two proteins are also known to be independently expressed by mucosal epithelium in specific states of acute inflammation.

[0057] In the case of appendicitis, the luminal obstruction and the resultant distention of the appendiceal wall triggers an inflammatory response. The circulating neutrophils are then recruited into the area, as are activated macrophages. While the expression of this protein complex is related to the activity of the macrophages in inflammation, the exact relationship between MRP-8/14 and cellular activity is not fully known. What is known is that the intracellular distribution of MRP-8/14 varies with the activation state of macrophages. Normal macrophages contain the complexes in the cytosol, but once stimulated, MRP-8/14 translocates from the cytosol to the cell membrane (specifically with the proteins of the cytoskeleton). This would imply that MRP-8/14 may be related to cell movement, phagocytosis or inflammatory signal transduction. The roles of cellular movement and signal transduction may also explain why MRP-8/14 is produced directly from vascular epithelium such as that lining the blood vessels within the appendix.

[0058] Regardless of its role in certain inflammatory conditions, MRP-8/14's abundance within cells of acute inflammation makes it an excellent detector and monitor of acute appendicitis. The first step in the inflammatory process is the recruitment of neutrophils and macrophages to a specific site. In our study, the specific site is the appendix, where those MRP-8/14-containing cells will engage the offending stimulus. This engagement will usually result in MRP-8/14 cell death and the liberation of MRP-8/14 from either the cytosol or cell membrane into the patient's circulation. At the same time, the mucosal linings of the appendix will start to produce and release MRP-8/14 to facilitate macrophage migration or inflammatory amplification. This process will then escalate as increasing amounts of MRP-8/14 cells are recruited by the appendicitis to ultimately release more MRP-8/14 into the circulation. Other examples of inflammatory states causing increases of extracellular MRP-8/14 and the tendency of these increases of MRP-8/14 to correlate with extent of inflammation are known. Specifically, chronic bronchitis, cystic fibrosis and rheumatoid arthritis are all associated with elevated serum levels of MRP-8/14 and the severity of these diseases is generally proportional to the serum levels of MRP-8/14 detected.

[0059] The physiological role of MRP-8/14 makes it an ideal clinical marker for acute appendicitis. As patients with appendicitis are generally young and healthy, they generally produce a vigorous inflammatory response. This vigorous response is believed to liberate MRP-8/14 in the earliest stages of the disease, which then escalates as appendicitis progresses. Additionally, the diseases known to be associated with elevated levels of MRP-8/14 are not common in this younger age group and usually do not produce symptomology similar to appendicitis. Finally, as MRP-8/14 is not located in nor associated with lymphocyte proliferation, this marker is not believed to be elevated in viral infections. This is an especially powerful advantage for diagnosing appendicitis, as viral infections are one of the most common imitators of appendicitis.

[0060] Haptoglobin was also identified in this invention as a useful marker for appendicitis. A differential proteomic

screen of depleted serum identified haptoglobin as a marker for appendicitis. A second differential screen of appendix tissue confirmed that haptoglobin is upregulated in the appendix tissue of patients with appendicitis. This finding was confirmed by western blotting of tissue protein. In particular the alpha subunit isoforms were present only in diseased tissue. Since haptoglobin is a plasma protein, it is highly valuable as a biomarker for appendicitis.

EXAMPLES

Example 1

MRP-8/14

[0061] The objective of this study was to identify a tissue-specific marker that could contribute to the decision matrix for diagnosing early acute appendicitis. A proteomic screen was used to identify a protein in the appendix specifically upregulated in acute appendicitis. MRP-8/14 was identified as present both in the diseased appendix and in serum of acute appendicitis patients.

[0062] Materials and Methods

[0063] Specimen and Serum Collection. All patients enrolled in this study were treated according to accepted standards of care as defined by their treating physicians. Prior to being approached for inclusion in our study, all patients were evaluated by a surgeon and diagnosed by that surgeon as having appendicitis. The treating surgeon's plans for these appendicitis patients included an immediate appendectomy. The specifics of all treatments such as use of antibiotics, operative technique (either open or laparoscopic) were determined by the individual surgeon.

[0064] Exclusion Criteria: Any patients with pre-existing chronic inflammatory diseases such as asthma, rheumatoid arthritis, inflammatory bowel disease, psoriasis, or neutropenia. Pregnancy was also considered an exclusion criterion.

[0065] An investigator counseled all patients about the study and informed consent was obtained. At the time of informed consent, the subject was assigned an identification number and non-personal demographic and clinical information was obtained (age, sex, race, duration of symptoms, white blood count (WBC), results of imaging studies, etc). At the time of surgery, following induction of general anesthesia, a whole blood sample (5-10 cc volume) was obtained via peripheral venopuncture. This blood specimen was then placed on ice. As soon as possible, a small sample (approximately 1 gram) of inflamed appendix was taken from the pathologic specimen and also placed on ice. The iced blood specimens were then centrifuged for 20 minutes at 3000 rpm and the separated serum isolated. This isolated serum and the piece of appendix tissue were then stored separately, frozen at -80°C .

[0066] Appendicitis Tissue Processing. Appendix tissue from appendectomy patients was harvested and stored at -80°C until processed. Individual tissue samples were ground into powder using a sterile mortar and pestle under liquid nitrogen. Protein was extracted from tissue powder by incubating at 37°C in Extraction Buffer (0.025M Tris-base, 200 mM Sodium Chloride, 5 mM EDTA, 0.1% Sodium Azide, pH 7.5). Samples were centrifuged for 10 minutes at 14K rpm. Supernatants were stored at -80°C until analysis.

[0067] 2D Gel Analysis of Extracted Tissue Samples. 2D gel analysis was performed on depleted serum samples and extracted tissue samples. Isoelectric focusing (IEF) and SDS-PAGE were performed according to the Zoom (Invitrogen) protocol for 2D Gel analysis. Equal quantities of protein were analyzed on each gel.

[0068] Comparisons between negative serum gel and positive serum gel were made to determine which proteins were present in positive samples and absent in negative samples. Candidate gel spots were identified and submitted to MALDI-TOF protein identification analysis (Linden Biosciences).

[0069] Western Blot Analysis of Extracted Appendix Tissue Samples. Samples (10 μ g) were subjected to standard Laemmli SDS-PAGE and proteins were transferred to nitrocellulose membrane for western blot analysis using standard techniques with chemiluminescent detection. Magic Mark Western Standard (Invitrogen) was used to determine molecular weight. MRP-8 (Calgranulin A C-1 g, Santa Cruz, SC-8112) was used in a 1:100 dilution in 0.5 $\times$ Uniblock (AspenBio, Inc) for primary antibody. The secondary antibody was Peroxidase anti-goat IgG (H+ L), affinity purified (Vector, PI-9500) in a 1:2000 dilution in Uniblock. MR-14 (Calgranulin B C-19, Santa Cruz, SC-8114) was used in a 1:100 dilution in 0.5 $\times$ Uniblock for primary antibody. The secondary antibody was Peroxidase anti-goat IgG (H+L), affinity purified (Vector, PI-9500) in a 1:2000 dilution in Uniblock.

[0070] Serum MRP-8/14 Determinations. Serum levels of MRP-8/14 were determined by ELISA using a commercially available ELISA (Buhmann S100-Cellion S100 A8/A9) according to the manufacturer's protocol.

[0071] Results

[0072] Identification of Proteins Present in Appendix Tissue from Appendicitis Patients. A differential proteomic analysis was performed on depleted serum samples with the goal of identifying proteins elevated in patients with acute appendicitis. The analysis involved comparing samples from normal patients versus patients with perforated appendices. Blood samples were obtained immediately prior to surgery. A normal patient in this study is one that presented with abdominal pain, underwent surgery, and was found to have a normal appendix. Normal and diseased appendix tissue was collected during surgery.

[0073] The proteomic approach was to compare a pool of 4 normal samples with a pool of 4 appendicitis samples using two-dimensional electrophoresis. FIG. 1 shows the 2D profile of proteins analyzed. Comparison between the gels was performed and the most obvious difference is indicated in FIG. 1B as AP-93. Based on the gel in FIG. 1, the molecular weight of AP-93 is approximately 14 kilodaltons. The corresponding gel slice was analyzed by MALDI-TOF and a positive identification was made. The identification

was based upon spectra of two tryptic peptides, NIETIINTFHQYSVK [SEQ ID NO:1] and LGHPDTLN-QGEFKELVR [SEQ ID NO:2]. The peptides correspond to the underlined residues in the following amino acid sequence of MRP-14 (GenBank Accession Number P06702):

[0074] MTCKMSQLERNIETIINTFHQYSVKLGHPDTLNQGEFKELVRKDLQNFLLKE NKNEKVIEHIMEDLDTNADKQLSFEEFIMLMARLTWASHEKMHEGDEGPGHHHKPGL GEGTP [SEQ ID NO:3].

[0075] The MALDI-TOF identification of AP-93 as MRP-14 was confirmed by the matching molecular weights. Based on this data, MRP-14 protein was more highly abundant in the diseased sample pool than in the normal sample pool.

[0076] Presence of MRP-14 and MRP-8 in Diseased Appendix Tissue. In order to confirm the presence of MRP-14 in diseased tissue, an anti-MRP-14 antibody was used in western blotting of tissue extracts from individual normal and diseased appendices. FIG. 2 shows the western blot data from 9 normal and 11 appendicitis samples. A 14 kilodalton band is present in every appendicitis sample. There is no detectable signal in the normal samples. This data confirms the proteomic screen data and shows that the protein is an indicator of diseased appendix tissue.

[0077] Since it is known that MRP-8 exists as a dimer with MRP-14, tissue specimens were also examined for the presence of MRP-8. FIG. 3 shows the western blot data using an anti-MRP-8 antibody on the normal and diseased tissue samples. As expected, MRP-8 is present in all of the diseased appendix samples and not detectable in the normal appendix tissue. These western blot data show that the MRP-8 and MRP-14 proteins are markedly more abundant in appendicitis than in normal appendix tissue.

[0078] Elevated Serum Levels of MRP-8/14 Patients with Acute Appendicitis. The high correlation between appendicitis and the presence of MRP-8/14 in the appendix led us to examine the MRP-8/14 levels in serum of those patients and other patients subsequently added to the study. The sera were collected before surgery, banked and analyzed after the disease status was known. MRP-8/14 levels were measured using a sandwich ELISA specific for the complex.

[0079] Table 1 lists serum MRP-8/14 levels for 39 patients as determined by an ELISA manufactured by Hycult (Netherlands) and available commercially through Cell Sciences, Canton, M A. The amounts are given as fractions compared to an average level for patients in the study without appendicitis. Note that all patients with appendicitis show a fold-increase of MRP-8/14 over average normal levels. The procedure was conducted according to instructions accompanying the ELISA product. The sample numbers do not correspond to the sample numbers shown in FIGS. 2 and 3 as the samples were renumbered.

TABLE 1

Sample Number	Clinical Diagnosis	Pathology	Grading	Fraction of Normal
1	Advanced Appendicitis	Mild Acute Appendicitis	2	2.80428
2	Normal Appy	Normal	1	0.960805
3	Advanced Appendicitis	Transmural Appendicitis	3	5.554904
4	Perforated Appy	Perforated Appy-Necrosis	4	6.53913
5	Early Appy	Mild Acute Appendicitis	2	4.562059

TABLE 1-continued

Sample Number	Clinical Diagnosis	Pathology	Grading	Fraction of Normal
6	Early Appy	Mild-Acute Appendicitis	1	2.881124
7	Horrible perforated	Perforated Appy-Necrosis	4	7.906886
8	Normal Appy	Mild Acute Appendicitis	2	3.971489
9	Early Appy	Transmural Appendicitis	3	3.83328
10	Advanced Appendicitis	Transmural Appendicitis	3	3.566665
11	Appendicitis	Mild Acute Appendicitis	2	3.205335
12	Appendicitis	Transmural Appendicitis	3	5.51224
13	Advanced Appendicitis	Transmural Appendicitis	3	2.92671
14	Advanced Appendicitis	Transmural Appendicitis w Necrosis	4	3.866306
15	PERFORATED	Transmural Appendicitis	3	4.54657
16	Advanced Appendicitis	Perforated Appy	4	7.01877
17	Advanced Appendicitis	Transmural Appendicitis	3	4.25998
18	Appendicitis	Transmural Appendicitis	3	6.90312
19	Normal Appy	Normal	1	0.838679
20	Normal Appy	Normal	1	0.590095
21	Early Appy	Appendicitis with Peri appy changers	3	1.682291
22	Normal Appy	Normal	1	1.128849
23	Advanced appendicitis	Transmural Appendicitis	4	2.338583
24	Normal Appy	Normal	1	2.478035
25	Hot appy			2.807046
26	Perforated	Perforated	4	4.954136
27	Normal	Normal	1	0.918438
28	Hot	Hot	2	4.387589
29	Early	Transmural Appendicitis	3	4.015013
30	Hot	Transmural Appendicitis	3	2.460902
31	Normal	Normal	1	0.594943
32	Hot	Transmural Appendicitis	3	4.211086
33	Perforated	Transmural Appendicitis	4	3.835219
34	Normal	Normal	1	1.968859
35	Perforated	Transmural Appendicitis	4	4.126198
36	Advanced	Transmural Appendicitis	3	2.423726
37	Hot Appy	Transmural Appendicitis	3	4.178647
38	Early	Transmural Appendicitis	3	9.584398
39	Normal	Transmural Appendicitis	2	2.835339

[0080] We have identified a protein complex that is present in the appendix and serum of appendicitis patients. Based on the western blot data, the presence of MRP-8/14 in appendix tissue is highly correlative with disease. Furthermore, levels of MRP-8/14 in serum are predictive of appendicitis. We presume that this increase is due to increased production of these proteins from systemic neutrophil infiltration of the appendix and possibly direct mucosal production of the proteins by the appendix itself. This study demonstrates that MRP-8/14 is a useful clinical marker for acute appendicitis. After our discovery that MRP-8/14 was a molecule differentially associated with appendicitis, our work was confirmed by the finding of Power, C. et al., 2004 and 2005, who reported detection of this molecule in feces of patients having acute appendicitis.

Example 2

Haptoglobin

[0081] Using a proteomic screen of serum and appendix tissue, we determined that haptoglobin is upregulated in patients with acute appendicitis. The alpha subunit of haptoglobin is an especially useful marker in screening for the disease.

[0082] Materials and Methods

[0083] Specimen and serum collection, appendicitis tissue processing, 2D gel analysis of extracted tissue samples, and

western blot analysis of extracted appendix tissue samples were as described above in Example 1, except that for the western blot, affinity-purified anti-human haptoglobin (Rockland, 600401-272) was used at a 1:5000 dilution in 0.5× uniblock for the primary antibody; and the secondary antibody was peroxidase anti-rabbit IgG (h+1), affinity purified (vector, pi-1000) in a 1:5000 dilution in uniblock.

[0084] Results

[0085] Identification of proteins present in appendix tissue from appendicitis patients. A differential proteomic analysis was performed on depleted serum samples with the goal of identifying proteins elevated in patients with acute appendicitis. The analysis involved comparing samples from normal patients versus patients with perforated appendices. Blood samples were obtained immediately prior to surgery. A normal patient in this study is one that presented with abdominal pain, underwent surgery, and was found to have a normal appendix. Normal and diseased appendix tissue was collected during surgery.

[0086] The proteomic approach was to compare a pool of 4 normal samples with a pool of 4 appendicitis samples using two-dimensional electrophoresis. FIG. 5 shows the 2D profile of proteins analyzed from serum depleted of IgG and albumin. Comparison between the gels was performed and the most obvious difference is indicated in FIG. 5B as AP-77. The protein in gel spot AP-77 was digested with trypsin and analyzed by MALDI-TOF. The resulting two

peptides have the following sequences: TEGDGVYTLN-NEKQWINK [SEQ ID NO:4] and AVGD-KLPECEADDGCPKPPEIAHGVEHSVR [SEQ ID NO:5]. The sequences were aligned with the alpha subunit of haptoglobin. The sequence of haptoglobin precursor (GenBank Accession Number NP005134) is shown below with the tryptic fragments underlined.

[SEQ ID NO:6]

MSALGAVIALLLWGQLFAVDSGNDVTDIADDGCPKPPEIAHGVEHSVR
YQCKNYYKLRTGEDGVYTLNDDKKQWINKAVGDKLPECEADDGCPKPPEI
AHGVEHSVRYQCKNYYKLRTGEDGVYTLNNEKQWINKAVGDKLPECEA
VCGKPKNPANPVQRILGGHLDAGSFPWQAKMVSHHNLTTGATLINEQW
LLTTAKNLFNLHSENAKADIAPTLTLYVGKKQLVEIEKVVLHPNYSQV
DIGLIKLRQKQVSVNERVMPICLPSKDYAEVGRVGVYSGWRNANFKFTD
HLKYVMLPVADQDCIRHYEGSTVPEKKTTPKSPVGVQPILNEHTFCAGM
SKYQEDTCYGDAGSAFAVHDLDEEDTWYATGILSFDKSCAVAEYGVYVKV
TSIQDWVQKTIEN.

[0087] FIG. 6 shows the two-dimensional electrophoresis profile comparison between diseased and normal appendix tissue proteins. Two spots, AP-91 and AP-93, were analyzed by MALDI-TOF and positive identifications were determined. AP-91 protein was determined to be identical to AP-77, haptoglobin-alpha.

[0088] Elevated haptoglobin in diseased appendix tissue. In order to confirm the presence of haptoglobin in diseased tissue, an anti-haptoglobin antibody was used in western blotting of tissue extracts from individual normal and diseased appendices. FIG. 7 shows the western blot data from 6 normal and 6 appendicitis samples. Nearly every sample contained some level of the 38 kd beta subunit, however, there appeared to be an elevated level in cases of appendicitis. A >20 kilodalton band is present in every appendicitis sample and absent from all of the normal tissue samples. This data confirms the proteomic screen data and shows that the protein is an indicator of diseased appendix tissue. The alpha subunit has higher specificity than the beta subunit.

Example 3

Method of Identifying Molecules Using Fluid Samples

[0089] In variations of this example, fluid samples can include whole blood, serum, or plasma. The samples are whole blood collected from human patients immediately prior to an appendectomy. The specimens are placed on ice and transported to the lab. The blood is then processed by centrifugation at 3000 rpm for 15 minutes. Plasma is then separated by pouring into another container

[0090] Upon performing an appendectomy, a patient is classified as having appendicitis (AP) or non-appendicitis (NAP). The classification is based on clinical evaluation, pathology, or both as known in the art. For cases of appendicitis, the clinical condition is also characterized as either perforated or non-perforated.

[0091] The samples from AP patients are optionally pooled and divided into aliquots. Optionally, a pooled ali-

quot is treated so as to remove selected components such as antibodies and serum albumin. Similarly, the samples from NAP patients are optionally pooled and divided into aliquots with optional treatment to remove the same selected components. Preferably the AP samples and NAP samples are processed in a similar manner.

[0092] Next, the pooled aliquots of AP and NAP samples are each subjected to two-dimensional gel electrophoresis as known in the art. The results of each sample type are compared with respect to the presence, absence, and relative expression levels of proteins. Preferably, one detects a signal corresponding to a protein derived from an AP sample that is either absent or expressed at relatively lower levels in a NAP sample. Further characterization is performed for such an AP protein.

[0093] The further characterization can include partial amino acid sequencing, mass spectrometry, and other analytical techniques as known in the art. A full length clone of the gene corresponding to the partial amino acid sequence can be isolated and expressed as a recombinant protein. The recombinant protein can be used as an antigen for detection. Alternatively, a partial or complete recombinant protein can be used to induce or otherwise generate a specific antibody reagent, polyclonal or monoclonal. The antibody reagent is used in the detection of antigen in a patient so as to aid in appendicitis diagnosis. A combination of antigenic molecules can be employed in appendicitis diagnosis.

Example 4

Method of Identifying Molecules Using Tissue Samples

[0094] Tissue samples are collected from appendicitis (AP) and non-appendicitis (NAP) patients. Preferably the tissue is the appendix. The AP or NAP tissues samples are optionally pooled so as to generate an AP tissue pool or an NAP tissue pool. The AP and NAP tissue samples are each used as a source for isolation of total RNA and/or mRNA. Upon isolation, the AP-RNA and NAP-RNA are maintained separately and used for preparation of cDNA.

[0095] A subtraction library is created using techniques available in the art. A cDNA library is optionally amplified. The cDNA library is treated so as to remove undesirable constituents such as highly redundant species and species expressed both in diseased and normal samples. Examples of the techniques include those described by Bonaldo et al. (1996) and Deichmann M et al. (2001).

[0096] Upon generation of the subtraction library, one analyzes, isolates, and sequences selected clones corresponding to sequences differentially expressed in the disease condition. Using molecular biology techniques, one selects candidates for recombinant expression of a partial or complete protein. Such a protein is then used as an antigen for detection. Alternatively, a partial or complete recombinant protein can be used to induce or otherwise generate a specific antibody reagent, polyclonal or monoclonal. The antibody reagent is used in the detection of antigen in a patient so as to aid in appendicitis diagnosis. It is envisioned that a combination of antigenic molecules can be employed in appendicitis diagnosis

Example 5

Method of Appendicitis Diagnosis by Evaluation of Plasma Sample Viscosity

[0097] Whole blood is drawn from a suspected appendicitis patient immediately prior to appendectomy. The specimens are placed on ice and transported to the clinical lab. The blood is processed by centrifugation at 3000 rpm for 15 minutes followed by separation of plasma from the sample by pouring into another container.

[0098] During the step of pouring, the samples are evaluated with respect to viscosity. Increased viscosity is indicative of appendicitis. Approximately 80% of samples corresponding to appendicitis cases demonstrate increased viscosity, whereas approximately none to less than 5% of samples corresponding to non-appendicitis cases demonstrate increased viscosity. It is noted that the degree of increased viscosity can correlate with the severity of appendicitis.

[0099] Viscosity measurements can be conducted by visual observation or by using techniques known in the art. For example, a Coulter Harkness capillary viscometer can be used (Harkness J., 1963) or other techniques (Haidekker M A, et al., 2002).

[0100] The presence of increased viscosity in plasma may be used in combination with other diagnostic techniques, for example with one or more of the following: physical examination, complete blood count (CBC) with or without differential, urinalysis (UA), computed tomography (CT), abdominal ultrasonography, and laparoscopy.

[0101] All references throughout this application, for example publications, patents, and patent documents, are incorporated by reference herein in their entireties, as though individually incorporated by reference, to the extent each reference is at not inconsistent with the disclosure in this application.

[0102] Where the terms “comprise”, “comprises”, “comprised”, or “comprising” are used herein, they are to be interpreted as specifying the presence of the stated features, integers, steps, or components referred to, but not to preclude the presence or addition of one or more other feature, integer, step, component, or group thereof.

[0103] The invention has been described with reference to various specific and preferred embodiments and techniques. However, it should be understood that many variations and modifications may be made while remaining within the spirit and scope of the invention. It will be apparent to one of ordinary skill in the art that compositions, methods, devices, device elements, materials, procedures, techniques, and embodiments, and variations respectively thereof, other than those specifically described herein can be applied to the practice of the invention as broadly disclosed herein without resort to undue experimentation. All art-known functional equivalents are intended to be encompassed by this invention. Whenever a range is disclosed, all subranges and individual values are intended to be encompassed. This invention is not to be limited by the embodiments disclosed, including any shown in the drawings or exemplified in the specification, which are given by way of example and not of limitation.

REFERENCES

- [0104] Aadland E, Fagerhol M K. Faecal calprotectin: a marker of inflammation throughout the intestinal tract. *Europ J Gastroenterol Hepatol* 2002; 14:1
- [0105] Ahlquist D A, Gilbert J A. Stool markers for colorectal screening: future considerations. *Dig Dis* 1996; 14(3):132-44.
- [0106] Alic M. Is fecal calprotectin the next standard in inflammatory bowel disease activity tests?. [Letter] *American Journal of Gastroenterology* 1999; 94(11):3370-1.
- [0107] Arnott I D R, Watts D, Ghosh S. Review article: is clinical remission the optimum therapeutic goal in the treatment of Crohn's disease? *Aliment Pharmacol Ther* 2002; 16:857.
- [0108] Arredouani M S, Kasran A, Vanoirbeek J A, Berger F G, Baumann H Ceuppens J L. 2005. Haptoglobin dampens endotoxin-induced inflammatory effects both in vitro and in vivo. *Immunology*. 114(2):263-71.
- [0109] Berger D, Bolke E, Seidelmann M, Beger H G. Time-scale of interleukin-6, myeloid related proteins (MRP), C reactive protein (CRP), and endotoxin plasma levels during the postoperative acute phase reaction. *Shock* 1997; 7(6):422-6.
- [0110] Berntzen H B, Endresen G K, Fagerhol M K, Spiechowicz J, Mowinkel P. Calprotectin (the L1 protein) during surgery in patients with rheumatoid arthritis. *Scand J Clin Lab Invest* 1991; 51(7):643-50.
- [0111] Berntzen H B, Fagerhol M K, Ostensen M, Mowinkel P, Hoyeraal H M. The L1 protein as a new indicator of inflammatory activity in patients with juvenile rheumatoid arthritis. *J Rheumatol* 1991; 18(1):133-8.
- [0112] Berntzen H B, Munthe E, Fagerhol M K. A longitudinal study of the leukocyte protein L1 as an indicator of disease activity in patients with rheumatoid arthritis. *J Rheumatol* 1989; 16(11):1416-20.
- [0113] Berntzen H B, Munthe E, Fagerhol M K. The major leukocyte protein L1 as an indicator of inflammatory joint disease. *Scand J Rheumatol* 1988; Supp 76:251-6.
- [0114] Berntzen H B, Olmez U, Fagerhol M K, Munthe E. The leukocyte protein L1 in plasma and synovial fluid from patients with rheumatoid arthritis and osteoarthritis. *Scand J Rheumatol* 1991; 20(2):74-82.
- [0115] Berstad A, Arsiand G, Folvik G. Relationship between intestinal permeability and calprotectin in gut lavage fluid. *Scand J Gastroenterol* 2000; 35(1):64-9.
- [0116] Bjarnason I, Sherwood R. Fecal calprotectin: a significant step in the noninvasive assessment of intestinal inflammation. *J Pediatr Gastroent Nutr* 2001; 33:11.
- [0117] Bjerke K, Halstensen T S, Jahnsen F, Pulford K, Brandtzaeg P. Distribution of macrophages and granulocytes expressing L1 protein (calprotectin) in human Peyer's patches compared with normal ileal lamina propria and mesenteric lymph nodes. *Gut* 1993; 34(10):1357-63.
- [0118] Bogumil T, Rieckmann P, Kubuschok B, Felgenhauer K, Bruck W. Serum levels of macrophage-derived

- protein MRP-8114 are elevated in active multiple sclerosis. *Neuroscience Letters* 1998; 247(2-3):195-7
- [0119] Bonaldo M F et al., 1996. Normalization and subtraction: Two approaches to facilitate Gene discovery. *Genome Res.* 6:791-806.
- [0120] Brandtzaeg P, Dale I, Fagerhol M K. Distribution of a formalin-resistant myelomonocytic antigen (L1) in human tissues. I. Comparison with other leukocyte markers by paired immunofluorescence and immunoenzyme staining. *Am J Clin Pathol* 1987; 87(6):681-99.
- [0121] Brandtzaeg P, Dale I, Fagerhol M K. Distribution of a formalin-resistant myelomonocytic antigen (L1) in human tissues. I. Normal and aberrant occurrence in various epithelia. *Am J Clin Pathol* 1987; 87(6):700-7.
- [0122] Brandtzaeg P, Dale I, Gabrielsen T O. The leukocyte protein L1 (calprotectin): usefulness as an immunohistochemical marker antigen and putative biological function. *Histopathol* 1992; 21(2):191-6.
- [0123] Brun J G, Cuida M, Jacobsen H, Kloster R, Johannesen A C, Hoyeraal H M, Jonsson R. Sjögren's syndrome in inflammatory rheumatic diseases: analysis of the leukocyte protein calprotectin in plasma and saliva. *Scand J Rheumatol* 1994; 23(3):114-8.
- [0124] Brun J G, Haga H J, Boe E, Kallay I, Lekven C, Berntzen H B, Fagerhol M K. Calprotectin in patients with rheumatoid arthritis: relation to clinical and laboratory variables of disease activity. *J Rheumatol* 1992; 19(6):859-62.
- [0125] Brydon W G, Campbell S S, Anderson N A, Wilson R G, Ghosh S. Faecal calprotectin levels and colorectal neoplasia. *Gut* 2001; 48(4):579-80.
- [0126] Bunn S K, Bisset W M, Main M J, Golden B E. Fecal calprotectin as a measure of disease activity in childhood inflammatory bowel disease. *J Pediatr Gastroenterol Nutr* 2001; 32(2):171-7
- [0127] Bunn S K, Bisset W M, Main M J C, Gray E S, Olson S, Golden B E. Fecal calprotectin: Validation as a noninvasive measure of bowel inflammation in childhood inflammatory bowel disease. *J Pediatr Gastroenterol Nutr* 2001; 33:11.
- [0128] Burkhardt K, Radespiel-Troger M, Rupprecht H D, Goppelt-Strube M, Riess R, Renders L, Hauser I A, and U Kunzendorf 2001. An increase in myeloid-related protein serum levels precedes acute renal allograft rejection. *J Am Soc Nephrol* 12:1947-57.
- [0129] Clark B R, Kelly S E, Fleming S. Calgranulin expression and association with the keratinocyte cytoskeleton. *J Pathol* 1990; 160(1):25-30
- [0130] Dale I, Brandtzaeg P, Fagerhol M K, Scott H. Distribution of a new myelomonocytic antigen (L1) in human peripheral blood leukocytes. Immunofluorescence and immunoperoxidase staining features in comparison with lysozyme and lactoferrin. *Am J Clin Pathol* 1985; 84(1):24-34.
- [0131] Dale I, Brandtzaeg P. Expression of the epithelial L1 antigen as an immunohistochemical marker of squamous cell carcinoma of the lung. *Histopathol* 1989; 14(5):493-502.
- [0132] Dale I. Plasma levels of the calcium-binding L1 leukocyte protein: standardization of blood collection and evaluation of reference intervals in healthy controls. *Scand J Clin Lab Invest* 1990; 50(8):837-41.
- [0133] Deichmann, M., Polychronidis, M., Wacker, J., Thome, M. & Naher, H., 2001. The protein phosphatase 2A subunit Bg gene is identified to be differentially expressed in malignant melanomas by subtractive suppression hybridization. *Melanoma Research* 2001(11):1-9.
- [0134] Dobryszczycka W. 1997. Biological functions of haptoglobin—new pieces to an old puzzle. *Eur J Clin Chem Clin Biochem* 35(9):647-54.
- [0135] Eversole L R, Miyasaki K T, Christensen R E. Keratinocyte expression of calprotectin in oral inflammatory mucosal diseases. *J Oral Pathol Med* 1993; 22(7):303-7.
- [0136] Eversole L R, Miyasaki K T, Christensen R E. The distribution of the antimicrobial protein, calprotectin, in normal oral keratinocytes. *Arch Oral Biol* 1992; 37(11):963-8.
- [0137] Fagerhol M K, Andersson K B, Naess-Andresen C F, Brandtzaeg P, Dale I. Calprotectin (The L1 Leukocyte Protein) In: VL Smith & JR Dedman (Eds): *Stimulus Response Coupling. The Role of Intracellular Calcium-Binding Proteins*, CRC Press, Boca Raton, Fla., USA, 1990, pp. 187-210.
- [0138] Fagerhol M K. Calprotectin, a faecal marker of organic gastrointestinal abnormality. *Lancet* 2000; 356(9244):17834.
- [0139] Flum D R et al., 2001. Has misdiagnosis of appendicitis decreased over time? A population-based analysis. *JAMA* 286 (14):1748-1753.
- [0140] Fosse E, Moen O., Johnson E, Semb G, Brockmeier V, Mollnes T E, Fagerhol M K, Venge P. Reduced complement and granulocyte activation with heparin-coated cardiopulmonary bypass. *Annals of Thoracic Surgery* 1994; 58(2):472-7.
- [0141] Frosch M, Strey A, Vogl T, Wulffraat N M, Kuis W, Sunderkotter C, Harms E, Sorg C, and J Roth 2000. Myeloid-related proteins 8 and 14 are specifically secreted during interaction of phagocytes and activated endothelium and are useful markers for monitoring disease activity in pauciarticular-onset juvenile rheumatoid arthritis. *Arthritis Rheum* 43:628-37.
- [0142] Gabrielsen T O, Brandtzaeg P, Hoel P S, Dale I. Epithelial distribution of a myelomonocytic antigen L1 in relation to cutaneous malignancies and melanocytic naevi. *Br J Dermatol* 1988; 118(1):59-67.
- [0143] Gabrielsen T O, Dale I, Brandtzaeg P, Hoel P S, Fagerhol M K, Larsen T E, Thune P O. Epidermal and dermal distribution of a myelomonocytic antigen (L1) shared by epithelial cells in various inflammatory skin diseases. *J Am Acad Dermatol* 1986; 15(2 Pt 1):173-9.
- [0144] Garred P, Fosse E, Fagerhol M K, Videm V, Mollnes T E. Calprotectin and complement activation during major operations with or without cardiopulmonary bypass. *Annals of Thoracic Surgery* 1993; 55(3):694-9.

- [0145] Gasché, C., 2005. Laboratory Tests—What Do They Tell Us?, Falk Symposium Abstract, Jun. 17-18, 2005, Munich, Germany.
- [0146] Gaya D R, Mackenzie J F. Faecal calprotectin: a bright future for assessing disease activity in Crohn's disease. *Q J Med* 2002; 95:557 (editorial).
- [0147] Gilbert J A, Ahlquist D A, Mahoney D W, Zinsmeister A R, Rubin J, Ellefson R D. Fecal marker variability in colorectal cancer: calprotectin versus hemoglobin. *Scand J Gastroenterol* 1996; 31(10):1001-5.
- [0148] Golden B E, Clohessy P A, Russell G, Fagerhol M K. Calprotectin as a marker of inflammation in cystic fibrosis. *Archives of Disease in Childhood* 1996; 74(2):136-9.
- [0149] Haga H J, Brun J G, Berntzen H B, Cervera R, Khamashta M, Hughes G R. Calprotectin in patients with systemic lupus erythematosus: relation to clinical and laboratory parameters of disease activity. *Lupus* 1993; 2(1):47-50.
- [0150] Haidekker M A, et al., 2002. A novel approach to blood plasma viscosity measurement using fluorescent molecular rotors. *Am J Physiol Heart Circ Physiol* 282:H1609-H1614.
- [0151] Hammer H B, Kvien T K, Glennas A, Melby K. A longitudinal study of calprotectin as an inflammatory marker in patients with reactive arthritis. *Clin Exp Rheumatol* 1995; 13(1):59-64.
- [0152] Hanai H, Takeuchi K, Iida T, Arai H, Kanaoka K, Iwasaki T, Nakamura A, Hosoda Y, Shirai N, Hirasawa K, Takahira K, Kataoka H, Sano M, Osawa M, Sugimoto S. Clinical significance of faecal calprotectin levels in patients with ulcerative colitis. *Nippon Shokakibyo Gakkaï Zasshi* 2003; 100:21.
- [0153] Harkness J., 1963. A new method for the measurement of plasma viscosity. *Lancet* 2:280-281.
- [0154] Hetland G, Berntzen H B, Fagerhol M K. Levels of calprotectin (leukocyte L1 protein) during apheresis. *Scand J Clin Lab Invest* 1992; 52(6):479-82.
- [0155] Homann C, Garred P, Graudal N, Hasselqvist P, Christiansen M, Fagerhol M K, Thomsen A C. Plasma calprotectin: a new prognostic marker of survival in alcohol-induced cirrhosis. *Hepatology* 1995; 21(4):979-85.
- [0156] Hsieh H L, Schafer B W, Weigle B, and C W Heizmann 2004 S100 protein translocation in response to extracellular S100 is mediated by receptor for advanced glycation endproducts in human endothelial cells. *Biochem Biophys Res Commun* 316:949-59.
- [0157] Hycult Biotechnology b.v., ELISA Test Kit for Human Calprotectin information sheet, Catalog No. HK325.
- [0158] Hycult Biotechnology b.v., Monoclonal Antibody to Human S100A8/A9 (MRP-8/MRP-14), calprotectin Clone 27E10 information sheet, Catalog No. HM2156.
- [0159] Ikemoto, M., et al. 2003. New ELISA System for Myeloid-related Protein Complex (MRP8/14) and its Clinical Significance as a Sensitive Marker for Inflammatory Responses Associated with Transplant Rejection, *Clin. Chem.* 49:594-600.
- [0160] John B, Fagerhol M K, Lyberg T, Prydz H, Brandtzaeg P, Naess-Andresen C F, Dale I. Functional and clinical aspects of the myelomonocyte protein calprotectin. *Molecular Pathology* 1997; 50(3): 113-23.
- [0161] John B, Kronborg O, Ton H, Kristinsson J, Fuglerud P. A new fecal calprotectin test for colorectal neoplasia. Clinical results and comparison with previous method. *Scand J Gastroenterol* 2001; 36(3):291-6.
- [0162] Katnik et al, 1989. Monoclonal Antibodies Against Human Haptoglobin. *Hybridoma* 8(5):551-560.
- [0163] Kelly S E, Hunter J A, Jones D B, Clark B R, Fleming S. Morphological evidence for calcium-dependent association of calgranulin with the epidermal cytoskeleton in inflammatory dermatoses. *Br J Dermatol* 1991; 124(5):403-9.
- [0164] Kelly S E, Jones D B, Fleming S. Calgranulin expression in inflammatory dermatoses. *J Pathol* 1989; 159(1):17-21.
- [0165] Kerkhoff C, Klempt M, Sorg C. Novel insights into structure and function of MRP8 (S100A8) and MRP14 (S100A9). *Biochimica et Biophysica Acta* 1998; 1448(2):200-11.
- [0166] Kjeldsen-Kragh J, Mellbye O J, Haugen M, Mollnes T E, Hammer H B, Sioud M, Forre O. Changes in laboratory variables in rheumatoid arthritis patients during a trial of fasting and one-year vegetarian diet. *Scand J Rheumatol* 1995; 24(2):85-93.
- [0167] Koike T, Kondo K, Makita T, Kajiyama K, Yoshida T, Morikawa M. Intracellular localization of migration inhibitory factor-related protein (MRP) and detection of cell surface MRP binding sites on human leukemia cell lines. *J Biochem* 1998; 123(6):1079-87.
- [0168] Kristinsson J, Armbruster C H, Ugstad M, Kriwanek S, Nygaard K, Ton H, Fuglerud P. Fecal excretion of calprotectin in colorectal cancer; relationship to tumor characteristics. *Scand J Gastroenterol* 2001; 36(2):202-7.
- [0169] Kristinsson J, Roseth A, Fagerhol M K, Aadland E, Schjonsby H, Bormer O P, Raknerud N, Nygaard K. Fecal calprotectin concentration in patients with colorectal carcinoma. *Diseases of the Colon & Rectum* 1998; 41(3):316-21.
- [0170] Kronborg O, Ugstad M, Fuglerud P, John B, Hardcastle J, Scholefield J H, Vellacott K, Moshakis V, Reynolds J R. Faecal calprotectin levels in a high risk population for colorectal neoplasia. *Gut* 2000; 46(6):795-800.
- [0171] Kumar, R. K., et al. 2001. Dimeric S100A8 in human neutrophils is diminished after phagocytosis. *J. Leukoc. Biol.* 70(1):59-64.
- [0172] Limburg P J, Ahlquist D A, Sandborn W J, Mahoney D W, Devens M E, Harrington J J and A R Zinsmeister 2000. Fecal calprotectin levels predict colorectal inflammation among patients with chronic diarrhea referred for colonoscopy. *Am J Gastroenterol*, 10:2831-7.

- [0173] Limburg P J, Ahlquist D A, Sandborn W J, Mahoney D W, Devens M E, Harrington J J, Zinsmeister A R. Faecal calprotectin levels predict colorectal inflammation among patients with chronic diarrhea referred for colonoscopy. *American Journal of Gastroenterology* 2000; 95(10):2831-7.
- [0174] Longbottom D, Sallénave J M, van Heyningen V. Subunit structure of calgranulins A and B obtained from sputum, plasma, granulocytes and cultured epithelial cells. *Biochimica et Biophysica Acta* 1992; 1120(2):215-22.
- [0175] Luger N, Stoll R, Schmid K W, Kucharzik T, Stein H, Burmeister G, Sorg C, Domschke W. The myeloid related protein MRP8/14 (27E10 antigen)-usefulness as a potential marker for disease activity in ulcerative colitis and putative biological function. *Europ J Clin Invest* 1995; 25(9):659-64.
- [0176] Meling T R, Aabakken L, Roseth A, Osnes M. Faecal calprotectin shedding after short-term treatment with non-steroidal anti-inflammatory drugs. *Scandinavian Journal of Gastroenterology* 1996; 31(4):339-44.
- [0177] Moen O, Fosse E, Braten J, Andersson C, Fagerhol M K, Venge P, Hogasen K, Mollnes T E. Roller and centrifugal pumps compared in vitro with regard to haemolysis, granulocyte and complement activation. *Perfusion* 1994; 9(2):109-17.
- [0178] Muller F, Froland S S, Aukrust P, Fagerhol M K. Elevated serum calprotectin levels in HIV-infected patients: the calprotectin response during ZDV treatment is associated with clinical events. *J Acq Immune Defic Syndr* 1994; 7(9):931-9.
- [0179] Muller, F., et al. 1994. Elevated serum calprotectin levels in HIV-infected patients: the calprotectin response during ZDV treatment is associated with clinical events, *J. Acqui. Immune Defic. Syndr.* 7(9):931-939.
- [0180] Neary, Walter, 2001. Press Release from University of Washington, Misdiagnosis of appendicitis continues despite new tools.
- [0181] Olafsdottir E, Aksnes L, Fluge G, Berstad A. Faecal calprotectin in infants with infantile colic, healthy infants, children with inflammatory bowel disease, children with recurrent abdominal pain and healthy children. *Acta Paediatr* 2002; 91:45.
- [0182] Pekna M, Borowiec J, Fagerhol M K, Venge P, Thelin S. Biocompatibility of heparin-coated circuits used in cardiopulmonary bypass. *Scand J Thorac Cardiovasc Surg* 1994; 28(1):5-11.
- [0183] Power, C. et al, 2005. Raised faecal calprotectin levels in patients presenting with right iliac fossa pain warrant mandatory laparoscopy: a non-invasive predictor of acute appendicitis, *Thieme connect, Endoscopy* 37:DOI:10.1055/2-2005-868524.
- [0184] Power, C., et al. 2004, Irish Society of Gastroenterology Winter Meeting Program Oral Presentation Raised Faecal Calprotectin Levels in Patients Presenting with Right Iliac Fossa Pain Warrant Mandatory Laparoscopy: A Non-invasive Predictor of Acute Appendicitis.
- [0185] Robinson M J, Tessier P, Poulsom R, and N. Hogg 2002 The S100 family heterodimer, MRP-8/14, binds with high affinity to heparin and heparan sulfate glycosaminoglycans on endothelial cells. *J Biol. Chem.* 277:3658-65.
- [0186] Roseth A G, Aadland E, Grzyb K. Normalization of faecal calprotectin: a predictor of mucosal healing in patients with inflammatory bowel disease. *Scand J Gastroenterol.* 2004 October;39(10): 1017-20.
- [0187] Roseth A G, Fagerhol M K, Aadland E, Schjonsby H. Assessment of the neutrophil dominating protein calprotectin in feces. A methodologic study. *Scand J Gastroenterol* 1992; 27(9):793-8.
- [0188] Roseth A G, Kristinsson J, Fagerhol M K, Schjonsby H, Aadland E, Nygaard K, Roald B. Faecal calprotectin: a novel test for the diagnosis of colorectal cancer? *Scand J Gastroenterol* 1993; 28(12):1073-6.
- [0189] Roseth A G, Aadland E, Jahnsen J, Raknerud N. Assessment of disease activity in ulcerative colitis by faecal calprotectin, a novel granulocyte marker protein. *Digestion* 1997; 58(2):176-80.
- [0190] Roseth A G, Fagerhol M K, Aadland E, Schjonsby H. Assessment of the neutrophil dominating protein calprotectin in feces. A methodologic study. *Scandinavian Journal of Gastroenterology* 1992; 27(9):793-8.
- [0191] Roseth A G, Schmidt P N, Fagerhol M K. Correlation between faecal excretion of indium-111-labelled granulocytes and calprotectin, a granulocyte marker protein, in patients with inflammatory bowel disease. *Scandinavian Journal of Gastroenterology* 1999; 34(1):50-4.
- [0192] Ryckman C, Vandal K, Rouleau P, Talbot M, and PA Tessier 2003 Proinflammatory activities of S100: proteins S100A8, S100A9, and S100A8/A9 induce neutrophil chemotaxis and adhesion. *J. Immunol.* 170:3233-42.
- [0193] Saintigny G, Schmidt R, Shroot B, Juhlin L, Reichert U, Michel S. Differential expression of calgranulin A and B in various epithelial cell lines and reconstructed epidermis. *J Invest Dermatol* 1992; 99(5):639-44.
- [0194] Sander J, Fagerhol M K, Bakken J S, Dale I. Plasma levels of the leucocyte L1 protein in febrile conditions: relation to aetiology, number of leucocytes in blood, blood sedimentation reaction and C-reactive protein. *Scand J Clin Lab Invest* 1984; 44(4):357-62.
- [0195] Semb A G, Gabrielsen T O, Halstensen T S, Fagerhol M K, Brandtzaeg P, Vaage J. Cardiac surgery and distribution of the leukocyte L1 protein-calprotectin. *Europ J Cardio-Thoracic Surgery* 1991; 5(7):363-7.
- [0196] Shanahan F. Inflammatory bowel disease: immunodiagnostics, immunotherapeutics, and ecotherapeutics. *Gastroenterol* 2001; 120:622.
- [0197] Stockley R A, Dale I, Hill S L, Fagerhol M K. Relationship of neutrophil cytoplasmic protein (L1) to acute and chronic lung disease. *Scand J Clin Lab Invest* 1984; 44(7):629-34.
- [0198] Striz, I. and I. Trebichavsky 2004. Calprotectin—a Pleiotropic Molecule in Acute and Chronic Inflammation. *Physiol Res.* 53:245-253.

- [0199] Thomas P. Rihani H. Roseth A. Sigthorsson G. Price A. Nicholls R J. Bjarnason I. Assessment of ileal pouch inflammation by single-stool calprotectin assay. *Diseases of the Colon & Rectum* 2000; 43(2):214-20.
- [0200] Tibble J, Sigthorsson G, Foster R, Fagerhol M K, Bjarnason I. Faecal calprotectin and faecal occult blood tests in the diagnosis of colorectal carcinoma and adenoma. *Gut* 2001; 49:402.
- [0201] Tibble J. Teahon K. Thjodleifsson B. Roseth A. Sigthorsson G. Bridger S. Foster R. Sherwood R. Fagerhol M. Bjarnason I. A simple method for assessing intestinal inflammation in Crohn's disease. *Gut* 2000; 47(4):506-13.
- [0202] Tibble J A, Bjarnason I. Department of Medicine, Guy's, King's, St Thomas's Medical School, Bessemer Road, London SE5 9PJ, UK. Non-invasive investigation of inflammatory bowel disease.
- [0203] Tibble J A, Bjarnason I. Department of Medicine, Guy's, King's, St. Thomas's Medical School, London, UK. Faecal calprotectin as an index of intestinal inflammation.
- [0204] Tibble J A, Bjarnason I. Markers of intestinal inflammation and predictors of clinical relapse in patients with quiescent IBD. *Medscape Gastroenterol* 2001; 3 (2).
- [0205] Tibble J A. Sigthorsson G. Bridger S. Fagerhol M K. Bjarnason I. Surrogate markers of intestinal inflammation are predictive of relapse in patients with inflammatory bowel disease. [Journal Article] *Gastroenterology* 2000; 119(1):15-22.
- [0206] Tibble J A. Sigthorsson G. Foster R. Scott D. Fagerhol M K. Roseth A. Bjarnason I. High prevalence of NSAID enteropathy as shown by a simple faecal test. *Gut* 1999; 45(3):362-6.
- [0207] Ton H. Brandsnes. Dale S. Holtund J. Skuibina E. Schjonsby H. John B. Improved assay for fecal calprotectin. *Clinica Chimica Acta* 2000; 292(1-2):41-54.
- [0208] Tungekar M F, Heryet A, Gatter K C. The L1 antigen and squamous metaplasia in the bladder. *Histopathol* 1991; 19(3):245-50.
- [0209] U.S. Pat. No. 5,350,687, Odink, et al., Sep. 27, 1994, Antibodies which bind to novel lymphokine related peptides.
- [0210] U.S. Pat. No. 5,455,160, Fagerhol, et al., Diagnostic test and kit for disease disorders in the digestive system.
- [0211] U.S. Pat. No. 5,552,295, Stanker, et al., Sep. 3, 1996, Monoclonal antibodies to bovine haptoglobin and methods for detecting serum haptoglobin levels.
- [0212] U.S. Pat. No. 6,451,550, Eckersall, Sep. 17, 2002, Haptoglobin assay.
- [0213] U.S. patent publication No. 20030224452, Colgin, et al., Pregnancy Detection.
- [0214] Wilkinson M M, Busuttill A, Hayward C, Brock D J, Dorin J R, Van Heyningen V. Expression pattern of two related cystic fibrosis-associated calcium-binding proteins in normal and abnormal tissues. *J Cell Science* 1988; 91 (Pt 2):221-30.
- [0215] Ye B, Cramer D W, Skates S J, Gygi S P, Pratomo V, Fu L, Horick N K, Licklider L J, Schorge J O, Berkowitz R S, Mok S C. 2003 Haptoglobin-alpha subunit as potential serum biomarker in ovarian cancer: identification and characterization using proteomic profiling and mass spectrometry. *Clin Cancer Res* 9(8):2904-11.
- [0216] Yerly S, Bouvier M, Rougemont A, Srivastava I, Perrin L H. 1990. Development of a haptoglobin ELISA. Its use as an indicator for malaria. *Acta Trop.* 1990 47(4):237-44.

 SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 6

<210> SEQ ID NO 1

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Artificial sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic peptide

<400> SEQUENCE: 1

Asn	Ile	Glu	Thr	Ile	Ile	Asn	Thr	Phe	His	Gln	Tyr	Ser	Val	Lys
1				5				10						15

<210> SEQ ID NO 2

<211> LENGTH: 17

<212> TYPE: PRT

<213> ORGANISM: Artificial sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic peptide

<400> SEQUENCE: 2

Leu	Gly	His	Pro	Asp	Thr	Leu	Asn	Gln	Gly	Glu	Phe	Lys	Glu	Leu	Val
1				5				10							15

-continued

Arg

<210> SEQ ID NO 3
<211> LENGTH: 114
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 3

Met Thr Cys Lys Met Ser Gln Leu Glu Arg Asn Ile Glu Thr Ile Ile
1 5 10 15
Asn Thr Phe His Gln Tyr Ser Val Lys Leu Gly His Pro Asp Thr Leu
20 25 30
Asn Gln Gly Glu Phe Lys Glu Leu Val Arg Lys Asp Leu Gln Asn Phe
35 40 45
Leu Lys Lys Glu Asn Lys Asn Glu Lys Val Ile Glu His Ile Met Glu
50 55 60
Asp Leu Asp Thr Asn Ala Asp Lys Gln Leu Ser Phe Glu Glu Phe Ile
65 70 75 80
Met Leu Met Ala Arg Leu Thr Trp Ala Ser His Glu Lys Met His Glu
85 90 95
Gly Asp Glu Gly Pro Gly His His His Lys Pro Gly Leu Gly Glu Gly
100 105 110

Thr Pro

<210> SEQ ID NO 4
<211> LENGTH: 18
<212> TYPE: PRT
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic peptide

<400> SEQUENCE: 4

Thr Glu Gly Asp Gly Val Tyr Thr Leu Asn Asn Glu Lys Gln Trp Ile
1 5 10 15

Asn Lys

<210> SEQ ID NO 5
<211> LENGTH: 31
<212> TYPE: PRT
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic peptide

<400> SEQUENCE: 5

Ala Val Gly Asp Lys Leu Pro Glu Cys Glu Ala Asp Asp Gly Cys Pro
1 5 10 15
Lys Pro Pro Glu Ile Ala His Gly Tyr Val Glu His Ser Val Arg
20 25 30

<210> SEQ ID NO 6
<211> LENGTH: 406
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 6

Met Ser Ala Leu Gly Ala Val Ile Ala Leu Leu Leu Trp Gly Gln Leu
1 5 10 15

Phe	Ala	Val	Asp	Ser	Gly	Asn	Asp	Val	Thr	Asp	Ile	Ala	Asp	Asp	Gly	
			20					25					30			
Cys	Pro	Lys	Pro	Pro	Glu	Ile	Ala	His	Gly	Tyr	Val	Glu	His	Ser	Val	
			35					40					45			
Arg	Tyr	Gln	Cys	Lys	Asn	Tyr	Tyr	Lys	Leu	Arg	Thr	Glu	Gly	Asp	Gly	
			50		55						60					
Val	Tyr	Thr	Leu	Asn	Asp	Lys	Lys	Gln	Trp	Ile	Asn	Lys	Ala	Val	Gly	
65			70						75			80				
Asp	Lys	Leu	Pro	Glu	Cys	Glu	Ala	Asp	Asp	Gly	Cys	Pro	Lys	Pro	Pro	
			85					90						95		
Glu	Ile	Ala	His	Gly	Tyr	Val	Glu	His	Ser	Val	Arg	Tyr	Gln	Cys	Lys	
			100					105						110		
Asn	Tyr	Tyr	Lys	Leu	Arg	Thr	Glu	Gly	Asp	Gly	Val	Tyr	Thr	Leu	Asn	
			115		120						125					
Asn	Glu	Lys	Gln	Trp	Ile	Asn	Lys	Ala	Val	Gly	Asp	Lys	Leu	Pro	Glu	
130			135						140			145				
Cys	Glu	Ala	Val	Cys	Gly	Lys	Pro	Lys	Asn	Pro	Ala	Asn	Pro	Val	Gln	
145			150						155			160				
Arg	Ile	Leu	Gly	Gly	His	Leu	Asp	Ala	Lys	Gly	Ser	Phe	Pro	Trp	Gln	
			165					170						175		
Ala	Lys	Met	Val	Ser	His	His	Asn	Leu	Thr	Thr	Gly	Ala	Thr	Leu	Ile	
			180					185						190		
Asn	Glu	Gln	Trp	Leu	Leu	Thr	Thr	Ala	Lys	Asn	Leu	Phe	Leu	Asn	His	
195						200						205				
Ser	Glu	Asn	Ala	Thr	Ala	Lys	Asp	Ile	Ala	Pro	Thr	Leu	Thr	Leu	Tyr	
210						215						220				
Val	Gly	Lys	Lys	Gln	Leu	Val	Glu	Ile	Glu	Lys	Val	Val	Leu	His	Pro	
225			230						235			240				
Asn	Tyr	Ser	Gln	Val	Asp	Ile	Gly	Leu	Ile	Lys	Leu	Lys	Gln	Lys	Val	
			245					250						255		
Ser	Val	Asn	Glu	Arg	Val	Met	Pro	Ile	Cys	Leu	Pro	Ser	Lys	Asp	Tyr	
			260					265						270		
Ala	Glu	Val	Gly	Arg	Val	Gly	Tyr	Val	Ser	Gly	Trp	Gly	Arg	Asn	Ala	
275						280						285				
Asn	Phe	Lys	Phe	Thr	Asp	His	Leu	Lys	Tyr	Val	Met	Leu	Pro	Val	Ala	
290						295						300				
Asp	Gln	Asp	Gln	Cys	Ile	Arg	His	Tyr	Glu	Gly	Ser	Thr	Val	Pro	Glu	
305			310						315			320				
Lys	Lys	Thr	Pro	Lys	Ser	Pro	Val	Gly	Val	Gln	Pro	Ile	Leu	Asn	Glu	
			325					330						335		
His	Thr	Phe	Cys	Ala	Gly	Met	Ser	Lys	Tyr	Gln	Glu	Asp	Thr	Cys	Tyr	
			340					345						350		
Gly	Asp	Ala	Gly	Ser	Ala	Phe	Ala	Val	His	Asp	Leu	Glu	Glu	Asp	Thr	
355						360						365				
Trp	Tyr	Ala	Thr	Gly	Ile	Leu	Ser	Phe	Asp	Lys	Ser	Cys	Ala	Val	Ala	
370						375						380				
Glu	Tyr	Gly	Val	Tyr	Val	Lys	Val	Thr	Ser	Ile	Gln	Asp	Trp	Val	Gln	
385			390						395			400				
Lys	Thr	Ile	Ala	Glu	Asn											
405																

1. A method for diagnosing appendicitis in a patient comprising:

identifying at least one symptom of appendicitis in said patient; and

identifying the presence of at least one molecule differentially associated with appendicitis in a fluid or tissue sample of said patient;

thereby diagnosing appendicitis in said patient.

2. The method of claim 1 wherein the patient is not known to have an interfering condition associated with the presence of said molecule in a sample of the type in which said molecule is identified.

3. The method of claim 1 wherein said interfering condition is selected from the group consisting of recent allograft; septicemia; meningitis; pneumonia; tuberculosis; rheumatoid arthritis; gastrointestinal cancer; inflammatory bowel disease; skin cancer, periodontitis, preeclampsia, and AIDS.

4. The method of claim 1 wherein said fluid is blood.

5. The method of claim 1 wherein said fluid is serum.

6. The method of claim 1 wherein said molecule is identified in tissue of said patient.

7. The method of claim 1 wherein two or more molecules differentially associated with appendicitis are identified.

8. The method of claim 1 wherein the presence of MRP-8/14 is identified.

9. The method of claim 1 wherein the presence of MRP-8/14 and haptoglobin are identified.

10. The method of claim 1 wherein said symptom of appendicitis is selected from the group consisting of: pain in the abdomen; pain that starts near the navel, then moves to the lower right quadrant of the abdomen; anorexia (loss of appetite); trouble eating accompanied by sleepiness; nausea starting after onset of pain; vomiting starting after onset of pain; vomiting accompanied by fatigue; constipation; small stools with mucus; diarrhea; inability to pass gas; low-grade fever; abdominal swelling; pain in abdomen worsening; tenesmus (feeling of needing to move the bowels); high fever; leukocytosis; and increased plasma viscosity.

11. The method of claim 1 wherein at least two symptoms of appendicitis are identified in said patient.

12. The method of claim 1 wherein the presence of MRP-8/14 or haptoglobin is identified and in addition at least one second molecule is identified selected from the group consisting of: unique structural proteins of the gastrointestinal tract, stress-related inflammatory mediators, immunologic factors, indicators of intestinal bacterial flora, Plasminogen Activator Inhibitor-1, fatty acid binding proteins, nuclear factor kappa beta (NFkB), specific appendix antigens (HLA-DR), inflammation associated antigens, and nucleic acids coding for any of the foregoing.

13. The method of claim 1 also comprising identifying at least one of said molecules by a process including:

obtaining a second fluid or tissue sample from a second patient, wherein the second patient has appendicitis;

obtaining a third fluid or tissue sample from a third patient wherein the third patient has a non-appendicitis condition characterized by at least one symptom of appendicitis;

analyzing said second and third samples so as to detect a molecule differentially associated with the appendicitis in the second patient, whereby said molecule is differentially associated with appendicitis; and

identifying the presence of the molecule, or presence of an amount of the molecule greater than that present in the first sample, thereby diagnosing appendicitis in said patient.

14. The method of claim 1 wherein said sample is a fluid sample and said identification of said molecule is performed in an immunological assay device.

15. The method of claim 14 wherein said sample is blood.

16. The method of claim 1 wherein said identification of said molecule is performed by detecting a nucleic acid sequence coding for said molecule in said sample.

17. A method for identifying a molecule differentially associated with appendicitis, said method comprising:

obtaining a from each of a plurality of patients who are undergoing surgery for suspected appendicitis;

determining during surgery whether each said patient has appendicitis or not; and

analyzing said samples for the presence of a molecule differentially associated with appendicitis.

18. The method of claim 17 wherein said samples are blood samples.

19. The method of claim 17 wherein said samples are appendix tissue samples.

20. The method of claim 19 wherein following identification of said molecule in tissue, it is also identified in plasma.

21. The method of claim 20 wherein samples of blood are also taken from said patients, and the amount of said molecule in plasma of said patients who have appendicitis is determined compared to the amount of said molecule in plasma of patients who do not have appendicitis.

22. The method of claim 17 wherein the molecule differentially associated with appendicitis is selected from the group consisting of unique structural proteins of the gastrointestinal tract, stress-related inflammatory mediators, immunologic factors, indicators of intestinal bacterial flora, Plasminogen Activator Inhibitor-1, fatty acid binding proteins, nuclear factor kappa beta (NFkB), specific appendix antigens (HLA-DR), inflammation associated antigens, and nucleic acids coding for any of the foregoing.

23. An immunoassay test device for detecting the presence of a molecule differentially associated with appendicitis in a sample from a patient suspected of having appendicitis, said device comprising:

a first antibody specific to said molecule;

a support for said first antibody;

means for contacting said first antibody with said sample; and

means for detecting binding of said first antibody with said molecule.

24. The device of claim 23 wherein said first antibody is an antibody to MRP-8/14.

25. The device of claim 23 wherein said first antibody is an antibody to haptoglobin.

26. The immunoassay test device of claim 23 also comprising means for lysing cells in said sample.

27. The immunoassay test device of claim 23 which is a lateral flow device comprising inlet means for flowing a liquid sample into contact with said first antibody.

28. The immunoassay test device of claim 27 also comprising a filter for filtering serum or plasma from a blood sample.

29. The device of claim 23 also comprising a housing.

30. The device of claim 23 in the format of a dipstick or a cartridge.

31. A kit comprising the device of claim 23 and instructions for using said device.

32. A method of detecting the presence of a molecule differentially associated with appendicitis comprising:

a) contacting an immunoassay test device of claim 23 with said sample; and

b) reading the results of said means for detecting binding.

* * * * *

专利名称(译)	用于诊断阑尾炎的方法和装置		
公开(公告)号	US20060024719A1	公开(公告)日	2006-02-02
申请号	US11/189120	申请日	2005-07-25
[标]申请(专利权)人(译)	COLGIN马克 BEALER约翰·F· DONNELLY RICHARD 纽曼DIANE		
申请(专利权)人(译)	COLGIN马克 BEALER约翰·F· DONNELLY RICHARD 纽曼DIANE		
当前申请(专利权)人(译)	COLGIN马克 BEALER约翰·F· DONNELLY RICHARD 纽曼DIANE		
[标]发明人	COLGIN MARK A BEALER JOHN F DONNELLY RICHARD NEWMAN DIANE		
发明人	COLGIN, MARK A. BEALER, JOHN F. DONNELLY, RICHARD NEWMAN, DIANE		
IPC分类号	C12Q1/68 G01N33/53 G01N33/558		
CPC分类号	C12Q1/6883 G01N33/6872 Y10S435/81 G01N2800/06 C12Q2600/158 G01N33/6893		
优先权	60/590631 2004-07-23 US		
其他公开文献	US7501256		
外部链接	Espacenet USPTO		

摘要(译)

提供了一种用于诊断患者的阑尾炎的方法，其包括鉴定患者中的阑尾炎的至少一种症状，并鉴定在所述患者的流体或组织样品中存在与阑尾炎差异相关的至少一种分子。MRP-8/14和触珠蛋白是与阑尾炎差异相关的分子的实例。还提供了用于进行本发明的阑尾炎测定的装置和试剂盒。在一个实施方案中，该装置采用流通免疫测定形式，用于测试血液样品。此外，提供了用于筛选与阑尾炎差异相关的分子的方法，其包括使用来自患者的样品用于疑似阑尾炎。

