



US 20110244489A1

(19) **United States**

(12) **Patent Application Publication**
DEVARAJAN et al.

(10) **Pub. No.: US 2011/0244489 A1**
(43) **Pub. Date: Oct. 6, 2011**

(54) **METHOD FOR THE EARLY DETECTION OF RENAL INJURY**

(76) Inventors: **Prasad DEVARAJAN**, Cincinnati, OH (US); **Jonathan M. Barasch**, New York, NY (US)

(21) Appl. No.: **12/785,220**

(22) Filed: **May 21, 2010**

Related U.S. Application Data

(63) Continuation of application No. 11/096,113, filed on Mar. 31, 2005.

(60) Provisional application No. 60/577,662, filed on Jun. 7, 2004.

Publication Classification

(51) **Int. Cl.**
G01N 33/53 (2006.01)
C07K 16/22 (2006.01)
(52) **U.S. Cl.** **435/7.92; 530/388.1**
(57) **ABSTRACT**

A method and kit for detecting the immediate or early onset of renal disease and injury, including renal tubular cell injury, utilizing NGAL as an immediate or early on-set biomarker in a sample of blood serum. NGAL is a small secreted polypeptide that is protease resistant and consequently readily detected in the blood serum following renal tubule cell injury. NGAL protein expression is detected predominantly in proximal tubule cells, in a punctuate cytoplasmic distribution reminiscent of a secreted protein. The appearance NGAL in the serum is related to the dose and duration of renal ischemia and nephrotoxemia, and is diagnostic of renal tubule cell injury and renal failure. NGAL detection is also a useful marker for monitoring the nephrotoxic side effects of drugs or other therapeutic agents.

Figure 1





Figure 2


Figure 3


Figure 4


Figure 5

Figure 6




Figure 7





Figure 8



Figure 9

Figure 10

METHOD FOR THE EARLY DETECTION OF RENAL INJURY

CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] This application is a continuation application of U.S. application Ser. No. 11/096,113, filed Mar. 31, 2005 (pending), which claims the benefit of U.S. provisional application 60/577,662, filed Jun. 7, 2004, the disclosures of which are incorporated herein by reference.

BACKGROUND OF THE INVENTION

[0002] Acute renal failure (ARF) secondary to a renal tubular cell injury, including an ischemic injury or a nephrotoxic injury remains a common and potentially devastating problem in clinical medicine and nephrology, with a persistently high rate of mortality and morbidity despite significant advances in supportive care. Pioneering studies over several decades have illuminated the roles of persistent vasoconstriction, tubular obstruction, cellular structural and metabolic alterations, and the inflammatory response in the pathogenesis of ARF. While these studies have suggested possible therapeutic approaches in animal models, translational research efforts in humans have yielded disappointing results. The reasons for this may include the multifaceted response of the kidney to ischemic injury and nephrotoxins, and a paucity of early biomarkers for ARF with a resultant delay in initiating therapy.

[0003] An individual is considered to have acute renal failure when the patient's serum creatinine value either (1) increased by at least 0.5 mg/dL when the baseline serum creatinine level was less than 2.0 mg/dL; (2) increased by at least 1.5 mg/dL when the baseline serum creatinine level was greater than or equal to 2.0 mg/dL; or (3) increased by at least 0.5 mg/dL, regardless of the baseline serum creatinine level, as a consequence of exposure to radiographic agents.

[0004] It is believed that introduction of therapy early in the disease process will reduce the mortality rate associated with ARF and shorten the time for treatment of various types of renal tubular cell injuries, including, but not limited to, ischemic and nephrotoxic renal injuries. The identification of a reliable, early biomarker for a renal tubular cell injury would be useful to facilitate early therapeutic intervention, and help guide pharmaceutical development by providing an indicator of nephrotoxicity.

[0005] The traditional laboratory approach for detection of renal disease involved determining the serum creatinine, blood urea nitrogen, creatinine clearance, urinary electrolytes, microscopic examination of the urine sediment, and radiological studies. These indicators are not only insensitive and nonspecific, but also do not allow for early detection of the disease. Indeed, while a rise in serum creatinine is widely considered as the "gold standard" for the detection of ARF, it is now clear that as much as 50% of the kidney function may already be lost by the time the serum creatinine changes.

[0006] A few urinary biomarkers for ischemic renal injury have been earlier described, including kidney injury molecule-1 (KIM-1) and cysteine rich protein 61 (Cyr61). KIM-1 is a putative adhesion molecule involved in renal regeneration. In a rat model of ischemia-reperfusion injury, KIM-1 was found to be upregulated 24-48 hours after the initial insult, rendering it a reliable but somewhat late marker of tubular cell damage. Recent studies have shown that KIM-1

can be detected in the kidney biopsy and urine of patients with ischemic acute tubular necrosis. However, this detection was documented in patients with established ischemic renal damage, late in the course of the illness. The utility of urinary KIM-1 measurement for the detection of early ARF or sub-clinical renal injury has thus far not been validated.

[0007] The protein Cyr61 was found to be a secreted cysteine-rich protein that is detectable in the urine 3-6 hours after ischemic renal injury in animal models. However, this detection required a bioaffinity purification and concentration step with heparin-sepharose beads, followed by a Western blotting protocol. Even after bioaffinity purification several non-specific cross-reacting peptides were apparent. Thus, the detection of Cyr61 in the urine is problematic with respect to specificity as well as the cumbersome nature of the procedure.

[0008] An older name for NGAL is HNL. Prior art U.S. Pat. No. 6,136,526 teaches a method for detecting HNL to distinguish a bacterial infection from a viral infection. Infections cause inflammation in the classical sense of induction of the immune system by attracting neutrophils and other immune cells to the site of infection. When the immune cells infiltrate the affected region, histamines and an array of proinflammatory cytokines are released in the intracellular spaces to induce phagocytosis and killing of the organisms. Activated neutrophils also secrete NGAL in response to bacterial but not viral infections. This differential response is likely to be due to a lipopolysaccharide (LPS) moiety on the surface of bacteria, since NGAL avidly binds LPS. NGAL will then diffuse into capillaries located close to an infected site and, when it reaches a sufficient level, can be detected in serum or plasma. It is not clear how soon neutrophils begin to secrete NGAL in response to bacterial infection, or how long it takes before NGAL released from neutrophils reaches detectable levels in serum.

[0009] Therefore, there remains an urgent need to identify improved biomarkers for immediate and early on-set detection and monitoring of ischemic and nephrotoxic renal injuries.

SUMMARY OF THE INVENTION

[0010] The present invention relates to a method for the immediate or early on-set detection of a renal tubular cell injury in a mammalian subject, comprising the steps of: 1) obtaining a blood serum sample from a mammalian subject; 2) determining from the serum sample the level of a biomarker selected from an immediate renal tubular cell injury biomarker, an early on-set renal tubular cell injury biomarker, and mixtures thereof, and 3) evaluating the renal tubular cell injury status of the subject.

[0011] The present invention also relates to a method for the immediate or early-onset detection of a renal tubular cell injury in a mammal, comprising the steps of: 1) obtaining a blood serum sample from a mammalian subject; 2) contacting the serum sample with an antibody for a renal tubular cell injury biomarker, the renal tubular cell injury biomarker comprising NGAL, to allow formation of a complex of the antibody and the renal tubular cell injury biomarker; and 3) detecting the antibody-biomarker complex.

[0012] The present invention relates to a method for monitoring the effectiveness of a treatment for renal tubular cell injury, comprising the steps of: 1) providing a treatment to a mammalian subject experiencing renal tubular cell injury; 2) obtaining at least one post-treatment serum sample from the subject; 3) determining from the post-treatment serum

sample the level of a biomarker selected from an immediate renal tubular cell injury biomarker, an early on-set renal tubular cell injury biomarker, and mixtures thereof, and 4) evaluating the renal tubular cell injury status of the subject.

[0013] The present invention also relates to a method of monitoring the effectiveness of a treatment for renal tubular cell injury comprising the steps of: 1) providing a treatment to a mammalian subject experiencing renal tubular cell injury; 2) obtaining at least one post-treatment serum sample from the subject; and 3) determining from the post-treatment serum sample the level of a biomarker for renal tubular cell injury selected from an immediate renal tubular cell injury biomarker, an early on-set renal tubular cell injury biomarker, and mixtures thereof.

[0014] The present invention relates to a kit for use in detecting the presence of an immediate or early onset biomarker for renal tubular cell injury, comprising: 1) a means for acquiring a quantity of a blood serum sample; and 2) an assay for the detection in the serum sample of the biomarker.

[0015] The invention further relates to a kit for use in detecting the presence of an immediate or early onset biomarker for renal tubular cell injury in the serum of a subject, comprising: 1) a means for acquiring a quantity of a blood serum sample; 2) a media having affixed thereto a capture antibody capable of complexing with a renal tubular cell injury biomarker selected from an immediate renal tubular cell injury biomarker, an early on-set renal tubular cell injury biomarker, and mixtures thereof; and 3) an assay for the detection of a complex of the renal tubular cell injury biomarker and the capture antibody.

[0016] The invention further relates to a method of identifying the extent of a renal tubular cell injury caused by an event, comprising: 1) obtaining at least one serum sample from a mammalian subject; 2) detecting in the serum sample the presence of a biomarker selected from an immediate renal tubular cell injury biomarker, an early-onset renal tubular cell injury biomarker, and mixtures thereof; and 3) determining the extent of renal tubular cell injury based on the time for on-set of the presence in the serum sample of the biomarker, relative to the time of the event.

[0017] The present invention relates to a method for the detection of a renal tubular cell injury in a mammalian subject, comprising the steps of: 1) obtaining a blood serum sample from a mammalian subject comprising up to 1 milliliter from a mammalian subject following a suspected renal tubular cell injury; 2) determining from the serum sample the level of a biomarker selected from an immediate renal tubular cell injury biomarker, an early on-set renal tubular cell injury biomarker, and mixtures thereof, and (c) evaluating the renal tubular cell injury status of the subject.

[0018] The present invention further relates to a method for the detection of a renal tubular cell injury in a mammalian subject, comprising the steps of: 1) obtaining a blood serum sample comprising up to 1 milliliter from a mammalian subject following a suspected a biomarker for a biomarker selected from an immediate renal tubular cell injury biomarker, an early on-set renal tubular cell injury biomarker, and mixtures thereof, to allow formation of a complex of the antibody and the biomarker; and 3) detecting the antibody-biomarker complex.

[0019] A preferred early on-set renal tubular cell injury biomarker is NGAL. A preferred immediate tubular cell renal injury biomarker is NGAL.

BRIEF DESCRIPTION OF THE DRAWINGS

[0020] FIG. 1 shows Western analysis of urine NGAL in (Left Panel) samples obtained at various times as shown after

cardiopulmonary bypass (CPB) from a subject who subsequently developed ARF, and (Right Panel) recombinant human NGAL standards. Molecular weights in kDa are along the left margin.

[0021] FIG. 2 shows urine NGAL (in ng/ml) at various times after CPB in patients who subsequently developed ARF (upper line, ARF) versus those who did not (lower line, No ARF). The bar represents the time when the initial rise in serum creatinine was detected.

[0022] FIG. 3 shows urine NGAL values of FIG. 2 corrected for urine creatinine excretion.

[0023] FIG. 4 shows urine NGAL (in ng/ml) at various times after CPB in patients who subsequently developed ARF (upper line, ARF) versus those who did not (lower line, No ARF), determined by ELISA. The bar represents the time when the initial rise in serum creatinine was detected.

[0024] FIG. 5 shows urine NGAL values of FIG. 4 corrected for urine creatinine excretion.

[0025] FIG. 6 shows a scatter graph of all urine NGAL measurements at 2 hours post CPB. An arbitrary dashed line at 50 ng/ml illustrates the separation of values in patients who developed ARF versus those with No ARF.

[0026] FIG. 7 shows serum NGAL (ng/ml) at various times after CPB in patients who subsequently developed ARF (upper line, ARF) versus those who did not (lower line, No ARF), determined by ELISA. The bar represents the time when the initial rise in serum creatinine was detected.

[0027] FIG. 8 shows a scatter graph of all serum NGAL measurements at 2 hours post CPB in patients who developed ARF versus those with No ARF.

[0028] FIG. 9 shows receiver operating characteristic (ROC) curves to determine the discriminatory power of NGAL measurements for the early diagnosis of acute renal injury, with an ROC curve for urine NGAL at 2 hours post CPB.

[0029] FIG. 10 shows receiver operating characteristic (ROC) curves to determine the discriminatory power of NGAL measurements for the early diagnosis of acute renal injury, with an ROC curve for serum NGAL at 2 hours post CPB.

DETAILED DESCRIPTION OF THE INVENTION

[0030] As used herein the expression “renal tubular cell injury” shall mean a renal or kidney failure or dysfunction, either sudden (acute) or slowly declining over time (chronic), that can be triggered by a number of disease or disorder processes, including (but not limited to) for renal tubular cell injury; ischemic renal injury (IRI), including acute ischemic injury and chronic ischemic injury; acute renal failure; acute nephrotoxic renal injury (NRI) toxicity, including sepsis (infection), shock, trauma, kidney stones, kidney infection, drug toxicity, poisons or toxins, or after injection with an iodinated contrast dye (adverse effect); and for chronic nephrotoxic renal injury: long-standing hypertension, diabetes, congestive heart failure, lupus, or sickle cell anemia. Both forms of renal failure can result in a life-threatening metabolic derangement.

[0031] As used herein the expression “immediate” in relation to a renal tubular cell biomarker is a biomarker protein that can appear in the blood serum within 2 hours of the onset of renal tubular cell injury.

[0032] As used herein the expression “early on-set” in relation to a renal tubular cell biomarker is a biomarker protein

that can appear in the blood serum within the first 24 hours, more typically within the first 6 hours, of the onset of renal tubular cell injury.

[0033] The present invention provides a method and kit for assaying the presence of a renal tubular cell injury biomarker (which will also be referred to as RTCI biomarker) present in the blood serum of a subject immediately after or at the early onset of renal tubular cell injury. Early detection of the onset of the injury can reduce the time for treatment of the injury, and can reduce the risk of developing clinical acute renal failure (ARF).

[0034] A simple point-of-care kit that uses principles similar to the widely-used blood glucose testing kits, for the rapid detection of serum NGAL at the bedside will allow the clinician to rapidly diagnose renal tubular cell injury (which will be referred to as RTCI), and to rapidly institute proven and effective therapeutic and preventive measures. The use of the kit can represent the standard of care for all patients who are at risk of developing RTCI, especially acute renal failure (or ARF), including use in cardiac surgery, kidney transplantation, stroke, trauma, sepsis, dehydration, and nephrotoxins (antibiotics, anti-inflammatory agents, radio-contrast agents, and chemotherapeutic agents). In current clinical practice, when ARF occurs in the setting of these predisposing conditions, the diagnosis is very delayed, and the associated mortality and morbidity unacceptably high. Ironically, even tragically, effective preventive and therapeutic measures are widely available, but almost never administered in a timely manner due to the lack of early biomarkers of RTCI. It is anticipated that multiple serial measurements of NGAL will be become indispensable not only for diagnosing and quantifying the initial kidney injury, but also for following the response to early treatment, and for predicting long term outcome.

[0035] The biomarker for RTCI can be an immediate RTCI biomarker, such as NGAL, which can appear in the blood serum within 2 hours of the onset of renal tubular cell injury. An immediate RTCI biomarker can, as in the case of NGAL, be present in the blood serum of a subject almost immediately after the onset of renal tubular cell injury. The RTCI biomarker can also be an early-onset RTCI biomarker that can appear within the first 24 hours, more typically within the first 6 hours, of the onset of renal tubular cell injury. As such, NGAL is also an example of an early-onset RTCI biomarker.

[0036] An effective RTCI biomarker is typically a secreted protein, whereby it can be excreted by the kidney into the urine or transported within the blood serum. An effective RTCI biomarker is also typically a protease-resistant protein, such as NGAL. Nevertheless, an RTCI biomarker can also be a protease-sensitive protein, so long as stable fragments of the protein can be detected in the urine or in the blood serum, such as by antibodies as described hereinafter for NGAL.

[0037] The RTCI biomarker can be an ischemic renal injury biomarker (IRI biomarker), a nephrotoxic renal injury biomarker (NRI biomarker), or a mixture thereof. NGAL is an example of both an IRI biomarker and an NRI biomarker.

[0038] The method of the invention can be used to detect the onset of renal tubular cell injury, and to monitor the treatment thereof, for a wide variety of events that can include all varieties of diminished blood supply to the kidneys, impaired heart function, surgical procedures, patients in intensive care units, and the administration of pharmaceuticals, radiocontrast dyes, or other medicament substances to a subject. The renal tubular cell injury can be an ischemic renal

injury, a nephrotoxic renal injury, or other injury that affects the tubular cells of the kidney. The event can include administration or ingestion of a large and wide variety of nephrotoxins, including, but not limited to cancer chemotherapy (cisplatin, cyclophosphamide, ifosfamide, methotrexate), antibiotics (gentamicin, vancomycin, tobramycin), antifungal agents (amphotericin), anti-inflammatory agents (NSAIDs), immunosuppressants (cyclosporins, tacrolimus), and radiocontrast agents. The method can be used to evaluate the nephrotoxicity of both newly-developed and well-known compounds.

[0039] The invention also provides a method and a kit for assessing the extent of renal injury based on a proportional relationship between the extent of injury, which can range from the very onset of renal tubular cell injury, to clinical ARF, with the quantity of NGAL present in the blood serum of the subject. The invention provides a means for a clinician to estimate the degree of renal injury at an initial assessment, and to monitor the change in status of the injury (worsening, improving, or remaining the same) based on the detected amount of NGAL in the blood serum.

[0040] Typically, the clinician would establish a protocol of collecting and analyzing a quantity of fresh blood samples from the patient at selected intervals. Typically the blood sample is obtained intermittently during a prescribed period. The period of time between intermittent sampling can be dictated by the condition of the subject, and can range from a sample each 24 hours to a sample taken continuously, more typically from each 4 hours to each 30 minutes. A serum sample is then typically isolated from the blood sample by well known means.

[0041] Using the methods and techniques described herein, the presence of the RTCI biomarker can be determined, and both a qualitative level of the RTCI biomarker present in the serum can be analyzed and estimated, and a quantitative level of RTCI biomarker present in the serum can be analyzed and measured. The clinician would select the qualitative method, the quantitative method, or both, depending upon the status of the patient. Typically, the quantity of blood serum to be collected is less than 1 milliliter, and more typically less than 10 μ l. A typical sample can range from about 1 μ l to about 1 ml. Typically the larger quantities of a blood serum sample (about 1 ml) are used for quantitative assays. Typically, these small amounts of serum are easily and readily available from clinical subjects who are either prone to developing ARF, or have developed ARF.

[0042] Once an indication of renal tubular cell injury or acute renal failure has been detected, and intervention and treatment of the disease or condition has commenced, the clinician can employ the method and kit of the invention to monitor the progress of the treatment or intervention. If a treatment or surgery that might cause renal tubular cell injury is planned, the clinician can obtain a pretreatment serum sample to determine a baseline value for an individual. Typically, one or more subsequent post-treatment serum samples will be taken and analyzed for the presence of the RTCI biomarker as the treatment of the renal injury commences and continues. If a baseline value was obtained, these post-treatment values can be compared to the baseline value to determine the relative condition of the patient. Detection of the immediate or early on-set biomarkers better relates the injury status of the subject, and can improve the responsiveness and the quality of the treatment options. The treatment is continued until the presence of the RTCI biomarker in subsequent

post-treatment serum samples is not detected. As the treatment and intervention ameliorate the condition, the expression of RTCI biomarker, and its presence in the serum, will be correspondingly reduced. The degree of amelioration will be expressed by a correspondingly reduced level of RTCI biomarker, such as NGAL, detected in a sample. As the renal injury nears complete healing, the method can be used to detect the complete absence of the RTCI biomarker, signaling the completion of the course of treatment. Studies with animal models of ischemic or nephrotoxic injury event demonstrated that NGAL is produced in renal tubular cells within minutes following the event. As shown in the examples of the present invention, the NGAL expressed by renal tubular cells rapidly accumulates in the blood, and can be detected far earlier than current diagnostic tests available to indicate renal cell damage. Since NGAL is easily detected in the serum within 2 hours of the renal injury or nephrotoxic treatment, the invention is suitable for use as an early-onset diagnostic. NGAL testing of serum samples from a subject can begin within 30 minutes of a suspected injury, since NGAL begins to appear in the serum at low levels, and continues to rise thereafter. Therefore, it is also of great value to initiate testing at any time within 2 hours of a suspected injury, when NGAL is clearly apparent in serum. Furthermore, it is of value to test at any other time during the first 24 hours following a suspected injury, since NGAL is a highly reliable and easily measured marker of injury that appears in the serum before changes in other parameters, such as creatinine, can be detected. The most highly preferred course of NGAL testing is to collect samples at intervals throughout the course of treatment to monitor real time changes in renal health status.

[0043] Both monoclonal and polyclonal antibodies that bind an RTCI biomarker are useful in the methods and kits of the present invention. The antibodies can be prepared by methods known in the art. Monoclonal antibodies for a preferred RTCI biomarker, NGAL, are described, for example, in "Characterization of two ELISAs for NGAL, a newly described lipocalin in human neutrophils", Lars Kjeldsen et al., (1996) *Journal of Immunological Methods*, Vol. 198, 155-16, herein incorporated by reference. Examples of monoclonal antibodies for NGAL can be obtained from the Antibody Shop, Copenhagen, Denmark, as HYB-211-01, HYB-211-02, and NYB-211-05. Typically, HYB-211-01 and HYB-211-02 can be used with NGAL in both its reduced and unreduced forms. An example of a polyclonal antibody for NGAL is described in "An Iron Delivery Pathway Mediated by a Lipocalin", Jun Yang et al., *Molecular Cell*, (2002), Vol. 10, 1045-1056, herein incorporated by reference. To prepare this polyclonal antibody, rabbits were immunized with recombinant gel-filtered NGAL protein. Sera were incubated with GST-Sepharose 4B beads to remove contaminants, yielding the polyclonal antibodies in serum, as described by the applicants in Jun Yang et al. *Molecular Cell* (2002).

[0044] Typically, the step of detecting the complex of the capture antibody and the RTCI biomarker comprises contacting the complex with a second antibody for detecting the biomarker.

[0045] The method for detecting the complex of the RTCI biomarker and the primary antibody comprises the steps of separating any unbound material of the serum sample from the capture antibody-biomarker complex; contacting the capture antibody-biomarker complex with a second antibody for detecting the RTCI biomarker, to allow formation of a complex between the RTCI biomarker and the second antibody;

separating any unbound second antibody from the RTCI biomarker-second antibody complex; and detecting the second antibody of the RTCI biomarker-second antibody complex.

[0046] A kit for use in the methods of the present invention typically comprises a media having affixed thereto the capture antibody, whereby the serum sample is contacted with the media to expose the capture antibody to NGAL contained in the sample. The kit includes an acquiring means that can comprise an implement, such as a spatula or a simple stick, having a surface comprising the media. The acquiring means can also comprise a container for accepting the serum sample, where the container has a serum-contacting surface that comprises the media. In an another typical embodiment, the assay for detecting the complex of the RTCI biomarker and the antibody can comprise an ELISA, and can be used to quantitate the amount of NGAL in a serum sample. In an alternative embodiment, the acquiring means can comprise an implement comprising a cassette containing the media.

[0047] Early detection of the RTCI biomarker can provide an indication of the presence of the protein in a serum sample in a short period of time. Generally, a method and a kit of the present invention can detect the RTCI biomarker in a sample of serum within four hours, more typically within two hours, and most typically within one hour, following renal tubular cell injury. Preferably, the RTCI biomarker can be detected within about 30 minutes following renal tubular cell injury.

[0048] A method and kit of the present invention for detecting the RTCI biomarker can be made by adapting the methods and kits known in the art for the rapid detection of other proteins and ligands in a biological sample. Examples of methods and kits that can be adapted to the present invention are described in U.S. Pat. No. 5,656,503, issued to May et al. on Aug. 12, 1997, U.S. Pat. No. 6,500,627, issued to O'Conner et al. on Dec. 31, 2002, U.S. Pat. No. 4,870,007, issued to Smith-Lewis on Sep. 26, 1989, U.S. Pat. No. 5,273,743, issued to Ahlem et al. on Dec. 28, 1993, and U.S. Pat. No. 4,632,901, issued to Valkers et al. on Dec. 30, 1986, all such references being hereby incorporated by reference.

[0049] A rapid one-step method of detecting the RTCI biomarker can reduce the time for detecting the renal tubular cell injury. A typical method can comprise the steps of: obtaining a blood serum sample suspected of containing the RTCI biomarker; mixing a portion of the sample with a detecting antibody which specifically binds to the RTCI biomarker, so as to initiate the binding of the detecting antibody to the RTCI biomarker in the sample; contacting the mixture of sample and detecting antibody with an immobilized capture antibody which specifically binds to the RTCI biomarker, which capture antibody does not cross-react with the detecting antibody, so as to bind the detecting antibody to the RTCI biomarker, and the RTCI biomarker to the capture antibody, to form a detectable complex; removing unbound detecting antibody and any unbound sample from the complex; and detecting the detecting antibody of the complex. The detectable antibody can be labeled with a detectable marker, such as a radioactive label, enzyme, biological dye, magnetic bead, or biotin, as is well known in the art. The detectable antibody can be attached to a supporting material, such as a membrane, plastic strip, plastic laboratory plate such as those used for ELISA or other high-throughput assays, or any other supporting material, such as those used in other diagnostic kits well known in the art.

[0050] To identify potential genes and their proteins that may accompany and mark the earliest onset of renal tubular cell injuries, such as ischemic and nephrotoxic renal injuries, a cDNA microarray assay can be used to detect which of a large number of potential gene targets are markedly upregulated. Utilizing this screening technique, neutrophil gelatinase-associated lipocalin (NGAL) was identified as a gene whose expression is upregulated more than 10 fold within the first few hours following an ischemic renal injury in a mouse model.

[0051] NGAL belongs to the lipocalin superfamily of over 20 structurally related secreted proteins that are thought to transport a variety of ligands within a β -barreled calyx. Human NGAL was originally identified as a 25 kDa protein covalently bound to gelatinase from human neutrophils, where it represents one of the neutrophil secondary granule proteins. Molecular cloning studies have revealed human NGAL to be similar to the mouse 24p3 gene first identified in primary cultures of mouse kidneys that were induced to proliferate. NGAL is expressed at very low levels in other human tissues, including kidney, trachea, lungs, stomach, and colon. NGAL expression is markedly induced in stimulated epithelia. For example, it is upregulated in colonic epithelial cells in areas of inflammation or neoplasia, but is absent from intervening uninvolved areas or within metastatic lesions. NGAL concentrations are elevated in the serum of patients with acute bacterial infections, the sputum of subjects with asthma or chronic obstructive pulmonary disease, and the bronchial fluid from the emphysematous lung. In all these cases, NGAL induction is postulated to be the result of interactions between inflammatory cells and the epithelial lining, with upregulation of NGAL expression being evident in both neutrophils and the epithelium.

[0052] It is believed that the detected NGAL induction represents a novel intrinsic response of the kidney proximal tubule cells to renal tubular cell injuries, including both ischemic and nephrotoxic injuries, and is not derived merely from activated neutrophils. First, the response is rapid, with NGAL appearing in the serum within 2 hours of the injury following renal artery occlusion, while renal neutrophil accumulation in this model of ischemic ARF is usually first noted at 4 hours after injury. Second, the temporal patterns of NGAL induction and neutrophil accumulation are divergent. NGAL mRNA and protein expression was maximally noted at 12 hours of reflow, whereas neutrophil accumulation peaks at 24 hours by which time NGAL expression has significantly declined. Third, no NGAL-expressing neutrophils were detectable by immunofluorescence in the kidney samples examined. Fourth, NGAL mRNA and protein induction was documented to occur in cultured human proximal tubule cells following *in vitro* ischemia, with NGAL secreted into the culture medium within 1 hour of ATP depletion, in a system where neutrophils are absolutely absent. Nevertheless, some contribution from infiltrating neutrophils to the observed NGAL upregulation may have occurred. It is possible that upregulation of NGAL in renal tubule cells may be induced by local release of cytokines from neutrophils trapped in the microcirculation early after ischemic injury.

[0053] While the prior art recognizes that NGAL can be used to distinguish a bacterial infection from a viral infection, this is in contrast to the present invention in several respects. First, there is little or no involvement of neutrophils or other immune cells in early ischemic or nephrotoxic injury. Second, ischemic or nephrotoxic injuries induce early and rapid

expression of NGAL in cells of the affected tissues, such as those lining the various nephron segments. Third, the injured cells of the kidney release NGAL directly into the urine, where it appears within minutes of the injury. Fourth, the inflammation that typically occurs 6-12 hours after ischemic or nephrotoxic injury is distinct from that caused by an infection. Cell death induced by ischemic or nephrotoxic injury induces infiltrates primarily comprising macrophages that secrete proinflammatory cytokines to promote phagocytosis of cellular debris in the damaged tissue. Fifth, although some neutrophil accumulation has been shown to occur in animal models of ischemic kidney injury, this starts to occur only about 4 hours after the injury and peaks at about 24 hours after the injury. In contrast, urine NGAL peaks at 2-4 hours after the injury and is significantly diminished by 24 hours (1-3). Thus, the different time courses of urinary NGAL excretion and neutrophil accumulation argue against an inflammatory source for the urine and serum NGAL following ischemic injury. Sixth, although neutrophil accumulation has been shown in animal models, this has never been shown or substantiated in human acute renal failure. Seventh, we have documented NGAL accumulation in cultured kidney tubule cells following ischemic injury *in vitro*, in a system where neutrophils are absolutely absent. See Rabb H and Star R. *Acute Renal Failure*, Molitoris B A and Finn W F (editors), WB Saunders, Philadelphia, 2001, pp 89-100; Chiao et al. *J Clin Invest* 1997; 99:1165-1172; and Rabb H et al. *Am J Physiol* 1996; 271 F408-P413.

[0054] An adequate explanation for the induction of NGAL by stimulated epithelia has been lacking, and whether NGAL is protective or proximate to injury or even an innocent bystander remains unclear. Recent evidence suggests that, at least in a subset of cell types, NGAL may represent a pro-apoptotic molecule. In the mouse pro-B lymphocytic cell line, cytokine withdrawal resulted in a marked induction of NGAL as well as onset of apoptosis. Purified NGAL produced the same pro-apoptotic response as cytokine deprivation, including activation of Bax, suggesting that NGAL is proximate to programmed cell death. NGAL has also been linked to apoptosis in reproductive tissues. Epithelial cells of the involuting mammary gland and uterus express high levels of NGAL, temporally coinciding with a period of maximal apoptosis. It is likely that NGAL regulates a subset of cell populations by inducing apoptosis. Stimulated epithelia may upregulate NGAL in order to induce apoptosis of infiltrating neutrophils, thereby allowing the resident cells to survive the ravages of the inflammatory response. Alternatively, epithelial cells may utilize this mechanism to regulate their own demise. However, it is interesting to note that induction of NGAL following renal ischemia-reperfusion injury occurs predominantly in the proximal tubule cells, and apoptosis under the same circumstances is primarily a distal tubule cell phenomenon.

[0055] Other recent studies have revealed that NGAL enhances the epithelial phenotype. NGAL is expressed by the penetrating rat ureteric bud, and triggers nephrogenesis by stimulating the conversion of mesenchymal cells into kidney epithelia. Another lipocalin, glycodeulin, has been shown to induce an epithelial phenotype when expressed in human breast carcinoma cells. These findings are especially pertinent to the mature kidney, in which one of the well-documented responses to ischemic injury is the remarkable appearance of dedifferentiated epithelial cells lining the proximal tubules. An important aspect of renal regeneration

and repair after ischemic injury involves the reacquisition of the epithelial phenotype, a process that recapitulates several aspects of normal development. This suggests that NGAL may be expressed by the damaged tubule in order to induce re-epithelialization. Support for this notion derives from the recent identification of NGAL as an iron transporting protein that is complementary to transferrin during nephrogenesis. It is well known that the delivery of iron into cells is crucial for cell growth and development, and this is presumably critical to postischemic renal regeneration just as it is during ontogeny. Since NGAL appears to bind and transport iron, it is also likely that NGAL may serve as a sink for iron that is shed from damaged proximal tubule epithelial cells. Because it has been observed that NGAL can be endocytosed by the proximal tubule, the protein could potentially recycle iron into viable cells. This might stimulate growth and development, as well as remove iron, a reactive molecule, from the site of tissue injury, thereby limiting iron-mediated cytotoxicity.

[0056] NGAL is a novel serum biomarker for cisplatin-induced nephrotoxic renal injury that is more sensitive than previously described biomarkers. One example is kidney injury molecule-1 or KIM-1, a putative adhesion molecule involved in renal regeneration. In a rat model of cisplatin nephrotoxicity, KIM-1 was qualitatively detectable 24-48 hours after the initial insult, rendering it a somewhat late marker of tubular cell damage. NGAL is believed to be readily and quantitatively detected within 3 hours following cisplatin administration at doses known to result in renal failure. In addition, urinary and serum NGAL detection precede the appearance of other markers in the urine such as NAG. Appearance of NGAL in the urine and serum also precede the increase in serum creatinine, that is widely used to diagnose nephrotoxic renal failure.

[0057] It is believed that serum NGAL is evident even after mild "sub-clinical" doses of cisplatin, in spite of normal serum creatinine levels. Thus, the invention has important implications for the clinical management of patients on cisplatin therapy. The efficacy of cisplatin is dose dependent, but the occurrence of nephrotoxicity frequently hinders the use of higher doses to maximize its antineoplastic potential. Nephrotoxicity following cisplatin treatment is common and may manifest after a single dose with acute renal failure. Although several therapeutic maneuvers have proven to be efficacious in the treatment of cisplatin-induced nephrotoxicity in animals, successful human experiences have remained largely anecdotal. One reason for this may be the lack of early markers for nephrotoxic acute renal failure, and hence a delay in initiating therapy. In current clinical practice, acute renal injury is typically diagnosed by measuring serum creatinine. However, it is well known that creatinine is an unreliable and delayed indicator during acute changes in kidney function. First, serum creatinine concentrations may not change until about 50% of kidney function has already been lost. Second, serum creatinine does not accurately depict kidney function until a steady state has been reached, which may require several days. Thus, the use of serum creatinine measurements impairs the ability to both detect and quantify renal damage during the early phases of renal injury. However, animal studies have suggested that while nephrotoxic acute renal failure can be prevented and/or treated, there is a narrow "window of opportunity" to accomplish this, and treatment must be instituted very early after the initial insult. The lack of immediate and early on-set biomarkers of renal injury has impaired the ability of clinicians to initiate potentially

effective therapies in a timely manner. The use of NGAL in an assay system would also be of value for testing existing or emerging therapeutic or preventive interventions, and for the early evaluation of the nephrotoxic potential of other pharmaceutical agents. NGAL detection is a non-invasive, early serum biomarker for cisplatin-induced kidney damage. Early detection may enable clinicians to administer timely therapeutic interventions, and to institute maneuvers that prevent progression to overt nephrotoxic renal failure.

[0058] The upregulation and serum transport of NGAL may represent a rapid response of renal tubule cells to a variety of insults, and the detection of NGAL in the serum may represent a widely applicable noninvasive clinical tool for the early diagnosis of tubule cell injury.

[0059] NGAL is a sensitive, noninvasive serum biomarker for renal tubular cell injuries, including renal ischemia and nephrotoxemia. The examination of the expression of NGAL in the serum of patients with acute, mild and early forms of renal tubular cell injury, using the rapid and simple detection methods and kits of the invention, can alert and enable clinicians to institute timely interventional efforts in patients experiencing acute renal failure, and to alert clinicians to institute maneuvers aimed at preventing progression in patients with subtle, subclinical renal tubular cell injuries (such as a nephrotoxins, kidney transplants, vascular surgery, and cardiovascular events) to overt ARF.

[0060] In the United States alone, there are approximately 16,000 kidney transplants performed every year. This number has been steadily increasing every year. About 10,000 of these are cadaveric kidney transplants, and are at risk for ARE. Each of these patients would benefit enormously from serial NGAL measurements, which could represent routine care.

[0061] Ischemic renal injury has also been associated with open heart surgery, due to the brief interruption in blood flow that is inherent in this procedure. The number of open heart surgeries performed annually can be estimated. In any moderately busy adult hospital, approximately 500 such operations are performed every year. Given that there are at least 400 such moderately busy hospitals in the United States alone, one can conservatively estimate that 200,000 open heart surgeries are performed every year. Again, serial NGAL measurements would be invaluable in these patients, and would represent the standard of care.

EXAMPLES OF THE INVENTION

[0062] In the following examples of the invention, 71 children undergoing CPB were studied. Serial urine and blood samples were analyzed by Western blots and ELISA for NGAL expression. The primary outcome variable was acute renal failure, defined as a 50% increase in serum creatinine from baseline. Twenty patients (28%) developed acute renal failure, but the diagnosis using serum creatinine was possible only 1-3 days after CPB. In contrast, urine NGAL rose from a baseline of 1.6 ± 0.3 ng/ml to 147 ± 23 ng/ml at 2 hours after CPB. Serum NGAL increased from a baseline of 3.2 ± 0.5 ng/ml to 61 ± 10 ng/ml at 2 hours after CPB. Univariate analysts showed a significant correlation between acute renal failure and the following: 2 hour urine NGAL, 2 hour serum NGAL, and CPB time. By multivariate analysis, the urine NGAL at 2 hours post CPB emerged as the most powerful independent predictor of acute renal failure. A ROC curve for the 2-hour urine NGAL revealed an area under the curve of 0.998, and a sensitivity of 1.00 and specificity of 0.98 for a cutoff value of 50 ng/ml. Urine and serum NGAL were novel,

sensitive, specific, highly predictive early biomarkers for acute renal failure following cardiac surgery.

[0063] Study Design: The investigation was approved by the Institutional Review Board of the Cincinnati Children's Hospital Medical Center. Written informed consent was obtained from the legal guardian of each patient before enrollment. All patients undergoing cardiopulmonary bypass (CPB) for surgical correction of congenital heart disease between January and November of 2004 were prospectively enrolled. Exclusion criteria included pre-existing renal insufficiency, diabetes mellitus, peripheral vascular disease, and the use of nephrotoxic agents before or during the study period. We therefore studied a homogeneous population of patients with very likely no major confounding variables in whom the only obvious renal insult would be the result of ischemia-reperfusion injury following CPB. To minimize post-operative volume depletion, all patients received at least 80% of their maintenance fluid requirements during the first 24 hours after surgery, and 100% maintenance subsequently. Spot urine and blood samples were collected at baseline and at frequent intervals for five days following CPB. Urine samples were obtained every two hours for the first 12 hours, and then once every 12 hours. Blood samples were collected at 2 hours post CPB, every 12 hours for the first day, and then once daily for five days. When the CPB time exceeded 2 hours, the first post-operative urine and serum samples were obtained at the end of CPB, and this sample was considered as the 2 hour collection. Urine and blood were also obtained from healthy adult volunteers for establishment of normal NGAL values. Samples were centrifuged at 2,000 g for 5 min. and the supernatants stored in aliquots at -80° C. Serum creatinine was measured at baseline, and routinely monitored in these critically ill children at least twice a day in the immediate post-operative period, and at least daily after post-operative day three.

[0064] Statistical Methods: All results are expressed as means \pm SE. The SAS 8.2 statistical software was utilized for the analysis. A two-sample t-test or Mann-Whitney Rank Sum Test was used to compare continuous variables, and the Chi-square test or Fisher's exact test as indicated were used to compare categorical variables. A conventional receiver operating characteristic (ROC) curve was generated for urine NGAL at 2 and 4 hours post CPB and for serum NGAL at 2 hours post CPB. These were utilized to determine the sensitivities and specificities at different cutoff levels for urine and serum NGAL. The area under the curve was calculated to determine the quality of NGAL as a biomarker. An area of 0.5 is no better than expected by chance, whereas a value of 1.0 signifies a perfect biomarker. Univariate and multivariate stepwise multiple logistic regression analyses were performed to assess predictors of acute renal failure. Potential independent predictor variables included age, gender, race, CPB time, previous heart surgery, urine output, urine NGAL at 2 hours post CPB, and serum NGAL at 2 hours post CPB. A p value of <0.05 was considered significant.

[0065] Patient Characteristics: The guardians of 100 patients provided their informed written consents for their children's participation in this study. Twenty nine patients were excluded, all because of nephrotoxin use (ibuprofen, ACE inhibitors, gentamicin, vancomycin) before or soon after the surgery. Thus, 71 patients were included in the study, whose demographic characteristics, diagnoses, and outcome variables are shown in Table 1, below. All subjects started with normal kidney function and essentially undetectable

levels of NGAL in the urine and serum, just like healthy controls. This study design allowed for the determination of the precise timing of NGAL appearance in the urine and serum following CPB. The results indicate that NGAL is not only a powerful immediate early biomarker for acute renal failure, preceding any increase in serum creatinine by 1-3 days, but is also a valid discriminatory marker over the entire duration of the study.

TABLE 1

Characteristic	Patient characteristics and clinical outcomes.	
	*p < 0.05 versus controls.	
Age (years)	4.0 \pm 0.7	2.1 \pm 1.2*
Gender (% males)	62%	65%
Race (% Caucasian)	88%	85%
Previous Heart Surgery	29%	25%
25% CPB Time (minutes)	105 \pm 8.6	179 \pm 13.6*
Change in serum creatinine (%)	7.7 \pm 1.8	99 \pm 9.3*
Diagnosis (n)		
Ventricular septal defect	9	3
Tetralogy of Fallot	3	9
Atrial Septal Defect	7	0
Coarctation of Aorta	5	1
Aortic Stenosis	6	0
Hypoplastic Left Heart	2	3
AV Canal	3	2
Pulmonic Stenosis	3	1
Transposition of the great arteries	4	0
Tricuspid atresia	3	0
Double-outlet right ventricle	2	0
Anomalous left coronary artery	1	0
Cor Triatriatum	0	1
LV Outflow Tract Obstruction	1	0
Mitral Regurgitation	1	0
Aortic Regurgitation	1	0

*p < 0.05 versus controls.

A major strength of this study is the prospective recruitment of a homogeneous cohort of children subjected to renal ischemia-reperfusion injury during surgical correction of congenital cardiac disease. The patients in these examples were devoid of common co-morbid variables such as atherosclerotic disease, diabetes, and nephrotoxin use, all of which can confound and vitiate the identification of early biomarkers for ischemic acute renal injury.

[0066] Clinical Outcomes: The primary outcome, acute renal failure, defined as a 50% or greater increase in serum creatinine from baseline, occurred in 20 out of 71 patients within a three-day period, yielding an incidence rate of 28%. Out of these, 8 patients displayed an increase in serum creatinine in the 24-48 hours post CPB, but in the other 12 patients, the increase was further delayed to the 48-72 hour period post CPB. Thus, the diagnosis of acute renal failure using currently accepted clinical practices could be made only days after the inciting event.

[0067] Based on the primary outcome, subjects were classified as "control" or "acute renal failure". There were no differences between the two groups in gender, race, or urine output. Other variables that were collected included age, CPB time, previous heart surgery, urine output, and urine creatinine. Children who developed acute renal failure tended to be younger and with longer CPB time, as shown above in TABLE 1. Acute renal failure was more common in patients with an underlying diagnosis of hypoplastic left heart, Tetral-

ogy of Fallot, and AV canal, and was less common or absent in patients with atrial septal defect, ventricular septal defect, or valvular heart disease. The primary outcome variable was the development of acute renal failure, defined as a 50% or greater increase in serum creatinine from baseline.

Example 1

[0068] Western Analysis For NGAL Expression And Quantitation: Equal aliquots (30 μ l) of each urine sample were boiled for 10 min in denaturing buffer and subjected to standard Western Blot analysis with an affinity purified goat polyclonal antibody raised against human NGAL (F-19, Santa Cruz Biotechnology). Simultaneous blots were prepared under identical conditions of transfer and exposure with known quantities of recombinant human NGAL, as standards for quantitation of urine NGAL as previously described by Mishra et al. in Am J Nephrol 2004; 24:307-315. The laboratory investigators were blinded to the sample sources and clinical outcomes until the end of the study.

[0069] Urine NGAL Measurements—Western Analysis: NGAL was virtually undetectable in the urine of all patients prior to surgery, and in healthy volunteers (n=10). FIG. 1 shows a Western Blot typical of that for a patient undergoing CPB. NGAL is not detected at 0 hours, or before CPB, but rapidly appears in the urine by 2 hours or less, and remains detectable by Western blot for at least 12 hours.

[0070] ELISA For NGAL Quantitation: A sensitive and reproducible ELISA for NGAL is an example of a method to provide accurate quantitation of the samples and to confirm the data obtained by Western analysis. Indeed, the ELISA results very closely paralleled those obtained by Western analysis, with a difference of less than 20%. The clinical utility of immunoblot-based techniques for the rapid detection of biomarkers for acute renal injury is limited by the time factor and variations in assay conditions. We modified previously published protocols for detection of NGAL derived from neutrophils as described by Kjeldsen et al. in J Immunol Methods 1996; 198:155-164. Briefly, microliter plates were coated overnight at 4° C. with a mouse monoclonal antibody raised against human NGAL (#HYB211-05, Antibody Shop). All subsequent steps were performed at room temperature. Plates were blocked with buffer containing 1% BSA, coated with 100 μ l of samples (urine or serum) or standards (NGAL concentrations ranging from 1-1000 ng/ml), and incubated with a biotinylated monoclonal antibody against human NGAL (#HYB211-01B, Antibody Shop) followed by avidin-conjugated HRP (Dako). TMB substrate (BD Biosciences) was added for color development, which was read after 30 min at 450 nm with a microplate reader (Benchmark Plus, BioRad). All measurements were made in triplicate, and in a blinded fashion.

Example 2

[0071] In patients who never developed acute renal injury, there was a small but statistically significant increase in urinary NGAL at 2 hours or the first available sample post CPB (4.9 \pm 1.5 ng/ml versus 0.9 \pm 0.3 ng/ml at baseline, p<0.05) and 4 hours post CPB (4.9 \pm 1.2 ng/ml, p<0.05 versus baseline). In marked contrast, patients who subsequently developed acute renal failure displayed a dramatic increase in urinary NGAL at all time points examined, as shown in FIG. 2. The pattern of urinary NGAL excretion was characterized by a peak very early after the precipitating event (2-6 hours following CPB),

followed by a lesser but sustained increase over the entire duration of the study. This overall pattern remained unchanged when urinary NGAL concentration was normalized for urinary creatinine excretion (FIG. 3).

Example 3

[0072] Urine NGAL levels were consistently low in healthy volunteers (2.2 \pm 0.5 ng/ml, n=10) and at baseline in all subjects (1.6 \pm 0.3 ng/ml, n=71). In patients who never developed acute renal failure, there was a small but statistically significant increase in urinary NGAL at 2 hours post CPB (5.9 \pm 1.4 ng/ml, p<0.05 versus baseline) and 4 hours post CPB (5.6 \pm 1.2 ng/ml, p<0.05 versus baseline). Patients who subsequently developed acute renal failure displayed a remarkable increase in urinary NGAL at all time points examined, as shown in FIG. 4. Urinary NGAL excretion peaked very early after CPB, followed by a lesser but sustained increase over the entire duration of the study. Urinary NGAL levels were 147 \pm 23 ng/ml at 2 hours or the first available sample, 179 \pm 30 ng/ml at 4 hours, and 150 \pm 30 ng/ml at 6 hours post CPB in the acute renal injury group. This overall pattern remained consistent when urinary NGAL concentration was normalized for urinary creatinine excretion, i.e. 138 \pm 28 ng/mg creatinine at 2 hours, 155 \pm 40 ng/mg at 4 hours, and 123 \pm 35 ng/mg at 6 hours post CPB (FIG. 5). A scatter plot of the first available post-operative urine NGAL measurements revealed that all 20 patients who subsequently developed acute renal failure displayed a level above an arbitrary cutoff value of 50 ng/ml, whereas only 1 out of 51 patients in the control group showed a urinary NGAL value above this arbitrary cutoff (FIG. 6).

Example 4

[0073] Serum NGAL Measurements—ELISA: Serum NGAL is a novel early biomarker of ischemic renal injury, similar to troponins in myocardial ischemia, and detection of serum NGAL is an example of the invention. Serum NGAL levels were consistently low in normal healthy volunteers (2.5 \pm 0.8, n=6) and all study subjects prior to surgery (3.2 \pm 0.5 ng/ml, n=71). Patients who never developed acute renal failure showed a small but statistically significant increase in serum NGAL at 2 hours or the first available sample post CPB (7.0 \pm 1.1 ng/ml, p<0.05 versus baseline) and 12 hours post CPB (5.2 \pm 0.8 ng/ml, p<0.05 versus baseline). Patients who subsequently developed acute renal failure displayed a striking increase in serum NGAL at all time points examined, as shown in FIG. 7. Similar to urine NGAL, the serum NGAL peaked very early after CPB, followed by a lesser but sustained increase over the entire duration of the study. Serum NGAL levels were 61 \pm 10 ng/ml at 2 hours, 54.7 \pm 7.9 ng/ml at 12 hours, and 47.4 \pm 7.9 ng/ml at 24 hours post CPB in the acute renal failure group. A scatter plot of all the earliest serum NGAL measurements (2 hours post CPB) revealed that none of the 51 patients in the control group displayed a level above an arbitrary cutoff value of 50 ng/ml, whereas the majority of patients who developed acute renal failure showed a serum NGAL value above this value (FIG. 8). The ELISA of the invention is an example of point-of-care diagnostic kits for NGAL.

Example 5

[0074] NGAL for Prediction of Acute Renal Failure: A univariate analysis of the data revealed that the following outcomes were not predictive of acute renal injury: age, gen-

der, race, previous surgery, and urine output. There was a significant correlation between acute renal failure (50% or greater in serum creatinine) and the following: urine NGAL at 2 hours or the first available sample post CPB ($R=0.79$, $p<0.001$), serum NGAL at 2 hours or the first available sample post CPB ($R=0.64$, $p<0.001$), and duration of CPB ($R=0.49$, $p<0.001$). However, by multiple stepwise regression analysis, only the urine NGAL at 2 hours post CPB emerged as the most powerful independent predictor of acute renal failure in this cohort ($R=0.76$, $p<0.001$).

[0075] An ROC curve was constructed to determine the discriminatory power of urine and serum NGAL measurements for the early diagnosis of acute renal failure. For urine NGAL, the area under the curve was 0.998 at 2 hours post CPB (FIG. 9), and 1.000 at 4 hours post CPB (not shown). For serum NGAL, the area under the curve was 0.906 at 2 hours post CPB (FIG. 10). These values indicate that both urine and serum NGAL are excellent tests for the early diagnosis of acute renal failure. The derived sensitivities, specificities, and predictive values at different cutoff levels are listed in TABLE 2. For urine NGAL, a cutoff of either 25 or 50 ng/ml yields outstanding sensitivity and specificity at both 2 hours and 4 hours post CPB. For serum NGAL at 2 hours post CPB, sensitivity and specificity are optimal at the 25 ng/ml cutoff.

TABLE 2

NGAL Test Characteristics at Various Cutoff Values.

	Sensitivity	Specificity	Positive Predictive Value	Negative Predictive Value
Cutoffs for 2 hr Urine NGAL (ng/ml)				
25	1.00	0.98	0.95	1.00
50	1.00	0.98	0.95	1.00
80	0.90	1.00	1.00	0.96
100	0.70	1.00	1.00	0.89
Cutoffs for 4 hr Urine NGAL (ng/ml)				
25	1.00	0.96	0.91	1.00
50	0.95	1.00	0.95	0.98
80	0.70	1.00	1.00	0.89
100	0.65	1.00	1.00	0.88
Cutoffs for 2 hr Serum NGAL (ng/ml)				
25	0.70	0.94	0.82	0.89
50	0.50	1.00	1.00	0.84
80	0.20	1.00	1.00	0.76

[0076] NGAL is normally expressed at very low levels in several human tissues, including kidney, trachea, lungs, stomach, and colon (Cowland et al., Genomics 1997; 45:17-23.). NGAL expression is markedly induced in injured epithelia. For example, NGAL concentrations are elevated in the serum of patients with acute bacterial infections, the sputum of subjects with asthma or chronic obstructive pulmonary disease, and the bronchial fluid from the emphysematous lung (Xu et al., Biochim Biophys Acta 2000; 1482:298-307). The invention described herein stemmed from observations in animal models that NGAL is one of the earliest and most robustly induced genes and proteins in the kidney after ischemic injury, and that NGAL is easily detected in the urine soon after ischemia. See Supavekin et al., Kidney Int 2003; 63:1714-1724; Mishra et al., J Am Soc Nephrol 2003;

4:2534-2543; and Devarajan et al., Mol Genet Metab 2003; 80:365-376, in the post-ischemic kidney. NGAL is markedly upregulated in several nephron segments and the protein accumulates predominantly in proximal tubules where it co-localizes with proliferating epithelial cells. These findings suggest that NGAL may be expressed by the damaged tubule in order to induce re-epithelialization. In support of this hypothesis is the recent identification of NGAL as a regulator of epithelial morphogenesis in cultured kidney tubule cells, and as an iron transporting protein during nephrogenesis (Gwira et al., J Biol Chem 2005, and Yang et al. Mol Cell 2002; 10:1045-1056). It is well known that the delivery of iron into cells is crucial for cell growth and development, and this is presumably also critical to renal regeneration following ischemic injury. Indeed, recent findings indicate that exogenously administered NGAL ameliorates ischemic acute renal injury in mice by tilting the balance of tubule cell fate towards survival (Mishra et al., J Am Soc Nephrol 2004; 15:3073-3082). Thus, NGAL has emerged as a center-stage player in the ARF field, not only as a novel biomarker but also as an innovative therapeutic maneuver.

[0077] While urinary diagnostics have several advantages, including the non-invasive nature of sample collection and the relatively few interfering proteins, some disadvantages also exist. These include the difficulty in obtaining urine samples from patients with severe oliguria, the potential changes in urinary biomarker concentration induced by the overall fluid status and diuretic therapy, and the fact that several urinary biomarkers have in the past shown insufficient sensitivity or specificity (Rabb H. Am J Kidney Dis 2003; 42:599-600.). Serum-based diagnostics have revolutionized intensive care medicine. Recent examples include the measurement of troponins for the early diagnosis of and timely interventions in acute myocardial infarction and the prognostic value of B-type natriuretic peptide in patients with acute coronary syndrome (Hamm et al. N Eng J Med 1997; 337: 1648-1653; and De Lemos et al. N Engl J Med 2001; 345: 1014-1021). To our knowledge, NGAL is the only biomarker that has been examined in both serum and urine for the early diagnosis of ischemic renal injury.

[0078] The methods and use of the invention compare favorably with or surpass the usefulness of several other biomarkers for ischemic renal injury, such as those discussed in Hewitt et al., J Am Soc Nephrol 2004; 15:1677-1689; Herget-Rosenthal et al., Kidney Int 2004; 66:1115-1122; and Rabb, Am J Kidney Dis 2003; 42:599-600). The majority of studies reported thus far have been retrospective, have examined biomarkers in the established phase of ARF, and have been restricted to only the urine and to only one method of detection. Several tubular proteins have been measured in the urine, with conflicting and unsatisfactory results (Westhuyzen et al. Nephrol Dial Transplant 2003; 18:543-551; Herget-Rosenthal et al. Clin Chem 2004; 50:552-558; Han et al., Kidney Int 2002; 62:237-244). Kidney injury molecule-1 (KIM-1), a novel kidney-specific adhesion molecule, is detectable by ELISA in the urine of patients with established acute tubular necrosis. Also, the sodium hydrogen exchanger isoform 3 (NHE3) has been shown by Western blots to be increased in the membrane fractions of urine from subjects with established ARF (du Chevron et al., Am J Kidney Dis 2003; 42:497-506). However, the sensitivity and specificity of these biomarkers for the detection of renal injury have not been reported. Of the inflammatory cytokines involved in ARF, elevated levels of urinary IL-6, IL-8 and IL-18 have

been demonstrated in patients with delayed graft function following cadaveric kidney transplants (35, 36). With the exception of NGAL, none of the biomarkers have been examined prospectively for appearance in the urine during the evolution of ischemic ARF. A recent prospective study has demonstrated that an increase in serum cystatin C precedes the increase in serum creatinine in a select patient population at high risk to develop ARF (Herget-Rosenthal et al., *Kidney Int* 2004; 66:1115-1122). However, the ARF in these subjects was multifactorial, due to a combination of ischemic, prerenal, nephrotoxic, and septic etiologies. Furthermore, since cystatin C is primarily a marker of glomerular filtration rate (GFR), it can be inferred that serum cystatin C levels will rise only after the GFR begins to fall. On the other hand, NGAL is rapidly induced in the kidney tubule cells in response to ischemic injury, and its early appearance in the urine and serum is independent of the GFR, but is highly predictive of a fall in GFR that may occur several days later. A small transient increase in urine and serum NGAL in patients who did not develop ARF was consistent with previous observations that cardiopulmonary bypass surgery leads to release of NGAL into the circulation, probably secondary to inflammatory activation of leukocytes initiated by the extracorporeal circuit (Herget-Rosenthal et al., *Kidney Int* 2004; 66:1115-1122).

[0079] While the invention has been described in conjunction with preferred embodiments, one of ordinary skill after reading the foregoing specification will be able to effect various changes, substitutions of equivalents, and alterations to the subject matter set forth herein. Hence, the invention can be practiced in ways other than those specifically described herein. It is therefore intended that the protection herein be limited only by the appended claims and equivalents thereof.

1.-21. (canceled)

22. A method of diagnosing, monitoring or assessing the severity of disease or injury to an organ, tissue or cell type in a mammal by measuring the concentration of at least one individual molecular form of neutrophil gelatinase-associated lipocalin (NGAL) in a sample of bodily fluid from the mammal and comparing said concentration with the range of concentrations of said individual molecular form that occur in individuals not known to be affected by said disease or injury, whereby a deviation of the concentration from said range determines the presence of disease or injury affecting said organ, tissue or cell type.

23. The method of claim **22** in which the individual molecular form of NGAL is free NGAL monomer.

24. The method of claim **22** in which the individual molecular form of NGAL is the NGAL homodimer or a higher oligomer.

25. The method of claim **22** in which the individual molecular form of NGAL is NGAL which has been glycosylated in a particular manner by its cells of origin, that particular manner being distinguishable from NGAL that has been glycosylated in another manner by different cells of origin.

26. A method of diagnosing, monitoring or assessing the severity of disease or injury to an organ, tissue or cell type in a mammal, said disease or injury being characterized by a certain pattern of concentrations of individual molecular forms of NGAL being present in a bodily fluid from said mammal, by measuring the concentrations of two or more individual molecular forms of NGAL in a sample of said bodily fluid and comparing the results with the pattern of said concentrations that occur in said disease or injury, whereby

the presence of a particular pattern of concentrations indicates the presence of the corresponding type of disease or injury.

27. The method of claim **26** in which the concentration ratio is determined between at least one of: a) free NGAL monomer and NGAL homodimer, b) free NGAL monomer and the sum of NGAL homodimer plus higher oligomers of NGAL, and c) molecular forms of NGAL that are free of matrix metalloproteinase 9 (MMP-9) and NGAL/MMP-9 complexes.

28. (canceled)

29. (canceled)

30. A method according to claim **22** of diagnosing, monitoring or assessing the severity of a renal injury, disease or disorder which has led to acute renal failure or which signifies an immediate risk of developing acute renal failure in a mammal, said method comprising the steps of i) determining the concentration of free NGAL monomer in a sample of bodily fluid from the mammal and ii) comparing said concentration value with a cutoff value determined from the range of free NGAL monomer concentrations in other individuals of that mammalian species which show no evidence of renal injury, disease or disorder, so that a value greater than the cutoff value indicates the presence of said renal injury, disease or disorder.

31. A method according to claim **22** of diagnosing, monitoring or determining the risk of developing acute renal failure in a mammal, wherein said method discriminates between an individual which does not have acute renal failure and is not at immediate risk of developing acute renal failure, and an individual which may have acute renal failure or is at risk of developing acute renal failure, said method comprising the steps of i) determining the concentration of free NGAL monomer in a sample of bodily fluid from the mammal, and ii) comparing said concentration with a predetermined cutoff value chosen so that a concentration of free NGAL monomer below the cutoff value categorizes the mammal as not having and not being at immediate risk of developing acute renal failure.

32. The method of claim **30**, comprising the further step of repeating steps i) and ii) of said claims one or more times.

33. The method of claim **30**, comprising the further step of repeating steps i) and ii) of claim **30** within 24 hours, e.g. within 12 hours, such as within 6 hours, e.g. within 3 hours, such as within 1 hour, e.g. within 30 minutes or within 15 minutes.

34. The method of claim **30**, comprising the further step of repeating steps i) and ii) of claim **30** after a treatment of acute renal failure has been initiated or completed.

35. The method of claim **30**, wherein the renal injury or the risk of developing acute renal failure is due to at least one of ischemia and a surgical intervention.

36. The method of claim **30**, wherein the renal injury or the risk of developing acute renal failure is due to at least one of a complication of an inflammatory, infective or neoplastic disease, critical illness of any cause requiring intensive care, the administration of a nephrotoxic agent, external physical or chemical causes and exposure to radiation.

37. (canceled)

38. (canceled)

39. (canceled)

40. (canceled)

41. (canceled)

42. The method of claim 22, wherein the bodily fluid is blood plasma.

43. The method of claim 22, wherein the bodily fluid is blood serum.

44. The method of claim 22, wherein the bodily fluid is urine.

45. The method of claim 22, wherein free monomer NGAL is measured by means of a molecule that binds specifically to free NGAL monomer and not to complexed forms of NGAL of the mammalian species to which the method is applied.

46. The method of claim 22, wherein NGAL homodimer or the sum of NGAL homodimer plus higher oligomers of

NGAL is measured by a combination of binding molecules that are incapable of binding at one and the same time to free NGAL monomer.

47. The method of claim 22, wherein the mammal is a human individual.

48. (canceled)

49. A method of selecting a binding molecule, including a monoclonal antibody, that is capable of binding specifically to NGAL monomer by selecting said molecule from a series of candidate binding molecules according to its ability to bind to a preparation of NGAL monomer and its inability to bind to NGAL homodimer.

* * * * *

专利名称(译)	早期发现肾损伤的方法		
公开(公告)号	US20110244489A1	公开(公告)日	2011-10-06
申请号	US12/785220	申请日	2010-05-21
[标]申请(专利权)人(译)	DEVARAJAN普拉萨德 BARASCH JONATHAN中号		
申请(专利权)人(译)	DEVARAJAN普拉萨德 BARASCH JONATHAN中号		
当前申请(专利权)人(译)	美国国立卫生研究院 (NIH) , 美国DEPT.卫生与人类服务部 (DHHS) , 美国政府的		
[标]发明人	DEVARAJAN PRASAD BARASCH JONATHAN M		
发明人	DEVARAJAN, PRASAD BARASCH, JONATHAN M.		
IPC分类号	G01N33/53 C07K16/22 G01N33/542 G01N33/68		
CPC分类号	G01N33/6893 G01N2333/82 G01N33/573 G01N2800/52 G01N2800/347		
优先权	60/577662 2004-06-07 US		
外部链接	Espacenet USPTO		

摘要(译)

一种用于检测肾病和损伤的立即或早期发作的方法和试剂盒，包括肾小管细胞损伤，利用NGAL作为血清样品中的即时或早期起始生物标志物。NGAL是一种小的分泌多肽，其具有蛋白酶抗性，因此在肾小管细胞损伤后容易在血清中检测到。NGAL蛋白表达主要在近端小管细胞中检测到，呈点状细胞质分布，让人联想到分泌的蛋白质。血清中NGAL的外观与肾缺血和肾毒性的剂量和持续时间有关，并且是肾小管细胞损伤和肾衰竭的诊断。NGAL检测也是监测药物或其他治疗剂的肾毒性副作用的有用标记物。

Figure 1

