



US007410756B2

(12) **United States Patent**  
**King**

(10) **Patent No.:** **US 7,410,756 B2**  
(45) **Date of Patent:** **Aug. 12, 2008**

(54) **METHODS OF MODULATING ANGIOGENESIS**

(75) Inventor: **George L. King**, Dover, MA (US)

(73) Assignee: **Joslin Diabetes Center, Inc.**, Boston, MA (US)

(\*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 602 days.

(21) Appl. No.: **10/464,720**

(22) Filed: **Jun. 17, 2003**

(65) **Prior Publication Data**  
US 2003/0211109 A1 Nov. 13, 2003

**Related U.S. Application Data**

(63) Continuation of application No. 09/907,255, filed on Jul. 17, 2001, now abandoned.

(60) Provisional application No. 60/219,245, filed on Jul. 18, 2000.

(51) **Int. Cl.**  
**C12Q 1/70** (2006.01)  
**G01N 33/53** (2006.01)

(52) **U.S. Cl.** ..... **435/5; 435/7.1**

(58) **Field of Classification Search** ..... None  
See application file for complete search history.

(56) **References Cited**

**U.S. PATENT DOCUMENTS**

5,959,096 A \* 9/1999 Bennett et al. .... 536/24.5

**OTHER PUBLICATIONS**

Fukumoto et al (J. Biol. Chem. May 1997;272(21):13816-13822).\*

Lu et al (Mol Cell Biol. Jun. 1997;17(6):3418-28).\*

Harrington et al., "Endothelial Proliferation, Migration, and . . .", Biochem Biophys Res Commun, vol. 271, No. 2, May 2000;499-508.

Kent et al., "Requirement for Protein Kinase C Activation in Basic . . .", Circ. Res., vol. 77, No. 2, Aug. 1995;231-238.

Koyama et al., "Tranilast inhibits protein kinase C-dependent signaling pathway . . .", Br. J. Pharmacol., vol. 127, No. 2, May 1999;537-45.

Yoshiji et al., "Protein Kinase C Lies on the Signaling Pathway . . .", Cancer Res., vol. 59, No. 17, Sep. 1999;4413-4418.

Gschwendt et al., "Differential Effects of Suramin on Protein . . .", Jan. 1998, FEBS Letters, vol. 421, No. 2;165-168.

Shih et al., "Role of Protein Kinase C Isoforms in Phorbol . . .", May 1999, J.Biol.Chem., vol. 274, No. 22;15407-15414.

U Jones et al Biochem Biophys Res Commun Aug. 1998; 249(1): 118-123.

V Jain N et al (J Biol Chem Aug. 20, 1999; 274(34): 24392-400).

W Carmeliet et al (Nature 2000; 407:249-257).

\* cited by examiner

*Primary Examiner*—Christopher Yaen

(74) *Attorney, Agent, or Firm*—Fish & Richardson P.C.

(57) **ABSTRACT**

Featured are methods of modulating angiogenesis, e.g., by modulating PKC delta. The methods are useful in the treatment of disorders characterized by aberrant and/or increased blood vessel formation and/or increased permeability. The methods are also useful for treatment a disorder characterized by abnormal, inadequate or unstable blood vessel formation or increased permeability. Also featured are diagnostic and screening methods.

**3 Claims, No Drawings**

## METHODS OF MODULATING ANGIOGENESIS

### RELATED APPLICATIONS

This application is a continuation of U.S. application Ser. No. 09/907,255, filed Jul. 17, 2001 now abandoned, which claims the benefit of U.S. Provisional Application Ser. No. 60/219,245, filed on Jul. 18, 2000, the contents of which is incorporated herein by reference in its entirety.

### BACKGROUND OF THE INVENTION

The process of angiogenesis is very complex involving multiple steps. These include proliferation of endothelial cells, degradation of basement membrane, migration of endothelial cells and the formation of tubes or capillaries. Each of these steps is regulated by multiple cytokine and signal transduction pathways. It is believed that abnormality in any of these steps could lead to vascular disease.

### SUMMARY OF THE INVENTION

The present invention is based, in part, on the discovery that PKC delta isoform regulates endothelial cell differentiation into tubes. Thus, PKC delta plays an important role in the final organization and integrity of a capillary, which is an important and final step in angiogenesis. Based on the discovery that PKC delta plays a role in tube formation, it is possible to use a PKC delta agonist to promote tube formation to reduce permeability. In addition, it is possible to prevent or reduce tube formation, thereby inhibiting angiogenesis, by using a PKC delta antagonist.

Accordingly, in one aspect, the invention features a method of modulating angiogenesis. The method includes modulating, e.g., increasing or decreasing, PKC delta expression or activity, to thereby modulate angiogenesis and/or vascular tube formation.

In one aspect, the invention features a method of promoting angiogenesis in a subject. The method includes increasing PKC delta expression and/or activity, to thereby promote angiogenesis.

In a preferred embodiment, PKC delta is increased by administering an agent which increases the level of PKC delta expression and/or activity. An agent which increases the level of PKC delta expression and/or activity can be one or more of: a PKC delta agonist; a PKC delta polypeptide or a functional fragment or analog thereof; a nucleotide sequence encoding a PKC delta polypeptide or functional fragment or analog thereof; an agent which increases PKC delta nucleic acid expression, e.g., a small molecule which binds to the promoter region of PKC delta and/or which stimulates PKC delta activity, e.g., PMA. In a preferred embodiment, PKC delta levels are increased by administering, e.g., introducing, a nucleotide sequence encoding a PKC delta polypeptide or functional fragment or analog thereof, into a particular cell, e.g., an endothelial cell, a kidney cell, etc., in the subject. The nucleotide sequence can be a genome sequence or a cDNA sequence. The nucleotide sequence can include: a PKC delta coding region; a promoter sequence, e.g., a promoter sequence from the PKC delta gene or from another gene; an enhancer sequence; untranslated regulatory sequences, e.g., a 5' untranslated region (UTR), e.g., a 5' UTR from the PKC delta gene or from another gene, a 3' UTR, e.g., a 3' UTR from the PKC delta gene or from another gene; a polyadenylation site; an insulator sequence. In another preferred embodiment, the level of PKC protein is increased by increasing the level of

expression of an endogenous PKC delta gene, e.g., by increasing transcription of the PKC delta gene. In a preferred embodiment, transcription of the PKC delta gene is increased by: altering the regulatory sequence of the endogenous PKC delta gene, e.g., by the addition of a positive regulatory element (such as an enhancer or a DNA-binding site for a transcriptional activator); the deletion of a negative regulatory element (such as a DNA-binding site for a transcriptional repressor) and/or replacement of the endogenous regulatory sequence, or elements therein, with that of another gene, thereby allowing the coding region of the PKC delta gene to be transcribed more efficiently.

In a preferred embodiment, PKC delta expression and/or activity is increased in a subject, e.g., a human or non-human subject, having a disorder characterized by abnormal, inadequate or unstable blood vessel formation, and/or increased permeability, e.g., a leaky blood vessels. Such disorders include, but are not limited to, diabetes, sickle cell anemia, edema and macular degeneration.

In another aspect, the invention features a method of reducing and/or inhibiting angiogenesis in a subject. The method includes decreasing PKC delta expression and/or activity, to thereby reduce or inhibit angiogenesis.

In a preferred embodiment, PKC delta is decreased by administering an agent which decreases the level of PKC delta expression and/or activity. An agent which decreases the level of PKC delta expression or activity can be one or more of: a PKC delta antagonist, e.g., a PKC delta binding protein which binds to PKC delta but does not activate the enzyme; a PKC delta nucleic acid molecule which can bind to a cellular PKC delta nucleic acid sequence, e.g., mRNA, and inhibit expression of the protein, e.g., an antisense molecule or PKC delta ribozyme; an antibody that specifically binds to PKC delta protein, e.g., an antibody that disrupts PKC delta's catalytic activity or an antibody that disrupts the ability of upstream activators to activate PKC delta; an agent which decreases PKC delta gene expression and/or activity, e.g., a small molecule which inhibits PKC delta, e.g., Rottlerin. In another preferred embodiment, PKC delta expression is inhibited by decreasing the level of expression of an endogenous PKC delta gene, e.g., by decreasing transcription of the PKC delta gene. In a preferred embodiment, transcription of the PKC delta gene can be decreased by: altering the regulatory sequences of the endogenous PKC delta gene, e.g., by the addition of a negative regulatory sequence (such as a DNA-binding site for a transcriptional repressor), or by the removal of a positive regulatory sequence (such as an enhancer or a DNA-binding site for a transcriptional activator).

In a preferred embodiment, angiogenesis is inhibited in a subject having a disorder characterized by aberrant and/or increased blood vessel formation and/or increased permeability. For example, angiogenesis can be inhibited to treat a subject having a disease and/or process that is mediated by angiogenesis including, but not limited to, hemangioma, solid tumors, leukemia, metastasis, telangiectasia psoriasis scleroderma, pyogenic granuloma, myocardial angiogenesis, plaque neovascularization, coronary collaterals, cerebral collaterals, arteriovenous malformations, ischemic limb angiogenesis, corneal diseases, rubeosis, neovascular glaucoma, diabetic retinopathy, retrolental fibroplasia, arthritis, diabetic neovascularization, macular degeneration, wound healing, peptic ulcer, fractures, keloids, vasculogenesis, hematopoiesis, ovulation, menstruation, and placentation.

In another aspect, the invention features a method of determining if a subject is at risk for a disorder characterized by abnormal or inadequate blood vessel formation, e.g., to determine if a subject is at risk for a disorder characterized by leaky

blood vessels. The method includes evaluating, e.g., detecting, the presence or absence of a genetic lesion in a PKC delta gene, or evaluating, e.g., detecting, misexpression of the PKC delta gene.

In a preferred embodiment, the method includes evaluating, e.g., detecting, a genetic lesion in a PKC delta gene, or evaluating, e.g., detecting, underexpression of the PKC delta gene, to thereby determine if a subject is at risk abnormal or inadequate blood vessel formation.

In preferred embodiment, the method includes evaluating in a sample of cells from the subject for the presence or absence of a genetic lesion, e.g., a mutation in the gene encoding a PKC delta protein. The presence of a genetic lesion is indicative of a risk.

In another preferred embodiment, the method includes evaluating in a sample of cells for the expression levels of the PKC delta to determine underexpression. Underexpression of PKC delta is indicative of a risk.

In a preferred embodiment, the genetic lesions is evaluated by contacting the sample with a nucleic acid probe capable of hybridizing to PKC delta mRNA, e.g., a labeled probe. In another preferred embodiment, expression of PKC delta is evaluated with an antibody capable of binding to PKC delta protein, e.g., a labeled antibody, or by detecting mRNA levels, e.g., using an amplification assay.

In another aspect, the invention features a method for identifying a compound capable of modulating angiogenesis.

In one embodiment, the method is used for identifying an agent which increases angiogenesis. The method includes: contacting a cell capable of expressing a PKC delta polypeptide with a test compound; and determining the level of PKC delta polypeptide or nucleic acid expression, wherein a compound capable of increasing PKC delta polypeptide or nucleic acid expression is indicative of a compound capable of promoting angiogenesis.

In a preferred embodiment, the compound is a PKC delta fragment or analog.

In a preferred embodiment, the method further includes evaluating a control cell, e.g., an identical cell which is not treated with the compound.

In a preferred embodiment, PKC delta nucleic acid expression is evaluated using a nucleic acid probe, e.g., a labeled probe, capable of hybridizing to a PKC delta nucleic acid molecule, e.g., PKC delta mRNA, and/or by using an amplification assay for PKC delta, e.g., RT-PCR to detect PKC delta mRNA levels. In a preferred embodiment, PKC delta polypeptide expression is evaluated using an anti-PKC delta antibody, e.g., a labeled anti-PKC delta antibody.

In a preferred embodiment, the agent which increases angiogenesis can further be evaluated by administering the agent to an animal model characterized by abnormal, inadequate or unstable blood vessel formation, and/or increased permeability, e.g., a leaky blood vessels

In another aspect, the invention features a method for identifying a compound capable of inhibiting angiogenesis. The method includes: contacting a cell capable of expressing a PKC delta polypeptide with a test compound; and determining the level of PKC delta polypeptide or nucleic acid expression in the presence and absence of the compound, wherein a compound capable of decreasing PKC delta polypeptide or nucleic acid expression is indicative of a compound capable of inhibiting angiogenesis.

In a preferred embodiment, PKC delta nucleic acid expression is evaluated using a nucleic acid probe, e.g., a labeled probe, capable of hybridizing to a PKC delta nucleic acid molecule, e.g., PKC delta mRNA, and/or by using an amplification assay for PKC delta, e.g., RT-PCR of PKC delta

mRNA. In a preferred embodiment, PKC delta polypeptide expression is evaluated using an anti-PKC delta antibody, e.g., a labeled anti-PKC delta antibody.

In a preferred embodiment, the agent which inhibits angiogenesis can further be evaluated by administering the agent to an animal model characterized by aberrant and/or increased blood vessel formation and/or increased permeability.

#### DETAILED DESCRIPTION OF THE INVENTION

In an attempt to understand the roles of PKC isoforms in the various stages of angiogenesis, full-length cDNA of PCK  $\alpha$ ,  $\beta$  $\frac{1}{2}$ ,  $\gamma$ ,  $\delta$ ,  $\epsilon$  and  $\zeta$ , as well as a dominant negative kinase mutant of each, were placed into an adenoviral vector. Using these adenovirus, retinal endothelial cells were infected in culture with an expression of 5 to 20 fold of protein of these PKC isoforms. In addition, it was shown that the wild type can increase specific PKC isoform activity and their dominant negative of the PKC mutant appears to inhibit specific isoform activity acting as a dominant mutant. Using this technology, it was shown that VEGF's effect on tube formation or endothelial cell differentiation can be prevented in endothelial cells culture models only with PKC delta dominant negative isoforms. Overexpression of PKC delta appears to improve the tube formation as stimulated by VEGF, whereas, overexpression of dominant negative form of PKC delta inhibit this effect of VEGF. Thus, the activation of PKC delta isoform is useful to stabilize capillaries and decrease permeability. In addition, inhibiting PKC delta decreases angiogenesis. Thus, diseases which have increased permeability and/or angiogenesis can be involved.

#### Protein Kinase C and Protein Kinase C Delta Isoform

Protein kinase C (PKC) is a membrane-associated enzyme that is regulated by a number of factors, including membrane phospholipids, calcium, and membrane lipids such as diacylglycerols that are liberated in response to the activities of phospholipases (Bell et al. *J. Biol. Chem.* 1991. 266:4661-4664; Nishizuka, *Science* 1992. 258:607-614. The protein kinase C isozymes, alpha, beta ( $\beta$ )-1, beta-2 and gamma, require membrane phospholipid, calcium and diacylglycerol/phorbol esters for full activation. The delta, epsilon, eta, and theta forms of PKC are calcium-independent in their mode of activation. The zeta and lambda forms of PKC are independent of both calcium and diacylglycerol and are believed to require only membrane phospholipid for their activation. PKC- and isozyme-specific (e.g., PKC delta specific) modulators are described, e.g., in Goekjian et al. *Current Medicinal Chemistry*, 1999, 6:877-903; Way et al., *Trends Pharmacol Sci*, 2000, 21:181-7, and in U.S. Pat. No. 5,843,935.

#### Diagnostic Assays

The level of mRNA corresponding to the PKC delta gene in a cell can be determined by in situ or in vitro formats.

The PKC delta mRNA probes can be used in hybridization or amplification assays that include, but are not limited to, Southern or Northern analyses, polymerase chain reaction analyses and probe arrays. One diagnostic method for the detection of mRNA levels involves contacting the mRNA with a nucleic acid molecule (probe) that can hybridize to the mRNA encoded by the gene being detected. The nucleic acid probe can be, for example, a full-length PKC delta nucleic acid or a portion thereof, such as an oligonucleotide of at least 7, 10, 15, 30, 50, 100, 250 or 500 nucleotides in length and sufficient to specifically hybridize under stringent conditions to PKC delta mRNA, cDNA, or portions thereof. As used herein, the term "hybridizes under stringent conditions" describes conditions for hybridization and washing. Stringent

conditions are known to those skilled in the art and can be found in Current Protocols in Molecular Biology, John Wiley & Sons, N.Y. (1989), 6.3.1-6.3.6. The probes can be labeled with a detectable reagent to facilitate identification of the probe. Useful reagents include, but are not limited to, radio-

activity, fluorescent dyes or enzymes capable of catalyzing a detectable product. In one format, mRNA (or cDNA) is immobilized on a surface and contacted with the probes, for example by running the isolated mRNA on an agarose gel and transferring the mRNA from the gel to a membrane, such as nitrocellulose. In an alternative format, the probes are immobilized on a surface and the mRNA (or cDNA) is contacted with the probes, for example, in a two-dimensional gene chip array. A skilled artisan can adapt known mRNA detection methods for use in detecting the level of mRNA encoded by the PKC delta gene.

The level of mRNA in a sample that is encoded by a PKC delta nucleic acid can be evaluated with nucleic acid amplification, e.g., by RT-PCR (U.S. Pat. No. 4,683,202), ligase chain reaction (Barany, Proc. Natl. Acad. Sci. USA 88:189-193, 1991), self sustained sequence replication (Guatelli et al., Proc. Natl. Acad. Sci. USA 87:1874-1878, 1990), transcriptional amplification system (Kwoh et al., Proc. Natl. Acad. Sci. USA 86:1173-1177, 1989), Q-Beta Replicase (Lizardi et al., Bio/Technology 6:1197, 1988), rolling circle replication (U.S. Pat. No. 5,854,033), or any other nucleic acid amplification method, followed by the detection of the amplified molecules using techniques known in the art. As used herein, amplification primers are defined as being a pair of nucleic acid molecules that can anneal to 5' or 3' regions of a gene (plus and minus strands, respectively, or vice-versa) and contain a short region in between. In general, amplification primers are from about 10 to 30 nucleotides in length and flank a region from about 50 to 200 nucleotides in length. Under appropriate conditions and with appropriate reagents, such primers permit the amplification of a nucleic acid molecule including the nucleotide sequence flanked by the primers.

In another embodiment, the methods further include contacting a control sample with a compound or agent capable of detecting PKC delta mRNA, or cDNA, and comparing the presence of PKC delta mRNA or cDNA in the control sample with the presence of PKC delta mRNA or cDNA in the test sample.

A variety of methods can be used to determine the level of protein encoded by the PKC delta gene. In general, these methods include contacting an agent that selectively binds to the protein, such as an antibody, with a sample to evaluate the level of protein in the sample. In one embodiment, the antibody bears a detectable label. Antibodies can be polyclonal or monoclonal. An intact antibody, or a fragment thereof (e.g., Fab or F(ab')<sub>2</sub>) can be used. The term "labeled," with regard to the probe or antibody, is intended to encompass direct labeling of the probe or antibody by coupling (i.e., physically linking) a detectable substance to the probe or antibody, as well as indirect labeling of the probe or antibody by reactivity with a detectable substance.

The detection methods can be used to detect PKC delta protein in a biological sample in vitro. In vitro techniques for detection of PKC delta protein include enzyme linked immunosorbent assays (ELISAs), immunoprecipitations, immunofluorescence, enzyme immunoassays (EIA), radioimmunoassays (RIA), and Western blot analysis.

In another embodiment, the methods further include contacting the control sample with a compound or agent capable of detecting PKC delta protein, and comparing the presence

of PKC delta protein in the control sample with the presence of PKC delta protein in the test sample.

The invention also provides methods for identifying modulators, i.e., candidate or test compounds or agents (e.g., proteins, peptides, peptidomimetics, peptoids, small molecules or other drugs) which have stimulatory or inhibitory effect on, for example, the expression or activity of PKC delta, thereby modulating angiogenesis. Compounds thus identified can be used to modulate the activity of PKC delta, in a method described herein.

#### Generation of Analogs: Production of Altered DNA and Peptide Sequences by Random Methods

Amino acid sequence variants of a protein, e.g., a PKC delta agonist or antagonist, can be prepared by random mutagenesis of DNA which encodes a protein or a particular domain or region of a protein. Useful methods include PCR mutagenesis and saturation mutagenesis. A library of random amino acid sequence variants can also be generated by the synthesis of a set of degenerate oligonucleotide sequences. (Methods for screening proteins in a library of variants, e.g., screening for PKC delta modulating activity, are elsewhere herein.)

#### PCR Mutagenesis

In PCR mutagenesis, reduced Taq polymerase fidelity is used to introduce random mutations into a cloned fragment of DNA (Leung et al., 1989, *Technique* 1:11-15). This is a very powerful and relatively rapid method of introducing random mutations. The DNA region to be mutagenized is amplified using the polymerase chain reaction (PCR) under conditions that reduce the fidelity of DNA synthesis by Taq DNA polymerase, e.g., by using a dGTP/dATP ratio of five and adding Mn<sup>2+</sup> to the PCR reaction. The pool of amplified DNA fragments are inserted into appropriate cloning vectors to provide random mutant libraries.

#### Saturation Mutagenesis

Saturation mutagenesis allows for the rapid introduction of a large number of single base substitutions into cloned DNA fragments (Mayers et al., 1985, *Science* 229:242). This technique includes generation of mutations, e.g., by chemical treatment or irradiation of single-stranded DNA in vitro, and synthesis of a complimentary DNA strand. The mutation frequency can be modulated by modulating the severity of the treatment, and essentially all possible base substitutions can be obtained. Because this procedure does not involve a genetic selection for mutant fragments both neutral substitutions, as well as those that alter function, are obtained. The distribution of point mutations is not biased toward conserved sequence elements.

#### Degenerate Oligonucleotides

A library of homologs can also be generated from a set of degenerate oligonucleotide sequences. Chemical synthesis of a degenerate sequences can be carried out in an automatic DNA synthesizer, and the synthetic genes then ligated into an appropriate expression vector. The synthesis of degenerate oligonucleotides is known in the art (see for example, Narang, SA (1983) *Tetrahedron* 39:3; Itakura et al. (1981) *Recombinant DNA, Proc 3rd Cleveland Sympos. Macromolecules*, ed. A G Walton, Amsterdam: Elsevier pp273-289; Itakura et al. (1984) *Annu. Rev. Biochem.* 53:323; Itakura et al. (1984) *Science* 198:1056; Ike et al. (1983) *Nucleic Acid Res.* 11:477. Such techniques have been employed in the directed evolution of other proteins (see, for example, Scott et al. (1990) *Science* 249:386-390; Roberts et al. (1992) *PNAS* 89:2429-2433; Devlin et al. (1990) *Science* 249: 404-406; Cwirla et al. (1990) *PNAS* 87: 6378-6382; as well as U.S. Pat. Nos. 5,223, 409, 5,198,346, and 5,096,815).

### Generation of Analogs: Production of Altered DNA and Peptide Sequences by Directed Mutagenesis

Non-random or directed, mutagenesis techniques can be used to provide specific sequences or mutations in specific regions. These techniques can be used to create variants which include, e.g., deletions, insertions, or substitutions, of residues of the known amino acid sequence of a protein. The sites for mutation can be modified individually or in series, e.g., by (1) substituting first with conserved amino acids and then with more radical choices depending upon results achieved, (2) deleting the target residue, or (3) inserting residues of the same or a different class adjacent to the located site, or combinations of options 1-3.

#### Alanine Scanning Mutagenesis

Alanine scanning mutagenesis is a useful method for identification of certain residues or regions of the desired protein that are preferred locations or domains for mutagenesis, Cunningham and Wells (*Science* 244:1081-1085, 1989). In alanine scanning, a residue or group of target residues are identified (e.g., charged residues such as Arg, Asp, His, Lys, and Glu) and replaced by a neutral or negatively charged amino acid (most preferably alanine or polyalanine). Replacement of an amino acid can affect the interaction of the amino acids with the surrounding aqueous environment in or outside the cell. Those domains demonstrating functional sensitivity to the substitutions are then refined by introducing further or other variants at or for the sites of substitution. Thus, while the site for introducing an amino acid sequence variation is predetermined, the nature of the mutation per se need not be predetermined. For example, to optimize the performance of a mutation at a given site, alanine scanning or random mutagenesis may be conducted at the target codon or region and the expressed desired protein subunit variants are screened for the optimal combination of desired activity.

#### Oligonucleotide-Mediated Mutagenesis

Oligonucleotide-mediated mutagenesis is a useful method for preparing substitution, deletion, and insertion variants of DNA, see, e.g., Adelman et al., (*DNA* 2:183, 1983). Briefly, the desired DNA is altered by hybridizing an oligonucleotide encoding a mutation to a DNA template, where the template is the single-stranded form of a plasmid or bacteriophage containing the unaltered or native DNA sequence of the desired protein. After hybridization, a DNA polymerase is used to synthesize an entire second complementary strand of the template that will thus incorporate the oligonucleotide primer, and will code for the selected alteration in the desired protein DNA. Generally, oligonucleotides of at least 25 nucleotides in length are used. An optimal oligonucleotide will have 12 to 15 nucleotides that are completely complementary to the template on either side of the nucleotide(s) coding for the mutation. This ensures that the oligonucleotide will hybridize properly to the single-stranded DNA template molecule. The oligonucleotides are readily synthesized using techniques known in the art such as that described by Crea et al. (*Proc. Natl. Acad. Sci.* (1978) USA, 75: 5765).

#### Cassette Mutagenesis

Another method for preparing variants, cassette mutagenesis, is based on the technique described by Wells et al. (*Gene*, 34:315[1985]). The starting material is a plasmid (or other vector) which includes the protein subunit DNA to be mutated. The codon(s) in the protein subunit DNA to be mutated are identified. There must be a unique restriction endonuclease site on each side of the identified mutation site(s). If no such restriction sites exist, they may be generated using the above-described oligonucleotide-mediated mutagenesis method to introduce them at appropriate locations in the desired protein subunit DNA. After the restriction

sites have been introduced into the plasmid, the plasmid is cut at these sites to linearize it. A double-stranded oligonucleotide encoding the sequence of the DNA between the restriction sites but containing the desired mutation(s) is synthesized using standard procedures. The two strands are synthesized separately and then hybridized together using standard techniques. This double-stranded oligonucleotide is referred to as the cassette. This cassette is designed to have 3' and 5' ends that are comparable with the ends of the linearized plasmid, such that it can be directly ligated to the plasmid. This plasmid now contains the mutated desired protein subunit DNA sequence.

#### Combinatorial Mutagenesis

Combinatorial mutagenesis can also be used to generate mutants. For example, the amino acid sequences for a group of homologs or other related proteins are aligned, preferably to promote the highest homology possible. All of the amino acids which appear at a given position of the aligned sequences can be selected to create a degenerate set of combinatorial sequences. The variegated library of variants is generated by combinatorial mutagenesis at the nucleic acid level, and is encoded by a variegated gene library. For example, a mixture of synthetic oligonucleotides can be enzymatically ligated into gene sequences such that the degenerate set of potential sequences are expressible as individual peptides, or alternatively, as a set of larger fusion proteins containing the set of degenerate sequences.

#### Primary High-Through-Put Methods for Screening Libraries of Peptide Fragments or Homologs

Various techniques are known in the art for screening generated mutant gene products. Techniques for screening large gene libraries often include cloning the gene library into replicable expression vectors, transforming appropriate cells with the resulting library of vectors, and expressing the genes under conditions in which detection of a desired activity, assembly into a trimeric molecules, binding to natural ligands, e.g., a receptor or substrates, facilitates relatively easy isolation of the vector encoding the gene whose product was detected. Each of the techniques described below is amenable to high throughput analysis for screening large numbers of sequences created, e.g., by random mutagenesis techniques.

#### Two Hybrid Systems

Two hybrid (interaction trap) assays can be used to identify a protein that interacts with a PKC, e.g., PKC delta. These may include agonists, superagonists, and antagonists of PKC or PKC delta. (The subject protein and a protein it interacts with are used as the bait protein and fish proteins.). These assays rely on detecting the reconstitution of a functional transcriptional activator mediated by protein-protein interactions with a bait protein. In particular, these assays make use of chimeric genes which express hybrid proteins. The first hybrid comprises a DNA-binding domain fused to the bait protein, e.g., a PKC delta molecule or a fragment thereof. The second hybrid protein contains a transcriptional activation domain fused to a "fish" protein, e.g. an expression library. If the fish and bait proteins are able to interact, they bring into close proximity the DNA-binding and transcriptional activator domains. This proximity is sufficient to cause transcription of a reporter gene which is operably linked to a transcriptional regulatory site which is recognized by the DNA binding domain, and expression of the marker gene can be detected and used to score for the interaction of the bait protein with another protein.

## Display Libraries

In one approach to screening assays, the candidate peptides are displayed on the surface of a cell or viral particle, and the ability of particular cells or viral particles to bind an appropriate receptor protein via the displayed product is detected in a "panning assay". For example, the gene library can be cloned into the gene for a surface membrane protein of a bacterial cell, and the resulting fusion protein detected by panning (Ladner et al., WO 88/06630; Fuchs et al. (1991) *Bio/Technology* 9:1370-1371; and Goward et al. (1992) *TIBS* 18:136-140). In a similar fashion, a detectably labeled ligand can be used to score for potentially functional peptide homologs. Fluorescently labeled ligands, e.g., receptors, can be used to detect homolog which retain ligand-binding activity. The use of fluorescently labeled ligands, allows cells to be visually inspected and separated under a fluorescence microscope, or, where the morphology of the cell permits, to be separated by a fluorescence-activated cell sorter.

A gene library can be expressed as a fusion protein on the surface of a viral particle. For instance, in the filamentous phage system, foreign peptide sequences can be expressed on the surface of infectious phage, thereby conferring two significant benefits. First, since these phage can be applied to affinity matrices at concentrations well over  $10^{13}$  phage per milliliter, a large number of phage can be screened at one time. Second, since each infectious phage displays a gene product on its surface, if a particular phage is recovered from an affinity matrix in low yield, the phage can be amplified by another round of infection. The group of almost identical *E. coli* filamentous phages M13, fd., and fl are most often used in phage display libraries. Either of the phage gIII or gVIII coat proteins can be used to generate fusion proteins without disrupting the ultimate packaging of the viral particle. Foreign epitopes can be expressed at the  $\text{NH}_2$ -terminal end of pIII and phage bearing such epitopes recovered from a large excess of phage lacking this epitope (Ladner et al. PCT publication WO 90/02909; Garrard et al., PCT publication WO 92/09690; Marks et al. (1992) *J. Biol Chem.* 267:16007-16010; Griffiths et al. (1993) *EMBO J* 12:725-734; Clackson et al. (1991) *Nature* 352:624-628; and Barbas et al. (1992) *PNAS* 89:4457-4461).

A common approach uses the maltose receptor of *E. coli* (the outer membrane protein, LamB) as a peptide fusion partner (Charbit et al. (1986) *EMBO* 5, 3029-3037). Oligonucleotides have been inserted into plasmids encoding the LamB gene to produce peptides fused into one of the extracellular loops of the protein. These peptides are available for binding to ligands, e.g., to antibodies, and can elicit an immune response when the cells are administered to animals. Other cell surface proteins, e.g., OmpA (Schorr et al. (1991) *Vaccines* 9, pp. 387-392), PhoE (Agterberg, et al. (1990) *Gene* 88, 37-45), and PAL (Fuchs et al. (1991) *Bio/Tech* 9, 1369-1372), as well as large bacterial surface structures have served as vehicles for peptide display. Peptides can be fused to pilin, a protein which polymerizes to form the pilus-a conduit for interbacterial exchange of genetic information (Thiry et al. (1989) *Appl. Environ. Microbiol.* 55, 984-993). Because of its role in interacting with other cells, the pilus provides a useful support for the presentation of peptides to the extracellular environment. Another large surface structure used for peptide display is the bacterial motive organ, the flagellum. Fusion of peptides to the subunit protein flagellin offers a dense array of many peptides copies on the host cells (Kuwajima et al. (1988) *Bio/Tech.* 6, 1080-1083). Surface proteins of other bacterial species have also served as peptide fusion partners. Examples include the *Staphylococcus* protein A and the outer membrane protease IgA of *Neisseria*

(Hansson et al. (1992) *J. Bacteriol.* 174, 4239-4245 and Klause et al. (1990) *EMBO J.* 9, 1991-1999).

In the filamentous phage systems and the LamB system described above, the physical link between the peptide and its encoding DNA occurs by the containment of the DNA within a particle (cell or phage) that carries the peptide on its surface. Capturing the peptide captures the particle and the DNA within. An alternative scheme uses the DNA-binding protein LacI to form a link between peptide and DNA (Cull et al. (1992) *PNAS USA* 89:1865-1869). This system uses a plasmid containing the LacI gene with an oligonucleotide cloning site at its 3'-end. Under the controlled induction by arabinose, a LacI-peptide fusion protein is produced. This fusion retains the natural ability of LacI to bind to a short DNA sequence known as LacO operator (LacO). By installing two copies of LacO on the expression plasmid, the LacI-peptide fusion binds tightly to the plasmid that encoded it. Because the plasmids in each cell contain only a single oligonucleotide sequence and each cell expresses only a single peptide sequence, the peptides become specifically and stably associated with the DNA sequence that directed its synthesis. The cells of the library are gently lysed and the peptide-DNA complexes are exposed to a matrix of immobilized receptor to recover the complexes containing active peptides. The associated plasmid DNA is then reintroduced into cells for amplification and DNA sequencing to determine the identity of the peptide ligands. As a demonstration of the practical utility of the method, a large random library of dodecapeptides was made and selected on a monoclonal antibody raised against the opioid peptide dynorphin B. A cohort of peptides was recovered, all related by a consensus sequence corresponding to a six-residue portion of dynorphin B. (Cull et al. (1992) *Proc. Natl. Acad. Sci. U.S.A.* 89-1869)

This scheme, sometimes referred to as peptides-on-plasmids, differs in two important ways from the phage display methods. First, the peptides are attached to the C-terminus of the fusion protein, resulting in the display of the library members as peptides having free carboxy termini. Both of the filamentous phage coat proteins, pIII and pVIII, are anchored to the phage through their C-termini, and the guest peptides are placed into the outward-extending N-terminal domains. In some designs, the phage-displayed peptides are presented right at the amino terminus of the fusion protein. (Cwirla, et al. (1990) *Proc. Natl. Acad. Sci. U.S.A.* 87, 6378-6382) A second difference is the set of biological biases affecting the population of peptides actually present in the libraries. The LacI fusion molecules are confined to the cytoplasm of the host cells. The phage coat fusions are exposed briefly to the cytoplasm during translation but are rapidly secreted through the inner membrane into the periplasmic compartment, remaining anchored in the membrane by their C-terminal hydrophobic domains, with the N-termini, containing the peptides, protruding into the periplasm while awaiting assembly into phage particles. The peptides in the LacI and phage libraries may differ significantly as a result of their exposure to different proteolytic activities. The phage coat proteins require transport across the inner membrane and signal peptidase processing as a prelude to incorporation into phage. Certain peptides exert a deleterious effect on these processes and are underrepresented in the libraries (Gallop et al. (1994) *J. Med. Chem.* 37(9):1233-1251). These particular biases are not a factor in the LacI display system.

The number of small peptides available in recombinant random libraries is enormous. Libraries of  $10^7$ - $10^9$  independent clones are routinely prepared. Libraries as large as  $10^{11}$  recombinants have been created, but this size approaches the practical limit for clone libraries. This limitation in library

size occurs at the step of transforming the DNA containing randomized segments into the host bacterial cells. To circumvent this limitation, an in vitro system based on the display of nascent peptides in polysome complexes has recently been developed. This display library method has the potential of producing libraries 3-6 orders of magnitude larger than the currently available phage/phagemid or plasmid libraries. Furthermore, the construction of the libraries, expression of the peptides, and screening, is done in an entirely cell-free format.

In one application of this method (Gallop et al. (1994) *J. Med. Chem.* 37(9):1233-1251), a molecular DNA library encoding  $10^{12}$  decapeptides was constructed and the library expressed in an *E. coli* S30 in vitro coupled transcription/translation system. Conditions were chosen to stall the ribosomes on the mRNA, causing the accumulation of a substantial proportion of the RNA in polysomes and yielding complexes containing nascent peptides still linked to their encoding RNA. The polysomes are sufficiently robust to be affinity purified on immobilized receptors in much the same way as the more conventional recombinant peptide display libraries are screened. RNA from the bound complexes is recovered, converted to cDNA, and amplified by PCR to produce a template for the next round of synthesis and screening. The polysome display method can be coupled to the phage display system. Following several rounds of screening, cDNA from the enriched pool of polysomes was cloned into a phagemid vector. This vector serves as both a peptide expression vector, displaying peptides fused to the coat proteins, and as a DNA sequencing vector for peptide identification. By expressing the polysome-derived peptides on phage, one can either continue the affinity selection procedure in this format or assay the peptides on individual clones for binding activity in a phage ELISA, or for binding specificity in a completion phage ELISA (Barret, et al. (1992) *Anal. Biochem.* 204,357-364). To identify the sequences of the active peptides one sequences the DNA produced by the phagemid host.

#### Secondary Screens

The high through-put assays described above can be followed by secondary screens in order to identify further biological activities which will, e.g., allow one skilled in the art to differentiate agonists from antagonists. The type of a secondary screen used will depend on the desired activity that needs to be tested. For example, an assay can be developed in which the ability to inhibit an interaction between a protein of interest (e.g., PKC delta) and a ligand (e.g., a PKC substrate) can be used to identify antagonists from a group of peptide fragments isolated through one of the primary screens described above.

Therefore, methods for generating fragments and analogs and testing them for activity are known in the art. Once the core sequence of interest is identified, it is routine to perform for one skilled in the art to obtain analogs and fragments.

#### Peptide Mimetics

The invention also provides for reduction of the protein binding domains of the subject polypeptides, e.g., PKC delta, to generate mimetics, e.g. peptide or non-peptide agents. See, for example, "Peptide inhibitors of human papillomavirus protein binding to retinoblastoma gene protein" European patent applications EP 0 412 762 and EP 0 031 080.

Non-hydrolyzable peptide analogs of critical residues can be generated using benzodiazepine (e.g., see Freidinger et al. in *Peptides: Chemistry and Biology*, G. R. Marshall ed., ESCOM Publisher: Leiden, Netherlands, 1988), azepine (e.g., see Huffman et al. in *Peptides: Chemistry and Biology*, G. R. Marshall ed., ESCOM Publisher: Leiden, Netherlands,

1988), substituted gamma lactam rings (Garvey et al. in *Peptides: Chemistry and Biology*, G. R. Marshall ed., ESCOM Publisher: Leiden, Netherlands, 1988), keto-methylene pseudopeptides (Ewenson et al. (1986) *J Med Chem* 29:295; and Ewenson et al. in *Peptides: Structure and Function* (Proceedings of the 9th American Peptide Symposium) Pierce Chemical Co. Rockland, Ill., 1985),  $\beta$ -turn dipeptide cores (Nagai et al. (1985) *Tetrahedron Lett* 26:647; and Sato et al. (1986) *J Chem Soc Perkin Trans* 1:1231), and  $\beta$ -aminoalcohols (Gordon et al. (1985) *Biochem Biophys Res Commun* 126:419; and Dann et al. (1986) *Biochem Biophys Res Commun* 134:71).

#### Antibodies

The invention also includes antibodies specifically reactive with a PKC delta described herein. Anti-protein/anti-peptide antisera or monoclonal antibodies can be made as described herein by using standard protocols (See, for example, *Antibodies: A Laboratory Manual* ed. by Harlow and Lane (Cold Spring Harbor Press: 1988)).

PKC delta, or a portion or fragment thereof, can be used as an immunogen to generate antibodies that bind the component using standard techniques for polyclonal and monoclonal antibody preparation. The full-length component protein can be used or, alternatively, antigenic peptide fragments of the component can be used as immunogens.

Typically, a peptide is used to prepare antibodies by immunizing a suitable subject, (e.g., rabbit, goat, mouse or other mammal) with the immunogen. An appropriate immunogenic preparation can contain, for example, a recombinant PKC peptide, or a chemically synthesized PKC peptide or analog. See, e.g., U.S. Pat. No. 5,460,959; and co-pending U.S. applications Ser. Nos. 08/334,797; 08/231,439; 08/334,455; and 08/928,881 which are hereby expressly incorporated by reference in their entirety. The nucleotide and amino acid sequences of PKC delta are known. The preparation can further include an adjuvant, such as Freund's complete or incomplete adjuvant, or similar immunostimulatory agent. Immunization of a suitable subject with an immunogenic PKC delta preparation induces a polyclonal anti-PKC antibody response.

Antibodies to PKC delta or fragments thereof, can be used to inhibit the levels of such a component, thereby increasing NOS activity. Examples of antibody fragments include F(v), Fab, Fab' and F(ab')<sub>2</sub> fragments which can be generated by treating the antibody with an enzyme such as pepsin. The term "monoclonal antibody" or "monoclonal antibody composition", as used herein, refers to a population of antibody molecules that contain only one species of an antigen binding site capable of immunoreacting with a particular epitope. A monoclonal antibody composition thus typically displays a single binding affinity for a particular protein with which it immunoreacts.

Additionally, antibodies produced by genetic engineering methods, such as chimeric and humanized monoclonal antibodies, comprising both human and non-human portions, which can be made using standard recombinant DNA techniques, can be used. Such chimeric and humanized monoclonal antibodies can be produced by genetic engineering using standard DNA techniques known in the art, for example using methods described in Robinson et al. International Application No. PCT/US86/02269; Akira, et al. European Patent Application 184,187; Taniguchi, M., European Patent Application 171,496; Morrison et al. European Patent Application 173,494; Neuberger et al. PCT International Publication No. WO 86/01533; Cabilly et al. U.S. Pat. No. 4,816,567; Cabilly et al. European Patent Application 125,023; Better et



al., *Science* 240:1041-1043, 1988; Liu et al., *PNAS* 84:3439-3443, 1987; Liu et al., *J. Immunol.* 139:3521-3526, 1987; Sun et al. *PNAS* 84:214-218, 1987; Nishimura et al., *Canc. Res.* 47:999-1005, 1987; Wood et al., *Nature* 314:446-449, 1985; and Shaw et al., *J. Natl. Cancer Inst.* 80:1553-1559, 1988); Morrison, S. L., *Science* 229:1202-1207, 1985; Oi et al., *BioTechniques* 4:214, 1986; Winter U.S. Pat. No. 5,225,539; Jones et al., *Nature* 321:552-525, 1986; Verhoeyan et al., *Science* 239:1534, 1988; and Beidler et al., *J. Immunol.* 141:4053-4060, 1988.

In addition, a human monoclonal antibody directed against a PKC described herein can be made using standard techniques. For example, human monoclonal antibodies can be generated in transgenic mice or in immune deficient mice engrafted with antibody-producing human cells. Methods of generating such mice are describe, for example, in Wood et al. PCT publication WO 91/00906, Kucherlapati et al. PCT publication WO 91/10741; Lonberg et al. PCT publication WO 92/03918; Kay et al. PCT publication WO 92/03917; Kay et al. PCT publication WO 93/12227; Kay et al. PCT publication 94/25585; Rajewsky et al. Pct publication WO 94/04667; Ditullio et al. PCT publication WO 95/17085; Lonberg, N. et al. (1994) *Nature* 368:856-859; Green, L. L. et al. (1994) *Nature Genet.* 7:13-21; Morrison, S. L. et al. (1994) *Proc. Natl. Acad. Sci. USA* 81:6851-6855; Bruggeman et al. (1993) *Year Immunol* 7:33-40; Choi et al. (1993) *Nature Genet.* 4:117-123; Tuailon et al. (1993) *PNAS* 90:3720-3724; Bruggeman et al. (1991) *Eur J Immunol* 21:1323-1326; Duchosal et al. PCT publication WO 93/05796; U.S. Pat. No. 5,411,749; McCune et al. (1988) *Science* 241:1632-1639), Kamel-Reid et al. (1988) *Science* 242:1706; Spanopoulou (1994) *Genes & Development* 8:1030-1042; Shinkai et al. (1992) *Cell* 68:855-868). A human antibody-transgenic mouse or an immune deficient mouse engrafted with human antibody-producing cells or tissue can be immunized with a PKC delta isoform described herein or an antigenic peptide thereof and splenocytes from these immunized mice can then be used to create hybridomas. Methods of hybridoma production are well known.

Human monoclonal antibodies against a PKC isoform described herein can also be prepared by constructing a combinatorial immunoglobulin library, such as a Fab phage display library or a scFv phage display library, using immunoglobulin light chain and heavy chain cDNAs prepared from mRNA derived from lymphocytes of a subject. See, e.g., McCafferty et al. PCT publication WO 92/01047; Marks et al. (1991) *J. Mol. Biol.* 222:581-597; and Griffiths et al. (1993) *EMBO J* 12:725-734. In addition, a combinatorial library of antibody variable regions can be generated by mutating a known human antibody. For example, a variable region of a human antibody known to bind PKC delta, can be mutated, by for example using randomly altered mutagenized oligonucleotides, to generate a library of mutated variable regions which can then be screened to bind to a PKC. Methods of inducing random mutagenesis within the CDR regions of immunoglobulin heavy and/or light chains, methods of crossing randomized heavy and light chains to form pairings and screening methods can be found in, for example, Barbas et al. PCT publication WO 96/07754; Barbas et al. (1992) *Proc. Nat'l Acad. Sci. USA* 89:4457-4461.

The immunoglobulin library can be expressed by a population of display packages, preferably derived from filamentous phage, to form an antibody display library. Examples of methods and reagents particularly amenable for use in generating antibody display library can be found in, for example, Ladner et al. U.S. Pat. No. 5,223,409; Kang et al. PCT publication WO 92/18619; Dower et al. PCT publication WO

91/17271; Winter et al. PCT publication WO 92/20791; Markland et al. PCT publication WO 92/15679; Breitling et al. PCT publication WO 93/01288; McCafferty et al. PCT publication WO 92/01047; Garrard et al. PCT publication WO 92/09690; Ladner et al. PCT publication WO 90/02809; Fuchs et al. (1991) *Bio/Technology* 9:1370-1372; Hay et al. (1992) *Hum Antibod Hybridomas* 3:81-85; Huse et al. (1989) *Science* 246:1275-1281; Griffiths et al. (1993) supra; Hawkins et al. (1992) *J Mol Biol* 226:889-896; Clackson et al. (1991) *Nature* 352:624-628; Gram et al. (1992) *PNAS* 89:3576-3580; Garrad et al. (1991) *Bio/Technology* 9:1373-1377; Hoogenboom et al. (1991) *Nuc Acid Res* 19:4133-4137; and Barbas et al. (1991) *PNAS* 88:7978-7982. Once displayed on the surface of a display package (e.g., filamentous phage), the antibody library is screened to identify and isolate packages that express an antibody that binds a PKC described herein. In a preferred embodiment, the primary screening of the library involves panning with an immobilized PKC described herein and display packages expressing antibodies that bind immobilized PKC described herein are selected.

#### Antisense Nucleic Acid Sequences

Nucleic acid molecules which are antisense to a nucleotide encoding a PKC described herein, e.g., PKC delta can be used as an agent which inhibits expression of the PKC. An "antisense" nucleic acid includes a nucleotide sequence which is complementary to a "sense" nucleic acid encoding the component, e.g., complementary to the coding strand of a double-stranded cDNA molecule or complementary to an mRNA sequence. Accordingly, an antisense nucleic acid can form hydrogen bonds with a sense nucleic acid. The antisense nucleic acid can be complementary to an entire coding strand, or to only a portion thereof. For example, an antisense nucleic acid molecule which antisense to the "coding region" of the coding strand of a nucleotide sequence encoding the component can be used.

The coding strand sequence encoding PKC delta is known. Given the coding strand sequences encoding these isozymes, antisense nucleic acids can be designed according to the rules of Watson and Crick base pairing. The antisense nucleic acid molecule can be complementary to the entire coding region of mRNA, but more preferably is an oligonucleotide which is antisense to only a portion of the coding or noncoding region of mRNA. For example, the antisense oligonucleotide can be complementary to the region surrounding the translation start site of the mRNA. An antisense oligonucleotide can be, for example, about 5, 10, 15, 20, 25, 30, 35, 40, 45 or 50 nucleotides in length. An antisense nucleic acid can be constructed using chemical synthesis and enzymatic ligation reactions using procedures known in the art. For example, an antisense nucleic acid (e.g., an antisense oligonucleotide) can be chemically synthesized using naturally occurring nucleotides or variously modified nucleotides designed to increase the biological stability of the molecules or to increase the physical stability of the duplex formed between the antisense and sense nucleic acids, e.g., phosphorothioate derivatives and acridine substituted nucleotides can be used. Examples of modified nucleotides which can be used to generate the antisense nucleic acid include 5-fluorouracil, 5-bromouracil, 5-chlorouracil, 5-iodouracil, hypoxanthine, xanthine, 4-acytylcytosine, 5-(carboxyhydroxymethyl) uracil, 5-carboxymethylaminomethyl-2-thiouridine, 5-carboxymethylaminomethyluracil, dihydrouracil, beta-D-galactosylqueosine, inosine, N6-isopentenyladenine, 1-methylguanine, 1-methylinosine, 2,2-dimethylguanine, 2-methyladenine, 2-methylguanine, 3-methylcytosine, 5-methylcytosine, N6-adenine, 7-methylguanine, 5-methylaminomethyluracil, 5-meth-



oxyaminomethyl-2-thiouracil, beta-D-mannosylqueosine, 5'-methoxycarboxymethyluracil, 5-methoxyuracil, 2-methylthio-N6-isopentenyladenine, uraci-5-oxyacetic acid (v), wybutoxosine, pseudouracil, queosine, 2-thiocytosine, 5-methyl-2-thiouracil, 2-thiouracil, 4-thiouracil, 5-methyluracil, uracil-5-oxyacetic acid methylester, uracil-5-oxyacetic acid (v), 5-methyl-2-thiouracil, 3-(3-amino-3-N-2-carboxypropyl) uracil, (acp3)w, and 2,6-diaminopurine. Alternatively, the antisense nucleic acid can be produced biologically using an expression vector into which a nucleic acid has been subcloned in an antisense orientation (i.e., RNA transcribed from the inserted nucleic acid will be of an antisense orientation to a target nucleic acid of interest.

#### Administration

An agent which modulates the level of expression of PKC delta as described herein can be administered to a subject by standard methods. For example, the agent can be administered by any of a number of different routes including intravenous, intradermal, subcutaneous, oral (e.g., inhalation), transdermal (topical), and transmucosal. In one embodiment, the PKC modulating agent can be administered orally. In another embodiment, the agent is administered by injection, e.g., intramuscularly, or intravenously.

The agent which modulates protein levels, e.g., nucleic acid molecules, polypeptides, fragments or analogs, modulators, and antibodies (also referred to herein as "active compounds") can be incorporated into pharmaceutical compositions suitable for administration to a subject, e.g., a human. Such compositions typically include the nucleic acid molecule, polypeptide, modulator, or antibody and a pharmaceutically acceptable carrier. As used herein the language "pharmaceutically acceptable carrier" is intended to include any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents, and the like, compatible with pharmaceutical administration. The use of such media and agents for pharmaceutically active substances are known. Except insofar as any conventional media or agent is incompatible with the active compound, such media can be used in the compositions of the invention. Supplementary active compounds can also be incorporated into the compositions.

A pharmaceutical composition can be formulated to be compatible with its intended route of administration. Solutions or suspensions used for parenteral, intradermal, or subcutaneous application can include the following components: a sterile diluent such as water for injection, saline solution, fixed oils, polyethylene glycols, glycerine, propylene glycol or other synthetic solvents; antibacterial agents such as benzyl alcohol or methyl parabens; antioxidants such as ascorbic acid or sodium bisulfite; chelating agents such as ethylenediaminetetraacetic acid; buffers such as acetates, citrates or phosphates and agents for the adjustment of tonicity such as sodium chloride or dextrose. pH can be adjusted with acids or bases, such as hydrochloric acid or sodium hydroxide. The parenteral preparation can be enclosed in ampoules, disposable syringes or multiple dose vials made of glass or plastic.

Pharmaceutical compositions suitable for injectable use include sterile aqueous solutions (where water soluble) or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersion. For intravenous administration, suitable carriers include physiological saline, bacteriostatic water, Cremophor EL™ (BASF, Parsippany, N.J.) or phosphate buffered saline (PBS). In all cases, the composition must be sterile and should be fluid to the extent that easy syringability exists. It must be stable under the conditions of manufacture and storage and

must be preserved against the contaminating action of microorganisms such as bacteria and fungi. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (for example, glycerol, propylene glycol, and liquid polyethylene glycol, and the like), and suitable mixtures thereof. The proper fluidity can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of dispersion and by the use of surfactants. Prevention of the action of microorganisms can be achieved by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, ascorbic acid, thimerosal, and the like. In many cases, it will be preferable to include isotonic agents, for example, sugars, polyalcohols such as manitol, sorbitol, sodium chloride in the composition. Prolonged absorption of the injectable compositions can be brought about by including in the composition an agent which delays absorption, for example, aluminum monostearate and gelatin.

Sterile injectable solutions can be prepared by incorporating the active compound (e.g., a PKC delta polypeptide or anti-PKC delta antibody) in the required amount in an appropriate solvent with one or a combination of ingredients enumerated above, as required, followed by filtered sterilization. Generally, dispersions are prepared by incorporating the active compound into a sterile vehicle which contains a basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum drying and freeze-drying which yields a powder of the active ingredient plus any additional desired ingredient from a previously sterile-filtered solution thereof.

Oral compositions generally include an inert diluent or an edible carrier. They can be enclosed in gelatin capsules or compressed into tablets. For the purpose of oral therapeutic administration, the active compound can be incorporated with excipients and used in the form of tablets, troches, or capsules. Oral compositions can also be prepared using a fluid carrier for use as a mouthwash, wherein the compound in the fluid carrier is applied orally and swished and expectorated or swallowed. Pharmaceutically compatible binding agents, and/or adjuvant materials can be included as part of the composition. The tablets, pills, capsules, troches and the like can contain any of the following ingredients, or compounds of a similar nature: a binder such as microcrystalline cellulose, gum tragacanth or gelatin; an excipient such as starch or lactose, a disintegrating agent such as alginic acid, Primogel, or corn starch; a lubricant such as magnesium stearate or Sterotes; a glidant such as colloidal silicon dioxide; a sweetening agent such as sucrose or saccharin; or a flavoring agent such as peppermint, methyl salicylate, or orange flavoring.

Systemic administration can also be by transmucosal or transdermal means. For transmucosal or transdermal administration, penetrants appropriate to the barrier to be permeated are used in the formulation. Such penetrants are generally known, and include, for example, for transmucosal administration, detergents, bile salts, and fusidic acid derivatives. Transmucosal administration can be accomplished through the use of nasal sprays or suppositories. For transdermal administration, the active compounds are formulated into ointments, salves, gels, or creams as generally known in the art.

In one embodiment, the active compounds are prepared with carriers that will protect the compound against rapid elimination from the body, such as a controlled release formulation, including implants and microencapsulated delivery systems. Biodegradable, biocompatible polymers can be

used, such as ethylene vinyl acetate, polyanhydrides, polyglycolic acid, collagen, polyorthoesters, and polylactic acid. Methods for preparation of such formulations will be apparent to those skilled in the art. The materials can also be obtained commercially from Alza Corporation and Nova Pharmaceuticals, Inc. Liposomal suspensions (including liposomes targeted to infected cells with monoclonal antibodies to viral antigens) can also be used as pharmaceutically acceptable carriers. These can be prepared according to methods known to those skilled in the art, for example, as described in U.S. Pat. No. 4,522,811.

The nucleic acid molecules described herein can be inserted into vectors and used as gene therapy vectors. Gene therapy vectors can be delivered to a subject by, for example, intravenous injection, local administration (see U.S. Pat. No. 5,328,470) or by stereotactic injection (see e.g., Chen et al., *PNAS* 91:3054-3057, 1994). The pharmaceutical preparation of the gene therapy vector can include the gene therapy vector in an acceptable diluent, or can include a slow release matrix in which the gene delivery vehicle is imbedded. Alternatively, where the complete gene delivery vector can be produced intact from recombinant cells, e.g. retroviral vectors, the pharmaceutical preparation can include one or more cells which produce the gene delivery system.

The pharmaceutical compositions can be included in a container, pack, or dispenser together with instructions for administration.

#### Gene Therapy

The nucleic acids described herein, e.g., a nucleic acid encoding PKC delta, or an antisense nucleic acid, can be incorporated into gene constructs to be used as a part of a gene therapy protocol to deliver nucleic acids encoding either an agonistic or antagonistic form of PKC delta. The invention features expression vectors for in vivo transfection and expression of PKC described herein in particular cell types so as to reconstitute the function of, or alternatively, antagonize the function of the component in a cell in which that polypeptide is misexpressed. Expression constructs of such components may be administered in any biologically effective carrier, e.g. any formulation or composition capable of effectively delivering the component gene to cells in vivo. Approaches include insertion of the subject gene in viral vectors including recombinant retroviruses, adenovirus, adeno-associated virus, and herpes simplex virus-1, or recombinant bacterial or eukaryotic plasmids. Viral vectors transfect cells directly; plasmid DNA can be delivered with the help of, for example, cationic liposomes (lipofectin) or derivatized (e.g. antibody conjugated), polylysine conjugates, gramicidin S, artificial viral envelopes or other such intracellular carriers, as well as direct injection of the gene construct or CaPO<sub>4</sub> precipitation carried out in vivo.

A preferred approach for in vivo introduction of nucleic acid into a cell is by use of a viral vector containing nucleic acid, e.g. a cDNA, encoding PKC delta. Infection of cells with a viral vector has the advantage that a large proportion of the targeted cells can receive the nucleic acid. Additionally, molecules encoded within the viral vector, e.g., by a cDNA contained in the viral vector, are expressed efficiently in cells which have taken up viral vector nucleic acid.

Retrovirus vectors and adeno-associated virus vectors can be used as a recombinant gene delivery system for the transfer of exogenous genes in vivo, particularly into humans. These vectors provide efficient delivery of genes into cells, and the transferred nucleic acids are stably integrated into the chromosomal DNA of the host. The development of specialized cell lines (termed "packaging cells") which produce only replication-defective retroviruses has increased the utility of

retroviruses for gene therapy, and defective retroviruses are characterized for use in gene transfer for gene therapy purposes (for a review see Miller, A. D. (1990) *Blood* 76:271). A replication defective retrovirus can be packaged into virions which can be used to infect a target cell through the use of a helper virus by standard techniques. Protocols for producing recombinant retroviruses and for infecting cells in vitro or in vivo with such viruses can be found in *Current Protocols in Molecular Biology*, Ausubel, F. M. et al. (eds.) Greene Publishing Associates, (1989), Sections 9.10-9.14 and other standard laboratory manuals. Examples of suitable retroviruses include pLJ, pZIP, pWE and pEM which are known to those skilled in the art. Examples of suitable packaging virus lines for preparing both ecotropic and amphotropic retroviral systems include \*Crip, \*Cre, \*2 and \*Am. Retroviruses have been used to introduce a variety of genes into many different cell types, including epithelial cells, in vitro and/or in vivo (see for example Eglitis, et al. (1985) *Science* 230:1395-1398; Danos and Mulligan (1988) *Proc. Natl. Acad. Sci. USA* 85:6460-6464; Wilson et al. (1988) *Proc. Natl. Acad. Sci. USA* 85:3014-3018; Armentano et al. (1990) *Proc. Natl. Acad. Sci. USA* 87:6141-6145; Huber et al. (1991) *Proc. Natl. Acad. Sci. USA* 88:8039-8043; Ferry et al. (1991) *Proc. Natl. Acad. Sci. USA* 88:8377-8381; Chowdhury et al. (1991) *Science* 254:1802-1805; van Beusechem et al. (1992) *Proc. Natl. Acad. Sci. USA* 89:7640-7644; Kay et al. (1992) *Human Gene Therapy* 3:641-647; Dai et al. (1992) *Proc. Natl. Acad. Sci. USA* 89:10892-10895; Hwu et al. (1993) *J. Immunol.* 150:4104-4115; U.S. Pat. No. 4,868,116; U.S. Pat. No. 4,980,286; PCT Application WO 89/07136; PCT Application WO 89/02468; PCT Application WO 89/05345; and PCT Application WO 92/07573).

Another viral gene delivery system useful in the present invention utilizes adenovirus-derived vectors. The genome of an adenovirus can be manipulated such that it encodes and expresses a gene product of interest but is inactivated in terms of its ability to replicate in a normal lytic viral life cycle. See, for example, Berkner et al. (1988) *BioTechniques* 6:616; Rosenfeld et al. (1991) *Science* 252:431-434; and Rosenfeld et al. (1992) *Cell* 68:143-155. Suitable adenoviral vectors derived from the adenovirus strain Ad type 5 d1324 or other strains of adenovirus (e.g., Ad2, Ad3, Ad7 etc.) are known to those skilled in the art. Recombinant adenoviruses can be advantageous in certain circumstances in that they are not capable of infecting nondividing cells and can be used to infect a wide variety of cell types, including epithelial cells (Rosenfeld et al. (1992) cited supra). Furthermore, the virus particle is relatively stable and amenable to purification and concentration, and as above, can be modified so as to affect the spectrum of infectivity. Additionally, introduced adenoviral DNA (and foreign DNA contained therein) is not integrated into the genome of a host cell but remains episomal, thereby avoiding potential problems that can occur as a result of insertional mutagenesis in situ where introduced DNA becomes integrated into the host genome (e.g., retroviral DNA). Moreover, the carrying capacity of the adenoviral genome for foreign DNA is large (up to 8 kilobases) relative to other gene delivery vectors (Berkner et al. cited supra; Haj-Ahmand and Graham (1986) *J. Virol.* 57:267).

Yet another viral vector system useful for delivery of the subject gene is the adeno-associated virus (AAV). Adeno-associated virus is a naturally occurring defective virus that requires another virus, such as an adenovirus or a herpes virus, as a helper virus for efficient replication and a productive life cycle. (For a review see Muzyczka et al. (1992) *Curr. Topics in Micro. and Immunol.* 158:97-129). It is also one of the few viruses that may integrate its DNA into non-dividing

cells, and exhibits a high frequency of stable integration (see for example Flotte et al. (1992) *Am. J. Respir. Cell. Mol. Biol.* 7:349-356; Samulski et al. (1989) *J. Virol.* 63:3822-3828; and McLaughlin et al. (1989) *J. Virol.* 62:1963-1973). Vectors containing as little as 300 base pairs of AAV can be packaged and can integrate. Space for exogenous DNA is limited to about 4.5 kb. An AAV vector such as that described in Tratschin et al. (1985) *Mol. Cell. Biol.* 5:3251-3260 can be used to introduce DNA into cells. A variety of nucleic acids have been introduced into different cell types using AAV vectors (see for example Hermonat et al. (1984) *Proc. Natl. Acad. Sci. USA* 81:6466-6470; Tratschin et al. (1985) *Mol. Cell. Biol.* 4:2072-2081; Wondisford et al. (1988) *Mol. Endocrinol.* 2:32-39; Tratschin et al. (1984) *J. Virol.* 51:611-619; and Flotte et al. (1993) *J. Biol. Chem.* 268:3781-3790).

In addition to viral transfer methods, such as those illustrated above, non-viral methods can also be employed to cause expression of a PKC described herein in the tissue of a subject. Most nonviral methods of gene transfer rely on normal mechanisms used by mammalian cells for the uptake and intracellular transport of macromolecules. In preferred embodiments, non-viral gene delivery systems of the present invention rely on endocytic pathways for the uptake of the subject gene by the targeted cell. Exemplary gene delivery systems of this type include liposomal derived systems, polylysine conjugates, and artificial viral envelopes. Other embodiments include plasmid injection systems such as are described in Meuli et al. (2001) *J Invest Dermatol.* 116(1):131-135; Cohen et al. (2000) *Gene Ther* 7(22):1896-905; or Tam et al. (2000) *Gene Ther* 7(21):1867-74.

In a representative embodiment, a gene encoding PKC delta can be entrapped in liposomes bearing positive charges on their surface (e.g., lipofectins) and (optionally) which are tagged with antibodies against cell surface antigens of the target tissue (Mizuno et al. (1992) *No Shinkei Geka* 20:547-551; PCT publication WO91/06309; Japanese patent application 1047381; and European patent publication EP-A-43075).

In clinical settings, the gene delivery systems for the therapeutic gene can be introduced into a patient by any of a number of methods, each of which is familiar in the art. For instance, a pharmaceutical preparation of the gene delivery system can be introduced systemically, e.g. by intravenous injection, and specific transduction of the protein in the target cells occurs predominantly from specificity of transfection provided by the gene delivery vehicle, cell-type or tissue-type expression due to the transcriptional regulatory sequences controlling expression of the receptor gene, or a combination thereof. In other embodiments, initial delivery of the recombinant gene is more limited with introduction into the animal being quite localized. For example, the gene delivery vehicle can be introduced by catheter (see U.S. Pat. No. 5,328,470) or by stereotactic injection (e.g. Chen et al. (1994) *PNAS* 91:3054-3057).

The pharmaceutical preparation of the gene therapy construct can consist essentially of the gene delivery system in an acceptable diluent, or can comprise a slow release matrix in which the gene delivery vehicle is imbedded. Alternatively, where the complete gene delivery system can be produced in tact from recombinant cells, e.g. retroviral vectors, the pharmaceutical preparation can comprise one or more cells which produce the gene delivery system.

#### Cell Therapy

A PKC described herein can also be increased in a subject by introducing into a cell, e.g., an endothelial cell, a nucleotide sequence that modulates the production of PKC delta, e.g., a nucleotide sequence encoding a PKC delta polypeptide

or functional fragment or analog thereof, a promoter sequence, e.g., a promoter sequence from a PKC delta gene or from another gene; an enhancer sequence, e.g., 5' untranslated region (UTR), e.g., a 5' UTR from a PKC delta gene or from another gene, a 3' UTR, e.g., a 3' UTR from a PKC delta gene or from another gene; a polyadenylation site; an insulator sequence; or another sequence that modulates the expression of PKC delta. The cell can then be introduced into the subject.

Primary and secondary cells to be genetically engineered can be obtained from a variety of tissues and include cell types which can be maintained propagated in culture. For example, primary and secondary cells include fibroblasts, keratinocytes, epithelial cells (e.g., mammary epithelial cells, intestinal epithelial cells), endothelial cells, glial cells, neural cells, formed elements of the blood (e.g., lymphocytes, bone marrow cells), muscle cells (myoblasts) and precursors of these somatic cell types. Primary cells are preferably obtained from the individual to whom the genetically engineered primary or secondary cells are administered. However, primary cells may be obtained for a donor (other than the recipient).

The term "primary cell" includes cells present in a suspension of cells isolated from a vertebrate tissue source (prior to their being plated i.e., attached to a tissue culture substrate such as a dish or flask), cells present in an explant derived from tissue, both of the previous types of cells plated for the first time, and cell suspensions derived from these plated cells. The term "secondary cell" or "cell strain" refers to cells at all subsequent steps in culturing. Secondary cells are cell strains which consist of secondary cells which have been passaged one or more times.

Primary or secondary cells of vertebrate, particularly mammalian, origin can be transfected with an exogenous nucleic acid sequence which includes a nucleic acid sequence encoding a signal peptide, and/or a heterologous nucleic acid sequence, e.g., encoding a PKC delta, or an agonist or antagonist thereof, and produce the encoded product stably and reproducibly in vitro and in vivo, over extended periods of time. A heterologous amino acid can also be a regulatory sequence, e.g., a promoter, which causes expression, e.g., inducible expression or upregulation, of an endogenous sequence. An exogenous nucleic acid sequence can be introduced into a primary or secondary cell by homologous recombination as described, for example, in U.S. Pat. No. 5,641,670, the contents of which are incorporated herein by reference. The transfected primary or secondary cells may also include DNA encoding a selectable marker which confers a selectable phenotype upon them, facilitating their identification and isolation.

Vertebrate tissue can be obtained by standard methods such as a punch biopsy or other surgical methods of obtaining a tissue source of the primary cell type of interest. For example, punch biopsy is used to obtain skin as a source of fibroblasts or keratinocytes. A mixture of primary cells is obtained from the tissue, using known methods, such as enzymatic digestion or explanting. If enzymatic digestion is used, enzymes such as collagenase, hyaluronidase, dispase, pronase, trypsin, elastase and chymotrypsin can be used.

The resulting primary cell mixture can be transfected directly or it can be cultured first, removed from the culture plate and resuspended before transfection is carried out. Primary cells or secondary cells are combined with exogenous nucleic acid sequence to, e.g., stably integrate into their genomes, and treated in order to accomplish transfection. As used herein, the term "transfection" includes a variety of techniques for introducing an exogenous nucleic acid into a cell including calcium phosphate or calcium chloride precipi-

tation, microinjection, DEAE-dextrin-mediated transfection, lipofection or electrophoration, all of which are routine in the art.

Transfected primary or secondary cells undergo sufficient number doubling to produce either a clonal cell strain or a heterogeneous cell strain of sufficient size to provide the therapeutic protein to an individual in effective amounts. The number of required cells in a transfected clonal heterogeneous cell strain is variable and depends on a variety of factors, including but not limited to, the use of the transfected cells, the functional level of the exogenous DNA in the transfected cells, the site of implantation of the transfected cells (for example, the number of cells that can be used is limited by the anatomical site of implantation), and the age, surface area, and clinical condition of the patient.

The transfected cells, e.g., cells produced as described herein, can be introduced into an individual to whom the product is to be delivered. Various routes of administration and various sites (e.g., renal sub capsular, subcutaneous, central nervous system (including intrathecal), intravascular, intrahepatic, intrasplanchnic, intraperitoneal (including intraomental), intramuscularly implantation) can be used. One implanted in individual, the transfected cells produce the product encoded by the heterologous DNA or are affected by the heterologous DNA itself. For example, an individual who suffers from an insulin related disorder is a candidate for implantation of cells producing an antagonist of PKC delta described herein.

An immunosuppressive agent e.g., drug, or antibody, can be administered to a subject at a dosage sufficient to achieve

the desired therapeutic effect (e.g., inhibition of rejection of the cells). Dosage ranges for immunosuppressive drugs are known in the art. See, e.g., Freed et al. (1992) N. Engl. J. Med. 327:1549; Spencer et al. (1992) N. Engl. J. Med. 327:1541; Widner et al. (1992) n. Engl. J. Med. 327:1556). Dosage values may vary according to factors such as the disease state, age, sex, and weight of the individual.

All patents and references cited herein are hereby incorporated by reference in their entirety. Other embodiments are within the following claims.

What is claimed is:

1. A method for identifying a compound capable of inhibiting angiogenesis, comprising:
  - contacting a cell capable of expressing a PKC delta polypeptide with a test compound;
  - determining a level of PKC delta polypeptide expression in the cell;
  - selecting a test compound if it is capable of inhibiting PKC delta polypeptide expression;
  - assaying an effect of the selected test compound in an assay of angiogenesis, and selecting the compound if it is capable of inhibiting angiogenesis in the assay.
2. The method of claim 1, wherein the assay of angiogenesis comprises assaying tube formation in endothelial cells.
3. The method of claim 1, wherein the assay of angiogenesis comprises administering the selected compound to an animal model characterized by increased blood vessel formation.

\* \* \* \* \*

专利名称(译)	调节血管生成的方法		
公开(公告)号	<a href="#">US7410756</a>	公开(公告)日	2008-08-12
申请号	US10/464720	申请日	2003-06-17
申请(专利权)人(译)	Joslin糖尿病中心INC.		
当前申请(专利权)人(译)	Joslin糖尿病中心 , INC.		
[标]发明人	KING GEORGE L		
发明人	KING, GEORGE L.		
IPC分类号	C12Q1/70 G01N33/53 A61K49/00 C12Q1/48		
CPC分类号	A61K49/0004 C12Q1/485 G01N2500/00		
代理机构(译)	FISH & RICHARDSON P.C.		
优先权	60/219245 2000-07-18 US		
其他公开文献	US20030211109A1		
外部链接	<a href="#">Espacenet</a> <a href="#">USPTO</a>		

#### 摘要(译)

特征在于调节血管生成的方法，例如通过调节PKC $\delta$ 。该方法可用于治疗以异常和/或血管形成增加和/或渗透性增加为特征的疾病。该方法还可用于治疗以异常，不充分或不稳定的血管形成或增加的渗透性为特征的病症。还有诊断和筛查方法。