



(19) **United States**

(12) **Patent Application Publication** (10) **Pub. No.: US 2003/0186344 A1**

Holmes et al. (43) **Pub. Date: Oct. 2, 2003**

(54) **DETECTION AND QUANTITATION OF 8-OH-ADENINE USING MONOCLONAL ANTIBODIES**

(52) **U.S. Cl. 435/7.92; 435/326; 530/388.1**

(75) **Inventors: Eric H. Holmes, Bothell, WA (US); Thomas G. Greene, Seattle, WA (US)**

(57) **ABSTRACT**

Correspondence Address:
**CHRISTENSEN, O'CONNOR, JOHNSON, KINDNESS, PLLC
1420 FIFTH AVENUE
SUITE 2800
SEATTLE, WA 98101-2347 (US)**

The present invention relates to methods and materials for the detection and quantitation 8-OH-Ade in biological specimens. Specifically, the present invention is directed to a group of highly specific monoclonal antibodies reactive with the modified nucleoside structure 8-OH-Ade, and to various immunoassays for 8-OH-Ade utilizing these monoclonal antibodies. The monoclonal antibodies of the present invention may be used in assays for diagnosing or monitoring the progression of certain types of cancer, in addition to a variety of other diseases associated with mutagenesis resulting from oxidative damage of DNA. Assays utilizing the monoclonal antibodies of the present invention may also be used to analyze or monitor toxicant exposure, such as from environmental sources. The monoclonal antibodies of the present invention were prepared with the immunogen 8-OH-adenosine coupled to keyhole limpet hemocyanin (KLH), not to 8-OH-Ade directly. It is believed that the monoclonal antibodies bind with the base portion of the structure (8-OH-Ade) and not the carbohydrate (ribose) or protein linkage region of the conjugate, because, as demonstrated, conjugates bound to nucleosides other than 8-OH-adenosine were unreactive with these antibodies. Therefore, the antibodies of the present invention can be used to detect and quantitate (by the use of a standard curve) the presence of 8-OH-Ade in biological specimens of DNA. Procedures for such an assay include immobilizing the DNA, denaturing it to disrupt the base-pairing scheme exposing the free base structures, and quantitating the amount of 8-OH-Ade present per amount of DAN in a quantitative immunoassay.

(73) **Assignee: CytoChem, Inc.**

(21) **Appl. No.: 10/375,697**

(22) **Filed: Feb. 25, 2003**

Related U.S. Application Data

(63) Continuation of application No. 09/645,005, filed on Aug. 23, 2000, now abandoned, which is a continuation of application No. 09/043,461, filed on Mar. 18, 1998, now Pat. No. 6,187,551, filed as 371 of international application No. PCT/US96/15034, filed on Sep. 19, 1996.

(60) Provisional application No. 60/003,976, filed on Sep. 19, 1995.

Publication Classification

(51) **Int. Cl.⁷ G01N 33/53; G01N 33/537; G01N 33/543; C12N 5/06; C07K 16/00**




Fig. 1A.




Fig. 1B.




Fig. 2.




Fig. 3.




Fig. 4.




Fig. 5.




Fig. 6.




Fig. 7.




Fig. 8.




Fig. 9.




Fig. 10.




Fig. 11.




Fig. 12.




Fig. 13.




Fig. 14.




Fig. 15.




Fig. 16.




Fig. 17.




Fig. 18.




Fig. 19.




Fig. 20.




Fig. 21.




Fig. 22.




Fig. 23.




Fig. 24.




Fig. 25.




Fig. 26.




Fig. 27.




Fig. 28.

DETECTION AND QUANTITATION OF 8-OH-ADENINE USING MONOCLONAL ANTIBODIES

[0001] This invention was made with United States government support under National Institutes of Health Grant No. R43 CA67163. The United States government has certain rights in the invention.

FIELD OF THE INVENTION

[0002] The present invention relates generally to assays for 8-hydroxyadenine (8-OH-Ade) and, more particularly, to improved immunoassays for the detection and quantitation of 8-OH-Ade in biological specimens. The present invention further relates to new monoclonal antibodies directed against 8-OH-Ade for use in the immunoassays.

BACKGROUND OF THE INVENTION

[0003] 8-OH-Ade is a modified nucleotide base resulting from single electron oxidation reactions. Steenken, "Purine bases, nucleosides and nucleotides: Aqueous solution redox chemistry and transformation of their radical cations \dot{e} and OH Adducts," *Chem. Rev.*, Vol. 89, pp. 503-520 (1989). The presence of this oxidized base has been detected in biological specimens in a manner consistent with toxicant exposure and carcinogenesis. Malins et al., "The Etiology of Cancer: hydroxyl radical-induced DNA lesions in histologically normal livers of fish from a population with liver tumors," *Aquat. Toxicol.*, Vol. 20, pp. 123-130 (1991) and Malins et al., "The Etiology of Breast Cancer," *Cancer* Vol. 71, No. 10, pp. 3036-3043 (1993). Previous methods of detecting and quantitating 8-OH-Ade have relied on analysis by high performance liquid chromatography-electrochemical detection (HPLC-ECD) (Shigenaga et al., "Urinary 8-hydroxy-2'-Deoxyguanosine as a Biological Marker of In Vivo Oxidative DNA Damage," *Proc. National. Acad. Sci. USA*, Vol. 86, pp. 9697-9701 (1989)) and gas chromatography-mass spectrometry with selected ion monitoring (GC-MS/SIM) (Malins et al. (1993)) procedures. Recently, assays utilizing polyclonal antibodies have also been used to detect 8-OH-Ade. West et al., "Radioimmunoassay of 7, 8-dihydro-8-oxoadenine (8-hydroxyadenine)," *Int. J. Radiat. Biol.*, Vol. 42, No. 5, pp. 481-490 (1982). The present invention provides new and improved materials and methods for detecting and quantitating the 8-OH-Ade base structure in a biological specimen comprising a group of highly specific monoclonal antibodies against 8-OH-Ade which may be used in a variety of immunoassays.

[0004] Oxygen-free radicals are the primary mediators of cellular free-radical reactions. They are produced in normal or pathological cell metabolism, and as a result of exposure to a variety of exogenous sources of oxidative stress such as tobacco smoke, fatty acids in foods, iron and copper ions, and ethanol. Furthermore, ultraviolet light and ionizing radiation can stimulate the generation of oxygen-free radicals.

[0005] Oxygen-free radicals cause constant damage which the body's antioxidant defense systems usually repair so that a dynamic equilibrium is maintained. However, occasionally an overabundance of oxygen free radicals in the body occurs. Oxidative stress refers to the condition in which there is an overproduction of oxygen-free radicals or a deficiency in the antioxidant defense and repair mecha-

nisms. Examples of short-term oxidative stress reactions include ischemia, reperfusion injury, acute inflammation and hyperoxia. Dreher et al., "Role of Oxygen Free Radicals in Cancer Development," *European Journal of Cancer*, Vol. 32A(1), pp. 30-38 (1996).

[0006] While short-term oxidative stress does not generally result in severe or debilitating illness, chronic oxidative stress can result in oxidative damage to an organisms DNA, which in turn has been associated with a variety of diseases. It has been established that reactive oxygen species play a significant role in mutagenesis, carcinogenesis and tumor promotion. Bhimani et al., "Inhibition of Oxidative Stress in HeLa Cells by Chemopreventive Agents," *Cancer Research*, Vol. 53, pp. 4528-4533 (1993). The dismutation of superoxide yields hydrogen peroxide, which is highly reactive in vivo. Hydrogen peroxide reacts with partially reduced metal ions to form the hydroxyl radical which can directly inflict DNA damage. Dreher et al. (1996). If hydroxyl radicals are generated close to DNA, they can attack the purine and pyrimidine bases, causing mutations. Halliwell, "Free Radicals, Antioxidants, and Human Disease: Curiosity, Cause, or Consequence?," *The Lancet*, Vol. 344, pp. 721-724 (1994). A discussion of the mechanisms of oxygen-free radical related mutagenesis resulting from DNA damage can be found in Dreher et al., Role of Oxygen-Free Radicals in Cancer Development, *European Journal of Cancer*, Vol. 32A, No. 1, pp. 30-38 (1996).

[0007] The involvement of reactive oxygen species in the development of cancer in humans is supported by the abundant presence of oxidative DNA modifications in cancer tissue. Loft et al., "Cancer Risk and Oxidative DNA Damage in Man," *J. Mol. Med.*, Vol. 74, pp. 297-312 (1996). For example, in breast cancer, base lesion concentrations have been found to be substantial. Base lesions previously reported include 8-OH-Ade, among others. Malins, et al. (1990). It is believed that these base lesions play a pivotal roll in oncogenesis and may further serve as early predictors of breast cancer risk, in addition to a variety of other cancers, due to their inherently mutagenic and carcinogenic effects. See Halliwell (1994) and Bhimani et al. (1993) and Dreher et al. (1996).

[0008] Reactive oxygen species have also been implicated in the etiology and pathophysiology of many other human diseases including cardiovascular disease, chronic inflammatory disease, neurodegenerative diseases, rheumatoid arthritis, systemic lupus, erythematosis and sickle cell anemia. Bhimani et al. (1993) and Halliwell (1994).

[0009] The etiology of these diseases and their progression, as well as toxicant exposure, can be studied by measuring levels of oxidized DNA bases in biological specimens. Levels of oxidized DNA bases, such as 8-OH-Ade, may also be used as a predictor of risk of disease, thereby allowing preventive intervention before the clinical disease develops. Strickland et al., "Methodologies for Measuring Carcinogen Adducts in Humans," *Cancer Epidemiology, Biomarkers and Prevention*, Vol. 2, pp. 607-619 (1993).

[0010] One of the most abundant and most studied oxidative modifications of DNA-bases is the C-8 hydroxylation of guanine. Loft et al., "Cancer Risk and Oxidative DNA Damage in Man," *J. Mol. Med.*, Vol. 74, pp. 297-312 (1996). Other abundant oxidatively modified purines and pyrimidines include 8-oxoadenine, 2-hydroxyadenine, Fapy-A,

5-hydroxy cytosine, and thymine glycol, among others. Loft et al. (1996). Of interest to the present invention in the modified purine 8-OH-Ade.

[0011] The presence of oxidative damage in genomic and mitochondrial DNA obtained from biological specimens such as tissues and isolated cells has been studied by a variety of methods, including most commonly gas chromatography/mass spectroscopy with selective ion monitoring (GC/MS-SIM) and high-performance liquid chromatographic separation (HPLC) followed by detection by UV or electrochemistry. Four primary methods used for monitoring and quantifying oxidative DNA damage in biological specimens are discussed in detail in the review article by Strickland et al. (1993) and include immunoassay techniques, post-labeling, fluorescence spectroscopy and GC/MS.

[0012] In view of the great interest in DNA adducts, including 8-OH-Ade, efforts have been made to make antibodies sensitive to these oxidized nucleosides. The use of antibodies to detect DNA adducts began in the mid-1970s. Strickland et al. (1993). A variety of assays utilizing the antibodies have been developed, including the competitive radioimmunoassay, solid phase enzyme immunoassay, competitive solid phase enzyme immunoassay, ELISA, and immunoaffinity chromatography. Strickland et al. (1993). Monoclonal antibodies have been used to detect, for example, 1-methyladenosine and pseudouridine in urine (Matsuda et al., "An Immunohistochemical Analysis for Cancer of the Esophagus Using Monoclonal Antibodies specific for Modified Nucleosides," *Cancer*, Vol. 72, pp. 3571-3578 (1993); various derivatives of guanosine and thymine (Adamkiewicz et al., "Monoclonal Antibody-Based Immunoanalytical Methods for Detection of Carcinogen-Modified DNA Components," *Arc Sc. Publ.*, Vol. 70, pp. 403-41 (1986); and 8-OH-guanine (Lee et al., "Identification of 8-Hydroxyguanine Glycosylase Activity in Mammalian Tissues Using 8-Hydroxyguanine Specific Monoclonal Antibody," *Biochemical and Biophysical Research Communications*, Vol. 196, No. 3, pp. 1545-1551 (1993) and Yin et al., "Determination of 8-Hydroxydeoxyguanosine by an Immunoaffinity Chromatography-Monoclonal Antibody-Base Elisa," *Free Rad. Biol. & Med.*, Vol. 18(6), pp. 1023-1032 (1995)). In the article "Radial Immunoassay of 7, 8-Dihydro-8 Oxoadenine (8-Hydroxyadenine)," West et al. (1982), the authors discuss a specific radioimmunoassay for 8 hydroxy-adenine, and its application in the study of irradiated adenine solutions as well as a preliminary measurements of the production of 8-hydroxyadenine, using polyclonal antibodies.

[0013] Although detection and quantitation of oxidized nucleoside basis by immunoassay is gaining popularity, the challenge is in producing a suitable antibody for a specific base. It is preferable to have a specific and highly sensitive antibody which exhibits little cross-reactivity with related molecules. Polyclonal antibodies lack specificity and assays using such antibodies lack sensitivity. Therefore, use of monoclonal antibodies are preferred. However, in practice, obtaining suitable monoclonal antibodies can be difficult. The technique of producing monoclonal antibodies by hybridoma technology is well known in the art. Nevertheless, the results obtained by this technique are unpredictable. Only by carrying out the process for making the monoclonal antibodies can the nature of the, monoclonal antibodies be determined and ascertained. To the best of the inventors

knowledge, no assay has been developed for the detection and quantitation of 8-OH-Ade in a biological specimen using monoclonal antibodies.

[0014] Therefore, what is needed in the art is a highly sensitive and specific assay for detecting the presence of and quantitating the amount of 8-OH-Ade present in a biological specimen.

SUMMARY OF THE INVENTION

[0015] The present invention relates to methods and materials for the detection and quantitation 8-OH-Ade in biological specimens. Specifically, the present invention is directed to a group of highly specific monoclonal antibodies reactive with the modified nucleoside structure 8-OH-Ade, and to various immunoassays for 8-OH-Ade utilizing these monoclonal antibodies. The monoclonal antibodies of the present invention may be used in assays for diagnosing or monitoring the progression of certain types of cancer, in addition to a variety of other diseases associated with mutagenesis resulting from oxidative damage of DNA. Assays utilizing the monoclonal antibodies of the present invention may also be used to analyze or monitor toxicant exposure, such as from environmental sources. The monoclonal antibodies of the present invention may also be used to detect and quantitate epitopes of 8-OH-Ade.

[0016] The monoclonal antibodies of the present invention were prepared with the immunogen 8-OH-adenosine coupled to keyhole limpet hemocyanin (KLH). It is believed that the monoclonal antibodies bind with the base portion of the structure (8-OH-Ade) and not the carbohydrate (ribose) or protein linkage region of the conjugate, because, as demonstrated, conjugates bound to nucleosides other than 8-OH-adenosine were unreactive with these antibodies. Therefore, the antibodies of the present invention can be used to detect and quantitate (by the use of a standard curve) the presence of 8-OH-Ade in biological specimens of DNA. Procedures for such an assay include, for example, immobilizing the DNA, denaturing it to disrupt the base-pairing scheme exposing the free base structures, and quantitating the amount of 8-OH-Ade present per amount of DNA in a quantitative immunoassay.

[0017] Thus, it is an object of the present invention to develop an immunoassay for 8-OH-Ade, and its epitopes, in a biological specimen.

[0018] It is another object of the present invention to develop monoclonal antibodies specific against 8-OH-Ade.

[0019] Yet another object of the present invention is to develop an immunoassay highly sensitive and specific for 8-OH-Ade, utilizing monoclonal antibodies against 8-OH-Ade.

[0020] It is a further object of the present invention to prepare antibody producing hybridoma cells characterized by their production of monoclonal antibodies against 8-OH-Ade.

[0021] Yet another object of the present is to provide an immunoassay for diagnosing diseases associated with oxidative DNA damage resulting in 8-OH-Ade formation.

[0022] It is a further object of the present invention to provide an immunoassay for diagnosing and monitoring the development of cancer in a patient.

[0023] Another object of the present invention is to develop an immunoassay for monitoring a patient's response to treatment for diseases associated with oxidative DNA damage resulting in 8-OH-Ade formation.

[0024] A further object of the present invention is to develop an immunoassay for monitoring the effects of toxicant exposure in a human or animal.

[0025] It is also an object of the present invention to develop an immunoassay for detecting toxicant exposure in a human or animal.

[0026] Other aspects and advantages of the present invention will be apparent upon consideration of the following detailed description.

BRIEF DESCRIPTION OF THE DRAWINGS

[0027] The foregoing aspects and many of the attendant advantages of this invention will become more readily appreciated as the same becomes better understood by reference to the following detailed description, when taken in conjunction with the accompanying drawings, wherein:

[0028] **FIGS. 1A and 1B** are graphs showing the GC-MS/SIM analysis of synthetic 8-OH-adenosine after hydrolysis using 60% formic acid and triethylsililation. **FIG. 1A** is a total ion chromatogram of the trimethylsililated derivative. The major peak eluted at 9.7 min from an HP-1 GC column. **FIG. 1B** is a mass spectrum of the major peak eluting at 9.7 min. The major ions at m/z 352 and 367, characteristic of 8-OH-Ade, appear in mass spectrum.

[0029] **FIG. 2** is a graph of the results of a solid phase immunoassay of antibody binding specificity by analysis of antibody binding to periodate oxidized nucleosides conjugated to BSA for the monoclonal antibody 8A1 of the present invention. The conditions of the assay were as described in Example 4. The protein conjugates were coated onto the plastic surface in 50 μ l of volume at a starting concentration of 50 μ g/ml and serially diluted 1:2 as indicated. The haptens conjugated to BSA were as indicated in the FIGURE legend.

[0030] **FIG. 3** is a graph of the results of a solid phase immunoassay of antibody binding specificity by analysis of antibody binding to periodate oxidized nucleosides conjugated to BSA for the monoclonal antibody 8A2 of the present invention. The conditions of the assay were as described in Example 4. The protein conjugates were coated onto the plastic surface in 50 μ l of volume at a starting concentration of 50 μ g/ml and serially diluted 1:2 as indicated. The haptens conjugated to BSA were as indicated in the FIGURE legend.

[0031] **FIG. 4** is a graph of the results of a solid phase immunoassay of antibody binding specificity by analysis of antibody binding to periodate oxidized nucleosides conjugated to BSA for the monoclonal antibody 8A3.E10 of the present invention. The conditions of the assay were as described in Example 4. The protein conjugates, were coated onto the plastic surface in 50 μ l of volume at a starting concentration of 50 μ g/ml and serially diluted 1:2 as indicated. The haptens conjugated to BSA were as indicated in the FIGURE legend.

[0032] **FIG. 5** is a graph of the results of a solid phase immunoassay of antibody binding specificity by analysis of

antibody binding to periodate oxidized nucleosides conjugated to BSA for the monoclonal antibody 8A3.E11 of the present invention. The conditions of the assay were as described in Example 4. The protein conjugates were coated onto the plastic surface in 50 μ l of volume at a starting concentration of 50 μ g/ml and serially diluted 1:2 as indicated. The haptens conjugated to BSA were as indicated in the FIGURE legend.

[0033] **FIG. 6** is a graph of the results of a solid phase immunoassay of antibody binding specificity by analysis of antibody binding to periodate oxidized nucleosides conjugated to BSA for the monoclonal antibody 8A4.B7 of the present invention. The conditions of the assay were as described in Example 4. The protein conjugates were coated onto the plastic surface in 50 μ l of volume at a starting concentration of 50 μ g/ml and serially diluted 1:2 as indicated. The haptens conjugated to BSA were as indicated in the FIGURE legend.

[0034] **FIG. 7** is a graph of the results of a solid phase immunoassay of antibody binding specificity by analysis of antibody binding to periodate oxidized nucleosides conjugated to BSA for the monoclonal antibody 8A4.G10 of the present invention. The conditions of the assay were as described in Example 4. The protein conjugates were coated onto the plastic surface in 50 μ l of volume at a starting concentration of 50 μ g/ml and serially diluted 1:2 as indicated. The haptens conjugated to BSA were as indicated in the FIGURE legend.

[0035] **FIG. 8** is a graph of the results of a solid phase immunoassay of antibody binding specificity by analysis of antibody binding to periodate oxidized nucleosides conjugated to BSA for the monoclonal antibody 8A5 of the present invention. The conditions of the assay were as described in Example 4. The protein conjugates were coated onto the plastic surface in 50 μ l of volume at a starting concentration of 50 μ g/ml and serially diluted 1:2 as indicated. The haptens conjugated to BSA were as indicated in the FIGURE legend.

[0036] **FIG. 9** is a graph of the results of a solid phase immunoassay of antibody binding specificity by analysis of antibody binding to periodate oxidized nucleosides conjugated to BSA for monoclonal antibody 8A6 of the present invention. The conditions of the assay were as described in Example 4. The protein conjugates were coated onto the plastic surface in 50 μ l of volume at a starting concentration of 50 μ g/ml and serially diluted 1:2 as indicated. The haptens conjugated to BSA were as indicated in the FIGURE legend.

[0037] **FIG. 10** is a graph of the results of a solid phase immunoassay of antibody binding specificity by analysis of antibody binding to periodate oxidized nucleosides conjugated to BSA for the monoclonal antibody 8A7 of the present invention. The conditions of the assay were as described in Example 4. The protein conjugates were coated onto the plastic surface in 50 μ l of volume at a starting concentration of 50 μ g/ml and serially diluted 1:2 as indicated. The haptens conjugated to BSA were as indicated in the FIGURE legend.

[0038] **FIG. 11** is a graph of the results of a solid phase immunoassay of antibody binding specificity by analysis of antibody binding to periodate oxidized nucleosides conjugated to BSA for the monoclonal antibody 8A8 of the

present invention. The conditions of the assay were as described in Example 4. The protein conjugates were coated onto the plastic surface in 50 μ l of volume at a starting concentration of 50 μ g/ml and serially diluted 1:2 as indicated. The haptens conjugated to BSA were as indicated in the FIGURE legend.

[0039] **FIG. 12** is a graph of the results of a solid phase immunoassay of antibody binding specificity by analysis of antibody binding to periodate oxidized nucleosides conjugated to BSA for the monoclonal antibody 8A9. The conditions of the assay were as described in Example 4. The protein conjugates were coated onto the plastic surface in 50 μ l of volume at a starting concentration of 50 μ g/ml and serially diluted 1:2 as indicated. The haptens conjugated to BSA were as indicated in the FIGURE legend.

[0040] **FIG. 13** is a graph showing inhibition of antibody binding to 8-OH-adenosine-BSA conjugates by soluble ligands for the monoclonal antibody 8A1 of the present invention. The data shows results of inhibition of antibody binding to 8-OH-adenosine conjugated BSA (50 μ l of 50 μ g of protein/ml as coating mixture) in a solid phase assay using commercial 8-OH-AMP (Sigma, St. Louis, Mo.) (prepared from 3',5'cyclic 8-OH-ANT after hydrolysis by nucleotidase), AMP, or 4,6-diamino-5-formamidopyrimidine (Fapy-A) as shown in the FIGURE legend. The antibody present in culture supernatant was incubated in the presence of the indicated final concentration of soluble inhibitor overnight prior to the solid phase assay.

[0041] **FIG. 14** is a graph showing inhibition of antibody binding to 8-OH-adenosine-BSA conjugates by soluble ligands for the monoclonal antibody 8A2 of the present invention. The data shows results of inhibition of antibody binding to 8-OH-adenosine conjugated BSA (50 μ l of 50 μ g of protein/ml as coating mixture) in a solid phase assay using commercial 8-OH-AMP (Sigma, St. Louis, Mo.) (prepared from 3',5'cyclic 8-OH-AMP after hydrolysis by nucleotidase), AMP, or 4,6-diamino-5-formamidopyrimidine (Fapy-A) as shown in the FIGURE legend. The antibody present in culture supernatant was incubated in the presence of the indicated final concentration of soluble inhibitor overnight prior to the solid phase assay.

[0042] **FIG. 15** is a graph showing inhibition of antibody binding to 8-OH-adenosine-BSA conjugates by soluble ligands for the monoclonal antibody 8A3.E10 of the present invention. The data shows results of inhibition of antibody binding to 8-OH-adenosine conjugated BSA (50 μ l of 50 μ g of protein/ml as coating mixture) in a solid phase assay using commercial 8-OH-AMP (Sigma, St. Louis, Mo.) (prepared from 3',5'cyclic 8-OH-AMP after hydrolysis by nucleotidase), AMP, or 4,6-diamino-5-formamidopyrimidine (Fapy-A) as shown in the FIGURE legend. The antibody present in culture supernatant was incubated in the presence of the indicated final concentration of soluble inhibitor overnight prior to the solid phase assay.

[0043] **FIG. 16** is a graph showing inhibition of antibody binding to 8-OH-adenosine-BSA conjugates by soluble ligands for the monoclonal antibody 8A3.E11. The data shows results of inhibition of antibody binding to 8-OH-adenosine conjugated BSA (50 μ l of 50 μ g of protein/ml as coating mixture) in a solid phase assay using commercial 8-OH-AMP (Sigma, St. Louis, Mo.) (prepared from 3',5'cyclic 8-OH-AMP after hydrolysis by nucleotidase), AMP, or

4,6-diamino-5-formamidopyrimidine (Fapy-A) as shown in the FIGURE legend. The antibody present in culture supernatant was incubated in the presence of the indicated final concentration of soluble inhibitor overnight prior to the solid phase assay.

[0044] **FIG. 17** is a graph showing inhibition of antibody binding to 8-OH-adenosine-BSA conjugates by soluble ligands for the monoclonal antibody 8A4.B7 of the present invention. The data shows results of inhibition of antibody binding to 8-OH-adenosine conjugated BSA (50 μ l of 50 μ g of protein/ml as coating mixture) in a solid phase assay using commercial 8-OH-AMP (Sigma, St. Louis, Mo.) (prepared from 3',5'cyclic 8-OH-AMP after hydrolysis by nucleotidase), AMP, or 4,6-diamino-5-formamidopyrimidine (Fapy-A) as shown in the FIGURE legend. The antibody present in culture supernatant was incubated in the presence of the indicated final concentration of soluble inhibitor overnight prior to the solid phase assay.

[0045] **FIG. 18** is a graph showing inhibition of antibody binding to 8-OH-adenosine-BSA conjugates by soluble ligands for the monoclonal antibody 8A4.G10 of the present invention. The data shows results of inhibition of antibody binding to 8-OH-adenosine conjugated BSA (50 μ l of 50 μ g of protein/ml as coating mixture) in a solid phase assay using commercial 8-OH-AMP (Sigma, St. Louis, Mo.) (prepared from 3',5'cyclic 8-OH-AMP after hydrolysis by nucleotidase), AMP, or 4,6-diamino-5-formamidopyrimidine (Fapy-A) as shown in the FIGURE legend. The antibody present in culture supernatant was incubated in the presence of the indicated final concentration of soluble inhibitor overnight prior to the solid phase assay.

[0046] **FIG. 19** is a graph showing inhibition of antibody binding to 8-OH-adenosine-BSA conjugates by soluble ligands for monoclonal antibody 8A5 of the present invention. The data shows results of inhibition of antibody binding to 8-OH-adenosine conjugated BSA (50 μ l of 50 μ g of protein/ml as coating mixture) in a solid phase assay using commercial 8-OH-AMP (Sigma, St. Louis, Mo.) (prepared from 3',5'cyclic 8-OH-AMP after hydrolysis by nucleotidase), AMP, or 4,6-diamino-5-formamidopyrimidine (Fapy-A) as shown in the FIGURE legend. The antibody present in culture supernatant was incubated in the presence of the indicated final concentration of soluble inhibitor overnight prior to the solid phase assay.

[0047] **FIG. 20** is a graph showing inhibition of antibody binding to 8-OH-adenosine-BSA conjugates by soluble ligands for the monoclonal antibody 8A6 of the present invention. The data shows results of inhibition of antibody binding to 8-OH-adenosine conjugated BSA (50 μ l of 50 μ g of protein/ml as coating mixture) in a solid phase assay using commercial 8-OH-AMP (Sigma, St. Louis, Mo.) (prepared from 3',5'cyclic 8-OH-AMP after hydrolysis by nucleotidase), AMP, or 4,6-diamino-5-formamidopyrimidine (Fapy-A) as shown in the FIGURE legend. The antibody present in culture supernatant was incubated in the presence of the indicated final concentration of soluble inhibitor overnight prior to the solid phase assay.

[0048] **FIG. 21** is a graph showing inhibition of antibody binding to 8-OH-adenosine-BSA conjugates by soluble ligands for the monoclonal antibody 8A7 of the present invention. The data shows results of inhibition of antibody binding to 8-OH-adenosine conjugated BSA (50 μ g of 50 μ g

of protein/ml as coating mixture) in a solid phase assay using commercial 8-OH-AMP (Sigma, St. Louis, Mo.) (prepared from 3',5'cyclic 8-OH-AMP after hydrolysis by nucleotidase), AMP, or 4,6-diamino-5-formamidopyrimidine (Fapy-A) as shown in the FIGURE legend. The antibody present in culture supernatant, was incubated in the presence of the indicated final concentration of soluble inhibitor overnight prior to the solid phase assay.

[0049] FIG. 22 is a graph showing inhibition of antibody binding to 8-OH-adenosine-BSA conjugates by soluble ligands or the monoclonal antibody 8A8 of the present invention. The data shows results of inhibition of antibody binding to 8-OH-adenosine conjugated BSA (50 μ l of 50 μ g of protein/ml as coating mixture) in a solid phase assay using commercial 8-OH-AMP (Sigma, St. Louis, Mo.) (prepared from 3',5'cyclic 8-OH-AMP after hydrolysis by nucleotidase), AMP, or 4,6-diamino-5-formamidopyrimidine (Fapy-A) as shown in the FIGURE legend. The antibody present in culture supernatant was incubated in the presence of the indicated final concentration of soluble inhibitor overnight prior to the solid phase assay.

[0050] FIG. 23 is a graph inhibition of antibody binding to 8-OH-adenosine-BSA conjugates by soluble ligands for the monoclonal antibody 8A9 of the present invention. The data shows results of inhibition of antibody binding to 8-OH-adenosine conjugated BSA (50 μ l of 50 μ g of protein/ml as coating mixture) in a solid phase assay using commercial 8-OH-AMP (Sigma, St. Louis, Mo.) (prepared from 3',5'cyclic 8-OH-AMP after hydrolysis by nucleotidase), AMP, or 4,6-diamino-5-formamidopyrimidine (Fapy-A) as shown in the FIGURE legend. The antibody present in culture supernatant was incubated in the presence of the indicated final concentration of soluble inhibitor overnight prior to the solid phase assay.

[0051] FIG. 24 is a sample antibody titration curve for determining dilution of antibody to be used in evaluating the ability of soluble ligands to inhibit antibody binding to immobilized 8-OH-adenosine conjugated BSA. The data shown is for antibody 8A2. The antibody dilution to be used based on this data was 1:32.

[0052] FIG. 25 is a graph showing inhibition of antibody binding to 8-OH-adenosine-BSA conjugates by soluble ligands for monoclonal antibody 8A6 of the present invention. The data shows results of inhibition of antibody binding to 8-OH-adenosine conjugated BSA (50 μ l of 50 μ g of protein/ml as coating mixture) in a solid phase assay using commercial 8-OH-AMP (Sigma, St. Louis, Mo.) (prepared from 3',5'cyclic 8-OH-AMP after hydrolysis by nucleotidase). The antibody present in culture supernatant was diluted 1:12 and incubated in the presence of the indicated final concentration of soluble inhibitor overnight prior to the solid phase assay.

[0053] FIG. 26 is an inhibition of antibody binding to 8-OH-adenosine-BSA conjugates by soluble ligands for monoclonal antibody 8A6 of the present invention. The data shows results of inhibition of antibody binding to 8-OH-adenosine conjugated BSA (50 μ l of 1.56 μ g of protein/ml as coating mixture) in a solid phase assay using commercial 8-OH-AMP (Sigma, St. Louis, Mo.) (prepared from 3',5'cyclic 8-OH-AMP after hydrolysis by nucleotidase). The antibody present in culture supernatant was diluted 1:12 and incubated in the presence of the indicated final concentration of soluble inhibitor overnight prior to the solid phase assay.

[0054] FIG. 27 is a graph showing inhibition of antibody binding to 8-OH-adenosine-BSA conjugates by soluble ligands for the monoclonal antibody 8A9 of the present invention. The data shows results of inhibition of antibody binding to 8-OH-adenosine conjugated BSA (50 μ l of 50 μ g of protein/ml as coating mixture) in a solid phase assay using commercial 8-OH-AMP (Sigma, St. Louis, Mo.) (prepared from 3',5'cyclic 8-OH-AMP after hydrolysis by nucleotidase). The antibody present in culture supernatant was diluted 1:25 and incubated in the presence of the indicated final concentration of soluble inhibitor overnight prior to the solid phase assay.

[0055] FIG. 28 is a graph showing inhibition of antibody binding to 8-OH-adenosine-BSA conjugates by soluble ligands by the monoclonal antibody 8A9 of the present invention. The data shows results of inhibition of antibody binding to 8-OH-adenosine conjugated BSA (50 μ l of 0.78 μ g of protein/ml as coating mixture) in a solid phase assay using commercial 8-OH-AMP (Sigma, St. Louis, Mo.) (prepared from 3',5'cyclic 8-OH-AMP after hydrolysis by nucleotidase). The antibody present in culture supernatant was diluted 1:25 and incubated in the presence of the indicated final concentration of soluble inhibitor overnight prior to the solid phase assay. The results are shown for antibody 8A9.

DETAILED DESCRIPTION OF THE INVENTION

[0056] As used herein, the term "8-OH-Ade" refers to both bound, e.g., incorporated into DNA or RNA, and free forms of the nucleotide base.

[0057] The present invention is directed to materials and methods for detecting 8-OH-Ade in a sample of nucleic acids obtained from a biological specimen using monoclonal antibodies. In one aspect of the present invention, monoclonal antibodies are provided that are characterized by their specific reactivity with 8-OH-Ade. Representative embodiments of this aspect of the invention are the monoclonal antibodies identified below, produced by the hybridomas also identified below:

TABLE I

Monoclonal Antibodies of the Present Invention		
Antibody	Hybridoma	Accession No.
8A1	8A1	
8A2	8A2	
8A3.E10	8A13.E10	
8A3.E11	8A3.E11	
8A4.B7	8A4.B7	
8A4.G10	8A4.G10	
8A5	8A5	
8A6	8A6	ATCCHB12189
8A7	8A7	
8A8	8A8	
8A9	8A9	ATCCHB12188

[0058] The hybridoma cells 8A6 and 8A9 were deposited with American Type Culture Collection located at 12301 Parklawn Drive, Rockville, Md. 20852 on Sep. 13, 1996 and assigned the accession numbers indicated above.

[0059] The preferred monoclonal antibodies for use in the assays of the present invention are antibodies 8A6 and 8A9, identified above. . Antibody 8A5 is reactive primarily with

8-OH-Ade, but was also identified as reactive with 8-OH-guanine. Thus, antibody 8A5 may be useful in detecting 8-OH-adducts of purines generally.

[0060] Monoclonal antibodies of this invention can be prepared according to conventional methods by using 8-OH-adenosine conjugated to a carrier protein as an immunogen, as described in *Current Protocols in Immunology*, John Wiley & Sons, Inc. New York, N.Y. (1994), incorporated herein by reference. The synthesis of 8-OH-adenosine from 8-BR-adenosine is described in Cho et al., "Structure of Oxidatively Damaged Nucleic Acid Adducts. 3 Tautomerism, Ionization and Protonation of 8-Hydroxy-adenosine Studied by ¹⁵NMR Spectroscopy," *Nucleic Acids Research*, Vol. 19, pp. 1041-1047 (1991), West et al. (1982) and in Example 1 herein. **FIG. 1** shows results of GC-MS/SIM analysis of synthetic 8-OH-adenosine prepared in accordance with the method of the present invention after hydrolysis using 60% formic acid and trimethylsilylation. The product elutes as a major peak with a retention time of 9.7 min. There are only minor additional peaks in the total ion chromatogram, indicating high purity of this material. The retention time and mass spectrum of the synthetic 8-OH-adenosine are indistinguishable from a pure standard 8-OH-Ade, and monoclonal antibodies against 8-OH-adenosine react virtually equivalently against 8-OH-Ade.

[0061] To prepare the immunogen the 8-OH-adenosine product can be readily coupled to carrier proteins through the available amino groups by methods known in the art, although other types of antigen carrying molecules may be used. In a preferred embodiment of the present invention, 8-OH-adenosine is coupled to keyhole limpet hemocyanin (KLH) by Schiff base formation with the lysine groups of KLH, followed by reduction with NaCNBH₃. KLH is a preferred conjugation protein for such a coupling reaction since it aids in stimulating an immune response from attached ligands. Harlow et al., *Antibodies*, Cold Spring Harbor Laboratory (1988). Because of this, such an antigen is not useful for hybridoma screening. Other antigen carrier molecules which may be suitable for practicing the present invention include, but are not limited to BSA, ovalbumin, nucleic acids, lipids, carbohydrates, and naturally occurring biological conjugates such as glucuronic acid conjugates.

[0062] Immunization may be carried out according to conventional methods well known to those skilled in the art, such as by subcutaneously, intravenously or intraperitoneally injecting the 8-OH-adenosine conjugated to a carrier protein into an animal. More specifically, the immunogen may be diluted with PBS or physiological saline to a suitable concentration, and then injected into the animal, together with a suitable adjuvant if necessary. The immunogen should be injected several times (3 to 5 times) at an interval of 7 to 10 days with 50 to 100 μ g of immunogen in 0.1 ml total volume until the total volume injected reaches 100 μ l per animal. A conventional carrier may be used for the injection. Spleen cells isolated from the animal three days after the completion of the injection of the 8-OH-adenine conjugate are desirable for use as immune cells.

[0063] In a preferred embodiment to the present invention balb/c mice were injected subcutaneously in multiple sites with 8-OH-adenosine-KLH conjugate, prepared by mixing the immunogen with PBS and Freund's incomplete adjuvant followed by emulsification. The mice were immunized four

times at ten day intervals with a total volume of 100 μ l of immunogen per animal. Three days after fusion, spleen cells were then isolated for use as immune cells.

[0064] The transformed mammalian cells immunized by an 8-OH-adenosine conjugate are then fused with mammal plasmacytoma to produce hybridomas. A clone recognizing 8-OH-adenosine is selected from the hybridomas and the target monoclonal antibody is then obtained from the clone. In the above process there are few limitations to the mammal cells to be transformed with the immune antigen. It is desired that the immune antigen be selected taking its compatibility with the mammal plasmacytoma be fused into consideration. Mice, rats, rabbits and the like are generally preferable for use.

[0065] Various known myeloma cells can be used as mammal plasmacytoma to be fused with the above immune cells. Such myeloma cells include, for example, p3 (pe/x63-Ag8) (*Nature*, 256:495-497 (1975)), P3-U1 (*Current Topics of Microbiology and Immunology*, 81:1-7 (1987)), NS-1 (*Eur. J. Immunol.*, 6:511-519 (1976)), MPC-11 (*Cell*, 8405-415 (1976)), PS2/0 (*Nature*, 276:269-270 (1978)), FO (*J. Immunol. Meth.*, 35:1-21 (1980)), x63, 6, 5, 3 (*J. Immunol.*, 123:1548-1550 (1979)), S194 (*J. Exp. Med.*, 148:313-323 (1978)), and R210 (*Nature*, 277:131-133 (1979)) of rat, and the like. In a preferred embodiment of the present invention, mouse X63 myeloma cells are used.

[0066] The fusion of the immune cell and the plasmacytoma can be carried out in accordance with known methods, (see Harlow et al. (1988)) in the presence of a fusion accelerator and in a conventional nutritious medium. Conventional fusion accelerators, such as polyethylene glycol (PEG) and sendai virus (HVJ) can be used. Optionally, adjuvants such as dimethylsulfoxide and the like may be used in order to promote the efficiency of the fusion. A conventional fusion ratio of about 1-10 immune cells per one plasmacytoma may be used. As a medium for the fusion, any medium used for the cultivation of the plasmacytoma, such as PRMI 1640 medium and MEM medium, as well as other various media used for the cultivation of this type of cell, can be used. Serum obtained by removing serum complement from fetal calf serum (FCS) is a typical example of the type of medium that may be used.

[0067] The fusion is carried out by thoroughly mixing a prescribed amount of the immune cells with the plasmacytoma and blending this mixture with a medium to which about 30-60% (w/v) of a PEG (e.g., PEG with an average molecular weight of 1,000-6,000) solution which has been heated to about 37° C. in advance is added. The cultivation in the HAT medium is continued for a period sufficient for cells other than hybridoma (such as unfused cells) to die, usually for several days to several weeks. The hybridoma obtained is then subjected to a conventional limiting dilution method to detect the target cell lines producing the antibody of interest. In a preferred method of the present invention, the hybridoma is subjected to limited dilution cloning containing 8×10^5 mouse thymocytes as feeder cells per well. RPMI medium with 10% FCS and 1 mM pyrovalate and 2 mM glutamine thymocytes were used as feeder cells both for original fusion and in subcloning.

[0068] The detection of the antibody-producing cell lines of the present invention may be carried out according to standard methods commonly used for the detection of anti-

bodies, as described in the laboratory manual by Harlow et al. (1988) cited elsewhere herein, for example. Standard methods commonly used include the ELISA method, the plaque method, the spot method, the agglomeration reaction method, the Ouchterlony method, the radio immunoassay (RIA), and the like. Use of 8-OH-adenosine-conjugated BSA as an antigen for the detection is desirable.

[0069] In accordance with a preferred embodiment of the present invention, supernatant was tested for binding to 8-OH-adenosine conjugated BSA in a solid phase binding assay. The initial differential screening of the fusion was conducted with an 8-OH-adenosine-BSA conjugate versus an adenosine-BSA conjugate, along with BSA alone. Reactivity of 8-OH-adenosine selected clones from two independent fusions with alternate antigens was analyzed. The number of independent clones reactive with BSA coupled to one of three antigens is shown in Table II below:

TABLE II

Initial Differential Screening of Antibodies			
Fusion #	8-OH-adenosine/BSA	Adenosine/BSA	BSA Control
1	23	0	0
2	38	0	0

[0070] As demonstrated, the 8-OH-adenosine hapten was highly immunogenic by virtue of the considerable number of positive specific wells obtained in both fusions.

[0071] Hybridomas that showed reactive antibodies were further analyzed for binding specificity by comparing the reactivity of BSA conjugates linked to 8-OH-adenosine, native base structure, alternate oxidative products, irrelevant bases and oxidized products and a negative control. In accordance with the present invention and methods known in the art, antibody-producing cell lines were screened to obtain those cell lines that generate antibody having binding specificity for 8-OH-adenosine. See Example 4 and FIGS. 2 to 12. Hybridomas producing target monoclonal antibodies of the invention can be cultivated over generations in conventional media and can be stored in liquid nitrogen.

[0072] Collection of monoclonal antibodies of the present invention from hybridomas of the invention can be performed by cultivating the hybridoma according to conventional methods and obtaining the monoclonal antibody as a supernatant, or by administering the hybridoma to a mammal with which the hybridoma is compatible, allowing the hybridoma to proliferate, and collecting the desired antibodies from the ascites fluid. The former method is adaptable to the production of high purity monoclonal antibody, and the latter to mass production of monoclonal antibody; Monoclonal antibodies thus obtained may be purified by means of salting, gel filtration, affinity chromatography, or in accordance with other methods.

[0073] The monoclonal antibodies of the present invention are identified in Table I above. The isotype of each monoclonal antibody was determined by an Isostrip assay (Boehringer-Mannheim, Indianapolis, Ind.). The results were as shown in Table III below:

TABLE III

Antibody	Isotype
8A1	IgM
8A2	IgG1
8A3.E10	IgG1
8A3.E11	IgG1
8A4.B7	IgG1
8A4.G10	IgG1
8A5	IgG1
8A6	IgG1
8A7	IgM
8A8	IgG1
8A9	IgG1

[0074] The antibodies of the present invention can be used to detect and quantitate (by the use of a standard curve) the presence of 8-OH-adenine in biological specimens of DNA. Procedures for doing this would include immobilizing the DNA, denaturing it to disrupt the base-pairing scheme exposing the free base structures, and quantitating the amount of 8-OH-Ade present per amount of DNA in a quantitative immunoassay similar to those described below.

[0075] The presence of 8-OH-Ade in a biological sample can be analyzed at a high sensitivity and precision and with a high specificity in a simple manner by the use of monoclonal antibodies of the invention in conventional immunoassay formats, such as enzymatic immunoassays (EIA), enzyme-linked immunosorbent assays (ELISA), radioimmunoassays (RIA), immunoturbidimetric assays, or others known in the prior art. The lab manual by Harlow et al. (1988) discusses many of these methods. Because the monoclonal antibodies of the present invention react with 8-OH-Ade with specificity they are useful for the determination of 8-OH-Ade in clinical samples by immunoassay, thus enabling screening for various diseases and exposure to toxicants associated with elevated levels of 8-OH-Ade and associated with mutagenesis resulting from oxidative DNA damage. Thus, the present invention further provides immunoassay methods for determining the presence or amount of 8-OH-Ade in a biological fluid sample using the monoclonal antibodies of the invention. The assay comprises immunochemical reagents for forming an immunoreaction product whose presence or amount relates, either directly or indirectly, to the presence or amount of 8-OH-Ade in the sample. Those skilled in the art will appreciate that there are numerous well known clinical diagnostic procedures in which the immunochemical reagents of this invention can be used to form an immunoreaction product whose presence and/or amount relates to the presence and/or amount of 8-OH-Ade present in a sample. While exemplary assay methods are described herein, the invention is not so limited. Various heterogeneous and homogenous protocols, either competitive or noncompetitive, can be employed in performing an assay of this invention.

[0076] For example, the monoclonal antibodies of the present invention can be used in a direct solid phase immunoassay of antigen present in a biological specimen. DNA can be extracted from a tissue, cell or urine, for example, and subjected to a solid phase assay under conditions where the results with known amounts of DNA (e.g. by weight) are compared to a standard curve containing known amounts of antigen. This methodology could also be applied to impure

DNA fractions or unfractionated biological specimens such as tissue, cells, or bodily fluid and the results normalized to another parameter such as protein concentration or nucleic acid using alternate means for determining the amount of nucleic acid present in the specimen (e.g. the amount of adenine present).

[0077] Furthermore, the monoclonal antibodies of the present invention can be used in a quantitative immunohistochemical analysis of cells and tissues. For example, cells or tissue sections can be immobilized on glass slides under conditions which denature the cellular DNA, such as heating or drying the specimen. Antibody analysis can be conducted with, for example, fluorescently labeled 8-OH-Ade specific antibodies under conditions where the fluorescence intensity of the stained sections is proportional to the amount of 8-OH-Ade present in the specimen. A similar assay was previously described for monoclonal antibodies reactive with other types of modified nucleotide bases by Matsuda et al. (1993).

[0078] Alternatively, the 8-OH-Ade specific antibodies can be immobilized and used to absorb or capture soluble antigen from known amounts of biological specimens such as cells, tissue, and fluids, including bodily fluids. This can be used as a concentration step prior to elution and a detection and quantitation step using other methodologies such as HPLC-ECM (Shigenaga et al. (1989)) or GC-MS/SIM (Malins et al. (1993)) procedures. In addition, a detection and quantitation step involving inhibition of antibody binding to antigen as discussed below could be applied.

[0079] Furthermore, soluble antigen present in known amounts of biological specimens, including bodily fluids, can be detected and quantitated either directly or after an initial concentration step by determining the amount of this material required to provide inhibition of antibody binding to immobilized antigen. In these procedures, the specimen would be combined with monoclonal antibody of the present invention and incubated for a period of time sufficient to allow antibody complexes to form with the soluble antigen. The resulting mixture would be incubated with immobilized antigen and the amount of antibody binding to the immobilized antigen determined. The concentration of antigen present in the specimen would be determined by comparison to the effect with known amounts of 8-OH-Ade containing soluble fractions in either single determinations or in serial dilutions of the specimen. The dilution state required to relieve the inhibition of binding to the immobilized antigen to a proscribed level would be proportional to the concentration of 8-OH-Ade present in the specimen.

[0080] In another illustrative embodiment, a double antibody or "sandwich" immunoassay format may be employed comprising the steps of (a) forming a first immunoreaction admixture by admixing a sample with a first antibody, e.g., a monoclonal antibody, wherein the antibody and 8-OH-Ade present in the sample are capable of forming a first immunoreaction product (the first antibody can be operatively linked to a solid matrix); (b) maintaining the first immunoreaction admixture so formed under biological assay conditions for a time period sufficient to form the first immunoreaction product (the first immunoreaction product can then be separated from the sample); (c) forming a second immunoreaction admixture by admixing the first immunoreaction product with a second antibody, monoclonal or poly-

clonal, which recognizes 8-OH-Ade; (d) maintaining the second immunoreaction admixture so formed under biological assay conditions for a period sufficient to form the second or "sandwich" immunoreaction product; and (e) determining the presence and, optionally, the amount of second immunoreaction product formed, and thereby the presence and, optionally, the amount of 8-OH-Ade in the sample. Preferably, the second antibody is labeled, preferably with an enzyme, and thus the second immunoreaction product formed will be a labeled product to facilitate determination of the second immunoreaction product.

[0081] In preferred double antibody assay methods, the amount of immunoreaction product determined is related to the amount of immunoreaction product similarly formed and determined using a standard sample in place of the biological sample wherein the standard sample contains a known amount of 8-OH-Ade in accordance with this invention. Alternatively, a synthetic secondary standard can be used.

[0082] It is also preferred that the second antibody be directed to a site on the 8-OH-Ade which is not the same as the site to which the first antibody is directed. For example, the first antibody can be directed to a site other than that which reacts with the monoclonal antibodies of the present invention.

[0083] In any of the illustrative assays, the biological sample can be provided as a known or unknown quantity of urine, semen, seminal fluid, saliva, tissue, blood, or a blood derived product such as serum or plasma. Samples for study of oxidative DNA damage generally come from two main sources: urinary excretions of oxidized nucleosides and bases from DNA isolated target tissue or cells, such as lymphocytes. First, the DNA in the specimen must be immobilized, and then denatured to disrupt the base pairing scheme, exposing the tree base structures. The amount of antibody used can be known or unknown. The admixture is maintained under biological assay conditions for a predetermined period of from about 1 hour to about 16 hours at a temperature of from about 4° C. to about 37° C., most preferably about 22° C.

[0084] Biological assay conditions are those that maintain the biological activity of the immunochemical reagents of this invention and the 8-OH-Ade. Those conditions can generally include a temperature range of from about 4° C. to about 37° C., a pH value range of from at least about 6.0 to about 8.0, with a preferred range of 7.0 to 7.4, and an ionic strength varying from about 50 mM to 500 mM. Upon routine experimentation, other biological assay conditions may be learned. Methods for optimizing such conditions are well known to those skilled in the art.

[0085] Another assay format that is may be used in practicing the present invention is the precipitation assay. In this embodiment, the process comprises formation of an immunoreaction admixture by admixing a DNA sample obtained from a biological specimen with a monoclonal antibody of the invention to yield a precipitous immunoreaction product. The antibody can be operatively linked to a solid particulate such as a microparticle or bead, such that when antibody-antigen cross-linking occurs, the particulate matter aggregates, indicating the presence of the target material.

[0086] Another method is immunoturbidimetry because of its adaptability to automatic analysis, enabling a large num-

ber of samples to be measured at one time. Specifically, an amount of 8-OH-Ade in a sample of DNA obtained from urine, blood, or the like can be determined by adding one or more of the monoclonal antibodies of the present invention to the sample for the reaction and by measuring changes in the absorbance before and after the reaction.

[0087] Many other types of assays within the scope of this invention will be readily apparent to those skilled in the art.

[0088] Furthermore, the monoclonal antibodies of the present invention may form part of a kit comprising the monoclonal antibody of the invention and a means for detecting an immunoreaction product comprising 8-OH-Ade and the monoclonal antibody. Instructions for use of a packaged immunochemical reagent are also typically included in such a kit.

[0089] As used herein, the term "packaged" can refer to the use of a solid matrix or material such as glass, plastic, paper, fiber, foil and the like capable of holding within fixed limits an antibody of this invention. Thus, for example, a package can be a glass vial used to contain monoclonal milligram quantities of antibody of the present invention, or it can be a microliter plate well to which microgram quantities of a contemplated antibody has been operatively affixed. Alternatively, a package could include antibody-coated microparticles entrapped within a porous membrane or embedded in a test strip or dipstick, etc. Alternatively, the antibody can be directly coated onto a membrane, test strip or dipstick, etc. which contacts the sample fluid. Many other possibilities exist and will be readily recognized by those skilled in this art.

[0090] Instructions for use typically include a tangible expression describing the reagent concentration or at least one assay method parameter such as the relative amounts of reagent and sample to be admixed, maintenance time periods for reagent/sample admixtures, temperature, buffer conditions and the like.

[0091] In preferred embodiments, a diagnostic system of the present invention further includes a label or indicating means capable of signaling the formation of a complex containing an antibody of the present invention.

[0092] The word "complex" as used herein refers to the product of a specific binding reaction such as an antibody-antigen or receptor-ligand reaction. Exemplary complexes are immunoreaction products.

[0093] As used herein, the terms "label" and "indicating means" in their various grammatical forms refer to single atoms and molecules that are either directly or indirectly involved in the production of a detectable signal to indicate the presence of a complex. Any label or indicating means can be linked to or incorporated in an expressed protein, peptide, or antibody molecule that is part of the present invention, or used separately, and those atoms or molecules can be used alone or in conjunction with additional reagents. Such labels are themselves well known in the diagnostic art.

[0094] The labeling means can be a fluorescent labeling agent that chemically binds to antibodies or antigens without denaturing them to form a fluorochrome (dye) that is a useful immunofluorescent tracer. Suitable fluorescent labeling agents are fluorochromes such as fluorescein isocyanate (FIC), fluorescein isothiocyanate (FITC), 5-diethylamine-1-

natphthalenesulfonyl chloride (DANSC), tetramethylrhodamine isothiocyanate (TRITC), lissamine, rhodamine 8200 sulphonyl chloride (RB 200 SC) and the like. A description of immunofluorescence analysis techniques is found in DeLuca, "Immunofluorescence Analysis," *Antibody As a Tool*, Marchalonis et al., Eds., John Wiley & Sons, Ltd., pp. 189-231 (1982).

[0095] The indicating group may also be an enzyme such as horseradish peroxidase (HRP), glucose oxidase, or the like. In such cases where the principle indicating group is an enzyme such as HRP or glucose oxidase, additional reagents are required to indicate that a receptor-ligand complex (immunoreactant) has formed. Such additional reagents for HRP include hydrogen peroxide and an oxidation dye precursor such as diaminobenzidine. An additional reagent useful with glucose oxidase is 2,2'-azino-di-(3-ethyl-benzthiazoline-G-sulfonic acid) (ABTS).

[0096] Radioactive elements are also useful labeling agents and may be used in practicing the present invention. An exemplary radiolabeling agent is a radioactive element that produces gamma ray emissions. Elements which themselves emit gamma rays, such as ^{124}I , ^{125}I , ^{128}I , ^{132}I and ^{51}Cr represent one class of gamma ray emission-producing radioactive element indicating groups. Particularly preferred is ^{125}I . Another group of useful labeling means are those elements such as ^{14}C , ^{18}F , ^{15}O and ^{13}N which themselves emit positrons. Also useful is a beta emitter, such as ^{113}In or ^3H .

[0097] The linking of labels, i.e., labeling of peptides and proteins, is well known in the art. For instance, monoclonal antibodies produced by a hybridoma can be labeled by metabolic incorporation of radioisotope-containing amino acids provided as a component in the culture medium. See, for example, Galfre et al., *Meth. Enzymol.*, 73:3-46 (1981). The techniques of protein conjugation or coupling through activated functional groups are also applicable. See, for example, Aurameas et al., *Scand. J. Immunol.*, 8(7):7-23 (1978); Rodwell et al., *Biotech.*, 3:889-894 (1984); and U.S. Pat. No. 4,493,795.

[0098] The diagnostic test kit can also include, preferably as a separate package, a "specific binding agent," which is a molecular entity capable of selectively binding an antibody of this invention or a complex containing such a species, but is not itself an antibody of this invention. Exemplary specific binding agents are second antibody molecules, complement, proteins or fragments thereof. Preferably the specific binding agent binds the antibody when it is present as part of a complex.

[0099] In preferred embodiments, the specific binding agent is labeled. However, when the diagnostic system includes a -specific binding agent that is not labeled, the agent is typically used as an amplifying means or reagent. In these embodiments, the labeled specific binding agent is capable of specifically binding the amplifying means when the amplifying means is bound to a complex.

[0100] The diagnostic kits of the present invention can be used in an "ELISA" format to detect the quantity of 8-OH-Ade in biological samples of DNA obtained from biological specimens such as cells, plasma, saliva, serum, semen, seminal fluid tissue, urine or blood. "ELISA" refers to an enzyme linked immunosorbent assay such as those dis-

cussed above, which employ an antibody or antigen bound to a solid phase and an enzyme-antigen or enzyme-antibody conjugate to detect and quantify the amount of an antigen present in a sample. A description of the ELISA technique is found in Harlow et al. (1988).

[0101] Thus, in preferred embodiments a monoclonal antibody with inherent specificity for 8-OH-Ade can be affixed to a solid matrix to form a solid support. A reagent is typically affixed to a solid matrix by adsorption from an aqueous medium, although other modes of affixation applicable to proteins and peptides well known to those skilled in the art can be used.

[0102] Useful solid matrices are also well known in the art. Such materials are water insoluble and include the cross-linked dextran available under the trademark SEPHADEX (Pharmacia Fine Chemicals, Piscataway, N.J.); agarose; polystyrene beads about 1 micron to about 5 millimeters in diameter; polyvinyl chloride, polystyrene, cross-linked polyacrylamide, nitrocellulose- or nylon-based webs such as sheets, strips or paddles; or tubes, plates or the wells of a microtiter plate such as those made from polystyrene or polyvinylchloride.

[0103] The immunoreagents of any diagnostic system described herein can be provided in solution, as a liquid dispersion or as a substantially dry powder, e.g., in lyophilized form. Where the indicating means is an enzyme, the enzyme's substrate can also be provided in a separate package. A solid support such as the above-described microtiter plate and one or more buffers can also be included as separately packaged elements in the diagnostic assay systems of this invention.

[0104] The packaging materials discussed herein in relation to diagnostic systems are those customarily utilized in diagnostic systems. Such materials include glass and plastic (e.g., polyethylene, polypropylene and polycarbonate) bottles, vials, plastic and plastic-foil laminated envelopes and the like.

[0105] Thus, the monoclonal antibodies of the present invention may be used in a variety of immunoassays to detect and quantitate 8-OH-Ade in biological specimens. These assays may be useful for research, diagnosis of disease, prognosis and tracking of response to treatment. The following examples illustrate a preferred method for making the monoclonal antibodies of the present invention.

EXAMPLE 1

Preparation of 8-OH-adenosine

[0106] 8-OH-adenosine was prepared according to the method previously described in Cho et al.(1991). 8-Br-Adenosine (14 mmol) (Sigma, St. Louis, Mo.) was dissolved in 40 ml of dry DMSO and added to a mixture containing sodium benzyloxide. This reagent was made by reacting 1 g of sodium metal with 35 ml of benzyl alcohol in 100 ml of DMSO. The reaction mixture was heated at 65° C. for 24 hours, cooled to room temperature, acidified with glacial acetic acid, and poured into 1 L of anhydrous ether. Preparation of 8-OH-adenosine is a single step reaction due to the acid labile nature of the initial C8-benzyloxylation product which yields 8-OH-adenosine directly. The precipitate was collected, dissolved in methanol, and absorbed onto 5 g of

silica gel (Merck, Darmstadt, Germany). The solvent was removed on a rotary evaporator, and the residue applied to a 500 ml bed volume silica gel column in a solvent composed of $\text{CHCl}_3:\text{CH}_3\text{OH}$ (4:1). The column was eluted with this solvent and fractions were collected. The elution of products was monitored by thin layer chromatography and appropriate fractions pooled. The fractions which contained 8-OH-adenosine were identified after GC-MS/SIM analysis of the purified components using standard methodology as described in Malins et al. (1993). The product was recrystallized from water and yielded about 0.5 g of pure 8-OH-adenosine as a white powder.

EXAMPLE 2

Coupling of 8-OH-adenosine to Keyhole Limpet Hemocyanin (KLH) and Bovine Serum Albumin (BSA)

[0107] The 8-OH-adenosine product can be readily coupled to carrier proteins through available amino groups. Coupling of this hapten to carrier protein was conducted through the ribose moiety of the nucleoside after mild oxidation by NaIO_4 at pH 4.5 using a sodium phosphate buffer. The progress of the oxidation reaction was followed by the change in mobility of the UV-absorbing spots after thin layer chromatography on silica gel plates using a solvent system composed of $\text{CHCl}_3:\text{CH}_3\text{OH}$ (2:1). The oxidized product migrates as a faster moving spot on the chromatogram. This introduces vicinal aldehyde groups capable of forming Schiff bases with primary amines.

[0108] The antigen for immunization was prepared by Schiff base formation with lysine groups of keyhole limpet hemocyanin (KLH) followed by reduction with NaCNBH_3 . The antigen for hybridoma screening was prepared by coupling the IO_4^- oxidized derivative to BSA. This placed the specific ligand of interest on a molecule that was ideal for solid-phase immunoassays to be used in the hybridoma screening process.

[0109] Hapten conjugates of BSA using a variety of negative control nucleotide base structures were prepared using these same procedures with commercially available periodate oxidized nucleosides (Sigma, St. Louis, Mo.). Commercially obtained periodate oxidized nucleosides used were adenosine, cytosine, uracil, and guanosine. In addition, 8-OH-guanosine, which was prepared according to the method described by Cho et al., *Chem. Res. Toxicol.*, Vol. 3, pp. 445-452, (1990), was also periodate oxidized and coupled to BSA. All BSA conjugates containing approximately 15 hapten molecules per protein subunit.

EXAMPLE 3

Preparation of Anti-8-OH-Ade Antibodies

[0110] The methods employed were standard procedures as described in the laboratory manual Harlow et al. (1988). Balb/c mice were injected subcutaneously in multiple sites with 8-OH-adenosine conjugated keyhole limpet hemocyanin. The immunogen was prepared by mixing 2 mg of 8-OH-adenosine-KLH conjugate in 900 μl of phosphate buffered saline (PBS), 100 μl of 2 mg/ml MDP (N-acetylmuramy 1-L-alanyl-D-isoglutamine (Pierce, Rockford, Ill.), and 1 ml of Freund's incomplete adjuvant (Sigma, St. Louis, Mo.) followed by emulsification. The mice were immunized

4 times at 10 day intervals with 100 μ l of immunogen, the last 3 days before fusion. Spleen cells were isolated and fused with mouse X63 myeloma cells.

EXAMPLE 4

Solid Phase Immunoassay

[0111] Once clones of hybridoma cells appeared after HAT selection, supernatant was tested for binding to 8-OH-adenosine conjugated BSA in a solid phase binding assay on 96-well Probind plates. Wells containing reactive antibodies were moved to 24-well plates and expanded for more detailed analysis of binding specificity. Each reactive antibody was tested for specificity in side-by-side comparison of reactivity with BSA conjugates linked to the specific antigen, native base structure, alternate oxidative products, irrelevant bases and oxidized products, and a negative control. Wells showing antibody of proper specificity were selected and cloned to yield monoclonal antibody producing cell lines.

[0112] The 8-OH-adenosine-conjugated BSA was deposited on 96-well Probind plates by coating each well with 50 μ l of a solution containing 50 μ g protein conjugate per ml of 50 mM sodium phosphate buffer, pH 7.5, 5 mM MgCl₂, 15 mM NaN₃, and incubated overnight. The plates were blocked with PBS containing 5% BSA for 2 hours, followed by incubation with antibody containing culture supernatant for 18 hours. The plates were then washed extensively with PBS, followed by incubation with 1:500 diluted rabbit anti-mouse whole Ig (ICN Immunobiologicals, Costa Mesa, Calif.) for 1 hour. The plates were again extensively washed with PBS and incubated with ¹²⁵I-protein A (90,000 cpm/well) for 1 hour. The plates were washed again with PBS and the amount of ¹²⁵I in each well was determined in a gamma counter.

[0113] FIGS. 2 through 12 show results of solid phase immunoassays using senior dilutions of nucleoside-BSA conjugates and cloned hybridoma culture supernatants as the antibody source. The results indicate that antibodies 8A1, 8A2, 8A3.E10, 8A3.E11, 8A4.B7, 8A4.G10, 8A6, and 8A9 of the present invention were absolutely specific for 8-OH-adenosine haptens coupled to BSA. No cross reactivity was observed with BSA conjugated to either adenosine, cytosine, uracil, guanosine, or 8-OH-guanosine. A broader specificity was observed with 8A5 (FIG. 8) in which substantial reactivity with 8-OH-guanosine was also observed, and with 8A7 (FIG. 10) and 8A8 (FIG. 11) which also reacted with guanosine and adenosine, and 8-OH-guanosine, respectively.

[0114] In experiments to titrate antigen reactivity, the conjugate protein was serially diluted in the above buffer alone. As shown in FIGS. 2 to 12, the antibodies were capable of detecting very low amounts of 8-OH-Ade. In particular, antibodies 8A1, 8A2, 8A6, 8A7, 8A8 and 8A9 were reactive with 8-OH-Ade levels present in quantities of less than 5 fmol per assay well.

EXAMPLE 5

Inhibition of Antibody Binding by Soluble Ligands

[0115] Culture supernatants containing hybridomas of known titer were mixed with soluble ligands which were

serially diluted 1:2 in a 96-well plate starting with a concentration of 1 mM down to 5 μ M and incubated overnight at 4° C. followed by incubation for 2 hours on an 8-OH-adenosine-BSA antigen coated plate (50 μ l of 50 μ g/ml protein conjugate). Conditions of the assay were otherwise the same as described in Example 4 above.

[0116] FIGS. 13 to 23 show results of inhibition of antibody binding to 8-OH-adenosine conjugated BSA (50 μ g of protein/ml as coating mixture) in a solid phase assay using commercial 8-OH-AMP (Sigma, St. Louis, Mo.) (Prepared from 3',5'-cyclic 8-OH-AMP after hydrolysis by nucleotidase), AMP, or 4,6-diamino-5-formamidopyridine (Fapy-A). Fapy-A is a ring-opening oxidation product from single electron oxidations and is an alternate reaction product of the same reaction which leads to 8-OH-adenosine. In these experiments the titer of antibody present in the culture supernatant used was adjusted to yield near maximal, but not saturating levels of antibody binding so that the antibody titer was not in large excess over the antigen coated onto the plate. This is demonstrated in FIG. 24. The results indicate that in the cases of antibodies 8A1, 8A2, 8A3.E10, 8A3.E11, 8A4.B7, 8A4.G10, 8A6, 8A7, 8A8, and 8A9 specific inhibition of antibody binding was observed only for 8-OH-AMP as its concentration was varied from 1 mM to 5 μ M. No apparent inhibition with any antigen under the concentration range used was observed for 8A5. The concentration of soluble 8-OH-AMP capable of inhibiting the solid phase antibody binding was variable among the antibodies, with 8A6 and 8A9 showing the lowest concentrations of the soluble ligand capable of binding inhibition.

[0117] To determine if changes in experimental conditions can modulate the concentration of soluble ligand capable of causing significant binding inhibition, the ratio of soluble 8-OH-AMP ligand to immobilized 8-OH-adenosine present in BSA conjugates was varied and the effect on the concentration of soluble ligand capable of inhibiting antibody binding determined. The results for antibody 8A6 is shown in FIGS. 25 and 26. FIG. 25 shows results under standard conditions with plates coated with the BSA conjugate at a concentration of 50 μ g/ml compared to an amount 32-fold lower shown in FIG. 26. These results indicate that this change in the ratio of soluble to bound ligand reduced the concentration of soluble ligand needed to provide a similar amount of binding inhibition by a factor of about 16. A similar study was also conducted with antibody 8A9 as shown in FIGS. 27 and 28 with more striking results. The concentration of soluble ligand capable of providing the same amount of inhibition in comparisons of the standard condition (50 μ l of 50 μ g/ml protein coated per well) versus a 64-fold dilution (i.e. 50 μ l of 0.78 μ g/ml protein) was itself 64-fold lower. Thus, the ratio of bound vs. soluble antigen in the inhibition assay can be adjusted to provide a more sensitive assay and the assay can be adjusted to detect binding inhibition caused by very low concentrations of soluble antigen. This is done by reducing the amount of bound antigen used in the assay and amplifying the signal due to primary antibody binding to it so that low concentrations of soluble antigen can effectively competitively bind to the antigen binding site on the antibody and result in detectable binding inhibition. Such an assay can be used to determine the presence and concentration of soluble antigen present in biological specimens and bodily fluids including, but not limited to, blood or urine.

EXAMPLE 6

Binding of Antibodies 8A6 and 8A9 to
Physiological DNA Specimens

[0118] The ability of this panel of monoclonal antibodies to detect 8-OH-Ade lesions in physiological DNA was tested. For convenience, antibodies 8A6 and 8A9 were selected for this analysis. DNA specimens were analyzed for the levels of 8-OH-Ade present using GC-MS/SIM methodology (Malins et al. (1993)). Control DNA contained 1.2 8-OH-Ade lesions per 10^5 normal bases. Test DNA contained 4.7 8-OH-Ade lesions per 10^5 normal bases.

[0119] DNA specimens (50 ng) were dissolved in 100 μ l of 1 M ammonium acetate and slot blotted onto nitrocellulose membrane (0.45 micron) using a Minifold II slot-blot system (Schleicher & Schuell, Keene, N.H.) according to manufacturer's instructions. The blots were removed and heated in an oven at 80° C. for 1 hour to denature the DNA. The blots were blocked using 25 mM Tris-HCl, pH 7.4, 150 mM NaCl, 0.1% Tween-20, 5% non-fat dry milk overnight at room temperature, followed by treatment with culture supernatants containing monoclonal antibody for 3 hours at room temperature. The blots were washed extensively with blocking buffer and followed by incubation with 1:1000 diluted rabbit anti-mouse whole Ig secondary antibody (ICN, Costa Mesa, Calif.) for 1 hour at room temperature. This antibody was diluted in blocking buffer without milk. After extensive washing with blocking buffer, the blots were incubated with 125 I-Protein (20,000 cpm per 50 μ l) for 1 hour at room temperature. 125 I-Protein A was also diluted in blocking buffer without milk. After extensive washing with blocking buffer, the blots were dried and exposed to x-ray film overnight to visualize spots. The labeled spots were cut out and counted in a gamma counter. Net cpm 125 I-Protein A bound was calculated after subtraction of background. Background counts were determined by counting an equal area of the blot which contained no DNA.

TABLE IV

Analysis of Binding of Antibodies 8A6 and 8A9 to Denatured DNA Specimens of Known 8-OH-Ade Concentration				
DNA Specimen	# 8-OH-Ade 105 bases	fmol 8-OH-Ade/ assay	Net cpm 125 I-Protein A Bound	
			Antibody 8A6	Antibody 8A9
Control DNA	1.2	0.5	882	910
Test DNA	4.7	1.8	3671	4602

[0120] The data obtained is shown in Table IV below. The amount of 8-OH-Ade present at each blot is shown, as is the net cpm of 125 I-Protein A bound. This value is proportional to the amount of 8A6 or 8A9 antibody bound to the DNA on the blot. This data shows increased antibody binding to test DNA compared to the control. In addition, each antibody is capable of producing a clear signal above background when 8-OH-Ade is present in the range of 2 fmol/assay. Thus, these data confirm that the antibodies bind directly to physiological DNA and are capable of detecting low fmol quantities of 8-OH-Ade per assay.

[0121] In summary, the present invention comprises a panel of highly specific monoclonal antibodies reactive with

the modified nucleoside structure 8-OH-adenosine. The results of the inventors' experiments further define that it is the base portion of the structure (8-OH-Ade) and not the carbohydrate (ribose) or protein linkage region of the conjugate which is involved in antibody binding. This is known because conjugates with alternate nucleosides other than 8-OH-adenosine were unreactive with these antibodies, as demonstrated in Example 4 and FIGS. 2 to 12. In addition, direct binding of antibodies to 8-OH-Ade present in physiological DNA was demonstrated (see Example 6). In some instances minor specificities of certain antibodies were found (all of which correlate with antibody binding to the base structure). Of particular interest in this regard is antibody 8A5 which is reactive primarily with 8-OH-Ade but also detects 8-OH-guanosine. Also of interest are antibodies 8A7 and 8A8 which show some reactivity to guanosine. Thus, monoclonal antibody 8A5 in particular (or one similar to it), and antibodies 8A7 and 8A8 may be useful in detecting 8-OH-adducts of purines generally.

[0122] The results obtained by the inventors are typical for binding specificities of monoclonal antibodies. That is, specificity is defined by certain epitope portions of a larger molecule. Beyond the specific epitope, there is no recognition and no bias as to the chemical form containing the epitope for antigen presentation to the antibody. Similar results were demonstrated by the inventors for anti-carbohydrate antibodies (Holmes and Greene, Arch. Biochem. Biophys., 288, 87-96, 1991). In those instances, the aglycone portion carrying the antigen (e.g. from lipid to protein) was demonstrated to be irrelevant and antibody binding independent from it.

[0123] The base ligand structure as the specific epitope for the antibodies of the present invention is further supported by antibody binding inhibition studies with soluble ligands. Soluble 8-OH-adenosine containing ligands specifically inhibited solid phase antibody binding, as demonstrated by the results shown in FIGS. 13 to 23. Soluble 8-OH-AMP (a single nucleotide moiety component of nucleic acids) inhibited antibody binding but adenine (from AMP) or Fapy-A structures did not. Thus, the antibodies of the present invention recognized antigen presented by both phosphorylated native nucleotides and periodate oxidized nucleosides. This last condition is suitable to cause ribose ring opening and, after reduction of a Schiff base with a protein amino group, yields a very different and non-physiological structure. Given these results, the only possible conclusion is that the monoclonal antibodies of the present invention are specific for the base structure (8-OH-Ade) alone and are independent of the nature of the group or structure the base is coupled to. Therefore, these antibodies are useful in detecting and quantitating the amount of 8-OH-Ade present in biological specimens by virtue of their ability to specifically bind to this ligand.

[0124] The foregoing illustrative examples relate to the measurement of 8-OH-Ade in a biological sample using monoclonal antibodies. While the preferred embodiment of the present invention has been described in terms of specific conditions and format, it is understood that numerous variations and modifications will occur to those skilled in the art upon consideration of the present invention without departing from the spirit and scope of the invention.

The embodiments of the invention in which an exclusive property or privilege is claimed are defined as follows:

1. A monoclonal antibody that is capable of binding 8-OH-adenine.

2. The monoclonal antibody of claim 1, wherein the monoclonal antibody is obtained from a hybridoma selected from the group consisting of hybridomas ATCCHB12188 and ATCCHB12189.

3. A monoclonal antibody that is capable of binding 8-OH-adenosine or 8-OH-deoxyadenosine.

4. The monoclonal antibody of claim 3, wherein the monoclonal antibody is obtained from a hybridoma selected from the group consisting of hybridomas ATCCHB12188 and ATCCHB12189.

5. A hybridoma that produces a monoclonal antibody that is capable of binding 8-OH-adenine.

6. The hybridoma of claim 5, wherein the hybridoma is selected from the group consisting of hybridomas ATCCHB12189 and ATCCHB12189.

7. A hybridoma that produces a monoclonal antibody that is capable of binding 8-OH-adenosine or 8-OH-deoxyadenosine.

8. The hybridoma of claim 7, wherein the hybridoma is selected from the group consisting of hybridomas ATCCHB12188 and ATCCHB12189.

9. A method for determining the presence or amount of 8-OH-adenine in a biological specimen comprising the steps of contacting the sample with at least one monoclonal antibody that is capable of binding 8-OH-adenine to form a complex of monoclonal antibody and 8-OH-adenine, and determining the presence or amount of complex formed as an indication of the presence or amount of 8-OH-adenine in the sample.

10. The method of claim 9, wherein the monoclonal antibody is obtained from a hybridoma selected from the group consisting of hybridomas ATCCHB12188 and ATCCHB12189.

11. The method of claim 9, wherein the monoclonal antibody is obtained from the hybridoma ATCCHB12188.

12. The method of claim 10, wherein the monoclonal antibody is obtained from the hybridoma ATCCHB12189.

13. The method of claim 9, wherein the biological specimen is selected from the group consisting of cells, tissue, blood, saliva, serum, plasma, semen, seminal fluid and urine.

14. A method for diagnosing or monitoring the status of a disease associated with elevated levels of 8-OH-adenine in a human or animal comprising the steps of:

a. obtaining a biological specimen acids from the human or animal;

b. contacting the specimen with at least one monoclonal antibody that is capable of binding 8-OH-adenine to form a complex of monoclonal antibody and 8-OH-adenine;

c. determining the amount of complex formed as a measure of the presence or amount of 8-OH-adenine in the specimen.

15. The method of claim 14, wherein the disease is cancer.

16. The method of claim 14, wherein the monoclonal antibody is obtained from a hybridoma selected from the group consisting of hybridomas ATCCHB12188 and ATCCHB12189.

17. The method of claim 14, wherein the monoclonal antibody is obtained from the hybridoma ATCCHB12188.

18. The method of claim 14, wherein the monoclonal antibody is obtained from the hybridoma ATCCBB12189.

19. The method of claim 14, wherein the biological specimen obtained from the human or animal is selected from the group consisting of cells, tissue, blood, saliva, serum, plasma, semen, seminal fluid and urine.

20. The method of claim 14, wherein the determination of monoclonal antibody bound to 8-OH-adenine is determined by a method selected from the group consisting of ELISA, radioimmunoassay, photochemical assay and fluorescence assay.

21. A method for determining and monitoring exposure to a toxicant associated with elevated levels of 8-OH-adenine by a human or animal comprising the steps of:

a. obtaining a biological specimen from the human or animal;

b. contacting the specimen with at least one monoclonal antibody that is capable of binding 8-OH-adenine to form a complex of monoclonal antibody and 8-OH-adenine;

c. determining the amount of complex formed as an indication of the presence or amount of 8-OH-adenine in the specimen.

22. The method of claim 20, wherein the monoclonal antibody is obtained from a hybridoma selected from the group consisting of hybridomas ATCCHB12188 and ATCCHB12189.

23. The method of claim 20, wherein the monoclonal antibody is obtained from the hybridoma ATCCHB12188.

24. The method of claim 20, wherein the monoclonal antibody is obtained from the hybridoma ATCCHB12189.

25. The method of claim 20, wherein the biological specimen obtained from the human or animal is selected from the group consisting of cells, tissue, blood, saliva, serum, plasma, semen, seminal fluid and urine.

26. The method of claim 20, wherein the determination of monoclonal antibody bound to 8-OH-adenine is determined by a method selected from the group consisting of ELISA, radioimmunoassay, photochemical assay and fluorescence assay.

27. A method for determining the amount of 8-OH-adenine in a biological specimen comprising:

a. contacting a monoclonal antibody obtained from a hybridoma selected from the group consisting of hybridomas ATCCHB12188 and ATCCHB12189 with the sample;

b. quantitating the amount of monoclonal antibody bound to the specimen; and

c. comparing the amount of monoclonal antibody bound to the specimen to the amount of monoclonal antibody bound to a known standard.

28. The method of claim 25, wherein the biological specimen obtained from a human or animal is selected from the group consisting of cells, tissue, blood, saliva, serum, plasma, semen, seminal fluid and urine.

29. The method of claim 25, wherein the quantitation of monoclonal antibody bound to 8-OH-adenine is determined by a method selected from the group consisting of ELISA, radioimmunoassay, photochemical assay and fluorescence assay.

30. A method for determining the presence or amount of 8-OH-adenine in a biological specimen comprising:

- a. contacting the specimen with at least one monoclonal antibody obtained from a hybridoma selected from the group consisting of hybridomas ATCCHB12188 and ATCCHB12189 to form a first complex comprising the monoclonal antibody and 8-OH-adenine;
- b. contacting the specimen containing the first complex with immobilized 8-OH-adenine;
- c. quantitating the amount of monoclonal antibody bound to the immobilized 8-OH-adenine; and
- d. comparing the amount of monoclonal antibody bound to the immobilized 8-OH-adenine to the amount of monoclonal antibody bound to a known standard.

31. The method of claim 28, wherein the biological specimen obtained from a human or animal is selected from the group consisting of cells, tissue, blood, saliva, serum, plasma, semen, seminal fluid and urine.

32. The method of claim 28, wherein the quantitation of monoclonal antibody bound to immobilized 8-OH-adenine is determined by a method selected from the group consisting of ELISA, radioimmunoassay, photochemical assay and fluorescence assay.

33. A method for determining the presence or amount of 8-OH-adenine in a biological specimen comprising the steps of:

- a. contacting the specimen with at least one monoclonal antibody obtained from a hybridoma selected from the group consisting of hybridomas ATCCHB12188 and ATCCHB12189 to form a first complex comprising the monoclonal antibody and 8-OH-adenine;

- b. contacting the specimen containing the first complex with at least one second antibody capable of binding to the first complex;

- c. determining the amount of second complex formed as a measure of the presence or amount of 8-OH-adenine in the specimen.

34. The method of claim 31, wherein the biological specimen from a human or animal is selected from the group consisting of cells, tissue, blood, saliva, serum, plasma, semen, seminal fluid and urine.

35. A method for determining the presence or amount of an epitope of 8-OH-adenine in a biological specimen comprising the steps of contacting the sample with at least one monoclonal antibody that is capable of binding 8-OH-adenine to form a complex of monoclonal antibody and epitope, and determining the presence or amount of complex formed as a measure of the presence or amount of epitope in the specimen.

36. The method of claim 33, wherein the monoclonal antibody is obtained from a hybridoma selected from the group consisting of hybridomas ATCCHB12188 and ATCCHB12189.

37. The method of claim 33, wherein the monoclonal antibody is obtained from the hybridoma ATCCHB12188.

38. The method of claim 33 wherein the monoclonal antibody is obtained from the hybridoma ATCCHB12189.

39. The method of claim 33, wherein the biological specimen from a human or animal is selected from the group consisting of cells, tissue, blood, saliva, serum, plasma, semen, seminal fluid and urine.

* * * * *

专利名称(译)	使用单克隆抗体检测和定量8-OH-腺嘌呤		
公开(公告)号	US20030186344A1	公开(公告)日	2003-10-02
申请号	US10/375697	申请日	2003-02-25
[标]申请(专利权)人(译)	CYTOCHEM		
申请(专利权)人(译)	CYTOCHEM INC.		
当前申请(专利权)人(译)	CYTOCHEM INC.		
[标]发明人	HOLMES ERIC H GREENE THOMAS G		
发明人	HOLMES, ERIC H. GREENE, THOMAS G.		
IPC分类号	G01N33/53 G01N33/574 G01N33/537 G01N33/543 C12N5/06 C07K16/00		
CPC分类号	G01N33/57484 G01N33/5308		
优先权	09/043461 1998-03-18 US PCT/US1996/015034 1996-09-19 WO 60/003976 1995-09-19 US		
其他公开文献	US6900291		
外部链接	Espacenet USPTO		

摘要(译)

本发明涉及用于检测和定量生物样本中8-OH-Ade的方法和材料。具体地，本发明涉及一组与修饰的核苷结构8-OH-Ade反应的高度特异性单克隆抗体，以及利用这些单克隆抗体的8-OH-Ade的各种免疫测定。除了与由DNA的氧化损伤引起的诱变相关的多种其他疾病之外，本发明的单克隆抗体还可以用于诊断或监测某些类型癌症进展的测定中。利用本发明的单克隆抗体的测定法也可用于分析或监测毒物暴露，例如来自环境来源。本发明的单克隆抗体用免疫原8-OH-腺苷与钥孔血蓝蛋白 (KLH) 偶联制备，而不是直接与8-OH-Ade偶联。据信单克隆抗体与结构的碱基部分 (8-OH-Ade) 结合而不与结合物的碳水化合物 (核糖) 或蛋白质连接区结合，因为如所证明的，结合物与8-以外的核苷结合。OH-腺苷与这些抗体不反应。因此，本发明的抗体可用于检测和定量 (通过使用标准曲线) DNA的生物样品中8-OH-Ade的存在。这种测定的程序包括固定DNA，使其变性以破坏暴露游离碱结构的碱基配对方案，并定量定量每量DAN存在的8-OH-Ade的量。免疫测定。




Fig. 1.A.




Fig. 1.B.