



US 20050142606A1

(19) **United States**

(12) **Patent Application Publication** (10) **Pub. No.: US 2005/0142606 A1**
Yuan (43) **Pub. Date: Jun. 30, 2005**

(54) **DETERMINATION OF IONS USING
ION-SENSITIVE ENZYMES**

Publication Classification

(76) Inventor: **Chong-Sheng Yuan**, San Diego, CA
(US)

(51) **Int. Cl.⁷** **G01N 33/53**; C07K 14/705;
C12N 9/22
(52) **U.S. Cl.** **435/7.1**; 530/350; 435/199

Correspondence Address:
MORRISON & FOERSTER LLP
3811 VALLEY CENTRE DRIVE
SUITE 500
SAN DIEGO, CA 92130-2332 (US)

(57) **ABSTRACT**

(21) Appl. No.: **10/665,883**

(22) Filed: **Sep. 19, 2003**

The invention relates generally to the field of sodium and lithium ion detection. In particular, the invention provides chimeric proteins, nucleic acids encoding chimeric proteins, methods and kits for assaying for sodium ions and for lithium ions in a sample, using inter alia, a 3'(2'),5'-bisphosphate nucleotidase.




Figure 1




Figure 2

DETERMINATION OF IONS USING ION-SENSITIVE ENZYMES

BACKGROUND OF THE INVENTION

[0001] Serum electrolytes play a critical role in regulating normal physiologic functioning within and between cells. The testing of serum electrolytes is one of the most common analytical tests performed within hospitals. Such measurements are employed for routine monitoring of a patient as well as in emergency and life-threatening situations. Because of the vital role of electrolytes in normal physiologic responses, it is important that the measurement of the serum levels of electrolytes can be performed efficiently and accurately.

[0002] Sodium is one serum electrolyte critical in the physiologic control of water movement between the intracellular fluid compartment and the extracellular fluid compartment, i.e., maintaining osmotic pressure. In the healthy individual, the serum level of sodium is 135-145 mEq/l. Small deviations from normal level can have severe health consequences. An increased serum sodium can result from dehydration due to diarrhea or vomiting or nephrogenic diabetes. Low sodium levels usually are a result of too much water in the body, a condition associated with congestive heart failure, cirrhosis, nephritic syndrome, chronic renal failure, and syndrome of inappropriate anti-diuretic hormone (IADH).

[0003] Another source of electrolytes affecting physiologic function can also be those ions exogenously administered for therapeutic purposes. One example of such an ion is lithium. Therapeutic administration of lithium, typically as lithium carbonate, is one of the most effective agents for the treatment of patients suffering from bipolar disorder (manic depressive psychosis). Lithium acts by altering intraneuronal metabolism of catecholamines, inhibition of noradrenaline sensitive adenylate cyclase, and reduction in synaptic transmission and increase in neuronal excitability without modification of central nervous system (CNS) amine levels. Recently, studies have shown that lithium also holds promise in the treatment of Alzheimer's disease. However, lithium has severe toxic side effects. Toxicity is closely related to serum lithium levels and can occur at doses close to therapeutic levels, making the timely and accurate monitoring of serum levels critical. For example, serum Li^+ levels over 1.5 mM (12 hours after a dose) usually indicate a significant risk of lithium toxicity.

[0004] Currently, the two most commonly used methods to detect serum sodium and lithium are ion-selective electrode (ISE) and flame photometry. ISE relies on ion-specific electrodes. Ideally, each electrode possesses a unique ion-selective property that allows it to respond to the desired ion. However, in practice, interference from other ions in the sample compromise the specificity of the detecting electrode, rendering the electrodes susceptible to false readings. The instrumentation for ISE is relatively expensive, requires routine maintenance that is sometimes cumbersome and time-consuming, and demands that the operating technician to have a considerable degree of skill and knowledge for accurate and consistent readings. Flame photometry relies on the principle that certain atoms, when energized by heat, become excited and emit a light of characteristic wavelength of radiant energy when returning to ground state. The

intensity of the characteristic wavelength of radiant energy produced by atoms in the flame is directly proportional to the number of atoms excited in the flames, which is directly proportional to the concentration of the substance of interest in the sample. Like ISE, the instrumentation required for this method is complex and relatively expensive. Moreover, flame photometry requires the use of combustible gas, introducing sometimes expensive hazard prevention measures.

[0005] Conventional methods to detect sodium and lithium ions in samples are limited by complex instrumentation, potentially expensive and cumbersome maintenance, additional hazards, and often time requirements not suitable to emergency situations. The present invention addresses these problems and is more user friendly in automated analyzers.

BRIEF SUMMARY OF THE INVENTION

[0006] In one aspect, the present invention is directed to an isolated chimeric protein, which chimeric protein comprises, from N-terminus to C-terminus: a) a first peptidyl fragment comprising a bacterial leader sequence from about 5 to about 30 amino acid residues; and b) a second peptidyl fragment comprising a 3'(2'),5'-bisphosphate nucleotidase.

[0007] In another aspect, the present invention is directed to an isolated nucleic acid comprising a nucleotide sequence encoding a chimeric protein, which chimeric protein comprises, from N-terminus to C-terminus: a) a first peptidyl fragment comprising a bacterial leader sequence from about 5 to about 30 amino acid residues; and b) a second peptidyl fragment comprising a 3'(2'),5'-bisphosphate nucleotidase. Recombinant cells comprising the nucleic acid and methods for producing the chimeric protein using the nucleic acid are also provided.

[0008] In still another aspect, the present invention is directed to a method for assaying for sodium ions in a sample, which method comprises: a) contacting the sample with a sodium-sensitive 3'(2'),5'-bisphosphate nucleotidase, wherein the nucleotidase consumes adenosine 3',5'-bisphosphate (PAP) and forms AMP and P_i ; and b) assessing the consumption of PAP or the formation of AMP and P_i in step a) to determine the presence or amount of sodium ions in the sample.

[0009] In yet another aspect, the present invention is directed to a method for assaying for sodium ions in a sample, which method comprises: a) contacting the sample with a first composition comprising adenosine 3',5'-bisphosphate (PAP); b) contacting the sample with a second composition comprising a sodium-sensitive 3'(2'),5'-bisphosphate nucleotidase; and c) assessing the production of AMP to determine the presence or amount of sodium ions in the sample. In one embodiment, the first composition further comprises 4-aminoantipyrene (4-AA), N-ethyl-N-(2-hydroxy-3-sulfopropyl)-3-m-toluidine (EHSPT), purine nucleoside phosphorylase, xanthine oxidase, and peroxidase, and the second composition further comprises adenosine deaminase, 5'-nucleotidase, and MgCl_2 . Kits for assaying for sodium ions using the methods are also provided.

[0010] In still another aspect, the present invention is directed to a method for assaying for lithium ions in a sample, which method comprises: a) contacting the sample

with a lithium-sensitive 3'(2'),5'-bisphosphate nucleotidase, wherein the nucleotidase consumes adenosine 3',5'-bisphosphate (PAP) and forms AMP and P_i; and b) assessing the amount of PAP consumed or AMP formed in step b) to determine the presence or absence of lithium ions in the sample. In one embodiment, the sample is first contacted with a sodium blocking agent. In a specific embodiment, the blocking agent is 3',5' bisphosphate nucleotidase.

[0011] In yet another aspect, the present invention is directed to a method for assaying for lithium ions in a sample, which method comprises: a) contacting the sample with a first composition comprising a adenosine 3',5'-bisphosphate (PAP); b) contacting the sample with a second composition comprising a lithium-sensitive 3'(2'),5'-bisphosphate nucleotidase; and c) assessing the production of a detectable product to determine the presence or absence of lithium ions in the sample. In one embodiment, the first composition further comprises 4-aminoantipyrine (4-AA), N-ethyl-N-(2-hydroxy-3-sulfoethyl)-3-m-toluidine (EHSPT), purine nucleoside phosphorylase, xanthine oxidase, and peroxidase, and the second composition further comprises adenosine deaminase, 5'-nucleotidase, and MgCl₂. In one embodiment, the sample is first contacted with a sodium blocking agent. In a preferred embodiment, the sodium blocking agent is 4,7,13,16,21-pentaoxa-1,10-diazabicyclo[8.8.5]-tricosane. Kits for assaying for lithium ions using the method are also provided.

BRIEF DESCRIPTION OF THE DRAWINGS

[0012] FIG. 1 is a serum sodium calibration curve. The calibration curve was generated using the methods disclosed in the Example 1. Briefly, the calibration curve was constructed by plotting the ΔA values of the standards against the corresponding sodium concentration.

[0013] FIG. 2 is a serum lithium calibration curve. The calibration curve was generated using the methods disclosed in the Example 2. Briefly, the calibration curve was constructed by plotting the ΔA values of the standards against the corresponding lithium concentration.

DETAILED DESCRIPTION OF THE INVENTION

[0014] For clarity of disclosure, and not by way of limitation, the detailed description of the invention is divided into the subsections that follow.

[0015] A. Definitions


[0016] Unless defined otherwise, all technical and scientific terms used herein have the same meaning as is commonly understood by one of ordinary skill in the art to which this invention belongs. All patents, applications, published applications and other publications referred to herein are incorporated by reference in their entirety. If a definition set forth in this section is contrary to or otherwise inconsistent with a definition set forth in the patents, applications, published applications and other publications that are herein incorporated by reference, the definition set forth in this section prevails over the definition that is incorporated herein by reference.

[0017] As used herein, "a" or "an" means "at least one" or "one or more."

[0018] As used herein, a "leader sequence" refers to a peptide sequence, when fused to a target peptide or protein, increases stability and/or expression level of the target

peptide or protein. Normally, a leader sequence increases stability and/or expression level of the target peptide or protein for at least 50%. Preferably, a leader sequence increases stability and/or expression level of the target peptide or protein for at least 1 fold, 2 folds, 5 folds, 10 folds or more than 10 folds. In the regulation of gene expression for enzymes concerned with amino acid synthesis in prokaryotes, the leader sequence codes for the leader peptide that contains several residues of the amino acid being regulated. Transcription is closely linked to translation, and if translation is retarded by limited supply of aminoacyl tRNA for the specific amino acid, the mode of transcription of the leader sequence permits full transcription of the operon genes; otherwise complete transcription of the leader sequence prematurely terminates transcription of the regulated gene.

[0019] As used herein, a "3'(2'),5'-bisphosphate nucleotidase" refers to an enzyme catalyzing the dephosphorylation of adenosine 3',5'-bisphosphate to yield corresponding adenosine 5'-phosphate (AMP) and P_i, as shown in the following reaction:



[0020] Other synonyms of 3'(2'),5'-bisphosphate nucleotidase include bisphosphate 3'-nucleotidase, HAL2 phosphatase, phosphoadenylate 3'-nucleotidase, 3'(2'),5'-bisphosphonucleoside, 3'(2')-phosphohydrolase, 3'-phosphoadenylylsulfate 3'-phosphatase, DPNase, and PAP phosphatase. For purposes herein, the name "3'(2'),5'-bisphosphate nucleotidase" is used herein, although all such chemical synonyms are contemplated. "3'(2'),5'-bisphosphate nucleotidase" also encompasses a functional fragment or a derivative that still substantially retain its enzymatic activity catalyzing the dephosphorylation of adenosine 3',5'-bisphosphate to yield corresponding AMP and P_i. Typically, a functional fragment or derivative retains at least 50% of its 3'(2'),5'-bisphosphate nucleotidase activity. Preferably, a functional fragment or derivative retains at least 60%, 70%, 80%, 90%, 95%, 99% or 100% of its 3'(2'),5'-bisphosphate nucleotidase activity. It is also intended that a 3'(2'),5'-bisphosphate nucleotidase can include conservative amino acid substitutions that do not substantially alter its activity. Suitable conservative substitutions of amino acids are known to those of skill in this art and may be made generally without altering the biological activity of the resulting molecule. Those of skill in this art recognize that, in general, single amino acid substitutions in non-essential regions of a polypeptide do not substantially alter biological activity (see, e.g., Watson, et al., *Molecular Biology of the Gene*, 4th Edition, 1987, The Bejacmin/Cummings Pub. Co., p. 224). Such exemplary substitutions are preferably made in accordance with those set forth in TABLE 1 as follows:

TABLE 1

Original residue	Conservative substitution
Ala (A)	Gly; Ser
Arg (R)	Lys
Asn (N)	Gln; His
Cys (C)	Ser
Gln (Q)	Asn

TABLE 1-continued

Original residue	Conservative substitution
Glu (E)	Asp
Gly (G)	Ala; Pro
His (H)	Asn; Gln
Ile (I)	Leu; Val
Leu (L)	Ile; Val
Lys (K)	Arg; Gln; Glu
Met (M)	Leu; Tyr; Ile
Phe (F)	Met; Leu; Tyr
Ser (S)	Thr
Thr (T)	Ser
Trp (W)	Tyr
Tyr (Y)	Trp; Phe
Val (V)	Ile; Leu

[0021] Other substitutions are also permissible and may, be determined empirically or in accord with known conservative substitutions.

[0022] As used herein, a “composition” refers to any mixture of two or more products or compounds. It may be a solution, a suspension, liquid, powder, a paste, aqueous, non-aqueous, or any combination thereof.

[0023] As used herein, a “combination” refers to any association between two or among more items.

[0024] As used herein, “biological sample” refers to any sample from a biologic source, including but not limited to blood, plasma, and serum samples.

[0025] As used herein, “plasma” refers to the fluid, non-cellular portion of the blood, distinguished from the serum obtained after coagulation.

[0026] As used herein, “serum” refers to the fluid portion of the blood obtained after removal of the fibrin clot and blood cells, distinguished from the plasma in circulating blood.

[0027] As used herein, “fluid” refers to any composition that can flow. Fluids thus encompass compositions that are in the form of semi-solids, pastes, solutions, aqueous mixtures, gels, lotions, creams, and other such compositions.

[0028] As used herein, “whole blood sample” refers to a blood sample containing both the cell and fluid portions of blood.

[0029] As used herein, “red blood cell sample” refers to the red blood cells portion of the blood obtained after removal of the serum portion of the blood.

[0030] As used herein, “peroxidase” refers to an enzyme that catalyzes a host of reactions in which hydrogen peroxide is a specific oxidizing agent and a wide range of substrates act as electron donors. It is intended to encompass a peroxidase with conservative amino acid substitutions that do not substantially alter its activity. The chief commercially available peroxidase is horseradish peroxidase.

[0031] As used herein, “5'-nucleotidase” refers to an enzyme that catalyzes the formation of adenosine and P_i from adenosine 5'-phosphate (AMP). It is intended to encompass 5'-nucleotidase with conservative amino acid substitutions that do not substantially alter its activity.

[0032] As used herein, “adenosine deaminase” refers to an enzyme that catalyzes the formation of inosine and NH_3 from adenosine. It is intended to encompass any adenosine deaminase with conservative amino acid substitutions that do not substantially alter its activity.

[0033] As used herein, “purine nucleoside phosphorylase” or “PNP” refers to an enzyme that catalyzes the formation of hypoxanthine and ribose-1-phosphate (RIP) from inosine and P_i . It is intended to encompass purine nucleoside phosphorylase with conservative amino acid substitutions that do not substantially alter its activity.

[0034] As used herein, “xanthine oxidase” refers to an enzyme that catalyzes the conversion of hypoxanthine to uric acid and H_2O_2 in the presence of H_2O and O_2 . Other synonyms include xanthine: O_2 oxide reductase. It is intended to encompass xanthine oxidase with conservative amino acid substitutions that do not substantially alter its activity.

[0035] As used herein, the abbreviations for any protective groups, amino acids and other compounds are in accord with their common usage, recognized abbreviations, or the IUPAC-IUB Commission on Biochemical Nomenclature, unless otherwise indicated (see *Biochemistry* 11: 1726 (1972)).

[0036] B. Chimeric Proteins Comprising a 3'(2'),5'-bisphosphate Nucleotidase and Nucleic Acids Encoding the Same

[0037] In one aspect, the present invention is directed to isolated chimeric protein, which chimeric protein comprises, from N-terminus to C-terminus: a) a first peptidyl fragment comprising a bacterial leader sequence from about 5 to about 30 amino acid residues; and b) a second peptidyl fragment comprising a 3'(2'),5'-bisphosphate nucleotidase.

[0038] Any suitable bacterial leader sequences can be used. As disclosed in U.S. Pat. No. 6,194,200, expression of the polypeptide of interest as a fused protein with a leader sequence from another gene has several advantages in addition to providing for stability. For example, the presence of the N-terminal amino acids provides a means for using general purification techniques for purification of any of a variety of polypeptides. For example, the N-terminal amino acids of the N-protein are predictably antigenic, and thus specific antibodies raised against the N-terminal amino acids of the N-protein may be used for the amino purification of the fusion proteins containing the N-terminus of the N-protein. Furthermore, the N-terminus of the N-protein has a high positive charge, which facilitates purification of the desired protein by ion-exchange chromatography, and the like.

[0039] The leader sequence can also be a hydrophobic amino acid sequence, which may additionally function as a signal sequence for secretion. See U.S. Pat. No. 6,194,200. A DNA sequence encoding the signal sequence is joined upstream from and in reading frame with the gene of interest. Typically, the signal sequence includes a cleavage site which is recognized by a signal sequence peptidase. Thus, positioning the polypeptide of interest directly after the signal sequence cleavage site will allow it to be specifically cleaved from the signal sequence and secreted as a mature polypeptide. Examples of hydrophobic amino acid sequences include the bacterial alkaline phosphatase signal

sequence; the OMP-A, B, C, D, E or F signal sequences; the LPP signal sequence, β -lactamase signal sequence; and toxin signal sequences.

[0040] Other leader sequences which can be used include hydrophilic sequences, for example the N-terminal 41 amino acid residues from amphiregulin which may provide for modification of the function of the polypeptide of interest. See U.S. Pat. No. 6,194,200. In addition, a cytotoxic agent such as a toxin A-chain fragment, ricin A-chain, snake venom growth arresting peptide, or a targeting molecule such as a hormone or antibody can be coupled covalently with the leader sequence with in most cases minimal effect on the biological activity of the gene product of interest. As with the other leader sequences, a DNA sequence encoding the leader sequence is joined upstream from and in reading frame with the gene of interest.

[0041] Where the leader sequence is not a signal sequence or does not contain a convenient natural cleavage site, additional amino acids may be inserted between the gene of interest and the leader sequence to provide an enzymatic or chemical cleavage site for cleavage of the leader peptide, following purification of the fusion protein, to allow for subsequent purification of the mature polypeptide. See U.S. Pat. No. 6,194,200. For example, introduction of acid-labile aspartyl-proline linkages between the two segments of the fusion protein facilitates their separation at low pH. This method is not suitable if the desired polypeptide is acid-labile. The fusion protein may be cleaved with, for example, cyanogen bromide, which is specific for the carboxy side of methionine residues. Positioning a methionine between the leader sequence and the desired polypeptide would allow for release of the desired polypeptide. This method is not suitable when the desired polypeptide contains methionine residues.

[0042] Other bacterial leader sequences disclosed in the following patents, patent application and references can also be used: WO 00/28041 and WO 89/03886; U.S. Pat. Nos. 5,914,250, 5,885,811, 5,171,670, 5,030,563, 4,948,729 and 4,588,684; EP Patent Nos. EP 0,196,864, EP 0,186,643 and EP 0,121,352; Michiels et al., *Trends Microbiol.*, 9(4): 164-8 (2001); Hobom et al., *Dev. Biol. Stand.*, 84: 255-62 (1995); Hardy and Randall, *J. Cell. Sci. Suppl.*, 11: 29-43 (1989); Saier et al., *FASEB J.*, 2(3): 199-208 (1988); and Peakman et al., *Nucleic Acids Res.*, 20(22): 6111-2 (1992). Preferably, the bacterial leader sequence is a leader sequence of an *E. coli* protein, e.g., the *E. coli* leader sequences disclosed in Roesser and Yanofsky, *Nucleic Acids Res.*, 19(4): 795-800 (1991); and Kuhn et al., *Mol. Gen. Genet.*, 167(3): 235-41 (1979). In one example, the leader sequence has at least 40% identity to the amino acid sequence set forth in SEQ ID NO: 1 (MGSGDDDDDLAL), in which the percentage identity is determined over an amino acid sequence of identical size to the amino acid sequence set forth in SEQ ID NO: 1. Preferably, the leader sequence has at least 50%, 60%, 70%, 80%, 90%, 95%, 99% or 100% identity to the amino acid sequence set forth in SEQ ID NO: 1, in which the percentage identity is determined over an amino acid sequence of identical size to the amino acid sequence set forth in SEQ ID NO: 1. Also preferably, the leader sequence binds to an antibody that specifically binds to an amino acid sequence set forth in SEQ ID NO: 1. Still preferably, the leader sequence comprises the amino acid sequence set forth in SEQ ID NO: 1.

[0043] The first peptidyl fragment can have any suitable length. For example, the first peptidyl fragment comprises

about 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29 or 30 amino acid residues. Preferably, the first peptidyl fragment comprises about 20 amino acid residues.

[0044] Any suitable 3',5' bisphosphate nucleotidase can be used. In one example, the 3',5' bisphosphate nucleotidase is of *Saccharomyces cerevisiae* origin (See e.g., Murguía et al., *J. Biol. Chem.*, 271(46): 29029-33 (1996)). This nucleotidase is also known as the HAL2 nucleotidase. Moreover, any suitable 3',5' bisphosphate nucleotidase catalyzing the reaction defined in Section B can be used in the present compositions and methods. The enzyme useful in the present compositions and methods is not limited those enzymes having only 3'(2'),5'-bisphosphate nucleotidase activity. For example, the enzyme may have dual enzymatic activity, e.g., Tol-1. Homologues of the HAL2 phosphatase are also contemplated. Useful enzymes capable of catalyzing the above reaction include, but are not limited to BPntase (see, e.g., Spiegelberg et al., *J. Biol. Chem.* 274(19): 13619-28 (1999)), HsPIP, RnPIP (see, e.g., López-Coronado, et al., *J. Biol. Chem.* 274(23): 16034-39 (1999), and Tol-1 (see, e.g., Amoto, et al., *J. Bacteriol.* 182(13): 3619-25 (2000)). Other useful 3'(2'),5'-bisphosphate nucleotidases, e.g., 3',5' bisphosphate nucleotidases are disclosed in Peng et al., *J. Biol. Chem.* 270(49): 29105-10 (1995), Dichtl et al., *EMBO J.*, 16(23): 7184-95 (1997), Gil-Mascarell et al., *The Plant J.* 17(4): 373-83 (1999), can also be used. A functional fragment or a derivative of a 3'(2'),5'-bisphosphate nucleotidase that still substantially retain its enzymatic activity catalyzing the dephosphorylation of adenosine 3',5'-bisphosphate to yield corresponding adenosine 5'-phosphate (AMP) and P_i can also be used.

[0045] Normally, a functional fragment or a derivative of a 3',5'-bisphosphate nucleotidase retain at least 50% of its enzymatic activity. Preferably, a functional fragment or a derivative of a 3'(2'),5'-bisphosphate nucleotidase retain at least 50%, 60%, 70%, 80%, 90%, 95%, 99% or 100% of its enzymatic activity.

[0046] The dephosphorylation of adenosine 3',5' bisphosphate (PAP) can be assessed by any suitable methods. For example, the dephosphorylation of adenosine 3',5' bisphosphate can be assessed by assessing consumption of the adenosine 3',5' bisphosphate in the dephosphorylation reaction or the formation of the AMP or P_i in the reaction.

[0047] Assays for enzymatic activities of 3'(2'),5'-bisphosphate nucleotidases are known in the art (See e.g., Murguía et al., *J. Biol. Chem.*, 271(46): 29029-33 (1996)). Exemplary methods for phosphatase activity include determining the formation of inorganic phosphate (P_i) and AMP include calorimetric methods (See, e.g., Gumber et al., *Plant Physiol.*, 76: 40-44 (1984); Baykov et al., *Anal. Biochem.* 171: 266-70 (1988)) and radioactive-labeled substrates (See, e.g., Spiegelberg et al., *J. Biol. Chem.* 274(19): 13619-28 (1999); Peng et al., *J. Biol. Chem.* 270(49): 29105-29110 (1995)).

[0048] In another example, the 3'(2'),5'-bisphosphate nucleotidase has at least 40% identity to the amino acid sequence set forth in SEQ ID NO:2 (ALER-ELLVATQAVRKASLLTKRIQSEVISHKD-STITTKNDNSPVTITGDYAAQTIIIN AIKSNFPDDKV-VGESSSGLSDAFVSGILNEIKANDEVYNKYNKKDD-FLFTNDQFP LKSLEDVRQIIDFGNYEGGRKGRFW-CLDPIDGTGKFLRGEQFAVCLALIVDGVVQ LGCIGCPNLVLSSYGAQDLKGHESFGYI-FRAVRGLGAFYSPSSDAESWTKIHRVHL

KDTKDMITLEGVEKGHSSHD-EQTAIKNKLNISKSLHLDLSDQAKYCLLALGLADVYLRLPIKLSYQEKIWDHHAAGNVIVHEAG-GIHTDAMEDVPLDFGNGRTLATKGVIASS GPRELH-DLVVSTSCDVQSRNA), in which the percentage of identity is determined over an amino acid sequence of identical size to the amino acid sequence set forth in SEQ ID NO:2.

[0049] The first and second peptidyl fragments can be linked via any suitable linkage. For example, the first and second peptidyl fragments can be linked via a cleavable linkage.

[0050] The isolated chimeric protein can further comprise, at its C-terminus, a third peptidyl fragment comprising a second bacterial leader sequence from about 5 to about 30 amino acid residues. Any suitable bacterial leader sequences, including the ones described above, can be used.

[0051] In one example, the second bacterial leader sequence is a leader sequence of an *E. coli* protein. In another example, the second bacterial leader sequence has at least 40% identity to the amino acid sequence set forth in SEQ ID NO:3 (KGELEGLPIPNLLRTG), in which the percentage identity is determined over an amino acid sequence of identical size to the amino acid sequence set

forth in SEQ ID NO:3. Preferably, the second bacterial leader sequence has at least 50%, 60%, 70%, 80%, 90%, 95%, 99% or 100% identity to the amino acid sequence set forth in SEQ ID NO:4, in which the percentage identity is determined over an amino acid sequence of identical size to the amino acid sequence set forth in SEQ ID NO:4. Also preferably, the second bacterial leader sequence binds to an antibody that specifically binds to an amino acid sequence set forth in SEQ ID NO:3. Also preferably, the second bacterial leader sequence comprises the amino acid sequence set forth in SEQ ID NO:3.

[0052] The third peptidyl fragment can have any suitable length. For example, the third peptidyl fragment comprises about 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29 or 30 amino acid residues. Preferably, the third peptidyl fragment comprises about 20 amino acid residues.

[0053] The isolated chimeric protein can further comprise, at its C-terminus, a third peptidyl fragment comprising a peptide tag. Any suitable tag can be used. For example, the tag can be FLAG, HA, HA1, c-Myc, 6-His, AU1, EE, T7, 4A6, ϵ , B, gE and Ty1 tag (See Table 2).

TABLE 2

Exemplary epitope tag systems			
Epitope	Peptide	SEQ ID Antibody	Reference
FLAG	AspTyrLysAspAspAspLys	11 4E11	Prickett ¹
HA	TyrProTyrAspValProAspTyrAla	12 12Ca5	Xie ²
HA1	CysGlnAspLeuProGlyAsnAspAsnSerThr	13 mouse MAb	Nagelkerken ³
c-Myc	GluGlnLysLeuIleSerGluGluAspLeu	14 9E10	Xie ²
6-His	HisHisHisHisHisHis	15 BABCO*	
AU1	AspThrTyrArgTyrIle	16 BABCO	
EE	GluTyrMetProMetGlu	17 anti-EE	Tolbert ⁴
T7	AlaSerMetThrGlyGlyGlnGlnMetGlyArg	18 Invitrogen	Chen ⁵
			Tseng ⁶
4A6	SerPheProGlnPheLysProGlnGluIle	19 4A6	Rudiger ⁷
ϵ	LysGlyPheSerTyrPheGlyGluAspLeuMetPro	20 anti-PKCe	Olah ⁸
B	GlnTyrProAlaLeuThr	21 D11, F10	Wang ⁹
gE	GlnArgGlnTyrGlyAspValPheLysGlyAsp	22 3B3	Grose ¹⁰
Ty1	GluValHisThrAsnGlnAspProLeuAsp	23 BB2, TYG5	Bastin ¹¹

¹Prickett, et al., Bio Techniques, 7(6): 580-584 (1989)

²Xie, et al., Endocrinology, 139(11): 4563-4567 (1998)

³Nagelkerke, et al., Electrophoresis, 18: 2694-2698 (1997)

⁴Tolbert and Lamah, J. Neurochem., 70: 113-119 (1998)

⁵Chen and Katz, Bio Techniques, 25(1): 22-24 (1998)

⁶Tseng and Verma, Gene, 169: 287-288 (1996)

⁷Rudiger, et al., Bio Techniques, 23(1): 96-97 (1997)

⁸Olah, et al., Biochem., 221: 94-102 (1994)

⁹Wang, et al., Gene, 169(1): 53-58 (1996)

¹⁰Grose, U.S. Pat. No. 5,710,248

¹¹Bastin, et al., Mol. Biochem. Parasitology, 77: 235-239 (1996)

Invitrogen, Sigma, Santa Cruz Biotech

[0054] In an example, the isolated chimeric protein comprises the amino acid sequence set forth in SEQ ID NO: 4

```
(mgsgdddlalALERELLVATQAVRKASLLTKRIQSEVISHKDSTTITKNDNSPVTTG
DYAAQTII INAIKSNFPDDKVVGEESSSGLSDAFVSGILNEIKANDEVYNKYK
DFLFTNDQFPLKSLEDVRQIIDFGNYEGGRKGRFWCLDPIDGTGKFLRGEQFAVCL
ALIVDGVVQLGCIGCPNLVLSYGAQDLKGHESFGYIFRAVRGLGAFYSPSSDAES
WTKIHVRLKDTKDMITLEGVEKGHSSHDEQTAIKNKLNIKSLHLDSQAKYCLL
ALGLADVYLRLPIKLSYQEKIWDHAAGNVIVHEAGGIHTDAMEDVPLDFGNRT
LATKGVIIASSGPRELHDLVVSTSCDVIQSRNAkgelelplpnp1l1rtghhhhh).
```

[0055] In another aspect, the present invention is directed to an isolated nucleic acid comprising a nucleotide sequence encoding a chimeric protein, which chimeric protein com-

[0056] In one example, the isolated nucleic acid comprises a nucleotide sequence encoding the chimeric protein com-

prising the amino acid sequence set forth in SEQ ID NO:4. In another example, the isolated nucleic acid comprises a nucleotide sequence set forth in SEQ ID NO: 5

```
(atgggcgatccggtgatgacgatgacctcgcccttGCATTGGAAGAGAATTATTGGTTGCAACT
CAAGCTGTACGAAAGCGCTCTTATTGACTAAGAGAATTCAATCTGAAGTGAT
TTCTCACAAGACTCCACTACTATTACCAAGAATGATAATCTCCAGTAACCA
CAGGTGATTATGCTGCACAAACGATCATCAAAATGCTATCAAGAGCAATTTT
CCTGATGATAAGGTAGTTGGTGAAGAATCCTCATCAGGATTGAGCGACGCATT
CGTCTCAGGAATTTTAAACGAAATAAAAGCCAATGACGAAGTTTATAACAAG
AATTATAAAAAGGATGATTTTCTGTTTACAAACGATCAGTTTCCGCTAAATC
TTTGGAGGACGTCAGGCAAATCATCGATTTTCGGCAATTACGAAGGTGGTAGAA
AAGGAAGATTTTGGTGTGGATCCTATTGACGGAACCAAGGGGTTTTTAAGA
GGTGAACAGTTTGCAGTATGCTGGCCTTAATTGTGACGGTGTGTTTCAGCTT
GGTTGATTGGATGCCCCAACTTAGTTTAAAGTTCTTATGGGGCCAAGATTTG
AAAGCCATGAGTCATTTGGTTATATCTTTCGTGCTGTTAGAGGTTTAGGTGCC
TTCTATTCTCCATCTTCAGATGCAGAGTCATGGACAAAATCCACGTTAGACA
CTTAAAGACACTAAAGACATGATTACTTTAGAGGGAGTTGAAAAGGGACAC
TCCTCTCATGATGAACAACTGCTATCAAAACAACTAAATATATCCAAATC
TTTGCACTTGGATTCTCAAGCCAAGTACTGTTTGTAGCATTGGGCTTAGCAGA
CGTATATTTACGTCTGCCTATCAAACTTTCTTACCAAGAAAAGATCTGGGACC
ATGCTGCAGGCAACGTTATTGTCCATGAAGCTGGAGGTATCCATACAGATGCC
ATGGAAGATGTTCTCTAGACTTCGGTAACGGTAGAACGCTAGCTACGAAGG
AGTTATAGCGTCAAGTGCCACGCGAGTTACATGACTTGGTGGTGTCTACAT
CATGCGATGTCATTAGTCAAGAAACGCCaagggcgagcttgaaggtttgcctatccctaaccctctc
ctccgtaccggtcatcatcaccatcaccattga).
```

prises, from N-terminus to C-terminus: a) a first peptidyl fragment comprising a bacterial leader sequence from about 5 to about 30 amino acid residues; and b) a second peptidyl fragment comprising a 3'(2'),5'-bisphosphate nucleotidase.

[0057] In still another example, the isolated nucleic acid comprising a nucleotide sequence complementary to the nucleotide sequence encoding a chimeric protein, which chimeric protein comprises, from N-terminus to C-terminus:

a) a first peptidyl fragment comprising a bacterial leader sequence from about 5 to about 30 amino acid residues; and b) a second peptidyl fragment comprising a 3'(2'),5'-bisphosphate nucleotidase.

[0058] In another example, a recombinant cell containing the nucleic acid, or a complementary strand thereof, encoding a chimeric protein, which chimeric protein comprises, from N-terminus to C-terminus: a) a first peptidyl fragment comprising a bacterial leader sequence from about 5 to about 30 amino acid residues; and b) a second peptidyl fragment comprising a 3'(2'),5'-bisphosphate nucleotidase, is contemplated.

[0059] A method of producing a chimeric protein is also contemplated, which method comprising growing a recombinant cell containing the nucleic acid encoding a chimeric protein, which chimeric protein comprises, from N-terminus to C-terminus: a) a first peptidyl fragment comprising a bacterial leader sequence from about 5 to about 30 amino acid residues; and b) a second peptidyl fragment comprising a 3'(2'),5'-bisphosphate nucleotidase, such that the encoded chimeric protein is expressed by the cell, and recovering the expressed chimeric protein. The product of the method is further contemplated.

[0060] The chimeric proteins and the nucleic acids encoding the chimeric proteins can be prepared by any suitable methods, e.g., chemical synthesis, recombinant production or a combination thereof (See e.g., CURRENT PROTOCOLS IN MOLECULAR BIOLOGY, Ausubel, et al. eds., John Wiley & Sons, Inc. (2000) and Sambrook, et al., MOLECULAR CLONING: A LABORATORY MANUAL, Cold Spring Harbor Laboratory press, (1989)).

[0061] C. Methods and Kits for Assaying for Sodium Ions Using a Chimeric Protein

[0062] In still another aspect, the present invention is directed to a method for assaying for sodium ions in a sample, which method comprises: a) contacting the sample with a sodium-sensitive 3'(2'),5'-bisphosphate nucleotidase, wherein the nucleotidase consumes adenosine 3',5'-bisphosphate (PAP) and forms AMP and P_i ; and b) assessing the consumption of PAP or the formation of AMP and P_i in step a) to determine the presence or amount of sodium ions in the sample.


[0063] In yet another aspect, the present invention is directed to a method for assaying for sodium ions in a sample, which method comprises: a) contacting the sample with a first composition comprising adenosine 3',5'-bisphosphate (PAP); b) contacting the sample with a second composition comprising a sodium-sensitive 3'(2'),5'-bisphosphate nucleotidase; and c) assessing the production of AMP to determine the presence or amount of sodium ions in the sample. In one embodiment, the first composition further comprises 4-aminoantipyrine (4-AA), N-ethyl-N-(2-hydroxy-3-sulfopropyl)-3-m-toluidine (EHSPT), purine nucleoside phosphorylase, xanthine oxidase, and peroxidase, and the second composition further comprises adenosine deaminase, 5'-nucleotidase, and $MgCl_2$.

[0064] The dephosphorylation of adenosine 3',5' bisphosphate (PAP) can be assessed by any suitable methods. For example, the dephosphorylation of adenosine 3',5' bisphosphate can be assessed by assessing consumption of the

adenosine 3',5' bisphosphate in the dephosphorylation reaction or the formation of the AMP or P_i in the reaction.

[0065] In one embodiment, the formation of AMP can be assessed using a combination of 5'-nucleotidase, adenosine deaminase, purine nucleoside phosphorylase, xanthine oxidase, and peroxidase. For example, the following series of coupled enzymatic reactions can result in the production of detectable quinone dye:


Reaction (I):




Reaction (II):




Reaction (III):



Reaction (IV):



Reaction (V):



[0066] Any suitable 3'(2'),5'-bisphosphate nucleotidase can be used. Any source or form known in the art that permits the production of P_i and AMP from PAP is contemplated. In particular, any suitable chimeric proteins, including the ones described in the above Section B, can be used in the present methods. In one example, the chimeric protein comprises the amino acid sequence set forth in SEQ ID NO:4. In another example, the chimeric protein is encoded by the nucleotide sequence set forth in SEQ ID NO:5.

[0067] Any suitable adenosine 3',5'-bisphosphate (PAP) may be used. PAP may be isolated, purified or recombinantly generated from any source known in the art, that is subject to the enzymatic activity of 3'(2'),5'-bisphosphate nucleotidase.

[0068] Any suitable 5'-nucleotidase, adenosine deaminase, purine nucleoside phosphorylase, and xanthine oxidase can be used. The enzymes can be derived from any source known in the art, including microbial and mammalian, that will permit the generation of a suitable detectable product. In one embodiment, ascorbate oxidase is also employed.

[0069] H_2O_2 formation can be assessed any suitable means. In one embodiment, the H_2O_2 formation is assessed by a peroxidase and Trinder reaction. Any suitable peroxidase can be used. More preferably, a horseradish peroxidase is used. For example, the horseradish peroxidases with the following GenBank accession Nos. can be used: E01651; D90116 (prxC3 gene); D90115 (prxC2 gene); J05552 (Synthetic isoenzyme C(HRP-C)); S14268 (neutral); OPRHC (C1 precursor); S00627 (C1C precursor); JH0150 (C3 precursor); S00626 (C1B precursor); JH0149 (C2 precursor); CAA00083 (*Armoracia rusticana*); and AAA72223 (synthetic horseradish peroxidase isoenzyme C(HRP-C)). Any suitable Tinder reagent can be used herein. Hydrogen peroxide can be quantitated by the quinone dye assay. See, e.g., Tamaokel, et al., *Chem. Pharm. Bull.* 30: 2497 (1982); Shimojo et al., *Clin. Chem.* 35(9): 1992-94 (1989). The amount of quinone dye formed is inversely related to the amount of sodium ions in the sample.

[0070] Any suitable chromagen may be employed. In one embodiment, the chromagen is a Tinder reagent. Any suitable Tinder reagent can be used herein. In a specific embodiment, the chromagen is N-ethyl-N-(2-hydroxy-3-sulfopropyl)-3-m-toluidine (EHSPT) in combination with 4-aminoantipyrine (4-AA). Exemplary chromagens include, but are not limited to the combinations of a coupler (e.g., 4-aminoantipyrine, 3-methyl-2-benzothiazolinone hydrazone, NCP-04, NCP-05, NCP-06, or NCP-07) and a phenol derivative (e.g., phenol, 2-chlorophenol, 4-chlorophenol, 2,4-dichlorophenol) or an aniline derivative (e.g., aniline, N,N-dimethyl-m-anisidine, N-ethyl-N-(3-methyl-phenyl)-N'-acetylenediamine, N-ethyl-N-(beta-hydroxyethyl)-m-toluidine, N-ethyl-N-(hydroxy-3-sulfopropyl)-m-toluidine, N-ethyl-N-sulfopropyl-m-toluidine, N-ethyl-N-sulfopropyl-3,5-dimethoxyaniline, N-ethyl-N-(2-hydroxy-3-sulfopropyl)-3,5-dimethoxyaniline, N-ethyl-N-sulfopropyl-m-anisidine, N-ethyl-N-(3-methylphenyl)-N'-succinylethylenediamine, and N-ethyl-N-(2-hydroxy-3-sulfopropyl)-m-anisidine. Leuco dyes (e.g., 10-N-methylcarbamoyl-3,7-dimethylamino-10H-phenothiazine, bis[3-bis(4-chlorophenyl)methyl-dimethylaminophenyl]amine, 4,4-bis(dimethylamino)diphenyl(2,7-dihydroxy-1-naphthyl)methane) also are contemplated as useful in the present methods. Other aniline derivatives include N,N-bis(4-sulfobutyl)-3-methylaniline (TODB), N,N-bis(4-sulfobutyl)-3,5-dimethylaniline (MADB), N-ethyl-N-(2-hydroxy-3-sulfopropyl)-3,5-dimethylaniline (MAOS), N-2-hydroxy-3-sulfopropyl)-3,5-dimethoxyaniline (HDAOS), N-(3-sulfopropyl)-3,5-dimethoxyaniline (HDAPS), N-ethyl-N-(3-sulfopropyl)-3-methoxyaniline (ADPS), and the like. Other suitable chromagens include N-(carboxymethylaminocarbonyl-4,4'-bis(dimethylamino)-diphenylamine (DA-64, $E_{727}^{nm}=9 \times 10^4$), 10-(carboxymethylaminocarbonyl)-3,7-bis(dimethylamino)-phenothiazine (DA-67, $E_{666}^{nm}=9 \times 10^4$).

[0071] The sample can be contacted with the 3'(2'),5'-bisphosphate nucleotidase and peroxidase sequentially or simultaneously. Likewise, any other enzymes used can be contacted with the 3'(2'),5'-bisphosphate nucleotidase sequentially or simultaneously in a fashion that permits the formation of a detectable product.

[0072] If desirable, interference of the assay can be countered. For example, ascorbate interference can be countered using a copper (II) compound, a cholic acid or a batho-

phenanthroline disulphonic acid or a mixture thereof. Bilirubin interference can be countered using a ferrocyanide salt.

[0073] The present methods can be used to assay any suitable sample. Preferably, the sample is a biological sample. In one example, the sample is a blood sample, e.g., a plasma, serum, red blood cell or whole blood sample.

[0074] The present methods can be used for any suitable purpose. Preferably, the method is used in prognosis or diagnosis of a disease or disorder. In particular, the present methods are useful in assessing the presence or amount of sodium ions in a sample.

[0075] Any suitable conditions for detection or measurement of sodium ions can be used. The reaction temperature is usually in the range from 10° C. to 40° C., with a preferred temperature of 25° C. or 37° C. The reaction time is preferably not more than 15 minutes, most preferably about 10 minutes or less.

[0076] Any suitable means of performing calorimetric analysis can be used. In one embodiment, the samples are analyzed for the presence of quinone dye in a Roche Cobas Mira Chemistry Analyzer.

[0077] Any suitable means for preparing the sample may be employed. In one embodiment, serum or plasma samples are treated with heparinate.

[0078] In yet another aspect, the present invention is directed to a kit of assaying for sodium ions in a sample, which kit comprises: a) a first composition comprising a sodium-sensitive 3'(2'),5'-bisphosphate nucleotidase that consumes adenosine 3',5'-bisphosphate and forms AMP and P_i ; and b) means for assessing the product formed or the substrate consumed by the nucleotidase to determine the presence or amount of the sodium ions in the sample. In one embodiment, the first composition further comprises adenosine deaminase, 5'-nucleotidase and $MgCl_2$. In one embodiment, the kit further comprises a second composition comprising 4-aminoantipyrine (4-AA), N-ethyl-N-(2-hydroxy-3-sulfopropyl)-3-m-toluidine (EHSPT), purine nucleoside phosphorylase, xanthine oxidase, and peroxidase, wherein the reaction of 4-AA and EHSPT in the presence of peroxidase is the means for assessing the product formed if sodium ions are not present. In some embodiments, the kit also comprises a low sodium serum standard and a high sodium serum standard. In a specific embodiment, the low sodium standard is 80 mM Na^+ and the high sodium standard is 180 mM Na^+ .

[0079] Any suitable means can be included in the present kits. For example, the means for assessing dephosphorylation of the adenosine 3',5' bisphosphate by the chimeric protein can comprise a peroxidase. Preferably, the chimeric protein and the peroxidase are formulated in a single composition.

[0080] Any suitable 3'(2'),5'-bisphosphate nucleotidase, including the ones described in the above Sections B, can be used in the present methods. For example, the 3'(2'),5'-bisphosphate nucleotidase can comprise a chimeric protein, which chimeric protein comprises, from N-terminus to C-terminus: a) a first peptidyl fragment comprising a bacterial leader sequence from about 5 to about 30 amino acid residues; and b) a second peptidyl fragment comprising an 3'(2'),5'-bisphosphate nucleotidase. Preferably, the chimeric

protein comprises the amino acid sequence set forth in SEQ ID NO:4. Also preferably, the chimeric protein is encoded by the nucleotide sequence set forth in SEQ ID NO:5.

[0081] The adenosine 3',5'-bisphosphate (PAP) to be used herein may be in any suitable form of a salt, so long as it contains no sodium ions. A preferred form is a potassium salt.

[0082] The compositions of the present invention may be formulated into a reagent having a pH adjusted by the addition of a buffer to pH 6 to 9. Any suitable buffer may be used. It is contemplated that such buffers contain no sodium ions. Exemplary buffers are Good's buffer, triethanolamine buffer, MES buffer, and tris buffer.

[0083] The compositions of the present invention may further contain any surfactant, preservative, stabilizer, and enzyme activator. Preferred examples of the surfactant are Triton-100. Preferred examples of the preservative include Thimerosal. Any suitable stabilizer can be used. In one embodiment, the stabilizer is a protein. In a specific embodiment, the protein is bovine serum albumin. Any suitable enzyme activator can be used. In one embodiment, the activator is Mg^{2+} or a salt thereof, e.g., $MgCl_2$.

[0084] Any suitable concentration of 3'(2'),5'-bisphosphate nucleotidase can be used in a composition for measurement of sodium ions. In a preferred embodiment, the concentration is in the range of 0.1-5 u/ml, more preferably, 0.5-3 u/ml, most preferably 2-3 u/ml. Any suitable concentration of 5'-nucleotidase can be used. In a preferred embodiment, the concentration is in the range of 0.1-5 u/ml, more preferably, 0.5-3 u/ml, most preferably 2-3 u/ml. Any suitable concentration of adenosine deaminase can be used. In a preferred embodiment, the concentration is in the range of 0.1-5 u/ml, more preferably, 0.5-3 u/ml, most preferably 1-2 u/ml. Any suitable concentration of peroxidase can be used. In a preferred embodiment, the concentration is in the range of 1-50 u/ml, more preferably, 5-30 u/ml, most preferably 5-10 u/ml. In one embodiment, ascorbate oxidase is employed.

[0085] The chromagen of the reduced type, N-ethyl-N-(2-hydroxy-3-sulfopropyl)-3-m-toluidine (EHSPT), and 4-aminoantipyrene (4-AA), or a salt thereof are used at any concentration suitable for measurement. The chromagen of the reduced type is preferably used at a concentration in the range of 0.01 to 10 mM. The N-ethyl-N-(2-hydroxy-3-sulfopropyl)-3-m-toluidine (EHSPT) or salt thereof is preferably used at a concentration of 4 mM. 4-aminoantipyrene (4-AA) or salt thereof is preferably used at a concentration of 2 mM.

[0086] In some embodiments, standards for calibration of the assay are included. In one embodiment, a low sodium serum standard and a high sodium standard are included. Preferably, the low sodium serum standard comprises 80-110 mM of sodium, preferably 80 mM, in serum and the high sodium serum standard comprises 160-180 mM of sodium, preferably 180 mM, in serum. In one embodiment, the presence or amount of sodium ions are calculated using a calibration curve. The amount of detectable chromagen is assessed at time 1 for a value of A_1 and at time 2 for a value

of A_2 . The resultant value is calculated in the following equation: $\Delta A = A_2 - A_1$. A calibration curve is generated by plotting the ΔA values of the standards. The amount of sodium in the samples are then determined by plotting the sample ΔA value on the calibration curve. In one embodiment, time 1 is 3 minutes after the addition of means to assess P_i production and time 2 is 8 minutes after time 1.

[0087] D. Methods and Kits for Assaying for Lithium Ions Using a Chimeric Protein

[0088] In still another aspect, the present invention is directed to a method for assaying for lithium ions in a sample, which method comprises: a) contacting the sample with a lithium-sensitive 3'(2'),5'-bisphosphate nucleotidase, wherein the nucleotidase consumes adenosine 3',5'-bisphosphate (PAP) and forms AMP and P_i ; and b) assessing the amount of PAP consumed or AMP formed in step b) to determine the presence or absence of lithium ions in the sample. In one embodiment, the sample is first contacted with a sodium blocking agent. In a specific embodiment, the sodium blocking agent is 4,7,13,16,21-pentaoxa-1,10-diazabicyclo[8.8.5]-tricosane.

[0089] In yet another aspect, the present invention is directed to a method for assaying for lithium ions in a sample, which method comprises: a) contacting the sample with a first composition comprising an adenosine 3',5'-bisphosphate (PAP); b) contacting the sample with a second composition comprising a lithium-sensitive 3'(2'),5'-bisphosphate nucleotidase; and c) assessing the production of a detectable product to determine the presence or absence of lithium ions in the sample. In one embodiment, the first composition further comprises 4-aminoantipyrene (4-AA), N-ethyl-N-(2-hydroxy-3-sulfopropyl)-3-m-toluidine (EHSPT), purine nucleoside phosphorylase, xanthine oxidase, and peroxidase, and, and the second composition further comprises adenosine deaminase, 5'-nucleotidase, and $MgCl_2$. In one embodiment, the sample is first contacted with a sodium blocking agent. In a specific embodiment, the sodium blocking agent is 4,7,13,16,21-pentaoxa-1,10-diazabicyclo[8.8.5]-tricosane.

[0090] Any suitable blocking agent may be used in the present methods. Exemplary blocking agents include, but are not limited to bis[(12-crown-4)methyl]-2-dodecyl-2-methylmalonate and 4,7,13,16,21-pentaoxa-1,10-diazabicyclo[8.8.5]-tricosane. In a preferred embodiment, the sodium blocking agent is 4,7,13,16,21-pentaoxa-1,10-diazabicyclo[8.8.5]-tricosane.

[0091] The dephosphorylation of adenosine 3',5' bisphosphate (PAP) can be assessed by any suitable methods. For example, the dephosphorylation of adenosine 3',5' bisphosphate can be assessed by assessing consumption of the adenosine 3',5' bisphosphate in the dephosphorylation reaction or the formation of the AMP or P_i in the reaction.

[0092] Any suitable 3'(2'),5'-bisphosphate nucleotidase can be used, as disclosed in Section C. Any source or form known in the art that permits the production of P_i and AMP from PAP is contemplated. In particular, any suitable chimeric proteins, including the ones described in the above Section B, can be used in the present methods. In one example, the chimeric protein comprises the amino acid sequence set forth in SEQ ID NO:4. In another example, the chimeric protein is encoded by the nucleotide sequence set forth in SEQ ID NO:5.

[0093] In one embodiment, the formation of AMP can be assessed using a combination of 5'-nucleotidase, adenosine deaminase, purine nucleoside phosphorylase, xanthine oxidase, and peroxidase. For example, the following series of coupled enzymatic reactions, as detailed in Section C, can result in the production of detectable quinone dye. In one embodiment, ascorbate oxidase is also employed.

[0094] Any suitable 5'-nucleotidase, adenosine deaminase, purine nucleoside phosphorylase, and xanthine oxidase can be used. The enzymes can be derived from any source known in the art, including microbial and mammalian, that will permit the generation of a detectable product if adenosine 3',5'-bisphosphate is consumed by the 3'(2'),5'-bisphosphate nucleotidase.

[0095] Any suitable means for assessing H_2O_2 formation may be employed as disclosed in Section C. Any suitable peroxidase can be used. More preferably, a horseradish peroxidase is used. Exemplary peroxidases, Trinder reagents, and other chromagens are those in Section C. In one embodiment, the amount of quinone dye formed is assessed to determine the presence or amount of Li^+ ions. Here, the amount of quinone dye formed is inversely related to the amount of lithium ions in the sample.

[0096] Any suitable means for preparing the sample may be employed. In one embodiment, serum or plasma samples are treated with heparinate.

[0097] The sample can be contacted with the 3'(2'),5'-bisphosphate nucleotidase and the peroxidase sequentially or simultaneously. Likewise, any other enzymes used can be contacted with the 3'(2'),5'-bisphosphate nucleotidase sequentially or simultaneously in a fashion that permits the formation of a suitable detectable product.

[0098] Any suitable conditions for detection or measurement of sodium ions can be used. The reaction temperature is usually in the range from 10° C. to 40° C., with a preferred temperature of 37° C. The reaction time is preferably not more than 15 minutes, most preferably about 9 minutes or less.

[0099] Any suitable means of performing colorimetric analysis can be used. In one embodiment, the samples are analyzed for the presence of quinone dye in a Roche Cobas Mira Chemistry Analyzer.

[0100] If desirable, interference of the assay can be countered. For example, ascorbate interference can be countered using a copper (II) compound, a cholic acid or a bathophenanthroline disulphonic acid or a mixture thereof. Bilirubin interference can be countered using a ferrocyanide salt.

[0101] The present methods can be used to assay any suitable sample. Preferably, the sample is a biological sample. In one example, the sample is a blood sample, e.g., a plasma, serum, red blood cell or whole blood sample.

[0102] Any suitable chimeric proteins, including the ones described in the above Section B, can be used in the present methods. In one example, the chimeric protein comprises the amino acid sequence set forth in SEQ ID NO:4. In another example, the chimeric protein is encoded by the nucleotide sequence set forth in SEQ ID NO:5.

[0103] The present methods can be used for any suitable purpose. Preferably, the method used in the prognosis or

diagnosis of a disease or disorder. In one embodiment, the present methods are used to detect the presence or amount of lithium ions in a serum sample.

[0104] In one aspect, the present invention is directed to a kit for assaying for lithium ion in a sample, which kit comprises: a) a first composition comprising a lithium-sensitive 3'(2'),5'-bisphosphate nucleotidase; and b) a means for assessing the adenosine 3',5'-bisphosphate consumed or the AMP formed by the 3'(2'),5'-bisphosphate nucleotidase to determine the presence or amount of said lithium ions in the sample. In one embodiment, the kit further comprises a sodium blocking agent. In one embodiment, the first composition further comprises adenosine deaminase, 5'-nucleotidase and $MgCl_2$. In one embodiment, the kit further comprises a second composition comprising 4-aminoantipyrine (4-AA), N-ethyl-N-(2-hydroxy-3-sulfoxypropyl)-3-m-toluidine (EHSPT), purine nucleoside phosphorylase, xanthine oxidase, and peroxidase, wherein the reaction of 4-AA and EHSPT in the presence of peroxidase is the means for assessing the product formed if lithium ions are not present. The kit can also further comprise a low lithium serum standard, a medium lithium standard, and a high lithium serum standard.

[0105] Any suitable blocking agent may be used in the present methods. In a preferred embodiment, the sodium blocking agent is 4,7,13,16,21-pentaoxa-1,10-diazabicyclo[8.8.5]-tricosane.

[0106] The adenosine 3',5'-bisphosphate (PAP) to be used herein may be in any suitable form of a salt, so long as it contains no lithium ions. A preferred form is potassium salt.

[0107] The compositions of the present invention may be formulated into a reagent having a pH adjusted by the addition of a buffer to pH 6 to 9. Any suitable buffer may be used. It is contemplated that such buffers contain no sodium ions. Exemplary buffers are Good's buffer, 2-[N-morpholino]ethane-sulfonic acid (MES) buffer, and tris buffer.

[0108] The compositions of the present invention may further contain any surfactant, preservative, stabilizer, and enzyme activator. Preferred examples of the surfactant are Triton-100. Preferred examples of the preservative include Thimerosal. Any suitable stabilizer can be used. In one embodiment, the stabilizer is a protein. In a specific embodiment, the protein is bovine serum albumin. Any suitable enzyme activator can be used. In one embodiment, the activator is Mg^{2+} or a salt thereof, e.g., $MgCl_2$.

[0109] Any suitable concentration of 3'(2'),(5')-bisphosphate nucleotidase can be used in a composition for measurement of sodium ions. In a preferred embodiment, the concentration is in the range of 0.1-5 u/ml, more preferably, 0.5-3 u/ml, most preferably 2-3 u/ml. Any suitable concentration of 5'-nucleotidase can be used. In a preferred embodiment, the concentration is in the range of 0.1-5 u/ml, more preferably, 0.5-3 u/ml, most preferably 2-3 u/ml. Any suitable concentration of adenosine deaminase can be used. In a preferred embodiment, the concentration is in the range of 0.1-5 u/ml, more preferably, 0.5-3 u/ml, most preferably 2-3 u/ml. Any suitable concentration of xanthine oxidase can be used. In a preferred embodiment, the concentration is in the range of 0.1-5 u/ml, more preferably, 0.5-3 u/ml, most preferably 2-3 u/ml. Any suitable concentration of peroxidase can be used. In a preferred embodiment, the concentration is in the range of 1-50 u/ml, more preferably, 1-30 u/ml, most preferably 5-10 u/ml.

[0110] Any suitable chromagen may be employed, particularly those in Section C. The chromagen of the reduced type,

N-ethyl-N-(2-hydroxy-3-sulfopropyl)-3-m-toluidine (EHSPT), and 4-aminoantipyrine (4-AA), or a salt thereof are used at any concentration suitable for measurement. The chromagen of the reduced type is preferably used at a concentration in the range of 0.01 to 10 mM. The N-ethyl-N-(2-hydroxy-3-sulfopropyl)-3-m-toluidine (EHSPT) or salt thereof is preferably used at a concentration of 4 mM. 4-aminoantipyrine (4-AA) or salt thereof is preferably used at a concentration of 2 mM.

peroxide (H_2O_2). H_2O_2 generated reacts with N-Ethyl-N-(2-hydroxy-3-sulfopropyl)-3-m-toluidine (EHSPT) and 4-aminoantipyrine (4-AA) in the presence of peroxidase (POD) to form a quinone dye which had maximal absorbance at 556 nm. The rate of the quinone dye formation was inversely proportion to the concentration of lithium in serum samples. The enzymatic coupling reaction scheme is shown below in Table 3:

TABLE 3

PAP + H_2O		Phosphatase	AMP + Pi	
AMP		5'-Nucleotidase/ADA deaminase/PPN	Hypoxanthine + Pi + NH_3 + R-1-P	
Hypoxanthine + $2H_2O$ + $2O_2$		Xanthine Oxidase	Uric Acid + $2H_2O_2$	
H_2O_2 + 4-AA + EHSPT		Peroxidase	$4H_2O$ + Quinone dye (λ max 556 nm)	

PAP: 3'-phosphoadenosine 5'-phosphate (adenosine 3', 5'-bisphosphate)

AMP: Adenosine-5'-phosphate

PPN: Purine Nucleoside Phosphorylase

4-AA: 4-Aminoantipyrine

EHSPT: N-Ethyl-N-(2-Hydroxy-3-Sulfopropyl)-m-Toluidine

[0111] In some embodiments, standards for calibration of the assay are included. In one embodiment, a low lithium serum standard, a medium lithium standard, and a high lithium standard are included. Preferably, the low lithium serum standard comprises 0 mM of lithium in serum, and the medium lithium serum standard comprises 0.5-1.5 mM of lithium in serum, preferably 1 mM, and the high lithium serum standard comprises 2.5-3.5 mM of lithium, preferably 3.0 mM, in serum. In one embodiment, the presence or amount of lithium ions are calculated using a calibration curve. The amount of detectable chromagen is assessed at time 1 for a value of A_1 and at time 2 for a value of A_2 . The resultant value is calculate in the following equation: $\Delta A = A_2 - A_1$. A calibration curve is generated by plotting the ΔA values of the standards. The amount of lithium in the samples are then determined by plotting the sample ΔA value on the calibration curve. In one embodiment, time 1 is 6 minutes after the addition of means to assess P_i production and time 2 is 3 minutes after time 1.

E. EXAMPLES

Example 1

Sodium Ion Detection Assay Kit

[0112] Intended Use. The exemplary assay kit is for the quantitative in vitro determination of sodium in serum and plasma.

[0113] Assay Principle. Sodium was determined spectrophotometrically through a kinetic coupling assay system involving the chimeric 3'(2'),5'-bisphosphate nucleotidase (as described in Section B) whose activity was sensitive to sodium concentration ($IC_{50}=20$ mM). Through enzymatic coupling, the phosphatase substrate, adenosine 3',5'-bisphosphate (PAP) was converted to hypoxanthine by a series of enzymatic reactions to generate uric acid and hydrogen

[0114] Key Assay Characteristics. The sodium enzymatic assay was a two reagent (R1 and R2) based kinetic assay system. The results were obtained in 10 min by measuring absorbance at 550 nm. No off line pretreatment was needed. The assay had a wide measuring range from 80 to 180 mmol/L. The assay offered excellent precision as shown in the table below:

TABLE 4

	130M Na^+	150 mM Na^+
Intra-assay	CV % = 3.8%	CV % = 4.8%
Inter-assay	CV % = 4.2%	CV % = 4.1%

[0115]

TABLE 5

Reagents
Reagent 1. Buffer/enzyme/substrates
Enzyme/substrate lyophilized powder containing
Good's buffer, PAP, $MgCl_2$, 4-AA, Enzymes and stabilizers
Reagent 2. Buffer/protein/substrate
Enzyme/substrate lyophilized powder containing
Good's buffer, Enzymes, $MgCl_2$, and stabilizers
Low sodium Serum Standard
High sodium Serum Standard

[0116] Reagent Preparation. One vial of Reagent 1 (R1) was reconstituted with 50 ml distilled water. The reagents were mixed gently by inversion and then allowed to stand a minimum of 10 min in an ice bath before use. The reconstituted R1 solution was stable for 1 week at 2-8° C. One vial of Reagent 2 (R2) was reconstituted with 25 ml of distilled water. The reagents were gently by inversion and then allowed to stand a minimum of 10 min in an ice bath before

use. The reconstituted R2 solution was stable for 1 week at 2-8° C. Standards included were ready to use and were stable up to expiration date when stored under 2-8° C.

[0117] Normal Values. The normal Na⁺ values in serum are 136-146 mM (313-336 mg/dL).

[0118] Test samples. Test samples were serum or plasma treated with heparinate.


[0119] Assay Procedure.

[0120] 1. Reconstituted R1 and R2 reagents as described in Reagent Preparation section and kept the reconstituted R1 and R2 reagents on ice bath.

[0121] 2. In a cuvette, mixed 180 μL of Reagent R1 and 20 μL of serum sample and incubated at 37° C. for 1 minute and then added 85 μL of Reagent 2.

[0122] 3. Read absorbance (550 nm) at 3 min (2 min after addition of Reagent 2) as A_1 . Incubated for a further 7 min and read the absorbance at 10 min as A_2 .

[0123] 4. Calculated $\Delta A = A_2 - A_1$



[0124] Calibration. This assay was calibrated daily using the enclosed low and high sodium standards. A calibration curve was constructed by plotting the ΔA values of the standards against the corresponding sodium concentrations.

at 0.25 mM; triglyceride at 250 mg/dl; ascorbic acid at 5 mM; and bilirubin at 10 mg/dl.

REFERENCES

[0126] 1. N. Tietz. TEXTBOOK OF CLINICAL CHEMISTRY, p. 1841. W.B. Saunders Company, Philadelphia (1986)

[0127] 2. L. Heppel and R. Hilmo. *J. Bio. Chem.* 188, 665-676 (1951)

[0128] 3. J. Murguia, J. Belles, and R. Serrano. *Science* 267, 232-234 (1995)

Example 2

Lithium Ion Detection Assay

[0129] Intended Use. The exemplary assay kit was for the quantitative in vitro determination of lithium in serum and plasma.

[0130] Assay Principle. Lithium was determined spectrophotometrically through a kinetic coupling assay system involving the chimeric 3'(2'),5'-bisphosphate nucleotidase, as described in Section B, whose activity was sensitive to lithium concentration ($IC_{50}=0.1$ mM). Through enzymatic coupling, the phosphatase substrate, adenosine 3',5'-bisphosphate (PAP) was converted to hypoxanthine by a series of enzymatic reactions to generate uric acid and hydrogen peroxide (H_2O_2). The H_2O_2 generated reacted with N-Ethyl-N-(2-hydroxy-3-sulfopropyl)-3-m-toluidine (EHSPT) and 4-aminoantipyrine (4-AA) in the presence of peroxidase (POD) to form a quinone dye which had maximal absorbance at 556 nm. The rate of the quinone dye formation was inversely proportion to the concentration of lithium in serum samples. The enzymatic coupling reaction scheme is shown below in Table 6.

TABLE 6

PAP + H ₂ O	Phosphatase	AMP + P _i
AMP	5'-Nucleotidase/ADA deaminase/PNP	Hypoxanthine + P _i + NH ₃ + R-1-P
Hypoxanthine + 2H ₂ O + 2O ₂	Xanthine Oxidase	Uric Acid + 2H ₂ O ₂
H ₂ O ₂ + 4AA + EHSPT	Peroxidase	4H ₂ O + Quinone dye (λ max 556 nm)

PAP: 3'-phosphoadenosine 5'-phosphate (adenosine 3',5'-bisphosphate)

AMP: Adenosine-5'-phosphate

PNP: Purine Nucleoside Phosphorylase

4-AA: 4-Aminoantipyrine

EHSPT: N-Ethyl-N-(2-Hydroxy-3-Sulfopropyl)-m-Toluidin

The sodium concentration of the sample was read from the calibration curve. A representative calibration curve is shown in **FIG. 1**.

[0125] Interference. The assay was not interfered by the following substances at indicated concentrations: NH_4Cl at 0.5 mM; KPi at 1.5 mM; CaCl_2 at 5 mM; NaCl at 200 mM; KCl at 10 mM; CuCl_2 at 0.25 mM; FeCl_3 at 0.25 mM; ZnCl_2

[0131] Key Assay Characteristics. The lithium enzymatic assay was a two reagent (R1 and R2) based kinetic assay system. The results were obtained in 10 min by measuring absorbance at 550 nm. No off line pretreatment was needed. The assay had a wide measuring range from 0 to 3 mmol/L. The assay offered excellent precision as shown in Table 7 below:

TABLE 7

	1 mM Li ⁺	2 mM Li ⁺
Intra-assay	CV % = 3.5%	CV % = 4.5%
Inter-assay	CV % = 4.8%	CV % = 4.2%

[0132] Reagent Preparation. One vial of Reagent 1 (R1) was reconstituted with 25 ml distilled water. The reagent was mixed gently by inversion and then allowed to stand for a minimum of 10 min in ice bath before use. The reconstituted R1 solution was stable for 1 week at 2-8° C. One vial of Reagent 2 (R2) was reconstituted with 12.5 ml of distilled water. The reagent was mixed gently by inversion and then allowed to stand for a minimum of 10 min in ice bath before use. The reconstituted R2 solution was stable for 1 week at 2-8° C.

TABLE 8

Reagents
Reagent 1 Buffer/enzyme/substrates
Enzyme/substrate lyophilized powder containing
Good's buffer, PAP, MgCl ₂ , 4-AA, Enzymes and stabilizers
Reagent 2 Buffer/protein/substrate
Enzyme/substrate lyophilized powder containing
Good's buffer, enzymes, MgCl ₂ , and stabilizers
Low lithium Serum Standard
Med. lithium Serum Standard
High lithium Serum Standard

[0133] Normal Values. Typically, the desirable serum lithium levels are 0.6 to 1.2 mEq/l.

[0134] Test Samples. The test samples were serum or plasma treated with heparin. Plasma containing EDTA-Na should not be used.


[0135] Assay Procedure.

[0136] 1. Reconstituted R1 and R2 reagents as described in Reagent Preparation section and kept the reconstituted R1 and R2 reagents on ice bath.

[0137] 2. In a cuvette, mixed 180 μ L of Reagent R1 and 5 μ L of serum sample and incubated at 37° C. for 1 minute and then added 85 μ L of Reagent 2.

[0138] 3. Read absorbance (550 nm) at 6 minutes as A₁. Incubated for 3 more minutes and read the absorbance at 9 minute as A₂.

[0139] 4. Calculated $\Delta A = A_2 - A_1$



[0140] Calibration and Quality Control. The assay was calibrated daily using the enclosed low and high lithium standards. The calibration curve was constructed by plotting the ΔA values of the standards against the corresponding lithium concentrations. The lithium concentration of the sample was read from the calibration curve. The assay should be calibrated daily.

[0141] Interference. The assay was not interfered by the following substances at indicated concentrations: Na⁺200 mM, NH₄⁺0.5 mM, Ca²⁺4.0 mM, Mg²⁺2.0 mM, ascorbic acid 5.0 mM, 0.25 mM Zn²⁺, 0.25 mM Fe³⁺, 0.25 mM Cu²⁺, 10 mM K⁺, and bilirubin 45 mg/dl.

[0142] Performance Features. The assay had a linear range from 0.1-3.0 mM. The intra assay % CV was 3.5%, and the inter assay % CV was 4.5%.

REFERENCES

- [0143] 1. N. Tietz. TEXTBOOK OF CLINICAL CHEMISTRY, p. 1841. W.B. Saunders Company, Philadelphia (1986)
- [0144] 2. L. Heppel and R. Hilmoe. *J. Bio. Chem.* 188, 665-676 (1951)
- [0145] 3. J. Murguia, J. Belles, and R. Serrano. *Science* 267, 232-234 (1995)

[0146] The above examples are included for illustrative purposes only and are not intended to limit the scope of the invention. Many variations to those described above are possible. Since modifications and variations to the examples described above will be apparent to those of skill in this art, it is intended that this invention be limited only by the scope of the appended claims.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 18

<210> SEQ ID NO 1

<211> LENGTH: 12

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: 40%-100% identity to leader sequence

<400> SEQUENCE: 1

-continued

Met Gly Gly Ser Gly Asp Asp Asp Asp Leu Ala Leu
1 5 10

<210> SEQ ID NO 2
<211> LENGTH: 356
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: 40%-100% identity to the biphosphate
nucleotidase

<400> SEQUENCE: 2

Ala Leu Glu Arg Glu Leu Leu Val Ala Thr Gln Ala Val Arg Lys Ala
1 5 10 15
Ser Leu Leu Thr Lys Arg Ile Gln Ser Glu Val Ile Ser His Lys Asp
20 25 30
Ser Thr Thr Ile Thr Lys Asn Asp Asn Ser Pro Val Thr Thr Gly Asp
35 40 45
Tyr Ala Ala Gln Thr Ile Ile Ile Asn Ala Ile Lys Ser Asn Phe Pro
50 55 60
Asp Asp Lys Val Val Gly Glu Glu Ser Ser Ser Gly Leu Ser Asp Ala
65 70 75 80
Phe Val Ser Gly Ile Leu Asn Glu Ile Lys Ala Asn Asp Glu Val Tyr
85 90 95
Asn Lys Asn Tyr Lys Lys Asp Asp Phe Leu Phe Thr Asn Asp Gln Phe
100 105 110
Pro Leu Lys Ser Leu Glu Asp Val Arg Gln Ile Ile Asp Phe Gly Asn
115 120 125
Tyr Glu Gly Gly Arg Lys Gly Arg Phe Trp Cys Leu Asp Pro Ile Asp
130 135 140
Gly Thr Lys Gly Phe Leu Arg Gly Glu Gln Phe Ala Val Cys Leu Ala
145 150 155 160
Leu Ile Val Asp Gly Val Val Gln Leu Gly Cys Ile Gly Cys Pro Asn
165 170 175
Leu Val Leu Ser Ser Tyr Gly Ala Gln Asp Leu Lys Gly His Glu Ser
180 185 190
Phe Gly Tyr Ile Phe Arg Ala Val Arg Gly Leu Gly Ala Phe Tyr Ser
195 200 205
Pro Ser Ser Asp Ala Glu Ser Trp Thr Lys Ile His Val Arg His Leu
210 215 220
Lys Asp Thr Lys Asp Met Ile Thr Leu Glu Gly Val Glu Lys Gly His
225 230 235 240
Ser Ser His Asp Glu Gln Thr Ala Ile Lys Asn Lys Leu Asn Ile Ser
245 250 255
Lys Ser Leu His Leu Asp Ser Gln Ala Lys Tyr Cys Leu Leu Ala Leu
260 265 270
Gly Leu Ala Asp Val Tyr Leu Arg Leu Pro Ile Lys Leu Ser Tyr Gln
275 280 285
Glu Lys Ile Trp Asp His Ala Ala Gly Asn Val Ile Val His Glu Ala
290 295 300
Gly Gly Ile His Thr Asp Ala Met Glu Asp Val Pro Leu Asp Phe Gly
305 310 315 320
Asn Gly Arg Thr Leu Ala Thr Lys Gly Val Ile Ala Ser Ser Gly Pro

-continued

325	330	335
-----	-----	-----

Arg Glu Leu His Asp Leu Val Val Ser Thr Ser Cys Asp Val Ile Gln
 340 345 350

Ser Arg Asn Ala
 355

<210> SEQ ID NO 3
 <211> LENGTH: 17
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: 40%-100% identity to the second bacterial leader sequence

<400> SEQUENCE: 3

Lys Gly Glu Leu Glu Gly Leu Pro Ile Pro Asn Pro Leu Leu Arg Thr
1 5 10 15

Gly

<210> SEQ ID NO 4
 <211> LENGTH: 356
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Chimeric protein

<400> SEQUENCE: 4

Ala Leu Glu Arg Glu Leu Leu Val Ala Thr Gln Ala Val Arg Lys Ala
1 5 10 15
Ser Leu Leu Thr Lys Arg Ile Gln Ser Glu Val Ile Ser His Lys Asp
20 25 30
Ser Thr Thr Ile Thr Lys Asn Asp Asn Ser Pro Val Thr Thr Gly Asp
35 40 45
Tyr Ala Ala Gln Thr Ile Ile Ile Asn Ala Ile Lys Ser Asn Phe Pro
50 55 60
Asp Asp Lys Val Val Gly Glu Glu Ser Ser Ser Gly Leu Ser Asp Ala
65 70 75 80
Phe Val Ser Gly Ile Leu Asn Glu Ile Lys Ala Asn Asp Glu Val Tyr
85 90 95
Asn Lys Asn Tyr Lys Lys Asp Asp Phe Leu Phe Thr Asn Asp Gln Phe
100 105 110
Pro Leu Lys Ser Leu Glu Asp Val Arg Gln Ile Ile Asp Phe Gly Asn
115 120 125
Tyr Glu Gly Gly Arg Lys Gly Arg Phe Trp Cys Leu Asp Pro Ile Asp
130 135 140
Gly Thr Lys Gly Phe Leu Arg Gly Glu Gln Phe Ala Val Cys Leu Ala
145 150 155 160
Leu Ile Val Asp Gly Val Val Gln Leu Gly Cys Ile Gly Cys Pro Asn
165 170 175
Leu Val Leu Ser Ser Tyr Gly Ala Gln Asp Leu Lys Gly His Glu Ser
180 185 190
Phe Gly Tyr Ile Phe Arg Ala Val Arg Gly Leu Gly Ala Phe Tyr Ser
195 200 205
Pro Ser Ser Asp Ala Glu Ser Trp Thr Lys Ile His Val Arg His Leu
210 215 220

-continued

Lys Asp Thr Lys Asp Met Ile Thr Leu Glu Gly Val Glu Lys Gly His
 225 230 235 240

Ser Ser His Asp Glu Gln Thr Ala Ile Lys Asn Lys Leu Asn Ile Ser
 245 250 255

Lys Ser Leu His Leu Asp Ser Gln Ala Lys Tyr Cys Leu Leu Ala Leu
 260 265 270

Gly Leu Ala Asp Val Tyr Leu Arg Leu Pro Ile Lys Leu Ser Tyr Gln
 275 280 285

Glu Lys Ile Trp Asp His Ala Ala Gly Asn Val Ile Val His Glu Ala
 290 295 300

Gly Gly Ile His Thr Asp Ala Met Glu Asp Val Pro Leu Asp Phe Gly
 305 310 315 320

Asn Gly Arg Thr Leu Ala Thr Lys Gly Val Ile Ala Ser Ser Gly Pro
 325 330 335

Arg Glu Leu His Asp Leu Val Val Ser Thr Ser Cys Asp Val Ile Gln
 340 345 350

Ser Arg Asn Ala
 355

<210> SEQ ID NO 5

<211> LENGTH: 1176

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Nucleotide sequence encoding a chimeric protein

<400> SEQUENCE: 5

```

atgggcggat ccggtgatga cgatgacctc gcccttgcat tggaaagaga attattggtt    60
gcaactcaag ctgtacgaaa ggcgtcttta ttgactaaga gaattcaatc tgaagtgatt    120
tctcacaagg actccactac tattaccaag aatgataatt ctccagtaac cacagggtgat    180
tatgctgcac aaacgatcat cataaatgct atcaagagca attttcctga tgataaggta    240
gttggtgaag aatcctcatc aggattgagc gacgcattcg tctcaggaat tttaaacgaa    300
ataaaagcca atgacgaagt ttataacaag aattataaaa aggatgattt tctgtttaca    360
aacgatcagt ttccgctaaa atctttggag gacgtcaggc aaatcatcga tttcggcaat    420
tacgaagggtg gtgaaaaagg aagattttgg tgtttggatc ctattgacgg aaccaagggg    480
tttttaagag gtgaacagtt tgcagtatgt ctggccttaa ttgtggacgg tgtgtttcag    540
cttggttgta ttggatgccc caacttagtt ttaagttctt atggggccca agatttgaaa    600
ggccatgagt cattttggtta tatctttcgt gctgttagag gtttaggtgc cttctattct    660
ccatcttcag atgcagagtc atggaccaa atccacgta gacacttaaa agacactaaa    720
gacatgatta ctttagaggg agttgaaaaa ggacactcct ctcagatga acaaactgct    780
atcaaaaaa aactaaatat atccaaatct ttgcacttgg attctcaagc caagtactgt    840
ttgttagcat tgggcttagc agacgtatat ttacgtctgc ctatcaaaact ttcttaccaa    900
gaaaagatct gggaccatgc tgcaggcaac gttattgtcc atgaagctgg aggtatccat    960
acagatgcca tggaagatgt tcctctagac ttcggtaacg gtagaacgct agctacgaag   1020
ggagttatag cgtcaagtgg cccacgcgag ttacatgact tgggtggtgc tacatcatgc   1080
gatgtcattc agtcaagaaa cgccaagggc gagcttgaag gtttgcctat ccctaaccct   1140
ctcctccgta ccggtcatca tcaccatcac cattga                               1176

```

-continued

<210> SEQ ID NO 6
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Exemplary epitope tag

<400> SEQUENCE: 6

Asp Tyr Lys Asp Asp Lys
1 5

<210> SEQ ID NO 7
<211> LENGTH: 9
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Exemplary epitope tag

<400> SEQUENCE: 7

Tyr Pro Tyr Asp Val Pro Asp Tyr Ala
1 5

<210> SEQ ID NO 8
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Exemplary epitope tag

<400> SEQUENCE: 8

Cys Gln Asp Leu Pro Gly Asn Asp Asn Ser Thr
1 5 10

<210> SEQ ID NO 9
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Exemplary epitope tag

<400> SEQUENCE: 9

Glu Gln Lys Leu Ile Ser Glu Glu Asp Leu
1 5 10

<210> SEQ ID NO 10
<211> LENGTH: 6
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Exemplary epitope tag

<400> SEQUENCE: 10

His His His His His His
1 5

<210> SEQ ID NO 11
<211> LENGTH: 6
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Exemplary epitope tag

<400> SEQUENCE: 11

-continued

Asp Thr Tyr Arg Tyr Ile
1 5

<210> SEQ ID NO 12
<211> LENGTH: 6
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Exemplary epitope tag

<400> SEQUENCE: 12

Glu Tyr Met Pro Met Glu
1 5

<210> SEQ ID NO 13
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Exemplary epitope tag

<400> SEQUENCE: 13

Ala Ser Met Thr Gly Gly Gln Gln Met Gly Arg
1 5 10

<210> SEQ ID NO 14
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Exemplary epitope tag

<400> SEQUENCE: 14

Ser Phe Pro Gln Phe Lys Pro Gln Glu Ile
1 5 10

<210> SEQ ID NO 15
<211> LENGTH: 12
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Exemplary epitope tag

<400> SEQUENCE: 15

Lys Gly Phe Ser Tyr Phe Gly Glu Asp Leu Met Pro
1 5 10

<210> SEQ ID NO 16
<211> LENGTH: 6
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Exemplary epitope tag

<400> SEQUENCE: 16

Gln Tyr Pro Ala Leu Thr
1 5

<210> SEQ ID NO 17
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Exemplary epitope tag

-continued

<400> SEQUENCE: 17

Gln Arg Gln Tyr Gly Asp Val Phe Lys Gly Asp
 1 5 10

<210> SEQ ID NO 18

<211> LENGTH: 10

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Exemplary epitope tag

<400> SEQUENCE: 18

Glu Val His Thr Asn Gln Asp Pro Leu Asp
 1 5 10

1. An isolated chimeric protein, which chimeric protein comprises, from N-terminus to C-terminus:

- a) a first peptidyl fragment comprising a bacterial leader sequence from about to about 30 amino acid residues; and
- b) a second peptidyl fragment comprising a 3'(2'),5'-bispophosphate nucleotidase.

2. The isolated chimeric protein of claim 1, wherein the bacterial leader sequence is a leader sequence of an *E. coli* protein.

3. The isolated chimeric protein of claim 1, wherein the leader sequence has at least 40% identity to the amino acid sequence set forth in SEQ ID NO: 1 (MGGSGDDDLAL), in which the percentage identity is determined over an amino acid sequence of identical size to the amino acid sequence set forth in SEQ ID NO:1.

4. The isolated chimeric protein of claim 1, wherein the leader sequence binds to an antibody that specifically binds to an amino acid sequence set forth in SEQ ID NO: 1.

5. The isolated chimeric protein of claim 1, wherein the leader sequence comprises the amino acid sequence set forth in SEQ ID NO: 1.

6. The isolated chimeric protein of claim 1, wherein the first peptidyl fragment comprises about 20 amino acid residues.

7. The isolated chimeric protein of claim 1, wherein the nucleotidase is 3'(2'),5'-bispophosphate nucleotidase.

8. The isolated chimeric protein of claim 7, wherein the 3'(2'),5'-bispophosphate nucleotidase is of yeast, bacterial, or mammalian origin.

9. The isolated chimeric protein of claim 1, wherein the nucleotidase has at least 40% identity to the amino acid sequence set forth in SEQ ID NO:2 (ALER-ELLVATQAVRKASLLTKRIQSEVISHKD-STTTTKNDNSPVTG DYAAQTHIN AIKSNEFPDDKV-VGEESSGLSDAFVSGILNEIKANDEVYNKKNYKKD-DLFTNDQFP LKSLDVDRIIDFGNYEGGRKGRFW-CLDPIDGTGKGLRGEQFAVCLALIVDGVVQ LGCIGCPNLVSSYGAQDLKGHESFGYI-FRAVRGLGAFYSPSSDAESWTKIHRHL KDTKD-MITLEGVEKGHSSHDEQTAIKNKLNISK-SLHLD SQAKYCLLALGLADVYL RLPIKLSYQEKIWDHAAAGNVIVHEAG-GIHTDAMEDVPLDFGNGRTLATKGVIASS GPRELH-

DLVVSTSCDVIQSRNA), in which the percentage of identity is determined over an amino acid sequence of identical size to the amino acid sequence set forth in SEQ ID NO:2.

10. The isolated chimeric protein of claim 1, wherein the nucleotidase binds to an antibody that specifically binds to an amino acid sequence set forth in SEQ ID NO:2.

11. The isolated chimeric protein of claim 1, wherein the nucleotidase comprises the amino acid sequence set forth in SEQ ID NO:2.

12. The isolated chimeric protein of claim 1, wherein the first and second peptidyl fragments are linked via a cleavable linkage.

13. The isolated chimeric protein of claim 1, which further comprises, at its C-terminus, a third peptidyl fragment comprising a second bacterial leader sequence from about 5 to about 30 amino acid residues.

14. The isolated chimeric protein of claim 13, wherein the second bacterial leader sequence is a leader sequence of an *E. coli* protein.

15. The isolated chimeric protein of claim 13, wherein the second bacterial leader sequence has at least 40% identity to the amino acid sequence set forth in SEQ ID NO:3 (KGELEGLPIPNPLLR TG), in which the percentage identity is determined over an amino acid sequence of identical size to the amino acid sequence set forth in SEQ ID NO:3.

16. The isolated chimeric protein of claim 13, wherein the second bacterial leader sequence binds to an antibody that specifically binds to an amino acid sequence set forth in SEQ ID NO:3.

17. The isolated chimeric protein of claim 13, wherein the second bacterial leader sequence comprises the amino acid sequence set forth in SEQ ID NO:3.

18. The isolated chimeric protein of claim 13, wherein the third peptidyl fragment comprises about 20 amino acid residues.

19. The isolated chimeric protein of claim 1, which further comprises at its C-terminus, a third peptidyl fragment comprising a peptide tag.

20. The isolated chimeric protein of claim 19, wherein the peptide tag is selected from the group consisting of FLAG, HA HA1, c-Myc, 6-His, AU1, EE, T7, 4A6, ϵ , B, gE, and Ty1 tag.

21. The isolated chimeric protein of claim 13, which further comprises, at its C-terminus a fourth peptidyl fragment comprising a peptide tag.

22. The isolated chimeric protein of claim 21, wherein the peptide tag is selected from the group consisting of FLAG, HA HA1, c-Myc, 6-His, AU1, EE, T7, 4A6, ϵ , B, gE, and Ty1 tag.

23. The isolated chimeric protein of claim 1, which comprises the amino acid sequence set forth in SEQ ID NO:4

24. An isolated nucleic acid comprising a nucleotide sequence encoding the chimeric protein of claim 1.

25. An isolated nucleic acid comprising a nucleotide sequence encoding the chimeric protein of claim 23.

26. The nucleic acid of claim 24, which comprises the nucleotide sequence set forth in SEQ ID NO:5

```
(mgsgsgdddlalALERELLVATQAVRKASLLTKRIQSEVISHKDSTTITKNDNSPVTTG
DYAAQTII INAIKSNFPDDKVVGEESSSGLSDAFVSGILNEIKANDEVYNKNYKKD
DFLFTNDQFPLKSLEDVRQIIDFGNYEGGRKGRFWCLDPIDGTKGFLRGEQFAVCL
ALIVDGVVQLGCIGCPNLVLSYGAQDLKGHESFGYIFRAVRGLGAFYSPSSDAES
WTKIHVRHLKDTKDMITLEGVEKGHSSHDEQTAIKNKLNIKSLHLDSQAKYCLL
ALGLADVYLRLPIKLSYQEKIWDHAAGNVIVHEAGGIHTDAMEDVPLDFGNRT
LATKGVIASSGPRELHDLVVSTSCDVIQSRNAkgelelplipnpllrthghhhhh).
```

```
(atgggcgatccggtgatgacgatgacctcgcccttGCATTGGAAGAGAATTATTGGTTGCAACT
CAAGCTGTACGAAAGCGTCTTTATTGACTAAGAGAATTCAATCTGAAGTGAT
TTCTCACAAGGACTCCACTACTATTACCAAGAATGATAATTCTCCAGTAACCA
CAGGTGATTATGCTGCACAAACGATCATCATAAATGCTATCAAGAGCAATTTT
CCTGATGATAAGGTAGTTGGTGAAGAATCCTCATCAGGATTGAGCGACGCATT
CGTCTCAGGAATTTTAAACGAAATAAAGCCAATGACGAAGTTTATAACAAG
AATTATAAAAAGGATGATTTTCTGTTTACAAACGATCAGTTTCCGCTAAAATC
TTTGAGGACGTCAGGCAAAATCATCGATTTTCGCAATTACGAAGGTGGTAGAA
AAGGAAGATTTGGTGTTTGGATCCTATTGACGGAACCAAGGGTTTTAAGA
GGTGAACAGTTTGCAGTATGCTGGCCTTAATTGTGGACGGTGTGTTTCAGCTT
GGTTGTATTGGATGCCCCAACTTAGTTTAAAGTTCTTATGGGGCCCAAGATTTG
AAAGGCCATGAGTCATTTGGTTATATCTTTTCGTGCTGTTAGAGGTTTAGTGCC
TTCTATTCTCCATCTTCAGATGCAGAGTCATGGACCAAAATCCACGTTAGACA
CTTAAAAGACACTAAAGACATGATTACTTTAGAGGGAGTTGAAAAGGGACAC
TCCTCTCATGATGAACAACTGCTATCAAAAACAACTAAATATATCCAAATC
TTTGCACTTGGATTCTCAAGCCAAGTACTGTTTGTAGCATTGGGCTTAGCAGA
CGTATATTACGTCTGCCTATCAAACTTTCTTACCAAGAAAAGATCTGGGACC
ATGCTGCAGGCAACGTTATTGTCCATGAAGCTGGAGGTATCCATACAGATGCC
ATGGAAGATGTTCTCTAGACTTCGGTAACGGTAGAACGCTAGCTACGAAGGG
AGTTATAGCGTCAAGTGGCCACGCGAGTTACATGACTTGGTGGTGTCTACAT
CATGCGATGTCATTGAGTCAAGAAACGCCAagggcgagcttgaaggtttgcctatccctaacctctc
ctccgtaccgggtcatcatcaccatcaccattga).
```

27. An isolated nucleic acid comprising a nucleotide sequence complementary to the nucleotide sequence of claim 24.

28. A recombinant cell containing the nucleic acid of claim 24.

29. A method of producing a chimeric protein comprising growing a recombinant cell containing the nucleic acid of claim 24 such that the encoded chimeric protein is expressed by the cell, and recovering the expressed chimeric protein.

30. The product of the method of claim 29.

31. A method for assaying for sodium ions in a sample, which method comprises:

a) contacting the sample with a sodium-sensitive 3'(2'), 5'-bisphosphate nucleotidase, wherein the nucleotidase consumes adenosine 3',5'-bisphosphate (PAP) and forms AMP and P_i; and

b) assessing the consumption of PAP or the formation of AMP and P_i in step a) to determine the presence or amount of sodium ions in the sample.


32. The method of claim 31, wherein the sample is a biological sample.

33. The method of claim 32, wherein the biological sample is a blood sample.

34. The method of claim 33, wherein the blood sample is a plasma, serum, red blood cell, or whole blood sample.

35. The method of claim 31, wherein the 3'(2'),5'-bisphosphate nucleotidase is the chimeric protein of claim 1.

36. The method of claim 31, wherein the 3'(2'),5'-bisphosphate nucleotidase is an enzyme that catalyzes the following reaction:



37. The method of claim 31, wherein the amount of AMP formed is inversely related to the amount of sodium ions in the sample.

38. The method of claim 31, which is used in prognosis or diagnosis of a disease or disorder.

39. A method for assaying for sodium ions in a sample, which method comprises:

a) contacting the sample with a first composition comprising adenosine 3',5'-bisphosphate (PAP);

b) contacting the sample with a second composition comprising a sodium-sensitive 3'(2'),5'-bisphosphate nucleotidase; and

c) assessing the production of AMP to determine the presence or amount of sodium ions in the sample.

40. The method of claim 39, wherein the sample is a biological sample.

41. The method of claim 40, wherein the biological sample is a blood sample.

42. The method of claim 41, wherein the blood sample is a plasma, serum, red blood cell, or whole blood sample.

43. The method of claim 39, wherein the 3'(2'),5'-bisphosphate nucleotidase is the chimeric protein of claim 1.

44. The method of claim 39, wherein the first composition further comprises 4-aminoantipyrine (4-AA), N-ethyl-N-(2-hydroxy-3-sulfopropyl)-3-m-toluidine (EHSPT), purine nucleoside phosphorylase, xanthine oxidase, and peroxi-

dase, and the second composition further comprises adenosine deaminase, 5'-nucleotidase, and MgCl₂.

45. A kit of assaying for sodium ions in a sample, which kit comprises

a) a first composition comprising a sodium-sensitive 3'(2'),5'-bisphosphate nucleotidase that consumes adenosine 3',5'-bisphosphate and forms AMP and P_i; and

b) means for assessing the product formed or the substrate consumed by the nucleotidase to determine the presence or amount of the sodium ions in the sample.

46. The kit of claim 45, wherein the first composition further comprises adenosine deaminase, 5'-nucleotidase and MgCl₂.

47. The kit of claim 45, further comprising a second composition comprising 4-aminoantipyrine (4-AA), N-ethyl-N-(2-hydroxy-3-sulfopropyl)-3-m-toluidine (EHSPT), purine nucleoside phosphorylase, xanthine oxidase, and peroxidase, wherein the reaction of 4-AA and EHSPT in the presence of peroxidase is the means for assessing the product formed if sodium ions are not present.

48. The kit of claim 45, which further comprises a low sodium serum standard and a high sodium serum standard.

49. The kit of claim 45, wherein the 3'(2'),5'-bisphosphate nucleotidase is the chimeric protein of claim 1.

50. A method for assaying for lithium ions in a sample, which method comprises:

a) contacting the sample with a lithium-sensitive 3'(2'), 5'-bisphosphate nucleotidase, wherein the nucleotidase consumes adenosine 3',5'-bisphosphate (PAP) and forms AMP and P_i; and

b) assessing the amount of PAP consumed or AMP formed in step b) to determine the presence or absence of lithium ions in the sample.

51. The method of claim 50 further comprising first contacting the sample with a sodium blocking agent.

52. The method of claim 51, wherein the sodium blocking agent is 4,7,13,16,21-pentaoxa-1,10-diazabicyclo[8.8.5]-tricosane.


53. The method of claim 51, wherein the sample is a biological sample.

54. The method of claim 53, wherein the biological sample is a blood sample.

55. The method of claim 54, wherein the blood sample is a plasma, serum, red blood cell, or whole blood sample.

56. The method of claim 51, wherein the 3'(2'),5'-bisphosphate nucleotidase is the chimeric protein of claim 1.

57. The method of claim 51, wherein the nucleotidase is an enzyme that catalyzes the following reaction:



58. The method of claim 51, wherein the amount of AMP formed is inversely correlated to the amount of lithium ions in the sample.

59. The method of claim 51, which is used in prognosis or diagnosis of a disease or disorder.

60. A method for assaying for lithium ions in a sample, which method comprises:

- a) contacting the sample with a first composition comprising an adenosine 3',5'-bisphosphate (PAP);
 - b) contacting the sample with a second composition comprising a lithium-sensitive 3'(2'),5'-bisphosphate nucleotidase; and
 - c) assessing the production of a detectable product to determine the presence or absence of lithium ions in the sample.
- 61.** The method of claim 60 further comprising first contacting the sample with a sodium blocking agent.
- 62.** The method of claim 61, wherein the sodium blocking agent is 4,7,13,16,21-pentaoxa-1,10-diazabicyclo[8.8.5]-tricosane.
- 63.** The method of claim 60, wherein the sample is a biological sample.
- 64.** The method of claim 63, wherein the biological sample is a blood sample.
- 65.** The method of claim 64, wherein the blood sample is a plasma, serum, red blood cell, or whole blood sample.
- 66.** The method of claim 60, wherein the 3'(2'),5'-bisphosphate nucleotidase is the chimeric protein of claim 1.
- 67.** The method of claim 60, wherein the first composition further comprises 4-aminoantipyrine (4-AA), N-ethyl-N-(2-hydroxy-3-sulfo-3-propyl)-3-m-toluidine (EHSPT), purine nucleoside phosphorylase, xanthine oxidase, and peroxidase, and the second composition further comprises adenosine deaminase, 5'-nucleotidase, and $MgCl_2$.
- 68.** A kit for assaying for lithium ion in a sample, which kit comprises:
- a) a first composition comprising a lithium-sensitive 3'(2'),5'-bisphosphate nucleotidase; and
 - b) a means for assessing the adenosine 3',5'-bisphosphate consumed or the AMP formed by the 3'(2'),5'-bisphosphate nucleotidase to determine the presence or amount of said lithium ions in the sample.
- 69.** The kit of claim 68 further comprising a sodium blocking agent.
- 70.** The kit of claim 68, wherein the first composition further comprises adenosine deaminase, 5'-nucleotidase and $MgCl_2$.
- 71.** The kit of claim 68, further comprising a second composition comprising 4-aminoantipyrine (4-AA), N-ethyl-N-(2-hydroxy-3-sulfo-3-propyl)-3-m-toluidine (EHSPT), purine nucleoside phosphorylase, xanthine oxidase, and peroxidase, wherein the reaction of 4-AA and EHSPT in the presence of peroxidase is the means for assessing the product formed if lithium ions are not present.
- 72.** The kit of claim 68, which further comprises a low lithium serum standard, a medium lithium sodium standard, and a high lithium serum standard.
- * * * * *

专利名称(译)	使用离子敏感酶测定离子		
公开(公告)号	US20050142606A1	公开(公告)日	2005-06-30
申请号	US10/665883	申请日	2003-09-19
[标]申请(专利权)人(译)	袁充盛		
申请(专利权)人(译)	袁庄胜		
当前申请(专利权)人(译)	袁庄胜		
[标]发明人	YUAN CHONG SHENG		
发明人	YUAN, CHONG-SHENG		
IPC分类号	A61B C07K14/705 C12N9/22 G01N33/53 G01N33/84		
CPC分类号	G01N33/84		
其他公开文献	US8187831		
外部链接	Espacenet USPTO		

摘要(译)

本发明一般涉及钠和锂离子检测领域。特别地，本发明提供了嵌合蛋白，编码嵌合蛋白的核酸，用于测定样品中钠离子和锂离子的方法和试剂盒，尤其是使用3' (2')，5'-二磷酸核苷酸酶。

