



US 20080267958A1

(19) **United States**

(12) **Patent Application Publication**
Jalkanen et al.

(10) **Pub. No.: US 2008/0267958 A1**

(43) **Pub. Date: Oct. 30, 2008**

(54) **COMMON LYMPHATIC ENDOTHELIAL AND VASCULAR ENDOTHELIAL RECEPTOR-1 (CLEVER-1) AND USES THEREOF**

(60) Provisional application No. 60/346,288, filed on Jan. 9, 2002.

Publication Classification

(75) Inventors: **Sirpa Jalkanen**, Piispanristi (FI); **Heikki Irjala**, Turku (FI); **Marko Salmi**, Turku (FI)

(51) **Int. Cl.**
C07K 16/28 (2006.01)
A61K 39/395 (2006.01)
C07K 16/30 (2006.01)
G01N 33/53 (2006.01)
C12N 5/06 (2006.01)
G01N 33/574 (2006.01)
C07K 14/705 (2006.01)

Correspondence Address:
STERNE, KESSLER, GOLDSTEIN & FOX P.L.
L.C.
1100 NEW YORK AVENUE, N.W.
WASHINGTON, DC 20005 (US)

(52) **U.S. Cl.** **424/133.1**; 530/395; 530/387.1; 530/388.22; 435/7.24; 435/7.23; 435/7.2; 435/325; 424/174.1; 424/143.1

(73) Assignee: **Faron Pharmaceuticals Oy**

(57) **ABSTRACT**


(21) Appl. No.: **12/073,451**

A novel protein Common Lymphatic Endothelial and Vascular Endothelial Receptor-1 (CLEVER-1) is described. CLEVER-1 mediates leukocyte and malignant cell binding to vascular and lymphoid endothelial cells. CLEVER-1 is the first protein that has been reported to mediate both influx into and efflux from the lymph nodes. Also provided are methods of treating inflammation and preventing metastasis of malignant cells by providing an inhibitor of CLEVER-1 binding.

(22) Filed: **Mar. 5, 2008**

Related U.S. Application Data

(62) Division of application No. 10/497,991, filed on Nov. 30, 2004, now Pat. No. 7,354,577, filed as application No. PCT/FI03/00010 on Jan. 8, 2003.






Fig. 1





Fig. 2

A Lymphocyte binding to HEVs:



B Lymphocyte binding to lymph endothelium:





Fig. 3

A Tumor cell binding to HEVs:



B Tumor cell binding to lymphatic endothelium:




Fig. 4




Fig. 5




Fig. 6




Fig. 7




Fig. 8

actctgtcct ggacagcgtg cccaccagcc atg gcg ggg ccc cgg ggc ctc ctc 54
cca ctc tgc ctc ctg gcc ttc tgc ctg gca ggc ttc agc ttc gtc agg 102
ggg cag gtg ctg ttc aaa ggc tgt gat gtg aaa acc acg ttt gtc act 150
cat gta ccc tgc acc tcg tgc gcg gcc atc aag aag cag acg tgt ccc 198
tca ggc tgg ctg cgg gag ctc ccg gat cag ata acc cag gac tgc cgc 246
tac gaa gta cag ctg ggg ggc tct atg gtg tcc atg agc ggc tgc aga 294
cgg aag tgc cgg aag caa gtg gtg cag aag gcc tgc tgc cct ggc tac 342
tgg ggt tcc cgg tgc cat gaa tgc cct ggg ggc gct gag acc cca tgc 390
aat ggc cac ggg acc tgc ttg gat ggc atg gac agg aat ggg acc tgt 438
gtg tgc cag gaa aac ttc cgc ggc tca gcc tgc cag gag tgc caa gac 486
ccc aac cgg ttc ggg cct gac tgc caa tcg gtg tgc agc tgt gtg cac 534
gga gtg tgc aac cat ggg cca cgt ggg gat gga agc tgc ctg tgc ttt 582
gct gga tac act ggc ccc cac tgt gat caa gag ctg ccc gtc tgc cag 630
gag ctg cgc tgt ccc cag aac acc cag tgc tcc gca gag gct ccc agc 678
tgc agg tgc ctg ccc ggc tac aca cag cag ggc agt gaa tgc cga gcc 726
ccc aac ccc tgc tgg cca tca ccc tgc tca ctg ctg gcc cag tgc tcg 774
gtg agc ccc aag ggg cag gct cag tgt cac tgc cct gag aac tac cat 822
ggc gat ggg atg gtg tgt ctg ccc aag gac cca tgc act gac aac ctt 870
ggt ggc tgc ccc agc aac tct act ttg tgt gtg tac cag aag ccg ggc 918
cag gcc ttc tgc acc tgc cgg cca ggc ctg gtc agc atc aac agc aac 966
gct tct gcg ggc tgc ttc gcc ttc tgc tcc ccc ttc tcc tgc gac cgg 1014
tct gcc act tgc cag gtg acc gct gat ggg aag acc agc tgt gtg tgc 1062
agg gaa agc gag gtg ggg gat ggg cgt gcc tgc tac gga cac ctg ctc 1110
cac gag gtg cag aag gcc acg cag aca ggc cgg gtg ttc ctg cag ctg 1158
agg gtc gcc gtg gcc atg atg gac cag ggc tgc cgg gaa atc ctt acc 1206
aca gcg ggc cct ttc acc gtg ctg gtg cca tcc gtc tcc tcc ttc tcc 1254
tcc agg acc atg aat gca tcc ctt gcc cag cag ctc tgt aga cag cac 1302
atc atc gca ggg cag cac atc ctg gag gac aca agg acc caa caa aca 1350
cga agg tgg tgg acg ctg gcc ggg cag gag atc acc gtc acc ttt aac 1398
caa ttc acg aaa tac tcc tac aag tac aaa gac cag ccc cag cag acg 1446
ttc aac atc tac aag gcc aac aac ata gca gct aat ggc gtc ttc cac 1494
gtg gtc act ggc ctg cgg tgg cag gcc ccc tct ggg acc cct ggg gat 1542
ccc aag aga act atc gga cag atc ctc gcc tct acc gag gcc ttc agc 1590
cgc ttt gaa acc atc ctg gag aac tgt ggg ctg ccc tcc atc ctg gac 1638

Fig. 9

gga cct ggg ccc ttc aca gtc ttt gcc cca agc aat gag gct gtg gac 1686
agc ttg cgt gac ggc cgc ctg atc tac ctc ttc aca gcg ggt ctc tct 1734
aaa ctg cag gag ttg gtg cgg tac cac atc tac aac cac ggc cag ctg 1782
acc gtt gag aag ctc atc tcc aag ggt cgg atc ctc acc atg gcg aac 1830
cag gtc ctg gct gtg aac att tct gag gag ggg cgc atc ctg ctg gga 1878
ccc gag ggg gtc ccg ctg cag agg gta gac gtg atg gcc gcc aat ggt 1926
gtg atc cac atg ctg gac ggc atc ctg ctg ccc ccg acc atc ctg ccc 1974
atc ctg ccc aag cac tgc agc gag gag cag cac aag att gtg gcg ggc 2022
tcc tgt gtg gac tgc caa gcc ctg aac acc agc acg tgt ccc ccc aac 2070
agt gtg aag ctg gac atc ttc ccc aag gag tgt gtc tac atc cat gac 2118
cca acg ggg ctc aat gtg cta aag aag ggc tgt gcc agc tac tgc aac 2166
caa acc atc atg gaa caa ggc tgc tgc aaa ggt ttt ttc ggg cct gac 2214
tgc acg cag tgt cct ggg ggc ttc tcc aac ccc tgc tat ggc aaa ggc 2262
aat tgc agt gat ggg atc cag ggc aat ggg gcc tgc ctc tgc ttc cca 2310
gac tac aag ggc atc gcc tgc cac atc tgc tgc aac cca aac aag cat 2358
gga gag caa tgc cag gaa gac tgc ggc tgt gtc cat ggt ctc tgc gac 2406
aac cgc cca ggc agt ggg ggg gtg tgc cag cag ggc acg tgt gcc cct 2454
ggc ttc agt ggc cgg ttc tgc aac gag tcc atg ggg gac tgt ggg ccc 2502
aca ggg ctg gcc cag cac tgc cac ctg cat gcc cgc tgt gtt agc cag 2550
gag ggt gtt gcc aga tgt cgc tgt ctt gat ggc ttt gag ggt gat ggc 2598
ttc tcc tgc aca cct agc aac ccc tgc tcc cac ccg gac cgt gga ggc 2646
tgc tca gag aat gct gag tgt gtc cct ggg tcc ctg ggc acc cac cac 2694
tgc aca tgc cac aaa ggc tgg agt ggg gat ggc cgc gtc tgt gtg gct 2742
att gac gag tgt gag ctg gac gtg aga ggt ggc tgc cac acc gat gcc 2790
ctc tgc agc tat gtg ggc ccc ggg cag agc cga tgc acc tgc aag ctg 2838
ggc ttt gcc ggg gat ggc tac cag tgc agc ccc atc gac ccc tgc cgg 2886
gca ggc aat ggc ggc tgc cac ggc ctg gcc acc tgc cgg gca gtg ggg 2934
gga ggt cag cgg gtc tgc acg tgc ccc cct ggc ttt ggg ggt gat ggc 2982
ttc agc tgt tat gga gac atc ttc cgg gag ctg gag gca aat gcc cac 3030
ttc tcc atc ttc tac caa tgg ctt aag agt gcc ggc atc acg ctt 3075
cct gcc gac cgc cga gtc aca gcc ctg gtg ccc tcc gag gct gca 3120
gtc cgt cag ctg agc ccc gag gac cga gct ttc tgg ctg cag cca 3165
agg acg ctg ccg aac ctg gtc agg gcc cat ttt ctc cag ggt gcc 3210
ctc ttc gag gag gag ctg gcc cgg ctg ggt ggg cag gaa gtg gcc 3255
acc ctg aac ccc acc aca cgc tgg gag att cgc aac att agt ggg 3300

Fig. 9 (cont.)

agg gtc tgg gtg cag aat gcc agc gtg gat gtg gct gac ctc ctt 3345
gcc acc aac ggt gtc cta cac atc ctc agc cag gtc tta ctg ccc 3390
ccc cga ggg gat gtg ccc ggt ggg cag ggg ttg ctg cag cag ctg 3435
gac ttg gtg cct gcc ttc agc ctc ttc cgg gaa ttg ctg cag cac 3480
cat ggg ttg gtg ccc cag att gag gct gcc act gcc tac acc atc 3525
ttt gtg ccc acc aac cgc tcc ctg gag gcc cag ggc aac agc agt 3570
cac ctg gac gca gac aca gtg cgg cac cat gtg gtc ctg ggg gag 3615
gcc ctc tcc atg gaa acc ctg cgg aag ggt gga cac cgc aac tcc 3660
ctc ctg ggc cct gcc cac tgg atc gtc ttc tac aac cac agt ggc 3705
cag cct gag gtg aac cat gtg cca ctg gaa ggc ccc atg ctg gag 3750
gcc cct ggc cgc tcg ctg att ggt ctg tcg ggg gtc ctg acg gtg 3795
ggc tca agt cgc tgc ctg cat agc cac gct gag gcc ctg cgg gag 3840
aaa tgt gta aac tgc acc agg aga ttc cgc tgc act cag ggc ttc 3885
cag ctg cag gac aca ccc agg aag agc tgt gtc tac cga tct ggc 3930
ttc tcc ttc tcc cgg ggc tgc tct tac aca tgt gcc aag aag atc 3975
cag gtg ccg gac tgc tgc cct ggt ttc ttt ggc acg ctg tgt gag 4020
cca tgc cca ggg ggt cta ggg ggg gtg tgc tca ggc cat ggg cag 4065
tgc cag gac agg ttc ctg ggc agc ggg gag tgc cac tgc cac gag 4110
ggc ttc cat gga acg gcc tgt gag gtg tgt gag ctg ggc cgc tac 4155
ggg ccc aac tgc acc gga gtg tgt gac tgt gcc cat ggg ctg tgc 4200
cag gag ggg ctg caa ggg gac gga agc tgt gtc tgt aac gtg ggc 4245
tgg cag ggc ctc cgc tgt gac cag aaa atc acc agc cct cag tgc 4290
cct agg aag tgc gac ccc aat gcc aac tgc gtg cag gac tcg gcc 4335
gga gcc tcc acc tgc gcc tgt gct gcg gga tac tcc ggc aat ggc 4380
atc ttc tgt tca gag gtg gac ccc tgc gcc cac ggc cat ggg ggc 4425
tgc tcc cct cat gcc aac tgt acc aag gtg gca cct ggg cag cgg 4470
aca tgc acc tgc cag gat ggc tac atg ggc gac ggg gag ctg tgc 4515
cag gaa att aac agc tgt ctc atc cac cac ggg ggc tgc cac att 4560
cac gcc gag tgc atc ccc act ggc ccc cag cag gtc tcc tgc agc 4605
tgc cgt gag ggt tac agc ggg gat ggc atc cgg acc tgc gag ctc 4650
ctg gac ccc tgc tct aag aac aat gga gga tgc agc cca tat gcc 4695
acc tgc aaa agc aca ggg gat ggc cag agg aca tgt acc tgc gac 4740
aca gcc cac acc gtg ggg gac ggc ctc acc tgc cgt gcc cga gtc 4785
ggc ctg gag ctc ctg agg gat aag cat gcc tca ttc ttc agc ctc 4830
cgc ctc ctg gaa tat aag gag ctc aag ggc gat ggg cct ttc acc 4875

Fig. 9 (cont.)

atc ttc gtg ccg cac	gca gat cta atg agc	aac ctg tcg cag gat	4920
gag ctg gcc cgg att	cgt gcg cat cgc cag	ctg gtg ttt cgc tac	4965
cac gtg gtt ggc tgt	cgg cgg ctg cgg agc	gag gac ctg ctg gag	5010
cag ggg tac gcc acg	gcc ctc tca ggg cac	cca ctg cgc ttc agc	5055
gag agg gag ggc agc	ata tac ctc aat gac	ttc gcg cgc gtg gtg	5100
agc agc gac cat gag	gcc gtg aac ggc atc	ctg cac ttc att gac	5145
cgt gtc ctg ctg ccc	ccc gag gcg ctg cac	tgg gag cct gat gat	5190
gct ccc atc ccg agg	aga aat gtc acc gcc	gcc gcc cag ggc ttc	5235
ggg tac aag atc ttc	agc ggc ctc ctg aag	gtg gcc ggc ctc ctg	5280
ccc ctg ctt cga gag	gca tcc cat agg ccc	ttc aca atg ctg tgg	5325
ccc aca gac gcc gcc	ttt cga gct ctg cct	ccg gat cgc cag gcc	5370
tgg ctg tac cat gag	gac cac cgt gac aag	cta gca gcc att ctg	5415
cgg ggc cac atg att	cgc aat gtc gag gcc	ttg gca tct gac ctg	5460
ccc aac ctg ggc cca	ctt cga acc atg cat	ggg acc ccc atc tct	5505
ttc tcc tgc agc cga	acg cgg ccc ggt gag	ctc atg gtg ggt gag	5550
gat gat gct cgc att	gtg cag cgg cac ttg	ccc ttt gag ggt ggc	5595
ctg gcc tat ggc atc	gac cag ctg ctg gag	cca cct ggc ctt ggt	5640
gct cgc tgt gac cac	ttt gag acc cgg ccc	ctg cga ctg aac acc	5685
tgc agc atc tgt ggg	ctg gag cca ccc tgt	cct gag ggg tca cag	5730
gag cag ggc agc cct	gag gcc tgc tgg cgc	ttc tac ccg aag ttc	5775
tgg acg tcc cct ccg	ctg cac tct ttg gga	tta cgc agc gtc tgg	5820
gtc cac ccc agc ctt	tgg ggt agg ccc caa	ggc ctg ggc agg ggc	5865
tgc cac cgc aat tgt	gtc acc acc acc tgg	aag ccc agc tgc tgc	5910
cct ggt cac tat ggc	agt gag tgc caa gct	tgc cct ggc ggc ccc	5955
agc agc cct tgt agt	gac cgt ggc gtg tgc	atg gac ggc atg agt	6000
ggc agt ggg cag tgt	ctg tgc cgt tca ggt	ttt gct ggg aca gcc	6045
tgt gaa ctc tgt gct	cct ggt gcc ttt ggg	ccc cat tgt caa gcc	6090
tgc cgc tgc act gtg	cat ggc cgc tgt gat	gag ggc ctt ggg ggc	6135
tct ggc tcc tgc ttc	tgt gat gaa ggc tgg	act ggg cca cgc tgt	6180
gag gtg caa ctg gag	ctg cag cct gtg tgt	acc cca ccc tgt gca	6225
ccc gag gct gtg tgc	cgt gca ggc aac agc	tgt gag tgc agc ctg	6270
ggc tat gaa ggg gat	ggc cgc gtg tgt aca	gtg gca gac ctg tgc	6315
cag gac ggg cat ggt	ggc tgc agt gag cac	gcc aac tgt agc cag	6360
gta gga aca atg gtc	act tgt acc tgc ctg	ccc gac tac gag ggt	6405
gat ggc tgg agc tgc	cgg gcc cgc aac ccc	tgc aca gat ggc cac	6450

Fig. 9 (cont.)

cgc ggg ggc tgc agc gag cac gcc aac tgc ttg agc acc ggc ctg 6495
aac aca cgg cgc tgt gag tgc cac gca ggc tac gta ggc gat gga 6540
ctg cag tgt ctg gag gag tcg gaa cca cct gtg gac cgc tgc ttg 6585
ggc cag cca ccg ccc tgc cac tca gat gcc atg tgc act gac ctg 6630
cac ttc cag gag aaa cgg gct ggc gtt ttc cac ctc cag gcc acc 6675
agc ggc cct tat ggt ctg aac ttt tcg gag gct gag gcg gca tgc 6720
gaa gca cag gga gcc gtc ctt gct tca ttc cct cag ctc tct gct 6765
gcc cag cag ctg ggc ttc cac ctg tgc ctc atg ggc tgg ctg gcc 6810
aat ggc tcc act gcc cac cct gtg gtt ttc cct gtg gcg gac tgt 6855
ggc aat ggt cgg gtg ggc gta gtc agc ctg ggt gcc cgc aag aac 6900
ctc tca gaa cgc tgg gat gcc tac tgc ttc cgt gtg caa gat gtg 6945
gcc tgc cga tgc cga aat ggc ttc gtg ggt gac ggg atc agc acg 6990
tgc aat ggg aag ctg ctg gat gtg ctg gct gcc act gcc aac ttc 7035
tcc acc ttc tat ggg atg cta ttg ggc tat gcc aat gcc acc cag 7080
cgg ggt ctc gac ttc ctg gac ttc ctg gat gat gag ctc acg tat 7125
aag aca ctc ttc gtc cct gtc aat gaa ggc ttt gtg gac aac atg 7170
acg ctg agt ggc cca gac ttg gag ctg cat gcc tcc aac gcc acc 7215
ctc cta agt gcc aac gcc agc cag ggg aag ttg ctt ccg gcc cac 7260
tca ggc ctc agc ctc atc atc agt gac gca ggc cct gac aac agt 7305
tcc tgg gcc cct gtg gcc cca ggg aca gtt gtg gtt agc cgt atc 7350
att gtg tgg gac atc atg gcc ttc aat ggc atc atc cat gct ctg 7395
gcc agc ccc ctc ctg gca ccc cca cag ccc cag gca gtg ctg gcg 7440
cct gaa gcc cca cct gtg gcg gca ggc gtg ggg gct gtg ctt gcc 7485
gct gga gca ctg ctt ggc ttg gtg gcc gga gct ctc tac ctc cgt 7530
gcc cga ggc aag ccc acg ggc ttt ggc ttc tct gcc ttc cag gcg 7575
gaa gat gat gct gac gac gac ttc tca ccg tgg caa gaa ggg acc 7620
aac ccc acc ctg gtc tct gtc ccc aac cct gtc ttt ggc agc gac 7665
acc ttt tgt gaa ccc ttc gat gac tca ctg ctg gag gag gac ttc 7710
cct gac acc cag agg atc ctc aca gtc aag tga cggagget ggggctgaaa 7760
gcagaagcat gcacagggag gagaccactt ttattgcttg tctgggtgga tggggcagga 7820
ggggctgagg gcctgtccca gacaataaag tgcctcagc ggatgtgggc catgtcacc 7879

Fig. 9 (cont.)

COMMON LYMPHATIC ENDOTHELIAL AND VASCULAR ENDOTHELIAL RECEPTOR-1 (CLEVER-1) AND USES THEREOF**BACKGROUND OF THE INVENTION**

[0001] 1. Field of the Invention

[0002] The invention is in the field of cell adhesion proteins. Specifically, the invention is in the field of CLEVER-1, a novel protein that facilitates the influx of leukocytes and malignant cells into the lymphatic system, and also the efflux of the same out of the lymph nodes.

[0003] 2. Background Art

[0004] Leukocytes are the major cellular components of inflammatory and immune responses. Leukocytes include lymphocytes, natural killer (NK) cells, monocytes, dendritic cells and granulocytes (neutrophils, eosinophils and basophils). See, HARRISON'S PRINCIPLES OF INTERNAL MEDICINE, Fauci, A. S. et al. eds. (14th ed. 1998). Lymphocytes are composed of B cells and T cells. B cells provide humoral immunity and are the precursors of plasma cells. T cells provide cell mediated immunity. In tissues monocytes differentiate further into macrophages. At sites of inflammation, blood monocytes can attach to inflamed endothelia. Macrophages recognize and take up a wide range of exogenous materials such as bacteria. Granulocytes also have critical roles in inflammation. They are needed to clear infections with extracellular bacteria. The immune response has an important role in the growth, differentiation, and mobilization of granulocytes.

[0005] Continuous lymphocyte recirculation between blood and lymphoid tissues forms a basis for the function of the immune system. However such lymphocyte recirculation inadvertently also facilitates at least two medical conditions: inflammation and metastasis.

[0006] Lymphocytes enter the lymphoid tissues by binding to vascular endothelial cells. Lymphocyte adherence to endothelial cells is mediated by complementary, surface expressed molecules on both cell types. The adhesion molecules and mechanisms of lymphocyte entrance into the tissues from the blood have been thoroughly characterized, but mechanisms controlling lymphocyte exit from the non-lymphoid and lymphoid tissues via lymphatics have remained unknown.

[0007] The majority of lymphocytes extravasate into the lymph nodes via specialized vessels called high endothelial venules, or HEV. The rest of the incoming lymphocytes enter the nodes via afferent lymphatics together with antigens and other types of hematopoietic cells such as dendritic cells, macrophages and granulocytes. However, only lymphocytes are able to leave the nodes via the efferent lymphatic system by first traversing the sinusoidal endothelium and then entering the efferent lymphatic vessel. To maintain the homeostasis in the lymph node the numbers of entering and exiting lymphocytes need to be well balanced. The molecular mechanisms involved in lymphocyte exit are unknown.

[0008] In addition to being of fundamental importance in normal lymphocyte recirculation, the lymphatics also regulate seeding of metastasizing cells in approximately 50% of cancers that use this type of vessel for spreading. Lymph nodes are often the first organ to develop metastases, especially in the case of carcinomas. The design of the lymphatic system makes it relatively easy for malignant tumor cells to enter (Sleeman, J. P., *Recent Results Cancer Res.* 157:55-81

(2000)), and thus compounds that prevent the entry and exit of malignant tumor cells from the lymphatics have tremendous therapeutic potential.

BRIEF SUMMARY OF THE INVENTION

[0009] Recognizing the need to control lymphocyte recirculation, the inventors initiated a study of the proteins of the efferent lymphatic vessels. These studies have culminated in the discovery of a novel protein, Common Lymphatic Endothelial and Vascular Endothelial Receptor-1 (CLEVER-1), a binding protein that mediates adhesion of lymphocytes (and malignant tumor cells) to endothelium in both the systemic vasculature and in the lymphatics. The inventors have discovered that by blocking the interaction of CLEVER-1 and its lymphocyte substrate, the artisan can, for the first time simultaneously, control lymphocyte recirculation and lymphocyte migration, and related conditions such as inflammation, at the site of lymphocyte influx into, and efflux from, the tissues. The inventors have also discovered that CLEVER-1 also mediates binding of other types of leukocytes such as monocytes and granulocytes to HEV-like vessels. Further, by blocking the interaction of CLEVER-1 and malignant tumor cells, the artisan can also, for the first time, control metastasis by preventing malignant cells that bind to CLEVER-1 from being taken up by the lymphatic vessels, and thus preventing spread of the malignancy into the lymph nodes.

[0010] Accordingly, in a first embodiment, the invention is directed to cellular and subcellular extracts that contain CLEVER-1.

[0011] In a further embodiment, the invention is directed to purified or isolated CLEVER-1.

[0012] The invention further provides a method for inhibiting CLEVER-1 mediated leukocyte (such as lymphocyte, monocyte, and granulocyte) adhesion in a subject in need of the same, such method comprising administering a CLEVER-1 binding agent, or soluble CLEVER-1, to such subject.

[0013] The invention further provides a method for preventing or reducing CLEVER-1 mediated inflammation in a subject in need of the same, such method comprising administering a CLEVER-1 binding agent, or soluble CLEVER-1, to such subject.

[0014] The invention further provides a method for inhibiting CLEVER-1 mediated metastasis of malignant cells in a subject in need of the same, especially metastasis into the lymph nodes, such method comprising blocking the interaction of CLEVER-1 and the metastasizing cell in such subject with a CLEVER-1 binding agent, or by administering soluble CLEVER-1, to such subject.

[0015] The invention further provides a method for extracting cells that bind to CLEVER-1 from a population of cells, such method comprising mixing a population of cells with a preparation that contains CLEVER-1 and purifying the CLEVER-1-cell complex away from the rest of the non-binding cells in the population. Such method can be used to enrich for a subpopulation of CLEVER-1 binding cells.

[0016] The invention further provides a method for identifying an agent that binds to CLEVER-1, such method comprising determining whether a substance will successfully or unsuccessfully compete with leukocytes, malignant cells or with CLEVER-1 antibodies for binding to CLEVER-1.

[0017] The invention further provides a method for identifying an agent that is capable of inhibiting CLEVER-1 mediated cell migration, such method comprising assaying

CLEVER-1 trafficking of lymphocytes into the afferent lymphatics or HEV, and/or out of the lymph nodes, in the presence of such agent, and identifying such agent on the basis of its ability to inhibit such trafficking. In another embodiment, CLEVER-1 trafficking of malignant cells is assayed.

[0018] The present invention also provides a method of stimulating CLEVER-1 binding, for example, in immunocompromised hosts to facilitate leukocyte (such as lymphocyte, monocyte and granulocyte) trafficking and the function of immune defense systems.

BRIEF DESCRIPTION OF THE DRAWINGS

[0019] FIGS. 1a-1i. Indirect immunoperoxidase staining showing that monoclonal antibodies 3-266 and 3-372 recognize endothelium both in afferent and efferent lymphatic systems and on HEV. FIGS. 1a-1c are from the skin, FIGS. 1d-1i are from a lymph node. FIGS. 1a, 1d and 1g show the staining with monoclonal antibody 3-266, FIGS. 1b, 1e and 1h show the staining with monoclonal antibody 3-372 and FIGS. 1c, 1f and 1i show the staining with a negative control antibody, 3G6. In FIGS. 1a, 1b and 1c, the arrows point to the epithelium and arrowheads to afferent lymphatics. In FIGS. 1d and 1e, the arrows point to the lymphatic vessels (lymphatic sinusoids that belong to the efferent lymphatic system) within the lymph node. In FIGS. 1g and 1h, the arrows point to HEV.

[0020] FIG. 2. Monoclonal antibodies 3-266 and 3-372 recognize an about 270-300 kDa molecule. Molecules in lymph node lysates were separated by SDS-PAGE, blotted to nitrocellulose sheets and probed with monoclonal antibody 3-266 and 3-372 or with a negative control antibody (3G6).

[0021] FIGS. 3A and 3B. CLEVER-1 is involved in lymphocyte binding to endothelial cells both in HEV and lymphatics. An adhesion assay was performed to measure lymphocyte binding to HEV (FIG. 3A) and to lymphatic endothelium (FIG. 3B). The sections were pre-incubated with monoclonal antibody 3-266 or 3-372, or negative control antibody anti-HLA ABC or 3G6 ("neg co") after which the sections were overlaid with normal lymphocytes. The results of three to four independent inhibition experiments are shown as mean percentage of maximal binding \pm SEM.

[0022] FIGS. 4A and 4B. CLEVER-1 is involved in binding of tumor cells to endothelial cells both in HEV and lymphatics. An adhesion assay was performed to measure binding of different tumor cell lines to HEV (FIG. 4A) and to lymphatic endothelium (FIG. 4B). The sections were pre-incubated with 3-372 or negative control antibody (anti-HLA ABC) after which the sections were overlaid with different tumor cells: Nu, NA, IBW4, KCA and CRL 1648. The results of three to four independent inhibition experiments are shown as mean percentage of maximal binding \pm SEM.

[0023] FIG. 5. CLEVER-1 is induced on HEV-like vessels at sites of inflammation in connection to infiltrations of inflammatory cells (FIGS. 5a-5c, synovium; FIGS. 5d-5f, skin). FIG. 5a. Fibrotic type of inflamed synovium without any marked infiltrations of inflammatory cells. Only the afferent lymphatics expressed CLEVER-1 (arrows). FIG. 5b. CLEVER-1 was upregulated on a HEV-like vessel (marked by a dashed line) within a heavy lymphocytic infiltration. FIG. 5c. Staining with a negative control antibody (3G6). FIG. 5d. In normal skin afferent lymphatics expressed CLEVER-1 (arrows), but in inflamed skin HEV-like vessels (dashed line) also expressed CLEVER-1 (FIG. 5e). Negative control staining (FIG. 5f). Arrowheads point to epidermis (FIGS. 5d-5f).

[0024] FIG. 6. CLEVER-1 mediates binding of monocytes and granulocytes to HEV-like vessels at sites of inflammation. Contribution of CLEVER-1 in binding of monocytes and granulocytes to inflamed synovial vessels and binding of granulocytes to tonsil was tested using Stamper-Woodruff type of binding assay. 3-372 and 3-266 (pooled) but not the class-matched control antibody (3G6) significantly inhibited binding of granulocytes and monocytes to HEV-like vessels in the organs tested. The results of four independent assays are shown as mean percentage of maximal binding (=100% in the presence of the control antibody) \pm SEM.

[0025] FIG. 7. Molecular characterization of CLEVER-1. (a) Antibodies 3-266 and 3-372 recognize a 270-300 kDa molecule in immunoblotting. 3G6 is a negative control antibody. (b) In gels run 48 hr for better resolution, at least three different isoforms of CLEVER-1 are seen, and enzymatic digestions with neuraminidase and O-glycans reveal the sialoglycoprotein nature of CLEVER-1. (c) Relative contribution of different isoforms of CLEVER-1 is different in tonsil, lymph nodes and synovium. (d) An alternative spliced form missing exon 27 is present in 1. lung, 2. brain, 3. placenta, 4. heart, 5. liver, 6. skeletal muscle, 7. kidney, 8. pancreas, 9. spleen, 10. thymus, 11. prostata, 12. testis, 13. ovary, 14. small testine, 15. colon, 16. lymph nodes. Water control negative (lane 17). The upper band represents the standard form and the lower one is the splice variant of CLEVER-1.

[0026] FIG. 8. CLEVER-1 is expressed on the surface of endothelium in vivo and inhibition of its function blocks lymphocyte trafficking. Intravenously given 3-372 antibody (a) but not a negative class-matched control antibody (b) localized on the surface of HEV in lymph nodes after a 5 min circulation. HEV is pointed out by arrows in a. (c) Anti-CLEVER antibody treatment significantly inhibits the increase of the size of the lymph nodes draining the footpads. (One lymph node of a 3-372 treated rabbit was not found). (d) Lymphatic sinusoids of 3-372 treated animals contained less lymphocytes than those of control treated rabbits (e).

[0027] FIG. 9. The nucleotide sequence (7879 nt) of CLEVER-1. Boxed in grey are the translation initiation codon, translation stop codon, the two RGDs, the potential polyadenylation signal and the four nucleotide differences compared to Genebank entry AJ 2752213 (stabilin-1), i.e., nucleotides 1131, 2767, 6629 and 6969. Underlined are the nucleotides corresponding to the alternatively spliced exons.

DETAILED DESCRIPTION OF THE INVENTION

[0028] The term "ameliorate" denotes a lessening of an effect. To ameliorate a condition or disease refers to a lessening of the symptoms of the condition or disease.

[0029] The term "modulate" means to control in a predictable fashion, either by increasing or by decreasing the targeted parameter, as indicated from the context.

[0030] The term "effective amount" refers to that amount of the indicated agent that is sufficient to achieve the desired effect.

[0031] The term "inflammatory condition" refers to a physiological or pathological condition that is accompanied by an inflammatory response in a subject, which includes, inter alia, an undesired accumulation of leukocytes at one or more sites in such subject. The inflammatory condition can be hyperacute, acute, subacute or chronic. The inflammatory condition can be localized at the site of the inflammatory lesion or diffuse throughout the subject.

[0032] The term “drug” denotes any pharmaceutical or physiological agent, composition, bioactive compound, or combinations thereof, useful in the diagnosis, cure, mitigation, treatment, or prevention of a disease, or for any other medical purpose. The term “drug” is intended to be interpreted broadly and is not limited in terms of chemical composition or biological activity.

[0033] The term “essentially free of contaminants” refers to a substance that is of, undesired or unnecessary substances that had been present during the *in vitro* or *in vivo* synthesis of the desired substance.

[0034] The term “treatment” or “treating” refers to the administration of an agent to a subject for purposes which can include prophylaxis, amelioration, prevention or cure of an undesired disorder. Such treatment need not necessarily completely ameliorate the disorder, for example, inflammation; it is sufficient that such treatment ameliorates the disorder to a degree that is beneficial to the subject to which it is administered. Further, such treatment can be used in conjunction with other traditional treatments, for example, alternative treatments for reducing the inflammatory condition, known to those of skill in the art and as desired by the practitioner.

[0035] By “systemic vasculature” is meant the vascular network of blood vessels throughout the body of an animal or human.

[0036] By “lymphatic system” is meant the specialized part of the circulatory system that consists of lymph, the lymphatics, and the lymph nodes. The lymph nodes are located along the paths of the lymph collecting vessels and in isolated nodules of lymphatic patches in the intestinal wall. Additionally, there are specialized lymphatic organs such as the tonsils, thymus and spleen. B lymphocytes begin their final stages of maturation within the germinal centers of the lymph nodes’ cortical nodules. Maturing lymphocytes are then pushed to the more densely packed outer layers as they mature, before being released into the efferent lymphatics. The lymph nodes that are located in the floor of the mouth are called the submental and submaxillary lymph nodes. The superficial cervical lymph nodes are located in the neck. The superficial cubital or supratroclear lymph nodes are located just above the bend in the elbow. The auxiliary lymph nodes are clustered deep within the underarm and upper chest region. Inguinal lymph nodes are located in the groin. By “lymphatics” is meant the vessels that return lymph to the blood. Lymph is the clear fluid that flow in the lymphatics. Lymph arises from plasma that filters into the interstitial spaces from blood flowing through the capillaries. Although most of this plasma is taken up and absorbed by cells or the blood, a small amount is not absorbed. The lymphatics act as drains to collect this excess fluid and return it to the venous blood just before it reaches the heart. The lymph nodes act as filters that collect the lymph from several different lymphatics and “percolate” the lymph through spaces termed sinuses before draining into a single efferent draining vessel.

[0037] By “afferent lymphatics” is meant the vessels through which antigens enter the lymph nodes. Lymphocytes can enter the lymph nodes via the afferent lymphatics or via the high endothelial venules (HEV).

[0038] By “high endothelial venules” (HEV) is meant a specialized cortical postcapillary venules whose endothelium is simple cuboidal to columnar instead of simple squamous. HEVs are located mainly in the paracortex of the lymph nodes. Lymphocytes cross the HEV, and thus “traffic” into the lymph nodes by diapedesis, that is, the lymphocytes stick to

the luminal surface of the HEV, and then squeeze into the space between two or more HEV cells.

[0039] By “efferent lymphatics” meant the vessels that drain the lymph nodules (nodes).

[0040] By “lymphocyte recirculation” is meant the continuous movement of lymphocytes throughout the circulatory and lymph system. Lymphocytes leave the lymph node and are first delivered via the lymph to venous system draining into the heart.

[0041] The lymphocytes then circulate throughout the body in the bloodstream. Most of the lymphocytes are redelivered to the spleen or to another lymph node. About 10% go to non-lymphoid organs. Lymphocytes that have never been activated cannot enter non-lymphoid organs.

[0042] Lymphocyte “trafficking” refers to lymphocyte cell movement to specific locations. Outside of the lymph nodes, the trafficking of circulating lymphocytes allows the lymphocyte to accumulate at sites of inflammation. Activated effector lymphocytes tend to home to areas of inflammation, resulting in a large influx of lymphocytes in areas of inflammation. At the inflamed site, lymphocytes attach to the endothelial cells that line the blood vessels. This attachment localizes the lymphocyte at the site of inflammation and allows for subsequent emigration of the cells into the surrounding tissues (extravasation).

The Identification and Purification of CLEVER-1

[0043] The basis of the invention is the discovery of a new molecule, a novel protein herein designated “Common Lymphatic Endothelial and Vascular Endothelial Receptor-1 in the systemic vasculature, and in the afferent and efferent lymphatics. It has been found that leukocytes such as lymphocytes, monocytes, and granulocytes, and malignant cells specifically bind to this protein. It has also been found that this protein acts as a receptor that facilitates entry of bound leukocytes and malignant cells through the walls of the systemic vasculature, into the lymph nodes and out of the lymph nodes.

[0044] To search for a protein that played a role in lymphocyte lymphatic efflux, the inventors first identified cell migration-associated lymphatic structures from isolated efferent lymphatic vessels of human lymph nodes. These structures were used to produce monoclonal antibodies. Hybridomas were screened on frozen sections of human lymph nodes using immunoperoxidase staining.

[0045] Two of the hybridomas produced antibodies (designated 3-266 and 3-372) that clearly stained lymphatic endothelium both in afferent and efferent lymphatic systems and vascular endothelium on HEV, while other structures remained unstained. This is consistent with the expected pattern for an antibody that recognizes a lymphocyte migration-associated structure. Cell culture of 3-266 (DSM ACC2519) and cell culture of 3-372 (DSM ACC2590) were both deposited under the terms of the Budapest Treaty on the International Recognition of the Deposit of Micro-organisms for the Purposes of Patent Procedure on Aug. 21, 2001, with DSMZ-Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH, Mascheroder Weg 1b, D-38124 Braunschweig.

[0046] In molecular weight determinations by immunoblotting, both antibodies recognized a molecule of the same size, about 270-300 kDa. Due to this and an identical staining pattern, these antibodies were considered to recognize the same antigen; that antigen was named Common Lymphatic Endothelial and Vascular Endothelial Receptor-1 (CLEVER-1).

[0047] CLEVER-1 was purified from CLEVER-1 containing lymph node preparations using affinity chromatography with the 3-372 antibody. The eluted material was subjected to SDS-PAGE analysis and silver staining. The specific band was excised, reduced, alkylated and digested with trypsin.

[0048] After cleavage with trypsin, mass spectrometric analyses yielded 27 peptides. Twenty-one (77%) of those had identical sequences with stabilin. These sequences covered altogether 268 amino acids (10% of the 2570 amino acids of stabilin-1) and spanned the amino acids between 53 and 2301 (Table 1). The peptide data suggest that CLEVER-1 has some homology with stabilin-1 at the structural level. No functional information regarding stabilin-1 can be found in the literature.

[0049] Peptide analysis of CLEVER-1 indicates no significant homology with any of the known endothelial homing-associated molecules, such as ICAM-1 (Intercellular Adhesion Molecule), ICAM-2, or VAP-1 (Vascular Adhesion Protein).

[0050] CLEVER-1 has several structural motifs that are associated with adhesive functions in other molecules. They include a proteoglycan link homology region important in CD44 for hyaluronan binding and two RGD motifs known to serve as integrin ligands in certain molecules, such as in fibronectin. In addition, CLEVER-1 has seven fasciclin domains also present in several molecules such as priostin, fasciclin and transforming growth factor- β -induced gene, big-h3 and in all of these cases it is essential for adhesive function of these molecules. Interestingly, twenty-two epidermal growth factor (EGF) repeats are also found in CLEVER-1. This structural domain is also present in all members of the selectin family. Although the lectin domains of selectins are of utmost importance for the transient interaction with their sialomucin ligands, EGF-repeats have been reported to functionally contribute with lectin domains to binding between leukocytes and endothelium. Based on the structural complexity of CLEVER-1 it may turn out have several ligand molecules and be multifunctional in its nature.

[0051] It has been discovered that CLEVER-1 is involved in the process of lymphocyte recirculation. CLEVER-1 is present on the endothelium of the systemic vasculature, especially on the HEV and also on the endothelium of both afferent and efferent lymphatic systems. CLEVER-1 is a protein adhesion molecule, and especially, a cell adhesion molecule (CAM), that mediates adhesion of lymphocytes and of malignant tumor cells to CLEVER-1 in the systemic vasculature and the lymphatic system. These sites are of utmost importance as control points in lymphocyte trafficking.

[0052] CLEVER-1 is the first molecule that has been identified to facilitate lymphocyte and malignant cell exit from the lymph nodes. Additionally, CLEVER-1 is the first molecule that has been identified that regulates both entrance of lymphocytes and tumor cells into the lymph nodes and exit of lymphocytes and tumor cells from the lymph nodes. CLEVER-1 has been found to also mediate binding of other leukocytes such as monocytes and granulocytes to HEV-like vessels.

[0053] By "CLEVER-1 mediated cell binding" is meant the specific association of CLEVER-1 with either a leukocyte, such as a lymphocyte, monocyte, or granulocyte, or a CLEVER-1-binding malignant cell. CLEVER-1 mediated cell binding can occur with CLEVER-1 in a soluble form or in

a particulate form (for example, when CLEVER-1 is present in a form that is membrane associated).

CLEVER-1 Mediated Binding of Leukocytes

[0054] According to the invention, adhesion of leukocytes, such as lymphocytes, monocytes, and granulocytes, to the endothelium (that is, to an endothelial cell(s)) in the systemic vasculature, especially the HEV, and to the endothelium in the afferent and efferent lymphatics can be blocked by blocking the binding between such endothelial cell's CLEVER-1 and leukocyte.

[0055] In the systemic vasculature, inhibiting or preventing endothelial cell CLEVER-1 mediated lymphocyte binding will inhibit or prevent lymphocytes, especially activated lymphocytes, from accumulating at such sites, and thus prevent or lessen inflammation at such sites. Thus, the invention provides a method of treating inflammation, by administering an agent that inhibits or prevents CLEVER-1 mediated endothelial cell binding to lymphocytes.

[0056] In the afferent lymphatics, inhibiting or preventing lymphocytes from binding to endothelial cell CLEVER-1 will inhibit or prevent such lymphocytes from entering the afferent lymphatics and thus the lymph nodes. Thus, the invention provides a method of treating inflammation, by administering an agent that inhibits or prevents CLEVER-1 mediated endothelial cell binding to lymphocytes in the afferent lymphatics and especially at HEV in lymph nodes or HEV-like venules at sites of inflammation. The invention also provides a method of inhibiting lymphocyte trafficking into the lymph nodes, by administering an agent that inhibits or prevents afferent lymphatic CLEVER-1 mediated endothelial cell binding, and especially HEV binding, to lymphocytes and other leukocytes.

[0057] In the efferent lymphatics, inhibiting or preventing lymphocytes from binding to endothelial cell CLEVER-1 will prevent the lymphocytes from exiting the lymph node and entering the blood. Thus, the invention provides a method of treating inflammation, by administering an agent that inhibits or prevents CLEVER-1 mediated endothelial cell binding to lymphocytes in the efferent lymphatics. The invention also provides a method of inhibiting lymphocyte trafficking out of the lymph nodes, by administering an agent that inhibits or prevents efferent lymphatic CLEVER-1 mediated endothelial cell binding to lymphocytes.

[0058] Therefore, CLEVER-1 binding with lymphocytes presents a unique, three-prong approach to treat diseases or conditions characterized by an undesired lymphocyte accumulation or trafficking in which the artisan can target lymphocyte entry into the lymph nodes, lymphocyte exit from the lymph nodes, and lymphocyte binding to the systemic vasculature, with the same agent.

[0059] The discovery of CLEVER-1 and its role has thus resulted in a new method to control lymphocyte migration by inhibiting CLEVER-1 mediated cell binding to such cells. Thus, the present invention provides a method of inhibiting undesired CLEVER-1 mediated lymphocyte trafficking, and thus blocking harmful or otherwise undesired lymphocyte migration, by preventing the association of CLEVER-1 with lymphocytes. Similarly, the invention provides a method of inhibiting undesired CLEVER-1 mediated binding of other leukocytes by preventing association of CLEVER-1 with the leukocytes.

[0060] The present invention also provides a method of stimulating CLEVER-1 binding, for example, in immuno-

compromised hosts to facilitate lymphocyte trafficking and other leukocyte binding and the function of immune defense systems.

CLEVER-1 Mediated Cell Binding to Malignant Cells

[0061] Because cancer cells often break away from a malignant tumor and enter the lymphatics, cancer cells travel to and establish themselves in the lymph nodes. According to the invention, the ability of a malignant tumor cell to establish itself in a lymph node can be inhibited or prevented by inhibiting or preventing CLEVER-1 binding to such malignant tumor cell.

[0062] The term “tumor” refers to a neoplasm, a tissue mass that is characteristic of a neoplasia. Neoplasia is distinguished from other forms of tissue growth, first, by the formation of a tissue mass, a neoplasm, or tumor. Second, neoplasia is considered to be an irreversible process. Third, neoplastic tissue tends to morphologically resemble its tissue of origin. Fourth, neoplastic tissue tends to functionally resemble its tissue of origin. Fifth, neoplasms grow and function somewhat independently of the homeostatic mechanisms that control normal tissue growth and function.

[0063] A neoplasm can be benign or malignant. A benign neoplasm consists of a discrete tissue mass that continues to grow. A benign neoplasm will simply push adjacent tissues out of its way as it grows.

[0064] The definitive features of a malignant neoplasm, a malignancy, are invasion and metastasis, that is, the spread of the neoplasm to a distant site. A malignant neoplasm will grow into the adjacent tissue, rather than pushing it away. The terms “malignant neoplasm,” “malignant tumor,” and “cancer” are synonymous.

[0065] Cancer cells typically invade thin-walled vessels such as small veins, venules, capillaries and lymphatics. The passage of cancer cells via lymphatics to lymph nodes, and via blood vessels to other organs and structures, and the subsequent implantation and growth of the cancer cells in those sites is referred to as “metastasis.” The lymph nodes are common sites for metastasis.

[0066] Cancer cells can also spread by seeding—shedding into, for example, the peritoneal fluid. The cells can be carried by the fluid to a distant site on the peritoneal surface where they can implant and form new foci of cancer growth.

[0067] Most neoplasms are one of four types: epithelial, non-epithelial, blastomas and teratomas. Malignant epithelial neoplasms are termed carcinomas. An adenocarcinoma is a carcinoma in which gland-like structures are present. Carcinomas can be papillary or cystic. Benign epithelial neoplasms are generally adenomas, polyps or papillomas.

[0068] Non-epithelial tumors can also be benign or malignant. They are generally named by a prefix that indicates the histologic type and a suffix. The suffix -oma generally means benign while the suffix -sarcoma means malignant. However, several malignant neoplasms have traditional names ending in -oma: for example, melanoma, hepatoma, and lymphoma.

[0069] Lymphomas are malignant neoplasms arising from cells of the lymphoid series. Blastomas and teratomas contain more than one type of tissue. Malignant teratomas are often termed teratocarcinomas.

[0070] A “leukemia” is a tumor of white blood cells that is present in the bone marrow and blood. A “lymphoma” is a tumor of white blood cells that is present in the lymph nodes and tissues.

[0071] According to the invention, the binding of CLEVER-1-binding malignant cells to the endothelium in the systemic vasculature, especially the HEV, and to the endothelium in the afferent and efferent lymphatics can be inhibited or prevented by inhibiting or preventing the binding between such endothelial cell’s CLEVER-1 and such malignant cell. Thus, the invention provides a method of treating cancer, and especially, a method of preventing metastasis, by administration of an agent that inhibits or prevents CLEVER-1 mediated malignant cell binding to the endothelium.

[0072] In the systemic vasculature, inhibiting or preventing CLEVER-1 mediated cell binding will inhibit or prevent the establishment of CLEVER-1 binding malignant cells at such sites, and thus lessen or prevent metastasis of such malignant cells. Thus, the invention provides a method of treating cancer, and especially, a method for preventing metastasis of a CLEVER-1 binding malignant cell, by administering an agent that inhibits or prevents CLEVER-1 mediated endothelial cell binding to CLEVER-1-binding tumor cells in the systemic vasculature. The invention also provides a method of inhibiting metastasis, by administering an agent that inhibits or prevents systemic vasculature CLEVER-1 mediated endothelial cell binding to such malignant cells.

[0073] In the afferent lymphatics, inhibiting or preventing CLEVER-1 binding malignant cells from binding to endothelial cell CLEVER-1 will inhibit or prevent such CLEVER-1 binding malignant cell from entering and establishing in the lymph node, and thus lessen or prevent metastasis of such cell to the lymph node or thus to other sites in the body. In this context it is worth to note that a metastasizing malignant cell cannot survive long times without matrix support—a condition present for example in blood. Thus, the invention provides a method of treating cancer, and especially, a method for preventing metastasis of a CLEVER-1 binding malignant cell, by administering an agent that inhibits or prevents CLEVER-1 mediated endothelial cell binding to CLEVER-1-binding malignant cells in the afferent lymphatics and at HEV in systemic vasculature. The invention also provides a method of inhibiting metastasis, by administering an agent that inhibits or prevents afferent lymphatic CLEVER-1 mediated endothelial cell binding, and especially HEV binding, to such malignant cells.

[0074] In the efferent lymphatics, inhibiting or preventing CLEVER-1 binding malignant cells from binding to endothelial cell CLEVER-1 will inhibit or prevent such CLEVER-1 binding malignant cell from leaving the lymph node, and thus lessen or prevent metastasis of such cell from the lymph node to other sites in the body. Thus, the invention provides a method of treating cancer, and especially, a method for preventing metastasis of a CLEVER-1 binding malignant cell, by administering an agent that inhibits or prevents CLEVER-1 mediated endothelial cell binding to CLEVER-1-binding malignant cells in the efferent lymphatics. The invention also provides a method of inhibiting metastasis, by administering an agent that inhibits or prevents efferent lymphatic CLEVER-1 mediated endothelial cell binding to such malignant cells.

[0075] CLEVER-1 interaction with CLEVER-1 binding malignant cells thus presents a unique, three-prong approach to inhibit or prevent metastasis in which not only can the artisan block such malignant cells from entering into and

exiting from the lymph system, but also, the artisan can block association of such malignant cell with CLEVER-1 in the vascular endothelium.

Agents that Block or Inhibit CLEVER-1 Mediated Cell Binding

[0076] Soluble CLEVER-1 and antibodies to CLEVER-1 can be provided to the host cell to block or inhibit CLEVER-1 mediated cell binding. Soluble CLEVER-1 can be used to “coat” the CLEVER-1 binding sites on the leukocyte, such as lymphocyte, monocyte, or granulocyte, or tumor cell and thus prevent the coated cell from association with the native CLEVER-1 on the HEV or afferent or efferent lymphatics.

[0077] CLEVER-1 antibodies can be administered to a patient in need of the same to coat CLEVER-1 that is present on the vascular endothelium or lymphatics, of such patient, especially the afferent lymphatics so as to prevent leukocyte or malignant cell binding to such CLEVER-1 in the patient. CLEVER-1 antibody producing cells can be administered directly to the patient so as to provide a source of the same.

[0078] Moreover, the present invention provides a method of identifying an agent that inhibits the binding of CLEVER-1 to cells by providing an agent to cells in the presence of CLEVER-1 and comparing the binding of CLEVER-1 to cells provided with the agent to binding of CLEVER-1 in the absence of the agent. Similarly, the invention provides a method of identifying an agent that stimulates the binding of CLEVER-1 to cells by providing an agent to cells in the presence of CLEVER-1 and comparing the binding of CLEVER-1 to cells provided with the agent to binding of CLEVER-1 in the absence of said agent.

[0079] The term “antibody” is used in the broadest sense and specifically covers single monoclonal antibodies (including agonist and antagonist antibodies), polyclonal antibodies, as well as antibody fragments and single chain antibodies (e.g., Fab, F(ab')₂, Fv), so long as they exhibit the desired biological activity.

[0080] Pepsin digestion of antibodies produces two identical antigen binding fragments, called Fab fragments, each with a single antigen binding site, and a residual “Fc” fragment, whose name reflects its ability to crystallize readily. Pepsin treatment yields an F(ab')₂ fragment that has two antigen combining sites and is still capable of cross-linking antigen.

[0081] Single chain “Fv” is the minimum antibody fragment which contains a complete antigen recognition and binding site. This region consists of a dimer of one heavy and one light chain variable domain in tight, non-covalent association. It is in this configuration that the three CDRs of each variable domain interact to define an antigen binding site on the surface of the V_H-V_L dimer. Collectively, the six CDRs confer antigen binding specificity to the antibody. However, even a single variable domain (or half of an Fv comprising only three CDRs specific for an antigen) has the ability to recognize and bind antigen, although at a lower affinity than the entire binding site. See, Ladner et al., U.S. Pat. No. 4,946,778, and Bird, R. E. et al., *Science*, 242:423-426 (1988).

[0082] The term “monoclonal antibody” as used herein refers to an antibody obtained from a population of substantially homogeneous antibodies, i.e., the individual antibodies comprising the population are identical except for possible naturally occurring mutations that may be present in minor amounts. Monoclonal antibodies are highly specific, being directed against a single antigenic site. Furthermore, in contrast to conventional (polyclonal) antibody preparations

which typically include different antibodies directed against different determinants (epitopes), each monoclonal antibody is directed against a single determinant on the antigen. In addition to their specificity, the monoclonal antibodies are advantageous in that they are synthesized by a hybridoma culture, uncontaminated by other immunoglobulins. The modifier “monoclonal” indicates the character of the antibody as being obtained from a substantially homogeneous population of antibodies, and is not to be construed as requiring production of the antibody by any particular method. For example, the monoclonal antibodies to be used in accordance with the present invention may be made by the hybridoma method first described by Kohler and Milstein, *Nature* 256:495 (1975), or may be made by recombinant DNA methods (e.g., Cabilly et al., U.S. Pat. No. 4,816,567).

[0083] Preparation of Immunizing Antigen, and Polyclonal and Monoclonal antibody production can be performed as described herein, or using other suitable techniques. A variety of methods have been described (see e.g., Kohler et al., *Nature* 256:495-497 (1975), and *Eur. J. Immunol.* 6:511-519 (1976); Milstein et al., *Nature* 266:550-552 (1977); Koprowski et al., U.S. Pat. No. 4,172,124; Harlow, E. and D. Lane, *Antibodies: A Laboratory Manual* (Cold Spring Harbor Laboratory: Cold Spring Harbor, N.Y., 1988); CURRENT PROTOCOLS IN MOLECULAR BIOLOGY, Vol. 2 (Supplement 27, 1994), Ausubel, F. M. et al., John Wiley & Sons, eds., New York, N.Y.), Chapter 11, (1991)). Generally, a hybridoma can be produced by fusing a suitable immortal cell line (e.g., a myeloma cell line such as SP2/0) with antibody producing cells. The antibody producing cell, preferably those of the spleen or lymph nodes, are obtained from animals immunized with the antigen of interest. The fused cells (hybridomas) can be isolated using selective culture conditions, and cloned by limiting dilution. Cells which produce antibodies with the desired binding properties can be selected by a suitable assay (e.g., ELISA).

[0084] The term “antibody” also includes chimeric, humanized or primatized (CDR-grafted) antibodies, as well as chimeric or CDR-grafted single chain antibodies, and the like, comprising portions derived from different species. “Chimeric” antibodies (immunoglobulins) have a portion of the heavy and/or light chain is identical with or homologous to corresponding sequences in antibodies derived from a particular species or belonging to a particular antibody class or subclass, while the remainder of the chain(s) is identical with or homologous to corresponding sequences in antibodies derived from another species or belonging to another antibody class or subclass, as well as fragments of such antibodies, so long as they exhibit the desired biological activity (Cabilly et al., U.S. Pat. No. 4,816,567; Morrison et al., *Proc. Natl. Acad. Sci. USA* 81:6851-6855 (1984)). The various portions of these antibodies can be joined together chemically by conventional techniques, or can be prepared as a contiguous protein using genetic engineering techniques. For example, nucleic acids encoding a chimeric or humanized chain can be expressed to produce a contiguous protein. See, e.g., Cabilly et al., U.S. Pat. No. 4,816,567; Cabilly et al., EP 0 125 023 B1; Boss et al., U.S. Pat. No. 4,816,397; Boss et al., EP 0 120 694 B1; Neuberger, M. S. et al., WO 86/01533; Neuberger, M. S. et al., EP 0 194 276 B1; Winter, U.S. Pat. No. 5,225,539; Winter, EP 0 239 400 B1; and Queen et al., U.S. Pat. Nos. 5,585,089, 5,698,761 and 5,698,762. See also, Newman, R. et al., *BioTechnology* 10: 1455-1460 (1992), regarding primatized antibody.

[0085] By “agonist antibody” is meant an antibody which is able to bind to CLEVER-1 and facilitate adhesion of lymphocytes (and malignant tumor cells) to endothelium. By “antagonist antibody” is meant an antibody that is able to bind to CLEVER-1 and inhibit adhesion of lymphocytes (and malignant tumor cells) to endothelium.

[0086] Anti-idiotypic antibodies are also provided. Anti-idiotypic antibodies recognize antigenic determinants associated with the antigen-binding site of another antibody. Anti-idiotypic antibodies can be prepared against second antibody by immunizing an animal of the same species, and preferably of the same strain, as the animal used to produce the second antibody. See e.g., U.S. Pat. No. 4,699,880.

In Vitro Adhesion Assay and Diagnostic Uses Thereof

[0087] In a further embodiment, the present invention is directed to an adhesion assay in which CLEVER-1 binding is used to assay for the presence of leukocytes or malignant cells that bind to HEV and lymphatic endothelium. Both static and non-static assays are possible. The adhesion assay is exemplified in Example 4. Both static and non-static assays can be used to study leukocyte binding to systemic vasculature.

[0088] In the static assay, a tissue section is exposed to leukocytes or malignant cells for a desired period of time, without continuous agitation or rotation of the preparation during the exposure. Static assays are preferred for examining the ability of leukocytes and malignant cells to bind to lymphatic endothelium, especially the efferent lymph vessels.

[0089] In the non-static assay, the CLEVER-1 containing tissue sample and leukocytes are constantly rotated during the time period in which the leukocytes are given to adhere to the CLEVER-1. Non-static assays are preferred for studying leukocyte and malignant cell binding to CLEVER-1 in the HEV. The non-static assay mimics adhesion to the systemic vasculature.

[0090] In another embodiment, the present invention relates to a method for detection of malignant tumor cells. As explained with detail in Example 1, CLEVER-1 antibodies reduce the binding of malignant tumor cells to the vascular and lymphoid endothelium, demonstrating CLEVER-1 is a receptor for such malignant tumor cells. CLEVER-1 protein, or fragments thereof, or CLEVER-1 binding compounds, including but not restricted to, antibodies against CLEVER-1, both monoclonal and polyclonal, antibodies against antigenic fragments of CLEVER-1, both monoclonal and polyclonal, antigenic polypeptides, small molecule inhibitors or drugs, can be used in both quantitative and qualitative assays to detect the presence of malignant tumor cells in a sample, said sample being tissue or blood from a human or animal.

[0091] CLEVER-1 protein, or fragments thereof, or the above-mentioned CLEVER-1 binding compounds can be attached to a solid support matrix, including but not limited to microtiter plates, agarose columns, or magnetic beads. The above-mentioned sample can then be exposed to said solid support matrix, and the percentage of cells retained by said solid support matrix determined. For example, a sample from a normal, healthy individual, said individual being either a human or an animal, would have a statistically predicted number of leukocytes that bind to CLEVER-1. A sample that contains both leukocytes and malignant tumor cells would have a detectably higher number of cells that bind CLEVER-1.

[0092] In a preferred embodiment, a blood or tissue sample that has been taken from a patient who is in need of treatment,

especially treatment for cancer, is used as the source of the CLEVER-1 binding cells in the in vitro adhesion assay. Such patient can be a patient being treated for a previously diagnosed malignancy, or a patient suspected of having a malignant tumor, or a patient who appears to be cured of such malignant tumor but is in need of monitoring for the reoccurrence of the same. Preferably, such blood or tissue sample is from a patient who is to be tested for the presence of malignant cells that bind to CLEVER-1 in such sample.

[0093] The blood or tissue sample that is to be examined in the in vitro assay of the invention can be processed, if desired, by methods known in the art so as to further extract or concentrate any CLEVER-1 binding cells that might be present in the sample, prior to the sample's being used in the in vitro adhesion assay of the invention.

[0094] Additionally, once the in vitro adhesion assay is complete, the adherent cells can be studied using other methods. For example, in the static assay, where the bound cells have been fixed, ability of the adherent cells to be recognized by a monoclonal antibody that is diagnostic for the type of tumor can be performed.

[0095] Detecting the binding of malignant cells to the CLEVER-1 containing lymphatic endothelium indicates that the patient is in need of treatment for such malignant cells, and especially to prevent the metastasis of such malignant cells

[0096] The present invention provides in this aspect a novel, efficient, and convenient assay for identifying antagonists, including but not limited to, monoclonal and polyclonal antibodies, peptides, protein fragments, small molecular inhibitors, drugs, and other agents, which can inhibit the adhesion of leukocytes and malignant tumor cells to the vascular and lymphatic endothelium.

[0097] For example, CLEVER-1 containing samples of lymph node sections can be incubated with and without the agent, and the number of bound lymphocytes and/or malignant cells determined. The antagonists can be pre-incubated with lymph node sections (a non-competitive assay) or simultaneously added with lymphocytes to the lymph node sections (a competitive assay).

[0098] Such a screen can also be used to customize an anti-metastasis treatment to an individual patient, and allows the practitioner to identify and select those agents or combinations thereof that have the best ability to inhibit CLEVER-1 malignant cell binding to vascular and/or lymphatic endothelium in such patient, and thus maximize the benefit of the treatment with such agents for such patient.

[0099] Additionally, such in vitro assay allows the practitioner to select for agents that provide a beneficial effect on disrupting malignant cell: CLEVER-1 containing endothelium interactions, nevertheless, minimize, if possible, the effect of such treatment on CLEVER-1 mediated leukocyte binding.

[0100] An antagonist can inhibit malignant cell or lymphocyte cell migration into or out of the lymph nodes. In a preferred embodiment, antagonists would inhibit both entrance and exit of an undesired cell into and out of the lymph nodes, respectively. As such, malignant tumor cells would preferably be prevented from entering a lymph node, and establishing there, and any that did enter the lymph nodes via an afferent lymph vessel independent mechanisms would be contained, thus slowing metastasis.

[0101] This assay can be used further to monitor the efficacy of chemotherapy treatments administered to an indi-

vidual, said individual being a human or an animal, in need thereof. Samples can be analyzed before, during, and after chemotherapy for the presence of malignant tumor cells that bind to CLEVER-1 or antigenic fragments thereof, or CLEVER-1 binding compounds.

[0102] In a further embodiment, purified CLEVER-1 protein, or fragments thereof can be used for high volume screening of antagonists that are capable of preventing or lowering the ability of a leukocyte or malignant cell to adhere to endothelial cell CLEVER-1. CLEVER-1 protein, or fragments thereof can be attached to a solid support matrix, including but not limited to a microtiter plate, an agarose column, or magnetic beads, using standard methods well known in the art. Antagonists can be screened for interaction with CLEVER-1 or fragments thereof, either in the absence or presence of leukocytes. Leukocytes or malignant cells can be labeled with fluorescent dyes such as, for example, bis-carboxyethyl carboxyfluorescein or fluorescein isothiocyanate and the number of bound cells in presence or absence of the antagonists can be analyzed by a fluorimager.

[0103] The high volume screen assay of this aspect of the invention can be used to screen combinatorial libraries for molecules that inhibit the binding of leukocytes and malignant tumor cells to CLEVER-1 or a fragments thereof. Antagonists that show strong affinity for purified CLEVER-1 protein or fragments thereof can be screened further using the in vitro adhesion assay described above.

[0104] Antibodies used in the methods of the invention as CLEVER-1 binding compounds are preferably antibodies with a specificity against CLEVER-1, or an antigenic fragment thereof. Such antibodies can be polyclonal or monoclonal.

[0105] Another potential CLEVER-1 antagonist is a peptide derivative of the CLEVER-1 polypeptide that are naturally or synthetically modified analogs of the polypeptides that have lost biological function yet still recognize and bind to the ligand of the polypeptides to thereby effectively block the interaction of said ligand with CLEVER-1. Examples of peptide derivatives include, but are not limited to, small peptides or peptide-like molecules.

[0106] Another potential human CLEVER-1 antagonist is a peptide derivative of the ligand polypeptides which are naturally or synthetically modified analogs of the polypeptides that have lost biological function yet still recognize and bind to CLEVER-1 to thereby effectively block CLEVER-1. Examples of peptide derivatives include, but are not limited to, small peptides or peptide-like molecules.

[0107] The present invention relates to a diagnostic method for the detection of cells that contain CLEVER-1, that is, CLEVER-1 positive cells, in samples taken from the human or animal body. Such a method would involve the use of CLEVER-1 binding compounds, including but not limited to, monoclonal and polyclonal antibodies, with specificity for CLEVER-1. Such compounds can be labeled with a substance, such as a colorimetric dye or radioactive molecule, to permit rapid and easy detection of binding of the compound to cells that express CLEVER-1.

Therapeutic Uses of CLEVER-1 Antagonists

[0108] In another embodiment, the present invention relates to a method of treating malignant carcinomas. It is common for carcinomas to metastasize first to the regional lymph nodes (Sleeman, J. P., *Recent Results Cancer Res.* 157:55-81 (2000)). As described herein, CLEVER-1 is

involved in the entrance and exit of malignant tumor cells to and from the lymph nodes. As such, antagonists that inhibit malignant tumor cell binding to CLEVER-1, including but not limited to, monoclonal and polyclonal antibodies, peptides, small molecule inhibitors, drugs, and other such agents can reduce metastasis and serve as effective chemotherapeutic agents.

[0109] In another aspect, the present invention relates to a method of treating disorders where the leukocyte-endothelial cell adhesion reaction is associated with acute or chronic inflammatory diseases such as skin inflammations, diabetes, connective tissue diseases (such as lupus, rheumatoid arthritis, osteoarthritis), obstructive and restrictive lung diseases (such as asthma, ARDS, sarcoidosis, idiopathic pulmonary fibrosis), inflammatory bowel diseases (such as ulcerative colitis and Crohn's disease), various nephritides, non-viral hepatitis, cirrhosis, cholangitis, atherosclerosis, vasculitis, thyroiditis, multiple sclerosis, myositis, ischemia reperfusion injury, transplantation rejection.

[0110] The antagonists can also be employed to treat histamine-mediated allergic reactions and immunological disorders including late phase allergic reactions, chronic urticaria, and atopic dermatitis. IgE-mediated allergic reactions such as allergic asthma, rhinitis, and eczema can also be treated.

[0111] The antagonists can also be employed to treat chronic and acute inflammation by preventing the extravasation of leukocytes to a wound area. They can also be employed to regulate normal pulmonary macrophage populations, since chronic and acute inflammatory pulmonary diseases are associated with sequestration of mononuclear phagocytes in the lung.

[0112] Antagonists can also be employed to treat rheumatoid arthritis by preventing the extravasation of leukocytes into synovial fluid in the joints of patients. Monocyte influx and activation plays a significant role in the pathogenesis of both degenerative and inflammatory arthropathies.

[0113] The antagonists can also be employed to treat asthma and allergy by preventing eosinophil accumulation in the lung. The antagonists can also be employed to treat sub-epithelial basement membrane fibrosis which is a prominent feature of the asthmatic lung.

[0114] The antagonists can also be employed for treating atherosclerosis, by preventing monocyte infiltration in the artery wall.

[0115] The antagonists can be employed in a composition with a pharmaceutically acceptable carrier, e.g., as hereinafter described.

Formulations of Compounds

[0116] The antagonists of CLEVER-1 can be used as therapeutic compositions. The antagonists of CLEVER-1 can be administered as a single dose or in multiple doses. The antagonists of the present invention can be administered either as an independent therapeutic regime or in combination with other therapeutic agents. The antagonists can be combined with conventional therapies, which can be administered simultaneously or sequentially.

[0117] Such therapeutic compositions can consist solely of the antagonist of CLEVER-1 although, preferably, the compositions will contain the antagonist of CLEVER-1 combined in admixture with a pharmaceutically acceptable carrier vehicle. Suitable vehicles and their formulation, inclusive of other human proteins, e.g., human serum albumin, are described for example in Remington: The Science and Prac-

tice of Pharmacy, Gennaro, Alfonso, 20th ed. (2000). In order to form a pharmaceutically acceptable composition that is suitable for effective administration to a patient in need of such composition, such compositions will contain an effective amount of the antagonist of CLEVER-1 together with a suitable amount of carrier vehicle.

[0118] Compositions containing antagonists of CLEVER-1 can be administered perorally, intravenously, intramuscularly, or sub-cutaneously at the appropriate dosages, which will depend upon the severity of the condition of the patient and upon such criteria as the patient's height, weight, sex, age, and medical history. The dose will also depend upon whether the compound of the invention is being administered to a human patient or in a veterinary setting to an animal, in need thereof.

[0119] For the purpose of parenteral administration, compositions containing the antagonists of CLEVER-1 are preferably dissolved in distilled water and the pH-value is preferably adjusted to about 6 to 8. In order to facilitate the lyophilization process resulting in a suitable product, lactose can be added to the solution. Preferably, the solution is then filtered sterilized, introduced into vials, and lyophilized. In a preferred embodiment, the compound of the invention is administered orally to a patient, at the time of eating or shortly thereafter. The concentration of the antagonists of CLEVER-1 in these compositions, whether oral or parenteral, can vary, e.g., from 10^{-12} M to 10^{-3} M.

[0120] Additional pharmaceutical methods can be employed to control the duration of action. Controlled release preparations can be achieved by the use of polymers to complex or adsorb the antagonists of CLEVER-1. The controlled delivery can be exercised by selecting appropriate macromolecules (for example, polyesters, polyamino acids, polyvinyl pyrrolidone, ethylenevinylacetate, methylcellulose, carboxymethylcellulose, and protamine sulfate) and the concentration of macromolecules as well as the methods of incorporation in order to control release. Another possible method to control the duration of action by controlled release preparations is to incorporate the antagonists of CLEVER-1 into particles of a polymeric material such as polyesters, polyamino acids, hydrogels, poly (lactic acid) or ethylene vinylacetate copolymers. Alternatively, instead of incorporating the CLEVER-1 antagonists into these polymeric particles, it is possible to entrap these derivatives in microcapsules prepared, for example, by coacervation techniques or by interfacial polymerization, for example, hydroxymethylcellulose or gelatin-microcapsules and poly (methylmethacrylate) microcapsules, respectively, or in colloidal drug delivery systems, for example, liposomes, albumin microspheres, microemulsions, nanoparticles, and nanocapsules or in macroemulsions. Such teachings are disclosed in Remington: The Science and Practice of Pharmacy, Gennaro, Alfonso, 20th ed. (2000).

[0121] The following Example serves only to illustrate the invention, and is not to be construed as in any way limiting the invention.

EXAMPLE 1

Production of Monoclonal Antibodies

[0122] Balb/c mice were immunized to footpads four times at one week intervals, with incomplete Freund's adjuvant containing suspension made from lymphatic vessels excised from human lymph nodes under stereo microscope. The sus-

pension was made by cuffing the vessels to small pieces by scissors and the pieces in phosphate buffered saline were then drawn back and forth into a syringe connected to a 21 g needle. The popliteal lymph node lymphocytes from the immunized mice were isolated by a glass homogenizer. The popliteal lymph node lymphocytes of the immunized mice were fused with Sp2/0 myeloma cells. Hybridoma supernatants were primarily tested on frozen sections of human lymph nodes using immunoperoxidase staining. The testing conditions were the same for antibodies 3-266 and 3-372 generated by two of the hybridomas.

[0123] Immunoperoxidase stainings were performed as described (Salmi, *Science* 257:1407-1409 (1992)). Briefly, acetone fixed 6 μ m frozen sections from different human tissues (lymph nodes, appendix, bronchus, cerebellum, epididymis, esophagus, heart, small and large intestine, kidney, liver, lung, normal and psoriatic skin, synovium, testis and tonsil) were stained with antibody 3-266, 3-372 or 3G6, a negative class matched control antibody for 3-266 and 3-372 (mouse IgG1) and 3,3-diaminobenzidine was used as a substrate. Procedures for tissue collection were approved by the Local and National Boards of Medicolegal Affairs in Finland.

[0124] Two of the hybridomas produced antibodies (3-266 (DSM ACC2519) and 3-372 (DSM ACC2590)) that clearly stained lymphatic endothelium both in afferent and efferent lymphatic systems and vascular endothelium on HEV, while the other structures remained unstained. The staining of the lymphatic endothelium is shown in FIG. 1. FIG. 1 is an indirect immunoperoxidase staining that shows that monoclonal antibodies 3-266 and 3-372 recognize endothelium both in afferent and efferent lymphatic systems and on HEV. FIGS. 1a-1c are from the skin. FIGS. 1d-1i are from a lymph node. FIGS. 1a, 1d and 1g show the staining with monoclonal antibody 3-266, FIGS. 1b, 1e and 1h show the staining with monoclonal antibody 3-372 and FIGS. 1c, 1f and 1i show the staining with a negative control antibody, 3G6. In FIGS. 1a, 1b and 1c, the arrows point to the epithelium and arrowheads to afferent lymphatics. In FIGS. 1d and 1e, the arrows point to the lymphatic vessels (lymphatic sinusoids that belong to the efferent lymphatic system) within the lymph node. In FIGS. 1g and 1h, the arrows point to HEV.

EXAMPLE 2

Determination of Molecular Weight of CLEVER-1

[0125] Molecular weight determination was performed by immunoblotting. One percent NP-40 lysates containing of human lymph nodes was analyzed using 5-12.5% sodium dodecyl sulphate-polyacrylamide gel electrophoresis (SDS-PAGE). SDS-PAGE was run in non-reducing conditions. The molecules in the gel were blotted overnight to nitrocellulose sheets and probed with 3-266, 3-372 or a negative control antibody (3G6) (Salmi, M. et al., *J. Exp. Med.* 183:569-579 (1996)). Peroxidase conjugated rabbit anti-mouse Ig was used as the second stage reagent. Detection was performed using enhanced chemiluminescence system according to the instructions of the manufacturer (Amersham).

[0126] Both antibodies recognized a molecule of the same size (about 270-300 kDa; FIG. 2). Due to this and an identical

staining pattern, these antibodies were assumed to recognize the same antigen and this antigen was named CLEVER-1.

EXAMPLE 3

Purification and Molecular Characterization of CLEVER-1

[0127] The molecule recognized by 3-372 antibody was purified from human lymph node lysate overnight (lysis buffer: 150 mM NaCl, 10 mM Trisbase, pH 7.2, 1.5 mM MgCl₂, 1% NP-40, 1% aprotinin, and 1 mM PMSF) as described in Smith, D. J. et al., *J. Exp. Med.* 188:17-27 (1998). After centrifugation, the lysate supernatant was applied sequentially to immunoaffinity columns containing CnBr-activated Sepharose beads armed with irrelevant mAbs and 3-372 (3 mg/ml beads). After washing with lysis buffer, the antigens recognized by 3-372 were eluted with 50 mM triethylamine, frozen and subsequently lyophilized. The eluted material was then subjected to SDS-PAGE analysis and silver staining (O'Connell, K. L. and Stults, J. T., *Electrophoresis* 18:349-359 (1997)). The specific band was excised, reduced, alkylated and digested with trypsin (Promega) overnight at +37° C. as described (Shevchenko, A. et al., *Anal. Chem.* 68:850-855 (1996); O'Connell, K. L. and Stults, J. T., *Electrophoresis* 18:349-359 (1997)). The peptides were analyzed using PerSeptive BioSystems Voyager DE-PRO mass spectrometer operated in the reflectron delayed-extraction mode. Calibration of the spectrum was performed internally by using autolysis products of trypsin or with added calibration mixture 2 (PerSeptive BioSystems). Database search was performed by MS-Fit algorithm (http://prospector.ucsf.edu/_ucsfhtml3.2/msfit.htm) of the University of California, San Francisco mass spectrometry facility.

[0128] After cleavage with trypsin, mass spectrometric analyses yielded 27 peptides. 21 (77%) of those had identical nucleotide sequences with two Genbank entries: AJ 275213, a submission for a cDNA clone called stabilin-1, and D87433, a cDNA clone KIAA0246 isolated from the cell line KG-1. The peptide sequences covered altogether 268 amino acids (10% of the 2570 amino acids of stabilin-1) and spanned the amino acids between 53 and 2301.

[0129] Next we designed primers based on the peptide sequences, the 5' end of the cDNA for stabilin-1 and the 3' end of the cDNA of KIAA0246 and used them to make several RT-PCR fragments that were then ligated together to clone the full-length 7879 bp cDNA (SEQ ID NO:1). Sequencing of the whole construct revealed a high homology with the existing 3' end KIAA0246 sequence available in the data bank. However, it contained 4 nucleotide differences, when compared to the Stabilin sequence. They all cause a change at the amino acid level. Two of these changes are identical with the genomic sequence data available from the HUGO project (AC 006208), but since the genomic clone only covers about half of the gene for this cDNA, the nature of the two other changes remains to be determined.

[0130] Sequencing of several different CLEVER-1 cDNA-clones also revealed the existence of at least two alternatively spliced isoforms of the molecule: the regions covered by exons 23 (nucleotides 2377-2562) and 27 (nucleotides 2914-3009) can be spliced out. We could confirm the existence of one of those splice variants (lacking exon 27) also at the mRNA level (FIG. 7) but the second one (lacking exon 23) that we cloned from a human peripheral lymph node library

was not visible in the system suggesting a low abundance/turnover of the mRNA encoding this form.

[0131] The sequence comparisons revealed significant homologies to proteoglycan link protein-like sequence, epidermal growth factor-like repeats and two RGD motifs being well in line with the adhesive properties of CLEVER-1.

EXAMPLE 4

In Vitro Adhesion Assay

[0132] Lymph node sections were first incubated with 3-266, 3-372 or control antibodies against human HLA ABC (HB-95, ATCC and 3G6 (against chicken T cells) and then overlaid with Ficoll gradient (Pharmacia) purified peripheral blood mononuclear cells or different human tumor cell lines (lymphoblastoid cell lines, KCA and IBW4; a Burkitt lymphoma CRL-1648; squamocellular carcinoma lines NA and NU). Thereafter, the sections were subjected to two different types of assays: 1. A non-static assay, which optimally measures binding of cells to HEV and is performed under rotatory conditions (60 rpm on orbital shaker for 30 min at +7° C.). 2. A static assay, in which the sections overlaid with cells are let to stay in static conditions for 15 min, followed by 5 min of rotation at 60 rpm and then again 15 min without rotation at 7° C. (Static conditions were needed to optimal binding to lymphatic endothelium). The adherent cells were fixed in 1% glutaraldehyde. The number of lymphocytes bound to HEV and to sinusoidal (lymphatic) endothelium was counted single blind under dark-field illumination in which setting the sinusoidal vessels are easily recognizable. The results of the inhibition assays are presented as percentage of control binding (the number of adherent cells/vessel in the presence of control mAb defines 100% adherence).

[0133] When the assay was performed in non-static conditions, mimicking the blood flow, lymphocyte binding to HEV was reduced 43.6% and 45.2% by 3-266 and 3-372, respectively (FIG. 3A). To mimic the conditions at sites of lymphocyte exit, the assay was performed in static conditions. In these assays lymphocyte binding to lymphatic endothelium was decreased 46.4% and 64% by 3-266 and 3-372, respectively (FIG. 3B). These data indicate that the molecule recognized by 3-266 and 3-372 indeed mediate lymphocyte binding both to HEV and to lymphatic endothelium.

[0134] To study the role of this molecule in migration of malignant cells, the assays were performed using three lymphoma cell lines (CRL 1648, KCA and IBW4) and two squamocellular carcinoma cell lines (NA and NU). For these assays 3-372 antibody was chosen because of its higher inhibitory capacity (FIG. 3). The results of these experiments clearly demonstrated that CLEVER-1 is also involved in binding of malignant cells to endothelium both at entrance and exit sites within the lymph nodes (FIGS. 4A and 3).

EXAMPLE 5

CLEVER-1 is Upregulated at Sites of Inflammation on HEV-Like Vessels

[0135] Synovial samples from 18 patients suffering from chronic arthritis and undergoing synovectomies, skin samples from diseased skin of patients suffering from psoriasis (n=5), lichen (n=1), mycosis fungoides (n=1), erythroder-

mia (n=2), exanthema (n=1), folliculitis (n=3) and normal skin samples from 15 individuals were studied for expression of CLEVER-1 using immunoperoxidase method as described above. Like in normal non-lymphoid tissues CLEVER-1 was present in afferent lymphatic vessels in inflamed synovial and both in normal and diseased skin samples. In addition, CLEVER-1 expression was induced on HEV-like vessels that appear at sites of inflammation and are surrounded by heavy infiltrations of inflammatory cells (FIG. 5). Table 2 illustrates complete correlation between the extent of inflammatory infiltration and upregulation of CLEVER-1 expression in synovial samples. The same phenomenon was observed in skin samples: all diseased skin samples had inflammatory infiltrations that contained CLEVER-1 positive HEV-like vessels. Those vessels were absent in normal skin samples.

EXAMPLE 6

Clever-1 also Mediates Binding of Monocytes and Granulocytes to HEV-Like Vessels

[0136] Human monocytes from peripheral blood were purified from Ficoll-gradient (Pharmacia) isolated mononuclear cells by letting them to adhere to plastic surfaces for an hour at +37° C. Granulocytes were purified from leukocyte rich buffy coats from human blood using Percoll-gradient (Pharmacia) centrifugation. Their binding was tested to HEV-like vessels in inflamed synovium. In addition, granulocyte binding was tested to tonsil HEV that brightly express CLEVER-1. (When tonsils are removed they always have variable extent of inflammation, although they as a lymphoid organ have HEV without any inflammation). Both granulocytes and monocytes bound efficiently to HEV-like vessels in inflamed synovium and granulocytes adhered avidly to HEV in tonsils. Their binding to these organs was significantly inhibited by the antibody pool containing 3-372 and 3-266 but not with the control antibody (FIG. 6).

EXAMPLE 7

CLEVER-1 Controls Lymphocyte Trafficking In Vivo

[0137] In order to verify that CLEVER-1 has a functional role in vivo, it was at first confirmed by intravenous injection of 3-372 antibody that rabbits express CLEVER-1 on the surface of endothelium in vivo. The presence of CLEVER-1 on HEV was detected after the 3-372 antibody had circulated 5 minutes in vivo using frozen sections and FITC labelled anti-mouse IgG second stage antibody after sacrifice (FIG. 8a). In this time frame the intravenously given 180 kDa immunoglobulin molecule does not have a possibility to leak and diffuse into the tissue. Based on these results antibody 3-372 (and a class-matched negative control antibody) was given to the rabbits immunized with keyhole limpet hemocyanin to footpads and the effects of the antibody treatment on the size and cellularity of the lymph nodes draining the footpads was analyzed.

[0138] Antibody treatment against CLEVER-1 significantly prevented increase of the size of the popliteal lymph nodes (FIG. 8c) indicating that CLEVER-1 has a functional role in lymphocyte traffic in vivo. Most likely it exerts its effects both at lymphocyte entrance in HEV and their exit in lymphatic sinusoids, because the rabbits treated with 3-372 antibody had only few lymphocytes in their lymphatic sinusoids when analyzed using histological section (FIG. 8d) Intravenously given 3-372 antibody was also detected to bind CLEVER-1 on lymphatic sinuses when tested at sacrifice 3 days after the final 3-372 dose. No signal was detected in rabbits which received a control antibody (data not shown).

[0139] All documents, e.g., scientific publications, patents and patent publications recited herein are hereby incorporated by reference in their entirety to the same extent as if each individual document was specifically and individually indicated to be incorporated by reference in its entirety. Where the document cited only provides the first page of the document, the entire document is intended, including the remaining pages of the document.

TABLE 1

Matches with CLEVER-1 and stabilin-1						
1. 21/27 matches (77%). 275350.0 Da, pI = 6.04. Acc. #6469374. HOMO SAPIENS. (AJ275213) stabilin-1.						
m/z submitted	MH ⁺ matched	Delta ppm	start	end	Peptide Sequence (Click for Fragment Ions)	Modifications
775.488	775.483	6.4034	372	377	(R) VFLQLR(V)	
787.36	787.3739	-17.6253	1299	1305	(R) SGFSFSR(G)	
799.495	799.5042	-11.4617	1585	1591	(R) VGLELLR(D)	
812.495	812.4994	-5.4309	1047	1053	(R) TLPNLVR(A)	
917.502	917.4997	2.4556	1040	1046	(R) AFWLQPR(T)	
1017.44	1017.425	14.4862	2295	2301	(R) WDAYCFR(V)	
1104.54	1104.526	12.6406	53	61	(K) QTCPSGWL(R)	
1212.7	1212.695	3.9482	1021	1032	(R) VTALVPS EA AVR(Q)	
1284.65	1284.622	21.4530	1678	1688	(R) EGS IY LND FA R(V)	
1291.79	1291.774	12.5434	613	624	(R) ILLGPEGVPLQR(V)	
1330.63	1330.575	41.7401	953	965	(R) AGNGGCHGLATCR(A)	

TABLE 1-continued

Matches with CLEVER-1 and stabilin-1						
1. 21/27 matches (77%). 275350.0 Da, pI = 6.04. Acc. #6469374. <i>HOMO SAPIENS</i> . (AJ275213) stabilin-1.						
m/z submitted	MH ⁺ matched	Delta ppm	start	end	Peptide Sequence (Click for Fragment Ions)	Modifications
1330.63	1330.633	-1.9754	1873	1882	(R) <u>CDHFETRPLR</u> (L)	
1374.66	1374.632	20.1117	62	72	(R) <u>ELPDQITQDCR</u> (Y)	
1456.79	1456.776	9.6229	1069	1082	(R) <u>LGGQEVATLNPTTR</u> (W)	
1493.8	1493.796	2.4215	508	521	(R) <u>TIGQILASTEAFSR</u> (F)	
1555.7	1555.663	23.5678	219	231	(R) <u>CLPGYTQGSSECR</u> (A)	
1678.94	1678.913	16.1953	1802	1817	(R) <u>NVEALASDLPLNLGPLR</u> (T)	
1730.89	1730.887	1.9674	1054	1068	(R) <u>AHFLOGALFEEELAR</u> (L)	
1912.82	1912.832	-6.3742	936	952	(K) <u>LGFAGDGYQCSPIIDPCR</u> (A)	
2057.05	2057.03	9.5484	1655	1673	(R) <u>SEDLLEQGYATALSGHPLR</u> (F)	
2165.11	2165.14	-13.6320	1707	1725	(R) <u>VLLPPEAIHWEPDDAPIPR</u> (R)	
2295.22	2295.224	-1.5853	389	410	(R) <u>EILTTAGPFTVLVPSVSSFSRR</u> (T)	

6 unmatched masses: 871.5410 949.4800 1360.6500 1538.6900 1787.9200 2008.1400

The matched peptides cover 10% (268/2570 AA's) of the protein.

Coverage Map for This Hit (MS-Digest index #): 427477

TABLE 2

CLEVER-1 Expression Is Induced Mainly on Vessels Surrounded by Lymphocytic Infiltrations in Inflamed Synovia		
Expression of CLEVER-1	The degree of inflammatory infiltration ²	
	on HEV-like vessels ¹	0/1 (n = 9)
-/+	100%	0
++/+++	0	100%

¹Intensity was scored as -, ±, + negative or weak, ++,+++ moderate or strong.

²Degree of the inflammatory cell infiltration in 18 synovial samples was scored as: 0/1, none or few lymphocytes around the vessels, 2/3 marked or massive lymphocytic infiltrations.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 23

<210> SEQ ID NO 1
 <211> LENGTH: 7879
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens
 <220> FEATURE:
 <221> NAME/KEY: exon
 <222> LOCATION: (31)..(7740)

<400> SEQUENCE: 1

actctgtect ggacagcgtg cccaccagcc atg gcg ggg ccc cgg ggc ctc ctc

54

-continued

Gln	Ala	Phe	Cys	Thr	Cys	Arg	Pro	Gly	Leu	Val	Ser	Ile	Asn	Ser	Asn		
			300					305					310				
gct	tct	gcg	ggc	tgc	ttc	gcc	ttc	tgc	tcc	ccc	ttc	tcc	tgc	gac	cgg		1014
Ala	Ser	Ala	Gly	Cys	Phe	Ala	Phe	Cys	Ser	Pro	Phe	Ser	Cys	Asp	Arg		
		315					320					325					
tct	gcc	act	tgc	cag	gtg	acc	gct	gat	ggg	aag	acc	agc	tgt	gtg	tgc		1062
Ser	Ala	Thr	Cys	Gln	Val	Thr	Ala	Asp	Gly	Lys	Thr	Ser	Cys	Val	Cys		
	330					335					340						
agg	gaa	agc	gag	gtg	ggg	gat	ggg	cgt	gcc	tgc	tac	gga	cac	ctg	ctc		1110
Arg	Glu	Ser	Glu	Val	Gly	Asp	Gly	Arg	Ala	Cys	Tyr	Gly	His	Leu	Leu		
	345				350					355					360		
cac	gag	gtg	cag	aag	gcc	acg	cag	aca	ggc	cgg	gtg	ttc	ctg	cag	ctg		1158
His	Glu	Val	Gln	Lys	Ala	Thr	Gln	Thr	Gly	Arg	Val	Phe	Leu	Gln	Leu		
			365						370					375			
agg	gtc	gcc	gtg	gcc	atg	atg	gac	cag	ggc	tgc	cgg	gaa	atc	ctt	acc		1206
Arg	Val	Ala	Val	Ala	Met	Met	Asp	Gln	Gly	Cys	Arg	Glu	Ile	Leu	Thr		
			380					385					390				
aca	gcg	ggc	cct	ttc	acc	gtg	ctg	gtg	cca	tcc	gtc	tcc	tcc	ttc	tcc		1254
Thr	Ala	Gly	Pro	Phe	Thr	Val	Leu	Val	Pro	Ser	Val	Ser	Ser	Phe	Ser		
		395					400					405					
tcc	agg	acc	atg	aat	gca	tcc	ctt	gcc	cag	cag	ctc	tgt	aga	cag	cac		1302
Ser	Arg	Thr	Met	Asn	Ala	Ser	Leu	Ala	Gln	Gln	Leu	Cys	Arg	Gln	His		
	410					415					420						
atc	atc	gca	ggg	cag	cac	atc	ctg	gag	gac	aca	agg	acc	caa	caa	aca		1350
Ile	Ile	Ala	Gly	Gln	His	Ile	Leu	Glu	Asp	Thr	Arg	Thr	Gln	Gln	Thr		
	425				430				435						440		
cga	agg	tgg	tgg	acg	ctg	gcc	ggg	cag	gag	atc	acc	gtc	acc	ttt	aac		1398
Arg	Arg	Trp	Trp	Thr	Leu	Ala	Gly	Gln	Glu	Ile	Thr	Val	Thr	Phe	Asn		
				445				450						455			
caa	ttc	acg	aaa	tac	tcc	tac	aag	tac	aaa	gac	cag	ccc	cag	cag	acg		1446
Gln	Phe	Thr	Lys	Tyr	Ser	Tyr	Lys	Tyr	Lys	Asp	Gln	Pro	Gln	Gln	Thr		
			460					465					470				
ttc	aac	atc	tac	aag	gcc	aac	aac	ata	gca	gct	aat	ggc	gtc	ttc	cac		1494
Phe	Asn	Ile	Tyr	Lys	Ala	Asn	Asn	Ile	Ala	Ala	Asn	Gly	Val	Phe	His		
		475				480						485					
gtg	gtc	act	ggc	ctg	cgg	tgg	cag	gcc	ccc	tct	ggg	acc	cct	ggg	gat		1542
Val	Val	Thr	Gly	Leu	Arg	Trp	Gln	Ala	Pro	Ser	Gly	Thr	Pro	Gly	Asp		
		490				495					500						
ccc	aag	aga	act	atc	gga	cag	atc	ctc	gcc	tct	acc	gag	gcc	ttc	agc		1590
Pro	Lys	Arg	Thr	Ile	Gly	Gln	Ile	Leu	Ala	Ser	Thr	Glu	Ala	Phe	Ser		
	505				510					515					520		
cgc	ttt	gaa	acc	atc	ctg	gag	aac	tgt	ggg	ctg	ccc	tcc	atc	ctg	gac		1638
Arg	Phe	Glu	Thr	Ile	Leu	Glu	Asn	Cys	Gly	Leu	Pro	Ser	Ile	Leu	Asp		
				525					530					535			
gga	cct	ggg	ccc	ttc	aca	gtc	ttt	gcc	cca	agc	aat	gag	gct	gtg	gac		1686
Gly	Pro	Gly	Pro	Phe	Thr	Val	Phe	Ala	Pro	Ser	Asn	Glu	Ala	Val	Asp		
			540					545					550				
agc	ttg	cgt	gac	ggc	cgc	ctg	atc	tac	ctc	ttc	aca	gcg	ggt	ctc	tct		1734
Ser	Leu	Arg	Asp	Gly	Arg	Leu	Ile	Tyr	Leu	Phe	Thr	Ala	Gly	Leu	Ser		
		555				560						565					
aaa	ctg	cag	gag	ttg	gtg	cgg	tac	cac	atc	tac	aac	cac	ggc	cag	ctg		1782
Lys	Leu	Gln	Glu	Leu	Val	Arg	Tyr	His	Ile	Tyr	Asn	His	Gly	Gln	Leu		
		570				575					580						
acc	gtt	gag	aag	ctc	atc	tcc	aag	ggt	cgg	atc	ctc	acc	atg	gcg	aac		1830
Thr	Val	Glu	Lys	Leu	Ile	Ser	Lys	Gly	Arg	Ile	Leu	Thr	Met	Ala	Asn		
				585		590				595					600		
cag	gtc	ctg	gct	gtg	aac	att	tct	gag	gag	ggg	cgc	atc	ctg	ctg	gga		1878

-continued

Gln	Val	Leu	Ala	Val	Asn	Ile	Ser	Glu	Glu	Gly	Arg	Ile	Leu	Leu	Gly	
				605					610					615		
ccc	gag	ggg	gtc	ccg	ctg	cag	agg	gta	gac	gtg	atg	gcc	gcc	aat	ggg	1926
Pro	Glu	Gly	Val	Pro	Leu	Gln	Arg	Val	Asp	Val	Met	Ala	Ala	Asn	Gly	
			620					625					630			
gtg	atc	cac	atg	ctg	gac	ggc	atc	ctg	ctg	ccc	ccg	acc	atc	ctg	ccc	1974
Val	Ile	His	Met	Leu	Asp	Gly	Ile	Leu	Leu	Pro	Pro	Thr	Ile	Leu	Pro	
		635					640					645				
atc	ctg	ccc	aag	cac	tgc	agc	gag	gag	cag	cac	aag	att	gtg	gcg	ggc	2022
Ile	Leu	Pro	Lys	His	Cys	Ser	Glu	Glu	Gln	His	Lys	Ile	Val	Ala	Gly	
		650				655					660					
tcc	tgt	gtg	gac	tgc	caa	gcc	ctg	aac	acc	agc	acg	tgt	ccc	ccc	aac	2070
Ser	Cys	Val	Asp	Cys	Gln	Ala	Leu	Asn	Thr	Ser	Thr	Cys	Pro	Pro	Asn	
				670						675				680		
agt	gtg	aag	ctg	gac	atc	ttc	ccc	aag	gag	tgt	gtc	tac	atc	cat	gac	2118
Ser	Val	Lys	Leu	Asp	Ile	Phe	Pro	Lys	Glu	Cys	Val	Tyr	Ile	His	Asp	
				685				690						695		
cca	acg	ggg	ctc	aat	gtg	cta	aag	aag	ggc	tgt	gcc	agc	tac	tgc	aac	2166
Pro	Thr	Gly	Leu	Asn	Val	Leu	Lys	Lys	Gly	Cys	Ala	Ser	Tyr	Cys	Asn	
			700					705					710			
caa	acc	atc	atg	gaa	caa	ggc	tgc	tgc	aaa	ggg	ttt	ttc	ggg	cct	gac	2214
Gln	Thr	Ile	Met	Glu	Gln	Gly	Cys	Cys	Lys	Gly	Phe	Phe	Gly	Pro	Asp	
		715					720					725				
tgc	acg	cag	tgt	cct	ggg	ggc	ttc	tcc	aac	ccc	tgc	tat	ggc	aaa	ggc	2262
Cys	Thr	Gln	Cys	Pro	Gly	Gly	Phe	Ser	Asn	Pro	Cys	Tyr	Gly	Lys	Gly	
		730				735					740					
aat	tgc	agt	gat	ggg	atc	cag	ggc	aat	ggg	gcc	tgc	ctc	tgc	ttc	cca	2310
Asn	Cys	Ser	Asp	Gly	Ile	Gln	Gly	Asn	Gly	Ala	Cys	Leu	Cys	Phe	Pro	
				745		750				755				760		
gac	tac	aag	ggc	atc	gcc	tgc	cac	atc	tgc	tgc	aac	cca	aac	aag	cat	2358
Asp	Tyr	Lys	Gly	Ile	Ala	Cys	His	Ile	Cys	Ser	Asn	Pro	Asn	Lys	His	
				765					770					775		
gga	gag	caa	tgc	cag	gaa	gac	tgc	ggc	tgt	gtc	cat	ggg	ctc	tgc	gac	2406
Gly	Glu	Gln	Cys	Gln	Glu	Asp	Cys	Gly	Cys	Val	His	Gly	Leu	Cys	Asp	
			780					785					790			
aac	cgc	cca	ggc	agt	ggg	ggg	gtg	tgc	cag	cag	ggc	acg	tgt	gcc	cct	2454
Asn	Arg	Pro	Gly	Ser	Gly	Gly	Val	Cys	Gln	Gln	Gly	Thr	Cys	Ala	Pro	
			795				800					805				
ggc	ttc	agt	ggc	cgg	ttc	tgc	aac	gag	tcc	atg	ggg	gac	tgt	ggg	ccc	2502
Gly	Phe	Ser	Gly	Arg	Phe	Cys	Asn	Glu	Ser	Met	Gly	Asp	Cys	Gly	Pro	
			810			815					820					
aca	ggg	ctg	gcc	cag	cac	tgc	cac	ctg	cat	gcc	cgc	tgt	gtt	agc	cag	2550
Thr	Gly	Leu	Ala	Gln	His	Cys	His	Leu	His	Ala	Arg	Cys	Val	Ser	Gln	
				825		830				835				840		
gag	ggg	gtt	gcc	aga	tgt	cgc	tgt	ctt	gat	ggc	ttt	gag	ggg	gat	ggc	2598
Glu	Gly	Val	Ala	Arg	Cys	Arg	Cys	Leu	Asp	Gly	Phe	Glu	Gly	Asp	Gly	
				845					850					855		
ttc	tcc	tgc	aca	cct	agc	aac	ccc	tgc	tcc	cac	ccg	gac	cgt	gga	ggc	2646
Phe	Ser	Cys	Thr	Pro	Ser	Asn	Pro	Cys	Ser	His	Pro	Asp	Arg	Gly	Gly	
			860					865					870			
tgc	tca	gag	aat	gct	gag	tgt	gtc	cct	ggg	tcc	ctg	ggc	acc	cac	cac	2694
Cys	Ser	Glu	Asn	Ala	Glu	Cys	Val	Pro	Gly	Ser	Leu	Gly	Thr	His	His	
			875				880						885			
tgc	aca	tgc	cac	aaa	ggc	tgg	agt	ggg	gat	ggc	cgc	gtc	tgt	gtg	gct	2742
Cys	Thr	Cys	His	Lys	Gly	Trp	Ser	Gly	Asp	Gly	Arg	Val	Cys	Val	Ala	
			890			895					900					
att	gac	gag	tgt	gag	ctg	gac	gtg	aga	ggg	ggc	tgc	cac	acc	gat	gcc	2790

-continued

Ile Asp Glu Cys Glu Leu Asp Val Arg Gly Gly Cys His Thr Asp Ala 905 910 915 920	
ctc tgc agc tat gtg ggc ccc ggg cag agc cga tgc acc tgc aag ctg Leu Cys Ser Tyr Val Gly Pro Gly Gln Ser Arg Cys Thr Cys Lys Leu 925 930 935	2838
ggc ttt gcc ggg gat ggc tac cag tgc agc ccc atc gac ccc tgc cgg Gly Phe Ala Gly Asp Gly Tyr Gln Cys Ser Pro Ile Asp Pro Cys Arg 940 945 950	2886
gca ggc aat ggc ggc tgc cac ggc ctg gcc acc tgc cgg gca gtg ggg Ala Gly Asn Gly Gly Cys His Gly Leu Ala Thr Cys Arg Ala Val Gly 955 960 965	2934
gga ggt cag cgg gtc tgc acg tgc ccc cct gcc ttt ggg ggt gat ggc Gly Gly Gln Arg Val Cys Thr Cys Pro Pro Gly Phe Gly Gly Asp Gly 970 975 980	2982
ttc agc tgt tat gga gac atc ttc cgg gag ctg gag gca aat gcc cac Phe Ser Cys Tyr Gly Asp Ile Phe Arg Glu Leu Glu Ala Asn Ala His 985 990 995 1000	3030
ttc tcc atc ttc tac caa tgg ctt aag agt gcc ggc atc acg ctt Phe Ser Ile Phe Tyr Gln Trp Leu Lys Ser Ala Gly Ile Thr Leu 1005 1010 1015	3075
cct gcc gac cgc cga gtc aca gcc ctg gtg ccc tcc gag gct gca Pro Ala Asp Arg Arg Val Thr Ala Leu Val Pro Ser Glu Ala Ala 1020 1025 1030	3120
gtc cgt cag ctg agc ccc gag gac cga gct ttc tgg ctg cag cca Val Arg Gln Leu Ser Pro Glu Asp Arg Ala Phe Trp Leu Gln Pro 1035 1040 1045	3165
agg acg ctg cgg aac ctg gtc agg gcc cat ttt ctc cag ggt gcc Arg Thr Leu Pro Asn Leu Val Arg Ala His Phe Leu Gln Gly Ala 1050 1055 1060	3210
ctc ttc gag gag gag ctg gcc cgg ctg ggt ggg cag gaa gtg gcc Leu Phe Glu Glu Glu Leu Ala Arg Leu Gly Gly Gln Glu Val Ala 1065 1070 1075	3255
acc ctg aac ccc acc aca cgc tgg gag att cgc aac att agt ggg Thr Leu Asn Pro Thr Thr Arg Trp Glu Ile Arg Asn Ile Ser Gly 1080 1085 1090	3300
agg gtc tgg gtg cag aat gcc agc gtg gat gtg gct gac ctc ctt Arg Val Trp Val Gln Asn Ala Ser Val Asp Val Ala Asp Leu Leu 1095 1100 1105	3345
gcc acc aac ggt gtc cta cac atc ctc agc cag gtc tta ctg ccc Ala Thr Asn Gly Val Leu His Ile Leu Ser Gln Val Leu Leu Pro 1110 1115 1120	3390
ccc cga ggg gat gtg ccc ggt ggg cag ggg ttg ctg cag cag ctg Pro Arg Gly Asp Val Pro Gly Gly Gln Gly Leu Leu Gln Gln Leu 1125 1130 1135	3435
gac ttg gtg cct gcc ttc agc ctc ttc cgg gaa ttg ctg cag cac Asp Leu Val Pro Ala Phe Ser Leu Phe Arg Glu Leu Leu Gln His 1140 1145 1150	3480
cat ggg ttg gtg ccc cag att gag gct gcc act gcc tac acc atc His Gly Leu Val Pro Gln Ile Glu Ala Ala Thr Ala Tyr Thr Ile 1155 1160 1165	3525
ttt gtg ccc acc aac cgc tcc ctg gag gcc cag gcc aac agc agt Phe Val Pro Thr Asn Arg Ser Leu Glu Ala Gln Gly Asn Ser Ser 1170 1175 1180	3570
cac ctg gac gca gac aca gtg cgg cac cat gtg gtc ctg ggg gag His Leu Asp Ala Asp Thr Val Arg His His Val Val Leu Gly Glu 1185 1190 1195	3615
gcc ctc tcc atg gaa acc ctg cgg aag ggt gga cac cgc aac tcc	3660

-continued

Ala	Leu	Ser	Met	Glu	Thr	Leu	Arg	Lys	Gly	Gly	His	Arg	Asn	Ser		
				1200					1205					1210		
ctc	ctg	ggc	cct	gcc	cac	tgg	atc	gtc	ttc	tac	aac	cac	agt	ggc		3705
Leu	Leu	Gly	Pro	Ala	His	Trp	Ile	Val	Phe	Tyr	Asn	His	Ser	Gly		
				1215					1220					1225		
cag	cct	gag	gtg	aac	cat	gtg	cca	ctg	gaa	ggc	ccc	atg	ctg	gag		3750
Gln	Pro	Glu	Val	Asn	His	Val	Pro	Leu	Glu	Gly	Pro	Met	Leu	Glu		
				1230					1235					1240		
gcc	cct	ggc	cgc	tgc	ctg	att	ggc	ctg	tgc	ggg	gtc	ctg	acg	gtg		3795
Ala	Pro	Gly	Arg	Ser	Leu	Ile	Gly	Leu	Ser	Gly	Val	Leu	Thr	Val		
				1245					1250					1255		
ggc	tca	agt	cgc	tgc	ctg	cat	agc	cac	gct	gag	gcc	ctg	cgg	gag		3840
Gly	Ser	Ser	Arg	Cys	Leu	His	Ser	His	Ala	Glu	Ala	Leu	Arg	Glu		
				1260					1265					1270		
aaa	tgt	gta	aac	tgc	acc	agg	aga	ttc	cgc	tgc	act	cag	ggc	ttc		3885
Lys	Cys	Val	Asn	Cys	Thr	Arg	Arg	Phe	Arg	Cys	Thr	Gln	Gly	Phe		
				1275					1280					1285		
cag	ctg	cag	gac	aca	ccc	agg	aag	agc	tgt	gtc	tac	cga	tct	ggc		3930
Gln	Leu	Gln	Asp	Thr	Pro	Arg	Lys	Ser	Cys	Val	Tyr	Arg	Ser	Gly		
				1290					1295					1300		
ttc	tcc	ttc	tcc	cgg	ggc	tgc	tct	tac	aca	tgt	gcc	aag	aag	atc		3975
Phe	Ser	Phe	Ser	Arg	Gly	Cys	Ser	Tyr	Thr	Cys	Ala	Lys	Lys	Ile		
				1305					1310					1315		
cag	gtg	ccg	gac	tgc	tgc	cct	ggc	ttc	ttt	ggc	acg	ctg	tgt	gag		4020
Gln	Val	Pro	Asp	Cys	Cys	Pro	Gly	Phe	Phe	Gly	Thr	Leu	Cys	Glu		
				1320					1325					1330		
cca	tgc	cca	ggg	ggc	cta	ggg	ggg	gtg	tgc	tca	ggc	cat	ggg	cag		4065
Pro	Cys	Pro	Gly	Gly	Leu	Gly	Gly	Val	Cys	Ser	Gly	His	Gly	Gln		
				1335					1340					1345		
tgc	cag	gac	agg	ttc	ctg	ggc	agc	ggg	gag	tgc	cac	tgc	cac	gag		4110
Cys	Gln	Asp	Arg	Phe	Leu	Gly	Ser	Gly	Glu	Cys	His	Cys	His	Glu		
				1350					1355					1360		
ggc	ttc	cat	gga	acg	gcc	tgt	gag	gtg	tgt	gag	ctg	ggc	cgc	tac		4155
Gly	Phe	His	Gly	Thr	Ala	Cys	Glu	Val	Cys	Glu	Leu	Gly	Arg	Tyr		
				1365					1370					1375		
ggg	ccc	aac	tgc	acc	gga	gtg	tgt	gac	tgt	gcc	cat	ggg	ctg	tgc		4200
Gly	Pro	Asn	Cys	Thr	Gly	Val	Cys	Asp	Cys	Ala	His	Gly	Leu	Cys		
				1380					1385					1390		
cag	gag	ggg	ctg	caa	ggg	gac	gga	agc	tgt	gtc	tgt	aac	gtg	ggc		4245
Gln	Glu	Gly	Leu	Gln	Gly	Asp	Gly	Ser	Cys	Val	Cys	Asn	Val	Gly		
				1395					1400					1405		
tgg	cag	ggc	ctc	cgc	tgt	gac	cag	aaa	atc	acc	agc	cct	cag	tgc		4290
Trp	Gln	Gly	Leu	Arg	Cys	Asp	Gln	Lys	Ile	Thr	Ser	Pro	Gln	Cys		
				1410					1415					1420		
cct	agg	aag	tgc	gac	ccc	aat	gcc	aac	tgc	gtg	cag	gac	tgc	gcc		4335
Pro	Arg	Lys	Cys	Asp	Pro	Asn	Ala	Asn	Cys	Val	Gln	Asp	Ser	Ala		
				1425					1430					1435		
gga	gcc	tcc	acc	tgc	gcc	tgt	gct	gcg	gga	tac	tcc	ggc	aat	ggc		4380
Gly	Ala	Ser	Thr	Cys	Ala	Cys	Ala	Ala	Gly	Tyr	Ser	Gly	Asn	Gly		
				1440					1445					1450		
atc	ttc	tgt	tca	gag	gtg	gac	ccc	tgc	gcc	cac	ggc	cat	ggg	ggc		4425
Ile	Phe	Cys	Ser	Glu	Val	Asp	Pro	Cys	Ala	His	Gly	His	Gly	Gly		
				1455					1460					1465		
tgc	tcc	cct	cat	gcc	aac	tgt	acc	aag	gtg	gca	cct	ggg	cag	cgg		4470
Cys	Ser	Pro	His	Ala	Asn	Cys	Thr	Lys	Val	Ala	Pro	Gly	Gln	Arg		
				1470					1475					1480		
aca	tgc	acc	tgc	cag	gat	ggc	tac	atg	ggc	gac	ggg	gag	ctg	tgc		4515

-continued

Thr	Cys	Thr	Cys	Gln	Asp	Gly	Tyr	Met	Gly	Asp	Gly	Glu	Leu	Cys	
				1485					1490					1495	
cag	gaa	att	aac	agc	tgt	ctc	atc	cac	cac	ggg	ggc	tgc	cac	att	4560
Gln	Glu	Ile	Asn	Ser	Cys	Leu	Ile	His	His	Gly	Gly	Cys	His	Ile	
				1500					1505					1510	
cac	gcc	gag	tgc	atc	ccc	act	ggc	ccc	cag	cag	gtc	tcc	tgc	agc	4605
His	Ala	Glu	Cys	Ile	Pro	Thr	Gly	Pro	Gln	Gln	Val	Ser	Cys	Ser	
				1515					1520					1525	
tgc	cgt	gag	ggt	tac	agc	ggg	gat	ggc	atc	cgg	acc	tgc	gag	ctc	4650
Cys	Arg	Glu	Gly	Tyr	Ser	Gly	Asp	Gly	Ile	Arg	Thr	Cys	Glu	Leu	
				1530					1535					1540	
ctg	gac	ccc	tgc	tct	aag	aac	aat	gga	gga	tgc	agc	cca	tat	gcc	4695
Leu	Asp	Pro	Cys	Ser	Lys	Asn	Asn	Gly	Gly	Cys	Ser	Pro	Tyr	Ala	
				1545					1550					1555	
acc	tgc	aaa	agc	aca	ggg	gat	ggc	cag	agg	aca	tgt	acc	tgc	gac	4740
Thr	Cys	Lys	Ser	Thr	Gly	Asp	Gly	Gln	Arg	Thr	Cys	Thr	Cys	Asp	
				1560					1565					1570	
aca	gcc	cac	acc	gtg	ggg	gac	ggc	ctc	acc	tgc	cgt	gcc	cga	gtc	4785
Thr	Ala	His	Thr	Val	Gly	Asp	Gly	Leu	Thr	Cys	Arg	Ala	Arg	Val	
				1575					1580					1585	
ggc	ctg	gag	ctc	ctg	agg	gat	aag	cat	gcc	tca	ttc	ttc	agc	ctc	4830
Gly	Leu	Glu	Leu	Leu	Arg	Asp	Lys	His	Ala	Ser	Phe	Phe	Ser	Leu	
				1590					1595					1600	
cgc	ctc	ctg	gaa	tat	aag	gag	ctc	aag	ggc	gat	ggg	cct	ttc	acc	4875
Arg	Leu	Leu	Glu	Tyr	Lys	Glu	Leu	Lys	Gly	Asp	Gly	Pro	Phe	Thr	
				1605					1610					1615	
atc	ttc	gtg	ccg	cac	gca	gat	cta	atg	agc	aac	ctg	tcg	cag	gat	4920
Ile	Phe	Val	Pro	His	Ala	Asp	Leu	Met	Ser	Asn	Leu	Ser	Gln	Asp	
				1620					1625					1630	
gag	ctg	gcc	egg	att	cgt	gcg	cat	cgc	cag	ctg	gtg	ttt	cgc	tac	4965
Glu	Leu	Ala	Arg	Ile	Arg	Ala	His	Arg	Gln	Leu	Val	Phe	Arg	Tyr	
				1635					1640					1645	
cac	gtg	ggt	ggc	tgt	cgg	egg	ctg	cgg	agc	gag	gac	ctg	ctg	gag	5010
His	Val	Val	Gly	Cys	Arg	Arg	Leu	Arg	Ser	Glu	Asp	Leu	Leu	Glu	
				1650					1655					1660	
cag	ggg	tac	gcc	acg	gcc	ctc	tca	ggg	cac	cca	ctg	cgc	ttc	agc	5055
Gln	Gly	Tyr	Ala	Thr	Ala	Leu	Ser	Gly	His	Pro	Leu	Arg	Phe	Ser	
				1665					1670					1675	
gag	agg	gag	ggc	agc	ata	tac	ctc	aat	gac	ttc	gcg	cgc	gtg	gtg	5100
Glu	Arg	Glu	Gly	Ser	Ile	Tyr	Leu	Asn	Asp	Phe	Ala	Arg	Val	Val	
				1680					1685					1690	
agc	agc	gac	cat	gag	gcc	gtg	aac	ggc	atc	ctg	cac	ttc	att	gac	5145
Ser	Ser	Asp	His	Glu	Ala	Val	Asn	Gly	Ile	Leu	His	Phe	Ile	Asp	
				1695					1700					1705	
cgt	gtc	ctg	ctg	ccc	ccc	gag	gcg	ctg	cac	tgg	gag	cct	gat	gat	5190
Arg	Val	Leu	Leu	Pro	Pro	Glu	Ala	Leu	His	Trp	Glu	Pro	Asp	Asp	
				1710					1715					1720	
gct	ccc	atc	ccg	agg	aga	aat	gtc	acc	gcc	gcc	gcc	cag	ggc	ttc	5235
Ala	Pro	Ile	Pro	Arg	Arg	Asn	Val	Thr	Ala	Ala	Ala	Gln	Gly	Phe	
				1725					1730					1735	
ggt	tac	aag	atc	ttc	agc	ggc	ctc	ctg	aag	gtg	gcc	ggc	ctc	ctg	5280
Gly	Tyr	Lys	Ile	Phe	Ser	Gly	Leu	Leu	Lys	Val	Ala	Gly	Leu	Leu	
				1740					1745					1750	
ccc	ctg	ctt	cga	gag	gca	tcc	cat	agg	ccc	ttc	aca	atg	ctg	tgg	5325
Pro	Leu	Leu	Arg	Glu	Ala	Ser	His	Arg	Pro	Phe	Thr	Met	Leu	Trp	
				1755					1760					1765	
ccc	aca	gac	gcc	gcc	ttt	cga	gct	ctg	cct	ccg	gat	cgc	cag	gcc	5370

-continued

Pro Thr Asp Ala Ala Phe Arg Ala Leu Pro Pro Asp Arg Gln Ala	1770	1775	1780
tgg ctg tac cat gag gac cac cgt gac aag cta gca gcc att ctg			5415
Trp Leu Tyr His Glu Asp His Arg Asp Lys Leu Ala Ala Ile Leu	1785	1790	1795
cgg ggc cac atg att cgc aat gtc gag gcc ttg gca tct gac ctg			5460
Arg Gly His Met Ile Arg Asn Val Glu Ala Leu Ala Ser Asp Leu	1800	1805	1810
ccc aac ctg ggc cca ctt cga acc atg cat ggg acc ccc atc tct			5505
Pro Asn Leu Gly Pro Leu Arg Thr Met His Gly Thr Pro Ile Ser	1815	1820	1825
ttc tcc tgc agc cga acg cgg ccc ggt gag ctc atg gtg ggt gag			5550
Phe Ser Cys Ser Arg Thr Arg Pro Gly Glu Leu Met Val Gly Glu	1830	1835	1840
gat gat gct cgc att gtg cag cgg cac ttg ccc ttt gag ggt ggc			5595
Asp Asp Ala Arg Ile Val Gln Arg His Leu Pro Phe Glu Gly Gly	1845	1850	1855
ctg gcc tat ggc atc gac cag ctg ctg gag cca cct ggc ctt ggt			5640
Leu Ala Tyr Gly Ile Asp Gln Leu Leu Glu Pro Pro Gly Leu Gly	1860	1865	1870
gct cgc tgt gac cac ttt gag acc cgg ccc ctg cga ctg aac acc			5685
Ala Arg Cys Asp His Phe Glu Thr Arg Pro Leu Arg Leu Asn Thr	1875	1880	1885
tgc agc atc tgt ggg ctg gag cca ccc tgt cct gag ggg tca cag			5730
Cys Ser Ile Cys Gly Leu Glu Pro Pro Cys Pro Glu Gly Ser Gln	1890	1895	1900
gag cag ggc agc cct gag gcc tgc tgg cgc ttc tac ccg aag ttc			5775
Glu Gln Gly Ser Pro Glu Ala Cys Trp Arg Phe Tyr Pro Lys Phe	1905	1910	1915
tgg acg tcc cct ccg ctg cac tct ttg gga tta cgc agc gtc tgg			5820
Trp Thr Ser Pro Pro Leu His Ser Leu Gly Leu Arg Ser Val Trp	1920	1925	1930
gtc cac ccc agc ctt tgg ggt agg ccc caa ggc ctg ggc agg ggc			5865
Val His Pro Ser Leu Trp Gly Arg Pro Gln Gly Leu Gly Arg Gly	1935	1940	1945
tgc cac cgc aat tgt gtc acc acc acc tgg aag ccc agc tgc tgc			5910
Cys His Arg Asn Cys Val Thr Thr Thr Trp Lys Pro Ser Cys Cys	1950	1955	1960
cct ggt cac tat ggc agt gag tgc caa gct tgc cct ggc ggc ccc			5955
Pro Gly His Tyr Gly Ser Glu Cys Gln Ala Cys Pro Gly Gly Pro	1965	1970	1975
agc agc cct tgt agt gac cgt ggc gtg tgc atg gac ggc atg agt			6000
Ser Ser Pro Cys Ser Asp Arg Gly Val Cys Met Asp Gly Met Ser	1980	1985	1990
ggc agt ggg cag tgt ctg tgc cgt tca ggt ttt gct ggg aca gcc			6045
Gly Ser Gly Gln Cys Leu Cys Arg Ser Gly Phe Ala Gly Thr Ala	1995	2000	2005
tgt gaa ctc tgt gct cct ggt gcc ttt ggg ccc cat tgt caa gcc			6090
Cys Glu Leu Cys Ala Pro Gly Ala Phe Gly Pro His Cys Gln Ala	2010	2015	2020
tgc cgc tgc act gtg cat ggc cgc tgt gat gag ggc ctt ggg ggc			6135
Cys Arg Cys Thr Val His Gly Arg Cys Asp Glu Gly Leu Gly Gly	2025	2030	2035
tct ggc tcc tgc ttc tgt gat gaa ggc tgg act ggg cca cgc tgt			6180
Ser Gly Ser Cys Phe Cys Asp Glu Gly Trp Thr Gly Pro Arg Cys	2040	2045	2050
gag gtg caa ctg gag ctg cag cct gtg tgt acc cca ccc tgt gca			6225

-continued

Glu Val Gln Leu	Glu	Leu Gln Pro Val Cys	Thr Pro Pro Cys Ala	
	2055		2060	2065
ccc gag gct gtg tgc	cgt gca ggc aac agc	tgt gag tgc agc ctg		6270
Pro Glu Ala Val Cys	Arg Ala Gly Asn Ser	Cys Glu Cys Ser Leu		
	2070	2075	2080	
ggc tat gaa ggg gat	ggc cgc gtg tgt aca	gtg gca gac ctg tgc		6315
Gly Tyr Glu Gly Asp	Gly Arg Val Cys Thr	Val Ala Asp Leu Cys		
	2085	2090	2095	
cag gac ggg cat ggt	ggc tgc agt gag cac	gcc aac tgt agc cag		6360
Gln Asp Gly His	Gly Cys Ser Glu His	Ala Asn Cys Ser Gln		
	2100	2105	2110	
gta gga aca atg gtc	act tgt acc tgc ctg	ccc gac tac gag ggt		6405
Val Gly Thr Met Val	Thr Cys Thr Cys Leu	Pro Asp Tyr Glu Gly		
	2115	2120	2125	
gat ggc tgg agc tgc	cgg gcc cgc aac ccc	tgc aca gat ggc cac		6450
Asp Gly Trp Ser	Arg Ala Arg Asn Pro	Cys Thr Asp Gly His		
	2130	2135	2140	
cgc ggg ggc tgc agc	gag cac gcc aac tgc	ttg agc acc ggc ctg		6495
Arg Gly Gly Cys Ser	Glu His Ala Asn Cys	Leu Ser Thr Gly Leu		
	2145	2150	2155	
aac aca cgg cgc tgt	gag tgc cac gca ggc	tac gta ggc gat gga		6540
Asn Thr Arg Arg Cys	Glu Cys His Ala Gly	Tyr Val Gly Asp Gly		
	2160	2165	2170	
ctg cag tgt ctg gag	gag tgg gaa cca cct	gtg gac cgc tgc ttg		6585
Leu Gln Cys Leu Glu	Glu Ser Glu Pro Pro	Val Asp Arg Cys Leu		
	2175	2180	2185	
ggc cag cca cgg ccc	tgc cac tca gat gcc	atg tgc act gac ctg		6630
Gly Gln Pro Pro Pro	Cys His Ser Asp Ala	Met Cys Thr Asp Leu		
	2190	2195	2200	
cac ttc cag gag aaa	cgg gct ggc gtt ttc	cac ctc cag gcc acc		6675
His Phe Gln Glu Lys	Arg Ala Gly Val Phe	His Leu Gln Ala Thr		
	2205	2210	2215	
agc ggc cct tat ggt	ctg aac ttt tgg gag	gct gag gcg gca tgc		6720
Ser Gly Pro Tyr Gly	Leu Asn Phe Ser Glu	Ala Glu Ala Ala Cys		
	2220	2225	2230	
gaa gca cag gga gcc	gtc ctt gct tca ttc	cct cag ctc tct gct		6765
Glu Ala Gln Gly Ala	Val Leu Ala Ser Phe	Pro Gln Leu Ser Ala		
	2235	2240	2245	
gcc cag cag ctg ggc	ttc cac ctg tgc ctc	atg ggc tgg ctg gcc		6810
Ala Gln Gln Leu Gly	Phe His Leu Cys Leu	Met Gly Trp Leu Ala		
	2250	2255	2260	
aat ggc tcc act gcc	cac cct gtg gtt ttc	cct gtg gcg gac tgt		6855
Asn Gly Ser Thr Ala	His Pro Val Val Phe	Pro Val Ala Asp Cys		
	2265	2270	2275	
ggc aat ggt cgg gtg	ggc gta gtc agc ctg	ggt gcc cgc aag aac		6900
Gly Asn Gly Arg Val	Gly Val Val Ser Leu	Gly Ala Arg Lys Asn		
	2280	2285	2290	
ctc tca gaa cgc tgg	gat gcc tac tgc ttc	cgt gtg caa gat gtg		6945
Leu Ser Glu Arg Trp	Asp Ala Tyr Cys Phe	Arg Val Gln Asp Val		
	2295	2300	2305	
gcc tgc cga tgc cga	aat ggc ttc gtg ggt	gac ggg atc agc acg		6990
Ala Cys Arg Cys Arg	Asn Gly Phe Val Gly	Asp Gly Ile Ser Thr		
	2310	2315	2320	
tgc aat ggg aag ctg	ctg gat gtg ctg gct	gcc act gcc aac ttc		7035
Cys Asn Gly Lys Leu	Leu Asp Val Leu Ala	Ala Thr Ala Asn Phe		
	2325	2330	2335	
tcc acc ttc tat ggg	atg cta ttg ggc tat	gcc aat gcc acc cag		7080

-continued

Ser	Thr	Phe	Tyr	Gly	Met	Leu	Leu	Gly	Tyr	Ala	Asn	Ala	Thr	Gln	
				2340					2345					2350	
cgg	ggt	ctc	gac	ttc	ctg	gac	ttc	ctg	gat	gat	gag	ctc	acg	tat	7125
Arg	Gly	Leu	Asp	Phe	Leu	Asp	Phe	Leu	Asp	Asp	Glu	Leu	Thr	Tyr	
				2355					2360					2365	
aag	aca	ctc	ttc	gtc	cct	gtc	aat	gaa	ggc	ttt	gtg	gac	aac	atg	7170
Lys	Thr	Leu	Phe	Val	Pro	Val	Asn	Glu	Gly	Phe	Val	Asp	Asn	Met	
				2370					2375					2380	
acg	ctg	agt	ggc	cca	gac	ttg	gag	ctg	cat	gcc	tcc	aac	gcc	acc	7215
Thr	Leu	Ser	Gly	Pro	Asp	Leu	Glu	Leu	His	Ala	Ser	Asn	Ala	Thr	
				2385					2390					2395	
ctc	cta	agt	gcc	aac	gcc	agc	cag	ggg	aag	ttg	ctt	ccg	gcc	cac	7260
Leu	Leu	Ser	Ala	Asn	Ala	Ser	Gln	Gly	Lys	Leu	Leu	Pro	Ala	His	
				2400					2405					2410	
tca	ggc	ctc	agc	ctc	atc	atc	agt	gac	gca	ggc	cct	gac	aac	agt	7305
Ser	Gly	Leu	Ser	Leu	Ile	Ile	Ser	Asp	Ala	Gly	Pro	Asp	Asn	Ser	
				2415					2420					2425	
tcc	tgg	gcc	cct	gtg	gcc	cca	ggg	aca	gtt	gtg	ggt	agc	cgt	atc	7350
Ser	Trp	Ala	Pro	Val	Ala	Pro	Gly	Thr	Val	Val	Val	Ser	Arg	Ile	
				2430					2435					2440	
att	gtg	tgg	gac	atc	atg	gcc	ttc	aat	ggc	atc	atc	cat	gct	ctg	7395
Ile	Val	Trp	Asp	Ile	Met	Ala	Phe	Asn	Gly	Ile	Ile	His	Ala	Leu	
				2445					2450					2455	
gcc	agc	ccc	ctc	ctg	gca	ccc	cca	cag	ccc	cag	gca	gtg	ctg	gcg	7440
Ala	Ser	Pro	Leu	Leu	Ala	Pro	Pro	Gln	Pro	Gln	Ala	Val	Leu	Ala	
				2460					2465					2470	
cct	gaa	gcc	cca	cct	gtg	gcg	gca	ggc	gtg	ggg	gct	gtg	ctt	gcc	7485
Pro	Glu	Ala	Pro	Pro	Val	Ala	Ala	Gly	Val	Gly	Ala	Val	Leu	Ala	
				2475					2480					2485	
gct	gga	gca	ctg	ctt	ggc	ttg	gtg	gcc	gga	gct	ctc	tac	ctc	cgt	7530
Ala	Gly	Ala	Leu	Leu	Gly	Leu	Val	Ala	Gly	Ala	Leu	Tyr	Leu	Arg	
				2490					2495					2500	
gcc	cga	ggc	aag	ccc	acg	ggc	ttt	ggc	ttc	tct	gcc	ttc	cag	gcg	7575
Ala	Arg	Gly	Lys	Pro	Thr	Gly	Phe	Gly	Phe	Ser	Ala	Phe	Gln	Ala	
				2505					2510					2515	
gaa	gat	gat	gct	gac	gac	gac	ttc	tca	ccg	tgg	caa	gaa	ggg	acc	7620
Glu	Asp	Asp	Ala	Asp	Asp	Asp	Phe	Ser	Pro	Trp	Gln	Glu	Gly	Thr	
				2520					2525					2530	
aac	ccc	acc	ctg	gtc	tct	gtc	ccc	aac	cct	gtc	ttt	ggc	agc	gac	7665
Asn	Pro	Thr	Leu	Val	Ser	Val	Pro	Asn	Pro	Val	Phe	Gly	Ser	Asp	
				2535					2540					2545	
acc	ttt	tgt	gaa	ccc	ttc	gat	gac	tca	ctg	ctg	gag	gag	gac	ttc	7710
Thr	Phe	Cys	Glu	Pro	Phe	Asp	Asp	Ser	Leu	Leu	Glu	Glu	Asp	Phe	
				2550					2555					2560	
cct	gac	acc	cag	agg	atc	ctc	aca	gtc	aag	tgacgaggct	ggggctgaaa				7760
Pro	Asp	Thr	Gln	Arg	Ile	Leu	Thr	Val	Lys						
				2565					2570						
gcagaagcat	gcacagggag	gagaccactt	ttattgcttg	tctgggtgga	tggggcagga										7820
ggggctgagg	gcctgtccca	gacaataaag	tgccctcagc	ggatgtgggc	catgtcacc										7879

<210> SEQ ID NO 2

<211> LENGTH: 6

<212> TYPE: PRT

<213> ORGANISM: homo sapiens

<400> SEQUENCE: 2

Val Phe Leu Gln Leu Arg

-continued

1 5

<210> SEQ ID NO 3
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: homo sapiens

<400> SEQUENCE: 3

Ser Gly Phe Ser Phe Ser Arg
1 5

<210> SEQ ID NO 4
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: homo sapiens

<400> SEQUENCE: 4

Val Gly Leu Glu Leu Leu Arg
1 5

<210> SEQ ID NO 5
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: homo sapiens

<400> SEQUENCE: 5

Thr Leu Pro Asn Leu Val Arg
1 5

<210> SEQ ID NO 6
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: homo sapiens

<400> SEQUENCE: 6

Ala Phe Trp Leu Gln Pro Arg
1 5

<210> SEQ ID NO 7
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: homo sapiens

<400> SEQUENCE: 7

Trp Asp Ala Tyr Cys Phe Arg
1 5

<210> SEQ ID NO 8
<211> LENGTH: 9
<212> TYPE: PRT
<213> ORGANISM: homo sapiens

<400> SEQUENCE: 8

Gln Thr Cys Pro Ser Gly Trp Leu Arg
1 5

<210> SEQ ID NO 9
<211> LENGTH: 12
<212> TYPE: PRT
<213> ORGANISM: homo sapiens

<400> SEQUENCE: 9

-continued

Val Thr Ala Leu Val Pro Ser Glu Ala Ala Val Arg
1 5 10

<210> SEQ ID NO 10
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: homo sapiens

<400> SEQUENCE: 10

Glu Gly Ser Ile Tyr Leu Asn Asp Phe Ala Arg
1 5 10

<210> SEQ ID NO 11
<211> LENGTH: 12
<212> TYPE: PRT
<213> ORGANISM: homo sapiens

<400> SEQUENCE: 11

Ile Leu Leu Gly Pro Glu Gly Val Pro Leu Gln Arg
1 5 10

<210> SEQ ID NO 12
<211> LENGTH: 13
<212> TYPE: PRT
<213> ORGANISM: homo sapiens

<400> SEQUENCE: 12

Ala Gly Asn Gly Gly Cys His Gly Leu Ala Thr Cys Arg
1 5 10

<210> SEQ ID NO 13
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: homo sapiens

<400> SEQUENCE: 13

Cys Asp His Phe Glu Thr Arg Pro Leu Arg
1 5 10

<210> SEQ ID NO 14
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: homo sapiens

<400> SEQUENCE: 14

Glu Leu Pro Asp Gln Ile Thr Gln Asp Cys Arg
1 5 10

<210> SEQ ID NO 15
<211> LENGTH: 14
<212> TYPE: PRT
<213> ORGANISM: homo sapiens

<400> SEQUENCE: 15

Leu Gly Gly Gln Glu Val Ala Thr Leu Asn Pro Thr Thr Arg
1 5 10

<210> SEQ ID NO 16
<211> LENGTH: 14
<212> TYPE: PRT
<213> ORGANISM: homo sapiens

<400> SEQUENCE: 16

-continued

Thr Ile Gly Gln Ile Leu Ala Ser Thr Glu Ala Phe Ser Arg
 1 5 10

<210> SEQ ID NO 17
 <211> LENGTH: 13
 <212> TYPE: PRT
 <213> ORGANISM: homo sapiens

<400> SEQUENCE: 17

Cys Leu Pro Gly Tyr Thr Gln Gln Gly Ser Glu Cys Arg
 1 5 10

<210> SEQ ID NO 18
 <211> LENGTH: 16
 <212> TYPE: PRT
 <213> ORGANISM: homo sapiens

<400> SEQUENCE: 18

Asn Val Glu Ala Leu Ala Ser Asp Leu Pro Asn Leu Gly Pro Leu Arg
 1 5 10 15

<210> SEQ ID NO 19
 <211> LENGTH: 15
 <212> TYPE: PRT
 <213> ORGANISM: homo sapiens

<400> SEQUENCE: 19

Ala His Phe Leu Gln Gly Ala Leu Phe Glu Glu Glu Leu Ala Arg
 1 5 10 15

<210> SEQ ID NO 20
 <211> LENGTH: 17
 <212> TYPE: PRT
 <213> ORGANISM: homo sapiens

<400> SEQUENCE: 20

Leu Gly Phe Ala Gly Asp Gly Tyr Gln Cys Ser Pro Ile Asp Pro Cys
 1 5 10 15

Arg

<210> SEQ ID NO 21
 <211> LENGTH: 19
 <212> TYPE: PRT
 <213> ORGANISM: homo sapiens

<400> SEQUENCE: 21

Ser Glu Asp Leu Leu Glu Gln Gly Tyr Ala Thr Ala Leu Ser Gly His
 1 5 10 15

Pro Leu Arg

<210> SEQ ID NO 22
 <211> LENGTH: 19
 <212> TYPE: PRT
 <213> ORGANISM: homo sapiens

<400> SEQUENCE: 22

Val Leu Leu Pro Pro Glu Ala Leu His Trp Glu Pro Asp Asp Ala Pro
 1 5 10 15

Ile Pro Arg

-continued

<210> SEQ ID NO 23
 <211> LENGTH: 22
 <212> TYPE: PRT
 <213> ORGANISM: homo sapiens

<400> SEQUENCE: 23

Glu Ile Leu Thr Thr Ala Gly Pro Phe Thr Val Leu Val Pro Ser Val
 1 5 10 15
 Ser Ser Phe Ser Ser Arg
 20

1. A purified lymphatic endothelial glycoprotein, CLEVER-1, having a molecular weight of 270-300 kD in SDS-PAGE under non-reducing conditions, recognisable by a monoclonal antibody selected from the group consisting of (a) monoclonal antibody, DSM ACC 2519; and (b) monoclonal antibody, DSM ACC 2590.

2. The glycoprotein according to claim 1, encoded by a nucleic acid selected from the group consisting of (a) sequence of SEQ ID NO: 1; (b) sequence of SEQ ID NO: 1 lacking nucleotides 2377-2562; and (c) sequence of SEQ ID NO: 1 lacking nucleotides 2914-3009.

3. A CLEVER-1 antibody.

4. The CLEVER-1 antibody of claim 3, which is monoclonal antibody 3-266 (DSM ACC2519).

5. The CLEVER-1 antibody of claim 3, which is monoclonal antibody 3-372 (DSM ACC2590).

6. A cell-free preparation comprising CLEVER-1.

7. A method of diagnosing inflammatory diseases in a patient, said method comprising: (a) exposing a blood or tissue sample from said patient to CLEVER-1 in vitro for a period of time and under conditions sufficient to allow binding of leukocytes if present in said sample; and (b) detecting leukocytes bound to said CLEVER-1 in said blood or tissue sample.

8. A method of detecting malignant cells in a patient, said method comprising: (a) exposing a blood or tissue sample from a patient to CLEVER-1 in vitro for a period of time and under conditions sufficient to allow binding of said malignant cells if present in said sample; and (b) detecting whether any malignant cells bound to said CLEVER-1 in said blood or tissue sample.

9. The method of claim 8, wherein said CLEVER-1 is on a solid support.

10. The method of claim 8, wherein said CLEVER-1 is provided on lymphoid tissue.

11. The method of claim 8, wherein said CLEVER-1 is present in the membrane of endothelial cells.

12. The method of claim 8, wherein said CLEVER-1 is in a soluble form.

13. The method of claim 8, wherein said detection step is performed by imaging.

14. A method of identifying an agent that inhibits the binding of CLEVER-1 to cells, said method comprising: (a) providing an agent to cells in the presence of CLEVER-1; and (b) comparing the binding of CLEVER-1 to cells in (a) to binding of CLEVER-1 in the absence of said agent.

15. A method of identifying an agent that stimulates the binding of CLEVER-1 to cells, said method comprising: (a) providing an agent to cells in the presence of CLEVER-1; and

(b) comparing the binding of CLEVER-1 to cells in (A) to binding of CLEVER-1 in the absence of said agent.

16. A method of removing malignant cells from a sample, said method comprising: (a) exposing said malignant cells to CLEVER-1 in vitro for a period of time and under conditions sufficient to allow binding of said malignant cells if present in said sample, and (b) separating said CLEVER-1 and the malignant cells that adhere thereto from said sample.

17. (canceled)

18. A method of inhibiting metastasis in a patient in need of the same, said method comprising administering an agent that inhibits CLEVER-1 mediated malignant cell binding to said patient, wherein said inhibiting agent is selected from the group consisting of: (a) CLEVER-1 antibodies or fragments thereof; and (b) soluble CLEVER-1 or fragments thereof that bind to said malignant cell.

19. The method of claim 18, wherein said CLEVER-1 mediated cell binding inhibits leukocyte binding to endothelial cells in said patient's systemic vasculature and lymphatic system.

20. The method of claim 18, wherein said CLEVER-1 mediated cell binding inhibits lymphocyte binding to endothelial cells in said patient's systemic vasculature and lymphatic system.

21. The method of claim 18, wherein said CLEVER-1 mediated cell binding inhibits monocyte binding to endothelial cells in said patient's systemic vasculature and lymphatic system.

22. The method of claim 18, wherein said CLEVER-1 mediated cell binding inhibits granulocyte binding to endothelial cells in said patient's systemic vasculature and lymphatic system.

23. The method of claim 18, wherein said CLEVER-1 mediated cell binding inhibits malignant cell binding to endothelial cells in said patient's systemic vasculature and lymphatic system.

24. The method of claim 18, wherein said CLEVER-1 binding agent is administered to a patient in need of an inhibition of CLEVER-1 mediated cell binding.

25. The method of claim 24, wherein said patient is in need of treatment of a malignancy.

26. The method of claim 24, wherein said patient is in need of treatment for a possible malignancy.

27. A method of stimulating CLEVER-1 binding in a patient in need of the same, said method comprising: administering an agent that stimulates CLEVER-1 mediated leukocyte binding to said patient.

28. A method of inhibiting metastasis in a patient in need of the same, said method comprising administering, to said

patient, an agent that inhibits CLEVER-1 mediated endothelial cell binding to CLEVER-1-binding malignant cells in said patient, wherein said agent is a CLEVER-1 antagonist antibody or a fragment thereof that contains an antigen binding site.

29. The method of claim **28**, wherein said antagonist antibody is a monoclonal antibody.

30. The method of claim **28**, wherein said CLEVER-1 antibody is antibody 3-266 produced by the hybridoma DSM ACC2519 or antibody 3-372 produced by the hybridoma DSM ACC2590, or said fragment thereof, or a chimeric, humanized or primatized antibody thereof.

31. The method of claim **30**, wherein said agent is antibody 3-266 produced by the hybridoma DSM ACC2519, or fragment thereof that contains an antigen binding site.

32. The method of claim **30**, wherein said agent is antibody 3-372 produced by the hybridoma DSM ACC2590, or fragment thereof that contains an antigen binding site.

33. The method of claim **30**, wherein said antibody is said antibody fragment.

34. The method of claim **33**, wherein said fragment is a Fab, F(ab')₂ or Fv antibody fragment.

35. The method of claim **30**, wherein said antibody is said chimeric, humanized or primatized antibody.

36. The method of claim **28**, wherein said agent inhibits CLEVER-1 mediated endothelial cell binding to CLEVER-1-binding tumor cells in the systemic vasculature of said patient.

37. The method of claim **28**, wherein said agent inhibits CLEVER-1 mediated endothelial cell binding to CLEVER-1-binding tumor cells in the afferent lymphatics of said patient.

38. The method of claim **28**, wherein said agent inhibits CLEVER-1 mediated endothelial cell binding to CLEVER-1-binding tumor cells in the efferent lymphatics of said patient.

* * * * *

专利名称(译)	常见的淋巴管内皮和血管内皮受体-1 (CLEVER-1) 及其用途		
公开(公告)号	US20080267958A1	公开(公告)日	2008-10-30
申请号	US12/073451	申请日	2008-03-05
申请(专利权)人(译)	法龙制药OY		
当前申请(专利权)人(译)	法龙制药OY		
[标]发明人	JALKANEN SIRPA IRJALA HEIKKI SALMI MARKO		
发明人	JALKANEN, SIRPA IRJALA, HEIKKI SALMI, MARKO		
IPC分类号	C07K16/28 A61K39/395 C07K16/30 G01N33/53 C12N5/06 G01N33/574 C07K14/705 G01N33/50 A61K45/00 A61P29/00 A61P35/00 A61P35/04 A61P43/00 C12N15/02 C12P21/08 C12Q1/02 G01N33 /15		
CPC分类号	A61K2039/505 C07K14/7056 C07K16/28 A61P29/00		
优先权	10/497991 2004-11-30 US PCT/FI2003/000010 2003-01-08 WO 60/346288 2002-01-09 US		
其他公开文献	US7910097		
外部链接	Espacenet USPTO		

摘要(译)

描述了一种新的蛋白质共同淋巴内皮和血管内皮细胞受体-1 (CLEVER-1)。CLEVER-1介导白细胞和恶性细胞与血管和淋巴内皮细胞的结合。CLEVER-1是第一种报道介导淋巴结流入和流出的蛋白质。还提供了通过提供CLEVER-1结合抑制剂来治疗炎症和预防恶性细胞转移的方法。

