



US 20090068676A1

(19) **United States**

(12) **Patent Application Publication**
Jackowski et al.

(10) **Pub. No.: US 2009/0068676 A1**
(43) **Pub. Date: Mar. 12, 2009**

(54) **DIAGNOSTIC METHODS FOR CONGESTIVE
HEART FAILURE**

(21) Appl. No.: **11/786,885**

(22) Filed: **Apr. 13, 2007**

Related U.S. Application Data

(76) Inventors: **George Jackowski**, Kettleby (CA);
Tracy Van Lieshout, Hamilton
(CA); **Brad Thatcher**, Casalnuovo
de Napoli (IT); **Rulin Zhang**,
Brampton (CA); **Jason Yantha**,
Toronto (CA); **Michele**
Rasamoelisololo, Winnipeg (CA)

(63) Continuation of application No. 10/706,599, filed on
Nov. 11, 2003, now abandoned.

Publication Classification

(51) **Int. Cl.**
G01N 33/53 (2006.01)

(52) **U.S. Cl.** **435/7.1**

(57) **ABSTRACT**

Correspondence Address:

**MINTZ, LEVIN, COHN, FERRIS, GLOVSKY
AND POPEO, P.C**
5 Palo Alto Square - 6th Floor, 3000 El Camino Real
PALO ALTO, CA 94306-2155 (US)

The invention provides an assay for the quantification of
circulating glycoporphin in biological fluid samples. The cir-
culating glycoporphin measured by this assay is a truncated
glycoporphin diagnostic for congestive heart failure (CHF).




Figure 1




Figure 2





Figure 3




Figure 4




Figure 5A




Figure 5B




Figure 5C




Figure 6

DIAGNOSTIC METHODS FOR CONGESTIVE HEART FAILURE

CROSS REFERENCE TO RELATED APPLICATION

[0001] This application is a continuation of U.S. patent application Ser. No. 10/706,599, filed on Nov. 11, 2003, the contents of which is herein incorporated by reference.

FIELD OF THE INVENTION

[0002] The instant invention relates generally to the field of immunology; particularly to the use of immunologic assays to diagnose abnormal or disease states and most particularly to a sandwich ELISA (enzyme-linked immunosorbent assay) assay for the quantification of a truncated glycoprotein circulating in biological fluid which is diagnostic for congestive heart failure (CHF).

BACKGROUND OF THE INVENTION

[0003] The diagnosis of a given disease requires standard agreed-upon observations usually made by the attending physician of the sick patient. For some diseases, a single test is available which gives nearly definitive results sufficient for a correct diagnosis, for example, the glucose tolerance test for diabetes. However, most diseases require a number of sophisticated tests to arrive at a probable diagnosis. At the present time, therapeutic interventions are frequently initiated at late stages of disease, often resulting in only modest improvements in the quality and length of the affected patients life. Disease prevention is easier and more effective than disease therapy. Earlier diagnosis decreases disease-associated morbidities, increases the quality and length of life of the patient and decreases overall costs of health care. Thus, it is a goal of biomedical researchers to develop diagnostic tests which can correctly diagnose disease at the early stages.

[0004] Early diagnosis of congestive heart failure (CHF) is particularly beneficial since the cardiac re-structuring which occurs with progressive disease may be slowed or prevented with early therapeutic intervention. However, early diagnosis has proven elusive since symptoms generally do not appear until the heart has already suffered structural changes.

[0005] CHF is a serious condition with a high mortality rate affecting approximately five million Americans (see U.S. Pat. No. 6,572,895 for a discussion of CHF). It is currently believed that CHF is not a distinct disease process in itself, but rather represents the effect of multiple abnormalities which interact together to ultimately produce the progressive loss of the ability of the heart to function as a circulatory pump. Major pathophysiologic abnormalities which occur in CHF are activation of the hypothalamic-pituitary-adrenal axis, systemic endothelial dysfunction and myocardial re-structuring. The progression of CHF can be initiated by an event such as myocardial infarction wherein the heart muscle is damaged or it can result from hypertension and/or cardiac malformations. Recently, it has been discovered that patients with certain conditions such as insulin resistance and Type II diabetes have a particularly high risk for heart failure and poor prognosis once they develop CHF (Solång et al. European Heart Journal 20:789-795 1999).

[0006] Disease processes, such as those which occur in diabetes and CHF, often result in cellular and/or tissue damage followed by the release of cellular and/or tissue specific biopolymer markers into the bodily fluids of an individual.

These biopolymer markers are harbingers of disease and/or disease progression. Association of such biopolymer markers with abnormal and/or disease states provides new diagnostic avenues which may allow identification of patients in the early stages of disease or patients at risk for developing disease. Identification of biopolymer markers diagnostic for CHF is particularly advantageous considering the progressive pathophysiology involved in CHF. What is lacking in the art is an efficient, easy to perform diagnostic method capable of identifying an individual suffering from CHF.

SUMMARY OF THE INVENTION

[0007] The instant invention provides an efficient, easy to perform diagnostic method capable of identifying an individual suffering from CHF. The method comprises a sandwich ELISA assay using mouse monoclonal antibodies (anti-glycoproteins) to quantify elevated glycoprotein in biological fluids. Glycoprotein is the major integral membrane protein of the mammalian red blood cell (RBC) and is highly glycosylated. The glycosylation of glycoprotein is responsible for the overall negative charge of the RBC cellular surface leading to the normal electrostatic repulsion among red blood cells. In the disease processes of diabetes and CHF the red blood cell (RBC) membrane proteins, including glycoproteins, are abnormally degraded, thus reducing the overall negative charge of the cellular surface leading to a decrease in the normal electrostatic repulsion among red blood cells. As a consequence, aggregation of red blood cells occurs in the pathogenesis of diabetes and CHF. Using the sandwich ELISA assay of the invention, the instant inventors identified an abnormal, circulating glycoprotein in the plasma of CHF patients. This glycoprotein had a lower molecular weight than that of normal glycoprotein, thus it is predicted to be a truncated fragment which has been cleaved from the RBC membrane surface during the disease process.

[0008] Three mouse monoclonal antibodies are used in the ELISA assay of the instant invention; 3F4, 6G4 and 5F4. Monoclonal antibody 3F4 recognizes amino acid residues 5-25 of SEQ ID NO:2 and SEQ ID NO:4 (glycoproteins A and B). Monoclonal antibody 6G4 recognizes amino acid residues 39-45 of SEQ ID NO:2 (glycoprotein A). Monoclonal antibody 5F4 recognizes amino acid residues 107-119 of SEQ ID NO:2 (glycoprotein A).

[0009] Accordingly, it is an objective of the instant invention to provide a sandwich ELISA assay using mouse anti-glycoprotein monoclonal antibodies 3F4, 6G4 and 5F4 for the quantification of an abnormal, truncated glycoprotein circulating in biological fluid.

[0010] It is another objective of the instant invention to identify a circulating, truncated glycoprotein diagnostic for congestive heart failure (CHF).

[0011] Other objectives and advantages of this invention will become apparent from the following description taken in conjunction with the accompanying drawings wherein are set forth, by way of illustration and example, certain embodiments of this invention. The drawings constitute a part of this specification and include exemplary embodiments of the present invention and illustrate various objects and features thereof.

BRIEF DESCRIPTION OF THE FIGURES

[0012] FIG. 1 shows the data resulting from the sandwich ELISA using monoclonal antibody 3F4.

[0013] FIG. 2 shows the data resulting from the sandwich ELISA using monoclonal antibodies 6G4, 5F4 and 3F4.

[0014] FIG. 3 shows the data resulting from the direct ELISA evaluating the presence of an autoantibody to glycoporphin.

[0015] FIG. 4 shows the results of immunoprecipitation of glycoporphin from the plasma of CHF patients.

[0016] FIGS. 5A-C show chromatograms; FIG. 5A shows captured glycoporphin from CHF patients; FIG. 5B shows captured glycoporphin from healthy patients and FIG. 5C shows captured purified glycoporphin.

[0017] FIG. 6 shows chromatograms after deglycosylation treatment; the top chromatograph shows purified glycoporphin; the middle chromatograph shows captured glycoporphin from CHF patients and the bottom chromatograph is a control run without a glycoporphin sample.

DEFINITIONS

[0018] The following list defines terms, phrases and abbreviations used throughout the instant specification. Although the terms, phrases and abbreviations are listed in the singular tense the definitions are intended to encompass all grammatical forms.

[0019] As used herein, the abbreviation "CHF" refers to congestive heart failure.

[0020] As used herein, the abbreviation "GP" refers to glycoporphin.

[0021] As used herein, the abbreviation "GPA" refers to glycoporphin A.

[0022] As used herein, the abbreviation "GPB" refers to glycoporphin B.

[0023] As used herein, the abbreviation "GPx2" refers to the dimerized form of glycoporphin A.

[0024] As used herein, the abbreviation "GPBx2" refers to the dimerized form of glycoporphin B.

[0025] As used herein, the abbreviation "ELISA" refers to enzyme-linked immunosorbent assay.

[0026] As used herein, the abbreviation "RBC" refers to red blood cell.

[0027] As used herein, the abbreviation "MoAb" refers to monoclonal antibody.

[0028] As used herein, the abbreviation "MS" refers to mass spectrometry.

[0029] As used herein, the abbreviation "SELDI" refers to a mass spectrometric technique; surface enhanced laser desorption ionization.

[0030] As used herein, the abbreviation "PBS" refers to phosphate buffered saline.

[0031] The terms "RBC", "red blood cell" and "erythrocyte" are used interchangeably herein.

[0032] As used herein, the term "glycoporphin" refers to the major integral glycoprotein of the mammalian erythrocyte membrane. Glycoporphin is highly glycosylated and occurs in isoforms A and B (see Concise Encyclopedia: Biochemistry and Molecular Biology, Third Edition, Revised and Expanded by Thomas A. Scott and E. Ian Mercer, Walter de Gruyter, Berlin-New York 1997, pages 201-202 and Instant Notes: BioChemistry, 2nd edition, B. D. Hames and N. M. Hooper, Springer-Verlag New York 2000, pages 125, 126 and 130 for an introduction to the RBC membrane and glycoporphins).

[0033] As used herein, the term "circulating, truncated glycoporphin" refers to the abnormal glycoporphin fragment identified by the assay of the instant invention in the serum of CHF

patients. The 3F4 mouse anti-glycoporphin monoclonal antibody which recognizes the extracellular portion of glycoporphin A and B binds to this circulating, truncated glycoporphin. This circulating, truncated glycoporphin is structurally different from the normal soluble glycoporphin and is theorized to be a fragment cleaved from the RBC surface during disease processes.

[0034] As used herein, the term "biological fluid" refers to any bodily fluid. Illustrative, albeit non-limiting examples are blood, blood products, urine, saliva, cerebrospinal fluid and lymphatic fluid.

[0035] As used herein, the term "subject" refers to any mammalian organism. A particularly preferred subject is a human.

[0036] As used herein, the term "corresponding" is used generally with reference to antibody-antigen binding complexes, for example, an antibody corresponding to an antigen will bind to the antigen under physiologic conditions. The bound antibody-antigen is referred to as an antibody-antigen binding complex.

[0037] As used herein, the term "signal generating substance" refers to any material which undergoes a measurable reaction. Illustrative, albeit non-limiting examples are fluorophores, enzymes and radioisotopes. A particularly preferred signal generating substance is peroxidase, the use of which is illustrated in the examples herein.

[0038] As used herein, the term "congestive heart failure" refers to a progressive, debilitating condition wherein the heart loses its ability to function as a circulatory pump.

[0039] As used herein, the term "antibody" refers to a protein secreted by B lymphocytes capable of binding specific molecules under physiologic conditions.

[0040] As used herein, the term "monoclonal antibody" refers to an antibody having single epitope specificity.

[0041] As used herein, the term "polyclonal antibody" refers to an antibody capable of binding with multiple epitopes.

[0042] As used herein, the term "antigen" broadly refers to any substance which induces an immune reaction; more particularly the term "antigen" refers to the corresponding binding partner of an antibody.

[0043] As used herein, the term "auto-antibody" refers to an antibody which recognizes self antigens, for example, antibodies produced by an organism which bind the organism's own proteins are referred to as auto-antibodies.

[0044] Specific antibodies can be used to quantify the amount of corresponding antigen in a biological sample. As used herein, the term "ELISA" refers to an enzyme-linked immunosorbent assay which can quickly detect and quantify minute amounts (less than a nanogram) of antigen in a biological sample. The test antibody is bound to an inert polymer support, such as a plastic tray with molded wells, and then exposed to the biological sample. Unbound proteins are washed away and a second antibody that reacts with the antigen at a different epitope than the test antibody reacts with is added. This second antibody has an enzyme attached to it that converts a colorless or nonfluorescent substrate into a colored or fluorescent product. The amount of second antibody bound, and hence the amount of protein antigen present in the original biological sample, is determined by the quantification of the intensity of color or fluorescence produced. This ELISA assay is also referred to as an "indirect ELISA" or a "sandwich ELISA". (see Instant Notes: BioChemistry, 2nd edition, B. D. Hames and N. M. Hooper, Springer-Verlag

New York 2000, pages 112-114 for an introduction to the general principles of ELISA assays). There is also a form of ELISA assay that is referred to as "direct" wherein the antigen is bound to an inert polymer support and exposed to a biological sample containing the corresponding antibody.

DETAILED DESCRIPTION OF THE INVENTION

[0045] As a result of disease processes, damage to cells and tissues of the body occurs at the cellular and sub-cellular levels. Initially, these processes may only cause damage to the outer membranes of cells, causing a sloughing off of portions of the exterior cellular matrices, which process is broadly defined as reversible damage. As the length of time and/or the severity of the disease condition increases, the outer membranes begin to break down, resulting in membrane rupture followed by the release of intra-cellular components, which process is broadly defined as irreversible damage. When such damage occurs (reversible or irreversible), biopolymer markers are released into the circulation, causing the immune system to become activated, since these biopolymer markers are not normally present in the bodily fluids. The immune system views these biopolymer markers as invading pathogens or foreign bodies whose threat must be neutralized. In an effort to persevere against this perceived threat, auto-antibodies are formed to these biopolymer markers. These auto-antibodies can be characterized as sequela which are indicative of the original damaging insult to the organism. The presence of the auto-antibodies validates the theory that cellular damage acts as an initiator of an immune response leading to a cascade of auto-antibody production which ultimately manifests itself in a characteristic and often predictable disease state. The appearance of these biopolymer markers and their associated auto-antibodies are harbingers of disease and/or disease progression and are useful for diagnostic purposes.

[0046] Damage to the red blood cell membrane is known to occur in disease processes such as diabetes and CHF. In these diseases there is an increase in enzyme production and/or activation (neutrophil proteases, metalloproteases, sialidases and endopeptidases) that directly and/or indirectly leads to abnormal degradation of red blood cell membrane proteins (Gaczynska et al. *Cytobios* 75:7-11 1993; Venerando et al. *Blood* 99(3):1064-1070 2002; Wegner et al. *Cardiovascular Research* 31:891-898 1996; Piwowar et al. *Clinical Chemistry Lab Medicine* 38(12):1257-1261 2000 and Santos-Silva et al. *Clinica Chimica Acta* 320:29-35 2002).

[0047] Additionally, it is well-documented that erythrocyte (RBC) aggregability is increased in diabetes and in vascular atherosclerotic disease (Caimi et al. *Thromb Haemost* 83:516-517 2000; Demiroglu et al. *Experimental Clinical Endocrinol Diabetes* 107(1):35-39 1999; Martinez et al. *Clinical Hemorheology and Microcirculation* 18:253-258 1998 and Ziegler et al. *Metabolism* 43(9):1182-1186 1994). Alterations in RBC membrane phospholipids are associated with RBC aggregability (Martinez et al. *Clinical Hemorheology and Microcirculation* 18:253-258 1998). Sphingomyelin is the main phospholipid of the outer membrane and has been shown to contain a greater percentage of saturated fatty acids in diabetic patients than in non-diabetic patients. This increase in saturation is thought to reduce electrostatic repulsion between red blood cells, which in turn increases their aggregability.

[0048] Loss of glycoporphins further reduces the electrostatic repulsion of red blood cells. Glycophorin is the major

RBC integral membrane glycoprotein. The high sialylation of glycoporphin is responsible for the negative surface charge which leads to the normal electrostatic repulsion between red blood cells (Eylar et al. *The Journal of Biological Chemistry* 237(6):1992-2000 1962). The increase in enzyme production and/or enzyme activation in disease processes such as diabetes results in the loss of glycoporphins from the RBC membrane. These glycoporphin fragments are released into the bodily fluids where they stimulate the production of auto-antibodies. The decrease in glycoporphin in turn leads to a decrease in the normal negative charge of the RBC membrane surface and thus decreases the overall electrostatic repulsion between blood cells. Loss of the electrostatic repulsion between red blood cells results with the aggregation of red blood cells seen in diabetes.

[0049] Without being bound by any particular theory, the instant inventors propose that the circulating, truncated glycoporphin identified in the plasma of CHF patients using the sandwich ELISA assay described herein is an extracellular glycoporphin fragment which has been cleaved from the RBC membrane during the disease process. This circulating, truncated glycoporphin is structurally different from the normal soluble form of glycoporphin. The mouse anti-glycoporphin 3F4 monoclonal antibody which recognizes amino acid residues 5-25 of SEQ ID NO:2 and SEQ ID NO:4 (glycoporphins A and B) also recognizes the circulating, truncated glycoporphin. The instant inventors have also shown by direct ELISA assay that CHF patients show an increase in anti-glycoporphin auto-antibodies. Thus, it is concluded that this circulating, truncated glycoporphin can be used as a new biopolymer marker for CHF diagnosis.

EXPERIMENTAL PROCEDURES

Sequences

[0050] *Homo sapiens* (human) glycoporphin A nucleic acid sequence is disclosed as SEQ ID NO:1 and translates into glycoporphin A protein disclosed as amino acid sequence SEQ ID NO:2. *Homo sapiens* (human) glycoporphin B nucleic acid sequence is disclosed as SEQ ID NO:3 and translates into glycoporphin B protein disclosed as amino acid sequence SEQ ID NO:4.

Antibodies

[0051] The mouse anti-glycoporphin monoclonal antibodies used in the following experiments were purchased from Bio-Atlantic (Nantes Cedex, France). Monoclonal antibody 6G4 recognizes amino acid residues 39-45 of SEQ ID NO:2 (glycoporphin A). Monoclonal antibody 5F4 recognizes the intracellular portion of glycoporphin A comprising amino acid residues 107-119 of SEQ ID NO:2. Monoclonal antibody 3F4 recognizes the extracellular portion of glycoporphins A and B amino acid residues 5-25 of SEQ ID NO:2 and SEQ ID NO:4. The binding of the 3F4 antibody to its epitope is sugar-dependent whereas the binding of the 6G4 antibody is not. These monoclonal antibodies are described in detail in Rasamoeliso et al. *Vox Sanguinis* 72:185-191 1997.

[0052] The mouse anti-glycoporphin 3F4 monoclonal antibody was deposited with the American Type Culture Collection (ATCC) on Apr. 23, 2000 as hybridoma NaM26-3F4D11A2 under Accession number PTA-5154. The American Type Culture Collection (ATCC) is located at 10801 University Boulevard, Manassas, Va. 20110-2209. Applicants submit that all restrictions on the availability to

the public of this deposited material will be irrevocably removed upon granting of a patent in the United States.

Quantification of Glycophorin by Sandwich ELISA

[0053] One μg of each MoAb in 100 μl of 50 mM carbonate pH 9.4 was coated on ELISA plates (Nuc, Denmark) and set overnight at +4° C. Plates were then washed 3 times with 0.01M phosphate buffer 150 mM NaCl pH 7.4 (PBS) purchased from Sigma containing 0.05% Tween 20 (PBST). Plates were then blocked with 200 μl of PBST containing 0.5% BSA (Sigma) for 30 minutes at 37° C. 100 μl of CHF patient plasma (PRAISE 2 study) and healthy control plasma (Intergen) diluted $\frac{1}{10}$ in PBST were then added per well in duplicate and incubated for 2 hours at room temperature. After 3 washes with PBST, 100 μl of rabbit polyclonal anti-glycophorin A+B (BioAtlantic) were added and incubated for 1 hour at room temperature followed by the addition of 100 μl of peroxidase labeled donkey polyclonal anti-rabbit IgG (H+L) diluted $\frac{1}{50,000}$ in PBST containing 0.5% BSA (Jackson ImmunoResearch). The presence of the captured glycophorins is detected by adding 100 μl of TMB (Moss, Inc.). The reaction was stopped with 50 μl of 1N H₂SO₄. Plates were then read at 450 nm on the BioRad microplate reader.

[0054] FIG. 1 shows the result of the sandwich ELISA using the 3F4 monoclonal antibody. The absorbance at 450 nm is shown on the Y axis. Glycophorin captured from the plasma of CHF patients is shown on the left and the glycophorin captured from normal plasma (control, n=36) is shown on the right. The signal is significantly higher in CHF plasma than in controls (p<0.001) calculated by an independent t-test indicating a higher amount of glycophorins in CHF plasma samples. The 3F4 MoAb recognizes the common sequence on both glycophorins A and B (amino acid residues 5-25 of SEQ ID NO:2 and SEQ ID NO:4). This binding is sugar-dependent since this fragment of glycophorin is highly glycosylated.

[0055] In order to ascertain whether the assay is specific to the extracellular polypeptide of glycophorin or the oligosaccharide chains, the MoAbs 6G4 (recognizes amino acid residues 39-45 of SEQ ID NO:2) and 5F4 (recognizes amino acid residues 112-129 of SEQ ID NO:2) were used. Both bind to the glycophorin A backbone independently of the sugar chains.

[0056] Eight CHF samples having the most elevated amount of glycophorin and 8 normal plasma samples having the lowest amount of glycophorin were analyzed and the result is shown in FIG. 2. FIG. 2 shows results from sandwich ELISA assays comparing the glycophorin captured in plasma from CHF patients and the glycophorin captured in normal control plasma (n=8). The top panel shows results using the 6G4 MoAb (p=0.001); the middle panel shows results using the 5F4 MoAb (p=0.36) and the bottom panel shows the results using the 3F4 MoAb (p=0.003). The Y axis represents the absorbance read at 450 nm. Glycophorin captured from the plasma of CHF patients is shown on the left and the glycophorin captured from normal plasma is shown on the right in all three panels. The result shows that 6G4 detects elevated amount of glycophorin in CHF samples, while 5F4 shows no significant difference between both CHF and normal human plasma. This result indicates that glycophorin may be cleaved from the red blood cell membrane during the progression of CHF since the fragments recognized by the antibodies are extracellular fragments. However, it is noted that a soluble form of glycophorin is present in normal as well

as CHF patient plasma that is detected by the 5F4 monoclonal anti-intracellular domain of glycophorin.

Detection of Auto-Antibody by Direct ELISA

[0057] 0.5 μg of purified glycophorin from blood group MM or asialoglycophorins from blood group MN (Sigma) in 50 mM carbonate buffer pH 9.4 was adsorbed onto ELISA plates overnight at +4° C. Plates washed 3 times with 0.01M Phosphate buffer 150 mM NaCl pH 7.4 (PBS) from Sigma containing 0.05% Tween 20 (PBST). Plates were then blocked with 200 μl of PBST containing 0.5% BSA (Sigma) for 30 minutes at 37° C. 100 μl of CHF plasma (PRAISE 2 study) and normal control plasma (Intergen) diluted $\frac{1}{100}$ in PBST were then added per well in duplicate and incubated for 2 hours at room temperature. After 3 washes with PBST, 100 μl of peroxidase labeled goat polyclonal anti-human IgG (H+L) diluted $\frac{1}{10,000}$ in PBST (Jackson ImmunoResearch) were added. The presence of auto-antibody anti-glycophorins was detected by adding 100 μl of TMB (Moss, Inc.) and the reaction was stopped with 50 μl of 1N H₂SO₄. Plates were read at 450 nm on the BioRad microplate reader.

[0058] Glycophorin is known to be highly immunogenic due to the presence of a high amount of sugar chains. Once found in plasma it may induce an immune response generating anti-glycophorin auto-antibody.

[0059] To demonstrate the presence of CHF-induced auto-antibody against glycophorin, glycophorins from blood group MM and asialo glycophorins from blood group MN were coated on ELISA plates and plasma from healthy donors or from CHF patients were added. FIG. 3 shows the results of the direct ELISA assay evaluating the presence of a CHF-induced auto-antibody in the plasma of normal and CHF patients (n=36). In the top panel, glycophorin from blood group MN was coated on the plate (p=0.01) and the bottom panel, desialylated glycophorin from blood group MN was coated on the plate (p=0.03). The Y axis represents the absorbance read at 450 nm. FIG. 3 shows the presence of auto-antibodies in CHF; independent to the blood group (M or N) and the heavy sialic acids on glycophorin.

Identification of Glycophorins in CHF Plasma by Immunoprecipitation and Detection by Immunoblotting

[0060] 1.2 ml of pooled CHF plasma from the PRAISE 2 study was diluted v/v with PBS containing 0.5% Triton X-100. Then 2 μl of 3F4 MoAb at 1.7 mg/ml were added. After overnight incubation at +4° C., 25 μl of goat IgG anti-mouse IgG (H+L) coupled to SEPHAROSE-4B beads (Zymed) were added. The mixture was incubated for 5 hours at +4° C. and then the beads were washed 3 times with PBS containing 0.05% Tween 20. The captured (glyco)protein was eluted with 100 μl of 0.1M glycine pH 2.5 then neutralized with 1M Tris pH 11. The eluate was concentrated on CentriVap Concentrator (Labconco), resuspended in 50 μl of SDS-PAGE sample buffer, boiled 5 minutes at 100° C. and then loaded on 10% SDS-PAGE gel. At the end of the electrophoresis, proteins were transferred onto a nitrocellulose membrane and stained with 3F4 MoAb anti-GPA+B followed by a peroxidase labeled goat polyclonal anti-mouse IgG (H+L) diluted $\frac{1}{50,000}$ in PBST (Jackson ImmunoResearch). The immunoblot was then developed using ECL (Amersham Pharmacia). To control the cross-reactivity of the secondary antibody to the 3F4 eluted from the column, the blot was incubated with the secondary antibody alone.

[0061] The molecules captured by 3F4-column were eluted and loaded on 10% SDS-PAGE gel and assessed on immunoblotting against the same MoAb. As shown in FIG. 4, the glycoporphins found in CHF plasma have a molecular weight of 75, 45 and 40 kDa (lane 2, blot incubated with 3F4). Usually glycoporphins run at 80-70-40-37 and 20 kDa as dimer form of GPA, dimer GPA/GPB, dimer form of GPB, monomer form of GPA and monomer form of GPB, respectively as shown on lane 1 loaded with normal glycoporphin purified from normal red blood cell membrane. Thus, the glycoporphins found in the plasma of CHF patients have different molecular weights as compared to the normal glycoporphin purified from RBC membranes. The immunoblot was incubated with the secondary antibody alone (control) or with the 3F4 antibody and then the secondary antibody. Lane 1 (in both blots) shows glycoporphin purified from RBC membranes and Lane 2 (both blots) shows glycoporphin from CHF patient plasma. Protein markers from 25 to 200 kDaltons are shown on the far left.

[0062] The IgG identified in control and 3F4 blots is the mouse monoclonal 3F4 used for the immunoprecipitation and released from the column. A band with a high MW>200 kDa is also detected. The instant inventors are not sure about the nature of this band. The band may be a complex form of IgM or IgG autoantibodies and the glycoporphins.

Identification of Glycoporphin in CHF Patient Sample by SELDI-TOF

[0063] The method of the instant invention can be carried out using the techniques of mass spectrometry. The PS20 chip (Ciphergen) was washed with pure Acetonitrile-190 (ACN) (Calcdon) and allowed to air dry. 50 µg of Protein G (Pierce) was dissolved in 50 µl UF water and 1 ul was loaded to each spot containing 1 µl of ACN. The mixture was incubated 1 hour in a humidity chamber and then the spot was blocked with 10 µl of 0.5M Tris-HCl pH 7.4 (Calcdon) for 15 minutes. The chip was then washed with UF water and allowed to air dry. Monoclonal antibody (MoAb) anti-GPA+GPB, the 3F4 at 1.7 mg/ml (BioAtlantic) was diluted 1/3 in PBS containing 0.1% TRITON X (Sigma) and 3 µl of the MoAb solution was loaded per spot and incubated for 1 hour in a humidity chamber. Unbound MoAb was washed away from the chip by washing with PBS.

[0064] Purified glycoporphin (Sigma), CHF plasma from PRAISE 2 study or normal plasma (Intergen) was added to the 3F4-coated chip as follows:

[0065] The glycoporphin at 1 mg/ml was diluted 1/3 in PBS; CHF and normal plasma samples were diluted 1/3 in PBS containing 0.05% Tween 20, and 2 µl of each were loaded per spot. The chip was then incubated for 1 hour in a humidity chamber and washed twice with UF water.

[0066] The captured glycoporphin was then treated with Endoproteinase GluC (Roche Diagnostics). For that, the GluC powder was dissolved in 50 µl of UF water and a 1/10 dilution in 50 mM Ammonium Carbonate pH 7.8 (BDH Laboratory Supplies) was prepared. 1 µl of the GluC solution was added to each spot and incubated 2 hours in a humidity chamber. The spot was then allowed to dry and was either treated using Calbiochem deglycosylation kit or directly analyzed on SELDI after adding 1 ul of saturated sinapinic acid (Sigma) in 0.5% TFA 50% ACN. The chip was then read on SELDI (Ciphergen) at a Sensitivity=10, Intensity=180-190, range of 0-5000 Da (optimized for 0-5000 Da).

[0067] The (glyco)protein captured on the 3F4 chip was treated by GluC. FIG. 5A shows data resulting from the on-chip treatment of the captured glycoporphin from CHF. FIG. 5B shows data resulting from the on-chip treatment of the normal plasma samples. FIG. 5C shows data resulting from the on-chip treatment of purified glycoporphin. As shown in FIGS. 5A-C, a (glyco)peptide with a m/z of 2361+H is found in both CHF and glycoporphin demonstrating that the (glyco)protein captured from CHF corresponds probably to the glycoporphin. It is interesting to note that the chromatograms (FIGS. 5A-C) obtained from the purified glycoporphin and the one from CHF plasma were not overlapped. This is due to the fact that the structure of the glycoporphin in CHF is maybe slightly modified.

[0068] To further prove that the captured (glyco)protein is related to glycoporphin, the captured (glyco)protein was deglycosylated on chip. FIG. 6 shows on-chip deglycosylation treatment of the glycopeptides captured from either purified glycoporphin or CHF plasma using the 3F4 monoclonal antibody coated on a PS20 chip. As shown in FIG. 6, at least 8 major peaks now matched to the peaks generated from the standard glycoporphin. Also, it is noted that a lot more peaks were detected, they correspond not only to the peptides but also to the sugar chains released after the deglycosylation treatment. In conclusion, the instant invention provides a sandwich ELISA assay for quantification of a truncated, glycoporphin circulating in biological fluid which is diagnostic for CHF. It is important to note that glycoporphin has not been previously recognized as a marker for congestive heart failure (CHF). The instant inventors are the first to document glycoporphin as a marker for CHF and the assay described herein provides an efficient, easy to perform diagnostic method capable of identifying an individual suffering from CHF.

[0069] All patents and publications mentioned in this specification are indicative of the levels of those skilled in the art to which the instant invention pertains. All patents and publications are herein incorporated by reference to the same extent as if each individual patent and publication was specifically and individually indicated to be incorporated by reference.

[0070] It is to be understood that while a certain form of the invention is illustrated, it is not to be limited to the specific form or arrangement of parts herein described and shown. It will be apparent to those skilled in the art that various changes may be made without departing from the scope of the invention and the invention is not to be considered limited to what is shown and described in the specification.

[0071] One skilled in the art will readily appreciate that the present invention is well adapted to carry out the objects and obtain the ends and advantages mentioned, as well as those inherent therein. The oligonucleotides, peptides, polypeptides, biologically related compounds, methods, procedures and techniques described herein are presently representative of the preferred embodiments, are intended to be exemplary and are not intended as limitations on the scope. Changes therein and other uses will occur to those skilled in the art which are encompassed within the spirit of the invention and are defined by the scope of the appended claims. Although the invention has been described in connection with specific preferred embodiments, it should be understood that the invention as claimed should not be unduly limited to such specific embodiments. Indeed various modifications of the described modes for carrying out the invention which are obvious to those skilled in the art are intended to be within the scope of the following claims.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 4

<210> SEQ ID NO 1

<211> LENGTH: 2593

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 1

agttgtcttt ggtagttttt tgcactaact tcaggaacca gctcatgac tcaggatgta 60
tggaaaaata atctttgtat tactattgtc agcaattgtg agcatatcag catcaagtac 120
cactgggtgtg gcaatgcaca ctccaacctc ttcttcagtc acaaaagatt acatctcatc 180
acagacaaat gatcgcaca aacgggacac atatgcagcc actcctagag ctcatgaagt 240
ttcagaaatt tctgttagaa ctgtttacc tccagaagag gaaaccggag aaagggtaca 300
acttgcccat catttctctg aaccagagat aacctcatt atttttgggg tgatggctgg 360
tgttattgga acgatcctct taattttcta cggatttcgc cgactgataa agaaaagccc 420
atctgatgta aaacctctcc cctcacctga cacagacgtg cctttaagtt ctgttgaaat 480
agaaaatcca gagacaagtg atcaatgaga atctgttcac caaaccaaat gtggaagaa 540
caciaaagag acataagact tcagtcaagt gaaaaattaa catgtggact ggacactcca 600
ataaattata tactgccta agttgtacaa tttcagaatg caattttcat tataatgagt 660
tccagtgact caatgatggg gaaaaaatc tctgctcatt aatatttcaa gataaagaac 720
aaatgtttcc ttgaatgctt gcttttgtgt gttagcataa tttttagaat tgtttgagaa 780
ttctgatcca aaactttagt tgaattcacc tacgtttggt taatattaac ttaacctatt 840
ctattgtatt ataatgatg ttctgtcaaa tgaaggctt gaaataccta gatgaagttt 900
agattttctt cctattgtaa acttttgagt ctggtttcat tgttttaaat aaattaaggg 960
gacactaaag tccatcatt cattccttca ttctgaacag gcaagatata atattacatg 1020
aatgattact atattttgtt cacactaata aagcttatgc tcagaaatgc catacacaca 1080
caciaaacaca cacatttacc atttaatgca taaatcaaca caaaaggttt tcccattaat 1140
atgaaatatt acatatata agtgccata tttaaaataa tttgtctaac agtagaacta 1200
tgtcggagca ctcactgaag cttcgatttc cactgaaag agttatttgt tgtaagtaga 1260
gttatcccg agaaggaaa agaacttacg acctttcttt ataacagaaa gctcaactct 1320
aaattcaaca agatgtgcaa accggacatg cagggaata ttttaatagg ttactataag 1380
gttctcaatt aaattcttta atctgtccag tccagtttc tcttattaat aaaactttgg 1440
aaattgcttt aaaccattta aaggaaattt ctagatatag aaactaaagg actgtgacta 1500
tacagtgtca ctcatttcta gtaaaactta aaaagcaaaa acaaaaaaca aaaaagacct 1560
tctctgtgata ctttatttcc gaactaataa aaatctatat gactttttat tattgtgtga 1620
taaccaagta aatgttttct attttcgata ttttcaggca tggtaacaga aatttacctt 1680
ttaataaatt aaaaaatcta aattttaacc tacttgtatg ttcggagagt gttttgtac 1740
tatattgact acttaaaata gagaatgaga ctaagaaggg aacatttctg ttgatacatg 1800
tttttataaa gtaattttta agagcattat taggttaatt taatccaatt aatgacccaa 1860
atgccaaagt aattttaaat ttacattttt aataaaagca acatgttgaa acaagagagg 1920

-continued

```

gtgagattaa ccttttct aaagtaattt acaagtcaaa gacaggaaga gatcagagtg 1980
aatgtgcctt cctaaccaga gctacagaat ttagtgaata attaaagtac aaactgcttt 2040
gacctccttg aacttttcca agcaatttct ctgtacttct atatatgaat gtcttagcca 2100
atthtctgct actataacag aatacgacag actgggtaat ttaaaaagaa aagaaattta 2160
ttttcttct agttctggag gctgggaagg cgaagggcat ggcaactgaca tctgccttgt 2220
aactgatgag aactttctta ctgcatgata acaaagcagc aaggcaagca aaagcgtaag 2280
atgaagagag aggaaatgaa gccaaacaca tcctttcatc agaagcccat tcctctata 2340
aggcgttact acatttatga gaatggagtc ctcacacact aatcgtgacc ttaaaggccc 2400
ctcccaacac tgttacaatg gcaattaaat ttcaacaaag gttccagagg tgacattcga 2460
atcagcaatg aaattttcat agttaaattt ggtattcgtg ggggaagaaa tgaccatttc 2520
ccttgatttt ttataattaa atcagcaaaa tattgtaata aagaatcttt tcctgtgaag 2580
ataccatgac ccc 2593

```

```

<210> SEQ ID NO 2
<211> LENGTH: 150
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 2

```

```

Met Tyr Gly Lys Ile Ile Phe Val Leu Leu Leu Ser Ala Ile Val Ser
1           5           10           15
Ile Ser Ala Ser Ser Thr Thr Gly Val Ala Met His Thr Ser Thr Ser
20           25           30
Ser Ser Val Thr Lys Ser Tyr Ile Ser Ser Gln Thr Asn Asp Thr His
35           40           45
Lys Arg Asp Thr Tyr Ala Ala Thr Pro Arg Ala His Glu Val Ser Glu
50           55           60
Ile Ser Val Arg Thr Val Tyr Pro Pro Glu Glu Glu Thr Gly Glu Arg
65           70           75           80
Val Gln Leu Ala His His Phe Ser Glu Pro Glu Ile Thr Leu Ile Ile
85           90           95
Phe Gly Val Met Ala Gly Val Ile Gly Thr Ile Leu Leu Ile Ser Tyr
100          105          110
Gly Ile Arg Arg Leu Ile Lys Lys Ser Pro Ser Asp Val Lys Pro Leu
115          120          125
Pro Ser Pro Asp Thr Asp Val Pro Leu Ser Ser Val Glu Ile Glu Asn
130          135          140
Pro Glu Thr Ser Asp Gln
145          150

```

```

<210> SEQ ID NO 3
<211> LENGTH: 526
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 3

```

```

agttgtcttt ggtagtcttt ttgcactaac ttcaggaacc agtcatgat ctcaggatgt 60
atggaaaaat aatctttgta ttactattgt cagaaattgt gagcatatca gcattaagta 120
ccactgaggt ggcaatgcac acttcaacct cttcttcagt cacaagagt tacatctcat 180

```

-continued

```

cacagacaaa tggagaaacg ggacaacttg tccatcgttt cactgtacca gctcctgtag    240
tgataaact cattaatttg tgtgtgatgg ctggtattat tggaacgata ctcttaattt    300
cttacactat tcgccgactg ataaaggcat gaggatgtgg cctgcatgct gctgtattg    360
cctgcatgct gctgtattt gcgataaacc getgcacctg ctgttctctt ctttatgcaa    420
actggctgca cctgctattc ctttgettat gccctaccc ctggctatcc taattccctg    480
ttctcctgcc tcaactattac tgtattctct acttctaaat aaaaaa                    526
    
```

```

<210> SEQ ID NO 4
<211> LENGTH: 91
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
    
```

<400> SEQUENCE: 4

```

Met Tyr Gly Lys Ile Ile Phe Val Leu Leu Leu Ser Glu Ile Val Ser
1           5           10           15
Ile Ser Ala Leu Ser Thr Thr Glu Val Ala Met His Thr Ser Thr Ser
                20           25           30
Ser Ser Val Thr Lys Ser Tyr Ile Ser Ser Gln Thr Asn Gly Glu Thr
            35           40           45
Gly Gln Leu Val His Arg Phe Thr Val Pro Ala Pro Val Val Ile Ile
50           55           60
Leu Ile Ile Leu Cys Val Met Ala Gly Ile Ile Gly Thr Ile Leu Leu
65           70           75           80
Ile Ser Tyr Thr Ile Arg Arg Leu Ile Lys Ala
            85           90
    
```

What is claimed is:

1. A method useful in determining whether a subject is suffering from congestive heart failure comprising:
 - analyzing a body fluid of the subject to detect a presence and concentration level of unbound and freely circulating glycoporphin or glycoporphin derived fragments in said body fluid;
 - wherein information obtained from said analyses is employed to determine whether the subject is suffering from congestive heart failure.
2. The method of claim 1, wherein the body fluid is blood plasma.
3. The method of claim 2, wherein the blood plasma is substantially free of red blood cells.

4. The method of claim 1, wherein the glycoporphin comprises a circulating, truncated glycoporphin.
5. A diagnostic kit useful in diagnosing congestive heart failure comprising a solid support having immobilized thereon at least one antibody that is specific for a freely circulating glycoporphin characterized as circulating in bodily fluid free from the cellular surface of a red blood cell.
6. The diagnostic kit of claim 5, wherein the bodily fluid is blood plasma.
7. The diagnostic kit of claim 6, wherein the blood plasma is substantially free of red blood cells.
8. The diagnostic kit of claim 5, wherein the glycoporphin comprises a circulating, truncated glycoporphin.

* * * * *

专利名称(译)	充血性心力衰竭的诊断方法		
公开(公告)号	US20090068676A1	公开(公告)日	2009-03-12
申请号	US11/786885	申请日	2007-04-13
[标]申请(专利权)人(译)	JACKOWSKI GEORGE VAN LIESHOUT TRACY 撒切尔BRAD 张汝林 YANTHA JASON RASAMOELISOLO MICHELE		
申请(专利权)人(译)	JACKOWSKI GEORGE VAN LIESHOUT TRACY 撒切尔BRAD 张儒林 YANTHA JASON RASAMOELISOLO MICHELE		
当前申请(专利权)人(译)	NEXUS DX , INC.		
[标]发明人	JACKOWSKI GEORGE VAN LIESHOUT TRACY THATCHER BRAD ZHANG RULIN YANTHA JASON RASAMOELISOLO MICHELE		
发明人	JACKOWSKI, GEORGE VAN LIESHOUT, TRACY THATCHER, BRAD ZHANG, RULIN YANTHA, JASON RASAMOELISOLO, MICHELE		
IPC分类号	G01N33/53 G01N33/543 G01N33/577 G01N33/68		
CPC分类号	G01N2800/325 G01N33/6893		
外部链接	Espacenet USPTO		

摘要(译)

本发明提供了用于定量生物流体样品中循环血型糖蛋白的测定法。通过该测定法测量的循环血型糖蛋白是截短的血型糖蛋白，其诊断为充血性心力衰竭 (CHF)。




Figure 1