



US 20090311726A1

(19) **United States**

(12) **Patent Application Publication**
Markowitz

(10) **Pub. No.: US 2009/0311726 A1**
(43) **Pub. Date: Dec. 17, 2009**

(54) **METHODS AND COMPOSITIONS FOR CATEGORIZING PATIENTS**

(76) Inventor: **Sanford D. Markowitz**, Pepper Pike, OH (US)

Correspondence Address:
ROPES & GRAY LLP
PATENT DOCKETING 39/41, ONE INTERNATIONAL PLACE
BOSTON, MA 02110-2624 (US)

(21) Appl. No.: **12/386,176**

(22) Filed: **Apr. 13, 2009**

Related U.S. Application Data

(63) Continuation of application No. 10/649,591, filed on Aug. 26, 2003, now abandoned, which is a continua-

tion-in-part of application No. 10/274,177, filed on Oct. 18, 2002, now Pat. No. 7,118,912, which is a continuation-in-part of application No. 10/229,345, filed on Aug. 26, 2002, now Pat. No. 7,081,516.

Publication Classification

(51) **Int. Cl.**
G01N 33/53 (2006.01)
(52) **U.S. Cl.** **435/7.92; 435/7.9; 435/7.1**

(57) **ABSTRACT**

The disclosure provides, among other things, molecular markers for categorizing the neoplastic state of a patient, methods for using the molecular markers in diagnostic tests, nucleic acid and amino acid sequences related to the molecular markers, reagents for detection of molecular markers, and methods for identifying candidate molecular markers in highly parallel gene expression data.

Figure 1A. Amino acid sequence of secreted ColoUp1 protein
(1) (SEQ ID NO: 1)

TVAAGCPDQSPPELQPWNPBGHDQDHHVHIGQGKTLTSSATVYSIHISEGGKLVIKDHD
EPIVLRTRHILIDNGGELHAGSALCPFQGNFTIILYGRADEGIQPDPPYGLKYIGVGGK
GALELHGQKLSWTFLNKTLHPGGMAEGGYFFERSWGHARGVIVHVIDPKSGTVIHSDF
DTYRSKESERLVQYLNAPDGRILSVAVNDEGSRNLDDMARKAMTKLGSKHFLHLGFR
HPWSFLTVMGNPSSSVEDHIEYHGHRGSAARVFKLFQTEHGEYFNVLSSEWVQDVEW
TEWFDHDKVSQTKGGEKISDLWKAHPGKICNRPIDIQATTMDGVNLSTEVVYKKGQDYR
FACYDRGRACRSYRVRFLCGKPVPRKLTVTIDTNVNSTILNLEDNVQSWKPGDTLVIAS
TDYSMYQAEFQVLPCRSCAPNQVKVAGKPMYLIHIGEEIDGVDMRAEVGLLSRNIIVMG
EMEDKCYPYRNHICNFFDFDTFGGHIKFALGFKAAHLEGTELKHMGGQLVGGQYPIHFHL
AGVDDERGGYDPPTYIRDLSIHHTFSRCVTVHGSNGLLIKDVVGYNSLGHCFFTEDGPE
ERNTFDHCLGLLVKSGTLLPSDRDSKMCKMITEDSYPGYIPKPRQDCNAVSTFWMANPN
NNLINCAAAGSEETGFWFI FHHVPTGPSVGMYS PGYSEHIPLGKFYNNRAHSNYRAGMI
IDNGVKTTEASAKDKRPFLSII SARYSPHQDADPLKPREPAIIRHFIAYKNQDHGAWLR
GGDVWLDSCRFA DNGIGLTLASGGTFPYDDGSKQEIKNLSLVGSGNVGTEMMDNRIWG
PGGLDHSGRTLPIGQNFPIRGIQLYDGPINIQNCTFRKFVALEGRHTSALAFRLNNAWQ
SCPHNNVTGIAFEDVPITSRVFFGEPGPWFNQLDMDGDKTSVFHDVDGVSVEYPGSYLT
KNDNWLVRHPDCINVPDWRGAI CSGCYAQMYIQAYKTSNLRMKI IKNDFPSHPLYLEGA
LTRSTHYQQYQPVVTLQKGYTIHWDQTAPAELAIWLINFNKGDWIRVGLCYPRGTTFSI
LSDVHNRLKQTSKTGVFVRTLQMDKVEQSYPGRSHYYWDEDSGLLFLKKAQNEREKF
AFCSMKGCERIKIKALI PKNAGVSDCTATAYPKFTERAVVDVPMPPKLFQSGLKTKDHF
LEVKMESKQHFFHLW NDFAYIEVDGKKYPSSE DGIQVVVIDGNQGRVVSHTSFRNSIL
QGI PWQLFNIVATI PDNSI VLMASKGRYVSRGPWTRVLEKLGADRGLKLEQMAFVGFK
GSFRPIWVTLDTEDHKAKI FQVVPI PVVKKKKL

Figure 1B. Amino acid sequence of secreted ColoUp1 protein (II) (SEQ ID NO: 2)

AGCPDQSPQLPWNPGHDQDHHVHIGQGKTLTLLTSSATVYSIHI SEGGKLVIKDHDEPI
VLRRRHILIDNGGELHAGSALCPFQGNFTI ILYGRADEGIQPDPPYGLKYIGVGKGGAL
ELHGQKKLSWTFNLKTLHPGMAEGGYFFERSWGHRGVI VHVIDPKSGTVIHSRDFD
RSKKESERLVQYLNAPDGRILSVAVNDEGSRNLDDMARKAMTKLGSKHFLHLGFRHPW
SFLTVKGNPSSSVEDHIEYHGHRGSAARVFKLFQTEHGEYFNVLSSEWVQDVEWTEW
FDHDKVSQTKGGEKISDLWKAHPGKICNRPIDIQATTMDGVNLSSTEVVYKKGQDYRFAC
YDRGRACRSYRVRFLCGKPVPRKLTVTIDTNVNSTILNLEDNVQSWKPGDTLVIASTDY
SMYQAEFFQVLPSCRSCAPNQVKVAGKPMYLIHIGEEIDGVMRAEVGLLSRNIIVMGEME
DKCYPYRNHICNFFDFDTFGGHIK FALGFKAHLEGTELKHMGGQLVGGYPIHFHLAGD
VDERGGYDPPTYIRDLSIHHTFSRCVTVHGSNGLLIKDVVGYNSLGHCFFTEGPEERN
TFDHCLGLLVKSGLTLLPSDRDSKMCKMITEDSYPGYI PKPRQDCNAVSTFWMANPNNNL
INCAAAGSEETGFWFI FHHVPTGPSVGMSPGYSEHI PLGKFYNNRAHSNYRAGMI IDN
GVKTTTEASAKDKRPFLSII SARYSPHQDADPLKPREPAIIRHF IAYKNQDHGAWLRGD
VWLDSCR FADNGIGLTLASGGTFPYDDGSKQEIKNLFLVGE SGNVGTMMDNRIWGP
LDHSGRTLPIGQNFPIRGIQLYDGPINIQNCTFRKFVALEGRHTSALAFRLNNAWQSCP
HNNVTGIAFEDVPITSRVFFGEPGPWFNQLDMDGDKTSVFHDVDGSVSEYPGSYLT
KNLWLRHPDCINVPDWRGAI CSGCYAQMYIQAYKTSNLRMKI IKNDFPSHPLYLEGALTR
STHYQQYQPVVTLQKGYTIHWDQTAPAELAIWLINFNKGDWIRVGLCYPRGTTFSILSD
VHNRLKQTSKTGVFVRTLQMDKVEQSYPGRSHYYWDEDSGLLFLKKAQNEREKFAFC
SMKGCERIKIKALIPKNAGVSDCTATAYPKFTERAVVDVPMPPKLFGLKTKDHFLEV
KMESSKQHFHLWDFAYIEVDGKKYPSSEDDGIQVVVIDGNQGRVVSHTSFRNSILQGI
PWQLFNYVATI PDNSIVLMASKGRYVSRGPWTRVLEKLGADRGLKKEQMAFVGFKGSF
RPIWVTLDTEDHKAKIFQVVPIPVVKKKKL

Figure 2. Amino acid sequence of secreted ColoUp2 protein
(SEQ ID NO: 3)

LQEVHVSKETIGKISAASKMMWCSAAVDIMFLLDGSNSVGKGSFERSKHFAITVCDGLD
ISPERRVVGAFQFSSTPHLEFPLDSFSTQQEVKARIKRMVFKGGRTETELALKYLLHRG
LPGGRNASVPQILIIIVTDGKSQGDVALPSKQLKERGVTVFAVGVRFPWEELHALASEP
RGQHVLLAEQVEDATNGLFSTLSSSAICSSATPDCRVEAHPCEHRTLEMVREFAGNAPC
WRGSRRTLAVLAAHCPFYSWKRVFLTHPATCYRTTCPGPCDSQPCQNGGTCVPEGLDGY
QCLCPLAFGGEANCALKLSLECRVDLLFLLDSSAGTTLDGFLRAKVFVKRFVRAVLSED
SRARVGVATYSRELLVAVPVGEYQDVPDLVWSLDGIPFRGGPTLTGSALRQAAERGFSG
ATRTGQDRPRRVVLLTESHSEDEVAGPARHARARELLLLGVGSEAVRAELEEEITGSPK
HVMVYSDPQDLFNQIPELQKLCRQRPGCRTQALDLVFMLDTSASVGPENFAQMOSFV
RSCALQFEVNPDVTQVGLVVYGSQVQTAFLDTPKTRAAMLRAISQAPYLGGVGSAGTA
LLHIYDKVMTVQRGARPGVPAVVVLTGGRGAEDAAVPAQKLRNNGISVLVVGVPVLS
EGLRRLAGPRDSLIVHVAAYADLRYHQDVLI EWLCGEAKQPVNLCCKPSPCMNEGSCVLQN
GSYRCKCRDGEWEGPHCENRFLRRP

Figure 3. Nucleic acid sequence of ColoUp1 (SEQ ID NO: 4)

CGTGACACTGTCTCGGCTACAGACCCAGAGGGAGCACACTGCCAGGATGGGAGCTGCTG
GGAGGCAGGACTTCTCTTCAAGGCCATGCTGACCATCAGCTGGCTCACTCTGACCTGC
TTCCCTGGGGCCACATCCACAGTGGCTGCTGGGTGCCCTGACCAGAGCCCTGAGTTGCA
ACCCTGGAACCTTGGCCATGACCAAGACCACCATGTGCATATCGGCCAGGGCAAGACAC
TGCTGCTCACCTCTTCTGCCACGGTCTATTCCATCCACATCTCAGAGGGAGGCAAGCTG
GTCATTAAGACCCAGACGAGCCGATTGTTTTGCGAACCCGGCACATCCTGATTGACAA
CGGAGGAGAGCTGCATGCTGGGAGTGCCCTCTGCCCTTCCAGGGCAATTTACCATCA
TTTTGTATGGAAGGGCTGATGAAGGTATTAGCCGGATCCTTACTATGGTCTGAAGTAC
ATTGGGGTTGGTAAAGGAGGCGCTCTTGAGTTGCATGGACAGAAAAGCTCTCCTGGAC
ATTTCTGAACAAGACCCTTACCCAGGTGGCATGGCAGAAGGAGGCTATTTTTTTGAAA
GGAGCTGGGGCCACCGTGGAGTTATTGTTTCATGTCATCGACCCAAATCAGGCACAGTC
ATCCATTCTGACCGTTTTGACACCTATAGATCCAAGAAAAGAGAGTGAACGTCTGGTCCA
GTATTTGAACGCGGTGCCCGATGGCAGGATCCTTTCTGTTGCAGTGAATGATGAAGGTT
CTCGAAATCTGGATGACATGGCCAGGAAGGCGATGACCAAATGGGAAGCAAACACTTC
CTGCACCTTGGATTTAGACACCCTTGGAGTTTTCTAACTGTGAAAGGAAATCCATCATC
TTCAGTGGAAGACCATATTGAATATCATGGACATCGAGGCTCTGCTGCTGCCCGGGTAT
TCAAATGTTCCAGACAGAGCATGGCGAATATTTCAATGTTTCTTTGTCCAGTGAGTGG
GTTCAAGACGTGGAGTGGACGGAGTGGTTTCGATCATGATAAAGTATCTCAGACTAAAGG
TGGGGAGAAAATTTAGACCTCTGAAAGCTCACCCAGGAAAAATATGCAATCGTCCCA
TTGATATACAGGCCACTACAATGGATGGAGTTAACCTCAGCACCGAGGTTGTCTACAAA
AAAGGCCAGGATTATAGGTTTGCTTGCTACGACCCGGGGCAGAGCCTGCCGGAGCTACCG
TGTACGGTTCCTCTGTGGGAAGCCTGTGAGGCCCAAACCTCACAGTCACCATTGACACCA
ATGTGAACAGCACCATTTCTGAACTTGGAGGATAATGTACAGTCATGGAACCTGGAGAT
ACCCTGGTCATTGCCAGTACTGATTAATCCATGTACCAGGCAGAAGAGTTCCAGGTGCT
TCCCTGCAGATCCTGCGCCCCAACCCAGGTCAAAGTGGCAGGGAAACCAATGTACCTGC
ACATCGGGGAGAGATAGACGGCGTGGACATCGGGCGGAGGTTGGGCTTTCAGCCGG
AAATCATAGTAGTATGGGGAGATGGAGGACAAATGCTACCCCTACAGAAACCATCATG
CAATTTCTTTGACTTCGATACCTTTGGGGGCCACATCAAGTTTGCTCTGGGATTTAAGG
CAGCACACTTGGAGGGCACGGAGCTGAAGCATATGGGACAGCAGCTGGTGGGTGAGTAC
CCGATTCACCTTCCACCTGGCCGGTGTGTAGACGAAAGGGGAGGTTATGACCCACCCAC
ATACATCAGGGACCTCTCCATCCATCATACTTCTCTCGCTGCGTCAAGTCCATGGCT
CCAATGGCTTGTGATCAAGGACGTTGTGGGCTATAACTCTTTGGGCCACTGCTTCTTC
ACGGAAGATGGGCCGGAGGAACGCAACACTTTTGACCACTGTCTTGGCCTCCTTGTCAA
GTCTGGAACCCCTCCTCCCCTCGGACCGTGCAGCAAGATGTGCAAGATGATCACAGAGG
ACTCCTACCCAGGTACATCCCCAAGCCCAGGCAAGACTGCAATGCTGTGTCCACCTTC
TGGATGGCCAATCCCAACAACAACCTCATCAACTGTGCCGCTGCAGGATCTGAGGAAAC
TGGATTTTGGTTTATTTTTTACCACGTACCAACGGGCCCTCCGTGGGAATGTACTCCC
CAGGTTATTAGAGCACATTCCACTGGGAAAATCTATAACAACCGAGCACATTCCAAC
TACCGGGCTGGCATGATCATAGACAACGGAGTCAAACACCCAGGCCTCTGCCAAGGA
CAAGCGGCCGTTCTCTCAATCATCTCTGCCAGATACAGCCCTCACCAGGACGCCGACC
CGCTGAAGCCCCGGGAGCCGGCCATCATCAGACACTTCATTGCCTACAAGAACCAGGAC
CACGGGGCTGGCTGCGCGGGGGGATGTGTGGCTGGACAGCTGCCGGTTTGTGACAA
TGGCATTGGCCTGACCTGGCCAGTGGTGGAACTTCCCGTATGACGACGGCTCCAAGC
AAGAGATAAAGAACAGCTTGTGTTGGCGAGAGTGGCAACGTGGGGACGGAAATGATG
GACAAATAGGATCTGGGGCCCTGGCGGCTTGGACCATAGCGGAAGGACCCTCCCTATAGG

CCAGAATTTTCCAATTAGAGGAATTCAGTTATATGATGGCCCCATCAACATCCAAAAC
GCACTTTCCGAAAGTTTGTGGCCCTGGAGGGCCGGCACACCAGCGCCCTGGCCTTCCGC
CTGAATAATGCCTGGCAGAGCTGCCCCATAACAACGTGACCGGCATTGCCTTTGAGGA
CGTTCGGATTACTTCCAGAGTGTCTTTCGGAGAGCCTGGGCCCTGGTTCAACCAGCTGG
ACATGGATGGGGATAAGACATCTGTGTTCCATGACGTGACCGGCTCCGTGTCCGAGTAC
CCTGGCTCCTACCTCACGAAGAATGACAACTGGCTGGTCCGGCACCCAGACTGCATCAA
TGTTCCCGACTGGAGAGGGGCCATTTGCAGTGGGTGCTATGCACAGATGTACATTC AAG
CCTACAAGACCAGTAACCTGCGAATGAAGATCATCAAGAATGACTTCCCCAGCCACCCT
CTTTACCTGGAGGGGGCGCTCACCAGGAGCACCCATTACCAGCAATACCAACCGGTTGT
CACCTGCAGAAGGGCTACACCATCCACTGGGACCAGACGGCCCCCGCGAACTCGCCA
TCTGGCTCATCAACTTCAACAAGGGCGACTGGATCCGAGTGGGGCTCTGCTACCCGCGA
GGCACCACATTCCTCATCCTCTCGGATGTTACAATCGCCTGCTGAAGCAAACGTCCAA
GACGGGCGTCTTCGTGAGGACCTTGCAGATGGACAAAGTGGAGCAGAGCTACCCTGGCA
GGAGCCACTACTACTGGGACGAGGACTCAGGGCTGTTGTTCTTGAAGCTGAAAGCTCAG
AACGAGAGAGAGAAGTTTGCTTTCTGCTCCATGAAAGGCTGTGAGAGGATAAAGATTAA
AGCTCTGATTCCAAAGAACGCAGGCGTCAGTGAAGTGCACAGCCACAGCTTACCCCAAGT
TCACCGAGAGGGCTGTCTGATAGACGTGCCGATGCCCAAGAAGCTCTTTGGTTCTCAGCTG
AAAACAAAGGACCATTTCTTGGAGGTGAAGATGGAGAGTTC AAGCAGCACTTCTTCCA
CCTCTGGAACGACTTCGCTTACATTGAAGTGGATGGGAAGAAGTACCCAGTTCGGAGG
ATGGCATCCAGGTGGTGGTGTGATTGACGGGAACCAAGGGCGCGTGGTGAGCCACACGAGC
TTCAGGAACTCCATTCTGCAAGGCATACCATGGCAGCTTTTCAACTATGTGGCGACCAT
CCCTGACAATTCATAGTGCTTATGGCATCAAAGGGAAGATACGTCTCCAGAGGCCCAT
GGACCAGAGTGCTGGAAAAGCTTGGGGCAGACAGGGGTCTCAAGTTGAAAGAGCAAATG
GCATTCGTTGGCTTCAAAGGCAGCTTCCGGCCCATCTGGGTGACACTGGACACTGAGGA
TCACAAAGCCAAAATCTTCCAAGTTGTGCCCATCCCTGTGGTGAAGAAGAAGAAGTTGT
GAGGACAGCTGCCGCCCGGTGCCACCTCGTGGTAGACTATG

Figure 4. Nucleic acid sequence of ColoUp2 (SEQ ID NO: 5)

GCCCCCTGGCCCGAGCCGCGCCCGGGTCTGTGAGTAGAGCCCGCCGGGCACCGAGCGCT
GGTCGCGCGCTCTCCTTCCGTTATATCAACATGCCCCCTTTCCTGTTGCTGGAAGCCGTC
TGTGTTTTTCTGTTTTCCAGAGTGCCCCCATCTCTCCCTCTCCAGGAAGTCCATGTAAG
CAAAGAAACCATCGGGAAGATTTT CAGCTGCCAGCAAATGATGTGGTGCTCGGCTGCAG
TGGACATCATGTTTCTGTTAGATGGGTCTAACAGCGTCGGGAAAGGGAGCTTTGAAAGG
TCCAAGCACTTTGCCATCACAGTCTGTGACGGTCTGGACATCAGCCCCGAGAGGGT CAG
AGTGGGAGCATTCAGTTCAGTTCACCTCCTCATCTGGAATTCCCCTTGGATTTCATTTT
CAACCCAACAGGAAGTGAAGGCAAGAATCAAGAGGATGGTTTTCAAAGGAGGGCGCACG
GAGACGGAACTTGCTCTGAAATACCTTCTGCACAGAGGGTTGCCTGGAGGCAGAAATGC
TTCTGTGCCCCAGATCCTCATCATCGTCACTGATGGGAAGTCCCAGGGGGATGTGGCAC
TGCCATCCAAGCAGCTGAAGGAAAGGGGTGTCACTGTGTTTGCTGTGGGGGT CAGGTTT
CCCAGGTGGGAGGAGCTGCATGCACTGGCCAGCGAGCCTAGAGGGCAGCACGTGCTGTT
GGCTGAGCAGGTGGAGGATGCCACCAACGGCCTCTTCAGCACCCCTCAGCAGCTCGGCCA
TCTGCTCCAGCGCCACGCCAGACTGCAGGGTTCGAGGCTCACCCCTGTGAGCACAGGACG
CTGGAGATGGTCCGGGAGTTCGCTGGCAATGCCCCATGCTGGAGAGGATCGCGGCGGAC
CCTTGCGGTGCTGGCTGCACACTGTCCCTTCTACAGCTGGAAGAGAGTGTTCCTAACCC
ACCCTGCCACCTGCTACAGGACCACCTGCCAGGCCCTGTGACTCGCAGCCCTGCCAG
AATGGAGGCACATGTGTTCCAGAAGGACTGGACGGCTACCAGTGCCTCTGCCCGCTGGC
CTTTGGAGGGGAGGCTAACTGTGCCCTGAAGCTGAGCCTGGAATGCAGGGTTCGACCTCC
TCTTCTGCTGGACAGCTCTGCGGGCACCCTCTGGACGGCTTCTGCGGGCCAAAGTC
TTCGTGAAGCGTTTTGTGCGGGCCGTGCTGAGCGAGGACTCTCGGGCCCGAGTGGGTGT
GGCCACATACAGCAGGGAGCTGCTGGTGGCGGTGCCTGTGGGGGAGTACCAGGATGTGC
CTGACCTGGTCTGGAGCCTCGATGGCATTCCCTTCCGTGGTGGCCCCACCCTGACGGGC
AGTGCCTTGCGGCAGGCGGCAGAGCGTGGCTTCGGGAGCGCCACCAGGACAGGCCAGGA
CCGGCCACGTAGAGTGGTGGTTTTGCTCACTGAGTCACTCCGAGGATGAGGTTGCGG
GCCCAGCGCGTACGCAAGGGCGCGAGAGCTGCTCCTGCTGGGTGTAGGCAGTGAGGCC
GTGCGGGCAGAGCTGGAGGAGATCACAGGCAGCCCAAAGCATGTGATGGTCTACTCGGA
TCCTCAGGATCTGTTCAACCAAATCCCTGAGCTGCAGGGGAAGCTGTGCAGCCGGCAGC
GGCCAGGGTGC CGGACACAAGCCCTGGACCTCGTCTTCATGTTGGACACCTCTGCCTCA
GTAGGGCCCCGAGAATTTTGCTCAGATGCAGAGCTTTGTGAGAAGCTGTGCCCTCCAGTT
TGAGGTGAACCTGACGTGACACAGGTTCGGCCTGGTGGTGTATGGCAGCCAGGTGCAGA
CTGCCTTCGGGCTGGACACCAAACCCACCCGGGCTGCGATGCTGCGGGCCATTAGCCAG
GCCCCCTACCTAGGTGGGGTGGGCTCAGCCGGCACCCGCCCTGCTGCACATCTATGACAA
AGTGATGACCGTCCAGAGGGGTGCCCGGCCTGGTGTCCCCAAAGCTGTGGTGGTGTCTCA
CAGGCGGGAGAGGCGCAGAGGATGCAGCCGTTCCCTGCCCAGAAGCTGAGGAACAATGGC
ATCTCTGTCTTGGTTCGTGGGCGTGGGGCCTGTCTAAGTGAGGGTCTGCGGAGGCTTGC
AGGTCCCCGGGATTCCCTGATCCACGTGGCAGCTTACGCCGACCTGCGGTACCACCAGG
ACGTGCTCATTGAGTGGCTGTGTGGAGAAGCCAAGCAGCCAGTCAACCTCTGCAAACCC
AGCCCGTGCATGAATGAGGGCAGCTGCGTCTGCAGAATGGGAGCTACCGCTGCAAGTG
TCGGGATGGCTGGGAGGGCCCCACTGCGAGAACCGATTCTTGAGACGCCCCTGAGGCA
CATGGCTCCCCTGCAGGAGGGCAGCAGCCGTACCCCTCCCAGCAACTACAGAGAAGGCC
TGGGCAC TGAAATGGTGCCTACCTTCTGGAATGTCTGTGCCCCAGGTCCCTTAGAATGTC
TGCTTCCC GCCGTGGCCAGGACCACTATTCTCACTGAGGGAGGAGGATGTCCCAACTGC
AGCCATGCTGCTTAGAGACAAGAAAGCAGCTGATGTCAACCACAAACGATGTTGTTGAA
AAGTTTTGATGTGTAAGTAAATACCCACTTTCTGTACCTGCTGTGCCTTGTGAGGCTA

TGTCATCTGCCACCTTTCCCTTGAGGATAAACAAGGGGTCCTGAAGACTTAAATTTAGC
GGCCTGACGTTCCCTTTGCACACAATCAATGCTCGCCAGAATGTTGTTGACACAGTAATG
CCCAGCAGAGGCCTTTACTAGAGCATCCTTTGGACGG

Figure 5. Nucleic acid sequence of Osteopontin (SEQ ID NO: 6)

GCAGAGCACAGCATCGTTCGGGACCAGACTCGTCTCAGGCCAGTTGCAGCCTTCTCAGCC
AAACGCCGACCAAGGAAAACCTCACTACCAATGAGAATTGCAGTGATTTGCTTTTGCCTCC
TAGGCATCACCTGTGCCATAACAGTTAAACAGGCTGATTCTGGAAGTTCTGAGGAAAAG
CAGCTTTACAACAAATACCCAGATGCTGTGGCCACATGGCTAAACCCTGACCCATCTCA
GAAGCAGAATCTCCTAGCCCCACAGACCCTTCCAAGTAAGTCCAACGAAAGCCATGACC
ACATGGATGATATGGATGATGAAGATGATGATGACCATGTGGACAGCCAGGACTCCATT
GACTCGAACGACTCTGATGATGTAGATGACACTGATGATTCTCACCAGTCTGATGAGTC
TCACCATTCTGATGAATCTGATGAAGTGGTCACTGATTTTCCCACGGACCTGCCAGCAA
CCGAAGTTTCACTCCAGTTGTCCCCACAGTAGACACATATGATGGCCGAGGTGATAGT
GTGGTTTATGGACTGAGGTCAAATCTAAGAAGTTTTCGACAGCCTGACATCCAGTACCC
TGATGCTACAGACGAGGACATCACCTCACACATGGAAAGCGAGGAGTTGAATGGTGCAT
ACAAGGCCATCCCCGTTGCCCAGGACCTGAACGCGCCTTCTGATTGGGACAGCCGTGGG
AAGGACAGTTATGAAACGAGTCAGCTGGATGACCAGAGTGCTGAAACCCACAGCCACAA
GCAGTCCAGATTATATAAGCGGAAAGCCAATGATGAGAGCAATGAGCATTCCGATGTGA
TTGATAGTCAGGAACCTTCCAAAGTCAGCCGTGAATTCCACAGCCATGAATTCACAGC
CATGAAGATATGCTGGTTGTAGACCCCAAAGTAAGGAAGAAGATAAACACCTGAAATT
TCGTATTTCTCATGAATTAGATAGTGCATCTTCTGAGGTCAATTAAAAAGGAGAAAAAAT
ACAATTTCTCACTTTGCATTTAGTCAAAAGAAAAAATGCTTTATAGCAAAATGAAAGAG
AACATGAAATGCTTCTTTCTCAGTTTATTGGTTGAATGTGTATCTATTTGAGTCTGGAA
ATAACTAATGTGTTTGATAATTAGTTTAGTTTGTGGCTTCATGGAAACTCCCTGTAAC
TAAAAGCTTCAGGGTTATGTCTATGTTCACTTCTATAGAAGAAATGCAAACATCACTGT
ATTTAATATTTGTTATTCTCTCATGAATAGAAATTTATGTAGAAGCAAACAAAATACT
TTTACCCACTTAAAAAGAGAAATAACATTTTATGTCACTATAATCTTTTGTTTTTTAA
GTTAGTGTATATTTTGTGTTGATTATCTTTTGTGGTGTGAATAAATCTTTTATCTTGA
ATGTAATAAGAATTTGGTGGTGTCAATTGCTTATTTGTTTTCCCACGGTTGTCCAGCAA
TTAATAAAACATAACCTTTTTTACTGCCTAAAAA

Figure 6. Nucleic acid sequence of ColoUp3 (SEQ ID NO: 7)

AAAGGGGCAAGAGCTGAGCGGAACACCGCCCCGCGTCGCGGCAGCTGCTTCACCCCTC
TCTCTGCAGCCATGGGGCTCCCTCGTGGACCTCTCGCGTCTCTCCTCCTTCTCCAGGTT
TGCTGGCTGCAGTGCAGCGGCCTCCGAGCCGTGCCGGGCGGTCTTCAGGGAGGCTGAAGT
GACCTTGAGGGCGGGAGGGCGCGGAGCAGGAGCCCGCCAGGCGCTGGGGAAAGTATTCA
TGGGCTGCCCTGGGCAAGAGCCAGCTCTGTTAGCACTGATAATGATGACTTCACTGTG
CGGAATGGCGAGACAGTCCAGGAAAGAAGGTCACCTGAAGGAAAGGAATCCATTGAAGAT
CTTCCCATCCAAACGTATCTTACGAAGACACAAGAGAGATTGGGTGGTTGCTCCAATAT
CTGTCCCTGAAAATGGCAAGGGTCCCTTCCCCCAGAGACTGAATCAGCTCAAGTCTAAT
AAAGATAGAGACACCAAGATTTTCTACAGCATCACGGGGCCGGGGGCAGACAGCCCCC
TGAGGTGTCTTCGCTGTAGAGAAGGAGACAGGCTGGTGTGTGTTGAATAAGCCACTGG
ACCGGGAGGAGATTGCCAAGTATGAGCTCTTTGGCCACGCTGTGTGTCAGAGAATGGTGCC
TCAGTGGAGACCCCATGAACATCTCCATCATCGTGACCGACCAGAATGACCACAAGCC
CAAGTTTACCCAGGACACCTTCCGAGGGAGTGTCTTAGAGGGAGTCTACCAGTACTT
CTGTGATGCAGGTGACAGCCACGGATGAGGATGATGCCATCTACACCTACAATGGGGTG
GTTGCTTACTCCATCCATAGCCAAGAACCAAAGGACCCACACGACCTCATGTTACCCAT
TCACCGGAGCACAGGCACCATCAGCGTCATCTCCAGTGGCCTGGACCGGGAAAAAGTCC
CTGAGTACACACTGACCATCCAGGCCACAGACATGGATGGGGACGGCTCCACCACCACG
GCAGTGGCAGTAGTGGAGATCCTTGATGCCAATGACAATGCTCCCATGTTTGACCCCCA
GAAGTACGAGGCCCATGTGCCTGAGAATGCAGTGGGCCATGAGGTGCAGAGGCTGACGG
TCACTGATCTGGACGCCCCAACTCACCAGCGTGGCGTGCCACCTACCTTATCATGGGC
GGTGACGACGGGGACCATTTTACCATCACCACCCACCCTGAGAGCAACCAGGGCATCCT
GACAACCAGGAAGGGTTTGGATTTGAGGCCAAAACCAGCACACCCTGTACGTTGAAG
TGACCAACGAGGCCCCCTTTTGTGCTGAAGCTCCCAACCTCCACAGCCACCATAGTGGTC
CACGTGGAGGATGTGAATGAGGCACCTGTGTTTGTCCCACCCTCCAAAGTCGTTGAGGT
CCAGGAGGGCATCCCCACTGGGGAGCCTGTGTGTGTCTACACTGCAGAAGACCCCTGACA
AGGAGAATCAAAGATCAGCTACCGCATCCTGAGAGACCCAGCAGGGTGGCTAGCCATG
GACCCAGACAGTGGGCAGGTACAGCTGTGGGCACCCTCGACCGTGAGGATGAGCAGTT
TGTGAGGAACAACATCTATGAAGTCATGGTCTTGGCCATGGACAATGGAAGCCCTCCA
CCACTGGACGGGAACCTTCTGCTAACACTGATTGATGTCAATGACCATGGCCCAGTC
CCTGAGCCCCGTCAGATCACCATCTGCAACCAAAGCCCTGTGCGCCAGGTGCTGAACAT
CACGGACAAGGACCTGTCTCCCCACACCTCCCCTTTCCAGGCCAGCTCACAGATGACT
CAGACATCTACTGGACGGCAGAGGTCAACGAGGAAGGTGACACAGTGGTCTTGTCCCTG
AAGAAGTTCCTGAAGCAGGATACATATGACGTGCACCTTTCTCTGTCTGACCATGGCAA
CAAAGAGCAGCTGACGGTGATCAGGGCCACTGTGTGCGACTGCCATGGCCATGTCGAAA
CCTGCCCTGGACCTGGAAGGGAGGTTTCATCCTCCCTGTGCTGGGGCTGTCTGGCT
CTGCTGTTCCCTCCTGCTGGTGTGCTTTTGTGTTGGTGAGAAAGAAGCGGAAGATCAAGGA
GCCCCCTCTACTCCAGAAGATGACACCCGTGACAACGTCTTCTACTATGGCGAAGAGG
GGGGTGGCGAAGAGGACCAGGACTATGACATCACCCAGCTCCACCGAGGTCTGGAGGCC
AGGCCGGAGGTGGTTCTCCGCAATGACGTGGCACCAACCATCATCCCCGACACCCATGTA
CCGTCTCGGCCAGCCAAACCAGATGAAATCGGCAACTTTATAATTGAGAACCTGAAGG
CGGCTAACACAGACCCACAGCCCCGCCCTACGACACCCTCTTGGTGTTCGACTATGAG
GGCAGCGGCTCCGACGCCCGCTCCCTGAGCTCCCTCACCTCCTCCGCCTCCGACCAAGA
CCAAGATTACGATTATCTGAACGAGTGGGGCAGCCGCTTCAAGAAGCTGGCAGACATGT
ACGGTGGCGGGGAGGACGACTAGGCGGCCCTGCCTGCAGGGCTGGGGACCAAACGTCAGG
CCACAGAGCATCTCAAGGGTCTCAGTTCCCCCTTCCAGCTGAGGACTTCGGAGCTTGT

CAGGAAGTGGCCGTAGCAACTTGGCGGAGACAGGCTATGAGTCTGACGTTAGAGTGGTT
GCTTCCTTAGCCTTTCAGGATGGAGGAATGTGGGCAGTTTGACTTCAGCACTGAAAACC
TCTCCACCTGGGCCAGGGTTGCCTCAGAGGCCAAGTTTCCAGAAGCCTCTTACCTGCCG
TAAAATGCTCAACCCTGTGTCCTGGGCCTGGGCCTGCTGTGACTGACCTACAGTGGACT
TTCTCTCTGGAATGGAACCTTCTTAGGCCTCCTGGTGCAACTTAATTTTTTTTTTTAAT
GCTATCTTCAAAACGTTAGAGAAAGTTCTTCAAAAGTGCAGCCCAGAGCTGCTGGGCCC
ACTGGCCGTCCTGCATTTCTGGTTTCCAGACCCCAATGCCTCCCATTCGGATGGATCTC
TGCGTTTTTATACTGAGTGTGCCTAGGTTGCCCTTATTTTTTATTTCCCTGTTGCGT
TGCTATAGATGAAGGTGAGGACAATCGTGTATATGTACTAGAACTTTTTTATTAAAGA
AACTTTCCAGAAAAAAA

Figure 7. Nucleic acid sequence of ColoUp4 (SEQ ID NO: 8)

ATGAAGCACCTGAAGCGGTGGTGGTTCGGCCGGCGGCCTCCTGCACCTCACCTCCT
 GCTGAGCTTGGCGGGCTCCGCGTAGACCTAGATCTTTACCTGCTGCTGCCGCCGCCA
 CCTGCTGCAGGACGAGCTGCTGTTCTGGGCGGCCCGCCAGCTCCGCTACGCGCTC
 AGCCCTTCTCGGCTCGGGAGGGTGGGGCGCGCGGGCCACTTGCACCCCAAGGGCCG
 GGAGCTGACCTGCCGCGCCCGCCGAGGGCCAGCTGCTCCGGGAGGTGCGCGCGCTCG
 GGTCCCCTTCGTCCCTCGCACCCAGCGTGGATGCATGGCTGGTGCACAGCGTGGCTGCC
 GGGAGCGCGGACGAGGCCACGGGCTGCTCGGCGCCGCCCGCCTCGTCCACCGGAGG
 AGCCGGCGCCAGCGTGGACGGCGGCAGCCAGGCTGTGCAGGGGGGGCGCGGGGACCCCC
 GAGCGGCTCGGAGTGGCCCCCTGGACGCCGGGAAGAGGAGAAGGCACCCGCGGAACCG
 ACGGCTCAGGTGCCGACGCTGGCGGATGTGCGAGCGAGGAGAATGGGGTACTAAGAGA
 AAAGCACGAAGCTGTGGATCATAGTTCACGATGAGGAAAATGAAGAAAGGGTGTGAG
 CCCAGAAGGAGAATCACTTCCAGCAGATGATGATGATGAAAACAAAATAGCAGAGAAA
 CCTGACTGGGAGGCAGAAAAGACCACTGAATCTAGAAAATGAGAGACATCTGAATGGGAC
 AGATACTTCTTTCTCTCTGGAAGACTTATTCCAGTTGCTTTTCATCACAGCTGAAAATT
 CACTGGAGGGCATCTCATTGGGAGATATTCTCTTCCAGGCAGTATCAGTGATGGCATG
 AATTCTTCAGCACATTATCATGTAACTTCAGCCAGGCTATAAGTCAGGATGTGAATCT
 TCATGAGGCCATCTTGCTTTGTCCCAACAATACATTTAGAAGAGATCCAACAGCAAGGA
 CTTACAGTCACAAGAACCATTTCTGCAGTTAAATTCTCATAACCAATCCTGAGCAA
 ACCCTTCTGGAACATAATTTGACAGGATTTCTTTACCGGTTGACAATCATATGAGGAA
 TAATGTCATTGGCCACAGAAGACAACCTTTGATCCAATCGATGTTTCTCAGCTTTTGT
 GAACCAGATTCTGATTCTGGCCTTTCTTTAGATTCAAGTCACAATAATACCTCTGTCT
 CAAGTCTAATTCCTCTCACTCTGTGTGTGATGAAGGTGCTATAGGTTATTGCACTGACC
 ATGAATCTAGTTCCCATCATGACTTAGAAGGTGCTGTAGGTGGCTACTACCAGAACCC
 AGTAAGCTTTGTCACTTGGATCAAAGTGATTCTGATTTCCATGGAGATCTTACATTTCA
 ACACGTATTTTATAACCACACTTACCCTTACAGCCAACTGCACCAGAATCTACTTCTG
 AACCTTTTCCGTGGCCTGGGAAGTCACAGAAGATAAGGAGTAGATACCTTGAAGACACA
 GATAGAACTTGAGCCGTGATGAACAGCGTGCTAAAGCTTTGCATATCCCTTTTTCTGT
 AGATGAAATTGTGGCATGCCTGTTGATTCTTTCAATAGCATGTTAAGTAGATATTATC
 TGACAGACCTACAAGTCTCACTTATCCGTGACATCAGACGAAGAGGGAAAAATAAAGTT
 GCTGCGCAGAAGTGTGTAACGCAAATTTGGACATAATTTTGAATTTAGAAGATGATGT
 ATGTAAGTTCGAAGCAAAGAAGGAACTCTTAAGAGAGAGCAAGCACAATGTAACAAAG
 CTATTAACATAATGAAACAGAACTGCATGACCTTTATCATGATATTTTAGTAGATTA
 AGAGATGACCAAGGTAGGCCAGTCAATCCAACCACTATGCTCTCCAGTGTACCCATGA
 TGGAAATCTTTGATAGTACCCAAAGAACTGGTGGCTCAGGCCACAAAAGGAAACCC
 AAAAGGGAAAGAGAAAGTGAAGAAGAACTGAAGATGGACTCTATTATGTGAAGTAGTAA
 TGTTTCAGAAACTGATTATTTGGATCAGAAAACCATTGAACTGCTTCAAGAATTGTATCT
 TTAAGTACTGCTACTTGAATAACTCAGTTAACGCTGTTTGAAGCTTACATGGACAAAT
 GTTtaggacttcaagatcacacttGTGGGCAATCTGGGGGAGCCACAACCTTTTCATGAA
 GTGCATTGTATACAAAATTCATAGTTATGTCCAAAGAATAGGTTAACATGAAAACCCAG
 TAAGACTTTCCATCTTGGCAGCCATCCTTTTTAAGAGTAAGTTGGTTACTTCAAAAAGA
 GCAAACACTGGGGATCAAATATTTTAAGAGGTATTTCAAGTTTTAAATGCAAAATAGCC
 TTATTTTTCATTTAGTTTGTAGCACTATAGTGAGCTTTCAAACACTATTTTAATCTTT
 ATATTTAACTTATAAATTTTGCTTTCTATGGAAATAAATTTTGTATTTGTATTAATAAAA
 AAAAAA

Figure 8. Nucleic acid sequence of ColoUp5 (SEQ ID NO: 9)

ATGAAGTTGGAGGTGTTCGTCCCTCGCGCGGCCACGGGGACAAGCAGGGCAGTGACCT
GGAGGGCGCGGGCGGCAGCGACGCGCCGTCCCCGCTGTCGCGCGCGGGAGACGACTCCC
TGGGCTCAGATGGGGACTGCGCGGCCAAGCCGTCCGCGGGCGCGGCCAGAGATACG
CAGGGCGACGGCGAACAGAGTGGCGGGAGGCGGGCCGGGCGCGGAGGAGGCGATCCCGGC
AGCAGCTGCTGCAGCGGTGGTGGCGGAGGGCGCGGAGGCCGGGGCGCGGGGGCCAGGGC
CGGGCGGCGCGGGGAGCGGCGAGGGTGCACGCAGCAAGCCATATACGCGGGCGGCCAAG
CCCCCTACTCGTACATCGCGCTCATCGCCATGGCCATCCGCGACTCGGCGGGCGGGCG
CTTGACGCTGGCGGAGATCAACGAGTACCTCATGGGCAAGTTCCCTTTTTCCGCGGCA
GCTACACGGGCTGGCGCAACTCCGTGCGCCACAACCTTTCGCTCAACGACTGCTTCGTC
AAGGTGCTGCGGACCCCTCGCGGCCCTGGGGCAAGGACAACACTACTGGATGCTCAACCC
CAACAGCGAGTACACCTTCGCCGACGGGGTCTTCGCCCGCCCGCAAGCGCCTCAGCC
ACCGCGCGCCGTCCCCGCGCCCGGGCTGCGGCCCGAGGAGGCCCGGGCCTCCCCGCC
GCCCCGCGCCCGCGCCCGCCGCCCCCGCCTCGCCCCGCATGCGCTCGCCCGCCGCCA
GGAGGAGCGCGCCAGCCCCGCGGCAAGTTCTCCAGCTCCTTCGCCATCGACAGCATCC
TGCGCAAGCCCTTCGCAGCCGTGCCTCAGGGACACGGCCCCGGGACGACGCTTCAG
TGGGGCGCCGCGCCCTGCCCGCGCTGCCCGCGTTCCCCGCGCTCCTCCCCGCGGCGCC
CTGCAGGGCCCTGCTGCCGCTCTGCGCGTACGGCGCGGGCGAGCCGGCGCGGCTGGGGC
CGCGCGAGGCCGAGGTGCCACCGACCGCGCCGCCCTCCTGCTTGACCTCTCCCGGC
GCGGCCCCCGCAAGCCACTCCGAGGCCCGGGCGGCGGGCGCGCACCTGTACTGCC
CCTGCGGCTGCCCGAGCCCTGCAGGCGGCCTTAGTCCGNCGTCTGGCCCGCACCTGT
CGTACCCGGTGGAGACGCTCCTAGCTTGA

Figure 9. Nucleic acid sequence of ColoUp6 (SEQ ID NO: 10)

GGCAGATGAAATATAAGATTCATCAACCACATTTGACAGCCCATGGCAGGTTTTCTGT
TTCCATCGTCCCTCTGCAGGTCACAGACACACAGAGCCCAGCCGTGGCAGGCTCAGCCG
GGTCCGGGGCTGCTAACAACGGCTACATTCCTCCCCAGGGCCAAGGGAAATCCTGAG
CGCAGGCCAGGTTGTTTGGTTTTGAGGTGTGCTGGGATGAAAGGCACCCTGGAAGTGG
AAGTTTCGGTCATTCATTAATTAATTACATCTATAATTGAGGGTTTTGTTCTTAAGAGCG
AGTCTTTTGAAGTACTTTCTTTCAAACAGTGAAGTCCACAAAGGCATCAGATATTCAC
CACCTTCTCGGCTGCCTCAGCACAGCAAGCTTTATTCTGGGACCTGAGATCCTGTTCTG
AGCTGGCTTTCCCTTCTCCAGGCTCGCTCACCTCCCTTTAGAGATAGTGGATGGTAAG
ATGACCAATGCTCAGATTATTCTTCTCATTGACAATGCCAGGATGGCAGTGGATGACTT
CAACCTCAAGAAATGGAGAAGCATCATGTGCCAAGTGAAGTCAATGTCAATGTGAAGGT
GGATACAGGTTCCAGGGAAGATCTGATTAAGGTCTGGAGGATATGAGACAAGAATATG
AGCTTATAATAAAGAAGAAGCATCGAGACTTGGACACTTGGTATAAAGAACAGTCTGCA
GCCATGTCCCAGGAGGCAGCCAGTCCAGCCACTGTGCAGAGCAGACAAGGTGACATCCA
CGAACTGAAGCGCACATTCAGGCCCTGGAGATTGACCTGCAGGCACAGTACAGCACGA
AATCTGCTTTGGAAAACATGTTATCCGAGACCCAGTCTCGGTACTCCTGCAAGCTCCAG
GACATGCAAGAGATCATCTCCCACTATGAGGAGGAAGTACGCAGCTACGCCACGAACT
GGAGCGGCAGAACAATGAATACCAAGTGTGCTGGGCATCAAAACCCACCTGGAGAAGG
AAATCACCACGTACCGACGGCTCCTGGAGGGAGAGAGTGAAGGGACACGGGAAGAATCA
AAGTCGAGCATGAAAGTGTCTGCAACTCCAAAGATCAAGGCCATAACCCAGGAGACCAT
CAACGGAAGATTAGTTCTTTGTCAAGTGAATGAAATCCAAAAGCACGCATGAGACCAAT
GAAAGTTTCCGCTGTTGTAAAATCTATTTTCCCCAAGGAAAGTCTTGCACAGACAC
CAGTGAAGTGTCTAAAAGATACCCTTGAATTATCAGACTCAGAACTTTTATTTTT
TTTTTCTGTAAACAGTCTCACCAGACTTCTCATAATGCTCTTAATATATTGCACTTTTCT
AATCAAAGTGGAGTTTATGAGGGTAAAGCTCTACTTTCTACTGCAGCCTTCAGATTC
TCATCATTTTGCATCTATTTTGTAGCCAATAAAACTCCGCACTAGCAAAAAAAAAAAAA

Figure 10. Nucleic acid sequence of ColoUp7 (SEQ ID NO: 11)

TTTTTTTTTTAAAAAAGAGGCTTGGTAAGTTTTTGATGCTTAGTTGACTTTTAGCATT
ATCCAGCATTGTATTATGAACCAGTGAGTACTGTAATTTTTCTTCCCTTTCAGAAAG
ACTCAAAGGGAACATATAAATGTTTCCTATTTTTAATGTGGCAATAGTGTAGCTAACAC
TGGTACAGACGGAATAAACACACCTCTAATATTCTCCTGAAGATTTGGTGATCCAGTTT
CAAATAAGGTATGGGAAAAACAGATGTTTTCATTATCGCCACTTAATCCTTACTTCCGA
TTATAATTATACATGTTTGGCTGTAATAACTATACTAAAGCATGCTTGTGAAAGTAGAC
TTCTACAAGGACAGAAAACCCACAACAACAAGATCGATCACGAAAGACAAGGCATA

Figure 11. Nucleic acid sequence of ColoUp8 (SEQ ID NO: 12)

CTTTTCTTCCGCACGGTTGGAGGAGGTTCGGCTGGTTATCGGGAGTTGGAGGGCTGAGGT
CGGGAGGGTGGTGTGTACAGAGCTCTAGGACTCACGCACCAGGCCAGTCGCGGATTTTG
GGCCGAGGCCTGGGTTACAAGCAGCAAGTGC GCGGTTGGGGCCACTGCGAGGCCGTTTT
AGAAAAC TGGTTAAAACAAGAGCAATTGATGGGATAAAATCAGGAATAGATTCTCTTGAC
CATGTGACATCTGATGCTGTGGAACCTGCAAATCGAAGTGATAACTCTTCTGATAGCAG
CTTATTTAAAAC TCAGTGTATCCCTTACTCACCTAAAAGGGGAGAAAAGAAACCCCATTC
GAAAATTTGTTTCGTACACCTGAAAGTGTTACGCAAGTGATTCATCAAGTGACTCATCT
TTTGAACCAATACCATTGACTATAAAAGCTATTTTTGAAAGATTCAAGAACAGGAAAAA
GAGATATAAAAAAAGAAAAAGAGGAGGTACCAGCCAACAGGAAGACCAGGGGAAAGAC
CAGAAGGAAGGAGAAATCCTATATACTCACTAATAGATAAGAAGAAAACAATTTAGAAGC
AGAGGATCTGGCTTCCCATTTTTAGAAATCAGAGAATGAAAAAACGCACCTTGGAGAAA
AATTTTAACTGTTGAGCAAGCTGTTGCAAGAGGATTTTTTAACTATATTGAAAAGCTGA
AGTATGAACACCACCTGAAAGAATCATTGAAGCAAATGAATGTTGGTGAAGATTTAGAA
AATGAAGATTTTGACAGTCGTAGATACAAATTTTTGGATGATGATGGATCCATTTCTCC
TATTGAGGAGTCAACAGCAGAGGATGAGGATGCAACACATCTTGAAGATAACGAATGTG
ATATCAAATTGGCAGGGGATAGTTTCATAGTAAGTTCTGAATTCCTGTAAGACTGAGT
GTATACTTAGAAGAAGAGGATATTACTGAAGAAGCTGCTTTGTCTAAAAAGAGAGCTAC
AAAAGCCAAAAATACTGGACAGAGAGGCCTGAAAATGTGACAGGATCATGAATGTCAA
GGCTTTTATCTTGAGAACATGGTGTCTGGAGTTAAAGGTATTGGCATACTCCACACATC
TGTACCATTCTTGAGTGATCGCTTAGGAATGAATGTGATTTGAACTCATTCATGTTGAG
AGGGTGTCAAATTGAGAACCAGGTAGATCCCCACCACCTACAGTAAAAAGGACCCTAAA
GTAAATTGGTTGAAGAAATTAGATCCCAAAGATTCTTGGTGAATTTTGAAGTCTTCATC
AGTATATCCATATTAACGAGATGACAGAAGCCAAAGTAATTATGGCAAGTAATGGTT
TTTATCTTAACTATAAGTTATTTGCTCAAGGGTGTAAATGGTCATTACCAAGGCTTTTAG
AATGCAGTTTCTCATTGCTGTGGACATGACCATAAAAAAAATTTCCAGTAGGTTTT
CTATCTGCTACGTTGCTAGCAATCAGCTTATTGGGAACAGTTGATTAAGTAAATAGAA
ATGCAATACAAATAAAATGTGAACCACATGTGATTTTTCTTTAAAATCAGTGAGATTTG
AAAATCTCCTAGATCTCTTGAATCATGCAAATTTGCTTTGCCTTTATATTGTAACCCT
TGTGGGTTGCTAATAACCAAGCAGTTTGTAGTAGAGTTAACTCAGGCTCGTTCTAGGGA
CTCATTATGTTCACTCACTGTACACTCATCTCTGGAAATGTAATAATTTACTTTTATAC
TATTGTTATGTAGGGCTGACAGGACAACCTGGATCAGTTTCATTAAAAAGGTATGTATGC
ATTAGAAAAGACATTTGTATGGGTCAATTTCAAAGAGGGCTTATGAGGCTGTGAAACCCA
GAGCTCTAACGCTGTGACCAAAGATGGAAGTTCTCTATAGGAAGCCATAGCACTCCTA
ATGTTTGGTGTATGTTTTCTGAGGAGATATAAACGTAATAATCCATGATTGTTGCC
ATGTGAGAGTTTTAAAGGTTAATCAAAATTTCTCTTCTCAGGGCAAACCTTGAAGATAA
ATCTTTGACTCCAGCTCTTTAGAGGATCTAAAGTGACCTTGATGGACAGTGAAGAAA
TCACAACATGGAATTCCTCGAATAACAATTTATTGACTTTAAATAATTTTGTCTAATGC
TACATATACACAATTA AAAAACCTTTACTACTATTTCTAGAAAAGTCAGCATGATTTTTG
GCTCGAAGTTTCTCTAGTGTCTTCTGTGGAAGGAATAAAAAATTTGAGTTTCAAAAAA
AAAAA

Figure 12. Amino acid sequence of full-length ColoUp1 protein (SEQ ID NO: 13)

MGAAGRQDFLFKAMLTISWLTLTCTFPGATSTVAAGCPDQSPQLPWNPGHDQDHHVHIG
QGKTLTLLTSSATVYSIHISEGGKLVIKDHDEPIVLRTRHILIDNGGELHAGSALCPFQG
NFTIILYGRADEGIQPDPIYGLKYIGVGKGALELHGQKLSWTFLNKTLHPGGMAEGG
YFFERSWHRGVIVHVIDPKSGTVIHSDFDITYRSKKESERLVQYLNVPDGRILSVAV
NDEGSRNLDDMARKAMTKLGSKHFLHLGFRHPWSFLTVMKGNPSSSVEDHIEYHGHRGSA
AARVFKLFQTEHGEYFNVSLSSEWVQDVEWTEWFDHDKVSQTKGGEKISDLWKAHPGKI
CNRPIDIQATTMDGVNLSDEVVYKKGQDYRFACYDRGRACRSYRVRFLCGKPVPRKLT
TIDTNVNSTILNLEDNVQSWKPGDTLVIASDYSMYQAEFFQVLPSCRSCAPNQVKVAGK
PMYLHIGEEIDGVMRAEVGLLSRNIIVMGEMEDKCYPYRNHICNFFDFDTFGGHIKFA
LGFKAAHLEGTELKHMGGQQLVGQYPIHFHLAGDVDERGGYDPPYIRDLSEIHTFSRCV
TVHGSNGLLIKDVVGYNSLGHCFFTEDGPEERNTFDHCLGLLVKSGTLLPSPDRDSKMCK
MITEDSYPGYIPKPRQDCNAVSTFWMANPNNNLINCAAAGSEETGFWFI FHHVPTGPSV
GMYSPIYSEHIPGKGFYNNRAHSNYRAGMIIDNGVKTTEASAKDKRPFLSII SARYSPH
QDADPLKPREPAIIRHFIAYKNQDHGAWLRGGDVWLDSCRFAFADNGIGLTLASGGTFPYD
DGSKQEIKNLSLVGSGNVGTEEMDNRIWGPGLDHSRGLPIGQNFPIRGIQLYDGPI
NIQNCTFRKFVALEGRHTSALAFRLNNAWQSCPHNNVTGIAFEDVPIITSRVFFGEPGPW
FNQLDMDGDKTSVFHDVDSVSEYPGSYLTKNDNWLVRHPDCINVPDWRGAI CSGCY AQ
MYIQAYKTSNLRMKI IKNDFPSHPLYLEGALTRSTHYQYQPVVTLQKGYTIHWDQTAP
AELAIWLINFNKGDWIRVGLCYPRGTTFSILSDVHNRLKQTSKTGVFVRTLQMDKVEQ
SYPGRSHYYWDEDSGLLFLKKAQNEREKFAFCMKGCIKIKALIPKNAGVSDCTAT
AYPKFTERAVDVPMPKFLGSQKTKDHFLEVKMESSKQHFHFWNDFAFIEVDGKKY
PSSDGIQVVVIDGNQGRVVSHTSFRNSILQGI PWQLFNYVATIPDNSIVLMASKGRYV
SRGPWTRVLEKLGADRGLKLEQMAFVGFKGSFRPIWVTLDTEDHKAKIFQVVPIPVVK
KKKL

Figure 13. Amino acid sequence of full-length ColoUp2 protein (SEQ ID NO: 14)

MPPFLLLEAVCVFLFSRVPPSLPLQEVHVSKETIGKISAASKMMWCSSAAVDIMFLLDGS
NSVGKGSFERSKHFAITVCDGLDISPERVRVGAFQFSSTPHLEFPLDSFSTQQEVKARI
KRMVFKGGRTETELALKYLLHRGLPGGRNASVPQILIIIVTDGKSQGDVALPSKQLKERG
VTVFAVGVRFPWEELHALASEPRGQHVLLAEQVEDATNGLFSTLSSSAICSSATPDCR
VEAHPCEHRTLEMVREFAGNAPCWGRSRRTLAVLAAHCPFYSWKRVFLTHPATCYRTTC
PGPCDSQPCQNGGTCVPEGLDGYQCLCPLAFGGEANCALKLSLECRVDLLFLLDSSAGT
TLDGFLRAKVFKRFVRAVLSEDSRARVGVATYSRELLVAVPVGEYQDVPDLVWSLDGI
PFRGGPTLTGSALRQAAERGFSGSATRTGQDRPRRVVLLTESHSEDEVAGPARHARARE
LLLLGVGSEAVRAELEETGSPKHVMVYSDPQDLFNQIPELQKLC SRQRP GCRTQALD
LVFMLDTSASVGPENFAQM QS FVRSCALQFEVNP DVTQVGLVVYGSQVQTAFGLDTKPT
RAAMLRAISQAPYLG GVSAGTALLHIYDKVMTVORGARPGVPKAVVVL TGGRGAEDAA
VPAQKLRNNGISVLVVGVPVLSEGLRRLAGPRDSL IHVAAYADLRYHQDVLI EWLCGE
AKQPVNLC KPS PCMNEGSCV LQNGSYRCKCRD GWEGPHCENRFLRRP

Figure 14. Amino acid sequence of full-length Osteopontin protein (SEQ ID NO: 15)

MRIAVICFLLGITCAIPVKQADSGSSEEKQLYNKYPDAVATWLNPDPSQKQNLLAPQT
LPSKSNESHDMDDMDEDDDDHVDSQDSIDSNSDDVDDTDDSHQSDESHHSDESDEL
VTDFPTDLPATEVFTPVVPTVDTYDGRGDSVVYGLRSKSKKFRRPDIQYPDATDEDITS
HMESEELNGAYKAIPVAQDLNAPSDWDSRGKDSYETSQLDDQSAETHSHKQSRLYKRKA
NDESNEHSDVIDSQELSKVSRFHSHEFHSHEDMLVVDPKSKEEDKHLKFRISHELDSA
SSEVN

Figure 15. Amino acid sequence of full-length ColoUp3 protein (SEQ ID NO: 16)

MGLPRGPLASLLLLQVCWLQCAASEPCRAVFREA EVTLEAGGAEQEPGQALGKVFMGCP
GQEPALFSTDND DFTVRNGETVQERRSLKERNPLKIFPSKRILRRHKRDWV VAPISVPE
NGKGFPPQRLNQLKSNKDRDTKIFYSITGPGADSPPEGVFAVEKETGWLLL NKPLDREE
IAKYELFGHAVSENGASVEDPMNISIIIVTDQNDHKPKFTQDTFRGSVLEGVLPGTSVMQ
VTATDEDDAIYTYNGVVAYSIHSQEPKDPHDLMFTIHRSTGTISVISSGLDREKVPEYT
LTIQATDMDGDGSTTTAVAVVEILDANDNAPMFD POKYEAHV PENAVGHEVQRLTVTDL
DAPNSPAWRATYLIMGGDDGDHFTITTHPESNQGILTRKGLDFEAKNQHTLYVEVTNE
APFVLKLP TSTATIVVHVEDVNEAPV FVPPSKVVEVQEGIPTGEPVCVYTAEDPDKENQ
KISYRILRDPAGWLAMD PDSGQVTAVGTL DREDEQFVRNNIYEVMLAMDNGSPPTTGT
GTL LLLTLIDVNDHGPVPEPRQITICNQSPVRQVLNITDKDLS PHTSPFQAQLTDDSDIY
WTAEVNEEGD TVVLSLKKFLKQD TYDVHLSLSDHGNKEQLTVIRATVCDCHGHVETCPG
PWKGGF ILPVLGAVLALLFLLL VLLLVRKKRKIKEPLLLPEDDTRDNV FYYGEEGGGE
EDQDYDITQLHRGLEARPEV VLRNDVAPTIIPTPMYRPRPANPDEIGNFIIENLKAANT
DPTAPPYDTLLVFDYEGSGSDAASLS SLSASDQDQDYDYLNEWGSRFKKLADMYGGG
EDD

Figure 16. Amino acid sequence of full-length ColoUp4 protein (SEQ ID NO: 17)

MKHLKRWSAGGGLLHLTLLLSLAGLRVDLDLYLLLPPPTLLQDELLFLGGPASSAYAL
SPFSASGGWGRAGHLHPKGRELDPAAPPEGQLLREVRALGVFPVPRTSVDAWLVHSVAA
GSADEAHGLLGAAAASSTGGAGASVDGGSQAVQGGGGDPRAARSGPLDAGEEEKAPAEP
TAQVPDAGGCASEENGLREKHEAVDHSQHEENEERVSAQKENSLOQNDDEKIAEK
PDWEAEKTTESRNERHLNGTDTFSLEDLFQLLSSQPENSLEGISLGDIPGPSISDGM
NSSAHYHVNFSAISQDVNLHEAILLCPNNTFRRDPTARTSQSQEPFLQLNSHTTNPEQ
TLPGTNLTGFLSPVDNHMRNLTSQDLLYDLINIFDEINLMSLATEDNFDPIDVSQLFD
EPDSDSGLSLDSSHNTSVIKSNSHSVCDEGAIGYCTDHESSSHDLEGAVGGYYPEP
SKLCHLDQSDSDFHGDLTFQHVFNHTYHLQPTAPESTSEPFPPWPGKSQKIRSRYLEDT
DRNLSRDEQRAKALHIPFSVDEIVGMPVDSFNSMLSRYYLTDLQVSLIRDIRRRGKKNKV
AAQNCRKRKLDIILNLEDDVCNLQAKKETLKREQAQCNKAINIMKQKLHDLYHDI FSRL
RDDQGRPVNPNHYALQCTHDGSILIVPKELVASGHKKETQKGKRK

Figure 17. Amino acid sequence of full-length ColoUp5 protein (SEQ ID NO: 18)

MKLEVFVPRAAHGDKQGS DLE GAGGSDAPSPLSAAGDDSLGSDGDCAAKPSAGGGARDT
QGDGEQSAGGGPGAEEAI PAAAAA AVVAEGAEAGAAGPGAGGAGSGEGARSKPYTRRPK
PPYSYIALIAMAIRDSAGGRLTLAEINEYLMGKFPFFRGSYTGWRNSVRHNLSLNDCFV
KVL RDPSRPWGKDNYWMLNPSEYTFADGVFRRRRKRLSHRAPVPAPGLRPEEAPGLPA
APPPAPAAPASPRMRSPARQEERAS PAGKFSSSFAIDSILRKPFRRRLRDTAPGTTLQ
WGAAPCPPLPAFPALLPAAPCRALLPLCAYGAGEPARLGAREAEVPPTAPLLLAPLPA
AAPAKPLRGPAAGGAHLYCPLRLPAALQAALVRRPGPHLSYPVETLLA

Figure 18. Amino acid sequence of full-length ColoUp6 protein (SEQ ID NO: 19)

MEKHHVPSDFNVNVKVDTPREDLIKVLEDMRQEYELIIKKKHRDLDTWYKEQSAAMSQ
EAASPATVQSRQGD I HELKRTFQALEIDLQAQYSTKSALENMLSETQSRYSCKLQDMQE
IISHYEEELTQLRH ELERQNEYQVLLGIKTHLEKEITTYRRLLEGESEGTREESKSSM
KVSATPKIKAITQETINGRLVLCQVNEIQKHA

Figure 19. Amino acid sequence of full-length ColoUp8 protein (SEQ ID NO: 20)

MDKSGIDSLDHVTSDAVELANRSDNSSDSSLFKTQCIPYSPKGEKRNPPIRKFVRTPEV
HADSSSDSSFEPILTIKAIERFKNRKKRYKKKKRRYQPTGRPRGRPEGRRNPIYS
LIDKKKQFRSRGSGFPFLESENEKNAPWRKILTFEQAVARGFFNYIEKLYEHLKESL
KQMNVEDLENEFDSSRYKFLDDDGSISPIEESTAEDEDATHLEDNECDIKLAGDSFI
VSSEFPVRLSVYLEEEDITEEAALSKKRATKAKNTGQRGLKM




Figure 20




Figure 21




Figure 22A




Figure 22B




Figure 22C




Figure 22D




Figure 23A




Figure 23B




Figure 23C




Figure 23D




Figure 24A




Figure 24B




Figure 24C




Figure 24D




Figure 25A



Figure 25B




Figure 25C




Figure 25D




Figure 26A




Figure 26B




Figure 26C




Figure 26D




Figure 27A




Figure 27B




Figure 27C




Figure 27D




Figure 28A




Figure 28B




Figure 28C




Figure 28D




Figure 29A




Figure 29B




Figure 29C




Figure 30A




Figure 30B




Figure 31




Figure 32




Figure 33




Figure 34

<i>human foxq1</i>	1	A T G A A G T T G G A G G T
<i>mouse foxq1</i>	1	A T G A A A T T G G A G G T
<i>rat foxq1</i>	1	A T G A A A T T G G A G G T
<i>human foxq1</i>	81	C G C G C C G T C C C G C
<i>mouse foxq1</i>	81	C G T G C C A T C T C C A C
<i>rat foxq1</i>	81	C G T G C C A T C T C C G C
<i>human foxq1</i>	161	G C G G C G G C G C C A G A
<i>mouse foxq1</i>	161	G C A G C G G C G C C G G G
<i>rat foxq1</i>	161	G C A G A G G C G C C G T G
<i>human foxq1</i>	238	G C A G C A G C T G C T G C
<i>mouse foxq1</i>	238	G A G G C A A C T G - - - -
<i>rat foxq1</i>	238	G A G G T G A C C G - - - -
<i>human foxq1</i>	318	C G A G G G T G C A C G C A
<i>mouse foxq1</i>	306	C G A G G G C G C G C G C A
<i>rat foxq1</i>	306	T G A G G G C G C G C G C A
<i>human foxq1</i>	398	G C G A C T C G G C G G G C
<i>mouse foxq1</i>	386	G C G A C T C C G C G G G C
<i>rat foxq1</i>	386	G C G A C T C C G C G G G C
<i>human foxq1</i>	478	T A C A C G G G C T G G C G
<i>mouse foxq1</i>	466	T A C A C G G G C T G G C G
<i>rat foxq1</i>	466	T A C A C G G G C T G G C G
<i>human foxq1</i>	558	G C C C T G G G G C A A G G
<i>mouse foxq1</i>	546	G C C C T G G G G C A A G G
<i>rat foxq1</i>	546	G C C C T G G G G C A A G G

Figure 35 (part 1)

<i>human foxq1</i>	638	G C A A G C G C C T C A G C
<i>mouse foxq1</i>	626	G C A A G C G C C T C A G C
<i>rat foxq1</i>	626	G C A A G C G C C T C A G C
<i>human foxq1</i>	715	C C G C C G C C C G C G C C
<i>mouse foxq1</i>	706	C C G C A G C C C G C G C C
<i>rat foxq1</i>	706	C C G C A G C C C G C G C C
<i>human foxq1</i>	795	C A A G T T C T C C A G C T
<i>mouse foxq1</i>	786	C A A G T T C T C C A G C T
<i>rat foxq1</i>	786	C A A G T T C T C C A G C T
<i>human foxq1</i>	875	G G A C G A C G C T T C A G
<i>mouse foxq1</i>	866	G G G T G C A G C T A C C C
<i>rat foxq1</i>	866	G G G T G C A G C T A C C C
<i>human foxq1</i>	955	G C C C T G C T G C C G C T
<i>mouse foxq1</i>	946	G C T C T G C T A C C G C T
<i>rat foxq1</i>	946	G C C C T G C T G C C G C T
<i>human foxq1</i>	1035	G C C G C C C C T C C T G C
<i>mouse foxq1</i>	1023	G G C G C C C C T T C T G C
<i>rat foxq1</i>	1023	G G C G C C C C T G T T G C
<i>human foxq1</i>	1112	A C C T G T A C T G C C C C
<i>mouse foxq1</i>	1103	A C C T G T A C T G C C C C
<i>rat foxq1</i>	1103	A C C T G T A C T G C C C C
<i>human foxq1</i>	1192	G T G G A G A C G C T C C T
<i>mouse foxq1</i>	1183	G T G G A G A C T C T G C T
<i>rat foxq1</i>	1183	G T G G A G A C G C T G C T

Figure 35 (part 2)




Figure 35 (part 3)




Figure 35 (part 4)




Figure 35 (part 5)




Figure 35 (part 6)

G A G G G C G C G G G C G G C A G C G A	80
G A G G G G G C C G G C A G C A G C G A	80
G A G G G G G C C G G C A G C A G C G A	80
G G C C A A C A G C C C G G C C G C G G	160
A G C C A A C A G C C C G G C G G C G G	160
A G C C A A C A G C C C G G C G G C G G	160
G C G C G G A G G A G G C G A T C C C G	237
G C G C C C A A G A - - C G G T C C - G	237
G C A C C C A A G A - - C G A T C C - C	237
G C G G G C G G C G C G G G G A G C G G	317
G C G G G C G G C G T G G G C G G C G G	305
G C G G G C A G C G T G G G C G G C G G	305
G C T C A T C G C C A T G G C C A T C C	397
T C T C A T C G C C A T G G C C A T C C	385
A C T C A T C G C C A T G G C C A T C C	385
T C C C C T T T T C C G C G G C A G C	477
T C C C C T T T T C C G G G G C A G C	465
T C C C C T T T T C C G G G G C A G C	465
G T G C T G C G C G A C C C C T C G C G	557
G T G C T G C G C G A C C C C T C G C G	545
G T G C T G C G C G A C C C C T C G C G	545
C G G G G T C T T C C G C C G C C G C C	637
C G G G G T C T T C C G C C G C C G C C	625
C G G G G T C T T C C G C C G C C G C C	625

Figure 35 (part 7)




Figure 35 (part 8)




Figure 36




Figure 37






Figure 38



time points(hr)	resistance	transfection efficiency
0	220	N/A
24	199	15%
48	184	25%
72	220	30%
96	214	30%

Figure 39



Day	Resistance	% GFP Exp.
0	300	0
1	415	10
2	260	20
3	206	25
4	218	25

Figure 40

Amino Acid Sequence of a Secreted C-terminal Portion of ColoUp2

AVLAAHCPFYSWKRVFLTHPATCYRTTCPGPCDSQPCQNGGTCVPEGLDGYQCL
CPLAFGGEANCALKLSLECRVDLLFLLDSSAGTTLDGFLRAKVFVKRFVRAVLSE
DSRARVGVATYSRELLVAVPVGEYQDVPDLVWSLDGIPFRGGPTLTGSALRQAA
ERGFSGATRTGQDRPRRVVLLTESHSEDEVAGPARHARARELLLLGVGSEAVR
AELEEITGSPKHVMVYSDPQDLFNQIPELQGKLCRQRPGCRTQALDLVFMLDTS
ASVGPENFAQMOSFVRSCALQFEVNPDVTQVGLVVYGSQVQTAFGLDTKPTR
AMLRAISQAPYLGGVGSAGTALLHIYDKVMTVQRGARPGVPKAVVVLTGGRGA
EDAAVPAQKLRNNGISVLVVGVPVLSEGLRRLAGPRDSLHVAAAYADLRYHQD
VLIEWLCGEAKQPVNLCCKPSPCMNEGSCVLQNGSYRCKCRDGWEGPHCENRFL
RRP (SEQ ID NO:21)

Figure 41

METHODS AND COMPOSITIONS FOR CATEGORIZING PATIENTS

CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] This application is a continuation-in-part of U.S. patent application Ser. No. 10/274,177, filed Oct. 18, 2002, which is a continuation-in-part of U.S. patent application Ser. No. 10/229,345, filed Aug. 26, 2002, and both of the aforementioned patent applications are incorporated herein by reference.

FUNDING

[0002] Work described herein was funded, in part, by grant number 1 U01 CA-88130-01 from the National Cancer Institute. The United States government has certain rights in the invention.

BACKGROUND

[0003] Colorectal cancer, also referred to herein as colon cancer, is the second leading cause of cancer mortality in the adult American population. An estimated 135,000 new cases of colon cancer occur each year. Although many people die of colon cancer, early stage colon cancers are often treatable by surgical removal (resection) of the affected tissue. Surgical treatment can be combined with chemotherapeutic agents to achieve an even higher survival rate in certain colon cancers. However, the survival rate drops to 5% or less over five years in patients with metastatic (late stage) colon cancer.

[0004] Effective screening and early identification of affected patients coupled with appropriate therapeutic intervention is proven to reduce the number of colon cancer mortalities. It is estimated that 74,000,000 older Americans would benefit from regular screening for colon cancer and precancerous colon adenomas (together, adenomas and colon cancers may be referred to as colon neoplasias). However, present systems for screening for colon neoplasia are inadequate. For example, the Fecal Occult Blood Test involves testing a stool sample from a patient for the presence of blood. This test is relatively simple and inexpensive, but it often fails to detect colon neoplasia (low sensitivity) and often even when blood is detected in the stool, a colon neoplasia is not present (low specificity). Flexible sigmoidoscopy involves the insertion of a short scope into the rectum to visually inspect the lower third of the colon. Because the sigmoidoscope is relatively short, it is also a relatively uncomplicated diagnostic method. However, nearly half of all colon neoplasia occurs in the upper portions of the colon that can not be viewed with the sigmoidoscope. Colonoscopy, in which a scope is threaded through the entire length of the colon, provides a very reliable method of detecting colon neoplasia in a subject, but colonoscopy is costly, time consuming and requires sedation of the patient.

[0005] Modern molecular biology has made it possible to identify proteins and nucleic acids that are specifically associated with certain physiological states. These molecular markers have revolutionized diagnostics for a variety of health conditions ranging from pregnancy to viral infections, such as HIV.

[0006] Researchers generally identify molecular markers for a health condition by searching for genes and proteins that are expressed at different levels in one health condition versus another (e.g. in pregnant women versus women who are not

pregnant). Traditional methods for pursuing this research, such as Northern blots and reverse transcriptase polymerase chain reaction, allow a researcher to study only a handful of potential molecular markers at a time. Microarrays, consisting of an ordered array of hundreds or thousands of probes for detection of hundreds or thousands of gene transcripts, allow researchers to gather data on many potential molecular markers in a single experiment. Researchers now face the challenge of sifting through large quantities of microarray-generated gene expression data to identify genes that may be of genuine use as molecular markers to distinguish different health conditions.

[0007] Improved systems for identifying high quality candidate molecular markers in large volumes of gene expression data may help to unlock the power of such tools and increase the likelihood of identifying a molecular marker for important disease states, such as colon neoplasia. Effective molecular markers for colon neoplasia could potentially revolutionize the diagnosis, management and overall health impact of colon cancer.

BRIEF SUMMARY

[0008] This application is based at least in part on the selection of useful molecular markers of colon neoplasia. Colon neoplasia is a multi-stage process involving progression from normal healthy tissues to the development of pre-cancerous colon adenomas to more invasive stages of colon cancer such as the Dukes A and Dukes B stages and finally to metastatic stages such as Dukes C and Dukes D stages of colon cancer.

[0009] In one aspect, this application provides molecular markers that are useful in the detection or diagnosis of colon neoplasia. In certain embodiments, molecular markers described in the application are helpful in distinguishing normal subjects from those who are likely to develop colon neoplasia or are likely to harbor a colon adenoma. In other aspects the invention provides molecular markers that may be useful in distinguishing subjects who are either normal or precancerous from those who have colon cancer. In another embodiment, the application provides markers that help in staging the colon cancer in patients. In still other embodiments the application contemplates the use of one or more of the molecular markers described herein for the detection, diagnosis, and staging of colon neoplasias.

[0010] In one aspect the application provides a method of screening a subject for a condition associated with increased levels of one or more molecular markers that are indicative of colon neoplasia such as for example ColoUp1-ColoUp8 and osteopontin. In a preferred embodiment, the application provides a method for screening a subject for conditions associated with secreted markers such as ColoUp1 or ColoUp2, by detecting in a biological sample an amount of ColoUp1 or ColoUp2 and comparing the amount of ColoUp1 and ColoUp2 found in the subject to one or more of the following: a predetermined standard, the amount of ColoUp1 or ColoUp2 detected in a normal sample from the subject, the subject's historical baseline level of ColoUp1 or ColoUp2, or the ColoUp1 or ColoUp2 level detected in a different, normal subject (a control subject). Detection of a level of ColoUp1 and ColoUp2 in the subject that is greater than that of the predetermined standard or that is increased from a subject's past baseline is indicative of a condition such as colon neoplasia. In certain aspects, an increase in the amount of ColoUp1 or ColoUp2 as compared to the subject's historical baseline would be indicative of a new neoplasm, or progres-

sion of an existing neoplasm. Similarly, a decrease in the amount of ColoUp1 or ColoUp2 as compared to the subject's historical baseline would be indicative of regression on an existing neoplasm

[0011] In one aspect the molecular markers described herein are encoded by a nucleic acid sequence that is at least 90%, 95%, 98%, 99%, 99.3%, 99.5% or 99.7% identical to the nucleic acid sequence of SEQ ID Nos: 4-12, and more preferably to the nucleic acid sequences as set forth in SEQ ID Nos: 4-5. In another aspect, the application provides markers that are encoded by a nucleic acid sequence that hybridizes under high stringency conditions to the nucleic acid sequences of SEQ ID Nos: 4-12, more preferably to the nucleic acid sequences as set forth in SEQ ID Nos: 4-5.

[0012] In another aspect the application provides molecular markers that are diagnostic of colon neoplasia, said markers having an amino acid sequence that is at least 90%, 95%, 98%, 99%, 99.3%, 99.5% or 99.7% identical to the amino acid sequence as set forth in SEQ ID Nos: 1-3 or 13-20, more preferably the amino acid sequence as set forth in SEQ ID Nos: 3 and 14.

[0013] In one aspect, the application provides methods for detecting secreted polypeptide forms of a ColoUp1-ColoUp8 polypeptide or osteopontin in biological samples. In other aspects, the application provides methods for imaging a colon neoplasm by targeting antibodies to any one of the markers ColoUp1 through ColoUp8 described herein, and in preferred embodiments, the antibodies are targeted to ColoUp3. In certain aspects, the application provides methods for administering an imaging agent comprising a targeting moiety and an active moiety. The targeting moiety may be an antibody, Fab, F(Ab)₂, a single chain antibody or other binding agent that interacts with an epitope specified by a polypeptide sequence having an amino acid sequence as set forth in SEQ ID Nos: 1-3 and 13-20. The active moiety may be a radioactive agent, such as radioactive technetium, radioactive indium, or radioactive iodine. The imaging agent is administered in an amount effective for diagnostic use in a mammal such as a human and the localization and accumulation of the imaging agent is then detected. The localization and accumulation of the imaging agent may be detected by radioscintigraphy, nuclear magnetic resonance imaging, computed tomography or positron emission tomography.

[0014] In a preferred embodiment, the application provides methods for detecting a polypeptide comprising an amino acid sequence as set forth in one of SEQ ID Nos: 1-3. As will be apparent to the skilled artisan, the molecular markers described herein may be detected in a number of ways such as by various assays, including antibody-based assays. Examples of antibody-based assays include immunoprecipitation assays, Western blots, radioimmunoassays or enzyme-linked immunosorbent assays (ELISAs). Molecular markers described herein may be detected by assays that do not employ an antibody, such as by methods employing two-dimensional gel electrophoresis, methods employing mass spectroscopy, methods employing suitable enzymatic activity assays, etc. In a preferred embodiment the application provides methods for the detection of secreted markers such as ColoUp1 or ColoUp2 polypeptides in blood, blood fractions (such as blood serum or blood plasma), urine or stool samples. Increased levels of these markers may be associated with a number of conditions such as for example colon neoplasia, including colon adenomas, colon cancer, and metastatic colon cancer. In certain aspects the application provides

methods including the detection of more than one marker that is indicative of colon neoplasia such as methods for detecting both ColoUp1 and ColoUp2. In yet another aspect, combinations of the ColoUp markers may be useful, for instance, a combination of tests including testing biological samples for secreted markers such as ColoUp1 or ColoUp2 in combination with testing for transmembrane markers such as ColoUp3 as targets for imaging agents.

[0015] In yet another aspect, the application provides a method of determining whether a subject is likely to develop colon cancer or is more likely to harbor a precancerous colon adenoma by detecting the presence or absence of the molecular markers as set forth in SEQ ID Nos: 1-3. Detection of combinations of these markers is also helpful in staging the colon neoplasias.

[0016] In yet another aspect, the application provides markers that are useful in distinguishing normal and precancerous subjects from those subjects having colon cancer. In certain embodiments, the application contemplates determining the levels of markers provided herein such as ColoUp1 through ColoUp8 and osteopontin. In one aspect, markers such as ColoUp6 and osteopontin are helpful in distinguishing between the category of patients that are normal or have precancerous colon adenomas and the category of patients having colon cancer. In another aspect, the application provides detection of one or more of said markers in determining the stages of colon neoplasia.

[0017] In certain aspect, the invention provides an immunoassay for determining the presence of any one of the polypeptides having an amino acid sequence as set forth in SEQ ID Nos: 1-3 and 13-20, more preferably any one of the polypeptides having an amino acid sequence as set forth in SEQ ID Nos: 1-3 in a biological sample. The method includes obtaining a biological sample and contacting the sample with an antibody specific for a polypeptide having an amino acid sequence as set forth in SEQ ID Nos: 1-3 and detecting the binding of the antibody.

[0018] In some aspects, the application provides methods for the detection of a molecular marker in a biological sample such as blood, including blood fractions such as serum or plasma. For instance, the blood sample obtained from a patient may be further processed such as by fractionation to obtain blood serum, and the serum may then be enriched for certain polypeptides. The serum so enriched is then contacted with an antibody that is reactive with an epitope of the desired marker polypeptide.

[0019] In yet another embodiment, the application provides methods for determining the appropriate therapeutic protocol for a subject. For example detection of a colon neoplasia provides the treating physician valuable information in determining whether intensive or invasive protocols such as colonoscopy, surgery or chemotherapy would be needed for effective diagnosis or treatment. Such detection would be helpful not only for patients not previously diagnosed with colon neoplasia but also in those cases where a patient has previously received or is currently receiving therapy for colon cancer, the presence or absence or a change in the level of the molecular markers set forth herein may be indicative that the subject is likely to have a relapse or a progressive, or a persistent colon cancer.

[0020] In certain aspects, the application provides molecular markers of colon neoplasia such as ColoUp1 through ColoUp8. In certain instances these markers are secreted proteins such as ColoUp1, ColoUp2 and osteopontin, and are

useful for detecting and diagnosing colon neoplasia. In other aspects, these markers may be transmembrane proteins such as ColoUp3 and may be useful as targets for imaging agents, e.g. as targets to label cells of a neoplasm.

[0021] In one aspect, the application provides isolated, purified or recombinant polypeptides having an amino acid sequence that is at least 90%, 95% or 98-99% identical to an amino acid sequence as set forth in SEQ ID Nos: 1-3 or an amino acid sequence as set forth in SEQ ID Nos: 13-20. In a more preferred embodiment, the application provides an amino acid sequence that is at least 90%, 95%, 98-99%, 99.3%, 99.5% or 99.7% identical to the amino acid sequence as set forth in SEQ ID No: 3 or SEQ ID No: 14. The application also provides fusion proteins comprising the ColoUp proteins described herein fused to a heterologous protein. In certain embodiments, such polypeptides are useful, for example, for generating antibodies or for use in screening assays to identify candidate therapeutics.

[0022] In other aspects the application provides for nucleic acid sequences encoding the polypeptides as set forth in SEQ ID Nos: 1-3 and 13-20. In one aspect the application provides nucleic acids comprising nucleic acid sequences that are at least 90%, 95%, 98-99%, 99.3%, 99.5% or 99.7% identical to the nucleic acid sequence in SEQ ID Nos: 4-12, more preferably 4-5. Also contemplated herein are vectors comprising the nucleic acid sequences set forth in SEQ ID Nos: 4-12, more preferably SEQ ID Nos: 4-5, and host cells expressing the nucleic acid sequences.

[0023] In another aspect, the application provides an antibody that interacts with an epitope specified by one of SEQ ID Nos: 1-3 and 13-20 or portions thereof, more preferably SEQ ID Nos: 1-3 or portions thereof. In a preferred embodiment the antibody is useful for detecting colon adenomas and interacts with an epitope specified by one of SEQ ID Nos: 1-3. In certain aspects the application provides for generating such antibodies, including methods for generating monoclonal and polyclonal antibodies, as well as methods for generating other types of antibodies. In other aspects, the application also provides a hybridoma cell line capable of producing an antibody that interacts with an epitope specified by SEQ ID Nos: 1-3 and 13-20, more preferably SEQ ID Nos: 1-3, or portions thereof. In yet other embodiments, the antibody may be a single chain antibody.

[0024] In yet other embodiments, the application provides a kit for detecting colon neoplasia in a biological sample. Such kits include one or more antibodies that are capable of interacting with an epitope specified by one of SEQ ID Nos: 1-3 and 13-20, more preferably with an epitope specified by one of SEQ ID Nos: 1-3. In more preferred embodiments, the antibodies may be detectably labeled, such as for example with an enzyme, a fluorescent substance, a chemiluminescent substance, a chromophore, a radioactive isotope or a complexing agent.

[0025] In certain embodiments, the application provides the identity of ColoUp1 and ColoUp2 polypeptides that are secreted into the serum in vivo, and that are secreted across the apical and basolateral cell surfaces in cultured intestinal cells. Accordingly, in certain embodiments, the application provides methods for detecting whether a subject is likely to have a colon neoplasia comprising: a) obtaining a biological sample from said subject; and b) detecting one or more polypeptides selected from among: one or more secreted ColoUp1 polypeptides and one or more secreted ColoUp2

polypeptides, wherein the presence of said one or more polypeptides is indicative of colon neoplasia.

[0026] In certain embodiments, a secreted ColoUp2 polypeptide is selected from among: a) a secreted polypeptide produced by the expression of a nucleic acid that is at least 95% identical to the amino acid sequence of SEQ ID No: 5; b) a secreted polypeptide produced by the expression of a nucleic acid that is a naturally occurring variant of SEQ ID No: 5; c) a secreted polypeptide produced by the expression of a nucleic acid that hybridizes under stringent conditions to a nucleic acid sequence of SEQ ID No: 5; d) a secreted polypeptide having a sequence that is at least 95% identical to the amino acid sequence of SEQ ID No: 3; and e) a secreted polypeptide having a sequence that is at least 95% identical to the amino acid sequence of SEQ ID No: 21. Optionally, the secreted ColoUp2 polypeptide is produced by the expression of a nucleic acid having the sequence of SEQ ID No: 5, and preferably the secreted ColoUp2 polypeptide is produced by the expression of a nucleic acid sequence that is at least 98%, 99% or 100% identical to the nucleic acid sequence of SEQ ID No: 5. In certain embodiments, the secreted ColoUp2 polypeptide has an amino acid sequence that is at least 98%, 99% or 100% identical to an amino acid sequence selected from among SEQ ID No: 3 and SEQ ID No: 21. In certain embodiments, the secreted ColoUp1 polypeptide is selected from among: a) a secreted polypeptide produced by the expression of a nucleic acid that is at least 95% identical to the amino acid sequence of SEQ ID No: 4; b) a secreted polypeptide produced by the expression of a nucleic acid that is a naturally occurring variant of SEQ ID No: 4; c) a secreted polypeptide produced by the expression of a nucleic acid that hybridizes under stringent conditions to a nucleic acid sequence of SEQ ID No: 4; d) a secreted polypeptide having a sequence that is at least 95% identical to the amino acid sequence of SEQ ID No: 1; and e) a secreted polypeptide having a sequence that is at least 95% identical to the amino acid sequence of SEQ ID No: 2. Optionally, the secreted ColoUp1 polypeptide is produced by the expression of a nucleic acid having a sequence that is at least 95%, 98, 99% or 100% identical to the nucleic acid sequence of SEQ ID No: 4. Preferably, the secreted ColoUp1 polypeptide has an amino acid sequence that is at least 95%, 98%, 99% or 100% identical to an amino acid sequence selected from among SEQ ID No: 1 and SEQ ID No: 2. Optionally, for detection of basolaterally secreted ColoUp1 or ColoUp2 polypeptides, the biological sample is a blood sample or a fraction derived from blood, such as serum, plasma, cells, or a fraction enriched for apically secreted ColoUp1 or ColoUp2 polypeptide. Optionally, for detection of basolaterally secreted ColoUp1 or ColoUp2 polypeptides, the biological sample is a urine sample or a fraction derived from urine. Optionally, for detection of apically secreted ColoUp1 or ColoUp2 polypeptides, the biological sample is derived from the inner wall and/or lumen of the intestinal tract, such as intestinal mucous or other fluid, excreted stool and stool removed from within the colon. In certain embodiments, the polypeptide is detected by an assay that employs an antibody, such as an immunoprecipitation assay, a Western blot, a radioimmunoassay or an enzyme-linked immunosorbent assay (ELISA). Optionally, an assay comprises contacting the biological sample with an antibody that interacts with a secreted ColoUp1 polypeptide or a secreted ColoUp2 polypeptide. An antibody may, for example, interact with an epitope of an amino acid sequence selected from among: SEQ ID No: 1 and SEQ ID No: 2. An

antibody may, for example, interact with an epitope of an amino acid sequence selected from among: SEQ ID No: 3 and SEQ ID No: 21. Optionally, the antibody is detectably labeled, such as with an enzyme, a fluorescent substance, a chemiluminescent substance, a chromophore, a radioactive isotope or a complexing agent. Optionally, the amount of at least one secreted ColoUp1 polypeptide and/or at least one secreted ColoUp2 polypeptide in the biological sample is compared to a predetermined standard (e.g., a known amount of purified ColoUp1 or ColoUp2 polypeptide). Optionally, the amount of at least one secreted ColoUp1 polypeptide and/or at least one secreted ColoUp2 polypeptide in the biological sample is compared to the subject's historical baseline. In certain embodiments, the presence of at least one secreted ColoUp1 polypeptide and/or at least one secreted ColoUp2 polypeptide is indicative that the subject is likely to harbor a colon adenoma or a colon cancer. In certain embodiments, the presence of at least one secreted ColoUp1 polypeptide and/or at least one secreted ColoUp2 polypeptide may be used in determining the therapeutic protocol to be administered to a subject having a colon neoplasia, and the subject may not have been previously diagnosed with colon cancer or the subject may have previously received or is currently receiving a therapy for colon cancer, wherein the presence of at least one secreted ColoUp1 polypeptide and/or at least one secreted ColoUp2 polypeptide indicates that the subject is likely to have a relapse or a persistent or progressive colon cancer. The detection of said secreted polypeptide may indicate the presence of a variety of neoplasias in a subject, such as a colon adenoma, a colon cancer and a metastatic colon cancer. Optionally, a method involves detecting both at least one secreted ColoUp1 polypeptide and at least one secreted ColoUp2 polypeptide in the biological sample.

[0027] In certain embodiments, the application provides kits for detecting one or more molecular markers of colon neoplasia in a biological sample. A kit may comprise a) an antibody which interacts with an epitope of a secreted ColoUp1 polypeptide or a secreted ColoUp2 polypeptide; and b) instructions for use. Optionally, the antibody interacts with an epitope of a polypeptide selected from among: the polypeptide of SEQ ID No:1, the polypeptide of SEQ ID No:2, the polypeptide of SEQ ID No:3 and the polypeptide of SEQ ID No:21. Optionally, the antibody is detectably labeled.

[0028] In certain embodiments, the application provides a novel purified polypeptide, which is a portion of ColoUp2 that is found in serum. Such a polypeptide may consist essentially of an amino acid sequence that is at least 95%, 98%, 99% or 100% identical to the sequence of SEQ ID No: 21. By "consisting essentially" is meant that there may be, in addition to the indicated amino acid sequence, a variety of modifications, such as phosphorylations, glycosylations, disulfide bonds, unusual or modified amino acids, etc.

[0029] In certain embodiments, the application provides novel fusion proteins comprising a first polypeptide domain and a second polypeptide domain, wherein the first polypeptide domain consists essentially of an amino acid sequence that is at least 95%, 98%, 99% or 100% identical to an amino acid sequence of SEQ ID No. 21. The second polypeptide domain may be a domain selected from the group consisting of: a detection domain, a purification domain and an antigenic domain.

[0030] In certain embodiments, the application provides antibodies that bind specifically to a ColoUp2 polypeptide consisting essentially of the amino acid sequence of SEQ ID

No: 21. The antibody may binds the ColoUp2 polypeptide with a dissociation constant of less than $10^{-6}M$, $10^{-7}M$, $10^{-8}M$ or $10^{-9}M$. The antibody may be essentially any type of antibody, including polyclonal, monoclonal, and single chain antibodies, or other fragments. For diagnostic use, there may be little benefit to having a humanized antibody, however, humanized antibodies are highly desirable for therapeutic uses. Preferably, a diagnostic antibody is effective for detecting the ColoUp2 polypeptide in a biological sample, such as a blood, stool or urine sample, or a fraction thereof. Optionally, the antibody is effective for detecting the ColoUp2 polypeptide in a sample comprising cells from a colon neoplasia. The application further provides methods for making such antibodies in a variety of ways. For example, a monoclonal antibody may be produced in a method comprising: (a) administering to a mouse an amount of an immunogenic composition comprising the ColoUp2 polypeptide effective to stimulate a detectable immune response; (b) obtaining antibody-producing cells from the mouse and fusing the antibody-producing cells with myeloma cells to obtain antibody-producing hybridomas; (c) testing the antibody-producing hybridomas to identify a preferred hybridoma, wherein the preferred hybridoma is a hybridoma that produces a monoclonal antibody that binds specifically to the ColoUp2 polypeptide; (d) culturing the preferred hybridoma cell culture that produces the monoclonal antibody that binds specifically to the ColoUp2 polypeptide; and (e) obtaining the monoclonal antibody that binds specifically to the ColoUp2 polypeptide from the cell culture. Optionally, the antibody-producing hybridomas comprises testing whether the antibody-producing hybridomas produce an antibody that binds to the ColoUp2 polypeptide in an assay selected from the group consisting of: an enzyme-linked immunosorbent assay, a Bia-core assay and an immunoprecipitation assay.

[0031] The embodiments and practices of the present invention, other embodiments, and their features and characteristics, will be apparent from the description, figures and claims that follow, with all of the claims hereby being incorporated by this reference into this Summary.

BRIEF DESCRIPTION OF THE DRAWINGS

[0032] FIG. 1 shows the amino acid sequences (SEQ ID NOs: 1 and 2) of secreted ColoUp1 protein. A. An N-terminal signal peptide is cleaved between amino acids 30-31 of the full-length ColoUp1 protein; B. An N-terminal signal peptide is cleaved between amino acids 33-34 of the full-length ColoUp1 protein.

[0033] FIG. 2 shows the amino acid sequence (SEQ ID NO: 3) of secreted ColoUp2 protein.

[0034] FIG. 3 shows the nucleic acid sequence (SEQ ID NO: 4) of ColoUp1.

[0035] FIG. 4 shows the nucleic acid sequence (SEQ ID NO: 5) of ColoUp2.

[0036] FIG. 5 shows the nucleic acid sequence (SEQ ID NO: 6) of Osteopontin.

[0037] FIG. 6 shows the nucleic acid sequence (SEQ ID NO: 7) of ColoUp3.

[0038] FIG. 7 shows the nucleic acid sequence (SEQ ID NO: 8) of ColoUp4.

[0039] FIG. 8 shows the nucleic acid sequence (SEQ ID NO: 9) of ColoUp5.

[0040] FIG. 9 shows the nucleic acid sequence (SEQ ID NO: 10) of ColoUp6.

[0041] FIG. 10 shows the nucleic acid sequence (SEQ ID NO: 11) of ColoUp7.

[0042] FIG. 11 shows the nucleic acid sequence (SEQ ID NO: 12) of ColoUp8.

[0043] FIG. 12 shows the amino acid sequence (SEQ ID NO: 13) of full-length ColoUp1 protein.

[0044] FIG. 13 shows the amino acid sequence (SEQ ID NO: 14) of full-length ColoUp2 protein.

[0045] FIG. 14 shows the amino acid sequence (SEQ ID NO: 15) of full-length Osteopontin protein.

[0046] FIG. 15 shows the amino acid sequence (SEQ ID NO: 16) of full-length ColoUp3 protein.

[0047] FIG. 16 shows the amino acid sequence (SEQ ID NO: 17) of full-length ColoUp4 protein.

[0048] FIG. 17 shows the amino acid sequence (SEQ ID NO: 18) of full-length ColoUp5 protein.

[0049] FIG. 18 shows the amino acid sequence (SEQ ID NO: 19) of full-length ColoUp6 protein.

[0050] FIG. 19 shows the amino acid sequence (SEQ ID NO: 20) of full-length ColoUp8 protein.

[0051] FIG. 20 is a graphical display of ColoUp1 expression levels measured by micro-array profiling in different samples. A. In normal colon epithelial strips, normal liver, and colonic muscle; B. In premalignant colon adenomas as well as in colon cancers of Dukes stages B, Dukes stage C, and Duke stages D; C. In colon cancer liver metastasis; D. In colon cancer cell lines, colon cancer xenografts grown in athymic mice, MSI cell lines, and V330 cell lines treated with TGF β .

[0052] FIG. 21 is a graphical display of ColoUp2 expression levels measured by micro-array profiling in different samples. A. In normal colon epithelial strips, normal liver, and colonic muscle; B. In premalignant colon adenomas as well as in colon cancers of Dukes stages B, Dukes stage C, and Duke stages D; C. In colon cancer liver metastasis; D. In colon cancer cell lines, colon cancer xenografts grown in athymic mice, MSI cell lines, and V330 cell lines treated with TGF β .

[0053] FIG. 22 is a graphical display of Osteopontin expression levels measured by micro-array profiling in different samples. A. In normal colon epithelial strips, normal liver, and colonic muscle; B. In premalignant colon adenomas as well as in colon cancers of Dukes stages B, Dukes stage C, and Duke stages D; C. In colon cancer liver metastasis; D. In colon cancer cell lines, colon cancer xenografts grown in athymic mice, MSI cell lines, and V330 cell lines treated with TGF β .

[0054] FIG. 23 is a graphical display of ColoUp3 expression levels measured by micro-array profiling in different samples. A. In normal colon epithelial strips, normal liver, and colonic muscle; B. In premalignant colon adenomas as well as in colon cancers of Dukes stages B, Dukes stage C, and Duke stages D; C. In colon cancer liver metastasis; D. In colon cancer cell lines, colon cancer xenografts grown in athymic mice, MSI cell lines, and V330 cell lines treated with TGF β .

[0055] FIG. 24 is a graphical display of ColoUp4 expression levels measured by micro-array profiling in different samples. A. In normal colon epithelial strips, normal liver, and colonic muscle; B. In premalignant colon adenomas as well as in colon cancers of Dukes stages B, Dukes stage C, and Duke stages D; C. In colon cancer liver metastasis; D. In

colon cancer cell lines, colon cancer xenografts grown in athymic mice, MSI cell lines, and V330 cell lines treated with TGF β .

[0056] FIG. 25 is a graphical display of ColoUp5 expression levels measured by micro-array profiling in different samples. A. In normal colon epithelial strips, normal liver, and colonic muscle; B. In premalignant colon adenomas as well as in colon cancers of Dukes stages B, Dukes stage C, and Duke stages D; C. In colon cancer liver metastasis; D. In colon cancer cell lines, colon cancer xenografts grown in athymic mice, MSI cell lines, and V330 cell lines treated with TGF β .

[0057] FIG. 26 is a graphical display of ColoUp6 expression levels measured by micro-array profiling in different samples. A. In normal colon epithelial strips, normal liver, and colonic muscle; B. In premalignant colon adenomas as well as in colon cancers of Dukes stages B, Dukes stage C, and Duke stages D; C. In colon cancer liver metastasis; D. In colon cancer cell lines, colon cancer xenografts grown in athymic mice, MSI cell lines, and V330 cell lines treated with TGF β .

[0058] FIG. 27 is a graphical display of ColoUp7 expression levels measured by micro-array profiling in different samples. A. In normal colon epithelial strips, normal liver, and colonic muscle; B. In premalignant colon adenomas as well as in colon cancers of Dukes stages B, Dukes stage C, and Duke stages D; C. In colon cancer liver metastasis; D. In colon cancer cell lines, colon cancer xenografts grown in athymic mice, MSI cell lines, and V330 cell lines treated with TGF β .

[0059] FIG. 28 is a graphical display of ColoUp8 expression levels measured by micro-array profiling in different samples. A. In normal colon epithelial strips, normal liver, and colonic muscle; B. In premalignant colon adenomas as well as in colon cancers of Dukes stages B, Dukes stage C, and Duke stages D; C. In colon cancer liver metastasis; D. In colon cancer cell lines, colon cancer xenografts grown in athymic mice, MSI cell lines, and V330 cell lines treated with TGF β .

[0060] FIG. 29 shows northern blot analysis of ColoUp1 mRNA levels in normal colon tissues and colon cancer cell lines or tissues. A. In normal colon tissue samples and a group of colon cancer cell lines; B. and C. In normal colon tissues and colon neoplasms from 15 individuals with colon cancers and one individual with a colon adenoma.

[0061] FIG. 30 shows detection of T7 epitope-tagged ColoUp1 protein levels in transfected FET cells and Vaco400 cells. A. Secretion of epitope-tagged ColoUp1 protein in V400 cell growth media by Western blot ("T" are transfectants with an epitope tagged ColoUp1 expression vector; "C" are transfectants with an empty control vector); B. Expression of T7 epitope-tagged ColoUp1 protein in transfected FET cells and V400 cells by Western blot (left panel), and secretion of epitope-tagged ColoUp1 protein in growth media by serial immunoprecipitation and Western blot (right panel) (Cell extract amounts loaded: FET=75 mg/well; V400=31.1 mg/well; Volume of media used for immuno-precipitation=1 ml of 20 ml).

[0062] FIG. 31 shows northern blot analysis of ColoUp2 mRNA levels in normal colon tissue samples and a group of colon cancer cell lines (top panel). The bottom panel shows the ethidium bromide stained gel corresponding to the blot.

[0063] FIG. 32 shows detection of V5 epitope-tagged ColoUp2 protein levels in transfected SW480 cells and

Vaco400 cells (24 hours and 48 hours after transfection). Expression of epitope-tagged ColoUp2 protein in transfected cells by Western blot (right panel), and secretion of epitope-tagged ColoUp2 protein in growth media by serial immunoprecipitation and Western blot (left panel).

[0064] FIG. 33 shows two northern blot analysis of ColoUp5 mRNA levels in normal colon tissues and a group of colon cancer cell lines (top panels). The bottom panels show the ethidium bromide stained gel corresponding to the blot.

[0065] FIG. 34 illustrates an alignment of the human, mouse, and rat ColoUp5 (FoxQ1) amino acid sequences.

[0066] FIG. 35 illustrates an alignment of the human, mouse, and rat ColoUp5 (FoxQ1) nucleic acid sequences.

[0067] FIG. 36 shows a western blot of V5 tagged ColoUp2 protein detected by anti-V5 antibody. Lane 1: media supernate from SW480 colon cancer cells transfected with an empty expression vector. Lane 2: media supernate from ColoUp2-V5 expressing cells. Lane 3: size markers. Lane 4 shows assay of serum from a mouse xenografted with control SW480 cells corresponding to lane 1. Lanes 5 and 6 show detection of circulating ColoUp2 proteins in blood from two mice bearing human colon cancer xenografts from ColoUp2-V5 expressing SW480 colon cells shown in lane 2. ColoUp2 is secreted as an 85 KD and a companion 55 KD size protein.

[0068] FIG. 37 shows a western blot with anti-V5 antibody of V5 tagged ColoUp1 protein. Lane 1: media supernate from SW480 colon cancer cells transfected with an empty expression vector. Lane 2: media supernate from ColoUp1-V5 expressing SW480 cells. Lane 3 shows assay of serum from a mouse xenografted with control SW480 cells corresponding to lane 1. Lanes 4 shows detection of circulating ColoUp1 proteins in blood from a mouse bearing tumor xenografts from ColoUp1-V5 expressing SW480 cells shown in lane 2. Lane 5: size markers.

[0069] FIG. 38 shows, in the upper panel, the purification of ColoUp2 protein. Shown is a Coomassie blue staining of 250 ng (lane 2a) and 500 ng (lane 3a) of a purified ColoUp2 protein preparation. Size markers are in lane 1a. In the lower panel is shown a Coomassie blue stained gel showing purification of His-tagged ColoUp1 protein on Ni-NTA beads. Lane 1: markers, Lane 2 media from mock transfected cells, Lane 3 purification of media from ColoUp1 transfected cells. Clearly shown is purification to homogeneity of the 180 kd ColoUp protein.

[0070] FIG. 39 shows, in the top panel, detection on an anti-V5 western of V5-tagged ColoUp2 protein. Lane 1: media from mock transfected Caco2 cells. Lane 2: detection of secreted ColoUp2 protein from transiently transfected Caco2 cells grown in standard culture dishes. Seen are the typical 85 KD and 55 KD secreted bands (the lane is heavily overloaded and minor degradation products are also visualized). Lane 3: molecular weight markers. Lanes 4-7: detection of ColoUp2 secreted into the basolateral compartment (lower chamber) of transiently transfected Caco2 grown as a monolayer on a transwell filter. Lanes 9-12 show the general absence of ColoUp2 in the corresponding apical compartment, with the exception of the 48 hour time point. The table shows the electrical resistance and transfection efficiency (gfp expression) measured at each time point. A dip in the electrical resistance at 48 hours suggests some leakiness of the monolayer at that time point.

[0071] FIG. 40: Top panel shows detection on anti-V5 western of V5-tagged ColoUp1 protein. Control lane shows detection of purified recombinant ColoUp1. Identical bands

are seen in media harvested on days 1-4 (lanes D1-D4) from both apical and basolateral compartments. The table shows the electrical resistance and transfection efficiency (gfp expression) measured at each time point.

[0072] FIG. 41 shows the amino acid sequence of the approximately 55 kDa C-terminal fragment of ColoUp2 that is a prominent secreted and serum form of ColoUp2.

DETAILED DESCRIPTION

1. Definitions:

[0073] For convenience, certain terms employed in the specification, examples, and appended claims are collected here. Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs.

[0074] The articles “a” and “an” are used herein to refer to one or to more than one (i.e., to at least one) of the grammatical object of the article. By way of example, “an element” means one element or more than one element.

[0075] The terms “adenoma”, “colon adenoma” and “polyp” are used herein to describe any precancerous neoplasia of the colon.

[0076] The term “antibody” as used herein is intended to include whole antibodies, e.g., of any isotype (IgG, IgA, IgM, IgE, etc), and includes fragments thereof which are also specifically reactive with a vertebrate, e.g., mammalian, protein. Antibodies can be fragmented using conventional techniques and the fragments screened for utility and/or interaction with a specific epitope of interest. Thus, the term includes segments of proteolytically-cleaved or recombinantly-prepared portions of an antibody molecule that are capable of selectively reacting with a certain protein. Non-limiting examples of such proteolytic and/or recombinant fragments include Fab, F(ab')₂, Fab', Fv, and single chain antibodies (scFv) containing a V[L] and/or V[H] domain joined by a peptide linker. The scFv's may be covalently or non-covalently linked to form antibodies having two or more binding sites. The term antibody also includes polyclonal, monoclonal, or other purified preparations of antibodies and recombinant antibodies.

[0077] The term “colon” as used herein is intended to encompass the right colon (including the cecum), the transverse colon, the left colon and the rectum.

[0078] The terms “colorectal cancer” and “colon cancer” are used interchangeably herein to refer to any cancerous neoplasia of the colon (including the rectum, as defined above).

[0079] The term “ColoUpX” (e.g. ColoUp1, ColoUp2 . . . ColoUp8) is used to refer to a nucleic acid encoding a ColoUp protein or a ColoUp protein itself, as well as distinguishable fragments of such nucleic acids and proteins, longer nucleic acids and polypeptides that comprise distinguishable fragments or full length nucleic acids or polypeptides, and variants thereof. Variants include polypeptides that are at least 90% identical to the relevant human ColoUp SEQ ID Nos. referred to in the application, and nucleic acids encoding such variant polypeptides. In addition, variants include different post-translational modifications, such as glycosylations, methylations, etc. Particularly preferred variants include any naturally occurring variants, such as allelic differences, mutations that occur in a neoplasia and secreted or processed forms. The terms “variants” and “fragments” are overlapping.

[0080] As used herein, the phrase “gene expression” or “protein expression” includes any information pertaining to the amount of gene transcript or protein present in a sample, as well as information about the rate at which genes or proteins are produced or are accumulating or being degraded (eg. reporter gene data, data from nuclear runoff experiments, pulse-chase data etc.). Certain kinds of data might be viewed as relating to both gene and protein expression. For example, protein levels in a cell are reflective of the level of protein as well as the level of transcription, and such data is intended to be included by the phrase “gene or protein expression information”. Such information may be given in the form of amounts per cell, amounts relative to a control gene or protein, in unitless measures, etc.; the term “information” is not to be limited to any particular means of representation and is intended to mean any representation that provides relevant information. The term “expression levels” refers to a quantity reflected in or derivable from the gene or protein expression data, whether the data is directed to gene transcript accumulation or protein accumulation or protein synthesis rates, etc.

[0081] The term “detection” is used herein to refer to any process of observing a marker, in a biological sample, whether or not the marker is actually detected. In other words, the act of probing a sample for a marker is a “detection” even if the marker is determined to be not present or below the level of sensitivity. Detection may be a quantitative, semi-quantitative or non-quantitative observation.

[0082] The terms “healthy”, “normal” and “non-neoplastic” are used interchangeably herein to refer to a subject or particular cell or tissue that is devoid (at least to the limit of detection) of a disease condition, such as a neoplasia, that is associated with increased expression of a ColoUp gene. These terms are often used herein in reference to tissues and cells of the colon. Thus, for the purposes of this application, a patient with severe heart disease but lacking a ColoUp-associated disease would be termed “healthy”.

[0083] The term “including” is used herein to mean, and is used interchangeably with, the phrase “including but not limited to”.

[0084] As used herein, the term “nucleic acid” refers to polynucleotides such as deoxyribonucleic acid (DNA), and, where appropriate, ribonucleic acid (RNA). The term should also be understood to include analogs of either RNA or DNA made from nucleotide analogs, and, as applicable to the embodiment being described, single-stranded (such as sense or antisense) and double-stranded polynucleotides.

[0085] The term “or” is used herein to mean, and is used interchangeably with, the term “and/or”, unless context clearly indicates otherwise.

[0086] The term “percent identical” refers to sequence identity between two amino acid sequences or between two nucleotide sequences. Identity can each be determined by comparing a position in each sequence which may be aligned for purposes of comparison. When an equivalent position in the compared sequences is occupied by the same base or amino acid, then the molecules are identical at that position; when the equivalent site occupied by the same or a similar amino acid residue (e.g., similar in steric and/or electronic nature), then the molecules can be referred to as homologous (similar) at that position. Expression as a percentage of homology/similarity or identity refers to a function of the number of identical or similar amino acids at positions shared by the compared sequences. Various alignment algorithms and/or programs may be used, including FASTA, BLAST or

ENTREZ. FASTA and BLAST are available as a part of the GCG sequence analysis package (University of Wisconsin, Madison, Wis.), and can be used with, e.g., default settings. ENTREZ is available through the National Center for Biotechnology Information, National Library of Medicine, National Institutes of Health, Bethesda, Md. In one embodiment, the percent identity of two sequences can be determined by the GCG program with a gap weight of 1, e.g., each amino acid gap is weighted as if it were a single amino acid or nucleotide mismatch between the two sequences.

[0087] The terms “polypeptide” and “protein” are used interchangeably herein.

[0088] The term “purified protein” refers to a preparation of a protein or proteins which are preferably isolated from, or otherwise substantially free of, other proteins normally associated with the protein(s) in a cell or cell lysate. The term “substantially free of other cellular proteins” (also referred to herein as “substantially free of other contaminating proteins”) is defined as encompassing individual preparations of each of the component proteins comprising less than 20% (by dry weight) contaminating protein, and preferably comprises less than 5% contaminating protein. Functional forms of each of the component proteins can be prepared as purified preparations by using a cloned gene as described in the attached examples. By “purified”, it is meant, when referring to component protein preparations used to generate a reconstituted protein mixture, that the indicated molecule is present in the substantial absence of other biological macromolecules, such as other proteins (particularly other proteins which may substantially mask, diminish, confuse or alter the characteristics of the component proteins either as purified preparations or in their function in the subject reconstituted mixture). The term “purified” as used herein preferably means at least 80% by dry weight, more preferably in the range of 85% by weight, more preferably 95-99% by weight, and most preferably at least 99.8% by weight, of biological macromolecules of the same type present (but water, buffers, and other small molecules, especially molecules having a molecular weight of less than 5000, can be present). The term “pure” as used herein preferably has the same numerical limits as “purified” immediately above.

[0089] A “recombinant nucleic acid” is any nucleic acid that has been placed adjacent to another nucleic acid by recombinant DNA techniques. A “recombinant nucleic acid” also includes any nucleic acid that has been placed next to a second nucleic acid by a laboratory genetic technique such as, for example, transformation and integration, transposon hopping or viral insertion. In general, a recombinant nucleic acid is not naturally located adjacent to the second nucleic acid.

[0090] The term “recombinant protein” refers to a protein that is produced by expression from a recombinant nucleic acid.

[0091] A “sample” includes any material that is obtained or prepared for detection of a molecular marker, or any material that is contacted with a detection reagent or detection device for the purpose of detecting a molecular marker.

[0092] A “subject” is any organism of interest, generally a mammalian subject, such as a mouse, and preferably a human subject.

2. Overview

[0093] In certain aspects, the invention relates to methods for determining whether a subject is likely or unlikely to have a colon neoplasia. In other aspects, the invention relates to

methods for determining whether a patient is likely or unlikely to have a colon cancer. In further aspects, the invention relates to methods for monitoring colon neoplasia in a subject. In further aspects, the invention relates to methods for staging a subject's colon neoplasia. A colon neoplasia is any cancerous or precancerous growth located in, or derived from, the colon. The colon is a portion of the intestinal tract that is roughly three feet in length, stretching from the end of the small intestine to the rectum. Viewed in cross section, the colon consists of four distinguishable layers arranged in concentric rings surrounding an interior space, termed the lumen, through which digested materials pass. In order, moving outward from the lumen, the layers are termed the mucosa, the submucosa, the muscularis propria and the subserosa. The mucosa includes the epithelial layer (cells adjacent to the lumen), the basement membrane, the lamina propria and the muscularis mucosae. In general, the "wall" of the colon is intended to refer to the submucosa and the layers outside of the submucosa. The "lining" is the mucosa.

[0094] Precancerous colon neoplasias are referred to as adenomas or adenomatous polyps. Adenomas are typically small mushroom-like or wart-like growths on the lining of the colon and do not invade into the wall of the colon. Adenomas may be visualized through a device such as a colonoscope or flexible sigmoidoscope. Several studies have shown that patients who undergo screening for and removal of adenomas have a decreased rate of mortality from colon cancer. For this and other reasons, it is generally accepted that adenomas are an obligate precursor for the vast majority of colon cancers.

[0095] When a colon neoplasia invades into the basement membrane of the colon, it is considered a colon cancer, as the term "colon cancer" is used herein. In describing colon cancers, this specification will generally follow the so-called "Dukes" colon cancer staging system. Other staging systems have been devised, and the particular system selected is, for the purposes of this disclosure, unimportant. The characteristics that describe a cancer are of greater significance than the particular term used to describe a recognizable stage. The most widely used staging systems generally use at least one of the following characteristics for staging: the extent of tumor penetration into the colon wall, with greater penetration generally correlating with a more dangerous tumor; the extent of invasion of the tumor through the colon wall and into other neighboring tissues, with greater invasion generally correlating with a more dangerous tumor; the extent of invasion of the tumor into the regional lymph nodes, with greater invasion generally correlating with a more dangerous tumor; and the extent of metastatic invasion into more distant tissues, such as the liver, with greater metastatic invasion generally correlating with a more dangerous disease state.

[0096] "Dukes A" and "Dukes B" colon cancers are neoplasias that have invaded into the wall of the colon but have not spread into other tissues. Dukes A colon cancers are cancers that have not invaded beyond the submucosa. Dukes B colon cancers are subdivided into two groups: "Dukes B1" and "Dukes B2". "Dukes B1" colon cancers are neoplasias that have invaded up to but not through the muscularis propria. Dukes B2 colon cancers are cancers that have breached completely through the muscularis propria. Over a five year period, patients with Dukes A cancer who receive surgical treatment (i.e. removal of the affected tissue) have a greater than 90% survival rate. Over the same period, patients with Dukes B1 and Dukes B2 cancer receiving surgical treatment

have a survival rate of about 85% and 75%, respectively. Dukes A, B1 and B2 cancers are also referred to as T1, T2 and T3-T4 cancers, respectively.

[0097] "Dukes C" colon cancers are cancers that have spread to the regional lymph nodes, such as the lymph nodes of the gut. Patients with Dukes C cancer who receive surgical treatment alone have a 35% survival rate over a five year period, but this survival rate is increased to 60% in patients that receive chemotherapy.

[0098] "Dukes D" colon cancers are cancers that have metastasized to other organs. The liver is the most common organ in which metastatic colon cancer is found. Patients with Dukes D colon cancer have a survival rate of less than 5% over a five year period, regardless of the treatment regimen.

[0099] As noted above, early detection of colon neoplasia, coupled with appropriate intervention, is important for increasing patient survival rates. Present systems for screening for colon neoplasia are deficient for a variety of reasons, including a lack of specificity or sensitivity (e.g. Fecal Occult Blood Test, flexible sigmoidoscopy) or a high cost and intensive use of medical resources (e.g. colonoscopy). Alternative systems for detection of colon neoplasia would be useful in a wide range of other clinical circumstances as well. For example, patients who receive surgical or pharmaceutical therapy for colon cancer may experience a relapse. It would be advantageous to have an alternative system for determining whether such patients have a recurrent or relapsed colon neoplasia. As a further example, an alternative diagnostic system would facilitate monitoring an increase, decrease or persistence of colon neoplasia in a patient known to have a colon neoplasia. A patient undergoing chemotherapy may be monitored to assess the effectiveness of the therapy.

[0100] Accordingly, in certain embodiments, the invention provides molecular markers that distinguish between cells that are not part of a colon neoplasia, referred to herein as "healthy cells", and cells that are part of a colon neoplasia (e.g. an adenoma or a colon cancer), referred to herein as "colon neoplasia cells". Certain molecular markers of the invention, including ColoUp1 and ColoUp2, are expressed at significantly higher levels in adenomas, Dukes A, Dukes B1, Dukes B2 and metastatic colon cancer of the liver (liver metastases) than in healthy colon tissue, healthy liver or healthy colon muscle. Certain molecular markers, including ColoUp1 and ColoUp2 are expressed at significantly higher levels in cell lines derived from colon cancer or cell lines engineered to imitate an aspect of a colon cancer cell. Particularly preferred molecular markers of the invention are markers that distinguish between healthy cells and cells of an adenoma. While not wishing to be bound to theory, it is contemplated that because adenomas are thought to be an obligate precursor for greater than 90% of colon cancers, markers that distinguish between healthy cells and cells of an adenoma are particularly valuable for screening apparently healthy patients to determine whether the patient is at increased risk for (predisposed to) developing a colon cancer. Furthermore, particularly preferred molecular markers are those that are actually present in the serum of an animal having a colon neoplasia, and in general, a secreted protein will generally occur in the serum only if it is secreted from a cell contacting a blood vessel, or a compartment in diffusional contact with a blood vessel. For example, protein secreted from a large or advanced colon cancer will generally be found in the blood stream, but a protein secreted from a colon adenoma may not be present in the blood unless it is secreted

from the basolateral face of the cell. Molecular markers that occur in the urine are generally derived from a polypeptide that is present in the blood. Optionally, a molecular marker is one that is present in the lumen of the colon (e.g., may be found in the intestinal mucous or in stool samples), and such a marker will generally be one that is secreted from the apical face of a cell.

[0101] In certain embodiments, the invention provides methods for using ColoUp molecular markers for determining whether a patient has or does not have a condition characterized by increased expression of one or more ColoUp nucleic acids or proteins described herein. In certain embodiments, the invention provides methods for determining whether a patient is or is not likely to have a colon neoplasia. In further embodiments, the invention provides methods for determining whether the patient is having a relapse or determining whether a patient's colon neoplasia is responding to treatment.

[0102] 3. Methods for Identifying Candidate Molecular Markers for Colon Neoplasia

[0103] In certain aspects, the invention relates to the observation that when gene expression data is analyzed using carefully selected criteria, the likelihood of identifying strong candidate molecular markers of a colon neoplasia is quite high. Accordingly, in certain embodiments, the invention provides methods and criteria for analyzing gene expression data to identify candidate molecular markers for colon neoplasia. Although methods and criteria of the invention may be applied to essentially any relevant gene expression data, the benefits of using the inventive methods and criteria are readily apparent when applied to the copious data produced by highly parallel gene expression measurement systems, such as microarray systems. The human genome is estimated to be capable of producing roughly 20,000 to 100,000 different gene transcripts, thousands of which may show a change in expression level in healthy cells versus colon neoplasia cells. It is relatively cost-effective to obtain large quantities of gene expression data and to use this data to identify thousands of candidate molecular markers. However, a significant amount of labor intensive experimentation is generally needed to move from the identification of a candidate molecular marker to an effective diagnostic test for a health condition of interest. In fact, as of the time of filing of this application, the resources required to generate a diagnostic test from a single candidate molecular marker identified by gene expression data are large enough that it is essentially impossible to extract commercially valuable and clinically useful diagnostics from a list of hundreds or thousands of genes whose expression levels change in a particular situation. Accordingly, there is a substantial practical value in being able to select a small number (e.g. ten or fewer) of high-quality molecular markers for further study.

[0104] In certain embodiments, candidate molecular markers for colon neoplasia may be selected by comparing gene expression in liver metastatic colon cancer samples ("liver mets"), normal (non-neoplastic) colon samples and normal liver samples. In this embodiment, candidate molecular markers are those genes (and their gene products) that have a level of expression in liver mets (assessed as a median expression level across the sample set) that is at least four times greater than the level of expression in normal colon samples (also assessed as a median expression level across the sample set). Furthermore, in this embodiment, the median level of expression in liver mets should be greater than the median

level of expression in normal liver samples. The criteria employed in this embodiment provide a high threshold to eliminate most lower quality markers and further eliminate contaminants from liver tissue.

[0105] In certain embodiments, candidate molecular markers for colon neoplasia may be selected by comparing gene expression in normal colon to gene expression in a plurality of different cell lines cultured from metastatic colon cancer samples. For example median metastatic colon cancer cell line gene expression may be calculated as the median of 8 colon cancer cell lines of the Vaco colon cancer cell line series (Markowitz, S. et al. *Science*. 268: 1336-1338, 1995), such as the following liver metastases-derived cell lines: V394, V576, V241, V9M, V400, V10M, V503, V786. In embodiments employing this criterion, candidate molecular markers are those genes (and their gene products) that have at least a three-fold higher median level of expression across the cell lines tested than in the normal colon tissue.

[0106] In certain embodiments, candidate molecular markers for colon neoplasia may be selected by comparing gene expression in normal colon to gene expression in a plurality of colon cancer xenografts grown in athymic mice ("xenografts"). In embodiments employing this criterion, candidate molecular markers are those genes (and their gene products) that have at least a four-fold higher median level of expression across the xenografts tested than in the normal colon tissue.

[0107] In certain embodiments, candidate molecular markers for colon neoplasia may be selected by comparing maximum gene expression in normal colon to minimum gene expression in liver mets. In these embodiments, candidate molecular markers are those genes (and their gene products) that have a minimum gene expression in liver mets that is at least equal to the maximum gene expression in normal colon. Furthermore, in this embodiment, the median level of expression in liver mets should be greater than the median level of expression in normal liver samples.

[0108] In a preferred embodiment, a list of candidate molecular markers for colon neoplasia is selected by first identifying a subset of genes having a four-fold greater median expression in liver mets than in normal colon and in normal liver. This subset is then further narrowed to a final list by identifying those genes that have a three-fold greater median expression across colon cancer cell lines than in normal colon. Optionally, a particularly preferred list may be generated by further selecting those genes having a minimum gene expression in liver mets that is greater than or equal to the maximum gene expression in normal colon. The gene products (e.g. proteins and nucleic acids) of the short list of genes generated in these preferred embodiments constitute a list of high-quality candidate molecular markers for colon cancer.

[0109] In another preferred embodiment, a list of candidate molecular markers for colon neoplasia is selected by first identifying a subset of genes having a four-fold greater median expression in liver mets than in normal colon and in normal liver. This subset is then further narrowed by identifying those genes that have a nine-fold greater median expression in liver mets than in normal colon. This subset is then further narrowed to a final list by identifying those genes that have a four-fold greater median expression across colon cancer cell lines than in normal colon. The gene products (e.g. proteins and nucleic acids) of the short list of genes generated

in these preferred embodiments constitute a list of high-quality candidate molecular markers for colon cancer.

[0110] Depending on the nature of the intended use for the molecular marker it may be desirable to add further criteria to any of the preceding embodiments. In certain embodiments, the invention relates to candidate molecular markers for categorizing a patient as likely to have or not likely to have a colon neoplasia (including adenomas and colon cancers), and in these embodiments, a high-quality candidate molecular marker will be expressed from a gene having an increased expression in both adenomas and liver mets relative to normal colon, and preferably in other colon cancer stages, including Dukes A, Dukes B1, Dukes B2 and Dukes C. In certain embodiments the invention relates to candidate molecular markers for categorizing a patient as likely to have or not likely to have a colon cancer (including metastatic and non-metastatic forms), and in these embodiments, a high-quality candidate molecular marker will be expressed from a gene having an increased expression in liver mets relative to adenomas and normal colon, and preferably there will be elevated expression in other colon cancer stages, including Dukes A, Dukes B1, Dukes B2 and Dukes C. In certain embodiments, the invention relates to candidate molecular markers for categorizing a patient as likely or not likely to have a metastatic colon cancer, and in such embodiments, a comparison to gene expression in other colon neoplasias (e.g. adenomas, Dukes A, Dukes B1, Dukes B2, Dukes C), while potentially useful, is not necessary, although it is noted that expression in non-metastatic states may indicate that a candidate molecular marker is not of high quality for distinguishing metastatic colon cancer from non-metastatic states.

[0111] Furthermore, in those embodiments pertaining to molecular markers to be used for detection in a body fluid, such as blood, a high quality molecular marker will preferably be a secreted protein. In those embodiments pertaining to neoplasia identification or targeting, a high quality molecular marker will preferably be a protein with a portion adherent to and exposed on the extracellular surface of a neoplasia, such as a transmembrane protein with a significant extracellular portion.

[0112] Gene expression data may be gathered using one or more of the many known and appropriate techniques that, in view of this specification, may be selected to one of skill in the art. In certain preferred embodiments, gene expression data is gathered by a highly parallel system, meaning a system that allows simultaneous or near-simultaneous collection of expression data for one hundred or more gene transcripts. Exemplary highly parallel systems include probe arrays ("arrays") that are often divided into microarrays and macroarrays, where microarrays have a much higher density of individual probe species per area. Arrays generally consist of a surface to which probes that correspond in sequence to gene products (e.g., cDNAs, mRNAs, oligonucleotides) are bound at known positions. The probes can be, e.g., a synthetic oligomer, a full-length cDNA, a less-than full length cDNA, or a gene fragment. Usually a microarray will have probes corresponding to at least 100 gene products and more preferably, 500, 1000, 4000 or more. Probes may be small oligomers or larger polymers, and there may be a plurality of overlapping or non-overlapping probes for each transcript.

[0113] The nucleic acids to be contacted with the microarray may be prepared in a variety of ways. Methods for preparing total and poly(A)⁺ RNA are well known and are described generally in Sambrook et al., *supra*. Labeled cDNA

may be prepared from mRNA by oligo dT-primed or random-primed reverse transcription, both of which are well known in the art (see e.g., Klug and Berger, 1987, *Methods Enzymol.* 152:316-325). cDNAs may be labeled by incorporation of labeled nucleotides or by labeling after synthesis. Preferred labels are fluorescent labels.

[0114] Nucleic acid hybridization and wash conditions are chosen so that the population of labeled nucleic acids will specifically hybridize to appropriate, complementary probes affixed to the matrix. Optimal hybridization conditions will depend on the length (e.g., oligomer versus polynucleotide greater than 200 bases) and type (e.g., RNA, DNA, PNA) of labeled nucleic acids and immobilized polynucleotide or oligonucleotide. General parameters for specific (i.e., stringent) hybridization conditions for nucleic acids are described in Sambrook et al., *supra*, and in Ausubel et al., 1987, *Current Protocols in Molecular Biology*, Greene Publishing and Wiley-Interscience, New York, which is incorporated in its entirety for all purposes. Non-specific binding of the labeled nucleic acids to the array can be decreased by treating the array with a large quantity of non-specific DNA—a so-called "blocking" step.

[0115] Signals, such as fluorescent emissions for each location on an array are generally recorded, quantitated and analyzed using a variety of computer software. Signal for any one gene product may be normalized by a variety of different methods. Arrays preferably include control and reference probes. Control probes are nucleic acids which serve to indicate that the hybridization was effective. Reference probes allow the normalization of results from one experiment to another, and to compare multiple experiments on a quantitative level. Reference probes are typically chosen to correspond to genes that are expressed at a relatively constant level across different cell types and/or across different culture conditions. Exemplary reference nucleic acids include house-keeping genes of known expression levels, e.g., GAPDH, hexokinase and actin.

[0116] Following the data gathering operation, the data will typically be reported to a data analysis system. To facilitate data analysis, the data obtained by the reader from the device will typically be analyzed using a digital computer. Typically, the computer will be appropriately programmed for receipt and storage of the data from the device, as well as for analysis and reporting of the data gathered, e.g., subtraction of the background, deconvolution multi-color images, flagging or removing artifacts, verifying that controls have performed properly, normalizing the signals, interpreting fluorescence data to determine the amount of hybridized target, normalization of background and single base mismatch hybridizations, and the like. Various analysis methods that may be employed in such a data analysis system, or by a separate computer are described herein.

[0117] A number of methods for constructing or using arrays are described in the following references. Schena et al., 1995, *Science* 270:467-470; DeRisi et al., 1996, *Nature Genetics* 14:457-460; Shalon et al., 1996, *Genome Res.* 6:639-645; Schena et al., 1995, *Proc. Natl. Acad. Sci. USA* 93:10539-11286; Fodor et al., 1991, *Science* 251:767-773; Pease et al., 1994, *Proc. Natl. Acad. Sci. USA* 91:5022-5026; Lockhart et al., 1996, *Nature Biotech* 14:1675; U.S. Pat. Nos. 6,051,380; 6,083,697; 5,578,832; 5,599,695; 5,593,839; 5,631,734; 5,556,752; 5,510,270; EP No. 0 799 897; PCT No.

WO 97/29212; PCT No. WO 97/27317; EP No. 0 785 280; PCT No. WO 97/02357; EP No. 0 728 520; EP No. 0 721 016; PCT No. WO 95/22058.

[0118] A variety of companies provide microarrays and software for extracting certain information from microarray data. Such companies include Affymetrix (Santa Clara, Calif.), GeneLogic (Gaithersburg, Md.) and Eos Biotechnology Inc. (South San Francisco, Calif.).

[0119] While the above discussion focuses on the use of arrays for the collection of gene expression data, such data may also be obtained through a variety of other methods, that, in view of this specification, are known to one of skill in the art. Such methods include the serial analysis of gene expression (SAGE) technique, first described in Velculescu et al. (1995) *Science* 270, 484-487. Reverse transcriptase-polymerase chain reaction (RT-PCR) may be used, and particularly in combination with fluorescent probe systems such as the Taqman™ fluorescent probe system. Numerous RT-PCR samples can be analyzed simultaneously by conducting parallel PCR amplification, e.g., by multiplex PCR. Further techniques include dotblot analysis and related methods (see, e.g., G. A. Beltz et al., in *Methods in Enzymology*, Vol. 100, Part B, R. Wu, L. Grossman, K. Moldave, Eds., Academic Press, New York, Chapter 19, pp. 266-308, 1985), Northern blots and in situ hybridization (probing a tissue sample directly).

[0120] The quality and biological relevance of gene expression data will be significantly affected by the quality of the biological material used to obtain gene expression. In preferred embodiments, the methods described herein for identifying candidate molecular markers for colon neoplasia employ tissue samples obtained with appropriate consent from human patients and rapidly frozen. At a point prior to gene expression analysis, the tissue sample is preferably prepared by carefully dissecting away as much heterogeneous tissue as is possible with the available tools. In other words, for a colon cancer sample, adherent non-cancerous tissue should be dissected away, to the extent that it is possible. In preferred embodiments, healthy tissue is obtained from a subject that has a colon neoplasia but is tissue that is not directly entangled in a neoplasia.

[0121] Example 1, below, illustrates the operation of a method of selecting high-quality molecular markers, and the following markers were selected, using criteria disclosed herein, from microarray expression data: ColoUp1, ColoUp2, ColoUp3, ColoUp4, ColoUp5, ColoUp6, ColoUp7 and ColoUp8. In addition, osteopontin was identified as having expression characteristics very similar to those identified using the selection criteria. Further experimentation (see Examples) demonstrated that these molecular markers fall into four categories: "secreted" (ColoUp1, ColoUp2 and osteopontin), "transmembrane" (ColoUp3), "transcription factors" (ColoUp4, ColoUp5) and "other" (ColoUp6, ColoUp7, ColoUp8). Further experimentation also demonstrated that ColoUp1, ColoUp2, ColoUp3, ColoUp5 and ColoUp7 are, generally speaking, expressed at higher levels in a variety of colon neoplasias (adenomas, Dukes B tumors, Dukes C tumors and liver mets) than in healthy cells. In addition, further experimentation demonstrated that osteopontin is overexpressed in colon cancers (Dukes B, Dukes C and liver mets) relative to adenomas and normal colon.

[0122] In certain embodiments, a preferred molecular marker for use in a diagnostic test that employs a body fluid sample, such as a blood or urine sample, or an excreted

sample material, such as stool, is a secreted protein, such as the secreted portion of a ColoUp1 protein, ColoUp2 protein or osteopontin protein.

[0123] In certain embodiments, a preferred molecular marker for a method that involves targeting or marking a colon neoplasia is a transmembrane protein, such as ColoUp3, and particularly the extracellular portion of ColoUp3. Transmembrane proteins are desirable for such methods because they are both anchored to the neoplastic cell and exposed to the extracellular surface.

[0124] In certain embodiments, a preferred molecular marker for use in a diagnostic test to distinguish subjects likely to have a colon neoplasia from those not likely to have a colon neoplasia is a gene product of the ColoUp1, ColoUp2, ColoUp3, ColoUp4 or ColoUp5 genes. Examples of suitable gene products include proteins, both secreted and not secreted and transcripts. In embodiments employing proteins that are not secreted, such as ColoUp3, ColoUp4 and ColoUp5, a preferred embodiment of the diagnostic test is a test for the presence of the protein or transcript in cells shed from the colon or colon neoplasia (which, in the case of metastases is not necessarily located in the colon) into a sample material, such as stool. In embodiments employing proteins that are secreted, such as ColoUp1 and ColoUp2, a preferred embodiment of the diagnostic test is a test for the presence of the protein in a body fluid, such as urine or blood or an excreted material, such as stool. It should be noted, however, that intracellular protein may be present in a body fluid if there is significant cell lysis or through some other process. Likewise, secreted proteins are likely to be adherent, even if at a relatively low level, to the cells in which they were produced.

[0125] In certain embodiments, a preferred molecular marker for distinguishing subjects having a colon cancer from those having an adenoma or a normal colon is gene product of the ColoUp6 and osteopontin genes. In embodiments preferably employing marker proteins that are secreted, such as a test using a body fluid sample, a preferred marker is a secreted osteopontin protein.

ColoUp1:

[0126] A human ColoUp1 nucleic acid sequence encodes a full-length protein of 1361 amino acids. SignalP V1.1 predicts that human ColoUp1 protein has an N-terminal signal peptide that is cleaved between either amino acids 30-31 (ATS-TV) or amino acids 33-34 (TVA-AG). Four potential glycosylation sites are identified in ColoUp1 protein. Further, ColoUp1 protein is predicted to have multiple serine, threonine, and tyrosine phosphorylation sites for kinases such as protein kinase C, cAMP- and cGMP-dependent protein kinases, casein kinase II, and tyrosine kinases. The ColoUp1 protein shares limited sequence homology to a human transmembrane protein 2 (See Scott et al. 2000 Gene 246:265-74). A mouse ColoUp1 homolog is identified in existing GenBank databases and is linked with mesoderm development (see Wines et al. 2001 Genomics. 88-98; GenBank entry AAG41062, AY007815 for the 1179 bp nucleic acid sequence entry, with 363/390 (93%) identities with human ColoUp1).

[0127] As demonstrated herein, ColoUp1 is secreted from both the basolateral and apical surfaces of intestinal cells.

ColoUp2:

[0128] The ColoUp2 nucleic acid sequence encodes a full-length protein of 755 amino acids. The application also dis-

closes certain polymorphisms that have been observed, for example at nucleotide 113 GCC→ACC (Ala-Thr); nt 480 GAA→GGA (Glu-Gly); and at nt 2220 CAG→CGG (Gln-Arg). The sequence of ColoUp2 protein is similar to that of alpha 3 type VI collagen, isoform 2 precursor. In addition, a few domains are identified in the ColoUp2 protein such as a von Willebrand factor type A domain (vWF) and an EGF-like domain. The vWF domain is found in various plasma proteins such as some complement factors, the integrins, certain collagen, and other extracellular proteins. Proteins with vWF domains participate in numerous biological events which involve interaction with a large array of ligands, for example, cell adhesion, migration, homing, pattern formation, and signal transduction. The EGF-like domain consisting of about 30-40 amino acid residues has been found many proteins. The functional significance of EGF domains is not yet clear. However, a common feature is that these EGF-like repeats are found in the extracellular domain of membrane-bound proteins or in proteins known to be secreted.

[0129] As demonstrated herein, ColoUp2 is secreted from both the apical and basolateral surfaces of intestinal cells, and can be found in the blood in two different forms, a full-length secreted form and a C-terminal fragment (approximately 55 kDa).

Osteopontin:

[0130] The Osteopontin nucleic acid sequence encodes a full-length protein of 300 amino acids. Osteopontin is an acidic glycoprotein and is produced primarily by osteoclasts, macrophages, T-cells, kidneys, and vascular smooth muscle cells. As a cytokine, Osteopontin is known to contribute substantially to metastasis formation by various cancers. In addition, it contributes to macrophage homing and cellular immunity, mediates neovascularization, inhibits apoptosis, and maintains the homeostasis of free calcium (see a review, Weber G F. 2001 *Biochim Biophys Acta*. 1552:61-85).

ColoUp3:

[0131] The ColoUp3 nucleic acid sequence encodes a full-length protein of 829 amino acids. ColoUp3 is referred to in the literature as P-cadherin (or cadherin 3, type 1). P-cadherin belongs to a cadherin family that includes E-cadherin and N-cadherin. P-cadherin is expressed in placenta and stratified squamous epithelia (see Shimoyama et al. 1989 *J Cell Biol*. 109:1787-94), but not in normal colon. P-cadherin null mice develop mammary gland hyperplasia, dysplasia, and abnormal lymphoid infiltration (see Radice et al. 1997 *J Cell Biol*. 139:1025-32), demonstrating that loss of normal P-cadherin expression leads to cellular and glandular abnormalities. It has been shown that P-cadherin is aberrantly expressed in inflamed and dysplastic colitic mucosa, with concomitant E-cadherin downregulation. Recently, aberrant P-cadherin expression is found as an early event in hyperplastic and dysplastic transformation in the colon (see Hardy et al. 2002 *Gut*. 50:513-514).

ColoUp4:

[0132] The ColoUp4 nucleic acid sequence encodes a full-length protein of 694 amino acids. ColoUp4 is referred to in the literature as NF-E2 related factor 3 (NRF3). NRF3 was identified and characterized as a novel Cap'n' collar (CNC) factor, with a basic region-leucine zipper domain highly homologous to those of other CNC proteins such as NRF1

and NRF2. These CNC factors bind to Maf recognition elements (MARE) through heterodimer formation with small Maf proteins. In vitro and in vivo analyses showed that NRF3 can heterodimerize with MafK and that this complex binds to the MARE in the chicken β -globin enhancer and can activate transcription. NRF3 mRNA is highly expressed in human placenta and B cell and monocyte lineage. (see Kobayashi et al. 1999 *J Biol Chem*. 274:6443-52).

ColoUp5:

[0133] The ColoUp5 nucleic acid sequence encodes a full-length protein of 402 amino acids. ColoUp5 is referred to in the literature as FoxQ1 (Forkhead box, subclass q, member 1, formerly known as Hfh-1). FoxQ1 is a member of the evolutionarily conserved winged helix/forkhead transcription factor gene family. The hallmark of this family is a conserved DNA binding region of approximately 110 amino acids (FOX domain). Members of the FOX gene family are found in a broad range of organisms from yeast to human. Human FoxQ1 gene is expressed in different tissues such as stomach, trachea, bladder, and salivary gland. FoxQ1 gene plays important roles in tissue-specific gene regulation and development, for example, embryonic development, cell cycle regulation, cell signaling, and tumorigenesis. The FoxQ1 gene is located on chromosome 6p23-25. Sequence analysis indicates that human FoxQ1 shows 82% homology with the mouse Foxq1 gene (formerly Hfh-1L) and with a revised sequence of the rat FoxQ1 gene (formerly Hfh-1). Mouse FoxQ1 was shown to regulate differentiation of hair in Satin mice. The DNA-binding motif (i.e., the FOX domain) is well conserved, showing 100% identity in human, mouse, and rat. The human FoxQ1 protein sequence contains two putative transcriptional activation domains, which share a high amino acid identity with the corresponding mouse and rat domains (see Bieller et al. 2001 *DNA Cell Biol*. 20:555-61).

ColoUp6:

[0134] The ColoUp6 nucleic acid sequence encodes a full-length protein of 209 amino acids. The ColoUp6 protein is 99% identical to the C-terminal portion of keratin 23 (or cytokeratin 23, or the type I intermediate filament cytokeratin), and accordingly the term ColoUp6 includes both the 209 amino acid protein (and related nucleic acids, fragments, variants, etc.) and the cytokeratin 23 amino acid sequence of GenBank entry BAA92054.1 (and related nucleic acids, fragments, variants, etc.). Keratin 23 mRNA was found highly induced in different pancreatic cancer cell lines in response to sodium butyrate. The keratin 23 protein has 422 amino acids, and has an intermediate filament signature sequence and extensive homology to type I keratins. It is suggested that keratin 23 is a novel member of the acidic keratin family that is induced in pancreatic cancer cells undergoing differentiation by a mechanism involving histone hyperacetylation (See Zhang et al. 2001 *Genes Chromosomes Cancer*. 30:123-35).

ColoUp7:

[0135] The ColoUp7 nucleic acid sequence is an EST sequence. No information relating to the function of the ColoUp7 gene is identified.

ColoUp8:

[0136] The ColoUp8 nucleic acid sequence encodes a full-length protein of 278 amino acids. No function has been suggested relating to the ColoUp8 gene.

[0137] Accordingly, in certain embodiments, the application provides isolated, purified or recombinant ColoUp1, ColoUp2, ColoUp3, ColoUp4, ColoUp5, ColoUp6, ColoUp7, ColoUp8 and osteopontin nucleic acids. In certain embodiments, such nucleic acids may encode a complete or partial ColoUp polypeptide or such nucleic acids may also be probes or primers useful for methods involving detection or amplification of ColoUp nucleic acids. In certain embodiments, a ColoUp nucleic acid is single-stranded or double-stranded and composed of natural nucleic acids, nucleotide analogs, or mixtures thereof. In certain embodiments, the application provides isolated, purified or recombinant nucleic acids comprising a nucleic acid sequence that is at least 90% identical to a nucleic acid sequence of any of SEQ ID Nos: 3-12, or a complement thereof, and optionally at least 95%, 97%, 98%, 99%, 99.3%, 99.5%, 99.7% or 100% identical to a nucleic acid of any of SEQ ID Nos: 3-12, or a complement thereof. In certain preferred embodiments, the application provides a isolated, purified or recombinant nucleic acids comprising a nucleic acid sequence that is at least 90%, 95%, 97%, 98%, 99%, 99.3%, 99.5%, 99.7% or 100% identical to a nucleic acid of any of SEQ ID Nos: 3-12, or a complement thereof. In certain embodiments, the application provides isolated, purified or recombinant nucleic acids comprising a nucleic acid sequence that encodes a polypeptide that is at least 90% identical to an amino acid sequence of any of SEQ ID Nos: 1-3 or 13-21, or a complement thereof, and optionally at least 95%, 97%, 98%, 99%, 99.3%, 99.5%, 99.7% or 100% identical to an amino acid sequence of any of SEQ ID Nos: 1-3 or 13-21, or a complement thereof. In certain preferred embodiments, the application provides isolated, purified or recombinant nucleic acids comprising a nucleic acid sequence that encodes a polypeptide that is at least 90% identical to an amino acid sequence of any of SEQ ID Nos: 3, 14 or 21, or a complement thereof, and optionally at least 95%, 97%, 98%, 99%, 99.3%, 99.5%, 99.7% or 100% identical to an amino acid sequence of any of SEQ ID Nos: 3, 14 or 21, or a complement thereof.

[0138] In further embodiments, the application provides expression constructs, vectors and cells comprising a ColoUp nucleic acid. Expression constructs are nucleic acid constructs that are designed to permit expression of an expressible nucleic acid (e.g. a ColoUp nucleic acid) in a suitable cell type or in vitro expression system. A variety of expression construct systems are, in view of this specification, well known in the art, and such systems generally include a promoter that is operably linked to the expressible nucleic acid. The promoter may be a constitutive promoter, as in the case of many viral promoters, or the promoter may be a conditional promoter, as in the case of the prokaryotic *lacI*-repressible, IPTG-inducible promoter and as in the case of the eukaryotic tetracycline-inducible promoter. Vectors refer to any nucleic acid that is capable of transporting another nucleic acid to which it has been linked between different cells or viruses. One type of vector is an episome, i.e., a nucleic acid capable of extra-chromosomal replication, such as a plasmid. Episome-type vectors typically carry an origin of replication that directs replication of the vector in a host cell. Another type of vector is an integrative vector that is designed to recombine with the genetic material of a host cell. Vectors may be both autonomously replicating and integrative, and the properties of a vector may differ depending on the cellular context (i.e. a vector may be autonomously replicating in one host cell type and purely integrative in another host cell type). Vectors

capable of directing the expression of genes to which they are operatively linked are referred to herein as "expression vectors". Vectors that carry an expression construct are generally expression vectors. Vectors have been designed for a variety of cell types. For example, in the bacterium *E. coli*, commonly used vectors include pUC plasmids, pBR322 plasmids, pBlueScript and M13 plasmids. In insect cells (e.g. SF-9, SF-21 and High-Five cells), commonly used vectors include BacPak6 (Clontech) and BaculoGold (Pharmingen) (both Clontech and Pharmingen are divisions of Becton, Dickinson and Co., Franklin Lakes, N.J.). In mammalian cells (e.g. Chinese hamster ovary (CHO) cells, Vaco cells and human embryonic kidney (HEK) cells), commonly used vectors include pCMV vectors (Stratagene, Inc., La Jolla, Calif.), and pRK vectors. In certain embodiments, the application provides cells that comprise a ColoUp nucleic acid, particularly a recombinant ColoUp nucleic acid, such as an expression construct or vector that comprises a ColoUp nucleic acid. Cells may be eukaryotic or prokaryotic, depending on the anticipated use. Prokaryotic cells, especially *E. coli*, are particularly useful for storing and replicating nucleic acids, particularly nucleic acids carried on plasmid or viral vectors. Bacterial cells are also particularly useful for expressing nucleic acids to produce large quantities of recombinant protein, but bacterial cells do not usually mimic eukaryotic post-translational modifications, such as glycosylations or lipid-modifications, and so will tend to be less suitable for production of proteins in which the post-translational modification state is significant. Eukaryotic cells, and especially cell types such as insect cells that work with baculovirus-based protein expression systems, and Chinese hamster ovary cells, are good systems for expressing eukaryotic proteins that have significant post-translational modifications. Eukaryotic cells are also useful for studying various aspects of the function of eukaryotic proteins. For example, colon cancer cell lines are good model systems for studying the role of ColoUp genes and proteins in colon cancers.

[0139] In certain aspects the application further provides methods for preparing ColoUp polypeptides. In general, such methods comprise obtaining a cell that comprises a nucleic acid encoding a ColoUp polypeptide, and culturing the cell under conditions that cause production of the ColoUp polypeptide. Polypeptides produced in this manner may be obtained from the appropriate cell or culture fraction. For example, secreted proteins are most readily obtained from the culture supernatant, soluble intracellular proteins are most readily obtained from the soluble fraction of a cell lysate, and membrane proteins are most readily obtained from a membrane fraction. However, proteins of each type can generally be found in all three types of cell or culture fraction. Crude cellular or culture fractions may be subjected to further purification procedures to obtain substantially purified ColoUp polypeptides. Common purification procedures include affinity purification (e.g. with hexahistidine-tagged polypeptides), ion exchange chromatography, reverse phase chromatography, gel filtration chromatography, etc.

[0140] In certain aspects the application provides recombinant, isolated, substantially purified or purified ColoUp1, ColoUp2, ColoUp3, ColoUp4, ColoUp5, ColoUp6, ColoUp7, ColoUp8 and osteopontin polypeptides. In certain embodiments, such polypeptides may encode a complete or partial ColoUp polypeptide. In certain embodiments, a ColoUp polypeptide is composed of natural amino acids, amino acid analogs, or mixtures thereof. ColoUp polypep-

tides may also include one or more post-translational modifications, such as glycosylation, phosphorylation, lipid modification, acetylation, etc. In certain embodiments, the application provides isolated, substantially purified, purified or recombinant polypeptides comprising an amino acid sequence that is at least 90% identical to an amino acid sequence of any of SEQ ID Nos: 1-3 or 13-21 and optionally at least 95%, 97%, 98%, 99%, 99.3%, 99.5% or 99.7% identical to a nucleic acid of any of SEQ ID Nos: 1-3 or 13-21. In certain preferred embodiments, the application provides a isolated, substantially purified, purified or recombinant polypeptide comprising an amino acid sequence that is at least 90%, 95%, 97%, 98%, 99%, 99.3%, 99.5% or 99.7% identical to a nucleic acid of any of SEQ ID Nos: 3, 14 or 21. In certain preferred embodiments, the application provides an isolated, substantially purified, purified or recombinant polypeptide comprising an amino acid sequence that differs from SEQ ID Nos. 3, 14 or 21 by no more than 4 amino acid substitutions, additions or deletions. Optionally, a polypeptide of the invention comprises an additional moiety, such as an additional polypeptide sequence or other added compound, with a particular function, such as an epitope tag that facilitates detection of the recombinant polypeptide with an antibody, a purification moiety that facilitates purification (e.g. by affinity purification), a detection moiety, that facilitates detection of the polypeptide in vivo or in vitro, or an antigenic moiety that increases the antigenicity of the polypeptide so as to facilitate antibody production. Often, a single moiety will provide multiple functionalities. For example, an epitope tag will generally also assist in purification, because an antibody that recognizes the epitope can be used in an affinity purification procedure as well. Examples of commonly used epitope tags are: an HA tag, a hexahistidine tag, a V5 tag, a Glu-Glu tag, a c-myc tag, a VSV-G tag, a FLAG tag, an enterokinase cleavage site tag and a T7 tag. Commonly used purification moieties include: a hexahistidine tag, a glutathione-S-transferase domain, a cellulose binding domain and a biotin tag. Commonly used detection moieties include fluorescent proteins (e.g. green fluorescent proteins), a biotin tag, and chromogenic/fluorogenic enzymes (e.g. beta-galactosidase and luciferase). Commonly used antigenic moieties include the keyhole limpet hemocyanin and serum albumins. Note that these moieties need not be polypeptides and need not be connected to the polypeptide by a traditional peptide bond.

4. Antibodies and Uses Therefor

[0141] Another aspect of the invention pertains to an antibody specifically reactive with a ColoUp polypeptide, preferably antibodies that are specifically reactive with ColoUp polypeptides such as ColoUp1 and ColoUp2 polypeptides. For example, by using immunogens derived from a ColoUp polypeptide, e.g., based on the cDNA sequences, anti-protein/anti-peptide antisera or monoclonal antibodies can be made by standard protocols (See, for example, *Antibodies: A Laboratory Manual* ed. by Harlow and Lane (Cold Spring Harbor Press: 1988)). A mammal, such as a mouse, a hamster or rabbit can be immunized with an immunogenic form of the peptide (e.g., a ColoUp polypeptide or an antigenic fragment which is capable of eliciting an antibody response, or a fusion protein). Techniques for conferring immunogenicity on a protein or peptide include conjugation to carriers or other techniques well known in the art. An immunogenic portion of a ColoUp polypeptide can be administered in the presence of

adjuvant. The progress of immunization can be monitored by detection of antibody titers in plasma or serum. Standard ELISA or other immunoassays can be used with the immunogen as antigen to assess the levels of antibodies. In a preferred embodiment, the subject antibodies are immunospecific for antigenic determinants of a ColoUp polypeptide of a mammal, e.g., antigenic determinants of a protein set forth in SEQ ID Nos: 1-3 and 13-21, more preferably SEQ ID Nos: 1-3 or 21.

[0142] In one embodiment, antibodies are specific for the secreted proteins as encoded by nucleic acid sequences as set forth in SEQ ID Nos: 4-5. In another embodiment, the antibodies are immunoreactive with one or more proteins having an amino acid sequence that is at least 80% identical to an amino acid sequence as set forth in SEQ ID Nos: 1-3 and 13-21, preferably SEQ ID Nos: 1-3 or 21. In other embodiments, an antibody is immunoreactive with one or more proteins having an amino acid sequence that is at least 85%, 90%, 95%, 98%, 99%, 99.3%, 99.5%, 99.7% identical or 100% identical to an amino acid sequence as set forth in SEQ ID Nos: 1-3 and 13-21. More preferably, the antibody is immunoreactive with one or more proteins having an amino acid sequence that is at least 85%, 90%, 95%, 98%, 99%, 99.3%, 99.5%, 99.7% or identical to an amino acid sequence as set forth in SEQ ID Nos: 1-3 or 21. In certain preferred embodiments, the invention provides an antibody that binds to an epitope including the C-terminal portion of the polypeptide of SEQ ID Nos: 3, 14 or 21. In certain preferred embodiments, the invention provides an antibody that binds to an epitope of a ColoUp2 polypeptide that is prevalent in the blood of an animal having a colon neoplasia, such SEQ ID No: 3 or 21.

[0143] Following immunization of an animal with an antigenic preparation of a ColoUp polypeptide, anti-ColoUp antisera can be obtained and, if desired, polyclonal anti-ColoUp antibodies can be isolated from the serum. To produce monoclonal antibodies, antibody-producing cells (lymphocytes) can be harvested from an immunized animal and fused by standard somatic cell fusion procedures with immortalizing cells such as myeloma cells to yield hybridoma cells. Such techniques are well known in the art, and include, for example, the hybridoma technique (originally developed by Kohler and Milstein, (1975) *Nature*, 256: 495-497), the human B cell hybridoma technique (Kozbar et al., (1983) *Immunology Today*, 4: 72), and the EBV-hybridoma technique to produce human monoclonal antibodies (Cole et al., (1985) *Monoclonal Antibodies and Cancer Therapy*, Alan R. Liss, Inc. pp. 77-96). Hybridoma cells can be screened immunochemically for production of antibodies specifically reactive with a mammalian ColoUp polypeptide of the present invention and monoclonal antibodies isolated from a culture comprising such hybridoma cells. In one embodiment anti-human ColoUp antibodies specifically react with the protein encoded by a nucleic acid having SEQ ID Nos: 4-12; more preferably the antibodies specifically react with the protein encoded by a nucleic acid having SEQ ID Nos: 4 or 5, and preferably a secreted protein that is produced by the expression of a nucleic acid having a sequence of SEQ ID Nos: 4 or 5.

[0144] The term antibody as used herein is intended to include fragments thereof which are also specifically reactive with one of the subject ColoUp polypeptides. Antibodies can be fragmented using conventional techniques and the fragments screened for utility in the same manner as described above for whole antibodies. For example, F(ab)₂ fragments

can be generated by treating antibody with pepsin. The resulting F(ab)₂ fragment can be treated to reduce disulfide bridges to produce Fab fragments. The antibody of the present invention is further intended to include bispecific, single-chain, and chimeric and humanized molecules having affinity for a ColoUp polypeptide conferred by at least one CDR region of the antibody. In preferred embodiments, the antibodies, the antibody further comprises a label attached thereto and able to be detected, (e.g., the label can be a radioisotope, fluorescent compound, enzyme or enzyme co-factor).

[0145] In certain preferred embodiments, an antibody of the invention is a monoclonal antibody, and in certain embodiments the invention makes available methods for generating novel antibodies. For example, a method for generating a monoclonal antibody that binds specifically to a ColoUp polypeptide, such as a ColoUp2 polypeptide may comprise administering to a mouse an amount of an immunogenic composition comprising the ColoUp2 polypeptide effective to stimulate a detectable immune response, obtaining antibody-producing cells (e.g. cells from the spleen) from the mouse and fusing the antibody-producing cells with myeloma cells to obtain antibody-producing hybridomas, and testing the antibody-producing hybridomas to identify a hybridoma that produces a monoclonal antibody that binds specifically to the ColoUp2 polypeptide. Once obtained, a hybridoma can be propagated in a cell culture, optionally in culture conditions where the hybridoma-derived cells produce the monoclonal antibody that binds specifically to the ColoUp2 polypeptide. The monoclonal antibody may be purified from the cell culture.

[0146] Anti-ColoUp antibodies can be used, e.g., to detect ColoUp polypeptides in biological samples and/or to monitor ColoUp polypeptide levels in an individual, for determining whether or not said patient is likely to develop colon cancer or is more likely to harbor colon adenomas, or allowing determination of the efficacy of a given treatment regimen for an individual afflicted with colon neoplasia, colon cancer, metastatic colon cancer and colon adenomas. The level of ColoUp polypeptide may be measured in a variety of sample types such as, for example, in cells, stools, and/or in bodily fluid, such as in whole blood samples, blood serum, blood plasma and urine. The adjective "specifically reactive with" as used in reference to an antibody is intended to mean, as is generally understood in the art, that the antibody is sufficiently selective between the antigen of interest (e.g. a ColoUp polypeptide) and other antigens that are not of interest that the antibody is useful for, at minimum, detecting the presence of the antigen of interest in a particular type of biological sample. In certain methods employing the antibody, a higher degree of specificity in binding may be desirable. For example, an antibody for use in detecting a low abundance protein of interest in the presence of one or more very high abundance protein that are not of interest may perform better if it has a higher degree of selectivity between the antigen of interest and other cross-reactants. Monoclonal antibodies generally have a greater tendency (as compared to polyclonal antibodies) to discriminate effectively between the desired antigens and cross-reacting polypeptides. In addition, an antibody that is effective at selectively identifying an antigen of interest in one type of biological sample (e.g. a stool sample) may not be as effective for selectively identifying the same antigen in a different type of biological sample (e.g. a blood sample). Likewise, an antibody that is effective at identifying an antigen of interest in a purified protein preparation that is devoid of other biological

contaminants may not be as effective at identifying an antigen of interest in a crude biological sample, such as a blood or urine sample. Accordingly, in preferred embodiments, the application provides antibodies that have demonstrated specificity for an antigen of interest (particularly, although not limited to, a ColoUp1 or ColoUp2 polypeptide) in a sample type that is likely to be the sample type of choice for use of the antibody. In a particularly preferred embodiment, the application provides antibodies that bind specifically to a ColoUp1 or ColoUp2 polypeptide in a protein preparation from blood (optionally serum or plasma) from a patient that has a colon neoplasia or that bind specifically in a crude blood sample (optionally a crude serum or plasma sample).

[0147] One characteristic that influences the specificity of an antibody:antigen interaction is the affinity of the antibody for the antigen. Although the desired specificity may be reached with a range of different affinities, generally preferred antibodies will have an affinity (a dissociation constant) of about 10^{-6} , 10^{-7} , 10^{-8} , 10^{-9} or less.

[0148] In addition, the techniques used to screen antibodies in order to identify a desirable antibody may influence the properties of the antibody obtained. For example, an antibody to be used for certain therapeutic purposes will preferably be able to target a particular cell type. Accordingly, to obtain antibodies of this type, it may be desirable to screen for antibodies that bind to cells that express the antigen of interest (e.g. by fluorescence activated cell sorting). Likewise, if an antibody is to be used for binding an antigen in solution, it may be desirable to test solution binding. A variety of different techniques are available for testing antibody:antigen interactions to identify particularly desirable antibodies. Such techniques include ELISAs, surface plasmon resonance binding assays (e.g. the Biacore binding assay, Biacore AB, Uppsala, Sweden), sandwich assays (e.g. the paramagnetic bead system of IGEN International, Inc., Gaithersburg, Md.), western blots, immunoprecipitation assays and immunohistochemistry.

[0149] Another application of anti-ColoUp antibodies of the present invention is in the immunological screening of cDNA libraries constructed in expression vectors such as gt11, gt18-23, ZAP, and ORF8. Messenger libraries of this type, having coding sequences inserted in the correct reading frame and orientation, can produce fusion proteins. For instance, gt11 will produce fusion proteins whose amino termini consist of β -galactosidase amino acid sequences and whose carboxy termini consist of a foreign polypeptide. Antigenic epitopes of a ColoUp polypeptide, e.g., other orthologs of a particular protein or other paralogs from the same species, can then be detected with antibodies, as, for example, reacting nitrocellulose filters lifted from infected plates with the appropriate anti-ColoUp antibodies. Positive phage detected by this assay can then be isolated from the infected plate. Thus, the presence of ColoUp homologs can be detected and cloned from other animals, as can alternate isoforms (including splice variants) from humans.

5. Methods for Detecting Molecular Markers in a Patient

[0150] In certain embodiments, the invention provides methods for detecting molecular markers, such as proteins or nucleic acid transcripts of the ColoUp markers described herein. In certain embodiments, a method of the invention comprises providing a biological sample and probing the biological sample for the presence of a ColoUp marker. Information regarding the presence or absence of the ColoUp

marker, and optionally the quantitative level of the ColoUp marker, may then be used to draw inferences about the nature of the biological sample and, if the biological sample was obtained from a subject, the health state of the subject.

[0151] Samples for use with the methods described herein may be essentially any biological material of interest. For example, a sample may be a tissue sample from a subject, a fluid sample from a subject, a solid or semi-solid sample from a subject, a primary cell culture or tissue culture of materials derived from a subject, cells from a cell line, or medium or other extracellular material from a cell or tissue culture, or a xenograft (meaning a sample of a colon cancer from a first subject, e.g. a human, that has been cultured in a second subject, e.g. an immunocompromised mouse). The term "sample" as used herein is intended to encompass both a biological material obtained directly from a subject (which may be described as the primary sample) as well as any manipulated forms or portions of a primary sample. For example, in certain embodiments, a preferred fluid sample is a blood sample. In this case, the term sample is intended to encompass not only the blood as obtained directly from the patient but also fractions of the blood, such as plasma, serum, cell fractions (e.g. platelets, erythrocytes, lymphocytes), protein preparations, nucleic acid preparations, etc. A sample may also be obtained by contacting a biological material with an exogenous liquid, resulting in the production of a lavage liquid containing some portion of the contacted biological material. Furthermore, the term "sample" is intended to encompass the primary sample after it has been mixed with one or more additive, such as preservatives, chelators, anti-clotting factors, etc. In certain embodiments, a fluid sample is a urine sample. In certain embodiments, a preferred solid or semi-solid sample is a stool sample. In certain embodiments, a preferred tissue sample is a biopsy from a tissue known to harbor or suspected of harboring a colon neoplasia. In certain embodiments, a preferred cell culture sample is a sample comprising cultured cells of a colon cancer cell line, such as a cell line cultured from a metastatic colon cancer tumor or a colon-derived cell line lacking a functional TGF- β , TGF- β receptor or TGF- β signaling pathway. A subject is preferably a human subject, but it is expected that the molecular markers disclosed herein, and particularly their homologs from other animals, are of similar utility in other animals. In certain embodiments, it may be possible to detect a marker directly in an organism without obtaining a separate portion of biological material. In such instances, the term sample is intended to encompass that portion of biological material that is contacted with a reagent or device involved in the detection process.

[0152] In certain embodiments, a method of the invention comprises detecting the presence of a ColoUp protein in a sample. Optionally, the method involves obtaining a quantitative measure of the ColoUp protein in the sample. In view of this specification, one of skill in the art will recognize a wide range of techniques that may be employed to detect and optionally quantitate the presence of a protein. In preferred embodiments, a ColoUp protein is detected with an antibody. Suitable antibodies are described in a separate section below. In many embodiments, an antibody-based detection assay involves bringing the sample and the antibody into contact so that the antibody has an opportunity to bind to proteins having the corresponding epitope. In many embodiments, an antibody-based detection assay also typically involves a system for detecting the presence of antibody-epitope complexes,

thereby achieving a detection of the presence of the proteins having the corresponding epitope. Antibodies may be used in a variety of detection techniques, including enzyme-linked immunosorbent assays (ELISAs), immunoprecipitations, Western blots. Antibody-independent techniques for identifying a protein may also be employed. For example, mass spectroscopy, particularly coupled with liquid chromatography, permits detection and quantification of large numbers of proteins in a sample. Two-dimensional gel electrophoresis may also be used to identify proteins, and may be coupled with mass spectroscopy or other detection techniques, such as N-terminal protein sequencing. RNA aptamers with specific binding for the protein of interest may also be generated and used as a detection reagent.

[0153] In certain preferred embodiments, methods of the invention involve detection of a secreted form of a ColoUp protein or osteopontin, particularly ColoUp1 protein or ColoUp2 protein.

[0154] Samples should generally be prepared in a manner that is consistent with the detection system to be employed. For example, a sample to be used in a protein detection system should generally be prepared in the absence of proteases. Likewise, a sample to be used in a nucleic acid detection system should generally be prepared in the absence of nucleases. In many instances, a sample for use in an antibody-based detection system will not be subjected to substantial preparatory steps. For example, urine may be used directly, as may saliva and blood, although blood will, in certain preferred embodiments, be separated into fractions such as plasma and serum.

[0155] In certain embodiments, a method of the invention comprises detecting the presence of a ColoUp expressed nucleic acid, such as an mRNA, in a sample. Optionally, the method involves obtaining a quantitative measure of the ColoUp expressed nucleic acid in the sample. In view of this specification, one of skill in the art will recognize a wide range of techniques that may be employed to detect and optionally quantitate the presence of a nucleic acid. Nucleic acid detection systems generally involve preparing a purified nucleic acid fraction of a sample, and subjecting the sample to a direct detection assay or an amplification process followed by a detection assay. Amplification may be achieved, for example, by polymerase chain reaction (PCR), reverse transcriptase (RT) and coupled RT-PCR. Detection of a nucleic acid is generally accomplished by probing the purified nucleic acid fraction with a probe that hybridizes to the nucleic acid of interest, and in many instances detection involves an amplification as well. Northern blots, dot blots, microarrays, quantitative PCR and quantitative RT-PCR are all well known methods for detecting a nucleic acid in a sample.

[0156] In certain embodiments, the invention provides nucleic acid probes that bind specifically to a ColoUp nucleic acid. Such probes may be labeled with, for example, a fluorescent moiety, a radionuclide, an enzyme or an affinity tag such as a biotin moiety. For example, the TaqMan® system employs nucleic acid probes that are labeled in such a way that the fluorescent signal is quenched when the probe is free in solution and bright when the probe is incorporated into a larger nucleic acid.

[0157] In certain embodiments, the application provides methods for imaging a colon neoplasm by targeting antibodies to any one of the markers ColoUp1 through ColoUp8 or osteopontin described herein, more preferably the antibodies

are targeted to ColoUp3. The markers described herein may be targeted using monoclonal antibodies which may be labeled with radioisotopes for clinical imaging of tumors or with toxic agents to destroy them.

[0158] In other embodiments, the application provides methods for administering a imaging agent comprising a targeting moiety and an active moiety. The targeting moiety may be an antibody, Fab, F(Ab)₂, a single chain antibody or other binding agent that interacts with an epitope specified by a polypeptide sequence having an amino acid sequence as set forth in SEQ ID Nos: 1-3 and 13-21, preferably an epitope specified by SEQ ID No: 16. The active moiety may be a radioactive agent, such as: radioactive heavy metals such as iron chelates, radioactive chelates of gadolinium or manganese, positron emitters of oxygen, nitrogen, iron, carbon, or gallium, ⁴³K, ⁵²Fe, ⁵⁷Co, ⁶⁷CU, ⁶⁷Ga, ⁶⁸Ga, ¹²³I, ¹²⁵I, ¹³¹I, ¹³²I, or ^{99m}Tc. The imaging agent is administered in an amount

examples, which are included merely for purposes of illustration of certain aspects and embodiments of the present invention, and are not intended to limit the invention.

Example 1

Selection of Eight Molecular Markers for Colon Neoplasia

[0161] Expression micro-array profiling was used to find genes whose expression was different between normal colon and metastatic colon cancer. Normal colon and metastatic colon cancer samples were analyzed for gene expression using DNA expression microarray techniques that profiled expression patterns of nearly 50,000 genes, ESTs and predicted exons. Analysis of the data identified eight molecular markers for colon neoplasia, as shown in Table 2.

TABLE 2

Eight Selected Molecular Markers for Colon Neoplasia						
Marker Name	Example Sequences (SEQ ID Nos.)	(Median Liver Mets)/ (Median Normal Colon)	(Median Liver Mets)/ (Median Normal Liver)	(Minimum Liver Mets)/ (Maximum Normal Colon)	(Median Met Cell Lines)/ (Median Normal Colon)	(Median Met Xenografts)/ (Median Normal Colon)
ColoUp1	1, 2, 4, 13	13.94	13.94	0.26	14.08	15.48
ColoUp2	3, 5, 14	5.70	5.70	1.00	5.32	1.24
ColoUp3	7, 16	16.36	16.36	0.80	21.50	15.68
ColoUp4	8, 17	4.68	4.68	1.00	4.88	1.56
ColoUp5	9, 18	4.58	4.74	1.15	4.82	4.63
ColoUp6	10, 19	9.52	9.52	0.52	11.58	1.92
ColoUp7	11	9.20	9.20	0.18	4.30	9.00
ColoUp8	12, 20	4.78	4.78	1.27	3.76	2.72

effective for diagnostic use in a mammal such as a human and the localization and accumulation of the imaging agent is then detected. The localization and accumulation of the imaging agent may be detected by radiosintigraphy, nuclear magnetic resonance imaging, computed tomography or positron emission tomography.

[0159] Immunoscintigraphy using monoclonal antibodies directed at the ColoUp markers may be used to detect and/or diagnose colon neoplasia. For example, monoclonal antibodies against the ColoUp marker such as ColoUp3 labeled with ^{99m}Technetium, ¹¹¹Indium, ¹²⁵Iodine-may be effectively used for such imaging. As will be evident to the skilled artisan, the amount of radioisotope to be administered is dependent upon the radioisotope. Those having ordinary skill in the art can readily formulate the amount of the imaging agent to be administered based upon the specific activity and energy of a given radionuclide used as the active moiety. Typically 0.1-100 millicuries per dose of imaging agent, preferably 1-10 millicuries, most often 2-5 millicuries are administered. Thus, compositions according to the present invention useful as imaging agents comprising a targeting moiety conjugated to a radioactive moiety comprise 0.1-100 millicuries, in some embodiments preferably 1-10 millicuries, in some embodiments preferably 2-5 millicuries, in some embodiments more preferably 1-5 millicuries.

Exemplification

[0160] The invention now being generally described, it will be more readily understood by reference to the following

[0162] Osteopontin was also identified as a molecular marker having similar characteristics (Example sequences SEQ ID Nos: 6, 15). Each of these molecular markers was subjected to additional analysis in various types of colon neoplasia. In the case of ColoUp1 and ColoUp2, the microarray expression was confirmed by Northern blot and secretion of the protein was established.

Example 2

Expression Pattern of ColoUp1 in Various Cell Types

[0163] Shown in FIG. 20 is a graphical display of ColoUp1 expression levels measured for different tissue samples. ColoUp1 transcript was essentially undetectable (AI expression levels less than 0) in normal colon epithelial strips (labeled colon epithelial), in normal liver and in colonic muscle (labeled c. muscle). In contrast ColoUp1 expression was clearly detected in premalignant colon adenomas as well as in 90% of Dukes stage B (early node negative colon cancers), Dukes stage C (node positive colon cancer), Dukes stage D (primary colon cancers with associated metastatic spread) and in colon cancer liver metastasis (labeled liver metastasis). ColoUp1 expression was also demonstrated in colon cancer cell lines (labeled colon cell lines) and in colon cancer xenografts grown in athymic mice (labeled xenografts). The

expression in cell lines and xenografts confirms that colon neoplasia cells are the source of ColoUp1 expression in the tumors.

[0164] The probe for ColoUp1 was designed to recognize transcripts corresponding to gene KIAA1199, Genbank entry AB033025, Unigene entry Hs.50081. A transcript corresponding to this gene was amplified by RT-PCR from colon cancer cell line Vaco-394. The sequence of this transcript is presented in FIG. 3.

Example 3

Confirmed Gene Expression Pattern of ColoUp1

[0165] FIG. 29 shows a northern analysis using the cloned ColoUp1 cDNA that identifies a transcript running above the large ribosomal subunit (to which the probe cross hybridizes) that is not expressed in normal colon tissue samples and is ubiquitously expressed in a group of colon cancer cell lines.

[0166] FIGS. 29B and 29C show the results of northern analysis of ColoUp1 in normal colon tissue and colon neoplasms from 15 individuals with colon cancers and one individual with a colon adenoma. No normal colon sample expresses ColoUp1. However, expression is seen in 13 of 15 colon cancers, and in the one colon adenoma. Expression is seen in cancers arising in both the right and left colon, and in cancers of Dukes Stage B2, C and D.

Example 4

ColoUp1 is a Secreted Protein

[0167] The cloned ColoUp1 colonic transcript was inserted into a cDNA expression vector with a C-terminal T7 epitope tag. FIG. 30A shows a summary of the behavior of the tagged protein expressed by transfection of the vector into Vaco400 cells. An anti T7 western blot shows expression of the transfected tagged protein detected in the lysate of a pellet of transfected cells (lane T of cell pellet) which is absent in cells transfected with a control empty expression vector (lane C of cell pellet). Moreover, serial immunoprecipitation and western blotting of T7 tagged protein from media in which V400 cells were growing (which had been clarified by centrifugation prior to immunoprecipitation) also clearly demonstrates secretion of ColoUp1 protein into the growth medium.

[0168] FIG. 30B shows the full gels demonstrating expression of tagged 409041 protein in V400 cells demonstrated by western analysis at left and shows detection of secreted 409041 protein in growth media as detected at right by serial immunoprecipitation and western analysis. (Antibody from the high level of serum in which FET cells are grown blocked the ability of staphA conjugated beads to precipitate anti-T7 bound to 409041 in growth media from FET cells).

Example 5

Expression Pattern of ColoUp2 in Various Cell Types

[0169] Shown in FIG. 21 is the graphical display of ColoUp2 expression levels measured for different samples analyzed. ColoUp2 transcript was essentially undetectable (AI expression levels less than 0) in normal colon epithelial strips (labeled colon epithelial), in normal liver and in colonic muscle (labeled c. muscle). In contrast ColoUp2 expression was clearly detected in premalignant colon adenomas as well as in 90% of Dukes stage B (early node negative colon cancers), Dukes stage C (node positive colon cancer), Dukes

stage D (primary colon cancers with associated metastatic spread) and in colon cancer liver metastasis (labeled liver metastasis). ColoUp2 expression was also demonstrated in colon cancer cell lines (labeled colon cell lines) and in colon cancer xenografts grown in athymic mice (labeled xenografts). The expression in cell lines and xenografts confirms that colon neoplasia cells are the source of ColoUp2 expression in the tumors.

[0170] Probe ColoUp2 was designed to recognize transcripts corresponding to a noncoding EST, Genbank entry AI357412, Unigene entry Hs.157601. By 5' RACE, database assembly, and ultimately RT-PCR, we cloned from a colon cancer cell line a novel protein encoding RNA transcript whose noncoding 3' UTR was shown to correspond to the ColoUp2 specified EST. This full length coding sequence was determined by RT-PCR amplification from colon cancer cell line Vaco503 and sequences are provided in FIG. 4.

[0171] ColoUp2 is a "class identifier" (that is, it is higher in all colon cancer samples than in all normal colon samples), it is not-expressed in normal body tissues and it contains a signal sequence predicting that the protein product will be secreted (as well as several other recognizable protein motifs including domains from the epidermal growth factor protein and from the Von Willebrands protein).

Example 6

Confirmed Gene Expression Pattern of ColoUp2

[0172] FIG. 31 shows a northern analysis using the cloned ColoUp2 cDNA that identifies a transcript running above the large ribosomal subunit (to which the probe cross hybridizes) that is not expressed in normal colon tissue samples and is expressed in the majority of group of colon cancer cell lines. Panel A of the figure shows the northern hybridization. The red arrow designates the ColoUp2 transcript. Above each lane is the name of the sample and the level (in parenthesis) of ColoUp2 expression recorded. The black arrow designates the cross hybridizing ribosomal large subunit. Panel B shows the ethidium bromide stained gel corresponding to the blot, and the black arrows designate the large and small ribosomal subunits.

Example 7

ColoUp2 is a Secreted Protein

[0173] The cloned ColoUp2 colonic transcript was inserted into a cDNA expression vector with a C-terminal V5 epitope tag. FIG. 32 shows a summary of the behavior of the tagged protein expressed by transfection of the vector into SW480 and Vaco400 cells. An anti V5 western blot shows (red arrows) expression of the transfected tagged protein detected in the lysate of a pellet of transfected cells (lysates western panel, lanes labeled ColoUp2/V5) which is absent in cells transfected with a control empty expression vector (lanes labeled pcDNA3.1). Moreover, serial immunoprecipitation and western blotting of V5 tagged protein from media in which V400 and SW480 cells were growing (which had been clarified by centrifugation prior to immunoprecipitation) also clearly demonstrates secretion of the ColoUp2 protein into the growth medium (panel labeled medium IP-western). Antibody bands from the immunoprecipitation are also present on

the IP-western blot. Detection of secreted ColoUp2 protein was shown in cells assayed both 24 hours and 48 hours after transfection.

Example 8

Expression Pattern of ColoUp3-ColoUp8 and Osteopontin in Various Cell Types

[0174] Shown in FIGS. 22-28 are the graphical displays of ColoUp3-ColoUp8 and osteopontin expression levels measured for different samples analyzed.

Example 9

Confirmed Gene Expression Pattern of ColoUp5

[0175] Shown in FIG. 33 is a northern blot showing that ColoUp5 is expressed in colon cancer cell lines and not expressed in non-neoplastic material. FIG. 33 shows two northern blot analysis of ColoUp5 mRNA levels in normal colon tissues and a group of colon cancer cell lines (top panels). The bottom panels show the ethidium bromide stained gel corresponding to the blot. Homologs for ColoUp5 are found in other mammals, including mouse and rat, and sequence alignments are shown in FIGS. 34 and 35.

Example 10

Detection of Xenograft Derived ColoUp1 and ColoUp2 Proteins Circulating in the Blood of Mice

[0176] To determine that ColoUp1 and ColoUp2 proteins are effective serologic markers of colon neoplasia, we derived transfected cell lines that stably expressed and secreted V5-epitope tagged ColoUp1 and ColoUp2 proteins. These cells lines were then injected into athymic mice and grown as tumor xenografts. Mice were sacrificed and serum was obtained. V5 tagged proteins were then precipitated from the serum using beads conjugated to anti-V5 antibodies. Precipitated serum proteins were run out on SDS-PAGE, and visualized by western blotting using HRP-conjugated anti-V5 antibodies (thereby eliminating visualization of any contaminating mouse immunoglobulin). FIG. 36 shows detection of circulating ColoUp2 protein in mouse serum. The ColoUp2 protein is secreted as 2 bands of 85 KD and 55 KD in size, of which the 55 KD band predominates in the serum. The 55 KD band is presumably a processed form of the 85 KD band. This observation demonstrates that, in this mouse model, ColoUp2 is indeed a secreted marker of colon cancers and adenomas, and that ColoUp2 can gain access to and circulate stably in patient serum. This observation provides the surprising result that a processed fragment of ColoUp2 is the predominant serum form of the protein and therefore detection reagents targeted to this portion would be particularly suitable for diagnostic testing.

[0177] A time course experiment showed that ColoUp2 protein was detectable in mouse blood at the earliest time assayed, 1 week after injection of ColoUp2 secreting colon cancer cells, at which time xenograft tumor volume as only 100 mm³.

[0178] Similar observations were also made for ColoUp1, as shown in FIG. 37.

Example 11

Purification of ColoUp1 and ColoUp2 Proteins

[0179] In order to develop monoclonal antibodies against native ColoUp1 and ColoUp2 proteins, we devised a protocol for purification on Ni-NTA agarose (QIAGEN) nickel beads of recombinant His tagged ColoUp1 and ColoUp2 proteins from the media supernate of SW480 cells engineered to express these proteins. Currently we have purified both ColoUp1 and ColoUp2 proteins to sufficient purity to generate antibodies. As shown in FIG. 38, a Coomassie blue stained gel of purified ColoUp2 shows only the 85 KD and 55 KD size bands that correspond to the tagged ColoUp2 proteins visualized on western blot. Similarly, a Coomassie blue stained gel of purified ColoUp1 shows the preparation is highly purified and composed of a single 180 KD band that corresponds perfectly to the size band seen on western blotting of the epitope tagged ColoUp1 protein. Thus we have purified ColoUp2 and ColoUp1 to sufficient homogeneity and yield. Scaled up purification of these proteins from a 50 liter media preparation should yield 2.5 mg of protein, more than adequate for immunizing mice and screening fusion supernates for development of monoclonal antibodies specific for native ColoUp1 and ColoUp2.

Example 12

Measuring Apical and Basolateral Secretion of ColoUp1 and ColoUp2

[0180] We expected that ColoUp2 will serve as a serologic marker detection not only of colon cancers but also of large colon adenomas that also express ColoUp2. Adenomas, unlike colon cancers, are non-invasive. Thus, for adenomas to move ColoUp2 proteins into the circulation they would need to secrete this protein from the basolateral cell surface facing capillaries and lymphatics, rather than from the apical cell surface facing the colon lumen. To determine the polarity of ColoUp2 secretion we transiently transfected a monolayer of polarized Caco2 colon cancer cells with an expression vector for V5-epitope tagged ColoUp2 protein. This cell monolayer was grown in transwell dishes on filters that separate an upper transwell chamber (representing media exposed to the apical surface of the monolayer) from a lower transwell chamber (representing media exposed to the basolateral surface of the monolayer). Integrity of the sealing of the monolayer was assayed by measuring electrical resistance across the filters, and efficiency of transient transfection was monitored by expression of a GFP marker. Media from upper and lower chambers was harvested at 24, 48, 72, and 96 hours post transfection, and secreted tagged ColoUp2 protein was detected by western analysis directed against the V5 epitope tag. As FIG. 39 shows, characteristic 85 KD and 55 KD secreted forms of ColoUp2 were detected in media sampling the basolateral monolayer compartment at all time points assayed. At a single time point, 48 hours, ColoUp2 was additionally detected in media representing the apical secretion face; however, a dip in the transfilter electrical resistance at 48 hours suggests the likelihood of some leaking across the monolayer at this time point. Certainly, the data clearly shows secretion of ColoUp2 into the basolateral monolayer com-

partment, and hence establishes ColoUp2 as demonstrating the requisite biology for a candidate serologic marker of colon adenomas.

[0181] As was done for ColoUp2, ColoUp1 expression vectors were used to transiently transfect Caco2 cell monolayers grown on transwell filters. Secretion of ColoUp1 was then assayed in media collected respectively from the upper and lower transwell chambers. Western blot assays demonstrated equal secretion of ColoUp1 from both apical and basolateral monolayer surfaces. Studies of ColoUp1 were done in parallel with those of ColoUp2, and electrical resistance of the ColoUp1 monolayers exceeded that of the ColoUp2 monolayers, supporting that the ColoUp1 transfected monolayers were well sealed. Additionally, levels of secreted ColoUp1 protein were similar to those of secreted ColoUp2, suggesting that ColoUp1 secretion by both apical and basolateral compartments was not simply due to overexpression. Accordingly, we predict that native ColoUp1 protein is likely secreted at least in part from the basolateral epithelial face, and hence should be detectable as a serologic marker of large colon adenomas.

Example 13

Determining the Sequence of the 55 kDa ColoUp2 Fragment

[0182] The protein sequence of C-terminal fragment of ColoUp2 that is secreted by human cell lines and detected as predominant fragment in blood (488 aa) was determined. As described above, we have found on western blots and on purified preparations of C-terminal epitope tagged (V5-His epitope) ColoUp2 protein secreted by transfected human colon cancer cells, both a full sized band of approximately 90 kDa and a smaller approximately 55 kDa C-terminal fragment (as demonstrated by the retention of the C-terminal

epitope tag). Moreover, when these cells were injected into athymic mice, the 55 kDa C-terminal tagged protein was the predominant species detected as circulating in the mouse blood, when mouse serum is analyzed by serial immunoprecipitation and western blot analysis directed against the V5 tag. The precise location of the cleavage site accounting for the C-terminal fragment was established by excising the acrylamide gel band containing the purified C-terminal fragment and performing mass spectroscopy analysis of tryptic fragments from the protein. A peptide of sequence AVLAAHCP-FYSWK was present only in the digest of the 55 KD fragment, but was absent from the digest of the full length protein, demonstrating that this peptide corresponded to the unique amino terminus of the 55 KD fragment. The complete sequence of the 55 KD C-terminal fragment is shown in FIG. 41.

Incorporation by Reference

[0183] All publications and patents mentioned herein are hereby incorporated by reference in their entirety as if each individual publication or patent was specifically and individually indicated to be incorporated by reference. In case of conflict, the present application, including any definitions herein, will control.

Equivalents

[0184] While specific embodiments of the subject invention have been discussed, the above specification is illustrative and not restrictive. Many variations of the invention will become apparent to those skilled in the art upon review of this specification and the claims below. The full scope of the invention should be determined by reference to the claims, along with their full scope of equivalents, and the specification, along with such variations.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 28

<210> SEQ ID NO 1

<211> LENGTH: 1331

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 1

```

Thr Val Ala Ala Gly Cys Pro Asp Gln Ser Pro Glu Leu Gln Pro Trp
 1           5           10          15
Asn Pro Gly His Asp Gln Asp His Val His Ile Gly Gln Gly Lys
 20          25
Thr Leu Leu Leu Thr Ser Ser Ala Thr Val Tyr Ser Ile His Ile Ser
 35          40          45
Glu Gly Gly Lys Leu Val Ile Lys Asp His Asp Glu Pro Ile Val Leu
 50          55          60
Arg Thr Arg His Ile Leu Ile Asp Asn Gly Gly Glu Leu His Ala Gly
 65          70          75          80
Ser Ala Leu Cys Pro Phe Gln Gly Asn Phe Thr Ile Ile Leu Tyr Gly
 85          90          95
Arg Ala Asp Glu Gly Ile Gln Pro Asp Pro Tyr Tyr Gly Leu Lys Tyr

```

-continued

100				105				110							
Ile	Gly	Val	Gly	Lys	Gly	Gly	Ala	Leu	Glu	Leu	His	Gly	Gln	Lys	Lys
	115						120					125			
Leu	Ser	Trp	Thr	Phe	Leu	Asn	Lys	Thr	Leu	His	Pro	Gly	Gly	Met	Ala
	130					135					140				
Glu	Gly	Gly	Tyr	Phe	Phe	Glu	Arg	Ser	Trp	Gly	His	Arg	Gly	Val	Ile
145					150					155					160
Val	His	Val	Ile	Asp	Pro	Lys	Ser	Gly	Thr	Val	Ile	His	Ser	Asp	Arg
				165					170					175	
Phe	Asp	Thr	Tyr	Arg	Ser	Lys	Lys	Glu	Ser	Glu	Arg	Leu	Val	Gln	Tyr
			180						185					190	
Leu	Asn	Ala	Val	Pro	Asp	Gly	Arg	Ile	Leu	Ser	Val	Ala	Val	Asn	Asp
		195					200						205		
Glu	Gly	Ser	Arg	Asn	Leu	Asp	Asp	Met	Ala	Arg	Lys	Ala	Met	Thr	Lys
	210					215					220				
Leu	Gly	Ser	Lys	His	Phe	Leu	His	Leu	Gly	Phe	Arg	His	Pro	Trp	Ser
225					230					235					240
Phe	Leu	Thr	Val	Lys	Gly	Asn	Pro	Ser	Ser	Ser	Val	Glu	Asp	His	Ile
				245					250					255	
Glu	Tyr	His	Gly	His	Arg	Gly	Ser	Ala	Ala	Ala	Arg	Val	Phe	Lys	Leu
			260						265					270	
Phe	Gln	Thr	Glu	His	Gly	Glu	Tyr	Phe	Asn	Val	Ser	Leu	Ser	Ser	Glu
			275				280						285		
Trp	Val	Gln	Asp	Val	Glu	Trp	Thr	Glu	Trp	Phe	Asp	His	Asp	Lys	Val
	290					295					300				
Ser	Gln	Thr	Lys	Gly	Gly	Glu	Lys	Ile	Ser	Asp	Leu	Trp	Lys	Ala	His
305					310					315					320
Pro	Gly	Lys	Ile	Cys	Asn	Arg	Pro	Ile	Asp	Ile	Gln	Ala	Thr	Thr	Met
				325					330					335	
Asp	Gly	Val	Asn	Leu	Ser	Thr	Glu	Val	Val	Tyr	Lys	Lys	Gly	Gln	Asp
			340						345				350		
Tyr	Arg	Phe	Ala	Cys	Tyr	Asp	Arg	Gly	Arg	Ala	Cys	Arg	Ser	Tyr	Arg
		355				360						365			
Val	Arg	Phe	Leu	Cys	Gly	Lys	Pro	Val	Arg	Pro	Lys	Leu	Thr	Val	Thr
	370					375					380				
Ile	Asp	Thr	Asn	Val	Asn	Ser	Thr	Ile	Leu	Asn	Leu	Glu	Asp	Asn	Val
385					390					395					400
Gln	Ser	Trp	Lys	Pro	Gly	Asp	Thr	Leu	Val	Ile	Ala	Ser	Thr	Asp	Tyr
				405					410					415	
Ser	Met	Tyr	Gln	Ala	Glu	Glu	Phe	Gln	Val	Leu	Pro	Cys	Arg	Ser	Cys
			420						425				430		
Ala	Pro	Asn	Gln	Val	Lys	Val	Ala	Gly	Lys	Pro	Met	Tyr	Leu	His	Ile
		435					440						445		
Gly	Glu	Glu	Ile	Asp	Gly	Val	Asp	Met	Arg	Ala	Glu	Val	Gly	Leu	Leu
	450					455					460				
Ser	Arg	Asn	Ile	Ile	Val	Met	Gly	Glu	Met	Glu	Asp	Lys	Cys	Tyr	Pro
465					470					475					480
Tyr	Arg	Asn	His	Ile	Cys	Asn	Phe	Phe	Asp	Phe	Asp	Thr	Phe	Gly	Gly
			485						490					495	
His	Ile	Lys	Phe	Ala	Leu	Gly	Phe	Lys	Ala	Ala	His	Leu	Glu	Gly	Thr
			500						505					510	

-continued

Glu Leu Lys His Met Gly Gln Gln Leu Val Gly Gln Tyr Pro Ile His
 515 520 525
 Phe His Leu Ala Gly Asp Val Asp Glu Arg Gly Gly Tyr Asp Pro Pro
 530 535 540
 Thr Tyr Ile Arg Asp Leu Ser Ile His His Thr Phe Ser Arg Cys Val
 545 550 555 560
 Thr Val His Gly Ser Asn Gly Leu Leu Ile Lys Asp Val Val Gly Tyr
 565 570 575
 Asn Ser Leu Gly His Cys Phe Phe Thr Glu Asp Gly Pro Glu Glu Arg
 580 585 590
 Asn Thr Phe Asp His Cys Leu Gly Leu Leu Val Lys Ser Gly Thr Leu
 595 600 605
 Leu Pro Ser Asp Arg Asp Ser Lys Met Cys Lys Met Ile Thr Glu Asp
 610 615 620
 Ser Tyr Pro Gly Tyr Ile Pro Lys Pro Arg Gln Asp Cys Asn Ala Val
 625 630 635 640
 Ser Thr Phe Trp Met Ala Asn Pro Asn Asn Asn Leu Ile Asn Cys Ala
 645 650 655
 Ala Ala Gly Ser Glu Glu Thr Gly Phe Trp Phe Ile Phe His His Val
 660 665 670
 Pro Thr Gly Pro Ser Val Gly Met Tyr Ser Pro Gly Tyr Ser Glu His
 675 680 685
 Ile Pro Leu Gly Lys Phe Tyr Asn Asn Arg Ala His Ser Asn Tyr Arg
 690 695 700
 Ala Gly Met Ile Ile Asp Asn Gly Val Lys Thr Thr Glu Ala Ser Ala
 705 710 715 720
 Lys Asp Lys Arg Pro Phe Leu Ser Ile Ile Ser Ala Arg Tyr Ser Pro
 725 730 735
 His Gln Asp Ala Asp Pro Leu Lys Pro Arg Glu Pro Ala Ile Ile Arg
 740 745 750
 His Phe Ile Ala Tyr Lys Asn Gln Asp His Gly Ala Trp Leu Arg Gly
 755 760 765
 Gly Asp Val Trp Leu Asp Ser Cys Arg Phe Ala Asp Asn Gly Ile Gly
 770 775 780
 Leu Thr Leu Ala Ser Gly Gly Thr Phe Pro Tyr Asp Asp Gly Ser Lys
 785 790 795 800
 Gln Glu Ile Lys Asn Ser Leu Phe Val Gly Glu Ser Gly Asn Val Gly
 805 810 815
 Thr Glu Met Met Asp Asn Arg Ile Trp Gly Pro Gly Gly Leu Asp His
 820 825 830
 Ser Gly Arg Thr Leu Pro Ile Gly Gln Asn Phe Pro Ile Arg Gly Ile
 835 840 845
 Gln Leu Tyr Asp Gly Pro Ile Asn Ile Gln Asn Cys Thr Phe Arg Lys
 850 855 860
 Phe Val Ala Leu Glu Gly Arg His Thr Ser Ala Leu Ala Phe Arg Leu
 865 870 875 880
 Asn Asn Ala Trp Gln Ser Cys Pro His Asn Asn Val Thr Gly Ile Ala
 885 890 895
 Phe Glu Asp Val Pro Ile Thr Ser Arg Val Phe Phe Gly Glu Pro Gly
 900 905 910

-continued

Pro Trp Phe Asn Gln Leu Asp Met Asp Gly Asp Lys Thr Ser Val Phe
 915 920 925
 His Asp Val Asp Gly Ser Val Ser Glu Tyr Pro Gly Ser Tyr Leu Thr
 930 935 940
 Lys Asn Asp Asn Trp Leu Val Arg His Pro Asp Cys Ile Asn Val Pro
 945 950 955 960
 Asp Trp Arg Gly Ala Ile Cys Ser Gly Cys Tyr Ala Gln Met Tyr Ile
 965 970 975
 Gln Ala Tyr Lys Thr Ser Asn Leu Arg Met Lys Ile Ile Lys Asn Asp
 980 985 990
 Phe Pro Ser His Pro Leu Tyr Leu Glu Gly Ala Leu Thr Arg Ser Thr
 995 1000 1005
 His Tyr Gln Gln Tyr Gln Pro Val Val Thr Leu Gln Lys Gly Tyr Thr
 1010 1015 1020
 Ile His Trp Asp Gln Thr Ala Pro Ala Glu Leu Ala Ile Trp Leu Ile
 1025 1030 1035 1040
 Asn Phe Asn Lys Gly Asp Trp Ile Arg Val Gly Leu Cys Tyr Pro Arg
 1045 1050 1055
 Gly Thr Thr Phe Ser Ile Leu Ser Asp Val His Asn Arg Leu Leu Lys
 1060 1065 1070
 Gln Thr Ser Lys Thr Gly Val Phe Val Arg Thr Leu Gln Met Asp Lys
 1075 1080 1085
 Val Glu Gln Ser Tyr Pro Gly Arg Ser His Tyr Tyr Trp Asp Glu Asp
 1090 1095 1100
 Ser Gly Leu Leu Phe Leu Lys Leu Lys Ala Gln Asn Glu Arg Glu Lys
 1105 1110 1115 1120
 Phe Ala Phe Cys Ser Met Lys Gly Cys Glu Arg Ile Lys Ile Lys Ala
 1125 1130 1135
 Leu Ile Pro Lys Asn Ala Gly Val Ser Asp Cys Thr Ala Thr Ala Tyr
 1140 1145 1150
 Pro Lys Phe Thr Glu Arg Ala Val Val Asp Val Pro Met Pro Lys Lys
 1155 1160 1165
 Leu Phe Gly Ser Gln Leu Lys Thr Lys Asp His Phe Leu Glu Val Lys
 1170 1175 1180
 Met Glu Ser Ser Lys Gln His Phe Phe His Leu Trp Asn Asp Phe Ala
 1185 1190 1195 1200
 Tyr Ile Glu Val Asp Gly Lys Lys Tyr Pro Ser Ser Glu Asp Gly Ile
 1205 1210 1215
 Gln Val Val Val Ile Asp Gly Asn Gln Gly Arg Val Val Ser His Thr
 1220 1225 1230
 Ser Phe Arg Asn Ser Ile Leu Gln Gly Ile Pro Trp Gln Leu Phe Asn
 1235 1240 1245
 Tyr Val Ala Thr Ile Pro Asp Asn Ser Ile Val Leu Met Ala Ser Lys
 1250 1255 1260
 Gly Arg Tyr Val Ser Arg Gly Pro Trp Thr Arg Val Leu Glu Lys Leu
 1265 1270 1275 1280
 Gly Ala Asp Arg Gly Leu Lys Leu Lys Glu Gln Met Ala Phe Val Gly
 1285 1290 1295
 Phe Lys Gly Ser Phe Arg Pro Ile Trp Val Thr Leu Asp Thr Glu Asp
 1300 1305 1310
 His Lys Ala Lys Ile Phe Gln Val Val Pro Ile Pro Val Val Lys Lys

-continued

1315	1320	1325
Lys Lys Leu 1330		
 <210> SEQ ID NO 2 <211> LENGTH: 1328 <212> TYPE: PRT <213> ORGANISM: Homo sapiens <400> SEQUENCE: 2		
Ala Gly Cys Pro Asp Gln Ser Pro Glu Leu Gln Pro Trp Asn Pro Gly 1 5 10 15		
His Asp Gln Asp His His Val His Ile Gly Gln Gly Lys Thr Leu Leu 20 25 30		
Leu Thr Ser Ser Ala Thr Val Tyr Ser Ile His Ile Ser Glu Gly Gly 35 40 45		
Lys Leu Val Ile Lys Asp His Asp Glu Pro Ile Val Leu Arg Thr Arg 50 55 60		
His Ile Leu Ile Asp Asn Gly Gly Glu Leu His Ala Gly Ser Ala Leu 65 70 75 80		
Cys Pro Phe Gln Gly Asn Phe Thr Ile Ile Leu Tyr Gly Arg Ala Asp 85 90 95		
Glu Gly Ile Gln Pro Asp Pro Tyr Tyr Gly Leu Lys Tyr Ile Gly Val 100 105 110		
Gly Lys Gly Gly Ala Leu Glu Leu His Gly Gln Lys Lys Leu Ser Trp 115 120 125		
Thr Phe Leu Asn Lys Thr Leu His Pro Gly Gly Met Ala Glu Gly Gly 130 135 140		
Tyr Phe Phe Glu Arg Ser Trp Gly His Arg Gly Val Ile Val His Val 145 150 155 160		
Ile Asp Pro Lys Ser Gly Thr Val Ile His Ser Asp Arg Phe Asp Thr 165 170 175		
Tyr Arg Ser Lys Lys Glu Ser Glu Arg Leu Val Gln Tyr Leu Asn Ala 180 185 190		
Val Pro Asp Gly Arg Ile Leu Ser Val Ala Val Asn Asp Glu Gly Ser 195 200 205		
Arg Asn Leu Asp Asp Met Ala Arg Lys Ala Met Thr Lys Leu Gly Ser 210 215 220		
Lys His Phe Leu His Leu Gly Phe Arg His Pro Trp Ser Phe Leu Thr 225 230 235 240		
Val Lys Gly Asn Pro Ser Ser Ser Val Glu Asp His Ile Glu Tyr His 245 250 255		
Gly His Arg Gly Ser Ala Ala Ala Arg Val Phe Lys Leu Phe Gln Thr 260 265 270		
Glu His Gly Glu Tyr Phe Asn Val Ser Leu Ser Ser Glu Trp Val Gln 275 280 285		
Asp Val Glu Trp Thr Glu Trp Phe Asp His Asp Lys Val Ser Gln Thr 290 295 300		
Lys Gly Gly Glu Lys Ile Ser Asp Leu Trp Lys Ala His Pro Gly Lys 305 310 315 320		
Ile Cys Asn Arg Pro Ile Asp Ile Gln Ala Thr Thr Met Asp Gly Val 325 330 335		

-continued

Asn Leu Ser Thr Glu Val Val Tyr Lys Lys Gly Gln Asp Tyr Arg Phe
 340 345 350

Ala Cys Tyr Asp Arg Gly Arg Ala Cys Arg Ser Tyr Arg Val Arg Phe
 355 360 365

Leu Cys Gly Lys Pro Val Arg Pro Lys Leu Thr Val Thr Ile Asp Thr
 370 375 380

Asn Val Asn Ser Thr Ile Leu Asn Leu Glu Asp Asn Val Gln Ser Trp
 385 390 395 400

Lys Pro Gly Asp Thr Leu Val Ile Ala Ser Thr Asp Tyr Ser Met Tyr
 405 410 415

Gln Ala Glu Glu Phe Gln Val Leu Pro Cys Arg Ser Cys Ala Pro Asn
 420 425 430

Gln Val Lys Val Ala Gly Lys Pro Met Tyr Leu His Ile Gly Glu Glu
 435 440 445

Ile Asp Gly Val Asp Met Arg Ala Glu Val Gly Leu Leu Ser Arg Asn
 450 455 460

Ile Ile Val Met Gly Glu Met Glu Asp Lys Cys Tyr Pro Tyr Arg Asn
 465 470 475 480

His Ile Cys Asn Phe Phe Asp Phe Asp Thr Phe Gly Gly His Ile Lys
 485 490 495

Phe Ala Leu Gly Phe Lys Ala Ala His Leu Glu Gly Thr Glu Leu Lys
 500 505 510

His Met Gly Gln Gln Leu Val Gly Gln Tyr Pro Ile His Phe His Leu
 515 520 525

Ala Gly Asp Val Asp Glu Arg Gly Gly Tyr Asp Pro Pro Thr Tyr Ile
 530 535 540

Arg Asp Leu Ser Ile His His Thr Phe Ser Arg Cys Val Thr Val His
 545 550 555 560

Gly Ser Asn Gly Leu Leu Ile Lys Asp Val Val Gly Tyr Asn Ser Leu
 565 570 575

Gly His Cys Phe Phe Thr Glu Asp Gly Pro Glu Glu Arg Asn Thr Phe
 580 585 590

Asp His Cys Leu Gly Leu Leu Val Lys Ser Gly Thr Leu Leu Pro Ser
 595 600 605

Asp Arg Asp Ser Lys Met Cys Lys Met Ile Thr Glu Asp Ser Tyr Pro
 610 615 620

Gly Tyr Ile Pro Lys Pro Arg Gln Asp Cys Asn Ala Val Ser Thr Phe
 625 630 635 640

Trp Met Ala Asn Pro Asn Asn Asn Leu Ile Asn Cys Ala Ala Ala Gly
 645 650 655

Ser Glu Glu Thr Gly Phe Trp Phe Ile Phe His His Val Pro Thr Gly
 660 665 670

Pro Ser Val Gly Met Tyr Ser Pro Gly Tyr Ser Glu His Ile Pro Leu
 675 680 685

Gly Lys Phe Tyr Asn Asn Arg Ala His Ser Asn Tyr Arg Ala Gly Met
 690 695 700

Ile Ile Asp Asn Gly Val Lys Thr Thr Glu Ala Ser Ala Lys Asp Lys
 705 710 715 720

Arg Pro Phe Leu Ser Ile Ile Ser Ala Arg Tyr Ser Pro His Gln Asp
 725 730 735

Ala Asp Pro Leu Lys Pro Arg Glu Pro Ala Ile Ile Arg His Phe Ile

-continued

740				745				750							
Ala	Tyr	Lys	Asn	Gln	Asp	His	Gly	Ala	Trp	Leu	Arg	Gly	Gly	Asp	Val
		755					760					765			
Trp	Leu	Asp	Ser	Cys	Arg	Phe	Ala	Asp	Asn	Gly	Ile	Gly	Leu	Thr	Leu
	770					775					780				
Ala	Ser	Gly	Gly	Thr	Phe	Pro	Tyr	Asp	Asp	Gly	Ser	Lys	Gln	Glu	Ile
785					790					795					800
Lys	Asn	Ser	Leu	Phe	Val	Gly	Glu	Ser	Gly	Asn	Val	Gly	Thr	Glu	Met
			805						810						815
Met	Asp	Asn	Arg	Ile	Trp	Gly	Pro	Gly	Gly	Leu	Asp	His	Ser	Gly	Arg
			820						825					830	
Thr	Leu	Pro	Ile	Gly	Gln	Asn	Phe	Pro	Ile	Arg	Gly	Ile	Gln	Leu	Tyr
		835					840					845			
Asp	Gly	Pro	Ile	Asn	Ile	Gln	Asn	Cys	Thr	Phe	Arg	Lys	Phe	Val	Ala
	850					855						860			
Leu	Glu	Gly	Arg	His	Thr	Ser	Ala	Leu	Ala	Phe	Arg	Leu	Asn	Asn	Ala
865					870					875					880
Trp	Gln	Ser	Cys	Pro	His	Asn	Asn	Val	Thr	Gly	Ile	Ala	Phe	Glu	Asp
				885					890						895
Val	Pro	Ile	Thr	Ser	Arg	Val	Phe	Phe	Gly	Glu	Pro	Gly	Pro	Trp	Phe
			900					905						910	
Asn	Gln	Leu	Asp	Met	Asp	Gly	Asp	Lys	Thr	Ser	Val	Phe	His	Asp	Val
		915					920						925		
Asp	Gly	Ser	Val	Ser	Glu	Tyr	Pro	Gly	Ser	Tyr	Leu	Thr	Lys	Asn	Asp
	930					935						940			
Asn	Trp	Leu	Val	Arg	His	Pro	Asp	Cys	Ile	Asn	Val	Pro	Asp	Trp	Arg
945					950					955					960
Gly	Ala	Ile	Cys	Ser	Gly	Cys	Tyr	Ala	Gln	Met	Tyr	Ile	Gln	Ala	Tyr
			965						970					975	
Lys	Thr	Ser	Asn	Leu	Arg	Met	Lys	Ile	Ile	Lys	Asn	Asp	Phe	Pro	Ser
			980					985					990		
His	Pro	Leu	Tyr	Leu	Glu	Gly	Ala	Leu	Thr	Arg	Ser	Thr	His	Tyr	Gln
		995					1000						1005		
Gln	Tyr	Gln	Pro	Val	Val	Thr	Leu	Gln	Lys	Gly	Tyr	Thr	Ile	His	Trp
	1010					1015						1020			
Asp	Gln	Thr	Ala	Pro	Ala	Glu	Leu	Ala	Ile	Trp	Leu	Ile	Asn	Phe	Asn
1025				1030						1035					1040
Lys	Gly	Asp	Trp	Ile	Arg	Val	Gly	Leu	Cys	Tyr	Pro	Arg	Gly	Thr	Thr
				1045					1050					1055	
Phe	Ser	Ile	Leu	Ser	Asp	Val	His	Asn	Arg	Leu	Leu	Lys	Gln	Thr	Ser
			1060					1065						1070	
Lys	Thr	Gly	Val	Phe	Val	Arg	Thr	Leu	Gln	Met	Asp	Lys	Val	Glu	Gln
	1075						1080					1085			
Ser	Tyr	Pro	Gly	Arg	Ser	His	Tyr	Tyr	Trp	Asp	Glu	Asp	Ser	Gly	Leu
	1090					1095						1100			
Leu	Phe	Leu	Lys	Leu	Lys	Ala	Gln	Asn	Glu	Arg	Glu	Lys	Phe	Ala	Phe
1105					1110					1115					1120
Cys	Ser	Met	Lys	Gly	Cys	Glu	Arg	Ile	Lys	Ile	Lys	Ala	Leu	Ile	Pro
			1125						1130					1135	
Lys	Asn	Ala	Gly	Val	Ser	Asp	Cys	Thr	Ala	Thr	Ala	Tyr	Pro	Lys	Phe
		1140						1145						1150	

-continued

Thr Glu Arg Ala Val Val Asp Val Pro Met Pro Lys Lys Leu Phe Gly
 1155 1160 1165
 Ser Gln Leu Lys Thr Lys Asp His Phe Leu Glu Val Lys Met Glu Ser
 1170 1175 1180
 Ser Lys Gln His Phe Phe His Leu Trp Asn Asp Phe Ala Tyr Ile Glu
 1185 1190 1195 1200
 Val Asp Gly Lys Lys Tyr Pro Ser Ser Glu Asp Gly Ile Gln Val Val
 1205 1210 1215
 Val Ile Asp Gly Asn Gln Gly Arg Val Val Ser His Thr Ser Phe Arg
 1220 1225 1230
 Asn Ser Ile Leu Gln Gly Ile Pro Trp Gln Leu Phe Asn Tyr Val Ala
 1235 1240 1245
 Thr Ile Pro Asp Asn Ser Ile Val Leu Met Ala Ser Lys Gly Arg Tyr
 1250 1255 1260
 Val Ser Arg Gly Pro Trp Thr Arg Val Leu Glu Lys Leu Gly Ala Asp
 1265 1270 1275 1280
 Arg Gly Leu Lys Leu Lys Glu Gln Met Ala Phe Val Gly Phe Lys Gly
 1285 1290 1295
 Ser Phe Arg Pro Ile Trp Val Thr Leu Asp Thr Glu Asp His Lys Ala
 1300 1305 1310
 Lys Ile Phe Gln Val Val Pro Ile Pro Val Val Lys Lys Lys Lys Leu
 1315 1320 1325

<210> SEQ ID NO 3
 <211> LENGTH: 732
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 3

Leu Gln Glu Val His Val Ser Lys Glu Thr Ile Gly Lys Ile Ser Ala
 1 5 10 15
 Ala Ser Lys Met Met Trp Cys Ser Ala Ala Val Asp Ile Met Phe Leu
 20 25 30
 Leu Asp Gly Ser Asn Ser Val Gly Lys Gly Ser Phe Glu Arg Ser Lys
 35 40 45
 His Phe Ala Ile Thr Val Cys Asp Gly Leu Asp Ile Ser Pro Glu Arg
 50 55 60
 Val Arg Val Gly Ala Phe Gln Phe Ser Ser Thr Pro His Leu Glu Phe
 65 70 75 80
 Pro Leu Asp Ser Phe Ser Thr Gln Gln Glu Val Lys Ala Arg Ile Lys
 85 90 95
 Arg Met Val Phe Lys Gly Gly Arg Thr Glu Thr Glu Leu Ala Leu Lys
 100 105 110
 Tyr Leu Leu His Arg Gly Leu Pro Gly Gly Arg Asn Ala Ser Val Pro
 115 120 125
 Gln Ile Leu Ile Ile Val Thr Asp Gly Lys Ser Gln Gly Asp Val Ala
 130 135 140
 Leu Pro Ser Lys Gln Leu Lys Glu Arg Gly Val Thr Val Phe Ala Val
 145 150 155 160
 Gly Val Arg Phe Pro Arg Trp Glu Glu Leu His Ala Leu Ala Ser Glu
 165 170 175
 Pro Arg Gly Gln His Val Leu Leu Ala Glu Gln Val Glu Asp Ala Thr

-continued

180					185					190					
Asn	Gly	Leu	Phe	Ser	Thr	Leu	Ser	Ser	Ser	Ala	Ile	Cys	Ser	Ser	Ala
	195						200					205			
Thr	Pro	Asp	Cys	Arg	Val	Glu	Ala	His	Pro	Cys	Glu	His	Arg	Thr	Leu
	210					215					220				
Glu	Met	Val	Arg	Glu	Phe	Ala	Gly	Asn	Ala	Pro	Cys	Trp	Arg	Gly	Ser
	225				230					235					240
Arg	Arg	Thr	Leu	Ala	Val	Leu	Ala	Ala	His	Cys	Pro	Phe	Tyr	Ser	Trp
				245					250						255
Lys	Arg	Val	Phe	Leu	Thr	His	Pro	Ala	Thr	Cys	Tyr	Arg	Thr	Thr	Cys
			260					265						270	
Pro	Gly	Pro	Cys	Asp	Ser	Gln	Pro	Cys	Gln	Asn	Gly	Gly	Thr	Cys	Val
		275					280					285			
Pro	Glu	Gly	Leu	Asp	Gly	Tyr	Gln	Cys	Leu	Cys	Pro	Leu	Ala	Phe	Gly
	290					295					300				
Gly	Glu	Ala	Asn	Cys	Ala	Leu	Lys	Leu	Ser	Leu	Glu	Cys	Arg	Val	Asp
	305				310					315					320
Leu	Leu	Phe	Leu	Leu	Asp	Ser	Ser	Ala	Gly	Thr	Thr	Leu	Asp	Gly	Phe
				325					330					335	
Leu	Arg	Ala	Lys	Val	Phe	Val	Lys	Arg	Phe	Val	Arg	Ala	Val	Leu	Ser
			340					345					350		
Glu	Asp	Ser	Arg	Ala	Arg	Val	Gly	Val	Ala	Thr	Tyr	Ser	Arg	Glu	Leu
		355					360					365			
Leu	Val	Ala	Val	Pro	Val	Gly	Glu	Tyr	Gln	Asp	Val	Pro	Asp	Leu	Val
	370					375					380				
Trp	Ser	Leu	Asp	Gly	Ile	Pro	Phe	Arg	Gly	Gly	Pro	Thr	Leu	Thr	Gly
	385				390					395					400
Ser	Ala	Leu	Arg	Gln	Ala	Ala	Glu	Arg	Gly	Phe	Gly	Ser	Ala	Thr	Arg
				405					410					415	
Thr	Gly	Gln	Asp	Arg	Pro	Arg	Arg	Val	Val	Val	Leu	Leu	Thr	Glu	Ser
			420					425					430		
His	Ser	Glu	Asp	Glu	Val	Ala	Gly	Pro	Ala	Arg	His	Ala	Arg	Ala	Arg
		435					440					445			
Glu	Leu	Leu	Leu	Leu	Gly	Val	Gly	Ser	Glu	Ala	Val	Arg	Ala	Glu	Leu
	450				455						460				
Glu	Glu	Ile	Thr	Gly	Ser	Pro	Lys	His	Val	Met	Val	Tyr	Ser	Asp	Pro
	465				470					475					480
Gln	Asp	Leu	Phe	Asn	Gln	Ile	Pro	Glu	Leu	Gln	Gly	Lys	Leu	Cys	Ser
				485					490					495	
Arg	Gln	Arg	Pro	Gly	Cys	Arg	Thr	Gln	Ala	Leu	Asp	Leu	Val	Phe	Met
			500					505					510		
Leu	Asp	Thr	Ser	Ala	Ser	Val	Gly	Pro	Glu	Asn	Phe	Ala	Gln	Met	Gln
		515					520					525			
Ser	Phe	Val	Arg	Ser	Cys	Ala	Leu	Gln	Phe	Glu	Val	Asn	Pro	Asp	Val
	530					535					540				
Thr	Gln	Val	Gly	Leu	Val	Val	Tyr	Gly	Ser	Gln	Val	Gln	Thr	Ala	Phe
	545				550					555					560
Gly	Leu	Asp	Thr	Lys	Pro	Thr	Arg	Ala	Ala	Met	Leu	Arg	Ala	Ile	Ser
				565					570					575	
Gln	Ala	Pro	Tyr	Leu	Gly	Gly	Val	Gly	Ser	Ala	Gly	Thr	Ala	Leu	Leu
			580					585					590		

-continued

His Ile Tyr Asp Lys Val Met Thr Val Gln Arg Gly Ala Arg Pro Gly
 595 600 605
 Val Pro Lys Ala Val Val Val Leu Thr Gly Gly Arg Gly Ala Glu Asp
 610 615 620
 Ala Ala Val Pro Ala Gln Lys Leu Arg Asn Asn Gly Ile Ser Val Leu
 625 630 635 640
 Val Val Gly Val Gly Pro Val Leu Ser Glu Gly Leu Arg Arg Leu Ala
 645 650 655
 Gly Pro Arg Asp Ser Leu Ile His Val Ala Ala Tyr Ala Asp Leu Arg
 660 665 670
 Tyr His Gln Asp Val Leu Ile Glu Trp Leu Cys Gly Glu Ala Lys Gln
 675 680 685
 Pro Val Asn Leu Cys Lys Pro Ser Pro Cys Met Asn Glu Gly Ser Cys
 690 695 700
 Val Leu Gln Asn Gly Ser Tyr Arg Cys Lys Cys Arg Asp Gly Trp Glu
 705 710 715 720
 Gly Pro His Cys Glu Asn Arg Phe Leu Arg Arg Pro
 725 730

<210> SEQ ID NO 4
 <211> LENGTH: 4171
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 4

cgtgacactg tctcggctac agaccacag ggagcacact gccaggatgg gagctgctgg 60
 gaggcaggac ttctcttcca aggccatgct gaccatcagc tggetcactc tgacctgctt 120
 ccctggggcc acatccacag tggctgctgg gtgccctgac cagagccctg agttgcaacc 180
 ctggaaccct ggccatgacc aagaccacca tgtgcatatc ggccagggca agacactgct 240
 gctcactctc tctgccacgg tctattccat ccacatctca gagggaggca agctgggtcat 300
 taaagaccac gacgagccga ttgttttgcg aaccggcac atcctgattg acaacggagg 360
 agagctgcat gctgggagtg ccctctgccc ttccagggc aattccacca tcattttgta 420
 tggaagggct gatgaaggta ttcagccgga tccttactat ggtctgaagt acattggggt 480
 tggtaaagga ggcgctcttg agttgcatgg acagaaaaag ctctctgga cattttctgaa 540
 caagaccctt caccagggtg gcatggcaga aggaggctat tttttgaaa ggagctgggg 600
 ccaccgtgga gttattgttc atgtcatoga ccccaaatca ggcacagtca tccattctga 660
 cgggtttgac acctatagat ccaagaaaga gactgaacgt ctggtccagt atttgaacgc 720
 ggtgcccgat ggcaggatcc tttctgttgc agtgaatgat gaaggttctc gaaatctgga 780
 tgacatggcc aggaaggcga tgaccaaatt ggaagcaaa cacttctctc accttgatt 840
 tagacaccct tggagttttc taactgtgaa aggaaatcca tcattctcag tggaagacca 900
 tattgaatat catggacatc gaggctctgc tctgcccgg gtattcaaat tgttccagac 960
 agagcatggc gaatatttca atgtttcttt gtccagtgag tgggttcaag acgtggagtg 1020
 gacggagtg ttcgatcatg ataaagtatc tcagactaaa ggtggggaga aaatttcaga 1080
 cctctggaaa gctcaccagg gaaaatatg caatcgtccc attgatatac aggccactac 1140
 aatggatgga gttaacctca gcaccagggt tgtctacaaa aaaggccagg attataggtt 1200

-continued

tgcttgctac gaccggggca gagcctgccc gagctaccgt gtaegggtcc tctgtgggaa	1260
gctgtgagg cccaaactca cagtcaccat tgacaccaat gtgaacagca ccattctgaa	1320
cttgagggat aatgtacagt catggaaacc tggagatacc ctggtcattg ccagtactga	1380
ttactccatg taccaggcag aagagttcca ggtgcttccc tgcagatcct gcgcccccaa	1440
ccaggtcaaa gtggcaggga aaccaatgta cctgcacatc ggggaggaga tagacggcgt	1500
ggacatgccc gcggaggttg ggcttctgag ccggaacatc atagtgatgg gggagatgga	1560
ggacaaatgc taccctaca gaaaccacat ctgcaatttc tttgacttcg atacctttgg	1620
ggccacatc aagtttgcct tgggatttaa ggcagcacac ttggagggca cggagctgaa	1680
gcatatggga cagcagctgg tgggtcagta cccgattcac ttccacctgg ccggtgatgt	1740
agacgaaagg ggaggttatg acccaccac atacatcagg gacctctcca tccatcatac	1800
attctctcgc tgcgtcacag tccatggctc caatggcttg ttgatcaagg acgtttgagg	1860
ctataactct ttgggacctc gcttcttcac ggaagatggg ccggaggaac gcaacacttt	1920
tgaccactgt cttggcctcc ttgtcaagtc tggaaacctc ctcccctcgg accgtgacag	1980
caagatgtgc aagatgatca cagaggactc ctaccagggt tacatcccca agcccaggca	2040
agactgcaat gctgtgtcca ccttctggat ggccaatccc aacaacaacc tcatcaactg	2100
tgccgctgca ggatctgagg aaactggatt ttggtttatt ttccaccacg taccaacggg	2160
cccctccgtg ggaatgtact ccccaggtta ttcagagcac attccactgg gaaaattcta	2220
taacaaccga gcacattcca actaccgggc tggcatgatc atagacaacg gagtcaaac	2280
caccgaggcc tctgccaaagg acaagcggcc gttcctctca atcatctctg ccagatacag	2340
ccctcaccag gacgccgacc cgctgaagcc ccgggagccg gccatcatca gacacttcat	2400
tgcctacaag aaccaggacc acggggcctg gctgcgcccg ggggatgtgt ggcctggacag	2460
ctgccggttt gctgacaatg gcattggcct gaccctggcc agtgggtgaa ccttcccgtg	2520
tgacgacggc tccaagcaag agataaagaa cagcttgttt gttggcgaga gtggcaactg	2580
ggggacggaa atgatggaca ataggatctg gggccctggc ggcttgacc atagcgggag	2640
gaccctccct ataggccaga atttccaat tagaggaatt cagttatatg atggcccat	2700
caacatccaa aactgcactt tccgaaagt tgtggccctg gagggccggc acaccagcgc	2760
cctggccttc cgctgaata atgcctggca gagctgcccc cataacaacg tgaccggcat	2820
tgcctttgag gacgttccga ttaactccag agtgttcttc ggagagcctg ggcctggtt	2880
caaccagctg gacatggatg gggataagac atctgtgttc catgacgtcg acggctccgt	2940
gtccgagtac cctggctcct acctcacgaa gaatgacaac tggctggtcc ggcaccaga	3000
ctgcatcaat gttcccact ggagaggggc catttgacgt ggggtgctatg cacagatgta	3060
cattcaagcc tacaagacca gtaacctgcg aatgaagatc atcaagaatg acttccccag	3120
ccaccctctt taectggagg gggcgctcac caggagcacc cattaccagc aataccaacc	3180
ggttgtcacc ctgcagaagg gctacaccat ccaactgggac cagacggccc ccgccaact	3240
cgccatctgg ctcatcaact tcaacaaggc cgactggatc cgagtggggc tctgctaccc	3300
gcgaggcacc acattcteca tccctctgga tgttcacaat cgcctgctga agcaaacgtc	3360
caagacgggc gtctctgta ggcacttga gatggacaaa gtggagcaga gctaccctgg	3420
caggagccac tactactggg acgaggactc agggctggtt ttcctgaagc tgaaagctca	3480

-continued

```

gaacgagaga gagaagtttg cttctgctc catgaaaggc tgtgagagga taaagattaa 3540
agctctgatt ccaaagaacg caggcgtcag tgactgcaca gccacagctt accccaagtt 3600
caccgagagg gctgtcgtag acgtgccgat gcccaagaag ctctttgggt ctcagctgaa 3660
aacaaggac catttcttgg aggtgaagat ggagagtcc aagcagcact tcttccacct 3720
ctggaacgac ttcgcttaca ttgaagtgga tgggaagaag taccocagtt cggaggatgg 3780
catccaggtg gtggtgattg acgggaacca agggcgcgtg gtgagccaca cgagcttcag 3840
gaactccatt ctgcaaggca taccatggca gcttttcaac tatgtggcga ccatccctga 3900
caattccata gtgcttatgg catcaaaggg aagatacgtc tccagaggcc catggaccag 3960
agtgtggaa aagcttgggg cagacagggg tctcaagttg aaagagcaaa tggcattcgt 4020
tggttcaaa ggcagcttcc ggcccatctg ggtgacactg gacactgagg atcacaagc 4080
caaaatctc caagttgtgc ccatccctgt ggtgaagaag aagaagttgt gaggacagct 4140
gccgcccggg gccacctcgt ggtagactat g 4171

```

<210> SEQ ID NO 5

<211> LENGTH: 2810

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 5

```

gccccctggc cegagccgcg cccgggtctg tgagtagagc cgccccggca cegagcgtg 60
gtcgccgctc tcttccgtt atatcaacat gcccccttc ctggtgctgg aagccgtctg 120
tgtttctctg ttttccagag tgccccatc tctccctctc caggaagtcc atgtaagcaa 180
agaaaccatc ggaagattt cagctgccag caaaatgatg tgggtctcgg ctgcagtgga 240
catcatgttt ctgtagatg ggtctaacag cgtcgggaaa gggagcttg aaaggtccaa 300
gcactttgcc atcacagtct gtgacggctt ggacatcagc cccgagaggg tcagagtggg 360
agcattccag ttcagtcca ctctcatct ggaattcccc ttggattcat tttcaacca 420
acaggaagtg aaggcaagaa tcaagaggat ggttttcaaa ggaggcgcga cggagacgga 480
acttctctg aaatacctc tgcaacagag gttgcctgga ggcagaaatg cttctgtgcc 540
ccagatctc atcatcgtca ctgatgggaa gtcccagggg gatgtggcac tgccatccaa 600
gcagctgaag gaaaggggtg tcaactgtgt tgctgtgggg gtcaggttcc ccaggtggga 660
ggagctgcat gcactggcca gcgagcctag agggcagcac gtgctgttg ctgagcaggt 720
ggaggatgcc accaacggcc tcttcagcac cctcagcagc tcggccatct gctccagcgc 780
cacgccagac tgcagggtcg aggtcaacc ctgtgagcac aggaagctgg agatggtccg 840
ggagtctgct ggcaatgccc catgctggag aggatcgcgg cggacccttg cgtgctggc 900
tgcaactgt ccttctaca gctggaagag agtgttccta acccaccctg ccacctgcta 960
caggaccacc tgcccaggcc cctgtgactc gcagccctgc cagaatggag gcacatgtgt 1020
tccagaagga ctggacggct accagtgcct ctgccgctg gcctttggag gggaggctaa 1080
ctgtgccctg aagctgagcc tggaaatgca ggtcgacctc ctcttctgc tggacagctc 1140
tgccggcacc actctggagc gcttctgctg ggccaaagtc ttcgtgaagc ggtttgtgctg 1200
ggccgtgctg agcaggactc ctccggcccg agtgggtgtg gccacatata gcagggagct 1260
gctggtggcg gtgcctgtgg gggagtacca ggatgtgcct gacctggtct ggagcctcga 1320

```

-continued

tggeattccc	ttccgtggtg	gccccaccct	gacgggcagt	gccttgccgc	aggcggcaga	1380
gogtggcttc	gggagcgcca	ccaggacagg	ccaggaccgg	ccacgtagag	tggtggtttt	1440
gctcactgag	tcacactccg	aggatgaggt	tgccggccca	gcccgtcacg	caagggcgcg	1500
agagctgctc	ctgctgggtg	taggcagtga	ggccgtgcgg	gcagagctgg	aggagatcac	1560
aggcagccca	aagcatgtga	tggtctactc	ggatcctcag	gatctgttca	accaaaatccc	1620
tgagctgcag	gggaagctgt	gcagccggca	gcggccaggg	tgccggacac	aagccctgga	1680
cctcgtcttc	atgttgga	cctctgcctc	agtagggccc	gagaattttg	ctcagatgca	1740
gagctttgtg	agaagctgtg	ccctccagtt	tgaggtgaac	cctgacgtga	cacaggtcgg	1800
cctggtgggtg	tatggcagcc	agggtcagac	tgccctcggg	ctggacacca	aaccaccccg	1860
ggctgcgatg	ctgcgggcca	ttagccaggc	cccctaccta	gggggggtgg	gctcagccgg	1920
caccgccctg	ctgcacatct	atgacaaagt	gatgaccgtc	cagaggggtg	cccggcctgg	1980
tgtcccaaaa	gctgtgggtg	tgctcacagg	cgggagaggc	gcagaggatg	cagccgttcc	2040
tgcccagaag	ctgaggaaca	atggcatctc	tgtcttggtc	gtggggctgg	ggcctgtcct	2100
aagtgagggt	ctgcggaggc	ttgcaggctc	ccgggattcc	ctgatccacg	tggcagctta	2160
cgccgacctg	cgttaccacc	aggacgtgct	cattgagtgg	ctgtgtggag	aagccaagca	2220
gccagtcaac	ctctgcaaac	ccagcccgtg	catgaatgag	ggcagctgcg	tcctgcagaa	2280
tgggagctac	cgtgcaagt	gtcgggatgg	ctgggagggc	cccactgcg	agaaccgatt	2340
cttgagacgc	ccctgaggca	catggctccc	gtgcaggagg	gcagcagccg	taccctccc	2400
agcaactaca	gagaaggcct	gggcactgaa	atgggtccta	cctctggaa	tgtctgtgcc	2460
ccaggtcctt	agaatgtctg	cttcccgcgc	tggccaggac	cactattctc	actgaggag	2520
gaggatgtcc	caactgcagc	catgctgctt	agagacaaga	aagcagctga	tgtcaccac	2580
aaacgatgtt	ggtgaaaaat	tttgatgtgt	aagtaataac	ccactttctg	tacctgctgt	2640
gccttgttga	ggctatgtca	tctgccacct	ttcccttgag	gataaacaag	gggtcctgaa	2700
gacttaaatt	tagccgacctg	acgttccctt	gcacacaatc	aatgctcgcc	agaatgttgt	2760
tgacacagta	atgccagca	gaggccttta	ctagagcatc	ctttggacgg		2810

<210> SEQ ID NO 6

<211> LENGTH: 1524

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 6

gcagagcaca	gcacgtcgcg	gaccagactc	gtctcaggcc	agttgcagcc	ttctcagcca	60
aacgcccacc	aaggaaaact	cactaccatg	agaattgcag	tgatttgctt	ttgcctccta	120
ggcatcacct	gtgccatacc	agttaaacag	gctgattctg	gaagttctga	ggaaaagcag	180
ctttacaaca	aataccaga	tgctgtggcc	acatggctaa	accctgaccc	atctcagaag	240
cagaatctcc	tagccccaca	gacccttcca	agtaagtcca	acgaaagcca	tgaccacatg	300
gatgatatgg	atgatgaaga	tgatgatgac	catgtggaca	gccaggactc	cattgactcg	360
aacgactctg	atgatgtaga	tgacactgat	gattctcacc	agtctgatga	gtctcaccat	420
tctgatgaat	ctgatgaact	ggctcactgat	tttcccacgg	acctgccagc	aaccgaagtt	480
ttcactccag	ttgtccccc	agtagacaca	tatgatggcc	gaggtgatag	tgtggtttat	540

-continued

```

ggactgaggt caaaatctaa gaagtttcgc agacctgaca tccagtaccc tgatgctaca 600
gacgaggaca tcacctcaca catggaaagc gaggagtga atggtgcata caaggccatc 660
cccgttcccc aggacctgaa cgcgccttct gattgggaca gccgtgggaa ggacagttat 720
gaaacgagtc agctggatga ccagagtgct gaaacccaca gccacaagca gtccagatta 780
tataagcgga aagccaatga tgagagcaat gagcattccg atgtgattga tagtcaggaa 840
ctttccaaag tcagccgtga attccacagc catgaatttc acagccatga agatatgctg 900
gttgtagacc ccaaaagtaa ggaagaagat aaacacctga aatttcgtat ttctcatgaa 960
ttagatagtg catcttctga ggtcaattaa aaggagaaaa aatacaattt ctcactttgc 1020
atthagtcaa aagaaaaaat gctttatagc aaaatgaaag agaacatgaa atgcttcttt 1080
ctcagtttat tggttgaatg tgtatctatt tgagtctgga aataactaat gtgtttgata 1140
attagtttag tttgtggctt catggaaact ccctgtaaac taaaagcttc agggttatgt 1200
ctatgttcat tctatagaag aaatgcaaac taccactgta ttttaattatt tgttattctc 1260
tcatgaatag aaatttatgt agaagcaaac aaaatacttt taccactta aaaagagaat 1320
ataacatttt atgtcactat aatcttttgt tttttaagtt agtgtatatt ttgttgatg 1380
tatctttttg tgggtggaat aaatctttta tcttgaatgt aataagaatt tgggtggtgc 1440
aattgcttat ttgttttccc acggttgtcc agcaattaat aaaacataac cttttttact 1500
gcctaaaaaa aaaaaaaaaa aaaa 1524

```

<210> SEQ ID NO 7

<211> LENGTH: 3205

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 7

```

aaaggggcaa gagctgagcg gaacaccggc ccgcccgtgc ggcagctgct tcaccctct 60
ctctgcagcc atggggctcc ctctggacc tctcgcgtct ctctccttc tccaggtttg 120
ctggctgcag tgcgcggcct ccgagccgtg ccgggcccgtc ttcagggagg ctgaagtgc 180
cttgaggcg ggaggcgcgg agcaggagcc cggccaggcg ctggggaaag tattcatggg 240
ctgccctggg caagagccag ctctgtttag cactgataat gatgactca ctgtgcggaa 300
tggcgagaca gtccaggaaa gaaggtcact gaaggaaagg aatccattga agatcttccc 360
atccaaaagt atcttacgaa gacacaagag agattgggtg gttgctcaa tatctgtccc 420
tgaaaatggc aagggtccct tccccagag actgaatcag ctcaagtcta ataaagatag 480
agacaccaag attttctaca goatcacggg gccgggggca gacagcccc ctgagggtgt 540
ctctcgtgta gagaaggaga caggctggtt gttgttgaat aagccactgg accgggagga 600
gattgccaag tatgagctct ttggccacgc tgtgtcagag aatgggtcct cagtggagga 660
ccccatgaac atctccatca tcgtgaccga ccagaatgac cacaagccca agtttaccca 720
ggacaccttc cgagggagtg tcttagaggg agtccctacca ggtacttctg tgatgcaggt 780
gacagccacg gatgaggatg atgcatceta cacctacaat ggggtggttg cttactccat 840
ccatagccaa gaaccaagg acccacacga cctcatgttc accattcacc ggagcacagg 900
caccatcagc gtcactccta ttggcctgga ccgggaaaaa gtccttgagt acacactgac 960
catccaggcc acagacatgg atggggacgg ctccaccacc acggcagtgg cagtagtgga 1020

```

-continued

gatccttgat gccaatgaca atgetcccat gtttgacccc cagaagtacg aggcccatgt	1080
gctgagaat gcagtgggcc atgaggtgca gaggctgacg gtcactgatc tggacgcccc	1140
caactcacca gcgtggcgtg ccacctacct tatcatgggc ggtgacgacg gggaccattt	1200
taccatcacc acccaccctg agagcaacca gggcatcctg acaaccagga agggtttggga	1260
ttttgaggcc aaaaaccagc acaccctgta cgttgaagtg accaacgagg ccccttttgt	1320
gctgaagctc ccaacctcca cagccaccat agtgggccac gtggaggatg tgaatgaggc	1380
acctgtgttt gtcccacct ccaaagtcgt tgaggccag gagggcatcc ccaactgggga	1440
gcctgtgtgt gtctacactg cagaagaccc tgacaaggag aatcaaaaga tcagctaccg	1500
catcctgaga gacccagcag ggtggctagc catggacca gacagtgggc aggtcacagc	1560
tgtgggcacc ctgcaccgtg aggatgagca gtttgtgagg aacaacatct atgaagtcac	1620
ggtcttgacc atggacaatg gaagccctcc caccactggc acgggaaccc ttctgctaac	1680
actgattgat gtcaatgacc atggcccagt ccctgagccc cgctcagatca ccatctgcaa	1740
ccaaagccct gtgcgccagg tgctgaacat cacggacaag gacctgtctc cccacacctc	1800
ccctttccag gccagctca cagatgactc agacatctac tggacggcag aggtcaacga	1860
ggaaggtgac acagtggtct tgtccctgaa gaagtccctg aagcaggata catatgacgt	1920
gcacccttct ctgtctgacc atggcaacaa agagcagctg acggtgatca gggccactgt	1980
gtgcgactgc catggccatg tcgaaacctg ccctggacce tggaaaggag gtttcatcct	2040
ccctgtgctg ggggctgtcc tggetctget gttcctcctg ctggtgctgc ttttgttgg	2100
gagaaagaag cggaaagatca agggcccct cctactccca gaagatgaca cccgtgacaa	2160
cgtctctac tatggcgaag aggggggtgg cgaagaggac caggactatg acatcaccca	2220
gctccaccga ggtctggagg ccaggccgga ggtggttctc cgcaatgacg tggcaccaac	2280
catcatcccg acaccatgt accgtcctcg gccagccaac ccagatgaaa tcggcaactt	2340
tataattgag aacctgaagg cggctaacac agaccccaca gcccgcctc acgacaccct	2400
cttggtgttc gactatgagg gcagcggctc cgacgcccgc tccctgagct cctcaccctc	2460
ctccgcctcc gaccaagacc aagattacga ttatctgaac gagtggggca gccgcttcaa	2520
gaagctggca gacatgtacg gtggcgggga ggacgactag gcggcctgcc tgcagggctg	2580
gggaccaaac gtcaggccac agagcatctc caaggggtct cagttcccc ttcagctgag	2640
gacttcggag cttgtcagga agtggccgta gcaacttggc ggagacaggc tatgagtctg	2700
acgttagagt ggttgettcc ttagccttcc aggatggagg aatgtgggca gtttgacttc	2760
agcactgaaa acctctccac ctgggcccag gttgcctcag agggcaagtt tccagaagcc	2820
tcttacctgc cgtaaaatgc tcaaccctgt gtccctgggc tgggcctgct gtgactgacc	2880
tacagtggac tttctctctg gaatggaacc ttcttaggcc tccgtgtgca acttaatttt	2940
tttttttaat gctatcttca aaacgtaga gaaagtctt caaaagtgca gccagagct	3000
gctgggcccc ctggccgtcc tgcatttctg gtttcagac cccaatgctc cccattcgga	3060
tggatctctg cgtttttata ctgagtgtgc ctaggttgcc ccttattttt tattttccct	3120
gttgcgttgc tatagatgaa gggtgaggac aatcgtgtat atgtactaga acttttttat	3180
taaagaaact tttcccagaa aaaaa	3205

<210> SEQ ID NO 8

-continued

```

<211> LENGTH: 2603
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 8
atgaagcacc tgaagcggtg gtggtcggcc ggcggcggcc tcctgcacct caccctcctg    60
ctgagcttgg cggggctccg cgtagacctg gatctttacc tgetgctgcc gccgcccacc    120
ctgctgcagg acgagctgct gttcctgggc ggcccggcca gctccgccta cgcgctcagc    180
cccttctcgg cctcgggagg gtggggggcg gcggggccact tgcaccccaa gggccgggag    240
ctggaccctg ccgcccggcc cgagggccag ctgctccggg aggtgcgcgc gctcggggtc    300
cccttcgtcc ctgcaccag cgtggatgca tggctggtgc acagcgtggc tgcggggagc    360
gcggacgagg cccaegggtc gctcggggcc gccgcccgct cgteccaccg aggagccggc    420
gccagcgtgg acggcggcag ccaggctgtg cagggggggc gcgggggacc ccgagcggct    480
cggagtggcc ccttggacgc cggggaagag gagaaggcac ccgcggaacc gacggctcag    540
gtgcgggacg ctggcggatg tgcgagcag gagaatgggg tactaagaga aaagcacgaa    600
gctgtggatc atagttccca gcatgaggaa aatgaagaaa ggggtgacgc ccagaaggag    660
aactcacttc agcagaatga tgatgatgaa aacaaaatag cagagaaacc tgactgggag    720
gcagaaaaga ccactgaatc tagaaatgag agacatctga atgggacaga tacttctttc    780
tctctggaag acttattcca gttgctttca tcacgcctg aaaattcact ggagggcacc    840
tcattgggag atattcctct tccaggcagt atcagtgatg gcattgaattc ttcagcacat    900
tatcatgtaa acttcaccga ggctataagt caggatgtga atcttcatga ggccatcttg    960
ctttgtccca acaatacatt tagaagagat ccaacagcaa ggacttcaca gtcacaagaa   1020
ccatttctgc agttaaattc tcataccacc aatcctgagc aaacccttcc tggaaactaat   1080
ttgacaggat ttctttcacc ggttgacaat catatgagga atctaacaag ccaagaccta   1140
ctgtatgacc ttgacataaa tatatttgat gagataaact taatgtcatt ggccacagaa   1200
gacaactttg atccaatcga tgtttctcag ctttttgatg aaccagattc tgattctggc   1260
ctttctttag attcaagtca caataatacc tctgtcatca agtctaattc ctctcactct   1320
gtgtgtgatg aaggtyctat aggttattgc actgaccatg aatctagttc ccatcatgac   1380
ttagaaggty ctgtaggtgg ctactaccga gaaccagta agctttgtca cttggatcaa   1440
agtgattctg atttccatgg agatcttaca tttcaacacg tatttcataa ccacacttac   1500
cacttacagc caactgcacc agaactctact tctgaacctt ttccgtggcc tgggaagtca   1560
cagaagataa ggagtagata ccttgaagac acagatagaa acttgagccg tgatgaacag   1620
cgtgctaaag ctttgcatat cccttttctt gtagatgaaa ttgtcggcat gectgttgat   1680
tctttcaata gcatgttaag tagatattat ctgacagacc tacaagtctc acttatccgt   1740
gacatcagac gaagagggaa aaataaagtt gctgctcaga actgtcgtaa acgcaaattg   1800
gacataatth tgaatttaga agatgatgta tgtaacttgc aagcaaagaa ggaactctt   1860
aagagagagc aagcacaatg taacaaagct attaacataa tgaacagaa actgcatgac   1920
ctttatcatg atatttttag tagattaaga gatgaccaag gtaggccagt caatcccaac   1980
cactatgctc tccagtgtac ccatgatgga agtatcttga tagtacccaa agaactggtg   2040
gcctcaggcc acaaaaagga aacccaaaag ggaaagagaa agtgagaaga aactgaagat   2100

```

-continued

```

ggactctatt atgtgaagta gtaatgttca gaaactgatt atttggatca gaaaccattg 2160
aaactgcttc aagaattgta tctttaagta ctgctacttg aataactcag ttaacgctgt 2220
tttgaagcctt acatggacaa atgttttagga cttcaagatc acacttgtgg gcaatctggg 2280
ggagccacaa cttttcatga agtgcattgt atacaaaatt catagtattg tccaaagaat 2340
aggttaacat gaaaaccagg taagactttc catcttgcca gccatccttt ttaagagtaa 2400
gttgggtact tcaaaaagag caaacactgg ggatcaaatt attttaagag gtatttcagt 2460
tttaaagtca aaatagcctt attttcattt agtttgtag cactatagtg agcttttcaa 2520
acactatctt aatctttata tttaacttat aaatcttctt ttctatggaa ataaatcttg 2580
tattgtatt aaaaaaaaaa aaa 2603

```

```

<210> SEQ ID NO 9
<211> LENGTH: 1209
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: 1161
<223> OTHER INFORMATION: n = A,T,C or G

```

```

<400> SEQUENCE: 9

```

```

atgaagtgg aggtgttcgt ccctcgcgcg gccacggggg acaagcaggg cagtgcactg 60
gagggcgcgg gggcagcaga cgcgcctgcc ccgctgtcgg cggcgggaga cgactccctg 120
ggctcagatg gggactgcgc ggccaagccg tccgcggggc gggcggccag agatacgcag 180
ggcgacggcg aacagagtgc gggaggcggg ccgggcgcgg aggaggcgat cccggcagca 240
gctgctgcag cgtgtgtggc ggagggcgcg gaggcggggg cggcggggcc aggcgcgggc 300
ggcgcgggga gcggcagagg tgacgcagc aagccatata cgcggcggcc caagccccc 360
tactcgtaca tcgcgctcat cgccatggcc atccgcgact cggcggggcg gcgcttgacg 420
ctggcggaga tcaacgagta cctcatgggc aagttcccct ttttcgcggc cagctacacg 480
ggctggcgcg actcctgtcg ccacaacctt tcgctcaacg actgcttcgt caaggtgctg 540
cgcgaccctc cgcggccctg gggcaaggac aactactgga tgctcaacc caacagcgag 600
tacaccttcg ccgacggggg ttccgcgcgc cgcgcgaagc gcctcagcca ccgcgcgcgg 660
gtccccgcgc ccggggtcgg gcccgaggag gccccgggccc tccccgcgc cccgcgcgcc 720
gcgcccgcgc ccccgccctc gccccgcgat cgcctgcccc cccgccagga ggagcgcgcc 780
agccccgcgg gcaagtcttc cagctccttc gccatcgaca gcctcctgcg caagcccttc 840
cgcagccgct gcctcagggg cacggccccc gggacgacgc ttcagtgggg cgcgcgcgcc 900
tgccccgcgc tgccccgctt ccccgcgctc ctccccgcgg cgccctgcag ggccctgctg 960
ccgctctgcg cgtacggcgc gggcgagccg gcgcgctgg gcgcgcgcga ggccgaggtg 1020
ccaccgaccg cgcgcgccct cctgcttgca cctctcccgg cggcggcccc cgccaagcca 1080
ctccgaggcc cggcggcccg cggcgcgcac ctgtactgcc cctcgggct gcccgagcc 1140
ctgcaggcgg ccttagtccg ncgtcctggc ccgcacctgt cgtaccgggt ggagcgcctc 1200
ctagcttga 1209

```

```

<210> SEQ ID NO 10
<211> LENGTH: 1474
<212> TYPE: DNA

```

-continued

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 10

```

ggcagatgaa atataagatt catcaaccac atttgacagc ccatggcagg tttcctgttt      60
tccatcgacc ctctgcaggt cacagacaca cagagcccag ccgtggcagg ctcagccggg    120
gtccggggct gctaacaacg gctacattcc tccccaggg ccaagggaaa tcctgagcgc     180
aggccagggt tgtttgttt tgaggtgtgc tgggatgaaa ggcaccctgg aagtggaagg     240
ttcggtcatt cattaattaa ttacatctat aattgagggt ttgttcttaa gagcgagtcc    300
tttgaaagta ctttccctca aacagtgact gccacaaagg catcagatat tcaccacctt    360
ctcggctgcc tcagcacagc aagctttatt ctgggacctg agatcctgtt ctgagctggc    420
tttcccttct ccaggctcgc tcaccctccc tttagagata gtggatgta agatgaccaa    480
tgctcagatt attcttctca ttgacaatgc caggatggca gtggatgact tcaacctcaa    540
gaaatggaga agcatcatgt gccaaagtgc ttcaatgtca atgtgaagggt ggatacaggt    600
cccaggaag atctgattaa ggtcctggag gatatgagac aagaatatga gcttataata    660
aagaagaagc atcgagactt ggacacttgg tataaagaac agtctgcagc catgtcccag    720
gaggcagcca gtccagccc tgctgcagagc agacaagggt acatccacga actgaaagcgc    780
acattccagg ccttgagatg tgacctgcag gcacagtaca gcacgaaatc tgctttggaa    840
aacatgttat ccgagaccca gtctcgttac tctgcaagc tccaggacat gcaagagatc    900
atctcccact atgaggagga actgacgcag ctacgccacg aactggagcg gcagaacaat    960
gaataccaag tgctgtctgg catcaaaacc cacctggaga aggaaatcac cacgtaccga   1020
cggctcctgg agggagagag tgaagggaca cgggaagaat caaagtcgag catgaaagtg   1080
tctgcaactc caaagatcaa ggccataacc caggagacca tcaacggaag attagttctt   1140
tgtcaagtga atgaaatcca aaagcacgca tgagaccaat gaaagtttcc gctgtttgta   1200
aaatctatct tcccccaagg aaagtcttg cacagacacc agtgagttag ttctaaaaga   1260
tacccttggg attatcagac tcagaaaactt ttattttttt tttctgtaac agtctacca   1320
gacttctcat aatgctctta atatattgca cttttctaat caaagtgcga gtttatgagg   1380
gtaaagctct actttctact tgcagccttc agattctcat ctttttgcac ctattttgta   1440
gccaataaaa ctccgcaact gcaaaaaaaaa aaaa                                     1474

```

<210> SEQ ID NO 11

<211> LENGTH: 411

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 11

```

ttttttttt aaaaaagag gcttggtgtaag tttttgatgc ttagttgact tttagcatta      60
tccagcattt gtattatgaa ccagtgagta ctgtaatttt tctttccctt tcagaaagac    120
tcaaagggaa catataaatg tttcctatct ttaatgtggc aatagtgtag ctaaacctgg    180
tacagacgga ataaacacac ctctaattatt ctctgaaga tttggtgatc cagtttcaaa    240
taaggatgag gaaaaacaga tgttttcatt atcgccactt aatccttact tccgattata    300
attatacatg tttggctgta ataactatac taaagcatgc ttgtgaaagt agacttctac    360
aaggacagaa aaccacacaac aacaagatgc gatcacgaaa gacaaggcat a              411

```

-continued

```

<210> SEQ ID NO 12
<211> LENGTH: 2336
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 12

cttttcttcc gcacggttgg aggaggtcgg ctggttatcg ggagttggag ggctgaggtc   60
gggaggggtg tgtgtacaga gctctaggac tcacgcacca ggccagtcgc ggattttggg   120
ccgaggcctg ggttacaagc agcaagtgcg cggttggggc cactgcgagg ccgttttaga   180
aaactgttta aaacaaagag caattgatgg ataaatcagg aatagattct cttgacctat   240
tgacatctga tctgtggaa cttgcaaatc gaagtataaa ctcttctgat agcagcttat   300
ttaaaactca gtgtatccct tactcaccta aaggggagaa aagaaacccc attcgaaaat   360
ttgttcgtac acctgaaagt gttcacgcaa gtgattcatc aagtgactca tcttttgaac   420
caataccatt gactataaaa gctatttttg aaagattcaa gaacaggaaa aagagatata   480
aaaaaaaaaa aaagaggaggg taccagccaa caggaagacc acggggaaga ccagaaggaa   540
ggagaaatcc tatatactca ctaatagata agaagaaaca atttagaagc agaggatctg   600
gcttcccatt tttagaatca gagaatgaaa aaaacgcacc ttggagaaaa attttaacgt   660
ttgagcaagc tgttgcaaga ggatttttta actatattga aaagctgaag tatgaacacc   720
acctgaaaga atcattgaag caaatgaatg ttgggtgaaga tttagaaaat gaagattttg   780
acagtcgtag atacaaatth ttggatgatg atggatccat ttctcctatt gaggagtcaa   840
cagcagagga tgaggatgca acacatcttg aagataacga atgtgatatc aaattggcag   900
gggatagttt catagtaagt tctgaattcc ctgtaagact gagtgatata ttagaagaag   960
aggatattac tgaagaagct gctttgtcta aaaagagagc tacaaaagcc aaaaatactg  1020
gacagagagg cctgaaaatg tgacaggatc atgaatgtca aaggctttta tcttgagaac  1080
atgggtgtctg gagttaaagg tattggcata ctccacacat ctgtaccatt cttgagtgat  1140
cgcttaggaa tgaatgtgat ttgaactcat tcatgttgag aggggtgcaa attgagaacc  1200
aggtagatcc ccaccaccta cagtaaaaag gaccctaaag taaattgggt gaagaaatta  1260
gatcccaaag attcctgggtg aattttgaag tcttcatcag tatatccata ttaaacgag  1320
atgacagaag ccaaagtaat tatggcaagt aatggttttt atcttaacta taagttatth  1380
gctcaagggt gtaatggtea ttaccaaggc ttttagaatg cagtttctca tttgctgtgg  1440
acatgaccaa aaaaaaaaaa ttcccagtag gttttctatc tgctacgttg ctagcaatca  1500
gottattggg aacagttgat taactgtaat agaaatgcaa tacaataaaa atgtgaacca  1560
catgtgattt ttctttaaaa tcagtgagat ttgaaaatc tcttagatct cttgaatcat  1620
gcaaatthgc tttgccttta tattgtaacc cttgtgggtt gctaataacc aagcagtttg  1680
tagtagagtt aactcagggt cgttctaggg actcattcat gttcactcac tgtacactca  1740
tctctggaaa tgtaaaatth acttttatac tattgttatg tagggctgac aggacaactg  1800
gatcagtttc attaaaaagg tatgtatgca ttgaaaaga catttgatg ggtcatttca  1860
aagagggcct atgaggctgt gaaaccaga gctcttaacg ctgtgaccaa agatggaagt  1920
tctctatagg aagccatagc actcctaagc tttgggtgcta tgttttctcg aggagatata  1980
aaacgtaata atccatgatt gttgccatgt gagagtttta aaggttaac aaaaatttctc  2040

```

-continued

```

ttcttcaggg caaacttgaa gataaatcctt ttgactccag ctcttttagag gatctaaagt 2100
gaccttgatg gacagtggaa gaaatcacaa catggaattc ctcgaataac aatttattga 2160
ctttaaataa tttgtctaa tgctacatat acacaattaa aaaaccttta cactatttct 2220
agaaagtcag catgtatttt tggctcgaag tttctctagt gttttctgtg gaaggaataa 2280
aaatttgagt ttcaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaa 2336

```

```

<210> SEQ ID NO 13
<211> LENGTH: 1361
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 13

```

```

Met Gly Ala Ala Gly Arg Gln Asp Phe Leu Phe Lys Ala Met Leu Thr
 1          5          10          15
Ile Ser Trp Leu Thr Leu Thr Cys Phe Pro Gly Ala Thr Ser Thr Val
 20          25          30
Ala Ala Gly Cys Pro Asp Gln Ser Pro Glu Leu Gln Pro Trp Asn Pro
 35          40          45
Gly His Asp Gln Asp His His Val His Ile Gly Gln Gly Lys Thr Leu
 50          55          60
Leu Leu Thr Ser Ser Ala Thr Val Tyr Ser Ile His Ile Ser Glu Gly
 65          70          75          80
Gly Lys Leu Val Ile Lys Asp His Asp Glu Pro Ile Val Leu Arg Thr
 85          90          95
Arg His Ile Leu Ile Asp Asn Gly Gly Glu Leu His Ala Gly Ser Ala
100          105          110
Leu Cys Pro Phe Gln Gly Asn Phe Thr Ile Ile Leu Tyr Gly Arg Ala
115          120          125
Asp Glu Gly Ile Gln Pro Asp Pro Tyr Tyr Gly Leu Lys Tyr Ile Gly
130          135          140
Val Gly Lys Gly Gly Ala Leu Glu Leu His Gly Gln Lys Lys Leu Ser
145          150          155          160
Trp Thr Phe Leu Asn Lys Thr Leu His Pro Gly Gly Met Ala Glu Gly
165          170          175
Gly Tyr Phe Phe Glu Arg Ser Trp Gly His Arg Gly Val Ile Val His
180          185          190
Val Ile Asp Pro Lys Ser Gly Thr Val Ile His Ser Asp Arg Phe Asp
195          200          205
Thr Tyr Arg Ser Lys Lys Glu Ser Glu Arg Leu Val Gln Tyr Leu Asn
210          215          220
Ala Val Pro Asp Gly Arg Ile Leu Ser Val Ala Val Asn Asp Glu Gly
225          230          235          240
Ser Arg Asn Leu Asp Asp Met Ala Arg Lys Ala Met Thr Lys Leu Gly
245          250          255
Ser Lys His Phe Leu His Leu Gly Phe Arg His Pro Trp Ser Phe Leu
260          265          270
Thr Val Lys Gly Asn Pro Ser Ser Ser Val Glu Asp His Ile Glu Tyr
275          280          285
His Gly His Arg Gly Ser Ala Ala Ala Arg Val Phe Lys Leu Phe Gln
290          295          300
Thr Glu His Gly Glu Tyr Phe Asn Val Ser Leu Ser Ser Glu Trp Val

```

-continued

305		310				315				320					
Gln	Asp	Val	Glu	Trp	Thr	Glu	Trp	Phe	Asp	His	Asp	Lys	Val	Ser	Gln
				325					330					335	
Thr	Lys	Gly	Gly	Glu	Lys	Ile	Ser	Asp	Leu	Trp	Lys	Ala	His	Pro	Gly
			340					345						350	
Lys	Ile	Cys	Asn	Arg	Pro	Ile	Asp	Ile	Gln	Ala	Thr	Thr	Met	Asp	Gly
		355					360						365		
Val	Asn	Leu	Ser	Thr	Glu	Val	Val	Tyr	Lys	Lys	Gly	Gln	Asp	Tyr	Arg
	370					375						380			
Phe	Ala	Cys	Tyr	Asp	Arg	Gly	Arg	Ala	Cys	Arg	Ser	Tyr	Arg	Val	Arg
385					390					395					400
Phe	Leu	Cys	Gly	Lys	Pro	Val	Arg	Pro	Lys	Leu	Thr	Val	Thr	Ile	Asp
				405					410						415
Thr	Asn	Val	Asn	Ser	Thr	Ile	Leu	Asn	Leu	Glu	Asp	Asn	Val	Gln	Ser
			420					425						430	
Trp	Lys	Pro	Gly	Asp	Thr	Leu	Val	Ile	Ala	Ser	Thr	Asp	Tyr	Ser	Met
		435					440						445		
Tyr	Gln	Ala	Glu	Glu	Phe	Gln	Val	Leu	Pro	Cys	Arg	Ser	Cys	Ala	Pro
	450					455						460			
Asn	Gln	Val	Lys	Val	Ala	Gly	Lys	Pro	Met	Tyr	Leu	His	Ile	Gly	Glu
465					470					475					480
Glu	Ile	Asp	Gly	Val	Asp	Met	Arg	Ala	Glu	Val	Gly	Leu	Leu	Ser	Arg
				485					490						495
Asn	Ile	Ile	Val	Met	Gly	Glu	Met	Glu	Asp	Lys	Cys	Tyr	Pro	Tyr	Arg
			500					505						510	
Asn	His	Ile	Cys	Asn	Phe	Phe	Asp	Phe	Asp	Thr	Phe	Gly	Gly	His	Ile
		515					520					525			
Lys	Phe	Ala	Leu	Gly	Phe	Lys	Ala	Ala	His	Leu	Glu	Gly	Thr	Glu	Leu
	530					535						540			
Lys	His	Met	Gly	Gln	Gln	Leu	Val	Gly	Gln	Tyr	Pro	Ile	His	Phe	His
545					550					555					560
Leu	Ala	Gly	Asp	Val	Asp	Glu	Arg	Gly	Gly	Tyr	Asp	Pro	Pro	Thr	Tyr
				565					570						575
Ile	Arg	Asp	Leu	Ser	Ile	His	His	Thr	Phe	Ser	Arg	Cys	Val	Thr	Val
			580					585						590	
His	Gly	Ser	Asn	Gly	Leu	Leu	Ile	Lys	Asp	Val	Val	Gly	Tyr	Asn	Ser
		595					600					605			
Leu	Gly	His	Cys	Phe	Phe	Thr	Glu	Asp	Gly	Pro	Glu	Glu	Arg	Asn	Thr
	610					615						620			
Phe	Asp	His	Cys	Leu	Gly	Leu	Leu	Val	Lys	Ser	Gly	Thr	Leu	Leu	Pro
625					630						635				640
Ser	Asp	Arg	Asp	Ser	Lys	Met	Cys	Lys	Met	Ile	Thr	Glu	Asp	Ser	Tyr
				645					650						655
Pro	Gly	Tyr	Ile	Pro	Lys	Pro	Arg	Gln	Asp	Cys	Asn	Ala	Val	Ser	Thr
			660					665						670	
Phe	Trp	Met	Ala	Asn	Pro	Asn	Asn	Asn	Leu	Ile	Asn	Cys	Ala	Ala	Ala
		675					680						685		
Gly	Ser	Glu	Glu	Thr	Gly	Phe	Trp	Phe	Ile	Phe	His	His	Val	Pro	Thr
	690					695					700				
Gly	Pro	Ser	Val	Gly	Met	Tyr	Ser	Pro	Gly	Tyr	Ser	Glu	His	Ile	Pro
705					710					715					720

-continued

Leu Gly Lys Phe Tyr Asn Asn Arg Ala His Ser Asn Tyr Arg Ala Gly
 725 730 735
 Met Ile Ile Asp Asn Gly Val Lys Thr Thr Glu Ala Ser Ala Lys Asp
 740 745 750
 Lys Arg Pro Phe Leu Ser Ile Ile Ser Ala Arg Tyr Ser Pro His Gln
 755 760 765
 Asp Ala Asp Pro Leu Lys Pro Arg Glu Pro Ala Ile Ile Arg His Phe
 770 775 780
 Ile Ala Tyr Lys Asn Gln Asp His Gly Ala Trp Leu Arg Gly Gly Asp
 785 790 795 800
 Val Trp Leu Asp Ser Cys Arg Phe Ala Asp Asn Gly Ile Gly Leu Thr
 805 810 815
 Leu Ala Ser Gly Gly Thr Phe Pro Tyr Asp Asp Gly Ser Lys Gln Glu
 820 825 830
 Ile Lys Asn Ser Leu Phe Val Gly Glu Ser Gly Asn Val Gly Thr Glu
 835 840 845
 Met Met Asp Asn Arg Ile Trp Gly Pro Gly Gly Leu Asp His Ser Gly
 850 855 860
 Arg Thr Leu Pro Ile Gly Gln Asn Phe Pro Ile Arg Gly Ile Gln Leu
 865 870 875 880
 Tyr Asp Gly Pro Ile Asn Ile Gln Asn Cys Thr Phe Arg Lys Phe Val
 885 890 895
 Ala Leu Glu Gly Arg His Thr Ser Ala Leu Ala Phe Arg Leu Asn Asn
 900 905 910
 Ala Trp Gln Ser Cys Pro His Asn Asn Val Thr Gly Ile Ala Phe Glu
 915 920 925
 Asp Val Pro Ile Thr Ser Arg Val Phe Phe Gly Glu Pro Gly Pro Trp
 930 935 940
 Phe Asn Gln Leu Asp Met Asp Gly Asp Lys Thr Ser Val Phe His Asp
 945 950 955 960
 Val Asp Gly Ser Val Ser Glu Tyr Pro Gly Ser Tyr Leu Thr Lys Asn
 965 970 975
 Asp Asn Trp Leu Val Arg His Pro Asp Cys Ile Asn Val Pro Asp Trp
 980 985 990
 Arg Gly Ala Ile Cys Ser Gly Cys Tyr Ala Gln Met Tyr Ile Gln Ala
 995 1000 1005
 Tyr Lys Thr Ser Asn Leu Arg Met Lys Ile Ile Lys Asn Asp Phe Pro
 1010 1015 1020
 Ser His Pro Leu Tyr Leu Glu Gly Ala Leu Thr Arg Ser Thr His Tyr
 1025 1030 1035 1040
 Gln Gln Tyr Gln Pro Val Val Thr Leu Gln Lys Gly Tyr Thr Ile His
 1045 1050 1055
 Trp Asp Gln Thr Ala Pro Ala Glu Leu Ala Ile Trp Leu Ile Asn Phe
 1060 1065 1070
 Asn Lys Gly Asp Trp Ile Arg Val Gly Leu Cys Tyr Pro Arg Gly Thr
 1075 1080 1085
 Thr Phe Ser Ile Leu Ser Asp Val His Asn Arg Leu Leu Lys Gln Thr
 1090 1095 1100
 Ser Lys Thr Gly Val Phe Val Arg Thr Leu Gln Met Asp Lys Val Glu
 1105 1110 1115 1120

-continued

Gln Ser Tyr Pro Gly Arg Ser His Tyr Tyr Trp Asp Glu Asp Ser Gly
 1125 1130 1135
 Leu Leu Phe Leu Lys Leu Lys Ala Gln Asn Glu Arg Glu Lys Phe Ala
 1140 1145 1150
 Phe Cys Ser Met Lys Gly Cys Glu Arg Ile Lys Ile Lys Ala Leu Ile
 1155 1160 1165
 Pro Lys Asn Ala Gly Val Ser Asp Cys Thr Ala Thr Ala Tyr Pro Lys
 1170 1175 1180
 Phe Thr Glu Arg Ala Val Val Asp Val Pro Met Pro Lys Lys Leu Phe
 1185 1190 1195 1200
 Gly Ser Gln Leu Lys Thr Lys Asp His Phe Leu Glu Val Lys Met Glu
 1205 1210 1215
 Ser Ser Lys Gln His Phe Phe His Leu Trp Asn Asp Phe Ala Tyr Ile
 1220 1225 1230
 Glu Val Asp Gly Lys Lys Tyr Pro Ser Ser Glu Asp Gly Ile Gln Val
 1235 1240 1245
 Val Val Ile Asp Gly Asn Gln Gly Arg Val Val Ser His Thr Ser Phe
 1250 1255 1260
 Arg Asn Ser Ile Leu Gln Gly Ile Pro Trp Gln Leu Phe Asn Tyr Val
 1265 1270 1275 1280
 Ala Thr Ile Pro Asp Asn Ser Ile Val Leu Met Ala Ser Lys Gly Arg
 1285 1290 1295
 Tyr Val Ser Arg Gly Pro Trp Thr Arg Val Leu Glu Lys Leu Gly Ala
 1300 1305 1310
 Asp Arg Gly Leu Lys Leu Lys Glu Gln Met Ala Phe Val Gly Phe Lys
 1315 1320 1325
 Gly Ser Phe Arg Pro Ile Trp Val Thr Leu Asp Thr Glu Asp His Lys
 1330 1335 1340
 Ala Lys Ile Phe Gln Val Val Pro Ile Pro Val Val Lys Lys Lys Lys
 1345 1350 1355 1360
 Leu

<210> SEQ ID NO 14

<211> LENGTH: 755

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 14

Met Pro Pro Phe Leu Leu Leu Glu Ala Val Cys Val Phe Leu Phe Ser
 1 5 10 15
 Arg Val Pro Pro Ser Leu Pro Leu Gln Glu Val His Val Ser Lys Glu
 20 25 30
 Thr Ile Gly Lys Ile Ser Ala Ala Ser Lys Met Met Trp Cys Ser Ala
 35 40 45
 Ala Val Asp Ile Met Phe Leu Leu Asp Gly Ser Asn Ser Val Gly Lys
 50 55 60
 Gly Ser Phe Glu Arg Ser Lys His Phe Ala Ile Thr Val Cys Asp Gly
 65 70 75 80
 Leu Asp Ile Ser Pro Glu Arg Val Arg Val Gly Ala Phe Gln Phe Ser
 85 90 95
 Ser Thr Pro His Leu Glu Phe Pro Leu Asp Ser Phe Ser Thr Gln Gln
 100 105 110

-continued

Glu Val Lys Ala Arg Ile Lys Arg Met Val Phe Lys Gly Gly Arg Thr
 115 120 125
 Glu Thr Glu Leu Ala Leu Lys Tyr Leu Leu His Arg Gly Leu Pro Gly
 130 135 140
 Gly Arg Asn Ala Ser Val Pro Gln Ile Leu Ile Ile Val Thr Asp Gly
 145 150 155 160
 Lys Ser Gln Gly Asp Val Ala Leu Pro Ser Lys Gln Leu Lys Glu Arg
 165 170 175
 Gly Val Thr Val Phe Ala Val Gly Val Arg Phe Pro Arg Trp Glu Glu
 180 185 190
 Leu His Ala Leu Ala Ser Glu Pro Arg Gly Gln His Val Leu Leu Ala
 195 200 205
 Glu Gln Val Glu Asp Ala Thr Asn Gly Leu Phe Ser Thr Leu Ser Ser
 210 215 220
 Ser Ala Ile Cys Ser Ser Ala Thr Pro Asp Cys Arg Val Glu Ala His
 225 230 235 240
 Pro Cys Glu His Arg Thr Leu Glu Met Val Arg Glu Phe Ala Gly Asn
 245 250 255
 Ala Pro Cys Trp Arg Gly Ser Arg Arg Thr Leu Ala Val Leu Ala Ala
 260 265 270
 His Cys Pro Phe Tyr Ser Trp Lys Arg Val Phe Leu Thr His Pro Ala
 275 280 285
 Thr Cys Tyr Arg Thr Thr Cys Pro Gly Pro Cys Asp Ser Gln Pro Cys
 290 295 300
 Gln Asn Gly Gly Thr Cys Val Pro Glu Gly Leu Asp Gly Tyr Gln Cys
 305 310 315 320
 Leu Cys Pro Leu Ala Phe Gly Gly Glu Ala Asn Cys Ala Leu Lys Leu
 325 330 335
 Ser Leu Glu Cys Arg Val Asp Leu Leu Phe Leu Leu Asp Ser Ser Ala
 340 345 350
 Gly Thr Thr Leu Asp Gly Phe Leu Arg Ala Lys Val Phe Val Lys Arg
 355 360 365
 Phe Val Arg Ala Val Leu Ser Glu Asp Ser Arg Ala Arg Val Gly Val
 370 375 380
 Ala Thr Tyr Ser Arg Glu Leu Leu Val Ala Val Pro Val Gly Glu Tyr
 385 390 395 400
 Gln Asp Val Pro Asp Leu Val Trp Ser Leu Asp Gly Ile Pro Phe Arg
 405 410 415
 Gly Gly Pro Thr Leu Thr Gly Ser Ala Leu Arg Gln Ala Ala Glu Arg
 420 425 430
 Gly Phe Gly Ser Ala Thr Arg Thr Gly Gln Asp Arg Pro Arg Arg Val
 435 440 445
 Val Val Leu Leu Thr Glu Ser His Ser Glu Asp Glu Val Ala Gly Pro
 450 455 460
 Ala Arg His Ala Arg Ala Arg Glu Leu Leu Leu Leu Gly Val Gly Ser
 465 470 475 480
 Glu Ala Val Arg Ala Glu Leu Glu Glu Ile Thr Gly Ser Pro Lys His
 485 490 495
 Val Met Val Tyr Ser Asp Pro Gln Asp Leu Phe Asn Gln Ile Pro Glu
 500 505 510
 Leu Gln Gly Lys Leu Cys Ser Arg Gln Arg Pro Gly Cys Arg Thr Gln

-continued

```

Asp Glu Ser Asp Glu Leu Val Thr Asp Phe Pro Thr Asp Leu Pro Ala
   115                               120                               125

Thr Glu Val Phe Thr Pro Val Val Pro Thr Val Asp Thr Tyr Asp Gly
   130                               135                               140

Arg Gly Asp Ser Val Val Tyr Gly Leu Arg Ser Lys Ser Lys Lys Phe
  145                               150                               155                               160

Arg Arg Pro Asp Ile Gln Tyr Pro Asp Ala Thr Asp Glu Asp Ile Thr
                               165                               170                               175

Ser His Met Glu Ser Glu Glu Leu Asn Gly Ala Tyr Lys Ala Ile Pro
   180                               185                               190

Val Ala Gln Asp Leu Asn Ala Pro Ser Asp Trp Asp Ser Arg Gly Lys
   195                               200                               205

Asp Ser Tyr Glu Thr Ser Gln Leu Asp Asp Gln Ser Ala Glu Thr His
  210                               215                               220

Ser His Lys Gln Ser Arg Leu Tyr Lys Arg Lys Ala Asn Asp Glu Ser
  225                               230                               235                               240

Asn Glu His Ser Asp Val Ile Asp Ser Gln Glu Leu Ser Lys Val Ser
                               245                               250                               255

Arg Glu Phe His Ser His Glu Phe His Ser His Glu Asp Met Leu Val
                               260                               265                               270

Val Asp Pro Lys Ser Lys Glu Glu Asp Lys His Leu Lys Phe Arg Ile
   275                               280                               285

Ser His Glu Leu Asp Ser Ala Ser Ser Glu Val Asn
   290                               295                               300
    
```

```

<210> SEQ ID NO 16
<211> LENGTH: 829
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
    
```

<400> SEQUENCE: 16

```

Met Gly Leu Pro Arg Gly Pro Leu Ala Ser Leu Leu Leu Leu Gln Val
  1                               5                               10                               15

Cys Trp Leu Gln Cys Ala Ala Ser Glu Pro Cys Arg Ala Val Phe Arg
   20                               25                               30

Glu Ala Glu Val Thr Leu Glu Ala Gly Gly Ala Glu Gln Glu Pro Gly
   35                               40                               45

Gln Ala Leu Gly Lys Val Phe Met Gly Cys Pro Gly Gln Glu Pro Ala
   50                               55                               60

Leu Phe Ser Thr Asp Asn Asp Asp Phe Thr Val Arg Asn Gly Glu Thr
   65                               70                               75                               80

Val Gln Glu Arg Arg Ser Leu Lys Glu Arg Asn Pro Leu Lys Ile Phe
   85                               90                               95

Pro Ser Lys Arg Ile Leu Arg Arg His Lys Arg Asp Trp Val Val Ala
  100                               105                               110

Pro Ile Ser Val Pro Glu Asn Gly Lys Gly Pro Phe Pro Gln Arg Leu
  115                               120                               125

Asn Gln Leu Lys Ser Asn Lys Asp Arg Asp Thr Lys Ile Phe Tyr Ser
  130                               135                               140

Ile Thr Gly Pro Gly Ala Asp Ser Pro Pro Glu Gly Val Phe Ala Val
  145                               150                               155                               160

Glu Lys Glu Thr Gly Trp Leu Leu Leu Asn Lys Pro Leu Asp Arg Glu
  165                               170                               175
    
```

-continued

Glu Ile Ala Lys Tyr Glu Leu Phe Gly His Ala Val Ser Glu Asn Gly
 180 185 190
 Ala Ser Val Glu Asp Pro Met Asn Ile Ser Ile Ile Val Thr Asp Gln
 195 200 205
 Asn Asp His Lys Pro Lys Phe Thr Gln Asp Thr Phe Arg Gly Ser Val
 210 215 220
 Leu Glu Gly Val Leu Pro Gly Thr Ser Val Met Gln Val Thr Ala Thr
 225 230 235 240
 Asp Glu Asp Asp Ala Ile Tyr Thr Tyr Asn Gly Val Val Ala Tyr Ser
 245 250 255
 Ile His Ser Gln Glu Pro Lys Asp Pro His Asp Leu Met Phe Thr Ile
 260 265 270
 His Arg Ser Thr Gly Thr Ile Ser Val Ile Ser Ser Gly Leu Asp Arg
 275 280 285
 Glu Lys Val Pro Glu Tyr Thr Leu Thr Ile Gln Ala Thr Asp Met Asp
 290 295 300
 Gly Asp Gly Ser Thr Thr Thr Ala Val Ala Val Val Glu Ile Leu Asp
 305 310 315 320
 Ala Asn Asp Asn Ala Pro Met Phe Asp Pro Gln Lys Tyr Glu Ala His
 325 330 335
 Val Pro Glu Asn Ala Val Gly His Glu Val Gln Arg Leu Thr Val Thr
 340 345 350
 Asp Leu Asp Ala Pro Asn Ser Pro Ala Trp Arg Ala Thr Tyr Leu Ile
 355 360 365
 Met Gly Gly Asp Asp Gly Asp His Phe Thr Ile Thr Thr His Pro Glu
 370 375 380
 Ser Asn Gln Gly Ile Leu Thr Thr Arg Lys Gly Leu Asp Phe Glu Ala
 385 390 395 400
 Lys Asn Gln His Thr Leu Tyr Val Glu Val Thr Asn Glu Ala Pro Phe
 405 410 415
 Val Leu Lys Leu Pro Thr Ser Thr Ala Thr Ile Val Val His Val Glu
 420 425 430
 Asp Val Asn Glu Ala Pro Val Phe Val Pro Pro Ser Lys Val Val Glu
 435 440 445
 Val Gln Glu Gly Ile Pro Thr Gly Glu Pro Val Cys Val Tyr Thr Ala
 450 455 460
 Glu Asp Pro Asp Lys Glu Asn Gln Lys Ile Ser Tyr Arg Ile Leu Arg
 465 470 475 480
 Asp Pro Ala Gly Trp Leu Ala Met Asp Pro Asp Ser Gly Gln Val Thr
 485 490 495
 Ala Val Gly Thr Leu Asp Arg Glu Asp Glu Gln Phe Val Arg Asn Asn
 500 505 510
 Ile Tyr Glu Val Met Val Leu Ala Met Asp Asn Gly Ser Pro Pro Thr
 515 520 525
 Thr Gly Thr Gly Thr Leu Leu Leu Thr Leu Ile Asp Val Asn Asp His
 530 535 540
 Gly Pro Val Pro Glu Pro Arg Gln Ile Thr Ile Cys Asn Gln Ser Pro
 545 550 555 560
 Val Arg Gln Val Leu Asn Ile Thr Asp Lys Asp Leu Ser Pro His Thr
 565 570 575

-continued

Ser Pro Phe Gln Ala Gln Leu Thr Asp Asp Ser Asp Ile Tyr Trp Thr
 580 585 590
 Ala Glu Val Asn Glu Glu Gly Asp Thr Val Val Leu Ser Leu Lys Lys
 595 600 605
 Phe Leu Lys Gln Asp Thr Tyr Asp Val His Leu Ser Leu Ser Asp His
 610 615 620
 Gly Asn Lys Glu Gln Leu Thr Val Ile Arg Ala Thr Val Cys Asp Cys
 625 630 635 640
 His Gly His Val Glu Thr Cys Pro Gly Pro Trp Lys Gly Gly Phe Ile
 645 650 655
 Leu Pro Val Leu Gly Ala Val Leu Ala Leu Leu Phe Leu Leu Leu Val
 660 665 670
 Leu Leu Leu Leu Val Arg Lys Lys Arg Lys Ile Lys Glu Pro Leu Leu
 675 680 685
 Leu Pro Glu Asp Asp Thr Arg Asp Asn Val Phe Tyr Tyr Gly Glu Glu
 690 695 700
 Gly Gly Gly Glu Glu Asp Gln Asp Tyr Asp Ile Thr Gln Leu His Arg
 705 710 715 720
 Gly Leu Glu Ala Arg Pro Glu Val Val Leu Arg Asn Asp Val Ala Pro
 725 730 735
 Thr Ile Ile Pro Thr Pro Met Tyr Arg Pro Arg Pro Ala Asn Pro Asp
 740 745 750
 Glu Ile Gly Asn Phe Ile Ile Glu Asn Leu Lys Ala Ala Asn Thr Asp
 755 760 765
 Pro Thr Ala Pro Pro Tyr Asp Thr Leu Leu Val Phe Asp Tyr Glu Gly
 770 775 780
 Ser Gly Ser Asp Ala Ala Ser Leu Ser Ser Leu Thr Ser Ser Ala Ser
 785 790 795 800
 Asp Gln Asp Gln Asp Tyr Asp Tyr Leu Asn Glu Trp Gly Ser Arg Phe
 805 810 815
 Lys Lys Leu Ala Asp Met Tyr Gly Gly Gly Glu Asp Asp
 820 825

<210> SEQ ID NO 17

<211> LENGTH: 694

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 17

Met Lys His Leu Lys Arg Trp Trp Ser Ala Gly Gly Gly Leu Leu His
 1 5 10 15
 Leu Thr Leu Leu Leu Ser Leu Ala Gly Leu Arg Val Asp Leu Asp Leu
 20 25 30
 Tyr Leu Leu Leu Pro Pro Pro Thr Leu Leu Gln Asp Glu Leu Leu Phe
 35 40 45
 Leu Gly Gly Pro Ala Ser Ser Ala Tyr Ala Leu Ser Pro Phe Ser Ala
 50 55 60
 Ser Gly Gly Trp Gly Arg Ala Gly His Leu His Pro Lys Gly Arg Glu
 65 70 75 80
 Leu Asp Pro Ala Ala Pro Pro Glu Gly Gln Leu Leu Arg Glu Val Arg
 85 90 95
 Ala Leu Gly Val Pro Phe Val Pro Arg Thr Ser Val Asp Ala Trp Leu
 100 105 110

-continued

Pro Phe Pro Trp Pro Gly Lys Ser Gln Lys Ile Arg Ser Arg Tyr Leu
 515 520 525
 Glu Asp Thr Asp Arg Asn Leu Ser Arg Asp Glu Gln Arg Ala Lys Ala
 530 535 540
 Leu His Ile Pro Phe Ser Val Asp Glu Ile Val Gly Met Pro Val Asp
 545 550 555 560
 Ser Phe Asn Ser Met Leu Ser Arg Tyr Tyr Leu Thr Asp Leu Gln Val
 565 570 575
 Ser Leu Ile Arg Asp Ile Arg Arg Arg Gly Lys Asn Lys Val Ala Ala
 580 585 590
 Gln Asn Cys Arg Lys Arg Lys Leu Asp Ile Ile Leu Asn Leu Glu Asp
 595 600 605
 Asp Val Cys Asn Leu Gln Ala Lys Lys Glu Thr Leu Lys Arg Glu Gln
 610 615 620
 Ala Gln Cys Asn Lys Ala Ile Asn Ile Met Lys Gln Lys Leu His Asp
 625 630 635 640
 Leu Tyr His Asp Ile Phe Ser Arg Leu Arg Asp Asp Gln Gly Arg Pro
 645 650 655
 Val Asn Pro Asn His Tyr Ala Leu Gln Cys Thr His Asp Gly Ser Ile
 660 665 670
 Leu Ile Val Pro Lys Glu Leu Val Ala Ser Gly His Lys Lys Glu Thr
 675 680 685
 Gln Lys Gly Lys Arg Lys
 690

<210> SEQ ID NO 18

<211> LENGTH: 402

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 18

Met Lys Leu Glu Val Phe Val Pro Arg Ala Ala His Gly Asp Lys Gln
 1 5 10 15
 Gly Ser Asp Leu Glu Gly Ala Gly Gly Ser Asp Ala Pro Ser Pro Leu
 20 25 30
 Ser Ala Ala Gly Asp Asp Ser Leu Gly Ser Asp Gly Asp Cys Ala Ala
 35 40 45
 Lys Pro Ser Ala Gly Gly Gly Ala Arg Asp Thr Gln Gly Asp Gly Glu
 50 55 60
 Gln Ser Ala Gly Gly Gly Pro Gly Ala Glu Glu Ala Ile Pro Ala Ala
 65 70 75 80
 Ala Ala Ala Ala Val Val Ala Glu Gly Ala Glu Ala Gly Ala Ala Gly
 85 90 95
 Pro Gly Ala Gly Gly Ala Gly Ser Gly Glu Gly Ala Arg Ser Lys Pro
 100 105 110
 Tyr Thr Arg Arg Pro Lys Pro Pro Tyr Ser Tyr Ile Ala Leu Ile Ala
 115 120 125
 Met Ala Ile Arg Asp Ser Ala Gly Gly Arg Leu Thr Leu Ala Glu Ile
 130 135 140
 Asn Glu Tyr Leu Met Gly Lys Phe Pro Phe Phe Arg Gly Ser Tyr Thr
 145 150 155 160
 Gly Trp Arg Asn Ser Val Arg His Asn Leu Ser Leu Asn Asp Cys Phe
 165 170 175

-continued

Val Lys Val Leu Arg Asp Pro Ser Arg Pro Trp Gly Lys Asp Asn Tyr
 180 185 190

Trp Met Leu Asn Pro Asn Ser Glu Tyr Thr Phe Ala Asp Gly Val Phe
 195 200 205

Arg Arg Arg Arg Lys Arg Leu Ser His Arg Ala Pro Val Pro Ala Pro
 210 215 220

Gly Leu Arg Pro Glu Glu Ala Pro Gly Leu Pro Ala Ala Pro Pro Pro
 225 230 235 240

Ala Pro Ala Ala Pro Ala Ser Pro Arg Met Arg Ser Pro Ala Arg Gln
 245 250 255

Glu Glu Arg Ala Ser Pro Ala Gly Lys Phe Ser Ser Ser Phe Ala Ile
 260 265 270

Asp Ser Ile Leu Arg Lys Pro Phe Arg Ser Arg Arg Leu Arg Asp Thr
 275 280 285

Ala Pro Gly Thr Thr Leu Gln Trp Gly Ala Ala Pro Cys Pro Pro Leu
 290 295 300

Pro Ala Phe Pro Ala Leu Leu Pro Ala Ala Pro Cys Arg Ala Leu Leu
 305 310 315 320

Pro Leu Cys Ala Tyr Gly Ala Gly Glu Pro Ala Arg Leu Gly Ala Arg
 325 330 335

Glu Ala Glu Val Pro Pro Thr Ala Pro Pro Leu Leu Leu Ala Pro Leu
 340 345 350

Pro Ala Ala Ala Pro Ala Lys Pro Leu Arg Gly Pro Ala Ala Gly Gly
 355 360 365

Ala His Leu Tyr Cys Pro Leu Arg Leu Pro Ala Ala Leu Gln Ala Ala
 370 375 380

Leu Val Arg Arg Pro Gly Pro His Leu Ser Tyr Pro Val Glu Thr Leu
 385 390 395 400

Leu Ala

<210> SEQ ID NO 19
 <211> LENGTH: 209
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 19

Met Glu Lys His His Val Pro Ser Asp Phe Asn Val Asn Val Lys Val
 1 5 10 15

Asp Thr Gly Pro Arg Glu Asp Leu Ile Lys Val Leu Glu Asp Met Arg
 20 25 30

Gln Glu Tyr Glu Leu Ile Ile Lys Lys Lys His Arg Asp Leu Asp Thr
 35 40 45

Trp Tyr Lys Glu Gln Ser Ala Ala Met Ser Gln Glu Ala Ala Ser Pro
 50 55 60

Ala Thr Val Gln Ser Arg Gln Gly Asp Ile His Glu Leu Lys Arg Thr
 65 70 75 80

Phe Gln Ala Leu Glu Ile Asp Leu Gln Ala Gln Tyr Ser Thr Lys Ser
 85 90 95

Ala Leu Glu Asn Met Leu Ser Glu Thr Gln Ser Arg Tyr Ser Cys Lys
 100 105 110

Leu Gln Asp Met Gln Glu Ile Ile Ser His Tyr Glu Glu Glu Leu Thr
 115 120 125

-continued

Gln Leu Arg His Glu Leu Glu Arg Gln Asn Asn Glu Tyr Gln Val Leu
 130 135 140
 Leu Gly Ile Lys Thr His Leu Glu Lys Glu Ile Thr Thr Tyr Arg Arg
 145 150 155 160
 Leu Leu Glu Gly Glu Ser Glu Gly Thr Arg Glu Glu Ser Lys Ser Ser
 165 170 175
 Met Lys Val Ser Ala Thr Pro Lys Ile Lys Ala Ile Thr Gln Glu Thr
 180 185 190
 Ile Asn Gly Arg Leu Val Leu Cys Gln Val Asn Glu Ile Gln Lys His
 195 200 205

Ala

<210> SEQ ID NO 20
 <211> LENGTH: 278
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 20

Met Asp Lys Ser Gly Ile Asp Ser Leu Asp His Val Thr Ser Asp Ala
 1 5 10 15
 Val Glu Leu Ala Asn Arg Ser Asp Asn Ser Ser Asp Ser Ser Leu Phe
 20 25 30
 Lys Thr Gln Cys Ile Pro Tyr Ser Pro Lys Gly Glu Lys Arg Asn Pro
 35 40 45
 Ile Arg Lys Phe Val Arg Thr Pro Glu Ser Val His Ala Ser Asp Ser
 50 55 60
 Ser Ser Asp Ser Ser Phe Glu Pro Ile Pro Leu Thr Ile Lys Ala Ile
 65 70 75 80
 Phe Glu Arg Phe Lys Asn Arg Lys Lys Arg Tyr Lys Lys Lys Lys Lys
 85 90 95
 Arg Arg Tyr Gln Pro Thr Gly Arg Pro Arg Gly Arg Pro Glu Gly Arg
 100 105 110
 Arg Asn Pro Ile Tyr Ser Leu Ile Asp Lys Lys Lys Gln Phe Arg Ser
 115 120 125
 Arg Gly Ser Gly Phe Pro Phe Leu Glu Ser Glu Asn Glu Lys Asn Ala
 130 135 140
 Pro Trp Arg Lys Ile Leu Thr Phe Glu Gln Ala Val Ala Arg Gly Phe
 145 150 155 160
 Phe Asn Tyr Ile Glu Lys Leu Lys Tyr Glu His His Leu Lys Glu Ser
 165 170 175
 Leu Lys Gln Met Asn Val Gly Glu Asp Leu Glu Asn Glu Asp Phe Asp
 180 185 190
 Ser Arg Arg Tyr Lys Phe Leu Asp Asp Asp Gly Ser Ile Ser Pro Ile
 195 200 205
 Glu Glu Ser Thr Ala Glu Asp Glu Asp Ala Thr His Leu Glu Asp Asn
 210 215 220
 Glu Cys Asp Ile Lys Leu Ala Gly Asp Ser Phe Ile Val Ser Ser Glu
 225 230 235 240
 Phe Pro Val Arg Leu Ser Val Tyr Leu Glu Glu Glu Asp Ile Thr Glu
 245 250 255
 Glu Ala Ala Leu Ser Lys Lys Arg Ala Thr Lys Ala Lys Asn Thr Gly
 260 265 270

-continued

Gln Arg Gly Leu Lys Met
275

<210> SEQ ID NO 21

<211> LENGTH: 488

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 21

Ala Val Leu Ala Ala His Cys Pro Phe Tyr Ser Trp Lys Arg Val Phe
1 5 10 15

Leu Thr His Pro Ala Thr Cys Tyr Arg Thr Thr Cys Pro Gly Pro Cys
20 25 30

Asp Ser Gln Pro Cys Gln Asn Gly Gly Thr Cys Val Pro Glu Gly Leu
35 40 45

Asp Gly Tyr Gln Cys Leu Cys Pro Leu Ala Phe Gly Gly Glu Ala Asn
50 55 60

Cys Ala Leu Lys Leu Ser Leu Glu Cys Arg Val Asp Leu Leu Phe Leu
65 70 75 80

Leu Asp Ser Ser Ala Gly Thr Thr Leu Asp Gly Phe Leu Arg Ala Lys
85 90 95

Val Phe Val Lys Arg Phe Val Arg Ala Val Leu Ser Glu Asp Ser Arg
100 105 110

Ala Arg Val Gly Val Ala Thr Tyr Ser Arg Glu Leu Leu Val Ala Val
115 120 125

Pro Val Gly Glu Tyr Gln Asp Val Pro Asp Leu Val Trp Ser Leu Asp
130 135 140

Gly Ile Pro Phe Arg Gly Gly Pro Thr Leu Thr Gly Ser Ala Leu Arg
145 150 155 160

Gln Ala Ala Glu Arg Gly Phe Gly Ser Ala Thr Arg Thr Gly Gln Asp
165 170 175

Arg Pro Arg Arg Val Val Val Leu Leu Thr Glu Ser His Ser Glu Asp
180 185 190

Glu Val Ala Gly Pro Ala Arg His Ala Arg Ala Arg Glu Leu Leu Leu
195 200 205

Leu Gly Val Gly Ser Glu Ala Val Arg Ala Glu Leu Glu Glu Ile Thr
210 215 220

Gly Ser Pro Lys His Val Met Val Tyr Ser Asp Pro Gln Asp Leu Phe
225 230 235 240

Asn Gln Ile Pro Glu Leu Gln Gly Lys Leu Cys Ser Arg Gln Arg Pro
245 250 255

Gly Cys Arg Thr Gln Ala Leu Asp Leu Val Phe Met Leu Asp Thr Ser
260 265 270

Ala Ser Val Gly Pro Glu Asn Phe Ala Gln Met Gln Ser Phe Val Arg
275 280 285

Ser Cys Ala Leu Gln Phe Glu Val Asn Pro Asp Val Thr Gln Val Gly
290 295 300

Leu Val Val Tyr Gly Ser Gln Val Gln Thr Ala Phe Gly Leu Asp Thr
305 310 315 320

Lys Pro Thr Arg Ala Ala Met Leu Arg Ala Ile Ser Gln Ala Pro Tyr
325 330 335

Leu Gly Gly Val Gly Ser Ala Gly Thr Ala Leu Leu His Ile Tyr Asp

-continued

145		150		155		160									
Thr	Gly	Trp	Arg	Asn	Ser	Val	Arg	His	Asn	Leu	Ser	Leu	Asn	Asp	Cys
				165					170					175	
Phe	Val	Lys	Val	Leu	Arg	Asp	Pro	Ser	Arg	Pro	Trp	Gly	Lys	Asp	Asn
			180						185				190		
Tyr	Trp	Met	Leu	Asn	Pro	Asn	Ser	Glu	Tyr	Thr	Phe	Ala	Asp	Gly	Val
		195					200					205			
Phe	Arg	Arg	Arg	Arg	Lys	Arg	Leu	Ser	His	Arg	Ala	Pro	Val	Pro	Ala
	210				215						220				
Pro	Gly	Leu	Arg	Pro	Glu	Glu	Ala	Pro	Gly	Leu	Pro	Ala	Ala	Pro	Pro
	225				230					235					240
Pro	Ala	Pro	Ala	Ala	Pro	Ala	Ser	Pro	Arg	Met	Arg	Ser	Pro	Ala	Arg
			245						250					255	
Gln	Glu	Glu	Arg	Ala	Ser	Pro	Ala	Gly	Lys	Phe	Ser	Ser	Ser	Phe	Ala
			260					265					270		
Ile	Asp	Ser	Ile	Leu	Arg	Lys	Pro	Phe	Arg	Ser	Arg	Arg	Leu	Arg	Asp
		275					280						285		
Thr	Ala	Pro	Gly	Thr	Thr	Leu	Gln	Trp	Gly	Ala	Ala	Pro	Cys	Pro	Pro
	290					295						300			
Leu	Pro	Ala	Phe	Pro	Ala	Leu	Leu	Pro	Ala	Ala	Pro	Cys	Arg	Ala	Leu
	305				310					315					320
Leu	Pro	Leu	Cys	Ala	Tyr	Gly	Ala	Gly	Glu	Pro	Ala	Arg	Leu	Gly	Ala
			325						330					335	
Arg	Glu	Ala	Glu	Val	Pro	Pro	Thr	Ala	Pro	Pro	Leu	Leu	Leu	Ala	Pro
			340					345					350		
Leu	Pro	Ala	Ala	Ala	Pro	Ala	Lys	Pro	Leu	Arg	Gly	Pro	Ala	Ala	Gly
		355					360					365			
Gly	Ala	His	Leu	Tyr	Cys	Pro	Leu	Arg	Leu	Pro	Ala	Ala	Leu	Gln	Ala
	370					375					380				
Ala	Ser	Val	Arg	Arg	Pro	Gly	Pro	His	Leu	Pro	Tyr	Pro	Val	Glu	Thr
	385				390					395					400

Leu Leu Ala

<210> SEQ ID NO 24
 <211> LENGTH: 400
 <212> TYPE: PRT
 <213> ORGANISM: Mus musculus

<400> SEQUENCE: 24

Met	Lys	Leu	Glu	Val	Phe	Val	Pro	Arg	Ala	Ala	His	Gly	Asp	Lys	Met
1				5					10					15	
Gly	Ser	Asp	Leu	Glu	Gly	Ala	Gly	Ser	Ser	Asp	Val	Pro	Ser	Pro	Leu
			20					25					30		
Ser	Ala	Ala	Gly	Asp	Asp	Ser	Leu	Gly	Ser	Asp	Gly	Asp	Cys	Ala	Ala
		35					40					45			
Asn	Ser	Pro	Ala	Ala	Gly	Ser	Gly	Ala	Gly	Asp	Leu	Glu	Gly	Gly	Gly
		50				55					60				
Gly	Glu	Arg	Asn	Ser	Ser	Gly	Gly	Pro	Ser	Ala	Gln	Asp	Gly	Pro	Glu
	65				70					75					80
Ala	Thr	Asp	Asp	Ser	Arg	Thr	Gln	Ala	Ser	Ala	Ala	Gly	Pro	Cys	Ala
			85						90					95	
Gly	Gly	Val	Gly	Gly	Gly	Glu	Gly	Ala	Arg	Ser	Lys	Pro	Tyr	Thr	Arg

-continued

	100							105						110			
Arg	Pro	Lys	Pro	Pro	Tyr	Ser	Tyr	Ile	Ala	Leu	Ile	Ala	Met	Ala	Ile		
	115						120					125					
Arg	Asp	Ser	Ala	Gly	Gly	Arg	Leu	Thr	Leu	Ala	Glu	Ile	Asn	Glu	Tyr		
	130					135					140						
Leu	Met	Gly	Lys	Phe	Pro	Phe	Phe	Arg	Gly	Ser	Tyr	Thr	Gly	Trp	Arg		
145				150					155						160		
Asn	Ser	Val	Arg	His	Asn	Leu	Ser	Leu	Asn	Asp	Cys	Phe	Val	Lys	Val		
			165					170						175			
Leu	Arg	Asp	Pro	Ser	Arg	Pro	Trp	Gly	Lys	Asp	Asn	Tyr	Trp	Met	Leu		
		180						185					190				
Asn	Pro	Asn	Ser	Glu	Tyr	Thr	Phe	Ala	Asp	Gly	Val	Phe	Arg	Arg	Arg		
	195						200					205					
Arg	Lys	Arg	Leu	Ser	His	Arg	Thr	Thr	Val	Ser	Ala	Ser	Gly	Leu	Arg		
	210				215						220						
Pro	Glu	Glu	Ala	Pro	Pro	Gly	Pro	Ala	Gly	Thr	Pro	Gln	Pro	Ala	Pro		
225					230					235					240		
Ala	Ala	Arg	Ser	Ser	Pro	Ile	Ala	Arg	Ser	Pro	Ala	Arg	Gln	Glu	Glu		
				245					250					255			
Arg	Ser	Ser	Pro	Ala	Ser	Lys	Phe	Ser	Ser	Ser	Phe	Ala	Ile	Asp	Ser		
			260					265					270				
Ile	Leu	Ser	Lys	Pro	Phe	Arg	Ser	Arg	Arg	Asp	Gly	Asp	Ser	Ala	Leu		
	275					280						285					
Gly	Val	Gln	Leu	Pro	Trp	Gly	Ala	Ala	Pro	Cys	Pro	Pro	Leu	Arg	Ala		
	290					295					300						
Tyr	Pro	Ala	Leu	Leu	Pro	Ala	Ala	Pro	Gly	Gly	Ala	Leu	Leu	Pro	Leu		
305					310				315					320			
Cys	Ala	Tyr	Gly	Ala	Ser	Glu	Pro	Thr	Leu	Leu	Ala	Ser	Arg	Gly	Thr		
			325					330						335			
Glu	Val	Gln	Pro	Ala	Ala	Pro	Leu	Leu	Leu	Ala	Pro	Leu	Ser	Thr	Ala		
			340				345						350				
Ala	Pro	Ala	Lys	Pro	Phe	Arg	Gly	Pro	Glu	Thr	Ala	Gly	Ala	Ala	His		
	355						360					365					
Leu	Tyr	Cys	Pro	Leu	Arg	Leu	Pro	Thr	Ala	Leu	Gln	Ala	Ala	Ala	Ala		
	370					375					380						
Cys	Gly	Pro	Gly	Pro	His	Leu	Ser	Tyr	Pro	Val	Glu	Thr	Leu	Leu	Ala		
385					390					395					400		

<210> SEQ ID NO 25
 <211> LENGTH: 400
 <212> TYPE: PRT
 <213> ORGANISM: Rattus rattus

<400> SEQUENCE: 25

Met	Lys	Leu	Glu	Val	Phe	Ala	Pro	Arg	Ala	Ala	His	Gly	Asp	Lys	Met
1				5					10					15	
Gly	Ser	Asp	Leu	Glu	Gly	Ala	Gly	Ser	Ser	Asp	Val	Pro	Ser	Pro	Leu
			20					25					30		
Ser	Ala	Ala	Gly	Asp	Asp	Ser	Leu	Gly	Ser	Asp	Gly	Asp	Cys	Ala	Ala
	35					40					45				
Asn	Ser	Pro	Ala	Ala	Gly	Arg	Gly	Ala	Val	Asp	Leu	Glu	Gly	Gly	Gly
	50					55					60				

-continued

Gly Glu Arg Asn Ser Ser Gly Gly Ala Ser Thr Gln Asp Asp Pro Glu
65 70 75 80

Val Thr Asp Gly Ser Arg Thr Gln Ala Ser Pro Val Gly Pro Cys Ala
85 90 95

Gly Ser Val Gly Gly Gly Glu Gly Ala Arg Ser Lys Pro Tyr Thr Arg
100 105 110

Arg Pro Lys Pro Pro Tyr Ser Tyr Ile Ala Leu Ile Ala Met Ala Ile
115 120 125

Arg Asp Ser Ala Gly Gly Arg Leu Thr Leu Ala Glu Ile Asn Glu Tyr
130 135 140

Leu Met Gly Lys Phe Pro Phe Phe Arg Gly Ser Tyr Thr Gly Trp Arg
145 150 155 160

Asn Ser Val Arg His Asn Leu Ser Leu Asn Asp Cys Phe Val Lys Val
165 170 175

Leu Arg Asp Pro Ser Arg Pro Trp Gly Lys Asp Asn Tyr Trp Met Leu
180 185 190

Asn Pro Asn Ser Glu Tyr Thr Phe Ala Asp Gly Val Phe Arg Arg Arg
195 200 205

Arg Lys Arg Leu Ser His Arg Thr Thr Val Ser Ala Ser Gly Leu Arg
210 215 220

Pro Glu Glu Ala Pro Pro Gly Pro Ala Gly Thr Pro Gln Pro Ala Pro
225 230 235 240

Thr Ala Gly Ser Ser Pro Ile Ala Arg Ser Pro Ala Arg Gln Glu Glu
245 250 255

Gly Ser Ser Pro Ala Ser Lys Phe Ser Ser Ser Phe Ala Ile Asp Ser
260 265 270

Ile Leu Ser Lys Pro Phe Arg Ser Arg Arg Asp Gly Asp Pro Ala Leu
275 280 285

Gly Val Gln Leu Pro Trp Ser Ala Ala Pro Cys Pro Pro Leu Arg Ala
290 295 300

Tyr Pro Ala Leu Leu Pro Ala Ser Ser Gly Gly Ala Leu Leu Pro Leu
305 310 315 320

Cys Ala Tyr Gly Ala Gly Glu Pro Thr Leu Leu Ala Ser Arg Gly Ala
325 330 335

Glu Val Gln Pro Ala Ala Pro Leu Leu Leu Ala Pro Leu Ser Thr Ala
340 345 350

Ala Pro Ala Lys Pro Phe Arg Gly Pro Glu Thr Ala Gly Ala Ala His
355 360 365

Leu Tyr Cys Pro Leu Arg Leu Pro Thr Ala Leu Gln Ala Ala Ala Ala
370 375 380

Cys Gly Pro Gly Pro His Leu Ser Tyr Arg Val Glu Thr Leu Leu Ala
385 390 395 400

<210> SEQ ID NO 26
<211> LENGTH: 1212
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 26

atgaagttgg aggtgttcgt ccctcgcgcg gccacggggg acaagcaggg cagtgcacctg 60
gagggcgcgg gcggcagcga cgcgccgtcc ccgetgtcgg cggcgggaga cgactccctg 120
ggctcagatg gggactgcgc ggccaacagc ccggccgcgg gcggcggcgc cagagatacg 180

-continued

```

cagggcgacg gcgaacagag tgcgggaggc gggccgggcg cggaggaggc gatcccggca 240
gcagctgctg cagcgggtgtt ggcgggagggc gcggaggccg gggcggcggg gccaggcgcg 300
ggcggcgcgg ggagcggcga ggggtgcaagc agcaagccat ataccggcgg gcccaagccc 360
ccctactcgt acatcgcgct catcgccatg gccatccgcg actcggcggg cgggcgcttg 420
acgctggcgg agatcaacga gtacctcatg ggcaagtcc cctttttccg cggcagctac 480
acgggctggc gcaactccgt gcgccacaac ctttcgctca acgactgctt cgtcaagggtg 540
ctgcgcgacc cctcgcggcc ctggggcaag gacaactact ggatgctcaa ccccaacagc 600
gagtacacct tcgcccagcg ggtcttcgcg cgcgcgcgca agcgcctcag ccaccgcgcg 660
ccggctcccc cgccccgggt gcggcccag gaggcgccgg gcctccccgc cccccgcgcg 720
cccgcgcgcg ccgcccccgc ctcgcccgcg atgcgctcgc ccgcccgcca ggaggagcgc 780
gccagccccg cgggcaagtt ctccagctcc ttgcgcatcg acagcatcct gcgcaagccc 840
ttccgcagcc gccgcctcag ggacacggcc cccgggacga cgttcagtg gggcgcgcg 900
ccctgcccgc cgtgcgcgcg gttccccgcg ctctccccgc cggcgccttg cagggccctg 960
ctgcgcctct gcgcgtacgg cgcggggcag ccggcgcggc tgggcgcgcg caggccgag 1020
gtgccaccga ccgcgcgcgc cctcctgctt gcacctctcc cggcggcggc ccccgccaaag 1080
ccaactccgag gccccggcggc cggcggcgcg cacctgtaet gcccctcgcg gctgcccgcg 1140
gccctgcagg cggcctcagt ccgcgcgcct ggcccgcacc tgccgtaacc ggtggagacg 1200
ctgctagctt ga 1212

```

```

<210> SEQ ID NO 27
<211> LENGTH: 1203
<212> TYPE: DNA
<213> ORGANISM: Mus musculus

```

```

<400> SEQUENCE: 27

```

```

atgaaattgg aggtgttctt cccacgcgca gcccaacgggg acaaaatggg cagcgatctg 60
gagggggcgc gcagcagcga cgtgccatct ccaactgtccg cggctggtga cgactcctta 120
ggctcagacg gggactgtgc agccaacagc ccggcggcgg gcagcggcgc cgggatctg 180
gaaggtggcg gcggcgagag gaattcagat ggcgggcccga gcgcccaga cggctccggag 240
gcaactgatg acagcagaac gcaggcctcc gcggcagggc cgtgcgcggg cggcgtgggc 300
ggcggcgagg gcgcgcgag caagccgtac acgcggcggc ccaagcccc atactcctac 360
atcgctctca tcgcatggc catccgcgac tccgcgggcg gacgcctgac actggccgag 420
atcaacgagt acctcatggg caagttcccc tttttccggg gcagctacac gggctggcgc 480
aactccgtgc gccacaacct ctcgctcaac gactgtttcg tcaaggtgct gcgcgacccc 540
tcgcgccctt ggggcaagga caactactgg atgctcaacc ccaacagcga atacacctc 600
gccgacgggg tcttccgcgc ccgcgcgaag gcctcagcc accggaccac agtctccgcg 660
tccgggctgc ggcgggagga agcccccccc ggacctgccc ggacccccga gcccgcgccc 720
gccgcccctt cctccccgat cgcgcgctcg ccggctcgc aggaggagcg ctccagccct 780
gcgagcaagt tctccagctc cttcgcctac gacagcatc tcagcaagcc ttttcgcagc 840
cgcgcgcagc gcgactcggc tctgggggtg cagctaccct ggggcgcgcg tccctgcccg 900
ccgctgcgcg cctatecccc gctccttccc gcggcgcccc gtggcgtctt gctaccgctc 960

```

-continued

```

tgtgcttacg gcgcaagcga gcctacgctg ctggcgctcg gcgggaccga ggtgcagccc 1020
gccgcgcccc ttctgtggc gccctctcc accgeggctc cagccaagcc attccgaggt 1080
ccggagaccg ccggcgcggc gcacctgtac tgccccctac ggctgcccac ggccctgcag 1140
gcggcagcgg cctgcggtcc cggtcgcac ctgtcctacc cggtgagac tctgctagct 1200
tga 1203

```

```

<210> SEQ ID NO 28
<211> LENGTH: 1203
<212> TYPE: DNA
<213> ORGANISM: Rattus rattus

```

```

<400> SEQUENCE: 28
atgaaattgg aggtatttgc cccacgcgca gccacgggg acaagatggg cagtgcactg 60
gagggggcgc gcagcagcga cgtgccatct ccgctgtccg cggctggcga cgactcctta 120
ggctctgacg gggactgtgc agccaacagc ccggcgcgcg gcagagcgc cgtggatctg 180
gaaggcgcgc gcggcgagag gaattcgagt ggcgggcgca gcaccaaga cgatcccag 240
gtgaccgatg gcagcagaac gcaggcctcc ccggtggggc cgtgcgcggg cagcgtgggc 300
ggcggtgagg gcgcgcgcag caagccgtac acgcgcggc ccaagcccc ctactcctac 360
atcgactca tcgccatggc catccgcgac tccgcgggcg gacgctgac gctggccgag 420
atcaacgagt acctcatggg caagtcccc ttttccggg gcagctaac gggctggcgc 480
aactccgtgc gccacaaact ctcgctcaac gactgtttcg tcaaggtgct gcgcgacccc 540
tcgcgccct ggggcaagga caattactgg atgctcaacc ccaacagcga atacacctc 600
gccgacgggg tcttcgcgcg ccgcccgaag cgctcagcc accggaccac agtctccgca 660
tcggggctac ggcgggagga agcccccccc ggacctgcgg ggacccccga gcccgcgccc 720
accgcccggc cctccccaat gcgcgcctcg cccgctcgcc agggagaggg ctccagcccg 780
gcgagcaagt tctccagctc cttcgccatc gacagcatcc tcagcaagcc gtttcgcagc 840
cgccgcgacg gcgaccgcgc tctgggggtg cagctaccct ggagcgtgc tccctgcccc 900
ccgctgcgcg cctateccgc gctccttccc gcgtgctccg cgggtgcctc gctgcgctc 960
tgtgcttacg gcgcgggcga gccacgctg ctggcgctcg gcggggccga ggtgcagccc 1020
gccgcgcccc tgtgtgtggc gccctctcc accgeggccc cagccaagcc atttcgaggt 1080
ccggagaccg ccggcgcggc gcacctgtac tgccccctac ggctgcccac ggccctgcag 1140
gcggcgcgcg cctgcggtcc cggtcgcac ctgtcctacc cggtgagac gctgctagct 1200
tga 1203

```

1-122. (canceled)

123. A method of detecting whether a subject is likely to have colon neoplasia comprising:

- (a) obtaining a biological sample from said subject;
- (b) detecting the presence or absence of a ColoUp2 polypeptide, wherein said ColoUp2 polypeptide is encoded by a nucleic acid sequence that hybridizes under stringent conditions to a nucleic acid sequence of SEQ ID NO: 5,

wherein the presence of said polypeptide is indicative of colon neoplasia.

124. The method of claim **123**, wherein said biological sample is selected from the group consisting of whole blood or a fraction thereof.

125. The method of claim **123**, wherein said biological sample is selected from the group consisting of urine or stool samples.

126. The method of claim **123**, wherein said biological sample is a blood sample.

127. The method of claim **126**, wherein said blood sample is fractionated to obtain blood serum and/or blood plasma.

128. The method of claim **127**, wherein said biological sample is enriched for ColoUp2.

129. The method of claim **123**, wherein the polypeptide is detected by an assay.

130. The method of claim **129**, wherein said assay employs an antibody.

131. The method of claim **130**, where said assay is selected from the group consisting of an immunoprecipitation assay, a Western blot, a radioimmunoassay, and an enzyme-linked immunosorbent assay (ELISA).

132. The method of claim **130**, wherein said assay comprises contacting the biological sample with an antibody that interacts with the ColoUp2 polypeptide.

133. The method of claim **132**, wherein the antibody is detectably labeled.

134. The method of claim **133**, wherein the label is selected from the group consisting of an enzyme, a fluorescent substance, a chemiluminescent substance, a chromophore, a radioactive isotope and a complexing agent.

* * * * *

专利名称(译)	用于分类患者的方法和组合物		
公开(公告)号	US20090311726A1	公开(公告)日	2009-12-17
申请号	US12/386176	申请日	2009-04-13
[标]申请(专利权)人(译)	MARKOWITZ SANFORD D		
申请(专利权)人(译)	MARKOWITZ SANFORD D		
当前申请(专利权)人(译)	凯斯西储大学		
[标]发明人	MARKOWITZ SANFORD D		
发明人	MARKOWITZ, SANFORD D.		
IPC分类号	G01N33/53		
CPC分类号	C07K14/4748 G01N2500/00 G01N33/57419		
其他公开文献	US8268568		
外部链接	Espacenet USPTO		

摘要(译)

本公开尤其提供了用于对患者的肿瘤状态进行分类的分子标记，在诊断测试中使用分子标记的方法，与分子标记相关的核酸和氨基酸序列，用于检测分子标记的试剂，以及方法。用于鉴定高度平行的基因表达数据中的候选分子标记。

Figure 1A. Amino acid sequence of secreted ColoUpl protein (I) (SEQ ID NO: 1)

```
TVAAGCPDQSPQLQPNPGHDQDHHVHIGQSKTLLLTSSATVYSIHSSEGGKLVKIDHD
EPIVLRTRHILIDNGGELHAGSALCPFGQNFITILYGRADEGIQPPYYGLKYIGVKGK
GALELHGQKLSWTFLNKTLHPGGMAEGGYFFERSWGHGRVIVHVIDPKSGTVIHSDFR
DTYRSKKESERLVQYLNAVDPGRILSVAVNDEGSRLDDMARKAMTKLGSKHFHLGFR
HPWSFLTVMGNPSSSVEDHIEYHGRGSAARVFKLPQTEHGEYFVNLSSSEWVQDVEW
TEWFDHDKVSQTKGGEKISDLWKAHPGKICNRPIDIQATMDGVNLSDEVVYKGGQYR
FACYDRGRACRSYRVRFLCGKPVKPLTVIDTNVNSTILNLEDNVQSWKPGDTLVIAS
TDYSMYQAEFPQVLPFCRSCAPNQVKVAGKPMYLIHIGEEIDGVDMRAEVLGSLRNIIVMG
EMEDKCYPIRNHICNFFDFTFGGHIKFFALGPKAAHLEGTGLKHMGGQLVGGYPIHFHL
AGDVDERGGYDPPYIRDLSTHHTFSRCVTVHGSNGLLIKDVVGYNSLGHCFPTEDGPE
ERNTFDHLCLGLLVKSGTLLPSDRDSKMCKMITEDSYPGYIPKPRQDCNAVSTFWMANPN
NNLINCAAAGSEETGFWPIPHHVPTGSPVGMSPGYSEHILGKPFYNNRAHSNYRAGMI
IDNGVKTTEASAKDKRPFLSIIISARYSPHQDADPLKPREPAIIRHPIAYKNQDHGAWLR
GGDVWLDSCRFAADNGIGLTLASGGTFPYDDGSKQEIKNSLFVGESGNVGTMMMDNRWIG
PGGLDHSGRTLPIGQNFPIRIGIQLYDGPINIQNCTFRKFLVLEGRHTSALAFRLNNAWQ
SCPHNVTGIAFEDVPIITSRVFPGEFPGWPNQDMDGDKTSVFHDVDSVSEYVPGSYLT
KNDNWLVRHPCINVPDWRGAI CSGCCYAQMVIQAYKTSNLRMKIIRKNDPFSHPPLYLEGA
LTRSTHYQQYQPVVTLQKGYTIHWDQTAPELAIWLIINFNKGDNIIRVGLCYPRGTTFSI
LSDVHNRLKQTSKTVGVFVRTLQMDKVRQSYPRGRSHYWDDEDSGLLFLKKAQNEREFK
AFCSMKGCERIKIKALIPKNAGVSDCTATAYPKFTERAVVDVPMPKKLFQSLKTKDHF
LEVKMESSKQHFHWNDFAYIEVDGKYPSSSEDIQVVVIDGNQGRVVVSHTSFRNSIL
QGIPWQLFNYVATIPDNSIVLMAKGRYVSRGFWTRVLEKLGADRGLKLEQMAFVGFK
GSFRPIWVTLTDETHKAKIFQVVPIPVVKKKL
```