



US 20020009738A1

(19) **United States**

(12) **Patent Application Publication**

Houghton et al.

(10) **Pub. No.: US 2002/0009738 A1**

(43) **Pub. Date: Jan. 24, 2002**

(54) **METHODS, COMPOSITIONS AND KITS FOR THE DETECTION AND MONITORING OF BREAST CANCER**

(76) Inventors: **Raymond L. Houghton**, Bothell, WA (US); **Davin C. Dillon**, Issaquah, WA (US); **David Molesh**, Kingston, WA (US); **Jiangchun Xu**, Bellevue, WA (US); **Barbara Zehentner**, Bainbridge Island, WA (US); **David H. Persing**, Redmond, WA (US)

Correspondence Address:
SEED INTELLECTUAL PROPERTY LAW GROUP PLLC
701 FIFTH AVE
SUITE 6300
SEATTLE, WA 98104-7092 (US)

(21) Appl. No.: **09/825,301**

(22) Filed: **Apr. 2, 2001**

Related U.S. Application Data

(63) Non-provisional of provisional application No. 60/194,241, filed on Apr. 3, 2000. Non-provisional of

provisional application No. 60/219,862, filed on Jul. 20, 2000. Non-provisional of provisional application No. 60/221,300, filed on Jul. 27, 2000. Non-provisional of provisional application No. 60/256,592, filed on Dec. 18, 2000.


Publication Classification

(51) **Int. Cl. 7** **C12Q 1/68; G01N 33/574**
(52) **U.S. Cl.** **435/6; 435/7.23**

(57)

ABSTRACT

Compositions and methods for the therapy and diagnosis of cancer, such as breast cancer, are disclosed. Compositions may comprise one or more breast tumor proteins, immunogenic portions thereof, or polynucleotides that encode such portions. Alternatively, a therapeutic composition may comprise an antigen presenting cell that expresses a breast tumor protein, or a T cell that is specific for cells expressing such a protein. Such compositions may be used, for example, for the prevention and treatment of diseases such as breast cancer. Diagnostic methods based on detecting a breast tumor protein, or mRNA encoding such a protein, in a sample are also provided.






Fig. 2




Fig. 3

GACAGCGGCTTCCTTGATCCTGCCACCCGCGACTGAACACCGACAGCAG 50
CAGCCTCACCATGAAGTTGCTGATGGTCCTCATGCTGGCGCCCTCTCCC 100
AGCACTGCTACGCAGGCTCTGGCTGCCCTTATTGGAGAATGTGATTCC 150
AAGACAATCAATCCACAAGTGTCTAAGACTGAATACAAAGAACTTCTCA 200
AGAGTTCATAGACGACAATGCCACTACAAATGCCATAGATGAATTGAAGG 250
AATGTTTCTTAACCAACGGATGAAACTCTGAGCAATGTTGAGGTGTTT 300
CTGCAATTAAATATGACAGCAGTCTTGTGATTATTAACTTCTGC 350
AAGACCTTGGCTCACAGAACTGCAGGGTATGGTAGAGAAACCAACTACGG 400
ATTGCTGCAAACACACCTCTCTTCTATGCTTTTACTACAAACTA 450
CAAGACAATTGTTGAAACCTGCTATACATGTTATTAAATAAAATTGATG 500
GCA 503

Fig. 4

CACTGCTACGCAGGCTCTGGCTGCCCTTATTGGAGAATGTGATTCCAA 50
GACAATCAATCCACAAGTGTCTAAGACTGAATACAAAGAACTTCTTCAG 100
AGTTCATAGACGACAATGCCACTACAAATGCCATAGATGAATTGAAGGAA 150
TGTTTCTTAACCAACGGATGAAACTCTGAGCAATGTTGAGGTGTTTAT 200
GCAATTAAATATGACAGCAGTCTTGTGATTATTGGCGGCCATCACC 250
ATCACCACATCACTAAGGTCCCGAGCTCGAATTCTGCAGATATCCATCACAC 300
T 301

Fig. 5

GGGACAGGGCTGAGGATGAGGAGAACCTGGGACCCAGAACAGCCGTGCCTGCCCGAAGTCCTGCCTGAGGCCTGAAGGACTTGCCCTAACAGAGCC 100
 TCAACAACTACCTGGTATTCTACTTCAGCCCTGGTGTAGCAGCTTCACATGAACATGAACACAGCCTCCATTGGCTCTGTGAGCTCTT
 CACTGAGAGGATGTGCATCCAGGGAGTCAGTTCAACGTCAGGTCGGCAGAAGT3ACAAGCTTCCCTGCCCTGGCTTGAGAACCTCACAGCAGGATAT
 AACAAATTCTCAGGCCAATTGGTGGAGAACCGTACAGATAGCGCTACTCTGGACATTGCAAGTATCTCTAGCATTAGAGAGAACATGGACT
 ACACAGCACCATACTCCGACAGCGCTGGATGGACCAAGCGGCTGGTGTGAAGGACAAAGAGCTCACTCTGGATGCCGCTCGTGGAGTTCC 500
 CTGGGTGCCAGATACTTACATTGGAGTCCAAGAAGTCCTCCTCCATGAAGTCAGTGTGGAAACAGGCTCATCCGCTCTCTCCAATGGCACGGTC
 CTGTATGCCCTCAGAACATCACGACAACGTGTGCATGTAACATGGATCTGTSTAATACCCCATGGACACACAGACATGCAAGTTGAGCTGGAAAGCTGG
 GCTATGATGGAAATGATGTGGAGTTACCTGGCTGAGAGGGAAACGACTCTGTGCGTGGACTGGAACACCTGCCGCTTGCTCAGTACACCATAGAGCGGTA
 TTTCACCTTAGTCACAGATCGCAGCAGGAGACAGGAATTACACTAGATTGGTCTACAGTTGAGCTTGGAGGAATGTTGTATTTCATTTGGAA
 ACCTACGTTCTCCACTTCTGGTGTGCTGGTTTCACTTGATCTCTCGATTCACTGCCAAGAACCTGCATTGGAGTGACGACCG 1000
 TGTATCAAAGACCACACTGATGATCGGGTCCGCACTCTCTCCAAACACCAACTGCTCATCAAGGCCA'CGATGTTACCTGGGATCTGCTTAG
 CTTTGTTGGGCTTGCTAGAATATGAGTTGCTACTACAGTTCCAGGAAAGATGGCAGGATGTTGATTATTCAAAACCCAGTAATGTTGA
 GTCAGTATTACTAATATCATCAACAGCTCATCTCAGTTAACCGGAAGATCAGTTGCCAGCATTGAAATTCCAGCGACAACGTTGACTACAGTG
 ACTTGACAATGAAAACAGCGACAAGTCAAGTTGCTTCCGAAAGATGGCAGGATGTTGATTATTCAAAACCCAGTAATGTTGA
 TCACTATTCCAAACTACTGTTCTTGTATTGCTAGCCAATGTTACTGGCATACTACATGTTTGAGTCATGTTAAATTCTTGCA 1500
 TGCCATAGGTCTCAACAGGACAAGATAATGATGTAATGGTATTAGGCAAGTGTGCAACCACATCCAATGGTGTACAAGTGTGACTAAATAATT
 TGAGTCTTCTGCTCAAAGAATGAAGCTCAACATTGTTCTAAGCTGTAGAAGTCCTAGATTAGGATCTGTAAATAGAAACATCAGTCATTCC
 TCTTCATCTTAATCAAGGACATTCCATGGAGGCCAAGATTACAATGTACTCAGGCTGTTATCGGTGGCTCCCTGGTTGCAATTACCTCATATA
 AAGAATGGGAAGGAGACCATTGGTAACCTCAAGTGTCAAGAGTTCTAAAGTAACTACATGTTTACTAAATCTGCAGTGTCTATAAAA
 TACATTGTTGCTATTAGGAGTAACATTAGTTCTGGTAAAATGAAATATGGGCTATGTCATTGCAACTTCAATTGGAAGTCATGCACTAAC 2000
 TCAATACCAAGATGAGTTAAATAATGAATTATTAAATACACAAACAGAATTATCCCAATTCCAAATAAGTCCTATCATTGAAAATTCAAATATA
 AGTGAAGAAAAATTAGTAGATCAACAATCTAAACAAATCCCTCGTTCTAAGATAACATGGATTCCCATCTGGAGGACTCTGAGGCTTATTCCCC
 CACTATGCATATCTTATCATTATTATACACACATCCATCTAAACTATACTAAAGCCTTTCCCATGTCATGGATGGAAATGGAAGATTGTTG
 TAACTTGTCTAGAAGTCTTAATATGGCTGTGCTGAAGGCTTGAGATTGAGTCATTCTAGCTGCTTATTACATAGTGTGGGTACTA
 AAAGTACTGGGTGACTCAGAGAGTCGCTGTCATTGCTACTCTAACACTGAGCAACACTCTCCAGTCGAGATCCCCTGTATCATTCC 2500
 AAGAGGGCATTCTCCCTTGTCTAATGTCAGGAATGATGTTTGTAGAAAACAAACTGCTTGACCCAGGAACAAGTGGCTTAGCTTAAGTAAACTT
 GGCTTGCTCAGATCCCTGATCCTCCAGCTGGCTGCTGAGTGGCTTATCCCGATGAGCAGGAGCGTGTGGCCCTGAGTACTGAACTTCTGAGT
 AACAAATGAGACACGTTACAGAACCTATGTTCAAGGTTGCGGGTGAGCTGCCCTCTCCAAATCAGGCCAGAGATGCACTTCCTCGGCCAGTCAGCCAAC
 AGTACCAAAAGTGTGTTGAGGTGCGCAGGGTAAGGCTTCAGCTCAGTTATTAGACAATCTGCCATCTTAAATTCTAGCTTCTGT
 TCTAATAATGACGGCTTACCTTCTGTCAAAATAACCAAGGCTCTAAAGATGATTCCCTCTGTAACCTCCAGGCCACAGGTTCTCATT
 CTTTCCCTTACTTCTCACAATTCACTGAGTTGATGACCTGATTGTTAAACAAATATTCTAACGGGAATGGTGGGAGTGTGGTGA 3000
 AAAGAGATGAAATGTGGTTGATGAGCCAATCATTTGTGATTGTTAAAGGAAATATCTGTTCTGAAACCCACTAACGATTGTT
 TTATATAAAAACAATGATAAGATGTGAACATGAAATAATACCATATTAGCTACCCACAAAAAA 3282

Fig. 6




Fig. 7




Fig. 8

Multiplex PCR assay:
Gene determination by amplicon size




Fig. 9




Fig. 10

METHODS, COMPOSITIONS AND KITS FOR THE DETECTION AND MONITORING OF BREAST CANCER

REFERENCE TO RELATED APPLICATIONS

[0001] This application claims the benefit of U.S. Provisional Application No. 60/194,241, filed Apr. 3, 2000; U.S. Provisional Application No. 60/219,862, filed Jul. 20, 2000; U.S. Provisional Application No. 60/221,300, filed Jul. 27, 2000; and U.S. Provisional Application No. 60/256,592, filed Dec. 18, 2000, each of which applications are incorporated herein by reference in their entirety.

GOVERNMENTAL SUPPORT

[0002] This work was supported in part by Grants CA-75794 and CA-80518 from the National Cancer Institute. The government may have certain rights in the invention.

TECHNICAL FIELD OF THE INVENTION

[0003] The present invention relates generally to the field of cancer diagnostics. More specifically, the present invention relates to methods, compositions and kits for the detection of cancer that employ oligonucleotide hybridization and/or amplification to simultaneously detect two or more tissue-specific polynucleotides in a biological sample suspected of containing cancer cells.

BACKGROUND OF THE INVENTION

[0004] Cancer remains a significant health problem throughout the world. The failure of conventional cancer treatment regimens can commonly be attributed, in part, to delayed disease diagnosis. Although significant advances have been made in the area of cancer diagnosis, there still remains a need for improved detection methodologies that permit early, reliable and sensitive determination of the presence of cancer cells.

[0005] Breast cancer is second only to lung cancer in mortality among women in the U.S., affecting more than 180,000 women each year and resulting in approximately 40,000-50,000 deaths annually. For women in North America, the life-time odds of getting breast cancer are one in eight.

[0006] Management of the disease currently relies on a combination of early diagnosis (through routine breast screening procedures) and aggressive treatment, which may include one or more of a variety of treatments such as surgery, radiotherapy, chemotherapy and hormone therapy. The course of treatment for a particular breast cancer is often selected based on a variety of prognostic parameters, including analysis of specific tumor markers. See, e.g., Porter-Jordan et al., *Breast Cancer* 8:73-100 (1994). The use of established markers often leads, however, to a result that is difficult to interpret; and the high mortality observed in breast cancer patients indicates that improvements are needed in the diagnosis of the disease.

[0007] The recent introduction of immunotherapeutic approaches to breast cancer treatment which are targeted to Her2/neu have provided significant motivation to identify additional breast cancer specific genes as targets for therapeutic antibodies and T-cell vaccines as well as for diagnosis

of the disease. To this end, mammaglobin, has been identified as one of the most breast-specific genes discovered to date, being expressed in approximately 70-80% of breast cancers. Because of its highly tissue-specific distribution, detection of mammaglobin gene expression has been used to identify micrometastatic lesions in lymph node tissues and, more recently, to detect circulating breast cancer cells in peripheral blood of breast cancer patients with known primary and metastatic lesions.

[0008] Mammaglobin is a homologue of a rabbit uteroglobin and the rat steroid binding protein subunit C3 and is a low molecular weight protein that is highly glycosylated. Watson et al., *Cancer Res.* 56:860-5 (1996); Watson et al., *Cancer Res.* 59:3028-3031 (1999); Watson et al., *Oncogene* 16:817-24 (1998). In contrast to its homologs, mammaglobin has been reported to be breast specific and overexpression has been described in breast tumor biopsies (23%), primary and metastatic breast tumors (~75%) with reports of the detection of mammaglobin mRNA expression in 91% of lymph nodes from metastatic breast cancer patients. Leygue et al., *J. Pathol.* 189:28-33 (1999) and Min et al., *Cancer Res.* 58:4581-4584 (1998).

[0009] Since mammaglobin gene expression is not a universal feature of breast cancer, the detection of this gene alone may be insufficient to permit the reliable detection of all breast cancers. Accordingly, what is needed in the art is a methodology that employs the detection of two or more breast cancer specific genes in order to improve the sensitivity and reliability of detection of micrometastases, for example, in lymph nodes and bone marrow and/or for recognition of anchorage-independent cells in the peripheral circulation.

[0010] The present invention achieves these and other related objectives by providing methods that are useful for the identification of tissue-specific polynucleotides, in particular tumor-specific polynucleotides, as well as methods, compositions and kits for the detection and monitoring of cancer cells in a patient afflicted with the disease.

SUMMARY OF THE INVENTION

[0011] By certain embodiments, the present invention provides methods for identifying one or more tissue-specific polynucleotides which methods comprise the steps of: (a) performing a genetic subtraction to identify a pool of polynucleotides from a tissue of interest; (b) performing a DNA microarray analysis to identify a first subset of said pool of polynucleotides of interest wherein each member polynucleotide of said first subset is at least two-fold over-expressed in said tissue of interest as compared to a control tissue; and (c) performing a quantitative polymerase chain reaction analysis on polynucleotides within said first subset to identify a second subset of polynucleotides that are at least two-fold over-expressed as compared to the control tissue. Preferred genetic subtractions are selected from the group consisting of differential display and cDNA subtraction and are described in further detail herein below.

[0012] Alternate embodiments of the present invention provide methods of identifying a subset of polynucleotides showing concordant and/or complementary tissue-specific expression profiles in a tissue of interest. Such methods comprise the steps of, (a) performing an expression analysis selected from the group consisting of DNA microarray and

quantitative PCR to identify a first polynucleotides that is at least two-fold over-expressed in a tissue of interest as compared to a control tissue; and (b) performing an expression analysis selected from the group consisting of DNA microarray and quantitative PCR to identify a first polynucleotides that is at least two-fold over-expressed in a tissue of interest as compared to a control tissue.

[0013] Further embodiments of the present invention provide methods for detecting the presence of a cancer cell in a patient. Such methods comprise the steps of: (a) obtaining a biological sample from the patient; (b) contacting the biological sample with a first oligonucleotide pair wherein the members of the first oligonucleotide pair hybridize, under moderately stringent conditions, to a first polynucleotide and the complement thereof, respectively; (c) contacting the biological sample with a second oligonucleotide pair wherein the members of the second oligonucleotide pair hybridize, under moderately stringent conditions, to a second polynucleotide and the complement thereof, respectively and wherein the first polynucleotide is unrelated in nucleotide sequence to the second polynucleotide; (d) amplifying the first polynucleotide and the second polynucleotide; and (e) detecting the amplified first polynucleotide and the amplified second polynucleotide; wherein the presence of the amplified first polynucleotide or amplified second polynucleotide indicates the presence of a cancer cell in the patient.

[0014] By some embodiments, detection of the amplified first and/or second polynucleotides may be preceded by a fractionation step such as, for example, gel electrophoresis. Alternatively or additionally, detection of the amplified first and/or second polynucleotides may be achieved by hybridization of a labeled oligonucleotide probe that hybridizes specifically, under moderately stringent conditions, to the first or second polynucleotide. Oligonucleotide labeling may be achieved by incorporating a radiolabeled nucleotide or by incorporating a fluorescent label.

[0015] In certain preferred embodiments, cells of a specific tissue type may be enriched from the biological sample prior to the steps of detection. Enrichment may be achieved by a methodology selected from the group consisting of cell capture and cell depletion. Exemplary cell capture methods include immunocapture and comprise the steps of: (a) adsorbing an antibody to a tissue-specific cell surface to cells said biological sample; (b) separating the antibody adsorbed tissue-specific cells from the remainder of the biological sample. Exemplary cell depletion may be achieved by cross-linking red cells and white cells followed by a subsequent fractionation step to remove the cross-linked cells.

[0016] Alternative embodiments of the present invention provide methods for determining the presence or absence of a cancer in a patient, comprising the steps of: (a) contacting a biological sample obtained from the patient with an oligonucleotide that hybridizes to a polynucleotide that encodes a breast tumor protein; (b) detecting in the sample a level of a polynucleotide (such as, for example, mRNA) that hybridizes to the oligonucleotide; and (c) comparing the level of polynucleotide that hybridizes to the oligonucleotide with a predetermined cut-off value, and therefrom determining the presence or absence of a cancer in the patient. Within certain embodiments, the amount of mRNA is detected via

polymerase chain reaction using, for example, at least one oligonucleotide primer that hybridizes to a polynucleotide encoding a polypeptide as recited above, or a complement of such a polynucleotide. Within other embodiments, the amount of mRNA is detected using a hybridization technique, employing an oligonucleotide probe that hybridizes to a polynucleotide that encodes a polypeptide as recited above, or a complement of such a polynucleotide.

[0017] In related aspects, methods are provided for monitoring the progression of a cancer in a patient, comprising the steps of: (a) contacting a biological sample obtained from a patient with an oligonucleotide that hybridizes to a polynucleotide that encodes a breast tumor protein; (b) detecting in the sample an amount of a polynucleotide that hybridizes to the oligonucleotide; (c) repeating steps (a) and (b) using a biological sample obtained from the patient at a subsequent point in time; and (d) comparing the amount of polynucleotide detected in step (c) with the amount detected in step (b) and therefrom monitoring the progression of the cancer in the patient.

[0018] Certain embodiments of the present invention provide that the step of amplifying said first polynucleotide and said second polynucleotide is achieved by the polymerase chain reaction (PCR).

[0019] Within certain embodiments, the cancer cell to be detected may be selected from the group consisting of prostate cancer, breast cancer, colon cancer, ovarian cancer, lung cancer head & neck cancer, lymphoma, leukemia, melanoma, liver cancer, gastric cancer, kidney cancer, bladder cancer, pancreatic cancer and endometrial cancer. Still further embodiments of the present invention provide that the biological sample is selected from the group consisting of blood, a lymph node and bone marrow. The lymph node may be a sentinel lymph node.

[0020] Within specific embodiments of present invention it is provided that the first polynucleotide is selected from the group consisting of gammaglobin, lipophilin B, GABA α (B899P), B726P, B511S, B533S, B305D and B311D. Other embodiments provide that the second polynucleotide is selected from the group consisting of gammaglobin, lipophilin B, GABA α (B899P), B726P, B511S, B533S, B305D and B311D.

[0021] Alternate embodiments of the present invention provide methods for detecting the presence or absence of a cancer in a patient, comprising the steps of: (a) contacting a biological sample obtained from a patient with a first oligonucleotide that hybridizes to a polynucleotide selected from the group consisting of gammaglobin and lipophilin B; (b) contacting the biological sample with a second oligonucleotide that hybridizes to a polynucleotide sequence selected from the group consisting of GABA α (B899P), B726P, B511S, B533S, B305D and B311D; (c) detecting in the sample an amount of a polynucleotide that hybridizes to at least one of the oligonucleotides; and (d) comparing the amount of polynucleotide that hybridizes to the oligonucleotide to a predetermined cut-off value, and therefrom determining the presence or absence of a cancer in the patient.

[0022] According to certain embodiments, oligonucleotides may be selected from those disclosed herein such as those presented in SEQ ID Nos:33-72. By other embodiments, the amount of polynucleotide that hybridizes to the

oligonucleotide is determined using a polymerase chain reaction. Alternatively, the amount of polynucleotide that hybridizes to the oligonucleotide may be determined using a hybridization assay.

[0023] Still other embodiments of the present invention provide methods for determining the presence or absence of a cancer cell in a patient, comprising the steps of: (a) contacting a biological sample obtained from a patient with a first oligonucleotide that hybridizes to a polynucleotide selected from the group consisting of a polynucleotide depicted in SEQ ID NO:73 and SEQ ID NO:74 or complement thereof; (b) contacting the biological sample with a second oligonucleotide that hybridizes to a polynucleotide depicted in SEQ ID NO:75 or complement thereof; (c) contacting the biological sample with a third oligonucleotide that hybridizes to a polynucleotide selected from the group consisting of a polynucleotide depicted in SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:6 and SEQ ID NO:7 or complement thereof; (d) contacting the biological sample with a fourth oligonucleotide that hybridizes to a polynucleotide selected from the group consisting of a polynucleotide depicted in SEQ ID NO:11 or complement thereof; (e) contacting the biological sample with a fifth oligonucleotide that hybridizes to a polynucleotide selected from the group consisting of a polynucleotide depicted in SEQ ID NO:13, 15 and 17 or complement thereof; (f) contacting the biological sample with a sixth oligonucleotide that hybridizes to a polynucleotide selected from the group consisting of a polynucleotide depicted in SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23 and SEQ ID NO:24 or complement thereof; (g) contacting the biological sample with a seventh oligonucleotide that hybridizes to a polynucleotide depicted in SEQ ID NO:30 or complement thereof; (h) contacting the biological sample with an eighth oligonucleotide that hybridizes to a polynucleotide depicted in SEQ ID NO:32 or complement thereof; (i) contacting the biological sample with a ninth oligonucleotide that hybridizes to a polynucleotide depicted in SEQ ID NO:76 or complement thereof; (j) detecting in the sample a hybridized oligonucleotide of any one of steps (a) through (i); and (j) comparing the amount of polynucleotide that hybridizes to the oligonucleotide to a predetermined cut-off value, wherein the presence of a hybridized oligonucleotide in any one of steps (a) through (i) in excess of the pre-determined cut-off value indicates the presence of a cancer cell in the biological sample of said patient.

[0024] Other related embodiments of the present invention provide methods for determining the presence or absence of a cancer cell in a patient, comprising the steps of: (a) contacting a biological sample obtained from a patient with a first oligonucleotide and a second oligonucleotide wherein said first and second oligonucleotides hybridize under moderately stringent conditions to a first and a second polynucleotide selected from the group selected from the group consisting of SEQ ID NO:73, SEQ ID NO:74, SEQ ID NO:75, SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:11, SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:30, SEQ ID NO:32, and SEQ ID NO:76 and wherein said first polynucleotide is unrelated structurally to said second polynucleotide; (b) detecting in the sample said first and said second hybridized oligonucleotides; and (c) comparing the amount of polynucleotide that

hybridizes to the oligonucleotide to a predetermined cut-off value, wherein the presence of a hybridized first oligonucleotide or a hybridized second oligonucleotide in excess of the pre-determined cut-off value indicates the presence of a cancer cell in the biological sample of said patient.

[0025] Other related embodiments of the present invention provide methods for determining the presence or absence of a cancer cell in a patient, comprising the steps of: (a) contacting a biological sample obtained from a patient with a first oligonucleotide and a second oligonucleotide wherein said first and second oligonucleotides hybridize under moderately stringent conditions to a first and a second polynucleotide are both tissue-specific polynucleotides of the cancer to be detected and wherein said first polynucleotide is unrelated structurally to said second polynucleotide; (b) detecting in the sample said first and said second hybridized oligonucleotides; and (c) comparing the amount of polynucleotide that hybridizes to the oligonucleotide to a predetermined cut-off value, wherein the presence of a hybridized first oligonucleotide or a hybridized second oligonucleotide in excess of the pre-determined cut-off value indicates the presence of a cancer cell in the biological sample of said patient.

[0026] In other related aspects, the present invention further provides compositions useful in the methods disclosed herein. Exemplary compositions comprise two or more oligonucleotide primer pairs each one of which specifically hybridizes to a distinct polynucleotide. Exemplary oligonucleotide primers suitable for compositions of the present invention are disclosed herein by SEQ ID NOs: 33-71. Exemplary polynucleotides suitable for compositions of the present invention are disclosed in SEQ ID NO:73, SEQ ID NO:74, SEQ ID NO:75, SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:11, SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:30, SEQ ID NO:32, and SEQ ID NO:76.

[0027] The present invention also provides kits that are suitable for performing the detection methods of the present invention. Exemplary kits comprise oligonucleotide primer pairs each one of which specifically hybridizes to a distinct polynucleotide. Within certain embodiments, kits according to the present invention may also comprise a nucleic acid polymerase and suitable buffer. Exemplary oligonucleotide primers suitable for kits of the present invention are disclosed herein by SEQ ID NOs: 33-71. Exemplary polynucleotides suitable for kits of the present invention are disclosed in SEQ ID NO:73, SEQ ID NO:74, SEQ ID NO:75, SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:11, SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:30, SEQ ID NO:32, and SEQ ID NO:76.

[0028] These and other aspects of the present invention will become apparent upon reference to the following detailed description and attached drawings. All references disclosed herein are hereby incorporated by reference in their entirety as if each was incorporated individually.

**BRIEF DESCRIPTION OF THE DRAWINGS
AND SEQUENCE IDENTIFIERS**

[0029] **FIG. 1** shows the mRNA expression profiles for B311D, B533S and B726P as determined using quantitative PCR (Taqman™). Abbreviations: B.T.: Breast tumor; B.M.: Bone marrow; B.R.: Breast reduction.

[0030] **FIG. 2** shows the relationship of B533S expression to pathological stage of tumor. Tissues from normal breast (8), benign breast disorders (3), and breast tumors stage I (5), stage II (6), stage III (7), stage IV (3) and metastases (1 lymph node and 3 pleural effusions) were tested in real-time PCR. The data is expressed as the mean copies/ng β -actin for each group tested and the line is the calculated trend line.

[0031] **FIGS. 3A and 3B** show the gene complementation of B305D C-form, B726P, GABA π and gammaglobin in metastases and primary tumors, respectively. The cut-off for each of the genes was 6.57, 1.65, 4.58 and 3.56 copies/ng β -Actin based on the mean of the negative normal tissues plus 3 standard deviations.

[0032] **FIG. 4** shows the full-length cDNA sequence for gammaglobin.

[0033] **FIG. 5** shows the determined cDNA sequence of the open reading frame encoding a gammaglobin recombinant polypeptide expressed in *E. coli*.

[0034] **FIG. 6** shows the full-length cDNA sequence for GABA π .

[0035] **FIG. 7** shows the mRNA expression levels for gammaglobin, GABA π , B305D (C form) and B726P in breast tumor and normal samples determined using real-time PCR and the SYBR detection system. Abbreviations: BT: Breast tumor; BR: Breast reduction; A. PBMC: Activated peripheral blood mononuclear cells; R. PBMC: resting PBMC; T. Gland: Thyroid gland; S. Cord: Spinal Cord; A. Gland: Adrenal gland; B. Marrow: Bone marrow; S. Muscle: Skeletal muscle.

[0036] **FIG. 8** is a bar graph showing a comparison between the LipophilinB alone and the LipophilinB-B899P-B305D-C-B726 multiplex assays tested on a panel of breast tumor samples. Abbreviations: BT: Breast tumor; BR: Breast reduction; SCID: severe combined immunodeficiency.

[0037] **FIG. 9** is a gel showing the unique band length of four amplification products of tumor genes of interest (gammaglobin, B305D, B899P, B726P) tested in a multiplex Real-time PCR assay.

[0038] **FIG. 10** shows a comparison of a multiplex assay using intron-exon border spanning primers (bottom panel) and those using non-optimized primers (top panel), to detect breast cancer cells in a panel of lymph node tissues.

[0039] SEQ ID NO:1 is the determined cDNA sequence for a first splice variant of B305D isoform A.

[0040] SEQ ID NO:2 is the amino acid sequence encoded by the sequence of SEQ ID NO:1.

[0041] SEQ ID NO:3 is the determined cDNA sequence for a second splice variant of B305D isoform A.

[0042] SEQ ID NO:4 is the amino acid sequence encoded by the sequence of SEQ ID NO:3.

[0043] SEQ ID NO:5-7 are the determined cDNA sequences for three splice variants of B305D isoform C.

[0044] SEQ ID NO:8-10 are the amino acid sequences encoded by the sequence of SEQ ID NO:5-7, respectively.

[0045] SEQ ID NO:11 is the determined cDNA sequence for B311D.

[0046] SEQ ID NO:12 is the amino acid sequence encoded by the sequence of SEQ ID NO:11.

[0047] SEQ ID NO:13 is the determined cDNA sequence of a first splice variant of B726P.

[0048] SEQ ID NO:14 is the amino acid sequence encoded by the sequence of SEQ ID NO:13.

[0049] SEQ ID NO:15 is the determined cDNA sequence of a second splice variant of B726P.

[0050] SEQ ID NO:16 is the amino acid sequence encoded by the sequence of SEQ ID NO:15.

[0051] SEQ ID NO:17 is the determined cDNA sequence of a third splice variant of B726P.

[0052] SEQ ID NO:18 is the amino acid sequence encoded by the sequence of SEQ ID NO:17.

[0053] SEQ ID NO:19-24 are the determined cDNA sequences of further splice variants of B726P.

[0054] SEQ ID NO:25-29 are the amino acid sequences encoded by SEQ ID NO: 19-24, respectively.

[0055] SEQ ID NO:30 is the determined cDNA sequence for B511S.

[0056] SEQ ID NO:31 is the amino acid sequence encoded by SEQ ID NO:30.

[0057] SEQ ID NO:32 is the determined cDNA sequence for B533S.

[0058] SEQ ID NO:33 is the DNA sequence of Lipophilin B forward primer.

[0059] SEQ ID NO:34 is the DNA sequence of Lipophilin B reverse primer.

[0060] SEQ ID NO:35 is the DNA sequence of Lipophilin B probe.

[0061] SEQ ID NO:36 is the DNA sequence of GABA (B899P) forward primer.

[0062] SEQ ID NO:37 is the DNA sequence of GABA (B899P) reverse primer.

[0063] SEQ ID NO:38 is the DNA sequence of GABA (B899P) probe.

[0064] SEQ ID NO:39 is the DNA sequence of B305D (C form) forward primer.

[0065] SEQ ID NO:40 is the DNA sequence of B305D (C form) reverse primer.

[0066] SEQ ID NO:41 is the DNA sequence of B305D (C form) probe.

[0067] SEQ ID NO:42 is the DNA sequence of B726P forward primer.

[0068] SEQ ID NO:43 is the DNA sequence of B726P reverse primer.

[0069] SEQ ID NO:44 is the DNA sequence of B726P probe.

[0070] SEQ ID NO:45 is the DNA sequence of Actin forward primer.

[0071] SEQ ID NO:46 is the DNA sequence of Actin reverse primer.

[0072] SEQ ID NO:47 is the DNA sequence of Actin probe.

[0073] SEQ ID NO:48 is the DNA sequence of Mammaglobin forward primer.

[0074] SEQ ID NO:49 is the DNA sequence of Mammaglobin reverse primer.

[0075] SEQ ID NO:50 is the DNA sequence of Mammaglobin probe.

[0076] SEQ ID NO:51 is the DNA sequence of a second GABA (B899P) reverse primer.

[0077] SEQ ID NO:52 is the DNA sequence of a second B726P forward primer.

[0078] SEQ ID NO:53 is the DNA sequence of a GABA B899P-INT forward primer.

[0079] SEQ ID NO:54 is the DNA sequence of a GABA B899P-INT reverse primer.

[0080] SEQ ID NO:55 is the DNA sequence of a GABA B899P-INT Taqman probe.

[0081] SEQ ID NO:56 is the DNA sequence of a B305D-INT forward primer.

[0082] SEQ ID NO:57 is the DNA sequence of a B305D-INT reverse primer.

[0083] SEQ ID NO:58 is the DNA sequence of a B305D-INT Taqman probe.

[0084] SEQ ID NO:59 is the DNA sequence of a B726-INT forward primer.

[0085] SEQ ID NO:60 is the DNA sequence of a B726-INT reverse primer.

[0086] SEQ ID NO:61 is the DNA sequence of a B726-INT Taqman probe.

[0087] SEQ ID NO:62 is the DNA sequence of a GABA B899P Taqman probe.

[0088] SEQ ID NO:63 is the DNA sequence of a B311D forward primer.

[0089] SEQ ID NO:64 is the DNA sequence of a B311D reverse primer.

[0090] SEQ ID NO:65 is the DNA sequence of a B311D Taqman probe.

[0091] SEQ ID NO:66 is the DNA sequence of a B533S forward primer.

[0092] SEQ ID NO:67 is the DNA sequence of a B533S reverse primer.

[0093] SEQ ID NO:68 is the DNA sequence of a B533S Taqman probe.

[0094] SEQ ID NO:69 is the DNA sequence of a B511S forward primer.

[0095] SEQ ID NO:70 is the DNA sequence of a B511S reverse primer.

[0096] SEQ ID NO:71 is the DNA sequence of a B511S Taqman probe.

[0097] SEQ ID NO:72 is the DNA sequence of a GABA π reverse primer.

[0098] SEQ ID NO:73 is the full-length cDNA sequence for mammaglobin.

[0099] SEQ ID NO:74 is the determined cDNA sequence of the open reading frame encoding a mammaglobin recombinant polypeptide expressed in *E. coli*.

[0100] SEQ ID NO:75 is the full-length cDNA sequence for GABA π .

[0101] SEQ ID NO:76 is the full-length cDNA sequence for lipophilin B.

[0102] SEQ ID NO:77 is the amino acid sequence encoded by the sequence of SEQ ID NO:76.

DETAILED DESCRIPTION OF THE INVENTION

[0103] As noted above, the present invention is directed generally to methods that are suitable for the identification of tissue-specific polynucleotides as well as to methods, compositions and kits that are suitable for the diagnosis and monitoring of cancer. While certain exemplary methods, compositions and kits disclosed herein are directed to the identification, detection and monitoring of breast cancer, in particular breast cancer-specific polynucleotides, it will be understood by those of skill in the art that the present invention is generally applicable to the identification, detection and monitoring of a wide variety of cancers, and the associated over-expressed polynucleotides, including, for example, prostate cancer, breast cancer, colon cancer, ovarian cancer, lung cancer, head & neck cancer, lymphoma, leukemia, melanoma, liver cancer, gastric cancer, kidney cancer, bladder cancer, pancreatic cancer and endometrial cancer. Thus, it will be apparent that the present invention is not limited solely to the identification of breast cancer-specific polynucleotides or to the detection and monitoring of breast cancer.

[0104] Identification of Tissue-specific Polynucleotides

[0105] Certain embodiments of the present invention provide methods, compositions and kits for the detection of a cancer cell within a biological sample. These methods comprise the step of detecting one or more tissue-specific polynucleotide(s) from a patient's biological sample the over-expression of which polynucleotides indicates the presence of a cancer cell within the patient's biological sample. Accordingly, the present invention also provides methods that are suitable for the identification of tissue-specific polynucleotides. As used herein, the phrases "tissue-specific polynucleotides" or "tumor-specific polynucleotides" are meant to include all polynucleotides that are at least two-fold over-expressed as compared to one or more control tissues. As discussed in further detail herein below, over-expression of a given polynucleotide may be assessed, for example, by microarray and/or quantitative real-time polymerase chain reaction (Real-time PCRTM) methodologies.

[0106] Exemplary methods for detecting tissue-specific polynucleotides may comprise the steps of: (a) performing a genetic subtraction to identify a pool of polynucleotides from a tissue of interest; (b) performing a DNA microarray analysis to identify a first subset of said pool of polynucleotides of interest wherein each member polynucleotide of said first subset is at least two-fold over-expressed in said tissue of interest as compared to a control tissue; and (c) performing a quantitative polymerase chain reaction analysis on polynucleotides within said first subset to identify a second subset of polynucleotides that are at least two-fold over-expressed as compared to said control tissue.

[0107] Polynucleotides Generally

[0108] As used herein, the term "polynucleotide" refers generally to either DNA or RNA molecules. Polynucleotides may be naturally occurring as normally found in a biological sample such as blood, serum, lymph node, bone marrow, sputum, urine and tumor biopsy samples. Alternatively, polynucleotides may be derived synthetically by, for example, a nucleic acid polymerization reaction. As will be recognized by the skilled artisan, polynucleotides may be single-stranded (coding or antisense) or double-stranded, and may be DNA (genomic, cDNA or synthetic) or RNA molecules. RNA molecules include HnRNA molecules, which contain introns and correspond to a DNA molecule in a one-to-one manner, and mRNA molecules, which do not contain introns. Additional coding or non-coding sequences may, but need not, be present within a polynucleotide of the present invention, and a polynucleotide may, but need not, be linked to other molecules and/or support materials.

[0109] Polynucleotides may comprise a native sequence (i.e. an endogenous sequence that encodes a tumor protein, such as a breast tumor protein, or a portion thereof) or may comprise a variant, or a biological or antigenic functional equivalent of such a sequence. Polynucleotide variants may contain one or more substitutions, additions, deletions and/or insertions, as further described below. The term "variants" also encompasses homologous genes of xenogenic origin.

[0110] When comparing polynucleotide or polypeptide sequences, two sequences are said to be "identical" if the sequence of nucleotides or amino acids in the two sequences is the same when aligned for maximum correspondence, as described below. Comparisons between two sequences are typically performed by comparing the sequences over a comparison window to identify and compare local regions of sequence similarity. A "comparison window" as used herein, refers to a segment of at least about 20 contiguous positions, usually 30 to about 75, 40 to about 50, in which a sequence may be compared to a reference sequence of the same number of contiguous positions after the two sequences are optimally aligned.

[0111] Optimal alignment of sequences for comparison may be conducted using the Megalign program in the Lasergene suite of bioinformatics software (DNASTAR, Inc., Madison, Wis.), using default parameters. This program embodies several alignment schemes described in the following references: Dayhoff, M. O. (1978) A model of evolutionary change in proteins—Matrices for detecting distant relationships. In Dayhoff, M. O. (ed.) *Atlas of Protein Sequence and Structure*, National Biomedical Research Foundation, Washington D.C. Vol. 5, Suppl. 3, pp. 345-358; Hein J. (1990) Unified Approach to Alignment and Phylo-

genes pp. 626-645 *Methods in Enzymology* vol. 183, Academic Press, Inc., San Diego, Calif.; Higgins, D. G. and Sharp, P. M. (1989) *CABIOS* 5:151-153; Myers, E. W. and Muller W. (1988) *CABIOS* 4:11-17; Robinson, E. D. (1971) *Comb. Theor* 11:105; Santou, N. Nes, M. (1987) *Mol. Biol. Evol.* 4:406-425; Sneath, P. H. A. and Sokal, R. R. (1973) *Numerical Taxonomy—the Principles and Practice of Numerical Taxonomy*, Freeman Press, San Francisco, Calif.; Wilbur, W. J. and Lipman, D. J. (1983) *Proc. Natl. Acad. Sci. USA* 80:726-730.

[0112] Alternatively, optimal alignment of sequences for comparison may be conducted by the local identity algorithm of Smith and Waterman (1981) *Add. APL. Math* 2:482, by the identity alignment algorithm of Needleman and Wunsch (1970) *J. Mol. Biol.* 48:443, by the search for similarity methods of Pearson and Lipman (1988) *Proc. Natl. Acad. Sci. USA* 85: 2444, by computerized implementations of these algorithms (GAP, BESTFIT, BLAST, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group (GCG), 575 Science Dr., Madison, Wis.), or by inspection.

[0113] One preferred example of algorithms that are suitable for determining percent sequence identity and sequence similarity are the BLAST and BLAST 2.0 algorithms, which are described in Altschul et al. (1977) *Nucl. Acids Res.* 25:3389-3402 and Altschul et al. (1990) *J. Mol. Biol.* 215:403-410, respectively. BLAST and BLAST 2.0 can be used, for example with the parameters described herein, to determine percent sequence identity for the polynucleotides and polypeptides of the invention. Software for performing BLAST analyses is publicly available through the National Center for Biotechnology Information. In one illustrative example, cumulative scores can be calculated using, for nucleotide sequences, the parameters M (reward score for a pair of matching residues; always >0) and N (penalty score for mismatching residues; always <0). For amino acid sequences, a scoring matrix can be used to calculate the cumulative score. Extension of the word hits in each direction are halted when: the cumulative alignment score falls off by the quantity X from its maximum achieved value; the cumulative score goes to zero or below, due to the accumulation of one or more negative-scoring residue alignments; or the end of either sequence is reached. The BLAST algorithm parameters W, T and X determine the sensitivity and speed of the alignment. The BLASTN program (for nucleotide sequences) uses as defaults a wordlength (W) of 11, and expectation (E) of 10, and the BLOSUM62 scoring matrix (see Henikoff and Henikoff (1989) *Proc. Natl. Acad. Sci. USA* 89:10915) alignments, (B) of 50, expectation (E) of 10, M=5, N=-4 and a comparison of both strands.

[0114] Preferably, the "percentage of sequence identity" is determined by comparing two optimally aligned sequences over a window of comparison of at least 20 positions, wherein the portion of the polynucleotide or polypeptide sequence in the comparison window may comprise additions or deletions (i.e., gaps) of 20 percent or less, usually 5 to 15 percent, or 10 to 12 percent, as compared to the reference sequences (which does not comprise additions or deletions) for optimal alignment of the two sequences. The percentage is calculated by determining the number of positions at which the identical nucleic acid bases or amino acid residue occurs in both sequences to yield the number of matched positions, dividing the number of matched positions by the

total number of positions in the reference sequence (i.e., the window size) and multiplying the results by 100 to yield the percentage of sequence identity.

[0115] Therefore, the present invention encompasses polynucleotide and polypeptide sequences having substantial identity to the sequences disclosed herein, for example those comprising at least 50% sequence identity, preferably at least 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% or higher, sequence identity compared to a polynucleotide or polypeptide sequence of this invention using the methods described herein, (e.g., BLAST analysis using standard parameters, as described below). One skilled in this art will recognize that these values can be appropriately adjusted to determine corresponding identity of proteins encoded by two nucleotide sequences by taking into account codon degeneracy, amino acid similarity, reading frame positioning and the like.

[0116] In additional embodiments, the present invention provides isolated polynucleotides and polypeptides comprising various lengths of contiguous stretches of sequence identical to or complementary to one or more of the sequences disclosed herein. For example, polynucleotides are provided by this invention that comprise at least about 15, 20, 30, 40, 50, 75, 100, 150, 200, 300, 400, 500 or 1000 or more contiguous nucleotides of one or more of the sequences disclosed herein as well as all intermediate lengths there between. It will be readily understood that "intermediate lengths", in this context, means any length between the quoted values, such as 16, 17, 18, 19, etc.; 21, 22, 23, etc.; 30, 31, 32, etc.; 50, 51, 52, 53, etc.; 100, 101, 102, 103, etc.; 150, 151, 152, 153, etc.; including all integers through 200-500; 500-1,000, and the like.

[0117] The polynucleotides of the present invention, or fragments thereof, regardless of the length of the coding sequence itself, may be combined with other DNA sequences, such as promoters, polyadenylation signals, additional restriction enzyme sites, multiple cloning sites, other coding segments, and the like, such that their overall length may vary considerably. It is therefore contemplated that a nucleic acid fragment of almost any length may be employed, with the total length preferably being limited by the ease of preparation and use in the intended recombinant DNA protocol. For example, illustrative DNA segments with total lengths of about 10,000, about 5000, about 3000, about 2,000, about 1,000, about 500, about 200, about 100, about 50 base pairs in length, and the like, (including all intermediate lengths) are contemplated to be useful in many implementations of this invention.

[0118] In other embodiments, the present invention is directed to polynucleotides that are capable of hybridizing under moderately stringent conditions to a polynucleotide sequence provided herein, or a fragment thereof, or a complementary sequence thereof. Hybridization techniques are well known in the art of molecular biology. For purposes of illustration, suitable moderately stringent conditions for testing the hybridization of a polynucleotide of this invention with other polynucleotides include prewashing in a solution of 5×SSC, 0.5% SDS, 1.0 mM EDTA (pH 8.0); hybridizing at 50° C.-65° C., 5×SSC, overnight; followed by washing twice at 65° C. for 20 minutes with each of 2×, 0.5× and 0.2×SSC containing 0.1% SDS.

[0119] Moreover, it will be appreciated by those of ordinary skill in the art that, as a result of the degeneracy of the

genetic code, there are many nucleotide sequences that encode a polypeptide as described herein. Some of these polynucleotides bear minimal homology to the nucleotide sequence of any native gene. Nonetheless, polynucleotides that vary due to differences in codon usage are specifically contemplated by the present invention. Further, alleles of the genes comprising the polynucleotide sequences provided herein are within the scope of the present invention. Alleles are endogenous genes that are altered as a result of one or more mutations, such as deletions, additions and/or substitutions of nucleotides. The resulting mRNA and protein may, but need not, have an altered structure or function. Alleles may be identified using standard techniques (such as hybridization, amplification and/or database sequence comparison).

[0120] Microarray Analyses

[0121] Polynucleotides that are suitable for detection according to the methods of the present invention may be identified, as described in more detail below, by screening a microarray of cDNAs for tissue and/or tumor-associated expression (e.g., expression that is at least two-fold greater in a tumor than in normal tissue, as determined using a representative assay provided herein). Such screens may be performed, for example, using a Synteni microarray (Palo Alto, Calif.) according to the manufacturer's instructions (and essentially as described by Schena et al., *Proc. Natl. Acad. Sci. USA* 93:10614-10619 (1996) and Heller et al., *Proc. Natl. Acad. Sci. USA* 94:2150-2155 (1997)).

[0122] Microarray is an effective method for evaluating large numbers of genes but due to its limited sensitivity it may not accurately determine the absolute tissue distribution of low abundance genes or may underestimate the degree of overexpression of more abundant genes due to signal saturation. For those genes showing overexpression by microarray expression profiling, further analysis was performed using quantitative RT-PCR based on Taqman™ probe detection, which comprises a greater dynamic range of sensitivity. Several different panels of normal and tumor tissues, distant metastases and cell lines were used for this purpose.

[0123] Quantitative Real-time Polymerase Chain Reaction

[0124] Suitable polynucleotides according to the present invention may be further characterized or, alternatively, originally identified by employing a quantitative PCR methodology such as, for example, the Real-time PCR methodology. By this methodology, tissue and/or tumor samples, such as, e.g., metastatic tumor samples, may be tested along side the corresponding normal tissue sample and/or a panel of unrelated normal tissue samples.

[0125] Real-time PCR (see Gibson et al., *Genome Research* 6:995-1001, 1996; Heid et al., *Genome Research* 6:986-994, 1996) is a technique that evaluates the level of PCR product accumulation during amplification. This technique permits quantitative evaluation of mRNA levels in multiple samples. Briefly, mRNA is extracted from tumor and normal tissue and cDNA is prepared using standard techniques.

[0126] Real-time PCR may, for example, be performed either on the ABI 7700 Prism or on a GeneAmp® 5700 sequence detection system (PE Biosystems, Foster City, Calif.). The 7700 system uses a forward and a reverse primer in combination with a specific probe with a 5' fluorescent

reporter dye at one end and a 3' quencher dye at the other end (Taqman T). When the Real-time PCR is performed using Taq DNA polymerase with 5'-3' nuclease activity, the probe is cleaved and begins to fluoresce allowing the reaction to be monitored by the increase in fluorescence (Real-time). The 5700 system uses SYBR® green, a fluorescent dye, that only binds to double stranded DNA, and the same forward and reverse primers as the 7700 instrument. Matching primers and fluorescent probes may be designed according to the primer express program (PE Biosystems, Foster City, Calif.). Optimal concentrations of primers and probes are initially determined by those of ordinary skill in the art. Control (e.g., β -actin) primers and probes may be obtained commercially from, for example, Perkin Elmer/Applied Biosystems (Foster City, Calif.).

[0127] To quantitate the amount of specific RNA in a sample, a standard curve is generated using a plasmid containing the gene of interest. Standard curves are generated using the Ct values determined in the real-time PCR, which are related to the initial cDNA concentration used in the assay. Standard dilutions ranging from 10-10⁶ copies of the gene of interest are generally sufficient. In addition, a standard curve is generated for the control sequence. This permits standardization of initial RNA content of a tissue sample to the amount of control for comparison purposes.

[0128] In accordance with the above, and as described further below, the present invention provides the illustrative breast tissue- and/or tumor-specific polynucleotides mammaglobin, lipophilin B, GABA π (B899P), B726P, B511S, B533S, B305D and B311D having sequences set forth in SEQ ID NO: 1, 3, 5-7, 11, 13, 15, 17, 19-24, 30, 32, and 73-76 illustrative polypeptides encoded thereby having amino acid sequences set forth in SEQ ID NO: 2, 4, 8-10, 12, 14, 16, 18, 25-29 and 31 and 77 that may be suitably employed in the detection of cancer, more specifically, breast cancer.

[0129] The methods disclosed herein will also permit the identification of additional and/or alternative polynucleotides that are suitable for the detection of a wide range of cancers including, but not limited to, prostate cancer, breast cancer, colon cancer, ovarian cancer, lung cancer head & neck cancer, lymphoma, leukemia, melanoma, liver cancer, gastric cancer, kidney cancer, bladder cancer, pancreatic cancer and endometrial cancer.

[0130] Methodologies for the Detection of Cancer

[0131] In general, a cancer cell may be detected in a patient based on the presence of one or more polynucleotides within cells of a biological sample (for example, blood, lymph nodes, bone marrow, sera, sputum, urine and/or tumor biopsies) obtained from the patient. In other words, such polynucleotides may be used as markers to indicate the presence or absence of a cancer such as, e.g., breast cancer.

[0132] As discussed in further detail herein, the present invention achieves these and other related objectives by providing a methodology for the simultaneous detection of more than one polynucleotide, the presence of which is diagnostic of the presence of cancer cells in a biological sample. Each of the various cancer detection methodologies disclosed herein have in common a step of hybridizing one or more oligonucleotide primers and/or probes, the hybrid-

ization of which is demonstrative of the presence of a tumor- and/or tissue-specific polynucleotide. Depending on the precise application contemplated, it may be preferred to employ one or more intron-spanning oligonucleotides that are inoperative against polynucleotide of genomic DNA and, thus, these oligonucleotides are effective in substantially reducing and/or eliminating the detection of genomic DNA in the biological sample.

[0133] Further disclosed herein are methods for enhancing the sensitivity of these detection methodologies by subjecting the biological samples to be tested to one or more cell capture and/or cell depletion methodologies.

[0134] By certain embodiments of the present invention, the presence of a cancer cell in a patient may be determined by employing the following steps: (a) obtaining a biological sample from said patient; (b) contacting said biological sample with a first oligonucleotide that hybridizes to a first polynucleotide said first polynucleotide selected from the group consisting of polynucleotides depicted in SEQ ID NO:73 and SEQ ID NO:74; (c) contacting said biological sample with a second oligonucleotide that hybridizes to a second polynucleotide selected from the group consisting of SEQ ID NO: 1, 3, 5-7, 11, 13, 15, 17, 19-24, 30, 32, and 75; (d) detecting in said sample an amount of a polynucleotide that hybridizes to at least one of the oligonucleotides; and (e) comparing the amount of the polynucleotide that hybridizes to said oligonucleotide to a predetermined cut-off value, and therefrom determining the presence or absence of a cancer in the patient.

[0135] Alternative embodiments of the present invention provide methods wherein the presence of a cancer cell in a patient is determined by employing the steps of: (a) obtaining a biological sample from said patient; (b) contacting said biological sample with a first oligonucleotide that hybridizes to a first polynucleotide said first polynucleotide depicted in SEQ ID NO:76; (c) contacting said biological sample with a second oligonucleotide that hybridizes to a second polynucleotide selected from the group consisting of SEQ ID NO: 1, 3, 5-7, 11, 13, 15, 17, 19-24, 30, 32, and 75; (d) detecting in said sample an amount of a polynucleotide that hybridizes to at least one of the oligonucleotides; and (e) comparing the amount of the polynucleotide that hybridizes to said oligonucleotide to a predetermined cut-off value, and therefrom determining the presence or absence of a cancer in the patient.

[0136] Other embodiments of the present invention provide methods for determining the presence or absence of a cancer in a patient. Such methods comprise the steps of: (a) obtaining a biological sample from said patient; (b) contacting said biological sample obtained from a patient with a first oligonucleotide that hybridizes to a polynucleotide sequence selected from the group consisting of polynucleotides depicted in SEQ ID NO:73, SEQ ID NO:74 and SEQ ID NO:76; (c) contacting said biological sample with a second oligonucleotide that hybridizes to a polynucleotide as depicted in SEQ ID NO:75; (d) contacting said biological sample with a third oligonucleotide that hybridizes to a polynucleotide selected from the group consisting of polynucleotides depicted in SEQ ID NO:5, SEQ ID NO:6 and SEQ ID NO:7; (e) contacting said biological sample with a fourth oligonucleotide that hybridizes to a polynucleotide selected from the group consisting of polynucleotides

depicted in SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23 and SEQ ID NO:24; (f) detecting in said biological sample an amount of a polynucleotide that hybridizes to at least one of said oligonucleotides; and (g) comparing the amount of polynucleotide that hybridizes to the oligonucleotide to a predetermined cut-off value, and therefrom determining the presence or absence of a cancer in the patient.

[0137] To permit hybridization under assay conditions, oligonucleotide primers and probes should comprise an oligonucleotide sequence that has at least about 60%, preferably at least about 75% and more preferably at least about 90%, identity to a portion of a polynucleotide encoding a breast tumor protein that is at least 10 nucleotides, and preferably at least 20 nucleotides, in length. Preferably, oligonucleotide primers hybridize to a polynucleotide encoding a polypeptide described herein under moderately stringent conditions, as defined above. Oligonucleotide primers which may be usefully employed in the diagnostic methods described herein preferably are at least 10-40 nucleotides in length. In a preferred embodiment, the oligonucleotide primers comprise at least 10 contiguous nucleotides, more preferably at least 15 contiguous nucleotides, of a DNA molecule having a sequence recited in SEQ ID NO: 1, 3, 5-7, 11, 13, 15, 17, 19-24, 30, 32 and 73-76. Techniques for both PCR based assays and hybridization assays are well known in the art (see, for example, Mullis et al., *Cold Spring Harbor Symp. Quant. Biol.*, 51:263, 1987; Erlich ed., *PCR Technology*, Stockton Press, NY, 1989).

[0138] The present invention also provides amplification-based methods for detecting the presence of a cancer cell in a patient. Exemplary methods comprise the steps of (a) obtaining a biological sample from a patient; (b) contacting the biological sample with a first oligonucleotide pair the first pair comprising a first oligonucleotide and a second oligonucleotide wherein the first oligonucleotide and the second oligonucleotide hybridize to a first polynucleotide and the complement thereof, respectively; (c) contacting the biological sample with a second oligonucleotide pair the second pair comprising a third oligonucleotide and a fourth oligonucleotide wherein the third and the fourth oligonucleotide hybridize to a second polynucleotide and the complement thereof, respectively, and wherein the first polynucleotide is unrelated in nucleotide sequence to the second polynucleotide; (d) amplifying the first polynucleotide and the second polynucleotide; and (e) detecting the amplified first polynucleotide and the amplified second polynucleotide; wherein the presence of the amplified first polynucleotide or the amplified second polynucleotide indicates the presence of a cancer cell in the patient.

[0139] Methods according to the present invention are suitable for identifying polynucleotides obtained from a wide variety of biological sample such as, for example, blood, serum, lymph node, bone marrow, sputum, urine and tumor biopsy sample. In certain preferred embodiments, the biological sample is either blood, a lymph node or bone marrow. In other embodiments of the present invention, the lymph node may be a sentinel lymph node.

[0140] It will be apparent that the present methods may be employed in the detection of a wide variety of cancers. Exemplary cancers include, but are not limited to, prostate

cancer, breast cancer, colon cancer, ovarian cancer, lung cancer head & neck cancer, lymphoma, leukemia, melanoma, liver cancer, gastric cancer, kidney cancer, bladder cancer, pancreatic cancer and endometrial cancer.

[0141] Certain exemplary embodiments of the present invention provide methods wherein the polynucleotides to be detected are selected from the group consisting of gammaglobin, lipophilin B, GABA α (B899P), B726P, B511S, B533S, B305D and B311D. Alternatively and/or additionally, polynucleotides to be detected may be selected from the group consisting of those depicted in SEQ ID NO:73, SEQ ID NO:74, SEQ ID NO:75, SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:11, SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:30, SEQ ID NO:32, and SEQ ID NO:76.

[0142] Suitable exemplary oligonucleotide probes and/or primers that may be used according to the methods of the present invention are disclosed herein by SEQ ID NOs:33-35 and 63-72. In certain preferred embodiments that eliminate the background detection of genomic DNA, the oligonucleotides may be intron spanning oligonucleotides. Exemplary intron spanning oligonucleotides suitable for the detection of various polynucleotides disclosed herein are depicted in SEQ ID NOs:36-62.

[0143] Depending on the precise application contemplated, the artisan may prefer to detect the tissue- and/or tumor-specific polynucleotides by detecting a radiolabel and detecting a fluorophore. More specifically, the oligonucleotide probe and/or primer may comprise a detectable moiety such as, for example, a radiolabel and/or a fluorophore.

[0144] Alternatively or additionally, methods of the present invention may also comprise a step of fractionation prior to detection of the tissue- and/or tumor-specific polynucleotides such as, for example, by gel electrophoresis.

[0145] In other embodiments, methods described herein may be used as to monitor the progression of cancer. By these embodiments, assays as provided for the diagnosis of a cancer may be performed over time, and the change in the level of reactive polypeptide(s) or polynucleotide(s) evaluated. For example, the assays may be performed every 24-72 hours for a period of 6 months to 1 year, and thereafter performed as needed. In general, a cancer is progressing in those patients in whom the level of polypeptide or polynucleotide detected increases over time. In contrast, the cancer is not progressing when the level of reactive polypeptide or polynucleotide either remains constant or decreases with time.

[0146] Certain in vivo diagnostic assays may be performed directly on a tumor. One such assay involves contacting tumor cells with a binding agent. The bound binding agent may then be detected directly or indirectly via a reporter group. Such binding agents may also be used in histological applications. Alternatively, polynucleotide probes may be used within such applications.

[0147] As noted above, to improve sensitivity, multiple breast tumor protein markers may be assayed within a given sample. It will be apparent that binding agents specific for different proteins provided herein may be combined within a single assay. Further, multiple primers or probes may be

used concurrently. The selection of tumor protein markers may be based on routine experiments to determine combinations that results in optimal sensitivity. In addition, or alternatively, assays for tumor proteins provided herein may be combined with assays for other known tumor antigens.

[0148] Cell Enrichment

[0149] In other aspects of the present invention, cell capture technologies may be used prior to polynucleotide detection to improve the sensitivity of the various detection methodologies disclosed herein.

[0150] Exemplary cell enrichment methodologies employ immunomagnetic beads that are coated with specific monoclonal antibodies to surface cell markers, or tetrameric antibody complexes, may be used to first enrich or positively select cancer cells in a sample. Various commercially available kits may be used, including Dynabeads® Epithelial Enrich (Dynal Biotech, Oslo, Norway), StemSep™ (Stem-Cell Technologies, Inc., Vancouver, BC), and RosetteSep (StemCell Technologies). The skilled artisan will recognize that other readily available methodologies and kits may also be suitably employed to enrich or positively select desired cell populations.

[0151] Dynabeads® Epithelial Enrich contains magnetic beads coated with mAbs specific for two glycoprotein membrane antigens expressed on normal and neoplastic epithelial tissues. The coated beads may be added to a sample and the sample then applied to a magnet, thereby capturing the cells bound to the beads. The unwanted cells are washed away and the magnetically isolated cells eluted from the beads and used in further analyses.

[0152] RosetteSep can be used to enrich cells directly from a blood sample and consists of a cocktail of tetrameric antibodies that target a variety of unwanted cells and crosslinks them to glycophorin A on red blood cells (RBC) present in the sample, forming rosettes. When centrifuged over Ficoll, targeted cells pellet along with the free RBC.

[0153] The combination of antibodies in the depletion cocktail determines which cells will be removed and consequently which cells will be recovered. Antibodies that are available include, but are not limited to: CD2, CD3, CD4, CD5, CD8, CD10, CD11b, CD14, CD15, CD16, CD19, CD20, CD24, CD25, CD29, CD33, CD34, CD36, CD38, CD41, CD45, CD45RA, CD45RO, CD56, CD66B, CD66e, HLA-DR, IgE, and TCR $\alpha\beta$. Additionally, it is contemplated in the present invention that mAbs specific for breast tumor antigens, can be developed and used in a similar manner. For example, mAbs that bind to tumor-specific cell surface antigens may be conjugated to magnetic beads, or formulated in a tetrameric antibody complex, and used to enrich or positively select metastatic breast tumor cells from a sample.

[0154] Once a sample is enriched or positively selected, cells may be further analysed. For example, the cells may be lysed and RNA isolated. RNA may then be subjected to RT-PCR analysis using breast tumor-specific multiplex primers in a Real-time PCR assay as described herein.

[0155] In another aspect of the present invention, cell capture technologies may be used in conjunction with Real-time PCR to provide a more sensitive tool for detection of metastatic cells expressing breast tumor antigens. Detection of breast cancer cells in bone marrow samples, peripheral

blood, and small needle aspiration samples is desirable for diagnosis and prognosis in breast cancer patients.

[0156] Probes and Primers

[0157] As noted above and as described in further detail herein, certain methods, compositions and kits according to the present invention utilize two or more oligonucleotide primer pairs for the detection of cancer. The ability of such nucleic acid probes to specifically hybridize to a sequence of interest will enable them to be of use in detecting the presence of complementary sequences in a biological sample.

[0158] Alternatively, in other embodiments, the probes and/or primers of the present invention may be employed for detection via nucleic acid hybridization. As such, it is contemplated that nucleic acid segments that comprise a sequence region of at least about 15 nucleotide long contiguous sequence that has the same sequence as, or is complementary to, a 15 nucleotide long contiguous sequence of a polynucleotide to be detected will find particular utility. Longer contiguous identical or complementary sequences, e.g., those of about 20, 30, 40, 50, 100, 200, 500, 1000 (including all intermediate lengths) and even up to full length sequences will also be of use in certain embodiments.

[0159] Oligonucleotide primers having sequence regions consisting of contiguous nucleotide stretches of 10-14, 15-20, 30, 50, or even of 100-200 nucleotides or so (including intermediate lengths as well), identical or complementary to a polynucleotide to be detected, are particularly contemplated as primers for use in amplification reactions such as, e.g., the polymerase chain reaction (PCR™). This would allow a polynucleotide to be analyzed, both in diverse biological samples such as, for example, blood, lymph nodes and bone marrow.

[0160] The use of a primer of about 15-25 nucleotides in length allows the formation of a duplex molecule that is both stable and selective. Molecules having contiguous complementary sequences over stretches greater than 15 bases in length are generally preferred, though, in order to increase stability and selectivity of the hybrid, and thereby improve the quality and degree of specific hybrid molecules obtained. One will generally prefer to design primers having gene-complementary stretches of 15 to 25 contiguous nucleotides, or even longer where desired.

[0161] Primers may be selected from any portion of the polynucleotide to be detected. All that is required is to review the sequence, such as those exemplary polynucleotides set forth in SEQ ID NO: 1, 3, 5-7, 11, 13, 15, 17, 19-24, 30, 32, 73-75 (FIGS. 3-6, respectively) and SEQ ID NO:76 (lipophilin B) or to any continuous portion of the sequence, from about 15-25 nucleotides in length up to and including the full length sequence, that one wishes to utilize as a primer. The choice of primer sequences may be governed by various factors. For example, one may wish to employ primers from towards the termini of the total sequence. The exemplary primers disclosed herein may optionally be used for their ability to selectively form duplex molecules with complementary stretches of the entire polynucleotide of interest such as those set forth in SEQ ID NO: 1, 3, 5-7, 11, 13, 15, 17, 19-24, 30, 32, 73-75 (FIGS. 3-6, respectively), and SEQ ID NO:76 (lipophilin B).

[0162] The present invention further provides the nucleotide sequence of various exemplary oligonucleotide primers and probes, set forth in SEQ ID NOs: 33-71, that may be used, as described in further detail herein, according to the methods of the present invention for the detection of cancer.

[0163] Oligonucleotide primers according to the present invention may be readily prepared routinely by methods commonly available to the skilled artisan including, for example, directly synthesizing the primers by chemical means, as is commonly practiced using an automated oligonucleotide synthesizer. Depending on the application envisioned, one will typically desire to employ varying conditions of hybridization to achieve varying degrees of selectivity of probe towards target sequence. For applications requiring high selectivity, one will typically desire to employ relatively stringent conditions to form the hybrids, e.g., one will select relatively low salt and/or high temperature conditions, such as provided by a salt concentration of from about 0.02 M to about 0.15 M salt at temperatures of from about 50° C. to about 70° C. Such selective conditions tolerate little, if any, mismatch between the probe and the template or target strand, and would be particularly suitable for isolating related sequences.

[0164] Polynucleotide Amplification Techniques

[0165] Each of the specific embodiments outlined herein for the detection of cancer has in common the detection of a tissue- and/or tumor-specific polynucleotide via the hybridization of one or more oligonucleotide primers and/or probes. Depending on such factors as the relative number of cancer cells present in the biological sample and/or the level of polynucleotide expression within each cancer cell, it may be preferred to perform an amplification step prior to performing the steps of detection. For example, at least two oligonucleotide primers may be employed in a polymerase chain reaction (PCR) based assay to amplify a portion of a breast tumor cDNA derived from a biological sample, wherein at least one of the oligonucleotide primers is specific for (i.e., hybridizes to) a polynucleotide encoding the breast tumor protein. The amplified cDNA may optionally be subjected to a fractionation step such as, for example, gel electrophoresis.

[0166] A number of template dependent processes are available to amplify the target sequences of interest present in a sample. One of the best known amplification methods is the polymerase chain reaction (PCR™) which is described in detail in U.S. Pat. Nos. 4,683,195, 4,683,202 and 4,800,159, each of which is incorporated herein by reference in its entirety. Briefly, in PCR™, two primer sequences are prepared which are complementary to regions on opposite complementary strands of the target sequence. An excess of deoxynucleoside triphosphates is added to a reaction mixture along with a DNA polymerase (e.g., Taq polymerase). If the target sequence is present in a sample, the primers will bind to the target and the polymerase will cause the primers to be extended along the target sequence by adding on nucleotides. By raising and lowering the temperature of the reaction mixture, the extended primers will dissociate from the target to form reaction products, excess primers will bind to the target and to the reaction product and the process is repeated. Preferably reverse transcription and PCR™ amplification procedure may be performed in order to quantify the amount of mRNA amplified. Polymerase chain reaction methodologies are well known in the art.

[0167] One preferred methodology for polynucleotide amplification employs RT-PCR, in which PCR is applied in conjunction with reverse transcription. Typically, RNA is extracted from a biological sample, such as blood, serum, lymph node, bone marrow, sputum, urine and tumor biopsy samples, and is reverse transcribed to produce cDNA molecules. PCR amplification using at least one specific primer generates a cDNA molecule, which may be separated and visualized using, for example, gel electrophoresis. Amplification may be performed on biological samples taken from a patient and from an individual who is not afflicted with a cancer. The amplification reaction may be performed on several dilutions of cDNA spanning two orders of magnitude. A two-fold or greater increase in expression in several dilutions of the test patient sample as compared to the same dilutions of the non-cancerous sample is typically considered positive.

[0168] Any of a variety of commercially available kits may be used to perform the amplification step. One such amplification technique is inverse PCR (see Triglia et al., *Nucl. Acids Res.* 16:8186, 1988), which uses restriction enzymes to generate a fragment in the known region of the gene. The fragment is then circularized by intramolecular ligation and used as a template for PCR with divergent primers derived from the known region. Within an alternative approach, sequences adjacent to a partial sequence may be retrieved by amplification with a primer to a linker sequence and a primer specific to a known region. The amplified sequences are typically subjected to a second round of amplification with the same linker primer and a second primer specific to the known region. A variation on this procedure, which employs two primers that initiate extension in opposite directions from the known sequence, is described in WO 96/38591. Another such technique is known as "rapid amplification of cDNA ends" or RACE. This technique involves the use of an internal primer and an external primer, which hybridizes to a polyA region or vector sequence, to identify sequences that are 5' and 3' of a known sequence. Additional techniques include capture PCR (Lagerstrom et al., *PCR Methods Appl.* 1:111-19, 1991) and walking PCR (Parker et al., *Nucl. Acids. Res.* 19:3055-60, 1991). Other methods employing amplification may also be employed to obtain a full length cDNA sequence.

[0169] Another method for amplification is the ligase chain reaction (referred to as LCR), disclosed in Eur. Pat. Appl. Publ. No. 320,308 (specifically incorporated herein by reference in its entirety). In LCR, two complementary probe pairs are prepared, and in the presence of the target sequence, each pair will bind to opposite complementary strands of the target such that they abut. In the presence of a ligase, the two probe pairs will link to form a single unit. By temperature cycling, as in PCR™, bound ligated units dissociate from the target and then serve as "target sequences" for ligation of excess probe pairs. U.S. Pat. No. 4,883,750, incorporated herein by reference in its entirety, describes an alternative method of amplification similar to LCR for binding probe pairs to a target sequence.

[0170] Qbeta Replicase, described in PCT Intl. Pat. Appl. Publ. No. PCT/US87/00880, incorporated herein by reference in its entirety, may also be used as still another amplification method in the present invention. In this method, a replicative sequence of RNA that has a region

complementary to that of a target is added to a sample in the presence of an RNA polymerase. The polymerase will copy the replicative sequence that can then be detected.

[0171] An isothermal amplification method, in which restriction endonucleases and ligases are used to achieve the amplification of target molecules that contain nucleotide 5'-[α -thio]triphosphates in one strand of a restriction site (Walker et al., 1992, incorporated herein by reference in its entirety), may also be useful in the amplification of nucleic acids in the present invention.

[0172] Strand Displacement Amplification (SDA) is another method of carrying out isothermal amplification of nucleic acids which involves multiple rounds of strand displacement and synthesis, i.e. nick translation. A similar method, called Repair Chain Reaction (RCR) is another method of amplification which may be useful in the present invention and is involves annealing several probes throughout a region targeted for amplification, followed by a repair reaction in which only two of the four bases are present. The other two bases can be added as biotinylated derivatives for easy detection. A similar approach is used in SDA.

[0173] Sequences can also be detected using a cyclic probe reaction (CPR). In CPR, a probe having a 3' and 5' sequences of non-target DNA and an internal or "middle" sequence of the target protein specific RNA is hybridized to DNA which is present in a sample. Upon hybridization, the reaction is treated with RNaseH, and the products of the probe are identified as distinctive products by generating a signal that is released after digestion. The original template is annealed to another cycling probe and the reaction is repeated. Thus, CPR involves amplifying a signal generated by hybridization of a probe to a target gene specific expressed nucleic acid.

[0174] Still other amplification methods described in Great Britain Pat. Appl. No. 2 202 328, and in PCT Intl. Pat. Appl. Publ. No. PCT/US89/01025, each of which is incorporated herein by reference in its entirety, may be used in accordance with the present invention. In the former application, "modified" primers are used in a PCR-like, template and enzyme dependent synthesis. The primers may be modified by labeling with a capture moiety (e.g., biotin) and/or a detector moiety (e.g., enzyme). In the latter application, an excess of labeled probes is added to a sample. In the presence of the target sequence, the probe binds and is cleaved catalytically. After cleavage, the target sequence is released intact to be bound by excess probe. Cleavage of the labeled probe signals the presence of the target sequence.

[0175] Other nucleic acid amplification procedures include transcription-based amplification systems (TAS) (Kwoh et al., 1989; PCT Intl. Pat. Appl. Publ. No. WO 88/10315, incorporated herein by reference in its entirety), including nucleic acid sequence based amplification (NASBA) and 3SR. In NASBA, the nucleic acids can be prepared for amplification by standard phenol/chloroform extraction, heat denaturation of a sample, treatment with lysis buffer and minispin columns for isolation of DNA and RNA or guanidinium chloride extraction of RNA. These amplification techniques involve annealing a primer that has sequences specific to the target sequence. Following polymerization, DNA/RNA hybrids are digested with RNase H while double stranded DNA molecules are heat-denatured again. In either case the single stranded DNA is made fully

double stranded by addition of second target-specific primer, followed by polymerization. The double stranded DNA molecules are then multiply transcribed by a polymerase such as T7 or SP6. In an isothermal cyclic reaction, the RNAs are reverse transcribed into DNA, and transcribed once again with a polymerase such as T7 or SP6. The resulting products, whether truncated or complete, indicate target-specific sequences.

[0176] Eur. Pat. Appl. Publ. No. 329,822, incorporated herein by reference in its entirety, disclose a nucleic acid amplification process involving cyclically synthesizing single-stranded RNA ("ssRNA"), ssDNA, and double-stranded DNA (dsDNA), which may be used in accordance with the present invention. The ssRNA is a first template for a first primer oligonucleotide, which is elongated by reverse transcriptase (RNA-dependent DNA polymerase). The RNA is then removed from resulting DNA:RNA duplex by the action of ribonuclease H (RNase H, an RNase specific for RNA in a duplex with either DNA or RNA). The resultant ssDNA is a second template for a second primer, which also includes the sequences of an RNA polymerase promoter (exemplified by T7 RNA polymerase) 5' to its homology to its template. This primer is then extended by DNA polymerase (exemplified by the large "Klenow" fragment of *E. coli* DNA polymerase I), resulting as a double-stranded DNA ("dsDNA") molecule, having a sequence identical to that of the original RNA between the primers and having additionally, at one end, a promoter sequence. This promoter sequence can be used by the appropriate RNA polymerase to make many RNA copies of the DNA. These copies can then re-enter the cycle leading to very swift amplification. With proper choice of enzymes, this amplification can be done isothermally without addition of enzymes at each cycle. Because of the cyclical nature of this process, the starting sequence can be chosen to be in the form of either DNA or RNA.

[0177] PCT Intl. Pat. Appl. Publ. No. WO 89/06700, incorporated herein by reference in its entirety, disclose a nucleic acid sequence amplification scheme based on the hybridization of a promoter/primer sequence to a target single-stranded DNA ("ssDNA") followed by transcription of many RNA copies of the sequence. This scheme is not cyclic; i.e. new templates are not produced from the resultant RNA transcripts. Other amplification methods include "RACE" (Frohman, 1990), and "one-sided PCR" (Ohara, 1989) which are well-known to those of skill in the art.

[0178] Compositions and Kits for the Detection of Cancer

[0179] The present invention further provides kits for use within any of the above diagnostic methods. Such kits typically comprise two or more components necessary for performing a diagnostic assay. Components may be compounds, reagents, containers and/or equipment. For example, one container within a kit may contain a monoclonal antibody or fragment thereof that specifically binds to a breast tumor protein. Such antibodies or fragments may be provided attached to a support material, as described above. One or more additional containers may enclose elements, such as reagents or buffers, to be used in the assay. Such kits may also, or alternatively, contain a detection reagent as described above that contains a reporter group suitable for direct or indirect detection of antibody binding.

[0180] The present invention also provides kits that are suitable for performing the detection methods of the present

invention. Exemplary kits comprise oligonucleotide primer pairs each one of which specifically hybridizes to a distinct polynucleotide. Within certain embodiments, kits according to the present invention may also comprise a nucleic acid polymerase and suitable buffer. Exemplary oligonucleotide primers suitable for kits of the present invention are disclosed herein by SEQ ID NOs: 33-71. Exemplary polynucleotides suitable for kits of the present invention are disclosed in SEQ ID NO:73, SEQ ID NO:74, SEQ ID NO:75, SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:11, SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:30, SEQ ID NO:32, and lipophilin B.

[0181] Alternatively, a kit may be designed to detect the level of mRNA encoding a breast tumor protein in a biological sample. Such kits generally comprise at least one oligonucleotide probe or primer, as described above, that hybridizes to a polynucleotide encoding a breast tumor protein. Such an oligonucleotide may be used, for example, within a PCR or hybridization assay. Additional components that may be present within such kits include a second oligonucleotide and/or a diagnostic reagent or container to facilitate the detection of a polynucleotide encoding a breast tumor protein.

[0182] In other related aspects, the present invention further provides compositions useful in the methods disclosed herein. Exemplary compositions comprise two or more oligonucleotide primer pairs each one of which specifically hybridizes to a distinct polynucleotide. Exemplary oligonucleotide primers suitable for compositions of the present invention are disclosed herein by SEQ ID NOs: 33-71. Exemplary polynucleotides suitable for compositions of the present invention are disclosed in SEQ ID NO:73, SEQ ID NO:74, SEQ ID NO:75, SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:11, SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:30, SEQ ID NO:32, and lipophilin B.

[0183] The following Examples are offered by way of illustration and not by way of limitation.

EXAMPLES

Example 1

Differential Display

[0184] This example discloses the use of differential display to enrich for polynucleotides that are over-expressed in breast tumor tissues.

[0185] Differential display was performed as described in the literature (see, e.g., Liang, P. et al., *Science* 257:967-971 (1993), incorporated herein by reference in its entirety) with the following modifications: (a) PCR amplification products were visualized on silver stained gels (b) genetically matched pairs of tissues were used to eliminate polymorphic variation (c) two different dilutions of cDNA were used as template to eliminate any dilutional effects (see, Mou, E. et al., *Biochem Biophys Res Commun.* 199:564-569 (1994), incorporated herein by reference in its entirety).

Example 2

Preparation of cDNA Subtraction library

[0186] This example discloses the preparation of a breast tumor cDNA subtraction library enriched in breast tumor specific polynucleotides.

[0187] cDNA library subtraction was performed as described with some modification. See, Hara, T. et al., *Blood* 84: 189-199 (1994), incorporated herein by reference in its entirety. The breast tumor library (tracer) that was made from a pool of three breast tumors was subtracted with normal breast library (driver) to identify breast tumor specific genes. More recent subtractions utilized 6-10 normal tissues as driver to subtract out common genes more efficiently, with an emphasis on essential tissues along with one "immunological" tissue (e.g., spleen, lymph node, or PBMC), to assist in the removal of cDNAs derived from lymphocyte infiltration in tumors. The breast tumor specific subtracted cDNA library was generated as follows: driver cDNA library was digested with EcoRI, NotI, and SfuI (SfuI cleaves the vector), filled in with DNA polymerase klenow fragment. After phenol-chloroform extraction and ethanol precipitation, the DNA was labeled with Photoprobe biotin and dissolved in H₂O. Tracer cDNA library was digested with BamHI and XhoI, phenol chloroform extracted, passed through Chroma spin-400 columns, ethanol precipitated, and mixed with driver DNA for hybridization at 68° C. for 20 hours [long hybridization (LH)]. The reaction mixture was then subjected to the streptavidin treatment followed by phenol/chloroform extraction for a total of four times. Subtracted DNA was precipitated and subjected to a hybridization at 68° C. for 2 hours with driver DNA again [short hybridization (SH)]. After removal of biotinylated double-stranded DNA, subtracted cDNA was ligated into BamHI/XhoI site of Chloramphenicol resistant pBCSK+and transformed into ElectroMax *E. coli* DH10B cells by electroporation to generate subtracted cDNA library. To clone less abundant breast tumor specific genes, cDNA library subtraction was repeated by subtracting the tracer cDNA library with the driver cDNA library plus abundant cDNAs from primary subtractions. This resulted in the depletion of these abundant sequences and the generation of subtraction libraries that contain less abundant sequences.

[0188] To analyze the subtracted cDNA library, plasmid DNA was prepared from 100-200 independent clones, which were randomly picked from the subtracted library, and characterized by DNA sequencing. The determined cDNA and expected amino acid sequences for the isolated cDNAs were compared to known sequences using the most recent Genbank and human EST databases.

Example 3

PCR-subtraction

[0189] This example discloses PCR subtraction to enrich for breast tumor specific polynucleotides.

[0190] PCR-subtraction was performed essentially as described in the literature. See, Diatchenko, L. et al., *Proc Natl Acad Sci USA.* 93:6025-6030 (1996) and Yang, G. P. et al., *Nucleic acids Res.* 27:1517-23 (1999), incorporated herein by reference in their entirety. Briefly, this type of subtraction works by ligating two different adapters to

different aliquots of a restriction enzyme digested tester (breast tumor) cDNA sample, followed by mixing of the testers separately with excess driver (without adapters). This first hybridization results in normalization of single stranded tester specific cDNA due to the second order kinetics of hybridization. These separate hybridization reactions are then mixed without denaturation, and a second hybridization performed which produces the target molecules; double stranded cDNA fragments containing both of the different adapters. Two rounds of PCR were performed, which results in the exponential amplification of the target population molecules (normalized tester specific cDNAs), while other fragments were either unamplified or only amplified in a linear manner. The subtractions performed included a pool of breast tumors subtracted with a pool of normal breast and a pool of breast tumors subtracted with a pool of normal tissues including PBMC, brain, pancreas, liver, small intestine, stomach, heart and kidney.

[0191] Prior to cDNA synthesis RNA was treated with DNase I (Ambion) in the presence of RNasin (Promega Biotech, Madison, Wis.) to remove DNA contamination. The cDNA for use in real-time PCR tissue panels was prepared using 25 μ l Oligo dT (Boehringer-Mannheim) primer with superscript II reverse transcriptase (Gibco BRL, Bethesda, Md.).

Example 4

Detection of Breast Cancer using Breast-specific Antigens

[0192] The isolation and characterization of the breast-specific antigens B511S and B533S is described in U.S. patent application Ser. No. 09/346,327, filed Jul. 2, 1999, the disclosure of which is hereby incorporated by reference in its entirety. The determined cDNA sequence for B511S is provided in SEQ ID NO: 30, with the corresponding amino acid sequence being provided in SEQ ID NO: 31. The determined cDNA sequence for B533S is provided in SEQ ID NO: 32. The isolation and characterization of the breast-specific antigen B726P is described in U.S. patent application Ser. No. 09/285,480, filed Apr. 2, 1999, and Ser. No. 09/433,826, filed Nov. 3, 1999, the disclosures of which are hereby incorporated by reference in their entirety.

[0193] The determined cDNA sequences for splice variants of B726P are provided in SEQ ID NO: 13, 15, 17 and 19-24, with the corresponding amino acid sequences being provided in SEQ ID NO: 14, 16, 18 and 25-29.

[0194] The isolation and characterization of the breast-specific antigen B305D forms A and C has been described in U.S. patent application Ser. No. 09/429,755, filed Oct. 28, 1999, the disclosure of which is hereby incorporated by reference in its entirety. Determined cDNA sequences for B305D isoforms A and C are provided in SEQ ID NO: 1, 3 and 5-7, with the corresponding amino acid sequences being provided in SEQ ID NO: 2, 4 and 8-10.

[0195] The isolation and characterization of the breast-specific antigen B311D has been described in U.S. patent application Ser. No. 09/289,198, filed Apr. 9, 1999, the disclosure of which is hereby incorporated by reference in its entirety. The determined cDNA sequence for B311D is provided in SEQ ID NO:11, with the corresponding amino acid sequence being provided in SEQ ID NO:12.

[0196] cDNA sequences for mammaglobin are provided in FIGS. 4 and 5, with the cDNA sequence for GABA π being provided in FIG. 6 and are disclosed in SEQ ID NOs: 73-75, respectively.

[0197] The isolation and characterization of the breast-specific antigen lipophilin B has been described in U.S. patent application Ser. No. 09/780,842, filed Feb. 8, 2001, the disclosure of which is hereby incorporated by reference in its entirety. The determined cDNA sequence for lipophilin B is provided in SEQ ID NO:76, with the corresponding amino acid sequence being provided in SEQ ID NO:77. The nucleotide sequences of several sequence variants of lipophilin B are also described in the Ser. No. 09/780,842 application.

Example 5

Microarray Analysis

[0198] This example discloses the use of microarray analyses to identify polynucleotides that are at least two-fold overexpressed in breast tumor tissue samples as compared to normal breast tissue samples.

[0199] mRNA expression of the polynucleotides of interest was performed as follows. cDNA for the different genes was prepared as described above and arrayed on a glass slide (Incyte, Palo Alto, Calif.). The arrayed cDNA was then hybridized with a 1:1 mixture of Cy3 or Cy5 fluorescent labeled first strand cDNAs obtained from polyA+RNA from breast tumors, normal breast and normal tissues and other tumors as described in Shalon, D. et al., *Genome Res.* 6:639-45 (1996), incorporated herein by reference in its entirety. Typically Cy3 (Probe 1) was attached to cDNAs from breast tumors and Cy5 (Probe 2) to normal breast tissue or other normal tissues. Both probes were allowed to compete with the immobilized gene specific cDNAs on the chip, washed then scanned for fluorescence intensity of the individual Cy3 and Cy5 fluorescence to determine extent of hybridization. Data were analyzed using GEMTOOLS software (Incyte, Palo Alto, Calif.) which enabled the overexpression patterns of breast tumors to be compared with normal tissues by the ratios of Cy3/Cy5. The fluorescence intensity was also related to the expression level of the individual genes. DNA microarray analyses was used primarily as a screening tool to determine tissue/tumor specificity of cDNA's recovered from the differential display, cDNA library and PCR subtractions, prior to more rigorous analysis by quantitative RT-PCR, northern blotting, and immunohistochemistry. Microarray analysis was performed on two microchips. A total of 3603 subtracted cDNA's and 197 differential display templates were evaluated to identify 40 candidates for further analysis by quantitative PCR.

[0200] From these candidates, several were chosen on the basis of favorable tissue specificity profiles, including B305D, B311D, B726P, B511S and B533S, indicating their overexpression profiles in breast tumors and/or normal breast versus other normal tissues. It was evident that the expression of these genes showed a high degree of specificity for breast tumors and/or breast tissue. In addition, these genes have in many cases complementary expression profiles.

[0201] The two known breast-specific genes, mammaglobin and γ -aminobutyrate type A receptor π subunit

(GABA π) were also subjected to microarray analysis. mRNA expression of mammaglobin has been previously described to be upregulated in proliferating breast tissue, including breast tumors. See, (Watson et al., *Cancer Res.*, 56: 860-5 (1996); Watson et al., *Cancer Res.*, 59: 3028-3031 (1999); Watson et al., *Oncogene*, 16:817-24 (1998), incorporated herein by reference in their entirety). The GABA π mRNA levels were over-expressed in breast tumors. Previous studies had demonstrated its overexpression in uterus and to some degree in prostate and lung (Hedblom et al., *J Biol. Chem.* 272:15346-15350 (1997)) but no previous study had indicated elevated levels in breast tumors.

other 3' end (TaqmanTM). During PCR the Taq DNA polymerase with its 5'-3' nuclease activity cleaved the probe which began to fluoresce, allowing the reaction to be monitored by the increase in fluorescence (Real-time). Holland et al., *Proc Natl Acad Sci U S A.* 88:7276-7280 (1991). The 5700 system used SYBR[®] green, a fluorescent dye, that only binds to double stranded DNA (Schneeberger et al., *PCR Methods Appl.* 4:234-8 (1995)), and the same forward and reverse primers as the 7700 instrument. No probe was needed. Matching primers and fluorescent probes were designed for each of the genes according to the Primer Express program (PE Biosystems, Foster City, Calif.).

TABLE 1

Primer and Probe Sequences for the Genes of Interest			
	Forward Primer	Reverse primer	Probe
Mammaglobin	TGCCATAGATGA ATTGAAGGAATG (SEQ ID NO: 48)	TGTCATATAITAATT GCATAAACACCTCA (SEQ ID NO: 49)	TCTTAACCAAACGG ATGAAAACCTTGAGC AATG (SEQ ID NO: 50)
B305D-C form	AAAGCAGATGGT GGTITGAGGIT (SEQ ID NO: 39)	CCTGAGACCAAATG GCITTCITC (SEQ ID NO: 40)	ATTCCATGCCGGCT GCTTCITCTG (SEQ ID NO: 41)
B311D	CCGCTTCTGACAA	CCTATAAGATGTT	CCCCCTCCCTCAGGG
	CACTAGAGATC (SEQ ID NO: 63)	ATGTACCAAAATG	TATGGCCC (SEQ ID NO: 65)
B726P	TCTGGTTTCTCA	AAGT (SEQ ID NO: 64)	CAACCCACGTGACA
	TTCTTTATTCAATT	TGCCAAGGAGCGGA	AACACTGGAATTAC
	TATT (SEQ ID NO: 42)	TTATCT (SEQ ID NO: 43)	AGG (SEQ ID NO: 44)
B533S	CCCTTTCTCACCC ACACACTGT (SEQ ID NO: 66)	TGCATTCTCTCATAT GTGGAAGCT (SEQ ID NO: 67)	CCGGGCTCAGGC ATATACTATCTAC TGTCTG (SEQ ID NO: 68)
GABA π	AAGCCTCAGAGT CCTTCCAGTATG (SEQ ID NO: 36)	AAATATAAGTGAAG AAAAAAATAGTAG	AATCCATTGTATCT TAGAACCGAGGGAA
	AT (SEQ ID NO: 72)	AT (SEQ ID NO: 72)	TTTGTTTAGA (SEQ ID NO: 38)
B511S	GACATTCCAGTTT TACCCAAATGG (SEQ ID NO: 69)	TGCAGAAAGACTCAA GCTGAITCC (SEQ ID NO: 70)	TCTCAGGGACACAC TCTACCATTCGGGA (SEQ ID NO: 71)

Example 6

Quantitative Real-time PCR Analysis

[0202] This example discloses the use of quantitative Real-time PCR to confirm the microarray identification polynucleotide that are at least two-fold overexpressed in breast tumor tissue samples as compared to normal breast tissue samples.

[0203] The tumor- and/or tissue-specificity of the polynucleotides identified by the microarray analyses disclosed herein in Example 5, were confirmed by quantitative PCR analyses. Breast metastases, breast tumors, benign breast disorders and normal breast tissue along with other normal tissues and tumors were tested in quantitative (Real time) PCR. This was performed either on the ABI 7700 Prism or on a GeneAmp[®] 5700 sequence detection system (PE Biosystems, Foster City, Calif.). The 7700 system uses a forward and a reverse primer in combination with a specific probe designed to anneal to sequence between the forward and reverse primer. This probe was conjugated at the 5'end with a fluorescent reporter dye and a quencher dye at the

[0204] [0204] The concentrations used in the quantitative PCR for the forward primers for mammaglobin, GABA π , B305D C form, B311D, B511S, B533S and B726P were 900, 900, 300, 900, 900, 300 and 300nM respectively. For the reverse primers they were 300, 900, 900, 900, 300, 900 and 900 nM respectively. Primers and probes so produced were used in the universal thermal cycling program in real-time PCR. They were titrated to determine the optimal concentrations using a checkerboard approach. A pool of cDNA from target tumors was used in this optimization process. The reaction was performed in 25 μ l volumes. The final probe concentration in all cases was 160 nM. dATP, dCTP and dGTP were at 0.2 mM and dUTP at 0.4 mM. AmpliTaq gold and Amperase UNG (PE Biosystems, Foster City, Calif.) were used at 0.625 units and 0.25 units per reaction. MgCl₂ was at a final concentration of 5 mM. Trace amounts of glycerol, gelatin and Tween 20 (Sigma Chem Co, St Louis, Mo.) were added to stabilize the reaction. Each reaction contained 2 μ l of diluted template. The cDNA from RT reactions prepared as above was diluted 1:10 for the gene of interest and 1:100 for P-Actin. Primers and probes for β -Actin (PE Biosystems, Foster City, Calif.) were used in a similar manner to quantitate the presence of β -actin in the samples. In the case of

the SYBR® green assay, the reaction mix (25 μ l) included 2.5 μ l of SYBR green buffer, 2 μ l of cDNA template and 2.5 μ l each of the forward and reverse primers for the gene of interest. This mix also contained 3 mM MgCl₂, 0.25units of AmpErase UNG, 0.625 units of AmpliTaq gold, 0.08% glycerol, 0.05% gelatin, 0.0001% Tween 20 and 1 mM dNTP mix. In both formats, 40 cycles of amplification were performed.

[0205] In order to quantitate the amount of specific cDNA (and hence initial mRNA) in the sample, a standard curve was generated for each run using the plasmid containing the gene of interest. Standard curves were generated using the Ct values determined in the real-time PCR which were related to the initial cDNA concentration used in the assay. Standard dilutions ranging from 20-2 \times 10⁶ copies of the gene of interest were used for this purpose. In addition, a standard curve was generated for the housekeeping gene β -actin ranging from 200 fg-200 pg to enable normalization to a constant amount of β -Actin. This allowed the evaluation of the over-expression levels seen with each of the genes.

of distant metastases derived from breast cancers all three genes reacted with 14/21 metastases and presented similar profiles. All three genes also exhibited increasing levels of expression as a function of pathological stage of the tumor, as shown for B533S in **FIG. 2**.

[0207] Mammaglobin is a homologue of a rabbit uteroglobin and the rat steroid binding protein subunit C3 and is a low molecular weight protein that is highly glycosylated. In contrast to its homologs, mammaglobin has been reported to be breast specific and over-expression has been described in breast tumor biopsies (23%) and primary and metastatic breast tumors (~75%) with reports of the detection of mammaglobin mRNA expression in 91% of lymph nodes from metastatic breast cancer patients. However, more rigorous analysis of mammaglobin gene expression by microarray and quantitative PCR as described above (panels (a) and (b) and a panel of other tumors and normal tissues and additional breast tumors), showed expression at significant levels in skin and salivary gland with much lower levels in esophagus and trachea, as shown in Table 2 below.

TABLE 2

Normalized Distribution of Mammaglobin and B511S mRNA in Various Tissues

Tissue	Mean Copies Mammaglobin /ng β -Actin ± SD	PCR Positive	Mean Copies B511S /ng β -Actin ± SD	PCR Positive	PCR Positive (Mammaglobin/ B511S)
Breast Tumors	1233.88 ± 3612 .74	31/42	1800.40 ± 3893.24	33/42	38/42
Breast tumor	1912.54 ± 4625 .85	14/24	3329.50 ± 10820.71	14/24	17/24
Metastases					
Benign Breast disorders	121.87 ± 78.63	3/3	524.66 ± 609.43	2/3	3/3
Normal breast	114.19 ± 94.40	11/11	517.64 ± 376.83	8/9	11/11
Breast reduction	231.50 ± 276.6 8	2/3	482.54 ± 680.28	½	2/3
Other tumors	0.13 ± 0.65	1/39	24.17 ± 36.00	5/23	
Salivary gland	435.65 ± 705.1 1	2/3	45766.61 ± 44342.43	3/3	
Skin	415.74 ± 376.1 4	7/9	7039.05 ± 7774.24	9/9	
Esophagus	4.45 ± 3.86	2/3	1.02 ± 0.14	0/3	
Bronchia	0.16	0/1	84.44 ± 53.31	2/2	
Other normal tissues	0.33 ± 1.07	0/85	5.49 ± 10.65	3/75	

[0206] The genes B311D, B533S and B726P were evaluated in quantitative PCR as described above on two different panels consisting of: (a) breast tumor, breast normal and normal tissues; and (b) breast tumor metastases (primarily lymph nodes), using the primers and probes shown above in Table 1. The data for panel (a) is shown in **FIG. 1** for all three genes. The three genes showed identical breast tissue expression profiles. However, the relative level of gene expression was very different in each case. B311D in general was expressed at lower levels than B533S and both less than B726P, but all three were restricted to breast tissue. The quantitative PCR thus confirmed there was a differential expression between normal breast tissue and breast tumors for all three genes, and that approximately 50% of breast tumors over-expressed these genes. When tested on a panel

[0208] The breast-specific gene B511S, while having a different profile of reactivity on breast tumors and normal breast tissue to mammaglobin, reacted with the same subset of normal tissues as mammaglobin. B511S by PSORT analysis is indicated to have an ORF of 90 aa and to be a secreted protein as is the case for mammaglobin. B511S has no evidence of a transmembrane domain but may harbor a cleavable signal sequence. Mammaglobin detected 14/24 of distant metastatic breast tumors, 31/42 breast tumors and exhibited ten-fold over-expression in tumors and metastases as compared to normal breast tissue. There was at least 300-fold over-expression in normal breast tissue versus other negative normal tissues and tumors tested, which were essentially negative for mammaglobin expression. B511S

detected 33/42 breast tumors and 14/24 distant metastases, while a combination of B511S with mammaglobin would be predicted to detect 38/42 breast tumors and 17/24 metastatic lesions (Table 2 above). The quantitative level of expression of B511S and mammaglobin were also in similar ranges, in concordance with the microarray profiles observed for these two genes. Other genes that were additive with mammaglobin are shown in Table 3.

TABLE 3

mRNA Complementation of Mammaglobin with Other Genes						
Mammaglobin Positive	Mammaglobin Negative					
	B305D	GABA α	B726P	B305D + GABA α	B305D + GABA α + B726P	
Breast Metastases	13/21	2/8	5/8	3/8	7/8	8/8
Breast tumors	18/25	3/7	4/7	5/7	7/7	7/7

[0209] B305D was shown to be highly over-expressed in breast tumors, prostate tumors, normal prostate tissue and testis compared to normal tissues, including normal breast tissue. Different splice variants of B305D have been identified with form A and C being the most abundant but all tested have similar tissue profiles in quantitative PCR. The A and C forms contain ORF's of 320 and 385 aa, respectively. B305D is predicted by PSORT to be a Type II membrane protein that comprises a series of ankyrin repeats. A known gene shown to be complementary with B305D, in breast tumors, was GABA α . This gene is a member of the GABA A receptor family and encodes a protein that has 30-40% amino acid homology with other family members, and has been shown by Northern blot analysis to be over-expressed in lung, thymus and prostate at low levels and highly over-expressed in uterus. Its expression in breast tissue has not been previously described. This is in contrast to other GABA A receptors that have appreciable expression in neuronal tissues. Tissue expression profiling of this gene showed it to be over-expressed in breast tumors in an inverse relationship to the B305D gene (Table 3). GABA α detected 15/25 tumors and 6/21 metastases including 4 tumors and 5 metastases missed by mammaglobin. In contrast, B305D detected 13/25 breast tumors and 8/21 metastases, again including 3 tumors and 2 metastases missed by mammaglobin. A combination of just B305D and the GABA α would be predicted to identify 22/25 breast tumors and 14/21 metastases. The combination of B305D and GABA α with mammaglobin in detecting breast metastases is shown in Table 3 above and **FIGS. 3A and 3B**. This combination detected 20/21 of the breast metastases as well as 25/25 breast tumors that were evaluated on the same panels for all three genes. The one breast metastasis that was negative for these three genes was strongly positive for B726P (**FIGS. 3A and 3B**).

[0210] To evaluate the presence of circulating tumor cells, an immunocapture (cell capture) method was employed to

first enrich for epithelial cells prior to RT-PCR analysis. Immunomagnetic polystyrene beads coated with specific monoclonal antibodies to two glycoproteins on the surface of epithelial cells were used for this purpose. Such an enrichment procedure increased the sensitivity of detection (~100 fold) as compared to direct isolation of poly A $^+$ RNA, as shown in Table 4.

TABLE 4

Extraction of Mammaglobin Positive Cells (MB415) Spiked into Whole Blood and Detection by Real-time PCR		
MB415 cells/ml Blood	Epithelial cell extraction (Poly A $^+$ RNA)	Direct Extraction (Poly A $^+$ RNA)
100000	54303.2	58527.1
10000	45761.9	925.9
1000	15421.2	61.6
100	368.0	5.1
10	282.0	1.1
1	110.2	0
0	0	0

[0211] Mammaglobin-positive cells (MB415) were spiked into whole blood at various concentrations and then extracted using either epithelial cell enrichment or direct isolation from blood. Using enrichment procedures, mammaglobin mRNA was found to be detectable at much lower levels than when direct isolation was used. Whole blood samples from patients with metastatic breast cancer were subsequently treated with the immunomagnetic beads. Poly A $^+$ RNA was then isolated, cDNA prepared and run in quantitative PCR using two gene specific primers (Table 1) and a fluorescent probe (TaqmanTM). As observed in breast cancer tissues, complementation was also seen in the detection of circulating tumor cells derived from breast cancers. Again, mammaglobin PCR detected circulating tumor cells in a high percentage of blood samples, albeit at low levels, from metastatic breast cancer patients (20/32) when compared to the normal blood samples (Table 5) but several of the other genes tested to date further increased this detection rate. This included B726P, B305D, B311D, B533S and GABA α . The detection level of mammaglobin in blood samples from metastatic breast cancer patients is higher than described previously (62 vs. 49%), despite testing smaller blood volumes, probably because of the use of epithelial marker-specific enrichment in our study. A combination of all the genes tested indicate that 27/32 samples were positive by one or more of these genes.

TABLE 5

Gene Complementation in Epithelial Cells Isolated from Blood of Normal Individuals and Metastatic Breast Cancer Patients

Sample ID	Mammaglobin	B305D	B311D	B533S	B726P	GABA π	Combo
2	+	-	-	+	-	-	+
3	+	-	-	+	-	-	+
5	+	+	-	-	+	-	+
6	+	-	-	+	+	-	+
8	-	-	+	-	-	-	+
9	+	+	+	-	+	-	+
10	+	-	+	-	+	-	+
11	-	-	-	-	-	-	-
12	-	+	+	-	-	-	+
13	-	-	-	+	-	-	+
15	-	-	-	-	-	-	-
18	+	-	-	-	-	-	+
19	+	-	-	-	-	+	+
21	+	-	-	-	-	-	+
22	-	-	-	-	-	-	-
23	+	-	-	-	-	-	+
24	+	-	-	-	-	-	+
25	-	+	-	-	-	-	+
26	-	-	-	-	-	-	-
29	+	-	+	+	+	-	+
31	+	-	-	+	-	-	+
32	-	-	-	-	-	±	±
33	-	-	-	-	+	-	+
34	+	-	-	-	-	+	+
35	+	-	-	-	+	-	+
36	-	-	-	-	-	+	+
37	+	-	-	+	-	-	+
38	-	-	-	-	-	-	-
40	+	-	-	-	-	-	+
41	+	-	-	+	-	-	+
42	+	-	-	-	-	-	+
43	-	-	-	-	-	+	+
Donor 104	-	-	-	-	-	+	+
Donor 348	-	-	-	-	-	Nd	-
Donor 392	-	-	-	-	-	Nd	-
Donor 408	-	-	-	-	-	Nd	-
Donor 244	-	-	-	-	-	-	-
Donor 355	-	-	-	-	-	-	-
Donor 264	-	-	-	-	-	-	-
Donor 232	-	-	-	-	-	Nd	-
Donor 12	-	-	-	-	-	-	-
Donor 415	-	-	-	-	-	Nd	-
Donor 35	-	-	-	-	-	-	-
Donor 415	-	-	-	-	-	Nd	-
Donor 35	-	-	-	-	-	-	-
Sensitivity	20/32	4/32	7/32	9/32	7/32	4/32	27/32

[0212] In further studies, mammaglobin, GABA π , B305D (C form) and B726P specific primers and specific Taqman probes were employed in different combinations to analyze their combined mRNA expression profile in breast metastases (B. met) and breast tumor (B. tumors) samples using real-time PCR. The forward and reverse primers and probes employed for mammaglobin, B305D (C form) and

B726P are shown in Table 1. The forward primer and probe employed for GABA π are shown in Table 1, with the reverse primer being as follows: TTCAAAATATAAGTGAA-GAAAAAAATTAGTAGATCAA (SEQ ID NO:51). As shown below in Table 6, a combination of mammaglobin, GABA π , B305D (C form) and B726P was found to detect 22/22 breast tumor samples, with an increase in expression being seen in 5 samples (indicated by ++).

TABLE 6

Real-time PCR Detection of Tumor Samples using Different Primer Combinations

Tumor sample	Mammaglobin	Mammaglobin + GABA	Mammaglobin + GABA + B305D	Mammaglobin + GABA + B305D + B726P
B. Met 316A		+	+	+
B. Met 317A	+	+	+	+

TABLE 6-continued

Real-time PCR Detection of Tumor Samples using Different Primer Combinations				
Tumor sample	Mammaglobin	Mammaglobin + GABA	Mammaglobin + GABA + B305D	Mammaglobin + GABA + B305D + B726P
B. Met 318A		+	+	+-
B. Met 595A	+	+	+	+
B. Met 611A	+	+	+	+
B. Met 612A	+	+	+	+
B. Met 614A		+	+	+
B. Met 616A		+	+	+
B. Met 618A	+	+	+	+
B. Met 620A	+	+	+	+
B. Met 621A	+	+	+	+
B. Met 624A	+	+	+	+
B. Met 625A			+	+
B. Met 627A	+		+	+
B. Met 629A		+	+	+
B. Met 631A	+	+	+	+
B. Tumor 154A	+	+	+	++
B. Tumor 155A	+	+	+	++
B. Tumor 81D			+	++
B. Tumor 209A	-		+	+
B. Tumor 208A		+	+	++
B. Tumor 10A	-	+	+	+

[0213] The increase of message signals by the addition of specific primers was further demonstrated in a one plate experiment employing the four tumor samples B. met 316A, B. met 317A, B. tumor 81D and B. tumor 209A.

[0214] Expression of a combination of mammaglobin, GABA π , B305D (C form) and B726P in a panel of breast tumor and normal tissue samples was also detected using real-time PCR with a SYBR Green detection system instead of the Taqman probe approach. The results obtained using this system are shown in FIG. 7.

Example 7

Quantitative PCR in Peripheral Blood of Breast Cancer Patients

[0215] The known genes evaluated in this study were mammaglobin and 7 aminobutyrate type A receptor π sub-unit (GABA π). In order to identify novel genes which are over-expressed in breast cancer we have used an improved version of the differential display RT-PCR (DDPCR) technique (Liang et al., *Science* 257:967-971 (1993); Mou et al., *Biochem Biophys Res Commun.* 199:564-569 (1994)); cDNA library extraction methods (Hara et al., *Blood* 84:189-199 (1994)) and PCR subtraction (Diatchenko et al., *Proc Natl Acad Sci USA*, 93:6025-6030 (1996); Yang et al., *Nucleic Acids Res.* 27:1517-23 (1999)).

[0216] Differential display resulted in the recovery of two cDNA fragments designated as B305D and B311D (Houghton et al., *Cancer Res.* 40: Abstract #217, 32-33, (1999). B511S and B533S are two cDNA fragments isolated using cDNA library subtraction approach (manuscript in preparation) while the B726P cDNA fragment was derived from PCR subtraction (Jiang et al., *Proceedings of the Amer Assoc Cancer Res.* 40:Abstract #216, 32 (1999); Xu et al., *Proceedings of the Amer Assoc Cancer Res.* 40:Abstract #2115, 319 (1999); and Molesh et al., *Proceedings of Amer Assoc Cancer Res.* 41:Abstract #4330, 681 (2000).

[0217] Three of the novel genes, B311D, B533S and B726P, showed identical breast tissue expression profile by

quantitative PCR analysis. These genes were evaluated in quantitative PCR on two different panels consisting of (a) breast tumor, breast normal and normal tissues and (b) panel of breast tumor metastases (primarily lymph nodes). Primers and probes used are shown in Table 1. The data for panel (a) is shown in FIG. 2 for all three genes. Overall, the expression profiles are comparable and are in the same rank order, however, the levels of expression are considerably different. B311D in general was expressed at lower levels than B533S and both less than B726P but all three were restricted to breast tissue. All three sequences were used to search against the Genbank database. Both B311D and B533S sequences contain different repetitive sequences and an ORF has not been identified for either. B726P is a novel gene, with mRNA splicing yielding several different putative ORF's.

[0218] The quantitative PCR confirmed there was a differential mRNA expression between normal breast tissue and breast tumors, with approximately 50% of breast tumors overexpressed these genes. When tested on a panel of distant metastases derived from breast cancers all three genes reacted with 14/21 metastases and presented similar profiles (data not shown). Interestingly, when tested on a prostate cancer panel, all three genes identified the same 3/24 prostate tumors but at much lower expression levels than in breast. This group of genes exhibited increasing levels of expression as a function of pathological stage of the tumor as shown for B533S.

[0219] More rigorous analysis of mammaglobin gene expression by microarray, and quantitative PCR showed expression at significant levels in skin and salivary gland and much lower levels in esophagus and trachea. B511S had a slightly different profile of reactivity on breast tumors and normal breast tissue when compared to mammaglobin, yet reacted with a similar subset of normal tissues as mammaglobin. Mammaglobin detected 14/24 of distant metastatic breast tumors, 31/42 breast tumors and exhibited ten-fold over-expression in tumors and metastases as compared to normal breast tissue. There was at least 300-fold over-expression of mammaglobin in normal breast tissue versus

other negative normal tissues and tumors tested. B511S detected 33/42 breast tumors and 14/24 distant metastases. A combination of B511S with mammaglobin would be predicted to detect 38/42 breast tumors and 17/24 metastatic lesions. The quantitative level of expression of B511S and mammaglobin were also in similar ranges, in concordance with the microarray profiles observed for these two genes.

[0220] Certain genes complemented mammaglobin's expression profile, i.e. were shown to express in tumors that mammaglobin did not. B305D was highly over-expressed in breast tumors, prostate tumors, normal prostate tissue and testis compared to normal tissues including normal breast tissue. Different splice variants of B305D were identified with the forms A and C being the most abundant. All forms tested had similar tissue profiles in quantitative PCR. The A and C forms contain ORF's of 320 and 385 aa, respectively. A known gene shown to be complementary with B305D, in breast tumors, was GABA π . This tissue expression profile is in contrast to other GABA α receptors that typically have appreciable expression in neuronal tissues. An additional observation was that tissue expression profiling of this gene showed it to be over-expressed in breast tumors in an inverse relationship to the B305D gene (Table 3). GABA π detected 15/25 tumors and 6/21 metastases including 4 tumors and 5 metastases missed by mammaglobin. In contrast, B305D detected 13/25 breast tumors and 8/21 metastases again including 3 tumors and 2 metastases missed by mammaglobin. A combination of just B305D and the GABA π would be predicted to identify 22/25 breast tumors and 14/21 metastases. This combination detected 20/21 of the breast metastases as well as 25/25 breast tumors that were evaluated on the same panels for all three genes. The one breast metastasis that was negative for these three genes was strongly positive for B726P.

[0221] The use of microarray analysis followed by quantitative PCR provided a methodology to accurately determine the expression of breast cancer genes both in breast tissues (tumor and normal) as well as in normal tissues and to assess their diagnostic and therapeutic potential. Five novel genes and two known genes were evaluated using these techniques. Three of these genes B311D, B533S and B726P exhibited concordant mRNA expression and collectively the data is consistent with coordinated expression of these three loci at the level of transcription control. All three genes showed differential expression in breast tumors versus normal breast tissue and the level of overexpression appeared related to the pathological stage of the tumor. In the case of mammaglobin, expression was found in other tissues apart from breast tissue. Expression was seen in skin, salivary gland and to a much lesser degree in trachea.

[0222] Expression of GABA π in breast tumors was also a novel observation. While the expression of several genes complemented that seen with mammaglobin, two genes in particular, B305D and GABA π added to the diagnostic sensitivity of mammaglobin detection. A combination of these three genes detected 45/46 (97.8%) breast tumors and metastases evaluated. Inclusion of B726P enabled the detection of all 25 of the breast tumors and 21 distant metastases.

Example 8

Enrichment of Circulating Breast Cancer Cells by Immunocapture

[0223] This example discloses the enhanced sensitivity achieved by use of the immunocapture cell capture methodology for enrichment of circulating breast cancer cells.

[0224] To evaluate the presence of circulating tumor cells an immunocapture method was adopted to first enrich for epithelial cells prior to RT-PCR analysis. Epithelial cells were enriched from blood samples with an immunomagnetic bead separation method (Dynal A.S, Oslo, Norway) utilizing magnetic beads coated with monoclonal antibodies specific for glycopolypeptide antigens on the surface of human epithelial cells. (Exemplary suitable cell-surface antigens are described, for example, in Momburg, F. et al., *Cancer Res.*, 41:2883-91 (1997); Naume, B. et al., *Journal of Hemotherapy*, 6:103-113 (1997); Naume, B. et al., *Int J Cancer*, 78:556-60 (1998); Martin, V. M. et al., *Exp Hematol.*, 26:252-64 (1998); Hildebrandt, M. et al., *Exp Hematol.*, 25:57-65 (1997); Eaton, M. C. et al., *Biotechniques* 22:100-5 (1997); Brandt, B. et al., *Clin Exp Metastases* 14:399-408 (1996), each of which are incorporated herein by reference in their entirety. Cells isolated this way were lysed and the magnetic beads removed. The lysate was then processed for poly A $^+$ mRNA isolation using magnetic beads (Dynabeads) coated with Oligo (dT) 25. After washing the beads in the kit buffer bead/polyA $^+$ RNA samples were finally suspended in 10 mM Tris HCl pH 8 and subjected to reverse transcription. The RNA was then subjected to Real time PCR using gene specific primers and probes with reaction conditions as outlined herein above. β -Actin content was also determined and used for normalization. Samples with gene of interest copies/ng β -actin greater than the mean of the normal samples+3 standard deviations were considered positive. Real time PCR on blood samples was performed exclusively using the TaqmanTM procedure but extending to 50 cycles.

[0225] Mammaglobin mRNA using enrichment procedures was found to be detectable at much lower levels than when direct isolation was used. Whole blood samples from patients with metastatic breast cancer were subsequently treated with the immunomagnetic beads, polyA $^+$ RNA was then isolated, cDNA made and run in quantitative PCR using two gene specific primers to mammaglobin and a fluorescent probe (TaqmanTM). As observed in breast cancer tissues, complementation was also seen in the detection of circulating tumor cells derived from breast cancers. Again, mammaglobin PCR detected circulating tumor cells in a high percentage of bloods, albeit at low levels, from metastatic breast cancer (20/32) when compared to the normal blood samples. Several of the other genes tested to date could further increase this detection rate; this includes B726P, B305D, B311D, B533S and GABA π . A combination of all the genes tested indicates that 27/32 samples were positive by one or more of these genes.

Example 9

Multiplex Detection of Breast Tumors

[0226] Additional Multiplex Real-time PCR assays were established in order to simultaneously detect the expression

of four breast cancer-specific genes: LipophilinB, Gaba (B899P), B305D-C and B726P. In contrast to detection approaches relying on expression analysis of single breast cancer-specific genes, this Multiplex assay was able to detect all breast tumor samples tested.

[0227] This Multiplex assay was designed to detect LipophilinB expression instead of Mammaglobin. Due to their similar expression profiles, LipophilinB can replace Mammaglobin in this Multiplex PCR assay for breast cancer detection. The assay was carried out as follows: LipophilinB, B899P (Gaba), B305D, and B726P specific primers, and specific Taqman probes, were used to analyze their combined mRNA expression profile in breast tumors. The primers and probes are shown below:

[0228] LipophilinB: Forward Primer (SEQ ID NO:33): 5' TGCCCCTCCGGAAAGCT. Reverse Primer (SEQ ID NO:34): 5' CGTTTCTGAAGGGACATCTGATC. Probe (SEQ ID NO:35) (FAM-5'-3'-TAMRA): TTGCAGC-CAAGTTAGGAGTGAAGAGATGCA.

[0229] GABA (B899P): Forward Primer (SEQ ID NO:36): 5' AAGCCTCAGAGTCCTCCAGTATG. Reverse Primer (SEQ ID NO:37): 5' TTCAAATATAAGT-GAAGAAAAAATTAGTAGATCAA. Probe (SEQ ID NO:38) (FAM-5'-3'-TAMRA): AATCCATTGTATCTT-GAACCGAGGGATTGTTAGA.

[0230] B305D (C form): Forward Primer (SEQ ID NO:39): 5' AAAGCAGATGGTGGTTGAGGTT. Reverse Primer (SEQ ID NO:40): 5' CCTGAGACCAAATGGCT-TCTTC. Probe (SEQ ID NO:41) (FAM-5'-3'-TAMRA): ATTCCATGCCGGCTGCTTCTTCTG.

[0231] B726P: Forward Primer (SEQ ID NO:42): 5' TCTGGTTTCTCAATTCTTATTCAATTATT. Reverse Primer (SEQ ID NO:43): 5' TGCCAAGGAGCGGAT-TATCT. Probe (SEQ ID NO:44) (FAM-5'-3'-TAMRA): CAACCACGTGACAAACACTGGAATTACAGG.

[0232] Actin: Forward Primer (SEQ ID NO:45): 5' ACTG-GAACGGTGAAAGGTGACA. Reverse Primer (SEQ ID NO 46): 5' CGGCCACATTGTGAACCTTG. Probe (SEQ ID NO:47): (FAM-5'-3'-TAMRA): CAGTCGGTTGGAGC-GAGCATCCC.

[0233] The assay conditions were:

[0234] Taqman protocol (7700 Perkin Elmer):

[0235] In 25 μ l final volume: 1x Buffer A, 5 mM MgCl, 0.2 mM dCTP, 0.2 mM dATP, 0.4 mM dUTP, 0.2 mM dGTP, 0.01 U/ μ l AmpErase UNG, 0.025 μ l TaqGold, 8% (v/v) Glycerol, 0.05% (v/v) Gelatin, 0.01% (v/v) Tween20, 4 pmol of each gene specific Taqman probe (LipophilinB+Gaba+B305D+B726P), 100 nM of B726P-F+B726P-R, 300 nM of Gaba-R, and 50 nM of LipophilinB-F+LipophilinB-R+B305D-R+Gaba-R, template cDNA (originating from 0.02 μ g polyA+RNA).

[0236] LipophilinB expression was detected in 14 out of 27 breast tumor samples. However, the Multiplex assay for LipophilinB, B899P, B305D-C and B726P detected an expression signal in 27 out of 27 tumors with the detection level above 10 mRNA copies/1000 pg actin in the majority of samples and above 100 mRNA copies/1000 pg actin in 5 out of the 27 samples tested (FIG. 8).

Example 10

Multiplex Detection Optimization

[0237] The Multiplex Real-time PCR assay described above was used to detect the expression of Mammaglobin (or LipophilinB), Gaba (B899P), B305D-C and B726P simultaneously. According to this Example, assay conditions and primer sequences were optimized to achieve parallel amplification of four PCR products with different lengths. Positive samples of this assay can be further characterized by gel electrophoresis and the expressed gene(s) of interest can be determined according to the detected amplicon size(s).

[0238] Mammaglobin (or LipophilinB), Gaba (B899P), B305D and B726P specific primers and specific Taqman probes were used to simultaneously detect their expression. The primers and probes used in this example are shown below.

[0239] Mammaglobin: Forward Primer (SEQ ID NO:48): 5' TGCCATAGATGAAITGAAGGAATG. Reverse Primer (SEQ ID NO:49): 5' TGTCAATATAAAITGCATAAA-CACCTCA. Probe (SEQ ID NO:50): (FAM-5'-3'-TAMRA): TCTTAACCAAACGGATGAAACTCTGAGCAATG.

[0240] GABA (B899P): Forward Primer (SEQ ID NO:36): 5' AAGCCTCAGAGTCCTCCAGTATG. Reverse Primer (SEQ ID NO:51): 5' ATCATTGAAAAT-TCAAATATAAGTGAAG. Probe (SEQ ID NO:38) (FAM-5'-3'-TAMRA): AATCCATTGTATCTTAGAACCGGAGG-GATTTGTTAGA.

[0241] B305D (C form): Forward Primer (SEQ ID NO:39): 5' AAAGCAGATGGTGGTTGAGGTT. Reverse Primer (SEQ ID NO:40): 5' CCTGAGACCAAATGGCT-TCTTC. Probe (SEQ ID NO:41): (FAM-5'-3'-TAMRA): ATTCCATGCCGGCTGCTTCTTCTG.

[0242] B726P: Forward Primer (SEQ ID NO:52): 5' GTAGTTGTGCATTGAAATAATTATCATTAT. Reverse Primer (SEQ ID NO:43): 5' TGCCAAGGAGCGGAT-TATCT. Probe (SEQ ID NO:44) (FAM-5'-3'-TAMRA): CAACCACGTGACAAACACTGGAATTACAGG.

[0243] Primer locations and assay conditions were optimized to achieve parallel amplification of four PCR products with different sizes. The assay conditions were:

[0244] Tagman protocol (7700 Perkin Elmer):

[0245] In 25 μ l final volume: 1x Buffer A, 5 mM MgCl, 0.2 mM dCTP, 0.2 mM dATP, 0.4 mM dUTP, 0.2 mM dGTP, 0.01 U/ μ l AmpErase UNG, 0.0375 U/ μ l TaqGold, 8% (v/v) Glycerol, 0.05% (v/v) Gelatin, 0.01% (v/v) Tween20, 4 pmol of each gene specific Taqman probe (Mammaglobin+Gaba+B305D+B726P), 300 nM of Gaba-R+Gaba-F, 100 nM of Mammaglobin-F+R; B726P-F+R, and 50 nM of B305D-F+R template cDNA (originating from 0.02 (μ g polyA+RNA).

[0246] PCR protocol:

[0247] 50° for 2': x 1, 95° for 10': X 1, and 95° for 15"/60° for 1'/68° for 1': x 50.

[0248] Since each primer set in the multiplex assay results in a band of unique length, expression signals of the four genes of interest can be measured individually by agarose

gel analysis (see, FIG. 9), or the combined expression signal of all four genes can be measured in real-time on an ABI 7700 Prism sequence detection system (PE Biosystems, Foster City, Calif.). The expression of LipophilinB can also be detected instead of Mammaglobin. Although specific primers have been described herein, different primer sequences, different primer or probe labeling and different detection systems could be used to perform this multiplex

[0250] Mammaglobin, Gaba (B899P), B305D and B726P specific primers and specific Taqman probes were used to simultaneously detect their expression (Table 7). Primer locations were optimized (Intron-Exon border spanning) to exclusively detect cDNA and to exclude genomic DNA from amplification. The identity of the expressed gene(s) was determined by gel electrophoresis.

TABLE 7

Intron-Exon border Spanning Primer and Probe Sequences for Breast Tumor Multiples Assay			
Gene	Forward Primer	Reverse Primer	Taqman probe (FAM-5'- 3'TAMRA)
Mammaglobin	tgccatagatgtaaatgg atg (SEQ ID NO:48)	tgtcatatataattgtcataaaacacct ca (SEQ ID NO:49)	tcttaaccaaacggatgaaactctgagca atg (SEQ ID NO:50)
B899P	aagcctcagacttcctccagta tg (SEQ ID NO:36)	tccaaatataatgtaaagaaaaaaatta gtatgtatcaa (SEQ ID NO:37)	aatccattgtatcttagaaccggaggatt gtt (SEQ ID NO:62)
B305D	aaagcagatgggtggatgt t (SEQ ID NO:39)	cctgagaccaaattgcgttcc (SEQ ID NO:40)	attccatgcggctgtcttcgt (SEQ ID NO:41)
B726P	tctggtttcatttttttttttttt tatt (SEQ ID NO:42)	tgcggaggcggattatct (SEQ ID NO:43)	caaccacgtgacaaacactggattaca gg (SEQ ID NO:44)
Actin	actggaaacggtgaaagggtgac a (SEQ ID NO:45)	cggccacatgtgaaacttg (SEQ ID NO:46)	cagtcgggtggagcggatcccc (SEQ ID NO:47)
B899P-INT	caattttggggagaaacccg (SEQ ID NO:53)	gtctgcggaggatataatgg (SEQ ID NO:54)	catttcagagatgaaatggactacaca (SEQ ID NO:55)
B305D-INT	tctgtataaaggccgtacaatg (SEQ ID NO:56)	tcaacacttgctttttgtc (SEQ ID NO:57)	atcaaaaacaagcatggctcacacca ct (SEQ ID NO:58)
B726P-INT	gcaagtgtccaaatgtacagagg (SEQ ID NO:59)	atatacgtcgtatcacact (SEQ ID NO:60)	tcccatcagaatccaaacaagaggaa tg (SEQ ID NO:61)

assay. For example, a second fluorogenic reporter dye could be incorporated for parallel detection of a reference gene by real-time PCR. Or, for example a SYBR Green detection system could be used instead of the Taqman probe approach.

Example 11

Design and use of Genomic DNA-excluding, Intron-exon Border Spanning Primer Rairs for Breast Cancer Multiplex Assay

[0249] The Multiplex Real-time PCR assay described herein can detect the expression of Mammaglobin, Gaba (B899P), B305D-C and B726P simultaneously. The combined expression levels of these genes is measured in real-time on an ABI 7700 Prism sequence detection system (PE Biosystems, Foster City, Calif.). Individually expressed genes can also be identified due to different amplicon sizes via gel electrophoresis. In order to use this assay with samples derived from non-DNase treated RNAs (e.g. lymph node cDNA) and to avoid DNase-treatment for small RNA-samples (e.g. from blood specimens, tumor and lymph node aspirates), intron-spanning primer pairs have been designed to exclude the amplification of genomic DNA and therefore to eliminate nonspecific and false positive signals. False positive signal is caused by genomic DNA contamination in cDNA specimens. The optimized Multiplex assay described herein excludes the amplification of genomic DNA and allows specific detection of target gene expression without the necessity of prior DNase treatment of RNA samples. Moreover the genomic match and the location of the Intron-Exon border could be verified with these primer sets.

[0251] Primer locations and assay conditions were optimized to achieve parallel amplification of the four PCR products. The assay conditions were as follows:

[0252] Tagman Protocol (7700 Perkin Elmer)

[0253] In 25 μ l final volume: 1x Buffer A, 5 mM MgCl₂, 0.2 mM dCTP, 0.2 mM dATP, 0.4 mM dUTP, 0.2 mM dGTP, 0.01 U/AmpErase UNG, 8% (v/v) Glycerol, 0.05% (v/v) Gelatin, 0.01% (v/v) Tween20, 4 pmol of each gene specific Taqman probe (Mammaglobin+B899P-INT+B305D-INT+B726P-INT), 300 nM of B305D-INT-F; B899P-INT-F, 100 nM of Mammaglobin-F+R; B726P-INT-F +R, 50 nM of B899P-INT-R; B305D-INT-R, template cDNA (originating from 0.02 μ g polyA+ RNA).

[0254] PCR Cycling Conditions

[0255] 1 cycle at 50° C. for 2 minutes, 1 cycle at 95° C. for 10 minutes, 50 cycles of 95° C. for 1 minute and 68° C. for 1 minute.

[0256] FIG. 10 shows a comparison of the multiplex assay using intron-exon border spanning primers (bottom panel) and the multiplex assay using non-optimized primers (top panel), to detect breast cancer cells in a panel of lymph node tissues. This experiment shows that reduction in background resulting from genomic DNA contamination in samples is achieved using the intron-exon spanning primers of the present invention.

Example 12

Multiplex Detection of Metastasized Breast Tumor Cells in Sentinel Lymph Node Biopsy Samples

[0257] Lymph node staging is important for determining appropriate adjuvant hormone and chemotherapy. In con-

trast to conventional axillary dissection a less invasive approach for staging of minimal residual disease is sentinel lymph node biopsy. Sentinel lymph node biopsy (SLNB) has the potential to improve detection of metastases and to provide prognostic values to lead to therapy with minimal morbidity associated with complete lymph node dissection. SLNB implements mapping of the one or two lymph nodes which primarily drain the tumor and therefore are most likely to harbor metastatic disease (the sentinel nodes). Routine pathological analysis of lymph nodes result in a high false-negative rate: one-third of women with pathologically negative lymph nodes develop recurrent disease [Bland: *The Breast*: Saunders 1991]. A more sensitive detection technique for tumor cells would be RT-PCR but its application is limited by lack of a single specific markers. The multimarker assay described above increases the likelihood of cancer detection across the population without producing false-positive results from normal lymph nodes.

[0258] As mentioned above, lymphatic afferents from a primary tumor drain into a single node, the sentinel lymph node, before drainage into the regional lymphatic basin occurs. Sentinel lymph nodes are located with dyes and/or radiolabelled colloid injected in the primary lesion site and sentinel lymph node biopsy allows pathological examination for micrometastatic deposits, staging of the axilla and therefore can avoid unnecessary axillary dissection. Nodal micrometastases can be located with staining (haematoxylin or eosin) or immunohistochemical analysis for cytokeratin proteins. Immunocytochemical staining techniques can produce frequent false-negative results by missing small metastatic foci due to inadequate sectioning of the node. Immunohistochemistry can result in false-positive results due to illegitimate expression of cytokeratins (reticulum cells) or in false-negative results when using the antibody Ber-Ep4 which corresponding antigen is not expressed on all tumor cells.

[0259] The multiplex assay described herein could provide a more sensitive detection tool for positive sentinel lymph nodes. Moreover the detection of breast cancer cells in bone marrow samples, peripheral blood and small needle aspiration samples is desirable for diagnosis and prognosis in breast cancer patients.

[0260] Twenty-two metastatic lymph node samples, in addition to 15 samples also previously analyzed and shown in **FIG. 3A**, were analyzed using the intron-exon border spanning multiplex PCR assay described herein. The results from this analysis are summarized in Table 8. Twenty-seven primary tumors were also analyzed and the results shown in Table 9. Twenty normal lymph node samples tested using this assay were all negative.

TABLE 8

Multiples Real-time PCR Analysis of 37 Metastatic Lymph Nodes

breast metastatic lymph node samples	Mamma-globin	B305D	B899P	B726P	Multiplex
B. Met 317A	++	+	+	+++	
B. Met 318A			++	+++	
B. Met 595A	+		+	+++	
B. Met 611A	+	+	+++	++	
B. Met 612A	++	++	+	++	
B. Met 614A		++	++	+++	

TABLE 8-continued

Multiples Real-time PCR Analysis of 37 Metastatic Lymph Nodes

breast metastatic lymph node samples	Mamma-globin	B305D	B899P	B726P	Multiplex
B. Met 616A				+	++
B. Met 618A	+++	+			+++
B. Met 620A	++	++		++	+++
B. Met 621A	+	---		+	+++
B. Met 624A				-+	+++
B. Met 625A		---		++	+
B. Met 627A		+		+	+
B. Met 629A	++				+++
B. Met 631A	+		++		+
1255	+++	++		++	++
1257	+++	+	+	+	++
769	+++			+	++
1258	+	+	+		+
1259		++	++		+++
1250	+++	+		+	+++
1726	+++	+		+	+++
786	---	+	+		+++
281-LI-r	+++				+++
289-L2	---	+			++
366-S	+				+
374-S+	+++	++			+++
376-S	++			+	++
381-S	+	+			+
383-Sx	+++	++			+++
496-M	+++	++			+++
591-SI-A	+	+			+
652-I		+	++		+++
772	-				+
777	+	+		++	++
778	+++				+++
779	+		++		++

[0261]

TABLE 9

Multiplex Real-time PCR Analysis of 27 Primary Breast Tumors

breast primary tumor samples	Mamma-globin	B305D	B899P	B726P	Multiplex
T443	+	++		+++	+++
T457		+	+		++
T395			++		++
T10A	-	+++		+++	+++
T446		+		++	++
T11C	+		+++		+++
T23B	+	++			+++
T207A		++			+
T437	+	+		++	+++
T391	+	++		+++	+++
T392	+	+			++
TS76	+	++			+++
T483	++	+			+++
T81G	+	+	++	++	+++
T430	+		++		++
T465	+	+		+	++
TS80				+	+
T469	+			+	---
T467	+			++	+++
T439		+			+
T387	++		+	+	++
T318				+	++
T154A				+	+
T387A	+++		+	+	+++
T155A	+		++	+	+
T209A			++		++
T208A		+		+	++

[0262] From the foregoing it will be appreciated that, although specific embodiments of the invention have been described herein for purposes of illustration, various modi-

fications may be made without deviating from the spirit and scope of the invention. Accordingly, the invention is not limited except as by the appended claims.

 SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 77

<210> SEQ ID NO 1
 <211> LENGTH: 1851
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapien

<400> SEQUENCE: 1

tcatcaccat	tgccagcagc	ggcaccgtta	gtcaggtttt	ctggaaatcc	cacatgagta	60
cttccgtgtt	cttcattctt	cttcaatagc	cataaaatctt	ctagctctgg	ctggctgttt	120
tcacttcctt	taagcctttg	tgactcttcc	tctgatgtca	gctttaagtc	ttgttctgga	180
ttgctgtttt	cagaagagat	ttttaacatc	tgtttttctt	tgtatgcaga	aagtaactgg	240
caaattacat	gatgtatgt	agaaacagca	tactctctgg	ccgtctttcc	agatcttgag	300
aagatacatc	aacatttgc	tcaagtagag	ggctgactat	acttgctgtat	ccacaacata	360
cagcaagtat	gagagcagtt	cttccatatac	tatccagcgc	atttaaattc	gctttttct	420
tgattaaaaa	tttaccact	tgcgtttttt	gctcatgtat	accaagtagc	agtgggtgtga	480
ggccatgctt	gttttttgc	tgcataatcag	caccgtataa	gagcagtgct	ttggccattt	540
atttatcttc	attgttagaca	gcatagtgt	gagtggattt	tccatactca	tctggaatat	600
ttggatcagt	gccatgttcc	agcaacattt	acgcacattc	atcttcctgg	cattgtacgg	660
cctttgtcag	agctgtcctc	ttttgttgtt	caaggacatt	aagttgacat	cgtctgtcca	720
gcacagttt	tactacttct	gaattccat	tggcagaggc	cagatgtaga	gcagtcctct	780
tttgcttgc	cctcttggtc	acatccgtgt	ccctgagcat	gacgtatgaga	tcctttctgg	840
ggactttacc	ccaccaggca	gctctgtgga	gcttgcctag	atcttctcca	tggacgttgt	900
acctgggatc	catgaaggcg	ctgtcatgt	agtctccca	agcgcaccacg	ttgctcttgc	960
cgcctccctg	cagcagggga	agcagtggca	gcaccacttgc	cacctcttgc	tcccaagcgt	1020
cttcacagag	gagtcgttgt	ggtctccaga	agtgcacccacg	ttgctcttgc	cgctccccct	1080
gtccatccag	ggaggaagaa	atgcaggaaa	tgaaagatgc	atgcacgtat	gtataactcct	1140
cagccatcaa	acttctggac	agcaggtcac	ttccagcaag	gtggagaaaag	ctgtccaccc	1200
acagaggatg	agatccagaa	accacaatat	ccattcacaa	acaaacactt	ttcagccaga	1260
cacaggtact	gaaatcatgt	catctgcggc	aacatggtgg	aacctaccca	atcacacatc	1320
aagagatgaa	gacactgcag	tatatctgca	caacgtaata	ctcttcatcc	ataacaaaat	1380
aatataat	tcctctggag	ccatatggat	gaactatgaa	ggaagaactc	cccgaaagaag	1440
ccagtcgcag	agaagccaca	ctgaagctct	gtcctcagcc	atcagcgcca	cgacacaggar	1500
tgtgtttctt	ccccagtgtat	gcagcctcaa	gttatccga	agctgccgca	gcacacggtg	1560
gctcctgaga	aacacccacg	ctcttccgg	ctaacacagg	caagtcataa	aatgtataa	1620
tcacataaaac	agaattaaaa	gcaaagtac	ataagcatct	caacagacac	agaaaaggca	1680
tttgacaaaa	tccagcatcc	ttgtat	ttgtgcagtt	ctcagaggaa	atgcttctaa	1740
ctttcccca	tttagtatta	tgtggctgt	gggcttgc	taggtggttt	ttattacttt	1800

-continued

```

aaggatatgtc ccttctatgc ctgtttgct gagggttttta attctcgatc c          1851

<210> SEQ ID NO 2
<211> LENGTH: 329
<212> TYPE: PRT
<213> ORGANISM: Homo sapien

<400> SEQUENCE: 2

Met Asp Ile Val Val Ser Gly Ser His Pro Leu Trp Val Asp Ser Phe
 1           5           10          15

Leu His Leu Ala Gly Ser Asp Leu Leu Ser Arg Ser Leu Met Ala Glu
20          25           30

Glu Tyr Thr Ile Val His Ala Ser Phe Ile Ser Cys Ile Ser Ser Ser
35           40           45

Leu Asp Gly Gln Gly Glu Arg Gln Glu Gln Arg Gly His Phe Trp Arg
50           55           60

Pro Gln Arg Leu Leu Cys Glu Asp Ala Trp Glu Gln Glu Val Gln Val
65           70           75           80

Val Leu Pro Leu Leu Pro Leu Leu Gln Gly Ser Gly Lys Ser Asn Val
85           90           95

Val Ala Trp Gly Asp Tyr Asp Asp Ser Ala Phe Met Asp Pro Arg Tyr
100          105          110

His Val His Gly Glu Asp Leu Asp Lys Leu His Arg Ala Ala Trp Trp
115          120          125

Gly Lys Val Pro Arg Lys Asp Leu Ile Val Met Leu Arg Asp Thr Asp
130          135          140

Val Asn Lys Arg Asp Lys Gln Lys Arg Thr Ala Leu His Leu Ala Ser
145          150          155          160

Ala Asn Gly Asn Ser Glu Val Val Lys Leu Val Asp Arg Arg Cys
165          170          175

Gln Leu Asn Val Leu Asp Asn Lys Lys Arg Thr Ala Leu Thr Lys Ala
180          185          190

Val Gln Cys Gln Glu Asp Glu Cys Ala Leu Met Leu Leu Glu His Gly
195          200          205

Thr Asp Pro Asn Ile Pro Asp Glu Tyr Gly Asn Thr Thr Leu His Tyr
210          215          220

Ala Val Tyr Asn Glu Asp Lys Leu Met Ala Lys Ala Leu Leu Tyr
225          230          235          240

Gly Ala Asp Ile Glu Ser Lys Asn Lys His Gly Leu Thr Pro Leu Leu
245          250          255

Leu Gly Ile His Glu Gln Lys Gln Val Val Lys Phe Leu Ile Lys
260          265          270

Lys Lys Ala Asn Leu Asn Ala Leu Asp Arg Tyr Gly Arg Thr Ala Leu
275          280          285

Ile Leu Ala Val Cys Cys Gly Ser Ala Ser Ile Val Ser Pro Leu Leu
290          295          300

Glu Gln Asn Val Asp Val Ser Ser Gln Asp Leu Glu Arg Arg Pro Glu
305          310          315          320

Ser Met Leu Phe Leu Val Ile Ile Met
325

```

<210> SEQ ID NO 3

-continued

<211> LENGTH: 1852

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 3

ggcacgagaa ttaaaaccct cagcaaaaca ggcatagaag ggacataacct taaagtaata	60
aaaaccacct atgacaagcc cacagccaaac ataataactaa atggggaaaa gttagaagca	120
tttcctctga gaactgcaac aataaaataca aggatgctgg attttgtcaa atgcctttc	180
tgtgtctgtt gagatgctta tgtgactttg cttttaattc tggatgtgtt attatcacat	240
ttattgactt gcctgtgtta gaccggaaga gctgggggtgt ttctcaggag ccaccgtgtg	300
ctgcggcagc ttccggataa cttgaggctg catcaactgg gaagaaacac aytccgtc	360
gtggcgctga tggctgagga cagagttca gtgtggcttc tctgcgactg gcttc	420
ggagtttttc cttcatagtt catccatatg gctccagagg aaaattatata tattttgtta	480
tggatgaaga gtattacgtt gtgcagatata actgcagtgt ctccatctct tgatgtgtga	540
ttgggtaggt tccaccatgt tgccgcagat gacatgattt cagtttgcgtt gtctggctga	600
aaagtgtttt tttgtgaatg gatattgtgg tttctggatc tcatcctctg tgggtggaca	660
gctttctcca ctttgctgaa agtgcactgc tgcacatgtt gttatgtgtt gaggagttata	720
ccatcgatcg tgcacatcttc atttcctgca tttcttcctc cttggatgaa cagggggagc	780
ggcaagagca acgtgggcac ttctggagac cacaacgact cctctgtgaa gacgcttggg	840
agcaagaggt gcaagtgggtg ctgcactgc ttccctgtt gcaggggggag cggcaagagc	900
aacgtggatcg cttggggaga ctacgatgac agcgcctca tggatcccag gtaccacgtc	960
catggagaag atctggacaa gctccacaga gctgcctggt ggggtaaagt ccccaaaaaa	1020
gatctcatcg tcatgctcg ggacacggat gtgaacaaga gggacaagca aaagaggact	1080
gctctacatc tggctctgc caatggaaat tcagaagtagt taaaactcgt gctggacaga	1140
cgtatgtcaac ttaatgtcct tgacaacaaa aagaggacag ctctgacaaa ggcgtacaa	1200
tgccaggaag atgaatgtgc gttatgttg ctggaaacatg gcactgatcc aaatattcca	1260
gatgagttatg gaaataaccac tctacactat gctgtctaca atgaagataa attaatggcc	1320
aaagcactgc tcttatacgg tgctgatatc gaatcaaaaa acaagcatgg cctcacacca	1380
ctgctacttg gtatacatga gcaaaaaacag caagtggtaa aatttttat caagaaaaaa	1440
gcgaatttaa atgcgctgaa tagatatgga agaactgctc tcatactcgt tggatgtgt	1500
ggatcagcaa gtatagtcg ccctctactt gagcaaaatg ttgatgtatc ttctcaagat	1560
ctggaaagac ggccagagag tatgtgtttt ctatgtatca tcattgtat tggcgttac	1620
tttctgacta caaagaaaaa cagatgttaa aaatctcttc tgaaaacagc aatccagaac	1680
aagacttaaa gctgacatca gaggaaaggt cacaagggt taaaggaagt gaaaacagcc	1740
agccagagct agaagattta tggctattga agaagaatga agaacacgaa agtactcatg	1800
tgggattccc agaaaacctg actaacggtg ccgctgtgg caatggatgat ga	1852

<210> SEQ ID NO 4

<211> LENGTH: 292

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 4

-continued

Met His Leu Ser Phe Pro Ala Phe Leu Pro Pro Trp Met Asp Arg Gly
 5 10 15

Ser Gly Lys Ser Asn Val Gly Thr Ser Gly Asp His Asn Asp Ser Ser
 20 25 30

Val Lys Thr Leu Gly Ser Lys Arg Cys Lys Trp Cys Cys His Cys Phe
 35 40 45

Pro Cys Cys Arg Gly Ser Gly Lys Ser Asn Val Val Ala Trp Gly Asp
 50 55 60

Tyr Asp Asp Ser Ala Phe Met Asp Pro Arg Tyr His Val His Gly Glu
 65 70 75 80

Asp Leu Asp Lys Leu His Arg Ala Ala Trp Trp Gly Lys Val Pro Arg
 85 90 95

Lys Asp Leu Ile Val Met Leu Arg Asp Thr Asp Val Asn Lys Arg Asp
 100 105 110

Lys Gln Lys Arg Thr Ala Leu His Leu Ala Ser Ala Asn Gly Asn Ser
 115 120 125

Glu Val Val Lys Leu Val Leu Asp Arg Arg Cys Gln Leu Asn Val Leu
 130 135 140

Asp Asn Lys Lys Arg Thr Ala Leu Thr Lys Ala Val Gln Cys Gln Glu
 145 150 155 160

Asp Glu Cys Ala Leu Met Leu Leu Glu His Gly Thr Asp Pro Asn Ile
 165 170 175

Pro Asp Glu Tyr Gly Asn Thr Thr Leu His Tyr Ala Val Tyr Asn Glu
 180 185 190

Asp Lys Leu Met Ala Lys Ala Leu Leu Tyr Gly Ala Asp Ile Glu
 195 200 205

Ser Lys Asn Lys His Gly Leu Thr Pro Leu Leu Leu Gly Ile His Glu
 210 215 220

Gln Lys Gln Gln Val Val Lys Phe Leu Ile Lys Lys Lys Ala Asn Leu
 225 230 235 240

Asn Ala Leu Asp Arg Tyr Gly Arg Thr Ala Leu Ile Leu Ala Val Cys
 245 250 255

Cys Gly Ser Ala Ser Ile Val Ser Pro Leu Leu Glu Gln Asn Val Asp
 260 265 270

Val Ser Ser Gln Asp Leu Glu Arg Arg Pro Glu Ser Met Leu Phe Leu
 275 280 285

Val Ile Ile Met
 290

<210> SEQ_ID NO 5

<211> LENGTH: 1155

<212> TYPE: DNA

<213> ORGANISM: Homo sapien

<400> SEQUENCE: 5

atgggttgtg aggttatttc catgccggct gcctttctg tgaagaagcc atttggctc 60
 aggagcaaga tggcaagtg gtgctgccgt tgcttccctt gctgcaggaa gagcggcaag 120
 agcaacgtgg gcacttctgg agaccacgac gactctgcta tgaagacact caggagcaag 180
 atggcaagt ggtgcccca ctgcttcccc tgctgcagg ggagtggcaa gagcaacgtg 240
 ggcgttctg gagaccacga cgactctgct atgaagacac tcaggaacaa gatggcaag 300
 tggtgctgcc actgcttccc ctgctgcagg gggagcggca agagcaaggt gggcgcttgg 360

-continued

ggagactacg atgacagtgc cttcatggag cccaggtacc acgtccgtgg agaagatctg	420
gacaagctcc acagagctgc ctgggtgggt aaagtccccca gaaaggatct catcgcatg	480
ctcaggaca ctgacgtgaa caagaaggac aagcaaaaga ggactgctct acatctggcc	540
tctgccaatg ggaattcaga agtagtaaaa ctcctgctgg acagacgtatg tcaacttaat	600
gtccttgaca aaaaaaagag gacagctctg ataaaggccg tacaatgccca ggaagatgaa	660
tgtgcgttaa tggcgttggaa acatggcact gatccaaata ttccagatga gtatggaaat	720
accactctgc actacgctat ctataatgaa gataaattaa tggccaaagc actgcttta	780
tatggtgctg atatcgaatc aaaaaacaag catggcctca caccactgtt acttgggtgt	840
catgagcaaa aacagcaagt cgtgaaattt ttaatcaaga aaaaagcgaa tttaaatgca	900
ctggatagat atgaaaggac tgctctcata cttgctgtat gttgtggatc agcaagtata	960
gtcagccttc tacttgagca aaatattgtat gtatcttcata aagatctatc tggacagacg	1020
gccagagagt atgctgttcc tagtcatcat catgttaattt gccagttact ttctgactac	1080
aaagaaaaac agatgctaaa aatctttctt gaaaacagca atccagaaaaa tgtctcaaga	1140
accagaaata aataa	1155

<210> SEQ ID NO 6
<211> LENGTH: 2000
<212> TYPE: DNA
<213> ORGANISM: Homo sapien

<400> SEQUENCE: 6

atgggtgttgc aggttggattc catgccccgt gccttttttgc tgaagaagcc atttgggttca	60
aggagcaaga tggcaagtg gtgtgtccgt tgcttccccct gctgcaggga gagcggcaag	120
agcaacgtgg gcacttctgg agaccacgac gactctgttca tgaagacact caggagcaag	180
atgggcaagt ggtgccgcca ctgtttccccct tgctgcagggg ggagtggcaa gagcaacgtg	240
ggcgcttctg gagaccacga cgactctgttca atgaagacac tcaggaacaaat gatgggcaag	300
tgggtgtgcc actgcttccc ctgtgcagg gggagcggca agagcaaggt gggcgcttgg	360
ggagactacg atgacagtgc cttcatggag cccaggtacc acgtccgtgg agaagatctg	420
gacaagctcc acagagctgc ctgggtgggt aaagtccccca gaaaggatct catcgcatg	480
ctcaggaca ctgacgtgaa caagaaggac aagcaaaaga ggactgctct acatctggcc	540
tctgccaatg ggaattcaga agtagtaaaa ctcctgctgg acagacgtatg tcaacttaat	600
gtccttgaca aaaaaaagag gacagctctg ataaaggccg tacaatgccca ggaagatgaa	660
tgtgcgttaa tggcgttggaa acatggcact gatccaaata ttccagatga gtatggaaat	720
accactctgc actacgctat ctataatgaa gataaattaa tggccaaagc actgcttta	780
tatggtgctg atatcgaatc aaaaaacaag catggcctca caccactgtt acttgggtgt	840
catgagcaaa aacagcaagt cgtgaaattt ttaatcaaga aaaaagcgaa tttaaatgca	900
ctggatagat atgaaaggac tgctctcata cttgctgtat gttgtggatc agcaagtata	960
gtcagccttc tacttgagca aaatattgtat gtatcttcata aagatctatc tggacagacg	1020
gccagagagt atgctgttcc tagtcatcat catgttaattt gccagttact ttctgactac	1080
aaagaaaaac agatgctaaa aatctttctt gaaaacagca atccagaaaaa agacttaag	1140
ctgacatcg aggaagagtc acaaagggttc aaaggcagtg aaaatagcca gccagagaaaa	1200

-continued

atgtctcaag aaccagaaaat aaataaggat ggtgatagag aggttgaaga agaaatgaag	1260
aagcatgaaa gtaataatgt gggattacta gaaaacctga ctaatggtgt cactgctggc	1320
aatgggtata atggattaaat tcctcaaagg aagagcagaa cacctgaaaa tcagcaattt	1380
cctgacaacg aaagtgaaga gtatcacaga atttgcgaat tagttctga ctacaaagaa	1440
aaacagatgc caaaaactc ttctgaaaac agcaacccag aacaagactt aaagctgaca	1500
tcagaggaag agtcacaaaag gcttgaggc agtggaaatg gccagccaga gctagaaaat	1560
tttatggcta tcgaagaaaat gaagaagcac ggaagtaactc atgtcggtt cccagaaaac	1620
ctgactaatg gtgccactgc tggcaatggt gatgatggat taattccctcc aaggaagagc	1680
agaacacctg aaagccagca atttctgac actgagaatg aagagtatca cagtgacgaa	1740
caaaaatgata ctcagaagca attttgcgaa gaacagaaca ctggaatattt acacgatgag	1800
attctgattc atgaagaaaa gcagatagaa gtgggtgaaa aaatgaattt tgagcttct	1860
ctttagttgtt agaaaagaaaa agacatcttgc catgaaaata gtacgttgcc ggaagaaattt	1920
gccatgctaa gactggagct agacacaatg aaacatcaga gccagctaaa aaaaaaaaaaa	1980
aaaaaaaaaa aaaaaaaaaaa	2000

<210> SEQ ID NO 7
 <211> LENGTH: 2040
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapien

<400> SEQUENCE: 7

atgggtgttgc aggttggattc catgccccgt gccttttttgc tgaagaagcc atttgggttgc	60
aggagcaaga tggcaagtg gtgtgtccgt tgcttccctt gctgcaggga gagcggcaag	120
agcaacgtgg gcacttctgg agaccacgac gactctgcta tgaagacact caggagcaag	180
atggggcaagt ggtggcccca ctgtttcccc tgctgcagggg ggagtggcaa gagcaacgtg	240
ggcgcttctg gagaccacga cgactctgct atgaagacac tcaggaacaa gatgggcaag	300
tgggtgtgcc actgtttccc ctgtgcagg gggagcggca agagcaaggt gggcgcttgg	360
ggagactacg atgacagtgc cttcatggag cccaggtacc acgtccgtgg agaagatctg	420
gacaagctcc acagagctgc ctgggggggt aaagtccccca gaaaggatct catcgatcg	480
ctcagggaca ctgacgtgaa caagaaggac aagcaaaaaga ggactgctct acatctggcc	540
tctgccaatg ggaattcaga agtagtaaaa ctcctgtgg acagacgtatg tcaacttaat	600
gtccctgaca acaaaaagag gacagctctg ataaaggccg tacaatgca ggaagatgaa	660
tgtgcgttaa tgggtgtggaa acatggact gatccaaata ttccagatga gtatggaaat	720
accactctgc actacgctat ctataatgaa gataaaattaa tggccaaagc actgcttta	780
tatgggtctg atatcgaatc aaaaaacaag catggccctca caccactgtt acttgggtta	840
catgagcaaa aacagcaagt cgtgaaattt ttaatcaaga aaaaagcgaa tttaaatgca	900
ctggatagat atgaaaggac tgctctcata cttgtgttat gttgtggatc agcaagtata	960
gtcagccttc tacttgagca aaatattgtat gtatcttctc aagatctatc tggacagacg	1020
gccagagagt atgctgtttc tagtcatcat catgttaattt gccagttact ttctgactac	1080
aaagaaaaac agatgctaaa aatcttttctt gaaaacagca atccagaaca agacttaag	1140
ctgacatcg aggaagagtc acaaagggttc aaaggcagtg aaaatagcca gccagagaaa	1200

-continued

atgtctcaag aaccagaaaat aaataaggat ggtgatagag aggttgaaga agaaatgaag	1260
aagcatgaaa gtaataatgt gggattacta gaaaacctga ctaatggtgt cactgctggc	1320
aatgggtata atggattaaat tcctcaaagg aagagcagaa cacctgaaaa tcagcaattt	1380
cctgacaacg aaagtgaaga gtatcacaga atttgcgaat tagttctga ctacaaagaa	1440
aaacagatgc caaaatactc ttctgaaaac agcaacccag aacaagactt aaagctgaca	1500
tcagaggaag agtcacaaaag gcttgaggc agtggaaatg gccagccaga gaaaagatct	1560
caagaaccag aaataaaataa ggatggtgat agagagctag aaaattttat ggctatcgaa	1620
gaaaatgaaga agcacggaag tactcatgtc ggattccag aaaacctgac taatgggcc	1680
actgctggca atggtgatga tggattaaatt cctccaagga agagcagaac acctgaaagc	1740
cagcaatttc ctgacactga gaatgaagag tatcacagtg acgaacaaaa tgatactcag	1800
aagcaatttt gtgaagaaca gaacactgga atattacacg atgagattct gattcatgaa	1860
gaaaagcaga tagaagtggt tgaaaaatg aattctgagc tttctcttag ttgtaaagaa	1920
gaaaaagaca tcttgcataa aaatagtacg ttgcggaaag aaattgccat gctaagactg	1980
gagctagaca caatgaaaca tcagagccag ctaaaaaaaaaaaaaaa aaaaaaaaaa	2040

<210> SEQ ID NO 8

<211> LENGTH: 384

<212> TYPE: PRT

<213> ORGANISM: Homo sapien

<400> SEQUENCE: 8

Met Val Val Glu Val Asp Ser Met Pro Ala Ala Ser Ser Val Lys Lys			
1	5	10	15
10	15		

Pro Phe Gly Leu Arg Ser Lys Met Gly Lys Trp Cys Cys Arg Cys Phe			
20	25	30	
30			

Pro Cys Cys Arg Glu Ser Gly Lys Ser Asn Val Gly Thr Ser Gly Asp			
35	40	45	
45			

His Asp Asp Ser Ala Met Lys Thr Leu Arg Ser Lys Met Gly Lys Trp			
50	55	60	
60			

Cys Arg His Cys Phe Pro Cys Cys Arg Gly Ser Gly Lys Ser Asn Val			
65	70	75	80
75	80		

Gly Ala Ser Gly Asp His Asp Asp Ser Ala Met Lys Thr Leu Arg Asn			
85	90	95	
95			

Lys Met Gly Lys Trp Cys Cys His Cys Phe Pro Cys Cys Arg Gly Ser			
100	105	110	
110			

Gly Lys Ser Lys Val Gly Ala Trp Gly Asp Tyr Asp Asp Ser Ala Phe			
115	120	125	
125			

Met Glu Pro Arg Tyr His Val Arg Gly Glu Asp Leu Asp Lys Leu His			
130	135	140	
140			

Arg Ala Ala Trp Trp Gly Lys Val Pro Arg Lys Asp Leu Ile Val Met			
145	150	155	160
155	160		

Leu Arg Asp Thr Asp Val Asn Lys Lys Asp Lys Gln Lys Arg Thr Ala			
165	170	175	
175			

Leu His Leu Ala Ser Ala Asn Gly Asn Ser Glu Val Val Lys Leu Leu			
180	185	190	
190			

Leu Asp Arg Arg Cys Gln Leu Asn Val Leu Asp Asn Lys Lys Arg Thr			
195	200	205	
205			

-continued

Ala Leu Ile Lys Ala Val Gln Cys Gln Glu Asp Glu Cys Ala Leu Met
 210 215 220

Leu Leu Glu His Gly Thr Asp Pro Asn Ile Pro Asp Glu Tyr Gly Asn
 225 230 235 240

Thr Thr Leu His Tyr Ala Ile Tyr Asn Glu Asp Lys Leu Met Ala Lys
 245 250 255

Ala Leu Leu Leu Tyr Gly Ala Asp Ile Glu Ser Lys Asn Lys His Gly
 260 265 270

Leu Thr Pro Leu Leu Leu Gly Val His Glu Gln Lys Gln Gln Val Val
 275 280 285

Lys Phe Leu Ile Lys Lys Ala Asn Leu Asn Ala Leu Asp Arg Tyr
 290 295 300

Gly Arg Thr Ala Leu Ile Leu Ala Val Cys Cys Gly Ser Ala Ser Ile
 305 310 315 320

Val Ser Leu Leu Leu Glu Gln Asn Ile Asp Val Ser Ser Gln Asp Leu
 325 330 335

Ser Gly Gln Thr Ala Arg Glu Tyr Ala Val Ser Ser His His His Val
 340 345 350

Ile Cys Gln Leu Leu Ser Asp Tyr Lys Glu Lys Gln Met Leu Lys Ile
 355 360 365

Ser Ser Glu Asn Ser Asn Pro Glu Asn Val Ser Arg Thr Arg Asn Lys
 370 375 380

<210> SEQ_ID NO 9
 <211> LENGTH: 656
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapien

<400> SEQUENCE: 9

Met Val Val Glu Val Asp Ser Met Pro Ala Ala Ser Ser Val Lys Lys
 1 5 10 15

Pro Phe Gly Leu Arg Ser Lys Met Gly Lys Trp Cys Cys Arg Cys Phe
 20 25 30

Pro Cys Cys Arg Glu Ser Gly Lys Ser Asn Val Gly Thr Ser Gly Asp
 35 40 45

His Asp Asp Ser Ala Met Lys Thr Leu Arg Ser Lys Met Gly Lys Trp
 50 55 60

Cys Arg His Cys Phe Pro Cys Cys Arg Gly Ser Gly Lys Ser Asn Val
 65 70 75 80

Gly Ala Ser Gly Asp His Asp Asp Ser Ala Met Lys Thr Leu Arg Asn
 85 90 95

Lys Met Gly Lys Trp Cys Cys His Cys Phe Pro Cys Cys Arg Gly Ser
 100 105 110

Gly Lys Ser Lys Val Gly Ala Trp Gly Asp Tyr Asp Asp Ser Ala Phe
 115 120 125

Met Glu Pro Arg Tyr His Val Arg Gly Glu Asp Leu Asp Lys Leu His
 130 135 140

Arg Ala Ala Trp Trp Gly Lys Val Pro Arg Lys Asp Leu Ile Val Met
 145 150 155 160

Leu Arg Asp Thr Asp Val Asn Lys Lys Asp Lys Gln Lys Arg Thr Ala
 165 170 175

Leu His Leu Ala Ser Ala Asn Gly Asn Ser Glu Val Val Lys Leu Leu
 180 185 190

-continued

Leu Asp Arg Arg Cys Gln Leu Asn Val Leu Asp Asn Lys Lys Arg Thr
 195 200 205
 Ala Leu Ile Lys Ala Val Gln Cys Gln Glu Asp Glu Cys Ala Leu Met
 210 215 220
 Leu Leu Glu His Gly Thr Asp Pro Asn Ile Pro Asp Glu Tyr Gly Asn
 225 230 235 240
 Thr Thr Leu His Tyr Ala Ile Tyr Asn Glu Asp Lys Leu Met Ala Lys
 245 250 255
 Ala Leu Leu Tyr Gly Ala Asp Ile Glu Ser Lys Asn Lys His Gly
 260 265 270
 Leu Thr Pro Leu Leu Leu Gly Val His Glu Gln Lys Gln Gln Val Val
 275 280 285
 Lys Phe Leu Ile Lys Lys Ala Asn Leu Asn Ala Leu Asp Arg Tyr
 290 295 300
 Gly Arg Thr Ala Leu Ile Leu Ala Val Cys Cys Gly Ser Ala Ser Ile
 305 310 315 320
 Val Ser Leu Leu Glu Gln Asn Ile Asp Val Ser Ser Gln Asp Leu
 325 330 335
 Ser Gly Gln Thr Ala Arg Glu Tyr Ala Val Ser Ser His His Val
 340 345 350
 Ile Cys Gln Leu Leu Ser Asp Tyr Lys Glu Lys Gln Met Leu Lys Ile
 355 360 365
 Ser Ser Glu Asn Ser Asn Pro Glu Gln Asp Leu Lys Leu Thr Ser Glu
 370 375 380
 Glu Glu Ser Gln Arg Phe Lys Gly Ser Glu Asn Ser Gln Pro Glu Lys
 385 390 395 400
 Met Ser Gln Glu Pro Glu Ile Asn Lys Asp Gly Asp Arg Glu Val Glu
 405 410 415
 Glu Glu Met Lys Lys His Glu Ser Asn Asn Val Gly Leu Leu Glu Asn
 420 425 430
 Leu Thr Asn Gly Val Thr Ala Gly Asn Gly Asp Asn Gly Leu Ile Pro
 435 440 445
 Gln Arg Lys Ser Arg Thr Pro Glu Asn Gln Gln Phe Pro Asp Asn Glu
 450 455 460
 Ser Glu Glu Tyr His Arg Ile Cys Glu Leu Val Ser Asp Tyr Lys Glu
 465 470 475 480
 Lys Gln Met Pro Lys Tyr Ser Ser Glu Asn Ser Asn Pro Glu Gln Asp
 485 490 495
 Leu Lys Leu Thr Ser Glu Glu Ser Gln Arg Leu Glu Gly Ser Glu
 500 505 510
 Asn Gly Gln Pro Glu Leu Glu Asn Phe Met Ala Ile Glu Glu Met Lys
 515 520 525
 Lys His Gly Ser Thr His Val Gly Phe Pro Glu Asn Leu Thr Asn Gly
 530 535 540
 Ala Thr Ala Gly Asn Gly Asp Asp Gly Leu Ile Pro Pro Arg Lys Ser
 545 550 555 560
 Arg Thr Pro Glu Ser Gln Gln Phe Pro Asp Thr Glu Asn Glu Glu Tyr
 565 570 575
 His Ser Asp Glu Gln Asn Asp Thr Gln Lys Gln Phe Cys Glu Glu Gln
 580 585 590

-continued

Asn Thr Gly Ile Leu His Asp Glu Ile Leu Ile His Glu Glu Lys Gln
 595 600 605

Ile Glu Val Val Glu Lys Met Asn Ser Glu Leu Ser Leu Ser Cys Lys
 610 615 620

Lys Glu Lys Asp Ile Leu His Glu Asn Ser Thr Leu Arg Glu Glu Ile
 625 630 635 640

Ala Met Leu Arg Leu Glu Leu Asp Thr Met Lys His Gln Ser Gln Leu
 645 650 655

<210> SEQ_ID NO 10

<211> LENGTH: 671

<212> TYPE: PRT

<213> ORGANISM: Homo sapien

<400> SEQUENCE: 10

Met Val Val Glu Val Asp Ser Met Pro Ala Ala Ser Ser Val Lys Lys
 1 5 10 15

Pro Phe Gly Leu Arg Ser Lys Met Gly Lys Trp Cys Cys Arg Cys Phe
 20 25 30

Pro Cys Cys Arg Glu Ser Gly Lys Ser Asn Val Gly Thr Ser Gly Asp
 35 40 45

His Asp Asp Ser Ala Met Lys Thr Leu Arg Ser Lys Met Gly Lys Trp
 50 55 60

Cys Arg His Cys Phe Pro Cys Cys Arg Gly Ser Gly Lys Ser Asn Val
 65 70 75 80

Gly Ala Ser Gly Asp His Asp Asp Ser Ala Met Lys Thr Leu Arg Asn
 85 90 95

Lys Met Gly Lys Trp Cys Cys His Cys Phe Pro Cys Cys Arg Gly Ser
 100 105 110

Gly Lys Ser Lys Val Gly Ala Trp Gly Asp Tyr Asp Asp Ser Ala Phe
 115 120 125

Met Glu Pro Arg Tyr His Val Arg Gly Glu Asp Leu Asp Lys Leu His
 130 135 140

Arg Ala Ala Trp Trp Gly Lys Val Pro Arg Lys Asp Leu Ile Val Met
 145 150 155 160

Leu Arg Asp Thr Asp Val Asn Lys Asp Lys Gln Lys Arg Thr Ala
 165 170 175

Leu His Leu Ala Ser Ala Asn Gly Asn Ser Glu Val Val Lys Leu Leu
 180 185 190

Leu Asp Arg Arg Cys Gln Leu Asn Val Leu Asp Asn Lys Lys Arg Thr
 195 200 205

Ala Leu Ile Lys Ala Val Gln Cys Gln Glu Asp Glu Cys Ala Leu Met
 210 215 220

Leu Leu Glu His Gly Thr Asp Pro Asn Ile Pro Asp Glu Tyr Gly Asn
 225 230 235 240

Thr Thr Leu His Tyr Ala Ile Tyr Asn Glu Asp Lys Leu Met Ala Lys
 245 250 255

Ala Leu Leu Leu Tyr Gly Ala Asp Ile Glu Ser Lys Asn Lys His Gly
 260 265 270

Leu Thr Pro Leu Leu Leu Gly Val His Glu Gln Lys Gln Val Val
 275 280 285

Lys Phe Leu Ile Lys Lys Ala Asn Leu Asn Ala Leu Asp Arg Tyr
 290 295 300

-continued

Gly Arg Thr Ala Leu Ile Leu Ala Val Cys Cys Gly Ser Ala Ser Ile
 305 310 315 320
 Val Ser Leu Leu Leu Glu Gln Asn Ile Asp Val Ser Ser Gln Asp Leu
 325 330 335
 Ser Gly Gln Thr Ala Arg Glu Tyr Ala Val Ser Ser His His His Val
 340 345 350
 Ile Cys Gln Leu Leu Ser Asp Tyr Lys Glu Lys Gln Met Leu Lys Ile
 355 360 365
 Ser Ser Glu Asn Ser Asn Pro Glu Gln Asp Leu Lys Leu Thr Ser Glu
 370 375 380
 Glu Glu Ser Gln Arg Phe Lys Gly Ser Glu Asn Ser Gln Pro Glu Lys
 385 390 395 400
 Met Ser Gln Glu Pro Glu Ile Asn Lys Asp Gly Asp Arg Glu Val Glu
 405 410 415
 Glu Glu Met Lys Lys His Glu Ser Asn Asn Val Gly Leu Leu Glu Asn
 420 425 430
 Leu Thr Asn Gly Val Thr Ala Gly Asn Gly Asp Asn Gly Leu Ile Pro
 435 440 445
 Gln Arg Lys Ser Arg Thr Pro Glu Asn Gln Gln Phe Pro Asp Asn Glu
 450 455 460
 Ser Glu Glu Tyr His Arg Ile Cys Glu Leu Val Ser Asp Tyr Lys Glu
 465 470 475 480
 Lys Gln Met Pro Lys Tyr Ser Ser Glu Asn Ser Asn Pro Glu Gln Asp
 485 490 495
 Leu Lys Leu Thr Ser Glu Glu Ser Gln Arg Leu Glu Gly Ser Glu
 500 505 510
 Asn Gly Gln Pro Glu Lys Arg Ser Gln Glu Pro Glu Ile Asn Lys Asp
 515 520 525
 Gly Asp Arg Glu Leu Glu Asn Phe Met Ala Ile Glu Glu Met Lys Lys
 530 535 540
 His Gly Ser Thr His Val Gly Phe Pro Glu Asn Leu Thr Asn Gly Ala
 545 550 555 560
 Thr Ala Gly Asn Gly Asp Asp Gly Leu Ile Pro Pro Arg Lys Ser Arg
 565 570 575
 Thr Pro Glu Ser Gln Gln Phe Pro Asp Thr Glu Asn Glu Glu Tyr His
 580 585 590
 Ser Asp Glu Gln Asn Asp Thr Gln Lys Gln Phe Cys Glu Glu Gln Asn
 595 600 605
 Thr Gly Ile Leu His Asp Glu Ile Leu Ile His Glu Glu Lys Gln Ile
 610 615 620
 Glu Val Val Glu Lys Met Asn Ser Glu Leu Ser Leu Ser Cys Lys Lys
 625 630 635 640
 Glu Lys Asp Ile Leu His Glu Asn Ser Thr Leu Arg Glu Glu Ile Ala
 645 650 655
 Met Leu Arg Leu Glu Leu Asp Thr Met Lys His Gln Ser Gln Leu
 660 665 670

<210> SEQ ID NO 11

<211> LENGTH: 800

<212> TYPE: DNA

<213> ORGANISM: Homo sapien

-continued

<400> SEQUENCE: 11

atkagttcc	gcttctgaca	acactagaga	tccctccctt	ccctcagggt	atggccctcc	60
acttcatttt	tgg tacataa	catcttataa	ggacagggggt	aaaatccaa	tactaacagg	120
agaatgctta	ggactctaac	agg ttttga	aatgtgttg	gtaagggcca	ctcaatccaa	180
ttttttttgg	tcctccttgc	ggtctaggag	gacaggcaag	ggtcagatt	ttcaagaatg	240
catcagtaag	ggccactaaa	tccgaccttc	ctcggtcctc	cttgggtct	gggaggaaaa	300
ctagtgttcc	tgttgctgt	tcaacttcc	gatcagcagg	gtccaggac	360	
cactgcagg	tcttgggcag	ggggagaaac	aaaacaaacc	aaaaccatgg	gcrgtttgc	420
ctttcagatg	ggaaacactc	aggcatcaac	aggctcacct	ttgaaatgca	tcctaagcca	480
atgggacaaa	tttgacccac	aaacccttgg	aaaagaggtg	gctcattttt	tttgcactat	540
ggcttggccc	caacattctc	tctctgtatgg	ggaaaaatgg	ccacctgagg	gaagtacaga	600
ttacaatact	atcctgcagc	ttgacccctt	ctgtaagagg	gaaggcaaat	ggagtgaaat	660
accttatgtc	caagctttct	tttcattgaa	ggagaataca	ctatgcaaag	cttgcatttt	720
acatcccaca	ggaggacctc	tca gttacc	cccatatcc	agcctcccta	tagctccct	780
tcctattagt	gataagcctc					800

<210> SEQ ID NO 12

<211> LENGTH: 102

<212> TYPE: PRT

<213> ORGANISM: Homo sapien

<220> FEATURE:

<221> NAME/KEY: VARIANT

<222> LOCATION: (1)...(102)

<223> OTHER INFORMATION: Xaa = Any Amino Acid

<400> SEQUENCE: 12

Met	Gly	Xaa	Phe	Val	Phe	Gln	Met	Gly	Asn	Thr	Gln	Ala	Ser	Thr	Gly
1				5			10					15			

Ser	Pro	Leu	Lys	Cys	Ile	Leu	Ser	Gln	Trp	Asp	Lys	Phe	Asp	Pro	Gln
					20			25			30				

Thr	Leu	Glu	Lys	Glu	Val	Ala	His	Phe	Phe	Cys	Thr	Met	Ala	Trp	Pro
					35		40			45					

Gln	His	Ser	Leu	Ser	Asp	Gly	Glu	Lys	Trp	Pro	Pro	Glu	Gly	Ser	Thr
					50		55		60						

Asp	Tyr	Asn	Thr	Ile	Leu	Gln	Leu	Asp	Leu	Phe	Cys	Lys	Arg	Glu	Gly
				65		70		75		80					

Lys	Trp	Ser	Glu	Ile	Pro	Tyr	Val	Gln	Ala	Phe	Phe	Ser	Leu	Lys	Glu
				85			90		95						

Asn	Thr	Leu	Cys	Lys	Ala
		100			

<210> SEQ ID NO 13

<211> LENGTH: 1206

<212> TYPE: DNA

<213> ORGANISM: Homo sapien

<400> SEQUENCE: 13

ggcacgagga	agt tttgtgt	actgaaaaag	aaactgtcag	aagcaaaaga	aataaaatca	60
cagttagaga	accaaaaagt	taaatggaa	caagagctc	gcagtgtgag	gtttctcaca	120
ctcatgaaaa	t gaaaattat	ctcttacatg	aaaattgc	ttgaaaaag	gaaattgc	180

-continued

tgctaaaact	ggaaatagcc	acactgaaac	accaataccca	ggaaaaggaa	aataaatact	240
ttgaggacat	taagatttta	aaagaaaaga	atgctgact	tcagatgacc	ctaaaactga	300
aagaggaatc	attaactaaa	agggcatctc	aatatagtgg	gcagcttaaa	gttctgatag	360
ctgagaacac	aatgctcact	tctaaattga	aggaaaaaca	agacaaagaa	atactagagg	420
cagaaaattga	atcacaccat	cctagactgg	cttctgctgt	acaagaccat	gatcaaattg	480
tgacatcaag	aaaaagtcaa	gaacctgctt	tccacattgc	aggagatgct	tgtttgcaaa	540
gaaaaatgaa	tgttgatgt	agtagtacga	tatataacaa	tgaggtgctc	catcaaccac	600
tttctgaagc	tcaaaggaaa	tccaaaagcc	taaaaattaa	tctcaattat	gccggagatg	660
ctctaaagaga	aaatacattg	gtttcagaac	atgcacaaag	agaccaacgt	gaaacacagt	720
gtcaaatgaa	ggaagctgaa	cacatgtatc	aaaacgaaca	agataatgtg	aacaaacaca	780
ctgaacagca	ggagtctcta	gatcagaaat	tatttcaact	acaaagcaaa	aatatgtggc	840
ttcaacagca	attagttcat	gcacataaga	aagctgacaa	caaaagcaag	ataacaattg	900
atattcattt	tcttgagagg	aaaatgcaac	atcatctcct	aaaagagaaa	aatgaggaga	960
tatttaat	caataaccat	ttaaaaaacc	gtatataatca	atatgaaaaa	gagaaagcag	1020
aaacagaagt	tatataatag	tataacactg	ccaaggagcg	gattatctca	tcttcattcct	1080
gtaattccag	tgtttgtcac	gtgggtgtt	aataaatgaa	taaagaatga	gaaaaccaga	1140
agctctgata	cataatcata	atgataatta	tttcaatgca	caactacggg	tggtgctgct	1200
cgtgcc						1206

<210> SEQ ID NO 14

<211> LENGTH: 317

<212> TYPE: PRT

<213> ORGANISM: Homo sapien

<400> SEQUENCE: 14

Met	Gly	Thr	Arg	Ala	Leu	Gln	Cys	Glu	Val	Ser	His	Thr	His	Glu	Asn
1				5				10				15			

Glu	Asn	Tyr	Leu	Leu	His	Glu	Asn	Cys	Met	Leu	Lys	Lys	Glu	Ile	Ala
			20				25				30				

Met	Leu	Lys	Leu	Glu	Ile	Ala	Thr	Leu	Lys	His	Gln	Tyr	Gln	Glu	Lys
	35				40				45						

Glu	Asn	Lys	Tyr	Phe	Glu	Asp	Ile	Lys	Ile	Leu	Lys	Glu	Lys	Asn	Ala
	50				55			60							

Glu	Leu	Gln	Met	Thr	Leu	Lys	Leu	Lys	Glu	Glu	Ser	Leu	Thr	Lys	Arg
	65				70			75				80			

Ala	Ser	Gln	Tyr	Ser	Gly	Gln	Leu	Lys	Val	Leu	Ile	Ala	Glu	Asn	Thr
	85					90				95					

Met	Leu	Thr	Ser	Lys	Leu	Lys	Glu	Lys	Gln	Asp	Lys	Glu	Ile	Leu	Glu
	100				105			110							

Ala	Glu	Ile	Glu	Ser	His	His	Pro	Arg	Leu	Ala	Ser	Ala	Val	Gln	Asp
	115				120			125							

His	Asp	Gln	Ile	Val	Thr	Ser	Arg	Lys	Ser	Gln	Glu	Pro	Ala	Phe	His
	130				135			140							

Ile	Ala	Gly	Asp	Ala	Cys	Leu	Gln	Arg	Lys	Met	Asn	Val	Asp	Val	Ser
	145				150			155			160				

Ser	Thr	Ile	Tyr	Asn	Asn	Glu	Val	Leu	His	Gln	Pro	Leu	Ser	Glu	Ala
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

-continued

165	170	175
Gln Arg Lys Ser Lys Ser Leu Lys Ile Asn Leu Asn Tyr Ala Gly Asp		
180	185	190
Ala Leu Arg Glu Asn Thr Leu Val Ser Glu His Ala Gln Arg Asp Gln		
195	200	205
Arg Glu Thr Gln Cys Gln Met Lys Glu Ala Glu His Met Tyr Gln Asn		
210	215	220
Glu Gln Asp Asn Val Asn Lys His Thr Glu Gln Gln Glu Ser Leu Asp		
225	230	235
Gln Lys Leu Phe Gln Leu Gln Ser Lys Asn Met Trp Leu Gln Gln Gln		
245	250	255
Leu Val His Ala His Lys Lys Ala Asp Asn Lys Ser Lys Ile Thr Ile		
260	265	270
Asp Ile His Phe Leu Glu Arg Lys Met Gln His His Leu Leu Lys Glu		
275	280	285
Lys Asn Glu Glu Ile Phe Asn Tyr Asn Asn His Leu Lys Asn Arg Ile		
290	295	300
Tyr Gln Tyr Glu Lys Glu Lys Ala Glu Thr Glu Val Ile		
305	310	315

<210> SEQ ID NO 15

<211> LENGTH: 1665

<212> TYPE: DNA

<213> ORGANISM: Homo sapien

<400> SEQUENCE: 15

gcaaacttcc aagcagagcc tcccgagaag ccatctgcct tcgagccctgc cattgaaatg	60
caaaaagtctg ttccaaataa agccttggaa ttgaagaatg aacaaacatt gagagcagat	120
cagatgttcc cttcagaatc aaaacaaaag aaggttgaag aaaattcttg ggattctgag	180
agtctccgtg agactgttcc acagaaggat gtgtgtgtac ccaaggctac acatcaaaaa	240
gaaatggata aaataagtgg aaaatttagaa gattcaacta gcctatcaaa aatcttggat	300
acagttcatt ctgtgaaag agcaaggaa cttcaaaaag atcaactgtga acaacgtaca	360
ggaaaaatgg aacaaatgaa aaagaagttt tgtgtactga aaaagaaact gtcagaagca	420
aaagaaataa aatcacagtt agagaaccaa aaagttaaat gggacaaga gctctgcagt	480
gtgaggtttc tcacactcat gaaaatgaaa attatcttt acatgaaaat tgcatgttga	540
aaaagggaaat tgccatgcta aaactggaaa tagccacact gaaacaccaa taccaggaaa	600
aggaaaaataa atacttttag gacattaaga ttttaaaaaga aaagaatgtc gaacttcaga	660
tgaccctaaa actgaaagag gaatcattaa ctaaaaggcc atctcaatat agtgggcagc	720
ttaaagtctt gatagtcttag aacacaatgc tcacttctaa attgaaggaa aaacaagaca	780
aagaaaatact agaggcagaa attgaatcac accatcctag actggcttct gctgtacaag	840
accatgatca aattgtgaca tcaagaaaaa gtcaagaacc tgctttccac attgcaggag	900
atgcttggttt gcaaagaaaa atgaatgttg atgtgagtag tacgatatat aacaatgagg	960
tgctccatca accactttct gaagctcaa ggaaatccaa aagcctaaaa attaatctca	1020
attatgccgg agatgtctca agagaaaaata cattggtttc agaacatgc caaagagacc	1080
aacgtgaaac acagtgtcaa atgaaggaag ctgaacacat gtatcaaaac gaacaagata	1140
atgtgaacaa acacactgaa cagcaggagt ctctagatca gaaatttattt caactacaaa	1200

-continued

gcaaaaatat	gtggcttcaa	cagcaattag	ttcatgcaca	taagaaagct	gacaacaaaa	1260
gcaagataac	aattgatatt	catttcttg	agaggaaaat	gcaacatcat	ctcctaaaag	1320
agaaaaatga	ggagatattt	aattacaata	accatttaaa	aaaccgtata	tatcaatatg	1380
aaaaagagaa	agcagaaaaca	gaaaactcat	gagagacaag	cagtaagaaa	cttctttgg	1440
agaaacaaca	gaccagatct	ttactcacaa	ctcatgctag	gaggccagtc	ctagcattac	1500
cttatgttga	aaatcttacc	aatagtctgt	gtcaacagaa	tacttatttt	agaagaaaaa	1560
ttcatgattt	cttcctgaag	cctggcgac	agagcgagac	tctgtctcaa	aaaaaaaaaa	1620
aaaaaaagaa	agaaagaaaat	gcctgtgctt	acttcgcttc	ccagg		1665

<210> SEQ ID NO 16

<211> LENGTH: 179

<212> TYPE: PRT

<213> ORGANISM: Homo sapien

<400> SEQUENCE: 16

Ala	Asn	Phe	Gln	Ala	Glu	Pro	Pro	Glu	Lys	Pro	Ser	Ala	Phe	Glu	Pro
1				5			10			15					

Ala	Ile	Glu	Met	Gln	Lys	Ser	Val	Pro	Asn	Lys	Ala	Leu	Glu	Leu	Lys
	20				25					30					

Asn	Glu	Gln	Thr	Leu	Arg	Ala	Asp	Gln	Met	Phe	Pro	Ser	Glu	Ser	Lys
	35			40				45							

Gln	Lys	Lys	Val	Glu	Glu	Asn	Ser	Trp	Asp	Ser	Glu	Ser	Leu	Arg	Glu
	50			55				60							

Thr	Val	Ser	Gln	Lys	Asp	Val	Cys	Val	Pro	Lys	Ala	Thr	His	Gln	Lys
	65			70			75		80						

Glu	Met	Asp	Lys	Ile	Ser	Gly	Lys	Leu	Glu	Asp	Ser	Thr	Ser	Leu	Ser
	85				90			95							

Lys	Ile	Leu	Asp	Thr	Val	His	Ser	Cys	Glu	Arg	Ala	Arg	Glu	Leu	Gln
	100				105			110							

Lys	Asp	His	Cys	Glu	Gln	Arg	Thr	Gly	Lys	Met	Glu	Gln	Met	Lys	Lys
	115			120				125							

Lys	Phe	Cys	Val	Leu	Lys	Lys	Leu	Ser	Glu	Ala	Lys	Glu	Ile	Lys
	130			135			140							

Ser	Gln	Leu	Glu	Asn	Gln	Lys	Val	Lys	Trp	Glu	Gln	Glu	Leu	Cys	Ser
	145			150			155		160						

Val	Arg	Phe	Leu	Thr	Leu	Met	Lys	Met	Lys	Ile	Ile	Ser	Tyr	Met	Lys
	165				170		175								

Ile Ala Cys

<210> SEQ ID NO 17

<211> LENGTH: 1681

<212> TYPE: DNA

<213> ORGANISM: Homo sapien

<400> SEQUENCE: 17

gatacagtca	ttcttgtcaa	agagcaaggg	aacttcaaaa	agatcactgt	gaacaacgta	60
cagaaaaat	ggaacaaaatg	aaaaagaagt	tttgcgtact	aaaaaagaaa	ctgtcagaag	120
caaaagaat	aaaatcacag	tttagagaacc	aaaaagttaa	atgggaacaa	gagctctgca	180
gtgtgagatt	gactttaaac	caagaagaag	agaagagaag	aaatgccat	atattaaatg	240

-continued

aaaaaaattag ggaagaatta ggaagaatcg aagagcagca taggaaagag tttagaagtga	300
aacaacaact tgaacaggct ctcagaatac aagatataga attgaagagt gtagaaagta	360
atttgaatca ggtttctcac actcatgaaa atgaaaatta tctcttacat gaaaattgca	420
tgttgaaaaa gggaaattgcc atgctaaaac tggaaatagc cacactgaaa caccaatacc	480
aggaaaaggaa aaataaaatac tttgaggaca ttaagatttt aaaagaaaaag aatgctgaac	540
ttcagatgac cctaaaaactg aaagaggaat cattaactaa aagggcatct caatatagtg	600
ggcagcttaa agttctgata gctgagaaca caatgctcac ttcttaattg aaggaaaaac	660
aagacaaaaga aatacttagag gcagaaaattg aatcacacca tcctagactg gcttctgctg	720
tacaagacca tgatcaaatt gtgacatcaa gaaaaagtca agaacctgct ttccacattg	780
caggagatgc ttgttgcaa agaaaaatga atgttgatgt gagtagtacg atatataaca	840
atgaggtgct ccatcaacca ctttctgaag ctcaaaggaa atccaaaagc ctaaaaatta	900
atctcaatta tgccggagat gctctaagag aaaatacatt ggtttcagaa catgcacaaa	960
gagaccaacgc tgaaacacag tgcataatga aggaagctg acacatgtat caaaacgaac	1020
aagataatgt gaacaaacac actgaacacgc aggagtctct agatcagaaa ttatttcaac	1080
tacaaagcaa aaatatgtgg cttcaacacgc aatttttca tgcacataag aaagctgaca	1140
acaaaaagcaa gataacaatt gatatttcatt ttcttgagag gaaaatgcaa catcatctcc	1200
taaaagagaa aaatgaggag atatttaatt acaataacca tttaaaaaac cgtatataatc	1260
aatatgaaaaa agagaaagca gaaacagaaa actcatgaga gacaagcagt aagaaacttc	1320
ttttggagaa acaacagacc agatcttac tcacaactca tgcttaggagg ccagtcctag	1380
cattacccta tggaaaaaa tcttaccaat agtctgtgtc aacagaatac ttattttaga	1440
agaaaaaatttc atgatttctt cctgaaggct acagacataa aataacagtg tgaagaatta	1500
cttggtcacg aattgcataa aagctgccc ggatttccat ctaccctgga tgatggcgg	1560
gacatcatc aatccaaacca gaatctcgct ctgtcactca ggctggagtg cagtgccgc	1620
aatctcggtc cactgcaact ctgcctccca ggttcacgccc attctctggc acagcctccc	1680
g	1681

<210> SEQ ID NO 18
 <211> LENGTH: 432
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapien

<400> SEQUENCE: 18

Asp Thr Val His Ser Cys Glu Arg Ala Arg Glu Leu Gln Lys Asp His	
1	5
10	15

Cys Glu Gln Arg Thr Gly Lys Met Glu Gln Met Lys Lys Lys Phe Cys	
20	25
30	

Val Leu Lys Lys Leu Ser Glu Ala Lys Glu Ile Lys Ser Gln Leu	
35	40
45	

Glu Asn Gln Lys Val Lys Trp Glu Gln Glu Leu Cys Ser Val Arg Leu	
50	55
60	

Thr Leu Asn Gln Glu Glu Lys Arg Arg Asn Ala Asp Ile Leu Asn	
65	70
75	80

Glu Lys Ile Arg Glu Glu Leu Gly Arg Ile Glu Glu Gln His Arg Lys	
85	90
95	

-continued

Glu Leu Glu Val Lys Gln Gln Leu Glu Gln Ala Leu Arg Ile Gln Asp
 100 105 110

Ile Glu Leu Lys Ser Val Glu Ser Asn Leu Asn Gln Val Ser His Thr
 115 120 125

His Glu Asn Glu Asn Tyr Leu Leu His Glu Asn Cys Met Leu Lys Lys
 130 135 140

Glu Ile Ala Met Leu Lys Leu Glu Ile Ala Thr Leu Lys His Gln Tyr
 145 150 155 160

Gln Glu Lys Glu Asn Lys Tyr Phe Glu Asp Ile Lys Ile Leu Lys Glu
 165 170 175

Lys Asn Ala Glu Leu Gln Met Thr Leu Lys Leu Lys Glu Glu Ser Leu
 180 185 190

Thr Lys Arg Ala Ser Gln Tyr Ser Gly Gln Leu Lys Val Leu Ile Ala
 195 200 205

Glu Asn Thr Met Leu Thr Ser Lys Leu Lys Glu Lys Gln Asp Lys Glu
 210 215 220

Ile Leu Glu Ala Glu Ile Glu Ser His His Pro Arg Leu Ala Ser Ala
 225 230 235 240

Val Gln Asp His Asp Gln Ile Val Thr Ser Arg Lys Ser Gln Glu Pro
 245 250 255

Ala Phe His Ile Ala Gly Asp Ala Cys Leu Gln Arg Lys Met Asn Val
 260 265 270

Asp Val Ser Ser Thr Ile Tyr Asn Asn Glu Val Leu His Gln Pro Leu
 275 280 285

Ser Glu Ala Gln Arg Lys Ser Lys Ser Leu Lys Ile Asn Leu Asn Tyr
 290 295 300

Ala Gly Asp Ala Leu Arg Glu Asn Thr Leu Val Ser Glu His Ala Gln
 305 310 315 320

Arg Asp Gln Arg Glu Thr Gln Cys Gln Met Lys Glu Ala Glu His Met
 325 330 335

Tyr Gln Asn Glu Gln Asp Asn Val Asn Lys His Thr Glu Gln Gln Glu
 340 345 350

Ser Leu Asp Gln Lys Leu Phe Gln Leu Gln Ser Lys Asn Met Trp Leu
 355 360 365

Gln Gln Gln Leu Val His Ala His Lys Lys Ala Asp Asn Lys Ser Lys
 370 375 380

Ile Thr Ile Asp Ile His Phe Leu Glu Arg Lys Met Gln His His Leu
 385 390 395 400

Leu Lys Glu Lys Asn Glu Glu Ile Phe Asn Tyr Asn Asn His Leu Lys
 405 410 415

Asn Arg Ile Tyr Gln Tyr Glu Lys Glu Lys Ala Glu Thr Glu Asn Ser
 420 425 430

<210> SEQ ID NO 19
 <211> LENGTH: 3681
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 19

```

tccgagctga ttacagacac caaggaagat gctgtaaaaga gtcagcagcc acagccctgg 60
ctagctggcc ctgtggcat ttattagtaa agttttaatg acaaaagctt tgagtcaaca 120
cacccgtggg taattaaacct ggtcatcccc accctggaga gccatcctgc ccatgggtga 180
  
```

-continued

tcaaagaagg aacatctgca ggaacacctg atgaggctgc acccttggcg gaaagaacac	240
ctgacacagc tgaaagcttg gtggaaaaaa cacctgatga ggcgcaccc ttggtgaaa	300
gaacacactga caoggctgaa agcttggtgg aaaaaacacc tcatgaggct gcatcctgg	360
tggagggAAC atctgacaaa attcaatgtt tggagaaagc gacatctgga aagttcgaac	420
agtcaGcaga agaaacacct agggaaatta cggatcctgc aaaagaaaca tctgagaaat	480
ttacgtggcc agcaaaagga agaccttagga agatcgcatg ggagaaaaaa gaagacacac	540
ctaggaaat tatgagtccc gcaaaaagaaa catctgagaa atttacgtgg gcagcaaaag	600
gaagacctag gaagatcgca tgggagaaaa aagaaacacc tggtaaagact ggatgcgtgg	660
caagagtaac atctaataaa actaaagttt tggaaaaagg aagatctaag atgattgcat	720
gtcctacaaa agaatcatct acaaaagcaa gtgcattatca tcagaggttc ccatcagaat	780
ccaaacaaga ggaagatgaa gaatattctt gtgattctcg gagtctctt gagagttctg	840
caaagattca agtgtgtata cctgagtcta tatataaaaa agtaatggag ataaatagag	900
aagtagaaga gcctcctaag aagccatctg cttcaagcc tgccattgaa atgcaaaact	960
ctgttccaaa taaagccttt gaattgaaga atgaacaaac attgagagca gatccgatgt	1020
tcccaccaga atccaaacaa aaggactatg aagaaaattc ttgggattct gagagttct	1080
gtgagactgt ttacacagaag gatgtgtgtt tacccaaggc tacacatcaa aaagaaatag	1140
ataaaaataaa tggaaaatta gaagagtctc ctaataaaaga tggcttctg aaggctacct	1200
gcggaatgaa agtttctatt ccaactaaag ctttagaatt gaaggacatg caaacttca	1260
aagcagagcc tccggggaaag ccatctgcct tcgagcctgc cactgaaatg caaaagtctg	1320
tcccaaataa agccttggaa ttgaaaaatg aacaaacatt gagagcagat gagataactcc	1380
catcagaatc caaacaAAAag gactatgaag aaagttcttgg ggattctgag agtctctgt	1440
agactgtttc acagaaggat gtgtgtttac ccaaggctrcc rcatcaaaaa gaaatagata	1500
aaataaaatgg aaaatttagaa gggatccttg ttaaagatgg tcttctgaag gctaaactgcg	1560
gaatgaaatgt ttctattcca actaaagcct tagaattgtt ggacatgcaaa acttcaag	1620
cagagcctcc cgagaagcca tctgccttcg agcctgcatt tgaaatgcaaa aagtctgtt	1680
caaataaaagc cttggaaattt aagaatgaac aaacatttag agcagatgag atactccat	1740
cagaatccaa acaaaaggac tatgaagaaa gttcttgggat ttctgagatg ctctgtgaga	1800
ctgtttcaca gaaggatgtg ttttaccca aggctrccat tcaaaaagaa atagataaaaa	1860
taaatggaaa attagaagag tctctgtata atgatggttt tctgagatgtt ccctgcagaa	1920
tgaaagtttc tattccaact aaagccttag aattgtatgg catgcaaaact ttcaaagcag	1980
agcctcccgaa gaagccatct gccttcgagc ctgcatttga aatgcaaaag tctgttccaa	2040
ataaaagcctt ggaatttgaag aatgaacaaa cattgagatg agatcagatg ttcccttcag	2100
aatcaaaaaca aaagaasgtt gaagaaaattt cttgggattt tgagatctc cgtgagactg	2160
tttcacagaa ggtatgtgtgtt gtacccaagg ctacacatca aaaagaaatgt gataaaaataa	2220
gtggaaaaattt agaagattca actagcctat caaaaatctt ggatacagtt cattttgt	2280
aaagagcaag ggaacttcaa aaagatcact gtgacacaacg tacagaaaaa atggaaacaaa	2340
tgaaaaagaa gttttgtgtt ctgaaaaaga aactgtcaga agcaaaagaa ataaaatcac	2400
agtttagagaa ccaaaaagttt aatggaaac aagagctctg cagtgtgagg tttctcacac	2460

-continued

tcatgaaaat	gaaaattatc	tcttacatga	aaattgcatg	ttgaaaaagg	aaattgccat	2520
gctaaaatcg	gaaatagcca	caactgaaaca	ccaataccag	gaaaaggaaa	ataaaatactt	2580
tgaggacatt	aagatttaa	aagaaaagaa	tgctgaactt	cagatgaccc	taaaactgaa	2640
agaggaaatca	ttaactaaaa	gggcatctca	atatagtggg	cagcttaaag	ttctgatagc	2700
tgagaacaca	atgctcaactt	ctaaattgaa	ggaaaaacaa	gacaaagaaa	tactagaggc	2760
agaaattgaa	tcacaccatc	ctagactggc	ttctgctgta	caagaccatg	atcaaattgt	2820
gacatcaaga	aaaagtcaag	aacatgcctt	ccacattgca	ggagatgctt	gttgcaaag	2880
aaaaatgaat	gttgatgtga	gttagtacgt	atataacaat	gaggtgcctc	atcaaccact	2940
ttctgaagct	caaaggaaat	ccaaaagcct	aaaaattaat	ctcaattatg	cmggagatgc	3000
tctaagagaa	aatacattgg	tttcagaaca	tgcacaaaaga	gaccaacgtg	aaacacagtg	3060
tcaaataatgaa	gaagctgaac	acatgtatca	aaacgaacaa	gataatgtga	acaaacacac	3120
tgaacacgac	gagtctctag	atcagaaatt	atttcaacta	caaagcaaaa	atatgtggct	3180
tcaacacgaa	ttagttcatg	cacataagaa	agctgacaac	aaaagcaaga	taacaattga	3240
tattcatttt	cttgagagga	aatgcaaca	tcatctccta	aaagagaaaa	atgaggagat	3300
atttaaattac	aataaccatt	taaaaaaccg	tatatatcaa	tatgaaaaag	agaaagcaga	3360
aacagaaaac	tcatgagaga	caagcagtaa	gaaacttctt	ttggagaaac	aacagaccag	3420
atcttactc	acaactcatg	ctaggaggcc	agtccttagca	tcacctttagt	ttgaaaatct	3480
taccaatagt	ctgtgtcaac	agaatactta	ttttagaaga	aaaattcatg	atttcttcct	3540
gaaggctaca	gacataaaat	aacagtgtga	agaattactt	gttcacgaat	tgcataaagc	3600
tgcacaggat	tcccatctac	cctgtatgt	cagcagacat	cattcaatcc	aaccagaatc	3660
tcgcctctgtc	actcaggctg	g				3681

<210> SEQ ID NO 20
<211> LENGTH: 1424
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 20

tccgagctga	ttacagacac	caaggaagat	gctgtaaaaga	gtcagcagcc	acagccctgg	60
ctagctggcc	ctgtggcat	ttattagtaa	agtttaatg	acaaaagctt	tgagtcaaca	120
cacccgtgg	taatattaccc	ggtcatcccc	accctggaga	gccatcctgc	ccatgggtga	180
tcaaagaagg	aacatctgca	ggaacacctg	atgaggctgc	acccttgcgc	gaaagaacac	240
ctgacacacg	tgaaagcttg	gtggaaaaaa	cacctgtatga	ggctgcaccc	ttggtgaaaa	300
gaacacctga	cacggctgaa	agcttggtgg	aaaaaacacc	tgtatgaggt	gcatccttgg	360
tggagggAAC	atctgacaaa	attcaatgtt	tggagaaagc	gacatctgga	aagttcgaac	420
agtcaacgaca	agaaacaccc	agggaaatta	cgagtccctgc	aaaagaaaca	tctgagaaat	480
ttacgtggcc	agccaaaaggaa	agaccttagga	agatcgatcg	ggagaaaaaa	gaagacacac	540
ctagggaaat	tatgagtccc	gcaaaagaaa	catctgagaa	atttacgtgg	gcagcaaaag	600
gaagaccttag	gaagatcgca	ttggagaaaa	aagaaacacc	tgtaaagact	ggatgcgtgg	660
caagagtaac	atctaataaa	actaaagttt	tggaaaaagg	aagatctaag	atgattgcat	720
gtcctacaaa	agaatcatct	acaaaagcaa	gtgccaatga	tcagaggttc	ccatcagaat	780

-continued

ccaaacaaga ggaagatgaa gaatattctt gtgattctcg gagtctctt gagagttctg	840
caaagattca agtgtgtata cctgagtcta tataatcaaaa agtaatggag ataaatagag	900
aagtagaaga gcotcctaag aagccatctg cttcaagcc tgccattgaa atgcaaaact	960
ctgttccaaa taaaggcttt gaattgaaga atgaacaaac attgagagca gatccgatgt	1020
tcccaccaga atccaaacaa aaggactatg aagaaaattc ttggattct gagagtctct	1080
gtgagactgt ttcacagaag gatgtgttt tacccaaggc tacacatcaa aaagaaatag	1140
ataaaataaa tggaaaatta gaaggtaaaa accgtttttt atttaaaaat cagttgaccg	1200
aatatttctc taaactgatg aggaggata tcctctagta gctgaagaaa attacccct	1260
aaatgcaaac catggaaaaaa aagagaagtg caatggcgt aagttgtatg tctcatcagg	1320
tgttggcaac agactatatt gagagtgcgt aaaaggagct gaattatttag tttgaattca	1380
agatattgca agacctgaga gaaaaaaaaaaaaaaa aaaaaaaaaaaaaaaa	1424

<210> SEQ ID NO 21

<211> LENGTH: 674

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 21

attccgagct gattacagac accaaggaag atgctgtaaa gagtcagcag ccacagccct	60
ggctagctgg ccctgtgggc atttattagt aaagttttaa tgacaaaagc tttgagtcaa	120
cacacccgtg ggttaattaac ctggtcatcc ccaccctgga gagccatcct gcccatgggt	180
gatcaaagaa ggaacatctg caggaacacc tgatgagggt gcacccttgg cgaaagaac	240
acctgacaca gctgaaagct tggggaaaaaacacctgtat gaggctgcac ctttgggatgg	300
aagaacacct gacacggctg aaagcttgggt ggaaaaaaca cctgtatgagg ctgcacccctt	360
gggtggggg acatctgaca aaattcaatg tttggagaaa gcgacatctg gaaagttcga	420
acagtcaagca gaagaaacac ctagggaaat tacgagtccct gcaaaagaaa catctgagaa	480
atttacgtgg ccagcaaaag gaagacctag gaagatcgca tgggagaaaa aagatgactc	540
agttaaggca aaaaaaaaaaaaaaaa aaaaaaaaaaaaaaaa aaaaaaaaaaaaaaaa aaaaaaaaaaaaa	600
aaaaaaaaaaaaaaa aaaaaaaaaaaaaaaa aaaaaaaaaaaaaaaa aaaaaaaaaaaaaaaa aaaaaaaaaaaaa	660
aaaaaaaaaaaaaaa aaaaaaaaaaaaaaaa aaaaaaaaaaaaaaaa aaaaaaaaaaaaaaaa aaaaaaaaaaaaa	660
aaaaaaaaaaaaaaa aaaaa	674

<210> SEQ ID NO 22

<211> LENGTH: 1729

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<220> FEATURE:

<221> NAME/KEY: unsure

<222> LOCATION: (11)

<223> OTHER INFORMATION: n=A,T,C or G

<221> NAME/KEY: unsure

<222> LOCATION: (1128)

<223> OTHER INFORMATION: n=A,T,C or G

<400> SEQUENCE: 22

gaaagttcga ncagtcaagca gaagaaacac ctagggaaat tacgagtccct gcaaaagaaa	60
catctgagaa atttacgtgg ccagcaaaag gaagacctag gaagatcgca tgggagaaaa	120
aagaagacac acctaggaa attatgagtc ccgcaaaaga aacatctgag aaatttacgt	180

-continued

gggcagcaaa aggaagacct aggaagatcg catgggagaa aaaagaaaca cctgtaaaga 240
ctggatgcgt ggcagaagta acatctaata aaactaaagt tttggaaaaa ggaagatcta 300
agatgattgc atgtcctaca aaagaatcat ctacaaaagc aagtgc当地 gatcagaggt 360
tcccatcaga atccaaacaa gaggaagatg aagaatattc ttgtgattct cggagtctct 420
ttgagagttc tgcaagattt caagtgtta tacctgagtc tatatatcaa aaagtaatgg 480
agataaataag agaagtagaa gagcctccta agaagccatc tgcctcaag cctgcccattg 540
aaatgc当地 ctcgtttcca aataaagcct ttgaattgaa gaatgaacaa acattgagag 600
cagatccgat gttccacca gaatccaaac aaaaggacta tgaagaaaat tcttgggatt 660
ctgagagtc ctgtgagact gttcacaga agatgtgtg ttacccaag gctacacatc 720
aaaaagaaaat agataaaaata aatggaaaat tagaagactc tcctaataaa gatggcttcc 780
tgaaggctac ctgc当地aaatg aaagtttcta ttccaaactaa agccttagaa ttgaaggaca 840
tgcaaaactt caaagcagag ctc当地gggaa agccatctgc ctgc当地gc gcaactgaaa 900
tgcaaaagtc tgc当地aaat aaagccttgg aattgaaaaa tgaacaaaca ttgagagcag 960
atgagatact cccatcagaa tccaaacaaa aggactatga agaaaattct tggatactg 1020
agagtctctg tgagactgtt tcacagaagg atgtgttgg accaaggct ggc当地caaa 1080
aagaaataga taaaataaaat ggaaaattag aagggtctcc tggtaanat ggtcttctga 1140
aggctaactg cggaatgaaa gtttcttattc caactaaagc ct当地aaattt atggacatgc 1200
aaactttcaa agcagagcct cccgagaagc catctgc当地t cc当地ctgc当地t attgaaatgc 1260
aaaagtctgt tccaaataaa gc当地tggaaat tgaagaatga acaaacattt agagcagatg 1320
agataactccc atcagaatcc aaacaaaagg actatgaaga aagtttcttgg gattctgaga 1380
gtctctgtga gactgtttca cagaaggatg tgc当地tacc caaggctgc当地t catcaaaaag 1440
aaatagataa aataaatgga aaattttagaag gtaagaaccg ttttttattt aaaaatcatt 1500
tgaccaataa tttctctaaa ttgtatgatggaa aggatattct ctatgtatgtt aagaaaattt 1560
cctccttaataa gcaaaaccatg gaaaaaaaaga gaagtgc当地t ggtc当地atgc tatgtgtctc 1620
atcaggcatt ggcaacagac tatattgtga gtgc当地t ggc当地tgaat tactatgtt 1680
aattcaagat attcaagac gtgaggaaaa tgagaaaaaa aaaaaaaaaa 1729

<210> SEQ ID NO 23
<211> LENGTH: 1337
<212> TYPE: DNA
<213> ORGANISM: *Homo sapiens*

<400> SEQUENCE: 23

aaaagaaaat agataaaata aatggaaaat tagaagggtc tcctgttaaa gatggtcttc	60
tgaaggctaa ctgcggaatg aaagtttcta ttccaaactaa agccttagaa ttgtatggaca	120
tgc当地acttt caaagcagag cctcccgaga agccatctgc ctgc当地gacct gccattgaaa	180
tgc当地aaagtc tggtccaaat aaagccttgg aattgaagaa tgaacaaaca ttgagagcag	240
atgagatact cccatcagaa tccaaacaaa aggactatga agaaagttct tggattctg	300
agagtctctg tgagactgtt tcacagaagg atgtgtgtt accaaggct gc当地atcaa	360
aagaaaataga taaaataaaat ggaaaattag aagagtctcc tgataatgat gttttctga	420
aqgctccctq cagaatqaaa qtttcttattc caactaaqc ct当地aattq atqgacatqc	480

-continued

aaacttcaa	agcagagcct	cccgagaagc	catctgcctt	cgagcctgcc	attgaaatgc	540
aaaagtctgt	tccaaataaa	gccttggat	tgaagaatga	acaaacattg	agagcagatc	600
agatgttccc	ttcagaatca	aaacaaaaga	aggttgaaga	aaattctgg	gattctgaga	660
gtctccgtga	gactgtttca	cagaaggatg	tgtgtgtacc	caaggctaca	catcaaaaag	720
aaatggataa	aataagtgg	aaatttagaag	attcaactag	cctatcaaaa	atcttggata	780
cagttcattc	tttgtgaaaga	gcaagggAAC	ttcaaaaaga	tcactgtgaa	caacgtacag	840
gaaaaatgg	acaaatggaa	aagaagttt	gtgtactgaa	aaagaaactg	tcagaagcaa	900
aagaaaataaa	atcacagtta	gagaacaaa	aagttaaatg	ggaacaagag	ctctgcagtg	960
tgagattgac	tttaaacc	gaagaagaga	agagaagaaa	tgccgatata	ttaatgaaa	1020
aaatttaggg	agaatttagga	agaatcgaag	agcagcata	gaaagagtt	gaagtggaaac	1080
aacaacttga	acaggctc	agaatacaag	atatagaatt	gaagagtgt	gaaagtaatt	1140
tgaatcaggt	tttcacact	catgaaaatg	aaaattatct	cttacatgaa	aattgcatgt	1200
tgaaaaagg	aattgccat	ctaaaactgg	aaatagccac	actgaaacac	caataccagg	1260
aaaaggaaaa	taaatactt	gaggacatta	agatttaaa	agaaaagaat	gctgaacttc	1320
agatgacccc	tcgtgcc					1337

<210> SEQ ID NO 24

<211> LENGTH: 2307

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 24

attgagagca	gatgagatac	tcccatcaga	atccaaacaa	aaggactatg	aagaaagttc	60
ttgggattct	gagagtctct	gtgagactgt	ttcacagaag	gatgtgtt	taccaaggc	120
tacacatcaa	aaagaaatag	ataaaataaa	tggaaaatta	gaagggtctc	ctgttaaaga	180
tggcttctg	aaggctaact	gcggatgaa	agtttctatt	ccaaactaaag	ccttagatt	240
gatggacatg	caaacttca	aagcagagcc	tcccgagaag	ccatctgcct	tcgaggctgc	300
cattgaaatg	caaaaagtctg	ttccaaataa	agccttggaa	ttgaagaatg	aacaaacatt	360
gagagcagat	gagatactcc	catcagaatc	caaacaaaag	gactatgaa	aaagttcttg	420
ggattctgag	agtctctgt	agactgtttc	acagaaggat	gtgtgttac	ccaaggctac	480
acataaaaaa	gaaatagata	aaataatgg	aaaattagaa	gagtctcctg	ataatgtgg	540
ttttctgaag	tctccctgca	gaatgaaagt	ttctattcca	actaaagcct	tagaattgat	600
ggacatgcaa	acttcaaag	cagagcctcc	cgagaagcca	tctgccttcg	agcctgccc	660
tgaaatgcaa	aagtctgttc	caaataaagc	cttggattt	aagaatgaa	aaacattgag	720
agcagatcg	atgttccctt	cagaatcaa	acaaaagaac	gttgaagaaa	attcttggg	780
ttctgagagt	ctccgtgaga	ctgtttcaca	gaaggatgt	tgtgtaccca	aggctacaca	840
tcaaaaaagaa	atggataaaa	taagtggaaa	attagaatg	tcaactagcc	tatcaaaaat	900
cttggataca	gttcatttt	gtgaaagagc	aaggaaactt	caaaaagatc	actgtgaaca	960
acgtacagga	aaaatggaac	aaatgaaaaa	gaagttttgt	gtactgaaaa	agaaactgtc	1020
agaagcaaaa	gaaataaaaat	cacagttaga	gaaccaaaaa	gttaaatggg	aacaagagct	1080
ctgcagtgtg	aggttctca	cactcatgaa	aatgaaaatt	atcttaca	tgaaaattgc	1140

-continued

atgttggaaa agggaaattgc catgctaaaa ctggaaatag ccacactgaa acaccaatac	1200
caggaaaagg aaaataaaata ctttggaggac attaagatt taaaagaaaa gaatgtgaa	1260
cttcagatga ccctaaaact gaaagaggaa tcattaacta aaagggcatc tcaatatagt	1320
gggcagctta aagttctgtat agctgagaac acaatgctca cttctaaatt gaaggaaaaa	1380
caagacaaag aaataactaga ggcagaaaatt gaatcacacc atcctagact ggcttctgct	1440
gtacaagacc atgatcaa at tttccacatt	1500
gcaggagatg ctgtttgc aagaaaaatg aatgttgatg tgagtagtac gatataaac	1560
aatgaggtgc tccatcaacc actttctgaa gctcaaagga aatccaaaag cctaaaaatt	1620
aatctcaatt atgcaggaga tgctctaaga gaaaatacat tggttcaga acatgcacaa	1680
agagaccaac gtgaaacaca gtgtcaa atg aaggaagctg aacacatgt a tcaaaaacgaa	1740
caagataatg tgaacaaaca cactgaacag caggagtctc tagatcagaa attatttcaa	1800
ctacaaaagca aaaatatgtg gcttcaacag caatttagttc atgcacataa gaaagctgac	1860
aacaaaagca agataacaat tgatattcat tttcttgaga ggaaaatgca acatcatctc	1920
ctaaaaagaga aaaatgagga gatatttaat tacaataacc attttaaaaa ccgtatata	1980
caatatgaaa aagagaaaagc agaaacagaa aactcatgag agacaagcag taagaaaactt	2040
cttttggaga aacaacagac cagatctta ctcacaactc atgctaggag gccagtccta	2100
gcatacacctt atgttggaaa tcttaccaat agtctgtgtc aacagaatac ttattttaga	2160
agaaaaattc atgatttctt cctgaagcct acagacataa aataacagtg tgaagaatta	2220
cttggcactc aattgcataa agctgcacag gattccatc taccctgtatg atgcagcaga	2280
catcattcaa tccaaaccaga atctcg	2307

<210> SEQ ID NO 25
 <211> LENGTH: 650
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens
 <220> FEATURE:
 <221> NAME/KEY: unsure
 <222> LOCATION: (310)
 <223> OTHER INFORMATION: Xaa = Any Amino Acid
 <221> NAME/KEY: unsure
 <222> LOCATION: (429)
 <223> OTHER INFORMATION: Xaa = Any Amino Acid
 <221> NAME/KEY: unsure
 <222> LOCATION: (522)
 <223> OTHER INFORMATION: Xaa = Any Amino Acid

<400> SEQUENCE: 25

Met Ser Pro Ala Lys Glu Thr Ser Glu Lys Phe Thr Trp Ala Ala Lys	
5	10
15	

Gly Arg Pro Arg Lys Ile Ala Trp Glu Lys Lys Glu Thr Pro Val Lys	
20	25
30	

Thr Gly Cys Val Ala Arg Val Thr Ser Asn Lys Thr Lys Val Leu Glu	
35	40
45	

Lys Gly Arg Ser Lys Met Ile Ala Cys Pro Thr Lys Glu Ser Ser Thr	
50	55
60	

Lys Ala Ser Ala Asn Asp Gln Arg Phe Pro Ser Glu Ser Lys Gln Glu	
65	70
75	80

Glu Asp Glu Glu Tyr Ser Cys Asp Ser Arg Ser Leu Phe Glu Ser Ser	
85	90
95	

-continued

Ala Lys Ile Gln Val Cys Ile Pro Glu Ser Ile Tyr Gln Lys Val Met
 100 105 110

Glu Ile Asn Arg Glu Val Glu Glu Pro Pro Lys Lys Pro Ser Ala Phe
 115 120 125

Lys Pro Ala Ile Glu Met Gln Asn Ser Val Pro Asn Lys Ala Phe Glu
 130 135 140

Leu Lys Asn Glu Gln Thr Leu Arg Ala Asp Pro Met Phe Pro Pro Glu
 145 150 155 160

Ser Lys Gln Lys Asp Tyr Glu Glu Asn Ser Trp Asp Ser Glu Ser Leu
 165 170 175

Cys Glu Thr Val Ser Gln Lys Asp Val Cys Leu Pro Lys Ala Thr His
 180 185 190

Gln Lys Glu Ile Asp Lys Ile Asn Gly Lys Leu Glu Glu Ser Pro Asn
 195 200 205

Lys Asp Gly Leu Leu Lys Ala Thr Cys Gly Met Lys Val Ser Ile Pro
 210 215 220

Thr Lys Ala Leu Glu Leu Lys Asp Met Gln Thr Phe Lys Ala Glu Pro
 225 230 235 240

Pro Gly Lys Pro Ser Ala Phe Glu Pro Ala Thr Glu Met Gln Lys Ser
 245 250 255

Val Pro Asn Lys Ala Leu Glu Leu Lys Asn Glu Gln Thr Leu Arg Ala
 260 265 270

Asp Glu Ile Leu Pro Ser Glu Ser Lys Gln Lys Asp Tyr Glu Glu Ser
 275 280 285

Ser Trp Asp Ser Glu Ser Leu Cys Glu Thr Val Ser Gln Lys Asp Val
 290 295 300

Cys Leu Pro Lys Ala Xaa His Gln Lys Glu Ile Asp Lys Ile Asn Gly
 305 310 315 320

Lys Leu Glu Gly Ser Pro Val Lys Asp Gly Leu Leu Lys Ala Asn Cys
 325 330 335

Gly Met Lys Val Ser Ile Pro Thr Lys Ala Leu Glu Leu Met Asp Met
 340 345 350

Gln Thr Phe Lys Ala Glu Pro Pro Glu Lys Pro Ser Ala Phe Glu Pro
 355 360 365

Ala Ile Glu Met Gln Lys Ser Val Pro Asn Lys Ala Leu Glu Leu Lys
 370 375 380

Asn Glu Gln Thr Leu Arg Ala Asp Glu Ile Leu Pro Ser Glu Ser Lys
 385 390 395 400

Gln Lys Asp Tyr Glu Glu Ser Ser Trp Asp Ser Glu Ser Leu Cys Glu
 405 410 415

Thr Val Ser Gln Lys Asp Val Cys Leu Pro Lys Ala Xaa His Gln Lys
 420 425 430

Glu Ile Asp Lys Ile Asn Gly Lys Leu Glu Glu Ser Pro Asp Asn Asp
 435 440 445

Gly Phe Leu Lys Ala Pro Cys Arg Met Lys Val Ser Ile Pro Thr Lys
 450 455 460

Ala Leu Glu Leu Met Asp Met Gln Thr Phe Lys Ala Glu Pro Pro Glu
 465 470 475 480

Lys Pro Ser Ala Phe Glu Pro Ala Ile Glu Met Gln Lys Ser Val Pro
 485 490 495

Asn Lys Ala Leu Glu Leu Lys Asn Glu Gln Thr Leu Arg Ala Asp Gln

-continued

500	505	510
Met Phe Pro Ser Glu Ser Lys Gln Lys Xaa Val Glu Glu Asn Ser Trp		
515	520	525
Asp Ser Glu Ser Leu Arg Glu Thr Val Ser Gln Lys Asp Val Cys Val		
530	535	540
Pro Lys Ala Thr His Gln Lys Glu Met Asp Lys Ile Ser Gly Lys Leu		
545	550	555
Glu Asp Ser Thr Ser Leu Ser Lys Ile Leu Asp Thr Val His Ser Cys		
565	570	575
Glu Arg Ala Arg Glu Leu Gln Lys Asp His Cys Glu Gln Arg Thr Gly		
580	585	590
Lys Met Glu Gln Met Lys Lys Phe Cys Val Leu Lys Lys Lys Leu		
595	600	605
Ser Glu Ala Lys Glu Ile Lys Ser Gln Leu Glu Asn Gln Lys Val Lys		
610	615	620
Trp Glu Gln Glu Leu Cys Ser Val Arg Phe Leu Thr Leu Met Lys Met		
625	630	635
Lys Ile Ile Ser Tyr Met Lys Ile Ala Cys		
645	650	

<210> SEQ ID NO 26
 <211> LENGTH: 228
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 26

Met Ser Pro Ala Lys Glu Thr Ser Glu Lys Phe Thr Trp Ala Ala Lys	5	10	15
Gly Arg Pro Arg Lys Ile Ala Trp Glu Lys Lys Glu Thr Pro Val Lys			
20	25	30	
Thr Gly Cys Val Ala Arg Val Thr Ser Asn Lys Thr Lys Val Leu Glu			
35	40	45	
Lys Gly Arg Ser Lys Met Ile Ala Cys Pro Thr Lys Glu Ser Ser Thr			
50	55	60	
Lys Ala Ser Ala Asn Asp Gln Arg Phe Pro Ser Glu Ser Lys Gln Glu			
65	70	75	80
Glu Asp Glu Glu Tyr Ser Cys Asp Ser Arg Ser Leu Phe Glu Ser Ser			
85	90	95	
Ala Lys Ile Gln Val Cys Ile Pro Glu Ser Ile Tyr Gln Lys Val Met			
100	105	110	
Glu Ile Asn Arg Glu Val Glu Glu Pro Pro Lys Lys Pro Ser Ala Phe			
115	120	125	
Lys Pro Ala Ile Glu Met Gln Asn Ser Val Pro Asn Lys Ala Phe Glu			
130	135	140	
Leu Lys Asn Glu Gln Thr Leu Arg Ala Asp Pro Met Phe Pro Pro Glu			
145	150	155	160
Ser Lys Gln Lys Asp Tyr Glu Glu Asn Ser Trp Asp Ser Glu Ser Leu			
165	170	175	
Cys Glu Thr Val Ser Gln Lys Asp Val Cys Leu Pro Lys Ala Thr His			
180	185	190	
Gln Lys Glu Ile Asp Lys Ile Asn Gly Lys Leu Glu Gly Lys Asn Arg			
195	200	205	

-continued

Phe Leu Phe Lys Asn Gln Leu Thr Glu Tyr Phe Ser Lys Leu Met Arg
210 215 220

Arg Asp Ile Leu
225

<210> SEQ ID NO 27
<211> LENGTH: 154
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: unsure
<222> LOCATION: (148)
<223> OTHER INFORMATION: Xaa = Any Amino Acid

<400> SEQUENCE: 27

Met Arg Leu His Pro Trp Arg Lys Glu His Leu Thr Gln Leu Lys Ala
5 10 15

Trp Trp Lys Lys His Leu Met Arg Leu His Pro Trp Trp Lys Glu His
20 25 30

Leu Thr Arg Leu Lys Ala Trp Trp Lys Lys His Leu Met Arg Leu His
35 40 45

Pro Trp Trp Arg Glu His Leu Thr Lys Phe Asn Val Trp Arg Lys Arg
50 55 60

His Leu Glu Ser Ser Asn Ser Gln Gln Lys Lys His Leu Gly Lys Leu
65 70 75 80

Arg Val Leu Gln Lys Lys His Leu Arg Asn Leu Arg Gly Gln Gln Lys
85 90 95

Glu Asp Leu Gly Arg Ser His Gly Arg Lys Lys Met Thr Gln Leu Arg
100 105 110

Gln Lys
115 120 125

Lys
130 135 140

Lys Lys Lys Xaa Lys Lys Lys Lys Lys Lys
145 150

<210> SEQ ID NO 28
<211> LENGTH: 466
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: unsure
<222> LOCATION: (329)
<223> OTHER INFORMATION: Xaa = Any Amino Acid

<400> SEQUENCE: 28

Met Ser Pro Ala Lys Glu Thr Ser Glu Lys Phe Thr Trp Ala Ala Lys
5 10 15

Gly Arg Pro Arg Lys Ile Ala Trp Glu Lys Lys Glu Thr Pro Val Lys
20 25 30

Thr Gly Cys Val Ala Arg Val Thr Ser Asn Lys Thr Lys Val Leu Glu
35 40 45

Lys Gly Arg Ser Lys Met Ile Ala Cys Pro Thr Lys Glu Ser Ser Thr
50 55 60

Lys Ala Ser Ala Asn Asp Gln Arg Phe Pro Ser Glu Ser Lys Gln Glu
65 70 75 80

Glu Asp Glu Glu Tyr Ser Cys Asp Ser Arg Ser Leu Phe Glu Ser Ser

-continued

85	90	95
Ala Lys Ile Gln Val Cys Ile Pro Glu Ser Ile Tyr Gln Lys Val Met		
100	105	110
Glu Ile Asn Arg Glu Val Glu Glu Pro Pro Lys Lys Pro Ser Ala Phe		
115	120	125
Lys Pro Ala Ile Glu Met Gln Asn Ser Val Pro Asn Lys Ala Phe Glu		
130	135	140
Leu Lys Asn Glu Gln Thr Leu Arg Ala Asp Pro Met Phe Pro Pro Glu		
145	150	155
Ser Lys Gln Lys Asp Tyr Glu Glu Asn Ser Trp Asp Ser Glu Ser Leu		
165	170	175
Cys Glu Thr Val Ser Gln Lys Asp Val Cys Leu Pro Lys Ala Thr His		
180	185	190
Gln Lys Glu Ile Asp Lys Ile Asn Gly Lys Leu Glu Ser Pro Asn		
195	200	205
Lys Asp Gly Leu Leu Lys Ala Thr Cys Gly Met Lys Val Ser Ile Pro		
210	215	220
Thr Lys Ala Leu Glu Leu Lys Asp Met Gln Thr Phe Lys Ala Glu Pro		
225	230	235
Pro Gly Lys Pro Ser Ala Phe Glu Pro Ala Thr Glu Met Gln Lys Ser		
245	250	255
Val Pro Asn Lys Ala Leu Glu Leu Lys Asn Glu Gln Thr Leu Arg Ala		
260	265	270
Asp Glu Ile Leu Pro Ser Glu Ser Lys Gln Lys Asp Tyr Glu Glu Asn		
275	280	285
Ser Trp Asp Thr Glu Ser Leu Cys Glu Thr Val Ser Gln Lys Asp Val		
290	295	300
Cys Leu Pro Lys Ala Ala His Gln Lys Glu Ile Asp Lys Ile Asn Gly		
305	310	315
Lys Leu Glu Gly Ser Pro Gly Lys Xaa Gly Leu Leu Lys Ala Asn Cys		
325	330	335
Gly Met Lys Val Ser Ile Pro Thr Lys Ala Leu Glu Leu Met Asp Met		
340	345	350
Gln Thr Phe Lys Ala Glu Pro Pro Glu Lys Pro Ser Ala Phe Glu Pro		
355	360	365
Ala Ile Glu Met Gln Lys Ser Val Pro Asn Lys Ala Leu Glu Leu Lys		
370	375	380
Asn Glu Gln Thr Leu Arg Ala Asp Glu Ile Leu Pro Ser Glu Ser Lys		
385	390	395
400		
Gln Lys Asp Tyr Glu Glu Ser Ser Trp Asp Ser Glu Ser Leu Cys Glu		
405	410	415
Thr Val Ser Gln Lys Asp Val Cys Leu Pro Lys Ala Ala His Gln Lys		
420	425	430
Glu Ile Asp Lys Ile Asn Gly Lys Leu Glu Gly Lys Asn Arg Phe Leu		
435	440	445
Phe Lys Asn His Leu Thr Lys Tyr Phe Ser Lys Leu Met Arg Lys Asp		
450	455	460
Ile Leu		
465		

-continued

<211> LENGTH: 445

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 29

Lys Glu Ile Asp Lys Ile Asn Gly Lys Leu Glu Gly Ser Pro Val Lys
5 10 15

Asp Gly Leu Leu Lys Ala Asn Cys Gly Met Lys Val Ser Ile Pro Thr
20 25 30

Lys Ala Leu Glu Leu Met Asp Met Gln Thr Phe Lys Ala Glu Pro Pro
35 40 45

Glu Lys Pro Ser Ala Phe Glu Pro Ala Ile Glu Met Gln Lys Ser Val
50 55 60

Pro Asn Lys Ala Leu Glu Leu Lys Asn Glu Gln Thr Leu Arg Ala Asp
65 70 75 80

Glu Ile Leu Pro Ser Glu Ser Lys Gln Lys Asp Tyr Glu Glu Ser Ser
85 90 95

Trp Asp Ser Glu Ser Leu Cys Glu Thr Val Ser Gln Lys Asp Val Cys
100 105 110

Leu Pro Lys Ala Ala His Gln Lys Glu Ile Asp Lys Ile Asn Gly Lys
115 120 125

Leu Glu Glu Ser Pro Asp Asn Asp Gly Phe Leu Lys Ala Pro Cys Arg
130 135 140

Met Lys Val Ser Ile Pro Thr Lys Ala Leu Glu Leu Met Asp Met Gln
145 150 155 160

Thr Phe Lys Ala Glu Pro Pro Glu Lys Pro Ser Ala Phe Glu Pro Ala
165 170 175

Ile Glu Met Gln Lys Ser Val Pro Asn Lys Ala Leu Glu Leu Lys Asn
180 185 190

Glu Gln Thr Leu Arg Ala Asp Gln Met Phe Pro Ser Glu Ser Lys Gln
195 200 205

Lys Lys Val Glu Glu Asn Ser Trp Asp Ser Glu Ser Leu Arg Glu Thr
210 215 220

Val Ser Gln Lys Asp Val Cys Val Pro Lys Ala Thr His Gln Lys Glu
225 230 235 240

Met Asp Lys Ile Ser Gly Lys Leu Glu Asp Ser Thr Ser Leu Ser Lys
245 250 255

Ile Leu Asp Thr Val His Ser Cys Glu Arg Ala Arg Glu Leu Gln Lys
260 265 270

Asp His Cys Glu Gln Arg Thr Gly Lys Met Glu Gln Met Lys Lys Lys
275 280 285

Phe Cys Val Leu Lys Lys Lys Leu Ser Glu Ala Lys Glu Ile Lys Ser
290 295 300

Gln Leu Glu Asn Gln Lys Val Lys Trp Glu Gln Glu Leu Cys Ser Val
305 310 315 320

Arg Leu Thr Leu Asn Gln Glu Glu Lys Arg Arg Asn Ala Asp Ile
325 330 335

Leu Asn Glu Lys Ile Arg Glu Glu Leu Gly Arg Ile Glu Glu Gln His
340 345 350

Arg Lys Glu Leu Glu Val Lys Gln Gln Leu Glu Gln Ala Leu Arg Ile
355 360 365

Gln Asp Ile Glu Leu Lys Ser Val Glu Ser Asn Leu Asn Gln Val Ser

-continued

370	375	380
His Thr His Glu Asn Glu Asn Tyr Leu Leu His Glu Asn Cys Met Leu		
385	390	395
Lys Lys Glu Ile Ala Met Leu Lys Leu Glu Ile Ala Thr Leu Lys His		
405	410	415
Gln Tyr Gln Glu Lys Glu Asn Lys Tyr Phe Glu Asp Ile Lys Ile Leu		
420	425	430
Lys Glu Lys Asn Ala Glu Leu Gln Met Thr Pro Arg Ala		
435	440	445

<210> SEQ ID NO 30

<211> LENGTH: 578

<212> TYPE: DNA

<213> ORGANISM: Human

<400> SEQUENCE: 30

cttgccttct cttaggcttt gaagcatttt tgtctgtgtt ccctgatctt caggtcacca	60
ccatgaagtt cttagcagtc ctggactct tgggagtttc catctttctg gtctctgccc	120
agaatccgac aacagctgct ccagctgaca cgtatccagc tactggcctt gctgatgatg	180
aagccctga tgcgtgaaacc actgctgctg caaccactgc gaccactgct gctcctacca	240
ctgcaaccac cgctgcttct accactgctc gtaaagacat tccagttta cccaaatggg	300
ttggggatct cccgaatggt agagtgtgtc cctgagatgg aatcagctt agtcttctgc	360
aattggtcac aactattcat gcttcctgtg atttcatcca actacttacc ttgcctacga	420
tatccccctt atctctaatac agtttatttt ctttcaaata aaaaataact atgagcaaca	480
aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa	540
aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa	578

<210> SEQ ID NO 31

<211> LENGTH: 90

<212> TYPE: PRT

<213> ORGANISM: Homo sapien

<400> SEQUENCE: 31

Met Lys Phe Leu Ala Val Leu Val Leu Leu Gly Val Ser Ile Phe Leu	1 5 10 15	
Val Ser Ala Gln Asn Pro Thr Thr Ala Ala Pro Ala Asp Thr Tyr Pro	20 25 30	
Ala Thr Gly Pro Ala Asp Asp Glu Ala Pro Asp Ala Glu Thr Thr Ala	35 40 45	
Ala Ala Thr Thr Ala Thr Thr Ala Ala Pro Thr Thr Ala Thr Thr Ala	50 55 60	
Ala Ser Thr Thr Ala Arg Lys Asp Ile Pro Val Leu Pro Lys Trp Val	65 70 75 80	
Gly Asp Leu Pro Asn Gly Arg Val Cys Pro	85 90	

<210> SEQ ID NO 32

<211> LENGTH: 3101

<212> TYPE: DNA

<213> ORGANISM: Homo sapien

<400> SEQUENCE: 32

-continued

tgttggggcc tcagcctccc aagtagctgg gactacaggt gcctgccacc acggccagct 60
aattttttgt atattttta gtagagacgg ggtttcaccg tggctcaat ctccgtaccc 120
cgtgatctgc cagccttggc ctcccaaagt gtattctatt tttatttatta ttattttttt 180
ttagatggag tctgtctcg tcgcccaggc tggagtgcag tggtgcatc tctgctcaact 240
gcaagctccg cctcctgggt tcatgccatt ctcctgcctc agcctcccgta gtagctggga 300
ctacaggccc ctgccaccac acccggtctaa tttttgtat ttttagtaga gacagggttt 360
caccatgtta gccagggtgg tctctatctt ctgacacctg gatccgcctg cctcgtctc 420
tcaaagtgtt gggattacag gcgtgagcca ccgcgaccag ccaactattt ctgttttattt 480
ttaaatatata tttaaagaaa caatttagatt tgtttcttt ctcattctt tacttctact 540
cttcatgtat gtataattat atttgtgttt tctattacct ttttccttt tactgtattt 600
gactataata attgtgctca ctaattctg ttcactaata ttatcagctt agataataact 660
ttaatttta acttatatat tgagtattaa attgatcagt tttatttga attatctatc 720
ttccgcctgg ctgaatataa cttcttaagc ttataacttc ttgttcttc catgttattt 780
ttttctttt ttaatgtat tgaattctt ctgacactca ttctagtaac tttttctcg 840
gtgtgcacacg taaggtaataa ttgttctc agatttggaa tctgccataa gtttggggct 900
ttatttttt ttttatggca agtcggacaa cctgcatgga tttggcatca 960
atgtagtac ccatatctaa gagcagcact tgcttcttag catgtgagt tgtttctgga 1020
ttgtttctt atttactta tattctggt agattctt atttccctt caactctatt 1080
cagcattta ggaattctt ggacttctg agaattttag cttctgtat taaatgtttt 1140
taatgagtagt tgcatctt caaaaagcac aaatataatc agtgtacaca tgaggaaaac 1200
tatataataa ttctgttgca gatgacagca ttcataaca aaatcctagt tacttcattt 1260
aaaagacagc ttcctccaa tataactatga ggtaacaaaa atttgttagt tgtaattttt 1320
ttaatattag aaaactcatc ttacattgtg cacaatttc tgaagtgata atacttcact 1380
gttttctat agaagtaact taatattggc aaaattactt atttgaattt aggtttggc 1440
tttcatcata tacttcctca ttaacatttc cctcaatcca taaatgcaat ctcatgttga 1500
atcttccatt taaccaggaa gttaattttt aaaaccttaa taaaatttga atgttagctag 1560
atattatttg ttggttacat attgtcaat aatttatattt acttacaatg atcagaaaat 1620
atgtatctgaa ttctgtgtt cataattca ataacgtatt ttggcctaa acctttccat 1680
ttcaaatcct tgggtctggt aattgaaaat aatcattatc tttgttttc tggccaaaaa 1740
tgctgcccatt tattttctat ccctaatttag tcaaaacttcc taataatgtt atttaacgtt 1800
aatgatgttt atttgctgt tggataactaa aaccatttagt ttctataatt taaatgtcac 1860
ctaataatgag tgaaaatgtg tcagaggctg gggagaatg tggatggaga aaggaaagg 1920
gttgcataaa aagtacccaa gtttcgtt cacaggaggc atgagattga tctgtgcac 1980
aaaatgtga gtataataaa taataatgca ctgtatattt tggaaattgtt aaaaatgtat 2040
ttaaaattgtt tttacataat attttacata tttataaaggc acatgcaata tggatgttaca 2100
tgtatagaat gtgcacatgat caagtcaggg tatctgtgtt atccaccact ttgagcattt 2160
atcgattcta tatgtcagga acatttcaag ttatctgttc tagcaaggaa atataaaaata 2220
cattataatgtt aactatqgccc tatctcataatgtt qcaactaaac actaattttt attcctttcc 2280

-continued

aactgtgggt ttgttattcat ttaccaccct cttttcattc cctttctcac ccacacactg	2340
tgccgggcct caggcatata ctattctact gtctgtctct gtaaggatta tcatttttagc	2400
ttccacatata gagagaatgc atgcaaagtt tttcttcca tgtctggctt atttcactta	2460
acaaaatgac ctccgcttcc atccatgtta tttatattac ccaatagtgt tcataaatat	2520
atatacacac atatatacca cattgcattt gtccaaattat tcattgacgg aaactgggta	2580
atgttatatac gttgctattt gtaatagtgc tgcaataaac acgcaagtgg ggatataatt	2640
tgaagagttt ttttggat gttccataca aatttttaaga ttgttttgc tatgttttg	2700
aaaatggcgt tagtattttc atagagattt cattgaatct gttagattgt ttgggtaagt	2760
atggttattt ttaggttattt aatttttca ttccatgaag atgagatgtc tttccattt	2820
tttggccttcc ttacattttc tttcatcaaa gttttgtgtt attttgaag tagatgttt	2880
tcacccata gatcaagtgtt attccctaaa tatttttattt ttgttagctat tgtagatgaa	2940
attgccttcc cgattttttt ttcaacttaat tcattttagt tgtagggaaa tgtagatggat	3000
ttttttttgt tggtttttaa tcaaaaactg tattaaactt agagttttt gtggagttt	3060
taagtttttc tagatataag atcatgacat ctacaaaaaa a	3101

<210> SEQ ID NO 33
 <211> LENGTH: 16
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: PCR primer

<400> SEQUENCE: 33

tgccctccg gaagct 16

<210> SEQ ID NO 34
 <211> LENGTH: 23
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: PCR primer

<400> SEQUENCE: 34

cgtttctgaa gggacatctg atc 23

<210> SEQ ID NO 35
 <211> LENGTH: 30
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: PCR primer

<400> SEQUENCE: 35

ttgcagccaa gttaggagtg aagagatgca 30

<210> SEQ ID NO 36
 <211> LENGTH: 24
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: PCR primer

<400> SEQUENCE: 36

aagcctcaga gtccttccag tatg 24

-continued

<210> SEQ ID NO 37
<211> LENGTH: 35
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: PCR primer

<400> SEQUENCE: 37

ttcaaatata agtgaagaaa aaatttagtag atcaa

35

<210> SEQ ID NO 38
<211> LENGTH: 37
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: PCR primer

<400> SEQUENCE: 38

aatccattgt atcttagaac cgagggattt gtttaga

37

<210> SEQ ID NO 39
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: PCR primer

<400> SEQUENCE: 39

aaagcagatg gtgggtgagg tt

22

<210> SEQ ID NO 40
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: PCR primer

<400> SEQUENCE: 40

cctgagacca aatggcttct tc

22

<210> SEQ ID NO 41
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: PCR primer

<400> SEQUENCE: 41

attccatgcc ggctgcttct tctg

24

<210> SEQ ID NO 42
<211> LENGTH: 30
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: PCR primer

<400> SEQUENCE: 42

tctggtttcc tcattcttta ttcatttttt

30

<210> SEQ ID NO 43
<211> LENGTH: 20
<212> TYPE: DNA

-continued

<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: PCR primer

<400> SEQUENCE: 43

tgccaaaggag cggattatct 20

<210> SEQ ID NO 44
<211> LENGTH: 30
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: PCR primer

<400> SEQUENCE: 44

caaccacgtg acaaacactg gaattacagg 30

<210> SEQ ID NO 45
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: PCR primer

<400> SEQUENCE: 45

actggaacgg tgaaggtgac a 21

<210> SEQ ID NO 46
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: PCR primer

<400> SEQUENCE: 46

cggccacatt gtgaactttg 20

<210> SEQ ID NO 47
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: PCR primer

<400> SEQUENCE: 47

cagtcggttg gagcgagcat ccc 23

<210> SEQ ID NO 48
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: PCR primer

<400> SEQUENCE: 48

tgcctatagat gaattgaagg aatg 24

<210> SEQ ID NO 49
<211> LENGTH: 29
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: PCR primer

-continued

<400> SEQUENCE: 49

tgtcatatat taattgcata aacacctca

29

<210> SEQ ID NO 50

<211> LENGTH: 32

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: PCR primer

<400> SEQUENCE: 50

tcttaaccaa acggatgaaa ctctgagcaa tg

32

<210> SEQ ID NO 51

<211> LENGTH: 28

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: PCR primer

<400> SEQUENCE: 51

atcattgaaa attcaaataat aagtgaag

28

<210> SEQ ID NO 52

<211> LENGTH: 30

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: PCR primer

<400> SEQUENCE: 52

gtagttgtgc attgaaataa ttatcattat

30

<210> SEQ ID NO 53

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: PCR Primer

<400> SEQUENCE: 53

caatttttgtt ggagaacccg

20

<210> SEQ ID NO 54

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: PCR Primer

<400> SEQUENCE: 54

gctgtcggag gtatatatggtg

20

<210> SEQ ID NO 55

<211> LENGTH: 28

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: PCR Primer

<400> SEQUENCE: 55

catttcagag agtaaacatgg actacaca

28

-continued

<210> SEQ ID NO 56
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: PCR Primer

<400> SEQUENCE: 56

tctgataaaag gccgtacaat g 21

<210> SEQ ID NO 57
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: PCR Primer

<400> SEQUENCE: 57

tcacgacttg ctgttttgc tc 22

<210> SEQ ID NO 58
<211> LENGTH: 30
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: PCR Primer

<400> SEQUENCE: 58

atcaaaaaaac aagcatggcc tcacaccact 30

<210> SEQ ID NO 59
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: PCR Primer

<400> SEQUENCE: 59

gcaagtgc当地 atgatcagag g 21

<210> SEQ ID NO 60
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: PCR Primer

<400> SEQUENCE: 60

atatagactc aggtatacac act 23

<210> SEQ ID NO 61
<211> LENGTH: 30
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: PCR Primer

<400> SEQUENCE: 61

tcccatcaga atccaaacaa gaggaagatg 30

<210> SEQ ID NO 62
<211> LENGTH: 34
<212> TYPE: DNA

-continued

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: PCR Primer

<400> SEQUENCE: 62

aatccattgt atcttagaac cgagggattt gttt

34

<210> SEQ ID NO 63

<211> LENGTH: 24

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: PCR Primer

<400> SEQUENCE: 63

ccgcttctga caacactaga gatc

24

<210> SEQ ID NO 64

<211> LENGTH: 32

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: PCR Primer

<400> SEQUENCE: 64

cctataaaga tggatgtac caaaaatgaa gt

32

<210> SEQ ID NO 65

<211> LENGTH: 22

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: PCR Primer

<400> SEQUENCE: 65

ccccccctc aggatggc cc

22

<210> SEQ ID NO 66

<211> LENGTH: 22

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: PCR Primer

<400> SEQUENCE: 66

cccttctca cccacacact gt

22

<210> SEQ ID NO 67

<211> LENGTH: 24

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: PCR Primer

<400> SEQUENCE: 67

tgcattctc catatgtgga agct

24

<210> SEQ ID NO 68

<211> LENGTH: 33

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: PCR Primer

-continued

<400> SEQUENCE: 68

ccgggcctca ggcataact attctactgt ctg 33

<210> SEQ ID NO 69
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: PCR Primer

<400> SEQUENCE: 69

gacattccag ttttacccaa atgg 24

<210> SEQ ID NO 70
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: PCR Primer

<400> SEQUENCE: 70

tgcagaagac tcaagctgat tcc 23

<210> SEQ ID NO 71
<211> LENGTH: 28
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: PCR Primer

<400> SEQUENCE: 71

tctcagggac acactctacc attcggga 28

<210> SEQ ID NO 72
<211> LENGTH: 30
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: PCR Primer

<400> SEQUENCE: 72

aaatataagt gaagaaaaaa attagtagat 30

<210> SEQ ID NO 73
<211> LENGTH: 503
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 73

gacagcggct tccttgatcc ttgccacccg cgactgaaca ccgacagcag cagcctcacc	60
atgaagttgc tggatggctt catgtggcg gccctctccc agcactgcta cgaggctct	120
ggctgcccct tattggagaa tgtgattcc aagacaatca atccacaatgt gtctaaact	180
gaataacaag aacttcttca agagttcata gacgacaatg ccactacaaa tgccatagat	240
gaattgaagg aatgtttct taaccaaactg gatgaaactc tgagcaatgt tgagggttt	300
ctgcaattaa tatatgacag cagttttgt gatttatttt aactttctgc aagacctttg	360
gctcacagaa ctgcagggta tggtgagaaa ccaactacgg attgctgca accacacctt	420
ctctttctta tgtctttta ctacaaacta caagacaatt gttgaaacct gctatacatg	480

-continued

tttattttaa taaattgatg gca	503
<p><210> SEQ ID NO 74 <211> LENGTH: 301 <212> TYPE: DNA <213> ORGANISM: Homo sapiens</p>	
<p><400> SEQUENCE: 74</p>	
cactgctacg caggctctgg ctgcccctta ttggagaatg tgattccaa gacaatcaat	60
ccacaagtgt ctaagactga atacaaagaa cttcttcaag agttcataga cgacaatgcc	120
actacaaatg ccatagatga attgaaggaa tgtttctta accaaacgga tgaaactctg	180
agcaatgttgc aggtgtttat gcaattaata tatgacagca gtctttgtga tttatggc	240
ggccatcacc atcaccatca ctaaggccc gagctcgaat tctgcagata tccatcacac	300
t	301
<p><210> SEQ ID NO 75 <211> LENGTH: 3282 <212> TYPE: DNA <213> ORGANISM: Homo sapiens</p>	
<p><400> SEQUENCE: 75</p>	
gggacacgggc tgaggatgag gagaaccctg gggacccaga agaccgtgcc ttgcccggaa	60
gtcctgcctg taggcctgaa ggacttgcctt taacagagcc tcaacaacta cctgggtatt	120
cctacttcag ccccttgggtg tgagcagctt ctcacatga actacagcct ccacttggcc	180
ttcgtgtgtc tgagtctt cactgagagg atgtgcatttcc aggggagtc gttcaacgtc	240
gaggtcggca gaagtgcacaa gctttccctg cctggctttt agaacctcac agcaggatata	300
aacaaatttc tcaggccaa ttttgggttga gaaccctgtc agatagcgtc gactctggac	360
attgcaagta tctcttagcat ttcaagagat aacatggact acacagccac catataccctc	420
cgacagcgctt ggttggatca gcccgttgc tttttttttt acaagagctt cactctggat	480
gccccctcg tggagttctt ctgggtgcata gatacttaca ttgtggagtc caagaagtcc	540
ttccctccatg aagtcaactgtt gggaaacagg ctcatccgc tcttctccaa tggcacggtc	600
ctgtatgcctt tcagaatcac gacaactgtt gcatgttacaa tggatctgtc taaatacccc	660
atggacacac agacatgcaaa gtttcagctg gaaagctgg gctatgtatgg aatgtatgt	720
gagttcacctt ggttggatca gacactctt gtgcgtggac tggacacacctt gcccgttgc	780
cagtacacca tagagcggta tttcacctt gtcaccatgtt cgcagcagga gacagggaaat	840
tacactagat tggcttaca gtttggatctt cggaggaatg ttctgtatccat cattttggaa	900
acctacgttc cttccactttt cctgggtgtt ttgtcctggg tttcatctt gatctcttc	960
gattcagtcc ctgcaagaac ctgcatttggaa gtgcacggcc tggatcttcaat gaccacactg	1020
atgtatcggtt cccgcacttc tcttccaaac accaactgtt tcatcaaggc catcgatgt	1080
tacctggggta tctgttttag ctgtgtgtttt ggggccttgc tagaatatgc agttgtcact	1140
tacagttccatc tacagcagat ggcagccaaa gatagggggca caacaaagga agtagaaagaa	1200
gtcagtttata ctaatatcat caacagctcc atctccagttt ttaaacggaa gatcagctt	1260
gccagcatttgc aaatttccatc cgacaacgtt gactacatgtt acttgacaaat gaaaaccagc	1320
gacaagttca agttgtctt ccgagaaaag atggcagga ttgttgcattttt tttccacaaatt	1380

-continued

caaaaaccca	gtaatgttga	tcactattcc	aaactactgt	ttcctttgat	ttttatgcta	1440						
gc	ccatgtat	tttactgggc	atactacatg	tat	ttttagag	tcaatgttaa	atttcttgca	1500				
tgccataggt	cttcaacagg	acaagataat	gatgtaaatg	gtat	tttagg	ccaagtgtgc	1560					
acccacatcc	aatggtgcta	caagtgactg	aaataatatt	tgagtcttcc	tgctcaaaga	1620						
atgaagctcc	aaccattgtt	ctaa	gctgtg	taga	agtcc	tttt	ttttgttgc	1680				
tagaaacatc	agtccattcc	tctt	tttcatct	taat	caagga	catt	cccattg	1740				
ttacaatgt	actcagg	ggct	tttattcgg	tgg	ctcc	cctt	gttgcattt	1800				
aagaatggga	aggagaccat	tgg	gttaaccc	tca	agtgtca	gaag	ttgttt	ctaaagtaac	1860			
tatacatgtt	ttttactaaa	tct	ctgc	act	tataaaaa	tacatt	tttg	cctattttagg	1920			
gagtaacatt	ttctag	tttt	tgtt	tctgg	taaaatgaa	tat	gggc	tta	tgtcaattca	1980		
ttggaagtca	atgc	acta	ac	tca	ataac	ca	gat	gat	tttt	atattattta	2040	
ataccacaac	agaatt	at	tatcc	cc	aaat	tcc	ata	gtc	ttt	ttcaaatata	2100	
agtgaagaaa	aaat	tttag	tag	atc	aaacaa	atc	cctcg	gtt	ttt	ttcaataat	2160	
ggattccccca	tact	gg	gaagg	act	ctgaggc	ttt	att	ccc	cact	atgc	tttatcat	2220
tttatttattt	ta	cac	acatc	cat	cctaa	tat	act	tttccc	atgc	atggat	2280	
ggaaatggaa	gat	ttttt	ttt	ta	actt	gtt	cc	ta	ttt	tttgc	atgcattga	2340
aggcttgcag	aat	ttagt	cc	tcc	tttgc	tttgc	tttgc	tttgc	tttgc	tttgc	tttgc	2400
aaagtactgg	gtt	gact	tcag	ag	atcg	ctg	tc	att	ctg	ct	taacact	2460
gagcaacact	ct	ccc	act	gg	cag	atccc	ct	atc	tttcc	tttcc	tttcc	2520
tgctcta	at	tc	at	atg	at	gat	ttt	tttgc	tttgc	tttgc	tttgc	2580
ggcttagctt	aa	gt	taa	act	tttgc	tttgc	tttgc	tttgc	tttgc	tttgc	tttgc	2640
tgagtggc	ttt	at	cc	cc	at	cc	at	cc	at	cc	at	2700
aacaatgaga	ca	cac	ttt	ttt	ttt	ttt	ttt	ttt	ttt	ttt	ttt	2760
ccagccagag	at	gc	ac	at	tc	cc	at	tc	cc	at	tc	2820
agtgtgccag	gg	taa	agg	gt	cag	ttc	at	ttt	tttgc	tttgc	tttgc	2880
ttaatttctt	ag	ttt	cct	gt	tta	ata	aa	at	tttgc	tttgc	tttgc	2940
accaaggctc	taaa	aa	atg	ttt	cc	tct	tttgc	tttgc	tttgc	tttgc	tttgc	3000
ctttccat	tata	actt	tctc	aca	attc	agt	tttgc	tttgc	tttgc	tttgc	tttgc	3060
acaaaatatt	tct	aac	acgg	at	gg	gg	gt	gt	gg	gg	gg	3120
tatgagccaa	tca	at	ttt	ttt	ttt	ttt	ttt	ttt	ttt	ttt	ttt	3180
aaccccactt	aac	gat	ttt	ttt	ttt	ttt	ttt	ttt	ttt	ttt	ttt	3240
aatataccat	at	ta	ccat	cc	cc	cc	cc	cc	cc	cc	cc	3282

<210> SEQ ID NO 76

<211> LENGTH: 463

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 76

tagaattcag	cggccgctt	attctagaag	tccaaatcac	tcattgttgc	tgaaagctga	60
gctcacagca	aaacaagcca	ccatgaagct	gtcggtgtgt	ctcctgc	tgcacgcgtgc	120

-continued

cctctgctgc taccaggcca atgcccagtt ctgcccagct cttgtttctg agctgttaga	180
cttccttcttc attagtgaac ctctgttcaa gttaaagtctt gccaaatttg atgcccctcc	240
ggaagctgtt gcagccaaatgtt taggagtgaa gagatgcacg gatcagatgt cccttcagaa	300
acgaaggcctc attgcggaaat tcctgggtgaa aatattgaag aatattgtatgt tggatgttgcacatgt	360
aaaaaacttc atcctggttt ccactgtctt tcaatgacac cctgtatcttc actgcagaat	420
gttaaagggttt caacgtcttg cttaataaa tcacttgctc tac	463

<210> SEQ ID NO 77

<211> LENGTH: 90

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 77

Met Lys Leu Ser Val Cys Leu Leu Leu Val Thr Leu Ala Leu Cys Cys	
1 5 10 15	
Tyr Gln Ala Asn Ala Glu Phe Cys Pro Ala Leu Val Ser Glu Leu Leu	
20 25 30	
Asp Phe Phe Phe Ile Ser Glu Pro Leu Phe Lys Leu Ser Leu Ala Lys	
35 40 45	
Phe Asp Ala Pro Pro Glu Ala Val Ala Ala Lys Leu Gly Val Lys Arg	
50 55 60	
Cys Thr Asp Gln Met Ser Leu Gln Lys Arg Ser Leu Ile Ala Glu Val	
65 70 75 80	
Leu Val Lys Ile Leu Lys Lys Cys Ser Val	
85 90	

We claim:

1. A method for identifying one or more tissue-specific polynucleotides, said method comprising the steps of:

- (a) performing a genetic subtraction to identify a pool of polynucleotides from a tissue of interest;
- (b) performing a DNA microarray analysis to identify a first subset of said pool of polynucleotides of interest wherein each member polynucleotide of said first subset is at least two-fold over-expressed in said tissue of interest as compared to a control tissue; and
- (c) performing a quantitative polymerase chain reaction (PCR) analysis on polynucleotides within said first subset to identify a second subset of polynucleotides that are at least two-fold over-expressed as compared to said control tissue;

wherein a polynucleotide is identified as tissue-specific if it is at least two-fold over-expressed by both microarray and quantitative PCR analyses.

2. The method of claim 1 wherein said genetic subtraction is selected from the group consisting of differential display and cDNA subtraction.

3. A method for identifying a subset of polynucleotides showing complementary tissue-specific expression profiles in a tissue of interest, said method comprising the steps of:

- (a) performing a first expression analysis selected from the group consisting of DNA microarray and quantitative PCR to identify a first polynucleotide that is at least

two-fold over-expressed in a first tissue sample of interest obtained from a first patient but not over-expressed in a second tissue sample of interest as compared to a control tissue; and

(b) performing a second expression analysis selected from the group consisting of DNA microarray and quantitative PCR to identify a second polynucleotide that is at least two-fold over-expressed in a second tissue sample of interest obtained from a second patient but not over-expressed in a first tissue sample of interest as compared to said control tissue;

wherein the first tissue sample and said second tissue sample are of the same tissue type, and wherein over-expression of said first polynucleotide in only said first tissue samples of interest and over-expression of said second polynucleotide in only said second tissue sample of interest indicates complementary tissue-specific expression of said first polynucleotide and said second polynucleotide.

4. A method for determining the presence of a cancer cell in a patient, said method comprising the steps of:

- (a) obtaining a biological sample from said patient;
- (b) contacting said biological sample with a first oligonucleotide that hybridizes to a first polynucleotide said first polynucleotide selected from the group consisting of polynucleotides depicted in SEQ ID NO:73, SEQ ID NO:74 and SEQ ID NO:76;

(c) contacting said biological sample with a second oligonucleotide that hybridizes to a second polynucleotide selected from the group consisting of SEQ ID NO:1, 3, 5-7, 11, 13, 15, 17, 19-24, 30, 32, and 75;

(d) detecting in said sample an amount of a polynucleotide that hybridizes to at least one of said oligonucleotides; and

(e) comparing the amount of the polynucleotide that hybridizes to said oligonucleotide to a predetermined cut-off value, and therefrom determining the presence or absence of a cancer in the patient.

5. A method for determining the presence or absence of a cancer in a patient, said method comprising the steps of:

(a) obtaining a biological sample from said patient;

(b) contacting said biological sample with a first oligonucleotide that hybridizes to a first polynucleotide selected from the group consisting of polynucleotides depicted in SEQ ID NO:73, SEQ ID NO:74 and SEQ ID NO:76;

(c) contacting said biological sample with a second oligonucleotide that hybridizes to a second polynucleotide as depicted in SEQ ID NO:75;

(d) contacting said biological sample with a third oligonucleotide that hybridizes to a third polynucleotide selected from the group consisting of polynucleotides depicted in SEQ ID NO:5, SEQ ID NO:6 and SEQ ID NO:7;

(e) contacting said biological sample with a fourth oligonucleotide that hybridizes to a fourth polynucleotide selected from the group consisting of polynucleotides depicted in SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23 and SEQ ID NO:24;

(f) detecting in said biological sample an amount of a polynucleotide that hybridizes to at least one of said oligonucleotides; and

(g) comparing the amount of polynucleotide that hybridizes to the oligonucleotide to a predetermined cut-off value, and therefrom determining the presence or absence of a cancer in the patient.

6. A method for determining the presence or absence of a cancer in a patient, said method comprising the steps of:

(a) obtaining a biological sample from said patient;

(b) contacting said biological sample with an oligonucleotide that hybridizes to a tissue-specific polynucleotide;

(c) detecting in the sample a level of a polynucleotide that hybridizes to the oligonucleotide; and

(d) comparing the level of polynucleotide that hybridizes to the oligonucleotide with a predetermined cut-off value, and therefrom determining the presence or absence of a cancer in the patient.

7. A method for monitoring the progression of a cancer in a patient, said method comprising the steps of:

(a) obtaining a first biological sample from said patient;

(b) contacting said biological sample with an oligonucleotide that hybridizes to a polynucleotide that encodes a breast tumor protein;

(c) detecting in the sample an amount of said polynucleotide that hybridizes to said oligonucleotide;

(d) repeating steps (b) and (c) using a second biological sample obtained from said patient at a subsequent point in time; and

(e) comparing the amount of polynucleotide detected in step (d) with the amount detected in step (c) and therefrom monitoring the progression of the cancer in the patient.

8. The method any one of claim 6 and claim 7 wherein said polynucleotide encodes a breast tumor protein selected from the group consisting of mammaglobin, lipophilin B, GABA π (B899P), B726P, B511S, B533S, B305D and B311D.

9. A method for detecting the presence of a cancer cell in a patient, said method comprising the steps of:

(a) obtaining a biological sample from said patient;

(b) contacting said biological sample with a first oligonucleotide that hybridizes to a first polynucleotide selected from the group consisting of mammaglobin and lipophilin B;

(c) contacting said biological sample with a second oligonucleotide that hybridizes to a second polynucleotide sequence selected from the group consisting of GABA π (B899P), B726P, B511S, B533S, B305D and B311D;

(d) detecting in said biological sample an amount of a polynucleotide that hybridizes to at least one of the oligonucleotides; and

(e) comparing the amount of polynucleotide that hybridizes to the oligonucleotide to a predetermined cut-off value, and therefrom determining the presence or absence of a cancer in the patient.

10. A method for determining the presence of a cancer cell in a patient, said method comprising the steps of:

(a) obtaining a biological sample from said patient;

(b) contacting said biological sample with a first oligonucleotide that hybridizes to a first polynucleotide selected from the group consisting of a polynucleotide depicted in SEQ ID NO:73 and SEQ ID NO:74 or complement thereof;

(c) contacting said biological sample with a second oligonucleotide that hybridizes to a second polynucleotide depicted in SEQ ID NO:75 or complement thereof;

(d) contacting said biological sample with a third oligonucleotide that hybridizes to a third polynucleotide selected from the group consisting of a polynucleotide depicted in SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:6 and SEQ ID NO:7 or complement thereof,

(e) contacting said biological sample with a fourth oligonucleotide that hybridizes to a fourth polynucleotide selected from the group consisting of a polynucleotide depicted in SEQ ID NO:11 or complement thereof;

- (f) contacting said biological sample with a fifth oligonucleotide that hybridizes to a fifth polynucleotide selected from the group consisting of a polynucleotide depicted in SEQ ID NO:13, 15 and 17 or complement thereof;
- (g) contacting said biological sample with a sixth oligonucleotide that hybridizes to a sixth polynucleotide selected from the group consisting of a polynucleotide depicted in SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23 and SEQ ID NO:24 or complement thereof;
- (h) contacting said biological sample with a seventh oligonucleotide that hybridizes to a seventh polynucleotide depicted in SEQ ID NO:30 or complement thereof;
- (i) contacting said biological sample with an eighth oligonucleotide that hybridizes to an eighth polynucleotide depicted in SEQ ID NO:32 or complement thereof;
- (j) contacting said biological sample with a ninth oligonucleotide that hybridizes to a polynucleotide depicted in SEQ ID NO:76 or complement thereof;
- (k) detecting in said biological sample a hybridized oligonucleotide of any one of steps (b) through (j) and comparing the amount of polynucleotide that hybridizes to the oligonucleotide to a predetermined cut-off value, wherein the presence of a hybridized oligonucleotide in any one of steps (b) through (j) in excess of the pre-determined cut-off value indicates the presence of a cancer cell in the biological sample of said patient.

11. A method for determining the presence of a cancer cell in a patient, said method comprising the steps of:

- (a) obtaining a biological sample from said patient;
- (b) contacting said biological sample with a first oligonucleotide and a second oligonucleotide;
 - i. wherein said first oligonucleotide and said second oligonucleotide hybridize to a first polynucleotide and a second polynucleotide, respectively;
 - ii. wherein said first polynucleotide and said second polynucleotide are selected from the group consisting of polynucleotides depicted in SEQ ID NO:73, SEQ ID NO:74, SEQ ID NO:75, SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:11, SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:30, SEQ ID NO:32, and SEQ ID NO:76; and
 - iii. wherein said first polynucleotide is unrelated in nucleotide sequence to said second polynucleotide;
- (c) detecting in said biological sample said hybridized first oligonucleotide and said hybridized second oligonucleotide; and
- (d) comparing the amount of said hybridized first oligonucleotide and said hybridized second hybridized oligonucleotide to a predetermined cut-off value; wherein an amount of said hybridized first oligonucleotide or said hybridized second oligonucleotide in excess of the

predetermined cut-off value indicates the presence of a cancer cell in the biological sample of said patient.

12. A method for determining the presence or absence of a cancer cell in a patient, said method comprising the steps of:

- (a) obtaining a biological sample from said patient;
- (b) contacting said biological sample with a first oligonucleotide and a second oligonucleotide;
 - i. wherein said first oligonucleotide and said second oligonucleotide hybridize to a first polynucleotide and a second polynucleotide, respectively;
 - ii. wherein said first polynucleotide and said second polynucleotide are both tissue-specific polynucleotides of the cancer cell to be detected; and
 - iii. wherein said first polynucleotide is unrelated in nucleotide sequence to said second polynucleotide;
- (c) detecting in said biological sample said first hybridized oligonucleotide and said second hybridized oligonucleotide; and
- (d) comparing the amount of polynucleotide that hybridizes to the oligonucleotide to a predetermined cut-off value, wherein the presence of a hybridized first oligonucleotide or a hybridized second oligonucleotide in excess of the pre-determined cut-off value indicates the presence of a cancer cell in the biological sample of said patient.

13. A method for detecting the presence of a cancer cell in a patient, said method comprising the steps of:

- (a) obtaining a biological sample from said patient;
- (b) contacting said biological sample with a first oligonucleotide pair said first pair comprising a first oligonucleotide and a second oligonucleotide wherein said first oligonucleotide and said second oligonucleotide hybridize to a first polynucleotide and the complement thereof, respectively;
- (c) contacting said biological sample with a second oligonucleotide pair said second pair comprising a third oligonucleotide and a fourth oligonucleotide wherein said third and said fourth oligonucleotide hybridize to a second polynucleotide and the complement thereof, respectively, and wherein said first polynucleotide is unrelated in nucleotide sequence to said second polynucleotide;
- (d) amplifying said first polynucleotide and said second polynucleotide; and
- (e) detecting said amplified first polynucleotide and said amplified second polynucleotide; wherein the presence of said amplified first polynucleotide or said amplified second polynucleotide indicates the presence of a cancer cell in said patient.

14. The method of any one of claims 4-7 and 9-13 wherein said biological sample is selected from the group consisting of blood, serum, lymph node, bone marrow, sputum, urine and tumor biopsy sample.

15. The method of claim 14 wherein said biological sample is selected from the group consisting of blood, a lymph node and bone marrow.

16. The method of claim 15 wherein said lymph node is a sentinel lymph node.

17. The method of any one of claims **4-7** and **9-13** wherein said cancer is selected from the group consisting of prostate cancer, breast cancer, colon cancer, ovarian cancer, lung cancer head & neck cancer, lymphoma, leukemia, melanoma, liver cancer, gastric cancer, kidney cancer, bladder cancer, pancreatic cancer and endometrial cancer.

18. The method of any one of claims **12** and **13** wherein said first polynucleotide and said second polynucleotide are selected from the group consisting of gammaglobin, lipophilin B, GABA α (B899P), B726P, B511S, B533S, B305D and B311D.

19. The method of any one of claims **12** and **13** wherein said first polynucleotide and said second polynucleotide are selected from the group consisting of polynucleotide depicted in SEQ ID NO:73, SEQ ID NO:74, SEQ ID NO:75, SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:11, SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:30, SEQ ID NO:32, and SEQ ID NO:76.

20. The method of any one of claims **12** and **13** wherein said oligonucleotides are selected from the group consisting of oligonucleotides depicted in SEQ ID NO:33-35 and 63-72.

21. The method of any one of claims **12** and **13** wherein the step of detection of said first amplified polynucleotide and said second polynucleotide comprises a step selected from the group consisting of detecting a radiolabel and detecting a fluorophore.

22. The method of any one of claims **4-7** and **9-13** wherein said step of detection comprises a step of fractionation.

23. The method of any one of claims **12** and **13** wherein said first and said oligonucleotides are intron spanning oligonucleotides.

24. The method of claim 23 wherein said intron spanning oligonucleotides are selected from the group consisting of oligonucleotides depicted in SEQ ID NO:36-62.

25. The method of claim 13 wherein detection of said amplified first or said second polynucleotide comprises contacting said amplified first or said second polynucleotide with a labeled oligonucleotide probe that hybridizes, under moderately stringent conditions, to said first or said second polynucleotide.

26. The method of claim 13 wherein said labeled oligonucleotide probe comprises a detectable moiety selected from the group consisting of a radiolabel and a fluorophore.

27. The method of any one of claims **4-7** and **9-13** further comprising a step of enriching said cancer cell from said biological sample prior to hybridizing said oligonucleotide primer(s).

28. The method of claim 27 wherein said step of enriching said cancer cell from said biological sample is achieved by a methodology selected from the group consisting of cell capture and cell depletion.

29. The method of claim 28 wherein cell capture is achieved by immunocapture, said immunocapture comprising the steps of:

- adsorbing an antibody to the surface of said cancer cells; and
- separating said antibody adsorbed cancer cells from the remainder of said biological sample.

30. The method of claim 29 wherein said antibody is directed to an antigen selected from the group consisting of CD2, CD3, CD4, CD5, CD8, CD10, CD11b, CD14, CD15, CD16, CD19, CD20, CD24, CD25, CD29, CD33, CD34, CD36, CD38, CD41, CD45, CD45RA, CD45RO, CD56, CD66B, CD66e, HLA-DR, IgE and TCR $\alpha\beta$.

31. The method of claim 29 wherein said antibody is directed to a breast tumor antigen.

32. The method of any one of claims **29-31** wherein said antibody is a monoclonal antibody.

33. The method of claim 29 wherein said antibody is conjugated to magnetic beads.

34. The method of claim 29 wherein said antibody is formulated in a tetrameric antibody complex.

35. The method of claim 28 wherein cell depletion is achieved by a method comprising the steps of:

- cross-linking red cells and white cells, and
- fractionating said cross-linked red and white cells from the remainder of said biological sample.

36. The method of claim 13 wherein said step of amplifying is achieved by a polynucleotide amplification methodology selected from the group consisting of reverse transcription polymerase chain reaction (RT-PCR), inverse PCR, RACE, ligase chain reaction (LCR), Qbeta Replicase, isothermal amplification, strand displacement amplification (SDA), rolling chain reaction (RCR), cyclic probe reaction (CPR), transcription-based amplification systems (TAS), nucleic acid sequence based amplification (NASBA) and 3SR.

37. A composition for detecting a cancer cell in a biological sample of a patient, said composition comprising:

- a first oligonucleotide; and
- a second oligonucleotide; wherein said first oligonucleotide and said second oligonucleotide hybridize to a first polynucleotide and to a second polynucleotide, respectively; wherein said first polynucleotide is unrelated in nucleotide sequence from said second polynucleotide; and wherein said first polynucleotide and said second polynucleotide are tissue-specific polynucleotides of the cancer cell to be detected.

38. The composition of claim 37 wherein said first polynucleotide and said second polynucleotide are complementary tissue-specific polynucleotides of the tissue-type of said cancer cell.

39. The composition of any one of claim 37 and claim 38 wherein said first polynucleotide and said second polynucleotide are selected from the group consisting of the polynucleotides depicted in SEQ ID NO:73, SEQ ID NO:74, SEQ ID NO:75, SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:11, SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:30, SEQ ID NO:32, and SEQ ID NO:76.

40. The composition of any one of claim 37 and claim 38 wherein said oligonucleotides are selected from the group consisting of oligonucleotides as disclosed in SEQ ID NO:33-72.

41. A composition for detecting a cancer cell in a biological sample of a patient, said composition comprising:

- (a) a first oligonucleotide pair; and
- (b) a second oligonucleotide pair;

wherein said first oligonucleotide pair and said second oligonucleotide pair hybridize to a first polynucleotide (or complement thereof) and to a second polynucleotide (or complement thereof), respectively; wherein said first polynucleotide is unrelated in nucleotide sequence from said second polynucleotide; and wherein said first polynucleotide and said second polynucleotide are tissue-specific polynucleotides of the cancer cell to be detected.

42. The composition of claim 41 wherein said first polynucleotide and said second polynucleotide are complementary tissue-specific polynucleotides of the tissue-type of said cancer cell.

43. The composition of any one of claim 41 and claim 42 wherein said first polynucleotide and said second polynucleotide are selected from the group consisting of the polynucleotides depicted in SEQ ID NO:73, SEQ ID NO:74,

SEQ ID NO:75, SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:11, SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:30, SEQ ID NO:32, and SEQ ID NO:76.

44. The composition of any one of claim 41 and claim 42 wherein said oligonucleotides are selected from the group consisting of oligonucleotides as disclosed in SEQ ID NO:33-72.

45. A composition comprising an oligonucleotide primer or probe of between 15 and 100 nucleotides that comprises an oligonucleotide selected from the group consisting of oligonucleotides depicted in SEQ ID NO:33-72.

46. The composition of claim 45 comprising an oligonucleotide primer or probe selected from the group consisting of oligonucleotides depicted in SEQ ID NO:33-72.

* * * * *

专利名称(译)	用于检测和监测乳腺癌的方法，组合物和试剂盒		
公开(公告)号	US20020009738A1	公开(公告)日	2002-01-24
申请号	US09/825301	申请日	2001-04-02
[标]申请(专利权)人(译)	HOUGHTON RAYMOND 大号 DILLON DAVIN C MOLESH DAVID 徐JIANGCHUN ZEHENTNER BARBARA PERSING David H制作		
申请(专利权)人(译)	HOUGHTON RAYMOND L. 狄龙DAVIN C. MOLESH DAVID 徐JIANGCHUN ZEHENTNER BARBARA PERSING DAVID H.		
当前申请(专利权)人(译)	HOUGHTON RAYMOND L. 狄龙DAVIN C. MOLESH DAVID 徐JIANGCHUN ZEHENTNER BARBARA PERSING DAVID H.		
[标]发明人	HOUGHTON RAYMOND L DILLON DAVIN C MOLESH DAVID XU JIANGCHUN ZEHENTNER BARBARA PERSING DAVID H		
发明人	HOUGHTON, RAYMOND L. DILLON, DAVIN C. MOLESH, DAVID XU, JIANGCHUN ZEHENTNER, BARBARA PERSING, DAVID H.		
IPC分类号	G01N33/53 C12N5/08 C12N15/09 C12Q1/68 G01N33/566 G01N33/574 G01N33/58		
CPC分类号	C12Q1/6809 C12Q1/6844 C12Q1/6851 C12Q1/6886 C12Q2565/501 C12Q2545/114 C12Q2531/113 C12Q2600/16		
优先权	60/194241 2000-04-03 US 60/221300 2000-07-27 US 60/219862 2000-07-20 US 60/256592 2000-12-18 US		
外部链接	Espacenet USPTO		

摘要(译)

公开了用于治疗和诊断癌症(例如乳腺癌)的组合物和方法。组合物可包含一种或多种乳腺肿瘤蛋白，其免疫原性部分或编码这些部分的多核苷酸。或者，治疗组合物可包含表达乳腺肿瘤蛋白的抗原呈递细胞，或对表达这种蛋白的细胞特异的T细胞。此类组合物可用于例如预防和治疗诸如乳腺癌的疾病。还提供了基于在样品中检测乳腺肿瘤蛋白或编码这种蛋白的mRNA的诊断方法。




Fig. 1