



US007897361B2

(12) **United States Patent**
Westbrook et al.

(10) **Patent No.:** **US 7,897,361 B2**
(45) **Date of Patent:** **Mar. 1, 2011**

(54) **METHODS OF DIAGNOSING ALZHEIMER'S DISEASE**

G01N 33/566 (2006.01)

G01N 33/567 (2006.01)

(52) **U.S. Cl.** **435/7.92**; 435/7.1; 435/7.9; 436/501;
436/503

(58) **Field of Classification Search** None
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

2002/0174447 A1 11/2002 Greenspan et al.
2003/0131364 A1 7/2003 Duff
2003/0211622 A1 11/2003 Roberts

OTHER PUBLICATIONS

Schenk et al. J. Mol. Chem., 1995, vol. 38, No. 21, pp. 4141-4154.*
Tillemann, Kelly et al., "Differential expression of brain proteins in glycogen synthase kinase-3beta transgenic mice: A proteomics point of view", Proteomics, 2(1): 94-104 (2002).
Andreasen, Niels et al., "Cerebrospinal fluid beta-amyloid(1-42) in Alzheimer disease: Differences between early- and late-onset Alzheimer disease and stability during the course of disease", Archives of Neurology, 56(6): 673-680 (1999).
Kanai et al., "Longitudinal Study of Cerebrospinal Fluid Levels of Tau, ABeta1-40, and ABeta1-42(43) in Alzheimer's Disease: A Study in Japan", Annals of Neurology, 44(1): 17-26 (1998).

* cited by examiner

Primary Examiner — Olga N Chernyshev

(74) *Attorney, Agent, or Firm* — Dann, Dorfman, Herrell and Skillman, P.C.; Patrick J. Hagan

(57) **ABSTRACT**

Methods and compositions relating to Alzheimer's disease are provided. Specifically, proteins that are differentially expressed in the Alzheimer's disease state relative to their expression in the normal state are provided. Proteins associated with Alzheimer's disease are identified and described. Methods of diagnosis of Alzheimer's disease using the differentially expressed proteins are also provided, as are methods for the identification and therapeutic use of compounds for the prevention and treatment of Alzheimer's disease.

8 Claims, 20 Drawing Sheets

(76) Inventors: **Jules Westbrook**, Dublin (IE); **Helen Byers**, Cobham Surrey (GB); **Malcolm Ward**, Cobham Surrey (GB); **Simon Lovestone**, London (GB); **Abdul Hye**, London (GB); **Stephen Lynham**, London (GB); **Richard Joubert**, Niedernhausen (DE); **Petra Prefot**, Wiesbaden (DE); **Karsten Kuhn**, Hofheim (DE); **Christian Baumann**, Offenbad (DE); **Juergen Schaefer**, Lauterbach (DE); **Thorsten Prinz**, Hofheim (DE); **Stefan Kienle**, Frankfurt (DE)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

(21) Appl. No.: **11/664,076**

(22) PCT Filed: **Sep. 29, 2005**

(86) PCT No.: **PCT/GB2005/003756**

§ 371 (c)(1),

(2), (4) Date: **Oct. 16, 2007**

(87) PCT Pub. No.: **WO2006/035237**

PCT Pub. Date: **Apr. 6, 2006**

(65) **Prior Publication Data**

US 2008/0070995 A1 Mar. 20, 2008

(30) **Foreign Application Priority Data**

Sep. 29, 2004 (GB) 0421639.6

(51) **Int. Cl.**

C12Q 1/68 (2006.01)

G01N 33/53 (2006.01)

Figure 1




Figure 2




Figure 3





Figure 4






Figure 5

Spot No.	Rank	p Value	Fold Difference	State Change	Protein I.D.	Accession No.	Search Log No.
196	1	0.00030199	1.78	↑ AD	Desmoplakin (DP) (250/210 kDa paraneoplastic pemphigus antigen) Ig kappa chain C region Ig kappa chain V-II region TEW Serum amyloid P-component precursor (SAP) (9.5S alpha-1-glycoprotein)	P15924 P01834 P01617 P02743	7495 7542 7542 7951
171	2	0.001255545	2.11	↑ AD	Ig kappa chain C region Serum albumin precursor Galectin-7 (Gal-7) (HKL-14) (P17) (p53-induced protein 1)	P01834 P02768 P47929	5623 7954 5623
2 (old)	3	0.001447694	13.75	↑ AD	Complement factor H precursor (H factor 1) Serum albumin precursor Alpha-2-macroglobulin precursor (Alpha-2-M) Ceruloplasmin precursor (EC 1.16.3.1) (Ferroxidase)	P08603 P02768 P01023 P00450	6672 6672 6672 6672
184	4	0.005360087	2.43	↑ AD	Ig lambda chain C regions Ig lambda chain V-III region LOI Serum albumin precursor Complement factor H-related protein 2 precursor (FHR-2)	P01842 P80748 P02768 P36980	7818 7818 7818 7818
177 (old)	5	0.005382883	1.92	↑ AD	Ig lambda chain C regions Serum albumin precursor Ig lambda chain V-III region LOI Ig kappa chain C region	P01842 P02768 P80748 P01834	5627 7955 7955 7955
4	6	0.005985336	8.83	↑ AD	Alpha-2-macroglobulin precursor (Alpha-2-M)	P01023	7827
170	7	0.01167553	1.98	↑ AD			
13	8	0.015500401	4.23	↓ AD	Inter-alpha-trypsin inhibitor heavy chain H4 precursor (ITI heavy chain H4) Ceruloplasmin precursor (EC 1.16.3.1) (Ferroxidase)	Q14624 P00450	7829 7829
165 (old)	9	0.018305158	1.58	↓ AD	Serum albumin precursor	P02768	5625
164	10	0.020647469	2.03	↓ AD	Complement C4 precursor [Contains: C4a anaphylatoxin; C4b]	P01028	7821

Figure 6

14 (old)	11	0.025004429	10.82	↓ AD	Ig gamma-1 chain C region	P01857	7821
					Serum albumin precursor	P02768	6227
					Histone H2B.a/g/h/k/l (H2B.1 A)	P62807	6227
					(H2B/a) (H2B/g) (H2B/h) (H2B/k) (H2B/l)		
126	12	0.028979402	1.6	↓ AD	CD5 antigen-like precursor (SP-alpha) (CT-2) (IgM-associated peptide)	O43866	7493
					Serum albumin precursor	P02768	7952
					Ig mu chain C region	P01871	7952
176	13	0.029106689	1.75	↑ AD	Ig lambda chain C regions	P01842	7816
					Serum albumin precursor	P02768	7816
					Ig lambda chain V-III region LOI	P80748	7816
123	14	0.031441346	1.36	↑ AD	Serum albumin precursor	P02768	7462
1	15	0.034723104	3.32	↑ AD	Alpha-2-macroglobulin precursor (Alpha-2-M)	P01023	7823
					Ig alpha-1 chain C region	P01876	7823

Figure 6 (continued)

Band No.	Protein I.D.	Species	Accession No.	Gal	MW (Da)	pI	No. Peptides Matched	Percentage Coverage	Error (ppm)	Search Log No.	Peptide Matched
SPI_1C	Haptoglobin precursor	Human	P00738	19300	45177	6.13	5	9%	16	5288	
SPI_2C	Transferrin	Human	g339685	18200	12835	5.33	8	87%	223	5275	
	Serum albumin precursor	Human	P02768	18200	69248	5.82	3	6%	236	5274	
	Complement C4 precursor	Human	P01028	18200	192650	6.65	1	0%	236	5274	GLEELQFSLGSK
	Fibrinogen alpha1alpha-E chain precursor	Human	P02671	18200	94914	5.7	1	3%	224	5274	ETWISDGDQPEAMDGLTSLGIGTLDGFR
SPI_3C	Clain A, Transferrin	Human	g343285	14900	13753	5.35	10	92%	102	5251	
	Apolipoprotein A-IV precursor (Apo-AIV)	Human	P06727	14900	43343	5.28	2	10%	100	5247	
	Serum albumin precursor	Human	P02768	14900	69321	5.92	1	1%	109	5247	
SPI_4C	Transferrin precursor	Human	P02766	14200	13877	5.52	5	60%	170	5260	
	Hemoglobin beta chain	Human	P02023	14200	13857	6.81	4	28%	12	5260	
	Serum albumin precursor	Human	P02768	14200	69321	5.92	4	7%	7	5260	
SPI_5C	Haptoglobin-related protein precursor	Human	P00739	12600	36983	6.42	5	10%	102	5294	
	Transferrin precursor	Human	P02766	12600	13877	5.52	2	16%	129	5294	
	Serum albumin precursor	Human	P02768	12600	69321	5.92	4	10%	120	5294	
	Apolipoprotein C-III precursor (Apo-CIII)	Human	P02656	12600	10845	5.23	1	16%	122	5294	DALSSVQESQVAQQAR
	Hemoglobin alpha	Human	P01922	12600	15227	9.84	2	17%	118	5294	
	Hemoglobin beta chain	Human	P02023	12600	13857	6.81	1	15%	131	5294	SAVTALWGKYNVDENGEALGE
SPI_6C	Serum albumin precursor	Human	P02768	11600	69321	5.92	6	11%	244	5280	
	Apolipoprotein C-III precursor (Apo-CIII)	Human	P02656	11600	10845	5.23	3	37%	244	5280	
	Haptoglobin precursor	Human	P00738	11600	45177	6.13	2	6%	233	5280	
	Vitronectin precursor (Serum spreading factor) (S-protein)	Human	P04004	11600	54271	5.55	1	3%	249	5280	STAQYWLGGPAPCHL

Figure 7

No	IPI Accession no	SWISS- PROT Accession no	Name	No of matched peptides	regulation (control / disease)	CV (%)
1	IPI00166866	P01876	MGC27165 PROTEIN	2	0,38	7
2	IPI00336074	P01876	IG ALPHA-1 CHAIN C REGION	2	0,35	4
3	IPI00423461	P01842	HYPOTHETICAL PROTEIN DKFZP686C0222 0 (FRAGMENT)	2	0,35	24
4	IPI00431645	P00738	HAPTOGLOBIN PRECURSOR	1	0,33	-
5	IPI00478493	P00738	HAPTOGLOBIN PRECURSOR	1	0,34	-

Figure 8

1 MFLKAVVLT~~L~~ ALVAVAGARA EVSADQVATV MWDYFSQLSN NAKEAVEHLQ
51 KSELTQQLNA LFQDKLGEVN TYAGDLQKKL VPFATELHER LAKDSEKLKE
101 EIGKELEELR ARLLPHANEV SQKIGDNLRE LQQRLEPYAD QLRTQVNTQA
151 EQLRRQLTPY AQRMERVLRE NADSLQASLR PHADELKAKI DQNVEELKGR
201 LTPYADEFKV KIDQTVEELR RSLAPYAQDT QEKLNHQLEG LTFQMKKNAE
251 ELKARISASA EELRQRLAPL AEDVRGNLKG NTEGLQKSLA ELGGHLDQQV
301 EEFRRRVEPY GENFNKALVQ QMEQLRQKLG PHAGDVEGHL SFLEKDLRDK
351 VNSFFSTFKE KESQDKTSL PELEQQQEEQ QEQQQEQVQM LAPLES

Figure 9

1 MRLLWGLIWA SSFFTLISLQK PRLLLFSPSV VHLGVPLSVG VQLQDVPRGQ
51 VVKGSVFLRN PSRNNVPCSP KVDFTLSSER DFALLSLQVP LKDAKSCGLH
101 QLLRGPEVQL VAHSPWLKDS LSRTTNIQGI NLLFS.SRRGH LFLQTDQPIY
151 NPGQVRVYRV FALDQKMRPS TDTITVMVEN SHGLRVRKKE VYMPSSIFQD
201 DFVIPDISEP GTWKISARFS DGLSNSSTQ FEVKKYVLPN FEVKITPGKP
251 YILTVPGHLD EMQLDIQARY IYGKPVQGVA YVRFGLLDED GKKTFFRGLE
301 SQTCLVNGQS HISLSKAEFQ DALEKLNMG I TDLQGLRLYV AAAIIESPGG
351 EMEEAELTSW YFVSSPFSLD LSKTKRHLVP GAPFLQLQALV REMSGSPASG
401 IPVKVSATVS SPGSVPEVQD IQQNTDGSQ VSIPI IIPQT ISELQLSVSA
451 GSPHPAIARL TVAAPPSGGP GFLSIERPDS RPPRVGDTLN LNLRAVGSGA
501 TFSHYYMIL SRGQIVFMNR EPKRTLTSVS VFVDHHLAPS FYFVAFYYHG
551 DHPVANSLRV DVQAGACEGK LELSVDGAKQ YRNGESVKLH LETDSLALVA
601 LGALDTALYA AGSKSHKPLN MGKVFEAMNS YDLGC GPGGG DSALQVFQAA
651 GLAFSDGDQW TLSRKRLSCP KEKTTRKKRN VNFQK AINEK LGQYASPTAK
701 RCCQDGVTRL PMMRSCEQRA ARVQQPDCRE PFLSCCQFAE SLRKKSRDKG
751 QAGLQRALEI LQEEDLIDED DIPVRSFFPE NWLWRVETVD RFQILTLWLP
801 DSLTTWEIHG LSLSKTKGLC VATPVQLRVF REFHLHLRLP MSVRRFEQLE
851 LRPVLYNYLD KNLTVSVHVS PVEGLCLAGG GGLAQQVLVP AGSARPVAFS
901 VVPTAAAVS LKVVARGSFE FPGDAVSKV LQIEKEGAIH REELVYELNP
951 LDHRGRTLEI PGNSDPNMIP DGDFNSYVRV TASDPLDTLG SEGALSPGGV
1001 ASLLRLPRGC GEQTMIIYLA TLAASRYLDK TEQWS'TLPPE TKDHAVDLIQ
1051 KGYMRIQQFR KADGSYAAWL SRDSSTWLTA FVLKVLSLAQ EQVGG SPEKL
1101 QETSNWLLSQ QQADGSFQDP CPVLDRSMQG GLVGNDETVA LTAFVTIALH
1151 HGLAVFQDEG AEPLKQVEA SISKANSFLG EKASAGLLGA HAAAITAYAL
1201 SLTKAPVDLL GVAHNNLMAM AQETGDNLYW GSVTG.SQSNA VSPTPAPRNP
1251 SDPMPQAPAL WIETTAYALL HLLLHEGKAE MADQASAWLT RQGSFQGGFR
1301 STQDTVIALD ALSAYWIAH TTEERGLNVT LSSTGRNGFK SHALQLNNRQ
1351 IRGLEEBELQF SLGSKINVKV GGNSKGT LKV LRTYNVLD MK NTTCQDLQIE
1401 VTVKGHVEYT MEANEDYEDY EYDELPKDD PDAPLQPVTP LQLFEGRRNR
1451 RRREAPKVVE EQESRVHYTV CIWRNGKVGL SGMAIADVT L LSGFHALRAD
1501 LEKLTSLSDR YVSHFETEGP HVLLYFDSVP TSREC'VGFEA VQEVVGLVQ
1551 PASATLYDYY NPERRCSV FY GAPSKSRLLA TLCSAEVCQC AEGKCPRQRR
1601 ALERGLQDED GYRMKFACY Y PRVEYGFQVK VLRED.SRAAF RLFETKITQV
1651 LHFTKDVKAA ANQMRNFLVR ASCRLRLPEG KEYLIMGLDG ATYDLEGHPQ
1701 YLLDSNSWIE EMPSERLCRS TRQRAACAQL NDFLQ'EYGTQ GCQV

Figure 10




Figure 11

Spot no.	Protein Name	Acc. No.	Norm. Vol Control	CV (%)	Norm. Vol Disease	CV (%)	Expressio n ratio	T-test (p)	Detection ratio	Theoretical Mr	pI	Cover -age (%)
174	alpha-2-macroglobulin precursor	P01023	0,36604	52	0,14510	57	0,40	2,50852E-06	28/21	160796 Da	5,95	13,8
178	alpha-2-macroglobulin precursor	P01023	0,29348	53	0,12475	57	0,43	1,25521E-05	27/20	160796 Da	5,95	11,7
232	Inter-alpha-trypsin inhibitor heavy chain H4 precursor	Q14624	0,32468	81	0,14481	64	0,45	0,001793752	28/27	103358 Da	6,51	25,6
712	Complement C3 precursor	P01024	0,81225	74	0,36446	62	0,45	0,001371446	28/24	184967 Da	6,00	14,9
712	Clusterin precursor	P10909	0,81225	74	0,36446	62	0,45	0,001371446	28/24	50062 Da	5,89	22,9
713	Complement C3 precursor	P01024	3,45803	67	1,44590	63	0,42	9,67927E-05	29/29	184967 Da	6,00	16,9
652	Complement C4 precursor	P01028	0,18067	122	0,39844	69	2,21	0,003595593	24/25	192771 Da	6,60	5,9
675	Actin cytoplasmic 2 (Gamma/beta actin)	P63261	0,21268	89	0,45443	74	2,14	0,002107165	25/28	41793 Da	5,31	49,1
702	Haptoglobin precursor	P00738	0,09210	86	0,40084	115	4,35	0,002920373	20/25	43349 Da	6,13	21,2
703	Haptoglobin precursor	P00738	1,54479	95	4,64500	76	3,01	0,000171685	24/28	43349 Da	6,13	23,9
706	Haptoglobin precursor	P00738	1,00814	112	3,26743	89	3,24	0,000607583	21/28	43349 Da	6,13	23,6
832	Complement C4 precursor	P01028	0,24743	126	0,61914	88	2,50	0,0030005256	28/28	192771 Da	6,60	4,5

Figure 12

1 MGKNKLLHPS LVLALLVLLP TDASVSGKPQ YMLVPSLLH TETTEKGCVL LSYLNETVTV
61 SASLESVRGN RSLETDLEAE NDVLHCVAFA VPKSSSNEEV MFLTVQVKGP TQEFKKRTTV
121 MVKNEDSLVF VQTDKSIYKP GQTVKFRVVS MDENFHPLNE LIPLVYIQDP KGNRIAQWQS
181 FQLEGGLKQF SFPLSSEPFQ GSYKVVQKK SGGRTEHPFT VEEFVLPKFE VQVTVPKIIT
241 ILEEMNVSV CGLYTYGKPV PGHVTVSICR KYSDASDCHG EDSQAFCEKF SGQLNSHGCF
301 YQQVKTTKVFQ LKRKEYEMKL HTEAQIQEEG TVVELTGRQS SEITRTITKL SFVKVDSHFR
361 QGIPFFGQVR LVDGKGVPIP NKVIFIRGNE ANYYSNATTD EHGLVQFSIN TTNVMGTSIT
421 VRVNYKDRSP CYGYQWVSEE HEEAHHTAYL VFSPSKSEFH LEPMSHELPC GHTQTVQAHY
481 ILNGGTLLGL KKLSTFYLLIM AKGGIVRTGT HGLLVKQEDM KGHFSISIPV KSDIAPVARL
541 LIYAVLPTGD VIGDSAKYDV ENCLANKVDL SFSPSQSLPA SHAHLRVTAA PQSVCALRAV
601 DQSVLLMKPD AELSASSVYN LLPEKDLTGF PGPLNDQDDE DCINRHNVI NGITYTPVSS
661 TNEKDMYSFL EDMGLKAFTN SKIRKPKMCP QLQQYEMHGP EGLRVGFYES DVMGRGHARL
721 VHVEEPHTET VRKYFPETWI WDLVVNSAG VAEVGVTVPD TITEWKAGAF CLSEDAGLGI
781 SSTASLRAFQ PFFVELTMPY SVIRGEAFTL KATVLNLYPK CIRVSVQLEA SPAFLAVPVE
841 KEQAPHCICA NGRQTVSWAV TPKSLGNVNF TVSAEALESQ ELCGTEVPSV PEHGRKDTVI
901 KPLLVEPEGL EKETTFFNSLL CPSSGGEVSEE LSLKLPPNVV EESARASVSV LGDILGSAMQ
961 NTQNLQMPY GCGEQNMVLF APNIYVLDYL NETQQLTPEV KSKAIGYLT GYQRQLNYKH
1021 YDGSYSTFGE RYGRNQGNTW LTAFVLKTFA QARAYIFIDE AHITQALIWL SQRQKDNCGF
1081 RSSGSLNNA IKGGVEDEV T LSAYITIALL EIPLTVTHPV VRNALFCLES AWKTAQEGDH
1141 GSHVYTKALL AYAFALAGNQ DKRKEVLKSL NEEAVKDNS VHWERPQKPK APVGHFYEPQ
1201 APSAEVEMTS YVLLAYLTAQ PAPTSEDITS ATNIVKWITK QQNAQGGFSS TQDTVVALHA
1261 LSKYGAATFT RTGKAAQVTI QSSGTFSSKF QVDNNRLLL QQVSLPELPG EYSMKVTGEG
1321 CVYLQTSKY NILPEKEEFP FALGVQTLPO TCDEPKAHTS FQISLSVSYT GSRASASMAI
1381 VDKMVSQFI PLKPTVKMLE RSNHVSRTVE SSNHVLIYLD KVSNTLSLF FTVLQDVPVR
1441 DLKPAIVKVY DYYETDEFAI AEYNAPCSKD LGNA

Figure 13

1 MKPPRPVRTC SKVLVLLSLL AIHQTTTAEK NGIDIYSLTV DSRVSSRFAH **TVVTSRVVNR**
61 **ANTVQEATFQ** **MELPKKAFIT** NFSMNIDGMT YPGIIKEKAE AQAQYSAAVA KGKSAGLVKA
121 TGRNMEQFQV SVSVAPNAKI TFELVYEELL KRRLGVYELL **LKVRPQQLVK** HLQMDIHIFE
181 PQGISFLETE STFMTNQLVD ALTTWQNKTK **AHIRFKPTLS** **QQQKSPEQQE** **TVLDGNLIIR**
241 YDVDRAISGG **SIQIENGYFV** **HYFAPEGLTT** MPKNVVFVID KSGSMSGRKI QQTREALIKI
301 LDDLSPRDQF NLIVFSTEAT QWRPSLVPAS AENVNKARSE AAGIQALGGT NINDAMLMAY
361 QLLDSSNQEE RLPEGSVSLI ILLTDGDPTV **GETNPRSIQN** NVREAVSGRY SLFCLGFGFD
421 VSYAFLEKLA LDNGGLARRI HEDSDSALQL QDFYQEVANP LLTAVTFEYP SNAVEEVTQN
481 NFRLLFKGSE MIVAGKLQDR **GPDVLTATVS** **GKLPTQNITF** QTESSVAEQE AEFQSPKYIF
541 **HNFMERLWAY** LTIQQLLEQT VSASDADQQA LRNQALNLSL AYSFVTPLTS MVVTKPDDQE
601 QSQVAEKPME GESRNRNVHS **GSTFFKYYLQ** GAKIPKPEAS **FSPRRGWNRQ** AGAAGSRMNF
661 **RPGVLSSRQL** **GLPGPPDVPD** **HAAYHPFRR** AILPASAPPA TSNPDPAVSR VMNMKIEETT
721 MTTQTPAPIQ APSAILPLPG QSVRLCVDV RHRQGPVNLL SDPEQGVEVT GQYEREKAGF
781 SWIEVTFKNP LVVWHASPEH VVTRNRSS AYKWKETLFS VMPGLKMTMD KTGLLLLLSDP
841 DKVTIGLLFW DGRGGLRLL LRDTDRESSH VGGTLGQFYQ EVLWGSPAAS DDGRRTLVRQ
901 GNDHSATRER RLDYQEGPPG VEISCWSVEL

Figure 14

1 MGPTSGPSLL LLLLTHLPLA LGSPMYSIIT PNILRLESEE TMVLEAHDAQ GDVPVTVTVH
61 DFPGKKLVLS SEKTVLTPAT NHMGNVTFIT PANREFKSEK GRNKFVTVQA TFGTQVVEKV
121 VLVSLQSGYL FIQTDKTIYT PGSTVLYRIF TVNHKLLPVG RTVMVNIENP EGIPVKQDSL
181 SSQNQLGVLP LSWDIPELVN MGQWKIRAYY ENSPQQVFST EFEVKEYVLP SFEVIVEPTE
241 KFYIYNEKG LEVTITARFL YGKKVEGTAF VIFGIQDGEQ RISLPESLKR IPIEDGSGEV
301 VLSRKVLLDG VQNLRAEDLV GKSLYVSATV ILHSGSDMVQ AERSGIPIVT SPYQIHFTKT
361 PKYFKPGMPF DLMVFVTNPD GSPAYRVPVA VQGEDTVQSL TQGDGVAKLS INTHPSQKPL
421 SITVRTKKQE LSEAEQATRT MQALPYSTVG NSNNYLHLSV LRTELRPGET LNVNELLRMD
481 RAHEAKIRYY TYLIMNKGRL LKAGRQVREP GQDLVVLPLS ITTDFIPSEF LVAYYTLIGA
541 SGQREVADS VWVDVKDSCV GSLVVKSGQS EDRQVPVPGQ MTLKIEGDHG ARVVLVAVDK
601 GVFVLNKKNK LTQSKIWDVV EKADIGCTPG SGKDYAGVFS DAGLTFTSSS GQQTAAQRAEL
661 QCPQPAARRR RSVQLTEKRM DKVGKYPKEL RKCCEDGMRE NPMRFSCQRR TRFISLGEAC
721 **KKVFLDCCNY** ITELRRQHAR ASHLGLARSN LDEDIIAEEN IVSRSEFPES WLWNVEDLKE
781 PPKNGISTKL MNIFLKDSIT TWEILAVSMS DKKGICVADP FEVTVMQDFE IDLRLPYSVV
841 RNEQVEIRAV LYNRYQNQEL KVRVELLHNP AFCSLATTKR RHQQTVTIPP KSSLSVPYVI
901 VPLKTGLQEV EVKAAVYHHE ISDGVKSLK VVEGIRMNK TVAVRTLDPE RLGREGV**QKE**
961 **DIPPADLSDQ** **VPDTESETRI** **LLQGTTPVAQM** **TEDAVDAERL** KHLIVTPSGC GEQNMIGMTP
1021 TVIAVHYLDE TEQWEKFGLE KRQGALELIK **KGYTQQLAFR** **QPSSAFAAFV** KRAPSTWLTA
1081 YVVKV**FS**LAV **NLIAIDSQVL** **CGAVKWILE** **KQKPDGVFQE** **DAPVIHQEMI** **GGLRNNNEKD**
1141 MALTAFLVIS **LQEAKDICEE** **QVNSLPGSIT** **KAGDFLEANY** **MNLQRSYTVA** **IAGYALAQMG**
1201 **RLKGPLLKNE** **LTTAKDKNRW** **EDPGKQLYNV** **EATSYALLAL** **LQLKDFDFVP** **PVVRWLNEQR**
1261 **YYGGGYGSTQ** **ATFMVFQALA** **QYQKADPDHQ** **ELNLDVSLQL** **PSRSSKITHR** **IHWESASLLR**
1321 **SEETKENEGF** **TVTAEGKGQC** **TLVVVTMYHA** **KAKDQLTCNK** **FDLKVTIKPA** **PETEKRPQDA**
1381 **KNTMILEICT** **RYRGDQDATM** **SILDISMMTG** **FAPDTDDLKQ** **LANGVDYRIS** **KYELDKAFSD**
1441 **RNTLIIYLDK** **VSHSEDDCLA** **FKVHQYFNVE** **LIQPGAVKVY** **AYYNLEESCT** **RFYHPEKEDG**
1501 **KLNLKCRDEL** **CRCAEENCFI** **QKSDDKVTLE** **ERLDKACEPG** **VDYVYKTRLV** **KVQLSNDFDE**
1561 **YIMAIEQTIK** **SGSDEVQVGQ** **QRTFISPIKC** **REALKLEEK** **HYLMWGLSSD** **FWGEKPNLSY**
1621 **IIGKDTWVEH** **WPEDDECQDE** **ENQKQCQDLG** **AFTESMVVFG** **CPN**

Figure 15

1 MMKTLLLFVG LLLTWESQV LGDQTVSDNE LQEMSNQGSK **YVNKEIQNAV** NGVKQIKTLI
61 EKTNEERKTL LSNLEEAKKK KEDALNETRE SETKLKELPG **VCNETMMALW** **EECKPCLKQT**
121 CMKFYARVCR SGSGLVGRQL EEFLNQSSPF YFWMNGDRID SLENDRQQT HMLDVMQDHF
181 **SRASSIIDEL** **FQDRFFTREP** **QDTYHYLPFS** **LPHRRPHFTF** **PKSRIVRSLM** PFSPIEPLNF
241 HAMFQPFLEM IHEAQQAMDI HFHSPAFQHP PTEFIREGDD DRTVCREIRH NSTGCLRMKD
301 QCDKCREILS VDCSTNNPSQ AKLRREIDES **LQVAERLTRK** YNELLKSYQW KMLNTSSLLE
361 QLNEQFNWVS RLANLTQGED QYYLRVTTVA SHTSDSDVPS GVTEVVVKLF **DSDPITVTVP**
421 **VEVSRKNPKF** METVAEKALQ EYRKKHREE

Figure 16

1 MRLLLWGLIWA SSFFTLSLQK PRLLLFSPSV VHLGVPLSVG VQLQDVPRGQ VVKGSVFLRN
61 PSRNNVPCSP KVDFTLSSER DFALLSLQVP LKDAKSCGLH QLLRGPEVQL VAHSPWLKDS
121 LSRTTNIQGI NLLFSSRRGH LFLQTDQPIY NPGQVRVRYRV FALDQKMRPS TDTITVMVEN
181 SHGLRVRKKE VYMPSSIFQD DFVIPDISEP GTWKISARFS DGLESNSSTQ FEVKKYVLPN
241 FEVKITPGKP YILTVPGHLD EMQLDIQARY IYGKPVQGVA YVRFGLLDED GKKTFFRGLE
301 SQTCLVNGQS HISLSKAEFQ DALEKLNMG I TDLOGLRLYV AAAIESPGC EMBEAEELTSW
361 YFVSSPFSLD LSKTKRHLVP GAPFLLQALV REMSGSPASG IPVKVSATVS SPGSVPEVQD
421 IQQNTDGSQ VSIPITIPQT ISELQLSVSA GSPHPAIARL TVAAPPSGGP GFLSIERPDS
481 RPPRVGDTLN LNLRAVGSGA TFSHYYMIL SRGQIVFMNR EPKRTLTSVS VFVDHHLAPS
541 FYFVAFYYHG DHPVANSLRV DVQAGACEGK LELSVDGAKQ YRNGESVKLH LETDSLALVA
601 LGALDTALYA AGSKSHKPLN MGKVFEAMNS YDLGCGPGGG DSALQVFQAA GLAFSDGDQW
661 TLSRKRLSCP **KEKT**TRKKRN VNFQKAIN EK **LGQYASPTAK** RCCQDGVTRL PMMRSCEQRA
721 ARVQQPDCRE PFLSCCQFAE SLRKKSRDKG QAGLQRALEI LQEEIDLDED DIPVRSFFPE
781 NWLWRVETVD RFQILTTLWLP DSLTTWEIHG LSLSKTKGLC **VATPVQLRVF** **REFHLHLRLP**
841 **MSVRRFEQLE** LRPVLYNYLD KNLTVSVHVS PVEGLCLAGG GGLAQQVLVP AGSARPVAFS
901 VVPTAAAVS LKVVARGSFE **FPVGD**AVSKV **LQIEKEGAIH** **REELVYELNP** **LDHRGR****TLEI**
961 **PGNSDPNMIP** **DGDFNSYVRV** **TASDPLDTLG** **SEGALSPGGV** **ASLLRLPRGC** **GEQ****TMIIY****LAP**
1021 **TLAASRY**LDK TEQWSTLPPE TKDHAVDLIQ KGYMRIQQFR KADGSYAAWL SRDSSTWLTA
1081 FVLKVLSLAQ EQVGGSP EKL QETSNWLLSQ QQADGSFQDP CPVLD RSMQG GLVGNDETVA
1141 LTAFTVIALH HGLAVFQDEG AEPLKQ RVEA SISKANSFLG EKASAGLLGA HAAAITAYAL
1201 SLTKAPVDLL GVAHNNLMAM AQETGDNLYW GSVTGSQSNA VSPTPAPRNP SDPMPQAPAL
1261 WIETTAYALL HLLHEGKAE MADQASAWLT **RQGSFQGGFR** STQDTVIALD ALSAYWIASH
1321 TTEERGLNVT LSSTGRNGFK SHALQLNNRQ IRGLEEELQF SLGSKINVKV GGNSKGT LKV
1381 LRTYNVLD MK NTTCQDLQIE VTVKGHVEYT MEANEDYEDY EYDELP AKDD PDAPLPVTP
1441 LQLFEGRRNR RRREAPKVVE EQESRVHYTV CIWRNGKVGL SGMAIADVT L LSGFHALRAD
1501 LEKLTSLSDR YVSHFETEGP HVLLYFDSVP TSRECVGF EA VQEV PVGLVQ PASATLYDYY
1561 NPERRCSV FY GAPSKSRLLA TLCSAEVCQC AEGKCPQR R ALERGLQDED GYRMKFACY
1621 PRVEYGFQVK VLREDSRAAF RLFETKITQV LHFTKDV KAA ANQMRNFLVR ASCRLRL EPG
1681 KEYLIMGLDG ATYDLEGHPQ YLLDSNSWIE EMPSERLCRS TRQRAACAQL NDFLOEYGTQ
1741 GCQV

Figure 17

1 MEEEEIAALVI DNGSGMCKAG FAGDDAPRAV **FPSIVGRPRH** QGVMVGMGQK **DSYVGDEAQS**
61 **KRGILTLKYP** IEHGIVTNWD DMEKIWHHTF YNELRVAPEE **HPVLLTEAPL** **NPKANREKMT**
121 QIMFETFNTF AMYVAIQAVL SLYASGR**TTG** **IVMDSGDGVT** **HTVPIYEGYA** **LPHAILRLDL**
181 **AGRDLDYLM** KILTERGYSF TTTAEREIVR DIKEKLCYVA LDFEQEMATA ASSSSLEKSY
241 ELPDGQVITI GNERFRCPEA LFQPSFLGME SCGIHETTFN SIMKCDVDIR **KDLYANTVLS**
301 **GGTTMYPGIA** DRMQKEITAL APSTM**KIKII** **APPERKYSVW** IGGSILASLS TFQQMWISKQ
361 **EYDESGPSIV** HRKCF

Figure 18

1 MSALGAVIAL LLWGQLFAVD SGNDVTDIAD DGCPKPPEIA HGYVEHSVRY QCKNYYKLRT
61 EGDGVYTLND KKQWINKAVG DKLPECEADD GCPKPPEIAH GYVEHSVRYQ CKNYYKL RTE
121 GDGVYTLNNE KQWINKAVGD KLPECEAVCG KPKNPANPVQ RILGGHLD AK GSFPWQAKMV
181 SHHNLT TGAT LINEQWLLTT AKNLFLNHSE NATAKD IAPT **LTLYVGKKQL** VEIEKVVLHP
241 NYSQVDIGLI KLKQKVS VNE RVMPICLPSK **DYAEVGRVGY** VSGWGRNANF **KFTDHLKYVM**
301 **LPVADQDQCI** **RHYEGSTVPE** **KKTPKSPVGV** **QPILNEHTFC** **AGMSKYQEDT** CYGDAGSAFA
361 VHDLEEDTWY **ATGILSFDKS** **CAVAEYGVYV** KVTSIQDWVQ KTIAEN

Figure 19

METHODS OF DIAGNOSING ALZHEIMER'S DISEASE

FIELD OF THE INVENTION

The present invention relates to methods and compositions relating to Alzheimer's disease. Specifically, the present invention identifies and describes proteins that are differentially expressed in the Alzheimer's disease state relative to their expression in the normal state and, in particular, identifies and describes proteins associated with Alzheimer's disease. Further, the present invention provides methods of diagnosis of Alzheimer's disease using the differentially expressed proteins. Still further, the present invention provides methods for the identification and therapeutic use of compounds for the prevention and treatment of Alzheimer's disease.

BACKGROUND OF THE INVENTION

Dementia is one of the major public health problems of the elderly, and in our ageing populations the increasing numbers of patients with dementia is imposing a major financial burden on health systems around the world. More than half of the patients with dementia have Alzheimer's disease (AD). The prevalence and incidence of AD have been shown to increase exponentially. The prevalence for AD in Europe is 0.3% for ages 60-69 years, 3.2% for ages 70-79 years, and 10.8% for ages 80-89 years (Rocca, Hofman et al. 1991). The survival time after the onset of AD is approximately from 5 to 12 years (Friedland 1993).

Alzheimer's disease (AD), the most common cause of dementia in older individuals, is a debilitating neurodegenerative disease for which there is currently no cure. It destroys neurons in parts of the brain, chiefly the hippocampus, which is a region involved in coding memories. Alzheimer's disease gives rise to an irreversible progressive loss of cognitive functions and of functional autonomy. The earliest signs of AD may be mistaken for simple forgetfulness, but in those who are eventually diagnosed with the disease, these initial signs inexorably progress to more severe symptoms of mental deterioration. While the time it takes for AD to develop will vary from person to person, advanced signs include severe memory impairment, confusion, language disturbances, personality and behaviour changes, and impaired judgement. Persons with AD may become non-communicative and hostile. As the disease ends its course in profound dementia, patients are unable to care for themselves and often require institutionalisation or professional care in the home setting. While some patients may live for years after being diagnosed with AD, the average life expectancy after diagnosis is eight years.

In the past, AD could only be definitively diagnosed by brain biopsy or upon autopsy after a patient died. These methods, which demonstrate the presence of the characteristic plaque and tangle lesions in the brain, are still considered the gold standard for the pathological diagnoses of AD. However, in the clinical setting brain biopsy is rarely performed and diagnosis depends on a battery of neurological, psychometric and biochemical tests, including the measurement of biochemical markers such as the ApoE and tau proteins or the beta-amyloid peptide in cerebrospinal fluid and blood.

Biomarkers may possibly possess the key in the next step for diagnosing AD and other dementias. A biological marker that fulfils the requirements for the diagnostic test for AD would have several advantages. An ideal biological marker would be one that identifies AD cases at a very early stage of

the disease, before there is degeneration observed in the brain imaging and neuropathological tests. A biomarker could be the first indicator for starting treatment as early as possible, and also very valuable in screening the effectiveness of new therapies, particularly those that are focussed on preventing the development of neuropathological changes. A biological marker would also be useful in the follow-up of the development of the disease.

Markers related to pathological characteristics of AD; plaques and tangles (A β and tau) have been the most extensively studied. The most promising has been from studies of CSF concentration of A β (1-40), A β (1-42) and tau or the combination of both proteins in AD. Many studies have reported a decrease in A β (1-42) in CSF, while the total A β protein or A β (1-40) concentration remain unchanged (Iida, Hartmann et al. 1996; Kanai, Matsubara et al. 1998; Andreasen, Hesse et al. 1999).

SUMMARY OF THE INVENTION

Broadly, the present invention relates to methods and compositions for the diagnosis of Alzheimer's disease. More specifically, the present invention identifies and describes proteins that are differentially expressed in the Alzheimer's disease state relative to their expression in the normal state.

In a first aspect, the invention provides a method of diagnosing Alzheimer's disease in a subject, the method comprising detecting one or more of a differentially expressed protein identified by the methods described herein in a tissue sample or body fluid sample from said subject. Preferably, the method is an in vitro method.

In all aspects, the methods of the present invention may also be used in relation to pre-Alzheimer's stages such as mild cognitive impairment (MCI) as well as advanced Alzheimer's disease.

In another aspect, the present invention provides a method of determining the nature or degree of Alzheimer's disease in a human or animal subject, the method comprising detecting one or more of a differentially expressed protein identified by the methods described herein in a tissue sample or body fluid sample from said subject. Thus, the methods of the present invention encompass methods of monitoring the progress of Alzheimer's disease or of disease progression from MCI to Alzheimer's disease. Also encompassed are prognostic methods, for example prognosis of likely progression from MCI to Alzheimer's disease, or prognosis of likely duration or severity of Alzheimer's disease.

In a preferred embodiment the method comprises:

- (a) establishing a paradigm in which at least one protein is differentially expressed in relevant tissue or body fluid sample from, or representative of, subjects having differential levels of Alzheimer's disease;
- (b) obtaining a sample of the tissue or body fluid sample from the subject;
- (c) determining the presence, absence or degree of expression of the differentially expressed protein or proteins in the sample; and
- (d) relating the determination to the nature or degree of the Alzheimer's disease by reference to a previous correlation between such a determination and clinical information.

In one embodiment, the progression of the disorder may be tracked by using the methods of the invention to determine the severity of the disorder, e.g. global dementia severity). In another embodiment, the duration of the disorder up to the point of assessment may be determined using the methods of the invention. For example, expression of an Ig lambda chain

C region (see spot 177, FIG. 6) may correlate with global dementia severity. Expression of a serum albumin precursor (see spot 165, FIG. 6) may show a negative correlation with the duration of the disease.

This method allows the type of Alzheimer's disease of a patient to be correlated to different types of prophylactic or therapeutic treatment available in the art, thereby enhancing the likely response of the patient to the therapy.

In some embodiments, more than one protein is differentially expressed, providing a multi-protein fingerprint of the nature or degree of the Alzheimer's disease. Preferably, at least four proteins are differentially expressed.

Conveniently, the patient sample used in the methods of the invention can be a tissue sample or body fluid sample such as a blood, plasma, serum or urine sample. Use of body fluids such as those listed is preferred because they can be more readily obtained from a subject. This has clear advantages in terms of cost, ease, speed and subject wellbeing. Blood, blood products such as plasma, and urine are particularly preferred.

The step of detecting the differentially expressed protein may be preceded by a depletion step to remove the most abundant proteins from the sample, as described below.

Preferably, at least one of the differentially expressed proteins is a protein shown in FIG. 6, FIG. 7 or FIG. 12. In preferred embodiments, the differentially expressed protein is apolipoprotein A-IV precursor, apolipoprotein C-III precursor, transthyretin, galectin 7, complement C4 precursor, alpha-2-macroglobulin precursor, Ig alpha-1 chain C, histone 2B, Ig lambda chain C region, fibrinogen gamma chain precursor, complement factor H, inter-alpha-trypsin heavy chain H4 precursor, complement C3 precursor, clusterin precursor, gamma or beta actin, haptoglobin precursor or the serum albumin precursor isoform found in spot ID no 2, 14, 15, 123, 165, 176 or 184 of FIG. 6 or fragments thereof. Preferred fragments are a C-terminal fragment of Apo-AIV or a C4 alpha region of complement C4 precursor lacking the anaphylatoxin domain. For example, the fragment may comprise amino acid residues 270-309 of apolipoprotein A-IV; residues 1446-1744 of complement C4, or may be an N-terminal fragment of apolipoprotein A-IV which migrates as a polypeptide of 10-16 kD or a polypeptide of 28 kD in SDS-PAGE, or a fragment of any of the proteins in FIG. 7 with a molecular weight of 6430, 14640, 27147 or 14646 Da. Other preferred fragments comprise the areas indicated in bold in FIGS. 9, 10, and 13 to 19.

Preferred fragments are less than 50, less than 100, less than 150, less than 200, less than 250, less than 300, less than 350, less than 400, less than 500, less than 600, less than 700, less than 800, less than 900, less than 1000, less than 1100, less than 1200, less than 1300, less than 1400, less than 1500, less than 1600, less than 1700, less than 1800, less than 1900 or less than 2000 amino acids in length.

The expression of certain differentially expressed proteins may be increased in subjects with Alzheimer's disease as compared to control subjects. The expression of other differentially expressed proteins may be decreased in subjects with Alzheimer's disease as compared to control subjects. FIGS. 6, 8 and 12 indicate whether the expression of the proteins disclosed therein is increased or decreased in Alzheimer's versus control subjects. It is thus clear from the figures whether an increase or decrease in expression is indicative of the disease state for all the proteins listed therein. Including the preferred proteins listed above.

Preferably, a differentially expressed protein shows a fold difference in expression of at least 1.5, at least 1.6, at least 1.7, at least 1.8, at least 1.9, at least 2.0, at least 2.5, at least 3, at

least 3.5, at least 4, at least 5, at least 10 or more between the level found in patients with Alzheimer's versus control subjects.

The differentially expressed protein may be detected using an antibody specific to that protein, for example in an ELISA assay or Western blotting. Alternatively, the differentially expressed protein may be detected by, amongst others, 2D gel electrophoresis or mass spectrometry techniques including LS/MS/MS, MALDI-TOF or SELDI-TOF. The sample may be immobilised on a solid support for analysis.

In one embodiment, a diagnosis may be made solely on the basis of the pattern of spots on a 2D gel prepared from a subject sample. The pattern of spots obtained from Alzheimer's disease or MCI subjects may be compared directly with the pattern obtained from control subject samples, without the need for identifying individual proteins.

In one embodiment, an antibody sandwich technique where antibodies specific for one or more of the biomarkers is added and the immobilised antibodies capture the biomarker protein. The captured proteins are then detected using a second antibody that may be directly labelled with a signal generating agent (enzyme, fluorescent tag, radiolabel etc.) or may be detected using further amplification (labelled secondary antibody, streptavidin/biotin systems with enzyme, fluorophore, radiolabel etc.). Other immunological methods may include one-dimensional or two-dimensional gel electrophoresis of patient samples followed by transfer to a solid surface using techniques such as Western blotting and subsequent detection using antibodies specific for the AD biomarkers.

In an alternative embodiment, autoantibodies to the biomarkers may be detected by using the Western blotting approach described above using either samples from a patient or representative of AD and then detecting the presence of antibodies specific for the biomarker that are present in the blood of AD patients but not in controls.

The method may further comprise determining an effective therapy for treating the Alzheimer's disease.

In a further aspect, the present invention provides a method of treatment by the use of an agent that will restore the expression of one or more differentially expressed proteins in the Alzheimer's disease state towards that found in the normal state in order to prevent the development or progression of Alzheimer's disease. Preferably, the expression of the protein is restored to that of the normal state.

In a further aspect, the present invention provides a method whereby the pattern of differentially expressed proteins in a tissue sample or body fluid sample or urine of an individual with Alzheimer's disease is used to predict the most appropriate and effective therapy to alleviate the Alzheimer's disease.

Also provided is a method of screening an agent to determine its usefulness in treating a Alzheimer's disease, the method comprising:

- (a) obtaining a sample of relevant tissue taken from, or representative of, a subject having Alzheimer's disease symptoms, who or which has been treated with the agent being screened;
- (b) determining the presence, absence or degree of expression of the differentially expressed protein or proteins in the tissue from, or representative of, the treated subject; and,
- (c) selecting or rejecting the agent according to the extent to which it changes the expression, activity or amount of the differentially expressed protein or proteins in the treated subject having Alzheimer's disease symptoms.

Optionally, the method may further comprise, prior to step (a), the step of establishing a paradigm in which at least one protein is differentially expressed in relevant tissue from, or representative of, subjects having Alzheimer's disease symptoms and normal subjects.

Preferably, the agent is selected if it converts the expression of the differentially expressed protein towards that of a normal subject. More preferably, the agent is selected if it converts the expression of the protein or proteins to that of the normal subject.

Also provided is a method of screening an agent to determine its usefulness in treating Alzheimer's disease, the method comprising:

- (a) obtaining over time samples of relevant tissue or body fluid taken from, or representative of, a subject having Alzheimer's disease symptoms, who or which has been treated with the agent being screened;
- (b) determining the presence, absence or degree of expression of a differentially expressed protein or proteins in said samples; and,
- (c) determining whether the agent affects the change over time in the expression of the differentially expression protein in the treated subject having Alzheimer's disease symptoms.

Optionally, the method may further comprise, prior to step (a), the step of establishing a paradigm in which at least one protein is differentially expressed in relevant tissue or body fluid from, or representative of, subjects having Alzheimer's disease symptoms and normal subjects; and/or

establishing that expression of said differentially expressed protein diverges over time in subjects having Alzheimer's disease symptoms and normal subjects.

Samples taken over time may be taken at intervals of weeks, months or years. For example, samples may be taken at monthly, two-monthly, three-monthly, four-monthly, six-monthly, eight-monthly or twelve-monthly intervals.

A change in expression over time may be an increase or decrease in expression, compared to the initial level of expression in samples from the subject and/or compared to the level of expression in samples from normal subjects. The agent is selected if it slows or stops the change of expression over time.

In the screening methods described above, subjects having differential levels of protein expression comprise:

- (a) normal subjects and subjects having Alzheimer's disease symptoms; and,
- (b) subjects having Alzheimer's disease symptoms which have not been treated with the agent and subjects Alzheimer's disease which have been treated with the agent.

In alternative embodiments, the subjects having differential levels of protein expression comprise:

- (a) normal subjects who have and have not been treated with the agent; and one or both of
- (b) subjects having mild cognitive impairment who have and have not been treated with the agent; and
- (c) subjects having Alzheimer's disease symptoms who have and have not been treated with the agent.

Preferably, the differential levels of protein expression are not observed in normal subjects who have and have not been treated with the agent.

The subjects having Alzheimer's disease symptoms are preferably human subjects with Alzheimer's disease.

Alternatively, the subjects having Alzheimer's disease symptoms may be an animal model such as mutant amyloid precursor protein (APP) transgenic mice, presenilin-1 (PS-1) transgenic mice, and/or double transgenic APP/PS-1 transgenic mice.

The tissue or body fluid samples may be, for example, brain tissue, blood, plasma, serum, saliva or cerebro-spinal fluid samples.

In one embodiment, the paradigm is established using two-dimensional (2D) gel electrophoresis carried out on the relevant tissue or a protein-containing extract thereof.

In another embodiment, the paradigm is established using SELDI analysis of the relevant tissue or a protein-containing extract thereof. Preferably, the tissue or extract is immobilised on a solid support, for example a chip.

Conveniently, a depletion step may be performed prior to 2D gel electrophoresis or SELDI analysis, to remove the most abundant proteins from the samples and reduce background.

The method may further comprise the step of isolating a differentially expressed protein identified in the method, and optionally the step of characterising the isolated protein.

Preferably, at least one of the differentially expressed proteins is a protein shown in FIG. 6, FIG. 7, FIG. 8 or FIG. 12 or a rodent equivalent thereof. In preferred embodiments, the differentially expressed protein is apolipoprotein A-IV precursor, apolipoprotein C-III precursor, transthyretin, galectin 7, complement C4 precursor, complement factor H, S100 calcium binding protein or ceruloplasmin, inter-alpha-trypsin heavy chain H4 precursor, complement C3 precursor, clusterin precursor, gamma or beta actin, haptoglobin precursor or fragments thereof. Preferred fragments are a C-terminal fragment of Apo-AIV or a C4 alpha region of complement C4 precursor lacking the anaphylatoxin domain. For example, the fragment may comprise amino acid residues 270-309 of apolipoprotein A-IV; residues 1446-1744 of complement C4.

Preferred fragments will comprise one or more of the sequences highlighted in FIGS. 9, 10 and 13-19.

In a further aspect, the invention provides a method of making a pharmaceutical composition which comprises having identified an agent using the method described above, the further step of manufacturing the agent and formulating it with an acceptable carrier to provide the pharmaceutical composition.

In a further aspect, the invention provides a method of identifying a protein which is differentially expressed in relevant tissue or body fluid sample from subjects with mild cognitive impairment and/or subjects with Alzheimer's disease and normal subjects, comprising:

- i) immobilising a tissue sample or body fluid sample or protein-containing extract thereof on a solid support
- ii) analysing the immobilised proteins by surface enhanced laser desorption time of flight mass spectroscopy
- iii) comparing the spectra obtained to detect differences in protein expression between Alzheimer's subjects and normal subjects.

Also provided is protein which is differentially expressed in relevant tissue from, or representative of subjects having differential levels of Alzheimer's disease symptoms and which is as obtainable by the methods described herein or by two-dimensional gel electrophoresis carried out on said tissue or a protein-containing extract thereof, the method comprising:

- (a) providing non-linear immobilized pH gradient (ILG) strips of acrylamide polymer 3 mm×180 mm;
- (b) rehydrating the IPG strips in a cassette containing 25 ml. of an aqueous solution of urea (8M), 3-[(cholamidopropyl)dimethylammonio]-1-propanesulphonate (CHAPS, 2% w/v), 0.5% IPG Pharmalyte and a trace of Bromophenol Blue;
- (c) emptying the cassette of liquid, transferring the strips to an electrophoretic tray fitted with humid electrode

- wicks, electrodes and sample cups, covering the strips and cups with low viscosity paraffin oil;
- (d) applying 200 micrograms of an aqueous solution of dried, powdered material of the relevant body tissue in urea (8M), CHAPS (4% w/v), Tris (40 mM), 0.5% IPG Pharmalyte and a trace of Bromophenol Blue to the sample cups, at the cathodic end of the IPG strips;
 - (e) carrying out isoelectric focusing on the gel at S1 500V step-n-hold (s/h) for 1 h; S2 500V s/h for 2 h; S3 1000V gradient (G) for 1 h; S4 1000V s/h for 2 h; S5 8000V G for 2 h and S6 8000V s/h for a time effective to enable the proteins to migrate in the strips to their pI-dependent final positions;
 - (f) equilibrating the strips within the tray with 100 ml of an aqueous solution containing Tris-HCl (50 mM) pH 6.8, urea (6M), glycerol (30% v/v), SDS (2% w/v) and DTT (10 mg/ml);
 - (g) replacing this solution by 100 ml. of an aqueous solution containing Tris-HCl (50 mM) pH 8.8, urea (6M), glycerol (30% v/v), SDS (2% w/v), iodoacetamide (25 mg/ml) and a trace of Bromophenol Blue and incubating for 20 minutes;
 - (h) providing a vertical gradient slab gel 160×200×1.5 mm of acrylamide/piperazine-diacrylyl cross-linker (9-16% T/2.6% C), polymerised in the presence of TEMED (0.5% w/v), ammonium persulphate (0.1% w/v) and sodium thiosulphate (5 mM), in Tris-HCl (0.375M) pH 8.8 as leading buffer;
 - (i) over-layering the gel with sec-butanol for about 2 hours, removing the overlay and replacing it with water;
 - (j) cutting the IPG gel strips to a size suitable for the second dimensional electrophoresis, removing 6 mm from the anode end and 14 mm from the cathode end;
 - (k) over-layering the slab gel with an aqueous solution of agarose (0.5% w/v) and Tris-glycine-SDS (25 mM-198 mM-0.1% w/v) as leading buffer, heated to 70° C. and loading the IPG gel strips onto the slab gel through this over-layered solution;
 - (l) running the second dimensional electrophoresis at a constant current of 40 mA at 8-12° C. for 5 hours; and
 - (m) washing the gel.

This invention is based, in part, on systematic search strategies involving sensitive detection of proteins by 2D-electrophoresis. To aid the identification of differentially expressed protein a standard marker set of proteins such as those available from Genomic Solutions may be run on an extra lane to 2D electrophoresis.

The examples presented below demonstrate the successful use of the experimental paradigms of the invention to identify target proteins associated with Alzheimer's disease.

DEFINITIONS

"Differential expression", as used herein, refers to at least one recognisable difference in tissue or body fluid protein expression. It may be a quantitatively measurable, semi-quantitatively estimatable or qualitatively detectable difference in tissue protein expression. Thus, a differentially expressed protein (herein DEP) may be strongly expressed in tissue in the normal state and less strongly expressed or not expressed at all in tissue in the Alzheimer's disease state. Conversely, it may be strongly expressed in tissue in the Alzheimer's disease state and less strongly expressed or not expressed at all in the normal state. Further, expression may be regarded as differential if the protein undergoes any recognisable change between the two states under comparison.

The term "paradigm" means a prototype example, test model or standard.

Wherever a differentially expressible protein is used in the screening procedure, it follows that there must have been at some time in the past a preliminary step of establishing a paradigm by which the differential expressibility of the protein was pre-determined. Once the paradigm has been established, it need not be re-established on every occasion that a screening procedure is carried out. The term "establishing a paradigm" is to be construed accordingly.

"Relevant tissue" means any tissue involved in brain function, in particular tissue involved in Alzheimer's disease.

"Tissue/Body fluid . . . representative of . . . subjects" means any tissue or body fluid in which the above-mentioned biological change can be simulated for laboratory purposes and includes, for example, a primary cell culture or cell line derived ultimately from relevant tissue.

The term "subjects" includes human and animal subjects.

The treatments referred to above can comprise the administration of one or more drugs or foodstuffs, and/or other factors such as diet or exercise.

The differentially expressed proteins (DEPs) include "fingerprint proteins", "target proteins" or "pathway proteins".

The term "fingerprint protein", as used herein, means a DEP, the expression of which can be used, alone or together with other DEPs, to monitor or assess the condition of a patient suspected of suffering from Alzheimer's disease. Since these proteins will normally be used in combination, especially a combination of four or more, they are conveniently termed "fingerprint proteins", without prejudice to the possibility that on occasions they may be used singly or along with only one or two other proteins for this purpose. Such a fingerprint protein or proteins can be used, for example, to diagnose a particular type of Alzheimer's disease and thence to suggest a specific treatment for it.

The term "diagnosis", as used herein, includes the provision of any information concerning the existence, non-existence or probability of the disorder in a patient. It further includes the provision of information concerning the type or classification of the disorder or of symptoms which are or may be experienced in connection with it. This may include, for example, diagnosis of the severity of the disorder. It encompasses prognosis of the medical course of the disorder, for example its duration, severity and the course of progression from MCI to Alzheimer's disease.

Currently disease status is assessed by duration of disease from inception to present (longer duration equals more severe disease) and clinical assessment measures. These assessment measures include clinical tests for memory and other cognitions, clinical tests for function (abilities of daily living) and clinical assessments of global severity. Trials of potential therapies in AD are currently evaluated against these measures. The FDA and other medicines approval bodies require as part of these assessments measures of both cognition and global function. The Global Dementia Scale is one such measure of global function. It is assessed by rater assessment of severity including cognition and function against a standardised set of severity criteria.

The term "target protein", as used herein, means a DEP, the level or activity of which can be modulated by treatment to alleviate Alzheimer's disease. Modulation of the level or activity of the target protein in a patient may be achieved, for example, by administering the target protein, another protein or gene which interacts with it or an agent which counteracts or reduces it, for example an antibody to the protein, com-

petitive inhibitor of the protein or an agent which acts in the process of transcription or translation of the corresponding gene.

The term "alleviate", as used herein, in relation to Alzheimer's disease means any form of reducing one or more undesired symptoms or effects thereof. Any amelioration of Alzheimer's disease of the patient falls within the term "alleviation". Amelioration may also include slowing down the progression of the disease.

Alternatively or additionally, the DEPs can interact with at least one other protein or with a gene involved in the regulation of brain function. Such other proteins are termed herein "pathway proteins" (PPs). The term is applied to the protein with which the DEP interacts, not to the DEP itself, although a pathway protein can be another DEP.

By way of example, embodiments of the present invention will now be described in more detail with reference to the accompanying figures.

BRIEF DESCRIPTION OF THE FIGURES

FIG. 1 shows spectra for the 6430 Da peak identified by SELDI analysis in normal (top) and Alzheimer's disease (bottom) subjects.

FIG. 2 shows spectra for the 14640 Da peak identified by SELDI analysis in normal (top) and Alzheimer's disease (bottom) subjects.

FIG. 3 shows spectra for the 27147 Da peak identified by SELDI analysis in normal (top) and Alzheimer's disease (bottom) subjects.

FIG. 4 shows spectra for the 14646 Da peak identified by SELDI analysis in pooled normal (top) and Alzheimer's disease (bottom) subjects.

FIG. 5 shows a silver stained gel obtained from material extracted from the chips used for SELDI analysis. The bands (1-6) excised and analysed by LC/MS/MS are indicated by arrows.

FIG. 6 shows differentially expressed proteins identified by 2D gel analysis and mass spectroscopy.

FIG. 7 shows differentially expressed proteins identified by SELDI and LC/MS/MS.

FIG. 8 shows differentially expressed proteins identified by qPST.

FIG. 9 shows the sequence coverage (indicated in bold) obtained for apolipoprotein A-IV (P06727) in the 14.6 kDa band isolated on the Q10 SAX2 SELDI chip. C-terminal residues 270-396 are underlined. The amino acid sequence is SEQ ID NO: 1.

FIG. 10 shows sequence coverage (indicated in bold) obtained for Complement C4 precursor (P01028) in 2DE spot 164. The amino acid sequence is SEQ ID NO: 2.

FIG. 11 shows a 2D gel obtained from the pre-depletion analysis. The differentially expressed spots are circled.

FIG. 12 lists the differentially expressed spots identified by the pre-depletion analysis. Column 3 gives the accession number for the human protein, column 4 the mean normalised spot volume in the control samples, column 6 the mean normalised spot volume in the disease samples, column 8 the mean normalised spot volume in the disease sample divided by that in the control sample, column 9 the significance (p-value) of the difference in spot volumes compared by Student's t-test, column 10 the number of gels in the control group in which the spot was detected. CV is coefficient of variation.

FIG. 13 shows sequence coverage (indicated in bold) obtained for alpha-2 macroglobulin (P01023) in the pre-

depletion analysis. The signal sequence is underlined. The amino acid sequence is SEQ ID NO: 3.

FIG. 14 shows sequence coverage (indicated in bold) obtained for inter-alpha trypsin inhibitor heavy chain H4 precursor (Q14624) in the pre-depletion analysis. The signal sequence is underlined. The amino acid sequence is SEQ ID NO: 4.

FIG. 15 shows sequence coverage (indicated in bold) obtained for complement C3 precursor (P01024) in the pre-depletion analysis. The signal sequence is underlined. The amino acid sequence is SEQ ID NO: 5.

FIG. 16 shows sequence coverage (indicated in bold) obtained for clusterin precursor (P10909) in the pre-depletion analysis. The signal sequence is underlined. The amino acid sequence is SEQ ID NO: 6.

FIG. 17 shows sequence coverage (indicated in bold for spot 832 and bold italic for spot 652) obtained for complement C4 precursor (P01028) in the pre-depletion analysis. The signal sequence is underlined. The amino acid sequence is SEQ ID NO: 7.

FIG. 18 shows sequence coverage (indicated in bold) obtained for gamma actin (P63261) in the pre-depletion analysis. The signal sequence is underlined. The amino acid sequence is SEQ ID NO: 8.

FIG. 19 shows sequence coverage (indicated in bold) obtained for haptoglobin precursor (P00738) in the pre-depletion analysis. The signal sequence is underlined. The amino acid sequence is SEQ ID NO: 9.

DETAILED DESCRIPTION

Methods and compositions for the treatment of Alzheimer's disease. Proteins termed 'target proteins' and/or fingerprint proteins are described which are differentially expressed in Alzheimer's disease states relative to their expression in normal states. Methods for the identification of such fingerprint and target proteins are also described.

'Differential expression' as used herein indicates that a protein is present at different levels in samples from normal and diseased subjects.

Also described below are methods for prognostic and diagnostic evaluation of Alzheimer's disease states and for the identification of subjects exhibiting a predisposition to Alzheimer's disease.

1. Identification of Differentially Expressed and Pathway Proteins

In one embodiment, the present invention concerns methods for the identification of proteins which are involved in Alzheimer's disease. Such proteins may represent proteins which are differentially expressed in Alzheimer's disease states relative to their expression in normal states. Such differentially expressed proteins may represent 'target' or 'fingerprint' proteins.

Methods for the identification of such proteins are described in Section 1. Methods for the further characterisation of such differentially expressed proteins and for their identification as target and/or fingerprint proteins are presented below in Section 1.1.

'Differential expression', as used herein, refers to both qualitative as well as quantitative differences in protein expression. Thus a differentially expressed protein may qualitatively have its expression activated or completely inactivated in normal versus Alzheimer's disease state. Such a qualitatively regulated protein will exhibit an expression pattern within a given tissue, cell type or body fluid sample which is detectable in either control or Alzheimer's disease subject, but not detectable in both. Alternatively, such a quali-

tatively regulated protein will exhibit an expression pattern within a given tissue, cell type or body fluid sample, which is detectable in either control or Alzheimer's disease subjects but not detectable in both. 'Detectable', as used herein, refers to a protein expression pattern, which are detectable using techniques described herein.

Alternatively, a differentially expressed protein may have its expression modulated, i.e. quantitatively increased or decreased, in normal versus Alzheimer's disease states. The degree to which expression differs in normal versus Alzheimer's disease states need only be large enough to be visualised via standard characterisation techniques, such as silver staining of 2D-electrophoretic gels. Other such standard characterisation techniques by which expression differences may be visualised are well known to those skilled in the art. These include successive chromatographic separations of fractions and comparisons of the peaks, capillary electrophoresis, separations using micro-channel networks, including on a micro-chip, SELDI analysis and qPST analysis.

Chromatographic separations can be carried out by high performance liquid chromatography as described in Pharmacia literature, the chromatogram being obtained in the form of a plot of absorbance of light at 280 nm against time of separation. The material giving incompletely resolved peaks is then re-chromatographed and so on.

Capillary electrophoresis is a technique described in many publications, for example in the literature "Total CE Solutions" supplied by Beckman with their P/ACE 5000 system. The technique depends on applying an electric potential across the sample contained in a small capillary tube. The tube has a charged surface, such as negatively charged silicate glass. Oppositely charged ions (in this instance, positive ions) are attracted to the surface and then migrate to the appropriate electrode of the same polarity as the surface (in this instance, the cathode). In this electroosmotic flow (EOF) of the sample, the positive ions move fastest, followed by uncharged material and negatively charged ions. Thus, proteins are separated essentially according to charge on them.

Micro-channel networks function somewhat like capillaries and can be formed by photocablation of a polymeric material. In this technique, a UV laser is used to generate high energy light pulses that are fired in bursts onto polymers having suitable UV absorption characteristics, for example polyethylene terephthalate or polycarbonate. The incident photons break chemical bonds with a confined space, leading to a rise in internal pressure, mini-explosions and ejection of the ablated material, leaving behind voids which form micro-channels. The micro-channel material achieves a separation based on EOF, as for capillary electrophoresis. It is adaptable to micro-chip form, each chip having its own sample injector, separation column and electrochemical detector: see J. S. Rossier et al., 1999, *Electrophoresis* 20: pages 727-731.

Surface enhanced laser desorption ionisation time of flight mass spectrometry (SELDI-TOF-MS) combined with ProteinChip technology can also provide a rapid and sensitive means of profiling proteins and is used as an alternative to 2D gel electrophoresis in a complementary fashion. The ProteinChip system consists of aluminium chips to which protein samples can be selectively bound on the surface chemistry of the chip (eg. anionic, cationic, hydrophobic, hydrophilic etc). Bound proteins are then co-crystallised with a molar excess of small energy-absorbing molecules. The chip is then analysed by short intense pulses of N2 320 nm UV laser with protein separation and detection being by time of flight mass spectrometry. Spectral profiles of each group within an experi-

ment are compared and any peaks of interest can be further analysed using techniques as described below to establish the identity of the protein.

Quantitative protein sequence tag (qPST) technology may also be used to detect differentially expressed proteins. Briefly, the proteins in the samples for comparison are labelled with a stable isotope tag. A different isotope is used for each sample. The proteins are enzymatically cleaved and the labelled peptides in each sample are quantified by mass spectrometry. In this way, expression of equivalent proteins in the different samples can be compared directly by comparing the intensities of their respective isotopic peaks.

Detection of differentially expressed proteins may be preceded by a depletion step to remove the most abundant proteins from the sample. The large majority of the protein composition of serum/plasma consists of just a few proteins. For example, albumin, which is present at a concentration of 35-50 mg/ml, represents approximately 54% of the total protein content with IgG adding other 16%. In contrast, proteins changing in response to disease, for example as a result of tissue leakage, may circulate at 10 ng/ml. This vast dynamic range of protein concentrations represents a major analytical challenge and to overcome the problem, a multiple affinity depletion column can be used to remove the most highly abundant proteins (eg the 5, 6, 7, 8, 9 or 10 most highly abundant proteins). This enables the detection of changes in lower abundance ranges because more starting material can be used and there is less interference from the highly abundant molecules. Such a depletion strategy can be applied before any detection method.

Differentially expressed proteins may be further described as target proteins and/or fingerprint proteins. 'Fingerprint proteins', as used herein, refer to a differentially expressed protein whose expression pattern may be utilised as part of a prognostic or diagnostic Alzheimer's disease evaluation or which, alternatively, may be used in methods for identifying compounds useful for the treatment of Alzheimer's disease. A fingerprint protein may also have characteristics of a target protein or a pathway protein.

'Target protein', as used herein, refers to a differentially expressed protein involved in Alzheimer's disease such that modulation of the level or activity of the protein may act to prevent the development of Alzheimer's disease. A target protein may also have the characteristics of a fingerprint protein or a pathway protein.

1.1 Characterisation of Differentially Expressed Proteins

Differentially expressed proteins, such as those identified via the methods discussed above, may be further characterised by using, for example, methods such as those discussed herein. Such proteins will be referred to herein as 'identified proteins'.

Analyses such as those described herein, yield information regarding the biological function of the identified proteins. An assessment of the biological function of the differentially expressed proteins, in addition, will allow for their designation as target and/or fingerprint proteins.

Specifically, any of the differentially expressed proteins whose further characterisation indicates that a modulation of the proteins expressed or a modulation of the proteins activity may ameliorate Alzheimer's disease will be designated 'target proteins', as defined above, in Section 1.

Any of the differentially expressed proteins whose expression pattern contributes to a protein 'fingerprint' profile correlative of Alzheimer's disease, will be designated a 'fingerprint protein'. 'Fingerprint profiles' will be more fully discussed below. It should be noted that each of the target proteins may also function as fingerprint proteins.

A variety of techniques can be utilised to further characterise the identified proteins. First, the corresponding nucleotide sequence of the identified protein may be obtained by utilising standard techniques well known to those of skill in the art, may, for example, be used to reveal homologies to one or more known sequence motifs which may yield information regarding the biological function of the identified protein.

Secondly, the biological function of the identified proteins may be more directly assessed by utilising relevant *in vivo* and *in vitro* systems. *In vivo* systems may include, but are not limited to, animal systems which naturally exhibit Alzheimer's disease-like symptoms and/or pathology, or ones which have been engineered to exhibit such symptom and/or pathology. Further, such systems may include systems for the further characterisation of Alzheimer's disease, and may include, but are not limited to, naturally occurring and transgenic animal systems.

In vitro systems may include cell lines derived from such animals or Alzheimer's disease subjects. Animal models may be used to generate cell lines, containing one or more cell types involved in Alzheimer's disease, that can be used as cell culture models for this disorder. While primary cultures derived from the transgenic animals of the invention may be utilised, the generation of continuous cell lines is preferred. For examples of techniques which may be used to derive a continuous cell line from the transgenic animals, see Small, et al., 1985, *Mol. Cell. Biol.* 5: 642-648.

Preferred transgenic animal models of Alzheimer's disease include mice overexpressing glycogen synthase kinase (GSK) (see Lucas et al (2001) *EMBO J.* 20, p 27-39), mice overexpressing mutant alleles of APP or PS1 and double (APP/PS1) transgenic mouse models overexpressing mutant alleles of both APP and PS1. Double transgenic mice resulting from a cross between a mutant APP line Tg2576 and a mutant PS1M146L transgenic line is reported in Holcomb et al., *Nat. Med.* 1998 January; 4(1):97-100).

In further characterising the biological function of the identified proteins, the expression of these proteins may be modulated within the *in vivo* and/or *in vitro* systems, i.e. either overexpressed or underexpressed in, for example, transgenic animals and/or cell lines, and its subsequent effect on the system then assayed. Alternatively, the activity of the identified protein may be modulated by either increasing or decreasing the level of activity in the *in vivo* and/or *in vitro* system of interest, and its subsequent effect then assayed.

The information obtained through such characterisations may suggest relevant methods for the treatment of Alzheimer's disease using the protein of interest. For example, treatment may include a modulation of protein expression and/or protein activity. Characterisation procedures such as those described herein may indicate where such modulation should involve an increase or a decrease in the expression or activity of the protein of interest. Such methods of treatment are discussed below in Section 4.

2. Differentially Expressed Proteins

Identified proteins, which include differentially expressed proteins such as those identified in Section 1 above, are described herein. Specifically, the amino acid sequences of such identified proteins are described. Further, antibodies directed against the identified protein, and cell- and animal-based models by which the identified proteins may be further characterised and utilised are also discussed in this Section.

2.1 Antibodies Specific for Differentially Expressed or Pathway Proteins

The present invention also relates to methods for the production of antibodies capable of specifically recognising one or more differentially expressed or pathway protein epitopes.

Such antibodies may include, but are not limited to, polyclonal antibodies, monoclonal antibodies (mAbs), humanised or chimeric antibodies, single chain antibodies, Fab fragments, F(ab')₂ fragments, fragments produced by a Fab expression library, anti-idiotypic (anti-Id) antibodies, and epitope-binding fragments of any of the above. Such antibodies may be utilised as part of Alzheimer's disease treatment methods, and/or may be used as part of diagnostic techniques whereby patients may be tested for abnormal levels of fingerprint, target, or pathway gene proteins, or for the presence of abnormal forms of such proteins.

For the production of antibodies to a differentially expressed or pathway protein, various host animals may be immunised by injection with a differentially expressed or pathway protein, or a portion thereof. Such host animals may include, but are not limited to, rabbits, mice and rats, to name but a few. Various adjuvants may be used to increase the immunological response, depending on the host species, including active substances such as lysolecithin, Pluronic polyols, polyanions, peptides, oil emulsions, keyhole limpet hemocyanin, dinitrophenol, and potentially useful human adjuvant such as BCG bacille Calmette-Fuerin) and *Corynebacterium parvum*.

Polyclonal antibodies are heterogeneous populations of antibody molecules derived from the sera of animals immunised with an antigen, such as target proteins, or an antigenic functional derivative thereof. For the production of polyclonal antibodies, host animals such as those described above, may be immunised by injection with differentially expressed or pathway protein supplemented with adjuvants as also described above.

Monoclonal antibodies, which are homogeneous populations of antibodies to a particular antigen, may be obtained by any technique, which provides for the production of antibody molecules by continuous cell lines in culture. These include, but are not limited to, the hybridoma technique of Kohler and Milstein (1975, *Nature* 256: 495-497; and U.S. Pat. No. 4,376,110), the human β -cell hybridoma technique (Kosbor, et al., 1983, *Immunology Today* 4: 72; Cole, et al., 1983, *Proc. Natl. Acad. Sci. USA* 80: 2026-2030), and the EBV-hybridoma technique (Cole, et al., 1985, *Monoclonal Antibodies and Cancer Therapy*, Alan R. Liss Inc., pp. 77-96). Such antibodies may be of any immunoglobulin class including IgG, IgM, IgE, IgA, IgD and any subclass thereof. The hybridoma producing the mAb of this invention may be cultivated *in vitro* or *in vivo*. Production of high titers of mAbs *in vivo* makes this the presently preferred method of production.

In addition, techniques developed for the production of 'chimeric antibodies' (Morrison, et al., 1984, *Proc. Natl. Acad. Sci.* 81: 6851-6855; Neuberger, et al., 1984, *Nature* 312: 604-608; Takeda, et al., 1985, *Nature* 314: 452-454) by splicing the genes from a mouse antibody molecule of appropriate antigen specificity together with genes from a human antibody molecule of appropriate biological activity can be used. A chimeric antibody is a molecule in which different portions are derived from different animal species, such as those having a variable region derived from a murine mAb and a human immunoglobulin constant region.

Alternatively, techniques described for the production of single chain antibodies (U.S. Pat. No. 4,946,778; Bird, 1988, *Science* 242: 423-426; Huston, et al., 1988, *Proc. Natl. Acad. Sci. USA* 85: 5879-5883; and Ward, et al., 1989, *Nature* 334: 544-546) can be adapted to produce differentially expressed or pathway protein-single chain antibodies. Single chain antibodies are formed by linking the heavy and light chain fragments of the Fv region via an amino acid bridge, resulting in a single chain polypeptide.

Antibody fragments, which recognise specific epitopes, may be generated by known techniques. For example, such fragments include, but are not limited to, the F(ab')₂ fragments which can be produced by pepsin digestion of the antibody molecule and the Fab fragments which can be generated by reducing the disulfide bridges of the F(ab')₂ fragments. Alternative, Fab expression libraries may be constructed (Huse, et al., 1989, Science 246: 1275-1281) to allow rapid and easy identification of monoclonal Fab fragments with the desired specificity.

3 Assays for Amelioration of Alzheimer's Disease Symptoms

The differentially expressed proteins described herein may be used to test compounds for the ability to prevent or ameliorate Alzheimer's disease.

Such compounds may be tested in human subjects in clinical trials. Any compound which restores the expression of a differentially expressed protein or proteins towards the normal level may be of potential use in treating Alzheimer's disease, i.e. reducing Alzheimer's disease symptoms or slowing the progression of Alzheimer's disease.

With regard to intervention, any treatments that restore or partially restore marker protein expression to normal levels should be considered as candidates for Alzheimer's disease therapeutic intervention. Dosages of test agents may be determined by deriving dose-response curves, as discussed in Section 6 below.

Similarly, any treatments that can prevent the development of Alzheimer's disease or prevent progression to levels of more advanced Alzheimer's disease should be considered as candidates for the Alzheimer's disease therapeutic intervention.

In addition, animal models of Alzheimer's disease, such as those described above, may be used to identify compounds capable of treating Alzheimer's disease symptoms. Such animal models may be used as test substrates for the identification of drugs, pharmaceuticals, therapies and interventions which may be effective in treating such disorders. The response of the animals to the exposure may be monitored by assessing the expression of the marker proteins and comparing it to that of wild-type mice.

Protein expression patterns may be utilised in conjunction with animal model systems to assess the ability of a compound to ameliorate Alzheimer's disease symptoms, or prevent the progression of Alzheimer's disease. For example, the expression pattern of one or more fingerprint proteins may form part of a fingerprint profile, which may then be used in such an assessment. Fingerprint profiles may be characterised for Alzheimer's disease states within the animal model systems. Subsequently, these known fingerprint profiles may be compared to ascertain the effect a test compound has to modify such fingerprint profiles, and to cause the profile to more closely resemble that of a more desirable fingerprint. For example, administration of a compound may cause the fingerprint profile of an Alzheimer's disease model system to more closely resemble the control system, or may prevent further changes in fingerprint profile. Administration of a compound may, alternatively, cause the fingerprint profile of a control system to begin to mimic an Alzheimer's disease state, which may, for example, be used in further characterising the compound of interest, or may be used in the generation of additional animal models.

4. Compounds and Methods for Treatment of Alzheimer's Disease

Described below are methods and compositions whereby Alzheimer's disease symptoms may be ameliorated or the progression of Alzheimer's disease slowed or halted. It is possible that Alzheimer's disease symptoms may be brought

about, at least in part, by an abnormal level of target protein, or by the presence of a target protein exhibiting an abnormal activity. As such, the reduction in the level and/or activity of such target protein would bring about the amelioration Alzheimer's disease symptoms. Techniques for the reduction of target protein gene expression levels or target protein activity levels are discussed in Section 4.1.

Alternatively, it is possible that Alzheimer's disease symptoms may be brought about, at least in part, by the absence or reduction of the level of target protein expression, or a reduction in the level of a target protein's activity. As such, an increase in the level of target protein gene expression and/or the activity of such proteins would bring about the amelioration Alzheimer's disease symptoms. Techniques for increasing target protein gene expression levels or target protein activity levels are discussed in Section 4.2.

4.1 Compounds that Inhibit Expression, Synthesis or Activity of Target Proteins

As discussed above, target proteins involved in Alzheimer's disease may cause such disorders via an increased level of target protein activity. A variety of techniques may be utilised to inhibit the expression, synthesis, or activity of such target genes and/or proteins.

For example, compounds which exhibit inhibitory activity, may be used in accordance with the invention to prevent mild cognitive impairment or Alzheimer's disease symptoms. Such molecules may include, but are not limited to, peptides (such as, for example, peptides representing soluble extracellular portions of target protein transmembrane receptors), phosphopeptides, small organic or inorganic molecules, or antibodies (including, for example, polyclonal, monoclonal, humanised, anti-idiotypic, chimeric or single chain antibodies, and FAb, F(ab')₂ and FAb expression library fragments, and epitope-binding fragments thereof). Techniques for determination of effective doses and administration of such compounds are described below, in Section 6.1. Inhibitory antibody techniques are further described below, in Section 4.1.2.

Further, antisense, siRNA and ribozyme molecules, which inhibit expression of the target protein gene, may also be used in accordance with the invention to inhibit the aberrant target protein gene activity. Such techniques are described below, in Section 4.1.1; triple helix molecules may be utilised in inhibiting the aberrant target protein gene activity.

4.1.1 Inhibitory Antisense, Ribozyme and Triple Helix Approaches

Antisense, ribozyme and triple helix molecules may be designed to reduce or inhibit either wild type, or if appropriate, mutant target protein gene activity. Techniques for the production and use of such molecules are well known to those of skill in the art.

Antisense RNA and DNA molecules act to directly block the translation of mRNA by hybridising to targeted mRNA and preventing protein translation. With respect to antisense DNA, oligodeoxy-ribonucleotides derived from the translation initiation site, e.g. between the -10 and +10 regions of the target gene nucleotide sequence of interest, are preferred.

Ribozymes are enzymatic RNA molecules capable of catalysing the specific cleavage of RNA. (For a review, see Rossi, J., 1994, Current Biology 4: 469-471). The mechanism of ribozyme action involves sequence specific hybridisation of the ribozyme molecule to complementary target RNA, followed by an endonucleolytic cleavage. The composition of ribozyme molecules must include one or more sequences complementary to the target protein mRNA, and must include the well known catalytic sequence responsible for mRNA cleavage. For this sequence, see U.S. Pat. No. 5,093,246. As

such, within the scope of the invention are engineered hammerhead motif ribozyme molecules that specifically and efficiently catalyse endonucleolytic cleavage of RNA sequences encoding target proteins.

Specific ribozyme cleavage sites within any potential RNA target are initially identified by scanning the molecule of interest for ribozyme cleavage sites which include the following sequences, GUA, GUU and GUC. Once identified, short TNA sequences of between 15 and 20 ribonucleotides corresponding to the region of the target protein gene, containing the cleavage site may be evaluated for predicted structural features, such as secondary structure, that may render the oligonucleotide sequence unsuitable. The suitability of candidate sequences may also be evaluated by testing their accessibility to hybridise with complementary oligonucleotides, using ribonuclease protection assays.

RNA interference (RNAi) is a process of sequence-specific, post-transcriptional gene silencing in animals and plants, initiated by double-stranded RNA (dsRNA) that is homologous in sequence to the silenced gene. RNAi is mediated by short double-stranded RNA molecules (small interfering RNAs or siRNAs). siRNAs may be introduced into a cell as short RNA oligonucleotides of 10-15 bp, or as longer dsRNAs which are subsequently cleaved to produce siRNAs. The RNA may be introduced into the cell as RNA, or may be transcribed from a DNA or RNA vector.

siRNA molecules may be synthesized using standard solid or solution phase synthesis techniques which are known in the art. Alternatively, siRNA molecules or longer dsRNA molecules may be made recombinantly by transcription of a nucleic acid sequence, preferably contained within a vector as described below.

Another alternative is the expression of a short hairpin RNA molecule (shRNA) in the cell. shRNAs are more stable than synthetic siRNAs. A shRNA consists of short inverted repeats separated by a small loop sequence. One inverted repeat is complementary to the gene target. The shRNA is then processed into an siRNA which degrades the target gene mRNA and suppresses expression. shRNAs can be produced within a cell by transfecting the cell with a DNA construct encoding the shRNA sequence under control of a RNA polymerase III promoter, such as the human H1 or 7SK promoter. Alternatively, the shRNA may be synthesised exogenously and introduced directly into the cell. Preferably, the shRNA sequence is between 40 and 100 bases in length, more preferably between 40 and 70 bases in length. The stem of the hairpin is preferably between 19 and 30 base pairs in length. The stem may contain G-U pairings to stabilise the hairpin structure.

Nucleic acid molecules to be used in triplex helix formation for the inhibition of transcription should be single stranded and composed of deoxynucleotides. The base composition of these oligonucleotides must be designed to promote triple helix formation via Hoogsteen base pairing rules, which generally require sizeable stretches of either purines or pyrimidines to be present on one strand of a duplex. Nucleotide sequences may be pyrimidine-based, which will result in TAT and CGC⁺ triplets across the three associated strands of the resulting triple helix. The pyrimidine-rich molecules provide base complementary to a purine-rich region of a single strand of the duplex in a parallel orientation to that strand. In addition, nucleic acid molecules may be chosen that are purine-rich, for example, containing a stretch of G residues. These molecules will form a triple helix with a DNA duplex that is rich in GC pairs, in which the majority of the

purine residues are located on a single strand of the targeted duplex, resulting in GGC triplets across the three strands in the triplex.

Alternatively, the potential sequences that can be targeted for triple helix formation may be increased by creating a so-called "switchback" nucleic acid molecule. Switchback molecules are synthesised in an alternating 5'-3',3'-5' manner, such that they base pair with first one strand of a duplex and then the other, eliminating the necessity for a sizeable stretch of either purines or pyrimidines to be present on one strand of a duplex.

Anti-sense RNA and DNA, siRNAs, ribozyme and triple helix molecules of the invention may be prepared by any method known in the art for the synthesis of DNA and RNA molecules. They include techniques for chemically synthesising oligodeoxyribonucleotides and oligo-ribonucleotides well known in the art such as, for example, solid phase phosphoramidite chemical synthesis. Alternatively, RNA molecules may be generated by in vitro and in vivo transcription of DNA sequences encoding the antisense RNA molecule. Such DNA sequences may be incorporated into a wide variety of vectors, which incorporate suitable RNA polymerase promoters such as the T7 or SP6 polymerase promoters. Alternatively, antisense cDNA constructs that synthesise antisense RNA constitutively inducibly, depending on the promoter used, can be introduced stably into cell lines.

4.1.2 Antibodies for the Inhibition of Target Protein

Antibodies that are both specific for target protein and interfere with its activity may be used to inhibit target protein function. Where desirable, antibodies specific for mutant target protein, which interferes with the activity of such mutant target product, may also be used. Such antibodies may be generated, using standard techniques described in Section 2. above, against the proteins themselves or against peptides corresponding to portions of the proteins. The antibodies include, but are not limited to, polyclonal, monoclonal, Fab fragments, single chain antibodies, chimeric antibodies, etc.

In instances where the target gene protein is intracellular and whole antibodies are used, internalising antibodies may be preferred. However, lipofectin or liposomes may be used to deliver the antibody or a fragment of the Fab region, which binds to the target protein epitope into cells. Where fragments of the antibody are used, the smallest inhibitory fragment, which binds to the target protein's binding domain, is preferred. For example, peptides having an amino acid sequence corresponding to the domain of the variable region of the antibody that binds to the target protein may be used. Such peptides may be synthesised chemically or produced via recombinant DNA technology using methods well known in the art (e.g. see Creighton, 1983, supra; and Sambrook et al, 1989, supra).

Alternatively, single chain neutralising antibodies, which bind to intracellular target protein epitopes, may also be administered. Such single chain antibodies may be administered, for example, by expressing nucleotide sequences encoding single-chain antibodies within the target cell population by utilising, for example, techniques such as those described in Marasco et al (Marasco, W. et al, 1993, Proc. Natl. Acad. Sci. USA, 90: 7889-7893).

In instances where the target protein is extracellular, or is a transmembrane protein, any of the administration techniques described below, in Section 6, which are appropriate for peptide administration may be utilised to effectively administer inhibitory target protein antibodies to their site of action.

4.2 Methods for Restoring Target Protein Activity

Target proteins that cause Alzheimer's disease may be underexpressed in Alzheimer's disease disorder situations.

Alternatively, the activity of target protein may be diminished, leading to the development of Alzheimer's disease symptoms. Described in this Section are methods whereby the level of target protein may be increased to levels wherein Alzheimer's disease symptoms are prevented or ameliorated. The level of target protein activity may be increased, for example, by either increasing the level of target protein present or by increasing the level of active target protein which is present.

For example, a target protein, at a level sufficient to ameliorate Alzheimer's disease symptoms may be administered to a patient exhibiting such symptoms. Any of the techniques discussed below, in Section 6, may be utilised for such administration. One of skill in the art will readily know how to determine the concentration of effective, non-toxic doses of the normal target protein, utilising techniques such as those described below.

Further, patients may be treated by gene replacement therapy. One or more copies of a normal target protein gene or a portion of the gene that directs the production of a normal target protein with target protein gene function, may be inserted into cells, using vectors which include, but are not limited to, adenovirus, adeno-associated virus, and retrovirus vectors, in addition to other particles that introduce DNA into cells, such as liposomes. Additionally, techniques such as those described above may be utilised for the introduction of normal target protein gene sequences into human cells.

Cells, preferably autologous cells, containing normal target protein gene sequences may then be introduced or reintroduced into the patient at positions which allow for the prevention or amelioration of Alzheimer's disease symptoms. Such cell replacement techniques may be preferred, for example, when the target protein is a secreted, extracellular protein.

Additionally, antibodies may be administered which specifically bind to a target protein and by binding, serve to, either directly or indirectly, activate the target protein function. Such antibodies can include, but are not limited to, polyclonal, monoclonal, FAb fragments, single chain antibodies, chimeric antibodies and the like. The antibodies may be generated using standard techniques such as those described above, in Section 2.3, and may be generated against the protein themselves or against proteins corresponding to portions of the proteins. The antibodies may be administered, for example, according to the techniques described above.

5. Pharmaceutical Preparations and Methods of Administration

The identified compounds, nucleic acid molecules and cells that affect target protein expression, synthesis and/or activity can be administered to a patient at therapeutically effective doses to prevent or to treat or to ameliorate Alzheimer's disease. A therapeutically effective dose refers to that amount of the compound sufficient to result in amelioration of symptoms Alzheimer's disease, or alternatively, to that amount of a nucleic acid molecule sufficient to express a concentration of protein which results in the amelioration of such symptoms.

5.1 Effective Dose

Toxicity and therapeutic efficacy of such compounds can be determined by standard pharmaceutical procedures in cell cultures or experimental animals, e.g. for determining by ED₅₀ (the dose therapeutically effective in 50% of the population) and by determining the ED₅₀ of any side-effects (toxicity—TD₅₀). The dose ratio between toxic and therapeutic effects is the therapeutic index and it can be expressed as the ratio TD₅₀/ED₅₀. Compounds, which exhibit large therapeutic indices, are preferred. While compounds that exhibit toxic

side effects may be used, care should be taken to design a delivery system that targets such compounds to the site of affected tissue in order to minimise potential damage to unaffected cells and, thereby, reduce side effects.

The data obtained from the animal studies can be used in formulating a range of dosage for use in humans. The dosage of such compounds lies preferably within a range of circulating concentrations that include the ED₅₀ with little or no toxicity. The dosage may vary within this range depending upon the dosage form employed and the route of administration utilised.

5.2 Formulations and Use

Pharmaceutical compositions for use in accordance with the present invention may be formulated in conventional manner using one or more physiologically acceptable carriers or excipients.

Thus, the compounds and their physiologically acceptable salts and solvates may be formulated for administration by inhalation or insufflation (either through the mouth or the nose) or oral, buccal, parenteral and rectal administration.

For oral administration, the pharmaceutical compositions may take the form of, for example, tablets or capsules prepared by conventional means with pharmaceutically acceptable excipients such as binding agents (e.g. pre-gelatinised maize starch, polyvinylpyrrolidone or hydroxypropyl methyl-cellulose); fillers (e.g. lactose, microcrystalline cellulose or calcium hydrogen phosphate); lubricants (e.g. magnesium stearate, talc or silica); disintegrants (e.g. potato starch or sodium starch glycollate); or wetting agents (e.g. sodium lauryl sulphate). The tablets may be coated by methods well known in the art. Liquid preparations for oral administration may take the form of, for example, solutions, syrups or suspensions, or they may be presented as a dry product for constitution with water or other suitable vehicle before use. Such liquid preparations may be prepared by conventional means with pharmaceutically acceptable additives such as suspending agents (e.g. sorbitol syrup, cellulose derivatives or hydrogenated edible fats); emulsifying agents (e.g. lecithin or acacia); and preservatives (e.g. methyl or propyl-p-hydroxybenzoates or sorbic acid). The preparations may also contain buffer salts, flavouring, colouring and sweetening agents as appropriate.

Preparations for oral administration may be suitably formulated to give controlled release of the active compound.

For buccal administration the compositions may take the form of tablets or lozenges formulated in conventional manner.

For administration by inhalation, the compounds for use according to the present invention are conveniently delivered in the form of an aerosol spray presentation from pressurised packs or a nebuliser, with the use of a suitable propellant, e.g. dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, carbon dioxide or other suitable gas. In the case of a pressurised aerosol the dosage unit may be determined by providing a valve to deliver a metered amount. Capsules and cartridges of, e.g. gelatin, for use in an inhaler or insufflator may be formulated containing a powder mix of the compound and a suitable powder base such as lactose or starch.

The compounds may be formulated for parenteral administration by injection, e.g. by bolus injection or continuous infusion. Formulations for injection may be presented in unit dosage form, e.g. in ampoules or in multi-dose containers, with an added preservative. The compositions may take such forms as suspensions, solutions or emulsions in oily or aqueous vehicles, and may contain formulatory agents such as suspending, stabilising and/or dispersing agents. Alterna-

tively, the active ingredient may be in powder form for constitution with a suitable vehicle, e.g. sterile pyrogen-free water, before use.

The compounds may also be formulated in rectal compositions such as suppositories or retention enemas, e.g. containing conventional suppository bases such as cocoa butter or other glycerides.

In addition to the formulations described previously, the compounds may also be formulated as a depot preparation. Such long acting formulations may be administered by implantation, for example, subcutaneously or intramuscularly) or by intramuscular injection. Thus, for example, the compounds may be formulated with suitable polymeric or hydrophobic materials (for example, as an emulsion in an acceptable oil) or ion exchange resins, or as sparingly soluble derivatives, for example, as a sparingly soluble salt.

The compositions may, if desired, be presented in a pack or dispenser device, which may contain one or more unit dosage forms containing the active ingredient. The pack may, for example, comprise metal or plastic foil, such as blister pack. The pack or dispenser device may be accompanied by instructions for administration.

6. Diagnosis of Alzheimer's Disease

A variety of methods may be employed for the diagnosis of Alzheimer's disease, monitoring progression of mild cognitive impairment and Alzheimer's disease, the predisposition to Alzheimer's disease, and for monitoring the efficacy of any Alzheimer's disease compounds during, for example, clinical trials and for monitoring patients undergoing clinical evaluation for the treatment of Alzheimer's disease. The differentially expressed and fingerprint proteins can also be used to define the nature or degree of Alzheimer's disease to aid in the identification and/or selection of treatments for the disorder.

Alzheimer's disease is characterised by a progressive, insidious onset, two or more deficits in cognitive function, and the absence of any other illnesses that could account for the dementia

In addition to memory loss, there may be disorientation, poor attention span, and language impairment. There is likely to be a decline in the activity of daily living, and possibly also impaired perception and personality changes. Behavioural symptoms include delusions, aggression, agitation, anger, wandering, hallucinations, and sleep disturbance.

A simple test assessing orientation, registration, calculations and attention, recall, language, and visual-spatial function may be used for an initial diagnosis.

Structural imaging by standard CT or MRI may also be used. Typically a non-contrast head CT scan suffices, but MRI is preferred for those who have hypertension or diabetes, who are at risk for cerebral vascular disease.

Alzheimer's disease may be confirmed histologically by autopsy or brain biopsy showing neurofibrillary tangles and senile plaques.

Identifying individuals at risk from Alzheimer's disease may involve diagnosis of mild cognitive impairment (MCI). (MCI) may be a transitional state between normal aging and dementia. There are different types of MCI. There may be cognitive impairment in multiple areas of cognitive function, in addition to memory. In some cases, memory is normal but some other domain of cognitive function is abnormal.

Amnesic MCI appears to be a risk state for the development of Alzheimer's disease. Amnesic impairment is defined by subjective memory complaints. These patients have poor memory performance for their age and education on formal testing when compared to age-matched peers. General cognitive functions and the ability to perform the activities of daily living should be entirely normal. The amnesic type of

MCI is associated with hippocampal atrophy, neurofibrillary tangles in the medial temporal lobes, and elevated levels of Tau in the cerebrospinal fluid (CSF).

Methods for diagnosing Alzheimer's disease or predisposition to Alzheimer's disease may also, for example, utilise reagents such as the differentially expressed and fingerprint proteins described above, and antibodies directed against differentially expressed, as described above. Specifically, such reagents may be used for the detection of either an over- or an under-abundance of target protein relative to the normal state.

The methods described herein may be performed, for example, by utilising pre-packaged diagnostic kits comprising at least one specific differentially expressed/fingerprint protein or anti-differentially expressed/fingerprint protein antibody reagent described herein, which may be conveniently used, e.g. in clinical settings, to diagnose patients exhibiting Alzheimer's disease symptoms.

Any cell type, tissue or body fluid in which the fingerprint protein is expressed may be utilised in the diagnostics described herein. Examples of suitable samples types include cell samples, tissue samples, and fluid samples such as blood, urine, serum, saliva, cerebrospinal fluid or plasma.

Among the methods which can be utilised herein, are methods for monitoring the efficacy of compounds in clinical trials for the treatment of Alzheimer's disease. Such compounds can, for example, be compounds such as those described above, in Section 4. Such a method comprises detecting, in a patient sample, a protein, which is differentially expressed in the Alzheimer's disease state relative to its expression in a normal state.

During clinical trials, for example, the expression of a single differentially expressed protein, or alternatively, a fingerprint pattern of a cell involved in Alzheimer's disease can be determined in the presence or absence of the compound being tested. The efficacy of the compound can be followed by comparing the expression data obtained to the corresponding known expression patterns in a normal state. Compounds exhibiting efficacy are those which alter the protein expression and/or the fingerprint pattern to more closely resemble that of the normal state, or which stabilise protein expression and/or the fingerprint pattern i.e. prevent progression of the disease.

The detection of the protein differentially expressed in the Alzheimer's disease state relative to their expression in a normal state can also be used for monitoring the efficacy of potential compounds for the treatment of Alzheimer's disease during clinical trials. During clinical trials, for example, the level and/or activity of the differentially expressed protein can be determined in relevant cells and/or tissues and/or body fluids in the presence or absence of the compound being tested. The efficacy of the compound can be followed by comparing the protein level and/or activity data obtained to the corresponding known levels/activities for the cells and/or tissues and/or body fluids in a normal state. Compounds exhibiting efficacy are those which alter the pattern of the cell and/or tissue sample and/or body fluid from an Alzheimer's disease subject to more closely resemble that of the normal state or which stabilise the pattern i.e. prevent progression of the disease.

EXPERIMENTAL

Subjects

The study population is derived from a large, longitudinally assessed, community based population of people with AD (NINCDS-ADRDA probable), other dementias and normal elderly persons. Samples are available on over 1000

subjects, all whom have detailed clinical assessment. Clinical research data includes systematic diagnostic, cognitive and behavioural assessments. Approximately 50 ml blood (4×10 ml in BD vacutainer K3E 15% tubes and 1×10 ml in exetainer) is drawn from each subject. Subjects have had no food or fluid intake for more than 2 hours prior to collection. One BD vacutainer K3E (plasma) and exetainer (serum) is used for proteomics study. The serum/plasma samples collected for proteomics are spun at 3000 rpm for 8 min within 2 h of collection.

Analysis

Serum/plasma samples were lysed and rehydrated in a 2D lysis buffer consisting of 8M Urea, 2% w/v CHAPS, 0.5% IPG Pharmalyte (pH 3-10; Amersham Biotech, UK). The lysed samples were then subjected to isoelectrofocusing 18 cm 3-10 NL Immobiline pH gradient strips. IPG electrofocusing of the rehydrated strips was carried out for 16 h using the following protocol: S1 500V step-n-hold (s/h; i.e. the electric current applied to the strip is gradually increased in steps holding at particular settings for the times indicated) for 1 h; S2 500V s/h for 2 h; S3 1000V gradient (G) for 1 h; S4 1000V s/h for 2 h; S5 8000V G for 2 h and a final step S6 8000V s/h for 8 h with the IPGphor™.

Electrofocused IPG strips were then equilibrated in a SDS equilibration buffer (50 mM Tris-HCl pH8.8, 6M urea, 30% (v/v) glycerol, 2% SDS, and trace amount of bromophenol blue) with 10 mg/ml dithiothreitol (DTT) for 20 min, followed by 20 min step with 25 mg/ml Iodoacetamide. The equilibrated strips were then separated on a 10% acrylamide second dimension electrophoresis gel using the Ettan Dalt II system.

Following the electrophoresis the gels were placed in separate staining boxes and fixed using 40% ethanol/10% acetic acid for 1 h at room temperature and then stained according to Hochstrasse protocol (Table 1). Gel analysis was performed using the Melanie 3 software and Mann and Whitney rank sum test and False Discovery Rate statistical analysis was carried out to compare subject groups.

TABLE 1

Hochstrasse staining protocol	
Staining step	Time
Fix 40% ethanol/10% acetic acid	1 h
Soak in 5% ethanol/5% acetic acid	3 hr or overnight
Wash in water	5 min
Soak in 0.5M Sodium acetate, 1% glutaraldehyde	1.5 h
Wash	4 × 15 min
Soak in 0.05% Naphthalene sulphonic acid	2 × 30 min
Rinse in water	4 × 15 min
Silver stain (12 g silver, 20 ml ammonium hydroxide and 3 ml 10M sodium hydroxide)	25 min
Wash	4 × 4 min
Develop (0.005% citric acid and 0.1% formaldehyde)	As required
Stop solution (5% tris and 2% acetic acid)	1-2 h
Storage solution (35% ethanol and 5% glycerol)	

Sample preparation In-gel reduction, alkylation and digestion (with trypsin) was performed prior to subsequent analysis by mass spectrometry. Cysteine residues were reduced with DTT and derivatized by treatment with iodoacetamide to form stable carbamidomethyl (CAM) derivatives. Trypsin digestion was carried out overnight at room temperature after an initial 1 hr incubation at 37° C.

MALDI-TOF Mass Spectrometry

The digested sample (3 µl) was desalted and concentrated using ZipTipC18 microtips (Millipore). Peptides were eluted in 4 µl 50% acetonitrile/0.1% trifluoroacetic acid. 0.5 µl was then loaded onto a target plate with 0.5 µl matrix (α -Cyano-4-hydroxy-cinnamic acid). Peptide mass fingerprints were acquired using a Voyager De-Pro, MALDI-TOF mass spectrometer (Applied Biosystems). The mass spectra were acquired in reflectron mode with delayed extraction. An autolytic tryptic peptide of mass 2163.0569 Da was then used to lock mass the acquired spectra, to achieve a mass accuracy of better than 30 ppm.

LC/MS/MS

Peptides were extracted from the gel pieces by a series of acetonitrile and aqueous washes. The extract was pooled with the initial supernatant and lyophilised. Each sample was then resuspended in 6 µL of 50 mM ammonium bicarbonate and analysed by LC/MS/MS. Chromatographic separations were performed using an Ultimate LC system (Dionex, UK). Peptides were resolved by reverse phase chromatography on a 75 µm C18 PepMap column. A gradient of acetonitrile in 0.05% formic acid was delivered to elute the peptides at a flow rate of 200 nl/min. Peptides were ionised by eLectrospray ionisation using a Z-spray source fitted to a QTOFmicro (Waters Corporation). The instrument was set to run in automated switching mode, selecting precursor ions based on their intensity, for sequencing by collision-induced fragmentation. The MS/MS analyses were conducted using collision energy profiles that were chosen based on the m/z and the charge state of the peptide.

Results

Analysis of all control group (n=50) and case group (n=50) 2D gel images and subjecting them to statistical analysis. A total of 16 protein spots show a significant result (p<0.05) (see FIG. 6).

The results shown in FIG. 6 are unambiguous matches as they are based on exact matching of multiple MS/MS spectra. The sequence of selected proteins showing the peptide coverage obtained is given in FIGS. 8 to 10.

Class Prediction Using Peptide Fingerprinting

A class prediction analysis was performed in order to determine whether the pattern of peptide spots on 2DGE could predict caseness as determined clinically. Support Vector Machines (SVM), a supervised machine learning algorithm for prediction of class set in a group based upon a training set of data¹³, was used. SVM is most typically used in microarray analyses. However the statistical challenges are similar for proteomics and SVM has previously been used as a class prediction model for various proteomic studies^{14,15}. Using GeneSpring (Silicon Genetics) the original 25 cases and 25 controls were designated as a training set and then the replication 25 cases and 25 controls designated as a test set. All identified proteins were used as possible identifiers and with the parameters Polynomial Dot Product Order 1 and Diagonal Scaling Factor 1; 34 of the 50 test-samples were correctly identified as being either cases or controls. Sensitivity was 56% and specificity 80% using SVM analysis of 2DGE data alone.

Identification of Peptides that Differentiate Between Cases and Controls

The normalised spot optical density in both the initial set of cases and controls and the replication set was compared. Mean differences between patients and controls at each spot were compared using the Wilcoxon rank-sum (Mann-Whitney) tests. The p-values for the nul hypothesis of no mean differences were saved, sorted by increasing value and ranked. A false discovery rate index (FDR) was computed as

the ratio of the rank number and the theoretical probability (which is the rank number divided by the total number of spots). Fifteen spots were identified to have a FDR of less than 0.50. These were then identified using LC-MS/MS (FIG. 6).

Correlation of Peptide Spots with Clinical Parameters

Although the cases and controls were similarly aged it was possible that the observed peptide or spot differences were due to an association with age, gender or APOE genotype. A correlation analysis was thus performed for the 15 spots that differed between cases and controls in all 100 subjects with age, gender and APOE genotype. Data was first scaled to unit variance so as to standardise the scales upon which the variables were compared (i.e. each value was divided by the standard deviation of all the values for that particular variable). The Pearson correlation coefficient was then calculated. Cases with missing values were excluded pairwise. There were no strong correlations of any spot with age, gender or APOE. Two spots weakly correlated with age, two with gender and one with APOE genotype.

An ideal biomarker would not only be different between cases and controls but would be a marker of disease progression. The 15 spots showing case-control differences in all 50 cases were thus correlated with duration of dementia and severity as measured by MMSE and GDS. Two spots correlated moderately and significantly with measures of disease progression and global dementia severity ($r^2=0.29$ with spot 177) and duration of disease ($r^2=0.29$ with spot 166). Thus, one peptide—an Ig lambda chain C region (spot 177) correlates with global dementia severity. The other marker of disease progression examined, duration, shows a negative correlation with albumin (Spot 165).

Pre-Depletion Analysis

In these experiments, human plasma samples were depleted to remove the 6 most abundant proteins before the 2D gel electrophoresis step.

Methods

60 human plasma samples (30 Controls and 30 disease subjects) were depleted using a removal column from Agilent. The samples were separated by 2D electrophoresis (pH 3-10 NL, 10% SDS-PAGE, 75 μ g protein load). Gels were silver-stained, scanned (8 bit, 200 dpi) and quantitatively analysed with Progenesis. To pick gel plugs from preparative gels, several control samples were mixed together and 3 gels run (2 gels with 205 μ g protein load and 1 gel with 350 μ g protein load). The same strategy was used with disease samples to run preparative gels. Protein spots were then destained, trypsinated and polypeptides were spotted onto MALDI target with Spot handling workstations (GE Amersham Biosciences). Peptide profiles generated were analysed with Ms-Fit programme in combination with the Swiss-Prot database.

Results

Gel images of proteins extracted from control and disease samples were analysed with Progenesis (v2005). Each group (Control and Disease) were based on 29 analytical gels. Spot detection, matching were performed with Progenesis, then the spot data were exported to Excel and a macro developed in-house was used to calculate coefficient of variation (CV %), T-Test and Regulation factor or change.

11 spots were selected for analysis based on the following selection criteria: spots have to be found within at least 60% of gels, 2-fold up/down regulation and p value < 0.005. FIG. 11 displays the location of these 11 spots in the reference gel. This image corresponds to the 2D profile of proteins extracted from a control sample. The normalised volumes of the 11 spots detected in gels was analysed and is given in FIG. 12.

All protein spots were picked from 3 to 6 different preparative gels and submitted to MS analyses. All protein spots were successfully identified as shown in FIG. 12. In the down regulated spots, we found two spots of alpha-2-macroglobulin precursor (174; 178), one spot of inter-alpha-trypsin inhibitor heavy chain H4 precursor (232), one spot of a mix of complement C3 precursor with clusterin precursor (712) as indicated in grey in FIG. 12 and one spot of complement C3 precursor (713). In the up regulated spots, we found two spots of complement C4 precursor (652; 832), one spot of actin (675) and three spots of haptoglobin precursor (702; 703; 706).

To estimate the coverage of proteins identified and to discriminate the different chains or isoforms, for each spot, a common list of peptide masses was established. This list regroups all peptide masses matched corresponding to the same spot picked in 3 to 6 preparative gels. The amino acids belonging to the peptides matched are underlined in FIGS. 13 to 19.

Discussion

The 11 spots analysed identified 7 regulated proteins between control- and disease samples, namely alpha-2-macroglobulin, inter-alpha-trypsin inhibitor heavy chain H4, complement C3, complement C4, actin cytoplasmic and haptoglobin.

Alpha-2-macroglobulin protein is able to inhibit all four classes of proteinases by a unique "trapping" mechanism. The observed molecular weight of the gel spots (~100 kDa, FIG. 11, spots 174; 178), matched by PMF, cover mainly the N-terminus of the protein (FIG. 14). The protein identified may thus correspond to a fragment of the full-length sequence of alpha-2-macroglobulin. As spots identified as alpha-2-macroglobulin belong to the same chain of spots (FIG. 11), it is possible that the difference between the two spots may be due to a post-translation modification.

There are two isoforms of inter-alpha-trypsin inhibitor heavy chain H4. Isoform 1 has 930 amino-acids and isoform 2 has 914 amino-acids. This protein is cleaved by plasma kallikrein to yield 100 kDa and a 35 kDa fragments. The resulting 100 kDa fragment is further converted to a 70 kDa fragment. The masses matched by PMF cover the sequence up to amino acid (aa) 688. This sequence corresponds to isoform 1 and may include the 70 kDa fragment and a short potentially active peptide. In this case, there is good agreement between the theoretical molecular weight and pI (74 kDa and 6.04 respectively) and the observed ones from the gel spot (see FIG. 11, spot 232).

Complement C3 precursor plays a central role in the activation of the complement system. This protein contains two chains (alpha and beta). We identified peptide masses covering the sequence from aa 714 to aa 1360 (FIG. 15), which corresponds to the alpha chain of complement C3. The theoretical molecular weight and pI of the alpha chain (115 kDa and 5.55 respectively) are not in agreement with the observed ones from the gel spots (see FIG. 11, spots 712, 713). The alpha chain is processed into different fragments. It appears that a temporary peptide appearing during the activation of complement system. As spots identified as complement C3 belong to the same chain of spots (FIG. 11), it is possible that the difference is due to a post-translational modification.

Complement C4 plays a central role in the activation of the classical pathway of the complement system. This protein contains three chains (alpha, beta and gamma). We identified peptide masses covering the alpha and beta chains for spot 832 and only alpha chain for spot 652 (FIG. 17). The theoretical molecular weights and pIs of these chains differ from the observed ones from the gel spots (see FIG. 11, spots 652;

832). As for complement C3, clusterin precursor protein contains two chains (alpha and beta). We identified peptide masses covering the alpha and beta chains (FIG. 16). The theoretical molecular weight and pI of clusterin (50 kDa and 5.89 respectively) are in agreement with those from the gel (FIG. 11, spot 712). It appears the full-length protein was identified.

Surface Enhanced Laser Desorption Ionisation Time of Flight Mass Spectrometry [SELDI-TOF-MS].

SELDI-TOF-MS and ProteinChip technology were combined to identify protein peaks differing between Alzheimer's and control subjects, followed by extraction of material from the chips to allow further characterisation of the material and identification of the components present.

METHOD (SELDI Analysis)

The SELDI analysis comprises of a comparison of AD cases and control samples and data has been obtained for both a set of individual samples as well as a pooled set of samples. In each case spectral profiles of sera from control and AD cases were compared.

A). Analysis of a Set of Individuals

Control and AD sera from individuals were run on Q10-SAX2 chips:

n=4 control

n=4 AD

Serum samples were prepared fresh by diluting 20 µl serum with 30 µl SELDI lysis buffer. Five microlitres of sample were spotted onto each spot as necessary.

The chips were processed using the following protocol:

Chip Preparation

A hydrophobic ring is drawn around each spot using a PAP pen and the PAP allowed to dry thoroughly by placing chip on the SELDI machine for up to 25 minutes.

Sample Preparation

Serum diluted in SELDI lysis buffer using a 40:60 ratio (40 µl serum+60 µl lysis buffer). Typically, this dilution will render the sample at a 20 mg/mL to 30 mg/mL concentration. Therefore, using a 5 µL aliquot of the lysis buffer sample will enable between 100 µg to 150 µg protein to be loaded on each spot.

Samples are vortexed and incubated on ice until ready to use, then briefly centrifuged samples immediately before use (30 secs, 14,000 rpm).

Chip Equilibration

The chip is placed in a 15 mL Falcon tube and 10-15 mL 100 mM Tris buffer pH 9 at room temperature added, then mixed on a rotary mixer for 5 minutes. The procedure is repeated twice.

Sample Application

After the last equilibration step, the chip is removed and dried carefully with soft tissue. 5 µL of sample is pipetted onto each spot, the chip is placed in a sealed humidity chamber and placed on a shaker for 30 minutes.

Chip Washing

After incubation, the sample is carefully removed from each spot and the chip replaced in the Falcon tube. 10-15 mL 100 mM Tris buffer pH 9 is added, and the Falcon tube mixed on a rotary mixer and for 5 minutes. This is repeat four more times, then the chip washed twice in double distilled water.

Chip Drying

After the last wash step, the chip is removed and dried carefully with soft tissue, then left to air-dry at room temperature for 25 minutes.

SPA Application

2×0.6 µL saturated SPA matrix (freshly made) is pipette onto each spot. The first application is allowed to dry before applying the second 0.6 µL aliquot. The SPA is then left to dry for up to 10 minutes on the SELDI machine.

The chips are then read on the SELDI machine.

The following criteria were applied for data analysis:

Clustering criteria: 5 s/n; 100% spectra; 0.3% mass; 2 s/n; add est. peaks.

Normalisation: Total ion count between 3,000 and 30,000

Da only.

Results

Spot to spot reproducibility between loadings of the same sample was very good. Good correlation was achieved.

Patient to patient variability in both control and dementia groups was observed. This may be due to differences in protein amount as well as idiosyncratic differences. Using very stringent clustering, 3 peaks were found to be statistically significant (p=0.05) and these were visually verified to check validity. The three peaks of interest (see FIGS. 1-3) are as follows:

Mr 6,430 Da abundance in AD	1.62 fold increase in p = 0.027
Mr 14,640 Da abundance in AD	2.29 fold increase in p = 0.036
Mr 27,147 Da abundance in AD	2.82 fold increase in p = 0.004574

B) Analysis of Pooled Sets

A set of pooled samples were analysed using exactly the same methods and criteria as described above. Here, however, we analysed 3 pooled AD samples versus 3 pooled controls where each pool contains serum from at least 25 individuals. In this manner we have encompassed samples from over 75 individuals with AD and compared them against a control cohort representing 75 number of individuals. Pooled groups are described as: AD Pool 1, 2 and 3 comprising of 25, 25 and 25 unique individuals respectively. Similarly, the pooled controls are described as: Control Pool 1, 2 and 3 comprising of 25, 25 and 25 unique individuals respectively.

Results

Using very stringent clustering, 1 peak was found to be statistically significant (p=0.05) and this was visually verified to check validity.

The peak of interest (see FIG. 4) is as follows:

Mr 14,646 Da abundance in AD	1.72 fold increase in p = 0.037
------------------------------	---------------------------------

SELDI Peak Identification Strategy

The differentially expressed proteins identified by SELDI analysis were further analysed by SDS-PAGE. Bands corresponding to the MW of differentially expressed proteins were excised for analysis by mass spectroscopy.

Material was extracted from chips Q10 854 & 855 ("individual" samples) by boiling for 10 minutes in Laemmli buffer and control and disease spots were pooled into separate Eppendorf tubes. Extracted material was separated using SDS-PAGE (18%, tris-glycine, Novex) and the gel was initially stained with Colloidal Coomassie Blue (CCB) but no bands were visualised. Subsequently the same gel was restained using modified (MS-compatible) silver stain (FIG. 5).

Six bands, between 11 and 20 kDa, were visualised and these were excised from the 1st control lane for analysis by LC/MS/MS as described above.

Identified proteins are shown in FIG. 7.

Further Analysis of Identified Proteins

Apolipoprotein A-IV

Sequence coverage obtained for apolipoprotein A-IV (P06727) is shown in FIG. 9 for the 14.6 kDa band isolated on the Q10 SAX2 SELDI chip

The molecular weight of the biomarker of interest observed within the SELDI profiling experiments was determined to be 14640+/-6 Da. The 14.6 kDa species is thought to be a

fragment of ApoA-IV based on the facts that the intact protein should be observed at higher mass (45 kDa) and that the peptides observed in the LC/MS/MS analysis only represented the C-terminal region of the protein. The observed molecular weight is in good agreement with the average molecular weight of 14636 Da predicted for residues 270-396 of the sequence defined for apolipoprotein A-IV within the Swiss Prot database entry P06727.

Both authentic full length apolipoprotein A-IV and a C-terminal fragment of apolipoprotein A-IV comprising of residues 270-396 may thus represent serum biomarkers of Alzheimer's disease.

Complement C4 Precursor

Sequence coverage obtained for Complement C4 precursor (P01028) in 2DE spot 164 is shown in FIG. 10. Spot 164 was identified on the basis of several peptides indicated in underlined bold and this defines the protein in Spot 164 as a C-Terminal fragment extending from residues 1466-1744.

Quantitative Protein Sequence Tag (qPST) Analysis

10 disease samples and 10 control samples were individually immunodepleted for the 6 most highly abundant proteins. 2 pools consisting of either the disease or the control samples were generated and applied to the qPST procedure (precleavage with CNBr, labelling with dimethylglycine, trypsin and fractionation by strong cation exchange). The obtained SCX fractions were analysed by LC-MS and LC-MS/MS using the QTOF-II instrument following the standard approach (LC-MS and LC-MS/MS by three different data acquisition methods)

Results

Identification of Proteins

As stated above, three different MS/MS acquisition strategies were employed:

1. Data Dependent Analysis to obtain as many as possible peptide ID's (1 mass window).

2. Data acquisition by an 'include list' containing regulated pairs, ie peptides whose intensity varied between disease and control samples (regulation criteria: $\geq 2/\geq 0.5$)

3. Data acquisition by an 'include list' containing non-paired MS-signals.

Taking all results from these three approaches into account and correcting them for redundancy, 88 protein IDs were obtained.

Directed Searching for Regulated Proteins by Include List (Pairs with a Regulation $\geq 2.0/\geq 0.5$) MS/MS Strategy and Crossmatching:

8 peptides were identified which could be crossmatched to regulations. These 8 peptides represent five proteins (the peptide grouping to obtain protein ID's was achieved by the ProteinProphet algorithm).

The ID's of these five proteins are shown in FIG. 8.

The 2 peptides which represent protein 1 also occur in Ig alpha-1 chain C region, so that the protein ID's 1 and 2 in fact represent one ID (Ig alpha-1 chain C region).

The hypothetical protein DKFZP686C02220 is a unique one (in fact, one peptide is unique, the second one can occur in several proteins). This protein has typical signatures of immunoglobulins (regarding InterPro entries), and the second peptide also occurs in Ig alpha-2 chain C region.

The proteins 4 and 5 represent one protein ID (haptoglobin precursor) because both peptides occur also in haptoglobin precursor, but the corresponding peptides were grouped as individual proteins by the algorithm used.

Validation of APO-AIV Fragments Using Western Blotting

Western blotting has been undertaken to confirm that the 14.6 kDa species was a fragment of APO-AIV.

Plasma samples were diluted 1:10 with double distilled water and assayed using a Bradford dye-binding method (diluted samples permit handling of suitably sized aliquot volumes).

SDS-PAGE was carried out using 20 µg sample per lane (2 µg if sample is a denatured primary or secondary antibody) on 16% acrylamide gels, 1.5 mm thick, 10 wells (NOVEX) for 1 hr 80 V; 1% hrs 125 V. This was followed by Western Blotting onto nitrocellulose membrane at 50 V for 1½ hrs. The blots were probed with the following antibodies:

Anti-ApoA-IV (N-terminal specific), Santa Cruz Biotechnology, Inc.

Anti-ApoA-IV (C-terminal specific), Santa Cruz Biotechnology, Inc.

Both antibodies are affinity purified goat polyclonals raised against a peptide mapping near the amino (N-terminal) or carboxy (C-terminal) terminus of ApoA-IV of human origin. These antibodies were chosen since probing for the N- and C-terminals should increase the chance of detection of the ApoA-IV protein and/or fragments.

Several bands were found that appear to be ApoA-IV specific and also discriminatory for AD. These bands do not appear in the secondary antibody-only control blot for control or AD samples.

Bands 3-6 which are observed in the 10-16 kDa region are discriminatory for AD bands 3-6, but also appear to align with bands in the denatured ApoA-IV antibody lanes. It has also been observed that bands 3-6 are much stronger on blots where the N-terminal specific anti-ApoA-IV antibody has been used.

Two other key bands are observed. Band 1 is observed at approximately 45 kDa and appears to correspond to the full length mature APO-AIV protein. Band 2 is observed at approximately 28 kDa and appears to be an N-terminal fragment of APO-AIV.

Complement Factor H Validation.

Methods

Sample Dilution

Plasma samples were diluted to 1 in 8 in Phosphate buffered saline (PBS). An equal volume of Laemmli 2x sample buffer was added and then boiled for 10 min until use.

Western Blot

SDS gel electrophoresis was performed using the Fisher Scientific 36 well, 1.5 mm gels (all solutions were purchased from National Diagnostics). Samples were separated on a 10% resolving gel with a 4% stacking gel (all solutions were purchased from National Diagnostics). Samples (20 µl) were separated initially for 30 min at 110V and then for 60 min at 150V until the dye front just began to enter the running buffer.

The gel was transferred to PVDF (Amersham Biosciences) using a Semi-dry transblot (Bio-Rad) for 45 min at 15V. The membrane was then blocked in 5% milk made in PBS-Tween and probed with Complement factor H primary antibody (Abcam, UK) overnight at 4° C. The bands were detected with a chemiluminescence Western detection kit (ECL+, Amersham Biosciences) and the membranes were scanned using Storm fluorescence scanner (Amersham Biosciences).

An immunoreactive band was observed at 139 kDa (CfH) and the optical density was quantified using the Image Quant (Amersham Biosciences) software. Analysis was by non-parametric Mann-Whitney using the SPSS package.

Results

Western blot data was acquired from plasma from 128 people with NINCDS-ADRDA probable AD and 78 normal healthy elderly controls. Cases with AD had a 32% increase in CFH (Mann-Whitney; table 2)

TABLE 2

Diagnosis	Number	Mean CFH	SD	SEM
Controls	128	65.6	65.5	5.8
Probable AD	78	96.0	96.8	11.0

There was a gender difference with a relatively higher CFH value in females overall relative to males ($p=0.05$). However CFH was higher in cases with AD relative to controls even when considering genders separately ($p<0.01$; table 3)

TABLE 3

Females only	Number	Mean CFH	SEM
Controls	78	73.0	8.9
Probable AD	64	102.7	13.0
Total	142	86.4	7.7

A receiver operator curve (ROC) analysis showed that CFH performs better than chance as a diagnostic test.

The references mentioned herein are all expressly incorporated by reference.

REFERENCES

- Andreasen, N., C. Hesse, et al. (1999). "Cerebrospinal fluid beta-amyloid(1-42) in Alzheimer disease: differences between early- and late-onset Alzheimer disease and stability during the course of disease." *Arch Neurol* 56(6): 673-80.
- Bosman, G. J., I. G. Bartholomeus, et al. (1991). "Erythrocyte membrane characteristics indicate abnormal cellular aging in patients with Alzheimer's disease." *Neurobiol Aging* 12(1): 13-8.
- Friedland, R. P. (1993). "Epidemiology, education, and the ecology of Alzheimer's disease." *Neurology* 43(2): 246-9.

- Ida, N., T. Hartmann, et al. (1996). "Analysis of heterogeneous A4 peptides in human cerebrospinal fluid and blood by a newly developed sensitive Western blot assay." *J Biol Chem* 271(37): 22908-14.
- Kanai, M., E. Matsubara, et al. (1998). "Longitudinal study of cerebrospinal fluid levels of tau, A beta1-40, and A beta1-42(43) in Alzheimer's disease: a study in Japan." *Ann Neurol* 44(1): 17-26.
- Kawarabayashi, T., L. H. Younkin, et al. (2001). "Age-dependent changes in brain, CSF, and plasma amyloid (beta) protein in the Tg2576 transgenic mouse model of Alzheimer's disease." *J Neurosci* 21(2): 372-81.
- Kosaka, T., M. Imagawa, et al. (1997). "The beta APP717 Alzheimer mutation increases the percentage of plasma amyloid-beta protein ending at A beta42(43)." *Neurology* 48(3): 741-5.
- Kuo, Y. M., T. A. Kokjohn, et al. (2000). "Elevated abeta42 in skeletal muscle of Alzheimer disease patients suggests peripheral alterations of AbetaPP metabolism." *Am J Pathol* 156(3): 797-805.
- Lindner, M. D., D. D. Gordon, et al. (1993). "Increased levels of truncated nerve growth factor receptor in urine of mildly demented patients with Alzheimer's disease." *Arch Neurol* 50(10): 1054-60.
- Pirttila, T., S. Mattinen, et al. (1992). "The decrease of CD8-positive lymphocytes in Alzheimer's disease." *J Neurol Sci* 107(2): 160-5.
- Rocca, W. A., A. Hofman, et al. (1991). "Frequency and distribution of Alzheimer's disease in Europe: a collaborative study of 1980-1990 prevalence findings. The EURO-DEM-Prevalence Research Group." *Ann Neurol* 30(3): 381-90.
- Scheuner, D., C. Eckman, et al. (1996). "Secreted amyloid beta-protein similar to that in the senile plaques of Alzheimer's disease is increased in vivo by the presenilin 1 and 2 and APP mutations linked to familial Alzheimer's disease." *Nat Med* 2(8): 864-70.
- Ueno, I., T. Sakai, et al. (2000). "Analysis of blood plasma proteins in patients with Alzheimer's disease by two-dimensional electrophoresis, sequence homology and immunodetection." *Electrophoresis* 21(9): 1832-45.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 9

<210> SEQ ID NO 1

<211> LENGTH: 396

<212> TYPE: PRT

<213> ORGANISM: Homo Sapiens

<400> SEQUENCE: 1

Met Phe Leu Lys Ala Val Val Leu Thr Leu Ala Leu Val Ala Val Ala
1 5 10 15

Gly Ala Arg Ala Glu Val Ser Ala Asp Gln Val Ala Thr Val Met Trp
20 25 30

Asp Tyr Phe Ser Gln Leu Ser Asn Asn Ala Lys Glu Ala Val Glu His
35 40 45

Leu Gln Lys Ser Glu Leu Thr Gln Gln Leu Asn Ala Leu Phe Gln Asp
50 55 60

Lys Leu Gly Glu Val Asn Thr Tyr Ala Gly Asp Leu Gln Lys Lys Leu
65 70 75 80

Val Pro Phe Ala Thr Glu Leu His Glu Arg Leu Ala Lys Asp Ser Glu
85 90 95

-continued

```

Lys Leu Lys Glu Glu Ile Gly Lys Glu Leu Glu Glu Leu Arg Ala Arg
    100                      105                      110

Leu Leu Pro His Ala Asn Glu Val Ser Gln Lys Ile Gly Asp Asn Leu
    115                      120                      125

Arg Glu Leu Gln Gln Arg Leu Glu Pro Tyr Ala Asp Gln Leu Arg Thr
    130                      135                      140

Gln Val Asn Thr Gln Ala Glu Gln Leu Arg Arg Gln Leu Thr Pro Tyr
    145                      150                      155                      160

Ala Gln Arg Met Glu Arg Val Leu Arg Glu Asn Ala Asp Ser Leu Gln
    165                      170                      175

Ala Ser Leu Arg Pro His Ala Asp Glu Leu Lys Ala Lys Ile Asp Gln
    180                      185                      190

Asn Val Glu Glu Leu Lys Gly Arg Leu Thr Pro Tyr Ala Asp Glu Phe
    195                      200                      205

Lys Val Lys Ile Asp Gln Thr Val Glu Glu Leu Arg Arg Ser Leu Ala
    210                      215                      220

Pro Tyr Ala Gln Asp Thr Gln Glu Lys Leu Asn His Gln Leu Glu Gly
    225                      230                      235                      240

Leu Thr Phe Gln Met Lys Lys Asn Ala Glu Glu Leu Lys Ala Arg Ile
    245                      250                      255

Ser Ala Ser Ala Glu Glu Leu Arg Gln Arg Leu Ala Pro Leu Ala Glu
    260                      265                      270

Asp Val Arg Gly Asn Leu Lys Gly Asn Thr Glu Gly Leu Gln Lys Ser
    275                      280                      285

Leu Ala Glu Leu Gly Gly His Leu Asp Gln Gln Val Glu Glu Phe Arg
    290                      295                      300

Arg Arg Val Glu Pro Tyr Gly Glu Asn Phe Asn Lys Ala Leu Val Gln
    305                      310                      315                      320

Gln Met Glu Gln Leu Arg Gln Lys Leu Gly Pro His Ala Gly Asp Val
    325                      330                      335

Glu Gly His Leu Ser Phe Leu Glu Lys Asp Leu Arg Asp Lys Val Asn
    340                      345                      350

Ser Phe Phe Ser Thr Phe Lys Glu Lys Glu Ser Gln Asp Lys Thr Leu
    355                      360                      365

Ser Leu Pro Glu Leu Glu Gln Gln Gln Glu Gln Gln Gln Glu Gln Gln
    370                      375                      380

Gln Glu Gln Val Gln Met Leu Ala Pro Leu Glu Ser
    385                      390                      395

```

```

<210> SEQ ID NO 2
<211> LENGTH: 1744
<212> TYPE: PRT
<213> ORGANISM: Homo Sapiens

```

```

<400> SEQUENCE: 2

```

```

Met Arg Leu Leu Trp Gly Leu Ile Trp Ala Ser Ser Phe Phe Thr Leu
  1           5           10           15

Ser Leu Gln Lys Pro Arg Leu Leu Leu Phe Ser Pro Ser Val Val His
    20           25           30

Leu Gly Val Pro Leu Ser Val Gly Val Gln Leu Gln Asp Val Pro Arg
    35           40           45

Gly Gln Val Val Lys Gly Ser Val Phe Leu Arg Asn Pro Ser Arg Asn
    50           55           60

Asn Val Pro Cys Ser Pro Lys Val Asp Phe Thr Leu Ser Ser Glu Arg
    65           70           75           80

```

Asp	Phe	Ala	Leu	Leu	Ser	Leu	Gln	Val	Pro	Leu	Lys	Asp	Ala	Lys	Ser	
				85					90					95		
Cys	Gly	Leu	His	Gln	Leu	Leu	Arg	Gly	Pro	Glu	Val	Gln	Leu	Val	Ala	
				100					105					110		
His	Ser	Pro	Trp	Leu	Lys	Asp	Ser	Leu	Ser	Arg	Thr	Thr	Asn	Ile	Gln	
				115					120					125		
Gly	Ile	Asn	Leu	Leu	Phe	Ser	Ser	Arg	Arg	Gly	His	Leu	Phe	Leu	Gln	
				130					135					140		
Thr	Asp	Gln	Pro	Ile	Tyr	Asn	Pro	Gly	Gln	Arg	Val	Arg	Tyr	Arg	Val	
				145					150					155		
Phe	Ala	Leu	Asp	Gln	Lys	Met	Arg	Pro	Ser	Thr	Asp	Thr	Ile	Thr	Val	
				165					170					175		
Met	Val	Glu	Asn	Ser	His	Gly	Leu	Arg	Val	Arg	Lys	Lys	Glu	Val	Tyr	
				180					185					190		
Met	Pro	Ser	Ser	Ile	Phe	Gln	Asp	Asp	Phe	Val	Ile	Pro	Asp	Ile	Ser	
				195					200					205		
Glu	Pro	Gly	Thr	Trp	Lys	Ile	Ser	Ala	Arg	Phe	Ser	Asp	Gly	Leu	Glu	
				210					215					220		
Ser	Asn	Ser	Ser	Thr	Gln	Phe	Glu	Val	Lys	Lys	Tyr	Val	Leu	Pro	Asn	
				225					230					235		
Phe	Glu	Val	Lys	Ile	Thr	Pro	Gly	Lys	Pro	Tyr	Ile	Leu	Thr	Val	Pro	
				245					250					255		
Gly	His	Leu	Asp	Glu	Met	Gln	Leu	Asp	Ile	Gln	Ala	Arg	Tyr	Ile	Tyr	
				260					265					270		
Gly	Lys	Pro	Val	Gln	Gly	Val	Ala	Tyr	Val	Arg	Phe	Gly	Leu	Leu	Asp	
				275					280					285		
Glu	Asp	Gly	Lys	Lys	Thr	Phe	Phe	Arg	Gly	Leu	Glu	Ser	Gln	Thr	Lys	
				290					295					300		
Leu	Val	Asn	Gly	Gln	Ser	His	Ile	Ser	Leu	Ser	Lys	Ala	Glu	Phe	Gln	
				305					310					315		
Asp	Ala	Leu	Glu	Lys	Leu	Asn	Met	Gly	Ile	Thr	Asp	Leu	Gln	Gly	Leu	
				325					330					335		
Arg	Leu	Tyr	Val	Ala	Ala	Ala	Ile	Ile	Glu	Ser	Pro	Gly	Gly	Glu	Met	
				340					345					350		
Glu	Glu	Ala	Glu	Leu	Thr	Ser	Trp	Tyr	Phe	Val	Ser	Ser	Pro	Phe	Ser	
				355					360					365		
Leu	Asp	Leu	Ser	Lys	Thr	Lys	Arg	His	Leu	Val	Pro	Gly	Ala	Pro	Phe	
				370					375					380		
Leu	Leu	Gln	Ala	Leu	Val	Arg	Glu	Met	Ser	Gly	Ser	Pro	Ala	Ser	Gly	
				385					390					395		
Ile	Pro	Val	Lys	Val	Ser	Ala	Thr	Val	Ser	Ser	Pro	Gly	Ser	Val	Pro	
				405					410					415		
Glu	Val	Gln	Asp	Ile	Gln	Gln	Asn	Thr	Asp	Gly	Ser	Gly	Gln	Val	Ser	
				420					425					430		
Ile	Pro	Ile	Ile	Ile	Pro	Gln	Thr	Ile	Ser	Glu	Leu	Gln	Leu	Ser	Val	
				435					440					445		
Ser	Ala	Gly	Ser	Pro	His	Pro	Ala	Ile	Ala	Arg	Leu	Thr	Val	Ala	Ala	
				450					455					460		
Pro	Pro	Ser	Gly	Gly	Pro	Gly	Phe	Leu	Ser	Ile	Glu	Arg	Pro	Asp	Ser	
				465					470					475		
Arg	Pro	Pro	Arg	Val	Gly	Asp	Thr	Leu	Asn	Leu	Asn	Leu	Arg	Ala		

-continued

930	935	940
Val Tyr Glu Leu Asn Pro Leu Asp His Arg Gly Arg Thr Leu Glu Ile 945 950 955 960		
Pro Gly Asn Ser Asp Pro Asn Met Ile Pro Asp Gly Asp Phe Asn Ser 965 970 975		
Tyr Val Arg Val Thr Ala Ser Asp Pro Leu Asp Thr Leu Gly Ser Glu 980 985 990		
Gly Ala Leu Ser Pro Gly Gly Val Ala Ser Leu Leu Arg Leu Pro Arg 995 1000 1005		
Gly Cys Gly Glu Gln Thr Met Ile Tyr Leu Ala Pro Thr Leu Ala Ala 1010 1015 1020		
Ser Arg Tyr Leu Asp Lys Thr Glu Gln Trp Ser Thr Leu Pro Pro Glu 1025 1030 1035 1040		
Thr Lys Asp His Ala Val Asp Leu Ile Gln Lys Gly Tyr Met Arg Ile 1045 1050 1055		
Gln Gln Phe Arg Lys Ala Asp Gly Ser Tyr Ala Ala Trp Leu Ser Arg 1060 1065 1070		
Asp Ser Ser Thr Trp Leu Thr Ala Phe Val Leu Lys Val Leu Ser Leu 1075 1080 1085		
Ala Gln Glu Gln Val Gly Gly Ser Pro Glu Lys Leu Gln Glu Thr Ser 1090 1095 1100		
Asn Trp Leu Leu Ser Gln Gln Gln Ala Asp Gly Ser Phe Gln Asp Pro 1105 1110 1115 1120		
Cys Pro Val Leu Asp Arg Ser Met Gln Gly Gly Leu Val Gly Asn Asp 1125 1130 1135		
Glu Thr Val Ala Leu Thr Ala Phe Val Thr Ile Ala Leu His His Gly 1140 1145 1150		
Leu Ala Val Phe Gln Asp Glu Gly Ala Glu Pro Leu Lys Gln Arg Val 1155 1160 1165		
Glu Ala Ser Ile Ser Lys Ala Asn Ser Phe Leu Gly Glu Lys Ala Ser 1170 1175 1180		
Ala Gly Leu Leu Gly Ala His Ala Ala Ala Ile Thr Ala Tyr Ala Leu 1185 1190 1195 1200		
Ser Leu Thr Lys Ala Pro Val Asp Leu Leu Gly Val Ala His Asn Asn 1205 1210 1215		
Leu Met Ala Met Ala Gln Glu Thr Gly Asp Asn Leu Tyr Trp Gly Ser 1220 1225 1230		
Val Thr Gly Ser Gln Ser Asn Ala Val Ser Pro Thr Pro Ala Pro Arg 1235 1240 1245		
Asn Pro Ser Asp Pro Met Pro Gln Ala Pro Ala Leu Trp Ile Glu Thr 1250 1255 1260		
Thr Ala Tyr Ala Leu Leu His Leu Leu Leu His Glu Gly Lys Ala Glu 1265 1270 1275 1280		
Met Ala Asp Gln Ala Ser Ala Trp Leu Thr Arg Gln Gly Ser Phe Gln 1285 1290 1295		
Gly Gly Phe Arg Ser Thr Gln Asp Thr Val Ile Ala Leu Asp Ala Leu 1300 1305 1310		
Ser Ala Tyr Trp Ile Ala Ser His Thr Thr Glu Glu Arg Gly Leu Asn 1315 1320 1325		
Val Thr Leu Ser Ser Thr Gly Arg Asn Gly Phe Lys Ser His Ala Leu 1330 1335 1340		
Gln Leu Asn Asn Arg Gln Ile Arg Gly Leu Glu Glu Glu Leu Gln Phe 1345 1350 1355 1360		

-continued

```

Ser Leu Gly Ser Lys Ile Asn Val Lys Val Gly Gly Asn Ser Lys Gly
    1365                      1370                      1375

Thr Leu Lys Val Leu Arg Thr Tyr Asn Val Leu Asp Met Lys Asn Thr
    1380                      1385                      1390

Thr Cys Gln Asp Leu Gln Ile Glu Val Thr Val Lys Gly His Val Glu
    1395                      1400                      1405

Tyr Thr Met Glu Ala Asn Glu Asp Tyr Glu Asp Tyr Glu Tyr Asp Glu
    1410                      1415                      1420

Leu Pro Ala Lys Asp Asp Pro Asp Ala Pro Leu Gln Pro Val Thr Pro
    1425                      1430                      1435                      1440

Leu Gln Leu Phe Glu Gly Arg Arg Asn Arg Arg Arg Arg Glu Ala Pro
    1445                      1450                      1455

Lys Val Val Glu Glu Gln Glu Ser Arg Val His Tyr Thr Val Cys Ile
    1460                      1465                      1470

Trp Arg Asn Gly Lys Val Gly Leu Ser Gly Met Ala Ile Ala Asp Val
    1475                      1480                      1485

Thr Leu Leu Ser Gly Phe His Ala Leu Arg Ala Asp Leu Glu Lys Leu
    1490                      1495                      1500

Thr Ser Leu Ser Asp Arg Tyr Val Ser His Phe Glu Thr Glu Gly Pro
    1505                      1510                      1515                      1520

His Val Leu Leu Tyr Phe Asp Ser Val Pro Thr Ser Arg Glu Cys Val
    1525                      1530                      1535

Gly Phe Glu Ala Val Gln Glu Val Pro Val Gly Leu Val Gln Pro Ala
    1540                      1545                      1550

Ser Ala Thr Leu Tyr Asp Tyr Tyr Asn Pro Glu Arg Arg Cys Ser Val
    1555                      1560                      1565

Phe Tyr Gly Ala Pro Ser Lys Ser Arg Leu Leu Ala Thr Leu Cys Ser
    1570                      1575                      1580

Ala Glu Val Cys Gln Cys Ala Glu Gly Lys Cys Pro Arg Gln Arg Arg
    1585                      1590                      1595                      1600

Ala Leu Glu Arg Gly Leu Gln Asp Glu Asp Gly Tyr Arg Met Lys Phe
    1605                      1610                      1615

Ala Cys Tyr Tyr Pro Arg Val Glu Tyr Gly Phe Gln Val Lys Val Leu
    1620                      1625                      1630

Arg Glu Asp Ser Arg Ala Ala Phe Arg Leu Phe Glu Thr Lys Ile Thr
    1635                      1640                      1645

Gln Val Leu His Phe Thr Lys Asp Val Lys Ala Ala Ala Asn Gln Met
    1650                      1655                      1660

Arg Asn Phe Leu Val Arg Ala Ser Cys Arg Leu Arg Leu Glu Pro Gly
    1665                      1670                      1675                      1680

Lys Glu Tyr Leu Ile Met Gly Leu Asp Gly Ala Thr Tyr Asp Leu Glu
    1685                      1690                      1695

Gly His Pro Gln Tyr Leu Leu Asp Ser Asn Ser Trp Ile Glu Glu Met
    1700                      1705                      1710

Pro Ser Glu Arg Leu Cys Arg Ser Thr Arg Gln Arg Ala Ala Cys Ala
    1715                      1720                      1725

Gln Leu Asn Asp Phe Leu Gln Glu Tyr Gly Thr Gln Gly Cys Gln Val
    1730                      1735                      1740

```

```

<210> SEQ ID NO 3
<211> LENGTH: 1474
<212> TYPE: PRT
<213> ORGANISM: Homo Sapiens

<400> SEQUENCE: 3

```

-continued

Met	Gly	Lys	Asn	Lys	Leu	Leu	His	Pro	Ser	Leu	Val	Leu	Leu	Leu	Leu	1	5	10	15
Val	Leu	Leu	Pro	Thr	Asp	Ala	Ser	Val	Ser	Gly	Lys	Pro	Gln	Tyr	Met	20	25	30	
Val	Leu	Val	Pro	Ser	Leu	Leu	His	Thr	Glu	Thr	Thr	Glu	Lys	Gly	Cys	35	40	45	
Val	Leu	Leu	Ser	Tyr	Leu	Asn	Glu	Thr	Val	Thr	Val	Ser	Ala	Ser	Leu	50	55	60	
Glu	Ser	Val	Arg	Gly	Asn	Arg	Ser	Leu	Phe	Thr	Asp	Leu	Glu	Ala	Glu	65	70	75	80
Asn	Asp	Val	Leu	His	Cys	Val	Ala	Phe	Ala	Val	Pro	Lys	Ser	Ser	Ser	85	90	95	
Asn	Glu	Glu	Val	Met	Phe	Leu	Thr	Val	Gln	Val	Lys	Gly	Pro	Thr	Gln	100	105	110	
Glu	Phe	Lys	Lys	Arg	Thr	Thr	Val	Met	Val	Lys	Asn	Glu	Asp	Ser	Leu	115	120	125	
Val	Phe	Val	Gln	Thr	Asp	Lys	Ser	Ile	Tyr	Lys	Pro	Gly	Gln	Thr	Val	130	135	140	
Lys	Phe	Arg	Val	Val	Ser	Met	Asp	Glu	Asn	Phe	His	Pro	Leu	Asn	Glu	145	150	155	160
Leu	Ile	Pro	Leu	Val	Tyr	Ile	Gln	Asp	Pro	Lys	Gly	Asn	Arg	Ile	Ala	165	170	175	
Gln	Trp	Gln	Ser	Phe	Gln	Leu	Glu	Gly	Gly	Leu	Lys	Gln	Phe	Ser	Phe	180	185	190	
Pro	Leu	Ser	Ser	Glu	Pro	Phe	Gln	Gly	Ser	Tyr	Lys	Val	Val	Val	Gln	195	200	205	
Lys	Lys	Ser	Gly	Gly	Arg	Thr	Glu	His	Pro	Phe	Thr	Val	Glu	Glu	Phe	210	215	220	
Val	Leu	Pro	Lys	Phe	Glu	Val	Gln	Val	Thr	Val	Pro	Lys	Ile	Ile	Thr	225	230	235	240
Ile	Leu	Glu	Glu	Glu	Met	Asn	Val	Ser	Val	Cys	Gly	Leu	Tyr	Thr	Tyr	245	250	255	
Gly	Lys	Pro	Val	Pro	Gly	His	Val	Thr	Val	Ser	Ile	Cys	Arg	Lys	Tyr	260	265	270	
Ser	Asp	Ala	Ser	Asp	Cys	His	Gly	Glu	Asp	Ser	Gln	Ala	Phe	Cys	Glu	275	280	285	
Lys	Phe	Ser	Gly	Gln	Leu	Asn	Ser	His	Gly	Cys	Phe	Tyr	Gln	Gln	Val	290	295	300	
Lys	Thr	Lys	Val	Phe	Gln	Leu	Lys	Arg	Lys	Glu	Tyr	Glu	Met	Lys	Leu	305	310	315	320
His	Thr	Glu	Ala	Gln	Ile	Gln	Glu	Glu	Gly	Thr	Val	Val	Glu	Leu	Thr	325	330	335	
Gly	Arg	Gln	Ser	Ser	Glu	Ile	Thr	Arg	Thr	Ile	Thr	Lys	Leu	Ser	Phe	340	345	350	
Val	Lys	Val	Asp	Ser	His	Phe	Arg	Gln	Gly	Ile	Pro	Phe	Phe	Gly	Gln	355	360	365	
Val	Arg	Leu	Val	Asp	Gly	Lys	Gly	Val	Pro	Ile	Pro	Asn	Lys	Val	Ile	370	375	380	
Phe	Ile	Arg	Gly	Asn	Glu	Ala	Asn	Tyr	Tyr	Ser	Asn	Ala	Thr	Thr	Asp	385	390	395	400
Glu	His	Gly	Leu	Val	Gln	Phe	Ser	Ile	Asn	Thr	Thr	Asn	Val	Met	Gly	405	410	415	
Thr	Ser	Leu	Thr	Val	Arg	Val	Asn	Tyr	Lys	Asp	Arg	Ser	Pro	Cys	Tyr	420	425	430	

-continued

Gly Tyr Gln Trp Val Ser Glu Glu His Glu Glu Ala His His Thr Ala
 435 440 445
 Tyr Leu Val Phe Ser Pro Ser Lys Ser Phe Val His Leu Glu Pro Met
 450 455 460
 Ser His Glu Leu Pro Cys Gly His Thr Gln Thr Val Gln Ala His Tyr
 465 470 475 480
 Ile Leu Asn Gly Gly Thr Leu Leu Gly Leu Lys Lys Leu Ser Phe Tyr
 485 490 495
 Tyr Leu Ile Met Ala Lys Gly Gly Ile Val Arg Thr Gly Thr His Gly
 500 505 510
 Leu Leu Val Lys Gln Glu Asp Met Lys Gly His Phe Ser Ile Ser Ile
 515 520 525
 Pro Val Lys Ser Asp Ile Ala Pro Val Ala Arg Leu Leu Ile Tyr Ala
 530 535 540
 Val Leu Pro Thr Gly Asp Val Ile Gly Asp Ser Ala Lys Tyr Asp Val
 545 550 555 560
 Glu Asn Cys Ile Ala Asn Lys Val Asp Leu Ser Phe Ser Pro Ser Gln
 565 570 575
 Ser Leu Pro Ala Ser His Ala His Leu Arg Val Thr Ala Ala Pro Gln
 580 585 590
 Ser Val Cys Ala Leu Arg Ala Val Asp Gln Ser Val Leu Leu Met Lys
 595 600 605
 Pro Asp Ala Glu Leu Ser Ala Ser Ser Val Tyr Asn Leu Leu Pro Glu
 610 615 620
 Lys Asp Leu Thr Gly Phe Pro Gly Pro Leu Asn Asp Gln Asp Asp Glu
 625 630 635 640
 Asp Cys Ile Asn Arg His Asn Val Tyr Ile Asn Gly Ile Thr Tyr Thr
 645 650 655
 Pro Val Ser Ser Thr Asn Glu Lys Asp Met Tyr Ser Phe Leu Glu Asp
 660 665 670
 Met Gly Leu Lys Ala Phe Thr Asn Ser Lys Ile Arg Lys Pro Lys Met
 675 680 685
 Cys Pro Gln Leu Gln Gln Tyr Glu Met His Gly Pro Glu Gly Leu Arg
 690 695 700
 Val Gly Phe Tyr Glu Ser Asp Val Met Gly Arg Gly His Ala Arg Leu
 705 710 715 720
 Val His Val Glu Glu Pro His Thr Glu Thr Val Arg Lys Tyr Phe Pro
 725 730 735
 Glu Thr Trp Ile Trp Asp Leu Val Val Val Asn Ser Ala Gly Val Ala
 740 745 750
 Glu Val Gly Val Thr Val Pro Asp Thr Ile Thr Glu Trp Lys Ala Gly
 755 760 765
 Ala Phe Cys Leu Ser Glu Asp Ala Gly Leu Gly Ile Ser Ser Thr Ala
 770 775 780
 Ser Leu Arg Ala Phe Gln Pro Phe Phe Val Glu Leu Thr Met Pro Tyr
 785 790 795 800
 Ser Val Ile Arg Gly Glu Ala Phe Thr Leu Lys Ala Thr Val Leu Asn
 805 810 815
 Tyr Leu Pro Lys Cys Ile Arg Val Ser Val Gln Leu Glu Ala Ser Pro
 820 825 830
 Ala Phe Leu Ala Val Pro Val Glu Lys Glu Gln Ala Pro His Cys Ile
 835 840 845
 Cys Ala Asn Gly Arg Gln Thr Val Ser Trp Ala Val Thr Pro Lys Ser

-continued

850	855	860
Leu Gly Asn Val Asn Phe Thr Val Ser Ala Glu Ala Leu Glu Ser Gln 865 870 875 880		
Glu Leu Cys Gly Thr Glu Val Pro Ser Val Pro Glu His Gly Arg Lys 885 890 895		
Asp Thr Val Ile Lys Pro Leu Leu Val Glu Pro Glu Gly Leu Glu Lys 900 905 910		
Glu Thr Thr Phe Asn Ser Leu Leu Cys Pro Ser Gly Gly Glu Val Ser 915 920 925		
Glu Glu Leu Ser Leu Lys Leu Pro Pro Asn Val Val Glu Glu Ser Ala 930 935 940		
Arg Ala Ser Val Ser Val Leu Gly Asp Ile Leu Gly Ser Ala Met Gln 945 950 955 960		
Asn Thr Gln Asn Leu Leu Gln Met Pro Tyr Gly Cys Gly Glu Gln Asn 965 970 975		
Met Val Leu Phe Ala Pro Asn Ile Tyr Val Leu Asp Tyr Leu Asn Glu 980 985 990		
Thr Gln Gln Leu Thr Pro Glu Val Lys Ser Lys Ala Ile Gly Tyr Leu 995 1000 1005		
Asn Thr Gly Tyr Gln Arg Gln Leu Asn Tyr Lys His Tyr Asp Gly Ser 1010 1015 1020		
Tyr Ser Thr Phe Gly Glu Arg Tyr Gly Arg Asn Gln Gly Asn Thr Trp 1025 1030 1035 1040		
Leu Thr Ala Phe Val Leu Lys Thr Phe Ala Gln Ala Arg Ala Tyr Ile 1045 1050 1055		
Phe Ile Asp Glu Ala His Ile Thr Gln Ala Leu Ile Trp Leu Ser Gln 1060 1065 1070		
Arg Gln Lys Asp Asn Gly Cys Phe Arg Ser Ser Gly Ser Leu Leu Asn 1075 1080 1085		
Asn Ala Ile Lys Gly Gly Val Glu Asp Glu Val Thr Leu Ser Ala Tyr 1090 1095 1100		
Ile Thr Ile Ala Leu Leu Glu Ile Pro Leu Thr Val Thr His Pro Val 1105 1110 1115 1120		
Val Arg Asn Ala Leu Phe Cys Leu Glu Ser Ala Trp Lys Thr Ala Gln 1125 1130 1135		
Glu Gly Asp His Gly Ser His Val Tyr Thr Lys Ala Leu Leu Ala Tyr 1140 1145 1150		
Ala Phe Ala Leu Ala Gly Asn Gln Asp Lys Arg Lys Glu Val Leu Lys 1155 1160 1165		
Ser Leu Asn Glu Glu Ala Val Lys Lys Asp Asn Ser Val His Trp Glu 1170 1175 1180		
Arg Pro Gln Lys Pro Lys Ala Pro Val Gly His Phe Tyr Glu Pro Gln 1185 1190 1195 1200		
Ala Pro Ser Ala Glu Val Glu Met Thr Ser Tyr Val Leu Leu Ala Tyr 1205 1210 1215		
Leu Thr Ala Gln Pro Ala Pro Thr Ser Glu Asp Leu Thr Ser Ala Thr 1220 1225 1230		
Asn Ile Val Lys Trp Ile Thr Lys Gln Gln Asn Ala Gln Gly Gly Phe 1235 1240 1245		
Ser Ser Thr Gln Asp Thr Val Val Ala Leu His Ala Leu Ser Lys Tyr 1250 1255 1260		
Gly Ala Ala Thr Phe Thr Arg Thr Gly Lys Ala Ala Gln Val Thr Ile 1265 1270 1275 1280		

-continued

Gln Ser Ser Gly Thr Phe Ser Ser Lys Phe Gln Val Asp Asn Asn Asn
1285 1290 1295

Arg Leu Leu Leu Gln Gln Val Ser Leu Pro Glu Leu Pro Gly Glu Tyr
1300 1305 1310

Ser Met Lys Val Thr Gly Glu Gly Cys Val Tyr Leu Gln Thr Ser Leu
1315 1320 1325

Lys Tyr Asn Ile Leu Pro Glu Lys Glu Glu Phe Pro Phe Ala Leu Gly
1330 1335 1340

Val Gln Thr Leu Pro Gln Thr Cys Asp Glu Pro Lys Ala His Thr Ser
1345 1350 1355 1360

Phe Gln Ile Ser Leu Ser Val Ser Tyr Thr Gly Ser Arg Ser Ala Ser
1365 1370 1375

Asn Met Ala Ile Val Asp Val Lys Met Val Ser Gly Phe Ile Pro Leu
1380 1385 1390

Lys Pro Thr Val Lys Met Leu Glu Arg Ser Asn His Val Ser Arg Thr
1395 1400 1405

Glu Val Ser Ser Asn His Val Leu Ile Tyr Leu Asp Lys Val Ser Asn
1410 1415 1420

Gln Thr Leu Ser Leu Phe Phe Thr Val Leu Gln Asp Val Pro Val Arg
1425 1430 1435 1440

Asp Leu Lys Pro Ala Ile Val Lys Val Tyr Asp Tyr Tyr Glu Thr Asp
1445 1450 1455

Glu Phe Ala Ile Ala Glu Tyr Asn Ala Pro Cys Ser Lys Asp Leu Gly
1460 1465 1470

Asn Ala

<210> SEQ ID NO 4
 <211> LENGTH: 930
 <212> TYPE: PRT
 <213> ORGANISM: Homo Sapiens

<400> SEQUENCE: 4

Met Lys Pro Pro Arg Pro Val Arg Thr Cys Ser Lys Val Leu Val Leu
1 5 10 15

Leu Ser Leu Leu Ala Ile His Gln Thr Thr Thr Ala Glu Lys Asn Gly
20 25 30

Ile Asp Ile Tyr Ser Leu Thr Val Asp Ser Arg Val Ser Ser Arg Phe
35 40 45

Ala His Thr Val Val Thr Ser Arg Val Val Asn Arg Ala Asn Thr Val
50 55 60

Gln Glu Ala Thr Phe Gln Met Glu Leu Pro Lys Lys Ala Phe Ile Thr
65 70 75 80

Asn Phe Ser Met Asn Ile Asp Gly Met Thr Tyr Pro Gly Ile Ile Lys
85 90 95

Glu Lys Ala Glu Ala Gln Ala Gln Tyr Ser Ala Ala Val Ala Lys Gly
100 105 110

Lys Ser Ala Gly Leu Val Lys Ala Thr Gly Arg Asn Met Glu Gln Phe
115 120 125

Gln Val Ser Val Ser Val Ala Pro Asn Ala Lys Ile Thr Phe Glu Leu
130 135 140

Val Tyr Glu Glu Leu Leu Lys Arg Arg Leu Gly Val Tyr Glu Leu Leu
145 150 155 160

Leu Lys Val Arg Pro Gln Gln Leu Val Lys His Leu Gln Met Asp Ile
165 170 175

His Ile Phe Glu Pro Gln Gly Ile Ser Phe Leu Glu Thr Glu Ser Thr

-continued

180							185					190				
Phe	Met	Thr	Asn	Gln	Leu	Val	Asp	Ala	Leu	Thr	Thr	Trp	Gln	Asn	Lys	
195							200					205				
Thr	Lys	Ala	His	Ile	Arg	Phe	Lys	Pro	Thr	Leu	Ser	Gln	Gln	Gln	Lys	
210							215					220				
Ser	Pro	Glu	Gln	Gln	Glu	Thr	Val	Leu	Asp	Gly	Asn	Leu	Ile	Ile	Arg	
225							230					235				
Tyr	Asp	Val	Asp	Arg	Ala	Ile	Ser	Gly	Gly	Ser	Ile	Gln	Ile	Glu	Asn	
245							250					255				
Gly	Tyr	Phe	Val	His	Tyr	Phe	Ala	Pro	Glu	Gly	Leu	Thr	Thr	Met	Pro	
260							265					270				
Lys	Asn	Val	Val	Phe	Val	Ile	Asp	Lys	Ser	Gly	Ser	Met	Ser	Gly	Arg	
275							280					285				
Lys	Ile	Gln	Gln	Thr	Arg	Glu	Ala	Leu	Ile	Lys	Ile	Leu	Asp	Asp	Leu	
290							295					300				
Ser	Pro	Arg	Asp	Gln	Phe	Asn	Leu	Ile	Val	Phe	Ser	Thr	Glu	Ala	Thr	
305							310					315				
Gln	Trp	Arg	Pro	Ser	Leu	Val	Pro	Ala	Ser	Ala	Glu	Asn	Val	Asn	Lys	
325							330					335				
Ala	Arg	Ser	Phe	Ala	Ala	Gly	Ile	Gln	Ala	Leu	Gly	Gly	Thr	Asn	Ile	
340							345					350				
Asn	Asp	Ala	Met	Leu	Met	Ala	Val	Gln	Leu	Leu	Asp	Ser	Ser	Asn	Gln	
355							360					365				
Glu	Glu	Arg	Leu	Pro	Glu	Gly	Ser	Val	Ser	Leu	Ile	Ile	Leu	Leu	Thr	
370							375					380				
Asp	Gly	Asp	Pro	Thr	Val	Gly	Glu	Thr	Asn	Pro	Arg	Ser	Ile	Gln	Asn	
385							390					395				
Asn	Val	Arg	Glu	Ala	Val	Ser	Gly	Arg	Tyr	Ser	Leu	Phe	Cys	Leu	Gly	
405							410					415				
Phe	Gly	Phe	Asp	Val	Ser	Tyr	Ala	Phe	Leu	Glu	Lys	Leu	Ala	Leu	Asp	
420							425					430				
Asn	Gly	Gly	Leu	Ala	Arg	Arg	Ile	His	Glu	Asp	Ser	Asp	Ser	Ala	Leu	
435							440					445				
Gln	Leu	Gln	Asp	Phe	Tyr	Gln	Glu	Val	Ala	Asn	Pro	Leu	Leu	Thr	Ala	
450							455					460				
Val	Thr	Phe	Glu	Tyr	Pro	Ser	Asn	Ala	Val	Glu	Glu	Val	Thr	Gln	Asn	
465							470					475				
Asn	Phe	Arg	Leu	Leu	Phe	Lys	Gly	Ser	Glu	Met	Val	Val	Ala	Gly	Lys	
485							490					495				
Leu	Gln	Asp	Arg	Gly	Pro	Asp	Val	Leu	Thr	Ala	Thr	Val	Ser	Gly	Lys	
500							505					510				
Leu	Pro	Thr	Gln	Asn	Ile	Thr	Phe	Gln	Thr	Glu	Ser	Ser	Val	Ala	Glu	
515							520					525				
Gln	Glu	Ala	Glu	Phe	Gln	Ser	Pro	Lys	Tyr	Ile	Phe	His	Asn	Phe	Met	
530							535					540				
Glu	Arg	Leu	Trp	Ala	Tyr	Leu	Thr	Ile	Gln	Gln	Leu	Leu	Glu	Gln	Thr	
545							550					555				
Val	Ser	Ala	Ser	Asp	Ala	Asp	Gln	Gln	Ala	Leu	Arg	Asn	Gln	Ala	Leu	
565							570					575				
Asn	Leu	Ser	Leu	Ala	Tyr	Ser	Phe	Val	Thr	Pro	Leu	Thr	Ser	Met	Val	
580							585					590				
Val	Thr	Lys	Pro	Asp	Asp	Gln	Glu	Gln	Ser	Gln	Val	Ala	Glu	Lys	Pro	
595							600					605				

-continued

Met	Glu	Gly	Glu	Ser	Arg	Asn	Arg	Asn	Val	His	Ser	Gly	Ser	Thr	Phe
610						615					620				
Phe	Lys	Tyr	Tyr	Leu	Gln	Gly	Ala	Lys	Ile	Pro	Lys	Pro	Glu	Ala	Ser
625					630					635					640
Phe	Ser	Pro	Arg	Arg	Gly	Trp	Asn	Arg	Gln	Ala	Gly	Ala	Ala	Gly	Ser
				645					650						655
Arg	Met	Asn	Phe	Arg	Pro	Gly	Val	Leu	Ser	Ser	Arg	Gln	Leu	Gly	Leu
			660					665					670		
Pro	Gly	Pro	Pro	Asp	Val	Pro	Asp	His	Ala	Ala	Tyr	His	Pro	Phe	Arg
			675				680					685			
Arg	Leu	Ala	Ile	Leu	Pro	Ala	Ser	Ala	Pro	Pro	Ala	Thr	Ser	Asn	Pro
	690					695					700				
Asp	Pro	Ala	Val	Ser	Arg	Val	Met	Asn	Met	Lys	Ile	Glu	Glu	Thr	Thr
705					710					715					720
Met	Thr	Thr	Gln	Thr	Pro	Ala	Pro	Ile	Gln	Ala	Pro	Ser	Ala	Ile	Leu
			725						730						735
Pro	Leu	Pro	Gly	Gln	Ser	Val	Glu	Arg	Leu	Cys	Val	Asp	Pro	Arg	His
			740					745					750		
Arg	Gln	Gly	Pro	Val	Asn	Leu	Leu	Ser	Asp	Pro	Glu	Gln	Gly	Val	Glu
		755					760					765			
Val	Thr	Gly	Gln	Tyr	Glu	Arg	Glu	Lys	Ala	Gly	Phe	Ser	Trp	Ile	Glu
	770					775					780				
Val	Thr	Phe	Lys	Asn	Pro	Leu	Val	Trp	Val	His	Ala	Ser	Pro	Glu	His
785					790					795					800
Val	Val	Val	Thr	Arg	Asn	Arg	Arg	Ser	Ser	Ala	Tyr	Lys	Trp	Lys	Glu
			805						810						815
Thr	Leu	Phe	Ser	Val	Met	Pro	Gly	Leu	Lys	Met	Thr	Met	Asp	Lys	Thr
			820					825					830		
Gly	Leu	Leu	Leu	Leu	Ser	Asp	Pro	Asp	Lys	Val	Thr	Ile	Gly	Leu	Leu
		835					840					845			
Phe	Trp	Asp	Gly	Arg	Gly	Glu	Gly	Leu	Arg	Leu	Leu	Leu	Arg	Asp	Thr
	850					855					860				
Asp	Arg	Phe	Ser	Ser	His	Val	Gly	Gly	Thr	Leu	Gly	Gln	Phe	Tyr	Gln
865					870				875						880
Glu	Val	Leu	Trp	Gly	Ser	Pro	Ala	Ala	Ser	Asp	Asp	Gly	Arg	Arg	Thr
			885						890						895
Leu	Arg	Val	Gln	Gly	Asn	Asp	His	Ser	Ala	Thr	Arg	Glu	Arg	Arg	Leu
		900						905					910		
Asp	Tyr	Gln	Glu	Gly	Pro	Pro	Gly	Val	Glu	Ile	Ser	Cys	Trp	Ser	Val
		915					920					925			
Glu	Leu														
	930														

<210> SEQ ID NO 5

<211> LENGTH: 1663

<212> TYPE: PRT

<213> ORGANISM: Homo Sapiens

<400> SEQUENCE: 5

Met	Gly	Pro	Thr	Ser	Gly	Pro	Ser	Leu	Leu	Leu	Leu	Leu	Leu	Thr	His
1				5				10						15	
Leu	Pro	Leu	Ala	Leu	Gly	Ser	Pro	Met	Tyr	Ser	Ile	Ile	Thr	Pro	Asn
		20						25					30		
Ile	Leu	Arg	Leu	Glu	Ser	Glu	Glu	Thr	Met	Val	Leu	Glu	Ala	His	Asp
		35						40					45		

-continued

Ala	Gln	Gly	Asp	Val	Pro	Val	Thr	Val	Thr	Val	His	Asp	Phe	Pro	Gly
50						55					60				
Lys	Lys	Leu	Val	Leu	Ser	Ser	Glu	Lys	Thr	Val	Leu	Thr	Pro	Ala	Thr
65					70					75					80
Asn	His	Met	Gly	Asn	Val	Thr	Phe	Thr	Ile	Pro	Ala	Asn	Arg	Glu	Phe
			85						90					95	
Lys	Ser	Glu	Lys	Gly	Arg	Asn	Lys	Phe	Val	Thr	Val	Gln	Ala	Thr	Phe
			100					105					110		
Gly	Thr	Gln	Val	Val	Glu	Lys	Val	Val	Leu	Val	Ser	Leu	Gln	Ser	Gly
		115					120					125			
Tyr	Leu	Phe	Ile	Gln	Thr	Asp	Lys	Thr	Ile	Tyr	Thr	Pro	Gly	Ser	Thr
	130					135					140				
Val	Leu	Tyr	Arg	Ile	Phe	Thr	Val	Asn	His	Lys	Leu	Leu	Pro	Val	Gly
145					150					155					160
Arg	Thr	Val	Met	Val	Asn	Ile	Glu	Asn	Pro	Glu	Gly	Ile	Pro	Val	Lys
			165					170					175		
Gln	Asp	Ser	Leu	Ser	Ser	Gln	Asn	Gln	Leu	Gly	Val	Leu	Pro	Leu	Ser
		180						185				190			
Trp	Asp	Ile	Pro	Glu	Leu	Val	Asn	Met	Gly	Gln	Trp	Lys	Ile	Arg	Ala
	195						200					205			
Tyr	Tyr	Glu	Asn	Ser	Pro	Gln	Gln	Val	Phe	Ser	Thr	Glu	Phe	Glu	Val
	210					215					220				
Lys	Glu	Tyr	Val	Leu	Pro	Ser	Phe	Glu	Val	Ile	Val	Glu	Pro	Thr	Glu
225					230					235					240
Lys	Phe	Tyr	Tyr	Ile	Tyr	Asn	Glu	Lys	Gly	Leu	Glu	Val	Thr	Ile	Thr
			245						250					255	
Ala	Arg	Phe	Leu	Tyr	Gly	Lys	Lys	Val	Glu	Gly	Thr	Ala	Phe	Val	Ile
		260						265					270		
Phe	Gly	Ile	Gln	Asp	Gly	Glu	Gln	Arg	Ile	Ser	Leu	Pro	Glu	Ser	Leu
	275						280					285			
Lys	Arg	Ile	Pro	Ile	Glu	Asp	Gly	Ser	Gly	Glu	Val	Val	Leu	Ser	Arg
	290					295					300				
Lys	Val	Leu	Leu	Asp	Gly	Val	Gln	Asn	Leu	Arg	Ala	Glu	Asp	Leu	Val
305				310						315					320
Gly	Lys	Ser	Leu	Tyr	Val	Ser	Ala	Thr	Val	Ile	Leu	His	Ser	Gly	Ser
			325						330					335	
Asp	Met	Val	Gln	Ala	Glu	Arg	Ser	Gly	Ile	Pro	Ile	Val	Thr	Ser	Pro
		340						345					350		
Tyr	Gln	Ile	His	Phe	Thr	Lys	Thr	Pro	Lys	Tyr	Phe	Lys	Pro	Gly	Met
	355						360					365			
Pro	Phe	Asp	Leu	Met	Val	Phe	Val	Thr	Asn	Pro	Asp	Gly	Ser	Pro	Ala
	370					375						380			
Tyr	Arg	Val	Pro	Val	Ala	Val	Gln	Gly	Glu	Asp	Thr	Val	Gln	Ser	Leu
385					390					395					400
Thr	Gln	Gly	Asp	Gly	Val	Ala	Lys	Leu	Ser	Ile	Asn	Thr	His	Pro	Ser
			405						410					415	
Gln	Lys	Pro	Leu	Ser	Ile	Thr	Val	Arg	Thr	Lys	Lys	Gln	Glu	Leu	Ser
			420					425					430		
Glu	Ala	Glu	Gln	Ala	Thr	Arg	Thr	Met	Gln	Ala	Leu	Pro	Tyr	Ser	Thr
	435						440					445			
Val	Gly	Asn	Ser	Asn	Asn	Tyr	Leu	His	Leu	Ser	Val	Leu	Arg	Thr	Glu
	450					455					460				
Leu	Arg	Pro	Gly	Glu	Thr	Leu	Asn	Val	Asn	Phe	Leu	Leu	Arg	Met	Asp
465					470					475					480

-continued

Arg Ala His Glu Ala Lys Ile Arg Tyr Tyr Thr Tyr Leu Ile Met Asn
 485 490 495
 Lys Gly Arg Leu Leu Lys Ala Gly Arg Gln Val Arg Glu Pro Gly Gln
 500 505 510
 Asp Leu Val Val Leu Pro Leu Ser Ile Thr Thr Asp Phe Ile Pro Ser
 515 520 525
 Phe Arg Leu Val Ala Tyr Tyr Thr Leu Ile Gly Ala Ser Gly Gln Arg
 530 535 540
 Glu Val Val Ala Asp Ser Val Trp Val Asp Val Lys Asp Ser Cys Val
 545 550 555 560
 Gly Ser Leu Val Val Lys Ser Gly Gln Ser Glu Asp Arg Gln Pro Val
 565 570 575
 Pro Gly Gln Gln Met Thr Leu Lys Ile Glu Gly Asp His Gly Ala Arg
 580 585 590
 Val Val Leu Val Ala Val Asp Lys Gly Val Phe Val Leu Asn Lys Lys
 595 600 605
 Asn Lys Leu Thr Gln Ser Lys Ile Trp Asp Val Val Glu Lys Ala Asp
 610 615 620
 Ile Gly Cys Thr Pro Gly Ser Gly Lys Asp Tyr Ala Gly Val Phe Ser
 625 630 635 640
 Asp Ala Gly Leu Thr Phe Thr Ser Ser Ser Gly Gln Gln Thr Ala Gln
 645 650 655
 Arg Ala Glu Leu Gln Cys Pro Gln Pro Ala Ala Arg Arg Arg Arg Ser
 660 665 670
 Val Gln Leu Thr Glu Lys Arg Met Asp Lys Val Gly Lys Tyr Pro Lys
 675 680 685
 Glu Leu Arg Lys Cys Cys Glu Asp Gly Met Arg Glu Asn Pro Met Arg
 690 695 700
 Phe Ser Cys Gln Arg Arg Thr Arg Phe Ile Ser Leu Gly Glu Ala Cys
 705 710 715 720
 Lys Lys Val Phe Leu Asp Cys Cys Asn Tyr Ile Thr Glu Leu Arg Arg
 725 730 735
 Gln His Ala Arg Ala Ser His Leu Gly Leu Ala Arg Ser Asn Leu Asp
 740 745 750
 Glu Asp Ile Ile Ala Glu Glu Asn Ile Val Ser Arg Ser Glu Phe Pro
 755 760 765
 Glu Ser Trp Leu Trp Asn Val Glu Asp Leu Lys Glu Pro Pro Lys Asn
 770 775 780
 Gly Ile Ser Thr Lys Leu Met Asn Ile Phe Leu Lys Asp Ser Ile Thr
 785 790 795 800
 Thr Trp Glu Ile Leu Ala Val Ser Met Ser Asp Lys Lys Gly Ile Cys
 805 810 815
 Val Ala Asp Pro Phe Glu Val Thr Val Met Gln Asp Phe Phe Ile Asp
 820 825 830
 Leu Arg Leu Pro Tyr Ser Val Val Arg Asn Glu Gln Val Glu Ile Arg
 835 840 845
 Ala Val Leu Tyr Asn Tyr Arg Gln Asn Gln Glu Leu Lys Val Arg Val
 850 855 860
 Glu Leu Leu His Asn Pro Ala Phe Cys Ser Leu Ala Thr Thr Lys Arg
 865 870 875 880
 Arg His Gln Gln Thr Val Thr Ile Pro Pro Lys Ser Ser Leu Ser Val
 885 890 895
 Pro Tyr Val Ile Val Pro Leu Lys Thr Gly Leu Gln Glu Val Glu Val

-continued

900							905					910				
Lys	Ala	Ala	Val	Tyr	His	His	Phe	Ile	Ser	Asp	Gly	Val	Arg	Lys	Ser	
	915						920					925				
Leu	Lys	Val	Val	Pro	Glu	Gly	Ile	Arg	Met	Asn	Lys	Thr	Val	Ala	Val	
	930					935					940					
Arg	Thr	Leu	Asp	Pro	Glu	Arg	Leu	Gly	Arg	Glu	Gly	Val	Gln	Lys	Glu	
945					950					955					960	
Asp	Ile	Pro	Pro	Ala	Asp	Leu	Ser	Asp	Gln	Val	Pro	Asp	Thr	Glu	Ser	
				965					970					975		
Glu	Thr	Arg	Ile	Leu	Leu	Gln	Gly	Thr	Pro	Val	Ala	Gln	Met	Thr	Glu	
			980					985					990			
Asp	Ala	Val	Asp	Ala	Glu	Arg	Leu	Lys	His	Leu	Ile	Val	Thr	Pro	Ser	
	995						1000					1005				
Gly	Cys	Gly	Glu	Gln	Asn	Met	Ile	Gly	Met	Thr	Pro	Thr	Val	Ile	Ala	
	1010					1015					1020					
Val	His	Tyr	Leu	Asp	Glu	Thr	Glu	Gln	Trp	Glu	Lys	Phe	Gly	Leu	Glu	
1025					1030					1035					1040	
Lys	Arg	Gln	Gly	Ala	Leu	Glu	Leu	Ile	Lys	Lys	Gly	Tyr	Thr	Gln	Gln	
				1045					1050					1055		
Leu	Ala	Phe	Arg	Gln	Pro	Ser	Ser	Ala	Phe	Ala	Ala	Glu	Val	Lys	Arg	
			1060					1065					1070			
Ala	Pro	Ser	Thr	Trp	Leu	Thr	Ala	Tyr	Val	Val	Lys	Val	Phe	Ser	Leu	
	1075						1080					1085				
Ala	Val	Asn	Leu	Ile	Ala	Ile	Asp	Ser	Gln	Val	Leu	Cys	Gly	Ala	Val	
	1090					1095					1100					
Lys	Trp	Leu	Ile	Leu	Glu	Lys	Gln	Lys	Pro	Asp	Gly	Val	Phe	Gln	Glu	
1105				1110						1115					1120	
Asp	Ala	Pro	Val	Ile	His	Gln	Glu	Met	Ile	Gly	Gly	Leu	Arg	Asn	Asn	
				1125					1130					1135		
Asn	Glu	Lys	Asp	Met	Ala	Leu	Thr	Ala	Phe	Val	Leu	Ile	Ser	Leu	Gln	
			1140					1145					1150			
Glu	Ala	Lys	Asp	Ile	Cys	Glu	Glu	Gln	Val	Asn	Ser	Leu	Pro	Gly	Ser	
	1155					1160						1165				
Ile	Thr	Lys	Ala	Gly	Asp	Phe	Leu	Glu	Ala	Asn	Tyr	Met	Asn	Leu	Gln	
	1170					1175						1180				
Arg	Ser	Tyr	Thr	Val	Ala	Ile	Ala	Gly	Tyr	Ala	Leu	Ala	Gln	Met	Gly	
1185					1190					1195					1200	
Arg	Leu	Lys	Gly	Pro	Leu	Leu	Asn	Lys	Phe	Leu	Thr	Thr	Ala	Lys	Asp	
				1205					1210					1215		
Lys	Asn	Arg	Trp	Glu	Asp	Pro	Gly	Lys	Gln	Leu	Tyr	Asn	Val	Glu	Ala	
			1220					1225					1230			
Thr	Ser	Tyr	Ala	Leu	Leu	Ala	Leu	Leu	Gln	Leu	Lys	Asp	Phe	Asp	Phe	
	1235						1240					1245				
Val	Pro	Pro	Val	Val	Arg	Trp	Leu	Asn	Glu	Gln	Arg	Tyr	Tyr	Gly	Gly	
	1250					1255					1260					
Gly	Tyr	Gly	Ser	Thr	Gln	Ala	Thr	Phe	Met	Val	Phe	Gln	Ala	Leu	Ala	
1265					1270					1275					1280	
Gln	Tyr	Gln	Lys	Asp	Ala	Pro	Asp	His	Gln	Glu	Leu	Asn	Leu	Asp	Val	
			1285						1290					1295		
Ser	Leu	Gln	Leu	Pro	Ser	Arg	Ser	Ser	Lys	Ile	Thr	His	Arg	Ile	His	
			1300					1305					1310			
Trp	Glu	Ser	Ala	Ser	Leu	Leu	Arg	Ser	Glu	Glu	Thr	Lys	Glu	Asn	Glu	
	1315						1320					1325				

-continued

Gly Phe Thr Val Thr Ala Glu Gly Lys Gly Gln Gly Thr Leu Ser Val		
1330	1335	1340
Val Thr Met Tyr His Ala Lys Ala Lys Asp Gln Leu Thr Cys Asn Lys		
1345	1350	1355 1360
Phe Asp Leu Lys Val Thr Ile Lys Pro Ala Pro Glu Thr Glu Lys Arg		
	1365 1370	1375
Pro Gln Asp Ala Lys Asn Thr Met Ile Leu Glu Ile Cys Thr Arg Tyr		
	1380 1385	1390
Arg Gly Asp Gln Asp Ala Thr Met Ser Ile Leu Asp Ile Ser Met Met		
	1395 1400	1405
Thr Gly Phe Ala Pro Asp Thr Asp Asp Leu Lys Gln Leu Ala Asn Gly		
	1410 1415	1420
Val Asp Arg Tyr Ile Ser Lys Tyr Glu Leu Asp Lys Ala Phe Ser Asp		
1425	1430	1435 1440
Arg Asn Thr Leu Ile Ile Tyr Leu Asp Lys Val Ser His Ser Glu Asp		
	1445 1450	1455
Asp Cys Leu Ala Phe Lys Val His Gln Tyr Phe Asn Val Glu Leu Ile		
	1460 1465	1470
Gln Pro Gly Ala Val Lys Val Tyr Ala Tyr Tyr Asn Leu Glu Glu Ser		
	1475 1480	1485
Cys Thr Arg Phe Tyr His Pro Glu Lys Glu Asp Gly Lys Leu Asn Lys		
	1490 1495	1500
Leu Cys Arg Asp Glu Leu Cys Arg Cys Ala Glu Glu Asn Cys Phe Ile		
1505	1510	1515 1520
Gln Lys Ser Asp Asp Lys Val Thr Leu Glu Glu Arg Leu Asp Lys Ala		
	1525 1530	1535
Cys Glu Pro Gly Val Asp Tyr Val Tyr Lys Thr Arg Leu Val Lys Val		
	1540 1545	1550
Gln Leu Ser Asn Asp Phe Asp Glu Tyr Ile Met Ala Ile Glu Gln Thr		
	1555 1560	1565
Ile Lys Ser Gly Ser Asp Glu Val Gln Val Gly Gln Gln Arg Thr Phe		
	1570 1575	1580
Ile Ser Pro Ile Lys Cys Arg Glu Ala Leu Lys Leu Glu Glu Lys Lys		
1585	1590	1595 1600
His Tyr Leu Met Trp Gly Leu Ser Ser Asp Phe Trp Gly Glu Lys Pro		
	1605 1610	1615
Asn Leu Ser Tyr Ile Ile Gly Lys Asp Thr Trp Val Glu His Trp Pro		
	1620 1625	1630
Glu Glu Asp Glu Cys Gln Asp Glu Glu Asn Gln Lys Gln Cys Gln Asp		
	1635 1640	1645
Leu Gly Ala Phe Thr Glu Ser Met Val Val Phe Gly Cys Pro Asn		
	1650 1655	1660

<210> SEQ ID NO 6

<211> LENGTH: 449

<212> TYPE: PRT

<213> ORGANISM: Homo Sapiens

<400> SEQUENCE: 6

Met Met Lys Thr Leu Leu Leu Phe Val Gly Leu Leu Leu Thr Trp Glu		
1	5	10 15
Ser Gly Gln Val Leu Gly Asp Gln Thr Val Ser Asp Asn Glu Leu Gln		
	20 25	30
Glu Met Ser Asn Gln Gly Ser Lys Tyr Val Asn Lys Glu Ile Gln Asn		
	35 40	45

-continued

Ala	Val	Asn	Gly	Val	Lys	Gln	Ile	Lys	Thr	Leu	Ile	Glu	Lys	Thr	Asn
50						55					60				
Glu	Glu	Arg	Lys	Thr	Leu	Leu	Ser	Asn	Leu	Glu	Glu	Ala	Lys	Lys	Lys
65					70					75					80
Lys	Glu	Asp	Ala	Leu	Asn	Glu	Thr	Arg	Glu	Ser	Glu	Thr	Lys	Leu	Lys
				85					90					95	
Glu	Leu	Pro	Gly	Val	Cys	Asn	Glu	Thr	Met	Met	Ala	Leu	Trp	Glu	Glu
			100					105					110		
Cys	Lys	Pro	Cys	Leu	Lys	Gln	Thr	Cys	Met	Lys	Phe	Tyr	Ala	Arg	Val
		115					120					125			
Cys	Arg	Ser	Gly	Ser	Gly	Leu	Val	Gly	Arg	Gln	Leu	Glu	Glu	Phe	Leu
	130					135					140				
Asn	Gln	Ser	Ser	Pro	Phe	Tyr	Phe	Trp	Met	Asn	Gly	Asp	Arg	Ile	Asp
145					150					155					160
Ser	Leu	Leu	Glu	Asn	Asp	Arg	Gln	Gln	Thr	His	Met	Leu	Asp	Val	Met
				165					170					175	
Gln	Asp	His	Phe	Ser	Arg	Ala	Ser	Ser	Ile	Ile	Asp	Glu	Leu	Phe	Gln
		180							185				190		
Asp	Arg	Phe	Phe	Thr	Arg	Glu	Pro	Gln	Asp	Thr	Tyr	His	Tyr	Leu	Pro
	195						200				205				
Phe	Ser	Leu	Pro	His	Arg	Arg	Pro	His	Phe	Phe	Phe	Pro	Lys	Ser	Arg
	210					215					220				
Ile	Val	Arg	Ser	Leu	Met	Pro	Phe	Ser	Pro	Tyr	Glu	Pro	Leu	Asn	Phe
225					230					235					240
His	Ala	Met	Phe	Gln	Pro	Phe	Leu	Glu	Met	Ile	His	Glu	Ala	Gln	Gln
			245						250					255	
Ala	Met	Asp	Ile	His	Phe	His	Ser	Pro	Ala	Phe	Gln	His	Pro	Pro	Thr
		260						265					270		
Glu	Phe	Ile	Arg	Glu	Gly	Asp	Asp	Asp	Arg	Thr	Val	Cys	Arg	Glu	Ile
	275					280						285			
Arg	His	Asn	Ser	Thr	Gly	Cys	Leu	Arg	Met	Lys	Asp	Gln	Cys	Asp	Lys
	290					295					300				
Cys	Arg	Glu	Ile	Leu	Ser	Val	Asp	Cys	Ser	Thr	Asn	Asn	Pro	Ser	Gln
305					310					315					320
Ala	Lys	Leu	Arg	Arg	Glu	Leu	Asp	Glu	Ser	Leu	Gln	Val	Ala	Glu	Arg
			325					330					335		
Leu	Thr	Arg	Lys	Tyr	Asn	Glu	Leu	Leu	Lys	Ser	Tyr	Gln	Trp	Lys	Met
		340					345					350			
Leu	Asn	Thr	Ser	Ser	Leu	Leu	Glu	Gln	Leu	Asn	Glu	Gln	Phe	Asn	Trp
	355						360					365			
Val	Ser	Arg	Leu	Ala	Asn	Leu	Thr	Gln	Gly	Glu	Asp	Gln	Tyr	Tyr	Leu
	370					375					380				
Arg	Val	Thr	Thr	Val	Ala	Ser	His	Thr	Ser	Asp	Ser	Asp	Val	Pro	Ser
385					390					395					400
Gly	Val	Thr	Glu	Val	Val	Val	Lys	Leu	Phe	Asp	Ser	Asp	Pro	Ile	Thr
		405						410					415		
Val	Thr	Val	Pro	Val	Glu	Val	Ser	Arg	Lys	Asn	Pro	Lys	Phe	Met	Glu
		420						425				430			
Thr	Val	Ala	Glu	Lys	Ala	Leu	Gln	Glu	Tyr	Arg	Lys	Lys	His	Arg	Glu
		435					440					445			

Glu

<210> SEQ ID NO 7

<211> LENGTH: 1744

-continued

<212> TYPE: PRT

<213> ORGANISM: Homo Sapiens

<400> SEQUENCE: 7

```

Met Arg Leu Leu Trp Gly Leu Ile Trp Ala Ser Ser Phe Phe Thr Leu
 1           5           10          15

Ser Leu Gln Lys Pro Arg Leu Leu Leu Phe Ser Pro Ser Val Val His
 20           25           30

Leu Gly Val Pro Leu Ser Val Gly Val Gln Leu Gln Asp Val Pro Arg
 35           40           45

Gly Gln Val Val Lys Gly Ser Val Phe Leu Arg Asn Pro Ser Arg Asn
 50           55           60

Asn Val Pro Cys Ser Pro Lys Val Asp Phe Thr Leu Ser Ser Glu Arg
 65           70           75           80

Asp Phe Ala Leu Leu Ser Leu Gln Val Pro Leu Lys Asp Ala Lys Ser
 85           90           95

Cys Gly Leu His Gln Leu Leu Arg Gly Pro Glu Val Gln Leu Val Ala
 100          105          110

His Ser Pro Trp Leu Lys Asp Ser Leu Ser Arg Thr Thr Asn Ile Gln
 115          120          125

Gly Ile Asn Leu Leu Phe Ser Ser Arg Arg Gly His Leu Phe Leu Gln
 130          135          140

Thr Asp Gln Pro Ile Tyr Asn Pro Gly Gln Arg Val Arg Tyr Arg Val
 145          150          155          160

Phe Ala Leu Asp Gln Lys Met Arg Pro Ser Thr Asp Thr Ile Thr Val
 165          170          175

Met Val Glu Asn Ser His Gly Leu Arg Val Arg Lys Lys Glu Val Tyr
 180          185          190

Met Pro Ser Ser Ile Phe Gln Asp Asp Phe Val Ile Pro Asp Ile Ser
 195          200          205

Glu Pro Gly Thr Trp Lys Ile Ser Ala Arg Phe Ser Asp Gly Leu Glu
 210          215          220

Ser Asn Ser Ser Thr Gln Phe Glu Val Lys Lys Tyr Val Leu Pro Asn
 225          230          235          240

Phe Glu Val Lys Ile Thr Pro Gly Lys Pro Tyr Ile Leu Thr Val Pro
 245          250          255

Gly His Leu Asp Glu Met Gln Leu Asp Ile Gln Ala Arg Tyr Ile Tyr
 260          265          270

Gly Lys Pro Val Gln Gly Val Ala Tyr Val Arg Phe Gly Leu Leu Asp
 275          280          285

Glu Asp Gly Lys Lys Thr Phe Phe Arg Gly Leu Glu Ser Gln Thr Lys
 290          295          300

Leu Val Asn Gly Gln Ser His Ile Ser Leu Ser Lys Ala Glu Phe Gln
 305          310          315          320

Asp Ala Leu Glu Lys Leu Asn Met Gly Ile Thr Asp Leu Gln Gly Leu
 325          330          335

Arg Leu Tyr Val Ala Ala Ala Ile Ile Glu Ser Pro Gly Gly Glu Met
 340          345          350

Glu Glu Ala Glu Leu Thr Ser Trp Tyr Phe Val Ser Ser Pro Phe Ser
 355          360          365

Leu Asp Leu Ser Lys Thr Lys Arg His Leu Val Pro Gly Ala Pro Phe
 370          375          380

Leu Leu Gln Ala Leu Val Arg Glu Met Ser Gly Ser Pro Ala Ser Gly
 385          390          395          400

```

-continued

Ile	Pro	Val	Lys	Val	Ser	Ala	Thr	Val	Ser	Ser	Pro	Gly	Ser	Val	Pro	405	410	415
Glu	Val	Gln	Asp	Ile	Gln	Gln	Asn	Thr	Asp	Gly	Ser	Gly	Gln	Val	Ser	420	425	430
Ile	Pro	Ile	Ile	Ile	Pro	Gln	Thr	Ile	Ser	Glu	Leu	Gln	Leu	Ser	Val	435	440	445
Ser	Ala	Gly	Ser	Pro	His	Pro	Ala	Ile	Ala	Arg	Leu	Thr	Val	Ala	Ala	450	455	460
Pro	Pro	Ser	Gly	Gly	Pro	Gly	Phe	Leu	Ser	Ile	Glu	Arg	Pro	Asp	Ser	465	470	475
Arg	Pro	Pro	Arg	Val	Gly	Asp	Thr	Leu	Asn	Leu	Asn	Leu	Arg	Ala	Val	485	490	495
Gly	Ser	Gly	Ala	Thr	Phe	Ser	His	Tyr	Tyr	Tyr	Met	Ile	Leu	Ser	Arg	500	505	510
Gly	Gln	Ile	Val	Phe	Met	Asn	Arg	Glu	Pro	Lys	Arg	Thr	Leu	Thr	Ser	515	520	525
Val	Ser	Val	Phe	Val	Asp	His	His	Leu	Ala	Pro	Ser	Phe	Tyr	Phe	Val	530	535	540
Ala	Phe	Tyr	Tyr	His	Gly	Asp	His	Pro	Val	Ala	Asn	Ser	Leu	Arg	Val	545	550	555
Asp	Val	Gln	Ala	Gly	Ala	Cys	Glu	Gly	Lys	Leu	Glu	Leu	Ser	Val	Asp	565	570	575
Gly	Ala	Lys	Gln	Tyr	Arg	Asn	Gly	Glu	Ser	Val	Lys	Leu	His	Leu	Glu	580	585	590
Thr	Asp	Ser	Leu	Ala	Leu	Val	Ala	Leu	Gly	Ala	Leu	Asp	Thr	Ala	Leu	595	600	605
Tyr	Ala	Ala	Gly	Ser	Lys	Ser	His	Lys	Pro	Leu	Asn	Met	Gly	Lys	Val	610	615	620
Phe	Glu	Ala	Met	Asn	Ser	Tyr	Asp	Leu	Gly	Cys	Gly	Pro	Gly	Gly	Gly	625	630	635
Asp	Ser	Ala	Leu	Gln	Val	Phe	Gln	Ala	Ala	Gly	Leu	Ala	Phe	Ser	Asp	645	650	655
Gly	Asp	Gln	Trp	Thr	Leu	Ser	Arg	Lys	Arg	Leu	Ser	Cys	Pro	Lys	Glu	660	665	670
Lys	Thr	Thr	Arg	Lys	Lys	Arg	Asn	Val	Asn	Phe	Gln	Lys	Ala	Ile	Asn	675	680	685
Glu	Lys	Leu	Gly	Gln	Tyr	Ala	Ser	Pro	Thr	Ala	Lys	Arg	Cys	Cys	Gln	690	695	700
Asp	Gly	Val	Thr	Arg	Leu	Pro	Met	Met	Arg	Ser	Cys	Glu	Gln	Arg	Ala	705	710	715
Ala	Arg	Val	Gln	Gln	Pro	Asp	Cys	Arg	Glu	Pro	Phe	Leu	Ser	Cys	Cys	725	730	735
Gln	Phe	Ala	Glu	Ser	Leu	Arg	Lys	Lys	Ser	Arg	Asp	Lys	Gly	Gln	Ala	740	745	750
Gly	Leu	Gln	Arg	Ala	Leu	Glu	Ile	Leu	Gln	Glu	Glu	Asp	Leu	Ile	Asp	755	760	765
Glu	Asp	Asp	Ile	Pro	Val	Arg	Ser	Phe	Phe	Pro	Glu	Asn	Trp	Leu	Trp	770	775	780
Arg	Val	Glu	Thr	Val	Asp	Arg	Phe	Gln	Ile	Leu	Thr	Leu	Trp	Leu	Pro	785	790	795
Asp	Ser	Leu	Thr	Thr	Trp	Glu	Ile	His	Gly	Leu	Ser	Leu	Ser	Lys	Thr	805	810	815
Lys	Gly	Leu	Cys	Val	Ala	Thr	Pro	Val	Gln	Leu	Arg	Val	Phe	Arg	Glu	820	825	830

-continued

Phe His Leu His Leu Arg Leu Pro Met Ser Val Arg Arg Phe Glu Gln
 835 840 845
 Leu Glu Leu Arg Pro Val Leu Tyr Asn Tyr Leu Asp Lys Asn Leu Thr
 850 855 860
 Val Ser Val His Val Ser Pro Val Glu Gly Leu Cys Leu Ala Gly Gly
 865 870 875 880
 Gly Gly Leu Ala Gln Gln Val Leu Val Pro Ala Gly Ser Ala Arg Pro
 885 890 895
 Val Ala Phe Ser Val Val Pro Thr Ala Ala Ala Ala Val Ser Leu Lys
 900 905 910
 Val Val Ala Arg Gly Ser Phe Glu Phe Pro Val Gly Asp Ala Val Ser
 915 920 925
 Lys Val Leu Gln Ile Glu Lys Glu Gly Ala Ile His Arg Glu Glu Leu
 930 935 940
 Val Tyr Glu Leu Asn Pro Leu Asp His Arg Gly Arg Thr Leu Glu Ile
 945 950 955 960
 Pro Gly Asn Ser Asp Pro Asn Met Ile Pro Asp Gly Asp Glu Asn Ser
 965 970 975
 Tyr Val Arg Val Thr Ala Ser Asp Pro Leu Asp Thr Leu Gly Ser Glu
 980 985 990
 Gly Ala Leu Ser Pro Gly Gly Val Ala Ser Leu Leu Arg Leu Pro Arg
 995 1000 1005
 Gly Cys Gly Glu Gln Thr Met Ile Tyr Leu Ala Pro Thr Leu Ala Ala
 1010 1015 1020
 Ser Arg Tyr Leu Asp Lys Thr Glu Gln Trp Ser Thr Leu Pro Pro Glu
 1025 1030 1035 1040
 Thr Lys Asp His Ala Val Asp Leu Ile Gln Lys Gly Tyr Met Arg Ile
 1045 1050 1055
 Gln Gln Phe Arg Lys Ala Asp Gly Ser Tyr Ala Ala Trp Leu Ser Arg
 1060 1065 1070
 Asp Ser Ser Thr Trp Leu Thr Ala Phe Val Leu Lys Val Leu Ser Leu
 1075 1080 1085
 Ala Gln Glu Gln Val Gly Gly Ser Pro Glu Lys Leu Gln Glu Thr Ser
 1090 1095 1100
 Asn Trp Leu Leu Ser Gln Gln Ala Asp Gly Ser Phe Gln Asp Pro
 1105 1110 1115 1120
 Cys Pro Val Leu Asp Arg Ser Met Gln Gly Gly Leu Val Gly Asn Asp
 1125 1130 1135
 Glu Thr Val Ala Leu Thr Ala Phe Val Thr Ile Ala Leu His His Gly
 1140 1145 1150
 Leu Ala Val Phe Gln Asp Glu Gly Ala Glu Pro Leu Lys Gln Arg Val
 1155 1160 1165
 Glu Ala Ser Ile Ser Lys Ala Asn Ser Phe Leu Gly Glu Lys Ala Ser
 1170 1175 1180
 Ala Gly Leu Leu Gly Ala His Ala Ala Ala Ile Thr Ala Tyr Ala Leu
 1185 1190 1195 1200
 Ser Leu Thr Lys Ala Pro Val Asp Leu Leu Gly Val Ala His Asn Asn
 1205 1210 1215
 Leu Met Ala Met Ala Gln Glu Thr Gly Asp Asn Leu Tyr Trp Gly Ser
 1220 1225 1230
 Val Thr Gly Ser Gln Ser Asn Ala Val Ser Pro Thr Pro Ala Pro Arg
 1235 1240 1245
 Asn Pro Ser Asp Pro Met Pro Gln Ala Pro Ala Leu Trp Ile Glu Thr

-continued

1250	1255	1260
Thr Ala Tyr Ala Leu Leu His Leu Leu Leu His Glu Gly Lys Ala Glu 1265	1270	1275 1280
Met Ala Asp Gln Ala Ser Ala Trp Leu Thr Arg Gln Gly Ser Phe Gln 1285	1290	1295
Gly Gly Glu Arg Ser Thr Gln Asp Thr Val Ile Ala Leu Asp Ala Leu 1300	1305	1310
Ser Ala Tyr Trp Ile Ala Ser His Thr Thr Glu Glu Arg Gly Leu Asn 1315	1320	1325
Val Thr Leu Ser Ser Thr Gly Arg Asn Gly Phe Lys Ser His Ala Leu 1330	1335	1340
Gln Leu Asn Asn Arg Gln Ile Arg Gly Leu Glu Glu Glu Leu Gln Phe 1345	1350	1355 1360
Ser Leu Gly Ser Lys Ile Asn Val Lys Val Gly Gly Asn Ser Lys Gly 1365	1370	1375
Thr Leu Lys Val Leu Arg Thr Tyr Asn Val Leu Asp Met Lys Asn Thr 1380	1385	1390
Thr Cys Gln Asp Leu Gln Ile Glu Val Thr Val Lys Gly His Val Glu 1395	1400	1405
Tyr Thr Met Glu Ala Asn Glu Asp Tyr Glu Asp Tyr Glu Tyr Asp Glu 1410	1415	1420
Leu Pro Ala Lys Asp Asp Pro Asp Ala Pro Leu Gln Pro Val Thr Pro 1425	1430	1435 1440
Leu Gln Leu Phe Glu Gly Arg Arg Asn Arg Arg Arg Glu Ala Pro 1445	1450	1455
Lys Val Val Glu Glu Gln Glu Ser Arg Val His Tyr Thr Val Cys Ile 1460	1465	1470
Trp Arg Asn Gly Lys Val Gly Leu Ser Gly Met Ala Ile Ala Asp Val 1475	1480	1485
Thr Leu Leu Ser Gly Phe His Ala Leu Arg Ala Asp Leu Glu Lys Leu 1490	1495	1500
Thr Ser Leu Ser Asp Arg Tyr Val Ser His Phe Glu Thr Glu Gly Pro 1505	1510	1515 1520
His Val Leu Leu Tyr Phe Asp Ser Val Pro Thr Ser Arg Glu Cys Val 1525	1530	1535
Gly Phe Glu Ala Val Gln Glu Val Pro Val Gly Leu Val Gln Pro Ala 1540	1545	1550
Ser Ala Thr Leu Tyr Asp Tyr Tyr Asn Pro Glu Arg Arg Cys Ser Val 1555	1560	1565
Phe Tyr Gly Ala Pro Ser Lys Ser Arg Leu Leu Ala Thr Leu Cys Ser 1570	1575	1580
Ala Glu Val Cys Gln Cys Ala Glu Gly Lys Cys Pro Arg Gln Arg Arg 1585	1590	1595 1600
Ala Leu Glu Arg Gly Leu Gln Asp Glu Asp Gly Tyr Arg Met Lys Phe 1605	1610	1615
Ala Cys Tyr Tyr Pro Arg Val Glu Tyr Gly Phe Gln Val Lys Val Leu 1620	1625	1630
Arg Glu Asp Ser Arg Ala Ala Phe Arg Leu Phe Glu Thr Lys Ile Thr 1635	1640	1645
Gln Val Leu His Phe Thr Lys Asp Val Lys Ala Ala Ala Asn Gln Met 1650	1655	1660
Arg Asn Phe Leu Val Arg Ala Ser Cys Arg Leu Arg Leu Glu Pro Gly 1665	1670	1675 1680

-continued

Lys Glu Tyr Leu Ile Met Gly Leu Asp Gly Ala Thr Tyr Asp Leu Glu
 1685 1690 1695
 Gly His Pro Gln Tyr Leu Leu Asp Ser Asn Ser Trp Ile Glu Glu Met
 1700 1705 1710
 Pro Ser Glu Arg Leu Cys Arg Ser Thr Arg Gln Arg Ala Ala Cys Ala
 1715 1720 1725
 Gln Leu Asn Asp Phe Leu Gln Glu Tyr Gly Thr Gln Gly Cys Gln Val
 1730 1735 1740

<210> SEQ ID NO 8
 <211> LENGTH: 375
 <212> TYPE: PRT
 <213> ORGANISM: Homo Sapiens

<400> SEQUENCE: 8

Met Glu Glu Glu Ile Ala Ala Leu Val Ile Asp Asn Gly Ser Gly Met
 1 5 10 15
 Cys Lys Ala Gly Phe Ala Gly Asp Asp Ala Pro Arg Ala Val Phe Pro
 20 25 30
 Ser Ile Val Gly Arg Pro Arg His Gln Gly Val Met Val Gly Met Gly
 35 40 45
 Gln Lys Asp Ser Tyr Val Gly Asp Glu Ala Gln Ser Lys Arg Gly Ile
 50 55 60
 Leu Thr Leu Lys Tyr Pro Ile Glu His Gly Ile Val Thr Asn Trp Asp
 65 70 75 80
 Asp Met Glu Lys Ile Trp His His Thr Phe Tyr Asn Glu Leu Arg Val
 85 90 95
 Ala Pro Glu Glu His Pro Val Leu Leu Thr Glu Ala Pro Leu Asn Pro
 100 105 110
 Lys Ala Asn Arg Glu Lys Met Thr Gln Ile Met Phe Glu Thr Phe Asn
 115 120 125
 Thr Pro Ala Met Tyr Val Ala Ile Gln Ala Val Leu Ser Leu Tyr Ala
 130 135 140
 Ser Gly Arg Thr Thr Gly Ile Val Met Asp Ser Gly Asp Gly Val Thr
 145 150 155 160
 His Thr Val Pro Ile Tyr Glu Gly Tyr Ala Leu Pro His Ala Ile Leu
 165 170 175
 Arg Leu Asp Leu Ala Gly Arg Asp Leu Thr Asp Tyr Leu Met Lys Ile
 180 185 190
 Leu Thr Glu Arg Gly Tyr Ser Phe Thr Thr Thr Ala Glu Arg Glu Ile
 195 200 205
 Val Arg Asp Ile Lys Glu Lys Leu Cys Tyr Val Ala Leu Asp Phe Glu
 210 215 220
 Gln Glu Met Ala Thr Ala Ala Ser Ser Ser Ser Leu Glu Lys Ser Tyr
 225 230 235 240
 Glu Leu Pro Asp Gly Gln Val Ile Thr Ile Gly Asn Glu Arg Phe Arg
 245 250 255
 Cys Pro Glu Ala Leu Phe Gln Pro Ser Phe Leu Gly Met Glu Ser Cys
 260 265 270
 Gly Ile His Glu Thr Thr Phe Asn Ser Ile Met Lys Cys Asp Val Asp
 275 280 285
 Ile Arg Lys Asp Leu Tyr Ala Asn Thr Val Leu Ser Gly Gly Thr Thr
 290 295 300
 Met Tyr Pro Gly Ile Ala Asp Arg Met Gln Lys Glu Ile Thr Ala Leu
 305 310 315 320

-continued

Lys	Lys	Thr	Pro	Lys	Ser	Pro	Val	Gly	Val	Gln	Pro	Ile	Leu	Asn	Glu
				325					330					335	
His	Thr	Phe	Cys	Ala	Gly	Met	Ser	Lys	Tyr	Gln	Glu	Asp	Thr	Cys	Tyr
			340					345					350		
Gly	Asp	Ala	Gly	Ser	Ala	Phe	Ala	Val	His	Asp	Leu	Glu	Glu	Asp	Thr
		355					360					365			
Trp	Tyr	Ala	Thr	Gly	Ile	Leu	Ser	Phe	Asp	Lys	Ser	Cys	Ala	Val	Ala
	370					375					380				
Glu	Tyr	Gly	Val	Tyr	Val	Lys	Val	Thr	Ser	Ile	Gln	Asp	Trp	Val	Gln
385					390					395					400
Lys	Thr	Ile	Ala	Glu	Asn										
				405											

The invention claimed is:

1. A method of diagnosing Alzheimer's disease in a subject, the method comprising detecting a differentially expressed protein, said protein being clusterin precursor protein (Swiss-PROT Accession number (SPN) P10909), in a sample of blood or plasma from said subject, wherein a decrease in the expression of said clusterin precursor protein compared to the expression of said protein in a control subject is indicative of Alzheimer's disease in the subject undergoing diagnosis.

2. A method according to claim 1, wherein the clusterin precursor protein is detected in the sample using (i) an antibody specific to said clusterin precursor protein; (ii) via detection of an autoantibody specific to said clusterin precursor protein, or by mass spectrometry.

3. A method according to claim 1, wherein the clusterin precursor protein is detected using 2D gel electrophoresis.

4. A method according to claim 2 wherein the sample is immobilised on a solid support.

5. A method according to claim 1, wherein said sample is a blood sample.

6. A method according to claim 1, further comprising detecting said clusterin precursor protein in combination with detection of an increase or decrease of the expression level of at least one other differentially expressed protein biomarker of Alzheimer's disease selected from the group consisting of the following proteins or a fragment thereof: apolipoprotein A-N precursor (SPN P06727), as determined by an increase in the expression thereof, apolipoprotein C-III precursor (SPN P02656), as determined by an increase in the expression

thereof, transthyretin (SPN P02766), as determined by an increase in the expression thereof, galectin 7 (SPN P47929), as determined by an increase in the expression thereof, complement C4 precursor (SPN P01028), as determined by an increase in the expression thereof, complement factor H (SPN P08603), as determined by an increase in the expression thereof, S100 calcium binding protein or ceruloplasmin precursor (SPN P00450), as determined by an increase in the expression thereof, histone H2B. a/g/h/k/l (SPN P62807), as determined by a decrease in the expression thereof, Ig lambda chain C region (SPN P01842), as determined by an increase in the expression thereof, inter-alpha-trypsin inhibitor heavy chain H4 precursor (SPN Q14624), as determined by an increase in the expression thereof, complement C3 precursor (SPN P01024), as determined by a decrease in the expression thereof, gamma/beta actin (SPN P63261), as determined by an increase in the expression thereof, haptoglobin precursor (SPN P00738), as determined by an increase in the expression thereof, alpha-2-macroglobulin (SPN P01023), as determined by an increase in the expression thereof and the serum albumin precursor isoform (SPN P02768), as determined by an increase in the expression thereof, said determination(s) being in comparison to the expression of the same protein(s) in a control subject.

7. A method according to claim 6 which comprises detecting more than one differentially expressed protein.

8. A method according to claim 7 which comprises detecting four or more differentially expressed proteins.

* * * * *

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 7,897,361 B2
APPLICATION NO. : 11/664076
DATED : March 1, 2011
INVENTOR(S) : Jules Westbrook et al.

Page 1 of 1


It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

Title Page; item (76);

Page 1, under "Inventors", 6th listed inventor, change "Stephen Lynham" to "Steven Lynham"; and

10th listed inventor, change city of residence from "Offenbad" to
"Offenbach".

Signed and Sealed this
Twenty-seventh Day of September, 2011

A handwritten signature in black ink that reads "David J. Kappos". The signature is written in a cursive, flowing style with a large initial 'D' and a stylized 'K'.

David J. Kappos
Director of the United States Patent and Trademark Office

专利名称(译)	诊断阿尔茨海默病的方法		
公开(公告)号	US7897361	公开(公告)日	2011-03-01
申请号	US11/664076	申请日	2005-09-29
[标]申请(专利权)人(译)	WESTBROOK JULES BYERS HELEN MALCOLM WARD LOVESTONE SIMON HYE ABDUL LYNHAM史蒂芬 JOUBERT RICHARD PREFOT PETRA 库恩KARSTEN BAUMANN CHRISTIAN SCHAFER JURGEN PRINZ托尔斯滕 KIENLE STEFAN		
申请(专利权)人(译)	WESTBROOK JULES BYERS HELEN MALCOLM WARD LOVESTONE SIMON HYE ABDUL LYNHAM史蒂芬 JOUBERT RICHARD PREFOT PETRA 库恩KARSTEN BAUMANN CHRISTIAN SCHAEFER JUERGEN PRINZ托尔斯滕 KIENLE STEFAN		
当前申请(专利权)人(译)	WESTBROOK JULES BYERS HELEN MALCOLM WARD LOVESTONE SIMON HYE ABDUL LYNHAM史蒂芬 JOUBERT RICHARD PREFOT PETRA 库恩KARSTEN BAUMANN CHRISTIAN SCHAEFER JUERGEN PRINZ托尔斯滕 KIENLE STEFAN		
[标]发明人	WESTBROOK JULES BYERS HELEN WARD MALCOLM LOVESTONE SIMON HYE ABDUL LYNHAM STEPHEN JOUBERT RICHARD		

PREFOT PETRA
KUHN KARSTEN
BAUMANN CHRISTIAN
SCHAEFER JUERGEN
PRINZ THORSTEN
KIENLE STEFAN

发明人
WESTBROOK, JULES
BYERS, HELEN
WARD, MALCOLM
LOVESTONE, SIMON
HYE, ABDUL
LYNHAM, STEPHEN
JOUBERT, RICHARD
PREFOT, PETRA
KUHN, KARSTEN
BAUMANN, CHRISTIAN
SCHAEFER, JUERGEN
PRINZ, THORSTEN
KIENLE, STEFAN

IPC分类号 C12Q1/68 G01N33/53 G01N33/566 G01N33/567 G01N33/68

CPC分类号 G01N33/6896 G01N2800/52 G01N2800/2821 A61P25/28

优先权 2004021639 2004-09-29 GB

其他公开文献 US20080070995A1

外部链接 [Espacenet](#) [USPTO](#)

摘要(译)

提供了与阿尔茨海默病有关的方法和组合物。具体地，提供了相对于它们在正常状态下的表达而在阿尔茨海默病状态中差异表达的蛋白质。鉴定并描述了与阿尔茨海默病相关的蛋白质。还提供了使用差异表达的蛋白质诊断阿尔茨海默病的方法，以及用于预防和治疗阿尔茨海默病的化合物的鉴定和治疗用途的方法。

