



US 20190310265A1

(19) **United States**

(12) **Patent Application Publication**

Van Eyk et al.

(10) **Pub. No.: US 2019/0310265 A1**

(43) **Pub. Date: Oct. 10, 2019**

(54) **CITRULLINATED BRAIN AND NEUROLOGICAL PROTEINS AS BIOMARKERS OF BRAIN INJURY OR NEURODEGENERATION**

(60) Provisional application No. 61/610,034, filed on Mar. 13, 2012.

Publication Classification

(71) Applicant: **THE JOHNS HOPKINS UNIVERSITY, BALTIMORE, MD (US)**

(51) **Int. Cl.**
G01N 33/68 (2006.01)
H01J 49/00 (2006.01)
G01N 33/53 (2006.01)

(72) Inventors: **Jennifer E. Van Eyk, Baltimore, MD (US); Allen Dale Everett, Baltimore, MD (US); Zhicheng Jin, Baltimore, MD (US)**

(52) **U.S. Cl.**
CPC *G01N 33/6848* (2013.01); *G01N 33/6842* (2013.01); *H01J 49/0027* (2013.01); *G01N 2800/28* (2013.01); *H01J 49/0077* (2013.01); *G01N 2440/18* (2013.01); *G01N 33/5306* (2013.01)

(73) Assignee: **THE JOHNS HOPKINS UNIVERSITY, BALTIMORE, MD (US)**

(57) **ABSTRACT**

(21) Appl. No.: **16/433,106**

(22) Filed: **Jun. 6, 2019**

The present invention relates to the field of biomarkers. More specifically, the present invention relates to biomarkers useful in diagnosing brain injury or neurodegeneration. In one embodiment, a method for diagnosing brain injury in a patient comprises the steps of (a) obtaining a sample from the patient; (b) determining the ratio of citrullinated to unmodified arginine residues at one or more arginine residues of one or more brain injury biomarker proteins; and (c) correlating the ratio to a patient having brain injury or to a patient not having brain injury, thereby providing the diagnosis.


Related U.S. Application Data

(60) Division of application No. 15/636,076, filed on Jun. 28, 2017, now Pat. No. 10,365,288, which is a continuation-in-part of application No. 14/384,848, filed on Sep. 12, 2014, now Pat. No. 9,709,573, filed as application No. PCT/US2013/031012 on Mar. 13, 2013.

Specification includes a Sequence Listing.

ZJ_110418_MBP_Lys-C #2699 RT: 29.05 AV: 1 NL: 2.47E4
T: ITMS + c NSI d Full ms2 746.91 @cid35.00 [195.00-1505.00]_1065.66

FIG. 1A






FIG. 1B





FIG. 1C



ZJ_110418_MBP_Lys-C #3147 RT: 31.67 AV: 1 NL: 7.87E4
T: ITMS + c NSI d Full ms2 688.32@cid35.00 [175.00-1390.00]




FIG. 2A




FIG. 2B

ZJ_110419_MSP_Lys-C #2332 RT: 26.96 AV: 1 NL: 3.44E4
T: ITMS + c NSI d Full ms2 626.64@3435.00 [160.00-1690.00]




FIG. 3A




FIG. 3B

Bovine: MDCCTE~~S~~ACS KPDDDILDIP LDDPCANAAA AKIQASFRGH MARKKIKSGE RGRKGGPGG
 60

Human: MDCCTE~~S~~ACS KPDDDILDIP LDDPCANAAA AKIQASFRGH MARKKIKSGE RGRKGGPGG
 60

Bovine: EGGAG~~S~~ARGG AGGGPSGD 78 (SEQ ID NO:6)

Human: EGGAG~~S~~ARGG AGGGPSGD 78 (SEQ ID NO:1)

FIG. 4




FIG. 5A





FIG. 5B

FIG. 6A






FIG. 6B




FIG. 7A




FIG. 7B

60 MHHHHHENLYFQGAIA MDCCTENACS KPDDDDI LDIP LDDPGANAAA AKIQASH **R**GHMA **R**KKIKSGE
1 10 30 50

60 **RGR**KGP GPGG PGGAGV **R**GGAGGGP SGDT **R**
78

FIG. 8




FIG. 9A




FIG. 9B

```

Human  MERRRITSAARRSYVSSGEMVVGGLAPGR  LGPGTRLSLARMPPEPLFTRVDFSLAGALNA  60
-----*-----*-----*-----*-----*-----*-----*-----*-----*
Bovine  MERRRVTSAPRRSYVSSSEMVG  GRR  LGPGTRLSLARMPPEPLPARVDFSLAGALNS  56

Human  GFKETRASERAEMMELNDPFASYIEKVRFL  EQQNFALAAELNQLRAKEFTKLADVYQAEI  120
-----*-----*-----*-----*-----*-----*-----*-----*-----*
Bovine  GFKETRASERAEMMELNDPFASYIEKVRFL  EQQNKALAAELNQLRAKEFTKLADVYQAEI  116

Human  RELRELDQLTANSRLEVERDNLAQQLAT  VPQKLQDETNLRLEAENNLAARQEADEAT  160
-----*-----*-----*-----*-----*-----*-----*-----*-----*
Bovine  RELKLELDQLTANSARLEVERDNLAQQLGT  LRQKLQDETNQRLEAENNLAARQEADEAT  176

Human  LARLDLERKIESLEEEIRFLRKIHSEEVRE  LQEQLARQQVHVVELDVAKPDLTAALKEIRT  240
-----*-----*-----*-----*-----*-----*-----*-----*-----*
Bovine  LARLDLERKIESLEEEIRFLRKIHSEEVRE  LQEQLAQQQVHVEMDVAKPDLTAALKEIRT  236

Human  QYEANASSRNHEAEWYRSKFADLTAAAR  HAELLRQAKHEANDYRRQLQSLTCDLESIR  300
-----*-----*-----*-----*-----*-----*-----*-----*-----*
Bovine  QYEAASSRNHEAEWYRSKFADLNDAAAR  HAELLRQAKHEANDYRRQLQALTCDESIR  296

Human  GTNESLERQMKEQEERHVREAASYQELAR  LEEBQQSLDENARHLQEYQDLLENVKLALD  360
-----*-----*-----*-----*-----*-----*-----*-----*-----*
Bovine  GTNESLERQMKEQEERHAREAAASYQELAR  LEEBQQSLDENARHLQEYQDLLENVKLALD  356

Human  IEIATYRKLLEGEENRITIPVQTFSNLQIR  ETSLDTSVSEGHKRNIVVKTVEKRDGEV  420
-----*-----*-----*-----*-----*-----*-----*-----*-----*
Bovine  IEIATYRKLLEGEENRITIPVQTFSNLQIR  ETSLDTSVSEGHKRNIIYKTVEMPDGEV  416

Human  IKESKQEHKD  VM  422  (SEQ ID NO:3)
-----*-----*-----*-----*-----*-----*-----*-----*-----*
Bovine  IKESKQEHKD  VM  428  (SEQ ID NO:7)
-----*-----*-----*-----*-----*-----*-----*-----*-----*

```

FIG. 10




FIG. 11

FIG. 12A





FIG. 12B






FIG. 13A




FIG. 13B




FIG. 13C




FIG. 14

**CITRULLINATED BRAIN AND
NEUROLOGICAL PROTEINS AS
BIOMARKERS OF BRAIN INJURY OR
NEURODEGENERATION**

**CROSS-REFERENCE TO RELATED
APPLICATIONS**

[0001] This application is a divisional of U.S. Ser. No. 15/636,076, filed on Jun. 28, 2017, which is a continuation of patent application U.S. Ser. No. 14/384,848, filed on Sep. 12, 2014, now U.S. Pat. No. 9,709,573, issued on Jul. 18, 2017, which is a U.S. National Stage Application under 35 U.S.C. § 371 of International PCT Application No. PCT/US2013/031012, having an international filing date of Mar. 13, 2013, which claims the benefit of U.S. Provisional Application No. 61/610,034, filed on Mar. 13, 2012, the contents of all of which are incorporated by reference herein in their entireties.

STATEMENT OF GOVERNMENTAL INTEREST

[0002] This invention was made with U.S. government support under grant nos. 1R01HL091759-02, 5U54HL090515-02, and NHLBI-HV-10-05(2). The government has certain rights in the invention.

FIELD OF THE INVENTION

[0003] The present invention relates to the field of biomarkers. More specifically, the present invention relates to biomarkers useful in diagnosing brain injury or neurodegeneration.

**INCORPORATION-BY-REFERENCE OF
MATERIAL SUBMITTED ELECTRONICALLY**

[0004] This application contains a Sequence Listing, which has been submitted electronically via EFS-Web as an ASCII text file entitled "P11929-02_Sequence_Listing.txt." The Sequence Listing is 75,410 bytes in size, and was created on Mar. 12, 2013. It is hereby incorporated by reference in its entirety.

BACKGROUND OF THE INVENTION

[0005] Clinical tools such as physical exam, and CNS imaging (CT Scan or MRI) are subjective, not widely available, not sensitive or specific enough and too costly to identify the infant, child or adult with CNS injury. This can include individuals on life support or cardiopulmonary bypass, trauma, loss of oxygen, etc, regardless of the initial injury or disease. There is a great clinical need to identify patients with CNS or brain injury and especially subclinical injury because these infants, children and adults are at significant risk of progressing to overt stroke and development of cognitive and motor loss, dementia and poor mental performance. In addition, accurate and sensitive identification of CNS injury by circulating biomarkers will provide an objective gold standard to test and compare new therapeutic modalities for efficacy.

SUMMARY OF THE INVENTION

[0006] The present invention is based, at least in part, on the discovery of unique post-translational modifications (e.g., citrullination) of CNS proteins neurogranin (NRGN), myelin basic protein (MBP), glial fibrillary acid protein

(GFAP), tubulin beta-4B chain, tubulin alpha-1B chain, CNPase, PPIA, Septin-7, Elongation factor1-alpha2, TPPP, TPPP3, Ermin Isoform 2, and NDRG2 Isoform 2. Other proteins include astrotactin 1 (ASTN1); brain angiogenesis inhibitor 3 (BAI3); carnosine dipeptidase 1 (CNDP1); ERMIN; glutamate receptor metabotropic 3 (GRM3); kelch-like protein 32 (KLH32); melanoma antigen family E,2 (MAGE2); myelin basic protein (MBP); neuregulin 3 (NRG3); oligodendrocyte myelin glycoprotein (OMG); solute carrier family 39 (zinc transporter); reticulon 1 (RTN1); and peptidylarginine deiminase (types 1-4 and 6) (PAD) (including PAD-2), are found circulating in patients with brain injury. Thus, the present invention provides methods for the detection and quantification of one or more modified protein or specific amino acid residues on one or more brain injury biomarker proteins. As described further herein, the present invention also provides methods for the detection and quantification of autoantibodies to the unmodified and/or modified brain injury biomarker proteins listed above.

[0007] Accordingly, the identification and development of assays for post-translation modifications and unmodified forms of brain specific circulating proteins such as NRGN, MBP and GFAP provide a new specific diagnostic of brain injury providing the exquisite brain specificity for an accurate and fast diagnosis. In addition, modified forms of NRGN, MBP and GFAP provide information on the biology of injury that can be useful for defining the phase of injury or recovery that can be used for targeting specific therapies to personalize and improve outcome. Ratios of citrullinated to non-citrullinated proteins could be used to determine the risk or progression of CNS injury, stroke, reduced mental capacity, and neurodegenerative diseases.

[0008] As described in certain embodiments, unmodified and citrullinated NRGN, MBP and/or GFAP represent a new diagnostic assay for brain injury. The present invention provides kits using antibodies, aptamers, and mass spectrometry based methods for detection of citrullinated proteins and or modified amino acid residues. The present invention can be used to detect and/or quantify brain injury biomarkers in various body fluids (including plasma, serum, cerebral spinal fluid) and tissue and cells.

[0009] In specific embodiments, the citrullination at residues at NRGN: R38, R43, R51, R53, R68; MBP: R31, R43, R49, R65, R107, R113, R162, R169; and GFAP: R88, R105, R124, R126, R136, R173, R217, R258, R270, R286, R287, R367, R406, and combinations of any of the foregoing can be used to diagnosis brain injury in a patient. In more specific embodiments, the detection and quantification can be directed at the ratio of modified to unmodified proteins or specific amino acid residues.

[0010] The present invention is applicable to the prediction of the susceptibility or presence of both subclinical and overt brain injury. The present invention further provides protein diagnostic(s)/prognostic(s) useful for identifying infants, children and adults with subclinical brain injury, in which routine clinical assessments are normal, to prevent progression to overt stroke. More specifically, protein diagnostic(s)/prognostic(s) of the present invention can be used to assess and monitor efficacy of therapies in infants, children and adults. In fact, the protein diagnostic(s)/prognostic(s) described herein can be used to identify brain injury in ill infants, children and adults at risk for brain injury and to predict outcomes.

[0011] In other embodiments, protein diagnostic(s)/prognostic(s) can be used to (1) identify brain injury in children and adults on life support or cardiopulmonary bypass including during surgery (regardless of initialing injury or disease) to assess potential neurological injury; (2) identify the permeability of the blood brain barrier; and/or (3) identify degenerative brain disease.

[0012] Furthermore, as citrullination of proteins has been proposed to increase immunogenicity of some proteins, the identification of citrullinated brain proteins may be very important for diagnosis of chronic neurodegenerative diseases such as chronic traumatic encephalopathy, Alzheimer's disease, multiple sclerosis and Parkinson's disease. It is possible that auto-antibodies against the various citrullinated forms of these proteins could act as auto-antigens following brain injury exacerbating brain injury. Thus, detection (or blocking) of these auto-antibodies could be used for improved prognosis, risk stratification, or therapy.

[0013] Accordingly, the present invention provides methods for diagnosing brain injury in a patient. In one embodiment, a method for diagnosing brain injury in a patient comprises the steps of (a) obtaining a sample from the patient; (b) determining the ratio of citrullinated to unmodified arginine residues at one or more arginine residues of one or more brain injury biomarker proteins; and (c) correlating the ratio to a patient having brain injury or to a patient not having brain injury, thereby providing the diagnosis.

[0014] The sample can be selected from the group consisting of blood, peripheral blood, serum, plasma, cerebrospinal fluid, urine, saliva, stool and synovial fluid. In a specific embodiment, the sample is blood, plasma serum, cerebrospinal fluid (CSF), or urine. In a more specific embodiment, the sample is CSF. In another specific embodiment, the sample is blood. In an alternative embodiment, the sample is serum.

[0015] In certain embodiments, the determining step is accomplished using mass spectrometry. In a specific embodiment, the determining step is accomplished using multiple reaction monitoring mass spectrometry (MRM-MS). In other embodiments, the determining step is accomplished using an immunoassay. In further embodiments, aptamers, peptoids, or other capture/detection systems are used.

[0016] In a specific embodiment, the one or more brain injury biomarker proteins is neurogranin (NRGN), myelin basic protein (MBP), glial fibrillary acid protein (GFAP), peptidylarginine deiminase (PAD), isoforms thereof, post-translationally modified forms thereof, or combinations of any of the foregoing. In a more specific embodiment, the brain injury biomarker protein is NRGN, isoforms thereof, or post-translationally modified forms thereof. In another specific embodiment, the brain injury biomarker protein is MBP, isoforms thereof, or post-translationally modified forms thereof. In yet another embodiment, the brain injury biomarker protein is GFAP, isoforms thereof, or post-translationally modified forms thereof. In a further embodiment, in the brain injury biomarker protein is PAD-2, isoforms thereof, or post-translationally modified forms thereof.

[0017] In yet another embodiment, the brain injury biomarker protein is one or more proteins selected from the group consisting of tubulin beta-4B chain; tubulin alpha-1B chain; CNPase; PPIA; Septin-7; Elongation factor1-alpha2; TPPP; TPPP3; Ermin Isoform 2; NDRG2 Isoform 2, astrocytactin 1 (ASTN1), brain angiogenesis inhibitor 3 (BAI3);

carosine dipeptidase 1 (CNDP1); ERMIN; glial fibrillary acidic protein (GFAP); glutamate receptor metabotropic 3 (GRM3); kelch-like protein 32 (KLH32); melanoma antigen family E,2 (MAGE2); myelin basic protein (MBP); neuregulin 3 (NRG3); neurogranin (NRGN); oligodendrocyte myelin glycoprotein (OMG); solute carrier family 39 (zinc transporter); reticulon 1 (RTN1); and peptidylarginine deiminase (types 1-4 and 6) (PAD); isoforms thereof; post-translationally modified forms thereof; or combinations of any of the foregoing.

[0018] The present invention also provides a method for diagnosing brain injury in a patient comprising the steps of (a) determining the ratio of one or more citrullinated peptides to the corresponding unmodified peptides in a sample collected from the patient using mass spectrometry; and (b) correlating the ratio to a patient having brain injury or to a patient not having brain injury, thereby providing the diagnosis. In an alternative embodiment, a method for diagnosing brain injury in a patient comprises the steps of (a) determining the ratio of one or more citrullinated peptides to the corresponding unmodified peptides in a sample collected from the patient using mass spectrometry; and (b) comparing the ratio with predefined ratios of the same peptides that correlate to a patient having brain injury and predefined ratios of the same peptides that correlate to a patient not having brain injury, wherein a correlation to one of the predefined ratios provides the diagnosis.

[0019] In such embodiments, the one or more peptides is selected from the group consisting of tubulin beta-4B chain, tubulin alpha-1B chain, CNPase, PPIA, Septin-7, Elongation factor1-alpha2, TPPP, TPPP3, Ermin Isoform 2, NDRG2 Isoform 2, ASTN1, BAI3, CNDP1, ERMIN, GFAP, GRM3, KLH32, MAGE2, MBP, NRG3, NRGN, OMG, SLC39A12, RTN1, MT3, PAD (including PAD-2), isoforms thereof, post-translationally modified forms thereof, or combinations of any of the foregoing. Alternatively, the one or more peptide is NRGN, MBP, GFAP, PAD or a combination thereof. In other embodiments, the one or more peptide is NRGN, MBP, GFAP, isoforms thereof, post-translationally modified forms thereof, or combinations of any of the foregoing.

[0020] In a specific embodiment, the present invention provides a method for diagnosing brain injury in a patient comprising the steps of (a) determining the ratio of one or more post-translationally modified peptides to the corresponding unmodified peptides in a sample collected from the patient using MRM-MS; and (b) correlating the ratio to a patient having brain injury or to a patient not having brain injury, thereby providing the diagnosis. In another specific embodiment, a method for diagnosing brain injury in a patient comprises the steps of (a) determining the ratio of one or more post-translationally modified peptides to the corresponding unmodified peptides in a sample collected from the patient using MRM-MS; and (b) comparing the one or more ratios with predefined ratios of the same post-translationally modified/unmodified peptides that correlate to a patient having brain injury and predefined ratios of the same post-translationally modified/unmodified peptides that correlate to a patient not having brain injury, wherein a correlation to one of the predefined ratios provides the diagnosis.

[0021] In such embodiments, the post-translational modification is citrullination, oxidation, methylation, phosphorylation, cysteinylolation s-nitrosation, s-glutathylation, or a

combination thereof. Moreover, the one or more peptides can be selected from the group consisting of tubulin beta-4B chain, tubulin alpha-1B chain, CNPase, PPIA, Septin-7, Elongation factor1-alpha2, TPPP, TPPP3, Ermin Isoform 2, NDRG2 Isoform 2, ASTN1, BAI3, CNDP1, ERMIN, GFAP, GRM3, KLH32, MAGE2, MBP, NRG3, NRGN, OMG, SLC39A12, RTN1, PAD, isoforms thereof, post-translationally modified forms thereof, or combinations of any of the foregoing. In a specific embodiment, the one or more peptide is NRGN, MBP, GFAP, PAD, isoforms thereof, post-translationally modified forms thereof, or combinations of any of the foregoing.

[0022] In another embodiment, a method for diagnosing brain injury in a patient comprises the steps of (a) determining the ratio of one or more citrullinated peptides to the corresponding unmodified peptides in a sample collected from the patient using mass spectrometry; (b) determining the ratio of one or more post-translationally modified peptides to the corresponding unmodified peptides in the same sample collected from the patient using MRM-MS; (c) comparing the one or more citrullinated:unmodified peptide ratios to one or more post-translationally modified:unmodified peptide ratios; and (d) correlating the compared ratios to a patient having brain injury or to a patient not having brain injury, thereby providing the diagnosis.

[0023] In another embodiment, a method for diagnosing brain injury in a patient comprises the steps of (a) determining the degree of citrullination of one or more arginine sites of one or more of NRGN, PAD, MBP GFAP, isoforms thereof, post-translationally modified forms thereof, or combinations of any of the foregoing; and (b) correlating the degree of citrullination to a patient having brain injury or to a patient not having brain injury, thereby providing the diagnosis.

[0024] In particular embodiments, the peptides are one or more peptides as shown in Tables 5-9, 11-12, and 14-17. In other embodiments, the peptides comprise an amino acid sequence of about 8 to about 45 amino acid residues of NRGN, MBP, GFAP, and PAD, isoforms thereof, post-translationally modified forms thereof, or combinations of any of the foregoing. In further embodiments, the peptides comprise an amino acid sequence of about 8 to about 45 amino acid residues of tubulin beta-4B chain, tubulin alpha-1B chain, CNPase, PPIA, Septin-7, Elongation factor1-alpha2, TPPP, TPPP3, Ermin Isoform 2, NDRG2 Isoform 2, ASTN1, BAI3, CNDP1, ERMIN, GFAP, GRM3, KLH32, MAGE2, MBP, NRG3, NRGN, OMG, SLC39A12, RTN1, PAD, isoforms thereof, post-translationally modified forms thereof, or combinations of any of the foregoing.

[0025] In another aspect, the present invention provides methods for determining the concentration or levels of citrullinated brain injury biomarker proteins/peptides. In one embodiment, a method for diagnosing brain injury in a patient comprises the steps of (a) measuring the level of one or more citrullinated brain injury biomarker proteins in a sample collected from the patient; and (b) comparing the level of the one or more biomarkers with predefined levels of the same biomarkers that correlate to a patient having brain injury and predefined levels of the same biomarkers that correlate to a patient not having brain injury, wherein a correlation to one of the predefined levels provides the diagnosis.

[0026] In particular embodiments, the one or more brain injury biomarker proteins is tubulin beta-4B chain, tubulin

alpha-1B chain, CNPase, PPIA, Septin-7, Elongation factor1-alpha2, TPPP, TPPP3, Ermin Isoform 2, NDRG2 Isoform 2, ASTN1, BAI3, CNDP1, ERMIN, GFAP, GRM3, KLH32, MAGE2, MBP, NRG3, NRGN, OMG, SLC39A12, RTN1, MT3, PAD, isoforms thereof, post-translationally modified forms thereof, or combinations of any of the foregoing. Alternatively, the one or more brain injury biomarker proteins is NRGN, MBP, GFAP, PAD, isoforms thereof, post-translationally modified forms thereof, or combinations of any of the foregoing. In a specific embodiment, the brain injury biomarker protein is NRGN, isoforms thereof, or post-translationally modified forms thereof. In another embodiment, the brain injury biomarker protein is MBP, isoforms thereof, or post-translationally modified forms thereof. In yet another embodiment, the brain injury biomarker protein is GFAP, isoforms thereof, or post-translationally modified forms thereof. In a further embodiment, the brain injury biomarker protein is PAD, isoforms thereof, or post-translationally modified forms thereof.

[0027] In such embodiments, the sample can be selected from the group consisting of blood, peripheral blood, serum, plasma, cerebrospinal fluid, urine, saliva, stool and synovial fluid. In a specific embodiment, the sample is blood, plasma serum, cerebrospinal fluid (CSF), or urine. In a more specific embodiment, the sample is CSF. In another specific embodiment, the sample is blood. In yet another embodiment, the sample is serum.

[0028] In certain embodiments, the determining step is accomplished using mass spectrometry. In a specific embodiment, the determining step is accomplished using multiple reaction monitoring mass spectrometry (MRM-MS). In other embodiments, the determining step is accomplished using an immunoassay. In further embodiments, aptamers, peptoids, or other capture/detection systems are used.

[0029] The present invention also provides a method for diagnosing brain injury in a patient comprising the steps of (a) collecting a sample from the patient; (b) measuring the levels of a panel of citrullinated brain injury biomarker proteins in the sample collected from the patient using mass spectrometry, wherein the panel of biomarkers comprises NRGN, MBP, GFAP, PAD, isoforms thereof, post-translationally modified forms thereof, or combinations of any of the foregoing; and (c) comparing the levels of the panel of biomarkers with predefined levels of the same panel of biomarkers that correlate to a patient having brain injury and predefined levels of the same panel of biomarkers that correlate to a patient not having brain injury, wherein a correlation to one of the predefined levels provides the diagnosis.

[0030] The panel of biomarkers can further comprise one or more citrullinated brain injury biomarker proteins selected from the group consisting of tubulin beta-4B chain, tubulin alpha-1B chain, CNPase, PPIA, Septin-7, Elongation factor1-alpha2, TPPP, TPPP3, Ermin Isoform 2, NDRG2 Isoform 2, ASTN1, BAI3, CNDP1, ERMIN, GRM3, KLH32, MAGE2, NRG3, OMG, SLC39A12, RTN1, MT3, isoforms thereof, post-translationally modified forms thereof, or combinations of any of the foregoing.

[0031] In another embodiment, a method for determining brain injury status in a patient comprises the steps of (a) collecting a sample from the patient; (b) measuring the levels of a panel of citrullinated brain injury biomarker proteins in the sample collected from the patient using

SRM-MS, wherein the panel of biomarkers comprises NRGN, MBP, GFAP, PAD, isoforms thereof, post-translationally modified forms thereof, or combinations of any of the foregoing; and (c) comparing the levels of the panel of biomarkers with predefined levels of the same panel of biomarkers that correlate to one or more brain injury statuses selected from the group consisting of having brain injury, not having brain injury, progressing brain injury, and regressing brain injury, wherein a correlation to one of the predefined levels determines the brain injury status of the patient. In another embodiment, the panel of biomarkers further comprises one or more citrullinated brain injury biomarker proteins selected from the group consisting of tubulin beta-4B chain, tubulin alpha-1B chain, CNPase, PPIA, Septin-7, Elongation factor1-alpha2, TPPP, TPPP3, Ermin Isoform 2, NDRG2 Isoform 2, ASTN1, BAI3, CNDP1, ERMIN, GRM3, KLH32, MAGE2, NRG3, OMG, SLC39A12, RTN1, MT3, isoforms thereof, post-translationally modified forms thereof, or combinations of any of the foregoing.

[0032] The present invention also provides diagnostic kits for use in diagnosing brain injury in a patient. In one embodiment, a diagnostic kit for diagnosing brain injury in a patient comprises (a) a substrate for collecting a biological sample from the patient; and (b) means for measuring the levels of one or more human citrullinated brain injury biomarker proteins selected from the group consisting of NRGN, MBP, GFAP, PAD, isoforms thereof, post-translationally modified forms thereof, or combinations of any of the foregoing. In another embodiment, the one or more human citrullinated brain injury biomarker proteins further comprises tubulin beta-4B chain, tubulin alpha-1B chain, CNPase, PPIA, Septin-7, Elongation factor1-alpha2, TPPP, TPPP3, Ermin Isoform 2, NDRG2 Isoform 2, ASTN1, BAI3, CNDP1, ERMIN, GRM3, KLH32, MAGE2, NRG3, OMG, SLC39A12, RTN1, MT3, isoforms thereof, post-translationally modified forms thereof, or combinations of any of the foregoing.

[0033] In certain embodiments, the ratio of one or more citrullinated peptides to the corresponding unmodified peptides in GFAP is one or more of SEQ ID NO:28: SEQ ID NO:27; SEQ ID NO:30: SEQ ID NO:29; SEQ ID NO:40: SEQ ID NO:41; SEQ ID NO:42: SEQ ID NO:43; SEQ ID NO:44: SEQ ID NO:45; SEQ ID NO:46: SEQ ID NO:47; SEQ ID NO:48: SEQ ID NO:49; SEQ ID NO:50: SEQ ID NO:51; SEQ ID NO:52: SEQ ID NO:53; SEQ ID NO:54: SEQ ID NO:55; SEQ ID NO:56: SEQ ID NO:57; SEQ ID NO:91: SEQ ID NO:92; SEQ ID NO:93: SEQ ID NO:94; SEQ ID NO:95: SEQ ID NO:96; SEQ ID NO:97: SEQ ID NO:27; SEQ ID NO:98: SEQ ID NO:160; SEQ ID NO:118: SEQ ID NO:119; SEQ ID NO:120: SEQ ID NO:121; SEQ ID NO:122: SEQ ID NO:123; and SEQ ID NO:124: SEQ ID NO:125.

[0034] In other embodiments, the ratio of one or more citrullinated peptides to the corresponding unmodified peptides in MBP is one or more of SEQ ID NO:32: SEQ ID NO:31; SEQ ID NO:34: SEQ ID NO:33; SEQ ID NO:75: SEQ ID NO:74; SEQ ID NO:77: SEQ ID NO:76; SEQ ID NO:68: SEQ ID NO:69; SEQ ID NO:79: SEQ ID NO:78; SEQ ID NO:80: SEQ ID NO:78; SEQ ID NO:104: SEQ ID NO:105; SEQ ID NO:106: SEQ ID NO:107; SEQ ID NO:108: SEQ ID NO:109; SEQ ID NO:110: SEQ ID

NO:111; SEQ ID NO:112: SEQ ID NO:113; SEQ ID NO:114: SEQ ID NO:115; and SEQ ID NO:116: SEQ ID NO:117.

[0035] In further embodiments, the ratio of one or more citrullinated peptides to the corresponding unmodified peptides in NRGN is one or more of SEQ ID NO:35: SEQ ID NO:36; SEQ ID NO:37: SEQ ID NO:157; SEQ ID NO:71: SEQ ID NO:73; SEQ ID NO:77: SEQ ID NO:159; and SEQ ID NO:99: SEQ ID NO:100.

[0036] In additional embodiments, the ratio of one or more citrullinated peptides to the corresponding unmodified peptides in the proteins listed in Table 17 is one or more of SEQ ID NO:128: SEQ ID NO:129; SEQ ID NO:130: SEQ ID NO:131; SEQ ID NO:132: SEQ ID NO:133; SEQ ID NO:134: SEQ ID NO:135; SEQ ID NO:136: SEQ ID NO:137; SEQ ID NO:138: SEQ ID NO:139; SEQ ID NO:140: SEQ ID NO:141; SEQ ID NO:142: SEQ ID NO:143; SEQ ID NO:144: SEQ ID NO:145; SEQ ID NO:146: SEQ ID NO:147; SEQ ID NO:148: SEQ ID NO:149; SEQ ID NO:150: SEQ ID NO:151; SEQ ID NO:152: SEQ ID NO:153; and SEQ ID NO:154: SEQ ID NO:155.

[0037] Furthermore, the present invention provides for the combination of any of the foregoing ratios of citrullinated or other post-translationally modified peptides to the corresponding unmodified peptides provided herein including, but not limited to, SEQ ID NO:28: SEQ ID NO:27; SEQ ID NO:30: SEQ ID NO:29; SEQ ID NO:40: SEQ ID NO:41; SEQ ID NO:42: SEQ ID NO:43; SEQ ID NO:44: SEQ ID NO:45; SEQ ID NO:46: SEQ ID NO:47; SEQ ID NO:48: SEQ ID NO:49; SEQ ID NO:50: SEQ ID NO:51; SEQ ID NO:52: SEQ ID NO:53; SEQ ID NO:54: SEQ ID NO:55; SEQ ID NO:56: SEQ ID NO:57; SEQ ID NO:91: SEQ ID NO:92; SEQ ID NO:93: SEQ ID NO:94; SEQ ID NO:95: SEQ ID NO:96; SEQ ID NO:97: SEQ ID NO:27; SEQ ID NO:98: SEQ ID NO:160; SEQ ID NO:118: SEQ ID NO:119; SEQ ID NO:120: SEQ ID NO:121; SEQ ID NO:122: SEQ ID NO:123; SEQ ID NO:124: SEQ ID NO:125; SEQ ID NO:32: SEQ ID NO:31; SEQ ID NO:34: SEQ ID NO:33; SEQ ID NO:75: SEQ ID NO:74; SEQ ID NO:77: SEQ ID NO:76; SEQ ID NO:68: SEQ ID NO:69; SEQ ID NO:79: SEQ ID NO:78; SEQ ID NO:80: SEQ ID NO:78; SEQ ID NO:104: SEQ ID NO:105; SEQ ID NO:106: SEQ ID NO:107; SEQ ID NO:108: SEQ ID NO:109; SEQ ID NO:110: SEQ ID NO:111; SEQ ID NO:112: SEQ ID NO:113; SEQ ID NO:114: SEQ ID NO:115; SEQ ID NO:116: SEQ ID NO:117; SEQ ID NO:35: SEQ ID NO:36; SEQ ID NO:37: SEQ ID NO:157; SEQ ID NO:71: SEQ ID NO:73; SEQ ID NO:77: SEQ ID NO:159; SEQ ID NO:99: SEQ ID NO:100; SEQ ID NO:128: SEQ ID NO:129; SEQ ID NO:130: SEQ ID NO:131; SEQ ID NO:132: SEQ ID NO:133; SEQ ID NO:134: SEQ ID NO:135; SEQ ID NO:136: SEQ ID NO:137; SEQ ID NO:138: SEQ ID NO:139; SEQ ID NO:140: SEQ ID NO:141; SEQ ID NO:142: SEQ ID NO:143; SEQ ID NO:144: SEQ ID NO:145; SEQ ID NO:146: SEQ ID NO:147; SEQ ID NO:148: SEQ ID NO:149; SEQ ID NO:150: SEQ ID NO:151; SEQ ID NO:152: SEQ ID NO:153; and SEQ ID NO:154: SEQ ID NO:155.

[0038] In another aspect, the present invention provides methods for diagnosing brain injury by detecting autoantibodies to one or more citrullinated biomarker peptides described herein. In one embodiment, a method for diag-

nosing brain injury in a subject comprises the steps of (a) collecting a sample from the subject; (b) detecting the presence of autoantibodies to citrullinated brain injury biomarker peptides in the sample collected from the subject; and (c) correlating the amount of autoantibodies to citrullinated brain injury biomarker peptides to a patient having brain injury or to a patient not having brain injury, thereby providing the diagnosis. In another embodiment, the detecting step comprises the steps of (a) contacting a biological sample taken from a subject with a citrullinated brain injury biomarker peptide; and (b) detecting the binding of the peptide with an autoantibody specific for the peptide, wherein the detection of binding is indicative of the presence of citrullinated brain injury biomarker peptide autoantibodies in the subject. In a specific embodiment, the binding is detected by enzyme-linked immunosorbent assay (ELISA), immunoprecipitation or immunoblotting. In certain embodiments, the citrullinated brain injury biomarker peptide is one or more of NRGN, MBP, GFAP, PAD-2, tubulin beta-4B chain, tubulin alpha-1B chain, CNPase, PPIA, Septin-7, Elongation factor1-alpha2, TPPP, TPPP3, Ermin Isoform 2, and NDRG2 Isoform 2. In other embodiments, the citrullinated brain injury biomarker peptide is one or more of ASTN1, BAI3, CNDP1, ERMIN, GFAP, GRM3, KLH32, MAGE2, MBP, NRG3, NRGN, OMG, SLC39A12, RTN1, MT3, PAD.

[0039] In another embodiment, a method for assessing efficacy of a brain injury treatment regimen in a subject comprises the steps of (a) establishing a baseline level of autoantibodies to cit-NRGN, cit-MBP, and cit-GFAP in a subject prior to brain injury treatment regimen; (b) monitoring the levels of autoantibodies to cit-NRGN, cit-MBP, and cit-GFAP using a cit-NRGN, cit-MBP, and cit-GFAP peptide at least at one point after initiation of the brain injury treatment regimen; and (c) comparing the observed level of autoantibodies to cit-NRGN, cit-MBP, and cit-GFAP to the baseline level of autoantibodies to cit-NRGN, cit-MBP, and cit-GFAP, wherein a decrease in the level of autoantibodies is indicative of the efficacy of the brain injury treatment regimen. In a specific embodiment, the level of autoantibodies to cit-NRGN, cit-MBP, and cit-GFAP is measured by ELISA, immunoprecipitation or immunoblotting. In other embodiments, the baseline level of autoantibodies is also established with respect to one of more of citrullinated tubulin beta-4B chain, tubulin alpha-1B chain, CNPase, PPIA, Septin-7, Elongation factor1-alpha2, TPPP, TPPP3, Ermin Isoform 2, and NDRG2 Isoform 2. In other embodiments, the baseline level of autoantibodies is also established with respect to one or more of ASTN1, BAI3, CNDP1, ERMIN, GFAP, GRM3, KLH32, MAGE2, MBP, NRG3, NRGN, OMG, SLC39A12, RTN1, MT3, PAD.

[0040] In a further embodiment, a method for qualifying brain injury status in a subject comprises the steps of (a) measuring the level of autoantibodies to cit-NRGN, cit-MBP, and cit-GFAP in a biological sample from the subject; and (b) correlating the measurement with brain injury status. In a specific embodiment, the level of autoantibodies to cit-NRGN, cit-MBP, and cit-GFAP is measured by ELISA, immunoprecipitation or immunoblotting. In particular embodiments, the brain injury status is selected from the group consisting of the risk of brain injury, the development of brain injury, the presence or absence of brain injury, the

stage of brain injury, the subtype of brain injury, the prognosis for the subject, and the effectiveness of treatment of brain injury.

[0041] In another aspect, the present invention relates to methods for detecting the presence of autoantibodies to citrullinated peptides in a subject (cit-peptide autoantibodies) in a subject. In certain embodiments, the citrullinated peptide is MBP, GFAP, tubulin beta-4B chain, tubulin alpha-1B chain, CNPase, PPIA, Septin-7, Elongation factor1-alpha2, TPPP, TPPP3, Ermin Isoform 2, NDRG2 Isoform 2, ASTN1, BAI3, CNDP1, ERMIN, GFAP, GRM3, KLH32, MAGE2, NRG3, NRGN, OMG, SLC39A12, RTN1, MT3, and PAD2, as well as the peptides listed in Tables 1-11 and 13-17. The present invention also relates to methods for detecting the presence of autoantibodies to unmodified peptides.

[0042] In particular embodiment, the method comprises contacting a biological sample taken from a subject with a polypeptide of the present invention and detecting the binding of the polypeptide with an autoantibody specific for the polypeptide, wherein the detection of binding is indicative of the presence of cit-polypeptide autoantibodies in the subject. In such methods, the binding can be detected by enzyme-linked immunosorbent assay (ELISA), immunoprecipitation or immunoblotting.

[0043] In yet another aspect, the present invention relates to methods for assessing efficacy of a brain injury treatment regimen in a subject. In particular embodiments, the methods comprise establishing a baseline level of cit-peptide autoantibodies in a subject prior to a brain injury treatment regimen; monitoring the level of cit-peptide autoantibodies using a polypeptide of the present invention at least at one point after initiation of the brain injury treatment regimen; and comparing the observed level of cit-peptide autoantibodies to the baseline level of cit-peptide autoantibodies, wherein a decrease in the level of peptide autoantibodies is indicative of the efficacy of the brain injury treatment regimen. The peptide can be one or more of peptide is MBP, GFAP, PAD-2, tubulin beta-4B chain, tubulin alpha-1B chain, CNPase, PPIA, Septin-7, Elongation factor1-alpha2, TPPP, TPPP3, Ermin Isoform 2, NDRG2 Isoform 2, ASTN1, BAI3, CNDP1, ERMIN, GFAP, GRM3, KLH32, MAGE2, NRG3, NRGN, OMG, SLC39A12, RTN1, MT3, and PAD2. In such methods, the binding can be detected by enzyme-linked immunosorbent assay (ELISA), immunoprecipitation or immunoblotting.

[0044] In a further aspect, the present invention relates to methods for qualifying brain injury status in a subject. Qualifying brain injury status can be qualifying the risk of brain injury, the development of brain injury, the presence or absence of brain injury, the stage of brain injury, the subtype of brain injury, the prognosis for the subject, and the effectiveness of treatment for brain injury. In certain embodiments, the methods comprise measuring the level of cit-peptide autoantibodies in a biological sample from the subject; and correlating the measurement with brain injury status. In one embodiment, the level of cit-peptide autoantibodies is measured using a polypeptide described herein. In another embodiment, the level of cit-peptide autoantibodies is measured by ELISA, immunoprecipitation or immunoblotting. The peptide can be one or more of peptide is MBP, GFAP, PAD-2, tubulin beta-4B chain, tubulin alpha-1B chain, CNPase, PPIA, Septin-7, Elongation factor1-alpha2, TPPP, TPPP3, Ermin Isoform 2, NDRG2 Isoform 2,

ASTN1, BAI3, CNBP1, ERMIN, GFAP, GRM3, KLH32, MAGE2, NRG3, NRGN, OMG, SLC39A12, RTN1, MT3, and PAD2.

[0045] Finally, the present invention provides methods of detecting, measuring, determining, and the like, the biomarkers described herein in terms of both unmodified and modified forms, as well as autoantibodies thereto, i.e., combinations of both protein form and autoantibodies thereto.

BRIEF DESCRIPTION OF THE FIGURES

[0046] FIGS. 1A-1D present MS/MS spectra and extracted ion chromatograms of citrullinated peptides of endogenous bovine MBP digested with Lys-C. FIG. 1A: LTQ MS/MS spectrum of doubly charged species of NIVTPR*TPPPSQGK (residue 91-104) at m/z 746.91. FIG. 1B: extracted ion chromatogram of doubly charged species of NIVTPR*TPPPSQGK and NIVTPRTTPPPSQGK. *: Likely Q deamination occurred for the peptide NIVTPRTTPPPSQ GK. FIG. 1C: high resolution MS spectrum of double charged species of NIVTPRTTPPPSQ GK. FIG. 1D: high resolution MS spectrum of double charged species of NIVTPR*TPPPSQ GK.

[0047] FIGS. 2A and 2B show MS/MS spectra and extracted ion chromatograms of citrullinated peptide of endogenous bovine MBP digested with Lys-C. FIG. 2A: LTQ MS/MS spectrum of doubly charged species of PGFGYGGGR*ASDYK (residue 122-134) at m/z 688.32. FIG. 2B: extracted ion chromatogram of doubly charged species of PGFGYGGGR*ASDYK (residues 122-134) and PGFGYGGGRASDYK.

[0048] FIGS. 3A and 3B show MS/MS spectra and extracted ion chromatograms of citrullinated peptide of endogenous bovine MBP digested with Lys-C. FIG. 3A: Panel A: LTQ MS/MS spectrum of triply charged species of DGHHAAR*TTHYGSLPQK (residue 57-73) at m/z 626.31. FIG. 3B: Panel B: extracted ion chromatogram of triply charged species of DGHHAARTTTHYGSLPQK (residues 122-134) and DGHHAAR*TTHYGSLPQK.

[0049] FIG. 4 shows the amino acid sequence alignment for bovine NRGN and human NRGN. The underline indicates the peptides observed in tryptic digests. The initial Met (1) is removed when the protein is translated, but is included in the numbering of the peptides. Shaded area indicated sequence difference of bovine and human NRGN. R68 (in bold, larger font) is the citrullination site in endogenous NRGN.

[0050] FIGS. 5A and 5B present MS/MS spectra and extracted ion chromatograms of a citrullinated tryptic peptide of bovine neurogranin. FIG. 5A: MS/MS spectrum of doubly charged species of KGPGPGGPGGAGGAR*GGAGGGPSGD (residues 54-78) at m/z 953.44. FIG. 5B: extracted ion chromatogram of the ion m/z 953.44.

[0051] FIGS. 6A and 6B show MS/MS spectra and extracted ion chromatograms of citrullinated peptide of endogenous bovine GFAP digested with Glu-C. FIG. 6A: LTQ MS/MS spectrum of doubly charged species of GHLKR*NIVVKTVE (residues 398-410) at m/z 747.44. FIG. 6B: extracted ion chromatogram of doubly charged species of GHLKR*NIVVKTVE and GHLKRNIVVKTVE.

[0052] FIGS. 7A and 7B show extracted ion chromatograms of citrullinated peptide in an endogenous human GFAP sample. FIG. 7A: extracted ion chromatogram of

doubly charged species of GHLKR*NIVVKTVE (residues 402-414) and GHLKRNIVVKTVE of human GFAP. FIG. 7B: extracted ion chromatogram of doubly charged species of NIVTPR*TPPPSQGK (residues 92-105) and NIVTPRTTPPPSQGK of human MBP isoform 5.

[0053] FIG. 8 shows the amino acid sequence of recombinant human protein NRGN. The initial Met (1) is removed when the protein is translated but is included in the numbering of this protein. Shaded amino acids at N-terminus, T79, and R80 were introduced during subcloning. Shaded amino acids at N-terminus were not included in the numbering of this protein. The underline indicates the peptides identified in Lys-C digests and dotted line shows the peptide observed in Glu-C digests. Endogenous citrullination site was marked in italics. New citrullination sites observed after PAD2 treatment were in bold, larger font.

[0054] FIGS. 9A and 9B present MS/MS spectra and extracted ion chromatograms of a citrullinated peptide IQASFR*GHMAR*K (SEQ ID NO:35) of recombinant human protein NRGN after PAD2 treatment and digestion with Lys-C. FIG. 9A: LTQ MS/MS spectrum of doubly charged species of IQASFR*GHMAR*K (residues 33-44) (SEQ ID NO:35) at m/z 702.36. FIG. 9B: extracted ion chromatogram of doubly charged species of IQASFR*GHMAR*K (SEQ ID NO:35).

[0055] FIG. 10 shows the amino acid sequence of human GFAP and identified peptides with citrullination sites. The underline indicates the modified peptides in Lys-C digests. The dotted line indicates modified peptide in Glu-C digests. The initial Met (1) is removed when the protein is translated but is included in the numbering of the protein. Endogenous citrullination site was marked with an arrow and citrullination sites observed after PAD2 treatment were in italicized, bold, larger font.

[0056] FIG. 11 presents the aligned amino acid sequence of the isoform 5 (18.5 kDa, 170 residues) of human MBP (SEQ ID NO:162) and bovine MBP (169 amino acids) (SEQ ID NO:5). The initial Met (1) is removed in the numbering of both proteins. The underline indicated the peptide observed in Lys-C digests and dotted line showed the peptides observed in tryptic digests. Double-headed black arrows indicate citrullination sites previously reported in bovine MBP. Black arrows indicate citrullination sites previously reported in human MBP. Dotted arrows are previously identified citrullination sites in human MBP after PAD4 treatment. For all citrullinated residues identified in this study, endogenous citrullination sites were marked in grey shading and citrullination sites observed after PAD2 treatment were in italicized, bold larger font.

[0057] FIGS. 12A and 12B present a comparison of CID and HCD spectra of a citrullinated peptide of human GFAP in Glu-C digests. FIG. 12A: CID spectrum of the citrullinated peptide, GHLKR*NIVVKTVE of human GFAP at m/z 747.44, z : +2; FIG. 12B: HCD spectrum of this citrullinated peptide at m/z 747.44.

[0058] FIGS. 13A-13C present extracted ion chromatograms of the in vivo citrullinated peptide and intact peptide of NRGN in Lys-C digests of the AD3 brain sample. FIG. 13A: HPLC peak of the unmodified peptide at m/z 909.9275 with mass tolerance of 10 ppm. Right: isotopic clusters of the +2 ion. FIG. 13B: HPLC peak of the citrullinated peptide at m/z 910.4195 with mass tolerance of 10 ppm. Right: isotopic clusters of the +2 ion. FIG. 13C: Overlap of HPLC peaks shown in FIGS. 13A and 13B. From integrated peak

area, occupancy rate of citrullinated peptide at R68 residue was 8.6% for the AD3 sample.

[0059] FIG. 14: Citrullination occupancy rate for NRGN at R68, MBP at R92 and R124. Occupancy rate was calculated based on integrated peak area of citrullinated peptide vs. the sum of citrullinated and intact peptides.

DETAILED DESCRIPTION OF THE INVENTION

[0060] It is understood that the present invention is not limited to the particular methods and components, etc., described herein, as these may vary. It is also to be understood that the terminology used herein is used for the purpose of describing particular embodiments only, and is not intended to limit the scope of the present invention. It must be noted that as used herein and in the appended claims, the singular forms “a,” “an,” and “the” include the plural reference unless the context clearly dictates otherwise. Thus, for example, a reference to a “protein” is a reference to one or more proteins, and includes equivalents thereof known to those skilled in the art and so forth.

[0061] Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. Specific methods, devices, and materials are described, although any methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention.

[0062] All publications cited herein are hereby incorporated by reference including all journal articles, books, manuals, published patent applications, and issued patents. In addition, the meaning of certain terms and phrases employed in the specification, examples, and appended claims are provided. The definitions are not meant to be limiting in nature and serve to provide a clearer understanding of certain aspects of the present invention.

[0063] Brain and spinal cord (central nervous system, CNS) injury takes many forms, including hemorrhagic or ischemic stroke, hypoxic-ischemic encephalopathy, traumatic, mass effect compression from tumors or indolent as with degenerative brain diseases. Release or secretion of proteins from cells of the injured CNS can be useful for diagnostic/prognostic assessment of patient viability, recovery and the effects of therapy to stabilize or prevent new or recurrent CNS injury in children and adults. It can also reflect compromise of the brain blood barrier. Thus detection of circulating CNS proteins in body fluids, including the peripheral blood, saliva, urine and CSF, could improve the diagnostic accuracy of CNS injury by identifying children and adults with subclinical and overt CNS injury. This can provide insight into stroke, brain injury following surgery or with life support, following trauma as well as providing the means to determine and validate new and existing CNS injury treatments for efficacy to improve outcomes. In addition to brain specificity for a circulating protein to identify brain injury, are specific protein post-translational modifications of brain proteins to increase the diagnostic specificity of acute brain injury. The present inventors have discovered that certain brain proteins are post-translationally modified. Such proteins include neurogranin (gene symbol: NRGN, Uniprot accession: Q92686), myelin basic protein (gene symbol: MBP, Uniprot accession: P02686) and glial fibrillary acidic protein (gene symbol GFAP, Uniprot accession: P14136), tubulin beta-4B chain, tubulin alpha-1B

chain, CNPase, PPIA, Septin-7, Elongation factor1-alpha2, TPPP, TPPP3, Ermin Isoform 2, and NDRG2 Isoform 2. They discovered that NRGN and GFAP are endogenously citrullinated on specific arginines and citrullinated on additional arginines that were identified in vitro in proteins treated by the enzyme peptidyl arginine deiminase, PAD. PADs are calcium-activated enzymes that deaminate arginine residues when they are part of a protein creating a citrulline moiety in the place of the specific arginine residue. The ratio of citrullinated and unmodified forms of NRGN, MBP and GFAP (or fragments) can be quantitated multiple ways including antibody and aptamer based approaches. As well, mass spectrometry based methods can be used and the present inventors have developed specific quantitative multiple reaction monitoring assays for modified and unmodified peptides representing the total (unmodified) protein concentration, and each potential citrullinated residue. The value that quantitation of citrulline modification adds to detection of brain specific proteins is that it provides a window into the biology of the injured brain. The calcium burst necessary to increase PAD activity is a consequence of CNS cell injury. Quantifying the amount of the degree of citrullination provides insights into the scale, timing and recovery of injury that is critical to personalizing and developing new therapies appropriate to the phase of injury/recovery. In addition, citrullination of some proteins is known to increase their antigenicity, thus identification of citrullinated brain proteins may be very important for diagnosis of chronic neurodegenerative diseases such as Alzheimer's disease, multiple sclerosis and Parkinson's Disease. Detection and quantification of autoantibodies to these modified proteins could also be used to assess long term brain injury. Taken together, assays of post-translation modifications or the ratio of modified to unmodified at one or more specific arginine residues of circulating brain proteins will provide more accurate and specific diagnostic information for diagnosing brain injury.

I. Definitions

[0064] As used herein, the term “antibody” is used in reference to any immunoglobulin molecule that reacts with a specific antigen. It is intended that the term encompass any immunoglobulin (e.g., IgG, IgM, IgA, IgE, IgD, etc.) obtained from any source (e.g., humans, rodents, non-human primates, caprines, bovines, equines, ovines, etc.). Specific types/examples of antibodies include polyclonal, monoclonal, humanized, chimeric, human, or otherwise-human-suitable antibodies. “Antibodies” also includes any fragment or derivative of any of the herein described antibodies.

[0065] As used herein, the term “antigen” is generally used in reference to any substance that is capable of reacting with an antibody. It is intended that this term encompass any antigen and “immunogen” (i.e., a substance which induces the formation of antibodies). Thus, in an immunogenic reaction, antibodies are produced in response to the presence of an antigen (immunogen) or portion of an antigen. More specifically, the terms are used herein to describe an antigen that elicits a humoral and/or cellular immune response (i.e., is immunogenic), such that administration of the immunogen to an animal (e.g., via a vaccine) mounts an antigen-specific immune response against the same or similar antigens that are encountered within the tissues of the animal. In

another embodiment, when it is desirable to suppress an immune response against a given antigen, an antigen may comprise a toleragen.

[0066] As used herein, the term “autoantibodies” refers to antibodies that are capable of reacting against an antigenic constituent of an individual’s own tissue or cells (e.g., the antibodies recognize and bind to “self” antigens). In certain embodiments, the term “neurogranin autoantibodies” refers to antibodies produced by an individual that are immunospecific to the individual’s own neurogranin protein. In other embodiments, the term “citrullinated neurogranin autoantibodies” or “cit-neurogranin autoantibodies” refers to antibodies produced by an individual that are immunospecific to the individual’s own citrullinated neurogranin protein.

[0067] The term “brain injury” refers to a condition in which the brain is damaged by injury caused by an event. As used herein, an “injury” is an alteration in cellular or molecular integrity, activity, level, robustness, state, or other alteration that is traceable to an event. For example, an injury includes a physical, mechanical, chemical, biological, functional, infectious, or other modulator of cellular or molecular characteristics. An event can include a physical trauma such as a single or repetitive impact (percussive) or a biological abnormality such as a stroke resulting from either blockade or leakage of a blood vessel. An event is optionally an infection by an infectious agent. A person of skill in the art recognizes numerous equivalent events that are encompassed by the terms injury or event.

[0068] More specifically, the term “brain injury” refers to a condition that results in central nervous system damage, irrespective of its pathophysiological basis. Among the most frequent origins of a “brain injury” are stroke and traumatic brain injury (TBI). A “stroke” is classified into hemorrhagic and non-hemorrhagic. Examples of hemorrhagic stroke include cerebral hemorrhage, subarachnoid hemorrhage, and intracranial hemorrhage secondary to cerebral arterial malformation, while examples of non-hemorrhagic stroke include cerebral infarction.

[0069] The term “traumatic brain injury” or “TBI” refer to traumatic injuries to the brain which occur when physical trauma causes brain damage. For example, TBI can result from a closed head injury or a penetrating head injury. A “non-traumatic brain injury” refers to brain injuries that do not involve ischemia or external mechanical force (e.g., stroke, Alzheimer’s disease, Parkinson’s disease, Huntington’s disease, multiple sclerosis, amyotrophic lateral sclerosis, brain hemorrhage, brain infections, brain tumor, among others).

[0070] The term “brain injury” also refers to subclinical brain injury, spinal cord injury, and anoxic-ischemic brain injury. The term “subclinical brain injury” (SCI) refers to brain injury without overt clinical evidence of brain injury. A lack of clinical evidence of brain injury when brain injury actually exists could result from degree of injury, type of injury, level of consciousness, medications particularly sedation and anesthesia.

[0071] The “spinal cord injury” refers to a condition in which the spinal cord receives compression/detrition due to a vertebral fracture or dislocation to cause dysfunction. As used herein, the term “anoxic-ischemic brain injury” refers to deprivation of oxygen supply to brain tissue resulting in compromised brain function and includes cerebral hypoxia. For example, anoxic-ischemic brain injury includes focal cerebral ischemia, global cerebral ischemia, hypoxic

hypoxia (i.e., limited oxygen in the environment causes reduced brain function, such as with divers, aviators, mountain climbers, and fire fighters, all of whom are at risk for this kind of cerebral hypoxia), obstructions in the lungs (e.g., hypoxia resulting from choking, strangulation, the crushing of the windpipe).

[0072] The term “brain injury biomarker” (BIB), “brain injury biomarker protein”, “brain injury biomarker peptide”, “brain injury biomarker polypeptide” and the like refer to a protein, including those described herein, that can be used in a method of the present invention, e.g., to diagnose brain injury in a patient. Brain injury biomarker proteins include, but are not limited to, neurogranin (NRGN), glial fibrillary acidic protein (GFAP) and myelin basic protein (MBP). The term further includes, but is not limited to, PAD-2, tubulin beta-4B chain, tubulin alpha-1B chain, CNPase, PPIA, Septin-7, Elongation factor1-alpha2, TPPP, TPPP3, Ermin Isoform 2, NDRG2 Isoform 2, astrotactin 1 (ASTN1), brain angiogenesis inhibitor 3 (BAI3); carnosine dipeptidase 1 (CNDP1); ERMIN; glutamate receptor metabotropic 3 (GRM3); kelch-like protein 32 (KLH32); melanoma antigen family E,2 (MAGE2); neuregulin 3 (NRG3); oligodendrocyte myelin glycoprotein (OMG); solute carrier family 39 (zinc transporter); reticulon 1 (RTN1); and peptidylarginine deiminase (types 1-4 and 6) (PAD). The term also includes other brain injury biomarker proteins known in the art. In addition, the term “brain injury biomarkers” also includes the isoforms and/or post-translationally modified forms of any of the foregoing. In further embodiments, the term includes autoantibodies to the foregoing. The present invention contemplates the detection, measurement, quantification, determination and the like of both unmodified and modified (e.g., citrullination or other post-translational modification) proteins/polypeptides/peptides as well as autoantibodies to any of the foregoing. In certain embodiments, it is understood that reference to the detection, measurement, determination, and the like, of a biomarker refers detection of the protein/polypeptide/peptide (modified and/or unmodified). In other embodiments, reference to the detection, measurement, determination, and the like, of a biomarker refers detection of autoantibodies of the protein/polypeptide/peptide.

[0073] As used herein, the term “comparing” refers to making an assessment of how the proportion, level or cellular localization of one or more biomarkers in a sample from a patient relates to the proportion, level or cellular localization of the corresponding one or more biomarkers in a standard or control sample. For example, “comparing” may refer to assessing whether the proportion, level, or cellular localization of one or more biomarkers in a sample from a patient is the same as, more or less than, or different from the proportion, level, or cellular localization of the corresponding one or more biomarkers in standard or control sample. More specifically, the term may refer to assessing whether the proportion, level, or cellular localization of one or more biomarkers in a sample from a patient is the same as, more or less than, different from or otherwise corresponds (or not) to the proportion, level, or cellular localization of predefined biomarker levels/ratios that correspond to, for example, a patient having brain injury, not having brain injury, is responding to treatment for brain injury, is not responding to treatment for brain injury, is/is not likely to respond to a particular brain injury treatment, or having/not having another disease or condition. In a specific embodi-

ment, the term “comparing” refers to assessing whether the level of one or more biomarkers of the present invention in a sample from a patient is the same as, more or less than, different from other otherwise correspond (or not) to levels/ratios of the same biomarkers in a control sample (e.g., predefined levels/ratios that correlate to uninfected individuals, standard brain injury levels/ratios, etc.).

[0074] In another embodiment, the term “comparing” refers to making an assessment of how the proportion, level or cellular localization of one or more biomarkers in a sample from a patient relates to the proportion, level or cellular localization of another biomarker in the same sample. For example, a ratio of one biomarker to another from the same patient sample can be compared. In another embodiment, a level of one biomarker in a sample (e.g., a post-translationally modified biomarker protein) can be compared to the level of the same biomarker (e.g., unmodified biomarker protein) in the sample. In a specific embodiment, the proportion of a citrullinated biomarker protein can be compared to the unmodified protein, both of which are measured in the same patient sample. Ratios of modified: unmodified biomarker proteins can be compared to other protein ratios in the same sample or to predefined reference or control ratios.

[0075] As used herein, the terms “indicates” or “correlates” (or “indicating” or “correlating,” or “indication” or “correlation,” depending on the context) in reference to a parameter, e.g., a modulated proportion, level, or cellular localization in a sample from a patient, may mean that the patient has a brain injury or is suffering from neurodegeneration. In specific embodiments, the parameter may comprise the level of one or more biomarkers of the present invention. A particular set or pattern of the amounts of one or more biomarkers may indicate that a patient has a brain injury (i.e., correlates to a patient having brain injury). In other embodiments, a correlation could be the ratio of a post-translationally modified protein to the unmodified protein indicates (or a change in the ratio over time or as compared to a reference/control ratio) could mean that the patient has a brain injury). In specific embodiments, a correlation could be the ratio of a citrullinated peptide to the non-citrullinated form, or any other combination in which a change in one peptide causes or is accompanied by a change in another.

[0076] In other embodiments, a particular set or pattern of the amounts of one or more biomarkers may be correlated to a patient being unaffected (i.e., indicates a patient does not have brain injury). In certain embodiments, “indicating,” or “correlating,” as used according to the present invention, may be by any linear or non-linear method of quantifying the relationship between levels/ratios of biomarkers to a standard, control or comparative value for the assessment of the diagnosis, prediction of brain injury or brain injury progression, assessment of efficacy of clinical treatment, identification of a patient that may respond to a particular treatment regime or pharmaceutical agent, monitoring of the progress of treatment, and in the context of a screening assay, for the identification of an anti-brain injury therapeutic.

[0077] The terms “patient,” “individual,” or “subject” are used interchangeably herein, and refer to a mammal, particularly, a human. The patient may have mild, intermediate or severe disease. The patient may be treatment naïve, responding to any form of treatment, or refractory. The patient may be an individual in need of treatment or in need

of diagnosis based on particular symptoms or family history. In some cases, the terms may refer to treatment in experimental animals, in veterinary application, and in the development of animal models for disease, including, but not limited to, rodents including mice, rats, and hamsters; and primates.

[0078] The terms “measuring” and “determining” are used interchangeably throughout, and refer to methods which include obtaining a patient sample and/or detecting the level of a biomarker(s) in a sample. In one embodiment, the terms refer to obtaining a patient sample and detecting the level of one or more biomarkers in the sample. In another embodiment, the terms “measuring” and “determining” mean detecting the level of one or more biomarkers in a patient sample. Measuring can be accomplished by methods known in the art and those further described herein. The term “measuring” is also used interchangeably throughout with the term “detecting.”

[0079] The terms “sample,” “patient sample,” “biological sample,” and the like, encompass a variety of sample types obtained from a patient, individual, or subject and can be used in a diagnostic or monitoring assay. The patient sample may be obtained from a healthy subject, a diseased patient or a patient having associated symptoms of brain injury. Moreover, a sample obtained from a patient can be divided and only a portion may be used for diagnosis. Further, the sample, or a portion thereof, can be stored under conditions to maintain sample for later analysis. The definition specifically encompasses blood and other liquid samples of biological origin (including, but not limited to, peripheral blood, serum, plasma, cerebrospinal fluid, urine, saliva, stool and synovial fluid), solid tissue samples such as a biopsy specimen or tissue cultures or cells derived therefrom and the progeny thereof. In certain embodiment, a sample comprises cerebrospinal fluid. In a specific embodiment, a sample comprises a blood sample. In another embodiment, a sample comprises a plasma sample. In yet another embodiment, a serum sample is used.

[0080] The definition of “sample” also includes samples that have been manipulated in any way after their procurement, such as by centrifugation, filtration, precipitation, dialysis, chromatography, treatment with reagents, washed, or enriched for certain cell populations. The terms further encompass a clinical sample, and also include cells in culture, cell supernatants, tissue samples, organs, and the like. Samples may also comprise fresh-frozen and/or formalin-fixed, paraffin-embedded tissue blocks, such as blocks prepared from clinical or pathological biopsies, prepared for pathological analysis or study by immunohistochemistry.

[0081] The terms “specifically binds to,” “specific for,” and related grammatical variants refer to that binding which occurs between such paired species as enzyme/substrate, receptor/agonist, antibody/antigen, and lectin/carbohydrate which may be mediated by covalent or non-covalent interactions or a combination of covalent and non-covalent interactions. When the interaction of the two species produces a non-covalently bound complex, the binding which occurs is typically electrostatic, hydrogen-bonding, or the result of lipophilic interactions. Accordingly, “specific binding” occurs between a paired species where there is interaction between the two which produces a bound complex having the characteristics of an antibody/antigen or enzyme/substrate interaction. In particular, the specific binding is characterized by the binding of one member of a pair to a

particular species and to no other species within the family of compounds to which the corresponding member of the binding member belongs. Thus, for example, an antibody typically binds to a single epitope and to no other epitope within the family of proteins. In some embodiments, specific binding between an antigen and an antibody will have a binding affinity of at least 10^{-6} M. In other embodiments, the antigen and antibody will bind with affinities of at least 10^{-7} M, 10^{-8} M to 10^{-9} M, 10^{-10} M, 10^{-11} M, or 10^{-12} M.

[0082] Various methodologies of the instant invention include a step that involves comparing a value, level, feature, characteristic, property, etc. to a “suitable control,” referred to interchangeably herein as an “appropriate control” or a “control sample.” A “suitable control,” “appropriate control” or a “control sample” is any control or standard familiar to one of ordinary skill in the art useful for comparison purposes. In one embodiment, a “suitable control” or “appropriate control” is a value, level, feature, characteristic, property, etc., determined in a cell, organ, or patient, e.g., a control or normal cell, organ, or patient, exhibiting, for example, normal traits. For example, the biomarkers of the present invention may be assayed for levels/ratios in a sample from an unaffected individual (UI) or a normal control individual (NC) (both terms are used interchangeably herein). In another embodiment, a “suitable control” or “appropriate control” is a value, level, feature, characteristic, property, ratio, etc. determined prior to performing a therapy (e.g., a brain injury treatment) on a patient. In yet another embodiment, a transcription rate, mRNA level, translation rate, protein level/ratio, biological activity, cellular characteristic or property, genotype, phenotype, etc., can be determined prior to, during, or after administering a therapy into a cell, organ, or patient. In a further embodiment, a “suitable control” or “appropriate control” is a predefined value, level, feature, characteristic, property, ratio, etc. A “suitable control” can be a profile or pattern of levels/ratios of one or more biomarkers of the present invention that correlates to brain injury, to which a patient sample can be compared. The patient sample can also be compared to a negative control, i.e., a profile that correlates to not having brain injury.

II. Detection of Brain Injury or Neurodegeneration Biomarkers

[0083] A. Detection by Mass Spectrometry

[0084] In one aspect, the biomarkers of the present invention may be detected by mass spectrometry, a method that employs a mass spectrometer to detect gas phase ions. Examples of mass spectrometers are time-of-flight, magnetic sector, quadrupole filter, ion trap, ion cyclotron resonance, Orbitrap, hybrids or combinations of the foregoing, and the like.

[0085] In particular embodiments, the biomarkers of the present invention are detected using selected reaction monitoring (SRM) mass spectrometry techniques. Selected reaction monitoring (SRM) is a non-scanning mass spectrometry technique, performed on triple quadrupole-like instruments and in which collision-induced dissociation is used as a means to increase selectivity. In SRM experiments two mass analyzers are used as static mass filters, to monitor a particular fragment ion of a selected precursor ion. The specific pair of mass-over-charge (m/z) values associated to the precursor and fragment ions selected is referred to as a “transition” and can be written as parent m/z →fragment m/z (e.g. 673.5→534.3). Unlike common MS based proteomics,

no mass spectra are recorded in a SRM analysis. Instead, the detector acts as counting device for the ions matching the selected transition thereby returning an intensity distribution over time. Multiple SRM transitions can be measured within the same experiment on the chromatographic time scale by rapidly toggling between the different precursor/fragment pairs (sometimes called multiple reaction monitoring, MRM). Typically, the triple quadrupole instrument cycles through a series of transitions and records the signal of each transition as a function of the elution time. The method allows for additional selectivity by monitoring the chromatographic coelution of multiple transitions for a given analyte. The terms SRM/MRM are occasionally used also to describe experiments conducted in mass spectrometers other than triple quadrupoles (e.g. in trapping instruments) where upon fragmentation of a specific precursor ion a narrow mass range is scanned in MS2 mode, centered on a fragment ion specific to the precursor of interest or in general in experiments where fragmentation in the collision cell is used as a means to increase selectivity. In this application the terms SRM and MRM or also SRM/MRM can be used interchangeably, since they both refer to the same mass spectrometer operating principle. As a matter of clarity, the term MRM is used throughout the text, but the term includes both SRM and MRM, as well as any analogous technique, such as e.g. highly-selective reaction monitoring, hSRM, LC-SRM or any other SRM/MRM-like or SRM/MRM-mimicking approaches performed on any type of mass spectrometer and/or, in which the peptides are fragmented using any other fragmentation method such as e.g. CAD (collision-activated dissociation (also known as CID or collision-induced dissociation), HCD (higher energy CID), ECD (electron capture dissociation), PD (photodissociation) or ETD (electron transfer dissociation).

[0086] In another specific embodiment, the mass spectrometric method comprises matrix assisted laser desorption/ionization time-of-flight (MALDI-TOF MS or MALDI-TOF). In another embodiment, method comprises MALDI-TOF tandem mass spectrometry (MALDI-TOF MS/MS). In yet another embodiment, mass spectrometry can be combined with another appropriate method(s) as may be contemplated by one of ordinary skill in the art. For example, MALDI-TOF can be utilized with trypsin digestion and tandem mass spectrometry as described herein.

[0087] In an alternative embodiment, the mass spectrometric technique comprises surface enhanced laser desorption and ionization or “SELDI,” as described, for example, in U.S. Pat. Nos. 6,225,047 and 5,719,060. Briefly, SELDI refers to a method of desorption/ionization gas phase ion spectrometry (e.g. mass spectrometry) in which an analyte (here, one or more of the biomarkers) is captured on the surface of a SELDI mass spectrometry probe. There are several versions of SELDI that may be utilized including, but not limited to, Affinity Capture Mass Spectrometry (also called Surface-Enhanced Affinity Capture (SEAC)), and Surface-Enhanced Neat Desorption (SEND) which involves the use of probes comprising energy absorbing molecules that are chemically bound to the probe surface (SEND probe). Another SELDI method is called Surface-Enhanced Photolabile Attachment and Release (SEPAR), which involves the use of probes having moieties attached to the surface that can covalently bind an analyte, and then release the analyte through breaking a photolabile bond in the moiety after exposure to light, e.g., to laser light (see, U.S.

Pat. No. 5,719,060). SEPAR and other forms of SELDI are readily adapted to detecting a biomarker or biomarker panel, pursuant to the present invention.

[0088] In another mass spectrometry method, the biomarkers can be first captured on a chromatographic resin having chromatographic properties that bind the biomarkers. For example, one could capture the biomarkers on a cation exchange resin, such as CM Ceramic HyperD F resin, wash the resin, elute the biomarkers and detect by MALDI. Alternatively, this method could be preceded by fractionating the sample on an anion exchange resin before application to the cation exchange resin. In another alternative, one could fractionate on an anion exchange resin and detect by MALDI directly. In yet another method, one could capture the biomarkers on an immuno-chromatographic resin that comprises antibodies that bind the biomarkers, wash the resin to remove unbound material, elute the biomarkers from the resin and detect the eluted biomarkers by MALDI or by SELDI.

[0089] B. Detection by Immunoassay

[0090] In other embodiments, the biomarkers of the present invention can be detected and/or measured by immunoassay. Immunoassay requires biospecific capture reagents, such as antibodies, to capture the biomarkers. Many antibodies are available commercially. Antibodies also can be produced by methods well known in the art, e.g., by immunizing animals with the biomarkers. Biomarkers can be isolated from samples based on their binding characteristics. Alternatively, if the amino acid sequence of a polypeptide biomarker is known, the polypeptide can be synthesized and used to generate antibodies by methods well-known in the art.

[0091] The present invention contemplates traditional immunoassays including, for example, sandwich immunoassays including ELISA or fluorescence-based immunoassays, immunoblots, Western Blots (WB), as well as other enzyme immunoassays. Nephelometry is an assay performed in liquid phase, in which antibodies are in solution. Binding of the antigen to the antibody results in changes in absorbance, which is measured. In a SELDI-based immunoassay, a biospecific capture reagent for the biomarker is attached to the surface of an MS probe, such as a pre-activated protein chip array. The biomarker is then specifically captured on the biochip through this reagent, and the captured biomarker is detected by mass spectrometry.

[0092] Although antibodies are useful because of their extensive characterization, any other suitable agent (e.g., a peptide, an aptamer, or a small organic molecule) that specifically binds a biomarker of the present invention is optionally used in place of the antibody in the above described immunoassays. For example, an aptamer that specifically binds all neurogranin and/or one or more of its breakdown products might be used. Aptamers are nucleic acid-based molecules that bind specific ligands. Methods for making aptamers with a particular binding specificity are known as detailed in U.S. Pat. Nos. 5,475,096; 5,670,637; 5,696,249; 5,270,163; 5,707,796; 5,595,877; 5,660,985; 5,567,588; 5,683,867; 5,637,459; and 6,011,020.

[0093] C. Detection by Electrochemiluminescent Assay

[0094] In several embodiments, the biomarker biomarkers of the present invention may be detected by means of an electrochemiluminescent assay developed by Meso Scale Discovery (Gaithersburg, Md.). Electrochemiluminescence detection uses labels that emit light when electrochemically

stimulated. Background signals are minimal because the stimulation mechanism (electricity) is decoupled from the signal (light). Labels are stable, non-radioactive and offer a choice of convenient coupling chemistries. They emit light at ~620 nm, eliminating problems with color quenching. See U.S. Pat. Nos. 7,497,997; 7,491,540; 7,288,410; 7,036,946; 7,052,861; 6,977,722; 6,919,173; 6,673,533; 6,413,783; 6,362,011; 6,319,670; 6,207,369; 6,140,045; 6,090,545; and 5,866,434. See also U.S. Patent Applications Publication 2009/0170121; No. 2009/006339; No. 2009/0065357; No. 2006/0172340; No. 2006/0019319; No. 2005/0142033; No. 2005/0052646; No. 2004/0022677; No. 2003/0124572; No. 2003/0113713; No. 2003/0003460; No. 2002/0137234; No. 2002/0086335; and No. 2001/0021534.

[0095] D. Other Methods for Detecting Biomarkers

[0096] The biomarkers of the present invention can be detected by other suitable methods. Detection paradigms that can be employed to this end include optical methods, electrochemical methods (voltametry and amperometry techniques), atomic force microscopy, and radio frequency methods, e.g., multipolar resonance spectroscopy. Illustrative of optical methods, in addition to microscopy, both confocal and non-confocal, are detection of fluorescence, luminescence, chemiluminescence, absorbance, reflectance, transmittance, and birefringence or refractive index (e.g., surface plasmon resonance, ellipsometry, a resonant mirror method, a grating coupler waveguide method or interferometry).

[0097] Furthermore, a sample may also be analyzed by means of a biochip. Biochips generally comprise solid substrates and have a generally planar surface, to which a capture reagent (also called an adsorbent or affinity reagent) is attached. Frequently, the surface of a biochip comprises a plurality of addressable locations, each of which has the capture reagent bound there. Protein biochips are biochips adapted for the capture of polypeptides. Many protein biochips are described in the art. These include, for example, protein biochips produced by CIPHERGEN Biosystems, Inc. (Fremont, Calif.), Invitrogen Corp. (Carlsbad, Calif.), Affymetrix, Inc. (Fremont, Calif.), Zyomyx (Hayward, Calif.), R&D Systems, Inc. (Minneapolis, Minn.), Biacore (Uppsala, Sweden) and Procognia (Berkshire, UK). Examples of such protein biochips are described in the following patents or published patent applications: U.S. Pat. Nos. 6,537,749; 6,329,209; 6,225,047; 5,242,828; PCT International Publication No. WO 00/56934; and PCT International Publication No. WO 03/048768.

III. Determination of a Patient's Brain Injury Status

[0098] A. The present invention relates to the use of biomarkers to diagnose brain injury or neurodegeneration. It is understood that, for the sake of brevity, the term "brain injury" is used throughout the specification, but it is understood that the methods and biomarkers described herein are applicable in the context of diagnosing neurodegeneration. More specifically, the biomarkers of the present invention can be used in diagnostic tests to determine, qualify, and/or assess brain injury or status, for example, to diagnose brain injury, in an individual, subject or patient. In particular embodiments, brain injury status can include determining a patient's brain injury status or brain injury status, for example, to diagnose brain injury, in an individual, subject or patient. More specifically, the biomarkers to be detected in diagnosing brain injury (e.g., subclinical brain injury)

include, but are not limited to, MBP, GFAP, tubulin beta-4B chain, tubulin alpha-1B chain, CNPase, PPIA, Septin-7, Elongation factor1-alpha2, TPPP, TPPP3, Ermin Isoform 2, NDRG2 Isoform 2, astrotactin 1 (ASTN1), brain angiogenesis inhibitor 3 (BAI3); carnosine dipeptidase 1 (CNDP1); ERMIN; glial fibrillary acidic protein (GFAP); glutamate receptor metabotropic 3 (GRM3); kelch-like protein 32 (KLH32); melanoma antigen family E,2 (MAGE2); neuregulin 3 (NRG3); neurogranin (NRGN); oligodendrocyte myelin glycoprotein (OMG); solute carrier family 39 (zinc transporter), member 12 (SLC39A12); reticulon 1 (RTN1); metallothionein (MT3), and peptidylarginine deiminase type-2 (PAD2). Tables 1-11 and 13-17 also list biomarkers (i.e., peptides) useful in the methods of the present invention. Other biomarkers known in the relevant art may be used in combination with the biomarkers described herein. The present invention further contemplates the detection, measurement, quantification, determination and the like of both unmodified and modified (e.g., citrullination or other post-translational modification) proteins/polypeptides/peptides as well as autoantibodies to any of the foregoing, determining a patient's brain injury status.

[0099] B. Biomarker Panels

[0100] The biomarkers of the present invention can be used in diagnostic tests to assess, determine, and/or qualify (used interchangeably herein) brain injury status in a patient. The phrase "brain injury status" includes any distinguishable manifestation of the condition, including not having brain injury. For example, brain injury status includes, without limitation, the presence or absence of brain injury in a patient, the risk of developing brain injury, the stage or severity of brain injury, the progress of brain injury (e.g., progress of brain injury over time) and the effectiveness or response to treatment of brain injury (e.g., clinical follow up and surveillance of brain injury after treatment). Based on this status, further procedures may be indicated, including additional diagnostic tests or therapeutic procedures or regimens.

[0101] The power of a diagnostic test to correctly predict status is commonly measured as the sensitivity of the assay, the specificity of the assay or the area under a receiver operated characteristic ("ROC") curve. Sensitivity is the percentage of true positives that are predicted by a test to be positive, while specificity is the percentage of true negatives that are predicted by a test to be negative. An ROC curve provides the sensitivity of a test as a function of 1-specificity. The greater the area under the ROC curve, the more powerful the predictive value of the test. Other useful measures of the utility of a test are positive predictive value and negative predictive value. Positive predictive value is the percentage of people who test positive that are actually positive. Negative predictive value is the percentage of people who test negative that are actually negative.

[0102] In particular embodiments, the biomarker panels of the present invention may show a statistical difference in different brain injury statuses of at least $p < 0.05$, $p < 10^{-1}$, $p < 10^{-3}$, $p < 10^{-4}$ or $p < 10^{-5}$. Diagnostic tests that use these biomarkers may show an ROC of at least 0.6, at least about 0.7, at least about 0.8, or at least about 0.9.

[0103] The biomarkers can be differentially present in UI (NC or non-brain injury) and brain injury, and, therefore, are useful in aiding in the determination of brain injury status. In certain embodiments, the biomarkers are measured in a patient sample using the methods described herein and

compared, for example, to predefined biomarker levels/ratios and correlated to brain injury status. In particular embodiments, the measurement(s) may then be compared with a relevant diagnostic amount(s), cut-off(s), or multivariate model scores that distinguish a positive brain injury status from a negative brain injury status. The diagnostic amount(s) represents a measured amount of a biomarker(s) above which or below which a patient is classified as having a particular brain injury status. For example, if the biomarker(s) is/are up-regulated compared to normal during brain injury, then a measured amount(s) above the diagnostic cutoff(s) provides a diagnosis of brain injury. Alternatively, if the biomarker(s) is/are down-regulated during brain injury, then a measured amount(s) at or below the diagnostic cutoff(s) provides a diagnosis of non-brain injury. As is well understood in the art, by adjusting the particular diagnostic cut-off(s) used in an assay, one can increase sensitivity or specificity of the diagnostic assay depending on the preference of the diagnostician. In particular embodiments, the particular diagnostic cut-off can be determined, for example, by measuring the amount of biomarkers in a statistically significant number of samples from patients with the different brain injury statuses, and drawing the cut-off to suit the desired levels of specificity and sensitivity.

[0104] In other embodiments, ratios of post-translationally modified biomarkers to the corresponding unmodified biomarkers are useful in aiding in the determination of brain injury status. In certain embodiments, the biomarker ratios are indicative of diagnosis. In other embodiments, a biomarker ratio can be compared to another biomarker ratio in the same sample or to a set of biomarker ratios from a control or reference sample.

[0105] Indeed, as the skilled artisan will appreciate there are many ways to use the measurements of two or more biomarkers in order to improve the diagnostic question under investigation. In a quite simple, but nonetheless often effective approach, a positive result is assumed if a sample is positive for at least one of the markers investigated.

[0106] Furthermore, in certain embodiments, the values measured for markers of a biomarker panel are mathematically combined and the combined value is correlated to the underlying diagnostic question. Biomarker values may be combined by any appropriate state of the art mathematical method. Well-known mathematical methods for correlating a marker combination to a disease status employ methods like discriminant analysis (DA) (e.g., linear-, quadratic-, regularized-DA), Discriminant Functional Analysis (DFA), Kernel Methods (e.g., SVM), Multidimensional Scaling (MDS), Nonparametric Methods (e.g., k-Nearest-Neighbor Classifiers), PLS (Partial Least Squares), Tree-Based Methods (e.g., Logic Regression, CART, Random Forest Methods, Boosting/Bagging Methods), Generalized Linear Models (e.g., Logistic Regression), Principal Components based Methods (e.g., SIMCA), Generalized Additive Models, Fuzzy Logic based Methods, Neural Networks and Genetic Algorithms based Methods. The skilled artisan will have no problem in selecting an appropriate method to evaluate a biomarker combination of the present invention. In one embodiment, the method used in a correlating a biomarker combination of the present invention, e.g. to diagnose brain injury, is selected from DA (e.g., Linear-, Quadratic-, Regularized Discriminant Analysis), DFA, Kernel Methods (e.g., SVM), MDS, Nonparametric Methods (e.g., k-Nearest-Neighbor Classifiers), PLS (Partial Least Squares), Tree-

Based Methods (e.g., Logic Regression, CART, Random Forest Methods, Boosting Methods), or Generalized Linear Models (e.g., Logistic Regression), and Principal Components Analysis. Details relating to these statistical methods are found in the following references: Ruczinski et al., 12 J. OF COMPUTATIONAL AND GRAPHICAL STATISTICS 475-511 (2003); Friedman, J. H., 84 J. OF THE AMERICAN STATISTICAL ASSOCIATION 165-75 (1989); Hastie, Trevor, Tibshirani, Robert, Friedman, Jerome, The Elements of Statistical Learning, Springer Series in Statistics (2001); Breiman, L., Friedman, J. H., Olshen, R. A., Stone, C. J. Classification and regression trees, California: Wadsworth (1984); Breiman, L., 45 MACHINE LEARNING 5-32 (2001); Pepe, M. S., The Statistical Evaluation of Medical Tests for Classification and Prediction, Oxford Statistical Science Series, 28 (2003); and Duda, R. O., Hart, P. E., Stork, D. G., Pattern Classification, Wiley Interscience, 2nd Edition (2001).

[0107] C. Determining Risk of Developing Brain Injury

[0108] In a specific embodiment, the present invention provides methods for determining the risk of developing brain injury in a patient. Biomarker percentages, ratios, amounts or patterns are characteristic of various risk states, e.g., high, medium or low. The risk of developing brain injury is determined by measuring the relevant biomarkers and then either submitting them to a classification algorithm or comparing them with a reference amount, i.e., a pre-defined level or pattern of biomarkers that is associated with the particular risk level.

[0109] D. Determining Brain Injury Severity

[0110] In another embodiment, the present invention provides methods for determining the severity of brain injury in a patient. Each grade or stage of brain injury likely has a characteristic level of a biomarker or relative levels/ratios of a set of biomarkers (a pattern or ratio). The severity of brain injury is determined by measuring the relevant biomarkers and then either submitting them to a classification algorithm or comparing them with a reference amount, i.e., a pre-defined level or pattern of biomarkers that is associated with the particular stage.

[0111] E. Determining Brain Injury Prognosis

[0112] In one embodiment, the present invention provides methods for determining the course of brain injury in a patient. Brain injury course refers to changes in brain injury status over time, including brain injury progression (worsening) and brain injury regression (improvement). Over time, the amount or relative amount (e.g., the pattern or ratio) of the biomarkers changes. For example, biomarker "X" may be increased with brain injury, while biomarker "Y" may be decreased with brain injury. Therefore, the trend of these biomarkers, either increased or decreased over time toward brain injury or non-brain injury indicates the course of the condition. Accordingly, this method involves measuring the level of one or more biomarkers in a patient at least two different time points, e.g., a first time and a second time, and comparing the change, if any. The course of brain injury is determined based on these comparisons.

[0113] F. Patient Management

[0114] In certain embodiments of the methods of qualifying brain injury status, the methods further comprise managing patient treatment based on the status. Such management includes the actions of the physician or clinician subsequent to determining brain injury status. For example, if a physician makes a diagnosis of brain injury, then a certain regime of monitoring would follow. An assessment

of the course of brain injury using the methods of the present invention may then require a certain brain injury therapy regimen. Alternatively, a diagnosis of non-brain injury might be followed with further testing to determine a specific disease that the patient might be suffering from. Also, further tests may be called for if the diagnostic test gives an inconclusive result on brain injury status.

[0115] G. Determining Therapeutic Efficacy of Pharmaceutical Drug

[0116] In another embodiment, the present invention provides methods for determining the therapeutic efficacy of a pharmaceutical drug. These methods are useful in performing clinical trials of the drug, as well as monitoring the progress of a patient on the drug. Therapy or clinical trials involve administering the drug in a particular regimen. The regimen may involve a single dose of the drug or multiple doses of the drug over time. The doctor or clinical researcher monitors the effect of the drug on the patient or subject over the course of administration. If the drug has a pharmacological impact on the condition, the amounts or relative amounts (e.g., the pattern, profile or ratio) of one or more of the biomarkers of the present invention may change toward a non-brain injury profile. Therefore, one can follow the course of one or more biomarkers in the patient during the course of treatment. Accordingly, this method involves measuring one or more biomarkers in a patient receiving drug therapy, and correlating the biomarker levels/ratios with the brain injury status of the patient (e.g., by comparison to predefined levels/ratios of the biomarkers that correspond to different brain injury statuses). One embodiment of this method involves determining the levels/ratios of one or more biomarkers for at least two different time points during a course of drug therapy, e.g., a first time and a second time, and comparing the change in levels/ratios of the biomarkers, if any. For example, the levels/ratios of one or more biomarkers can be measured before and after drug administration or at two different time points during drug administration. The effect of therapy is determined based on these comparisons. If a treatment is effective, then the level/ratio of one or more biomarkers will trend toward normal, while if treatment is ineffective, the level/ratio of one or more biomarkers will trend toward brain injury indications.

[0117] H. Generation of Classification Algorithms for Qualifying Brain Injury Status

[0118] In some embodiments, data that are generated using samples such as "known samples" can then be used to "train" a classification model. A "known sample" is a sample that has been pre-classified. The data that are used to form the classification model can be referred to as a "training data set." The training data set that is used to form the classification model may comprise raw data or pre-processed data. Once trained, the classification model can recognize patterns in data generated using unknown samples. The classification model can then be used to classify the unknown samples into classes. This can be useful, for example, in predicting whether or not a particular biological sample is associated with a certain biological condition (e.g., diseased versus non-diseased).

[0119] Classification models can be formed using any suitable statistical classification or learning method that attempts to segregate bodies of data into classes based on objective parameters present in the data. Classification methods may be either supervised or unsupervised. Examples of supervised and unsupervised classification pro-

cesses are described in Jain, "Statistical Pattern Recognition: A Review", IEEE Transactions on Pattern Analysis and Machine Intelligence, Vol. 22, No. 1, January 2000, the teachings of which are incorporated by reference.

[0120] In supervised classification, training data containing examples of known categories are presented to a learning mechanism, which learns one or more sets of relationships that define each of the known classes. New data may then be applied to the learning mechanism, which then classifies the new data using the learned relationships. Examples of supervised classification processes include linear regression processes (e.g., multiple linear regression (MLR), partial least squares (PLS) regression and principal components regression (PCR)), binary decision trees (e.g., recursive partitioning processes such as CART), artificial neural networks such as back propagation networks, discriminant analyses (e.g., Bayesian classifier or Fischer analysis), logistic classifiers, and support vector classifiers (support vector machines).

[0121] Another supervised classification method is a recursive partitioning process. Recursive partitioning processes use recursive partitioning trees to classify data derived from unknown samples. Further details about recursive partitioning processes are provided in U.S. Patent Application No. 2002 0138208 A1 to Paulse et al., "Method for analyzing mass spectra."

[0122] In other embodiments, the classification models that are created can be formed using unsupervised learning methods. Unsupervised classification attempts to learn classifications based on similarities in the training data set, without pre-classifying the spectra from which the training data set was derived. Unsupervised learning methods include cluster analyses. A cluster analysis attempts to divide the data into "clusters" or groups that ideally should have members that are very similar to each other, and very dissimilar to members of other clusters. Similarity is then measured using some distance metric, which measures the distance between data items, and clusters together data items that are closer to each other. Clustering techniques include the MacQueen's K-means algorithm and the Kohonen's Self-Organizing Map algorithm.

[0123] Learning algorithms asserted for use in classifying biological information are described, for example, in PCT International Publication No. WO 01/31580 (Barnhill et al., "Methods and devices for identifying patterns in biological systems and methods of use thereof"), U.S. Patent Application Publication No. 2002/0193950 (Gavin et al. "Method or analyzing mass spectra"), U.S. Patent Application Publication No. 2003/0004402 (Hitt et al., "Process for discriminating between biological states based on hidden patterns from biological data"), and U.S. Patent Application Publication No. 2003/0055615 (Zhang and Zhang, "Systems and methods for processing biological expression data").

[0124] The classification models can be formed on and used on any suitable digital computer. Suitable digital computers include micro, mini, or large computers using any standard or specialized operating system, such as a Unix, Windows® or Linux™ based operating system. In embodiments utilizing a mass spectrometer, the digital computer that is used may be physically separate from the mass spectrometer that is used to create the spectra of interest, or it may be coupled to the mass spectrometer.

[0125] The training data set and the classification models according to embodiments of the invention can be embodied

by computer code that is executed or used by a digital computer. The computer code can be stored on any suitable computer readable media including optical or magnetic disks, sticks, tapes, etc., and can be written in any suitable computer programming language including R, C, C++, visual basic, etc.

[0126] The learning algorithms described above are useful both for developing classification algorithms for the biomarkers already discovered, and for finding new biomarker biomarkers. The classification algorithms, in turn, form the base for diagnostic tests by providing diagnostic values (e.g., cut-off points) for biomarkers used singly or in combination.

IV. Kits for the Detection of Brain Injury Biomarkers

[0127] In another aspect, the present invention provides kits for qualifying brain injury status, which kits are used to detect the biomarkers described herein. In a specific embodiment, the kit is provided as an ELISA kit comprising antibodies to the biomarkers of the present invention including, but not limited to, MBP, GFAP, PAD-2, tubulin beta-4B chain, tubulin alpha-1B chain, CNPase, PPIA, Septin-7, Elongation factor1-alpha2, TPPP, TPPP3, Ermin Isoform 2, NDRG2 Isoform 2, ASTN1, BAI3, CNDP1, ERMIN, GFAP, GRM3, KLH32, MAGE2, NRG3, NRGN, OMG, SLC39A12, RTN1, MT3, and PAD2, as well as the peptides listed in Tables 1-11 and 13-17. In a specific embodiment, the antibodies specifically bind to the modified or unmodified forms of NRGN or peptides thereof.

[0128] The ELISA kit may comprise a solid support, such as a chip, microtiter plate (e.g., a 96-well plate), bead, or resin having biomarker capture reagents attached thereon. The kit may further comprise a means for detecting the biomarkers, such as antibodies, and a secondary antibody-signal complex such as horseradish peroxidase (HRP)-conjugated goat anti-rabbit IgG antibody and tetramethyl benzidine (TMB) as a substrate for HRP.

[0129] The kit for qualifying brain injury status may be provided as an immuno-chromatography strip comprising a membrane on which the antibodies are immobilized, and a means for detecting, e.g., gold particle bound antibodies, where the membrane, includes NC membrane and PVDF membrane. The kit may comprise a plastic plate on which a sample application pad, gold particle bound antibodies temporally immobilized on a glass fiber filter, a nitrocellulose membrane on which antibody bands and a secondary antibody band are immobilized and an absorbent pad are positioned in a serial manner, so as to keep continuous capillary flow of blood serum.

[0130] In certain embodiments, a patient can be diagnosed by adding blood or blood serum from the patient to the kit and detecting the relevant biomarkers conjugated with antibodies, specifically, by a method which comprises the steps of: (i) collecting blood or blood serum from the patient; (ii) separating blood serum from the patient's blood; (iii) adding the blood serum from patient to a diagnostic kit; and, (iv) detecting the biomarkers conjugated with antibodies. In this method, the antibodies are brought into contact with the patient's blood. If the biomarkers are present in the sample, the antibodies will bind to the sample, or a portion thereof. In other kit and diagnostic embodiments, blood or blood serum need not be collected from the patient (i.e., it is

already collected). Moreover, in other embodiments, the sample may comprise a tissue sample or a clinical sample.

[0131] The kit can also comprise a washing solution or instructions for making a washing solution, in which the combination of the capture reagents and the washing solution allows capture of the biomarkers on the solid support for subsequent detection by, e.g., antibodies or mass spectrometry. In a further embodiment, a kit can comprise instructions for suitable operational parameters in the form of a label or separate insert. For example, the instructions may inform a consumer about how to collect the sample, how to wash the probe or the particular biomarkers to be detected, etc. In yet another embodiment, the kit can comprise one or more containers with biomarker samples, to be used as standard(s) for calibration.

IV. Citrullinated Polypeptides

[0132] The present invention provides citrullinated polypeptides. As used herein, the terms “polypeptide,” “peptide,” and “protein” are used interchangeably herein to refer to polymers of amino acids of any length. Thus, peptides, oligopeptides, and proteins are included within the definition of polypeptide. This term also encompasses post expression modifications of the polypeptide, for example, glycosylations, acetylations, phosphorylations and the like. Included within the definition are, for example, polypeptides containing one or more analogues of an amino acid (including, for example, unnatural amino acids, etc.), polypeptides with substituted linkages, as well as other modifications known in the art, both naturally occurring and non-naturally occurring.

[0133] In a particular aspect, the citrullinated polypeptide sequences are human amino acid sequences. The present invention also encompasses both the full length amino acid sequences having at least one of the arginine amino acids converted to a citrulline, or fragments thereof. The limitation being that any fragment of any desired length has at least one citrulline that specifically binds to autoantibodies. Indeed, in a specific example, the present invention encompasses any NRGN peptide that is demonstrated to be a potential target of cit-NRGN auto antibodies in brain injury patients. The full length amino acid sequence for human NRGN is shown at SEQ ID NO:1. As a further example, the amino acid sequences for MBP, GFAP, and PAD-2 are shown in SEQ ID NOS:2-4, respectively. The amino acid sequences of the other biomarker proteins described herein, namely, tubulin beta-4B chain, tubulin alpha-1B chain, CNPase, PPIA, Septin-7, Elongation factor1-alpha2, TPPP, TPPP3, Ermin Isoform 2, NDRG2 Isoform 2, ASTN1, BAI3, CNDP1, ERMIN, GRM3, KLH32, MAGE2, NRG3, OMG, SLC39A12, RTN1, MT3, are publicly available.

[0134] In a particular aspect, the citrullinated peptides of the invention bind with high affinity to cit-peptide autoantibodies (the term “peptide” is used generally and can refer to a protein biomarker described herein including NRGN, MBP, etc.). It is understood by one of skill in the art, that “high affinity” is used synonymously with the terms “specifically binds to” and “specific for” and refers to the capability of the citrullinated NRGN peptides (for example) to bind with higher or increased affinity to a cit-NRGN autoantibody as compared with a non-citrullinated NRGN antibody. It is further understood that such binding affinity can be readily established for example in vitro using a peptide binding assay in which a sample peptide is used to displace a standard peptide.

[0135] Accordingly, in a specific embodiment, the present invention provides a NRGN polypeptide comprising one or more citrullinated arginine sites. In another embodiments, the present invention provides peptides comprising one or more citrullinated arginine sites of MBP, GFAP, PAD-2, tubulin beta-4B chain, tubulin alpha-1B chain, CNPase, PPIA, Septin-7, Elongation factor1-alpha2, TPPP, TPPP3, Ermin Isoform 2, NDRG2 Isoform 2, ASTN1, BAI3, CNDP1, ERMIN, GRM3, KLH32, MAGE2, NRG3, OMG, SLC39A12, RTN1, MT3, isoforms thereof, post-translationally modified forms thereof. Reference is made to the Tables and sequence listing provided herein for the citrullinated arginine residues.

[0136] Methods for generating peptides are well known in the art. Therefore, additional citrullinated polypeptides are within the scope of the invention. The replacement of one or more arginine residues with a citrulline residue to generate a multitude of citrullinated polypeptides (of any length) and subsequent analyses thereof can be carried out without undue experimentation.

[0137] Certain of these polypeptide sequences may contain additional arginines that may be converted to citrulline. Generally, the polypeptides of the present invention may comprise any suitable length for specific recognition by cit-peptide autoantibodies. In one embodiment, the citrullinated polypeptide is a full length protein. In other embodiments, the citrullinated polypeptide may comprise at least about 2 amino acids to about several hundred amino acids in length. More specifically, the citrullinated polypeptides may comprise at least about 8 or 9 amino acids to several hundred amino acids in length. Even more specifically, the citrullinated polypeptides may comprise at least about 10-20 amino acids in length to at least about 100 amino acids in length. Furthermore the citrullinated polypeptides of the present invention may comprise about 9 to about 50 amino acids in length and include any ranges of length therein (i.e., 9-50, 9-45, 9-40, 9-35, 9-30, 9-25, 9-20, 9-15, etc.) as is understood by one of skill in the art. Peptides of over about 50 amino acids in length are also encompassed by the present invention. The length of polypeptide is only restricted by its binding capability to specifically bind cit-polypeptide autoantibodies.

[0138] The polypeptides of the present invention may also include multi-peptides. In the context of the present invention, a multi-peptide is a molecule comprised of at least two antigenic peptide units, i.e. combinations of peptide units that may or may not be linked by a covalent bond. Such multi-peptides may be comprised of linear, branched, cyclic peptide units or a combination of these. Multi-peptides may be comprised both of peptide units having the same amino acid sequence, and of peptide units having different amino acid sequences. A multi-peptide according to the invention comprises at least 7, preferably at least 10 amino acids, i.e. the peptide units may overlap. In specific embodiments, the present invention may comprise cyclic versions of citrullinated polypeptides.

[0139] In addition, the citrullinated polypeptides of the present invention may also include dimers and trimers of the peptides as well as additional stabilizing flanking sequences as is understood by those of skill in the art and described for example in U.S. Pat. Nos. 6,184,204 and 5,824,315. A multimer according to the invention can either be a homomer, consisting of a multitude of the same peptide, or a heteromer consisting of different peptides. As stated, the

amino acid sequences of the polypeptides according to the invention can be flanked by random amino acid sequences. Preferred are flanking sequences that have a stabilizing effect on the polypeptides, thus increasing their biological availability. In addition, other peptidomimetics are also useful in the polypeptides of the present invention. For a general review, see A. F. Spatola, in *CHEMISTRY AND BIOCHEMISTRY OF AMINO ACIDS, PEPTIDES AND PROTEINS*, B. Weinstein, eds., Marcel Dekker, New York, p. 267 (1983). The polypeptides of the invention also encompass polypeptides that have been modified by, for example, phosphorylation, glycosylation or lipidation.

[0140] Furthermore, the polypeptides of the present invention may also encompass “functionally equivalent variants” or “analogues” of the polypeptides. As such, this would include but not be limited to polypeptides with partial sequence homology, polypeptides having one or more specific conservative and/or non-conservative amino acid changes and polypeptide conjugates which do not alter the biological or structural properties of the polypeptide.

[0141] In terms of “functional analogues”, it is well understood by those skilled in the art, that inherent in the definition of a biologically functional polypeptide analogue is the concept that there is a limit to the number of changes that may be made within a defined portion of the molecule and still result in a molecule with an acceptable level of equivalent biological activity. A plurality of distinct polypeptides with different substitutions may easily be made and used in accordance with the invention. It is also understood that certain residues are particularly important to the biological or structural properties of a polypeptide, and such residues may not generally be exchanged.

[0142] Functional analogues can be generated by conservative or non-conservative amino acid substitutions. Amino acid substitutions are generally based on the relative similarity of the amino acid side-chain substituents, for example, their hydrophobicity, hydrophilicity, charge, size and the like. Thus, within the scope of the invention, conservative amino acid changes means, an amino acid change at a particular position which is of the same type as originally present; i.e. a hydrophobic amino acid exchanged for a hydrophobic amino acid, a basic amino acid for a basic amino acid, etc. Examples of conservative substitutions include the substitution of non-polar (hydrophobic) residues such as isoleucine, valine, leucine or methionine for another, the substitution of one polar (hydrophilic) residue for another such as between arginine and lysine, between glutamine and asparagine, between glycine and serine, the substitution of one basic residue such as lysine, arginine or histidine for another, or the substitution of one acidic residue, such as aspartic acid or glutamic acid for another, the substitution of a branched chain amino acid, such as isoleucine, leucine, or valine for another, the substitution of one aromatic amino acid, such as phenylalanine, tyrosine or tryptophan for another. Such amino acid changes result in functional analogues in that they do not significantly alter the overall charge and/or configuration of the polypeptide. Examples of such conservative changes are well-known to the skilled artisan and are within the scope of the present invention. Conservative substitution also includes the use of a chemically derivatized residue in place of a non-derivatized residue provided that the resulting polypeptide is a biologically functional equivalent to the polypeptides of the invention. Therefore, the citrullinated polypeptides of the

present invention encompass a polypeptide having an amino acid sequence that differs from the sequences provided herein by one or more conservative amino acid substitutions. The citrullinated polypeptides of the invention also encompass a polypeptide having an amino acid sequence that differs from the sequences provided herein by a single mutation, where the single mutation represents a single amino acid deletion, insertion or substitution.

[0143] The present invention further provides citrullinated peptides. The citrullinated peptides of the present invention may be made by methods known to those of skill in the art most notably and preferably by chemical synthesis using techniques well known in the chemistry of proteins such as solid phase synthesis (Merrifield et al., 65 *J. AM. CHEM. ASSOC.* 2149 (1964); Merrifield et al., 85 *J. AMER. CHEM. SOC.* 2149 (1963); and Merrifield et al., 35 *INT. J. PEPTIDE PROTEIN RES.* 161-214 (1990)) or synthesis in homogenous solution (*METHoDs OF ORGANIC CHEMISTRY*, E. Wansch (Ed.) Vol. 15, pts. I and II, Thieme, Stuttgart (1987)) to generate synthetic peptides. Citrulline is a post-translationally modified arginine that is created through the process of deimination which is catalyzed by the enzyme peptidylarginine deiminase 4 (PAD-4) that removes a positive charge from arginine and makes the resulting citrulline polar in nature.

[0144] In one embodiment, citrullinated peptides of the invention can be made from known commercially available sources. In this aspect, the lyophilized protein is reconstituted in an appropriate buffer to which the enzyme peptidylarginine deiminase 4 is added. Alternatively, Ca^{2+} is added to PAD-4 in solution. The solution is allowed to stand at an appropriate temperature for a time sufficient to cause modification of arginine residues to citrulline and thus create a citrullinated protein. The citrullinated protein is then isolated by the removal of the enzyme using a high molecular weight membrane to separate the enzyme or other methods of chromatography. One of skill in the art will understand that the temperature of incubation, buffer condition and time of incubation may vary depending on the protein that is being deiminated (Masson-Bessiere et al., 166 *J. IMMUNOL.* 4177-4184 (2001)).

[0145] The citrullinated proteins may be further isolated and purified by methods selected on the basis of properties revealed by its sequence. Purification can be achieved by protein purification procedures such as chromatography methods (gel-filtration, ion-exchange and immunoaffinity), by high-performance liquid chromatography (HPLC, RP-HPLC, ion-exchange HPLC, size-exclusion HPLC, high-performance chromatofocusing and hydrophobic interaction chromatography) or by precipitation (immunoprecipitation). Polyacrylamide gel electrophoresis can also be used to isolate the citrullinated proteins based on the molecular weight of the protein, charge properties and hydrophobicity. The purified citrullinated proteins can be used in further biochemical analyses to establish secondary and tertiary structure which may aid in the design of pharmaceuticals to interact with the protein, alter the protein charge configuration or charge interaction with other proteins or alter its function.

V. Assays of Autoantibodies to Citrullinated Peptides

[0146] The present invention provides compositions and methods for using citrullinated peptides. In several embodiments, the citrullinated peptides described herein can be

used to assay for the presence of corresponding autoantibodies. In a specific embodiment, a method for detecting the presence of autoantibodies to citrullinated NRG1 in a subject comprises contacting a biological sample taken from a subject with a citrullinated NRG1 polypeptide, and detecting the binding of the polypeptide with an autoantibody specific for the polypeptide, wherein the detection of binding is indicative of the presence of citrullinated NRG1 autoantibodies in the subject. The present invention contemplates the detection of autoantibodies to tubulin beta-4B chain, tubulin alpha-1B chain, CNPase, PPIA, Septin-7, Elongation factor1-alpha2, TPPP, TPPP3, Ermin Isoform 2, NDRG2 Isoform 2, ASTN1, BAI3, CNDP1, ERMIN, GRM3, KLH32, MAGE2, NRG3, OMG, SLC39A12, RTN1, MT3, isoforms thereof, post-translationally modified forms thereof, or combinations of any of the foregoing. More specifically, the present invention contemplates the detection/quantification/measurement of autoantibodies against the modified and/or unmodified protein/peptide.

[0147] Methods for assaying such autoantibodies are described herein in and known to those of ordinary skill in the art. For example, an immunoassay can be used to detect and analyze autoantibodies in a biological sample. As used herein, the term "immunoassay" is used in reference to any method in which antibodies are used in the detection of an antigen. It is contemplated that a range of immunoassay formats be encompassed by this definition, including but not limited to, direct immunoassays, indirect immunoassays, and "sandwich immunoassays." However, it is not intended that the present invention be limited to any particular format. It is contemplated that other formats, including radioimmunoassays (RIA), immunofluorescent assays (IFA), and other assay formats, including, but not limited to, variations on the ELISA, RIA and/or IFA methods will be useful in the methods of the present invention. The term also includes immunoprecipitation and immunoblotting.

[0148] Thus, in one aspect, the methods of the present invention include using a sandwich assay to detect the cit-NRG1 autoantibodies. It is understood that in the context of this section, the term NRG1 is used as a specific example out of convenience. The assays and peptides described below are applicable to the other proteins/peptides/biomarkers described herein including, but not limited to, tubulin beta-4B chain, tubulin alpha-1B chain, CNPase, PPIA, Septin-7, Elongation factor1-alpha2, TPPP, TPPP3, Ermin Isoform 2, NDRG2 Isoform 2, ASTN1, BAI3, CNDP1, ERMIN, GRM3, KLH32, MAGE2, NRG3, OMG, SLC39A12, RTN1, MT3, isoforms thereof, post-translationally modified forms thereof, or combinations of any of the foregoing. Sandwich assays generally involve the use of two binding agents, e.g., antibodies, each capable of binding to a different portion, or epitope, of the protein(s) to be detected and/or quantified. In a sandwich assay, the analyte is typically bound by a first binding agent which is immobilized on a solid support, and thereafter a second binding agent binds to the analyte, thus forming an insoluble complex. See, e.g., U.S. Pat. No. 4,376,110. Alternatively, the sandwich assay may be performed in solution, also referred to as a homogeneous assay. See, e.g., U.S. Pat. No. 7,413,862.

[0149] In some embodiments of these methods, a capture probe including a first binding agent is capable of specifically binding to a brain injury-associated antigen, e.g., a NRG1 polypeptide, which is bound to one or more autoantibodies. In turn, the detection probe including a second

binding agent binds to the autoantibodies. Thus, in this particular example, a four-part complex is formed between: (1) the capture probe, (2) the disease-associated antigen, (3) the autoantibody, and (4) the detection probe. In an alternative embodiment, the positions of the first and second binding agents are reversed, such that the capture probe attached to the solid support is capable of specifically binding to the autoantibodies and the detection probe is capable of specifically binding to the brain injury-associated antigen.

[0150] As stated above, the methods can be performed using any immunological technique known to those skilled in the art of immunochemistry. As examples, ELISA, immunofluorescence, radioimmunoassays or similar techniques may be utilized. In general, an appropriate capture probe is immobilized on a solid surface and the sample to be tested (e.g., human serum) is brought into contact with the capture probe. For example, modified glass substrates that covalently or non-covalently bind proteins can be used to bind the disease-associated antigen. The substrate may be treated with suitable blocking agents to minimize non-specific binding. If the autoantibody is present in the sample, a complex between the autoantibody and the capture probe is formed. A detection probe is then added, which specifically recognizes an epitope of a human immunoglobulin (Ig), if present. The anti-human immunoglobulin detection probe may be directed against the Fc region of the human antibody and with as little cross-reactivity as possible against the capture antibody species.

[0151] In another embodiment, the methods comprise contacting a sample with a capture probe including an antibody capable of binding to a disease-associated antigen. The sample is also contacted with a detection probe including anti-human Ig antibodies. The presence, absence, and/or amount of the complex may be detected, wherein the presence or absence of the complex is indicative of the presence or absence of the autoantibodies.

[0152] The complex can then be detected or quantitatively measured using methods well-known in the art. The detection probe may be labeled with biochemical markers such as, for example, a nanoparticle, horseradish peroxidase (HRP) or alkaline phosphatase (AP), and detection of the complex can be achieved by the addition of a substrate for the enzyme which generates a calorimetric, chemiluminescent or fluorescent product. Alternatively, the presence of the complex may be determined by addition of a marker protein labeled with a detectable label, for example an appropriate enzyme. In this case, the amount of enzymatic activity measured is inversely proportional to the quantity of complex formed and a negative control is needed as a reference to determine the presence of antigen in the sample. Another method for detecting the complex may utilize antibodies or antigens that have been labeled with radioisotopes followed by measure of radioactivity.

[0153] The sample may be contacted with the detection probe before, after, or simultaneously with the capture probe. In one embodiment, the sample is first contacted with the detection probe so that autoantibodies present in the sample bind to the detection probe to form a target analyte complex. The mixture is then contacted with the substrate having capture probes bound thereto so that the target analyte complex binds to the capture probe on the substrate. In another embodiment, the sample is first contacted with the substrate so that a target analyte complex present in the

sample binds to a capture probe, and the target analyte complex bound to the capture probe is then contacted with the detection probe so that the autoantibodies bind to the detection probe. In another embodiment, the sample, the detection probe and the capture probe on the substrate are contacted simultaneously.

[0154] The present invention further provides kits for commercial sale. In certain embodiments, the kit may comprise at least one cit-NRGN polypeptide. The kit may comprise the equipment, solutions and and/or instructions necessary for all steps in the process of creating cit-NRGN polypeptides, detecting cit-NRGN autoantibodies, and the like. Furthermore, the kit can further comprise peptides of MBP, GFAP, PAD-2, tubulin beta-4B chain, tubulin alpha-1B chain, CNPase, PPIA, Septin-7, Elongation factor1-alpha2, TPPP, TPPP3, Ermin Isoform 2, NDRG2 Isoform 2, ASTN1, BAI3, CNDP1, ERMIN, GRM3, KLH32, MAGE2, NRG3, OMG, SLC39A12, RTN1, MT3, isoforms thereof, post-translationally modified forms thereof, for use in detecting autoantibodies thereto.

[0155] Without further elaboration, it is believed that one skilled in the art, using the preceding description, can utilize the present invention to the fullest extent. The following examples are illustrative only, and not limiting of the remainder of the disclosure in any way whatsoever.

EXAMPLES

[0156] The following examples are put forth so as to provide those of ordinary skill in the art with a complete disclosure and description of how the compounds, compositions, articles, devices, and/or methods described and claimed herein are made and evaluated, and are intended to be purely illustrative and are not intended to limit the scope of what the inventors regard as their invention. Efforts have been made to ensure accuracy with respect to numbers (e.g., amounts, temperature, etc.) but some errors and deviations should be accounted for herein. Unless indicated otherwise, parts are parts by weight, temperature is in degrees Celsius or is at ambient temperature, and pressure is at or near atmospheric. There are numerous variations and combinations of reaction conditions, e.g., component concentrations, desired solvents, solvent mixtures, temperatures, pressures and other reaction ranges and conditions that can be used to optimize the product purity and yield obtained from the described process. Only reasonable and routine experimentation will be required to optimize such process conditions.

Materials and Methods

[0157] Reagents and Chemicals.

[0158] Bovine myelin basic protein (MBP) and peptidylarginine deiminase type 2 (PAD2) were obtained from Sigma-Aldrich (St. Louis, Mo.). Bovine neurogranin (NRGN) and bovine glial fibrillary acidic protein (GFAP) were identified by mass spectrometry (MS) within the bovine MBP sample. Recombinant human NRGN was expressed in bacteria (Rosetta 2 DE3, pEX—N-His-NRGN plasmid) and purified with Ni-NTA agarose beads. Additional amino acid residues were introduced in the recombinant protein NRGN, including His tag at N-terminus, T79 and R80 at C-terminus (FIG. 8). Native human glial fibrillary acidic protein (GFAP) from human brain was purchased from Calbiochem (EMD Chemical, Inc., Gibbstown, N.J.). Human MBP was identified by mass spectrometry within the

human GFAP sample. Sequencing grade modified trypsin and endoproteinase Glu-C were purchased from Promega (Madison, Wis.). Endoproteinase Lys-C was obtained from Roche Diagnostics (Indianapolis, Ind.). RapiGest surfactant was purchased from Waters (Milford, Mass.). Other chemicals were from Sigma-Aldrich (St. Louis, Mo.).

[0159] Peptidylargininedeiminase Treatment and Proteolysis.

[0160] Bovine MBP, human GFAP, and human recombinant protein NRGN were untreated or treated with PAD2 to convert peptidylarginine to peptidylcitrulline. Protein concentration was determined using a Pierce bicinchoninic acid (BCA) protein assay kit (Pierce, Rockford, Ill.). Each protein (2 μ g) was treated with 0.15 μ g of PAD2 (specific activity, 0.254 unit/ μ g; one unit will produce 1 μ mole of N- α -benzoylcitrulline ethyl ester from N-benzoyl-L-arginine ethyl ester (BAEE) per hr at 55° C. at pH 7.2) in a total volume of 40 μ L of a buffer composed of 20mM CaCl₂, 200 mM Tris-HCl pH 7.5, 10 mM dithiothreitol, at 55° C. for 2 hours. Untreated samples were dissolved in the same buffer. Samples with and without PAD2 treatment were denatured with 0.1% RapiGest surfactant, reduced with 5 mM tris(2-carboxyethyl)phosphine at 50° C. for 30 minutes, and alkylated with 10 mM iodoacetamide at room temperature for 30 min (in the dark). Sequencing grade modified trypsin, Lys-C, or Glu-C was added to protein samples at a ratio of 1:20 (enzyme to substrate). The samples were incubated at 37° C. for 16 hours. Samples were desalted by solid phase extraction with Oasis reverse-phase HLB cartridges (30 mg/30 μ m, Waters, Milford, Mass.).

[0161] Protein Extraction from Human Brain Tissue Samples.

[0162] Human spinal cord tissue was obtained with permission from two patients of the School of Medicine Hospital at Johns Hopkins University. The tissue sample (~60 mg) was homogenized in 0.5 mL lysis buffer (8 M urea, Amberlite IRN 150L, 2 M thiourea, 4% Chaps, 1% DTT) and proteins were extracted with Sample Grinding Kit (GE Healthcare, Piscataway, N.J.) according to the product protocol. The homogenate was centrifuged at 12,000 rpm for 10 min and the supernatant was saved in -80° C. freezer. The supernatant (100 μ L) was cleaned with 2-D clean-up kit (GE Healthcare, Piscataway, N.J.) following the instruction. After precipitation and centrifugation, the pellet was dissolved in 6 M urea. The protein concentration of two preparations was determined using Pierce BCA protein assay kit (Pierce, Rockford, Ill.). The protein extract was reduced, alkylated, and digested with trypsin, Lys-C, or Glu-C as described above.

[0163] LC-MS/MS and Data Analysis.

[0164] Digested protein samples (100 ng) were analyzed by nanoflow HPLC on a LTQ-Orbitrap mass spectrometer (Thermo Fisher Scientific, San Jose, Calif.). An Agilent 1200 series nanoflow LC system (Agilent, Santa Clara, Calif.) was used for chromatographic separation with Solvent A (0.1% formic acid) and Solvent B (90% acetonitrile in 0.1% formic acid). Samples were loaded onto reverse-phase capillary columns, 75 μ m inner diameter PicoFrit-Self/P column (New Objective, Woburn, Mass.), in house packed with 10 cm of Magic C18AQ packing material (5 μ m diameter particles, 200 A pore size) from Michrom Bioresources, Inc. (Auburn, Calif.). Separation started with 2% solvent B for 8 min at a flow rate of 2 μ L/min. A 36 min

linear gradient from 10% to 45% solvent B was followed by a 10 min linear gradient from 45% to 95% solvent B at a flow rate of 300 nL/min.

[0165] The LTQ-Orbitrap with a nanoelectrospray ionization source was controlled by Xcalibur 2.0.7 and LTQ-Orbitrap MS SP2 (Thermo Fisher Scientific, San Jose, Calif.). The LTQ-Orbitrap mass spectrometer was operated in positive mode with the electrospray voltage at +1.7 kV. An MS survey scan (from m/z 250-1800) was acquired in the Orbitrap with a resolution $r=60,000$ at m/z 400. The top five most intense ions were isolated and fragmented by CID (normalized collision energy: 35%) in the linear ion trap. The MS/MS spectra were acquired in the linear ion trap at unit resolution sequentially. Dynamic exclusion was enabled with exclusion duration of 30 sec.

[0166] Peptides and proteins were initially identified using the Sorcerer 2 SEQUEST (version 3.5, Sage-N Research, Milpitas, Calif.) search engine with postsearch analysis using Scaffold 3 (Proteome Software Inc., Portland, Oreg.). MS/MS spectra were searched against the International Protein Index human sequence database (v3.79) or International Protein Index bovine sequence database (v3.68). Search parameters included carbamidomethylation at cysteines, semi-enzymatic digestion with up to three missed cleavages, asparagines or glutamine deamination as variable modifications. Mass tolerance was 50 ppm for precursor ions and 1.00 Da for fragment ions. Manual verification was applied to MS/MS spectra of citrullinated peptides and their unmodified counterpart. First, MS/MS spectra were compared with theoretical fragment ion peaks. Second, m/z values of precursor ions were compared to theoretical values obtained from MS-Product (<http://prospector.ucsf.edu>). Third, HPLC peaks of citrullinated peptides and their unmodified counterpart were identified from all protein samples.

Results

Example 1: LC-MS/MS Analysis of Bovine MBP Protein

[0167] It was reported previously that bovine MBP has two citrullination sites. Because bovine MBP was readily available commercially, bovine MBP was selected as a model protein to establish a LC-MS/MS method for the identification of citrullinated peptides from complex mixtures. Both Lys-C and trypsin were used as endoproteinase to generate peptide fragments from bovine MBP sample.

[0168] A total of five arginine residues were found citrullinated naturally in bovine MBP (Table 1). The citrullinated peptides and related unmodified peptides were identified by initial database searching. Each LTQ-MS/MS spectrum (FIGS. 1A, 2A, and 3A) was manually verified by comparing it with theoretical fragment ion peaks. Citrullination of a peptide results in mass increment increase of 0.98402amu. The peptide sequences were also confirmed by high-resolution Orbitrap MS analysis (Table 1 and FIGS. 1C, 1D). Natural citrullination at R41, R47, and R63 residues in MBP protein were first identified. Citrullination at R129 residue (R130 in human MBP) was previously reported in human MBP protein and citrullination of R96 residue was reported previously in bovine MBP. These citrullinated arginine residues are conserved in human. The citrullinated peptide NIVTPR*TPPPSQGK (residue 91-104) (SEQ ID NO:17), PGFGYGGGR*ASDYK (residue 122-134) (SEQ ID NO:10), and DGHHAAR*TTHYGSLPQK (residue 57-73) (SEQ ID NO: 13) were eluted later than their unmodified peptides on the reversed-phase HPLC column (FIGS. 1B, 2B and 3B). These results can be explained by the loss of positive charged functional group of arginine after citrullination. As shown in FIG. 2A, neutral loss of isocyanic acid (HN=CO) in the MS/MS spectra of citrullinated peptides is a typical fragmentation pathway under collision induced dissociation (CID). This unique fragment ion was monitored in the identification and verification of citrullinated peptides.

TABLE 1

Citrullinated Peptide and Related Unmodified Peptide of MBP Identified in Bovine MBP Digested with Lys-C or Trypsin; New Sites are Underlined.							
Peptide Sequence	Sequence Number ^a	Enzyme	Theoretical m/z ^b	Experimental M/z	z ^c	R.T. (min)	Citrullinated amino acid residue
NIVTPRTPPPSQGK (SEQ ID NO: 8)	91-104 ^d	Lys-C	746.4181	746.4188	+2	27.9	NA ^e
NIVTPR*TPPPSQGK (SEQ ID NO: 9)	91-104	Lys-C	746.9095	746.9110	+2	29.1	R96 ^f
PGFGYGGGRASDYK (SEQ ID NO: 10)	122-134	Lys-C	687.8261	687.8264	+2	29.9	NA
PGFGYGGGR*ASDYK (SEQ ID NO: 11)	122-134	Lys-C	688.3181	688.3188	+2	31.7	R129 ^g
DGHHAAR*TTHYGSLPQK (SEQ ID NO: 12)	57-73	Lys-C	625.9785	625.9796	+3	26.2	NA
DGHHAAR*TTHYGSLPQK (SEQ ID NO: 13)	57-73	Lys-C	626.3065	626.3068	+3	27.0	R63 ^g
DTGILDLSLGR*FFGSDR (SEQ ID NO: 14)	32-47	Trypsin	878.9287	878.9311	+2	47.7	R41 ^g
DTGILDLSLGR*FFGSDR*GAPK (SEQ ID NO: 15)	32-51	Trypsin	1056.0238	1056.0277	+2	46.7	R41, R47 ^g

TABLE 1-continued

Citrullinated Peptide and Related Unmodified Peptide of MBP Identified in Bovine MBP Digested with Lys-C or Trypsin; New Sites are Underlined.							
Peptide Sequence	Sequence Number ^a	Enzyme	Theoretical m/z ^b	Experimental M/z	z ^c	R.T. (min)	Citrullinated amino acid residue
NIVTPR (SEQ ID NO: 16)	91-96	Trypsin	350.2110	350.2110	+2	26.1	NA
NIVTPR*TPPPSQGK (SEQ ID NO: 9)	91-104	Trypsin	746.9095	746.9102	+2	28.4	R96 ^f
PGFGYGGR (SEQ ID NO: 17)	122-129	Trypsin	405.6983	405.6981	+2	28.6	NA
PGFGYGGR*ASDYK (SEQ ID NO: 11)	122-134	Trypsin	688.3181	688.3188	+2	30.8	R129 ^f

^asequence number, the initiating amino acid at position 1 is alanine for bovine MBP

^bmass-to-charge ratio of the peptide

^ccharge state of selected precursor ions

^dpeptides with or without phosphorylation at T97 were both detected

^eNA, not applicable, as no citrullinated residues was identified

^fPeptide sequence is conserved in human.

^gCitrullinated arginine residue is conserved in human, but peptide sequence is different.

[0169] These citrullinated amino acid residues were confirmed in tryptic digests of bovine MBP. The observation of citrullinated peptides NIVTPR*TPPPSQGK (residue 91-104) (SEQ ID NO:9), PGFGYGGR*ASDYK (residue 122-134) (SEQ ID NO:11), and their unmodified counterpart NIVTPR (residue 91-96) (SEQ ID NO:16), PGFGYGGR (residue 122-129) (SEQ ID NO:17) indicated that citrullination inhibited proteolytic digestion by trypsin (Table 1). These results were different from a previous report that citrullination seemed to not affect tryptic cleavage. The peptide NIVTPRTPPPSQGK (SEQ ID NO:8) likely underwent deamination at Q102 (FIG. 1B). This deamination was also reported previously in the chicken MBP protein.

[0170] It was reported previously that the R106 residue of bovine MBP can be mono- or dimethylated. The peptide, GRGLSLSRFSWGAEGQK (residue 105-121) (SEQ ID NO:18) without modification, with mono- or di-methylation at R106 were observed. After PAD2 treatment, GR*GLSLSR*FSWGAEGQK (residue 105-121) (SEQ ID

NO:19), peptide GRGLSLSR*FSWGAEGQK (SEQ ID NO:20) with mono- and di-methylation at R106 were observed. These results indicated that mono- or dimethylated arginine may not be citrullinated by PAD2 enzyme.

Example 2: Bovine NRGN and GFAP in MBP Sample

[0171] A new protein, neurogranin (NRGN), was found citrullinated in bovine MBP sample. A citrullinated peptide of GFAP was also found. This is the first time that citrullinated residue of GFAP was identified. Both citrullinated peptides and their unmodified counterpart of bovine NRGN and GFAP were identified in the bovine MBP sample (Table 2). The sequence of the citrullinated peptides was confirmed by comparing MS/MS spectra with theoretical fragment ions, unique fragmentation pattern of citrullinated peptide (FIGS. 5A, 5B, 6A and 6B), and high resolution mass-to-charge ratio of precursor ions.

TABLE 2

Other Citrullinated Peptides and Proteins Identified in Bovine MBP Sample Digested with Lys-C, Glu-C, or Trypsin; New Sites are Underlined.							
Protein	Peptide Sequence	Sequence Number ^a	Enzyme	Theoretical m/z ^b	Experimental m/z	z ^c	Citrullinated amino acid residue
NRGN	GPGPGGPGGAGGARGGAGG	55-78	Lys-C	888.9040	888.9058	+2	NA ^d
	GPSGD (SEQ ID NO: 21)						
	GPGPGGPGGAGGAR*GGAG	55-78	Lys-C	889.3960	889.3983	+2	<u>R68</u> ^e
	GGPSGD (SEQ ID NO: 22)						
	GPGPGGPGGAGGAR	55-68	Trypsin	532.7652	532.7656	+2	NA
	(SEQ ID NO: 23)						
	KGPGPGGPGGAGGAR*GGA	54-78	Trypsin	953.4435	953.4429	+2	<u>R68</u> ^e
	GGGPSGD (SEQ ID NO: 24)						

TABLE 2-continued

Other Citrullinated Peptides and Proteins Identified in Bovine MBP Sample Digested with Lys-C, Glu-C, or Trypsin; New Sites are Underlined.						
Protein	Peptide Sequence	Sequence Number ^a	Enzyme	Theoretical m/z ^b	Experimental m/z	Citrullinated amino acid z ^c residue
GFAP	GHLKRNIIVVKTVE (SEQ ID NO: 25)	398-410	Glu-C	746.9515	746.9504	+2 NA
	GHLKR*NIVVKTVE (SEQ ID NO: 26)	398-410	Glu-C	747.4435	747.4420	+2 <u>R402</u> ^f

^a sequence number includes the initiating Met at position 1 for all proteins except for MBP, bovine. Met is cleaved from in the mature protein.

^b mass-to-charge ratio of the peptide

^c z is the charge state of selected precursor ions

^d NA, not applicable, as no citrullinated residues was identified

^e Citrullinated arginine residue is conserved in human, but peptide sequence is different.

^f Peptide sequence is conserved in human.

[0172] R68 residue of NRGN was identified as a natural citrullination site present in bovine NRGN. The peptide sequences were confirmed in the digests of Lys-C and trypsin. Because the primary sequence of NRGN is highly conserved across species (FIG. 4), this unique post-translational modification may also be present in human NRGN.

[0173] Citrullination at R402 residue of GFAP was identified in endogenous GFAP. Citrullinated peptide GHLKR*NIVVKTVE (residues 398-410) (SEQ ID NO: 26) and the unmodified counterpart (SEQ ID NO: 25) were observed in the Glu-C digests of bovine MBP (FIGS. 6A and 6B).

Example 3: Citrullinated Protein Identified from Human GFAP Sample

[0174] It was reported previously that GFAP and MBP protein were citrullinated in the human autoimmune disease, multiple sclerosis (MS). In the previous studies, the citrullinated GFAP and MBP were detected using an antibody against chemically modified citrulline. However, the citrullinated residue(s) of GFAP was not identified.

[0175] In this investigation, two natural citrullination sites, R270 and R406 residues, were found in an endogenous human GFAP sample digested with Lys-C or Glu-C (Table 3). LC peaks of citrullinated peptide GHLKR*NIVVKTVE (402-414) (SEQ ID NO: 30) and the unmodified counterpart are shown in FIG. 7A. MS/MS spectrum of the citrullinated peptide GHLKR*NIVVKTVE (402-414) (SEQ ID NO: 30) was the same as FIG. 6A and was not shown here. Natural citrullination of R406 residue (R402 in bovine) was also found in bovine GFAP as described above. However, relative abundance of the citrullinated peptide GHLKR*NIVVKTVE (402-414) (SEQ ID NO:30) versus unmodified peptide (SEQ ID NO:29) was much higher in the human GFAP sample (FIGS. 6B and 7A). This result indicated that higher percentage of endogenous GFAP protein was citrullinated in human sample (without considering ionization efficiencies of the two peptides).

[0176] Human MBP protein was identified in the human GFAP sample digested with Lys-C (Table 3). This was not unexpected because the GFAP sample was extracted from human brain tissue. Proteins that bind to GFAP may also be

TABLE 3

Citrullinated Peptide and Unmodified Peptide Identified in Human GFAP Samples Digested with Lys-C, Glu-C; New Sites are Underlined.						
Protein	Peptide Sequence	Sequence Number ^a	Enzyme	Theoretical m/z ^b	Experimental m/z	Citrullinated amino acid z ^c residue
GFAP, human	FADLTDAARNAE LLR QAK (SEQ ID NO: 27)	261-279	Lys-C	692.0395	692.0394	+3 36.6 NA ^d
	FADLTDAAR*NAELL RQAK (SEQ ID NO: 28)	261-279	Lys-C	692.3675	692.3679	+3 38.0 <u>R270</u>
	GHLKRNIIVVKTVE (SEQ ID NO: 29)	402-414	Glu-C	746.9515	746.9543	+2 29.5 NA
	GHLKR*NIVVKTVE (SEQ ID NO: 30)	402-414	Glu-C	747.4435	747.4464	+2 29.5 <u>R406</u>
MBP, human	NIVTPRTPPPSQGK (SEQ ID NO: 31)	92-105	Lys-C	746.4181	746.4174	+2 26.7 NA
	NIVTPR*TPPPSQGK (SEQ ID NO: 32)	92-105	Lys-C	746.9095	746.9084	+2 27.7 R97

^a sequence number includes the initiating Met at position 1 for GFAP. Met is removed in the mature protein.

^b mass-to-charge ratio of the peptide

^c z is the charge state of selected precursor ions

^d NA, not applicable, as no citrullinated residues was identified

isolated during protein purification. Both citrullinated peptide NIVTPR*TPPPSQGK (residues 226-239) (SEQ ID NO:33) and the unmodified counterpart (SEQ ID NO:34) were identified. MS/MS spectra of the citrullinated peptide NIVTPR*TPPPSQ GK (residues 226-239) (SEQ ID NO:33) is the same as FIG. 1A and was not shown here. The peptide sequences were also confirmed by high resolution precursor ion MS analysis (less than ± 5 ppm). LC peaks of the citrullinated peptide NIVTPR*TPPPSQGK (residues 226-239) (SEQ ID NO:33) and the unmodified peptide were shown in FIG. 7B. Relative ratio of the citrullinated peptide versus unmodified counterpart was higher in human MBP than that in bovine MBP protein (FIG. 7B and FIG. 1B).

Example 4: LC-MS/MS Analysis of NRGN, GFAP, and MBP after PAD2 Treatment

[0177] To locate all possible citrullination sites in NRGN, GFAP, and MBP, these three proteins were treated with PAD2 enzyme. Lys-C and Glu-C were used to generate appropriate peptides for mass spectrometry analysis. The recombinant human NRGN has additional amino acids at the N- and C-terminals introduced as a consequence of subcloning (FIG. 8).

[0178] The R68, natural citrullination site found in bovine NRGN, was also citrullinated in human recombinant NRGN after PAD2 treatment. Five additional citrullination sites

were observed for recombinant protein NRGN (Table 4 and FIGS. 9A and 9B), including the R80 amino acid introduced during subcloning. These results demonstrated that all five arginine residues of human NRGN may be citrullinated. MS/MS spectrum and extracted ion chromatogram of citrullinated peptide IQASFR*GHMAR*K (residues 33-44) (SEQ ID NO: 35) at m/z 702.36 of human recombinant NRGN in Lys-C digests were shown in FIGS. 9A and 9B. This citrullinated peptide was also observed in the Lys-C digests of bovine NRGN after PAD2 treatment. The unmodified peptide IQASFRGHMARK (residues 33-44) (SEQ ID NO: 36) of NRGN was not observed after PAD2 treatment. This result indicated that PAD2 enzyme catalyzed complete conversion of peptidylarginine to peptidylcitrulline.

[0179] For human GFAP protein, a total of thirteen citrullinated arginine residues were identified (Table 4). Two arginine residues, R270 and R406, were found citrullinated naturally in human GFAP protein (Table 3). Eleven additional citrullinated residues were found in PAD2 treated human GFAP in either Lys-C or Glu-C digestions (Table 4 and FIG. 10). The peptide sequences were confirmed by high resolution MS analysis of precursor ions and their CID spectra. Considering the fact that all arginine residues of NRGN can be citrullinated, it's possible that there are additional citrullination sites in the remaining 34 arginine residues of human GFAP.

TABLE 4

Citrullination Sites Identified from GFAP, MBP and NRGN after PAD2 Treatment Followed by Lys-C or Glu-C Digestion; New Sites are Underlined.						
Protein	Citrullinated peptide	Sequence number ^a	Theoretical m/z ^b	Experimental m/z	Enzyme	Citrullinated amino acid residue
NRGN human	IQASFR*GHMAR*K (SEQ ID NO: 35)	33-44	702.3642	702.3629	Lys-C	<u>R38</u> , <u>R43</u>
	GPGPGGPGGAGVAR*GGAGGG PSGDTR* (SEQ ID NO: 37)	55-80	1039.4859	1039.4839	Lys-C	<u>R68</u> , R80 ^c
NRGN bovine	IQASFR*GHMAR*K (SEQ ID NO: 38)	33-44	702.3642	702.3647	Lys-C	<u>R38</u> , <u>R43</u>
	GPGPGGPGGAGGAR*GGAGGG PSGD (SEQ ID NO: 39)	55-78	889.3960	889.3967	Lys-C	<u>R68</u>
GFAP human	VR*FLEQQNK (SEQ ID NO: 40)	87-95	581.8144	581.8152	Lys-C	<u>R88</u>
	ALAAELNQLR*AK (SEQ ID NO: 42)	96-107	649.8750	649.8754	Lys-C	<u>R105</u>
	LR*LR*LDQLTANSAR*LE (SEQ ID NO: 44)	123-138	936.5129	936.5170	Glu-C	<u>R124</u> , <u>R126</u> , <u>R136</u>
	AENNLAAYR*QEADE (SEQ ID NO: 46)	165-178	797.8526	797.8555	Glu-C	<u>R173</u>
	LQEQLAR*QQV HVE (SEQ ID NO: 48)	211-223	789.9154	789.9181	Glu-C	<u>R217</u>
	WYR*SKFADLTDAAR*NAE (SEQ ID NO: 50)	256-273	1043.9996	1043.9951	Glu-C	<u>R258</u> , <u>R270</u>
	FADLTDAAR*NAELLRQAK (SEQ ID NO: 28)	261-279	692.3675	692.3679	Lys-C	<u>R270</u>
	ANDYR*R*QLQSLTCDLE (SEQ ID NO: 52)	282-297	992.4575	992.4612	Glu-C	<u>R286</u> , <u>R287</u>
	LALDIEIATYR*K (SEQ ID NO: 54)	357-368	703.8981	703.8985	Lys-C	<u>R367</u>
	GHLKR*NIVVKTVE (SEQ ID NO: 56)	402-414	747.4435	747.4455	Glu-C	<u>R406</u>
MBP human	NIVTPR*TPPPSQGK (SEQ ID NO: 32)	92-105	746.9095	746.9089	Lys-C	R97

TABLE 4-continued

Citrullination Sites Identified from GFAP, MBP and NRGN after PAD2 Treatment Followed by Lys-C or Glu-C Digestion; New Sites are Underlined.						
Protein	Citrullinated peptide	Sequence number ^a	Theoretical m/z ^b	Experimental m/z	Enzyme	Citrullinated amino acid residue
MBP bovine	YLASASTMDHAR*HGFLPR*HR (SEQ ID NO: 58)	12-31	775.7150 (z: +3)	775.7166 (z: +3)	trypsin	R23, <u>R29</u>
	YLASASTMDHAR*HGFLPR*HR * (SEQ ID NO: 60)	12-31	776.0430 (z: +3)	776.0431 (z: +3)	trypsin	R23, <u>R29</u> , R31
	DTGILDLSLGR*FFGSDR*GAPK (SEQ ID NO: 61)	32-51	1056.0238	1056.0266	trypsin	<u>R41</u> , <u>R47</u>
	R*GSGKDGHHAAR*TTHYGSLP QK (SEQ ID NO: 63)	52-73	591.5454 (z: +4)	591.5466 (z: +4)	Lys-C	R52, <u>R63</u>
	DGHHAAR*TTHYGSLPQK (SEQ ID NO: 13)	57-73	938.9561	938.9566	Lys-C	<u>R63</u>
	AQGHR*PQDENPVVHFFK (SEQ ID NO: 65)	74-90	1003.9952	1003.9971	Lys-C	R78
	NIVTPR*TPPPSQGK ^d (SEQ ID NO: 9)	91-104	746.9095	746.9109	Lys-C	R96
	GR*GLSLSR*FSWGAEGQK ^e (SEQ ID NO: 66)	105-121	919.4632	919.4651	Lys-C	<u>R106</u> , <u>R112</u>
	PGFYGGR*ASDYK (SEQ ID NO: 11)	122-134	688.3181	688.3184	Lys-C	R129
	LGGR*DSR*SGSPMAR*R* (SEQ ID NO: 68)	155-169	803.8837	803.8842	Lys-C	R158, <u>R161</u> , <u>R168</u> , R169

^a sequence number includes the initiating Met at position 1 for all proteins except for MBP, bovine

^b mass-to-charge ratio of doubly charged ions; charge state of precursor ions is +2 usually

^c T79 and R80 are introduced into recombinant protein NRGN during subcloning

^d peptides with or without phosphorylation at T97 were both detected

^e mono- and di-methylation of R106 were observed on this peptide

[0180] It was reported previously that 11 arginine residues of human MBP may be citrullinated. Six arginine residues were endogenous citrullination sites (FIG. 11) and five sites were citrullinated after PAD4 treatment. Bovine MBP was reported to have at least two citrullination sites (FIG. 11).

[0181] It was found herein that residue R97 was naturally citrullinated in human MBP (isoform 5, 170 residues) as described above. In bovine MBP protein, three new natural citrullination sites, R41, R47, and R63, were identified which were not reported previously. Two natural citrullination sites at the residues R96 and R129 (R130 in human MBP) were confirmed. In the bovine MBP sample treated with PAD2 enzyme, five new sites were identified, and eleven citrullination sites were confirmed (Table 4 and FIG. 11). The C-terminal peptide LGGR*DSR*SGSPMAR*R* contains four citrullination sites. The sequences of all peptides were confirmed by high resolution MS analysis of precursor ions and their CID spectra. These results showed that all arginine residues of MBP may be citrullinated by PAD enzyme. As shown above, eight new citrullination sites have been identified in bovine MBP: R29, R41, R47, R63, R106, R112, R161, and R168. The corresponding residues in human MBP are R31, R43, R49, R65, R107, R113, R162, and R169.

Example 5: Citrullinated Protein in Human Tissue Samples

[0182] MBP, GFAP, and NRGN were identified in the human spinal brain tissue sample in both trypsin and lys-C digests. Protein sequence coverage was 64%, 54%, and 32% respectively. For MBP, eleven endogenous citrullination sites were confirmed in human tissue sample (FIG. 11). The

citrullination sites include five novel arginine residues (R43, R49, R65, R162, and R169) that were identified in this invention. This result proves that citrullinated forms of neurological proteins are present in human samples. The modified form of the peptides and proteins can be detected in complex mixtures using our current method.

Example 6: Multiple Reactions Monitor (MRM) Assays for NRGN, GFAP, and MBP

[0183] Based on these discoveries, the present inventors have developed MRM assays for the quantification of endogenous human proteins, NRGN, GFAP, MBP, and PAD2 (Tables 5-9, 11-12, and 14-17). The MRM assays were only a fraction of possible example bioanalytical assays based on citrullinated and unmodified peptides described in Table 1-4. These are not a full list of all possible MRM assays. Alternative MRM assays can be developed based on peptides ions of different charge states, peptides with other modifications (for example, oxidation, methylation, and phosphorylation), peptides with combination of citrullinated residues and/or modifications, and peptides generated using other endoprotease or chemical reagents. Additional citrullination sites may be present for the human GFAP protein. Similar MRM assays can be developed for these additional modified peptides of human GFAP protein. MRM assays could be developed for citrullinated peptides or proteins that were chemically modified.

[0184] Other bioanalytical assays can also be developed targeting at either citrullinated peptides and/or their unmodified counterparts. For example, antibodies could be produced to the modified proteins or peptides and be used as assay reagents. Additional assays include but are not limited

to, ELISA assays, HPLC with various detectors, capillary electrophoresis (CE) coupled with a mass spectrometer or any other detectors. Similar bioanalytical assays could be developed for citrullinated peptides or proteins that were chemically modified. For high sensitive assays, enrichment method could be developed to improve recovery of modified proteins or peptides. The enrichment methods could be based on chemical reactions that are specific to ureido group

of citrulline. Other enrichment methods could antibody based, for example, antibodies against conjugated citrulline, modified peptides, or modified proteins.

[0185] One or more of these modified proteins could be used in MRM assays or other bioanalytical assays. Any or combination of modified residues, peptides of one or more of these modified proteins could be used in a bioanalytical assays.

TABLE 5

Developed MRM Assay for Endogenous Human NRG1					
Protein	Peptide	Enzyme	Q1, m/z	Charge	Q3, m/z
NRGN, human	GPGPGGPGGAGVAR (SEQ ID NO: 70)	Trypsin	553.79	+2	476.75
			553.79	+2	448.24
			553.79	+2	952.50
			553.79	+2	684.38
			553.79	+2	741.40
	GPGPGGPGGAGVAR*GGAGGGPSGD (SEQ ID NO: 71)	Trypsin	910.42	+2	888.92
			910.42	+2	731.29
			910.42	+2	1089.54
			910.42	+2	674.27
			910.42	+2	546.21
	KGPDPGPGGAGVAR*GGAGGGPSGD (SEQ ID NO: 72)	Trypsin	974.47	+2	952.97
			974.47	+2	731.29
			974.47	+2	833.43
			974.47	+2	1060.55
			974.47	+2	546.21
	GPGPGGPGGAGVARGGAGGGPSGD (SEQ ID NO: 73)	Lys-C	909.93	+2	804.87
			909.93	+2	833.38
			909.93	+2	1115.51
			909.93	+2	546.24
			909.93	+2	1058.49
	GPGPGGPGGAGVAR*GGAGGGPSGD (SEQ ID NO: 71)	Lys-C	910.42	+2	888.92
			910.42	+2	731.29
			910.42	+2	1089.54
			910.42	+2	674.27
			910.42	+2	546.21
	IQASFRGHMARK (SEQ ID NO: 36)	Lys-C	467.92	+3	580.81
			467.92	+3	644.84
			467.92	+3	545.29
			467.92	+3	456.47
			467.92	+3	516.74
	IQASFR*GHMAR*K (SEQ ID NO: 35)	Lys-C	468.58	+3	454.48
			468.58	+3	581.79
			468.58	+3	645.82
			468.58	+3	607.81
			468.58	+3	546.27

TABLE 6

Developed MRM Assay for Endogenous Human GFAP					
Protein	Peptide	Enzyme	Q1, m/z	Charge	Q3, m/z
GFAP human	VRFLEQQNK (SEQ ID NO: 41)	Lys-C	581.32	+2	517.27
			581.32	+2	645.37
			581.32	+2	773.43
			581.32	+2	901.49
			581.32	+2	1015.53
	VR*FLEQQNK (SEQ ID NO: 40)	Lys-C	581.81	+2	560.31
			581.81	+2	404.23
			581.81	+2	906.47
			581.81	+2	759.40
			581.81	+2	646.32

TABLE 6-continued

Developed MRM Assay for Endogenous Human GFAP					
Protein	Peptide	Enzyme	Q1, m/z	Charge	Q3, m/z
	ALAAELNQLRAK (SEQ ID NO: 43)	Lys-C	649.38	+2	374.25
			649.38	+2	487.33
			649.38	+2	924.51
			649.38	+2	1080.62
			649.38	+2	1151.65
	ALAAELNQLR*AK (SEQ ID NO: 42)	Lys-C	649.88	+2	628.38
			649.88	+2	375.24
			649.88	+2	488.32
			649.88	+2	730.42
			649.88	+2	924.51
	LALDIEIATYRK (SEQ ID NO: 55)	Lys-C	703.41	+2	466.28
			703.41	+2	567.33
			703.41	+2	638.36
			703.41	+2	751.45
			703.41	+2	880.49
	LALDIEIATYR*K (SEQ ID NO: 54)	Lys-C	703.90	+2	682.40
			703.90	+2	839.49
			703.90	+2	881.47
			703.90	+2	752.43
			703.90	+2	568.31
	GHLKRNIIVKTV (SEQ ID NO: 57)	Glu-C	498.31	+3	486.81
			498.31	+3	573.37
			498.31	+3	673.43
			498.31	+3	593.37
			498.31	+3	649.91
	GHLKR*NIVKTV (SEQ ID NO: 56)	Glu-C	498.63	+3	484.30
			498.63	+3	487.30
			498.63	+3	410.74
			498.63	+3	575.34
			498.63	+3	674.41

TABLE 7

Developed MRM Assay for Endogenous Human MBP					
Protein	Peptide	Enzyme	Q1, m/z	Charge	Q3, m/z
MBP	NIVTPRTPPPSQGK (SEQ ID NO: 31)	Lys-C	746.42	+2	583.32
human			746.42	+2	632.86
			746.42	+2	710.38
			746.42	+2	1064.58
			746.42	+2	1165.63
	NIVTPR*TPPPSQGK (SEQ ID NO: 32)	Lys-C	746.91	+2	725.41
			746.91	+2	710.38
			746.91	+2	1065.57
			746.91	+2	1166.62
			746.91	+2	533.29
	SHGRTQDENPVVHFFK (SEQ ID NO: 74)	Lys-C	949.47	+2	782.35
			949.47	+2	802.88
			949.47	+2	911.40
			949.47	+2	987.54
			949.47	+2	1116.58
	SHGR*TQDENPVVHFFK (SEQ ID NO: 75)	Lys-C	949.96	+2	928.46
			949.96	+2	987.54
			949.96	+2	873.50
			949.96	+2	677.38
			949.96	+2	578.31
	DSHHPARTAHYGLPQK (SEQ ID NO: 76)	Lys-C	951.47	+2	372.22
			951.47	+2	792.43
			951.47	+2	893.96
			951.47	+2	929.48
			951.47	+2	1110.52
	DSHHPAR*TAHYGLPQK (SEQ ID NO: 77)	Lys-C	951.96	+2	930.46
			951.96	+2	759.36
			951.96	+2	872.95

TABLE 7-continued

Developed MRM Assay for Endogenous Human MBP					
Protein	Peptide	Enzyme	Q1, m/z	Charge	Q3, m/z
			951.96	+2	894.45
			951.96	+2	1101.57
	LGGRDSRSGSPMARR (SEQ ID NO: 69)	Lys-C	534.95	+3	515.28
			534.95	+3	523.34
			534.95	+3	552.79
			534.95	+3	714.86
			534.95	+3	499.23
	LGGR*DSR*SGSPMAR*R* (SEQ ID NO: 68)	Lys-C	803.88	+2	782.39
			803.88	+2	760.89
			803.88	+2	632.32
			803.88	+2	975.32
			803.88	+2	863.40
MBP human	GRGLSLSRFSWGAEQQRPGFGYGRASDYK ^a (SEQ ID NO: 78)	Lys-C	1073.87	+3	512.23
			1073.87	+3	796.39
			1073.87	+3	827.48
			1073.87	+3	974.55
			1073.87	+3	1061.59
	GRGLSLSRFSWGAEQQRPGFGYGR*ASDYK (SEQ ID NO: 79)	Lys-C	1074.19	+3	1059.86
			1074.19	+3	797.38
			1074.19	+3	854.4
			1074.19	+3	1017.46
			1074.19	+3	827.48
	GR*GLSLSR*FSWGAEQQRPGFGYGR*ASDYK (SEQ ID NO: 80)	Lys-C	1074.85	+3	1060.52
			1074.85	+3	797.38
			1074.85	+3	854.40
			1074.85	+3	585.34
			1074.85	+3	829.45
	GRGLSLSRFSWGAEQQRPGFGYGRASDYK ^a (SEQ ID NO: 78)	Lys-C	1073.87	+3	512.23

^aWhen Lys-C is used, this peptide is longer for human MBP protein because of sequence difference.

TABLE 8

Developed MRM Assay for Human PAD2					
Protein	Peptide	Enzyme	Q1, m/z	Charge	Q3, m/z
PAD2 human	GFPVVLDSFR (SEQ ID NO: 81)	Trypsin	543.80	+2	441.76
			543.80	+2	882.50
			543.80	+2	587.31
			543.80	+2	686.38
			543.80	+2	474.23
	WIQDEIEFGYIEAPHK (SEQ ID NO: 82)	Trypsin	987.98	+2	914.47
			987.98	+2	694.39
			987.98	+2	1061.54
			987.98	+2	1190.58
			987.98	+2	452.26

Example 6: In Vivo Analysis of Human GFAP Using CID Triggered HCD

[0186] To assess the potential for the citrullination of GFAP, the purified human protein was treated with the PAD2 enzyme in presence of activating calcium. Trypsin, Lys-C or Glu-C were used to generate appropriate peptides of the untreated and treated sample prior to mass spectrometry analysis. In the untreated sample, there were 5 natural citrullination sites, residues R30, R36, R270, R406, and R416, identified for human GFAP protein (Table 9; MS/MS spectra of the citrullinated peptides (data not shown)). Each MS/MS spectrum was manually verified by comparing it with theoretical fragment ions. The peptide sequences were also confirmed by high-resolution Orbitrap MS analysis

(Table 12). Neutral loss of isocyanic acid (HNCO) in the MS/MS spectra of citrullinated peptides is a typical fragmentation pathway upon collision induced dissociation. This unique fragment ion was monitored for identification and verification of citrullinated peptides and was observed in majority of samples.

[0187] With HCD fragmentation for the same untreated sample the peptide amide backbone cleavage dominates MS/MS spectra of citrullinated peptides as illustrated from a representative typical HCD spectrum of a citrullinated GFAP peptide shown in FIG. 12A and FIG. 12B. The relative abundance of the neutral loss fragment ion was reduced significantly. The formation of sequence-informative b and y ions with HCD greatly facilitates the identification of citrullinated peptides and proteins from complex protein digests. For the identification of citrullinated peptides, HCD spectra are complementary to CID spectra.

[0188] After PAD2 treatment of GFAP, a total of seven citrullinated arginine residues were identified (Table 9). Note this includes site that were only detected with Lys-C or Glu-C digestion and not trypsin. These included the five in vivo citrullination sites of GFAP observed in the untreated sample as well as an additional twelve citrullinated residues. The peptide sequences were confirmed by CID spectra (FIGS. 13A-13C) and the high accurate m/z values of the precursor ions.

Example 7: LC-MS/MS Analysis of Bovine MBP Protein

[0189] A total of five arginine residues of bovine MBP were found citrullinated in the untreated sample whether

digested with trypsin (Table 9) or digested with Lys-C (Table 13). The observation of citrullinated peptides NIVTPR*TPPPSQGK (residue 91-104) (SEQ ID NO:9), PGFGYGGR*ASDYK (residue 122-134) (SEQ ID NO:11), and their unmodified counterpart NIVTPR (residue 91-96), PGFGYGGR (residue 122-129) (SEQ ID NO:17) indicated that citrullination may inhibit proteolytic digestion by trypsin at the C-terminal of citrulline residue. However, one citrullinated peptide of MBP does contain citrulline at the C-terminal (Table 9 and Supplement Table 13). The peptide NIVTPRTPPPSQGK (SEQ ID NO:8) can also undergo deamination at residue Q102. This deamination was reported previously in chicken MBP protein. A neutral loss peak can be used as a diagnostic fragment ion to differentiate this peptide with a N or Q deamination from the peptide with citrullination at residue R96. However, MS/MS spectrum of peptides with N or Q deamination may not be collected due to low ion intensities. In this case, high resolution HCD spectrum was helpful in identifying the peptides with citrullination at arginine residues.

[0190] In the bovine MBP sample treated in vitro with the PAD2 enzyme, five novel (not detected previously) citrullination sites were identified, and all eleven citrullination sites that had previously reported were confirmed (Table 9). The CID spectra of all peptides were shown in Table 12 and FIG. 14. The C-terminal peptide LGGR*DSR*SGSPMAR*R* (SEQ ID NO: 83) contains four potential citrullination sites and truncated C-terminal peptides were also observed (Table 9).

Example 8: Citrullinated Peptide of NRGN

[0191] Bovine neurogranin was identified within in the commercial bovine MBP sample, as a contaminant which co-purified with MBP. This bovine NRGN protein was found both as the R68 unmodified and citrullinated forms (Table 14). The sequence of the citrullinated peptides containing R68 were confirmed by MS/MS spectra upon CID (data not shown) and accurate m/z value of precursor ions. To locate all possible citrullination sites in human NRGN, which is highly conserved with bovine, recombinant human NRGN was treated with PAD2 enzyme and the untreated and treated samples were digested with either trypsin, Lys-C, or Glu-C. Not unexpected, residue R68 of human NRGN was also citrullinated in human NRGN after PAD2 treatment. Five additional citrullination sites were observed for recombinant protein NRGN after treatment (Table 9 and Table 15). This included an additional R80 amino acid not found in the endogenous protein but had been introduced during cloning of the NRGN. These results demonstrated that all five arginine residues of human NRGN can be citrullinated in vitro. Data of MS/MS spectra of the citrullinated peptides of human NRGN not shown.

Example 9: Citrullinated Proteins in Human Brain Samples

[0192] Endogenous citrullinated sites of three brain proteins GFAP, MBP, and NRGN were identified and quantified in brain tissue samples obtained from non-Alzheimer controls (n=3) and with Alzheimer's disease (n=3) (Table 2 for clinical description). GFAP, MBP, and NRGN were identified in all of the human brain tissue samples with maximum amino acid sequence coverage of 64%, 54%, and 44% respectively. The peptide sequences and the citrullinated residues are listed in Table 11. In vivo, GFAP contained five citrullination sites at residues R30, R36, R270, R406, and R416 while MBP had fourteen residues citrullinated, including six novel arginine residues (R32, R44, R50, R92, R189, and R196). Mapping of citrullinated arginine residues demonstrated conserved in vivo citrullination of MBP R92 and R124, GFAP R406 and 416 in both the non-AZ and AZ samples and MBP R32 and GFAP R270 detected only in AZ samples. Residue R68 of NRGN was found citrullinated (representative chromatographic peak of the citrullinated and unmodified peptides containing residue R68 of human NRGN; see, FIGS. 13A-13C). With mass tolerance of 10 ppm, the citrullinated peptide can be differentiated from the C13 peak of the unmodified peptide based on the m/z value of the precursor ion.

[0193] Based on integrated peak area (FIG. 13C), occupancy rate of citrullination can be estimated by calculating the ratio of modified peptide vs. the sum of modified and unmodified peptides. It was estimated that occupancy rate of citrullinated peptide at R68 residue was 8.6% for the AD3 brain tissue sample without considering different ionization efficiency of the two peptides (FIG. 14). The occupancy rate of the endogenous citrullination site of NRGN (residue R68) and MBP (residues R92 and R124) in all brain tissue samples were determined for citrullinated peptides identified in Lys-C digests. For peptide DSHHPAR*TAHYGSLPQK of MBP that contains residue R92, peak area of both charge+4 and charge+3 ions are included. The occupancy rates of three sites were significantly higher for AD3 and Control_1 samples. Citrullination of NRGN at residue R68 was detected in AD1, AD3, and C1 control samples. The higher ratio of citrullination may be age related.

Example 10: Citrullinated Proteins in Human Brain Samples with Stroke

[0194] Endogenous citrullination of a number of additional brain proteins were identified in brain tissue samples from a single patient from an area of normal brain and from the penumbra region of an infarcted area from a clinical stroke. Several of these citrullinated forms were only identified in the stroke tissue (Tubulin beta-4B chain, Tubulin alpha-1B chain, TPPP3, and NDRG2, Isoform 2) suggesting ischemic specific modification of these proteins.

TABLE 9

Citrullination sites of MBP, GFAP, and NRGN					
Protein, UniProt ID	Peptide	Sequence number ^a	Enzyme	In vitro site	In vivo site
MBP, bovine P02687	YLASASTMDHAR*HGFLPR (SEQ ID NO: 84)	12-29	Trypsin	R23	
	YLASASTMDHAR*HGFLPR*HR (SEQ ID NO: 58)	12-31	Trypsin	R23, R29	

TABLE 9-continued

Protein, UniProt ID		Peptide	Sequence number ^a	Enzyme	In vitro site	In vivo site
		YLASASTMDHAR*HGFLPR*HR* (SEQ ID NO: 60)	12-31	Trypsin	R23, R29, R31	
		DTGILDSLGR*FFGSDR (SEQ ID NO: 14)	32-47	Trypsin	R41	R41
		FFGSDR*GAPK (SEQ ID NO: 85)	42-51	Trypsin	R47	R47
		R*GSGKDGHHAAR*TTHYGSLPQK (SEQ ID NO: 63)	52-73	Lys-C	R52, R63	R63
		DGHHAAR*TTHYGSLPQK (SEQ ID NO: 13)	57-73	Trypsin, Lys-C	R63	R63
		AQGHR*PQDENPVVHFFK (SEQ ID NO: 65)	74-90	Trypsin, Lys-C	R78	
		NIVTPR*TPPPSQGK (SEQ ID NO: 32)	91-104	Trypsin, Lys-C	R96	R96
		GR*GLSLSR*FSWGAEGQK (SEQ ID NO: 66)	105-121	Trypsin, Lys-C	R106, R112	
		PGFGYGGGR*ASDYK (SEQ ID NO: 11)	122-134	Trypsin, Lys-C	R129	R129
		LGGR*DSR*SGSPMA(-) ^c (SEQ ID NO: 87)	155-167	Trypsin, Lys-C	R158, R161	
		LGGR*DSR*SGSPMAR*(-) (SEQ ID NO: 89)	155-168	Trypsin, Lys-C	R158, R161, R168	
		LGGR*DSR*SGSPMAR*R*(-) (SEQ ID NO: 83)	155-169	Trypsin, Lys-C	R158, R161, R168, R169	
GFAP, Human P14136		R*LPGGTR*LSLAR (SEQ ID NO: 158)	30-41	Trypsin	R30, R36	R30, R36
		LPGGTR*LSLAR (SEQ ID NO: 93)	31-41	Trypsin	R36	R36
		VR*FLEQQNK (SEQ ID NO: 40)	87-95	Lys-C	R88	
		ALAAELNQLR*AK (SEQ ID NO: 42)	96-107	Lys-C	R105	
		LR*LR*LDQLTANSAR*LE (SEQ ID NO: 44)	123-138	Glu-C	R124, R126, R136	
		AENNLAAAYR*QEADE (SEQ ID NO: 46)	165-178	Glu-C	R173	
		LQEQLAR*QQVHVE (SEQ ID NO: 48)	211-223	Glu-C	R217	
		WYR*SKFADLTDAAR*NAE (SEQ ID NO: 50)	256-273	Glu-C	R258, R270	
		FADLTDAAR*NAELLR (SEQ ID NO: 95)	261-276	Trypsin	R270	R270
		FADLTDAAR*NAELLR*QAK (SEQ ID NO: 97)	261-279	Lys-C	R270, R276	
		ANDYR*R*QLQSLTCDLE (SEQ ID NO: 52)	282-297	Glu-C	R286, R287	
		LALDIEIATYR*K (SEQ ID NO: 54)	357-368	Lys-C	R367	
		GHLKR*NIVVKTVE (SEQ ID NO: 56)	402-414	Glu-C	R406	R406
		TVEMR*DGEVIK (SEQ ID NO: 98)	412-422	Trypsin	R416	R416
NRGN, human, Q92686		GPSPGGPGGAGVAR*GGAGGGPSGD (-) (SEQ ID NO: 71)	55-78	Trypsin, Lys-C	R68	R68
		RGRKGPGGPGGAGVAR*GGAGGG PSGD(-) (SEQ ID NO: 99)	51-78	Glu-C	R68	R68
NRGN, recombinant protein ^b		IQASFR*GHMAR*K (SEQ ID NO: 101)	33-44	Lys-C	R38, R43	
		GPSPGGPGGAGVAR*GGAGGGPSGD	55-80	Lys-C	R68, R80 ^b	
		TR*(-) (SEQ ID NO: 102)				
		R*GR*KGPSPGGPGGAGVAR*GGAG GGPSGDTR*(-) (SEQ ID NO: 103)	51-80	Glu-C	R51, R53 R68, R80 ^b	

^a sequence number includes the initiating Met at position 1 for all proteins

^b T79 and R80 are introduced into recombinant protein NRGN during subcloning

^c (-) indicates the C-terminal of this protein

TABLE 10

Summary of human brain tissue samples							
Samples	CERAD ^a	BRAAK			PMD		
		Stages	Age	Sex	Race	Score	From
Control 1		1	91	F	W	8	Occ ^b
Control 2		0	80	F	W	8	Occ
Control 3	1	2	68	F	W	12	Occ
AD 1	C	6	68	F	W	5.5	Occ

TABLE 10-continued

Summary of human brain tissue samples							
Samples	CERAD ^a	BRAAK			PMD		
		Stages	Age	Sex	Race	Score	From
AD 2	C	6	79	F	W	4	Occ
AD 3	C	6	92	F	W	7	Occ

^aCERAD: Consortium to establish a registry for Alzheimer's Disease^bOcc: occipital brain section

TABLE 11

In vivo citrullination sites of human MBP, GFAP, and NRG1										
Protein, UniProt ID	Peptide	Enzyme	In vivo site	Control ^a AD ^a						
				1	2	3	1	2	3	
mBp, human P02686-3	YLATASTMDHAR*HGFLPR (SEQ ID NO: 104)	Trypsin	R26	+	+		+	+		
	YLATASTMDHAR*HGFLPR* HR (SEQ ID NO: 106)	Trypsin	R26, R32				+	+		
	HR*DTGILDSIGR (SEQ ID NO: 108)	Trypsin	R34	+	+	+	+	+	+	
	DTGILDSIGR*FPGGDR (SEQ ID NO: 110)	Trypsin	R44	+	+		+	+		
	FPGGDR*GAPK (SEQ ID NO: 112)	Trypsin	R50	+	+		+	+	+	
	DSHHPAR*TAHYGSLPQK (SEQ ID NO: 77)	Trypsin, R92 Lys-C		+	+	+	+	+	+	
	SHGR*TQDENPVVHFFK (SEQ ID NO: 75)	Trypsin, R106 Lys-C		+	+	+	+	+		
	NIVTPR*TPPPSQGK (SEQ ID NO: 32)	Trypsin, R124 Lys-C		+	+	+	+	+	+	
	FSWGAEGQR*PGFGYGGGR (SEQ ID NO: 114)	Trypsin	R149	+	+		+	+		
	PGFGYGGGR*ASDYK (SEQ ID NO: 116)	Trypsin	R157	+	+		+	+	+	
	LGGR*DSR*SGSPMAR*R*(-) (SEQ ID NO: 68)	Trypsin, R186, R189 Lys-C	R196, R197	+		+	+	+	+	
	GFAP, Human P14136	R*LPGPTR*LSLAR (SEQ ID NO: 118)	Trypsin	R30, R36	+		+	+	+	
		LGPTR*LSLAR (SEQ ID NO: 120)	Trypsin	R36	+		+	+	+	
		FADLTDAAR*NAELLR (SEQ ID NO: 122)	Trypsin	R270					+	+
GHLKR*NIVVKTVE (SEQ ID NO: 30)		Glu-C	R406	+	+	+	+	+	+	
TVEMR*DGEVIK (SEQ ID NO: 124)		Trypsin	R416	+	+	+	+	+	+	
NRG1, Human Q92686	GPGPGGPGGAGVAR*GGAG GGPSGD(-) (SEQ ID NO: 71)	Trypsin, R68 Lys-C		+			+	+		
	RGRKGGPGGPGGAGVAR* GGAGGPGSGD(-) (SEQ ID NO: 99)	Glu-C	R68	+				+		

^a Refer to Table 2 for the details of brain tissue samples

TABLE 12

Citrullinated peptides of human GFAP after PAD2 treatment followed by trypsin, Lys-C or Glu-C digestion						
Protein, UniProt ID	Citrullinated peptide	Sequence number ^a	Theoretical m/z ^b	Experimental m/z	Enzyme	Residue
	LGPTR*LSLAR (SEQ ID NO: 120)	31-41	571.3380	571.3375	Trypsin	R36

TABLE 12-continued

Citrullinated peptides of human GFAP after PAD2 treatment followed by trypsin, Lys-C or Glu-C digestion						
Protein, UniProt ID	Citrullinated peptide	Sequence number ^a	Theoretical m/z ^b	Experimental m/z	Enzyme	Residue
	VR*FLEQQNK (SEQ ID NO: 40)	87-95	581.8144	581.8152	Lys-C	R88
	ALAAELNQLR*AK (SEQ ID NO: 42)	96-107	649.8750	649.8754	Lys-C	R105
	LR*LR*LDQLTANSAR*LE (SEQ ID NO: 44)	123-138	936.5129	936.5170	Glu-C	R124, R126, R136
	AENNLAAAYR*QEADE (SEQ ID NO: 46)	165-178	797.8526	797.8555	Glu-C	R173
	LQEQLAR*QQVHVE (SEQ ID NO: 48)	211-223	789.9154	789.9181	Glu-C	R217
	WYR*SKFADLTDAAR*N AE (SEQ ID NO: 50)	256-273	1043.9996	1043.9951	Glu-C	R258, R270
	FADLTDAAR*NAELLR (SEQ ID NO: 28)	261-276	874.4523	874.4540	Trypsin	R270
	FADLTDAAR*NAELLR* QAK (SEQ ID NO: 97)	261-279	1038.5396	1038.5411	Lys-C	R270, R276
	ANDYR*R*QLQSLTCDLE (SEQ ID NO: 52)	282-297	992.4575	992.4612	Glu-C	R286, R287
	LALDIEIATYR*K (SEQ ID NO: 54)	357-368	703.8981	703.8985	Lys-C	R367
	GHLKR*NIVVKTVE (SEQ ID NO: 30)	402-414	747.4435	747.4455	Glu-C	R406
	TVEMR*DGEVIK (SEQ ID NO: 124)	412-422	639.3239	639.3220	Trypsin	R416

^a sequence number includes the initiating Met at position 1^b mass-to-charge ratio of doubly charged ions

TABLE 13

Citrullinated peptides of bovine MBP after PAD treatment followed by trypsin or Lys-C digestion						
Protein, UniProt ID	Citrullinated peptide	Sequence number ^a	Theoretical m/z ^b	Experimental m/z	Enzyme	Residue
MBP, bovine P02687	YLASASTMDHAR*HGFLPR (SEQ ID NO: 84)	12-29	677.6670 (z: +3)	677.6676 (z: +3)	Trypsin	R23
	YLASASTMDHAR*HGFLPR*HR (SEQ ID NO: 58)	12-31	775.7150 (z: +3)	775.7166 (z: +3)	Trypsin	R23, R29
	YLASASTMDHAR*HGFLPR*HR * (SEQ ID NO: 60)	12-31	776.0430 (z: +3)	776.0431 (z: +3)	Trypsin	R23, R29, R31
	DTGILDSLGR*FFGSDR (SEQ ID NO: 14)	32-47	878.9287	878.9320	Trypsin	R41
	FFGSDR*GAPK (SEQ ID NO: 85)	42-51	541.7669	541.7678	Trypsin	R47
	R*GSGKDGHAAR*TTHYGSLP QK (SEQ ID NO: 63)	52-73	591.5454 (z: +4)	591.5466 (z: +4)	Lys-C	R52, R63
	DGHHAAR*TTHYGSLPQK (SEQ ID NO: 13)	57-73	938.9561	938.9566	Trypsin, Lys-C	R63
	AQGHR*PQDENPVVHFFK (SEQ ID NO: 65)	74-90	1003.9952	1003.9971	Trypsin, Lys-C	R78
	NIVTPR*TPPPSQGK ^c (SEQ ID NO: 32)	91-104	746.9095	746.9109	Trypsin, Lys-C	R96
	GR*GLSLSR*FSWGAEQK ^d (SEQ ID NO: 66)	105-121	919.4632	919.4651	Trypsin, Lys-C	R106, R112
	PGPGYGGR*ASDYK (SEQ ID NO: 11)	122-134	688.3181	688.3184	Trypsin, Lys-C	R129
	LGGR*DSR*SGSPMA (-) ^e (SEQ ID NO: 87)	155-167	646.7986	646.7998	Trypsin, Lys-C	R158, R161
	LGGR*DSR*SGSPMAR* (-) (SEQ ID NO: 89)	155-168	725.3411	725.3438	Trypsin, Lys-C	R158, R161, R168

TABLE 13-continued

Citrullinated peptides of bovine MBP after PAD treatment followed by trypsin or Lys-C digestion						
Protein, UniProt ID	Citrullinated peptide	Sequence number ^a	Theoretical m/z ^b	Experimental m/z	Enzyme	Residue
	LGGR*DSR*SGSPMAR*R*(-) (SEQ ID NO: 83)	155-169	803.8837	803.8842	Trypsin, Lys-C	R158, R161, R168, R169

^a sequence number does not include the initiating Met at position 1^b charge state of precursor ion is +2 unless otherwise specified^c peptides with or without phosphorylation at T97 were both detected^d peptides with mono- and di-methylation at residue R106 were observed^e (-) indicates the C-terminal of this protein

TABLE 14

Citrullinated peptides and related unmodified peptide of neurogranin						
Protein, UniProt ID	Peptide Sequence	Seq. Number	^a Enzyme	Theoretical m/z ^b	Exp. m/z	Residues
NRGN, bovine P35722	GPGPGGPGGAGGARGGAGGG	55-78	Lys-C	888.9040	888.9058	NA ^c
	PSGD(-) (SEQ ID NO: 21)					
	GPGPGGPGGAGGAR*GGAGG	55-78	Lys-C	889.3960	889.3983	R68
	GPSGD(-) (SEQ ID NO: 22)					
NRGN, human Q92686	GPGPGGPGGAGGAR	55-68	Trypsin	532.7652	532.7656	NA
	(SEQ ID NO: 23)					
	KGPGPGGPGGAGGAR*GGAG	54-78	Trypsin	953.4435	953.4429	R68
	GGPSGD(-) (SEQ ID NO: 23)					
NRGN, human Q92686	GPGPGGPGGAGVARGGAGGG	54-78	Lys-C	909.9275	909.9316	NA
	PSGD(-) (SEQ ID NO: 24)					
	GPGPGGPGGAGVAR*GGAGG	55-78	Trypsin,	910.4195	910.4236	R68
	GPSGD(-) (SEQ ID NO: 73)		Lys-C			
	RGRKGPGGPGGAGVARGG	51-78	Glu-C	772.7270	772.7285	NA
	AGGGPSGD(-) (SEQ ID NO: 71)			(z: +3)	(z: +3)	
	RGRKGPGGPGGAGVAR*G	51-78	Glu-C	773.0550	773.0557	R68
GAGGGPSGD(-) (SEQ ID NO: 100)			(z: +3)	(z: +3)		
NRGN, recombinant protein	IQASFR*GHMAR*K	33-44	Lys-C	702.3642	702.3629	R38, R43
	(SEQ ID NO: 101)					
	GPGPGGPGGAGVAR*GGAGG	55-80	Lys-C	1039.4859	1039.4839	R68, R80
	GPSGDTR*(-) (SEQ ID NO: 102)					
NRGN, recombinant protein	R*GR*KGPGPGGAGVAR*	51-80	Glu-C	859.7553	859.7544	R51, R53,
	GGAGGGPSGDTR*(-) (SEQ ID NO: 103)			(z: +3)	(z: +3)	R68, R80 ^b

^a sequence number includes the initiating Met at position 1. Met is cleaved from in the mature protein.^b mass-to-charge ratio of the peptide^c NA, not applicable for unmodified peptides

TABLE 15

In vivo citrullinated peptides of human MBP						
Protein, UniProt ID	Citrullinated peptide	Sequence number ^a	Theoretical m/z ^b	Experimental m/z	Enzyme	Residues
MBP, human P02686-3	YLATASTMDHAR*HGFLPR	15-32	1023.0047	1023.0049	Trypsin	R26
	(SEQ ID NO: 104)					
	YLATASTMDHAR*HGFLPR*HR	15-34	780.3869	780.3881	Trypsin	R26, R32
	(SEQ ID NO: 106)		(z: +3)	(z: +3)		
	HR*DTGILDSIGR	33-44	670.8495	670.8501	Trypsin	R34
	(SEQ ID NO: 108)					
	DTGILDSIGR*FFGGDR	35-50	863.9234	863.9266	Trypsin	R44
	(SEQ ID NO: 110)					
	FFGGDR*GAPK	45-54	526.7616	526.7618	Trypsin	R50
	(SEQ ID NO: 112)					
	DSHHPAR*TAHYGSLPQK	86-102	951.9639	951.9655	Trypsin,	R92
	(SEQ ID NO: 77)				Lys-C	

TABLE 15-continued

In vivo citrullinated peptides of human MBP						
Protein, UniProt ID	Citrullinated peptide	Sequence number ^a	Theoretical m/z ^b	Experimental m/z	Enzyme	Residues
	SHGR*TQDENPVVHFFK (SEQ ID NO: 75)	103-118	949.9608	949.9587	Trypsin, Lys-C	R106
	NIVTPR*TPPPSQGK (SEQ ID NO: 75)	119-132	746.9095	746.9106	Trypsin, Lys-C	R124
	FSWGAEGQR*PGFGYGGR (SEQ ID NO: 114)	141-157	915.4213	915.4208	Trypsin	R149
	PGFGYGGR*ASDYK (SEQ ID NO: 116)	150-162	688.3184	688.3177	Trypsin	R157
	LGGR*DSR*SGSPMAR*R* (SEQ ID NO: 68)	783-197	803.8837	803.8853	Trypsin, Lys-C	R186, R189 R196, R197

^a sequence number does not include the initiating Met at position 1

^b charge state of precursor ion is +2 unless otherwise specified

TABLE 16

In vivo citrullination sites of human MBP and GFAP in brain tissue				
Protein (UniProt ID)	Peptide	Enzyme	In vivo site	Brain Sample ¹
MBP, human (P02686-3)	YLATASTMDHAR*HGFLPR (SEQ ID NO: 104)	Trypsin	R26	C, S
	HR*DTGILDSIGR (SEQ ID NO: 108)	Trypsin	R34	C, S
	FFGGDR*GAPK (SEQ ID NO: 112)	Trypsin	R50	C, S
	DSHHPAR*TAHYGSLPQK (SEQ ID NO: 77)	Trypsin, Lys-C	R92	C, S
	SHGR*TQDENPVVHFFK (SEQ ID NO: 75)	Trypsin, Lys-C	R106	C, S
	NIVTPR*TPPPSQGK (SEQ ID NO: 32)	Trypsin, Lys-C	R124	C, S
	FSWGAEGQR*PGFGYGGR (SEQ ID NO: 114)	Trypsin	R149	C, S
	PGFGYGGR*ASDYK (SEQ ID NO: 116)	Trypsin	R157	C, S
	LGGR*DSR*SGSPMAR*R* (SEQ ID NO: 68)	Lys-C	R186 R189 R196 R197	C, S C, S C, S C, S
	GFAP, Human (P14136)	R*LGPGR*LSLAR (SEQ ID NO: 118)	Trypsin	R30 R36
GHLKR*NIVVKTVE (SEQ ID NO: 30)		Glu-C	R406	C, S,
TVEMR*DGEVIK (SEQ ID NO: 124)		Trypsin	R416	C, S,
NRGN, Human (Q92686)		IQASFR (SEQ ID NO: 126)	Trypsin	NA
	GPGGGGGGAGVAR (SEQ ID NO: 70)	Trypsin	NA	C, S
	KGPGGGGGAGVAR (SEQ ID NO: 127)	Trypsin	NA	C, S
	GPGGGGGGAGVARGGAGG	Lys-C	NA	C, S
	GPSGD (SEQ ID NO: 73)			
	RGRKGGGGGGAGVARG GAGGGPSGD (SEQ ID NO:)	Glu-C	NA	C, S

Brain tissue sample from one patient. S, brain tissue from stroke region; C: brain tissue from normal region. Extent of modification may differ between control and disease.

TABLE 17

In vivo citrullination sites of other proteins in human brain tissue (not patented yet)						
Proteins	UniProt ID	Peptide	start	stop	Residue	Brain Sample ¹
Tubulin beta-4B chain	P68371	IRREYPPDR*IMNTF (SEQ ID NO: 128)	155	167	R162	S
Tubulin alpha-1B chain	P68363	YMACLLLYR*GDVVPK (SEQ ID NO: 130)	312	326	R320	C, S
		VR*TGTYR*QLFHPE (SEQ ID NO: 132)	78	90	R79, R84	S
CNPase	P09543	STLAR*VIVDK (SEQ ID NO: 134)	64	73	R68	C, S
		ITPGAR*GAFSEEYK (SEQ ID NO: 136)	88	101	R93	C, S
PPIA	P62937	TAENFR*ALSTGEK (SEQ ID NO: 138)	32	44	R37	C, S
Septin-7	Q16181	ILEQQNSSR*TLEK (SEQ ID NO: 140)	417	429	R425	C, S
Elongation factor 1-alpha 2	Q05639	PLR*LPLQDVYK (SEQ ID NO: 142)	245	255	R247	C, S
		VYKIGGIGTVPVGR*VE (SEQ ID NO: 144)	253	268	R266	C, S
TPPP	O94811	AISSPTVSR*LTDTTK (SEQ ID NO: 146)	157	171	R165	C, S
TPPP3	Q9BW30	TGGAVDRLTDTSR*YTGSHK (SEQ ID NO: 148)	118	136	R130	S
		GIAGR*QDILDDSGYVSAYK (SEQ ID NO: 150)	147	165	R151	S
Ermin, Isoform 2	Q8TAM6-2	LTDVDSPLPHYR*VEPSLE (SEQ ID NO: 152)	46	63	R57	C
NDRG2, Isoform 2	Q9UN36-2	TASLTSAAASVDGNR*SR (SEQ ID NO: 154)	316	331	R329	S

Brain tissue sample from one patient. S, brain tissue from stroke region; C, brain tissue from normal region.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 162

<210> SEQ ID NO 1

<211> LENGTH: 78

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<220> FEATURE:

<221> NAME/KEY: MISC_FEATURE

<222> LOCATION: (1)..(78)

<223> OTHER INFORMATION: Neurogranin (NRGN) amino acid sequence

<400> SEQUENCE: 1

Met Asp Cys Cys Thr Glu Asn Ala Cys Ser Lys Pro Asp Asp Asp Ile
1 5 10 15

Leu Asp Ile Pro Leu Asp Asp Pro Gly Ala Asn Ala Ala Ala Lys
20 25 30

Ile Gln Ala Ser Phe Arg Gly His Met Ala Arg Lys Lys Ile Lys Ser
35 40 45

Gly Glu Arg Gly Arg Lys Gly Pro Gly Pro Gly Gly Pro Gly Ala
50 55 60

Gly Val Ala Arg Gly Gly Ala Gly Gly Pro Ser Gly Asp

-continued

<220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (1)..(432)
 <223> OTHER INFORMATION: Glial Fibrillary Acidic Protein (GFAP) amino acid sequence

<400> SEQUENCE: 3

Met Glu Arg Arg Arg Ile Thr Ser Ala Ala Arg Arg Ser Tyr Val Ser
 1 5 10 15

Ser Gly Glu Met Met Val Gly Gly Leu Ala Pro Gly Arg Arg Leu Gly
 20 25 30

Pro Gly Thr Arg Leu Ser Leu Ala Arg Met Pro Pro Pro Leu Pro Thr
 35 40 45

Arg Val Asp Phe Ser Leu Ala Gly Ala Leu Asn Ala Gly Phe Lys Glu
 50 55 60

Thr Arg Ala Ser Glu Arg Ala Glu Met Met Glu Leu Asn Asp Arg Phe
 65 70 75 80

Ala Ser Tyr Ile Glu Lys Val Arg Phe Leu Glu Gln Gln Asn Lys Ala
 85 90 95

Leu Ala Ala Glu Leu Asn Gln Leu Arg Ala Lys Glu Pro Thr Lys Leu
 100 105 110

Ala Asp Val Tyr Gln Ala Glu Leu Arg Glu Leu Arg Leu Arg Leu Asp
 115 120 125

Gln Leu Thr Ala Asn Ser Ala Arg Leu Glu Val Glu Arg Asp Asn Leu
 130 135 140

Ala Gln Asp Leu Ala Thr Val Arg Gln Lys Leu Gln Asp Glu Thr Asn
 145 150 155 160

Leu Arg Leu Glu Ala Glu Asn Asn Leu Ala Ala Tyr Arg Gln Glu Ala
 165 170 175

Asp Glu Ala Thr Leu Ala Arg Leu Asp Leu Glu Arg Lys Ile Glu Ser
 180 185 190

Leu Glu Glu Glu Ile Arg Phe Leu Arg Lys Ile His Glu Glu Glu Val
 195 200 205

Arg Glu Leu Gln Glu Gln Leu Ala Arg Gln Gln Val His Val Glu Leu
 210 215 220

Asp Val Ala Lys Pro Asp Leu Thr Ala Ala Leu Lys Glu Ile Arg Thr
 225 230 235 240

Gln Tyr Glu Ala Met Ala Ser Ser Asn Met His Glu Ala Glu Glu Trp
 245 250 255

Tyr Arg Ser Lys Phe Ala Asp Leu Thr Asp Ala Ala Ala Arg Asn Ala
 260 265 270

Glu Leu Leu Arg Gln Ala Lys His Glu Ala Asn Asp Tyr Arg Arg Gln
 275 280 285

Leu Gln Ser Leu Thr Cys Asp Leu Glu Ser Leu Arg Gly Thr Asn Glu
 290 295 300

Ser Leu Glu Arg Gln Met Arg Glu Gln Glu Glu Arg His Val Arg Glu
 305 310 315 320

Ala Ala Ser Tyr Gln Glu Ala Leu Ala Arg Leu Glu Glu Glu Gly Gln
 325 330 335

Ser Leu Lys Asp Glu Met Ala Arg His Leu Gln Glu Tyr Gln Asp Leu
 340 345 350

Leu Asn Val Lys Leu Ala Leu Asp Ile Glu Ile Ala Thr Tyr Arg Lys
 355 360 365

-continued

Leu Leu Glu Gly Glu Glu Asn Arg Ile Thr Ile Pro Val Gln Thr Phe
 370 375 380

Ser Asn Leu Gln Ile Arg Glu Thr Ser Leu Asp Thr Lys Ser Val Ser
 385 390 395 400

Glu Gly His Leu Lys Arg Asn Ile Val Val Lys Thr Val Glu Met Arg
 405 410 415

Asp Gly Glu Val Ile Lys Glu Ser Lys Gln Glu His Lys Asp Val Met
 420 425 430

<210> SEQ ID NO 4
 <211> LENGTH: 665
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (1)..(665)
 <223> OTHER INFORMATION: Protein-arginine deiminase type-2 (PAD2) amino
 acid sequence

<400> SEQUENCE: 4

Met Leu Arg Glu Arg Thr Val Arg Leu Gln Tyr Gly Ser Arg Val Glu
 1 5 10 15

Ala Val Tyr Val Leu Gly Thr Tyr Leu Trp Thr Asp Val Tyr Ser Ala
 20 25 30

Ala Pro Ala Gly Ala Gln Thr Phe Ser Leu Lys His Ser Glu His Val
 35 40 45

Trp Val Glu Val Val Arg Asp Gly Glu Ala Glu Glu Val Ala Thr Asn
 50 55 60

Gly Lys Gln Arg Trp Leu Leu Ser Pro Ser Thr Thr Leu Arg Val Thr
 65 70 75 80

Met Ser Gln Ala Ser Thr Glu Ala Ser Ser Asp Lys Val Thr Val Asn
 85 90 95

Tyr Tyr Asp Glu Glu Gly Ser Ile Pro Ile Asp Gln Ala Gly Leu Phe
 100 105 110

Leu Thr Ala Ile Glu Ile Ser Leu Asp Val Asp Ala Asp Arg Asp Gly
 115 120 125

Val Val Glu Lys Asn Asn Pro Lys Lys Ala Ser Trp Thr Trp Gly Pro
 130 135 140

Glu Gly Gln Gly Ala Ile Leu Leu Val Asn Cys Asp Arg Glu Thr Pro
 145 150 155 160

Trp Leu Pro Lys Glu Asp Cys Arg Asp Glu Lys Val Tyr Ser Lys Glu
 165 170 175

Asp Leu Lys Asp Met Ser Gln Met Ile Leu Arg Thr Lys Gly Pro Asp
 180 185 190

Arg Leu Pro Ala Gly Tyr Glu Ile Val Leu Tyr Ile Ser Met Ser Asp
 195 200 205

Ser Asp Lys Val Gly Val Phe Tyr Val Glu Asn Pro Phe Phe Gly Gln
 210 215 220

Arg Tyr Ile His Ile Leu Gly Arg Arg Lys Leu Tyr His Val Val Lys
 225 230 235 240

Tyr Thr Gly Gly Ser Ala Glu Leu Leu Phe Phe Val Glu Gly Leu Cys
 245 250 255

Phe Pro Asp Glu Gly Phe Ser Gly Leu Val Ser Ile His Val Ser Leu
 260 265 270

-continued

Leu Glu Tyr Met Ala Gln Asp Ile Pro Leu Thr Pro Ile Phe Thr Asp
 275 280 285
 Thr Val Ile Phe Arg Ile Ala Pro Trp Ile Met Thr Pro Asn Ile Leu
 290 295 300
 Pro Pro Val Ser Val Phe Val Cys Cys Met Lys Asp Asn Tyr Leu Phe
 305 310 315 320
 Leu Lys Glu Val Lys Asn Leu Val Glu Lys Thr Asn Cys Glu Leu Lys
 325 330 335
 Val Cys Phe Gln Tyr Leu Asn Arg Gly Asp Arg Trp Ile Gln Asp Glu
 340 345 350
 Ile Glu Phe Gly Tyr Ile Glu Ala Pro His Lys Gly Phe Pro Val Val
 355 360 365
 Leu Asp Ser Pro Arg Asp Gly Asn Leu Lys Asp Phe Pro Val Lys Glu
 370 375 380
 Leu Leu Gly Pro Asp Phe Gly Tyr Val Thr Arg Glu Pro Leu Phe Glu
 385 390 395 400
 Ser Val Thr Ser Leu Asp Ser Phe Gly Asn Leu Glu Val Ser Pro Pro
 405 410 415
 Val Thr Val Asn Gly Lys Thr Tyr Pro Leu Gly Arg Ile Leu Ile Gly
 420 425 430
 Ser Ser Phe Pro Leu Ser Gly Gly Arg Arg Met Thr Lys Val Val Arg
 435 440 445
 Asp Phe Leu Lys Ala Gln Gln Val Gln Ala Pro Val Glu Leu Tyr Ser
 450 455 460
 Asp Trp Leu Thr Val Gly His Val Asp Glu Phe Met Ser Phe Val Pro
 465 470 475 480
 Ile Pro Gly Thr Lys Lys Phe Leu Leu Leu Met Ala Ser Thr Ser Ala
 485 490 495
 Cys Tyr Lys Leu Phe Arg Glu Lys Gln Lys Asp Gly His Gly Glu Ala
 500 505 510
 Ile Met Phe Lys Gly Leu Gly Gly Met Ser Ser Lys Arg Ile Thr Ile
 515 520 525
 Asn Lys Ile Leu Ser Asn Glu Ser Leu Val Gln Glu Asn Leu Tyr Phe
 530 535 540
 Gln Arg Cys Leu Asp Trp Asn Arg Asp Ile Leu Lys Lys Glu Leu Gly
 545 550 555 560
 Leu Thr Glu Gln Asp Ile Ile Asp Leu Pro Ala Leu Phe Lys Met Asp
 565 570 575
 Glu Asp His Arg Ala Arg Ala Phe Phe Pro Asn Met Val Asn Met Ile
 580 585 590
 Val Leu Asp Lys Asp Leu Gly Ile Pro Lys Pro Phe Gly Pro Gln Val
 595 600 605
 Glu Glu Glu Cys Cys Leu Glu Met His Val Arg Gly Leu Leu Glu Pro
 610 615 620
 Leu Gly Leu Glu Cys Thr Phe Ile Asp Asp Ile Ser Ala Tyr His Lys
 625 630 635 640
 Phe Leu Gly Glu Val His Cys Gly Thr Asn Val Arg Arg Lys Pro Phe
 645 650 655
 Thr Phe Lys Trp Trp His Met Val Pro
 660 665

-continued

<210> SEQ ID NO 5
 <211> LENGTH: 169
 <212> TYPE: PRT
 <213> ORGANISM: Bos taurus
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (1)..(169)
 <223> OTHER INFORMATION: Bovine Myelin Basic Protein (MBP) amino acid sequence

<400> SEQUENCE: 5

Ala Ala Gln Lys Arg Pro Ser Gln Arg Ser Lys Tyr Leu Ala Ser Ala
 1 5 10 15
 Ser Thr Met Asp His Ala Arg His Gly Phe Leu Pro Arg His Arg Asp
 20 25 30
 Thr Gly Ile Leu Asp Ser Leu Gly Arg Phe Phe Gly Ser Asp Arg Gly
 35 40 45
 Ala Pro Lys Arg Gly Ser Gly Lys Asp Gly His His Ala Ala Arg Thr
 50 55 60
 Thr His Tyr Gly Ser Leu Pro Gln Lys Ala Gln Gly His Arg Pro Gln
 65 70 75 80
 Asp Glu Asn Pro Val Val His Phe Phe Lys Asn Ile Val Thr Pro Arg
 85 90 95
 Thr Pro Pro Pro Ser Gln Gly Lys Gly Arg Gly Leu Ser Leu Ser Arg
 100 105 110
 Phe Ser Trp Gly Ala Glu Gly Gln Lys Pro Gly Phe Gly Tyr Gly Gly
 115 120 125
 Arg Ala Ser Asp Tyr Lys Ser Ala His Lys Gly Leu Lys Gly His Asp
 130 135 140
 Ala Gln Gly Thr Leu Ser Lys Ile Phe Lys Leu Gly Gly Arg Asp Ser
 145 150 155 160
 Arg Ser Gly Ser Pro Met Ala Arg Arg
 165

<210> SEQ ID NO 6
 <211> LENGTH: 78
 <212> TYPE: PRT
 <213> ORGANISM: Bos taurus
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (1)..(78)
 <223> OTHER INFORMATION: Bovine Neurogranin (NRGN) amino acid sequence

<400> SEQUENCE: 6

Met Asp Cys Cys Thr Glu Ser Ala Cys Ser Lys Pro Asp Asp Asp Ile
 1 5 10 15
 Leu Asp Ile Pro Leu Asp Asp Pro Gly Ala Asn Ala Ala Ala Ala Lys
 20 25 30
 Ile Gln Ala Ser Phe Arg Gly His Met Ala Arg Lys Lys Ile Lys Ser
 35 40 45
 Gly Glu Arg Gly Arg Lys Gly Pro Gly Pro Gly Gly Pro Gly Ala
 50 55 60
 Gly Gly Ala Arg Gly Gly Ala Gly Gly Gly Pro Ser Gly Asp
 65 70 75

<210> SEQ ID NO 7
 <211> LENGTH: 428

-continued

```

<212> TYPE: PRT
<213> ORGANISM: Bos taurus
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (1)..(428)
<223> OTHER INFORMATION: Bovine Glial Fibrillary Acidic Protein (GFAP)
        amino acid sequence

```

```

<400> SEQUENCE: 7

```

```

Met Glu Arg Arg Arg Val Thr Ser Ala Thr Arg Arg Ser Tyr Val Ser
1           5           10           15

Ser Ser Glu Met Val Val Gly Gly Arg Arg Leu Gly Pro Gly Thr Arg
20           25           30

Leu Ser Leu Ala Arg Met Pro Pro Pro Leu Pro Ala Arg Val Asp Phe
35           40           45

Ser Leu Ala Gly Ala Leu Asn Ser Gly Phe Lys Glu Thr Arg Ala Ser
50           55           60

Glu Arg Ala Glu Met Met Glu Leu Asn Asp Arg Phe Ala Ser Tyr Ile
65           70           75           80

Glu Lys Val Arg Phe Leu Glu Gln Gln Asn Lys Ala Leu Ala Ala Glu
85           90           95

Leu Asn Gln Leu Arg Ala Lys Glu Pro Thr Lys Leu Ala Asp Val Tyr
100          105          110

Gln Ala Glu Leu Arg Glu Leu Arg Leu Arg Leu Asp Gln Leu Thr Ala
115          120          125

Asn Ser Ala Arg Leu Glu Val Glu Arg Asp Asn Leu Ala Gln Asp Leu
130          135          140

Gly Thr Leu Arg Gln Lys Leu Gln Asp Glu Thr Asn Gln Arg Leu Glu
145          150          155          160

Ala Glu Asn Asn Leu Ala Ala Tyr Arg Gln Glu Ala Asp Glu Ala Thr
165          170          175

Leu Ala Arg Leu Asp Leu Glu Arg Lys Ile Glu Ser Leu Glu Glu Glu
180          185          190

Ile Arg Phe Leu Arg Lys Ile His Glu Glu Glu Val Arg Glu Leu Gln
195          200          205

Glu Gln Leu Ala Gln Gln Gln Val His Val Glu Met Asp Val Ala Lys
210          215          220

Pro Asp Leu Thr Ala Ala Leu Arg Glu Ile Arg Thr Gln Tyr Glu Ala
225          230          235          240

Val Ala Ser Ser Asn Met His Glu Ala Glu Glu Trp Tyr Arg Ser Lys
245          250          255

Phe Ala Asp Leu Asn Asp Ala Ala Ala Arg Asn Ala Glu Leu Leu Arg
260          265          270

Gln Ala Lys His Glu Ala Asn Asp Tyr Arg Arg Gln Leu Gln Ala Leu
275          280          285

Thr Cys Asp Leu Glu Ser Leu Arg Gly Thr Asn Glu Ser Leu Glu Arg
290          295          300

Gln Met Arg Glu Gln Glu Glu Arg His Ala Arg Glu Ala Ala Ser Tyr
305          310          315          320

Gln Glu Ala Leu Ala Arg Leu Glu Glu Glu Gly Gln Ser Leu Lys Asp
325          330          335

Glu Met Ala Arg His Leu Gln Glu Tyr Gln Asp Leu Leu Asn Val Lys
340          345          350

```


-continued

<210> SEQ ID NO 12
<211> LENGTH: 17
<212> TYPE: PRT
<213> ORGANISM: Bos taurus
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (1)..(17)
<223> OTHER INFORMATION: Bovine MBP peptide, amino acids 57-73

<400> SEQUENCE: 12

Asp Gly His His Ala Ala Arg Thr Thr His Tyr Gly Ser Leu Pro Gln
1 5 10 15

Lys

<210> SEQ ID NO 13
<211> LENGTH: 17
<212> TYPE: PRT
<213> ORGANISM: Bos taurus
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (7)..(7)
<223> OTHER INFORMATION: Bovine MBP peptide, amino acids 57-73, 7th
residue citrullinated (Arg63)

<400> SEQUENCE: 13

Asp Gly His His Ala Ala Arg Thr Thr His Tyr Gly Ser Leu Pro Gln
1 5 10 15

Lys

<210> SEQ ID NO 14
<211> LENGTH: 16
<212> TYPE: PRT
<213> ORGANISM: Bos taurus
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: Bovine MBP peptide, amino acids 32-47, 10th
residue citrullinated (Arg41)

<400> SEQUENCE: 14

Asp Thr Gly Ile Leu Asp Ser Leu Gly Arg Phe Phe Gly Ser Asp Arg
1 5 10 15

<210> SEQ ID NO 15
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: Bos taurus
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: Bovine MBP peptide, amino acids 32-51, 10th
residue citrullinated (Arg41)

<220> FEATURE:

<221> NAME/KEY: MOD_RES

<222> LOCATION: (16)..(16)

<223> OTHER INFORMATION: Bovine MBP peptide, amino acids 32-51, 16th
residue citrullinated (Arg47)

<400> SEQUENCE: 15

Asp Thr Gly Ile Leu Asp Ser Leu Gly Arg Phe Phe Gly Ser Asp Arg
1 5 10 15

Gly Ala Pro Lys
20

-continued

<210> SEQ ID NO 16
<211> LENGTH: 6
<212> TYPE: PRT
<213> ORGANISM: Bos taurus
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (1)..(6)
<223> OTHER INFORMATION: Bovine MBP peptide, amino acids 91-96

<400> SEQUENCE: 16

Asn Ile Val Thr Pro Arg
1 5

<210> SEQ ID NO 17
<211> LENGTH: 8
<212> TYPE: PRT
<213> ORGANISM: Bos taurus
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (1)..(8)
<223> OTHER INFORMATION: Bovine MBP peptide, amino acids 122-129

<400> SEQUENCE: 17

Pro Gly Phe Gly Tyr Gly Gly Arg
1 5

<210> SEQ ID NO 18
<211> LENGTH: 17
<212> TYPE: PRT
<213> ORGANISM: Bos taurus
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (1)..(17)
<223> OTHER INFORMATION: Bovine MBP peptide, amino acids 105-121

<400> SEQUENCE: 18

Gly Arg Gly Leu Ser Leu Ser Arg Phe Ser Trp Gly Ala Glu Gly Gln
1 5 10 15

Lys

<210> SEQ ID NO 19
<211> LENGTH: 17
<212> TYPE: PRT
<213> ORGANISM: Bos taurus
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (2)..(2)
<223> OTHER INFORMATION: Bovine MBP peptide, amino acids 101-121, 2nd
residue citrullinated (Arg106)
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Bovine MBP peptide, amino acids 101-121, 8th
residue citrullinated (Arg112)

<400> SEQUENCE: 19

Gly Arg Gly Leu Ser Leu Ser Arg Phe Ser Trp Gly Ala Glu Gly Gln
1 5 10 15

Lys

<210> SEQ ID NO 20
<211> LENGTH: 17
<212> TYPE: PRT
<213> ORGANISM: Bos taurus
<220> FEATURE:

-continued

<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Bovine MBP peptide, amino acids 105-121), 8th
residue citrullinated (Arg112)

<400> SEQUENCE: 20

Gly Arg Gly Leu Ser Leu Ser Arg Phe Ser Trp Gly Ala Glu Gly Gln
1 5 10 15

Lys

<210> SEQ ID NO 21
<211> LENGTH: 24
<212> TYPE: PRT
<213> ORGANISM: Bos taurus
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (1)..(24)
<223> OTHER INFORMATION: Bovine Neurogranin (NRGN) peptide, amino acids
55-78

<400> SEQUENCE: 21

Gly Pro Gly Pro Gly Gly Pro Gly Gly Ala Gly Gly Ala Arg Gly Gly
1 5 10 15

Ala Gly Gly Gly Pro Ser Gly Asp
20

<210> SEQ ID NO 22
<211> LENGTH: 24
<212> TYPE: PRT
<213> ORGANISM: Bos taurus
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (14)..(14)
<223> OTHER INFORMATION: Bovine NRGN peptide, amino acids 55-78, 14th
residue citrullinated

<400> SEQUENCE: 22

Gly Pro Gly Pro Gly Gly Pro Gly Gly Ala Gly Gly Ala Arg Gly Gly
1 5 10 15

Ala Gly Gly Gly Pro Ser Gly Asp
20

<210> SEQ ID NO 23
<211> LENGTH: 14
<212> TYPE: PRT
<213> ORGANISM: Bos taurus
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (1)..(14)
<223> OTHER INFORMATION: Bovine NRGN peptide, amino acids 55-68

<400> SEQUENCE: 23

Gly Pro Gly Pro Gly Gly Pro Gly Gly Ala Gly Gly Ala Arg
1 5 10

<210> SEQ ID NO 24
<211> LENGTH: 25
<212> TYPE: PRT
<213> ORGANISM: Bos taurus
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (15)..(15)
<223> OTHER INFORMATION: Bovine NRGN peptide, amino acids 54-78, 15th
residue citrullinated (Arg68)

-continued

<400> SEQUENCE: 24

Lys Gly Pro Gly Pro Gly Gly Pro Gly Gly Ala Gly Gly Ala Arg Gly
 1 5 10 15
 Gly Ala Gly Gly Gly Pro Ser Gly Asp
 20 25

<210> SEQ ID NO 25
 <211> LENGTH: 13
 <212> TYPE: PRT
 <213> ORGANISM: Bos taurus
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (1)..(13)
 <223> OTHER INFORMATION: Bovine GFAP peptide, amino acids 398-410

<400> SEQUENCE: 25

Gly His Leu Lys Arg Asn Ile Val Val Lys Thr Val Glu
 1 5 10

<210> SEQ ID NO 26
 <211> LENGTH: 13
 <212> TYPE: PRT
 <213> ORGANISM: Bos taurus
 <220> FEATURE:
 <221> NAME/KEY: MOD_RES
 <222> LOCATION: (5)..(5)
 <223> OTHER INFORMATION: Bovine GFAP peptide, amino acids 398-410, 5th
 residue citrullinated (Arg402)

<400> SEQUENCE: 26

Gly His Leu Lys Arg Asn Ile Val Val Lys Thr Val Glu
 1 5 10

<210> SEQ ID NO 27
 <211> LENGTH: 19
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (1)..(19)
 <223> OTHER INFORMATION: Human GFAP peptide, amino acids 261-279

<400> SEQUENCE: 27

Phe Ala Asp Leu Thr Asp Ala Ala Ala Arg Asn Ala Glu Leu Leu Arg
 1 5 10 15

Gln Ala Lys

<210> SEQ ID NO 28
 <211> LENGTH: 19
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens
 <220> FEATURE:
 <221> NAME/KEY: MOD_RES
 <222> LOCATION: (10)..(10)
 <223> OTHER INFORMATION: Human GFAP peptide, amino acids 261-279, 10th
 residue citrullinated (Arg270)

<400> SEQUENCE: 28

Phe Ala Asp Leu Thr Asp Ala Ala Ala Arg Asn Ala Glu Leu Leu Arg
 1 5 10 15

Gln Ala Lys

<210> SEQ ID NO 29

-continued

<211> LENGTH: 13
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (1)..(13)
<223> OTHER INFORMATION: Human GFAP peptide, amino acids 402-414

<400> SEQUENCE: 29

Gly His Leu Lys Arg Asn Ile Val Val Lys Thr Val Glu
1 5 10

<210> SEQ ID NO 30
<211> LENGTH: 13
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (5)..(5)
<223> OTHER INFORMATION: Human GFAP peptide, amino acids 402-414, 5th
residue citrullinated (Arg406)

<400> SEQUENCE: 30

Gly His Leu Lys Arg Asn Ile Val Val Lys Thr Val Glu
1 5 10

<210> SEQ ID NO 31
<211> LENGTH: 14
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (1)..(14)
<223> OTHER INFORMATION: Human MBP peptide, amino acids 92-105

<400> SEQUENCE: 31

Asn Ile Val Thr Pro Arg Thr Pro Pro Pro Ser Gln Gly Lys
1 5 10

<210> SEQ ID NO 32
<211> LENGTH: 14
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (6)..(6)
<223> OTHER INFORMATION: Human MBP peptide, amino acids 92-105, 6th
residue citrullinated (Arg97)

<400> SEQUENCE: 32

Asn Ile Val Thr Pro Arg Thr Pro Pro Pro Ser Gln Gly Lys
1 5 10

<210> SEQ ID NO 33
<211> LENGTH: 14
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (6)..(6)
<223> OTHER INFORMATION: Human MBP peptide, amino acids 226-239, 6th
residue citrullinated (Arg231)

<400> SEQUENCE: 33

Asn Ile Val Thr Pro Arg Thr Pro Pro Pro Ser Gln Gly Lys
1 5 10

-continued

<210> SEQ ID NO 34
 <211> LENGTH: 14
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (1)..(14)
 <223> OTHER INFORMATION: Human MBP peptide, amino acids 226-249

<400> SEQUENCE: 34

Asn Ile Val Thr Pro Arg Thr Pro Pro Pro Ser Gln Gly Lys
 1 5 10

<210> SEQ ID NO 35
 <211> LENGTH: 12
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens
 <220> FEATURE:
 <221> NAME/KEY: MOD_RES
 <222> LOCATION: (6)..(6)
 <223> OTHER INFORMATION: Human NRGN peptide, amino acids 33-44, 6th
 residue citrullinated (Arg38)
 <220> FEATURE:
 <221> NAME/KEY: MOD_RES
 <222> LOCATION: (11)..(11)
 <223> OTHER INFORMATION: Human NRGN peptide, amino acids 33-44, 11th
 residue citrullinated (Arg43)

<400> SEQUENCE: 35

Ile Gln Ala Ser Phe Arg Gly His Met Ala Arg Lys
 1 5 10

<210> SEQ ID NO 36
 <211> LENGTH: 12
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (1)..(12)
 <223> OTHER INFORMATION: Human NRGN peptide, amino acids 33-44

<400> SEQUENCE: 36

Ile Gln Ala Ser Phe Arg Gly His Met Ala Arg Lys
 1 5 10

<210> SEQ ID NO 37
 <211> LENGTH: 26
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens
 <220> FEATURE:
 <221> NAME/KEY: MOD_RES
 <222> LOCATION: (14)..(14)
 <223> OTHER INFORMATION: Human NRGN peptide, amino acids 55-80, 14th
 residue citrullinated (Arg68)
 <220> FEATURE:
 <221> NAME/KEY: MOD_RES
 <222> LOCATION: (26)..(26)
 <223> OTHER INFORMATION: Human NRGN peptide, amino acids 55-80, 26th
 residue citrullinated (Arg80)

<400> SEQUENCE: 37

Gly Pro Gly Pro Gly Gly Pro Gly Gly Ala Gly Val Ala Arg Gly Gly
 1 5 10 15

Ala Gly Gly Gly Pro Ser Gly Asp Thr Arg
 20 25

-continued

<210> SEQ ID NO 38
<211> LENGTH: 12
<212> TYPE: PRT
<213> ORGANISM: Bos taurus
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (6)..(6)
<223> OTHER INFORMATION: Bovine NRGN peptide, amino acids 33-44, 6th
residue citrullinated (Arg38)
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (11)..(11)
<223> OTHER INFORMATION: Bovine NRGN peptide, amino acids 33-44, 11th
residue citrullinated (Arg43)
<400> SEQUENCE: 38

Ile Gln Ala Ser Phe Arg Gly His Met Ala Arg Lys
1 5 10

<210> SEQ ID NO 39
<211> LENGTH: 24
<212> TYPE: PRT
<213> ORGANISM: Bos taurus
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (14)..(14)
<223> OTHER INFORMATION: Bovine NRGN peptide, amino acids 55-78, 14th
residue citrullinated (Arg68)
<400> SEQUENCE: 39

Gly Pro Gly Pro Gly Gly Pro Gly Gly Ala Gly Gly Ala Arg Gly Gly
1 5 10 15
Ala Gly Gly Gly Pro Ser Gly Asp
20

<210> SEQ ID NO 40
<211> LENGTH: 9
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (2)..(2)
<223> OTHER INFORMATION: Human GFAP peptide, amino acids 87-95, 2nd
residue citrullinated (Arg 88)
<400> SEQUENCE: 40

Val Arg Phe Leu Glu Gln Gln Asn Lys
1 5

<210> SEQ ID NO 41
<211> LENGTH: 9
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (1)..(9)
<223> OTHER INFORMATION: Human GFAP peptide, amino acids 87-95
<400> SEQUENCE: 41

Val Arg Phe Leu Glu Gln Gln Asn Lys
1 5

<210> SEQ ID NO 42
<211> LENGTH: 12
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:

-continued

<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: Human GFAP peptides, amino acids 96-107, 10th
residue citrullinated (Arg105)

<400> SEQUENCE: 42

Ala Leu Ala Ala Glu Leu Asn Gln Leu Arg Ala Lys
1 5 10

<210> SEQ ID NO 43
<211> LENGTH: 12
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (1)..(12)
<223> OTHER INFORMATION: Human GFAP peptide, amino acids 96-107

<400> SEQUENCE: 43

Ala Leu Ala Ala Glu Leu Asn Gln Leu Arg Ala Lys
1 5 10

<210> SEQ ID NO 44
<211> LENGTH: 16
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (2)..(2)
<223> OTHER INFORMATION: Human GFAP peptide, amino acids 123-138, 2nd
residue citrullinated (Arg124)
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Human GFAP peptide, amino acids 123-138, 4th
residue citrullinated (Arg126)
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (14)..(14)
<223> OTHER INFORMATION: Human GFAP peptide, amino acids 123-138, 14th
residue citrullinated (Arg136)

<400> SEQUENCE: 44

Leu Arg Leu Arg Leu Asp Gln Leu Thr Ala Asn Ser Ala Arg Leu Glu
1 5 10 15

<210> SEQ ID NO 45
<211> LENGTH: 16
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (1)..(16)
<223> OTHER INFORMATION: Human GFAP peptide, amino acids 123-138

<400> SEQUENCE: 45

Leu Arg Leu Arg Leu Asp Gln Leu Thr Ala Asn Ser Ala Arg Leu Glu
1 5 10 15

<210> SEQ ID NO 46
<211> LENGTH: 14
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Human GFAP peptide, amino acids 165-178, 9th
residue citrullinated (Arg173)

-continued

<400> SEQUENCE: 46

Ala Glu Asn Asn Leu Ala Ala Tyr Arg Gln Glu Ala Asp Glu
1 5 10

<210> SEQ ID NO 47

<211> LENGTH: 14

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<220> FEATURE:

<221> NAME/KEY: MISC_FEATURE

<222> LOCATION: (1)..(14)

<223> OTHER INFORMATION: Human GFAP peptide, amino acids 165-178

<400> SEQUENCE: 47

Ala Glu Asn Asn Leu Ala Ala Tyr Arg Gln Glu Ala Asp Glu
1 5 10

<210> SEQ ID NO 48

<211> LENGTH: 13

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<220> FEATURE:

<221> NAME/KEY: MOD_RES

<222> LOCATION: (7)..(7)

<223> OTHER INFORMATION: Human GFAP peptide, amino acids 211-223, 7th residue citrullinated (Arg117)

<400> SEQUENCE: 48

Leu Gln Glu Gln Leu Ala Arg Gln Gln Val His Val Glu
1 5 10

<210> SEQ ID NO 49

<211> LENGTH: 13

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<220> FEATURE:

<221> NAME/KEY: MISC_FEATURE

<222> LOCATION: (1)..(13)

<223> OTHER INFORMATION: Human GFAP peptide, amino acids 211-223

<400> SEQUENCE: 49

Leu Gln Glu Gln Leu Ala Arg Gln Gln Val His Val Glu
1 5 10

<210> SEQ ID NO 50

<211> LENGTH: 18

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<220> FEATURE:

<221> NAME/KEY: MOD_RES

<222> LOCATION: (3)..(3)

<223> OTHER INFORMATION: Human GFAP peptide, amino acids 256-273, 3rd residue citrullinated (Arg258)

<220> FEATURE:

<221> NAME/KEY: MOD_RES

<222> LOCATION: (15)..(15)

<223> OTHER INFORMATION: Human GFAP peptide, amino acids 256-273, 15th residue citrullinated (Arg270)

<400> SEQUENCE: 50

Trp Tyr Arg Ser Lys Phe Ala Asp Leu Thr Asp Ala Ala Ala Arg Asn
1 5 10 15

Ala Glu

-continued

<210> SEQ ID NO 51
<211> LENGTH: 18
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (1)..(18)
<223> OTHER INFORMATION: Human GFAP peptide, amino acids 256-273

<400> SEQUENCE: 51

Trp Tyr Arg Ser Lys Phe Ala Asp Leu Thr Asp Ala Ala Ala Arg Asn
1 5 10 15

Ala Glu

<210> SEQ ID NO 52
<211> LENGTH: 16
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (5)..(5)
<223> OTHER INFORMATION: Human GFAP peptide, amino acids 282-297, 5th
residue citrullinated (Arg286)
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (6)..(6)
<223> OTHER INFORMATION: Human GFAP peptide, amino acids 282-297, 6th
residue citrullinated (Arg287)

<400> SEQUENCE: 52

Ala Asn Asp Tyr Arg Arg Gln Leu Gln Ser Leu Thr Cys Asp Leu Glu
1 5 10 15

<210> SEQ ID NO 53
<211> LENGTH: 16
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (1)..(16)
<223> OTHER INFORMATION: Human GFAP peptide, amino acids 282-297

<400> SEQUENCE: 53

Ala Asn Asp Tyr Arg Arg Gln Leu Gln Ser Leu Thr Cys Asp Leu Glu
1 5 10 15

<210> SEQ ID NO 54
<211> LENGTH: 12
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (11)..(11)
<223> OTHER INFORMATION: Human GFAP peptide, amino acids 357-368, 11th
residue citrullinated (Arg367)

<400> SEQUENCE: 54

Leu Ala Leu Asp Ile Glu Ile Ala Thr Tyr Arg Lys
1 5 10

<210> SEQ ID NO 55
<211> LENGTH: 12
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (1)..(12)

-continued

<223> OTHER INFORMATION: Human GFAP peptide, amino acids 357-368

<400> SEQUENCE: 55

Leu Ala Leu Asp Ile Glu Ile Ala Thr Tyr Arg Lys
1 5 10

<210> SEQ ID NO 56

<211> LENGTH: 13

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<220> FEATURE:

<221> NAME/KEY: MOD_RES

<222> LOCATION: (5)..(5)

<223> OTHER INFORMATION: Human GFAP peptide, amino acids 402-414, 5th
residue citrullinated (Arg406)

<400> SEQUENCE: 56

Gly His Leu Lys Arg Asn Ile Val Val Lys Thr Val Glu
1 5 10

<210> SEQ ID NO 57

<211> LENGTH: 13

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<220> FEATURE:

<221> NAME/KEY: MISC_FEATURE

<222> LOCATION: (1)..(13)

<223> OTHER INFORMATION: Human GFAP peptide, amino acids 402-414

<400> SEQUENCE: 57

Gly His Leu Lys Arg Asn Ile Val Val Lys Thr Val Glu
1 5 10

<210> SEQ ID NO 58

<211> LENGTH: 20

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<220> FEATURE:

<221> NAME/KEY: MOD_RES

<222> LOCATION: (12)..(12)

<223> OTHER INFORMATION: Human MBP peptide, amino acids 12-31, 12th
residue citrullinated (Arg23)

<220> FEATURE:

<221> NAME/KEY: MOD_RES

<222> LOCATION: (18)..(18)

<223> OTHER INFORMATION: Human MBP peptide, amino acids 12-31, 18th
residue citrullinated (Arg29)

<400> SEQUENCE: 58

Tyr Leu Ala Ser Ala Ser Thr Met Asp His Ala Arg His Gly Phe Leu
1 5 10 15

Pro Arg His Arg
 20

<210> SEQ ID NO 59

<211> LENGTH: 20

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<220> FEATURE:

<221> NAME/KEY: MISC_FEATURE

<222> LOCATION: (1)..(20)

<223> OTHER INFORMATION: Human MBP peptide, amino acids 12-31

<400> SEQUENCE: 59

Tyr Leu Ala Ser Ala Ser Thr Met Asp His Ala Arg His Gly Phe Leu
1 5 10 15

-continued

Pro Arg His Arg
20

<210> SEQ ID NO 60
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (12)..(12)
<223> OTHER INFORMATION: Human MBP peptide, amino acids 12-29, 12th
residue citrullinated (Arg23)
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (18)..(18)
<223> OTHER INFORMATION: Human MBP peptide, amino acids 12-29, 18th
residue citrullinated (Arg29)
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (20)..(20)
<223> OTHER INFORMATION: Human MBP peptide, amino acids 12-29, 20th
residue citrullinated (Arg31)

<400> SEQUENCE: 60

Tyr Leu Ala Ser Ala Ser Thr Met Asp His Ala Arg His Gly Phe Leu
1 5 10 15

Pro Arg His Arg
20

<210> SEQ ID NO 61
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: Human MBP peptide, amino acids 32-51, 10th
residue citrullinated (Arg41)
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (16)..(16)
<223> OTHER INFORMATION: Human MBP peptide, amino acids 32-51, 16th
residue citrullinated (Arg47)

<400> SEQUENCE: 61

Asp Thr Gly Ile Leu Asp Ser Leu Gly Arg Phe Phe Gly Ser Asp Arg
1 5 10 15

Gly Ala Pro Lys
20

<210> SEQ ID NO 62
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (1)..(20)
<223> OTHER INFORMATION: Human MBP peptide, amino acids 32-51

<400> SEQUENCE: 62

Asp Thr Gly Ile Leu Asp Ser Leu Gly Arg Phe Phe Gly Ser Asp Arg
1 5 10 15

Gly Ala Pro Lys
20

-continued

<210> SEQ ID NO 63
<211> LENGTH: 22
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Human MBP peptide, amino acids 52-73, 1st
residue citrullinated (Arg52)

<400> SEQUENCE: 63

Arg Gly Ser Gly Lys Asp Gly His His Ala Ala Arg Thr Thr His Tyr
1 5 10 15

Gly Ser Leu Pro Gln Lys
20

<210> SEQ ID NO 64
<211> LENGTH: 22
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (1)..(22)
<223> OTHER INFORMATION: Human MBP peptide, amino acids 52-73

<400> SEQUENCE: 64

Arg Gly Ser Gly Lys Asp Gly His His Ala Ala Arg Thr Thr His Tyr
1 5 10 15

Gly Ser Leu Pro Gln Lys
20

<210> SEQ ID NO 65
<211> LENGTH: 17
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (5)..(5)
<223> OTHER INFORMATION: Human MBP peptide, amino acids 74-90, 5th
residue citrullinated (Arg78)

<400> SEQUENCE: 65

Ala Gln Gly His Arg Pro Gln Asp Glu Asn Pro Val Val His Phe Phe
1 5 10 15

Lys

<210> SEQ ID NO 66
<211> LENGTH: 17
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (2)..(2)
<223> OTHER INFORMATION: Human MBP peptide, amino acids 105-121, 2nd
residue citrullinated (Arg106)

<220> FEATURE:

<221> NAME/KEY: MOD_RES

<222> LOCATION: (8)..(8)

<223> OTHER INFORMATION: Human MBP peptide, amino acids 105-121, 8th
residue citrullinated (Arg112)

<400> SEQUENCE: 66

Gly Arg Gly Leu Ser Leu Ser Arg Phe Ser Trp Gly Ala Glu Gly Gln
1 5 10 15

Lys

-continued

<210> SEQ ID NO 67
<211> LENGTH: 17
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (1)..(17)
<223> OTHER INFORMATION: Human MBP peptide, amino acids 105-121

<400> SEQUENCE: 67

Gly Arg Gly Leu Ser Leu Ser Arg Phe Ser Trp Gly Ala Glu Gly Gln
1 5 10 15

Lys

<210> SEQ ID NO 68
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Human MBP peptide, amino acids 155-169, 4th
residue citrullinated (Arg158)
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (7)..(7)
<223> OTHER INFORMATION: Human MBP peptide, amino acids 155-169, 7th
residue citrullinated (Arg161)
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (14)..(14)
<223> OTHER INFORMATION: Human MBP peptide, amino acids 155-169, 14th
residue citrullinated (Arg168)
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (15)..(15)
<223> OTHER INFORMATION: Human MBP peptide, amino acids 155-169, 15th
residue citrullinated (Arg169)

<400> SEQUENCE: 68

Leu Gly Gly Arg Asp Ser Arg Ser Gly Ser Pro Met Ala Arg Arg
1 5 10 15

<210> SEQ ID NO 69
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (1)..(15)
<223> OTHER INFORMATION: Human MBP peptide, amino acids 155-169

<400> SEQUENCE: 69

Leu Gly Gly Arg Asp Ser Arg Ser Gly Ser Pro Met Ala Arg Arg
1 5 10 15

<210> SEQ ID NO 70
<211> LENGTH: 14
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (1)..(14)
<223> OTHER INFORMATION: Human NRGN peptide

<400> SEQUENCE: 70

-continued

Gly Pro Gly Pro Gly Gly Pro Gly Gly Ala Gly Val Ala Arg
1 5 10

<210> SEQ ID NO 71
 <211> LENGTH: 24
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens
 <220> FEATURE:
 <221> NAME/KEY: MOD_RES
 <222> LOCATION: (14)..(14)
 <223> OTHER INFORMATION: Human NRGN peptide, 14th residue citrullinated

<400> SEQUENCE: 71

Gly Pro Gly Pro Gly Gly Pro Gly Gly Ala Gly Val Ala Arg Gly Gly
1 5 10 15

Ala Gly Gly Gly Pro Ser Gly Asp
20

<210> SEQ ID NO 72
 <211> LENGTH: 25
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens
 <220> FEATURE:
 <221> NAME/KEY: MOD_RES
 <222> LOCATION: (15)..(15)
 <223> OTHER INFORMATION: Human NRGN peptide, 15th residue citrullinated

<400> SEQUENCE: 72

Lys Gly Pro Gly Pro Gly Gly Pro Gly Gly Ala Gly Val Ala Arg Gly
1 5 10 15

Gly Ala Gly Gly Gly Pro Ser Gly Asp
20 25

<210> SEQ ID NO 73
 <211> LENGTH: 24
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (1)..(24)
 <223> OTHER INFORMATION: Human NRGN peptide

<400> SEQUENCE: 73

Gly Pro Gly Pro Gly Gly Pro Gly Gly Ala Gly Val Ala Arg Gly Gly
1 5 10 15

Ala Gly Gly Gly Pro Ser Gly Asp
20

<210> SEQ ID NO 74
 <211> LENGTH: 16
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (1)..(16)
 <223> OTHER INFORMATION: Human MBP peptide

<400> SEQUENCE: 74

Ser His Gly Arg Thr Gln Asp Glu Asn Pro Val Val His Phe Phe Lys
1 5 10 15

<210> SEQ ID NO 75
 <211> LENGTH: 16
 <212> TYPE: PRT

-continued

<213> ORGANISM: Homo sapiens
 <220> FEATURE:
 <221> NAME/KEY: MOD_RES
 <222> LOCATION: (4)..(4)
 <223> OTHER INFORMATION: Human MBP peptide, 4th residue citrullinated

<400> SEQUENCE: 75

Ser His Gly Arg Thr Gln Asp Glu Asn Pro Val Val His Phe Phe Lys
 1 5 10 15

<210> SEQ ID NO 76
 <211> LENGTH: 17
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (1)..(17)
 <223> OTHER INFORMATION: Human MBP peptide

<400> SEQUENCE: 76

Asp Ser His His Pro Ala Arg Thr Ala His Tyr Gly Ser Leu Pro Gln
 1 5 10 15

Lys

<210> SEQ ID NO 77
 <211> LENGTH: 17
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens
 <220> FEATURE:
 <221> NAME/KEY: MOD_RES
 <222> LOCATION: (7)..(7)
 <223> OTHER INFORMATION: Human MBP peptide, 7th residue citrullinated

<400> SEQUENCE: 77

Asp Ser His His Pro Ala Arg Thr Ala His Tyr Gly Ser Leu Pro Gln
 1 5 10 15

Lys

<210> SEQ ID NO 78
 <211> LENGTH: 30
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (1)..(30)
 <223> OTHER INFORMATION: Human MBP peptide

<400> SEQUENCE: 78

Gly Arg Gly Leu Ser Leu Ser Arg Phe Ser Trp Gly Ala Glu Gly Gln
 1 5 10 15

Arg Pro Gly Phe Gly Tyr Gly Gly Arg Ala Ser Asp Tyr Lys
 20 25 30

<210> SEQ ID NO 79
 <211> LENGTH: 30
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens
 <220> FEATURE:
 <221> NAME/KEY: MOD_RES
 <222> LOCATION: (25)..(25)
 <223> OTHER INFORMATION: Human MBP peptide, 25th residue citrullinated

<400> SEQUENCE: 79

Gly Arg Gly Leu Ser Leu Ser Arg Phe Ser Trp Gly Ala Glu Gly Gln

-continued

```

1           5           10           15
Arg Pro Gly Phe Gly Tyr Gly Gly Arg Ala Ser Asp Tyr Lys
           20           25           30

```

```

<210> SEQ ID NO 80
<211> LENGTH: 30
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (2)..(2)
<223> OTHER INFORMATION: Human MBP peptide, 2nd residue citrullinated
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Human MBP peptide, 8th residue citrullinated
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (25)..(25)
<223> OTHER INFORMATION: Human MBP peptide, 25th residue citrullinated

```

```

<400> SEQUENCE: 80

```

```

Gly Arg Gly Leu Ser Leu Ser Arg Phe Ser Trp Gly Ala Glu Gly Gln
1           5           10           15

```

```

Arg Pro Gly Phe Gly Tyr Gly Gly Arg Ala Ser Asp Tyr Lys
           20           25           30

```

```

<210> SEQ ID NO 81
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (1)..(10)
<223> OTHER INFORMATION: Human PAD2 peptide

```

```

<400> SEQUENCE: 81

```

```

Gly Phe Pro Val Val Leu Asp Ser Pro Arg
1           5           10

```

```

<210> SEQ ID NO 82
<211> LENGTH: 16
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (1)..(16)
<223> OTHER INFORMATION: Human PAD2 peptide

```

```

<400> SEQUENCE: 82

```

```

Trp Ile Gln Asp Glu Ile Glu Phe Gly Tyr Ile Glu Ala Pro His Lys
1           5           10           15

```

```

<210> SEQ ID NO 83
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Bos taurus
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Bovine MBP peptide, 4th residue citrullinated
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (7)..(7)
<223> OTHER INFORMATION: Bovine MBP peptide, 7th residue citrullinated
<220> FEATURE:
<221> NAME/KEY: MOD_RES

```

-continued

<222> LOCATION: (14)..(14)
<223> OTHER INFORMATION: Bovine MBP peptide, 14th residue citrullinated
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (15)..(15)
<223> OTHER INFORMATION: Bovine MBP peptide, 15th residue citrullinated

<400> SEQUENCE: 83

Leu Gly Gly Arg Asp Ser Arg Ser Gly Ser Pro Met Ala Arg Arg
1 5 10 15

<210> SEQ ID NO 84
<211> LENGTH: 18
<212> TYPE: PRT
<213> ORGANISM: Bos taurus
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (12)..(12)
<223> OTHER INFORMATION: Bovine MBP peptide, amino acids 12-29, 12th residue citrullinated (Arg23)

<400> SEQUENCE: 84

Tyr Leu Ala Ser Ala Ser Thr Met Asp His Ala Arg His Gly Phe Leu
1 5 10 15

Pro Arg

<210> SEQ ID NO 85
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Bos taurus
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (6)..(6)
<223> OTHER INFORMATION: Bovine MBP peptide, amino acids 42-51, 6th residue citrullinated (Arg47)

<400> SEQUENCE: 85

Phe Phe Gly Ser Asp Arg Gly Ala Pro Lys
1 5 10

<210> SEQ ID NO 86
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Bos taurus
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (1)..(10)
<223> OTHER INFORMATION: Bovine MBP peptide, amino acids 42-51

<400> SEQUENCE: 86

Phe Phe Gly Ser Asp Arg Gly Ala Pro Lys
1 5 10

<210> SEQ ID NO 87
<211> LENGTH: 13
<212> TYPE: PRT
<213> ORGANISM: Bos taurus
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Bovine MBP, amino acids 155-167, 4th residue citrullinated (Arg158)
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (7)..(7)
<223> OTHER INFORMATION: Bovine MBP, amino acids 155-167, 7th residue citrullinated (Arg161)

-continued

<210> SEQ ID NO 92
<211> LENGTH: 12
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (1)..(12)
<223> OTHER INFORMATION: Human GFAP peptide, amino acids 30-41

<400> SEQUENCE: 92

Arg Leu Gly Pro Gly Thr Arg Leu Ser Leu Ala Arg
1 5 10

<210> SEQ ID NO 93
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (6)..(6)
<223> OTHER INFORMATION: Human GFAP peptide, amino acids 31-41, 6th
residue citrullinated (Arg36)

<400> SEQUENCE: 93

Leu Gly Pro Gly Thr Arg Leu Ser Leu Ala Arg
1 5 10

<210> SEQ ID NO 94
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (1)..(11)
<223> OTHER INFORMATION: Human GFAP peptide, amino acids 31-41

<400> SEQUENCE: 94

Leu Gly Pro Gly Thr Arg Leu Ser Leu Ala Arg
1 5 10

<210> SEQ ID NO 95
<211> LENGTH: 16
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: Human GFAP peptide, amino acids 261-276, 10th
residue citrullinated (Arg270)

<400> SEQUENCE: 95

Phe Ala Asp Leu Thr Asp Ala Ala Ala Arg Asn Ala Glu Leu Leu Arg
1 5 10 15

<210> SEQ ID NO 96
<211> LENGTH: 16
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (1)..(16)
<223> OTHER INFORMATION: Human GFAP peptide, amino acids 261-276

<400> SEQUENCE: 96

Phe Ala Asp Leu Thr Asp Ala Ala Ala Arg Asn Ala Glu Leu Leu Arg

-continued

1 5 10 15

<210> SEQ ID NO 97
 <211> LENGTH: 19
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens
 <220> FEATURE:
 <221> NAME/KEY: MOD_RES
 <222> LOCATION: (10)..(10)
 <223> OTHER INFORMATION: Human GFAP peptide, amino acids 261-279, 10th
 residue citrullinated (Arg270)
 <220> FEATURE:
 <221> NAME/KEY: MOD_RES
 <222> LOCATION: (16)..(16)
 <223> OTHER INFORMATION: Human GFAP peptide, amino acids 261-279, 16th
 residue citrullinated (Arg276)

<400> SEQUENCE: 97

Phe Ala Asp Leu Thr Asp Ala Ala Ala Arg Asn Ala Glu Leu Leu Arg
 1 5 10 15

Gln Ala Lys

<210> SEQ ID NO 98
 <211> LENGTH: 11
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens
 <220> FEATURE:
 <221> NAME/KEY: MOD_RES
 <222> LOCATION: (5)..(5)
 <223> OTHER INFORMATION: Human GFAP peptide, amino acids 412-422, 5th
 residue citrullinated (Arg416)

<400> SEQUENCE: 98

Thr Val Glu Met Arg Asp Gly Glu Val Ile Lys
 1 5 10

<210> SEQ ID NO 99
 <211> LENGTH: 28
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens
 <220> FEATURE:
 <221> NAME/KEY: MOD_RES
 <222> LOCATION: (18)..(18)
 <223> OTHER INFORMATION: Human NRGN peptide, amino acids 51-78, 18th
 residue citrullinated (Arg68)

<400> SEQUENCE: 99

Arg Gly Arg Lys Gly Pro Gly Pro Gly Gly Pro Gly Gly Ala Gly Val
 1 5 10 15

Ala Arg Gly Gly Ala Gly Gly Gly Pro Ser Gly Asp
 20 25

<210> SEQ ID NO 100
 <211> LENGTH: 28
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (1)..(28)
 <223> OTHER INFORMATION: Human NRGN peptide, amino acids 51-78

<400> SEQUENCE: 100

Arg Gly Arg Lys Gly Pro Gly Pro Gly Gly Pro Gly Gly Ala Gly Val
 1 5 10 15

Ala Arg Gly Gly Ala Gly Gly Gly Pro Ser Gly Asp

-continued

<223> OTHER INFORMATION: Recombinant NRGN peptide, amino acids 51-80,
30th residue citrullinated (Arg80)

<400> SEQUENCE: 103

Arg Gly Arg Lys Gly Pro Gly Pro Gly Gly Pro Gly Gly Ala Gly Val
1 5 10 15
Ala Arg Gly Gly Ala Gly Gly Gly Pro Ser Gly Asp Thr Arg
 20 25 30

<210> SEQ ID NO 104

<211> LENGTH: 18

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<220> FEATURE:

<221> NAME/KEY: MOD_RES

<222> LOCATION: (12)..(12)

<223> OTHER INFORMATION: Human MBP peptide, 12th residue citrullinated
(Arg26)

<400> SEQUENCE: 104

Tyr Leu Ala Thr Ala Ser Thr Met Asp His Ala Arg His Gly Phe Leu
1 5 10 15

Pro Arg

<210> SEQ ID NO 105

<211> LENGTH: 18

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<220> FEATURE:

<221> NAME/KEY: MISC_FEATURE

<222> LOCATION: (1)..(18)

<223> OTHER INFORMATION: Human MBP peptide

<400> SEQUENCE: 105

Tyr Leu Ala Thr Ala Ser Thr Met Asp His Ala Arg His Gly Phe Leu
1 5 10 15

Pro Arg

<210> SEQ ID NO 106

<211> LENGTH: 20

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<220> FEATURE:

<221> NAME/KEY: MOD_RES

<222> LOCATION: (12)..(12)

<223> OTHER INFORMATION: Human MBP peptide, 12th residue citrullinated
(Arg 26)

<220> FEATURE:

<221> NAME/KEY: MOD_RES

<222> LOCATION: (18)..(18)

<223> OTHER INFORMATION: Human MBP peptide, 18th residue citrullinated
(Arg 232)

<400> SEQUENCE: 106

Tyr Leu Ala Thr Ala Ser Thr Met Asp His Ala Arg His Gly Phe Leu
1 5 10 15

Pro Arg His Arg
 20

<210> SEQ ID NO 107

<211> LENGTH: 20

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<220> FEATURE:

-continued

<221> NAME/KEY: MISC_FEATURE

<222> LOCATION: (1)..(20)

<223> OTHER INFORMATION: Human MBP peptide

<400> SEQUENCE: 107

Tyr Leu Ala Thr Ala Ser Thr Met Asp His Ala Arg His Gly Phe Leu
1 5 10 15

Pro Arg His Arg
20

<210> SEQ ID NO 108

<211> LENGTH: 12

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<220> FEATURE:

<221> NAME/KEY: MOD_RES

<222> LOCATION: (2)..(2)

<223> OTHER INFORMATION: Human MBP peptide, 2nd residue citrullinated
(Arg34)

<400> SEQUENCE: 108

His Arg Asp Thr Gly Ile Leu Asp Ser Ile Gly Arg
1 5 10

<210> SEQ ID NO 109

<211> LENGTH: 12

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<220> FEATURE:

<221> NAME/KEY: MISC_FEATURE

<222> LOCATION: (1)..(12)

<223> OTHER INFORMATION: Human MBP peptide

<400> SEQUENCE: 109

His Arg Asp Thr Gly Ile Leu Asp Ser Ile Gly Arg
1 5 10

<210> SEQ ID NO 110

<211> LENGTH: 16

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<220> FEATURE:

<221> NAME/KEY: MOD_RES

<222> LOCATION: (10)..(10)

<223> OTHER INFORMATION: Human MBP peptide, 10th residue citrullinated
(Arg44)

<400> SEQUENCE: 110

Asp Thr Gly Ile Leu Asp Ser Ile Gly Arg Phe Phe Gly Gly Asp Arg
1 5 10 15

<210> SEQ ID NO 111

<211> LENGTH: 16

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<220> FEATURE:

<221> NAME/KEY: MISC_FEATURE

<222> LOCATION: (1)..(16)

<223> OTHER INFORMATION: Human MBP peptide

<400> SEQUENCE: 111

Asp Thr Gly Ile Leu Asp Ser Ile Gly Arg Phe Phe Gly Gly Asp Arg
1 5 10 15

<210> SEQ ID NO 112

-continued

<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (6)..(6)
<223> OTHER INFORMATION: Human MBP peptide, 6th residue citrullinated
(Arg50)

<400> SEQUENCE: 112

Phe Phe Gly Gly Asp Arg Gly Ala Pro Lys
1 5 10

<210> SEQ ID NO 113
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (1)..(10)
<223> OTHER INFORMATION: Human MBP peptide

<400> SEQUENCE: 113

Phe Phe Gly Gly Asp Arg Gly Ala Pro Lys
1 5 10

<210> SEQ ID NO 114
<211> LENGTH: 17
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Human MBP peptide, 9th residue citrullinated
(Arg149)

<400> SEQUENCE: 114

Phe Ser Trp Gly Ala Glu Gly Gln Arg Pro Gly Phe Gly Tyr Gly Gly
1 5 10 15

Arg

<210> SEQ ID NO 115
<211> LENGTH: 17
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (1)..(17)
<223> OTHER INFORMATION: Human MBP peptide

<400> SEQUENCE: 115

Phe Ser Trp Gly Ala Glu Gly Gln Arg Pro Gly Phe Gly Tyr Gly Gly
1 5 10 15

Arg

<210> SEQ ID NO 116
<211> LENGTH: 13
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Human MBP peptide, 8th residue citrullinated
(Arg157)

<400> SEQUENCE: 116

-continued

Pro Gly Phe Gly Tyr Gly Gly Arg Ala Ser Asp Tyr Lys
1 5 10

<210> SEQ ID NO 117
<211> LENGTH: 13
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (1)..(13)
<223> OTHER INFORMATION: Human MBP peptide

<400> SEQUENCE: 117

Pro Gly Phe Gly Tyr Gly Gly Arg Ala Ser Asp Tyr Lys
1 5 10

<210> SEQ ID NO 118
<211> LENGTH: 12
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Human GFAP peptide, 1st residue citrullinated
(Arg30)
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (7)..(7)
<223> OTHER INFORMATION: Human GFAP peptide, 7th residue citrullinated
(Arg36)

<400> SEQUENCE: 118

Arg Leu Gly Pro Gly Thr Arg Leu Ser Leu Ala Arg
1 5 10

<210> SEQ ID NO 119
<211> LENGTH: 12
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (1)..(12)
<223> OTHER INFORMATION: Human GFAP peptide

<400> SEQUENCE: 119

Arg Leu Gly Pro Gly Thr Arg Leu Ser Leu Ala Arg
1 5 10

<210> SEQ ID NO 120
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (6)..(6)
<223> OTHER INFORMATION: Human GFAP peptide, 6th residue citrullinated
(Arg36)

<400> SEQUENCE: 120

Leu Gly Pro Gly Thr Arg Leu Ser Leu Ala Arg
1 5 10

<210> SEQ ID NO 121
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

-continued

<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (1)..(11)
<223> OTHER INFORMATION: Human GFAP peptide

<400> SEQUENCE: 121

Leu Gly Pro Gly Thr Arg Leu Ser Leu Ala Arg
1 5 10

<210> SEQ ID NO 122
<211> LENGTH: 16
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: Human GFAP peptide, 10th residue citrullinated
(Arg270)

<400> SEQUENCE: 122

Phe Ala Asp Leu Thr Asp Ala Ala Ala Arg Asn Ala Glu Leu Leu Arg
1 5 10 15

<210> SEQ ID NO 123
<211> LENGTH: 16
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (1)..(16)
<223> OTHER INFORMATION: Human GFAP peptide

<400> SEQUENCE: 123

Phe Ala Asp Leu Thr Asp Ala Ala Ala Arg Asn Ala Glu Leu Leu Arg
1 5 10 15

<210> SEQ ID NO 124
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (5)..(5)
<223> OTHER INFORMATION: Human GFAP peptide, 5th residue citrullinated
(Arg416)

<400> SEQUENCE: 124

Thr Val Glu Met Arg Asp Gly Glu Val Ile Lys
1 5 10

<210> SEQ ID NO 125
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (1)..(11)
<223> OTHER INFORMATION: Human GFAP peptide

<400> SEQUENCE: 125

Thr Val Glu Met Arg Asp Gly Glu Val Ile Lys
1 5 10

<210> SEQ ID NO 126
<211> LENGTH: 6
<212> TYPE: PRT

-continued

<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (1)..(6)
<223> OTHER INFORMATION: Human NRGN peptide

<400> SEQUENCE: 126

Ile Gln Ala Ser Phe Arg
1 5

<210> SEQ ID NO 127
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (1)..(15)
<223> OTHER INFORMATION: Human NRGN peptide

<400> SEQUENCE: 127

Lys Gly Pro Gly Pro Gly Gly Pro Gly Gly Ala Gly Val Ala Arg
1 5 10 15

<210> SEQ ID NO 128
<211> LENGTH: 13
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Human Tubulin beta-4B chain peptide, 8th
residue citrullinated (Arg162)

<400> SEQUENCE: 128

Ile Arg Glu Glu Tyr Pro Asp Arg Ile Met Asn Thr Phe
1 5 10

<210> SEQ ID NO 129
<211> LENGTH: 13
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (1)..(13)
<223> OTHER INFORMATION: Human Tubulin beta-4B chain peptide

<400> SEQUENCE: 129

Ile Arg Glu Glu Tyr Pro Asp Arg Ile Met Asn Thr Phe
1 5 10

<210> SEQ ID NO 130
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Human Tubulin alpha-1B chain peptide, 9th
residue citrullinated (Arg320)

<400> SEQUENCE: 130

Tyr Met Ala Cys Cys Leu Leu Tyr Arg Gly Asp Val Val Pro Lys
1 5 10 15

<210> SEQ ID NO 131
<211> LENGTH: 15

-continued

<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (1)..(15)
<223> OTHER INFORMATION: Human tubulin alpha-1B chain peptide

<400> SEQUENCE: 131

Tyr Met Ala Cys Cys Leu Leu Tyr Arg Gly Asp Val Val Pro Lys
1 5 10 15

<210> SEQ ID NO 132
<211> LENGTH: 13
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (2)..(2)
<223> OTHER INFORMATION: Human tubulin alpha-1B chain peptide, 1st
residue citrullinated (Arg79)
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (7)..(7)
<223> OTHER INFORMATION: Human tubulin alpha-1B chain peptide, 7th
residue citrullinated (Arg84)

<400> SEQUENCE: 132

Val Arg Thr Gly Thr Tyr Arg Gln Leu Phe His Pro Glu
1 5 10

<210> SEQ ID NO 133
<211> LENGTH: 13
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (1)..(13)
<223> OTHER INFORMATION: Human tubulin alpha-1B chain

<400> SEQUENCE: 133

Val Arg Thr Gly Thr Tyr Arg Gln Leu Phe His Pro Glu
1 5 10

<210> SEQ ID NO 134
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (5)..(5)
<223> OTHER INFORMATION: Human CNPase peptide, 5th residue citrullinated
(Arg68)

<400> SEQUENCE: 134

Ser Thr Leu Ala Arg Val Ile Val Asp Lys
1 5 10

<210> SEQ ID NO 135
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (1)..(10)
<223> OTHER INFORMATION: Human CNPase peptide

<400> SEQUENCE: 135

-continued

Ser Thr Leu Ala Arg Val Ile Val Asp Lys
1 5 10

<210> SEQ ID NO 136
<211> LENGTH: 14
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (6)..(6)
<223> OTHER INFORMATION: Human CNPase peptide, 6th residue citrullinated
(Arg93)

<400> SEQUENCE: 136

Ile Thr Pro Gly Ala Arg Gly Ala Phe Ser Glu Glu Tyr Lys
1 5 10

<210> SEQ ID NO 137
<211> LENGTH: 14
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (1)..(14)
<223> OTHER INFORMATION: Human CNPase peptide

<400> SEQUENCE: 137

Ile Thr Pro Gly Ala Arg Gly Ala Phe Ser Glu Glu Tyr Lys
1 5 10

<210> SEQ ID NO 138
<211> LENGTH: 13
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (6)..(6)
<223> OTHER INFORMATION: Human PPIA peptide, 6th residue citrullinated
(Arg37)

<400> SEQUENCE: 138

Thr Ala Glu Asn Phe Arg Ala Leu Ser Thr Gly Glu Lys
1 5 10

<210> SEQ ID NO 139
<211> LENGTH: 13
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (1)..(13)
<223> OTHER INFORMATION: Human PPIA peptide

<400> SEQUENCE: 139

Thr Ala Glu Asn Phe Arg Ala Leu Ser Thr Gly Glu Lys
1 5 10

<210> SEQ ID NO 140
<211> LENGTH: 13
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Human Septin-7 peptide, 10th residue
citrullinated (Arg425)

-continued

<400> SEQUENCE: 140

Ile Leu Glu Gln Gln Asn Ser Ser Arg Thr Leu Glu Lys
1 5 10

<210> SEQ ID NO 141
<211> LENGTH: 13
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (1)..(13)
<223> OTHER INFORMATION: Human Septin-7 peptide

<400> SEQUENCE: 141

Ile Leu Glu Gln Gln Asn Ser Ser Arg Thr Leu Glu Lys
1 5 10

<210> SEQ ID NO 142
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (3)..(3)
<223> OTHER INFORMATION: Human Elongation factor 1-alpha 2 peptide, 3rd
residue citrullinated (Arg247)

<400> SEQUENCE: 142

Pro Leu Arg Leu Pro Leu Gln Asp Val Tyr Lys
1 5 10

<210> SEQ ID NO 143
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (1)..(11)
<223> OTHER INFORMATION: Human elongation factor1-alpha 2

<400> SEQUENCE: 143

Pro Leu Arg Leu Pro Leu Gln Asp Val Tyr Lys
1 5 10

<210> SEQ ID NO 144
<211> LENGTH: 16
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (14)..(14)
<223> OTHER INFORMATION: Human elongation factor 1-alpha 2 peptide, 14th
residue citrullinated (Arg266)

<400> SEQUENCE: 144

Val Tyr Lys Ile Gly Gly Ile Gly Thr Val Pro Val Gly Arg Val Glu
1 5 10 15

<210> SEQ ID NO 145
<211> LENGTH: 16
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (1)..(16)
<223> OTHER INFORMATION: Human elongation factor 1-alpha 2 peptide

-continued

<400> SEQUENCE: 145

Val Tyr Lys Ile Gly Gly Ile Gly Thr Val Pro Val Gly Arg Val Glu
1 5 10 15

<210> SEQ ID NO 146

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<220> FEATURE:

<221> NAME/KEY: MOD_RES

<222> LOCATION: (9)..(9)

<223> OTHER INFORMATION: Human TPPP peptide, 9th residue citrullinated
(Arg165)

<400> SEQUENCE: 146

Ala Ile Ser Ser Pro Thr Val Ser Arg Leu Thr Asp Thr Thr Lys
1 5 10 15

<210> SEQ ID NO 147

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<220> FEATURE:

<221> NAME/KEY: MISC_FEATURE

<222> LOCATION: (1)..(15)

<223> OTHER INFORMATION: Human TPPP peptide

<400> SEQUENCE: 147

Ala Ile Ser Ser Pro Thr Val Ser Arg Leu Thr Asp Thr Thr Lys
1 5 10 15

<210> SEQ ID NO 148

<211> LENGTH: 19

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<220> FEATURE:

<221> NAME/KEY: MOD_RES

<222> LOCATION: (13)..(13)

<223> OTHER INFORMATION: Human TPPP3 peptide, 13th residue citrullinated
(Arg130)

<400> SEQUENCE: 148

Thr Gly Gly Ala Val Asp Arg Leu Thr Asp Thr Ser Arg Tyr Thr Gly
1 5 10 15

Ser His Lys

<210> SEQ ID NO 149

<211> LENGTH: 19

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<220> FEATURE:

<221> NAME/KEY: MISC_FEATURE

<222> LOCATION: (1)..(19)

<223> OTHER INFORMATION: Human TPPP3 peptide

<400> SEQUENCE: 149

Thr Gly Gly Ala Val Asp Arg Leu Thr Asp Thr Ser Arg Tyr Thr Gly
1 5 10 15

Ser His Lys

<210> SEQ ID NO 150

<211> LENGTH: 19

<212> TYPE: PRT

-continued

<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (5)..(5)
<223> OTHER INFORMATION: Human TPPP3 peptide, 5th residue citrullinated
(Arg151)

<400> SEQUENCE: 150

Gly Ile Ala Gly Arg Gln Asp Ile Leu Asp Asp Ser Gly Tyr Val Ser
1 5 10 15

Ala Tyr Lys

<210> SEQ ID NO 151
<211> LENGTH: 19
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (1)..(19)
<223> OTHER INFORMATION: Human TPPP3 peptide

<400> SEQUENCE: 151

Gly Ile Ala Gly Arg Gln Asp Ile Leu Asp Asp Ser Gly Tyr Val Ser
1 5 10 15

Ala Tyr Lys

<210> SEQ ID NO 152
<211> LENGTH: 18
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (12)..(12)
<223> OTHER INFORMATION: Human Ermin, Isoform 2, 12th residue
citrullinated (Arg57)

<400> SEQUENCE: 152

Leu Thr Asp Val Asp Ser Pro Leu Pro His Tyr Arg Val Glu Pro Ser
1 5 10 15

Leu Glu

<210> SEQ ID NO 153
<211> LENGTH: 18
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (1)..(18)
<223> OTHER INFORMATION: Human Ermin, Isoform2

<400> SEQUENCE: 153

Leu Thr Asp Val Asp Ser Pro Leu Pro His Tyr Arg Val Glu Pro Ser
1 5 10 15

Leu Glu

<210> SEQ ID NO 154
<211> LENGTH: 16
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (14)..(14)
<223> OTHER INFORMATION: Human NDRG2, Isoform2 peptide, 14th residue
citrullinated (Arg329)

-continued

<400> SEQUENCE: 154

Thr Ala Ser Leu Thr Ser Ala Ala Ser Val Asp Gly Asn Arg Ser Arg
 1 5 10 15

<210> SEQ ID NO 155

<211> LENGTH: 16

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<220> FEATURE:

<221> NAME/KEY: MISC_FEATURE

<222> LOCATION: (1)..(16)

<223> OTHER INFORMATION: Human NDRG2, Isoform 2

<400> SEQUENCE: 155

Thr Ala Ser Leu Thr Ser Ala Ala Ser Val Asp Gly Asn Arg Ser Arg
 1 5 10 15

<210> SEQ ID NO 156

<211> LENGTH: 17

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<220> FEATURE:

<221> NAME/KEY: MISC_FEATURE

<222> LOCATION: (1)..(17)

<223> OTHER INFORMATION: Human MBP peptide, amino acids 74-90

<400> SEQUENCE: 156

Ala Gln Gly His Arg Pro Gln Asp Glu Asn Pro Val Val His Phe Phe
 1 5 10 15

Lys

<210> SEQ ID NO 157

<211> LENGTH: 26

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<220> FEATURE:

<221> NAME/KEY: MISC_FEATURE

<222> LOCATION: (1)..(26)

<223> OTHER INFORMATION: Human NRG1 peptide, amino acids 55-80

<400> SEQUENCE: 157

Gly Pro Gly Pro Gly Gly Pro Gly Gly Ala Gly Val Ala Arg Gly Gly
 1 5 10 15

Ala Gly Gly Gly Pro Ser Gly Asp Thr Arg
 20 25

<210> SEQ ID NO 158

<211> LENGTH: 12

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<220> FEATURE:

<221> NAME/KEY: MOD_RES

<222> LOCATION: (1)..(1)

<223> OTHER INFORMATION: Human GFAP peptide, amino acids 30-41, 1st residue citrullinated (Arg30)

<220> FEATURE:

<221> NAME/KEY: MOD_RES

<222> LOCATION: (7)..(7)

<223> OTHER INFORMATION: Human GFAP peptide, amino acids 30-41, 7th residue citrullinated (Arg36)

<400> SEQUENCE: 158

Arg Leu Gly Pro Gly Thr Arg Leu Ser Leu Ala Arg
 1 5 10

-continued

<210> SEQ ID NO 159
 <211> LENGTH: 25
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (1)..(25)
 <223> OTHER INFORMATION: Human NRGN peptide

<400> SEQUENCE: 159

Lys Gly Pro Gly Pro Gly Gly Pro Gly Gly Ala Gly Val Ala Arg Gly
 1 5 10 15
 Gly Ala Gly Gly Gly Pro Ser Gly Asp
 20 25

<210> SEQ ID NO 160
 <211> LENGTH: 11
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (1)..(11)
 <223> OTHER INFORMATION: Human GFAP peptide, amino acids 412-422

<400> SEQUENCE: 160

Thr Val Glu Met Arg Asp Gly Glu Val Ile Lys
 1 5 10

<210> SEQ ID NO 161
 <211> LENGTH: 97
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Recombinant human NRGN protein

<400> SEQUENCE: 161

Met His His His His His His Glu Asn Leu Tyr Phe Gln Gly Ala Ile
 1 5 10 15
 Ala Met Asp Cys Cys Thr Glu Asn Ala Cys Ser Lys Pro Asp Asp Asp
 20 25 30
 Ile Leu Asp Ile Pro Leu Asp Asp Pro Gly Ala Asn Ala Ala Ala Ala
 35 40 45
 Lys Ile Gln Ala Ser Phe Arg Gly His Met Ala Arg Lys Lys Ile Lys
 50 55 60
 Ser Gly Glu Arg Gly Arg Lys Gly Pro Gly Pro Gly Gly Pro Gly Gly
 65 70 75 80
 Ala Gly Val Ala Arg Gly Gly Ala Gly Gly Gly Pro Ser Gly Asp Thr
 85 90 95

Arg

<210> SEQ ID NO 162
 <211> LENGTH: 171
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (1)..(171)
 <223> OTHER INFORMATION: Human MBP Isoform 5

<400> SEQUENCE: 162

-continued

Met	Ala	Ser	Gln	Lys	Arg	Pro	Ser	Gln	Arg	His	Gly	Ser	Lys	Tyr	Leu	1	5	10	15
Ala	Thr	Ala	Ser	Thr	Met	Asp	His	Ala	Arg	His	Gly	Phe	Leu	Pro	Arg	20	25	30	
His	Arg	Asp	Thr	Gly	Ile	Leu	Asp	Ser	Ile	Gly	Arg	Phe	Phe	Gly	Gly	35	40	45	
Asp	Arg	Gly	Ala	Pro	Lys	Arg	Gly	Ser	Gly	Lys	Asp	Ser	His	His	Pro	50	55	60	
Ala	Arg	Thr	Ala	His	Tyr	Gly	Ser	Leu	Pro	Gln	Lys	Ser	His	Gly	Arg	65	70	75	80
Thr	Gln	Asp	Glu	Asn	Pro	Val	Val	His	Phe	Phe	Lys	Asn	Ile	Val	Thr	85	90	95	
Pro	Arg	Thr	Pro	Pro	Pro	Ser	Gln	Gly	Lys	Gly	Arg	Gly	Leu	Ser	Leu	100	105	110	
Ser	Arg	Phe	Ser	Trp	Gly	Ala	Glu	Gly	Gln	Arg	Pro	Gly	Phe	Gly	Tyr	115	120	125	
Gly	Gly	Arg	Ala	Ser	Asp	Tyr	Lys	Ser	Ala	His	Lys	Gly	Phe	Lys	Gly	130	135	140	
Val	Asp	Ala	Gln	Gly	Thr	Leu	Ser	Lys	Ile	Phe	Lys	Leu	Gly	Gly	Arg	145	150	155	160
Asp	Ser	Arg	Ser	Gly	Ser	Pro	Met	Ala	Arg	Arg	165	170							

What is claimed is:

1. A method of detecting treatment efficacy in a subject who is being treated for brain injury, the method comprising:

- measuring at a first time point the level of autoantibodies to at least two citrullinated brain injury biomarker proteins selected from the group consisting of neurogranin (cit-NRGN), glial fibrillary acidic protein (cit-GFAP), and citrullinated myelin basic protein (cit-MBP), in a sample obtained from the subject prior to initiating brain injury treatment in the subject;
- measuring the levels of autoantibodies to cit-NRGN, cit-GFAP and/or cit-MBP biomarker proteins using a cit-NRGN-specific peptide, a cit-GFAP-specific peptide and/or a cit-MBP-specific peptide that specifically binds the autoantibodies in a sample obtained from the subject at one or more time points after initiating brain injury treatment in the subject; and
- detecting efficacy of the brain injury treatment by measuring decreased levels of autoantibodies that bind to the cit-NRGN, cit-GFAP and/or cit-MBP peptide in the sample in step (b) relative to the levels of autoantibodies to cit-NRGN, cit-GFAP and/or cit-MBP measured in the sample at the first time point in step (a).

2. The method of claim 1, wherein the citrullinated NRGN is citrullinated human NRGN which is citrullinated at one or more arginine (R) amino acid residues selected from R38, R43, R51, R53, and R68 in the human NRGN polypeptide of SEQ ID NO: 1.

3. The method of claim 1, wherein the citrullinated GFAP is citrullinated human GFAP, which is citrullinated at one or more arginine (R) amino acid residues selected from R30, R36, R88, R105, R124, R126, R136, R173, R217, R258, R270, R276, R286, R287, R367, R406, and R416 of the human GFAP polypeptide of SEQ ID NO: 3.

4. The method of claim 1, wherein the citrullinated MBP is citrullinated human MBP, which is citrullinated at one or more arginine (R) amino acid residues selected from R26, R32, R34, R44, R50, R92, R106, R124, R157, R186, R189, R196, and R197 in the human MBP polypeptide of SEQ ID NO: 163.

5. The method of claim 1, further comprising contacting the sample with one or more citrullinated brain injury biomarker proteins selected from the group consisting of tubulin beta-4B chain, tubulin alpha-1B chain, CNPase, PPIA, Septin-7, Elongation factor1-alpha2, TPPP, TPPP3, Ermin Isoform 2, NDRG2 Isoform 2, astrotactin 1 (ASTN1), brain angiogenesis inhibitor 3 (BAI3); carnosine dipeptidase 1 (CNDP1); ERMIN; glutamate receptor metabotropic 3 (GRM3); kelch-like protein 32 (KLH32); melanoma antigen family E,2 (MAGE2); neuregulin 3 (NRG3); oligodendrocyte myelin glycoprotein (OMG); solute carrier family 39 (zinc transporter); reticulon 1 (RTN1); MT3, and peptidylarginine deiminase, types 1-4 and 6 (PAD1-4 or PAD6).

6. The method of claim 1, wherein the sample is blood, serum, plasma, cerebrospinal fluid (CSF), urine, saliva, stool and synovial fluid.

7. The method of claim 6, wherein the sample is blood, serum or plasma.

8. The method of claim 1, wherein binding is detected using a detection method selected from mass spectrometry, immunoassay, electrochemical luminescent assay, electrochemical voltammetry, electrochemical amperometry, atomic force microscopy, radio frequency multipolar resonance spectroscopy, confocal microscopy, non-confocal microscopy, fluorescence optical detection, luminescence optical detection, chemiluminescence optical detection, absorbance optical detection, reflectance optical detection, transmittance optical detection, birefringence optical detection, refractive

index detection, surface plasmon resonance, ellipsometry, resonant mirror detection, grating coupler waveguide detection, or interferometry.

9. The method of claim 8, wherein the detecting by mass spectrometry is accomplished using multiple reaction monitoring mass spectrometry (MRM-MS).

10. The method of claim 1, wherein the subject has or is suspected of having a neurodegenerative disease.

11. The method of claim 10, wherein the neurodegenerative disease is selected from hemorrhagic stroke, ischemic stroke, chronic traumatic encephalopathy, Alzheimer's disease, or Parkinson's Disease.

12. A method of detecting progress of brain injury treatment in a subject having or suspected of having brain injury, the method comprising:

- a. measuring at a first time point prior to treatment a baseline level of autoantibodies to at least two citrullinated brain injury biomarker proteins selected from the group consisting of neurogranin (cit-NRGN), glial fibrillary acidic protein (cit-GFAP), and citrullinated myelin basic protein (cit-MBP), in a sample obtained from a subject in need thereof;
- b. measuring the level of autoantibodies to cit-NRGN, cit-GFAP and/or cit-MBP biomarker proteins using a cit-NRGN-specific peptide, a cit-GFAP-specific peptide and/or a cit-MBP-specific peptide that specifically binds the autoantibodies in a sample obtained from the subject at least one additional time point at after treatment initiation; and
- c. detecting progress of the brain injury treatment by measuring decreased levels of autoantibodies that bind to the cit-NRGN, cit-GFAP and/or cit-MBP peptide in the sample in step (b) relative to the baseline level of autoantibodies to cit-NRGN, cit-GFAP and/or cit-MBP measured in the sample at the first time point in step (a).

13. The method of claim 12, wherein the citrullinated NRGN is citrullinated human NRGN which is citrullinated at one or more arginine (R) amino acid residues selected from R38, R43, R51, R53, and R68 in the human NRGN polypeptide of SEQ ID NO: 1.

14. The method of claim 12, wherein the citrullinated GFAP is citrullinated human GFAP, which is citrullinated at one or more arginine (R) amino acid residues selected from R30, R36, R88, R105, R124, R126, R136, R173, R217, R258, R270, R276, R286, R287, R367, R406, and R416 of the human GFAP polypeptide of SEQ ID NO: 3.

15. The method of claim 12, wherein the citrullinated MBP is citrullinated human MBP, which is citrullinated at one or more arginine (R) amino acid residues selected from R26, R32, R34, R44, R50, R92, R106, R124 R157, R186, R189, R196, and R197 in the human MBP polypeptide of SEQ ID NO: 163.

16. The method of claim 12, further comprising contacting the sample with one or more citrullinated brain injury biomarker proteins selected from the group consisting of tubulin beta-4B chain, tubulin alpha-1B chain, CNPase, PPIA, Septin-7, Elongation factor1-alpha2, TPPP, TPPP3, Ermin Isoform 2, NDRG2 Isoform 2, astrotactin 1 (ASTN1), brain angiogenesis inhibitor 3 (BAI3); carnosine dipeptidase 1 (CNDP1); ERMIN; glutamate receptor metabotropic 3 (GRM3); kelch-like protein 32 (KLH32); melanoma antigen family E,2 (MAGE2); neuregulin 3 (NRG3); oligodendrocyte myelin glycoprotein (OMG); solute carrier family 39 (zinc transporter); reticulon 1 (RTN1); MT3, and peptidyl-arginine deiminase, types 1-4 and 6 (PAD1-4 or PAD6).

17. The method of claim 12, wherein the sample is blood, serum, plasma, cerebrospinal fluid (CSF), urine, saliva, stool and synovial fluid.

18. The method of claim 17, wherein the sample is blood, serum or plasma.

* * * * *

专利名称(译)	瓜氨酸化的大脑和神经蛋白作为脑损伤或神经变性的生物标志物		
公开(公告)号	US20190310265A1	公开(公告)日	2019-10-10
申请号	US16/433106	申请日	2019-06-06
[标]申请(专利权)人(译)	约翰霍普金斯大学		
申请(专利权)人(译)	约翰·霍普金斯大学		
当前申请(专利权)人(译)	约翰·霍普金斯大学		
[标]发明人	VAN EYK JENNIFER E EVERETT ALLEN DALE JIN ZHICHENG		
发明人	VAN EYK, JENNIFER E. EVERETT, ALLEN DALE JIN, ZHICHENG		
IPC分类号	G01N33/68 H01J49/00 G01N33/53		
CPC分类号	G01N2440/18 G01N33/5306 G01N33/6848 G01N33/6842 H01J49/0027 G01N2800/28 H01J49/0077		
优先权	14/384848 2014-09-12 US PCT/US2013/031012 2013-03-13 WO 61/610034 2012-03-13 US		
外部链接	Espacenet USPTO		

摘要(译)

本发明涉及生物标志物领域。更具体地，本发明涉及可用于诊断脑损伤或神经变性的生物标志物。在一个实施例中，一种用于诊断患者的脑损伤的方法包括以下步骤：(a) 从患者获得样品；(b) 确定在一种或多种脑损伤生物标志物蛋白的一个或多个精氨酸残基上瓜氨酸化与未修饰的精氨酸残基的比例；(c) 将该比例与脑损伤患者或非脑损伤患者相关联，从而提供诊断。

