



US008298776B2

(12) **United States Patent**
Raitano et al.(10) **Patent No.: US 8,298,776 B2**
(45) **Date of Patent: *Oct. 30, 2012**(54) **ANTIBODIES TO TUMOR ASSOCIATED
PROTEINS**(75) Inventors: **Arthur B. Raitano**, Los Angeles, CA
(US); **Karen Jane Meyrick Morrison**,
Santa Monica, CA (US); **Wangmao Ge**,
Tampa, FL (US); **Pia M. Challita-Eid**,
Encino, CA (US); **Aya Jakobovits**,
Beverly Hills, CA (US)(73) Assignee: **Agensys, Inc.**, Santa Monica, CA (US)(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.This patent is subject to a terminal dis-
claimer.(21) Appl. No.: **12/829,256**(22) Filed: **Jul. 1, 2010**(65) **Prior Publication Data**

US 2010/0273174 A1 Oct. 28, 2010

Related U.S. Application Data(63) Continuation of application No. 12/694,209, filed on
Jan. 26, 2010, now Pat. No. 8,124,741, which is a
continuation of application No. 11/842,016, filed on
Aug. 20, 2007, now Pat. No. 7,858,750, which is a
continuation of application No. 11/518,610, filed on
Sep. 8, 2006, now Pat. No. 7,935,491, which is a
continuation of application No. 10/306,631, filed on
Nov. 27, 2002, now Pat. No. 7,244,827, which is a
continuation-in-part of application No. 09/547,789,
filed on Apr. 12, 2000, now Pat. No. 6,943,235.(60) Provisional application No. 60/128,858, filed on Apr.
12, 1999.(51) **Int. Cl.**
G01N 33/53 (2006.01)(52) **U.S. Cl.** **435/7.1**(58) **Field of Classification Search** None
See application file for complete search history.(56) **References Cited****U.S. PATENT DOCUMENTS**

4,742,000	A	5/1988	Greene
5,693,762	A	12/1997	Queen et al.
6,312,922	B1	11/2001	Edwards et al.
6,551,795	B1	4/2003	Rubenfield et al.
6,682,736	B1	1/2004	Hanson et al.
6,747,137	B1	6/2004	Weinstock et al.
6,913,919	B2	7/2005	Botstein et al.
6,930,170	B2	8/2005	Desnoyers et al.
6,943,235	B1	9/2005	Afar et al.
6,953,836	B2	10/2005	Desnoyers et al.
6,956,108	B2	10/2005	Desnoyers et al.
6,972,185	B2	12/2005	Desnoyers et al.
7,018,811	B2	3/2006	Botstein et al.
7,019,116	B2	3/2006	Desnoyers et al.
7,029,873	B2	4/2006	Desnoyers et al.
7,034,106	B2	4/2006	Desnoyers et al.

7,034,122	B2	4/2006	Desnoyers et al.
7,034,136	B2	4/2006	Goddard et al.
7,244,827	B2	7/2007	Raitano et al.
7,288,251	B2	10/2007	Bedian et al.
7,303,895	B1	12/2007	O'Regan et al.
7,378,492	B2	5/2008	Chawla et al.
7,935,491	B2 *	5/2011	Raitano et al. 435/7.1
2002/0022248	A1	2/2002	Xu et al.
2002/0103125	A1	8/2002	Ashkenazi et al.
2002/0119130	A1	8/2002	Eaton et al.
2002/0123463	A1	9/2002	Ashkenazi et al.
2002/0127576	A1	9/2002	Ashkenazi et al.
2002/0132252	A1	9/2002	Ashkenazi et al.
2002/0142961	A1	10/2002	Ashkenazi et al.
2002/0160384	A1	10/2002	Ashkenazi et al.
2002/0177164	A1	11/2002	Ashkenazi et al.
2002/0182638	A1	12/2002	Eaton et al.
2002/0183493	A1	12/2002	Eaton et al.
2002/0183494	A1	12/2002	Eaton et al.
2002/0192763	A1	12/2002	Xu et al.
2002/0193299	A1	12/2002	Ashkenazi et al.
2002/0193300	A1	12/2002	Ashkenazi et al.
2002/0197615	A1	12/2002	Ashkenazi et al.
2002/0198148	A1	12/2002	Ashkenazi et al.
2003/0003531	A1	1/2003	Ashkenazi et al.
2003/0008297	A1	1/2003	Ashkenazi et al.
2003/0009012	A1	1/2003	Eaton et al.
2003/0009013	A1	1/2003	Eaton et al.
2003/0013855	A1	1/2003	Eaton et al.

(Continued)

FOREIGN PATENT DOCUMENTSCA 2369413 10/2000
(Continued)**OTHER PUBLICATIONS**

Fadlon et al (British Journal of Cancer, 1996, 74: 400-405).*

Alberts et al., Molecular Biology of the Cell, 3rd. ed. (1994) p. 465.

Alzari et al., Annual Rev Immunol (1988) 6:555-580.

Benedict et al., J. Exp. Medicine (2001) 193(1):89-99.

Boehringer Mannheim Biochemicals, 1994 Catalog, p. 93.

Bowie et al., Science (1990) 247:1306-1310.

Brennan et al., Journal of Autoimmunity (1989) 2(suppl.):177-186.

Bruggemann et al., PNAS USA (1989) 86:6709-6713.

Busken et al., Digestive Disease Week Abstracts and Itinerary Plan-
ner (2003) Abstract No. 850.

Casset et al., Biochem. Biophys. Res. Commun. (2003) 307:198-205.

Chang et al., The Journal of Histochemistry and Cytochemistry
(1991) 39(9):1281-1287.

(Continued)

Primary Examiner — Sean Aeder(74) *Attorney, Agent, or Firm* — Morrison & Foerster LLP(57) **ABSTRACT**

A novel gene 024P4C12 (also designated 24P4C12) and its encoded protein, and variants thereof, are described wherein 24P4C12 exhibits tissue specific expression in normal adult tissue, and is aberrantly expressed in the cancers listed in Table I. Consequently, 24P4C12 provides a diagnostic, prognostic, prophylactic and/or therapeutic target for cancer. The 24P4C12 gene or fragment thereof, or its encoded protein, or variants thereof, or a fragment thereof, can be used to elicit a humoral or cellular immune response; antibodies or T cells reactive with 24P4C12 can be used in active or passive immunization.

3 Claims, 58 Drawing Sheets

U.S. PATENT DOCUMENTS

2003/0017476 A1 1/2003 Ashkenazi et al.
 2003/0017981 A1 1/2003 Ashkenazi et al.
 2003/0017982 A1 1/2003 Ashkenazi et al.
 2003/0018168 A1 1/2003 Eaton et al.
 2003/0018172 A1 1/2003 Eaton et al.
 2003/0018173 A1 1/2003 Eaton et al.
 2003/0018183 A1 1/2003 Eaton et al.
 2003/0022187 A1 1/2003 Ashkenazi et al.
 2003/0023042 A1 1/2003 Eaton et al.
 2003/0027162 A1 2/2003 Ashkenazi et al.
 2003/0027163 A1 2/2003 Ashkenazi et al.
 2003/0027212 A1 2/2003 Eaton et al.
 2003/0027754 A1 2/2003 Ashkenazi et al.
 2003/0027985 A1 2/2003 Ashkenazi et al.
 2003/0027986 A1 2/2003 Eaton et al.
 2003/0027992 A1 2/2003 Eaton et al.
 2003/0027993 A1 2/2003 Eaton et al.
 2003/0032023 A1 2/2003 Ashkenazi et al.
 2003/0036634 A1 2/2003 Eaton et al.
 2003/0040473 A1 2/2003 Ashkenazi et al.
 2003/0044806 A1 3/2003 Ashkenazi et al.
 2003/0045463 A1 3/2003 Ashkenazi et al.
 2003/0045684 A1 3/2003 Eaton et al.
 2003/0049638 A1 3/2003 Ashkenazi et al.
 2003/0049681 A1 3/2003 Ashkenazi et al.
 2003/0049682 A1 3/2003 Ashkenazi et al.
 2003/0049735 A1 3/2003 Eaton et al.
 2003/0050462 A1 3/2003 Eaton et al.
 2003/0050465 A1 3/2003 Eaton et al.
 2003/0054359 A1 3/2003 Ashkenazi et al.
 2003/0054403 A1 3/2003 Ashkenazi et al.
 2003/0054404 A1 3/2003 Ashkenazi et al.
 2003/0054987 A1 3/2003 Ashkenazi et al.
 2003/0059780 A1 3/2003 Ashkenazi et al.
 2003/0059782 A1 3/2003 Ashkenazi et al.
 2003/0059783 A1 3/2003 Ashkenazi et al.
 2003/0059831 A1 3/2003 Ashkenazi et al.
 2003/0059832 A1 3/2003 Ashkenazi et al.
 2003/0059833 A1 3/2003 Ashkenazi et al.
 2003/0060407 A1 3/2003 Ashkenazi et al.
 2003/0060600 A1 3/2003 Eaton et al.
 2003/0060601 A1 3/2003 Eaton et al.
 2003/0060602 A1 3/2003 Eaton et al.
 2003/0064375 A1 4/2003 Ashkenazi et al.
 2003/0065143 A1 4/2003 Eaton et al.
 2003/0065161 A1 4/2003 Eaton et al.
 2003/0068623 A1 4/2003 Ashkenazi et al.
 2003/0068647 A1 4/2003 Ashkenazi et al.
 2003/0069394 A1 4/2003 Eaton et al.
 2003/0069403 A1 4/2003 Ashkenazi et al.
 2003/0073090 A1 4/2003 Ashkenazi et al.
 2003/0073623 A1 4/2003 Drmanac et al.
 2003/0078387 A1 4/2003 Eaton et al.
 2003/0082546 A1 5/2003 Ashkenazi et al.
 2003/0083461 A1 5/2003 Ashkenazi et al.
 2003/0083473 A1 5/2003 Eaton et al.
 2003/0087304 A1 5/2003 Ashkenazi et al.
 2003/0087305 A1 5/2003 Ashkenazi et al.
 2003/0099974 A1 5/2003 Lillie et al.
 2003/0147904 A1 8/2003 Afar et al.
 2003/0211100 A1 11/2003 Bedian et al.
 2004/0228858 A1 11/2004 Hanson et al.
 2005/0019870 A1 1/2005 Afar et al.
 2007/0004910 A1 1/2007 Sexton et al.
 2007/0027308 A1 2/2007 Edwards et al.
 2007/0160617 A1 7/2007 Ma et al.
 2008/0311107 A1 12/2008 Bollinger et al.

FOREIGN PATENT DOCUMENTS

EP 1 033 401 9/2000
 EP 1 074 617 2/2001
 WO WO-99/06548 2/1999
 WO WO-99/06549 2/1999
 WO WO-99/06550 2/1999
 WO WO-99/40189 8/1999
 WO WO-99/63088 12/1999
 WO WO-00/04149 1/2000

WO WO 00/61746 * 10/2000
 WO WO-00/61746 10/2000
 WO WO-00/73454 12/2000
 WO WO-00/77021 12/2000
 WO WO-01/16318 3/2001
 WO WO-01/25272 4/2001
 WO WO-01/34802 5/2001
 WO WO-01/46258 6/2001
 WO WO-01/51628 7/2001
 WO WO-01/51633 7/2001
 WO WO-01/53836 7/2001
 WO WO-01/57270 8/2001
 WO WO-01/57271 8/2001
 WO WO-01/57272 8/2001
 WO WO-01/57273 8/2001
 WO WO-01/57274 8/2001
 WO WO-01/57275 8/2001
 WO WO-01/57276 8/2001
 WO WO-01/57277 8/2001
 WO WO-01/57278 8/2001
 WO WO-01/60860 8/2001
 WO WO-01/73027 10/2001
 WO WO-01/73032 10/2001
 WO WO-01/75067 10/2001
 WO WO-01/86003 11/2001
 WO WO-01/90304 11/2001
 WO WO-01/96388 12/2001
 WO WO-01/96390 12/2001
 WO WO-02/12328 2/2002
 WO WO-02/58534 8/2002
 WO WO-02/74961 9/2002
 WO WO-02/83876 10/2002
 WO WO-02/89747 11/2002
 WO WO-02/97031 12/2002
 WO WO-2009/033094 3/2009

OTHER PUBLICATIONS

Craft et al., *Cancer Res.* (1999) 59:5030-5036.
 Dermer, *Bio/Technology* (1994) 12:320.
 Dictionary Definition of "coupled to" from Merriam-Webster online, visited Jan. 16, 2010.
 Drexler et al., *Leukemia and Lymphoma* (1993) 9:1-25.
 Dulcert et al., *Accession No. AAY12282*, 1999.
 Ericksson et al., *Diabetologia* (1992) 35:143-147.
 Ezzell, *Journal of NIH Research* (1995) 7:46-49.
 Freshney, *Culture of Animal Cells, A Manual of Basic Technique*, Alan R. Liss, Inc., New York (1983) p. 4.
 Fu et al., *EMBO Journal* (1996) 15:4392-4401.
 Greenbaum et al., *Genome Biology* (2003) 4(9):117.1-117.8.
 Gress et al., *Oncogene* (1996) 13:1818-1830.
 Gura, *Science* (1997) 278:1041-1042.
 Gussow and Seemann, *Methods in Enzymology* (1991) 203:99-121.
 Harlow and Lane (eds.), *Antibodies: A Laboratory Manual*, Cold Spring Harbor (1988) pp. 591-598.
 Hirashima et al., *Int. Arch. Allergy Immunol.* (2000) Suppl.1:6-9.
 Hsu, in *Tissue Culture Methods and Applications*, Kruse and Patterson, eds. (1973) Academic Press, Abstract p. 764.
 Huang, G.M. et al., "Prostate cancer expression profiling by cDNA sequencing analysis," EMBL Database entry AI557659, *Accession No. AI557659*, Mar. 25, 1999, EP002144281, & Huang, G.M. et al., *Genomics*, vol. 59, No. 2, Jul. 1999, pp. 178-186.
 Huang, Guyang Matthew, Aug. 9, 1999, dbEST Id 2373824, GenBank Acc. AI557660.
 Hubert et al., *PNAS USA* (1999) 96(25):14523-14528.
 Huston et al., *PNAS USA* (1988) 85:5879-5883.
 Inoko, Hidetoshi, Mar. 30, 2000, NCBI Accession No. AP000502.
 International Search Report for PCT/US02/38264, mailed on Oct. 20, 2004, 3 pages.
 Jakobovits, *Expert Opinion on Investigational Drugs* (1998) 7(4):607-614.
 Jiang et al., *JBC* (2003) 278(7):4763-4769.
 Johnstone and Thorpe, *Immunocytochemistry in Practice*, 2nd edition, Blackwell Scientific Publications, Oxford (1987) pp. 113-130.
 Kerlavage, A. R., Apr. 21, 1997, dbEST Id 1008183, GenBank Acc. AA 366876.
 Kilty and Amara, *Curr. Opin. Biotechnology* (1992) 3:675-682.

Figure 2C-3

```
1921 TGCCCATCATGACCTCCATCCTGGGGGCCTATGTCATCGCCAGCGGCTTCTTCAGCGTTT
    660  G M C V D T L F L C F L E D L E R N N G
1981 TCGGCATGTGTGTGGACACGCTCTTCTCTGCTTCCTGGAAGACCTGGAGCGGAACAACG
    680  S L D R P Y Y M S K S L L K I L G K K N
2041 GCTCCCTGGACCGGCCCTACTACATGTCCAAGAGCCTTCTAAAGATTCTGGGCAAGAAGA
    700  E A F P D N K K R K K *
2101 ACGAGGCGCCCCCGGACAACAAGAAGAGGAAGAAGTGAcagctccggccctgatccagga
2161 ctgcacccccacccccaccgtccagccatccaacctcacttcgccttacaggtctccattt
2221 tgtggtaaaaaaaggtttttaggccaggcgccgtggctcacgcctgtaatccaacactttg
2281 agaggttgaggcgggcggtacacctgagtcaggagttcgagaccagcctggccaacatgg
2341 tgaacctccgtctctattataaaatacaaaaattagccgagagtgggtggcatgcacctgt
2401 catcccagctactcgggaggtgaggcaggagaatcgcttgaaacccgggaggcagaggtt
2461 gcagtgagccgagatcgcgccactgcactccaacctgggtgacagactctgtctccaaaa
2521 caaaacaaacaaacaaaaagatttttattaaagatatatttgtaactcagtaaaaaaaaaa
2581 aaaaaaa
```

Figure 2D. The cDNA (SEQ ID. NO. : 8) and amino acid sequence (SEQ ID. NO. : 9) of 24P4C12 v.4.

The start methionine is underlined. The open reading frame extends from nucleic acid 6-2138 including the stop codon.

Figure 2D-1

```
1      M G G K Q R D E D D E A Y G K P V K Y
1 gagccATGGGGGAAAGCAGCGGGACGAGGATGACGAGGCCTACGGGAAGCCAGTCAAAT
20   D P S F R G P I K N R S C T D V I C C V
61 ACGACCCCTCCTTTCGAGGCCCCATCAAGAACAGAAGCTGCACAGATGTCATCTGCTGCG
40   L F L L F I L G Y I V V G I V A W L Y G
121 TCCTCTTCCTGCTCTTCATTCTAGGTTACATCGTGGTGGGGATTGTGGCCTGGTTGTATG
60   D F R Q V L Y P R N S T G A Y C G M G E
181 GAGACCCCGGCAAGTCCTCTACCCCAAGAACTCTACTGGGGCCTACTGTGGCATGGGGG
80   N K D K P Y L L Y F N I F S C I L S S N
241 AGAACAAAGATAAGCCGTATCTCCTGTACTTCAACATCTTCAGCTGCATCCTGTCCAGCA
100  I I S V A E N G L Q C P T P Q V C V S S
301 ACATCATCTCAGTTGCTGAGAACGGCCTACAGTGCCCCACACCCAGGTGTGTGTGTCCT
120  C P E D P W T V G K N E F S Q T V G E V
361 CCTGCCCCGAGGACCCATGGACTGTGGGAAAAAACGAGTTCTCACAGACTGTTGGGGAAG
140  F Y T K N R N F C L P G V P W N M T V I
421 TCTTCTATACAAAAACAGGAACCTTTTGTCTGCCAGGGGTACCCTGGAATATGACGGTGA
160  T S L Q Q E L C P S F L L P S A P A L G
481 TCACAAGCCTGCAACAGGAACCTTGCCCCAGTTTCCTCCTCCCTCTGCTCCAGCTCTGG
180  R C F P W T N V T P P A L P G I T N D T
541 GGCGCTGCTTTCCATGGACCAACGTTACTCCACCGGCGCTCCAGGGATCACCAATGACA
```

protein specifically binds to 24P4C12 and blocks 24P4C12 interaction with a binding partner. Such dimeric fusion proteins are further combined into multimeric proteins using known antibody linking technologies.

XII.C.) Inhibition of 24P4C12 Transcription or Translation

The present invention also comprises various methods and compositions for inhibiting the transcription of the 24P4C12 gene. Similarly, the invention also provides methods and compositions for inhibiting the translation of 24P4C12 mRNA into protein.

In one approach, a method of inhibiting the transcription of the 24P4C12 gene comprises contacting the 24P4C12 gene with a 24P4C12 antisense polynucleotide. In another approach, a method of inhibiting 24P4C12 mRNA translation comprises contacting a 24P4C12 mRNA with an antisense polynucleotide. In another approach, a 24P4C12 specific ribozyme is used to cleave a 24P4C12 message, thereby inhibiting translation. Such antisense and ribozyme based methods can also be directed to the regulatory regions of the 24P4C12 gene, such as 24P4C12 promoter and/or enhancer elements. Similarly, proteins capable of inhibiting a 24P4C12 gene transcription factor are used to inhibit 24P4C12 mRNA transcription. The various polynucleotides and compositions useful in the aforementioned methods have been described above. The use of antisense and ribozyme molecules to inhibit transcription and translation is well known in the art.

Other factors that inhibit the transcription of 24P4C12 by interfering with 24P4C12 transcriptional activation are also useful to treat cancers expressing 24P4C12. Similarly, factors that interfere with 24P4C12 processing are useful to treat cancers that express 24P4C12. Cancer treatment methods utilizing such factors are also within the scope of the invention.

XII.D.) General Considerations for Therapeutic Strategies

Gene transfer and gene therapy technologies can be used to deliver therapeutic polynucleotide molecules to tumor cells synthesizing 24P4C12 (i.e., antisense, ribozyme, polynucleotides encoding intrabodies and other 24P4C12 inhibitory molecules). A number of gene therapy approaches are known in the art. Recombinant vectors encoding 24P4C12 antisense polynucleotides, ribozymes, factors capable of interfering with 24P4C12 transcription, and so forth, can be delivered to target tumor cells using such gene therapy approaches.

The above therapeutic approaches can be combined with any one of a wide variety of surgical, chemotherapy or radiation therapy regimens. The therapeutic approaches of the invention can enable the use of reduced dosages of chemotherapy (or other therapies) and/or less frequent administration, an advantage for all patients and particularly for those that do not tolerate the toxicity of the chemotherapeutic agent well.

The anti-tumor activity of a particular composition (e.g., antisense, ribozyme, intrabody), or a combination of such compositions, can be evaluated using various in vitro and in vivo assay systems. In vitro assays that evaluate therapeutic activity include cell growth assays, soft agar assays and other assays indicative of tumor promoting activity, binding assays capable of determining the extent to which a therapeutic composition will inhibit the binding of 24P4C12 to a binding partner, etc.

In vivo, the effect of a 24P4C12 therapeutic composition can be evaluated in a suitable animal model. For example, xenogenic prostate cancer models can be used, wherein human prostate cancer explants or passaged xenograft tissues are introduced into immune compromised animals, such as nude or SCID mice (Klein et al., 1997, *Nature Medicine* 3: 402-408). For example, PCT Patent Application WO98/

16628 and U.S. Pat. No. 6,107,540 describe various xenograft models of human prostate cancer capable of recapitulating the development of primary tumors, micrometastasis, and the formation of osteoblastic metastases characteristic of late stage disease. Efficacy can be predicted using assays that measure inhibition of tumor formation, tumor regression or metastasis, and the like.

In vivo assays that evaluate the promotion of apoptosis are useful in evaluating therapeutic compositions. In one embodiment, xenografts from tumor bearing mice treated with the therapeutic composition can be examined for the presence of apoptotic foci and compared to untreated control xenograft-bearing mice. The extent to which apoptotic foci are found in the tumors of the treated mice provides an indication of the therapeutic efficacy of the composition.

The therapeutic compositions used in the practice of the foregoing methods can be formulated into pharmaceutical compositions comprising a carrier suitable for the desired delivery method. Suitable carriers include any material that when combined with the therapeutic composition retains the anti-tumor function of the therapeutic composition and is generally non-reactive with the patient's immune system. Examples include, but are not limited to, any of a number of standard pharmaceutical carriers such as sterile phosphate buffered saline solutions, bacteriostatic water, and the like (see, generally, Remington's *Pharmaceutical Sciences* 16th Edition, A. Osal., Ed., 1980).

Therapeutic formulations can be solubilized and administered via any route capable of delivering the therapeutic composition to the tumor site. Potentially effective routes of administration include, but are not limited to, intravenous, parenteral, intraperitoneal, intramuscular, intratumor, intradermal, intraorgan, orthotopic, and the like. A preferred formulation for intravenous injection comprises the therapeutic composition in a solution of preserved bacteriostatic water, sterile unpreserved water, and/or diluted in polyvinylchloride or polyethylene bags containing 0.9% sterile Sodium Chloride for Injection, USP. Therapeutic protein preparations can be lyophilized and stored as sterile powders, preferably under vacuum, and then reconstituted in bacteriostatic water (containing for example, benzyl alcohol preservative) or in sterile water prior to injection.

Dosages and administration protocols for the treatment of cancers using the foregoing methods will vary with the method and the target cancer, and will generally depend on a number of other factors appreciated in the art.

XIII.) IDENTIFICATION, CHARACTERIZATION AND USE OF MODULATORS OF 24P4C12

Methods to Identify and Use Modulators

In one embodiment, screening is performed to identify modulators that induce or suppress a particular expression profile, suppress or induce specific pathways, preferably generating the associated phenotype thereby. In another embodiment, having identified differentially expressed genes important in a particular state; screens are performed to identify modulators that alter expression of individual genes, either increase or decrease. In another embodiment, screening is performed to identify modulators that alter a biological function of the expression product of a differentially expressed gene. Again, having identified the importance of a gene in a particular state, screens are performed to identify agents that bind and/or modulate the biological activity of the gene product.

In addition, screens are done for genes that are induced in response to a candidate agent. After identifying a modulator

(one that suppresses a cancer expression pattern leading to a normal expression pattern, or a modulator of a cancer gene that leads to expression of the gene as in normal tissue) a screen is performed to identify genes that are specifically modulated in response to the agent. Comparing expression profiles between normal tissue and agent-treated cancer tissue reveals genes that are not expressed in normal tissue or cancer tissue, but are expressed in agent treated tissue, and vice versa. These agent-specific sequences are identified and used by methods described herein for cancer genes or proteins. In particular these sequences and the proteins they encode are used in marking or identifying agent-treated cells. In addition, antibodies are raised against the agent-induced proteins and used to target novel therapeutics to the treated cancer tissue sample.

Modulator-Related Identification and Screening Assays: Gene Expression-Related Assays

Proteins, nucleic acids, and antibodies of the invention are used in screening assays. The cancer-associated proteins, antibodies, nucleic acids, modified proteins and cells containing these sequences are used in screening assays, such as evaluating the effect of drug candidates on a "gene expression profile," expression profile of polypeptides or alteration of biological function. In one embodiment, the expression profiles are used, preferably in conjunction with high throughput screening techniques to allow monitoring for expression profile genes after treatment with a candidate agent (e.g., Davis, G F, et al, J Biol Screen 7:69 (2002); Zlokarnik, et al., Science 279:84-8 (1998); Heid, Genome Res 6:986-94, 1996).

The cancer proteins, antibodies, nucleic acids, modified proteins and cells containing the native or modified cancer proteins or genes are used in screening assays. That is, the present invention comprises methods for screening for compositions which modulate the cancer phenotype or a physiological function of a cancer protein of the invention. This is done on a gene itself or by evaluating the effect of drug candidates on a "gene expression profile" or biological function. In one embodiment, expression profiles are used, preferably in conjunction with high throughput screening techniques to allow monitoring after treatment with a candidate agent, see Zlokarnik, supra.

A variety of assays are executed directed to the genes and proteins of the invention. Assays are run on an individual nucleic acid or protein level. That is, having identified a particular gene as up regulated in cancer, test compounds are screened for the ability to modulate gene expression or for binding to the cancer protein of the invention. "Modulation" in this context includes an increase or a decrease in gene expression. The preferred amount of modulation will depend on the original change of the gene expression in normal versus tissue undergoing cancer, with changes of at least 10%, preferably 50%, more preferably 100-300%, and in some embodiments 300-1000% or greater. Thus, if a gene exhibits a 4-fold increase in cancer tissue compared to normal tissue, a decrease of about four-fold is often desired; similarly, a 10-fold decrease in cancer tissue compared to normal tissue a target value of a 10-fold increase in expression by the test compound is often desired. Modulators that exacerbate the type of gene expression seen in cancer are also useful, e.g., as an upregulated target in further analyses.

The amount of gene expression is monitored using nucleic acid probes and the quantification of gene expression levels, or, alternatively, a gene product itself is monitored, e.g., through the use of antibodies to the cancer protein and standard immunoassays. Proteomics and separation techniques also allow for quantification of expression.

Expression Monitoring to Identify Compounds that Modify Gene Expression

In one embodiment, gene expression monitoring, i.e., an expression profile, is monitored simultaneously for a number of entities. Such profiles will typically involve one or more of the genes of FIG. 2. In this embodiment, e.g., cancer nucleic acid probes are attached to biochips to detect and quantify cancer sequences in a particular cell. Alternatively, PCR can be used. Thus, a series, e.g., wells of a microtiter plate, can be used with dispensed primers in desired wells. A PCR reaction can then be performed and analyzed for each well.

Expression monitoring is performed to identify compounds that modify the expression of one or more cancer-associated sequences, e.g., a polynucleotide sequence set out in FIG. 2. Generally, a test modulator is added to the cells prior to analysis. Moreover, screens are also provided to identify agents that modulate cancer, modulate cancer proteins of the invention, bind to a cancer protein of the invention, or interfere with the binding of a cancer protein of the invention and an antibody or other binding partner.

In one embodiment, high throughput screening methods involve providing a library containing a large number of potential therapeutic compounds (candidate compounds). Such "combinatorial chemical libraries" are then screened in one or more assays to identify those library members (particular chemical species or subclasses) that display a desired characteristic activity. The compounds thus identified can serve as conventional "lead compounds," as compounds for screening, or as therapeutics.

In certain embodiments, combinatorial libraries of potential modulators are screened for an ability to bind to a cancer polypeptide or to modulate activity. Conventionally, new chemical entities with useful properties are generated by identifying a chemical compound (called a "lead compound") with some desirable property or activity, e.g., inhibiting activity, creating variants of the lead compound, and evaluating the property and activity of those variant compounds. Often, high throughput screening (HTS) methods are employed for such an analysis.

As noted above, gene expression monitoring is conveniently used to test candidate modulators (e.g., protein, nucleic acid or small molecule). After the candidate agent has been added and the cells allowed to incubate for a period, the sample containing a target sequence to be analyzed is, e.g., added to a biochip.

If required, the target sequence is prepared using known techniques. For example, a sample is treated to lyse the cells, using known lysis buffers, electroporation, etc., with purification and/or amplification such as PCR performed as appropriate. For example, an in vitro transcription with labels covalently attached to the nucleotides is performed. Generally, the nucleic acids are labeled with biotin-FITC or PE, or with cy3 or cy5.

The target sequence can be labeled with, e.g., a fluorescent, a chemiluminescent, a chemical, or a radioactive signal, to provide a means of detecting the target sequence's specific binding to a probe. The label also can be an enzyme, such as alkaline phosphatase or horseradish peroxidase, which when provided with an appropriate substrate produces a product that is detected. Alternatively, the label is a labeled compound or small molecule, such as an enzyme inhibitor, that binds but is not catalyzed or altered by the enzyme. The label also can be a moiety or compound, such as, an epitope tag or biotin which specifically binds to streptavidin. For the example of biotin, the streptavidin is labeled as described above, thereby,

providing a detectable signal for the bound target sequence. Unbound labeled streptavidin is typically removed prior to analysis.

As will be appreciated by those in the art, these assays can be direct hybridization assays or can comprise "sandwich assays", which include the use of multiple probes, as is generally outlined in U.S. Pat. Nos. 5,681,702; 5,597,909; 5,545,730; 5,594,117; 5,591,584; 5,571,670; 5,580,731; 5,571,670; 5,591,584; 5,624,802; 5,635,352; 5,594,118; 5,359,100; 5,124,246; and 5,681,697. In this embodiment, in general, the target nucleic acid is prepared as outlined above, and then added to the biochip comprising a plurality of nucleic acid probes, under conditions that allow the formation of a hybridization complex.

A variety of hybridization conditions are used in the present invention, including high, moderate and low stringency conditions as outlined above. The assays are generally run under stringency conditions which allow formation of the label probe hybridization complex only in the presence of target. Stringency can be controlled by altering a step parameter that is a thermodynamic variable, including, but not limited to, temperature, formamide concentration, salt concentration, chaotropic salt concentration pH, organic solvent concentration, etc. These parameters may also be used to control non-specific binding, as is generally outlined in U.S. Pat. No. 5,681,697. Thus, it can be desirable to perform certain steps at higher stringency conditions to reduce non-specific binding.

The reactions outlined herein can be accomplished in a variety of ways. Components of the reaction can be added simultaneously, or sequentially, in different orders, with preferred embodiments outlined below. In addition, the reaction may include a variety of other reagents. These include salts, buffers, neutral proteins, e.g. albumin, detergents, etc. which can be used to facilitate optimal hybridization and detection, and/or reduce nonspecific or background interactions. Reagents that otherwise improve the efficiency of the assay, such as protease inhibitors, nuclease inhibitors, anti-microbial agents, etc., may also be used as appropriate, depending on the sample preparation methods and purity of the target. The assay data are analyzed to determine the expression levels of individual genes, and changes in expression levels as between states, forming a gene expression profile.

Biological Activity-Related Assays

The invention provides methods identify or screen for a compound that modulates the activity of a cancer-related gene or protein of the invention. The methods comprise adding a test compound, as defined above, to a cell comprising a cancer protein of the invention. The cells contain a recombinant nucleic acid that encodes a cancer protein of the invention. In another embodiment, a library of candidate agents is tested on a plurality of cells.

In one aspect, the assays are evaluated in the presence or absence or previous or subsequent exposure of physiological signals, e.g. hormones, antibodies, peptides, antigens, cytokines, growth factors, action potentials, pharmacological agents including chemotherapeutics, radiation, carcinogenics, or other cells (i.e., cell-cell contacts). In another example, the determinations are made at different stages of the cell cycle process. In this way, compounds that modulate genes or proteins of the invention are identified. Compounds with pharmacological activity are able to enhance or interfere with the activity of the cancer protein of the invention. Once identified, similar structures are evaluated to identify critical structural features of the compound.

In one embodiment, a method of modulating (e.g., inhibiting) cancer cell division is provided; the method comprises

administration of a cancer modulator. In another embodiment, a method of modulating (e.g., inhibiting) cancer is provided; the method comprises administration of a cancer modulator. In a further embodiment, methods of treating cells or individuals with cancer are provided; the method comprises administration of a cancer modulator.

In one embodiment, a method for modulating the status of a cell that expresses a gene of the invention is provided. As used herein status comprises such art-accepted parameters such as growth, proliferation, survival, function, apoptosis, senescence, location, enzymatic activity, signal transduction, etc. of a cell. In one embodiment, a cancer inhibitor is an antibody as discussed above. In another embodiment, the cancer inhibitor is an antisense molecule. A variety of cell growth, proliferation, and metastasis assays are known to those of skill in the art, as described herein.

High Throughput Screening to Identify Modulators

The assays to identify suitable modulators are amenable to high throughput screening. Preferred assays thus detect enhancement or inhibition of cancer gene transcription, inhibition or enhancement of polypeptide expression, and inhibition or enhancement of polypeptide activity.

In one embodiment, modulators evaluated in high throughput screening methods are proteins, often naturally occurring proteins or fragments of naturally occurring proteins. Thus, e.g., cellular extracts containing proteins, or random or directed digests of proteinaceous cellular extracts, are used. In this way, libraries of proteins are made for screening in the methods of the invention. Particularly preferred in this embodiment are libraries of bacterial, fungal, viral, and mammalian proteins, with the latter being preferred, and human proteins being especially preferred. Particularly useful test compound will be directed to the class of proteins to which the target belongs, e.g., substrates for enzymes, or ligands and receptors.

Use of Soft Agar Growth and Colony Formation to Identify and Characterize Modulators

Normal cells require a solid substrate to attach and grow. When cells are transformed, they lose this phenotype and grow detached from the substrate. For example, transformed cells can grow in stirred suspension culture or suspended in semi-solid media, such as semi-solid or soft agar. The transformed cells, when transfected with tumor suppressor genes, can regenerate normal phenotype and once again require a solid substrate to attach to and grow. Soft agar growth or colony formation in assays are used to identify modulators of cancer sequences, which when expressed in host cells, inhibit abnormal cellular proliferation and transformation. A modulator reduces or eliminates the host cells' ability to grow suspended in solid or semisolid media, such as agar.

Techniques for soft agar growth or colony formation in suspension assays are described in Freshney, *Culture of Animal Cells a Manual of Basic Technique* (3rd ed., 1994). See also, the methods section of Garkavtsev et al. (1996), supra.

Evaluation of Contact Inhibition and Growth Density Limitation to Identify and Characterize Modulators

Normal cells typically grow in a flat and organized pattern in cell culture until they touch other cells. When the cells touch one another, they are contact inhibited and stop growing. Transformed cells, however, are not contact inhibited and continue to grow to high densities in disorganized foci. Thus, transformed cells grow to a higher saturation density than corresponding normal cells. This is detected morphologically by the formation of a disoriented monolayer of cells or cells in foci. Alternatively, labeling index with (³H)-thymidine at saturation density is used to measure density limitation of growth, similarly an MTT or Alamar blue assay will reveal

proliferation capacity of cells and the ability of modulators to affect same. See Freshney (1994), *supra*. Transformed cells, when transfected with tumor suppressor genes, can regenerate a normal phenotype and become contact inhibited and would grow to a lower density.

In this assay, labeling index with 3H-thymidine at saturation density is a preferred method of measuring density limitation of growth. Transformed host cells are transfected with a cancer-associated sequence and are grown for 24 hours at saturation density in non-limiting medium conditions. The percentage of cells labeling with (³H)-thymidine is determined by incorporated cpm.

Contact independent growth is used to identify modulators of cancer sequences, which had led to abnormal cellular proliferation and transformation. A modulator reduces or eliminates contact independent growth, and returns the cells to a normal phenotype.

Evaluation of Growth Factor or Serum Dependence to Identify and Characterize Modulators

Transformed cells have lower serum dependence than their normal counterparts (see, e.g., Temin, *J. Natl. Cancer Inst.* 37:167-175 (1966); Eagle et al., *J. Exp. Med.* 131:836-879 (1970)); Freshney, *supra*. This is in part due to release of various growth factors by the transformed cells. The degree of growth factor or serum dependence of transformed host cells can be compared with that of control. For example, growth factor or serum dependence of a cell is monitored in methods to identify and characterize compounds that modulate cancer-associated sequences of the invention.

Use of Tumor-Specific Marker Levels to Identify and Characterize Modulators

Tumor cells release an increased amount of certain factors (hereinafter "tumor specific markers") than their normal counterparts. For example, plasminogen activator (PA) is released from human glioma at a higher level than from normal brain cells (see, e.g., Gullino, *Angiogenesis, Tumor Vascularization, and Potential Interference with Tumor Growth*, in *Biological Responses in Cancer*, pp. 178-184 (Mihich (ed.) 1985)). Similarly, Tumor Angiogenesis Factor (TAF) is released at a higher level in tumor cells than their normal counterparts. See, e.g., Folkman, *Angiogenesis and Cancer*, *Sem Cancer Biol.* (1992)), while bFGF is released from endothelial tumors (Ensoli, B et al).

Various techniques which measure the release of these factors are described in Freshney (1994), *supra*. Also, see, Unkless et al., *J. Biol. Chem.* 249:4295-4305 (1974); Strickland & Beers, *J. Biol. Chem.* 251:5694-5702 (1976); Whur et al., *Br. J. Cancer* 42:305 312 (1980); Gullino, *Angiogenesis, Tumor Vascularization, and Potential Interference with Tumor Growth*, in *Biological Responses in Cancer*, pp. 178-184 (Mihich (ed.) 1985); Freshney, *Anticancer Res.* 5:111-130 (1985). For example, tumor specific marker levels are monitored in methods to identify and characterize compounds that modulate cancer-associated sequences of the invention.

Invasiveness into Matrigel to Identify and Characterize Modulators

The degree of invasiveness into Matrigel or an extracellular matrix constituent can be used as an assay to identify and characterize compounds that modulate cancer associated sequences. Tumor cells exhibit a positive correlation between malignancy and invasiveness of cells into Matrigel or some other extracellular matrix constituent. In this assay, tumorigenic cells are typically used as host cells. Expression of a tumor suppressor gene in these host cells would decrease invasiveness of the host cells. Techniques described in *Cancer Res.* 1999; 59:6010; Freshney (1994), *supra*, can be used.

Briefly, the level of invasion of host cells is measured by using filters coated with Matrigel or some other extracellular matrix constituent. Penetration into the gel, or through to the distal side of the filter, is rated as invasiveness, and rated histologically by number of cells and distance moved, or by prelabeling the cells with ¹²⁵I and counting the radioactivity on the distal side of the filter or bottom of the dish. See, e.g., Freshney (1984), *supra*.

Evaluation of Tumor Growth In Vivo to Identify and Characterize Modulators

Effects of cancer-associated sequences on cell growth are tested in transgenic or immune-suppressed organisms. Transgenic organisms are prepared in a variety of art-accepted ways. For example, knock-out transgenic organisms, e.g., mammals such as mice, are made, in which a cancer gene is disrupted or in which a cancer gene is inserted. Knock-out transgenic mice are made by insertion of a marker gene or other heterologous gene into the endogenous cancer gene site in the mouse genome via homologous recombination. Such mice can also be made by substituting the endogenous cancer gene with a mutated version of the cancer gene, or by mutating the endogenous cancer gene, e.g., by exposure to carcinogens.

To prepare transgenic chimeric animals, e.g., mice, a DNA construct is introduced into the nuclei of embryonic stem cells. Cells containing the newly engineered genetic lesion are injected into a host mouse embryo, which is re-implanted into a recipient female. Some of these embryos develop into chimeric mice that possess germ cells some of which are derived from the mutant cell line. Therefore, by breeding the chimeric mice it is possible to obtain a new line of mice containing the introduced genetic lesion (see, e.g., Capecchi et al., *Science* 244:1288 (1989)). Chimeric mice can be derived according to U.S. Pat. No. 6,365,797, issued 2 Apr. 2002; U.S. Pat. No. 6,107,540 issued 22 Aug. 2000; Hogan et al., *Manipulating the Mouse Embryo: A laboratory Manual*, Cold Spring Harbor Laboratory (1988) and *Teratocarcinomas and Embryonic Stem Cells: A Practical Approach*, Robertson, ed., IRL Press, Washington, D.C., (1987).

Alternatively, various immune-suppressed or immune-deficient host animals can be used. For example, a genetically athymic "nude" mouse (see, e.g., Giovanella et al., *J. Natl. Cancer Inst.* 52:921 (1974)), a SCID mouse, a thymectomized mouse, or an irradiated mouse (see, e.g., Bradley et al., *Br. J. Cancer* 38:263 (1978); Selby et al., *Br. J. Cancer* 41:52 (1980)) can be used as a host. Transplantable tumor cells (typically about 10⁶ cells) injected into isogenic hosts produce invasive tumors in a high proportion of cases, while normal cells of similar origin will not. In hosts which developed invasive tumors, cells expressing cancer-associated sequences are injected subcutaneously or orthotopically. Mice are then separated into groups, including control groups and treated experimental groups) e.g. treated with a modulator. After a suitable length of time, preferably 4-8 weeks, tumor growth is measured (e.g., by volume or by its two largest dimensions, or weight) and compared to the control. Tumors that have statistically significant reduction (using, e.g., Student's T test) are said to have inhibited growth.

In Vitro Assays to Identify and Characterize Modulators

Assays to identify compounds with modulating activity can be performed in vitro. For example, a cancer polypeptide is first contacted with a potential modulator and incubated for a suitable amount of time, e.g., from 0.5 to 48 hours. In one embodiment, the cancer polypeptide levels are determined in vitro by measuring the level of protein or mRNA. The level of protein is measured using immunoassays such as Western blotting, ELISA and the like with an antibody that selectively

binds to the cancer polypeptide or a fragment thereof. For measurement of mRNA, amplification, e.g., using PCR, LCR, or hybridization assays, e.g., Northern hybridization, RNase protection, dot blotting, are preferred. The level of protein or mRNA is detected using directly or indirectly labeled detection agents, e.g., fluorescently or radioactively labeled nucleic acids, radioactively or enzymatically labeled antibodies, and the like, as described herein.

Alternatively, a reporter gene system can be devised using a cancer protein promoter operably linked to a reporter gene such as luciferase, green fluorescent protein, CAT, or P-gal. The reporter construct is typically transfected into a cell. After treatment with a potential modulator, the amount of reporter gene transcription, translation, or activity is measured according to standard techniques known to those of skill in the art (Davis G F, *supra*; Gonzalez, J. & Negulescu, P. *Curr. Opin. Biotechnol.* 1998: 9:624).

As outlined above, in vitro screens are done on individual genes and gene products. That is, having identified a particular differentially expressed gene as important in a particular state, screening of modulators of the expression of the gene or the gene product itself is performed.

In one embodiment, screening for modulators of expression of specific gene(s) is performed. Typically, the expression of only one or a few genes is evaluated. In another embodiment, screens are designed to first find compounds that bind to differentially expressed proteins. These compounds are then evaluated for the ability to modulate differentially expressed activity. Moreover, once initial candidate compounds are identified, variants can be further screened to better evaluate structure activity relationships.

Binding Assays to Identify and Characterize Modulators

In binding assays in accordance with the invention, a purified or isolated gene product of the invention is generally used. For example, antibodies are generated to a protein of the invention, and immunoassays are run to determine the amount and/or location of protein. Alternatively, cells comprising the cancer proteins are used in the assays.

Thus, the methods comprise combining a cancer protein of the invention and a candidate compound such as a ligand, and determining the binding of the compound to the cancer protein of the invention. Preferred embodiments utilize the human cancer protein; animal models of human disease of can also be developed and used. Also, other analogous mammalian proteins also can be used as appreciated by those of skill in the art. Moreover, in some embodiments variant or derivative cancer proteins are used.

Generally, the cancer protein of the invention, or the ligand, is non-diffusibly bound to an insoluble support. The support can, e.g., be one having isolated sample receiving areas (a microtiter plate, an array, etc.). The insoluble supports can be made of any composition to which the compositions can be bound, is readily separated from soluble material, and is otherwise compatible with the overall method of screening. The surface of such supports can be solid or porous and of any convenient shape.

Examples of suitable insoluble supports include microtiter plates, arrays, membranes and beads. These are typically made of glass, plastic (e.g., polystyrene), polysaccharide, nylon, nitrocellulose, or Teflon™, etc. Microtiter plates and arrays are especially convenient because a large number of assays can be carried out simultaneously, using small amounts of reagents and samples. The particular manner of binding of the composition to the support is not crucial so long as it is compatible with the reagents and overall methods of the invention, maintains the activity of the composition and is nondiffusible. Preferred methods of binding include the

use of antibodies which do not sterically block either the ligand binding site or activation sequence when attaching the protein to the support, direct binding to "sticky" or ionic supports, chemical crosslinking, the synthesis of the protein or agent on the surface, etc. Following binding of the protein or ligand/binding agent to the support, excess unbound material is removed by washing. The sample receiving areas may then be blocked through incubation with bovine serum albumin (BSA), casein or other innocuous protein or other moiety.

Once a cancer protein of the invention is bound to the support, and a test compound is added to the assay. Alternatively, the candidate binding agent is bound to the support and the cancer protein of the invention is then added. Binding agents include specific antibodies, non-natural binding agents identified in screens of chemical libraries, peptide analogs, etc.

Of particular interest are assays to identify agents that have a low toxicity for human cells. A wide variety of assays can be used for this purpose, including proliferation assays, cAMP assays, labeled in vitro protein-protein binding assays, electrophoretic mobility shift assays, immunoassays for protein binding, functional assays (phosphorylation assays, etc.) and the like.

A determination of binding of the test compound (ligand, binding agent, modulator, etc.) to a cancer protein of the invention can be done in a number of ways. The test compound can be labeled, and binding determined directly, e.g., by attaching all or a portion of the cancer protein of the invention to a solid support, adding a labeled candidate compound (e.g., a fluorescent label), washing off excess reagent, and determining whether the label is present on the solid support. Various blocking and washing steps can be utilized as appropriate.

In certain embodiments, only one of the components is labeled, e.g., a protein of the invention or ligands labeled. Alternatively, more than one component is labeled with different labels, e.g., I^{125} , for the proteins and a fluorophor for the compound. Proximity reagents, e.g., quenching or energy transfer reagents are also useful.

Competitive Binding to Identify and Characterize Modulators

In one embodiment, the binding of the "test compound" is determined by competitive binding assay with a "competitor." The competitor is a binding moiety that binds to the target molecule (e.g., a cancer protein of the invention). Competitors include compounds such as antibodies, peptides, binding partners, ligands, etc. Under certain circumstances, the competitive binding between the test compound and the competitor displaces the test compound. In one embodiment, the test compound is labeled. Either the test compound, the competitor, or both, is added to the protein for a time sufficient to allow binding. Incubations are performed at a temperature that facilitates optimal activity, typically between four and 40° C. Incubation periods are typically optimized, e.g., to facilitate rapid high throughput screening; typically between zero and one hour will be sufficient. Excess reagent is generally removed or washed away. The second component is then added, and the presence or absence of the labeled component is followed, to indicate binding.

In one embodiment, the competitor is added first, followed by the test compound. Displacement of the competitor is an indication that the test compound is binding to the cancer protein and thus is capable of binding to, and potentially modulating, the activity of the cancer protein. In this embodiment, either component can be labeled. Thus, e.g., if the competitor is labeled, the presence of label in the post-test compound wash solution indicates displacement by the test

compound. Alternatively, if the test compound is labeled, the presence of the label on the support indicates displacement.

In an alternative embodiment, the test compound is added first, with incubation and washing, followed by the competitor. The absence of binding by the competitor indicates that the test compound binds to the cancer protein with higher affinity than the competitor. Thus, if the test compound is labeled, the presence of the label on the support, coupled with a lack of competitor binding, indicates that the test compound binds to and thus potentially modulates the cancer protein of the invention.

Accordingly, the competitive binding methods comprise differential screening to identify agents that are capable of modulating the activity of the cancer proteins of the invention. In this embodiment, the methods comprise combining a cancer protein and a competitor in a first sample. A second sample comprises a test compound, the cancer protein, and a competitor. The binding of the competitor is determined for both samples, and a change, or difference in binding between the two samples indicates the presence of an agent capable of binding to the cancer protein and potentially modulating its activity. That is, if the binding of the competitor is different in the second sample relative to the first sample, the agent is capable of binding to the cancer protein.

Alternatively, differential screening is used to identify drug candidates that bind to the native cancer protein, but cannot bind to modified cancer proteins. For example the structure of the cancer protein is modeled and used in rational drug design to synthesize agents that interact with that site, agents which generally do not bind to site-modified proteins. Moreover, such drug candidates that affect the activity of a native cancer protein are also identified by screening drugs for the ability to either enhance or reduce the activity of such proteins.

Positive controls and negative controls can be used in the assays. Preferably control and test samples are performed in at least triplicate to obtain statistically significant results. Incubation of all samples occurs for a time sufficient to allow for the binding of the agent to the protein. Following incubation, samples are washed free of non-specifically bound material and the amount of bound, generally labeled agent determined. For example, where a radiolabel is employed, the samples can be counted in a scintillation counter to determine the amount of bound compound.

A variety of other reagents can be included in the screening assays. These include reagents like salts, neutral proteins, e.g. albumin, detergents, etc. which are used to facilitate optimal protein-protein binding and/or reduce non-specific or background interactions. Also reagents that otherwise improve the efficiency of the assay, such as protease inhibitors, nuclease inhibitors, anti-microbial agents, etc., can be used. The mixture of components is added in an order that provides for the requisite binding.

Use of Polynucleotides to Down-regulate or Inhibit a Protein of the Invention.

Polynucleotide modulators of cancer can be introduced into a cell containing the target nucleotide sequence by formation of a conjugate with a ligand-binding molecule, as described in WO 91/04753. Suitable ligand-binding molecules include, but are not limited to, cell surface receptors, growth factors, other cytokines, or other ligands that bind to cell surface receptors. Preferably, conjugation of the ligand binding molecule does not substantially interfere with the ability of the ligand binding molecule to bind to its corresponding molecule or receptor, or block entry of the sense or antisense oligonucleotide or its conjugated version into the cell. Alternatively, a polynucleotide modulator of cancer can be introduced into a cell containing the target nucleic acid

sequence, e.g., by formation of a polynucleotide-lipid complex, as described in WO 90/10448. It is understood that the use of antisense molecules or knock out and knock in models may also be used in screening assays as discussed above, in addition to methods of treatment.

Inhibitory and Antisense Nucleotides

In certain embodiments, the activity of a cancer-associated protein is down-regulated, or entirely inhibited, by the use of antisense polynucleotide or inhibitory small nuclear RNA (snRNA), i.e., a nucleic acid complementary to, and which can preferably hybridize specifically to, a coding mRNA nucleic acid sequence, e.g., a cancer protein of the invention, mRNA, or a subsequence thereof. Binding of the antisense polynucleotide to the mRNA reduces the translation and/or stability of the mRNA.

In the context of this invention, antisense polynucleotides can comprise naturally occurring nucleotides, or synthetic species formed from naturally occurring subunits or their close homologs. Antisense polynucleotides may also have altered sugar moieties or inter-sugar linkages. Exemplary among these are the phosphorothioate and other sulfur containing species which are known for use in the art. Analogs are comprised by this invention so long as they function effectively to hybridize with nucleotides of the invention. See, e.g., Isis Pharmaceuticals, Carlsbad, Calif.; Sequitor, Inc., Natick, Mass.

Such antisense polynucleotides can readily be synthesized using recombinant means, or can be synthesized in vitro. Equipment for such synthesis is sold by several vendors, including Applied Biosystems. The preparation of other oligonucleotides such as phosphorothioates and alkylated derivatives is also well known to those of skill in the art.

Antisense molecules as used herein include antisense or sense oligonucleotides. Sense oligonucleotides can, e.g., be employed to block transcription by binding to the anti-sense strand. The antisense and sense oligonucleotide comprise a single stranded nucleic acid sequence (either RNA or DNA) capable of binding to target mRNA (sense) or DNA (antisense) sequences for cancer molecules. Antisense or sense oligonucleotides, according to the present invention, comprise a fragment generally at least about 12 nucleotides, preferably from about 12 to 30 nucleotides. The ability to derive an antisense or a sense oligonucleotide, based upon a cDNA sequence encoding a given protein is described in, e.g., Stein & Cohen (Cancer Res. 48:2659 (1988 and van der Krol et al. (BioTechniques 6:958 (1988)).

Ribozymes

In addition to antisense polynucleotides, ribozymes can be used to target and inhibit transcription of cancer-associated nucleotide sequences. A ribozyme is an RNA molecule that catalytically cleaves other RNA molecules. Different kinds of ribozymes have been described, including group I ribozymes, hammerhead ribozymes, hairpin ribozymes, RNase P, and axhead ribozymes (see, e.g., Castanotto et al., Adv. in Pharmacology 25: 289-317 (1994) for a general review of the properties of different ribozymes).

The general features of hairpin ribozymes are described, e.g., in Hampel et al., Nucl. Acids Res. 18:299-304 (1990); European Patent Publication No. 0360257; U.S. Pat. No. 5,254,678. Methods of preparing are well known to those of skill in the art (see, e.g., WO 94/26877; Ojwang et al., Proc. Natl. Acad. Sci. USA 90:6340-6344 (1993); Yamada et al., Human Gene Therapy 1:39-45 (1994); Leavitt et al., Proc. Natl. Acad. Sci. USA 92:699-703 (1995); Leavitt et al., Human Gene Therapy 5: 1151-120 (1994); and Yamada et al., Virology 205: 121-126 (1994)).

Use of Modulators in Phenotypic Screening

In one embodiment, a test compound is administered to a population of cancer cells, which have an associated cancer expression profile. By "administration" or "contacting" herein is meant that the modulator is added to the cells in such a manner as to allow the modulator to act upon the cell, whether by uptake and intracellular action, or by action at the cell surface. In some embodiments, a nucleic acid encoding a proteinaceous agent (i.e., a peptide) is put into a viral construct such as an adenoviral or retroviral construct, and added to the cell, such that expression of the peptide agent is accomplished, e.g., PCT US97/01019. Regulatable gene therapy systems can also be used. Once the modulator has been administered to the cells, the cells are washed if desired and are allowed to incubate under preferably physiological conditions for some period. The cells are then harvested and a new gene expression profile is generated. Thus, e.g., cancer tissue is screened for agents that modulate, e.g., induce or suppress, the cancer phenotype. A change in at least one gene, preferably many, of the expression profile indicates that the agent has an effect on cancer activity. Similarly, altering a biological function or a signaling pathway is indicative of modulator activity. By defining such a signature for the cancer phenotype, screens for new drugs that alter the phenotype are devised. With this approach, the drug target need not be known and need not be represented in the original gene/protein expression screening platform, nor does the level of transcript for the target protein need to change. The modulator inhibiting function will serve as a surrogate marker.

As outlined above, screens are done to assess genes or gene products. That is, having identified a particular differentially expressed gene as important in a particular state, screening of modulators of either the expression of the gene or the gene product itself is performed.

Use of Modulators to Affect Peptides of the Invention

Measurements of cancer polypeptide activity, or of the cancer phenotype are performed using a variety of assays. For example, the effects of modulators upon the function of a cancer polypeptide(s) are measured by examining parameters described above. A physiological change that affects activity is used to assess the influence of a test compound on the polypeptides of this invention. When the functional outcomes are determined using intact cells or animals, a variety of effects can be assessed such as, in the case of a cancer associated with solid tumors, tumor growth, tumor metastasis, neovascularization, hormone release, transcriptional changes to both known and uncharacterized genetic markers (e.g., by Northern blots), changes in cell metabolism such as cell growth or pH changes, and changes in intracellular second messengers such as cGNIP.

Methods of Identifying Characterizing Cancer-Associated Sequences

Expression of various gene sequences is correlated with cancer. Accordingly, disorders based on mutant or variant cancer genes are determined. In one embodiment, the invention provides methods for identifying cells containing variant cancer genes, e.g., determining the presence of, all or part, the sequence of at least one endogenous cancer gene in a cell. This is accomplished using any number of sequencing techniques. The invention comprises methods of identifying the cancer genotype of an individual, e.g., determining all or part of the sequence of at least one gene of the invention in the individual. This is generally done in at least one tissue of the individual, e.g., a tissue set forth in Table I, and may include the evaluation of a number of tissues or different samples of the same tissue. The method may include comparing the sequence of the sequenced gene to a known cancer gene, i.e.,

a wild-type gene to determine the presence of family members, homologies, mutations or variants. The sequence of all or part of the gene can then be compared to the sequence of a known cancer gene to determine if any differences exist. This is done using any number of known homology programs, such as BLAST, Bestfit, etc. The presence of a difference in the sequence between the cancer gene of the patient and the known cancer gene correlates with a disease state or a propensity for a disease state, as outlined herein.

In a preferred embodiment, the cancer genes are used as probes to determine the number of copies of the cancer gene in the genome. The cancer genes are used as probes to determine the chromosomal localization of the cancer genes. Information such as chromosomal localization finds use in providing a diagnosis or prognosis in particular when chromosomal abnormalities such as translocations, and the like are identified in the cancer gene locus.

XIV.) KITS/ARTICLES OF MANUFACTURE

For use in the diagnostic and therapeutic applications described herein, kits are also within the scope of the invention. Such kits can comprise a carrier, package or container that is compartmentalized to receive one or more containers such as vials, tubes, and the like, each of the container(s) comprising one of the separate elements to be used in the method. For example, the container(s) can comprise a probe that is or can be detectably labeled. Such probe can be an antibody or polynucleotide specific for a FIG. 2-related protein or a FIG. 2 gene or message, respectively. Where the method utilizes nucleic acid hybridization to detect the target nucleic acid, the kit can also have containers containing nucleotide(s) for amplification of the target nucleic acid sequence and/or a container comprising a reporter-means, such as a biotin-binding protein, such as avidin or streptavidin, bound to a reporter molecule, such as an enzymatic, fluorescent, or radioisotope label. The kit can include all or part of the amino acid sequences in FIG. 2 or FIG. 3 or analogs thereof, or a nucleic acid molecules that encodes such amino acid sequences.

The kit of the invention will typically comprise the container described above and one or more other containers comprising materials desirable from a commercial and user standpoint, including buffers, diluents, filters, needles, syringes; carrier, package, container, vial and/or tube labels listing contents and/or instructions for use, and package inserts with instructions for use.

A label can be present on the container to indicate that the composition is used for a specific therapy or non-therapeutic application, such as a diagnostic or laboratory application, and can also indicate directions for either in vivo or in vitro use, such as those described herein. Directions and or other information can also be included on an insert(s) or label(s) which is included with or on the kit.

The terms "kit" and "article of manufacture" can be used as synonyms.

In another embodiment of the invention, an article(s) of manufacture containing compositions, such as amino acid sequence(s), small molecule(s), nucleic acid sequence(s), and/or antibody(s), e.g., materials useful for the diagnosis, prognosis, prophylaxis and/or treatment of neoplasias of tissues such as those set forth in Table I is provided. The article of manufacture typically comprises at least one container and at least one label. Suitable containers include, for example, bottles, vials, syringes, and test tubes. The containers can be formed from a variety of materials such as glass or plastic. The container can hold amino acid sequence(s), small mol-

ecule(s), nucleic acid sequence(s), and/or antibody(s), in one embodiment the container holds a polynucleotide for use in examining the mRNA expression profile of a cell, together with reagents used for this purpose.

The container can alternatively hold a composition which is effective for treating, diagnosis, prognosing or prophylaxing a condition and can have a sterile access port (for example the container can be an intravenous solution bag or a vial having a stopper pierceable by a hypodermic injection needle). The active agents in the composition can be an antibody capable of specifically binding 24P4C12 and modulating the function of 24P4C12.

The label can be on or associated with the container. A label can be on a container when letters, numbers or other characters forming the label are molded or etched into the container itself; a label can be associated with a container when it is present within a receptacle or carrier that also holds the container, e.g., as a package insert. The label can indicate that the composition is used for diagnosing, treating, prophylaxing or prognosing a condition, such as a neoplasia of a tissue set forth in Table I. The article of manufacture can further comprise a second container comprising a pharmaceutically-acceptable buffer, such as phosphate-buffered saline, Ringer's solution and/or dextrose solution. It can further include other materials desirable from a commercial and user standpoint, including other buffers, diluents, filters, stirrers, needles, syringes, and/or package inserts with indications and/or instructions for use.

EXAMPLES

Various aspects of the invention are further described and illustrated by way of the several examples that follow, none of which are intended to limit the scope of the invention.

Example 1

SSH-Generated Isolation of cDNA Fragment of the 24P4C12 Gene

Suppression Subtractive Hybridization (SSH) was used to identify cDNAs corresponding to genes that may be differentially expressed in prostate cancer. The SSH reaction utilized cDNA from the LAPC-9 AD prostate cancer xenograft. The gene 24P4C12 was derived from an LAPC-9 AD minus benign prostatic hyperplasia experiment.

The 24P4C12 SSH cDNA of 160 bp is listed in FIG. 1. The full length 24P4C12 cDNAs and ORFs are described in FIG. 2 with the protein sequences listed in FIG. 3.

Materials and Methods

Human Tissues:

The patient cancer and normal tissues were purchased from different sources such as the NDRI (Philadelphia, Pa.). mRNA for some normal tissues were purchased from Clontech, Palo Alto, Calif.

RNA Isolation:

Tissues were homogenized in Trizol reagent (Life Technologies, Gibco BRL) using 10 ml/g tissue isolate total RNA. Poly A RNA was purified from total RNA using Qiagen's Oligotex mRNA Mini and Midi kits. Total and mRNA were quantified by spectrophotometric analysis (O.D. 260/280 nm) and analyzed by gel electrophoresis.

Oligonucleotides:

The following HPLC purified oligonucleotides were used.

DPNCDN (cDNA synthesis primer):	(SEQ ID NO: 33)
5' TTTTGATCAAGCTT ₃₀ 3'	
Adaptor 1:	(SEQ ID NO: 34)
5' CTAATACGACTCACTATAGGGCTCGAGCGGCCGCCGGCAG3'	(SEQ ID NO: 35)
3' GGCCCGTCCTAG5'	
Adaptor 2:	(SEQ ID NO: 36)
5' GTAATACGACTCACTATAGGGCAGCGTGGTCGCGGCCGAG3'	(SEQ ID NO: 37)
3' CGGCTCCTAG5'	
PCR primer 1:	(SEQ ID NO: 38)
5' CTAATACGACTCACTATAGGGC3'	
Nested primer (NP)1:	(SEQ ID NO: 39)
5' TCGAGCGGCCGCCCGGGCAGGA3'	
Nested primer (NP)2:	(SEQ ID NO: 40)
5' AGCGTGGTCGCGGCCGAGGA3'	

Suppression Subtractive Hybridization:

Suppression Subtractive Hybridization (SSH) was used to identify cDNAs corresponding to genes that may be differentially expressed in prostate cancer. The SSH reaction utilized cDNA from prostate cancer and normal tissues.

The gene 24P4C12 sequence was derived from LAPC-4AD prostate cancer xenograft minus benign prostatic hyperplasia cDNA subtraction. The SSH DNA sequence (FIG. 1) was identified.

The cDNA derived from a pool of normal tissues and benign prostatic hyperplasia was used as the source of the "driver" cDNA, while the cDNA from LAPC-4AD xenograft was used as the source of the "tester" cDNA. Double stranded cDNAs corresponding to tester and driver cDNAs were synthesized from 2 µg of poly(A)⁺ RNA isolated from the relevant xenograft tissue, as described above, using CLONTECH's PCR-Select cDNA Subtraction Kit and 1 ng of oligonucleotide DPNCDN as primer. First- and second-strand synthesis were carried out as described in the Kit's user manual protocol (CLONTECH Protocol No. PT1117-1, Catalog No. K1804-1). The resulting cDNA was digested with Dpn II for 3 hrs at 37° C. Digested cDNA was extracted with phenol/chloroform (1:1) and ethanol precipitated.

Driver cDNA was generated by combining in a 1:1 ratio Dpn II digested cDNA from the relevant tissue source (see above) with a mix of digested cDNAs derived from the nine normal tissues: stomach, skeletal muscle, lung, brain, liver, kidney, pancreas, small intestine, and heart.

Tester cDNA was generated by diluting 1 µl of Dpn II digested cDNA from the relevant tissue source (see above) (400 ng) in 5 µl of water. The diluted cDNA (2 µl, 160 ng) was then ligated to 2 µl of Adaptor 1 and Adaptor 2 (10 µM), in separate ligation reactions, in a total volume of 10 µl at 16° C. overnight, using 400 u of T4 DNA ligase (CLONTECH). Ligation was terminated with 1 µl of 0.2 M EDTA and heating at 72° C. for 5 min.

The first hybridization was performed by adding 1.5 µl (600 ng) of driver cDNA to each of two tubes containing 1.5 µl (20 ng) Adaptor 1- and Adaptor 2-ligated tester cDNA. In a final volume of 4 µl, the samples were overlaid with mineral oil, denatured in an MJ Research thermal cycler at 98° C. for 1.5 minutes, and then were allowed to hybridize for 8 hrs at 68° C. The two hybridizations were then mixed together with

an additional 1 µl of fresh denatured driver cDNA and were allowed to hybridize overnight at 68° C. The second hybridization was then diluted in 200 µl of 20 mM Hepes, pH 8.3, 50 mM NaCl, 0.2 mM EDTA, heated at 70° C. for 7 min. and stored at -20° C.

PCR Amplification, Cloning and Sequencing of Gene Fragments Generated from SSH:

To amplify gene fragments resulting from SSH reactions, two PCR amplifications were performed. In the primary PCR reaction 1 µl of the diluted final hybridization mix was added to 1 µl of PCR primer 1 (10 µM), 0.5 µl dNTP mix (10 µM), 2.5 µl 10× reaction buffer (CLONTECH) and 0.5 µl 50× Advantage cDNA polymerase Mix (CLONTECH) in a final volume of 25 µl. PCR 1 was conducted using the following conditions: 75° C. for 5 min., 94° C. for 25 sec., then 27 cycles of 94° C. for 10 sec, 66° C. for 30 sec, 72° C. for 1.5 min. Five separate primary PCR reactions were performed for each experiment. The products were pooled and diluted 1:10 with water. For the secondary PCR reaction, 1 µl from the pooled and diluted primary PCR reaction was added to the same reaction mix as used for PCR 1, except that primers NP1 and NP2 (10 µM) were used instead of PCR primer 1. PCR 2 was performed using 10-12 cycles of 94° C. for 10 sec, 68° C. for 30 sec, and 72° C. for 1.5 minutes. The PCR products were analyzed using 2% agarose gel electrophoresis.

The PCR products were inserted into pCR2.1 using the T/A vector cloning kit (Invitrogen). Transformed *E. coli* were subjected to blue/white and ampicillin selection. White colonies were picked and arrayed into 96 well plates and were grown in liquid culture overnight. To identify inserts, PCR amplification was performed on 1 µl of bacterial culture using the conditions of PCR1 and NP1 and NP2 as primers. PCR products were analyzed using 2% agarose gel electrophoresis.

Bacterial clones were stored in 20% glycerol in a 96 well format. Plasmid DNA was prepared, sequenced, and subjected to nucleic acid homology searches of the GenBank, dBest, and NCI-CGAP databases.

RT-PCR Expression Analysis:

First strand cDNAs can be generated from 1 µg of mRNA with oligo (dT) 12-18 priming using the Gibco-BRL Superscript Preamplification system. The manufacturer's protocol was used which included an incubation for 50 min at 42° C. with reverse transcriptase followed by RNase H treatment at 37° C. for 20 min. After completing the reaction, the volume can be increased to 200 µl with water prior to normalization. First strand cDNAs from 16 different normal human tissues can be obtained from Clontech.

Normalization of the first strand cDNAs from multiple tissues was performed by using the primers 5'atcgcgcgcgtcgtcgcaaa3' (SEQ ID NO: 41) and 5'agccacacgcagctcattgtagaagg 3' (SEQ ID NO: 42) to amplify β-actin. First strand cDNA (5 µl) were amplified in a total volume of 50 µl containing 0.4 µM primers, 0.2 µM each dNTPs, 1×PCR buffer (Clontech, 10 mM Tris-HCL, 1.5 mM MgCl₂, 50 mM KCl, pH8.3) and 1× KlenTaq DNA polymerase (Clontech). Five µl of the PCR reaction can be removed at 18, 20, and 22 cycles and used for agarose gel electrophoresis. PCR was performed using an MJ Research thermal cycler under the following conditions: Initial denaturation can be at 94° C. for 15 sec, followed by a 18, 20, and 22 cycles of 94° C. for 15, 65° C. for 2 min, 72° C. for 5 sec. A final extension at 72° C. was carried out for 2 min. After agarose gel electrophoresis, the band intensities of the 283 b.p. β-actin bands from multiple tissues were compared by visual inspection. Dilution factors for the first strand cDNAs were calculated to result in equal β-actin band intensities in all tissues after 22 cycles of

PCR. Three rounds of normalization can be required to achieve equal band intensities in all tissues after 22 cycles of PCR.

To determine expression levels of the 24P4C12 gene, 5 µl of normalized first strand cDNA were analyzed by PCR using 26, and 30 cycles of amplification. Semi-quantitative expression analysis can be achieved by comparing the PCR products at cycle numbers that give light band intensities. The primers used for RT-PCR were designed using the 24P4C12 SSH sequence and are listed below:

24P4C12.1
5' -AGATGAGGAGGAGGACAAAGGTG-3' (SEQ ID NO: 43)

24P4C12.2
5' -ACTGCTGGGAGGAGTACCGAGTG-3' (SEQ ID NO: 44)

Example 2

Isolation of Full Length 24P4C12 Encoding cDNA

The 24P4C12 SSH cDNA sequence was derived from a subtraction consisting of LAPC-4AD xenograft minus benign prostatic hyperplasia. The SSH cDNA sequence (FIG. 1) was designated 24P4C12.

The isolated gene fragment of 160 bp encodes a putative open reading frame (ORF) of 53 amino acids and exhibits significant homology to an EST derived from a colon tumor library. Two larger cDNA clones were obtained by gene trap experiments, GTE9 and GTF8. The ORF revealed a significant homology to the mouse gene NG22 and the *C. elegans* gene CEESB82F. NG22 was recently identified as one of many ORFs within a genomic BAC clone that encompasses the MHC class III in the mouse genome. Both NG22 and CEESB82F appear to be genes that contain 12 transmembrane domains. This suggests that the gene encoding 24P4C12 contains 12 transmembrane domains and is the human homologue of mouse NG22 and *C. elegans* CEESB82F. Functional studies in *C. elegans* may reveal the biological role of these homologs. If 24P4C12 is a cell surface marker, then it may have an application as a potential imaging reagent and/or therapeutic target in prostate cancer.

The 24P4C12 v.1 of 2587 bp codes for a protein of 710 amino acids (FIG. 2 and FIG. 3). Other variants of 24P4C12 were also identified and these are listed in FIGS. 2 and 3. 24P4C12 v.1, v.3, v.5 and v.6 proteins are 710 amino acids in length and differ from each other by one amino acid as shown in FIG. 11. 24P4C12 v.2 and v.4 code for the same protein as 24P4C12 v.1. 24P4C12 v.7, v.8 and v.9 are alternative splice variants and code for proteins of 598, 722 and 712 amino acids in length, respectively.

Example 3

Chromosomal Mapping of 24P4C12

Chromosomal localization can implicate genes in disease pathogenesis. Several chromosome mapping approaches are available including fluorescent in situ hybridization (FISH), human/hamster radiation hybrid (RH) panels (Walter et al., 1994; Nature Genetics 7:22; Research Genetics, Huntsville Ala.), human-rodent somatic cell hybrid panels such as is available from the Coriell Institute (Camden, N.J.), and genomic viewers utilizing BLAST homologies to sequenced and mapped genomic clones (NCBI, Bethesda, Md.). 24P4C12 maps to chromosome 6p21.3 using 24P4C12

sequence and the NCBI BLAST tool located on the World Wide Web at (ncbi.nlm.nih.gov/genome/seq/page.cgi?F=HsBlast.html&&ORG=Hs).

Example 4

Expression Analysis of 24P4C12

Expression analysis by RT-PCR demonstrated that 24P4C12 is strongly expressed in prostate and ovary cancer patient specimens (FIG. 14). First strand cDNA was generated from vital pool 1 (kidney, liver and lung), vital pool 2 (colon, pancreas and stomach), a pool of prostate cancer xenografts (LAPC-4AD, LAPC-4AI, LAPC-9AD and LAPC-9AI), prostate cancer pool, bladder cancer pool, kidney cancer pool, colon cancer pool, ovary cancer pool, breast cancer pool, and cancer metastasis pool. Normalization was performed by PCR using primers to actin. Semi-quantitative PCR, using primers to 24P4C12, was performed at 26 and 30 cycles of amplification. Results show strong expression of 24P4C12 in prostate cancer pool and ovary cancer pool. Expression was also detected in prostate cancer xenografts, bladder cancer pool, kidney cancer pool, colon cancer pool, breast cancer pool, cancer metastasis pool, vital pool 1, and vital pool 2.

Extensive northern blot analysis of 24P4C12 in multiple human normal tissues is shown in FIG. 15. Two multiple tissue northern blots (Clontech) both with 2 µg of mRNA/lane were probed with the 24P4C12 SSH sequence. Expression of 24P4C12 was detected in prostate, kidney and colon. Lower expression is detected in pancreas, lung and placenta amongst all 16 normal tissues tested.

Expression of 24P4C12 was tested in prostate cancer xenografts and cell lines. RNA was extracted from a panel of cell lines and prostate cancer xenografts (PrEC, LAPC-4AD, LAPC-4AI, LAPC-9AD, LAPC-9AI, LNCaP, PC-3, DU145, TsuPr, and LAPC-4CL). Northern blot with 10 µg of total RNA/lane was probed with 24P4C12 SSH sequence. Size standards in kilobases (kb) are indicated on the side. The 24P4C12 transcript was detected in LAPC-4AD, LAPC-4AI, LAPC-9AD, LAPC-9AI, LNCaP, and LAPC-4 CL.

Expression of 24P4C12 in patient cancer specimens and human normal tissues is shown in FIG. 16. RNA was extracted from a pool of prostate cancer specimens, bladder cancer specimens, colon cancer specimens, ovary cancer specimens, breast cancer specimens and cancer metastasis specimens, as well as from normal prostate (NP), normal bladder (NB), normal kidney (NK), and normal colon (NC). Northern blot with 10 µg of total RNA/lane was probed with 24P4C12 SSH sequence. Size standards in kilobases (kb) are indicated on the side. Strong expression of 24P4C12 transcript was detected in the patient cancer pool specimens, and in normal prostate but not in the other normal tissues tested.

Expression of 24P4C12 was also detected in individual prostate cancer patient specimens (FIG. 17). RNA was extracted from normal prostate (N), prostate cancer patient tumors (T) and their matched normal adjacent tissues (Nat). Northern blots with 10 µg of total RNA were probed with the 24P4C12 SSH fragment. Size standards in kilobases are on the side. Results show expression of 24P4C12 in normal prostate and all prostate patient tumors tested.

Expression of 24P4C12 in colon cancer patient specimens is shown in FIG. 18. RNA was extracted from colon cancer cell lines (CL: Colo 205, LoVo, and SK—CO—), normal colon (N), colon cancer patient tumors (T) and their matched normal adjacent tissues (Nat). Northern blots with 10 µg of total RNA were probed with the 24P4C12 SSH fragment.

Size standards in kilobases are on the side. Results show expression of 24P4C12 in normal colon and all colon patient tumors tested. Expression was detected in the cell lines Colo 205 and SK—CO—, but not in LoVo.

FIG. 20 displays expression results of 24P4C12 in lung cancer patient specimens. Ma was extracted from lung cancer cell lines (CL: CALU-1, A427, NCI-H82, NCI-H146), normal lung (N), lung cancer patient tumors (T) and their matched normal adjacent tissues (Nat). Northern blots with 10 µg of total RNA were probed with the 24P4C12 SSH fragment. Size standards in kilobases are on the side. Results show expression of 24P4C12 in lung patient tumors tested, but not in normal lung. Expression was also detected in CALU-1, but not in the other cell lines A427, NCI-H82, and NCI-H₁₄₆.

24P4C12 was assayed in a panel of human stomach and breast cancers (T) and their respective matched normal tissues (N) on RNA dot blots. 24P4C12 expression was seen in both stomach and breast cancers. The expression detected in normal adjacent tissues (isolated from diseased tissues) but not in normal tissues (isolated from healthy donors) may indicate that these tissues are not fully normal and that 24P4C12 may be expressed in early stage tumors.

The level of expression of 24P4C12 was analyzed and quantitated in a panel of patient cancer tissues. First strand cDNA was prepared from a panel of ovary patient cancer specimens (A), uterus patient cancer specimens (B), prostate cancer specimens (C), bladder cancer patient specimens (D), lung cancer patient specimens (E), pancreas cancer patient specimens (F), colon cancer specimens (G), and kidney cancer specimens (H). Normalization was performed by PCR using primers to actin. Semi-quantitative PCR, using primers to 24P4C12, was performed at 26 and 30 cycles of amplification. Samples were run on an agarose gel, and PCR products were quantitated using the AlphaImager software. Expression was recorded as absent, low, medium or strong. Results show expression of 24P4C12 in the majority of patient cancer specimens tested, 73.3% of ovary patient cancer specimens, 83.3% of uterus patient cancer specimens, 95.0% of prostate cancer specimens, 61.1% of bladder cancer patient specimens, 80.6% of lung cancer patient specimens, 87.5% of pancreas cancer patient specimens, 87.5% of colon cancer specimens, 68.4% of clear cell renal carcinoma, 100% of papillary renal cell carcinoma. The restricted expression of 24P4C12 in normal tissues and the expression detected in prostate cancer, ovary cancer, bladder cancer, colon cancer, lung cancer pancreas cancer, uterus cancer, kidney cancer, stomach cancer and breast cancer suggest that 24P4C12 is a potential therapeutic target and a diagnostic marker for human cancers.

Example 5

Transcript Variants of 24P4C12

Transcript variants are variants of mature mRNA from the same gene which arise by alternative transcription or alternative splicing. Alternative transcripts are transcripts from the same gene but start transcription at different points. Splice variants are mRNA variants spliced differently from the same transcript. In eukaryotes, when a multi-exon gene is transcribed from genomic DNA, the initial RNA is spliced to produce functional mRNA, which has only exons and is used for translation into an amino acid sequence. Accordingly, a given gene can have zero to many alternative transcripts and each transcript can have zero to many splice variants. Each transcript variant has a unique exon makeup, and can have

Figure 2D-2

200 T I Q Q G I S G L I D S L N A R D I S V
601 CCACCATACAGCAGGGGATCAGCGGTCTTATTGACAGCCTCAATGCCCCGAGACATCAGTG
220 K I F E D F A Q S W Y W I L V A L G V A
661 TTAAGATCTTTGAAGATTTTGCCAGTCCTGGTATTGGATTCTTGTGGCCCTGGGGGTGG
240 L V L S L L F I L L L R L V A G F L V L
721 CTCGGTCTTGAGCCTACTGTTTATCTTGCTTCGCGCCTGGTGGCTGGGCCCCCTGGTGC
260 V L I L G V L G V L A Y G I Y Y C W E E
781 TGGTGTGATCCTGGGAGTGTGGGCGTGTGGCATATGGCATCTACTACTGTGGGAGG
280 Y R V L R D K G A S I S Q L G F T T N L
841 AGTACCGAGTGTGCGGGACAAGGGCGCTCCATCTCCCAGCTGGGTTTCACCACCAACC
300 S A Y Q S V Q E I W L A A L I V L A V L
901 TCAGTGCCTACCAGAGCGTGCAGGAGACCTGGCTGGCCGCCCTGATCGTGTGGCGGTGC
320 E A I L L L M L I F L R Q R I R I A I A
961 TTGAAGCCATCCTGCTGCTGATGCTCATCTTCCTGCGGCAGCGGATTCTGATTGCCATCG
340 L L K E A S K A V G Q M M S T M F Y P L
1021 CCCTCCTGAAGGAGGCCAGCAAGGCTGTGGGACAGATGATGTCTACCATGTTCTACCCAC
360 V T F V L L L I C I A Y W A M T A L Y L
1081 TGGTCACCTTTGTCTCCTCCTCATCTGCATTGCCTACTGGGCCATGACTGCTCTGTACC
380 A T S G Q P Q Y V L W A S N I S S P G C
1141 TGGCTACATCGGGGCAACCCAGTATGTGCTCTGGGCATCCAACATCAGCTCCCCGGCT
400 E K V P I N T S C N P T A H L V N S S C
1201 GTGAGAAAGTGCCAATAAATACATCATGCAACCCACGGCCACCTTGTGAACCTCCTCGT
420 F G L M C V F Q G Y S S K G L I Q R S V
1261 GCCCCAGGGCTGATGTGCGTCTTCCAGGGCTACTCATCCAAAGGCCTAATCCAACGTTCTG
440 F N L Q I Y G V L G L F W T L N W V L A
1321 TCTTCAATCTGCAAATCTATGGGGTCTGGGGCTCTTCTGGACCCCTTAAGTGGGTACTGG
460 L G Q C V L A G A F A S F Y W A F H K P
1381 CCCTGGGCCAATGCGTCCCTCGCTGGAGCCTTTGCCTCCTTCTACTGGGCCTTCCACAAGC
480 Q D I P T F P L I S A F I R T L R Y H T
1441 CCCAGGACATCCCTACCTTCCCCTTAATCTCTGCCTTCATCCGCACACTCCGTTACCACA
500 G S L A F G A L I L T L V Q I A F V I L
1501 CIGGGTCATTGGCATTGGAGCCCTCATCTGACCCTTGTGCAGATAGCCCGGGTCATCT
520 E Y I D H K L R G V Q N P V A R C I M C
1561 TGGAGTATATTGACCACAAGCTCAGAGGAGTGCAGAACCCIGTAGCCCGCTGCATCATGT
540 C F K C C L W C L E K F I K F L N R N A
1621 GCTGTTTCAAGTGTGCTGCCCTCGGTGTCTGGAAAAATTTATCAAGTTCCTAAACCGCAATG
560 Y I M I A I Y G K N F C V S A K N A F M
1681 CATACATCATGATCGCCAICTACGGGAAGAATTTCTGTGTCTCAGCCAAAAATGCGTTCA
580 L L M R N I V R V V V L D K V T D L L L
1741 TGCTACTCATGCGAAACATTGTGAGGGTGGTGTCTCTGGACAAAGTCACAGACCTGTGTC

different coding and/or non-coding (5' or 3' end) portions, from the original transcript. Transcript variants can code for similar or different proteins with the same or a similar function or can encode proteins with different functions, and can be expressed in the same tissue at the same time, or in different tissues at the same time, or in the same tissue at different times, or in different tissues at different times. Proteins encoded by transcript variants can have similar or different cellular or extracellular localizations, e.g., secreted versus intracellular.

Transcript variants are identified by a variety of art-accepted methods. For example, alternative transcripts and splice variants are identified by full-length cloning experiment, or by use of full-length transcript and EST sequences. First, all human ESTs were grouped into clusters which show direct or indirect identity with each other. Second, ESTs in the same cluster were further grouped into sub-clusters and assembled into a consensus sequence. The original gene sequence is compared to the consensus sequence(s) or other full-length sequences. Each consensus sequence is a potential splice variant for that gene. Even when a variant is identified that is not a full-length clone, that portion of the variant is very useful for antigen generation and for further cloning of the full-length splice variant, using techniques known in the art.

Moreover, computer programs are available in the art that identify transcript variants based on genomic sequences. Genomic-based transcript variant identification programs include FgenesH (A. Salamov and V. Solovyev, "Ab initio gene finding in *Drosophila* genomic DNA," Genome Research. 2000 April; 10(4):516-22); Grail (URL at comp-bio.ornl.gov/Grail-bin/EmptyGrailForm) and GenScan (URL at genes.mit.edu/GENSCAN.html). For a general discussion of splice variant identification protocols see, e.g., Southan, C., A genomic perspective on human proteases, FEBS Lett. 2001 Jun. 8; 498(2-3):214-8; de Souza, S. J., et al., Identification of human chromosome 22 transcribed sequences with ORF expressed sequence tags, Proc. Natl Acad Sci USA. 2000 Nov. 7; 97(23):12690-3.

To further confirm the parameters of a transcript variant, a variety of techniques are available in the art, such as full-length cloning, proteomic validation, PCR-based validation, and 5' RACE validation, etc. (see e.g., Proteomic Validation: Brennan, S. O., et al., Albumin banks peninsula: a new termination variant characterized by electrospray mass spectrometry, Biochem Biophys Acta. 1999 Aug. 17; 1433(1-2): 321-6; Ferranti P, et al., Differential splicing of pre-messenger RNA produces multiple forms of mature caprine alpha(s1)-casein, Eur J Biochem. 1997 Oct. 1; 249(1):1-7. For PCR-based Validation: Wellmann S, et al., Specific reverse transcription-PCR quantification of vascular endothelial growth factor (VEGF) splice variants by LightCycler technology, Clin Chem. 2001 April; 47(4):654-60; Jia, H. P., et al., Discovery of new human beta-defensins using a genomics-based approach, Gene. 2001 Jan. 24; 263(1-2): 211-8. For PCR-based and 5' RACE Validation: Brigle, K. E., et al., Organization of the murine reduced folate carrier gene and identification of variant splice forms, Biochem Biophys Acta. 1997 Aug. 7; 1353(2): 191-8).

It is known in the art that genomic regions are modulated in cancers. When the genomic region to which a gene maps is modulated in a particular cancer, the alternative transcripts or splice variants of the gene are modulated as well. Disclosed herein is that 24P4C12 has a particular expression profile related to cancer. Alternative transcripts and splice variants of 24P4C12 may also be involved in cancers in the same or different tissues, thus serving as tumor-associated markers/antigens.

The exon composition of the original transcript, designated as 24P4C12 v.1, is shown in Table LI. Using the full-length gene and EST sequences, three transcript variants were identified, designated as 24P4C12 v.7, v.8 and v.9. Compared with 24P4C12 v.1, transcript variant 24P4C12 v.7 has spliced out exons 10 and 11 from variant 24P4C12 v.1, as shown in FIG. 12. Variant 24P4C12 v.8 inserted 36 bp in between 1931 and 1932 of variant 24P4C12 v.1 and variant 24P4C12 v.9 replaced with 36 bp the segment 1136-1163 of variant 24P4C12 v.1. Theoretically, each different combination of exons in spatial order, e.g. exons 2 and 3, is a potential splice variant. FIG. 12 shows the schematic alignment of exons of the four transcript variants.

Tables LII through LXIII are set forth on a variant by variant basis. Tables LII, LVI, and LX show nucleotide sequences of the transcript variant. Tables LIII, LVII, and LXI show the alignment of the transcript variant with the nucleic acid sequence of 24P4C12 v.1. Tables LIV, LVIII, and LXII lay out the amino acid translation of the transcript variant for the identified reading frame orientation. Tables LV, LIX, and LXIII display alignments of the amino acid sequence encoded by the splice variant with that of 24P4C12 v.1.

Example 6

Single Nucleotide Polymorphisms of 24P4C12

A Single Nucleotide Polymorphism (SNP) is a single base pair variation in a nucleotide sequence at a specific location. At any given point of the genome, there are four possible nucleotide base pairs: A/T, C/G, G/C and T/A. Genotype refers to the specific base pair sequence of one or more locations in the genome of an individual. Haplotype refers to the base pair sequence of more than one location on the same DNA molecule (or the same chromosome in higher organisms), often in the context of one gene or in the context of several tightly linked genes. SNPs that occur on a cDNA are called cSNPs. These cSNPs may change amino acids of the protein encoded by the gene and thus change the functions of the protein. Some SNPs cause inherited diseases; others contribute to quantitative variations in phenotype and reactions to environmental factors including diet and drugs among individuals. Therefore, SNPs and/or combinations of alleles (called haplotypes) have many applications, including diagnosis of inherited diseases, determination of drug reactions and dosage, identification of genes responsible for diseases, and analysis of the genetic relationship between individuals (P. Nowotny, J. M. Kwon and A. M. Goate, "SNP analysis to dissect human traits," Curr. Opin. Neurobiol. 2001 October; 11(5):637-641; M. Pirmohamed and B. K. Park, "Genetic susceptibility to adverse drug reactions," Trends Pharmacol. Sci. 2001 June; 22(6):298-305; J. H. Riley, C. J. Allan, E. Lai and A. Roses, "The use of single nucleotide polymorphisms in the isolation of common disease genes," Pharmacogenomics. 2000 February; 1(1):39-47; R. Judson, J. C. Stephens and A. Windemuth, "The predictive power of haplotypes in clinical response," Pharmacogenomics. 2000 feb; 1(1):15-26).

SNPs are identified by a variety of art-accepted methods (P. Bean, "The promising voyage of SNP target discovery," Am. Clin. Lab. 2001 October-November; 20(9):18-20; K. M. Weiss, "In search of human variation," Genome Res. 1998 July; 8(7):691-697; M. M. She, "Enabling large-scale pharmacogenetic studies by high-throughput mutation detection and genotyping technologies," Clin. Chem. 2001 February; 47(2):164-172). For example, SNPs are identified by sequencing DNA fragments that show polymorphism by gel-based methods such as restriction fragment length polymor-

phism (RFLP) and denaturing gradient gel electrophoresis (DGGE). They can also be discovered by direct sequencing of DNA samples pooled from different individuals or by comparing sequences from different DNA samples. With the rapid accumulation of sequence data in public and private databases, one can discover SNPs by comparing sequences using computer programs (Z. Gu, L. Hillier and P. Y. Kwok, "Single nucleotide polymorphism hunting in cyberspace," *Hum. Mutat.* 1998; 12(4):221-225). SNPs can be verified and genotype or haplotype of an individual can be determined by a variety of methods including direct sequencing and high throughput microarrays (P. Y. Kwok, "Methods for genotyping single nucleotide polymorphisms," *Annu. Rev. Genomics Hum. Genet.* 2001; 2:235-258; M. Kokoris, K. Dix, K. Moynihan, J. Mathis, B. Erwin, P. Grass, B. Hines and A. Dueterhoeft, "High-throughput SNP genotyping with the Masscode system," *Mol. Diagn.* 2000 December; 5(4):329-340).

Using the methods described above, five SNPs were identified in the original transcript, 24P4C12 v.1, at positions 542 (G/A), 564 (G/A), 818 (C/T), 981 (A/G) and 1312 (A/C). The transcripts or proteins with alternative alleles were designated as variants 24P4C12 v.2, v.3, v.4, v.5 and v.6, respectively. FIG. 10 shows the schematic alignment of the SNP variants. FIG. 11 shows the schematic alignment of protein variants, corresponding to nucleotide variants. Nucleotide variants that code for the same amino acid sequence as variant 1 are not shown in FIG. 11. These alleles of the SNPs, though shown separately here, can occur in different combinations (haplotypes) and in any one of the transcript variants (such as 24P4C12 v.7) that contains the sequence context of the SNPs.

Example 7

Production of Recombinant 24P4C12 in Prokaryotic Systems

To express recombinant 24P4C12 and 24P4C12 variants in prokaryotic cells, the full or partial length 24P4C12 and 24P4C12 variant cDNA sequences are cloned into any one of a variety of expression vectors known in the art. The full length cDNA, or any 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30 or more contiguous amino acids from 24P4C12, variants, or analogs thereof are used.

A. In vitro transcription and translation constructs:

PCRII: To generate 24P4C12 sense and anti-sense RNA probes for RNA in situ investigations, pCRII constructs (Invitrogen, Carlsbad Calif.) are generated encoding either all or fragments of the 24P4C12 cDNA. The pCRII vector has Sp6 and T7 promoters flanking the insert to drive the transcription of 24P4C12 RNA for use as probes in RNA in situ hybridization experiments. These probes are used to analyze the cell and tissue expression of 24P4C12 at the RNA level. Transcribed 24P4C12 RNA representing the cDNA amino acid coding region of the 24P4C12 gene is used in in vitro translation systems such as the TnT™ Coupled Reticulolysate System (Promega, Corp., Madison, Wis.) to synthesize 24P4C12 protein.

B. Bacterial Constructs:

pGEX Constructs: To generate recombinant 24P4C12 proteins in bacteria that are fused to the Glutathione S-transferase (GST) protein, all or parts of the 24P4C12 cDNA or variants are cloned into the GST-fusion vector of the pGEX family (Amersham Pharmacia Biotech, Piscataway, N.J.). These constructs allow controlled expression of recombinant 24P4C12 protein sequences with GST fused at the amino-

terminus and a six histidine epitope (6×His) at the carboxyl-terminus. The GST and 6× His tags permit purification of the recombinant fusion protein from induced bacteria with the appropriate affinity matrix and allow recognition of the fusion protein with anti-GST and anti-His antibodies. The 6× His tag is generated by adding 6 histidine codons to the cloning primer at the 3' end, e.g., of the open reading frame (ORF). A proteolytic cleavage site, such as the PreScission™ recognition site in pGEX-6P-1, may be employed such that it permits cleavage of the GST tag from 24P4C12-related protein. The ampicillin resistance gene and pBR322 origin permits selection and maintenance of the pGEX plasmids in *E. coli*.

pMAL Constructs: To generate, in bacteria, recombinant 24P4C12 proteins that are fused to maltose-binding protein (MBP), all or parts of the 24P4C12 cDNA protein coding sequence are fused to the MBP gene by cloning into the pMAL-c2X and pMAL-p2X vectors (New England Biolabs, Beverly, Mass.). These constructs allow controlled expression of recombinant 24P4C12 protein sequences with MBP fused at the amino-terminus and a 6×His epitope tag at the carboxyl-terminus. The MBP and 6×His tags permit purification of the recombinant protein from induced bacteria with the appropriate affinity matrix and allow recognition of the fusion protein with anti-MBP and anti-His antibodies. The 6×His epitope tag is generated by adding 6 histidine codons to the 3' cloning primer. A Factor Xa recognition site permits cleavage of the pMAL tag from 24P4C12. The pMAL-c2X and pMAL-p2X vectors are optimized to express the recombinant protein in the cytoplasm or periplasm respectively. Periplasm expression enhances folding of proteins with disulfide bonds.

pET Constructs: To express 24P4C12 in bacterial cells, all or parts of the 24P4C12 cDNA protein coding sequence are cloned into the pET family of vectors (Novagen, Madison, Wis.). These vectors allow tightly controlled expression of recombinant 24P4C12 protein in bacteria with and without fusion to proteins that enhance solubility, such as NusA and thioredoxin (Trx), and epitope tags, such as 6×His and S-Tag™ that aid purification and detection of the recombinant protein. For example, constructs are made utilizing pET NusA fusion system 43.1 such that regions of the 24P4C12 protein are expressed as amino-terminal fusions to NusA.

C. Yeast Constructs:

pESC Constructs: To express 24P4C12 in the yeast species *Saccharomyces cerevisiae* for generation of recombinant protein and functional studies, all or parts of the 24P4C12 cDNA protein coding sequence are cloned into the pESC family of vectors each of which contain 1 of 4 selectable markers, HIS3, TRP1, LEU2, and URA3 (Stratagene, La Jolla, Calif.). These vectors allow controlled expression from the same plasmid of up to 2 different genes or cloned sequences containing either Flag™ or Myc epitope tags in the same yeast cell. This system is useful to confirm protein-protein interactions of 24P4C12. In addition, expression in yeast yields similar post-translational modifications, such as glycosylations and phosphorylations, that are found when expressed in eukaryotic cells.

pESP Constructs: To express 24P4C12 in the yeast species *Saccharomyces pombe*, all or parts of the 24P4C12 cDNA protein coding sequence are cloned into the pESP family of vectors. These vectors allow controlled high level of expression of a 24P4C12 protein sequence that is fused at either the amino terminus or at the carboxyl terminus to GST which aids purification of the recombinant protein. A Flag™ epitope tag allows detection of the recombinant protein with anti-Flag™ antibody.

Production of Recombinant 24P4C12 in Higher Eukaryotic Systems

A. Mammalian Constructs:

To express recombinant 24P4C12 in eukaryotic cells, the full or partial length 24P4C12 cDNA sequences can be cloned into any one of a variety of expression vectors known in the art. One or more of the following regions of 24P4C12 are expressed in these constructs, amino acids 1 to 710, or any 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30 or more contiguous amino acids from 24P4C12 v.1 through v.6; amino acids 1 to 598, or any 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30 or more contiguous amino acids from 24P4C12 v.7, amino acids 1 to 722, or any 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30 or more contiguous amino acids from 24P4C12 v.8, amino acids 1 to 712, or any 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30 or more contiguous amino acids from 24P4C12 v.9, variants, or analogs thereof.

The constructs can be transfected into any one of a wide variety of mammalian cells such as 293T cells. Transfected 293T cell lysates can be probed with the anti-24P4C12 polyclonal serum, described herein.

pcDNA3.1/MycHis Constructs: To express 24P4C12 in mammalian cells, a 24P4C12 ORF, or portions thereof, of 24P4C12 with a consensus Kozak translation initiation site was cloned into pcDNA3.1/MycHis Version A (Invitrogen, Carlsbad, Calif.). Protein expression is driven from the cytomegalovirus (CMV) promoter. The recombinant proteins have the myc epitope and 6xHis epitope fused to the carboxyl-terminus. The pcDNA3.1/MycHis vector also contains the bovine growth hormone (BGH) polyadenylation signal and transcription termination sequence to enhance mRNA stability, along with the SV40 origin for episomal replication and simple vector rescue in cell lines expressing the large T antigen. The Neomycin resistance gene can be used, as it allows for selection of mammalian cells expressing the protein and the ampicillin resistance gene and ColE1 origin permits selection and maintenance of the plasmid in *E. coli*. Figure Art-1 demonstrates expression of 24P4C12 from the pcDNA3.1/MycHis construct in transiently transfected 293T cells.

pcDNA4/HisMax Constructs: To express 24P4C12 in mammalian cells, a 24P4C12 ORF, or portions thereof, of 24P4C12 are cloned into pcDNA4/HisMax Version A (Invitrogen, Carlsbad, Calif.). Protein expression is driven from the cytomegalovirus (CMV) promoter and the SP16 translational enhancer. The recombinant protein has XpreSS™ and six histidine (6xHis) epitopes fused to the amino-terminus. The pcDNA4/HisMax vector also contains the bovine growth hormone (BGH) polyadenylation signal and transcription termination sequence to enhance mRNA stability along with the SV40 origin for episomal replication and simple vector rescue in cell lines expressing the large T antigen. The Zeocin resistance gene allows for selection of mammalian cells expressing the protein and the ampicillin resistance gene and ColE1 origin permits selection and maintenance of the plasmid in *E. coli*.

pcDNA3.1/CT-GFP-TOPO Construct: To express 24P4C12 in mammalian cells and to allow detection of the recombinant proteins using fluorescence, a 24P4C12 ORF, or portions thereof, with a consensus Kozak translation initiation site are cloned into pcDNA3.1/CT-GFP-TOPO (Invitrogen, Calif.). Protein expression is driven from the cytomega-

lovirus (CMV) promoter. The recombinant proteins have the Green Fluorescent Protein (GFP) fused to the carboxyl-terminus facilitating non-invasive, in vivo detection and cell biology studies. The pcDNA3.1 CT-GFP-TOPO vector also contains the bovine growth hormone (BGH) polyadenylation signal and transcription termination sequence to enhance mRNA stability along with the SV40 origin for episomal replication and simple vector rescue in cell lines expressing the large T antigen. The Neomycin resistance gene allows for selection of mammalian cells that express the protein and the ampicillin resistance gene and ColE1 origin permits selection and maintenance of the plasmid in *E. coli*. Additional constructs with an amino-terminal GFP fusion are made in pcDNA3.1/NT-GFP-TOPO spanning the entire length of a 24P4C12 protein.

pTag5: A 24P4C12 ORF, or portions thereof, were cloned into pTag-5. This vector is similar to pAptag but without the alkaline phosphatase fusion. This construct generates 24P4C12 protein with an amino-terminal IgGk signal sequence and myc and 6xHis epitope tags at the carboxyl-terminus that facilitate detection and affinity purification. The resulting recombinant 24P4C12 protein were optimized for secretion into the media of transfected mammalian cells, and is used as immunogen or ligand to identify proteins such as ligands or receptors that interact with the 24P4C12 proteins. Protein expression is driven from the CMV promoter. The Zeocin resistance gene present in the vector allows for selection of mammalian cells expressing the protein, and the ampicillin resistance gene permits selection of the plasmid in *E. coli*. Figure Art-3 shows expression of 24P4C12 from two different pTag5 constructs.

PAPtag: A 24P4C12 ORF, or portions thereof, is cloned into pAptag-5 (GenHunter Corp. Nashville, Tenn.). This construct generates an alkaline phosphatase fusion at the carboxyl-terminus of a 24P4C12 protein while fusing the IgGk signal sequence to the amino-terminus. Constructs are also generated in which alkaline phosphatase with an amino-terminal IgGk signal sequence is fused to the amino-terminus of a 24P4C12 protein. The resulting recombinant 24P4C12 proteins are optimized for secretion into the media of transfected mammalian cells and can be used to identify proteins such as ligands or receptors that interact with 24P4C12 proteins. Protein expression is driven from the CMV promoter and the recombinant proteins also contain myc and 6xHis epitopes fused at the carboxyl-terminus that facilitates detection and purification. The Zeocin resistance gene present in the vector allows for selection of mammalian cells expressing the recombinant protein and the ampicillin resistance gene permits selection of the plasmid in *E. coli*.

PsecFc: A 24P4C12 ORF, or portions thereof, is also cloned into psecFc. The psecFc vector was assembled by cloning the human immunoglobulin G1 (IgG) Fc (hinge, CH2, CH3 regions) into pSecTag2 (Invitrogen, California). This construct generates an IgG1 Fc fusion at the carboxyl-terminus of the 24P4C12 proteins, while fusing the IgGk signal sequence to N-terminus. 24P4C12 fusions utilizing the murine IgG1 Fc region are also used. The resulting recombinant 24P4C12 proteins are optimized for secretion into the media of transfected mammalian cells, and can be used as immunogens or to identify proteins such as ligands or receptors that interact with 24P4C12 protein. Protein expression is driven from the CMV promoter. The hygromycin resistance gene present in the vector allows for selection of mammalian cells that express the recombinant protein, and the ampicillin resistance gene permits selection of the plasmid in *E. coli*.

pSRα Constructs: To generate mammalian cell lines that express 24P4C12 constitutively, 24P4C12 ORF, or portions

thereof, of 24P4C12 were cloned into pSR α constructs. Amphotrophic and ecotropic retroviruses were generated by transfection of pSR α constructs into the 293T-10A1 packaging line or co-transfection of pSR α and a helper plasmid (containing deleted packaging sequences) into the 293 cells, respectively. The retrovirus is used to infect a variety of mammalian cell lines, resulting in the integration of the cloned gene, 24P4C12, into the host cell-lines. Protein expression is driven from a long terminal repeat (LTR). The Neomycin resistance gene present in the vector allows for selection of mammalian cells that express the protein, and the ampicillin resistance gene and ColE1 origin permit selection and maintenance of the plasmid in *E. coli*. The retroviral vectors can thereafter be used for infection and generation of various cell lines using, for example, PC3, NIH 3T3, TsuPr1, 293 or rat-1 cells. FIG. 23 shows RNA expression of 24P4C12 driven from the 24P4C12.pSR α construct in stably transduced PC3, 3T3 and 300.19 cells. Figure Art-2 shows 24P4C12 protein expression in PC3 cells stably transduced with 24P4C12.pSR α construct.

Additional pSR α constructs are made that fuse an epitope tag such as the FLAGTM tag to the carboxyl-terminus of 24P4C12 sequences to allow detection using anti-Flag antibodies. For example, the FLAGTM sequence 5' gat tac aag gat gac gac gat aag 3' (SEQ ID NO: 45) is added to cloning primer at the 3' end of the ORF. Additional pSR α constructs are made to produce both amino-terminal and carboxyl-terminal GFP and myc/6xHis fusion proteins of the full-length 24P4C12 proteins.

Additional Viral Vectors: Additional constructs are made for viral-mediated delivery and expression of 24P4C12. High virus titer leading to high level expression of 24P4C12 is achieved in viral delivery systems such as adenoviral vectors and herpes amplicon vectors. A 24P4C12 coding sequences or fragments thereof are amplified by PCR and subcloned into the AdEasy shuttle vector (Stratagene). Recombination and virus packaging are performed according to the manufacturer's instructions to generate adenoviral vectors. Alternatively, 24P4C12 coding sequences or fragments thereof are cloned into the HSV-1 vector (Imgenex) to generate herpes viral vectors. The viral vectors are thereafter used for infection of various cell lines such as PC3, NIH 3T3, 293 or rat-1 cells.

Regulated Expression Systems: To control expression of 24P4C12 in mammalian cells, coding sequences of 24P4C12, or portions thereof, are cloned into regulated mammalian expression systems such as the T-Rex System (Invitrogen), the GeneSwitch System (Invitrogen) and the tightly-regulated Ecdysone System (Stratagene). These systems allow the study of the temporal and concentration dependent effects of recombinant 24P4C12. These vectors are thereafter used to control expression of 24P4C12 in various cell lines such as PC3, NIH 3T3, 293 or rat-1 cells.

B. Baculovirus Expression Systems

To generate recombinant 24P4C12 proteins in a baculovirus expression system, 24P4C12 ORF, or portions thereof, are cloned into the baculovirus transfer vector pBlueBac 4.5 (Invitrogen), which provides a His-tag at the N-terminus. Specifically, pBlueBac-24P4C12 is co-transfected with helper plasmid pBac-N-Blue (Invitrogen) into SF9 (*Spodoptera frugiperda*) insect cells to generate recombinant baculovirus (see Invitrogen instruction manual for details). Baculovirus is then collected from cell supernatant and purified by plaque assay.

Recombinant 24P4C12 protein is then generated by infection of HighFive insect cells (Invitrogen) with purified baculovirus. Recombinant 24P4C12 protein can be detected using anti-24P4C12 or anti-His-tag antibody. 24P4C12 protein can

be purified and used in various cell-based assays or as immunogen to generate polyclonal and monoclonal antibodies specific for 24P4C12.

Example 9

Antigenicity Profiles and Secondary Structure

FIGS. 5-9 depict graphically five amino acid profiles of the 24P4C12 variant 1, assessment available by accessing the ProtScale website located on the World Wide Web at (expasy.ch/cgi-bin/protscale.pl) on the ExPasy molecular biology server.

These profiles: FIG. 5, Hydrophilicity, (Hopp T. P., Woods K. R., 1981. Proc. Natl. Acad. Sci. U.S.A. 78:3824-3828); FIG. 6, Hydropathicity, (Kyte J., Doolittle R. F., 1982. J. Mol. Biol. 157:105-132); FIG. 7, Percentage Accessible Residues (Janin J., 1979 Nature 277:491-492); FIG. 8, Average Flexibility, (Bhaskaran R., and Ponnuswamy P. K., 1988. Int. J. Pept. Protein Res. 32:242-255); FIG. 9, Beta-turn (Deleage, G., Roux B. 1987 Protein Engineering 1:289-294); and optionally others available in the art, such as on the ProtScale website, were used to identify antigenic regions of the 24P4C12 protein. Each of the above amino acid profiles of 24P4C12 were generated using the following ProtScale parameters for analysis: 1) A window size of 9; 2) 100% weight of the window edges compared to the window center; and, 3) amino acid profile values normalized to lie between 0 and 1.

Hydrophilicity (FIG. 5), Hydropathicity (FIG. 6) and Percentage Accessible Residues (FIG. 7) profiles were used to determine stretches of hydrophilic amino acids (i.e., values greater than 0.5 on the Hydrophilicity and Percentage Accessible Residues profile, and values less than 0.5 on the Hydropathicity profile). Such regions are likely to be exposed to the aqueous environment, be present on the surface of the protein, and thus available for immune recognition, such as by antibodies.

Average Flexibility (FIG. 8) and Beta-turn (FIG. 9) profiles determine stretches of amino acids (i.e., values greater than 0.5 on the Beta-turn profile and the Average Flexibility profile) that are not constrained in secondary structures such as beta sheets and alpha helices. Such regions are also more likely to be exposed on the protein and thus accessible to immune recognition, such as by antibodies.

Antigenic sequences of the 24P4C12 protein and of the variant proteins indicated, e.g., by the profiles set forth in FIG. 5, FIG. 6, FIG. 7, FIG. 8, and/or FIG. 9 are used to prepare immunogens, either peptides or nucleic acids that encode them, to generate therapeutic and diagnostic anti-24P4C12 antibodies. The immunogen can be any 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 30, 35, 40, 45, 50 or more than 50 contiguous amino acids, or the corresponding nucleic acids that encode them, from the 24P4C12 protein variants listed in FIGS. 2 and 3. In particular, peptide immunogens of the invention can comprise, a peptide region of at least 5 amino acids of FIGS. 2 and 3 in any whole number increment that includes an amino acid position having a value greater than 0.5 in the Hydrophilicity profile of FIG. 5; a peptide region of at least 5 amino acids of FIGS. 2 and 3 in any whole number increment that includes an amino acid position having a value less than 0.5 in the Hydropathicity profile of FIG. 6; a peptide region of at least 5 amino acids of FIGS. 2 and 3 in any whole number increment that includes an amino acid position having a value greater than 0.5 in the Percent Accessible Residues profile of FIG. 7; a peptide region of at least 5 amino acids of FIGS. 2 and 3 in any whole number

increment that includes an amino acid position having a value greater than 0.5 in the Average Flexibility profile on FIG. 8; and, a peptide region of at least 5 amino acids of FIGS. 2 and 3 in any whole number increment that includes an amino acid position having a value greater than 0.5 in the Beta-turn profile of FIG. 9. Peptide immunogens of the invention can also comprise nucleic acids that encode any of the foregoing.

All immunogens of the invention, peptide or nucleic acid, can be embodied in human unit dose form, or comprised by a composition that includes a pharmaceutical excipient compatible with human physiology.

The secondary structure of 24P4C12 variant 1, namely the predicted presence and location of alpha helices, extended strands, and random coils, are predicted from the respective primary amino acid sequences using the HNN—Hierarchical Neural Network method (Guerneur, 1997, http://pbil.ibcp.fr/cgi-bin/npsa_automat.pl?page=npsa_nn.html), accessed from the ExPasy molecular biology server (<http://www.expasy.ch/tools/>). The analysis indicates that 24P4C12 variant 1 is composed of 53.94% alpha helix, 9.44% extended strand, and 36.62% random coil (FIG. 13a). Analysis for the potential presence of transmembrane domains in 24P4C12 variants were carried out using a variety of transmembrane prediction algorithms accessed from the ExPasy molecular biology server (<http://www.expasy.ch/tools/>). Shown graphically are the results of analysis of variant 1 depicting the presence and location of 10 transmembrane domains using the TMpred program (FIG. 13b) and TMHMM program (FIG. 13c). The results of each program, namely the amino acids encoding the transmembrane domains are summarized in Table L.

Example 10

Generation of 24P4C12 Polyclonal Antibodies

Polyclonal antibodies can be raised in a mammal, for example, by one or more injections of an immunizing agent and, if desired, an adjuvant. Typically, the immunizing agent and/or adjuvant will be injected in the mammal by multiple subcutaneous or intraperitoneal injections. In addition to immunizing with the full length 24P4C12 protein, computer algorithms are employed in design of immunogens that, based on amino acid sequence analysis contain characteristics of being antigenic and available for recognition by the immune system of the immunized host (see the Example entitled "Antigenicity Profiles"). Such regions would be predicted to be hydrophilic, flexible, in beta-turn conformations, and be exposed on the surface of the protein (see, e.g., FIG. 5, FIG. 6, FIG. 7, FIG. 8, or FIG. 9 for amino acid profiles that indicate such regions of 24P4C12 and variants).

For example, 24P4C12 recombinant bacterial fusion proteins or peptides containing hydrophilic, flexible, beta-turn regions of 24P4C12 variant proteins are used as antigens to generate polyclonal antibodies in New Zealand White rabbits. For example, such regions include, but are not limited to, amino acids 1-34, amino acids 118-135, amino acids 194-224, amino acids 280-290, and amino acids 690-710, of 24P4C12 variants 1. It is useful to conjugate the immunizing agent to a protein known to be immunogenic in the mammal being immunized. Examples of such immunogenic proteins include, but are not limited to, keyhole limpet hemocyanin (KLH), serum albumin, bovine thyroglobulin, and soybean trypsin inhibitor. In one embodiment, a peptide encoding amino acids 1-14 of 24P4C12 variant 1 was conjugated to KLH and used to immunize a rabbit. This antiserum exhibited a high titer to the peptide (>10,000) and recognized 24P4C12 in transfected 293T cells by Western blot and flow cytometry

(FIG. 24) and in stable recombinant PC3 cells by Western blot and immunohistochemistry (FIG. 25). Alternatively the immunizing agent may include all or portions of the 24P4C12 variant proteins, analogs or fusion proteins thereof. For example, the 24P4C12 variant 1 amino acid sequence can be fused using recombinant DNA techniques to any one of a variety of fusion protein partners that are well known in the art, such as glutathione-S-transferase (GST) and HIS tagged fusion proteins. Such fusion proteins are purified from induced bacteria using the appropriate affinity matrix.

In one embodiment, a GST-fusion protein encoding amino acids 379-453, encompassing the third predicted extracellular loop of variant 1, is produced, purified, and used as immunogen. Other recombinant bacterial fusion proteins that may be employed include maltose binding protein, LacZ, thioredoxin, NusA, or an immunoglobulin constant region (see the section entitled "Production of 24P4C12 in Prokaryotic Systems" and Current Protocols In Molecular Biology, Volume 2, Unit 16, Frederick M. Ausubul et al. eds., 1995; Linsley, P.S., Brady, W., Urnes, M., Grosmaire, L., Damle, N., and Ledbetter, L. (1991) J. Exp. Med. 174, 561-566).

In addition to bacterial derived fusion proteins, mammalian expressed protein antigens are also used. These antigens are expressed from mammalian expression vectors such as the Tag5 and Fc-fusion vectors (see the Example entitled "Production of Recombinant 24P4C12 in Eukaryotic Systems"), and retains post-translational modifications such as glycosylations found in native protein. In two embodiments, the predicted 1st and third extracellular loops of variant 1, amino acids 59-227 and 379-453 respectively, were each cloned into the Tag5 mammalian secretion vector and expressed in 293T cells (FIG. 26). Each recombinant protein is then purified by metal chelate chromatography from tissue culture supernatants and/or lysates of 293T cells stably expressing the recombinant vector. The purified Tag5 24P4C12 protein is then used as immunogen.

During the immunization protocol, it is useful to mix or emulsify the antigen in adjuvants that enhance the immune response of the host animal. Examples of adjuvants include, but are not limited to, complete Freund's adjuvant (CFA) and MPL-TDM adjuvant (monophosphoryl Lipid A, synthetic trehalose dicorynomycolate).

In a typical protocol, rabbits are initially immunized subcutaneously with up to 200 µg, typically 100-200 µg, of fusion protein or peptide conjugated to KLH mixed in complete Freund's adjuvant (CFA). Rabbits are then injected subcutaneously every two weeks with up to 200 µg, typically 100-200 µg, of the immunogen in incomplete Freund's adjuvant (IFA). Test bleeds are taken approximately 7-10 days following each immunization and used to monitor the titer of the antiserum by ELISA.

To test reactivity and specificity of immune serum, such as the rabbit serum derived from immunization with a KLH-conjugated peptide encoding amino acids 1-14 of variant 1, the full-length 24P4C12 variant 1 cDNA is cloned into pcDNA 3.1 myc-his or retroviral expression vectors (Invitrogen, see the Example entitled "Production of Recombinant 24P4C12 in Eukaryotic Systems"). After transfection of the constructs into 293T cells or transduction of PC3 with 24P4C12 retrovirus, cell lysates are probed with the anti-24P4C12 serum and with anti-His antibody (Santa Cruz Biotechnologies, Santa Cruz, Calif.) to determine specific reactivity to denatured 24P4C12 protein using the Western blot technique. As shown in FIGS. 24 and 25 the antiserum specifically recognizes 24P4C12 protein in 293T and PC3 cells. In addition, the immune serum is tested by fluorescence microscopy, flow cytometry, and immunohistochemistry

(FIG. 25) and immunoprecipitation against 293T and other recombinant 24P4C12-expressing cells to determine specific recognition of native protein. Western blot, immunoprecipitation, fluorescent microscopy, immunohistochemistry and flow cytometric techniques using cells that endogenously express 24P4C12 are also carried out to test reactivity and specificity.

Anti-serum from rabbits immunized with 24P4C12 variant fusion proteins, such as GST and MBP fusion proteins, are purified by depletion of antibodies reactive to the fusion partner sequence by passage over an affinity column containing the fusion partner either alone or in the context of an irrelevant fusion protein. For example, antiserum derived from a GST-24P4C12 fusion protein encoding amino acids 379-453 of variant 1 is first purified by passage over a column of GST protein covalently coupled to AffiGel matrix (BioRad, Hercules, Calif.). The antiserum is then affinity purified by passage over a column composed of a MBP-fusion protein also encoding amino acids 379-453 covalently coupled to Affigel matrix. The serum is then further purified by protein G affinity chromatography to isolate the IgG fraction. Sera from other His-tagged antigens and peptide immunized rabbits as well as fusion partner depleted sera are affinity purified by passage over a column matrix composed of the original protein immunogen or free peptide.

Example 11

Generation of 24P4C12 Monoclonal Antibodies (mAbs)

In one embodiment, therapeutic mAbs to 24P4C12 variants comprise those that react with epitopes specific for each variant protein or specific to sequences in common between the variants that would disrupt or modulate the biological function of the 24P4C12 variants, for example those that would disrupt the interaction with ligands and substrates or disrupt its biological activity. Immunogens for generation of such mAbs include those designed to encode or contain the entire 24P4C12 protein variant sequence, regions of the 24P4C12 protein variants predicted to be antigenic from computer analysis of the amino acid sequence (see, e.g., FIG. 5, FIG. 6, FIG. 7, FIG. 8, or FIG. 9, and the Example entitled "Antigenicity Profiles"). Immunogens include peptides, recombinant bacterial proteins, and mammalian expressed Tag 5 proteins and human and murine IgG FC fusion proteins. In addition, cells engineered to express high levels of a respective 24P4C12 variant, such as 293T-24P4C12 variant 1 or 300.19-24P4C12 variant 1 murine Pre-B cells, are used to immunize mice.

To generate mAbs to a 24P4C12 variant, mice are first immunized intraperitoneally (IP) with, typically, 10-50 µg of protein immunogen or 10⁷ 24P4C12-expressing cells mixed in complete Freund's adjuvant. Mice are then subsequently immunized IP every 2-4 weeks with, typically, 10-50 µg of protein immunogen or 10⁷ cells mixed in incomplete Freund's adjuvant. Alternatively, MPL-TDM adjuvant is used in immunizations. In one embodiment, mice were immunized as above with 300.19-24P4C12 cells in complete and then incomplete Freund's adjuvant, and subsequently sacrificed and the spleens harvested and used for fusion and hybridoma generation. As is can be seen in FIG. 27, 2 hybridomas were generated whose antibodies specifically recognize 24P4C12 protein expressed in 293T cells by flow cytometry. In addition to the above protein and cell-based immunization strategies, a DNA-based immunization protocol is employed in which a mammalian expression vector encoding a 24P4C12 variant

sequence is used to immunize mice by direct injection of the plasmid DNA. In one embodiment, a Tag5 mammalian secretion vector encoding amino acids 59-227 of the variant 1 sequence (FIG. 26) was used to immunize mice. Subsequent booster immunizations are then carried out with the purified protein. In another example, the same amino acids are cloned into an Fc-fusion secretion vector in which the 24P4C12 variant 1 sequence is fused at the amino-terminus to an IgK leader sequence and at the carboxyl-terminus to the coding sequence of the human or murine IgG Fc region. This recombinant vector is then used as immunogen. The plasmid immunization protocols are used in combination with purified proteins as above and with cells expressing the respective 24P4C12 variant.

During the immunization protocol, test bleeds are taken 7-10 days following an injection to monitor titer and specificity of the immune response. Once appropriate reactivity and specificity is obtained as determined by ELISA, Western blotting, immunoprecipitation, fluorescence microscopy, immunohistochemistry, and flow cytometric analyses, fusion and hybridoma generation is then carried out with established procedures well known in the art (see, e.g., Harlow and Lane, 1988).

In one embodiment for generating 24P4C12 variant 8 specific monoclonal antibodies, a peptide encoding amino acids 643-654 (RNPITPTIGHVFQ) (SEQ ID NO: 46) of 24P4C12 variant 8 is synthesized, coupled to KLH and used as immunogen. Balb C mice are initially immunized intraperitoneally with 25 µg of the KLH-24P4C12 variant 8 peptide mixed in complete Freund's adjuvant. Mice are subsequently immunized every two weeks with 25 µg of the antigen mixed in incomplete Freund's adjuvant for a total of three immunizations. ELISA using the free peptide determines the reactivity of serum from immunized mice. Reactivity and specificity of serum to full length 24P4C12 variant 8 protein is monitored by Western blotting, immunoprecipitation and flow cytometry using 293T cells transfected with an expression vector encoding the 24P4C12 variant 8 cDNA compared to cells transfected with the other 24P4C12 variants (see e.g., the Example entitled "Production of Recombinant 24P4C12 in Eukaryotic Systems"). Other recombinant 24P4C12 variant 8-expressing cells or cells endogenously expressing 24P4C12 variant 8 are also used. Mice showing the strongest specific reactivity to 24P4C12 variant 8 are rested and given a final injection of antigen in PBS and then sacrificed four days later. The spleens of the sacrificed mice are harvested and fused to SPO/2 myeloma cells using standard procedures (Harlow and Lane, 1988). Supernatants from HAT selected growth wells are screened by ELISA, Western blot, immunoprecipitation, fluorescent microscopy, and flow cytometry to identify 24P4C12 variant 8-specific antibody-producing clones. A similar strategy is also used to derive 24P4C12 variant 9-specific antibodies using a peptide encompassing amino acids 379-388 (PLPTOPATLG) (SEQ ID NO: 47).

The binding affinity of a 24P4C12 monoclonal antibody is determined using standard technologies. Affinity measurements quantify the strength of antibody to epitope binding and are used to help define which 24P4C12 monoclonal antibodies preferred for diagnostic or therapeutic use, as appreciated by one of skill in the art. The BIAcore system (Uppsala, Sweden) is a preferred method for determining binding affinity. The BIAcore system uses surface plasmon resonance (SPR, Welford K. 1991, Opt. Quant. Elect. 23:1; Morton and Myszk, 1998, Methods in Enzymology 295: 268) to monitor bimolecular interactions in real time. BIAcore analysis con-

veniently generates association rate constants, dissociation rate constants, equilibrium dissociation constants, and affinity constants.

Example 12

HLA Class I and Class II Binding Assays

HLA class I and class II binding assays using purified HLA molecules are performed in accordance with disclosed protocols (e.g., PCT publications WO 94/20127 and WO 94/03205; Sidney et al., *Current Protocols in Immunology* 18.3.1 (1998); Sidney, et al., *J. Immunol.* 154:247 (1995); Sette, et al., *Mol. Immunol.* 31:813 (1994)). Briefly, purified MHC molecules (5 to 500 nM) are incubated with various unlabeled peptide inhibitors and 1-10 nM Q¹²⁵I-radiolabeled probe peptides as described. Following incubation, MHC-peptide complexes are separated from free peptide by gel filtration and the fraction of peptide bound is determined. Typically, in preliminary experiments, each MHC preparation is titrated in the presence of fixed amounts of radiolabeled peptides to determine the concentration of HLA molecules necessary to bind 10-20% of the total radioactivity. All subsequent inhibition and direct binding assays are performed using these HLA concentrations.

Since under these conditions $[label] \ll [HLA]$ and $IC_{50} \gg [HLA]$, the measured IC_{50} values are reasonable approximations of the true K_D values. Peptide inhibitors are typically tested at concentrations ranging from 120 µg/ml to 1.2 ng/ml, and are tested in two to four completely independent experiments. To allow comparison of the data obtained in different experiments, a relative binding figure is calculated for each peptide by dividing the IC_{50} of a positive control for inhibition by the IC_{50} for each tested peptide (typically unlabeled versions of the radiolabeled probe peptide). For database purposes, and inter-experiment comparisons, relative binding values are compiled. These values can subsequently be converted back into IC_{50} nM values by dividing the IC_{50} nM of the positive controls for inhibition by the relative binding of the peptide of interest. This method of data compilation is accurate and consistent for comparing peptides that have been tested on different days, or with different lots of purified MHC.

Binding assays as outlined above may be used to analyze HLA supermotif and/or HLA motif-bearing peptides (see Table IV).

Example 13

Identification of HLA Supermotif- and Motif-Bearing CTL Candidate Epitopes

HLA vaccine compositions of the invention can include multiple epitopes. The multiple epitopes can comprise multiple HLA supermotifs or motifs to achieve broad population coverage. This example illustrates the identification and confirmation of supermotif- and motif-bearing epitopes for the inclusion in such a vaccine composition. Calculation of population coverage is performed using the strategy described below.

Computer Searches and Algorithms for Identification of Supermotif and/or Motif-Bearing Epitopes

The searches performed to identify the motif-bearing peptide sequences in the Example entitled "Antigenicity Profiles" and Tables VIII-XXI and XXII-XLIX employ the protein sequence data from the gene product of 24P4C12 set

forth in FIGS. 2 and 3, the specific search peptides used to generate the tables are listed in Table VII.

Computer searches for epitopes bearing HLA Class I or Class II supermotifs or motifs are performed as follows. All translated 24P4C12 protein sequences are analyzed using a text string search software program to identify potential peptide sequences containing appropriate HLA binding motifs; such programs are readily produced in accordance with information in the art in view of known motif/supermotif disclosures. Furthermore, such calculations can be made mentally.

Identified A2-, A3-, and DR-supermotif sequences are scored using polynomial algorithms to predict their capacity to bind to specific HLA-Class I or Class II molecules. These polynomial algorithms account for the impact of different amino acids at different positions, and are essentially based on the premise that the overall affinity (or AG) of peptide-HLA molecule interactions can be approximated as a linear polynomial function of the type:

$$-\Delta G = a_{1i} \times a_{2j} \times a_{3k} \dots \times a_{ni}$$

where a_{ji} is a coefficient which represents the effect of the presence of a given amino acid (j) at a given position (i) along the sequence of a peptide of n amino acids. The crucial assumption of this method is that the effects at each position are essentially independent of each other (i.e., independent binding of individual side-chains). When residue j occurs at position i in the peptide, it is assumed to contribute a constant amount i to the free energy of binding of the peptide irrespective of the sequence of the rest of the peptide.

The method of derivation of specific algorithm coefficients has been described in Gulukota et al., *J. Mol. Biol.* 267:1258-126, 1997; (see also Sidney et al., *Human Immunol.* 45:79-93, 1996; and Southwood et al., *J. Immunol.* 160:3363-3373, 1998). Briefly, for all i positions, anchor and non-anchor alike, the geometric mean of the average relative binding (ARB) of all peptides carrying j is calculated relative to the remainder of the group, and used as the estimate of j_i . For Class II peptides, if multiple alignments are possible, only the highest scoring alignment is utilized, following an iterative procedure. To calculate an algorithm score of a given peptide in a test set, the ARB values corresponding to the sequence of the peptide are multiplied. If this product exceeds a chosen threshold, the peptide is predicted to bind. Appropriate thresholds are chosen as a function of the degree of stringency of prediction desired.

Selection of HLA-A2 Supertype Cross-Reactive Peptides

Protein sequences from 24P4C12 are scanned utilizing motif identification software, to identify 8-, 9- 10- and 11-mer sequences containing the HLA-A2-supermotif main anchor specificity. Typically, these sequences are then scored using the protocol described above and the peptides corresponding to the positive-scoring sequences are synthesized and tested for their capacity to bind purified HLA-A*0201 molecules in vitro (HLA-A*0201 is considered a prototype A2 supertype molecule).

These peptides are then tested for the capacity to bind to additional A2-superotype molecules (A*0202, A*0203, A*0206, and A*6802). Peptides that bind to at least three of the five A2-superotype alleles tested are typically deemed A2-superotype cross-reactive binders. Preferred peptides bind at an affinity equal to or less than 500 nM to three or more HLA-A2 supertype molecules.

Selection of HLA-A3 Supermotif-Bearing Epitopes

The 24P4C12 protein sequence(s) scanned above is also examined for the presence of peptides with the HLA-A3-supermotif primary anchors. Peptides corresponding to the HLA A3 supermotif-bearing sequences are then synthesized

and tested for binding to HLA-A*0301 and HLA-A*1101 molecules, the molecules encoded by the two most prevalent A3-supertype alleles. The peptides that bind at least one of the two alleles with binding affinities of ≤ 500 nM, often ≤ 200 nM, are then tested for binding cross-reactivity to the other common A3-supertype alleles (e.g., A*3101, A*3301, and A*6801) to identify those that can bind at least three of the five HLA-A3-supertype molecules tested.

Selection of HLA-B7 Supermotif Bearing Epitopes

The 24P4C12 protein(s) scanned above is also analyzed for the presence of 8-, 9-, 10-, or 11-mer peptides with the HLA-B7-supermotif. Corresponding peptides are synthesized and tested for binding to HLA-B*0702, the molecule encoded by the most common B7-supertype allele (i.e., the prototype B7 supertype allele). Peptides binding B*0702 with IC_{50} of ≤ 500 nM are identified using standard methods. These peptides are then tested for binding to other common B7-supertype molecules (e.g., B*3501, B*5101, B*5301, and B*5401). Peptides capable of binding to three or more of the five B7-supertype alleles tested are thereby identified.

Selection of A1 and A24 Motif-Bearing Epitopes

To further increase population coverage, HLA-A1 and -A24 epitopes can also be incorporated into vaccine compositions. An analysis of the 24P4C12 protein can also be performed to identify HLA-A1- and A24-motif-containing sequences.

High affinity and/or cross-reactive binding epitopes that bear other motif and/or supermotifs are identified using analogous methodology.

Example 14

Confirmation of Immunogenicity

Cross-reactive candidate CTLA2-supermotif-bearing peptides that are identified as described herein are selected to confirm in vitro immunogenicity. Confirmation is performed using the following methodology:

Target Cell Lines for Cellular Screening:

The 0.221A2.1 cell line, produced by transferring the HLA-A2.1 gene into the HLA-A, -B, -C null mutant human B-lymphoblastoid cell line 721.221, is used as the peptide-loaded target to measure activity of HLA-A2.1-restricted CTL. This cell line is grown in RPMI-1640 medium supplemented with antibiotics, sodium pyruvate, nonessential amino acids and 10% (v/v) heat inactivated FCS. Cells that express an antigen of interest, or transfectants comprising the gene encoding the antigen of interest, can be used as target cells to confirm the ability of peptide-specific CTLs to recognize endogenous antigen.

Primary CTL Induction Cultures:

Generation of Dendritic Cells (DC): PBMCs are thawed in RPMI with 30 μ g/ml DNase, washed twice and resuspended in complete medium (RPMI-1640 plus 5% AB human serum, non-essential amino acids, sodium pyruvate, L-glutamine and penicillin/streptomycin). The monocytes are purified by plating 10×10^6 PBMC/well in a 6-well plate. After 2 hours at 37° C., the non-adherent cells are removed by gently shaking the plates and aspirating the supernatants. The wells are washed a total of three times with 3 ml RPMI to remove most of the non-adherent and loosely adherent cells. Three ml of complete medium containing 50 ng/ml of GM-CSF and 1,000 U/ml of IL-4 are then added to each well. TNF α is added to the DCs on day 6 at 75 ng/ml and the cells are used for CTL induction cultures on day 7.

Induction of CTL with DC and Peptide: CD8+ T-cells are isolated by positive selection with Dynal immunomagnetic

beads (Dynabeads® M-450) and the Detacha-Bead® reagent. Typically about $200\text{--}250 \times 10^6$ PBMC are processed to obtain 24×10^6 CD8+ T-cells (enough for a 48-well plate culture). Briefly, the PBMCs are thawed in RPMI with 30 μ g/ml DNase, washed once with PBS containing 1% human AB serum and resuspended in PBS/1% AB serum at a concentration of 20×10^6 cells/ml. The magnetic beads are washed 3 times with PBS/AB serum, added to the cells (140μ l beads/ 20×10^6 cells) and incubated for 1 hour at 4° C. with continuous mixing. The beads and cells are washed 4 $\times$ with PBS/AB serum to remove the nonadherent cells and resuspended at 100×10^6 cells/ml (based on the original cell number) in PBS/AB serum containing 100 μ l/ml Detacha-Bead® reagent and 30 μ g/ml DNase. The mixture is incubated for 1 hour at room temperature with continuous mixing. The beads are washed again with PBS/AB/DNase to collect the CD8+ T-cells. The DC are collected and centrifuged at 1300 rpm for 5-7 minutes, washed once with PBS with 1% BSA, counted and pulsed with 40 μ g/ml of peptide at a cell concentration of $1\text{--}2 \times 10^6$ /ml in the presence of 3 μ g/ml β_2 -microglobulin for 4 hours at 20° C. The DC are then irradiated (4,200 rads), washed 1 time with medium and counted again.

Setting up induction cultures: 0.25 ml cytokine-generated DC (at 1×10^5 cells/ml) are co-cultured with 0.25 ml of CD8+ T-cells (at 2×10^6 cell/ml) in each well of a 48-well plate in the presence of 10 ng/ml of IL-7. Recombinant human IL-10 is added the next day at a final concentration of 10 ng/ml and rhuman IL-2 is added 48 hours later at 10 IU/ml.

Restimulation of the induction cultures with peptide-pulsed adherent cells: Seven and fourteen days after the primary induction, the cells are restimulated with peptide-pulsed adherent cells. The PBMCs are thawed and washed twice with RPMI and DNase. The cells are resuspended at 5×10^6 cells/ml and irradiated at ~ 4200 rads. The PBMCs are plated at 2×10^6 in 0.5 ml complete medium per well and incubated for 2 hours at 37° C. The plates are washed twice with RPMI by tapping the plate gently to remove the nonadherent cells and the adherent cells pulsed with 1 μ g/ml of peptide in the presence of 3 μ g/ml B2 microglobulin in 0.25 ml RPMI/5% AB per well for 2 hours at 37° C. Peptide solution from each well is aspirated and the wells are washed once with RPMI. Most of the media is aspirated from the induction cultures (CD8+ cells) and brought to 0.5 ml with fresh media. The cells are then transferred to the wells containing the peptide-pulsed adherent cells. Twenty four hours later recombinant human IL-10 is added at a final concentration of 10 ng/ml and recombinant human IL2 is added the next day and again 2-3 days later at 50 IU/ml (Tsai et al., *Critical Reviews in Immunology* 18 (1-2):65-75, 1998). Seven days later, the cultures are assayed for CTL activity in a ^{51}Cr release assay. In some experiments the cultures are assayed for peptide-specific recognition in the in situ IFN γ ELISA at the time of the second restimulation followed by assay of endogenous recognition 7 days later. After expansion, activity is measured in both assays for a side-by-side comparison.

Measurement of CTL Lytic Activity by ^{51}Cr Release.

Seven days after the second restimulation, cytotoxicity is determined in a standard (5 hr) ^{51}Cr release assay by assaying individual wells at a single E:T. Peptide-pulsed targets are prepared by incubating the cells with 10 μ g/ml peptide overnight at 37° C.

Adherent target cells are removed from culture flasks with trypsin-EDTA. Target cells are labeled with 200 μ Ci of ^{51}Cr sodium chromate (Dupont, Wilmington, Del.) for 1 hour at 37° C. Labeled target cells are resuspended at 10^6 per ml and diluted 1:10 with K562 cells at a concentration of 3.3×10^6 /ml (an NK-sensitive erythroleukemia cell line used to reduce

non-specific lysis). Target cells (100 μ l) and effectors (100 μ l) are plated in 96 well round-bottom plates and incubated for 5 hours at 37° C. At that time, 100 μ l of supernatant are collected from each well and percent lysis is determined according to the formula:

$$\frac{[(\text{cpm of the test sample} - \text{cpm of the spontaneous } ^{51}\text{Cr release sample}) / (\text{cpm of the maximal } ^{51}\text{Cr release sample} - \text{cpm of the spontaneous } ^{51}\text{Cr release sample})] \times 100}{}$$

Maximum and spontaneous release are determined by incubating the labeled targets with 1% Triton X-100 and media alone, respectively. A positive culture is defined as one in which the specific lysis (sample-background) is 10% or higher in the case of individual wells and is 15% or more at the two highest E:T ratios when expanded cultures are assayed.

In situ Measurement of Human IFN γ Production as an Indicator of Peptide-Specific and Endogenous Recognition

Immulon 2 plates are coated with mouse anti-human IFN γ monoclonal antibody (4 μ g/ml 0.1M NaHCO $_3$, pH8.2) overnight at 4° C. The plates are washed with Ca $^{2+}$, Mg $^{2+}$ -free PBS/0.05% Tween 20 and blocked with PBS/10% FCS for two hours, after which the CTLs (100 μ l/well) and targets (100 μ l/well) are added to each well, leaving empty wells for the standards and blanks (which received media only). The target cells, either peptide-pulsed or endogenous targets, are used at a concentration of 1 $\times$ 10 ^{6} cells/ml. The plates are incubated for 48 hours at 37° C. with 5% CO $_2$.

Recombinant human IFN-gamma is added to the standard wells starting at 400 pg or 1200 pg/100 microliter/well and the plate incubated for two hours at 37° C. The plates are washed and 100 μ l of biotinylated mouse anti-human IFN-gamma monoclonal antibody (2 microgram/ml in PBS/3% FCS/0.05% Tween 20) are added and incubated for 2 hours at room temperature. After washing again, 100 microliter HRP-streptavidin (1:4000) are added and the plates incubated for one hour at room temperature. The plates are then washed 6 $\times$ with wash buffer, 100 microliter/well developing solution (TMB 1:1) are added, and the plates allowed to develop for 5-15 minutes. The reaction is stopped with 50 microliter/well 1 M H $_3$ PO $_4$ and read at OD450. A culture is considered positive if it measured at least 50 pg of IFN-gamma/well above background and is twice the background level of expression.

CTL Expansion.

Those cultures that demonstrate specific lytic activity against peptide-pulsed targets and/or tumor targets are expanded over a two week period with anti-CD3. Briefly, 5 $\times$ 10 ^{4} CD8 $^{+}$ cells are added to a T25 flask containing the following: 1 $\times$ 10 ^{6} irradiated (4,200 rad) PBMC (autologous or allogeneic) per ml, 2 $\times$ 10 ^{5} irradiated (8,000 rad) EBV-transformed cells per ml, and OKT3 (anti-CD3) at 30 ng per ml in RPMI-1640 containing 10% (v/v) human AB serum, non-essential amino acids, sodium pyruvate, 25 μ M 2-mercaptoethanol, L-glutamine and penicillin/streptomycin. Recombinant human IL2 is added 24 hours later at a final concentration of 200 IU/ml and every three days thereafter with fresh media at 50 IU/ml. The cells are split if the cell concentration exceeds 1 $\times$ 10 ^{6} /ml and the cultures are assayed between days 13 and 15 at E:T ratios of 30, 10, 3 and 1.1 in the ^{51} Cr release assay or at 1 $\times$ 10 ^{6} /ml in the in situ IFN γ assay using the same targets as before the expansion.

Cultures are expanded in the absence of anti-CD3 $^{+}$ as follows. Those cultures that demonstrate specific lytic activity against peptide and endogenous targets are selected and 5 $\times$ 10 ^{4} CD8 $^{+}$ cells are added to a T25 flask containing the following: 1 $\times$ 10 ^{6} autologous PBMC per ml which have been peptide-pulsed with 10 μ g/ml peptide for two hours at 37° C.

and irradiated (4,200 rad); 2 $\times$ 10 ^{5} irradiated (8,000 rad) EBV-transformed cells per ml RPMI-1640 containing 10% (v/v) human AB serum, non-essential AA, sodium pyruvate, 25 mM 2-ME, L-glutamine and gentamicin.

Immunogenicity of A2 Supermotif-Bearing Peptides

A2-supermotif cross-reactive binding peptides are tested in the cellular assay for the ability to induce peptide-specific CTL in normal individuals. In this analysis, a peptide is typically considered to be an epitope if it induces peptide-specific CTLs in at least individuals, and preferably, also recognizes the endogenously expressed peptide.

Immunogenicity can also be confirmed using PBMCs isolated from patients bearing a tumor that expresses 24P4C12. Briefly, PBMCs are isolated from patients, re-stimulated with peptide-pulsed monocytes and assayed for the ability to recognize peptide-pulsed target cells as well as transfected cells endogenously expressing the antigen.

Evaluation of A*03/A11 Immunogenicity

HLA-A3 supermotif-bearing cross-reactive binding peptides are also evaluated for immunogenicity using methodology analogous for that used to evaluate the immunogenicity of the HLA-A2 supermotif peptides.

Evaluation of B7 Immunogenicity

Immunogenicity screening of the B7-supertype cross-reactive binding peptides identified as set forth herein are confirmed in a manner analogous to the confirmation of A2- and A3-supermotif-bearing peptides.

Peptides bearing other supermotifs/motifs, e.g., HLA-A1, HLA-A24 etc. are also confirmed using similar methodology

Example 15

Implementation of the Extended Supermotif to Improve the Binding Capacity of Native Epitopes by Creating Analogs

HLA motifs and supermotifs (comprising primary and/or secondary residues) are useful in the identification and preparation of highly cross-reactive native peptides, as demonstrated herein. Moreover, the definition of HLA motifs and supermotifs also allows one to engineer highly cross-reactive epitopes by identifying residues within a native peptide sequence which can be analogized to confer upon the peptide certain characteristics, e.g. greater cross-reactivity within the group of HLA molecules that comprise a supertype, and/or greater binding affinity for some or all of those HLA molecules. Examples of analogizing peptides to exhibit modulated binding affinity are set forth in this example.

Analoging at Primary Anchor Residues

Peptide engineering strategies are implemented to further increase the cross-reactivity of the epitopes. For example, the main anchors of A2-supermotif-bearing peptides are altered, for example, to introduce a preferred L, I, V, or M at position 2, and I or V at the C-terminus.

To analyze the cross-reactivity of the analog peptides, each engineered analog is initially tested for binding to the prototype A2 supertype allele A*0201, then, if A*0201 binding capacity is maintained, for A2-supertype cross-reactivity.

Alternatively, a peptide is confirmed as binding one or all supertype members and then analogized to modulate binding affinity to any one (or more) of the supertype members to add population coverage.

The selection of analogs for immunogenicity in a cellular screening analysis is typically further restricted by the capacity of the parent wild type (WT) peptide to bind at least weakly, i.e., bind at an IC $_{50}$ of 500 nM or less, to three or more A2 supertype alleles. The rationale for this requirement is that

the WT peptides must be present endogenously in sufficient quantity to be biologically relevant. Analogous peptides have been shown to have increased immunogenicity and cross-reactivity by T cells specific for the parent epitope (see, e.g., Parkhurst et al., *J. Immunol.* 157:2539, 1996; and Pogue et al., *Proc. Natl. Acad. Sci. USA* 92:8166, 1995).

In the cellular screening of these peptide analogs, it is important to confirm that analog-specific CTLs are also able to recognize the wild-type peptide and, when possible, target cells that endogenously express the epitope.

Analoging of HLA-A3 and B7-Supermotif-Bearing Peptides

Analogues of HLA-A3 supermotif-bearing epitopes are generated using strategies similar to those employed in analoging HLA-A2 supermotif-bearing peptides. For example, peptides binding to 3/5 of the A3-supertype molecules are engineered at primary anchor residues to possess a preferred residue (V, S, M, or A) at position 2.

The analog peptides are then tested for the ability to bind A*03 and A*11 (prototype A3 supertype alleles). Those peptides that demonstrate ≤ 500 nM binding capacity are then confirmed as having A3-supertype cross-reactivity.

Similarly to the A2- and A3-motif bearing peptides, peptides binding 3 or more B7-supertype alleles can be improved, where possible, to achieve increased cross-reactive binding or greater binding affinity or binding half life. B7 supermotif-bearing peptides are, for example, engineered to possess a preferred residue (V, I, L, or F) at the C-terminal primary anchor position, as demonstrated by Sidney et al. (*J. Immunol.* 157:3480-3490, 1996).

Analoging at primary anchor residues of other motif and/or supermotif-bearing epitopes is performed in a like manner.

The analog peptides are then be confirmed for immunogenicity, typically in a cellular screening assay. Again, it is generally important to demonstrate that analog-specific CTLs are also able to recognize the wild-type peptide and, when possible, targets that endogenously express the epitope.

Analoging at Secondary Anchor Residues

Moreover, HLA supermotifs are of value in engineering highly cross-reactive peptides and/or peptides that bind HLA molecules with increased affinity by identifying particular residues at secondary anchor positions that are associated with such properties. For example, the binding capacity of a B7 supermotif-bearing peptide with an F residue at position 1 is analyzed. The peptide is then analoged to, for example, substitute L for F at position 1. The analoged peptide is evaluated for increased binding affinity, binding half life and/or increased cross-reactivity. Such a procedure identifies analoged peptides with enhanced properties.

Engineered analogs with sufficiently improved binding capacity or cross-reactivity can also be tested for immunogenicity in HLA-B7-transgenic mice, following for example, IFA immunization or lipopeptide immunization. Analoged peptides are additionally tested for the ability to stimulate a recall response using PBMC from patients with 24P4C12-expressing tumors.

Other Analoging Strategies

Another form of peptide analoging, unrelated to anchor positions, involves the substitution of a cysteine with α -amino butyric acid. Due to its chemical nature, cysteine has the propensity to form disulfide bridges and sufficiently alter the peptide structurally so as to reduce binding capacity. Substitution of α -amino butyric acid for cysteine not only alleviates this problem, but has been shown to improve binding and crossbinding capabilities in some instances (see, e.g., the review by Sette et al., In: *Persistent Viral Infections*, Eds. R. Ahmed and I. Chen, John Wiley & Sons, England, 1999).

Thus, by the use of single amino acid substitutions, the binding properties and/or cross-reactivity of peptide ligands for HLA supertype molecules can be modulated.

Example 16

Identification and Confirmation of 24P4C12-Derived Sequences with HLA-DR Binding Motifs

Peptide epitopes bearing an HLA class II supermotif or motif are identified and confirmed as outlined below using methodology similar to that described for HLA Class I peptides.

Selection of HLA-DR-Supermotif-Bearing Epitopes.

To identify 24P4C12-derived, HLA class II HTL epitopes, a 24P4C12 antigen is analyzed for the presence of sequences bearing an HLA-DR-motif or supermotif. Specifically, 15-mer sequences are selected comprising a DR-supermotif, comprising a 9-mer core, and three-residue N- and C-terminal flanking regions (15 amino acids total).

Protocols for predicting peptide binding to DR molecules have been developed (Southwood et al., *J. Immunol.* 160: 3363-3373, 1998). These protocols, specific for individual DR molecules, allow the scoring, and ranking, of 9-mer core regions. Each protocol not only scores peptide sequences for the presence of DR-supermotif primary anchors (i.e., at position 1 and position 6) within a 9-mer core, but additionally evaluates sequences for the presence of secondary anchors. Using allele-specific selection tables (see, e.g., Southwood et al., *ibid.*), it has been found that these protocols efficiently select peptide sequences with a high probability of binding a particular DR molecule. Additionally, it has been found that performing these protocols in tandem, specifically those for DR1, DR4w4, and DR7, can efficiently select DR cross-reactive peptides.

The 24P4C12-derived peptides identified above are tested for their binding capacity for various common HLA-DR molecules. All peptides are initially tested for binding to the DR molecules in the primary panel: DR1, DR4w4, and DR7. Peptides binding at least two of these three DR molecules are then tested for binding to DR2w2 β 1, DR2w2 β 2, DR6w19, and DR9 molecules in secondary assays. Finally, peptides binding at least two of the four secondary panel DR molecules, and thus cumulatively at least four of seven different DR molecules, are screened for binding to DR4w15, DR5w11, and DR8w2 molecules in tertiary assays. Peptides binding at least seven of the ten DR molecules comprising the primary, secondary, and tertiary screening assays are considered cross-reactive DR binders. 24P4C12-derived peptides found to bind common HLA-DR alleles are of particular interest.

Selection of DR3 Motif Peptides

Because HLA-DR3 is an allele that is prevalent in Caucasian, Black, and Hispanic populations, DR3 binding capacity is a relevant criterion in the selection of HTL epitopes. Thus, peptides shown to be candidates may also be assayed for their DR3 binding capacity. However, in view of the binding specificity of the DR3 motif, peptides binding only to DR3 can also be considered as candidates for inclusion in a vaccine formulation.

To efficiently identify peptides that bind DR3, target 24P4C12 antigens are analyzed for sequences carrying one of the two DR3-specific binding motifs reported by Geluk et al. (*J. Immunol.* 152:5742-5748, 1994). The corresponding peptides are then synthesized and confirmed as having the ability to bind DR3 with an affinity of 1 μ M or better, i.e., less than

Figure 2D-3

```
600 F F G K L L V V G G V G V L S F F F F S
1801 TGTTCCTTTGGGAAGCTGCTGGTGGTCGGAGGCGTGGGGTCTGTCCTTCTTTTTTCT
620 G R I F G L G K D F K S P H L N Y Y W L
1861 CCGGTCGCATCCCGGGGCTGGGTAAAGACTTTAAGAGCCCCACCTCAACTATTACTGGC
640 P I M T S I L G A Y V I A S G F F S V F
1921 TGCCCATCATGACCTCCATCCTGGGGGCTATGTCATCGCCAGCGGCTTCTTCAGCGTTT
660 G M C V D T L F L C F L E D L E R N N G
1981 TCGGCATGTGTGTGGACACGCTCTTCCTCTGCTTCCTGGAAGACCTGGAGCGGAACAACG
680 S L D R P Y Y M S K S L L K I L G K K N
2041 GCTCCCTGGACCGGCCCTACTACATGTCCAAGAGCCTTCTAAAGATTCTGGGCAAGAAGA
700 E A P P D N K K R K K *
2101 ACGAGGCGCCCCCGGACAACAAGAAGAGGAAGAAGTGACagctccggccctgatccagga
2161 ctgcacccccacccccaccgtccagccatccaacctcacttcgccttacaggtctccattt
2221 tgtggtaaaaaaagggttttaggccaggcgccgtggctcacgcctgtaatccaacatttg
2281 agaggctgaggcgggcggtacacctgagtcaggagttcgagaccagcctggccaacatgg
2341 tgaaacctccgtctctattaaaaatacaaaaattagccgagagtgggtggcatgcacctgt
2401 catcccagctactcgggaggtgaggcaggagaatcgcttgaacccgggaggcagaggtt
2461 gcagtgagccgagatcgcgccactgcactccaacctgggtgacagactctgtctccaaaa
2521 caaaacaaacaaacaaaaagattttattaaagatatatttgtaactcagtaaaaaaaaaa
2581 aaaaaaa
```

Figure 2E. The cDNA (SEQ ID. NO. : 10) and amino acid sequence (SEQ ID. NO. : 11) of 24P4C12 v.5.

The start methionine is underlined. The open reading frame extends from nucleic acid 6-2138 including the stop codon.

Figure 2E-1

```
1 M G G K Q R D E D D E A Y G K F V K Y
1 gagccATGGGGGAAAGCAGCGGGACGAGGATGACGAGGCCTACGGGAAGCCAGTCAAAT
20 D P S F R G P I K N R S C T D V I C C V
61 ACGACCCCTCCTTTCGAGGCCCCATCAAGAACAGAAGCTGCACAGATGTCATCTGCTGCG
40 L F L L F I L G Y I V V G I V A W L Y G
121 TCCTCTTCTGCTCTTTCATTCTAGGTTACATCGTGGTGGGGATTGTGGCCTGGTTGTATG
60 D P R Q V L Y P R N S T G A Y C G M G E
181 GAGACCCCGGCAAGTCCTCTACCCCGAAGTCTACTGGGGCCTACTGTGGCATGGGGG
80 N K D K P Y L L Y F N I F S C I L S S N
241 AGAACAAAGATAAGCCGTATCTCCTGTACTTCAACATCTTCAGCTGCATCCTGTCCAGCA
100 I I S V A E N G L Q C F T P Q V C V S S
301 ACATCATCTCAGTTGCTGAGAACGGCCTACAGTGCCCCACACCCAGGTGTGTGTGTCTCT
120 C P E D P W T V G K N E F S Q T V G E V
361 CCTGCCCCGAGGACCCATGGACTGTGGGAAAAAACGAGTTCTCACAGACTGTTGGGGAAG
140 F Y T K N R N F C L P G V P W N M T V I
```

1 μ M. Peptides are found that meet this binding criterion and qualify as HLA class II high affinity binders.

DR3 binding epitopes identified in this manner are included in vaccine compositions with DR supermotif-bearing peptide epitopes.

Similarly to the case of HLA class I motif-bearing peptides, the class II motif-bearing peptides are analogized to improve affinity or cross-reactivity. For example, aspartic acid at position 4 of the 9-mer core sequence is an optimal residue for DR3 binding, and substitution for that residue often improves DR 3 binding.

Example 17

Immunogenicity of 24P4C12-Derived HTL Epitopes

This example determines immunogenic DR supermotif- and DR3 motif-bearing epitopes among those identified using the methodology set forth herein.

Immunogenicity of HTL epitopes are confirmed in a manner analogous to the determination of immunogenicity of CTL epitopes, by assessing the ability to stimulate HTL responses and/or by using appropriate transgenic mouse models. Immunogenicity is determined by screening for: 1.) in vitro primary induction using normal PBMC or 2.) recall responses from patients who have 24P4C12-expressing tumors.

Example 18

Calculation of Phenotypic Frequencies of HLA-Supertypes in Various Ethnic Backgrounds to Determine Breadth of Population Coverage

This example illustrates the assessment of the breadth of population coverage of a vaccine composition comprised of multiple epitopes comprising multiple supermotifs and/or motifs.

In order to analyze population coverage, gene frequencies of HLA alleles are determined. Gene frequencies for each HLA allele are calculated from antigen or allele frequencies utilizing the binomial distribution formulae $gf=1-(\text{SQRT}(1-af))$ (see, e.g., Sidney et al., *Human Immunol.* 45:79-93, 1996). To obtain overall phenotypic frequencies, cumulative gene frequencies are calculated, and the cumulative antigen frequencies derived by the use of the inverse formula $[af=1-(1-Cgf)^2]$.

Where frequency data is not available at the level of DNA typing, correspondence to the serologically defined antigen frequencies is assumed. To obtain total potential supertype population coverage no linkage disequilibrium is assumed, and only alleles confirmed to belong to each of the superotypes are included (minimal estimates). Estimates of total potential coverage achieved by inter-loci combinations are made by adding to the A coverage the proportion of the non-A covered population that could be expected to be covered by the B alleles considered (e.g., $\text{total}=A+B*(1-A)$). Confirmed members of the A3-like supertype are A3, A11, A31, A*3301, and A*6801. Although the A3-like supertype may also include A34, A66, and A*7401, these alleles were not included in overall frequency calculations. Likewise, confirmed members of the A2-like supertype family are A*0201, A*0202, A*0203, A*0204, A*0205, A*0206, A*0207, A*6802, and A*6901. Finally, the B7-like supertype-confirmed alleles are: B7, B*3501-03, B51, B*5301, B*5401, B*5501-2, B*5601, B*6701, and B*7801 (potentially also B*1401, B*3504-06, B*4201, and B*5602).

Population coverage achieved by combining the A2-, A3- and B7-supertypes is approximately 86% in five major ethnic groups. Coverage may be extended by including peptides bearing the A1 and A24 motifs. On average, A1 is present in 12% and A24 in 29% of the population across five different major ethnic groups (Caucasian, North American Black, Chinese, Japanese, and Hispanic). Together, these alleles are represented with an average frequency of 39% in these same ethnic populations. The total coverage across the major ethnicities when A1 and A24 are combined with the coverage of the A2-, A3- and B7-supertype alleles is >95%, see, e.g., Table IV (G). An analogous approach can be used to estimate population coverage achieved with combinations of class II motif-bearing epitopes.

Immunogenicity studies in humans (e.g., Bertoni et al., *J. Clin. Invest.* 100:503, 1997; Doolan et al., *Immunity* 7:97, 1997; and Threlkeld et al., *J. Immunol.* 159:1648, 1997) have shown that highly cross-reactive binding peptides are almost always recognized as epitopes. The use of highly cross-reactive binding peptides is an important selection criterion in identifying candidate epitopes for inclusion in a vaccine that is immunogenic in a diverse population.

With a sufficient number of epitopes (as disclosed herein and from the art), an average population coverage is predicted to be greater than 95% in each of five major ethnic populations. The game theory Monte Carlo simulation analysis, which is known in the art (see e.g., Osborne, M. J. and Rubinstein, A. "A course in game theory" MIT Press, 1994), can be used to estimate what percentage of the individuals in a population comprised of the Caucasian, North American Black, Japanese, Chinese, and Hispanic ethnic groups would recognize the vaccine epitopes described herein. A preferred percentage is 90%. A more preferred percentage is 95%.

Example 19

CTL Recognition of Endogenously Processed Antigens After Priming

This example confirms that CTL induced by native or analogized peptide epitopes identified and selected as described herein recognize endogenously synthesized, i.e., native antigens.

Effector cells isolated from transgenic mice that are immunized with peptide epitopes, for example HLA-A2 supermotif-bearing epitopes, are re-stimulated in vitro using peptide-coated stimulator cells. Six days later, effector cells are assayed for cytotoxicity and the cell lines that contain peptide-specific cytotoxic activity are further re-stimulated. An additional six days later, these cell lines are tested for cytotoxic activity on ^{51}Cr labeled Jurkat-A2.1/ K^b target cells in the absence or presence of peptide, and also tested on ^{51}Cr labeled target cells bearing the endogenously synthesized antigen, i.e. cells that are stably transfected with 24P4C12 expression vectors.

The results demonstrate that CTL lines obtained from animals primed with peptide epitope recognize endogenously synthesized 24P4C12 antigen. The choice of transgenic mouse model to be used for such an analysis depends upon the epitope(s) that are being evaluated. In addition to HLA-A*0201/ K^b transgenic mice, several other transgenic mouse models including mice with human A11, which may also be used to evaluate A3 epitopes, and B7 alleles have been characterized and others (e.g., transgenic mice for HLA-A1 and

A24) are being developed. HLA-DR1 and HLA-DR3 mouse models have also been developed, which may be used to evaluate HTL epitopes.

Example 20

Activity of CTL-HTL Conjugated Epitopes in Transgenic Mice

This example illustrates the induction of CTLs and HTLs in transgenic mice, by use of a 24P4C12-derived CTL and HTL peptide vaccine compositions. The vaccine composition used herein comprise peptides to be administered to a patient with a 24P4C12-expressing tumor. The peptide composition can comprise multiple CTL and/or HTL epitopes. The epitopes are identified using methodology as described herein. This example also illustrates that enhanced immunogenicity can be achieved by inclusion of one or more HTL epitopes in a CTL vaccine composition; such a peptide composition can comprise an HTL epitope conjugated to a CTL epitope. The CTL epitope can be one that binds to multiple HLA family members at an affinity of 500 nM or less, or analogs of that epitope. The peptides may be lipidated, if desired.

Immunization procedures: Immunization of transgenic mice is performed as described (Alexander et al., *J. Immunol.* 159:4753-4761, 1997). For example, A2/K^b mice, which are transgenic for the human HLA A2.1 allele and are used to confirm the immunogenicity of HLA-A*0201 motif- or HLA-A2 supermotif-bearing epitopes, and are primed subcutaneously (base of the tail) with a 0.1 ml of peptide in Incomplete Freund's Adjuvant, or if the peptide composition is a lipidated CTL/HTL conjugate, in DMSO/saline, or if the peptide composition is a polypeptide, in PBS or Incomplete Freund's Adjuvant. Seven days after priming, splenocytes obtained from these animals are restimulated with syngenic irradiated LPS-activated lymphoblasts coated with peptide.

Cell lines: Target cells for peptide-specific cytotoxicity assays are Jurkat cells transfected with the HLA-A2.1/K^b chimeric gene (e.g., Vitiello et al., *J. Exp. Med.* 173:1007, 1991).

In vitro CTL activation: One week after priming, spleen cells (30×10⁶ cells/flask) are co-cultured at 37° C. with syngenic, irradiated (3000 rads), peptide coated lymphoblasts (10×10⁶ cells/flask) in 10 ml of culture medium/T25 flask. After six days, effector cells are harvested and assayed for cytotoxic activity.

Assay for cytotoxic activity: Target cells (1.0 to 1.5×10⁶) are incubated at 37° C. in the presence of 200 µl of ⁵¹Cr. After 60 minutes, cells are washed three times and resuspended in R10 medium. Peptide is added where required at a concentration of 1 µg/ml. For the assay, 10⁴ ⁵¹Cr-labeled target cells are added to different concentrations of effector cells (final volume of 200 µl) in U-bottom 96-well plates. After a six hour incubation period at 37° C., a 0.1 ml aliquot of supernatant is removed from each well and radioactivity is determined in a Micromedic automatic gamma counter. The percent specific lysis is determined by the formula: percent specific release=100×(experimental release–spontaneous release)/(maximum release–spontaneous release). To facilitate comparison between separate CTL assays run under the same conditions, % ⁵¹Cr release data is expressed as lytic units/10⁶ cells. One lytic unit is arbitrarily defined as the number of effector cells required to achieve 30% lysis of 10,000 target cells in a six hour ⁵¹Cr release assay. To obtain specific lytic units/10⁶, the lytic units/10⁶ obtained in the absence of peptide is subtracted from the lytic units/10⁶ obtained in the

presence of peptide. For example, if 30% ⁵¹Cr release is obtained at the effector (E):target (T) ratio of 50:1 (i.e., 5×10⁵ effector cells for 10,000 targets) in the absence of peptide and 5:1 (i.e., 5×10⁴ effector cells for 10,000 targets) in the presence of peptide, the specific lytic units would be: [(1/50,000)–(1/500,000)]×10⁶=18 LU.

The results are analyzed to assess the magnitude of the CTL responses of animals injected with the immunogenic CTL/HTL conjugate vaccine preparation and are compared to the magnitude of the CTL response achieved using, for example, CTL epitopes as outlined above in the Example entitled "Confirmation of Immunogenicity." Analyses similar to this may be performed to confirm the immunogenicity of peptide conjugates containing multiple CTL epitopes and/or multiple HTL epitopes. In accordance with these procedures, it is found that a CTL response is induced, and concomitantly that an HTL response is induced upon administration of such compositions.

Example 21

Selection of CTL and HTL Epitopes for Inclusion in a 24P4C12-Specific Vaccine

This example illustrates a procedure for selecting peptide epitopes for vaccine compositions of the invention. The peptides in the composition can be in the form of a nucleic acid sequence, either single or one or more sequences (i.e., minigene) that encodes peptide(s), or can be single and/or polypeptidic peptides.

The following principles are utilized when selecting a plurality of epitopes for inclusion in a vaccine composition. Each of the following principles is balanced in order to make the selection.

Epitopes are selected which, upon administration, mimic immune responses that are correlated with 24P4C12 clearance. The number of epitopes used depends on observations of patients who spontaneously clear 24P4C12. For example, if it has been observed that patients who spontaneously clear 24P4C12-expressing cells generate an immune response to at least three (3) epitopes from 24P4C12 antigen, then at least three epitopes should be included for HLA class I. A similar rationale is used to determine HLA class II epitopes.

Epitopes are often selected that have a binding affinity of an IC₅₀ of 500 nM or less for an HLA class I molecule, or for class II, an IC₅₀ of 1000 nM or less; or HLA Class I peptides with high binding scores from the BIMAS web site, at URL bimas.dcrt.nih.gov/.

In order to achieve broad coverage of the vaccine through out a diverse population, sufficient supermotif bearing peptides, or a sufficient array of allele-specific motif bearing peptides, are selected to give broad population coverage. In one embodiment, epitopes are selected to provide at least 80% population coverage. A Monte Carlo analysis, a statistical evaluation known in the art, can be employed to assess breadth, or redundancy, of population coverage.

When creating polyepitopic compositions, or a minigene that encodes same, it is typically desirable to generate the smallest peptide possible that encompasses the epitopes of interest. The principles employed are similar, if not the same, as those employed when selecting a peptide comprising nested epitopes. For example, a protein sequence for the vaccine composition is selected because it has maximal number of epitopes contained within the sequence, i.e., it has a high concentration of epitopes. Epitopes may be nested or overlapping (i.e., frame shifted relative to one another). For example, with overlapping epitopes, two 9-mer epitopes and

one 10-mer epitope can be present in a 10 amino acid peptide. Each epitope can be exposed and bound by an HLA molecule upon administration of such a peptide. A multi-epitopic, peptide can be generated synthetically, recombinantly, or via cleavage from the native source. Alternatively, an analog can be made of this native sequence, whereby one or more of the epitopes comprise substitutions that alter the cross-reactivity and/or binding affinity properties of the polypeptidic peptide. Such a vaccine composition is administered for therapeutic or prophylactic purposes. This embodiment provides for the possibility that an as yet undiscovered aspect of immune system processing will apply to the native nested sequence and thereby facilitate the production of therapeutic or prophylactic immune response-inducing vaccine compositions. Additionally such an embodiment provides for the possibility of motif-bearing epitopes for an HLA makeup that is presently unknown. Furthermore, this embodiment (absent the creating of any analogs) directs the immune response to multiple peptide sequences that are actually present in 24P4C12, thus avoiding the need to evaluate any junctional epitopes. Lastly, the embodiment provides an economy of scale when producing nucleic acid vaccine compositions. Related to this embodiment, computer programs can be derived in accordance with principles in the art, which identify in a target sequence, the greatest number of epitopes per sequence length.

A vaccine composition comprised of selected peptides, when administered, is safe, efficacious, and elicits an immune response similar in magnitude to an immune response that controls or clears cells that bear or overexpress 24P4C12.

Example 22

Construction of "Minigene" Multi-Epitope DNA Plasmids

This example discusses the construction of a minigene expression plasmid. Minigene plasmids may, of course, contain various configurations of B cell, CTL and/or HTL epitopes or epitope analogs as described herein.

A minigene expression plasmid typically includes multiple CTL and HTL peptide epitopes. In the present example, HLA-A2, -A3, -B7 supermotif-bearing peptide epitopes and HLA-A1 and -A24 motif-bearing peptide epitopes are used in conjunction with DR supermotif-bearing epitopes and/or DR3 epitopes. HLA class I supermotif or motif-bearing peptide epitopes derived 24P4C12, are selected such that multiple supermotifs/motifs are represented to ensure broad population coverage. Similarly, HLA class II epitopes are selected from 24P4C12 to provide broad population coverage, i.e. both HLA DR-1-4-7 supermotif-bearing epitopes and HLA DR-3 motif-bearing epitopes are selected for inclusion in the minigene construct. The selected CTL and HTL epitopes are then incorporated into a minigene for expression in an expression vector.

Such a construct may additionally include sequences that direct the HTL epitopes to the endoplasmic reticulum. For example, the Ii protein may be fused to one or more HTL epitopes as described in the art, wherein the CLIP sequence of the Ii protein is removed and replaced with an HLA class II epitope sequence so that HLA class II epitope is directed to the endoplasmic reticulum, where the epitope binds to an HLA class II molecules.

This example illustrates the methods to be used for construction of a minigene-bearing expression plasmid. Other expression vectors that may be used for minigene compositions are available and known to those of skill in the art.

The minigene DNA plasmid of this example contains a consensus Kozak sequence and a consensus murine kappa Ig-light chain signal sequence followed by CTL and/or HTL epitopes selected in accordance with principles disclosed herein. The sequence encodes an open reading frame fused to the Myc and H is antibody epitope tag coded for by the pcDNA 3.1 Myc-His vector.

Overlapping oligonucleotides that can, for example, average about 70 nucleotides in length with 15 nucleotide overlaps, are synthesized and HPLC-purified. The oligonucleotides encode the selected peptide epitopes as well as appropriate linker nucleotides, Kozak sequence, and signal sequence. The final multi-epitope minigene is assembled by extending the overlapping oligonucleotides in three sets of reactions using PCR. A Perkin/Elmer 9600 PCR machine is used and a total of 30 cycles are performed using the following conditions: 95° C. for 15 sec, annealing temperature (5° below the lowest calculated Tm of each primer pair) for 30 sec, and 72° C. for 1 min.

For example, a minigene is prepared as follows. For a first PCR reaction, 5 µg of each of two oligonucleotides are annealed and extended: In an example using eight oligonucleotides, i.e., four pairs of primers, oligonucleotides 1+2, 3+4, 5+6, and 7+8 are combined in 100 µl reactions containing Pfu polymerase buffer (1×=10 mM KCL, 10 mM (NH₄)₂SO₄, 20 mM Tris-chloride, pH 8.75, 2 mM MgSO₄, 0.1% Triton X-100, 100 µg/ml BSA), 0.25 mM each dNTP, and 2.5 U of Pfu polymerase. The full-length dimer products are gel-purified, and two reactions containing the product of 1+2 and 3+4, and the product of 5+6 and 7+8 are mixed, annealed, and extended for 10 cycles. Half of the two reactions are then mixed, and 5 cycles of annealing and extension carried out before flanking primers are added to amplify the full length product. The full-length product is gel-purified and cloned into pCR-blunt (Invitrogen) and individual clones are screened by sequencing.

Example 23

The Plasmid Construct and the Degree to which it Induces Immunogenicity

The degree to which a plasmid construct, for example a plasmid constructed in accordance with the previous Example, is able to induce immunogenicity is confirmed in vitro by determining epitope presentation by APC following transduction or transfection of the APC with an epitope-expressing nucleic acid construct. Such a study determines "antigenicity" and allows the use of human APC. The assay determines the ability of the epitope to be presented by the APC in a context that is recognized by a T cell by quantifying the density of epitope-HLA class I complexes on the cell surface. Quantitation can be performed by directly measuring the amount of peptide eluted from the APC (see, e.g., Sijts et al., *J. Immunol.* 156:683-692, 1996; Demotz et al., *Nature* 342:682-684, 1989); or the number of peptide-HLA class I complexes can be estimated by measuring the amount of lysis or lymphokine release induced by diseased or transfected target cells, and then determining the concentration of peptide necessary to obtain equivalent levels of lysis or lymphokine release (see, e.g., Kageyama et al., *J. Immunol.* 154:567-576, 1995).

Alternatively, immunogenicity is confirmed through in vivo injections into mice and subsequent in vitro assessment of CTL and HTL activity, which are analyzed using cytotoxicity and proliferation assays, respectively, as detailed e.g., in Alexander et al., *Immunity* 1:751-761, 1994.

For example, to confirm the capacity of a DNA minigene construct containing at least one HLA-A2 supermotif peptide to induce CTLs in vivo, HLA-A2.1/K^b transgenic mice, for example, are immunized intramuscularly with 100 µg of naked cDNA. As a means of comparing the level of CTLs induced by cDNA immunization, a control group of animals is also immunized with an actual peptide composition that comprises multiple epitopes synthesized as a single polypeptide as they would be encoded by the minigene.

Splenocytes from immunized animals are stimulated twice with each of the respective compositions (peptide epitopes encoded in the minigene or the polypeptidic peptide), then assayed for peptide-specific cytotoxic activity in a ⁵¹Cr release assay. The results indicate the magnitude of the CTL response directed against the A2-restricted epitope, thus indicating the in vivo immunogenicity of the minigene vaccine and polypeptidic vaccine.

It is, therefore, found that the minigene elicits immune responses directed toward the HLA-A2 supermotif peptide epitopes as does the polypeptidic peptide vaccine. A similar analysis is also performed using other HLA-A3 and HLA-B7 transgenic mouse models to assess CTL induction by HLA-A3 and HLA-B7 motif or supermotif epitopes, whereby it is also found that the minigene elicits appropriate immune responses directed toward the provided epitopes.

To confirm the capacity of a class II epitope-encoding minigene to induce HTLs in vivo, DR transgenic mice, or for those epitopes that cross react with the appropriate mouse MHC molecule, I-Ab-restricted mice, for example, are immunized intramuscularly with 100 µg of plasmid DNA. As a means of comparing the level of HTLs induced by DNA immunization, a group of control animals is also immunized with an actual peptide composition emulsified in complete Freund's adjuvant. CD4+ T cells, i.e. HTLs, are purified from splenocytes of immunized animals and stimulated with each of the respective compositions (peptides encoded in the minigene). The HTL response is measured using a ³H-thymidine incorporation proliferation assay, (see, e.g., Alexander et al. *Immunity* 1:751-761, 1994). The results indicate the magnitude of the HTL response, thus demonstrating the in vivo immunogenicity of the minigene.

DNA minigenes, constructed as described in the previous Example, can also be confirmed as a vaccine in combination with a boosting agent using a prime boost protocol. The boosting agent can consist of recombinant protein (e.g., Barnett et al., *Aids Res. and Human Retroviruses* 14, Supplement 3:S299-S309, 1998) or recombinant vaccinia, for example, expressing a minigene or DNA encoding the complete protein of interest (see, e.g., Hanke et al., *Vaccine* 16:439-445, 1998; Sedegah et al., *Proc. Natl. Acad. Sci. USA* 95:7648-53, 1998; Hanke and McMichael, *Immunol. Letters* 66:177-181, 1999; and Robinson et al., *Nature Med.* 5:526-34, 1999).

For example, the efficacy of the DNA minigene used in a prime boost protocol is initially evaluated in transgenic mice. In this example, A2.1/K^b transgenic mice are immunized IM with 100 µg of a DNA minigene encoding the immunogenic peptides including at least one HLA-A2 supermotif-bearing peptide. After an incubation period (ranging from 3-9 weeks), the mice are boosted IP with 10⁷ pfu/mouse of a recombinant vaccinia virus expressing the same sequence encoded by the DNA minigene. Control mice are immunized with 100 µg of DNA or recombinant vaccinia without the minigene sequence, or with DNA encoding the minigene, but without the vaccinia boost. After an additional incubation period of two weeks, splenocytes from the mice are immediately assayed for peptide-specific activity in an ELISPOT assay. Additionally, splenocytes are stimulated in vitro with the

A2-restricted peptide epitopes encoded in the minigene and recombinant vaccinia, then assayed for peptide-specific activity in an alpha, beta and/or gamma IFN ELISA.

It is found that the minigene utilized in a prime-boost protocol elicits greater immune responses toward the HLA-A2 supermotif peptides than with DNA alone. Such an analysis can also be performed using HLA-A11 or HLA-B7 transgenic mouse models to assess CTL induction by HLA-A3 or HLA-B7 motif or supermotif epitopes. The use of prime boost protocols in humans is described below in the Example entitled "Induction of CTL Responses Using a Prime Boost Protocol."

Example 24

Peptide Compositions for Prophylactic Uses

Vaccine compositions of the present invention can be used to prevent 24P4C12 expression in persons who are at risk for tumors that bear this antigen. For example, a polypeptidic peptide epitope composition (or a nucleic acid comprising the same) containing multiple CTL and HTL epitopes such as those selected in the above Examples, which are also selected to target greater than 80% of the population, is administered to individuals at risk for a 24P4C12-associated tumor.

For example, a peptide-based composition is provided as a single polypeptide that encompasses multiple epitopes. The vaccine is typically administered in a physiological solution that comprises an adjuvant, such as Incomplete Freund's Adjuvant. The dose of peptide for the initial immunization is from about 1 to about 50,000 g, generally 100-5,000 µg, for a 70 kg patient. The initial administration of vaccine is followed by booster dosages at 4 weeks followed by evaluation of the magnitude of the immune response in the patient, by techniques that determine the presence of epitope-specific CTL populations in a PBMC sample. Additional booster doses are administered as required. The composition is found to be both safe and efficacious as a prophylaxis against 24P4C12-associated disease.

Alternatively, a composition typically comprising transfecting agents is used for the administration of a nucleic acid-based vaccine in accordance with methodologies known in the art and disclosed herein.

Example 25

Polyepitopic Vaccine Compositions Derived from Native 24P4C12 Sequences

A native 24P4C12 polypeptide sequence is analyzed, preferably using computer algorithms defined for each class I and/or class II supermotif or motif, to identify "relatively short" regions of the polypeptide that comprise multiple epitopes. The "relatively short" regions are preferably less in length than an entire native antigen. This relatively short sequence that contains multiple distinct or overlapping, "nested" epitopes can be used to generate a minigene construct. The construct is engineered to express the peptide, which corresponds to the native protein sequence. The "relatively short" peptide is generally less than 250 amino acids in length, often less than 100 amino acids in length, preferably less than 75 amino acids in length, and more preferably less than 50 amino acids in length. The protein sequence of the vaccine composition is selected because it has maximal number of epitopes contained within the sequence, i.e., it has a high concentration of epitopes. As noted herein, epitope motifs may be nested or overlapping (i.e., frame shifted rela-

tive to one another). For example, with overlapping epitopes, two 9-mer epitopes and one 10-mer epitope can be present in a 10 amino acid peptide. Such a vaccine composition is administered for therapeutic or prophylactic purposes.

The vaccine composition will include, for example, multiple CTL epitopes from 24P4C12 antigen and at least one HTL epitope. This polyepitopic native sequence is administered either as a peptide or as a nucleic acid sequence which encodes the peptide. Alternatively, an analog can be made of this native sequence, whereby one or more of the epitopes comprise substitutions that alter the cross-reactivity and/or binding affinity properties of the polyepitopic peptide.

The embodiment of this example provides for the possibility that an as yet undiscovered aspect of immune system processing will apply to the native nested sequence and thereby facilitate the production of therapeutic or prophylactic immune response-inducing vaccine compositions. Additionally, such an embodiment provides for the possibility of motif-bearing epitopes for an HLA makeup(s) that is presently unknown. Furthermore, this embodiment (excluding an analogized embodiment) directs the immune response to multiple peptide sequences that are actually present in native 24P4C12, thus avoiding the need to evaluate any junctional epitopes. Lastly, the embodiment provides an economy of scale when producing peptide or nucleic acid vaccine compositions.

Related to this embodiment, computer programs are available in the art which can be used to identify in a target sequence, the greatest number of epitopes per sequence length.

Example 26

Polyepitopic Vaccine Compositions from Multiple Antigens

The 24P4C12 peptide epitopes of the present invention are used in conjunction with epitopes from other target tumor-associated antigens, to create a vaccine composition that is useful for the prevention or treatment of cancer that expresses 24P4C12 and such other antigens. For example, a vaccine composition can be provided as a single polypeptide that incorporates multiple epitopes from 24P4C12 as well as tumor-associated antigens that are often expressed with a target cancer associated with 24P4C12 expression, or can be administered as a composition comprising a cocktail of one or more discrete epitopes. Alternatively, the vaccine can be administered as a minigene construct or as dendritic cells which have been loaded with the peptide epitopes in vitro.

Example 27

Use of Peptides to Evaluate an Immune Response

Peptides of the invention may be used to analyze an immune response for the presence of specific antibodies, CTL or HTL directed to 24P4C12. Such an analysis can be performed in a manner described by Ogg et al., *Science* 279: 2103-2106, 1998. In this Example, peptides in accordance with the invention are used as a reagent for diagnostic or prognostic purposes, not as an immunogen.

In this example highly sensitive human leukocyte antigen tetrameric complexes ("tetramers") are used for a cross-sectional analysis of, for example, 24P4C12 HLA-A*0201-specific CTL frequencies from HLA A*0201-positive individuals at different stages of disease or following immunization comprising a 24P4C12 peptide containing an A*0201 motif.

Tetrameric complexes are synthesized as described (Musey et al., *N. Engl. J. Med.* 337:1267, 1997). Briefly, purified HLA heavy chain (A*0201 in this example) and β 2-microglobulin are synthesized by means of a prokaryotic expression system.

5 The heavy chain is modified by deletion of the transmembrane-cytosolic tail and COOH-terminal addition of a sequence containing a BirA enzymatic biotinylation site. The heavy chain, β 2-microglobulin, and peptide are refolded by dilution. The 45-kD refolded product is isolated by fast protein liquid chromatography and then biotinylated by BirA in the presence of biotin (Sigma, St. Louis, Mo.), adenosine 5' triphosphate and magnesium. Streptavidin-phycoerythrin conjugate is added in a 1:4 molar ratio, and the tetrameric product is concentrated to 1 mg/ml. The resulting product is referred to as tetramer-phycoerythrin.

For the analysis of patient blood samples, approximately one million PBMCs are centrifuged at 300 g for 5 minutes and resuspended in 50 μ l of cold phosphate-buffered saline. Tricolor analysis is performed with the tetramer-phycoerythrin, along with anti-CD8-Tricolor, and anti-CD38. The PBMCs are incubated with tetramer and antibodies on ice for 30 to 60 min and then washed twice before formaldehyde fixation. Gates are applied to contain >99.98% of control samples. Controls for the tetramers include both A*0201-negative individuals and A*0201-positive non-diseased donors. The percentage of cells stained with the tetramer is then determined by flow cytometry. The results indicate the number of cells in the PBMC sample that contain epitope-restricted CTLs, thereby readily indicating the extent of immune response to the 24P4C12 epitope, and thus the status of exposure to 24P4C12, or exposure to a vaccine that elicits a protective or therapeutic response.

Example 28

Use of Peptide Epitopes to Evaluate Recall Responses

The peptide epitopes of the invention are used as reagents to evaluate T cell responses, such as acute or recall responses, in patients. Such an analysis may be performed on patients who have recovered from 24P4C12-associated disease or who have been vaccinated with a 24P4C12 vaccine.

For example, the class I restricted CTL response of persons who have been vaccinated may be analyzed. The vaccine may be any 24P4C12 vaccine. PBMC are collected from vaccinated individuals and HLA typed. Appropriate peptide epitopes of the invention that, optimally, bear supermotifs to provide cross-reactivity with multiple HLA supertype family members, are then used for analysis of samples derived from individuals who bear that HLA type.

PBMC from vaccinated individuals are separated on Ficoll-Histopaque density gradients (Sigma Chemical Co., St. Louis, Mo.), washed three times in HBSS (GIBCO Laboratories), resuspended in RPMI-1640 (GIBCO Laboratories) supplemented with L-glutamine (2 mM), penicillin (50 U/ml), streptomycin (50 μ g/ml), and Hepes (10 mM) containing 10% heat-inactivated human AB serum (complete RPMI) and plated using microculture formats. A synthetic peptide comprising an epitope of the invention is added at 10 μ g/ml to each well and HBV core 128-140 epitope is added at 1 μ g/ml to each well as a source of T cell help during the first week of stimulation.

In the microculture format, 4×10^5 PBMC are stimulated with peptide in 8 replicate cultures in 96-well round bottom plate in 100 μ l/well of complete RPMI. On days 3 and 10, 100 μ l of complete RPMI and 20 U/ml final concentration of rIL-2

are added to each well. On day 7 the cultures are transferred into a 96-well flat-bottom plate and restimulated with peptide, rIL-2 and 10^5 irradiated (3,000 rad) autologous feeder cells. The cultures are tested for cytotoxic activity on day 14. A positive CTL response requires two or more of the eight replicate cultures to display greater than 10% specific ^{51}Cr release, based on comparison with non-diseased control subjects as previously described (Rehermann, et al., *Nature Med.* 2:1104, 1108, 1996; Rehermann et al., *J. Clin. Invest.* 97:1655-1665, 1996; and Rehermann et al. *J. Clin. Invest.* 98:1432-1440, 1996).

Target cell lines are autologous and allogeneic EBV-transformed B-LCL that are either purchased from the American Society for Histocompatibility and Immunogenetics (ASHI, Boston, Mass.) or established from the pool of patients as described (Guilhot, et al. *J. Virol.* 66:2670-2678, 1992).

Cytotoxicity assays are performed in the following manner. Target cells consist of either allogeneic HLA-matched or autologous EBV-transformed B lymphoblastoid cell line that are incubated overnight with the synthetic peptide epitope of the invention at 10 μM , and labeled with 100 μCi of ^{51}Cr (Amersham Corp., Arlington Heights, Ill.) for 1 hour after which they are washed four times with HBSS.

Cytolytic activity is determined in a standard 4-h, split well ^{51}Cr release assay using U-bottomed 96 well plates containing 3,000 targets/well. Stimulated PBMC are tested at effector/target (E/T) ratios of 20-50:1 on day 14. Percent cytotoxicity is determined from the formula: $100 \times [(\text{experimental release} - \text{spontaneous release}) / \text{maximum release} - \text{spontaneous release}]$. Maximum release is determined by lysis of targets by detergent (2% Triton X-100; Sigma Chemical Co., St. Louis, Mo.). Spontaneous release is <25% of maximum release for all experiments.

The results of such an analysis indicate the extent to which HLA-restricted CTL populations have been stimulated by previous exposure to 24P4C12 or a 24P4C12 vaccine.

Similarly, Class II restricted HTL responses may also be analyzed. Purified PBMC are cultured in a 96-well flat bottom plate at a density of 1.5×10^5 cells/well and are stimulated with 10 $\mu\text{g}/\text{ml}$ synthetic peptide of the invention, whole 24P4C12 antigen, or PHA. Cells are routinely plated in replicates of 4-6 wells for each condition. After seven days of culture, the medium is removed and replaced with fresh medium containing 10 U/ml IL-2. Two days later, 1 μCi ^3H -thymidine is added to each well and incubation is continued for an additional 18 hours. Cellular DNA is then harvested on glass fiber mats and analyzed for ^3H -thymidine incorporation. Antigen-specific T cell proliferation is calculated as the ratio of ^3H -thymidine incorporation in the presence of antigen divided by the ^3H -thymidine incorporation in the absence of antigen.

Example 29

Induction of Specific CTL Response in Humans

A human clinical trial for an immunogenic composition comprising CTL and HTL epitopes of the invention is set up as an IND Phase I, dose escalation study and carried out as a randomized, double-blind, placebo-controlled trial. Such a trial is designed, for example, as follows:

A total of about 27 individuals are enrolled and divided into 3 groups:

Group I: 3 subjects are injected with placebo and 6 subjects are injected with 5 μg of peptide composition;

Group II: 3 subjects are injected with placebo and 6 subjects are injected with 50 μg peptide composition;

Group III: 3 subjects are injected with placebo and 6 subjects are injected with 500 μg of peptide composition.

After 4 weeks following the first injection, all subjects receive a booster inoculation at the same dosage.

The endpoints measured in this study relate to the safety and tolerability of the peptide composition as well as its immunogenicity. Cellular immune responses to the peptide composition are an index of the intrinsic activity of this the peptide composition, and can therefore be viewed as a measure of biological efficacy. The following summarize the clinical and laboratory data that relate to safety and efficacy endpoints.

Safety: The incidence of adverse events is monitored in the placebo and drug treatment group and assessed in terms of degree and reversibility.

Evaluation of Vaccine Efficacy: For evaluation of vaccine efficacy, subjects are bled before and after injection. Peripheral blood mononuclear cells are isolated from fresh heparinized blood by Ficoll-Hypaque density gradient centrifugation, aliquoted in freezing media and stored frozen. Samples are assayed for CTL and HTL activity.

The vaccine is found to be both safe and efficacious.

Example 30

Phase II Trials in Patients Expressing 24P4C12

Phase II trials are performed to study the effect of administering the CTL-HTL peptide compositions to patients having cancer that expresses 24P4C12. The main objectives of the trial are to determine an effective dose and regimen for inducing CTLs in cancer patients that express 24P4C12, to establish the safety of inducing a CTL and HTL response in these patients, and to see to what extent activation of CTLs improves the clinical picture of these patients, as manifested, e.g., by the reduction and/or shrinking of lesions. Such a study is designed, for example, as follows:

The studies are performed in multiple centers. The trial design is an open-label, uncontrolled, dose escalation protocol wherein the peptide composition is administered as a single dose followed six weeks later by a single booster shot of the same dose. The dosages are 50, 500 and 5,000 micrograms per injection. Drug-associated adverse effects (severity and reversibility) are recorded.

There are three patient groupings. The first group is injected with 50 micrograms of the peptide composition and the second and third groups with 500 and 5,000 micrograms of peptide composition, respectively. The patients within each group range in age from 21-65 and represent diverse ethnic backgrounds. All of them have a tumor that expresses 24P4C12.

Clinical manifestations or antigen-specific T-cell responses are monitored to assess the effects of administering the peptide compositions. The vaccine composition is found to be both safe and efficacious in the treatment of 24P4C12-associated disease.

Example 31

Induction of CTL Responses Using a Prime Boost Protocol

A prime boost protocol similar in its underlying principle to that used to confirm the efficacy of a DNA vaccine in transgenic mice, such as described above in the Example entitled "The Plasmid Construct and the Degree to Which It Induces Immunogenicity," can also be used for the adminis-

tration of the vaccine to humans. Such a vaccine regimen can include an initial administration of, for example, naked DNA followed by a boost using recombinant virus encoding the vaccine, or recombinant protein/polypeptide or a peptide mixture administered in an adjuvant.

For example, the initial immunization may be performed using an expression vector, such as that constructed in the Example entitled "Construction of "Minigene" Multi-Epitope DNA Plasmids" in the form of naked nucleic acid administered IM (or SC or ID) in the amounts of 0.5-5 mg at multiple sites. The nucleic acid (0.1 to 1000 μ g) can also be administered using a gene gun. Following an incubation period of 3-4 weeks, a booster dose is then administered. The booster can be recombinant fowlpox virus administered at a dose of 5×10^7 to 5×10^9 pfu. An alternative recombinant virus, such as an MVA, canarypox, adenovirus, or adeno-associated virus, can also be used for the booster, or the polyepitopic protein or a mixture of the peptides can be administered. For evaluation of vaccine efficacy, patient blood samples are obtained before immunization as well as at intervals following administration of the initial vaccine and booster doses of the vaccine. Peripheral blood mononuclear cells are isolated from fresh heparinized blood by Ficoll-Hypaque density gradient centrifugation, aliquoted in freezing media and stored frozen. Samples are assayed for CTL and HTL activity.

Analysis of the results indicates that a magnitude of response sufficient to achieve a therapeutic or protective immunity against 24P4C12 is generated.

Example 32

Administration of Vaccine Compositions Using Dendritic Cells (DC)

Vaccines comprising peptide epitopes of the invention can be administered using APCs, or "professional" APCs such as DC. In this example, peptide-pulsed DC are administered to a patient to stimulate a CTL response in vivo. In this method, dendritic cells are isolated, expanded, and pulsed with a vaccine comprising peptide CTL and HTL epitopes of the invention. The dendritic cells are infused back into the patient to elicit CTL and HTL responses in vivo. The induced CTL and HTL then destroy or facilitate destruction, respectively, of the target cells that bear the 24P4C12 protein from which the epitopes in the vaccine are derived.

For example, a cocktail of epitope-comprising peptides is administered ex vivo to PBMC, or isolated DC therefrom. A pharmaceutical to facilitate harvesting of DC can be used, such as Progenipoiectin™ (Monsanto, St. Louis, Mo.) or GM-CSF/IL-4. After pulsing the DC with peptides, and prior to reinfusion into patients, the DC are washed to remove unbound peptides.

As appreciated clinically, and readily determined by one of skill based on clinical outcomes, the number of DC reinfused into the patient can vary (see, e.g., *Nature Med.* 4:328, 1998; *Nature Med.* 2:52, 1996 and *Prostate* 32:272, 1997). Although 2×10^6 DC per patient are typically administered, larger number of DC, such as 10^7 or 10^8 can also be provided. Such cell populations typically contain between 50-90% DC.

In some embodiments, peptide-loaded PBMC are injected into patients without purification of the DC. For example, PBMC generated after treatment with an agent such as Progenipoiectin™ are injected into patients without purification of the DC. The total number of PBMC that are administered often ranges from 10^8 to 10^{10} . Generally, the cell doses injected into patients is based on the percentage of DC in the

blood of each patient, as determined, for example, by immunofluorescence analysis with specific anti-DC antibodies. Thus, for example, if Progenipoiectin™ mobilizes 2% DC in the peripheral blood of a given patient, and that patient is to receive 5×10^6 DC, then the patient will be injected with a total of 2.5×10^8 peptide-loaded PBMC. The percent DC mobilized by an agent such as Progenipoiectin™ is typically estimated to be between 2-10%, but can vary as appreciated by one of skill in the art.

Ex Vivo Activation of CTL/HTL Responses

Alternatively, ex vivo CTL or HTL responses to 24P4C12 antigens can be induced by incubating, in tissue culture, the patient's, or genetically compatible, CTL or HTL precursor cells together with a source of APC, such as DC, and immunogenic peptides. After an appropriate incubation time (typically about 7-28 days), in which the precursor cells are activated and expanded into effector cells, the cells are infused into the patient, where they will destroy (CTL) or facilitate destruction (HTL) of their specific target cells, i.e., tumor cells.

Example 33

An Alternative Method of Identifying and Confirming Motif-Bearing Peptides

Another method of identifying and confirming motif-bearing peptides is to elute them from cells bearing defined MHC molecules. For example, EBV transformed B cell lines used for tissue typing have been extensively characterized to determine which HLA molecules they express. In certain cases these cells express only a single type of HLA molecule. These cells can be transfected with nucleic acids that express the antigen of interest, e.g. 24P4C12. Peptides produced by endogenous antigen processing of peptides produced as a result of transfection will then bind to HLA molecules within the cell and be transported and displayed on the cell's surface. Peptides are then eluted from the HLA molecules by exposure to mild acid conditions and their amino acid sequence determined, e.g., by mass spectral analysis (e.g., Kubo et al., *J. Immunol.* 152:3913, 1994). Because the majority of peptides that bind a particular HLA molecule are motif-bearing, this is an alternative modality for obtaining the motif-bearing peptides correlated with the particular HLA molecule expressed on the cell.

Alternatively, cell lines that do not express endogenous HLA molecules can be transfected with an expression construct encoding a single HLA allele. These cells can then be used as described, i.e., they can then be transfected with nucleic acids that encode 24P4C12 to isolate peptides corresponding to 24P4C12 that have been presented on the cell surface. Peptides obtained from such an analysis will bear motif(s) that correspond to binding to the single HLA allele that is expressed in the cell.

As appreciated by one in the art, one can perform a similar analysis on a cell bearing more than one HLA allele and subsequently determine peptides specific for each HLA allele expressed. Moreover, one of skill would also recognize that means other than transfection, such as loading with a protein antigen, can be used to provide a source of antigen to the cell.

Example 34

Complementary Polynucleotides

Sequences complementary to the 24P4C12-encoding sequences, or any parts thereof, are used to detect, decrease,

or inhibit expression of naturally occurring 24P4C12. Although use of oligonucleotides comprising from about 15 to 30 base pairs is described, essentially the same procedure is used with smaller or with larger sequence fragments. Appropriate oligonucleotides are designed using, e.g., OLIGO 4.06 software (National Biosciences) and the coding sequence of 24P4C12. To inhibit transcription, a complementary oligonucleotide is designed from the most unique 5' sequence and used to prevent promoter binding to the coding sequence. To inhibit translation, a complementary oligonucleotide is designed to prevent ribosomal binding to a 24P4C12-encoding transcript.

Example 35

Purification of Naturally-occurring or Recombinant 24P4C12 Using 24P4C12-Specific Antibodies

Naturally occurring or recombinant 24P4C12 is substantially purified by immunoaffinity chromatography using antibodies specific for 24P4C12. An immunoaffinity column is constructed by covalently coupling anti-24P4C12 antibody to an activated chromatographic resin, such as CNBr-activated SEPHAROSE (Amersham Pharmacia Biotech). After the coupling, the resin is blocked and washed according to the manufacturer's instructions.

Media containing 24P4C12 are passed over the immunoaffinity column, and the column is washed under conditions that allow the preferential absorbance of 24P4C12 (e.g., high ionic strength buffers in the presence of detergent). The column is eluted under conditions that disrupt antibody/24P4C12 binding (e.g., a buffer of pH 2 to pH 3, or a high concentration of a chaotrope, such as urea or thiocyanate ion), and GCR.P is collected.

Example 36

Identification of Molecules which Interact with 24P4C12

24P4C12, or biologically active fragments thereof, are labeled with 121 I Bolton-Hunter reagent. (See, e.g., Bolton et al. (1973) *Biochem. J.* 133:529.) Candidate molecules previously arrayed in the wells of a multi-well plate are incubated with the labeled 24P4C12, washed, and any wells with labeled 24P4C12 complex are assayed. Data obtained using different concentrations of 24P4C12 are used to calculate values for the number, affinity, and association of 24P4C12 with the candidate molecules.

Example 37

In Vivo Assay for 24P4C12 Tumor Growth Promotion

The effect of the 24P4C12 protein on tumor cell growth is evaluated in vivo by evaluating tumor development and growth of cells expressing or lacking 24P4C12. For example, SCID mice are injected subcutaneously on each flank with 1×10^6 of either 3T3, prostate, colon, ovary, lung, or bladder cancer cell lines (e.g. PC3, Caco, PA-1, CaLu or J82 cells) containing tkNeo empty vector or 24P4C12. At least two strategies may be used: (1) Constitutive 24P4C12 expression under regulation of a promoter, such as a constitutive promoter obtained from the genomes of viruses such as polyoma virus, fowlpox virus (UK 2,211,504 published 5 Jul. 1989), adenovirus (such as Adenovirus 2), bovine papilloma virus,

avian sarcoma virus, cytomegalovirus, a retrovirus, hepatitis-B virus and Simian Virus 40 (SV40), or from heterologous mammalian promoters, e.g., the actin promoter or an immunoglobulin promoter, provided such promoters are compatible with the host cell systems, and (2) Regulated expression under control of an inducible vector system, such as ecdysone, tetracycline, etc., provided such promoters are compatible with the host cell systems. Tumor volume is then monitored by caliper measurement at the appearance of palpable tumors and followed over time to determine if 24P4C12-expressing cells grow at a faster rate and whether tumors produced by 24P4C12-expressing cells demonstrate characteristics of altered aggressiveness (e.g. enhanced metastasis, vascularization, reduced responsiveness to chemotherapeutic drugs). As shown in FIG. 31 and FIG. 32, 24P4C12 has a profound effect on tumor growth in SCID mice. The prostate cancer cells PC3 and PC3-24P4C12 were injected subcutaneously in the right flank of SCID mice. Tumor growth was evaluated by caliper measurements. An increase in tumor growth was observed in PC3-24P4C12 tumors within 47 days of injection (FIG. 31). In addition, subcutaneous injection of 3T3-24P4C12 induced tumor formation in SCID mice (FIG. 32). This finding is significant as control 3T3 cells fail to form tumors, indicating that 24P4C12 has several tumor enhancing capabilities, including transformation, as well as tumor initiation and promotion.

Example 38

24P4C12 Monoclonal Antibody-mediated Inhibition of Prostate Tumors In Vivo

The significant expression of 24P4C12 in cancer tissues, together with its restrictive expression in normal tissues and cell surface localization, make 24P4C12 a good target for antibody therapy. Similarly, 24P4C12 is a target for T cell-based immunotherapy. Thus, the therapeutic efficacy of anti-24P4C12 mAbs in human prostate cancer xenograft mouse models is evaluated by using recombinant cell lines such as PC3-24P4C12, and 3T3-24P4C12 (see, e.g., Kaighn, M. E., et al., *Invest Urol*, 1979, 17(1): p. 16-23), as well as human prostate xenograft models such as LAPC9 (Saffran et al., *Proc Natl Acad Sci USA*, 2001, 98:2658). Similarly, anti-24P4C12 mAbs are evaluated in xenograft models of human bladder cancer, colon cancer, ovarian cancer or lung cancer using recombinant cell lines such as J82-24P4C12, Caco-24P4C12, PA-24P4C1 or CaLu-24P4C12, respectively.

Antibody efficacy on tumor growth and metastasis formation is studied, e.g., in a mouse orthotopic bladder cancer xenograft model, and a mouse prostate cancer xenograft model. The antibodies can be unconjugated, as discussed in this Example, or can be conjugated to a therapeutic modality, as appreciated in the art. Anti-24P4C12 mAbs inhibit formation of prostate and bladder xenografts. Anti-24P4C12 mAbs also retard the growth of established orthotopic tumors and prolonged survival of tumor-bearing mice. These results indicate the utility of anti-24P4C12 mAbs in the treatment of local and advanced stages of prostate, colon, ovarian, lung and bladder cancer. (See, e.g., Saffran, D., et al., *PNAS* 10:1073-1078 or www.pnas.org/cgi/doi/10.1073/pnas.051624698).

Administration of the anti-24P4C12 mAbs led to retardation of established orthotopic tumor growth and inhibition of metastasis to distant sites, resulting in a significant prolongation in the survival of tumor-bearing mice. These studies indicate that 24P4C12 as an attractive target for immunotherapy and demonstrate the therapeutic potential of anti-

24P4C12 mAbs for the treatment of local and metastatic cancer. This example demonstrates that unconjugated 24P4C12 monoclonal antibodies are effective to inhibit the growth of human prostate, colon, ovarian, lung and bladder cancer tumor xenografts grown in SCID mice; accordingly a combination of such efficacious monoclonal antibodies is also effective.

Tumor Inhibition using Multiple Unconjugated 24P4C12 mAbs

Materials and Methods

24P4C12 Monoclonal Antibodies:

Monoclonal antibodies are raised against 24P4C12 as described in the Example entitled "Generation of 24P4C12 Monoclonal Antibodies (mAbs)." The antibodies are characterized by ELISA, Western blot, FACS, and immunoprecipitation for their capacity to bind 24P4C12. Epitope mapping data for the anti-24P4C12 mAbs, as determined by ELISA and Western analysis, recognize epitopes on the 24P4C12 protein. Immunohistochemical analysis of prostate cancer tissues and cells with these antibodies is performed.

The monoclonal antibodies are purified from ascites or hybridoma tissue culture supernatants by Protein-G Sepharose chromatography, dialyzed against PBS, filter sterilized, and stored at -20°C . Protein determinations are performed by a Bradford assay (Bio-Rad, Hercules, Calif.). A therapeutic monoclonal antibody or a cocktail comprising a mixture of individual monoclonal antibodies is prepared and used for the treatment of mice receiving subcutaneous or orthotopic injections of SCABER, J82, A498, 769P, CaOv1 or PA1 tumor xenografts.

Cell Lines

The prostate, colon, ovarian, lung and bladder cancer carcinoma cell lines, Caco, PA-1, CaLu or J82 cells as well as the fibroblast line NIH 3T3 (American Type Culture Collection) are maintained in media supplemented with L-glutamine and 10% FBS.

PC3-24P4C12, Caco-24P4C12, PA-24P4C12, CaLu-24P4C12 or J82-24P4C12 cells and 3T3-24P4C12 cell populations are generated by retroviral gene transfer as described in Hubert, R. S., et al., *Proc Natl Acad Sci USA*, 1999. 96(25): 14523.

Xenograft Mouse Models.

Subcutaneous (s.c.) tumors are generated by injection of 1×10^6 cancer cells mixed at a 1:1 dilution with Matrigel (Collaborative Research) in the right flank of male SCID mice. To test antibody efficacy on tumor formation, i.p. antibody injections are started on the same day as tumor-cell injections. As a control, mice are injected with either purified mouse IgG (ICN) or PBS; or a purified monoclonal antibody that recognizes an irrelevant antigen not expressed in human cells. Tumor sizes are determined by caliper measurements, and the tumor volume is calculated as: Length $\times$ Width $\times$ Height. Mice with s.c. tumors greater than 1.5 cm in diameter are sacrificed.

Orthotopic injections are performed under anesthesia by using ketamine/xylazine. For bladder orthotopic studies, an incision is made through the abdomen to expose the bladder, and tumor cells (5×10^5) mixed with Matrigel are injected into the bladder wall in a 10- μl volume. To monitor tumor growth, mice are palpated and blood is collected on a weekly basis to measure BTA levels. For prostate orthotopic models, an incision is made through the abdominal muscles to expose the bladder and seminal vesicles, which then are delivered through the incision to expose the dorsal prostate. Tumor cells e.g. LAPC-9 cells (5×10^5) mixed with Matrigel are injected into the prostate in a 10- μl volume (Yoshida Y et al., *Anticancer Res.* 1998, 18:327; Ahn et al., *Tumour Biol.* 2001,

22:146). To monitor tumor growth, blood is collected on a weekly basis measuring PSA levels. Similar procedures are followed for lung and ovarian xenograft models. The mice are segregated into groups for the appropriate treatments, with anti-24P4C12 or control mAbs being injected i.p.

Anti-24P4C12 mAbs Inhibit Growth of 24P4C12-Expressing Xenograft-Cancer Tumors

The effect of anti-24P4C12 mAbs on tumor formation is tested on the growth and progression of bladder, and prostate cancer xenografts using PC3-24P4C12, Caco-24P4C12, PA-24P4C12, CaLu-24P4C12 or J82-24P4C12 orthotopic models. As compared with the s.c. tumor model, the orthotopic model, which requires injection of tumor cells directly in the mouse prostate, colon, ovary, lung and bladder, respectively, results in a local tumor growth, development of metastasis in distal sites, deterioration of mouse health, and subsequent death (Saffran, D., et al., *PNAS supra*; Fu, X., et al., *Int J Cancer*, 1992. 52(6): p. 987-90; Kubota, T., *J Cell Biochem*, 1994. 56(1): p. 4-8). The features make the orthotopic model more representative of human disease progression and allowed us to follow the therapeutic effect of mAbs on clinically relevant end points.

Accordingly, tumor cells are injected into the mouse organs, and 2 days later, the mice are segregated into two groups and treated with either: a) 200-500 μg , of anti-24P4C12 Ab, or b) PBS three times per week for two to five weeks.

A major advantage of the orthotopic cancer models is the ability to study the development of metastases. Formation of metastasis in mice bearing established orthotopic tumors is studies by IHC analysis on lung sections using an antibody against a tumor-specific cell-surface protein such as anti-CK20 for bladder cancer, anti-STEAP-1 for prostate cancer models (Lin S et al, *Cancer Detect Prev.* 2001; 25:202; Saffran, D., et al., *PNAS supra*).

Mice bearing established orthotopic tumors are administered 1000 μg injections of either anti-24P4C12 mAb or PBS over a 4-week period. Mice in both groups are allowed to establish a high tumor burden, to ensure a high frequency of metastasis formation in mouse lungs. Mice then are killed and their bladders, livers, bone and lungs are analyzed for the presence of tumor cells by IHC analysis.

These studies demonstrate a broad anti-tumor efficacy of anti-24P4C12 antibodies on initiation and progression of prostate and kidney cancer in xenograft mouse models. Anti-24P4C12 antibodies inhibit tumor formation of tumors as well as retarding the growth of already established tumors and prolong the survival of treated mice. Moreover, anti-24P4C12 mAbs demonstrate a dramatic inhibitory effect on the spread of local bladder and prostate tumor to distal sites, even in the presence of a large tumor burden. Thus, anti-24P4C12 mAbs are efficacious on major clinically relevant end points (tumor growth), prolongation of survival, and health.

Example 39

Therapeutic and Diagnostic use of Anti-24P4C12 Antibodies in Humans

Anti-24P4C12 monoclonal antibodies are safely and effectively used for diagnostic, prophylactic, prognostic and/or therapeutic purposes in humans. Western blot and immunohistochemical analysis of cancer tissues and cancer xenografts with anti-24P4C12 mAb show strong extensive staining in carcinoma but significantly lower or undetectable levels in normal tissues. Detection of 24P4C12 in carcinoma and in metastatic disease demonstrates the usefulness of the

mAb as a diagnostic and/or prognostic indicator. Anti-24P4C12 antibodies are therefore used in diagnostic applications such as immunohistochemistry of kidney biopsy specimens to detect cancer from suspect patients.

As determined by flow cytometry, anti-24P4C12 mAb specifically binds to carcinoma cells. Thus, anti-24P4C12 antibodies are used in diagnostic whole body imaging applications, such as radioimmunoscintigraphy and radioimmunotherapy, (see, e.g., Potamianos S., et. al. *Anti-cancer Res* 20(2A):925-948 (2000)) for the detection of localized and metastatic cancers that exhibit expression of 24P4C12. Shedding or release of an extracellular domain of 24P4C12 into the extracellular milieu, such as that seen for alkaline phosphodiesterase B10 (Meerson, N. R., *Hepatology* 27:563-568 (1998)), allows diagnostic detection of 24P4C12 by anti-24P4C12 antibodies in serum and/or urine samples from suspect patients.

Anti-24P4C12 antibodies that specifically bind 24P4C12 are used in therapeutic applications for the treatment of cancers that express 24P4C12. Anti-24P4C12 antibodies are used as an unconjugated modality and as conjugated form in which the antibodies are attached to one of various therapeutic or imaging modalities well known in the art, such as a prodrugs, enzymes or radioisotopes. In preclinical studies, unconjugated and conjugated anti-24P4C12 antibodies are tested for efficacy of tumor prevention and growth inhibition in the SCID mouse cancer xenograft models, e.g., kidney cancer models AGS-K3 and AGS-K6, (see, e.g., the Example entitled "24P4C12 Monoclonal Antibody-mediated Inhibition of Bladder and Lung Tumors In Vivo"). Either conjugated and unconjugated anti-24P4C12 antibodies are used as a therapeutic modality in human clinical trials either alone or in combination with other treatments as described in following Examples.

Example 40

Human Clinical Trials for the Treatment and Diagnosis of Human Carcinomas through use of Human Anti-24P4C12 Antibodies In Vivo

Antibodies are used in accordance with the present invention which recognize an epitope on 24P4C12, and are used in the treatment of certain tumors such as those listed in Table I. Based upon a number of factors, including 24P4C12 expression levels, tumors such as those listed in Table I are presently preferred indications. In connection with each of these indications, three clinical approaches are successfully pursued.

I.) Adjunctive therapy: In adjunctive therapy, patients are treated with anti-24P4C12 antibodies in combination with a chemotherapeutic or antineoplastic agent and/or radiation therapy. Primary cancer targets, such as those listed in Table I, are treated under standard protocols by the addition anti-24P4C12 antibodies to standard first and second line therapy. Protocol designs address effectiveness as assessed by reduction in tumor mass as well as the ability to reduce usual doses of standard chemotherapy. These dosage reductions allow additional and/or prolonged therapy by reducing dose-related toxicity of the chemotherapeutic agent. Anti-24P4C12 antibodies are utilized in several adjunctive clinical trials in combination with the chemotherapeutic or antineoplastic agents adriamycin (advanced prostrate carcinoma), cisplatin (advanced head and neck and lung carcinomas), taxol (breast cancer), and doxorubicin (preclinical).

II.) Monotherapy: In connection with the use of the anti-24P4C12 antibodies in monotherapy of tumors, the antibodies are administered to patients without a chemotherapeutic

or antineoplastic agent. In one embodiment, monotherapy is conducted clinically in end stage cancer patients with extensive metastatic disease. Patients show some disease stabilization. Trials demonstrate an effect in refractory patients with cancerous tumors.

III.) Imaging Agent: Through binding a radionuclide (e.g., iodine or yttrium (I^{131} , Y^{90}) to anti-24P4C12 antibodies, the radiolabeled antibodies are utilized as a diagnostic and/or imaging agent. In such a role, the labeled antibodies localize to both solid tumors, as well as, metastatic lesions of cells expressing 24P4C12. In connection with the use of the anti-24P4C12 antibodies as imaging agents, the antibodies are used as an adjunct to surgical treatment of solid tumors, as both a pre-surgical screen as well as a post-operative follow-up to determine what tumor remains and/or returns. In one embodiment, a (^{111}In)-24P4C12 antibody is used as an imaging agent in a Phase I human clinical trial in patients having a carcinoma that expresses 24P4C12 (by analogy see, e.g., Divgi et al. *J. Natl. Cancer Inst.* 83:97-104 (1991)). Patients are followed with standard anterior and posterior gamma camera. The results indicate that primary lesions and metastatic lesions are identified.

Dose and Route of Administration

As appreciated by those of ordinary skill in the art, dosing considerations can be determined through comparison with the analogous products that are in the clinic. Thus, anti-24P4C12 antibodies can be administered with doses in the range of 5 to 400 mg/m², with the lower doses used, e.g., in connection with safety studies. The affinity of anti-24P4C12 antibodies relative to the affinity of a known antibody for its target is one parameter used by those of skill in the art for determining analogous dose regimens. Further, anti-24P4C12 antibodies that are fully human antibodies, as compared to the chimeric antibody, have slower clearance; accordingly, dosing in patients with such fully human anti-24P4C12 antibodies can be lower, perhaps in the range of 50 to 300 mg/m², and still remain efficacious. Dosing in mg/m², as opposed to the conventional measurement of dose in mg/kg, is a measurement based on surface area and is a convenient dosing measurement that is designed to include patients of all sizes from infants to adults.

Three distinct delivery approaches are useful for delivery of anti-24P4C12 antibodies. Conventional intravenous delivery is one standard delivery technique for many tumors. However, in connection with tumors in the peritoneal cavity, such as tumors of the ovaries, biliary duct, other ducts, and the like, intraperitoneal administration may prove favorable for obtaining high dose of antibody at the tumor and to also minimize antibody clearance. In a similar manner, certain solid tumors possess vasculature that is appropriate for regional perfusion. Regional perfusion allows for a high dose of antibody at the site of a tumor and minimizes short term clearance of the antibody.

Clinical Development Plan (CDP)

Overview: The CDP follows and develops treatments of anti-24P4C12 antibodies in connection with adjunctive therapy, monotherapy, and as an imaging agent. Trials initially demonstrate safety and thereafter confirm efficacy in repeat doses. Trials are open label comparing standard chemotherapy with standard therapy plus anti-24P4C12 antibodies. As will be appreciated, one criteria that can be utilized in connection with enrollment of patients is 24P4C12 expression levels in their tumors as determined by biopsy.

As with any protein or antibody infusion-based therapeutic, safety concerns are related primarily to (i) cytokine release syndrome, i.e., hypotension, fever, shaking, chills; (ii) the development of an immunogenic response to the material

Figure 2E-2

421 TCTTCTATACAAAAACAGGAACCTTTGTCTGCCAGGGGTACCTTGAATATGACGGTGA
160 T S L Q Q E L C P S F L L P S A P A L G
481 TCACAAGCCTGCAACAGGAACCTCTGCCCCAGTTTCCTCCTCCCTCTGCTCCAGCTCTGG
180 R C F P W T N V T P P A L P G I T N D T
541 GGCGTCTGCTTTCCATGGACCAACGTTACTCCACCGGCGCTCCCAGGGATCACCAATGACA
200 I I Q Q G I S G L I D S L N A R D I S V
601 CCACCATACAGCAGGGGATCAGCGGTCTTATTGACAGCCTCAATGCCCCGAGACATCAGTG
220 K I F E D F A Q S W Y W I L V A L G V A
661 TTAAGATCTTTGAAGATTTTGCCAGTCCTGGTATTGGATTCTTGTGGCCTGGGGGTGG
240 L V L S L L F I L L L R L V A G P L V L
721 CTCTGGTCTTGAGCCTACTGTTTATCTTGCTTCTGCGCCTGGTGGCTGGGCCCCCTGGTGC
260 V L I L G V L G V L A Y G I Y Y C W E E
781 TGGTCTGATCCTGGGAGTCTGGGCGTCTGGCATAACGGCATCTACTACTGCTGGGAGG
280 Y R V L R D K G A S I S Q L G F T I N L
841 AGTACCGAGTCTGCGGGACAAGGCGCCTCCATCTCCCAGCTGGGTTTCACCACCAACC
300 S A Y Q S V Q E T W L A A L I V L A V L
901 TCAGTGCCTACCAGAGCGTGCAGGAGACCTGGCTGGCCGCCCTGATCGTGTGGCGGTGC
320 E A I L L L V L I F L R Q R I R I A I A
961 TTGAAGCCATCCTGCTGCTGGTGTCTATCTTCTGCGGCAGCGGATTCTGATTGCCATCG
340 L L K E A S K A V G Q M M S T M F Y P L
1021 CCCTCCTGAAGGAGGCCAGCAAGGCTGTGGGACAGATGATGTCTACCATGTTCTACCCAC
360 V T F V L L L I C I A Y W A M T A L Y L
1081 TGGTACCTTTGTCTCCTCCTCATCTGCATTGCCTACTGGGCCATGACTGCTCTGTACC
380 A T S G Q P Q Y V L W A S N I S S P G C
1141 TGGCTACATCGGGGCAACCCCAGTATGTGCTCTGGGCATCCAACATCAGCTCCCCCGGT
400 E K V P I N T S C N P T A H L V N S S C
1201 GTGAGAAAGTGCCAATAAATACATCATGCAACCCACGGCCACCTTGTGAACTCCTCGT
420 P G L M C V F Q G Y S S K G L I Q R S V
1261 GCCCAGGGCTGATGTGCGTCTTCCAGGGCTACTCATCCAAGGCCTAATCCAACGTTCTG
440 F N L Q I Y G V L G L F W I L N W V L A
1321 TCTTCAATCTGCAAATCTATGGGGTCTTCTGGGGCTCTTCTGGACCCTTAAGTGGTACTGG
460 L G Q C V L A G A F A S F Y W A F H K P
1381 CCCTGGGCCAATGCGTCTCGCTGGAGCCTTTGCCTCCTTCTACTGGGCCTTCCACAAGC
480 Q D I P T F P L I S A F I R T L R Y R T
1441 CCCAGGACATCCCTACCTTCCCCTTAATCTCTGCCTTCATCCGCACACTCCGTTACCACA
500 G S L A F G A L I L T L V Q I A R V I L
1501 CTGGGTCAATTGGCATTGGAGCCCTCATCTGACCCTTGTGAGATAGCCCGGTCTATCT
520 E Y I D H K L R G V Q N P V A R C I M C
1561 TGGAGTATATTGACCACAAGCTCAGAGGAGTGCAGAACCTGTAGCCCGCTGCATCATGT
540 C F R C C L W C L E K F I R F L N R N A

139

(i.e., development of human antibodies by the patient to the antibody therapeutic, or HAHA response); and, (iii) toxicity to normal cells that express 24P4C12. Standard tests and follow-up are utilized to monitor each of these safety concerns. Anti-24P4C12 antibodies are found to be safe upon human administration.

Example 41

Human Clinical Trial Adjunctive Therapy with
Human Anti-24P4C12 Antibody and
Chemotherapeutic Agent

A phase I human clinical trial is initiated to assess the safety of six intravenous doses of a human anti-24P4C12 antibody in connection with the treatment of a solid tumor, e.g., a cancer of a tissue listed in Table I. In the study, the safety of single doses of anti-24P4C12 antibodies when utilized as an adjunctive therapy to an antineoplastic or chemotherapeutic agent as defined herein, such as, without limitation: cisplatin, topotecan, doxorubicin, adriamycin, taxol, or the like, is assessed. The trial design includes delivery of six single doses of an anti-24P4C12 antibody with dosage of antibody escalating from approximately about 25 mg/m² to about 275 mg/m² over the course of the treatment in accordance with the following schedule:

	Day 0	Day 7	Day 14	Day 21	Day 28	Day 35
mAb Dose	25 mg/m ²	75 mg/m ²	125 mg/m ²	175 mg/m ²	225 mg/m ²	275 mg/m ²
Chemotherapy (standard dose)	+	+	+	+	+	+

Patients are closely followed for one-week following each administration of antibody and chemotherapy. In particular, patients are assessed for the safety concerns mentioned above: (i) cytokine release syndrome, i.e., hypotension, fever, shaking, chills; (ii) the development of an immunogenic response to the material (i.e., development of human antibodies by the patient to the human antibody therapeutic, or HAHA response); and, (iii) toxicity to normal cells that express 24P4C12. Standard tests and follow-up are utilized to monitor each of these safety concerns. Patients are also assessed for clinical outcome, and particularly reduction in tumor mass as evidenced by MRI or other imaging.

The anti-24P4C12 antibodies are demonstrated to be safe and efficacious, Phase II trials confirm the efficacy and refine optimum dosing.

Example 42

Human Clinical Trial: Monotherapy with Human
Anti-24P4C12 Antibody

Anti-24P4C12 antibodies are safe in connection with the above-discussed adjunctive trial, a Phase II human clinical trial confirms the efficacy and optimum dosing for monotherapy. Such trial is accomplished, and entails the same safety and outcome analyses, to the above-described adjunctive trial with the exception being that patients do not receive chemotherapy concurrently with the receipt of doses of anti-24P4C12 antibodies.

140

Example 43

Human Clinical Trial: Diagnostic Imaging with
Anti-24P4C12 Antibody

Once again, as the adjunctive therapy discussed above is safe within the safety criteria discussed above, a human clinical trial is conducted concerning the use of anti-24P4C12 antibodies as a diagnostic imaging agent. The protocol is designed in a substantially similar manner to those described in the art, such as in Divgi et al. *J. Natl. Cancer Inst.* 83:97-104 (1991). The antibodies are found to be both safe and efficacious when used as a diagnostic modality.

Example 44

Homology Comparison of 24P4C12 to Known
Sequences

The 24P4C12 protein of FIG. 3 has 710 amino acids with calculated molecular weight of 79.3 kDa, and pl of 8.9. Several variants of 24P4C12 have been identified, including 4 SNPs (namely v.1, v.3, v.5, v.6) and 3 splice variants (namely v.7, v.8 and v.9) (FIGS. 10 and 11). 24P4C12 variants v.3, v.5, and v.6 differ from 24P4C12 v.1 by 1 amino acid each, at aa positions 187, 326 and 436, respectively. Variant v.7 carries a

deletion of 111 aa long starting at aa 237, while variant v.8 and v.9 contain insertions at aa 642 and 378, respectively. The 24P4C12 protein exhibits homology to a previously cloned human gene, namely NG22 also known as choline transporter-like protein 4 (gi 14249468). It shows 99% identity and 99% homology to the CTL4 protein over the length of that protein (FIG. 4). 24P4C12 is a multi-transmembrane protein, predicted to carry 10, 11 or 13 transmembrane domains. Bioinformatic analysis indicates that the 24P4C12 protein localizes to the plasma membrane with some endoplasmic reticulum localization (see Table L). Recent evidence indicates that the 24P4C12 protein is a 10 transmembrane protein that localizes to the cell surface (O'Regan S et al PNAS 2000, 97:1835).

Choline as an essential component of cell membranes that plays an important role in cell integrity, growth and survival of normal and tumor cells. Choline accumulates at increased concentration in tumor cells relative to their normal counterparts and as such constitutes a tool for the detection of cancer cells by magnetic resonance imaging (Kurhanewicz J et al, *J Magn Reson Imaging*. 2002.). In addition to its role in maintaining membrane integrity, choline mediates signal transduction event from the membrane to the nucleus (Spiegel S, Milstien S. *J Membr Biol*. 1995, 146:225). Choline metabolites include sphingosylphosphorylcholine and lysophosphatidylcholine, both of which activate G-protein coupled receptors (Xu F et al *Biochim Biophys Acta* 2002, 1582:81). In addition, choline results in the activation of kinase pathways including Raf-1 (Lee M, Han S S, *Cell Signal* 2002, 14:373.). Choline also plays a role in regulating DNA methylation and regulation of gene expression. For example, choline methanolites regulate the expression of cytokines and

chemokines essential for tumor growth (Schwartz B M et al., *Gynecol Oncol.* 2001, 81:291; Denda A et al., *Carcinogenesis.* 2002, 23:245). Due to its effect on cell signaling and gene expression, choline controls cell growth and survival (Holmes-McNary M Q et al., *J Biol Chem.* 2001, 276: 41197; Albright et al., *FASEB* 1996, 10:510). Choline deficiency results in cell death, apoptosis and transformation, while accumulation of choline is associated with tumor growth (Zeisel S et al., *Carcinogenesis* 1997, 18:731). Accordingly, when 24P4C12 functions as a regulator of tumor formation, cell proliferation, invasion or cell signaling, 24P4C12 is used for therapeutic, diagnostic, prognostic and/or preventative purposes.

Example 45

Identification and Confirmation of Potential Signal Transduction Pathways

Many mammalian proteins have been reported to interact with signaling molecules and to participate in regulating signaling pathways. (*J Neurochem.* 2001; 76:217-223). In particular, choline have been reported to activate MAK cascades as well as G proteins, and been associated with the DAG and ceramide and sphingophosphorylcholine signaling pathway (Cummings et al., above). In addition, choline transmit its signals by regulating choline-kinase and phospholipase activity, resulting in enhance tumorigenic effect (Ramirez et al., *Oncogene.* 2002, 21:4317; Lucas et al., *Oncogene.* 2001, 20:1110; Chung T et al., *Cell Signal.* 2000, 12:279).

Using immunoprecipitation and Western blotting techniques, proteins are identified that associate with 24P4C12 and mediate signaling events. Several pathways known to play a role in cancer biology can be regulated by 24P4C12, including phospholipid pathways such as PI3K, AKT, etc, adhesion and migration pathways, including FAK, Rho, Rac-1, etc, as well as mitogenic/survival cascades such as ERK, p38, etc (*Cell Growth Differ.* 2000, 11:279; *J Biol Chem.* 1999, 274:801; *Oncogene.* 2000, 19:3003; *J. Cell Biol.* 1997, 138:913). Using Western blotting and other techniques, the ability of 24P4C12 to regulate these pathways is confirmed. Cells expressing or lacking 24P4C12 are either left untreated or stimulated with cytokines, androgen and anti-integrin antibodies. Cell lysates are analyzed using anti-phospho-specific antibodies (Cell Signaling, Santa Cruz Biotechnology) in order to detect phosphorylation and regulation of ERK, p38, AKT, PI3K, PLC and other signaling molecules.

To confirm that 24P4C12 directly or indirectly activates known signal transduction pathways in cells, luciferase (luc) based transcriptional reporter assays are carried out in cells expressing individual genes. These transcriptional reporters contain consensus-binding sites for known transcription factors that lie downstream of well-characterized signal transduction pathways. The reporters and examples of these associated transcription factors, signal transduction pathways, and activation stimuli are listed below.

1. NFkB-luc, NFkB/Rel; Ik-kinase/SAPK; growth/apoptosis/stress
2. SRE-luc, SRF/TCF/ELK1; MAPK/SAPK; growth/differentiation
3. AP-1-luc, FOS/JUN; MAPK/SAPK/PKC; growth/apoptosis/stress
4. ARE-luc, androgen receptor; steroids/MAPK; growth/differentiation/apoptosis
5. p53-luc, p53; SAPK; growth/differentiation/apoptosis
6. CRE-luc, CREB/ATF2; PKA/p38; growth/apoptosis/stress
7. TCF-luc, TCF/Lef; $\square$ -catenin, Adhesion/invasion

Gene-mediated effects can be assayed in cells showing mRNA expression. Luciferase reporter plasmids can be introduced by lipid-mediated transfection (TFX-50, Promega). Luciferase activity, an indicator of relative transcriptional activity, is measured by incubation of cell extracts with luciferin substrate and luminescence of the reaction is monitored in a luminometer.

Signaling pathways activated by 24P4C12 are mapped and used for the identification and validation of therapeutic targets. When 24P4C12 is involved in cell signaling, it is used as target for diagnostic, prognostic, preventative and/or therapeutic purposes.

Example 46

24P4C12 Functions as a Choline Transporter

Sequence and homology analysis of 24P4C12 indicate that 24P4C12 carries a transport domain and that 24P4C12 functions as a choline transporter. In order to confirm that 24P4C12 transports choline, primary and tumor cells, including prostate, colon, bladder and lung lines, are grown in the presence and absence of ^3H -choline. Radioactive choline uptake is measured by counting incorporated counts per minutes (cpm). Parental 24P4C12 negative cells are compared to 24P4C12-expressing cells using this and similar assays. Similarly, parental and 24P4C12-expressing cells can be compared for choline content using NMR spectroscopy. These assay systems can be used to identify small molecules and antibodies that interfere with choline uptake and/or with the function of 24P4C12.

Thus, compounds and small molecules designed to inhibit 24P4C12 function and downstream signaling events are used for therapeutic diagnostic, prognostic and/or preventative purposes.

Example 47

Regulation of Transcription

The cell surface localization of 24P4C12 and its ability to regulate DNA methylation indicate that it is effectively used as a modulator of the transcriptional regulation of eukaryotic genes. Regulation of gene expression is confirmed, e.g., by studying gene expression in cells expressing or lacking 24P4C12. For this purpose, two types of experiments are performed.

In the first set of experiments, RNA from parental and 24P4C12-expressing cells are extracted and hybridized to commercially available gene arrays (Clontech) (Smid-Koopman E et al. *Br J Cancer.* 2000, 83:246). Resting cells as well as cells treated with FBS, pheromones, or growth factors are compared. Differentially expressed genes are identified in accordance with procedures known in the art. The differentially expressed genes are then mapped to biological pathways (Chen K et al. *Thyroid.* 2001, 11:41.).

In the second set of experiments, specific transcriptional pathway activation is evaluated using commercially available (Stratagene) luciferase reporter constructs including: NFkB-luc, SRE-luc, ELK1-luc, ARE-luc, p53-luc, and CRE-luc. These transcriptional reporters contain consensus binding sites for known transcription factors that lie downstream of well-characterized signal transduction pathways, and represent a good tool to ascertain pathway activation and screen for positive and negative modulators of pathway activation.

Thus, 24P4C12 plays a role in gene regulation, and it is used as a target for diagnostic, prognostic, preventative and/or therapeutic purposes.

Example 48

Involvement in Tumor Progression

The 24P4C12 gene can contribute to the growth of cancer cells. The role of 24P4C12 in tumor growth is confirmed in a variety of primary and transfected cell lines including prostate, and bladder cell lines, as well as NIH 3T3 cells engineered to stably express 24P4C12. Parental cells lacking 24P4C12 and cells expressing 24P4C12 are evaluated for cell growth using a well-documented proliferation assay (Fraser S P, et al., *Prostate* 2000; 44:61, Johnson D E, Ochieng J, Evans S L. *Anticancer Drugs*. 1996, 7:288). Such a study was performed on prostate cancer cells and the results are shown in FIG. 28. The growth of parental PC3 and PC3-24P4C12 cells was compared in low (0.1%) and 10% FBS. Expression of 24P4C12 imparted a growth advantage to PC3 cells grown in 10% FBS. Similarly, expression of 24P4C12 in NIH-3T3 cells enhances the proliferation of these cells relative to control 3T3-neo cells. The effect of 24P4C12 can also be observed on cell cycle progression. Control and 24P4C12-expressing cells are grown in low serum overnight, and treated with 10% FBS for 48 and 72 hrs. Cells are analyzed for BrdU and propidium iodide incorporation by FACS analysis.

To confirm the role of 24P4C12 in the transformation process, its effect in colony forming assays is investigated. Parental NIH-3T3 cells lacking 24P4C12 are compared to NIH-3T3 cells expressing 24P4C12, using a soft agar assay under stringent and more permissive conditions (Song Z. et al. *Cancer Res.* 2000; 60:6730).

To confirm the role of 24P4C12 in invasion and metastasis of cancer cells, a well-established assay is used. A non-limiting example is the use of an assay which provides a basement membrane or an analog thereof used to detect whether cells are invasive (e.g., a Transwell Insert System assay (Becton Dickinson) (*Cancer Res.* 1999; 59:6010)). Control cells, including prostate, and bladder cell lines lacking 24P4C12 are compared to cells expressing 24P4C12. Cells are loaded with the fluorescent dye, calcein, and plated in the top well of a support structure coated with a basement membrane analog (e.g. the Transwell insert) and used in the assay. Invasion is determined by fluorescence of cells in the lower chamber relative to the fluorescence of the entire cell population.

24P4C12 can also play a role in cell cycle and apoptosis. Parental cells and cells expressing 24P4C12 are compared for differences in cell cycle regulation using a well-established BrdU assay (Abdel-Malek Z A. *J Cell Physiol.* 1988, 136: 247). In short, cells are grown under both optimal (full serum) and limiting (low serum) conditions are labeled with BrdU and stained with anti-BrdU Ab and propidium iodide. Cells are analyzed for entry into the G1, S, and G2M phases of the cell cycle. Alternatively, the effect of stress on apoptosis is evaluated in control parental cells and cells expressing 24P4C12, including normal and tumor prostate, colon and lung cells. Engineered and parental cells are treated with various chemotherapeutic agents, such as etoposide, flutamide, etc. and protein synthesis inhibitors, such as cycloheximide. Cells are stained with annexin V-FITC and cell death is measured by FACS analysis. The modulation of cell death by 24P4C12 can play a critical role in regulating tumor progression and tumor load.

When 24P4C12 plays a role in cell growth, transformation, invasion or apoptosis, it is used as a target for diagnostic, prognostic, preventative and/or therapeutic purposes.

Example 49

Involvement in Angiogenesis

Angiogenesis or new capillary blood vessel formation is necessary for tumor growth (Hanahan D, Folkman J. *Cell*. 1996, 86:353; Folkman J. *Endocrinology*. 1998 139:441). Based on the effect of phosphodiesterase inhibitors on endothelial cells, 24P4C12 plays a role in angiogenesis (DeFouw L et al, *Microvasc Res* 2001, 62:263). Several assays have been developed to measure angiogenesis in vitro and in vivo, such as the tissue culture assays endothelial cell tube formation and endothelial cell proliferation. Using these assays as well as in vitro neo-vascularization, the role of 24P4C12 in angiogenesis, enhancement or inhibition, is confirmed.

For example, endothelial cells engineered to express 24P4C12 are evaluated using tube formation and proliferation assays. The effect of 24P4C12 is also confirmed in animal models in vivo. For example, cells either expressing or lacking 24P4C12 are implanted subcutaneously in immunocompromised mice. Endothelial cell migration and angiogenesis are evaluated 5-15 days later using immunohistochemistry techniques. 24P4C12 affects angiogenesis and it is used as a target for diagnostic, prognostic, preventative and/or therapeutic purposes.

Example 50

Involvement in Adhesion

Cell adhesion plays a critical role in tissue colonization and metastasis. The presence of leucine rich and cysteine rich motifs in 24P4C12 is indicative of its role in cell adhesion. To confirm that 24P4C12 plays a role in cell adhesion, control cells lacking 24P4C12 are compared to cells expressing 24P4C12, using techniques previously described (see, e.g., Haier et al, *Br. J. Cancer*. 1999, 80:1867; Lehr and Pienta, J. *Natl. Cancer Inst.* 1998, 90:11). Briefly, in one embodiment, cells labeled with a fluorescent indicator, such as calcein, are incubated on tissue culture wells coated with media alone or with matrix proteins. Adherent cells are detected by fluorimetric analysis and percent adhesion is calculated. This experimental system can be used to identify proteins, antibodies and/or small molecules that modulate cell adhesion to extracellular matrix and cell-cell interaction. Since cell adhesion plays a critical role in tumor growth, progression, and, colonization, the gene involved in this process can serve as a diagnostic, preventative and therapeutic modality.

Example 51

Detection of 24P4C12 Protein in Cancer Patient Specimens

To determine the expression of 24P4C12 protein, specimens were obtained from various cancer patients and stained using an affinity purified polyclonal rabbit antibody raised against the peptide encoding amino acids 1-14 of 24P4C12 variant 1 and conjugated to KLH (See, Example 10: Generation of 24P4C12 Polyclonal Antibodies.) This antiserum exhibited a high titer to the peptide (>10,000) and recognized 24P4C12 in transfected 293T cells by Western blot and flow

cytometry (FIG. 24) and in stable recombinant PC3 cells by Western blot and immunohistochemistry (FIG. 25). Formalin fixed, paraffin embedded tissues were cut into 4 micron sections and mounted on glass slides. The sections were dewaxed, rehydrated and treated with antigen retrieval solution (0.1 M Tris, pH10) at high temperature. Sections were then incubated in polyclonal rabbit anti-24P4C12 antibody for 3 hours. The slides were washed three times in buffer and further incubated with DAKO EnVision+™ peroxidase-conjugated goat anti-rabbit immunoglobulin secondary antibody (DAKO Corporation, Carpinteria, Calif.) for 1 hour. The sections were then washed in buffer, developed using the DAB kit (SIGMA Chemicals), counterstained using hematoxylin, and analyzed by bright field microscopy. The results showed expression of 24P4C12 in cancer patients' tissue (FIGS. 29 and 30). Tissue from prostate cancer patients showed expression of 24P4C12 in the tumor cells and in the prostate epithelium of tissue normal adjacent to tumor (FIG. 29). Generally, expression of 24P4C12 was high in all prostate tumors and was expressed mainly around the cell membrane indicating that 24P4C12 is membrane associated in prostate tissues. All of the prostate samples tested were positive for 24P4C12. Other tumors that were positive for 24P4C12 included colon adenocarcinoma, breast ductal carcinoma, pancreatic adenocarcinoma, lung adenocarcinoma, bladder transitional cell carcinoma and renal clear cell carcinoma (FIG. 30). Normal tissues investigated for expression of 24P4C12 included heart, skeletal muscle, liver, brain, spinal cord, skin, adrenal, lymph node, spleen, salivary gland, small intestine and placenta. None demonstrated any expression of 24P4C12 by immunohistochemistry. Normal adjacent to tumor tissues were also studied to determine the presence of 24P4C12 protein by immunohistochemistry. These included breast, lung, colon, ileum, bladder, kidney and pancreas. In some of the tissues from these organs there was weak expression of 24P4C12. This expression may relate to the fact that the samples were not truly normal and may indicate a precancerous change. The ability to identify malignancy in tissue that has not undergone obvious morphological changes is an important diagnostic modality for cancerous and precancerous conditions.

These results indicate that 24P4C12 is a target for diagnostic, prophylactic, prognostic and therapeutic applications in cancer.

Throughout this application, various website data content, publications, patent applications and patents are referenced. (Websites are referenced by their Uniform Resource Locator, or URL, addresses on the World Wide Web.) The disclosures of each of these references are hereby incorporated by reference herein in their entireties.

The present invention is not to be limited in scope by the embodiments disclosed herein, which are intended as single illustrations of individual aspects of the invention, and any that are functionally equivalent are within the scope of the invention. Various modifications to the models and methods of the invention, in addition to those described herein, will become apparent to those skilled in the art from the foregoing description and teachings, and are similarly intended to fall within the scope of the invention. Such modifications or other embodiments can be practiced without departing from the true scope and spirit of the invention.

Tables:

a.

TABLE I

Tissues that Express 24P4C12:
Malignant Tissues

Prostate
Bladder
Kidney
Lung
Colon
Ovary
Breast
Uterus
Stomach

TABLE II

Amino Acid Abbreviations

SINGLE LETTER	THREE LETTER	FULL NAME
F	Phe	phenylalanine
L	Leu	leucine
S	Ser	serine
Y	Tyr	tyrosine
C	Cys	cysteine
W	Trp	tryptophan
P	Pro	proline
H	His	histidine
Q	Gln	glutamine
R	Arg	arginine
I	Ile	isoleucine
M	Met	methionine
T	Thr	threonine
N	Asn	asparagine
K	Lys	lysine
V	Val	valine
A	Ala	alanine
D	Asp	aspartic acid
E	Glu	glutamic acid
G	Gly	glycine

TABLE III

Amino Acid Substitution Matrix

Adapted from the GCG Software 9.0 BLOSUM62 amino acid substitution matrix (block substitution matrix). The higher the value, the more likely a substitution is found in related, natural proteins. (See world wide web URL ikp.unibe.ch/manual/blosum62.html)

A	C	D	E	F	G	H	I	K	L	M	N	P	Q	R	S	T	V	W	Y	.
4	0	-2	-1	-2	0	-2	-1	-1	-1	-1	-2	-1	-1	-1	1	0	0	-3	-2	A
	9	-3	-4	-2	-3	-3	-1	-3	-1	-1	-3	-3	-3	-3	-1	-1	-1	-2	-2	C
		6	2	-3	-1	-1	-3	-1	-4	-3	1	-1	0	-2	0	-1	-3	-4	-3	D
			5	-3	-2	0	-3	1	-3	-2	0	-1	2	0	0	-1	-2	-3	-2	E

TABLE III-continued

Amino Acid Substitution Matrix
Adapted from the GCG Software 9.0 BLOSUM62 amino acid substitution matrix (block substitution matrix). The higher the value, the more likely a substitution is found in related, natural proteins. (See world wide web URL ikp.unibe.ch/manual/blosum62.html)

[illegible]

TABLE IV

HLA Class I/II Motifs/Supermotifs			
TABLE IV (A): HLA Class I Supermotifs/Motifs			
	POSITION 2 (Primary Anchor)	POSITION 3 (Primary Anchor)	POSITION C Terminus (Primary Anchor)
SUPERMOTIF			
A1	TILVMS		FWY
A2	LIVMATQ		IVMATL
A3	VSMATLI		RK
A24	YFWVLMT		FIYWL
B7	P		VILEMWHYA
B27	RHK		FYLWMIVA
B44	ED		FWYLMVA
B58	ATS		FWYLIVMA
B62	QLIVMP		FWYMIVLA
MOTIFS			
A1	TSM		Y
A1		DEAS	Y
A2.1	LMVQIAT		VLMAT
A3	LMVISATFCGD		KYRHFA
A11	VTMLISAGNCDF		KRYH
A24	YFWM		FLIW
A*3101	MVTALIS		RK
A*3301	MVALFIST		RK
A*6801	AVTMSLI		RK
B*0702	P		LMFWYAIY
B*3501	P		LMFWYIVA

TABLE IV-continued

B51	P	LIVFWYAM
B*5301	P	IMFWYALV
B*5401	P	ATIVLMFWY

Bolded residues are preferred, italicized residues are less preferred:

A peptide is considered motif-bearing if it has primary anchors at each primary anchor position for a motif or supermotif as specified in the above table.

TABLE IV (B): HLA Class II Supermotif

1	6	9
W, F, Y, V, I, L	A, V, I, L, P, C, S, T	A, V, I, L, C, S, T, M, Y

TABLE IV (C): HLA Class II Motifs

MOTIFS	1° anchor 1	2	3	4	5	1° anchor 6	7	8	9
DR4	preferred deleterious	FMYLIVW	M	T	W	I	VSTCPALIM	MH R	MH WDE
DR1	preferred deleterious	MFLIVWY	C	CH	PAMQ FD	CWD	VMATSPLIC	M GDE	AVM D
DR7	preferred deleterious	MFLIVWY	M C	W	A G		IVMSACTPL	M GRD	IV G
DR3	MOTIFS	1° anchor 1	2	3	1° anchor 4	5	1° anchor 6		
	Motif a preferred	LIVMFY			D				
	Motif b preferred	LIVMFAY			DNQEST			KRH	
	DR Supermotif	MFLIVWY						VMSTACPLI	

Italicized residues indicate less preferred or "tolerated" residues

TABLE IV (D): HLA Class I Supermotifs

SUPER-		POSITION:								
MOTIFS		1	2	3	4	5	6	7	8	C-terminus
A1			<u>1°Anchor</u> TILVMS							<u>1°Anchor</u> FWY
A2			<u>1°Anchor</u> LIVMATQ							<u>1°Anchor</u> LIVMAT
A3	Preferred		<u>1°Anchor</u> VSMATLI	YFW (4/5)		YFW (3/5)		YFW (4/5)	P (4/5)	<u>1°Anchor</u> RK
	deleterious	DE (3/5); P (5/5)		DE (4/5)						
A24			<u>1°Anchor</u> YFWTVLMT							<u>1°Anchor</u> FIYWLM
B7	Preferred	FWY (5/5) LIVM (3/5)	<u>1°Anchor</u> P	FWY (4/5)					FWY (3/5)	<u>1°Anchor</u> VILFMWYA
	deleterious	DE (3/5); P (5/5); G (4/5); A (3/5); QN (3/5)				DE (3/5)	G (4/5)	QN (4/5)	DE (4/5)	
B27			<u>1°Anchor</u> RHK							<u>1°Anchor</u> FYLWMIVA
B44			<u>1°Anchor</u> ED							<u>1°Anchor</u> FWYLIMVA
B58			<u>1°Anchor</u> ATS							<u>1°Anchor</u> FWYLIVMA
B62			<u>1°Anchor</u> QLIVMP							<u>1°Anchor</u> FWYMIVLA

Italicized residues indicate less preferred or "tolerated" residues

TABLE IV (E): HLA Class I Motifs

		POSITION				
		1	2	3	4	5
A1 9-mer	preferred	GFYW	1°Anchor STM	DEA	YFW	
	deleterious	DE		RHKLIVMP	A	G
A1 9-mer	preferred	GRHK	ASTCLIVM	1°Anchor DEAS	GSTC	
	deleterious	A	RHKDEPYFW		DE	PQN
A1 10-mer	preferred	YFW	1°Anchor STM	DEAQN	A	YFWQN
	deleterious	GP		RHKGLIVM	DE	RHK

TABLE IV-continued

A1 10- mer	preferred	YFW	STCLIVM	<u>1°Anchor</u> DEAS	A	YFW
A2.1 9-mer	deleterious preferred	RHK YFW	RHKDEPYFW <u>1°Anchor</u> LMIVQAT	YFW	STC	P YFW
A2.1 10- mer	deleterious preferred	DEP AYFW	<u>1°Anchor</u> LMIVQAT	DERKH LVIM	G	
A3	deleterious preferred	DEP RHK	<u>1°Anchor</u> LMVISATFCGD	DE YFW	RKHA PRHKYF W	P A
A11	deleterious preferred	DEP A	<u>1°Anchor</u> VTLMISAGNCDF	DE YFW	YFW	A
A24 9-mer	deleterious preferred	DEP YFWRHK	<u>1°Anchor</u> YFWM		STC	
A24 10- mer	deleterious Preferred	DEG	<u>1°Anchor</u> YFWM	DE	G P	QNP YFWP
A3101	Deleterious Preferred	RHK	<u>1°Anchor</u> MVTALLIS	GDE YFW	QN P	RHK
A3301	Deleterious Preferred	DEP	<u>1°Anchor</u> MVALFIST	DE YFW		ADE
A6801	Deleterious Preferred	GP YFWSTC	<u>1°Anchor</u> AVTMSLI	DE		YFWLIV M
B0702	deleterious Preferred	GP RHKFWY	<u>1°Anchor</u> P	DEG RHK		RHK RHK
A1 9-mer	preferred	GFYW	<u>1°Anchor</u> STM	DEA	YFW	
A1 9-mer	deleterious preferred	DE GRHK	ASTCLIVM	RHKLIVMP <u>1°Anchor</u> DEAS	A GSTC	G
B3501	deleterious Preferred	A DEQNP FWYLIVM	RHKDEPYFW <u>1°Anchor</u> P	DEP FWY	DE DE	PQN DE
B51	deleterious Preferred	AGP LIVMFWY	<u>1°Anchor</u> P	FWY	STC	G FWY
B5301	deleterious preferred	AGPDER HKSTC LIVMFWY	<u>1°Anchor</u> P	FWY	STC	DE FWY
B5401	deleterious preferred	AGPQN FWY	<u>1°Anchor</u> P	FWYLIVM		LIVM
	deleterious	GPQNDE		GDESTC		RHKDE
POSITION						
		6	7	8	9 or C-terminus	C- terminus
A1 9-mer	preferred	P	DEQN	YFW	<u>1°Anchor</u> Y	
A1 9-mer	deleterious preferred	A ASTC	LIVM	DE	<u>1°Anchor</u> Y	
A1 10- mer	deleterious preferred	RHK	PG PASTC	GP GDE	P	<u>1°Anchor</u> Y
A1 10- mer	deleterious preferred	QNA	RHKYFW PG	RHK G	A YFW	<u>1°Anchor</u> Y
	deleterious	G		PRHK	QN	

TABLE IV-continued

A2.1 9-mer	preferred		A	P	<u>1°Anchor</u> <u>VLIMAT</u>	
A2.1 10- mer	deleterious preferred	RKH G	DERKH	FYWL VIM		<u>1°Anchor</u> <u>VLIMAT</u>
A3	deleterious preferred	YFW	RKH	DERKH P	RKH <u>1°Anchor</u> <u>KYRHEA</u>	
A11	deleterious preferred	YFW	YFW	P	<u>1°Anchor</u> <u>KRYH</u>	
A24 9-mer	deleterious preferred		A YFW	G YFW	<u>1°Anchor</u> <u>FLIW</u>	
A24 10- mer	deleterious Preferred	DERHK	G P	AQN		<u>1°Anchor</u> <u>FLIW</u>
A3101	Deleterious Preferred	DE YFW	A YFW	QN AP	DEA <u>1°Anchor</u> RK	
A3301	Deleterious Preferred	DE	DE AYFW	DE	<u>1°Anchor</u> RK	
A6801	Deleterious Preferred		YFW	P	<u>1°Anchor</u> RK	
B0702	deleterious Preferred	RHK	RHK	A PA	<u>1°Anchor</u> <u>LMFWYALV</u>	
A1 9-mer	preferred	P	DEQN	YFW	<u>1°Anchor</u> Y	
A1 9-mer	deleterious preferred	A ASTC	LIVM	DE	<u>1°Anchor</u> Y	
B3501	deleterious deleterious Preferred	RHK GDE	PG QN FWY	GP DE		<u>1°Anchor</u> <u>LMFWYIVA</u>
B51	deleterious Preferred	G	G	FWY	<u>1°Anchor</u> <u>LIVFWYAM</u>	
B5301	deleterious preferred	G	DEQN LIVMFWY	GDE FWY	<u>1°Anchor</u> <u>IMFWYALV</u>	
B5401	deleterious preferred	G	RHKQN ALIVM	DE FWYAP	<u>1°Anchor</u> <u>ATIVLMF</u> <u>WY</u>	
	deleterious	DE	QNDGE	DE		

TABLE IV (F): Summary of HLA-supertypes
Overall phenotypic frequencies of HLA-supertypes in different ethnic populations

Specificity			Phenotypic frequency					
Supertype	Position 2	C-Terminus	Caucasian	N.A. Black	Japanese	Chinese	Hispanic	Average
B7	P	AILMVFWY	43.2	55.1	57.1	43.0	49.3	49.5
A3	AILMVST	RK	37.5	42.1	45.8	52.7	43.1	44.2
A2	AILMVT	AILMVT	45.8	39.0	42.4	45.9	43.0	42.2
A24	YF (WIVLMT)	FI (YWLM)	23.9	38.9	58.6	40.1	38.3	40.0
B44	E (D)	FWYLIMVA	43.0	21.2	42.9	39.1	39.0	37.0
A1	TI (LVMS)	FWY	47.1	16.1	21.8	14.7	26.3	25.2
B27	RHK	FYL (WMI)	28.4	26.1	13.3	13.9	35.3	23.4
B62	QL (IVMP)	FWY (MIV)	12.6	4.8	36.5	25.4	11.1	18.1
B58	ATS	FWY (LIV)	10.0	25.1	1.6	9.0	5.9	10.3

TABLE IV (G):
Calculated population coverage afforded by different HLA-supertype combinations

		Phenotypic frequency					
HLA-supertypes		Caucasian	N.A Blacks	Japanese	Chinese	Hispanic	Average
A2, A3 and B7		83.0	86.1	87.5	88.4	86.3	86.2
A2, A3, B7, A24, B44 and A1		99.5	98.1	100.0	99.5	99.4	99.3
		99.9	99.6	100.0	99.8	99.9	99.8

TABLE IV-continued

A2, A3, B7, A24,
B44, A1, B27, B62,
and B 58

Motifs indicate the residues defining supertype specificities.

The motifs incorporate residues determined on the basis of published data to be recognized by multiple alleles within the supertype.

Residues within brackets are additional residues also predicted to be tolerated by multiple alleles within the supertype.

TABLE V

Frequently Occurring Motifs			
Name	avrg. % identity	Description	Potential Function
zf-C2H2	34%	Zinc finger, C2H2 type	Nucleic acid-binding protein functions as transcription factor, nuclear location probable
cytochrome_b_N	68%	Cytochrome b(N-terminal)/b6/petB	membrane bound oxidase, generate superoxide
Ig	19%	Immunoglobulin domain	domains are one hundred amino acids long and include a conserved intradomain disulfide bond.
WD40	18%	WD domain, G-beta repeat	tandem repeats of about 40 residues, each containing a Trp-Asp motif. Function in signal transduction and protein interaction
PDZ	23%	PDZ domain	may function in targeting signaling molecules to sub-membranous sites
LRR	28%	Leucine Rich Repeat	short sequence motifs involved in protein-protein interactions
Pkinase	23%	Protein kinase domain	conserved catalytic core common to both serine/threonine and tyrosine protein kinases containing an ATP binding site and a catalytic site
PH	16%	PH domain	pleckstrin homology involved in intracellular signaling or as constituents of the cytoskeleton
EGF	34%	EGF-like domain	30-40 amino-acid long found in the extracellular domain of membrane-bound proteins or in secreted proteins
Rvt	49%	Reverse transcriptase (RNA-dependent DNA polymerase)	
Ank	25%	Ank repeat	Cytoplasmic protein, associates integral membrane proteins to the cytoskeleton
Oxidored_q1	32%	NADH-Ubiquinone/plastoquinone (complex I), various chains	membrane associated. Involved in proton translocation across the membrane
Efhand	24%	EF hand	calcium-binding domain, consists of a 12 residue loop flanked on both sides by a 12 residue alpha-helical domain

TABLE V-continued

Frequently Occurring Motifs			
Name	avrg. % identity	Description	Potential Function
Rvp	79%	Retroviral aspartyl protease	Aspartyl or acid proteases, centered on a catalytic aspartyl residue
Collagen	42%	Collagen triple helix repeat (20 copies)	extracellular structural proteins involved in formation of connective tissue. The sequence consists of the G-X-Y and the polypeptide chains forms a triple helix.
Fn3	20%	Fibronectin type III domain	Located in the extracellular ligand-binding region of receptors and is about 200 amino acid residues long with two pairs of cysteines involved in disulfide bonds
7tm_1	19%	7 trans-membrane receptor (rhodopsin family)	seven hydrophobic trans-membrane regions, with the N-terminus located extracellularly while the C-terminus is cytoplasmic. Signal through G proteins

TABLE VI

Motifs and Post-translational Modifications of 24P4C12			
N-glycosylation site			
29-32	NRSC	(SEQ ID NO: 48)	
69-72	NSTG	(SEQ ID NO: 49)	
155-158	NMTV	(SEQ ID NO: 50)	
197-200	NDTT	(SEQ ID NO: 51)	
298-301	NLSA	(SEQ ID NO: 52)	
393-396	NISS	(SEQ ID NO: 53)	
405-408	NTSC	(SEQ ID NO: 54)	
416-419	NSSC	(SEQ ID NO: 55)	
678-681	NGSL	(SEQ ID NO: 56)	
Protein kinase C phosphorylation site			
22-24	SfR		
218-220	SvK		
430-432	SsK		

TABLE VI-continued

Motifs	and	Post-translational Modifications of 24P4C12	
494-496	TLR		5
573-575	SaK		
619-621	SgR		
Casein kinase II phosphorylation site			10
31-34	SCTD (SEQ ID NO: 57)		
102-105	SVAE (SEQ ID NO: 58)		
119-122	SCPE (SEQ ID NO: 59)		
135-138	TVGE (SEQ ID NO: 60)		15
304-307	SVQE (SEQ ID NO: 61)		
Tyrosine kinase phosphorylation site			20
6-13	RDEDEAY (SEQ ID NO: 62)		
N-myristoylation site			
72-77	GAYCGM (SEQ ID NO: 63)		
76-81	GMGENK (SEQ ID NO: 64)		

TABLE VI-continued

Motifs	and	Post-translational Modifications of 24P4C12	
151-156	GVPWNM (SEQ ID NO: 65)		
207-212	GLIDSL (SEQ ID NO: 66)		
272-277	GIYYCW (SEQ ID NO: 67)		
287-292	GASISQ (SEQ ID NO: 68)		
349-354	GQMMST (SEQ ID NO: 69)		
449-454	GLFWTL (SEQ ID NO: 70)		
467-472	GAFASF (SEQ ID NO: 71)		
Amidation site			
695-698	IGKK (SEQ ID NO: 72)		
Leucine zipper pattern			
245-266	LFILLRLRLVAGPLVLVLILGVL (SEQ ID NO: 73)		
Cysteine-rich region			
536-547	CIMCCFKCCLWC (SEQ ID NO: 74)		25

TABLE VII

Search Peptides
Variant 1, 9-mers, 10-mers, 15-mers (SEQ ID NO: 75)
MGGKQRDEDD EAYGKPKYD PSFRGPIKNR SCTDVICCVL FLLFILGYIV VGIVAWLYGD PRQVLYPRNS TGAYCGMGEM KDKPYLLYFN IFSCILSSNI ISVAENGLQC PTPQVCVSSC PEDPWTVGKN EFSQTVGEVF YTKNRNFCPL GVPWNMTVIT SLQQELCPSP LLPSAPALGR CFPWTNVTPP ALPGITNDTT IQQGISGLID SLNARDISVK IFEDFAQSWY WILVALGVAL VLSLLFILLL RLVAGPLVLV LILGVLGVL YGIYYCWEEY RVLKDKGAST SQLGFTTNLS AYQSVQETWL AALIVLAVLE AILLMLIFL RQRIRIAIAL LKEASKAVGQ MMSTMFYPLV TFVLLLICIA YWAMTALYLA TSGQPQYVLW ASNISSPGCE KVPINTSCNP TAHLVNSSCP GLMCVFQGY S KGLIQRVSF NLQIYGVVLG FWTLNWVLAL GQCVLAGAFA SFYWAFHKPQ DIPTFPLISA FIRTLYHTG SLAFGALILT LVQIARVILE YIDHKLRGVQ NPVARCIMCC FKCCLWCLEK FIKFLNRNAY IMIAIYGKNF CVSAKNAFML LMRNIVRVV LDKVTDLLLF FGKLLVVGGV GVLSFFFFSG RIPGLGKDFK SPHLNYYWLP IMTSILGAYV IASGFFSVFG MCDVTLFLCF LEDLERNNGS LDRPYMSKS LLKILGKKNE APPDNKKRK
Variant 3: 9-mers
GRCFPWTNITPPALPGI (SEQ ID NO: 76) 10-mers
LGRCFPWTNITPPALPGIT (SEQ ID NO: 77) 15-mers
PSAPALGRCFPWTNITPPALPGITNDTTI (SEQ ID NO: 78)
Variant 5: 9-mers
VLEAILLVLIFLRQRI (SEQ ID NO: 79) 10-mers
AVLEAILLVLIFLRQRI (SEQ ID NO: 80) 15-mers
ALIVLAVLEAILLVLIFLRQRI (SEQ ID NO: 81)
Variant 6: 9-mers
GYSSKGLIPRSVFNLQI (SEQ ID NO: 82)

Figure 2E-3

```
1621 GCTGTTTCAAGTGCTGCCTCTGGTGTCTGGAAAAATTATCAAGTTCCTAAACCGCAATG
560 Y I M I A I Y G K N F C V S A K N A F M
1681 CATACATCATGATCGCCATCTACGGGAAGAATTTCTGTGTCTCAGCCAAAAATGCGTTCA
580 L L M R N I V R V V V L D K V T D L L L
1741 TGCTACTCATGCGAAACATTGTCAGGGTGGTCTGCTGGACAAAGTCACAGACCTGCTGC
600 F F G K L L V V G G V G V L S F F F F S
1801 TGTTCTTTGGGAAGCTGCTGGTGGTGGGAGGCGTGGGGTCTGTCTTCTTTTTTTTCT
620 G R I P G L G K D F K S P H L N Y Y W L
1861 CCGGTCGCATCCCGGGGCTGGGTAAAGACTTTAAGAGCCCCACCTCAACTATTACTGGC
640 P I M T S I L G A Y V I A S G F F S V F
1921 TGCCCATCATGACCTCCATCCTGGGGGCCATGTCTCATCGCCAGCGGCTTCTTCAGCGTTT
660 G M C V D I L F L C F L E D L E R N N G
1981 TCGGCATGTGTGTGGACACGCTCTTCTCTGCTTCCTGGAAGACCTGGAGCGGAACAACG
680 S L D R P Y Y M S K S L L K I L G K K N
2041 GCTCCCTGGACCGGCCCTACTACATGTCCAAGAGCCTTCTAAAGATTCTGGGCAAGAAGA
700 E A P P D N K K R K K *
2101 ACGAGCGCCCCCGGACAACAAGAAGAGGAAGAAAGTGAcagctccggccctgatccagga
2161 ctgcacccccacccccaccgtccagccatccaacctcacttcgccttacaggtctccattt
2221 tgtggttaaaaaaggttttaggccaggcgccgtggctcacgcctgtaatccaacactttg
2281 agaggctgaggcgggcgggatcacctgagtcaggagttcgagaccagcctggccaacatgg
2341 tgaaacctccgtctctattataaaatacaaaaattagccgagagtggtggcatgcacctgt
2401 catcccagctactcgggaggtgaggcaggagaatcgcttgaaacccgggaggcagaggtt
2461 gcagtgagccgagatcgcgccactgcaactccaacctgggtgacagactctgtotccaaa
2521 caaaacaaacaaacaaaagattttattaaagatattttgtaaactcagtaaaaaaaaaa
2581 aaaaaaa
```

Figure 2F. The cDNA (SEQ ID. NO. : 12) and amino acid sequence (SEQ ID. NO. : 13) of 24P4C12 v.6.

The start methionine is underlined. The open reading frame extends from nucleic acid 6-2138 including the stop codon.

Figure 2F-1

```
1 M G G K Q F D E D D E A Y G K P V K Y
1 gagccATGGGGGAAAGCAGCGGGACGAGGATGACGAGGCCTACGGGAAGCCAGTCAAAT
20 D P S F R S P I K N R S C T D V I C C V
61 ACGACCCCTCCTTTCGAGGCCCATCAAGAACAGAAGCTGCACAGATGTCATCTGCTGCG
40 L F L L F I L G Y I V V G I V A W L Y G
121 TCCTCTTCTGCTCTTCATTCTAGGTTACATCGTGGTGGGGATTGTGGCCTGGTTGTATG
60 D P R Q V L Y P R N S T G A Y C G M G E
181 GAGACCCCCGCAAGTCCTCTACCCAGGAACCTCTACTGGGGCTACTGTGGCATGGGGG
80 N K D K P Y L L Y F N I F S C I L S S N
241 AGAACAAAGATAAGCCGTATCTCCTGTACTTCAACATCTTCAGCTGCATCCTGTCCAGCA
```

TABLE VII-continued

Search Peptides	
10-mers	
QGYSSKGLIPRSVFNLQIY (SEQ ID NO: 83)	
15-mers	
LMCVFQGYSSKGLIPRSVFNLQIYGVGL (SEQ ID NO: 84)	
Variant 7	
9-mers	
SWYWILVAVGQMMSTM (SEQ ID NO: 85)	
10-mers	
QSWYWILVAVGQMMSTM (SEQ ID NO: 86)	
15-mers	
FEDFAQSWYWILVAVGQMMSTMFYPLVT (SEQ ID NO: 87)	
Variant 8	
9-mers	
NYYWLPIMRNPITPTGHVFQTSILGAYV (SEQ ID NO: 88)	
10-mers	
LNYYWLPIMRNPITPTGHVFQTSILGAYVI (SEQ ID NO: 89)	
15-mers	
FKSPHLNYYWLPIMRNPITPTGHVFQTSILGAYVIASGFF (SEQ ID NO: 90)	
Variant 9	
9-mers	
YWAMTALYPLPTQPATLGYVLWASNI (SEQ ID NO: 91)	
10-mers	
AYWAMTALYPLPTQPATLGYVLWASNIS (SEQ ID NO: 92)	
15-mers	
LLICIAYWAMTALYPLPTQPATLGYVLWASNISSPGCE (SEQ ID NO: 93)	

Tables VII-XXI:

TABLE VIII		
V1-HLA-A1-9mers-24P4C12		
Each peptide is a portion of SEQ		
ID NO: 3; each start position is		
specified, the length of peptide is		
9 amino acids, and the end		
position for each peptide is the		
start position plus eight.		
Start	Subsequence	Score
58	YGDPRQVLY	125.000
662	CVDTLFLCF	25.000
77	MGENKDKPY	11.250
594	VTDLLLFFG	6.250
698	KNEAPPDNK	4.500
318	VLEAILLLM	4.500
363	VLLLICIAI	2.500
489	SAFIRTLRY	2.500
267	GVLAYGIYY	2.500
689	KSLKILGK	1.500
470	ASFYWAFHK	1.500
222	FEDFAQSWY	1.250
32	CTDVICCVL	1.250
5	QRDEDEAY	1.250
121	PEDPWTVGK	1.000
379	LATSGQPQY	1.000
700	EAPPDNKKR	1.000
558	NAYIMIAIY	1.000
542	KCCLWCLEK	1.000
7	DEDEAYGK	1.000
11	EAYGKPVKY	1.000

TABLE VIII-continued

670	FLEDLERNN	0.900
276	CWEEYRVLR	0.900
518	ILEYIDHKL	0.900
417	SSCPGLMCV	0.750
437	RSVFNLQIY	0.750
80	NKDKPYLLY	0.625
263	LGVLVGLAY	0.625
546	WCLEKFIKF	0.500
243	SLLFILLLR	0.500
238	VALVLSLLF	0.500
579	MLLMRNIVR	0.500
465	LAGAFASFY	0.500
421	GLMCVFQGY	0.500
508	ILTLVQIAR	0.500
593	KVTDLLLFF	0.500
321	AILLLMLIF	0.500
36	ICCVLFLLF	0.500
50	VVGIVAWLY	0.500
186	NVTPPALPG	0.500
609	GVGVLSFFF	0.500
287	GASISQLGF	0.500
187	VTPPALPGI	0.500
668	LCFLEDLER	0.500
323	LLMLIFLR	0.500
272	GIYYCWEEY	0.500
521	YIDHKLRGV	0.500
253	VAGPLVLVL	0.500
398	GCEKVPINT	0.450
560	YIMIAIYGK	0.400
338	IALLKEASK	0.400
135	TVGEVFTYK	0.400
349	QMMSTMIFY	0.375
118	SSCPEDPWT	0.300

161

TABLE VIII-continued

305	VQETWLAAL	0.270	
629	FKSPHLNYY	0.250	
214	ARDISVKIF	0.250	
702	PPDNKKRKK	0.250	5
641	IMTSILGAY	0.250	
678	NGSLDRPYY	0.250	
513	QIARVILEY	0.250	
483	PTFPLISAF	0.250	
120	CPEDPWTVG	0.225	
129	KNEFSQTVG	0.225	10
136	VGVEVFTYKN	0.225	
170	FLLPSAPAL	0.200	
147	FCLPGVPWN	0.200	
393	NISSPGCEK	0.200	
464	VLAGAFASF	0.200	
517	VILEYIDHK	0.200	
424	CVFQGYSSK	0.200	15
394	ISSPGCEKV	0.150	
133	SQTVGEVVFY	0.150	
613	LSFFFFSGR	0.150	
132	FSQTVGEVF	0.150	
488	ISAFIRTLR	0.150	
163	QQELCPSFL	0.135	20
199	TTTQQGISG	0.125	
485	FPLISAFIR	0.125	
607	VGGVGVLSF	0.125	
134	QTVGEVFTY	0.125	
575	KNAFMLLMR	0.125	
266	LGVLAYGIY	0.125	25
40	LFLLFILGY	0.125	
196	TNDTTTQQG	0.125	
610	VGVLSPFFF	0.125	
360	VTFVLLLIC	0.125	
156	MTVITSLQQ	0.125	
677	NNGSLDRPY	0.125	30
498	HTGSLAFGA	0.125	
172	LPSAPALGR	0.125	
195	ITNDTTTQQ	0.125	
452	WTLNWWLAL	0.125	
353	STMFYPLVT	0.125	
443	QIYGVGLGF	0.100	35
543	CCLWCLEKF	0.100	
207	GLIDSLNAR	0.100	
407	SCNPATAHLV	0.100	
180	RCFPWTNVT	0.100	
354	TMFYPLVTF	0.100	
V3I-HLA-A1-9mers-24P4C12			40
Each peptide is a portion of SEQ ID NO: 7; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.			
Start	Subsequence	Score	
9	ITPPALPGI	0.500	
8	NITPPALPG	0.500	
2	RCFPWTNIT	0.100	50
6	WTNITPPAL	0.050	
7	TNITPPALP	0.001	
1	GRCFPWTNI	0.001	
3	CFPWTNITP	0.000	
5	PWTNITPPA	0.000	
4	FPWTNITPP	0.000	55
V5-HLA-A1-9mers-24P4C12			
Each peptide is a portion of SEQ ID NO: 3; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.			60
Start	Subsequence	Score	
1	VLEAILLLV	4.500	
6	LLLVLIFLR	0.500	65
4	AILLLVLIF	0.500	

162

TABLE VIII-continued

8	LVLIFLRQR	0.100	
7	LLVLIFLRQ	0.050	
5	ILLVLIFL	0.050	
3	EAILLLVLI	0.020	
9	VLIFLRQRI	0.010	
2	LEAILLLVL	0.003	
V6-HLA-A1-9mers-24P4C12			
Each peptide is a portion of SEQ ID NO: 13; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.			
Start	Subsequence	Score	
6	GLIPRSVFN	0.200	
2	YSSKGLIPR	0.075	
5	KGLIPRSVF	0.025	
7	LIPRSVFNL	0.005	
3	SSKGLIPRS	0.003	20
4	SKGLIPRSV	0.001	
9	PRSVFNLQI	0.000	
8	IPRSVFNLQ	0.000	
1	GYSSKGLIP	0.000	
V7-HLA-A1-9mers-24P4C12			
Each peptide is a portion of SEQ ID NO: 15; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.			25
Start	Subsequence	Score	
7	VAVGQMMST	0.050	
6	LVAVGQMM	0.050	
8	AVGQMMSTM	0.010	35
5	ILVAVGQMM	0.010	
4	WILVAVGQM	0.010	
3	YWILVAVGQ	0.001	
1	SWYWILVAV	0.001	
2	WYWILVAVG	0.000	
V8-HLA-A1-9mers-24P4C12			40
Each peptide is a portion of SEQ ID NO: 17; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.			
Start	Subsequence	Score	
11	PITPTGHVF	0.100	
19	FQTSILGAY	0.075	50
20	QTSILGAYV	0.050	
17	HVFQTSILG	0.050	
12	ITPTGHVFQ	0.050	
1	NYWLPIMR	0.025	
13	TPTGHVFQT	0.013	
8	MRNPITPTG	0.010	55
4	WLPIMRNPI	0.010	
5	LPIMRNPI	0.005	
18	VFQTSILGA	0.003	
10	NPITPTGHV	0.003	
15	TGHVFQTSI	0.003	60
9	RNPITPTGH	0.003	
14	PTGHVFQTS	0.003	
7	IMRNPIPT	0.001	
3	YWLPIMRNP	0.001	
16	GHVFQTSIL	0.001	65
2	YWLPIMRN	0.000	
6	PIMRNPIPT	0.000	

163

TABLE VIII-continued

V9-HLA-A1-9mers-24P4C12			
Each peptide is a portion of SEQ ID NO: 19; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.			5
Start	Subsequence	Score	
11	PTQPATLGY	6.250	10
4	MTALYPLPT	0.125	
15	ATLGYVLWA	0.125	
8	YPLPTQPAT	0.050	
5	TALYPLPTQ	0.020	
2	WAMTALYPL	0.020	15
16	TLGYVLWAS	0.010	
6	ALYPLPTQP	0.010	
13	QPATLGYVL	0.005	
17	LGYVLWASN	0.005	
10	LPTQPATLG	0.003	
9	PLPTQPATL	0.002	20
14	PATLGYVLW	0.002	
12	TQPATLGYV	0.002	
3	AMTALYPLP	0.001	
18	GYVLWASNI	0.001	
7	LYPLPTQPA	0.001	
1	YWAMTALYP	0.000	25

TABLE IX

V1-HLA-A1-10mers- 24P4C12			30
Each peptide is a portion of SEQ ID NO: 3; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.			
Start	Subsequence	Score	
594	VTDLLLFFGK	125.000	
32	CTDVICCVLF	25.000	40
120	CPEDPWTVGK	9.000	
518	ILEYIDHKLK	9.000	
680	SLDRPYYSK	5.000	
698	KNEAPPDNKK	4.500	
318	VLEAILLLML	4.500	
488	ISAFIRTLRY	3.750	
39	VLFLLFILGY	2.500	45
262	ILGVVLGVLAY	2.500	
362	FVLLLICIAIY	2.500	
136	VEGVFYTKNR	2.250	
221	IFEDFAQSWY	2.250	
700	EAPPDNKKRK	2.000	
9	DDEAYGKPVK	1.800	50
6	RDEDEAYGK	1.800	
417	SSCPGLMCMVF	1.500	
132	FSQTVGEVFIY	1.500	
134	QTVGEVFIYTK	1.000	
469	FASFYWFPHK	1.000	
369	IAYWAMTALY	1.000	55
378	YLATSGQPQY	1.000	
670	FLEDLERNNG	0.900	
103	VAENGLQCPT	0.900	
277	WEYRVLRLDK	0.900	
242	LSLLFILLLR	0.750	
163	QQELCPSFLL	0.675	
58	YGDPRQVLYP	0.625	60
266	LGVLAYGIYIY	0.625	
348	VGQMMSTMPFIY	0.625	
171	LLPSAPALGR	0.500	
507	LILTLVQIAR	0.500	
237	GVALVLSLFIY	0.500	
320	EAILLLMLFIY	0.500	65
208	LIDSLNARDIY	0.500	

164

TABLE IX-continued

609	GVGVLSFFFF	0.500
353	STMFYPLVTF	0.500
464	VLGAFASFY	0.500
322	ILLMLIFLR	0.500
35	VICCVLFLLF	0.500
606	VVGGVGVLSF	0.500
521	YIDHKLKRGVQ	0.500
662	CVDTLFLCFL	0.500
661	MCVDTLFLCF	0.500
265	VLGVLAYGIY	0.500
49	IVVGIVAWLY	0.500
667	FLCFLEDLER	0.500
407	SCNPTAHLVN	0.500
165	ELCPSFLLPS	0.500
77	MGENKDKPYL	0.450
547	CLEKFIKFLN	0.450
337	AIALLKEASK	0.400
512	VQIARVILEY	0.375
689	KSLLKILGKK	0.300
305	VQETWLAALI	0.270
18	KYDPSFRGPI	0.250
76	GMGENKDKPY	0.250
557	RNAYIMIAIY	0.250
590	VLDKVTDLLE	0.250
677	NNGSLDRPYY	0.250
578	FMLLMRNIVR	0.250
187	VTPPALPGIT	0.250
463	CVLAGAFASF	0.200
516	RVILEYIDHK	0.200
74	YCGMGENDK	0.200
72	GAYCGMGENDK	0.200
423	MCVFQGYSSK	0.200
621	RIPGLGKDFK	0.200
170	FLLPSAPALG	0.200
211	SLNARDISVK	0.200
161	SLQQELCPSF	0.200
253	VAGPLVLVLI	0.200
186	NVTPPALPGI	0.200
618	FSGRIPGLGK	0.150
173	PSAPALGRCF	0.150
118	SSCPEDPWTV	0.150
125	WTVGKNEFSQ	0.125
676	RNNGSLDRPY	0.125
608	GGGVLSFFFF	0.125
286	KGASISQLGF	0.125
80	NKDKPYLLYF	0.125
360	VTFVLLLCI	0.125
196	TNDTTIQQGI	0.125
198	DTTIQQGISG	0.125
293	LGFTTNLSAY	0.125
271	YGIYYCWEEY	0.125
382	SGQPQYVLWA	0.125
467	GAFASFYWAF	0.100
487	LISAFIRTLR	0.100
650	VIASGFFSVF	0.100
64	VLYPRNSTGA	0.100
347	AVGQMMSTMF	0.100
272	GIYYCWEEYR	0.100
333	RIRIAIALLK	0.100
612	VLSFFFFSGR	0.100
147	FCLPGVPWNM	0.100
216	DISVKIFEDF	0.100
53	IVAWLYGDPR	0.100
326	MLIFLRQIR	0.100
544	CLWCLEKFIK	0.100

V3-HLA-A1-10mers-
24P4C12

Each peptide is a portion of SEQ
ID NO: 7; each start position is
specified, the length of peptide is
10 amino acids, and the end
position for each peptide is the
start position plus nine.

Start	Subsequence	Score
10	ITPPALPGIT	0.250
9	NITPPALPGI	0.200
3	RCFPWTNITP	0.050

165

TABLE IX-continued

8	TNITPPALPG	0.013
7	WTNITPPALP	0.005
5	FPWTNITPPA	0.001
2	GRCFPWNTNI	0.001
1	LGRCFPWTNI	0.000
6	PWTNITPPAL	0.000
4	CFPWTNITPP	0.000
V5-HLA-A1-10mers- 24P4C12		
Each peptide is a portion of SEQ ID NO: 11; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.		
Start	Subsequence	Score
2	VLEAILLLVL	4.500
6	ILLVLIFLR	0.500
4	EAILLLVLIF	0.500
8	LLVLIFLRQR	0.100
10	VLIFLRQRIR	0.100
7	LLVLIFLRQ	0.050
1	AVLEAILLLV	0.050
5	AILLLVLIFL	0.050
9	LVLIFLRQRI	0.010
3	LEAILLLVLI	0.001
V6-HLA-A1-10mers- 24P4C12		
Each peptide is a portion of SEQ ID NO: 13; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.		
Start	Subsequence	Score
7	GLIPRSVFNL	0.500
2	GYSSKGLIPR	0.025
6	KGLIPRSVFN	0.005
5	SKGLIPRSVF	0.005
3	YSSKGLIPRS	0.003
10	PRSVFNLQIY	0.003
4	SSKGLIPRSV	0.002
9	IPRSVFNLQI	0.001
1	QGYSSKGLIP	0.001
8	LIPRSVFNLQ	0.001
V7-HLA-A1-10mers- 24P4C12		
Each peptide is a portion of SEQ ID NO: 15; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.		
Start	Subsequence	Score
9	AVGQMMSTMF	0.100
6	ILVAVGQMMS	0.050
7	LVAVGQMMST	0.050
8	VAVGQMMSTM	0.010
5	WILVAVGQMM	0.010
1	QSWYWILVAV	0.003
2	SWYWILVAVG	0.001
4	YWILVAVGQM	0.001
3	WYWILVAVGQ	0.000

166

TABLE IX-continued

V8-HLA-A1-10mers- 24P4C12		
Each peptide is a portion of SEQ ID NO: 17; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.		
Start	Subsequence	Score
1	LNYYWLPIMR	0.125
13	ITPTGHVFQT	0.125
21	QTSILGAYVI	0.050
18	HVFQTSILGA	0.050
11	NPITPTGHVF	0.025
19	VFQTSILGAY	0.025
12	PITPTGHVFQ	0.020
5	WLPIMRNPIT	0.020
4	YWLPIMRNPI	0.005
9	MRNPITPTGH	0.005
20	FQTSILGAYV	0.003
15	PTGHVFQTSI	0.003
14	TPTGHVFQTS	0.003
10	RNPITPTGHV	0.003
2	NYYWLPIMRN	0.003
16	TGHVFQTSIL	0.003
17	GHVFQTSILG	0.003
6	LPIMRNPITP	0.001
8	IMRNPITPTG	0.001
7	PIMRNPITPT	0.000
3	YYWLPIMRNP	0.000
V9-HLA-A1-10mers- 24P4C12		
Each peptide is a portion of SEQ ID NO: 19; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.		
Start	Subsequence	Score
11	LPTQPATLGY	0.625
7	ALYPLPTQPA	0.100
9	YPLPTQPATL	0.050
5	MTALYPLPTQ	0.050
12	PTQPATLGYV	0.025
4	AMTALYPLPT	0.025
16	ATLGYVLWAS	0.025
17	TLGYVLWASN	0.020
15	PATLGYVLWA	0.005
14	QPATLGYVLW	0.005
13	TQPATLGYVL	0.003
18	LGYVLWASNI	0.003
3	WAMTALYPLP	0.002
2	YWAMTALYPL	0.001
10	PLPTQPATLG	0.001
6	TALYPLPTQP	0.001
8	LYPLPTQPAT	0.001
19	GYVLWASNIS	0.001
1	AYWAMTALYP	0.000

167

TABLE X

V1-HLA-A0201-9mers- 24P4C12		
Each peptide is a portion of SEQ ID NO: 3; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.		
Start	Subsequence	Score
449	GLFWTLNWV	3255.381
322	ILLMLIFL	1699.774
580	LLMRNIVRV	1006.209
597	LLFFFGKLL	510.604
544	CLWCLEKFI	476.257
598	LLFFGKLLV	437.482
170	FLLPSAPAL	363.588
86	LLYFNIPSC	360.526
578	FMLLMRNIV	350.529
244	LLFILLRL	309.050
41	FLLFILGYI	292.008
95	ILSSNIISV	271.948
260	VLILGVLGV	271.948
56	WLYGDPQV	204.761
42	LLFILGYIV	179.368
650	VIASGPFVS	179.161
564	AIYGKNFCV	177.497
239	ALVLSLLFI	131.975
604	LLVVGGVGV	118.238
589	VVLDKVTDL	110.872
268	VLAYGIYYC	106.837
456	WVLALGQCV	103.580
537	IMCCFKCCL	99.667
446	GVLGLFWTL	98.554
257	LVLVLILGV	88.043
660	GMCVDTLFL	84.856
686	YMSKSLKI	79.718
177	ALGRCPFWS	77.873
211	SLNARDISV	69.552
107	GLQCPTPQV	69.552
241	VLSLLFILL	69.001
434	LIQRSVFNL	66.613
35	VICCVLFLL	66.613
547	CLEKFIKFL	65.721
317	AVLEAILLL	65.219
240	LVLSSLFIL	64.306
302	YQSVQETWL	54.798
309	WLAALIVLA	52.561
351	MMSTMFYPL	49.834
365	LLICIAIYA	46.451
45	ILGYIVVGI	40.792
638	WLPIMTSIL	40.289
49	IVVGIVAWL	40.197
38	CVLFLLFIL	37.827
148	CLPGVPWNM	37.260
232	ILVALGVAL	36.316
291	SQLGFTTNL	30.453
85	YLLYFNIFS	26.508
506	ALILTLVQI	23.995
252	LVAGPLVLV	23.795
233	LVALGVALV	23.795
525	KLRGVQNPV	18.501
339	ALLKEASKA	18.382
265	VLGVLAYGI	17.736
326	MLIFLRQRI	17.736
340	LLKEASKAV	16.967
445	YGVGLFWT	16.418
315	VLAIVLEAIL	14.890
457	VLAGQCVCL	14.890
509	LTIVQIARV	13.975
119	SCPEDPWTV	13.961
366	LICIAIWAM	13.064
226	AQSWYWILV	11.988
452	WTLNWLVAL	11.615
426	FQGYSSKGL	9.963
554	FLNRNAYIM	9.370
642	MTSILGAYV	9.032
164	QELCPSFLL	8.914
693	KILGKKNEA	8.846

168

TABLE X-continued

251	RLVAGPLVL	8.759
501	SLAFGALIL	8.759
487	LISAFIRTL	8.729
442	LQIYGVGLL	8.469
262	ILGVLGVLA	8.446
521	YIDHKLRGV	8.094
373	AMTALYLAT	8.073
242	LSLLFILLL	7.666
134	QTVGEVFTT	7.594
191	ALPGITNDT	7.452
590	VLDKVTDDL	7.118
362	FVLLLICIA	6.977
200	TIQQGISGL	6.756
83	KPYLLYFNI	6.636
314	IVLAVLEAI	6.471
383	GQPQYVLWA	6.372
225	FAQSWYWIL	6.295
289	SISQLGFTT	5.943
364	LLLICIAYW	5.929
596	DLFFFPGKL	5.564
611	GVLSFFFFS	5.557
282	VLRDKGASI	5.526
154	WNMTVITSL	5.459
380	ATSGQPQYV	5.313
612	VLSFFFFSG	5.305
100	IISVAENGL	4.993
158	VITSLQQEL	4.993
504	FGALILTLV	4.804
536	CIMCCFKCC	4.802
246	FILLRLVLA	4.767
357	YPLVTFVLL	4.510
V3-HLA-A0201-9mers- 24P4C12		
Each peptide is a portion of SEQ ID NO: 7; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.		
Start	Subsequence	Score
6	WTNITPPAL	1.365
9	ITPPALPGI	0.567
2	RCFPWTNIT	0.074
8	NITPPALPG	0.010
4	FPWTNITPP	0.009
1	GRCPFWTNI	0.002
7	TNITPPALP	0.000
5	PWTNITPPA	0.000
3	CFPWTNITP	0.000
V5-HLA-A0201-9mers- 24P4C12		
Each peptide is a portion of SEQ ID NO: 11; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.		
Start	Subsequence	Score
5	ILLVLIFL	1699.774
9	VLIFLRQRI	17.736
1	VLEAILLLV	5
6	LLVLIFLRL	1.251
2	LEAILLLVL	0.666
7	LLVLIFLRQ	0.048
4	AILLVLVIF	0.036
3	EAILLLVLI	0.025
8	LVLIFLRQR	0.014

169

TABLE X-continued

V6-HLA-A0201-9mers- 24P4C12		
Each peptide is a portion of SEQ ID NO: 13; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.		
Start	Subsequence	Score
7	LIPRSVFNL	66.613
6	GLIPRSVFN	0.410
4	SKGLIPRSV	0.019
5	KGLIPRSVF	0.003
2	YSSKGLIPR	0.001
9	PRSVFNLQI	0.000
3	SSKGLIPRS	0.000
8	IPRSVFNLQ	0.000
1	GYSSKGLIP	0.000
V7-HLA-A0201-9mers- 24P4C12		
Each peptide is a portion of SEQ ID NO: 15; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.		
Start	Subsequence	Score
5	ILVAVGQMM	8.446
4	WILVAVGQM	3.476
8	AVGQMMSTM	1.000
7	VAVGQMMST	0.405
1	SWYWILVAV	0.071
6	LVAVGQMS	0.011
2	WYWILVAVG	0.000
3	YWILVAVGQ	0.000
V8-HLA-A0201-9mers- 24P4C12		
Each peptide is a portion of SEQ ID NO: 17; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.		
Start	Subsequence	Score
4	WLPIMRNPI	47.991
20	QTSILGAYV	5.313
7	IMRNPITPT	1.599
13	TPTGHVFQT	0.649
15	TGHVFQTSI	0.259
10	NPITPTGHV	0.059
5	LPIMRNPI	0.034
18	VFQTSILGA	0.013
19	FQTSILGAY	0.010
16	GHVFQTSIL	0.006
12	ITPTGHVFQ	0.002
2	YYWLPIMRN	0.001
17	HVFQTSILG	0.001
9	RNPITPTGH	0.000
6	PIMRNPI	0.000
11	PITPTGHVF	0.000
14	PTGHVFQTS	0.000
8	MRNPITPTG	0.000
3	YWLPIMRNP	0.000
1	NYWLPIMR	0.000

170

TABLE X-continued

V9-HLA-A0201-9mers- 24P4C12		
Each peptide is a portion of SEQ ID NO: 19; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.		
Start	Subsequence	Score
2	WAMTALYPL	11.615
12	TQPATLGYV	11.597
15	ATLGYVLWA	3.230
16	TLGYVLWAS	1.285
8	YPLPTQPAT	0.828
9	PLPTQPATL	0.470
4	MTALYPLPT	0.176
13	QPATLGYVL	0.057
6	ALYPLPTQP	0.048
3	AMTALYPLP	0.016
17	LGYYVLWASN	0.004
5	TALYPLPTQ	0.002
18	GYVLWASNI	0.001
7	LYPLPTQPA	0.001
10	LPTQPATLG	0.001
1	YWAMTALYP	0.000
14	PATLGYVLW	0.000
11	PTQPATLGY	0.000

TABLE XI

V1-HLA-A0201-10mers- 24P4C12		
Each peptide is a portion of SEQ ID NO: 3; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.		
Start	Subsequence	Score
354	TMFYPLVTFV	2351.109
85	YLLYFNIFSC	1127.969
579	MLLMRNIVRV	1006.209
603	KLLVVGGVGV	900.698
309	WLAALIVLAV	735.860
351	MMSTMFYPLV	486.748
41	FLLFILGYIV	485.348
641	IMTSILGAYV	469.669
546	WCLEKFIKFL	467.771
597	LLFFFGKLLV	437.482
598	LLFFGKLLVV	412.546
665	TLFLCFLEDL	338.500
241	VLSLLFILL	317.403
649	YVIASGFFSV	308.501
433	GLIQRSVFNL	284.974
508	ILTLVQIARV	271.948
232	ILVALGVALV	271.948
42	LLFILGYIVV	269.051
339	ALLKEASKAV	257.342
449	GLFWTLNWWL	243.051
244	LLFILLRLV	201.242
243	SLLFILLRL	181.794
364	LLLCIAYWA	171.868
48	YIVVGIVAWL	170.923
251	RLVAGPLVLV	159.970
321	AILLMLIFL	137.482
56	WLYGDPRQVL	128.926
239	ALVLSLLFIL	116.840
350	QMMSTMFYPL	108.462
86	LLYFNIFSCI	107.833
365	LLLCIAYWAM	95.013
259	LVLILGVLVG	88.043
162	LQQELCPSFL	83.030
580	LLMRNIVRVV	82.509
94	CILSSNIISV	81.385

171

TABLE XI-continued

517	VILEYIDHKL	75.751	
554	FLNRNAYIMI	71.986	
686	YMSKSLKIL	66.925	
44	FILGYIVVGI	56.155	5
133	SQTVGEVFT	55.435	
438	SVFNLQIYGV	51.790	
231	WILVALGVVAL	49.993	
235	ALGVALVLSL	49.134	
441	NLQIYGVGL	49.134	
660	GMCVDTLFLC	47.864	10
325	LMILIFLRQRI	47.394	
536	CIMCCFKCCL	41.299	
315	VLAVLEAILL	36.316	
448	LGLFWTLNWW	36.126	
662	CVDTLFLCFL	35.941	
64	VLYPRNSTGA	27.026	
589	VVLDDKVTDL	23.620	15
596	DLLFFFGKLL	22.527	
240	LVLSSLFILL	22.339	
357	YPLVTFVLL	20.744	
267	GVLAYGIYYC	20.346	
304	SVQETWLAAL	17.627	20
248	LLRLRVAGPL	17.468	
302	YQSVQETWLA	17.378	
501	SLAFGALILT	17.140	
317	AVLEAILLLM	15.167	
590	VLDKVTDL	14.526	
45	ILGYIVVGIV	14.495	
659	FGMCVDTLFL	13.054	25
456	WVLALGQCVL	13.044	
148	CLPGVPWNMT	12.668	
108	LQCPTPQVCV	11.988	
478	KPQDIPTFPL	11.606	
238	VALVLSLLFI	11.529	
312	ALIVLAVLEA	11.426	30
459	ALGQCVLAGA	11.426	
571	CVSAKNAFML	10.841	
563	IAIYGNKFCV	9.525	
445	YGVGLFWTL	9.141	
379	LATSGQPQYV	9.032	
327	LIFLRQIRI	9.023	35
249	LLRLVAGPLV	8.986	
539	CCPKCCLWCL	8.900	
513	QIARVILEYI	8.892	
510	TLVQIARVIL	8.759	
457	VVALGQCVLA	8.446	
95	ILSSNIISVA	7.964	
657	SVFGMCVDTL	7.794	40
225	FAQSWYWL	7.554	
588	VVLDKVTDL	7.309	
593	KVTDLLFFFG	6.865	
368	CIAYWAMTAL	6.756	
562	MIAYGNKFC	6.387	
363	VLLLCIAYW	5.929	45
36	ICCVLFLFI	5.565	
318	VLEAILLLML	5.346	
292	QLGFTTNLSA	4.968	
314	IVLAVLEAIL	4.821	
393	NISSPGCEKV	4.686	
506	ALILTLVQIA	4.685	50
260	VLILGVGLV	4.452	
604	LLVVGVGVL	4.452	
261	LILGVGLVLA	4.297	
502	LAFGALITL	4.292	
147	FCLPGVPWNM	4.140	
V3-HLA-A0201-10mers- 24P4C12			55
Each peptide is a portion of SEQ ID NO: 7; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.			
Start	Subsequence	Score	
9	NITPPALPGI	3.299	
5	FPWTNITPPA	1.238	
1	LGRCFPWTNI	0.015	60
10	ITPPALPGIT	0.009	

172

TABLE XI-continued

7	WTNITPPALP	0.001	
8	TNITPPALPG	0.000	
2	GRCFPWTNIT	0.000	
3	RCFPWTNITP	0.000	
6	PWTNITPPAL	0.000	
4	CFPWTNITPP	0.000	
V5-HLA-A0201-10mers- 24P4C12			10
Each peptide is a portion of SEQ ID NO: 11; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.			
Start	Subsequence	Score	
1	AVLEAILLLV	212.340	
5	AILLVLIFL	137.482	
9	LVLIFLRQRI	5.742	20
2	VLEAILLLVL	2.192	
6	ILLVLIFLR	1.251	
3	LEAILLLVLI	0.793	
7	LLVLIFLRQ	0.178	
8	LLVLIFLRQR	0.044	25
10	VLIFLRQRIR	0.002	
4	EAILLLVLIF	0.000	
V6-HLA-A0201-10mers- 24P4C12			30
Each peptide is a portion of SEQ ID NO: 13; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.			
Start	Subsequence	Score	
7	GLIPRSVFNL	284.974	
6	KGLIPRSVFN	0.035	
9	IPRSVFNLQI	0.033	
8	LIPRSVFNLQ	0.007	40
3	YSSKGLIPRS	0.005	
4	SSKGLIPRSV	0.003	
1	QGYSSKGLIP	0.000	
5	SKGLIPRSVF	0.000	
2	GYSSKGLIPR	0.000	45
10	PRSVFNLQIY	0.000	
V7-HLA-A0201-10mers- 24P4C12			50
Each peptide is a portion of SEQ ID NO: 15; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.			
Start	Subsequence	Score	
5	WILVAVGQMM	11.626	
1	QSWYWILVAV	8.667	
7	LAVVGQMMST	2.550	60
8	VAVGQMMSTM	0.270	
6	ILVAVGQMMS	0.127	
9	AVGQMMSTMF	0.007	
4	YWILVAVGQM	0.001	
3	WYWILVAVGQ	0.000	65
2	SWYWILVAVG	0.000	

173

TABLE XI-continued

V8-HLA-A0201-10mers- 24P4C12		
Each peptide is a portion of SEQ ID NO: 17; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.		
Start	Subsequence	Score
20	FQTSILGAYV	178.411
5	WLPIMRNPIT	14.054
13	ITPTGHVFQT	2.347
7	PIMRNPITPT	0.192
18	HVFQTSILGA	0.126
21	QTSILGAYVI	0.059
10	RNPITPTGHV	0.059
16	TGHVFQTSIL	0.057
4	YWLPIMRNPI	0.025
15	PTGHVFQTSI	0.012
8	IMRNPITPTG	0.007
14	TPTGHVFQTS	0.001
1	LNYYWLPIMR	0.001
12	PITPTGHVFQ	0.000
11	NPITPTGHVF	0.000
6	LPIMRNPITP	0.000
2	NYWLPIMRN	0.000
17	GHVFQTSILG	0.000
3	YYWLPIMRNP	0.000
19	VFQTSILGAY	0.000
9	MRNPITPTGH	0.000
V9-HLA-A0201-10mers- 24P4C12		
Each peptide is a portion of SEQ ID NO: 19; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.		
Start	Subsequence	Score
7	ALYPLPTQPA	15.898
4	AMTALYPLPT	5.382
9	YPLPTQPATL	2.373
13	TQPATLGTVL	0.888
18	LGYVLWASNI	0.370
17	TLGYVLWASN	0.127
16	ATLGTVLWAS	0.066
12	PTQPATLGTV	0.035
2	YWAMTALYPL	0.031
15	PATLGTVLWA	0.019
3	WAMTALYPLP	0.005
8	LYPLPTQPAT	0.002
10	PLPTQPATLG	0.002
11	LPTQPATLGY	0.001
5	MTALYPLPTQ	0.001
6	TALYPLPTQP	0.001
14	QPATLGTVLV	0.001
1	AYWAMTALYP	0.000
19	GYVLWASNIS	0.000

TABLE XII

V1-HLA-A3-9mers- 24P4C12		
Each peptide is a portion of SEQ ID NO: 3; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.		
Start	Subsequence	Score
421	GLMCVFQGY	81.000
135	TVGEVPYTK	40.500

174

TABLE XII-continued

207	GLIDSLNAR	27.000
323	LLLMLIFLR	27.000
243	SLLFILLLR	27.000
354	TMFYPLVTF	22.500
690	SLLKILGKK	20.250
517	VILEYIDHK	20.250
363	VLLLCIAY	18.000
585	IVRVVLDK	18.000
560	YIMAIYVK	13.500
508	ILTLVQIAR	12.000
579	MLLMRNIVR	12.000
267	GVLAYGIYY	10.800
424	CVFQGYSSK	10.000
244	LLFILLRL	9.000
464	VLAGAFASF	9.000
272	GIYYCWEEY	6.000
351	MMSTMFYPL	5.400
470	ASFYWAFFK	4.500
449	GLFWTLNWX	4.500
86	LLYFNIFSC	4.500
446	GVGLGLFWTL	3.645
660	GMCVDTLFL	3.600
633	HLNYYWLP	3.600
542	KCCLCLEK	3.600
241	VLSLLFILL	3.600
42	LLFILGYIV	3.000
393	NISSPGCEK	3.000
325	LMLIFLRQR	2.700
45	ILGYIVVGI	2.700
322	ILLMLIFL	2.700
239	ALVLSLLFI	2.700
641	IMTSILGAY	2.700
598	LLFFGKLLV	2.000
260	VLILGLVGV	1.800
265	VLGVLAYGI	1.800
513	QIARVILEY	1.800
609	GVGVLSFFF	1.800
537	IMCCFKCCL	1.800
50	VVGIVAWLY	1.800
686	YMSKSLKKI	1.800
251	RLVAGPLVL	1.800
593	KVTDLLFFF	1.800
358	PLVTFVLLL	1.620
544	CLWCLEKFI	1.500
689	KSLLKILGK	1.350
525	KLRGVQNPV	1.350
170	FLLPSAPAL	1.350
547	CLEKFIKFL	1.350
597	LLLFFGKLL	1.350
365	LLICIAIWA	1.350
506	ALILTLVQI	1.350
148	CLPGVPWNM	1.350
501	SLAFGALIL	1.200
662	CVDTLFLCF	1.200
349	GQMMSTMFY	1.080
443	QIYGVGLGF	1.012
321	AILLLMLIF	0.900
590	VLDKVTDLL	0.900
326	MLIFLRQRI	0.900
268	VLAYGIYYC	0.900
107	GLQCPTPOV	0.900
613	LSFFFFSGR	0.900
318	VLEAILLLM	0.900
232	ILVALGVAL	0.900
518	ILEYIDHKL	0.900
452	WTLNWLAL	0.810
596	DLLFFGKLL	0.729
645	ILGAYVIAS	0.720
258	VLVLILGVL	0.608
49	IVVGIVAWL	0.608
41	FLLFILGYI	0.608
54	VAWLYGDPR	0.600
665	TLFLCFLED	0.600
95	ILSSNIISV	0.600
457	VLAGQCQVL	0.600
282	VLRDKGASI	0.600
554	FLNRNAYIM	0.600
39	VLFLFLILG	0.600
315	VLAVLEAIL	0.600
638	WLPIMTSIL	0.600

175

TABLE XII-continued

434	LIQRSVFNL	0.540
612	VLSFFFFSG	0.540
611	GVLSFFFFS	0.486
647	GAYVIASGF	0.450
580	LLMRNIVRV	0.450
364	LLLICIAYW	0.450
564	AIYGKNFCV	0.450
237	GVALVLSLL	0.405
38	CVLFLLFIL	0.405
204	GISGLIDSL	0.405
35	VICCVLFLL	0.405
317	AVLEAILL	0.405
240	LVLVLLFIL	0.405
668	LCFLEDLER	0.400
388	VLWASNISS	0.400
489	SAFIRTLRY	0.400
211	SLNARDISV	0.400
85	YLLYFNIFS	0.360

V3-HLA-A3-9mers-
24P4C12

Each peptide is a portion of SEQ
ID NO: 7; each start position is
specified, the length of peptide is
9 amino acids, and the end
position for each peptide is the
start position plus eight.

Start	Subsequence	Score
9	ITPPALPGI	0.068
6	WTNITPPAL	0.030
2	RCFPWTNIT	0.022
8	NITPPALPG	0.009
1	GRCFPWTNI	0.003
4	FPWTNITPP	0.002
7	TNITPPALP	0.000
3	CFPWTNITP	0.000
5	PWTNITPPA	0.000

V5-HLA-A3-9mers-
24P4C12

Each peptide is a portion of SEQ
ID NO: 11; each start position is
specified, the length of peptide is
9 amino acids, and the end
position for each peptide is the
start position plus eight.

Start	Subsequence	Score
6	LLLVLIFLR	27.000
5	ILLVLIFL	4.050
4	AILLVLIF	1.800
9	VLIFLRQRI	0.900
1	VLEAILLLV	0.900
7	LLVLIFLRQ	0.270
8	LVLIFLRQR	0.270
2	LEAILLVL	0.005
3	EAILLLVLI	0.004

V6-HLA-A3-9mers-
24P4C12

Each peptide is a portion of SEQ
ID NO: 13; each start position is
specified, the length of peptide is
9 amino acids, and the end
position for each peptide is the
start position plus eight.

Start	Subsequence	Score
7	LIPRSVFNL	0.540
6	GLIPRSVFN	0.135
2	YSSKGLIPR	0.060
5	KGLIPRSVF	0.013
8	IPRSVFNLQ	0.001
3	SSKGLIPRS	0.000
9	PRSVFNLQI	0.000

176

TABLE XII-continued

1	GYSSKGLIP	0.000
4	SKGLIPRSV	0.000

V7-HLA-A3-9mers-
24P4C12

Each peptide is a portion of SEQ
ID NO: 15; each start position is
specified, the length of peptide is
9 amino acids, and the end
position for each peptide is the
start position plus eight.

Start	Subsequence	Score
5	ILVAVGQMM	0.450
8	AVGQMMSTM	0.030
4	WILVAVGQM	0.027
6	LVAVGQMMS	0.008
7	VAVGQMMST	0.007
1	SWYWILVAV	0.002
2	WYWILVAVG	0.000
3	YWILVAVGQ	0.000

V8-HLA-A3-9mers-
24P4C12

Each peptide is a portion of SEQ
ID NO: 17; each start position is
specified, the length of peptide is
9 amino acids, and the end
position for each peptide is the
start position plus eight.

Start	Subsequence	Score
4	WLPIMRNPI	0.600
7	IMRNPITPT	0.225
19	FQTSILGAY	0.081
1	NYWLPIMR	0.040
11	PITPTGHVF	0.030
17	HVFQTSILG	0.020
13	TPTGHVFQT	0.013
20	QTSILGAYV	0.010
16	GHVFQTSIL	0.003
15	TGHVFQTSI	0.002
5	LPIMRNPIT	0.002
10	NPITPTGHV	0.001
12	ITPTGHVFQ	0.001
14	PTGHVFPQS	0.001
18	VFQTSILGA	0.001
6	PIMRNPITP	0.001
2	YYWLPIMRN	0.000
9	RNPITPTGH	0.000
8	MRNPITPTG	0.000
3	YWLPIMRNP	0.000

V9-HLA-A3-9mers-
24P4C12

Each peptide is a portion of SEQ
ID NO: 19; each start position is
specified, the length of peptide is
9 amino acids, and the end
position for each peptide is the
start position plus eight.

Start	Subsequence	Score
15	ATLGYVLWA	0.405
16	TLGYVLWAS	0.270
6	ALYPLPTQP	0.150
11	PTQPATLGY	0.060
9	PLPTQPATL	0.060
2	WAMTALYPL	0.041
4	MTALYPLPT	0.030
3	AMTALYPLP	0.020
13	QPATLGYVL	0.018
18	GYVLWASNI	0.008
12	TQPATLGYV	0.003
8	YPLPTQPAT	0.002
5	TALYPLPTQ	0.001
7	LYPLPTQPA	0.000

177

TABLE XII-continued

10	LPTQPATLG	0.000	
14	PATLGVVLW	0.000	
17	LGTVLWASN	0.000	
1	YWAMTALYP	0.000	5

TABLE XIII

V1-HLA-A3-10mers-
24P4C12

Each peptide is a portion of SEQ
ID NO: 3; each start position is
specified, the length of peptide is
10 amino acids, and the end
position for each peptide is the
start position plus nine.

Start	Subsequence	Score	
544	CLWCLEKFIK	300.000	
39	VLFLLLFILGY	180.000	
680	SLDRPYYSK	120.000	
612	VLSFFFFSGR	36.000	
134	QTVGEVFTK	30.375	
211	SLNARDISVK	30.000	
449	GLFWTLNWL	27.000	
322	ILLMLIFLR	27.000	
584	NIVRVVLDK	27.000	
433	GLIQRSVFNL	24.300	
262	ILGVLGVLAY	24.000	
272	GIYYCWEEYR	18.000	
464	VLGAFASFY	18.000	
665	TLFLCFLEDL	13.500	
516	RVILEYIDHK	13.500	
86	LLYFNIFSCI	13.500	
171	LLPSAPALGR	12.000	
578	FMLLMRNIVR	12.000	
76	GMGENDKPY	9.000	
594	VTDLFFFSGK	9.000	
350	QMMSTMFYPL	8.100	
667	FLCFLEDLER	8.000	
56	WLYGDPQVL	6.750	
333	RIRIAIALLK	6.000	
609	GVGVLSFFFF	5.400	
241	VLSLLFILL	5.400	
561	IMIAIYGKNF	4.500	
239	ALVLSLLFIL	4.050	
49	IVVGIVAWLY	4.050	
378	YLATSGQPQY	4.000	
441	NLQIYGVGL	3.600	
235	ALGVALVLSL	3.600	
598	LLFFGKLLV	3.000	
621	RIPGLGKDFK	3.000	
354	TMFYPLVTFV	3.000	
72	GAYCGMGENK	3.000	
324	LLMLIFLRQR	2.700	
660	GMCVDTLFLC	2.700	
467	GAFASFYWAF	2.700	
243	SLLFILLRL	2.700	
83	KPYLLYFNIF	2.700	
42	LLFILGYIVV	2.000	
518	ILEYIDHKLR	2.000	
161	SLQQELCPSP	2.000	
337	AIALKEASK	2.000	
362	FVLLICIAY	1.800	
650	VIASGFFSVF	1.800	
606	VVGGVGVLSF	1.800	
507	LILTLVQIAR	1.800	
329	FLRQRIRIAI	1.800	
318	VLEAILLLML	1.800	
624	GLGKDFKSPH	1.800	
309	WLAALIVLAV	1.800	
312	ALIVLAVLEA	1.800	
469	FASFYWAFHK	1.800	
64	VLYPRNSTGA	1.500	
364	LLLICIAIYWA	1.350	
657	SVFGMCDVTL	1.350	
85	YLLYFNIFSC	1.350	

178

TABLE XIII-continued

220	KIPEDFAQSW	1.350	
264	GVLGVLAYGI	1.215	
315	VLAVLEAILL	1.200	
237	GVALVLSLFL	1.200	
554	FLNRNAYIMI	1.200	
590	VLDKVTDLLL	1.200	
265	VLGVLAYGIY	1.200	
35	VICCVLFLFL	1.200	
53	IVAWLYGDP	1.200	
447	VLGLFWTLNW	1.200	
268	VLAYGIYYCW	0.900	
413	HLVNSSCPGL	0.900	
275	YCWEEYRVLR	0.900	
232	ILVALGVALV	0.900	
325	LMLIFLRQRI	0.900	
463	CVLAGAFASF	0.900	
525	KLRGVQNPVA	0.900	
506	ALILTLVQIA	0.900	
603	KLLVVGGVGV	0.900	
633	HLNYWLPIM	0.900	
510	TLVQIARVIL	0.900	
365	LLICIAYWAM	0.900	
41	FLLFILGYIV	0.900	
512	VQIARVILEY	0.810	
604	LLVVGGVGV	0.810	
251	RLVAGPLVLV	0.675	
260	VLILGVLVGL	0.608	
44	FILGYIVVGI	0.608	
107	GLQCPTPQVC	0.600	
327	LIFLRQIRIR	0.600	
326	MLIFLRQIRIR	0.600	
597	LLFFFGKLLV	0.600	
487	LISAFIRTLR	0.600	
120	CPEDPWTVGK	0.600	
351	MMSTMFYPLV	0.600	
240	LVLSSLFILL	0.540	
252	LVAGPLVLVL	0.540	
360	VTFVLLLCI	0.450	
363	VLLLCIAIYW	0.450	
579	MLLMRNIVRV	0.450	
95	ILSSNIISVA	0.450	

Start	Subsequence	Score	
9	NITPPALPGI	0.135	
5	FPWNTITPPA	0.015	
3	RCFPWTNITP	0.003	
10	ITPPALPGIT	0.002	
7	WTNITPPALP	0.002	
1	LGRCFPWTNI	0.001	
2	GRCFPWTNIT	0.001	
8	TNITPPALPG	0.000	
6	PWTNITPPAL	0.000	
4	CFPWTNITPP	0.000	

Start	Subsequence	Score	
6	ILLVLIFLRL	27.000	
8	LLVLIFLRQR	2.700	
2	VLEAILLLVL	1.800	
10	VLIFLRQIRIR	0.600	
5	AILLLVLIFL	0.405	

Figure 2F-2

100 I I S V A E N G L Q C P T P Q V C V S S
301 ACATCATCTCAGTTGCTGAGAACGGCTACAGTGCCCCACACCCAGGTGTGTGTGTCTCT
120 C P E D P W T V G K N E F S Q T V G E V
361 CCTGCCCCGAGGACCCATGGACTGTGGGAAAAACGAGTTCTCACAGACIGTTGGGGAAG
140 F Y T K N R N F C L P G V P W N M T V I
421 TCTTCTATACAAAAACAGGAACTTTGTCTGCCAGGGGTACCCTGGAAIATGACGGTGA
160 T S L Q Q E L C P S F L L P S A P A L G
481 TCACAAGCCTGCAACAGGAACCTGCCCCAGTTTCTCTCCCTCTGCTCCAGCTCTGG
180 R C F P W T N V T P P A L P G I T N D T
541 GCGCTGCTTTCCATGGACCAACGTTACTCCACCGGCGTCCCAGGGATCACCATGACA
200 T I Q Q G I S G L I D S L N A R D I S V
601 CCACCATACAGCAGGGGATCAGCGGTCTTATTGACAGCCTCAATGCCCGAGACATCAGTG
220 K I F E D F A Q S W Y W I L V A L G V A
661 TTAAGATCTTTGAAGATTTTGCCAGTCCTGGTATTGGATTCTTGTGCCCCGGGGGTGG
240 L V L S L L F I L L L R L V A G P L V L
721 CTCGGTCTTGAGCCTACIGTTTATCTTGCTTCGCGCCTGGTGGCTGGGCCCTGGTGC
260 V L I L G V L G V L A Y G I Y Y C W E E
781 TGGTGCTGATCCTGGGAGTGCTGGGCGTGCTGGCATAACGGCATCTACTACTGCTGGGAGG
280 Y R V L R D K G A S I S Q L G F T T N L
841 AGTACCGAGTGCTGCGGACAAGGGCGCCTCCATCTCCCAGCTGGGTTCACCACCAACC
300 S A Y Q S V Q E T W L A A L I V L A V L
901 TCAGTGCCTACCAGAGCGTGCAGGAGACCTGGCTGGCCGCCCTGATCGTGTGGCGGTGC
320 E A I L L L M L I F L R Q R I R I A I A
961 TTGAAGCCATCCTGCTGCTGATGCTCATCTTCTGCGGCAGCGGATTCTGATTGCCATCG
340 L L K E A S K A V G Q M M S T M F Y F L
1021 CCCTCCTGAAGGAGGCCAGCAAGGCTGTGGGACAGATGATGTCTACCATGTTCTACCCAC
360 V T F V L L L I C I A Y W A M T A L Y L
1081 TGGTCACCTTTGTCTCTCTCTCATCTGCATTGCCTACTGGGCCATGACTGCTCTGTACC
380 A T S G Q P Q Y V L W A S N I S S P G C
1141 TGGCTACATCGGGGCAACCCAGTATGTGCTCTGGGCATCCAACATCAGCTCCCCCGGCT
400 E K V P I N T S C N P T A R L V N S S C
1201 GTGAGAAAGTGCCAATAAATACATCAAGCAACCCACGGCCACCTTGGAACCTCTCGT
420 P G L M C V F Q G Y S S K G L I P R S V
1261 GCCCAGGGCTGATGTGCGTCTTCCAGGGCTACTCATCCAAAGGCCTAATCCCACGTTCTG
440 F N L Q I Y G V L G L F W T L N W V L A
1321 TCTTCAATCTGCAAATCTATGGGGTCTGGGGCTCTTCTGGACCCTTAACTGGGTACTGG
460 L G Q C V L A G A F A S F Y W A F R R P
1381 CCCTGGGCCAATGCGTCTCGCTGGAGCCTTTGCCTCCTTCTACTGGGCCTTCCACAAGC
480 Q D I P T F P L I S A F I R T L R Y H T
1441 CCCAGGACATCCCTACCTTCCCTTAATCTCTGCCTTCATCCGCACACTCCGTTACCACA

179

TABLE XIII-continued

7	LLLVLIFLRQ	0.270
1	AVLEAILLLV	0.203
9	LVLIFLRQRI	0.090
4	EAILLLVLIF	0.054
3	LEAILLLVLI	0.003
V6-HLA-A3-10mers- 24P4C12		
Each peptide is a portion of SEQ ID NO: 13; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.		
Start	Subsequence	Score
7	GLIPRSVFNL	36.450
2	GYSSKGLIPR	0.036
9	IPRSVFNLQI	0.036
8	LIPRSVFNLQ	0.009
5	SKGLIPRSVF	0.003
10	PRSVFNLQIY	0.001
3	YSSKGLIPRS	0.000
4	YSSKGLIPRSV	0.000
1	QGYSSKGLIP	0.000
6	KGLIPRSVFN	0.000
V7-HLA-A3-10mers- 24P4C12		
Each peptide is a portion of SEQ ID NO: 15; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.		
Start	Subsequence	Score
9	AVGQMMSTMF	0.200
6	ILVAVGQMMS	0.120
5	WILVAVGQMM	0.045
7	LVAVGQMMST	0.030
1	QSWYWILVAV	0.011
8	VAVGQMMSTM	0.007
2	SWYWILVAVG	0.000
4	YWILVAVGQM	0.000
3	WYWILVAVGQ	0.000
V8-HLA-A3-10mers- 24P4C12		
Each peptide is a portion of SEQ ID NO: 17; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.		
Start	Subsequence	Score
18	HVFQTSILGA	0.300
5	WLPIMRNPI	0.100
21	QTSILGAYVI	0.090
1	LNYYWLPIMR	0.080
13	ITPTGHVFQT	0.045
11	NPITPTGHVF	0.030
8	IMRNPITPTG	0.030
15	PTGHVFQTSI	0.009
20	FQTSILGAYV	0.006
7	PIMRNPIPT	0.003
14	TPTGHVFQTS	0.003
19	VFQTSILGAY	0.003
4	YWLPIMRNPI	0.001
6	LPIMRNPIPT	0.001
16	TGHVFQTSIL	0.001
2	NYWLPIMRN	0.000
9	MRNPITPTGH	0.000
12	PITPTGHVFQ	0.000
17	GHVFQTSILG	0.000

180

TABLE XIII-continued

10	RNPITPTGHV	0.000
3	YYWLPIMRN	0.000
V9-HLA-A3-10mers- 24P4C12		
Each peptide is a portion of SEQ ID NO: 19; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.		
Start	Subsequence	Score
7	ALYPLPTQPA	2.250
4	AMTALYPLPT	0.600
11	LPTQPATLGY	0.080
13	TQPATLGYVL	0.054
16	ATLGYVLWAS	0.030
17	TLGYVLWASN	0.020
9	YPLPTQPATL	0.013
18	LGTVLWASNI	0.009
15	PATLGYVLWA	0.004
10	PLPTQPATLG	0.003
2	YWAMTALYPL	0.003
5	MTALYPLPTQ	0.002
14	QPATLGYVLW	0.002
12	PTQPATLGYV	0.001
6	TALYPLPTQP	0.000
3	WAMTALYPLP	0.000
1	AYWAMTALYP	0.000
19	GYVLWASNIS	0.000
8	LYPLPTQPAT	0.000

TABLE XIV

V1-HLA-A1101-9mers- 24P4C12		
Each peptide is a portion of SEQ ID NO: 3; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus nine.		
Start	Subsequence	Score
135	TVGEVFTYTK	4.000
585	IVRVVVLDK	4.000
424	CVFQGYSSK	4.000
560	YIMIAIYKG	1.600
685	YYMSKSLK	1.600
542	KCCLWCLEK	1.200
690	SLLKILGKK	0.600
517	VILEYIDHK	0.600
73	AYCGMGENK	0.400
393	NISSPGCEK	0.400
207	GLIDSLNAR	0.360
323	LLMLIFLR	0.360
338	IALLKEASK	0.300
579	MLLMRNIVR	0.240
243	SLLFILLR	0.240
622	IPGLGKDFK	0.200
689	KSLKILGK	0.180
516	RVILEYIDH	0.180
609	GVGVLSFFF	0.180
485	FPLISAFIR	0.180
446	GVGLFWTL	0.180
267	GVLAYGIYY	0.180
273	IYYCWEEYR	0.160
508	ILTLVQIAR	0.160
668	LCFLEDLER	0.160
698	KNEAPPDNK	0.120
470	ASFYWAFHK	0.120
593	KVTDLLFFF	0.120
701	APPDNKKRK	0.100
595	TDLLFFGK	0.090

181

TABLE XIV-continued

38	CVLFLLFIL	0.090
240	LVLSSLFIL	0.090
54	VAWLYGDPR	0.080
172	LPSAPALGR	0.080
349	GQMMSTMFY	0.072
334	IRIAIALLK	0.060
545	LWCLEKFIK	0.060
567	GKNFCVSAK	0.060
317	AVLEAIIILL	0.060
699	NEAPPDNKK	0.060
151	GVPWNMTVI	0.060
237	GVALVLSLL	0.060
257	LVLVLILGV	0.060
20	DPSFRGPIK	0.060
575	KNAFMLLMR	0.048
212	LNARDISVK	0.040
359	LVTFVLLLI	0.040
16	PVKYDPSFR	0.040
304	SVQETWLAA	0.040
619	SGRIPGLGK	0.040
50	VVGIVAWLY	0.040
681	LDRPYYSK	0.040
662	CVDTLFLCF	0.040
7	DEDDAYGK	0.036
83	KPYLLYFNI	0.036
47	GYIVVGIVA	0.036
251	RLVAGPLVL	0.036
383	GQPQYVLWA	0.036
49	IVVGIVAWL	0.030
314	IVLAVLEAI	0.030
456	WVLALGQCV	0.030
589	VVLDKVTDL	0.030
452	WTLNWWLAL	0.030
141	YTKNRNFCL	0.030
498	HTGSLAFGA	0.030
605	LVVGGVGV	0.030
362	FVLLLICIA	0.030
611	GVLSFFFFS	0.027
137	GEVFTKNR	0.027
564	AIYGKNFCV	0.024
272	GIYYCWEEY	0.024
60	DPRQVLYPR	0.024
421	GLMCVFQGY	0.024
467	GAFASFYWA	0.024
449	GLFWTLNVV	0.024
660	GMCVDTLFL	0.024
496	RYHTGSLAF	0.024
511	LVQIARVIL	0.020
218	SVKIFEDFA	0.020
233	LVALGVALV	0.020
22	SFRGPIKNR	0.020
75	CGMGENKDK	0.020
414	LVNSSCPGL	0.020
252	LVAGPLVLV	0.020
571	CVSAKNAFM	0.020
347	AVGQMMSTM	0.020
534	ARCIMCCFK	0.020
527	RGVQNPVAR	0.018
34	DVICCVLFL	0.018
693	KILGKKNEA	0.018
461	GQCVLAGAF	0.018
4	KQRDEDEDA	0.018
331	RQRIRIAIA	0.018
10	DEAYGKPVK	0.018
442	LQIYGVGL	0.018
255	GPLVLVLIL	0.018
598	LLFFGKLLV	0.016
42	LLFILGYIV	0.016
244	LLFILLRL	0.016
327	LIFLRQRIR	0.016

182

TABLE XIV-continued

V3-HLA-A1101-9mers- 24P4C12		
Each peptide is a portion of SEQ ID NO: 7; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.		
Start	Subsequence	Score
9	ITPPALPGI	0.010
6	WTNITPPAL	0.010
2	RCFPWTNIT	0.001
8	NITPPALPG	0.001
1	GRCFPWTNI	0.001
4	FPWTNITPP	0.000
3	CFPWTNITP	0.000
7	TNITPPALP	0.000
5	PWTNITPPA	0.000
V5-HLA-A1101-9mers- 24P4C12		
Each peptide is a portion of SEQ ID NO: 11; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.		
Start	Subsequence	Score
6	LLLVLIFLR	0.360
8	LVLIFLRQR	0.060
4	AILLLVLIF	0.012
5	ILLVLIFLR	0.012
1	VLEAIIILLV	0.008
9	VLIFLRQRI	0.006
7	LLVLIFLRQ	0.001
2	LEAIIILLVL	0.001
3	EATILLVLI	0.001
V6-HLA-A1101-9mers- 24P4C12		
Each peptide is a portion of SEQ ID NO: 13; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.		
Start	Subsequence	Score
7	LIPRSVFNL	0.012
2	YSSKGLIPR	0.008
1	GYSSKGLIP	0.002
6	GLIPRSVFN	0.002
5	KGLIPRSVF	0.001
8	IPRSVFNLQ	0.000
9	PRSVFNLQI	0.000
4	SKGLIPRSV	0.000
3	SSKGLIPRS	0.000
V7-HLA-A1101-9mers- 24P4C12		
Each peptide is a portion of SEQ ID NO: 15; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.		
Start	Subsequence	Score
8	AVGQMMSTM	0.020
5	ILVAVGQMM	0.006
4	WILVAVGQM	0.006
6	LVAVGQMMS	0.004
2	WYWILVAVG	0.001
7	VAVGQMMST	0.001

183

TABLE XIV-continued

1	SWYWILVAV	0.000
3	YWILVAVGQ	0.000
V8-HLA-A1101-9mers- 24P4C12		
Each peptide is a portion of SEQ ID NO: 17; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.		
Start	Subsequence	Score
1	NYYWLPIMR	0.320
20	QTSILGAYV	0.010
17	HVPQTSILG	0.008
19	FQTSILGAY	0.006
18	VFQTSILGA	0.004
4	WLPIMRNPI	0.004
10	NPITPTGHV	0.003
2	YYWLPIMRN	0.002
9	RNPITPTGH	0.001
12	ITPTGHVFQ	0.001
16	GHVFQTSIL	0.001
13	TPTGHVFQT	0.001
11	PITPTGHVF	0.000
7	IMRNPITPT	0.000
5	LPIMRNPI	0.000
15	TGHVFQTSI	0.000
6	PIMRNPI	0.000
14	PTGHVFQTS	0.000
8	MRNPITPTG	0.000
3	YWLPIMRNP	0.000
V9-HLA-A1101-9mers- 24P4C12		
Each peptide is a portion of SEQ ID NO: 19; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.		
Start	Subsequence	Score
15	ATLGYVLWA	0.030
18	GYVLWASNI	0.018
2	WAMTALYPL	0.008
12	TQPATLGYV	0.006
7	LYPLPTQPA	0.004
13	QPATLGYVL	0.004
4	MTALYPLPT	0.002
11	PTQPATLGY	0.002
6	ALYPLPTQP	0.001
16	TLGYVLWAS	0.001
3	AMTALYPLP	0.000
9	PLPTQPATL	0.000
8	YPLPTQPAT	0.000
5	TALYPLPTQ	0.000
10	LPTQPATLG	0.000
14	PATLGYVLW	0.000
17	LGYVLWASN	0.000
1	YWAMTALYP	0.000

184

TABLE XV

V1-A1101-10mers- 24P4C12		
Each peptide is a portion of SEQ ID NO: 3; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.		
Start	Subsequence	Score
516	RVILEYIDHK	9.000
594	VTDLLFFFGK	3.000
134	QTVGEVFFYTK	3.000
333	RIRIAIALLK	2.400
544	CLWCLEKFIK	2.400
621	RIPGLGKDFK	1.200
559	AYIMIAIYGK	1.200
72	GAYCGMGENK	1.200
584	NIVRVVVLDK	1.200
680	SLDRPYYSK	0.800
469	FASFYWAFHK	0.600
272	GIYYCWEEYR	0.480
428	GYSSKGLIQR	0.480
337	AIALLKKEASK	0.400
53	IVAWLYGDPR	0.400
211	SLNARDISVK	0.400
322	ILLMLLIFLR	0.360
423	MCVFQGYSSK	0.300
507	LILTLVQIAR	0.240
578	FMLLMRNIVR	0.240
120	CPEDPWTVGK	0.200
533	VARCIMCCFK	0.200
15	KPVKYDPSFR	0.180
264	GVLGVLAYGI	0.180
609	GVGVLSFFFF	0.180
684	PYYMSKSLK	0.160
667	FLCFLEDLER	0.160
171	LLPSAPALGR	0.160
6	RDEDEAYGK	0.120
698	KNEAPPDNKK	0.120
484	TFPLISAFIR	0.120
237	GVALVLSLLF	0.120
74	YCGMGENKDK	0.100
689	KSLLKILGKK	0.090
649	YVIASGFFSV	0.090
281	RVLRDKGASI	0.090
612	VLSFFFFSGR	0.080
487	LISAFIRTLR	0.080
275	YCWEYRVLR	0.080
438	SVFNLQIYGV	0.080
697	KKNEAPPDNK	0.060
392	SNISSPGCEK	0.060
571	CVSAKNAFML	0.060
259	LVLILGVLVG	0.060
49	IVVGIVAWLY	0.060
240	LVLSLLFILL	0.060
317	AVLEAILLLM	0.060
362	FVLLLCIAY	0.060
433	GLIQRSVFNL	0.054
449	GLFWTLNWVL	0.048
493	RTLRYHTGSL	0.045
518	ILEYIDHKLR	0.040
252	LVAGPLVLVL	0.040
618	FSGRIPGLGK	0.040
688	KSLLKILGK	0.040
606	VVGVGVLVSF	0.040
541	FKCCLWCLEK	0.040
657	SVFGMCVDTL	0.040
360	VPFVLLLCI	0.040
233	LVALGVALVL	0.040
331	RQRIRIAIAL	0.036
589	VVLDKVTDLL	0.030
157	TVITSLQQEL	0.030
463	CVLAGAFASF	0.030
588	VVLDKVTDL	0.030
700	EAPPDNKKRK	0.030
314	IVLAVLEAIL	0.030
456	WVLALGQCVL	0.030
257	LVLVLILGVL	0.030

185

TABLE XV-continued

34	DVICCVLFL	0.027
611	GVLSFFFFSG	0.027
59	GDPRQVLYPR	0.024
220	KIFEDFAQSW	0.024
654	GFESVFGMCV	0.024
548	LEKFIKFLNR	0.024
467	GAFASFYWAF	0.024
674	LERNNGLDR	0.024
347	AVGQMMSTMF	0.020
566	YGKNFCVSAK	0.020
353	STMFYPLVTF	0.020
585	IVRVVVLDKV	0.020
701	APPDNKKRKK	0.020
304	SVQETWLAAL	0.020
380	ATSGQPQYVL	0.020
662	CVDTLFLCFL	0.020
414	LVNSSCPGLM	0.020
19	YDPSFRGPIK	0.020
116	CVSSCPEDPW	0.020
186	NVTPPALPGI	0.020
642	MTSILGAYVI	0.020
512	VQIARVILEY	0.018
478	KPQDIPTFPL	0.018
47	GYIVVGIVAW	0.018
461	GQCVLAGAFA	0.018
239	ALVLSLLFIL	0.018
4	KQRDEDEAY	0.018
603	KLLVVGGVGV	0.018
553	KFLNRNAYIM	0.018
163	QQLCPSFLL	0.018
267	GVLAYGIYYC	0.018
V3-HLA-A1101- 10mers-24P4C12		
Each peptide is a portion of SEQ ID NO: 7; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.		
Start	Subsequence	Score
9	NITPPALPGI	0.004
5	FPWTNITPPA	0.004
3	RCPFWNITP	0.002
7	WTNITPPALP	0.001
10	ITPPALPGIT	0.001
4	CFPWTNITP	0.000
1	LGRCFPWTNI	0.000
8	TNITPPALPG	0.000
2	GRCFPWTNIT	0.000
6	PWTNITPPAL	0.000
V5-HLA-A1101- 10mers-24P4C12		
Each peptide is a portion of SEQ ID NO: 11; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.		
Start	Subsequence	Score
6	ILLVLVIFLR	0.360
1	AVLEAILLLV	0.060
9	LVLIFLRQRI	0.030
8	LLVLIFLRQR	0.012
10	VLIFLRQRI	0.012
5	AILLLVLIFL	0.012
2	VLEAILLLVL	0.008
4	EAILLLVLIF	0.002
7	LLVLVIFLRQ	0.001
3	LEAILLLVLI	0.001

186

TABLE XV-continued

V6-HLA-A1101- 10mers-24P4C12		
Each peptide is a portion of SEQ ID NO: 13; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.		
Start	Subsequence	Score
2	GYSSKGLIPR	0.480
7	GLIPRSVFNL	0.054
9	IPRSVFNLQI	0.004
8	LIPRSVFNLQ	0.000
5	SKGLIPRSVF	0.000
6	KGLIPRSVFN	0.000
1	QGYSSKGLIP	0.000
10	PRSVFNLQIY	0.000
4	SSKGLIPRSV	0.000
3	YSSKGLIPRS	0.000
V7-HLA-A1101- 10mers-24P4C12		
Each peptide is a portion of SEQ ID NO: 15; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.		
Start	Subsequence	Score
9	AVGQMMSTMF	0.020
5	WILVAVGQMM	0.006
7	LVAVGQMMST	0.004
8	VAVGQMMSTM	0.003
6	ILVAVGQMMS	0.001
3	WYWILVAVGQ	0.001
1	QSWYWILVAV	0.000
4	YWILVAVGQM	0.000
2	SWYWILVAVG	0.000
V8-HLA-A1101- 10mers-24P4C12		
Each peptide is a portion of SEQ ID NO: 17; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.		
Start	Subsequence	Score
18	HVFQTSILGA	0.080
1	LNYYWLPIMR	0.032
21	QTSILGAYVI	0.020
20	FQTSILGAYV	0.006
11	NPITPTGHVF	0.003
13	ITPTGHVFQT	0.003
19	VFQTSILGAY	0.002
2	NYWLPIMRN	0.002
10	RNPITPTGHV	0.001
15	PTGHVFQTSI	0.001
6	LPIMRNPITP	0.001
8	IMRNPITPTG	0.000
5	WLPIMRNPIT	0.000
4	YWLPIMRNPI	0.000
9	MRNPITPTGH	0.000
14	TPTGHVFQTS	0.000
16	TGHVFQTSIL	0.000
17	GHVFQTSILG	0.000
7	PIMRNPITPT	0.000
3	YWLPIMRNP	0.000
12	PITPTGHVFQ	0.000

187

TABLE XV-continued

V9-HLA-A1101- 10mers-24P4C12		
Each peptide is a portion of SEQ ID NO: 19; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.		
Start	Subsequence	Score
13	TQPATLGYVL	0.012
7	ALYPLPTQPA	0.008
11	LPTQPATLGY	0.004
9	YPLPTQPATL	0.003
16	ATLGYVLWAS	0.003
14	QPATLGYVLW	0.002
19	GYVLWASNIS	0.002
1	AYWAMTALYP	0.002
12	PTQPATLGYV	0.001
5	MTALYPLPTQ	0.001
4	AMTALYPLPT	0.001
8	LYPLPTQPAT	0.000
18	LGYVLWASNI	0.000
15	PATLGYVLWA	0.000
17	TLGYVLWASN	0.000
3	WAMTALYPLP	0.000
2	YWAMTALYPL	0.000
6	TALYPLPTQP	0.000
10	PLPTQPATLG	0.000

TABLE XVI

V1-HLA-A24-9mers- 24P4C12		
Each peptide is a portion of SEQ ID NO: 3; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.		
Start	Subsequence	Score
356	FYPLVTFVL	420.000
57	LYGDPRQVL	288.000
496	RYHTGSLAF	200.000
648	AYVIASGFF	150.000
87	LYFNIFSCI	84.000
386	QYVLWASNI	75.000
88	YFNIFSCIL	30.000
666	LFLCFLEDL	30.000
450	LFWTLNWVL	24.000
503	AFGALILTL	24.000
84	PYLLYFNIF	21.600
540	CFKCCLWCL	20.000
684	PYYMSKSL	20.000
617	FFSGRIPGL	20.000
658	VFGMCVDTL	20.000
553	KFLNRNAYI	15.000
251	RLVAGPLVL	12.000
583	RNIVRVVVL	12.000
484	TFPLISAFI	10.500
47	GYIVVGIVA	10.500
301	AYQSVQETW	10.500
468	AFASFYWAF	10.000
139	VFYTKNRNF	10.000
518	ILEYIDHKL	9.240
361	TFVLLLCI	9.000
577	AFMLLMRNI	9.000
446	GVLGLFWTL	8.640
258	VLVLILGVL	8.400
49	IVVGIVAWL	8.400
154	WNMTVITSL	8.400
311	AALIVLAVL	8.400
261	LILGVLGVL	8.400
440	FNLQIYGVL	8.400

188

TABLE XVI-continued

234	VALGVALVL	8.400	5
683	RPYYMSKSL	8.000	
333	RIRIAIAL	8.000	
596	DLLEFFGKL	7.920	
65	LYPRNSTGA	7.500	
328	IFLRQRIRI	7.500	10
317	AVLEAILLL	7.200	
255	GPLVLVLIL	7.200	
38	CVLFLLFIL	7.200	
240	LVLSLLFIL	7.200	
232	ILVALGVAL	7.200	15
589	VVLDKVTDL	7.200	
170	FLLPSAPAL	7.200	
357	YPLVTFVLL	7.200	
236	LGVALVLSL	7.200	
621	RIPGLGKDF	7.200	20
158	VITSLQQEL	6.336	
305	VQETWLAAL	6.000	
15	KPVKYDPSF	6.000	
547	CLEKFIKFL	6.000	
597	LLLEFFGKLL	6.000	25
565	IYGKNFCVS	6.000	
34	DVICCVLFL	6.000	
308	TWLAALIVL	6.000	
184	WTNVTTPPAL	6.000	
316	LAVLEAILL	6.000	30
200	TIQQGISGL	6.000	
635	NYWLPIMT	6.000	
140	FYTKNRNFC	6.000	
673	DLERNNGSL	6.000	
442	LQIYGVGL	6.000	35
414	LVNSSCPGL	6.000	
444	IYGVGLGLFW	6.000	
452	WTLNWLAL	6.000	
242	LSLLFILL	6.000	
605	LVVGGVGV	6.000	40
638	WLPIMTSIL	6.000	
511	LVQIARVIL	6.000	
163	QQELCPSFL	6.000	
291	SQLGFTTNL	6.000	
434	LIQRSVFNL	6.000	45
432	KGLIQRSVF	6.000	
225	FAQSWYWL	6.000	
322	ILLMLIFL	6.000	
593	KVTDLLFF	5.760	
241	VLSLLFILL	5.760	50
253	VAGPLVLVL	5.760	
237	GVALVLSLL	5.600	
228	SWYWILVAL	5.600	
249	LLRLVAGPL	5.600	
35	VICCVLFLL	5.600	55
32	CTDVICCVL	5.600	
590	VLDKVTDLL	5.600	
217	ISVKIFEDF	5.040	
224	DFAQSWYWI	5.000	
614	SFFFFSGRI	5.000	60
274	YYCWEEYRV	5.000	
636	YYWLPIMTS	5.000	
370	AYWAMTALY	5.000	
573	SAKNAFMLL	4.800	
351	MMSTMFYPL	4.800	65
315	VLAVLEAIL	4.800	
100	IISVAENGL	4.800	
204	GISGLIDSL	4.800	
687	MSKSLLKIL	4.800	
244	LLFILLRL	4.800	65
499	TGSLAFGAL	4.800	

189

TABLE XVI-continued

V3-HLA-A24-9mers- 24P4C12		
Each peptide is a portion of SEQ ID NO: 7; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.		
Start	Subsequence	Score
6	WTNITPPAL	6.000
9	ITPPALPGI	1.800
2	RCFPWTNIT	0.288
1	GRCFPWTNI	0.100
3	CFPWTNITP	0.075
7	TNITPPALP	0.015
5	PWTNITPPA	0.014
8	NITPPALPG	0.012
4	FPWTNITPP	0.010
V5-HLA-A24-9mers- 24P4C12		
Each peptide is a portion of SEQ ID NO: 11; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.		
Start	Subsequence	Score
5	ILLVLIFL	8.400
4	AILLVLIF	3.600
9	VLIFLRQRI	2.160
3	EAILLLVLI	1.800
2	LEAILLLVL	0.480
1	VLEAILLLV	0.210
7	LLVLIFLRQ	0.025
6	LLVLIFLFR	0.018
8	LVLIFLRQR	0.015
V6-HLA-A24-9mers- 24P4C12		
Each peptide is a portion of SEQ ID NO: 13; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.		
Start	Subsequence	Score
5	KGLIPRSVF	6.000
7	LIPRSVFNL	6.000
1	GYSSKGLIP	0.500
6	GLIPRSVFN	0.180
3	SSKGLIPRS	0.120
8	IPRSVFNLQ	0.020
4	SKGLIPRSV	0.014
2	YSSKGLIPR	0.010
9	PRSVFNLQI	0.010
V7-HLA-A24-9mers- 24P4C12		
Each peptide is a portion of SEQ ID NO: 15; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.		
Start	Subsequence	Score
5	ILVAVGQMM	1.260
4	WILVAVGQM	0.750
2	WYWILVAVG	0.600
8	AVGQMMSTM	0.500
7	VAVGQMMST	0.150
1	SWYWILVAV	0.140

190

TABLE XVI-continued

6	LVAVGQMMMS	0.100
3	YWILVAVGQ	0.021
V8-HLA-A24-9mers- 24P4C12		
Each peptide is a portion of SEQ ID NO: 17; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.		
Start	Subsequence	Score
2	YYWLPIMRN	5.000
4	WLPIMRNPI	1.800
15	TGHVFTQSI	1.000
18	VFQTSILGA	0.750
16	GHVFQTSIL	0.600
1	NYWLPIMR	0.600
11	PITPTGHVF	0.240
10	NPITPTGHV	0.150
5	LPIMRNPIT	0.150
19	FQTSILGAY	0.140
20	QTSILGAYV	0.120
13	TPTGHVFTQ	0.100
7	IMRNPITPT	0.100
9	RNPITPTGH	0.030
3	YWLPIMRNP	0.025
14	PTGHVFTQS	0.020
12	ITPTGHVFQ	0.015
17	HVFQTSILG	0.010
8	MRNPITPTG	0.003
6	PIMRNPITP	0.002
V9-HLA-A24-9mers- 24P4C12		
Each peptide is a portion of SEQ ID NO: 19; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.		
Start	Subsequence	Score
18	GYVLWASNI	75.000
7	LYPLPTQPA	9.000
2	WAMTALYPL	6.000
13	QPATLGIVL	4.800
9	PLPTQPATL	0.600
8	YPLPTQPAT	0.180
15	ATLGIVLWA	0.150
12	TQPATLGIV	0.150
16	TLGYVLWAS	0.140
17	LGYVLWASN	0.120
4	MTALYPLPT	0.100
11	PTQPATLG	0.018
5	TALYPLPTQ	0.015
6	ALYPLPTQP	0.014
3	AMTALYPLP	0.012
10	LPTQPATLG	0.010
14	PATLGIVLW	0.010
1	YWAMTALYP	0.010

191

TABLE XVII

V1-HLA-A24-10mers- 24P4C12		
Each peptide is a portion of SEQ ID NO: 3; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.		
Start	Subsequence	Score
356	FYPLVTFVLL	360.000
301	AYQSVQETWL	300.000
87	LYFNIFSCIL	200.000
140	FYTKNRNFCL	200.000
274	YYCWEEYRVL	200.000
370	AYWAMTALYL	200.000
18	KYDPSFRGPI	120.000
685	YYMSKSLKI	82.500
636	YYWLPIMTSI	70.000
439	VFNLQIYGVL	42.000
355	MFYPLVTFVL	33.600
169	SFLLPSAPAL	30.000
425	VFQGYSSKGL	30.000
616	FFFSGRIPGL	20.000
224	DFAQSWYWIL	20.000
478	KPQDITPFPL	14.400
131	EFSTVGEVF	14.000
658	VFGMCDVTLF	14.000
569	NFCVSAKNAF	12.000
630	KSPHLNYYWL	12.000
493	RTLRYHTGSL	12.000
331	RQRIRIAIAL	11.200
517	VILEYIDHKL	11.088
40	LFLLFILGYI	10.500
589	VVLDKVTDLL	10.080
157	TVITSLQQEL	9.504
520	EYIDHKLGRV	9.000
386	QYVLWASNIS	9.000
445	YGVGLGFWTL	8.640
240	LVLSLLFILL	8.640
248	LLLRLVAGPL	8.400
257	LVLVLILGVL	8.400
48	YIVVGIVAWL	8.400
260	VLILGVLGVL	8.400
236	LGVALVLSLL	8.400
34	DVICCVLFL	8.400
683	RPYMSKSL	8.000
648	AYVIASGFFS	7.500
47	GYIVVGIVAW	7.500
65	LYPRNSTGAY	7.500
553	KFLNRNAYIM	7.500
254	AGPLVLVLIL	7.200
304	SVQETWLAAL	7.200
231	WILVALGVAL	7.200
637	YWLPIMTSIL	7.200
162	LQQELCPSFL	7.200
239	ALVLSLLFIL	7.200
318	VLEAILLLML	7.200
314	IVLAVLEAIL	7.200
37	CCVLFLLFIL	7.200
546	WCLEKFIKFL	7.200
350	QMMSTMFYPL	7.200
99	NIISVAENGL	7.200
203	QGISGLIDSL	7.200
243	SLLFILLRL	7.200
229	WYWILVALGV	7.000
31	SCTDVICCVL	6.720
441	NLQIYGVGL	6.000
357	YPLVTFVLL	6.000
604	LLVVGVGVL	6.000
510	TLVQIARVIL	6.000
596	DLLFFGKLL	6.000
536	CIMCCFKCCL	6.000
588	VVLDKVTDL	6.000
433	GLIQRSVFNL	6.000
659	FGMCDVTLFL	6.000
456	WVLALGQCVL	6.000
413	HLVNSSCPGL	6.000
290	ISQLGFTTNL	6.000

192

TABLE XVII-continued

321	AILLLMLIFL	6.000
316	LAVLEAILLL	6.000
57	LYGDPRQVLY	6.000
91	IFSCILSSNI	6.000
77	MGENKDKPYL	6.000
163	QQELCPSFLL	6.000
199	TTIQQGISGL	6.000
500	GSLAFGALIL	6.000
83	KPYLLYFNIF	5.760
310	LAALIVLAVL	5.600
233	LVALGVALVL	5.600
227	QSWYWILVAL	5.600
661	MCVDTLFLCF	5.184
565	IYGNFCVSA	5.000
279	EYRVLRDKGA	5.000
635	NIYWLPIMTS	5.000
273	IYYCWEEYRV	5.000
444	IYGVGLFWT	5.000
686	YMSKSLLKIL	4.800
56	WLYGDPRQVL	4.800
235	ALGVALVLSL	4.800
252	LVAGPLVLVL	4.800
449	GLFWTLNWWL	4.800
502	LAFGALILTL	4.800
625	LGKDFKSPHL	4.800
498	HTGSLAFAGL	4.800
572	VSAKNAFMLL	4.800
542	KCCCLWCLEKF	4.400
442	LQIYGVGLGF	4.200
368	CIAYWAMTAL	4.000
241	VLSLLFILL	4.000
V3-HLA-A24-10mers- 24P4C12		
Each peptide is a portion of SEQ ID NO: 7; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.		
Start	Subsequence	Score
9	NITPPALPGI	1.200
1	LGRCFPWTNI	1.000
6	PWTNITPPAL	0.400
10	ITPPALPGIT	0.216
5	FPWTNITPPA	0.140
4	CFPWTNITPP	0.075
3	RCFPWTNITP	0.024
8	TNITPPALPG	0.015
7	WTNITPPALP	0.015
2	GRCFPWTNIT	0.012
V5-HLA-A24-10mers- 24P4C12		
Each peptide is a portion of SEQ ID NO: 11; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.		
Start	Subsequence	Score
5	AILLLVLIFL	8.400
2	VLEAILLLVL	7.200
4	EAILLLVLIF	3.600
9	LVLIFLRQRI	2.160
1	AVLEAILLLV	0.252
3	LEAILLLVLI	0.120
7	LLLVLIFLRQ	0.025
6	ILLVLVIFLR	0.018
10	VLIFLRQRI	0.015
8	LLVLIFLRQR	0.015

193

TABLE XVII-continued

V6-HLA-A24-10mers- 24P4C12		
Each peptide is a portion of SEQ ID NO: 13; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.		
Start	Subsequence	Score
7	GLIPRSVFNL	7.200
9	IPRSVFNLQI	1.000
2	GYSSKGLIPR	0.500
6	KGLIPRSVFN	0.300
5	SKGLIPRSVF	0.200
4	SSKGLIPRSV	0.140
3	YSSKGLIPRS	0.120
8	LIPRSVFNLQ	0.030
1	QGYSSKGLIP	0.010
10	PRSVFNLQIY	0.001
V7-HLA-A24-10mers- 24P4C12		
Each peptide is a portion of SEQ ID NO: 15; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.		
Start	Subsequence	Score
9	AVGQMMSTMF	2.000
5	WILVAVGQMM	1.260
4	YWILVAVGQM	0.750
8	VAVGQMMSTM	0.750
3	WYWILVAVGQ	0.700
6	ILVAVGQMMS	0.150
1	QSWYWILVAV	0.140
7	LVAVGQMMST	0.100
2	SWYWILVAVG	0.012
V8-HLA-A24-10mers- 24P4C12		
Each peptide is a portion of SEQ ID NO: 17; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.		
Start	Subsequence	Score
2	NYWLPIMRN	5.000
16	TGHVFQTSIL	4.000
11	NPITPTGHVF	3.000
4	YWLPIMRNP	2.160
19	VFQTSILGAY	1.050
21	QTSILGAYVI	1.000
3	YYWLPIMRNP	0.700
10	RNPITPTGHV	0.300
14	TPTGHVFQTS	0.202
5	WLPIMRNPIT	0.150
13	ITPTGHVFQT	0.150
20	FQTSILGAYV	0.120
18	HVFQTSILGA	0.100
15	PTGHVFQTSI	0.100
6	LPIMRNPITP	0.015
7	PIMRNPITPT	0.015
8	IMRNPITPTG	0.014
1	LNYYWLPIMR	0.012
9	MRNPITPTGH	0.002
17	GHVFQTSILG	0.002
12	PITPTGHVFQ	0.001

194

TABLE XVII-continued

V9-HLA-A24-10mers- 24P4C12		
Each peptide is a portion of SEQ ID NO: 19; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.		
Start	Subsequence	Score
19	GYVLWASNIS	9.000
8	LYPLPTQPAT	7.500
13	TQPATLGYVL	7.200
9	YPLPTQPATL	7.200
2	YWAMTALYPL	4.000
18	LGYYVLWASNI	1.000
1	AYWAMTALYP	0.500
16	ATLGYVLWAS	0.210
7	ALYPLPTQPA	0.144
17	TLGYVLWASN	0.120
4	AMTALYPLPT	0.100
14	QPATLGYVVLW	0.100
11	LPTQPATLGY	0.100
12	PTQPATLGYV	0.018
6	TALYPLPTQP	0.018
3	WAMTALYPLP	0.018
15	PATLGYVLWA	0.010
5	MTALYPLPTQ	0.010
10	PLPTQPATLG	0.002

TABLE XVIII		
V1-HLA-B7-9mers-24P4C12		
Each peptide is a portion of SEQ ID NO: 3; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.		
Start	Subsequence	Score
255	GPLVLVLIL	80.000
631	SPHLNYYWL	80.000
357	YPLVTFVLL	80.000
683	RPYYMSKSL	80.000
317	AVLEAILLL	60.000
249	LLRLVAGPL	40.000
494	TLRYHTGSL	40.000
333	RIRIAIAL	40.000
311	AALIVLAVL	36.000
511	LVQIARVIL	30.000
414	LVNSSCPGL	20.000
38	CVLFLLFIL	20.000
49	IVVGIVAWL	20.000
446	GVGLGFWTL	20.000
237	GVALVLSLL	20.000
240	LVLSSLFIL	20.000
605	LVVGGVGVL	20.000
34	DVICCVLFL	20.000
589	VVLDKVTDL	20.000
347	AVGQMMSTM	15.000
573	SAKNAFMLL	12.000
253	VAGPLVLVL	12.000
369	IAYWAMTAL	12.000
225	FAQSWYWIL	12.000
213	NARDISVKI	12.000
514	IARVILEYI	12.000
154	WNMTVITSL	12.000
316	LAVLEAILL	12.000
234	VALGVALVL	12.000
396	SPGCEKVP	8.000
83	KPYLLYFNI	8.000
406	TSCNPTAHL	6.000
381	TSGQPQYVL	6.000
571	CVSAKNAFM	5.000

195

TABLE XVIII-continued

261	LILGVLGV	4.000
315	VLAVLEAIL	4.000
291	SQLGFTTNL	4.000
638	WLPIMTSIL	4.000
258	VLVLILGV	4.000
452	WTLNWWLAL	4.000
28	KNRSCTDVI	4.000
241	VLSLLFILL	4.000
236	LGVALVLSL	4.000
440	FNLQIYGV	4.000
184	WTNVTTPAL	4.000
597	LLLFFGKLL	4.000
583	RNIVRVVVL	4.000
275	YCWEERYVL	4.000
170	FLLPSAPAL	4.000
596	DLLLFFGKL	4.000
282	VLKDKGASI	4.000
158	VITSLQOEL	4.000
537	IMCCFKCCL	4.000
660	GMCVDLFL	4.000
457	VLALGQCVL	4.000
499	TGSLAFGAL	4.000
66	YPRNSTGAY	4.000
141	YTKNRNFC	4.000
555	LNRNAYIMI	4.000
426	FQGYSSKGL	4.000
244	LLFILLRL	4.000
242	LSLLFILL	4.000
487	LISAFIRTL	4.000
79	ENKDKPYLL	4.000
351	MMSTMFPYPL	4.000
442	LQIYGVGL	4.000
200	TIQQGISGL	4.000
434	LIQSVFNL	4.000
501	SLAFGALIL	4.000
322	ILLMLIFL	4.000
251	RLVAGPLVL	4.000
204	GISGLIDSL	4.000
572	VSAKNAFML	4.000
687	MSKSLKIL	4.000
100	IISVAENGL	4.000
232	ILVALGVAL	4.000
302	YQSVQETWL	4.000
35	VICCVLFLL	4.000
25	GPIKNRSCT	3.000
482	IPTFPLISA	3.000
344	ASKAVGQMM	3.000
343	EASKAVGQM	3.000
149	LPGVPWNMT	3.000
581	LMRNIVRVV	2.000
152	VPWNMTVIT	2.000
531	NPVARCIM	2.000
188	TPPALPGIT	2.000
112	TPQVCVSSC	2.000
60	DPRQVLYPR	2.000
525	KLRGVQNPV	2.000
314	IVLAVLEAI	2.000
167	CPSFLLPSA	2.000
151	GVPWNMTVI	2.000
192	LPGITNDTT	2.000
359	LVTFVLLLI	2.000
252	LVAGPLVLV	1.500
491	FIRTLRYHT	1.500
530	QNPVARCIM	1.500
239	ALVLSLLFI	1.200
305	VQETWLAAL	1.200
V3-HLA-B7-9mers- 24P4C12		
Each peptide is a portion of SEQ ID NO: 7; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.		
Start	Subsequence	Score
6	WTNITPPAL	4.000
9	ITPPALPGI	0.400

196

TABLE XVIII-continued

4	FPWTNITPP	0.200
2	RCPFWTNIT	0.100
1	GRCFPWTNI	0.060
7	TNITPPALP	0.015
8	NITPPALPG	0.015
3	CFPWTNITP	0.001
5	PWTNITPPA	0.001
V5-HLA-B7-9mers- 24P4C12		
Each peptide is a portion of SEQ ID NO: 11; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.		
Start	Subsequence	Score
5	ILLVLIFL	4.000
3	EAILLLVLI	1.200
9	VLIFLRQRI	0.600
2	LEAILLLVL	0.400
4	AILLVLIF	0.060
1	VLEAILLLV	0.060
8	LVLIFLRQR	0.050
7	LLVLIFLRQ	0.010
6	LLLVLIFLR	0.010
V6-HLA-B7-9mers- 24P4C12		
Each peptide is a portion of SEQ ID NO: 13; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.		
Start	Subsequence	Score
7	LIPRSVFNL	4.000
8	IPRSVFNLQ	2.000
5	KGLIPRSVF	0.045
6	GLIPRSVFN	0.020
4	SKGLIPRSV	0.020
3	SSKGLIPRS	0.020
2	YSSKGLIPR	0.010
9	PRSVFNLQI	0.004
1	GYSSKGLIP	0.001
V7-HLA-B7-9mers- 24P4C12		
Each peptide is a portion of SEQ ID NO: 15; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.		
Start	Subsequence	Score
8	AVGQMMSTM	15.000
5	ILVAVGQMM	1.000
4	WILVAVGQM	1.000
7	VAVGQMMST	0.300
6	LVAVGQMMS	0.100
1	SWYWILVAV	0.020
3	YWILVAVGQ	0.001
2	WYWILVAVG	0.001

197

TABLE XVIII-continued

V8-HLA-B7-9mers- 24P4C12			5	
Each peptide is a portion of SEQ ID NO: 17; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.				
Start	Subsequence	Score		10
10	NPITPTGHV	PTA 6.000		
5	LPIMRNPIT	2.000		
13	TPTGHVFQT	2.000		
7	IMRNPITPT	1.500		
4	WLPIMRNPI	0.600	15	
15	TGHVVFQTSI	0.400		
16	GHVVFQTSIL	0.400		
20	QTSILGAYV	0.200		
17	HVFQTSILG	0.050		
19	FQTSILGAY	0.020	20	
18	VFQTSILGA	0.010		
12	ITPTGHVFQ	0.010		
9	RNPITPTGH	0.010		
6	PIMRNPITP	0.003		
2	YYWLPIMRN	0.003	25	
11	PITPTGHVF	0.002		
14	PTGHVVFQTS	0.002		
3	YWLPIMRNP	0.001		
8	MRNPITPTG	0.001		
1	NYWLPIMR	0.001		
V9-HLA-B7-9mers- 24P4C12			30	
Each peptide is a portion of SEQ ID NO: 19; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.				
Start	Subsequence	Score		35
13	QPATLGYYVL	80.000		
2	WAMTALYPL	36.000		
8	YPLPTQPAT	2.000		
9	PLPTQPATL	0.400		
10	LPTQPATLG	0.300	40	
15	ATLGYYVLWA	0.300		
12	TQPATLGYYV	0.200		
4	MTALYPLPT	0.100		
5	TALYPLPTQ	0.045		
18	GYVLWASNI	0.040	45	
3	AMTALYPLP	0.030		
6	ALYPLPTQP	0.030		
17	LGYVLWASN	0.020		
16	TLGYVLWAS	0.020		
7	LYPLPTQPA	0.015	50	
14	PATLGYYVLW	0.006		
11	PTQPATLGY	0.002		
1	YWAMTALYP	0.001		

TABLE XIX

V1-HLA-B7-10mers- 24P4C12			60
Each peptide is a portion of SEQ ID NO: 3; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.			
Start	Subsequence	Score	65
478	KPQDIPTFPL	120.000	
683	RPYYMSKSL	80.000	

198

TABLE XIX-continued

357	YPLVTFVLLL	80.000	80
331	RQRIRIAIAL	40.000	
571	CVSAKNAFML	20.000	85
257	LVLVLILGVL	20.000	
456	WVLALGQCVL	20.000	90
588	VVVLDKVTDL	20.000	
157	TVITSLQQEL	20.000	95
240	LVLSLLFILL	20.000	
66	YPRNSTGAYC	20.000	100
314	IVLAVLEAIL	20.000	
589	VVLDKVTDLL	20.000	105
252	LVAGPLVLVL	20.000	
657	SVFGMCVDTL	20.000	110
304	SVQETWLAAL	20.000	
34	DVICCVLFLL	20.000	115
233	LVALGVALVL	20.000	
380	ATSGQPQYVL	18.000	120
317	AVLEAILLLM	15.000	
321	AILLMLIFL	12.000	125
502	LAFGALITL	12.000	
239	ALVLSLLFIL	12.000	130
659	FGMCVDTLFL	12.000	
254	AGPLVLVLIL	12.000	135
350	QMMSTMFYPL	12.000	
235	ALGVALVLSL	12.000	140
536	CIMCCFKCCL	12.000	
316	LAVLEAILLL	12.000	145
310	LAALIVLAVL	12.000	
585	IVRVVVLDKV	10.000	150
56	WLYGDPKQVL	9.000	
192	LPGITNDTTI	8.000	155
510	TLVQIARVIL	6.000	
662	CVDTLFLCFL	6.000	160
329	FLRQRIRIAI	6.000	
405	NTSCNPTAHL	6.000	165
414	LVNSSCPGLM	5.000	
413	HLVNSSCPGL	4.000	170
203	QGISGLIDSL	4.000	
368	CIAYWAMTAL	4.000	175
686	YMSKSLKIL	4.000	
99	NIISVAENGL	4.000	180
665	TLFLCFLEDL	4.000	
290	ISQLGFTTNL	4.000	185
441	NLQIYGVGL	4.000	
630	KSPHLNYYWL	4.000	190
315	VLAVLEAILL	4.000	
236	LGVALVLSLL	4.000	195
596	DLLEFFGKLL	4.000	
60	DPRQVLYPRN	4.000	200
243	SLLFILLRL	4.000	
37	CCVLFLLFIL	4.000	205
449	GLFWTLNWL	4.000	
162	LQQELCPSEL	4.000	210
625	LGKDFKSPHL	4.000	
227	QSWYILVAL	4.000	215
498	HTGSLAFGAL	4.000	
48	YIVVGIVAWL	4.000	220
604	LLVVGVGVL	4.000	
149	LPGVPMNMTV	4.000	225
260	VLILGVLGVL	4.000	
493	RTLRYHTGSL	4.000	230
248	LLLRLVAGPL	4.000	
231	WILVALGVAL	4.000	235
500	GSLAFGALIL	4.000	
546	WCLEKFIKFL	4.000	240
241	VLSLLFILL	4.000	
539	CCFKCCLWCL	4.000	245
445	YGVGLFWTL	4.000	
307	ETWLAALIVL	4.000	250
435	IQRSVFNLQI	4.000	
572	VSAKNAFMLL	4.000	255
433	GLIQRSVFNL	4.000	
517	VILEYIDHKL	4.000	260
199	TTIQQGISGL	4.000	
31	SCTDVICCVL	4.000	265
178	LGRCFPWTNV	3.000	
343	EASKAVGQMM	3.000	270
346	KAVGQMMSTM	3.000	
581	LMRNIVRVVV	3.000	275
573	SAKNAFMLLM	3.000	

Figure 2F-3

```
500  G S L A F G A L I L T L V Q I A R V I L
1501 CTGGGTCATTGGCATTGGAGCCCTCATCCTGACCCTTGTGCAGATAGCCCGGGTCATCT
520  E Y I D H K L R G V Q N P V A R C I M C
1561 TGGAGTATATTGACCACAAGCTCAGAGGAGTGCAGAACCCCTGTAGCCCGCTGCATCATGT
540  C F K C C L W C L E K F I K F L N R N A
1621 GCTGTTTCAAGTGCIGCCICIGGTGTCTGGAAAAATTTATCAAGTTCCTAAACCGCAATG
560  Y I M I A I Y G K N F C V S A R N A F M
1681 CATAATCATGATCGCCAATCTACGGGAAGAATTTCTGTGTCTCAGCCAAAAATGCGTTCA
580  L L M R N I V R V V V L D K V T D L L L
1741 TGCTACTCATGCGAAACATTGTCAGGGTGGTCTGCTCTGGACAAAGTCACAGACCTGCTGC
600  F F G K L L V V G G V G V L S F F F F S
1801 TGTTCTTTGGGAAGCTGCTGGTGGTCGGAGGCGTGGGGGTCTGTCTTCTTTTTTTTCT
620  G R I P G L G K D F K S P H L N Y Y W L
1861 CCGGTCGCATCCCGGGGCTGGGTAAAGACTTTAAGAGCCCCACCTCAACTATTACTGGC
640  P I M T S I L G A Y V I A S G F F S V F
1921 TGCCCATCATGACCTCCAATCTGGGGGCTATGTCATCGCCAGCGGCTTCTTCAGCGTTT
660  G M C V D T L F L C F L E D L E R N N G
1981 TCGGCATGIGTGGACACGCTCTTCTCTGCTTCTGGAAGACCTGGAGCGGAACAACG
680  S L D R P Y Y M S K S L L K I L G K K N
2041 GCTCCCTGGACCGCCCTACTACATGTCCAAGAGCCTTCTAAAGATTCTGGGCAAGAAGA
700  E A P P D N K K R K K *
2101 ACGAGGCGCCCCGGACAACAAGAAGAGGAAGAAGTGAacagctcgggcctgatccagga
2161 ctgcacccccacccccaccgtccagccatccaacctcacttcgccttacaggtctccattt
2221 tgtggtataaaaaagggttttaggccaggcgccgtggtctcacgcctgtaatccaacatttg
2281 agaggctgaggcgggcggtatcacctgagtcaggagttcgagaccagcctggccaacatgg
2341 tgaaacctccgtctctattataaaaatacaaaaattagccgagagtggtggcatgcacctgt
2401 catcccagctactcgggaggctgaggcaggagaatcgcttgaaacccgggaggcagagggt
2461 gcagtgagccgagatcgcgccactgcactccaacctgggtgacagactctgtctcaaaa
2521 caaaacaaacaaacaaaaagattttattaaagatatatttgtttaactcagtaaaaaaaaaa
2581 aaaaaaa
```

Figure 2G. The cDNA (SEQ ID. NO. : 14) and amino acid sequence (SEQ ID. NO. : 15) of 24P4C12 v.7.

The start methionine is underlined. The open reading frame extends from nucleic acid 6-1802 including the stop codon.

Figure 2G-1

```
1      M G G K Q R D E D D E A Y G K P V K Y
1 gagccATGGGGGGAAAGCAGCGGGACGAGGATGACGAGGCCTACGGGAAGCCAGTCAAAT
20   D P S F R G P I K N R S C T D V I C C V
61 ACGACCCCTCCTTTGAGGCCCCATCAAGAACAGAAGCTGCACAGATGTCATCTGCTGCG
```

199

TABLE XIX-continued

652	ASGFFSVFGM	3.000
402	VPINTSCNPT	2.000
182	FPWTNVTTPA	2.000
528	GVQNPVARI	2.000
281	RVLKDKGASI	2.000
186	NVTTPALPGI	2.000
143	KNRNFCLPGV	2.000
639	LPIMTSILGA	2.000
249	LLRLVAGPLV	2.000
172	LPSAPALGRC	2.000
485	FPLISAFIRT	2.000
264	GVLGVLAYGI	2.000
531	NPVARCIMCC	2.000
163	QQELCPSFLL	1.800
529	VQNPVARCIM	1.500
576	NAFMLLMRNI	1.200
370	AYWAMTALYL	1.200
318	VLEAILLLML	1.200

V3-HLA-B7-10mers-
24P4C12

Each peptide is a portion of SEQ
ID NO: 7; each start position is
specified, the length of peptide is
10 amino acids, and the end
position for each peptide is the
start position plus nine.

Start	Subsequence	Score
1	LGRCPWTNI	6.000
5	FPWTNITPPA	2.000
9	NITPPALPGI	0.400
10	ITPPALPGIT	0.100
6	PWTNITPPAL	0.040
8	TNITPPALPG	0.015
7	WTNITPPALP	0.015
3	RCPFWTNITP	0.010
2	GRCFPWTNIT	0.010
4	CFPWTNITPP	0.001

V5-HLA-B7-10mers-
24P4C12

Each peptide is a portion of SEQ ID
NO: 11; each start position is
specified, the length of peptide is 10
amino acids, and the end position for
each peptide is the start position plus
nine.

Start	Subsequence	Score
5	AILLVLIFL	12.000
9	LVLIFLRQRI	3.000
1	AVLEAILLLV	3.000
2	VLEAILLLVL	1.200
4	EAILLLVLIF	0.060
3	LEAILLLVLI	0.040
7	LLLVLIFLRQ	0.010
6	ILLVLIFLR	0.010
10	VLIFLRQRI	0.010
8	LLVLIFLRQR	0.010

V6-HLA-B7-10mers-
24P4C12

Each peptide is a portion of SEQ
ID NO: 13; each start position is
specified, the length of peptide is
10 amino acids, and the end
position for each peptide is the
start position plus nine.

Start	Subsequence	Score
9	IPRSVFNLQI	80.000
7	GLIPRSVFNL	4.000
4	SSKGLIPRSV	0.200
6	KGLIPRSVFN	0.020
3	YSSKGLIPRS	0.020
8	LIPRSVFNLQ	0.010

200

TABLE XIX-continued

1	QGYSSKGLIP	0.010
5	SKGLIPRSVF	0.005
2	GYSSKGLIPR	0.001
10	PRSVFNLQIY	0.000

V7-HLA-B7-10mers-
24P4C12

Each peptide is a portion of SEQ
ID NO: 15; each start position is
specified, the length of peptide is
10 amino acids, and the end
position for each peptide is the
start position plus nine.

Start	Subsequence	Score
8	VAVGQMMSTM	3.000
5	WILVAVGQMM	1.000
7	LVAVGQMMST	0.500
9	AVGQMMSTMF	0.300
1	QSWYWLIVAV	0.200
4	YWILVAVGQM	0.100
6	ILVAVGQMMS	0.020
2	SWYWLIVAVG	0.001
3	WYWLIVAVGQ	0.001

V8-HLA-B7-10mers-
24P4C12

Each peptide is a portion of SEQ
ID NO: 17; each start position is
specified, the length of peptide is
10 amino acids, and the end
position for each peptide is the
start position plus nine.

Start	Subsequence	Score
16	TGHVFQTSIL	4.000
18	HVFQTSILGA	0.500
11	NPITPTGHVF	0.400
21	QTSILGAYVI	0.400
14	TPTGHVFQTS	0.400
10	RNPITPTGHV	0.300
20	FQTSILGAYV	0.200
6	LPIMRNPITP	0.200
13	ITPTGHVFQT	0.100
8	IMRNPITPTG	0.100
5	WLPIMRNPIT	0.100
4	YWLPIMRNPI	0.060
7	PIMRNPITPT	0.045
15	PTGHVFQTSI	0.040
1	LNYYWLPIMR	0.010
2	NYWLPIMRN	0.003
19	VFQTSILGAY	0.002
17	GHVFQTSILG	0.001
3	YYWLPIMRNP	0.001
12	PITPTGHVFQ	0.001
9	MRNPITPTGH	0.001

V9-HLA-B7-10mers-
24P4C12

Each peptide is a portion of SEQ ID
NO: 19; each start position is
specified, the length of peptide is
10 amino acids, and the end
position for each peptide is the start
position plus nine.

Start	Subsequence	Score
9	YPLPTQPATL	80.000
13	TQPATLGYVL	4.000
7	ALYPLPTQPA	0.450
11	LPTQPATLGY	0.400
14	QPATLGYVLW	0.400
2	YWAMTALYPL	0.400
18	LGYYLWASNI	0.400
4	AMTALYPLPT	0.300
3	WAMTALYPLP	0.090
16	ATLGYVLWAS	0.060

201

TABLE XIX-continued

6	TALYPLPTQP	0.030	5
15	PATLGYVLWA	0.030	
12	PTQPATLGYV	0.020	
17	TLGYVLWASN	0.020	
5	MTALYPLPTQ	0.015	10
8	LYPLPTQPAT	0.010	
1	AYWAMTALYP	0.003	
19	GYVLWASNIS	0.002	
10	PLPTQPATLG	0.002	

TABLE XX

V1-HLA-B35-9mers- 24P4C12		
Each peptide is a portion of SEQ ID NO: 3; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.		
Start	Subsequence	Score
66	YPRNSTGAY	120.000
683	RPYYMSKSL	40.000
15	KPVKYDPSF	40.000
344	ASKAVGQMM	30.000
437	RSVFNLQIY	20.000
679	GSLDRPYYM	20.000
357	YPLVTFVLL	20.000
255	GPLVLVLIL	20.000
631	SPHLNYYWL	20.000
83	KPYLLYFNI	16.000
687	MSKSLLKIL	15.000
396	SPGCEKVPI	12.000
69	NSTGAYCGM	10.000
573	SAKNAPMLL	9.000
533	VARCIMCCF	9.000
213	NARDISVKI	7.200
465	LAGAFASFY	6.000
11	EAYGKPVKY	6.000
333	RIRIAIAL	6.000
343	EASKAVGQM	6.000
489	SAFIRTLRY	6.000
79	ENKDKPYLL	6.000
379	LATSGQPQY	6.000
558	NAYIMIAIY	6.000
630	KSPHLNYYW	5.000
381	TSGQPQYVL	5.000
217	ISVKIFEDF	5.000
132	FSQTVGEVF	5.000
242	LSLLFILL	5.000
406	TSCNPTAHL	5.000
572	VSAKNAFML	5.000
316	LAVLEAILL	4.500
593	KVTDLLFF	4.000
514	IARVILEYI	3.600
287	GASISQLGF	3.000
238	VALVLSLLF	3.000
311	AALIVLAVL	3.000
275	YCWEEYRVL	3.000
253	VAGPLVLVL	3.000
651	IASGFFSVF	3.000
647	GAYVIASGF	3.000
225	FAQSWYWIL	3.000
174	SAPALGRCF	3.000
234	VALGVALVL	3.000
369	IAYWAMTAL	3.000

202

TABLE XX-continued

141	YTKNRNFCL	3.000
494	TLRYHTGSL	3.000
678	NGSLDRPYY	3.000
249	LLRLVAGPL	3.000
117	VSSCPEDPW	2.500
282	VLRDKGASI	2.400
28	KNRSCTDVI	2.400
317	AVLEAILLL	2.000
266	LGVLAYGIY	2.000
363	VLLLCIAY	2.000
267	GVLAYGIYY	2.000
25	GPIKNRSCT	2.000
415	VNSSCPGLM	2.000
50	VVGIVAWLY	2.000
589	VVLDKVTDL	2.000
272	GIYYCWEEY	2.000
188	TPPALPGIT	2.000
432	KGLIQRSVF	2.000
152	VPWNMTVIT	2.000
192	LPGITNDTT	2.000
531	NPVARCIMC	2.000
583	RNIVRVVVL	2.000
366	LICIAYWAM	2.000
546	WCLEKFIKF	2.000
554	FLNRNAYIM	2.000
513	QIARVILEY	2.000
92	FSCILSSNI	2.000
530	QNPVARCIM	2.000
133	SQTVGEVFF	2.000
251	RLVAGPLVL	2.000
409	NPTAHLVNS	2.000
347	AVGQMMSTM	2.000
634	LNYYWLPIM	2.000
621	RIPGLGKDF	2.000
643	TSILGAYVI	2.000
482	IPTFPLISA	2.000
110	CPTPQVCVS	2.000
641	IMTSILGAY	2.000
677	NNGSLDRPY	2.000
421	GLMCVFPQY	2.000
162	LQQELCPSF	2.000
149	LPGVPPWMT	2.000
263	LGVLGVLAY	2.000
167	CPSFLLPSA	2.000
148	CLPGVPWNM	2.000
571	CVSAKNAFM	2.000
112	TPQVCVSSC	2.000
384	QPQYVLWAS	2.000
500	GSLAFGALI	2.000
349	GQMMSTMFY	2.000
653	SGFFSVFGM	2.000
4	KQRDEDDAA	1.800
660	GMCVDTLFL	1.500
30	RSCTDVICC	1.500
430	SSKGLIQRS	1.500

V3-HLA-B35-9mers-
24P4C12

Each peptide is a portion of SEQ ID NO: 7; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.			60
Start	Subsequence	Score	
6	WTNITPPAL	1.000	
9	ITPPALPGI	0.400	
4	FPWTNITPP	0.200	65
2	RCFPWTNIT	0.200	
1	GRCFPWTNI	0.040	
7	TNITPPALP	0.010	
8	NITPPALPG	0.010	
3	CFPWTNITP	0.001	
5	PWTNITPPA	0.001	

203

TABLE XX-continued

V5-HLA-B35-9mers- 24P4C12		
Each peptide is a portion of SEQ ID NO: 11; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.		
Start	Subsequence	Score
3	EAILLLVLI	1.200
5	ILLVLIFL	1.000
4	AILLLVLIF	1.000
9	VLIFLRQRI	0.400
2	LEAILLLVL	0.100
1	VLEAILLLV	0.060
6	LLLVLIFLR	0.010
7	LLVLIFLRQ	0.010
8	LVLIFLRQR	0.010
V6-HLA-B35-9mers- 24P4C12		
Each peptide is a portion of SEQ ID NO: 13; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.		
Start	Subsequence	Score
5	KGLIPRSVF	2.000
3	SSKGLIPRS	1.500
7	LIPRSVFNL	1.000
8	IPRSVFNLQ	0.600
6	GLIPRSVFN	0.100
2	YSSKGLIPR	0.050
4	SKGLIPRSV	0.020
9	PRSVFNLQI	0.004
1	GYSSKGLIP	0.001
V7-HLA-B35-9mers- 24P4C12		
Each peptide is a portion of SEQ ID NO: 15; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.		
Start	Subsequence	Score
8	AVGQMMSTM	2.000
5	ILVAVGQMM	2.000
4	WILVAVGQM	2.000
7	VAVGQMMST	0.300
6	LVAVGQMMS	0.100
1	SWYWILVAV	0.020
3	YWILVAVGQ	0.001
2	WYWILVAVG	0.001
V8-HLA-B35-9mers- 24P4C12		
Each peptide is a portion of SEQ ID NO: 17; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.		
Start	Subsequence	Score
10	NPITPTGHV	4.000
13	TPTGHVFPQT	2.000
19	FQTSILGAY	2.000
5	LPIMRNPIT	2.000
15	TGHVFQTSI	0.400
4	WLPIMRNPI	0.400
7	IMRNPITPT	0.300

204

TABLE XX-continued

20	QTSILGAYV	0.200
11	PITPTGHVF	0.100
16	GHVFQTSIL	0.100
9	RNPITPTGH	0.020
2	YYWLPIMRN	0.010
18	VFQTSILGA	0.010
14	PTGHVFQTS	0.010
17	HVFQTSILG	0.010
12	ITPTGHVFQ	0.010
6	PIMRNPITP	0.001
3	YWLPIMRNP	0.001
8	MRNPITPTG	0.001
1	NYYWLPIMR	0.001

V9-HLA-B35-9mers-
24P4C12

Each peptide is a portion of SEQ ID
NO: 19; each start position is
specified, the length of peptide is 9
amino acids, and the end position for
each peptide is the start position plus
eight.

Start	Subsequence	Score
13	QPATLGYVL	20.000
2	WAMTALYPL	3.000
8	YPLPTQPAT	2.000
11	PTQPATLGY	0.200
12	TQPATLGYV	0.200
10	LPTQPATLG	0.200
14	PATLGYVLW	0.150
15	ATLGYVLWA	0.100
4	MTALYPLPT	0.100
16	TLGYVLWAS	0.100
17	LGYYVLWAS	0.100
9	PLPTQPATL	0.100
18	GYVLWASNI	0.040
5	TALYPLPTQ	0.030
7	LYPLPTQPA	0.010
3	AMTALYPLP	0.010
6	ALYPLPTQP	0.010
1	YWAMTALYP	0.001

TABLE XXI

V1-HLA-B35-10mers- 24P4C12		
Each peptide is a portion of SEQ ID NO: 3; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.		
Start	Subsequence	Score
478	KPQDIPTFPL	80.000
83	KPYLLYFNIF	40.000
683	RPYYMSKSLL	40.000
4	KQRDEDEAY	36.000
123	DPWTVGKNEF	20.000
482	IPTFPLISAF	20.000
357	YPLVTFVLLL	20.000
213	NARDISVKIF	18.000
573	SAKNAPMLLM	18.000
346	KAVGQMMSTM	12.000
79	ENKDKPYLLY	12.000
652	ASGFFSVFGM	10.000
488	ISAFIRTLRY	10.000
132	FSQTVGEV FY	10.000
175	APALGRCFPW	10.000
630	KSPHLNYYWL	10.000
192	LPGITNDTTI	8.000
551	FIKFLNRNAY	6.000
625	LGKDFKSPHL	6.000
331	RQRIRIAIAL	6.000

205

TABLE XXI-continued

343	EASKAVGQMM	6.000
60	DPRQVLYPRN	6.000
66	YPRNSTGAYC	6.000
369	IAYWAMTALY	6.000
572	VSAKNAPMLL	5.000
227	QSWYWILVAL	5.000
500	GSLAFGALIL	5.000
417	SSCPGLMCVF	5.000
290	ISQLGPTTNL	5.000
76	GMGENKDKPY	4.000
68	RNSTGAYCGM	4.000
317	AVLEAILLLM	4.000
557	RNAYIMIAIY	4.000
149	LPGVPMNMTV	4.000
676	RNNGSLDRPY	4.000
310	LAALIVLAVL	3.000
316	LAVLEAILLL	3.000
320	EAILLLMLIF	3.000
467	GAFASFYWAF	3.000
395	SSPGCEKVPI	3.000
647	GAYVIASGFF	3.000
677	NNGSLDRPY	3.000
502	LAFGALILTL	3.000
430	SSKGLIQRSV	3.000
381	TSGQPQYVLW	2.500
362	FVLLLCIAY	2.000
39	VLFLLFILGY	2.000
188	TPPALPGITN	2.000
152	VPWNMTVITS	2.000
348	VGQMMSTMFY	2.000
31	SCTDVICCVL	2.000
384	QPQYVLWASN	2.000
409	NPTAHLVNSS	2.000
613	LSFFFFSGRI	2.000
220	KIPEDFAQSW	2.000
110	CPTPQVCVSS	2.000
546	WCLEKFIKFL	2.000
271	YGIYYCWEEY	2.000
30	RSCTDVICCV	2.000
172	LPSAPALGRC	2.000
162	LQQELCPSFL	2.000
396	SPGCEKVPIN	2.000
266	LGVLAYGIYY	2.000
402	VPINTSCNPT	2.000
378	YLATSGQPQY	2.000
365	LLICIAYWAM	2.000
293	LGFTTNLSAY	2.000
262	ILGVLGVLAY	2.000
286	KGASISQLGF	2.000
529	VQNPVARCIM	2.000
678	NGSLDRPYM	2.000
49	IVVGIVAWLY	2.000
147	FCLPGVPWNM	2.000
265	VLGVLAYGIY	2.000
304	SVQETWLAAL	2.000
464	VLAGAFASFY	2.000
20	DPSFRGPIKN	2.000
661	MCVDTLFLCF	2.000
92	FSCILSSNII	2.000
512	VQIARVILEY	2.000
182	FPWTNVTTPA	2.000
639	LPIMTSILGA	2.000
570	FCVSAKNAFM	2.000
493	RTLRYHTGSL	2.000
633	HLNYYWLPIM	2.000
531	NPVARCIMCC	2.000
622	IPGLGKDFKS	2.000
485	FPLISAFIRT	2.000
542	KCLWCLEKF	2.000
589	VVLDKVTDLL	2.000
517	VILEYIDHKL	2.000

206

TABLE XXI-continued

414	LVNSSCPGLM	2.000
344	ASKAVGQMMS	1.500
465	LAGAFASFYW	1.500
300	SAYQSVQETW	1.500
659	FGMCVDTLFL	1.500
315	VLAVLEAILL	1.500
118	SSCPEDPWTV	1.500
576	NAFMLLMRNI	1.200
435	IQRSVFNLQI	1.200
V3-HLA-B35-10mers- 24P4C12		
Each peptide is a portion of SEQ ID NO: 7; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.		
Start	Subsequence	Score
5	FPWTNITPPA	2.000
1	LGRCFPWTNI	1.200
9	NITPPALPGI	0.400
10	ITPPALPGIT	0.100
3	RCFPWTNITP	0.020
8	TNITPPALPG	0.010
6	PWTNITPPAL	0.010
7	WTNITPPALP	0.010
2	GRCFPWTNIT	0.010
4	CFPWTNITPP	0.001
V5-HLA-B35-10mers- 24P4C12		
Each peptide is a portion of SEQ ID NO: 11; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.		
Start	Subsequence	Score
4	EAILLLVLIF	3.000
5	AILLLVLIFL	1.000
9	LVLIFLRQRI	0.400
1	AVLEAILLLV	0.400
2	VLEAILLLVL	0.300
3	LEAILLLVLI	0.040
6	ILLVLIFLIR	0.010
10	VLIFLRQRI	0.010
7	LLVLIFLRQ	0.010
8	LLVLIFLRQR	0.010
V6-HLA-B35-10mers- 24P4C12		
Each peptide is a portion of SEQ ID NO: 13; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.		
Start	Subsequence	Score
9	IPRSVFNLQI	24.000
4	SSKGLIPRSV	3.000
7	GLIPRSVFNL	1.000
3	YSSKGLIPRS	0.500
6	KGLIPRSVFN	0.200
5	SKGLIPRSVF	0.100
10	PRSVFNLQIY	0.020
8	LIPRSVFNLQ	0.010
1	QGYSSKGLIP	0.010
2	GYSSKGLIPR	0.001

207

TABLE XXI-continued

V7-HLA-B35-10mers- 24P4C12		
Each peptide is a portion of SEQ ID NO: 15; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.		
Start	Subsequence	Score
8	VAVGQMMSTM	6.000
5	WILVAVGQMM	2.000
9	AVGQMMSTM	1.000
1	QSWYWLIVAV	1.000
4	YWILVAVGQM	0.200
6	ILVAVGQMMS	0.100
7	LVAVGQMMST	0.100
2	SWYMILVAVG	0.001
3	WYWILVAVGQ	0.001
V8-HLA-B35-10mers- 24P4C12		
Each peptide is a portion of SEQ ID NO: 17; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.		
Start	Subsequence	Score
11	NPITPTGHVF	20.000
14	TPTGHVFPQTS	2.000
16	TGHVFPQTSIL	1.000
21	QTSILGAYVI	0.400
10	RNPITPTGHV	0.400
6	LPIMRNPITP	0.200
20	FQTSILGAVV	0.200
19	VFQTSILGAY	0.200
13	ITPTGHVFPQT	0.100
18	HVFQTSILGA	0.100
5	WLPIMRNPIT	0.100
15	PTGHVFPQTSI	0.040
4	YWLPIMRNPI	0.040
8	IMRNPITPTG	0.030
2	NYWLPIMRN	0.010
7	PIMRNPITPT	0.010
1	LNYYWLPIMR	0.010
17	GHVFPQTSILG	0.001
3	YYWLPIMRNP	0.001
12	PITPTGHVFPQ	0.001
9	MRNPITPTGH	0.001
V9-HLA-B35-10mers- 24P4C12		
Each peptide is a portion of SEQ ID NO: 19; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.		
Start	Subsequence	Score
2	YWAMTALYPL	1.000
13	TQPATLGYVL	1.000
9	YPLPTQPATL	1.000
18	LGYYLWASNI	1.000
14	QPATLGYVLW	0.500
11	LPTQPATLGY	0.200
16	ATLGYVLWAS	0.150
19	GYVLWASNIS	0.100
4	AMTALYPLPT	0.100
8	LYPLPTQPAT	0.100
17	TLGYVLWASN	0.100
7	ALYPLPTQPA	0.100
3	WAMTALYPLP	0.050
12	PTQPATLGYV	0.020
6	TALYPLPTQP	0.010

208

TABLE XXI-continued

5	MTALYPLPTQ	0.010
15	PATLGYVLWA	0.010
1	AYWAMTALYP	0.010
10	PLPTQPATLG	0.005

10 Tables XXII-XLIX:

TABLE XXII

V1-HLA-A1-9mers- 24P4C12										
Each peptide is a portion of SEQ ID NO: 3; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.										
Pos	1	2	3	4	5	6	7	8	9	score
80	N	<u>K</u>	D	K	P	Y	<u>L</u>	L	Y	34
58	Y	<u>G</u>	D	P	R	Q	<u>V</u>	L	Y	33
222	F	<u>E</u>	D	F	A	Q	<u>S</u>	W	Y	26
5	Q	<u>R</u>	D	E	D	D	<u>E</u>	A	Y	25
77	M	<u>G</u>	E	N	K	D	<u>K</u>	P	Y	25
263	L	<u>G</u>	V	L	G	V	<u>L</u>	A	Y	24
489	S	<u>A</u>	F	I	R	T	<u>L</u>	R	Y	23
513	Q	<u>I</u>	A	R	V	I	<u>L</u>	E	Y	23
628	D	<u>F</u>	K	S	P	H	<u>L</u>	N	Y	22
40	L	<u>F</u>	L	L	F	I	<u>L</u>	G	Y	21
267	G	<u>V</u>	L	A	Y	G	<u>I</u>	Y	Y	21
363	V	<u>L</u>	L	L	I	C	<u>I</u>	A	Y	21
421	G	<u>L</u>	M	C	V	F	<u>Q</u>	G	Y	21
50	V	<u>V</u>	G	I	V	A	<u>W</u>	L	Y	20
318	V	<u>L</u>	E	A	I	L	<u>L</u>	L	M	20
629	F	<u>K</u>	S	P	H	L	<u>N</u>	Y	Y	20
133	S	<u>Q</u>	T	V	G	E	<u>V</u>	F	Y	19
437	R	<u>S</u>	V	F	N	L	<u>Q</u>	I	Y	19
662	C	<u>V</u>	D	T	L	F	<u>L</u>	C	F	19
11	E	<u>A</u>	Y	G	K	P	<u>V</u>	K	Y	18
370	A	<u>Y</u>	W	A	M	T	<u>A</u>	L	Y	18
18	K	<u>Y</u>	D	P	S	F	<u>R</u>	G	P	17
32	C	<u>T</u>	D	V	I	C	<u>C</u>	V	L	17
66	Y	<u>P</u>	R	N	S	T	<u>G</u>	A	Y	17
277	W	<u>E</u>	E	Y	R	V	<u>L</u>	R	D	17
379	L	<u>A</u>	T	S	G	Q	<u>P</u>	Q	Y	17
594	V	<u>T</u>	D	L	L	L	<u>F</u>	F	G	17
165	E	<u>L</u>	C	P	S	F	<u>L</u>	L	P	16
353	S	<u>T</u>	M	F	Y	P	<u>L</u>	V	T	16
398	G	<u>C</u>	E	K	V	P	<u>I</u>	N	T	16
552	I	<u>K</u>	F	L	N	R	<u>N</u>	A	Y	16
590	V	<u>L</u>	D	K	V	T	<u>D</u>	L	L	16
678	N	<u>G</u>	S	L	D	R	<u>P</u>	Y	Y	16
V3-HLA-A1-9mers- 24P4C12										
Each peptide is a portion of SEQ ID NO: 7; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.										
Pos	1	2	3	4	5	6	7	8	9	score
8	N	<u>I</u>	T	P	P	A	<u>L</u>	P	G	11
9	I	<u>T</u>	P	P	A	L	<u>P</u>	G	I	10
6	W	<u>T</u>	N	I	T	P	<u>P</u>	A	L	6
3	C	<u>F</u>	P	W	T	N	<u>I</u>	T	P	5

209

TABLE XXII-continued

V5-HLA-A1-9mers- 24P4C12										
Each peptide is a portion of SEQ ID NO: 11; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.										
Pos	1	2	3	4	5	6	7	8	9	score
1	V	<u>L</u>	E	A	I	L	<u>L</u>	L	V	20
7	L	<u>L</u>	V	L	I	F	<u>L</u>	R	Q	10

V6-HLA-A1-9mers- 24P4C12										
Each peptide is a portion of SEQ ID NO: 13; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.										
Pos	1	2	3	4	5	6	7	8	9	score
2	Y	<u>S</u>	S	K	G	L	<u>I</u>	P	R	12
1	G	<u>Y</u>	S	S	K	G	<u>L</u>	I	P	7
3	S	<u>S</u>	K	G	L	I	<u>P</u>	R	S	7
8	I	<u>P</u>	R	S	V	F	<u>N</u>	L	Q	7
9	P	<u>R</u>	S	V	F	N	<u>L</u>	Q	I	7
6	G	<u>L</u>	I	P	R	S	<u>V</u>	F	N	5

V7-HLA-A1-9mers- 24P4C12										
Each peptide is a portion of SEQ ID NO: 15; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.										
Pos	1	2	3	4	5	6	7	8	9	score
5	I	<u>L</u>	V	A	V	G	<u>Q</u>	M	M	5
3	Y	<u>W</u>	I	L	V	A	<u>V</u>	G	Q	4
7	V	<u>A</u>	V	G	Q	M	<u>M</u>	S	T	4
6	L	<u>V</u>	A	V	G	Q	<u>M</u>	M	S	3
1	S	<u>W</u>	Y	W	I	L	<u>V</u>	A	V	2
2	W	<u>Y</u>	W	I	L	V	<u>A</u>	V	G	2

V8-HLA-A1-9mers- 24P4C12										
Each peptide is a portion of SEQ ID NO: 17; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.										
Pos	1	2	3	4	5	6	7	8	9	score
19	F	<u>Q</u>	T	S	I	L	<u>G</u>	A	Y	16
14	P	<u>T</u>	G	H	V	F	<u>Q</u>	T	S	11
12	I	<u>T</u>	P	T	G	H	<u>V</u>	F	Q	8
18	V	<u>F</u>	Q	T	S	I	<u>L</u>	G	A	7
20	Q	<u>T</u>	S	I	L	G	<u>A</u>	Y	V	7

210

TABLE XXII-continued

V9-HLA-A1-9mers- 24P4C12										
Each peptide is a portion of SEQ ID NO: 19; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.										
Pos	1	2	3	4	5	6	7	8	9	score
11	P	<u>T</u>	Q	P	A	T	<u>L</u>	G	Y	31
15	A	<u>T</u>	L	G	Y	V	<u>L</u>	W	A	16

TABLE XXIII

V1-HLA-A0201- 9mers-24P4C12										
Each peptide is a portion of SEQ ID NO: 3; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.										
Pos	1	2	3	4	5	6	7	8	9	score
260	V	L	I	L	G	<u>V</u>	L	G	V	31
244	L	L	F	I	L	<u>L</u>	L	R	L	29
580	L	L	M	R	N	<u>I</u>	V	R	V	29
95	I	L	S	S	N	<u>I</u>	I	S	V	28
204	G	I	S	G	L	<u>I</u>	D	S	L	28
261	L	I	L	G	V	<u>L</u>	G	V	L	28
322	I	L	L	L	M	<u>L</u>	I	F	L	28
506	A	L	I	L	T	<u>L</u>	V	Q	I	28
170	F	L	L	P	S	<u>A</u>	P	A	L	27
252	L	V	A	G	P	<u>L</u>	V	L	V	27
449	G	L	F	W	T	<u>L</u>	N	W	V	27
487	L	I	S	A	F	<u>I</u>	R	T	L	27
604	L	L	V	V	G	<u>G</u>	V	G	V	27
45	I	L	G	Y	I	<u>V</u>	V	G	I	26
232	I	L	V	A	L	<u>G</u>	V	A	L	26
233	L	V	A	L	G	<u>V</u>	A	L	V	26
315	V	L	A	V	L	<u>E</u>	A	I	L	26
501	S	L	A	F	G	<u>A</u>	L	I	L	26
521	Y	I	D	H	K	<u>L</u>	R	G	V	26
42	L	L	F	I	L	<u>G</u>	Y	I	V	25
107	G	L	Q	C	P	<u>T</u>	P	Q	V	25
200	T	I	Q	Q	G	<u>I</u>	S	G	L	25
211	S	L	N	A	R	<u>D</u>	I	S	V	25
239	A	L	V	L	S	<u>L</u>	L	F	I	25
257	L	V	L	V	L	<u>I</u>	L	G	V	25
258	V	L	V	L	I	<u>L</u>	G	V	L	25
282	V	L	R	D	K	<u>G</u>	A	S	I	25
317	A	V	L	E	A	<u>I</u>	L	L	L	25
457	V	L	A	L	G	<u>Q</u>	C	V	L	25
598	L	L	F	F	G	<u>K</u>	L	L	V	25
650	V	I	A	S	G	<u>F</u>	F	S	V	25
686	Y	M	S	K	S	<u>L</u>	L	K	I	25
41	F	L	L	F	I	<u>L</u>	G	Y	I	24
49	I	V	V	G	I	<u>V</u>	A	W	L	24
310	L	A	A	L	I	<u>V</u>	L	A	V	24
311	A	A	L	I	V	<u>L</u>	A	V	L	24
333	R	I	R	I	A	<u>I</u>	A	L	L	24
434	L	I	Q	R	S	<u>V</u>	F	N	L	24
509	L	T	L	V	Q	<u>I</u>	A	R	V	24
525	K	L	R	G	V	<u>Q</u>	N	P	V	24
564	A	I	Y	G	K	<u>N</u>	F	C	V	24
581	L	M	R	N	I	<u>V</u>	R	V	V	24
596	D	L	L	L	F	<u>F</u>	G	K	L	24
605	L	V	V	G	G	<u>V</u>	G	V	L	24
35	V	I	C	C	V	<u>L</u>	F	L	L	23
56	W	L	Y	G	D	<u>P</u>	R	Q	V	23
240	L	V	L	S	L	<u>L</u>	F	I	L	23
251	R	L	V	A	G	<u>P</u>	L	V	L	23

211

TABLE XXIII-continued

253	V	A	G	P	L	<u>V</u>	L	V	L	23
309	W	L	A	A	L	<u>I</u>	V	L	A	23
340	L	L	K	E	A	<u>S</u>	K	A	V	23
358	P	L	V	T	F	<u>V</u>	L	L	L	23
494	T	L	R	Y	H	<u>T</u>	G	S	L	23
518	I	L	E	Y	I	<u>D</u>	H	K	L	23
547	C	L	E	K	F	<u>I</u>	K	F	L	23
589	V	V	L	D	K	<u>V</u>	T	D	L	23
590	V	L	D	K	V	<u>T</u>	D	L	L	23
597	L	L	L	F	F	<u>G</u>	K	L	L	23
100	I	I	S	V	A	<u>E</u>	N	G	L	22
241	V	L	S	L	L	<u>F</u>	I	L	L	22
248	L	L	L	R	L	<u>V</u>	A	G	P	22
249	L	L	R	L	V	<u>A</u>	G	P	L	22
265	V	L	G	V	L	<u>A</u>	Y	G	I	22
446	G	V	L	G	L	<u>F</u>	W	T	L	22
452	W	T	L	N	W	<u>V</u>	L	A	L	22
578	F	M	L	L	M	<u>R</u>	N	I	V	22
638	W	L	P	I	M	<u>T</u>	S	I	L	22
660	G	M	C	V	D	<u>T</u>	L	F	L	22
158	V	I	T	S	L	<u>Q</u>	Q	E	L	21
187	V	T	P	S	A	<u>L</u>	P	G	I	21
191	A	L	P	G	I	<u>T</u>	N	D	T	21
237	G	V	A	L	V	<u>L</u>	S	L	L	21
247	I	L	L	R	L	<u>V</u>	A	G		21
313	L	I	V	L	A	<u>V</u>	L	E	A	21
314	I	V	L	A	V	<u>L</u>	E	A	I	21
442	L	Q	I	Y	G	<u>V</u>	L	G	L	21
507	L	I	L	T	L	<u>V</u>	Q	I	A	21
537	I	M	C	C	F	<u>K</u>	C	C	L	21
599	L	F	F	G	K	<u>L</u>	L	V	V	21
693	K	I	L	G	K	<u>K</u>	N	E	A	21
34	D	V	I	C	C	<u>V</u>	L	F	L	20
38	C	V	L	F	L	<u>L</u>	F	I	L	20
44	F	I	L	G	Y	<u>I</u>	V	V	G	20
207	G	L	I	D	S	<u>L</u>	N	A	R	20
228	S	W	Y	W	I	<u>L</u>	V	A	L	20
234	V	A	L	G	V	<u>A</u>	L	V	L	20
236	L	G	V	A	L	<u>V</u>	L	S	L	20
242	L	S	L	L	F	<u>I</u>	L	L	L	20
319	L	E	A	I	L	<u>L</u>	L	M	L	20
326	M	L	I	F	L	<u>R</u>	Q	R	I	20
339	A	L	L	K	E	<u>A</u>	S	K	A	20
364	L	L	L	I	C	<u>I</u>	A	Y	W	20
417	S	S	C	P	G	<u>L</u>	M	C	V	20
503	A	F	G	A	L	<u>I</u>	L	T	L	20
633	H	L	N	Y	Y	<u>W</u>	L	P	I	20
644	S	I	L	G	A	<u>V</u>	V	I	A	20
673	D	L	E	R	N	<u>N</u>	G	S	L	20
690	S	L	L	K	I	<u>L</u>	G	K	K	20
48	Y	I	V	V	G	<u>I</u>	V	A	W	19
245	L	F	I	L	L	<u>L</u>	R	L	V	19
255	G	P	L	V	L	<u>V</u>	L	I	L	19
262	I	L	G	V	L	<u>G</u>	V	L	A	19
268	V	L	A	Y	G	<u>I</u>	Y	Y	C	19
291	S	Q	L	G	F	<u>T</u>	N	L		19
318	V	L	E	A	I	<u>L</u>	L	L	M	19
323	L	L	L	M	L	<u>I</u>	F	L	R	19
329	F	L	R	Q	R	<u>I</u>	R	I	A	19
351	M	M	S	T	M	<u>E</u>	Y	P	L	19
365	L	L	I	C	I	<u>A</u>	Y	W	A	19
414	L	V	N	S	S	<u>C</u>	P	G	L	19
464	V	L	A	G	A	<u>E</u>	A	S	F	19
544	C	L	W	C	L	<u>E</u>	K	F	I	19
617	F	F	S	G	R	<u>I</u>	P	G	L	19
666	L	F	L	C	F	<u>L</u>	E	D	L	19
86	L	L	Y	F	N	<u>I</u>	F	S	C	18
231	W	I	L	V	A	<u>L</u>	G	V	A	18
235	A	L	G	V	A	<u>L</u>	V	L	S	18
243	S	L	L	F	I	<u>L</u>	L	L	R	18
336	I	A	I	A	L	<u>L</u>	K	E	A	18
355	M	F	Y	P	A	<u>V</u>	T	F	V	18
369	I	A	Y	W	A	<u>M</u>	T	A	L	18
380	A	T	S	G	Q	<u>P</u>	Q	Y	V	18
394	I	S	S	P	G	<u>C</u>	E	K	V	18
439	V	F	N	L	Q	<u>I</u>	Y	G	V	18
459	A	L	G	Q	C	<u>V</u>	L	A	G	18
510	T	L	V	Q	I	<u>A</u>	R	V	I	18
511	L	V	Q	I	A	<u>R</u>	V	I	L	18
514	I	A	R	V	I	<u>L</u>	E	Y	I	18

212

TABLE XXIII-continued

517	V	I	L	E	Y	<u>I</u>	D	H	K	18
583	R	N	I	V	R	<u>V</u>	V	V	L	18
602	G	K	L	L	V	<u>V</u>	G	G	V	18
645	I	L	G	A	Y	<u>V</u>	I	A	S	18
46	L	G	Y	I	V	<u>V</u>	G	I	V	17
128	G	K	N	E	F	<u>S</u>	Q	T	V	17
154	W	N	M	T	V	<u>I</u>	T	S	L	17
177	A	L	G	R	C	<u>F</u>	P	W	T	17
184	W	T	N	V	T	<u>P</u>	P	A	L	17
213	N	A	R	D	I	<u>S</u>	V	K	I	17
246	F	I	L	L	L	<u>R</u>	L	V	A	17
289	S	I	S	Q	L	<u>G</u>	F	T	T	17
300	S	A	Y	Q	S	<u>V</u>	Q	E	T	17
305	V	Q	E	T	W	<u>L</u>	A	A	L	17
312	A	L	I	V	L	<u>A</u>	V	L	E	17
325	L	M	L	I	F	<u>L</u>	R	Q	R	17
335	R	I	A	I	A	<u>L</u>	L	K	E	17
354	T	M	F	Y	P	<u>L</u>	V	T	F	17
359	L	V	T	F	V	<u>L</u>	L	L	I	17
453	T	L	N	W	V	<u>L</u>	A	L	G	17
456	W	V	L	A	L	<u>G</u>	Q	C	V	17
502	L	A	F	G	A	<u>L</u>	I	L	T	17
504	F	G	A	L	I	<u>L</u>	T	L	V	17
513	Q	I	A	R	V	<u>I</u>	L	E	Y	17
554	F	L	N	R	N	<u>A</u>	Y	I	M	17
560	Y	I	M	I	A	<u>I</u>	Y	G	K	17
586	V	R	V	V	V	<u>L</u>	D	K	V	17
642	M	T	S	I	L	<u>G</u>	A	Y	V	17
658	V	F	G	M	C	<u>V</u>	D	T	L	17
31	S	C	T	D	V	<u>I</u>	C	C	V	16
43	L	F	I	L	G	<u>Y</u>	I	V	V	16
64	V	L	Y	P	R	<u>N</u>	S	T	G	16
90	N	I	F	S	C	<u>I</u>	L	S	S	16
119	S	C	P	E	D	<u>P</u>	W	T	V	16
144	N	R	N	F	C	<u>L</u>	P	G	V	16
148	C	L	P	G	V	<u>P</u>	W	N	M	16
161	S	L	Q	Q	E	<u>L</u>	C	P	S	16
230	Y	W	I	L	V	<u>A</u>	L	G	V	16
254	A	G	P	L	V	<u>L</u>	V	L	I	16
308	T	W	L	A	A	<u>L</u>	I	V	L	16
316	L	A	V	L	E	<u>A</u>	I	L	L	16
320	E	A	I	L	L	<u>L</u>	M	L	I	16
357	Y	P	L	V	T	<u>F</u>	V	L	L	16
362	F	V	L	L	L	<u>I</u>	C	I	A	16
373	A	M	T	A	L	<u>Y</u>	L	A	T	16
376	A	L	Y	L	A	<u>T</u>	S	G	Q	16
407	S	C	N	P	T	<u>A</u>	H	L	V	16
458	L	A	L	G	Q	<u>C</u>	V	L	A	16
637	Y	W	L	P	I	<u>M</u>	T	S	I	16
640	P	I	M	T	S	<u>I</u>	L	G	A	16
52	G	I	V	A	W	<u>L</u>	Y	G	D	15
141	Y	T	K	N	R	<u>N</u>	F	C	L	15
225	F	A	Q	S	W	<u>Y</u>	W	I	L	15
250	L	R	L	V	A	<u>G</u>	P	L	V	15
264	G	V	L	G	V	<u>L</u>	A	Y	G	15
275	Y	C	W	E	E	<u>Y</u>	R	V	L	15
366	L	I	C	I	A	<u>Y</u>	W	A	M	15
368	C	I	A	Y	W	<u>A</u>	M	T	A	15
371	Y	W	A	M	T	<u>A</u>	L	Y	L	15
374	M	T	A	L	Y	<u>L</u>	A	T	S	15
406	T	S	C	N	P	<u>T</u>	A	H	L	15
433	G	L	I	Q	R	<u>S</u>	V	F	N	15
443	Q	I	Y	G	V	<u>L</u>	G	L	F	15
491	F	I	R	T	L	<u>R</u>	Y	H	T	15
573	S	A	K	N	A	<u>F</u>	M	L	T	15
657	S	V	F	G	M	<u>C</u>	V	D	T	15
663	V	D	T	L	F	<u>L</u>	C	F	L	15

213

TABLE XXIII--continued

V3-HLA-A0201- 9mers-24P4C12										
Each peptide is a portion of SEQ ID NO: 7; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.										
Pos	1	2	3	4	5	6	7	8	9	score
9	I	T	P	P	A	<u>L</u>	P	G	I	22
6	W	T	N	I	T	<u>P</u>	P	A	L	17
8	N	I	T	P	P	<u>A</u>	L	P	G	11
2	R	C	F	P	W	<u>T</u>	N	I	T	10
V5-HLA-A0201- 9mers-24P4C12										
Each peptide is a portion of SEQ ID NO: 11; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.										
Pos	1	2	3	4	5	6	7	8	9	score
5	I	L	L	L	V	<u>L</u>	I	F	L	28
1	V	L	E	A	I	<u>L</u>	L	L	V	25
9	V	L	I	F	L	<u>R</u>	Q	R	I	21
2	L	E	A	I	L	<u>L</u>	L	V	L	20
6	L	L	L	V	L	<u>I</u>	F	L	R	19
3	E	A	I	L	L	<u>L</u>	V	L	I	18
4	A	I	L	L	L	<u>V</u>	L	I	F	18
7	L	L	V	L	I	<u>F</u>	L	R	Q	13
8	L	V	L	I	F	<u>L</u>	R	Q	R	13
V6-HLA-A0201- 9mers-24P4C12										
Each peptide is a portion of SEQ ID NO: 13; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.										
Pos	1	2	3	4	5	6	7	8	9	score
2	Y	S	S	K	G	<u>L</u>	I	P	R	12
1	G	Y	S	S	K	<u>G</u>	L	I	P	7
3	S	S	K	G	L	<u>I</u>	P	R	S	7
8	I	P	R	S	V	<u>F</u>	N	L	Q	7
9	P	R	S	V	F	<u>N</u>	L	Q	I	7
6	G	L	I	P	R	<u>S</u>	V	F	N	5
V7-HLA-A0201- 9mers-24P4C12										
Each peptide is a portion of SEQ ID NO: 15; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.										
Pos	1	2	3	4	5	6	7	8	9	score
1	S	W	Y	W	I	<u>L</u>	V	A	V	20
4	W	I	L	V	A	<u>V</u>	G	Q	M	18
5	I	L	V	A	V	<u>G</u>	Q	M	M	16
7	V	A	V	G	Q	<u>M</u>	M	S	T	13
8	A	V	G	Q	M	<u>M</u>	S	T	M	12
6	L	V	A	V	G	<u>Q</u>	M	M	S	10

214

TABLE XXIII--continued

	V8-HLA-A0201- 9mers-24P4C12										
5	Each peptide is a portion of SEQ ID NO: 17; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.										
10	Pos	1	2	3	4	5	6	7	8	9	score
	4	W	L	P	I	M	<u>R</u>	N	P	I	19
	7	I	M	R	N	P	<u>I</u>	T	P	T	19
	20	Q	T	S	I	L	<u>G</u>	A	Y	V	17
15	10	N	P	I	T	P	<u>T</u>	G	H	V	15
	16	G	H	V	F	Q	<u>T</u>	S	I	L	12
	15	T	G	H	V	F	<u>Q</u>	T	S	I	11
	18	V	F	Q	T	S	<u>I</u>	L	G	A	11
	12	I	T	P	T	G	<u>H</u>	V	F	Q	10
	5	L	P	I	M	R	<u>N</u>	P	I	T	9
20	13	T	P	T	G	H	<u>V</u>	F	Q	T	9
	V9-HLA-A0201- 9mers-24P4C12										
25	Each peptide is a portion of SEQ ID NO: 19; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.										
	Pos	1	2	3	4	5	6	7	8	9	score
30	9	P	L	P	T	Q	<u>P</u>	A	T	L	21
	2	W	A	M	T	A	<u>L</u>	Y	P	L	20
	15	A	T	L	G	Y	<u>V</u>	L	W	A	20
	6	A	L	Y	P	L	<u>P</u>	T	Q	P	16
	12	T	Q	P	A	T	<u>L</u>	G	Y	V	14
35	13	Q	P	A	T	L	<u>G</u>	Y	V	L	14
	16	T	L	G	Y	V	<u>L</u>	W	A	S	14
	5	T	A	L	Y	P	<u>L</u>	P	T	Q	13
	4	M	T	A	L	Y	<u>P</u>	L	P	T	12
	8	Y	P	L	P	T	<u>Q</u>	P	A	T	12
40	3	A	M	T	A	L	<u>Y</u>	P	L	P	11
	TableXXIV-V1-HLA-A0203-9mers-24P4C12										
45	Pos	1234567890								score	
	NoResultsFound.										
	TableXXIV-V3-HLA-A0203-9mers-24P4C12										
50	Pos	1234567890								score	
	NoResultsFound.										
	TableXXIV-V5-HLA-A0203-9mers-24P4C12										
55	Pos	1234567890								score	
	NoResultsFound.										
	TableXXIV-V6-HLA-A0203-9mers-24P4C12										
60	Pos	1234567890								score	
	NoResultsFound.										
	TableXXIV-V7-HLA-A0203-9mers-24P4C12										
65	Pos	1234567890								score	

215

-continued

NoResultsFound.										
TableXXIV-V8-HLA-A0203-9mers-24P4C12										
Pos	1234567890								score	
NoResultsFound.										
TableXXIV-V9-HLA-A0203-9mers-24P4C12										
Pos	1234567890								score	
NoResultsFound.										
TABLE XXV										
V1-HLA-A3-9mers-24P										
Each peptide is a portion of SEQ ID NO: 3; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.										
Pos	1	2	3	4	5	6	7	8	9	score
585	I	V	R	V	V	V	L	D	K	29
424	C	V	F	Q	G	Y	S	S	K	27
4	V	L	Y	P	R	N	S	T	G	26
135	T	V	G	E	V	F	Y	T	K	26
251	R	L	V	A	G	P	L	V	L	26
506	A	L	I	L	T	L	V	Q	I	24
513	Q	I	A	R	V	I	L	E	Y	24
603	K	L	L	V	V	G	G	V	G	24
690	S	L	L	K	I	L	G	K	K	24
267	G	V	L	A	Y	G	I	Y	Y	23
282	V	L	R	D	K	G	A	S	I	23
312	A	L	I	V	L	A	V	L	E	23
334	I	R	I	A	I	A	L	L	K	23
102	S	V	A	E	N	G	L	Q	C	22
232	I	L	V	A	L	G	V	A	L	22
247	I	L	L	L	R	L	V	A	G	22
443	Q	I	Y	G	V	L	G	L	F	22
464	V	L	A	G	A	F	A	S	F	22
516	R	V	I	L	E	Y	I	D	H	22
579	M	L	L	M	R	N	I	V	R	22
50	V	V	G	I	V	A	W	L	Y	21
212	L	N	A	R	D	I	S	V	K	21
281	R	V	L	R	D	K	G	A	S	21
321	A	I	L	L	L	M	L	I	F	21
338	I	A	L	L	K	E	A	S	K	21
339	A	L	L	K	E	A	S	K	A	21
376	A	L	Y	L	A	T	S	G	Q	21
393	N	I	S	S	P	G	C	E	K	21
517	V	I	L	E	Y	I	D	H	K	21
593	K	V	T	D	L	L	L	F	F	21
619	S	G	R	I	P	G	L	G	K	21
621	R	I	P	G	L	G	K	D	F	21
44	F	I	L	G	Y	I	V	V	G	20
56	W	L	Y	G	D	P	R	Q	V	20
243	S	L	L	F	I	L	L	L	R	20
259	L	V	L	I	L	G	V	L	G	20
347	A	V	G	Q	M	M	S	T	M	20
363	V	L	L	L	I	C	I	A	Y	20
463	C	V	L	A	G	A	F	A	S	20
501	S	L	A	F	G	A	L	I	L	20
606	V	V	G	G	V	G	V	L	S	20
689	K	S	L	L	K	I	L	G	K	20
16	P	V	K	Y	D	P	S	F	R	19
170	F	L	L	P	S	A	P	A	L	19
186	N	V	T	P	P	A	L	P	G	19
207	G	L	I	D	S	L	N	A	R	19
246	F	I	L	L	L	R	L	V	A	19
249	L	L	R	L	V	A	G	P	L	19
260	V	L	I	L	G	V	L	G	V	19

216

TABLE XXV-continued

262	I	L	G	V	L	G	V	L	A	19
298	N	L	S	A	Y	Q	S	V	Q	19
317	A	V	L	E	A	I	L	L	L	19
333	R	I	R	I	A	I	A	L	L	19
433	G	L	I	Q	R	S	V	F	N	19
508	I	L	T	L	V	Q	I	A	R	19
525	K	L	R	G	V	Q	N	P	V	19
560	Y	I	M	I	A	I	Y	G	K	19
588	V	V	V	L	D	K	V	T	D	19
604	L	L	V	V	G	G	V	G	V	19
605	L	V	V	G	G	V	G	V	L	19
681	L	D	R	P	Y	Y	M	S	K	19
11	E	A	Y	G	K	P	V	K	Y	18
49	I	V	V	G	I	V	A	W	L	18
73	A	Y	C	G	M	G	E	N	K	18
220	K	I	F	E	D	F	A	Q	S	18
248	L	L	L	R	L	V	A	G	P	18
261	L	I	L	G	V	L	G	V	L	18
264	G	V	L	G	V	L	A	Y	G	18
272	G	I	Y	Y	C	W	E	E	Y	18
278	E	E	Y	R	V	L	R	D	K	18
314	I	V	L	A	V	L	E	A	I	18
432	K	G	L	I	Q	R	S	V	F	18
441	N	L	Q	I	Y	G	V	L	G	18
446	G	V	L	G	L	F	W	T	L	18
457	V	L	A	L	G	Q	C	V	L	18
564	A	I	Y	G	K	N	F	C	V	18
587	R	V	V	V	L	D	K	V	T	18
649	Y	V	I	A	S	G	F	F	S	18
10	D	E	A	Y	G	K	P	V	K	17
63	Q	V	L	Y	P	R	N	S	T	17
121	P	E	D	P	W	T	V	G	K	17
177	A	L	G	R	C	F	P	W	T	17
211	S	L	N	A	R	D	I	S	V	17
233	L	V	A	L	G	V	A	L	V	17
235	A	L	G	V	A	L	V	L	S	17
239	A	L	V	L	S	L	L	F	I	17
252	L	V	A	G	P	L	V	L	V	17
309	W	L	A	A	L	I	V	L	A	17
335	R	I	A	I	A	L	L	K	E	17
365	L	L	I	C	I	A	Y	W	A	17
368	C	I	A	Y	W	A	M	T	A	17
401	K	V	P	I	N	T	S	C	N	17
421	G	L	M	C	V	F	Q	G	Y	17
456	W	V	L	A	L	G	Q	C	V	17
459	A	L	G	Q	C	V	L	A	G	17
510	T	L	V	Q	I	A	R	V	I	17
542	K	C	C	L	W	C	L	E	K	17
562	M	I	A	I	Y	G	K	N	F	17
580	L	L	M	R	N	I	V	V	R	17
583	R	N	I	V	R	V	V	V	L	17
644	S	I	L	G	A	Y	V	I	A	17
657	S	V	F	G	M	C	V	D	T	17
662	C	V	D	T	L	F	L	C	F	17
26	P	I	K	N	R	S	C	T	D	16
34	D	V	I	C	C	V	L	F	L	16
45	I	L	G	Y	I	V	V	G	I	16
86	L	L	Y	F	N	I	F	S	C	16
157	T	V	I	T	S	L	Q	Q	E	16
165	E	L	C	P	S	F	L	L	P	16
237	G	V	A	L	V	L	S	L	L	16
258	V	L	V	L	I	L	G	V	L	16
289	S	I	S	Q	L	G	F	T	T	16
304	S	V	Q	E	T	W	L	A	A	16
323	L	L	L	M	L	I	F	L	R	16
364	L	L	L	I	C	I	A	Y	W	16
470	A	S	F	Y	W	A	F	H	K	16
494	T	L	R	Y	H	T	G	S	L	16
511	L	V	Q	I	A	R	V	I	L	16
554	F	L	N	R	N	A	Y	I	M	16
571	C	V	S	A	K	N	A	F	M	16
584	N	I	V	R	V	V	V	L	D	16
673	D	L	E	R	N	N	G	S	L	16
693	K	I	L	G	K	K	N	E	A	16
698	K	N	E	A	P	P	D	N	K	16
20	D	P	S	F	R	G	P	I	K	15
48	Y	I	V	V	G	I	V	A	W	15
58	Y	G	D	P	R	Q	V	L	Y	15
99	N	I	I	S	V	A	E	N	G	15
151	G	V	P	W	N	M	T	V	I	15

217

TABLE XXV-continued

191	A	L	P	G	I	T	N	D	T	15
231	W	I	L	V	A	L	G	V	A	15
234	V	A	L	G	V	A	L	V	L	15
257	L	V	L	V	L	I	L	G	V	15
318	V	L	E	A	I	L	L	L	M	15
322	I	L	L	L	M	L	I	F	L	15
327	L	I	F	L	R	Q	R	I	R	15
329	F	L	R	Q	R	I	R	I	A	15
532	P	V	A	R	C	I	M	C	C	15
589	V	V	L	D	K	V	T	D	L	15
597	L	L	L	F	F	G	K	L	L	15
598	L	L	F	F	G	K	L	L	V	15
622	I	P	G	L	G	K	D	F	K	15
645	I	L	G	A	Y	V	I	A	S	15
651	I	A	S	G	F	F	S	V	F	15
680	S	L	D	R	P	Y	Y	M	S	15
691	L	L	K	I	L	G	K	K	N	15
7	D	E	D	I	D	E	A	Y	G	14
42	L	L	F	I	L	G	Y	I	V	14
53	I	V	A	W	L	Y	G	D	P	14
81	K	D	K	P	Y	L	L	Y	F	14
95	I	L	S	S	N	I	I	S	V	14
148	C	L	P	G	V	P	W	N	M	14
171	L	L	P	S	A	P	A	L	G	14
244	L	L	F	I	L	L	L	R	L	14
311	A	A	L	I	V	L	A	V	L	14
315	V	L	A	V	L	E	A	I	L	14
324	M	L	M	L	I	F	L	R	Q	14
326	M	L	I	F	L	R	Q	R	I	14
337	A	I	A	L	L	K	E	A	S	14
359	L	V	T	F	V	L	L	L	I	14
370	A	Y	W	A	M	T	A	L	Y	14
378	Y	L	A	T	S	G	Q	P	Q	14
388	V	L	W	A	S	N	I	S	S	14
453	T	L	N	W	V	L	A	L	G	14
465	L	A	G	A	F	A	S	F	Y	14
487	L	I	S	A	F	I	R	T	L	14
496	R	Y	H	T	G	S	L	A	F	14
523	D	H	K	L	R	G	V	Q	N	14
527	R	G	V	Q	N	P	V	A	R	14
528	G	V	Q	N	P	V	A	R	C	14
534	A	R	C	I	M	C	C	F	K	14
558	N	A	Y	I	M	I	A	I	Y	14
567	G	K	N	F	C	V	S	A	K	14
596	D	L	L	L	F	F	G	K	L	14
609	G	V	G	V	L	S	F	F	F	14
638	W	L	P	I	M	T	S	I	L	14
647	G	A	Y	V	I	A	S	G	F	14
665	T	L	F	L	C	F	L	E	D	14
685	Y	Y	M	S	K	S	L	L	K	14
694	I	L	G	K	K	N	E	A	P	14
699	N	E	A	P	P	D	N	K	K	14
701	A	P	P	D	N	K	K	R	K	14

V3-HLA-A3-9mers-
24P4C12
Each peptide is a portion of
SEQ ID NO: 7; each start
position is specified, the length
of peptide is 9 amino acids, and
the end position for each
peptide is the start position plus
eight.

Pos	1	2	3	4	5	6	7	8	9	score
8	N	I	T	P	P	A	L	P	G	17

V5-HLA-A3-9mers-
24P4C12
Each peptide is a portion of SEQ
ID NO: 11; each start position is
specified, the length of peptide is
9 amino acids, and the end
position for each peptide is the
start position plus eight.

Pos	1	2	3	4	5	6	7	8	9	score
4	A	I	L	L	L	V	L	I	F	21
8	L	V	L	I	F	L	R	Q	R	20

218

TABLE XXV-continued

5	I	L	<u>L</u>	L	V	<u>L</u>	<u>I</u>	F	L	16
6	L	L	<u>L</u>	V	L	<u>I</u>	<u>F</u>	L	R	16
1	V	L	<u>E</u>	A	I	<u>L</u>	<u>L</u>	L	V	15
7	L	L	<u>V</u>	L	I	<u>F</u>	<u>L</u>	R	Q	14
9	V	L	<u>I</u>	F	L	<u>R</u>	<u>Q</u>	R	I	14

V6-HLA-A3-9mers- 24P4C12										
Each peptide is a portion of SEQ ID NO: 13; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.										
Pos	1	2	3	4	5	6	7	8	9	score
6	G	L	<u>I</u>	P	R	<u>S</u>	<u>V</u>	F	N	22
5	K	G	<u>L</u>	I	P	<u>R</u>	<u>S</u>	V	F	18
7	L	I	<u>P</u>	R	S	<u>V</u>	<u>F</u>	N	L	11

V7-HLA-A3-9mers- 24P4C12										
Each peptide is a portion of SEQ ID NO: 15; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.										
Pos	1	2	3	4	5	6	7	8	9	score
8	A	V	<u>G</u>	Q	M	<u>M</u>	<u>S</u>	T	M	20
5	I	L	<u>V</u>	A	V	<u>G</u>	<u>Q</u>	M	M	19
6	L	V	<u>A</u>	V	G	<u>Q</u>	<u>M</u>	M	S	15
4	W	I	<u>L</u>	V	A	<u>V</u>	<u>G</u>	Q	M	14
3	Y	W	<u>I</u>	L	V	<u>A</u>	<u>V</u>	G	Q	12
1	S	W	<u>Y</u>	W	I	<u>L</u>	<u>V</u>	A	V	10

V8-HLA-A3-9mers- 24P4C12										
Each peptide is a portion of SEQ ID NO: 17; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.										
Pos	1	2	3	4	5	6	7	8	9	score
11	P	I	<u>T</u>	P	T	<u>G</u>	<u>H</u>	V	F	22
6	P	I	<u>M</u>	R	N	<u>P</u>	<u>I</u>	T	P	16
4	W	L	<u>P</u>	I	M	<u>R</u>	<u>N</u>	P	I	12
9	R	N	<u>P</u>	I	T	<u>P</u>	<u>T</u>	G	H	11
1	N	Y	<u>Y</u>	W	L	<u>P</u>	<u>I</u>	M	R	10
17	H	V	<u>F</u>	Q	T	<u>S</u>	<u>I</u>	L	G	10

V9-HLA-A3-9mers- 24P4C12										
Each peptide is a portion of SEQ ID NO: 19; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.										
Pos	1	2	3	4	5	6	7	8	9	score
6	A	L	<u>Y</u>	P	L	<u>P</u>	<u>T</u>	Q	P	25
9	P	L	<u>P</u>	T	Q	<u>P</u>	<u>A</u>	T	L	18
11	P	T	<u>Q</u>	P	A	<u>T</u>	<u>L</u>	G	Y	12
16	T	L	<u>G</u>	Y	V	<u>L</u>	<u>W</u>	A	S	12

Figure 2G-2

40 L F L L F I L G Y I V V G I V A W L Y G
121 TCCTCTTCCTGCTCTTCATTCTAGGTTACATCGTGGTGGGGATTGTGGCCTGGTTGTATG
60 D P R Q V L Y F R N S T G A Y C G M G E
181 GAGACCCCCGGCAAGTCCTCTACCCCAGGAAGTCTACTGGGGCCTACTGTGGCATGGGGG
80 N K D K P Y L L Y F N I F S C I L S S N
241 AGAACAAAGATAAGCCGTATCTCCGTACTTCAACATCTTCAGCTGCATCCTGTCCAGCA
100 I I S V A E N G L Q C P T P Q V C V S S
301 ACATCATCTCAGTTGCTGAGAACGGCCTACAGTGCCCCACACCCAGGTGTGTGTCTCT
120 C P E D P W T V G K N E F S Q T V G E V
361 CCTGCCCCGAGGACCCATGGACTGTGGGAAAAACGAGTTCTCACAGACTGTTGGGGAAAG
140 F Y T K N R N F C L P G V P W N M T V I
421 TCTTCTATACAAAAACAGGAACCTTTGTCTGCCAGGGGTACCCCTGGAATATGACGGTGA
160 T S L Q Q E L C P S F L L P S A P A L G
481 TCACAAGCCTGCAACAGGAAGTCTGCCCCAGTTTCTCCTCCCTCTGCTCCAGCTCTGG
180 R C F P W T N V T P P A L P G I T N D T
541 GGCGCTGCTTTCCATGGACCAACGTTACTCCACCGGCGCTCCCAGGGATCACCAATGACA
200 T I Q Q G I S G L I D S L N A R D I S V
601 CCACCATACAGCAGGGGATCAGCGGTCTTATTGACAGCCTCAATGCCCGAGACATCAGTG
220 K I F E D F A Q S W Y W I L V A V G Q M
661 TTAAGATCTTTGAAGATTTTGCCAGTCCTGGTATTGGATTCTGTGGCTGTGGGACAGA
240 M S I M F Y P L V T F V L L L I C I A Y
721 TGATGTCTACCATGTTCTACCCACTGGTCACCTTTGTCTCCTCCTCATCTGCATTGCCT
260 W A M T A L Y L A T S G Q P Q Y V L W A
781 ACTGGGCCATGACTGCTCTGTACCTGGCTACATCGGGGCAACCCAGTATGTCTCTGGG
280 S N I S S P G C E K V P I N T S C N P T
841 CATCCAACATCAGCTCCCCCGGCTGTGAGAAAGTGCCAATAAATACATCATGCAACCCCA
300 A H L V N S S C P G L M C V F Q G Y S S
901 CGGCCCACCTTGTGAAGTCTCTGTCGCCAGGGCTGATGTGCGTCTTCCAGGGCTACTCAT
320 K G L I Q R S V F N L Q I Y G V L G L F
961 CCAAAGGCCTAATCCAACGTTCTGTCTTCAATCTGCAAATCTATGGGGTCTGGGGCTCT
340 W T L N W V L A L G Q C V L A G A F A S
1021 TCTGGACCCCTAACTGGGTACTGGCCCTGGGCCAATGCGTCCTCGCTGGAGCCTTTGCCT
360 F Y W A F H K P Q D I P T F P L I S A F
1081 CCTTCTACTGGGCCTTCCACAAGCCCCAGGACATCCCTACCTTCCCTTAATCTCTGCCT
380 I R T L R Y H T G S L A F G A L I L T L
1141 TCATCCGCACACTCCGTTACACACTGGGTGATTGGCATTGGAGCCCTCATCTGACCC
400 V Q I A R V I L E Y I D H K L R G V Q N
1201 TTGTGCAGATAGCCCGGGTCATCTTGAGTATATTGACCACAAGCTCAGAGGAGTGCAGA
420 P V A R C I M C C F R C C L W C L E K F
1261 ACCCTGTAGCCCGCTGCATCATGTGCTGTTTCAAGTGCTGCCTCTGGTGTCTGGAAAAAT

219

TABLE XXVI

V1-HLA-A26-9mers-24P4C12										
Each peptide is a portion of SEQ ID NO: 3; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.										
Pos	1	2	3	4	5	6	7	8	9	score
34	D	V	I	C	C	V	L	F	L	35
49	I	V	V	G	I	V	A	W	L	28
483	P	T	F	P	L	I	S	A	F	28
605	L	V	V	G	G	V	G	V	L	27
593	K	V	T	D	L	L	L	F	F	26
317	A	V	L	E	A	I	L	L	L	25
592	D	K	V	T	D	L	L	L	F	25
138	E	V	F	Y	T	K	N	R	N	24
240	L	V	L	S	L	L	F	I	L	24
589	V	V	L	D	K	V	T	D	L	24
38	C	V	L	F	L	L	F	I	L	23
237	G	V	A	L	V	L	S	L	L	23
11	E	A	Y	G	K	P	V	K	Y	22
267	G	V	L	A	Y	G	I	Y	Y	22
285	D	K	G	A	S	I	S	Q	L	22
452	W	T	L	N	W	V	L	A	L	22
50	V	V	G	I	V	A	W	L	Y	20
79	E	N	K	D	K	P	Y	L	L	20
157	T	V	I	T	S	L	Q	Q	E	20
263	L	G	V	L	G	V	L	A	Y	20
446	G	V	L	G	L	F	W	T	L	20
628	D	F	K	S	P	H	L	N	Y	20
641	I	M	T	S	I	L	G	A	Y	20
662	C	V	D	T	L	F	L	C	F	20
236	L	G	V	A	L	V	L	S	L	19
258	V	L	V	L	I	L	G	V	L	19
307	E	T	W	L	A	A	L	I	V	19
320	E	A	I	L	L	L	M	L	I	19
414	L	V	N	S	S	C	P	G	L	19
437	R	S	V	F	N	L	Q	I	Y	19
513	Q	I	A	R	V	I	L	E	Y	19
609	G	V	G	V	L	S	F	F	F	19
673	D	L	E	R	N	N	G	S	L	19
32	C	T	D	V	I	C	C	V	L	18
198	D	T	T	I	Q	Q	G	I	S	18
200	T	I	Q	Q	G	I	S	G	L	18
204	G	I	S	G	L	I	D	S	L	18
244	L	L	F	I	L	L	L	R	L	18
294	G	F	T	T	N	L	S	A	Y	18
354	T	M	F	Y	P	L	V	T	F	18
360	V	T	F	V	L	L	L	I	C	18
400	E	K	V	P	I	N	T	S	C	18
511	L	V	Q	I	A	R	V	I	L	18
596	D	L	L	L	F	F	G	K	L	18
102	S	V	A	E	N	G	L	Q	C	17
184	W	T	N	V	T	P	P	A	L	17
216	D	I	S	V	K	I	F	E	D	17
261	L	I	L	G	V	L	G	V	L	17
358	P	L	V	T	F	V	L	L	L	17
438	S	V	F	N	L	Q	I	Y	G	17
442	L	Q	I	Y	G	V	L	G	L	17
443	Q	I	Y	G	V	L	G	L	F	17
487	L	I	S	A	G	I	R	T	L	17
608	G	G	V	G	V	L	S	F	F	17
664	D	T	L	F	L	C	F	L	E	17

220

TABLE XXVI-continued

V3-HLA-A26-9mers-24P4C12										
Each peptide is a portion of SEQ ID NO: 7; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.										
Pos	1	2	3	4	5	6	7	8	9	score
6	W	T	N	I	T	P	P	A	L	17
9	I	T	P	P	A	L	P	G	I	13
V5-HLA-A26-9mers-24P4C12										
Each peptide is a portion of SEQ ID NO: 11; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.										
Pos	1	2	3	4	5	6	7	8	9	score
3	E	A	I	L	L	L	V	L	I	19
4	A	I	L	L	L	V	L	I	F	18
8	L	V	L	I	F	L	R	Q	R	15
25	2	L	E	A	I	L	L	V	L	14
5	I	L	L	L	V	L	I	F	L	13
V6-HLA-A26-9mers-24P4C12										
Each peptide is a portion of SEQ ID NO: 13; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.										
Pos	1	2	3	4	5	6	7	8	9	score
7	L	I	P	R	S	V	F	N	L	16
5	K	G	L	I	P	R	S	V	F	9
V7-HLA-A26-9mers-24P4C12										
Each peptide is a portion of SEQ ID NO: 15; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.										
Pos	1	2	3	4	5	6	7	8	9	score
8	A	V	G	Q	M	M	S	T	M	12
6	L	V	A	V	G	Q	M	M	S	11
4	W	I	L	V	A	V	G	Q	M	10
1	S	W	Y	W	I	L	V	A	V	8
5	I	L	V	A	V	G	Q	M	M	6
2	W	Y	W	I	L	V	A	V	G	5
7	V	A	V	G	Q	M	M	S	T	5
V8-HLA-A26-9mers-24P4C12										
Each peptide is a portion of SEQ ID NO: 17; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.										
Pos	1	2	3	4	5	6	7	8	9	score
19	F	Q	T	S	I	L	G	A	Y	20
11	P	I	T	P	T	G	H	V	F	15
17	H	V	F	Q	T	S	I	L	G	15
16	G	H	V	F	Q	T	S	I	L	13

221

TABLE XXVI-continued

20	Q	T	S	I	L	G	A	Y	V	10
14	P	T	G	H	V	F	Q	T	S	9
V9-HLA-A26-9mers- 24P4C12										
Each peptide is a portion of SEQ ID NO: 19; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.										
Pos	1	2	3	4	5	6	7	8	9	score
11	P	T	Q	P	A	T	L	G	Y	20
15	A	T	L	G	Y	V	L	W	A	13
2	W	A	M	T	A	L	Y	P	L	12
13	Q	P	A	T	L	G	Y	V	L	10
4	M	T	A	L	Y	P	L	P	T	9
9	P	L	P	T	Q	P	A	T	L	9

TABLE XXVII

V1-HLA-B0702- 9mers-24P4C12										
Each peptide is a portion of SEQ ID NO: 3; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.										
Pos	1	2	3	4	5	6	7	8	9	score
255	G	P	L	V	L	V	L	I	L	23
357	Y	P	L	V	T	F	V	L	L	23
683	R	P	Y	Y	M	S	K	S	L	21
149	L	P	G	V	P	W	N	M	T	20
396	S	P	G	C	E	K	V	P	I	20
482	I	P	T	F	P	L	I	S	A	20
631	S	P	H	L	N	Y	Y	W	L	20
15	K	P	V	K	Y	D	P	S	F	19
152	V	P	W	N	M	T	V	I	T	19
167	C	P	S	F	L	L	P	S	A	19
25	G	P	I	K	N	R	S	C	T	18
172	L	P	S	A	P	A	L	G	R	18
83	K	P	Y	L	L	Y	F	N	I	17
188	T	P	P	A	L	P	G	I	T	17
192	L	P	G	I	T	N	D	T	T	17
57	L	Y	G	D	P	R	Q	V	L	16
232	I	L	V	A	L	G	V	A	L	16
253	V	A	G	P	L	V	L	V	L	16
479	P	Q	D	I	P	T	F	P	L	16
503	A	F	G	A	L	I	L	T	L	16
49	I	V	V	G	I	V	A	W	L	15
120	C	P	E	D	P	W	T	V	G	15
175	A	P	A	L	G	R	C	F	P	15
189	P	P	A	L	P	G	I	T	N	15
234	V	A	L	G	V	A	L	V	L	15
251	R	L	V	A	G	P	L	V	L	15
381	T	S	G	Q	P	Q	Y	V	L	15
406	T	S	C	N	P	T	A	H	L	15
583	R	N	I	V	R	V	V	V	L	15
617	F	F	S	G	R	I	P	G	L	15
20	D	P	S	F	R	G	P	I	K	14
34	D	V	I	C	C	V	L	F	L	14
66	Y	P	R	N	S	T	G	A	Y	14
204	G	I	S	G	L	I	D	S	L	14
236	L	G	V	A	L	V	L	S	L	14
252	L	V	A	G	P	L	V	L	V	14
291	S	Q	L	G	F	T	T	N	L	14
311	A	A	L	I	V	L	A	V	L	14
317	A	V	L	E	A	I	L	L	L	14
333	R	I	R	I	A	I	A	L	L	14
351	M	M	S	T	M	F	Y	P	L	14
419	C	P	G	L	M	C	V	F	Q	14
452	W	T	L	N	W	V	L	A	L	14

222

TABLE XXVII-continued

499	T	G	S	L	A	F	G	A	L	14
605	L	V	V	G	G	V	G	V	L	14
660	G	M	C	V	D	T	L	F	L	14
60	D	P	R	Q	V	L	Y	P	R	13
100	I	I	S	V	A	E	N	G	L	13
110	C	P	T	P	Q	V	C	V	S	13
164	Q	E	L	C	P	S	F	L	L	13
170	F	L	L	P	S	A	P	A	L	13
182	F	P	W	T	N	V	T	P	P	13
228	S	W	Y	W	I	L	V	A	L	13
241	V	L	S	L	L	F	I	L	L	13
249	L	L	R	L	V	A	G	P	L	13
261	L	I	L	G	V	L	G	V	L	13
302	Y	Q	S	V	Q	E	T	W	L	13
319	L	E	A	I	L	L	L	M	L	13
358	P	L	V	T	F	V	L	L	L	13
369	I	A	Y	W	A	M	T	A	L	13
371	Y	W	A	M	T	A	L	Y	L	13
409	N	P	T	A	H	L	V	N	S	13
442	L	Q	I	Y	G	V	L	G	L	13
446	G	V	L	G	L	F	W	T	L	13
478	K	P	Q	D	I	P	T	F	P	13
487	L	I	S	A	F	I	R	T	L	13
494	T	L	R	Y	H	T	G	S	L	13
501	S	L	A	F	G	A	L	I	L	13
511	L	V	Q	I	A	R	V	I	L	13
590	V	L	D	K	V	T	D	L	L	13
622	I	P	G	L	G	K	D	F	K	13
651	I	A	S	G	F	F	S	V	F	13
32	C	T	D	V	I	C	C	V	L	12
78	G	E	N	K	D	K	P	Y	L	12
154	W	N	M	T	V	I	T	S	L	12
184	W	T	N	V	T	P	P	A	L	12
242	L	S	L	L	F	I	L	L	L	12
244	L	L	F	I	L	L	L	R	L	12
285	D	K	G	A	S	I	S	Q	L	12
305	V	Q	E	T	W	L	A	A	L	12
308	T	W	L	A	A	L	I	V	L	12
315	V	L	A	V	L	E	A	I	L	12
322	I	L	L	L	M	L	I	F	L	12
356	F	Y	P	L	V	T	F	V	L	12
373	A	M	T	A	L	Y	L	A	T	12
380	A	T	S	G	Q	P	Q	Y	V	12
457	V	L	A	L	G	Q	C	V	L	12
525	K	L	R	G	V	Q	N	P	V	12
547	C	L	E	K	F	I	K	F	L	12
572	V	S	A	K	N	A	F	M	L	12
589	V	V	L	D	K	V	T	D	L	12
591	L	D	K	V	T	D	L	L	L	12
626	G	K	D	F	K	S	P	H	L	12
658	V	F	G	M	C	V	D	T	L	12
701	A	P	P	D	N	K	K	R	K	12
28	K	N	R	S	C	T	D	V	I	11
45	I	L	G	Y	I	V	V	G	I	11
79	E	N	K	D	K	P	Y	L	L	11
104	A	E	N	G	L	Q	C	P	T	11
107	G	L	Q	C	P	T	P	Q	V	11
109	Q	C	P	T	P	Q	V	C	V	11
112	T	P	Q	V	C	V	S	S	C	11
123	D	P	W	T	V	G	K	N	E	11
163	Q	Q	E	L	C	P	S	F	L	11
169	S	F	L	L	P	S	A	P	A	11
177	A	L	G	R	C	F	P	W	T	11
191	A	L	P	G	I	T	N	D	L	11
237	G	V	A	L	V	L	S	L	L	11
239	A	L	V	L	S	L	L	F	I	11
258	V	L	V	L	I	L	G	V	L	11
262	I	L	G	V	L	G	V	L	A	11
275	Y	C	W	E	E	Y	R	V	L	11
310	L	A	A	L	I	V	L	A	V	11
332	Q	R	I	R	I	A	I	A	L	11
343	E	A	S	K	A	V	G	Q	M	11
354	T	M	F	Y	P	L	V	T	F	11
384	Q	P	Q	Y	V	L	W	A	S	11
414	L	V	N	S	S	C	P	G	L	11
426	F	Q	G	Y	S	S	C	K	G	11
434	L	I	Q	R	S	V	F	N	L	11
440	F	N	L	Q	I	Y	G	V	L	11
450	L	F	W	T	L	N	W	V	L	11
464	V	L	A	G	A	F	A	S	F	11

223

TABLE XXVII-continued

518	I	L	E	Y	I	D	H	K	L	11
531	N	P	V	A	R	C	I	M	C	11
537	I	M	C	C	F	K	C	C	L	11
571	C	V	S	A	K	N	A	F	M	11
573	S	A	K	N	A	F	M	L	L	11
574	A	K	N	A	F	M	L	L	M	11
596	D	L	L	L	F	F	G	K	L	11
597	L	L	L	F	F	G	K	L	L	11
599	L	F	F	G	K	L	L	V	V	11
638	W	L	P	I	M	T	S	I	L	11
663	V	D	T	L	F	L	C	F	L	11
686	Y	M	S	K	S	L	L	K	I	11
702	P	P	D	N	K	K	R	K	K	11

V3-HLA-B0702-
9mers-24P4C12

Each peptide is a portion of SEQ ID
NO: 7; each start position is
specified, the length of peptide is 9
amino acids, and the end position
for each peptide is the start position
plus eight.

Pos	1	2	3	4	5	6	7	8	9	score
4	F	P	W	T	N	I	T	P	P	12
6	W	T	N	I	T	P	P	A	L	12
1	G	R	C	F	P	W	T	N	I	10
2	R	C	F	P	W	T	N	I	T	9
5	P	W	T	N	I	T	P	P	A	9
9	I	T	P	P	A	L	P	G	I	9
8	N	I	T	P	P	A	L	P	G	7

V5-HLA-B0702-
9mers-24P4C12

Each peptide is a portion of SEQ ID
NO: 11; each start position is
specified, the length of peptide is 9
amino acids, and the end position
for each peptide is the start position
plus eight.

Pos	1	2	3	4	5	6	7	8	9	score
2	L	E	A	I	L	L	L	V	L	14
5	I	L	L	L	V	L	I	F	L	12
4	A	I	L	L	L	V	L	I	F	11
1	V	L	E	A	I	L	L	L	V	9
3	E	A	I	L	L	L	V	L	I	9
9	V	L	I	F	L	R	Q	R	I	7

V6-HLA-B0702-
9mers-24P4C12

Each peptide is a portion of SEQ ID
NO: 13; each start position is
specified, the length of peptide is 9
amino acids, and the end position
for each peptide is the start position
plus eight.

Pos	1	2	3	4	5	6	7	8	9	score
8	I	P	R	S	V	F	N	L	Q	14
5	K	G	L	I	P	R	S	V	F	12
7	L	I	P	R	S	V	F	N	L	11
9	P	R	S	V	F	N	L	Q	I	10
4	S	K	G	L	I	P	R	S	V	7

224

TABLE XXVII-continued

V7-HLA-B0702-
9mers-24P4C12

Each peptide is a portion of SEQ ID
NO: 15; each start position is
specified, the length of peptide is 9
amino acids, and the end position
for each peptide is the start position
plus eight.

Pos	1	2	3	4	5	6	7	8	9	score
1	S	W	Y	W	I	L	V	A	V	9
5	I	L	V	A	V	G	Q	M	M	9
8	A	V	G	Q	M	M	S	T	M	9
7	V	A	V	G	Q	M	M	S	T	8
4	W	I	L	V	A	V	G	Q	M	7

V8-HLA-B0702-9mers-
24P4C12

Each peptide is a portion of SEQ ID
NO: 17; each start position is
specified, the length of peptide is 9
amino acids, and the end position
for each peptide is the start position
plus eight.

Pos	1	2	3	4	5	6	7	8	9	score
19	F	Q	T	S	I	L	G	A	Y	20
11	P	I	T	P	T	G	H	V	F	15
17	H	V	F	Q	T	S	I	L	G	15
16	G	H	V	F	Q	T	S	I	L	13
20	Q	T	S	I	L	G	A	Y	V	10
14	P	T	G	H	V	F	Q	T	S	9

V9-HLA-B0702-9mers-
24P4C12

Each peptide is a portion of SEQ ID
NO: 19; each start position is
specified, the length of peptide is 9
amino acids, and the end position
for each peptide is the start position
plus eight.

Pos	1	2	3	4	5	6	7	8	9	score
13	Q	P	A	T	L	G	Y	V	L	23
8	Y	P	L	P	T	Q	P	A	T	19
10	L	P	T	Q	P	A	T	L	G	14
15	A	T	L	G	Y	V	L	W	A	13
2	W	A	M	T	A	L	Y	P	L	12
7	L	Y	P	L	P	T	Q	P	A	11
9	P	L	P	T	Q	P	A	T	L	11

TABLE XXVIII

V1-HLA-B08-9mers

Each peptide is a portion of SEQ
ID NO: 3; each start position is
specified, the length of peptide is 9
amino acids, and the end position
for each peptide is the start
position plus eight.

Pos	1	2	3	4	5	6	7	8	9	score
79	E	N	K	D	K	P	Y	L	L	32
141	Y	T	K	N	R	N	F	C	L	29
282	V	L	R	D	K	G	A	S	I	29
573	S	A	K	N	A	F	M	L	L	26
249	L	L	R	L	V	A	G	P	L	23
494	T	L	R	Y	H	T	G	S	L	23
26	P	I	K	N	R	S	C	T	D	22
329	F	L	R	Q	R	I	R	I	A	22
589	V	V	L	D	K	V	T	D	L	22
333	R	I	R	I	A	I	A	L	L	21
583	R	N	I	V	R	V	V	V	L	21

225

TABLE XXVIII-continued

591	L	D	K	V	T	D	L	L	L	21
626	G	K	D	F	K	S	P	H	L	21
687	M	S	K	S	L	L	K	I	L	21
340	L	L	K	E	A	S	K	A	V	20
474	W	A	F	H	K	P	Q	D	I	20
523	D	H	K	L	R	G	V	Q	N	20
540	C	F	K	C	C	L	W	C	L	20
617	F	F	S	G	R	I	P	G	L	20
2	G	G	K	Q	R	D	E	D	D	19
232	I	L	V	A	L	G	V	A	L	19
255	G	P	L	V	L	V	L	I	L	19
631	S	P	H	L	N	Y	Y	W	L	19
694	I	L	G	K	K	N	E	A	P	19
139	V	F	Y	T	K	N	R	N	F	18
170	F	L	L	P	S	A	P	A	L	18
241	V	L	S	L	L	F	I	L	L	18
247	I	L	L	L	R	L	V	A	G	18
258	V	L	V	L	I	L	G	V	L	18
315	V	L	A	V	L	E	A	I	L	18
322	I	L	L	L	M	L	I	F	L	18
357	Y	P	L	V	T	F	V	L	L	18
457	V	L	A	L	G	Q	C	V	L	18
501	S	L	A	F	G	A	L	I	L	18
514	I	A	R	V	I	L	E	Y	I	18
518	I	L	E	Y	I	D	H	K	L	18
546	W	C	L	E	K	F	I	K	F	18
547	C	L	E	K	F	I	K	F	L	18
683	R	P	Y	M	S	K	S	L		18
11	E	A	Y	G	K	P	V	K	Y	17
213	N	A	R	D	I	S	V	K	I	17
216	D	I	S	V	K	I	F	E	D	17
358	P	L	V	T	F	V	L	L	L	17
533	V	A	R	C	I	M	C	C	F	17
590	V	L	D	K	V	T	D	L	L	17
596	D	L	L	L	F	F	G	K	L	17
597	L	L	L	F	F	G	K	L	L	17
673	D	L	E	R	N	N	G	S	L	17
691	L	L	K	I	L	G	K	K	N	17
45	I	L	G	Y	I	V	V	G	I	16
64	V	L	Y	P	R	N	S	T	G	16
81	K	D	K	P	Y	L	L	Y	F	16
100	I	I	S	V	A	E	N	G	L	16
158	V	I	T	S	L	Q	Q	E	L	16
204	G	I	S	G	L	I	D	S	L	16
211	S	L	N	A	R	D	I	S	V	16
244	L	L	F	I	L	L	L	R	L	16
251	R	L	V	A	G	P	L	V	L	16
253	V	A	G	P	L	V	L	V	L	16
338	I	A	L	L	K	E	A	S	K	16
369	I	A	Y	W	A	M	T	A	L	16
433	G	L	I	Q	R	S	V	F	N	16
551	F	I	K	F	L	N	R	N	A	16
638	W	L	P	I	M	T	S	I	L	16
702	P	P	D	N	K	K	R	K	K	16
35	V	I	C	C	V	L	F	L	L	15
200	T	I	Q	Q	G	I	S	G	L	15
225	F	A	Q	S	W	Y	W	I	L	15
234	V	A	L	G	V	A	L	V	L	15
316	L	A	V	L	E	A	I	L	L	15
331	R	Q	R	I	R	I	A	I	A	15
396	S	P	G	C	E	K	V	P	I	15
434	L	I	Q	R	S	V	F	N	L	15
487	L	I	S	A	F	I	R	T	L	15
553	K	F	L	N	R	N	A	Y	I	15
564	A	I	Y	G	K	N	F	C	V	15
579	M	L	L	M	R	N	I	V	R	15
693	K	I	L	G	K	K	N	E	A	15

226

TABLE XXVIII-continued

V3-HLA-B08-9mers- 24P4C12											
5	Each peptide is a portion of SEQ ID NO: 7; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.										
10	Pos	1	2	3	4	5	6	7	8	9	score
	6	W	T	N	I	T	P	P	A	L	11
	4	F	P	W	T	N	I	T	P	P	8
15	1	G	R	C	F	P	W	T	N	I	7
	9	I	T	P	P	A	L	P	G	I	7
V5-B08-9mers- 24P4C12											
20	Each peptide is a portion of SEQ ID NO: 11; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.										
25	Pos	1	2	3	4	5	6	7	8	9	score
	5	I	L	L	L	V	L	I	F	L	18
	3	E	A	I	L	L	L	V	L	I	14
30	9	V	L	I	F	L	R	Q	R	I	13
	4	A	I	L	L	L	V	L	I	F	12
	2	L	E	A	I	L	L	L	V	L	10
	6	L	L	L	V	L	I	F	L	R	8
V6-HLA-B08-9mers- 24P4C12											
35	Each peptide is a portion of SEQ ID NO: 13; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.										
40	Pos	1	2	3	4	5	6	7	8	9	score
45	6	G	L	I	P	R	S	V	F	N	16
	7	L	I	P	R	S	V	F	N	L	15
	3	S	S	K	G	L	I	P	R	S	13
	8	I	P	R	S	V	F	N	L	Q	13
	1	G	Y	S	S	K	G	L	I	P	11
50	9	P	R	S	V	F	N	L	Q	I	8
V7-HLA-B08-9mers- 24P4C12											
55	Each peptide is a portion of SEQ ID NO: 15; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.										
60	Pos	1	2	3	4	5	6	7	8	9	score
	5	I	L	V	A	V	G	Q	M	M	7
	4	W	I	L	V	A	V	G	Q	M	6
	7	V	A	V	G	Q	M	M	S	T	5
65	1	S	W	Y	W	I	L	V	A	V	4

227

TABLE XXVIII-continued

V8-HLA-B08-9mers- 24P4C12										
Each peptide is a portion of SEQ ID NO: 17; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.										
Pos	1	2	3	4	5	6	7	8	9	score
5	L	P	I	M	R	N	P	I	T	15
4	W	L	P	I	M	R	N	P	I	12
16	G	H	V	F	Q	T	S	I	L	11
11	P	I	T	P	T	G	H	V	F	10
7	I	M	R	N	P	I	T	P	T	8
13	T	P	T	G	H	V	F	Q	T	7
15	T	G	H	V	F	Q	T	S	I	7

V9-HLA-B08- 9mers-24P4C12										
Each peptide is a portion of SEQ ID NO: 19; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.										
Pos	1	2	3	4	5	6	7	8	9	score
9	P	L	P	T	Q	P	A	T	L	16
13	Q	P	A	T	L	G	Y	V	L	16
2	W	A	M	T	A	L	Y	P	L	14
16	T	L	G	Y	V	L	W	A	S	8
18	G	Y	V	L	W	A	S	N	I	8
8	Y	P	L	P	T	Q	P	A	T	7

TABLE XXIX

V1-HLA-B1510-9mers-24P4C12										
Each peptide is a portion of SEQ ID NO: 3; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.										
Pos	1	2	3	4	5	6	7	8	9	score
275	Y	C	W	E	E	Y	R	V	L	16
583	R	N	I	V	R	V	V	V	L	16
57	L	Y	G	D	P	R	Q	V	L	15
232	I	L	V	A	L	G	V	A	L	15
253	V	A	G	P	L	V	L	V	L	15
381	T	S	G	Q	P	Q	Y	V	L	15
487	L	I	S	A	F	I	R	T	L	15
605	L	V	V	G	G	V	G	V	L	15
49	I	V	V	G	I	V	A	W	L	14
78	G	E	N	K	D	K	P	Y	L	14
100	I	I	S	V	A	E	N	G	L	14
170	F	L	L	P	S	A	P	A	L	14
184	W	T	N	V	T	P	P	A	L	14
200	T	I	Q	Q	G	I	S	G	L	14
204	G	I	S	G	L	I	D	S	L	14
251	R	L	V	A	G	P	L	V	L	14
357	Y	P	L	V	T	F	V	L	L	14
369	I	A	Y	W	A	M	T	A	L	14
457	V	L	A	L	G	Q	C	V	L	14
617	F	F	S	G	R	I	P	G	L	14
32	C	T	D	V	I	C	C	V	L	13
79	E	N	K	D	K	P	Y	L	L	13
228	S	W	Y	W	I	L	V	A	L	13
234	V	A	L	G	V	A	L	V	L	13
255	G	P	L	V	L	V	L	I	L	13
261	L	I	L	G	V	L	G	V	L	13
302	Y	Q	S	V	Q	E	T	W	L	13
308	T	W	L	A	A	L	I	V	L	13

228

TABLE XXIX-continued

440	F	N	L	Q	I	Y	G	V	L	13
446	G	V	L	G	L	F	W	T	L	13
499	T	G	S	L	A	F	G	A	L	13
511	L	V	Q	I	A	R	V	I	L	13
518	I	L	E	Y	I	D	H	K	L	13
537	I	M	C	C	F	K	C	C	L	13
547	C	L	E	K	F	I	K	F	L	13
572	V	S	A	K	N	A	F	M	L	13
163	Q	Q	E	L	C	P	S	F	L	12
237	G	V	A	L	V	L	S	L	L	12
244	L	L	F	I	L	L	L	R	L	12
258	V	L	V	L	I	L	G	V	L	12
305	V	Q	E	T	W	L	A	A	L	12
311	A	A	L	I	V	L	A	V	L	12
315	V	L	A	V	L	E	A	I	L	12
317	A	V	L	E	A	I	L	L	L	12
322	I	L	L	L	M	L	I	F	L	12
356	F	Y	P	L	V	T	F	V	L	12
371	Y	W	A	M	T	A	L	Y	L	12
406	T	S	C	N	P	T	A	H	L	12
412	A	H	L	V	N	S	S	C	P	12
442	L	Q	I	Y	G	V	L	G	L	12
450	L	F	W	T	L	N	W	V	L	12
452	W	T	L	N	W	V	L	A	L	12
476	F	H	K	P	Q	D	I	P	T	12
497	Y	H	T	G	S	L	A	F	G	12
501	S	L	A	F	G	A	L	I	L	12
503	A	F	G	A	L	I	L	T	L	12
523	D	H	K	L	R	G	V	Q	N	12
589	V	V	L	D	K	V	T	D	L	12
626	G	K	D	F	K	S	P	H	L	12
651	I	A	S	G	F	F	S	V	F	12
658	V	F	G	M	C	V	D	T	L	12
660	G	M	C	V	D	T	L	F	L	12
673	D	L	E	R	N	N	G	S	L	12
34	D	V	I	C	C	V	L	F	L	11
88	Y	F	N	I	F	S	C	I	L	11
141	Y	T	K	N	R	N	F	C	L	11
154	W	N	M	T	V	I	T	S	L	11
158	V	I	T	S	L	Q	Q	E	L	11
164	Q	E	L	C	P	S	F	L	L	11
236	L	G	V	A	L	V	L	S	L	11
241	V	L	S	L	L	F	I	L	L	11
242	L	S	L	L	F	I	L	L	L	11
285	D	K	G	A	S	I	S	Q	L	11
291	S	Q	L	G	F	T	T	N	L	11
319	L	E	A	I	L	L	L	M	L	11
332	Q	R	I	R	I	A	I	A	L	11
333	R	I	R	I	A	I	A	L	L	11
351	M	M	S	T	M	F	Y	P	L	11
354	T	M	F	Y	P	L	V	T	F	11
358	P	L	V	T	F	V	L	L	L	11
414	L	V	N	S	S	C	P	G	L	11
434	L	I	Q	R	S	V	F	N	L	11
479	P	Q	D	I	P	T	F	P	L	11
494	T	L	R	Y	H	T	G	S	L	11
590	V	L	D	K	V	T	D	L	L	11
591	L	D	K	V	T	D	L	L	L	11
631	S	P	H	L	N	Y	Y	W	L	11
684	P	Y	Y	M	S	K	S	L	L	11
35	V	I	C	C	V	L	F	L	L	10
38	C	V	L	F	L	L	F	I	L	10
124	P	W	T	V	G	K	N	E	F	10
225	F	A	Q	S	W	Y	F	I	L	10
240	L	V	L	S	L	L	W	I	L	10
249	L	L	R	L	V	A	G	P	L	10
316	L	A	V	L	E	A	I	L	L	10
343	E	A	S	K	A	V	G	Q	M	10
418	S	C	P	G	L	M	C	V	F	10
426	F	Q	G	Y	S	S	K	G	L	10
477	H	K	P	Q	D	I	P	T	F	10
483	P	T	F	P	L	I	S	A	F	10
540	C	F	K	C	C	L	W	C	L	10
573	S	A	K	N	A	F	M	L	L	10
596	D	L	L	L	F	F	G	K	L	10
597	L	L	L	F	F	G	K	L	L	10
632	P	H	L	N	Y	Y	W	L	P	10
638	W	L	P	I	M	T	S	I	L	10
65	V	D	T	L	F	L	C	F	L	10
666	L	F	L	C	F	L	E	D	L	10

229

TABLE XXIX-continued

683	R	P	Y	Y	M	S	K	S	L	10
687	M	S	K	S	L	L	K	I	L	10
33	T	D	V	I	C	C	V	L	F	9
36	I	C	C	V	L	F	L	L	F	9
217	I	S	V	K	I	F	E	D	F	9
347	A	V	G	Q	M	M	S	T	M	9
432	K	G	L	I	Q	R	S	V	F	9
461	G	Q	C	V	L	A	G	A	F	9
607	V	G	G	V	G	V	L	S	F	9
679	G	S	L	D	R	P	Y	Y	M	9
15	K	P	V	K	Y	D	P	S	F	8
81	K	D	K	P	Y	L	L	Y	F	8
132	F	S	Q	T	V	G	E	V	F	8
139	V	F	Y	T	K	N	R	N	F	8
148	C	L	P	G	V	P	W	N	M	8
162	L	Q	Q	E	L	C	P	S	F	8
174	S	A	P	A	L	G	R	C	F	8
287	G	A	S	I	S	Q	L	G	F	8
415	V	N	S	S	C	P	G	L	M	8
464	V	L	A	G	A	F	A	S	F	8
468	A	F	A	S	F	Y	W	A	F	8
496	R	Y	H	T	G	S	L	A	F	8
530	Q	N	P	V	A	R	C	I	M	8
570	F	C	V	S	A	K	N	A	F	8
608	G	G	V	G	V	L	S	F	F	8
609	G	V	G	V	L	S	F	F	F	8
647	G	A	Y	V	I	A	S	G	F	8
48	Y	I	V	V	G	I	V	A	W	7
69	N	S	T	G	A	Y	C	G	M	7
214	A	R	D	I	S	V	K	I	F	7
238	V	A	L	V	L	S	L	L	F	7
318	V	L	E	A	I	L	L	L	M	7
321	A	I	L	L	L	M	L	I	F	7
366	L	I	C	I	A	Y	W	A	M	7
443	Q	I	Y	G	V	L	G	L	F	7
533	V	A	R	C	I	M	C	C	F	7
546	W	C	L	E	K	F	I	K	F	7
554	F	L	N	R	N	A	Y	I	M	7
562	M	I	A	I	Y	G	K	N	F	7
571	C	V	S	A	K	N	A	F	M	7
574	A	K	N	A	F	M	L	L	M	7
593	K	V	T	D	L	L	L	F	F	7
621	R	I	P	G	L	G	K	D	F	7
634	L	N	Y	Y	W	L	P	I	M	7
653	S	G	F	F	S	V	F	G	M	7

V3-HLA-B1510-9mers-24P4C12
Each peptide is a portion of SEQ ID NO: 7; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.

Pos	1	2	3	4	5	6	7	8	9	score
6	W	T	N	I	T	P	P	A	L	13

V5-B1510-9mers-24P4C12
Each peptide is a portion of SEQ ID NO: 11; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.

Pos	1	2	3	4	5	6	7	8	9	score
2	L	E	A	I	L	L	L	V	L	13
5	I	L	L	L	V	L	I	F	L	12

V6-B1510-9mers-24P4C12
Each peptide is a portion of SEQ ID NO: 13; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.

Pos	1	2	3	4	5	6	7	8	9	score

230

TABLE XXIX-continued

7	L	I	P	R	S	V	F	N	L	11
5	K	G	L	I	P	R	S	V	F	10
3	S	S	K	G	L	I	P	R	S	5
6	G	L	I	P	R	S	V	F	N	5

V7-B1510-9mers-24P4C12
Each peptide is a portion of SEQ ID NO: 15; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.

Pos	1	2	3	4	5	6	7	8	9	score
8	A	V	G	Q	M	M	S	T	M	9
4	W	I	L	V	A	V	G	Q	M	8
5	I	L	V	A	V	G	Q	M	M	8
1	S	W	Y	W	I	L	V	A	V	3
2	W	Y	W	I	L	V	A	V	G	3
3	Y	W	I	L	V	A	V	G	Q	3
6	L	V	A	V	G	Q	M	M	S	3

V8-B1510-9mers-24P4C12
Each peptide is a portion of SEQ ID NO: 17; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.

Pos	1	2	3	4	5	6	7	8	9	score
16	G	H	V	F	Q	T	S	I	L	21
11	P	I	T	P	T	G	H	V	F	10
13	Q	P	A	T	L	G	Y	V	L	13
9	P	L	P	T	Q	P	A	T	L	12
2	W	A	M	T	A	L	Y	P	L	10

V9-B1510-9mers-24P4C12
Each peptide is a portion of SEQ ID NO: 19; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.

Pos	1	2	3	4	5	6	7	8	9	score
13	Q	P	A	T	L	G	Y	V	L	13
9	P	L	P	T	Q	P	A	T	L	12
2	W	A	M	T	A	L	Y	P	L	10

TABLE XXX

V1-HLA-B2705-9mers-24P4C12
Each peptide is a portion of SEQ ID NO: 3; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.

Pos	1	2	3	4	5	6	7	8	9	score
334	I	R	I	A	I	A	L	L	K	26
332	Q	R	I	R	I	A	I	A	L	25
675	E	R	N	N	G	S	L	D	R	24
214	A	R	D	I	S	V	K	I	F	23
534	A	R	C	I	M	C	C	F	K	21
620	G	R	I	P	G	L	G	K	D	21
5	Q	R	D	E	D	D	E	A	Y	20
204	G	I	S	G	L	I	D	S	L	20
446	G	V	L	G	L	F	W	T	L	20
689	K	S	L	L	K	I	L	G	K	20
251	R	L	V	A	G	P	L	V	L	19
424	C	V	F	Q	G	Y	S	S	K	19
436	Q	R	S	V	F	N	L	Q	I	19
483	P	T	F	P	L	I	S	A	F	19

231

TABLE XXX-continued

583	R	N	I	V	R	V	V	V	L	19
608	G	G	V	G	V	L	S	F	F	19
15	K	P	V	K	Y	D	P	S	F	18
22	S	F	R	G	P	I	K	N	R	18
179	G	R	C	F	P	W	T	N	V	18
200	T	I	Q	Q	G	I	S	G	L	18
207	G	L	I	D	S	L	N	A	R	18
234	V	A	L	G	V	A	L	V	L	18
244	L	L	F	I	L	L	L	R	L	18
255	G	P	L	V	L	V	L	I	L	18
291	S	Q	L	G	F	T	T	N	L	18
317	A	V	L	E	A	I	L	L	L	18
330	L	R	Q	R	I	R	I	A	I	18
333	R	I	R	I	A	I	A	L	L	18
496	R	Y	H	T	G	S	L	A	F	18
527	R	G	V	Q	N	P	V	A	R	18
647	G	A	Y	V	I	A	S	G	F	18
668	L	C	F	L	E	D	L	E	R	18
683	R	P	Y	Y	M	S	K	S	L	18
690	S	L	L	K	I	L	G	K	K	18
49	I	V	V	G	I	V	A	W	L	17
78	G	E	N	K	D	K	P	Y	L	17
154	W	N	M	T	V	I	T	S	L	17
237	G	V	A	L	V	L	S	L	L	17
242	L	S	L	L	F	I	L	L	L	17
261	L	I	L	G	V	L	G	V	L	17
287	G	A	S	I	S	Q	L	G	F	17
311	A	A	L	I	V	L	A	V	L	17
338	I	A	L	L	K	E	A	S	K	17
354	T	M	F	Y	P	L	V	T	F	17
381	T	S	G	Q	P	Q	Y	V	L	17
429	Y	S	S	K	G	L	I	Q	R	17
477	H	K	P	Q	D	I	P	T	F	17
503	A	F	G	A	L	I	L	T	L	17
516	R	V	I	L	E	Y	I	D	H	17
546	W	C	L	E	K	F	I	K	F	17
549	E	K	F	I	K	F	L	N	R	17
605	L	V	V	G	G	V	G	V	L	17
621	R	I	P	G	L	G	K	D	F	17
11	E	A	Y	G	K	P	V	K	Y	16
23	F	R	G	P	I	K	N	R	S	16
137	G	E	V	F	Y	T	K	N	R	16
139	V	F	Y	T	K	N	R	N	F	16
170	F	L	L	P	S	A	P	A	L	16
283	L	R	D	K	G	A	S	I	S	16
285	D	K	G	A	S	I	S	Q	L	16
321	A	I	L	L	L	M	L	I	F	16
322	I	L	L	L	M	L	I	F	L	16
323	L	L	L	M	L	I	F	L	R	16
327	L	I	F	L	R	Q	R	I	R	16
432	K	G	L	I	Q	R	S	V	F	16
440	F	N	L	Q	I	Y	G	V	L	16
442	L	Q	I	Y	G	V	L	G	L	16
443	Q	I	Y	G	V	L	G	L	F	16
457	V	L	A	L	G	Q	C	V	L	16
508	I	L	T	L	V	Q	I	A	R	16
517	V	I	L	E	Y	I	D	H	K	16
589	V	V	L	D	K	V	T	D	L	16
617	F	F	S	G	R	I	P	G	L	16
626	G	K	D	F	K	S	P	H	L	16
699	N	E	A	P	P	D	N	K	K	16
10	D	E	A	Y	G	K	P	V	K	15
40	L	F	L	L	F	I	L	G	Y	15
60	D	P	R	Q	V	L	Y	P	R	15
73	A	Y	C	G	M	G	E	N	K	15
81	K	D	K	P	Y	L	L	Y	F	15
124	P	W	T	V	G	K	N	E	F	15
212	L	N	A	R	D	I	S	V	K	15
217	I	S	V	K	I	F	E	D	F	15
228	S	W	Y	W	I	L	V	A	L	15
236	L	G	V	A	L	V	L	S	L	15
238	V	A	L	V	L	S	L	L	F	15
243	S	L	L	F	I	L	L	L	R	15
253	V	A	G	P	L	V	L	V	L	15
258	V	L	V	L	I	L	G	V	L	15
308	T	W	L	A	A	L	I	V	L	15
316	L	A	V	L	E	A	I	L	L	15
369	I	A	Y	W	A	M	T	A	L	15
461	G	Q	C	V	L	A	G	A	F	15
470	A	S	F	Y	W	A	F	H	K	15

232

TABLE XXX-continued

518	I	L	E	Y	I	D	H	K	L	15
542	K	C	C	L	W	C	L	E	K	15
543	C	C	L	W	C	L	E	K	F	15
547	C	L	E	K	F	I	K	F	L	15
567	G	K	N	F	C	V	S	A	K	15
579	M	L	L	M	R	N	I	V	R	15
586	V	R	V	V	V	L	D	K	V	15
593	K	V	T	D	L	L	L	F	F	15
596	D	L	L	L	F	F	G	K	L	15
607	V	G	G	V	G	V	L	S	F	15
609	G	V	G	V	L	S	F	F	F	15
622	I	P	G	L	G	K	D	F	K	15
651	I	A	S	G	F	F	S	V	F	15
684	P	Y	Y	M	S	K	S	L	L	15
698	K	N	E	A	P	P	D	N	K	15
34	D	V	I	C	C	V	L	F	L	14
38	C	V	L	F	L	L	F	I	L	14
61	P	R	Q	V	L	Y	P	R	N	14
75	C	G	M	G	E	N	K	D	K	14
83	K	P	Y	L	L	Y	F	N	I	14
84	P	Y	L	L	Y	F	N	I	F	14
135	T	V	G	E	V	F	Y	T	K	14
148	C	L	P	G	V	P	W	N	M	14
158	V	I	T	S	L	Q	Q	E	L	14
162	L	Q	Q	E	L	C	P	S	F	14
164	Q	E	L	C	P	S	F	L	L	14
232	I	L	V	A	L	G	V	A	L	14
240	L	V	L	S	L	L	F	I	L	14
263	L	G	V	L	G	V	L	A	Y	14
267	G	V	L	A	Y	G	I	Y	Y	14
272	G	I	Y	Y	C	W	E	E	Y	14
278	E	E	Y	R	V	L	R	D	K	14
325	L	M	L	I	F	L	R	Q	R	14
379	L	A	T	S	G	Q	P	Q	Y	14
418	S	C	P	G	L	M	C	V	F	14
434	L	I	Q	R	S	V	F	N	L	14
437	R	S	V	F	N	L	Q	I	Y	14
450	L	F	W	T	L	N	W	V	L	14
452	W	T	L	N	W	V	L	A	L	14
464	V	L	A	G	A	F	A	S	F	14
485	F	P	L	I	S	A	F	I	R	14
487	L	I	S	A	F	I	R	T	L	14
488	I	S	A	F	I	R	T	L	R	14
489	S	A	F	I	R	T	L	R	Y	14
501	S	L	A	F	G	A	L	I	L	14
513	Q	I	A	R	V	I	L	E	Y	14
515	A	R	V	I	L	E	Y	I	D	14
552	I	K	F	L	N	R	N	A	Y	14
556	N	R	N	A	Y	I	M	I	A	14
558	N	A	Y	I	M	I	A	I	Y	14
560	Y	I	M	I	A	I	Y	G	K	14
575	K	N	A	F	M	L	L	M	R	14
585	I	V	R	V	V	V	L	D	K	14
595	T	D	L	L	L	F	F	G	K	14
613	L	S	F	F	F	F	S	G	R	14
643	T	S	I	L	G	A	Y	V	I	14
659	F	G	M	C	V	D	T	L	F	14
660	G	M	C	V	D	T	L	F	L	14
679	G	S	L	D	R	P	Y	Y	M	14
700	E	A	P	P	D	N	K	K	R	14
701	A	P	P	D	N	K	K	K	R	14
702	P	P	D	N	K	K	R	K	K	14
7	D	E	D	D	E	A	Y	G	K	13
36	I	C	C	V	L	F	L	L	F	13
172	L	P	S	A	P	A	L	G	R	13
241	V	L	S	L	L	F	I	L	L	13
249	L	L	R	L	V	A	G	P	L	13
250	L	R	L	V	A	G	P	L	V	13
273	I	Y	Y	C	W	E	E	Y	R	13
275	Y	C	W	E	E	Y	R	V	L	13
280	Y	R	V	L	R	D	K	G	A	13
294	G	F	T	T	N	L	S	A	Y	13
319	L	E	A	I	L	L	L	M	L	13
347	A	V	G	Q	M	M	S	T	M	13
348	V	G	Q	M	M	S	T	M	F	13
349	G	Q	M	M	S	T	M	F	Y	13
356	F	Y	P	L	V	T	F	V	L	13
357	Y	P	L	V	T	F	V	L	L	13
358	P	L	V	T	F	V	L	L	L	13
363	V	L	L	L	I	C	I	A	Y	13

233

TABLE XXX-continued

492	I	R	T	L	R	Y	H	T	G	13
495	L	R	Y	H	T	G	S	L	A	13
506	A	L	I	L	T	L	V	Q	I	13
526	L	R	G	V	Q	N	P	V	A	13
545	L	W	C	L	E	K	F	I	K	13
570	F	C	V	S	A	K	N	A	F	13
572	V	S	A	K	N	A	F	M	L	13
582	M	R	N	I	V	R	V	V	V	13
590	V	L	D	K	V	T	D	L	L	13
592	D	K	V	T	D	L	L	L	F	13
610	V	G	V	L	S	F	F	F	F	13
637	Y	W	L	P	I	M	T	S	I	13
648	A	Y	V	I	A	S	G	F	F	13
653	S	G	F	F	S	V	F	G	M	13
666	L	F	L	C	F	L	E	D	L	13
681	L	D	R	P	Y	Y	M	S	K	13
682	D	R	P	Y	Y	M	S	K	S	13
685	Y	Y	M	S	K	S	L	L	K	13
686	Y	M	S	K	S	L	L	K	I	13
29	N	R	S	C	T	D	V	I	C	12
32	C	T	D	V	I	C	C	V	L	12
33	T	D	V	I	C	C	V	L	F	12
35	V	I	C	C	V	L	F	L	L	12
57	L	Y	G	D	P	R	Q	V	L	12
58	Y	G	D	P	R	Q	V	L	Y	12
79	E	N	K	D	K	P	Y	L	L	12
80	N	K	D	K	P	Y	L	L	Y	12
93	S	C	I	L	S	S	N	I	I	12
100	I	I	S	V	A	E	N	G	L	12
121	P	E	D	P	W	T	V	G	K	12
132	F	S	Q	T	V	G	E	V	F	12
144	N	R	N	F	C	L	P	G	V	12
151	G	V	P	W	N	M	T	V	I	12
163	Q	Q	E	L	C	P	S	F	L	12
190	P	A	L	P	G	I	T	N	D	12
193	P	G	I	T	N	D	T	T	I	12
239	A	L	V	L	S	L	L	F	I	12
276	C	W	E	E	Y	R	V	L	R	12
302	Y	Q	S	V	Q	E	T	W	L	12
305	V	Q	E	T	W	L	A	A	L	12
315	V	L	A	V	L	E	A	I	L	12
320	E	A	I	L	L	L	M	L	I	12
328	I	F	L	R	Q	R	I	R	I	12
343	E	A	S	K	A	V	G	Q	M	12
371	Y	W	A	M	T	A	L	Y	L	12
386	Q	Y	V	L	W	A	S	N	I	12
393	N	I	S	S	P	G	C	E	K	12
406	T	S	C	N	P	T	A	H	L	12
414	L	V	N	S	S	C	P	G	L	12
421	G	L	M	C	V	F	Q	G	Y	12
426	F	Q	G	Y	S	S	K	G	L	12
468	A	F	A	S	F	Y	W	A	F	12
490	A	F	I	R	T	L	R	Y	H	12
500	G	S	L	A	F	G	A	L	I	12
510	T	L	V	Q	I	A	R	V	I	12
519	L	E	Y	I	D	H	K	L	R	12
537	I	M	C	C	F	K	C	C	L	12
540	C	F	K	C	C	L	W	C	L	12
553	K	F	L	N	R	N	A	Y	I	12
557	R	N	A	Y	I	M	I	A	I	12
562	M	I	A	I	Y	G	K	N	F	12
591	L	D	K	V	T	D	L	L	L	12
597	L	L	L	F	F	G	K	L	L	12
614	S	F	F	F	F	S	G	R	I	12
619	S	G	R	I	P	G	L	G	K	12
628	D	F	K	S	P	H	L	N	Y	12
631	S	P	H	L	N	Y	Y	W	L	12
634	L	N	Y	Y	W	L	P	I	M	12
658	V	F	G	M	C	V	D	T	L	12
662	C	V	D	T	L	F	L	C	F	12
663	V	D	T	L	F	L	C	F	L	12
673	D	L	E	R	N	N	G	S	L	12
687	M	S	K	S	L	L	K	I	L	12

234

TABLE XXX-continued

V3-HLA-B2705-9mers-24P4C12										
Each peptide is a portion of SEQ ID										
NO: 7; each start position is specified,										
the length of peptide is 9 amino										
acids, and the end position for each										
peptide is the start position plus eight.										
Pos	1	2	3	4	5	6	7	8	9	score
1	G	R	C	F	P	W	T	N	I	24
6	W	T	N	I	T	P	P	A	L	11
V5-HLA-B2705-9mers-24P4C12										
Each peptide is a portion of SEQ ID										
NO: 11; each start position is										
specified, the length of peptide is 9										
amino acids, and the end position										
for each peptide is the start position										
plus eight.										
Pos	1	2	3	4	5	6	7	8	9	score
4	A	I	L	L	L	V	L	I	F	17
5	I	L	L	L	V	L	I	F	L	17
6	L	L	L	V	L	I	F	L	R	16
2	L	E	A	I	L	L	L	V	L	14
8	L	V	L	I	F	L	R	Q	R	14
3	E	A	I	L	L	L	V	L	I	12
9	V	L	I	F	L	R	Q	R	I	11
V6-HLA-B2705-9mers-24P4C12										
Each peptide is a portion of SEQ										
ID NO: 13; each start position is										
specified, the length of peptide is 9										
amino acids, and the end position										
for each peptide is the start										
position plus eight.										
Pos	1	2	3	4	5	6	7	8	9	score
9	P	R	S	V	F	N	L	Q	I	19
5	K	G	L	I	P	R	S	V	F	17
2	Y	S	S	K	G	L	I	P	R	16
7	L	I	P	R	S	V	F	N	L	14
3	S	S	K	G	L	I	P	R	S	9
V7-HLA-B2705-9mers-24P4C12										
Each peptide is a portion of SEQ										
ID NO: 15; each start position is										
specified, the length of peptide is										
9 amino acids, and the end										
position for each peptide is the										
start position plus eight.										
Pos	1	2	3	4	5	6	7	8	9	score
8	A	V	G	Q	M	M	S	T	M	13
4	W	I	L	V	A	V	G	Q	M	12
5	I	L	V	A	V	G	Q	M	M	11
3	Y	W	I	L	V	A	V	G	Q	6
V8-HLA-B2705-9mers-24P4C12										
Each peptide is a portion of SEQ ID										
NO: 17; each start position is										
specified, the length of peptide is 9										
amino acids, and the end position										
for each peptide is the start position										
plus eight.										
Pos	1	2	3	4	5	6	7	8	9	score
16	G	H	V	F	Q	T	S	I	L	15
1	N	Y	Y	W	L	P	I	M	R	14
8	M	R	N	P	I	T	P	T	G	14
9	R	N	P	I	T	P	T	G	H	14
11	P	I	T	P	T	G	H	V	F	12
15	T	G	H	V	F	Q	T	S	I	11
19	F	Q	T	S	I	L	G	A	Y	10
2	Y	Y	W	L	P	I	M	R	N	8
4	W	L	P	I	M	R	N	P	I	7

235

TABLE XXX-continued

7	I	M	R	N	P	I	T	P	T	7
17	H	V	F	Q	T	S	I	L	G	7

V9-HLA-B2705-9mers-24P4C12
Each peptide is a portion of SEQ
ID NO: 19; each start position is
specified, the length of peptide is 9
amino acids, and the end position
for each peptide is the start
position plus eight.

Pos	1	2	3	4	5	6	7	8	9	score
18	G	Y	V	L	W	A	S	N	I	15
13	Q	P	A	T	L	G	Y	V	L	13
2	W	A	M	T	A	L	Y	P	L	12
9	P	L	P	T	Q	P	A	T	L	12
11	P	T	Q	P	A	T	L	G	Y	10
6	A	L	Y	P	L	P	T	Q	P	8
15	A	T	L	G	Y	V	L	W	A	7

TABLE XXXI

V1-HLA-B2709-9merse-24P4C12 Each peptide is a portion of SEQ ID NO: 3; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.										
Pos	1	2	3	4	5	6	7	8	9	score
332	Q	R	I	R	I	A	I	A	L	23
179	G	R	C	F	P	W	T	N	V	22
250	L	R	L	V	A	G	P	L	V	21
214	A	R	D	I	S	V	K	I	F	20
436	Q	R	S	V	F	N	L	Q	I	20
144	N	R	N	F	C	L	P	G	V	19
330	L	R	Q	R	I	R	I	A	I	19
582	M	R	N	I	V	R	V	V	V	19
586	V	R	V	V	V	L	D	K	V	19
255	G	P	L	V	L	V	L	I	L	17
583	R	N	I	V	R	V	V	V	L	17
251	R	L	V	A	G	P	L	V	L	16
683	R	P	Y	Y	M	S	K	S	L	16
78	G	E	N	K	D	K	P	Y	L	15
170	F	L	L	P	S	A	P	A	L	15
334	I	R	I	A	I	A	L	L	K	15
446	G	V	L	G	L	F	W	T	L	15
620	G	R	I	P	G	L	G	K	D	15
647	G	A	Y	V	I	A	S	G	F	15
660	G	M	C	V	D	T	L	F	L	15
49	I	V	V	G	I	V	A	W	L	14
228	S	W	Y	W	I	L	V	A	L	14
234	V	A	L	G	V	A	L	V	L	14
244	L	L	F	I	L	L	L	R	L	14
317	A	V	L	E	A	I	L	L	L	14
333	R	I	R	I	A	I	A	L	L	14
452	W	T	L	N	W	V	L	A	L	14
602	G	K	L	F	V	V	G	G	V	14
626	G	K	D	L	K	S	P	H	L	14
679	G	S	L	D	R	P	Y	Y	M	14
23	F	R	G	P	I	K	N	R	S	13
34	D	V	I	C	C	V	L	F	L	13
83	K	P	Y	L	L	Y	F	N	I	13
107	G	L	Q	C	P	T	P	Q	V	13
204	G	I	S	G	L	I	D	S	L	13
232	I	L	V	A	L	G	V	A	L	13
236	L	G	V	A	L	V	L	S	L	13
237	G	V	A	L	V	L	S	L	L	13
240	L	V	L	S	L	L	F	I	L	13
242	L	S	L	L	F	I	L	L	L	13
253	V	A	G	P	L	V	L	V	L	13
291	S	Q	L	G	F	T	T	N	L	13
311	A	A	L	I	V	L	A	V	L	13
322	I	L	L	L	M	L	I	F	L	13

236

TABLE XXXI-continued

357	Y	P	L	V	T	F	V	L	L	13
358	P	L	V	T	F	V	L	L	L	13
369	I	A	Y	W	A	M	T	A	L	13
440	F	N	L	Q	I	Y	G	V	L	13
442	L	Q	I	Y	G	V	L	G	L	13
449	G	L	F	W	T	L	N	W	V	13
496	R	Y	H	T	G	S	L	A	F	13
500	G	S	L	A	F	G	A	L	I	13
515	A	R	V	I	L	E	Y	I	D	13
557	R	N	A	Y	I	M	I	A	I	13
589	V	V	L	D	K	V	T	D	L	13
15	K	P	V	K	Y	D	P	S	F	12
38	C	V	L	F	L	L	F	I	L	12
45	I	L	G	Y	I	V	V	G	I	12
56	W	L	Y	G	D	P	R	Q	V	12
61	P	R	Q	V	L	Y	P	R	N	12
81	K	D	K	P	Y	L	L	Y	F	12
158	V	I	T	S	L	Q	Q	E	L	12
164	Q	E	L	C	P	S	F	L	L	12
258	V	L	V	L	I	L	G	V	L	12
261	L	I	L	G	V	L	G	V	L	12
287	G	A	S	I	S	Q	L	G	F	12
308	T	W	L	A	A	L	I	V	L	12
316	L	A	V	L	E	A	I	L	L	12
321	A	I	L	L	L	M	L	I	F	12
328	I	F	L	R	Q	R	I	R	I	12
355	M	F	Y	P	L	V	T	F	V	12
371	Y	W	A	M	T	A	L	Y	L	12
414	L	V	N	S	S	C	P	G	L	12
432	K	G	L	I	Q	R	S	V	F	12
434	L	I	Q	R	S	V	F	N	L	12
461	G	Q	C	V	L	A	G	A	F	12
492	I	R	T	L	R	Y	H	T	G	12
495	L	R	Y	H	T	G	S	L	A	12
501	S	L	A	F	G	A	L	I	L	12
503	A	F	G	A	L	I	L	T	L	12
506	A	L	I	L	T	L	V	Q	I	12
518	I	L	E	Y	I	D	H	K	L	12
553	K	F	L	N	R	N	A	Y	I	12
593	K	V	T	D	L	L	L	F	F	12
596	D	L	L	L	F	F	G	K	L	12
597	L	L	L	F	F	G	K	L	L	12
605	L	V	V	G	G	V	G	V	L	12
608	G	G	V	G	V	L	S	F	F	12
621	R	I	P	G	L	G	K	D	F	12
637	Y	W	L	P	I	M	T	S	I	12
666	L	F	L	C	F	L	E	D	L	12
684	P	Y	Y	M	S	K	S	L	L	12
5	Q	R	D	E	D	D	E	A	Y	11
28	K	N	R	S	C	T	D	V	I	11
29	N	R	S	C	T	D	V	I	C	11
32	C	T	D	V	I	C	C	V	L	11
41	F	L	L	F	I	L	G	Y	I	11
42	L	L	F	I	L	G	Y	I	V	11
46	L	G	Y	I	V	V	G	I	V	11
67	P	R	N	S	T	G	A	Y	C	11
79	E	N	K	D	K	P	Y	L	L	11
87	L	Y	F	N	I	F	S	C	I	11
100	I	I	S	V	A	E	N	G	L	11
128	G	K	N	E	F	S	Q	T	V	11
139	V	F	Y	T	K	N	R	N	F	11
151	G	V	P	W	N	M	T	V	I	11
184	W	T	N	V	T	P	P	A	L	11
217	I	S	V	K	I	F	W	D	F	11
225	F	A	Q	S	W	Y	W	I	L	11
230	Y	W	I	L	V	A	L	G	V	11
238	V	A	L	V	L	S	L	L	F	11
239	A	L	V	L	S	L	L	F	I	11
249	L	L	R	L	V	A	G	P	L	11
257	L	V	L	V	L	I	L	G	V	11
260	V	L	I	L	G	V	L	G	V	11
280	Y	R	V	L	R	D	K	G	A	11
283	L	R	D	K	G	A	S	I	S	11
285	D	K	G	A	S	I	S	Q	L	11
297	T	N	L	S	A	Y	Q	S	V	11
310	L	A	A	L	I	V	L	A	V	11
314	I	V	L	A	V	L	E	A	I	11
319	L	E	A	I	L	L	L	M	L	11
351	M	M	S	T	M	F	Y	P	L	11
354	T	M	F	Y	P	L	V	T	F	11

237

TABLE XXXI-continued

381	T	S	G	Q	P	Q	Y	V	L	11
386	Q	Y	V	L	W	A	S	N	I	11
427	Q	G	Y	S	S	K	G	L	I	11
480	Q	D	I	P	T	F	P	L	I	11
483	P	T	F	P	L	I	S	A	F	11
509	L	T	L	V	Q	I	A	R	V	11
510	T	L	V	Q	I	A	R	V	I	11
511	L	V	Q	I	A	R	V	I	L	11
526	L	R	G	V	Q	N	P	V	A	11
534	A	R	C	I	M	C	C	F	K	11
537	I	M	C	C	F	K	C	C	L	11
564	A	I	Y	G	K	N	F	C	V	11
572	V	S	A	K	N	A	F	M	L	11
591	L	D	K	V	T	D	L	L	L	11
592	D	K	V	T	D	L	L	L	F	11
598	L	L	F	F	G	K	L	L	V	11
599	L	F	F	G	K	L	L	V	V	11
609	G	V	G	V	L	S	F	F	F	11
614	S	F	F	F	F	S	G	R	I	11
617	F	F	S	G	R	I	P	G	L	11
631	S	P	H	L	N	Y	Y	W	L	11
634	L	N	Y	Y	W	L	P	I	M	11
643	T	S	I	L	G	A	Y	V	I	11
653	S	G	F	F	S	V	F	G	M	11
658	V	F	G	M	C	V	D	T	L	11
663	V	D	T	L	F	L	C	F	L	11
675	E	R	N	N	G	S	L	D	R	11
687	M	S	K	S	L	L	K	I	L	11

V3-HLA-B2709-9mers-24P4C12
Each peptide is a portion of SEQ ID NO: 7; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.

Pos	1	2	3	4	5	6	7	8	9	score
1	G	R	C	F	P	W	T	N	I	22
6	W	T	N	I	T	P	P	A	L	11
9	I	T	P	P	A	L	P	G	I	11

V5-B2709-9mers-24P4C12
Each peptide is a portion of SEQ ID NO: 11; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.

Pos	1	2	3	4	5	6	7	8	9	score
4	A	I	L	L	L	V	L	I	F	13
5	I	L	L	L	V	L	I	F	L	13
2	L	E	A	I	L	L	L	V	L	11
1	V	L	E	A	I	L	L	L	V	10
3	E	A	I	L	L	L	V	L	I	10
9	V	L	I	F	L	R	Q	R	I	10

V6-HLA-B2709-9mers-24P4C12
Each peptide is a portion of SEQ ID NO: 13; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.

Pos	1	2	3	4	5	6	7	8	9	score
9	P	R	S	V	F	N	L	Q	I	20
5	K	G	L	I	P	R	S	V	F	12
7	L	I	P	R	S	V	F	N	L	12
4	S	K	G	L	I	P	R	S	V	9

238

TABLE XXXI-continued

V7-HLA-B2709-9mers-24P4C12 Each peptide is a portion of SEQ ID NO: 15; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.										
5	Pos	1	2	3	4	5	6	7	8	9 score
	1	S	W	Y	W	I	L	V	A	V 12
	4	W	I	L	V	A	V	G	Q	M 12
	5	I	L	V	A	V	G	Q	M	M 10
	8	A	V	G	Q	M	M	S	T	M 9
V8-HLA-B2709-9mers-24P4C12 Each peptide is a portion of SEQ ID NO: 17; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.										
10	Pos	1	2	3	4	5	6	7	8	9 score
	16	G	H	V	F	Q	T	S	I	L 14
	8	M	R	N	P	I	T	P	T	G 13
	11	P	I	T	P	T	G	H	V	F 10
	10	N	P	I	T	P	T	G	H	V 9
15	4	W	L	P	I	M	R	N	P	I 8
	15	T	G	H	V	F	Q	T	S	I 8
	20	Q	T	S	I	L	G	A	Y	V 8
V9-HLA-B2709-9mers-24P4C12 Each peptide is a portion of SEQ ID NO: 19; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.										
20	Pos	1	2	3	4	5	6	7	8	9 score
	18	G	Y	V	L	W	A	S	N	I 14
	2	W	A	M	T	A	L	Y	P	L 11
	13	Q	P	A	T	L	G	Y	V	L 11
	9	P	L	P	T	Q	P	A	T	L 10
25	12	T	Q	P	A	T	L	G	Y	V 8
TABLE XXXII										
V1-HLA-B4402-9mers-24P4C12 Each peptide is a portion of SEQ ID NO: 3; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.										
30	Pos	1	2	3	4	5	6	7	8	9 score
	164	Q	E	L	C	P	S	F	L	L 22
	319	L	E	A	I	L	L	L	M	L 22
	222	F	E	D	F	A	Q	S	W	Y 21
	78	G	E	N	K	D	K	P	Y	L 20
35	306	Q	E	T	W	L	A	A	L	I 20
	483	P	T	F	P	L	I	S	A	F 20
	317	A	V	L	E	A	I	L	L	L 19
	332	Q	R	I	R	I	A	I	A	L 19
	503	A	F	G	A	L	I	L	T	L 18
40	506	A	L	I	L	T	L	V	Q	I 18
	552	I	K	F	L	N	R	N	A	Y 18
	58	Y	G	D	P	R	Q	V	L	Y 17
	170	F	L	L	P	S	A	P	A	L 17
	214	A	R	D	I	S	V	K	I	F 17

Figure 2G-3

440 I K F L N R N A Y I M I A I Y G K N F C
1321 TTATCAAGTTCCTAAACCGCAATGCATACATCATGATCGCCATCTACGGGAAGAATTTCT
460 V S A K N A F M L L M R N I V R V V V L
1381 GTGTCTCAGCCAAAAATGCGTTCATGCTACTCATGCGAAACATTGTCAGGGTGGTCGTCC
480 D K V T D L L L F F G K L L V V G G V G
1441 TGGACAAAGTCACAGACCTGCTGCTGTTCTTTGGGAAGCTGCTGGTGGTCGGAGGCGTGG
500 V L S F F F F S G R I P G L G R D F K S
1501 GGGTCCTGTCCTTCTTTTTTTCTCCGGTCGCATCCCGGGCTGGGTAAAGACTTTAAGA
520 P H L N Y Y W L P I M T S I L G A Y V I
1561 GCCCCCACCTCAACTATTACTGGCTGCCCATCATGACCTCCATCCTGGGGGCCTATGTCA
540 A S G F F S V F G M C V D I L F L C F L
1621 TCGCCAGCGGCTTCTTCAGCGTTTTTCGGCATGTGTGTGGACACGCTCTTCCTCTGCTTCC
560 E D L E R N N G S L D R P Y Y M S K S L
1681 TGGAAGACCTGGAGCGGAACAACGGCTCCCTGGACCGGCCCTACTACATGTCCAAGAGCC
580 L K I L G K K N E A P P D N K K R K K *
1741 TTCTAAAGATTCTGGGCAAGAAGAACGAGGCGCCCCGGACAACAAGAAGAGGAAGAAGT
1801 GAcagctccggccctgatccaggactgcacccccccccaccgtccagccatccaacctc
1861 acttcgccttacaggtctccatTTTTgtggtaaaaaaaggTTTTtaggccaggcgccgtggc
1921 tcacgcctgtaatccaacactttgagaggctgaggcgggaggatcacctgagtcaggagt
1981 tcgagaccagcctggccaacatggtgaaacctccgtctctattaaaaatacaaaaattag
2041 ccgagagtggtggcatgcacctgtcatcccagctactcgggagggtgaggcaggagaatc
2101 gcttgaacccgggaggcagaggttgagtgagccgagatcgcgccactgcactccaacct
2161 ggggtgacagactctgtctccaaaacaaaacaaaacaaaagattttattaaagatat
2221 tttgttaactcagtaaaaaaaaaaaaaaaaaa

239

TABLE XXXII-continued

242	L	S	L	L	F	I	L	L	17
583	R	N	I	V	R	V	V	V	17
11	E	A	Y	G	K	P	V	K	16
40	L	F	L	L	F	I	L	G	16
48	Y	I	V	V	G	I	V	A	16
81	K	D	K	P	Y	L	L	Y	16
121	P	E	D	P	W	T	V	G	16
228	S	W	Y	W	I	L	V	A	16
253	V	A	G	P	L	V	L	V	16
254	A	G	P	L	V	L	V	L	16
311	A	A	L	I	V	L	A	V	16
320	E	A	I	L	L	L	M	L	16
321	A	I	L	L	L	M	L	I	16
363	V	L	L	L	I	C	I	A	16
382	S	G	Q	P	Q	Y	V	L	16
452	W	T	L	N	W	V	L	A	16
480	Q	D	I	P	T	F	P	L	16
487	L	I	S	A	F	I	R	T	16
489	S	A	F	I	R	T	L	R	16
617	F	F	S	G	R	I	P	G	16
629	F	K	S	P	H	L	N	Y	16
699	N	E	A	P	P	D	N	K	16
34	D	V	I	C	C	V	L	F	15
79	E	N	K	D	K	P	Y	L	15
130	N	E	F	S	Q	T	V	G	15
154	W	N	M	T	V	I	T	S	15
204	G	I	S	G	L	I	D	S	15
234	V	A	L	G	V	A	L	V	15
241	V	L	S	L	L	F	I	L	15
263	L	G	V	L	G	V	L	A	15
278	E	E	Y	R	V	L	R	D	15
294	G	F	T	T	N	L	S	A	15
354	T	M	F	Y	P	L	V	T	15
370	A	Y	W	A	M	T	A	L	15
399	C	E	K	V	P	I	N	T	15
442	L	Q	I	Y	G	V	L	G	15
468	A	F	A	S	F	Y	W	A	15
477	H	K	P	Q	D	I	P	T	15
499	T	G	S	L	A	F	G	A	15
513	Q	I	A	R	V	I	L	E	15
547	C	L	E	K	F	I	K	F	15
66	Y	P	R	N	S	T	G	A	14
80	N	K	D	K	P	Y	L	L	14
84	P	Y	L	L	Y	F	N	I	14
93	S	C	I	L	S	S	N	I	14
104	A	E	N	G	L	Q	C	P	14
193	P	G	I	T	N	D	T	T	14
223	E	D	F	A	Q	S	W	Y	14
239	A	L	V	L	S	L	L	F	14
244	L	L	F	I	L	L	L	R	14
258	V	L	V	L	I	L	G	V	14
261	L	I	L	G	V	L	G	V	14
285	D	K	G	A	S	I	S	Q	14
291	S	Q	L	G	F	T	T	N	14
301	A	Y	Q	S	V	Q	E	T	14
305	V	Q	E	T	W	L	A	A	14
308	T	W	L	A	A	L	I	V	14
316	L	A	V	L	E	A	I	L	14
322	I	L	L	L	M	L	I	F	14
330	L	R	Q	R	I	R	I	A	14
333	R	I	R	I	A	I	A	L	14
356	F	Y	P	L	V	T	F	V	14
357	Y	P	L	V	T	F	V	L	14
358	P	L	V	T	F	V	L	L	14
364	L	L	L	I	C	I	A	Y	14
418	S	C	P	G	L	M	C	V	14
432	K	G	L	I	Q	R	S	V	14
446	G	V	L	G	L	F	W	T	14
496	R	Y	H	T	G	S	L	A	14
546	W	C	L	E	K	F	I	K	14
558	N	A	Y	I	M	I	A	I	14
573	S	A	K	N	A	F	M	L	14
577	A	F	M	L	L	M	R	N	14
592	D	K	V	T	D	L	L	L	14
593	K	V	T	D	L	L	L	F	14
596	D	L	L	L	F	F	G	K	14
597	L	L	L	F	F	G	K	L	14
621	R	I	P	G	L	G	K	D	14
641	I	M	T	S	I	L	G	A	14
643	T	S	I	L	G	A	Y	V	14

240

TABLE XXXII-continued

651	I	A	S	G	F	F	S	V	F	14
662	C	V	D	T	L	F	L	C	F	14
671	L	E	D	L	E	R	N	N	G	14
678	N	G	S	L	D	R	P	Y	Y	14
5	Q	R	D	E	D	D	E	A	Y	13
7	D	E	D	D	E	A	Y	G	K	13
32	C	T	D	V	I	C	C	V	L	13
36	I	C	C	V	L	F	L	L	F	13
49	I	V	V	G	I	V	A	W	L	13
57	L	Y	G	D	P	R	Q	V	L	13
77	M	G	E	N	K	D	K	P	Y	13
87	L	Y	F	N	I	F	S	C	I	13
137	G	E	V	F	Y	T	K	N	R	13
146	N	F	C	L	P	G	V	P	W	13
174	S	A	P	A	L	G	R	C	F	13
176	P	A	L	G	R	C	F	P	W	13
184	W	T	N	V	T	P	P	A	L	13
187	V	T	P	P	A	L	P	G	I	13
200	T	I	Q	Q	G	I	S	G	L	13
209	I	D	S	L	N	A	R	D	I	13
213	N	A	R	D	I	S	V	K	I	13
232	I	L	V	A	L	G	V	A	L	13
237	G	V	A	L	V	L	S	L	L	13
238	V	A	L	V	L	S	L	L	F	13
251	R	L	V	A	G	P	L	V	L	13
255	G	P	L	V	L	V	L	I	L	13
277	W	E	E	Y	R	V	L	R	D	13
342	K	E	A	S	K	A	V	G	Q	13
351	M	M	S	T	M	F	Y	P	L	13
440	F	N	L	Q	I	Y	G	V	L	13
443	Q	I	Y	G	V	L	G	L	F	13
448	L	G	L	F	W	T	L	N	W	13
461	G	Q	C	V	L	A	G	A	F	13
466	A	G	A	F	A	S	F	Y	W	13
501	S	L	A	F	G	A	L	I	L	13
518	I	L	E	Y	I	D	H	K	L	13
519	L	E	Y	I	D	H	K	L	R	13
529	V	Q	N	P	V	A	R	C	I	13
543	C	C	L	W	C	L	E	K	F	13
570	F	C	V	S	A	K	N	A	F	13
589	V	V	L	D	K	V	T	D	L	13
590	V	L	D	K	V	T	D	L	L	13
605	L	V	V	G	G	V	G	V	L	13
631	S	P	H	L	N	Y	Y	W	L	13
637	Y	W	L	P	I	M	T	S	I	13
648	A	Y	V	I	A	S	G	F	F	13
674	L	E	R	N	N	G	S	L	D	13
687	M	S	K	S	L	L	K	I	L	13
33	T	D	V	I	C	C	V	L	F	12
35	V	I	C	C	V	L	F	L	L	12
38	C	V	L	F	L	L	F	I	L	12
50	V	V	G	I	V	A	W	L	Y	12
100	I	I	S	V	A	E	N	G	L	12
132	F	S	Q	T	V	G	E	V	F	12
133	S	Q	T	V	G	E	V	F	Y	12
139	V	F	Y	T	K	N	R	N	F	12
141	Y	T	K	N	R	N	F	C	L	12
163	Q	Q	E	L	C	P	S	F	L	12
217	I	S	V	K	I	F	E	D	F	12
221	I	F	E	D	F	A	Q	S	W	12
236	L	G	V	A	L	V	L	S	L	12
240	L	V	L	S	L	L	F	I	L	12
249	L	L	R	L	V	A	G	P	L	12
267	G	V	L	A	Y	G	I	Y	Y	12
269	L	A	Y	G	I	Y	Y	C	W	12
275	Y	C	W	E	E	Y	R	V	L	12
287	G	A	S	I	S	Q	L	G	F	12
314	I	V	L	A	V	L	E	A	I	12
326	M	L	I	F	L	R	Q	R	I	12
328	I	F	L	R	Q	R	I	R	I	12
349	G	Q	M	M	S	T	M	F	Y	12
369	I	A	Y	W	A	M	T	A	L	12
371	Y	W	A	M	T	A	L	Y	L	12
406	T	S	C	N	P	T	A	H	L	12
421	G	L	M	C	V	F	Q	G	Y	12
426	F	Q	G	Y	S	S	K	G	L	12
434	L	I	Q	R	S	V	F	N	L	12
437	R	S	V	F	N	L	Q	I	Y	12
450	L	F	W	T	L	N	W	V	L	12
457	V	L	A	L	G	Q	C	V	L	12

TABLE XXXII-continued

464	V	L	A	G	A	F	A	S	F	12
479	P	Q	D	I	P	T	F	P	L	12
510	T	L	V	Q	I	A	R	V	I	12
511	L	V	Q	I	A	R	V	I	L	12
548	L	E	K	F	I	K	F	L	N	12
553	K	F	L	N	R	N	A	Y	I	12
557	R	N	A	Y	I	M	I	A	I	12
562	M	I	A	I	Y	G	K	N	F	12
572	V	S	A	K	N	A	F	M	L	12
591	L	D	K	V	T	D	L	L	L	12
607	V	G	G	V	G	V	L	S	F	12
608	G	G	V	G	V	L	S	F	F	12
610	V	G	V	L	S	F	F	F	F	12
630	K	S	P	H	L	N	Y	Y	W	12
638	W	L	P	I	M	T	S	I	L	12
647	G	A	Y	V	I	A	S	G	F	12
658	V	F	G	M	C	V	D	T	L	12
659	F	G	M	C	V	D	T	L	F	12
660	G	M	C	V	D	T	L	F	L	12
663	V	D	T	L	F	L	C	F	L	12
666	L	F	L	C	F	L	E	D	L	12
673	D	L	E	R	N	N	G	S	L	12
677	N	N	G	S	L	D	R	P	Y	12
683	R	P	Y	Y	M	S	K	S	L	12
686	Y	M	S	K	S	L	L	K	I	12
10	D	E	A	Y	G	K	P	V	K	11
15	K	P	V	K	Y	D	P	S	F	11
28	K	N	R	S	C	T	D	V	I	11
37	C	C	V	L	F	L	L	F	I	11
41	F	L	L	F	I	L	G	Y	I	11
45	I	L	G	Y	I	V	V	G	I	11
117	V	S	S	C	P	E	D	P	W	11
124	P	W	T	V	G	K	N	E	F	11
151	G	V	P	W	N	M	T	V	I	11
197	N	D	T	T	I	Q	Q	G	I	11
201	I	Q	Q	G	I	S	G	L	I	11
266	L	G	V	L	A	Y	G	I	Y	11
302	Y	Q	S	V	Q	E	T	W	L	11
359	L	V	T	F	V	L	L	L	I	11
361	T	F	V	L	L	L	I	C	I	11
379	L	A	T	S	G	Q	P	Q	Y	11
381	T	S	G	Q	P	Q	Y	V	L	11
436	Q	R	S	V	F	N	L	Q	I	11
444	I	Y	G	V	L	G	L	F	W	11
465	L	A	G	A	F	A	S	F	Y	11
474	W	A	F	H	K	P	Q	D	I	11
484	T	F	P	L	I	S	A	F	I	11
494	T	L	R	Y	H	T	G	S	L	11
533	V	A	R	C	I	M	C	C	F	11
538	M	C	C	F	K	C	C	L	W	11
540	C	F	K	C	C	L	W	C	L	11
614	S	F	F	F	F	S	G	R	I	11
626	G	K	D	F	K	S	P	H	L	11
628	D	F	K	S	P	H	L	N	Y	11
684	P	Y	Y	M	S	K	S	L	L	11
19	Y	D	P	S	F	R	G	P	I	10
83	K	P	Y	L	L	Y	F	N	I	10
88	Y	F	N	I	F	S	C	I	L	10
158	V	I	T	S	L	Q	Q	E	L	10
162	L	Q	Q	E	L	C	P	S	F	10
225	F	A	Q	S	W	Y	W	I	L	10
272	G	I	Y	Y	C	W	E	E	Y	10
315	V	L	A	V	L	E	A	I	L	10
348	V	G	Q	M	M	S	T	M	F	10
386	Q	Y	V	L	W	A	S	N	I	10
396	S	P	G	C	E	K	V	P	I	10
414	L	V	N	S	S	C	P	G	L	10
500	G	S	L	A	F	G	A	L	I	10
514	I	A	R	V	I	L	E	Y	I	10
537	I	M	C	C	F	K	C	C	L	10
544	C	L	W	C	L	E	K	F	I	10
555	L	N	R	N	A	Y	I	M	I	10
609	G	V	G	V	L	S	F	F	F	10

TABLE XXXII-continued

V3-HLA-B4402-9mers-24P4C12										
Each peptide is a portion of SEQ ID NO: 7; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.										
Pos	1	2	3	4	5	6	7	8	9	score
6	W	T	N	I	T	P	P	A	L	13
9	I	T	P	P	A	L	P	G	I	13
1	G	R	C	F	P	W	T	N	I	8
2	R	C	F	P	W	T	N	I	T	7
7	T	N	I	T	P	P	A	L	P	6
8	N	I	T	P	P	A	L	P	G	6
V5-HLA-B4402-9mers-24P4C12										
Each peptide is a portion of SEQ ID NO: 11; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.										
Pos	1	2	3	4	5	6	7	8	9	score
2	L	E	A	I	L	L	L	V	L	23
3	E	A	I	L	L	L	V	L	I	17
4	A	I	L	L	L	V	L	I	F	17
5	I	L	L	L	V	L	I	F	L	14
9	V	L	I	F	L	R	Q	R	I	12
V6-HLA-B4402-9mers-24P4C12										
Each peptide is a portion of SEQ ID NO: 13; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.										
Pos	1	2	3	4	5	6	7	8	9	score
5	K	G	L	I	P	R	S	V	F	14
7	L	I	P	R	S	V	F	N	L	13
9	P	R	S	V	F	N	L	Q	I	11
6	G	L	I	P	R	S	V	F	N	8
V7-HLA-B4402-9mers-24P4C12										
Each peptide is a portion of SEQ ID NO: 15; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.										
Pos	1	2	3	4	5	6	7	8	9	score
1	S	W	Y	W	I	L	V	A	V	6
3	Y	W	I	L	V	A	V	G	Q	6
8	A	V	G	Q	M	M	S	T	M	4
4	W	I	L	V	A	V	G	Q	M	3
2	W	Y	W	I	L	V	A	V	G	2
V8-HLA-B4402-9mers-24P4C12										
Each peptide is a portion of SEQ ID NO: 17; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.										
Pos	1	2	3	4	5	6	7	8	9	score
11	P	I	T	P	T	G	H	V	F	15
19	F	Q	T	S	I	L	G	A	Y	14
4	W	L	P	I	M	R	N	P	I	11

243

TABLE XXXII-continued

16	G	H	V	F	Q	T	S	I	L	11
15	T	G	H	V	F	Q	T	S	I	8

V9-HLA-B4402-9mers-24P4C12

Each peptide is a portion of SEQ ID NO: 19; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.

Pos	1	2	3	4	5	6	7	8	9	score
11	P	T	Q	P	A	T	L	G	Y	15
9	P	L	P	T	Q	P	A	T	L	14
2	W	A	M	T	A	L	Y	P	L	13
14	P	A	T	L	G	Y	V	L	W	13
13	Q	P	A	T	L	G	Y	V	L	12
18	G	Y	V	L	W	A	S	N	I	10
6	A	L	Y	P	L	P	T	Q	P	8
15	A	T	L	G	Y	V	L	W	A	7

TABLE XXXIII

V1-HLA-B5101-9mers-24P4C12										
Each peptide is a portion of SEQ ID NO: 3; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.										
Pos	1	2	3	4	5	6	7	8	9	score
234	V	A	L	G	V	A	L	V	L	27
213	N	A	R	D	I	S	V	K	I	25
46	L	G	Y	I	V	V	G	I	V	24
83	K	P	Y	L	L	Y	F	N	I	24
311	A	A	L	I	V	L	A	V	L	24
253	V	A	G	P	L	V	L	V	L	23
310	L	A	A	L	I	V	L	A	V	23
357	Y	P	L	V	T	F	V	L	L	23
369	I	A	Y	W	A	M	T	A	L	23
474	W	A	F	H	K	P	Q	D	I	23
514	I	A	R	V	I	L	E	Y	I	23
683	R	P	Y	Y	M	S	K	S	L	22
254	A	G	P	L	V	L	V	L	I	21
255	G	P	L	V	L	V	L	I	L	21
320	E	A	I	L	L	L	M	L	I	21
396	S	P	G	C	E	K	V	P	I	21
427	Q	G	Y	S	S	K	G	L	I	21
11	E	A	Y	G	K	P	V	K	Y	20
193	P	G	I	T	N	D	T	T	I	20
316	L	A	V	L	E	A	I	L	L	20
123	D	P	W	T	V	G	K	N	E	19
236	L	G	V	A	L	V	L	S	L	18
314	I	V	L	A	V	L	E	A	I	18
599	L	F	F	G	K	L	L	V	V	18
686	Y	M	S	K	S	L	L	K	I	18
60	D	P	R	Q	V	L	Y	P	R	17
125	P	G	V	P	W	N	M	T	V	17
225	F	A	Q	S	W	Y	W	I	L	17
261	L	I	L	G	V	L	G	V	L	17
269	L	A	Y	G	I	Y	Y	C	W	17
300	S	A	Y	Q	S	V	Q	E	T	17
504	F	G	A	L	I	L	T	L	V	17
558	N	A	Y	I	M	I	A	I	Y	17
573	S	A	K	N	A	F	M	L	L	17
651	I	A	S	G	F	F	S	V	F	17
182	F	P	W	T	N	V	T	P	P	16
192	L	P	G	I	T	N	D	T	T	16
328	I	F	L	R	Q	R	I	R	I	16
355	M	F	Y	P	L	V	T	F	V	16
359	L	V	T	F	V	L	L	L	I	16
458	L	A	L	G	Q	C	V	L	A	16
502	L	A	F	G	A	L	I	L	T	16
505	G	A	L	I	L	T	L	V	Q	16

244

TABLE XXXIII-continued

510	T	L	V	Q	I	A	R	V	I	16
581	L	M	R	N	I	V	R	V	V	16
631	S	P	H	L	N	Y	Y	W	L	16
9	D	D	E	A	Y	G	K	P	V	15
45	I	L	G	Y	I	V	V	G	I	15
56	W	L	Y	G	D	P	R	Q	V	15
110	C	P	T	P	Q	V	C	V	S	15
120	C	P	E	D	P	W	T	V	G	15
151	G	V	P	W	N	M	T	V	I	15
172	L	P	S	A	P	A	L	G	R	15
224	D	F	A	Q	S	W	Y	W	I	15
275	Y	C	W	E	E	Y	R	V	L	15
308	T	W	L	A	A	L	I	V	L	15
336	I	A	I	A	L	L	K	E	A	15
338	I	A	L	L	K	E	A	S	K	15
375	T	A	L	Y	L	A	T	S	G	15
485	F	P	L	I	S	A	F	I	R	15
529	V	Q	N	P	V	A	R	C	I	15
564	A	I	Y	G	K	N	F	C	V	15
582	M	R	N	I	V	R	V	V	V	15
596	D	L	L	L	F	F	G	K	L	15
637	Y	W	L	P	I	M	T	S	I	15
643	T	S	I	L	G	A	Y	V	I	15
647	G	A	Y	V	I	A	S	G	F	15
700	E	A	P	P	D	N	K	K	R	15
20	D	P	S	F	R	G	P	I	K	14
41	F	L	L	F	I	L	G	Y	I	14
43	L	F	I	L	G	Y	I	V	V	14
72	G	A	Y	C	G	M	G	E	N	14
87	L	Y	F	N	I	F	S	C	I	14
119	S	C	P	E	D	P	W	T	V	14
152	V	P	W	N	M	T	V	I	T	14
188	T	P	P	A	L	P	G	I	T	14
190	P	A	L	P	G	I	T	N	D	14
209	I	D	S	L	N	A	R	D	I	14
230	Y	W	I	L	V	A	L	G	V	14
238	V	A	L	V	L	S	L	L	F	14
257	L	V	L	V	L	I	L	G	V	14
409	N	P	T	A	H	L	V	N	S	14
411	T	A	H	L	V	N	S	S	C	14
450	L	F	W	T	L	N	W	V	L	14
465	L	A	G	A	F	A	S	F	Y	14
467	G	A	F	A	S	F	Y	W	A	14
482	I	P	T	F	P	L	I	S	A	14
499	T	G	S	L	A	F	G	A	L	14
509	L	T	L	V	Q	I	A	R	V	14
576	N	A	F	M	L	L	M	R	N	14
586	V	R	V	V	V	L	D	K	V	14
589	V	V	L	D	K	V	T	D	L	14
602	G	K	L	L	V	V	G	G	V	14
605	L	V	V	G	G	V	G	V	L	14
639	L	P	I	M	T	S	I	L	G	14
701	A	P	P	D	N	K	K	R	K	14
702	P	P	D	N	K	K	R	K	K	14
19	Y	D	P	S	F	R	G	P	I	13
28	K	N	R	S	C	T	D	V	I	13
34	D	V	I	C	C	V	L	F	L	13
54	V	A	W	L	Y	G	D	P	R	13
66	Y	P	R	N	S	T	G	A	Y	13
112	T	P	Q	V	C	V	S	S	C	13
149	L	P	G	V	P	W	N	M	T	13
174	S	A	P	A	L	G	R	C	F	13
176	P	A	L	G	R	C	F	P	W	13
187	V	T	P	P	A	L	P	G	I	13
189	P	P	A	L	P	G	I	T	N	13
201	I	Q	Q	G	I	S	G	L	I	13
239	A	L	V	L	S	L	G	F	I	13
252	L	V	A	G	P	L	V	L	V	13
282	V	L	R	D	K	G	A	S	I	13
285	D	K	G	A	S	I	S	Q	L	13
293	L	G	F	T	T	N	L	S	A	13
322	I	L	L	L	M	L	I	F	L	13
330	L	R	Q	R	I	R	I	A	I	13
340	L	L	K	E	A	S	K	A	V	13
343	E	A	S	K	A	V	G	Q	M	13
356	F	Y	P	L	V	T	F	V	L	13
361	T	F	V	L	L	L	I	C	I	13
384	Q	P	Q	Y	V	L	W	A	S	13
478	K	P	Q	D	I	P	T	F	P	13
487	L	I	S	A	F	I	R	T	L	13

245

TABLE XXXIII-continued

489	S	A	F	I	R	T	L	R	Y	13
500	G	S	L	A	F	G	A	L	I	13
506	A	L	I	L	T	L	V	Q	I	13
521	Y	I	D	H	K	L	R	G	V	13
531	N	P	V	A	R	C	I	M	C	13
553	K	F	L	N	R	N	A	Y	I	13
555	L	N	R	N	A	Y	I	M	I	13
563	I	A	I	Y	G	K	N	F	C	13
578	F	M	L	L	M	R	N	I	V	13
580	L	L	M	R	N	I	V	R	V	13

V3-HLA-B5101-9mers-24P4C12
Each peptide is a portion of SEQ ID NO: 7; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.

Pos	1	2	3	4	5	6	7	8	9	score
4	F	P	W	T	N	I	T	P	P	15
9	I	T	P	P	A	L	P	G	I	14
1	G	R	C	F	P	W	T	N	I	11
6	W	T	N	I	T	P	P	A	L	8

V5-HLA-B5101-9mers-24P4C12
Each peptide is a portion of SEQ ID NO: 11; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.

Pos	1	2	3	4	5	6	7	8	9	score
3	E	A	I	L	L	L	V	L	I	22
5	I	L	L	L	V	L	I	F	L	14
2	L	E	A	I	L	L	L	V	L	13
1	V	L	E	A	I	L	L	L	V	12
9	V	L	I	F	L	R	Q	R	I	12

V6-HLA-B5101-9mers-24P4C12
Each peptide is a portion of SEQ ID NO: 13; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.

Pos	1	2	3	4	5	6	7	8	9	score
8	I	P	R	S	V	F	N	L	Q	16
7	L	I	P	R	S	V	F	N	L	12
9	P	R	S	V	F	N	L	Q	I	12
5	K	G	L	I	P	R	S	V	F	11
4	S	K	G	L	I	P	R	S	V	10

V7-HLA-B5101-9mers-24P4C12
Each peptide is a portion of SEQ ID NO: 15; each start position is specified, the length of peptide is 9 amino acids, and the end position for each peptide is the start position plus eight.

Pos	1	2	3	4	5	6	7	8	9	score
1	S	W	Y	W	I	L	V	A	V	14
7	V	A	V	G	Q	M	M	S	T	12
2	W	Y	W	I	L	V	A	V	G	6
3	Y	W	I	L	V	A	V	G	Q	6

246

TABLE XXXIII-continued

	V8-HLA-B5101-9mers-24P4C12											
	Each peptide is a portion of SEQ ID											
5	NO: 17; each start position is											
	specified, the length of peptide is 9											
	amino acids, and the end position for											
	each peptide is the start position plus											
	eight.											
	Pos	1	2	3	4	5	6	7	8	9	score	
10	10	N	P	I	T	P	T	G	H	V	21	
	15	T	G	H	V	F	Q	T	S	I	18	
	13	T	P	T	G	H	V	F	Q	T	14	
	4	W	L	P	I	M	R	N	P	I	13	
	5	L	P	I	M	R	N	P	I	T	13	
15	V9-HLA-B5101-9mers-24P4C12											
	Each peptide is a portion of SEQ											
	ID NO: 19; each start position is											
	specified, the length of peptide is 9											
	amino acids, and the end position											
20	for each peptide is the start											
	position plus eight.											
	Pos	1	2	3	4	5	6	7	8	9	score	
25	13	Q	P	A	T	L	G	Y	V	L	20	
	2	W	A	M	T	A	L	Y	P	L	18	
	5	T	A	L	Y	P	L	P	T	Q	16	
	8	Y	P	L	P	T	Q	P	A	T	15	
	10	L	P	T	Q	P	A	T	L	G	14	
	12	T	Q	P	A	T	L	G	Y	V	13	
	17	L	G	Y	V	L	W	A	S	N	12	
30	9	P	L	P	T	Q	P	A	T	L	11	
	14	P	A	T	L	G	Y	V	L	W	11	
	18	G	Y	V	L	W	A	S	N	I	11	
35	TABLE XXXIV											
	V1-HLA-A1-10mers-24P4C12											
	Each peptide is a portion of SEQ											
	ID NO: 3; each start position is											
	specified, the length of peptide is											
40	10 amino acids, and the end											
	position for each peptide is the											
	start position plus nine.											
	Pos	1	2	3	4	5	6	7	8	9	0	score
45	221	I	<u>F</u>	E	D	F	A	<u>Q</u>	S	W	Y	25
	488	I	<u>S</u>	A	F	I	R	<u>T</u>	L	R	Y	25
	39	V	<u>L</u>	F	L	L	F	<u>I</u>	L	G	Y	23
	58	Y	<u>G</u>	D	P	R	Q	<u>V</u>	L	Y	P	23
	79	E	<u>N</u>	K	D	K	P	<u>Y</u>	L	L	Y	23
	262	I	<u>L</u>	G	V	L	G	<u>V</u>	L	A	Y	23
	512	V	<u>Q</u>	I	A	R	V	<u>I</u>	L	E	Y	22
50	627	K	<u>D</u>	F	K	S	P	<u>H</u>	L	N	Y	21
	132	F	<u>S</u>	Q	T	V	G	<u>E</u>	V	F	Y	20
	266	L	<u>G</u>	V	L	A	Y	<u>G</u>	I	Y	Y	20
	362	F	<u>V</u>	L	L	L	I	<u>C</u>	I	A	Y	20
	590	V	<u>L</u>	D	K	V	T	<u>D</u>	L	L	L	20
	594	V	<u>T</u>	D	L	L	L	<u>F</u>	F	G	K	20
55	318	V	<u>L</u>	E	A	I	L	<u>L</u>	L	M	L	19
	32	C	<u>T</u>	D	V	I	C	<u>C</u>	V	L	F	18
	49	I	<u>V</u>	V	G	I	V	<u>A</u>	W	L	Y	18
	378	Y	<u>L</u>	A	T	S	G	<u>Q</u>	P	Q	Y	18
	420	P	<u>G</u>	L	M	C	V	<u>F</u>	Q	G	Y	18
	464	V	<u>L</u>	A	G	A	F	<u>A</u>	S	F	Y	18
60	10	D	<u>E</u>	A	Y	G	K	<u>P</u>	V	K	Y	17
	57	L	<u>Y</u>	G	D	P	R	<u>Q</u>	V	L	Y	17
	121	P	<u>E</u>	D	P	W	T	<u>V</u>	G	K	N	17
	265	V	<u>L</u>	G	V	L	A	<u>Y</u>	G	I	Y	17
	271	Y	<u>G</u>	I	Y	Y	C	<u>W</u>	E	E	Y	17
	276	C	<u>W</u>	E	E	Y	R	<u>V</u>	L	R	D	17
	369	I	<u>A</u>	Y	W	A	M	<u>T</u>	A	L	Y	17
65	551	F	<u>I</u>	K	F	L	N	<u>R</u>	N	A	Y	17
	80	N	<u>K</u>	D	K	P	Y	<u>L</u>	L	Y	F	16

247

TABLE XXXIV-continued

348	V	<u>G</u>	Q	M	M	S	<u>T</u>	M	F	Y	16
676	R	<u>N</u>	N	G	S	L	<u>D</u>	R	P	Y	16
677	N	<u>N</u>	G	S	L	D	<u>R</u>	P	Y	Y	16
4	K	<u>Q</u>	R	D	E	D	<u>D</u>	E	A	Y	15
18	K	<u>Y</u>	D	P	S	F	<u>R</u>	G	P	I	15
65	L	<u>Y</u>	P	R	N	S	<u>T</u>	G	A	Y	15
76	G	<u>M</u>	G	E	N	K	<u>D</u>	K	P	Y	15
214	A	<u>R</u>	D	I	S	V	<u>K</u>	I	F	E	15
293	L	<u>G</u>	F	T	N	L	<u>S</u>	A	Y	15	
436	Q	<u>R</u>	S	V	F	N	<u>L</u>	Q	I	Y	15
479	P	<u>Q</u>	D	I	P	T	<u>F</u>	P	L	I	15
557	R	<u>N</u>	A	Y	I	M	<u>I</u>	A	I	Y	15
628	D	<u>F</u>	K	S	P	H	<u>L</u>	N	Y	Y	15
640	P	<u>I</u>	M	T	S	I	<u>L</u>	G	A	Y	15
664	D	<u>T</u>	L	F	L	C	<u>F</u>	L	E	D	15
283	L	<u>R</u>	D	K	G	A	<u>S</u>	I	S	Q	14
521	Y	<u>I</u>	D	H	K	L	<u>R</u>	G	V	Q	14
673	D	<u>L</u>	E	R	N	N	<u>G</u>	S	L	D	14
141	Y	<u>T</u>	K	N	R	N	<u>F</u>	C	L	P	13
305	V	<u>Q</u>	E	T	W	L	<u>A</u>	A	L	I	13
382	S	<u>G</u>	Q	P	Q	Y	<u>V</u>	L	W	A	13
407	S	<u>C</u>	N	P	T	A	<u>H</u>	L	V	N	13
518	I	<u>L</u>	E	Y	I	D	<u>H</u>	K	L	R	13
547	C	<u>L</u>	E	K	F	I	<u>K</u>	F	L	N	13
670	F	<u>L</u>	E	D	L	E	<u>R</u>	N	N	G	13
680	S	<u>L</u>	D	R	P	Y	<u>Y</u>	M	S	K	13
7	D	<u>E</u>	D	D	E	A	<u>Y</u>	G	K	P	12
35	V	<u>I</u>	C	C	V	L	<u>F</u>	L	L	F	12
159	I	<u>T</u>	S	L	Q	Q	<u>E</u>	L	C	P	12
163	Q	<u>Q</u>	E	L	C	P	<u>S</u>	F	L	L	12
242	L	<u>S</u>	L	L	F	I	<u>L</u>	L	L	R	12
618	F	<u>S</u>	G	R	I	P	<u>G</u>	L	G	K	12
626	G	<u>K</u>	D	F	K	S	<u>P</u>	H	L	N	12
698	K	<u>N</u>	E	A	P	P	<u>D</u>	N	K	K	12

V3-HLA-A1-10mers-24P4C12
Each peptide is a portion of SEQ ID NO: 7; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.

Pos	1	2	3	4	5	6	7	8	9	0	score
10	I	<u>T</u>	P	P	A	L	<u>P</u>	G	I	T	10
3	R	<u>C</u>	F	P	W	T	<u>N</u>	I	T	P	9
7	W	<u>T</u>	N	I	T	P	<u>P</u>	A	L	P	8
8	T	<u>N</u>	I	T	P	P	<u>A</u>	L	P	G	6
9	N	<u>I</u>	T	P	P	A	<u>L</u>	P	G	I	4

V5-HLA-A1-10mers-24P4C12
Each peptide is a portion of SEQ ID NO: 11; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.

Pos	1	2	3	4	5	6	7	8	9	0	score
2	V	<u>L</u>	E	A	I	L	<u>L</u>	L	V	L	19
7	L	<u>L</u>	L	V	L	I	<u>F</u>	L	R	Q	10
1	A	<u>V</u>	L	E	A	I	<u>L</u>	L	L	V	9

V6-HLA-A1-10mers-24P4C12
Each peptide is a portion of SEQ ID NO: 13; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.

Pos	1	2	3	4	5	6	7	8	9	0	score
10	P	<u>R</u>	S	V	F	N	<u>L</u>	Q	I	Y	15
1	Q	<u>G</u>	Y	S	S	K	<u>G</u>	L	I	P	7

248

TABLE XXXIV-continued

4	S	<u>S</u>	K	G	L	I	<u>P</u>	R	S	V	7
9	I	<u>P</u>	R	S	V	F	<u>N</u>	L	Q	I	7

V7-HLA-A1-10mers-24P4C12
Each peptide is a portion of SEQ ID NO: 15; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.

Pos	1	2	3	4	5	6	7	8	9	0	score
1	Q	<u>S</u>	W	Y	W	I	<u>L</u>	V	A	V	4
2	S	<u>W</u>	Y	W	I	L	<u>V</u>	A	V	G	4
4	Y	<u>W</u>	I	L	V	A	<u>V</u>	G	Q	M	3
5	W	<u>I</u>	L	V	A	V	<u>G</u>	Q	M	M	2
6	I	<u>L</u>	V	A	V	G	<u>Q</u>	M	M	S	2
8	V	<u>A</u>	V	G	Q	M	<u>M</u>	S	T	M	2
9	A	<u>V</u>	G	Q	M	M	<u>S</u>	T	M	F	2

V8-HLA-A1-10mers-24P4C12
Each peptide is a portion of SEQ ID NO: 17; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.

Pos	1	2	3	4	5	6	7	8	9	0	score
19	V	<u>F</u>	Q	T	S	I	<u>L</u>	G	A	Y	16
4	Y	<u>W</u>	L	P	I	M	<u>R</u>	N	P	I	7
13	I	<u>T</u>	P	T	G	H	<u>V</u>	F	Q	T	7
21	Q	<u>T</u>	S	I	L	G	<u>A</u>	Y	V	I	7

V9-HLA-A1-10mers-24P4C12
Each peptide is a portion of SEQ ID NO: 19; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.

Pos	1	2	3	4	5	6	7	8	9	0	score
11	L	<u>P</u>	T	Q	P	A	<u>T</u>	L	G	Y	21
12	P	<u>T</u>	Q	P	A	T	<u>L</u>	G	Y	V	10

TABLE XXXV

V1-HLA-A0201-10mers-24P4C12
Each peptide is a portion of SEQ ID NO: 3; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.

Pos	1	2	3	4	5	6	7	8	9	0	score
235	A	L	G	V	A	<u>L</u>	V	L	S	L	29
44	F	I	L	G	Y	<u>I</u>	V	V	G	I	28
232	I	L	V	A	L	<u>G</u>	V	A	L	V	28
243	S	L	L	F	I	<u>L</u>	L	L	R	L	28
309	W	L	A	A	L	<u>I</u>	V	L	A	V	28
579	M	L	L	M	R	<u>N</u>	I	V	R	V	28
244	L	L	F	I	L	<u>L</u>	L	R	L	V	27
260	V	L	I	L	G	<u>V</u>	L	G	V	L	27
433	G	L	I	Q	R	<u>S</u>	V	F	N	L	27
508	I	L	T	L	V	<u>Q</u>	I	A	R	V	27
580	L	L	M	R	N	<u>I</u>	V	R	V	V	27
598	L	L	F	F	G	<u>K</u>	L	L	V	V	27
48	Y	I	V	V	G	<u>I</u>	V	A	W	L	26
94	C	I	L	S	S	<u>N</u>	I	I	S	V	26
239	A	L	V	L	S	<u>L</u>	L	F	I	L	26
241	V	L	S	L	L	<u>F</u>	I	L	L	L	26

TABLE XXXV-continued

251	R	L	V	A	G	<u>P</u>	L	V	L	V	26
321	A	I	L	L	L	<u>M</u>	L	I	F	L	26
441	N	L	Q	I	Y	<u>G</u>	V	L	G	L	26
502	L	A	F	G	A	<u>L</u>	I	L	T	L	26
517	V	I	L	E	Y	<u>I</u>	D	H	K	L	26
603	K	L	L	V	V	<u>G</u>	V	G	V	L	26
604	L	L	V	V	G	<u>G</u>	V	G	V	L	26
45	I	L	G	Y	I	<u>V</u>	V	G	I	V	25
252	L	V	A	G	P	<u>L</u>	V	L	V	L	25
304	S	V	Q	E	T	<u>W</u>	L	A	A	L	25
312	A	L	I	V	L	<u>A</u>	V	L	E	A	25
318	V	L	E	A	I	<u>L</u>	L	L	M	L	25
486	P	L	I	S	A	<u>F</u>	I	R	T	L	25
657	S	V	F	G	M	<u>C</u>	V	D	T	L	25
665	T	L	F	L	C	<u>F</u>	L	E	D	L	25
248	L	L	L	R	L	<u>V</u>	A	G	P	L	24
259	L	V	L	I	L	<u>G</u>	V	L	G	V	24
310	L	A	A	L	I	<u>V</u>	L	A	V	L	24
339	A	L	L	K	E	<u>A</u>	S	K	A	V	24
597	L	L	L	P	F	<u>G</u>	K	L	L	V	24
41	F	L	L	F	I	<u>L</u>	G	Y	I	V	23
42	L	L	F	I	L	<u>G</u>	Y	I	V	V	23
56	W	L	Y	G	D	<u>P</u>	R	Q	V	L	23
231	W	I	L	V	A	<u>L</u>	G	V	A	L	23
249	L	L	R	L	V	<u>A</u>	G	P	L	V	23
256	P	L	V	L	V	<u>L</u>	I	L	G	V	23
313	L	I	V	L	A	<u>V</u>	L	E	A	I	23
315	V	L	A	V	L	<u>E</u>	A	I	L	L	23
438	S	V	F	N	L	<u>Q</u>	I	Y	G	V	23
459	A	L	G	Q	C	<u>V</u>	L	A	G	A	23
686	Y	M	S	K	S	<u>L</u>	L	K	I	L	23
99	N	I	I	S	V	<u>A</u>	E	N	G	L	22
257	L	V	L	V	L	<u>I</u>	L	G	V	L	22
354	T	M	F	Y	P	<u>L</u>	V	T	F	V	22
413	H	L	V	N	S	<u>S</u>	C	P	G	L	22
449	G	L	F	W	T	<u>L</u>	N	W	V	L	22
506	A	L	I	L	T	<u>L</u>	V	Q	I	A	22
510	T	L	V	Q	I	<u>A</u>	R	V	I	L	22
513	Q	I	A	R	V	<u>I</u>	L	E	Y	I	22
581	L	M	R	N	I	<u>V</u>	R	V	V	V	22
585	I	V	R	V	V	<u>V</u>	L	D	K	V	22
590	V	L	D	K	V	<u>T</u>	D	L	L	L	22
199	T	T	I	Q	Q	<u>G</u>	I	S	G	L	21
247	I	L	L	L	R	<u>L</u>	V	A	G	P	21
253	V	A	G	P	L	<u>V</u>	L	V	L	I	21
316	L	A	V	L	E	<u>A</u>	I	L	L	L	21
501	S	L	A	F	G	<u>A</u>	L	I	L	T	21
505	G	A	L	I	L	<u>T</u>	L	V	Q	I	21
641	I	M	T	S	I	<u>L</u>	G	A	Y	V	21
86	L	L	Y	F	N	<u>I</u>	F	S	C	I	20
95	I	L	S	S	N	<u>I</u>	I	S	V	A	20
191	A	L	P	G	I	<u>T</u>	N	D	T	T	20
238	V	A	L	V	L	<u>S</u>	L	L	F	I	20
261	L	I	L	G	V	<u>L</u>	G	V	L	A	20
314	I	V	L	A	V	<u>L</u>	E	A	I	L	20
325	L	M	L	I	F	<u>L</u>	R	Q	R	I	20
329	F	L	R	Q	R	<u>I</u>	R	I	A	I	20
350	Q	M	M	S	T	<u>M</u>	F	Y	P	L	20
358	P	L	V	T	F	<u>V</u>	L	L	L	I	20
368	C	I	A	Y	W	<u>A</u>	M	T	A	L	20
393	N	I	S	S	P	<u>G</u>	C	E	K	V	20
554	F	L	N	R	N	<u>A</u>	Y	I	M	I	20
596	D	L	L	L	F	<u>E</u>	G	K	L	L	20
645	I	L	G	A	Y	<u>G</u>	I	A	S	G	20
649	Y	V	I	A	S	<u>V</u>	F	A	S	V	20
34	D	V	I	C	C	<u>V</u>	L	F	L	L	19
64	V	L	Y	P	R	<u>N</u>	S	T	G	A	19
85	Y	L	L	Y	F	<u>N</u>	I	F	S	C	19
186	N	V	T	P	P	<u>A</u>	L	P	G	I	19
233	L	V	A	L	G	<u>V</u>	A	L	V	L	19
264	G	V	L	G	V	<u>L</u>	A	Y	G	I	19
317	A	V	L	E	A	<u>I</u>	L	L	L	M	19
327	L	I	F	L	R	<u>Q</u>	R	I	R	I	19
335	R	I	A	I	A	<u>L</u>	L	K	E	A	19
351	M	M	S	T	M	<u>F</u>	Y	P	L	V	19
357	Y	P	L	V	T	<u>F</u>	V	L	L	L	19
363	V	L	L	L	I	<u>C</u>	I	A	Y	W	19
364	L	L	L	I	C	<u>I</u>	A	Y	W	A	19
365	L	L	I	C	I	<u>A</u>	Y	W	A	M	19
380	A	T	S	G	Q	<u>P</u>	Q	Y	V	L	19

TABLE XXXV-continued

457	V	L	A	L	G	<u>Q</u>	C	V	L	A	19
536	C	I	M	C	C	<u>F</u>	K	C	C	L	19
588	V	V	V	L	D	<u>K</u>	V	T	D	L	19
633	H	L	N	Y	Y	<u>W</u>	L	P	I	M	19
644	S	I	L	G	A	<u>Y</u>	V	I	A	S	19
39	V	L	F	L	L	<u>F</u>	I	L	G	Y	18
157	T	V	I	T	S	<u>L</u>	Q	Q	E	L	18
203	Q	G	I	S	G	<u>L</u>	I	D	S	L	18
208	L	I	D	S	L	<u>N</u>	A	R	D	I	18
240	L	V	L	S	L	<u>L</u>	F	I	L	L	18
246	F	I	L	L	L	<u>R</u>	L	V	A	G	18
262	I	L	G	V	L	<u>G</u>	V	L	A	Y	18
281	R	V	L	R	D	<u>K</u>	G	A	S	I	18
322	I	L	L	L	M	<u>L</u>	I	F	L	R	18
332	Q	R	I	R	I	<u>A</u>	I	A	L	L	18
360	V	T	F	V	L	<u>L</u>	L	I	C	I	18
388	V	L	W	A	S	<u>N</u>	I	S	S	P	18
448	L	G	L	F	W	<u>T</u>	L	N	W	V	18
493	R	T	L	R	Y	<u>H</u>	T	G	S	L	18
525	K	L	R	G	V	<u>Q</u>	N	P	V	A	18
589	V	V	L	D	K	<u>V</u>	T	D	L	L	18
616	F	F	F	S	G	<u>R</u>	I	P	G	L	18
662	C	V	D	T	L	<u>F</u>	L	C	F	L	18
685	Y	Y	M	S	K	<u>S</u>	L	L	K	I	18
130	N	E	F	S	Q	<u>T</u>	V	G	E	V	17
143	K	N	R	N	F	<u>C</u>	L	P	G	V	17
148	C	L	P	G	V	<u>P</u>	W	N	M	T	17
170	F	L	L	P	S	<u>A</u>	P	A	L	G	17
211	S	L	N	A	R	<u>D</u>	I	S	V	K	17
227	Q	S	W	Y	W	<u>I</u>	L	V	A	L	17
254	A	G	P	L	V	<u>L</u>	V	L	I	L	17
296	T	T	N	L	S	<u>A</u>	Y	Q	S	V	17
324	L	L	M	L	I	<u>F</u>	L	R	Q	R	17
373	A	M	T	A	L	<u>Y</u>	L	A	T	S	17
481	D	I	P	T	F	<u>P</u>	L	I	S	A	17
546	W	C	L	E	K	<u>F</u>	I	K	F	L	17
563	I	A	I	Y	G	<u>K</u>	N	F	C	V	17
582	M	R	N	I	V	<u>R</u>	V	V	V	L	17
40	L	F	L	L	F	<u>I</u>	L	G	Y	I	16
108	L	Q	C	P	T	<u>P</u>	Q	V	C	V	16
118	S	S	C	P	E	<u>D</u>	P	W	T	V	16
169	S	F	L	L	P	<u>S</u>	A	P	A	L	16
200	T	I	Q	Q	G	<u>I</u>	S	G	L	I	16
207	G	L	I	D	S	<u>L</u>	N	A	R	D	16
212	L	N	A	R	D	<u>I</u>	S	V	K	I	16
236	L	G	V	A	L	<u>V</u>	L	S	L	L	16
292	Q	L	G	F	T	<u>T</u>	N	L	S	A	16
307	E	T	W	L	A	<u>A</u>	L	I	V	L	16
319	L	E	A	I	L	<u>L</u>	L	M	L	I	16
337	A	I	A	L	L	<u>K</u>	E	A	S	K	16
366	L	I	C	I	A	<u>Y</u>	W	A	M	T	16
405	N	T	S	C	N	<u>P</u>	T	A	H	L	16
451	F	W	T	L	N	<u>W</u>	V	L	A	L	16
456	W	V	L	A	L	<u>G</u>	Q	C	V	L	16
458	L	A	L	G	Q	<u>C</u>	V	L	A	G	16
503	A	F	G	A	L	<u>I</u>	L	T	L	V	16
509	L	T	L	V	Q	<u>I</u>	A	R	V	I	16
637	Y	W	L	P	I	<u>M</u>	T	S	I	L	16
33	T	D	V	I	C	<u>C</u>	V	L	F	L	15
36	I	C	C	V	L	<u>F</u>	L	L	F	I	15
90	N	I	F	S	C	<u>I</u>	L	S	S	N	15
161	S	L	Q	Q	E	<u>L</u>	C	P	S	F	15
225	F	A	Q	S	W	<u>Y</u>	W	I	L	V	15
234	V	A	L	G	V	<u>A</u>	L	V	L	S	15
250	L	R	L	V	A	<u>G</u>	P	L	V	L	15
284	R	D	K	G	A	<u>S</u>	I	S	Q	L	15
323	L	L	L	M	L	<u>I</u>	F	L	R	Q	15
340	L	L	K	E	A	<u>S</u>	K	A	V	G	15
378	Y	L	A	T	S	<u>G</u>	Q	P	Q	Y	15
379	L	A	T	S	G	<u>Q</u>	P	Q	Y	V	15
430	S	S	K	G	L	<u>I</u>	Q	R	S	V	15
464	V	L	A	G	A	<u>F</u>	A	S	F	Y	15
498	H	T	G	S	L	<u>A</u>	F	G	A	L	15
520	E	Y	I	D	H	<u>K</u>	L	R	G	V	15
539	C	C	F	K	C	<u>C</u>	L	W	C	L	15
601	F	G	K	L	L	<u>V</u>	V	G	G	V	15
690	S	L	L	K	I	<u>L</u>	G	K	K	N	15
26	P	I	K	N	R	<u>S</u>	C	T	D	V	14
30	R	S	C	T	D	<u>V</u>	I	C	C	V	14
37	C	C	V	L	F	<u>L</u>	L	F	I	L	14

TABLE XXXV-continued

102	S	V	A	E	N	<u>G</u>	L	Q	C	P	14
149	L	P	G	V	P	<u>W</u>	N	M	T	V	14
153	P	W	N	M	T	<u>V</u>	I	T	S	L	14
162	L	Q	Q	E	L	<u>C</u>	P	S	F	L	14
165	E	L	C	P	S	<u>F</u>	L	L	P	S	14
171	L	L	P	S	A	<u>P</u>	A	L	G	R	14
177	A	L	G	R	C	<u>F</u>	P	W	T	N	14
220	K	I	F	E	D	<u>F</u>	A	Q	S	W	14
273	I	Y	Y	C	W	<u>E</u>	E	Y	R	V	14
338	I	A	L	L	K	<u>E</u>	A	S	K	A	14
353	S	T	M	F	Y	<u>P</u>	L	V	T	F	14
370	A	Y	W	A	M	<u>T</u>	A	L	Y	L	14
395	S	S	P	G	C	<u>E</u>	K	V	P	I	14
416	N	S	S	C	P	<u>G</u>	L	M	C	V	14
445	Y	G	V	L	G	<u>L</u>	F	W	T	L	14
483	P	T	F	P	L	<u>I</u>	S	A	F	I	14
500	G	S	L	A	F	<u>G</u>	A	L	I	L	14
571	C	V	S	A	K	<u>N</u>	A	F	M	L	14
577	A	F	M	L	L	<u>M</u>	R	N	I	V	14
595	T	D	L	L	L	<u>F</u>	F	G	K	L	14
606	V	V	G	G	V	<u>G</u>	V	L	S	F	14
639	L	P	I	M	T	<u>S</u>	I	L	G	A	14
680	S	L	D	R	P	<u>Y</u>	Y	M	S	K	14
693	K	I	L	G	K	<u>K</u>	N	E	A	P	14
694	I	L	G	K	K	<u>N</u>	E	A	P	P	14

V3-HLA-A0201-10mers-24P4C12											
Each peptide is a portion of SEQ											
ID NO: 7; each start position is											
specified, the length of peptide is											
10 amino acids, and the end											
position for each peptide is the											
start position plus nine.											
Pos	1	2	3	4	5	6	7	8	9	0	score
9	N	I	T	P	P	<u>A</u>	L	P	G	I	23
10	I	T	P	P	A	<u>L</u>	P	G	I	T	12

V5-HLA-A0201-10mers-24P4C12											
Each peptide is a portion of SEQ											
ID NO: 11; each start position is											
specified, the length of peptide is											
10 amino acids, and the end											
position for each peptide is the											
start position plus nine.											
Pos	1	2	3	4	5	6	7	8	9	0	score
5	A	I	L	L	L	<u>V</u>	L	I	F	L	26
1	A	V	L	E	A	<u>I</u>	L	L	L	V	25
2	V	L	E	A	I	<u>L</u>	L	L	V	L	25
3	L	E	A	I	L	<u>L</u>	L	V	L	I	18
6	I	L	L	L	V	<u>L</u>	I	F	L	R	18
8	L	L	V	L	I	<u>F</u>	L	R	Q	R	16
9	L	V	L	I	F	<u>L</u>	R	Q	R	I	16
7	L	L	L	V	L	<u>I</u>	F	L	R	Q	15
10	V	L	I	F	L	<u>R</u>	Q	R	I	R	12

V6-HLA-A0201-10mers-24P4C12											
Each peptide is a portion of SEQ											
ID NO: 13; each start position is											
specified, the length of peptide is											
10 amino acids, and the end											
position for each peptide is the											
start position plus nine.											
Pos	1	2	3	4	5	6	7	8	9	0	score
7	G	L	I	P	R	<u>S</u>	V	F	N	L	29
4	S	S	K	G	L	<u>I</u>	P	R	S	V	15

TABLE XXXV-continued

V7-HLA-A0201-10mers-24P4C12											
Each peptide is a portion of SEQ											
ID NO: 15; each start position is											
specified, the length of peptide is											
10 amino acids, and the end											
position for each peptide is the											
start position plus nine.											
Pos	1	2	3	4	5	6	7	8	9	0	score
1	Q	S	W	Y	W	<u>I</u>	L	V	A	V	4
2	S	W	Y	W	I	<u>L</u>	V	A	V	G	4
4	Y	W	I	L	V	<u>A</u>	V	G	Q	M	3
5	W	I	L	V	A	<u>V</u>	G	Q	M	M	2
6	I	L	V	A	V	<u>G</u>	Q	M	M	S	2
8	V	A	V	G	Q	<u>M</u>	M	S	T	M	2
9	A	V	G	Q	M	<u>M</u>	S	T	M	F	2

V8-HLA-A0201-10mers-24P4C12											
Each peptide is a portion of SEQ											
ID NO: 17; each start position is											
specified, the length of peptide											
is 10 amino acids, and the end											
position for each peptide is the											
start position plus nine.											
Pos	1	2	3	4	5	6	7	8	9	0	score
4	Y	W	L	P	I	<u>M</u>	R	N	P	I	15
5	W	L	P	I	M	<u>R</u>	N	P	I	T	15
18	H	V	F	Q	T	<u>S</u>	I	L	G	A	15
7	P	I	M	R	N	<u>P</u>	I	T	P	T	14
13	I	T	P	T	G	<u>H</u>	V	F	Q	T	14
8	I	M	R	N	P	<u>I</u>	T	P	T	G	13
21	Q	T	S	I	L	<u>G</u>	A	Y	V	I	13
20	F	Q	T	S	I	<u>L</u>	G	A	Y	V	12
15	P	T	G	H	V	<u>F</u>	Q	T	S	I	11
10	R	N	P	I	T	<u>P</u>	T	G	H	V	10
16	T	G	H	V	F	<u>Q</u>	T	S	I	L	10
12	P	I	T	P	T	<u>G</u>	H	V	F	Q	8

V9-HLA-A0201-10mers-24P4C12											
Each peptide is a portion of SEQ											
ID NO: 19; each start position is											
specified, the length of peptide is											
10 amino acids, and the end											
position for each peptide is the											
start position plus nine.											
Pos	1	2	3	4	5	6	7	8	9	0	score
9	Y	P	L	P	T	<u>Q</u>	P	A	T	L	20
2	Y	W	A	M	T	<u>A</u>	L	Y	P	L	19
7	A	L	Y	P	L	<u>P</u>	T	Q	P	A	19
12	P	T	Q	P	A	<u>T</u>	L	G	Y	V	17
16	A	T	L	G	Y	<u>V</u>	L	W	A	S	15
4	A	M	T	A	L	<u>Y</u>	P	L	P	T	14
5	M	T	A	L	Y	<u>P</u>	L	P	T	Q	13
17	T	L	G	Y	V	<u>L</u>	W	A	S	N	13
13	T	Q	P	A	T	<u>L</u>	G	Y	V	L	11
18	L	G	Y	V	L	<u>W</u>	A	S	N	I	11
15	P	A	T	L	G	<u>Y</u>	V	L	W	A	9

V1-HLA-A0203-10mers-24P4C12											
Each peptide is a portion of SEQ ID											
NO: 3; each start position is											
specified, the length of peptide is 10											
amino acids, and the end position											
for each peptide is the start position											
plus nine.											
Pos	1	2	3	4	5	6	7	8	9	0	score
303	Q	<u>S</u>	V	Q	E	T	<u>W</u>	L	A	A	19
168	P	<u>S</u>	F	L	L	P	<u>S</u>	A	P	A	18

TABLE XXXVI

TABLE XXXVI-continued

330	L	<u>R</u>	Q	R	I	R	<u>I</u>	A	I	A	18
459	A	<u>L</u>	G	Q	C	V	<u>L</u>	A	G	A	18
461	G	<u>Q</u>	C	V	L	A	<u>G</u>	A	F	A	18
304	S	<u>V</u>	Q	E	T	W	<u>L</u>	A	A	L	17
3	G	<u>K</u>	Q	R	D	E	<u>D</u>	D	E	A	10
46	L	<u>G</u>	Y	I	V	V	<u>G</u>	I	V	A	10
64	V	<u>L</u>	Y	P	R	N	<u>S</u>	T	G	A	10
95	I	<u>L</u>	S	S	N	I	<u>I</u>	S	V	A	10
166	L	<u>C</u>	P	S	F	L	<u>L</u>	P	S	A	10
182	F	<u>P</u>	W	T	N	V	<u>T</u>	P	P	A	10
205	I	<u>S</u>	G	L	I	D	<u>S</u>	L	N	A	10
217	I	<u>S</u>	V	K	I	F	<u>E</u>	D	F	A	10
226	A	<u>Q</u>	S	W	Y	W	<u>I</u>	L	V	A	10
230	Y	<u>W</u>	I	L	V	A	<u>L</u>	G	V	A	10
245	L	<u>F</u>	I	L	L	L	<u>R</u>	L	V	A	10
261	L	<u>I</u>	L	G	V	L	<u>G</u>	V	L	A	10
279	E	<u>Y</u>	R	V	L	R	<u>D</u>	K	G	A	10
292	Q	<u>L</u>	G	F	T	T	<u>N</u>	L	S	A	10
302	Y	<u>Q</u>	S	V	Q	E	<u>T</u>	W	L	A	10
308	T	<u>W</u>	L	A	A	L	<u>I</u>	V	L	A	10
312	A	<u>L</u>	I	V	L	A	<u>V</u>	L	E	A	10
328	I	<u>F</u>	L	R	Q	R	<u>I</u>	R	I	A	10
335	R	<u>I</u>	A	I	A	L	<u>L</u>	K	E	A	10
338	I	<u>A</u>	L	L	K	E	<u>A</u>	S	K	A	10
361	T	<u>F</u>	V	L	L	L	<u>I</u>	C	I	A	10
364	L	<u>L</u>	L	I	C	I	<u>A</u>	Y	W	A	10
367	I	<u>C</u>	I	A	Y	W	<u>A</u>	M	T	A	10
371	Y	<u>W</u>	A	M	T	A	<u>L</u>	Y	L	A	10
382	S	<u>G</u>	Q	P	Q	Y	<u>V</u>	L	W	A	10
403	P	<u>I</u>	N	T	S	C	<u>N</u>	P	T	A	10
450	L	<u>F</u>	W	T	L	N	<u>W</u>	V	L	A	10
457	V	<u>L</u>	A	L	G	Q	<u>C</u>	V	L	A	10
466	A	<u>G</u>	A	F	A	S	<u>F</u>	Y	W	A	10
481	D	<u>I</u>	P	T	F	P	<u>L</u>	I	S	A	10
494	T	<u>L</u>	R	Y	H	T	<u>G</u>	S	L	A	10
497	Y	<u>H</u>	T	G	S	L	<u>A</u>	F	G	A	10
506	A	<u>L</u>	I	L	T	L	<u>V</u>	Q	I	A	10
525	K	<u>L</u>	R	G	V	Q	<u>N</u>	P	V	A	10
550	K	<u>F</u>	I	K	F	L	<u>N</u>	R	N	A	10
555	L	<u>N</u>	R	N	A	Y	<u>I</u>	M	I	A	10
565	I	<u>Y</u>	G	K	N	F	<u>C</u>	V	S	A	10
568	K	<u>N</u>	F	C	V	S	<u>A</u>	K	N	A	10
639	L	<u>P</u>	I	M	T	S	<u>I</u>	L	G	A	10
643	T	<u>S</u>	I	L	G	A	<u>Y</u>	V	I	A	10
692	L	<u>K</u>	I	L	G	K	<u>K</u>	N	E	A	10
4	K	<u>Q</u>	R	D	E	D	<u>D</u>	E	A	Y	9
47	G	<u>Y</u>	I	V	V	G	<u>I</u>	V	A	W	9
65	L	<u>Y</u>	P	R	N	S	<u>T</u>	G	A	Y	9
96	L	<u>S</u>	S	N	I	I	<u>S</u>	V	A	E	9
167	C	<u>P</u>	S	F	L	L	<u>P</u>	S	A	P	9
169	S	<u>F</u>	L	L	P	S	<u>A</u>	P	A	L	9
183	P	<u>W</u>	T	N	V	T	<u>P</u>	P	A	L	9
206	S	<u>G</u>	L	I	D	S	<u>L</u>	N	A	R	9
218	S	<u>V</u>	K	I	F	E	<u>D</u>	F	A	Q	9
227	Q	<u>S</u>	W	Y	W	I	<u>L</u>	V	A	L	9
231	W	<u>I</u>	L	V	A	L	<u>G</u>	V	A	L	9
246	F	<u>I</u>	L	L	L	R	<u>L</u>	V	A	G	9
262	I	<u>L</u>	G	V	L	G	<u>V</u>	L	A	Y	9
280	Y	<u>R</u>	V	L	R	D	<u>K</u>	G	A	S	9
293	L	<u>G</u>	F	T	T	N	<u>L</u>	S	A	Y	9
309	W	<u>L</u>	A	A	L	I	<u>V</u>	L	A	V	9
313	L	<u>I</u>	V	L	A	V	<u>L</u>	E	A	I	9
329	F	<u>L</u>	R	Q	R	I	<u>R</u>	I	A	I	9
331	R	<u>A</u>	R	I	A	L	<u>A</u>	I	A	S	9
336	I	<u>Q</u>	I	A	R	L	<u>K</u>	E	A	L	9
339	A	<u>L</u>	L	K	E	A	<u>S</u>	K	A	V	9
362	F	<u>V</u>	L	L	L	I	<u>C</u>	I	A	Y	9
365	L	<u>L</u>	I	C	I	A	<u>Y</u>	W	A	M	9
368	C	<u>I</u>	A	Y	W	A	<u>M</u>	T	A	L	9
372	W	<u>A</u>	M	T	A	L	<u>Y</u>	L	A	T	9
383	G	<u>Q</u>	P	Q	Y	V	<u>L</u>	W	A	S	9
404	I	<u>N</u>	T	S	C	N	<u>P</u>	T	A	H	9
451	F	<u>W</u>	T	L	N	W	<u>V</u>	L	A	L	9
458	L	<u>A</u>	L	G	Q	C	<u>V</u>	L	A	G	9
460	L	<u>G</u>	Q	C	V	L	<u>A</u>	G	A	F	9
462	Q	<u>C</u>	V	L	A	G	<u>A</u>	F	A	S	9
467	G	<u>A</u>	F	A	S	F	<u>Y</u>	W	A	F	9
482	I	<u>P</u>	T	F	P	L	<u>I</u>	S	A	F	9
495	L	<u>R</u>	Y	H	T	G	<u>S</u>	L	A	F	9
498	H	<u>T</u>	G	S	L	A	<u>F</u>	G	A	L	9

TABLE XXXVI-continued

507	L	<u>I</u>	L	T	L	V	<u>Q</u>	I	A	R	9
526	L	<u>R</u>	G	V	Q	N	<u>P</u>	V	A	R	9
551	F	<u>I</u>	K	F	L	N	<u>R</u>	N	A	Y	9
556	N	<u>R</u>	N	A	Y	I	<u>M</u>	I	A	I	9
566	Y	<u>G</u>	K	N	F	C	<u>V</u>	S	A	K	9
569	N	<u>F</u>	C	V	S	A	<u>K</u>	N	A	F	9
640	P	<u>I</u>	M	T	S	I	<u>L</u>	G	A	Y	9
644	S	<u>I</u>	L	G	A	Y	<u>V</u>	I	A	S	9
693	K	<u>I</u>	L	G	K	K	<u>N</u>	E	A	P	9
V3-HLA-A0203-10mers-24P4C12											
Each peptide is a portion of SEQ ID											
NO: 7; each start position is specified,											
the length of peptide is 10 amino											
acids, and the end position for each											
peptide is the start position plus nine.											
Pos	1	2	3	4	5	6	7	8	9	0	score
5	F	<u>P</u>	W	T	N	I	<u>T</u>	P	P	A	10
6	P	<u>W</u>	T	N	I	T	<u>P</u>	P	A	L	9
7	W	<u>T</u>	N	I	T	P	<u>P</u>	A	L	P	8
V5-HLA-A0203-10mers-24P4C12											
Pos	1	2	3	4	5	6	7	8	9	0	score
No Results Found.											
V6-HLA-A0203-10mers-24P4C12											
Pos	1	2	3	4	5	6	7	8	9	0	score
No Results Found.											
V7-HLA-A0203-10mers-24P4C12											
Each peptide is a portion of SEQ ID											
NO: 15; each start position is specified,											
the length of peptide is 10 amino acids,											
and the end position for each peptide is											
the start position plus nine.											
Pos	1	2	3	4	5	6	7	8	9	0	score
1	Q	<u>S</u>	W	Y	W	I	<u>L</u>	V	A	V	9
2	S	<u>W</u>	Y	W	I	L	<u>V</u>	A	V	G	8
V8-HLA-A0203-10mers-24P4C12											
Each peptide is a portion of SEQ ID											
NO: 17; each start position is											
specified, the length of peptide is 10											
amino acids, and the end position for											
each peptide is the start position plus											
nine.											
Pos	1	2	3	4	5	6	7	8	9	0	score
18	H	<u>V</u>	F	Q	T	S	<u>I</u>	L	G	A	10
19	V	<u>F</u>	Q	T	S	I	<u>L</u>	G	A	Y	9
20	F	<u>Q</u>	T	S	I	L	<u>G</u>	A	Y	V	8
V9-HLA-A0203-10mers-24P4C12											
Each peptide is a portion of SEQ ID											
NO: 19; each start position is specified,											
the length of peptide is 10 amino											
acids, and the end position for each											
peptide is the start position plus nine.											
Pos	1	2	3	4	5	6	7	8	9	0	score
7	A	<u>L</u>	Y	P	L	P	<u>T</u>	Q	P	A	10
15	P	<u>A</u>	T	L	G	Y	<u>V</u>	L	W	A	10
8	L	<u>Y</u>	P	L	P	T	<u>Q</u>	P	A	T	9
16	A	<u>T</u>	L	G	Y	V	<u>L</u>	W	A	S	9
9	Y	<u>P</u>	L	P	T	Q	<u>P</u>	A	T	L	8
17	T	<u>L</u>	G	Y	V	L	<u>W</u>	A	S	N	8

255

TABLE XXXVII

V1-HLA-A3-10mers-24P4C12											
Each peptide is a portion of SEQ ID											
NO: 3; each start position is specified,											
the length of peptide is 10 amino											
acids, and the end position for each											
peptide is the start position plus nine.											
Pos	1	2	3	4	5	6	7	8	9	0	score
333	R	I	R	I	A	I	A	L	L	K	32
211	S	L	N	A	R	D	I	S	V	K	30
337	A	I	A	L	L	K	E	A	S	K	28
516	R	V	I	L	E	Y	I	D	H	K	28
281	R	V	L	R	D	K	G	A	S	I	27
680	S	L	D	R	P	Y	Y	M	S	K	27
464	V	L	A	G	A	F	A	S	F	Y	25
584	N	I	V	R	V	V	V	L	D	K	24
621	R	I	P	G	L	G	K	D	F	K	24
49	I	V	V	G	I	V	A	W	L	Y	23
463	C	V	L	A	G	A	F	A	S	F	23
233	L	V	A	L	G	V	A	L	V	L	22
262	I	L	G	V	L	G	V	L	A	Y	22
376	A	L	Y	L	A	T	S	G	Q	P	22
443	Q	I	Y	G	V	L	G	L	F	W	22
525	K	L	R	G	V	Q	N	P	V	A	22
587	R	V	V	V	L	D	K	V	T	D	22
603	K	L	L	V	V	G	G	V	G	V	22
56	W	L	Y	G	D	P	R	Q	V	L	21
63	Q	V	L	Y	P	R	N	S	T	G	21
177	A	L	G	R	C	F	P	W	T	N	21
564	A	I	Y	G	K	N	E	C	V	S	21
606	V	V	G	G	V	G	V	L	S	F	21
39	V	L	F	L	L	F	I	L	G	Y	20
53	I	V	A	W	L	Y	G	D	P	R	20
171	L	L	P	S	A	P	A	L	G	R	20
251	R	L	V	A	G	P	L	V	L	V	20
252	L	V	A	G	P	L	V	L	V	L	20
282	V	L	R	D	K	G	A	S	I	S	20
362	F	V	L	L	L	I	C	I	A	Y	20
378	Y	L	A	T	S	G	Q	P	Q	Y	20
544	C	L	W	C	L	E	K	F	I	K	20
650	V	I	A	S	G	E	F	S	V	F	20
95	I	L	S	S	N	I	I	S	V	A	19
170	F	L	L	P	S	A	P	A	L	G	19
191	A	L	P	G	I	T	N	D	T	T	19
237	G	V	A	L	V	L	S	L	L	F	19
248	L	L	L	R	L	V	A	G	P	L	19
260	V	L	I	L	G	V	L	G	V	L	19
261	L	I	L	G	V	L	G	V	L	A	19
298	N	L	S	A	Y	Q	S	V	Q	E	19
312	A	L	I	V	L	A	V	L	E	A	19
314	I	V	L	A	V	L	E	A	I	L	19
317	A	V	L	E	A	I	L	L	L	M	19
322	I	L	L	L	M	L	I	F	L	R	19
340	L	L	K	E	A	S	K	A	V	G	19
347	A	V	G	Q	M	M	S	T	M	F	19
494	T	L	R	Y	H	T	G	S	L	A	19
605	L	V	V	G	G	V	G	V	L	S	19
618	F	S	G	R	I	P	G	L	G	K	19
645	I	L	G	A	Y	V	I	A	S	G	19
673	D	L	E	R	N	N	G	S	L	D	19
6	R	D	E	D	D	E	A	Y	G	K	18
64	V	L	Y	P	R	N	S	T	G	A	18
134	Q	T	V	G	E	V	F	Y	T	K	18
231	W	I	L	V	A	L	G	V	A	L	18
235	A	L	G	V	A	L	V	L	S	L	18
247	I	L	L	L	R	L	V	A	G	P	18
258	V	L	V	L	I	L	G	V	L	G	18
324	L	L	M	L	I	F	L	R	Q	R	18
456	W	V	L	A	L	G	Q	C	V	L	18
532	P	V	A	R	C	I	M	C	C	F	18
72	G	A	Y	C	G	M	G	E	N	K	17
86	L	L	Y	F	N	I	E	S	C	I	17
161	S	L	Q	Q	E	L	C	P	S	F	17
207	G	L	I	D	S	L	N	A	R	D	17
220	K	I	F	E	D	F	A	Q	S	W	17
232	I	L	V	A	L	G	V	A	L	V	17
249	L	L	R	L	V	A	G	P	L	V	17
257	L	V	L	V	L	I	L	G	V	L	17
264	G	V	L	G	V	L	A	Y	G	I	17

256

TABLE XXXVII-continued

265	V	L	G	V	L	A	Y	G	I	Y	17
292	Q	L	G	F	T	T	N	L	S	A	17
309	W	L	A	A	L	I	V	L	A	V	17
326	M	L	I	F	L	R	Q	R	I	R	17
364	L	L	L	I	C	I	A	Y	W	A	17
388	V	L	W	A	S	N	I	S	S	P	17
392	S	N	I	A	S	P	G	C	E	K	17
486	P	L	I	S	A	F	I	R	T	L	17
506	A	L	I	L	T	L	V	Q	I	A	17
551	F	I	K	F	L	N	R	N	A	Y	17
580	L	L	M	R	N	I	V	R	V	V	17
598	L	L	F	F	G	K	L	L	V	V	17
612	V	L	S	F	F	F	F	S	G	R	17
624	G	L	G	K	D	E	K	S	P	H	17
649	Y	V	I	A	S	G	F	F	S	V	17
657	S	V	F	G	M	C	V	D	T	L	17
667	F	L	C	F	L	E	D	L	E	R	17
684	P	Y	Y	M	S	K	S	L	L	K	17
689	K	S	L	L	K	I	L	G	K	K	17
9	D	D	E	A	Y	G	K	P	V	K	16
44	F	I	L	G	Y	I	V	V	G	I	16
126	T	V	G	K	N	E	F	S	Q	T	16
165	E	L	C	P	S	F	L	L	P	S	16
243	S	L	L	F	I	L	L	L	R	L	16
246	F	I	L	L	L	R	L	V	A	G	16
259	L	V	L	I	L	G	V	L	G	V	16
272	G	I	Y	Y	C	W	E	E	Y	R	16
304	S	V	Q	E	T	W	L	A	A	L	16
318	V	L	E	A	I	L	L	L	M	L	16
339	A	L	L	K	E	A	S	K	A	V	16
363	V	L	L	L	I	C	I	A	Y	W	16
453	T	L	N	W	V	L	A	L	G	Q	16
457	V	L	A	L	G	Q	C	V	L	A	16
459	A	L	G	Q	C	V	L	A	G	A	16
487	L	I	S	A	F	I	R	T	L	R	16
508	I	L	T	L	V	Q	I	A	R	V	16
518	I	L	E	Y	I	D	H	K	L	R	16
559	A	Y	I	M	I	A	I	Y	G	K	16
566	Y	G	K	N	F	C	V	S	A	K	16
571	C	V	S	A	K	N	A	F	M	L	16
579	M	L	L	M	R	N	I	V	R	V	16
596	D	L	L	L	F	E	G	K	L	L	16
640	P	I	M	T	S	I	L	G	A	Y	16
690	S	L	L	K	I	L	G	K	K	N	16
693	K	I	L	G	K	K	N	E	A	P	16
35	V	I	C	C	V	L	F	L	L	F	15
41	F	L	L	F	I	L	G	Y	I	V	15
42	L	L	F	I	L	G	Y	I	V	V	15
107	G	L	Q	C	P	T	P	Q	V	C	15
120	C	P	E	D	P	W	T	V	G	K	15
180	R	C	F	P	W	T	N	V	T	P	15
323	L	L	L	M	L	I	F	L	R	Q	15
329	F	L	R	Q	R	I	R	I	A	I	15
367	I	C	I	A	Y	W	A	M	T	A	15
369	I	A	Y	W	A	M	T	A	L	Y	15
423	M	C	V	F	Q	G	Y	S	S	K	15
446	G	V	L	G	L	F	W	T	L	N	15
491	F	I	R	T	L	R	Y	H	T	G	15
507	L	I	L	T	L	V	Q	I	A	R	15
510	T	L	V	Q	I	A	R	V	I	L	15
585	I	V	R	V	V	V	L	D	K	V	15
597	L	L	L	F	F	G	K	L	L	V	15
604	L	L	V	V	G	G	V	G	V	L	15
688	S	K	S	L	L	K	I	L	G	K	15
694	I	L	G	K	K	N	E	A	P	P	15
697	K	K	N	E	A	P	P	D	N	K	15
698	K	N	E	A	P	P	D	N	K	K	15
V3-HLA-A3-10mers-24P4C12											
Each peptide is a portion of SEQ ID											
NO: 7; each start position is specified,											
the length of peptide is 10 amino											
acids, and the end position for each											
peptide is the start position plus nine.											
Pos	1	2	3	4	5	6	7	8	9	0	score
3	R	C	F	P	W	T	N	I	T	P	11
9	N	I	T	P	P	A	L	P	G	I	11
8	T	N	I	T	P	P	A	L	P	G	9

257

TABLE XXXVII-continued

10	I	T	<u>P</u>	P	A	<u>L</u>	<u>P</u>	G	I	T	7
7	W	T	<u>N</u>	I	T	<u>P</u>	<u>P</u>	A	L	P	5

V5-HLA-A3-10mers-24P4C12

Each peptide is a portion of SEQ ID
NO: 11; each start position is specified,
the length of peptide is 10 amino acids,
and the end position for each peptide is
the start position plus nine.

Pos	1	2	3	4	5	6	7	8	9	0	score
1	A	V	<u>L</u>	E	A	<u>I</u>	<u>L</u>	L	L	V	19
2	V	L	<u>E</u>	A	I	<u>L</u>	<u>L</u>	L	V	L	19
6	I	L	<u>L</u>	L	V	<u>L</u>	<u>I</u>	F	L	R	19
8	L	L	<u>V</u>	L	I	<u>F</u>	<u>L</u>	R	Q	R	18
10	V	L	<u>I</u>	F	L	<u>R</u>	<u>Q</u>	R	I	R	17
7	L	L	<u>L</u>	V	L	<u>I</u>	<u>F</u>	L	R	Q	15
5	A	I	<u>L</u>	L	L	<u>V</u>	<u>L</u>	I	F	L	14
9	L	V	<u>L</u>	I	F	<u>L</u>	<u>R</u>	Q	R	I	14
4	E	A	<u>I</u>	L	L	<u>L</u>	<u>V</u>	L	I	F	11

V6-HLA-A3-10mers-24P4C12

Each peptide is a portion of SEQ ID
NO: 13; each start position is specified,
the length of peptide is 10 amino acids,
and the end position for each
peptide is the start position plus nine.

Pos	1	2	3	4	5	6	7	8	9	0	score
7	G	L	<u>I</u>	P	R	<u>S</u>	<u>V</u>	F	N	L	16
5	S	K	<u>G</u>	L	I	<u>P</u>	<u>R</u>	S	V	F	14
1	Q	G	<u>Y</u>	S	S	<u>K</u>	<u>G</u>	L	I	P	12
8	L	I	<u>P</u>	R	S	<u>V</u>	<u>F</u>	N	L	Q	11
9	I	P	<u>R</u>	S	V	<u>F</u>	<u>N</u>	L	Q	I	11
6	K	G	<u>L</u>	I	P	<u>R</u>	<u>S</u>	V	F	N	10
4	S	S	<u>K</u>	G	L	<u>I</u>	<u>P</u>	R	S	V	7

V7-HLA-A3-10mers-24P4C12

Each peptide is a portion of SEQ ID
NO: 15; each start position is specified,
the length of peptide is 10 amino acids,
and the end position for each
peptide is the start position plus nine.

Pos	1	2	3	4	5	6	7	8	9	0	score
9	A	V	<u>G</u>	Q	M	<u>M</u>	<u>S</u>	T	M	F	19
6	I	L	<u>V</u>	A	V	<u>G</u>	<u>Q</u>	M	M	S	16
5	W	I	<u>L</u>	V	A	<u>V</u>	<u>G</u>	Q	M	M	14
7	L	V	<u>A</u>	V	G	<u>Q</u>	<u>M</u>	M	S	T	14
2	S	W	<u>Y</u>	W	I	<u>L</u>	<u>V</u>	A	V	G	12
8	V	A	<u>V</u>	G	Q	<u>M</u>	<u>M</u>	S	T	M	9

V8-HLA-A3-10mers-24P4C12

Each peptide is a portion of SEQ ID
NO: 17; each start position is
specified, the length of peptide is 10
amino acids, and the end position for
each peptide is the start position plus
nine.

Pos	1	2	3	4	5	6	7	8	9	0	score
12	P	I	<u>T</u>	P	T	<u>G</u>	<u>H</u>	V	F	Q	15
11	N	P	<u>I</u>	T	P	<u>T</u>	<u>G</u>	H	V	F	14
18	H	V	<u>F</u>	Q	T	<u>S</u>	<u>I</u>	L	G	A	13
7	P	I	<u>M</u>	R	N	<u>P</u>	<u>I</u>	T	P	T	12
5	W	L	<u>P</u>	I	M	<u>R</u>	<u>N</u>	P	I	T	11
1	L	N	<u>Y</u>	Y	W	<u>L</u>	<u>P</u>	I	M	R	10
8	I	M	<u>R</u>	N	P	<u>I</u>	<u>T</u>	P	T	G	10
21	Q	T	<u>S</u>	I	L	<u>G</u>	<u>A</u>	Y	V	I	10
9	M	R	<u>N</u>	P	I	<u>T</u>	<u>P</u>	T	G	H	9
6	L	P	<u>I</u>	M	R	<u>N</u>	<u>P</u>	I	T	P	8
19	V	F	<u>Q</u>	T	S	<u>I</u>	<u>L</u>	G	A	Y	8

258

TABLE XXXVII-continued

V9-HLA-A3-10mers-24P4C12											
Each peptide is a portion of SEQ ID											
NO: 19; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.											
Pos	1	2	3	4	5	6	7	8	9	0	score
7	A	L	<u>Y</u>	P	L	<u>P</u>	<u>T</u>	Q	P	A	20
17	T	L	<u>G</u>	Y	V	<u>L</u>	<u>W</u>	A	S	N	15
10	P	L	<u>P</u>	T	Q	<u>P</u>	<u>A</u>	T	L	G	14
9	Y	P	<u>L</u>	P	T	<u>Q</u>	<u>P</u>	A	T	L	13
1	A	Y	<u>W</u>	A	M	<u>T</u>	<u>A</u>	L	Y	P	11
18	L	G	<u>Y</u>	V	L	<u>W</u>	<u>A</u>	S	N	I	10
4	A	M	<u>T</u>	A	L	<u>Y</u>	<u>P</u>	L	P	T	9
11	L	P	<u>T</u>	Q	P	<u>A</u>	<u>T</u>	L	G	Y	9
13	T	Q	<u>P</u>	A	T	<u>L</u>	<u>G</u>	Y	V	L	9

TABLE XXXVIII											
V1-HLA-A26-10mers-24P4C12											
Each peptide is a portion of SEQ ID											
NO: 3; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.											
Pos	1	2	3	4	5	6	7	8	9	0	score
34	D	V	I	C	C	V	L	F	L	L	34
138	E	V	F	Y	T	K	N	R	N	F	32
307	E	T	W	L	A	A	L	I	V	L	31
657	S	V	F	G	M	C	V	D	T	L	28
199	T	T	I	Q	Q	G	I	S	G	L	26
304	S	V	Q	E	T	W	L	A	A	L	26
588	V	V	V	L	D	K	V	T	D	L	26
592	D	K	V	T	D	L	L	L	F	F	25
49	I	V	V	G	I	V	A	W	L	Y	24
606	V	V	G	G	V	G	V	L	S	F	24
157	T	V	I	T	S	L	Q	Q	E	L	23
252	L	V	A	G	P	L	V	L	V	L	23
257	L	V	L	V	L	I	L	G	V	L	23
320	E	A	I	L	L	L	M	L	I	F	23
628	D	F	K	S	P	H	L	N	Y	Y	23
79	E	N	K	D	K	P	Y	L	L	Y	22
353	S	T	M	F	Y	P	L	V	T	F	22
362	F	V	L	L	L	I	C	I	A	Y	22
662	C	V	D	T	L	F	L	C	F	L	22
672	E	D	L	E	R	N	N	G	S	L	22
48	Y	I	V	V	G	I	V	A	W	L	20
198	D	T	T	I	Q	Q	G	I	S	G	20
216	D	I	S	V	K	I	F	E	D	F	20
240	L	V	L	S	L	L	F	I	L	L	20
293	L	G	F	T	T	N	L	S	A	Y	20
640	P	I	M	T	S	I	L	G	A	Y	20
10	D	E	A	Y	G	K	P	V	K	Y	19
39	V	L	F	L	L	F	I	L	G	Y	19
131	E	F	S	Q	T	V	G	E	V	F	19
233	L	V	A	L	G	V	A	L	V	L	19
237	G	V	A	L	V	L	S	L	L	F	19
347	A	V	G	Q	M	M	S	T	M	F	19
438	S	V	F	N	L	Q	I	Y	G	V	19
463	C	V	L	A	G	A	F	A	S	F	19
498	H	T	G	S	L	A	F	G	A	L	19
512	V	Q	I	A	R	V	I	L	E	Y	19
520	E	Y	I	D	H	K	L	R	G	V	19
571	C	V	S	A	K	N	A	F	M	L	19
589	V	V	L	D	K	V	T	D	L	L	19
33	T	D	V	I	C	C	V	L	F	L	18
203	Q	G	I	S	G	L	I	D	S	L	18
314	I	V	L	A	V	L	E	A	I	L	18
456	W	V	L	A	L	G	Q	C	V	L	18
481	D	I	P	T	F	P	L	I	S	A	18
486	P	L	I	S	A	F	I	R	T	L	18
493	R	T	L	R	Y	H	T	G	S	L	18

Figure 2H. The cDNA (SEQ ID. NO. : 16) and amino acid sequence (SEQ ID. NO. : 17) of 24P4C12 v.8. The start methionine is underlined. The open reading frame extends from nucleic acid 6-2174 including the stop codon.

Figure 2H-1

```
1      M G G K Q R D E D D E A Y G K P V K Y
1 gagccATGGGGGAAAGCAGCGGGACGAGGATGACGAGGCCTACGGGAAGCCAGTCAAAT
20    D P S F R G P I K N R S C T D V I C C V
61    ACGACCCCTCCTTTTCGAGGCCCCATCAAGAACAGAAGCTGCACAGATGTCATCTGCTGCG
40    L F L L F I L G Y I V V G I V A W L Y G
121   TCCTCTTCCTGCTCTTCATTCTAGGTTACATCGTGGTGGGATTGTGGCCTGGTTGTATG
60    D P R Q V L Y P R N S T G A Y C G M G E
181   GAGACCCCCGGCAAGTCCTCTACCCAGGAACCTCTACTGGGGCCTACTGTGGCATGGGGG
80    N K D K P Y L L Y F N I F S C I L S S N
241   AGAACAAAGATAAGCCGTATCTCCTGTACTTCAACATCTTCAGCTGCATCCTGTCCAGCA
100   I I S V A E N G L Q C P T P Q V C V S S
301   ACATCATCTCAGTTGCTGAGAACGGCCTACAGTGCCCCACACCCAGGTGTGTGTGTCTCT
120   C P E D P W T V G K N E F S Q I V G E V
361   CCTGCCCGGAGGACCCATGGACTGTGGGAAAAACGAGTTCTCACAGACTGTGGGGAAG
140   F Y T K N R N F C L P G V P W N M T V I
421   TCTTCTATACAAAAACAGGAACTTTTGTCTGCCAGGGGTACCCTGGAATATGACGGTGA
160   T S L Q Q E L C P S F L L P S A P A L G
481   TCACAAGCCTGCAACAGGAACCTCTGCCCCAGTTTCCTCCTCCCTCTGCTCCAGCTCTGG
180   R C F P W T N V T P P A L P G I T N D T
541   GCGCTGCTTTCCATGGACCAACGTTACTCCACCGCGCTCCAGGGATCACCAATGACA
200   T I Q Q G I S G L I D S L N A R D I S V
601   CCACCATACAGCAGGGGATCAGCGGTCTTATTGACAGCCTCAATGCCCGAGACATCAGTG
220   K I F E D F A Q S W Y W I L V A L G V A
661   TTAAGATCTTTGAAGATTTTGCCAGTCCTGGTATTGGATTCTTGTGGCCCTGGGGGTGG
240   L V L S L L F I L L L R L V A G P L V L
721   CTCGTGCTTGAGCCTACTGTTTATCTTGCTTCTGCGCCTGGTGGCTGGGCCCTGGTGC
260   V L I L G V L G V L A Y G I Y Y C W E E
781   TGGTGTGATCCTGGGAGTGTCTGGCGTGCTGGCATAACGGCATCTACTACTGCTGGGAGG
280   Y R V L R D K G A S I S Q L G F T T N L
841   AGTACCGAGTGCTGCGGGACAAGGGCGCCTCCATCTCCAGCTGGGTTTACCACCAACC
300   S A Y Q S V Q E T W L A A L I V L A V L
901   TCAGTGCCTACCAGAGCGTGCAGGAGACCTGGCTGGCCGCCCTGATCGTGTGGCGGTGC
320   E A I L L L M L I F L R Q R I R I A I A
961   TTGAAGCCATCCTGCTGCTGATGCTCATCTTCCTGCGGCAGCGGATTCTGATTGCCATCG
340   L L K E A S K A V G Q M M S T M F Y P L
1021  CCCTCTGAAGGAGGCCAGCAAGGCTGTGGGACAGATGATGTCTACCATGTTCTACCCAC
360   V T F V L L L I C I A Y W A M T A L Y L
```

259

TABLE XXXVIII-continued

502	L	A	F	G	A	L	I	L	T	L	18
516	R	V	I	L	E	Y	I	D	H	K	18
532	P	V	A	R	C	I	M	C	C	F	18
549	E	K	F	I	K	F	L	N	R	N	18
609	G	V	G	V	L	S	F	F	F	F	18
99	N	I	I	S	V	A	E	N	G	L	17
102	S	V	A	E	N	G	L	Q	C	P	17
156	M	T	V	I	T	S	L	Q	Q	E	17
236	L	G	V	A	L	V	L	S	L	L	17
260	V	L	I	L	G	V	L	G	V	L	17
316	L	A	V	L	E	A	I	L	L	L	17
317	A	V	L	E	A	I	L	L	L	M	17
321	A	I	L	L	M	L	I	F	L	L	17
360	V	T	F	V	L	L	L	I	C	I	17
442	L	Q	I	Y	G	V	L	G	L	F	17
596	D	L	L	L	F	F	G	K	L	L	17
604	L	L	V	V	G	G	V	G	V	L	17
616	F	F	F	S	G	R	I	P	G	L	17
664	D	T	L	F	L	C	F	L	E	D	17
665	T	L	F	L	C	F	L	E	D	L	17
682	D	R	P	Y	Y	M	S	K	S	L	17
32	C	T	D	V	I	C	C	V	L	F	16
37	C	C	V	L	F	L	L	F	I	L	16
123	D	P	W	T	V	G	K	N	E	F	16
165	E	L	C	P	S	F	L	L	P	S	16
186	N	V	T	P	P	A	L	P	G	I	16
224	D	F	A	Q	S	W	Y	W	I	L	16
239	A	L	V	L	S	L	L	F	I	L	16
262	I	L	G	V	L	G	V	L	A	Y	16
266	L	G	V	L	A	Y	G	I	Y	Y	16
332	Q	R	I	R	I	A	I	A	L	L	16
359	L	V	T	F	V	L	L	L	I	C	16
380	A	T	S	G	Q	P	Q	Y	V	L	16
400	E	K	V	P	I	N	T	S	C	N	16
405	N	T	S	C	N	P	T	A	H	L	16
424	C	V	F	Q	G	Y	S	S	K	G	16
433	G	L	I	Q	R	S	V	F	N	L	16
539	C	C	F	K	C	C	L	W	C	L	16
593	K	V	T	D	L	L	L	F	F	G	16
V3-HLA-A26-10mers-24P4C12											
Each peptide is a portion of SEQ ID NO: 7; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.											
Pos	1	2	3	4	5	6	7	8	9	0	score
6	P	W	T	N	I	T	P	P	A	L	10
9	N	I	T	P	P	A	L	P	G	I	10
10	I	T	P	P	A	L	P	G	I	T	10
7	W	T	N	I	T	P	P	A	L	P	8
3	R	C	F	P	W	T	N	I	T	P	7
8	T	N	I	T	P	P	A	L	P	G	6
4	C	F	P	W	T	N	I	T	P	P	4
V5-HLA-A26-10mers-24P4C12											
Each peptide is a portion of SEQ ID NO: 11; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.											
Pos	1	2	3	4	5	6	7	8	9	0	score
4	E	A	I	L	L	L	V	L	I	F	27
1	A	V	L	E	A	I	L	L	L	V	17
5	A	I	L	L	L	V	L	I	F	L	17
2	V	L	E	A	I	L	L	L	V	L	13

260

TABLE XXXVIII-continued

V6-HLA-A26-10mers-24P4C12											
Each peptide is a portion of SEQ ID NO: 13; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.											
Pos	1	2	3	4	5	6	7	8	9	0	score
7	G	L	I	P	R	S	V	F	N	L	17
10	P	R	S	V	F	N	L	Q	I	Y	14
5	S	K	G	L	I	P	R	S	V	F	10
V7-HLA-A26-10mers-24P4C12											
Each peptide is a portion of SEQ ID NO: 15; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.											
Pos	1	2	3	4	5	6	7	8	9	0	score
9	A	V	G	Q	M	M	S	T	M	F	19
7	L	V	A	V	G	Q	M	M	S	T	11
4	Y	W	I	L	V	A	V	G	Q	M	10
V8-HLA-A26-10mers-24P4C12											
Each peptide is a portion of SEQ ID NO: 17; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.											
Pos	1	2	3	4	5	6	7	8	9	0	score
18	H	V	F	Q	T	S	I	L	G	A	19
19	V	F	Q	T	S	I	L	G	A	Y	16
11	N	P	I	T	P	T	G	H	V	F	13
13	I	T	P	T	G	H	V	F	Q	T	13
16	T	G	H	V	F	Q	T	S	I	L	10
15	P	T	G	H	V	F	Q	T	S	I	9
V9-HLA-A26-10mers-24P4C12											
Each peptide is a portion of SEQ ID NO: 19; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.											
Pos	1	2	3	4	5	6	7	8	9	0	score
12	P	T	Q	P	A	T	L	G	Y	V	14
5	M	T	A	L	Y	P	L	P	T	Q	13
16	A	T	L	G	Y	V	L	W	A	S	13
2	Y	W	A	M	T	A	L	Y	P	L	12
11	L	P	T	Q	P	A	T	L	G	Y	12
9	Y	P	L	P	T	Q	P	A	T	L	10
13	T	Q	P	A	T	L	G	Y	V	L	10
15	P	A	T	L	G	Y	V	L	W	A	6

261

TABLE XXXIX

V1-HLA-B0702- 10mers-24P4C12											
Each peptide is a portion of SEQ ID NO: 3; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.											
Pos	1	2	3	4	5	6	7	8	9	0	score
357	Y	P	L	V	T	F	V	L	L	L	23
478	K	P	Q	D	I	P	T	F	P	L	23
683	R	P	Y	Y	M	S	K	S	L	L	21
182	F	P	W	T	N	V	T	P	P	A	19
83	K	P	Y	L	L	Y	F	N	I	F	18
192	L	P	G	I	T	N	D	T	T	I	18
482	I	P	T	F	P	L	I	S	A	F	18
639	L	P	I	M	T	S	I	L	G	A	18
149	L	P	G	V	P	W	N	M	T	V	17
252	L	V	A	G	P	L	V	L	V	L	17
380	A	T	S	G	Q	P	Q	Y	V	L	17
402	V	P	I	N	T	S	C	N	P	T	17
485	F	P	L	I	S	A	F	I	R	T	17
123	D	P	W	T	V	G	K	N	E	F	16
235	A	L	G	V	A	L	V	L	S	L	16
254	A	G	P	L	V	L	V	L	I	L	15
370	A	Y	W	A	M	T	A	L	Y	L	15
659	F	G	M	C	V	D	T	L	F	L	15
33	T	D	V	I	C	C	V	L	F	L	14
56	W	L	Y	G	D	P	R	Q	V	L	14
175	A	P	A	L	G	R	C	F	P	W	14
233	L	V	A	L	G	V	A	L	V	L	14
241	V	L	S	L	G	F	I	L	L	L	14
331	R	Q	R	I	R	I	A	I	A	L	14
405	N	T	S	C	N	P	T	A	H	L	14
451	F	W	T	L	N	W	V	L	A	L	14
502	L	A	F	G	A	L	I	L	T	L	14
582	M	R	N	I	V	R	V	V	V	L	14
590	V	L	D	K	V	T	D	L	L	L	14
15	K	P	V	K	Y	D	P	S	F	R	13
60	D	P	R	Q	V	L	Y	P	R	N	13
66	Y	P	R	N	S	T	G	A	Y	C	13
110	C	P	T	P	Q	V	C	V	S	S	13
120	C	P	E	D	P	W	T	V	G	K	13
167	C	P	S	F	L	L	P	S	A	P	13
172	L	P	S	A	P	A	L	G	R	C	13
226	A	Q	S	W	Y	W	I	L	V	A	13
227	Q	S	W	Y	W	I	L	V	A	L	13
231	W	I	L	V	A	L	G	V	A	L	13
250	L	R	L	V	A	G	P	L	V	L	13
284	R	D	K	G	A	S	I	S	Q	L	13
290	I	S	Q	L	G	F	T	T	N	L	13
301	A	Y	Q	S	V	Q	E	T	W	L	13
310	L	A	A	L	I	V	L	A	V	L	13
314	I	V	L	A	V	L	E	A	I	L	13
318	V	L	E	A	I	L	L	L	M	L	13
321	A	I	L	L	L	M	L	I	F	L	13
350	Q	M	M	S	T	M	F	Y	P	L	13
355	M	F	Y	P	L	V	T	F	V	L	13
356	F	Y	P	L	V	T	F	V	L	L	13
368	C	I	A	Y	W	A	M	T	A	L	13
396	S	P	G	C	E	K	V	P	I	N	13
441	N	T	Q	I	Y	G	V	L	G	L	13
498	H	T	G	S	L	A	F	G	A	L	13
500	G	S	L	A	F	G	A	L	I	L	13
510	T	L	V	Q	I	A	R	V	I	L	13
525	K	L	R	G	V	Q	N	P	V	A	13
571	C	V	S	A	K	N	A	F	M	L	13
572	V	S	A	K	N	A	F	M	L	L	13
657	S	V	F	G	M	C	V	D	T	L	13
686	Y	M	S	K	S	L	L	K	I	L	13
20	D	P	S	F	R	G	P	I	K	N	12
48	Y	I	V	V	G	I	V	A	W	L	12
169	S	F	L	L	P	S	A	P	A	L	12
183	P	W	T	N	V	T	P	P	A	L	12
189	P	P	A	L	P	G	I	T	N	D	12
239	A	L	V	L	S	L	L	F	I	L	12
243	S	L	L	F	I	L	L	L	R	L	12
304	S	V	Q	E	T	W	L	A	A	L	12

262

TABLE XXXIX-continued

307	E	T	W	L	A	A	L	I	V	L	12
309	W	L	A	A	L	I	V	L	A	V	12
316	L	A	V	L	E	A	I	L	L	L	12
409	N	P	T	A	H	L	V	N	S	S	12
419	C	P	G	L	M	C	V	F	Q	G	12
425	V	F	Q	G	Y	S	S	K	G	L	12
456	W	V	L	A	L	G	Q	C	V	L	12
493	R	T	L	R	Y	H	T	G	S	L	12
581	L	M	R	N	I	V	R	V	V	V	12
588	V	V	V	L	D	K	V	T	D	L	12
604	L	L	V	V	G	G	V	G	V	L	12
606	V	V	G	G	V	G	V	L	S	F	12
622	I	P	G	L	G	K	D	F	K	S	12
637	Y	W	L	P	I	M	T	S	I	L	12
662	C	V	D	T	L	F	L	C	F	L	12
701	A	P	P	D	N	K	K	R	K	K	12
18	K	Y	D	P	S	F	R	G	P	I	11
25	G	P	I	K	N	R	S	C	T	D	11
31	S	C	T	D	V	I	C	C	V	L	11
44	F	I	L	G	Y	I	V	V	G	I	11
77	M	G	E	N	K	D	K	P	Y	L	11
78	G	E	N	K	D	K	P	Y	L	L	11
140	F	Y	T	K	N	R	N	F	C	L	11
152	V	P	W	N	M	T	V	I	T	S	11
153	P	W	N	M	T	V	I	T	S	L	11
162	L	Q	Q	E	L	C	P	S	F	L	11
188	T	P	P	A	L	P	G	I	T	N	11
224	D	F	A	Q	S	W	Y	W	I	L	11
236	L	G	V	A	L	V	L	S	L	L	11
240	L	V	L	S	L	L	F	I	L	L	11
248	L	L	L	R	L	V	A	G	P	L	11
257	L	V	L	V	L	I	L	G	V	L	11
260	V	L	I	L	G	V	L	G	V	L	11
274	Y	Y	C	W	E	E	Y	R	V	L	11
312	A	L	I	V	L	A	V	L	E	A	11
315	V	L	A	V	L	E	A	I	L	L	11
332	Q	R	I	R	I	A	I	A	L	L	11
384	Q	P	Q	Y	V	L	W	A	S	N	11
395	S	S	P	G	C	E	K	V	P	I	11
413	H	L	V	N	S	S	C	P	G	L	11
433	G	L	I	Q	R	S	V	F	N	L	11
435	I	Q	R	S	V	F	N	L	Q	I	11
439	V	F	N	L	Q	I	Y	G	V	L	11
445	Y	G	V	L	G	L	F	W	T	L	11
449	G	L	F	W	T	L	N	W	V	L	11
503	A	F	G	A	L	I	L	T	L	V	11
531	N	P	V	A	R	C	I	M	C	C	11
536	C	I	M	C	C	F	K	C	C	L	11
539	C	C	F	K	C	C	L	W	C	L	11
546	W	C	L	E	K	F	I	K	F	L	11
565	I	Y	G	K	N	F	C	V	S	A	11
589	V	V	L	D	K	V	T	D	L	L	11
595	T	D	L	L	L	F	F	G	K	L	11
616	F	F	F	S	G	R	I	P	G	L	11
625	L	G	K	D	F	K	S	P	H	L	11
630	K	S	P	H	L	N	Y	Y	W	L	11
672	E	D	L	E	R	N	N	G	S	L	11
V3-HLA-B0702- 10mers-24P4C12											
Each peptide is a portion of SEQ ID NO: 7; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.											
Pos	1	2	3	4	5	6	7	8	9	0	score
5	F	P	W	T	N	I	T	P	P	A	19
6	P	W	T	N	I	T	P	P	A	L	12
1	L	G	R	C	F	P	W	T	N	I	9

263

TABLE XXXIX-continued

V5-HLA-B*0702- 10mers-24P4C12 Each peptide is a portion of SEQ ID NO: 11; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.											
Pos	1	2	3	4	5	6	7	8	9	0	score
2	V	L	E	A	I	L	L	L	V	L	14
5	A	I	L	L	L	V	L	I	F	L	13
1	A	V	L	E	A	I	L	L	L	V	10
4	E	A	I	L	L	L	V	L	I	F	10
3	L	E	A	I	L	L	L	V	L	I	9
9	L	V	L	I	F	L	R	Q	R	I	7
V6-HLA-B*0702- 10mers-24P4C12 Each peptide is a portion of SEQ ID NO: 13; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.											
Pos	1	2	3	4	5	6	7	8	9	0	score
9	I	P	R	S	V	F	N	L	Q	I	21
7	G	L	I	P	R	S	V	F	N	L	12
V7-HLA-B*0702- 10mers-24P4C12 Each peptide is a portion of SEQ ID NO: 15; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.											
Pos	1	2	3	4	5	6	7	8	9	0	score
9	A	V	G	Q	M	M	S	T	M	F	10
1	Q	S	W	Y	W	I	L	V	A	V	9
8	V	A	V	G	Q	M	M	S	T	M	8
4	Y	W	I	L	V	A	V	G	Q	M	7
1	L	V	A	V	G	Q	M	M	S	T	7
5	W	I	L	V	A	V	G	Q	M	M	6
V8-HLA-B*0702- 10mers-24P4C12 Each peptide is a portion of SEQ ID NO: 17; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.											
Pos	1	2	3	4	5	6	7	8	9	0	score
11	N	P	I	T	P	T	G	H	V	F	17
14	T	P	T	G	H	V	F	Q	T	S	13
16	T	G	H	V	F	Q	T	S	I	L	11
6	L	P	I	M	R	N	P	I	T	P	10
4	Y	W	L	P	I	M	R	N	P	I	9
7	P	I	M	R	N	P	I	T	P	T	9
21	Q	T	S	I	L	G	A	Y	V	I	9
10	R	N	P	I	T	P	T	G	H	V	8
13	I	T	P	T	G	H	V	F	Q	T	8
15	P	T	G	H	V	F	Q	T	S	I	8
18	H	V	F	Q	T	S	I	L	G	A	8

264

TABLE XXXIX-continued

5	V9-HLA-B*0702-10mers-24P4C12 Each peptide is a portion of SEQ ID NO: 19; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.												
10	Pos	1	2	3	4	5	6	7	8	9	0	score	
	9	Y	P	L	P	T	Q	P	A	T	L	22	
	11	L	P	T	Q	P	A	T	L	G	Y	13	
	14	Q	P	A	T	L	G	Y	V	L	W	13	
	2	Y	W	A	M	T	A	L	Y	P	L	12	
15	4	A	M	T	A	L	Y	P	L	P	T	12	
	13	T	Q	P	A	T	L	G	Y	V	L	12	
	7	A	L	Y	P	L	P	T	Q	P	A	11	
20	TableXL-V1-HLA-B*08-10mers-24P4C12												
	Pos	1234567890									score		
25	NoResultsFound.												
	TableXL-V3-HLA-B*08-10mers-24P4C12												
	Pos	1234567890									score		
30	NoResultsFound.												
	TableXL-V5-HLA-B*08-10mers-24P4C12												
	Pos	1234567890									score		
35	NoResultsFound.												
	TableXL-V6-HLA-B*08-10mers-24P4C12												
	Pos	1234567890									score		
40	NoResultsFound.												
	TableXL-V7-HLA-B*08-10mers-24P4C12												
	Pos	1234567890									score		
45	NoResultsFound.												
	TableXL-V8-HLA-B*08-10mers-24P4C12												
	Pos	1234567890									score		
50	NoResultsFound.												
	TableXL-V9-HLA-B*08-10mers-24P4C12												
	Pos	1234567890									score		
55	NoResultsFound.												
60	TableXLI-V1-HLA-B*1510-10mers-24P4C12												
	Pos	1234567890									score		
	NoResultsFound.												
	TableXLI-V3-HLA-B*1510-10mers-24P4C12												
65	Pos	1234567890									score		

265

-continued

NoResultsFound.			
TableXLI-V5-HLA-B1510-10mers-24P4C12			5
Pos	1234567890	score	
NoResultsFound.			
TableXLI-V6-HLA-B1510-10mers-24P4C12			10
Pos	1234567890	score	
NoResultsFound.			
TableXLI-V7-HLA-B1510-10mers-24P4C12			15
Pos	1234567890	score	
NoResultsFound.			
TableXLI-V8-HLA-B1510-10mers-24P4C12			20
Pos	1234567890	score	
NoResultsFound.			
TableXLI-V9-HLA-B1510-10mers-24P4C12			25
Pos	1234567890	score	
NoResultsFound.			
			30
TableXLII-V1-HLA-B2705-10mers-24P4C12			
Pos	1234567890	score	35
NoResultsFound.			
TableXLII-V3-HLA-B2705-10mers-24P4C12			
Pos	1234567890	score	40
NoResultsFound.			
TableXLII-V5-HLA-B2705-10mers-24P4C12			
Pos	1234567890	score	45
NoResultsFound.			
TableXLII-V6-HLA-B2705-10mers-24P4C12			
Pos	1234567890	score	50
NoResultsFound.			
TableXLII-V7-HLA-B2705-10mers-24P4C12			
Pos	1234567890	score	55
NoResultsFound.			
TableXLII-V8-HLA-B2705-10mers-24P4C12			
Pos	1234567890	score	60
NoResultsFound.			
TableXLII-V9-HLA-B2705-10mers-24P4C12			
Pos	1234567890	score	65
NoResultsFound.			

266

TableXLIII-V1-HLA-B2709-10mers-24P4C12		
Pos	1234567890	score
NoResultsFound.		
TableXLIII-V5-HLA-B2709-10mers-24P4C12		
Pos	1234567890	score
NoResultsFound.		
TableXLIII-V6-HLA-B2709-10mers-24P4C12		
Pos	1234567890	score
NoResultsFound.		
TableXLIII-V7-HLA-B2709-10mers-24P4C12		
Pos	1234567890	score
NoResultsFound.		
TableXLIII-V8-HLA-B2709-10mers-24P4C12		
Pos	1234567890	score
NoResultsFound.		
TableXLIII-V9-HLA-B2709-10mers-24P4C12		
Pos	1234567890	score
NoResultsFound.		

TABLE XLIV

V1-HLA-B4402-10mers-24P4C12												
Each peptide is a portion of SEQ ID NO: 3; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.												
Pos	1	2	3	4	5	6	7	8	9	0	score	
10	D	E	A	Y	G	K	P	V	K	Y	23	
78	G	E	N	K	D	K	P	Y	L	L	22	
222	F	E	D	F	A	Q	S	W	Y	W	21	
319	L	E	A	I	L	L	L	M	L	I	20	
47	G	Y	I	V	V	G	I	V	A	W	19	
332	Q	R	I	R	I	A	I	A	L	L	18	
486	P	L	I	S	A	F	I	R	T	L	18	
502	L	A	F	G	A	L	I	L	T	L	18	
620	G	R	I	P	G	L	G	K	D	F	18	
39	V	L	F	L	L	F	I	L	G	Y	17	
241	V	L	S	L	L	F	I	L	L	L	17	
254	A	G	P	L	V	L	V	L	I	L	17	
320	E	A	I	L	L	L	M	L	I	F	17	
321	A	I	L	L	L	M	L	I	F	L	17	
476	F	H	K	P	Q	D	I	P	T	F	17	
512	V	Q	I	A	R	V	I	L	E	Y	17	
699	N	E	A	P	P	D	N	K	K	R	17	
121	P	E	D	P	W	T	V	G	K	N	16	
169	S	F	L	L	P	S	A	P	A	L	16	
199	T	T	I	Q	Q	G	I	S	G	L	16	
203	Q	G	I	S	G	L	I	D	S	L	16	
260	V	L	I	L	G	V	L	G	V	L	16	
293	L	G	F	T	T	N	L	S	A	Y	16	
307	E	T	W	L	A	A	L	I	V	L	16	
316	L	A	V	L	E	A	I	L	L	L	16	
380	A	T	S	G	Q	P	Q	Y	V	L	16	
546	W	C	L	E	K	F	I	K	F	L	16	
657	S	V	F	G	M	C	V	D	T	L	16	

267

TABLE XLIV-continued

34	D	V	I	C	C	V	L	F	L	L	15
65	L	Y	P	R	N	S	T	G	A	Y	15
79	E	N	K	D	K	P	Y	L	L	Y	15
99	N	I	I	S	V	A	E	N	G	L	15
104	A	E	N	G	L	Q	C	P	T	P	15
138	E	V	F	Y	T	K	N	R	N	F	15
213	N	A	R	D	I	S	V	K	I	F	15
235	A	L	G	V	A	L	V	L	S	L	15
239	A	L	V	L	S	L	L	F	I	L	15
278	E	E	Y	R	V	L	R	D	K	G	15
284	R	D	K	G	A	S	I	S	Q	L	15
353	S	T	M	F	Y	P	L	V	T	F	15
355	M	F	Y	P	L	V	T	F	V	L	15
356	F	Y	P	L	V	T	F	V	L	L	15
362	F	V	L	L	L	I	C	I	A	Y	15
363	V	L	L	L	I	C	I	A	Y	W	15
370	A	Y	W	A	M	T	A	L	Y	L	15
417	S	S	C	P	G	L	M	C	V	F	15
442	L	Q	I	Y	G	V	L	G	L	F	15
451	F	W	T	L	N	W	V	L	A	L	15
482	I	P	T	F	P	L	I	S	A	F	15
561	I	M	I	A	I	Y	G	K	N	F	15
596	D	L	L	L	F	F	G	K	L	L	15
616	F	F	F	S	G	R	I	P	G	L	15
637	Y	W	L	P	I	M	T	S	I	L	15
640	P	I	M	T	S	I	L	G	A	Y	15
4	K	Q	R	D	E	D	D	E	A	Y	14
18	K	Y	D	P	S	F	R	G	P	I	14
80	N	K	D	K	P	Y	L	L	Y	F	14
83	K	P	Y	L	L	Y	F	N	I	F	14
130	N	E	F	S	Q	T	V	G	E	V	14
131	E	F	S	Q	T	V	G	E	V	F	14
157	T	V	I	T	S	L	Q	Q	E	L	14
164	Q	E	L	C	P	S	F	L	L	P	14
173	P	S	A	P	A	L	G	R	C	F	14
175	A	P	A	L	G	R	C	F	P	W	14
183	P	W	T	N	V	T	P	P	A	L	14
220	K	I	F	E	D	F	A	Q	S	W	14
227	Q	S	W	Y	W	I	L	V	A	L	14
231	W	I	L	V	A	L	G	V	A	L	14
233	L	V	A	L	G	V	A	L	V	L	14
240	L	V	L	S	L	L	F	I	L	L	14
243	S	L	L	F	I	L	L	L	R	L	14
250	L	R	L	V	A	G	P	L	V	L	14
252	L	V	A	G	P	L	V	L	V	L	14
253	V	A	G	P	L	V	L	V	L	I	14
262	I	L	G	V	L	G	V	L	A	Y	14
304	S	V	Q	E	T	W	L	A	A	L	14
331	R	Q	R	I	R	I	A	I	A	L	14
357	Y	P	L	V	T	F	V	L	L	L	14
431	S	K	G	L	I	Q	R	S	V	F	14
433	G	L	I	Q	R	S	V	F	N	L	14
467	G	A	F	A	S	F	Y	W	A	F	14
542	K	C	C	L	W	C	L	E	K	F	14
545	L	W	C	L	E	K	F	I	K	F	14
551	F	I	K	F	L	N	R	N	A	Y	14
569	N	F	C	V	S	A	K	N	A	F	14
589	V	V	L	D	K	V	T	D	L	L	14
595	T	D	L	L	L	F	F	G	K	L	14
627	K	D	F	K	S	P	H	L	N	Y	14
629	F	K	S	P	H	L	N	Y	Y	W	14
665	T	L	F	L	C	F	L	E	D	L	14
686	Y	M	S	K	S	L	L	K	I	L	14
7	D	E	D	D	E	A	Y	C	G	K	13
31	S	C	T	D	V	I	C	C	V	L	13
32	C	T	D	V	I	C	C	V	L	F	13
35	V	I	C	C	V	L	F	L	L	F	13
49	I	V	V	G	I	V	A	W	L	Y	13
56	W	L	Y	G	D	P	R	Q	V	L	13
57	L	Y	G	D	P	R	Q	V	L	Y	13
87	L	Y	F	N	I	F	S	C	I	L	13
145	R	N	F	C	T	P	G	V	P	W	13
153	P	W	N	M	T	V	I	T	S	L	13
186	N	V	T	P	P	A	L	P	G	I	13
237	G	V	A	L	V	L	S	L	L	F	13
248	L	L	L	R	L	V	A	G	P	L	13
257	L	V	L	V	L	I	L	G	V	L	13
271	Y	G	I	Y	Y	C	W	E	E	Y	13
301	A	Y	Q	S	V	Q	E	T	W	L	13
310	L	A	A	L	I	V	L	A	V	L	13

268

TABLE XLIV-continued

315	V	L	A	V	L	E	A	I	L	L	13
327	L	I	F	L	R	Q	R	I	R	I	13
342	K	E	A	S	K	A	V	G	Q	M	13
347	A	V	G	Q	M	M	S	T	M	F	13
405	N	T	S	C	N	P	T	A	H	L	13
425	V	F	Q	G	Y	S	S	K	G	L	13
441	N	L	Q	I	Y	G	V	L	G	L	13
445	Y	G	V	L	G	L	F	W	T	L	13
447	V	L	G	L	F	W	T	L	N	W	13
449	G	L	F	W	T	L	N	W	V	L	13
460	L	G	Q	C	V	L	A	G	A	F	13
478	K	P	Q	D	I	P	T	F	P	L	13
483	P	T	F	P	L	I	S	A	F	I	13
493	R	T	L	R	Y	H	T	G	S	L	13
495	L	R	Y	H	T	G	S	L	A	F	13
498	H	T	G	S	L	A	F	G	A	L	13
500	G	S	L	A	F	G	A	L	I	L	13
517	V	I	L	E	Y	I	D	H	K	L	13
539	C	C	F	K	C	C	L	W	C	L	13
557	R	N	A	Y	I	M	I	A	I	Y	13
582	M	R	N	I	V	R	V	V	V	L	13
590	V	L	D	K	V	T	D	L	L	L	13
591	L	D	K	V	T	D	L	L	L	F	13
592	D	K	V	T	D	L	L	L	F	F	13
606	V	V	G	G	V	G	V	L	S	F	13
659	F	G	M	C	V	D	T	L	F	L	13
661	M	C	V	D	T	L	F	L	C	F	13
662	C	V	D	T	L	F	L	C	F	L	13
671	L	E	D	L	E	R	N	N	G	S	13
672	E	D	L	E	R	N	N	G	S	L	13
682	D	R	P	Y	Y	M	S	K	S	L	13
33	T	D	V	I	C	C	V	L	F	L	12
37	C	C	V	L	F	L	L	F	I	L	12
44	F	I	L	G	Y	I	V	V	G	I	12
76	G	M	G	E	N	K	D	K	P	Y	12
123	D	P	W	T	V	G	K	N	E	F	12
132	F	S	Q	T	V	G	E	V	F	Y	12
150	P	G	V	P	W	N	M	T	V	I	12
163	Q	Q	E	L	C	P	S	F	L	L	12
216	D	I	S	V	K	I	F	E	D	F	12
223	E	D	F	A	Q	S	W	Y	W	I	12
236	L	G	V	A	L	V	L	S	L	L	12
266	L	G	V	L	A	Y	G	I	Y	Y	12
274	Y	Y	C	W	E	E	Y	R	V	L	12
277	W	E	E	Y	R	V	L	R	D	K	12
286	K	G	A	S	I	S	Q	L	G	F	12
290	I	S	Q	L	G	F	T	T	N	L	12
300	S	A	Y	Q	S	V	Q	E	T	W	12
306	Q	E	T	W	L	A	A	L	I	V	12
313	L	I	V	L	A	V	L	E	A	I	12
318	V	L	E	A	I	L	L	L	M	L	12
329	F	L	R	Q	R	I	R	I	A	I	12
350	Q	M	M	S	T	M	F	Y	P	L	12
358	P	L	V	T	F	V	L	L	L	I	12
360	V	T	F	V	L	L	L	I	C	I	12
368	C	I	A	Y	W	A	M	T	A	L	12
369	I	A	Y	W	A	M	T	A	L	Y	12
378	Y	L	A	T	S	G	Q	P	Q	Y	12
381	T	S	G	Q	P	Q	Y	V	L	W	12
395	S	S	P	G	C	E	K	V	P	I	12
420	P	G	L	M	C	V	F	Q	G	Y	12
436	Q	R	S	V	F	N	L	Q	I	Y	12
439	V	F	N	L	Q	I	Y	G	V	L	12
443	Q	I	Y	G	V	L	G	L	F	W	12
456	W	V	L	A	L	G	Q	C	V	L	12
463	C	V	L	A	G	A	F	A	S	F	12
464	V	L	A	G	A	F	A	S	F	Y	12
488	I	S	A	F	I	R	T	L	R	Y	12
505	G	A	L	I	L	T	L	V	Q	I	12
509	L	T	L	V	Q	I	A	R	V	I	12
510	T	L	V	Q	I	A	R	V	I	L	12
548	L	E	K	F	I	K	F	L	N	R	12
556	N	R	N	A	Y	I	M	I	A	I	12
571	C	V	S	A	K	N	A	F	M	L	12
572	V	S	A	K	N	A	F	M	L	L	12
576	N	A	F	M	L	L	M	R	N	I	12
588	V	V	V	L	D	K	V	T	D	L	12
604	L	L	V	V	G	V	G	V	L	L	12
628	D	F	K	S	P	H	L	N	Y	Y	12
630	K	S	P	H	L	N	Y	Y	W	L	12

269

TABLE XLIV-continued

650	V	I	A	S	G	F	F	S	V	F	12
674	L	E	R	N	N	G	S	L	D	R	12
676	R	N	N	G	S	L	D	R	P	Y	12
677	N	N	G	S	L	D	R	P	Y	Y	12
685	Y	Y	M	S	K	S	L	L	K	I	12
14	G	K	P	V	K	Y	D	P	S	F	11
27	I	K	N	R	S	C	T	D	V	I	11
40	L	F	L	L	F	I	L	G	Y	I	11
48	Y	I	V	V	G	I	V	A	W	L	11
77	M	G	E	N	K	D	K	P	Y	L	11
116	C	V	S	S	C	P	E	D	P	W	11
137	G	E	V	F	Y	T	K	N	R	N	11
161	S	L	Q	Q	E	L	C	P	S	F	11
162	L	Q	Q	E	L	C	P	S	F	L	11
208	L	I	D	S	L	N	A	R	D	I	11
212	L	N	A	R	D	I	S	V	K	I	11
221	I	F	E	D	F	A	Q	S	W	Y	11
238	V	A	L	V	L	S	L	L	F	I	11
264	G	V	L	G	V	L	A	Y	G	I	11
305	V	Q	E	T	W	L	A	A	L	I	11
314	I	V	L	A	V	L	E	A	I	L	11
348	V	G	Q	M	M	S	T	M	F	Y	11
413	H	L	V	N	S	S	C	P	G	L	11
479	P	Q	D	I	P	T	F	P	L	I	11
499	T	G	S	L	A	F	G	A	L	I	11
519	L	E	Y	I	D	H	K	L	R	G	11
528	G	V	Q	N	P	V	A	R	C	I	11
532	P	V	A	R	C	I	M	C	C	F	11
536	C	I	M	C	C	F	K	C	C	L	11
537	I	M	C	C	F	K	C	C	L	W	11
543	C	C	L	W	C	L	E	K	F	I	11
552	I	K	F	L	N	R	N	A	Y	I	11
607	V	G	G	V	G	V	L	S	F	F	11
608	G	G	V	G	V	L	S	F	F	F	11
609	G	V	G	V	L	S	F	F	F	F	11
625	L	G	K	D	F	K	S	P	H	L	11
632	P	H	L	N	Y	Y	W	L	P	I	11
642	M	T	S	I	L	G	A	Y	V	I	11
646	L	G	A	Y	V	I	A	S	G	F	11
658	V	F	G	M	C	V	D	T	L	F	11
683	R	P	Y	Y	M	S	K	S	L	L	11

V3-HLA-B4402-10mers-24P4C12

Each peptide is a portion of SEQ ID NO: 7; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.

Pos	1	2	3	4	5	6	7	8	9	0	score
6	P	W	T	N	I	T	P	P	A	L	14
9	N	I	T	P	P	A	L	P	G	I	13
1	L	G	R	C	F	P	W	T	N	I	8
3	R	C	F	P	W	T	N	I	T	P	7
8	T	N	I	T	P	P	A	L	P	G	6

V5-HLA-B4402-10mers-24P4C12

Each peptide is a portion of SEQ ID NO: 11; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.

Pos	1	2	3	4	5	6	7	8	9	0	score
3	L	E	A	I	L	L	L	V	L	I	21
4	E	A	I	L	L	L	V	L	I	F	18
5	A	I	L	E	L	V	L	I	F	L	17
2	V	L	E	A	I	L	L	V	L	L	13
9	L	V	L	I	F	L	R	Q	R	I	10

270

TABLE XLIV-continued

V6-HLA-B4402- 10mers-24P4C12											
Each peptide is a portion of SEQ ID NO: 13; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.											
Pos	1	2	3	4	5	6	7	8	9	0	score
7	G	L	I	P	R	S	V	F	N	L	17
5	S	K	G	L	I	P	R	S	V	F	14
10	P	R	S	V	F	N	L	Q	I	Y	12
9	I	P	R	S	V	F	N	L	Q	I	10
V7-HLA-B4402- 10mers-24P4C12											
Each peptide is a portion of SEQ ID NO: 15; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.											
Pos	1	2	3	4	5	6	7	8	9	0	score
9	A	V	G	Q	M	M	S	T	M	F	13
4	Y	W	I	L	V	A	V	G	Q	M	6
V8-HLA-B4402- 10mers-24P4C12											
Each peptide is a portion of SEQ ID NO: 17; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.											
Pos	1	2	3	4	5	6	7	8	9	0	score
11	N	P	I	T	P	T	G	H	V	F	17
4	Y	W	L	P	I	M	R	N	P	I	14
19	V	F	Q	T	S	I	L	G	A	Y	14
16	T	G	H	V	F	Q	T	S	I	L	11
21	Q	T	S	I	L	G	A	Y	V	I	11
15	P	T	G	H	V	F	Q	T	S	I	8
V9-HLA-B4402- 10mers-24P4C12											
Each peptide is a portion of SEQ ID NO: 19; each start position is specified, the length of peptide is 10 amino acids, and the end position for each peptide is the start position plus nine.											
Pos	1	2	3	4	5	6	7	8	9	0	score
9	Y	P	L	P	T	Q	P	A	T	L	16
14	Q	P	A	T	L	G	Y	V	L	W	13
11	L	P	T	Q	P	A	T	L	G	Y	12
13	T	Q	P	A	T	L	G	Y	V	L	12
2	Y	W	A	M	T	A	L	Y	P	L	11
18	L	G	Y	V	L	W	A	S	N	I	9
16	A	T	L	G	Y	V	L	W	A	S	8
7	A	L	Y	P	L	P	T	Q	P	A	7
TableXLV-V1-HLA-B5101-10mers-24P4C12											
Pos	1234567890										score
NoResultsFound.											

271

-continued

TableXLV-V3-HLA-B5101-10mers-24P4C12		
Pos	1234567890	score
NoResultsFound.		
TableXLV-V5-HLA-B5101-10mers-24P4C12		
Pos	1234567890	score
NoResultsFound.		
TableXLV-V6-HLA-B5101-10mers-24P4C12		
Pos	1234567890	score
NoResultsFound.		
TableXLV-V7-HLA-B5101-10mers-24P4C12		
Pos	1234567890	score
NoResultsFound.		
TableXLV-V8-HLA-B5101-10mers-24P4C12		
Pos	1234567890	score
NoResultsFound.		
TableXLV-V9-HLA-B5101-10mers-24P4C12		
Pos	1234567890	score
NoResultsFound.		

TABLE XLVI

V1-HLA-DRB1-0101-
15mers-24P4C12

Each peptide is a portion of SEQ ID
NO: 3; each start position is specified,
the length of peptide is 15 amino
acids, and the end position for each
peptide is the start position plus
fourteen.

Pos	1	2	3	4	5	6	7	8	9	0	1	2	3	4	5	score
227	Q	S	W	Y	W	I	L	V	A	L	G	V	A	L	V	39
206	S	G	L	I	D	S	L	N	A	R	D	I	S	V	K	33
247	I	L	L	R	L	V	A	G	P	L	V	L	V	L	33	
313	L	I	V	L	A	V	L	E	A	I	L	L	L	M	L	33
601	F	G	K	L	L	V	V	G	G	V	G	V	L	S	F	33
246	F	I	L	L	R	L	V	A	G	P	L	V	L	V	32	
262	I	L	G	V	L	G	V	L	A	Y	G	I	Y	Y	C	32
353	S	T	M	F	Y	P	L	V	T	F	V	L	L	L	I	32
368	C	I	A	Y	W	A	M	T	A	L	Y	L	A	T	S	32
652	A	S	G	F	F	S	V	F	G	M	C	V	D	T	L	32
39	V	L	F	L	L	F	I	L	G	Y	I	V	V	G	I	31
181	C	F	P	W	T	N	V	T	P	P	A	L	P	G	I	31
277	W	E	E	Y	R	V	L	R	D	K	G	A	S	I	S	31
559	A	Y	I	M	I	A	I	Y	G	K	N	F	C	V	S	31
639	L	P	I	M	T	S	I	L	G	A	Y	V	I	A	S	31
85	Y	L	L	Y	F	N	I	F	S	C	I	L	S	S	N	30
89	F	N	I	F	S	C	I	L	S	S	N	I	I	S	V	30
257	L	V	L	V	L	I	L	G	V	L	G	V	L	A	Y	30
259	L	V	L	I	L	G	V	L	G	V	L	A	Y	G	I	30
635	N	Y	Y	W	L	P	I	M	T	S	I	L	G	A	Y	30
646	L	G	A	Y	V	I	A	S	G	F	F	S	V	F	G	30
235	A	L	G	V	A	L	V	L	S	L	L	F	I	L	L	29
345	S	K	A	V	G	Q	M	M	S	T	M	F	Y	P	L	29
40	L	F	L	L	F	I	L	G	Y	I	V	V	G	I	V	28
242	L	S	L	L	F	I	L	L	L	R	L	V	A	G	P	28
359	L	V	T	F	V	L	L	L	I	C	I	A	Y	W	A	28
453	T	L	N	W	V	L	A	L	G	Q	C	V	L	A	G	28
612	V	L	S	F	F	F	F	S	G	R	I	P	G	L	G	28

272

TABLE XLVI-continued

5	640	P	I	M	T	S	I	L	G	A	Y	V	I	A	S	G	28
	167	C	P	S	F	L	L	P	S	A	P	A	L	G	R	C	27
	243	S	L	L	F	I	L	L	L	R	L	V	A	G	P	L	27
	280	Y	R	V	L	R	D	K	G	A	S	I	S	Q	L	G	27
	362	F	V	L	L	L	I	C	I	A	Y	W	A	M	T	A	27
10	423	M	C	V	F	Q	G	Y	S	S	K	G	L	I	Q	R	27
	501	S	L	A	F	G	A	L	I	L	T	L	V	Q	I	A	27
	575	K	N	A	F	M	L	L	M	R	N	I	V	R	V	V	27
	129	K	N	E	F	S	Q	T	V	G	E	V	F	V	T	K	26
	230	Y	W	I	L	V	A	L	G	V	A	L	V	L	S	L	26
15	254	A	G	P	L	V	L	V	L	I	L	G	V	L	G	V	26
	384	Q	P	Q	Y	V	L	W	A	S	N	I	S	S	P	G	26
	436	Q	R	S	V	F	N	L	Q	I	Y	G	V	L	G	L	26
	437	R	S	V	F	N	L	Q	I	Y	G	V	L	G	L	F	26
	448	L	G	L	F	W	T	L	N	W	V	L	A	L	G	Q	26
20	492	I	R	T	L	R	Y	H	T	G	S	L	A	F	G	A	26
	551	F	I	K	F	L	N	R	N	A	Y	I	M	I	A	I	26
	594	V	T	D	L	L	L	F	F	G	K	L	L	V	V	G	26
	633	H	L	N	Y	Y	W	L	P	I	M	T	S	I	L	G	26
	688	S	K	S	L	L	K	I	L	G	K	K	N	E	A	P	26
25	44	F	I	L	G	Y	I	V	V	G	I	V	A	W	L	Y	25
	53	I	V	A	W	L	Y	G	D	P	R	Q	V	L	Y	P	25
	62	R	Q	V	L	Y	P	R	N	S	T	G	A	Y	C	G	25
	90	N	I	F	S	C	I	L	S	S	N	I	I	S	V	A	25
	228	S	W	Y	W	I	L	V	A	L	G	V	A	L	V	L	25
30	231	W	I	L	V	A	L	G	V	A	L	V	L	S	L	L	25
	239	A	L	V	L	S	L	L	F	I	L	L	R	L	V	25	
	293	L	G	F	T	T	N	L	S	A	Y	Q	S	V	Q	E	25
	299	L	S	A	Y	Q	S	V	Q	E	T	W	L	A	A	L	25
	304	S	V	Q	E	T	W	L	A	A	L	I	V	L	A	V	25
35	319	L	E	A	I	L	L	L	M	L	I	F	L	R	Q	R	25
	326	M	L	I	F	L	R	Q	R	I	R	I	A	I	A	L	25
	337	A	I	A	L	L	K	E	A	S	K	A	V	G	Q	M	25
	354	T	M	F	Y	P	L	V	T	F	V	L	L	L	I	C	25
	371	Y	W	A	M	T	A	L	Y	L	A	T	S	G	Q	P	25
40	399	C	E	K	V	P	I	N	T	S	C	N	P	T	A	H	25
	451	F	W	T	L	N	W	V	L	A	L	G	Q	C	V	L	25
	454	L	N	W	V	L	A	L	G	Q	C	V	L	A	G	A	25
	471	S	F	Y	W	A	F	H	K	P	Q	D	I	P	T	F	25
	482	I	P	T	F	P	L	I	S	A	F	I	R	T	L	R	25
45	526	L	R	G	V	Q	N	P	V	A	R	C	I	M	C	C	25
	583	R	N	I	V	R	V	V	V	L	D	K	V	T	D	L	25
	603	K	L	L	V	V	G	G	V	G	V	L	S	F	F	F	25
	51	V	G	I	V	A	W	L	Y	G	D	P	R	Q	V	L	24
	97	S	S	N	I	I	S	V	A	E	N	G	R	Q	C	P	24
50	229	W	Y	W	I	L	V	A	L	G	V	A	L	V	L	S	24
	238	V	A	L	V	L	S	L	L	F	I	L	L	L	R	L	24
	255	G	P	L	V	L	V	L	I	L	G	V	L	G	V	L	24
	256	P	L	V	L	V	L	I	L	G	V	L	G	V	L	A	24
	279	E	Y	R	V	L	R	D	K	G	A	S	I	S	Q	L	24
55	307	E	T	W	L	A	A	L	I	V	L	A	V	L	E	A	24
	310	L	A	A	L	I	V	L	A	V	L	E	A	I	L	L	24
	383	G	Q	P	Q	Y	V	L	W	A	S	N	I	S	S	P	24
	420	P	G	L	M	C	V	F	Q	G	Y	S	S	K	G	L	24
	459	A	L	G	Q	C	V	L	A	G	A	F	A	S	F	Y	24
60	506	A	L	I	L	T	L	V	Q	I	A	R	V	I	L	E	24
	523	D	H	K	L	R	G	V	Q	N	P	V	A	R	C	I	24
	569	N	F	C	V	S	A	K	N	A	F	M	L	L	M	R	24
	579	M	L	L	M	R	N	I	V	R	V	V	V	L	D	K	24
	588	V	V	V	L	D	K	V	T	D	L	L	L	F	F	G	24
65	607	V	G	G	V	G	V	L	S	F	F	F	F	S	G	R	24
	644	S	I	L	G	A	Y	V	I	A	S	G	F	F	S	V	24
	660	G	M	C	V	D	T	L	F	L	C	F	L	E	D	L	24
	47	G	Y	I	V	V	G	I	V	A	W	L	Y	G	D	P	23
	59	G	D	P	R	Q	V	L	Y	P	R	N	S	T	G	A	23
70	165	E	L	C	P	S	F	L	L	P	S	A	P	A	L	G	23
	166	L	C	P	S	F	L	L	P	S	A	P	A	L	G	R	23
	241	V	L	S	L	L	F	I	L	L	R	L	V	A	G	23	
	374	M	T	A	L	Y	L	A	T	S	G	Q	P	Q	Y	V	23
	412	A	H	L	V	N	S	S	C	P	G	L	M	C	V	F	23
75	507	L	I	L	T	L	V	Q	I	A	R	V	I	L	E	Y	23
	508	I	L	T	L	V	Q	I	A	R	V	I	L	E	Y	I	23
	566	Y	G	K	N	F	C	V	S	A	K	N	A	F	M	L	23
	604	L	F	V	V	G	G	V	V	G	V	L	S	F	F	F	23
	636	Y	Y	W	L	P	I	M	T	S	I	L	G	A	Y	V	23
80	33	T	D	V	I	C	C	V	L	F	L	F	I	A	W	G	22
	43	L	F	I	L	G	Y	I	V	V	G	I	V	I	L	G	22
	86	L	L	Y	F	N	I	F	S	C	I	L	S	S	N	I	22
	160	T	S	L	Q	Q	E	L	C	P	S	F	L	L	P	S	22
	198	D	T	T	I	Q	Q	G	I	S	G	L	I	D	S	L	22

273

TABLE XLVI-continued

312	A	L	I	V	L	A	V	L	E	A	I	L	L	L	M	22
316	L	A	V	L	E	A	I	L	L	L	M	L	I	F	L	22
349	G	Q	M	M	S	T	M	F	Y	P	L	V	T	F	V	22
363	V	L	L	L	I	C	I	A	Y	W	A	M	T	A	L	22
419	C	P	G	L	M	C	V	F	Q	G	Y	S	S	K	G	22
439	V	F	N	L	Q	I	Y	G	V	L	G	L	F	W	T	22
441	N	L	Q	I	Y	G	V	L	G	L	F	W	T	L	N	22
458	L	A	L	G	Q	C	V	L	A	G	A	F	A	S	F	22
481	D	I	P	T	F	P	L	I	S	A	F	I	R	T	L	22
511	L	V	Q	I	A	R	V	I	L	E	Y	I	D	H	K	22
587	R	V	V	V	L	D	K	V	T	D	L	L	L	F	F	22
598	L	L	F	F	G	K	L	L	V	V	G	G	V	G	V	22
655	F	F	S	V	F	G	M	C	V	D	T	L	F	L	C	22
689	K	S	L	L	K	I	L	G	K	K	N	E	A	P	P	22
138	E	V	F	Y	T	K	N	R	N	F	C	L	P	G	V	21
151	G	V	P	W	N	M	T	V	I	T	S	L	Q	Q	E	21
153	P	W	N	M	T	V	I	T	S	L	Q	Q	E	L	C	21
203	Q	G	I	S	G	L	I	D	S	L	N	A	R	D	I	21
300	S	A	Y	Q	S	V	Q	E	T	W	L	A	A	L	I	21
329	F	L	R	Q	R	I	R	I	A	I	A	L	K	E	A	21
331	R	Q	R	I	R	I	A	I	A	L	K	E	A	S	A	21
409	N	P	T	A	H	L	V	N	S	S	C	P	G	L	M	21
518	I	L	E	Y	I	D	H	K	L	R	G	V	Q	N	P	21
548	L	E	K	F	I	K	F	L	N	R	N	A	Y	I	M	21
606	V	G	G	V	G	V	L	S	F	F	F	F	S	G	G	21
10	D	E	A	Y	G	K	P	V	K	Y	D	P	S	F	R	20
20	D	P	S	F	R	G	P	I	K	N	R	S	C	T	D	20
272	G	I	Y	Y	C	W	E	E	Y	R	V	L	R	D	K	20
333	R	I	R	I	A	I	A	L	L	K	E	A	S	K	A	20
449	G	L	F	W	T	L	N	W	V	L	A	L	G	Q	C	20
476	F	H	K	P	Q	D	I	P	T	F	P	L	I	S	A	20
543	C	C	L	W	C	L	E	K	F	I	K	F	L	N	R	20
563	I	A	I	Y	G	K	N	F	C	V	S	A	K	N	A	20
599	L	F	F	G	K	L	L	V	V	G	G	V	G	V	L	20
614	S	F	F	F	F	S	G	R	I	P	G	L	G	K	D	20
634	L	N	Y	Y	W	L	P	I	M	T	S	I	L	G	A	20
645	I	L	G	A	Y	V	I	A	S	G	F	F	S	V	F	20
656	F	S	V	F	G	M	C	V	D	T	L	F	L	C	F	20
657	S	V	F	G	M	C	V	D	T	L	F	L	C	F	L	20
37	C	C	V	L	F	L	L	F	I	L	G	Y	I	V	V	19
38	C	V	L	F	L	L	F	I	L	G	Y	I	V	V	G	19
82	D	K	P	Y	L	L	Y	F	N	I	F	S	C	I	L	19
122	E	D	P	W	T	V	G	K	N	E	F	S	Q	T	V	19
179	G	R	C	F	P	W	T	N	V	T	P	P	A	L	P	19
184	W	T	N	V	T	P	P	A	L	P	G	I	T	N	D	19
245	L	F	I	L	L	R	L	V	E	A	G	P	L	V	L	19
271	Y	G	I	Y	Y	C	W	E	E	Y	R	V	L	R	D	19
317	A	V	L	E	A	I	L	L	L	M	L	I	F	L	R	19
323	L	L	L	M	L	I	F	L	R	Q	R	I	R	I	A	19
336	I	A	I	A	L	L	K	E	A	S	K	A	V	G	Q	19
369	I	A	Y	W	A	M	T	A	L	Y	L	A	T	S	G	19
411	T	A	H	L	V	N	S	S	C	P	G	L	M	C	V	19
442	L	Q	I	Y	G	V	N	S	G	L	F	W	T	L	N	19
460	L	G	Q	C	V	L	A	G	A	F	A	S	F	Y	W	19
495	L	R	Y	H	T	G	S	L	A	F	G	A	L	I	L	19
503	A	F	G	A	L	I	L	T	L	V	Q	I	A	R	V	19
557	R	N	A	Y	I	M	I	A	I	Y	G	K	N	F	C	19
586	V	R	V	V	V	L	D	K	V	T	D	L	L	L	F	19
683	R	P	Y	Y	M	S	K	S	L	L	K	I	L	G	K	19
684	P	Y	Y	M	S	K	S	L	L	K	I	L	G	K	K	19

V3-HLA-DRB1-0101-
15mers-24P4C12

Each peptide is a portion of SEQ ID
NO: 7; each start position is specified,
the length of peptide is 15 amino acids,
and the end position for each peptide is
the start position plus fourteen.

Pos	1	2	3	4	5	6	7	8	9	0	1	2	3	4	5	score
9	C	F	P	W	T	N	I	T	P	P	A	L	P	G	I	31
7	G	R	C	F	P	W	T	N	I	T	P	P	A	L	P	19
12	W	T	N	I	T	P	P	A	L	P	G	I	T	N	D	19
10	F	P	W	T	N	I	T	P	P	A	L	P	G	I	T	18
14	N	I	T	P	P	A	L	P	G	I	T	N	D	T	T	16

274

TABLE XLVI-continued

V5-HLA-DRB1-0101- 15mers-24P4C12																	
Each peptide is a portion of SEQ ID NO: 11; each start position is specified, the length of peptide is 15 amino acids, and the end position for each peptide is the start position plus fourteen.																	
Pos	1	2	3	4	5	6	7	8	9	0	1	2	3	4	5	score	
2	L	I	V	L	A	V	L	E	A	I	L	L	L	V	L	33	
8	L	E	A	I	L	L	L	V	L	I	F	L	R	Q	R	25	
15	V	L	I	F	L	R	Q	R	I	R	I	A	I	A	L	25	
1	A	L	I	V	L	A	V	L	E	A	I	L	L	L	V	22	
5	L	A	V	L	E	A	I	L	L	V	L	I	I	F	L	22	
6	A	V	L	E	A	I	L	L	L	V	L	I	F	L	R	19	
12	L	L	L	V	L	I	F	L	R	Q	R	I	R	I	A	19	
13	L	L	V	L	I	F	L	R	Q	R	I	R	I	A	I	18	
7	V	L	E	A	I	L	L	V	L	I	F	L	R	Q	17		
11	I	L	L	L	V	L	I	F	L	R	Q	R	I	R	I	17	
14	L	V	L	I	F	L	R	Q	R	I	R	I	A	I	A	17	
4	V	L	A	V	L	E	A	I	L	L	V	L	I	F	16		
10	A	I	L	L	L	V	L	I	F	L	R	Q	R	I	R	16	

V6-HLA-DRB1-0101-
15mers-24P4C12

Each peptide is a portion of SEQ ID
NO: 13; each start position is specified,
the length of peptide is 15 amino
acids, and the end position for each
peptide is the start position plus
fourteen.

Pos	1	2	3	4	5	6	7	8	9	0	1	2	3	4	5	score
2	M	C	V	F	Q	G	Y	S	S	K	G	L	I	P	R	27
15	P	R	S	V	F	N	L	Q	I	Y	G	V	L	G	L	26
7	G	Y	S	S	K	G	L	I	P	R	S	V	F	N	L	24
4	V	F	Q	G	Y	S	S	K	G	L	I	P	R	S	V	16
10	S	K	G	L	I	P	R	S	V	F	N	L	Q	I	Y	16
12	G	L	I	P	R	S	V	F	N	L	Q	I	Y	G	V	16
1	L	M	C	V	F	Q	G	Y	S	K	G	L	I	P	15	
8	Y	S	S	K	G	L	I	P	R	S	V	F	N	L	Q	15

V7-HLA-DRB1-0101-
15mers-24P4C12

Each peptide is a portion of SEQ ID
NO: 15; each start position is specified,
the length of peptide is 15 amino acids,
and the end position for each peptide is
the start position plus fourteen.

Pos	1	2	3	4	5	6	7	8	9	0	1	2	3	4	5	score
6	Q	S	W	Y	W	I	L	V	A	V	G	Q	M	M	S	31
12	L	V	A	V	G	Q	M	M	S	T	M	F	Y	P	L	29
7	S	W	Y	W	I	L	V	A	V	G	Q	M	M	S	T	25
8	W	Y	W	I	L	V	A	V	G	Q	M	M	S	T	M	24
9	Y	W	I	L	V	A	V	G	Q	M	M	S	T	M	F	24
1	F	E	D	F	A	Q	S	W	Y	W	I	L	V	A	V	18
5	A	Q	S	W	Y	W	I	L	V	A	V	G	Q	M	M	16
11	I	L	V	A	V	G	Q	M	M	S	T	M	F	Y	P	15

V8-HLA-DRB1-0101-
15mers-24P4C12

Each peptide is a portion of SEQ ID
NO: 17; each start position is specified,
the length of peptide is 15 amino acids,
and the end position for each peptide is
the start position plus fourteen.

Pos	1	2	3	4	5	6	7	8	9	0	1	2	3	4	5	score
24	V	F	Q	T	S	I	L	G	A	Y	V	I	A	S	G	28
7	N	Y	Y	W	L	P	I	M	R	N	P	I	T	P	T	24
23	H	V	F	Q	T	S	I	L	G	A	Y	V	I	A	S	23
6	L	N	Y	Y	W	L	P	I	M	R	N	P	I	T	P	20
5	H	L	N	Y	Y	W	L	P	I	M	R	N	P	I	T	18
21	T	G	H	V	F	Q	T	S	I	L	G	A	Y	V	I	18
3	S	P	H	L	N	Y	Y	W	L	P	I	M	R	N	P	17

275

TABLE XLVI-continued

8	Y	Y	W	L	P	I	M	R	N	P	I	T	P	T	G	17
13	I	M	R	N	P	I	T	P	T	G	H	V	F	Q	T	17
11	L	P	I	M	R	N	P	I	T	P	T	G	H	V	F	16
12	P	I	M	R	N	P	I	T	P	T	G	H	V	F	Q	16
14	M	R	N	P	I	T	P	T	G	H	V	F	Q	T	S	16
26	Q	T	S	I	L	G	A	Y	V	I	A	S	G	F	F	16
9	Y	W	L	P	I	M	R	N	P	I	T	P	T	G	H	15
18	I	T	P	T	G	H	V	F	Q	T	S	I	L	G	A	15
19	T	P	T	G	H	V	F	Q	T	S	I	L	G	A	Y	14
20	P	T	G	H	V	F	Q	T	S	I	L	G	A	Y	V	14

V9-HLA-DRB1-0101-
15mers-24P4C12

Each peptide is a portion of SEQ ID
NO: 19; each start position is specified,
the length of peptide is 15 amino acids,
and the end position for each peptide is
the start position plus fourteen.

Pos	1	2	3	4	5	6	7	8	9	0	1	2	3	4	5	score
4	C	I	A	Y	W	A	M	T	A	L	Y	P	L	P	T	32
10	M	T	A	L	Y	P	L	P	T	Q	P	A	T	L	G	30
22	T	L	G	Y	V	L	W	A	S	N	I	S	S	P	G	26
21	A	T	L	G	Y	V	L	W	A	S	N	I	S	S	P	24
7	Y	W	A	M	T	A	L	Y	P	L	P	T	Q	P	A	23
13	L	Y	P	L	P	T	Q	P	A	T	L	G	Y	V	L	23
5	I	A	Y	W	A	M	T	A	L	Y	P	L	P	T	Q	19
2	L	I	C	I	A	Y	W	A	M	T	A	L	Y	P	L	17
1	L	L	I	C	I	A	Y	W	A	M	T	A	L	Y	P	16
16	L	P	T	Q	P	A	T	L	G	Y	V	L	W	A	S	16
23	L	G	Y	V	L	W	A	S	N	I	S	S	P	G	C	16
24	G	Y	V	L	W	A	S	N	I	S	S	P	G	C	E	16
9	A	M	T	A	L	Y	P	L	P	T	Q	P	A	T	L	15

TABLE XLVII

V1-HLA-DRB1-0301-
15mers-24P4C12

Each peptide is a portion of SEQ ID
NO: 3; each start position is specified, the
length of peptide is 15 amino acids, and
the end position for each peptide is the
start position plus fourteen.

Pos	1	2	3	4	5	6	7	8	9	0	1	2	3	4	5	score
54	V	A	W	L	Y	G	D	P	R	Q	V	L	Y	P	R	36
586	V	R	V	V	V	L	D	K	V	T	D	L	L	L	F	31
667	F	L	C	F	L	E	D	L	E	R	N	N	G	S	L	29
312	A	L	I	V	L	A	V	L	E	A	I	L	L	L	M	28
97	S	S	N	I	I	S	V	A	E	N	G	L	Q	C	P	27
155	N	M	T	V	I	T	S	L	Q	Q	E	L	C	P	S	27
454	L	N	W	V	L	A	L	G	Q	C	V	L	A	G	A	27
549	E	K	F	I	K	F	L	N	R	N	A	Y	I	M	I	27
136	V	G	E	V	F	Y	T	K	N	R	N	F	C	L	P	26
508	I	L	T	L	V	Q	I	A	R	V	I	L	E	Y	I	26
622	I	P	G	L	G	K	D	F	K	S	P	H	L	N	Y	26
376	A	L	Y	L	A	T	S	G	Q	P	Q	Y	V	L	W	25
447	V	L	G	L	F	W	T	L	N	W	V	L	A	L	G	25
279	E	Y	R	V	L	R	D	K	G	A	S	I	S	Q	L	24
534	A	R	C	I	M	C	F	K	G	C	C	L	W	C	L	24
567	G	K	N	F	C	V	S	A	K	N	A	F	M	L	L	24
229	W	Y	W	I	L	V	A	L	G	V	A	L	V	L	S	23
238	V	A	L	V	L	S	L	L	F	I	L	L	L	R	L	23
14	G	K	P	V	K	Y	D	P	S	F	R	G	P	I	K	22
218	S	V	K	I	F	E	D	F	A	Q	S	W	Y	W	I	22
219	V	K	I	F	E	D	F	A	Q	S	W	Y	W	I	L	22
235	A	L	G	V	A	L	V	L	S	L	L	F	I	L	L	22
241	V	L	S	L	L	F	I	L	L	L	R	L	V	A	G	22
360	V	T	F	V	L	L	L	I	C	I	A	Y	W	A	M	22
515	A	R	V	I	L	E	Y	I	D	H	K	L	R	G	V	22
594	V	T	D	L	L	L	F	F	G	K	L	L	V	V	G	22
33	T	D	V	I	C	C	V	L	F	L	L	F	I	L	G	21
167	C	P	S	F	L	L	P	S	A	P	A	L	G	R	C	21
192	L	P	G	I	T	N	D	T	T	I	Q	Q	G	I	S	21
237	G	V	A	L	V	L	S	L	L	F	I	L	L	L	R	21

276

TABLE XLVII-continued

5	239	A	L	V	L	S	L	L	F	I	L	L	L	R	L	V	21	
	260	V	L	I	L	G	V	L	G	V	L	A	Y	L	G	I	Y	21
	302	Y	Q	S	V	Q	E	T	W	L	A	A	L	L	R	I	V	21
	319	L	E	A	I	L	L	L	M	L	I	F	L	L	R	Q	R	21
	431	S	K	G	L	I	Q	R	S	V	F	N	L	Q	I	Y	21	
10	461	G	Q	C	V	L	A	G	A	F	A	S	F	Y	W	A	21	
	587	R	V	V	V	L	D	K	V	T	D	L	L	F	F	F	21	
	590	V	L	D	K	V	T	D	L	L	L	F	F	G	K	L	21	
	595	T	D	L	L	L	F	F	G	K	L	L	V	V	G	G	21	
	658	V	F	G	M	C	V	D	T	L	F	L	C	F	L	E	21	
15	32	C	T	D	V	I	C	C	V	V	L	F	L	L	F	I	L	20
	37	C	C	V	L	F	L	L	F	I	L	G	Y	I	V	V	20	
	46	L	G	Y	I	V	V	G	I	V	A	W	L	Y	G	D	20	
	47	G	Y	I	V	V	G	I	V	A	W	L	Y	G	D	P	20	
	74	Y	C	G	M	G	E	N	K	D	K	P	Y	L	L	Y	20	
20	76	G	M	G	E	N	K	D	K	P	Y	L	L	Y	F	N	20	
	231	W	I	L	V	A	L	G	V	A	L	V	L	S	L	L	20	
	233	L	V	A	L	G	V	A	L	V	L	S	L	L	F	I	20	
	246	F	I	L	L	L	R	L	V	A	G	P	L	V	L	V	20	
	250	L	R	L	V	A	G	P	L	V	L	V	L	I	L	G	20	
25	255	G	P	L	V	L	V	L	I	L	G	V	L	G	V	L	20	
	258	V	L	V	L	I	L	G	V	L	G	V	L	A	Y	G	20	
	313	L	I	V	L	A	V	L	E	A	I	L	L	L	M	L	20	
	316	L	A	V	L	E	A	I	L	L	L	M	L	I	F	L	20	
	323	L	L	L	M	L	I	F	L	R	Q	R	I	R	I	A	20	
30	338	I	A	L	L	K	E	A	S	K	A	V	G	Q	M	M	20	
	411	T	A	H	L	V	N	S	S	C	P	G	L	M	C	V	20	
	439	V	F	N	L	Q	I	Y	G	V	L	G	L	R	F	W	20	
	484	T	F	P	L	I	S	A	F	I	R	T	L	R	Y	H	20	
	559	A	Y	I	M	I	A	I	Y	G	K	N	F	C	V	S	20	
35	588	V	V	V	L	D	K	V	T	D	L	L	L	F	F	G	20	
	602	G	K	L	L	V	V	G	G	V	G	V	L	S	F	F	20	
	604	L	L	V	V	G	G	V	G	V	L	S	F	F	F	F	20	
	691	L	L	K	I	L	G	K	K	N	E	A	P	P	D	N	20	
	156	M	T	V	I	T	S	L	Q	Q	E	L	C	P	S	F	19	
40	159	I	T	S	L	Q	Q	E	L	C	P	S	F	L	L	P	19	
	205	I	S	G	L	I	D	S	L	N	A	R	D	I	S	V	19	
	335	R	I	A	I	A	L	L	K	E	A	S	K	A	V	G	19	
	348	V	G	Q	M	M	S	T	M	F	Y	P	L	V	T	F	19	
	366	L	I	C	I	A	Y	W	A	M	T	A	L	Y	L	A	19	
45	385	P	Q	Y	V	L	W	A	S	N	I	S	S	P	G	C	19	
	505	G	A	L	I	L	T	L	V	Q	I	A	R	V	I	L	19	
	576	N	A	F	M	L	L	M	R	N	I	V	R	V	V	V	19	
	607	V	G	G	V	G	V	L	S	F	F	F	F	S	G	R	19	
	626	G	K	D	F	K	S	P	H	L	N	Y	Y	W	L	P	19	
50	638	W	L	P	I	M	T	S	I	L	G	A	Y	V	I	A	19	
	648	A	Y	V	I	A	S	G	F	F	S	V	F	G	M	C	19	
	663	V	D	T	L	F	L	C	F	L	E	D	L	E	R	N	19	
	668	L	C	F	L	E	D	L	E	R	N	N	G	S	L	D	19	
	684	P	Y	Y	M	S	K	S	L	L	K	I	L	G	K	K	19	
55	689	K	S	L	L	K	I	L	G	K	K	N	E	A	P	P	19	
	3	G	K	Q	R	D	E	D	D	E	A	Y	G	K	P	V	18	
	61	P	R	Q	V	L	Y	P	R	N	S	T	G	A	Y	C	18	
	98	S	N	I	I	S	V	A	E	N	G	L	Q	C	P	T	18	
	114	Q	V	C	V	S	S	C	P	E	D	P	W	T	V	G	18	
60	214	A	R	D	I	S	V	K	I	F	E	D	F	A	Q	S	18	
	243	S	L	L	F	I	L	L	L	R	L	V	A	G	P	L	18	
	263	L	G	V	L	G	V	L	A	Y	G	I	Y	Y	C	W	18	
	327	L	I	F	L	R	Q	R	I	R	I	A	I	A	L	L	18	
	345	S	K	A	V	G	Q	M	M	S	T	M	F	Y	P	L	18	
65	462	Q	C	V	L	A	G	A	F	A	S	F	Y	W	A	F	18	
	530	Q	N	P	V	A	R	C	I	M	C	C	F	K	C	C	18	
	560	Y	I	M	I	A	I	Y	G	K	N	F	C	V	S	A	18	
	569	N	F	C	V	S	A	K	N	A	F	M	L	L	M	R	18	
	579	M	L	L	M	R	N	I	V	R	V	V	V	L	D	K	18	
65	585	I	V	R	V	V	V	L	D	K	V	T	D	L	L	L	18	
	655	F	F	S	V	F	G	M	C	V	D	T	L	F	L	C	18	
	656	F	S	V	F	G	M	C	V	D	T	L	F	L	C	F	18	
	660	G	M	C	V	D	T	L	F	L	C	F	L	E	D	L	18	
	664	D	T	L	F	L	C	F	L	E	D	L	E	R	T	N	18	
60	284	R	D	K	G	A	S	I	S	Q	L	G	F	E	R	T	17	
	290	I	S	Q	L	G	F	T	T	N	L	S	A	Y	Q	S	17	
	324	L	L	M	L	I	F	L	R	Q	R	I	R	I	A	I	17	
	325	L	M	L	I	F	L	R	Q	R	I	R	I	A	I	A	17	
	353	S	T	M	F	Y	P	L	V	T	F	V	L	L	L	I	17	
65	423	M	C	V	F	Q	G	Y	S	S	K	G	L	I	Q	R	17	
	437	R	S	V	F	N	L	Q	I	Y	G	V	L	L	G	F	17	
	485	F	P	L	I	S	A	F	I	R	T	L	R	Y	H	T	17	
	517	V	I	L	E	Y	I	D	H	K	L	R	G	V	Q	N	17	
	519	L	E	Y	I	D	H	K	L	R	G	V	Q	N	P	V	17	

277

TABLE XLVII-continued

523	D	H	K	L	R	G	V	Q	N	P	V	A	R	C	I	17
542	K	C	C	L	W	C	L	E	K	F	I	K	F	L	N	17
545	L	W	C	L	E	K	F	I	K	F	L	N	R	N	A	17
548	L	E	K	F	I	K	F	L	N	R	N	A	Y	I	M	17
614	S	F	F	F	F	S	G	R	I	P	G	L	G	K	D	17
619	S	G	R	I	P	G	L	G	K	D	F	K	S	P	H	17
670	F	L	E	D	L	E	R	N	N	G	S	L	D	R	P	17
692	L	K	I	L	G	K	K	N	E	A	P	P	D	N	K	17

V3-HLA-DRB1-0301-15mers- 24P4C12																
Each peptide is a portion of SEQ ID NO: 7; each start position is specified, the length of peptide is 15 amino acids, and the end position for each peptide is the start position plus fourteen.																
Pos	1	2	3	4	5	6	7	8	9	0	1	2	3	4	5	score
12	W	T	N	I	T	P	P	A	L	P	G	I	T	N	D	12
3	A	P	A	L	G	R	C	F	P	W	T	N	I	T	P	10
9	C	F	P	W	T	N	I	T	P	P	A	L	P	G	I	10
7	G	R	C	F	P	W	T	N	I	T	P	P	A	L	P	8
6	L	G	R	C	F	P	W	T	N	I	T	P	P	A	L	7

V5-HLA-DRB1-0301- 15mers-24P4C12																
Each peptide is a portion of SEQ ID NO: 11; each start position is specified, the length of peptide is 15 amino acids, and the end position for each peptide is the start position plus fourteen.																
Pos	1	2	3	4	5	6	7	8	9	0	1	2	3	4	5	score
1	A	L	I	V	L	A	V	L	E	A	I	L	L	L	V	28
8	L	E	A	I	L	L	L	V	L	I	F	L	R	Q	R	21
2	L	I	V	L	A	V	L	E	A	I	L	L	L	V	L	20
5	L	A	V	L	E	A	I	L	L	L	V	L	I	F	L	20
12	L	L	L	V	L	I	F	L	R	Q	R	I	R	I	A	20
13	L	L	V	L	I	F	L	R	Q	R	I	R	I	A	I	17
14	L	V	L	I	F	L	R	Q	R	I	R	I	A	I	A	17
4	V	L	A	V	L	E	A	I	L	L	L	V	L	I	F	15
9	E	A	I	L	L	L	V	L	I	F	L	R	Q	R	I	15
10	A	I	L	L	L	V	L	I	F	L	R	Q	R	I	R	13

V6-HLA-DRB1-0301- 15mers-24P4C12																
Each peptide is a portion of SEQ ID NO: 13; each start position is specified, the length of peptide is 15 amino acids, and the end position for each peptide is the start position plus fourteen.																
Pos	1	2	3	4	5	6	7	8	9	0	1	2	3	4	5	score
10	S	K	G	L	I	P	R	S	V	F	N	L	Q	I	Y	22
2	M	C	V	F	Q	G	Y	S	S	K	G	L	I	P	R	17
8	Y	S	S	K	G	L	I	P	R	S	V	F	N	L	Q	16
11	K	G	L	I	P	R	S	V	F	N	L	Q	I	Y	G	12
1	L	M	C	V	F	Q	G	Y	S	S	K	G	L	I	P	11
15	P	R	S	V	F	N	L	Q	I	Y	G	V	L	G	L	10

V7-HLA-DRB1-0301-15mers- 24P4C12																
Each peptide is a portion of SEQ ID NO: 15; each start position is specified, the length of peptide is 15 amino acids, and the end position for each peptide is the start position plus fourteen.																
Pos	1	2	3	4	5	6	7	8	9	0	1	2	3	4	5	score
9	Y	W	I	L	V	A	V	G	Q	M	M	S	T	M	F	18
12	L	V	A	V	G	Q	M	M	S	T	M	F	Y	P	L	18
1	F	E	D	F	A	Q	S	W	Y	W	I	L	V	A	V	16
8	W	Y	W	I	L	V	A	V	G	Q	M	M	S	T	M	13
10	W	I	L	V	A	V	G	Q	M	M	S	T	M	F	Y	10
13	V	A	V	G	Q	M	M	S	T	M	F	Y	P	L	V	10

278

TABLE XLVII-continued

V8-HLA-DRB1-0301-15mers- 24P4C12																	
Each peptide is a portion of SEQ ID NO: 17; each start position is specified, the length of peptide is 15 amino acids, and the end position for each peptide is the start position plus fourteen.																	
Pos	1	2	3	4	5	6	7	8	9	0	1	2	3	4	5	score	
10	22	G	H	V	F	Q	T	S	I	L	G	A	Y	V	I	A	17
	8	Y	Y	W	L	P	I	M	R	N	P	I	T	P	T	G	16
	15	R	N	P	I	T	P	T	G	H	V	F	Q	T	S	I	14
	26	Q	T	S	I	L	G	A	Y	V	I	A	S	G	F	F	13
	21	T	G	H	V	F	Q	T	S	I	L	G	A	Y	V	I	12
15	10	W	L	P	I	M	R	N	P	I	T	P	T	G	H	V	11
	11	L	P	I	M	R	N	P	I	T	P	T	G	H	V	F	11
	3	S	P	H	L	N	Y	Y	W	L	P	I	M	R	N	P	10
	7	N	Y	Y	W	L	P	I	M	R	N	P	I	T	P	T	10
	14	M	R	N	P	I	T	P	T	G	H	V	F	Q	T	S	9
19	T	P	T	G	H	V	F	Q	T	S	I	L	G	A	Y	8	
V9-HLA-DRB1-0301-15mers- 24P4C12																	
Each peptide is a portion of SEQ ID NO: 19; each start position is specified, the length of peptide is 15 amino acids, and the end position for each peptide is the start position plus fourteen.																	
Pos	1	2	3	4	5	6	7	8	9	0	1	2	3	4	5	score	
30	2	L	I	C	I	A	Y	W	A	M	T	A	L	Y	P	L	19
	23	L	G	Y	V	L	W	A	S	N	I	S	S	P	G	C	19
	10	M	T	A	L	Y	P	L	P	T	Q	P	A	T	L	G	13
	7	Y	W	A	M	T	A	L	Y	P	L	P	T	Q	P	A	12
	12	A	L	Y	P	L	P	T	Q	P	A	T	L	G	Y	V	12
35	13	L	Y	P	L	P	T	Q	P	A	T	L	G	Y	V	L	12
	20	P	A	T	L	G	Y	V	L	W	A	S	N	I	S	S	12
	3	I	C	I	A	Y	W	A	M	T	A	L	Y	P	L	P	10
	14	Y	P	L	P	T	Q	P	A	T	L	G	Y	V	L	W	10
	24	G	Y	V	L	W	A	S	N	I	S	S	P	G	C	E	10
5	I	A	Y	W	A	M	T	A	L	Y	P	L	P	T	Q	9	
16	L	P	T	Q	P	A	T	L	G	Y	V	L	W	A	S	9	
TABLE XLVIII																	
V1-DR1-0401-15mers- 24P4C12																	
Each peptide is a portion of SEQ ID NO: 3; each start position is specified, the length of peptide is 15 amino acids, and the end position for each peptide is the start position plus fourteen.																	
Pos	1	2	3	4	5	6	7	8	9	0	1	2	3	4	5	score	
50	85	Y	L	L	Y	F	N	I	F	S	C	I	L	S	S	N	28
	89	F	N	I	F	S	C	I	L	S	S	N	I	I	S	V	28
	243	S	L	L	F	I	L	L	L	R	L	V	A	G	P	L	28
	353	S	T	M	F	Y	P	L	V	T	F	V	L	L	L	I	28
	469	F	A	S	F	Y	W	A	F	H	K	P	Q	D	I	P	28
55	548	L	E	K	F	I	K	F	L	N	R	N	A	Y	I	M	28
	575	K	N	A	F	M	L	L	M	R	N	I	V	R	V	V	28
	635	N	Y	Y	W	L	P	I	M	T	S	I	L	G	A	Y	28
	54	V	A	W	L	Y	G	D	P	R	Q	V	L	Y	P	R	26
	98	S	N	I	I	S	V	A	E	N	G	L	Q	C	P	T	26
60	153	P	W	N	M	T	V	I	T	S	L	Q	Q	E	L	C	26
	189	P	P	A	L	P	G	I	T	N	D	T	T	I	Q	Q	26
	192	L	P	G	I	T	N	D	T	T	I	Q	Q	G	I	S	26
	323	L	L	L	M	L	I	F	L	R	Q	R	I	R	I	A	26
	337	A	I	A	L	L	K	E	A	S	K	A	V	G	Q	M	26
65	385	P	Q	Y	V	L	W	A	S	N	I	S	S	P	G	C	26
	419	C	P	G	L	M	C	V	F	Q	G	Y	S	S	K	G	26
	454	L	N	W	V	L	A	L	G	Q	C	V	L	A	G	A	26
	508	I	L	T	L	V	Q	I	A	R	V	I	L	E	Y	I	26

- Klein, Immunology: the Science of Self-Nonself Discrimination, John Wiley & Sons, New York, (1982) p. 355.
- Klein et al., Nature Med. (1997) 3:402-408.
- Kohler and Milstein, Nature (1975) 256:495-497.
- Lewin et al., Genes VI, Oxford University Press, Inc., New York, (1997) Chapter 29.
- Mallampalli et al., Biochem. J. (1996) 318:333-341.
- McClellan and Hill, Eur. J. Cancer (1993) 29A:2243-2248.
- Morrison et al., PNAS USA (1984) 81:6851-6855.
- Morton and Myszka, Methods in Enzymology (1998) 295:268.
- Muller et al., MCB (1991) 11:1785.
- Muller-Pillasch et al., Gene (1998) 208:25-30.
- NCI-CGAP, "National Cancer Institute, Cancer Genome Anatomy Project (CGAP)", Oct. 7, 1997, EMBL Database Entry AA612666, Accession No. AA612666, XP002144282.
- Pascalis et al., Journal of Immunology (2002) 169:3076-3084.
- Paul, W.E., ed. Fundamental Immunology, Raven Press (1984) pp. 614-619.
- Pemberton et al., J. of Histochemistry and Cytochemistry (1997) 45:1697-1706.
- Pinto et al., Clin Cancer Res (1996) 2(9):1445-1451.
- Reiter, Robert E. et al., Prostate stem cell antigen: A cell surface marker overexpressed in prostate cancer., Proc. Natl. Acad. Sci., Feb. 1998, vol. 95, pp. 1735-1740, XP-002078363.
- Rowen, L. et al., "Sequence of the human major histocompatibility complex class III region", Mar. 29, 1999, EMBL Database Entry AF134726, Accession No. AF134726, XP002144283.
- Rowen, L. et al., "Sequence of the human major histocompatibility complex class III region", Nov. 1, 1999, EMBL Database Entry Q9Y332, XP002144284.
- Rowen, L., Mar. 24, 1999, NCBI Accession No. AAD21813.
- Rudikoff et al., PNAS USA (1982) 79:1979-1983.
- Sequence Search Result (enablement), searched Jan. 24, 2010.
- Sequence Search Result (ODP-'827), searched Jan. 18, 2010.
- Sequence Search Result (ODP-'823), searched Jan. 24, 2010.
- Sequence Search Result (ODP-'138), searched Feb. 2, 2010.
- Sequence Search Result (Ashkenazi), searched Jan. 11, 2010.
- Simpson, A.J.G., Mar. 16, 2000, dbEST Id 4011587, GenBank Acc. AW579065.
- Simpson, A. J. G., Mar. 23, 2000, dbEST Id 4036649, GenBank Acc. AW603383.
- Simpson, A. J. G., Mar. 23, 2000, dbEST Id 4035408, GenBank Acc. AW602142.
- Simpson, A.J.G., Feb. 4, 2000, dbEST Id 3787048, GenBank Acc. AW393065.
- Slootstra et al., Molecular Diversity (1996) 2:156-164.
- Spitler, Cancer Biotherapy (1995) 10:1-3.
- Stites et al., Basic and Clinical Immunology, Seventh Edition (1991), Appleton and Lange, East Norwalk, CT, p. 102.
- Storrie et al., Methods Enzymol. (1990) 182:203-225.
- Strausberg, Robert, Sep. 6, 1999, dbEST Id 3075200, GenBank Acc. AI951815.
- Strausberg, Robert, Mar. 9, 2000, dbEST Id 3079479, GenBank Acc. AI956094.
- Strausberg, Robert, Jun. 21, 1999, dbEST Id 2655196, GenBank Acc. AI745450.
- Strausberg, Robert, Mar. 7, 2000, dbEST Id 2893738, GenBank Acc. AI813886.
- Strausberg, Robert, Aug. 14, 1997, dbEST Id 1112901, GenBank Acc. AA468365.
- Strausberg, Robert, Aug. 21, 1997, dbEST Id 1178186, GenBank Acc. AA533783.
- Strausberg, Robert, Feb. 16, 1999, dbEST Id 2101871, GenBank Acc. AI318311.
- Strausberg, Robert, Oct. 30, 1999, dbEST Id 3291479, GenBank Acc. AW139432.
- Strausberg, Robert, Feb. 24, 2000, dbEST Id 3880006, GenBank Acc. AW469133.
- Strausberg, Robert, Mar. 7, 2000, dbEST Id 2947457, GenBank Acc. AI858987.
- Strausberg, Robert, May 13, 1999, dbEST Id 2376359, GenBank Acc. AI560195.
- Strausberg, Robert, Mar. 7, 2000, dbEST Id 2946846, GenBank Acc. AI858299.
- Strausberg, Robert, May 14, 1999, dbEST Id 2390443, GenBank Acc. AI572115.
- Strausberg, Robert, May 14, 1999, dbEST Id 2381301, GenBank Acc. AI565097.
- Strausberg, Robert, Dec. 14, 1999, dbEST Id 2443929, GenBank Acc. AI625125.
- Strausberg, Robert, Mar. 8, 2000, dbEST Id 3055029, GenBank Acc. AI932443.
- Su et al., PNAS USA (1996) 93:7252-7257.
- Takeda, Jun, Sep. 9, 1997, dbEST Id 1241269, GenBank Acc. C75094.
- Tockman et al., Cancer Res (1992) 52:2711s-2718s.
- Welch et al., Int. J. Cancer (1989) 43:449-457.
- Welford, Opt. Quant. Elect. (1991) 23:1.
- White et al., Ann. Rev. Med. (2001) 52:125-145.
- Wilson, R. K., Jun. 10, 1999, dbEST Id 2629269, GenBank Acc. AI721101.
- Wilson, R. K., Jul. 7, 1995, dbEST Id 285541, GenBank Acc. H25030.
- Wilson, R. K., Apr. 20, 1995, dbEST Id 194186, GenBank Acc. R24141.
- Zellner et al., Clin. Can. Res. (1998) 4:1797-1802.
- Zimmer, Cell Motility and the Cytoskeleton (1991) 20:325-337.
- International Search Report for PCT/US08/75488, mailed on Feb. 13, 2009, 5 pages.
- Written Opinion of the International Searching Authority for PCT/US08/75488, mailed on Feb. 13, 2009, 7 pages.
- Supplementary European Search Report for 02789937.6, mailed May 25, 2009, 7 pages.
- International Search Report for PCT/US10/26429, mailed on Jun. 10, 2010, 3 pages.
- Written Opinion of the International Searching Authority for PCT/US10/26429, mailed on Jun. 10, 2010, 3 pages.
- Correale et al., J. Natl. Cancer Inst. (1997) 89(4):293-300.
- McNeel et al., Cancer Res (2001) 61(13):5161-5167.
- Office Action for Canadian Patent Application 2,503,346, mailed Mar. 28, 2011, 3 pages.
- European Search Report for EP 10185550.0, mailed Jun. 15, 2012, 11 pages.
- Gress et al., "A pancreatic cancer-specific expression profile," Oncogene (1996) 13:1819-1830.
- O'Regan et al., "An electric lobe suppressor for a yeast choline transport mutation belongs to a new family of transporter-like proteins," PNAS (2000) 97(4):1835-1840.

* cited by examiner

Figure 2H-2

1081 TGGTCACCTTTGTCCTCCTCCTCATCTGCATTGCCTACTGGGCCAIGACTGCTCTGTACC
380 A T S G Q P Q Y V L W A S N I S S P G C
1141 TGGCTACATCGGGGCAACCCCAAGTATGTGCTCTGGGCATCCAACATCAGCTCCCCGGCT
400 E K V P I N T S C N F T A H L V N S S C
1201 GTGAGAAAGTGCCAATAAATACATCATGCAACCCCAAGGCCACCTTGTGAACCTCCTCGT
420 P G L M C V F Q G Y S S K G L I Q R S V
1261 GCCCAGGGCTGATGTGCGTCTTCCAGGGCTACTCAICCAAAGGCCTAATCCAACGTTCTG
440 F N L Q I Y G V L G L F W T L N W V L A
1321 TCTTCAATCTGCAAATCTATGGGGTCTGGGGCTCTTCTGGACCCCTAACTGGGTACTGG
460 L G Q C V L A G A F A S F Y W A F H K P
1381 CCCTGGGCCAATGCGTCCTCGCTGGAGCCTTTGCCTCCTTCTACTGGGCCTTCCACAAGC
480 Q D I P T F P L I S A F I R T L R Y H T
1441 CCCAGGACATCCCTACCTTCCCCTTAATCTCTGCCTTCATCCGCACACTCCGTTACCACA
500 G S L A F G A L I L T L V Q I A R V I L
1501 CTGGGTCATTGGCATTGGAGCCCTCATCCTGACCCTTGTGCAGATAGCCCGGGTCATCT
520 E Y I D H K L R G V Q N P V A R C I M C
1561 TGGAGTATATTGACCACAAGCTCAGAGGAGTGCAGAACCCCTGTAGCCCGCTGCATCATGT
540 C F R C C L W C L E K F I K F L N R N A
1621 GCTGTTTCAAGTGCTGCCTCTGGTGTCTGGAAAAATTTATCAAGTTCCTAAACGCAATG
560 Y I M I A I Y G K N F C V S A R N A F M
1681 CATACTCATGATCGCCATCTACGGGAAGAATTTCTGTGTCTCAGCCAAAAATGCGTTCA
580 L L M R N I V R V V V L D K V I D L L L
1741 TGCTACTCATGCGAAACATTGTGAGGGTGGTCTGCTCGGACAAAGTCACAGACCTGCTGC
600 F F G K L L V V G G V G V L S F F F F S
1801 TGTTCCTTTGGGAAGCTGCTGGTGGTCTGGAGGCGTGGGGTCTGTCTTCTTTTTTTTCT
620 G R I P G L G K D F R S P H L N Y Y W L
1861 CCGGTCGCATCCCGGGGCTGGGTAAAGACTTTAAGAGCCCCACCTCAACTATTACTGGC
640 P I M R N P I T P T G H V F Q T S I L G
1921 TGCCCATCATGAGGAACCCAATAACCCCAACGGGTCATGTCTTCCAGACCTCCATCCTGG
660 A Y V I A S G F F S V F G M C V D T L F
1981 GGGCCTATGTCATCGCCAGCGGCTTCTTCAGCGTTTTTCGGCATGTGTGTGGACACGCTCT
680 L C F L E D L E R N N G S L D R P Y Y M
2041 TCCTCTGCTTCTGGAAGACCTGGAGCGGAACAACGGCTCCCTGGACCGGCCCTACTACA
700 S K S L L K I L G K K N E A P P D N K K
2101 TGTCCAAGAGCCTTCTAAAGATTCTGGGCAAGAAGAACGAGGCGCCCCCGGACAACAAGA
720 R K K *
2161 AGAGGAAGAAGTGAacagctccggccctgatccaggactgcacccacccccaccgtccag
2221 ccatccaacctcacttcgccttacaggtctccatthttgtggtaaaaaaggttttaggcc
2281 aggcgcctggctcacgcctgtaatccaacactttgagaggctgaggcgggcgatcacc

TABLE XLVIII-continued

523	D	H	K	L	R	G	V	Q	N	P	V	A	R	C	I	26
579	M	L	M	R	D	P	S	I	V	R	V	V	L	D	N	26
16	P	V	K	Y	D	P	S	F	R	R	G	P	I	K	N	22
38	C	V	L	F	L	L	F	I	L	G	Y	I	V	V	G	22
82	D	K	P	Y	L	L	Y	F	N	I	F	S	C	S	I	22
86	L	L	Y	F	N	I	L	F	S	C	I	L	S	S	I	22
122	E	D	P	W	T	V	G	K	N	E	F	S	Q	T	V	22
138	E	V	F	Y	T	K	N	R	N	F	C	L	P	G	V	22
181	C	F	P	W	T	N	V	T	P	P	A	L	P	G	I	22
219	V	K	I	F	E	D	F	A	Q	S	W	Y	W	I	L	22
227	Q	S	W	Y	W	I	L	V	A	L	G	V	A	L	V	22
228	S	W	Y	W	I	L	V	A	L	G	V	A	L	V	L	22
272	G	I	Y	Y	C	W	E	R	E	Y	R	V	L	R	D	22
277	W	E	E	Y	R	V	L	R	D	K	G	A	S	I	S	22
292	Q	L	G	F	T	T	N	L	S	A	Y	Q	S	A	V	22
299	L	S	A	Y	Q	S	V	Q	E	T	W	L	A	A	L	22
306	Q	E	T	W	L	A	A	L	I	V	L	A	V	L	E	22
354	T	M	F	Y	P	L	L	T	F	V	L	L	L	I	C	22
359	L	V	T	F	V	L	L	L	I	C	I	A	S	Y	P	22
384	Q	P	Q	Y	V	L	W	A	S	N	I	S	S	P	G	22
423	M	C	V	F	Q	G	Y	S	S	K	G	L	I	Q	R	22
442	L	Q	I	Y	G	V	L	G	L	F	V	L	A	L	N	22
448	L	G	L	F	W	T	L	N	W	V	L	A	L	G	Q	22
453	T	L	N	W	V	L	A	L	L	G	Q	C	V	L	A	22
488	I	S	A	F	I	R	T	L	R	Y	H	T	G	S	L	22
501	S	L	A	F	I	G	A	L	I	L	T	L	V	Q	I	22
557	R	N	A	Y	I	M	I	A	I	Y	G	K	N	F	C	22
633	H	L	N	Y	Y	W	L	P	I	M	T	S	I	L	G	22
646	L	G	A	Y	V	I	A	S	G	F	F	S	V	F	G	22
652	A	S	G	F	F	S	V	F	L	G	M	C	V	D	T	22
667	F	L	C	F	L	E	D	L	E	R	N	N	G	S	L	22
682	D	R	P	Y	Y	M	S	K	S	L	L	K	I	L	G	22
14	G	K	P	V	K	Y	D	P	S	F	R	G	P	I	K	20
39	V	L	F	L	L	F	I	L	G	Y	I	V	V	G	I	20
40	L	F	L	L	F	I	L	G	Y	I	V	V	G	I	V	20
43	L	F	I	L	G	Y	I	V	V	G	I	V	A	W	L	20
97	S	S	N	I	I	S	V	A	E	N	G	L	Q	C	P	20
133	S	Q	T	V	G	E	V	F	Y	T	K	N	R	N	F	20
146	N	F	C	L	P	G	V	P	W	N	M	T	V	I	T	20
149	L	P	G	V	P	W	N	M	T	V	I	T	S	L	Q	20
155	N	M	T	V	I	T	S	L	Q	Q	E	L	C	P	S	20
156	M	T	V	I	T	S	L	Q	Q	E	L	C	P	S	F	20
198	D	T	T	I	Q	Q	G	I	S	G	L	I	D	S	L	20
202	Q	Q	G	I	S	G	L	I	D	S	L	N	A	R	D	20
206	S	G	L	I	D	S	L	N	E	A	R	D	I	S	V	20
216	D	I	S	V	K	I	F	E	D	F	A	Q	S	W	Y	20
229	W	Y	W	I	L	V	A	L	G	V	A	L	V	L	S	20
230	Y	W	I	L	V	A	L	G	V	A	L	V	L	S	L	20
233	L	V	A	L	G	V	A	L	V	L	S	L	L	F	I	20
235	A	L	G	V	A	L	V	L	S	L	L	F	I	L	L	20
238	V	A	L	V	L	S	L	L	F	I	L	L	L	R	L	20
239	A	L	V	L	S	L	L	F	I	L	L	L	R	L	V	20
241	V	L	S	L	L	F	I	L	L	L	R	L	V	A	G	20
242	L	S	L	L	F	I	L	L	L	R	L	V	A	G	P	20
246	F	I	L	L	L	R	L	V	A	G	P	L	V	L	V	20
247	I	L	L	L	R	L	V	A	G	P	L	V	L	V	L	20
254	A	G	P	L	V	L	V	L	I	L	G	V	L	G	V	20
255	G	P	L	V	L	V	L	I	L	G	V	L	G	V	L	20
257	L	V	L	V	L	I	L	G	V	L	G	V	L	A	Y	20
259	L	V	L	I	L	G	V	L	G	V	L	A	Y	G	I	20
262	I	L	G	V	L	R	G	V	L	A	Y	G	I	Y	Y	20
279	E	Y	R	V	L	R	D	K	G	A	S	I	S	Q	L	20
287	G	A	S	I	S	Q	L	G	F	T	T	N	L	S	A	20
290	I	S	Q	L	G	F	T	I	N	L	S	A	Y	Q	S	20
307	E	T	W	L	A	A	L	I	V	L	A	V	L	E	A	20
310	L	A	A	L	I	V	L	A	V	L	E	A	I	L	L	20
311	A	A	L	I	V	L	A	V	L	E	A	I	L	L	L	20
312	A	L	I	V	L	A	V	L	E	A	I	L	L	L	M	20
313	L	I	V	L	A	V	L	E	A	I	L	L	L	L	M	20
315	V	L	A	V	L	E	A	I	L	L	L	M	L	I	F	20
316	L	A	V	L	E	A	I	L	L	L	M	L	I	F	L	20
319	L	E	A	I	L	L	L	M	L	I	F	L	R	Q	R	20
321	A	I	L	L	L	M	L	I	F	L	R	Q	R	I	R	20
324	L	L	M	L	I	F	L	R	Q	R	I	R	I	A	I	20
331	R	Q	R	I	R	I	A	I	A	L	L	K	E	A	S	20
333	R	I	R	I	A	I	A	L	L	K	E	A	S	K	A	20
335	R	I	A	I	A	A	L	L	K	E	A	S	K	A	V	20
356	F	Y	P	L	V	T	F	V	L	L	I	C	I	A	A	20
363	V	L	L	L	I	C	I	A	Y	W	A	M	T	A	L	20
364	L	L	L	I	C	I	A	Y	W	A	M	T	A	L	Y	20

TABLE XLVIII-continued

5	371	Y	W	A	M	T	A	L	Y	L	A	T	S	G	Q	P	20	
	374	M	T	A	L	Y	L	A	T	S	G	Q	P	Q	Y	V	20	
	401	K	V	P	I	N	T	S	C	N	P	T	A	H	L	V	20	
	420	P	G	L	M	C	V	F	Q	G	Y	S	S	K	G	L	20	
	436	Q	R	S	V	F	N	L	Q	I	Y	G	V	L	G	L	20	
10	444	I	Y	G	V	L	G	L	F	W	T	L	N	W	V	L	20	
	445	Y	G	V	L	G	L	F	W	T	L	N	W	V	L	A	20	
	447	V	L	G	L	F	W	T	L	N	W	V	L	A	L	G	20	
	451	F	W	T	L	N	W	V	L	A	L	G	Q	C	V	L	20	
	479	P	Q	D	I	P	T	F	P	L	I	S	A	F	I	R	20	
15	484	T	F	P	L	I	S	A	F	I	R	T	L	R	Y	H	20	
	485	F	P	L	I	S	A	F	I	R	T	L	R	Y	H	T	20	
	505	G	A	L	I	L	T	L	V	Q	I	A	R	V	I	L	20	
	506	A	L	I	L	T	L	V	Q	I	A	R	V	I	L	E	20	
	511	L	V	Q	I	A	R	V	I	L	E	Y	I	D	H	K	20	
20	514	I	A	R	V	I	L	E	Y	I	D	H	K	L	R	G	20	
	516	R	V	I	L	E	Y	I	D	H	K	L	R	G	V	Q	20	
	542	K	C	C	L	W	C	L	E	K	F	I	K	F	L	N	20	
	545	L	W	C	L	E	K	F	I	K	F	L	N	R	N	A	20	
	549	E	K	F	I	K	F	L	N	R	N	A	Y	I	M	I	20	
25	558	N	A	Y	I	M	I	A	I	Y	G	K	N	F	C	V	20	
	582	M	R	N	I	V	R	V	V	V	L	D	K	V	T	D	20	
	583	R	N	I	V	R	V	V	V	V	L	D	K	V	T	D	20	
	586	V	R	V	V	V	L	D	K	V	T	D	L	L	L	F	20	
	588	V	V	V	L	D	K	V	T	D	L	L	L	F	F	G	20	
30	594	V	T	D	L	L	L	F	F	G	K	L	L	V	V	G	20	
	595	T	D	L	L	L	F	F	G	K	L	L	V	V	G	G	20	
	601	F	G	K	L	L	V	V	G	G	V	G	V	V	L	S	20	
	619	S	G	R	I	P	G	L	G	K	D	F	K	S	P	H	20	
	639	L	P	I	M	T	S	I	L	G	A	Y	V	I	A	S	20	
35	642	M	T	S	I	L	G	A	Y	V	I	A	S	G	F	F	20	
	660	G	M	C	V	D	T	L	F	L	C	F	L	E	D	L	20	
	668	L	C	F	L	E	D	L	E	R	N	N	G	S	L	D	20	
	688	S	K	S	L	L	K	I	L	G	K	K	N	E	A	P	20	
	90	N	I	F	S	C	I	L	S	S	N	I	I	S	V	A	18	
40	125	W	T	V	G	K	N	E	F	S	Q	T	V	G	E	V	18	
	152	V	P	W	N	M	T	V	I	T	S	L	Q	Q	E	L	18	
	166	L	C	P	S	F	L	L	P	S	A	P	A	L	G	R	18	
	195	I	T	N	D	T	T	I	Q	Q	G	I	S	G	L	I	18	
	203	Q	G	I	S	G	L	I	D	S	L	N	A	R	D	I	18	
45	210	D	S	L	N	A	R	D	I	S	V	K	I	F	E	D	18	
	289	S	I	S	Q	L	G	F	T	T	N	L	S	A	Y	Q	18	
	295	F	T	T	N	L	S	A	Y	Q	S	V	Q	E	T	W	18	
	342	K	E	A	S	K	A	V	G	Q	M	M	S	T	M	F	18	
	373	A	M	T	A	L	Y	L	A	T	S	G	Q	P	Q	Y	18	
50	398	G	C	E	K	V	P	I	N	T	S	C	N	P	T	A	18	
	428	G	Y	S	S	K	G	L	I	Q	R	S	V	I	N	L	18	
	433	G	L	I	Q	R	S	V	F	N	L	Q	I	Y	G	V	18	
	476	F	H	K	P	Q	D	I	P	T	F	P	L	I	S	A	18	
	481	D	I	P	T	F	P	L	I	S	T	F	Q	I	R	T	18	
55	502	L	A	F	G	A	L	I	L	T	L	V	Q	I	A	R	18	
	527	R	G	V	Q	N	P	V	A	R	C	I	M	C	C	F	18	
	568	K	N	F	C	V	S	A	K	N	A	F	M	L	L	M	18	
	611	G	V	L	S	F	F	F	F	S	G	R	I	P	G	L	18	
	623	P	G	L	G	K	D	F	K	S	P	H	L	N	Y	Y	18	
60	657	S	V	F	G	M	C	V	D	T	L	F	L	C	F	L	18	
	669	C	F	L	E	D	L	E	R	N	N	G	S	L	D	R	18	
	20	D	P	S	F	R	G	P	I	K	N	R	S	C	T	D	16	
	45	I	L	G	Y	I	V	V	G	I	V	A	W	L	Y	G	16	
	53	I	V	A	W	L	Y	G	D	P	R	Q	V	L	Y	P	16	
65	55	A	W	L	Y	G	D	P	R	Q	V	L	Y	P	C	R	16	
	63	Q	V	L	Y	P	R	N	S	T	G	A	Y	C	G	N	16	
	144	N	R	N	F	C	L	P	G	V	P	W	N	M	T	V	16	
	151	G	V	P	W	N	M	T	V	I	T	S	L	Q	Q	E	16	
	167	C	P	S	F	L	L	P	S	A	P	A	L	G	R	C	16	
70	222	F	E	D	F	A	Q	S	W	Y	W	I	L	V	A	L	16	
	226	A	G	S	W	Y	W	I	L	V	A	L	G	V	A	L	16	
	271	Y	G	I	Y	Y	C	W	E	E	Y	R	V	L	R	D	16	
	326	M	L	I	F	L	R	Q	R	I	R	I	A	I	A	L	16	
	368	C	I	A	Y	W	A	M	T	A	L	Y	L	A	T	S	16	
75	369	I	A	Y	W	A	M	T	A	L	Y	L	A	T	S	G	16	
	375	T	A	L	Y	L	A	T	S	G	Q	P	Q	Y	V	L	16	
	387	Y	S	V	L	W	A	S	N	I	S	S	P	G	C	E	K	16
	437	R	V	V	F	N	L	Q	I	Y	G	V	L	G	L	F	16	
	449	G	L	F	W	T	L	N	W	V	L	A	L	G	Q	C	16	
80	466	A	G	A	F	A	S	F	Y	W	A	F	H	K	P	Q	16	
	470	A	S	F	Y	W	A	F	H	K	P	Q	D	I	P	T	16	
	471	S	F	Y	W	A	F	H	K	P	Q	D	I	P	T	F	16	
	473	Y	W	A	F	H	K	P	Q	D	I	P	T	F	P	L	16	
	482	I	P	T	F	P	L	I	S	A	F	I	R	T	L	R	16	

281

TABLE XLVIII-continued

518	I	L	E	Y	I	D	H	K	L	R	G	V	Q	N	P	16
543	C	C	L	W	C	L	E	K	F	I	K	F	L	N	R	16
563	I	A	I	Y	G	K	N	F	C	V	S	A	K	N	A	16
598	L	L	F	F	G	K	L	L	V	V	G	G	V	N	G	16
612	V	L	S	F	F	F	F	S	G	R	I	P	G	L	G	16
613	L	S	F	F	F	F	S	G	R	I	P	G	L	G	K	16
614	S	F	F	F	F	F	S	G	R	I	P	G	L	G	K	16
634	L	N	Y	Y	W	L	P	I	M	T	S	I	L	G	A	16
653	S	G	F	F	S	V	F	I	G	M	C	V	D	T	L	16
664	D	T	L	F	F	L	C	F	L	E	D	L	E	R	N	16
62	R	Q	V	L	I	Y	P	R	R	N	S	T	G	A	Y	15
325	L	M	L	I	F	L	R	Q	R	I	R	I	A	I	A	15
327	L	I	F	L	R	Q	R	I	R	I	A	I	A	L	L	15
519	L	E	Y	I	D	H	K	L	R	G	V	Q	N	P	V	15
587	R	V	V	V	V	L	D	K	V	T	D	L	L	L	F	15
32	C	T	D	V	I	C	C	C	V	L	F	L	L	F	I	14
33	T	D	V	I	C	C	V	L	L	F	L	L	L	F	I	14
36	I	C	C	V	L	F	L	L	F	L	L	L	G	Y	I	14
37	C	C	V	L	F	L	L	L	F	I	L	G	Y	I	V	14
42	L	L	F	I	L	G	Y	I	V	V	G	I	V	A	W	14
46	L	G	Y	I	V	V	V	G	I	V	A	W	L	Y	G	14
47	G	Y	I	V	V	G	I	V	A	W	L	Y	G	D	P	14
48	Y	I	V	V	G	I	V	A	W	L	Y	G	D	P	R	14
51	V	G	I	V	A	W	L	Y	G	D	P	R	Q	V	L	14
61	P	R	Q	V	L	Y	P	R	N	S	T	G	A	Y	C	14
83	K	P	Y	L	L	Y	F	N	I	F	S	C	I	L	S	14
84	P	Y	L	L	Y	F	N	I	F	S	C	I	L	S	S	14
88	Y	F	N	I	F	S	C	S	I	L	S	S	N	I	I	14
92	F	S	C	I	L	S	S	N	I	I	S	V	A	E	N	14
93	S	C	I	L	S	S	N	I	I	S	V	A	E	N	G	14
124	P	W	T	V	G	K	N	E	F	S	Q	T	V	C	G	14
136	V	G	E	V	F	Y	T	K	N	R	N	F	C	L	P	14
159	I	T	S	L	Q	Q	E	F	L	C	P	S	F	L	P	14
163	Q	Q	E	L	C	P	S	F	L	L	P	S	A	P	A	14
169	S	F	L	L	P	S	A	P	A	L	G	R	C	F	P	14
175	A	P	A	L	G	R	C	F	A	P	W	T	N	V	T	14
184	W	T	N	V	T	P	P	A	L	P	G	I	T	N	D	14
205	I	S	G	L	I	D	S	L	N	A	R	D	I	S	V	14
218	S	V	K	I	F	E	D	F	A	Q	S	W	Y	S	L	14
231	W	I	L	V	A	L	D	G	V	A	L	V	L	S	I	14
237	G	V	A	L	V	L	S	L	L	F	I	L	L	S	L	14
244	L	L	F	I	L	L	L	R	L	V	A	G	P	L	V	14
249	L	L	R	L	V	A	G	P	L	V	L	V	L	I	L	14
250	L	R	L	V	A	G	P	L	V	L	V	L	I	L	G	14
256	P	L	V	L	V	L	L	I	L	G	V	L	G	V	L	14
258	V	L	V	L	I	L	G	V	L	G	V	L	A	Y	G	14
260	V	L	I	L	G	V	L	G	V	L	A	Y	G	I	Y	14
263	L	G	V	L	G	V	L	A	Y	G	I	Y	Y	C	W	14
296	T	T	N	L	S	A	Y	Q	S	V	Q	E	T	W	L	14
302	Y	Q	S	V	Q	E	T	W	L	A	A	L	I	V	L	14
322	I	L	L	L	M	L	I	F	L	R	Q	R	I	R	I	14
338	I	A	L	L	K	E	A	S	K	A	V	G	Q	M	M	14
345	S	K	A	V	G	Q	M	M	S	T	M	F	Y	P	L	14
348	V	G	Q	M	M	S	T	M	M	F	Y	P	L	V	T	14
349	G	Q	M	M	S	T	M	F	Y	P	L	V	T	F	V	14
352	M	S	T	M	F	Y	P	L	V	T	F	V	L	L	Y	14
357	Y	P	L	V	T	F	V	L	L	L	I	C	I	A	Y	14
360	V	T	F	V	L	L	L	I	C	I	A	Y	W	A	M	14
361	T	F	V	L	L	L	I	C	I	A	Y	W	A	M	T	14
362	F	V	L	L	L	L	I	C	I	A	Y	W	A	M	T	14
366	L	I	C	I	A	Y	W	A	M	T	A	L	Y	L	A	14
376	A	L	Y	L	A	T	S	G	Q	P	Q	Y	V	L	W	14
391	A	S	N	I	S	S	P	G	C	E	K	V	P	I	N	14
399	C	E	K	V	P	I	N	T	S	C	N	P	T	A	H	14
411	T	A	H	L	V	N	S	S	C	P	G	L	M	C	V	14
412	A	H	L	V	N	S	S	C	P	G	L	M	C	V	F	14
422	L	M	C	V	F	Q	G	Y	S	S	K	G	L	I	Q	14
432	K	G	L	I	Q	R	S	V	S	N	L	Q	I	Y	G	14
439	V	F	N	L	Q	I	Y	G	V	L	G	L	F	W	T	14
441	N	L	Q	I	Y	G	V	L	G	L	F	W	T	L	N	14
455	N	W	V	L	A	L	G	Q	C	V	L	A	G	A	F	14
457	V	L	A	L	G	Q	C	V	L	A	G	A	F	A	S	14
462	Q	C	V	L	A	G	A	F	A	S	F	Y	W	S	A	14
489	S	A	F	I	R	T	L	R	Y	H	T	G	S	L	A	14
492	I	R	T	L	R	Y	H	T	G	S	L	A	F	G	A	14
499	T	G	S	L	A	F	G	A	L	I	L	T	L	V	Q	14
504	F	G	A	L	I	L	T	L	V	Q	I	A	R	V	I	14
509	L	T	L	V	Q	I	A	R	V	I	L	E	Y	I	D	14
515	A	R	V	I	L	E	Y	I	D	H	K	L	R	G	V	14
526	L	R	G	V	Q	N	P	V	A	R	C	I	M	C	C	14

282

TABLE XLVIII-continued

5	534	A	R	C	I	M	C	C	F	K	C	C	L	W	C	L	14	
	535	R	C	I	M	C	C	F	K	C	C	L	W	C	L	E	14	
	552	I	K	F	L	N	R	N	A	Y	I	M	I	A	I	Y	14	
	559	A	Y	I	M	I	A	I	Y	G	K	N	F	C	V	S	14	
	576	N	A	F	M	L	L	M	R	N	I	V	R	V	V	V	14	
	578	F	M	L	L	M	R	N	I	V	R	V	V	V	L	D	14	
	585	I	V	R	V	V	V	L	D	K	V	T	D	L	L	L	14	
	591	L	D	K	V	T	D	L	L	L	F	F	G	K	L	L	14	
	596	D	L	L	L	F	F	G	K	L	L	V	V	G	G	V	14	
	602	G	K	L	L	V	V	G	G	V	G	V	L	S	F	F	14	
10	603	K	L	L	V	V	G	G	V	G	V	L	S	F	F	14		
	604	L	L	V	V	G	G	V	G	V	L	S	F	F	F	14		
	607	V	G	G	V	G	V	L	S	F	F	S	F	S	G	R	14	
	609	G	V	G	V	L	S	F	F	F	F	S	G	R	I	P	14	
	610	V	G	V	L	S	F	F	F	F	S	G	R	I	P	G	14	
	622	I	P	G	L	G	K	D	F	K	S	P	H	L	N	Y	14	
	631	S	P	H	L	N	Y	Y	W	L	P	I	M	T	S	I	14	
	636	Y	Y	W	L	P	I	M	T	S	I	L	G	A	Y	V	14	
	647	G	A	Y	V	I	A	S	G	F	F	S	V	F	G	M	14	
	655	F	F	S	V	F	G	M	C	V	D	T	L	F	L	C	14	
20	658	V	F	G	M	C	V	D	T	L	F	L	C	F	L	E	14	
	663	V	D	T	L	F	L	C	F	L	E	D	L	E	R	N	14	
	665	T	L	F	L	C	F	L	E	D	L	E	R	N	N	G	14	
	678	N	G	S	L	D	R	P	Y	Y	M	S	K	S	L	L	14	
	684	P	Y	Y	M	S	K	S	L	L	K	I	L	G	K	K	14	
	689	K	S	L	L	K	I	L	G	K	K	N	E	A	P	P	14	
	V3-HLA-DR1-0401-15mers-24P4C12																	
	Each peptide is a portion of SEQ ID NO: 7; each start position is specified, the length of peptide is 15 amino acids, and the end position for each peptide is the start position plus fourteen.																	
	30	Pos	1	2	3	4	5	6	7	8	9	0	1	2	3	4	5	score
		35	9	C	F	P	W	T	N	I	T	P	P	A	L	P	G	I
		3	A	P	A	L	G	R	C	F	P	W	T	N	I	T	P	14
		12	W	T	N	I	T	P	P	A	L	P	G	I	T	N	D	14
		4	P	A	L	G	R	C	F	P	W	T	N	I	T	P	P	12
		5	A	L	G	R	C	F	P	W	T	N	I	T	P	P	A	12
40		8	R	C	F	P	W	T	N	I	T	P	P	A	L	P	G	12
		13	T	N	I	T	P	P	A	L	P	G	I	T	N	D	T	12
		14	N	I	T	P	P	A	L	P	G	I	T	N	D	T	T	12
		7	G	R	C	F	P	W	T	N	I	T	P	P	A	L	P	10
45	V5-DR1-0401-15mers-24P4C12																	
	Each peptide is a portion of SEQ ID NO: 11; each start position is specified, the length of peptide is 15 amino acids, and the end position for each peptide is the start position plus fourteen.																	
	50	Pos	1	2	3	4	5	6	7	8	9	0	1	2	3	4	5	score
	55	12	L	L	L	V	L	I	F	L	R	Q	R	I	R	I	A	26
		1	A	L	I	V	L	A	V	L	E	A	I	L	L	L	V	20
		2	L	I	V	L	A	V	L	E	A	I	L	L	L	V	L	20
		4	V	L	A	V	L	E	A	I	L	L	L	V	L	I	F	20
		5	L	A	V	L	E	A	I	L	L	V	L	I	I	F	L	20
		8	L	E	A	I	L	L	L	V	L	I	F	L	R	Q	R	20
		10	A	I	L	L	L	V	L	I	F	L	R	Q	R	I	R	20
60	60	13	L	V	L	I	F	L	R	Q	R	I	R	I	A	I	20	
		15	V	L	I	F	L	R	Q	R	I	R	I	A	I	A	16	
		14	L	V	L	I	F	L	R	Q	R	I	R	I	A	I	A	15
		9	E	A	I	L	L	V	L	I	F	L	R	Q	R	I	14	
		11	I	L	L	V	L	I	F	L	R	Q	R	I	R	I	14	
		3	I	V	L	A	V	L	E	A	I	L	L	V	L	I	12	
	65	6	A	V	L	E	A	I	L	L	L	V	L	I	F	L	R	12

283

TABLE XLVIII-continued

V6-HLA-DR1-0401-15mers- 24P4C12																
Each peptide is a portion of SEQ ID NO: 13; each start position is specified, the length of peptide is 15 amino acids, and the end position for each peptide is the start position plus fourteen.																
Pos	1	2	3	4	5	6	7	8	9	0	1	2	3	4	5	score
2	M	C	V	F	Q	G	Y	S	S	K	G	L	I	P	R	22
15	P	R	S	V	F	N	L	Q	I	Y	G	V	L	G	L	20
12	G	L	I	P	R	S	V	F	N	L	Q	I	Y	G	V	18
1	L	M	C	V	F	Q	G	Y	S	S	K	G	L	I	P	14
11	K	G	L	I	P	R	S	V	F	N	L	Q	I	Y	G	14
7	G	Y	S	S	K	G	L	I	P	R	S	V	F	N	L	12
8	Y	S	S	K	G	L	I	P	R	S	V	F	N	L	Q	12
9	S	S	K	G	L	I	P	R	S	V	F	N	L	Q	I	12
V7-HLA-DR1-0401-15mers- 24P4C12																
Each peptide is a portion of SEQ ID NO: 15; each start position is specified, the length of peptide is 15 amino acids, and the end position for each peptide is the start position plus fourteen.																
Pos	1	2	3	4	5	6	7	8	9	0	1	2	3	4	5	score
9	Y	W	I	L	V	A	V	G	Q	M	M	S	T	M	F	26
6	Q	S	W	Y	W	I	L	V	A	V	G	Q	M	M	S	22
7	S	W	Y	W	I	L	V	A	V	G	Q	M	M	S	T	22
8	W	Y	W	I	L	V	A	V	G	Q	M	M	S	T	M	20
1	F	E	D	F	A	Q	S	W	Y	W	I	L	V	A	V	16
5	A	Q	S	W	Y	W	I	L	V	A	V	G	Q	M	M	16
10	W	I	L	V	A	V	G	Q	M	M	S	T	M	F	Y	14
12	L	V	A	V	G	Q	M	M	S	T	M	F	Y	P	L	14
V8-HLA-DR1-0401- 15mers-24P4C12																
Each peptide is a portion of SEQ ID NO: 3; each start position is specified, the length of peptide is 17 amino acids, and the end position for each peptide is the start position plus fourteen.																
Pos	1	2	3	4	5	6	7	8	9	0	1	2	3	4	5	score
7	N	Y	Y	W	L	P	I	M	R	N	P	I	T	P	T	28
5	H	L	N	Y	Y	W	L	P	I	M	R	N	P	I	T	22
8	Y	Y	W	L	P	I	M	R	N	P	I	T	P	T	G	20
15	R	N	P	I	T	P	T	G	H	V	F	Q	T	S	I	20
26	Q	T	S	I	L	G	A	Y	V	I	A	S	G	F	F	20
18	I	T	P	T	G	H	V	F	Q	T	S	I	L	G	A	18
19	T	P	T	G	H	V	F	Q	T	S	I	L	G	A	Y	18
3	S	P	H	L	N	Y	Y	W	L	P	I	M	R	N	P	14
10	W	L	P	I	M	R	N	P	I	T	P	T	G	H	V	14
11	L	P	I	M	R	N	P	I	T	P	T	G	H	V	F	14
21	T	G	H	V	F	Q	T	S	I	L	G	A	Y	V	I	14
V9-HLA-DR1-0401- 15mers-24P4C12																
Each peptide is a portion of SEQ ID NO: 19; each start position is specified, the length of peptide is 15 amino acids, and the end position for each peptide is the start position plus fourteen.																
Pos	1	2	3	4	5	6	7	8	9	0	1	2	3	4	5	score
10	M	T	A	L	Y	P	L	P	T	Q	P	A	T	L	G	26
23	L	G	Y	V	L	W	A	S	N	I	S	S	P	G	C	26
11	T	A	L	Y	P	L	P	T	Q	P	A	T	L	G	Y	22
22	T	L	G	Y	V	L	W	A	S	N	I	S	S	P	G	22
7	Y	W	A	M	T	A	L	Y	P	L	P	T	Q	P	A	20
20	P	A	T	L	G	Y	V	L	W	A	S	N	I	S	S	20
5	I	A	Y	W	A	M	T	A	L	Y	P	L	P	T	Q	16
2	L	I	C	I	A	Y	W	A	M	T	A	L	Y	P	L	14
3	I	C	I	A	Y	W	A	M	T	A	L	Y	P	L	P	12

284

TABLE XLVIII-continued

15	P	L	P	T	Q	P	A	T	L	G	Y	V	L	W	A	12
21	A	T	L	G	Y	V	L	W	A	S	N	I	S	S	P	12

TABLE XLIX

V1-DRB1-1101-15mers-
24P4C12

Each peptide is a portion of SEQ ID
NO: 3; each start position is specified,
the length of peptide is 15 amino acids,
and the end position for each peptide is
the start position plus fourteen.

Pos	1	2	3	4	5	6	7	8	9	0	1	2	3	4	5	score
243	S	L	L	F	I	L	L	L	R	L	V	A	G	P	L	31
10	D	E	A	Y	G	K	P	V	K	Y	D	P	S	F	R	26
20	D	P	S	F	R	G	P	I	K	N	R	S	C	T	D	26
668	L	C	F	L	E	D	L	E	R	N	N	G	S	L	D	26
575	K	N	A	F	M	L	L	M	R	N	I	V	R	V	V	25
613	L	S	F	F	F	F	S	G	R	I	P	G	L	G	K	25
226	A	Q	S	W	Y	W	I	L	V	A	L	G	V	A	L	23
228	S	W	Y	W	I	L	V	A	L	G	V	A	L	V	L	23
277	W	E	E	Y	R	V	L	R	D	K	G	A	S	I	S	23
359	L	V	T	F	V	L	L	L	I	C	I	A	Y	W	A	23
448	L	G	L	F	W	T	L	N	W	V	L	A	L	G	Q	23
579	M	L	L	M	R	N	I	V	R	V	V	V	L	D	K	23
598	L	L	F	F	G	K	L	L	V	V	G	G	V	G	V	22
633	H	L	N	Y	Y	W	L	P	I	M	T	S	I	L	G	22
276	C	W	E	E	Y	R	V	L	R	D	K	G	A	S	I	21
338	I	A	L	L	K	E	A	S	K	A	V	G	Q	M	M	21
508	I	L	T	L	V	Q	I	A	R	V	I	L	E	Y	I	21
516	R	V	I	L	E	Y	I	D	H	K	L	R	G	V	Q	21
542	K	C	C	L	W	C	L	E	K	F	I	K	F	L	N	21
585	I	V	R	V	V	V	L	D	K	V	T	D	L	L	L	21
685	Y	Y	M	S	K	S	L	L	K	I	L	G	K	K	N	21
172	L	P	S	A	P	A	L	G	R	C	F	P	W	T	N	20
334	I	R	I	A	I	A	L	L	K	E	A	S	K	A	V	20
371	Y	W	A	M	T	A	L	Y	L	A	T	S	G	Q	P	20
549	E	K	F	I	K	F	L	N	R	N	A	Y	I	M	I	20
591	L	D	K	V	T	D	L	L	L	F	F	G	K	L	L	20
619	S	G	R	I	P	G	L	G	K	D	F	K	S	P	H	20
689	K	S	L	L	K	I	L	G	K	K	N	E	A	P	P	20
36	I	C	C	V	L	F	L	L	F	I	L	G	Y	I	V	19
122	E	D	P	W	T	V	G	K	N	E	F	S	Q	T	V	19
256	P	L	V	L	V	L	I	L	G	V	L	G	V	L	A	19
259	L	V	L	I	L	G	V	L	G	V	L	A	Y	G	I	19
310	L	A	A	L	I	V	L	A	V	L	E	A	I	L	L	19
353	S	T	M	F	Y	P	L	V	T	F	V	L	L	L	I	19
523	D	H	K	L	R	G	V	Q	N	P	V	A	R	C	I	19
567	G	K	N	F	C	V	S	A	K	N	A	F	M	L	L	19
612	V	L	S	F	F	F	F	S	G	R	I	P	G	L	G	19
636	Y	Y	W	L	P	I	M	T	S	I	L	G	A	Y	V	19
16	P	V	K	Y	D	P	S	F	R	G	P	I	K	N	R	18
48	Y	I	V	V	G	I	V	A	W	L	Y	G	D	P	R	18
85	Y	L	L	Y	F	N	I	F	S	C	I	L	S	S	N	18
137	G	E	V	F	Y	T	K	N	R	N	F	C	L	P	G	18
181	C	F	P	W	T	N	V	T	P	P	A	L	P	G	I	18
227	Q	S	W	Y	W	I	L	V	A	L	G	V	A	L	V	18
244	L	L	F	I	L	L	L	R	L	V	A	G	P	L	V	18
326	M	L	I	F	L	R	Q	R	I	R	I	A	I	A	L	18
419	C	P	G	L	M	C	V	F	Q	G	Y	S	S	K	G	18
469	F	A	S	F	Y	W	A	F	H	K	P	Q	D	I	P	18
470	A	S	F	Y	W	A	F	H	K	P	Q	D	I	P	T	18
488	I	S	A	F	I	R	T	L	R	Y	H	T	G	S	L	18
489	S	A	F	I	R	T	L	R	Y	H	T	G	S	L	A	18
597	L	L	L	F	F	G	K	L	L	V	V	G	G	V	G	18
41	F	L	L	F	I	L	G	Y	I	V	V	G	I	V	A	17
45	I	L	G	Y	I	V	V	G	I	V	A	W	L	Y	G	17
71	T	G	A	Y	C	G	M	G	E	N	K	D	K	P	Y	17
86	L	L	Y	F	N	I	F	S	C	I	L	S	S	N	I	17
306	Q	E	T	W	L	A	A	L	I	V	L	A	V	L	E	17
325	L	M	L	I	F	L	R	Q	R	I	R	I	A	I	A	17
354	T	M	F	Y	P	L	V	T	F	V	L	L	L	I	C	17
369	I	A	Y	W	A	M	T	A	L	Y	L	A	T	S	G	17
384	Q	P	Q	Y	V	L	W	A	S	N	I	S	S	P	G	17
442	L	Q	I	Y	G	V	L	G	L	F	W	T	L	N	W	17

285

TABLE XLIX-continued

482	I	P	T	F	P	L	I	S	A	F	I	R	T	L	R	17
501	S	L	A	F	G	A	L	I	L	T	L	V	Q	I	A	17
548	L	E	K	F	I	K	F	L	N	R	N	A	Y	I	M	17
615	F	F	F	F	S	G	R	I	P	G	L	G	K	D	F	17
635	N	Y	Y	W	L	P	I	M	T	S	I	L	G	A	Y	17
652	A	S	G	F	F	S	V	F	G	M	C	V	D	T	L	17
82	D	K	P	Y	L	L	Y	F	N	I	F	S	C	I	L	16
89	F	N	I	F	S	C	I	L	S	S	N	I	I	S	V	16
179	G	R	C	F	P	W	T	N	V	T	P	P	A	L	P	16
253	V	A	G	P	L	V	L	V	L	I	L	G	V	L	G	16
299	L	S	A	Y	Q	S	V	Q	E	T	W	L	A	A	L	16
323	L	L	L	M	L	I	F	L	R	Q	R	I	R	I	A	16
368	C	I	A	Y	W	A	M	T	A	L	Y	L	A	T	S	16
387	Y	V	L	W	A	S	N	I	S	S	P	G	C	E	K	16
490	A	F	I	R	T	L	R	Y	H	T	G	S	L	A	F	16
494	T	L	R	Y	H	T	G	S	L	A	F	G	A	L	I	16
506	A	L	I	L	T	L	V	Q	I	A	R	V	I	L	E	16
517	V	I	L	E	Y	I	D	H	K	L	R	G	V	Q	N	16
557	R	N	A	Y	I	M	I	A	I	Y	G	K	N	F	C	16
563	I	A	I	Y	G	K	N	F	C	V	S	A	K	N	A	16
583	R	N	I	V	R	V	V	V	L	D	K	V	T	D	L	16
646	L	G	A	Y	V	I	A	S	G	F	F	S	V	F	G	16
43	L	F	I	L	G	Y	I	V	V	G	I	V	A	W	L	15
44	F	I	L	G	Y	I	V	V	G	I	V	A	W	L	Y	15
47	G	Y	I	V	V	G	I	V	A	W	L	Y	G	D	P	15
54	V	A	W	L	Y	G	D	P	R	Q	V	L	Y	P	R	15
73	A	Y	C	G	M	G	E	N	K	D	K	P	Y	L	L	15
153	P	W	N	M	T	V	I	T	S	L	Q	Q	E	L	C	15
156	M	T	V	I	T	S	L	Q	Q	E	L	C	P	S	F	15
195	I	T	N	D	T	T	I	Q	Q	G	I	S	G	L	I	15
207	G	L	I	D	S	L	N	A	R	D	I	S	V	K	I	15
242	L	S	L	L	F	I	L	L	L	R	L	V	A	G	P	15
357	Y	P	L	V	T	F	V	L	L	L	I	C	I	A	Y	15
429	Y	S	S	K	G	L	I	Q	R	S	V	F	N	L	Q	15
485	F	P	L	I	S	A	F	I	R	T	L	R	Y	H	T	15
519	L	E	Y	I	D	H	K	L	R	G	V	Q	N	P	V	15
527	R	G	V	Q	N	P	V	A	R	C	I	M	C	C	F	15
545	L	W	C	L	E	K	F	I	K	F	L	N	R	N	A	15
595	T	D	L	L	L	F	F	G	K	F	L	V	V	G	G	15
600	F	F	G	K	L	L	V	V	G	G	V	G	V	L	S	15
603	K	L	L	V	V	G	G	V	G	V	L	S	F	F	F	15
681	L	D	R	P	Y	Y	M	S	K	S	L	L	K	I	L	15

V3-HLA-DRB1-1101-

15mers-24P4C12

Each peptide is a portion of SEQ ID NO: 7; each start position is specified, the length of peptide is 15 amino acids, and the end position for each peptide is the start position plus fourteen.

Pos	1	2	3	4	5	6	7	8	9	0	1	2	3	4	5	score
9	C	F	P	W	T	N	I	T	P	P	A	L	P	G	I	18
7	G	R	C	F	P	W	T	N	I	T	P	P	A	L	P	16
12	W	T	N	I	T	P	P	A	L	P	G	I	T	N	D	8

V5-HLA-DRB1-1101-

15mers-24P4C12

Each peptide is a portion of SEQ ID NO: 11; each start position is specified, the length of peptide is 15 amino acids, and the end position for each peptide is the start position plus fourteen.

Pos	1	2	3	4	5	6	7	8	9	0	1	2	3	4	5	score
15	V	L	I	F	L	R	Q	R	I	R	I	A	I	A	L	18
14	L	V	L	I	F	L	R	Q	R	I	R	I	A	I	A	17
12	L	L	L	V	L	I	F	L	R	Q	R	I	R	I	A	16
10	A	I	L	L	V	L	V	L	I	F	L	R	Q	R	I	15
2	L	I	V	L	A	V	L	E	A	I	L	L	L	V	L	14
8	L	E	A	I	L	L	V	L	I	F	L	R	Q	R	14	
13	L	L	V	L	I	F	L	R	Q	R	I	R	I	A	I	14
1	A	L	I	V	L	A	V	L	E	A	I	L	L	L	V	13
5	L	A	V	L	E	A	I	L	L	L	V	L	I	F	L	13
9	E	A	I	L	L	V	L	I	I	F	L	R	Q	R	I	13
11	I	L	L	L	V	L	I	F	L	R	Q	R	I	R	I	13

286

TABLE XLIX-continued

V6-HLA-DRB1-1101-

15mers-24P4C12

Each peptide is a portion of SEQ ID NO: 13; each start position is specified, the length of peptide is 15 amino acids, and the end position for each peptide is the start position plus fourteen.

10	Pos	1	2	3	4	5	6	7	8	9	0	1	2	3	4	5	score
15	8	Y	S	S	K	G	L	I	P	R	S	V	F	N	L	Q	15
	1	L	M	C	V	F	Q	G	Y	S	S	K	G	L	I	P	14
	15	P	R	S	V	F	N	L	Q	I	Y	G	V	L	G	L	13
	2	M	C	V	F	Q	G	Y	S	S	K	G	L	I	P	R	10
	5	F	Q	G	Y	S	S	K	G	L	I	P	R	S	V	F	10
	3	C	V	F	Q	G	Y	S	S	K	G	L	I	P	R	S	9
	11	K	G	L	I	P	R	S	V	F	N	L	Q	I	Y	G	9
	6	Q	G	Y	S	S	K	G	L	I	P	R	S	V	F	N	8
	4	V	F	Q	G	Y	S	S	K	G	L	I	P	R	S	V	7
7	G	Y	S	S	K	G	L	I	P	R	S	V	F	N	L	7	

V7-HLA-DRB1-1101-

15mers-24P4C12

Each peptide is a portion of SEQ ID NO: 15; each start position is specified, the length of peptide is 15 amino acids, and the end position for each peptide is the start position plus fourteen.

	Pos	1	2	3	4	5	6	7	8	9	0	1	2	3	4	5	score
30	5	A	Q	S	W	Y	W	I	L	V	A	V	G	Q	M	M	23
	6	Q	S	W	Y	W	I	L	V	A	V	G	Q	M	M	S	18
	9	Y	W	I	L	V	A	V	G	Q	M	M	S	T	M	F	18
	7	S	W	Y	W	I	L	V	A	V	G	Q	M	M	S	T	16
	12	L	V	A	V	G	Q	M	M	S	T	M	F	Y	P	L	12
	1	F	E	D	F	A	Q	M	S	W	Y	W	I	L	V	A	V

V8-HLA-DRB1-1101-15mers-

24P4C12

Each peptide is a portion of SEQ ID NO: 17; each start position is specified, the length of peptide is 15 amino acids, and the end position for each peptide is the start position plus fourteen.

	Pos	1	2	3	4	5	6	7	8	9	0	1	2	3	4	5	score
45	7	N	Y	Y	W	L	P	I	M	R	N	P	I	T	P	T	24
	5	H	L	N	Y	Y	W	L	P	I	M	R	N	P	I	T	18
	6	L	N	Y	Y	W	L	P	I	M	R	N	P	I	T	P	17
	15	R	N	P	I	T	P	T	G	H	V	F	Q	T	S	I	16
	8	Y	Y	W	L	P	I	M	R	N	P	I	T	P	T	G	13
	21	T	G	H	V	F	O	T	S	I	L	G	A	Y	V	I	13

V9-HLA-DRB1-1101-15mers-

24P4C12

Each peptide is a portion of SEQ ID NO: 19; each start position is specified, the length of peptide is 15 amino acids, and the end position for each peptide is the start position plus fourteen.

55

Pos	1	2	3	4	5	6	7	8	9	0	1	2	3	4	5	score	
60	4	C	I	A	Y	W	A	M	T	A	L	Y	P	L	P	T	22
	10	M	T	A	L	Y	P	L	P	T	Q	P	A	T	L	G	18
	22	T	L	G	Y	V	L	W	A	S	N	I	S	S	P	G	17
	7	Y	W	A	M	T	A	L	Y	P	L	P	T	Q	P	A	14
	13	L	Y	P	L	P	T	Q	P	A	T	L	G	Y	V	L	13
	20	P	A	T	L	G	Y	V	L	W	A	S	N	I	S	S	12
	23	L	G	Y	V	L	W	A	S	N	I	S	S	P	G	C	12
	24	G	Y	V	L	W	A	S	N	I	S	S	P	G	C	E	12
65	5	I	A	Y	W	A	M	T	A	L	Y	P	L	P	T	Q	10
	11	T	A	L	Y	P	L	P	T	Q	P	A	T	L	G	Y	10

TABLE L

Properties of 24P4C12			
	Bioinformatic Program	URL	Outcome
ORF	ORF finder		6 to 2138
Protein length			710aa
Transmembrane region	TM Pred	http://www.ch.embnet.org/	11TM, 39-59, 86-104, 231-250, 252-273, 309-330, 360-380, 457-474, 497-515, 559-581, 604-626, 641-663
	HMMTop	http://www.enzim.hu/hmmtop/	11TM, 35-59 84-104 231-250 257-277 308-330 355-377 456-475 500-519 550-572 597-618 649-671
	Sosui	http://www.genome.ad.jp/SOSui/	13TM, 34-65, 86-108, 145-167, 225-247, 307-329, 357-379, 414-436, 447-469, 501-523, 564-586, 600-622, 644-666
	TMHMM	http://www.cbs.dtu.dk/services/TMHMM	10TM, 36-58, 228-250, 252-274, 308-330, 356-378, 454-476, 497-519, 559-581, 597-619
Signal Peptide	Signal P	http://www.cbs.dtu.dk/services/SignalP/	no
pI	pI/MW tool	http://www.expasy.ch/tools/	8.9 pI
Molecular weight	pI/MW tool	http://www.expasy.ch/tools/	79.3 kD
Localization	PSORT	http://psort.nibb.ac.jp/	80% Plasma Membrane, 40% Golgi
	PSORT II	http://psort.nibb.ac.jp/	65% Plasma Membrane, 38% endoplasmic reticulum
Motifs	Pfam	http://www.sanger.ac.uk/Pfam/	DUF580, unknown function
	Prints	http://www.biochem.ucl.ac.uk/	
	Blocks	http://www.blocks.fhcrc.org/	Anion exchanger family 313-359
	Prosite	http://www.prosite.org/	CYS-RICH 536-547

TABLE LI

Exon compositions of 24P4C12 v. 1				35
Exon number	Start	End	Length	
1	1	45	45	40
2	46	94	49	
3	95	168	74	
4	169	247	79	45
5	248	347	100	
6	348	473	126	
7	474	534	61	
8	535	622	88	50
9	623	706	84	
10	707	942	236	
11	943	1042	100	
12	1043	1135	93	55
13	1136	1238	103	
14	1239	1492	254	
15	1493	1587	95	
16	1588	1691	104	
17	1692	1765	74	60
18	1766	1836	71	
19	1837	1931	95	
20	1932	2016	85	
21	2017	2573	557	65

TABLE LII

Nucleotide sequence of transcript variant 24P4C12 v.7 (SEQ ID NO: 94)		
gagccatggg gggaaagcag cgggacgagg atgacgaggc ctacgggaag ccagtcaaat	60	
acgacccctc ctttcgaggc cccatcaaga acagaagctg cacagatgtc atctgtctgcg	120	
tcctcttctc gctcttcatt ctaggttaca tcgtggtggg gattgtggcc tggttgtatg	180	
gagaccccg gcaagtccctc taccocagga actctactgg ggctactgt ggcatggggg	240	
agaacaaaga taagccgtat ctctgtact tcaacatctt cagctgcac ctgtccagca	300	
acatcatctc agttgtctgag aacggcctac agtgccccac accccagggtg tgtgtgtcct	360	
cctgcccga ggacccatgg actgtgggaa aaaacgagtt ctacagact gttggggaag	420	
tcctctatac aaaaaacagg aacttttgct tgccagggt accctggaat atgacggtga	480	
tcacaagcct gcaacaggaa ctctgcccc gtttctctct cccctctgct ccagctctgg	540	
ggcgtgctt tccatggacc aacgttactc caccggcgt cccagggtac accaatgaca	600	
ccaccatata gcaggggatc agcgggtctta ttgacagcct caatgcccga gacatcagtg	660	
ttaagatctt tgaagatctt gccagctctc ggtattggat tctgtgggt gtgggacaga	720	
tgatgtctac catgttctac ccactggta cctttgtcct cctctctac tgcattgcct	780	
actgggcat gactgtctct tacctggcta catcggggca accccagtat gtgtctctggg	840	
catccaacat cagctcccc ggctgtgaga aagtgcctat aaatacatca tgcaaccca	900	
cggccacct tgtgaactcc tcgtgccag ggctgatgtg cgtcttcag ggtactcat	960	
ccaaggcct aatccaacgt tctgtcttca atctgcaat ctatgggtc ctggggctct	1020	
tctggacct taactgggtg ctggccctgg gccaatgct cctcgctgga gcctttgct	1080	
ccttctactg ggcttccac aagcccagg acatccctac ctccctta atctctgcct	1140	
tcacccgac actccgttac cacactgggt cattggcatt tggagccctc atcctgaccc	1200	
ttgtgcagat agcccggtc atcttgagat atattgacca caagtcaga ggagtcaga	1260	
accctgtagc ccgctgcac atgtgtgtt tcaagtgtg cctctggtgt ctggaaaaat	1320	
ttatcaagtt cctaaaccgc aatgcataca tcatgatcgc catctacggg aagaatttct	1380	
gtgtctcagc caaaaatgct ttcattgtac tcatgcgaaa cattgtcagg gtggtcgtcc	1440	
tggacaaagt cacagacctg ctgctgttct ttgggaagct gctggtggc ggaggcgtgg	1500	
gggtcctgtc cttctttttt ttctccggtc gcaccccggt gctgggtaaa gactttaaga	1560	
gccccacct caactattac tggctgcaca tcatgacctc catcctgggt gcctatgtca	1620	
tcgccagcgg cttcttcagc gttttcgga tgtgtgtgga cacgtcttc ctctgttcc	1680	
tggagacct ggagcggaac aacggctccc tggaccggc ctactacatg tccaagagcc	1740	
ttctaaagat tctgggcaag aagaacgagg cgtcccgga caacaagaag aggaagaagt	1800	
gacagctccg gccctgatcc aggactgcac cccacccca ccgtccagcc atccaacctc	1860	
acttcgctt acaggtctcc atttgtggt aaaaaaggt tttaggccag gcgctgtgc	1920	
tcacgcctgt aatccaacac tttgagaggc tgaggcgggc ggatcacctg agtcaggagt	1980	
tcgagaccag cctggccaac atggtgaaac ctccgtctct attaaaaata caaaaattag	2040	
ccgagagtgg tgcatgcac ctgtcatccc agctactcgg gaggtgagg caggagaatc	2100	
gcttgaacc gggaggcaga ggttgagtg agccagatc ggcactgc actccaacct	2160	
gggtgacaga ctctgtctcc aaaaacaaac aaacaacaa aaagatttta ttaagatat	2220	
tttgtaact cagtaaaaa aaaaaaaaa a	2251	

TABLE LIII

Nucleotide sequence alignment of 24P4C12v.1 v.1 (SEQ ID NO: 95) and 24P4C12 v.7 (SEQ ID NO: 96).				
Score = 1358 bits (706), Expect = 0.0Identities = 706/706 (100%) Strand = Plus/Plus				
24P4C12v.1: 1	gagccatgggggaaagcagcgggacgaggatgacgaggcctacgggaagccagtcaa	60		
24P4C12v.7: 1	gagccatgggggaaagcagcgggacgaggatgacgaggcctacgggaagccagtcaa	60		
24P4C12v.1: 61	acgaccctcctttcgaggcccatcaagaacagaagctgcacagatgtcatctgtgcg	120		
24P4C12v.7: 61	acgaccctcctttcgaggcccatcaagaacagaagctgcacagatgtcatctgtgcg	120		
24P4C12v.1: 121	acgaccctcctttcgaggcccatcaagaacagaagctgcacagatgtcatctgtgcg	180		
24P4C12v.7: 121	acgaccctcctttcgaggcccatcaagaacagaagctgcacagatgtcatctgtgcg	180		
24P4C12v.1: 181	gagaccccggaagtcctctacccaggaaactctactgggctactgtggcatgggg	240		
24P4C12v.7: 181	gagaccccggaagtcctctacccaggaaactctactgggctactgtggcatgggg	240		
24P4C12v.1: 241	agaacaaagataagccgtatctcctgtacttcaacatcttcagctgcatcctgtccagca	300		
24P4C12v.7: 241	agaacaaagataagccgtatctcctgtacttcaacatcttcagctgcatcctgtccagca	300		
24P4C12v.1: 301	acatcatctcagttgctgagaacggcctacagtgtccacacccaggtgtgtgtgtcct	360		
24P4C12v.7: 301	acatcatctcagttgctgagaacggcctacagtgtccacacccaggtgtgtgtgtcct	360		
24P4C12v.1: 361	cctgcccggaggacccatggactgtgggaaaaaacgagttctcacagactgttggggaag	420		
24P4C12v.7: 361	cctgcccggaggacccatggactgtgggaaaaaacgagttctcacagactgttggggaag	420		
24P4C12v.1: 421	tcttctatacaaaaaacaggaacttttgtctgccaggggtaccctggaatatgacgggtga	480		
24P4C12v.7: 421	tcttctatacaaaaaacaggaacttttgtctgccaggggtaccctggaatatgacgggtga	480		
24P4C12v.1: 481	tcacaagcctgcaacaggaactctgccccagtttctcctcctcctctgtcctcagctctgg	540		
24P4C12v.7: 481	tcacaagcctgcaacaggaactctgccccagtttctcctcctcctctgtcctcagctctgg	540		
24P4C12v.1: 541	ggcgctgctttccatggaccaacgttactccacggcgctcccagggtacaccaatgaca	600		
24P4C12v.7: 541	ggcgctgctttccatggaccaacgttactccacggcgctcccagggtacaccaatgaca	600		
24P4C12v.1: 601	ccaccatacagcaggggatcagcgggtcttattgacagcctcaatgccgagacatcagt	660		
24P4C12v.7: 601	ccaccatacagcaggggatcagcgggtcttattgacagcctcaatgccgagacatcagt	660		
24P4C12v.1: 661	ttaagatctttgaagattttgccagtcctggatttgattcttgt	706		
24P4C12v.7: 661	ttaagatctttgaagattttgccagtcctggatttgattcttgt	706		
Score = 2971 bits (1545), Expect = 0.0Identities = 1545/1545 (100%) Strand = Plus/Plus				
24P4C12v.1: 1043	ggctgtgggacagatgatgtctaccatgttctacccactggtcacctttgtcctcctcct	1102		
24P4C12v.7: 707	ggctgtgggacagatgatgtctaccatgttctacccactggtcacctttgtcctcctcct	766		
24P4C12v.1: 1103	catctgcattgcctactgggcatgactgtctgtacctggctacatcggggcaacccca	1162		
24P4C12v.7: 767	catctgcattgcctactgggcatgactgtctgtacctggctacatcggggcaacccca	826		

TABLE LIII-continued

Nucleotide sequence alignment of 24P4C12v.1 v.1 (SEQ ID NO: 95) and 24P4C12 v.7 (SEQ ID NO: 96).				
24P4C12v.1: 1163	gtatgtgctctg	ggcatccaacatcagctccccggctgtgagaaagtgccataataac	1222	
24P4C12v.7: 827	gtatgtgctctg	ggcatccaacatcagctccccggctgtgagaaagtgccataataac	886	
24P4C12v.1: 1223	atcatgcaacccc	caggccaccttgtaactcctcg	tgccagggtgatgtgcgtctt	1282
24P4C12v.7: 887	atcatgcaacccc	caggccaccttgtaactcctcg	tgccagggtgatgtgcgtctt	946
24P4C12v.1: 1283	ccagggtactcat	ccaaaggcctaataccaacgttctgtcttcaatctgcaaatctatgg	1342	
24P4C12v.7: 947	ccagggtactcat	ccaaaggcctaataccaacgttctgtcttcaatctgcaaatctatgg	1006	
24P4C12v.1: 1343	ggctctggggct	cttctggacccttaactgggtactggcctgggccaatgcgtcctcgc	1402	
24P4C12v.7: 1007	ggctctggggct	cttctggacccttaactgggtactggcctgggccaatgcgtcctcgc	1066	
24P4C12v.1: 1403	tggagcctttgc	ctccttctactgggccttcacaagccccaggacatccctaccttccc	1462	
24P4C12v.7: 1067	tggagcctttgc	ctccttctactgggccttcacaagccccaggacatccctaccttccc	1126	
24P4C12v.1: 1463	cttaatctctgc	cttcatccgcacactccgttaccacactgggtcattggcatttggagc	1522	
24P4C12v.7: 1127	cttaatctctgc	cttcatccgcacactccgttaccacactgggtcattggcatttggagc	1186	
24P4C12v.1: 1523	cctcatcctgac	ccttgtgcagatagcccggtcatcttggagtattgaccacaagct	1582	
24P4C12v.7: 1187	cctcatcctgac	ccttgtgcagatagcccggtcatcttggagtattgaccacaagct	1246	
24P4C12v.1: 1583	cagaggagtgc	cagaacctgtagcccgcgtgcacatcatgtgctgtttcaagtgtgcctctg	1642	
24P4C12v.7: 1247	cagaggagtgc	cagaacctgtagcccgcgtgcacatcatgtgctgtttcaagtgtgcctctg	1306	
24P4C12v.1: 1643	gtgtctggaaaa	atttatcaagttcctaaccgcaatgcatacatcatgatcgccatcta	1702	
24P4C12v.7: 1307	gtgtctggaaaa	atttatcaagttcctaaccgcaatgcatacatcatgatcgccatcta	1366	
24P4C12v.1: 1703	cgggaagaattt	ctgtgtctcagccaaaaatgcgttcattgctactcatgcgaaacattgt	1762	
24P4C12v.7: 1367	cgggaagaattt	ctgtgtctcagccaaaaatgcgttcattgctactcatgcgaaacattgt	1426	
24P4C12v.1: 1763	cagggtggctgc	tcttggaacaaagtacagacctgctgctgttctttgggaagctgctggt	1822	
24P4C12v.7: 1427	cagggtggctgc	tcttggaacaaagtacagacctgctgctgttctttgggaagctgctggt	1486	
24P4C12v.1: 1823	ggtcggaggcgt	gggggtcctgtccttctttttttctccggtcgcacccggggctggg	1882	
24P4C12v.7: 1487	ggtcggaggcgt	gggggtcctgtccttctttttttctccggtcgcacccggggctggg	1546	
24P4C12v.1: 1883	taaagacttta	agagccccacctcaactattactgggtgcccatcatgacctccatcct	1942	
24P4C12v.7: 1547	taaagacttta	agagccccacctcaactattactgggtgcccatcatgacctccatcct	1606	
24P4C12v.1: 1943	gggggcctatgt	catcgccagcggtctcttcagcggttttcggcatgtgtgtggacacgct	2002	
24P4C12v.7: 1607	gggggcctatgt	catcgccagcggtctcttcagcggttttcggcatgtgtgtggacacgct	1666	

TABLE LIII-continued

Nucleotide sequence alignment of 24P4C12v.1 v.1 (SEQ ID NO: 95) and 24P4C12 v.7 (SEQ ID NO: 96).			
24P4C12v.1: 2003	cttctctgcttctctggaagacctggagcggaacaacggctccctggaccggccctacta	2062	
24P4C12v.7: 1667	cttctctgcttctctggaagacctggagcggaacaacggctccctggaccggccctacta	1726	
24P4C12v.1: 2063	catgtccaagagccttctaagattctgggcaagaagaacgaggcgccccggacaacaa	2122	
24P4C12v.7: 1727	catgtccaagagccttctaagattctgggcaagaagaacgaggcgccccggacaacaa	1786	
24P4C12v.1: 2123	gaagaggaagaagtgcacagctccggccctgatccaggactgcacccccacccccaccgtcc	2182	
24P4C12v.7: 1787	gaagaggaagaagtgcacagctccggccctgatccaggactgcacccccacccccaccgtcc	1846	
24P4C12v.1: 2183	agccatccaacctcacttcgccttacagggtctccattttgtggtaaaaaagggttttagg	2242	
24P4C12v.7: 1847	agccatccaacctcacttcgccttacagggtctccattttgtggtaaaaaagggttttagg	1906	
24P4C12v.1: 2243	ccaggcgccgtggctcacgcctgtaatccaacctttgagaggctgaggcgggcggtatca	2302	
24P4C12v.7: 1907	ccaggcgccgtggctcacgcctgtaatccaacctttgagaggctgaggcgggcggtatca	1966	
24P4C12v.1: 2303	cctgagtcaggagttcgagaccagcctggccaacatggtgaaacctccgtctctattaaa	2362	
24P4C12v.7: 1967	cctgagtcaggagttcgagaccagcctggccaacatggtgaaacctccgtctctattaaa	2026	
24P4C12v.1: 2363	aatacaaaaattagccgagagtggtggcatgcacctgtcatcccagctactcgggaggct	2422	
24P4C12v.7: 2027	aatacaaaaattagccgagagtggtggcatgcacctgtcatcccagctactcgggaggct	2086	
24P4C12v.1: 2423	gaggcaggagaatcgcttgaacccgggaggcagaggttgacgtgagccgagatcgcgcca	2482	
24P4C12v.7: 2087	gaggcaggagaatcgcttgaacccgggaggcagaggttgacgtgagccgagatcgcgcca	2146	
24P4C12v.1: 2483	ctgcactccaacctgggtgacagactctgtctccaaaacaaaacaaaacaaaagat	2542	
24P4C12v.7: 2147	ctgcactccaacctgggtgacagactctgtctccaaaacaaaacaaaacaaaagat	2206	
24P4C12v.1: 2543	tttattaaagatatattttgttaactcagtaaaaaaaaaaaaaaaaaa	2587	
24P4C12v.7: 2207	tttattaaagatatattttgttaactcagtaaaaaaaaaaaaaaaaaa	2251	

40

TABLE LIV

Peptide sequences of protein coded by 24P4C12 v.7 (SEQ ID NO: 97)	
MGGKQRDEDD EAYGKPVKYD PSFRGPIKNR SCTDVICCVL FLLFILGYIV VGIVAWLYGD	60
PRQVLYPRNS TGAYCGMGEN KDKPYLLYFN IFSCILSSNI ISVAENGLQC PTPQVCVSSC	120
PEDPWTVGKN EFSQTVGEVF YTKNRNFCLP GVPWNMTVIT SLQQELCPSE LLPSAPALGR	180
CFPWTNVTTP ALPGITNDTT IQQGISGLID SLNARDISVK IPEDFAQSWY WILVAVGQMM	240
STMFYPLVTF VLLLICIAIW AMTALYLATS GQPQYVLWAS NISSPGCEKV PINTSCNPTA	300
HLVNSSCPGL MCVFQGYSSK GLIQRVFNEL QIYGVLGLFW TLNWVLALGQ CVLAGAFASF	360
YWAFHKPQDI PTFPLISAFI RTLRYHTGSL AFGALILTLV QIARVILEYI DHKLRGVQNP	420
VARCIMCCFK CCLWCLEKFI KFLNRNAYIM IAIYGKNFCV SAKNAFMLLM RNIVRVVULD	480
KVTDLLLFPG KLLVVGGVGV LSFFFFSGRI PGLGKDFKSP HLNYYWLPIM TSILGAYVIA	540
SGFFSVFGMC VDTLFLCFLE DLERNNGSLD RPYYSKSLK KILGKKNEAP PDNKKRKK	598

TABLE LV

Amino acid sequence alignment of 24P4C12v.1 v.1 (SEQ ID NO: 98) and 24P4C12 v.7 (SEQ ID NO: 99).		
Score = 1195 bits (3091), Expect = 0.0Identities = 598/710 (84%), Positives = 598/710 (84%), Gaps = 112/710 (15%)		
24P4C12v.1:	1MGGKQRDEDDDEAYGKPVKYDPSFRGPIKNRSCDVI CCVLFLFLFILGYIVVGIVAWLYGD	60
	MGKKQRDEDDDEAYGKPVKYDPSFRGPIKNRSCDVI CCVLFLFLFILGYIVVGIVAWLYGD	
24P4C12v.7:	1MGGKQRDEDDDEAYGKPVKYDPSFRGPIKNRSCDVI CCVLFLFLFILGYIVVGIVAWLYGD	60
24P4C12v.1:	61PRQVLYPRNSTGAYCGMGENKDKPYLLYFNIFSCILSSNII SVAENGLQCPTPQVCVSSC	120
	PRQVLYPRNSTGAYCGMGENKDKPYLLYFNIFSCILSSNII SVAENGLQCPTPQVCVSSC	
24P4C12v.7:	61PRQVLYPRNSTGAYCGMGENKDKPYLLYFNIFSCILSSNII SVAENGLQCPTPQVCVSSC	120
24P4C12v.1:	121PEDPWTVGKNEFSQTVGEVFYTKNRNFCPLPGVPWNMTVITSLQQELCP SFLLPSAPALGR	180
	PEDPWTVGKNEFSQTVGEVFYTKNRNFCPLPGVPWNMTVITSLQQELCP SFLLPSAPALGR	
24P4C12v.7:	121PEDPWTVGKNEFSQTVGEVFYTKNRNFCPLPGVPWNMTVITSLQQELCP SFLLPSAPALGR	180
24P4C12v.1:	181CFPWTNVTTPPALPGITNDTTIQQGISGLIDSLNARDISVKI FEDFAQSWYWILVALGVAL	240
	CFPWTNVTTPPALPGITNDTTIQQGISGLIDSLNARDISVKI FEDFAQSWYWILVA	
24P4C12v.7:	181CFPWTNVTTPPALPGITNDTTIQQGISGLIDSLNARDISVKI FEDFAQSWYWILVA-----	235
24P4C12v.1:	241VLSLLFILLRLVAGPLVLVLILGLVGLAYGIYYCWEYRVL RDKGASISQLGFTTNLS	300
24P4C12v.7:	235-----	235
24P4C12v.1:	301AYQSQETWLAALIVLAVLEAILLLMLIFLRQRIRIAIAL LKEASKAVGQMMSTMFYPLV	360
	VGQMMSTMFYPLV	
24P4C12v.7:	236-----VGQMMSTMFYPLV	248
24P4C12v.1:	361TFVLLLLICIAYWAMTALYLATSGQPQYVLWASNIS SPGCEKVPINTSCNPTAHLVNSSCP	420
	TFVLLLLICIAYWAMTALYLATSGQPQYVLWASNIS SPGCEKVPINTSCNPTAHLVNSSCP	
24P4C12v.7:	249TFVLLLLICIAYWAMTALYLATSGQPQYVLWASNIS SPGCEKVPINTSCNPTAHLVNSSCP	308
24P4C12v.1:	421GLMCVFQGYSSKGLIQRSVFNLQIYGVGLFWTLNWLAL GQCVLAGAFASFYWAFHKPQ	480
	GLMCVFQGYSSKGLIQRSVFNLQIYGVGLFWTLNWLAL GQCVLAGAFASFYWAFHKPQ	
24P4C12v.7:	309GLMCVFQGYSSKGLIQRSVFNLQIYGVGLFWTLNWLAL GQCVLAGAFASFYWAFHKPQ	368
24P4C12v.1:	481DIPTFPLISAFIRTLRYHTGSLAFGALILTLVQIARVILEYID HKLRGVQNPVARCIMCC	540
	DIPTFPLISAFIRTLRYHTGSLAFGALILTLVQIARVILEYID HKLRGVQNPVARCIMCC	
24P4C12v.7:	369DIPTFPLISAFIRTLRYHTGSLAFGALILTLVQIARVILEYID HKLRGVQNPVARCIMCC	428
24P4C12v.1:	541FKCCLWCLEKFIKFLNRNAYIMIAIYGKNFCVSAKNAFMLLM RNIVRVVVLDKVTDLLLF	600
	FKCCLWCLEKFIKFLNRNAYIMIAIYGKNFCVSAKNAFMLLM RNIVRVVVLDKVTDLLLF	
24P4C12v.7:	429FKCCLWCLEKFIKFLNRNAYIMIAIYGKNFCVSAKNAFMLLM RNIVRVVVLDKVTDLLLF	488
24P4C12v.1:	601FGKLLVVGGVGVLSFFFFSGRIPGLGKDFKSPHLNYYWLPIMT SILGAYVIASGFFSVFG	660
	FGKLLVVGGVGVLSFFFFSGRIPGLGKDFKSPHLNYYWLPIMT SILGAYVIASGFFSVFG	
24P4C12v.7:	489FGKLLVVGGVGVLSFFFFSGRIPGLGKDFKSPHLNYYWLPIMT SILGAYVIASGFFSVFG	548
24P4C12v.1:	661MCVDTLFLCFLEDLERNNGSLDRPYYMSKSLKILGKKNEAPD NKRRKK	710
	MCVDTLFLCFLEDLERNNGSLDRPYYMSKSLKILGKKNEAPD NKRRKK	
24P4C12v.7:	549MCVDTLFLCFLEDLERNNGSLDRPYYMSKSLKILGKKNEAPD NKRRKK	598

TABLE LVI

Nucleotide sequence of transcript variant 24P4C12 v.8 (SEQ ID NO: 100)		
gagccatggg	gggaaagcag cgggacgagg atgacgaggc ctacgggaag ccagtcaa	60
acgacccctc	ctttcgaggc cccatcaaga acagaagctg cacagatgtc atctgtctgc	120
tcctcttctc	gctcttcatt ctaggttaca tcgtggtggg gattgtggcc tggttgtatg	180
gagacccccg	gcaagtcttc taccacagga actctactgg ggcctactgt ggcattgggg	240
agaacaaaga	taagccgtat ctctctgtact tcaacatctt cagctgcac	300
acatcatctc	agttgctgag aacggcctac agtgccccac accccaggtg tgtgtgtcct	360
cctgcccga	ggacccatgg actgtgggaa aaaacagatt ctcacagact gttgggggaa	420
tcttctatac	aaaaaacagg aacttttgtc tgccaggggt accctggaat atgacggtga	480
tcacaagcct	gcaacaggaa ctctgcccc gtttcctcct cccctctgct ccagctctgg	540
ggcgctgctt	tccatggacc aacgttactc caccggcgct cccagggatc accaatgaca	600
ccaccataca	gcaggggatc agcggtctta ttgacagcct caatgcccc gacatcagtg	660

Figure 2H-3

```
2341  tgagtcaggaggttcgagaccagcctggccaacatggtgaaacctccgtctctattaaaaa
2401  tacaaaaattagccgagagtggatggcatgcacctgtcatccagctactcgaggaggtga
2461  ggcaggagaatcgcttgaacccgggaggcagaggttgacgtgagccgagatcgcgccact
2521  gcactccaacctgggtgacagactctgtctccaaaacaaaacaaaacaaaagattt
2581  tattaaagatattttgttaactcagtaaaaaaaaaaaaaaaaaa
```

Figure 2I. The cDNA (SEQ ID. NO. : 18) and amino acid sequence (SEQ ID. NO. : 19) of 24P4C12 v.9.

The start methionine is underlined. The open reading frame extends from nucleic acid 6-2144 including the stop codon.

Figure 2I-1

```
1      M G G K Q R D E D D E A Y G K P V K Y
1 gagccATGGGGGGAAAGCAGCGGGACGAGGATGACGAGGCCTACGGGAAGCCAGTCAAAT
20   D P S F R G P I K N R S C T D V I C C V
61 ACGACCCCTCCTTTCGAGGCCCCATCAAGAACAGAAGCTGCACAGATGTCAICTGCTGCG
40   L F L L F I L G Y I V V G I V A W L Y G
121 TCCTCTTCCTGCTCTTCATTCTAGGTTACATCGTGGTGGGGATTGTGGCCIGGTTGTATG
60   D P R Q V L Y P R N S T G A Y C G M G E
181 GAGACCCCGGCAAGTCCTCTACCCCAGGAACCTCTACGCGGCCTACTGTGGCAITGGGGG
80   N K D K P Y L L Y F N I F S C I L S S N
241 AGAACAAGAGATAAGCCGIATCTCTGTACTTCAACATCTTCAGCTGCATCCTGTCCAGCA
100  I I S V A E N G L Q C P T P Q V C V S S
301 ACATCATCTCAGTTGCTGAGAACGGCCTACAGTGCCCCACACCCAGGTGTGTGTCTCT
120  C P E D P W T V G K N E F S Q T V G E V
361 CCTGCCCGGAGGACCCATGGACTGTGGGAAAAAACGAGTTCTCACAGACTGTITGGGGAAG
140  F Y I K N R N F C L P G V P W N M T V I
421 TCTTCTATACAAAAACAGGAACCTTTTGTCTGCCAGGGGTACCCCTGGAATAIGACGGTGA
160  T S L Q Q E L C P S F L L P S A P A L G
481 TCACAAGCCTGCAACAGGAACCTCTGCCCCAGTTTCTCTCTCCCTCTGCTCCAGCTCTGG
180  R C F P W T N V T P P A L P G I T N D T
541 GCGCTGCTTTCCATGGACCAACGTTACTCCACCGCGCTCCCAGGGATCACCAATGACA
200  T I Q Q G I S G L I D S L N A R D I S V
601 CCACCATACAGCAGGGGATCAGCGGTCTTATTGACAGCCTCAATGCCCGAGACATCAGTG
220  K I F E D F A Q S W Y W I L V A L G V A
661 TTAAGATCTTTGAAGATTTTGGCCAGTCTGGTATTGGATTCTTGTGCCCCTGGGGGTGG
240  L V L S L L F I L L L R L V A G P L V L
721 CTCTGGTCTTGAGCTACTGTTTATCTTGTCTTCTGCGCTGGTGGCTGGGCCCTGGTGC
260  V L I L G V L G V L A Y G I Y Y C W E E
781 TGGTGTGATCCTGGGAGTGCTGGGCGTGTGGCATACGGCATCTACTACTGCTGGGAGG
280  Y R V L R D K G A S I S Q L G F T T N L
841 AGTACCGAGTGCTGCGGGACAAGGGCGCCTCCATCTCCAGCTGGGTTTACCACCAAC
```

TABLE LVI-continued

Nucleotide sequence of transcript variant 24P4C12 v.8 (SEQ ID NO: 100)	
ttaagatctt tgaagatctt gccagctctt ggtattggat tcttggtgcc ctgggggttg	720
ctctggctct gagcctactg tttatcttgc ttctgcgcct ggtggctggg cccctggctg	780
tgggtctgat cctgggagtg ctgggcgtgc tggcatacgg catctactac tgctgggagg	840
agtaccgagt gctgcgggac aagggcgcct ccatctccca gctgggttcc accaccaacc	900
tcagtgccta ccagagcgtg caggagacct ggtggccgcg cctgatcgtg ttggcggctg	960
ttgaagccat cctgctgctg atgctcatct tcctgcggca gcggattcgt attgccatcg	1020
ccctcctgaa ggaggccagc aaggctgtgg gacagatgat gtctaccatg ttctaccac	1080
tggcacctt tgcctcctc ctcatctgca ttgcctactg ggccatgact gctctgtacc	1140
tggctacatc ggggcaacc cagtatgtgc tctgggcata caacatcagc tccccggct	1200
gtgagaaagt gccaataaat acatcatgca accccacggc ccacctgtg aactcctcgt	1260
gccagggtg gatgtgcgtc ttccagggtc actcatccaa aggcctaata caacgttctg	1320
tctcaatct gcaaatctat ggggtcctgg ggtcttctg gaccttaac tgggtactgg	1380
ccctgggcca atgcgtcctc gctggagcct ttgcctcctt ctactgggccc ttccacaagc	1440
cccaggacat cctacacctc cccttaatct ctgccttcat ccgcacactc cgttaccaca	1500
ctgggtcatt ggcatttggg gccctcatcc tgaccttgtg gcagatagcc cgggtcatct	1560
tggagtatat tgaccacaag ctacagaggg tgcagaaccc tgtagccgcg tgcacatgt	1620
gctgtttcaa gtgctgcctc tgggtgtctg aaaaatttat caagttccta aaccgcaatg	1680
catacatcat gatcgccata tacgggaaga atttctgtgt ctacgcaaaa aatgcgttca	1740
tgctactcat gcgaaacatt gtcagggtgg tcgtcctgga caaagtcaca gacctgctgc	1800
tgttctttgg gaagctgctg gtggctggag gcgtgggggt cctgtccttc tttttttct	1860
ccggtcgcct cccggggctg ggtaagact ttaagagccc ccacctcaac tattactggc	1920
tgcccatcat gaggaacca ataaccctaa cgggtcatgt cttccagacc tccatcctgg	1980
gggcctatgt catcgccagc ggcttcttca gcgttttcgg catgtgtgtg gacacgctct	2040
tcctctgctt cctggaagac ctggagcgga acaacggctc cctggaccgg ccctactaca	2100
tgtccaagag ccttctaaag attctgggca agaagaacga ggcgcccccg gacaacaaga	2160
agaggaagaa gtgacagctc cggccctgat ccaggactgc accccacccc caccgtccag	2220
ccatccaacc tcacttcgct ttacaggtct ccattttgtg gtaaaaaaag gttttaggcc	2280
aggcgccgtg gctcacgcct gtaatccaac actttgagag gctgaggcgg gcggatcacc	2340
tgagtcagga gttcgagacc agcctggcca acatggtgaa acctcgtct ctattaaaaa	2400
tacaaaaatt agccgagagt ggtggcatgc acctgtcatc ccagctactc gggaggctga	2460
ggcaggagaa tcgcttgaac ccgggaggca gaggttgcat tgagccgaga tcgcgccact	2520
gcactccaac ctgggtgaca gactctgtct ccaaaacaaa acaaacaaac aaaaagattt	2580
tattaaagat attttgttaa ctacgtaaaa aaaaaaaaaa aaa	2623

TABLE LVII

Nucleotide sequence alignment of 24P4C12v.1 v.1 (SEQ ID NO: 101) and 24P4C12 v.8
(SEQ ID NO: 102)

Score = 3715 bits (1932), Expect = 0.0 Identites = 1932/1932 (100%) Strand = Plus/Plus

24P4C12v.1: 1	gagccatgggggaaagcagcgggacgaggatgacgaggcctacgggaagccagtcaa	60
24P4C12v.8: 1	gagccatgggggaaagcagcgggacgaggatgacgaggcctacgggaagccagtcaa	60
24P4C12v.1: 61	acgacccctcctttcgaggcccatcaagaacagaagctgcacagatgtcatctgtgcg	120
24P4C12v.8: 61	acgacccctcctttcgaggcccatcaagaacagaagctgcacagatgtcatctgtgcg	120
24P4C12v.1: 121	tctcttctctgctcttcatcttaggtacatcgtggtgggattgtggcctggtgtatg	180
24P4C12v.8: 121	tctcttctctgctcttcatcttaggtacatcgtggtgggattgtggcctggtgtatg	180
24P4C12v.1: 181	gagacccccggcaagtctctacccaggaaactctactgggcctactgtggcatggggg	240
24P4C12v.8: 181	gagacccccggcaagtctctacccaggaaactctactgggcctactgtggcatggggg	240
24P4C12v.1: 241	agaacaaagataagccgtatctcctgtacttcaacatcttcagctgcatcctgtccagca	300
24P4C12v.8: 241	agaacaaagataagccgtatctcctgtacttcaacatcttcagctgcatcctgtccagca	300
24P4C12v.1: 301	acatcatctcagttgctgagaacggcctacagtgcacacacccaggtgtgtgtcct	360
24P4C12v.8: 301	acatcatctcagttgctgagaacggcctacagtgcacacacccaggtgtgtgtcct	360
24P4C12v.1: 361	cctgcccggaggaccatggactgtgggaaaaaacaggttctcacagactgttggggaag	420
24P4C12v.8: 361	cctgcccggaggaccatggactgtgggaaaaaacaggttctcacagactgttggggaag	420
24P4C12v.1: 421	tcttctatacaaaaaacaggaacttttgtctgccaggggtaccctggaatatgacggtga	480
24P4C12v.8: 421	tcttctatacaaaaaacaggaacttttgtctgccaggggtaccctggaatatgacggtga	480
24P4C12v.1: 481	tcacaagcctgcaacaggaactctgccccagtttctcctcctccctctgctccagctctgg	540
24P4C12v.8: 481	tcacaagcctgcaacaggaactctgccccagtttctcctcctccctctgctccagctctgg	540
24P4C12v.1: 541	ggcgctgctttccatggaccaacgttactccaccggcgctccagggtacaccaatgaca	600
24P4C12v.8: 541	ggcgctgctttccatggaccaacgttactccaccggcgctccagggtacaccaatgaca	600
24P4C12v.1: 601	ccaccatacagcaggggatcagcgggtcttattgacagcctcaatgcccgagacatcagtg	660
24P4C12v.8: 601	ccaccatacagcaggggatcagcgggtcttattgacagcctcaatgcccgagacatcagtg	660
24P4C12v.1: 661	ttaagatctttgaagattttggccagtcctggtattggattcttgttgccctgggggtgg	720
24P4C12v.8: 661	ttaagatctttgaagattttggccagtcctggtattggattcttgttgccctgggggtgg	720
24P4C12v.1: 721	ctctgggtcttgagcctactgtttatcttgcttctgcgcctgggtgggcccctgggtgc	780
24P4C12v.8: 721	ctctgggtcttgagcctactgtttatcttgcttctgcgcctgggtgggcccctgggtgc	780
24P4C12v.1: 781	tgggtgctgatcctgggagtgtggcggtgctggcatacgcatctactactgctgggagg	840
24P4C12v.8: 781	tgggtgctgatcctgggagtgtggcggtgctggcatacgcatctactactgctgggagg	840

TABLE LVII-continued

Nucleotide sequence alignment of 24P4C12v.1 v.1 (SEQ ID NO: 101) and 24P4C12 v.8 (SEQ ID NO: 102)				
24P4C12v.1:	841	agtagcagtgctgcgggacaagggcgccctccatctcccagctgggtttcaccaccaacc	900	
24P4C12v.8:	841	agtagcagtgctgcgggacaagggcgccctccatctcccagctgggtttcaccaccaacc	900	
24P4C12v.1:	901	tcagtgccctaccagagcgtgcaggagacctggctggcgccctgatcggttggcggtgc	960	
24P4C12v.8:	901	tcagtgccctaccagagcgtgcaggagacctggctggcgccctgatcggttggcggtgc	960	
24P4C12v.1:	961	ttgaagccatcctgctgctgatgctcatcttctggcgagcggattcgattgccatcg	1020	
24P4C12v.8:	961	ttgaagccatcctgctgctgatgctcatcttctggcgagcggattcgattgccatcg	1020	
24P4C12v.1:	1021	ccctcctgaaggaggccagcaaggctgtgggacagatgatgtctaccatgttctaccac	1080	
24P4C12v.8:	1021	ccctcctgaaggaggccagcaaggctgtgggacagatgatgtctaccatgttctaccac	1080	
24P4C12v.1:	1081	tggtcacctttgtcctcctcctcatctgcattgcctactgggccatgactgctctgtacc	1140	
24P4C12v.8:	1081	tggtcacctttgtcctcctcctcatctgcattgcctactgggccatgactgctctgtacc	1140	
24P4C12v.1:	1141	tggtacatcggggcaacccagtatgtgctctgggcatccaacatcagctccccggct	1200	
24P4C12v.8:	1141	tggtacatcggggcaacccagtatgtgctctgggcatccaacatcagctccccggct	1200	
24P4C12v.1:	1201	gtgagaaagtgccataaatacatcatgcaacccacggccaccttgtgaactcctcgt	1260	
24P4C12v.8:	1201	gtgagaaagtgccataaatacatcatgcaacccacggccaccttgtgaactcctcgt	1260	
24P4C12v.1:	1261	gcccagggtgatgtgctcttccagggtactcatccaaaggcctaatacgaacttctg	1320	
24P4C12v.8:	1261	gcccagggtgatgtgctcttccagggtactcatccaaaggcctaatacgaacttctg	1320	
24P4C12v.1:	1321	tcttcaatctgcaaatctatggggctcctggggtcttctggacccttaactgggtactgg	1380	
24P4C12v.8:	1321	tcttcaatctgcaaatctatggggctcctggggtcttctggacccttaactgggtactgg	1380	
24P4C12v.1:	1381	ccctgggccaatgcgtcctcgctggagcctttgcctccttctactgggccttccacaagc	1440	
24P4C12v.8:	1381	ccctgggccaatgcgtcctcgctggagcctttgcctccttctactgggccttccacaagc	1440	
24P4C12v.1:	1441	cccaggacatccctaccttcccttaatactctgcttcatccgcacactccgttaccaca	1500	
24P4C12v.8:	1441	cccaggacatccctaccttcccttaatactctgcttcatccgcacactccgttaccaca	1500	
24P4C12v.1:	1501	ctgggtcattggcatttggagccctcatcctgacccttgtagatagccgggtcatct	1560	
24P4C12v.8:	1501	ctgggtcattggcatttggagccctcatcctgacccttgtagatagccgggtcatct	1560	
24P4C12v.1:	1561	tggagtatattgaccacaagctcagaggagtgcagaacccgtgagccgctgcatcatgt	1620	
24P4C12v.8:	1561	tggagtatattgaccacaagctcagaggagtgcagaacccgtgagccgctgcatcatgt	1620	
24P4C12v.1:	1621	gctgtttcaagtgcctctgggtgctggaaaaatttatcaagttcctaaaccgcaatg	1680	
24P4C12v.8:	1621	gctgtttcaagtgcctctgggtgctggaaaaatttatcaagttcctaaaccgcaatg	1680	

TABLE LVII-continued

Nucleotide sequence alignment of 24P4C12v.1 v.1 (SEQ ID NO: 101) and 24P4C12 v.8 (SEQ ID NO: 102)				
24P4C12v.1: 1681	catacatcatgatcgccatctacgggaagaatttctgtgtctcagccaaaaatgcgttca	1740		
24P4C12v.8: 1681	catacatcatgatcgccatctacgggaagaatttctgtgtctcagccaaaaatgcgttca	1740		
24P4C12v.1: 1741	tgctactcatgcgaacattgtcagggtggtcgtcctggacaaagtcacagacctgctgc	1800		
24P4C12v.8: 1741	tgctactcatgcgaacattgtcagggtggtcgtcctggacaaagtcacagacctgctgc	1800		
24P4C12v.1: 1801	tgttctttgggaagctgctgggtggtcggaggcgtgggggtcctgtccttctttttttct	1860		
24P4C12v.8: 1801	tgttctttgggaagctgctgggtggtcggaggcgtgggggtcctgtccttctttttttct	1860		
24P4C12v.1: 1861	ccgggtcgcatcccggggctgggtaaagactttaagagccccacctaactattactggc	1920		
24P4C12v.8: 1861	ccgggtcgcatcccggggctgggtaaagactttaagagccccacctaactattactggc	1920		
24P4C12v.1: 1921	tgcccatcatga	1932		
24P4C12v.8: 1921	tgcccatcatga	1932		
Score = 1263 bits (657), Expect = 0.0Identities = 657/657 (100%) Strand = Plus/Plus				
24P4C12v.1: 1931	gacctccatcctgggggcctatgtcatcgccagcggtcttcttcagcgttttcggcatgtg	1990		
24P4C12v.8: 1967	gacctccatcctgggggcctatgtcatcgccagcggtcttcttcagcgttttcggcatgtg	2026		
24P4C12v.1: 1991	tgtggacacgctcttctctgcttcttggaagacctggagcggaacaacggctccctgga	2050		
24P4C12v.8: 2027	tgtggacacgctcttctctgcttcttggaagacctggagcggaacaacggctccctgga	2086		
24P4C12v.1: 2051	ccggccctactacatgtccaagagccttctaaagattctgggcaagaagaacgaggcgcc	2110		
24P4C12v.8: 2087	ccggccctactacatgtccaagagccttctaaagattctgggcaagaagaacgaggcgcc	2146		
24P4C12v.1: 2111	cccggaacaagaagaggaagaagtacagctccggccctgatccaggactgcacccca	2170		
24P4C12v.8: 2147	cccggaacaagaagaggaagaagtacagctccggccctgatccaggactgcacccca	2206		
24P4C12v.1: 2171	ccccaccgtccagccatccaacctcacttcgccttacagggtctccattttgtggtaaaa	2230		
24P4C12v.8: 2207	ccccaccgtccagccatccaacctcacttcgccttacagggtctccattttgtggtaaaa	2266		
24P4C12v.1: 2231	aaagggttttaggccaggcgccgtggctcacgcctgtaatccaacactttgagaggctgag	2290		
24P4C12v.8: 2267	aaagggttttaggccaggcgccgtggctcacgcctgtaatccaacactttgagaggctgag	2326		
24P4C12v.1: 2291	gcgggcggtacacctgagtcaggagttcgagaccagcctggccaacatggtgaaacctcc	2350		
24P4C12v.8: 2327	gcgggcggtacacctgagtcaggagttcgagaccagcctggccaacatggtgaaacctcc	2386		
24P4C12v.1: 2351	gtctctattaaaaatacaaaaattagccgagagtgggtggcatgcacctgtcatcccagct	2410		
24P4C12v.8: 2387	gtctctattaaaaatacaaaaattagccgagagtgggtggcatgcacctgtcatcccagct	2446		
24P4C12v.1: 2411	actcgggaggctgaggcaggagaatcgcttgaacccgggaggcagaggttgagtgagcc	2470		
24P4C12v.8: 2447	actcgggaggctgaggcaggagaatcgcttgaacccgggaggcagaggttgagtgagcc	2506		
24P4C12v.1: 2471	gagatcgcgccactgcactccaacctgggtgacagactctgtctccaaaacaaaacaaac	2530		
24P4C12v.8: 2507	gagatcgcgccactgcactccaacctgggtgacagactctgtctccaaaacaaaacaaac	2566		
24P4C12v.1: 2531	aaacaaaaagattttattaaagatattttgttaactcagtaaaaaaaaaaaaaaaaaaa	2587		
24P4C12v.8: 2567	aaacaaaaagattttattaaagatattttgttaactcagtaaaaaaaaaaaaaaaaaaa	2623		

TABLE LVIII

Peptide sequences of protein coded by 24P4C12 v.8 (SEQ ID NO: 103)	
MGGKQRDEDD EAYGKPVKYD PSFRGPIKNR SCTDVICCVL FLLFILGYIV VGIVAWLYGD	60
PRQVLYPRNS TGAYCGMGEN KDKPYLLYFN IFSCILSSNI ISVAENGLQC PTPQVCVSSC	120
PEDPWTVGKN EFSQTVGEVF YTKNRNFCPL GVPWNMTVIT SLQQELCPSF LLPSAPALGR	180
CFPWTNVTTP ALPGITNDTT IQQGISGLID SLNARDISVK IFEDFAQSWY WILVALGVAL	240
VLSLLFILLL RLVAGPLVLV LILGVLGVLA YGIYCWEEY RVLDRKGASI SQLGFTTNLS	300
AYQSQVETWL AALIVLAVLE AILLMLIFL RQRIRIAIAL LKEASKAVGQ MMSTMFYPLV	360
TFVLLLICIA YWAMTALYLA TSGQPQYVLW ASNISSPGCE KVPINTSCNP TAHLVNSSCP	420
GLMCVFQGY SSKGLIQRSVF NLQIYGVGL FWTLNWVLAL GQCVLAGAFA SFYWAFHKPQ	480
DIPTFPLISA FIRTLYHTG SLAFGALILT LVQIARVILE YIDHKLRGVQ NPVARCIMCC	540
FKCCLWCLEK FIKFLNRNAY IMIAIYGKNF CVSAKNAFML LMRNIVRVVV LDKVTDLLLF	600
FGKLLVVGGV GVLSFFFFSG RIPGLGKDFK SPHLNYYWLP IMRNPITPTG HVPQTSILGA	660
YVIASGFFSV FGMCVDTLFL CFLEDLERNN GSLDRPYYS KSLKILGKK NEAPPDNKKR	720
KK	722

TABLE LIX

Amino acid sequence alignment of 24P4C12v.1 v.1 (SEQ ID NO: 104) and 24P4C12 v.8 (SEQ ID NO: 105)	
Score = 1438 bits (3722), Expect = 0.0Identities = 710/722 (98%), Positives = 710/722 (98%), Gaps = 12/722 (1%)	
24P4C12v.1: 1MGGKQRDEDD EAYGKPVKYDPSFRGPIKNRSCTDVICCVL FLLFILGYIVVGIVAWLYGD	60
24P4C12v.8: 1MGGKQRDEDD EAYGKPVKYDPSFRGPIKNRSCTDVICCVL FLLFILGYIVVGIVAWLYGD	60
24P4C12v.1: 61PRQVLYPRNSTGAYCGMGENKDKPYLLYFNIFSCILSSNIISVAENGLQCPTPQVCVSSC	120
24P4C12v.8: 61PRQVLYPRNSTGAYCGMGENKDKPYLLYFNIFSCILSSNIISVAENGLQCPTPQVCVSSC	120
24P4C12v.1: 121PEDPWTVGKNEFSQTVGEVFYTKNRNFCPLPGVPWNMTVITSLQQELCPSFLLPSAPALGR	180
24P4C12v.8: 121PEDPWTVGKNEFSQTVGEVFYTKNRNFCPLPGVPWNMTVITSLQQELCPSFLLPSAPALGR	180
24P4C12v.1: 181CFPWTNVTTPPALPGITNDTTIQQGISGLIDSLNARDISVKI FEDFAQSWY WILVALGVAL	240
24P4C12v.8: 181CFPWTNVTTPPALPGITNDTTIQQGISGLIDSLNARDISVKI FEDFAQSWY WILVALGVAL	240
24P4C12v.1: 241VLSLLFILLRLVAGPLVLVLILGVLGVLAYGIYCWEEYRVLDRKGASISQLGFTTNLS	300
24P4C12v.8: 241VLSLLFILLRLVAGPLVLVLILGVLGVLAYGIYCWEEYRVLDRKGASISQLGFTTNLS	300
24P4C12v.1: 301AYQSQVETWLAALIVLAVLEAILLMLIFLRQRIRIAIAL LKEASKAVGQMMSTMFYPLV	360
24P4C12v.8: 301AYQSQVETWLAALIVLAVLEAILLMLIFLRQRIRIAIAL LKEASKAVGQMMSTMFYPLV	360
24P4C12v.1: 361TFVLLLICIA YWAMTALYLATSGQPQYVLWASNISSPGCEKVPINTSCNPTAHLVNSSCP	420
24P4C12v.8: 361TFVLLLICIA YWAMTALYLATSGQPQYVLWASNISSPGCEKVPINTSCNPTAHLVNSSCP	420
24P4C12v.1: 421GLMCVFQGYSSKGLIQRSVFNLQIYGVGLFWTLNWVLALGQCVLAGAFASFYWAFHKPQ	480
24P4C12v.8: 421GLMCVFQGYSSKGLIQRSVFNLQIYGVGLFWTLNWVLALGQCVLAGAFASFYWAFHKPQ	480
24P4C12v.1: 481DIPTFPLISAFIRTLYHTGSLAFGALILTLVQIARVILEYIDHKLRGVQNPVARCIMCC	540
24P4C12v.8: 481DIPTFPLISAFIRTLYHTGSLAFGALILTLVQIARVILEYIDHKLRGVQNPVARCIMCC	540
24P4C12v.1: 541FKCCLWCLEKFIKFLNRNAYIMIAIYGKNFCVSAKNAFMLLMRNIVRVVVLDDKVTDLLEF	600
24P4C12v.8: 541FKCCLWCLEKFIKFLNRNAYIMIAIYGKNFCVSAKNAFMLLMRNIVRVVVLDDKVTDLLEF	600
24P4C12v.1: 601FGKLLVVGGVGVLSFFFFSGRIPGLGKDFKSPHLNYYWLPIM-----TSILGA	648
FGKLLVVGGVGVLSFFFFSGRIPGLGKDFKSPHLNYYWLPIM	TSILGA

TABLE LIX-continued

Amino acid sequence alignment of 24P4C12v.1 v.1 (SEQ ID NO: 104) and 24P4C12 v.8 (SEQ ID NO: 105)		
24P4C12v.8: 601FGKLLVVGVGVLSTFFFSGRIPGLGKDFKSPHLNYYWLPIMRNPITPTGHVFQTSILGA		660
24P4C12v.1: 649YVIASGFFSVFGMCVDTLFLCFLEDLERNNGSLDRPYMSKSLKILGKKNEAPPDNKKR		708
YVIASGFFSVFGMCVDTLFLCFLEDLERNNGSLDRPYMSKSLKILGKKNEAPPDNKKR		
24P4C12v.8: 661YVIASGFFSVFGMCVDTLFLCFLEDLERNNGSLDRPYMSKSLKILGKKNEAPPDNKKR		720
24P4C12v.1: 709KK		710
KK		
24P4C12v.8: 721KK		722

TABLE LX

Nucleotide sequence of transcript variant 24P4C12 v.9 (SEQ ID NO: 106)	
gagccatggg gggaaagcag cgggacgagg atgacgaggc ctacgggaag ccagtcaa	60
acgacccctc ctttcgaggc cccatcaaga acagaagctg cacagatgtc atctgctg	120
tcctcttctc gctcttcatt ctaggttaca tcgtggtggg gattgtggcc tggttgat	180
gagacccccg gcaagtcctc taccacagga actctactgg ggcctactgt ggcattgg	240
agaacaaaga taagccgtat ctctgttact tcaacatctt cagctgcac	300
acatcatctc agttgctgag aacggcctac agtgccccac acccaggtg tgtgtg	360
cctgcccga ggacccatgg actgtggga aaaacagatt ctacagact gttgggga	420
tcttctatc aaaaaacagg aacttttgc tgccaggggt accctggaat atgacggt	480
tcacaagcct gcaacaggaa ctctgcccc gtttctctc cccctctgct ccagctct	540
ggcgtgctt tccatggacc aacgttactc caccggcgct cccagggatc accaatga	600
ccaccatata gcaggggac agcgggtctt ttgacagcct caatgcccga gacatcag	660
ttaaagatct tgaagatttt gccagtcct ggtattggat tctgttgcc ctgggggt	720
ctctggctct gaggcctact tttatcttgc ttctgcgct ggtggctggg cccctg	780
tggtgctgat cctgggagtg ctgggcgtgc tggcatacgg catctactac tctggg	840
agtaccgagt gctgcccga aagggcgct ccactctcca gctgggttc accacca	900
tcagtgccta ccagagcgtg caggagacct ggctggccgc cctgatcgtg ttggcg	960
ttgaagccat cctgctgctg atgctcatct tctgcccga gcggattcgt attgccat	1020
ccctcctgaa ggaggccagc aaggctgtgg gacagatgat gtctaccatg ttctacc	1080
tggtcacctt tgcctctc ctcactgca ttgectactg ggcatgact gctctgtat	1140
ctctgcccac gcagccagcc actcttggat atgtgctctg ggcatccaac atcagct	1200
ccggctgtga gaaagtgcata ataaatacat catgcaacc caggcccac cttgtga	1260
cctctgccc agggctgatg tgcgtcttcc agggctactc atccaaggc ctaatcca	1320
gttctgtctt caatctgcaa atctatggg tcttggggct cttctggacc cttactg	1380
tactggccct gggccaatgc gtctctgctg gagectttgc ctctctctac tgggct	1440
acaagcccca ggacatccct accttccct taatctctgc cttcatccgc acactcc	1500
accacactgg gtcattggca tttggagccc tcatcctgac cctgtgacg atagccc	1560
tcatcttga gtatattgac cacaagctca gaggagtga gaacctgta gcccgctg	1620
tcatgtgctg tttcaagtgc tgccctgggt gtctggaaaa atttatcaag ttctaa	1680
gcaatgcata catcatgac gccatctacg ggaagaattt ctgtgtctca gcaaaa	1740
cgttcatgct actcatgca aacattgtca ggggtgctg cctggacaaa gtcacag	1800

TABLE LX-continued

Nucleotide sequence of transcript variant 24P4C12 v.9 (SEQ ID NO: 106)		
tgctgctggt ctttggaag ctgctgggtg tcggaggcgt ggggggtcctg tccttctttt	1860	
ttttctccgg tcgcatcccg gggctgggta aagactttaa gagccccac ctcaactatt	1920	
actggctgcc catcatgacc tccatcctgg gggcctatgt catcgccagc ggcttcttca	1980	
gcgttttcgg catgtgtgtg gacacgctct tcctctgctt cctggaagac ctggagcgga	2040	
acaacggctc cctggaccgg ccctactaca tgtccaagag ccttctaaag attctgggca	2100	
agaagaacga ggcgcccccg gacaacaaga agaggaagaa gtgacagctc cggccctgat	2160	
ccaggactgc accccacccc caccgtccag ccatccaacc tcaactcgcc ttacaggtct	2220	
ccattttgtg gtaaaaaaag gttttaggcc aggcgcctg gctcacgcct gtaatccaac	2280	
actttgagag gctgaggcgg gcggatcacc tgagtcagga gttcgagacc agcctggcca	2340	
acatggtgaa acctccgtct ctattaaaaa tacaaaaatt agccgagagt ggtggcatgc	2400	
acctgtcatc ccagctactc gggaggctga ggcaggagaa tcgcttgaac ccgggaggca	2460	
gagggttcag tgagccgaga tcgcgccact gcactccaac ctgggtgaca gactctgtct	2520	
ccaaaacaaa acaaacaaa aaaaagattt tattaagat attttgtaa ctcagtaaaa	2580	
aaaaaaaaaa aaa	2593	

TABLE LXI

Nucleotide sequence alignment of 24P4C12v.1 (SEQ ID NO: 107) and 24P4C12 v.9 (SEQ ID NO: 108)		
Score = 2188 bits (1138), Expect = 0.0Identites = 1138/1138 (100%) Strand = Plus/Plus		
24P4C12v.1: 1	gagccatgggggaaagcagcgggacgaggatgacgaggcctacgggaagccagtcaaat	60
24P4C12v.9: 1	gagccatgggggaaagcagcgggacgaggatgacgaggcctacgggaagccagtcaaat	60
24P4C12v.1: 61	acgacccctcctttcgaggcccatcaagaacagaagctgcacagatgtcatctgctgcg	120
24P4C12v.9: 61	acgacccctcctttcgaggcccatcaagaacagaagctgcacagatgtcatctgctgcg	120
24P4C12v.1: 121	tcctcttctgctcttctattctaggttacatcggtggggattgtggcctggtgtatg	180
24P4C12v.9: 121	tcctcttctgctcttctattctaggttacatcggtggggattgtggcctggtgtatg	180
24P4C12v.1: 181	gagaccccggaagtcctctaccccaggaactctactggggcctactgtggcatggggg	240
24P4C12v.9: 181	gagaccccggaagtcctctaccccaggaactctactggggcctactgtggcatggggg	240
24P4C12v.1: 241	agaacaaagataagccgtatctcctgtacttcaacatcttcagctgcatcctgtccagca	300
24P4C12v.9: 241	agaacaaagataagccgtatctcctgtacttcaacatcttcagctgcatcctgtccagca	300
24P4C12v.1: 301	acatcatctcagttgctgagaacggcctacagtgcacacacccaggtgtgtgtcct	360
24P4C12v.9: 301	acatcatctcagttgctgagaacggcctacagtgcacacacccaggtgtgtgtcct	360
24P4C12v.1: 361	cctgcccggaggaccatggactgtgggaaaaaacgagttctcacagactgttggggaag	420
24P4C12v.9: 361	cctgcccggaggaccatggactgtgggaaaaaacgagttctcacagactgttggggaag	420
24P4C12v.1: 421	tcttctatacaaaaaacaggaacttttgtctgccaggggtaccctggaatatgacggtga	480
24P4C12v.9: 421	tcttctatacaaaaaacaggaacttttgtctgccaggggtaccctggaatatgacggtga	480
24P4C12v.1: 481	tcacaagcctgcaacaggaactctgccccagtttctcctccctctgctccagctctgg	540
24P4C12v.9: 481	tcacaagcctgcaacaggaactctgccccagtttctcctccctctgctccagctctgg	540

TABLE LXI-continued

Nucleotide sequence alignment of 24P4C12v.1 v.1 (SEQ ID NO: 107) and 24P4C12 v.9 (SEQ ID NO: 108)				
24P4C12v.1:	541	ggcgctgctttccatggaccaacgttactccaccggcgctcccagggatcaccaatgaca	600	
24P4C12v.9:	541	ggcgctgctttccatggaccaacgttactccaccggcgctcccagggatcaccaatgaca	600	
24P4C12v.1:	601	ccaccatacagcaggggatcagcgggtcttattgacagcctcaatgccgagacatcagtg	660	
24P4C12v.9:	601	ccaccatacagcaggggatcagcgggtcttattgacagcctcaatgccgagacatcagtg	660	
24P4C12v.1:	661	ttaagatctttgaagattttgccagtcctggatttggattcttgttgccctgggggtgg	720	
24P4C12v.9:	661	ttaagatctttgaagattttgccagtcctggatttggattcttgttgccctgggggtgg	720	
24P4C12v.1:	721	ctctgggtcttgagcctactgtttatcttgcctctgcgcctgggtggctgggcccctgggtgc	780	
24P4C12v.9:	721	ctctgggtcttgagcctactgtttatcttgcctctgcgcctgggtggctgggcccctgggtgc	780	
24P4C12v.1:	781	tggtgctgatcctgggagtgtggcgctgctggcatacggcatctactactgctgggagg	840	
24P4C12v.9:	781	tggtgctgatcctgggagtgtggcgctgctggcatacggcatctactactgctgggagg	840	
24P4C12v.1:	841	agtaccgagtgtgcgggacaagggcgccctccatctcccagctgggtttcaccaccaacc	900	
24P4C12v.9:	841	agtaccgagtgtgcgggacaagggcgccctccatctcccagctgggtttcaccaccaacc	900	
24P4C12v.1:	901	tcagtgccctaccagagcgtgcaggagacctggctggcgccctgatcgtgttggcggtgc	960	
24P4C12v.9:	901	tcagtgccctaccagagcgtgcaggagacctggctggcgccctgatcgtgttggcggtgc	960	
24P4C12v.1:	961	ttgaagccatcctgctgctgatgctcatcttctgcggcagcggattcgtattgccatcg	1020	
24P4C12v.9:	961	ttgaagccatcctgctgctgatgctcatcttctgcggcagcggattcgtattgccatcg	1020	
24P4C12v.1:	1021	ccctcctgaaggaggccagcaaggctgtgggacagatgatgtctaccatgttctaccac	1080	
24P4C12v.9:	1021	ccctcctgaaggaggccagcaaggctgtgggacagatgatgtctaccatgttctaccac	1080	
24P4C12v.1:	1081	tggtcaacctttgtcctcctcctcatctgcattgcctactgggccatgactgctctgta	1138	
24P4C12v.9:	1081	tggtcaacctttgtcctcctcctcatctgcattgcctactgggccatgactgctctgta	1138	
Score = 2738 bits (1424), Expect = 0.0Identities = 1424/1424 (100%) Strand = Plus/Plus				
24P4C12v.1:	1164	tatgtgctctgggcatccaacatcagctccccggctgtgagaaagtgccataaataca	1223	
24P4C12v.9:	1170	tatgtgctctgggcatccaacatcagctccccggctgtgagaaagtgccataaataca	1229	
24P4C12v.1:	1224	tcatgcaacccccagggcccaccttgtgaactcctcgctgccagggtgatgtgcgtcttc	1283	
24P4C12v.9:	1230	tcatgcaacccccagggcccaccttgtgaactcctcgctgccagggtgatgtgcgtcttc	1289	
24P4C12v.1:	1284	cagggtactcatccaaggcctaataccaacgttctgtcttcaatctgcaaatctatggg	1343	
24P4C12v.9:	1290	cagggtactcatccaaggcctaataccaacgttctgtcttcaatctgcaaatctatggg	1349	
24P4C12v.1:	1344	gtcctggggctcttctggacccttaactgggtactggccctgggccaatgcgtcctcgct	1403	
24P4C12v.9:	1350	gtcctggggctcttctggacccttaactgggtactggccctgggccaatgcgtcctcgct	1409	
24P4C12v.1:	1404	ggagcctttgctccttctactgggccttcacaaagccccaggacatccctaccttcccc	1463	
24P4C12v.9:	1410	ggagcctttgctccttctactgggccttcacaaagccccaggacatccctaccttcccc	1469	
24P4C12v.1:	1464	ttaatctctgccttcatccgcacactcogttaaccacactgggtcattggcatttggagcc	1523	
24P4C12v.9:	1470	ttaatctctgccttcatccgcacactcogttaaccacactgggtcattggcatttggagcc	1529	

TABLE LXI-continued

Nucleotide sequence alignment of 24P4C12v.1 v.1 (SEQ ID NO: 107) and 24P4C12 v.9 (SEQ ID NO: 108)				
24P4C12v.1:	1524	ctcatcctgacccttgtgcagatagccccgggtcatcttggagtatttgaccacaagctc	1583	
24P4C12v.9:	1530	ctcatcctgacccttgtgcagatagccccgggtcatcttggagtatttgaccacaagctc	1589	
24P4C12v.1:	1584	agaggagtgcagaacctgtagcccgctgcacatgtgctgtttcaagtgtgcctctgg	1643	
24P4C12v.9:	1590	agaggagtgcagaacctgtagcccgctgcacatgtgctgtttcaagtgtgcctctgg	1649	
24P4C12v.1:	1644	tgtctggaataatttatcaagttcctaaaccgcaatgcatacatcatgatcgccatctac	1703	
24P4C12v.9:	1650	tgtctggaataatttatcaagttcctaaaccgcaatgcatacatcatgatcgccatctac	1709	
24P4C12v.1:	1704	gggaagaatttctgtgtctcagccaaaaatgcgttcatgctactcatgcgaacattgtc	1763	
24P4C12v.9:	1710	gggaagaatttctgtgtctcagccaaaaatgcgttcatgctactcatgcgaacattgtc	1769	
24P4C12v.1:	1764	aggggtggtcgtcctggacaaagtcacagacctgctgctgttcttgggaagctgctggg	1823	
24P4C12v.9:	1770	aggggtggtcgtcctggacaaagtcacagacctgctgctgttcttgggaagctgctggg	1829	
24P4C12v.1:	1824	gtcggaggcgtgggggtcctgtccttcttttttctccgggtcgcatcccggggctgggt	1883	
24P4C12v.9:	1830	gtcggaggcgtgggggtcctgtccttcttttttctccgggtcgcatcccggggctgggt	1889	
24P4C12v.1:	1884	aaagactttaagagccccacctcaactattactggctgcccatcatgacctccatcctg	1943	
24P4C12v.9:	1890	aaagactttaagagccccacctcaactattactggctgcccatcatgacctccatcctg	1949	
24P4C12v.1:	1944	ggggcctatgtcatcgccagcggtcttctcagcggtttcggcattgtgtggacacgctc	2003	
24P4C12v.9:	1950	ggggcctatgtcatcgccagcggtcttctcagcggtttcggcattgtgtggacacgctc	2009	
24P4C12v.1:	2004	ttcctctgcttctctggaagacctggagcgggaacaacgggtccctggaccggccctactac	2063	
24P4C12v.9:	2010	ttcctctgcttctctggaagacctggagcgggaacaacgggtccctggaccggccctactac	2069	
24P4C12v.1:	2064	atgtccaagagccttctaaagattctgggcaagaagaacgaggcgccccggacaacaag	2123	
24P4C12v.9:	2070	atgtccaagagccttctaaagattctgggcaagaagaacgaggcgccccggacaacaag	2129	
24P4C12v.1:	2124	aagaggaagaagtgcacagctccggccctgatccaggactgcacccacccccaccgtcca	2183	
24P4C12v.9:	2130	aagaggaagaagtgcacagctccggccctgatccaggactgcacccacccccaccgtcca	2189	
24P4C12v.1:	2184	gccatccaacctcacttcgccttacagggtctccattttgtggtaaaaaagggttttaggc	2243	
24P4C12v.9:	2190	gccatccaacctcacttcgccttacagggtctccattttgtggtaaaaaagggttttaggc	2249	
24P4C12v.1:	2244	caggcgccgtgggtcacgcctgtaatccaacactttgagaggctgaggcggggcggtcac	2303	
24P4C12v.9:	2250	caggcgccgtgggtcacgcctgtaatccaacactttgagaggctgaggcggggcggtcac	2309	
24P4C12v.1:	2304	ctgagtcaggagttcgagaccagcctggccaacatggtgaaacctccgtctctattaaaa	2363	
24P4C12v.9:	2310	ctgagtcaggagttcgagaccagcctggccaacatggtgaaacctccgtctctattaaaa	2369	
24P4C12v.1:	2364	atacaaaaattagccgagagtgggtggcatgcacctgtcatcccagctactcgggaggctg	2423	
24P4C12v.9:	2370	atacaaaaattagccgagagtgggtggcatgcacctgtcatcccagctactcgggaggctg	2429	
24P4C12v.1:	2424	aggcaggagaatcgcttgaacccgggagggcagaggttgacagtgcagccgagatcgcgccac	2483	
24P4C12v.9:	2430	aggcaggagaatcgcttgaacccgggagggcagaggttgacagtgcagccgagatcgcgccac	2489	
24P4C12v.1:	2484	tgcactccaacctgggtgacagactctgtctccaaaacaaaacaaacaaaaaagatt	2543	
24P4C12v.9:	2490	tgcactccaacctgggtgacagactctgtctccaaaacaaaacaaacaaaaaagatt	2549	
24P4C12v.1:	2544	ttatttaaagatatatttggtaactcagtaaaaaaaaaaaaaaaaaa	2587	
24P4C12v.9:	2550	ttatttaaagatatatttggtaactcagtaaaaaaaaaaaaaaaaaa	2593	

TABLE LXII

Peptide sequences of protein coded by 24P4C12 v.9 (SEQ ID NO: 109)	
MGGKQRDEDD EAYGKPVKYD PSFRGPIKNR SCTDVICCVL FLLFILGYIV VGIVAWLYGD	60
PRQVLYPRNS TGAYCGMGEN KDKPYLLYFN IFSCILSSNI ISVAENGLQC PTPQVCVSSC	120
PEDPWTVGKN EFSQTVGEVF YTKNRNFCPL GVPWNMTVIT SLQQELCPSF LLPSAPALGR	180
CFPWTNVTTP ALPGITNDTT IQQGISGLID SLNARDISVK IFEDFAQSWY WILVALGVAL	240
VLSLLFILLL RLVAGPLVLV LILGVLGVLA YGIYYCWEEY RVLDRKGASI SQLGFTTNLS	300
AYQSVQETWL AALIVLAVLE AILLMLIFL RQRIRIAIAL LKEASKAVGQ MMSTMFYPLV	360
TFVLLLICIA YWAMTALYPL PTQPATLGYV LWASNISSPG CEKVPINTSC NPTAHLVNSS	420
CPGLMCFVQG YSSKGLIQRS VFNLQIYGV LGLFWTLNWV ALGQCVLAGA FASFYWAFHK	480
PQDIPTFPLI SAFIRTLRYH TGSALFGALI LTLVQIARVI LEYIDHKLRG VQNPVARCIM	540
CCFKCCLWCL EKFIFLNRN AYIMIAIYK NFCVSAKNAF MLLMRNIVRV VVLDKVTDL	600
LLFPGKLLVVG GVGVLSPFFF SGRIPGLGKD FKSPHLNYYW LPIMTSILGA YVIASGFFSV	660
PGMCVDTLFL CFLEDLERNN GSLDRPYMS KSLKILGKK NEAPPDNKKR KK	712

TABLE LXIII

Amino acid sequence alignment of 24P4C12v.1 v.1 (SEQ ID NO: 110) and 24P4C12 v.9 (SEQ ID NO: 111)	
Score = 1424 bits (3686), Expect = 0.0Identities = 704/713 (98%), Positives = 705/713 (98%), Gaps = 4/713 (0%)	
24P4C12v.1: 1MGGKQRDEDD EAYGKPVKYDPSFRGPIKNRSCDVI CCVLFLLFILGYIVVGIVAWLYGD	60
24P4C12v.9: 1MGGKQRDEDD EAYGKPVKYDPSFRGPIKNRSCDVI CCVLFLLFILGYIVVGIVAWLYGD	60
24P4C12v.1: 61PRQVLYPRNSTGAYCGMGENKDKPYLLYFNIFSCILSSNIISVAENGLQCPTPQVCVSSC	120
24P4C12v.9: 61PRQVLYPRNSTGAYCGMGENKDKPYLLYFNIFSCILSSNIISVAENGLQCPTPQVCVSSC	120
24P4C12v.1: 121PEDPWTVGKNEFSQTVGEVFYTKNRNFCPLPGVPWNMTVITSLQQELCPSFLLPSAPALGR	180
24P4C12v.9: 121PEDPWTVGKNEFSQTVGEVFYTKNRNFCPLPGVPWNMTVITSLQQELCPSFLLPSAPALGR	180
24P4C12v.1: 181CFPWTNVTTPPALPGITNDTTIQQGISGLIDSLNARDISVKIFEDFAQSWYWILVALGVAL	240
24P4C12v.9: 181CFPWTNVTTPPALPGITNDTTIQQGISGLIDSLNARDISVKIFEDFAQSWYWILVALGVAL	240
24P4C12v.1: 241VLSLLFILLRLVAGPLVLVLILGVLGVLAYGIYYCWEEYRVLDRKGASISQLGFTTNLS	300
24P4C12v.9: 241VLSLLFILLRLVAGPLVLVLILGVLGVLAYGIYYCWEEYRVLDRKGASISQLGFTTNLS	300
24P4C12v.1: 301AYQSVQETWLAALIVLAVLEAILLMLIFLRQRIRIAIALKEASKAVGQMMSTMFYPLV	360
24P4C12v.9: 301AYQSVQETWLAALIVLAVLEAILLMLIFLRQRIRIAIALKEASKAVGQMMSTMFYPLV	360
24P4C12v.1: 361TFVLLLICIA YWAMTALYLATSGQPQ---YVLWASNISSPGCEKVPINTSCNPTAHLVNS	417
24P4C12v.9: 361TFVLLLICIA YWAMTALYPLPT-QPATLGYVLWASNISSPGCEKVPINTSCNPTAHLVNS	419
24P4C12v.1: 418SCPGLMCFVQGYSSKGLIQRSVFNLQIYGV LGLFWTLNWV LALGQCVLAGAFASFYWAFH	477
24P4C12v.9: 420SCPGLMCFVQGYSSKGLIQRSVFNLQIYGV LGLFWTLNWV LALGQCVLAGAFASFYWAFH	479
24P4C12v.1: 478KPQDIPTFPLISAFIRTLRYHTGSLAFGALILTLVQIARVILEYIDHKLRGVQNPVARI	537
24P4C12v.9: 480KPQDIPTFPLISAFIRTLRYHTGSLAFGALILTLVQIARVILEYIDHKLRGVQNPVARI	539
24P4C12v.1: 538MCCFKCCLWCLEKFIKFLNRNAYIMIAIYGNFCVSAKNAFMLLMRNIVRVVLDKVTDL	597
24P4C12v.9: 540MCCFKCCLWCLEKFIKFLNRNAYIMIAIYGNFCVSAKNAFMLLMRNIVRVVLDKVTDL	599
24P4C12v.1: 598LLFPGKLLVVG VGVLSPFFFSGRIPGLGKDFKSPHLNYYWLPIMTSILGAYVIASGFFS	657
24P4C12v.9: 600LLFPGKLLVVG VGVLSPFFFSGRIPGLGKDFKSPHLNYYWLPIMTSILGAYVIASGFFS	659

Figure 2I-2

300 S A Y Q S V Q E T W L A A L I V L A V L
901 TCAGTGCCTACCAGAGCGTGCAGGAGACCTGGCTGGCCGCCCTGATCGTGTGGCGGTGC
320 E A I L L L M L I F L R Q R I R I A I A
961 TTGAAGCCATCCTGCTGCTGATGCTCATCTTCCTGCGGCAGCGGATTTCGTATTGCCATCG
340 L L K E A S K A V G Q M M S T M F Y P L
1021 CCCTCCTGAAGGAGGCCAGCAAGGCTGTGGGACAGATGATGTCTACCATGTTCTACCCAC
360 V T F V L L L I C I A Y W A M T A L Y P
1081 TGGTCACCTTTGTCTCCTCCTCATCTGCATTGCCTACTGGGCCATGACTGCTCTGTATC
380 L P T Q P A T L G Y V L W A S N I S S P
1141 CTCTGCCCACGCAGCCAGCCACTCTTGATATGTGCTCTGGGCATCCAACATCAGTCCCC
400 G C E K V F I N I S C N P T A H L V N S
1201 CCGGCTGTGAGAAAGTGCCAATAAATACATCATGAACCCACGGCCACCTTGTGAAC
420 S C P G L M C V F Q G Y S S K G L I Q R
1261 CCTCGTGCCAGGGCTGATGTGCGTCTTCCAGGGCTACTCATCAAAGGCCTAATCCAAC
440 S V F N L Q I Y G V L G L F W T L N W V
1321 GTTCTGTCTTCAATCTGCAAATCTATGGGGTCTGGGGCTCTTCTGGACCCCTAACTGGG
460 L A L G Q C V L A G A F A S F Y W A F H
1381 TACTGGCCCTGGGCCAATGCGTCTCCTGCTGGAGCCTTTGCCCTCCTTCTACTGGGCCTTCC
480 K P Q D I P T F F L I S A F I R T L R Y
1441 ACAAGCCCCAGGACATCCCTACCTTCCCTTAAATCTCTGCCTTCATCCGCACACTCCGTT
500 H T G S L A F G A L I L T L V Q I A R V
1501 ACCACACTGGGTCATTGGCATTGGAGCCCTCATCCTGACCCTTGTGCAGATAGCCCCGGG
520 I L E Y I D H K L R G V Q N P V A R C I
1561 TCATCTTGGAGTATATTGACCACAAGCTCAGAGGAGTGCAGAACCCTGTAGCCCGCTGCA
540 M C C F K C C L W C L E K F I K F L N R
1621 TCATGTGCTGTTTCAAGTGTGCTGCTGCTGGTGTCTGGAAAAATTTATCAAGTTCTTAAACC
560 N A Y I M I A I Y G K N F C V S A K N A
1681 GCAATGCATACATCATGATCGCCATCTACGGGAAGAATTTCTGTGTCTCAGCCAAAAATG
580 F M L L M R N I V R V V V L D R V T D L
1741 CGTTCATGCTACTCATGCGAAACATTGTCAGGGTGGTCTGCTGGACAAAGTCACAGACC
600 L L F F G K L L V V G G V G V L S F F F
1801 TGCTGCTGTTCTTTGGGAAGCTGCTGGTGGTCTGGAGGCGTGGGGTCTGTCTTCTTTT
620 F S G R I P G L G K D F K S P H L N Y Y
1861 TTTTCTCCGGTCGCATCCCGGGGCTGGGTAAAGACTTTAAGAGCCCCACCTCAACTATT
640 W L P I M T S I L G A Y V I A S G F F S
1921 ACTGGCTGCCCATCATGACCTCCATCCTGGGGGCTATGTCATCGCCAGCGGCTTCTTCA
660 V F G M C V D T L F L C F L E D L E R N
1981 GCGTTTTTCGGCATGTGTGTGGACACGCTCTTCTCTGCTTCTGGAAGACCTGGAGCGGA
680 N G S L D R P Y Y M S K S L L R I L G K
2041 ACAACGGCTCCCTGGACCGGCCCTACTACATGTCCAAGAGCCTTCTAAAGATTCTGGGCA

TABLE LXIII-continued

Amino acid sequence alignment of 24P4C12v.1 v.1 (SEQ ID NO: 110) and 24P4C12 v.9 (SEQ ID NO: 111)	
24P4C12v.1: 658VFGMCVDTLFLCFLEDLERNNGSLDRPYMSKSLKILGKKNEAPPDNKKRKK	710
VFGMCVDTLFLCFLEDLERNNGSLDRPYMSKSLKILGKKNEAPPDNKKRKK	
24P4C12v.9: 660VFGMCVDTLFLCFLEDLERNNGSLDRPYMSKSLKILGKKNEAPPDNKKRKK	712

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 112

<210> SEQ ID NO 1

<211> LENGTH: 160

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 1

gatcagggcg gccagccagg tctcctgcac gctctggtag gcaactgaggt tgggtgtgaa 60
 acccagctgg gagatggagg cgccctcgtc ccgcagcact cgggtactcct cccagcagta 120
 gtagatgccca tatgccagca cgcccagcac tcccaggatc 160

<210> SEQ ID NO 2

<211> LENGTH: 2587

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 2

gagccatggg gggaaagcag cgggacgagg atgacgaggc ctacgggaag ccagtcaaat 60
 acgacccctc ctttcgaggc cccatcaaga acagaagctg cacagatgtc atctgtctgcg 120
 tcctcttctc gctcttcatt ctagggtaca tcgtggtggg gattgtggcc tggttgtatg 180
 gagacccccg gcaagtcctc taccacagga actctactgg ggcctactgt ggcatggggg 240
 agaacaaaga taagccgat ctctgtact tcaacatctt cagctgcac ctgtccagca 300
 acatcatctc agttgtctgag aacggcctac agtgccccac acccaggtg tgtgtgtcct 360
 cctgcccga ggacccatgg actgtggga aaaacgagtt ctcacagact gttggggaag 420
 tcttctatac aaaaaacagg aacttttgc tgccaggggt accctggaat atgacggtga 480
 tcacaagcct gcaacaggaa ctctgcccc gtttctctc cccctctgct ccagctctgg 540
 ggcgtctgct tccatggacc aacgttactc caccggcgt cccagggatc accaatgaca 600
 ccaccatata gcaggggatc agcgggtctta ttgacagcct caatgcccga gacatcagtg 660
 ttaagatctt tgaagatctt gccagtcct ggtattggat tcttgttgcc ctgggggtgg 720
 ctctggtctt gagcctactg tttatcttgc ttctgcgcct ggtggctggg ccctcggtgc 780
 tgggtgctgat cctgggagtg ctgggcgtgc tggcatacgg catctactac tgctgggagg 840
 agtaccgagt gctgctggac aagggcgcct ccatctccca gctgggttcc accaccaacc 900
 tcagtgccta ccagagcgtg caggagacct ggctggccgc cctgatcgtg ttggcggtgc 960
 ttgaagccat cctgctgctg atgctcatct tcctgcggca gcggattcgt attgccatcg 1020
 ccctctgaa ggaggccagc aaggctgtgg gacagatgat gtctaccatg ttctaccac 1080
 tggtcacott tgcctcctc ctcatctgca ttgcctactg ggcatgact gctctgtacc 1140
 tggctacatc ggggcaacct cagtatgtgc tctgggcata caacatcagc tccccgggt 1200
 gtgagaaagt gccaataaat acatcatgca accccacggc ccacctgtg aactcctcgt 1260

-continued

```

gcccagggct gatgtgcgtc ttccagggct actcaccacaa aggcctaate caacgttctg 1320
tcttcaatct gcaaatctat ggggtcctgg ggctcttctg gaccttaac tgggtactgg 1380
ccctgggcca atgcgtctct gctggagcct ttgcctcctt ctactgggccc ttccacaagc 1440
cccaggacat cctacacctc cccttaatct ctgccttcat ccgcacactc cgttaccaca 1500
ctgggtcatt ggcatttga gccctcatcc tgaccttgt gcagatagcc cgggtcatct 1560
tggagtatat tgaccacaag ctacagaggag tgcagaaccc ttagaccgcg tgcacatgt 1620
gctgtttcaa gtgcgtcctc tgggtgtctgg aaaaatttat caagttccta aaccgcaatg 1680
catacatcat gatcgccatc tacgggaaga atttctgtgt ctacgcaaaa aatgcgttca 1740
tgctactcat gcgaaacatt gtcagggtgg tcgtcctgga caaagtcaca gacctgctgc 1800
tgttctttgg gaagctgctg tgggtcggag gcgtgggggt cctgtccttc tttttttct 1860
ccggtcgcac cccggggctg ggtaaagact ttaagagccc ccacctcaac tattactggc 1920
tgcccatcat gacctccatc ctgggggctc atgtcatcgc cagcggttc ttcagcgttt 1980
tcggcatgtg tgtggacacg ctcttctctt gcttcctgga agacctggag cggaacaacg 2040
gctccctgga ccggccctac tacatgtcca agagccttct aaagattctg ggcaagaaga 2100
acgaggcgcc cccggacaac aagaagagga agaagtgaac gtcgggccc tgatccagga 2160
ctgcacccca cccccaccgt ccagccatcc aacctcactt cgccttacag gtctccattt 2220
tgtggtaaaa aaagggttta ggccaggcgc cgtggctcac gcctgtaac caacactttg 2280
agaggctgag gcggggcgat cacctgagtc aggagtctga gaccagctg gccaacatgg 2340
tgaaacctcc gtctctatta aaaatacaaa aattagccga gagtggtggc atgcacctgt 2400
catcccagct actcgggagg ctgaggcagg agaatcgctt gaacccggga ggcagagggt 2460
gcagtgaacc gagatcgcc cactgcactc caacctgggt gacagactct gtctccaaaa 2520
caaaacaaac aaacaaaaag attttattaa agatatattt ttaactcagt aaaaaaaaaa 2580
aaaaaaa 2587

```

```

<210> SEQ ID NO 3
<211> LENGTH: 710
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 3

```

```

Met Gly Gly Lys Gln Arg Asp Glu Asp Asp Glu Ala Tyr Gly Lys Pro
 1           5           10          15
Val Lys Tyr Asp Pro Ser Phe Arg Gly Pro Ile Lys Asn Arg Ser Cys
 20          25          30
Thr Asp Val Ile Cys Cys Val Leu Phe Leu Leu Phe Ile Leu Gly Tyr
 35          40          45
Ile Val Val Gly Ile Val Ala Trp Leu Tyr Gly Asp Pro Arg Gln Val
 50          55          60
Leu Tyr Pro Arg Asn Ser Thr Gly Ala Tyr Cys Gly Met Gly Glu Asn
 65          70          75          80
Lys Asp Lys Pro Tyr Leu Leu Tyr Phe Asn Ile Phe Ser Cys Ile Leu
 85          90          95
Ser Ser Asn Ile Ile Ser Val Ala Glu Asn Gly Leu Gln Cys Pro Thr
100         105         110
Pro Gln Val Cys Val Ser Ser Cys Pro Glu Asp Pro Trp Thr Val Gly
115         120         125
Lys Asn Glu Phe Ser Gln Thr Val Gly Glu Val Phe Tyr Thr Lys Asn
130         135         140

```

-continued

Arg	Asn	Phe	Cys	Leu	Pro	Gly	Val	Pro	Trp	Asn	Met	Thr	Val	Ile	Thr	145	150	155	160
Ser	Leu	Gln	Gln	Glu	Leu	Cys	Pro	Ser	Phe	Leu	Leu	Pro	Ser	Ala	Pro	165	170	175	
Ala	Leu	Gly	Arg	Cys	Phe	Pro	Trp	Thr	Asn	Val	Thr	Pro	Pro	Ala	Leu	180	185	190	
Pro	Gly	Ile	Thr	Asn	Asp	Thr	Thr	Ile	Gln	Gln	Gly	Ile	Ser	Gly	Leu	195	200	205	
Ile	Asp	Ser	Leu	Asn	Ala	Arg	Asp	Ile	Ser	Val	Lys	Ile	Phe	Glu	Asp	210	215	220	
Phe	Ala	Gln	Ser	Trp	Tyr	Trp	Ile	Leu	Val	Ala	Leu	Gly	Val	Ala	Leu	225	230	235	240
Val	Leu	Ser	Leu	Leu	Phe	Ile	Leu	Leu	Leu	Arg	Leu	Val	Ala	Gly	Pro	245	250	255	
Leu	Val	Leu	Val	Leu	Ile	Leu	Gly	Val	Leu	Gly	Val	Leu	Ala	Tyr	Gly	260	265	270	
Ile	Tyr	Tyr	Cys	Trp	Glu	Glu	Tyr	Arg	Val	Leu	Arg	Asp	Lys	Gly	Ala	275	280	285	
Ser	Ile	Ser	Gln	Leu	Gly	Phe	Thr	Thr	Asn	Leu	Ser	Ala	Tyr	Gln	Ser	290	295	300	
Val	Gln	Glu	Thr	Trp	Leu	Ala	Ala	Leu	Ile	Val	Leu	Ala	Val	Leu	Glu	305	310	315	320
Ala	Ile	Leu	Leu	Leu	Met	Leu	Ile	Phe	Leu	Arg	Gln	Arg	Ile	Arg	Ile	325	330	335	
Ala	Ile	Ala	Leu	Leu	Lys	Glu	Ala	Ser	Lys	Ala	Val	Gly	Gln	Met	Met	340	345	350	
Ser	Thr	Met	Phe	Tyr	Pro	Leu	Val	Thr	Phe	Val	Leu	Leu	Leu	Ile	Cys	355	360	365	
Ile	Ala	Tyr	Trp	Ala	Met	Thr	Ala	Leu	Tyr	Leu	Ala	Thr	Ser	Gly	Gln	370	375	380	
Pro	Gln	Tyr	Val	Leu	Trp	Ala	Ser	Asn	Ile	Ser	Ser	Pro	Gly	Cys	Glu	385	390	395	400
Lys	Val	Pro	Ile	Asn	Thr	Ser	Cys	Asn	Pro	Thr	Ala	His	Leu	Val	Asn	405	410	415	
Ser	Ser	Cys	Pro	Gly	Leu	Met	Cys	Val	Phe	Gln	Gly	Tyr	Ser	Ser	Lys	420	425	430	
Gly	Leu	Ile	Gln	Arg	Ser	Val	Phe	Asn	Leu	Gln	Ile	Tyr	Gly	Val	Leu	435	440	445	
Gly	Leu	Phe	Trp	Thr	Leu	Asn	Trp	Val	Leu	Ala	Leu	Gly	Gln	Cys	Val	450	455	460	
Leu	Ala	Gly	Ala	Phe	Ala	Ser	Phe	Tyr	Trp	Ala	Phe	His	Lys	Pro	Gln	465	470	475	480
Asp	Ile	Pro	Thr	Phe	Pro	Leu	Ile	Ser	Ala	Phe	Ile	Arg	Thr	Leu	Arg	485	490	495	
Tyr	His	Thr	Gly	Ser	Leu	Ala	Phe	Gly	Ala	Leu	Ile	Leu	Thr	Leu	Val	500	505	510	
Gln	Ile	Ala	Arg	Val	Ile	Leu	Glu	Tyr	Ile	Asp	His	Lys	Leu	Arg	Gly	515	520	525	
Val	Gln	Asn	Pro	Val	Ala	Arg	Cys	Ile	Met	Cys	Cys	Phe	Lys	Cys	Cys	530	535	540	
Leu	Trp	Cys	Leu	Glu	Lys	Phe	Ile	Lys	Phe	Leu	Asn	Arg	Asn	Ala	Tyr	545	550	555	560
Ile	Met	Ile	Ala	Ile	Tyr	Gly	Lys	Asn	Phe	Cys	Val	Ser	Ala	Lys	Asn				

-continued

565	570	575
Ala Phe Met Leu Leu Met Arg Asn Ile Val Arg Val Val Val Leu Asp 580 585 590		
Lys Val Thr Asp Leu Leu Leu Phe Phe Gly Lys Leu Leu Val Val Gly 595 600 605		
Gly Val Gly Val Leu Ser Phe Phe Phe Phe Ser Gly Arg Ile Pro Gly 610 615 620		
Leu Gly Lys Asp Phe Lys Ser Pro His Leu Asn Tyr Tyr Trp Leu Pro 625 630 635 640		
Ile Met Thr Ser Ile Leu Gly Ala Tyr Val Ile Ala Ser Gly Phe Phe 645 650 655		
Ser Val Phe Gly Met Cys Val Asp Thr Leu Phe Leu Cys Phe Leu Glu 660 665 670		
Asp Leu Glu Arg Asn Asn Gly Ser Leu Asp Arg Pro Tyr Tyr Met Ser 675 680 685		
Lys Ser Leu Leu Lys Ile Leu Gly Lys Lys Asn Glu Ala Pro Pro Asp 690 695 700		
Asn Lys Lys Arg Lys Lys 705 710		
<210> SEQ ID NO 4		
<211> LENGTH: 2587		
<212> TYPE: DNA		
<213> ORGANISM: Homo sapiens		
<400> SEQUENCE: 4		
gagccatggg gggaaagcag cgggacgagg atgacgaggc ctacgggaag ccagtcaa	60	
acgacccctc ctttcgaggg cccatcaaga acagaagctg cacagatgtc atctgtctgc	120	
tcctcttctc gctcttcatt ctagggtaca tcgtggtggg gattgtggcc tggttgtatg	180	
gagacccccc gcaagtccct taccocagga actctactgg ggcctactgt ggcattgggg	240	
agaacaaaga taagccgtat ctctgttact tcaacatctt cagctgcac cgtgccagca	300	
acatcatctc agttgctgag aacggcctac agtgccccac accccagggtg tgtgtgtcct	360	
cctgcccgga ggacccatgg actgtgggaa aaaacgagtt ctcacagact gttgggggag	420	
tctctatata aaaaaacagg aacttttgtc tgccaggggt accctggaat atgacgggtg	480	
tcacaagcct gcaacaggaa ctctgcccc gtttctctct cccctctgct ccagctctgg	540	
gacgtgctt tccatggacc aacgttactc caccggcgct cccagggatc accaatgaca	600	
ccaccatata gcaggggatc agcgggtctta ttgacagcct caatgcccgga gacatcagtg	660	
ttaagatctt tgaagatttt gccagctctt ggtattggat tcttgttgcc ctgggggtgg	720	
ctctggctct gaggcctactg tttatcttgc ttctgcgct ggtggctggg cccctgggtc	780	
tgggtctgat cctgggagtg ctgggcgtgc tggcatacgg catctactac tgcctggagg	840	
agtaccgagt gctgcgggac aagggcgct ccatctccca gctgggttcc accaccaacc	900	
tcagtgccta ccagagcgtg caggagacct ggctggccgc cctgatcgtg ttggcgggtc	960	
ttgaagccat cctgctgctg atgctcatct tcctgcggca gcggattcgt attgccatcg	1020	
ccctcctgaa ggaggccagc aaggctgtgg gacagatgat gtctaccatg ttctaccac	1080	
tggtcacctt tgcctcctc ctcacttgca ttgcctactg ggccatgact gctctgtacc	1140	
tggctacatc ggggcaaccc cagtatgtgc tctgggcac caacatcagc tccccgggt	1200	
gtgagaaagt gccataaat acatcatgca accccacggc ccacctgtg aactcctcgt	1260	
gcccagggt gatgtgcgtc ttccagggt actcatccaa aggcctaata caacgttctg	1320	

-continued

```

tcttcaatct gcaaatctat ggggtcctgg ggtctttctg gaccottaac tgggtactgg 1380
ccctggggcca atcgctcctc gctggagcct ttgcctcctt ctactggggc ttcacaagc 1440
cccaggacat ccttaccttc cccttaatct ctgccttcat ccgcacactc cgttaccaca 1500
ctgggtcatt ggcatttga gccctcatcc tgacccttgt gcagatagcc cgggtcatct 1560
tggagtatat tgaccacaag ctacaggagag tgcagaaccc tgtagcccg cgcacatcatgt 1620
gctgtttcaa gtgcgcctc tgggtgtctgg aaaaatttat caagtcccta aaccgcaatg 1680
catacatcat gatcgccatc tacgggaaga atttctgtgt ctacgcaaaa aatgcgttca 1740
tgctactcat gcgaaacatt gtcagggtgg tcgtcctgga caaagtcaca gacctgctgc 1800
tgttctttgg gaagctgctg gtggtcggag gcgtgggggt cctgtccttc ttttttttct 1860
ccggtcgcac cccggggctg ggtaagagact ttaagagccc ccacctcaac tattactggc 1920
tgcccatcat gacctccatc ctgggggacct atgtcatcgc cagcggett ctcagcgttt 1980
tcggcatgtg tgtggacacg ctcttcctct gcttcctgga agacctggag cggaacaacg 2040
gctccctgga ccggccctac tacatgtcca agagccttct aaagattctg ggcaagaaga 2100
acgaggcgcc cccggacaac aagaagagga agaagtgaca gctccggccc tgatccagga 2160
ctgcaccoca cccccaccgt ccagccatcc aacctcaatt cgccttacag gtctccattt 2220
tgtggtaaaa aaaggtttta ggcaggcgcc cgtggctcac gctgtaac caacactttg 2280
agaggctgag gcggggcgat cacctgagtc aggagtctga gaccagcctg gccaacatgg 2340
tgaaacctcc gtctctatta aaaatacaaa aattagccga gagtgggtggc atgcacctgt 2400
catcccagct actcgggagg ctgaggcagg agaatcgctt gaaccggga gccagagggt 2460
gcagtgagcc gagatcgcc cactgcactc caacctgggt gacagactct gtctccaaaa 2520
caaaacaaac aaacaaaaag attttattaa agatatatttg ttaactcagt aaaaaaaaaa 2580
aaaaaaa

```

<210> SEQ ID NO 5

<211> LENGTH: 710

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 5

```

Met Gly Gly Lys Gln Arg Asp Glu Asp Asp Glu Ala Tyr Gly Lys Pro
 1             5             10             15
Val Lys Tyr Asp Pro Ser Phe Arg Gly Pro Ile Lys Asn Arg Ser Cys
 20            25            30
Thr Asp Val Ile Cys Cys Val Leu Phe Leu Leu Phe Ile Leu Gly Tyr
 35            40            45
Ile Val Val Gly Ile Val Ala Trp Leu Tyr Gly Asp Pro Arg Gln Val
 50            55            60
Leu Tyr Pro Arg Asn Ser Thr Gly Ala Tyr Cys Gly Met Gly Glu Asn
 65            70            75            80
Lys Asp Lys Pro Tyr Leu Leu Tyr Phe Asn Ile Phe Ser Cys Ile Leu
 85            90            95
Ser Ser Asn Ile Ile Ser Val Ala Glu Asn Gly Leu Gln Cys Pro Thr
100           105           110
Pro Gln Val Cys Val Ser Ser Cys Pro Glu Asp Pro Trp Thr Val Gly
115           120           125
Lys Asn Glu Phe Ser Gln Thr Val Gly Glu Val Phe Tyr Thr Lys Asn
130           135           140

```

-continued

Arg	Asn	Phe	Cys	Leu	Pro	Gly	Val	Pro	Trp	Asn	Met	Thr	Val	Ile	Thr	145	150	155	160
Ser	Leu	Gln	Gln	Glu	Leu	Cys	Pro	Ser	Phe	Leu	Leu	Pro	Ser	Ala	Pro	165	170	175	
Ala	Leu	Gly	Arg	Cys	Phe	Pro	Trp	Thr	Asn	Val	Thr	Pro	Pro	Ala	Leu	180	185	190	
Pro	Gly	Ile	Thr	Asn	Asp	Thr	Thr	Ile	Gln	Gln	Gly	Ile	Ser	Gly	Leu	195	200	205	
Ile	Asp	Ser	Leu	Asn	Ala	Arg	Asp	Ile	Ser	Val	Lys	Ile	Phe	Glu	Asp	210	215	220	
Phe	Ala	Gln	Ser	Trp	Tyr	Trp	Ile	Leu	Val	Ala	Leu	Gly	Val	Ala	Leu	225	230	235	240
Val	Leu	Ser	Leu	Leu	Phe	Ile	Leu	Leu	Leu	Arg	Leu	Val	Ala	Gly	Pro	245	250	255	
Leu	Val	Leu	Val	Leu	Ile	Leu	Gly	Val	Leu	Gly	Val	Leu	Ala	Tyr	Gly	260	265	270	
Ile	Tyr	Tyr	Cys	Trp	Glu	Glu	Tyr	Arg	Val	Leu	Arg	Asp	Lys	Gly	Ala	275	280	285	
Ser	Ile	Ser	Gln	Leu	Gly	Phe	Thr	Thr	Asn	Leu	Ser	Ala	Tyr	Gln	Ser	290	295	300	
Val	Gln	Glu	Thr	Trp	Leu	Ala	Ala	Leu	Ile	Val	Leu	Ala	Val	Leu	Glu	305	310	315	320
Ala	Ile	Leu	Leu	Leu	Met	Leu	Ile	Phe	Leu	Arg	Gln	Arg	Ile	Arg	Ile	325	330	335	
Ala	Ile	Ala	Leu	Leu	Lys	Glu	Ala	Ser	Lys	Ala	Val	Gly	Gln	Met	Met	340	345	350	
Ser	Thr	Met	Phe	Tyr	Pro	Leu	Val	Thr	Phe	Val	Leu	Leu	Leu	Ile	Cys	355	360	365	
Ile	Ala	Tyr	Trp	Ala	Met	Thr	Ala	Leu	Tyr	Leu	Ala	Thr	Ser	Gly	Gln	370	375	380	
Pro	Gln	Tyr	Val	Leu	Trp	Ala	Ser	Asn	Ile	Ser	Ser	Pro	Gly	Cys	Glu	385	390	395	400
Lys	Val	Pro	Ile	Asn	Thr	Ser	Cys	Asn	Pro	Thr	Ala	His	Leu	Val	Asn	405	410	415	
Ser	Ser	Cys	Pro	Gly	Leu	Met	Cys	Val	Phe	Gln	Gly	Tyr	Ser	Ser	Lys	420	425	430	
Gly	Leu	Ile	Gln	Arg	Ser	Val	Phe	Asn	Leu	Gln	Ile	Tyr	Gly	Val	Leu	435	440	445	
Gly	Leu	Phe	Trp	Thr	Leu	Asn	Trp	Val	Leu	Ala	Leu	Gly	Gln	Cys	Val	450	455	460	
Leu	Ala	Gly	Ala	Phe	Ala	Ser	Phe	Tyr	Trp	Ala	Phe	His	Lys	Pro	Gln	465	470	475	480
Asp	Ile	Pro	Thr	Phe	Pro	Leu	Ile	Ser	Ala	Phe	Ile	Arg	Thr	Leu	Arg	485	490	495	
Tyr	His	Thr	Gly	Ser	Leu	Ala	Phe	Gly	Ala	Leu	Ile	Leu	Thr	Leu	Val	500	505	510	
Gln	Ile	Ala	Arg	Val	Ile	Leu	Glu	Tyr	Ile	Asp	His	Lys	Leu	Arg	Gly	515	520	525	
Val	Gln	Asn	Pro	Val	Ala	Arg	Cys	Ile	Met	Cys	Cys	Phe	Lys	Cys	Cys	530	535	540	
Leu	Trp	Cys	Leu	Glu	Lys	Phe	Ile	Lys	Phe	Leu	Asn	Arg	Asn	Ala	Tyr	545	550	555	560
Ile	Met	Ile	Ala	Ile	Tyr	Gly	Lys	Asn	Phe	Cys	Val	Ser	Ala	Lys	Asn	565	570	575	

-continued

Ala Phe Met Leu Leu Met Arg Asn Ile Val Arg Val Val Val Leu Asp
580 585 590

Lys Val Thr Asp Leu Leu Leu Phe Phe Gly Lys Leu Leu Val Val Gly
595 600 605

Gly Val Gly Val Leu Ser Phe Phe Phe Ser Gly Arg Ile Pro Gly
610 615 620

Leu Gly Lys Asp Phe Lys Ser Pro His Leu Asn Tyr Tyr Trp Leu Pro
625 630 635 640

Ile Met Thr Ser Ile Leu Gly Ala Tyr Val Ile Ala Ser Gly Phe Phe
645 650 655

Ser Val Phe Gly Met Cys Val Asp Thr Leu Phe Leu Cys Phe Leu Glu
660 665 670

Asp Leu Glu Arg Asn Asn Gly Ser Leu Asp Arg Pro Tyr Tyr Met Ser
675 680 685

Lys Ser Leu Leu Lys Ile Leu Gly Lys Lys Asn Glu Ala Pro Pro Asp
690 695 700

Asn Lys Lys Arg Lys Lys
705 710

<210> SEQ ID NO 6

<211> LENGTH: 2587

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 6

```

gagccatggg gggaaagcag cgggacgagg atgacgaggc ctacgggaag ccagtcaaat    60
acgacccctc ctttcgaggc cccatcaaga acagaagctg cacagatgtc atctgtctgcg    120
tcctcttctc gctcttcatt ctagggtaca tcgtggtggg gattgtggcc tggttgtatg    180
gagacccccg gcaagtcttc taccacagga actctactgg ggcctactgt ggcctggggg    240
agaacaaaga taagccgtat ctctgttact tcaacatctt cagctgcctc ctgtccagca    300
acatcatctc agttgtctgag aacggcctac agtgccccac accccagggtg tgtgtgtcct    360
cctgccccga ggacccatgg actgtgggaa aaaacgagtt ctcacagact gttggggaag    420
tcttctatac aaaaaacagg aacttttgtc tgccaggggt accctggaat atgacgggtga    480
tcacaagcct gcaacaggaa ctctgccccg gtttctctct cccctctgct ccagctctgg    540
ggcgctgctt tccatggacc aacattactc caccggcgct cccagggatc accaatgaca    600
ccaccataca gcaggggatc agcgggtctta ttgacagcct caatgccga gacatcagtg    660
ttaagatctt tgaagatctt gccagtcctt ggtattggat tcttgttgcc ctgggggtgg    720
ctctgggtctt gagectactg tttatcttgc ttctgcgcct ggtggctggg cccctgggtgc    780
tggtgctgat cctgggagtg ctgggcgtgc tggcatacgg catctactac tgetgggagg    840
agtaccgagt gctgcgggac aaggggcgct ccatctccca gctgggttcc accaccaacc    900
tcagtgccta ccagagcgtg caggagacct ggctggccgc cctgatcgtg ttggcgggtgc    960
ttgaagccat cctgctgctg atgctcatct tcttgcggca gcggattcgt attgccatcg   1020
ccctctgtaa ggaggccagc aaggctgtgg gacagatgat gtctaccatg ttctaccac    1080
tggtcacctt tgtctctctc ctcatctgca ttgcctactg ggccatgact gctctgtacc   1140
tggtctacac ggggcaaccc cagtatgtgc tctgggcata caacatcagc tccccgggt    1200
gtgagaaagt gccaaataat acatcatgca accccacggc ccaccttggt aactcctcgt   1260
gcccagggct gatgtgcgtc ttccagggct actcatccaa aggcctaata caacgttctg   1320

```

-continued

```

tcttcaatct gcaaatctat ggggtcctgg ggctcttctg gaccttaac tgggtactgg 1380
ccctgggcca atgcgccttc gctggagcct ttgcctcctt ctactgggcc ttccacaagc 1440
cccaggacat ccctaccttc cccttaatct ctgccttcat ccgcacactc cgttaccaca 1500
ctgggtcatt ggcatttgga gccctcatcc tgaccttgt gcagatagcc cgggtcatct 1560
tggagtatat tgaccacaag ctcagaggag tgcagaaccc tgtagccgc tgcacatgt 1620
gctgtttcaa gtgctgcctc tgggtgtctgg aaaaatttat caagttccta aaccgcaatg 1680
catacatcat gatcgccatc tacgggaaga atttctgtgt ctcagccaaa aatgcgttca 1740
tgctactcat gcgaacatt gtcagggtgg tcgtcttgga caaagtcaca gacctgctgc 1800
tgttctttgg gaagctgctg gtggtcggag gcgtgggggt cctgtccttc tttttttct 1860
ccggtcgcat cccggggctg ggtaaagact ttaagagccc ccacctcaac tattactggc 1920
tgcccatcat gacctccatc ctgggggct atgtcatgc cagcggttc ttcagcgttt 1980
tcggcatgtg tgtggacacg ctcttctctt gcttctgga agacctggag cggaacaacg 2040
gtccctgga ccggccctac tacatgtcca agagccttct aaagattctg ggcaagaaga 2100
acgaggcgcc cccggacaac aagaagagga agaagtgaca gctccggccc tgatccagga 2160
ctgcacccca cccccaccgt ccagccatcc aacctcactt cgccttacag gtctccattt 2220
tgtggtaaaa aaaggtttta ggcaggcgc cgtggctcac gcctgtaac caacactttg 2280
agaggctgag gcggggcgat cactgagtc aggagtcca gaccagcctg gccaacatgg 2340
tgaaacctcc gtctctatta aaaatacaaa aattagccga gagtgggtggc atgcacctgt 2400
catcccagct actcgggagg ctgaggcagg agaategctt gaaccggga ggcagaggtt 2460
gcagtgagcc gagatcgcg cactgcactc caacctgggt gacagactct gtctccaaaa 2520
caaaacaaac aaacaaaaag attttattaa agatatattt ttaactcagt aaaaaaaaaa 2580
aaaaaaa

```

```

<210> SEQ ID NO 7
<211> LENGTH: 710
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 7

```

```

Met Gly Gly Lys Gln Arg Asp Glu Asp Asp Glu Ala Tyr Gly Lys Pro
 1             5             10            15
Val Lys Tyr Asp Pro Ser Phe Arg Gly Pro Ile Lys Asn Arg Ser Cys
      20            25            30
Thr Asp Val Ile Cys Cys Val Leu Phe Leu Leu Phe Ile Leu Gly Tyr
      35            40            45
Ile Val Val Gly Ile Val Ala Trp Leu Tyr Gly Asp Pro Arg Gln Val
      50            55            60
Leu Tyr Pro Arg Asn Ser Thr Gly Ala Tyr Cys Gly Met Gly Glu Asn
      65            70            75            80
Lys Asp Lys Pro Tyr Leu Leu Tyr Phe Asn Ile Phe Ser Cys Ile Leu
      85            90            95
Ser Ser Asn Ile Ile Ser Val Ala Glu Asn Gly Leu Gln Cys Pro Thr
      100           105           110
Pro Gln Val Cys Val Ser Ser Cys Pro Glu Asp Pro Trp Thr Val Gly
      115           120           125
Lys Asn Glu Phe Ser Gln Thr Val Gly Glu Val Phe Tyr Thr Lys Asn
      130           135           140
Arg Asn Phe Cys Leu Pro Gly Val Pro Trp Asn Met Thr Val Ile Thr

```


-continued

145	150	155	160
Ser Leu Gln Gln Glu Leu Cys Pro Ser Phe Leu Leu Pro Ser Ala Pro	165	170	175
Ala Leu Gly Arg Cys Phe Pro Trp Thr Asn Ile Thr Pro Pro Ala Leu	180	185	190
Pro Gly Ile Thr Asn Asp Thr Thr Ile Gln Gln Gly Ile Ser Gly Leu	195	200	205
Ile Asp Ser Leu Asn Ala Arg Asp Ile Ser Val Lys Ile Phe Glu Asp	210	215	220
Phe Ala Gln Ser Trp Tyr Trp Ile Leu Val Ala Leu Gly Val Ala Leu	225	230	235
Val Leu Ser Leu Leu Phe Ile Leu Leu Leu Arg Leu Val Ala Gly Pro	245	250	255
Leu Val Leu Val Leu Ile Leu Gly Val Leu Gly Val Leu Ala Tyr Gly	260	265	270
Ile Tyr Tyr Cys Trp Glu Glu Tyr Arg Val Leu Arg Asp Lys Gly Ala	275	280	285
Ser Ile Ser Gln Leu Gly Phe Thr Thr Asn Leu Ser Ala Tyr Gln Ser	290	295	300
Val Gln Glu Thr Trp Leu Ala Ala Leu Ile Val Leu Ala Val Leu Glu	305	310	315
Ala Ile Leu Leu Leu Met Leu Ile Phe Leu Arg Gln Arg Ile Arg Ile	325	330	335
Ala Ile Ala Leu Leu Lys Glu Ala Ser Lys Ala Val Gly Gln Met Met	340	345	350
Ser Thr Met Phe Tyr Pro Leu Val Thr Phe Val Leu Leu Leu Ile Cys	355	360	365
Ile Ala Tyr Trp Ala Met Thr Ala Leu Tyr Leu Ala Thr Ser Gly Gln	370	375	380
Pro Gln Tyr Val Leu Trp Ala Ser Asn Ile Ser Ser Pro Gly Cys Glu	385	390	395
Lys Val Pro Ile Asn Thr Ser Cys Asn Pro Thr Ala His Leu Val Asn	405	410	415
Ser Ser Cys Pro Gly Leu Met Cys Val Phe Gln Gly Tyr Ser Ser Lys	420	425	430
Gly Leu Ile Gln Arg Ser Val Phe Asn Leu Gln Ile Tyr Gly Val Leu	435	440	445
Gly Leu Phe Trp Thr Leu Asn Trp Val Leu Ala Leu Gly Gln Cys Val	450	455	460
Leu Ala Gly Ala Phe Ala Ser Phe Tyr Trp Ala Phe His Lys Pro Gln	465	470	475
Asp Ile Pro Thr Phe Pro Leu Ile Ser Ala Phe Ile Arg Thr Leu Arg	485	490	495
Tyr His Thr Gly Ser Leu Ala Phe Gly Ala Leu Ile Leu Thr Leu Val	500	505	510
Gln Ile Ala Arg Val Ile Leu Glu Tyr Ile Asp His Lys Leu Arg Gly	515	520	525
Val Gln Asn Pro Val Ala Arg Cys Ile Met Cys Cys Phe Lys Cys Cys	530	535	540
Leu Trp Cys Leu Glu Lys Phe Ile Lys Phe Leu Asn Arg Asn Ala Tyr	545	550	555
Ile Met Ile Ala Ile Tyr Gly Lys Asn Phe Cys Val Ser Ala Lys Asn	565	570	575

-continued

Ala Phe Met Leu Leu Met Arg Asn Ile Val Arg Val Val Val Leu Asp
580 585 590

Lys Val Thr Asp Leu Leu Leu Phe Phe Gly Lys Leu Leu Val Val Gly
595 600 605

Gly Val Gly Val Leu Ser Phe Phe Phe Ser Gly Arg Ile Pro Gly
610 615 620

Leu Gly Lys Asp Phe Lys Ser Pro His Leu Asn Tyr Tyr Trp Leu Pro
625 630 635 640

Ile Met Thr Ser Ile Leu Gly Ala Tyr Val Ile Ala Ser Gly Phe Phe
645 650 655

Ser Val Phe Gly Met Cys Val Asp Thr Leu Phe Leu Cys Phe Leu Glu
660 665 670

Asp Leu Glu Arg Asn Asn Gly Ser Leu Asp Arg Pro Tyr Tyr Met Ser
675 680 685

Lys Ser Leu Leu Lys Ile Leu Gly Lys Lys Asn Glu Ala Pro Pro Asp
690 695 700

Asn Lys Lys Arg Lys Lys
705 710

<210> SEQ ID NO 8

<211> LENGTH: 2587

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 8

```

gagccatggg gggaaagcag cgggacgagg atgacgaggc ctacgggaag ccagtcaaat    60
acgacccctc ctttcgaggc cccatcaaga acagaagctg cacagatgtc atctgtctgcg    120
tcctcttctc gctcttcatt ctagggttaca tcgtggtggg gattgtggcc tggttgtatg    180
gagacccccg gcaagtcctc taccacagga actctactgg ggcctactgt ggcatggggg    240
agaacaaaga taagccgtat ctctgttact tcaacatctt cagctgcata ctgtccagca    300
acatcatctc agttgctgag aacggcctac agtgccccac acccaggtg tgtgtgtcct    360
cctgcccgga ggacccatgg actgtgggaa aaaacgagtt ctcacagact gttggggaag    420
tcttctatata aaaaaacagg aacttttgtc tgccaggggt accctggaat atgacggtga    480
tcacaagcct gcaacaggaa ctctgcccga gtttctctct cccctctgct ccagctcttg    540
ggcgctgctt tccatggacc aacgttactc caccggcgct cccagggatc accaatgaca    600
ccaccatata gcaggggatc agcgggtctta ttgacagcct caatgcccga gacatcagtg    660
ttaagatctt tgaagatttt gcccagtcct ggtattggat tcttggtgcc ctgggggtgg    720
ctctggtctt gagcctactg tttatcttgc ttctgcccct ggtggctggg ccctggtgc    780
tggtgctgat cctgggagtg ctgggcgtgc tggcatatgg catctactac tgctgggagg    840
agtaccgagt gctgctggac aaggggcctt ccatctccca gctgggtttc accaccaacc    900
tcagtgccta ccagagcgtg caggagacct ggctggccgc cctgatcgtg ttggcggtgc    960
ttgaagccat cctgctgctg atgctcatct tcctgcggca gcggattcgt attgccatcg   1020
ccctctgaa ggaggccagc aaggctgtgg gacagatgat gtctaccatg ttctaccac    1080
tggtcacctt tgcctcctc ctcatctgca ttgcctactg ggccatgact gctctgtacc    1140
tggctacatc ggggcaacco cagtatgtgc tctgggcata caacatcagc tccccggct    1200
gtgagaaagt gccaaataat acatcatgca accccacggc ccacctgtg aactcctcgt    1260
gccagggtgat gctgctgctc ttccagggtc actcatccaa aggcctaata caacgttctg    1320
tcttcaatct gcaaatctat ggggtcctgg ggetcttctg gacccttaac tgggtactgg    1380

```

Figure 2I-3

700 K N E A P F D N K K R K K *

2101 AGAAGAACGAGGCGCCCCGGACAACAAGAAGAGGAAGAAGTGAacagctccggccctgat

2161 ccaggactgcacccccacccccaccgtccagccatccaacctcacttcgccttacaggtct

2221 ccattttgtggtaaaaaaagggttttaggccaggcgccgtgggtcacgcctgtaatccaac

2281 actttgagaggctgaggcgggcggtacacctgagtcaggagttcgagaccagcctggcca

2341 acatggtgaaacctcgtctctattaaaaatacaaaaattagccgagagtggtggcatgc

2401 acctgtcatcccagctactcgggaggctgaggcaggagaatcgcttgaacccgggaggca

2461 gaggttgacgtgagccgagatcgcgccactgcactccaacctgggtgacagactctgtct

2521 ccaaaacaaaacaaacaaaacaaaagattttattaaagatatatttgttactcagtaaaa

2581 aaaaaaaaaaaaaa

-continued

```

ccctggggcca atcgctcctc gctggagcct ttgcctcctt ctactggggc ttcacaaagc 1440
cccaggacat ccctaccttc cccttaatct ctgccttcat ccgcacactc cgttaccaca 1500
ctgggtcatt ggcatttga gccctcatcc tgacccttgt gcagatagcc cgggtcatct 1560
tggagtatat tgaccacaag ctgagaggag tgcagaaccc tgtagcccg cgcacatgt 1620
gctgtttcaa gtgctgcctc tgggtgtctgg aaaaatttat caagttccta aaccgcaatg 1680
catacatcat gatcgccatc tacgggaaga atttctgtgt ctgagccaaa aatgcgttca 1740
tgctactcat gcgaaacatt gtcagggtgg tcgtcctgga caaagtcaca gacctgctgc 1800
tgttctttgg gaagctgctg tgggtcggag gcgtgggggt cctgtccttc tttttttct 1860
ccggtcgcac ccgggggctg ggtaaaagact ttaagagccc ccacatcaac tattactggc 1920
tgcccatcat gacctccatc ctgggggct atgtcatcgc cagcggttc ttcagcgttt 1980
tcggcatgtg tgtggacacg ctcttctct gcttctgga agacctggag cggaacaacg 2040
gctccctgga ccggccctac tacatgtcca agagccttct aaagattctg ggcaagaaga 2100
acgaggcgcc ccgggacaac aagaagagga agaagtgaca gctccggccc tgatccagga 2160
ctgcaccca cccccccgt ccagccatcc aacctcactt cgccttacag gtctccattt 2220
tgtggtaaaa aaaggtttta ggccaggcgc cgtggctcac gcctgtaac caacactttg 2280
agaggctgag gcggggcgat cacctgagtc aggagtctga gaccagctg gccaacatgg 2340
tgaaacctcc gtctctatta aaaatacaaa aattagccga gagtggtggc atgcacctgt 2400
catcccagct actcgggagg ctgaggcagg agaatcgctt gaaccgggga ggcagaggtt 2460
gcagtgagcc gagatcgcc cactgcactc caacctgggt gacagactct gtctccaaaa 2520
caaaaacaac aaacaaaaag attttattaa agatattttg ttaactcagt aaaaaaaaaa 2580
aaaaaaa 2587

```

<210> SEQ ID NO 9

<211> LENGTH: 710

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 9

```

Met Gly Gly Lys Gln Arg Asp Glu Asp Glu Ala Tyr Gly Lys Pro
 1           5           10          15
Val Lys Tyr Asp Pro Ser Phe Arg Gly Pro Ile Lys Asn Arg Ser Cys
 20          25          30
Thr Asp Val Ile Cys Cys Val Leu Phe Leu Leu Phe Ile Leu Gly Tyr
 35          40          45
Ile Val Val Gly Ile Val Ala Trp Leu Tyr Gly Asp Pro Arg Gln Val
 50          55          60
Leu Tyr Pro Arg Asn Ser Thr Gly Ala Tyr Cys Gly Met Gly Glu Asn
 65          70          75          80
Lys Asp Lys Pro Tyr Leu Leu Tyr Phe Asn Ile Phe Ser Cys Ile Leu
 85          90          95
Ser Ser Asn Ile Ile Ser Val Ala Glu Asn Gly Leu Gln Cys Pro Thr
100         105         110
Pro Gln Val Cys Val Ser Ser Cys Pro Glu Asp Pro Trp Thr Val Gly
115         120         125
Lys Asn Glu Phe Ser Gln Thr Val Gly Glu Val Phe Tyr Thr Lys Asn
130         135         140
Arg Asn Phe Cys Leu Pro Gly Val Pro Trp Asn Met Thr Val Ile Thr
145         150         155         160

```

-continued

Ser Leu Gln Gln Glu Leu Cys Pro Ser Phe Leu Leu Pro Ser Ala Pro
 165 170 175
 Ala Leu Gly Arg Cys Phe Pro Trp Thr Asn Val Thr Pro Pro Ala Leu
 180 185 190
 Pro Gly Ile Thr Asn Asp Thr Thr Ile Gln Gln Gly Ile Ser Gly Leu
 195 200 205
 Ile Asp Ser Leu Asn Ala Arg Asp Ile Ser Val Lys Ile Phe Glu Asp
 210 215 220
 Phe Ala Gln Ser Trp Tyr Trp Ile Leu Val Ala Leu Gly Val Ala Leu
 225 230 235 240
 Val Leu Ser Leu Leu Phe Ile Leu Leu Leu Arg Leu Val Ala Gly Pro
 245 250 255
 Leu Val Leu Val Leu Ile Leu Gly Val Leu Gly Val Leu Ala Tyr Gly
 260 265 270
 Ile Tyr Tyr Cys Trp Glu Glu Tyr Arg Val Leu Arg Asp Lys Gly Ala
 275 280 285
 Ser Ile Ser Gln Leu Gly Phe Thr Thr Asn Leu Ser Ala Tyr Gln Ser
 290 295 300
 Val Gln Glu Thr Trp Leu Ala Ala Leu Ile Val Leu Ala Val Leu Glu
 305 310 315 320
 Ala Ile Leu Leu Leu Met Leu Ile Phe Leu Arg Gln Arg Ile Arg Ile
 325 330 335
 Ala Ile Ala Leu Leu Lys Glu Ala Ser Lys Ala Val Gly Gln Met Met
 340 345 350
 Ser Thr Met Phe Tyr Pro Leu Val Thr Phe Val Leu Leu Leu Ile Cys
 355 360 365
 Ile Ala Tyr Trp Ala Met Thr Ala Leu Tyr Leu Ala Thr Ser Gly Gln
 370 375 380
 Pro Gln Tyr Val Leu Trp Ala Ser Asn Ile Ser Ser Pro Gly Cys Glu
 385 390 395 400
 Lys Val Pro Ile Asn Thr Ser Cys Asn Pro Thr Ala His Leu Val Asn
 405 410 415
 Ser Ser Cys Pro Gly Leu Met Cys Val Phe Gln Gly Tyr Ser Ser Lys
 420 425 430
 Gly Leu Ile Gln Arg Ser Val Phe Asn Leu Gln Ile Tyr Gly Val Leu
 435 440 445
 Gly Leu Phe Trp Thr Leu Asn Trp Val Leu Ala Leu Gly Gln Cys Val
 450 455 460
 Leu Ala Gly Ala Phe Ala Ser Phe Tyr Trp Ala Phe His Lys Pro Gln
 465 470 475 480
 Asp Ile Pro Thr Phe Pro Leu Ile Ser Ala Phe Ile Arg Thr Leu Arg
 485 490 495
 Tyr His Thr Gly Ser Leu Ala Phe Gly Ala Leu Ile Leu Thr Leu Val
 500 505 510
 Gln Ile Ala Arg Val Ile Leu Glu Tyr Ile Asp His Lys Leu Arg Gly
 515 520 525
 Val Gln Asn Pro Val Ala Arg Cys Ile Met Cys Cys Phe Lys Cys Cys
 530 535 540
 Leu Trp Cys Leu Glu Lys Phe Ile Lys Phe Leu Asn Arg Asn Ala Tyr
 545 550 555 560
 Ile Met Ile Ala Ile Tyr Gly Lys Asn Phe Cys Val Ser Ala Lys Asn
 565 570 575
 Ala Phe Met Leu Leu Met Arg Asn Ile Val Arg Val Val Val Leu Asp

-continued

580	585	590
Lys Val Thr Asp Leu Leu Leu Phe Phe Gly Lys Leu Leu Val Val Gly		
595	600	605
Gly Val Gly Val Leu Ser Phe Phe Phe Phe Ser Gly Arg Ile Pro Gly		
610	615	620
Leu Gly Lys Asp Phe Lys Ser Pro His Leu Asn Tyr Tyr Trp Leu Pro		
625	630	635
Ile Met Thr Ser Ile Leu Gly Ala Tyr Val Ile Ala Ser Gly Phe Phe		
645	650	655
Ser Val Phe Gly Met Cys Val Asp Thr Leu Phe Leu Cys Phe Leu Glu		
660	665	670
Asp Leu Glu Arg Asn Asn Gly Ser Leu Asp Arg Pro Tyr Tyr Met Ser		
675	680	685
Lys Ser Leu Leu Lys Ile Leu Gly Lys Lys Asn Glu Ala Pro Pro Asp		
690	695	700
Asn Lys Lys Arg Lys Lys		
705	710	

<210> SEQ ID NO 10

<211> LENGTH: 2587

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 10

gagccatggg gggaaagcag cgggacgagg atgacgaggc ctacgggaag ccagtcaa	60
acgacccctc ctttcgaggc cccatcaaga acagaagctg cacagatgtc atctgctg	120
tcctcttctc gctcttcatt ctagggtaca tcgtggtggg gattgtggcc tggttgtatg	180
gagacccccg gcaagtccctc taccccagga actctactgg ggcctactgt ggcattgggg	240
agaacaaaga taagccgtat ctctgtact tcaacatctt cagctgcac cgtccagca	300
acatcatctc agttgctgag aacggcctac agtgccccac acccagggtg tgtgtgtcct	360
cctgcccgga ggacccatgg actgtgggaa aaaacgagtt ctcacagact gttggggaag	420
tcttctatc aaaaaacagg aacttttgtc tgccaggggt accctggaat atgacggtga	480
tcacaagcct gcaacaggaa ctctgcccc gtttctctc cccctctgct ccagctctgg	540
ggcgctgctt tccatggacc aacgttactc caccggcgct cccagggtac accaatgaca	600
ccaccatata gcaggggac agcgggtctta ttgacagcct caatgccga gacatcagtg	660
ttaagatctt tgaagatttt gccagtcct ggtattggat tcttgttgcc ctgggggtgg	720
ctctggctct gagcctactg tttatcttgc ttctgcgcct ggtggctggg cccctgggtc	780
tgggtgctgat cctgggagtg ctgggctgctc tggcatacgg catctactac tgtgggagg	840
agtaccgagt gctgcgggac aagggcgcct ccatctccca gctgggttcc accaccaacc	900
tcagtgccta ccagagcgtg caggagacct ggctggccgc cctgatcgtg ttggcggtgc	960
ttgaagccat cctgctgctg gtgctcatct tcctgcggca gcgattcgt attgccatcg	1020
ccctcctgaa ggaggccagc aaggctgtgg gacagatgat gtctaccatg ttctaccac	1080
tggteacctt tgtctctctc ctcatctgca ttgctactg ggcattgact gctctgtacc	1140
tggctacatc ggggcaaccc cagtatgtgc tctgggcac caacatcagc tccccggct	1200
gtgagaaagt gccataaat acatcatgca accccacggc ccaccttggt aactcctcgt	1260
gccagggtgat gatgtcgctc ttccagggt actcatccaa aggcctaata caacgttctg	1320
tcttcaatct gcaaatctat ggggtcctgg ggctcttctg gaccottaac tgggtactgg	1380

-continued

```

cctctgggcca atcgctctc gctggagcct ttgcctcctt ctactgggcc ttccacaage 1440
cccaggacat cctacacctt ccttaatct ctgccttcat cgcacactc cgttaccaca 1500
ctgggtcatt ggcatttga gccctcatcc tgaccttgt gcagatagcc cgggtcatct 1560
tggagtatat tgaccacaag ctgagaggag tgcagaaccc ttagcccccgc tgcacatgt 1620
gctgtttcaa gtgctgcctc tgggtgtctgg aaaaatttat caagtcccta aaccgcaatg 1680
catacatcat gatcgccatc tacgggaaga atttctgtgt ctgagccaaa aatgcgttca 1740
tgctactcat gcgaaacatt gtcagggtgg tcgtcctgga caaagtcaca gacctgctgc 1800
tgttcttttg gaagctgctg gtggtcggag gcgtgggggt cctgtccttc ttttttttct 1860
ccggtcgcac cccggggctg ggtaaagact ttaagagccc ccacctcaac tattactggc 1920
tgcccatcat gacctccatc ctgggggcct atgtcatcgc cagcggttcc ttcagcggtt 1980
tcggcatgtg tgtggcacg ctcttctctt gtttctgga agacctggag cggaacaacg 2040
gctccctgga cgggcctac tacatgtoca agagccttct aaagattctg ggcaagaaga 2100
acgaggcgcc cccggacaac aagaagagga agaagtgaca gtcgagccc tgatccagga 2160
ctgcacccca cccccaccgt ccagccatcc aacctcactt cgccttacag gtctccattt 2220
tgtggtaaaa aaaggtttta ggcaggcgc cgtggctcac gctgtaac caacactttg 2280
agaggctgag gcggggcgat cacctgagtc aggagtctga gaccagcctg gccaacatgg 2340
tgaaacctcc gtctctatta aaaatacaaa aattagccga gagggtggc atgcacctgt 2400
catcccagct actcgggagg ctgaggcagg agaatcgctt gaacccggga ggcagaggtt 2460
gcagtgagcc gagatcgcc cactgcactc caacctgggt gacagactct gtctccaaaa 2520
caaaacaaac aaacaaaaag attttattaa agatattttg ttaactcagt aaaaaaaaaa 2580
aaaaaaa 2587

```

<210> SEQ ID NO 11

<211> LENGTH: 710

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 11

```

Met Gly Gly Lys Gln Arg Asp Glu Asp Asp Glu Ala Tyr Gly Lys Pro
 1             5             10             15
Val Lys Tyr Asp Pro Ser Phe Arg Gly Pro Ile Lys Asn Arg Ser Cys
 20             25             30
Thr Asp Val Ile Cys Cys Val Leu Phe Leu Leu Phe Ile Leu Gly Tyr
 35             40             45
Ile Val Val Gly Ile Val Ala Trp Leu Tyr Gly Asp Pro Arg Gln Val
 50             55             60
Leu Tyr Pro Arg Asn Ser Thr Gly Ala Tyr Cys Gly Met Gly Glu Asn
 65             70             75             80
Lys Asp Lys Pro Tyr Leu Leu Tyr Phe Asn Ile Phe Ser Cys Ile Leu
 85             90             95
Ser Ser Asn Ile Ile Ser Val Ala Glu Asn Gly Leu Gln Cys Pro Thr
 100            105            110
Pro Gln Val Cys Val Ser Ser Cys Pro Glu Asp Pro Trp Thr Val Gly
 115            120            125
Lys Asn Glu Phe Ser Gln Thr Val Gly Glu Val Phe Tyr Thr Lys Asn
 130            135            140
Arg Asn Phe Cys Leu Pro Gly Val Pro Trp Asn Met Thr Val Ile Thr
 145            150            155            160

```

Ser	Leu	Gln	Gln	Glu	Leu	Cys	Pro	Ser	Phe	Leu	Leu	Pro	Ser	Ala	Pro
				165					170					175	Pro
Ala	Leu	Gly	Arg	Cys	Phe	Pro	Trp	Thr	Asn	Val	Thr	Pro	Pro	Ala	Leu
				180					185					190	
Pro	Gly	Ile	Thr	Asn	Asp	Thr	Thr	Ile	Gln	Gln	Gly	Ile	Ser	Gly	Leu
				195					200					205	
Ile	Asp	Ser	Leu	Asn	Ala	Arg	Asp	Ile	Ser	Val	Lys	Ile	Phe	Glu	Asp
				210					215					220	
Phe	Ala	Gln	Ser	Trp	Tyr	Trp	Ile	Leu	Val	Ala	Leu	Gly	Val	Ala	Leu
				225					230					235	Leu
Val	Leu	Ser	Leu	Leu	Phe	Ile	Leu	Leu	Leu	Arg	Leu	Val	Ala	Gly	Pro
				245					250					255	
Leu	Val	Leu	Val	Leu	Ile	Leu	Gly	Val	Leu	Gly	Val	Leu	Ala	Tyr	Gly
				260					265					270	
Ile	Tyr	Tyr	Cys	Trp	Glu	Glu	Tyr	Arg	Val	Leu	Arg	Asp	Lys	Gly	Ala
				275					280					285	
Ser	Ile	Ser	Gln	Leu	Gly	Phe	Thr	Thr	Asn	Leu	Ser	Ala	Tyr	Gln	Ser
				290					295					300	
Val	Gln	Glu	Thr	Trp	Leu	Ala	Ala	Leu	Ile	Val	Leu	Ala	Val	Leu	Glu
				305					310					315	320
Ala	Ile	Leu	Leu	Leu	Val	Leu	Ile	Phe	Leu	Arg	Gln	Arg	Ile	Arg	Ile
				325					330					335	
Ala	Ile	Ala	Leu	Leu	Lys	Glu	Ala	Ser	Lys	Ala	Val	Gly	Gln	Met	Met
				340					345					350	
Ser	Thr	Met	Phe	Tyr	Pro	Leu	Val	Thr	Phe	Val	Leu	Leu	Leu	Ile	Cys
				355					360					365	
Ile	Ala	Tyr	Trp	Ala	Met	Thr	Ala	Leu	Tyr	Leu	Ala	Thr	Ser	Gly	Gln
				370					375					380	
Pro	Gln	Tyr	Val	Leu	Trp	Ala	Ser	Asn	Ile	Ser	Ser	Pro	Gly	Cys	Glu
				385					390					395	400
Lys	Val	Pro	Ile	Asn	Thr	Ser	Cys	Asn	Pro	Thr	Ala	His	Leu	Val	Asn
				405					410					415	
Ser	Ser	Cys	Pro	Gly	Leu	Met	Cys	Val	Phe	Gln	Gly	Tyr	Ser	Ser	Lys
				420					425					430	
Gly	Leu	Ile	Gln	Arg	Ser	Val	Phe	Asn	Leu	Gln	Ile	Tyr	Gly	Val	Leu
				435					440					445	
Gly	Leu	Phe	Trp	Thr	Leu	Asn	Trp	Val	Leu	Ala	Leu	Gly	Gln	Cys	Val
				450					455					460	
Leu	Ala	Gly	Ala	Phe	Ala	Ser	Phe	Tyr	Trp	Ala	Phe	His	Lys	Pro	Gln
				465					470					475	480
Asp	Ile	Pro	Thr	Phe	Pro	Leu	Ile	Ser	Ala	Phe	Ile	Arg	Thr	Leu	Arg
				485					490					495	
Tyr	His	Thr	Gly	Ser	Leu	Ala	Phe	Gly	Ala	Leu	Ile	Leu	Thr	Leu	Val
				500					505					510	
Gln	Ile	Ala	Arg	Val	Ile	Leu	Glu	Tyr	Ile	Asp	His	Lys	Leu	Arg	Gly
				515					520					525	
Val	Gln	Asn	Pro	Val	Ala	Arg	Cys	Ile	Met	Cys	Cys	Phe	Lys	Cys	Cys
				530					535					540	
Leu	Trp	Cys	Leu	Glu	Lys	Phe	Ile	Lys	Phe	Leu	Asn	Arg	Asn	Ala	Tyr
				545					550					555	560
Ile	Met	Ile	Ala	Ile	Tyr	Gly	Lys	Asn	Phe	Cys	Val	Ser	Ala	Lys	Asn

-continued

Lys Val Thr Asp Leu Leu Leu Phe Phe Gly Lys Leu Leu Val Val Gly
 595 600 605
 Gly Val Gly Val Leu Ser Phe Phe Phe Phe Ser Gly Arg Ile Pro Gly
 610 615 620
 Leu Gly Lys Asp Phe Lys Ser Pro His Leu Asn Tyr Tyr Trp Leu Pro
 625 630 635 640
 Ile Met Thr Ser Ile Leu Gly Ala Tyr Val Ile Ala Ser Gly Phe Phe
 645 650 655
 Ser Val Phe Gly Met Cys Val Asp Thr Leu Phe Leu Cys Phe Leu Glu
 660 665 670
 Asp Leu Glu Arg Asn Asn Gly Ser Leu Asp Arg Pro Tyr Tyr Met Ser
 675 680 685
 Lys Ser Leu Leu Lys Ile Leu Gly Lys Lys Asn Glu Ala Pro Pro Asp
 690 695 700
 Asn Lys Lys Arg Lys Lys
 705 710

<210> SEQ ID NO 12

<211> LENGTH: 2587

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 12

gagccatggg gggaaagcag cgggacgagg atgacgaggc ctacgggaag ccagtcaaat 60
 acgacccctc ctttcgaggc cccatcaaga acagaagctg cacagatgtc atctgtgtcg 120
 tcctcttctc gctcttcatt ctagggtaca tcgtggtggg gattgtggcc tggttgtatg 180
 gagacccccg gcaagtcttc taccocagga actctactgg ggctactgt ggcattgggg 240
 agaacaaga taagccgtat ctctgtact tcaacatctt cagctgcac ctgtccagca 300
 acatcatctc agttgtctgag aacggcctac agtgccccc accccagggtg tgtgtgtect 360
 cctgcccgga ggacccatgg actgtgggaa aaaacgagtt ctacagact gttggggaag 420
 tctctatac aaaaaacagg aacttttgtc tgccaggggt accctggaat atgacgggtga 480
 tcacaagcct gcaacaggaa ctctgcccc gtttctctct cccctctgct ccagctctgg 540
 ggcgtgctt tccatggacc aacgttactc caccggcgct cccagggatc accaatgaca 600
 ccaccataca gcaggggatc agcgggtctta ttgacagcct caatgcccga gacatcagtg 660
 ttaagatctt tgaagatctt gccagctct ggtattggat tcttgttgcc ctgggggtgg 720
 ctctggtctt gacgctactg tttatcttgc ttctgcgcct ggtggctggg cccctggtgc 780
 tgggtgctgat cctgggagtg ctgggcgtgc tggcatacgg catctactac tgcctggagg 840
 agtaccgagt gctgcgggac aaggggcgct ccatctccca gctgggttcc accaccaacc 900
 tcagtgccta ccagagcgtg caggagacct ggctggccgc cctgatcgtg ttggcgggtgc 960
 ttgaagccat cctgctgctg atgctcatct tcctgcggca gcggattcgt attgccatcg 1020
 ccctcctgaa ggaggccagc aaggctgtgg gacagatgat gtctaccatg ttctaccacc 1080
 tggteacett tgtcctctc ctcatctgca ttgcctaact ggccatgact gctctgtacc 1140
 tggctacatc ggggcaaccc cagtatgtgc tctgggcata caacatcagc tccccgggt 1200
 gtgagaaagt gccaaataat acatcatgca accccacggc ccacctgtg aactcctcgt 1260
 gcccagggt gatgtgcgtc ttccagggt actcatccaa aggcctaata ccacgttctg 1320
 tcttcaatct gcaaatctat ggggtcctgg ggctcttctg gaccctaac tgggtactgg 1380
 ccttgggcca atgcgtcctc gctggagcct ttgcctcctt ctactgggcc ttccacaagc 1440

-continued

```

cccaggacat cctaccttc cccttaatct ctgccttcac cgcacactc cgttaccaca 1500
ctgggtcatt ggcatttga gccctcatcc tgacccttgt gcagatagcc cgggtcatct 1560
tggagtatat tgaccacaag ctgagaggag tgcagaaccc tgtagcccgc tgcacatgt 1620
gctgtttcaa gtctgcctc tgggtgtctgg aaaaatttat caagttccta aaccgcaatg 1680
catacatcat gatcgccatc tacgggaaga atttctgtgt ctacgcaaaa aatgcgttca 1740
tgctactcat gcgaaacatt gtcagggtgg tcgtcttgga caaagtcaca gacctgctgc 1800
tgttcttttg gaagctgctg gtggtcggag gcgtgggggt cctgtccttc ttttttttct 1860
ccggtcgcac cccggggctg ggtaaagact ttaagagccc ccacctcaac tattactggc 1920
tgcccatcat gacctccatc ctgggggect atgtcatcgc cagcggcttc ttcagcgttt 1980
tcggcatgtg tgtggacacg ctcttctctt gcttcttgga agacctggag cggaacaacg 2040
gctccctgga ccggccctac tacatgtcca agagccttct aaagattctg ggcaagaaga 2100
acgaggcgcc cccggacaac aagaagagga agaagtgaca gctccggccc tgatccagga 2160
ctgcacccca cccccaccgt ccagccatcc aacctcactt cgccttacag gtctccattt 2220
tgtggtaaaa aaaggtttta ggccaggcgc cgtggctcac gcctgtaatc caacactttg 2280
agaggctgag gcggggcgat cacctgagtc aggagtctga gaccagctg gccaacatgg 2340
tgaaacctcc gtctctatta aaaatacaaa aattagccga gagtgggtgc atgcacctgt 2400
catcccagct actcgggagg ctgaggcagg agaatcgctt gaaccgggga ggcagaggtt 2460
gcagtgagcc gagatcgcgc cactgcactc caacctgggt gacagactct gtctccaaaa 2520
caaaacaaac aaacaaaaag attttattaa agatatattg ttaactcagt aaaaaaaaaa 2580
aaaaaaa

```

```

<210> SEQ ID NO 13
<211> LENGTH: 710
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 13

```

```

Met Gly Gly Lys Gln Arg Asp Glu Asp Asp Glu Ala Tyr Gly Lys Pro
 1             5             10             15
Val Lys Tyr Asp Pro Ser Phe Arg Gly Pro Ile Lys Asn Arg Ser Cys
 20             25             30
Thr Asp Val Ile Cys Cys Val Leu Phe Leu Leu Phe Ile Leu Gly Tyr
 35             40             45
Ile Val Val Gly Ile Val Ala Trp Leu Tyr Gly Asp Pro Arg Gln Val
 50             55             60
Leu Tyr Pro Arg Asn Ser Thr Gly Ala Tyr Cys Gly Met Gly Glu Asn
 65             70             75             80
Lys Asp Lys Pro Tyr Leu Leu Tyr Phe Asn Ile Phe Ser Cys Ile Leu
 85             90             95
Ser Ser Asn Ile Ile Ser Val Ala Glu Asn Gly Leu Gln Cys Pro Thr
100            105            110
Pro Gln Val Cys Val Ser Ser Cys Pro Glu Asp Pro Trp Thr Val Gly
115            120            125
Lys Asn Glu Phe Ser Gln Thr Val Gly Glu Val Phe Tyr Thr Lys Asn
130            135            140
Arg Asn Phe Cys Leu Pro Gly Val Pro Trp Asn Met Thr Val Ile Thr
145            150            155            160
Ser Leu Gln Gln Glu Leu Cys Pro Ser Phe Leu Leu Pro Ser Ala Pro

```

-continued

165							170					175				
Ala	Leu	Gly	Arg	Cys	Phe	Pro	Trp	Thr	Asn	Val	Thr	Pro	Pro	Ala	Leu	
			180					185					190			
Pro	Gly	Ile	Thr	Asn	Asp	Thr	Thr	Ile	Gln	Gln	Gly	Ile	Ser	Gly	Leu	
		195					200					205				
Ile	Asp	Ser	Leu	Asn	Ala	Arg	Asp	Ile	Ser	Val	Lys	Ile	Phe	Glu	Asp	
	210					215					220					
Phe	Ala	Gln	Ser	Trp	Tyr	Trp	Ile	Leu	Val	Ala	Leu	Gly	Val	Ala	Leu	
	225				230					235					240	
Val	Leu	Ser	Leu	Leu	Phe	Ile	Leu	Leu	Leu	Arg	Leu	Val	Ala	Gly	Pro	
			245						250					255		
Leu	Val	Leu	Val	Leu	Ile	Leu	Gly	Val	Leu	Gly	Val	Leu	Ala	Tyr	Gly	
		260					265						270			
Ile	Tyr	Tyr	Cys	Trp	Glu	Glu	Tyr	Arg	Val	Leu	Arg	Asp	Lys	Gly	Ala	
	275						280					285				
Ser	Ile	Ser	Gln	Leu	Gly	Phe	Thr	Thr	Asn	Leu	Ser	Ala	Tyr	Gln	Ser	
	290					295					300					
Val	Gln	Glu	Thr	Trp	Leu	Ala	Ala	Leu	Ile	Val	Leu	Ala	Val	Leu	Glu	
	305				310					315					320	
Ala	Ile	Leu	Leu	Leu	Met	Leu	Ile	Phe	Leu	Arg	Gln	Arg	Ile	Arg	Ile	
			325					330						335		
Ala	Ile	Ala	Leu	Leu	Lys	Glu	Ala	Ser	Lys	Ala	Val	Gly	Gln	Met	Met	
		340						345						350		
Ser	Thr	Met	Phe	Tyr	Pro	Leu	Val	Thr	Phe	Val	Leu	Leu	Leu	Ile	Cys	
		355					360					365				
Ile	Ala	Tyr	Trp	Ala	Met	Thr	Ala	Leu	Tyr	Leu	Ala	Thr	Ser	Gly	Gln	
	370					375						380				
Pro	Gln	Tyr	Val	Leu	Trp	Ala	Ser	Asn	Ile	Ser	Ser	Pro	Gly	Cys	Glu	
	385				390					395					400	
Lys	Val	Pro	Ile	Asn	Thr	Ser	Cys	Asn	Pro	Thr	Ala	His	Leu	Val	Asn	
			405						410					415		
Ser	Ser	Cys	Pro	Gly	Leu	Met	Cys	Val	Phe	Gln	Gly	Tyr	Ser	Ser	Lys	
			420					425					430			
Gly	Leu	Ile	Pro	Arg	Ser	Val	Phe	Asn	Leu	Gln	Ile	Tyr	Gly	Val	Leu	
		435					440					445				
Gly	Leu	Phe	Trp	Thr	Leu	Asn	Trp	Val	Leu	Ala	Leu	Gly	Gln	Cys	Val	
	450					455						460				
Leu	Ala	Gly	Ala	Phe	Ala	Ser	Phe	Tyr	Trp	Ala	Phe	His	Lys	Pro	Gln	
	465				470					475					480	
Asp	Ile	Pro	Thr	Phe	Pro	Leu	Ile	Ser	Ala	Phe	Ile	Arg	Thr	Leu	Arg	
			485					490						495		
Tyr	His	Thr	Gly	Ser	Leu	Ala	Phe	Gly	Ala	Leu	Ile	Leu	Thr	Leu	Val	
		500						505					510			
Gln	Ile	Ala	Arg	Val	Ile	Leu	Glu	Tyr	Ile	Asp	His	Lys	Leu	Arg	Gly	
		515					520					525				
Val	Gln	Asn	Pro	Val	Ala	Arg	Cys	Ile	Met	Cys	Cys	Phe	Lys	Cys	Cys	
	530					535						540				
Leu	Trp	Cys	Leu	Glu	Lys	Phe	Ile	Lys	Phe	Leu	Asn	Arg	Asn	Ala	Tyr	
	545				550					555					560	
Ile	Met	Ile	Ala	Ile	Tyr	Gly	Lys	Asn	Phe	Cys	Val	Ser	Ala	Lys	Asn	
			565					570						575		
Ala	Phe	Met	Leu	Leu	Met	Arg	Asn	Ile	Val	Arg	Val	Val	Val	Leu	Asp	
			580					585						590		

-continued

Lys Val Thr Asp Leu Leu Leu Phe Phe Gly Lys Leu Leu Val Val Gly
 595 600 605

 Gly Val Gly Val Leu Ser Phe Phe Phe Phe Ser Gly Arg Ile Pro Gly
 610 615 620

 Leu Gly Lys Asp Phe Lys Ser Pro His Leu Asn Tyr Tyr Trp Leu Pro
 625 630 635 640

 Ile Met Thr Ser Ile Leu Gly Ala Tyr Val Ile Ala Ser Gly Phe Phe
 645 650 655

 Ser Val Phe Gly Met Cys Val Asp Thr Leu Phe Leu Cys Phe Leu Glu
 660 665 670

 Asp Leu Glu Arg Asn Asn Gly Ser Leu Asp Arg Pro Tyr Tyr Met Ser
 675 680 685

 Lys Ser Leu Leu Lys Ile Leu Gly Lys Lys Asn Glu Ala Pro Pro Asp
 690 695 700

 Asn Lys Lys Arg Lys Lys
 705 710

<210> SEQ ID NO 14

<211> LENGTH: 2251

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 14

gagccatggg gggaaagcag cgggacgagg atgacgaggc ctacgggaag ccagtcaa	60
acgacccctc ctttcgaggc cccatcaaga acagaagctg cacagatgtc atctgtgtcg	120
tcctcttctc gctcttcatt ctagggtaca tcgtggtggg gattgtggcc tggttgtatg	180
gagacccccg gcaagtcctc taccacagga actctactgg ggcctactgt ggcacggggg	240
agaacaaaga taagccgtat ctctgtact tcaacatctt cagctgcac cgttcacagca	300
acatcatctc agttgtgtag aacggcctac agtgccccac acccaggtg tgtgtgtcct	360
cctgcccga ggacccatgg actgtgggaa aaaacgagtt ctcacagact gttggggaag	420
tcctctatac aaaaaacagg aacttttgtc tgccaggggg accctggaat atgacgggtga	480
tcacaagcct gcaacaggaa ctctgcccc gtttctctct cccctctgct ccagctctgg	540
ggcgctgctt tccatggacc aacgttactc caccggcgct ccaggggac accaatgaca	600
ccaccatata gcaggggac agcgggtctta ttgacagcct caatgcccga gacatcagtg	660
ttaagatctt tgaagatctt gccagtcct ggtattggat tctgtgggt gtgggacaga	720
tgatgtctac catgtttctc ccaactggta cctttgtctc cctctctc tgcattgcct	780
actgggcat gactgtctg tacctggcta catcggggca acccagtat gtgctctggg	840
catccaacat cagctcccc ggctgtgaga aagtgccaat aaatacatca tgcaaccca	900
cgccccact tgtgaactcc tcgtgccag ggetgatgtg cgtcttcag ggetactcat	960
ccaaaggcct aatccaacgt tctgtcttca atctgcaaat ctatggggtc ctggggctct	1020
tctggacct taactgggta ctggccctgg gccaatgctg cctcgtgga gcctttgcct	1080
cctctactg ggccttccac aagccccagg acatccctac ctccccctta atctctgcct	1140
tcatecgac actccgttac cacactgggt cattggcatt tggagccctc atcctgaccc	1200
ttgtgcagat agccccgggc atcttgagat atattgacca caagctcaga ggagtgcaga	1260
acctgtagc ccgtgcac atgtgtgtt tcaagtgtg cctctggtgt ctggaaaaat	1320
ttatcaagtt cctaaaccgc aatgcataca tcatgatcgc catctacggg aagaatttct	1380
gtgtctcagc caaaaatgcg ttcagtctac tcatgcgaaa cattgtcagg gtggctgtcc	1440

-continued

```

tggacaaagt cacagacctg ctgctgttct ttggaagct gctggtggtc ggaggcgtgg 1500
gggtcctgtc cttctttttt ttctccggtc gcatcccggg gctgggtaaa gactttaaga 1560
gccccacact caactattac tggctgccc tcatgacctc catcctgggg gcctatgtca 1620
tcgccagcgg cttcttcagc gttttcggca tgtgtgtgga cacgtcttcc ctctgettcc 1680
tggaaagacct ggagcggaac aacggctccc tggaccggcc ctactacatg tccaagagcc 1740
ttctaaagat tctgggcaag aagaacgagg cgcctccgga caacaagaag aggaagaagt 1800
gacagctccg gcctgatcc aggactgcac cccaccccca ccgtccagcc atccaacctc 1860
acttcgctt acaggtctcc atttgtggt aaaaaaagg tttaggccag gcgccgtggc 1920
tcacgcctgt aatccaacac tttgagaggc tgaggcgggc ggatcacctg agtcaggagt 1980
tcgagaccag cctggccaac atggtgaaac ctccgtctct attaaaaata caaaaattag 2040
ccgagagtgg tggcatgcac ctgtcatccc agctactcgg gaggtgagg caggagaatc 2100
gcttgaaccc gggaggcaga ggttgacgtg agccgagatc gcgccactgc actccaacct 2160
gggtgacaga ctctgtctcc aaaacaaaac aaacaacaa aaagatttta ttaaagatat 2220
tttgtaact cagtaaaaaa aaaaaaaaaa a 2251

```

<210> SEQ ID NO 15

<211> LENGTH: 598

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 15

```

Met Gly Gly Lys Gln Arg Asp Glu Asp Asp Glu Ala Tyr Gly Lys Pro
 1             5             10             15
Val Lys Tyr Asp Pro Ser Phe Arg Gly Pro Ile Lys Asn Arg Ser Cys
 20            25            30
Thr Asp Val Ile Cys Cys Val Leu Phe Leu Leu Phe Ile Leu Gly Tyr
 35            40            45
Ile Val Val Gly Ile Val Ala Trp Leu Tyr Gly Asp Pro Arg Gln Val
 50            55            60
Leu Tyr Pro Arg Asn Ser Thr Gly Ala Tyr Cys Gly Met Gly Glu Asn
 65            70            75            80
Lys Asp Lys Pro Tyr Leu Leu Tyr Phe Asn Ile Phe Ser Cys Ile Leu
 85            90            95
Ser Ser Asn Ile Ile Ser Val Ala Glu Asn Gly Leu Gln Cys Pro Thr
100           105           110
Pro Gln Val Cys Val Ser Ser Cys Pro Glu Asp Pro Trp Thr Val Gly
115           120           125
Lys Asn Glu Phe Ser Gln Thr Val Gly Glu Val Phe Tyr Thr Lys Asn
130           135           140
Arg Asn Phe Cys Leu Pro Gly Val Pro Trp Asn Met Thr Val Ile Thr
145           150           155           160
Ser Leu Gln Gln Glu Leu Cys Pro Ser Phe Leu Leu Pro Ser Ala Pro
165           170           175
Ala Leu Gly Arg Cys Phe Pro Trp Thr Asn Val Thr Pro Pro Ala Leu
180           185           190
Pro Gly Ile Thr Asn Asp Thr Thr Ile Gln Gln Gly Ile Ser Gly Leu
195           200           205
Ile Asp Ser Leu Asn Ala Arg Asp Ile Ser Val Lys Ile Phe Glu Asp
210           215           220
Phe Ala Gln Ser Trp Tyr Trp Ile Leu Val Ala Val Gly Gln Met Met
225           230           235           240

```

Figure 3:

Figure 3A. Amino acid sequence of 24P4C12 v.1 (SEQ ID. NO. : 20). The 24P4C12 v.1 protein has 710 amino acids.

```
1  MGGKQRDEDD EAYGKPVKYD PSFRGPIKNR SCTDVICCVL FLLFILGYIV VGIVAWLYGD
61  PRQVLYPRNS TGAYCGMGEN KDKPYLLYFN IFSCILSSNI ISVAENGLQC PTPQVCVSSC
121 PEDPWTVGKN EFSQTVGEVF YTKNRNFCLP GVPWNMTVIT SLQQELCPSF LLPSAPALGR
181 CFPWTNVTTP ALPGITNDTT IQQGISGLID SLNARDISVK IFEDFAQSWY WILVALGVAL
241 VLSLLFILLL RLVAGPLVLV LILGVLGVLA YGIYYCWEEY RVLDRKGASI SQLGFTTNLS
301 AYQSVQETWL AALIVLAVLE AILLMLIFL RQRIRIAIAL LKEASKAVGQ MMSTMFYPLV
361 TFVLLLICIA YWAMTALYLA TSGQPQYVLW ASNISSPGCE KVPINTSCNP TAHLVNSSCP
421 GLMCVFQGYG SKGLIQRSVF NLQIYGVGLG FWTLNWLAL GQCVLAGAFA SFYWAFHKPQ
481 DIPTFPLISA FIRTLRYHTG SLAFGALILT LVQIARVILE YIDHKLRGVQ NPVARCIMCC
541 FKCLWCLEK FIKFLNRNAY IMIAIYGKNF CVSAKNAFML LMRNIVRVVV LDKVTDLLLF
601 FGKLLVVGCV GVLSFFFFSG RIPGLGKDFK SPHLNYYWLP IMTSILGAYV IASGFFSVFG
661 MCVDTLFLCF LEDLERNNGS LDRPYMSKS LLKILGKKNE APPDNKKRKK
```

Figure 3B. Amino acid sequence of 24P4C12 v.3 (SEQ ID. NO. : 21). The 24P4C12 v.3 protein has 710 amino acids.

```
1  MGGKQRDEDD EAYGKPVKYD PSFRGPIKNR SCTDVICCVL FLLFILGYIV VGIVAWLYGD
61  PRQVLYPRNS TGAYCGMGEN KDKPYLLYFN IFSCILSSNI ISVAENGLQC PTPQVCVSSC
121 PEDPWTVGKN EFSQTVGEVF YTKNRNFCLP GVPWNMTVIT SLQQELCPSF LLPSAPALGR
181 CFPWTNITPP ALPGITNDTT IQQGISGLID SLNARDISVK IFEDFAQSWY WILVALGVAL
241 VLSLLFILLL RLVAGPLVLV LILGVLGVLA YGIYYCWEEY RVLDRKGASI SQLGFTTNLS
301 AYQSVQETWL AALIVLAVLE AILLMLIFL RQRIRIAIAL LKEASKAVGQ MMSTMFYPLV
361 TFVLLLICIA YWAMTALYLA TSGQPQYVLW ASNISSPGCE KVPINTSCNP TAHLVNSSCP
421 GLMCVFQGYG SKGLIQRSVF NLQIYGVGLG FWTLNWLAL GQCVLAGAFA SFYWAFHKPQ
481 DIPTFPLISA FIRTLRYHTG SLAFGALILT LVQIARVILE YIDHKLRGVQ NPVARCIMCC
541 FKCLWCLEK FIKFLNRNAY IMIAIYGKNF CVSAKNAFML LMRNIVRVVV LDKVTDLLLF
601 FGKLLVVGCV GVLSFFFFSG RIPGLGKDFK SPHLNYYWLP IMTSILGAYV IASGFFSVFG
661 MCVDTLFLCF LEDLERNNGS LDRPYMSKS LLKILGKKNE APPDNKKRKK
```

Figure 3C. Amino acid sequence of 24P4C12 v.5 (SEQ ID. NO. : 22). The 24P4C12 v.5 protein has 710 amino acids.

Figure 3C-1

```
1  MGGKQRDEDD EAYGKPVKYD PSFRGPIKNR SCTDVICCVL FLLFILGYIV VGIVAWLYGD
61  PRQVLYPRNS TGAYCGMGEN KDKPYLLYFN IFSCILSSNI ISVAENGLQC PTPQVCVSSC
121 PEDPWTVGKN EFSQTVGEVF YTKNRNFCLP GVPWNMTVIT SLQQELCPSF LLPSAPALGR
181 CFPWTNVTTP ALPGITNDTT IQQGISGLID SLNARDISVK IFEDFAQSWY WILVALGVAL
241 VLSLLFILLL RLVAGPLVLV LILGVLGVLA YGIYYCWEEY RVLDRKGASI SQLGFTTNLS
301 AYQSVQETWL AALIVLAVLE AILLVLIFL RQRIRIAIAL LKEASKAVGQ MMSTMFYPLV
361 TFVLLLICIA YWAMTALYLA TSGQPQYVLW ASNISSPGCE KVPINTSCNP TAHLVNSSCP
421 GLMCVFQGYG SKGLIQRSVF NLQIYGVGLG FWTLNWLAL GQCVLAGAFA SFYWAFHKPQ
```

-continued

Ser Thr Met Phe Tyr Pro Leu Val Thr Phe Val Leu Leu Leu Ile Cys
 245 250 255
 Ile Ala Tyr Trp Ala Met Thr Ala Leu Tyr Leu Ala Thr Ser Gly Gln
 260 265 270
 Pro Gln Tyr Val Leu Trp Ala Ser Asn Ile Ser Ser Pro Gly Cys Glu
 275 280 285
 Lys Val Pro Ile Asn Thr Ser Cys Asn Pro Thr Ala His Leu Val Asn
 290 295 300
 Ser Ser Cys Pro Gly Leu Met Cys Val Phe Gln Gly Tyr Ser Ser Lys
 305 310 315 320
 Gly Leu Ile Gln Arg Ser Val Phe Asn Leu Gln Ile Tyr Gly Val Leu
 325 330 335
 Gly Leu Phe Trp Thr Leu Asn Trp Val Leu Ala Leu Gly Gln Cys Val
 340 345 350
 Leu Ala Gly Ala Phe Ala Ser Phe Tyr Trp Ala Phe His Lys Pro Gln
 355 360 365
 Asp Ile Pro Thr Phe Pro Leu Ile Ser Ala Phe Ile Arg Thr Leu Arg
 370 375 380
 Tyr His Thr Gly Ser Leu Ala Phe Gly Ala Leu Ile Leu Thr Leu Val
 385 390 395 400
 Gln Ile Ala Arg Val Ile Leu Glu Tyr Ile Asp His Lys Leu Arg Gly
 405 410 415
 Val Gln Asn Pro Val Ala Arg Cys Ile Met Cys Cys Phe Lys Cys Cys
 420 425 430
 Leu Trp Cys Leu Glu Lys Phe Ile Lys Phe Leu Asn Arg Asn Ala Tyr
 435 440 445
 Ile Met Ile Ala Ile Tyr Gly Lys Asn Phe Cys Val Ser Ala Lys Asn
 450 455 460
 Ala Phe Met Leu Leu Met Arg Asn Ile Val Arg Val Val Val Leu Asp
 465 470 475 480
 Lys Val Thr Asp Leu Leu Leu Phe Phe Gly Lys Leu Leu Val Val Gly
 485 490 495
 Gly Val Gly Val Leu Ser Phe Phe Phe Phe Ser Gly Arg Ile Pro Gly
 500 505 510
 Leu Gly Lys Asp Phe Lys Ser Pro His Leu Asn Tyr Tyr Trp Leu Pro
 515 520 525
 Ile Met Thr Ser Ile Leu Gly Ala Tyr Val Ile Ala Ser Gly Phe Phe
 530 535 540
 Ser Val Phe Gly Met Cys Val Asp Thr Leu Phe Leu Cys Phe Leu Glu
 545 550 555 560
 Asp Leu Glu Arg Asn Asn Gly Ser Leu Asp Arg Pro Tyr Tyr Met Ser
 565 570 575
 Lys Ser Leu Leu Lys Ile Leu Gly Lys Lys Asn Glu Ala Pro Pro Asp
 580 585 590
 Asn Lys Lys Arg Lys Lys
 595

<210> SEQ ID NO 16
 <211> LENGTH: 2623
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 16

gagccatggg gggaaagcag cgggacgagg atgacgaggc ctacgggaag ccagtcaaat

60

-continued

acgacccctc ctttcgaggc cccatcaaga acagaagctg cacagatgtc atctgtctgcg	120
tcctcttctc gctcttcatt ctaggttaca tcgtggtggg gattgtggcc tggttgtatg	180
gagacccccg gcaagtcttc taccocagga actctactgg ggctactgt ggcacggggg	240
agaacaaaga taagccgtat ctctgtact tcaacatctt cagctgcac cgtccagca	300
acatcatctc agttgctgag aacggcctac agtgccccac accccagggtg tgtgtgtect	360
cctgcccgga ggacccatgg actgtgggaa aaaacgagtt ctacacagact gttggggaag	420
tcttctatac aaaaaacagg aacttttgtc tgccagggtt accctggaat atgacgggtga	480
tcacaagcct gcaacaggaa ctctgcccc gtttctctc cccctctgct ccagctctgg	540
ggcgctgctt tccatggacc aacgttactc caccggcgct ccagggtat accaatgaca	600
ccaccataca gcaggggatc agcgggtctta ttgacagcct caatgcccga gacatcagtg	660
ttaagatctt tgaagatctt gccagtcct ggtattggat tctgtgtgcc ctgggggtgg	720
ctctggtctt gacgctactg tttatcttgc ttctgcgctt ggtggctggg cccctggtgc	780
tgggtctgat cctgggagtg ctgggcgtgc tggcatacgg catctactac tgcctggagg	840
agtaccgagt gctgcgggac aaggcgccct ccactcccca gctgggttcc accaccaacc	900
tcagtgccta ccagagcgtg caggagacct ggctggccgc cctgatcgtg ttggcgggtgc	960
ttgaagccat cctgctgctg atgctcatct tcctgcggca gcggtatcgt attgccatcg	1020
ccctcctgaa ggaggccagc aaggctgtgg gacagatgat gtctaccatg ttctaccac	1080
tggtcacett tgcctcctc ctcatctgca ttgcctactg ggccatgact gctctgtacc	1140
tggctacatc ggggcaaccc cagtatgtgc tctgggcac caacatcagc tccccgggt	1200
gtgagaaagt gccataaat acatcatgca accccacggc ccaccttggt aactcctcgt	1260
gcccagggtc gatgtgcgtc ttccagggtc actcatccaa aggcctaacc caacgttctg	1320
tcttcaatct gcaaatctat ggggtcctgg ggctcttctg gaccctaac tgggtactgg	1380
ccctgggcca atgcgtcctc gctggagcct ttgcctcctt ctactgggcc ttccacaagc	1440
cccaggacat ccctaccttc cccttaatct ctgccttcat ccgcacactc cgttaccaca	1500
ctgggtcatt ggcatttggg gccctcatcc tgaccttgt gcagatagcc cgggtcatct	1560
tggagtatat tgaccacaag ctacagaggag tgcagaaccc tgtagccgc tgcacatgt	1620
gctgtttcaa gtgctgcctc tgggtgtctgg aaaaatttat caagttccta aaccgcaatg	1680
catacatcat gatcgccatc tacgggaaga atttctgtgt ctacgcaaaa aatgcgttca	1740
tgtactcatc gcgaacatt gtcagggtgg tcgtcctgga caaagtcaca gacctgctgc	1800
tgttcttttg gaagctgctg gtggtcggag gcgtgggggt cctgtccttc tttttttct	1860
ccggtgcgat cccgggggtg ggtaagact ttaagagccc ccacctcaac tattactggc	1920
tgcccatcat gaggaaccca ataaccacca cgggtcatgt ctccagacc tccatcctgg	1980
gggcctatgt catcgccagc ggcttcttca gcgttttcgg catgtgtgtg gacacgtct	2040
tcctctgctt cctggaagac ctggagcgga acaacggctc cctggaccgg ccctactaca	2100
tgtccaagag ccttctaaag attctgggca agaagaacga ggcgcccccg gacaacaaga	2160
agaggaagaa gtgacagctc cggccctgat ccaggactgc accccacccc caccgtcag	2220
ccatccaacc tcacttcgcc ttacaggctc ccattttgtg gtaaaaaaag gttttaggcc	2280
aggcgccgtg gctcacgctt gtaatccaac actttgagag gctgaggcgg gcggtaccc	2340
tgagtcagga gtctgagacc agcctggcca acatggtgaa acctcgtct ctattaaaaa	2400
tacaaaaatt agccgagagt ggtggcatgc acctgtcatc ccagctactc gggaggctga	2460

-continued

```

ggcaggagaa tcgcttgaac ccgggaggca gaggttgacg tgagccgaga tcgcgccact 2520
gcactccaac ctgggtgaca gactctgtct ccaaaacaaa acaaacaaac aaaagattt 2580
tattaaagat attttgtaa ctcagtaaaa aaaaaaaaaa aaa 2623

```

<210> SEQ ID NO 17

<211> LENGTH: 722

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 17

```

Met Gly Gly Lys Gln Arg Asp Glu Asp Asp Glu Ala Tyr Gly Lys Pro
 1           5           10           15

Val Lys Tyr Asp Pro Ser Phe Arg Gly Pro Ile Lys Asn Arg Ser Cys
 20           25           30

Thr Asp Val Ile Cys Cys Val Leu Phe Leu Leu Phe Ile Leu Gly Tyr
 35           40           45

Ile Val Val Gly Ile Val Ala Trp Leu Tyr Gly Asp Pro Arg Gln Val
 50           55           60

Leu Tyr Pro Arg Asn Ser Thr Gly Ala Tyr Cys Gly Met Gly Glu Asn
 65           70           75           80

Lys Asp Lys Pro Tyr Leu Leu Tyr Phe Asn Ile Phe Ser Cys Ile Leu
 85           90           95

Ser Ser Asn Ile Ile Ser Val Ala Glu Asn Gly Leu Gln Cys Pro Thr
100           105           110

Pro Gln Val Cys Val Ser Ser Cys Pro Glu Asp Pro Trp Thr Val Gly
115           120           125

Lys Asn Glu Phe Ser Gln Thr Val Gly Glu Val Phe Tyr Thr Lys Asn
130           135           140

Arg Asn Phe Cys Leu Pro Gly Val Pro Trp Asn Met Thr Val Ile Thr
145           150           155           160

Ser Leu Gln Gln Glu Leu Cys Pro Ser Phe Leu Leu Pro Ser Ala Pro
165           170           175

Ala Leu Gly Arg Cys Phe Pro Trp Thr Asn Val Thr Pro Pro Ala Leu
180           185           190

Pro Gly Ile Thr Asn Asp Thr Thr Ile Gln Gln Gly Ile Ser Gly Leu
195           200           205

Ile Asp Ser Leu Asn Ala Arg Asp Ile Ser Val Lys Ile Phe Glu Asp
210           215           220

Phe Ala Gln Ser Trp Tyr Trp Ile Leu Val Ala Leu Gly Val Ala Leu
225           230           235           240

Val Leu Ser Leu Leu Phe Ile Leu Leu Leu Arg Leu Val Ala Gly Pro
245           250           255

Leu Val Leu Val Leu Ile Leu Gly Val Leu Gly Val Leu Ala Tyr Gly
260           265           270

Ile Tyr Tyr Cys Trp Glu Glu Tyr Arg Val Leu Arg Asp Lys Gly Ala
275           280           285

Ser Ile Ser Gln Leu Gly Phe Thr Thr Asn Leu Ser Ala Tyr Gln Ser
290           295           300

Val Gln Glu Thr Trp Leu Ala Ala Leu Ile Val Leu Ala Val Leu Glu
305           310           315           320

Ala Ile Leu Leu Leu Met Leu Ile Phe Leu Arg Gln Arg Ile Arg Ile
325           330           335

Ala Ile Ala Leu Leu Lys Glu Ala Ser Lys Ala Val Gly Gln Met Met
340           345           350

```

-continued

Ser	Thr	Met	Phe	Tyr	Pro	Leu	Val	Thr	Phe	Val	Leu	Leu	Leu	Ile	Cys
		355					360					365			
Ile	Ala	Tyr	Trp	Ala	Met	Thr	Ala	Leu	Tyr	Leu	Ala	Thr	Ser	Gly	Gln
	370					375					380				
Pro	Gln	Tyr	Val	Leu	Trp	Ala	Ser	Asn	Ile	Ser	Ser	Pro	Gly	Cys	Glu
385					390					395					400
Lys	Val	Pro	Ile	Asn	Thr	Ser	Cys	Asn	Pro	Thr	Ala	His	Leu	Val	Asn
			405						410					415	
Ser	Ser	Cys	Pro	Gly	Leu	Met	Cys	Val	Phe	Gln	Gly	Tyr	Ser	Ser	Lys
			420						425				430		
Gly	Leu	Ile	Gln	Arg	Ser	Val	Phe	Asn	Leu	Gln	Ile	Tyr	Gly	Val	Leu
	435						440					445			
Gly	Leu	Phe	Trp	Thr	Leu	Asn	Trp	Val	Leu	Ala	Leu	Gly	Gln	Cys	Val
	450					455					460				
Leu	Ala	Gly	Ala	Phe	Ala	Ser	Phe	Tyr	Trp	Ala	Phe	His	Lys	Pro	Gln
465					470					475					480
Asp	Ile	Pro	Thr	Phe	Pro	Leu	Ile	Ser	Ala	Phe	Ile	Arg	Thr	Leu	Arg
				485					490					495	
Tyr	His	Thr	Gly	Ser	Leu	Ala	Phe	Gly	Ala	Leu	Ile	Leu	Thr	Leu	Val
			500					505					510		
Gln	Ile	Ala	Arg	Val	Ile	Leu	Glu	Tyr	Ile	Asp	His	Lys	Leu	Arg	Gly
	515						520					525			
Val	Gln	Asn	Pro	Val	Ala	Arg	Cys	Ile	Met	Cys	Cys	Phe	Lys	Cys	Cys
	530					535					540				
Leu	Trp	Cys	Leu	Glu	Lys	Phe	Ile	Lys	Phe	Leu	Asn	Arg	Asn	Ala	Tyr
545					550					555					560
Ile	Met	Ile	Ala	Ile	Tyr	Gly	Lys	Asn	Phe	Cys	Val	Ser	Ala	Lys	Asn
			565					570						575	
Ala	Phe	Met	Leu	Leu	Met	Arg	Asn	Ile	Val	Arg	Val	Val	Val	Leu	Asp
			580				585						590		
Lys	Val	Thr	Asp	Leu	Leu	Leu	Phe	Phe	Gly	Lys	Leu	Leu	Val	Val	Gly
		595					600					605			
Gly	Val	Gly	Val	Leu	Ser	Phe	Phe	Phe	Phe	Ser	Gly	Arg	Ile	Pro	Gly
	610					615					620				
Leu	Gly	Lys	Asp	Phe	Lys	Ser	Pro	His	Leu	Asn	Tyr	Tyr	Trp	Leu	Pro
625					630					635					640
Ile	Met	Arg	Asn	Pro	Ile	Thr	Pro	Thr	Gly	His	Val	Phe	Gln	Thr	Ser
			645						650					655	
Ile	Leu	Gly	Ala	Tyr	Val	Ile	Ala	Ser	Gly	Phe	Phe	Ser	Val	Phe	Gly
			660					665					670		
Met	Cys	Val	Asp	Thr	Leu	Phe	Leu	Cys	Phe	Leu	Glu	Asp	Leu	Glu	Arg
		675					680					685			
Asn	Asn	Gly	Ser	Leu	Asp	Arg	Pro	Tyr	Tyr	Met	Ser	Lys	Ser	Leu	Leu
	690					695					700				
Lys	Ile	Leu	Gly	Lys	Lys	Asn	Glu	Ala	Pro	Pro	Asp	Asn	Lys	Lys	Arg
705					710					715					720
Lys	Lys														

<210> SEQ ID NO 18

<211> LENGTH: 2593

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 18

gagccatggg gggaaagcag cgggacgagg atgacgaggc ctacgggaag ccagtcaaat

60

-continued

acgacccctc ctttcgaggc cccatcaaga acagaagctg cacagatgtc atctgctgcg	120
tcctcttctc gctcttcatt ctagggtaca tcgtggtggg gattgtggcc tggttgtatg	180
gagacccccg gcaagtcctc taccccagga actctactgg ggctactgt ggcatggggg	240
agaacaaaga taagccgtat ctctgtact tcaacatctt cagctgcac cgttcacga	300
acatcatctc agttgtgag aacggcctac agtgcctcac acccagggtg tgtgtgtcct	360
cctgcccga ggaccatgg actgtggga aaaacgagtt ctacacagact gttggggaag	420
tctctatcac aaaaaacagg aacttttgc tgccagggt accctggaat atgacggtga	480
tcacaagcct gcaacaggaa ctctgcccc gtttctctc cccctctgct ccagctctgg	540
ggcgtgctt tccatggacc aacgttact caccggcgt cccagggtc accaatgaca	600
ccaccatata gcaggggatc agcgggtctta ttgacagcct caatgcccga gacatcagtg	660
ttaagatctt tgaagatttt gccagtcct ggtattggat tcttgttgcc ctgggggtgg	720
ctctggtctt gagcctactg tttatcttgc ttctgcgct ggtggctggg ccctggtgc	780
tgggtgctgat cctgggagtg ctgggctgct tggcatacgg catctactac tctgggagg	840
agtaccgagt gctgcgggac aagggcgct ccatctccca gctgggttc accaccaacc	900
tcagtgccta ccagagcgtg caggagacct ggctggcgc cctgatcgtg ttggcggtgc	960
ttgaagccat cctgctgctg atgctcatct tcctgcggca gcggattcgt attgccatcg	1020
ccctcctgaa ggaggccagc aaggctgtgg gacagatgat gtctaccatg ttctaccac	1080
tggtcacctt tgtctctc ctcactgca ttgctactg ggccatgact gctctgtatc	1140
ctctgccac gcagccagcc actcttgat atgtgctctg ggcatccaac atcagctccc	1200
ccgctgtga gaaagtcca ataaatacat catgcaacc caccggccac cttgtgaact	1260
cctcgtgcc agggctgatg tgcgtcttc agggctact atccaaagg ctaatccaac	1320
gttctgtctt caatctgcaa atctatggg tcctggggt cttctggacc cttaactggg	1380
tactggcctt gggccaatgc gtctcgtg gagcctttgc ctctctctac tgggccttc	1440
acaagcccca ggacatccct accctccct taatctctgc ctctatccgc acactccgtt	1500
accacactgg gtcattggca tttggagccc tcactctgac cctgtgacag atagccggg	1560
tcactcttga gtatattgac cacaagctca gaggagtga gaacctgta gcccgctga	1620
tcactgtctg tttcaagtgc tgctctggt gtctggaaaa atttatcaag ttctaaacc	1680
gcaatgcata catcatgat gccatctac ggaagaattt ctgtgtctca gcaaaaatg	1740
cgttcagtct actcatgca aacattgtca ggtggtcgt cctggacaaa gtcacagacc	1800
tgtcgtgctt ctttgggaag ctgctggtg tcggaggcgt gggggtcctg tcctctttt	1860
ttttctccg tcgcatccc gggctggga aagacttta gagccccac ctcaactatt	1920
actggctgcc catcatgac tccatcctgg gggcctatgt catcgccagc ggcttcttca	1980
gcgttttcgg catgtgtgtg gacacgtct tcctctgctt cctggaagac ctggagcgga	2040
acaacggctc cctggaccgg ccctactaca tgtccaagag ccttctaaag attctgggca	2100
agaagaacga ggcgccccg gacaacaaga agaggaagaa gtgacagctc cggccctgat	2160
ccaggactgc accccaccc caccgtccag ccatccaacc tcacttcgcc ttacagggtt	2220
ccattttgtg gtaaaaaaag gttttaggcc aggcgcctg gctcacgct gtaatccaac	2280
actttgagag gctgaggcgg cgggatcacc tgagtcagga gttcgagacc agcctggcca	2340
acatggtgaa acctccgtct ctattaaaaa tacaaaaatt agccgagagt ggtggcatgc	2400
acctgtcatc ccagctactc gggaggctga ggcaggagaa tcgcttgaac ccgggagga	2460

-continued

```

gaggttgacg tgagccgaga tcgcgccact gcactccaac ctgggtgaca gactctgtct 2520
ccaaaacaaa acaaacacaa aaaaagattt tattaagat atttgttaa ctcagtaaaa 2580
aaaaaaaaaa aaa 2593

```

```

<210> SEQ ID NO 19
<211> LENGTH: 712
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 19

```

```

Met Gly Gly Lys Gln Arg Asp Glu Asp Asp Glu Ala Tyr Gly Lys Pro
 1             5             10             15
Val Lys Tyr Asp Pro Ser Phe Arg Gly Pro Ile Lys Asn Arg Ser Cys
 20             25             30
Thr Asp Val Ile Cys Cys Val Leu Phe Leu Leu Phe Ile Leu Gly Tyr
 35             40             45
Ile Val Val Gly Ile Val Ala Trp Leu Tyr Gly Asp Pro Arg Gln Val
 50             55             60
Leu Tyr Pro Arg Asn Ser Thr Gly Ala Tyr Cys Gly Met Gly Glu Asn
 65             70             75             80
Lys Asp Lys Pro Tyr Leu Leu Tyr Phe Asn Ile Phe Ser Cys Ile Leu
 85             90             95
Ser Ser Asn Ile Ile Ser Val Ala Glu Asn Gly Leu Gln Cys Pro Thr
 100            105            110
Pro Gln Val Cys Val Ser Ser Cys Pro Glu Asp Pro Trp Thr Val Gly
 115            120            125
Lys Asn Glu Phe Ser Gln Thr Val Gly Glu Val Phe Tyr Thr Lys Asn
 130            135            140
Arg Asn Phe Cys Leu Pro Gly Val Pro Trp Asn Met Thr Val Ile Thr
 145            150            155            160
Ser Leu Gln Gln Glu Leu Cys Pro Ser Phe Leu Leu Pro Ser Ala Pro
 165            170            175
Ala Leu Gly Arg Cys Phe Pro Trp Thr Asn Val Thr Pro Pro Ala Leu
 180            185            190
Pro Gly Ile Thr Asn Asp Thr Thr Ile Gln Gln Gly Ile Ser Gly Leu
 195            200            205
Ile Asp Ser Leu Asn Ala Arg Asp Ile Ser Val Lys Ile Phe Glu Asp
 210            215            220
Phe Ala Gln Ser Trp Tyr Trp Ile Leu Val Ala Leu Gly Val Ala Leu
 225            230            235            240
Val Leu Ser Leu Leu Phe Ile Leu Leu Leu Arg Leu Val Ala Gly Pro
 245            250            255
Leu Val Leu Val Leu Ile Leu Gly Val Leu Gly Val Leu Ala Tyr Gly
 260            265            270
Ile Tyr Tyr Cys Trp Glu Glu Tyr Arg Val Leu Arg Asp Lys Gly Ala
 275            280            285
Ser Ile Ser Gln Leu Gly Phe Thr Thr Asn Leu Ser Ala Tyr Gln Ser
 290            295            300
Val Gln Glu Thr Trp Leu Ala Ala Leu Ile Val Leu Ala Val Leu Glu
 305            310            315            320
Ala Ile Leu Leu Leu Met Leu Ile Phe Leu Arg Gln Arg Ile Arg Ile
 325            330            335
Ala Ile Ala Leu Leu Lys Glu Ala Ser Lys Ala Val Gly Gln Met Met
 340            345            350

```

-continued

Ser Thr Met Phe Tyr Pro Leu Val Thr Phe Val Leu Leu Leu Ile Cys
 355 360 365
 Ile Ala Tyr Trp Ala Met Thr Ala Leu Tyr Pro Leu Pro Thr Gln Pro
 370 375 380
 Ala Thr Leu Gly Tyr Val Leu Trp Ala Ser Asn Ile Ser Ser Pro Gly
 385 390 395 400
 Cys Glu Lys Val Pro Ile Asn Thr Ser Cys Asn Pro Thr Ala His Leu
 405 410 415
 Val Asn Ser Ser Cys Pro Gly Leu Met Cys Val Phe Gln Gly Tyr Ser
 420 425 430
 Ser Lys Gly Leu Ile Gln Arg Ser Val Phe Asn Leu Gln Ile Tyr Gly
 435 440 445
 Val Leu Gly Leu Phe Trp Thr Leu Asn Trp Val Leu Ala Leu Gly Gln
 450 455 460
 Cys Val Leu Ala Gly Ala Phe Ala Ser Phe Tyr Trp Ala Phe His Lys
 465 470 475 480
 Pro Gln Asp Ile Pro Thr Phe Pro Leu Ile Ser Ala Phe Ile Arg Thr
 485 490 495
 Leu Arg Tyr His Thr Gly Ser Leu Ala Phe Gly Ala Leu Ile Leu Thr
 500 505 510
 Leu Val Gln Ile Ala Arg Val Ile Leu Glu Tyr Ile Asp His Lys Leu
 515 520 525
 Arg Gly Val Gln Asn Pro Val Ala Arg Cys Ile Met Cys Cys Phe Lys
 530 535 540
 Cys Cys Leu Trp Cys Leu Glu Lys Phe Ile Lys Phe Leu Asn Arg Asn
 545 550 555 560
 Ala Tyr Ile Met Ile Ala Ile Tyr Gly Lys Asn Phe Cys Val Ser Ala
 565 570 575
 Lys Asn Ala Phe Met Leu Leu Met Arg Asn Ile Val Arg Val Val Val
 580 585 590
 Leu Asp Lys Val Thr Asp Leu Leu Leu Phe Phe Gly Lys Leu Leu Val
 595 600 605
 Val Gly Gly Val Gly Val Leu Ser Phe Phe Phe Phe Ser Gly Arg Ile
 610 615 620
 Pro Gly Leu Gly Lys Asp Phe Lys Ser Pro His Leu Asn Tyr Tyr Trp
 625 630 635 640
 Leu Pro Ile Met Thr Ser Ile Leu Gly Ala Tyr Val Ile Ala Ser Gly
 645 650 655
 Phe Phe Ser Val Phe Gly Met Cys Val Asp Thr Leu Phe Leu Cys Phe
 660 665 670
 Leu Glu Asp Leu Glu Arg Asn Asn Gly Ser Leu Asp Arg Pro Tyr Tyr
 675 680 685
 Met Ser Lys Ser Leu Leu Lys Ile Leu Gly Lys Lys Asn Glu Ala Pro
 690 695 700
 Pro Asp Asn Lys Lys Arg Lys Lys
 705 710

<210> SEQ ID NO 20

<211> LENGTH: 710

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 20

Met Gly Gly Lys Gln Arg Asp Glu Asp Asp Glu Ala Tyr Gly Lys Pro
 1 5 10 15

-continued

Val	Lys	Tyr	Asp	Pro	Ser	Phe	Arg	Gly	Pro	Ile	Lys	Asn	Arg	Ser	Cys	
	20							25				30				
Thr	Asp	Val	Ile	Cys	Cys	Val	Leu	Phe	Leu	Leu	Phe	Ile	Leu	Gly	Tyr	
	35						40					45				
Ile	Val	Val	Gly	Ile	Val	Ala	Trp	Leu	Tyr	Gly	Asp	Pro	Arg	Gln	Val	
	50					55					60					
Leu	Tyr	Pro	Arg	Asn	Ser	Thr	Gly	Ala	Tyr	Cys	Gly	Met	Gly	Glu	Asn	
	65				70					75					80	
Lys	Asp	Lys	Pro	Tyr	Leu	Leu	Tyr	Phe	Asn	Ile	Phe	Ser	Cys	Ile	Leu	
				85					90					95		
Ser	Ser	Asn	Ile	Ile	Ser	Val	Ala	Glu	Asn	Gly	Leu	Gln	Cys	Pro	Thr	
			100						105				110			
Pro	Gln	Val	Cys	Val	Ser	Ser	Cys	Pro	Glu	Asp	Pro	Trp	Thr	Val	Gly	
		115					120					125				
Lys	Asn	Glu	Phe	Ser	Gln	Thr	Val	Gly	Glu	Val	Phe	Tyr	Thr	Lys	Asn	
	130					135					140					
Arg	Asn	Phe	Cys	Leu	Pro	Gly	Val	Pro	Trp	Asn	Met	Thr	Val	Ile	Thr	
	145				150					155					160	
Ser	Leu	Gln	Gln	Glu	Leu	Cys	Pro	Ser	Phe	Leu	Leu	Pro	Ser	Ala	Pro	
				165					170					175		
Ala	Leu	Gly	Arg	Cys	Phe	Pro	Trp	Thr	Asn	Val	Thr	Pro	Pro	Ala	Leu	
			180					185					190			
Pro	Gly	Ile	Thr	Asn	Asp	Thr	Thr	Ile	Gln	Gln	Gly	Ile	Ser	Gly	Leu	
		195				200						205				
Ile	Asp	Ser	Leu	Asn	Ala	Arg	Asp	Ile	Ser	Val	Lys	Ile	Phe	Glu	Asp	
	210				215						220					
Phe	Ala	Gln	Ser	Trp	Tyr	Trp	Ile	Leu	Val	Ala	Leu	Gly	Val	Ala	Leu	
	225				230					235					240	
Val	Leu	Ser	Leu	Leu	Phe	Ile	Leu	Leu	Leu	Arg	Leu	Val	Ala	Gly	Pro	
			245						250					255		
Leu	Val	Leu	Val	Leu	Ile	Leu	Gly	Val	Leu	Gly	Val	Leu	Ala	Tyr	Gly	
			260					265					270			
Ile	Tyr	Tyr	Cys	Trp	Glu	Glu	Tyr	Arg	Val	Leu	Arg	Asp	Lys	Gly	Ala	
		275					280					285				
Ser	Ile	Ser	Gln	Leu	Gly	Phe	Thr	Thr	Asn	Leu	Ser	Ala	Tyr	Gln	Ser	
	290					295					300					
Val	Gln	Glu	Thr	Trp	Leu	Ala	Ala	Leu	Ile	Val	Leu	Ala	Val	Leu	Glu	
	305				310					315					320	
Ala	Ile	Leu	Leu	Leu	Met	Leu	Ile	Phe	Leu	Arg	Gln	Arg	Ile	Arg	Ile	
			325						330				335			
Ala	Ile	Ala	Leu	Leu	Lys	Glu	Ala	Ser	Lys	Ala	Val	Gly	Gln	Met	Met	
		340						345					350			
Ser	Thr	Met	Phe	Tyr	Pro	Leu	Val	Thr	Phe	Val	Leu	Leu	Leu	Ile	Cys	
		355					360					365				
Ile	Ala	Tyr	Trp	Ala	Met	Thr	Ala	Leu	Tyr	Leu	Ala	Thr	Ser	Gly	Gln	
	370					375					380					
Pro	Gln	Tyr	Val	Leu	Trp	Ala	Ser	Asn	Ile	Ser	Ser	Pro	Gly	Cys	Glu	
	385				390					395					400	
Lys	Val	Pro	Ile	Asn	Thr	Ser	Cys	Asn	Pro	Thr	Ala	His	Leu	Val	Asn	
			405						410					415		
Ser	Ser	Cys	Pro	Gly	Leu	Met	Cys	Val	Phe	Gln	Gly	Tyr	Ser	Ser	Lys	
		420						425					430			
Gly	Leu	Ile	Gln	Arg	Ser	Val	Phe	Asn	Leu	Gln	Ile	Tyr	Gly	Val	Leu	

-continued

435	440	445
Gly Leu Phe Trp Thr Leu Asn Trp Val Leu Ala Leu Gly Gln Cys Val		
450	455	460
Leu Ala Gly Ala Phe Ala Ser Phe Tyr Trp Ala Phe His Lys Pro Gln		
465	470	475
Asp Ile Pro Thr Phe Pro Leu Ile Ser Ala Phe Ile Arg Thr Leu Arg		
	485	490
Tyr His Thr Gly Ser Leu Ala Phe Gly Ala Leu Ile Leu Thr Leu Val		
	500	505
Gln Ile Ala Arg Val Ile Leu Glu Tyr Ile Asp His Lys Leu Arg Gly		
	515	520
Val Gln Asn Pro Val Ala Arg Cys Ile Met Cys Cys Phe Lys Cys Cys		
	530	535
Leu Trp Cys Leu Glu Lys Phe Ile Lys Phe Leu Asn Arg Asn Ala Tyr		
545	550	555
Ile Met Ile Ala Ile Tyr Gly Lys Asn Phe Cys Val Ser Ala Lys Asn		
	565	570
Ala Phe Met Leu Leu Met Arg Asn Ile Val Arg Val Val Val Leu Asp		
	580	585
Lys Val Thr Asp Leu Leu Leu Phe Phe Gly Lys Leu Leu Val Val Gly		
	595	600
Gly Val Gly Val Leu Ser Phe Phe Phe Phe Ser Gly Arg Ile Pro Gly		
	610	615
Leu Gly Lys Asp Phe Lys Ser Pro His Leu Asn Tyr Tyr Trp Leu Pro		
625	630	635
Ile Met Thr Ser Ile Leu Gly Ala Tyr Val Ile Ala Ser Gly Phe Phe		
	645	650
Ser Val Phe Gly Met Cys Val Asp Thr Leu Phe Leu Cys Phe Leu Glu		
	660	665
Asp Leu Glu Arg Asn Asn Gly Ser Leu Asp Arg Pro Tyr Tyr Met Ser		
	675	680
Lys Ser Leu Leu Lys Ile Leu Gly Lys Lys Asn Glu Ala Pro Pro Asp		
	690	695
Asn Lys Lys Arg Lys Lys		
705	710	

<210> SEQ ID NO 21

<211> LENGTH: 710

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 21

Met Gly Gly Lys Gln Arg Asp Glu Asp Asp Glu Ala Tyr Gly Lys Pro		
1	5	10
Val Lys Tyr Asp Pro Ser Phe Arg Gly Pro Ile Lys Asn Arg Ser Cys		
	20	25
Thr Asp Val Ile Cys Cys Val Leu Phe Leu Leu Phe Ile Leu Gly Tyr		
	35	40
Ile Val Val Gly Ile Val Ala Trp Leu Tyr Gly Asp Pro Arg Gln Val		
	50	55
Leu Tyr Pro Arg Asn Ser Thr Gly Ala Tyr Cys Gly Met Gly Glu Asn		
65	70	75
Lys Asp Lys Pro Tyr Leu Leu Tyr Phe Asn Ile Phe Ser Cys Ile Leu		
	85	90
Ser Ser Asn Ile Ile Ser Val Ala Glu Asn Gly Leu Gln Cys Pro Thr		

-continued

100								105				110					
Pro	Gln	Val	Cys	Val	Ser	Ser	Cys	Pro	Glu	Asp	Pro	Trp	Thr	Val	Gly		
115								120				125					
Lys	Asn	Glu	Phe	Ser	Gln	Thr	Val	Gly	Glu	Val	Phe	Tyr	Thr	Lys	Asn		
130								135				140					
Arg	Asn	Phe	Cys	Leu	Pro	Gly	Val	Pro	Trp	Asn	Met	Thr	Val	Ile	Thr		
145								150				155					
Ser	Leu	Gln	Gln	Glu	Leu	Cys	Pro	Ser	Phe	Leu	Leu	Pro	Ser	Ala	Pro		
165								170				175					
Ala	Leu	Gly	Arg	Cys	Phe	Pro	Trp	Thr	Asn	Ile	Thr	Pro	Pro	Ala	Leu		
180								185				190					
Pro	Gly	Ile	Thr	Asn	Asp	Thr	Thr	Ile	Gln	Gln	Gly	Ile	Ser	Gly	Leu		
195								200				205					
Ile	Asp	Ser	Leu	Asn	Ala	Arg	Asp	Ile	Ser	Val	Lys	Ile	Phe	Glu	Asp		
210								215				220					
Phe	Ala	Gln	Ser	Trp	Tyr	Trp	Ile	Leu	Val	Ala	Leu	Gly	Val	Ala	Leu		
225								230				235					
Val	Leu	Ser	Leu	Leu	Phe	Ile	Leu	Leu	Leu	Arg	Leu	Val	Ala	Gly	Pro		
245								250				255					
Leu	Val	Leu	Val	Leu	Ile	Leu	Gly	Val	Leu	Gly	Val	Leu	Ala	Tyr	Gly		
260								265				270					
Ile	Tyr	Tyr	Cys	Trp	Glu	Glu	Tyr	Arg	Val	Leu	Arg	Asp	Lys	Gly	Ala		
275								280				285					
Ser	Ile	Ser	Gln	Leu	Gly	Phe	Thr	Thr	Asn	Leu	Ser	Ala	Tyr	Gln	Ser		
290								295				300					
Val	Gln	Glu	Thr	Trp	Leu	Ala	Ala	Leu	Ile	Val	Leu	Ala	Val	Leu	Glu		
305								310				315					
Ala	Ile	Leu	Leu	Leu	Met	Leu	Ile	Phe	Leu	Arg	Gln	Arg	Ile	Arg	Ile		
325								330				335					
Ala	Ile	Ala	Leu	Leu	Lys	Glu	Ala	Ser	Lys	Ala	Val	Gly	Gln	Met	Met		
340								345				350					
Ser	Thr	Met	Phe	Tyr	Pro	Leu	Val	Thr	Phe	Val	Leu	Leu	Leu	Ile	Cys		
355								360				365					
Ile	Ala	Tyr	Trp	Ala	Met	Thr	Ala	Leu	Tyr	Leu	Ala	Thr	Ser	Gly	Gln		
370								375				380					
Pro	Gln	Tyr	Val	Leu	Trp	Ala	Ser	Asn	Ile	Ser	Ser	Pro	Gly	Cys	Glu		
385								390				395					
Lys	Val	Pro	Ile	Asn	Thr	Ser	Cys	Asn	Pro	Thr	Ala	His	Leu	Val	Asn		
405								410				415					
Ser	Ser	Cys	Pro	Gly	Leu	Met	Cys	Val	Phe	Gln	Gly	Tyr	Ser	Ser	Lys		
420								425				430					
Gly	Leu	Ile	Gln	Arg	Ser	Val	Phe	Asn	Leu	Gln	Ile	Tyr	Gly	Val	Leu		
435								440				445					
Gly	Leu	Phe	Trp	Thr	Leu	Asn	Trp	Val	Leu	Ala	Leu	Gly	Gln	Cys	Val		
450								455				460					
Leu	Ala	Gly	Ala	Phe	Ala	Ser	Phe	Tyr	Trp	Ala	Phe	His	Lys	Pro	Gln		
465								470				475					
Asp	Ile	Pro	Thr	Phe	Pro	Leu	Ile	Ser	Ala	Phe	Ile	Arg	Thr	Leu	Arg		
485								490				495					
Tyr	His	Thr	Gly	Ser	Leu	Ala	Phe	Gly	Ala	Leu	Ile	Leu	Thr	Leu	Val		
500								505				510					
Gln	Ile	Ala	Arg	Val	Ile	Leu	Glu	Tyr	Ile	Asp	His	Lys	Leu	Arg	Gly		
515								520				525					

Figure 3C-2

```
481 DIPTFPLISA FIRTLYHTG SLAFGALILT LVQIARVILE YIDHKLRGVQ NPVARCIMCC
541 FKCCCLWCLEK FIKFLNRNAY IMIAIYGKNE CVSAKNAFML LMRNIVRVVV LDKVTDLLLF
601 FGKLLVVGCV GVLSEFFFSG RIPGLGKDFK SPHLNYYWLP IMTSILGAYV IASGFFSVFG
661 MCVDTLFLCF LEDLERNNGS LDRPYMSKS LLKILGKKNE APPDNKKRKK
```

Figure 3D. Amino acid sequence of 24P4C12 v.6 (SEQ ID. NO. : 23). The 24P4C12 v.6 protein has 710 amino acids.

```
1 MGGKQRDEDD EAYGKPVKYD PSFRGPIKNR SCTDVICCVL FLLFILGYIV VGIVAWLYGD
61 PRQVLYPRNS TGAYCGMGEN KDKPYLLYFN IFSCILSSNI ISVAENGLQC PTPQVCVSSC
121 PEDPWTVGKN EFSQTVGEVF YTKNRNFCLP GVPWNMTVIT SLQQELCPSF LLPSAPALGR
181 CFPWTNVTTP ALPGITNDTT IQQGISGLID SLNARDISVK IFEDFAQSWY WILVALGVAL
241 VLSLLFILLL RLVAGPLVLV LILGVLGVLA YGIYYCWEEY RVLDRDKGASI SQLGFTTNLS
301 AYQSVQETWL AALIVLAVLE AILLMLIFL RQRIRIAIAL LKEASKAVGQ MMSTMEYPLV
361 TFVLLLICIA YWAMTALYLA TSGQPQYVLW ASNISSPGCE KVPINTSCNP TAHLVNSSCP
421 GLMCVFQGYG SKGLIPRSVF NLQIYGVGLG FWTLNWLAL GQCVLAGAFA SFYWAFHKPQ
481 DIPTFPLISA FIRTLYHTG SLAFGALILT LVQIARVILE YIDHKLRGVQ NPVARCIMCC
541 FKCCCLWCLEK FIKFLNRNAY IMIAIYGKNE CVSAKNAFML LMRNIVRVVV LDKVTDLLLF
601 FGKLLVVGCV GVLSEFFFSG RIPGLGKDFK SPHLNYYWLP IMTSILGAYV IASGFFSVFG
661 MCVDTLFLCF LEDLERNNGS LDRPYMSKS LLKILGKKNE APPDNKKRKK
```

Figure 3E. Amino acid sequence of 24P4C12 v.7 (SEQ ID. NO. : 24). The 24P4C12 v.7 protein has 598 amino acids.

```
1 MGGKQRDEDD EAYGKPVKYD PSFRGPIKNR SCTDVICCVL FLLFILGYIV VGIVAWLYGD
61 PRQVLYPRNS TGAYCGMGEN KDKPYLLYFN IFSCILSSNI ISVAENGLQC PTPQVCVSSC
121 PEDPWTVGKN EFSQTVGEVF YTKNRNFCLP GVPWNMTVIT SLQQELCPSF LLPSAPALGR
181 CFPWTNVTTP ALPGITNDTT IQQGISGLID SLNARDISVK IFEDFAQSWY WILVAVGQMM
241 STMFYPLVTF VLLLICIAIW AMTALYLATS GQPQYVLWAS NISSPGCEKV PINTSCNPTA
301 HLVNSSCPGL MCVFQGYSSK GLIQRSVFNL QIYGVGLFW TLNWLALGQ CVLAGAFASF
361 YWAFHKPQDI PTFPLISAFI RTRYHTGSL AFGALILTLV QIARVILEYI DHKLRGVQNP
421 VARCIMCCFK CCLWCLEKFI KFLNRNAYIM IAIYGKNFCV SAKNAFMLLM RNIVRVVLD
481 KVIDLLFFFG KLLVVGCVGV LSFFFSGRIP PGLGKDFKSP HLNYYWLPIM TSILGAYVIA
541 SGFFSVFGMC VDTLFLCFLE DLERNNGSLD RPYYMSKSLL KILGKKNEAP PDNKKRKK
```

-continued

Val	Gln	Asn	Pro	Val	Ala	Arg	Cys	Ile	Met	Cys	Cys	Phe	Lys	Cys	Cys
530						535						540			
Leu	Trp	Cys	Leu	Glu	Lys	Phe	Ile	Lys	Phe	Leu	Asn	Arg	Asn	Ala	Tyr
545					550					555					560
Ile	Met	Ile	Ala	Ile	Tyr	Gly	Lys	Asn	Phe	Cys	Val	Ser	Ala	Lys	Asn
				565					570					575	
Ala	Phe	Met	Leu	Leu	Met	Arg	Asn	Ile	Val	Arg	Val	Val	Val	Leu	Asp
			580					585					590		
Lys	Val	Thr	Asp	Leu	Leu	Leu	Phe	Phe	Gly	Lys	Leu	Leu	Val	Val	Gly
		595					600					605			
Gly	Val	Gly	Val	Leu	Ser	Phe	Phe	Phe	Phe	Ser	Gly	Arg	Ile	Pro	Gly
	610					615					620				
Leu	Gly	Lys	Asp	Phe	Lys	Ser	Pro	His	Leu	Asn	Tyr	Tyr	Trp	Leu	Pro
625					630					635					640
Ile	Met	Thr	Ser	Ile	Leu	Gly	Ala	Tyr	Val	Ile	Ala	Ser	Gly	Phe	Phe
				645					650					655	
Ser	Val	Phe	Gly	Met	Cys	Val	Asp	Thr	Leu	Phe	Leu	Cys	Phe	Leu	Glu
			660					665					670		
Asp	Leu	Glu	Arg	Asn	Asn	Gly	Ser	Leu	Asp	Arg	Pro	Tyr	Tyr	Met	Ser
		675					680					685			
Lys	Ser	Leu	Leu	Lys	Ile	Leu	Gly	Lys	Lys	Asn	Glu	Ala	Pro	Pro	Asp
		690				695					700				
Asn	Lys	Lys	Arg	Lys	Lys										
705					710										

<210> SEQ ID NO 22

<211> LENGTH: 710

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 22

Met	Gly	Gly	Lys	Gln	Arg	Asp	Glu	Asp	Asp	Glu	Ala	Tyr	Gly	Lys	Pro
1				5					10					15	
Val	Lys	Tyr	Asp	Pro	Ser	Phe	Arg	Gly	Pro	Ile	Lys	Asn	Arg	Ser	Cys
			20					25					30		
Thr	Asp	Val	Ile	Cys	Cys	Val	Leu	Phe	Leu	Leu	Phe	Ile	Leu	Gly	Tyr
		35					40					45			
Ile	Val	Val	Gly	Ile	Val	Ala	Trp	Leu	Tyr	Gly	Asp	Pro	Arg	Gln	Val
	50					55					60				
Leu	Tyr	Pro	Arg	Asn	Ser	Thr	Gly	Ala	Tyr	Cys	Gly	Met	Gly	Glu	Asn
65				70					75					80	
Lys	Asp	Lys	Pro	Tyr	Leu	Leu	Tyr	Phe	Asn	Ile	Phe	Ser	Cys	Ile	Leu
			85					90						95	
Ser	Ser	Asn	Ile	Ile	Ser	Val	Ala	Glu	Asn	Gly	Leu	Gln	Cys	Pro	Thr
		100						105					110		
Pro	Gln	Val	Cys	Val	Ser	Ser	Cys	Pro	Glu	Asp	Pro	Trp	Thr	Val	Gly
		115					120					125			
Lys	Asn	Glu	Phe	Ser	Gln	Thr	Val	Gly	Glu	Val	Phe	Tyr	Thr	Lys	Asn
	130				135						140				
Arg	Asn	Phe	Cys	Leu	Pro	Gly	Val	Pro	Trp	Asn	Met	Thr	Val	Ile	Thr
145				150						155					160
Ser	Leu	Gln	Gln	Glu	Leu	Cys	Pro	Ser	Phe	Leu	Leu	Pro	Ser	Ala	Pro
			165						170					175	
Ala	Leu	Gly	Arg	Cys	Phe	Pro	Trp	Thr	Asn	Val	Thr	Pro	Pro	Ala	Leu
			180					185						190	

-continued

Pro Gly Ile Thr Asn Asp Thr Thr Ile Gln Gln Gly Ile Ser Gly Leu	195	200	205
Ile Asp Ser Leu Asn Ala Arg Asp Ile Ser Val Lys Ile Phe Glu Asp	210	215	220
Phe Ala Gln Ser Trp Tyr Trp Ile Leu Val Ala Leu Gly Val Ala Leu	225	230	235
Val Leu Ser Leu Leu Phe Ile Leu Leu Leu Arg Leu Val Ala Gly Pro	245	250	255
Leu Val Leu Val Leu Ile Leu Gly Val Leu Gly Val Leu Ala Tyr Gly	260	265	270
Ile Tyr Tyr Cys Trp Glu Glu Tyr Arg Val Leu Arg Asp Lys Gly Ala	275	280	285
Ser Ile Ser Gln Leu Gly Phe Thr Thr Asn Leu Ser Ala Tyr Gln Ser	290	295	300
Val Gln Glu Thr Trp Leu Ala Ala Leu Ile Val Leu Ala Val Leu Glu	305	310	315
Ala Ile Leu Leu Leu Val Leu Ile Phe Leu Arg Gln Arg Ile Arg Ile	325	330	335
Ala Ile Ala Leu Leu Lys Glu Ala Ser Lys Ala Val Gly Gln Met Met	340	345	350
Ser Thr Met Phe Tyr Pro Leu Val Thr Phe Val Leu Leu Leu Ile Cys	355	360	365
Ile Ala Tyr Trp Ala Met Thr Ala Leu Tyr Leu Ala Thr Ser Gly Gln	370	375	380
Pro Gln Tyr Val Leu Trp Ala Ser Asn Ile Ser Ser Pro Gly Cys Glu	385	390	395
Lys Val Pro Ile Asn Thr Ser Cys Asn Pro Thr Ala His Leu Val Asn	405	410	415
Ser Ser Cys Pro Gly Leu Met Cys Val Phe Gln Gly Tyr Ser Ser Lys	420	425	430
Gly Leu Ile Gln Arg Ser Val Phe Asn Leu Gln Ile Tyr Gly Val Leu	435	440	445
Gly Leu Phe Trp Thr Leu Asn Trp Val Leu Ala Leu Gly Gln Cys Val	450	455	460
Leu Ala Gly Ala Phe Ala Ser Phe Tyr Trp Ala Phe His Lys Pro Gln	465	470	475
Asp Ile Pro Thr Phe Pro Leu Ile Ser Ala Phe Ile Arg Thr Leu Arg	485	490	495
Tyr His Thr Gly Ser Leu Ala Phe Gly Ala Leu Ile Leu Thr Leu Val	500	505	510
Gln Ile Ala Arg Val Ile Leu Glu Tyr Ile Asp His Lys Leu Arg Gly	515	520	525
Val Gln Asn Pro Val Ala Arg Cys Ile Met Cys Cys Phe Lys Cys Cys	530	535	540
Leu Trp Cys Leu Glu Lys Phe Ile Lys Phe Leu Asn Arg Asn Ala Tyr	545	550	555
Ile Met Ile Ala Ile Tyr Gly Lys Asn Phe Cys Val Ser Ala Lys Asn	565	570	575
Ala Phe Met Leu Leu Met Arg Asn Ile Val Arg Val Val Val Leu Asp	580	585	590
Lys Val Thr Asp Leu Leu Leu Phe Phe Gly Lys Leu Leu Val Val Gly	595	600	605
Gly Val Gly Val Leu Ser Phe Phe Phe Phe Ser Gly Arg Ile Pro Gly	610	615	620

-continued

Leu Gly Lys Asp Phe Lys Ser Pro His Leu Asn Tyr Tyr Trp Leu Pro
 625 630 635 640
 Ile Met Thr Ser Ile Leu Gly Ala Tyr Val Ile Ala Ser Gly Phe Phe
 645 650 655
 Ser Val Phe Gly Met Cys Val Asp Thr Leu Phe Leu Cys Phe Leu Glu
 660 665 670
 Asp Leu Glu Arg Asn Asn Gly Ser Leu Asp Arg Pro Tyr Tyr Met Ser
 675 680 685
 Lys Ser Leu Leu Lys Ile Leu Gly Lys Lys Asn Glu Ala Pro Pro Asp
 690 695 700
 Asn Lys Lys Arg Lys Lys
 705 710

<210> SEQ ID NO 23
 <211> LENGTH: 710
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 23

Met Gly Gly Lys Gln Arg Asp Glu Asp Asp Glu Ala Tyr Gly Lys Pro
 1 5 10 15
 Val Lys Tyr Asp Pro Ser Phe Arg Gly Pro Ile Lys Asn Arg Ser Cys
 20 25 30
 Thr Asp Val Ile Cys Cys Val Leu Phe Leu Leu Phe Ile Leu Gly Tyr
 35 40 45
 Ile Val Val Gly Ile Val Ala Trp Leu Tyr Gly Asp Pro Arg Gln Val
 50 55 60
 Leu Tyr Pro Arg Asn Ser Thr Gly Ala Tyr Cys Gly Met Gly Glu Asn
 65 70 75 80
 Lys Asp Lys Pro Tyr Leu Leu Tyr Phe Asn Ile Phe Ser Cys Ile Leu
 85 90 95
 Ser Ser Asn Ile Ile Ser Val Ala Glu Asn Gly Leu Gln Cys Pro Thr
 100 105 110
 Pro Gln Val Cys Val Ser Ser Cys Pro Glu Asp Pro Trp Thr Val Gly
 115 120 125
 Lys Asn Glu Phe Ser Gln Thr Val Gly Glu Val Phe Tyr Thr Lys Asn
 130 135 140
 Arg Asn Phe Cys Leu Pro Gly Val Pro Trp Asn Met Thr Val Ile Thr
 145 150 155 160
 Ser Leu Gln Gln Glu Leu Cys Pro Ser Phe Leu Leu Pro Ser Ala Pro
 165 170 175
 Ala Leu Gly Arg Cys Phe Pro Trp Thr Asn Val Thr Pro Pro Ala Leu
 180 185 190
 Pro Gly Ile Thr Asn Asp Thr Thr Ile Gln Gln Gly Ile Ser Gly Leu
 195 200 205
 Ile Asp Ser Leu Asn Ala Arg Asp Ile Ser Val Lys Ile Phe Glu Asp
 210 215 220
 Phe Ala Gln Ser Trp Tyr Trp Ile Leu Val Ala Leu Gly Val Ala Leu
 225 230 235 240
 Val Leu Ser Leu Leu Phe Ile Leu Leu Leu Arg Leu Val Ala Gly Pro
 245 250 255
 Leu Val Leu Val Leu Ile Leu Gly Val Leu Gly Val Leu Ala Tyr Gly
 260 265 270
 Ile Tyr Tyr Cys Trp Glu Glu Tyr Arg Val Leu Arg Asp Lys Gly Ala
 275 280 285

-continued

Ser Ile Ser Gln Leu Gly Phe Thr Thr Asn Leu Ser Ala Tyr Gln Ser
 290 295 300
 Val Gln Glu Thr Trp Leu Ala Ala Leu Ile Val Leu Ala Val Leu Glu
 305 310 315 320
 Ala Ile Leu Leu Leu Met Leu Ile Phe Leu Arg Gln Arg Ile Arg Ile
 325 330 335
 Ala Ile Ala Leu Leu Lys Glu Ala Ser Lys Ala Val Gly Gln Met Met
 340 345 350
 Ser Thr Met Phe Tyr Pro Leu Val Thr Phe Val Leu Leu Leu Ile Cys
 355 360 365
 Ile Ala Tyr Trp Ala Met Thr Ala Leu Tyr Leu Ala Thr Ser Gly Gln
 370 375 380
 Pro Gln Tyr Val Leu Trp Ala Ser Asn Ile Ser Ser Pro Gly Cys Glu
 385 390 395 400
 Lys Val Pro Ile Asn Thr Ser Cys Asn Pro Thr Ala His Leu Val Asn
 405 410 415
 Ser Ser Cys Pro Gly Leu Met Cys Val Phe Gln Gly Tyr Ser Ser Lys
 420 425 430
 Gly Leu Ile Pro Arg Ser Val Phe Asn Leu Gln Ile Tyr Gly Val Leu
 435 440 445
 Gly Leu Phe Trp Thr Leu Asn Trp Val Leu Ala Leu Gly Gln Cys Val
 450 455 460
 Leu Ala Gly Ala Phe Ala Ser Phe Tyr Trp Ala Phe His Lys Pro Gln
 465 470 475 480
 Asp Ile Pro Thr Phe Pro Leu Ile Ser Ala Phe Ile Arg Thr Leu Arg
 485 490 495
 Tyr His Thr Gly Ser Leu Ala Phe Gly Ala Leu Ile Leu Thr Leu Val
 500 505 510
 Gln Ile Ala Arg Val Ile Leu Glu Tyr Ile Asp His Lys Leu Arg Gly
 515 520 525
 Val Gln Asn Pro Val Ala Arg Cys Ile Met Cys Cys Phe Lys Cys Cys
 530 535 540
 Leu Trp Cys Leu Glu Lys Phe Ile Lys Phe Leu Asn Arg Asn Ala Tyr
 545 550 555 560
 Ile Met Ile Ala Ile Tyr Gly Lys Asn Phe Cys Val Ser Ala Lys Asn
 565 570 575
 Ala Phe Met Leu Leu Met Arg Asn Ile Val Arg Val Val Val Leu Asp
 580 585 590
 Lys Val Thr Asp Leu Leu Leu Phe Phe Gly Lys Leu Leu Val Val Gly
 595 600 605
 Gly Val Gly Val Leu Ser Phe Phe Phe Phe Ser Gly Arg Ile Pro Gly
 610 615 620
 Leu Gly Lys Asp Phe Lys Ser Pro His Leu Asn Tyr Tyr Trp Leu Pro
 625 630 635 640
 Ile Met Thr Ser Ile Leu Gly Ala Tyr Val Ile Ala Ser Gly Phe Phe
 645 650 655
 Ser Val Phe Gly Met Cys Val Asp Thr Leu Phe Leu Cys Phe Leu Glu
 660 665 670
 Asp Leu Glu Arg Asn Asn Gly Ser Leu Asp Arg Pro Tyr Tyr Met Ser
 675 680 685
 Lys Ser Leu Leu Lys Ile Leu Gly Lys Lys Asn Glu Ala Pro Pro Asp
 690 695 700
 Asn Lys Lys Arg Lys Lys

-continued

705

710

<210> SEQ ID NO 24
 <211> LENGTH: 598
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 24

Met Gly Gly Lys Gln Arg Asp Glu Asp Asp Glu Ala Tyr Gly Lys Pro
 1 5 10 15
 Val Lys Tyr Asp Pro Ser Phe Arg Gly Pro Ile Lys Asn Arg Ser Cys
 20 25 30
 Thr Asp Val Ile Cys Cys Val Leu Phe Leu Leu Phe Ile Leu Gly Tyr
 35 40 45
 Ile Val Val Gly Ile Val Ala Trp Leu Tyr Gly Asp Pro Arg Gln Val
 50 55 60
 Leu Tyr Pro Arg Asn Ser Thr Gly Ala Tyr Cys Gly Met Gly Glu Asn
 65 70 75 80
 Lys Asp Lys Pro Tyr Leu Leu Tyr Phe Asn Ile Phe Ser Cys Ile Leu
 85 90 95
 Ser Ser Asn Ile Ile Ser Val Ala Glu Asn Gly Leu Gln Cys Pro Thr
 100 105 110
 Pro Gln Val Cys Val Ser Ser Cys Pro Glu Asp Pro Trp Thr Val Gly
 115 120 125
 Lys Asn Glu Phe Ser Gln Thr Val Gly Glu Val Phe Tyr Thr Lys Asn
 130 135 140
 Arg Asn Phe Cys Leu Pro Gly Val Pro Trp Asn Met Thr Val Ile Thr
 145 150 155 160
 Ser Leu Gln Gln Glu Leu Cys Pro Ser Phe Leu Leu Pro Ser Ala Pro
 165 170 175
 Ala Leu Gly Arg Cys Phe Pro Trp Thr Asn Val Thr Pro Pro Ala Leu
 180 185 190
 Pro Gly Ile Thr Asn Asp Thr Thr Ile Gln Gln Gly Ile Ser Gly Leu
 195 200 205
 Ile Asp Ser Leu Asn Ala Arg Asp Ile Ser Val Lys Ile Phe Glu Asp
 210 215 220
 Phe Ala Gln Ser Trp Tyr Trp Ile Leu Val Ala Val Gly Gln Met Met
 225 230 235 240
 Ser Thr Met Phe Tyr Pro Leu Val Thr Phe Val Leu Leu Leu Ile Cys
 245 250 255
 Ile Ala Tyr Trp Ala Met Thr Ala Leu Tyr Leu Ala Thr Ser Gly Gln
 260 265 270
 Pro Gln Tyr Val Leu Trp Ala Ser Asn Ile Ser Ser Pro Gly Cys Glu
 275 280 285
 Lys Val Pro Ile Asn Thr Ser Cys Asn Pro Thr Ala His Leu Val Asn
 290 295 300
 Ser Ser Cys Pro Gly Leu Met Cys Val Phe Gln Gly Tyr Ser Ser Lys
 305 310 315 320
 Gly Leu Ile Gln Arg Ser Val Phe Asn Leu Gln Ile Tyr Gly Val Leu
 325 330 335
 Gly Leu Phe Trp Thr Leu Asn Trp Val Leu Ala Leu Gly Gln Cys Val
 340 345 350
 Leu Ala Gly Ala Phe Ala Ser Phe Tyr Trp Ala Phe His Lys Pro Gln
 355 360 365
 Asp Ile Pro Thr Phe Pro Leu Ile Ser Ala Phe Ile Arg Thr Leu Arg

-continued

370	375	380
Tyr His Thr Gly Ser Leu Ala Phe Gly Ala Leu Ile Leu Thr Leu Val 385 390 395 400		
Gln Ile Ala Arg Val Ile Leu Glu Tyr Ile Asp His Lys Leu Arg Gly 405 410 415		
Val Gln Asn Pro Val Ala Arg Cys Ile Met Cys Cys Phe Lys Cys Cys 420 425 430		
Leu Trp Cys Leu Glu Lys Phe Ile Lys Phe Leu Asn Arg Asn Ala Tyr 435 440 445		
Ile Met Ile Ala Ile Tyr Gly Lys Asn Phe Cys Val Ser Ala Lys Asn 450 455 460		
Ala Phe Met Leu Leu Met Arg Asn Ile Val Arg Val Val Val Leu Asp 465 470 475 480		
Lys Val Thr Asp Leu Leu Leu Phe Phe Gly Lys Leu Leu Val Val Gly 485 490 495		
Gly Val Gly Val Leu Ser Phe Phe Phe Phe Ser Gly Arg Ile Pro Gly 500 505 510		
Leu Gly Lys Asp Phe Lys Ser Pro His Leu Asn Tyr Tyr Trp Leu Pro 515 520 525		
Ile Met Thr Ser Ile Leu Gly Ala Tyr Val Ile Ala Ser Gly Phe Phe 530 535 540		
Ser Val Phe Gly Met Cys Val Asp Thr Leu Phe Leu Cys Phe Leu Glu 545 550 555 560		
Asp Leu Glu Arg Asn Asn Gly Ser Leu Asp Arg Pro Tyr Tyr Met Ser 565 570 575		
Lys Ser Leu Leu Lys Ile Leu Gly Lys Lys Asn Glu Ala Pro Pro Asp 580 585 590		
Asn Lys Lys Arg Lys Lys 595		

<210> SEQ ID NO 25

<211> LENGTH: 722

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 25

Met Gly Gly Lys Gln Arg Asp Glu Asp Asp Glu Ala Tyr Gly Lys Pro 1 5 10 15
Val Lys Tyr Asp Pro Ser Phe Arg Gly Pro Ile Lys Asn Arg Ser Cys 20 25 30
Thr Asp Val Ile Cys Cys Val Leu Phe Leu Leu Phe Ile Leu Gly Tyr 35 40 45
Ile Val Val Gly Ile Val Ala Trp Leu Tyr Gly Asp Pro Arg Gln Val 50 55 60
Leu Tyr Pro Arg Asn Ser Thr Gly Ala Tyr Cys Gly Met Gly Glu Asn 65 70 75 80
Lys Asp Lys Pro Tyr Leu Leu Tyr Phe Asn Ile Phe Ser Cys Ile Leu 85 90 95
Ser Ser Asn Ile Ile Ser Val Ala Glu Asn Gly Leu Gln Cys Pro Thr 100 105 110
Pro Gln Val Cys Val Ser Ser Cys Pro Glu Asp Pro Trp Thr Val Gly 115 120 125
Lys Asn Glu Phe Ser Gln Thr Val Gly Glu Val Phe Tyr Thr Lys Asn 130 135 140
Arg Asn Phe Cys Leu Pro Gly Val Pro Trp Asn Met Thr Val Ile Thr

-continued

145	150	155	160
Ser Leu Gln Gln Glu Leu Cys Pro Ser Phe Leu Leu Pro Ser Ala Pro	165	170	175
Ala Leu Gly Arg Cys Phe Pro Trp Thr Asn Val Thr Pro Pro Ala Leu	180	185	190
Pro Gly Ile Thr Asn Asp Thr Thr Ile Gln Gln Gly Ile Ser Gly Leu	195	200	205
Ile Asp Ser Leu Asn Ala Arg Asp Ile Ser Val Lys Ile Phe Glu Asp	210	215	220
Phe Ala Gln Ser Trp Tyr Trp Ile Leu Val Ala Leu Gly Val Ala Leu	225	230	235
Val Leu Ser Leu Leu Phe Ile Leu Leu Leu Arg Leu Val Ala Gly Pro	245	250	255
Leu Val Leu Val Leu Ile Leu Gly Val Leu Gly Val Leu Ala Tyr Gly	260	265	270
Ile Tyr Tyr Cys Trp Glu Glu Tyr Arg Val Leu Arg Asp Lys Gly Ala	275	280	285
Ser Ile Ser Gln Leu Gly Phe Thr Thr Asn Leu Ser Ala Tyr Gln Ser	290	295	300
Val Gln Glu Thr Trp Leu Ala Ala Leu Ile Val Leu Ala Val Leu Glu	305	310	315
Ala Ile Leu Leu Leu Met Leu Ile Phe Leu Arg Gln Arg Ile Arg Ile	325	330	335
Ala Ile Ala Leu Leu Lys Glu Ala Ser Lys Ala Val Gly Gln Met Met	340	345	350
Ser Thr Met Phe Tyr Pro Leu Val Thr Phe Val Leu Leu Leu Ile Cys	355	360	365
Ile Ala Tyr Trp Ala Met Thr Ala Leu Tyr Leu Ala Thr Ser Gly Gln	370	375	380
Pro Gln Tyr Val Leu Trp Ala Ser Asn Ile Ser Ser Pro Gly Cys Glu	385	390	395
Lys Val Pro Ile Asn Thr Ser Cys Asn Pro Thr Ala His Leu Val Asn	405	410	415
Ser Ser Cys Pro Gly Leu Met Cys Val Phe Gln Gly Tyr Ser Ser Lys	420	425	430
Gly Leu Ile Gln Arg Ser Val Phe Asn Leu Gln Ile Tyr Gly Val Leu	435	440	445
Gly Leu Phe Trp Thr Leu Asn Trp Val Leu Ala Leu Gly Gln Cys Val	450	455	460
Leu Ala Gly Ala Phe Ala Ser Phe Tyr Trp Ala Phe His Lys Pro Gln	465	470	475
Asp Ile Pro Thr Phe Pro Leu Ile Ser Ala Phe Ile Arg Thr Leu Arg	485	490	495
Tyr His Thr Gly Ser Leu Ala Phe Gly Ala Leu Ile Leu Thr Leu Val	500	505	510
Gln Ile Ala Arg Val Ile Leu Glu Tyr Ile Asp His Lys Leu Arg Gly	515	520	525
Val Gln Asn Pro Val Ala Arg Cys Ile Met Cys Cys Phe Lys Cys Cys	530	535	540
Leu Trp Cys Leu Glu Lys Phe Ile Lys Phe Leu Asn Arg Asn Ala Tyr	545	550	555
Ile Met Ile Ala Ile Tyr Gly Lys Asn Phe Cys Val Ser Ala Lys Asn	565	570	575

-continued

Ala Phe Met Leu Leu Met Arg Asn Ile Val Arg Val Val Val Leu Asp
580 585 590

Lys Val Thr Asp Leu Leu Leu Phe Phe Gly Lys Leu Leu Val Val Gly
595 600 605

Gly Val Gly Val Leu Ser Phe Phe Phe Phe Ser Gly Arg Ile Pro Gly
610 615 620

Leu Gly Lys Asp Phe Lys Ser Pro His Leu Asn Tyr Tyr Trp Leu Pro
625 630 635 640

Ile Met Arg Asn Pro Ile Thr Pro Thr Gly His Val Phe Gln Thr Ser
645 650 655

Ile Leu Gly Ala Tyr Val Ile Ala Ser Gly Phe Phe Ser Val Phe Gly
660 665 670

Met Cys Val Asp Thr Leu Phe Leu Cys Phe Leu Glu Asp Leu Glu Arg
675 680 685

Asn Asn Gly Ser Leu Asp Arg Pro Tyr Tyr Met Ser Lys Ser Leu Leu
690 695 700

Lys Ile Leu Gly Lys Lys Asn Glu Ala Pro Pro Asp Asn Lys Lys Arg
705 710 715 720

Lys Lys

<210> SEQ ID NO 26
<211> LENGTH: 712
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 26

Met Gly Gly Lys Gln Arg Asp Glu Asp Asp Glu Ala Tyr Gly Lys Pro
1 5 10 15

Val Lys Tyr Asp Pro Ser Phe Arg Gly Pro Ile Lys Asn Arg Ser Cys
20 25 30

Thr Asp Val Ile Cys Cys Val Leu Phe Leu Leu Phe Ile Leu Gly Tyr
35 40 45

Ile Val Val Gly Ile Val Ala Trp Leu Tyr Gly Asp Pro Arg Gln Val
50 55 60

Leu Tyr Pro Arg Asn Ser Thr Gly Ala Tyr Cys Gly Met Gly Glu Asn
65 70 75 80

Lys Asp Lys Pro Tyr Leu Leu Tyr Phe Asn Ile Phe Ser Cys Ile Leu
85 90 95

Ser Ser Asn Ile Ile Ser Val Ala Glu Asn Gly Leu Gln Cys Pro Thr
100 105 110

Pro Gln Val Cys Val Ser Ser Cys Pro Glu Asp Pro Trp Thr Val Gly
115 120 125

Lys Asn Glu Phe Ser Gln Thr Val Gly Glu Val Phe Tyr Thr Lys Asn
130 135 140

Arg Asn Phe Cys Leu Pro Gly Val Pro Trp Asn Met Thr Val Ile Thr
145 150 155 160

Ser Leu Gln Gln Glu Leu Cys Pro Ser Phe Leu Leu Pro Ser Ala Pro
165 170 175

Ala Leu Gly Arg Cys Phe Pro Trp Thr Asn Val Thr Pro Pro Ala Leu
180 185 190

Pro Gly Ile Thr Asn Asp Thr Thr Ile Gln Gln Gly Ile Ser Gly Leu
195 200 205

Ile Asp Ser Leu Asn Ala Arg Asp Ile Ser Val Lys Ile Phe Glu Asp
210 215 220

Phe Ala Gln Ser Trp Tyr Trp Ile Leu Val Ala Leu Gly Val Ala Leu

-continued

225	230	235	240
Val Leu Ser Leu Leu Phe Ile Leu Leu Leu Arg Leu Val Ala Gly Pro	245	250	255
Leu Val Leu Val Leu Ile Leu Gly Val Leu Gly Val Leu Ala Tyr Gly	260	265	270
Ile Tyr Tyr Cys Trp Glu Glu Tyr Arg Val Leu Arg Asp Lys Gly Ala	275	280	285
Ser Ile Ser Gln Leu Gly Phe Thr Thr Asn Leu Ser Ala Tyr Gln Ser	290	295	300
Val Gln Glu Thr Trp Leu Ala Ala Leu Ile Val Leu Ala Val Leu Glu	305	310	315
Ala Ile Leu Leu Leu Met Leu Ile Phe Leu Arg Gln Arg Ile Arg Ile	325	330	335
Ala Ile Ala Leu Leu Lys Glu Ala Ser Lys Ala Val Gly Gln Met Met	340	345	350
Ser Thr Met Phe Tyr Pro Leu Val Thr Phe Val Leu Leu Leu Ile Cys	355	360	365
Ile Ala Tyr Trp Ala Met Thr Ala Leu Tyr Pro Leu Pro Thr Gln Pro	370	375	380
Ala Thr Leu Gly Tyr Val Leu Trp Ala Ser Asn Ile Ser Ser Pro Gly	385	390	395
Cys Glu Lys Val Pro Ile Asn Thr Ser Cys Asn Pro Thr Ala His Leu	405	410	415
Val Asn Ser Ser Cys Pro Gly Leu Met Cys Val Phe Gln Gly Tyr Ser	420	425	430
Ser Lys Gly Leu Ile Gln Arg Ser Val Phe Asn Leu Gln Ile Tyr Gly	435	440	445
Val Leu Gly Leu Phe Trp Thr Leu Asn Trp Val Leu Ala Leu Gly Gln	450	455	460
Cys Val Leu Ala Gly Ala Phe Ala Ser Phe Tyr Trp Ala Phe His Lys	465	470	475
Pro Gln Asp Ile Pro Thr Phe Pro Leu Ile Ser Ala Phe Ile Arg Thr	485	490	495
Leu Arg Tyr His Thr Gly Ser Leu Ala Phe Gly Ala Leu Ile Leu Thr	500	505	510
Leu Val Gln Ile Ala Arg Val Ile Leu Glu Tyr Ile Asp His Lys Leu	515	520	525
Arg Gly Val Gln Asn Pro Val Ala Arg Cys Ile Met Cys Cys Phe Lys	530	535	540
Cys Cys Leu Trp Cys Leu Glu Lys Phe Ile Lys Phe Leu Asn Arg Asn	545	550	555
Ala Tyr Ile Met Ile Ala Ile Tyr Gly Lys Asn Phe Cys Val Ser Ala	565	570	575
Lys Asn Ala Phe Met Leu Leu Met Arg Asn Ile Val Arg Val Val Val	580	585	590
Leu Asp Lys Val Thr Asp Leu Leu Phe Phe Gly Lys Leu Leu Val	595	600	605
Val Gly Gly Val Gly Val Leu Ser Phe Phe Phe Phe Ser Gly Arg Ile	610	615	620
Pro Gly Leu Gly Lys Asp Phe Lys Ser Pro His Leu Asn Tyr Tyr Trp	625	630	635
Leu Pro Ile Met Thr Ser Ile Leu Gly Ala Tyr Val Ile Ala Ser Gly	645	650	655

-continued

Phe Phe Ser Val Phe Gly Met Cys Val Asp Thr Leu Phe Leu Cys Phe
 660 665 670
 Leu Glu Asp Leu Glu Arg Asn Asn Gly Ser Leu Asp Arg Pro Tyr Tyr
 675 680 685
 Met Ser Lys Ser Leu Leu Lys Ile Leu Gly Lys Lys Asn Glu Ala Pro
 690 695 700
 Pro Asp Asn Lys Lys Arg Lys Lys
 705 710

<210> SEQ ID NO 27
 <211> LENGTH: 710
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 27

Met Gly Gly Lys Gln Arg Asp Glu Asp Asp Glu Ala Tyr Gly Lys Pro
 1 5 10 15
 Val Lys Tyr Asp Pro Ser Phe Arg Gly Pro Ile Lys Asn Arg Ser Cys
 20 25 30
 Thr Asp Val Ile Cys Cys Val Leu Phe Leu Leu Phe Ile Leu Gly Tyr
 35 40 45
 Ile Val Val Gly Ile Val Ala Trp Leu Tyr Gly Asp Pro Arg Gln Val
 50 55 60
 Leu Tyr Pro Arg Asn Ser Thr Gly Ala Tyr Cys Gly Met Gly Glu Asn
 65 70 75 80
 Lys Asp Lys Pro Tyr Leu Leu Tyr Phe Asn Ile Phe Ser Cys Ile Leu
 85 90 95
 Ser Ser Asn Ile Ile Ser Val Ala Glu Asn Gly Leu Gln Cys Pro Thr
 100 105 110
 Pro Gln Val Cys Val Ser Ser Cys Pro Glu Asp Pro Trp Thr Val Gly
 115 120 125
 Lys Asn Glu Phe Ser Gln Thr Val Gly Glu Val Phe Tyr Thr Lys Asn
 130 135 140
 Arg Asn Phe Cys Leu Pro Gly Val Pro Trp Asn Met Thr Val Ile Thr
 145 150 155 160
 Ser Leu Gln Gln Glu Leu Cys Pro Ser Phe Leu Leu Pro Ser Ala Pro
 165 170 175
 Ala Leu Gly Arg Cys Phe Pro Trp Thr Asn Val Thr Pro Pro Ala Leu
 180 185 190
 Pro Gly Ile Thr Asn Asp Thr Thr Ile Gln Gln Gly Ile Ser Gly Leu
 195 200 205
 Ile Asp Ser Leu Asn Ala Arg Asp Ile Ser Val Lys Ile Phe Glu Asp
 210 215 220
 Phe Ala Gln Ser Trp Tyr Trp Ile Leu Val Ala Leu Gly Val Ala Leu
 225 230 235 240
 Val Leu Ser Leu Leu Phe Ile Leu Leu Arg Leu Val Ala Gly Pro
 245 250 255
 Leu Val Leu Val Leu Ile Leu Gly Val Leu Gly Val Leu Ala Tyr Gly
 260 265 270
 Ile Tyr Tyr Cys Trp Glu Glu Tyr Arg Val Leu Arg Asp Lys Gly Ala
 275 280 285
 Ser Ile Ser Gln Leu Gly Phe Thr Thr Asn Leu Ser Ala Tyr Gln Ser
 290 295 300
 Val Gln Glu Thr Trp Leu Ala Ala Leu Ile Val Leu Ala Val Leu Glu
 305 310 315 320

Figure 3F. Amino acid sequence of 24P4C12 v.8 (SEQ ID. NO. : 25). The 24P4C12 v.8 protein has 722 amino acids.

```
1  MGGKQRDEDD EAYGKPVKYD PSFRGPIKNR SCTDVICCVL FLLFILGYIV VGIVAWLYGD
61  PRQVLYPRNS TGAYCGMGEN KDKPYLLYFN IFSCILSSNI ISVAENGLQC PTPQVCVSSC
121 PEDPWTVGKN EFSQTVGEVF YTKNRNFCLP GVPWNMTVIT SLQQELCPSE LLPSAPALGR
181 CFPWNTVTPP ALPGITNDTT IQQGISGLID SLNARDISVK IFEDFAQSWY WILVALGVAL
241 VLSLLFILLL RLVAGPLVLV LILGVLGVLA YGIYYCWEEY RVLDRKGASI SQLGFTTNLS
301 AYQSVQETWL AALIVLAVLE AILLMLLIFL RQRIRIAIAL LKEASKAVGQ MMSTMFYPLV
361 TFVLLLLICIA YWAMTALYLA TSGQPQYVLW ASNISSPGCE KVPINTSCNP TAHLVNSSCP
421 GLMCFVQGYG SKGLIQRSVF NLQIYGVGL FWTLNWVLAL GQCVLAGAFA SFYWAFHKPQ
481 DIPTFPLISA FIRTLYHTG SLAFGALILT LVQIARVILE YIDHKLRGVQ NPVARCIMCC
541 FKCCWLCEK FIKFLNRNAY IMIAIYGKNE CVSAKNAFML LMRNIVRVV LDKVTDLLLF
601 FGKLLVGGV GVLSEFFFSG RIPGLGKDFK SPHLNYYWLP IMRNPITPTG HVFQTSILGA
661 YVIASGFFSV FGMCVDTLFL CFLEDLERNN GSLDRPYYS KSLKILGKK NEAPPDNKKR
721 KK
```

Figure 3G. Amino acid sequence of 24P4C12 v.9 (SEQ ID. NO. : 26). The 24P4C12 v.9 protein has 712 amino acids.

```
1  MGGKQRDEDD EAYGKPVKYD PSFRGPIKNR SCTDVICCVL FLLFILGYIV VGIVAWLYGD
61  PRQVLYPRNS TGAYCGMGEN KDKPYLLYFN IFSCILSSNI ISVAENGLQC PTPQVCVSSC
121 PEDPWTVGKN EFSQTVGEVF YTKNRNFCLP GVPWNMTVIT SLQQELCPSE LLPSAPALGR
181 CFPWNTVTPP ALPGITNDTT IQQGISGLID SLNARDISVK IFEDFAQSWY WILVALGVAL
241 VLSLLFILLL RLVAGPLVLV LILGVLGVLA YGIYYCWEEY RVLDRKGASI SQLGFTTNLS
301 AYQSVQETWL AALIVLAVLE AILLMLLIFL RQRIRIAIAL LKEASKAVGQ MMSTMFYPLV
361 TFVLLLLICIA YWAMTALYPL PTQPATLGYV LWASNISSPG CEKVPINTSC NPTAHLVNSS
421 CPGLMCFVQG YSSKGLIORS VFNLQIYGV GLFWTLNWVL ALGQCVLAGA FAFYWAFHK
481 PQDIPTFPLI SAFIRTLYH TGLAFGALI LTLVQIARVI LEYIDHKLRG VQNPVARCIM
541 CCFKCCWLCL EKFIKFLNRN AYIMIAIYGK NCVSAKNAF MLLMRNIVRV VVLDKVIDLL
601 LFFGKLLVVG GVGVLSEFFF SGRIPGLGKD FKSPHLNYYW LPIMTSILGA YVIASGFFSV
661 FGMCVDTLFL CFLEDLERNN GSLDRPYYS KSLKILGKK NEAPPDNKKR KK
```

-continued

Ala	Ile	Leu	Leu	Leu	Met	Leu	Ile	Phe	Leu	Arg	Gln	Arg	Ile	Arg	Ile	
				325					330					335		
Ala	Ile	Ala	Leu	Leu	Lys	Glu	Ala	Ser	Lys	Ala	Val	Gly	Gln	Met	Met	
			340					345					350			
Ser	Thr	Met	Phe	Tyr	Pro	Leu	Val	Thr	Phe	Val	Leu	Leu	Leu	Ile	Cys	
		355					360					365				
Ile	Ala	Tyr	Trp	Ala	Met	Thr	Ala	Leu	Tyr	Leu	Ala	Thr	Ser	Gly	Gln	
	370					375					380					
Pro	Gln	Tyr	Val	Leu	Trp	Ala	Ser	Asn	Ile	Ser	Ser	Pro	Gly	Cys	Glu	
385					390					395					400	
Lys	Val	Pro	Ile	Asn	Thr	Ser	Cys	Asn	Pro	Thr	Ala	His	Leu	Val	Asn	
			405						410					415		
Ser	Ser	Cys	Pro	Gly	Leu	Met	Cys	Val	Phe	Gln	Gly	Tyr	Ser	Ser	Lys	
			420					425					430			
Gly	Leu	Ile	Gln	Arg	Ser	Val	Phe	Asn	Leu	Gln	Ile	Tyr	Gly	Val	Leu	
	435						440					445				
Gly	Leu	Phe	Trp	Thr	Leu	Asn	Trp	Val	Leu	Ala	Leu	Gly	Gln	Cys	Val	
	450					455					460					
Leu	Ala	Gly	Ala	Phe	Ala	Ser	Phe	Tyr	Trp	Ala	Phe	His	Lys	Pro	Gln	
465					470					475					480	
Asp	Ile	Pro	Thr	Phe	Pro	Leu	Ile	Ser	Ala	Phe	Ile	Arg	Thr	Leu	Arg	
			485						490					495		
Tyr	His	Thr	Gly	Ser	Leu	Ala	Phe	Gly	Ala	Leu	Ile	Leu	Thr	Leu	Val	
			500					505					510			
Gln	Ile	Ala	Arg	Val	Ile	Leu	Glu	Tyr	Ile	Asp	His	Lys	Leu	Arg	Gly	
	515						520					525				
Val	Gln	Asn	Pro	Val	Ala	Arg	Cys	Ile	Met	Cys	Cys	Phe	Lys	Cys	Cys	
	530					535					540					
Leu	Trp	Cys	Leu	Glu	Lys	Phe	Ile	Lys	Phe	Leu	Asn	Arg	Asn	Ala	Tyr	
545					550				555						560	
Ile	Met	Ile	Ala	Ile	Tyr	Gly	Lys	Asn	Phe	Cys	Val	Ser	Ala	Lys	Asn	
			565					570						575		
Ala	Phe	Met	Leu	Leu	Met	Arg	Asn	Ile	Val	Arg	Val	Val	Val	Leu	Asp	
			580					585						590		
Lys	Val	Thr	Asp	Leu	Leu	Leu	Phe	Phe	Gly	Lys	Leu	Leu	Val	Val	Gly	
		595					600					605				
Gly	Val	Gly	Val	Leu	Ser	Phe	Phe	Phe	Phe	Ser	Gly	Arg	Ile	Pro	Gly	
	610					615					620					
Leu	Gly	Lys	Asp	Phe	Lys	Ser	Pro	His	Leu	Asn	Tyr	Tyr	Trp	Leu	Pro	
625					630					635					640	
Ile	Met	Thr	Ser	Ile	Leu	Gly	Ala	Tyr	Val	Ile	Ala	Ser	Gly	Phe	Phe	
			645					650						655		
Ser	Val	Phe	Gly	Met	Cys	Val	Asp	Thr	Leu	Phe	Leu	Cys	Phe	Leu	Glu	
			660					665					670			
Asp	Leu	Glu	Arg	Asn	Asn	Gly	Ser	Leu	Asp	Arg	Pro	Tyr	Tyr	Met	Ser	
	675						680					685				
Lys	Ser	Leu	Leu	Lys	Ile	Leu	Gly	Lys	Lys	Asn	Glu	Ala	Pro	Pro	Asp	
	690					695					700					
Asn	Lys	Lys	Arg	Lys	Lys											
705					710											

<210> SEQ ID NO 28

<211> LENGTH: 710

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

-continued

<400> SEQUENCE: 28

```

Met Gly Gly Lys Gln Arg Asp Glu Asp Asp Glu Ala Tyr Gly Lys Pro
 1           5           10           15

Val Lys Tyr Asp Pro Ser Phe Arg Gly Pro Ile Lys Asn Arg Ser Cys
 20           25           30

Thr Asp Val Ile Cys Cys Val Leu Phe Leu Leu Phe Ile Leu Gly Tyr
 35           40           45

Ile Val Val Gly Ile Val Ala Trp Leu Tyr Gly Asp Pro Arg Gln Val
 50           55           60

Leu Tyr Pro Arg Asn Ser Thr Gly Ala Tyr Cys Gly Met Gly Glu Asn
 65           70           75           80

Lys Asp Lys Pro Tyr Leu Leu Tyr Phe Asn Ile Phe Ser Cys Ile Leu
 85           90           95

Ser Ser Asn Ile Ile Ser Val Ala Glu Asn Gly Leu Gln Cys Pro Thr
100           105           110

Pro Gln Val Cys Val Ser Ser Cys Pro Glu Asp Pro Trp Thr Val Gly
115           120           125

Lys Asn Glu Phe Ser Gln Thr Val Gly Glu Val Phe Tyr Thr Lys Ser
130           135           140

Ser Asn Phe Cys Leu Pro Gly Val Pro Trp Asn Met Thr Val Ile Thr
145           150           155           160

Ser Leu Gln Gln Glu Leu Cys Pro Ser Phe Leu Leu Pro Ser Ala Pro
165           170           175

Ala Leu Gly Arg Cys Phe Pro Trp Thr Asn Ile Thr Pro Pro Ala Leu
180           185           190

Pro Gly Ile Thr Asn Asp Thr Thr Ile Gln Gln Gly Ile Ser Gly Leu
195           200           205

Ile Asp Ser Leu Asn Ala Arg Asp Ile Ser Val Lys Ile Phe Glu Asp
210           215           220

Phe Ala Gln Ser Trp Tyr Trp Ile Leu Val Ala Leu Gly Val Ala Leu
225           230           235           240

Val Leu Ser Leu Leu Phe Ile Leu Leu Leu Arg Leu Val Ala Gly Pro
245           250           255

Leu Val Leu Val Leu Ile Leu Gly Val Leu Gly Val Leu Ala Tyr Gly
260           265           270

Ile Tyr Tyr Cys Trp Glu Glu Tyr Arg Val Leu Arg Asp Lys Gly Ala
275           280           285

Ser Ile Ser Gln Leu Gly Phe Thr Thr Asn Leu Ser Ala Tyr Gln Ser
290           295           300

Val Gln Glu Thr Trp Leu Ala Ala Leu Ile Val Leu Ala Val Leu Glu
305           310           315           320

Ala Ile Leu Leu Leu Val Leu Ile Phe Leu Arg Gln Arg Ile Arg Ile
325           330           335

Ala Ile Ala Leu Leu Lys Glu Ala Ser Lys Ala Val Gly Gln Met Met
340           345           350

Ser Thr Met Phe Tyr Pro Leu Val Thr Phe Val Leu Leu Leu Ile Cys
355           360           365

Ile Ala Tyr Trp Ala Met Thr Ala Leu Tyr Leu Ala Thr Ser Gly Gln
370           375           380

Pro Gln Tyr Val Leu Trp Ala Ser Asn Ile Ser Ser Pro Gly Cys Glu
385           390           395           400

Lys Val Pro Ile Asn Thr Ser Cys Asn Pro Thr Ala His Leu Val Asn
405           410           415

```

-continued

Ser Ser Cys Pro Gly Leu Met Cys Val Phe Gln Gly Tyr Ser Ser Lys
 420 425 430
 Gly Leu Ile Gln Arg Ser Val Phe Asn Leu Gln Ile Tyr Gly Val Leu
 435 440 445
 Gly Leu Phe Trp Thr Leu Asn Trp Val Leu Ala Leu Gly Gln Cys Val
 450 455 460
 Leu Ala Gly Ala Phe Ala Ser Phe Tyr Trp Ala Phe His Lys Pro Gln
 465 470 475 480
 Asp Ile Pro Thr Phe Pro Leu Ile Ser Ala Phe Ile Arg Thr Leu Arg
 485 490 495
 Tyr His Thr Gly Ser Leu Ala Phe Gly Ala Leu Ile Leu Thr Leu Val
 500 505 510
 Gln Ile Ala Arg Val Ile Leu Glu Tyr Ile Asp His Lys Leu Arg Gly
 515 520 525
 Val Gln Asn Pro Val Ala Arg Cys Ile Met Cys Cys Phe Lys Cys Cys
 530 535 540
 Leu Trp Cys Leu Glu Lys Phe Ile Lys Phe Leu Asn Arg Asn Ala Tyr
 545 550 555 560
 Ile Met Ile Ala Ile Tyr Gly Lys Asn Phe Cys Val Ser Ala Lys Asn
 565 570 575
 Ala Phe Met Leu Leu Met Arg Asn Ile Val Arg Val Val Val Leu Asp
 580 585 590
 Lys Val Thr Asp Leu Leu Leu Phe Phe Gly Lys Leu Leu Val Val Gly
 595 600 605
 Gly Val Gly Val Leu Ser Phe Phe Phe Ser Gly Arg Ile Pro Gly
 610 615 620
 Leu Gly Lys Asp Phe Lys Ser Pro His Leu Asn Tyr Tyr Trp Leu Pro
 625 630 635 640
 Ile Met Thr Ser Ile Leu Gly Ala Tyr Val Ile Ala Ser Gly Phe Phe
 645 650 655
 Ser Val Phe Gly Met Cys Val Asp Thr Leu Phe Leu Cys Phe Leu Glu
 660 665 670
 Asp Leu Glu Arg Asn Asn Gly Ser Leu Asp Arg Pro Tyr Tyr Met Ser
 675 680 685
 Lys Ser Leu Leu Lys Ile Leu Gly Lys Lys Asn Glu Ala Pro Pro Asp
 690 695 700
 Asn Lys Lys Arg Lys Lys
 705 710

<210> SEQ ID NO 29
 <211> LENGTH: 14
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: synthetically constructed tetanus toxoid

<400> SEQUENCE: 29

Gln Tyr Ile Lys Ala Asn Ser Lys Phe Ile Gly Ile Thr Glu
 1 5 10

<210> SEQ ID NO 30
 <211> LENGTH: 21
 <212> TYPE: PRT
 <213> ORGANISM: Plasmodium falciparum

<400> SEQUENCE: 30

Asp Ile Glu Lys Lys Ile Ala Lys Met Glu Lys Ala Ser Ser Val Phe

-continued

1	5	10	15
Asn Val Val Asn Ser			
20			
<210> SEQ ID NO 31 <211> LENGTH: 16 <212> TYPE: PRT <213> ORGANISM: Streptococcus aureus <400> SEQUENCE: 31			
Gly Ala Val Asp Ser Ile Leu Gly Gly Val Ala Thr Tyr Gly Ala Ala			
1	5	10	15
<210> SEQ ID NO 32 <211> LENGTH: 13 <212> TYPE: PRT <213> ORGANISM: Artificial Sequence <220> FEATURE: <221> NAME/KEY: VARIANT <222> LOCATION: 3 <223> OTHER INFORMATION: Xaa = cyclohexylalanine, phenylalanine, or tyrosine <220> FEATURE: <221> NAME/KEY: VARIANT <222> LOCATION: 1,13 <223> OTHER INFORMATION: Xaa = D-alanine or L-alanine <220> FEATURE: <223> OTHER INFORMATION: PAN DR-binding epitope <400> SEQUENCE: 32			
Xaa Lys Xaa Val Ala Ala Trp Thr Leu Lys Ala Ala Xaa			
1	5	10	
<210> SEQ ID NO 33 <211> LENGTH: 14 <212> TYPE: DNA <213> ORGANISM: Artificial Sequence <220> FEATURE: <223> OTHER INFORMATION: Primer <400> SEQUENCE: 33			
ttttgatcaa gctt			14
<210> SEQ ID NO 34 <211> LENGTH: 42 <212> TYPE: DNA <213> ORGANISM: Artificial Sequence <220> FEATURE: <223> OTHER INFORMATION: Primer <400> SEQUENCE: 34			
ctaatacgac tcactatagg gctcgagcgg ccgcccgggc ag			42
<210> SEQ ID NO 35 <211> LENGTH: 12 <212> TYPE: DNA <213> ORGANISM: Artificial Sequence <220> FEATURE: <223> OTHER INFORMATION: Primer <400> SEQUENCE: 35			
gacctgccc gg			12
<210> SEQ ID NO 36 <211> LENGTH: 40 <212> TYPE: DNA <213> ORGANISM: Artificial Sequence <220> FEATURE:			

-continued

<223> OTHER INFORMATION: Primer

<400> SEQUENCE: 36

gtaatacgac tcactatagg gcagcgtggt cgcggccgag 40

<210> SEQ ID NO 37

<211> LENGTH: 10

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Primer

<400> SEQUENCE: 37

gatacctcggc 10

<210> SEQ ID NO 38

<211> LENGTH: 22

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Primer

<400> SEQUENCE: 38

ctaatacgac tcactatagg gc 22

<210> SEQ ID NO 39

<211> LENGTH: 22

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Primer

<400> SEQUENCE: 39

tcgagcggcc gcccgggcag ga 22

<210> SEQ ID NO 40

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Primer

<400> SEQUENCE: 40

agcgtggtcg cggccgagga 20

<210> SEQ ID NO 41

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Primer

<400> SEQUENCE: 41

atatcgccgc gctcgtcgtc gacaa 25

<210> SEQ ID NO 42

<211> LENGTH: 26

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Primer

<400> SEQUENCE: 42

agccacacgc agtcattgt agaagg 26

<210> SEQ ID NO 43

-continued

<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Primer

<400> SEQUENCE: 43

agatgaggag gaggacaaag gtg 23

<210> SEQ ID NO 44
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Primer

<400> SEQUENCE: 44

actgctggga ggagtaccga gtg 23

<210> SEQ ID NO 45
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Primer

<400> SEQUENCE: 45

gattacaagg atgacgacga taag 24

<210> SEQ ID NO 46
<211> LENGTH: 12
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 46

Arg Asn Pro Ile Thr Pro Thr Gly His Val Phe Gln
1 5 10

<210> SEQ ID NO 47
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 47

Pro Leu Pro Thr Gln Pro Ala Thr Leu Gly
1 5 10

<210> SEQ ID NO 48
<211> LENGTH: 4
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 48

Asn Arg Ser Cys
1

<210> SEQ ID NO 49
<211> LENGTH: 4
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 49

Asn Ser Thr Gly
1

<210> SEQ ID NO 50

-continued

<211> LENGTH: 4
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 50

Asn Met Thr Val
1

<210> SEQ ID NO 51
<211> LENGTH: 4
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 51

Asn Asp Thr Thr
1

<210> SEQ ID NO 52
<211> LENGTH: 4
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 52

Asn Leu Ser Ala
1

<210> SEQ ID NO 53
<211> LENGTH: 4
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 53

Asn Ile Ser Ser
1

<210> SEQ ID NO 54
<211> LENGTH: 4
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 54

Asn Thr Ser Cys
1

<210> SEQ ID NO 55
<211> LENGTH: 4
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 55

Asn Ser Ser Cys
1

<210> SEQ ID NO 56
<211> LENGTH: 4
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 56

Asn Gly Ser Leu
1

<210> SEQ ID NO 57
<211> LENGTH: 4
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

-continued

<400> SEQUENCE: 57

Ser Cys Thr Asp
1<210> SEQ ID NO 58
<211> LENGTH: 4
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 58

Ser Val Ala Glu
1<210> SEQ ID NO 59
<211> LENGTH: 4
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 59

Ser Cys Pro Glu
1<210> SEQ ID NO 60
<211> LENGTH: 4
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 60

Thr Val Gly Glu
1<210> SEQ ID NO 61
<211> LENGTH: 4
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 61

Ser Val Gln Glu
1<210> SEQ ID NO 62
<211> LENGTH: 8
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 62

Arg Asp Glu Asp Asp Glu Ala Tyr
1 5<210> SEQ ID NO 63
<211> LENGTH: 6
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 63

Gly Ala Tyr Cys Gly Met
1 5<210> SEQ ID NO 64
<211> LENGTH: 6
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 64

-continued

Gly Met Gly Glu Asn Lys
1 5

<210> SEQ ID NO 65
<211> LENGTH: 6
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 65

Gly Val Pro Trp Asn Met
1 5

<210> SEQ ID NO 66
<211> LENGTH: 6
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 66

Gly Leu Ile Asp Ser Leu
1 5

<210> SEQ ID NO 67
<211> LENGTH: 6
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 67

Gly Ile Tyr Tyr Cys Trp
1 5

<210> SEQ ID NO 68
<211> LENGTH: 6
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 68

Gly Ala Ser Ile Ser Gln
1 5

<210> SEQ ID NO 69
<211> LENGTH: 6
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 69

Gly Gln Met Met Ser Thr
1 5

<210> SEQ ID NO 70
<211> LENGTH: 6
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 70

Gly Leu Phe Trp Thr Leu
1 5

<210> SEQ ID NO 71
<211> LENGTH: 6
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 71

Gly Ala Phe Ala Ser Phe
1 5

-continued

<210> SEQ ID NO 72
 <211> LENGTH: 3
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 72

Gly Lys Lys
 1

<210> SEQ ID NO 73
 <211> LENGTH: 22
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 73

Leu Phe Ile Leu Leu Leu Arg Leu Val Ala Gly Pro Leu Val Leu Val
 1 5 10 15

Leu Ile Leu Gly Val Leu
 20

<210> SEQ ID NO 74
 <211> LENGTH: 12
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 74

Cys Ile Met Cys Cys Phe Lys Cys Cys Leu Trp Cys
 1 5 10

<210> SEQ ID NO 75
 <211> LENGTH: 710
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 75

Met Gly Gly Lys Gln Arg Asp Glu Asp Asp Glu Ala Tyr Gly Lys Pro
 1 5 10 15

Val Lys Tyr Asp Pro Ser Phe Arg Gly Pro Ile Lys Asn Arg Ser Cys
 20 25 30

Thr Asp Val Ile Cys Cys Val Leu Phe Leu Leu Phe Ile Leu Gly Tyr
 35 40 45

Ile Val Val Gly Ile Val Ala Trp Leu Tyr Gly Asp Pro Arg Gln Val
 50 55 60

Leu Tyr Pro Arg Asn Ser Thr Gly Ala Tyr Cys Gly Met Gly Glu Asn
 65 70 75 80

Lys Asp Lys Pro Tyr Leu Leu Tyr Phe Asn Ile Phe Ser Cys Ile Leu
 85 90 95

Ser Ser Asn Ile Ile Ser Val Ala Glu Asn Gly Leu Gln Cys Pro Thr
 100 105 110

Pro Gln Val Cys Val Ser Ser Cys Pro Glu Asp Pro Trp Thr Val Gly
 115 120 125

Lys Asn Glu Phe Ser Gln Thr Val Gly Glu Val Phe Tyr Thr Lys Asn
 130 135 140

Arg Asn Phe Cys Leu Pro Gly Val Pro Trp Asn Met Thr Val Ile Thr
 145 150 155 160

Ser Leu Gln Gln Glu Leu Cys Pro Ser Phe Leu Leu Pro Ser Ala Pro
 165 170 175

Ala Leu Gly Arg Cys Phe Pro Trp Thr Asn Val Thr Pro Pro Ala Leu
 180 185 190

Figure 4: Alignment of 24P4C12 (SEQ ID NO: 27) with human choline transporter-like protein 4 (CTL4) (gil14249468) (SEQ ID NO: 28)

Score = 1323 bits (3423), Expect = 0.0
Identities = 706/710 (99%), Positives = 709/710 (99%)

24P4C12: 1 MGGKQRDEDEAYGKPKVYDPSFRGPIKNRSCDVIICCVLFLFLFILGYIVVGIVAWLYGD 60
MGGKQRDEDEAYGKPKVYDPSFRGPIKNRSCDVIICCVLFLFLFILGYIVVGIVAWLYGD
CTL-4 : 1 MGGKQRDEDEAYGKPKVYDPSFRGPIKNRSCDVIICCVLFLFLFILGYIVVGIVAWLYGD 60

Query: 61 PRQVLYPRNSTGAYCGMGENKDKPYLLYFNIFSCILSSNIISVAENGLQCPTPQVCVSSC 120
PRQVLYPRNSTGAYCGMGENKDKPYLLYFNIFSCILSSNIISVAENGLQCPTPQVCVSSC
Sbjct: 61 PRQVLYPRNSTGAYCGMGENKDKPYLLYFNIFSCILSSNIISVAENGLQCPTPQVCVSSC 120

Query: 121 PEDPWTIVGKNEFSQTVGEVFTKRNRFCLPGVPWNMTVITSLQQELCPSFLLPSAPALGR 180
PEDPWTIVGKNEFSQTVGEVFTKRNRFCLPGVPWNMTVITSLQQELCPSFLLPSAPALGR
Sbjct: 121 PEDPWTIVGKNEFSQTVGEVFTKRNRFCLPGVPWNMTVITSLQQELCPSFLLPSAPALGR 180

Query: 181 CFPWTNVTTPALPGITNDTTIQQGISGLIDSLNARDISVKIFEDFAQSWYWLVALGVAL 240
CFPWTN+TPALPGITNDTTIQQGISGLIDSLNARDISVKIFEDFAQSWYWLVALGVAL
Sbjct: 181 CFPWTNVTTPALPGITNDTTIQQGISGLIDSLNARDISVKIFEDFAQSWYWLVALGVAL 240

Query: 241 VLSLLFILLRLVAGPLVLVLILGLVGLVAYGIYYCWEEYRVLRDKGASISQLGFTTNLS 300
VLSLLFILLRLVAGPLVLVLILGLVGLVAYGIYYCWEEYRVLRDKGASISQLGFTTNLS
Sbjct: 241 VLSLLFILLRLVAGPLVLVLILGLVGLVAYGIYYCWEEYRVLRDKGASISQLGFTTNLS 300

Query: 301 AYQSVQETWLAALIVLAVLEAILLLMLIFLRQRIRIAIALKEASKAVGQMMSTMFYPLV 360
AYQSVQETWLAALIVLAVLEAILLL+LIFLRQRIRIAIALKEASKAVGQMMSTMFYPLV
Sbjct: 301 AYQSVQETWLAALIVLAVLEAILLLVLIFLRQRIRIAIALKEASKAVGQMMSTMFYPLV 360

Query: 361 TFVLLLLICIAYWAMTALYLATSGQPQYVLWASNISSPGCEKVPINTSCNPTAHLVNSSCP 420
TFVLLLLICIAYWAMTALYLATSGQPQYVLWASNISSPGCEKVPINTSCNPTAHLVNSSCP
Sbjct: 361 TFVLLLLICIAYWAMTALYLATSGQPQYVLWASNISSPGCEKVPINTSCNPTAHLVNSSCP 420

Query: 421 GLMCVFQGYSSKGLIQRSVFNLQIYGVLGFWTLNWWLALGQCVLAFASFYWAFHKPQ 480
GLMCVFQGYSSKGLIQRSVFNLQIYGVLGFWTLNWWLALGQCVLAFASFYWAFHKPQ
Sbjct: 421 GLMCVFQGYSSKGLIQRSVFNLQIYGVLGFWTLNWWLALGQCVLAFASFYWAFHKPQ 480

Query: 481 DIPTFPLISAFIRILRYHTGSLAFGALILTLVQIARVILEYIDHKLRGVQNPVARCIMCC 540
DIPTFPLISAFIRILRYHTGSLAFGALILTLVQIARVILEYIDHKLRGVQNPVARCIMCC
Sbjct: 481 DIPTFPLISAFIRILRYHTGSLAFGALILTLVQIARVILEYIDHKLRGVQNPVARCIMCC 540

Query: 541 FKCLWCLEKFIKFLNRNAYIMIAIYGKNFCVSAKNAFMLLMRNIVRVVLDKVIDLLE 600
FKCLWCLEKFIKFLNRNAYIMIAIYGKNFCVSAKNAFMLLMRNIVRVVLDKVIDLLE
Sbjct: 541 FKCLWCLEKFIKFLNRNAYIMIAIYGKNFCVSAKNAFMLLMRNIVRVVLDKVIDLLE 600

Query: 601 FGKLLVVGGVGLSFFFFSGRIPGLGKDFKSPHLNYYWLPIMTSILGAYVIASGFFSVFG 660
FGKLLVVGGVGLSFFFFSGRIPGLGKDFKSPHLNYYWLPIMTSILGAYVIASGFFSVFG
Sbjct: 601 FGKLLVVGGVGLSFFFFSGRIPGLGKDFKSPHLNYYWLPIMTSILGAYVIASGFFSVFG 660

Query: 661 MCVDTLFLCFLEDLERNNGSLDRPYMSKSLKILGKKNEAPPDNKKRKK 710
MCVDTLFLCFLEDLERNNGSLDRPYMSKSLKILGKKNEAPPDNKKRKK
Sbjct: 661 MCVDTLFLCFLEDLERNNGSLDRPYMSKSLKILGKKNEAPPDNKKRKK 710

-continued

Pro Gly Ile Thr Asn Asp Thr Thr Ile Gln Gln Gly Ile Ser Gly Leu	195	200	205
Ile Asp Ser Leu Asn Ala Arg Asp Ile Ser Val Lys Ile Phe Glu Asp	210	215	220
Phe Ala Gln Ser Trp Tyr Trp Ile Leu Val Ala Leu Gly Val Ala Leu	225	230	235
Val Leu Ser Leu Leu Phe Ile Leu Leu Leu Arg Leu Val Ala Gly Pro	245	250	255
Leu Val Leu Val Leu Ile Leu Gly Val Leu Gly Val Leu Ala Tyr Gly	260	265	270
Ile Tyr Tyr Cys Trp Glu Glu Tyr Arg Val Leu Arg Asp Lys Gly Ala	275	280	285
Ser Ile Ser Gln Leu Gly Phe Thr Thr Asn Leu Ser Ala Tyr Gln Ser	290	295	300
Val Gln Glu Thr Trp Leu Ala Ala Leu Ile Val Leu Ala Val Leu Glu	305	310	315
Ala Ile Leu Leu Leu Met Leu Ile Phe Leu Arg Gln Arg Ile Arg Ile	325	330	335
Ala Ile Ala Leu Leu Lys Glu Ala Ser Lys Ala Val Gly Gln Met Met	340	345	350
Ser Thr Met Phe Tyr Pro Leu Val Thr Phe Val Leu Leu Leu Ile Cys	355	360	365
Ile Ala Tyr Trp Ala Met Thr Ala Leu Tyr Leu Ala Thr Ser Gly Gln	370	375	380
Pro Gln Tyr Val Leu Trp Ala Ser Asn Ile Ser Ser Pro Gly Cys Glu	385	390	395
Lys Val Pro Ile Asn Thr Ser Cys Asn Pro Thr Ala His Leu Val Asn	405	410	415
Ser Ser Cys Pro Gly Leu Met Cys Val Phe Gln Gly Tyr Ser Ser Lys	420	425	430
Gly Leu Ile Gln Arg Ser Val Phe Asn Leu Gln Ile Tyr Gly Val Leu	435	440	445
Gly Leu Phe Trp Thr Leu Asn Trp Val Leu Ala Leu Gly Gln Cys Val	450	455	460
Leu Ala Gly Ala Phe Ala Ser Phe Tyr Trp Ala Phe His Lys Pro Gln	465	470	475
Asp Ile Pro Thr Phe Pro Leu Ile Ser Ala Phe Ile Arg Thr Leu Arg	485	490	495
Tyr His Thr Gly Ser Leu Ala Phe Gly Ala Leu Ile Leu Thr Leu Val	500	505	510
Gln Ile Ala Arg Val Ile Leu Glu Tyr Ile Asp His Lys Leu Arg Gly	515	520	525
Val Gln Asn Pro Val Ala Arg Cys Ile Met Cys Cys Phe Lys Cys Cys	530	535	540
Leu Trp Cys Leu Glu Lys Phe Ile Lys Phe Leu Asn Arg Asn Ala Tyr	545	550	555
Ile Met Ile Ala Ile Tyr Gly Lys Asn Phe Cys Val Ser Ala Lys Asn	565	570	575
Ala Phe Met Leu Leu Met Arg Asn Ile Val Arg Val Val Val Leu Asp	580	585	590
Lys Val Thr Asp Leu Leu Leu Phe Phe Gly Lys Leu Leu Val Val Gly	595	600	605
Gly Val Gly Val Leu Ser Phe Phe Phe Phe Ser Gly Arg Ile Pro Gly	610	615	620

-continued

Leu Gly Lys Asp Phe Lys Ser Pro His Leu Asn Tyr Tyr Trp Leu Pro
 625 630 635 640
 Ile Met Thr Ser Ile Leu Gly Ala Tyr Val Ile Ala Ser Gly Phe Phe
 645 650 655
 Ser Val Phe Gly Met Cys Val Asp Thr Leu Phe Leu Cys Phe Leu Glu
 660 665 670
 Asp Leu Glu Arg Asn Asn Gly Ser Leu Asp Arg Pro Tyr Tyr Met Ser
 675 680 685
 Lys Ser Leu Leu Lys Ile Leu Gly Lys Lys Asn Glu Ala Pro Pro Asp
 690 695 700
 Asn Lys Lys Arg Lys Lys
 705 710

<210> SEQ ID NO 76
 <211> LENGTH: 17
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 76

Gly Arg Cys Phe Pro Trp Thr Asn Ile Thr Pro Pro Ala Leu Pro Gly
 1 5 10 15

Ile

<210> SEQ ID NO 77
 <211> LENGTH: 19
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 77

Leu Gly Arg Cys Phe Pro Trp Thr Asn Ile Thr Pro Pro Ala Leu Pro
 1 5 10 15

Gly Ile Thr

<210> SEQ ID NO 78
 <211> LENGTH: 29
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 78

Pro Ser Ala Pro Ala Leu Gly Arg Cys Phe Pro Trp Thr Asn Ile Thr
 1 5 10 15

Pro Pro Ala Leu Pro Gly Ile Thr Asn Asp Thr Thr Ile
 20 25

<210> SEQ ID NO 79
 <211> LENGTH: 17
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 79

Val Leu Glu Ala Ile Leu Leu Leu Val Leu Ile Phe Leu Arg Gln Arg
 1 5 10 15

Ile

<210> SEQ ID NO 80
 <211> LENGTH: 19
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 80

-continued

Ala Val Leu Glu Ala Ile Leu Leu Leu Val Leu Ile Phe Leu Arg Gln
 1 5 10 15

Arg Ile Arg

<210> SEQ ID NO 81
 <211> LENGTH: 29
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 81

Ala Leu Ile Val Leu Ala Val Leu Glu Ala Ile Leu Leu Leu Val Leu
 1 5 10 15

Ile Phe Leu Arg Gln Arg Ile Arg Ile Ala Ile Ala Leu
 20 25

<210> SEQ ID NO 82
 <211> LENGTH: 17
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 82

Gly Tyr Ser Ser Lys Gly Leu Ile Pro Arg Ser Val Phe Asn Leu Gln
 1 5 10 15

Ile

<210> SEQ ID NO 83
 <211> LENGTH: 19
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 83

Gln Gly Tyr Ser Ser Lys Gly Leu Ile Pro Arg Ser Val Phe Asn Leu
 1 5 10 15

Gln Ile Tyr

<210> SEQ ID NO 84
 <211> LENGTH: 29
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 84

Leu Met Cys Val Phe Gln Gly Tyr Ser Ser Lys Gly Leu Ile Pro Arg
 1 5 10 15

Ser Val Phe Asn Leu Gln Ile Tyr Gly Val Leu Gly Leu
 20 25

<210> SEQ ID NO 85
 <211> LENGTH: 16
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 85

Ser Trp Tyr Trp Ile Leu Val Ala Val Gly Gln Met Met Ser Thr Met
 1 5 10 15

<210> SEQ ID NO 86
 <211> LENGTH: 18
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 86

Gln Ser Trp Tyr Trp Ile Leu Val Ala Val Gly Gln Met Met Ser Thr
 1 5 10 15

-continued

Met Phe

<210> SEQ ID NO 87
 <211> LENGTH: 28
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 87

Phe Glu Asp Phe Ala Gln Ser Trp Tyr Trp Ile Leu Val Ala Val Gly
 1 5 10 15

Gln Met Met Ser Thr Met Phe Tyr Pro Leu Val Thr
 20 25

<210> SEQ ID NO 88
 <211> LENGTH: 28
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 88

Asn Tyr Tyr Trp Leu Pro Ile Met Arg Asn Pro Ile Thr Pro Thr Gly
 1 5 10 15

His Val Phe Gln Thr Ser Ile Leu Gly Ala Tyr Val
 20 25

<210> SEQ ID NO 89
 <211> LENGTH: 30
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 89

Leu Asn Tyr Tyr Trp Leu Pro Ile Met Arg Asn Pro Ile Thr Pro Thr
 1 5 10 15

Gly His Val Phe Gln Thr Ser Ile Leu Gly Ala Tyr Val Ile
 20 25 30

<210> SEQ ID NO 90
 <211> LENGTH: 40
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 90

Phe Lys Ser Pro His Leu Asn Tyr Tyr Trp Leu Pro Ile Met Arg Asn
 1 5 10 15

Pro Ile Thr Pro Thr Gly His Val Phe Gln Thr Ser Ile Leu Gly Ala
 20 25 30

Tyr Val Ile Ala Ser Gly Phe Phe
 35 40

<210> SEQ ID NO 91
 <211> LENGTH: 26
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 91

Tyr Trp Ala Met Thr Ala Leu Tyr Pro Leu Pro Thr Gln Pro Ala Thr
 1 5 10 15

Leu Gly Tyr Val Leu Trp Ala Ser Asn Ile
 20 25

<210> SEQ ID NO 92
 <211> LENGTH: 28
 <212> TYPE: PRT

-continued

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 92

Ala Tyr Trp Ala Met Thr Ala Leu Tyr Pro Leu Pro Thr Gln Pro Ala
 1 5 10 15

Thr Leu Gly Tyr Val Leu Trp Ala Ser Asn Ile Ser
 20 25

<210> SEQ ID NO 93

<211> LENGTH: 38

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 93

Leu Leu Ile Cys Ile Ala Tyr Trp Ala Met Thr Ala Leu Tyr Pro Leu
 1 5 10 15

Pro Thr Gln Pro Ala Thr Leu Gly Tyr Val Leu Trp Ala Ser Asn Ile
 20 25 30

Ser Ser Pro Gly Cys Glu
 35

<210> SEQ ID NO 94

<211> LENGTH: 2251

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 94

gagccatggg gggaaagcag cgggacgagg atgacgaggc ctacgggaag ccagtcaaat 60
 acgacccttc ctttcgaggc cccatcaaga acagaagctg cacagatgtc atctgtctgcg 120
 tcctcttctc gctcttcatt ctagggtaca tcgtggtggg gattgtggcc tggttgtatg 180
 gagacccccg gcaagtcctc taccacagga actctactgg ggcctactgt ggcattggggg 240
 agaacaaaga taagccgtat ctctgtact tcaacatctt cagctgcac ctgtccagca 300
 acatcatctc agttgctgag aacggcctac agtgccccc accccagggtg tgtgtgtcct 360
 cctgcccgga ggacccatgg actgtgggaa aaaacgagtt ctcacagact gttggggaag 420
 tcttctatca aaaaaacagg aacttttgtc tgccaggggt accctggaat atgacgggtga 480
 tcacaagcct gcaacaggaa ctctgcccga gtttctctct cccctctgct ccagctcttg 540
 ggcgctgctt tccatggacc aacgttactc caccggcgct cccagggatc accaatgaca 600
 ccaccatata gcaggggatc agcgggtctta ttgacagcct caatgcccga gacatcagtg 660
 ttaagatctt tgaagatttt gcccagtcct ggtattggat tctgtggct gtgggacaga 720
 tgatgtctac catgttctac ccaactggta cctttgtcct cctcctcacc tgcattgcct 780
 actgggccat gactgctctg tacctggcta catcggggca accccagtat gtgctctggg 840
 catccaacat cagctccccc ggctgtgaga aagtgcgaat aaatacatca tgcaacccca 900
 cggccacact tgtgaactcc tcgtgccag ggctgatgtg cgtcttcag ggcactcat 960
 ccaaaggcct aatccaacgt tctgtcttca atctgcaaat ctatgggggc ctggggctct 1020
 tctggacct taactgggta ctggccctgg gccaatgcgt cctcgctgga gcctttgcct 1080
 ccttctactg ggccttcac aagccccagg acatccctac cttccctta atctctgcct 1140
 tcacccgcac actccgttac cacactgggt cattggcatt tggagccctc atcctgaccc 1200
 ttgtgcagat agccccgggc atcttgagat atattgacca caagctcaga ggagtgcaga 1260
 accctgtagc ccgctgcacc atgtgtgttt tcaagtgtg cctctggtgt ctggaaaaat 1320
 ttatcaagtt cctaaaccgc aatgcataca tcatgatcgc catctacggg aagaatttct 1380

-continued

```

gtgtctcagc caaaaatgcg ttcatgctac tcatgcgaaa cattgtcagg gtggtegtcc 1440
tggacaaagt cacagacctg ctgctgttct ttgggaagct gctgggtggc ggaggcgtgg 1500
gggtcctgtc cttctttttt ttctccgggc gcatcccggg gctgggtaaa gactttaaga 1560
gccccacct caactattac tggtgcccc tcatgacctc catcctgggg gcctatgtca 1620
tcgccagcgg cttcttcagc gttttcggca tgtgtgtgga cacgtcttc ctctgtctcc 1680
tggaaagacct ggagcggaac aacgggtccc tggaccggcc ctactacatg tccaagagcc 1740
ttctaaagat tctgggcaag aagaacgagg cgtcccgga caacaagaag aggaagaagt 1800
gacagctccg gcctgatcc aggaactgcac cccaccccca ccgtccagcc atccaacctc 1860
acttcgctt acaggtctcc attttgggt aaaaaaggt tttaggccag gcgccgtggc 1920
tcacgcctgt aatccaacac tttgagaggc tgaggcgggc ggatcacctg agtcaggagt 1980
tcgagaccag cctggccaac atggtgaaac ctccgtctct attaaaaata caaaaattag 2040
ccgagagtgg tggcatgcac ctgcatccc agctactcgg gaggtgagg caggagaatc 2100
gcttgaaccc gggaggcaga ggttgagtg agccgagatc gcgccactgc actccaacct 2160
gggtgacaga ctctgtctcc aaacaaaaa aaacaacaa aaagatttta ttaaagatat 2220
ttgttaact cagtaaaaa aaaaaaaaa a 2251

```

<210> SEQ ID NO 95

<211> LENGTH: 2587

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 95

```

gagccatggg gggaaagcag cgggacgagg atgacgaggc ctacgggaag ccagtcaa 60
acgaccctc ctttcgaggc cccatcaaga acagaagctg cacagatgtc atctgtgcg 120
tcctcttct gctcttcatt ctagggtaca tcgtggtggg gattgtggcc tggttgtatg 180
gagaccccg gcaagtcctc taccacagga actctactgg ggctactgt ggcattgggg 240
agaacaaaga taagccgtat ctctgtact tcaacatct cagctgcac ctgtccagca 300
acatcatctc agttgctgag aacggcctac agtgccccc accccagggtg tgtgtgtcct 360
cctgcccgga ggacccatgg actgtgggaa aaaacgagtt ctacagact gttggggaag 420
tcttctatac aaaaaacagg aacttttgtc tgccgggggt accctggaat atgacgggtga 480
tcacaagcct gcaacaggaa ctctgcccc gtttctctct cccctctgct ccagctctgg 540
ggcgctgctt tccatggacc aacgttactc caccggcgct ccagggatc accaatgaca 600
ccaccatata gcaggggcat agcgggtctta ttgacagcct caatgcccga gacatcagt 660
ttaagatctt tgaagatttt gccagtcct ggtattggat tctgttgcc ctgggggtgg 720
ctctggctct gagectactg tttatcttgc ttctgcgct ggtgggtggg cccctgggtg 780
tggtgctgat cctgggagtg ctgggcgtgc tggcatacgg catctactac tgctgggagg 840
agtaccgagt gctgcgggac aaggggcgct ccactctcca gctgggttcc accaccaacc 900
tcagtgccta ccagagcgtg caggagacct ggctggccgc cctgatcgtg ttggcggtgc 960
ttgaagccat cctgctgctg atgctcatct tctgcgga gcggattcgt attgccatcg 1020
ccctcctgaa ggaggccagc aaggctgtgg gacagatgat gtctaccatg ttctaccac 1080
tggtcacctt tgtcctctc ctcatctgca ttgcctactg ggccatgact gctctgtacc 1140
tggctacatc ggggcaaccc cagtatgtgc tctgggcac caacatcagc tccccggct 1200
gtgagaaagt gccaataaat acatcatgca accccacggc ccacctgtg aactcctcgt 1260

```

-continued

```

gccagggct gatgtgcgtc ttccagggct actcatccaa aggcctaata caacgtttctg 1320
tcttcaatct gcaaatctat ggggtcctgg ggctcttctg gacccttaac tgggtactgg 1380
ccctgggcca atcggtcctc gctggagcct ttgcctcctt ctactgggce ttccacaagc 1440
cccaggacat cctacette ccttaatct ctgccttcat cgcacactc cgttaccaca 1500
ctgggtcatt ggcatttggg gccctcatcc tgacccttgt gcagatagcc cgggtcatct 1560
tggagtatat tgaccacaag ctacagaggag tgcagaaccc tgtagccgc tgcacatgt 1620
gctgtttcaa gtgctgcctc tgggtgtctgg aaaaatttat caagtccca aaccgcaatg 1680
catacatcat gatcgccatc tacgggaaga atttctgtgt ctacgcaaaa aatgcgttca 1740
tgctactcat gcgaaacatt gtcaggggtg tcgtcctgga caaagtcaca gacctgctgc 1800
tgttctttgg gaagctgctg gtggtcggag gcgtgggggt cctgtccttc ttttttttct 1860
ccggtcgcac cccggggctg ggtaaaagct ttaagagccc ccacctcaac tattactggc 1920
tgcccatcat gaectccatc ctgggggctc atgtcatcgc cagcggttc ttcagegttt 1980
tcggcatgtg tgtggacacg ctcttctctc gcttctgga agacctggag cggaacaacg 2040
gctccctgga ccggccctac tacatgtcca agagccttct aaagattctg ggcaagaaga 2100
acgaggcgcc ccgggacaac aagaagagga agaagtgaca gctccggccc tgatccagga 2160
ctgcacccca cccccaccgt ccagccatcc aacctcactt cgccttacag gtctccattt 2220
tgtggtaaaa aaaggtttta ggcaggcgc cgtggctcac gcctgtaac caacactttg 2280
agaggctgag gcgggcgcat cactgagtc aggagtcca gaccagcctg gccaacatgg 2340
tgaaacctcc gtctctatta aaaatacaaa aattagccga gagtgggtggc atgcacctgt 2400
catcccagct actcgggagg ctgaggcagg agaatcgctt gaaccggga ggcagaggtt 2460
gcagtgagcc gagatcgcc cactgcactc caacctgggt gacagactct gtctccaaaa 2520
caaaacaaac aaacaaaaag attttattaa agatatatttg ttaactcagt aaaaaaaaaa 2580
aaaaaaa 2587

```

```

<210> SEQ ID NO 96
<211> LENGTH: 2251
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 96

```

```

gagccatggg gggaaagcag cgggacgagg atgacgaggc ctacgggaag ccagtcaaat 60
acgacccctc ctttcgaggc cccatcaaga acagaagctg cacagatgtc atctgctgag 120
tcctcttctc gctcttcatt ctagggtaca tcgtgggtgg gattgtggcc tggttgtatg 180
gagacccccg gcaagtccct taccccagga actctactgg ggctactgt ggcattgggg 240
agaacaaaga taagccgtat ctctgtact tcaacatctt cagctgcac cgttcagca 300
acatcatctc agttgctgag aacggcctac agtgccccc acccagggtg tgtgtgtcct 360
cctgcccgga ggaccatgg actgtgggaa aaaacgagtt ctacagact gttggggaag 420
tcttctatac aaaaaacagg aacttttgct tgccagggtg accctggaat atgacggtga 480
tcacaagcct gcaacaggaa ctctgcccca gtttctctct cccctctgct ccagctctgg 540
ggcgtgctt tccatggacc aacgttaact caccggcgct cccagggatc accaatgaca 600
ccaccataca gcaggggatc agcgggtctta ttgacagcct caatgccga gacatcagtg 660
ttaagatctt tgaagatttt gccagtcct ggtattggat tcttggtggt gtgggacaga 720
tgatgtctac catgttctac ccaatggcca cctttgtcct cctcctcacc tgcattgcct 780

```

-continued

```

actgggccaat gactgctctg tacctggcta catcggggca accccagtat gtgctctggg      840
catccaacat cagctcccc  ggctgtgaga aagtgccaat aaatacatca tgcaacccca      900
cggcccacct tgtgaactcc tcgtgccag ggctgatgtg cgtcttcag ggctactcat      960
ccaaaggcct aatccaacgt tctgtcttca atctgcaaat ctatggggtc ctggggctct 1020
tctggaccct taactgggta ctggccctgg gccaatgcgt cctcgctgga gcctttgcct 1080
ccttctactg ggctctccac aagccccagg acatccctac cttccctta atctctgcct 1140
tcateccgac actccgttac cacactgggt cattggcatt tggagccctc atcctgaccc 1200
ttgtgcagat agccccgggc atcttgagat atattgacca caagctcaga ggagtgcaga 1260
accctgtagc ccgtgcgcat atgtgctgtt tcaagtgtg cctctggtgt ctggaaaaat 1320
ttatcaagtt cctaaaccgc aatgcataca tcatgatcgc catctacggg aagaatttct 1380
gtgtctcagc caaaaatgcy ttcatgctac tcatgcgaaa cattgtcagg gtggtcgtcc 1440
tggacaaaagt cacagacctg ctgctgttct ttgggaagct gctgggtggtc ggaggcgtgg 1500
gggtcctgtc cttctttttt ttctccgggc gcatccgggg gctgggtaaa gactttaaga 1560
gccccacact caactattac tggctgcccc tcatgacctc catcctgggg gcctatgtca 1620
tcgccagcgg cttcttcagc gttttcggca tgtgtgtgga cacgtcttc ctctgcttcc 1680
tggaagacct ggagcggaac aacggctccc tggaccgggc ctactacatg tccaagagcc 1740
ttctaagat tctgggcaag aagaacgagg cgcctccgga caacaagaag aggaagaagt 1800
gacagctccg gccctgatcc aggactgcac cccaccccca ccgtccagcc atccaacctc 1860
acttcgctt acaggtctcc attttgtgtt aaaaaaagg tttaggccag gcgccgtggc 1920
tcacgcctgt aatccaacac tttgagaggc tgaggcgggc ggatcacctg agtcaggagt 1980
tcgagaccag cctggccaac atggtgaaac ctccgtctct attaaaaata caaaaattag 2040
ccgagagtgg tggcatgcac ctgtcatccc agctactcgg gaggtgagg caggagaatc 2100
gcttgaaccc gggaggcaga gggtgcagtg agccgagatc gcgccactgc actccaacct 2160
gggtgacaga ctctgtctcc aaaacaaaac aaacaacaa aaagatttta ttaaagatat 2220
tttgtaact cagtaaaaaa aaaaaaaaaa a                                     2251

```

<210> SEQ ID NO 97

<211> LENGTH: 598

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 97

```

Met Gly Gly Lys Gln Arg Asp Glu Asp Asp Glu Ala Tyr Gly Lys Pro
 1             5             10             15

Val Lys Tyr Asp Pro Ser Phe Arg Gly Pro Ile Lys Asn Arg Ser Cys
 20             25             30

Thr Asp Val Ile Cys Cys Val Leu Phe Leu Leu Phe Ile Leu Gly Tyr
 35             40             45

Ile Val Val Gly Ile Val Ala Trp Leu Tyr Gly Asp Pro Arg Gln Val
 50             55             60

Leu Tyr Pro Arg Asn Ser Thr Gly Ala Tyr Cys Gly Met Gly Glu Asn
 65             70             75             80

Lys Asp Lys Pro Tyr Leu Leu Tyr Phe Asn Ile Phe Ser Cys Ile Leu
 85             90             95

Ser Ser Asn Ile Ile Ser Val Ala Glu Asn Gly Leu Gln Cys Pro Thr
100             105             110

```

Pro	Gln	Val	Cys	Val	Ser	Ser	Cys	Pro	Glu	Asp	Pro	Trp	Thr	Val	Gly
115							120					125			
Lys	Asn	Glu	Phe	Ser	Gln	Thr	Val	Gly	Glu	Val	Phe	Tyr	Thr	Lys	Asn
130						135					140				
Arg	Asn	Phe	Cys	Leu	Pro	Gly	Val	Pro	Trp	Asn	Met	Thr	Val	Ile	Thr
145					150					155					160
Ser	Leu	Gln	Gln	Glu	Leu	Cys	Pro	Ser	Phe	Leu	Leu	Pro	Ser	Ala	Pro
				165					170					175	
Ala	Leu	Gly	Arg	Cys	Phe	Pro	Trp	Thr	Asn	Val	Thr	Pro	Pro	Ala	Leu
			180					185				190			
Pro	Gly	Ile	Thr	Asn	Asp	Thr	Thr	Ile	Gln	Gln	Gly	Ile	Ser	Gly	Leu
195						200					205				
Ile	Asp	Ser	Leu	Asn	Ala	Arg	Asp	Ile	Ser	Val	Lys	Ile	Phe	Glu	Asp
210						215					220				
Phe	Ala	Gln	Ser	Trp	Tyr	Trp	Ile	Leu	Val	Ala	Val	Gly	Gln	Met	Met
225					230					235					240
Ser	Thr	Met	Phe	Tyr	Pro	Leu	Val	Thr	Phe	Val	Leu	Leu	Leu	Ile	Cys
				245					250					255	
Ile	Ala	Tyr		Ala	Met	Thr	Ala	Leu	Tyr	Leu	Ala	Thr	Ser	Gly	Gln
260								265					270		
Pro	Gln	Tyr	Val	Leu	Trp	Ala	Ser	Asn	Ile	Ser	Ser	Pro	Gly	Cys	Glu
275						280						285			
Lys	Val	Pro	Ile	Asn	Thr	Ser	Cys	Asn	Pro	Thr	Ala	His	Leu	Val	Asn
290						295					300				
Ser	Ser	Cys	Pro	Gly	Leu	Met	Cys	Val	Phe	Gln	Gly	Tyr	Ser	Ser	Lys
305					310					315					320
Gly	Leu	Ile	Gln	Arg	Ser	Val	Phe	Asn	Leu	Gln	Ile	Tyr	Gly	Val	Leu
				325					330					335	
Gly	Leu	Phe	Trp	Thr	Leu	Asn	Trp	Val	Leu	Ala	Leu	Gly	Gln	Cys	Val
				340				345					350		
Leu	Ala	Gly	Ala	Phe	Ala	Ser	Phe	Tyr	Trp	Ala	Phe	His	Lys	Pro	Gln
355							360				365				
Asp	Ile	Pro	Thr	Phe	Pro	Leu	Ile	Ser	Ala	Phe	Ile	Arg	Thr	Leu	Arg
370						375					380				
Tyr	His	Thr	Gly	Ser	Leu	Ala	Phe	Gly	Ala	Leu	Ile	Leu	Thr	Leu	Val
385					390					395					400
Gln	Ile	Ala	Arg	Val	Ile	Leu	Glu	Tyr	Ile	Asp	His	Lys	Leu	Arg	Gly
			405						410					415	
Val	Gln	Asn	Pro	Val	Ala	Arg	Cys	Ile	Met	Cys	Cys	Phe	Lys	Cys	Cys
			420					425					430		
Leu	Trp	Cys	Leu	Glu	Lys	Phe	Ile	Lys	Phe	Leu	Asn	Arg	Asn	Ala	Tyr
435							440					445			
Ile	Met	Ile	Ala	Ile	Tyr	Gly	Lys	Asn	Phe	Cys	Val	Ser	Ala	Lys	Asn
450						455					460				
Ala	Phe	Met	Leu	Leu	Met	Arg	Asn	Ile	Val	Arg	Val	Val	Val	Leu	Asp
465					470										

-continued

Ser Val Phe Gly Met Cys Val Asp Thr Leu Phe Leu Cys Phe Leu Glu
545 550 555 560

Asp Leu Glu Arg Asn Asn Gly Ser Leu Asp Arg Pro Tyr Tyr Met Ser
565 570 575

Lys Ser Leu Leu Lys Ile Leu Gly Lys Lys Asn Glu Ala Pro Pro Asp
580 585 590

Asn Lys Lys Arg Lys Lys
595

<210> SEQ ID NO 98
 <211> LENGTH: 710
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 98

Met Gly Gly Lys Gln Arg Asp Glu Asp Asp Glu Ala Tyr Gly Lys Pro
1 5 10 15

Val Lys Tyr Asp Pro Ser Phe Arg Gly Pro Ile Lys Asn Arg Ser Cys
20 25 30

Thr Asp Val Ile Cys Cys Val Leu Phe Leu Leu Phe Ile Leu Gly Tyr
35 40 45

Ile Val Val Gly Ile Val Ala Trp Leu Tyr Gly Asp Pro Arg Gln Val
50 55 60

Leu Tyr Pro Arg Asn Ser Thr Gly Ala Tyr Cys Gly Met Gly Glu Asn
65 70 75 80

Lys Asp Lys Pro Tyr Leu Leu Tyr Phe Asn Ile Phe Ser Cys Ile Leu
85 90 95

Ser Ser Asn Ile Ile Ser Val Ala Glu Asn Gly Leu Gln Cys Pro Thr
100 105 110

Pro Gln Val Cys Val Ser Ser Cys Pro Glu Asp Pro Trp Thr Val Gly
115 120 125

Lys Asn Glu Phe Ser Gln Thr Val Gly Glu Val Phe Tyr Thr Lys Asn
130 135 140

Arg Asn Phe Cys Leu Pro Gly Val Pro Trp Asn Met Thr Val Ile Thr
145 150 155 160

Ser Leu Gln Gln Glu Leu Cys Pro Ser Phe Leu Leu Pro Ser Ala Pro
165 170 175

Ala Leu Gly Arg Cys Phe Pro Trp Thr Asn Val Thr Pro Pro Ala Leu
180 185 190

Pro Gly Ile Thr Asn Asp Thr Thr Ile Gln Gln Gly Ile Ser Gly Leu
195 200 205

Ile Asp Ser Leu Asn Ala Arg Asp Ile Ser Val Lys Ile Phe Glu Asp
210 215 220

Phe Ala Gln Ser Trp Tyr Trp Ile Leu Val Ala Leu Gly Val Ala Leu
225 230 235 240

Val Leu Ser Leu Leu Phe Ile Leu Leu Leu Arg Leu Val Ala Gly Pro
245 250 255


Leu Val Leu Val Leu Ile Leu Gly Val Leu Gly Val Leu Ala Tyr Gly
260 265 270

Ile Tyr Tyr Cys Trp Glu Glu Tyr Arg Val Leu Arg Asp Lys Gly Ala
275 280 285

Ser Ile Ser Gln Leu Gly Phe Thr Thr Asn Leu Ser Ala Tyr Gln Ser
290 295 300

Val Gln Glu Thr Trp Leu Ala Ala Leu Ile Val Leu Ala Val Leu Glu
305 310 315 320

Figure 5: 24P4C12 variant 1 Hydrophilicity profile
(Hopp T.P., Woods K.R., 1981.
Proc. Natl. Acad. Sci. U.S.A. 78:3824-3828)



-continued

Ala Ile Leu Leu Leu Met Leu Ile Phe Leu Arg Gln Arg Ile Arg Ile
 325 330 335
 Ala Ile Ala Leu Leu Lys Glu Ala Ser Lys Ala Val Gly Gln Met Met
 340 345 350
 Ser Thr Met Phe Tyr Pro Leu Val Thr Phe Val Leu Leu Leu Ile Cys
 355 360 365
 Ile Ala Tyr Trp Ala Met Thr Ala Leu Tyr Leu Ala Thr Ser Gly Gln
 370 375 380
 Pro Gln Tyr Val Leu Trp Ala Ser Asn Ile Ser Ser Pro Gly Cys Glu
 385 390 395 400
 Lys Val Pro Ile Asn Thr Ser Cys Asn Pro Thr Ala His Leu Val Asn
 405 410 415
 Ser Ser Cys Pro Gly Leu Met Cys Val Phe Gln Gly Tyr Ser Ser Lys
 420 425 430
 Gly Leu Ile Gln Arg Ser Val Phe Asn Leu Gln Ile Tyr Gly Val Leu
 435 440 445
 Gly Leu Phe Trp Thr Leu Asn Trp Val Leu Ala Leu Gly Gln Cys Val
 450 455 460
 Leu Ala Gly Ala Phe Ala Ser Phe Tyr Trp Ala Phe His Lys Pro Gln
 465 470 475 480
 Asp Ile Pro Thr Phe Pro Leu Ile Ser Ala Phe Ile Arg Thr Leu Arg
 485 490 495
 Tyr His Thr Gly Ser Leu Ala Phe Gly Ala Leu Ile Leu Thr Leu Val
 500 505 510
 Gln Ile Ala Arg Val Ile Leu Glu Tyr Ile Asp His Lys Leu Arg Gly
 515 520 525
 Val Gln Asn Pro Val Ala Arg Cys Ile Met Cys Cys Phe Lys Cys Cys
 530 535 540
 Leu Trp Cys Leu Glu Lys Phe Ile Lys Phe Leu Asn Arg Asn Ala Tyr
 545 550 555 560
 Ile Met Ile Ala Ile Tyr Gly Lys Asn Phe Cys Val Ser Ala Lys Asn
 565 570 575
 Ala Phe Met Leu Leu Met Arg Asn Ile Val Arg Val Val Val Leu Asp
 580 585 590
 Lys Val Thr Asp Leu Leu Leu Phe Phe Gly Lys Leu Leu Val Val Gly
 595 600 605
 Gly Val Gly Val Leu Ser Phe Phe Phe Phe Ser Gly Arg Ile Pro Gly
 610 615 620
 Leu Gly Lys Asp Phe Lys Ser Pro His Leu Asn Tyr Tyr Trp Leu Pro
 625 630 635 640
 Ile Met Thr Ser Ile Leu Gly Ala Tyr Val Ile Ala Ser Gly Phe Phe
 645 650 655
 Ser Val Phe Gly Met Cys Val Asp Thr Leu Phe Leu Cys Phe Leu Glu
 660 665 670
 Asp Leu Glu Arg Asn Asn Gly Ser Leu Asp Arg Pro Tyr Tyr Met Ser
 675 680 685
 Lys Ser Leu Leu Lys Ile Leu Gly Lys Lys Asn Glu Ala Pro Pro Asp
 690 695 700
 Asn Lys Lys Arg Lys Lys
 705 710

<210> SEQ ID NO 99

<211> LENGTH: 598

<212> TYPE: PRT

-continued

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 99

```

Met Gly Gly Lys Gln Arg Asp Glu Asp Asp Glu Ala Tyr Gly Lys Pro
 1          5          10          15
Val Lys Tyr Asp Pro Ser Phe Arg Gly Pro Ile Lys Asn Arg Ser Cys
 20          25          30
Thr Asp Val Ile Cys Cys Val Leu Phe Leu Leu Phe Ile Leu Gly Tyr
 35          40          45
Ile Val Val Gly Ile Val Ala Trp Leu Tyr Gly Asp Pro Arg Gln Val
 50          55          60
Leu Tyr Pro Arg Asn Ser Thr Gly Ala Tyr Cys Gly Met Gly Glu Asn
 65          70          75          80
Lys Asp Lys Pro Tyr Leu Leu Tyr Phe Asn Ile Phe Ser Cys Ile Leu
 85          90          95
Ser Ser Asn Ile Ile Ser Val Ala Glu Asn Gly Leu Gln Cys Pro Thr
100          105          110
Pro Gln Val Cys Val Ser Ser Cys Pro Glu Asp Pro Trp Thr Val Gly
115          120          125
Lys Asn Glu Phe Ser Gln Thr Val Gly Glu Val Phe Tyr Thr Lys Asn
130          135          140
Arg Asn Phe Cys Leu Pro Gly Val Pro Trp Asn Met Thr Val Ile Thr
145          150          155          160
Ser Leu Gln Gln Glu Leu Cys Pro Ser Phe Leu Leu Pro Ser Ala Pro
165          170          175
Ala Leu Gly Arg Cys Phe Pro Trp Thr Asn Val Thr Pro Pro Ala Leu
180          185          190
Pro Gly Ile Thr Asn Asp Thr Thr Ile Gln Gln Gly Ile Ser Gly Leu
195          200          205
Ile Asp Ser Leu Asn Ala Arg Asp Ile Ser Val Lys Ile Phe Glu Asp
210          215          220
Phe Ala Gln Ser Trp Tyr Trp Ile Leu Val Ala Val Gly Gln Met Met
225          230          235          240
Ser Thr Met Phe Tyr Pro Leu Val Thr Phe Val Leu Leu Leu Ile Cys
245          250          255
Ile Ala Tyr Trp Ala Met Thr Ala Leu Tyr Leu Ala Thr Ser Gly Gln
260          265          270
Pro Gln Tyr Val Leu Trp Ala Ser Asn Ile Ser Ser Pro Gly Cys Glu
275          280          285
Lys Val Pro Ile Asn Thr Ser Cys Asn Pro Thr Ala His Leu Val Asn
290          295          300
Ser Ser Cys Pro Gly Leu Met Cys Val Phe Gln Gly Tyr Ser Ser Lys
305          310          315          320
Gly Leu Ile Gln Arg Ser Val Phe Asn Leu Gln Ile Tyr Gly Val Leu
325          330          335
Gly Leu Phe Trp Thr Leu Asn Trp Val Leu Ala Leu Gly Gln Cys Val
340          345          350
Leu Ala Gly Ala Phe Ala Ser Phe Tyr Trp Ala Phe His Lys Pro Gln
355          360          365
Asp Ile Pro Thr Phe Pro Leu Ile Ser Ala Phe Ile Arg Thr Leu Arg
370          375          380
Tyr His Thr Gly Ser Leu Ala Phe Gly Ala Leu Ile Leu Thr Leu Val
385          390          395          400
Gln Ile Ala Arg Val Ile Leu Glu Tyr Ile Asp His Lys Leu Arg Gly

```

-continued

405	410	415
Val Gln Asn Pro Val Ala Arg Cys Ile Met Cys Cys Phe Lys Cys Cys 420 425 430		
Leu Trp Cys Leu Glu Lys Phe Ile Lys Phe Leu Asn Arg Asn Ala Tyr 435 440 445		
Ile Met Ile Ala Ile Tyr Gly Lys Asn Phe Cys Val Ser Ala Lys Asn 450 455 460		
Ala Phe Met Leu Leu Met Arg Asn Ile Val Arg Val Val Val Leu Asp 465 470 475 480		
Lys Val Thr Asp Leu Leu Leu Phe Phe Gly Lys Leu Leu Val Val Gly 485 490 495		
Gly Val Gly Val Leu Ser Phe Phe Phe Phe Ser Gly Arg Ile Pro Gly 500 505 510		
Leu Gly Lys Asp Phe Lys Ser Pro His Leu Asn Tyr Tyr Trp Leu Pro 515 520 525		
Ile Met Thr Ser Ile Leu Gly Ala Tyr Val Ile Ala Ser Gly Phe Phe 530 535 540		
Ser Val Phe Gly Met Cys Val Asp Thr Leu Phe Leu Cys Phe Leu Glu 545 550 555 560		
Asp Leu Glu Arg Asn Asn Gly Ser Leu Asp Arg Pro Tyr Tyr Met Ser 565 570 575		
Lys Ser Leu Leu Lys Ile Leu Gly Lys Lys Asn Glu Ala Pro Pro Asp 580 585 590		
Asn Lys Lys Arg Lys Lys 595		
<210> SEQ ID NO 100		
<211> LENGTH: 2623		
<212> TYPE: DNA		
<213> ORGANISM: Homo sapiens		
<400> SEQUENCE: 100		
gagccatggg gggaaagcag cgggacgagg atgacgaggc ctacgggaag ccagtcaaat	60	
acgacccctc ctttcgaggc cccatcaaga acagaagctg cacagatgtc atctgtgtgcg	120	
tcctcttctc gctcttcatt ctagggttaca tcgtggtggg gattgtggcc tggttgtatg	180	
gagacccccg gcaagtccct taccacagga actctactgg ggcctactgt ggcattggggg	240	
agaacaaaga taagccgtat ctctgttact tcaacatctt cagctgcac cgtgccagca	300	
acatcatctc agttgctgag aacggcctac agtgccccac acccagggtg tgtgtgtcct	360	
cctgcccga ggacccatgg actgtgggaa aaaacgagtt ctcacagact gttggggaag	420	
tcctctatc aaaaaacagg aacttttgtc tgccaggggt accctggaat atgacggtga	480	
tcacaagcct gcaacaggaa ctctgcccc gtttctctct cccctctgct ccagctctgg	540	
ggcgctgctt tccatggacc aacgttactc caccggcgct ccagggtat accaatgaca	600	
ccaccatata gcaggggatc agcgggtctta ttgacagcct caatgcccga gacatcagtg	660	
ttaagatctt tgaagatctt gccagtcct ggtattggat tctgtgtgcc ctgggggtgg	720	
ctctggtctt gagectactg tttatcttgc ttctgcgcct ggtggtggg cccctggtgc	780	
tgggtgctgat cctgggagtg ctgggcgtgc tggcatacgg catctactac tgctgggagg	840	
agtaccgagt gctgcgggac aaggggcct ccatctccca gctgggtttc accaccaacc	900	
tcagtgccta ccagagcgtg caggagacct ggctggccgc cctgatcgtg ttggcggtgc	960	
ttgaagccat cctgctgctg atgctcatct tcctgcggca gcggattcgt attgccatcg	1020	

-continued

ccctcctgaa ggagggccagc aaggetgtgg gacagatgat gtctaccatg ttctaccacac	1080
tggtcacett tgtcctcctc ctcatctgca ttgcctactg ggccatgact gctctgtacc	1140
tggctacatc ggggcaaccc cagtattgtc tctgggcac ccaacatcagc tcccccggt	1200
gtgagaaagt gccaataaat acatcatgca accccacggc ccaccttggt aactcctcgt	1260
gcccagggt gatgtgcgtc ttccagggt actcatccaa aggcctaata caacgtttctg	1320
tcttcaatct gcaaatctat ggggtcctgg ggctcttctg gaccttaac tgggtactgg	1380
ccctgggcca atgcgtcctc gctggagcct ttgcctcctt ctactgggccc ttccacaagc	1440
cccaggacat ccttacette ccttaatct ctgccttcat ccgcacactc cgttaccaca	1500
ctgggtcatt ggcatcttga gccctcatcc tgaccttgt gcagatagcc cgggtcatct	1560
tggagtatat tgaccacaag ctacagaggag tgcagaaccc tgtagcccg tgcacatgt	1620
gctgtttcaa gtgtgcctc tgggtgtctgg aaaaatttat caagttccta aaccgcaatg	1680
catacatcat gatcgccatc tacgggaaga atttctgtgt ctacgcaaaa aatgcgttca	1740
tgtactcat gcgaaacatt gtcagggtgg tcgtcctgga caaagtcaca gacctgctgc	1800
tgttctttgg gaagctgctg tgggtcggag gcgtgggggt cctgtccttc tttttttct	1860
ccggtcgcac cccggggctg ggtaagact ttaagagccc ccacctcaac tattactggc	1920
tgcccatcat gaggaaccca ataaccacca cgggtcatgt ctccagacc tccatcctgg	1980
gggcctatgt catcgccagc ggcttcttca gcgttttcgg catgtgtgtg gacacgctct	2040
tcctctgctt cctggaagac ctggagcggg acaacggctc cctggaccgg ccctactaca	2100
tgtccaagag ccttctaaag attctgggca agaagaacga ggcgccccg gacaacaaga	2160
agaggaagaa gtgacagctc cggccctgat ccaggactgc accccacccc caccgtccag	2220
ccatccaacc tcacttcgcc ttacaggtct ccattttgtg gtaaaaaaag gttttaggcc	2280
aggcgccgtg gctcacgctt gtaatccaac actttgagag gctgaggcgg gcggatcacc	2340
tgagtcagga gttcgagacc agcctggcca acatggtgaa acctcctct ctattaaaaa	2400
tacaaaaatt agccgagagt ggtggcatgc acctgtcatc ccagctactc gggaggctga	2460
ggcaggagaa tcgcttgaa cccggaggca gaggttgacg tgagccgaga tcgcgccact	2520
gcactccaac ctgggtgaca gactctgtct ccaaaacaaa acaaaacaaac aaaaagattt	2580
tattaaagat attttgttaa ctacgtaaaa aaaaaaaaaa aaa	2623

<210> SEQ ID NO 101

<211> LENGTH: 2587

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 101

gagccatggg gggaaagcag cgggacgagg atgacgaggc ctacgggaag ccagtcaaat	60
acgacccctc ctttcgaggc cccatcaaga acagaagctg cacagatgct atctgctgag	120
tcctcttctc gctcttcatt ctagggtaca tcgtggtggg gattgtggcc tggttgtatg	180
gagacccccg gcaagtcctc taccccagga actctactgg ggctactgt ggcattgggg	240
agaacaaaga taagccgtat ctctgtact tcaacatctt cagctgcac ctgtccagca	300
acatcatctc agttgtgag aacggcctac agtgccccac acccagggtg tgtgtgtcct	360
cctgcccgga ggacccatgg actgtgggaa aaaacgagtt ctacagact gttggggaag	420
tctctctata aaaaaacagg aacttttctg tgccagggtg acctggaat atgacggtga	480
tcacaagcct gcaacaggaa ctctgcccc gtttctctct cccctctgct ccagctctgtg	540

-continued

ggcgctgctt tccatggacc aacgttactc caccggcgct cccagggatc accaatgaca	600
ccaccatata gcaggggatac agcgggtctta ttgacagcct caatgcccga gacatcagtg	660
ttaagatctt tgaagatttt gccagtcctt ggtattggat tcttggtgcc ctgggggttg	720
ctctggctct gaggcctactg tttatcttgc ttctgcgcct ggtggctggg cccctggctg	780
tgggtgctgat cctgggagtg ctgggcgtgc tggcatacgg catctactac tgctgggagg	840
agtaccgagt gctgcgggac aagggcgctt ccatctccca gctgggtttc accaccaacc	900
tcagtgccta ccagagcgtg caggagacct ggctggcgcg cctgatcgtg ttggcggtgc	960
ttgaagccat cctgctgctg atgctcatct tcttgcggca gcggattcgt attgccatcg	1020
ccctcctgaa ggaggccagc aaggctgtgg gacagatgat gtctaccatg ttctaccac	1080
tggtcacctt tgtcctctc ctcatctgca ttgcctactg ggccatgact gctctgtacc	1140
tgggtacatc ggggcaaccc cagtatgtgc tctgggcata caacatcagc tccccggct	1200
gtgagaaagt gccaaataat acatcatgca accccacggc ccacctgtg aactcctcgt	1260
gccagggtct gatgtgcgtc ttccagggtt actcatccaa aggcctaata caacgttctg	1320
tcttcaatct gcaaatctat ggggtcctgg ggctcttctg gaccttaac tgggtactgg	1380
ccctgggcca atgcgtctc gctggagcct ttgcctcctt ctactgggca ttccacaagc	1440
cccaggacat ccttaccttc cccttaatct ctgccttcat cgcacactc cgttaccaca	1500
ctgggtcatt ggcatttggg gccctcatcc tgaccttgt gcagatagcc cgggtcatct	1560
tggagtatat tgaccacaag ctcagaggag tgcagaaccc tgtagccgcg tgcacatgt	1620
gctgtttcaa gtgctgcctc tgggtgtctg aaaaatttat caagttccta aaccgcaatg	1680
catacatcat gatcgccatc tacgggaaga atttctgtgt ctcagccaaa aatgcgttca	1740
tgctactcat gcgaaacatt gtcagggtgg tctcctgga caaagtcaca gacctgctgc	1800
tggtcttttg gaagctgctg gtggtcggag gcgtgggggt cctgtccttc tttttttct	1860
ccggtcgcct cccggggctg ggtaaaagact ttaagagccc ccacctcaac tattactggc	1920
tgcccatcat gacctccatc ctgggggcct atgtcatcgc cagcggttc ttcagcgttt	1980
tcggcatgtg tgtggacacg ctcttctctt gtttctgga agacctggag cggaacaacg	2040
gctccctgga ccggccctac tacatgtoca agagccttct aaagattctg ggcaagaaga	2100
acgaggcgcc cccggacaac aagaagagga agaagtgaca gctccggccc tgatccagga	2160
ctgcacccca cccccaccgt ccagccatcc aacctcaatt cgccttacag gtctccattt	2220
tgtggtaaaa aaaggtttta ggcaggcgc cgtggctcac gctgtaate caacactttg	2280
agaggctgag gcggggcgat cacctgagtc aggagttcga gaccagcctg gccaacatgg	2340
tgaacctcc gtctctatta aaaatacaaa aattagccga gagggtggc atgcacctgt	2400
catcccagct actcgggagg ctgaggcagg agaatcgctt gaacccggga ggcagaggtt	2460
gcagttagcc gagatcgcc cactgcactc caacctgggt gacagactct gtctccaaaa	2520
caaaacaaac aaacaaaaag attttattaa agatattttg ttaactcagt aaaaaaaaaa	2580
aaaaaaa	2587

<210> SEQ ID NO 102
 <211> LENGTH: 2623
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens
 <400> SEQUENCE: 102

gagccatggg gggaaagcag cgggacgagg atgacgagc ctacgggaag ccagtcaaat	60

-continued

acgacccctc ctttcgaggc cccatcaaga acagaagctg cacagatgtc atctgtctgcg	120
tcctcttctc gctcttcatt ctaggttaca tcgtggtggg gattgtggcc tggttgtatg	180
gagacccccg gcaagtcttc taccocagga actctactgg ggctactgt ggcacggggg	240
agaacaaaga taagccgtat ctctgtact tcaacatctt cagctgcac cgtccagca	300
acatcatctc agttgctgag aacggcctac agtgccccc accccagggtg tgtgtgtctc	360
cctgcccgga ggacccatgg actgtgggaa aaaacgagtt ctacacagact gttggggaag	420
tcttctatac aaaaaacagg aacttttgtc tgccaggggt accctggaat atgacgggtga	480
tcacaagcct gcaacaggaa ctctgcccc gtttctctc cccctctgct ccagctctgg	540
ggcgctgctt tccatggacc aacgttactc caccggcgct ccagggtac accaatgaca	600
ccaccataca gcaggggatc agcgggtctta ttgacagcct caatgcccga gacatcagtg	660
ttaagatctt tgaagatctt gccagtcct ggtattggat tctgtgtgcc ctgggggtgg	720
ctctggtctt gacgctactg tttatcttgc ttctgcgctt ggtggctggg cccctggtgc	780
tgggtctgat cctgggagtg ctgggcgtgc tggcatacgg catctactac tgcctggagg	840
agtaccgagt gctgcgggac aagggcgcct ccatctccca gctgggttcc accaccaacc	900
tcagtgccta ccagagcgtg caggagacct ggctggccgc cctgatcgtg ttggcgggtc	960
ttgaagccat cctgctgctg atgctcatct tcctgcggca gcggatcgt attgccatcg	1020
ccctcctgaa ggaggccagc aaggctgtgg gacagatgat gtctaccatg ttctaccac	1080
tggtcacett tgcctcctc ctcatctgca ttgcctactg ggccatgact gctctgtacc	1140
tggctacatc ggggcaaccc cagtatgtgc tctgggcac caacatcagc tccccgggt	1200
gtgagaaagt gccataaat acatcatgca accccacggc ccaccttggt aactcctcgt	1260
gccagggtct gatgtgcgtc ttccagggt actcatccaa aggcctaacc caacgttctg	1320
tcttcaatct gcaaatctat ggggtcctgg ggctcttctg gaccctaac tgggtactgg	1380
ccctgggcca atgcgtcctc gctggagcct ttgcctcct ctactgggc ttccacaagc	1440
cccaggacat ccctaccttc cccttaatct ctgccttcat ccgcacactc cgttaccaca	1500
ctgggtcatt ggcatttggg gccctcatcc tgaccttgt gcagatagcc cgggtcatct	1560
tggagtatat tgaccacaag ctacagaggag tgcagaaccc tgtagccgc tgcacatgt	1620
gctgtttcaa gtgctgcctc tgggtgtctgg aaaaatttat caagttccta aaccgcaatg	1680
catacatcat gatcgccatc tacgggaaga atttctgtgt ctacgcaaaa aatgcgttca	1740
tgtactcat gcgaacatt gtcagggtgg tcgtcctgga caaagtcaca gacctgctgc	1800
tgttcttttg gaagctgctg gtggtcggag gcgtgggggt cctgtccttc tttttttct	1860
ccggtgcgat cccgggggtg ggtaagact ttaagagccc ccacctcaac tattactggc	1920
tgcccatcat gaggaaccca ataaccacca cgggtcatgt ctccagacc tccatcctgg	1980
gggcctatgt catcgccagc ggcttcttca gcgttttctg catgtgtgtg gacacgtct	2040
tcctctgctt cctggaagac ctggagcgga acaacggctc cctggaccgg ccctactaca	2100
tgtccaagag ctttctaaag attctgggca agaagaacga ggcgcccccg gacaacaaga	2160
agaggaagaa gtgacagctc cggccctgat ccaggactgc accccacccc caccgtcag	2220
ccatccaacc tcacttcgct ttacaggtct ccattttgtg gtaaaaaaag gttttaggcc	2280
aggcgccgtg gctcacgctt gtaatccaac actttgagag gctgaggcgg gcggatcacc	2340
tgagtcagga gtctgagacc agcctggcca acatggtgaa acctcgtct ctattaaaaa	2400
tacaaaaatt agccgagagt ggtggcatgc acctgtcatc ccagctactc gggaggctga	2460

-continued

```

ggcaggagaa tcgcttgaac ccgggaggca gaggttgacg tgagccgaga tcgcgccact 2520
gcactccaac ctgggtgaca gactctgtct ccaaaacaaa acaaacaaac aaaagattt 2580
tattaaagat attttgtaa ctcagtaaaa aaaaaaaaaa aaa 2623

```

<210> SEQ ID NO 103

<211> LENGTH: 722

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 103

```

Met Gly Gly Lys Gln Arg Asp Glu Asp Asp Glu Ala Tyr Gly Lys Pro
 1           5           10           15

Val Lys Tyr Asp Pro Ser Phe Arg Gly Pro Ile Lys Asn Arg Ser Cys
 20           25           30

Thr Asp Val Ile Cys Cys Val Leu Phe Leu Leu Phe Ile Leu Gly Tyr
 35           40           45

Ile Val Val Gly Ile Val Ala Trp Leu Tyr Gly Asp Pro Arg Gln Val
 50           55           60

Leu Tyr Pro Arg Asn Ser Thr Gly Ala Tyr Cys Gly Met Gly Glu Asn
 65           70           75           80

Lys Asp Lys Pro Tyr Leu Leu Tyr Phe Asn Ile Phe Ser Cys Ile Leu
 85           90           95

Ser Ser Asn Ile Ile Ser Val Ala Glu Asn Gly Leu Gln Cys Pro Thr
100           105           110

Pro Gln Val Cys Val Ser Ser Cys Pro Glu Asp Pro Trp Thr Val Gly
115           120           125

Lys Asn Glu Phe Ser Gln Thr Val Gly Glu Val Phe Tyr Thr Lys Asn
130           135           140

Arg Asn Phe Cys Leu Pro Gly Val Pro Trp Asn Met Thr Val Ile Thr
145           150           155           160

Ser Leu Gln Gln Glu Leu Cys Pro Ser Phe Leu Leu Pro Ser Ala Pro
165           170           175

Ala Leu Gly Arg Cys Phe Pro Trp Thr Asn Val Thr Pro Pro Ala Leu
180           185           190

Pro Gly Ile Thr Asn Asp Thr Thr Ile Gln Gln Gly Ile Ser Gly Leu
195           200           205

Ile Asp Ser Leu Asn Ala Arg Asp Ile Ser Val Lys Ile Phe Glu Asp
210           215           220

Phe Ala Gln Ser Trp Tyr Trp Ile Leu Val Ala Leu Gly Val Ala Leu
225           230           235           240

Val Leu Ser Leu Leu Phe Ile Leu Leu Leu Arg Leu Val Ala Gly Pro
245           250           255

Leu Val Leu Val Leu Ile Leu Gly Val Leu Gly Val Leu Ala Tyr Gly
260           265           270

Ile Tyr Tyr Cys Trp Glu Glu Tyr Arg Val Leu Arg Asp Lys Gly Ala
275           280           285

Ser Ile Ser Gln Leu Gly Phe Thr Thr Asn Leu Ser Ala Tyr Gln Ser
290           295           300

Val Gln Glu Thr Trp Leu Ala Ala Leu Ile Val Leu Ala Val Leu Glu
305           310           315           320

Ala Ile Leu Leu Leu Met Leu Ile Phe Leu Arg Gln Arg Ile Arg Ile
325           330           335

Ala Ile Ala Leu Leu Lys Glu Ala Ser Lys Ala Val Gly Gln Met Met
340           345           350

```

Ser	Thr	Met	Phe	Tyr	Pro	Leu	Val	Thr	Phe	Val	Leu	Leu	Leu	Ile	Cys
355						360			365						
Ile	Ala	Tyr	Trp	Ala	Met	Thr	Ala	Leu	Tyr	Leu	Ala	Thr	Ser	Gly	Gln
370						375			380						
Pro	Gln	Tyr	Val	Leu	Trp	Ala	Ser	Asn	Ile	Ser	Ser	Pro	Gly	Cys	Glu
385			390						395			400			
Lys	Val	Pro	Ile	Asn	Thr	Ser	Cys	Asn	Pro	Thr	Ala	His	Leu	Val	Asn
			405						410			415			
Ser	Ser	Cys	Pro	Gly	Leu	Met	Cys	Val	Phe	Gln	Gly	Tyr	Ser	Ser	Lys
			420			425						430			
Gly	Leu	Ile	Gln	Arg	Ser	Val	Phe	Asn	Leu	Gln	Ile	Tyr	Gly	Val	Leu
435						440						445			
Gly	Leu	Phe	Trp	Thr	Leu	Asn	Trp	Val	Leu	Ala	Leu	Gly	Gln	Cys	Val
450						455			460						
Leu	Ala	Gly	Ala	Phe	Ala	Ser	Phe	Tyr	Trp	Ala	Phe	His	Lys	Pro	Gln
465			470						475			480			
Asp	Ile	Pro	Thr	Phe	Pro	Leu	Ile	Ser	Ala	Phe	Ile	Arg	Thr	Leu	Arg
			485						490			495			
Tyr	His	Thr	Gly	Ser	Leu	Ala	Phe	Gly	Ala	Leu	Ile	Leu	Thr	Leu	Val
			500			505						510			
Gln	Ile	Ala	Arg	Val	Ile	Leu	Glu	Tyr	Ile	Asp	His	Lys	Leu	Arg	Gly
515						520						525			
Val	Gln	Asn	Pro	Val	Ala	Arg	Cys	Ile	Met	Cys	Cys	Phe	Lys	Cys	Cys
530						535			540						
Leu	Trp	Cys	Leu	Glu	Lys	Phe	Ile	Lys	Phe	Leu	Asn	Arg	Asn	Ala	Tyr
545			550			555						560			
Ile	Met	Ile	Ala	Ile	Tyr	Gly	Lys	Asn	Phe	Cys	Val	Ser	Ala	Lys	Asn
			565						570			575			
Ala	Phe	Met	Leu	Leu	Met	Arg	Asn	Ile	Val	Arg	Val	Val	Val	Leu	Asp
580						585						590			
Lys	Val	Thr	Asp	Leu	Leu	Leu	Phe	Phe	Gly	Lys	Leu	Leu	Val	Val	Gly
595						600						605			
Gly	Val	Gly	Val	Leu	Ser	Phe	Phe	Phe	Phe	Ser	Gly	Arg	Ile	Pro	Gly
610			615						620						
Leu	Gly	Lys	Asp	Phe	Lys	Ser	Pro	His	Leu	Asn	Tyr	Tyr	Trp	Leu	Pro
625			630			635						640			
Ile	Met	Arg	Asn	Pro	Ile	Thr	Pro	Thr	Gly	His	Val	Phe	Gln	Thr	Ser
			645			650						655			
Ile	Leu	Gly	Ala	Tyr	Val	Ile	Ala	Ser	Gly	Phe	Phe	Ser	Val	Phe	Gly
			660			665						670			
Met	Cys	Val	Asp	Thr	Leu	Phe	Leu	Cys	Phe	Leu	Glu	Asp	Leu	Glu	Arg
675			680			685						690			
Asn	Asn	Gly	Ser	Leu	Asp	Arg	Pro	Tyr	Tyr	Met	Ser	Lys	Ser	Leu	Leu
690			695						700						
Lys	Ile	Leu	Gly	Lys	Lys	Asn	Glu	Ala	Pro	Pro	Asp	Asn	Lys	Lys	Arg
705			710			715						720			
Lys			Lys												

Met Gly Gly Lys Gln Arg Asp Glu Asp Asp Glu Ala Tyr Gly Lys Pro

-continued

1	5	10	15
Val Lys Tyr Asp	Pro Ser Phe Arg	Gly Pro Ile Lys	Asn Arg Ser Cys
20		25	30
Thr Asp Val Ile	Cys Cys Val Leu	Phe Leu Leu Phe	Ile Leu Gly Tyr
35		40	45
Ile Val Val Gly	Ile Val Ala Trp	Leu Tyr Gly Asp	Pro Arg Gln Val
50		55	60
Leu Tyr Pro Arg	Asn Ser Thr Gly	Ala Tyr Cys Gly	Met Gly Glu Asn
65		70	75
Lys Asp Lys Pro	Tyr Leu Leu Tyr	Phe Asn Ile Phe	Ser Cys Ile Leu
	85	90	95
Ser Ser Asn Ile	Ile Ser Val Ala	Glu Asn Gly Leu	Gln Cys Pro Thr
	100	105	110
Pro Gln Val Cys	Val Ser Ser Cys	Pro Glu Asp Pro	Trp Thr Val Gly
	115	120	125
Lys Asn Glu Phe	Ser Gln Thr Val	Gly Glu Val Phe	Tyr Thr Lys Asn
	130	135	140
Arg Asn Phe Cys	Leu Pro Gly Val	Pro Trp Asn Met	Thr Val Ile Thr
	145	150	155
Ser Leu Gln Gln	Glu Leu Cys Pro	Ser Phe Leu Leu	Pro Ser Ala Pro
	165	170	175
Ala Leu Gly Arg	Cys Phe Pro Trp	Thr Asn Val Thr	Pro Pro Ala Leu
	180	185	190
Pro Gly Ile Thr	Asn Asp Thr Thr	Ile Gln Gln Gly	Ile Ser Gly Leu
	195	200	205
Ile Asp Ser Leu	Asn Ala Arg Asp	Ile Ser Val Lys	Ile Phe Glu Asp
	210	215	220
Phe Ala Gln Ser	Trp Tyr Trp Ile	Leu Val Ala Leu	Gly Val Ala Leu
	225	230	235
Val Leu Ser Leu	Leu Phe Ile Leu	Leu Leu Arg Leu	Val Ala Gly Pro
	245	250	255
Leu Val Leu Val	Leu Ile Leu Gly	Val Leu Gly Val	Leu Ala Tyr Gly
	260	265	270
Ile Tyr Tyr Cys	Trp Glu Glu Tyr	Arg Val Leu Arg	Asp Lys Gly Ala
	275	280	285
Ser Ile Ser Gln	Leu Gly Phe Thr	Thr Asn Leu Ser	Ala Tyr Gln Ser
	290	295	300
Val Gln Glu Thr	Trp Leu Ala Ala	Leu Ile Val Leu	Ala Val Leu Glu
	305	310	315
Ala Ile Leu Leu	Leu Met Leu Ile	Phe Leu Arg Gln	Arg Ile Arg Ile
	325	330	335
Ala Ile Ala Leu	Leu Lys Glu Ala	Ser Lys Ala Val	Gly Gln Met Met
	340	345	350
Ser Thr Met Phe	Tyr Pro Leu Val	Thr Phe Val Leu	Leu Leu Ile Cys
	355	360	365
Ile Ala Tyr Trp	Ala Met Thr Ala	Leu Tyr Leu Ala	Thr Ser Gly Gln
	370	375	380
Pro Gln Tyr Val	Leu Trp Ala Ser	Asn Ile Ser Ser	Pro Gly Cys Glu
	385	390	395
Lys Val Pro Ile	Asn Thr Ser Cys	Asn Pro Thr Ala	His Leu Val Asn
	405	410	415
Ser Ser Cys Pro	Gly Leu Met Cys	Val Phe Gln Gly	Tyr Ser Ser Lys
	420	425	430

-continued

Gly Leu Ile Gln Arg Ser Val Phe Asn Leu Gln Ile Tyr Gly Val Leu
 435 440 445
 Gly Leu Phe Trp Thr Leu Asn Trp Val Leu Ala Leu Gly Gln Cys Val
 450 455 460
 Leu Ala Gly Ala Phe Ala Ser Phe Tyr Trp Ala Phe His Lys Pro Gln
 465 470 475 480
 Asp Ile Pro Thr Phe Pro Leu Ile Ser Ala Phe Ile Arg Thr Leu Arg
 485 490 495
 Tyr His Thr Gly Ser Leu Ala Phe Gly Ala Leu Ile Leu Thr Leu Val
 500 505 510
 Gln Ile Ala Arg Val Ile Leu Glu Tyr Ile Asp His Lys Leu Arg Gly
 515 520 525
 Val Gln Asn Pro Val Ala Arg Cys Ile Met Cys Cys Phe Lys Cys Cys
 530 535 540
 Leu Trp Cys Leu Glu Lys Phe Ile Lys Phe Leu Asn Arg Asn Ala Tyr
 545 550 555 560
 Ile Met Ile Ala Ile Tyr Gly Lys Asn Phe Cys Val Ser Ala Lys Asn
 565 570 575
 Ala Phe Met Leu Leu Met Arg Asn Ile Val Arg Val Val Val Leu Asp
 580 585 590
 Lys Val Thr Asp Leu Leu Leu Phe Phe Gly Lys Leu Leu Val Val Gly
 595 600 605
 Gly Val Gly Val Leu Ser Phe Phe Phe Phe Ser Gly Arg Ile Pro Gly
 610 615 620
 Leu Gly Lys Asp Phe Lys Ser Pro His Leu Asn Tyr Tyr Trp Leu Pro
 625 630 635 640
 Ile Met Thr Ser Ile Leu Gly Ala Tyr Val Ile Ala Ser Gly Phe Phe
 645 650 655
 Ser Val Phe Gly Met Cys Val Asp Thr Leu Phe Leu Cys Phe Leu Glu
 660 665 670
 Asp Leu Glu Arg Asn Asn Gly Ser Leu Asp Arg Pro Tyr Tyr Met Ser
 675 680 685
 Lys Ser Leu Leu Lys Ile Leu Gly Lys Lys Asn Glu Ala Pro Pro Asp
 690 695 700
 Asn Lys Lys Arg Lys Lys
 705 710

<210> SEQ ID NO 105

<211> LENGTH: 722


<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 105

Met Gly Gly Lys Gln Arg Asp Glu Asp Asp Glu Ala Tyr Gly Lys Pro
 1 5 10 15
 Val Lys Tyr Asp Pro Ser Phe Arg Gly Pro Ile Lys Asn Arg Ser Cys
 20 25 30
 Thr Asp Val Ile Cys Cys Val Leu Phe Leu Leu Phe Ile Leu Gly Tyr
 35 40 45
 Ile Val Val Gly Ile Val Ala Trp Leu Tyr Gly Asp Pro Arg Gln Val
 50 55 60
 Leu Tyr Pro Arg Asn Ser Thr Gly Ala Tyr Cys Gly Met Gly Glu Asn
 65 70 75 80
 Lys Asp Lys Pro Tyr Leu Leu Tyr Phe Asn Ile Phe Ser Cys Ile Leu
 85 90 95

Figure 6: 24P4C12 variant 1 Hydropathicity Profile
(Kyte J., Doolittle R.F., 1982. J. Mol. Biol. 157:105-132)



-continued

Ser	Ser	Asn	Ile	Ile	Ser	Val	Ala	Glu	Asn	Gly	Leu	Gln	Cys	Pro	Thr
		100						105					110		
Pro	Gln	Val	Cys	Val	Ser	Ser	Cys	Pro	Glu	Asp	Pro	Trp	Thr	Val	Gly
	115						120					125			
Lys	Asn	Glu	Phe	Ser	Gln	Thr	Val	Gly	Glu	Val	Phe	Tyr	Thr	Lys	Asn
	130					135					140				
Arg	Asn	Phe	Cys	Leu	Pro	Gly	Val	Pro	Trp	Asn	Met	Thr	Val	Ile	Thr
145					150					155					160
Ser	Leu	Gln	Gln	Glu	Leu	Cys	Pro	Ser	Phe	Leu	Leu	Pro	Ser	Ala	Pro
				165						170				175	
Ala	Leu	Gly	Arg	Cys	Phe	Pro	Trp	Thr	Asn	Val	Thr	Pro	Pro	Ala	Leu
			180					185					190		
Pro	Gly	Ile	Thr	Asn	Asp	Thr	Thr	Ile	Gln	Gln	Gly	Ile	Ser	Gly	Leu
	195					200						205			
Ile	Asp	Ser	Leu	Asn	Ala	Arg	Asp	Ile	Ser	Val	Lys	Ile	Phe	Glu	Asp
	210					215					220				
Phe	Ala	Gln	Ser	Trp	Tyr	Trp	Ile	Leu	Val	Ala	Leu	Gly	Val	Ala	Leu
225					230					235					240
Val	Leu	Ser	Leu	Leu	Phe	Ile	Leu	Leu	Leu	Arg	Leu	Val	Ala	Gly	Pro
			245						250					255	
Leu	Val	Leu	Val	Leu	Ile	Leu	Gly	Val	Leu	Gly	Val	Leu	Ala	Tyr	Gly
			260					265					270		
Ile	Tyr	Tyr	Cys	Trp	Glu	Glu	Tyr	Arg	Val	Leu	Arg	Asp	Lys	Gly	Ala
	275						280					285			
Ser	Ile	Ser	Gln	Leu	Gly	Phe	Thr	Thr	Asn	Leu	Ser	Ala	Tyr	Gln	Ser
	290					295					300				
Val	Gln	Glu	Thr	Trp	Leu	Ala	Ala	Leu	Ile	Val	Leu	Ala	Val	Leu	Glu
305					310					315					320
Ala	Ile	Leu	Leu	Leu	Met	Leu	Ile	Phe	Leu	Arg	Gln	Arg	Ile	Arg	Ile
			325						330					335	
Ala	Ile	Ala	Leu	Leu	Lys	Glu	Ala	Ser	Lys	Ala	Val	Gly	Gln	Met	Met
			340					345					350		
Ser	Thr	Met	Phe	Tyr	Pro	Leu	Val	Thr	Phe	Val	Leu	Leu	Leu	Ile	Cys
		355					360					365			
Ile	Ala	Tyr	Trp	Ala	Met	Thr	Ala	Leu	Tyr	Leu	Ala	Thr	Ser	Gly	Gln
	370					375					380				
Pro	Gln	Tyr	Val	Leu	Trp	Ala	Ser	Asn	Ile	Ser	Ser	Pro	Gly	Cys	Glu
385					390					395					400
Lys	Val	Pro	Ile	Asn	Thr	Ser	Cys	Asn	Pro	Thr	Ala	His	Leu	Val	Asn
			405						410					415	
Ser	Ser	Cys	Pro	Gly	Leu	Met	Cys	Val	Phe	Gln	Gly	Tyr	Ser	Ser	Lys
			420					425					430		
Gly	Leu	Ile	Gln	Arg	Ser	Val	Phe	Asn	Leu	Gln	Ile	Tyr	Gly	Val	Leu
			435				440					445			
Gly	Leu	Phe	Trp	Thr	Leu	Asn	Trp	Val	Leu	Ala	Leu	Gly	Gln	Cys	Val
	450					455						460			
Leu	Ala	Gly	Ala	Phe	Ala	Ser	Phe	Tyr	Trp	Ala	Phe	His	Lys	Pro	Gln
465					470					475					480
Asp	Ile	Pro	Thr	Phe	Pro	Leu	Ile	Ser	Ala	Phe	Ile	Arg	Thr	Leu	Arg
			485						490					495	
Tyr	His	Thr	Gly	Ser	Leu	Ala	Phe	Gly	Ala	Leu	Ile	Leu	Thr	Leu	Val
			500					505					510		
Gln	Ile	Ala	Arg	Val	Ile	Leu	Glu	Tyr	Ile	Asp	His	Lys	Leu	Arg	Gly
			515				520					525			

-continued

Val Gln Asn Pro Val Ala Arg Cys Ile Met Cys Cys Phe Lys Cys Cys
530 535 540

Leu Trp Cys Leu Glu Lys Phe Ile Lys Phe Leu Asn Arg Asn Ala Tyr
545 550 555 560

Ile Met Ile Ala Ile Tyr Gly Lys Asn Phe Cys Val Ser Ala Lys Asn
565 570 575

Ala Phe Met Leu Leu Met Arg Asn Ile Val Arg Val Val Val Leu Asp
580 585 590

Lys Val Thr Asp Leu Leu Leu Phe Phe Gly Lys Leu Leu Val Val Gly
595 600 605

Gly Val Gly Val Leu Ser Phe Phe Phe Phe Ser Gly Arg Ile Pro Gly
610 615 620

Leu Gly Lys Asp Phe Lys Ser Pro His Leu Asn Tyr Tyr Trp Leu Pro
625 630 635 640

Ile Met Arg Asn Pro Ile Thr Pro Thr Gly His Val Phe Gln Thr Ser
645 650 655

Ile Leu Gly Ala Tyr Val Ile Ala Ser Gly Phe Phe Ser Val Phe Gly
660 665 670

Met Cys Val Asp Thr Leu Phe Leu Cys Phe Leu Glu Asp Leu Glu Arg
675 680 685

Asn Asn Gly Ser Leu Asp Arg Pro Tyr Tyr Met Ser Lys Ser Leu Leu
690 695 700

Lys Ile Leu Gly Lys Lys Asn Glu Ala Pro Pro Asp Asn Lys Lys Arg
705 710 715 720

Lys Lys

<210> SEQ ID NO 106

<211> LENGTH: 2593

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 106

gagccatggg gggaaagcag cgggacgagg atgacgaggc ctacgggaag ccagtcaaat 60

acgacccctc ctttcgaggc cccatcaaga acagaagctg cacagatgtc atctgtgtcg 120

tcctcttctc gctcttcatt ctagggtaca tcgtggtggg gattgtggcc tggttgtatg 180

gagacccccg gcaagtcttc taccacagga actctactgg ggctactgt ggcattgggg 240

agaacaaaga taagccgtat ctctgtact tcaacatctt cagctgcac ctgtccagca 300

acatcatctc agttgtgag aacggcctac agtgccccac acccagggtg tgtgtgtcct 360

cctgcccgga ggacccatgg actgtgggaa aaaacgagtt ctacagact gttggggaag 420

tcttctatac aaaaaacagg aacttttgtc tgccaggggt accctggaat atgacgggtga 480

tcacaagcct gcaacaggaa ctctgcccc gtttctctc cccctctgct ccagctctgg 540

ggcgctgctt tccatggacc aacgttactc caccggcgct ccagggatc accaatgaca 600

ccaccataca gcaggggatc agcgggtctta ttgacagcct caatgcccga gacatcagtg 660

ttaagatctt tgaagatctt gccagctct ggtattggat tcttgttgcc ctgggggtgg 720

ctctggtctt gacgctactg tttatcttgc ttctgcgct ggtggctggg ccctgggtgc 780

tggtgctgat cctgggagtg ctgggcgtgc tggcatacgg catctactac tgctgggagg 840

agtaccgagt gctgcgggac aagggcgct ccatctccca gctgggttcc accaccaacc 900

tcagtgccta ccagagcgtg caggagacct ggctggccgc cctgatcgtg ttggcggtgc 960

ttgaagccat cctgctgctg atgctcatct tcctgcggca gcggattcgt attgcatcgt 1020

-continued

```

cctcctgaa ggaggccagc aaggctgtgg gacagatgat gtctaccatg ttctacccac 1080
tggtcacctt tgcctcctc ctcatctgca ttgcctactg ggccatgact gctctgtatc 1140
ctctgcccac gcagccagcc actcttggat atgtgctctg ggcatccaac atcagctccc 1200
ccggtgtga gaaagtgcc ataaatacat catgcaaccc cacggcccac cttgtgaact 1260
cctcgtgcc agggctgatg tgcgtcttcc agggctactc atccaaaggc ctaatccaac 1320
gttctgtctt caatctgcaa atctatgggg tcttggggct cttctggacc cttactggg 1380
tactggccct ggccaatgc gtctcgtcg gagcctttgc ctcttctac tgggccttcc 1440
acaagcccca ggacatccct accttccct taatctctgc cttcatccgc aactecgtt 1500
accacactgg gtcattggca tttggagccc tcctcctgac ccttgtcag atagcccggy 1560
tcactctgga gtatattgac cacaagctca gaggagtga gaacctgta gcccgctgca 1620
tcagtgtctg tttcaagtgc tgcctctggt gtctggaaaa atttatcaag ttctaaacc 1680
gcaatgcata catcatgac gccatctacg ggaagaattt ctgtgtctca gcaaaaaatg 1740
cgttcctgct actcatgca aacattgtca ggggtgctgt cctggacaaa gtcacagacc 1800
tgctgctgtt ctttgggaag ctgctggagg tggaggcgt gggggtcctg tccttctttt 1860
ttttctcgg tcgcatccc gggtctggga aagaacttaa gagccccac ctcaactatt 1920
actggctgcc catcatgacc tccatcctgg ggcctatgt catcgccagc ggcttcttca 1980
gcgttttcgg catgtgtgtg gacacgctct tctctgctt cctggaagac ctggagcgga 2040
acaacggctc cctggaccgg ccctactaca tgccaagag ccttctaaag attctgggca 2100
agaagaacga ggcgccccg gacaacaaga agaggaagaa gtgacagctc cggccctgat 2160
ccaggactgc accccacccc caccgtccag ccatccaacc tcacttcgcc ttacaggctc 2220
ccattttgtg gtaaaaaaag gttttaggcc aggcgcctg gctcacgct gtaatccaac 2280
actttgagag gctgaggcgg cgggatcacc tgagtcagga gttcgagacc agcctggcca 2340
acatggtgaa acctccgtct ctattaaaa taaaaaatt agccgagagt ggtggcatgc 2400
acctgtcatc ccagctactc gggaggctga ggcaggagaa tcgcttgaac ccgggaggca 2460
gaggttgcat tgagccgaga tcgcgcact gactccaac ctgggtgaca gactctgtct 2520
ccaaaacaaa acaaacaaac aaaaagattt tattaagat attttgttaa ctacagtaaaa 2580
aaaaaaaaaa aaa 2593

```

<210> SEQ ID NO 107

<211> LENGTH: 2587

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 107

```

gagccatggg gggaaagcag cgggacgagg atgacgagc ctacgggaag ccagtcaaat 60
acgacccctc ctttcgaggc cccatcaaga acagaagctg cacagatgct atctgtctgc 120
tcctcttctc gctcttcatt ctagggtaca tcgtggtggg gattgtggcc tggttgtatg 180
gagacccccg gcaagtcctc taccacagga actctactgg ggcctactgt ggcatggggg 240
agaacaaaga taagccgtat ctctgtact tcaacatctt cagctgcac ctgtccagca 300
acatcatctc agttgtgag aacggcctac agtgcaccac acccagggtg tgtgtgtcct 360
cctgcccgga ggacccatgg actgtgggaa aaaacgagtt ctacagact gttggggaag 420
tcttctatc aaaaaacagg aacttttgtc tgccaggggt accctggaat atgacggtga 480
tcacaagcct gcaacaggaa ctctgcccc gtttctctc cccctctgct ccagctctgg 540

```


-continued

```

ggcgctgctt tccatggacc aacgttactc caccggcgct cccagggatc accaatgaca 600
ccaccatata gcaggggata agcgggtctta ttgacagcct caatgcccga gacatcagtg 660
ttaagatctt tgaagatctt gccagtcctt ggtattggat tcttggtgcc ctgggggtgg 720
ctctggtctt gagectactg tttatcttgc ttctgcgcct ggtggetggg cccctggtgc 780
tgggtgctgat cctgggagtg ctgggcgtgc tggcatacgg catctactac tgctgggagg 840
agtaccgagt gctgcgggac aaggggcgct ccatctccca gctgggttcc accaccaacc 900
tcagtgccta ccagagcgtg caggagacct ggcggccgc cctgacgtg ttggcgggtgc 960
ttgaagccat cctgctgctg atgctcatct tctgcgga gccgattcgt attgccatcg 1020
ccctcctgaa ggaggccagc aaggctgtgg gacagatgat gtctaccatg ttctaccac 1080
tggtcacctt tgcctcctc ctcatctgca ttgcctactg ggccatgact gctctgtacc 1140
tggctacatc ggggcaaccc cagtatgtgc tctgggcac caacatcagc tccccggct 1200
gtgagaaagt gccaaataat acatcatgca accccacggc ccacctgtg aactcctcgt 1260
gccagggtg gatgtgcgtc ttccagggt actcatccaa aggcctaacc caacgttctg 1320
tcttcaatct gcaaatctat ggggtcctgg ggcctctctg gaccttaac tgggtactgg 1380
ccctgggcca atgcgtcctc gctggagcct ttgcctcct ctactgggcc ttccacaagc 1440
cccaggacat ccttaccttc ccttaatct ctgccttcat ccgcacactc cgttaccaca 1500
ctgggtcatt ggcatttga gccctcatcc tgaccttgt gcagatagcc cgggtcatct 1560
tggagtatat tgaccacaag ctgagaggag tgcagaaccc tgtagccgc tgcacatgt 1620
gctgtttcaa gtgctgcctc tgggtgtctg aaaaatttat caagtctta aaccgcaatg 1680
catacatcat gatcgccatc tacgggaaga atttctgtgt ctgagccaaa aatgcgttca 1740
tgctactcat gcgaaacatt gtcagggtgg tctctctgga caaagtcaca gacctgctgc 1800
tgttctttgg gaagctgctg tgggtcggag gcgtgggggt cctgtccttc tttttttct 1860
ccggtcgcac cccggggctg ggtaaaagact ttaagagccc ccacctcaac tattactggc 1920
tgcccatcat gacctccatc ctgggggcct atgtcatcgc cagcggcttc ttcagcgttt 1980
tcggcatgtg tgtggacacg ctcttctct gcttctctga agacctggag cggaacaacg 2040
gctccctgga ccggccctac tacatgtcca agagccttct aaagattctg ggcaagaaga 2100
acgaggcgcc cccggacaac aagaagagga agaagtgaca gctccggccc tgatccagga 2160
ctgcaccca cccccaccgt ccagccatcc aaactcaatt cgccttacag gtctccattt 2220
tgtggtaaaa aaagggttta ggccaggcgc cgtggctcac gcctgtaac caacactttg 2280
agaggctgag gcgggaggat cactgagtc aggagtctga gaccagcctg gccaacatgg 2340
tgaaacctcc gtctctatta aaaatacaaa aattagccga gagtgggtgg atgcacctgt 2400
catcccagct actcgggagg ctgaggcagg agaatcgctt gaaccgggga ggcagagggt 2460
gcagttagcc gagatcgcc cactgcactc caacctgggt gacagactct gtctccaaaa 2520
caaaacaaac aaacaaaaag attttattaa agatatattg ttaactcagt aaaaaaaaaa 2580
aaaaaaa 2587

```

<210> SEQ ID NO 108

<211> LENGTH: 2593

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 108

```

gagccatggg gggaaagcag cgggacgagg atgacgaggc ctacgggaag ccagtcaa 60

```

-continued

acgacccctc ctttcgaggc cccatcaaga acagaagctg cacagatgtc atctgctgcg	120
tcctcttctc gctcttcatt ctagggtaca tcgtggtggg gattgtggcc tggttgtatg	180
gagacccccg gcaagtcctc taccccagga actctactgg ggctactgt ggcatggggg	240
agaacaaaga taagccgtat ctctgtact tcaacatctt cagctgcac cgttcacga	300
acatcatctc agttgtgag aacggcctac agtgcctcac acccagggtg tgtgtgtcct	360
cctgcccga ggacccatgg actgtggga aaaacgagtt ctcacagact gttggggaag	420
tcttctatac aaaaaacagg aacttttgct tgccagggt accctggaat atgacggtga	480
tcacaagcct gcaacaggaa ctctgcccc gtttctctc cccctctgct ccagctctgg	540
ggcgtgctt tccatggacc aacgttact caccggcgt cccagggtc accaatgaca	600
ccaccataca gcaggggatc agcgggtctt ttgacagcct caatgcccga gacatcagtg	660
ttaagatctt tgaagatctt gccagtcct ggtattggat tcttgttgcc ctgggggtgg	720
ctctggtctt gagcctactg tttatcttgc ttctgcgct ggtggctggg ccctggtgc	780
tgggtgctgat cctgggagtg ctgggctgct tggcatacgg catctactac tctgggagg	840
agtaccgagt gctgcccga aaggcgctc ccatctcca gctgggttc accaccaacc	900
tcagtgccta ccagagcgtg caggagacct ggctggcgc cctgatcgtg ttggcggtgc	960
ttgaagccat cctgctgctg atgctcatct tcctgcggca gggattcgt attgccatcg	1020
ccctcctgaa ggaggccagc aaggctgtgg gacagatgat gtctaccatg ttctaccac	1080
tggtcacctt tgtctctc ctcactgca ttgctactg ggccatgact gctctgtatc	1140
ctctgccac gcagccagcc actcttgat atgtgctctg ggcatccaac atcagctccc	1200
ccgctgtga gaaagtcca ataatatcat catgcaacc caccggccac cttgtgaact	1260
cctcgtgcc agggctgatg tgctcttcc agggctact atccaaagg ctaatccaac	1320
gttctgtctt caatctgaa atctatggg tcctggggt cttctggacc cttaactggg	1380
tactggcctt gggccaatgc gtctcgtg gagcctttgc ctctctctac tgggccttcc	1440
acaagccca ggacatcctt accctccct taatctctgc ctctatcgc acactccgtt	1500
accacactgg gtcattggca tttggagccc tcactctgac cctgtgacg atagccggg	1560
tcactcttga gtatattgac cacaagctca gaggagtga gaacctgta gcccgctga	1620
tcactgtctg tttcaagtgc tgctctggt gtctggaaaa atttatcaag ttctaaacc	1680
gcaatgcata catcatgat gccatctac ggaagaattt ctgtgtctca gcaaaaatg	1740
cgttcagtct actcatgca aacattgtca ggtggtcgt cctggacaaa gtcacagacc	1800
tgtcgtgctt ctttgggaag ctgctggtg tcggaggcgt gggggtcctg tccttctttt	1860
ttttctccg tcgcatccc gggctggga aagacttta gagcccccac ctcaactatt	1920
actggctgcc catcatgacc tccatcctgg gggcctatgt catcgccagc ggcttcttca	1980
gcgttttcgg catgtgtgtg gacacgtct tcctctgctt cctggaagac ctggagcgga	2040
acaacggctc cctggaccgg ccctactaca tgtccaagag ccttctaaag attctgggca	2100
agaagaacga ggcgccccg gacaacaaga agaggaagaa gtgacagctc cggccctgat	2160
ccaggactgc accccaccc caccgtccag ccatccaacc tcacttcgcc ttacaggtct	2220
ccattttgtg gtaaaaaaag gttttaggcc aggcgcctg gctcacgct gtaatccaac	2280
actttgagag gctgaggcgg cgggatcacc tgagtcagga gttcgagacc agcctggcca	2340
acatggtgaa acctccgtct ctattaaaa tacaaaaatt agccgagagt ggtggcatgc	2400
acctgtcatc ccagctactc gggaggctga ggcaggagaa tcgcttgaac ccgggagga	2460

-continued

```

gaggttgacg tgagccgaga tcgcgccact gcactccaac ctgggtgaca gactctgtct 2520
ccaaaacaaa acaaacacaa aaaaagattt tattaagat atttgttaa ctcagtaaaa 2580
aaaaaaaaaa aaa 2593

```

```

<210> SEQ ID NO 109
<211> LENGTH: 712
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 109

```

```

Met Gly Gly Lys Gln Arg Asp Glu Asp Asp Glu Ala Tyr Gly Lys Pro
1          5          10          15
Val Lys Tyr Asp Pro Ser Phe Arg Gly Pro Ile Lys Asn Arg Ser Cys
20          25          30
Thr Asp Val Ile Cys Cys Val Leu Phe Leu Leu Phe Ile Leu Gly Tyr
35          40          45
Ile Val Val Gly Ile Val Ala Trp Leu Tyr Gly Asp Pro Arg Gln Val
50          55          60
Leu Tyr Pro Arg Asn Ser Thr Gly Ala Tyr Cys Gly Met Gly Glu Asn
65          70          75          80
Lys Asp Lys Pro Tyr Leu Leu Tyr Phe Asn Ile Phe Ser Cys Ile Leu
85          90          95
Ser Ser Asn Ile Ile Ser Val Ala Glu Asn Gly Leu Gln Cys Pro Thr
100         105         110
Pro Gln Val Cys Val Ser Ser Cys Pro Glu Asp Pro Trp Thr Val Gly
115         120         125
Lys Asn Glu Phe Ser Gln Thr Val Gly Glu Val Phe Tyr Thr Lys Asn
130         135         140
Arg Asn Phe Cys Leu Pro Gly Val Pro Trp Asn Met Thr Val Ile Thr
145         150         155         160
Ser Leu Gln Gln Glu Leu Cys Pro Ser Phe Leu Leu Pro Ser Ala Pro
165         170         175
Ala Leu Gly Arg Cys Phe Pro Trp Thr Asn Val Thr Pro Pro Ala Leu
180         185         190
Pro Gly Ile Thr Asn Asp Thr Thr Ile Gln Gln Gly Ile Ser Gly Leu
195         200         205
Ile Asp Ser Leu Asn Ala Arg Asp Ile Ser Val Lys Ile Phe Glu Asp
210         215         220
Phe Ala Gln Ser Trp Tyr Trp Ile Leu Val Ala Leu Gly Val Ala Leu
225         230         235         240
Val Leu Ser Leu Leu Phe Ile Leu Leu Leu Arg Leu Val Ala Gly Pro
245         250         255
Leu Val Leu Val Leu Ile Leu Gly Val Leu Gly Val Leu Ala Tyr Gly
260         265         270
Ile Tyr Tyr Cys Trp Glu Glu Tyr Arg Val Leu Arg Asp Lys Gly Ala
275         280         285
Ser Ile Ser Gln Leu Gly Phe Thr Thr Asn Leu Ser Ala Tyr Gln Ser
290         295         300
Val Gln Glu Thr Trp Leu Ala Ala Leu Ile Val Leu Ala Val Leu Glu
305         310         315         320
Ala Ile Leu Leu Leu Met Leu Ile Phe Leu Arg Gln Arg Ile Arg Ile
325         330         335
Ala Ile Ala Leu Leu Lys Glu Ala Ser Lys Ala Val Gly Gln Met Met
340         345         350

```

-continued

Ser Thr Met Phe Tyr Pro Leu Val Thr Phe Val Leu Leu Leu Ile Cys
 355 360 365
 Ile Ala Tyr Trp Ala Met Thr Ala Leu Tyr Pro Leu Pro Thr Gln Pro
 370 375 380
 Ala Thr Leu Gly Tyr Val Leu Trp Ala Ser Asn Ile Ser Ser Pro Gly
 385 390 395 400
 Cys Glu Lys Val Pro Ile Asn Thr Ser Cys Asn Pro Thr Ala His Leu
 405 410 415
 Val Asn Ser Ser Cys Pro Gly Leu Met Cys Val Phe Gln Gly Tyr Ser
 420 425 430
 Ser Lys Gly Leu Ile Gln Arg Ser Val Phe Asn Leu Gln Ile Tyr Gly
 435 440 445
 Val Leu Gly Leu Phe Trp Thr Leu Asn Trp Val Leu Ala Leu Gly Gln
 450 455 460
 Cys Val Leu Ala Gly Ala Phe Ala Ser Phe Tyr Trp Ala Phe His Lys
 465 470 475 480
 Pro Gln Asp Ile Pro Thr Phe Pro Leu Ile Ser Ala Phe Ile Arg Thr
 485 490 495
 Leu Arg Tyr His Thr Gly Ser Leu Ala Phe Gly Ala Leu Ile Leu Thr
 500 505 510
 Leu Val Gln Ile Ala Arg Val Ile Leu Glu Tyr Ile Asp His Lys Leu
 515 520 525
 Arg Gly Val Gln Asn Pro Val Ala Arg Cys Ile Met Cys Cys Phe Lys
 530 535 540
 Cys Cys Leu Trp Cys Leu Glu Lys Phe Ile Lys Phe Leu Asn Arg Asn
 545 550 555 560
 Ala Tyr Ile Met Ile Ala Ile Tyr Gly Lys Asn Phe Cys Val Ser Ala
 565 570 575
 Lys Asn Ala Phe Met Leu Leu Met Arg Asn Ile Val Arg Val Val Val
 580 585 590
 Leu Asp Lys Val Thr Asp Leu Leu Leu Phe Phe Gly Lys Leu Leu Val
 595 600 605
 Val Gly Gly Val Gly Val Leu Ser Phe Phe Phe Phe Ser Gly Arg Ile
 610 615 620
 Pro Gly Leu Gly Lys Asp Phe Lys Ser Pro His Leu Asn Tyr Tyr Trp
 625 630 635 640
 Leu Pro Ile Met Thr Ser Ile Leu Gly Ala Tyr Val Ile Ala Ser Gly
 645 650 655
 Phe Phe Ser Val Phe Gly Met Cys Val Asp Thr Leu Phe Leu Cys Phe
 660 665 670
 Leu Glu Asp Leu Glu Arg Asn Asn Gly Ser Leu Asp Arg Pro Tyr Tyr
 675 680 685
 Met Ser Lys Ser Leu Leu Lys Ile Leu Gly Lys Lys Asn Glu Ala Pro
 690 695 700
 Pro Asp Asn Lys Lys Arg Lys Lys
 705 710

<210> SEQ ID NO 110

<211> LENGTH: 710

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 110

Met Gly Gly Lys Gln Arg Asp Glu Asp Asp Glu Ala Tyr Gly Lys Pro
 1 5 10 15

-continued

Val	Lys	Tyr	Asp	Pro	Ser	Phe	Arg	Gly	Pro	Ile	Lys	Asn	Arg	Ser	Cys	
	20							25				30				
Thr	Asp	Val	Ile	Cys	Cys	Val	Leu	Phe	Leu	Leu	Phe	Ile	Leu	Gly	Tyr	
	35						40					45				
Ile	Val	Val	Gly	Ile	Val	Ala	Trp	Leu	Tyr	Gly	Asp	Pro	Arg	Gln	Val	
	50					55					60					
Leu	Tyr	Pro	Arg	Asn	Ser	Thr	Gly	Ala	Tyr	Cys	Gly	Met	Gly	Glu	Asn	
	65				70					75					80	
Lys	Asp	Lys	Pro	Tyr	Leu	Leu	Tyr	Phe	Asn	Ile	Phe	Ser	Cys	Ile	Leu	
				85					90					95		
Ser	Ser	Asn	Ile	Ile	Ser	Val	Ala	Glu	Asn	Gly	Leu	Gln	Cys	Pro	Thr	
			100					105					110			
Pro	Gln	Val	Cys	Val	Ser	Ser	Cys	Pro	Glu	Asp	Pro	Trp	Thr	Val	Gly	
		115					120					125				
Lys	Asn	Glu	Phe	Ser	Gln	Thr	Val	Gly	Glu	Val	Phe	Tyr	Thr	Lys	Asn	
	130					135					140					
Arg	Asn	Phe	Cys	Leu	Pro	Gly	Val	Pro	Trp	Asn	Met	Thr	Val	Ile	Thr	
	145				150				155						160	
Ser	Leu	Gln	Gln	Glu	Leu	Cys	Pro	Ser	Phe	Leu	Leu	Pro	Ser	Ala	Pro	
			165						170					175		
Ala	Leu	Gly	Arg	Cys	Phe	Pro	Trp	Thr	Asn	Val	Thr	Pro	Pro	Ala	Leu	
		180					185					190				
Pro	Gly	Ile	Thr	Asn	Asp	Thr	Thr	Ile	Gln	Gln	Gly	Ile	Ser	Gly	Leu	
	195					200						205				
Ile	Asp	Ser	Leu	Asn	Ala	Arg	Asp	Ile	Ser	Val	Lys	Ile	Phe	Glu	Asp	
	210				215						220					
Phe	Ala	Gln	Ser	Trp	Tyr	Trp	Ile	Leu	Val	Ala	Leu	Gly	Val	Ala	Leu	
	225				230					235					240	
Val	Leu	Ser	Leu	Leu	Phe	Ile	Leu	Leu	Leu	Arg	Leu	Val	Ala	Gly	Pro	
			245						250				255			
Leu	Val	Leu	Val	Leu	Ile	Leu	Gly	Val	Leu	Gly	Val	Leu	Ala	Tyr	Gly	
			260				265					270				
Ile	Tyr	Tyr	Cys	Trp	Glu	Glu	Tyr	Arg	Val	Leu	Arg	Asp	Lys	Gly	Ala	
	275						280					285				
Ser	Ile	Ser	Gln	Leu	Gly	Phe	Thr	Thr	Asn	Leu	Ser	Ala	Tyr	Gln	Ser	
	290					295					300					
Val	Gln	Glu	Thr	Trp	Leu	Ala	Ala	Leu	Ile	Val	Leu	Ala	Val	Leu	Glu	
	305				310					315					320	
Ala	Ile	Leu	Leu	Leu	Met	Leu	Ile	Phe	Leu	Arg	Gln	Arg	Ile	Arg	Ile	
			325						330				335			
Ala	Ile	Ala	Leu	Leu	Lys	Glu	Ala	Ser	Lys	Ala	Val	Gly	Gln	Met	Met	
		340						345					350			
Ser	Thr	Met	Phe	Tyr	Pro	Leu	Val	Thr	Phe	Val	Leu	Leu	Leu	Ile	Cys	
		355				360						365				
Ile	Ala	Tyr	Trp	Ala	Met	Thr	Ala	Leu	Tyr	Leu	Ala	Thr	Ser	Gly	Gln	
	370					375					380					
Pro	Gln	Tyr	Val	Leu	Trp	Ala	Ser	Asn	Ile	Ser	Ser	Pro	Gly	Cys	Glu	
	385				390					395					400	
Lys	Val	Pro	Ile	Asn	Thr	Ser	Cys	Asn	Pro	Thr	Ala	His	Leu	Val	Asn	
			405						410				415			
Ser	Ser	Cys	Pro	Gly	Leu	Met	Cys	Val	Phe	Gln	Gly	Tyr	Ser	Ser	Lys	
		420						425				430				
Gly	Leu	Ile	Gln	Arg	Ser	Val	Phe	Asn	Leu	Gln	Ile	Tyr	Gly	Val	Leu	

-continued

435	440	445
Gly Leu Phe Trp Thr Leu Asn Trp Val Leu Ala Leu Gly Gln Cys Val 450 455 460		
Leu Ala Gly Ala Phe Ala Ser Phe Tyr Trp Ala Phe His Lys Pro Gln 465 470 475 480		
Asp Ile Pro Thr Phe Pro Leu Ile Ser Ala Phe Ile Arg Thr Leu Arg 485 490 495		
Tyr His Thr Gly Ser Leu Ala Phe Gly Ala Leu Ile Leu Thr Leu Val 500 505 510		
Gln Ile Ala Arg Val Ile Leu Glu Tyr Ile Asp His Lys Leu Arg Gly 515 520 525		
Val Gln Asn Pro Val Ala Arg Cys Ile Met Cys Cys Phe Lys Cys Cys 530 535 540		
Leu Trp Cys Leu Glu Lys Phe Ile Lys Phe Leu Asn Arg Asn Ala Tyr 545 550 555 560		
Ile Met Ile Ala Ile Tyr Gly Lys Asn Phe Cys Val Ser Ala Lys Asn 565 570 575		
Ala Phe Met Leu Leu Met Arg Asn Ile Val Arg Val Val Val Leu Asp 580 585 590		
Lys Val Thr Asp Leu Leu Leu Phe Phe Gly Lys Leu Leu Val Val Gly 595 600 605		
Gly Val Gly Val Leu Ser Phe Phe Phe Phe Ser Gly Arg Ile Pro Gly 610 615 620		
Leu Gly Lys Asp Phe Lys Ser Pro His Leu Asn Tyr Tyr Trp Leu Pro 625 630 635 640		
Ile Met Thr Ser Ile Leu Gly Ala Tyr Val Ile Ala Ser Gly Phe Phe 645 650 655		
Ser Val Phe Gly Met Cys Val Asp Thr Leu Phe Leu Cys Phe Leu Glu 660 665 670		
Asp Leu Glu Arg Asn Asn Gly Ser Leu Asp Arg Pro Tyr Tyr Met Ser 675 680 685		
Lys Ser Leu Leu Lys Ile Leu Gly Lys Lys Asn Glu Ala Pro Pro Asp 690 695 700		
Asn Lys Lys Arg Lys Lys 705 710		

<210> SEQ ID NO 111

<211> LENGTH: 712

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 111

Met Gly Gly Lys Gln Arg Asp Glu Asp Asp Glu Ala Tyr Gly Lys Pro 1 5 10 15
Val Lys Tyr Asp Pro Ser Phe Arg Gly Pro Ile Lys Asn Arg Ser Cys 20 25 30
Thr Asp Val Ile Cys Cys Val Leu Phe Leu Leu Phe Ile Leu Gly Tyr 35 40 45
Ile Val Val Gly Ile Val Ala Trp Leu Tyr Gly Asp Pro Arg Gln Val 50 55 60
Leu Tyr Pro Arg Asn Ser Thr Gly Ala Tyr Cys Gly Met Gly Glu Asn 65 70 75 80
Lys Asp Lys Pro Tyr Leu Leu Tyr Phe Asn Ile Phe Ser Cys Ile Leu 85 90 95
Ser Ser Asn Ile Ile Ser Val Ala Glu Asn Gly Leu Gln Cys Pro Thr


-continued

100				105				110			
Pro	Gln	Val	Cys	Val	Ser	Ser	Cys	Pro	Glu	Asp	Pro
	115						120				125
Lys	Asn	Glu	Phe	Ser	Gln	Thr	Val	Gly	Glu	Val	Phe
	130					135					140
Arg	Asn	Phe	Cys	Leu	Pro	Gly	Val	Pro	Trp	Asn	Met
	145				150					155	
Ser	Leu	Gln	Gln	Glu	Leu	Cys	Pro	Ser	Phe	Leu	Leu
				165					170		
Ala	Leu	Gly	Arg	Cys	Phe	Pro	Trp	Thr	Asn	Val	Thr
			180					185			
Pro	Gly	Ile	Thr	Asn	Asp	Thr	Thr	Ile	Gln	Gln	Gly
	195						200				205
Ile	Asp	Ser	Leu	Asn	Ala	Arg	Asp	Ile	Ser	Val	Lys
	210					215					220
Phe	Ala	Gln	Ser	Trp	Tyr	Trp	Ile	Leu	Val	Ala	Leu
	225				230					235	
Val	Leu	Ser	Leu	Leu	Phe	Ile	Leu	Leu	Arg	Leu	Val
			245						250		
Leu	Val	Leu	Val	Leu	Ile	Leu	Gly	Val	Leu	Gly	Val
			260					265			
Ile	Tyr	Tyr	Cys	Trp	Glu	Glu	Tyr	Arg	Val	Leu	Arg
	275						280				
Ser	Ile	Ser	Gln	Leu	Gly	Phe	Thr	Thr	Asn	Leu	Ser
	290					295					300
Val	Gln	Glu	Thr	Trp	Leu	Ala	Ala	Leu	Ile	Val	Leu
	305				310					315	
Ala	Ile	Leu	Leu	Leu	Met	Leu	Ile	Phe	Leu	Arg	Gln
			325						330		
Ala	Ile	Ala	Leu	Leu	Lys	Glu	Ala	Ser	Lys	Ala	Val
			340					345			
Ser	Thr	Met	Phe	Tyr	Pro	Leu	Val	Thr	Phe	Val	Leu
		355					360				365
Ile	Ala	Tyr	Trp	Ala	Met	Thr	Ala	Leu	Tyr	Pro	Leu
	370					375					380
Ala	Thr	Leu	Gly	Tyr	Val	Leu	Trp	Ala	Ser	Asn	Ile
	385				390					395	
Cys	Glu	Lys	Val	Pro	Ile	Asn	Thr	Ser	Cys	Asn	Pro
			405						410		
Val	Asn	Ser	Ser	Cys	Pro	Gly	Leu	Met	Cys	Val	Phe
			420					425			
Ser	Lys	Gly	Leu	Ile	Gln	Arg	Ser	Val	Phe	Asn	Leu
	435						440				445
Val	Leu	Gly	Leu	Phe	Trp	Thr	Leu	Asn	Trp	Val	Leu
	450					455					460
Cys	Val	Leu	Ala	Gly	Ala	Phe	Ala	Ser	Phe	Tyr	Trp
	465				470					475	
Pro	Gln	Asp	Ile	Pro	Thr	Phe	Pro	Leu	Ile	Ser	Ala
			485						490		
Leu	Arg	Tyr	His	Thr	Gly	Ser	Leu	Ala	Phe	Gly	Ala
			500					505			
Leu	Val	Gln	Ile	Ala	Arg	Val	Ile	Leu	Glu	Tyr	Ile
	515						520				525

Figure 1: 24P4C12 SSH sequence of 160 nucleotides. (SEQ ID NO: 1)

```
1  GATCAGGGCG GCCAGCCAGG TCTCCTGCAC GCTCTGGTAG GCACTGAGGT TGGTGGTGAA
61  ACCCAGCTGG GAGATGGAGG CGCCCTCGTC CCGCAGCACT CGGTACTCCT CCCAGCAGTA
121 GTAGATGCCA TATGCCAGCA CGCCAGCAC TCCCAGGATC
```


Figure 7: 24P4C12 variant 1 % Accessible Residues
Profile (Janin J., 1979. Nature 277:491-492)



-continued

Arg	Gly	Val	Gln	Asn	Pro	Val	Ala	Arg	Cys	Ile	Met	Cys	Cys	Phe	Lys
530						535					540				
Cys	Cys	Leu	Trp	Cys	Leu	Glu	Lys	Phe	Ile	Lys	Phe	Leu	Asn	Arg	Asn
545					550					555					560
Ala	Tyr	Ile	Met	Ile	Ala	Ile	Tyr	Gly	Lys	Asn	Phe	Cys	Val	Ser	Ala
			565						570					575	
Lys	Asn	Ala	Phe	Met	Leu	Leu	Met	Arg	Asn	Ile	Val	Arg	Val	Val	Val
		580						585					590		
Leu	Asp	Lys	Val	Thr	Asp	Leu	Leu	Phe	Phe	Gly	Lys	Leu	Leu	Val	
	595						600				605				
Val	Gly	Gly	Val	Gly	Val	Leu	Ser	Phe	Phe	Phe	Phe	Ser	Gly	Arg	Ile
	610					615					620				
Pro	Gly	Leu	Gly	Lys	Asp	Phe	Lys	Ser	Pro	His	Leu	Asn	Tyr	Tyr	Trp
625					630					635					640
Leu	Pro	Ile	Met	Thr	Ser	Ile	Leu	Gly	Ala	Tyr	Val	Ile	Ala	Ser	Gly
			645						650					655	
Phe	Phe	Ser	Val	Phe	Gly	Met	Cys	Val	Asp	Thr	Leu	Phe	Leu	Cys	Phe
			660					665					670		
Leu	Glu	Asp	Leu	Glu	Arg	Asn	Asn	Gly	Ser	Leu	Asp	Arg	Pro	Tyr	Tyr
	675						680					685			
Met	Ser	Lys	Ser	Leu	Leu	Lys	Ile	Leu	Gly	Lys	Lys	Asn	Glu	Ala	Pro
	690					695					700				
Pro	Asp	Asn	Lys	Lys	Arg	Lys	Lys								
705					710										

<210> SEQ ID NO 112

<211> LENGTH: 710

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 112

Met	Gly	Gly	Lys	Gln	Arg	Asp	Glu	Asp	Asp	Glu	Ala	Tyr	Gly	Lys	Pro
1				5					10					15	
Val	Lys	Tyr	Asp	Pro	Ser	Phe	Arg	Gly	Pro	Ile	Lys	Asn	Arg	Ser	Cys
			20					25					30		
Thr	Asp	Val	Ile	Cys	Cys	Val	Leu	Phe	Leu	Leu	Phe	Ile	Leu	Gly	Tyr
		35					40					45			
Ile	Val	Val	Gly	Ile	Val	Ala	Trp	Leu	Tyr	Gly	Asp	Pro	Arg	Gln	Val
	50					55					60				
Leu	Tyr	Pro	Arg	Asn	Ser	Thr	Gly	Ala	Tyr	Cys	Gly	Met	Gly	Glu	Asn
65				70					75					80	
Lys	Asp	Lys	Pro	Tyr	Leu	Leu	Tyr	Phe	Asn	Ile	Phe	Ser	Cys	Ile	Leu
			85						90					95	
Ser	Ser	Asn	Ile	Ile	Ser	Val	Ala	Glu	Asn	Gly	Leu	Gln	Cys	Pro	Thr
		100						105					110		
Pro	Gln	Val	Cys	Val	Ser	Ser	Cys	Pro	Glu	Asp	Pro	Trp	Thr	Val	Gly
		115					120					125			
Lys	Asn	Glu	Phe	Ser	Gln	Thr	Val	Gly	Glu	Val	Phe	Tyr	Thr	Lys	Asn
	130					135					140				
Arg	Asn	Phe	Cys	Leu	Pro	Gly	Val	Pro	Trp	Asn	Met	Thr	Val	Ile	Thr
145				150						155					160
Ser	Leu	Gln	Gln	Glu	Leu	Cys	Pro	Ser	Phe	Leu	Leu	Pro	Ser	Ala	Pro
			165						170					175	
Ala	Leu	Gly	Arg	Cys	Phe	Pro	Trp	Thr	Asn	Val	Thr	Pro	Pro	Ala	Leu
			180					185						190	

-continued

Pro Gly Ile Thr Asn Asp Thr Thr Ile Gln Gln Gly Ile Ser Gly Leu	195	200	205
Ile Asp Ser Leu Asn Ala Arg Asp Ile Ser Val Lys Ile Phe Glu Asp	210	215	220
Phe Ala Gln Ser Trp Tyr Trp Ile Leu Val Ala Leu Gly Val Ala Leu	225	230	235
Val Leu Ser Leu Leu Phe Ile Leu Leu Leu Arg Leu Val Ala Gly Pro	245	250	255
Leu Val Leu Val Leu Ile Leu Gly Val Leu Gly Val Leu Ala Tyr Gly	260	265	270
Ile Tyr Tyr Cys Trp Glu Glu Tyr Arg Val Leu Arg Asp Lys Gly Ala	275	280	285
Ser Ile Ser Gln Leu Gly Phe Thr Thr Asn Leu Ser Ala Tyr Gln Ser	290	295	300
Val Gln Glu Thr Trp Leu Ala Ala Leu Ile Val Leu Ala Val Leu Glu	305	310	315
Ala Ile Leu Leu Leu Met Leu Ile Phe Leu Arg Gln Arg Ile Arg Ile	325	330	335
Ala Ile Ala Leu Leu Lys Glu Ala Ser Lys Ala Val Gly Gln Met Met	340	345	350
Ser Thr Met Phe Tyr Pro Leu Val Thr Phe Val Leu Leu Leu Ile Cys	355	360	365
Ile Ala Tyr Trp Ala Met Thr Ala Leu Tyr Leu Ala Thr Ser Gly Gln	370	375	380
Pro Gln Tyr Val Leu Trp Ala Ser Asn Ile Ser Ser Pro Gly Cys Glu	385	390	395
Lys Val Pro Ile Asn Thr Ser Cys Asn Pro Thr Ala His Leu Val Asn	405	410	415
Ser Ser Cys Pro Gly Leu Met Cys Val Phe Gln Gly Tyr Ser Ser Lys	420	425	430
Gly Leu Ile Gln Arg Ser Val Phe Asn Leu Gln Ile Tyr Gly Val Leu	435	440	445
Gly Leu Phe Trp Thr Leu Asn Trp Val Leu Ala Leu Gly Gln Cys Val	450	455	460
Leu Ala Gly Ala Phe Ala Ser Phe Tyr Trp Ala Phe His Lys Pro Gln	465	470	475
Asp Ile Pro Thr Phe Pro Leu Ile Ser Ala Phe Ile Arg Thr Leu Arg	485	490	495
Tyr His Thr Gly Ser Leu Ala Phe Gly Ala Leu Ile Leu Thr Leu Val	500	505	510
Gln Ile Ala Arg Val Ile Leu Glu Tyr Ile Asp His Lys Leu Arg Gly	515	520	525
Val Gln Asn Pro Val Ala Arg Cys Ile Met Cys Cys Phe Lys Cys Cys	530	535	540
Leu Trp Cys Leu Glu Lys Phe Ile Lys Phe Leu Asn Arg Asn Ala Tyr	545	550	555
Ile Met Ile Ala Ile Tyr Gly Lys Asn Phe Cys Val Ser Ala Lys Asn	565	570	575
Ala Phe Met Leu Leu Met Arg Asn Ile Val Arg Val Val Val Leu Asp	580	585	590
Lys Val Thr Asp Leu Leu Leu Phe Phe Gly Lys Leu Leu Val Val Gly	595	600	605
Gly Val Gly Val Leu Ser Phe Phe Phe Phe Ser Gly Arg Ile Pro Gly	610	615	620

[illegible]

* * * * *

Figure 8: 24P4C12 variant 1 Average Flexibility Profile
(Bhaskaran R., Ponnuswamy P.K., 1988.
Int. J. Pept. Protein Res. 32:242-255)




Figure 9: 24P4C12 variant 1 Beta-turn Profile
(Deleage, G., Roux B. 1987. Protein Engineering 1:289-294)





Figure 10


Figure 11

Figure 12




**Fig. 13A: Secondary structure prediction
of 24P4C12 variant 1**

10 20 30 40 50 60 70
| | | | | | |
MGGKQRDEDDDEAYGKPVKYDPSFRGPIKNRSCTDVICCVLFLLFILGYIVVGIVAWLYGDPRQVLYPRNS
ccccccccccccccccccccccccccccccccccccchhhhhhhhhhhhhhhhhhhhhhhhhheccccceeecccc
TGAYCGMGENKDKPYLLYFNIFSCILSSNIISVAENGLQCPTPQVCVSSCPEDPWTVGKNEFSQTVGEVF
ccccccccccccccccceehhhhhhhhhccchhhhhccccccccccccceccccccccccccchchccceee
YTKNRNFCCLPGVPWNMTVITSLQQELCPSFLLPSAPALGRCFPWTNVTTPALPGITNDTTIQGISGLID
eeccccceccccccccceeeehccccccccccccccccccccccccccccccccccccchhhhhhhhhh
SLNARDISVKIFEDFAQSWYWLVALGVALVLSLLFILLRLVAGPLVLVLILGVLGVLAYGIYYCWEEY
hccccchhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhchhhhhhhhhhhhhhhcheeechh
RVLDRKGASISQLGFTTNLSAYQSVQETWLAALIVLAVLEAILLLMLIFLRQIRIAIALKEASKAVGQ
hhhhhccccceeeccccccccchhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhh
MMSTMFYPLVTFVLLLICIAYWAMTALYLATSGQPQYVLWASNISSPGCEKVPINTSCNPTAHLVNSSCP
hhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhccccceeecccccccccccccccccccccecccc
GLMCFVQGYSSKGLIQRSVFNLIQYGVGLGFWTLNWLALGQCVLAGAFASFYWAFHKPQDIPTFPLISA
ceeececcccchhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhccccccccchhhh
FIRTLRYHTGSLAFGALILTLVQIARVILEYIDHKLRGVQNPVARCIMCCFKCCLWCLEKFIKFLNRNAY
hhhhhhhccccchhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhccce
IMIAIYGKNFCVSAKNAFMLLMRNIVRVVLDKVTDLLLFFGKLLVVGGVGLSFFFFSGRIPGLGKDFK
eeeeeeccchhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhccccceccccchheeecccccccccccc
SPHLNYYWLPIMTSILGAYVIASGFFSVFGMCDTLFLCFLEDLERNNGSLDRPYYSKSLKILGKKNE
ccccchhhhhhhhhhhchhehhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhccccccccchhhhhhhhhhhcccc
APPDNKKRKK
cccccccccc

Alpha helix (h) : 53.94%
Extended strand (e) : 9.44%
Random coil (c) : 36.62%

Figure 13B: Transmembrane prediction for 24P4C12 variant 1




Figure 13C: Transmembrane prediction for 24P4C12 variant 1
TMHMM posterior probabilities for Sequence




Figure 14: 24P4C12 Expression by RT-PCR




Figure 2:

Figure 2A. The cDNA (SEQ ID. NO. : 2) and amino acid sequence (SEQ ID. NO. : 3) of 24P4C12 v.1.

The start methionine is underlined. The open reading frame extends from nucleic acid 6-2138 including the stop codon.

Figure 2A-1

```
1      M G G K Q E D E D D E A Y G K P V K Y
1 gagccATGGGGGGAAAGCAGCGGGACGAGGATGACGAGGCCTACGGGAAGCCAGTCAAAT
20   D P S F R G P I K N R S C I D V I C C V
61   ACGACCCCTCCTTTCGAGGCCCCATCAAGAACAGAAGCTGCACAGATGTCATCTGCTGCG
40   L F L L F I L G Y I V V G I V A W L Y G
121  TCCTCTTCTGCTCTTTCATTCTAGGTTACATCGTGGTGGGATTGTGGCCTGGTTGTATG
60   D P R Q V L Y P R N S T G A Y C G M G E
181  GAGACCCCCGGCAAGTCTCTACCCAGGAACCTCTACTGGGGCCTACTGTGGCATGGGGG
80   N K D K P Y L L Y F N I F S C I L S S N
241  AGAACAAAGATAAGCCGTATCTCCTGTACTTCAACATCTTCAGCTGCATCCTGTCCAGCA
100  I I S V A E N G L Q C P T F Q V C V S S
301  ACATCATCTCAGTTGCTGAGAACGGCCTACAGTGCCCCACACCCAGGTGTGTGTCTCT
120  C P E D P W T V G K N E F S Q T V G E V
361  CCTGCCCGGAGGACCCATGGACTGTGGGAAAAACGAGTTCTCACAGACTGTTGGGGAAG
140  F Y T K N R N F C L P G V P W N M T V I
421  TCTTCTATACAAAAACAGGAACCTTTGTCTGCCAGGGGTACCCTGGAATATGACGGTGA
160  I S L Q Q E L C P S F L L F S A P A L G
481  TCACAAGCCTGCAACAGGAACCTTGCCCCAGTTTCCTCCTCCCCTCTGCTCCAGCTCTGG
180  R C F P W T N V T P P A L P G I T N D T
541  GCGCTGCTTTCCATGGACCAACGTTACTCCACCGGCGCTCCCAGGGATCACCAATGACA
200  T I Q Q G I S G L I D S L N A R D I S V
601  CCACCATACAGCAGGGGATCAGCGGTCTTATTGACAGCCTCAATGCCCGAGACATCAGTG
220  K I F E D F A Q S W Y W I L V A L G V A
661  TTAAGATCTTTGAAGATTTTGCCAGTCCTGGTATTGGATTCTTGTGGCCTGGGGGTGG
240  L V L S L L F I L L L R L V A G P L V L
721  CTCTGGTCTTGAGCCTACTGTTTATCTTGCTTCTGCGCCTGGTGGCTGGGCCCCTGGTGC
260  V L I L G V L G V L A Y G I Y Y C W E E
781  TGGTGTGATCCTGGGAGTGCTGGGCGTGCTGGCATAACGGCATCTACTACTGCTGGGAGG
280  Y R V L R D K G A S I S Q L G F T T N L
841  AGTACCGAGTGCTGCGGGACAAGGGCGCCTCCATCTCCCAGCTGGGTTTCACCACCAACC
300  S A Y Q S V Q E T W L A A L I V L A V L
901  TCAGTGCTTACCAGAGCGTGCAGGAGACCTGGCTGGCCGCCCTGATCGTGTGGCGGTGC
320  E A I L L L M L I F L R Q R I R I A I A
961  TTGAAGCCATCCTGCTGCTGATGCTCATCTTCTGCGGCAGCGGATTCTGATTGCCATCG
340  L L K E A S K A V G Q M M S T M F Y P L
```

Figure 15: Expression of 24P4C12 in Normal Tissues




Figure 16: Expression of 24P4C12 in Prostate Cancer Xenografts and Cell Lines




Figure 17: Expression of 24P4C12 in Human Patient Cancer Specimens




Figure 18: Expression of 24P4C12 in Prostate Cancer Patient Specimens




Figure 19: Expression of 24P4C12 in Colon Cancer Patient Samples




Figure 20: Expression of 24P4C12 in Lung Cancer Patient Samples





**Figure 21: Expression of 24P4C12 in
Patient Cancer Specimens**



T = tumor RNA

N = normal adjacent tissue RNA

Figure 22-1: 24P4C12 Expression in Patient Cancer Specimens

A.

	Diagnosis	Grade	24P4C12
1	Serous CA	3	
2	Mucinous CystadenoCA		
3	Mucinous CystadenoCA		
4	Mucinous CystadenoCA		
5	Granulosa	2	
6	Papillary serous CA	Fig0 Ia1	
7	Endometrioid AdenoCA	3	
8	Endometrioid AdenoCA	2	
9	Papillary serous CA	3	
10	ND		
11	Squamous		
12	Adeno CA Met to Oment.	3	
13	Papillary serous CA		
14	Endometrioid AdenoCA	Fig0 III	
15	Met to omentum		
	Met to Intestine	Undiff.	
Percentage positive			73.3%

B.

	Diagnosis	Stage/Grade	24P4C12
1	AdenoCA	T1cN0M0	
2	AdenoCA	Grade II	
3	AdenoCA Well Diff.	Grade 2	
4	Sarcoma		
5	Endometrioid CA	Grade 2	
6	Endometrioid CA	T1bNxMx	
7	Endometrioid CA	T1bNxMx	
8	Intraepithelial Neopl.	T3aN0Mx	
9	Sarcoma	Grade III	
10	Endometrioid CA		
11	Endometrioid CA	Grade II	
12	Endometrioid CA	Grade 2	
Percentage Positive			83.3%



ND = Not Determined

Figure 22-2: 24P4C12 Expression in Patient Cancer Specimens




Figure 22-3: 24P4C12 Expression in Patient Cancer Specimens

E.

	Pathology	Grade	24P4C12
1	Adeno	3	
2	Adeno	I	
3	Adeno	IB	
4	Adeno	IB	
5	Adeno	IIIA	
6	Adeno	IIIA	
7	Adeno	Mod Diff	
8	Adeno	Mod Diff	
9	Bronchioalv.	IA	
10	Large Cell	I	
11	Large Cell	IIB	
12	Large Cell	IIIA	
13	Large Cell	IV	
14	Non-small cell	Not Malignant	
15	Papillary	I	
16	Papillary	IB	
17	Papillary	IV	
18	Small Cell	I	
19	Small Cell	I	
20	Small Cell	I	
21	Small Cell	IIB	
22	Squamous	IB	
23	Squamous	IB	
24	Squamous	IB	
25	Squamous	IIB	
26	Squamous	IIB	
27	Squamous	IIIA	
28	Squamous	IIIA	
29	Squamous	ND	
30	Squamous	ND	
31	Squamous	ND	
Percentage positive			80.6%

F.

	Pathology	24P4C12
1	AdenoCA, ductal type, poorly diff.	
2	AdenoCA, invasive, T2N0Mx	
3	Pseudopapillary CA	
4	Small cell anaplastic, mets to LN	
5	Undifferentiated CA	
6	AdenoCA met	
7	AdenoCA met to ovary	
8	AdenoCA, mod to poorly diff.	
Percentage Positive		87.5%



Figure 2A-2

1021 CCCTCCTGAAGGAGGCCAGCAAGGCTGTGGGACAGATGATGTCTACCATGTTCTACCCAC
360 V T F V L L L I C I A Y W A M T A L Y L
1081 TGGTCACCTTTGTCTCCTCCTCATCTGCATTGCCTACTGGGCCATGACTGCTCTGTACC
380 A T S G Q P Q Y V L W A S N I S S P G C
1141 TGGCTACATCGGGGCAACCCCAAGTATGTGCTCTGGGCATCCAACATCAGCTCCCCCGGCT
400 E K V P I N T S C N P T A H L V N S S C
1201 GTGAGAAAGTGCCAATAAATACATCATGCAACCCACGGCCACCTTGTGAACCTCCTCGT
420 P G L M C V F Q G Y S S K G L I Q R S V
1261 GCCCAGGGCTGATGTGCGTCTTCCAGGGCTACTCATCCAAAGGCCTAATCCAACGTTCTG
440 F N L Q I Y G V L G L F W T L N W V L A
1321 TCTTCAATCTGCAAATCTATGGGGTCTGGGGCTCTTCTGGACCCCTTAACCTGGGTACTGG
460 L G Q C V L A G A F A S F Y W A F H K P
1381 CCCTGGGCCAATGCGTCTCGCTGGAGCCTTTGCCTCCTTCTACTGGGCCTTCCACAAGC
480 Q D I P T F F L I S A F I R T L R Y H T
1441 CCCAGGACATCCCTACCTTCCCCTTAATCTCTGCCTTCATCCGCACACTCCGTTACCACA
500 G S L A F G A L I L T L V Q I A R V I L
1501 CTGGGTCATTGGCATTGGAGCCCTCATCTGACCCTTGTGCAGATAGCCCGGGTCACT
520 E Y I D H K L R G V Q N P V A R C I M C
1561 TGGAGTATATTGACCACAAGCTCAGAGGAGTGCAGAACCCTGTAGCCCGCTGCATCATGT
540 C F K C C L W C L E R F I K F L N R N A
1621 GCTGTTTCAAGTGCTGCCTCTGGTGTCTGGAAAAATTTATCAAGTTCCTAAACCGCAATG
560 Y I M I A I Y G K N F C V S A K N A F M
1681 CATACTCATGATCGCCATCTACGGGAAGAATTTCTGTGTCTCAGCCAAAAATGCGTTCA
580 L L M R N I V R V V V L D K V T D L L L
1741 TGCTACTCATGCGAAACATTGTGAGGGTGGTCTGCTGGACAAAGTCACAGACCTGCTGC
600 F F G K L L V V G G V G V L S F F F F S
1801 TGTTCTTTGGGAAGCTGCTGGTGGTGGAGGCGTGGGGTCTGTCCTTCTTTTTTTTCT
620 G R I P G L G K D F K S P H L N Y Y W L
1861 CCGGTGCGATCCCGGGGCTGGGTAAAGACTTTAAGAGCCCCCACCTCAACTATTACTGGC
640 F I M T S I L G A Y V I A S G F F S V F
1921 TGCCCATCATGACCTCCATCCTGGGGGCTATGTCATCGCCAGCGGCTTCTCAGCGTTT
660 G M C V D T L F L C F L E D L E R N N G
1981 TCGGCATGTGTGTGGACACGCTCTTCTCTGCTTCTGGAAGACCTGGAGCGGAACAACG
680 S L D R P Y Y M S K S L L K I L G K K N
2041 GCTCCCTGGACCGGCCCTACTACATGTCCAAGAGCCTTCTAAAGATTCTGGGCAAGAAGA
700 E A P P D N K K R K R *
2101 ACGAGGCGCCCCCGGACAACAAGAAGAGGAAGAAGTGAcagctccggccctgatccagga
2161 ctgcacccccacccccaccgtccagccatccaacctcacttcgccttacaggctctccattt
2221 tgtggtataaaaaaggttttaggccaggcgccgtggctcacgcctgtaatccaacactttg
2281 agaggctgaggcgggcggatcacctgagtcaggagttcgagaccagcctggccaacatgg

Figure 22-4: 24P4C12 Expression in Patient Cancer Specimens




Figure 23: Expression of 24P4C12 in Transduced Cells




Figure 24: Expression of 24P4C12 in 293T cells




Figure 25: Expression and detection of 24P4C12 in stably transduced PC3 cells




**Figure 26: Expression of recombinant
24P4C12 antigens
in 293T cells**







Figure 27: Monoclonal antibodies detect 24P4C12 protein expression in 293T cells by flow cytometry





- PC3 and 3T3 were grown overnight in low FBS. Cells were then incubated in low or 10% FBS as indicated. Proliferation was measured by Alamar Blue.
- Experiments were performed in triplicate

Figure 29: Detection of 24P4C12 protein by immunohistochemistry in prostate cancer patient specimens.




Figure 30: Detection of 24P4C12 protein by immunohistochemistry in various cancer patient specimens.





Figure 31: 24P4C12 Enhances Tumor Growth in SCID Mice



•1 x 10⁶ PC3-24P4C12 cells were mixed with Matrigel and injected on the right and left subcutaneous flanks of 4 male SCID mice per group.

•Each data point represents mean tumor volume (n=8).

Figure 2A-3

```
2341 tgaacacctcgctctctattaaaaatacaaaaattagccgagagtggtggcatgcacctgt
2401 catcccagctactcgggaggtgagggcaggagaatcgcttgaacccgggaggcagaggtt
2461 gcagtgagccgagatcgcgccactgcactccaacctgggtgacagactctgtctccaaaa
2521 caaaacaaacaaacaaaaagattttattaaagatatTTTgttaactcagtaaaaaaaaaa
2581 aaaaaaa
```


Figure 2B. The cDNA (SEQ ID. NO. : 4) and amino acid sequence (SEQ ID. NO. : 5) of 24P4C12 v.2.

The start methionine is underlined. The open reading frame extends from nucleic acid 6-2138 including the stop codon.

Figure 2B-1

```
1      M G G K Q R D E D D E A Y G R P V K Y
1 gagccATGGGGGAAAGCAGCGGGACGAGGATGACGAGGCCTACGGGAAGCCAGTCAAAT
20    D P S F R G P I K N R S C T D V I C C V
61 ACGACCCCTCCTTTTCGAGGCCCATCAAGAACAGAAGCTGCACAGATGTCATCTGCTGCG
40    L F L L F I L G Y I V V G I V A W L Y G
121 TCCTCTTCCTGCTCTTCATTCIAGGTTACATCGTGGTGGGGATTGTGGCCTGGTTGTATG
60    D P R Q V L Y P R N S T G A Y C G M G E
181 GAGACCCCGGCAAGTCCTCTACCCAGGAACCTCTACTGGGCTTACTGTGGCATGGGGG
80    N K D K P Y L L Y F N I F S C I L S S N
241 AGAACAAAGATAAGCCGTATCTCCTGTACTTCAACATCTTCAGCTGCATCCTGTCCAGCA
100   I I S V A E N G L Q C P T P Q V C V S S
301 ACATCATCTCAGTTGCTGAGAACGGCCTACAGTGCCCCACACCCAGGTGTGTGTCTCT
120   C P E D P W T V G R N E P S Q T V G E V
361 CCTGCCCGGAGGACCAATGGACTGTGGGAAAAAACGAGTTCACAGACTGTTGGGGAAG
140   F Y T K N R N F C L P G V P W N M T V I
421 TCTTCTATACAAAAACAGGAACCTTTGTCTGCCAGGGGTACCCTGGAATATGACGGTGA
160   T S L Q Q E L C P S F L L P S A F A L G
481 TCACAAGCCTGCAACAGGAACCTTGCCCCAGTTTCCTCCTCCCTCTGCTCCAGCTCTGG
180   R C F P W T N V T P P A L P G I T N D T
541 GACGCTGCTTTCATGGACCAACGTTACTCCACCGCGCTCCAGGGATCACCAATGACA
200   T I Q Q G I S G L I D S L N A R D I S V
601 CCACCATACAGCAGGGGATCAGCGGTCTTATTGACAGCCTCAATGCCCCGAGACATCAGTG
220   R I F E D F A Q S W Y W I L V A L G V A
661 TTAAGATCTTTGAAGATTTTGGCCAGTCCTGGTATTGGATTCTTGTGGCCCTGGGGGTGG
240   L V L S L L F I L L L R L V A G P L V L
721 CTCGGTCTTGAGCCTACTGTTTATCTTGCTTCTGCGCCTGGTGGCTGGGCCCTGGTGC
260   V L I L G V L G V L A Y G I Y Y C W E E
781 TGGTGTGATCCTGGGAGTGCTGGGCGTGCTGGCATAACGGCATCTACTACTGCTGGGAGG
280   Y R V L R D K G A S I S Q L G F I T N L
841 AGTACCGAGTGCTGCGGGACAAGGGCGCCTCCATCTCCAGCTGGGTTTACCACCAACC
```

Figure 32: 24P4C12 Enhances Tumor Growth in SCID Mice



• 1×10^6 3T3-24P4C12 cells were mixed with Matrigel and injected on the right subcutaneous flanks of 7 male SCID mice per group.

• Each data point represents mean tumor volume ($n=6$).

ANTIBODIES TO TUMOR ASSOCIATED PROTEINS

CROSS-REFERENCE TO RELATED APPLICATIONS

This application is a continuation of pending U.S. patent application Ser. No. 12/694,209, filed 26 Jan. 2010, which is a continuation of U.S. patent application Ser. No. 11/842,016, filed 20 Aug. 2007, now allowed, which is a continuation of pending U.S. patent application Ser. No. 11/518,610, filed 8 Sep. 2006, which is a continuation of U.S. patent application Ser. No. 10/306,631, filed 27 Nov. 2002, now U.S. Pat. No. 7,244,827, issued 17 Jul. 2007, which is a continuation-in-part of U.S. patent application Ser. No. 09/547,789, filed 12 Apr. 2000, now U.S. Pat. No. 6,943,235, issued 13 Sep. 2005, and claims priority to U.S. provisional patent application No. 60/128,858, filed 12 Apr. 1999. The contents of these applications are incorporated herein by reference.

STATEMENT OF RIGHTS TO INVENTIONS MADE UNDER FEDERALLY SPONSORED RESEARCH

Not applicable.

REFERENCE TO SEQUENCE LISTING SUBMITTED VIA EFS-WEB

The entire content of the following electronic submission of the sequence listing via the USPTO EFS-WEB server, as authorized and set forth in MPEP §1730 II.B.2(a)(C), is incorporated herein by reference in its entirety for all purposes. The sequence listing is identified on the electronically filed text file as follows:

File Name	Date of Creation	Size (bytes)
511582001107Seqlist.txt	June 22, 2010	236,574

FIELD OF THE INVENTION

The invention described herein relates to a gene and its encoded protein, termed 24P4C12, expressed in certain cancers, and to diagnostic and therapeutic methods and compositions useful in the management of cancers that express 24P4C12.

BACKGROUND OF THE INVENTION

Cancer is the second leading cause of human death next to coronary disease. Worldwide, millions of people die from cancer every year. In the United States alone, as reported by the American Cancer Society, cancer causes the death of well over a half-million people annually, with over 1.2 million new cases diagnosed per year. While deaths from heart disease have been declining significantly, those resulting from cancer generally are on the rise. In the early part of the next century, cancer is predicted to become the leading cause of death.

Worldwide, several cancers stand out as the leading killers. In particular, carcinomas of the lung, prostate, breast, colon, pancreas, and ovary represent the primary causes of cancer death. These and virtually all other carcinomas share a common lethal feature. With very few exceptions, metastatic dis-

ease from a carcinoma is fatal. Moreover, even for those cancer patients who initially survive their primary cancers, common experience has shown that their lives are dramatically altered. Many cancer patients experience strong anxieties driven by the awareness of the potential for recurrence or treatment failure. Many cancer patients experience physical debilitations following treatment. Furthermore, many cancer patients experience a recurrence.

Worldwide, prostate cancer is the fourth most prevalent cancer in men. In North America and Northern Europe, it is by far the most common cancer in males and is the second leading cause of cancer death in men. In the United States alone, well over 30,000 men die annually of this disease—second only to lung cancer. Despite the magnitude of these figures, there is still no effective treatment for metastatic prostate cancer. Surgical prostatectomy, radiation therapy, hormone ablation therapy, surgical castration and chemotherapy continue to be the main treatment modalities. Unfortunately, these treatments are ineffective for many and are often associated with undesirable consequences.

On the diagnostic front, the lack of a prostate tumor marker that can accurately detect early-stage, localized tumors remains a significant limitation in the diagnosis and management of this disease. Although the serum prostate specific antigen (PSA) assay has been a very useful tool, however its specificity and general utility is widely regarded as lacking in several important respects.

Progress in identifying additional specific markers for prostate cancer has been improved by the generation of prostate cancer xenografts that can recapitulate different stages of the disease in mice. The LAPC (Los Angeles Prostate Cancer) xenografts are prostate cancer xenografts that have survived passage in severe combined immune deficient (SCID) mice and have exhibited the capacity to mimic the transition from androgen dependence to androgen independence (Klein et al., 1997, Nat. Med. 3:402). More recently identified prostate cancer markers include PCTA-1 (Su et al., 1996, Proc. Natl. Acad. Sci. USA 93: 7252), prostate-specific membrane (PSM) antigen (Pinto et al., Clin Cancer Res 1996 Sep. 2 (9): 1445-51), STEAP (Hubert, et al., Proc Natl Acad Sci USA. 1999 Dec. 7; 96(25): 14523-8) and prostate stem cell antigen (PSCA) (Reiter et al., 1998, Proc. Natl. Acad. Sci. USA 95: 1735).

While previously identified markers such as PSA, PSM, PCTA and PSCA have facilitated efforts to diagnose and treat prostate cancer, there is need for the identification of additional markers and therapeutic targets for prostate and related cancers in order to further improve diagnosis and therapy.

Renal cell carcinoma (RCC) accounts for approximately 3 percent of adult malignancies. Once adenomas reach a diameter of 2 to 3 cm, malignant potential exists. In the adult, the two principal malignant renal tumors are renal cell adenocarcinoma and transitional cell carcinoma of the renal pelvis or ureter. The incidence of renal cell adenocarcinoma is estimated at more than 29,000 cases in the United States, and more than 11,600 patients died of this disease in 1998. Transitional cell carcinoma is less frequent, with an incidence of approximately 500 cases per year in the United States.

Surgery has been the primary therapy for renal cell adenocarcinoma for many decades. Until recently, metastatic disease has been refractory to any systemic therapy. With recent developments in systemic therapies, particularly immunotherapies, metastatic renal cell carcinoma may be approached aggressively in appropriate patients with a possibility of durable responses. Nevertheless, there is a remaining need for effective therapies for these patients.

Of all new cases of cancer in the United States, bladder cancer represents approximately 5 percent in men (fifth most common neoplasm) and 3 percent in women (eighth most common neoplasm). The incidence is increasing slowly, concurrent with an increasing older population. In 1998, there was an estimated 54,500 cases, including 39,500 in men and 15,000 in women. The age-adjusted incidence in the United States is 32 per 100,000 for men and eight per 100,000 in women. The historic male/female ratio of 3:1 may be decreasing related to smoking patterns in women. There were an estimated 11,000 deaths from bladder cancer in 1998 (7,800 in men and 3,900 in women). Bladder cancer incidence and mortality strongly increase with age and will be an increasing problem as the population becomes more elderly.

Most bladder cancers recur in the bladder. Bladder cancer is managed with a combination of transurethral resection of the bladder (TUR) and intravesical chemotherapy or immunotherapy. The multifocal and recurrent nature of bladder cancer points out the limitations of TUR. Most muscle-invasive cancers are not cured by TUR alone. Radical cystectomy and urinary diversion is the most effective means to eliminate the cancer but carry an undeniable impact on urinary and sexual function. There continues to be a significant need for treatment modalities that are beneficial for bladder cancer patients.

An estimated 130,200 cases of colorectal cancer occurred in 2000 in the United States, including 93,800 cases of colon cancer and 36,400 of rectal cancer. Colorectal cancers are the third most common cancers in men and women. Incidence rates declined significantly during 1992-1996 (-2.1% per year). Research suggests that these declines have been due to increased screening and polyp removal, preventing progression of polyps to invasive cancers. There were an estimated 56,300 deaths (47,700 from colon cancer, 8,600 from rectal cancer) in 2000, accounting for about 11% of all U.S. cancer deaths.

At present, surgery is the most common form of therapy for colorectal cancer, and for cancers that have not spread, it is frequently curative. Chemotherapy, or chemotherapy plus radiation, is given before or after surgery to most patients whose cancer has deeply perforated the bowel wall or has spread to the lymph nodes. A permanent colostomy (creation of an abdominal opening for elimination of body wastes) is occasionally needed for colon cancer and is infrequently required for rectal cancer. There continues to be a need for effective diagnostic and treatment modalities for colorectal cancer.

There were an estimated 164,100 new cases of lung and bronchial cancer in 2000, accounting for 14% of all U.S. cancer diagnoses. The incidence rate of lung and bronchial cancer is declining significantly in men, from a high of 86.5 per 100,000 in 1984 to 70.0 in 1996. In the 1990s, the rate of increase among women began to slow. In 1996, the incidence rate in women was 42.3 per 100,000.

Lung and bronchial cancer caused an estimated 156,900 deaths in 2000, accounting for 28% of all cancer deaths. During 1992-1996, mortality from lung cancer declined significantly among men (-1.7% per year) while rates for women were still significantly increasing (0.9% per year). Since 1987, more women have died each year of lung cancer than breast cancer, which, for over 40 years, was the major cause of cancer death in women. Decreasing lung cancer incidence and mortality rates most likely resulted from decreased smoking rates over the previous 30 years; however, decreasing smoking patterns among women lag behind those of men. Of concern, although the declines in adult tobacco use have slowed, tobacco use in youth is increasing again.

Treatment options for lung and bronchial cancer are determined by the type and stage of the cancer and include surgery, radiation therapy, and chemotherapy. For many localized cancers, surgery is usually the treatment of choice. Because the disease has usually spread by the time it is discovered, radiation therapy and chemotherapy are often needed in combination with surgery. Chemotherapy alone or combined with radiation is the treatment of choice for small cell lung cancer; on this regimen, a large percentage of patients experience remission, which in some cases is long lasting. There is however, an ongoing need for effective treatment and diagnostic approaches for lung and bronchial cancers.

An estimated 182,800 new invasive cases of breast cancer were expected to occur among women in the United States during 2000. Additionally, about 1,400 new cases of breast cancer were expected to be diagnosed in men in 2000. After increasing about 4% per year in the 1980s, breast cancer incidence rates in women have leveled off in the 1990s to about 110.6 cases per 100,000.

In the U.S. alone, there were an estimated 41,200 deaths (40,800 women, 400 men) in 2000 due to breast cancer. Breast cancer ranks second among cancer deaths in women. According to the most recent data, mortality rates declined significantly during 1992-1996 with the largest decreases in younger women, both white and black. These decreases were probably the result of earlier detection and improved treatment.

Taking into account the medical circumstances and the patients preferences, treatment of breast cancer may involve lumpectomy (local removal of the tumor) and removal of the lymph nodes under the arm; mastectomy (surgical removal of the breast) and removal of the lymph nodes under the arm; radiation therapy; chemotherapy; or hormone therapy. Often, two or more methods are used in combination. Numerous studies have shown that, for early stage disease, long-term survival rates after lumpectomy plus radiotherapy are similar to survival rates after modified radical mastectomy. Significant advances in reconstruction techniques provide several options for breast reconstruction after mastectomy. Recently, such reconstruction has been done at the same time as the mastectomy.

Local excision of ductal carcinoma in situ (DCIS) with adequate amounts of surrounding normal breast tissue may prevent the local recurrence of the DCIS. Radiation to the breast and/or tamoxifen may reduce the chance of DCIS occurring in the remaining breast tissue. This is important because DCIS, if left untreated, may develop into invasive breast cancer. Nevertheless, there are serious side effects or sequelae to these treatments. There is, therefore, a need for efficacious breast cancer treatments.

There were an estimated 23,100 new cases of ovarian cancer in the United States in 2000. It accounts for 4% of all cancers among women and ranks second among gynecologic cancers. During 1992-1996, ovarian cancer incidence rates were significantly declining. Consequent to ovarian cancer, there were an estimated 14,000 deaths in 2000. Ovarian cancer causes more deaths than any other cancer of the female reproductive system.

Surgery, radiation therapy, and chemotherapy are treatment options for ovarian cancer. Surgery usually includes the removal of one or both ovaries, the fallopian tubes (salpingo-oophorectomy), and the uterus (hysterectomy). In some very early tumors, only the involved ovary will be removed, especially in young women who wish to have children. In advanced disease, an attempt is made to remove all intra-abdominal disease to enhance the effect of chemotherapy.

There continues to be an important need for effective treatment options for ovarian cancer.

There were an estimated 28,300 new cases of pancreatic cancer in the United States in 2000. Over the past 20 years, rates of pancreatic cancer have declined in men. Rates among women have remained approximately constant but may be beginning to decline. Pancreatic cancer caused an estimated 28,200 deaths in 2000 in the United States. Over the past 20 years, there has been a slight but significant decrease in mortality rates among men (about -0.9% per year) while rates have increased slightly among women.

Surgery, radiation therapy, and chemotherapy are treatment options for pancreatic cancer. These treatment options can extend survival and/or relieve symptoms in many patients but are not likely to produce a cure for most. There is a significant need for additional therapeutic and diagnostic options for pancreatic cancer.

SUMMARY OF THE INVENTION

The present invention relates to a gene, designated 24P4C12, that has now been found to be over-expressed in the cancer(s) listed in Table I. Northern blot expression analysis of 24P4C12 gene expression in normal tissues shows a restricted expression pattern in adult tissues. The nucleotide (FIG. 2) and amino acid (FIG. 2, and FIG. 3) sequences of 24P4C12 are provided. The tissue-related profile of 24P4C12 in normal adult tissues, combined with the over-expression observed in the tissues listed in Table I, shows that 24P4C12 is aberrantly over-expressed in at least some cancers, and thus serves as a useful diagnostic, prophylactic, prognostic, and/or therapeutic target for cancers of the tissue(s) such as those listed in Table I.

The invention provides polynucleotides corresponding or complementary to all or part of the 24P4C12 genes, mRNAs, and/or coding sequences, preferably in isolated form, including polynucleotides encoding 24P4C12-related proteins and fragments of 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or more than 25 contiguous amino acids; at least 30, 35, 40, 45, 50, 55, 60, 65, 70, 80, 85, 90, 95, 100 or more than 100 contiguous amino acids of a 24P4C12-related protein, as well as the peptides/proteins themselves; DNA, RNA, DNA/RNA hybrids, and related molecules, polynucleotides or oligonucleotides complementary or having at least a 90% homology to the 24P4C12 genes or mRNA sequences or parts thereof, and polynucleotides or oligonucleotides that hybridize to the 24P4C12 genes, mRNAs, or to 24P4C12-encoding polynucleotides. Also provided are means for isolating cDNAs and the genes encoding 24P4C12. Recombinant DNA molecules containing 24P4C12 polynucleotides, cells transformed or transduced with such molecules, and host-vector systems for the expression of 24P4C12 gene products are also provided. The invention further provides antibodies that bind to 24P4C12 proteins and polypeptide fragments thereof, including polyclonal and monoclonal antibodies, murine and other mammalian antibodies, chimeric antibodies, humanized and fully human antibodies, and antibodies labeled with a detectable marker or therapeutic agent. In certain embodiments, there is a proviso that the entire nucleic acid sequence of FIG. 2 is not encoded and/or the entire amino acid sequence of FIG. 2 is not prepared. In certain embodiments, the entire nucleic acid sequence of FIG. 2 is encoded and/or the entire amino acid sequence of FIG. 2 is prepared, either of which are in respective human unit dose forms.

The invention further provides methods for detecting the presence and status of 24P4C12 polynucleotides and proteins

in various biological samples, as well as methods for identifying cells that express 24P4C12. A typical embodiment of this invention provides methods for monitoring 24P4C12 gene products in a tissue or hematology sample having or suspected of having some form of growth dysregulation such as cancer.

The invention further provides various immunogenic or therapeutic compositions and strategies for treating cancers that express 24P4C12 such as cancers of tissues listed in Table I, including therapies aimed at inhibiting the transcription, translation, processing or function of 24P4C12 as well as cancer vaccines. In one aspect, the invention provides compositions, and methods comprising them, for treating a cancer that expresses 24P4C12 in a human subject wherein the composition comprises a carrier suitable for human use and a human unit dose of one or more than one agent that inhibits the production or function of 24P4C12. Preferably, the carrier is a uniquely human carrier. In another aspect of the invention, the agent is a moiety that is immunoreactive with 24P4C12 protein. Non-limiting examples of such moieties include, but are not limited to, antibodies (such as single chain, monoclonal, polyclonal, humanized, chimeric, or human antibodies), functional equivalents thereof (whether naturally occurring or synthetic), and combinations thereof. The antibodies can be conjugated to a diagnostic or therapeutic moiety. In another aspect, the agent is a small molecule as defined herein.

In another aspect, the agent comprises one or more than one peptide which comprises a cytotoxic T lymphocyte (CTL) epitope that binds an HLA class I molecule in a human to elicit a CTL response to 24P4C12 and/or one or more than one peptide which comprises a helper T lymphocyte (HTL) epitope which binds an HLA class II molecule in a human to elicit an HTL response. The peptides of the invention may be on the same or on one or more separate polypeptide molecules. In a further aspect of the invention, the agent comprises one or more than one nucleic acid molecule that expresses one or more than one of the CTL or HTL response stimulating peptides as described above. In yet another aspect of the invention, the one or more than one nucleic acid molecule may express a moiety that is immunologically reactive with 24P4C12 as described above. The one or more than one nucleic acid molecule may also be, or encodes, a molecule that inhibits production of 24P4C12. Non-limiting examples of such molecules include, but are not limited to, those complementary to a nucleotide sequence essential for production of 24P4C12 (e.g. antisense sequences or molecules that form a triple helix with a nucleotide double helix essential for 24P4C12 production) or a ribozyme effective to lyse 24P4C12 mRNA.

Note that to determine the starting position of any peptide set forth in Tables VIII-XXI and XXII to XLIX (collectively HLA Peptide Tables) relative to its parental protein, e.g., variant 1, variant 2, etc., reference is made to three factors: the particular variant, the length of the peptide in an HLA Peptide Table, and the Search Peptides in Table VII. Generally, a unique Search Peptide is used to obtain HLA peptides of a particular for a particular variant. The position of each Search Peptide relative to its respective parent molecule is listed in Table VII. Accordingly, if a Search Peptide begins at position "X", one must add the value "X-1" to each position in Tables VIII-XXI and XXII to XLIX to obtain the actual position of the HLA peptides in their parental molecule. For example, if a particular Search Peptide begins at position 150 of its parental molecule, one must add 150-1, i.e., 149 to each HLA peptide amino acid position to calculate the position of that amino acid in the parent molecule.

One embodiment of the invention comprises an HLA peptide, that occurs at least twice in Tables VIII-XXI and XXII to XLIX collectively, or an oligonucleotide that encodes the HLA peptide. Another embodiment of the invention comprises an HLA peptide that occurs at least once in Tables VIII-XXI and at least once in tables XXII to XLIX, or an oligonucleotide that encodes the HLA peptide.

Another embodiment of the invention is antibody epitopes, which comprise a peptide regions, or an oligonucleotide encoding the peptide region, that has one two, three, four, or five of the following characteristics:

i) a peptide region of at least 5 amino acids of a particular peptide of FIG. 3, in any whole number increment up to the full length of that protein in FIG. 3, that includes an amino acid position having a value equal to or greater than 0.5, 0.6, 0.7, 0.8, 0.9, or having a value equal to 1.0, in the Hydrophilicity profile of FIG. 5;

ii) a peptide region of at least 5 amino acids of a particular peptide of FIG. 3, in any whole number increment up to the full length of that protein in FIG. 3, that includes an amino acid position having a value equal to or less than 0.5, 0.4, 0.3, 0.2, 0.1, or having a value equal to 0.0, in the Hydropathicity profile of FIG. 6;

iii) a peptide region of at least 5 amino acids of a particular peptide of FIG. 3, in any whole number increment up to the full length of that protein in FIG. 3, that includes an amino acid position having a value equal to or greater than 0.5, 0.6, 0.7, 0.8, 0.9, or having a value equal to 1.0, in the Percent Accessible Residues profile of FIG. 7;

iv) a peptide region of at least 5 amino acids of a particular peptide of FIG. 3, in any whole number increment up to the full length of that protein in FIG. 3, that includes an amino acid position having a value equal to or greater than 0.5, 0.6, 0.7, 0.8, 0.9, or having a value equal to 1.0, in the Average Flexibility profile of FIG. 8; or

v) a peptide region of at least 5 amino acids of a particular peptide of FIG. 3, in any whole number increment up to the full length of that protein in FIG. 3, that includes an amino acid position having a value equal to or greater than 0.5, 0.6, 0.7, 0.8, 0.9, or having a value equal to 1.0, in the Beta-turn profile of FIG. 9.

BRIEF DESCRIPTION OF THE FIGURES

FIG. 1. The 24P4C12 SSH sequence of 160 nucleotides.

FIG. 2. A) The cDNA and amino acid sequence of 24P4C12 variant 1 (also called "24P4C12 v.1" or "24P4C12 variant 1") is shown in FIG. 2A. The start methionine is underlined. The open reading frame extends from nucleic acid 6-2138 including the stop codon.

B) The cDNA and amino acid sequence of 24P4C12 variant 2 (also called "24P4C12 v.2") is shown in FIG. 2B. The codon for the start methionine is underlined. The open reading frame extends from nucleic acid 6-2138 including the stop codon.

C) The cDNA and amino acid sequence of 24P4C12 variant 3 (also called "24P4C12 v.3") is shown in FIG. 2C. The codon for the start methionine is underlined. The open reading frame extends from nucleic acid 6-2138 including the stop codon.

D) The cDNA and amino acid sequence of 24P4C12 variant 4 (also called "24P4C12 v.4") is shown in FIG. 2D. The codon for the start methionine is underlined. The open reading frame extends from nucleic acid 6-2138 including the stop codon.

E) The cDNA and amino acid sequence of 24P4C12 variant 5 (also called "24P4C12 v.5") is shown in FIG. 2E. The codon

for the start methionine is underlined. The open reading frame extends from nucleic acid 6-2138 including the stop codon.

F) The cDNA and amino acid sequence of 24P4C12 variant 6 (also called "24P4C12 v.6") is shown in FIG. 2F. The codon for the start methionine is underlined. The open reading frame extends from nucleic acid 6-2138 including the stop codon.

G) The cDNA and amino acid sequence of 24P4C12 variant 7 (also called "24P4C12 v.7") is shown in FIG. 2G. The codon for the start methionine is underlined. The open reading frame extends from nucleic acid 6-1802 including the stop codon.

H) The cDNA and amino acid sequence of 24P4C12 variant 8 (also called "24P4C12 v.8") is shown in FIG. 2H. The codon for the start methionine is underlined. The open reading frame extends from nucleic acid 6-2174 including the stop codon.

I) The cDNA and amino acid sequence of 24P4C12 variant 9 (also called "24P4C12 v.9") is shown in FIG. 2I. The codon for the start methionine is underlined. The open reading frame extends from nucleic acid 6-2144 including the stop codon.

FIG. 3.

A) Amino acid sequence of 24P4C12 v.1 is shown in FIG. 3A; it has 710 amino acids.

B) The amino acid sequence of 24P4C12 v.3 is shown in FIG. 3B; it has 710 amino acids.

C) The amino acid sequence of 24P4C12 v.5 is shown in FIG. 3C; it has 710 amino acids.

D) The amino acid sequence of 24P4C12 v.6 is shown in FIG. 3D; it has 710 amino acids.

E) The amino acid sequence of 24P4C12 v.7 is shown in FIG. 3E; it has 598 amino acids.

F) The amino acid sequence of 24P4C12 v.8 is shown in FIG. 3F; it has 722 amino acids.

G) The amino acid sequence of 24P4C12 v.9 is shown in FIG. 3G; it has 712 amino acids. As used herein, a reference to 24P4C12 includes all variants thereof, including those shown in FIGS. 2, 3, 10, and 11, unless the context clearly indicates otherwise.

FIG. 4. Alignment of 24P4C12 with human choline transporter-like protein 4 (CTL4) (gil 14249468).

FIG. 5. Hydrophilicity amino acid profile of 24P4C12 determined by computer algorithm sequence analysis using the method of Hopp and Woods (Hopp T. P., Woods K. R., 1981. Proc. Natl. Acad. Sci. U.S.A. 78:3824-3828) accessed on the ProtScale website located on the World Wide Web at (.expasy.ch/cgi-bin/protscale.pl) through the ExPasy molecular biology server.

FIG. 6. Hydropathicity amino acid profile of 24P4C12 determined by computer algorithm sequence analysis using the method of Kyte and Doolittle (Kyte J., Doolittle R. F., 1982. J. Mol. Biol. 157:105-132) accessed on the ProtScale website located on the World Wide Web at (.expasy.ch/cgi-bin/protscale.pl) through the ExPasy molecular biology server.

FIG. 7. Percent accessible residues amino acid profile of 24P4C12 determined by computer algorithm sequence analysis using the method of Janin (Janin J., 1979 Nature 277:491-492) accessed on the ProtScale website located on the World Wide Web at (.expasy.ch/cgi-bin/protscale.pl) through the ExPasy molecular biology server.

FIG. 8. Average flexibility amino acid profile of 24P4C12 determined by computer algorithm sequence analysis using the method of Bhaskaran and Ponnuswamy (Bhaskaran R., and Ponnuswamy P. K., 1988. Int. J. Pept. Protein Res. 32:242-255) accessed on the ProtScale website located on the World Wide Web at (.expasy.ch/cgi-bin/protscale.pl) through the ExPasy molecular biology server.

FIG. 9. Beta-turn amino acid profile of 24P4C12 determined by computer algorithm sequence analysis using the method of Deleage and Roux (Deleage, G., Roux B. 1987 Protein Engineering 1:289-294) accessed on the ProtScale website located on the World Wide Web at (expasy.ch/cgi-bin/protscale.pl) through the ExPasy molecular biology server.

FIG. 10. Schematic alignment of SNP variants of 24P4C12. Variants 24P4C12 v.2 through v.6 are variants with single nucleotide differences. Though these SNP variants are shown separately, they could also occur in any combinations and in any transcript variants that contains the base pairs. Numbers correspond to those of 24P4C12 v.1. Black box shows the same sequence as 24P4C12 v.1. SNPs are indicated above the box.

FIG. 11. Schematic alignment of protein variants of 24P4C12. Protein variants correspond to nucleotide variants. Nucleotide variants 24P4C12 v.2, v.4 in FIG. 10 code for the same protein as 24P4C12 v.1. Nucleotide variants 24P4C12 v.7, v.8 and v.9 are splice variants of v.1, as shown in FIG. 12. Single amino acid differences were indicated above the boxes. Black boxes represent the same sequence as 24P4C12 v.1. Numbers underneath the box correspond to 24P4C12 v.1.

FIG. 12. Exon compositions of transcript variants of 24P4C12. Variant 24P4C12 v.7, v.8 and v.9 are transcript variants of 24P4C12 v.1. Variant 24P4C12 v.7 does not have exons 10 and 11 of variant 24P4C12 v.1. Variant 24P4C12 v.8 extended 36 bp at the 3' end of exon 20 of variant 24P4C12 v.1. Variant 24P4C12 v.9 had a longer exon 12 and shorter exon 13 as compared to variant 24P4C12 v.1. Numbers in “()” underneath the boxes correspond to those of 24P4C12 v.1. Lengths of introns and exons are not proportional.

FIG. 13. Secondary structure and transmembrane domains prediction for 24P4C12 protein variant 1 (SEQ ID NO: 112). A: The secondary structure of 24P4C12 protein variant 1 was predicted using the HNN—Hierarchical Neural Network method (Guermeur, 1997, http://pbil.ibcp.fr/cgi-bin/npsa_automat.pl?page=npsa_nn.html), accessed from the ExPasy molecular biology server (<http://www.expasy.ch/tools/>). This method predicts the presence and location of alpha helices, extended strands, and random coils from the primary protein sequence. The percent of the protein in a given secondary structure is also listed. B: Schematic representation of the probability of existence of transmembrane regions and orientation of 24P4C12 variant 1 based on the TMPred algorithm of Hofmann and Stoffel which utilizes TMBASE (K. Hofmann, W. Stoffel. TMBASE—A database of membrane spanning protein segments Biol. Chem. Hoppe-Seyler 374:166, 1993). C: Schematic representation of the probability of the existence of transmembrane regions and the extracellular and intracellular orientation of 24P4C12 variant 1 based on the TMHMM algorithm of Sonnhammer, von Heijne, and Krogh (Erik L. L. Sonnhammer, Gunnar von Heijne, and Anders Krogh: A hidden Markov model for predicting transmembrane helices in protein sequences. In Proc. of Sixth Int. Conf. on Intelligent Systems for Molecular Biology, p 175-182 Ed J. Glasgow, T. Littlejohn, F. Major, R. Lathrop, D. Sankoff, and C. Sensen Menlo Park, Calif.: AAAI Press, 1998). The TMPred and TMHMM algorithms are accessed from the ExPasy molecular biology server (<http://www.expasy.ch/tools/>).

FIG. 14. 24P4C12 Expression by RT-PCR. First strand cDNA was generated from vital pool 1 (kidney, liver and lung), vital pool 2 (colon, pancreas and stomach), a pool of prostate cancer xenografts (LAPC-4AD, LAPC-4AI, LAPC-9AD and LAPC-9AI), prostate cancer pool, bladder cancer pool, kidney cancer pool, colon cancer pool, ovary cancer

pool, breast cancer pool, and cancer metastasis pool. Normalization was performed by PCR using primers to actin. Semi-quantitative PCR, using primers to 24P4C12, was performed at 26 and 30 cycles of amplification. Results show strong expression of 24P4C12 in prostate cancer pool and ovary cancer pool. Expression was also detected in prostate cancer xenografts, bladder cancer pool, kidney cancer pool, colon cancer pool, breast cancer pool, cancer metastasis pool, vital pool 1, and vital pool 2.

FIG. 15. Expression of 24P4C12 in normal tissues. Two multiple tissue northern blots (Clontech) both with 2 ug of mRNA/lane were probed with the 24P4C12 sequence. Size standards in kilobases (kb) are indicated on the side. Results show expression of 24P4C12 in prostate, kidney and colon. Lower expression is detected in pancreas, lung and placenta amongst all 16 normal tissues tested.

FIG. 16. Expression of 24P4C12 in Prostate Cancer Xenografts and Cell Lines. RNA was extracted from a panel of cell lines and prostate cancer xenografts (PrEC, LAPC-4AD, LAPC-4AI, LAPC-9AD, LAPC-9AI, LNCaP, PC-3, DU145, TsuPr, and LAPC-4CL). Northern blot with 10 ug of total RNA/lane was probed with 24P4C12 SSH sequence. Size standards in kilobases (kb) are indicated on the side. The 24P4C12 transcript was detected in LAPC-4AD, LAPC-4AI, LAPC-9AD, LAPC-9AI, LNCaP, and LAPC-4 CL.

FIG. 17. Expression of 24P4C12 in Patient Cancer Specimens and Normal Tissues. RNA was extracted from a pool of prostate cancer specimens, bladder cancer specimens, colon cancer specimens, ovary cancer specimens, breast cancer specimens and cancer metastasis specimens, as well as from normal prostate (NP), normal bladder (NB), normal kidney (NK), and normal colon (NC). Northern blot with 10 ug of total RNA/lane was probed with 24P4C12 SSH sequence. Size standards in kilobases (kb) are indicated on the side. Strong expression of 24P4C12 transcript was detected in the patient cancer pool specimens, and in normal prostate but not in the other normal tissues tested.

FIG. 18. Expression of 24P4C12 in Prostate Cancer Patient Specimens. RNA was extracted from normal prostate (N), prostate cancer patient tumors (T) and their matched normal adjacent tissues (Nat). Northern blots with 10 ug of total RNA were probed with the 24P4C12 SSH fragment. Size standards in kilobases are on the side. Results show expression of 24P4C12 in normal prostate and all prostate patient tumors tested.

FIG. 19. Expression of 24P4C12 in Colon Cancer Patient Specimens. RNA was extracted from colon cancer cell lines (CL: Colo 205, LoVo, and SK—CO—), normal colon (N), colon cancer patient tumors (T) and their matched normal adjacent tissues (Nat). Northern blots with 10 ug of total RNA were probed with the 24P4C12 SSH fragment. Size standards in kilobases are on the side. Results show expression of 24P4C12 in normal colon and all colon patient tumors tested. Expression was detected in the cell lines Colo 205 and SK—CO—, but not in LoVo.

FIG. 20. Expression of 24P4C12 in Lung Cancer Patient Specimens. RNA was extracted from lung cancer cell lines (CL: CALU-1, A427, NCI-H82, NCI-H146), normal lung (N), lung cancer patient tumors (T) and their matched normal adjacent tissues (Nat). Northern blots with 10 ug of total RNA were probed with the 24P4C12 SSH fragment. Size standards in kilobases are on the side. Results show expression of 24P4C12 in lung patient tumors tested, but not in normal lung. Expression was also detected in CALU-1, but not in the other cell lines A427, NCI-H82, and NCI-H146.

FIG. 21. Expression of 24P4C12 in breast and stomach human cancer specimens. Expression of 24P4C12 was

assayed in a panel of human stomach and breast cancers (T) and their respective matched normal tissues (N) on RNA dot blots. 24P4C12 expression was seen in both stomach and breast cancers. The expression detected in normal adjacent tissues (isolated from diseased tissues) but not in normal tissues (isolated from healthy donors) may indicate that these tissues are not fully normal and that 24P4C12 may be expressed in early stage tumors.

FIG. 22. 24P4C12 Expression in a large panel of Patient Cancer Specimens. First strand cDNA was prepared from a panel of ovary patient cancer specimens (A), uterus patient cancer specimens (B), prostate cancer specimens (C), bladder cancer patient specimens (D), lung cancer patient specimens (E), pancreas cancer patient specimens (F), colon cancer specimens (G), and kidney cancer specimens (H). Normalization was performed by PCR using primers to actin. Semi-quantitative PCR, using primers to 24P4C12, was performed at 26 and 30 cycles of amplification. Samples were run on an agarose gel, and PCR products were quantitated using the Alphasampler software. Expression was recorded as absent, low, medium or strong. Results show expression of 24P4C12 in the majority of patient cancer specimens tested, 73.3% of ovary patient cancer specimens, 83.3% of uterus patient cancer specimens, 95.0% of prostate cancer specimens, 61.1% of bladder cancer patient specimens, 80.6% of lung cancer patient specimens, 87.5% of pancreas cancer patient specimens, 87.5% of colon cancer specimens, 68.4% of clear cell renal carcinoma, 100% of papillary renal cell carcinoma.

FIG. 23. 24P4C12 expression in transduced cells. PC3 prostate cancer cells, NIH-3T3 mouse cells and 300.19 mouse cells were transduced with 24P4C12.pSRa retroviral vector. Cells were selected in neomycin for the generation of stable cell lines. RNA was extracted following selection in neomycin. Northern blots with 10 ug of total RNA were probed with the 24P4C12 SSH fragment. Results show strong expression of 24P4C12 in 24P4C12.pSRa transduced PC3, 3T3 and 300.19 cells, but not in the control cells transduced with the parental pSRa construct.

FIG. 24. Expression of 24P4C12 in 293T cells. 293T cell were transiently transfected with either pcDNA3.1 Myc-His tagged expression vector, the pSR $\square$ expression vector each encoding the 24P4C12 variant 1 cDNA or a control neo vector. Cells were harvested 2 days later and analyzed by Western blot with anti-24P4C12 pAb (A) or by Flow cytometry (B) on fixed and permeabilized 293T cells with either the anti-24P4C12 pAb or anti-His pAb followed by a PE-conjugated anti-rabbit IgG secondary Ab. Shown is expression of the monomeric and aggregated forms of 24P4C12 by Western blot and a fluorescent shift of 24P4C12-293T cells compared to control neo cells when stained with the anti-24P4C12 and anti-His pAbs which are directed to the intracellular NH3 and COOH termini, respectively.

FIG. 25. Expression and detection of 24P4C12 in stably transduced PC3 cells. PC3 cells were infected with retrovirus encoding the 24P4C12 variant 1 cDNA and stably transduced cells were derived by G418 selection. Cells were then analyzed by Western blot (A) or immunohistochemistry (B) with anti-24P4C12 pAb. Shown with an arrow on the Western blot is expression of a ~94 kD band representing 24P4C12 expressed in PC3-24P4C12 cells but not in control neo cells. Immunohistochemical analysis shows specific staining of 24P4C12-PC3 cells and not PC3-neo cells which is competed away competitor peptide to which the pAb was derived.

FIG. 26. Expression of recombinant 24P4C12 antigens in 293T cells. 293T cells were transiently transfected with Tag5 His-tagged expression vectors encoding either amino acids 59-227 or 319-453 of 24P4C12 variant 1 or a control vector.

2 days later supernatants were collected and cells harvested and lysed. Supernatants and lysates were then subjected to Western blot analysis using an anti-His pAb. Shown is expression of the recombinant Tag5 59-227 protein in both the supernatant and lysate and the Tag5 319-453 protein in the cell lysate. These proteins are purified and used as antigens for generation of 24P4C12-specific antibodies.

FIG. 27. Monoclonal antibodies detect 24P4C12 protein expression in 293T cells by flow cytometry. 293T cells were transfected with either pcDNA 3.1 His-tagged expression vector for 24P4C12 or a control neo vector and harvested 2 days later. Cells were fixed, permeabilized, and stained with a 1:2 dilution of supernatants of the indicated hybridomas generated from mice immunized with 300.19-24P4C12 cells or with anti-His pAb. Cells were then stained with a PE-conjugated secondary Ab and analyzed by flow cytometry. Shown is a fluorescent shift of 293T-24P4C12 cells but not control neo cells demonstrating specific recognition of 24P4C12 protein by the hybridoma supernatants.

FIG. 28. Shows expression of 24P4C12 Enhances Proliferation. PC3 and 3T3 were grown overnight in low FBS. Cells were then incubated in low or 10% FBS as indicated. Proliferation was measured by Alamar Blue.

FIG. 29. Detection of 24P4C12 protein by immunohistochemistry in prostate cancer patient specimens. Prostate adenocarcinoma tissue and its matched normal adjacent tissue were obtained from prostate cancer patients. The results showed strong expression of 24P4C12 in the tumor cells and normal epithelium of the prostate cancer patients' tissue (panels (A) low grade prostate adenocarcinoma, (B) high grade prostate adenocarcinoma, (C) normal tissue adjacent to tumor). The expression was detected mostly around the cell membrane indicating that 24P4C12 is membrane associated in prostate tissues.

FIG. 30. Detection of 24P4C12 protein by immunohistochemistry in various cancer patient specimens. Tissue was obtained from patients with colon adenocarcinoma, breast ductal carcinoma, lung adenocarcinoma, bladder transitional cell carcinoma, renal clear cell carcinoma and pancreatic adenocarcinoma. The results showed expression of 24P4C12 in the tumor cells of the cancer patients' tissue (panel (A) colon adenocarcinoma, (B) lung adenocarcinoma, (C) breast ductal carcinoma, (D) bladder transitional carcinoma, (E) renal clear cell carcinoma, (F) pancreatic adenocarcinoma).

FIG. 31. Shows 24P4C12 Enhances Tumor Growth in SCID Mice. 1×10^6 PC3-24P4C12 cells were mixed with Matrigel and injected on the right and left subcutaneous flanks of 4 male SCID mice per group. Each data point represents mean tumor volume ($n=8$).

FIG. 32. Shows 24P4C12 Enhances Tumor Growth in SCID Mice. 1×10^6 3T3-24P4C12 cells were mixed with Matrigel and injected on the right subcutaneous flanks of 7 male SCID mice per group. Each data point represents mean tumor volume ($n=6$).

DETAILED DESCRIPTION OF THE INVENTION

Outline of Sections

I.) Definitions

II.) 24P4C12 Polynucleotides

II.A.) Uses of 24P4C12 Polynucleotides

II.A.1.) Monitoring of Genetic Abnormalities

II.A.2.) Antisense Embodiments

II.A.3.) Primers and Primer Pairs

II.A.4.) Isolation of 24P4C12-Encoding Nucleic Acid Molecules

- II.A.5.) Recombinant Nucleic Acid Molecules and Host-Vector Systems
- III.) 24P4C12-related Proteins
 - III.A.) Motif-bearing Protein Embodiments
 - III.B.) Expression of 24P4C12-related Proteins
 - III.C.) Modifications of 24P4C12-related Proteins
 - III.D.) Uses of 24P4C12-related Proteins
- IV.) 24P4C12 Antibodies
- V.) 24P4C12 Cellular Immune Responses
- VI.) 24P4C12 Transgenic Animals
- VII.) Methods for the Detection of 24P4C12
- VIII.) Methods for Monitoring the Status of 24P4C12-related Genes and Their Products
- IX.) Identification of Molecules That Interact With 24P4C12
 - X.) Therapeutic Methods and Compositions
 - X.A.) Anti-Cancer Vaccines
 - X.B.) 24P4C12 as a Target for Antibody-Based Therapy
 - X.C.) 24P4C12 as a Target for Cellular Immune Responses
 - X.C.1. Minigene Vaccines
 - X.C.2. Combinations of CTL Peptides with Helper Peptides
 - X.C.3. Combinations of CTL Peptides with T Cell Priming Agents
 - X.C.4. Vaccine Compositions Comprising DC Pulsed with CTL and/or HTL Peptides
 - X.D.) Adoptive Immunotherapy
 - X.E.) Administration of Vaccines for Therapeutic or Prophylactic Purposes
 - XI.) Diagnostic and Prognostic Embodiments of 24P4C12
 - XII.) Inhibition of 24P4C12 Protein Function
 - XII.A.) Inhibition of 24P4C12 With Intracellular Antibodies
 - XII.B.) Inhibition of 24P4C12 with Recombinant Proteins
 - XII.C.) Inhibition of 24P4C12 Transcription or Translation
 - XII.D.) General Considerations for Therapeutic Strategies
 - XIII.) Identification, Characterization and Use of Modulators of 24P4C12
 - XIV.) KITS/Articles of Manufacture

I.) DEFINITIONS

Unless otherwise defined, all terms of art, notations and other scientific terms or terminology used herein are intended to have the meanings commonly understood by those of skill in the art to which this invention pertains. In some cases, terms with commonly understood meanings are defined herein for clarity and/or for ready reference, and the inclusion of such definitions herein should not necessarily be construed to represent a substantial difference over what is generally understood in the art. Many of the techniques and procedures described or referenced herein are well understood and commonly employed using conventional methodology by those skilled in the art, such as, for example, the widely utilized molecular cloning methodologies described in Sambrook et al., *Molecular Cloning: A Laboratory Manual* 2nd. edition (1989) Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. As appropriate, procedures involving the use of commercially available kits and reagents are generally carried out in accordance with manufacturer defined protocols and/or parameters unless otherwise noted.

The terms “advanced prostate cancer”, “locally advanced prostate cancer”, “advanced disease” and “locally advanced disease” mean prostate cancers that have extended through the prostate capsule, and are meant to include stage C disease under the American Urological Association (AUA) system, stage C1-C2 disease under the Whitmore-Jewett system, and

stage T3-T4 and N+ disease under the TNM (tumor, node, metastasis) system. In general, surgery is not recommended for patients with locally advanced disease, and these patients have substantially less favorable outcomes compared to patients having clinically localized (organ-confined) prostate cancer. Locally advanced disease is clinically identified by palpable evidence of induration beyond the lateral border of the prostate, or asymmetry or induration above the prostate base. Locally advanced prostate cancer is presently diagnosed pathologically following radical prostatectomy if the tumor invades or penetrates the prostatic capsule, extends into the surgical margin, or invades the seminal vesicles.

“Altering the native glycosylation pattern” is intended for purposes herein to mean deleting one or more carbohydrate moieties found in native sequence 24P4C12 (either by removing the underlying glycosylation site or by deleting the glycosylation by chemical and/or enzymatic means), and/or adding one or more glycosylation sites that are not present in the native sequence 24P4C12. In addition, the phrase includes qualitative changes in the glycosylation of the native proteins, involving a change in the nature and proportions of the various carbohydrate moieties present.

The term “analog” refers to a molecule which is structurally similar or shares similar or corresponding attributes with another molecule (e.g. a 24P4C12-related protein). For example, an analog of a 24P4C12 protein can be specifically bound by an antibody or T cell that specifically binds to 24P4C12.

The term “antibody” is used in the broadest sense. Therefore, an “antibody” can be naturally occurring or man-made such as monoclonal antibodies produced by conventional hybridoma technology. Anti-24P4C12 antibodies comprise monoclonal and polyclonal antibodies as well as fragments containing the antigen-binding domain and/or one or more complementarity determining regions of these antibodies.

An “antibody fragment” is defined as at least a portion of the variable region of the immunoglobulin molecule that binds to its target, i.e., the antigen-binding region. In one embodiment it specifically covers single anti-24P4C12 antibodies and clones thereof (including agonist, antagonist and neutralizing antibodies) and anti-24P4C12 antibody compositions with polyepitopic specificity.

The term “codon optimized sequences” refers to nucleotide sequences that have been optimized for a particular host species by replacing any codons having a usage frequency of less than about 20%. Nucleotide sequences that have been optimized for expression in a given host species by elimination of spurious polyadenylation sequences, elimination of exon/intron splicing signals, elimination of transposon-like repeats and/or optimization of GC content in addition to codon optimization are referred to herein as an “expression enhanced sequences.”

A “combinatorial library” is a collection of diverse chemical compounds generated by either chemical synthesis or biological synthesis by combining a number of chemical “building blocks” such as reagents. For example, a linear combinatorial chemical library, such as a polypeptide (e.g., mutin) library, is formed by combining a set of chemical building blocks called amino acids in every possible way for a given compound length (i.e., the number of amino acids in a polypeptide compound). Numerous chemical compounds are synthesized through such combinatorial mixing of chemical building blocks (Gallop et al., *J. Med. Chem.* 37(9): 1233-1251 (1994)).

Preparation and screening of combinatorial libraries is well known to those of skill in the art. Such combinatorial chemical libraries include, but are not limited to, peptide libraries

(see, e.g., U.S. Pat. No. 5,010,175, Furka, Pept. Prot. Res. 37:487-493 (1991), Houghton et al., Nature, 354:84-88 (1991)), peptoids (PCT Publication No WO 91/19735), encoded peptides (PCT Publication WO 93/20242), random bio-oligomers (PCT Publication WO 92/00091), benzodiazepines (U.S. Pat. No. 5,288,514), diversomers such as hydantoins, benzodiazepines and dipeptides (Hobbs et al., Proc. Nat. Acad. Sol. USA 90:6909-6913 (1993)), vinylogous polypeptides (Hagihara et al., J. Amer. Chem. Soc. 114:6568 (1992)), nonpeptidal peptidomimetics with a Beta-D-Glucose scaffolding (Hirschmann et al., J. Amer. Chem. Soc. 114:9217-9218 (1992)), analogous organic syntheses of small compound libraries (Chen et al., J. Amer. Chem. Soc. 116:2661 (1994)), oligocarbamates (Cho, et al., Science 261:1303 (1993)), and/or peptidyl phosphonates (Campbell et al., J. Org. Chem. 59:658 (1994)). See, generally, Gordon et al., J. Med. Chem. 37:1385 (1994), nucleic acid libraries (see, e.g., Stratagene, Corp.), peptide nucleic acid libraries (see, e.g., U.S. Pat. No. 5,539,083), antibody libraries (see, e.g., Vaughn et al., Nature Biotechnology 14(3): 309-314 (1996), and PCT/US96/10287), carbohydrate libraries (see, e.g., Liang et al., Science 274:1520-1522 (1996), and U.S. Pat. No. 5,593,853), and small organic molecule libraries (see, e.g., benzodiazepines, Baum, C&EN, January 18, page 33 (1993); isoprenoids, U.S. Pat. No. 5,569,588; thiazolidinones and metathiazanones, U.S. Pat. No. 5,549,974; pyrrolidines, U.S. Pat. Nos. 5,525,735 and 5,519,134; morpholino compounds, U.S. Pat. No. 5,506,337; benzodiazepines, U.S. Pat. No. 5,288,514; and the like).

Devices for the preparation of combinatorial libraries are commercially available (see, e.g., 357 NIPS, 390 NIPS, Advanced Chem Tech, Louisville Ky.; Symphony, Rainin, Woburn, Mass.; 433A, Applied Biosystems, Foster City, Calif.; 9050, Plus, Millipore, Bedford, NIA). A number of well-known robotic systems have also been developed for solution phase chemistries. These systems include automated workstations such as the automated synthesis apparatus developed by Takeda Chemical Industries, LTD. (Osaka, Japan) and many robotic systems utilizing robotic arms (Zymate H, Zymark Corporation, Hopkinton, Mass.; Orca, Hewlett-Packard, Palo Alto, Calif.), which mimic the manual synthetic operations performed by a chemist. Any of the above devices are suitable for use with the present invention. The nature and implementation of modifications to these devices (if any) so that they can operate as discussed herein will be apparent to persons skilled in the relevant art. In addition, numerous combinatorial libraries are themselves commercially available (see, e.g., ComGenex, Princeton, N.J.; Asinex, Moscow, R U; Tripos, Inc., St. Louis, Mo.; ChemStar, Ltd, Moscow, R U; 3D Pharmaceuticals, Exton, Pa.; Martek Biosciences, Columbia, Md.; etc.).

The term "cytotoxic agent" refers to a substance that inhibits or prevents the expression activity of cells, function of cells and/or causes destruction of cells. The term is intended to include radioactive isotopes chemotherapeutic agents, and toxins such as small molecule toxins or enzymatically active toxins of bacterial, fungal, plant or animal origin, including fragments and/or variants thereof. Examples of cytotoxic agents include, but are not limited to auristatins, auromycins, maytansinoids, yttrium, bismuth, ricin, ricin A-chain, combrestatin, duocarmycins, dolostatins, doxorubicin, daunorubicin, taxol, cisplatin, cc1065, ethidium bromide, mitomycin, etoposide, tenoposide, vincristine, vinblastine, colchicine, dihydroxy anthracin dione, actinomycin, diphtheria toxin, *Pseudomonas* exotoxin (PE) A, PE40, abrin, abrin A chain, modeccin A chain, alpha-sarcin, gelonin, mitogellin, retsictocin, phenomycin, enomycin, curicin, croton, calicheamicin,

Sapaonaria officinalis inhibitor, and glucocorticoid and other chemotherapeutic agents, as well as radioisotopes such as At²¹¹, I¹³¹, I¹²⁵, Y⁹⁰, Re¹⁸⁶, Re¹⁸⁸, Sm¹⁵³, Bi^{212 or 213}, P³² and radioactive isotopes of Lu including Lu¹⁷⁷. Antibodies may also be conjugated to an anti-cancer pro-drug activating enzyme capable of converting the pro-drug to its active form.

The "gene product" is sometimes referred to herein as a protein or mRNA. For example, a "gene product of the invention" is sometimes referred to herein as a "cancer amino acid sequence", "cancer protein", "protein of a cancer listed in Table I", a "cancer mRNA", "mRNA of a cancer listed in Table I", etc. In one embodiment, the cancer protein is encoded by a nucleic acid of FIG. 2. The cancer protein can be a fragment, or alternatively, be the full-length protein to the fragment encoded by the nucleic acids of FIG. 2. In one embodiment, a cancer amino acid sequence is used to determine sequence identity or similarity. In another embodiment, the sequences are naturally occurring allelic variants of a protein encoded by a nucleic acid of FIG. 2. In another embodiment, the sequences are sequence variants as further described herein.

"High throughput screening" assays for the presence, absence, quantification, or other properties of particular nucleic acids or protein products are well known to those of skill in the art. Similarly, binding assays and reporter gene assays are similarly well known. Thus, e.g., U.S. Pat. No. 5,559,410 discloses high throughput screening methods for proteins; U.S. Pat. No. 5,585,639 discloses high throughput screening methods for nucleic acid binding (i.e., in arrays); while U.S. Pat. Nos. 5,576,220 and 5,541,061 disclose high throughput methods of screening for ligand/antibody binding.

In addition, high throughput screening systems are commercially available (see, e.g., Amersham Biosciences, Piscataway, N.J.; Zymark Corp., Hopkinton, Mass.; Air Technical Industries, Mentor, Ohio; Beckman Instruments, Inc. Fullerton, Calif.; Precision Systems, Inc., Natick, Mass.; etc.). These systems typically automate entire procedures, including all sample and reagent pipetting, liquid dispensing, timed incubations, and final readings of the microplate in detector(s) appropriate for the assay. These configurable systems provide high throughput and rapid start up as well as a high degree of flexibility and customization. The manufacturers of such systems provide detailed protocols for various high throughput systems. Thus, e.g., Zymark Corp. provides technical bulletins describing screening systems for detecting the modulation of gene transcription, ligand binding, and the like.

The term "homolog" refers to a molecule which exhibits homology to another molecule, by for example, having sequences of chemical residues that are the same or similar at corresponding positions.

"Human Leukocyte Antigen" or "HLA" is a human class I or class II Major Histocompatibility Complex (MHC) protein (see, e.g., Stites, et al., IMMUNOLOGY, 8TH Ed., Lange Publishing, Los Altos, Calif. (1994)).

The terms "hybridize", "hybridizing", "hybridizes" and the like, used in the context of polynucleotides, are meant to refer to conventional hybridization conditions, preferably such as hybridization in 50% formamide/6xSSC/0.1% SDS/100 µg/ml ssDNA, in which temperatures for hybridization are above 37 degrees C. and temperatures for washing in 0.1xSSC/0.1% SDS are above 55 degrees C.

The phrases "isolated" or 'biologically pure' refer to material which is substantially or essentially free from components which normally accompany the material as it is found in its native state. Thus, isolated peptides in accordance with the

invention preferably do not contain materials normally associated with the peptides in their in situ environment. For example, a polynucleotide is said to be "isolated" when it is substantially separated from contaminant polynucleotides that correspond or are complementary to genes other than the 24P4C12 genes or that encode polypeptides other than 24P4C12 gene product or fragments thereof. A skilled artisan can readily employ nucleic acid isolation procedures to obtain an isolated 24P4C12 polynucleotide. A protein is said to be "isolated," for example, when physical, mechanical or chemical methods are employed to remove the 24P4C12 proteins from cellular constituents that are normally associated with the protein. A skilled artisan can readily employ standard purification methods to obtain an isolated 24P4C12 protein. Alternatively, an isolated protein can be prepared by chemical means.

The term "mammal" refers to any organism classified as a mammal, including mice, rats, rabbits, dogs, cats, cows, horses and humans. In one embodiment of the invention, the mammal is a mouse. In another embodiment of the invention, the mammal is a human.

The terms "metastatic prostate cancer" and "metastatic disease" mean prostate cancers that have spread to regional lymph nodes or to distant sites, and are meant to include stage D disease under the AUA system and stage T_xN_xM₊ under the TNM system. As is the case with locally advanced prostate cancer, surgery is generally not indicated for patients with metastatic disease, and hormonal (androgen ablation) therapy is a preferred treatment modality. Patients with metastatic prostate cancer eventually develop an androgen-refractory state within 12 to 18 months of treatment initiation. Approximately half of these androgen-refractory patients die within 6 months after developing that status. The most common site for prostate cancer metastasis is bone. Prostate cancer bone metastases are often osteoblastic rather than osteolytic (i.e., resulting in net bone formation). Bone metastases are found most frequently in the spine, followed by the femur, pelvis, rib cage, skull and humerus. Other common sites for metastasis include lymph nodes, lung, liver and brain. Metastatic prostate cancer is typically diagnosed by open or laparoscopic pelvic lymphadenectomy, whole body radionuclide scans, skeletal radiography, and/or bone lesion biopsy.

The term "modulator" or "test compound" or "drug candidate" or grammatical equivalents as used herein describe any molecule, e.g., protein, oligopeptide, small organic molecule, polysaccharide, polynucleotide, etc., to be tested for the capacity to directly or indirectly alter the cancer phenotype or the expression of a cancer sequence, e.g., a nucleic acid or protein sequences, or effects of cancer sequences (e.g., signaling, gene expression, protein interaction, etc.) In one aspect, a modulator will neutralize the effect of a cancer protein of the invention. By "neutralize" is meant that an activity of a protein is inhibited or blocked, along with the consequent effect on the cell. In another aspect, a modulator will neutralize the effect of a gene, and its corresponding protein, of the invention by normalizing levels of said protein. In preferred embodiments, modulators alter expression profiles, or expression profile nucleic acids or proteins provided herein, or downstream effector pathways. In one embodiment, the modulator suppresses a cancer phenotype, e.g. to a normal tissue fingerprint. In another embodiment, a modulator induced a cancer phenotype. Generally, a plurality of assay mixtures is run in parallel with different agent concentrations to obtain a differential response to the various concentrations. Typically, one of these concentrations serves as a negative control, i.e., at zero concentration or below the level of detection.

Modulators, drug candidates or test compounds encompass numerous chemical classes, though typically they are organic molecules, preferably small organic compounds having a molecular weight of more than 100 and less than about 2,500 Daltons. Preferred small molecules are less than 2000, or less than 1500 or less than 1000 or less than 500 D. Candidate agents comprise functional groups necessary for structural interaction with proteins, particularly hydrogen bonding, and typically include at least an amine, carbonyl, hydroxyl or carboxyl group, preferably at least two of the functional chemical groups. The candidate agents often comprise cyclical carbon or heterocyclic structures and/or aromatic or polyaromatic structures substituted with one or more of the above functional groups. Modulators also comprise biomolecules such as peptides, saccharides, fatty acids, steroids, purines, pyrimidines, derivatives, structural analogs or combinations thereof. Particularly preferred are peptides. One class of modulators are peptides, for example of from about five to about 35 amino acids, with from about five to about 20 amino acids being preferred, and from about 7 to about 15 being particularly preferred. Preferably, the cancer modulatory protein is soluble, includes a non-transmembrane region, and/or, has an N-terminal Cys to aid in solubility. In one embodiment, the C-terminus of the fragment is kept as a free acid and the N-terminus is a free amine to aid in coupling, i.e., to cysteine. In one embodiment, a cancer protein of the invention is conjugated to an immunogenic agent as discussed herein. In one embodiment, the cancer protein is conjugated to BSA. The peptides of the invention, e.g., of preferred lengths, can be linked to each other or to other amino acids to create a longer peptide/protein. The modulatory peptides can be digests of naturally occurring proteins as is outlined above, random peptides, or "biased" random peptides. In a preferred embodiment, peptide/protein-based modulators are antibodies, and fragments thereof, as defined herein.

Modulators of cancer can also be nucleic acids. Nucleic acid modulating agents can be naturally occurring nucleic acids, random nucleic acids, or "biased" random nucleic acids. For example, digests of prokaryotic or eukaryotic genomes can be used in an approach analogous to that outlined above for proteins.

The term "monoclonal antibody" refers to an antibody obtained from a population of substantially homogeneous antibodies, i.e., the antibodies comprising the population are identical except for possible naturally occurring mutations that are present in minor amounts.

A "motif", as in biological motif of a 24P4C12-related protein, refers to any pattern of amino acids forming part of the primary sequence of a protein, that is associated with a particular function (e.g. protein-protein interaction, protein-DNA interaction, etc) or modification (e.g. that is phosphorylated, glycosylated or amidated), or localization (e.g. secretory sequence, nuclear localization sequence, etc.) or a sequence that is correlated with being immunogenic, either humorally or cellularly. A motif can be either contiguous or capable of being aligned to certain positions that are generally correlated with a certain function or property. In the context of HLA motifs, "motif" refers to the pattern of residues in a peptide of defined length, usually a peptide of from about 8 to about 13 amino acids for a class I HLA motif and from about 6 to about 25 amino acids for a class II HLA motif, which is recognized by a particular HLA molecule. Peptide motifs for HLA binding are typically different for each protein encoded by each human HLA allele and differ in the pattern of the primary and secondary anchor residues.

Figure 2B-2

300 S A Y Q S V Q E T W L A A L I V L A V L
901 TCAGTGCCTACCAGAGCGTGCAGGAGACCTGGCTGGCCGCCCTGATCGTGTGGCGGTGC
320 E A I L L L M L I F L F Q R I R I A I A
961 TTGAAGCCATCCTGCTGCTGATGCTCATCTTCCIGCGGCAGCGGATTTCGTATTGCCATCG
340 L L K E A S K A V G Q M M S T M F Y P L
1021 CCCTCCTGAAGGAGGCCAGCAAGGCTGTGGGACAGATGATGTCTACCATGTTCTACCCAC
360 V T F V L L L I C I A Y W A M T A L Y L
1081 TGGTCACCTTTGTCTCTCTCTCATCTGCATTGCTACTGGGCCATGACTGCTCTGTACC
380 A T S G Q P Q Y V L W A S N I S S P G C
1141 TGGCTACATCGGGGCAACCCAGTATGTGCTCTGGGCATCCAACATCAGCTCCCCCGGCT
400 E K V P I N T S C N P T A H L V N S S C
1201 GTGAGAAAGTGCCAATAAATACATCATGCAACCCACGGCCACCTTGIGAACCTCCTCGT
420 P G L M C V F Q G Y S S K G L I Q R S V
1261 GCCCAGGGCTGATGTGCGTCTTCCAGGGCTACTCATCCAAAGGCCTAATCCAACGTTCTG
440 F N L Q I Y G V L G L F W T L N W V L A
1321 TCTTCAATCTGCAAATCTATGGGGTCTGGGGCTCTTCTGGACCTTAAGTGGGTACTGG
460 L G Q C V L A G A F A S F Y W A F H K P
1381 CCCTGGGCCAAIGCGTCTCGTGGAGCCTTGGCTCTTCTACTGGGCCTTCCACAAGC
480 Q D I P T F P L I S A F I R T L R Y H T
1441 CCCAGGACATCCCTACCTTCCCCCTTAATCTCTGCCTTCATCCGCACACTCCGTTACCACA
500 G S L A F G A L I L T L V Q I A R V I L
1501 CTGGGTCATTGGCATTGGAGCCCTCATCCTGACCTTGTGCGAGATAGCCCGGGTCACTCT
520 E Y I D H K L R G V Q N P V A R C I M C
1561 TGGAGTATATTGACCACAAGCTCAGAGGAGTGCAGAACCCTGTAGCCCGCTGCATCATGT
540 C F R C C L W C L E K F I K F L N R N A
1621 GCTGTTTCAAGTGCTGCCTCTGGTGTCTGGAAAAATTTATCAAGTTCTTAAACCGCAATG
560 Y I M I A I Y G K N F C V S A R N A F M
1681 CATACTCATGATCGCCATCTACGGGAAGAATTTCTGTGTCTCAGCCAAAAATGCGTTCA
580 L L M R N I V R V V V L D K V T D L L L
1741 TGCTACTCATGCGAAACATTGTGAGGGTGGTCTGCTCGGACAAAGTCACAGACCTGCTGC
600 F F G K L L V V G G V G V L S F F F F S
1801 TGTTCCTTTGGGAAGCTGCTGGTGGTCTGGAGGCGTGGGGTCTGTCTCTCTTTTTTTTCT
620 G R I P G L G K D F R S P H L N Y Y W L
1861 CCGGTGCGATCCCGGGGCTGGGTAAAGACTTTAAGAGCCCCCACCTCAACTATTACTGGC
640 P I M T S I L G A Y V I A S G F F S V F
1921 TGCCCATCATGACCTCCATCCTGGGGGCTATGTCATCGCCAGCGGCTTCTTCAGCGTTT
660 G M C V D T L F L C F L E D L E R N N G
1981 TCGGCATGTGTGTGGACACGCTCTTCTCTGCTTCTGGAAGACCTGGAGCGGAACAACG
530 S L D R P Y Y M S R S L L K I L G K K N
2041 GCTCCCTGGACCGGCCCTACTACATGTCCAAGAGCCTTCTAAAGATTCTGGGCAAGAAGA

A “pharmaceutical excipient” comprises a material such as an adjuvant, a carrier, pH-adjusting and buffering agents, tonicity adjusting agents, wetting agents, preservative, and the like.

“Pharmaceutically acceptable” refers to a non-toxic, inert, and/or composition that is physiologically compatible with humans or other mammals.

The term “polynucleotide” means a polymeric form of nucleotides of at least 10 bases or base pairs in length, either ribonucleotides or deoxynucleotides or a modified form of either type of nucleotide, and is meant to include single and double stranded forms of DNA and/or RNA. In the art, this term is often used interchangeably with “oligonucleotide”. A polynucleotide can comprise a nucleotide sequence disclosed herein wherein thymidine (T), as shown for example in FIG. 2, can also be uracil (U); this definition pertains to the differences between the chemical structures of DNA and RNA, in particular the observation that one of the four major bases in RNA is uracil (U) instead of thymidine (T).

The term “polypeptide” means a polymer of at least about 4, 5, 6, 7, or 8 amino acids. Throughout the specification, standard three letter or single letter designations for amino acids are used. In the art, this term is often used interchangeably with “peptide” or “protein”.

An HLA “primary anchor residue” is an amino acid at a specific position along a peptide sequence which is understood to provide a contact point between the immunogenic peptide and the HLA molecule. One to three, usually two, primary anchor residues within a peptide of defined length generally defines a “motif” for an immunogenic peptide. These residues are understood to fit in close contact with peptide binding groove of an HLA molecule, with their side chains buried in specific pockets of the binding groove. In one embodiment, for example, the primary anchor residues for an HLA class I molecule are located at position 2 (from the amino terminal position) and at the carboxyl terminal position of a 8, 9, 10, 11, or 12 residue peptide epitope in accordance with the invention. Alternatively, in another embodiment, the primary anchor residues of a peptide binds an HLA class II molecule are spaced relative to each other, rather than to the termini of a peptide, where the peptide is generally of at least 9 amino acids in length. The primary anchor positions for each motif and supermotif are set forth in Table IV. For example, analog peptides can be created by altering the presence or absence of particular residues in the primary and/or secondary anchor positions shown in Table IV. Such analogs are used to modulate the binding affinity and/or population coverage of a peptide comprising a particular HLA motif or supermotif.

“Radioisotopes” include, but are not limited to the following (non-limiting exemplary uses are also set forth):

Examples of Medical Isotopes:

Isotope	Description of use
Actinium-225 (Ac-225)	See Thorium-229 (Th-229)
Actinium-227 (Ac-227)	Parent of Radium-223 (Ra-223) which is an alpha emitter used to treat metastases in the skeleton resulting from cancer (i.e., breast and prostate cancers), and cancer radioimmunotherapy
Bismuth-212 (Bi-212)	See Thorium-228 (Th-228)
Bismuth-213 (Bi-213)	See Thorium-229 (Th-229)
Cadmium-109 (Cd-109)	Cancer detection

-continued

Isotope	Description of use
5 Cobalt-60 (Co-60)	Radiation source for radiotherapy of cancer, for food irradiators, and for sterilization of medical supplies
Copper-64 (Cu-64)	A positron emitter used for cancer therapy and SPECT imaging
Copper-67 (Cu-67)	Beta/gamma emitter used in cancer radioimmunotherapy and diagnostic studies (i.e., breast and colon cancers, and lymphoma)
10 Dysprosium-166 (Dy-166)	Cancer radioimmunotherapy
Erbium-169 (Er-169)	Rheumatoid arthritis treatment, particularly for the small joints associated with fingers and toes
Europium-152 (Eu-152)	Radiation source for food irradiation and for sterilization of medical supplies
15 Europium-154 (Eu-154)	Radiation source for food irradiation and for sterilization of medical supplies
Gadolinium-153 (Gd-153)	Osteoporosis detection and nuclear medical quality assurance devices
Gold-198 (Au-198)	Implant and intracavity therapy of ovarian, prostate, and brain cancers
20 Holmium-166 (Ho-166)	Multiple myeloma treatment in targeted skeletal therapy, cancer radioimmunotherapy, bone marrow ablation, and rheumatoid arthritis treatment
Iodine-125 (I-125)	Osteoporosis detection, diagnostic imaging, tracer drugs, brain cancer treatment, radiolabeling, tumor imaging, mapping of receptors in the brain, interstitial radiation therapy, brachytherapy for treatment of prostate cancer, determination of glomerular filtration rate (GFR), determination of plasma volume, detection of deep vein thrombosis of the legs
25 Iodine-131 (I-131)	Thyroid function evaluation, thyroid disease detection, treatment of thyroid cancer as well as other non-malignant thyroid diseases (i.e., Graves disease, goiters, and hyperthyroidism), treatment of leukemia, lymphoma, and other forms of cancer (e.g., breast cancer) using radioimmunotherapy
30 Iridium-192 (Ir-192)	Brachytherapy, brain and spinal cord tumor treatment, treatment of blocked arteries (i.e., arteriosclerosis and restenosis), and implants for breast and prostate tumors
35 Lutetium-177 (Lu-177)	Cancer radioimmunotherapy and treatment of blocked arteries (i.e., arteriosclerosis and restenosis)
Molybdenum-99 (Mo-99)	Parent of Technetium-99m (Tc-99m) which is used for imaging the brain, liver, lungs, heart, and other organs. Currently, Tc-99m is the most widely used radioisotope used for diagnostic imaging of various cancers and diseases involving the brain, heart, liver, lungs; also used in detection of deep vein thrombosis of the legs
40 Osmium-194 (Os-194)	Cancer radioimmunotherapy
Palladium-103 (Pd-103)	Prostate cancer treatment
45 Platinum-195m (Pt-195m)	Studies on biodistribution and metabolism of cisplatin, a chemotherapeutic drug
Phosphorus-32 (P-32)	Polycythemia rubra vera (blood cell disease) and leukemia treatment, bone cancer diagnosis/treatment; colon, pancreatic, and liver cancer treatment; radiolabeling nucleic acids for in vitro research, diagnosis of superficial tumors, treatment of blocked arteries (i.e., arteriosclerosis and restenosis), and intracavity therapy
50 Phosphorus-33 (P-33)	Leukemia treatment, bone disease diagnosis/treatment, radiolabeling, and treatment of blocked arteries (i.e., arteriosclerosis and restenosis)
55 Radium-223 (Ra-223)	See Actinium-227 (Ac-227)
Rhenium-186 (Re-186)	Bone cancer pain relief, rheumatoid arthritis treatment, and diagnosis and treatment of lymphoma and bone, breast, colon, and liver cancers using radioimmunotherapy
60 Rhenium-188 (Re-188)	Cancer diagnosis and treatment using radioimmunotherapy, bone cancer pain relief, treatment of rheumatoid arthritis, and treatment of prostate cancer
Rhodium-105 (Rh-105)	Cancer radioimmunotherapy
Samarium-145 (Sm-145)	Ocular cancer treatment
65 Samarium-153 (Sm-153)	Cancer radioimmunotherapy and bone cancer pain relief

-continued

Isotope	Description of use
Scandium-47 (Sc-47)	Cancer radioimmunotherapy and bone cancer pain relief
Selenium-75 (Se-75)	Radiotracer used in brain studies, imaging of adrenal cortex by gamma-scintigraphy, lateral locations of steroid secreting tumors, pancreatic scanning, detection of hyperactive parathyroid glands, measure rate of bile acid loss from the endogenous pool
Strontium-85 (Sr-85)	Bone cancer detection and brain scans
Strontium-89 (Sr-89)	Bone cancer pain relief, multiple myeloma treatment, and osteoblastic therapy
Technetium-99m (Tc-99m)	See Molybdenum-99 (Mo-99)
Thorium-228 (Th-228)	Parent of Bismuth-212 (Bi-212) which is an alpha emitter used in cancer radioimmunotherapy
Thorium-229 (Th-229)	Parent of Actinium-225 (Ac-225) and grandparent of Bismuth-213 (Bi-213) which are alpha emitters used in cancer radioimmunotherapy
Thulium-170 (Tm-170)	Gamma source for blood irradiators, energy source for implanted medical devices
Tin-117m (Sn-117m)	Cancer immunotherapy and bone cancer pain relief
Tungsten-188 (W-188)	Parent for Rhenium-188 (Re-188) which is used for cancer diagnostics/treatment, bone cancer pain relief, rheumatoid arthritis treatment, and treatment of blocked arteries (i.e., arteriosclerosis and restenosis)
Xenon-127 (Xe-127)	Neuroimaging of brain disorders, high resolution SPECT studies, pulmonary function tests, and cerebral blood flow studies
Ytterbium-175 (Yb-175)	Cancer radioimmunotherapy
Yttrium-90 (Y-90)	Microseeds obtained from irradiating Yttrium-89 (Y-89) for liver cancer treatment
Yttrium-91 (Y-91)	A gamma-emitting label for Yttrium-90 (Y-90) which is used for cancer radioimmunotherapy (i.e., lymphoma, breast, colon, kidney, lung, ovarian, prostate, pancreatic, and inoperable liver cancers)

By "randomized" or grammatical equivalents as herein applied to nucleic acids and proteins is meant that each nucleic acid and peptide consists of essentially random nucleotides and amino acids, respectively. These random peptides (or nucleic acids, discussed herein) can incorporate any nucleotide or amino acid at any position. The synthetic process can be designed to generate randomized proteins or nucleic acids, to allow the formation of all or most of the possible combinations over the length of the sequence, thus forming a library of randomized candidate bicactive proteinaceous agents.

In one embodiment, a library is "fully randomized," with no sequence preferences or constants at any position. In another embodiment, the library is a "biased random" library. That is, some positions within the sequence either are held constant, or are selected from a limited number of possibilities. For example, the nucleotides or amino acid residues are randomized within a defined class, e.g., of hydrophobic amino acids, hydrophilic residues, sterically biased (either small or large) residues, towards the creation of nucleic acid binding domains, the creation of cysteines, for cross-linking, prolines for SH-3 domains, serines, threonines, tyrosines or histidines for phosphorylation sites, etc., or to purines, etc.

A "recombinant" DNA or RNA molecule is a DNA or RNA molecule that has been subjected to molecular manipulation in vitro.

Non-limiting examples of small molecules include compounds that bind or interact with 24P4C12, ligands including hormones, neuropeptides, chemokines, odorants, phospholipids, and functional equivalents thereof that bind and preferably inhibit 24P4C12 protein function. Such non-limiting small molecules preferably have a molecular weight of less

than about 10 kDa, more preferably below about 9, about 8, about 7, about 6, about 5 or about 4 kDa. In certain embodiments, small molecules physically associate with, or bind, 24P4C12 protein; are not found in naturally occurring metabolic pathways; and/or are more soluble in aqueous than non-aqueous solutions

"Stringency" of hybridization reactions is readily determinable by one of ordinary skill in the art, and generally is an empirical calculation dependent upon probe length, washing temperature, and salt concentration. In general, longer probes require higher temperatures for proper annealing, while shorter probes need lower temperatures. Hybridization generally depends on the ability of denatured nucleic acid sequences to reanneal when complementary strands are present in an environment below their melting temperature. The higher the degree of desired homology between the probe and hybridizable sequence, the higher the relative temperature that can be used. As a result, it follows that higher relative temperatures would tend to make the reaction conditions more stringent, while lower temperatures less so. For additional details and explanation of stringency of hybridization reactions, see Ausubel et al., *Current Protocols in Molecular Biology*, Wiley Interscience Publishers, (1995).

"Stringent conditions" or "high stringency conditions", as defined herein, are identified by, but not limited to, those that: (1) employ low ionic strength and high temperature for washing, for example 0.015 M sodium chloride/0.0015 M sodium citrate/0.1% sodium dodecyl sulfate at 50° C.; (2) employ during hybridization a denaturing agent, such as formamide, for example, 50% (v/v) formamide with 0.1% bovine serum albumin/0.1% Ficoll/0.1% polyvinylpyrrolidone/50 mM sodium phosphate buffer at pH 6.5 with 750 mM sodium chloride, 75 mM sodium citrate at 42° C.; or (3) employ 50% formamide, 5×SSC (0.75 M NaCl, 0.075 M sodium citrate), 50 mM sodium phosphate (pH 6.8), 0.1% sodium pyrophosphate, 5×Denhardt's solution, sonicated salmon sperm DNA (50 µg/ml), 0.1% SDS, and 10% dextran sulfate at 42° C., with washes at 42° C. in 0.2×SSC (sodium chloride/sodium citrate) and 50% formamide at 55° C., followed by a high-stringency wash consisting of 0.1×SSC containing EDTA at 55° C. "Moderately stringent conditions" are described by, but not limited to, those in Sambrook et al., *Molecular Cloning: A Laboratory Manual*, New York: Cold Spring Harbor Press, 1989, and include the use of washing solution and hybridization conditions (e.g., temperature, ionic strength and % SDS) less stringent than those described above. An example of moderately stringent conditions is overnight incubation at 37° C. in a solution comprising: 20% formamide, 5×SSC (150 mM NaCl, 15 mM trisodium citrate), 50 mM sodium phosphate (pH 7.6), 5×Denhardt's solution, 10% dextran sulfate, and 20 mg/mL denatured sheared salmon sperm DNA, followed by washing the filters in 1×SSC at about 37-50° C. The skilled artisan will recognize how to adjust the temperature, ionic strength, etc. as necessary to accommodate factors such as probe length and the like.

An HLA "supermotif" is a peptide binding specificity shared by HLA molecules encoded by two or more HLA alleles. Overall phenotypic frequencies of HLA-supertypes in different ethnic populations are set forth in Table IV (F). The non-limiting constituents of various supertypes are as follows:

A2: A*0201, A*0202, A*0203, A*0204, A*0205, A*0206, A*6802, A*6901, A*0207

A3: A3, A11, A31, A*3301, A*6801, A*0301, A*1101, A*3101

B7: B7, B*3501-03, B*51, B*5301, B*5401, B*5501, B*5502, B*5601, B*6701, B*7801, B*0702, B*5101, B*5602

B44: B*3701, B*4402, B*4403, B*60 (B*4001), B61 (B*4006)

A1: A*0102, A*2604, A*3601, A*4301, A*8001

A24: A*24, A*30, A*2403, A*2404, A*3002, A*3003

B27: B*1401-02, B*1503, B*1509, B*1510, B*1518, B*3801-02, B*3901, B*3902, B*3903-04, B*4801-02, B*7301, B*2701-08

B58: B*1516, B*1517, B*5701, B*5702, B58

B62: B*4601, B52, B*1501 (B62), B*1502 (B75), B*1513 (B77)

Calculated population coverage afforded by different HLA-supertype combinations are set forth in Table IV (G).

As used herein “to treat” or “therapeutic” and grammatically related terms, refer to any improvement of any consequence of disease, such as prolonged survival, less morbidity, and/or a lessening of side effects which are the byproducts of an alternative therapeutic modality; full eradication of disease is not required.

A “transgenic animal” (e.g., a mouse or rat) is an animal having cells that contain a transgene, which transgene was introduced into the animal or an ancestor of the animal at a prenatal, e.g., an embryonic stage. A “transgene” is a DNA that is integrated into the genome of a cell from which a transgenic animal develops.

As used herein, an HLA or cellular immune response “vaccine” is a composition that contains or encodes one or more peptides of the invention. There are numerous embodiments of such vaccines, such as a cocktail of one or more individual peptides; one or more peptides of the invention comprised by a polypeptidic peptide; or nucleic acids that encode such individual peptides or polypeptides, e.g., a minigene that encodes a polypeptidic peptide. The “one or more peptides” can include any whole unit integer from 1-150 or more, e.g., at least 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 105, 110, 115, 120, 125, 130, 135, 140, 145, or 150 or more peptides of the invention. The peptides or polypeptides can optionally be modified, such as by lipidation, addition of targeting or other sequences. HLA class I peptides of the invention can be admixed with, or linked to, HLA class II peptides, to facilitate activation of both cytotoxic T lymphocytes and helper T lymphocytes. HLA vaccines can also comprise peptide-pulsed antigen presenting cells, e.g., dendritic cells.

The term “variant” refers to a molecule that exhibits a variation from a described type or norm, such as a protein that has one or more different amino acid residues in the corresponding position(s) of a specifically described protein (e.g. the 24P4C12 protein shown in FIG. 2 or FIG. 3. An analog is an example of a variant protein. Splice isoforms and single nucleotides polymorphisms (SNPs) are further examples of variants.

The “24P4C12-related proteins” of the invention include those specifically identified herein, as well as allelic variants, conservative substitution variants, analogs and homologs that can be isolated/generated and characterized without undue experimentation following the methods outlined herein or readily available in the art. Fusion proteins that combine parts of different 24P4C12 proteins or fragments thereof, as well as fusion proteins of a 24P4C12 protein and a heterologous polypeptide are also included. Such 24P4C12 proteins are collectively referred to as the 24P4C12-related proteins, the proteins of the invention, or 24P4C12. The term “24P4C12-

related protein” refers to a polypeptide fragment or a 24P4C12 protein sequence of 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or more than 25 amino acids; or, at least 30, 35, 40, 45, 50, 55, 60, 65, 70, 80, 85, 90, 95, 100, 105, 110, 115, 120, 125, 130, 135, 140, 145, 150, 155, 160, 165, 170, 175, 180, 185, 190, 195, 200, 225, 250, 275, 300, 325, 350, 375, 400, 425, 450, 475, 500, 525, 550, 575, 600, 625, 650, or 664 or more amino acids.

II.) 24P4C12 POLYNUCLEOTIDES

One aspect of the invention provides polynucleotides corresponding or complementary to all or part of a 24P4C12 gene, mRNA, and/or coding sequence, preferably in isolated form, including polynucleotides encoding a 24P4C12-related protein and fragments thereof, DNA, RNA, DNA/RNA hybrid, and related molecules, polynucleotides or oligonucleotides complementary to a 24P4C12 gene or mRNA sequence or a part thereof, and polynucleotides or oligonucleotides that hybridize to a 24P4C12 gene, mRNA, or to a 24P4C12 encoding polynucleotide (collectively, “24P4C12 polynucleotides”). In all instances when referred to in this section, T can also be U in FIG. 2.

Embodiments of a 24P4C12 polynucleotide include: a 24P4C12 polynucleotide having the sequence shown in FIG. 2, the nucleotide sequence of 24P4C12 as shown in FIG. 2 wherein T is U; at least 10 contiguous nucleotides of a polynucleotide having the sequence as shown in FIG. 2; or, at least 10 contiguous nucleotides of a polynucleotide having the sequence as shown in FIG. 2 where T is U. For example, embodiments of 24P4C12 nucleotides comprise, without limitation:

- (I) a polynucleotide comprising, consisting essentially of, or consisting of a sequence as shown in FIG. 2, wherein T can also be U;
- (II) a polynucleotide comprising, consisting essentially of, or consisting of the sequence as shown in FIG. 2A, from nucleotide residue number 6 through nucleotide residue number 2138, including the stop codon, wherein T can also be U;
- (III) a polynucleotide comprising, consisting essentially of, or consisting of the sequence as shown in FIG. 2B, from nucleotide residue number 6 through nucleotide residue number 2138, including the stop codon, wherein T can also be U;
- (IV) a polynucleotide comprising, consisting essentially of, or consisting of the sequence as shown in FIG. 2C, from nucleotide residue number 6 through nucleotide residue number 2138, including the a stop codon, wherein T can also be U;
- (V) a polynucleotide comprising, consisting essentially of, or consisting of the sequence as shown in FIG. 2D, from nucleotide residue number 6 through nucleotide residue number 2138, including the stop codon, wherein T can also be U;
- (VI) a polynucleotide comprising, consisting essentially of, or consisting of the sequence as shown in FIG. 2E, from nucleotide residue number 6 through nucleotide residue number 2138, including the stop codon, wherein T can also be U;
- (VII) a polynucleotide comprising, consisting essentially of, or consisting of the sequence as shown in FIG. 2F, from nucleotide residue number 6 through nucleotide residue number 2138, including the stop codon, wherein T can also be U;
- (VIII) a polynucleotide comprising, consisting essentially of, or consisting of the sequence as shown in FIG. 2G,

- from nucleotide residue number 6 through nucleotide residue number 1802, including the stop codon, wherein T can also be U;
- (IX) a polynucleotide comprising, consisting essentially of, or consisting of the sequence as shown in FIG. 2H, from nucleotide residue number 6 through nucleotide residue number 2174, including the stop codon, wherein T can also be U;
- (X) a polynucleotide comprising, consisting essentially of, or consisting of the sequence as shown in FIG. 2I, from nucleotide residue number 6 through nucleotide residue number 2144, including the stop codon, wherein T can also be U;
- (XI) a polynucleotide that encodes a 24P4C12-related protein that is at least 90, 91, 92, 93, 94, 95, 96, 97, 98, 99 or 100% homologous to an entire amino acid sequence shown in FIGS. 2A-I;
- (XII) a polynucleotide that encodes a 24P4C12-related protein that is at least 90, 91, 92, 93, 94, 95, 96, 97, 98, 99 or 100% identical to an entire amino acid sequence shown in FIGS. 2A-I;
- (XIII) a polynucleotide that encodes at least one peptide set forth in Tables VIII-XXI and XXII-XLIX;
- (XIV) a polynucleotide that encodes a peptide region of at least 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35 amino acids of a peptide of FIG. 3A-D in any whole number increment up to 710 that includes at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35 amino acid position(s) having a value greater than 0.5 in the Hydrophilicity profile of FIG. 5;
- (XV) a polynucleotide that encodes a peptide region of at least 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35 amino acids of a peptide of FIG. 3A-D in any whole number increment up to 710 that includes 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35 amino acid position(s) having a value less than 0.5 in the Hydrophaticity profile of FIG. 6;
- (XVI) a polynucleotide that encodes a peptide region of at least 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35 amino acids of a peptide of FIG. 3A-D in any whole number increment up to 710 that includes 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35 amino acid position(s) having a value greater than 0.5 in the Percent Accessible Residues profile of FIG. 7;
- (XVII) a polynucleotide that encodes a peptide region of at least 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35 amino acids of a peptide of FIG. 3A-D in any whole number increment up to 710 that includes 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35 amino acid position(s) having a value greater than 0.5 in the Average Flexibility profile of FIG. 8;
- (XVIII) a polynucleotide that encodes a peptide region of at least 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35 amino acids of a peptide of FIG. 3A-D in any whole number increment up to 710 that includes 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22,

- 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35 amino acid position(s) having a value greater than 0.5 in the Beta-turn profile of FIG. 9;
- (XIX) a polynucleotide that encodes a peptide region of at least 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35 amino acids of a peptide of FIG. 3E in any whole number increment up to 598 that includes 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35 amino acid position(s) having a value greater than 0.5 in the Hydrophaticity profile of FIG. 5;
- (XX) a polynucleotide that encodes a peptide region of at least 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35 amino acids of a peptide of FIG. 3E in any whole number increment up to 598 that includes 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35 amino acid position(s) having a value less than 0.5 in the Hydrophaticity profile of FIG. 6;
- (XXI) a polynucleotide that encodes a peptide region of at least 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35 amino acids of a peptide of FIG. 3E in any whole number increment up to 598 that includes 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35 amino acid position(s) having a value greater than 0.5 in the Percent Accessible Residues profile of FIG. 7;
- (XXII) a polynucleotide that encodes a peptide region of at least 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35 amino acids of a peptide of FIG. 3E in any whole number increment up to 598 that includes 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35 amino acid position(s) having a value greater than 0.5 in the Average Flexibility profile of FIG. 8;
- (XXIII) a polynucleotide that encodes a peptide region of at least 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35 amino acids of a peptide of FIG. 3E in any whole number increment up to 598 that includes 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35 amino acid position(s) having a value greater than 0.5 in the Beta-turn profile of FIG. 9;
- (XXIV) a polynucleotide that encodes a peptide region of at least 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35 amino acids of a peptide of FIG. 3F in any whole number increment up to 722 that includes 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35 amino acid position(s) having a value greater than 0.5 in the Hydrophilicity profile of FIG. 5;
- (XXV) a polynucleotide that encodes a peptide region of at least 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35 amino acids of a peptide of FIG. 3F in any whole number increment up to 722 that includes 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35 amino acid position(s) having a value less than 0.5 in the Hydrophaticity profile of FIG. 6;

- (XXVI) a polynucleotide that encodes a peptide region of at least 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35 amino acids of a peptide of FIG. 3F in any whole number increment up to 722 that includes 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35 amino acid position(s) having a value greater than 0.5 in the Percent Accessible Residues profile of FIG. 7;
- (XXVII) a polynucleotide that encodes a peptide region of at least 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35 amino acids of a peptide of FIG. 3F in any whole number increment up to 722 that includes 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35 amino acid position(s) having a value greater than 0.5 in the Average Flexibility profile of FIG. 8;
- (XXVIII) a polynucleotide that encodes a peptide region of at least 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35 amino acids of a peptide of FIG. 3F in any whole number increment up to 722 that includes 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35 amino acid position(s) having a value greater than 0.5 in the Beta-turn profile of FIG. 9
- (XXIX) a polynucleotide that encodes a peptide region of at least 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35 amino acids of a peptide of FIG. 3G in any whole number increment up to 712 that includes 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35 amino acid position(s) having a value greater than 0.5 in the Hydrophilicity profile of FIG. 5;
- (XXX) a polynucleotide that encodes a peptide region of at least 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35 amino acids of a peptide of FIG. 3G in any whole number increment up to 712 that includes 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35 amino acid position(s) having a value less than 0.5 in the Hydrophobicity profile of FIG. 6;
- (XXXI) a polynucleotide that encodes a peptide region of at least 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35 amino acids of a peptide of FIG. 3G in any whole number increment up to 712 that includes 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35 amino acid position(s) having a value greater than 0.5 in the Percent Accessible Residues profile of FIG. 7;
- (XXXII) a polynucleotide that encodes a peptide region of at least 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35 amino acids of a peptide of FIG. 3G in any whole number increment up to 712 that includes 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35 amino acid position(s) having a value greater than 0.5 in the Average Flexibility profile of FIG. 8;
- (XXXIII) a polynucleotide that encodes a peptide region of at least 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35 amino acids of a peptide of FIG. 3G in any whole

- number increment up to 712 that includes 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35 amino acid position(s) having a value greater than 0.5 in the Beta-turn profile of FIG. 9
- (XXXIV) a polynucleotide that is fully complementary to a polynucleotide of any one of (I)-(XXXIII).
- (XXXV) a peptide that is encoded by any of (I) to (XXXIII); and
- (XXXVI) a composition comprising a polynucleotide of any of (I)-(XXXIV) or peptide of (XXXV) together with a pharmaceutical excipient and/or in a human unit dose form.
- (XXXVII) a method of using a polynucleotide of any (I)-(XXXIV) or peptide of (XXXV) or a composition of (XXXVI) in a method to modulate a cell expressing 24P4C12,
- (XXXVIII) a method of using a polynucleotide of any (I)-(XXXIV) or peptide of (XXXV) or a composition of (XXXVI) in a method to diagnose, prophylax, prognose, or treat an individual who bears a cell expressing 24P4C12
- (XXXIX) a method of using a polynucleotide of any (I)-(XXXIV) or peptide of (XXXV) or a composition of (XXXVI) in a method to diagnose, prophylax, prognose, or treat an individual who bears a cell expressing 24P4C12, said cell from a cancer of a tissue listed in Table I;
- (XL) a method of using a polynucleotide of any (I)-(XXXIV) or peptide of (XXXV) or a composition of (XXXVI) in a method to diagnose, prophylax, prognose, or treat a cancer;
- (XLI) a method of using a polynucleotide of any (I)-(XXXIV) or peptide of (XXXV) or a composition of (XXXVI) in a method to diagnose, prophylax, prognose, or treat a cancer of a tissue listed in Table I; and,
- (XLII) a method of using a polynucleotide of any (I)-(XXXIV) or peptide of (XXXV) or a composition of (XXXVI) in a method to identify or characterize a modulator of a cell expressing 24P4C12.
- As used herein, a range is understood to disclose specifically all whole unit positions thereof.
- Typical embodiments of the invention disclosed herein include 24P4C12 polynucleotides that encode specific portions of 24P4C12 mRNA sequences (and those which are complementary to such sequences) such as those that encode the proteins and/or fragments thereof, for example:
- (a) 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 105, 110, 115, 120, 125, 130, 135, 140, 145, 150, 155, 160, 165, 170, 175, 180, 185, 190, 195, 200, 225, 250, 275, 300, 325, 350, 375, 400, 425, 450, 475, 500, 525, 550, 575, 600, 625, 650, 675, 700, 710 or more contiguous amino acids of 24P4C12 variant 1; the maximal lengths relevant for other variants are: variant 3, 710 amino acids; variant 5, 710 amino acids, variant 6, 710 amino acids, variant 7, 598 amino acids, variant 8, 722 amino acids, and variant 9, 712 amino acids.
- For example, representative embodiments of the invention disclosed herein include: polynucleotides and their encoded peptides themselves encoding about amino acid 1 to about amino acid 10 of the 24P4C12 protein shown in FIG. 2 or FIG. 3, polynucleotides encoding about amino acid 10 to about amino acid 20 of the 24P4C12 protein shown in FIG. 2 or FIG. 3, polynucleotides encoding about amino acid 20 to about amino acid 30 of the 24P4C12 protein shown in FIG. 2 or FIG. 3, polynucleotides encoding about amino acid 30 to about

amino acid 40 of the 24P4C12 protein shown in FIG. 2 or FIG. 3, polynucleotides encoding about amino acid 40 to about amino acid 50 of the 24P4C12 protein shown in FIG. 2 or FIG. 3, polynucleotides encoding about amino acid 50 to about amino acid 60 of the 24P4C12 protein shown in FIG. 2 or FIG. 3, polynucleotides encoding about amino acid 60 to about amino acid 70 of the 24P4C12 protein shown in FIG. 2 or FIG. 3, polynucleotides encoding about amino acid 70 to about amino acid 80 of the 24P4C12 protein shown in FIG. 2 or FIG. 3, polynucleotides encoding about amino acid 80 to about amino acid 90 of the 24P4C12 protein shown in FIG. 2 or FIG. 3, polynucleotides encoding about amino acid 90 to about amino acid 100 of the 24P4C12 protein shown in FIG. 2 or FIG. 3, in increments of about 10 amino acids, ending at the carboxyl terminal amino acid set forth in FIG. 2 or FIG. 3. Accordingly, polynucleotides encoding portions of the amino acid sequence (of about 10 amino acids), of amino acids, 100 through the carboxyl terminal amino acid of the 24P4C12 protein are embodiments of the invention. Wherein it is understood that each particular amino acid position discloses that position plus or minus five amino acid residues.

Polynucleotides encoding relatively long portions of a 24P4C12 protein are also within the scope of the invention. For example, polynucleotides encoding from about amino acid 1 (or 20 or 30 or 40 etc.) to about amino acid 20, (or 30, or 40 or 50 etc.) of the 24P4C12 protein "or variant" shown in FIG. 2 or FIG. 3 can be generated by a variety of techniques well known in the art. These polynucleotide fragments can include any portion of the 24P4C12 sequence as shown in FIG. 2.

Additional illustrative embodiments of the invention disclosed herein include 24P4C12 polynucleotide fragments encoding one or more of the biological motifs contained within a 24P4C12 protein "or variant" sequence, including one or more of the motif-bearing subsequences of a 24P4C12 protein "or variant" set forth in Tables VIII-XXI and XXII-XLIX. In another embodiment, typical polynucleotide fragments of the invention encode one or more of the regions of 24P4C12 protein or variant that exhibit homology to a known molecule. In another embodiment of the invention, typical polynucleotide fragments can encode one or more of the 24P4C12 protein or variant N-glycosylation sites, cAMP and cGMP-dependent protein kinase phosphorylation sites, casein kinase II phosphorylation sites or N-myristoylation site and amidation sites.

Note that to determine the starting position of any peptide set forth in Tables VIII-XXI and Tables XXII to XLIX (collectively HLA Peptide Tables) respective to its parental protein, e.g., variant 1, variant 2, etc., reference is made to three factors: the particular variant, the length of the peptide in an HLA Peptide Table, and the Search Peptides listed in Table LVII. Generally, a unique Search Peptide is used to obtain HLA peptides for a particular variant. The position of each Search Peptide relative to its respective parent molecule is listed in Table VII. Accordingly, if a Search Peptide begins at position "X", one must add the value "X minus 1" to each position in Tables VIII-XXI and Tables XXII-IL to obtain the actual position of the HLA peptides in their parental molecule. For example if a particular Search Peptide begins at position 150 of its parental molecule, one must add 150-1, i.e., 149 to each HLA peptide amino acid position to calculate the position of that amino acid in the parent molecule.

II.A.) Uses of 24P4C12 Polynucleotides

II.A.1.) Monitoring of Genetic Abnormalities

The polynucleotides of the preceding paragraphs have a number of different specific uses. The human 24P4C12 gene maps to the chromosomal location set forth in the Example

entitled "Chromosomal Mapping of 24P4C12." For example, because the 24P4C12 gene maps to this chromosome, polynucleotides that encode different regions of the 24P4C12 proteins are used to characterize cytogenetic abnormalities of this chromosomal locale, such as abnormalities that are identified as being associated with various cancers. In certain genes, a variety of chromosomal abnormalities including rearrangements have been identified as frequent cytogenetic abnormalities in a number of different cancers (see e.g. Krajnovic et al., *Mutat. Res.* 382(3-4): 81-83 (1998); Johansson et al., *Blood* 86(10): 3905-3914 (1995) and Finger et al., *P.N.A.S.* 85(23): 9158-9162 (1988)). Thus, polynucleotides encoding specific regions of the 24P4C12 proteins provide new tools that can be used to delineate, with greater precision than previously possible, cytogenetic abnormalities in the chromosomal region that encodes 24P4C12 that may contribute to the malignant phenotype. In this context, these polynucleotides satisfy a need in the art for expanding the sensitivity of chromosomal screening in order to identify more subtle and less common chromosomal abnormalities (see e.g. Evans et al., *Am. J. Obstet. Gynecol.* 171(4): 1055-1057 (1994)).

Furthermore, as 24P4C12 was shown to be highly expressed in bladder and other cancers, 24P4C12 polynucleotides are used in methods assessing the status of 24P4C12 gene products in normal versus cancerous tissues. Typically, polynucleotides that encode specific regions of the 24P4C12 proteins are used to assess the presence of perturbations (such as deletions, insertions, point mutations, or alterations resulting in a loss of an antigen etc.) in specific regions of the 24P4C12 gene, such as regions containing one or more motifs. Exemplary assays include both RT-PCR assays as well as single-strand conformation polymorphism (SSCP) analysis (see, e.g., Marrogi et al., *J. Cutan. Pathol.* 26(8): 369-378 (1999)), both of which utilize polynucleotides encoding specific regions of a protein to examine these regions within the protein.

II.A.2.) Antisense Embodiments

Other specifically contemplated nucleic acid related embodiments of the invention disclosed herein are genomic DNA, cDNAs, ribozymes, and antisense molecules, as well as nucleic acid molecules based on an alternative backbone, or including alternative bases, whether derived from natural sources or synthesized, and include molecules capable of inhibiting the RNA or protein expression of 24P4C12. For example, antisense molecules can be RNAs or other molecules, including peptide nucleic acids (PNAs) or non-nucleic acid molecules such as phosphorothioate derivatives that specifically bind DNA or RNA in a base pair-dependent manner. A skilled artisan can readily obtain these classes of nucleic acid molecules using the 24P4C12 polynucleotides and polynucleotide sequences disclosed herein.

Antisense technology entails the administration of exogenous oligonucleotides that bind to a target polynucleotide located within the cells. The term "antisense" refers to the fact that such oligonucleotides are complementary to their intracellular targets, e.g., 24P4C12. See for example, Jack Cohen, *Oligodeoxynucleotides, Antisense Inhibitors of Gene Expression*, CRC Press, 1989; and *Synthesis* 1:1-5 (1988). The 24P4C12 antisense oligonucleotides of the present invention include derivatives such as S-oligonucleotides (phosphorothioate derivatives or S-oligos, see, Jack Cohen, supra), which exhibit enhanced cancer cell growth inhibitory action. S-oligos (nucleoside phosphorothioates) are isoelectronic analogs of an oligonucleotide (O-oligo) in which a nonbridging oxygen atom of the phosphate group is replaced by a sulfur atom. The S-oligos of the present invention can be

prepared by treatment of the corresponding O-oligos with 3H-1,2-benzodithiol-3-one-1,1-dioxide, which is a sulfur transfer reagent. See, e.g., Iyer, R. P. et al., *J. Org. Chem.* 55:4693-4698 (1990); and Iyer, R. P. et al., *J. Am. Chem. Soc.* 112:1253-1254 (1990). Additional 24P4C12 antisense oligonucleotides of the present invention include morpholino antisense oligonucleotides known in the art (see, e.g., Partridge et al., 1996, *Antisense & Nucleic Acid Drug Development* 6: 169-175).

The 24P4C12 antisense oligonucleotides of the present invention typically can be RNA or DNA that is complementary to and stably hybridizes with the first 100 5' codons or last 100 3' codons of a 24P4C12 genomic sequence or the corresponding mRNA. Absolute complementarity is not required, although high degrees of complementarity are preferred. Use of an oligonucleotide complementary to this region allows for the selective hybridization to 24P4C12 mRNA and not to mRNA specifying other regulatory subunits of protein kinase. In one embodiment, 24P4C12 antisense oligonucleotides of the present invention are 15 to 30-mer fragments of the antisense DNA molecule that have a sequence that hybridizes to 24P4C12 mRNA. Optionally, 24P4C12 antisense oligonucleotide is a 30-mer oligonucleotide that is complementary to a region in the first 10 5' codons or last 10 3' codons of 24P4C12. Alternatively, the antisense molecules are modified to employ ribozymes in the inhibition of 24P4C12 expression, see, e.g., L. A. Couture & D. T. Stinchcomb; *Trends Genet* 12: 510-515 (1996).

II.A.3.) Primers and Primer Pairs

Further specific embodiments of these nucleotides of the invention include primers and primer pairs, which allow the specific amplification of polynucleotides of the invention or of any specific parts thereof, and probes that selectively or specifically hybridize to nucleic acid molecules of the invention or to any part thereof. Probes can be labeled with a detectable marker, such as, for example, a radioisotope, fluorescent compound, bioluminescent compound, a chemiluminescent compound, metal chelator or enzyme. Such probes and primers are used to detect the presence of a 24P4C12 polynucleotide in a sample and as a means for detecting a cell expressing a 24P4C12 protein.

Examples of such probes include polypeptides comprising all or part of the human 24P4C12 cDNA sequence shown in FIG. 2. Examples of primer pairs capable of specifically amplifying 24P4C12 mRNAs are also described in the Examples. As will be understood by the skilled artisan, a great many different primers and probes can be prepared based on the sequences provided herein and used effectively to amplify and/or detect a 24P4C12 mRNA.

The 24P4C12 polynucleotides of the invention are useful for a variety of purposes, including but not limited to their use as probes and primers for the amplification and/or detection of the 24P4C12 gene(s), mRNA(s), or fragments thereof; as reagents for the diagnosis and/or prognosis of prostate cancer and other cancers; as coding sequences capable of directing the expression of 24P4C12 polypeptides; as tools for modulating or inhibiting the expression of the 24P4C12 gene(s) and/or translation of the 24P4C12 transcript(s); and as therapeutic agents.

The present invention includes the use of any probe as described herein to identify and isolate a 24P4C12 or 24P4C12 related nucleic acid sequence from a naturally occurring source, such as humans or other mammals, as well as the isolated nucleic acid sequence per se, which would comprise all or most of the sequences found in the probe used.

II.A.4.) Isolation of 24P4C12-Encoding Nucleic Acid Molecules

The 24P4C12 cDNA sequences described herein enable the isolation of other polynucleotides encoding 24P4C12 gene product(s), as well as the isolation of polynucleotides encoding 24P4C12 gene product homologs, alternatively spliced isoforms, allelic variants, and mutant forms of a 24P4C12 gene product as well as polynucleotides that encode analogs of 24P4C12-related proteins. Various molecular cloning methods that can be employed to isolate full length cDNAs encoding a 24P4C12 gene are well known (see, for example, Sambrook, J. et al., *Molecular Cloning: A Laboratory Manual*, 2d edition, Cold Spring Harbor Press, New York, 1989; *Current Protocols in Molecular Biology*, Ausubel et al., Eds., Wiley and Sons, 1995). For example, lambda phage cloning methodologies can be conveniently employed, using commercially available cloning systems (e.g., Lambda ZAP Express, Stratagene). Phage clones containing 24P4C12 gene cDNAs can be identified by probing with a labeled 24P4C12 cDNA or a fragment thereof. For example, in one embodiment, a 24P4C12 cDNA (e.g., FIG. 2) or a portion thereof can be synthesized and used as a probe to retrieve overlapping and full-length cDNAs corresponding to a 24P4C12 gene. A 24P4C12 gene itself can be isolated by screening genomic DNA libraries, bacterial artificial chromosome libraries (BACs), yeast artificial chromosome libraries (YACs), and the like, with 24P4C12 DNA probes or primers.

II.A.5.) Recombinant Nucleic Acid Molecules and Host-Vector Systems

The invention also provides recombinant DNA or RNA molecules containing a 24P4C12 polynucleotide, a fragment, analog or homologue thereof, including but not limited to phages, plasmids, phagemids, cosmids, YACs, BACs, as well as various viral and non-viral vectors well known in the art, and cells transformed or transfected with such recombinant DNA or RNA molecules. Methods for generating such molecules are well known (see, for example, Sambrook et al., 1989, *supra*).

The invention further provides a host-vector system comprising a recombinant DNA molecule containing a 24P4C12 polynucleotide, fragment, analog or homologue thereof within a suitable prokaryotic or eukaryotic host cell. Examples of suitable eukaryotic host cells include a yeast cell, a plant cell, or an animal cell, such as a mammalian cell or an insect cell (e.g., a baculovirus-infectible cell such as an Sf9 or HighFive cell). Examples of suitable mammalian cells include various prostate cancer cell lines such as DU145 and TsuPr1, other transfectable or transducible prostate cancer cell lines, primary cells (PrEC), as well as a number of mammalian cells routinely used for the expression of recombinant proteins (e.g., COS, CHO, 293, 293T cells). More particularly, a polynucleotide comprising the coding sequence of 24P4C12 or a fragment, analog or homologue thereof can be used to generate 24P4C12 proteins or fragments thereof using any number of host-vector systems routinely used and widely known in the art.

A wide range of host-vector systems suitable for the expression of 24P4C12 proteins or fragments thereof are available, see for example, Sambrook et al., 1989, *supra*; *Current Protocols in Molecular Biology*, 1995, *supra*). Preferred vectors for mammalian expression include but are not limited to pcDNA 3.1 myc-His-tag (Invitrogen) and the retroviral vector pSRatKneo (Muller et al., 1991, *MCB* 11:1785). Using these expression vectors, 24P4C12 can be expressed in several prostate cancer and non-prostate cell lines, including for example 293, 293T, rat-1, NIH 3T3 and TsuPr1. The host-vector systems of the invention are useful

for the production of a 24P4C12 protein or fragment thereof. Such host-vector systems can be employed to study the functional properties of 24P4C12 and 24P4C12 mutations or analogs.

Recombinant human 24P4C12 protein or an analog or homolog or fragment thereof can be produced by mammalian cells transfected with a construct encoding a 24P4C12-related nucleotide. For example, 293T cells can be transfected with an expression plasmid encoding 24P4C12 or fragment, analog or homolog thereof, a 24P4C12-related protein is expressed in the 293T cells, and the recombinant 24P4C12 protein is isolated using standard purification methods (e.g., affinity purification using anti-24P4C12 antibodies). In another embodiment, a 24P4C12 coding sequence is subcloned into the retroviral vector pSR α MSVtkneo and used to infect various mammalian cell lines, such as NIH 3T3, TsuPr1, 293 and rat-1 in order to establish 24P4C12 expressing cell lines. Various other expression systems well known in the art can also be employed. Expression constructs encoding a leader peptide joined in frame to a 24P4C12 coding sequence can be used for the generation of a secreted form of recombinant 24P4C12 protein.

As discussed herein, redundancy in the genetic code permits variation in 24P4C12 gene sequences. In particular, it is known in the art that specific host species often have specific codon preferences, and thus one can adapt the disclosed sequence as preferred for a desired host. For example, preferred analog codon sequences typically have rare codons (i.e., codons having a usage frequency of less than about 20% in known sequences of the desired host) replaced with higher frequency codons. Codon preferences for a specific species are calculated, for example, by utilizing codon usage tables available on the INTERNET such as at URL dna.affrc.go.jp/~nakamura/codon.html.

Additional sequence modifications are known to enhance protein expression in a cellular host. These include elimination of sequences encoding spurious polyadenylation signals, exon/intron splice site signals, transposon-like repeats, and/or other such well-characterized sequences that are deleterious to gene expression. The GC content of the sequence is adjusted to levels average for a given cellular host, as calculated by reference to known genes expressed in the host cell. Where possible, the sequence is modified to avoid predicted hairpin secondary mRNA structures. Other useful modifications include the addition of a translational initiation consensus sequence at the start of the open reading frame, as described in Kozak, *Mol. Cell Biol.*, 9:5073-5080 (1989). Skilled artisans understand that the general rule that eukaryotic ribosomes initiate translation exclusively at the 5' proximal AUG codon is abrogated only under rare conditions (see, e.g., Kozak PNAS 92(7): 2662-2666, (1995) and Kozak NAR 15(20): 8125-8148 (1987)).

III.) 24P4C12-RELATED PROTEINS

Another aspect of the present invention provides 24P4C12-related proteins. Specific embodiments of 24P4C12 proteins comprise a polypeptide having all or part of the amino acid sequence of human 24P4C12 as shown in FIG. 2 or FIG. 3. Alternatively, embodiments of 24P4C12 proteins comprise variant, homolog or analog polypeptides that have alterations in the amino acid sequence of 24P4C12 shown in FIG. 2 or FIG. 3.

Embodiments of a 24P4C12 polypeptide include: a 24P4C12 polypeptide having a sequence shown in FIG. 2, a peptide sequence of a 24P4C12 as shown in FIG. 2 wherein T is U; at least 10 contiguous nucleotides of a polypeptide

having the sequence as shown in FIG. 2; or, at least 10 contiguous peptides of a polypeptide having the sequence as shown in FIG. 2 where T is U. For example, embodiments of 24P4C12 peptides comprise, without limitation:

- (I) a protein comprising, consisting essentially of, or consisting of an amino acid sequence as shown in FIG. 2A-I or FIGS. 3A-G;
- (II) a 24P4C12-related protein that is at least 90, 91, 92, 93, 94, 95, 96, 97, 98, 99 or 100% homologous to an entire amino acid sequence shown in FIGS. 2A-I;
- (III) a 24P4C12-related protein that is at least 90, 91, 92, 93, 94, 95, 96, 97, 98, 99 or 100% identical to an entire amino acid sequence shown in FIG. 2A-I or 3A-G;
- (IV) a protein that comprises at least one peptide set forth in Tables VIII to XLIX, optionally with a proviso that it is not an entire protein of FIG. 2;
- (V) a protein that comprises at least one peptide set forth in Tables VIII-XXI, collectively, which peptide is also set forth in Tables XXII to XLIX, collectively, optionally with a proviso that it is not an entire protein of FIG. 2;
- (VI) a protein that comprises at least two peptides selected from the peptides set forth in Tables VIII-XLIX, optionally with a proviso that it is not an entire protein of FIG. 2;
- (VII) a protein that comprises at least two peptides selected from the peptides set forth in Tables VIII to XLIX collectively, with a proviso that the protein is not a contiguous sequence from an amino acid sequence of FIG. 2;
- (VIII) a protein that comprises at least one peptide selected from the peptides set forth in Tables VIII-XXI; and at least one peptide selected from the peptides set forth in Tables XXII to XLIX, with a proviso that the protein is not a contiguous sequence from an amino acid sequence of FIG. 2;
- (IX) a polypeptide comprising at least 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35 amino acids of a protein of FIG. 3A, 3B, 3C, 3D, 3E, 3F, or 3G in any whole number increment up to 710, 710, 710, 710, 598, 722, or 712 respectively that includes at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35 amino acid position(s) having a value greater than 0.5 in the Hydrophilicity profile of FIG. 5;
- (X) a polypeptide comprising at least 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35 amino acids of a protein of FIG. 3A, 3B, 3C, 3D, 3E, 3F, or 3G in any whole number increment up to 710, 710, 710, 710, 598, 722, or 712 respectively, that includes at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35 amino acid position(s) having a value less than 0.5 in the Hydrophobicity profile of FIG. 6;
- (XI) a polypeptide comprising at least 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35 amino acids of a protein of FIG. 3A, 3B, 3C, 3D, 3E, 3F, or 3G in any whole number increment up to 710, 710, 710, 710, 598, 722, or 712 respectively, that includes at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35 amino acid position(s) having a value greater than 0.5 in the Percent Accessible Residues profile of FIG. 7;
- (XII) a polypeptide comprising at least 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35 amino acids of a protein

- of FIG. 3A, 3B, 3C, 3D, 3E, 3F, or 3G in any whole number increment up to 710, 710, 710, 710, 598, 722, or 712 respectively, that includes at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35-amino acid position(s) having a value greater than 0.5 in the Average Flexibility profile of FIG. 8;
- (XIII) a polypeptide comprising at least 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35 amino acids of a protein of FIG. 3A, 3B, 3C, 3D, 3E, 3F, or 3G in any whole number increment up to 710, 710, 710, 710, 598, 722, or 712 respectively, that includes at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35 amino acid position(s) having a value greater than 0.5 in the Beta-turn profile of FIG. 9;
- (XIV) a peptide that occurs at least twice in Tables VIII-XXI and XXII to XLIX, collectively;
- (XV) a peptide that occurs at least three times in Tables VIII-XXI and XXII to XLIX, collectively;
- (XVI) a peptide that occurs at least four times in Tables VIII-XXI and XXII to XLIX, collectively;
- (XVII) a peptide that occurs at least five times in Tables VIII-XXI and XXII to XLIX, collectively;
- (XVIII) a peptide that occurs at least once in Tables VIII-XXI, and at least once in tables XXII to XLIX;
- (XIX) a peptide that occurs at least once in Tables VIII-XXI, and at least twice in tables XXII to XLIX;
- (XX) a peptide that occurs at least twice in Tables VIII-XXI, and at least once in tables XXII to XLIX;
- (XXI) a peptide that occurs at least twice in Tables VIII-XXI, and at least twice in tables XXII to XLIX;
- (XXII) a peptide which comprises one two, three, four, or five of the following characteristics, or an oligonucleotide encoding such peptide:
- a region of at least 5 amino acids of a particular peptide of FIG. 3, in any whole number increment up to the full length of that protein in FIG. 3, that includes an amino acid position having a value equal to or greater than 0.5, 0.6, 0.7, 0.8, 0.9, or having a value equal to 1.0, in the Hydrophilicity profile of FIG. 5;
 - a region of at least 5 amino acids of a particular peptide of FIG. 3, in any whole number increment up to the full length of that protein in FIG. 3, that includes an amino acid position having a value equal to or less than 0.5, 0.4, 0.3, 0.2, 0.1, or having a value equal to 0.0, in the Hydrophobicity profile of FIG. 6;
 - a region of at least 5 amino acids of a particular peptide of FIG. 3, in any whole number increment up to the full length of that protein in FIG. 3, that includes an amino acid position having a value equal to or greater than 0.5, 0.6, 0.7, 0.8, 0.9, or having a value equal to 1.0, in the Percent Accessible Residues profile of FIG. 7;
 - a region of at least 5 amino acids of a particular peptide of FIG. 3, in any whole number increment up to the full length of that protein in FIG. 3, that includes an amino acid position having a value equal to or greater than 0.5, 0.6, 0.7, 0.8, 0.9, or having a value equal to 1.0, in the Average Flexibility profile of FIG. 8; or;
 - a region of at least 5 amino acids of a particular peptide of FIG. 3, in any whole number increment up to the full length of that protein in FIG. 3, that includes an amino acid position having a value equal to or

- greater than 0.5, 0.6, 0.7, 0.8, 0.9, or having a value equal to 1.0, in the Beta-turn profile of FIG. 9;
- (XXIII) a composition comprising a peptide of (I)-(XXII) or an antibody or binding region thereof together with a pharmaceutical excipient and/or in a human unit dose form.
- (XXIV) a method of using a peptide of (I)-(XXII), or an antibody or binding region thereof or a composition of (XXIII) in a method to modulate a cell expressing 24P4C12,
- (XXV) a method of using a peptide of (I)-(XXII) or an antibody or binding region thereof or a composition of (XXIII) in a method to diagnose, prophylax, prognose, or treat an individual who bears a cell expressing 24P4C12
- (XXVI) a method of using a peptide of (I)-(XXII) or an antibody or binding region thereof or a composition (XXIII) in a method to diagnose, prophylax, prognose, or treat an individual who bears a cell expressing 24P4C12, said cell from a cancer of a tissue listed in Table I;
- (XXVII) a method of using a peptide of (I)-(XXII) or an antibody or binding region thereof or a composition of (XXIII) in a method to diagnose, prophylax, prognose, or treat a cancer;
- (XXVIII) a method of using a peptide of (I)-(XXII) or an antibody or binding region thereof or a composition of (XXIII) in a method to diagnose, prophylax, prognose, or treat a cancer of a tissue listed in Table I; and,
- (XXIX) a method of using a peptide of (I)-(XXII) or an antibody or binding region thereof or a composition (XXIII) in a method to identify or characterize a modulator of a cell expressing 24P4C12.

As used herein, a range is understood to specifically disclose all whole unit positions thereof.

Typical embodiments of the invention disclosed herein include 24P4C12 polynucleotides that encode specific portions of 24P4C12 mRNA sequences (and those which are complementary to such sequences) such as those that encode the proteins and/or fragments thereof, for example:

(a) 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 105, 110, 115, 120, 125, 130, 135, 140, 145, 150, 155, 160, 165, 170, 175, 180, 185, 190, 195, 200, 225, 250, 275, 300, 325, 350, 375, 400, 425, 450, 475, 500, 525, 550, 575, 600, 625, 650, 675, 700, 710 or more contiguous amino acids of 24P4C12 variant 1; the maximal lengths relevant for other variants are: variant 3, 710 amino acids; variant 5, 710 amino acids, variant 6, 710, variant 7, 598 amino acids, variant 8, 722 amino acids, and variant 9, 712 amino acids.

In general, naturally occurring allelic variants of human 24P4C12 share a high degree of structural identity and homology (e.g., 90% or more homology). Typically, allelic variants of a 24P4C12 protein contain conservative amino acid substitutions within the 24P4C12 sequences described herein or contain a substitution of an amino acid from a corresponding position in a homologue of 24P4C12. One class of 24P4C12 allelic variants are proteins that share a high degree of homology with at least a small region of a particular 24P4C12 amino acid sequence, but further contain a radical departure from the sequence, such as a non-conservative substitution, truncation, insertion or frame shift. In comparisons of protein sequences, the terms, similarity, identity, and homology each have a distinct meaning as appreciated in the field of genetics. Moreover, orthology and paralogy can be important concepts describing the relationship of members of

a given protein family in one organism to the members of the same family in other organisms.

Amino acid abbreviations are provided in Table II. Conservative amino acid substitutions can frequently be made in a protein without altering either the conformation or the function of the protein. Proteins of the invention can comprise 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15 conservative substitutions. Such changes include substituting any of isoleucine (I), valine (V), and leucine (L) for any other of these hydrophobic amino acids; aspartic acid (D) for glutamic acid (E) and vice versa; glutamine (Q) for asparagine (N) and vice versa; and serine (S) for threonine (T) and vice versa. Other substitutions can also be considered conservative, depending on the environment of the particular amino acid and its role in the three-dimensional structure of the protein. For example, glycine (G) and alanine (A) can frequently be interchangeable, as can alanine (A) and valine (V). Methionine (M), which is relatively hydrophobic, can frequently be interchanged with leucine and isoleucine, and sometimes with valine. Lysine (K) and arginine (R) are frequently interchangeable in locations in which the significant feature of the amino acid residue is its charge and the differing pK's of these two amino acid residues are not significant. Still other changes can be considered "conservative" in particular environments (see, e.g. Table III herein; pages 13-15 "Biochemistry" 2nd ED. Lubert Stryer ed (Stanford University); Henikoff et al., PNAS 1992 Vol 89 10915-10919; Lei et al., J Biol Chem 1995 May 19; 270(20):11882-6).

Embodiments of the invention disclosed herein include a wide variety of art-accepted variants or analogs of 24P4C12 proteins such as polypeptides having amino acid insertions, deletions and substitutions. 24P4C12 variants can be made using methods known in the art such as site-directed mutagenesis, alanine scanning, and PCR mutagenesis. Site-directed mutagenesis (Carter et al., *Nucl. Acids Res.*, 13:4331 (1986); Zoller et al., *Nucl. Acids Res.*, 10:6487 (1987)), cassette mutagenesis (Wells et al., *Gene*, 34:315 (1985)), restriction selection mutagenesis (Wells et al., *Philos. Trans. R. Soc. London Ser.A*, 317:415 (1986)) or other known techniques can be performed on the cloned DNA to produce the 24P4C12 variant DNA.

Scanning amino acid analysis can also be employed to identify one or more amino acids along a contiguous sequence that is involved in a specific biological activity such as a protein-protein interaction. Among the preferred scanning amino acids are relatively small, neutral amino acids. Such amino acids include alanine, glycine, serine, and cysteine. Alanine is typically a preferred scanning amino acid among this group because it eliminates the side-chain beyond the beta-carbon and is less likely to alter the main-chain conformation of the variant. Alanine is also typically preferred because it is the most common amino acid. Further, it is frequently found in both buried and exposed positions (Creighton, *The Proteins*, (W.H. Freeman & Co., N.Y.); Chothia, J. Mol. Biol., 150:1 (1976)). If alanine substitution does not yield adequate amounts of variant, an isosteric amino acid can be used.

As defined herein, 24P4C12 variants, analogs or homologs, have the distinguishing attribute of having at least one epitope that is "cross reactive" with a 24P4C12 protein having an amino acid sequence of FIG. 3. As used in this sentence, "cross reactive" means that an antibody or T cell that specifically binds to a 24P4C12 variant also specifically binds to a 24P4C12 protein having an amino acid sequence set forth in FIG. 3. A polypeptide ceases to be a variant of a protein shown in FIG. 3, when it no longer contains any epitope capable of being recognized by an antibody or T cell

that specifically binds to the starting 24P4C12 protein. Those skilled in the art understand that antibodies that recognize proteins bind to epitopes of varying size, and a grouping of the order of about four or five amino acids, contiguous or not, is regarded as a typical number of amino acids in a minimal epitope. See, e.g., Nair et al., *J. Immunol.* 2000 165(12): 6949-6955; Hebbes et al., *Mol Immunol* (1989) 26(9):865-73; Schwartz et al., *J Immunol* (1985) 135(4):2598-608.

Other classes of 24P4C12-related protein variants share 70%, 75%, 80%, 85% or 90% or more similarity with an amino acid sequence of FIG. 3, or a fragment thereof. Another specific class of 24P4C12 protein variants or analogs comprises one or more of the 24P4C12 biological motifs described herein or presently known in the art. Thus, encompassed by the present invention are analogs of 24P4C12 fragments (nucleic or amino acid) that have altered functional (e.g. immunogenic) properties relative to the starting fragment. It is to be appreciated that motifs now or which become part of the art are to be applied to the nucleic or amino acid sequences of FIG. 2 or FIG. 3.

As discussed herein, embodiments of the claimed invention include polypeptides containing less than the full amino acid sequence of a 24P4C12 protein shown in FIG. 2 or FIG. 3. For example, representative embodiments of the invention comprise peptides/proteins having any 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15 or more contiguous amino acids of a 24P4C12 protein shown in FIG. 2 or FIG. 3.

Moreover, representative embodiments of the invention disclosed herein include polypeptides consisting of about amino acid 1 to about amino acid 10 of a 24P4C12 protein shown in FIG. 2 or FIG. 3, polypeptides consisting of about amino acid 10 to about amino acid 20 of a 24P4C12 protein shown in FIG. 2 or FIG. 3, polypeptides consisting of about amino acid 20 to about amino acid 30 of a 24P4C12 protein shown in FIG. 2 or FIG. 3, polypeptides consisting of about amino acid 30 to about amino acid 40 of a 24P4C12 protein shown in FIG. 2 or FIG. 3, polypeptides consisting of about amino acid 40 to about amino acid 50 of a 24P4C12 protein shown in FIG. 2 or FIG. 3, polypeptides consisting of about amino acid 50 to about amino acid 60 of a 24P4C12 protein shown in FIG. 2 or FIG. 3, polypeptides consisting of about amino acid 60 to about amino acid 70 of a 24P4C12 protein shown in FIG. 2 or FIG. 3, polypeptides consisting of about amino acid 70 to about amino acid 80 of a 24P4C12 protein shown in FIG. 2 or FIG. 3, polypeptides consisting of about amino acid 80 to about amino acid 90 of a 24P4C12 protein shown in FIG. 2 or FIG. 3, polypeptides consisting of about amino acid 90 to about amino acid 100 of a 24P4C12 protein shown in FIG. 2 or FIG. 3, etc. throughout the entirety of a 24P4C12 amino acid sequence. Moreover, polypeptides consisting of about amino acid 1 (or 20 or 30 or 40 etc.) to about amino acid 20, (or 130, or 140 or 150 etc.) of a 24P4C12 protein shown in FIG. 2 or FIG. 3 are embodiments of the invention. It is to be appreciated that the starting and stopping positions in this paragraph refer to the specified position as well as that position plus or minus 5 residues.

24P4C12-related proteins are generated using standard peptide synthesis technology or using chemical cleavage methods well known in the art. Alternatively, recombinant methods can be used to generate nucleic acid molecules that encode a 24P4C12-related protein. In one embodiment, nucleic acid molecules provide a means to generate defined fragments of a 24P4C12 protein (or variants, homologs or analogs thereof).

III.A.) Motif-Bearing Protein Embodiments

Additional illustrative embodiments of the invention disclosed herein include 24P4C12 polypeptides comprising the

Figure 2B-3

```
700   E A P P D N K K R K K *
2101 ACGAGGCGCCCCCGGACAACAAGAAGAGGAAGAAGTGAacagctccggccctgatccagga
2161 ctgcacccccacccccaccgtccagccatccaacctcacttcgccttacagggtctccattt
2221 tgtggtaaaaaaagggttttaggccaggcgccgtgggtcacgcctgtaatccaacactttg
2281 agaggctgaggcgggaggatcacctgagtcaggagttcgagaccagcctggccaacatgg
2341 tgaaacctccgtctctattataaaaatacaaaaattagccgagagtgggtggcatgcacctgt
2401 catcccagctactcgggaggctgaggcaggagaatcgcttgaacccgggaggcagagggtt
2461 gcagtgagccgagatcgcgccactgcactccaacctgggtgacagactctgtctccaaaa
2521 caaaacaacaacaacaaaagattttattaaagatattttgttaactcagtaaaaaaaaaa
2581 aaaaaaa
```

Figure 2C. The cDNA (SEQ ID. NO. : 6) and amino acid sequence (SEQ ID. NO. : 7) of 24P4C12 v.3.

The start methionine is underlined. The open reading frame extends from nucleic acid 6-2138 including the stop codon.

Figure 2C-1

```
1       M G G K Q R D E D D E A Y G K P V K Y
1 gagccATGGGGGAAAGCAGCGGGACGAGGATGACGAGGCCTACGGGAAGCCAGTCAAAT
20    D P S F R G P I R N R S C T D V I C C V
61 ACGACCCCTCCTTTCGAGGCCCCATCAAGAACAGAAGCTGCACAGATGTCATCTGCTGCG
40    L F L L F I L G Y I V V G I V A W L Y G
121 TCCTCTCTCTGCTCTTTCATTCTAGGTTACATCGTGGTGGGGATTGTGGCCTGTTGTATG
60    D P R Q V L Y P R N S T G A Y C G M G E
181 GAGACCCCGGCAAGTCTCTTACCCAGGAACCTCTACTGGGGCCTACTGTGGCATGGGGG
80    N K D K P Y L L Y F N I F S C I L S S N
241 AGAACAAAGATAAGCCGTAICTCTCTGTACTTCAACATCTTCAGTGCATCCIGTCCAGCA
100   I I S V A E N G L Q C P T P Q V C V S S
301 ACAICATCTCAGTTGCTGAGAACGGCCTACAGTGCCCCACACCCAGGTGTGTGTGTCCT
120   C P E D P W I V G K N E F S Q I V G E V
361 CCTGCCCCGAGGACCCATGGACTGTGGGAAAAACGAGTTCTCAGACTGTTGGGGAAG
140   F Y T K N R N F C L P G V P W N M T V I
421 TCTTCTATACAAAAACAGGAACCTTTTGTCTGCCAGGGGTACCCTGGAATATGACGGTGA
160   T S L Q Q E L C P S F L L P S A P A L G
481 TCACAAGCCTGCAACAGGAACCTGCCCCAGTTTCTCTCTCCCTCTGCTCCAGCTCTGG
180   R C F P W T N I T P P A L P G I T N D T
541 GGCGCTGCTTTTCCATGGACCAACATTACTCCACCGGCGCTCCAGGGATCACCAATGACA
200   T I Q Q G I S G L I D S L N A R D I S V
601 CCACCATACAGCAGGGGATCAGCGGTCTTATTGACAGCCTCAATGCCCCGAGACATCAGTG
220   K I F E D F A Q S W Y W I L V A L G V A
661 TTAAGATCTTTGAAGATTTTCCCCAGTCCCTGGTATTGGATTCTTGTGCCCCTGGGGGTGG
240   L V L S L L F I L L L R L V A G P L V L
```

amino acid residues of one or more of the biological motifs contained within a 24P4C12 polypeptide sequence set forth in FIG. 2 or FIG. 3. Various motifs are known in the art, and a protein can be evaluated for the presence of such motifs by a number of publicly available Internet sites (see, e.g., URL addresses: pfam.wustl.edu/; searchlauncher.bcm.tmc.edu/seq-search/struc-predict.html; psort.ims.u-tokyo.ac.jp/; cbs.dtu.dk/; ebi.ac.uk/interpro/scan.html; expasy.ch/tools/scnpsit1.html; Epimatrix™ and Epimer™, Brown University, brown.edu/Research/TB-HIV_Lab/epimatrix/epimatrix.html; and BIMAS, bimas.dcrn.nih.gov/).

Motif bearing subsequences of all 24P4C12 variant proteins are set forth and identified in Tables VIII-XXI and XXII-XLIX.

Table V sets forth several frequently occurring motifs based on pfam searches (see URL address pfam.wustl.edu/). The columns of Table V list (1) motif name abbreviation, (2) percent identity found amongst the different member of the motif family, (3) motif name or description and (4) most common function; location information is included if the motif is relevant for location.

Polypeptides comprising one or more of the 24P4C12 motifs discussed above are useful in elucidating the specific characteristics of a malignant phenotype in view of the observation that the 24P4C12 motifs discussed above are associated with growth dysregulation and because 24P4C12 is over-expressed in certain cancers (See, e.g., Table I). Casein kinase II, cAMP and camp-dependent protein kinase, and Protein Kinase C, for example, are enzymes known to be associated with the development of the malignant phenotype (see e.g. Chen et al., *Lab Invest.*, 78(2): 165-174 (1998); Gaiddon et al., *Endocrinology* 136(10): 4331-4338 (1995); Hall et al., *Nucleic Acids Research* 24(6): 1119-1126 (1996); Peterziel et al., *Oncogene* 18(46): 6322-6329 (1999) and O'Brian, *Oncol. Rep.* 5(2): 305-309 (1998)). Moreover, both glycosylation and myristoylation are protein modifications also associated with cancer and cancer progression (see e.g. Dennis et al., *Biochem. Biophys. Acta* 1473(1):21-34 (1999); Raju et al., *Exp. Cell Res.* 235(1): 145-154 (1997)). Amidation is another protein modification also associated with cancer and cancer progression (see e.g. Treston et al., *J. Natl. Cancer Inst. Monogr.* (13): 169-175 (1992)).

In another embodiment, proteins of the invention comprise one or more of the immunoreactive epitopes identified in accordance with art-accepted methods, such as the peptides set forth in Tables VIII-XXI and XXII-XLIX. CTL epitopes can be determined using specific algorithms to identify peptides within a 24P4C12 protein that are capable of optimally binding to specified HLA alleles (e.g., Table IV; Epimatrix™ and Epimer™, Brown University, URL brown.edu/Research/TB-HIV_Lab/epimatrix/epimatrix.html; and BIMAS, URL bimas.dcrn.nih.gov/.) Moreover, processes for identifying peptides that have sufficient binding affinity for HLA molecules and which are correlated with being immunogenic epitopes, are well known in the art, and are carried out without undue experimentation. In addition, processes for identifying peptides that are immunogenic epitopes, are well known in the art, and are carried out without undue experimentation either in vitro or in vivo.

Also known in the art are principles for creating analogs of such epitopes in order to modulate immunogenicity. For example, one begins with an epitope that bears a CTL or HTL motif (see, e.g., the HLA Class I and HLA Class II motifs/supermotifs of Table IV). The epitope is analogized by substituting out an amino acid at one of the specified positions, and replacing it with another amino acid specified for that position. For example, on the basis of residues defined in Table IV,

one can substitute out a deleterious residue in favor of any other residue, such as a preferred residue; substitute a less-preferred residue with a preferred residue; or substitute an originally-occurring preferred residue with another preferred residue. Substitutions can occur at primary anchor positions or at other positions in a peptide; see, e.g., Table IV.

A variety of references reflect the art regarding the identification and generation of epitopes in a protein of interest as well as analogs thereof. See, for example, WO 97/33602 to Chesnut et al.; Sette, *Immunogenetics* 1999 50(3-4): 201-212; Sette et al., *J. Immunol.* 2001 166(2): 1389-1397; Sidney et al., *Hum. Immunol.* 1997 58(1): 12-20; Kondo et al., *Immunogenetics* 1997 45(4): 249-258; Sidney et al., *J. Immunol.* 1996 157(8): 3480-90; and Falk et al., *Nature* 351: 290-6 (1991); Hunt et al., *Science* 255:1261-3 (1992); Parker et al., *J. Immunol.* 149:3580-7 (1992); Parker et al., *J. Immunol.* 152:163-75 (1994); Kast et al., 1994 152(8): 3904-12; Borras-Cuesta et al., *Hum. Immunol.* 2000 61(3): 266-278; Alexander et al., *J. Immunol.* 2000 164(3): 1625-1633; Alexander et al., PMID: 7895164, UI: 95202582; O'Sullivan et al., *J. Immunol.* 1991 147(8): 2663-2669; Alexander et al., *Immunity* 1994 1(9): 751-761 and Alexander et al., *Immunol. Res.* 1998 18(2): 79-92.

Related embodiments of the invention include polypeptides comprising combinations of the different motifs set forth in Table VI, and/or, one or more of the predicted CTL epitopes of Tables VIII-XXI and XXII-XLIX, and/or, one or more of the predicted HTL epitopes of Tables XLVI-XLIX, and/or, one or more of the T cell binding motifs known in the art. Preferred embodiments contain no insertions, deletions or substitutions either within the motifs or within the intervening sequences of the polypeptides. In addition, embodiments which include a number of either N-terminal and/or C-terminal amino acid residues on either side of these motifs may be desirable (to, for example, include a greater portion of the polypeptide architecture in which the motif is located). Typically, the number of N-terminal and/or C-terminal amino acid residues on either side of a motif is between about 1 to about 100 amino acid residues, preferably 5 to about 50 amino acid residues.

24P4C12-related proteins are embodied in many forms, preferably in isolated form. A purified 24P4C12 protein molecule will be substantially free of other proteins or molecules that impair the binding of 24P4C12 to antibody, T cell or other ligand. The nature and degree of isolation and purification will depend on the intended use. Embodiments of a 24P4C12-related proteins include purified 24P4C12-related proteins and functional, soluble 24P4C12-related proteins. In one embodiment, a functional, soluble 24P4C12 protein or fragment thereof retains the ability to be bound by antibody, T cell or other ligand.

The invention also provides 24P4C12 proteins comprising biologically active fragments of a 24P4C12 amino acid sequence shown in FIG. 2 or FIG. 3. Such proteins exhibit properties of the starting 24P4C12 protein, such as the ability to elicit the generation of antibodies that specifically bind an epitope associated with the starting 24P4C12 protein; to be bound by such antibodies; to elicit the activation of HTL or CTL; and/or, to be recognized by HTL or CTL that also specifically bind to the starting protein.

24P4C12-related polypeptides that contain particularly interesting structures can be predicted and/or identified using various analytical techniques well known in the art, including, for example, the methods of Chou-Fasman, Garnier-Robson, Kyte-Doolittle, Eisenberg, Karplus-Schultz or Jameson-Wolf analysis, or based on immunogenicity. Fragments that contain such structures are particularly useful in

generating subunit-specific anti-24P4C12 antibodies or T cells or in identifying cellular factors that bind to 24P4C12. For example, hydrophilicity profiles can be generated, and immunogenic peptide fragments identified, using the method of Hopp, T. P. and Woods, K. R., 1981, *Proc. Natl. Acad. Sci. U.S.A.* 78:3824-3828. Hydrophobicity profiles can be generated, and immunogenic peptide fragments identified, using the method of Kyte, J. and Doolittle, R. F., 1982, *J. Mol. Biol.* 157:105-132. Percent (%) Accessible Residues profiles can be generated, and immunogenic peptide fragments identified, using the method of Janin J., 1979, *Nature* 277:491-492. Average Flexibility profiles can be generated, and immunogenic peptide fragments identified, using the method of Bhaskaran R., Ponnuswamy P. K., 1988, *Int. J. Pept. Protein Res.* 32:242-255. Beta-turn profiles can be generated, and immunogenic peptide fragments identified, using the method of Deleage, G., Roux B., 1987, *Protein Engineering* 1:289-294.

CTL epitopes can be determined using specific algorithms to identify peptides within a 24P4C12 protein that are capable of optimally binding to specified HLA alleles (e.g., by using the SYFPEITHI site at World Wide Web URL syfpeithi.bmi-heidelberg.com/; the listings in Table IV(A)-(E); Epimatrix™ and Epimer™, Brown University, URL (brown.edu/Research/TB-HIV_Lab/epimatrix/epimatrix.html); and BIMAS, URL bimas.dcrf.nih.gov/). Illustrating this, peptide epitopes from 24P4C12 that are presented in the context of human MHC Class I molecules, e.g., HLA-A1, A2, A3, A11, A24, B7 and B35 were predicted (see, e.g., Tables VIII-XXI, XXII-XLIX). Specifically, the complete amino acid sequence of the 24P4C12 protein and relevant portions of other variants, i.e., for HLA Class I predictions 9 flanking residues on either side of a point mutation or exon junction, and for HLA Class II predictions 14 flanking residues on either side of a point mutation or exon junction corresponding to that variant, were entered into the HLA Peptide Motif Search algorithm found in the Bioinformatics and Molecular Analysis Section (BIMAS) web site listed above; in addition to the site SYFPEITHI, at URL syfpeithi.bmi-heidelberg.com/.

The HLA peptide motif search algorithm was developed by Dr. Ken Parker based on binding of specific peptide sequences in the groove of HLA Class I molecules, in particular HLA-A2 (see, e.g., Falk et al., *Nature* 351: 290-6 (1991); Hunt et al., *Science* 255:1261-3 (1992); Parker et al., *J. Immunol.* 149:3580-7 (1992); Parker et al., *J. Immunol.* 152:163-75 (1994)). This algorithm allows location and ranking of 8-mer, 9-mer, and 10-mer peptides from a complete protein sequence for predicted binding to HLA-A2 as well as numerous other HLA Class I molecules. Many HLA class I binding peptides are 8-, 9-, 10 or 11-mers. For example, for Class I HLA-A2, the epitopes preferably contain a leucine (L) or methionine (M) at position 2 and a valine (V) or leucine (L) at the C-terminus (see, e.g., Parker et al., *J. Immunol.* 149: 3580-7 (1992)). Selected results of 24P4C12 predicted binding peptides are shown in Tables VIII-XXI and XXII-XLIX herein. In Tables VIII-XXI and XXII-XLVII, selected candidates, 9-mers and 10-mers, for each family member are shown along with their location, the amino acid sequence of each specific peptide, and an estimated binding score. In Tables XLVI-XLIX, selected candidates, 15-mers, for each family member are shown along with their location, the amino acid sequence of each specific peptide, and an estimated binding score. The binding score corresponds to the estimated half time of dissociation of complexes containing the peptide at 37° C. at pH 6.5. Peptides with the highest binding score are predicted to be the most tightly bound to

HLA Class I on the cell surface for the greatest period of time and thus represent the best immunogenic targets for T-cell recognition.

Actual binding of peptides to an HLA allele can be evaluated by stabilization of HLA expression on the antigen-processing defective cell line T2 (see, e.g., Xue et al., *Prostate* 30:73-8 (1997) and Peshwa et al., *Prostate* 36:129-38 (1998)). Immunogenicity of specific peptides can be evaluated in vitro by stimulation of CD8+ cytotoxic T lymphocytes (CTL) in the presence of antigen presenting cells such as dendritic cells.

It is to be appreciated that every epitope predicted by the BIMAS site, Epimer™ and Epimatrix™ sites, or specified by the HLA class I or class II motifs available in the art or which become part of the art such as set forth in Table IV (or determined using World Wide Web site URL syfpeithi.bmi-heidelberg.com/, or BIMAS, bimas.dcrf.nih.gov/) are to be "applied" to a 24P4C12 protein in accordance with the invention. As used in this context "applied" means that a 24P4C12 protein is evaluated, e.g., visually or by computer-based patterns finding methods, as appreciated by those of skill in the relevant art. Every subsequence of a 24P4C12 protein of 8, 9, 10, or 11 amino acid residues that bears an HLA Class I motif, or a subsequence of 9 or more amino acid residues that bear an HLA Class II motif are within the scope of the invention.

III.B.) Expression of 24P4C12-Related Proteins

In an embodiment described in the examples that follow, 24P4C12 can be conveniently expressed in cells (such as 293T cells) transfected with a commercially available expression vector such as a CMV-driven expression vector encoding 24P4C12 with a C-terminal 6xHis and MYC tag (pcDNA3.1/mycHis, Invitrogen or Tag5, GenHunter Corporation, Nashville Tenn.). The Tag5 vector provides an IgGK secretion signal that can be used to facilitate the production of a secreted 24P4C12 protein in transfected cells. The secreted HIS-tagged 24P4C12 in the culture media can be purified, e.g., using a nickel column using standard techniques.

III.C.) Modifications of 24P4C12-Related Proteins

Modifications of 24P4C12-related proteins such as covalent modifications are included within the scope of this invention. One type of covalent modification includes reacting targeted amino acid residues of a 24P4C12 polypeptide with an organic derivatizing agent that is capable of reacting with selected side chains or the N- or C-terminal residues of a 24P4C12 protein. Another type of covalent modification of a 24P4C12 polypeptide included within the scope of this invention comprises altering the native glycosylation pattern of a protein of the invention. Another type of covalent modification of 24P4C12 comprises linking a 24P4C12 polypeptide to one of a variety of nonproteinaceous polymers, e.g., polyethylene glycol (PEG), polypropylene glycol, or polyoxyalkylenes, in the manner set forth in U.S. Pat. No. 4,640,835; 4,496,689; 4,301,144; 4,670,417; 4,791,192 or 4,179,337.

The 24P4C12-related proteins of the present invention can also be modified to form a chimeric molecule comprising 24P4C12 fused to another, heterologous polypeptide or amino acid sequence. Such a chimeric molecule can be synthesized chemically or recombinantly. A chimeric molecule can have a protein of the invention fused to another tumor-associated antigen or fragment thereof. Alternatively, a protein in accordance with the invention can comprise a fusion of fragments of a 24P4C12 sequence (amino or nucleic acid) such that a molecule is created that is not, through its length, directly homologous to the amino or nucleic acid sequences shown in FIG. 2 or FIG. 3. Such a chimeric molecule can comprise multiples of the same subsequence of 24P4C12. A chimeric molecule can comprise a fusion of a 24P4C12-

related protein with a polyhistidine epitope tag, which provides an epitope to which immobilized nickel can selectively bind, with cytokines or with growth factors. The epitope tag is generally placed at the amino- or carboxyl-terminus of a 24P4C12 protein. In an alternative embodiment, the chimeric molecule can comprise a fusion of a 24P4C12-related protein with an immunoglobulin or a particular region of an immunoglobulin. For a bivalent form of the chimeric molecule (also referred to as an "immunoadhesin"), such a fusion could be to the Fc region of an IgG molecule. The Ig fusions preferably include the substitution of a soluble (transmembrane domain deleted or inactivated) form of a 24P4C12 polypeptide in place of at least one variable region within an Ig molecule. In a preferred embodiment, the immunoglobulin fusion includes the hinge, CH2 and CH3, or the hinge, CH1, CH2 and CH3 regions of an IgG1 molecule. For the production of immunoglobulin fusions see, e.g., U.S. Pat. No. 5,428,130 issued Jun. 27, 1995.

III.D.) Uses of 24P4C12-Related Proteins

The proteins of the invention have a number of different specific uses. As 24P4C12 is highly expressed in prostate and other cancers, 24P4C12-related proteins are used in methods that assess the status of 24P4C12 gene products in normal versus cancerous tissues, thereby elucidating the malignant phenotype. Typically, polypeptides from specific regions of a 24P4C12 protein are used to assess the presence of perturbations (such as deletions, insertions, point mutations etc.) in those regions (such as regions containing one or more motifs). Exemplary assays utilize antibodies or T cells targeting 24P4C12-related proteins comprising the amino acid residues of one or more of the biological motifs contained within a 24P4C12 polypeptide sequence in order to evaluate the characteristics of this region in normal versus cancerous tissues or to elicit an immune response to the epitope. Alternatively, 24P4C12-related proteins that contain the amino acid residues of one or more of the biological motifs in a 24P4C12 protein are used to screen for factors that interact with that region of 24P4C12.

24P4C12 protein fragments/subsequences are particularly useful in generating and characterizing domain-specific antibodies (e.g., antibodies recognizing an extracellular or intracellular epitope of a 24P4C12 protein), for identifying agents or cellular factors that bind to 24P4C12 or a particular structural domain thereof, and in various therapeutic and diagnostic contexts, including but not limited to diagnostic assays, cancer vaccines and methods of preparing such vaccines.

Proteins encoded by the 24P4C12 genes, or by analogs, homologs or fragments thereof, have a variety of uses, including but not limited to generating antibodies and in methods for identifying ligands and other agents and cellular constituents that bind to a 24P4C12 gene product. Antibodies raised against a 24P4C12 protein or fragment thereof are useful in diagnostic and prognostic assays, and imaging methodologies in the management of human cancers characterized by expression of 24P4C12 protein, such as those listed in Table I. Such antibodies can be expressed intracellularly and used in methods of treating patients with such cancers. 24P4C12-related nucleic acids or proteins are also used in generating HTL or CTL responses.

Various immunological assays useful for the detection of 24P4C12 proteins are used, including but not limited to various types of radioimmunoassays, enzyme-linked immunosorbent assays (ELISA), enzyme-linked immunofluorescent assays (ELIFA), immunocytochemical methods, and the like. Antibodies can be labeled and used as immunological imaging reagents capable of detecting 24P4C12-expressing cells (e.g., in radioscintigraphic imaging methods). 24P4C12 pro-

teins are also particularly useful in generating cancer vaccines, as further described herein.

IV.) 24P4C12 ANTIBODIES

Another aspect of the invention provides antibodies that bind to 24P4C12-related proteins. Preferred antibodies specifically bind to a 24P4C12-related protein and do not bind (or bind weakly) to peptides or proteins that are not 24P4C12-related proteins. For example, antibodies that bind 24P4C12 can bind 24P4C12-related proteins such as the homologs or analogs thereof.

24P4C12 antibodies of the invention are particularly useful in cancer (see, e.g., Table I) diagnostic and prognostic assays, and imaging methodologies. Similarly, such antibodies are useful in the treatment, diagnosis, and/or prognosis of other cancers, to the extent 24P4C12 is also expressed or overexpressed in these other cancers. Moreover, intracellularly expressed antibodies (e.g., single chain antibodies) are therapeutically useful in treating cancers in which the expression of 24P4C12 is involved, such as advanced or metastatic prostate cancers.

The invention also provides various immunological assays useful for the detection and quantification of 24P4C12 and mutant 24P4C12-related proteins. Such assays can comprise one or more 24P4C12 antibodies capable of recognizing and binding a 24P4C12-related protein, as appropriate. These assays are performed within various immunological assay formats well known in the art, including but not limited to various types of radioimmunoassays, enzyme-linked immunosorbent assays (ELISA), enzyme-linked immunofluorescent assays (ELIFA), and the like.

Immunological non-antibody assays of the invention also comprise T cell immunogenicity assays (inhibitory or stimulatory) as well as major histocompatibility complex (MHC) binding assays.

In addition, immunological imaging methods capable of detecting prostate cancer and other cancers expressing 24P4C12 are also provided by the invention, including but not limited to radioscintigraphic imaging methods using labeled 24P4C12 antibodies. Such assays are clinically useful in the detection, monitoring, and prognosis of 24P4C12 expressing cancers such as prostate cancer.

24P4C12 antibodies are also used in methods for purifying a 24P4C12-related protein and for isolating 24P4C12 homologues and related molecules. For example, a method of purifying a 24P4C12-related protein comprises incubating a 24P4C12 antibody, which has been coupled to a solid matrix, with a lysate or other solution containing a 24P4C12-related protein under conditions that permit the 24P4C12 antibody to bind to the 24P4C12-related protein; washing the solid matrix to eliminate impurities; and eluting the 24P4C12-related protein from the coupled antibody. Other uses of 24P4C12 antibodies in accordance with the invention include generating anti-idiotypic antibodies that mimic a 24P4C12 protein.

Various methods for the preparation of antibodies are well known in the art. For example, antibodies can be prepared by immunizing a suitable mammalian host using a 24P4C12-related protein, peptide, or fragment, in isolated or immunconjugated form (Antibodies: A Laboratory Manual, CSH Press, Eds., Harlow, and Lane (1988); Harlow, Antibodies, Cold Spring Harbor Press, NY (1989)). In addition, fusion proteins of 24P4C12 can also be used, such as a 24P4C12 GST-fusion protein. In a particular embodiment, a GST fusion protein comprising all or most of the amino acid sequence of FIG. 2 or FIG. 3 is produced, then used as an

immunogen to generate appropriate antibodies. In another embodiment, a 24P4C12-related protein is synthesized and used as an immunogen.

In addition, naked DNA immunization techniques known in the art are used (with or without purified 24P4C12-related protein or 24P4C12 expressing cells) to generate an immune response to the encoded immunogen (for review, see Donnelly et al., 1997, *Ann. Rev. Immunol.* 15: 617-648).

The amino acid sequence of a 24P4C12 protein as shown in FIG. 2 or FIG. 3 can be analyzed to select specific regions of the 24P4C12 protein for generating antibodies. For example, hydrophobicity and hydrophilicity analyses of a 24P4C12 amino acid sequence are used to identify hydrophilic regions in the 24P4C12 structure. Regions of a 24P4C12 protein that show immunogenic structure, as well as other regions and domains, can readily be identified using various other methods known in the art, such as Chou-Fasman, Garnier-Robson, Kyte-Doolittle, Eisenberg, Karplus-Schultz or Jameson-Wolf analysis. Hydrophilicity profiles can be generated using the method of Hopp, T. P. and Woods, K. R., 1981, *Proc. Natl. Acad. Sci. U.S.A.* 78:3824-3828. Hydropathicity profiles can be generated using the method of Kyte, J. and Doolittle, R.F., 1982, *J. Mol. Biol.* 157:105-132. Percent (%) Accessible Residues profiles can be generated using the method of Janin J., 1979, *Nature* 277:491-492. Average Flexibility profiles can be generated using the method of Bhaskaran R., Ponuswamy P. K., 1988, *Int. J. Pept. Protein Res.* 32:242-255. Beta-turn profiles can be generated using the method of Deleage, G., Roux B., 1987, *Protein Engineering* 1:289-294. Thus, each region identified by any of these programs or methods is within the scope of the present invention. Methods for the generation of 24P4C12 antibodies are further illustrated by way of the examples provided herein. Methods for preparing a protein or polypeptide for use as an immunogen are well known in the art. Also well known in the art are methods for preparing immunogenic conjugates of a protein with a carrier, such as BSA, KLH or other carrier protein. In some circumstances, direct conjugation using, for example, carbodiimide reagents are used; in other instances linking reagents such as those supplied by Pierce Chemical Co., Rockford, Ill., are effective. Administration of a 24P4C12 immunogen is often conducted by injection over a suitable time period and with use of a suitable adjuvant, as is understood in the art. During the immunization schedule, titers of antibodies can be taken to determine adequacy of antibody formation.

24P4C12 monoclonal antibodies can be produced by various means well known in the art. For example, immortalized cell lines that secrete a desired monoclonal antibody are prepared using the standard hybridoma technology of Kohler and Milstein or modifications that immortalize antibody-producing B cells, as is generally known. Immortalized cell lines that secrete the desired antibodies are screened by immunoassay in which the antigen is a 24P4C12-related protein. When the appropriate immortalized cell culture is identified, the cells can be expanded and antibodies produced either from in vitro cultures or from ascites fluid.

The antibodies or fragments of the invention can also be produced, by recombinant means. Regions that bind specifically to the desired regions of a 24P4C12 protein can also be produced in the context of chimeric or complementarity-determining region (CDR) grafted antibodies of multiple species origin. Humanized or human 24P4C12 antibodies can also be produced, and are preferred for use in therapeutic contexts. Methods for humanizing murine and other non-human antibodies, by substituting one or more of the non-human antibody CDRs for corresponding human antibody

sequences, are well known (see for example, Jones et al., 1986, *Nature* 321: 522-525; Riechmann et al., 1988, *Nature* 332: 323-327; Verhoeven et al., 1988, *Science* 239: 1534-1536). See also, Carter et al., 1993, *Proc. Natl. Acad. Sci. USA* 89: 4285 and Sims et al., 1993, *J. Immunol.* 151: 2296.

Methods for producing fully human monoclonal antibodies include phage display and transgenic methods (for review, see Vaughan et al., 1998, *Nature Biotechnology* 16: 535-539). Fully human 24P4C12 monoclonal antibodies can be generated using cloning technologies employing large human Ig gene combinatorial libraries (i.e., phage display) (Griffiths and Hoogenboom, *Building an in vitro immune system: human antibodies from phage display libraries*. In: *Protein Engineering of Antibody Molecules for Prophylactic and Therapeutic Applications in Man*, Clark, M. (Ed.), Nottingham Academic, pp 45-64 (1993); Burton and Barbas, *Human Antibodies from combinatorial libraries*. Id., pp 65-82). Fully human 24P4C12 monoclonal antibodies can also be produced using transgenic mice engineered to contain human immunoglobulin gene loci as described in PCT Patent Application WO98/24893, Kucherlapati and Jakobovits et al., published Dec. 3, 1997 (see also, Jakobovits, 1998, *Exp. Opin. Invest. Drugs* 7(4): 607-614; U.S. Pat. Nos. 6,162,963 issued 19 Dec. 2000; 6,150,584 issued 12 Nov. 2000; and, 6,114,598 issued 5 Sep. 2000). This method avoids the in vitro manipulation required with phage display technology and efficiently produces high affinity authentic human antibodies.

Reactivity of 24P4C12 antibodies with a 24P4C12-related protein can be established by a number of well known means, including Western blot, immunoprecipitation, ELISA, and FACS analyses using, as appropriate, 24P4C12-related proteins, 24P4C12-expressing cells or extracts thereof. A 24P4C12 antibody or fragment thereof can be labeled with a detectable marker or conjugated to a second molecule. Suitable detectable markers include, but are not limited to, a radioisotope, a fluorescent compound, a bioluminescent compound, chemiluminescent compound, a metal chelator or an enzyme. Further, bi-specific antibodies specific for two or more 24P4C12 epitopes are generated using methods generally known in the art. Homodimeric antibodies can also be generated by cross-linking techniques known in the art (e.g., Wolff et al., *Cancer Res.* 53: 2560-2565).

V.) 24P4C12 CELLULAR IMMUNE RESPONSES

The mechanism by which T cells recognize antigens has been delineated. Efficacious peptide epitope vaccine compositions of the invention induce a therapeutic or prophylactic immune responses in very broad segments of the world-wide population. For an understanding of the value and efficacy of compositions of the invention that induce cellular immune responses, a brief review of immunology-related technology is provided.

A complex of an HLA molecule and a peptidic antigen acts as the ligand recognized by HLA-restricted T cells (Buus, S. et al., *Cell* 147:1071, 1986; Babbitt, B. P. et al., *Nature* 317: 359, 1985; Townsend, A. and Bodmer, H., *Annu. Rev. Immunol.* 7:601, 1989; Germain, R. N., *Annu. Rev. Immunol.* 11:403, 1993). Through the study of single amino acid substituted antigen analogs and the sequencing of endogenously bound, naturally processed peptides, critical residues that correspond to motifs required for specific binding to HLA antigen molecules have been identified and are set forth in Table IV (see also, e.g., Southwood, et al., *J. Immunol.* 160: 3363, 1998; Rammensee, et al., *Immunogenetics* 41:178, 1995; Rammensee et al., SYFPEITHI, access via World Wide Web at URL (134.2.96.221/scripts.hlaserver.dll/home.htm);

Sette, A. and Sidney, J. *Curr. Opin. Immunol.* 10:478, 1998; Engelhard, V. H., *Curr. Opin. Immunol.* 6:13, 1994; Sette, A. and Grey, H. M., *Curr. Opin. Immunol.* 4:79, 1992; Sinigaglia, F. and Hammer, J. *Curr. Biol.* 6:52, 1994; Ruppert et al., *Cell* 74:929-937, 1993; Kondo et al., *J. Immunol.* 155: 4307-4312, 1995; Sidney et al., *J. Immunol.* 157:3480-3490, 1996; Sidney et al., *Human Immunol.* 45:79-93, 1996; Sette, A. and Sidney, J. *Immunogenetics* 1999 November; 50(3-4): 201-12, (Review).

Furthermore, x-ray crystallographic analyses of HLA-peptide complexes have revealed pockets within the peptide binding cleft/groove of HLA molecules which accommodate, in an allele-specific mode, residues borne by peptide ligands; these residues in turn determine the HLA binding capacity of the peptides in which they are present. (See, e.g., Madden, D. R. *Annu. Rev. Immunol.* 13:587, 1995; Smith, et al., *Immunity* 4:203, 1996; Fremont et al., *Immunity* 8:305, 1998; Stern et al., *Structure* 2:245, 1994; Jones, E. Y. *Curr. Opin. Immunol.* 9:75, 1997; Brown, J. H. et al., *Nature* 364:33, 1993; Guo, H. C. et al., *Proc. Natl. Acad. Sci. USA* 90:8053, 1993; Guo, H. C. et al., *Nature* 360:364, 1992; Silver, M. L. et al., *Nature* 360:367, 1992; Matsumura, M. et al., *Science* 257:927, 1992; Madden et al., *Cell* 70:1035, 1992; Fremont, D. H. et al., *Science* 257:919, 1992; Saper, M. A., Bjorkman, P. J. and Wiley, D.C., *J. Mol. Biol.* 219:277, 1991.)

Accordingly, the definition of class I and class II allele-specific HLA binding motifs, or class I or class II supermotifs allows identification of regions within a protein that are correlated with binding to particular HLA antigen(s).

Thus, by a process of HLA motif identification, candidates for epitope-based vaccines have been identified; such candidates can be further evaluated by HLA-peptide binding assays to determine binding affinity and/or the time period of association of the epitope and its corresponding HLA molecule. Additional confirmatory work can be performed to select, amongst these vaccine candidates, epitopes with preferred characteristics in terms of population coverage, and/or immunogenicity.

Various strategies can be utilized to evaluate cellular immunogenicity, including:

1) Evaluation of primary T cell cultures from normal individuals (see, e.g., Wentworth, P. A. et al., *Mol. Immunol.* 32:603, 1995; Celis, E. et al., *Proc. Natl. Acad. Sci. USA* 91:2105, 1994; Tsai, V. et al., *J. Immunol.* 158:1796, 1997; Kawashima, I. et al., *Human Immunol.* 59:1, 1998). This procedure involves the stimulation of peripheral blood lymphocytes (PBL) from normal subjects with a test peptide in the presence of antigen presenting cells in vitro over a period of several weeks. T cells specific for the peptide become activated during this time and are detected using, e.g., a lymphokine- or ⁵¹Cr-release assay involving peptide sensitized target cells.

2) Immunization of HLA transgenic mice (see, e.g., Wentworth, P. A. et al., *J. Immunol.* 26:97, 1996; Wentworth, P. A. et al., *Int. Immunol.* 8:651, 1996; Alexander, J. et al., *J. Immunol.* 159:4753, 1997). For example, in such methods peptides in incomplete Freund's adjuvant are administered subcutaneously to HLA transgenic mice. Several weeks following immunization, splenocytes are removed and cultured in vitro in the presence of test peptide for approximately one week. Peptide-specific T cells are detected using, e.g., a ⁵¹Cr-release assay involving peptide sensitized target cells and target cells expressing endogenously generated antigen.

3) Demonstration of recall T cell responses from immune individuals who have been either effectively vaccinated and/or from chronically ill patients (see, e.g., Rehmann, B. et al., *J. Exp. Med.* 181:1047, 1995; Doolan, D. L. et al., *Immunity*

7:97, 1997; Bertoni, R. et al., *J. Clin. Invest* 100:503, 1997; Threlkeld, S. C. et al., *J. Immunol.* 159:1648, 1997; Diepolder, H. M. et al., *J. Virol.* 71:6011, 1997). Accordingly, recall responses are detected by culturing PBL from subjects that have been exposed to the antigen due to disease and thus have generated an immune response "naturally", or from patients who were vaccinated against the antigen. PBL from subjects are cultured in vitro for 1-2 weeks in the presence of test peptide plus antigen presenting cells (APC) to allow activation of "memory" T cells, as compared to "naive" T cells. At the end of the culture period, T cell activity is detected using assays including ⁵¹Cr release involving peptide-sensitized targets, T cell proliferation, or lymphokine release.

VI.) 24P4C12 TRANSGENIC ANIMALS

Nucleic acids that encode a 24P4C12-related protein can also be used to generate either transgenic animals or "knock out" animals that, in turn, are useful in the development and screening of therapeutically useful reagents. In accordance with established techniques, cDNA encoding 24P4C12 can be used to clone genomic DNA that encodes 24P4C12. The cloned genomic sequences can then be used to generate transgenic animals containing cells that express DNA that encode 24P4C12. Methods for generating transgenic animals, particularly animals such as mice or rats, have become conventional in the art and are described, for example, in U.S. Pat. Nos. 4,736,866 issued 12 Apr. 1988, and 4,870,009 issued 26 Sep. 1989. Typically, particular cells would be targeted for 24P4C12 transgene incorporation with tissue-specific enhancers.

Transgenic animals that include a copy of a transgene encoding 24P4C12 can be used to examine the effect of increased expression of DNA that encodes 24P4C12. Such animals can be used as tester animals for reagents thought to confer protection from, for example, pathological conditions associated with its overexpression. In accordance with this aspect of the invention, an animal is treated with a reagent and a reduced incidence of a pathological condition, compared to untreated animals that bear the transgene, would indicate a potential therapeutic intervention for the pathological condition.

Alternatively, non-human homologues of 24P4C12 can be used to construct a 24P4C12 "knock out" animal that has a defective or altered gene encoding 24P4C12 as a result of homologous recombination between the endogenous gene encoding 24P4C12 and altered genomic DNA encoding 24P4C12 introduced into an embryonic cell of the animal. For example, cDNA that encodes 24P4C12 can be used to clone genomic DNA encoding 24P4C12 in accordance with established techniques. A portion of the genomic DNA encoding 24P4C12 can be deleted or replaced with another gene, such as a gene encoding a selectable marker that can be used to monitor integration. Typically, several kilobases of unaltered flanking DNA (both at the 5' and 3' ends) are included in the vector (see, e.g., Thomas and Capecchi, *Cell*, 51:503 (1987) for a description of homologous recombination vectors). The vector is introduced into an embryonic stem cell line (e.g., by electroporation) and cells in which the introduced DNA has homologously recombined with the endogenous DNA are selected (see, e.g., Li et al., *Cell*, 69:915 (1992)). The selected cells are then injected into a blastocyst of an animal (e.g., a mouse or rat) to form aggregation chimeras (see, e.g., Bradley, in *Teratocarcinomas and Embryonic Stem Cells: A Practical Approach*, E. J. Robertson, ed. (IRL, Oxford, 1987), pp. 113-152). A chimeric embryo can then be implanted into a

suitable pseudopregnant female foster animal, and the embryo brought to term to create a "knock out" animal. Progeny harboring the homologously recombined DNA in their germ cells can be identified by standard techniques and used to breed animals in which all cells of the animal contain the homologously recombined DNA. Knock out animals can be characterized, for example, for their ability to defend against certain pathological conditions or for their development of pathological conditions due to absence of a 24P4C12 polypeptide.

VII.) METHODS FOR THE DETECTION OF 24P4C12

Another aspect of the present invention relates to methods for detecting 24P4C12 polynucleotides and 24P4C12-related proteins, as well as methods for identifying a cell that expresses 24P4C12. The expression profile of 24P4C12 makes it a diagnostic marker for metastasized disease. Accordingly, the status of 24P4C12 gene products provides information useful for predicting a variety of factors including susceptibility to advanced stage disease, rate of progression, and/or tumor aggressiveness. As discussed in detail herein, the status of 24P4C12 gene products in patient samples can be analyzed by a variety of protocols that are well known in the art including immunohistochemical analysis, the variety of Northern blotting techniques including in situ hybridization, RT-PCR analysis (for example on laser capture micro-dissected samples), Western blot analysis and tissue array analysis.

More particularly, the invention provides assays for the detection of 24P4C12 polynucleotides in a biological sample, such as serum, bone, prostate, and other tissues, urine, semen, cell preparations, and the like. Detectable 24P4C12 polynucleotides include, for example, a 24P4C12 gene or fragment thereof, 24P4C12 mRNA, alternative splice variant 24P4C12 mRNAs, and recombinant DNA or RNA molecules that contain a 24P4C12 polynucleotide. A number of methods for amplifying and/or detecting the presence of 24P4C12 polynucleotides are well known in the art and can be employed in the practice of this aspect of the invention.

In one embodiment, a method for detecting a 24P4C12 mRNA in a biological sample comprises producing cDNA from the sample by reverse transcription using at least one primer; amplifying the cDNA so produced using a 24P4C12 polynucleotides as sense and antisense primers to amplify 24P4C12 cDNAs therein; and detecting the presence of the amplified 24P4C12 cDNA. Optionally, the sequence of the amplified 24P4C12 cDNA can be determined.

In another embodiment, a method of detecting a 24P4C12 gene in a biological sample comprises first isolating genomic DNA from the sample; amplifying the isolated genomic DNA using 24P4C12 polynucleotides as sense and antisense primers; and detecting the presence of the amplified 24P4C12 gene. Any number of appropriate sense and antisense probe combinations can be designed from a 24P4C12 nucleotide sequence (see, e.g., FIG. 2) and used for this purpose.

The invention also provides assays for detecting the presence of a 24P4C12 protein in a tissue or other biological sample such as serum, semen, bone, prostate, urine, cell preparations, and the like. Methods for detecting a 24P4C12-related protein are also well known and include, for example, immunoprecipitation, immunohistochemical analysis, Western blot analysis, molecular binding assays, ELISA, ELIFA and the like. For example, a method of detecting the presence of a 24P4C12-related protein in a biological sample comprises first contacting the sample with a 24P4C12 antibody, a

24P4C12-reactive fragment thereof, or a recombinant protein containing an antigen-binding region of a 24P4C12 antibody; and then detecting the binding of 24P4C12-related protein in the sample.

Methods for identifying a cell that expresses 24P4C12 are also within the scope of the invention. In one embodiment, an assay for identifying a cell that expresses a 24P4C12 gene comprises detecting the presence of 24P4C12 mRNA in the cell. Methods for the detection of particular mRNAs in cells are well known and include, for example, hybridization assays using complementary DNA probes (such as in situ hybridization using labeled 24P4C12 riboprobes, Northern blot and related techniques) and various nucleic acid amplification assays (such as RT-PCR using complementary primers specific for 24P4C12, and other amplification type detection methods, such as, for example, branched DNA, SISBA, TMA and the like). Alternatively, an assay for identifying a cell that expresses a 24P4C12 gene comprises detecting the presence of 24P4C12-related protein in the cell or secreted by the cell. Various methods for the detection of proteins are well known in the art and are employed for the detection of 24P4C12-related proteins and cells that express 24P4C12-related proteins.

24P4C12 expression analysis is also useful as a tool for identifying and evaluating agents that modulate 24P4C12 gene expression. For example, 24P4C12 expression is significantly upregulated in prostate cancer, and is expressed in cancers of the tissues listed in Table I. Identification of a molecule or biological agent that inhibits 24P4C12 expression or over-expression in cancer cells is of therapeutic value. For example, such an agent can be identified by using a screen that quantifies 24P4C12 expression by RT-PCR, nucleic acid hybridization or antibody binding.

VIII.) METHODS FOR MONITORING THE STATUS OF 24P4C12-RELATED GENES AND THEIR PRODUCTS

Oncogenesis is known to be a multistep process where cellular growth becomes progressively dysregulated and cells progress from a normal physiological state to precancerous and then cancerous states (see, e.g., Alers et al., *Lab Invest.* 77(5): 437-438 (1997) and Isaacs et al., *Cancer Surv.* 23: 19-32 (1995)). In this context, examining a biological sample for evidence of dysregulated cell growth (such as aberrant 24P4C12 expression in cancers) allows for early detection of such aberrant physiology, before a pathologic state such as cancer has progressed to a stage that therapeutic options are more limited and/or the prognosis is worse. In such examinations, the status of 24P4C12 in a biological sample of interest can be compared, for example, to the status of 24P4C12 in a corresponding normal sample (e.g. a sample from that individual or alternatively another individual that is not affected by a pathology). An alteration in the status of 24P4C12 in the biological sample (as compared to the normal sample) provides evidence of dysregulated cellular growth. In addition to using a biological sample that is not affected by a pathology as a normal sample, one can also use a predetermined normative value such as a predetermined normal level of mRNA expression (see, e.g., Grever et al., *J. Comp. Neurol.* 1996 Dec. 9; 376(2): 306-14 and U.S. Pat. No. 5,837,501) to compare 24P4C12 status in a sample.

The term "status" in this context is used according to its art accepted meaning and refers to the condition or state of a gene and its products. Typically, skilled artisans use a number of parameters to evaluate the condition or state of a gene and its products. These include, but are not limited to the location of

expressed gene products (including the location of 24P4C12 expressing cells) as well as the level, and biological activity of expressed gene products (such as 24P4C12 mRNA, polynucleotides and polypeptides). Typically, an alteration in the status of 24P4C12 comprises a change in the location of 24P4C12 and/or 24P4C12 expressing cells and/or an increase in 24P4C12 mRNA and/or protein expression.

24P4C12 status in a sample can be analyzed by a number of means well known in the art, including without limitation, immunohistochemical analysis, in situ hybridization, RT-PCR analysis on laser capture micro-dissected samples, Western blot analysis, and tissue array analysis. Typical protocols for evaluating the status of a 24P4C12 gene and gene products are found, for example in Ausubel et al. eds., 1995, *Current Protocols In Molecular Biology*, Units 2 (Northern Blotting), 4 (Southern Blotting), 15 (Immunoblotting) and 18 (PCR Analysis). Thus, the status of 24P4C12 in a biological sample is evaluated by various methods utilized by skilled artisans including, but not limited to genomic Southern analysis (to examine, for example perturbations in a 24P4C12 gene), Northern analysis and/or PCR analysis of 24P4C12 mRNA (to examine, for example alterations in the polynucleotide sequences or expression levels of 24P4C12 mRNAs), and, Western and/or immunohistochemical analysis (to examine, for example alterations in polypeptide sequences, alterations in polypeptide localization within a sample, alterations in expression levels of 24P4C12 proteins and/or associations of 24P4C12 proteins with polypeptide binding partners). Detectable 24P4C12 polynucleotides include, for example, a 24P4C12 gene or fragment thereof, 24P4C12 mRNA, alternative splice variants, 24P4C12 mRNAs, and recombinant DNA or RNA molecules containing a 24P4C12 polynucleotide.

The expression profile of 24P4C12 makes it a diagnostic marker for local and/or metastasized disease, and provides information on the growth or oncogenic potential of a biological sample. In particular, the status of 24P4C12 provides information useful for predicting susceptibility to particular disease stages, progression, and/or tumor aggressiveness. The invention provides methods and assays for determining 24P4C12 status and diagnosing cancers that express 24P4C12, such as cancers of the tissues listed in Table I. For example, because 24P4C12 mRNA is so highly expressed in prostate and other cancers relative to normal prostate tissue, assays that evaluate the levels of 24P4C12 mRNA transcripts or proteins in a biological sample can be used to diagnose a disease associated with 24P4C12 dysregulation, and can provide prognostic information useful in defining appropriate therapeutic options.

The expression status of 24P4C12 provides information including the presence, stage and location of dysplastic, precancerous and cancerous cells, predicting susceptibility to various stages of disease, and/or for gauging tumor aggressiveness. Moreover, the expression profile makes it useful as an imaging reagent for metastasized disease. Consequently, an aspect of the invention is directed to the various molecular prognostic and diagnostic methods for examining the status of 24P4C12 in biological samples such as those from individuals suffering from, or suspected of suffering from a pathology characterized by dysregulated cellular growth, such as cancer.

As described above, the status of 24P4C12 in a biological sample can be examined by a number of well-known procedures in the art. For example, the status of 24P4C12 in a biological sample taken from a specific location in the body can be examined by evaluating the sample for the presence or absence of 24P4C12 expressing cells (e.g. those that express

24P4C12 mRNAs or proteins). This examination can provide evidence of dysregulated cellular growth, for example, when 24P4C12-expressing cells are found in a biological sample that does not normally contain such cells (such as a lymph node), because such alterations in the status of 24P4C12 in a biological sample are often associated with dysregulated cellular growth. Specifically, one indicator of dysregulated cellular growth is the metastases of cancer cells from an organ of origin (such as the prostate) to a different area of the body (such as a lymph node). In this context, evidence of dysregulated cellular growth is important for example because occult lymph node metastases can be detected in a substantial proportion of patients with prostate cancer, and such metastases are associated with known predictors of disease progression (see, e.g., Murphy et al., *Prostate* 42(4): 315-317 (2000); Su et al., *Semin. Surg. Oncol.* 18(1): 17-28 (2000) and Freeman et al., *J Urol* 1995 August 154 (2 Pt 1):474-8).

In one aspect, the invention provides methods for monitoring 24P4C12 gene products by determining the status of 24P4C12 gene products expressed by cells from an individual suspected of having a disease associated with dysregulated cell growth (such as hyperplasia or cancer) and then comparing the status so determined to the status of 24P4C12 gene products in a corresponding normal sample. The presence of aberrant 24P4C12 gene products in the test sample relative to the normal sample provides an indication of the presence of dysregulated cell growth within the cells of the individual.

In another aspect, the invention provides assays useful in determining the presence of cancer in an individual, comprising detecting a significant increase in 24P4C12 mRNA or protein expression in a test cell or tissue sample relative to expression levels in the corresponding normal cell or tissue. The presence of 24P4C12 mRNA can, for example, be evaluated in tissues including but not limited to those listed in Table I. The presence of significant 24P4C12 expression in any of these tissues is useful to indicate the emergence, presence and/or severity of a cancer, since the corresponding normal tissues do not express 24P4C12 mRNA or express it at lower levels.

In a related embodiment, 24P4C12 status is determined at the protein level rather than at the nucleic acid level. For example, such a method comprises determining the level of 24P4C12 protein expressed by cells in a test tissue sample and comparing the level so determined to the level of 24P4C12 expressed in a corresponding normal sample. In one embodiment, the presence of 24P4C12 protein is evaluated, for example, using immunohistochemical methods. 24P4C12 antibodies or binding partners capable of detecting 24P4C12 protein expression are used in a variety of assay formats well known in the art for this purpose.

In a further embodiment, one can evaluate the status of 24P4C12 nucleotide and amino acid sequences in a biological sample in order to identify perturbations in the structure of these molecules. These perturbations can include insertions, deletions, substitutions and the like. Such evaluations are useful because perturbations in the nucleotide and amino acid sequences are observed in a large number of proteins associated with a growth dysregulated phenotype (see, e.g., Marrogi et al., 1999, *J. Cutan. Pathol.* 26(8):369-378). For example, a mutation in the sequence of 24P4C12 may be indicative of the presence or promotion of a tumor. Such assays therefore have diagnostic and predictive value where a mutation in 24P4C12 indicates a potential loss of function or increase in tumor growth.

A wide variety of assays for observing perturbations in nucleotide and amino acid sequences are well known in the art. For example, the size and structure of nucleic acid or

amino acid sequences of 24P4C12 gene products are observed by the Northern, Southern, Western, PCR and DNA sequencing protocols discussed herein. In addition, other methods for observing perturbations in nucleotide and amino acid sequences such as single strand conformation polymorphism analysis are well known in the art (see, e.g., U.S. Pat. Nos. 5,382,510 issued 7 Sep. 1999, and 5,952,170 issued 17 Jan. 1995).

Additionally, one can examine the methylation status of a 24P4C12 gene in a biological sample. Aberrant demethylation and/or hypermethylation of CpG islands in gene 5' regulatory regions frequently occurs in immortalized and transformed cells, and can result in altered expression of various genes. For example, promoter hypermethylation of the p1-class glutathione S-transferase (a protein expressed in normal prostate but not expressed in >90% of prostate carcinomas) appears to permanently silence transcription of this gene and is the most frequently detected genomic alteration in prostate carcinomas (De Marzo et al., *Am. J. Pathol.* 155(6): 1985-1992 (1999)). In addition, this alteration is present in at least 70% of cases of high-grade prostatic intraepithelial neoplasia (PIN) (Brooks et al., *Cancer Epidemiol. Biomarkers Prev.*, 1998, 7:531-536). In another example, expression of the LAGE-I tumor specific gene (which is not expressed in normal prostate but is expressed in 25-50% of prostate cancers) is induced by deoxy-azacytidine in lymphoblastoid cells, suggesting that tumoral expression is due to demethylation (Lethe et al., *Int. J. Cancer* 76(6): 903-908 (1998)). A variety of assays for examining methylation status of a gene are well known in the art. For example, one can utilize, in Southern hybridization approaches, methylation-sensitive restriction enzymes that cannot cleave sequences that contain methylated CpG sites to assess the methylation status of CpG islands. In addition, MSP (methylation specific PCR) can rapidly profile the methylation status of all the CpG sites present in a CpG island of a given gene. This procedure involves initial modification of DNA by sodium bisulfite (which will convert all unmethylated cytosines to uracil) followed by amplification using primers specific for methylated versus unmethylated DNA. Protocols involving methylation interference can also be found for example in *Current Protocols In Molecular Biology*, Unit 12, Frederick M. Ausubel et al. eds., 1995.

Gene amplification is an additional method for assessing the status of 24P4C12. Gene amplification is measured in a sample directly, for example, by conventional Southern blotting or Northern blotting to quantitate the transcription of mRNA (Thomas, 1980, *Proc. Natl. Acad. Sci. USA*, 77:5201-5205), dot blotting (DNA analysis), or in situ hybridization, using an appropriately labeled probe, based on the sequences provided herein. Alternatively, antibodies are employed that recognize specific duplexes, including DNA duplexes, RNA duplexes, and DNA-RNA hybrid duplexes or DNA-protein duplexes. The antibodies in turn are labeled and the assay carried out where the duplex is bound to a surface, so that upon the formation of duplex on the surface, the presence of antibody bound to the duplex can be detected.

Biopsied tissue or peripheral blood can be conveniently assayed for the presence of cancer cells using for example, Northern, dot blot or RT-PCR analysis to detect 24P4C12 expression. The presence of RT-PCR amplifiable 24P4C12 mRNA provides an indication of the presence of cancer. RT-PCR assays are well known in the art. RT-PCR detection assays for tumor cells in peripheral blood are currently being evaluated for use in the diagnosis and management of a number of human solid tumors. In the prostate cancer field, these include RT-PCR assays for the detection of cells expressing

PSA and PSM (Verkaik et al., 1997, *Urol. Res.* 25:373-384; Ghossein et al., 1995, *J. Clin. Oncol.* 13:1195-2000; Heston et al., 1995, *Clin. Chem.* 41:1687-1688).

A further aspect of the invention is an assessment of the susceptibility that an individual has for developing cancer. In one embodiment, a method for predicting susceptibility to cancer comprises detecting 24P4C12 mRNA or 24P4C12 protein in a tissue sample, its presence indicating susceptibility to cancer, wherein the degree of 24P4C12 mRNA expression correlates to the degree of susceptibility. In a specific embodiment, the presence of 24P4C12 in prostate or other tissue is examined, with the presence of 24P4C12 in the sample providing an indication of prostate cancer susceptibility (or the emergence or existence of a prostate tumor). Similarly, one can evaluate the integrity 24P4C12 nucleotide and amino acid sequences in a biological sample, in order to identify perturbations in the structure of these molecules such as insertions, deletions, substitutions and the like. The presence of one or more perturbations in 24P4C12 gene products in the sample is an indication of cancer susceptibility (or the emergence or existence of a tumor).

The invention also comprises methods for gauging tumor aggressiveness. In one embodiment, a method for gauging aggressiveness of a tumor comprises determining the level of 24P4C12 mRNA or 24P4C12 protein expressed by tumor cells, comparing the level so determined to the level of 24P4C12 mRNA or 24P4C12 protein expressed in a corresponding normal tissue taken from the same individual or a normal tissue reference sample, wherein the degree of 24P4C12 mRNA or 24P4C12 protein expression in the tumor sample relative to the normal sample indicates the degree of aggressiveness. In a specific embodiment, aggressiveness of a tumor is evaluated by determining the extent to which 24P4C12 is expressed in the tumor cells, with higher expression levels indicating more aggressive tumors. Another embodiment is the evaluation of the integrity of 24P4C12 nucleotide and amino acid sequences in a biological sample, in order to identify perturbations in the structure of these molecules such as insertions, deletions, substitutions and the like. The presence of one or more perturbations indicates more aggressive tumors.

Another embodiment of the invention is directed to methods for observing the progression of a malignancy in an individual over time. In one embodiment, methods for observing the progression of a malignancy in an individual over time comprise determining the level of 24P4C12 mRNA or 24P4C12 protein expressed by cells in a sample of the tumor, comparing the level so determined to the level of 24P4C12 mRNA or 24P4C12 protein expressed in an equivalent tissue sample taken from the same individual at a different time, wherein the degree of 24P4C12 mRNA or 24P4C12 protein expression in the tumor sample over time provides information on the progression of the cancer. In a specific embodiment, the progression of a cancer is evaluated by determining 24P4C12 expression in the tumor cells over time, where increased expression over time indicates a progression of the cancer. Also, one can evaluate the integrity 24P4C12 nucleotide and amino acid sequences in a biological sample in order to identify perturbations in the structure of these molecules such as insertions, deletions, substitutions and the like, where the presence of one or more perturbations indicates a progression of the cancer.

The above diagnostic approaches can be combined with any one of a wide variety of prognostic and diagnostic protocols known in the art. For example, another embodiment of the invention is directed to methods for observing a coincidence between the expression of 24P4C12 gene and 24P4C12

gene products (or perturbations in 24P4C12 gene and 24P4C12 gene products) and a factor that is associated with malignancy, as a means for diagnosing and prognosticating the status of a tissue sample. A wide variety of factors associated with malignancy can be utilized, such as the expression of genes associated with malignancy (e.g. PSA, PSCA and PSM expression for prostate cancer etc.) as well as gross cytological observations (see, e.g., Bocking et al., 1984, *Anal. Quant. Cytol.* 6(2):74-88; Epstein, 1995, *Hum. Pathol.* 26(2): 223-9; Thorson et al., 1998, *Mod. Pathol.* 11(6):543-51; Baisden et al., 1999, *Am. J. Surg. Pathol.* 23(8):918-24). Methods for observing a coincidence between the expression of 24P4C12 gene and 24P4C12 gene products (or perturbations in 24P4C12 gene and 24P4C12 gene products) and another factor that is associated with malignancy are useful, for example, because the presence of a set of specific factors that coincide with disease provides information crucial for diagnosing and prognosticating the status of a tissue sample.

In one embodiment, methods for observing a coincidence between the expression of 24P4C12 gene and 24P4C12 gene products (or perturbations in 24P4C12 gene and 24P4C12 gene products) and another factor associated with malignancy entails detecting the overexpression of 24P4C12 mRNA or protein in a tissue sample, detecting the overexpression of PSA mRNA or protein in a tissue sample (or PSCA or PSM expression), and observing a coincidence of 24P4C12 mRNA or protein and PSA mRNA or protein overexpression (or PSCA or PSM expression). In a specific embodiment, the expression of 24P4C12 and PSA mRNA in prostate tissue is examined, where the coincidence of 24P4C12 and PSA mRNA overexpression in the sample indicates the existence of prostate cancer, prostate cancer susceptibility or the emergence or status of a prostate tumor.

Methods for detecting and quantifying the expression of 24P4C12 mRNA or protein are described herein, and standard nucleic acid and protein detection and quantification technologies are well known in the art. Standard methods for the detection and quantification of 24P4C12 mRNA include in situ hybridization using labeled 24P4C12 riboprobes, Northern blot and related techniques using 24P4C12 polynucleotide probes, RT-PCR analysis using primers specific for 24P4C12, and other amplification type detection methods, such as, for example, branched DNA, SISBA, TMA and the like. In a specific embodiment, semi-quantitative RT-PCR is used to detect and quantify 24P4C12 mRNA expression. Any number of primers capable of amplifying 24P4C12 can be used for this purpose, including but not limited to the various primer sets specifically described herein. In a specific embodiment, polyclonal or monoclonal antibodies specifically reactive with the wild-type 24P4C12 protein can be used in an immunohistochemical assay of biopsied tissue.

IX.) IDENTIFICATION OF MOLECULES THAT INTERACT WITH 24P4C12

The 24P4C12 protein and nucleic acid sequences disclosed herein allow a skilled artisan to identify proteins, small molecules and other agents that interact with 24P4C12, as well as pathways activated by 24P4C12 via any one of a variety of art accepted protocols. For example, one can utilize one of the so-called interaction trap systems (also referred to as the "two-hybrid assay"). In such systems, molecules interact and reconstitute a transcription factor which directs expression of a reporter gene, whereupon the expression of the reporter gene is assayed. Other systems identify protein-protein interactions in vivo through reconstitution of a eukaryotic transcriptional activator, see, e.g., U.S. Pat. Nos. 5,955,280 issued

21 Sep. 1999, 5,925,523 issued 20 Jul. 1999, 5,846,722 issued 8 Dec. 1998 and 6,004,746 issued 21 Dec. 1999. Algorithms are also available in the art for genome-based predictions of protein function (see, e.g., Marcotte, et al., *Nature* 402: 4 Nov. 1999, 83-86).

Alternatively one can screen peptide libraries to identify molecules that interact with 24P4C12 protein sequences. In such methods, peptides that bind to 24P4C12 are identified by screening libraries that encode a random or controlled collection of amino acids. Peptides encoded by the libraries are expressed as fusion proteins of bacteriophage coat proteins, the bacteriophage particles are then screened against the 24P4C12 protein(s).

Accordingly, peptides having a wide variety of uses, such as therapeutic, prognostic or diagnostic reagents, are thus identified without any prior information on the structure of the expected ligand or receptor molecule. Typical peptide libraries and screening methods that can be used to identify molecules that interact with 24P4C12 protein sequences are disclosed for example in U.S. Pat. Nos. 5,723,286 issued 3 Mar. 1998 and 5,733,731 issued 31 Mar. 1998.

Alternatively, cell lines that express 24P4C12 are used to identify protein-protein interactions mediated by 24P4C12. Such interactions can be examined using immunoprecipitation techniques (see, e.g., Hamilton B. J., et al. *Biochem. Biophys. Res. Commun.* 1999, 261:646-51). 24P4C12 protein can be immunoprecipitated from 24P4C12-expressing cell lines using anti-24P4C12 antibodies. Alternatively, antibodies against His-tag can be used in a cell line engineered to express fusions of 24P4C12 and a His-tag (vectors mentioned above). The immunoprecipitated complex can be examined for protein association by procedures such as Western blotting, ³⁵S-methionine labeling of proteins, protein microsequencing, silver staining and two-dimensional gel electrophoresis.

Small molecules and ligands that interact with 24P4C12 can be identified through related embodiments of such screening assays. For example, small molecules can be identified that interfere with protein function, including molecules that interfere with 24P4C12's ability to mediate phosphorylation and de-phosphorylation, interaction with DNA or RNA molecules as an indication of regulation of cell cycles, second messenger signaling or tumorigenesis. Similarly, small molecules that modulate 24P4C12-related ion channel, protein pump, or cell communication functions are identified and used to treat patients that have a cancer that expresses 24P4C12 (see, e.g., Hille, B., *Ionic Channels of Excitable Membranes* 2nd Ed., Sinauer Assoc., Sunderland, Mass., 1992). Moreover, ligands that regulate 24P4C12 function can be identified based on their ability to bind 24P4C12 and activate a reporter construct. Typical methods are discussed for example in U.S. Pat. No. 5,928,868 issued 27 Jul. 1999, and include methods for forming hybrid ligands in which at least one ligand is a small molecule. In an illustrative embodiment, cells engineered to express a fusion protein of 24P4C12 and a DNA-binding protein are used to co-express a fusion protein of a hybrid ligand/small molecule and a cDNA library transcriptional activator protein. The cells further contain a reporter gene, the expression of which is conditioned on the proximity of the first and second fusion proteins to each other, an event that occurs only if the hybrid ligand binds to target sites on both hybrid proteins. Those cells that express the reporter gene are selected and the unknown small molecule or the unknown ligand is identified. This method provides a means of identifying modulators, which activate or inhibit 24P4C12.

An embodiment of this invention comprises a method of screening for a molecule that interacts with a 24P4C12 amino acid sequence shown in FIG. 2 or FIG. 3, comprising the steps of contacting a population of molecules with a 24P4C12 amino acid sequence, allowing the population of molecules and the 24P4C12 amino acid sequence to interact under conditions that facilitate an interaction, determining the presence of a molecule that interacts with the 24P4C12 amino acid sequence, and then separating molecules that do not interact with the 24P4C12 amino acid sequence from molecules that do. In a specific embodiment, the method further comprises purifying, characterizing and identifying a molecule that interacts with the 24P4C12 amino acid sequence. The identified molecule can be used to modulate a function performed by 24P4C12. In a preferred embodiment, the 24P4C12 amino acid sequence is contacted with a library of peptides.

X.) THERAPEUTIC METHODS AND COMPOSITIONS

The identification of 24P4C12 as a protein that is normally expressed in a restricted set of tissues, but which is also expressed in prostate and other cancers, opens a number of therapeutic approaches to the treatment of such cancers. As contemplated herein, 24P4C12 functions as a transcription factor involved in activating tumor-promoting genes or repressing genes that block tumorigenesis.

Accordingly, therapeutic approaches that inhibit the activity of a 24P4C12 protein are useful for patients suffering from a cancer that expresses 24P4C12. These therapeutic approaches generally fall into two classes. One class comprises various methods for inhibiting the binding or association of a 24P4C12 protein with its binding partner or with other proteins. Another class comprises a variety of methods for inhibiting the transcription of a 24P4C12 gene or translation of 24P4C12 mRNA.

X.A.) Anti-Cancer Vaccines

The invention provides cancer vaccines comprising a 24P4C12-related protein or 24P4C12-related nucleic acid. In view of the expression of 24P4C12, cancer vaccines prevent and/or treat 24P4C12-expressing cancers with minimal or no effects on non-target tissues. The use of a tumor antigen in a vaccine that generates humoral and/or cell-mediated immune responses as anti-cancer therapy is well known in the art and has been employed in prostate cancer using human PSMA and rodent PAP immunogens (Hodge et al., 1995, *Int. J. Cancer* 63:231-237; Fong et al., 1997, *J. Immunol.* 159:3113-3117).

Such methods can be readily practiced by employing a 24P4C12-related protein, or a 24P4C12-encoding nucleic acid molecule and recombinant vectors capable of expressing and presenting the 24P4C12 immunogen (which typically comprises a number of antibody or T cell epitopes). Skilled artisans understand that a wide variety of vaccine systems for delivery of immunoreactive epitopes are known in the art (see, e.g., Heryln et al., *Ann Med* 1999 February 31(1):66-78; Maruyama et al., *Cancer Immunol Immunother* 2000 June 49(3):123-32). Briefly, such methods of generating an immune response (e.g. humoral and/or cell-mediated) in a mammal, comprise the steps of: exposing the mammal's immune system to an immunoreactive epitope (e.g. an epitope present in a 24P4C12 protein shown in FIG. 3 or analog or homolog thereof) so that the mammal generates an immune response that is specific for that epitope (e.g. generates antibodies that specifically recognize that epitope). In a preferred method, a 24P4C12 immunogen contains a biological motif, see e.g., Tables VIII-XXI and XXII-XLIX, or a

peptide of a size range from 24P4C12 indicated in FIG. 5, FIG. 6, FIG. 7, FIG. 8, and FIG. 9.

The entire 24P4C12 protein, immunogenic regions or epitopes thereof can be combined and delivered by various means. Such vaccine compositions can include, for example, lipopeptides (e.g., Vitiello, A. et al., *J. Clin. Invest.* 95:341, 1995), peptide compositions encapsulated in poly(DL-lactide-co-glycolide) ("PLG") microspheres (see, e.g., Eldridge, et al., *Molec. Immunol.* 28:287-294, 1991; Alonso et al., *Vaccine* 12:299-306, 1994; Jones et al., *Vaccine* 13:675-681, 1995), peptide compositions contained in immune stimulating complexes (ISCOMS) (see, e.g., Takahashi et al., *Nature* 344:873-875, 1990; Hu et al., *Clin Exp Immunol.* 113:235-243, 1998), multiple antigen peptide systems (MAPs) (see e.g., Tam, J. P., *Proc. Natl. Acad. Sci. U.S.A.* 85:5409-5413, 1988; Tam, J.P., *J. Immunol. Methods* 196:17-32, 1996), peptides formulated as multivalent peptides; peptides for use in ballistic delivery systems, typically crystallized peptides, viral delivery vectors (Perkus, M. E. et al., In: *Concepts in vaccine development*, Kaufmann, S. H. E., ed., p. 379, 1996; Chakrabarti, S. et al., *Nature* 320:535, 1986; Hu, S. L. et al., *Nature* 320:537, 1986; Kieny, M.-P. et al., *AIDS Bio/Technology* 4:790, 1986; Top, F. H. et al., *J. Infect. Dis.* 124:148, 1971; Chanda, P. K. et al., *Virology* 175:535, 1990), particles of viral or synthetic origin (e.g., Kofler, N. et al., *J. Immunol. Methods* 192:25, 1996; Eldridge, J. H. et al., *Sem. Hematol.* 30:16, 1993; Faló, L. D., Jr. et al., *Nature Med.* 7:649, 1995), adjuvants (Warren, H. S., Vogel, F. R., and Chedid, L. A. *Annu. Rev. Immunol.* 4:369, 1986; Gupta, R. K. et al., *Vaccine* 11:293, 1993), liposomes (Reddy, R. et al., *J. Immunol.* 148:1585, 1992; Rack, K. L., *Immunol. Today* 17:131, 1996), or, naked or particle absorbed cDNA (Ulmer, J. B. et al., *Science* 259:1745, 1993; Robinson, H. L., Hunt, L. A., and Webster, R. G., *Vaccine* 11:957, 1993; Shiver, J. W. et al., In: *Concepts in vaccine development*, Kaufmann, S. H. E., ed., p. 423, 1996; Cease, K. B., and Berzofsky, J. A., *Annu. Rev. Immunol.* 12:923, 1994 and Eldridge, J. H. et al., *Sem. Hematol.* 30:16, 1993). Toxin-targeted delivery technologies, also known as receptor mediated targeting, such as those of Avant Immunotherapeutics, Inc. (Needham, Mass.) may also be used.

In patients with 24P4C12-associated cancer, the vaccine compositions of the invention can also be used in conjunction with other treatments used for cancer, e.g., surgery, chemotherapy, drug therapies, radiation therapies, etc. including use in combination with immune adjuvants such as IL-2, IL-12, GM-CSF, and the like.

Cellular Vaccines:

CTL epitopes can be determined using specific algorithms to identify peptides within 24P4C12 protein that bind corresponding HLA alleles (see e.g., Table IV; Epimer™ and Epimatrix™, Brown University (URL brown.edu/Research/TB-HIV_Lab/epimatrix/epimatrix.html); and, BIMAS, (URL bimas.dcrf.nih.gov/; SYFPEITHI at URL syfpeithi.bmi-heidelberg.com/). In a preferred embodiment, a 24P4C12 immunogen contains one or more amino acid sequences identified using techniques well known in the art, such as the sequences shown in Tables VIII-XXI and XXII-XLIX or a peptide of 8, 9, 10 or 11 amino acids specified by an HLA Class I motif/supermotif (e.g., Table IV (A), Table IV (D), or Table IV (E)) and/or a peptide of at least 9 amino acids that comprises an HLA Class II motif/supermotif (e.g., Table IV (B) or Table IV (C)). As is appreciated in the art, the HLA Class I binding groove is essentially closed ended so that peptides of only a particular size range can fit into the groove and be bound, generally HLA Class I epitopes are 8, 9, 10, or 11 amino acids long. In contrast, the HLA Class II binding groove is essentially open ended; therefore a peptide of about

Figure 2C-2

721 CTCTGGTCTTGAGCCTACTGTTTATCTTGCTTCTGCGCCTGGTGGCTGGGCCCCCTGGTGC
260 V L I L G V L G V L A Y G I Y Y C W E E
781 TGGTGTGATCCIGGGAGTGTCTGGGCGTGTGGCATAACGGCATCTACTACTGCTGGGAGG
280 Y R V L R D K G A S I S Q L G F T T N L
841 AGTACCGAGTGTCTGCGGGACAAGGGCGCCTCCATCTCCCAGCTGGGTTTCACCACCAACC
300 S A Y Q S V Q E T W L A A L I V L A V L
901 TCAGTGCCCTACCAGAGCGTGCAGGAGACCTGGCTGGCCGCCCTGATCGTGTGGCGGTGC
320 E A I L L L M L I F L R Q R I R I A I A
961 TTGAAGCCATCCTGCTGCTGATGCTCATCTTCCTGCGGCAGCGGATTTCGTATTGCCATCG
340 L L K E A S K A V G Q M M S T M F Y P L
1021 CCCTCCIGAAGGAGGCCAGCAAGGCTGTGGGACAGATGATGTCTACCATGTTCTACCCAC
360 V T F V L L L I C I A Y W A M T A L Y L
1081 TGGTCACCTTTGTCTCTCTCATCTGCATTGCCTACTGGGCCATGACTGCTCTGTACC
380 A T S G Q F Q Y V L W A S N I S S P G C
1141 TGGCTACATCGGGCAACCCAGTATGTGCTCTGGGCATCCAACATCAGCTCCCCCGGCT
400 E K V F I N T S C N P T A H L V N S S C
1201 GTGAGAAAGTGCCAATAAATACATCATGCAACCCACGGCCACCTTGTGAACCTCTCGT
420 P G L M C V F Q G Y S S K G L I Q R S V
1261 GCCCAGGGCTGATGTGCGTCTTCCAGGGCTACTCATCAAAGGCCTAATCCAACGTTCTG
440 F N L Q I Y G V L G L F W T L N W V L A
1321 TCTTCAATCTGCAAATCTATGGGGTCTGGGGCTCTTCTGGACCCTTAACCTGGGTACTGG
460 L G Q C V L A G A F A S F Y W A F H K P
1381 CCCTGGGCCAATGCGTCTCTGCTGGAGCCITTGCCTCCTTCTACTGGGCCCTCCACAAGC
480 Q D I P T F P L I S A F I R T L R Y H T
1441 CCCAGGACATCCCTACCTTCCCCCTTAATCTCTGCCTTCATCCGCACACTCCGTTACCACA
500 G S L A F G A L I L T L V Q I A R V I L
1501 CTGGGTCAITGGCATTGGAGCCCTCATCTGACCCTTGTGCAGATAGCCCGGTCACTCT
520 E Y I D H K L R G V Q N P V A R C I M C
1561 TGGAGTATATTGACCACAAGCTCAGAGGAGTGCAGAACCCTGTAGCCCGCTGCATCATGT
540 C F K C C L W C L E K F I K F L N R N A
1621 GCTGTTTCAAGTGTGCTCTGGTGTCTGGAAAAATTTATCAAGTTCTTAAACCGCAATG
560 Y I M I A I Y G K N F C V S A R N A F M
1681 CATACTCATGATCGCCATCTACGGGAAGAATTTCTGTGTCTCAGCCAAAAATGCGTTCA
580 L L M R N I V R V V V L D K V T D L L L
1741 TGCTACTCATGCGAAACATTGTGAGGGTGGTCTGTCCTGGACAAAGTCACAGACCTGTGC
600 F F G K L L V V G G V G V L S F F F F S
1801 TGTTCTTTGGGAAGCTGCTGGTGGTGGGAGGCGTGGGGTCTGTCTCTCTTTTTTTTCT
620 G R I F G L G K D F K S P H L N Y Y W L
1861 CCGGTGCGATCCCGGGGCTGGGTAAAGACTTAAGAGCCCCACCTCAACTATTACTGGC
640 P I M T S I L G A Y V I A S G F F S V F

9 or more amino acids can be bound by an HLA Class II molecule. Due to the binding groove differences between HLA Class I and II, HLA Class I motifs are length specific, i.e., position two of a Class I motif is the second amino acid in an amino to carboxyl direction of the peptide. The amino acid positions in a Class II motif are relative only to each other, not the overall peptide, i.e., additional amino acids can be attached to the amino and/or carboxyl termini of a motif-bearing sequence. HLA Class II epitopes are often 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, or 25 amino acids long, or longer than 25 amino acids.

Antibody-Based Vaccines

A wide variety of methods for generating an immune response in a mammal are known in the art (for example as the first step in the generation of hybridomas). Methods of generating an immune response in a mammal comprise exposing the mammal's immune system to an immunogenic epitope on a protein (e.g. a 24P4C12 protein) so that an immune response is generated. A typical embodiment consists of a method for generating an immune response to 24P4C12 in a host, by contacting the host with a sufficient amount of at least one 24P4C12 B cell or cytotoxic T-cell epitope or analog thereof; and at least one periodic interval thereafter re-contacting the host with the 24P4C12 B cell or cytotoxic T-cell epitope or analog thereof. A specific embodiment consists of a method of generating an immune response against a 24P4C12-related protein or a man-made multiepitopic peptide comprising: administering 24P4C12 immunogen (e.g. a 24P4C12 protein or a peptide fragment thereof, a 24P4C12 fusion protein or analog etc.) in a vaccine preparation to a human or another mammal. Typically, such vaccine preparations further contain a suitable adjuvant (see, e.g., U.S. Pat. No. 6,146,635) or a universal helper epitope such as a PADRET peptide (Epi-mune Inc., San Diego, Calif.; see, e.g., Alexander et al., *J. Immunol.* 2000 164(3): 164(3): 1625-1633; Alexander et al., *Immunity* 1994 1(9): 751-761 and Alexander et al., *Immunol. Res.* 1998 18(2): 79-92). An alternative method comprises generating an immune response in an individual against a 24P4C12 immunogen by: administering in vivo to muscle or skin of the individual's body a DNA molecule that comprises a DNA sequence that encodes a 24P4C12 immunogen, the DNA sequence operatively linked to regulatory sequences which control the expression of the DNA sequence; wherein the DNA molecule is taken up by cells, the DNA sequence is expressed in the cells and an immune response is generated against the immunogen (see, e.g., U.S. Pat. No. 5,962,428). Optionally a genetic vaccine facilitator such as anionic lipids; saponins; lectins; estrogenic compounds; hydroxylated lower alkyls; dimethyl sulfoxide; and urea is also administered. In addition, an antiidiotypic antibody can be administered that mimics 24P4C12, in order to generate a response to the target antigen.

Nucleic Acid Vaccines:

Vaccine compositions of the invention include nucleic acid-mediated modalities. DNA or RNA that encode protein(s) of the invention can be administered to a patient. Genetic immunization methods can be employed to generate prophylactic or therapeutic humoral and cellular immune responses directed against cancer cells expressing 24P4C12. Constructs comprising DNA encoding a 24P4C12-related protein/immunogen and appropriate regulatory sequences can be injected directly into muscle or skin of an individual, such that the cells of the muscle or skin take-up the construct and express the encoded 24P4C12 protein/immunogen. Alternatively, a vaccine comprises a 24P4C12-related protein. Expression of the 24P4C12-related protein immunogen results in the generation of prophylactic or therapeutic

humoral and cellular immunity against cells that bear a 24P4C12 protein. Various prophylactic and therapeutic genetic immunization techniques known in the art can be used (for review, see information and references published at Internet address genweb.com). Nucleic acid-based delivery is described, for instance, in Wolff et. al., *Science* 247:1465 (1990) as well as U.S. Pat. Nos. 5,580,859; 5,589,466; 5,804,566; 5,739,118; 5,736,524; 5,679,647; WO 98/04720. Examples of DNA-based delivery technologies include "naked DNA", facilitated (bupivacaine, polymers, peptide-mediated) delivery, cationic lipid complexes, and particle-mediated ("gene gun") or pressure-mediated delivery (see, e.g., U.S. Pat. No. 5,922,687).

For therapeutic or prophylactic immunization purposes, proteins of the invention can be expressed via viral or bacterial vectors. Various viral gene delivery systems that can be used in the practice of the invention include, but are not limited to, vaccinia, fowlpox, canarypox, adenovirus, influenza, poliovirus, adeno-associated virus, lentivirus, and sindbis virus (see, e.g., Restifo, 1996, *Curr. Opin. Immunol.* 8:658-663; Tsang et al. *J. Natl. Cancer Inst.* 87:982-990 (1995)). Non-viral delivery systems can also be employed by introducing naked DNA encoding a 24P4C12-related protein into the patient (e.g., intramuscularly or intradermally) to induce an anti-tumor response.

Vaccinia virus is used, for example, as a vector to express nucleotide sequences that encode the peptides of the invention. Upon introduction into a host, the recombinant vaccinia virus expresses the protein immunogenic peptide, and thereby elicits a host immune response. Vaccinia vectors and methods useful in immunization protocols are described in, e.g., U.S. Pat. No. 4,722,848. Another vector is BCG (Bacille Calmette Guerin). BCG vectors are described in Stover et al., *Nature* 351:456-460 (1991). A wide variety of other vectors useful for therapeutic administration or immunization of the peptides of the invention, e.g. adeno and adeno-associated virus vectors, retroviral vectors, *Salmonella typhi* vectors, detoxified anthrax toxin vectors, and the like, will be apparent to those skilled in the art from the description herein.

Thus, gene delivery systems are used to deliver a 24P4C12-related nucleic acid molecule. In one embodiment, the full-length human 24P4C12 cDNA is employed. In another embodiment, 24P4C12 nucleic acid molecules encoding specific cytotoxic T lymphocyte (CTL) and/or antibody epitopes are employed.

Ex Vivo Vaccines

Various ex vivo strategies can also be employed to generate an immune response. One approach involves the use of antigen presenting cells (APCs) such as dendritic cells (DC) to present 24P4C12 antigen to a patient's immune system. Dendritic cells express MHC class I and II molecules, B7 co-stimulator, and IL-12, and are thus highly specialized antigen presenting cells. In prostate cancer, autologous dendritic cells pulsed with peptides of the prostate-specific membrane antigen (PSMA) are being used in a Phase I clinical trial to stimulate prostate cancer patients' immune systems (Tjoa et al., 1996, *Prostate* 28:65-69; Murphy et al., 1996, *Prostate* 29:371-380). Thus, dendritic cells can be used to present 24P4C12 peptides to T cells in the context of MHC class I or II molecules. In one embodiment, autologous dendritic cells are pulsed with 24P4C12 peptides capable of binding to MHC class I and/or class II molecules. In another embodiment, dendritic cells are pulsed with the complete 24P4C12 protein. Yet another embodiment involves engineering the overexpression of a 24P4C12 gene in dendritic cells using various implementing vectors known in the art, such as adenovirus (Arthur et al., 1997, *Cancer Gene Ther.* 4:17-25), ret-

rovirus (Henderson et al., 1996, *Cancer Res.* 56:3763-3770), lentivirus, adeno-associated virus, DNA transfection (Ribas et al., 1997, *Cancer Res.* 57:2865-2869), or tumor-derived RNA transfection (Ashley et al., 1997, *J. Exp. Med.* 186:1177-1182). Cells that express 24P4C12 can also be engineered to express immune modulators, such as GM-CSF, and used as immunizing agents.

X.B.) 24P4C12 as a Target for Antibody-Based Therapy

24P4C12 is an attractive target for antibody-based therapeutic strategies. A number of antibody strategies are known in the art for targeting both extracellular and intracellular molecules (see, e.g., complement and ADCC mediated killing as well as the use of intrabodies). Because 24P4C12 is expressed by cancer cells of various lineages relative to corresponding normal cells, systemic administration of 24P4C12-immunoreactive compositions are prepared that exhibit excellent sensitivity without toxic, non-specific and/or non-target effects caused by binding of the immunoreactive composition to non-target organs and tissues. Antibodies specifically reactive with domains of 24P4C12 are useful to treat 24P4C12-expressing cancers systemically, either as conjugates with a toxin or therapeutic agent, or as naked antibodies capable of inhibiting cell proliferation or function.

24P4C12 antibodies can be introduced into a patient such that the antibody binds to 24P4C12 and modulates a function, such as an interaction with a binding partner, and consequently mediates destruction of the tumor cells and/or inhibits the growth of the tumor cells. Mechanisms by which such antibodies exert a therapeutic effect can include complement-mediated cytotoxicity, antibody-dependent cellular cytotoxicity, modulation of the physiological function of 24P4C12, inhibition of ligand binding or signal transduction pathways, modulation of tumor cell differentiation, alteration of tumor angiogenesis factor profiles, and/or apoptosis.

Those skilled in the art understand that antibodies can be used to specifically target and bind immunogenic molecules such as an immunogenic region of a 24P4C12 sequence shown in FIG. 2 or FIG. 3. In addition, skilled artisans understand that it is routine to conjugate antibodies to cytotoxic agents (see, e.g., Slevers et al. *Blood* 93:11 3678-3684 (Jun. 1, 1999)). When cytotoxic and/or therapeutic agents are delivered directly to cells, such as by conjugating them to antibodies specific for a molecule expressed by that cell (e.g. 24P4C12), the cytotoxic agent will exert its known biological effect (i.e. cytotoxicity) on those cells.

A wide variety of compositions and methods for using antibody-cytotoxic agent conjugates to kill cells are known in the art. In the context of cancers, typical methods entail administering to an animal having a tumor a biologically effective amount of a conjugate comprising a selected cytotoxic and/or therapeutic agent linked to a targeting agent (e.g. an anti-24P4C12 antibody) that binds to a marker (e.g. 24P4C12) expressed, accessible to binding or localized on the cell surfaces. A typical embodiment is a method of delivering a cytotoxic and/or therapeutic agent to a cell expressing 24P4C12, comprising conjugating the cytotoxic agent to an antibody that immunospecifically binds to a 24P4C12 epitope, and, exposing the cell to the antibody-agent conjugate. Another illustrative embodiment is a method of treating an individual suspected of suffering from metastasized cancer, comprising a step of administering parenterally to said individual a pharmaceutical composition comprising a therapeutically effective amount of an antibody conjugated to a cytotoxic and/or therapeutic agent.

Cancer immunotherapy using anti-24P4C12 antibodies can be done in accordance with various approaches that have been successfully employed in the treatment of other types of

cancer, including but not limited to colon cancer (Arlen et al., 1998, *Crit. Rev. Immunol.* 18:133-138), multiple myeloma (Ozaki et al., 1997, *Blood* 90:3179-3186, Tsunenari et al., 1997, *Blood* 90:2437-2444), gastric cancer (Kasprzyk et al., 1992, *Cancer Res.* 52:2771-2776), B-cell lymphoma (Funakoshi et al., 1996, *J. Immunother. Emphasis Tumor Immunol.* 19:93-101), leukemia (Zhong et al., 1996, *Leuk. Res.* 20:581-589), colorectal cancer (Moun et al., 1994, *Cancer Res.* 54:6160-6166; Velders et al., 1995, *Cancer Res.* 55:4398-4403), and breast cancer (Shepard et al., 1991, *J. Clin. Immunol.* 11:117-127). Some therapeutic approaches involve conjugation of naked antibody to a toxin or radioisotope, such as the conjugation of Y⁹¹ or I¹³¹ to anti-CD20 antibodies (e.g., ZevalinTM, IDEC Pharmaceuticals Corp. or BexxarTM, Coulter Pharmaceuticals), while others involve co-administration of antibodies and other therapeutic agents, such as HerceptinTM (trastuzumab) with paclitaxel (Genentech, Inc.). The antibodies can be conjugated to a therapeutic agent. To treat prostate cancer, for example, 24P4C12 antibodies can be administered in conjunction with radiation, chemotherapy or hormone ablation. Also, antibodies can be conjugated to a toxin such as calicheamicin (e.g., MylotargTM, Wyeth-Ayerst, Madison, N.J., a recombinant humanized IgG₄ kappa antibody conjugated to antitumor antibiotic calicheamicin) or a maytansinoid (e.g., taxane-based Tumor-Activated Prodrug, TAP, platform, ImmunoGen, Cambridge, Mass., also see e.g., U.S. Pat. No. 5,416,064).

Although 24P4C12 antibody therapy is useful for all stages of cancer, antibody therapy can be particularly appropriate in advanced or metastatic cancers. Treatment with the antibody therapy of the invention is indicated for patients who have received one or more rounds of chemotherapy. Alternatively, antibody therapy of the invention is combined with a chemotherapeutic or radiation regimen for patients who have not received chemotherapeutic treatment. Additionally, antibody therapy can enable the use of reduced dosages of concomitant chemotherapy, particularly for patients who do not tolerate the toxicity of the chemotherapeutic agent very well. Fan et al. (*Cancer Res.* 53:4637-4642, 1993), Prewett et al. (*International J. of Onco.* 9:217-224, 1996), and Hancock et al. (*Cancer Res.* 51:4575-4580, 1991) describe the use of various antibodies together with chemotherapeutic agents.

Although 24P4C12 antibody therapy is useful for all stages of cancer, antibody therapy can be particularly appropriate in advanced or metastatic cancers. Treatment with the antibody therapy of the invention is indicated for patients who have received one or more rounds of chemotherapy. Alternatively, antibody therapy of the invention is combined with a chemotherapeutic or radiation regimen for patients who have not received chemotherapeutic treatment. Additionally, antibody therapy can enable the use of reduced dosages of concomitant chemotherapy, particularly for patients who do not tolerate the toxicity of the chemotherapeutic agent very well.

Cancer patients can be evaluated for the presence and level of 24P4C12 expression, preferably using immunohistochemical assessments of tumor tissue, quantitative 24P4C12 imaging, or other techniques that reliably indicate the presence and degree of 24P4C12 expression. Immunohistochemical analysis of tumor biopsies or surgical specimens is preferred for this purpose. Methods for immunohistochemical analysis of tumor tissues are well known in the art.

Anti-24P4C12 monoclonal antibodies that treat prostate and other cancers include those that initiate a potent immune response against the tumor or those that are directly cytotoxic. In this regard, anti-24P4C12 monoclonal antibodies (mAbs) can elicit tumor cell lysis by either complement-mediated or antibody-dependent cell cytotoxicity (ADCC) mechanisms,

both of which require an intact Fc portion of the immunoglobulin molecule for interaction with effector cell Fc receptor sites on complement proteins. In addition, anti-24P4C12 mAbs that exert a direct biological effect on tumor growth are useful to treat cancers that express 24P4C12. Mechanisms by which directly cytotoxic mAbs act include: inhibition of cell growth, modulation of cellular differentiation, modulation of tumor angiogenesis factor profiles, and the induction of apoptosis. The mechanism(s) by which a particular anti-24P4C12 mAb exerts an anti-tumor effect is evaluated using any number of *in vitro* assays that evaluate cell death such as ADCC, ADMMC, complement-mediated cell lysis, and so forth, as is generally known in the art.

In some patients, the use of murine or other non-human monoclonal antibodies, or human/mouse chimeric mAbs can induce moderate to strong immune responses against the non-human antibody. This can result in clearance of the antibody from circulation and reduced efficacy. In the most severe cases, such an immune response can lead to the extensive formation of immune complexes which, potentially, can cause renal failure. Accordingly, preferred monoclonal antibodies used in the therapeutic methods of the invention are those that are either fully human or humanized and that bind specifically to the target 24P4C12 antigen with high affinity but exhibit low or no antigenicity in the patient.

Therapeutic methods of the invention contemplate the administration of single anti-24P4C12 mAbs as well as combinations, or cocktails, of different mAbs. Such mAb cocktails can have certain advantages inasmuch as they contain mAbs that target different epitopes, exploit different effector mechanisms or combine directly cytotoxic mAbs with mAbs that rely on immune effector functionality. Such mAbs in combination can exhibit synergistic therapeutic effects. In addition, anti-24P4C12 mAbs can be administered concomitantly with other therapeutic modalities, including but not limited to various chemotherapeutic agents, androgen-blockers, immune modulators (e.g., IL-2, GM-CSF), surgery or radiation. The anti-24P4C12 mAbs are administered in their "naked" or unconjugated form, or can have a therapeutic agent(s) conjugated to them.

Anti-24P4C12 antibody formulations are administered via any route capable of delivering the antibodies to a tumor cell. Routes of administration include, but are not limited to, intravenous, intraperitoneal, intramuscular, intratumor, intradermal, and the like. Treatment generally involves repeated administration of the anti-24P4C12 antibody preparation, via an acceptable route of administration such as intravenous injection (IV), typically at a dose in the range of about 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, or 25 mg/kg body weight. In general, doses in the range of 10-1000 mg mAb per week are effective and well tolerated.

Based on clinical experience with the Herceptin™ mAb in the treatment of metastatic breast cancer, an initial loading dose of approximately 4 mg/kg patient body weight IV, followed by weekly doses of about 2 mg/kg IV of the anti-24P4C12 mAb preparation represents an acceptable dosing regimen. Preferably, the initial loading dose is administered as a 90-minute or longer infusion. The periodic maintenance dose is administered as a 30 minute or longer infusion, provided the initial dose was well tolerated. As appreciated by those of skill in the art, various factors can influence the ideal dose regimen in a particular case. Such factors include, for example, the binding affinity and half life of the Ab or mAbs used, the degree of 24P4C12 expression in the patient, the extent of circulating shed 24P4C12 antigen, the desired steady-state antibody concentration level, frequency of treat-

ment, and the influence of chemotherapeutic or other agents used in combination with the treatment method of the invention, as well as the health status of a particular patient.

Optionally, patients should be evaluated for the levels of 24P4C12 in a given sample (e.g. the levels of circulating 24P4C12 antigen and/or 24P4C12 expressing cells) in order to assist in the determination of the most effective dosing regimen, etc. Such evaluations are also used for monitoring purposes throughout therapy, and are useful to gauge therapeutic success in combination with the evaluation of other parameters (for example, urine cytology and/or ImmunoCyt levels in bladder cancer therapy, or by analogy, serum PSA levels in prostate cancer therapy).

Anti-idiotypic anti-24P4C12 antibodies can also be used in anti-cancer therapy as a vaccine for inducing an immune response to cells expressing a 24P4C12-related protein. In particular, the generation of anti-idiotypic antibodies is well known in the art; this methodology can readily be adapted to generate anti-idiotypic anti-24P4C12 antibodies that mimic an epitope on a 24P4C12-related protein (see, for example, Wagner et al., 1997, *Hybridoma* 16: 33-40; Foon et al., 1995, *J. Clin. Invest.* 96:334-342; Herlyn et al., 1996, *Cancer Immunol. Immunother.* 43:65-76). Such an anti-idiotypic antibody can be used in cancer vaccine strategies.

X.C.) 24P4C12 as a Target for Cellular Immune Responses

Vaccines and methods of preparing vaccines that contain an immunogenically effective amount of one or more HLA-binding peptides as described herein are further embodiments of the invention. Furthermore, vaccines in accordance with the invention encompass compositions of one or more of the claimed peptides. A peptide can be present in a vaccine individually. Alternatively, the peptide can exist as a homopolymer comprising multiple copies of the same peptide, or as a heteropolymer of various peptides. Polymers have the advantage of increased immunological reaction and, where different peptide epitopes are used to make up the polymer, the additional ability to induce antibodies and/or CTLs that react with different antigenic determinants of the pathogenic organism or tumor-related peptide targeted for an immune response. The composition can be a naturally occurring region of an antigen or can be prepared, e.g., recombinantly or by chemical synthesis.

Carriers that can be used with vaccines of the invention are well known in the art, and include, e.g., thyroglobulin, albumins such as human serum albumin, tetanus toxoid, polyamino acids such as poly L-lysine, poly L-glutamic acid, influenza, hepatitis B virus core protein, and the like. The vaccines can contain a physiologically tolerable (i.e., acceptable) diluent such as water, or saline, preferably phosphate buffered saline. The vaccines also typically include an adjuvant. Adjuvants such as incomplete Freund's adjuvant, aluminum phosphate, aluminum hydroxide, or alum are examples of materials well known in the art. Additionally, as disclosed herein, CTL responses can be primed by conjugating peptides of the invention to lipids, such as tripalmitoyl-S-glycerylcysteinylserine (P₃CSS). Moreover, an adjuvant such as a synthetic cytosine-phosphorothiolated-guanine-containing (CpG) oligonucleotides has been found to increase CTL responses 10- to 100-fold. (see, e.g. Davila and Celis, *J. Immunol.* 165:539-547 (2000))

Upon immunization with a peptide composition in accordance with the invention, via injection, aerosol, oral, transdermal, transmucosal, intrapleural, intrathecal, or other suitable routes, the immune system of the host responds to the vaccine by producing large amounts of CTLs and/or HTLs specific for the desired antigen. Consequently, the host becomes at least partially immune to later development of

cells that express or overexpress 24P4C12 antigen, or derives at least some therapeutic benefit when the antigen was tumor-associated.

In some embodiments, it may be desirable to combine the class I peptide components with components that induce or facilitate neutralizing antibody and or helper T cell responses directed to the target antigen. A preferred embodiment of such a composition comprises class I and class II epitopes in accordance with the invention. An alternative embodiment of such a composition comprises a class I and/or class II epitope in accordance with the invention, along with a cross reactive HTL epitope such as PADRE™ (Epimmune, San Diego, Calif.) molecule (described e.g., in U.S. Pat. No. 5,736,142).

A vaccine of the invention can also include antigen-presenting cells (APC), such as dendritic cells (DC), as a vehicle to present peptides of the invention. Vaccine compositions can be created in vitro, following dendritic cell mobilization and harvesting, whereby loading of dendritic cells occurs in vitro. For example, dendritic cells are transfected, e.g., with a minigene in accordance with the invention, or are pulsed with peptides. The dendritic cell can then be administered to a patient to elicit immune responses in vivo. Vaccine compositions, either DNA- or peptide-based, can also be administered in vivo in combination with dendritic cell mobilization whereby loading of dendritic cells occurs in vivo.

Preferably, the following principles are utilized when selecting an array of epitopes for inclusion in a polyepitopic composition for use in a vaccine, or for selecting discrete epitopes to be included in a vaccine and/or to be encoded by nucleic acids such as a minigene. It is preferred that each of the following principles be balanced in order to make the selection. The multiple epitopes to be incorporated in a given vaccine composition may be, but need not be, contiguous in sequence in the native antigen from which the epitopes are derived.

1.) Epitopes are selected which, upon administration, mimic immune responses that have been observed to be correlated with tumor clearance. For HLA Class I this includes 3-4 epitopes that come from at least one tumor associated antigen (TAA). For HLA Class II a similar rationale is employed; again 3-4 epitopes are selected from at least one TAA (see, e.g., Rosenberg et al., *Science* 278:1447-1450). Epitopes from one TAA may be used in combination with epitopes from one or more additional TAAs to produce a vaccine that targets tumors with varying expression patterns of frequently-expressed TMs.

2.) Epitopes are selected that have the requisite binding affinity established to be correlated with immunogenicity: for HLA Class I an IC_{50} of 500 nM or less, often 200 nM or less; and for Class II an IC_{50} of 1000 nM or less.

3.) Sufficient supermotif bearing-peptides, or a sufficient array of allele-specific motif-bearing peptides, are selected to give broad population coverage. For example, it is preferable to have at least 80% population coverage. A Monte Carlo analysis, a statistical evaluation known in the art, can be employed to assess the breadth, or redundancy of, population coverage.

4.) When selecting epitopes from cancer-related antigens it is often useful to select analogs because the patient may have developed tolerance to the native epitope.

5.) Of particular relevance are epitopes referred to as "nested epitopes." Nested epitopes occur where at least two epitopes overlap in a given peptide sequence. A nested peptide sequence can comprise B cell, HLA class I and/or HLA class II epitopes. When providing nested epitopes, a general objective is to provide the greatest number of epitopes per sequence. Thus, an aspect is to avoid providing a peptide that

is any longer than the amino terminus of the amino terminal epitope and the carboxyl terminus of the carboxyl terminal epitope in the peptide. When providing a multi-epitopic sequence, such as a sequence comprising nested epitopes, it is generally important to screen the sequence in order to insure that it does not have pathological or other deleterious biological properties.

6.) If a polyepitopic protein is created, or when creating a minigene, an objective is to generate the smallest peptide that encompasses the epitopes of interest. This principle is similar, if not the same as that employed when selecting a peptide comprising nested epitopes. However, with an artificial polyepitopic peptide, the size minimization objective is balanced against the need to integrate any spacer sequences between epitopes in the polyepitopic protein. Spacer amino acid residues can, for example, be introduced to avoid junctional epitopes (an epitope recognized by the immune system, not present in the target antigen, and only created by the man-made juxtaposition of epitopes), or to facilitate cleavage between epitopes and thereby enhance epitope presentation. Junctional epitopes are generally to be avoided because the recipient may generate an immune response to that non-native epitope. Of particular concern is a junctional epitope that is a "dominant epitope." A dominant epitope may lead to such a zealous response that immune responses to other epitopes are diminished or suppressed.

7.) Where the sequences of multiple variants of the same target protein are present, potential peptide epitopes can also be selected on the basis of their conservancy. For example, a criterion for conservancy may define that the entire sequence of an HLA class I binding peptide or the entire 9-mer core of a class II binding peptide be conserved in a designated percentage of the sequences evaluated for a specific protein antigen.

X.C.1. Minigene Vaccines

A number of different approaches are available which allow simultaneous delivery of multiple epitopes. Nucleic acids encoding the peptides of the invention are a particularly useful embodiment of the invention. Epitopes for inclusion in a minigene are preferably selected according to the guidelines set forth in the previous section. A preferred means of administering nucleic acids encoding the peptides of the invention uses minigene constructs encoding a peptide comprising one or multiple epitopes of the invention.

The use of multi-epitope minigenes is described below and in, Ishioka et al., *J. Immunol.* 162:3915-3925, 1999; An, L. and Whitton, J. L., *J. Virol.* 71:2292, 1997; Thomson, S. A. et al., *J. Immunol.* 157:822, 1996; Whitton, J. L. et al., *J. Virol.* 67:348, 1993; Hanke, R. et al., *Vaccine* 16:426, 1998. For example, a multi-epitope DNA plasmid encoding supermotif- and/or motif-bearing epitopes derived 24P4C12, the PADRE® universal helper T cell epitope or multiple HTL epitopes from 24P4C12 (see e.g., Tables VIII-XXI and XXII to XLIX), and an endoplasmic reticulum-translocating signal sequence can be engineered. A vaccine may also comprise epitopes that are derived from other TAAs.

The immunogenicity of a multi-epitopic minigene can be confirmed in transgenic mice to evaluate the magnitude of CTL induction responses against the epitopes tested. Further, the immunogenicity of DNA-encoded epitopes in vivo can be correlated with the in vitro responses of specific CTL lines against target cells transfected with the DNA plasmid. Thus, these experiments can show that the minigene serves to both: 1.) generate a CTL response and 2.) that the induced CTLs recognized cells expressing the encoded epitopes.

For example, to create a DNA sequence encoding the selected epitopes (minigene) for expression in human cells,

the amino acid sequences of the epitopes may be reverse translated. A human codon usage table can be used to guide the codon choice for each amino acid. These epitope-encoding DNA sequences may be directly adjoined, so that when translated, a continuous polypeptide sequence is created. To optimize expression and/or immunogenicity, additional elements can be incorporated into the minigene design. Examples of amino acid sequences that can be reverse translated and included in the minigene sequence include: HLA class I epitopes, HLA class II epitopes, antibody epitopes, a ubiquitination signal sequence, and/or an endoplasmic reticulum targeting signal. In addition, HLA presentation of CTL and HTL epitopes may be improved by including synthetic (e.g. poly-alanine) or naturally-occurring flanking sequences adjacent to the CTL or HTL epitopes; these larger peptides comprising the epitope(s) are within the scope of the invention.

The minigene sequence may be converted to DNA by assembling oligonucleotides that encode the plus and minus strands of the minigene. Overlapping oligonucleotides (30-100 bases long) may be synthesized, phosphorylated, purified and annealed under appropriate conditions using well known techniques. The ends of the oligonucleotides can be joined, for example, using T4 DNA ligase. This synthetic minigene, encoding the epitope polypeptide, can then be cloned into a desired expression vector.

Standard regulatory sequences well known to those of skill in the art are preferably included in the vector to ensure expression in the target cells. Several vector elements are desirable: a promoter with a down-stream cloning site for minigene insertion; a polyadenylation signal for efficient transcription termination; an *E. coli* origin of replication; and an *E. coli* selectable marker (e.g. ampicillin or kanamycin resistance). Numerous promoters can be used for this purpose, e.g., the human cytomegalovirus (hCMV) promoter. See, e.g., U.S. Pat. Nos. 5,580,859 and 5,589,466 for other suitable promoter sequences.

Additional vector modifications may be desired to optimize minigene expression and immunogenicity. In some cases, introns are required for efficient gene expression, and one or more synthetic or naturally-occurring introns could be incorporated into the transcribed region of the minigene. The inclusion of mRNA stabilization sequences and sequences for replication in mammalian cells may also be considered for increasing minigene expression.

Once an expression vector is selected, the minigene is cloned into the polylinker region downstream of the promoter. This plasmid is transformed into an appropriate *E. coli* strain, and DNA is prepared using standard techniques. The orientation and DNA sequence of the minigene, as well as all other elements included in the vector, are confirmed using restriction mapping and DNA sequence analysis. Bacterial cells harboring the correct plasmid can be stored as a master cell bank and a working cell bank.

In addition, immunostimulatory sequences (ISSs or CpGs) appear to play a role in the immunogenicity of DNA vaccines. These sequences may be included in the vector, outside the minigene coding sequence, if desired to enhance immunogenicity.

In some embodiments, a bi-cistronic expression vector which allows production of both the minigene-encoded epitopes and a second protein (included to enhance or decrease immunogenicity) can be used. Examples of proteins or polypeptides that could beneficially enhance the immune response if co-expressed include cytokines (e.g., IL-2, IL-12, GM-CSF), cytokine-inducing molecules (e.g., LfL), costimulatory molecules, or for HTL responses, pan-DR

binding proteins (PADRE™, Epimmune, San Diego, Calif.). Helper (HTL) epitopes can be joined to intracellular targeting signals and expressed separately from expressed CTL epitopes; this allows direction of the HTL epitopes to a cell compartment different than that of the CTL epitopes. If required, this could facilitate more efficient entry of HTL epitopes into the HLA class II pathway, thereby improving HTL induction. In contrast to HTL or CTL induction, specifically decreasing the immune response by co-expression of immunosuppressive molecules (e.g. TGF-β) may be beneficial in certain diseases.

Therapeutic quantities of plasmid DNA can be produced for example, by fermentation in *E. coli*, followed by purification. Aliquots from the working cell bank are used to inoculate growth medium, and grown to saturation in shaker flasks or a bioreactor according to well-known techniques. Plasmid DNA can be purified using standard bioseparation technologies such as solid phase anion-exchange resins supplied by QIAGEN, Inc. (Valencia, Calif.). If required, supercoiled DNA can be isolated from the open circular and linear forms using gel electrophoresis or other methods.

Purified plasmid DNA can be prepared for injection using a variety of formulations. The simplest of these is reconstitution of lyophilized DNA in sterile phosphate-buffer saline (PBS). This approach, known as "naked DNA," is currently being used for intramuscular (IM) administration in clinical trials. To maximize the immunotherapeutic effects of minigene DNA vaccines, an alternative method for formulating purified plasmid DNA may be desirable. A variety of methods have been described, and new techniques may become available. Cationic lipids, glycolipids, and fusogenic liposomes can also be used in the formulation (see, e.g., as described by WO 93/24640; Mannino & Gould-Fogerite, *Bio Techniques* 6(7): 682 (1988); U.S. Pat. No. 5,279,833; WO 91/06309; and Feigner, et al., *Proc. Nat'l Acad. Sci. USA* 84:7413 (1987). In addition, peptides and compounds referred to collectively as protective, interactive, non-condensing compounds (PINC) could also be complexed to purified plasmid DNA to influence variables such as stability, intramuscular dispersion, or trafficking to specific organs or cell types.

Target cell sensitization can be used as a functional assay for expression and HLA class I presentation of minigene-encoded CTL epitopes. For example, the plasmid DNA is introduced into a mammalian cell line that is suitable as a target for standard CTL chromium release assays. The transfection method used will be dependent on the final formulation. Electroporation can be used for "naked" DNA, whereas cationic lipids allow direct in vitro transfection. A plasmid expressing green fluorescent protein (GFP) can be co-transfected to allow enrichment of transfected cells using fluorescence activated cell sorting (FACS). These cells are then chromium-51 (⁵¹Cr) labeled and used as target cells for epitope-specific CTL lines; cytotoxicity, detected by ⁵¹Cr release, indicates both production of, and HLA presentation of, minigene-encoded CTL epitopes. Expression of HTL epitopes may be evaluated in an analogous manner using assays to assess HTL activity.

In vivo immunogenicity is a second approach for functional testing of minigene DNA formulations. Transgenic mice expressing appropriate human HLA proteins are immunized with the DNA product. The dose and route of administration are formulation dependent (e.g., IM for DNA in PBS, intraperitoneal (i.p.) for lipid-complexed DNA). Twenty-one days after immunization, splenocytes are harvested and restimulated for one week in the presence of peptides encoding each epitope being tested. Thereafter, for CTL effector cells, assays are conducted for cytotoxicity of peptide-loaded,

⁵¹Cr-labeled target cells using standard techniques. Lysis of target cells that were sensitized by HLA loaded with peptide epitopes, corresponding to minigene-encoded epitopes, demonstrates DNA vaccine function for in vivo induction of CTLs. Immunogenicity of HTL epitopes is confirmed in transgenic mice in an analogous manner.

Alternatively, the nucleic acids can be administered using ballistic delivery as described, for instance, in U.S. Pat. No. 5,204,253. Using this technique, particles comprised solely of DNA are administered. In a further alternative embodiment, DNA can be adhered to particles, such as gold particles.

Minigenes can also be delivered using other bacterial or viral delivery systems well known in the art, e.g., an expression construct encoding epitopes of the invention can be incorporated into a viral vector such as vaccinia.

X.C.2. Combinations of CTL Peptides with Helper Peptides

Vaccine compositions comprising CTL peptides of the invention can be modified, e.g., analoged, to provide desired attributes, such as improved serum half life, broadened population coverage or enhanced immunogenicity.

For instance, the ability of a peptide to induce CTL activity can be enhanced by linking the peptide to a sequence which contains at least one epitope that is capable of inducing a T helper cell response. Although a CTL peptide can be directly linked to a T helper peptide, often CTL epitope/HTL epitope conjugates are linked by a spacer molecule. The spacer is typically comprised of relatively small, neutral molecules, such as amino acids or amino acid mimetics, which are substantially uncharged under physiological conditions. The spacers are typically selected from, e.g., Ala, Gly, or other neutral spacers of nonpolar amino acids or neutral polar amino acids. It will be understood that the optionally present spacer need not be comprised of the same residues and thus may be a hetero- or homo-oligomer. When present, the spacer will usually be at least one or two residues, more usually three to six residues and sometimes 10 or more residues. The CTL peptide epitope can be linked to the T helper peptide epitope either directly or via a spacer either at the amino or carboxy terminus of the CTL peptide. The amino terminus of either the immunogenic peptide or the T helper peptide may be acylated.

In certain embodiments, the T helper peptide is one that is recognized by T helper cells present in a majority of a genetically diverse population. This can be accomplished by selecting peptides that bind to many, most, or all of the HLA class II molecules. Examples of such amino acid bind many HLA Class II molecules include sequences from antigens such as tetanus toxoid at positions 830-843 (QYIKANSKFIGITE; SEQ ID NO: 29), *Plasmodium falciparum* circumsporozoite (CS) protein at positions 378-398 (DIEKKIAKMEKASS-VFNVVNS; SEQ ID NO: 30), and *Streptococcus* 18 kD protein at positions 116-131 (GAVDSILGGVATYGAA; SEQ ID NO: 31). Other examples include peptides bearing a DR 1-4-7 supermotif, or either of the DR3 motifs.

Alternatively, it is possible to prepare synthetic peptides capable of stimulating T helper lymphocytes, in a loosely HLA-restricted fashion, using amino acid sequences not found in nature (see, e.g., PCT publication WO 95/07707). These synthetic compounds called Pan-DR-binding epitopes (e.g., PADRE™, Epimmune, Inc., San Diego, Calif.) are designed, most preferably, to bind most HLA-DR (human HLA class II) molecules. For instance, a pan-DR-binding epitope peptide having the formula: AKXVAAWTLKAAA (SEQ ID NO: 32), where "X" is either cyclohexylalanine, phenylalanine, or tyrosine, and a is either D-alanine or L-alanine, has been found to bind to most HLA-DR alleles, and to

stimulate the response of T helper lymphocytes from most individuals, regardless of their HLA type. An alternative of a pan-DR binding epitope comprises all "L" natural amino acids and can be provided in the form of nucleic acids that encode the epitope.

HTL peptide epitopes can also be modified to alter their biological properties. For example, they can be modified to include D-amino acids to increase their resistance to proteases and thus extend their serum half life, or they can be conjugated to other molecules such as lipids, proteins, carbohydrates, and the like to increase their biological activity. For example, a T helper peptide can be conjugated to one or more palmitic acid chains at either the amino or carboxyl termini.

X.C.3. Combinations of CTL Peptides with T Cell Priming Agents

In some embodiments it may be desirable to include in the pharmaceutical compositions of the invention at least one component which primes B lymphocytes or T lymphocytes. Lipids have been identified as agents capable of priming CTL in vivo. For example, palmitic acid residues can be attached to the ϵ - and α -amino groups of a lysine residue and then linked, e.g., via one or more linking residues such as Gly, Gly-Gly-, Ser, Ser-Ser, or the like, to an immunogenic peptide. The lipidated peptide can then be administered either directly in a micelle or particle, incorporated into a liposome, or emulsified in an adjuvant, e.g., incomplete Freund's adjuvant. In a preferred embodiment, a particularly effective immunogenic composition comprises palmitic acid attached to ϵ - and α -amino groups of Lys, which is attached via linkage, e.g., Ser-Ser, to the amino terminus of the immunogenic peptide.

As another example of lipid priming of CTL responses, *E. coli* lipoproteins, such as tripalmitoyl-S-glycerylcysteinylserine (P₃CSS) can be used to prime virus specific CTL when covalently attached to an appropriate peptide (see, e.g., Deres, et al., *Nature* 342:561, 1989). Peptides of the invention can be coupled to P₃CSS, for example, and the lipopeptide administered to an individual to prime specifically an immune response to the target antigen. Moreover, because the induction of neutralizing antibodies can also be primed with P₃CSS-conjugated epitopes, two such compositions can be combined to more effectively elicit both humoral and cell-mediated responses.

X.C.4. Vaccine Compositions Comprising DC Pulsed with CTL and/or HTL Peptides

An embodiment of a vaccine composition in accordance with the invention comprises ex vivo administration of a cocktail of epitope-bearing peptides to PBMC, or isolated DC therefrom, from the patient's blood. A pharmaceutical to facilitate harvesting of DC can be used, such as Progenipoint™ (Pharmacia-Monsanto, St. Louis, Mo.) or GM-CSF/IL-4. After pulsing the DC with peptides and prior to reinfusion into patients, the DC are washed to remove unbound peptides. In this embodiment, a vaccine comprises peptide-pulsed DCs which present the pulsed peptide epitopes complexed with HLA molecules on their surfaces.

The DC can be pulsed ex vivo with a cocktail of peptides, some of which stimulate CTL responses to 24P4C12. Optionally, a helper T cell (HTL) peptide, such as a natural or artificial loosely restricted HLA Class II peptide, can be included to facilitate the CTL response. Thus, a vaccine in accordance with the invention is used to treat a cancer which expresses or overexpresses 24P4C12.

X.D. Adoptive Immunotherapy

Antigenic 24P4C12-related peptides are used to elicit a CTL and/or HTL response ex vivo, as well. The resulting CTL or HTL cells, can be used to treat tumors in patients that do not respond to other conventional forms of therapy, or will

not respond to a therapeutic vaccine peptide or nucleic acid in accordance with the invention. Ex vivo CTL or HTL responses to a particular antigen are induced by incubating in tissue culture the patient's, or genetically compatible, CTL or HTL precursor cells together with a source of antigen-presenting cells (APC), such as dendritic cells, and the appropriate immunogenic peptide. After an appropriate incubation time (typically about 7-28 days), in which the precursor cells are activated and expanded into effector cells, the cells are infused back into the patient, where they will destroy (CTL) or facilitate destruction (HTL) of their specific target cell (e.g., a tumor cell). Transfected dendritic cells may also be used as antigen presenting cells.

X.E. Administration of Vaccines for Therapeutic or Prophylactic Purposes

Pharmaceutical and vaccine compositions of the invention are typically used to treat and/or prevent a cancer that expresses or overexpresses 24P4C12. In therapeutic applications, peptide and/or nucleic acid compositions are administered to a patient in an amount sufficient to elicit an effective B cell, CTL and/or HTL response to the antigen and to cure or at least partially arrest or slow symptoms and/or complications. An amount adequate to accomplish this is defined as "therapeutically effective dose." Amounts effective for this use will depend on, e.g., the particular composition administered, the manner of administration, the stage and severity of the disease being treated, the weight and general state of health of the patient, and the judgment of the prescribing physician.

For pharmaceutical compositions, the immunogenic peptides of the invention, or DNA encoding them, are generally administered to an individual already bearing a tumor that expresses 24P4C12. The peptides or DNA encoding them can be administered individually or as fusions of one or more peptide sequences. Patients can be treated with the immunogenic peptides separately or in conjunction with other treatments, such as surgery, as appropriate.

For therapeutic use, administration should generally begin at the first diagnosis of 24P4C12-associated cancer. This is followed by boosting doses until at least symptoms are substantially abated and for a period thereafter. The embodiment of the vaccine composition (i.e., including, but not limited to embodiments such as peptide cocktails, polyepitopic polypeptides, minigenes, or TAA-specific CTLs or pulsed dendritic cells) delivered to the patient may vary according to the stage of the disease or the patient's health status. For example, in a patient with a tumor that expresses 24P4C12, a vaccine comprising 24P4C12-specific CTL may be more efficacious in killing tumor cells in patient with advanced disease than alternative embodiments.

It is generally important to provide an amount of the peptide epitope delivered by a mode of administration sufficient to stimulate effectively a cytotoxic T cell response; compositions which stimulate helper T cell responses can also be given in accordance with this embodiment of the invention.

The dosage for an initial therapeutic immunization generally occurs in a unit dosage range where the lower value is about 1, 5, 50, 500, or 1,000 µg and the higher value is about 10,000; 20,000; 30,000; or 50,000 µg. Dosage values for a human typically range from about 500 µg to about 50,000 µg per 70 kilogram patient. Boosting dosages of between about 1.0 µg to about 50,000 µg of peptide pursuant to a boosting regimen over weeks to months may be administered depending upon the patient's response and condition as determined by measuring the specific activity of CTL and HTL obtained from the patient's blood. Administration should continue until at least clinical symptoms or laboratory tests indicate

that the neoplasia, has been eliminated or reduced and for a period thereafter. The dosages, routes of administration, and dose schedules are adjusted in accordance with methodologies known in the art.

In certain embodiments, the peptides and compositions of the present invention are employed in serious disease states, that is, life-threatening or potentially life threatening situations. In such cases, as a result of the minimal amounts of extraneous substances and the relative nontoxic nature of the peptides in preferred compositions of the invention, it is possible and may be felt desirable by the treating physician to administer substantial excesses of these peptide compositions relative to these stated dosage amounts.

The vaccine compositions of the invention can also be used purely as prophylactic agents. Generally the dosage for an initial prophylactic immunization generally occurs in a unit dosage range where the lower value is about 1, 5, 50, 500, or 1000 µg and the higher value is about 10,000; 20,000; 30,000; or 50,000 µg. Dosage values for a human typically range from about 500 µg to about 50,000 µg per 70 kilogram patient. This is followed by boosting dosages of between about 1.0 µg to about 50,000 µg of peptide administered at defined intervals from about four weeks to six months after the initial administration of vaccine. The immunogenicity of the vaccine can be assessed by measuring the specific activity of CTL and HTL obtained from a sample of the patient's blood.

The pharmaceutical compositions for therapeutic treatment are intended for parenteral, topical, oral, nasal, intrathecal, or local (e.g. as a cream or topical ointment) administration. Preferably, the pharmaceutical compositions are administered parentally, e.g., intravenously, subcutaneously, intradermally, or intramuscularly. Thus, the invention provides compositions for parenteral administration which comprise a solution of the immunogenic peptides dissolved or suspended in an acceptable carrier, preferably an aqueous carrier.

A variety of aqueous carriers may be used, e.g., water, buffered water, 0.8% saline, 0.3% glycine, hyaluronic acid and the like. These compositions may be sterilized by conventional, well-known sterilization techniques, or may be sterile filtered. The resulting aqueous solutions may be packaged for use as is, or lyophilized, the lyophilized preparation being combined with a sterile solution prior to administration.

The compositions may contain pharmaceutically acceptable auxiliary substances as required to approximate physiological conditions, such as pH-adjusting and buffering agents, tonicity adjusting agents, wetting agents, preservatives, and the like, for example, sodium acetate, sodium lactate, sodium chloride, potassium chloride, calcium chloride, sorbitan monolaurate, triethanolamine oleate, etc.

The concentration of peptides of the invention in the pharmaceutical formulations can vary widely, i.e., from less than about 0.1%, usually at or at least about 2% to as much as 20% to 50% or more by weight, and will be selected primarily by fluid volumes, viscosities, etc., in accordance with the particular mode of administration selected.

A human unit dose form of a composition is typically included in a pharmaceutical composition that comprises a human unit dose of an acceptable carrier, in one embodiment an aqueous carrier, and is administered in a volume/quantity that is known by those of skill in the art to be used for administration of such compositions to humans (see, e.g., Remington's Pharmaceutical Sciences, 17th Edition, A. Gennaro, Editor, Mack Publishing Co., Easton, Pa., 1985). For example a peptide dose for initial immunization can be from about 1 to about 50,000 µg, generally 100-5,000 µg, for a 70 kg patient. For example, for nucleic acids an initial

immunization may be performed using an expression vector in the form of naked nucleic acid administered IM (or SC or ID) in the amounts of 0.5-5 mg at multiple sites. The nucleic acid (0.1 to 1000 µg) can also be administered using a gene gun. Following an incubation period of 3-4 weeks, a booster dose is then administered. The booster can be recombinant fowlpox virus administered at a dose of $5 \cdot 10^7$ to $5 \cdot 10^9$ pfu.

For antibodies, a treatment generally involves repeated administration of the anti-24P4C12 antibody preparation, via an acceptable route of administration such as intravenous injection (IV), typically at a dose in the range of about 0.1 to about 10 mg/kg body weight. In general, doses in the range of 10-500 mg mAb per week are effective and well tolerated. Moreover, an initial loading dose of approximately 4 mg/kg patient body weight IV, followed by weekly doses of about 2 mg/kg IV of the anti-24P4C12 mAb preparation represents an acceptable dosing regimen. As appreciated by those of skill in the art, various factors can influence the ideal dose in a particular case. Such factors include, for example, half life of a composition, the binding affinity of an Ab, the immunogenicity of a substance, the degree of 24P4C12 expression in the patient, the extent of circulating shed 24P4C12 antigen, the desired steady-state concentration level, frequency of treatment, and the influence of chemotherapeutic or other agents used in combination with the treatment method of the invention, as well as the health status of a particular patient. Non-limiting preferred human unit doses are, for example, 500 µg-1 mg, 1 mg-50 mg, 50 mg-100 mg, 100 mg-200 mg, 200 mg-300 mg, 400 mg-500 mg, 500 mg-600 mg, 600 mg-700 mg, 700 mg-800 mg, 800 mg-900 mg, 900 mg-1 g, or 1 mg-700 mg. In certain embodiments, the dose is in a range of 2-5 mg/kg body weight, e.g., with follow on weekly doses of 1-3 mg/kg; 0.5 mg, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 mg/kg body weight followed, e.g., in two, three or four weeks by weekly doses; 0.5-10 mg/kg body weight, e.g., followed in two, three or four weeks by weekly doses; 225, 250, 275, 300, 325, 350, 375, 400 mg m² of body area weekly; 1-600 mg m² of body area weekly; 225-400 mg m² of body area weekly; these doses can be followed by weekly doses for 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12 or more weeks.

In one embodiment, human unit dose forms of polynucleotides comprise a suitable dosage range or effective amount that provides any therapeutic effect. As appreciated by one of ordinary skill in the art a therapeutic effect depends on a number of factors, including the sequence of the polynucleotide, molecular weight of the polynucleotide and route of administration. Dosages are generally selected by the physician or other health care professional in accordance with a variety of parameters known in the art, such as severity of symptoms, history of the patient and the like. Generally, for a polynucleotide of about 20 bases, a dosage range may be selected from, for example, an independently selected lower limit such as about 0.1, 0.25, 0.5, 1, 2, 5, 10, 20, 30, 40, 50, 60, 70, 80, 90, 100, 200, 300, 400 or 500 mg/kg up to an independently selected upper limit, greater than the lower limit, of about 60, 80, 100, 200, 300, 400, 500, 750, 1000, 1500, 2000, 3000, 4000, 5000, 6000, 7000, 8000, 9000 or 10,000 mg/kg. For example, a dose may be about any of the following: 0.1 to 100 mg/kg, 0.1 to 50 mg/kg, 0.1 to 25 mg/kg, 0.1 to 10 mg/kg, 1 to 500 mg/kg, 100 to 400 mg/kg, 200 to 300 mg/kg, 1 to 100 mg/kg, 100 to 200 mg/kg, 300 to 400 mg/kg, 400 to 500 mg/kg, 500 to 1000 mg/kg, 500 to 5000 mg/kg, or 500 to 10,000 mg/kg. Generally, parenteral routes of administration may require higher doses of polynucleotide compared to more direct application to the nucleotide to diseased tissue, as do polynucleotides of increasing length.

In one embodiment, human unit dose forms of T-cells comprise a suitable dosage range or effective amount that provides any therapeutic effect. As appreciated by one of ordinary skill in the art, a therapeutic effect depends on a number of factors. Dosages are generally selected by the physician or other health care professional in accordance with a variety of parameters known in the art, such as severity of symptoms, history of the patient and the like. A dose may be about 10^4 cells to about 10^6 cells, about 10^6 cells to about 10^8 cells, about 10^8 to about 10^{11} cells, or about 10^8 to about $5 \cdot 10^{10}$ cells. A dose may also be about 10^6 cells/m² to about 10^{10} cells/m², or about 10^6 cells/m² to about 10^8 cells/m².

Proteins(s) of the invention, and/or nucleic acids encoding the protein(s), can also be administered via liposomes, which may also serve to: 1) target the proteins(s) to a particular tissue, such as lymphoid tissue; 2) to target selectively to diseases cells; or, 3) to increase the half-life of the peptide composition. Liposomes include emulsions, foams, micelles, insoluble monolayers, liquid crystals, phospholipid dispersions, lamellar layers and the like. In these preparations, the peptide to be delivered is incorporated as part of a liposome, alone or in conjunction with a molecule which binds to a receptor prevalent among lymphoid cells, such as monoclonal antibodies which bind to the CD45 antigen, or with other therapeutic or immunogenic compositions. Thus, liposomes either filled or decorated with a desired peptide of the invention can be directed to the site of lymphoid cells, where the liposomes then deliver the peptide compositions. Liposomes for use in accordance with the invention are formed from standard vesicle-forming lipids, which generally include neutral and negatively charged phospholipids and a sterol, such as cholesterol. The selection of lipids is generally guided by consideration of, e.g., liposome size, acid liability and stability of the liposomes in the blood stream. A variety of methods are available for preparing liposomes, as described in, e.g., Szoka, et al., *Ann. Rev. Biophys. Bloeng.* 9:467 (1980), and U.S. Pat. Nos. 4,235,871, 4,501,728, 4,837,028, and 5,019,369.

For targeting cells of the immune system, a ligand to be incorporated into the liposome can include, e.g., antibodies or fragments thereof specific for cell surface determinants of the desired immune system cells. A liposome suspension containing a peptide may be administered intravenously, locally, topically, etc. in a dose which varies according to, inter alia, the manner of administration, the peptide being delivered, and the stage of the disease being treated.

For solid compositions, conventional nontoxic solid carriers may be used which include, for example, pharmaceutical grades of mannitol, lactose, starch, magnesium stearate, sodium saccharin, talcum, cellulose, glucose, sucrose, magnesium carbonate, and the like. For oral administration, a pharmaceutically acceptable nontoxic composition is formed by incorporating any of the normally employed excipients, such as those carriers previously listed, and generally 10-95% of active ingredient, that is, one or more peptides of the invention, and more preferably at a concentration of 25%-75%.

For aerosol administration, immunogenic peptides are preferably supplied in finely divided form along with a surfactant and propellant. Typical percentages of peptides are about 0.01%-20% by weight, preferably about 1%-10%. The surfactant must, of course, be nontoxic, and preferably soluble in the propellant. Representative of such agents are the esters or partial esters of fatty acids containing from about 6 to 22 carbon atoms, such as caproic, octanoic, lauric, palmitic, stearic, linoleic, linolenic, oleosteric and oleic acids with an aliphatic polyhydric alcohol or its cyclic anhydride. Mixed

esters, such as mixed or natural glycerides may be employed. The surfactant may constitute about 0.1%-20% by weight of the composition, preferably about 0.25-5%. The balance of the composition is ordinarily propellant. A carrier can also be included, as desired, as with, e.g., lecithin for intranasal delivery.

XI.) DIAGNOSTIC AND PROGNOSTIC EMBODIMENTS OF 24P4C12

As disclosed herein, 24P4C12 polynucleotides, polypeptides, reactive cytotoxic T cells (CTL), reactive helper T cells (HTL) and anti-polypeptide antibodies are used in well known diagnostic, prognostic and therapeutic assays that examine conditions associated with dysregulated cell growth such as cancer, in particular the cancers listed in Table I (see, e.g., both its specific pattern of tissue expression as well as its overexpression in certain cancers as described for example in the Example entitled "Expression analysis of 24P4C12 in normal tissues, and patient specimens").

24P4C12 can be analogized to a prostate associated antigen PSA, the archetypal marker that has been used by medical practitioners for years to identify and monitor the presence of prostate cancer (see, e.g., Merrill et al., *J. Urol.* 163(2): 503-5120 (2000); Polascik et al., *J. Urol.* August; 162(2):293-306 (1999) and Fortier et al., *J. Nat. Cancer Inst.* 91(19): 1635-1640 (1999)). A variety of other diagnostic markers are also used in similar contexts including p53 and K-ras (see, e.g., Tulchinsky et al., *Int J Mol Med* 1999 July 4(1):99-102 and Minimoto et al., *Cancer Detect Prev* 2000; 24(1):1-12). Therefore, this disclosure of 24P4C12 polynucleotides and polypeptides (as well as 24P4C12 polynucleotide probes and anti-24P4C12 antibodies used to identify the presence of these molecules) and their properties allows skilled artisans to utilize these molecules in methods that are analogous to those used, for example, in a variety of diagnostic assays directed to examining conditions associated with cancer.

Typical embodiments of diagnostic methods which utilize the 24P4C12 polynucleotides, polypeptides, reactive T cells and antibodies are analogous to those methods from well-established diagnostic assays, which employ, e.g., PSA polynucleotides, polypeptides, reactive T cells and antibodies. For example, just as PSA polynucleotides are used as probes (for example in Northern analysis, see, e.g., Sharief et al., *Biochem. Mol. Biol. Int.* 33(3):567-74 (1994)) and primers (for example in PCR analysis, see, e.g., Okegawa et al., *J. Urol.* 163(4): 1189-1190 (2000)) to observe the presence and/or the level of PSA mRNAs in methods of monitoring PSA overexpression or the metastasis of prostate cancers, the 24P4C12 polynucleotides described herein can be utilized in the same way to detect 24P4C12 overexpression or the metastasis of prostate and other cancers expressing this gene. Alternatively, just as PSA polypeptides are used to generate antibodies specific for PSA which can then be used to observe the presence and/or the level of PSA proteins in methods to monitor PSA protein overexpression (see, e.g., Stephan et al., *Urology* 55(4):560-3 (2000)) or the metastasis of prostate cells (see, e.g., Alanen et al., *Pathol. Res. Pract.* 192(3):233-7 (1996)), the 24P4C12 polypeptides described herein can be utilized to generate antibodies for use in detecting 24P4C12 overexpression or the metastasis of prostate cells and cells of other cancers expressing this gene.

Specifically, because metastases involves the movement of cancer cells from an organ of origin (such as the lung or prostate gland etc.) to a different area of the body (such as a lymph node), assays which examine a biological sample for the presence of cells expressing 24P4C12 polynucleotides

and/or polypeptides can be used to provide evidence of metastasis. For example, when a biological sample from tissue that does not normally contain 24P4C12-expressing cells (lymph node) is found to contain 24P4C12-expressing cells such as the 24P4C12 expression seen in LAPC4 and LAPC9, xenografts isolated from lymph node and bone metastasis, respectively, this finding is indicative of metastasis.

Alternatively 24P4C12 polynucleotides and/or polypeptides can be used to provide evidence of cancer, for example, when cells in a biological sample that do not normally express 24P4C12 or express 24P4C12 at a different level are found to express 24P4C12 or have an increased expression of 24P4C12 (see, e.g., the 24P4C12 expression in the cancers listed in Table I and in patient samples etc. shown in the accompanying Figures). In such assays, artisans may further wish to generate supplementary evidence of metastasis by testing the biological sample for the presence of a second tissue restricted marker (in addition to 24P4C12) such as PSA, PSCA etc. (see, e.g., Alanen et al., *Pathol. Res. Pract.* 192(3): 233-237 (1996)).

Just as PSA polynucleotide fragments and polynucleotide variants are employed by skilled artisans for use in methods of monitoring PSA, 24P4C12 polynucleotide fragments and polynucleotide variants are used in an analogous manner. In particular, typical PSA polynucleotides used in methods of monitoring PSA are probes or primers which consist of fragments of the PSA cDNA sequence. Illustrating this, primers used to PCR amplify a PSA polynucleotide must include less than the whole PSA sequence to function in the polymerase chain reaction. In the context of such PCR reactions, skilled artisans generally create a variety of different polynucleotide fragments that can be used as primers in order to amplify different portions of a polynucleotide of interest or to optimize amplification reactions (see, e.g., Caetano-Anolles, G. *Biotechniques* 25(3): 472-476, 478-480 (1998); Robertson et al., *Methods Mol. Biol.* 98:121-154 (1998)). An additional illustration of the use of such fragments is provided in the Example entitled "Expression analysis of 24P4C12 in normal tissues, and patient specimens," where a 24P4C12 polynucleotide fragment is used as a probe to show the expression of 24P4C12 RNAs in cancer cells. In addition, variant polynucleotide sequences are typically used as primers and probes for the corresponding mRNAs in PCR and Northern analyses (see, e.g., Sawai et al., *Fetal Diagn. Ther.* 1996 November-December 11(6):407-13 and *Current Protocols In Molecular Biology*, Volume 2, Unit 2, Frederick M. Ausubel et al. eds., 1995)). Polynucleotide fragments and variants are useful in this context where they are capable of binding to a target polynucleotide sequence (e.g., a 24P4C12 polynucleotide shown in FIG. 2 or variant thereof) under conditions of high stringency.

Furthermore, PSA polypeptides which contain an epitope that can be recognized by an antibody or T cell that specifically binds to that epitope are used in methods of monitoring PSA. 24P4C12 polypeptide fragments and polypeptide analogs or variants can also be used in an analogous manner. This practice of using polypeptide fragments or polypeptide variants to generate antibodies (such as anti-PSA antibodies or T cells) is typical in the art with a wide variety of systems such as fusion proteins being used by practitioners (see, e.g., *Current Protocols In Molecular Biology*, Volume 2, Unit 16, Frederick M. Ausubel et al. eds., 1995). In this context, each epitope(s) functions to provide the architecture with which an antibody or T cell is reactive. Typically, skilled artisans create a variety of different polypeptide fragments that can be used in order to generate immune responses specific for different portions of a polypeptide of interest (see, e.g., U.S. Pat. No.

5,840,501 and U.S. Pat. No. 5,939,533). For example it may be preferable to utilize a polypeptide comprising one of the 24P4C12 biological motifs discussed herein or a motif-bearing subsequence which is readily identified by one of skill in the art based on motifs available in the art. Polypeptide fragments, variants or analogs are typically useful in this context as long as they comprise an epitope capable of generating an antibody or T cell specific for a target polypeptide sequence (e.g. a 24P4C12 polypeptide shown in FIG. 3).

As shown herein, the 24P4C12 polynucleotides and polypeptides (as well as the 24P4C12 polynucleotide probes and anti-24P4C12 antibodies or T cells used to identify the presence of these molecules) exhibit specific properties that make them useful in diagnosing cancers such as those listed in Table I. Diagnostic assays that measure the presence of 24P4C12 gene products, in order to evaluate the presence or onset of a disease condition described herein, such as prostate cancer, are used to identify patients for preventive measures or further monitoring, as has been done so successfully with PSA. Moreover, these materials satisfy a need in the art for molecules having similar or complementary characteristics to PSA in situations where, for example, a definite diagnosis of metastasis of prostatic origin cannot be made on the basis of a test for PSA alone (see, e.g., Alanen et al., *Pathol. Res. Pract.* 192(3): 233-237 (1996)), and consequently, materials such as 24P4C12 polynucleotides and polypeptides (as well as the 24P4C12 polynucleotide probes and anti-24P4C12 antibodies used to identify the presence of these molecules) need to be employed to confirm a metastases of prostatic origin.

Finally, in addition to their use in diagnostic assays, the 24P4C12 polynucleotides disclosed herein have a number of other utilities such as their use in the identification of oncogenetic associated chromosomal abnormalities in the chromosomal region to which the 24P4C12 gene maps (see the Example entitled "Chromosomal Mapping of 24P4C12" below). Moreover, in addition to their use in diagnostic assays, the 24P4C12-related proteins and polynucleotides disclosed herein have other utilities such as their use in the forensic analysis of tissues of unknown origin (see, e.g., Takahama K *Forensic Sci Int* 1996 Jun. 28; 80(1-2): 63-9).

Additionally, 24P4C12-related proteins or polynucleotides of the invention can be used to treat a pathologic condition characterized by the over-expression of 24P4C12. For example, the amino acid or nucleic acid sequence of FIG. 2 or FIG. 3, or fragments of either, can be used to generate an immune response to a 24P4C12 antigen. Antibodies or other molecules that react with 24P4C12 can be used to modulate the function of this molecule, and thereby provide a therapeutic benefit.

XII.) INHIBITION OF 24P4C12 PROTEIN FUNCTION

The invention includes various methods and compositions for inhibiting the binding of 24P4C12 to its binding partner or its association with other protein(s) as well as methods for inhibiting 24P4C12 function.

XII.A.) Inhibition of 24P4C12 with Intracellular Antibodies

In one approach, a recombinant vector that encodes single chain antibodies that specifically bind to 24P4C12 are introduced into 24P4C12 expressing cells via gene transfer technologies. Accordingly, the encoded single chain anti-24P4C12 antibody is expressed intracellularly, binds to 24P4C12 protein, and thereby inhibits its function. Methods for engineering such intracellular single chain antibodies are well known. Such intracellular antibodies, also known as

"intrabodies", are specifically targeted to a particular compartment within the cell, providing control over where the inhibitory activity of the treatment is focused. This technology has been successfully applied in the art (for review, see Richardson and Marasco, 1995, *TIBTECH* vol. 13). Intrabodies have been shown to virtually eliminate the expression of otherwise abundant cell surface receptors (see, e.g., Richardson et al., 1995, *Proc. Natl. Acad. Sci. USA* 92: 3137-3141; Beerli et al., 1994, *J. Biol. Chem.* 269: 23931-23936; Deshane et al., 1994, *Gene Ther.* 1: 332-337).

Single chain antibodies comprise the variable domains of the heavy and light chain joined by a flexible linker polypeptide, and are expressed as a single polypeptide. Optionally, single chain antibodies are expressed as a single chain variable region fragment joined to the light chain constant region. Well-known intracellular trafficking signals are engineered into recombinant polynucleotide vectors encoding such single chain antibodies in order to target precisely the intrabody to the desired intracellular compartment. For example, intrabodies targeted to the endoplasmic reticulum (ER) are engineered to incorporate a leader peptide and, optionally, a C-terminal ER retention signal, such as the KDEL amino acid motif. Intrabodies intended to exert activity in the nucleus are engineered to include a nuclear localization signal. Lipid moieties are joined to intrabodies in order to tether the intrabody to the cytosolic side of the plasma membrane. Intrabodies can also be targeted to exert function in the cytosol. For example, cytosolic intrabodies are used to sequester factors within the cytosol, thereby preventing them from being transported to their natural cellular destination.

In one embodiment, intrabodies are used to capture 24P4C12 in the nucleus, thereby preventing its activity within the nucleus. Nuclear targeting signals are engineered into such 24P4C12 intrabodies in order to achieve the desired targeting. Such 24P4C12 intrabodies are designed to bind specifically to a particular 24P4C12 domain. In another embodiment, cytosolic intrabodies that specifically bind to a 24P4C12 protein are used to prevent 24P4C12 from gaining access to the nucleus, thereby preventing it from exerting any biological activity within the nucleus (e.g., preventing 24P4C12 from forming transcription complexes with other factors).

In order to specifically direct the expression of such intrabodies to particular cells, the transcription of the intrabody is placed under the regulatory control of an appropriate tumor-specific promoter and/or enhancer. In order to target intrabody expression specifically to prostate, for example, the PSA promoter and/or promoter/enhancer can be utilized (See, for example, U.S. Pat. No. 5,919,652 issued 6 Jul. 1999).

XII.B.) Inhibition of 24P4C12 with Recombinant Proteins

In another approach, recombinant molecules bind to 24P4C12 and thereby inhibit 24P4C12 function. For example, these recombinant molecules prevent or inhibit 24P4C12 from accessing/binding to its binding partner(s) or associating with other protein(s). Such recombinant molecules can, for example, contain the reactive part(s) of a 24P4C12 specific antibody molecule. In a particular embodiment, the 24P4C12 binding domain of a 24P4C12 binding partner is engineered into a dimeric fusion protein, whereby the fusion protein comprises two 24P4C12 ligand binding domains linked to the Fc portion of a human IgG, such as human IgG1. Such IgG portion can contain, for example, the C_H2 and C_H3 domains and the hinge region, but not the C_H1 domain. Such dimeric fusion proteins are administered in soluble form to patients suffering from a cancer associated with the expression of 24P4C12, whereby the dimeric fusion

专利名称(译)	肿瘤相关蛋白的抗体		
公开(公告)号	US8298776	公开(公告)日	2012-10-30
申请号	US12/829256	申请日	2010-07-01
[标]申请(专利权)人(译)	RAITANO ARTHUR乙 MORRISON KAREN JANE蛾 GE WANGMAO CHALLITA EID PIA中号 JAKOBOVITS AYA		
申请(专利权)人(译)	RAITANO ARTHUR乙 MORRISON KAREN JANE蛾 GE WANGMAO CHALLITA-EID PIA中号 JAKOBOVITS AYA		
当前申请(专利权)人(译)	AGENSYS INC.		
[标]发明人	RAITANO ARTHUR B MORRISON KAREN JANE MEYRICK GE WANGMAO CHALLITA EID PIA M JAKOBOVITS AYA		
发明人	RAITANO, ARTHUR B. MORRISON, KAREN JANE MEYRICK GE, WANGMAO CHALLITA-EID, PIA M. JAKOBOVITS, AYA		
IPC分类号	G01N33/53 A61K38/17 A61K48/00 C07K14/705 G01N33/574		
CPC分类号	A61K38/1709 C07K16/18 G01N33/57492 C07K14/705 A01K2217/05 A01K2267/03 A61K48/00 C12N2799/026 C12N2799/027		
代理机构(译)	美富律师事务所		
优先权	60/128858 1999-04-12 US		
其他公开文献	US20100273174A1		
外部链接	Espacenet USPTO		

摘要(译)

描述了新基因024P4C12 (也称为24P4C12) 及其编码的蛋白质及其变体, 其中24P4C12在正常成人组织中表现出组织特异性表达, 并且在表I中列出的癌症中异常表达。因此, 24P4C12提供诊断, 癌症的预后, 预防和/或治疗靶标。 24P4C12基因或其片段, 或其编码的蛋白质, 或其变体, 或其片段, 可用于引发体液或细胞免疫应答;与24P4C12反应的抗体或T细胞可用于主动或被动免疫。


```
Score = 1358 bits (706), Expect = 0.0Identites = 706/706 (100%) Strand = Plus/Plus
24P4C12v.1: 1 gagccatgggggaaagcagcggaacgagatgacagagccctacgggaagccagtcgaat 60
24P4C12v.7: 1 gagccatgggggaaagcagcggaacgagatgacagagccctacgggaagccagtcgaat 60
24P4C12v.1: 61 acgacccctcctttcgaggccccatcaagaacagaagctgcacagatgtcatctgtctgg 120
24P4C12v.7: 61 acgacccctcctttcgaggccccatcaagaacagaagctgcacagatgtcatctgtctgg 120
24P4C12v.1: 121 acgacccctcctttcgaggccccatcaagaacagaagctgcacagatgtcatctgtctgg 180
24P4C12v.7: 121 acgacccctcctttcgaggccccatcaagaacagaagctgcacagatgtcatctgtctgg 180
24P4C12v.1: 181 gagacccccggcaagtccttacccccaggaaactctactggggcctactgtggcatgggg 240
24P4C12v.7: 181 gagacccccggcaagtccttacccccaggaaactctactggggcctactgtggcatgggg 240
24P4C12v.1: 241 agaacaaagataagcgcgtatctctgtgacttcaacatcttcacagtgcatctctgtcagg 300
24P4C12v.7: 241 agaacaaagataagcgcgtatctctgtgacttcaacatcttcacagtgcatctctgtcagg 300
24P4C12v.1: 301 acatcatctcagttgtctgagaacggcctacagtgccccacacccccaggtgtgtgtctct 360
24P4C12v.7: 301 acatcatctcagttgtctgagaacggcctacagtgccccacacccccaggtgtgtgtctct 360
24P4C12v.1: 361 cctgccccggaggcccatggacgtgtgggaaaaaacgagttctcacagacgtgttggggag 420
24P4C12v.7: 361 cctgccccggaggcccatggacgtgtgggaaaaaacgagttctcacagacgtgttggggag 420
24P4C12v.1: 421 tctttatatacaaaaaacagggaacttttctgtgccaggggtaacctgggaatatgacgggtg 480
24P4C12v.7: 421 tctttatatacaaaaaacagggaacttttctgtgccaggggtaacctgggaatatgacgggtg 480
24P4C12v.1: 481 tcacaaagcctgcacaaaggaaactctgccccagtttctctctcccccctctgtctcagctgg 540
24P4C12v.7: 481 tcacaaagcctgcacaaaggaaactctgccccagtttctctctcccccctctgtctcagctgg 540
24P4C12v.1: 541 ggcgctgctttccatggacccaacgttactccaacggcgctcccaggggatcacccatgaaca 600
24P4C12v.7: 541 ggcgctgctttccatggacccaacgttactccaacggcgctcccaggggatcacccatgaaca 600
24P4C12v.1: 601 ccaccatacagccagggaatcagcggctcttattgacagcctcaatgcccgagacatcagtg 660
24P4C12v.7: 601 ccaccatacagccagggaatcagcggctcttattgacagcctcaatgcccgagacatcagtg 660
24P4C12v.1: 661 ttaagatctttgaaattttggccagctcctggtattggattctgtg 706
24P4C12v.7: 661 ttaagatctttgaaattttggccagctcctggtattggattctgtg 706
Score = 2971 bits (1545), Expect = 0.0Identites = 1545/1545 (100%) Strand = Plus/Plus
24P4C12v.1: 1043 ggctgtgggacagatgatgtgtacacatgtgtacacacatggctacatttctctctctct 1102
24P4C12v.7: 1043 ggctgtgggacagatgatgtgtacacatgtgtacacacatggctacatttctctctctct 1102
24P4C12v.1: 1103 catctgcatttgcctacttgggccaatgactgtctgtacctgggtacatcggggcaacccca 1162
24P4C12v.7: 1103 catctgcatttgcctacttgggccaatgactgtctgtacctgggtacatcggggcaacccca 1162
```