



US 20040072286A1

(19) **United States**

(12) **Patent Application Publication** (10) **Pub. No.: US 2004/0072286 A1**

Heus et al. (43) **Pub. Date: Apr. 15, 2004**

(54) **EXTRACELLULAR MATRIX PROTEIN**

(30) **Foreign Application Priority Data**

(76) Inventors: **Hendrik Cornelis Heus**, Rueil
Malmaison (FR); **Robert Louis Hubert**
Nelissen, Oss (NL); **Cornelis Maria**
Leonardus Meeuwisse, Eindhoven
(NL)

Jul. 13, 2000 (EP) 00202495.8

Publication Classification

(51) **Int. Cl.⁷** **A61K 38/17**; C12P 21/02;
C12N 5/06; C07K 14/47

(52) **U.S. Cl.** **435/69.1**; 435/320.1; 435/325;
514/12; 530/350; 536/23.5

Correspondence Address:
William M Blackstone
Akzo Nobel Intervet Inc
Patent Department
405 State Street
Millsboro, DE 19966 (US)

(57) **ABSTRACT**

(21) Appl. No.: **10/332,929**

(22) PCT Filed: **Jul. 9, 2001**

(86) PCT No.: **PCT/EP01/07888**

The invention describes a novel protein and variants thereof as well as the genes coding for these proteins. The protein is useful in the prevention of inflammatory diseases, more specifically in the induction of T cell tolerance to the protein in patients suffering from rheumatoid arthritis.

1 2 3 4 5 6 7 8 9 10




Figure 1




Figure 2

Encoded carboxy-terminal ends of SCIM-1 polypeptide
Variant A ↓ aa598 ...NEDGTACV GTLGQSPGPRPTTPTAAAATAAAAAA AGAATAAPVLVDGDLNLGSVVKESCEPSC (aa653)
Variant B ...NEDGTACV AQVAFLGGYSSAASRISEPLSRASYLSLGLGLCLQLYAL aa637)
Variant C ...NEDGTACV GWWSPVLKIVTPQVGKSLGP (aa618)
Variant D ...NEDGTACV ERTLLLGLCNLLGK (aa612)

Figure 3




Figure 4

EXTRACELLULAR MATRIX PROTEIN

[0001] The current invention relates to the isolation of a new polynucleotide molecule encoding a novel extracellular matrix protein, i.e. SCIM-1. The encoded protein or derivatives thereof, are useful for diagnosing, preventing or treating medical conditions. They can be used as modulatory agents for the treatment of autoimmune diseases, and more specifically rheumatoid arthritis.

[0002] The primary functional role of the immune system is to protect the individual against invading pathogens bearing foreign, that is non-self, antigens. In order to fulfil this function in a safe and effective manner, a mechanism is required to discriminate between foreign antigens and autoantigens derived from the individuals own body. Failure of this process of self-non-self discrimination, that is loss of immune tolerance to self-antigens, may lead to immune reactivity to autoantigens resulting in autoimmune disease, involving tissue damage and loss of organ function.

[0003] Autoimmune diseases are a major problem in human health care. Some autoimmune diseases may be the result of an immunological process directed at one antigen or antigenic complex whereas in others the autoimmune reaction may involve many types of antigens that may be present in multiple organs. Several lines of evidence have indicated that the immune system is involved in the pathology of autoimmune diseases. First, the chances of individuals to develop an autoimmune disease are closely linked to their genetic backgrounds: genes encoding major histocompatibility complex (MHC) class II molecules that present (auto)antigens to responding T cells which recognise MHC-peptide complexes show a strong genetic linkage to disease susceptibility. Second, cells of the immune system such as monocyte/macrophages and T cells infiltrate target organs. Third, T cells of patients with autoimmune diseases proliferate in vitro in response to potentially involved autoantigens. Fourth, studies in animal models of autoimmunity have unequivocally demonstrated that cells of the immune system such as monocyte/macrophages and T cells are involved in induction and expression of disease activity.

[0004] A disease as rheumatoid arthritis (RA) can illustrate the immunopathology that may occur in case of an autoimmune disease. RA presents itself as a chronic multi-system disease in which the common clinical manifestation is the persistent inflammatory synovitis accompanied by proliferation of synovial cells, pannus formation, cartilage degradation and bone erosion, and ultimately joint deformity resulting in loss of function.

[0005] Existing therapies for the treatment of autoimmune disorders, such as RA, in which the immune system generates an unwanted and undesirable inflammatory response, are inadequate. Treatment has focused on relief of symptoms of autoimmune disease rather than on its cause. Most drugs used in the treatment of autoimmune diseases, e.g. steroids and non-steroidal anti-inflammatory compounds, are non-specific and have significant toxic side effects. This is especially problematic since autoimmune diseases are chronic conditions, which require the prolonged administration of drugs.

[0006] Antigen-driven, non-toxic immunomodulation therapy provides a very attractive alternative for the non-specific immunosuppression. This antigen-specific therapy

involves the treatment of patients with the target (auto)antigen or with synthetic T cell-reactive peptides derived from the (auto)antigen. These synthetic peptides correspond to T cell epitopes of the (auto)antigen and can be used to induce specific T cell tolerance both to themselves and to the (auto)antigen. The controlled administration of the target (auto)antigen can be very effective in desensitisation of the immune system. Desensitisation or immune tolerance of the immune system is based on the long-observed phenomenon that animals which have been fed or have inhaled an antigen or epitope are less capable of developing a systemic immune response towards said antigen or epitope when said antigen or epitope is introduced via a systemic route.

[0007] With regard to the use of an antigen for immune therapy, the human cartilage (IC) gp-39 protein has very recently been reported to be effective in induction of systemic immune tolerance (Joosten et al. *Arthritis Rheum.* 43:645-655, 2000) for the treatment of arthritis. Upon intranasal administration HC gp-39 interfered with development of collagen type II-induced arthritis in mice. Both disease activity and joint destruction ameliorated as a result of the treatment.

[0008] In a search for proteins that are relevant for autoimmune diseases such as rheumatoid arthritis, we have identified a novel gene SCIM-1 (Synovium/Cartilage Inflammation-linked Messenger 1). The protein that is encoded by the SCIM-1 mRNA has not been reported earlier. Upon identification of the SCIM-1 mRNA it was established that only fragments of the mRNA were present in public domain databases as expressed sequence tags (ESTs). From the public ESTs it was not possible to reconstruct the identified SCIM-1 cDNA.

[0009] The best homology with the SCIM-1 protein was found with the EGF-like domains of fibulin-1, an extracellular protein containing 9 EGF-like domains. Fibulin-1 can bridge the extracellular matrix of the vessel wall with integrin- α Ib β 3 on platelets, via an interaction with fibrinogen (Godyna et al. *Blood* 88:2569-2577,1996). Via its EGF-like domains fibulin-1 binds to nidogen, fibronectin, laminin, fibrinogen, NOVH, aggrecan, and versican (Tran et al. *J. Biol. Chem.* 272:22600-22606, 1997; Perbal et al. *Proc. Natl. Acad. Sci. USA* 96:869-874, 1999; Barth et al. *Matrix Biol.* 17:635-646, 1998). High expression of fibulin-1 seems to interfere with tumor formation and invasion (Qing et al. *Oncogene* 15:2159-2168, 1997). It is conceivable that SCIM-1 has similar capabilities, involving its RGD-motif en EGF-like domain.

[0010] Based on the report of Sarkissian and Lafyatis (*J. Immunol.* 162:1772-1779, 1999), the putative interaction of SCIM-1 with integrins could be relevant for RA, since proliferation of fibroblast-like synoviocytes (FLS), e.g. from RA patients, required a signal provided by integrin-binding extracellular matrix proteins in addition to a mitogenic cytokine, e.g. PDGF. Wang et al. (*Arthritis Rheum.* 40:1298-1307, 1997) suggested that α v, α 4, and α 5 integrins are required for IL-1 β -stimulated invasion of articular cartilage by RA-FLS.

[0011] The SCIM-1 gene appeared to be expressed relatively high in the synovial and cartilage tissues. Based on a partial cDNA sequence, a full-length cDNA was isolated from human chondrocytes. Probing multiple tissue northern blots, it was shown that the SCIM-1 gene was expressed in

primary chondrocytes from healthy human donors, while some expression was also observed in brain and lung, although the SCIM-1 mRNAs in these tissues seemed to differ in length. Only low or no expression was observed in other human tissues such as heart, placenta, liver, skeletal muscle, spleen, kidney and colon or cell lines of monocytes, arterial endothelial cells, and (cultured) synovial fibroblasts.

[0012] SCIM-1 expression appears to be positively correlated to a differentiated chondrocyte phenotype, since expression was only detected in primary chondrocytes but not in human primary chondrocytes transformed with SV40. Furthermore, it was shown that expression of the SCIM-1 gene in human chondrocytes was enhanced in the presence of the RA-related cytokines TNF α and IL β . In chondrocytes, transcription of the SCIM-1 gene results in a 2.7 kb mRNA which encodes a protein of 653 amino acids. Based on the multiple tissue Northern blot data and the sequences of Incyte and public domain databases, It was found that the SCIM-1 transcript undergoes a different splicing in tissues of the joint, brain, and lungs.

[0013] Based on these predicted protein motifs the SCIM-1 protein is expected to be an extracellular matrix protein or a protein that is localized on the cell surface of chondrocytes and synoviocytes and possibly some other cell types.

[0014] The main problem in (auto)immune diseases (such as e.g. RA) is that the precise targets or antigens that the immune system is adversely reacting to are largely unknown, implicating that modulating a disease entity in an antigen-specific fashion may not be possible.

[0015] It would be an important advantage, however, if an antigen-driven, non-toxic form of immunomodulation therapy could be utilised without knowledge of the antigen(s) that are involved as a target in the (auto)immune response. Such an antigen-driven therapy would involve the generation of antigen-specific modulator cells with the use of an antigen that is expected to be released or produced during the autoimmune process. Such an antigen would become available during inflammation or tissue destruction. In case of an autoimmune disease, the locally produced autoantigen should then activate or reactivate modulator cells induced with such an antigen.

[0016] To effectively use tolerance induction therapy to treat T cell mediated cartilage destruction, there is a great need to identify T cell-reactive (poly)peptides which can desensitise patients against the autoantigen that is activating the T cells responsible for the inflammatory process.

[0017] It is an object of the invention to provide a (poly)peptide which is capable of inducing systemic immune tolerance, more in particular specific T cell tolerance, preferably to the responsible cartilage antigen in patients suffering from T cell-mediated cartilage destruction. It has been found that SCIM-1 fulfils the above mentioned requirements and can be used as an effective toleragen in modulating the immune system.

[0018] In the present invention under induction of systemic immune tolerance is to be understood the stimulation of antigen specific lymphocytes by antigen presenting cells (APC) in such a way that the lymphocytes acquire a state in which they produce anti-inflammatory cytokines. Anti-inflammatory cytokines may for example be IL-4, IL-10,

and/or TGF- β . Lymphocytes brought to tolerance by APC are able to impose their anti-inflammatory state to other sites of the body, e.g. sites of ongoing inflammation.

[0019] The immune system protects individuals against foreign antigens and responds to exposure to a foreign antigen by activating specific cells such as T- and B-lymphocytes and producing soluble factors like interleukins, antibodies and complement factors. The antigen to which the immune system responds is degraded by the antigen presenting cells (APCs) and a fragment of the antigen is expressed on the cell surface associated with a major histocompatibility complex (NC) class II glycoprotein. The MHC-glycoprotein-antigen-fragment complex is presented to a T cell, which by virtue of its T cell receptor recognises the antigen fragment conjointly with the MHC class II protein to which it is bound. The T cell becomes activated, i.e. proliferates and/or produces interleukins, resulting in the expansion of the activated lymphocytes directed to the antigen under attack (Grey et al., *Sci. Am.*, 261:38-46, 1989).

[0020] Self-antigens are also continuously processed and presented as antigen fragments by the MHC glycoproteins to T cells (Jardetsky et al., *Nature* 353:326-329, 1991). Self recognition thus is intrinsic to the immune system. Under normal circumstances the immune system is tolerant to self-antigens and activation of the immune response by these self-antigens is avoided. When tolerance to self-antigens is lost, the immune system becomes activated against one or more self-antigens, resulting in the activation of autoreactive T cells and sometimes also the production of autoantibodies. This phenomenon is referred to as autoimmunity. As the immune response in general is destructive, i.e. meant to destroy the invasive foreign antigen, autoimmune responses can cause destruction of the body's own tissue.

[0021] It will thus be clear that fragments of the SCIM-1 protein will be expressed by the APC and that therefore also fragments of the SCIM-1 protein are capable of evoking an immune response. Also proteins of other species having a similar function or at least being structurally closely related to the human SCIM-1 protein might perform the same toleragenic effect. Thus, also homologous polypeptides or parts thereof are included in the invention.

[0022] The proteins according to the present invention include the polypeptide comprising SEQ ID NO: 2 but also polypeptides with a similarity of 70%, preferably 90%, more preferably 95%, 98%, most preferably 99%. Also portions of such polypeptides still capable of conferring the toleragenic effects are included. Such portions may be functional per se, e.g. in solubilized form or they might be linked to other polypeptides, either by known biotechnological ways or by chemical synthesis, to obtain chimeric proteins.

[0023] As used herein the term similarity is as defined in NCBI-BLAST 2.0.10 [Aug. 26, 1999] (Altschul et al., *Nucleic Acids Res.* 25:3389-3402, 1997).

[0024] The fragments of the SCIM-1 protein or homologous polypeptides are to be understood subsequences of the protein. These subsequences can modulate lymphocyte functioning. Preferably they have the following functional immunomodulating characteristics: i) peptides can be bound by the disease-associated MHC molecules, preferably HLA-DRB1*0101, DRB1*0401, DRB1*0404, DRB1*0408,

DRB1*0405, DQB*0301, or DQB*0302, and ii) peptides must be able to provoke a T cell response in humans, preferably autoimmune patients, more preferably RA patients. Such a response can for example be measured in an *in vitro* T cell proliferation assay or in an assay for the detection of T cell cytokine production (e.g. ELISA or ELISPOT) (Coligan et al., Current Protocols in Immunology. John Wiley & Sons, Inc., 1998). Preferably the peptides must also be recognized by T cells in animals transgenic for the relevant human MHC class II molecules, as mentioned above, and human CD4 upon immunization with a SCIM-1 (poly)peptide.

[0025] The length of these sub-sequences is not important provided that it comprises the epitope to be recognized by the relevant MHC molecule. Preferably these peptides have an amino acid sequence of 9-55 amino acid residues. More preferably the peptides as have an amino acid sequence of 9-35, in particular 9-25 amino acid residues. Much more preferred are peptides having an amino acid sequence of 9-15 amino acid residues. Highly preferred are peptides having an amino acid sequence of 13 or 14 amino acid residues.

[0026] Variations that can occur in a sequence, especially of smaller peptides, may be demonstrated by (an) amino acid difference(s) in the overall sequence or by deletions, substitutions, insertions, inversions or additions of (an) amino acid(s) in said sequence. Amino acid substitutions that are expected not to essentially alter biological and immunological activities have been described. Amino acid replacements between related amino acids or replacements which have occurred frequently in evolution are, *inter alia* Ser/Ala, Ser/Gly, Asp/Gly, Asp/Asn, Ile/Val (see Dayhof, M. D., Atlas of protein sequence and structure, Nat. Biomed. Res. Found., Washington D.C., 1978, vol. 5, suppl. 3). Based on this information Lipman and Pearson developed a method for rapid and sensitive protein comparison (Science, 227:1435-1441, 1985) and determining the functional similarity between homologous polypeptides.

[0027] Also within the scope of the invention are multimers of the peptides such as for example a dimer or trimer of the peptides according to the invention. A multimer according to the invention can either be a homomer, consisting of a multitude of the same peptide, or a heteromer consisting of different peptides.

[0028] It will be clear to those skilled in the art that the (poly)peptides may be extended at either side of the peptide or at both sides and still exert the same immunological function. The extended part may be an amino acid sequence similar to the natural sequence of the protein. However, the (poly)peptide might also be extended by non-natural sequences. It will be clear that the (poly)peptide need not to exert its original function and as such might be inactive while still performing its immunological function according to the invention. The (poly)peptide according to the invention might be connected to MHC molecules, such that the binding groove is occupied by the peptide. A flexible linker molecule, preferably also consisting of amino acid sequences might connect the peptide. The MHC molecules need not to possess their constant domains and might consist of their variable domains only, either directly connected to each other or connected through a flexible linker. The advantage of such a complex is that it might exist in a soluble form and can directly be recognised by T cells.

[0029] Thus, according to the invention the (poly)peptides, said (poly)peptides resembling the MHC Class II restricted T-cell epitopes present on the antigen comprising the polypeptide of SEQ ID NO: 2 or fragments thereof comprising these epitopes are very suitable for use in a therapy to induce systemic immune tolerance to said antigen in mammals, more specifically humans, suffering from lymphocyte or T-cell mediated cartilage destruction, such as for example arthritis, more specifically rheumatoid arthritis. Optionally such a treatment can be combined with the administration of other medicaments such as DMARDs (Disease Modifying Anti-Rheumatic Drugs e.g. sulfasalazine, anti-malarials (chloroquine, hydroxychloroquine) injectable or oral gold, methotrexate, D-penicillamine, azathioprine, cyclosporine, mycophenolate), NSAIDs (non steroidal anti inflammatory drugs), corticosteroids or other drugs known to influence the course of the disease in autoimmune patients.

[0030] The polypeptides according to the invention can also be used to modulate lymphocytes that are reactive to antigens other than said antigen but are present in the same tissue as the antigen i.e. proteins or parts thereof comprising the polypeptide according to SEQ ID NO: 2. By the induction of antigen-specific T-cell tolerance, autoimmune disorders can be treated by systemic immune tolerance. More in general, the cells to be modulated are hematopoietic cells. In general, in order to function as a toleragen the peptide must fulfil at least two conditions i.e. it must possess an immune modulating capacity and it must be expressed locally usually as part of a larger protein.

[0031] Thus, the present invention provides a method to treat patients suffering from inflammatory autoimmune diseases, by administration of a pharmaceutical preparation comprising the (poly)peptide according to the invention. The (poly)peptide comprises T-cell epitopes, which are recognised by and are able to stimulate autoreactive T-cells.

[0032] These T cells may be found e.g. in the blood of patients suffering from inflammatory disorders. Such patients may suffer from diseases like Graves' diseases, juvenile arthritis, primary glomerulonephritis, polyarthritis, osteoarthritis, Sjögren's syndrome, myasthenia gravis, rheumatoid arthritis, Addison's disease, primary biliary sclerosis, uveitis, systemic lupus erythematosus, inflammatory bowel disease, multiple sclerosis or diabetes. The polypeptides according to the present invention therefore can be used in the preparation of a pharmaceutical to prevent inflammatory diseases. Administration of SCIM-1 will induce systemic immune tolerance. More specifically the polypeptides can be used in the preparation of a pharmaceutical to induce specific immune tolerance in patients suffering from inflammatory diseases, preferably immune-cell mediated cartilage destruction. The immune cell, preferably is a T cell. The most preferred disease is arthritis, more preferably rheumatoid arthritis.

[0033] Treatment of autoimmune disorders with the peptides according to the invention makes use of the fact that systemic immune tolerance is induced to unrelated but co-localised antigens. The regulatory cells secrete in an antigen specific fashion pleiotropic proteins such as cytokines which may downmodulate the immune response.

[0034] The polypeptides according to the invention can be prepared by recombinant DNA techniques.

[0035] The present invention provides for such a DNA sequence encoding a protein or polypeptide according to the invention.

[0036] The invention also includes the entire mRNA sequence part of which is indicated in SEQ ID NO: 1. A complete coding DNA sequence is shown in SEQ ID NO: 1 nucleotides 59-2017. Furthermore, to accommodate codon variability, the invention also includes sequences coding for the same amino acid sequences as the sequences disclosed herein. Also portions of the coding sequences coding for individual polypeptides having the same immunological function are part of the invention as well as allelic and species variations thereof. Sometimes, a gene is expressed in a certain tissue as a splicing variant, resulting in an altered 5' or 3' mRNA or the inclusion of an additional exon sequence. These sequences as well as the proteins encoded by these sequences all are expected to perform the same or similar functions and form also part of the invention.

[0037] In particular, SEQ ID NO: 16, SEQ ID NO: 17 and SEQ ID NO: 18 represent specific splice variants which differ from SEQ ID NO: 1 in the sequence downstream of nucleotide 1852 of SEQ ID NO: 1. Translation of this splice variants leads to a truncated version of the protein in SEQ ID NO: 2, as shown in SEQ ID NO: 19, SEQ ID NO: 20 and SEQ ID NO: 21, respectively. The coding sequences of these latter nucleic acids run from positions 59-1969, 59-1912 and 59-1894, respectively and have the first 598 amino acids of SEQ ID NO: 2 in common. The complete proteins preferably are mature proteins. The signal sequence most likely is 19 amino acids but the length may differ slightly.

[0038] The sequence information as provided herein should not be so narrowly construed as to require inclusion of erroneously identified bases. The specific sequence disclosed herein can be readily used to isolate the complete genes which in turn can easily be subjected to further sequence analyses thereby identifying sequencing errors.

[0039] Thus, the present invention provides also for isolated polynucleotides encoding SCIM-1, truncated versions or fragments thereof.

[0040] The DNA according to the invention may be obtained from cDNA. The tissues preferably are from human origin. Preferably ribonucleic acids are isolated from fetal brain, fetal liver, fetal spleen, placenta or other tissues. Alternatively, the coding sequence might be genomic DNA, or prepared using DNA synthesis techniques. The polynucleotide may also be in the form of RNA. If the polynucleotide is DNA, it may be in single stranded or double stranded form. The single strand might be the coding strand or the non-coding (anti-sense) strand.

[0041] The present invention further relates to polynucleotides having slight variations or have polymorphic sites. Polynucleotides having slight variations encode polypeptides which retain the same biological function or activity as the natural, mature protein. Polymorphic sites are useful for diagnostic purposes. Such polynucleotides can be identified by hybridization under preferably highly stringent conditions. According to the present invention the term "stringent" means washing conditions of 1×SSC, 0.1% SDS at a temperature of 65° C.; highly stringent conditions refer to a reduction in SSC towards 0.3×SSC, more preferably to 0.1×SSC. Preferably the first two washings are subsequently

carried out twice each during 15-30 minutes. If there is a need to wash under highly stringent conditions an additional wash with 0.1×SSC is performed once during 15 minutes. Hybridization can be performed e.g. overnight in 0.5M phosphate buffer pH7.5/7% SDS at 65° C.

[0042] Alternatively, also fragments of the above mentioned polynucleotides which code for polypeptides having the same immunological function are embodied in the invention.

[0043] Also the encoded proteins or polypeptides derived from the proteins form part of the invention.

[0044] A nucleic acid sequence coding for the protein, a peptide according to the invention, a multimer of said peptides or a chimeric peptide is inserted into an expression vector. Suitable expression vectors comprise the necessary control regions for replication and expression. The expression vector can be brought to expression in a host cell. Suitable host cells are, for instance, bacteria, yeast cells and mammalian cells. Such techniques are well known in the art, see for instance Sambrook et al., *Molecular Cloning: a Laboratory Manual*, Cold Spring Harbor laboratory Press, Cold Spring Harbor, 1989.

[0045] The (smaller) (poly)peptides according to the invention can also be prepared by well known organic chemical methods for peptide synthesis such as, for example, solid-phase peptide synthesis described for instance in *J. Amer. Chem. Soc.* 85:2149 (1963) and *Int. J. Peptide Protein Res.* 35:161-214 (1990).

[0046] The (poly) peptides may be stabilised by C- and/or N-terminal modifications, which will decrease exopeptidase catalysed hydrolysis. The modifications may include: C-terminal acylation, (e.g. acylation=Ac-peptide), N-terminal amide introduction, (e.g. peptide-NH₂) combinations of acylation and amide introduction (e.g. Ac-peptide-NH₂) and introduction of D-amino acids instead of L-amino acids (Powell et al., *J. Pharm. Sci.*, 81:731-735, 1992).

[0047] Other modifications are focussed on the prevention of hydrolysis by endopeptidases. Examples of these modifications are: introduction of D-amino acids instead of L-amino acids, modified amino acids, cyclisation within the peptide, introduction of modified peptide bonds, e.g. reduced peptide bonds $\psi[\text{CH}_2\text{NH}]$ and e.g. peptoids (N-alkylated glycine derivatives) (Adang et al., *Recl. Trav. Chim. Pays-Bas*, 113:63-78, 1994 and Simon et al., *Proc. Natl. Acad. Sci. USA*, 89:9367-9371, 1992).

[0048] The tolerogenic peptides according to the invention can be identified by using a method comprising the steps of

[0049] a) introducing into a suitable host cell a DNA fragment encoding SCIM-1 peptide fragments;

[0050] b) culturing the host cells under conditions to allow expression of the introduced DNA sequence;

[0051] c) bringing the expression product in contact with lymphocytes; and

[0052] d) establishing the lymphocyte activity.

[0053] Preferably, the expression product is contacted with lymphocytes in vivo e.g. by administration of the product to animals. The lymphocyte activity can e.g. be measured by determination of the anti-inflammatory cytokines.

[0054] Alternatively, the expression product of the host cells under step b might be isolated, brought subsequently into contact with antigen presenting cells and T cells and the T cell activity might be established.

[0055] It will be clear that in the above described screening methods the peptides to be tested might also be synthesized chemically.

[0056] The peptides thus identified can be used for the formulation of a pharmaceutical composition comprising mixing the peptide with a pharmaceutically acceptable carrier.

[0057] According to the invention, patients suffering from T-cell mediated destruction of the articular cartilage can be treated with a therapeutical composition comprising one or more peptides according to the invention and a pharmaceutical acceptable carrier. Administration of the pharmaceutical composition according to the invention will induce systemic immune tolerance, in particular tolerance of the specific autoreactive T cells of these patients, to the autoantigenic proteins in the articular cartilage under attack and other self antigens which display the identified MHC Class II binding T cell epitopes characterised or mimicked by the amino acid sequences of one or more of the peptides according to the invention. The induced tolerance thus will lead to a reduction of the local inflammatory response in the articular cartilage under attack.

[0058] The (poly)peptides according to the invention have the advantage that they have a specific effect on the autoreactive T cells thus leaving the other components of the immune system intact as compared to the non-specific suppressive effect of immunosuppressive drugs.

[0059] Systemic immune tolerance can be attained by administering high or low doses of peptides according to the invention. The amount of peptide will depend on the route of administration, the time of administration, the age of the patient as well as general health conditions and diet.

[0060] In general, a dosage of 0.01 to 10000 μg of peptide per kg body weight, preferably 0.05 to 500 μg , more preferably 0.1 to 100 μg of peptide can be used.

[0061] Pharmaceutical acceptable carriers are well known to those skilled in the art and include, for example, sterile saline, lactose, sucrose, calcium phosphate, gelatin, dextrin, agar, pectin, peanut oil, olive oil, sesame oil and water. Other carriers may be, for example MHC class II molecules, if desired embedded in liposomes.

[0062] In addition the pharmaceutical composition according to the invention may comprise one or more adjuvants. Suitable adjuvants include, amongst others, aluminium hydroxide, aluminium phosphate, amphigen, tocopherols, monophosphoryl lipid A, muramyl dipeptide and saponins such as Quill A. Preferably, the adjuvants to be used in the tolerance therapy according to the invention are mucosal adjuvants such as the cholera toxin B-subunit or carbomers, which bind to the mucosal epithelium. The amount of adjuvant depends on the nature of the adjuvant itself.

[0063] Furthermore the pharmaceutical composition according to the invention may comprise one or more stabilisers such as, for example, carbohydrates including

sorbitol, mannitol, starch, sucrodedextrin and glucose, proteins such as albumin or casein, and buffers like alkaline phosphates.

[0064] Suitable administration routes are e.g. intramuscular injections, subcutaneous injections, intravenous injections or intraperitoneal injections, oral administration and nasal administration such as sprays.

[0065] For testing the ability of the (poly)peptides to modulate (auto)immune responses several murine models have been shown to be suitable, such as collagen induced arthritis in mice (CIA), adjuvant arthritis in rats, experimental allergic encephalomyelitis in mice and non-obese diabetes in mice (NOD) or in transgenic mice. Antigen may be administered intravenously, intraperitoneally, orally or nasally in such models (review by Liblau et al., Immunol. Today 18:599-603, 1997). To facilitate the read-out in these models, it is of importance to increase the confidence interval. According to the present invention it has been found that incidence and clinical score in arthritis models can be improved by combining the original trigger of arthritis, e.g. collagen type II in CIA with a peptide derived from the extracellular matrix protein aggrecan. This peptide might preferably be administered simultaneously with the original trigger although a separate administration might also be possible.

[0066] SCIM-1 or the peptides according to the invention are also very suitable for use in a diagnostic method to detect the presence of activated autoreactive T cells involved in the chronic inflammation of the articular cartilage.

[0067] The diagnostic method according to the invention comprises the following steps:

[0068] a) isolation of the peripheral blood mononuclear cells (PBMC) from a blood sample of an individual,

[0069] b) culture said PBMC under suitable conditions,

[0070] c) incubation of said PBMC culture in the presence of the autoantigen or one or more peptides derived thereof according to the invention, and

[0071] d) detection of a response of T cells, for example a proliferative response, indicating the presence of activated autoreactive T cells in the individual.

[0072] In case of detection of a response by measuring the proliferative response of the autoreactive T cells, the incorporation of a radioisotope such as for example ^3H -thymidine is a measure for the proliferation. A response of the autoreactive T cells present in the PBMC can also be detected by measuring the cytokine release with cytokine-specific ELISA, or the cytotoxicity with ^{51}Cr Chromium release. Another detection method is the measurement of expression of activation markers by FACS analysis, for example of II-2R. A diagnostic composition comprising one or more of the peptides according to the invention and a suitable detecting agent thus forms part of the invention. Depending on the type of detection, the detection agent can be a radioisotope, an enzyme, or antibodies specific for cell surface or activation markers.

[0073] Also within the scope of the invention are test kits which comprise one or more peptides according to the

invention. These test kits are suitable for use in a diagnostic method according to the invention.

[0074] Thus, the present invention provides for a method to detect whether autoaggressive T cells reactive towards SCIM-1 are present in patients suffering from T-cell mediated cartilage destruction such as for example arthritis, in particular rheumatoid arthritis. If SCIM-1-specific T cells are present, tolerization of these T cells with a pharmaceutical composition comprising SCIM-1 or peptides according to the present invention or combinations thereof can delay or suppress arthritis development.

[0075] The following examples are illustrative for the invention and should in no way be interpreted as limiting the scope of the invention.

LEGENDS TO THE FIGURES

[0076] FIG. 1

[0077] RT-PCR on human cartilage cDNA, using oligonucleotides specific for SCIM-1 and GAPDH. Detection of SCIM-1 gene expression is shown for healthy cartilage (knee, male; traffic accident; lane 1), and cartilage from 4 RA patients (knee/hip, 3 female and 1 male; lanes 2-5). Detection of GAPDH gene expression for the same 4 RA samples is shown in lanes 7-10. As a DNA fragment length marker the 100 bp-ladder (Gibco-BRL) was run in parallel in lane 6 (the major band in the middle of the lane corresponds to 600 bp).

[0078] FIG. 2

[0079] SCIM-1 probe (nt560-1048 of SEQ ID NO: 1) hybridized on custom-made joint Northern blot (upper panel). Represented human cells/tissues: U937 monocytes+PMA-ionomycin (lane 1), HAEC endothelial cells±TNF α (lanes 2-3, respectively), SCRO14.SF primary synovial fibroblasts+TNF α (lane 4), PCG SV40.04 transformed chondrocytes±TNF α (lanes 5-6, respectively), and primary chondrocytes±TNF α (lanes 7-8, respectively). Per lane 1-2 μ g of poly(A+) RNA was loaded. As a control the blot was probed with a ³²P-labelled cDNA derived from the human β -actin gene (lower panel).

[0080] FIG. 3

[0081] Alternative carboxy terminal ends as encoded by 4 different SCIM-1 mRNA species, i.e. variants A-D. The encoded C-termini are indicated and amino acids are shown in one-letter code. Sequences are different as from amino acid 599. Encoded amino acid sequences of variants A-D are represented by SEQ ID NO: 20-23, respectively, and the corresponding nucleic acids are represented by SEQ ID NO: 16-19, respectively.

[0082] FIG. 4

[0083] Detection of SCIM-1 variant A/His6 expression in transfected CHO cells. Samples of culture supernatant of CHO cells transfected with pNGVI (lane 1) or pNGV1.SCIM1.variantA(His6) (lanes 2 and 3) were separated on an SDS-polyacrylamide gel and subsequently Western blotted. Proteins containing a (His6)-tail were detected using an anti(His6)-tag monoclonal antibody (Dianova GMBH). A protein molecular weight marker was run in lane 4 (M).

EXAMPLES

Example 1

[0084] Identification of the SCIM-1 Gene

[0085] The cDNA libraries that were available from the LifeSeq database (Incyte Pharmaceuticals) were clustered based on their gene content in order to define custom tissue categories that represent RA relevant tissues such as joint tissues. Distances between cDNA libraries were calculated using the squared Euclidian distance measurement, and the results were clustered into a tree using an unweighted average linkage method (Fry 1993, Biological Data Analysis, Oxford University Press, New York). Two of the resulting clusters contained cDNAs of 5 libraries each derived from non-tumor synovial or cartilage tissues. For each of the synovial cDNA libraries data of 4038 to 5627 sequence reactions were available, and 1554 to 7231 for each of cartilage cDNA libraries. These sequence data were probed for genes of which expression seemed to be relatively restricted to these clusters. Two partial cDNA fragments, each composed of overlapping Incyte-est sequences and parts of SEQ ID NO: 1, were identified as positive hits. Since their expression patterns appeared to be similar, it was assumed that they were derived from the same gene.

Example 2

[0086] SCIM-1 Gene expression in RA Cartilage.

[0087] Expression of the SCIM-1 gene in diseased tissue was detected via RT-PCR with SCIM-1-specific oligonucleotides (SEQ ID NO: 3, 5'-TTGCCAATTACGCCTACGGT and SEQ ID NO: 4, 5'-CCTGGTCATTGTCAAAGTCGG) on cDNA that was derived from cartilage samples of 4 RA patients and 1 healthy donor. The arthritic cartilage was obtained during joint replacement surgery of the knee. Chondrocytes were isolated enzymatically from the cartilage (Cornelissen et al., 1993, J. Tiss. Cult. Meth. 15:139-146) upon which RNA was isolated using Trizol (Gibco-BRL) or RNAzol B (Campro Scientific). With 1 μ g of total RNA the synthesis of cDNA was performed using SuperscriptTM II (Gibco-BRL) in a total volume of 20 μ l. For RT-PCR on SCIM-1 and on housekeeping gene GAPDH, as positive control, 0.5 μ l cDNA per reaction was used. PCR was performed in a Perkin Elmer 9600: 1 cycle 5 min 94° C., 35 cycles 30 sec 94° C./30 sec 55° C./1 min 72° C., 1 cycle 5 min 72° C. with 50 ng/primer, 200 μ M dNTPs, and 2.5 u Taq polymerase (Pharmacia, #27-0799) in 25 μ l total volume. Oligonucleotides specific for GAPDH were SEQ ID NO: 5 (5'-CCCTTCATTGACCTCAACTACATGG) and SEQ ID NO: 6 (5'-GGTCCACCACCTGTTGCTGTAGCC). PCR samples were analysed on agarose gel (FIG. 1). Lanes 1-5 show clear signals of SCIM-1 cDNA amplification product of the expected length for the healthy cartilage (macroscopically) and 4/4 of the arthritis patients, while GAPDH-specific amplification signals are in the same order of magnitude among the RA cDNA preparations (lanes 7-10). The RT-PCR data indicate that the SCIM-1 gene is expressed in diseased tissue, i.e. afflicted knee cartilage, of 4/4 RA patients tested. It is likely that the SCIM-1 gene indeed is expressed in diseased articular cartilage of at least a considerable percentage of RA patients. Consequently, it is to be expected that the SCIM-1 protein is synthesised in diseased cartilage of RA patients.

Example 3

[0088] SCIM-1 Gene Expression in Joint-Related Cells

[0089] To determine tissue distribution of SCIM-1 gene expression multiple tissue Northern blots were probed with SCIM-1 cDNA fragments. Probes were generated via random primed labeling of SCIM-1 cDNA fragments nt(560-1048) and (1983-2427) (numbering according to SEQ ID NO: 1) using Ready-To-Go beads (Pharmacia) and is ³²P- α -dCTP (Amersham). Labeled probe was separated from free nucleotides on a 1 ml Sephadex G50-medium column and about 3 \times 10⁶ cpm/ml of labeled probe was added to the hybridization mix (0.5 M phosphate buffer pH 7.0, 7% SDS, 1 mM EDTA) and hybridized to the Northern blot for 16 h at -65° C. Blots were washed up to 0.5 \times SSC at 65° C. and exposed to a STORMTM840 Phosphor screen. Probe nt 560-1048 relative to SEQ ID NO: 1 hybridized weakly to a -2.4 kb mRNA in brain and a -3.0 kb mRNA in lung (Clontech human multiple tissue Northern blots H2 and H1, cat #7759-1 and 7760-1, respectively). Except for a clear signal at -2.6 kb in primary chondrocytes (**FIG. 2**, upper panel lanes 7-8) no signals were detected in other joint-related, cultured cells. Upon correction for the β -actin hybridization signal, as housekeeping gene control (**FIG. 2**, lower panel), the signal for SCIM-1 mRNA appeared to be enhanced with a factor 3-4 by TNF α (24 h, 10 ng/ml). With another SCIM-1 probe, i.e. nt(1983-2427), no signals were detected on Clontech's multiple tissue Northern blots H3 (cat #7767-1) and H2. Hybridization with probe nt(1983-2427) on RNA of primary chondrocytes yielded a signal for the same 2.6 kb mRNA as with probe nt(560-1048) (not shown), including the 3-4 fold upregulation of SCIM-1 gene expression by TNF α . Hybridization of a cDNA microarray with probes derived of chondrocytes incubated with or without IL-1, showed a 3-fold IL-1-dependent upregulation of SCIM-1 expression. Also for probe nt(1983-2427) no signals were observed on Northern blot for SV40-transformed human chondrocytes, which are likely to represent dedifferentiated cells. It was concluded that i) SCIM-1 gene expression displays a restricted tissue distribution, ii) the different lengths of detected mRNAs in brain, lung and chondrocytes could represent splice variants, and iii) SCIM-1 gene expression is enhanced by pro-inflammatory factors such as TNF and IL-1.

Example 4

[0090] Isolation of the Full Length SCIM-1 cDNA

[0091] The complete coding sequence of human SCIM-1 was identified via SMART-RACE (#K1811-1, Clontech), using SCIM-1-specific oligonucleotides on RACE-cDNA that, was generated from 1 μ g total RNA of TNF α -stimulated primary human chondrocytes. The chondrocytes were enzymatically (collagenase) isolated from healthy cartilage (knee), grown for about 3 weeks in monolayer culture (Hamm's/F12, 10% FCS) and subsequently stimulated with 10 ng/ml TNF α for 24 h. Oligonucleotides were designed based on sequences that were available for the two gene fragments identified from the Incyte database (Example 1). For the first gene fragment primers were: 5'RACE primer SEQ ID NO: 7 (5'GGGTCCATTGTACCCCGCCACGACG), and nested primer SEQ ID NO: 8 (5'CTCAAAGTCCCCATCATGGTCC), 3'RACE primer SEQ ID NO: 9 (5'CTCAGCCGCTGTCCGTCTTCCGG),

and nested primer SEQ ID NO: 10 (5'GCTTCAACAA-CAACTGGCTGCG). For the second gene fragment primers were: 5'RACE primer SEQ ID NO: 11 (5'GGATGGGCT-TGGGGAGGGTCTAGCTC), and nested primer SEQ ID NO: 12 (5'GCAGCAGCACAAGCCCACCTTTC), 3'RACE primer SEQ ID NO: 13 (5'GTGCCAGGGAGGTGGTGT-CACTG), and nested primer SEQ ID NO: 14 (5'GCACAG-GAAGTATGAGGACTTTAGTG). SMART-RACE PCRs were performed according to the Clontech manual PT3269-1 (March 1999). Sequencing of a number of RACE cDNA clones yielded the contig cDNA of SEQ ID NO: 1, which indicated that the two gene fragments as identified in Example 1, corresponded to the same gene, designated SCIM-1. A translation start codon was found at nt 59 and an open reading frame was found to encode 653 aa. With regard to the observed mRNA length of 2.6 kb in chondrocytes, the identified SCIM-1 cDNA of 2589 bp was considered very likely to contain the complete coding region of the corresponding gene.

[0092] Once the complete SCIM-1 cDNA had become available, a comparison was made with nucleic acid and protein sequences in Incyte and public domain databases, via blastsearches (NCBI-BLAST 2.0.10 Aug. 26, 1999; Altschul et al., *Nucleic Acids Res.* 25:3389-3402, 1997). Several public est sequences and a ~1100 bp cDNA from patent application WO99/58660 were found to be identical to parts of the SCIM-1 cDNA. As a result of incomplete overlap, the complete SCIM-1 gene could not be reconstructed from these public cDNA fragments. Based on our complete encoded SCIM-1 amino acid sequence (SEQ ID NO: 2) a prediction was made as to functional protein domains of SCIM-1. A signal sequence was identified at aa 1-19 or 1-21, an integrin-binding RGD motif at aa 263-265, a calcium-binding EGF-like domain at aa 551-598, a putative hydrophobic region at aa 614-635, and MHC class II DR4Dw4 binding motifs at aa 12-20, 39-47, 149-157, and 323-331.

Example 5

[0093] SCIM-1 Splice Variants and their Tissue Distribution

[0094] From multiple alignments (DNAMAN version 4.11) involving the full length SCIM-1 cDNA and all Incyte and public cDNAs that contained sequences identical to the SCIM-1 cDNA, it was discovered that the population of cDNAs was heterogeneous downstream of nucleotide 1852 (numbering according to SEQ ID NO: 1). This downstream region encodes for the C-terminal end of the protein, directly following the EGF-like domain. Four different C-terminal regions were identified for SCIM-1, i.e. variants A-D (Table 1; SEQ ID NO: 19-21, respectively, the C-terminal differences starting at amino acid position 599), deduced from cDNAs that were derived from different tissues (SEQ ID NO: 16-18, respectively; the 3'end differs as from nucleotide position 1853) and are expected to be the result of alternative splicing events. For the variants A, B, and C it is most likely that SEQ ID NO: 16-18 comprises the 3' ends of the corresponding mRNAs, respectively, since each contains a 3' poly(A) tail preceded by a putative poly-adenylation signal AATAAAA. The different lengths of the 3'UTR regions for variants A and B are in agreement with the lengths of the SCIM-1 mRNAs that were detected on Northern blots for cartilage and brain tissues, respectively (see example 3). The

number of cDNA clones that were found in the various databases to encode each of the variants A-D, strongly suggests that SCIM-1 variant A is predominantly expressed in joint tissues (cartilage/synovium) and to some extent in tumours and some other tissues, whereas expression of variant B seems to be restricted to tissues of the central nervous system (mostly brain). The suggested tissue-restricted expression was corroborated by an RT-PCR with oligonucleotides specific for the SCIM-1 variant B (SEQ ID NO: 9 and 15) on cDNA of total brain (Clontech human brain Quick-clone cDNA, cat. 7187-1/lot 9070843) and on cDNA of primary chondrocytes (cultured for 24 h with TNF- α , see example 4). Synthesis of cDNA and PCR conditions are as described in example 2. As a control PCR was performed with oligonucleotides specific for house-keeping gene GAPDH (SEQ ID NO: 5 and 6). A strong amplification signal of the expected length for the SCIM-1 variant B mRNA in brain tissue was seen whereas expression in chondrocytes hardly could be detected. The control PCR on GAPDH gene expression yielded a similar signal for both tissues, indicating that the input amounts of the corresponding cDNAs in the PCR were comparable. The finding that expression of variant B, represented by a 2.4 kb mRNA, was hardly detectable in the primary chondrocytes (24 h/TNF- α) is in agreement with the data of examples 3 and 4. On Northern blot only a single SCIM-1 mRNA variant of 2.6 kb was detected in primary chondrocytes (24 h/TNF- α) (FIG. 2, lane 8) which was concluded to represent variant A as 3' RACE amplification on chondrocyte cDNA yielded only variant A sequences (see SEQ ID NO: 1).

Example 6

[0095] Expression of Recombinant SCIM-1

[0096] The SCIM-1 (variant A) cDNA was cloned into cloning vector pCR2.1TOPO (Invitrogen) and subcloned as a Eco RI fragment to eukaryotic expression vector pNGV1 (EMBL accession number X99274). Consequently, the cDNA is situated behind the SV40 early promoter and a Kozak translation initiation sequence. Upstream of the translational stopcodon 18 nucleotides were inserted encoding a His6-tag. In order to produce recombinant SCIM-1(His6) protein in eukaryotic cells, CHO cells (ATCC CCL61) were cultured in DMEM/Hamm's F12 containing 5% FCS (Harlan sera lab). The pNGV1-SCIM-1(His6) construct was transfected to CHO-K1 cells using Transfectam (Promega) and selection medium DMEAMamm's F12 containing 5% FCS and 0.8 mg/ml neomycin (G418 sulphate Gibco BRL Life technology, filter sterilised using a 0.22 μ M Millipore SLGVO25BS filter). To detect recombinant protein, cells were cultured o/n in serum-free medium (containing 0.8 mg/ml neomycin) and expression was allowed to continue for one day. Serum-free culture supernatant was analysed on SDS-PAGE followed by Western blotting and subsequent detection with anti-His6 monoclonal antibody. The blot was blocked with 5% skimmed milk in P13S/0.05% Tween-20 and developed with a mouse-anti(His6)-tag antibody (Dianova GMBH cat. no. Dia 900. After three times washing with PBST for five minutes at RT, the blot was incubated with anti-mouse-IgG-HRP (Promega catno 3624512). After three times washing with PBST for five minutes at RT, detection was performed using a chromogenic substrate for HRP, i.e. diaminobezidine tetrahydrochloride in PBS, 2.5 MM CoCl₂ and 0.006% H₂O₂).

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 21

<210> SEQ ID NO 1

<211> LENGTH: 2589

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<220> FEATURE:

<221> NAME/KEY: misc_feature

<222> LOCATION: (2498)..(2498)

<223> OTHER INFORMATION: n is a, c, g, or t

<400> SEQUENCE: 1

```

aagcagtggg aacaacgcag agtacgogg ggtataaagc gagcgcgctg accccggcat    60
gtccaggatg ttaccgttcc tgctgctgct ctggtttctg cccatcaactg aggggtccca    120
gcgggctgaa cccatgttca ctgcagtcac caactcagtt ctgcctcctg actatgacag    180
taatcccacc cagctcaact atgggtgtggc agttactgat gtggaccatg atggggactt    240
tgagatcgtc gtggcggggg acaatggacc caacctgggt ctgaagtatg accggggcca    300
gaagcggctg gtgaacatcg cggtcgatga gcgcagctca ccctactacg cgctgcggga    360
ccggcagggg aacgcocatg gggtcacagc ctgcgacatc gacggggacg gccgggagga    420
gatctacttc ctcaacacca ataatgocct ctcggggggtg gccacgtaca ccgacaagtt    480
gttcaagttc cgcaataacc ggtgggaaga catcctgagc gatgaggtca acgtggcccc    540

```

-continued

tgggtgtggc agcctctttg ccggacgctc tgtggcctgt gtggacagaa agggctctgg	600
acgctactct atctacattg ccaattacgc ctacggtaat gtgggcccctg atgccctcat	660
tgaaatggac cctgaggcca gtgacctctc ccggggcatt ctggcgctca gagatgtggc	720
tgctgaggct ggggtcagca aatatacagg gggccgaggc gtcagcgtgg gccccatcct	780
cagcagcagt gcctcggata tctctctgca caatgagaat gggcctaact tccttttcca	840
caaccggggc gatggcacct ttgtggacgc tgcggccagt gctggtgtgg acgaccccca	900
ccagcatggg cgagggtgctg ccctggctga cttcaaccgt gatggcaaag tggacatcgt	960
ctatggcaac tggaaatggcc cccaccgctc ctatctgcaa atgagcacc atgggaaggt	1020
ccgcttccgg gacatcgctt cacccaagtt ctccatgccc tcccctgtcc gcacggtcat	1080
caccgccgac ttgacaatg accaggagct ggagatcttc ttcaacaaca ttgcctaccg	1140
cagctcctca gccaaaccgc tcttccgctg catccgtaga gacacggag accccctcat	1200
cgaggagctc aatcccggcg acgccttggg gcctgagggc cggggcacag ggggtgtggt	1260
gaccgacttc gacggagacg ggatgctgga cctcatcttg tcccatggag agtccatggc	1320
tcagccgctg tccgtcttc ggggcaatca gggcttcaac aacaactggc tgcgagtggg	1380
gccacgcacc cggtttgggg cctttgccag gggagctaag gtctgtctct acaccaagaa	1440
gagtggggcc caactgagga tcatcgacgg gggctcaggc tacctgtgtg agatggagcc	1500
cgtggcacac tttggcctgg ggaaggatga agccagcagt gtggaggatga cgtggccaga	1560
tggcaagatg gtgagccgga acgtggccag cggggagatg aactcagtgc tggagatcct	1620
ctacccccg gatgaggaca cacttcagga cccagccca ctggagtgtg gccaaaggatt	1680
ctcccagcag gaaaatggcc attgcatgga caccaatgaa tgcattccagt tcccattcgt	1740
gtgcctcga gacaagccc tatgtgtcaa cacctatgga agctacaggg gccggacca	1800
caagaagtgc agtcggggct acgagcccaa cgaggatggc acagcctgctg tggggactct	1860
cggccagtca ccgggcccc gccccaccac ccccaccgct gctgctgcca ctgccgctgc	1920
tgctgccgct gctggagctg ccaactgtgc accggtcctc gtagatggag atctcaatct	1980
ggggtcgggt gttaaaggaga gctgcgagcc cagctgctga gcaggggtgg gacatgaacc	2040
agcggatgga gtccagcagg ggagtgggaa agtgggcttg tgctgctgcc tagacagtag	2100
ggatgtaaa gcctgggagc tagaccctcc ccaagcccat ccatgcacat tacttagcta	2160
acaattaggg agactcgtaa ggccaggccc tgtgctggc acatagctgt gatcacagca	2220
gacaggtctg ctgccctgat ggcgcttaca ttccagtggg tetaatgacc atatcttagg	2280
acacagatgt gccagggag gtggtgtcac tgcacaggaa gtatgaggac tttagtgtcc	2340
tgagttcaaa tcctgattca ggaactcaca aagctatgtg accttacacc agtcaactaa	2400
cttgtagcc atccattatc gcatctgcaa aatggggatt aagaatagaa tcttgggggt	2460
agtgtggaga ttagattaaa tgtatgtaag acacttanca caaacctgg cacatagtaa	2520
aggctcaata aaaacaagt cctctcactg ggctttgtca acacgtcaaa aaaaaaaaa	2580
aaaaaaaa	2589

<210> SEQ ID NO 2
 <211> LENGTH: 653
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

-continued

<400> SEQUENCE: 2

Met Ser Arg Met Leu Pro Phe Leu Leu Leu Leu Trp Phe Leu Pro Ile
1 5 10 15
Thr Glu Gly Ser Gln Arg Ala Glu Pro Met Phe Thr Ala Val Thr Asn
20 25 30
Ser Val Leu Pro Pro Asp Tyr Asp Ser Asn Pro Thr Gln Leu Asn Tyr
35 40 45
Gly Val Ala Val Thr Asp Val Asp His Asp Gly Asp Phe Glu Ile Val
50 55 60
Val Ala Gly Tyr Asn Gly Pro Asn Leu Val Leu Lys Tyr Asp Arg Ala
65 70 75 80
Gln Lys Arg Leu Val Asn Ile Ala Val Asp Glu Arg Ser Ser Pro Tyr
85 90 95
Tyr Ala Leu Arg Asp Arg Gln Gly Asn Ala Ile Gly Val Thr Ala Cys
100 105 110
Asp Ile Asp Gly Asp Gly Arg Glu Glu Ile Tyr Phe Leu Asn Thr Asn
115 120 125
Asn Ala Phe Ser Gly Val Ala Thr Tyr Thr Asp Lys Leu Phe Lys Phe
130 135 140
Arg Asn Asn Arg Trp Glu Asp Ile Leu Ser Asp Glu Val Asn Val Ala
145 150 155 160
Arg Gly Val Ala Ser Leu Phe Ala Gly Arg Ser Val Ala Cys Val Asp
165 170 175
Arg Lys Gly Ser Gly Arg Tyr Ser Ile Tyr Ile Ala Asn Tyr Ala Tyr
180 185 190
Gly Asn Val Gly Pro Asp Ala Leu Ile Glu Met Asp Pro Glu Ala Ser
195 200 205
Asp Leu Ser Arg Gly Ile Leu Ala Leu Arg Asp Val Ala Ala Glu Ala
210 215 220
Gly Val Ser Lys Tyr Thr Gly Gly Arg Gly Val Ser Val Gly Pro Ile
225 230 235 240
Leu Ser Ser Ser Ala Ser Asp Ile Phe Cys Asp Asn Glu Asn Gly Pro
245 250 255
Asn Phe Leu Phe His Asn Arg Gly Asp Gly Thr Phe Val Asp Ala Ala
260 265 270
Ala Ser Ala Gly Val Asp Asp Pro His Gln His Gly Arg Gly Val Ala
275 280 285
Leu Ala Asp Phe Asn Arg Asp Gly Lys Val Asp Ile Val Tyr Gly Asn
290 295 300
Trp Asn Gly Pro His Arg Leu Tyr Leu Gln Met Ser Thr His Gly Lys
305 310 315 320
Val Arg Phe Arg Asp Ile Ala Ser Pro Lys Phe Ser Met Pro Ser Pro
325 330 335
Val Arg Thr Val Ile Thr Ala Asp Phe Asp Asn Asp Gln Glu Leu Glu
340 345 350
Ile Phe Phe Asn Asn Ile Ala Tyr Arg Ser Ser Ser Ala Asn Arg Leu
355 360 365
Phe Arg Val Ile Arg Arg Glu His Gly Asp Pro Leu Ile Glu Glu Leu
370 375 380
Asn Pro Gly Asp Ala Leu Glu Pro Glu Gly Arg Gly Thr Gly Gly Val
385 390 395 400

-continued

Val Thr Asp Phe Asp Gly Asp Gly Met Leu Asp Leu Ile Leu Ser His
 405 410 415

Gly Glu Ser Met Ala Gln Pro Leu Ser Val Phe Arg Gly Asn Gln Gly
 420 425 430

Phe Asn Asn Asn Trp Leu Arg Val Val Pro Arg Thr Arg Phe Gly Ala
 435 440 445

Phe Ala Arg Gly Ala Lys Val Val Leu Tyr Thr Lys Lys Ser Gly Ala
 450 455 460

His Leu Arg Ile Ile Asp Gly Gly Ser Gly Tyr Leu Cys Glu Met Glu
 465 470 475 480

Pro Val Ala His Phe Gly Leu Gly Lys Asp Glu Ala Ser Ser Val Glu
 485 490 495

Val Thr Trp Pro Asp Gly Lys Met Val Ser Arg Asn Val Ala Ser Gly
 500 505 510

Glu Met Asn Ser Val Leu Glu Ile Leu Tyr Pro Arg Asp Glu Asp Thr
 515 520 525

Leu Gln Asp Pro Ala Pro Leu Glu Cys Gly Gln Gly Phe Ser Gln Gln
 530 535 540

Glu Asn Gly His Cys Met Asp Thr Asn Glu Cys Ile Gln Phe Pro Phe
 545 550 555 560

Val Cys Pro Arg Asp Lys Pro Val Cys Val Asn Thr Tyr Gly Ser Tyr
 565 570 575

Arg Cys Arg Thr Asn Lys Lys Cys Ser Arg Gly Tyr Glu Pro Asn Glu
 580 585 590

Asp Gly Thr Ala Cys Val Gly Thr Leu Gly Gln Ser Pro Gly Pro Arg
 595 600 605

Pro Thr Thr Pro Thr Ala Ala Ala Thr Ala Ala Ala Ala Ala Ala
 610 615 620

Ala Gly Ala Ala Thr Ala Ala Pro Val Leu Val Asp Gly Asp Leu Asn
 625 630 635 640

Leu Gly Ser Val Val Lys Glu Ser Cys Glu Pro Ser Cys
 645 650

<210> SEQ ID NO 3
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 3

ttgccaatta cgctacggt

20

<210> SEQ ID NO 4
 <211> LENGTH: 21
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 4

cctggtcatt gtcaaagtcg g

21

<210> SEQ ID NO 5
 <211> LENGTH: 25
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 5

-continued

cccttcattg acctcaacta catgg 25

<210> SEQ ID NO 6
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 6

ggtccaccac cctggtgctg tagcc 25

<210> SEQ ID NO 7
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 7

gggtccattg taccocgcca cgacg 25

<210> SEQ ID NO 8
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 8

ctcaaagtcc ccatcatggt cc 22

<210> SEQ ID NO 9
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 9

ctcagccgct gtccgtcttc cgg 23

<210> SEQ ID NO 10
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 10

gcttcaacaa caactggctg cg 22

<210> SEQ ID NO 11
<211> LENGTH: 26
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 11

ggatgggctt ggggagggtc tagctc 26

<210> SEQ ID NO 12
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 12

gcagcagcac aagcccactt tc 22

<210> SEQ ID NO 13
<211> LENGTH: 24

-continued

<212> TYPE: DNA
 <213> ORGANISM: Homo sapiens
 <400> SEQUENCE: 13
 gtgcccaggg aggtggtgct actg 24

<210> SEQ ID NO 14
 <211> LENGTH: 26
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens
 <400> SEQUENCE: 14
 gcacaggaag tatgaggact ttagtg 26

<210> SEQ ID NO 15
 <211> LENGTH: 27
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens
 <400> SEQUENCE: 15
 gggatttaa gtgcatataa ctgaagg 27

<210> SEQ ID NO 16
 <211> LENGTH: 2039
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens
 <400> SEQUENCE: 16
 aagcagtgtt aacaacgcag agtacgcggg ggtataaagc gagcgcgctg accccggcat 60
 gtccaggatg ttaccgttcc tgctgctgct ctggtttctg cccatcactg aggggtccca 120
 gcgggctgaa cccatgttca ctgcagtcac caactcagtt ctgcctoctg actatgacag 180
 taatcccacc cagctcaact atggtgtggc agttactgat gtggaccatg atggggactt 240
 tgagatcgtc gtggcggggt acaatggacc caacctggtt ctgaaagtatg accgggcca 300
 gaagcggctg gtgaacatcg cggtcgatga gcgcagctca ccctactacg cgtgcgggga 360
 ccggcagggg aacgccatcg gggtcacagc ctgcgacatc gacggggacg gccgggagga 420
 gatctacttc ctcaacacca ataatgcctt ctccgggggtg gccacgtaca ccgacaagtt 480
 gttcaagtcc cgcaataacc ggtgggaaga catcctgagc gatgaggcca acgtggcccg 540
 tgggtgtggc agcctctttg ccggacgctc tgtggcctgt gtggacagaa agggctcttg 600
 acgctactct atctacattg ccaattacgc ctacggtaat gtgggcctctg atgocctcat 660
 tgaaatggac cctgaggcca gtgacctctc ccggggcatt ctggcgctca gagatgtggc 720
 tgctgaggct ggggtacgca aatatacagg gggccgagga gtcagcgtgg gccccatcct 780
 cagcagcagt gcctcggata tcttctgcga caatgagaat gggcctaact tccttttcca 840
 caaccggggc gatggcacct ttgtggacgc tgcggccagt gctggtgtgg acgaccccca 900
 ccagcatggg cgaggtgtcg ccttggtgta cttcaaccgt gatggcaaag tggacatcgt 960
 ctatggcaac tggaatggcc cccaccgctt ctatctgcaa atgagcacc atgggaaggt 1020
 ccgcttccgg gacatcgcct caccgaagtt ctccatgccc tcccctgtcc gcaaggatc 1080
 caccgccgac ttgacaatg accaggagct ggagatcttc ttcaacaaca ttgocctacc 1140
 cagctcctca gccaacgcc tcttccgcgt catccgtaga gagcacggag accccctcat 1200
 cgaggagctc aatcccggcg acgccttggg gcctgagggc cggggcacag ggggtgtggt 1260

-continued

```

gaccgacttc gacggagacg ggatgctgga cctcatcttg tcccatggag agtccatggc 1320
tcagccgctg tccgtcttcc ggggcaatca gggcttcaac aacaactggc tgcgagtggc 1380
gccacgcacc cggtttgggg cctttgccag gggagctaag gtogtgctct acaccaagaa 1440
gagtggggcc cacctgagga tcacgacgg gggctcaggc tacctgtgtg agatggagcc 1500
cgtggcacac tttggcctgg ggaaggatga agccagcagt gtggaggtga cgtggccaga 1560
tggcaagatg gtgagccgga acgtggccag cggggagatg aactcagtgc tggagatcct 1620
ctacccccgg gatgaggaca cacttcagga cccagcccca ctggagtgtg gccaaaggatt 1680
ctcccagcag gaaaatggcc attgcatgga caccaatgaa tgcacccagt tcccattcgt 1740
gtgcctctga gacaagccc tatgtgtcaa cacctatgga agctacaggt gccggaccaa 1800
caagaagtgc agtcggggct acgagcccaa cgaggatggc acagcctgcg tggctcaagt 1860
ggccttttta ggtgggtatt cttcagccgc ctctagaato tctgagcctc tctctcgggc 1920
ctcatatctt tctctagccc ttggactttg ccttcagtta tatgcacttt aaatcccatc 1980
aataaaggaa aaaacaaaac aaaactaaca gcctttgtgg aaaactaaaa aaaaaaaaaa 2039

```

<210> SEQ ID NO 17

<211> LENGTH: 2089

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 17

```

aagcagtggc aacaacgcag agtacgcggg ggtataaagc gagcgcgctg accccggcat 60
gtccaggatg ttaccgttcc tgctgctgct ctggtttctg cccatcactg aggggtccca 120
gctggctgaa cccatgttca ctgcagtcac caactcagtt ctgcctcctg actatgacag 180
taatcccacc cagctcaact atggtgtggc agttactgat gtggaccatg atggggactt 240
tgagatcgtc gtggcggggc acaatggacc caacctggtt ctgaaagtatg accgggcca 300
gaagcggctg gtgaacatcg cggtcgatga gcgcagctca ccctactacg cgtgctggga 360
ccggcagggg aacgccatcg gggtcacagc ctgcgacatc gacggggacg gccgggagga 420
gatctacttc ctcaacacca ataatgcctt ctccgggggtg gccacgtaca ccgacaagtt 480
gttcaagttc cgcaataacc ggtgggaaga catcctgagc gatgaggcca acgtggccc 540
tgggtgtggc agcctctttg ccggacgctc tgtggcctgt gtggacagaa agggctctgg 600
acgctactct atctacattg ccaattacgc ctacggtaat gtgggcctctg atgocctcat 660
tgaaatggac cctgaggcca gtgacctctc ccggggcatt ctggcgctca gagatgtggc 720
tgctgaggct ggggtacgca aatatacagg gggccgaggg gtcagcgtgg gccccatcct 780
cagcagcagt gcctcggata tcttctgcga caatgagaat gggcctaact tccttttcca 840
caaccggggc gatggcacct ttgtggacgc tgcggccagt gctggtgtgg acgaccccca 900
ccagcatggg cgaggtgtcg ccttggtgta cttcaaccgt gatggcaaag tggacatcgt 960
ctatggcaac tggaatggcc cccaccgctt ctatctgcaa atgagcacc atgggaaggt 1020
ccgcttccgg gacatcgcct cacccaagtt ctccatgccc tcccctgtcc gcaaggatc 1080
caccgccgac ttgacaatg accaggagct ggagatcttc ttcaacaaca ttgocctacc 1140
cagctcctca gccaacccgc tcttccgcgt catccgtaga gagcacggag accccctcat 1200
cgaggagctc aatcccggcg acgccttggg gcctgagggc cggggcacag ggggtgtgg 1260

```

-continued

```

gaccgacttc gacggagacg ggatgctgga cctcatcttg tcccatggag agtccatggc 1320
tcagccgctg tccgtcttcc ggggcaatca gggcttcaac aacaactggc tgcgagtggc 1380
gccacgcacc cggtttgggg cctttgccag gggagctaag gtogtgctct acaccaagaa 1440
gagtggggcc cacctgagga tcacgacgg gggctcaggc tacctgtgtg agatggagcc 1500
cgtggcacac tttggcctgg ggaaggatga agccagcagt gtggaggtga cgtggccaga 1560
tggcaagatg gtgagccgga acgtggccag cggggagatg aactcagtgc tggagatcct 1620
ctacccccgg gatgaggaca cacttcagga cccagcccca ctggagtgtg gccaaaggatt 1680
ctcccagcag gaaaatggcc attgcatgga caccaatgaa tgcacccagt tcccattcgt 1740
gtgccctcga gacaagccc tatgtgtcaa cacctatgga agctacaggt gccggaccaa 1800
caagaagtgc agtcggggct acgagcccaa cgaggatggc acagcctgcg tgggctggtg 1860
gagccctgtg ttgaagatag tgacaccaca agttgggaag agccttggtc cctgaatcac 1920
tgaatcactg ccttgaatca ccgctggaa tacctgttga tcaggaacac ttacctggaa 1980
cttactgag caggatacaa acttctattg tattaagcta ttaatacatt aagatttggg 2040
ggtgtacct tacataataa attcccattt cctcttgaia aaaaaaaaa 2089

```

<210> SEQ ID NO 18

<211> LENGTH: 2145

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 18

```

aagcagtggc aacaacgcag agtacgcggg ggtataaagc gagcgcgctg accccggcat 60
gtccaggatg ttaccgttcc tgctgctgct ctggtttctg cccatcactg aggggtccca 120
gctggctgaa cccatgttca ctgcagtcac caactcagtt ctgcctcctg actatgacag 180
taatcccacc cagctcaact atggtgtggc agttactgat gtggaccatg atggggactt 240
tgagatgctc gtggcggggc acaatggacc caacctggtt ctgaagtatg accgggcccc 300
gaagcggctg gtgaacatcg cggtcgatga ggcagctca ccctactacg cgtgctggga 360
ccggcagggg aacgccatcg ggtcaccagc ctgacgacac gacggggacg gccgggagga 420
gatctacttc ctcaacacca ataatgcctt ctgctgggtg gccacgtaca ccgacaagtt 480
gttcaagttc cgcaataacc ggtgggaaga catcctgagc gatgaggtca acgtggcccc 540
tgggtgtggc agcctctttg ccggacgctc tgtggcctgt gtggacagaa agggctctgg 600
acgctactct atctacattg ccaattacgc ctacggtaat gtgggcccctg atgccctcat 660
tgaaatggac cctgaggcca gtgacctctc ccggggcatt ctggcctca gagatgtggc 720
tgctgaggct ggggtcagca aatatacagg gggccgaggc gtcagcgtgg gccccatcct 780
cagcagcagt gcctcggata tcttctgcga caatgagaat gggcctaact tccttttcca 840
caaccggggc gatggaccct ttgtggacgc tgcggccagt gctggtgtgg acgaccccca 900
ccagcatggg cgaggtgtgc ccctggctga cttcaaccgt gatggcaaag tggacatcgt 960
ctatggcaac tggaatggcc cccaccgcct ctatctgcaa atgagcacc atgggaaggt 1020
ccgcttccgg gacatcgcct caccgaagtt ctccatgccc tcccctgtcc gcacggtcat 1080
caccgccgac tttgacaatg accaggagct ggagatcttc ttcaacaaca ttgcctaccg 1140
cagctcctca gccaacgcgc tcttccgctg catccgtaga gagcacggag accccctcat 1200

```

-continued

```

cgaggagctc aatccccggcg acgccttgga gcttgagggc cggggcacag ggggtgtggt 1260
gaccgacttc gacggagacg ggatgctgga cctcatcttg tcccatggag agtccatggc 1320
tcagccgctg tccgtcttcc ggggcaatca gggcttcaac aacaactggc tgcgagtggg 1380
gccacgcacc cggtttgggg cctttgccag gggagctaag gtcgtgctct acaccaagaa 1440
gagtggggcc cacctgagga tcacgcacgg gggctcaggc taactgtgtg agatggagcc 1500
cgtggcacac ttggcctgg ggaaggatga agccagcagt gtggaggatga cgtggccaga 1560
tggcaagatg gtgagccgga acgtggccag cggggagatg aactcagtgc tggagatcct 1620
ctacccccgg gatgaggaca cacttcagga cccagcccca ctggagtgtg gccaaggatt 1680
ctcccagcag gaaaatggcc attgcatgga caccaatgaa tgcattcagt tcccattcgt 1740
gtgcctcga gacaagcccg tatgtgtcaa cacctatgga agctacaggt gccggaccaa 1800
caagaagtgc agtcggggct acgagcccaa cgaggatggc acagcctgcg tggaaagAAC 1860
cttgctactt ggtctctgca atctccttgg aaaatagaat gctcctgtag ccagaacact 1920
ctttgaaat gatgtaacgc cacagaatat gctcagcacc agcctcagta gtgaaggctt 1980
gttataaaac atagatctat tcttcatggg gaaaaaaagt catcccagga tataataaaa 2040
ttacacttaa gaatgcagca acccttttaa aaacatattc aacaaaaaaa agcccctgcc 2100
gttttctgat atcatagtaa tactcctaataaatcacaaa atatg 2145

```

<210> SEQ ID NO 19

<211> LENGTH: 637

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 19

```

Met Ser Arg Met Leu Pro Phe Leu Leu Leu Leu Trp Phe Leu Pro Ile
1           5           10           15
Thr Glu Gly Ser Gln Arg Ala Glu Pro Met Phe Thr Ala Val Thr Asn
20           25           30
Ser Val Leu Pro Pro Asp Tyr Asp Ser Asn Pro Thr Gln Leu Asn Tyr
35           40           45
Gly Val Ala Val Thr Asp Val Asp His Asp Gly Asp Phe Glu Ile Val
50           55           60
Val Ala Gly Tyr Asn Gly Pro Asn Leu Val Leu Lys Tyr Asp Arg Ala
65           70           75           80
Gln Lys Arg Leu Val Asn Ile Ala Val Asp Glu Arg Ser Ser Pro Tyr
85           90           95
Tyr Ala Leu Arg Asp Arg Gln Gly Asn Ala Ile Gly Val Thr Ala Cys
100          105          110
Asp Ile Asp Gly Asp Gly Arg Glu Glu Ile Tyr Phe Leu Asn Thr Asn
115          120          125
Asn Ala Phe Ser Gly Val Ala Thr Tyr Thr Asp Lys Leu Phe Lys Phe
130          135          140
Arg Asn Asn Arg Trp Glu Asp Ile Leu Ser Asp Glu Val Asn Val Ala
145          150          155          160
Arg Gly Val Ala Ser Leu Phe Ala Gly Arg Ser Val Ala Cys Val Asp
165          170          175
Arg Lys Gly Ser Gly Arg Tyr Ser Ile Tyr Ile Ala Asn Tyr Ala Tyr
180          185          190

```

-continued

Gly Asn Val Gly Pro Asp Ala Leu Ile Glu Met Asp Pro Glu Ala Ser
 195 200 205
 Asp Leu Ser Arg Gly Ile Leu Ala Leu Arg Asp Val Ala Ala Glu Ala
 210 215 220
 Gly Val Ser Lys Tyr Thr Gly Gly Arg Gly Val Ser Val Gly Pro Ile
 225 230 235 240
 Leu Ser Ser Ser Ala Ser Asp Ile Phe Cys Asp Asn Glu Asn Gly Pro
 245 250 255
 Asn Phe Leu Phe His Asn Arg Gly Asp Gly Thr Phe Val Asp Ala Ala
 260 265 270
 Ala Ser Ala Gly Val Asp Asp Pro His Gln His Gly Arg Gly Val Ala
 275 280 285
 Leu Ala Asp Phe Asn Arg Asp Gly Lys Val Asp Ile Val Tyr Gly Asn
 290 295 300
 Trp Asn Gly Pro His Arg Leu Tyr Leu Gln Met Ser Thr His Gly Lys
 305 310 315 320
 Val Arg Phe Arg Asp Ile Ala Ser Pro Lys Phe Ser Met Pro Ser Pro
 325 330 335
 Val Arg Thr Val Ile Thr Ala Asp Phe Asp Asn Asp Gln Glu Leu Glu
 340 345 350
 Ile Phe Phe Asn Asn Ile Ala Tyr Arg Ser Ser Ser Ala Asn Arg Leu
 355 360 365
 Phe Arg Val Ile Arg Arg Glu His Gly Asp Pro Leu Ile Glu Glu Leu
 370 375 380
 Asn Pro Gly Asp Ala Leu Glu Pro Glu Gly Arg Gly Thr Gly Gly Val
 385 390 395 400
 Val Thr Asp Phe Asp Gly Asp Gly Met Leu Asp Leu Ile Leu Ser His
 405 410 415
 Gly Glu Ser Met Ala Gln Pro Leu Ser Val Phe Arg Gly Asn Gln Gly
 420 425 430
 Phe Asn Asn Asn Trp Leu Arg Val Val Pro Arg Thr Arg Phe Gly Ala
 435 440 445
 Phe Ala Arg Gly Ala Lys Val Val Leu Tyr Thr Lys Lys Ser Gly Ala
 450 455 460
 His Leu Arg Ile Ile Asp Gly Gly Ser Gly Tyr Leu Cys Glu Met Glu
 465 470 475 480
 Pro Val Ala His Phe Gly Leu Gly Lys Asp Glu Ala Ser Ser Val Glu
 485 490 495
 Val Thr Trp Pro Asp Gly Lys Met Val Ser Arg Asn Val Ala Ser Gly
 500 505 510
 Glu Met Asn Ser Val Leu Glu Ile Leu Tyr Pro Arg Asp Glu Asp Thr
 515 520 525
 Leu Gln Asp Pro Ala Pro Leu Glu Cys Gly Gln Gly Phe Ser Gln Gln
 530 535 540
 Glu Asn Gly His Cys Met Asp Thr Asn Glu Cys Ile Gln Phe Pro Phe
 545 550 555 560
 Val Cys Pro Arg Asp Lys Pro Val Cys Val Asn Thr Tyr Gly Ser Tyr
 565 570 575
 Arg Cys Arg Thr Asn Lys Lys Cys Ser Arg Gly Tyr Glu Pro Asn Glu
 580 585 590

-continued

Asp Gly Thr Ala Cys Val Ala Gln Val Ala Phe Leu Gly Gly Tyr Ser
595 600 605

Ser Ala Ala Ser Arg Ile Ser Glu Pro Leu Ser Arg Ala Ser Tyr Leu
610 615 620

Ser Leu Gly Leu Gly Leu Cys Leu Gln Leu Tyr Ala Leu
625 630 635

<210> SEQ ID NO 20
 <211> LENGTH: 618
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 20

Met Ser Arg Met Leu Pro Phe Leu Leu Leu Leu Trp Phe Leu Pro Ile
1 5 10 15

Thr Glu Gly Ser Gln Arg Ala Glu Pro Met Phe Thr Ala Val Thr Asn
20 25 30

Ser Val Leu Pro Pro Asp Tyr Asp Ser Asn Pro Thr Gln Leu Asn Tyr
35 40 45

Gly Val Ala Val Thr Asp Val Asp His Asp Gly Asp Phe Glu Ile Val
50 55 60

Val Ala Gly Tyr Asn Gly Pro Asn Leu Val Leu Lys Tyr Asp Arg Ala
65 70 75 80

Gln Lys Arg Leu Val Asn Ile Ala Val Asp Glu Arg Ser Ser Pro Tyr
85 90 95

Tyr Ala Leu Arg Asp Arg Gln Gly Asn Ala Ile Gly Val Thr Ala Cys
100 105 110

Asp Ile Asp Gly Asp Gly Arg Glu Glu Ile Tyr Phe Leu Asn Thr Asn
115 120 125

Asn Ala Phe Ser Gly Val Ala Thr Tyr Thr Asp Lys Leu Phe Lys Phe
130 135 140

Arg Asn Asn Arg Trp Glu Asp Ile Leu Ser Asp Glu Val Asn Val Ala
145 150 155 160

Arg Gly Val Ala Ser Leu Phe Ala Gly Arg Ser Val Ala Cys Val Asp
165 170 175

Arg Lys Gly Ser Gly Arg Tyr Ser Ile Tyr Ile Ala Asn Tyr Ala Tyr
180 185 190

Gly Asn Val Gly Pro Asp Ala Leu Ile Glu Met Asp Pro Glu Ala Ser
195 200 205

Asp Leu Ser Arg Gly Ile Leu Ala Leu Arg Asp Val Ala Ala Glu Ala
210 215 220

Gly Val Ser Lys Tyr Thr Gly Gly Arg Gly Val Ser Val Gly Pro Ile
225 230 235 240

Leu Ser Ser Ser Ala Ser Asp Ile Phe Cys Asp Asn Glu Asn Gly Pro
245 250 255

Asn Phe Leu Phe His Asn Arg Gly Asp Gly Thr Phe Val Asp Ala Ala
260 265 270

Ala Ser Ala Gly Val Asp Asp Pro His Gln His Gly Arg Gly Val Ala
275 280 285

Leu Ala Asp Phe Asn Arg Asp Gly Lys Val Asp Ile Val Tyr Gly Asn
290 295 300

Trp Asn Gly Pro His Arg Leu Tyr Leu Gln Met Ser Thr His Gly Lys
305 310 315 320

-continued

Val Arg Phe Arg Asp Ile Ala Ser Pro Lys Phe Ser Met Pro Ser Pro
 325 330 335

Val Arg Thr Val Ile Thr Ala Asp Phe Asp Asn Asp Gln Glu Leu Glu
 340 345 350

Ile Phe Phe Asn Asn Ile Ala Tyr Arg Ser Ser Ser Ala Asn Arg Leu
 355 360 365

Phe Arg Val Ile Arg Arg Glu His Gly Asp Pro Leu Ile Glu Glu Leu
 370 375 380

Asn Pro Gly Asp Ala Leu Glu Pro Glu Gly Arg Gly Thr Gly Gly Val
 385 390 395 400

Val Thr Asp Phe Asp Gly Asp Gly Met Leu Asp Leu Ile Leu Ser His
 405 410 415

Gly Glu Ser Met Ala Gln Pro Leu Ser Val Phe Arg Gly Asn Gln Gly
 420 425 430

Phe Asn Asn Asn Trp Leu Arg Val Val Pro Arg Thr Arg Phe Gly Ala
 435 440 445

Phe Ala Arg Gly Ala Lys Val Val Leu Tyr Thr Lys Lys Ser Gly Ala
 450 455 460

His Leu Arg Ile Ile Asp Gly Gly Ser Gly Tyr Leu Cys Glu Met Glu
 465 470 475 480

Pro Val Ala His Phe Gly Leu Gly Lys Asp Glu Ala Ser Ser Val Glu
 485 490 495

Val Thr Trp Pro Asp Gly Lys Met Val Ser Arg Asn Val Ala Ser Gly
 500 505 510

Glu Met Asn Ser Val Leu Glu Ile Leu Tyr Pro Arg Asp Glu Asp Thr
 515 520 525

Leu Gln Asp Pro Ala Pro Leu Glu Cys Gly Gln Gly Phe Ser Gln Gln
 530 535 540

Glu Asn Gly His Cys Met Asp Thr Asn Glu Cys Ile Gln Phe Pro Phe
 545 550 555 560

Val Cys Pro Arg Asp Lys Pro Val Cys Val Asn Thr Tyr Gly Ser Tyr
 565 570 575

Arg Cys Arg Thr Asn Lys Lys Cys Ser Arg Gly Tyr Glu Pro Asn Glu
 580 585 590

Asp Gly Thr Ala Cys Val Gly Trp Trp Ser Pro Val Leu Lys Ile Val
 595 600 605

Thr Pro Gln Val Gly Lys Ser Leu Gly Pro
 610 615

<210> SEQ ID NO 21
 <211> LENGTH: 612
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 21

Met Ser Arg Met Leu Pro Phe Leu Leu Leu Leu Trp Phe Leu Pro Ile
 1 5 10 15

Thr Glu Gly Ser Gln Arg Ala Glu Pro Met Phe Thr Ala Val Thr Asn
 20 25 30

Ser Val Leu Pro Pro Asp Tyr Asp Ser Asn Pro Thr Gln Leu Asn Tyr
 35 40 45

Gly Val Ala Val Thr Asp Val Asp His Asp Gly Asp Phe Glu Ile Val

-continued

50				55				60							
Val 65	Ala	Gly	Tyr	Asn 70	Gly	Pro	Asn	Leu	Val	Leu 75	Lys	Tyr	Asp	Arg	Ala 80
Gln	Lys	Arg	Leu	Val 85	Asn	Ile	Ala	Val	Asp 90	Glu	Arg	Ser	Ser	Pro	Tyr 95
Tyr	Ala	Leu	Arg	Asp 100	Arg	Gln	Gly	Asn 105	Ala	Ile	Gly	Val	Thr 110	Ala	Cys
Asp	Ile	Asp 115	Gly	Asp	Gly	Arg	Glu	Glu 120	Ile	Tyr	Phe	Leu 125	Asn	Thr	Asn
Asn	Ala	Phe 130	Ser	Gly	Val	Ala 135	Thr	Tyr	Thr	Asp 140	Lys	Leu	Phe	Lys	Phe
Arg	Asn	Asn	Arg	Trp	Glu 150	Asp	Ile	Leu	Ser	Asp 155	Glu	Val	Asn	Val	Ala 160
Arg	Gly	Val	Ala	Ser 165	Leu	Phe	Ala	Gly	Arg 170	Ser	Val	Ala	Cys	Val	Asp 175
Arg	Lys	Gly	Ser	Gly 180	Arg	Tyr	Ser	Ile 185	Tyr	Ile	Ala	Asn	Tyr	Ala	Tyr 190
Gly	Asn	Val 195	Gly	Pro	Asp	Ala	Leu	Ile 200	Glu	Met	Asp	Pro	Glu	Ala	Ser 205
Asp	Leu	Ser	Arg	Gly	Ile	Leu 215	Ala	Leu	Arg	Asp 220	Val	Ala	Ala	Glu	Ala
Gly	Val	Ser	Lys	Tyr	Thr 230	Gly	Gly	Arg	Gly	Val 235	Ser	Val	Gly	Pro	Ile 240
Leu	Ser	Ser	Ser	Ala 245	Ser	Asp	Ile	Phe	Cys 250	Asp	Asn	Glu	Asn	Gly	Pro 255
Asn	Phe	Leu	Phe	His 260	Asn	Arg	Gly	Asp 265	Gly	Thr	Phe	Val	Asp 270	Ala	Ala
Ala	Ser	Ala 275	Gly	Val	Asp	Asp	Pro	His 280	Gln	His	Gly	Arg 285	Gly	Val	Ala
Leu	Ala	Asp 290	Phe	Asn	Arg	Asp 295	Gly	Lys	Val	Asp 300	Ile	Val	Tyr	Gly	Asn
Trp	Asn	Gly	Pro	His 310	Arg	Leu	Tyr	Leu	Gln	Met 315	Ser	Thr	His	Gly	Lys 320
Val	Arg	Phe	Arg	Asp 325	Ile	Ala	Ser	Pro	Lys 330	Phe	Ser	Met	Pro	Ser	Pro 335
Val	Arg	Thr 340	Val	Ile	Thr	Ala	Asp	Phe 345	Asp	Asn	Asp	Gln	Glu	Leu	Glu 350
Ile	Phe	Phe 355	Asn	Asn	Ile	Ala	Tyr	Arg 360	Ser	Ser	Ser	Ala	Asn	Arg	Leu 365
Phe	Arg	Val 370	Ile	Arg	Arg	Glu	His	Gly 375	Asp	Pro	Leu	Ile	Glu	Glu	Leu 380
Asn	Pro	Gly	Asp	Ala 390	Leu	Glu	Pro	Glu	Gly	Arg 395	Gly	Thr	Gly	Gly	Val 400
Val	Thr	Asp	Phe	Asp 405	Gly	Asp	Gly	Met	Leu 410	Asp	Leu	Ile	Leu	Ser	His 415
Gly	Glu	Ser	Met	Ala 420	Gln	Pro	Leu	Ser 425	Val	Phe	Arg	Gly	Asn	Gln	Gly 430
Phe	Asn	Asn	Asn	Trp 435	Leu	Arg	Val	Val 440	Pro	Arg	Thr	Arg 445	Phe	Gly	Ala
Phe	Ala	Arg	Gly	Ala 450	Lys	Val	Val	Leu	Tyr	Thr	Lys	Lys	Ser	Gly	Ala 460

-continued

His Leu Arg Ile Ile Asp Gly Gly Ser Gly Tyr Leu Cys Glu Met Glu
 465 470 475 480
 Pro Val Ala His Phe Gly Leu Gly Lys Asp Glu Ala Ser Ser Val Glu
 485 490 495
 Val Thr Trp Pro Asp Gly Lys Met Val Ser Arg Asn Val Ala Ser Gly
 500 505 510
 Glu Met Asn Ser Val Leu Glu Ile Leu Tyr Pro Arg Asp Glu Asp Thr
 515 520 525
 Leu Gln Asp Pro Ala Pro Leu Glu Cys Gly Gln Gly Phe Ser Gln Gln
 530 535 540
 Glu Asn Gly His Cys Met Asp Thr Asn Glu Cys Ile Gln Phe Pro Phe
 545 550 555 560
 Val Cys Pro Arg Asp Lys Pro Val Cys Val Asn Thr Tyr Gly Ser Tyr
 565 570 575
 Arg Cys Arg Thr Asn Lys Lys Cys Ser Arg Gly Tyr Glu Pro Asn Glu
 580 585 590
 Asp Gly Thr Ala Cys Val Glu Arg Thr Leu Leu Leu Gly Leu Cys Asn
 595 600 605
 Leu Leu Gly Lys
 610

1. A polynucleotide encoding a polypeptide comprising the amino acid sequence extending from amino acid number 1-598 of SEQ ID NO: 2.

2. The polynucleotide according to claim 1 said polynucleotide encoding the amino acid sequence of SEQ ID NO: 2, SEQ ID NO: 19, SEQ ID NO: 20 or SEQ ID NO: 21.

3. The polynucleotide according to claim 1 or 2 said polynucleotide comprising the sequence extending from nucleotides 59-1852 of SEQ ID NO: 1.

4. The polynucleotide according to claims 1-3, said polynucleotide comprising the sequences SEQ ID NO: 1, SEQ ID NO: 16, SEQ ID NO: 17 or SEQ ID NO: 18 or the sequence extending from nucleotides 59-2017 from SEQ ID NO: 1, nucleotides 59-1969 from SEQ ID NO: 16, nucleotides 59-1912 from SEQ ID NO: 17 or nucleotides 59-1894 from SEQ ID NO: 18.

5. A recombinant expression vector comprising the DNA according to claims 1-4.

6. A polypeptide encoded by the polynucleotide according to claims 1-4, the expression vector according to claim 5 or immunomodulating fragments of SEQ ID NO: 2, SEQ ID NO: 19, SEQ ID NO: 20 or SEQ ID NO: 21.

7. Pharmaceutical composition comprising the polypeptide according to claim 6, and a pharmaceutical acceptable carrier.

8. The polypeptide according to claim 6 for use in therapy.

9. Use of the peptide according to claim 6 for the manufacture of a pharmaceutical preparation against inflammatory diseases.

10. Use of the polypeptide according to claims 6 for the manufacture of a pharmaceutical preparation for the induction of immunological tolerance to an autoantigen in patients suffering from autoimmune disorders, more specifically rheumatoid arthritis.

11. A cell transfected with DNA according to claims 1-4 or the expression vector according to claim 5.

12. A cell according to claim 11 which is a stable transfected cell which expresses the polypeptide according to claims 6.

13. Use of a DNA according to claims 1-4 or an expression vector according to claim 5, a cell according to claims 11 or 12 or a polypeptide according to claim 6 in a screening assay for identification of tolerogenic (poly)peptides.

14. A method for identifying tolerogenic peptides, said method comprising the steps of

- a) introducing into a suitable host cell a fragment of the polynucleotide according to claims 1-4;
- b) culturing the host cells under conditions to allow expression of the introduced sequence;
- c) bringing the expression product in contact with lymphocytes; and
- d) establishing the lymphocyte activity.

15. A method for identifying tolerogenic peptides, said method comprising the steps of

- a) introducing into a suitable host cell a fragment of the polynucleotide according to claims 1-4;
- b) culturing the host cells under conditions to allow expression of the introduced DNA sequence;
- c) isolating the expression product;
- d) bringing the expression product in contact with antigen presenting cells and T cells; and
- e) establishing the T cell activity.

16. A method for identifying tolerogenic peptides, said method comprising the steps of

- a) chemically synthesizing a fragment of SEQ ID NO: 2, SEQ ID NO: 19, SEQ ID NO: 20 or SEQ ID NO: 21;
- b) bringing the peptide fragment in contact with antigen presenting cells and T cells; and
- c) establishing the T cell activity.

17. A method for the formulation of a pharmaceutical composition comprising the method of claims **14-16** and mixing the peptide identified with a pharmaceutically acceptable carrier.

18. Use of a peptide as identified in the method of claims **14-16** for the preparation of a pharmaceutical suitable as a tolerogenic agent.

19. Diagnostic composition comprising one or more of the polypeptide according to claim 6 and a detection agent.

20. A diagnostic method for the detection of activated autoreactive T cells comprising the following steps:

- a) isolation of the peripheral blood mononuclear cells (PBMC) from a blood sample of an individual,
- b) culture of said PBMC under suitable conditions,
- c) incubation of said PBMC culture in the presence of SCIM-1, fragments thereof and/or one or more peptides according to claim 6, and
- d) detection of a response of T cells, indicating the presence of activated autoreactive T cells in the individual.

21. Test kit for the detection of activated autoreactive T cells, said test kit comprising SCIM-1 or one or more of the peptides according to claims 6.

* * * * *

专利名称(译)	细胞外基质蛋白		
公开(公告)号	US20040072286A1	公开(公告)日	2004-04-15
申请号	US10/332929	申请日	2001-07-09
[标]申请(专利权)人(译)	HEUS HENDRIK CORNELIS NELISSEN 罗伯特·路易斯 HUBERT MEEUWISSE CORNELIS MARIA LEONARDUS		
申请(专利权)人(译)	HEUS HENDRIK CORNELIS NELISSEN 罗伯特·路易斯 HUBERT MEEUWISSE CORNELIS MARIA LEONARDUS		
当前申请(专利权)人(译)	HEUS HENDRIK CORNELIS NELISSEN 罗伯特·路易斯 HUBERT MEEUWISSE CORNELIS MARIA LEONARDUS		
[标]发明人	HEUS HENDRIK CORNELIS NELISSEN ROBERT LOUIS HUBERT MEEUWISSE CORNELIS MARIA LEONARDUS		
发明人	HEUS, HENDRIK CORNELIS NELISSEN, ROBERT LOUIS HUBERT MEEUWISSE, CORNELIS MARIA LEONARDUS		
IPC分类号	G01N33/50 A61K38/00 A61K38/17 A61P19/02 A61P29/00 A61P37/02 A61P37/06 C07K14/47 C12N1/15 C12N1/19 C12N1/21 C12N5/10 C12N15/09 C12N15/12 C12N15/85 C12Q1/02 C12Q1/68 G01N33/15 G01N33/53 G01N33/68 C12P21/02 C12N5/06		
CPC分类号	C07K14/4713 A61K38/00 A61P19/02 A61P29/00 A61P37/00 A61P37/02 A61P37/06		
优先权	2000202495 2000-07-13 EP		
外部链接	Espacenet USPTO		

摘要(译)

本发明描述了一种新蛋白质及其变体以及编码这些蛋白质的基因。该蛋白质可用于预防炎性疾病，更具体地说，可用于在患有类风湿性关节炎的患者中诱导T细胞对蛋白质的耐受性。

1 2 3 4 5 6 7 8 9 10




Figure 1