



US 20100104501A1

(19) **United States**

(12) **Patent Application Publication**
Golz et al.

(10) **Pub. No.: US 2010/0104501 A1**
(43) **Pub. Date: Apr. 29, 2010**

(54) **PRSS23 AS A BIOMARKER, THERAPEUTIC AND DIAGNOSTIC TARGET**

(76) Inventors: **Stefan Golz**, Essen (DE); **Holger Summer**, Wuppertal (DE); **Andreas Geerts**, Wuppertal (DE); **Ulf Brüggemeier**, Leichlingen (DE); **Barbara Albrecht**, Wulfrath (DE); **Martina Klein**, Dusseldorf (DE); **Sonja Steppan**, Neu-Isenburg (DE); **Peter Ellinghaus**, Melle (DE); **Donatella D'urso**, Duesseldorf (DE); **Michael Seewald**, Langenfeld (DE); **Hendrik Milting**, Bad Oeynhausen (DE)

Correspondence Address:
BANNER & WITCOFF, LTD.
1100 13th STREET, N.W., SUITE 1200
WASHINGTON, DC 20005-4051 (US)

(21) Appl. No.: **12/445,494**
(22) PCT Filed: **Oct. 2, 2007**
(86) PCT No.: **PCT/EP07/08562**
§ 371 (c)(1),
(2), (4) Date: **Nov. 10, 2009**

(30) **Foreign Application Priority Data**

Oct. 16, 2006 (EP) 06021606.6

Publication Classification

(51) **Int. Cl.**
A61K 51/00 (2006.01)
C12Q 1/68 (2006.01)
G01N 33/53 (2006.01)
A61K 49/00 (2006.01)
(52) **U.S. Cl.** **424/1.11**; 435/6; 435/7.1; 424/9.1; 424/9.5

(57) **ABSTRACT**

The invention provides PRSS23, which is associated with cardiovascular diseases, hematological diseases, neurological diseases, cancer, endocrinological diseases, and urological diseases. The invention also provides assays for the identification of compounds useful in the treatment or prevention of cardiovascular diseases, hematological diseases, neurological diseases, cancer, endocrinological diseases, and urological diseases. The invention also features compounds which bind to and/or activate or inhibit the activity of PRSS23 as well as pharmaceutical compositions comprising such compounds. The invention also provides PRSS23 as a biomarker for diseases such as cardiovascular diseases, hematological diseases, neurological diseases, cancer, endocrinological diseases, and urological diseases.

Fig. 2

SEQ ID NO: 2

CGCGCGGTCTCGGGGCGCCACACCTGTCTGAGCGGCGCACGGCCGCGGCCCCGGCGGG
CTGCTCCACGCGGTAGCACTCAGCATGGCTGGAATCCCAGGACTCCTCATCCTTCTCCTTG
TCCTGCTCTGTGTGTTTCATGCAGGTGAGTCCCTACAACGTTCCGTGGAAACCAACATGGCC
GGCTTACCGCCTCCCTATAGTCTTGCCCTCAGTCTACCCCTCAAGTTAGCCAAAACCAGACTTC
GATGCCAAAGCAAATTTGGAGGTGTCCTCCTCATGTGGACCCCAAGTGTACAAAGGGAACAC
CGCTGCCCACCTACGAAGAGGCCAAGCAGTACCTTTCTTATGAAACCCTTTATGCCAACGG
CAGCCGCACAGAGACTCAGGTGGGCATCTACATCCTCAGCAACGGTGAAGGCAGGGCACGC
AGCAGAGACTCGGAGGCCGAGGGAAATCTCGCAGGAAGAGGCAGATTTATGGCTATGATG
GCAGATTTAGCATTTTCGGGAAGGACTTCCTGCTCAACTATCCCTTCTCAACATCGGTGAA
GTTGTCTACTGGCTGCACCGGCACCCTGGTGGCAGAGAAGCACGTCTCCTACTGCTGCCAC
TGCATACACGATGGCAAACCTATGTGAAAGGAACACAGAACTCCGAGTGGGCTTCTTGA
AGCCCAAGTATAAAGATGGTGCCGGAGGGGACAACAGTTCAAGCTCAGCTCTGGTAGAGAA
GATGAAATTTAGTGGATCCGTGTGAAACGCACCCATGTGCCCAAGGGGTGGATCAAGGGC
AATGCCAACGACATTGGCATGGATTATGACTACGCCCTTCTGGAACCTCAAGAAACCCACA
AAAGAAAGTTCATGAAGATGGTGTGAGTCCCTCCAGCGAAGCAGCTCCCGGGGGGCAGGAT
CCACTTCTCAGGCTACGACAATGACCGGCCCGGCAATTTGGTGTACCGCTTCTGTGATGTC
AAAGATGAGACATACGACCTTCTTTATCAGCAGTGTGATGCCAGCCCGGGGCCAGTGGTT
CAGGGGTCTATGTGAGGATGTGGAAGAGACCACAGCAGAAATGGGAAAGGAAAATTTATCGG
CATCTTTTACAGGGCACCAGTGGGTGGACATGAATGGCTCTCCGCAGGATTTCAACGTGGCA
GTTAGAATCACACCTCTTAAATATGCCAGATTTGCTATTGGATTAAAGGAACTACCTGG
ATTGCAGGGAGGGGTGACATGCGTCTTCTTGGCAGCACCGATGGTTCGTTTTGCACTCATTG
TAGGAGAGAGGCCAGCTTTATATCATTGACTCTTGTGGTGTGTGTCACATAGTATCTTTTA
CCTAATATTCTTCAAATAGCAAACCTATTGGTCTATTATTTGAAAATTGTTGTGTGCGTT
ATAGCATTTAAGCAGTCTGAAAGCATACTTTTTACATAGAGACTTTAAAATATTGATGTTT
GGTTAATAGGGGATATTTGGCAAGTAAATTAACTTTCAGCTTTTGGAGAGCTCTGATTT
TTGCCCTTATCCAAACATGCTTCAGAGGTTTATATCAAATATGTGACATACCGGAATATGG
ATTCCTTATGTTTGTATATGTATATGTCTTCTGAGAGTCATACATATATTGATATTTTTTTA
ATGTGTGGTTATAGTGTTCCAAATGACAATAGCAAAGCTTGCAATGGGCAATCATAATGT
TTCCGATTCAAACATTTACATAGTAGTCTTGAAGAGAAACAGTAATTTATTGGCTCTATT
GATACACATGTAAAGCTGTATCTTACAGTGCAGAGAATTTCCAGGCTGCTTTTAGTTTTGA
AAATAAACTCTGTCTTCTAAAAA

Fig. 3

SEQ ID NO: 3

MAGIPGLLFLFFLLCAVGQVSPYSAPWKPTWPA YRLPVVLPQSTLNLAKPDFGAE
AKLEVSSSCGPQCHKGTPLPTYEEAKQYLSYETLYANGSRTETQVGIYILSSSGDGA
QHRDSGSSGKSRRKRQIYGYDSRFSIFGKDFLLNYPFSTSVKLSGCTGTLVAEKHVL
TAAHCIHDGKTYVKGTQKLRVGF LKPKFKDGGRGANDSTSAMPEQMKFQWIRVK
RTHVPGWIKGNANDIGMDYDYALLELKKPHKRKFMKIGVSPPAKQLPGGRIHFS
GYDNDRPGNLVYRFCDVKDETYD LLYQQCDAQPGASGSGVYVRMWKRPQQKW
ERKIIIGIFSGHQWVDMNGSPQDFNVA VRITPLKYAQICYWIKGNYLDCREG

Fig. 4

SEQ ID NO: 4

MAGIPGLLILLVLLCVFMQVSPYNVPWKPTWPA YRLPIVLPQSTLKLAKPDFDAK
AKLEVSSSCGPQCHKGTPLPTYEEAKQYLSYETLYANGSRTETQVGIYILSNGEGR
ARSRDSEAAGKSRRKRQIYGYDGRFSIFGKDFLLNYPFSTSVKLSTGCTGTLVAEK
HVLTAHCIHDGKTYVKGTQKLRVGF LKPKYKDGAGDNNSSSSALVEKMKFQWI
RVKRTHVPGWIKGNANDIGMDYDYALLELKKPHKRKFMKIGVSPPAKQLPGGRI
HFSGYDNDRPGNLVYRFCDVKDETYD LLYQQCDAQPGASGSGVYVRMWKRPQQ
KWERKIIIGIFSGHQWVDMNGSPQDFNVA VRITPLKYAQICYWIKGNYLDCREG

Fig. 5

SEQ ID NO: 5

5' CGCCACACCTGTCTGA 3'

Fig. 6

SEQ ID NO: 6

5' TGGAATCCCTGCCATGC 3'

Fig. 7

SEQ ID NO: 7

5' CTCGGCGCGGAACAGTGCTC 3'

Fig. 8

SEQ ID NO: 8

5' CGCCCACACCTGTCTGA 3'

Fig. 9

SEQ ID NO: 9

5' TGGGATTCCAGCCATGC 3'

Fig. 10

SEQ ID NO: 10

5' CTGCTCCACGCGGTAGCACTCA 3'

Fig. 11

Real-time expression data of PRSS23 in rat heart (DOCA model)




Fig. 12

Real-time expression data of PRSS23 in rat heart (occlusion model)




Fig. 13

Real-time expression data of PRSS23 in rat heart (monocrotalin model)




Fig. 14

Microarray expression data of PRSS23 in rat heart (DOCA model)




Fig. 15

Microarray expression data of PRSS23 in rat heart (occlusion model)




Fig. 16

Microarray expression data of PRSS23 in rat heart (monocrotalin model)





Fig. 17

Microarray expression data of PRSS23 in human heart



PRSS23 AS A BIOMARKER, THERAPEUTIC AND DIAGNOSTIC TARGET

TECHNICAL FIELD OF THE INVENTION

[0001] The present invention is in the field of molecular biology, more particularly, the present invention relates to nucleic acid sequences and amino acid sequences of a human and rat PRSS23 and its regulation for the treatment, diagnostic and use as a biomarker of cardiovascular diseases, hematological diseases, neurological diseases, respiratory diseases, gastroenterological diseases and urological diseases in mammals.

BACKGROUND OF THE INVENTION

[0002] Proteases play a role in carefully controlled processes, such as blood coagulation, fibrinolysis, complement activation, fertilization, and hormone production. These enzymes are also used in a variety of diagnostic, therapeutic, and industrial contexts. PRSS23 is a member of the group of protease enzymes [DesGroseillers et al. (2001)].

[0003] Proteases were recognized very early in the history of biochemistry. In the nineteenth century, one primary focus of research was on digestive proteases, like pepsin and trypsin. Proteases belong systematically to the C—N Hydrolases. More specifically, proteases catalyze the hydrolytic cleavage of a peptide bond and are therefore called peptidases as well.

[0004] Proteases can be classified according to several criteria, e.g. by localisation. Digestive proteases are located in the gastro-intestinal tract. These proteases are responsible for the digestion of food proteins.

[0005] Peptidases located extracellularly in the blood or other extracellular compartments of the body play often regulatory roles in processes like for example blood clotting, fibrinolysis, or activation of complement constituents.

[0006] Intracellularly located proteases exhibit a wide variety of roles. They are found in compartments like the ER, the Golgi apparatus, or the lysosomes. Their functions include for example activation of peptide hormones, ubiquitin mediated proteolysis, among others.

[0007] Proteases are most commonly classified according to their mechanism of action, or to specific active groups that are present in the active center. The following groups can be distinguished:

- [0008] 1. Serin-peptidases, 2. cystein-peptidases, 3. aspartyl- or acidic-peptidases, 4. metallo-peptidases, or 5. peptidases with yet unclear reaction mechanism.

Serine Peptidases

[0009] Serine proteases exhibit a serine in the catalytic site which forms a covalent ester intermediate during the catalytic reaction pathway by a nucleophilic attack on the carboxy carbon of the peptide bond. In the active site of serine proteases a catalytic triad comprised of an aspartate, a histidine and the above mentioned serine is found. This triad functions in the reaction mechanism as a charge relay system.

[0010] To the large family of serine protease belong, for example, the digestive enzymes trypsin and chymotrypsin, components of the complement cascade, enzymes involved in the blood clotting cascade, as well as enzymes that function in degradation, rebuilding and maintenance of constituents of the extracellular matrix.

[0011] One feature of the serine protease family is the broad range of substrate specificity. Members of the trypase subgroup cleave after arginine or lysine, chymases after phenylalanine or leucine, aspartases after aspartate, metases after methionine and serases after serine.

Cysteine Proteases

[0012] During the catalytic reaction of cysteine proteases a covalent thioester intermediate is formed by a nucleophilic attack of the cysteine on the carboxy carbon of the peptide bond. Similar to the serine peptidases a catalytic triad comprised of the cysteine, a histidine and an asparagine is found which functions as a charge relay system to facilitate the formation of the thioester intermediate.

[0013] Members of the Cysteine protease family have roles in many different cellular processes, e.g. processing of precursors or intracellular degradation. Examples for cysteine proteases include lysosomal cathepsins, and cytosolic calpains.

Aspartyl- or Acidic Peptidase

[0014] The catalytic site of aspartyl proteases is composed of two aspartate residues. At the pH optimum of aspartyl proteases (2-3) one of the aspartyl carboxy groups is ionized and the other is neutral, which is important for the catalytic reaction to occur. Examples for aspartyl proteases are gastric pepsins A and C, chymosin, as well as mammalian renin.

Metallo-Peptidases

[0015] Metallo-peptidases are proteases, whose proteolytic activity depends on the presence of divalent cations in the active center. Examples of members of this class are carboxypeptidase A, which represents a pancreatic digestive enzyme, the Angiotension Converting Enzymes (ACE), which are responsible for the conversion of angiotensin I to angiotensin II, or the Extracellular Matrix Metalloproteinases.

[0016] In summary, a huge number of proteases play a central role in several important cellular and intracellular processes. Furthermore, the value as pharmaceutical targets has been proven for several proteases. For example, the protease encoded by the HIV genome is used as a target for drugs for the treatment of HIV infections, the proteasom complex has been discovered as an anti-cancer target, or Cys-proteases have been implemented as drug targets for inflammatory disorders. Selective inhibitors have been developed as therapeutic agents for diseases such as HIV. Thus, the identification of further disease implications of protease species and their splice variants may lead to the development of specific inhibitors or modulators, or suggest new utilities for known compounds affecting proteases. That in turn will provide additional pharmacological approaches to treat diseases and conditions in which protease activities are involved. These diseases may include, but are not limited to, infections such as bacterial, fungal, protozoan, and viral infections, particularly those caused by HIV viruses, cancers, allergies including asthma, cardiovascular diseases including acute heart failure, hypotension, hypertension, angina pectoris, myocardial infarction, hematological diseases, genito-urinary diseases including urinary incontinence and benign prostate hyperplasia, osteoporosis, peripheral and central nervous system disorders including pain, Alzheimer's disease and Parkinson's disease, respiratory diseases, metabolic diseases, inflamma-

tory diseases, gastro-enterological diseases, diseases of the endocrine system, dermatological diseases, diseases of muscles or the skeleton, immunological diseases, developmental diseases or diseases of the reproductive system.

Taqman-Technology/Expression Profiling

[0017] TaqMan is a recently developed technique, in which the release of a fluorescent reporter dye from a hybridisation probe in real-time during a polymerase chain reaction (PCR) is proportional to the accumulation of the PCR product. Quantification is based on the early, linear part of the reaction, and by determining the threshold cycle (CT), at which fluorescence above background is first detected.

[0018] Gene expression technologies may be useful in several areas of drug discovery and development, such as target identification, lead optimization, and identification of mechanisms of action. The TaqMan technology can be used to compare differences between expression profiles of normal tissue and diseased tissue. Expression profiling has been used in identifying genes, which are up- or downregulated in a variety of diseases. An interesting application of expression profiling is temporal monitoring of changes in gene expression during disease progression and drug treatment or in patients versus healthy individuals. The premise in this approach is that changes in pattern of gene expression in response to physiological or environmental stimuli (e.g., drugs) may serve as indirect clues about disease-causing genes or drug targets. Moreover, the effects of drugs with established efficacy on global gene expression patterns may provide a guidepost, or a genetic signature, against which a new drug candidate can be compared.

PRSS23

[0019] The nucleotide sequence of PRSS23 is accessible in the databases by the accession number AF015287 (human) and BC079179 (rat). The sequences are given in SEQ ID NO:1 (human) and SEQ ID NO:2 (rat). The amino acid sequence of PRSS23 depicted in SEQ ID NO:3 (human) and SEQ ID NO:4 (rat).

[0020] PRSS23 is described as a secreted protease with synonyms putative secreted protein ZSIG13; serine protease, umbilical endothelium; MGC5107 and SPUVE. PRSS23 mRNA expression is described as regulated in cancer tissues [Ifon et al. (2005); Pentecost, (2005)].

[0021] PRSS23 is published (but not limited to) in patents WO02081745, EP1612281 and EP1560025.

SUMMARY OF THE INVENTION

[0022] The invention relates to the use of PRSS23 polypeptides and polynucleotides as a biomarker in cardiovascular diseases, hematological diseases, neurological diseases, respiratory diseases, gastroenterological diseases and urological diseases. The invention also relates to novel disease associations of PRSS23 polypeptides and polynucleotides. The invention also relates to the use of PRSS23 as a biomarker for cardiovascular diseases, hematological diseases, neurological diseases, respiratory diseases, gastroenterological diseases and urological diseases. The invention also relates to novel methods of screening for therapeutic agents for the treatment of cardiovascular diseases, hematological diseases, neurological diseases, respiratory diseases, gastroenterological diseases and urological diseases in a mammal. The invention also relates to pharmaceutical compositions for the treat-

ment of cardiovascular diseases, hematological diseases, neurological diseases, respiratory diseases, gastroenterological diseases and urological diseases in a mammal comprising a PRSS23 polypeptide, a PRSS23 polynucleotide, or regulators of PRSS23 or modulators of PRSS23 activity. The invention further comprises methods of diagnosing cardiovascular diseases, hematological diseases, neurological diseases, respiratory diseases, gastro-enterological diseases and urological diseases in a mammal.

BRIEF DESCRIPTION OF THE DRAWINGS

[0023] FIG. 1 shows the nucleotide sequence of a PRSS23 polynucleotide human (SEQ ID NO:1).

[0024] FIG. 2 shows the nucleotide sequence of a PRSS23 polynucleotide rat (SEQ ID NO:2).

[0025] FIG. 3 shows the amino acid sequence of a PRSS23 polypeptide human (SEQ ID NO:3).

[0026] FIG. 4 shows the amino acid sequence of a PRSS23 polypeptide rat (SEQ ID NO:4).

[0027] FIG. 5 shows the nucleotide sequence of a primer useful for the invention (SEQ ID NO:5).

[0028] FIG. 6 shows the nucleotide sequence of a primer useful for the invention (SEQ ID NO:6).

[0029] FIG. 7 shows a nucleotide sequence useful as a probe to detect proteins of the invention (SEQ ID NO:7).

[0030] FIG. 8 shows the nucleotide sequence of a primer useful for the invention (SEQ ID NO:8).

[0031] FIG. 9 shows the nucleotide sequence of a primer useful for the invention (SEQ ID NO:9).

[0032] FIG. 10 shows a nucleotide sequence useful as a probe to detect proteins of the invention (SEQ ID NO:10).

[0033] FIG. 11 shows the results of real-time expression analysis of PRSS23 in rat hearts (DOCA). X axis: treatment; Y axis: relative expression; A: control; B: control/placebo; C1: compound P/concentration 1; C2: compound P/concentration 2; C3: compound P/concentration. The expression of PRSS23 is disease state, treatment- and dose-dependent regulated in the animal model.

[0034] FIG. 12 shows the results of real-time expression analysis of PRSS23 in rat hearts (occlusion). X axis: treatment; Y axis: relative expression; A: control; B: control/placebo; C1: compound P/concentration 1; C2: compound P/concentration 2; C3: compound P/concentration. The expression of PRSS23 is disease state, treatment- and dose-dependent regulated in the animal model.

[0035] FIG. 13 shows the results of real-time expression analysis of PRSS23 in rat hearts (monocrotalin). X axis: treatment; Y axis: relative expression; A: control; B: control/placebo; C1: compound P/concentration 1; C2: compound P/concentration 2. The expression of PRSS23 is disease state, treatment- and dose-dependent regulated in the animal model.

[0036] FIG. 14 shows the results of microarray expression analysis of PRSS23 in rat hearts (Doca). X axis: treatment; Y axis: relative expression; A: control; B: control/placebo; C: compound P/concentration 1; D: compound P/concentration 2. The expression of PRSS23 is disease state, treatment- and dose-dependent regulated in the animal model.

[0037] FIG. 15 shows the results of microarray expression analysis of PRSS23 in rat hearts (Occlusion). X axis: treatment; Y axis: relative expression; A: control; B: control/placebo; C: compound P/concentration 1; D: compound

P/concentration 2. The expression of PRSS23 is disease state, treatment- and dose-dependent regulated in the animal model.

[0038] FIG. 16 shows the results of microarray expression analysis of PRSS23 in rat hearts (monocrotalin). X axis: treatment; Y axis: relative expression; A: control; B: control/placebo; C: compound P/concentration 1; D: compound P/concentration 2. The expression of PRSS23 is disease state, treatment- and dose-dependent regulated in the animal model.

[0039] FIG. 17 shows the results of microarray expression analysis of PRSS23 in human hearts. X axis: disease status; Y axis: relative expression; N: non-failure; P: pre-LVAD. The expression of PRSS23 is disease state regulated in human heart.

DETAILED DESCRIPTION OF THE INVENTION

Definition of Terms

[0040] An “oligonucleotide” is a stretch of nucleotide residues which has a sufficient number of bases to be used as an oligomer, amplicon or probe in a polymerase chain reaction (PCR). Oligo-nucleotides are prepared from genomic or cDNA sequence and are used to amplify, reveal, or confirm the presence of a similar DNA or RNA in a particular cell or tissue. Oligonucleotides or oligomers comprise portions of a DNA sequence having at least about 10 nucleotides and as many as about 35 nucleotides, preferably about 25 nucleotides.

[0041] “Probes” may be derived from naturally occurring or recombinant single- or double-stranded nucleic acids or may be chemically synthesized. They are useful in detecting the presence of identical or similar sequences. Such probes may be labeled with reporter molecules using nick translation, Klenow fill-in reaction, PCR or other methods well known in the art. Nucleic acid probes may be used in southern, northern or in situ hybridizations to determine whether DNA or RNA encoding a certain protein is present in a cell type, tissue, or organ.

[0042] A “fragment of a polynucleotide” is a nucleic acid that comprises all or any part of a given nucleotide molecule, the fragment having fewer nucleotides than about 6 kb, preferably fewer than about 1 kb.

[0043] “Reporter molecules” are radionuclides, enzymes, fluorescent, chemiluminescent, or chromogenic agents which associate with a particular nucleotide or amino acid sequence, thereby establishing the presence of a certain sequence, or allowing for the quantification of a certain sequence.

[0044] “Chimeric” molecules may be constructed by introducing all or part of the nucleotide sequence of this invention into a vector containing additional nucleic acid sequence which might be expected to change any one or several of the following PRSS23 characteristics: cellular location, distribution, ligand-binding affinities, interchain affinities, degradation/turnover rate, signaling, etc.

[0045] “Active”, with respect to a PRSS23 polypeptide, refers to those forms, fragments, or domains of a PRSS23 polypeptide which retain the biological and/or antigenic activity of a PRSS23 polypeptide.

[0046] “Naturally occurring PRSS23 polypeptide” refers to a polypeptide produced by cells which have not been genetically engineered and specifically contemplates various polypeptides arising from post-translational modifications of

the polypeptide including but not limited to acetylation, carboxylation, glycosylation, phosphorylation, lipidation and acylation.

[0047] “Derivative” refers to polypeptides which have been chemically modified by techniques such as ubiquitination, labeling (see above), pegylation (derivatization with polyethylene glycol), and chemical insertion or substitution of amino acids such as ornithine which do not normally occur in human proteins.

[0048] “Conservative amino acid substitutions” result from replacing one amino acid with another having similar structural and/or chemical properties, such as the replacement of a leucine with an isoleucine or valine, an aspartate with a glutamate, or a threonine with a serine.

[0049] “Insertions” or “deletions” are typically in the range of about 1 to 5 amino acids. The variation allowed may be experimentally determined by producing the peptide synthetically while systematically making insertions, deletions, or substitutions of nucleotides in the sequence using recombinant DNA techniques.

[0050] A “signal sequence” or “leader sequence” can be used, when desired, to direct the polypeptide through a membrane of a cell. Such a sequence may be naturally present on the polypeptides of the present invention or provided from heterologous sources by recombinant DNA techniques.

[0051] An “oligopeptide” is a short stretch of amino acid residues and may be expressed from an oligo-nucleotide. Oligopeptides comprise a stretch of amino acid residues of at least 3, 5, 10 amino acids and at most 10, 15, 25 amino acids, typically of at least 9 to 13 amino acids, and of sufficient length to display biological and/or antigenic activity.

[0052] “Inhibitor” is any substance which retards or prevents a chemical or physiological reaction or response. Common inhibitors include but are not limited to antisense molecules, antibodies, and antagonists.

[0053] “Biomarker” are measurable and quantifiable biological parameters (e.g. specific enzyme concentration, specific hormone concentration, specific gene phenotype distribution in a population, presence of biological substances) which serve as indices for health—and physiology related assessments, such as disease risk, psychiatric disorders, environmental exposure and its effects, disease diagnosis, metabolic processes, substance abuse, pregnancy, cell line development, epidemiologic studies, etc. Parameter that can be used to identify a toxic effect in an individual organism and can be used in extrapolation between species. Indicator signalling an event or condition in a biological system or sample and giving a measure of exposure, effect, or susceptibility.

[0054] Biological markers can reflect a variety of disease characteristics, including the level of exposure to an environmental or genetic trigger, an element of the disease process itself, an intermediate stage between exposure and disease onset, or an independent factor associated with the disease state but not causative of pathogenesis. Depending on the specific characteristic, biomarkers can be used to identify the risk of developing an illness (antecedent biomarkers), aid in identifying disease (diagnostic biomarkers), or predict future disease course, including response to therapy (prognostic biomarkers).

[0055] “Standard expression” is a quantitative or qualitative measurement for comparison. It is based on a statistically appropriate number of normal samples and is created to use as a basis of comparison when performing diagnostic assays, running clinical trials, or following patient treatment profiles.

[0056] “Animal” as used herein may be defined to include human, domestic (e.g., cats, dogs, etc.), agricultural (e.g., cows, horses, sheep, etc.) or test species (e.g., mouse, rat, rabbit, etc.).

[0057] A “PRSS23 polynucleotide”, within the meaning of the invention, shall be understood as being a nucleic acid molecule selected from a group consisting of

[0058] (i) nucleic acid molecules encoding a polypeptide comprising the amino acid sequence of SEQ ID NO: 3 or SEQ ID NO: 4,

[0059] (ii) nucleic acid molecules comprising the sequence of SEQ ID NO: 1 or SEQ ID NO: 2,

[0060] (iii) nucleic acid molecules having the sequence of SEQ ID NO: 1 or SEQ ID NO: 2,

[0061] (iv) nucleic acid molecules the complementary strand of which hybridizes under stringent conditions to a nucleic acid molecule of (i), (ii), or (iii),

[0062] (v) nucleic acid molecules the sequence of which differs from the sequence of a nucleic acid molecule of (iii) due to the degeneracy of the genetic code,

[0063] (vi) nucleic acid molecules which have a sequence identity of at least 80%, 85%, 90%, 95%, 98% or 99%; and

[0064] (vii) wherein the polypeptide encoded by said nucleic acid molecules of (i)-(vi) have PRSS23 activity

[0065] A “PRSS23 polypeptide”, within the meaning of the invention, shall be understood as being a polypeptide selected from a group consisting of

[0066] (i) polypeptides having the sequence of SEQ ID NO: 3 or 4,

[0067] (ii) polypeptides comprising the sequence of SEQ ID NO: 3 or 4,

[0068] (iii) polypeptides encoded by PRSS23 polynucleotides; and

[0069] (iv) polypeptides which show at least 99%, 98%, 95%, 90%, or 80% identity with a polypeptide of (i), (ii), or (iii);

wherein said polypeptide has PRSS23 activity.

[0070] The nucleotide sequences encoding a PRSS23 (or their complement) have numerous applications in techniques known to those skilled in the art of molecular biology. These techniques include use as hybridization probes, use in the construction of oligomers for PCR, use for chromosome and gene mapping, use in the recombinant production of PRSS23, and use in generation of antisense DNA or RNA, their chemical analogs and the like. Uses of nucleotides encoding a PRSS23 disclosed herein are exemplary of known techniques and are not intended to limit their use in any technique known to a person of ordinary skill in the art. Furthermore, the nucleotide sequences disclosed herein may be used in molecular biology techniques that have not yet been developed, provided the new techniques rely on properties of nucleotide sequences that are currently known, e.g., the triplet genetic code, specific base pair interactions, etc.

[0071] It will be appreciated by those skilled in the art that as a result of the degeneracy of the genetic code, a multitude of PRSS23— encoding nucleotide sequences may be produced. Some of these will only bear minimal homology to the nucleotide sequence of the known and naturally occurring PRSS23. The invention has specifically contemplated each and every possible variation of nucleotide sequence that could be made by selecting combinations based on possible codon choices. These combinations are made in accordance with the standard triplet genetic code as applied to the nucle-

otide sequence of naturally occurring PRSS23, and all such variations are to be considered as being specifically disclosed.

[0072] Although the nucleotide sequences which encode a PRSS23, its derivatives or its variants are preferably capable of hybridizing to the nucleotide sequence of the naturally occurring PRSS23 polynucleotide under stringent conditions, it may be advantageous to produce nucleotide sequences encoding PRSS23 polypeptides or its derivatives possessing a substantially different codon usage. Codons can be selected to increase the rate at which expression of the peptide occurs in a particular prokaryotic or eukaryotic expression host in accordance with the frequency with which particular codons are utilized by the host. Other reasons for substantially altering the nucleotide sequence encoding a PRSS23 polypeptide and/or its derivatives without altering the encoded amino acid sequence include the production of RNA transcripts having more desirable properties, such as a greater half-life, than transcripts produced from the naturally occurring sequence.

[0073] Nucleotide sequences encoding a PRSS23 polypeptide may be joined to a variety of other nucleotide sequences by means of well established recombinant DNA techniques. Useful nucleotide sequences for joining to PRSS23 polynucleotides include an assortment of cloning vectors such as plasmids, cosmids, lambda phage derivatives, phagemids, and the like. Vectors of interest include expression vectors, replication vectors, probe generation vectors, sequencing vectors, etc. In general, vectors of interest may contain an origin of replication functional in at least one organism, convenient restriction endonuclease sensitive sites, and selectable markers for one or more host cell systems.

[0074] Another aspect of the subject invention is to provide for PRSS23-specific hybridization probes capable of hybridizing with naturally occurring nucleotide sequences encoding PRSS23. Such probes may also be used for the detection of similar protease encoding sequences and should preferably show at least 40% nucleotide identity to PRSS23 polynucleotides. The hybridization probes of the subject invention may be derived from the nucleotide sequence presented as SEQ ID NO: 1 or from genomic sequences including promoter, enhancers or introns of the native gene. Hybridization probes may be labelled by a variety of reporter molecules using techniques well known in the art.

[0075] It will be recognized that many deletional or mutational analogs of PRSS23 polynucleotides will be effective hybridization probes for PRSS23 polynucleotides. Accordingly, the invention relates to nucleic acid sequences that hybridize with such PRSS23 encoding nucleic acid sequences under stringent conditions.

[0076] “Stringent conditions” refers to conditions that allow for the hybridization of substantially related nucleic acid sequences. For instance, such conditions will generally allow hybridization of sequence with at least about 85% sequence identity, preferably with at least about 90% sequence identity, more preferably with at least about 95% sequence identity. Hybridization conditions and probes can be adjusted in well-characterized ways to achieve selective hybridization of human-derived probes. Stringent conditions, within the meaning of the invention are 68° C. in a buffer containing 0.2xSSC (1x standard saline-citrate=150 mM NaCl, 15 mM Trisodiumcitrat) [Sambrook et al., (1989)].

[0077] Nucleic acid molecules that will hybridize to PRSS23 polynucleotides under stringent conditions can be identified functionally. Without limitation, examples of the

uses for hybridization probes include: histochemical uses such as identifying tissues that express PRSS23; measuring mRNA levels, for instance to identify a sample's tissue type or to identify cells that express abnormal levels of PRSS23; and detecting polymorphisms of PRSS23.

[0078] PCR provides additional uses for oligonucleotides based upon the nucleotide sequence which encodes PRSS23. Such probes used in PCR may be of recombinant origin, chemically synthesized, or a mixture of both. Oligomers may comprise discrete nucleotide sequences employed under optimized conditions for identification of PRSS23 in specific tissues or diagnostic use. The same two oligomers, a nested set of oligomers, or even a degenerate pool of oligomers may be employed under less stringent conditions for identification of closely related DNAs or RNAs.

[0079] Rules for designing polymerase chain reaction (PCR) primers are now established, as reviewed by PCR Protocols. Degenerate primers, i.e., preparations of primers that are heterogeneous at given sequence locations, can be designed to amplify nucleic acid sequences that are highly homologous to, but not identical with PRSS23. Strategies are now available that allow for only one of the primers to be required to specifically hybridize with a known sequence. For example, appropriate nucleic acid primers can be ligated to the nucleic acid sought to be amplified to provide the hybridization partner for one of the primers. In this way, only one of the primers need be based on the sequence of the nucleic acid sought to be amplified.

[0080] PCR methods for amplifying nucleic acid will utilize at least two primers. One of these primers will be capable of hybridizing to a first strand of the nucleic acid to be amplified and of priming enzyme-driven nucleic acid synthesis in a first direction. The other will be capable of hybridizing the reciprocal sequence of the first strand (if the sequence to be amplified is single stranded, this sequence will initially be hypothetical, but will be synthesized in the first amplification cycle) and of priming nucleic acid synthesis from that strand in the direction opposite the first direction and towards the site of hybridization for the first primer. Conditions for conducting such amplifications, particularly under preferred stringent hybridization conditions, are well known.

[0081] Other means of producing specific hybridization probes for PRSS23 include the cloning of nucleic acid sequences encoding PRSS23 or PRSS23 derivatives into vectors for the production of mRNA probes. Such vectors are known in the art, are commercially available and may be used to synthesize RNA probes in vitro by means of the addition of the appropriate RNA polymerase as T7 or SP6 RNA polymerase and the appropriate reporter molecules.

[0082] It is possible to produce a DNA sequence, or portions thereof, entirely by synthetic chemistry. After synthesis, the nucleic acid sequence can be inserted into any of the many available DNA vectors and their respective host cells using techniques which are well known in the art. Moreover, synthetic chemistry may be used to introduce mutations into the nucleotide sequence. Alternately, a portion of sequence in which a mutation is desired can be synthesized and recombined with longer portion of an existing genomic or recombinant sequence.

[0083] PRSS23 polynucleotides may be used to produce a purified oligo- or polypeptide using well known methods of recombinant DNA technology. The oligopeptide may be expressed in a variety of host cells, either prokaryotic or eukaryotic. Host cells may be from the same species from

which the nucleotide sequence was derived or from a different species. Advantages of producing an oligo-nucleotide by recombinant DNA technology include obtaining adequate amounts of the protein for purification and the availability of simplified purification procedures.

Quantitative Determinations of Nucleic Acids

[0084] An important step in the molecular genetic analysis of human disease is often the enumeration of the copy number of a nucleus acid or the relative expression of a gene in particular tissues.

[0085] Several different approaches are currently available to make quantitative determinations of nucleic acids. Chromosome-based techniques, such as comparative genomic hybridization (CGH) and fluorescent in situ hybridization (FISH) facilitate efforts to cytogenetically localize genomic regions that are altered in tumor cells. Regions of genomic alteration can be narrowed further using loss of heterozygosity analysis (LOH), in which disease DNA is analyzed and compared with normal DNA for the loss of a heterozygous polymorphic marker. The first experiments used restriction fragment length polymorphisms (RFLPs) [Johnson, (1989)], or hypervariable mini-satellite DNA [Barnes, 2000]. In recent years LOH has been performed primarily using PCR amplification of microsatellite markers and electrophoresis of the radio labelled [Jeffreys, (1985)] or fluorescently labelled PCR products [Weber, (1990)] and compared between paired normal and disease DNAs.

[0086] A number of other methods have also been developed to quantify nucleic acids [Gergen, (1992)]. More recently, PCR and RT-PCR methods have been developed which are capable of measuring the amount of a nucleic acid in a sample. One approach, for example, measures PCR product quantity in the log phase of the reaction before the formation of reaction products plateaus [Thomas, (1980)].

[0087] A gene sequence contained in all samples at relatively constant quantity is typically utilized for sample amplification efficiency normalization. This approach, however, suffers from several drawbacks. The method requires that each sample has equal input amounts of the nucleic acid and that the amplification efficiency between samples is identical until the time of analysis. Furthermore, it is difficult using the conventional methods of PCR quantitation such as gel electrophoresis or plate capture hybridization to determine that all samples are in fact analyzed during the log phase of the reaction as required by the method.

[0088] Another method called quantitative competitive (QC)-PCR, as the name implies, relies on the inclusion of an internal control competitor in each reaction [Piatak, (1993), BioTechniques]. The efficiency of each reaction is normalized to the internal competitor. A known amount of internal competitor is typically added to each sample. The unknown target PCR product is compared with the known competitor PCR product to obtain relative quantitation. A difficulty with this general approach lies in developing an internal control that amplifies with the same efficiency than the target molecule.

5' Fluorogenic Nuclease Assays

[0089] Fluorogenic nuclease assays are a real time quantitation method that uses a probe to monitor formation of amplification product. The basis for this method of monitoring the formation of amplification product is to measure con-

tinuously PCR product accumulation using a dual-labelled fluorogenic oligonucleotide probe, an approach frequently referred to in the literature simply as the "TaqMan method" [Piatak, (1993), Science; Heid, (1996); Gibson, (1996); Holland, (1991)].

[0090] The probe used in such assays is typically a short (about 20-25 bases) oligonucleotide that is labeled with two different fluorescent dyes. The 5' terminus of the probe is attached to a reporter dye and the 3' terminus is attached to a quenching dye, although the dyes could be attached at other locations on the probe as well. The probe is designed to have at least substantial sequence complementarity with the probe binding site. Upstream and downstream PCR primers which bind to flanking regions of the locus are added to the reaction mixture. When the probe is intact, energy transfer between the two fluorophors occurs and the quencher quenches emission from the reporter. During the extension phase of PCR, the probe is cleaved by the 5' nuclease activity of a nucleic acid polymerase such as Taq polymerase, thereby releasing the reporter from the oligonucleotide-quencher and resulting in an increase of reporter emission intensity which can be measured by an appropriate detector.

[0091] One detector which is specifically adapted for measuring fluorescence emissions such as those created during a fluorogenic assay is the ABI 7700 or 4700 HT manufactured by Applied Biosystems, Inc. in Foster City, Calif. The ABI 7700 uses fiber optics connected with each well in a 96- or 384 well PCR tube arrangement. The instrument includes a laser for exciting the labels and is capable of measuring the fluorescence spectra intensity from each tube with continuous monitoring during PCR amplification. Each tube is re-examined every 8.5 seconds.

[0092] Computer software provided with the instrument is capable of recording the fluorescence intensity of reporter and quencher over the course of the amplification. The recorded values will then be used to calculate the increase in normalized reporter emission intensity on a continuous basis. The increase in emission intensity is plotted versus time, i.e., the number of amplification cycles, to produce a continuous measure of amplification. To quantify the locus in each amplification reaction, the amplification plot is examined at a point during the log phase of product accumulation. This is accomplished by assigning a fluorescence threshold intensity above background and determining the point at which each amplification plot crosses the threshold (defined as the threshold cycle number or Ct). Differences in threshold cycle number are used to quantify the relative amount of PCR target contained within each tube. Assuming that each reaction functions at 100% PCR efficiency, a difference of one Ct represents a two-fold difference in the amount of starting template. The fluorescence value can be used in conjunction with a standard curve to determine the amount of amplification product present.

Non-Probe-Based Detection Methods

[0093] A variety of options are available for measuring the amplification products as they are formed. One method utilizes labels, such as dyes, which only bind to double stranded DNA. In this type of approach, amplification product (which is double stranded) binds dye molecules in solution to form a complex. With the appropriate dyes, it is possible to distinguish between dye molecules free in solution and dye molecules bound to amplification product. For example, certain dyes fluoresce only when bound to amplification product.

Examples of dyes which can be used in methods of this general type include, but are not limited to, Syber Green™ and Pico Green from Molecular Probes, Inc. of Eugene, Oreg., ethidium bromide, propidium iodide, chromomycin, acridine orange, Hoechst 33258, Toto-1, Yoyo-1, DAPI (4',6'-diamidino-2-phenylindole hydrochloride).

[0094] Another real time detection technique measures alteration in energy fluorescence energy transfer between fluorophors conjugated with PCR primers [Livak, (1995)].

Probe-Based Detection Methods

[0095] These detection methods involve some alteration to the structure or conformation of a probe hybridized to the locus between the amplification primer pair. In some instances, the alteration is caused by the template-dependent extension catalyzed by a nucleic acid polymerase during the amplification process. The alteration generates a detectable signal which is an indirect measure of the amount of amplification product formed.

[0096] For example, some methods involve the degradation or digestion of the probe during the extension reaction. These methods are a consequence of the 5'-3' nuclease activity associated with some nucleic acid polymerases. Polymerases having this activity cleave mononucleotides or small oligonucleotides from an oligonucleotide probe annealed to its complementary sequence located within the locus.

[0097] The 3' end of the upstream primer provides the initial binding site for the nucleic acid polymerase. As the polymerase catalyzes extension of the upstream primer and encounters the bound probe, the nucleic acid polymerase displaces a portion of the 5' end of the probe and through its nuclease activity cleaves mononucleotides or oligonucleotides from the probe.

[0098] The upstream primer and the probe can be designed such that they anneal to the complementary strand in close proximity to one another. In fact, the 3' end of the upstream primer and the 5' end of the probe may abut one another. In this situation, extension of the upstream primer is not necessary in order for the nucleic acid polymerase to begin cleaving the probe. In the case in which intervening nucleotides separate the upstream primer and the probe, extension of the primer is necessary before the nucleic acid polymerase encounters the 5' end of the probe. Once contact occurs and polymerization continues, the 5'-3' exonuclease activity of the nucleic acid polymerase begins cleaving mononucleotides or oligonucleotides from the 5' end of the probe. Digestion of the probe continues until the remaining portion of the probe dissociates from the complementary strand.

[0099] In solution, the two end sections can hybridize with each other to form a hairpin loop. In this conformation, the reporter and quencher dye are in sufficiently close proximity that fluorescence from the reporter dye is effectively quenched by the quencher dye. Hybridized probe, in contrast, results in a linearized conformation in which the extent of quenching is decreased. Thus, by monitoring emission changes for the two dyes, it is possible to indirectly monitor the formation of amplification product.

Probes

[0100] The labeled probe is selected so that its sequence is substantially complementary to a segment of the test locus or a reference locus. As indicated above, the nucleic acid site to

which the probe binds should be located between the primer binding sites for the upstream and downstream amplification primers.

Primers

[0101] The primers used in the amplification are selected so as to be capable of hybridizing to sequences at flanking regions of the locus being amplified. The primers are chosen to have at least substantial complementarity with the different strands of the nucleic acid being amplified. When a probe is utilized to detect the formation of amplification products, the primers are selected in such that they flank the probe, i.e. are located upstream and downstream of the probe.

[0102] The primer must have sufficient length so that it is capable of priming the synthesis of extension products in the presence of an agent for polymerization. The length and composition of the primer depends on many parameters, including, for example, the temperature at which the annealing reaction is conducted, proximity of the probe binding site to that of the primer, relative concentrations of the primer and probe and the particular nucleic acid composition of the probe. Typically the primer includes 15-30 nucleotides. However, the length of the primer may be more or less depending on the complexity of the primer binding site and the factors listed above.

Labels for Probes and Primers

[0103] The labels used for labeling the probes or primers of the current invention and which can provide the signal corresponding to the quantity of amplification product can take a variety of forms. As indicated above with regard to the 5' fluorogenic nuclease method, a fluorescent signal is one signal which can be measured. However, measurements may also be made, for example, by monitoring radioactivity, colorimetry, absorption, magnetic parameters, or enzymatic activity. Thus, labels which can be employed include, but are not limited to, fluorophores, chromophores, radioactive isotopes, electron dense reagents, enzymes, and ligands having specific binding partners (e.g., biotin-avidin).

[0104] Monitoring changes in fluorescence is a particularly useful way to monitor the accumulation of amplification products. A number of labels useful for attachment to probes or primers are commercially available including fluorescein and various fluorescein derivatives such as FAM, HEX, TET and JOE (all which are available from Applied Biosystems, Foster City, Calif.); lucifer yellow, and coumarin derivatives.

[0105] Labels may be attached to the probe or primer using a variety of techniques and can be attached at the 5' end, and/or the 3' end and/or at an internal nucleotide. The label can also be attached to spacer arms of various sizes which are attached to the probe or primer. These spacer arms are useful for obtaining a desired distance between multiple labels attached to the probe or primer.

[0106] In some instances, a single label may be utilized; whereas, in other instances, such as with the 5' fluorogenic nuclease assays for example, two or more labels are attached to the probe. In cases wherein the probe includes multiple labels, it is generally advisable to maintain spacing between the labels which is sufficient to permit separation of the labels during digestion of the probe through the 5'-3' nuclease activity of the nucleic acid polymerase.

Microarray

[0107] Nucleic acid arrays that have been used in the present invention are those that are commercially available

from Affymetrix (Santa Clara, Calif.) under the brand name GeneChip Human Genome U133 Plus 2.0 Array.® or Rat Genome U230 plus 2.0 Array respectively which represents the complete coverage of the Human Genome U133 Set plus 9921 probe sets representing approximately 6,500 new genes (with a total of approximately 56 000 transcripts) or the Rat Genome respectively. Affymetrix (Santa Clara, Calif.) GeneChip technology platform which consists of high-density microarrays and tools to help process and analyze those arrays, including standardized assays and reagents, instrumentation, and data management and analysis tools.

[0108] GeneChip microarrays consist of small DNA fragments (referred to as probes), chemically synthesized at specific locations on a coated quartz surface. By extracting and labeling nucleic acids from experimental samples, and then hybridizing those prepared samples to the array, the amount of label can be monitored enabling a measurement of gene regulation

[0109] The GeneChip human genome arrays include a set of human maintenance genes to facilitate the normalization and scaling of array experiments and to perform data comparison. This set of normalization genes shows consistent levels of expression over a diverse set of tissues.

Patients Exhibiting Symptoms of Disease

[0110] A number of diseases are associated with changes in the copy number of a certain gene. For patients having symptoms of a disease, the real-time PCR method can be used to determine if the patient has copy number alterations which are known to be linked with diseases that are associated with the symptoms the patient has.

PRSS23 Expression

PRSS23 Fusion Proteins

[0111] Fusion proteins are useful for generating antibodies against PRSS23 polypeptides and for use in various assay systems. For example, fusion proteins can be used to identify proteins which interact with portions of PRSS23 polypeptides. Protein affinity chromatography or library-based assays for protein-protein interactions, such as the yeast two-hybrid or phage display systems, can be used for this purpose. Such methods are well known in the art and also can be used as drug screens.

[0112] A PRSS23 fusion protein comprises two polypeptide segments fused together by means of a peptide bond. The first polypeptide segment can comprise at least 54, 75, 100, 125, 139, 150, 175, 200, 225, 250, 275, 300, 325 or 350 contiguous amino acids of SEQ ID NO: 3 or 4 or of a variant, such as those described above. The first polypeptide segment also can comprise full-length PRSS23.

[0113] The second polypeptide segment can be a full-length protein or a protein fragment. Proteins commonly used in fusion protein construction include, but are not limited to p galactosidase, β -glucuronidase, green fluorescent protein (GFP), autofluorescent proteins, including blue fluorescent protein (BFP), glutathione-S-transferase (GST), luciferase, horseradish peroxidase (HRP), and chloramphenicol acetyltransferase (CAT). Additionally, epitope tags are used in fusion protein constructions, including histidine (His) tags, FLAG tags, influenza hemagglutinin (HA) tags, Myc tags, VSV-G tags, and thioredoxin (Trx) tags. Other fusion constructions can include maltose binding protein (MBP), S-tag, Lex a DNA binding domain (DBD) fusions, GAL4 DNA

binding domain fusions, and herpes simplex virus (HSV) BP16 protein fusions. A fusion protein also can be engineered to contain a cleavage site located adjacent to the PRSS23.

Preparation of Polynucleotides

[0114] A naturally occurring PRSS23 polynucleotide can be isolated free of other cellular components such as membrane components, proteins, and lipids. Polynucleotides can be made by a cell and isolated using standard nucleic acid purification techniques, or synthesized using an amplification technique, such as the polymerase chain reaction (PCR), or by using an automatic synthesizer. Methods for isolating polynucleotides are routine and are known in the art. Any such technique for obtaining a polynucleotide can be used to obtain isolated PRSS23 polynucleotides. For example, restriction enzymes and probes can be used to isolate polynucleotide fragments which comprise PRSS23 nucleotide sequences. Isolated polynucleotides are in preparations which are free or at least 70, 80, or 90% free of other molecules.

[0115] PRSS23 cDNA molecules can be made with standard molecular biology techniques, using PRSS23 mRNA as a template. PRSS23 cDNA molecules can thereafter be replicated using molecular biology techniques known in the art. An amplification technique, such as PCR, can be used to obtain additional copies of polynucleotides of the invention, using either human genomic DNA or cDNA as a template.

[0116] Alternatively, synthetic chemistry techniques can be used to synthesize PRSS23 polynucleotides. The degeneracy of the genetic code allows alternate nucleotide sequences to be synthesized which will encode PRSS23 having, for example, an amino acid sequence shown in SEQ ID NO: 2 or a biologically active variant thereof.

Extending Polynucleotides

[0117] Various PCR-based methods can be used to extend nucleic acid sequences encoding human PRSS23, for example to detect upstream sequences of PRSS23 gene such as promoters and regulatory elements. For example, restriction-site PCR uses universal primers to retrieve unknown sequence adjacent to a known locus. Genomic DNA is first amplified in the presence of a primer to a linker sequence and a primer specific to the known region. The amplified sequences are then subjected to a second round of PCR with the same linker primer and another specific primer internal to the first one. Products of each round of PCR are transcribed with an appropriate RNA polymerase and sequenced using reverse transcriptase.

[0118] Inverse PCR also can be used to amplify or extend sequences using divergent primers based on a known region. Primers can be designed using commercially available software, such as OLIGO 4.06 Primer Analysis software (National Biosciences Inc., Plymouth, Minn.), to be 22-30 nucleotides in length, to have a GC content of 50% or more, and to anneal to the target sequence at temperatures about 68-72° C. The method uses several restriction enzymes to generate a suitable fragment in the known region of a gene. The fragment is then circularized by intramolecular ligation and used as a PCR template.

[0119] Another method which can be used is capture PCR, which involves PCR amplification of DNA fragments adjacent to a known sequence in human and yeast artificial chromosome DNA. In this method, multiple restriction enzyme

digestions and ligations also can be used to place an engineered double-stranded sequence into an unknown fragment of the DNA molecule before performing PCR.

[0120] When screening for full-length cDNAs, it is preferable to use libraries that have been size-selected to include larger cDNAs. Randomly-primed libraries are preferable, in that they will contain more sequences which contain the 5' regions of genes. Use of a randomly primed library may be especially preferable for situations in which an oligo d(T) library does not yield a full-length cDNA. Genomic libraries can be useful for extension of sequence into 5' non-transcribed regulatory regions.

[0121] Commercially available capillary electrophoresis systems can be used to analyze the size or confirm the nucleotide sequence of PCR or sequencing products. For example, capillary sequencing can employ flowable polymers for electrophoretic separation, four different fluorescent dyes (one for each nucleotide) which are laser activated, and detection of the emitted wavelengths by a charge coupled device camera. Output/light intensity can be converted to electrical signal using appropriate equipment and software (e.g., GENOTYPER and Sequence NAVIGATOR, Perkin Elmer), and the entire process from loading of samples to computer analysis and electronic data display can be computer controlled. Capillary electrophoresis is especially preferable for the sequencing of small pieces of DNA which might be present in limited amounts in a particular sample.

Obtaining Polypeptides

[0122] PRSS23 can be obtained, for example, by purification from human cells, by expression of PRSS23 polynucleotides, or by direct chemical synthesis.

Protein Purification

[0123] PRSS23 can be purified from any human cell which expresses the enzyme, including those which have been transfected with expression constructs which express PRSS23. A purified PRSS23 is separated from other compounds which normally associate with PRSS23 in the cell, such as certain proteins, carbohydrates, or lipids, using methods well-known in the art. Such methods include, but are not limited to, size exclusion chromatography, ammonium sulfate fractionation, ion exchange chromatography, affinity chromatography, and preparative gel electrophoresis.

Expression of PRSS23 Polynucleotides

[0124] To express PRSS23, PRSS23 polynucleotides can be inserted into an expression vector which contains the necessary elements for the transcription and translation of the inserted coding sequence. Methods which are well known to those skilled in the art can be used to construct expression vectors containing sequences encoding PRSS23 and appropriate transcriptional and translational control elements. These methods include in vitro recombinant DNA techniques, synthetic techniques, and in vivo genetic recombination.

[0125] A variety of expression vector/host systems can be utilized to contain and express sequences encoding PRSS23. These include, but are not limited to, microorganisms, such as bacteria transformed with recombinant bacteriophage, plasmid, or cosmid DNA expression vectors; yeast transformed with yeast expression vectors, insect cell systems infected with virus expression vectors (e.g., baculovirus), plant cell

systems transformed with virus expression vectors (e.g., cauliflower mosaic virus, CaMV; tobacco mosaic virus, TMV) or with bacterial expression vectors (e.g., Ti or pBR322 plasmids), or animal cell systems.

[0126] The control elements or regulatory sequences are those non-translated regions of the vector-enhancers, promoters, 5' and 3' untranslated regions—which interact with host cellular proteins to carry out transcription and translation. Such elements can vary in their strength and specificity. Depending on the vector system and host utilized, any number of suitable transcription and translation elements, including constitutive and inducible promoters, can be used. For example, when cloning in bacterial systems, inducible promoters such as the hybrid lacZ promoter of the BLUE-SCRIPT phagemid (Stratagene, LaJolla, Calif.) or pSPORT1 plasmid (Life Technologies) and the like can be used. The baculovirus polyhedrin promoter can be used in insect cells. Promoters or enhancers derived from the genomes of plant cells (e.g., heat shock, RUBISCO, and storage protein genes) or from plant viruses (e.g., viral promoters or leader sequences) can be cloned into the vector. In mammalian cell systems, promoters from mammalian genes or from mammalian viruses are preferable. If it is necessary to generate a cell line that contains multiple copies of a nucleotide sequence encoding PRSS23, vectors based on SV40 or EBV can be used with an appropriate selectable marker.

Bacterial and Yeast Expression Systems

[0127] In bacterial systems, a number of expression vectors can be selected. For example, when a large quantity of PRSS23 is needed for the induction of antibodies, vectors which direct high level expression of fusion proteins that are readily purified can be used. Such vectors include, but are not limited to, multifunctional *E. coli* cloning and expression vectors such as BLUE-SCRIPT (Stratagene). In a BLUE-SCRIPT vector, a sequence encoding PRSS23 can be ligated into the vector in frame with sequences for the amino-terminal Met and the subsequent 7 residues of β -galactosidase so that a hybrid protein is produced. pIN vectors or pGEX vectors (Promega, Madison, Wis.) also can be used to express foreign polypeptides as fusion proteins with glutathione S-transferase (GST). In general, such fusion proteins are soluble and can easily be purified from lysed cells by adsorption to glutathione-agarose beads followed by elution in the presence of free glutathione. Proteins made in such systems can be designed to include heparin, thrombin, or factor Xa protease cleavage sites so that the cloned polypeptide of interest can be released from the GST moiety at will.

Plant and Insect Expression Systems

[0128] If plant expression vectors are used, the expression of sequences encoding PRSS23 can be driven by any of a number of promoters. For example, viral promoters such as the 35S and 19S promoters of CaMV can be used alone or in combination with the omega leader sequence from TMV. Alternatively, plant promoters such as the small subunit of RUBISCO or heat shock promoters can be used. These constructs can be introduced into plant cells by direct DNA transformation or by pathogen-mediated transfection.

[0129] An insect system also can be used to express PRSS23. For example, in one such system *Autographa californica* nuclear polyhedrosis virus (AcNPV) is used as a vector to express foreign genes in *Spodoptera frugiperda*

cells or in *Trichoplusia larvae*. Sequences encoding PRSS23 can be cloned into a non-essential region of the virus, such as the polyhedrin gene, and placed under control of the polyhedrin promoter. Successful insertion of PRSS23 will render the polyhedrin gene inactive and produce recombinant virus lacking coat protein. The recombinant viruses can then be used to infect *S. frugiperda* cells or *Trichoplusia larvae* in which PRSS23 can be expressed.

Mammalian Expression Systems

[0130] A number of viral-based expression systems can be used to express PRSS23 in mammalian host cells. For example, if an adenovirus is used as an expression vector, sequences encoding PRSS23 can be ligated into an adenovirus transcription/translation complex comprising the late promoter and tripartite leader sequence. Insertion in a non-essential E1 or E3 region of the viral genome can be used to obtain a viable virus which is capable of expressing PRSS23 in infected host cells [Engelhard, (1994)]. If desired, transcription enhancers, such as the Rous sarcoma virus (RSV) enhancer, can be used to increase expression in mammalian host cells.

[0131] Human artificial chromosomes (HACs) also can be used to deliver larger fragments of DNA than can be contained and expressed in a plasmid. HACs of 6M to 10M are constructed and delivered to cells via conventional delivery methods (e.g., liposomes, polycationic amino polymers, or vesicles). Specific initiation signals also can be used to achieve more efficient translation of sequences encoding PRSS23. Such signals include the ATG initiation codon and adjacent sequences. In cases where sequences encoding PRSS23, its initiation codon, and upstream sequences are inserted into the appropriate expression vector, no additional transcriptional or translational control signals may be needed. However, in cases where only coding sequence, or a fragment thereof, is inserted, exogenous translational control signals (including the ATG initiation codon) should be provided. The initiation codon should be in the correct reading frame to ensure translation of the entire insert. Exogenous translational elements and initiation codons can be of various origins, both natural and synthetic.

Host Cells

[0132] A host cell strain can be chosen for its ability to modulate the expression of the inserted sequences or to process the expressed PRSS23 in the desired fashion. Such modifications of the polypeptide include, but are not limited to, acetylation, carboxylation, glycosylation, phosphorylation, lipidation, and acylation. Post-translational processing which cleaves a “prepro” form of the polypeptide also can be used to facilitate correct insertion, folding and/or function. Different host cells which have specific cellular machinery and characteristic mechanisms for post-translational activities (e.g., CHO, HeLa, MDCK, HEK293, and WI38), are available from the American Type Culture Collection (ATCC; 10801 University Boulevard, Manassas, Va. 20110-2209) and can be chosen to ensure the correct modification and processing of the foreign protein.

[0133] Stable expression is preferred for long-term, high-yield production of recombinant proteins. For example, cell lines which stably express PRSS23 can be transformed using expression vectors which can contain viral origins of replication and/or endogenous expression elements and a selectable

marker gene on the same or on a separate vector. Following the introduction of the vector, cells can be allowed to grow for 1-2 days in an enriched medium before they are switched to a selective medium. The purpose of the selectable marker is to confer resistance to selection, and its presence allows growth and recovery of cells which successfully express the introduced PRSS23 sequences. Resistant clones of stably transformed cells can be proliferated using tissue culture techniques appropriate to the cell type. Any number of selection systems can be used to recover transformed cell lines. These include, but are not limited to, the herpes simplex virus thymidine kinase [Logan, (1984)] and adenine phosphoribosyltransferase [Wigler, (1977)] genes which can be employed in tk⁻ or apt⁻ cells, respectively. Also, antimetabolite, antibiotic, or herbicide resistance can be used as the basis for selection. For example, dhfr confers resistance to methotrexate [Lowy, (1980)], npt confers resistance to the aminoglycosides, neomycin and G-418 [Wigler, (1980)], and als and pat confer resistance to chlorsulfuron and phosphinotricin acetyltransferase, respectively [Colbere-Garapin, 1981]. Additional selectable genes have been described. For example, trpB allows cells to utilize indole in place of tryptophan, or hisD, which allows cells to utilize histinol in place of histidine. Visible markers such as anthocyanins, β -glucuronidase and its substrate GUS, and luciferase and its substrate luciferin, can be used to identify transformants and to quantify the amount of transient or stable protein expression attributable to a specific vector system

Detecting Polypeptide Expression

[0134] Although the presence of marker gene expression suggests that a PRSS23 polynucleotide is also present, its presence and expression may need to be confirmed. For example, if a sequence encoding PRSS23 is inserted within a marker gene sequence, transformed cells containing sequences which encode PRSS23 can be identified by the absence of marker gene function. Alternatively, a marker gene can be placed in tandem with a sequence encoding PRSS23 under the control of a single promoter. Expression of the marker gene in response to induction or selection usually indicates expression of PRSS23 polynucleotide.

[0135] Alternatively, host cells which contain a PRSS23 polynucleotide and which express PRSS23 can be identified by a variety of procedures known to those of skill in the art. These procedures include, but are not limited to, DNA-DNA or DNA-RNA hybridizations and protein bioassay or immunoassay techniques which include membrane, solution, or chip-based technologies for the detection and/or quantification of nucleic acid or protein. For example, the presence of a polynucleotide sequence encoding PRSS23 can be detected by DNA-DNA or DNA-RNA hybridization or amplification using probes or fragments or fragments of polynucleotides encoding PRSS23. Nucleic acid amplification-based assays involve the use of oligonucleotides selected from sequences encoding PRSS23 to detect transformants which contain a PRSS23 poly-nucleotide.

[0136] A variety of protocols for detecting and measuring the expression of PRSS23, using either polyclonal or monoclonal antibodies specific for the polypeptide, are known in the art. Examples include enzyme-linked immunosorbent assay (ELISA), radioimmunoassay (RIA), and fluorescence activated cell sorting (FACS). A two-site, monoclonal-based immunoassay using monoclonal antibodies reactive to two

non-interfering epitopes on PRSS23 can be used, or a competitive binding assay can be employed.

[0137] A wide variety of labels and conjugation techniques are known by those skilled in the art and can be used in various nucleic acid and amino acid assays. Means for producing labeled hybridization or PCR probes for detecting sequences related to polynucleotides encoding PRSS23 include oligolabeling, nick translation, end-labeling, or PCR amplification using a labeled nucleotide. Alternatively, sequences encoding PRSS23 can be cloned into a vector for the production of an mRNA probe. Such vectors are known in the art, are commercially available, and can be used to synthesize RNA probes in vitro by addition of labeled nucleotides and an appropriate RNA polymerase such as T7, T3, or SP6. These procedures can be conducted using a variety of commercially available kits (Amersham Pharmacia Biotech, Promega, and US Biochemical). Suitable reporter molecules or labels which can be used for ease of detection include radionuclides, enzymes, and fluorescent, chemiluminescent, or chromogenic agents, as well as substrates, cofactors, inhibitors, magnetic particles, and the like.

Expression and Purification of Polypeptides

[0138] Host cells transformed with PRSS23 polynucleotides can be cultured under conditions suitable for the expression and recovery of the protein from cell culture. The polypeptide produced by a transformed cell can be secreted or contained intracellularly depending on the sequence and/or the vector used. As will be understood by those of skill in the art, expression vectors containing PRSS23 polynucleotides can be designed to contain signal sequences which direct secretion of soluble PRSS23 through a prokaryotic or eukaryotic cell membrane or which direct the membrane insertion of membrane-bound PRSS23.

[0139] As discussed above, other constructions can be used to join a sequence encoding PRSS23 to a nucleotide sequence encoding a polypeptide domain which will facilitate purification of soluble proteins. Such purification facilitating domains include, but are not limited to, metal chelating peptides such as histidine-tryptophan modules that allow purification on immobilized metals, protein A domains that allow purification on immobilized immunoglobulin, and the domain utilized in the FLAG extension/affinity purification system (Immunex Corp., Seattle, Wash.). Inclusion of cleavable linker sequences such as those specific for Factor XA or enterokinase (Invitrogen, San Diego, Calif.) between the purification domain and PRSS23 also can be used to facilitate purification. One such expression vector provides for expression of a fusion protein containing PRSS23 and 6 histidine residues preceding a thioredoxin or an enterokinase cleavage site. The histidine residues facilitate purification by IMAC (immobilized metal ion affinity chromatography) Maddox, (1983)], while the enterokinase cleavage site provides a means for purifying PRSS23 from the fusion protein [Porath, (1992)].

Chemical Synthesis

[0140] Sequences encoding PRSS23 can be synthesized, in whole or in part, using chemical methods well known in the art. Alternatively, PRSS23 itself can be produced using chemical methods to synthesize its amino acid sequence, such as by direct peptide synthesis using solid-phase techniques. Protein synthesis can either be performed using manual tech-

niques or by automation. Automated synthesis can be achieved, for example, using Applied Biosystems 431A Peptide Synthesizer (Perkin Elmer). Optionally, fragments of PRSS23 can be separately synthesized and combined using chemical methods to produce a full-length molecule.

[0141] The newly synthesized peptide can be substantially purified by preparative high performance liquid chromatography. The composition of a synthetic PRSS23 can be confirmed by amino acid analysis or sequencing. Additionally, any portion of the amino acid sequence of PRSS23 can be altered during direct synthesis and/or combined using chemical methods with sequences from other proteins to produce a variant polypeptide or a fusion protein.

Production of Altered Polypeptides

[0142] As will be understood by those of skill in the art, it may be advantageous to produce PRSS23 polynucleotides possessing non-naturally occurring codons. For example, codons preferred by a particular prokaryotic or eukaryotic host can be selected to increase the rate of protein expression or to produce an RNA transcript having desirable properties, such as a half-life which is longer than that of a transcript generated from the naturally occurring sequence.

[0143] The nucleotide sequences referred to herein can be engineered using methods generally known in the art to alter PRSS23 polynucleotides for a variety of reasons, including but not limited to, alterations which modify the cloning, processing, and/or expression of the polypeptide or mRNA product. DNA shuffling by random fragmentation and PCR reassembly of gene fragments and synthetic oligonucleotides can be used to engineer the nucleotide sequences. For example, site-directed mutagenesis can be used to insert new restriction sites, alter glycosylation patterns, change codon preference, produce splice variants, introduce mutations, and so forth.

PRSS23 Analogs

[0144] One general class of PRSS23 analogs are variants having an amino acid sequence that is a mutation of the amino acid sequence disclosed herein. Another general class of PRSS23 analogs is provided by anti-idiotypic antibodies, and fragments thereof, as described below. Moreover, recombinant antibodies comprising anti-idiotypic variable domains can be used as analogs (see, for example, [Monfardini et al., (1996)]). Since the variable domains of anti-idiotypic PRSS23 antibodies mimic PRSS23, these domains can provide PRSS23 enzymatic activity. Methods of producing anti-idiotypic catalytic antibodies are known to those of skill in the art [Joron et al., (1992), Friboulet et al. (1994), Avalle et al., (1998)].

[0145] Another approach to identifying PRSS23 analogs is provided by the use of combinatorial libraries. Methods for constructing and screening phage display and other combinatorial libraries are provided, for example, by [Kay et al., Phage Display of Peptides and Proteins (Academic Press 1996), U.S. Pat. No. 5,783,384, U.S. Pat. No. 5,747,334, and U.S. Pat. No. 5,723,323].

[0146] One illustrative in vitro use of PRSS23 and its analogs is the production of labeled peptides from a labeled protein substrate. Proteases can also be used in detergents and cleaning solutions. For example, serine proteases are used in solutions to clean and to disinfect contact lenses (see, for example, [U.S. Pat. No. 5,985,629]). Another use for a serine protease is in the formulation of vaccines (see, for example,

[U.S. Pat. No. 5,885,814]). Those of skill in the art can devise other uses for molecules having PRSS23 activity.

Antibodies

[0147] Any type of antibody known in the art can be generated to bind specifically to an epitope of PRSS23.

[0148] "Antibody" as used herein includes intact immunoglobulin molecules, as well as fragments thereof, such as Fab, F(ab')₂, and Fv, which are capable of binding an epitope of PRSS23. Typically, at least 6, 8, 10, or 12 contiguous amino acids are required to form an epitope. However, epitopes which involve non-contiguous amino acids may require more, e.g., at least 15, 25, or 50 amino acid. An antibody which specifically binds to an epitope of PRSS23 can be used therapeutically, as well as in immunochemical assays, such as Western blots, ELISAs, radioimmuno-assays, immunohistochemical assays, immunoprecipitations, or other immunochemical assays known in the art. Various immunoassays can be used to identify antibodies having the desired specificity. Numerous protocols for competitive binding or immunoradiometric assays are well known in the art. Such immunoassays typically involve the measurement of complex formation between an immunogen and an antibody which specifically binds to the PRSS23 immunogen.

[0149] Typically, an antibody which specifically binds to PRSS23 provides a detection signal at least 5-, 10-, or 20-fold higher than a detection signal provided with other proteins when used in an immunochemical assay. Preferably, antibodies which specifically bind to PRSS23 do not detect other proteins in immunochemical assays and can immunoprecipitate PRSS23 from solution.

[0150] PRSS23 can be used to immunize a mammal, such as a mouse, rat, rabbit, guinea pig, monkey, or human, to produce polyclonal antibodies. If desired, PRSS23 can be conjugated to a carrier protein, such as bovine serum albumin, thyroglobulin, and keyhole limpet hemocyanin. Depending on the host species, various adjuvants can be used to increase the immunological response. Such adjuvants include, but are not limited to, Freund's adjuvant, mineral gels (e.g., aluminum hydroxide), and surface active substances (e.g., lysolécithin, pluronic polyols, polyanions, peptides, oil emulsions, keyhole limpet hemocyanin, and dinitrophenol). Among adjuvants used in humans, BCG (*Bacillus Calmette-Guerin*) and *Corynebacterium parvum* are especially useful.

[0151] Monoclonal antibodies which specifically bind to PRSS23 can be prepared using any technique which provides for the production of antibody molecules by continuous cell lines in culture. These techniques include, but are not limited to, the hybridoma technique, the human B-cell hybridoma technique, and the EBV-hybridoma technique [Roberge, (1995)].

[0152] In addition, techniques developed for the production of "chimeric antibodies", the splicing of mouse antibody genes to human antibody genes to obtain a molecule with appropriate antigen specificity and biological activity, can be used. Monoclonal and other antibodies also can be "humanized" to prevent a patient from mounting an immune response against the antibody when it is used therapeutically. Such antibodies may be sufficiently similar in sequence to human antibodies to be used directly in therapy or may require alteration of a few key residues. Sequence differences between rodent antibodies and human sequences can be minimized by replacing residues which differ from those in the human sequences by site directed mutagenesis of individual residues

or by grating of entire complementarity determining regions. Antibodies which specifically bind to PRSS23 can contain antigen binding sites which are either partially or fully humanized, as disclosed in U.S. Pat. No. 5,565,332.

[0153] Alternatively, techniques described for the production of single chain antibodies can be adapted using methods known in the art to produce single chain antibodies which specifically bind to PRSS23. Antibodies with related specificity, but of distinct idiotypic composition, can be generated by chain shuffling from random combinatorial immunoglobulin libraries. Single-chain antibodies also can be constructed using a DNA amplification method, such as PCR, using hybridoma cDNA as a template. Single-chain antibodies can be mono- or bispecific, and can be bivalent or tetravalent. Construction of tetravalent, bispecific single-chain antibodies is taught. A nucleotide sequence encoding a single-chain antibody can be constructed using manual or automated nucleotide synthesis, cloned into an expression construct using standard recombinant DNA methods, and introduced into a cell to express the coding sequence, as described below. Alternatively, single-chain antibodies can be produced directly using, for example, filamentous phage technology.

[0154] Antibodies which specifically bind to PRSS23 also can be produced by inducing in vivo production in the lymphocyte population or by screening immunoglobulin libraries or panels of highly specific binding reagents. Other types of antibodies can be constructed and used therapeutically in methods of the invention. For example, chimeric antibodies can be constructed as disclosed in WO 93/03151. Binding proteins which are derived from immunoglobulins and which are multivalent and multispecific, such as the "diabodies" described in WO 94/13804, also can be prepared.

[0155] Antibodies according to the invention can be purified by methods well known in the art. For example, antibodies can be affinity purified by passage over a column to which PRSS23 is bound. The bound antibodies can then be eluted from the column using a buffer with a high salt concentration.

Antisense Oligonucleotides

[0156] Antisense oligonucleotides are nucleotide sequences which are complementary to a specific DNA or RNA sequence. Once introduced into a cell, the complementary nucleotides combine with natural sequences produced by the cell to form complexes and block either transcription or translation. Preferably, an antisense oligonucleotide is at least 11 nucleotides in length, but can be at least 12, 15, 20, 25, 30, 35, 40, 45, or 50 or more nucleotides long. Longer sequences also can be used. Antisense oligonucleotide molecules can be provided in a DNA construct and introduced into a cell as described above to decrease the level of PRSS23 gene products in the cell.

[0157] Antisense oligonucleotides can be deoxyribonucleotides, ribonucleotides, or a combination of both. Oligonucleotides can be synthesized manually or by an automated synthesizer, by covalently linking the 5' end of one nucleotide with the 3' end of another nucleotide with non-phosphodiester internucleotide linkages such as alkylphosphonates, phosphorothioates, phosphorodithioates, alkylphosphonothioates, alkylphosphonates, phosphoramidates, phosphate esters, carbamates, acetamidate, carboxymethyl esters, carbonates, and phosphate triesters.

[0158] Modifications of PRSS23 gene expression can be obtained by designing antisense oligonucleotides which will form duplexes to the control, 5', or regulatory regions of the

PRSS23 gene. Oligo-nucleotides derived from the transcription initiation site, e.g., between positions -10 and +10 from the start site, are preferred. Similarly, inhibition can be achieved using "triple helix" base-pairing methodology. Triple helix pairing is useful because it causes inhibition of the ability of the double helix to open sufficiently for the binding of polymerases, transcription factors, or chaperons. Therapeutic advances using triplex DNA have been described in the literature [Nicholls, (1993)]. An antisense oligonucleotide also can be designed to block translation of mRNA by preventing the transcript from binding to ribosomes.

[0159] Precise complementarity is not required for successful complex formation between an antisense oligonucleotide and the complementary sequence of a PRSS23 polynucleotide. Antisense oligo-nucleotides which comprise, for example, 2, 3, 4, or 5 or more stretches of contiguous nucleotides which are precisely complementary to a PRSS23 polynucleotide, each separated by a stretch of contiguous nucleotides which are not complementary to adjacent PRSS23 nucleotides, can provide sufficient targeting specificity for PRSS23 mRNA. Preferably, each stretch of complementary contiguous nucleotides is at least 4, 5, 6, 7, or 8 or more nucleotides in length. Non-complementary intervening sequences are preferably 1, 2, 3, or 4 nucleotides in length. One skilled in the art can easily use the calculated melting point of an antisense-sense pair to determine the degree of mismatching which will be tolerated between a particular antisense oligonucleotide and a particular PRSS23 polynucleotide sequence. Antisense oligonucleotides can be modified without affecting their ability to hybridize to a PRSS23 polynucleotide. These modifications can be internal or at one or both ends of the antisense molecule. For example, internucleoside phosphate linkages can be modified by adding cholesteryl or diamine moieties with varying numbers of carbon residues between the amino groups and terminal ribose. Modified bases and/or sugars, such as arabinose instead of ribose, or a 3',5'-substituted oligonucleotide in which the 3' hydroxyl group or the 5' phosphate group are substituted, also can be employed in a modified antisense oligonucleotide. These modified oligonucleotides can be prepared by methods well known in the art.

Ribozymes

[0160] Ribozymes are RNA molecules with catalytic activity [Uhlmann, (1987)]. Ribozymes can be used to inhibit gene function by cleaving an RNA sequence, as is known in the art. The mechanism of ribozyme action involves sequence-specific hybridization of the ribozyme molecule to complementary target RNA, followed by endonucleolytic cleavage. Examples include engineered hammerhead motif ribozyme molecules that can specifically and efficiently catalyze endonucleolytic cleavage of specific nucleotide sequences. The coding sequence of a PRSS23 polynucleotide can be used to generate ribozymes which will specifically bind to mRNA transcribed from a PRSS23 polynucleotide. Methods of designing and constructing ribozymes which can cleave other RNA molecules in trans in a highly sequence specific manner have been developed and described in the art. For example, the cleavage activity of ribozymes can be targeted to specific RNAs by engineering a discrete "hybridization" region into the ribozyme. The hybridization region contains a sequence complementary to the target RNA and thus specifically hybridizes with the target RNA.

[0161] Specific ribozyme cleavage sites within a PRSS23 RNA target can be identified by scanning the target molecule for ribozyme cleavage sites which include the following sequences: GUA, GUU, and GUC. Once identified, short RNA sequences of between 15 and 20 ribonucleotides corresponding to the region of the target RNA containing the cleavage site can be evaluated for secondary structural features which may render the target inoperable. Suitability of candidate PRSS23 RNA targets also can be evaluated by testing accessibility to hybridization with complementary oligonucleotides using ribonuclease protection assays. The nucleotide sequences shown in SEQ ID NO: 1 and its complement provide sources of suitable hybridization region sequences. Longer complementary sequences can be used to increase the affinity of the hybridization sequence for the target. The hybridizing and cleavage regions of the ribozyme can be integrally related such that upon hybridizing to the target RNA through the complementary regions, the catalytic region of the ribozyme can cleave the target.

[0162] Ribozymes can be introduced into cells as part of a DNA construct. Mechanical methods, such as microinjection, liposome-mediated transfection, electroporation, or calcium phosphate precipitation, can be used to introduce a ribozyme-containing DNA construct into cells in which it is desired to decrease PRSS23 expression. Alternatively, if it is desired that the cells stably retain the DNA construct, the construct can be supplied on a plasmid and maintained as a separate element or integrated into the genome of the cells, as is known in the art. A ribozyme-encoding DNA construct can include transcriptional regulatory elements, such as a promoter element, an enhancer or UAS element, and a transcriptional terminator signal, for controlling transcription of ribozymes in the cells (U.S. Pat. No. 5,641,673). Ribozymes also can be engineered to provide an additional level of regulation, so that destruction of mRNA occurs only when both a ribozyme and a target gene are induced in the cells.

Screening/Screening Assays

Regulators

[0163] Regulators as used herein, refer to compounds that affect the activity of PRSS23 in vivo and/or in vitro. Regulators can be agonists and antagonists of PRSS23 polypeptide and can be compounds that exert their effect on the PRSS23 activity via the enzymatic activity, expression, post-translational modifications or by other means. Agonists of PRSS23 are molecules which, when bound to PRSS23, increase or prolong the activity of PRSS23. Agonists of PRSS23 include proteins, nucleic acids, carbohydrates, small molecules, or any other molecule which activate PRSS23. Antagonists of PRSS23 are molecules which, when bound to PRSS23, decrease the amount or the duration of the activity of PRSS23. Antagonists include proteins, nucleic acids, carbohydrates, antibodies, small molecules, or any other molecule which decrease the activity of PRSS23.

[0164] The term "modulate", as it appears herein, refers to a change in the activity of PRSS23 poly-peptide. For example, modulation may cause an increase or a decrease in enzymatic activity, binding characteristics, or any other biological, functional, or immunological properties of PRSS23.

[0165] As used herein, the terms "specific binding" or "specifically binding" refer to that interaction between a protein or peptide and an agonist, an antibody, or an antagonist. The interaction is dependent upon the presence of a particular

structure of the protein recognized by the binding molecule (i.e., the antigenic determinant or epitope). For example, if an antibody is specific for epitope "A" the presence of a polypeptide containing the epitope A, or the presence of free unlabeled A, in a reaction containing free labeled A and the antibody will reduce the amount of labeled A that binds to the antibody.

[0166] The invention provides methods (also referred to herein as "screening assays") for identifying compounds which can be used for the treatment of diseases related to PRSS23. The methods entail the identification of candidate or test compounds or agents (e.g., peptides, peptidomimetics, small molecules or other molecules) which bind to PRSS23 and/or have a stimulatory or inhibitory effect on the biological activity of PRSS23 or its expression and then determining which of these compounds have an effect on symptoms or diseases related to PRSS23 in an in vivo assay.

[0167] Candidate or test compounds or agents which bind to PRSS23 and/or have a stimulatory or inhibitory effect on the activity or the expression of PRSS23 are identified either in assays that employ cells which express PRSS23 (cell-based assays) or in assays with isolated PRSS23 (cell-free assays). The various assays can employ a variety of variants of PRSS23 (e.g., full-length PRSS23, a biologically active fragment of PRSS23, or a fusion protein which includes all or a portion of PRSS23). Moreover, PRSS23 can be derived from any suitable mammalian species (e.g., human PRSS23, rat PRSS23 or murine PRSS23). The assay can be a binding assay entailing direct or indirect measurement of the binding of a test compound or a known PRSS23 ligand to PRSS23. The assay can also be an activity assay entailing direct or indirect measurement of the activity of PRSS23. The assay can also be an expression assay entailing direct or indirect measurement of the expression of PRSS23 mRNA or PRSS23 protein. The various screening assays are combined with an in vivo assay entailing measuring the effect of the test compound on the symptoms of diseases related to PRSS23.

[0168] The present invention includes biochemical, cell free assays that allow the identification of inhibitors and agonists of proteases suitable as lead structures for pharmacological drug development. Such assays involve contacting a form of PRSS23 (e.g., full-length PRSS23, a biologically active fragment of PRSS23, or a fusion protein comprising all or a portion of PRSS23) with a test compound and determining the ability of the test compound to act as an antagonist (preferably) or an agonist of the enzymatic activity of PRSS23.

[0169] The activity of PRSS23 molecules of the present invention can be measured using a variety of assays that measure PRSS23 activity. For example, PRSS23 enzyme activity can be assessed by a standard in vitro serine/metallo/ . . . protease assay (see, for example, [U.S. Pat. No. 5,057, 414]). Those of skill in the art are aware of a variety of substrates suitable for in vitro assays, such as SucAla-Ala-Pro-Phe-pNA, fluorescein mono-p-guanidinobenzoate hydrochloride, benzyloxycarbonyl-L-Arginyl-5-benzylester, Nalpha-Benzoyl-L-arginine ethyl ester hydrochloride, and the like. In addition, protease assay kits available from commercial sources, such as Calbiochem™ (San Diego, Calif.). For general references, see Barrett (Ed.), *Methods in Enzymology, Proteolytic Enzymes: Serine and Cysteine Peptidases* (Academic Press Inc. 1994), and Barrett et al., (Eds.), *Handbook of Proteolytic Enzymes* (Academic Press Inc. 1998).

[0170] Solution in vitro assays can be used to identify a PRSS23 substrate or inhibitor. Solid phase systems can also be used to identify a substrate or inhibitor of a PRSS23 polypeptide. For example, a PRSS23 polypeptide or PRSS23 fusion protein can be immobilized onto the surface of a receptor chip of a commercially available biosensor instrument (BIACORE, Biacore AB; Uppsala, Sweden). The use of this instrument is disclosed, for example, by [Karlsson, (1991), and Cunningham and Wells, (1993)].

[0171] In brief, a PRSS23 polypeptide or fusion protein is covalently attached, using amine or sulfhydryl chemistry, to dextran fibers that are attached to gold film within a flow cell. A test sample is then passed through the cell. If a PRSS23 substrate or inhibitor is present in the sample, it will bind to the immobilized polypeptide or fusion protein, causing a change in the refractive index of the medium, which is detected as a change in surface plasmon resonance of the gold film. This system allows the determination on- and off-rates, from which binding affinity can be calculated, and assessment of the stoichiometry of binding, as well as the kinetic effects of PRSS23 mutation. This system can also be used to examine antibody-antigen interactions, and the interactions of other complement/anti-complement pairs.

[0172] In one embodiment, the invention provides assays for screening candidate or test compounds which bind to or modulate the activity of PRSS23. Such assays can employ full-length PRSS23, a biologically active fragment of PRSS23, or a fusion protein which includes all or a portion of PRSS23. As described in greater detail below, the test compound can be obtained by any suitable means, e.g., from conventional compound libraries.

[0173] Determining the ability of the test compound to modulate the activity of PRSS23 can be accomplished, for example, by determining the ability of PRSS23 to bind to or interact with a target molecule. The target molecule can be a molecule with which PRSS23 binds or interacts with in nature. The target molecule can be a component of a signal transduction pathway which facilitates transduction of an extracellular signal. The target PRSS23 molecule can be, for example, a second intracellular protein which has catalytic activity or a protein which facilitates the association of downstream signaling molecules with PRSS23.

[0174] Determining the ability of PRSS23 to bind to or interact with a target molecule can be accomplished by one of the methods described above for determining direct binding. In one embodiment, determining the ability of a polypeptide of the invention to bind to or interact with a target molecule can be accomplished by determining the activity of the target molecule. For example, the activity of the target molecule can be determined by detecting induction of a cellular second messenger of the target (e.g., intracellular Ca^{2+} , diacylglycerol, IP_3 , etc.), detecting catalytic/enzymatic activity of the target on an appropriate substrate, detecting the induction of a reporter gene (e.g., a regulatory element that is responsive to a polypeptide of the invention operably linked to a nucleic acid encoding a detectable marker, e.g., luciferase), or detecting a cellular response.

[0175] In various embodiments of the above assay methods of the present invention, it may be desirable to immobilize PRSS23 (or a PRSS23 target molecule) to facilitate separation of complexed from uncomplexed forms of one or both of the proteins, as well as to accommodate automation of the assay. Binding of a test compound to PRSS23, or interaction of PRSS23 with a target molecule in the presence and absence

of a candidate compound, can be accomplished in any vessel suitable for containing the reactants. Examples of such vessels include microtitre plates, test tubes, and micro-centrifuge tubes. In one embodiment, a fusion protein can be provided which adds a domain that allows one or both of the proteins to be bound to a matrix. For example, glutathione-S-transferase (GST) fusion proteins or glutathione-S-transferase fusion proteins can be adsorbed onto glutathione sepharose beads (Sigma Chemical; St. Louis, Mo.) or glutathione derivatized microtitre plates, which are then combined with the test compound or the test compound and either the non-adsorbed target protein or PRSS23, and the mixture incubated under conditions conducive to complex formation (e.g., at physiological conditions for salt and pH). Following incubation, the beads or microtitre plate wells are washed to remove any unbound components and complex formation is measured either directly or indirectly, for example, as described above. Alternatively, the complexes can be dissociated from the matrix, and the level of binding or activity of PRSS23 can be determined using standard techniques.

[0176] Other techniques for immobilizing proteins on matrices can also be used in the screening assays of the invention. For example, either PRSS23 or its target molecule can be immobilized utilizing conjugation of biotin and streptavidin. Biotinylated polypeptide of the invention or target molecules can be prepared from biotin-NHS (N-hydroxy-succinimide) using techniques well known in the art (e.g., biotinylation kit, Pierce Chemicals; Rockford, Ill.), and immobilized in the wells of streptavidin-coated plates (Pierce Chemical). Alternatively, antibodies reactive with PRSS23 or target molecules but which do not interfere with binding of the polypeptide of the invention to its target molecule can be derivatized to the wells of the plate, and unbound target or polypeptide of the invention trapped in the wells by antibody conjugation. Methods for detecting such complexes, in addition to those described above for the GST-immobilized complexes, include immunodetection of complexes using antibodies reactive with PRSS23 or target molecule, as well as enzyme-linked assays which rely on detecting an enzymatic activity associated with PRSS23 or target molecule.

[0177] Another technique for drug screening which may be used provides for high throughput screening of compounds having suitable binding affinity to the protein of interest as described in published PCT application WO84/03564. In this method, large numbers of different small test compounds are synthesized on a solid substrate, such as plastic pins or some other surface. The test compounds are reacted with PRSS23, or fragments thereof, and washed. Bound PRSS23 is then detected by methods well known in the art. Purified PRSS23 can also be coated directly onto plates for use in the aforementioned drug screening techniques. Alternatively, non-neutralizing antibodies can be used to capture the peptide and immobilize it on a solid support.

[0178] In another embodiment, one may use competitive drug screening assays in which neutralizing antibodies capable of binding PRSS23 specifically compete with a test compound for binding PRSS23. In this manner, antibodies can be used to detect the presence of any peptide which shares one or more antigenic determinants with PRSS23.

[0179] The screening assay can also involve monitoring the expression of PRSS23. For example, regulators of expression of PRSS23 can be identified in a method in which a cell is contacted with a candidate compound and the expression of PRSS23 protein or mRNA in the cell is determined. The level

of expression of PRSS23 protein or mRNA the presence of the candidate compound is compared to the level of expression of PRSS23 protein or mRNA in the absence of the candidate compound. The candidate compound can then be identified as a regulator of expression of PRSS23 based on this comparison. For example, when expression of PRSS23 protein or mRNA protein is greater (statistically significantly greater) in the presence of the candidate compound than in its absence, the candidate compound is identified as a stimulator of PRSS23 protein or mRNA expression. Alternatively, when expression of PRSS23 protein or mRNA is less (statistically significantly less) in the presence of the candidate compound than in its absence, the candidate compound is identified as an inhibitor of PRSS23 protein or mRNA expression. The level of PRSS23 protein or mRNA expression in the cells can be determined by methods described below.

Binding Assays

[0180] For binding assays, the test compound is preferably a small molecule which binds to and occupies the active site of PRSS23 polypeptide, thereby making the ligand binding site inaccessible to substrate such that normal biological activity is prevented. Examples of such small molecules include, but are not limited to, small peptides or peptide-like molecules. Potential ligands which bind to a polypeptide of the invention include, but are not limited to, the natural ligands of known PRSS23 proteases and analogues or derivatives thereof.

[0181] In binding assays, either the test compound or the PRSS23 polypeptide can comprise a detectable label, such as a fluorescent, radioisotopic, chemiluminescent, or enzymatic label, such as horseradish peroxidase, alkaline phosphatase, or luciferase. Detection of a test compound which is bound to PRSS23 polypeptide can then be accomplished, for example, by direct counting of radioemission, by scintillation counting, or by determining conversion of an appropriate substrate to a detectable product. Alternatively, binding of a test compound to a PRSS23 polypeptide can be determined without labeling either of the interactants. For example, a microphysiometer can be used to detect binding of a test compound with a PRSS23 polypeptide. A micro-physiometer (e.g., Cytosensor™) is an analytical instrument that measures the rate at which a cell acidifies its environment using a light-addressable potentiometric sensor (LAPS). Changes in this acidification rate can be used as an indicator of the interaction between a test compound and PRSS23 [Haseloff, (1988)].

[0182] Determining the ability of a test compound to bind to PRSS23 also can be accomplished using a technology such as real-time Bimolecular Interaction Analysis (BIA) [McCormell, (1992); Sjolander, (1991)]. BIA is a technology for studying biospecific interactions in real time, without labeling any of the interactants (e.g., BIAcore™). Changes in the optical phenomenon surface plasmon resonance (SPR) can be used as an indication of real-time reactions between biological molecules.

[0183] In yet another aspect of the invention, a PRSS23-like polypeptide can be used as a "bait protein" in a two-hybrid assay or three-hybrid assay [Szabo, (1995); U.S. Pat. No. 5,283,317], to identify other proteins which bind to or interact with PRSS23 and modulate its activity.

[0184] The two-hybrid system is based on the modular nature of most transcription factors, which consist of separable DNA-binding and activation domains. Briefly, the assay utilizes two different DNA constructs. For example, in one

construct, polynucleotide encoding PRSS23 can be fused to a polynucleotide encoding the DNA binding domain of a known transcription factor (e.g., GAL-4). In the other construct a DNA sequence that encodes an unidentified protein ("prey" or "sample") can be fused to a polynucleotide that codes for the activation domain of the known transcription factor. If the "bait" and the "prey" proteins are able to interact in vivo to form an protein-dependent complex, the DNA-binding and activation domains of the transcription factor are brought into close proximity. This proximity allows transcription of a reporter gene (e.g., LacZ), which is operably linked to a transcriptional regulatory site responsive to the transcription factor. Expression of the reporter gene can be detected, and cell colonies containing the functional transcription factor can be isolated and used to obtain the DNA sequence encoding the protein which interacts with PRSS23.

[0185] It may be desirable to immobilize either the PRSS23 (or polynucleotide) or the test compound to facilitate separation of the bound form from unbound forms of one or both of the interactants, as well as to accommodate automation of the assay. Thus, either the PRSS23-like polypeptide (or polynucleotide) or the test compound can be bound to a solid support. Suitable solid supports include, but are not limited to, glass or plastic slides, tissue culture plates, microtiter wells, tubes, silicon chips, or particles such as beads (including, but not limited to, latex, polystyrene, or glass beads). Any method known in the art can be used to attach PRSS23-like polypeptide (or polynucleotide) or test compound to a solid support, including use of covalent and non-covalent linkages, passive absorption, or pairs of binding moieties attached respectively to the polypeptide (or polynucleotide) or test compound and the solid support. Test compounds are preferably bound to the solid support in an array, so that the location of individual test compounds can be tracked. Binding of a test compound to PRSS23 (or a polynucleotide encoding for PRSS23) can be accomplished in any vessel suitable for containing the reactants. Examples of such vessels include microtiter plates, test tubes, and microcentrifuge tubes.

[0186] In one embodiment, PRSS23 is a fusion protein comprising a domain that allows binding of PRSS23 to a solid support. For example, glutathione-S-transferase fusion proteins can be adsorbed onto glutathione sepharose beads (Sigma Chemical, St. Louis, Mo.) or glutathione derivatized microtiter plates, which are then combined with the test compound or the test compound and the non-adsorbed PRSS23; the mixture is then incubated under conditions conducive to complex formation (e.g., at physiological conditions for salt and pH). Following incubation, the beads or microtiter plate wells are washed to remove any unbound components. Binding of the interactants can be determined either directly or indirectly, as described above. Alternatively, the complexes can be dissociated from the solid support before binding is determined.

[0187] Other techniques for immobilizing proteins or polynucleotides on a solid support also can be used in the screening assays of the invention. For example, either PRSS23 (or a polynucleotide encoding PRSS23) or a test compound can be immobilized utilizing conjugation of biotin and streptavidin. Biotinylated PRSS23 (or a polynucleotide encoding biotinylated PRSS23) or test compounds can be prepared from biotin-NHS (N-hydroxysuccinimide) using techniques well known in the art (e.g., biotinylation kit, Pierce Chemicals, Rockford, Ill.) and immobilized in the wells of streptavidin-coated plates (Pierce Chemical). Alternatively, antibodies

which specifically bind to PRSS23, polynucleotide, or a test compound, but which do not interfere with a desired binding site, such as the active site of PRSS23, can be derivatized to the wells of the plate. Unbound target or protein can be trapped in the wells by antibody conjugation.

[0188] Methods for detecting such complexes, in addition to those described above for the GST-immobilized complexes, include immunodetection of complexes using antibodies which specifically bind to PRSS23 polypeptide or test compound, enzyme-linked assays which rely on detecting an activity of PRSS23 polypeptide, and SDS gel electrophoresis under non-reducing conditions.

[0189] Screening for test compounds which bind to a PRSS23 polypeptide or polynucleotide also can be carried out in an intact cell. Any cell which comprises a PRSS23 polypeptide or polynucleotide can be used in a cell-based assay system. A PRSS23 polynucleotide can be naturally occurring in the cell or can be introduced using techniques such as those described above. Binding of the test compound to PRSS23 or a polynucleotide encoding PRSS23 is determined as described above.

Functional Assays

[0190] Test compounds can be tested for the ability to increase or decrease PRSS23 activity of a PRSS23 polypeptide. The PRSS23 activity can be measured, for example, using methods described in the specific examples, below. PRSS23 activity can be measured after contacting either a purified PRSS23 or an intact cell with a test compound. A test compound which decreases PRSS23 activity by at least about 10, preferably about 50, more preferably about 75, 90, or 100% is identified as a potential agent for decreasing PRSS23 activity. A test compound which increases PRSS23 activity by at least about 10, preferably about 50, more preferably about 75, 90, or 100% is identified as a potential agent for increasing PRSS23 activity.

Gene Expression

[0191] In another embodiment, test compounds which increase or decrease PRSS23 gene expression are identified. As used herein, the term "correlates with expression of a polynucleotide" indicates that the detection of the presence of nucleic acids, the same or related to a nucleic acid sequence encoding PRSS23, by northern analysis or realtime PCR is indicative of the presence of nucleic acids encoding PRSS23 in a sample, and thereby correlates with expression of the transcript from the polynucleotide encoding PRSS23. The term "microarray", as used herein, refers to an array of distinct polynucleotides or oligonucleotides arrayed on a substrate, such as paper, nylon or any other type of membrane, filter, chip, glass slide, or any other suitable solid support. A PRSS23 polynucleotide is contacted with a test compound, and the expression of an RNA or polypeptide product of PRSS23 polynucleotide is determined. The level of expression of appropriate mRNA or polypeptide in the presence of the test compound is compared to the level of expression of mRNA or polypeptide in the absence of the test compound. The test compound can then be identified as a regulator of expression based on this comparison. For example, when expression of mRNA or polypeptide is greater in the presence of the test compound than in its absence, the test compound is identified as a stimulator or enhancer of the mRNA or polypeptide expression. Alternatively, when expression of the

mRNA or polypeptide is less in the presence of the test compound than in its absence, the test compound is identified as an inhibitor of the mRNA or polypeptide expression.

[0192] The level of PRSS23 mRNA or polypeptide expression in the cells can be determined by methods well known in the art for detecting mRNA or polypeptide. Either qualitative or quantitative methods can be used. The presence of polypeptide products of PRSS23 polynucleotide can be determined, for example, using a variety of techniques known in the art, including immuno-chemical methods such as radio-immunoassay, Western blotting, and immunohistochemistry. Alternatively, polypeptide synthesis can be determined *in vivo*, in a cell culture, or in an *in vitro* translation system by detecting incorporation of labelled amino acids into PRSS23.

[0193] Such screening can be carried out either in a cell-free assay system or in an intact cell. Any cell which expresses PRSS23 polynucleotide can be used in a cell-based assay system. The PRSS23 polynucleotide can be naturally occurring in the cell or can be introduced using techniques such as those described above. Either a primary culture or an established cell line can be used.

Test Compounds

[0194] Suitable test compounds for use in the screening assays of the invention can be obtained from any suitable source, e.g., conventional compound libraries. The test compounds can also be obtained using any of the numerous approaches in combinatorial library methods known in the art, including: biological libraries; spatially addressable parallel solid phase or solution phase libraries; synthetic library methods requiring deconvolution; the "one-bead one-compound" library method; and synthetic library methods using affinity chromatography selection. The biological library approach is limited to peptide libraries, while the other four approaches are applicable to peptide, non-peptide oligomer or small molecule libraries of compounds [Lam, (1997)]. Examples of methods for the synthesis of molecular libraries can be found in the art. Libraries of compounds may be presented in solution or on beads, bacteria, spores, plasmids or phage.

Modeling of Regulators

[0195] Computer modeling and searching technologies permit identification of compounds, or the improvement of already identified compounds, that can modulate PRSS23 expression or activity. Having identified such a compound or composition, the active sites or regions are identified. Such sites might typically be the enzymatic active site, regulator binding sites, or ligand binding sites. The active site can be identified using methods known in the art including, for example, from the amino acid sequences of peptides, from the nucleotide sequences of nucleic acids, or from study of complexes of the relevant compound or composition with its natural ligand. In the latter case, chemical or X-ray crystallographic methods can be used to find the active site by finding where on the factor the complexed ligand is found.

[0196] Next, the three dimensional geometric structure of the active site is determined. This can be done by known methods, including X-ray crystallography, which can determine a complete molecular structure. On the other hand, solid or liquid phase NMR can be used to determine certain intramolecular distances. Any other experimental method of structure determination can be used to obtain partial or com-

plete geometric structures. The geometric structures may be measured with a complexed ligand, natural or artificial, which may increase the accuracy of the active site structure determined.

[0197] If an incomplete or insufficiently accurate structure is determined, the methods of computer based numerical modeling can be used to complete the structure or improve its accuracy. Any recognized modeling method may be used, including parameterized models specific to particular biopolymers such as proteins or nucleic acids, molecular dynamics models based on computing molecular motions, statistical mechanics models based on thermal ensembles, or combined models. For most types of models, standard molecular force fields, representing the forces between constituent atoms and groups, are necessary, and can be selected from force fields known in physical chemistry. The incomplete or less accurate experimental structures can serve as constraints on the complete and more accurate structures computed by these modeling methods.

[0198] Finally, having determined the structure of the active site, either experimentally, by modeling, or by a combination, candidate modulating compounds can be identified by searching databases containing compounds along with information on their molecular structure. Such a search seeks compounds having structures that match the determined active site structure and that interact with the groups defining the active site. Such a search can be manual, but is preferably computer assisted. These compounds found from this search are potential PRSS23 modulating compounds.

[0199] Alternatively, these methods can be used to identify improved modulating compounds from an already known modulating compound or ligand. The composition of the known compound can be modified and the structural effects of modification can be determined using the experimental and computer modeling methods described above applied to the new composition. The altered structure is then compared to the active site structure of the compound to determine if an improved fit or interaction results. In this manner systematic variations in composition, such as by varying side groups, can be quickly evaluated to obtain modified modulating compounds or ligands of improved specificity or activity.

[0200] Therapeutic Indications and Methods

[0201] It was found by the present applicant that PRSS23 is expressed in various human tissues.

Cardiovascular Disorders

[0202] The human Serine protease 23 precursor is highly expressed in the following cardiovascular related tissues: heart, heart, heart, heart myocardial infarction, heart myocardial infarction, heart myocardial infarction, pericardium, heart atrium (left), heart ventricle (left), heart ventricle (right), heart apex, Purkinje fibers, interventricular septum, fetal aorta, aorta, aorta valve, artery, coronary artery, coronary artery, pulmonary artery, carotid artery, mesenteric artery, arteria radialis, vein, pulmonic valve, vein (saphena magna), (caval) vein, coronary artery endothel cells, coronary artery smooth muscle primary cells, aortic smooth muscle cells, pulmonary artery smooth muscle cells, aortic endothel cells, HUVEC cells, pulmonary artery endothel cells, iliac artery endothel cells, fetal liver, liver, liver liver cirrhosis, liver tumor, thrombocytes, fetal kidney, kidney, kidney tumor, renal epithelial cells, HEK 293 cells. Expression in the above mentioned tissues demonstrates that the human Serine protease 23 precursor or mRNA can be utilized to diagnose of

cardiovascular diseases. Additionally the activity of the human Serine protease 23 precursor can be modulated to treat cardiovascular diseases.

[0203] The human Serine protease 23 precursor is highly expressed in liver tissues: fetal liver, liver, liver liver cirrhosis, liver tumor. Expression in liver tissues demonstrates that the human Serine protease 23 precursor or mRNA can be utilized to diagnose of dyslipidemia disorders as a cardiovascular disorder. Additionally the activity of the human Serine protease 23 precursor can be modulated to treat but not limited to dyslipidemia disorders.

[0204] The human Serine protease 23 precursor is highly expressed in kidney tissues: fetal kidney, kidney, kidney tumor, HEK 293 cells. Expression in kidney tissues demonstrates that the human Serine protease 23 precursor or mRNA can be utilized to diagnose of blood pressure disorders as a cardiovascular disorder. Additionally the activity of the human Serine protease 23 precursor can be modulated to treat but not limited to blood pressure disorders as hypertension or hypotension.

[0205] The human Serine protease 23 precursor is highly expressed in heart tissues: heart, heart, heart, heart myocardial infarction, heart myocardial infarction, heart myocardial infarction, pericardium, heart atrium (left), heart ventricle (left), heart ventricle (right), heart apex, Purkinje fibers, interventricular septum. Expression in heart tissues demonstrates that the human Serine protease 23 precursor or mRNA can be utilized to diagnose of heart disorders as a cardiovascular disorder. Additionally the activity of the human Serine protease 23 precursor can be modulated to treat but not limited to heart disorders as heart failure, myocardial infarction or ischemic diseases.

[0206] Heart failure is defined as a pathophysiological state in which an abnormality of cardiac function is responsible for the failure of the heart to pump blood at a rate commensurate with the requirement of the metabolizing tissue. It includes all forms of pumping failures such as high output and low output, acute and chronic, right sided or left sided, systolic or diastolic, independent of the underlying cause.

[0207] Myocardial infarction (MI) is generally caused by an abrupt decrease in coronary blood flow that follows a thrombotic occlusion of a coronary artery previously narrowed by arteriosclerosis. MI prophylaxis (primary and secondary prevention) is included as well as the acute treatment of MI and the prevention of complications.

[0208] Ischemic diseases are conditions in which the coronary flow is restricted resulting in a perfusion which is inadequate to meet the myocardial requirement for oxygen. This group of diseases includes stable angina, unstable angina and asymptomatic ischemia.

[0209] Arrhythmias include all forms of atrial and ventricular tachyarrhythmias, atrial tachycardia, atrial flutter, atrial fibrillation, atrio ventricular reentrant tachycardia, preexcitation syndrome, ventricular tachycardia, ventricular flutter, ventricular fibrillation, as well as bradycardic forms of arrhythmias.

[0210] Hypertensive vascular diseases include primary as well as all kinds of secondary arterial hypertension, renal, endocrine, neurogenic, others. The genes may be used as drug targets for the treatment of hypertension as well as for the prevention of all complications arising from cardiovascular diseases.

[0211] Peripheral vascular diseases are defined as vascular diseases in which arterial and/or venous flow is reduced

resulting in an imbalance between blood supply and tissue oxygen demand. It includes chronic peripheral arterial occlusive disease (PAOD), acute arterial thrombosis and embolism, inflammatory vascular disorders, Raynaud's phenomenon and venous disorders.

[0212] Atherosclerosis is a cardiovascular disease in which the vessel wall is remodeled, compromising the lumen of the vessel. The atherosclerotic remodeling process involves accumulation of cells, both smooth muscle cells and monocyte/macrophage inflammatory cells, in the intima of the vessel wall. These cells take up lipid, likely from the circulation, to form a mature atherosclerotic lesion. Although the formation of these lesions is a chronic process, occurring over decades of an adult human life, the majority of the morbidity associated with atherosclerosis occurs when a lesion ruptures, releasing thrombogenic debris that rapidly occludes the artery. When such an acute event occurs in the coronary artery, myocardial infarction can ensue, and in the worst case, can result in death.

[0213] The formation of the atherosclerotic lesion can be considered to occur in five overlapping stages such as migration, lipid accumulation, recruitment of inflammatory cells, proliferation of vascular smooth muscle cells, and extracellular matrix deposition. Each of these processes can be shown to occur in man and in animal models of atherosclerosis, but the relative contribution of each to the pathology and clinical significance of the lesion is unclear.

[0214] Thus, a need exists for therapeutic methods and agents to treat cardiovascular pathologies, such as atherosclerosis and other conditions related to coronary artery disease.

[0215] Cardiovascular diseases include but are not limited to disorders of the heart and the vascular system like congestive heart failure, myocardial infarction, ischemic diseases of the heart, all kinds of atrial and ventricular arrhythmias, hypertensive vascular diseases, peripheral vascular diseases, and atherosclerosis.

[0216] To high or to low levels of fats in the bloodstream, especially cholesterol, can cause long term problems. The risk to develop atherosclerosis and coronary artery or carotid artery disease (and thus the risk of having a heart attack or stroke) increases with the total cholesterol level increasing. Nevertheless, extremely low cholesterol levels may not be healthy. Examples of disorders of lipid metabolism are hyperlipidemia (abnormally high levels of fats (cholesterol, triglycerides, or both) in the blood, may be caused by family history of hyperlipidemia, obesity, a high fat diet, lack of exercise, moderate to high alcohol consumption, cigarette smoking, poorly controlled diabetes, and an underactive thyroid gland), hereditary hyperlipidemias (type I hyperlipoproteinemia (familial hyperchylomicronemia), type II hyperlipoproteinemia (familial hypercholesterolemia), type III hyperlipoproteinemia, type IV hyperlipoproteinemia, or type V hyperlipoproteinemia), hypolipoproteinemia, lipidoses (caused by abnormalities in the enzymes that metabolize fats), Gaucher's disease, Niemann Pick disease, Fabry's disease, Wolman's disease, cerebrotendinous xanthomatosis, sitosterolemia, Refsum's disease, or Tay Sachs disease.

[0217] Kidney disorders may lead to hyper or hypotension. Examples for kidney problems possibly leading to hypertension are renal artery stenosis, pyelonephritis, glomerulonephritis, kidney tumors, polycystic kidney disease, injury to the

kidney, or radiation therapy affecting the kidney. Excessive urination may lead to hypotension.

Hematological Disorders

[0218] The human Serine protease 23 precursor is highly expressed in the following tissues of the hematological system: leukocytes (peripheral blood), erythrocytes, lymphnode, thymus, thrombocytes, bone marrow stromal cells, cord blood CD71+ cells, T-cells peripheral blood CD8+, spleen. The expression in the above mentioned tissues demonstrates that the human Serine protease 23 precursor or mRNA can be utilized to diagnose of hematological diseases. Additionally the activity of the human Serine protease 23 precursor can be modulated to treat hematological disorders.

[0219] Hematological disorders comprise diseases of the blood and all its constituents as well as diseases of organs involved in the generation or degradation of the blood. They include but are not limited to 1) Anemias, 2) Myeloproliferative Disorders, 3) Hemorrhagic Disorders, 4) Leukopenia, 5) Eosinophilic Disorders, 6) Leukemias, 7) Lymphomas, 8) Plasma Cell Dyscrasias, 9) Disorders of the Spleen in the course of hematological disorders, Disorders according to 1) include, but are not limited to anemias due to defective or deficient hem synthesis, deficient erythropoiesis. Disorders according to 2) include, but are not limited to polycythemia vera, tumor-associated erythrocytosis, myelofibrosis, thrombocytopenia. Disorders according to 3) include, but are not limited to vasculitis, thrombocytopenia, heparin-induced thrombocytopenia, thrombotic thrombocytopenic purpura, hemolytic-uremic syndrome, hereditary and acquired disorders of platelet function, hereditary coagulation disorders. Disorders according to 4) include, but are not limited to neutropenia, lymphocytopenia. Disorders according to 5) include, but are not limited to hypereosinophilia, idiopathic hypereosinophilic syndrome. Disorders according to 6) include, but are not limited to acute myeloid leukemia, acute lymphoblastic leukemia, chronic myelocytic leukemia, chronic lymphocytic leukemia, myelodysplastic syndrome. Disorders according to 7) include, but are not limited to Hodgkin's disease, non-Hodgkin's lymphoma, Burkitt's lymphoma, mycosis fungoides cutaneous T-cell lymphoma. Disorders according to 8) include, but are not limited to multiple myeloma, macroglobulinemia, heavy chain diseases. In extension of the preceding idiopathic thrombocytopenic purpura, iron deficiency anemia, megaloblastic anemia (vitamin B12 deficiency), aplastic anemia, thalassemia, malignant lymphoma bone marrow invasion, malignant lymphoma skin invasion, haemolytic uraemic syndrome, giant platelet disease are considered to be hematological diseases too.

Disorders Related to Pulmology

[0220] The human Serine protease 23 precursor is highly expressed in the following tissues of the respiratory system: pulmonary artery, pulmonary artery smooth muscle cells, leukocytes (peripheral blood), fetal lung, fetal lung fibroblast IMR-90 cells, fetal lung fibroblast MRC-5 cells, lung, lung, lung tumor, lung COPD, trachea, primary bronchia, secondary bronchia, bronchial epithelial cells, bronchial smooth muscle cells, small airway epithelial cells. The expression in the above mentioned tissues and in particular the differential expression between diseased tissue fetal lung fibroblast IMR-90 cells and healthy tissue fetal lung, between diseased tissue fetal lung fibroblast MRC-5 cells and healthy tissue fetal

lung, between diseased tissue lung COPD and healthy tissue lung demonstrates that the human Serine protease 23 precursor or mRNA can be utilized to diagnose of respiratory diseases. Additionally the activity of the human Serine protease 23 precursor can be modulated to treat those diseases. Allergy is a complex process in which environmental antigens induce clinically adverse reactions. Asthma can be understood as an basically allergic disease of the lung and its tissues. The asthma inducing antigens, called allergens, typically elicit a specific IgE response and, although in most cases the allergens themselves have little or no intrinsic toxicity, they induce pathology when the IgE response in turn elicits an IgE-dependent or T cell-dependent hypersensitivity reaction. Hypersensitivity reactions can be local or systemic and typically occur within minutes after allergen exposure in individuals who have previously been sensitized to the respective allergen. The hypersensitivity reaction of allergy develops when the allergen is recognized by IgE antibodies bound to specific receptors on the surface of effector cells, such as mast cells, basophils, or eosinophils, which causes the activation of the effector cells and the release of mediators that produce the acute signs and symptoms of the reactions. Allergic diseases include asthma, allergic rhinitis (hay fever), atopic dermatitis, and anaphylaxis.

[0221] Asthma is thought to arise as a result of interactions between multiple genetic and environmental factors and is characterized by three major features: 1) intermittent and reversible airway obstruction caused by bronchoconstriction, increased mucus production, and thickening of the walls of the airways that leads to a narrowing of the airways, 2) airway hyperresponsiveness, and 3) airway inflammation. Certain cells are critical to the inflammatory reaction of asthma and they include T cells and antigen presenting cells, B cells that produce IgE, and mast cells, basophils, eosinophils, and other cells that bind IgE. These effector cells accumulate at the site of allergic reaction in the airways and release toxic products that contribute to the acute pathology and eventually to tissue destruction related to the disorder. Other resident cells, such as smooth muscle cells, lung epithelial cells, mucus-producing cells, and nerve cells may also be abnormal in individuals with asthma and may contribute to its pathology. While the airway obstruction of asthma, presenting clinically as an intermittent wheeze and shortness of breath, is generally the most pressing symptom of the disease requiring immediate treatment, the inflammation and tissue destruction associated with the disease can lead to irreversible changes that eventually makes asthma a chronic and disabling disorder requiring long-term management.

[0222] Despite recent important advances in our understanding of the pathophysiology of allergies and asthma, they appear to be increasing in prevalence and severity [Cawkwell et al. (1993)]. It is estimated that 30-40% of the population suffer with atopic allergy, and 15% of children and 5% of adults in the population suffer from asthma. Thus, an enormous burden is placed on our health care resources. However, both diagnosis and treatment of asthma are difficult. The severity of lung tissue inflammation is not easy to measure and the symptoms of the disease are often indistinguishable from those of respiratory infections, chronic respiratory inflammatory disorders, allergic rhinitis, or other respiratory disorders. Often, the inciting allergen cannot be determined, making removal of the causative environmental agent difficult. Current pharmacological treatments suffer their own set of disadvantages. Commonly used therapeutic agents, such as

beta agonists, can act as symptom relievers to transiently improve pulmonary function, but do not affect the underlying inflammation. Agents that can reduce the underlying inflammation, such as anti-inflammatory steroids, may have major drawbacks which range from immunosuppression to bone loss. In addition, many of the present therapies, such as inhaled corticosteroids, are short-lasting, inconvenient to use, and must be used often on a regular, in some cases lifelong basis, making failure of patients to comply with the treatment a major problem and thereby reducing their effectiveness as a treatment. Because of the problems associated with conventional therapies, alternative treatment strategies have been evaluated. Glycophorin A, cyclosporin and a nonapeptide fragment of IL-2 all inhibit interleukin-2 dependent T lymphocyte proliferation; however, they are known to have many other effects. For example, cyclosporin is used as an immunosuppressant after organ transplantation. While these agents may represent alternatives to steroids in the treatment of asthmatics, they inhibit interleukin-2 dependent T lymphocyte proliferation and potentially critical immune functions associated with homeostasis. Other treatments that block the release or activity of mediators of bronchoconstriction, such as cromones or anti-leukotrienes, have recently been introduced for the treatment of mild asthma, but they are expensive and not effective in all patients and it is unclear whether they affect the chronic changes associated with asthmatic inflammation at all. What is needed in the art is the identification of a treatment that can act on pathways critical to the development of asthma and that both blocks the episodic attacks of the disorder and which dampens the hyperactive allergic immune response without immunocompromising the patient.

[0223] Chronic obstructive pulmonary (or airways) disease (COPD) is a condition defined physiologically as airflow obstruction that generally results from a mixture of emphysema and peripheral airway obstruction due to chronic bronchitis [Botstein et al. (1980)]. Emphysema is characterized by destruction of alveolar walls leading to abnormal enlargement of the air spaces of the lung. Chronic bronchitis is defined clinically as the presence of chronic productive cough for three months in each of two successive years. In COPD, airflow obstruction is usually progressive and is only partially reversible. By far the most important risk factor for development of COPD is cigarette smoking, although the disease does also occur in non-smokers.

[0224] Chronic inflammation of the airways is a key pathological feature of COPD. The inflammatory cell population comprises increased numbers of macrophages, neutrophils and CD8+ lymphocytes. Inhaled irritants such as cigarette smoke activate macrophages resident in the respiratory tract as well as epithelial cells leading to release of chemokines (e.g., interleukin-8) and other chemotactic factors which act to increase the neutrophil/monocyte trafficking from the blood into lung tissue and airways. Neutrophils and monocytes recruited into the airways can release a variety of potentially damaging mediators such as proteolytic enzymes and reactive oxygen species. Matrix degradation and emphysema, along with airway wall thickening, surfactant dysfunction and mucus hypersecretion are all potential sequelae of this inflammatory response that lead to impaired airflow and gas exchange.

Disorders Related to Urology

[0225] Genitourinary disorders comprise benign and malign disorders of the organs constituting the genitourinary

system of female and male, renal diseases like acute or chronic renal failure, immunologically mediated renal diseases like renal transplant rejection, lupus nephritis, immune complex renal diseases, glomerulopathies, nephritis, toxic nephropathy, obstructive uropathies like benign prostatic hyperplasia (BPH), neurogenic bladder syndrome, urinary incontinence like urge-, stress-, or overflow incontinence, pelvic pain, and erectile dysfunction.

[0226] The human PRSS23 protein is highly expressed in the following urological tissues: dorsal root ganglia, spinal cord, prostate, prostate BPH, prostate tumor, bladder, bladder, fetal kidney, kidney. The expression in the above mentioned tissues and in particular the differential expression between diseased tissue prostate BPH and healthy tissue prostate demonstrates that the human PRSS23 protein or mRNA can be utilized to diagnose of urological disorders. Additionally the activity of the human PRSS23 protein can be modulated to treat urological disorders.

[0227] The human PRSS23 protein is highly expressed in spinal cord tissues: spinal cord. Expression in spinal cord tissues demonstrates that the human PRSS23 protein or mRNA can be utilized to diagnose of incontinence as an urological disorder. The spinal cord tissues are involved in the neuronal regulation of the urological system. Additionally the activity of the human PRSS23 protein can be modulated to treat—but not limited to—incontinence.

Gastrointestinal and Liver Diseases

[0228] The human Serine protease 23 precursor is highly expressed in the following tissues of the gastroenterological system: esophagus, esophagus tumor, stomach, stomach tumor, colon, colon tumor, small intestine, ileum, ileum tumor, Caco 2 cells, rectum, rectum tumor, fetal liver, liver, liver liver cirrhosis, liver tumor, HuH 7 cells. The expression in the above mentioned tissues and in particular the differential expression between diseased tissue liver liver cirrhosis and healthy tissue liver demonstrates that the human Serine protease 23 precursor or mRNA can be utilized to diagnose of gastroenterological disorders. Additionally the activity of the human Serine protease 23 precursor can be modulated to treat gastroenterological disorders.

[0229] Gastrointestinal diseases comprise primary or secondary, acute or chronic diseases of the organs of the gastrointestinal tract which may be acquired or inherited, benign or malignant or metaplastic, and which may affect the organs of the gastrointestinal tract or the body as a whole. They comprise but are not limited to 1) disorders of the esophagus like achalasia, vigorous achalasia, dysphagia, cricopharyngeal incoordination, pre esophageal dysphagia, diffuse esophageal spasm, globus sensation, Barrett's metaplasia, gastroesophageal reflux, 2) disorders of the stomach and duodenum like functional dyspepsia, inflammation of the gastric mucosa, gastritis, stress gastritis, chronic erosive gastritis, atrophy of gastric glands, metaplasia of gastric tissues, gastric ulcers, duodenal ulcers, neoplasms of the stomach, 3) disorders of the pancreas like acute or chronic pancreatitis, insufficiency of the exocrine or endocrine tissues of the pancreas like steatorrhea, diabetes, neoplasms of the exocrine or endocrine pancreas like 3.1) multiple endocrine neoplasia syndrome, ductal adenocarcinoma, cystadenocarcinoma, islet cell tumors, insulinoma, gastrinoma, carcinoid tumors, glucagonoma, Zollinger Ellison syndrome, Vipoma syndrome, malabsorption syndrome, 4) disorders of the bowel like chronic inflammatory diseases of the bowel, Crohn's

disease, ileus, diarrhea and constipation, colonic inertia, megacolon, malabsorption syndrome, ulcerative colitis, 4.1) functional bowel disorders like irritable bowel syndrome, 4.2) neoplasms of the bowel like familial polyposis, adenocarcinoma, primary malignant lymphoma, carcinoid tumors, Kaposi's sarcoma, polyps, cancer of the colon and rectum.

[0230] Liver diseases comprise primary or secondary, acute or chronic diseases or injury of the liver which may be acquired or inherited, benign or malignant, and which may affect the liver or the body as a whole. They comprise but are not limited to disorders of the bilirubin metabolism, jaundice, syndroms of Gilbert's, Crigler Najjar, Dubin Johnson and Rotor; intrahepatic cholestasis, hepatomegaly, portal hypertension, ascites, Budd Chiari syndrome, portal systemic encephalopathy, fatty liver, steatosis, Reye's syndrome, liver diseases due to alcohol, alcoholic hepatitis or cirrhosis, fibrosis and cirrhosis, fibrosis and cirrhosis of the liver due to inborn errors of metabolism or exogenous substances, storage diseases, syndroms of Gaucher's, Zellweger's, Wilson's disease, acute or chronic hepatitis, viral hepatitis and its variants, inflammatory conditions of the liver due to viruses, bacteria, fungi, protozoa, helminths; drug induced disorders of the liver, chronic liver diseases like primary sclerosing cholangitis, alpha1 antitrypsin deficiency, primary biliary cirrhosis, postoperative liver disorders like postoperative intrahepatic cholestasis, hepatic granulomas, vascular liver disorders associated with systemic disease, benign or malignant neoplasms of the liver, disturbance of liver metabolism in the new born or prematurely born.

Applications

[0231] The present invention provides PRSS23 for prophylactic, therapeutic and diagnostic methods for cardiovascular diseases, hematological diseases, neurological diseases, respiratory diseases, gastroenterological diseases and urological diseases.

[0232] The regulatory method of the invention involves contacting a cell with an agent that modulates one or more of the activities of PRSS23. An agent that modulates activity can be an agent as described herein, such as a nucleic acid or a protein, a naturally-occurring cognate ligand of the polypeptide, a peptide, a peptidomimetic, or any small molecule. In one embodiment, the agent stimulates one or more of the biological activities of PRSS23. Examples of such stimulatory agents include the active PRSS23 and nucleic acid molecules encoding a portion of PRSS23. In another embodiment, the agent inhibits one or more of the biological activities of PRSS23. Examples of such inhibitory agents include antisense nucleic acid molecules and antibodies. These regulatory methods can be performed in vitro (e.g., by culturing the cell with the agent) or, alternatively, in vivo (e.g., by administering the agent to a subject). As such, the present invention provides methods of treating an individual afflicted with a disease or disorder characterized by unwanted expression or activity of PRSS23 or a protein in the PRSS23 signaling pathway. In one embodiment, the method involves administering an agent like any agent identified or being identifiable by a screening assay as described herein, or combination of such agents that modulate say upregulate or down-regulate the expression or activity of PRSS23 or of any protein in the PRSS23 signaling pathway. In another embodiment, the method involves administering a regulator of PRSS23 as

therapy to compensate for reduced or undesirably low expression or activity of PRSS23 or a protein in the PRSS23 signaling pathway.

[0233] Stimulation of activity or expression of PRSS23 is desirable in situations in which enzymatic activity or expression is abnormally low and in which increased activity is likely to have a beneficial effect. Conversely, inhibition of enzymatic activity or expression of PRSS23 is desirable in situations in which activity or expression of PRSS23 is abnormally high and in which decreasing its activity is likely to have a beneficial effect.

[0234] The present invention provides for the use of PRSS23 or fragments of PRSS23 as a biomarker for cardiovascular diseases, hematological diseases, neurological diseases, respiratory diseases, gastroenterological diseases and urological diseases.

[0235] This invention is further illustrated by the following examples which should not be construed as limiting. The contents of all references, patents and published patent applications cited throughout this application are hereby incorporated by reference.

Pharmaceutical Compositions

[0236] This invention further pertains to novel agents identified by the above-described screening assays and uses thereof for treatments as described herein.

[0237] The nucleic acid molecules, polypeptides, and antibodies (also referred to herein as "active compounds") of the invention can be incorporated into pharmaceutical compositions suitable for administration. Such compositions typically comprise the nucleic acid molecule, protein, or antibody and a pharmaceutically acceptable carrier. As used herein the language "pharmaceutically acceptable carrier" is intended to include any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents, and the like, compatible with pharmaceutical administration. The use of such media and agents for pharmaceutically active substances is well known in the art. Except insofar as any conventional media or agent is incompatible with the active compound, use thereof in the compositions is contemplated. Supplementary active compounds can also be incorporated into the compositions.

[0238] The invention includes pharmaceutical compositions comprising a regulator of PRSS23 expression or activity (and/or a regulator of the activity or expression of a protein in the PRSS23 signaling pathway) as well as methods for preparing such compositions by combining one or more such regulators and a pharmaceutically acceptable carrier. Also within the invention are pharmaceutical compositions comprising a regulator identified using the screening assays of the invention packaged with instructions for use. For regulators that are antagonists of PRSS23 activity or which reduce PRSS23 expression, the instructions would specify use of the pharmaceutical composition for treatment of cardiovascular diseases, hematological diseases, neurological diseases, respiratory diseases, gastroenterological diseases and urological diseases. For regulators that are agonists of PRSS23 activity or increase PRSS23 expression, the instructions would specify use of the pharmaceutical composition for treatment of cardiovascular diseases, hematological diseases, neurological diseases, respiratory diseases, gastroenterological diseases and urological diseases.

[0239] An inhibitor of PRSS23 may be produced using methods which are generally known in the art. In particular,

purified PRSS23 may be used to produce antibodies or to screen libraries of pharmaceutical agents to identify those which specifically bind PRSS23. Antibodies to PRSS23 may also be generated using methods that are well known in the art. Such antibodies may include, but are not limited to, polyclonal, monoclonal, chimeric, single chain antibodies, Fab fragments, and fragments produced by a Fab expression library. Neutralizing antibodies like those which inhibit dimer formation are especially preferred for therapeutic use.

[0240] In another embodiment of the invention, the polynucleotides encoding PRSS23, or any fragment or complement thereof, may be used for therapeutic purposes. In one aspect, the complement of the polynucleotide encoding PRSS23 may be used in situations in which it would be desirable to block the transcription of the mRNA. In particular, cells may be transformed with sequences complementary to polynucleotides encoding PRSS23. Thus, complementary molecules or fragments may be used to modulate PRSS23 activity, or to achieve regulation of gene function. Such technology is now well known in the art, and sense or antisense oligonucleotides or larger fragments can be designed from various locations along the coding or control regions of sequences encoding PRSS23.

[0241] Expression vectors derived from retroviruses, adenoviruses, or herpes or vaccinia viruses, or from various bacterial plasmids, may be used for delivery of nucleotide sequences to the targeted organ, tissue, or cell population. Methods which are well known to those skilled in the art can be used to construct vectors which will express nucleic acid sequence complementary to the polynucleotides of the gene encoding PRSS23. These techniques are described, for example, in [Scott and Smith (1990)].

[0242] Any of the therapeutic methods described above may be applied to any subject in need of such therapy, including, for example, mammals such as dogs, cats, cows, horses, rabbits, monkeys, and most preferably, humans.

[0243] An additional embodiment of the invention relates to the administration of a pharmaceutical composition containing PRSS23 in conjunction with a pharmaceutically acceptable carrier, for any of the therapeutic effects discussed above. Such pharmaceutical compositions may consist of PRSS23, antibodies to PRSS23, and mimetics, agonists, antagonists, or inhibitors of PRSS23. The compositions may be administered alone or in combination with at least one other agent, such as a stabilizing compound, which may be administered in any sterile, biocompatible pharmaceutical carrier including, but not limited to, saline, buffered saline, dextrose, and water. The compositions may be administered to a patient alone, or in combination with other agents, drugs or hormones.

[0244] A pharmaceutical composition of the invention is formulated to be compatible with its intended route of administration. Examples of routes of administration include parenteral, e.g., intravenous, intradermal, subcutaneous, oral (e.g., inhalation), transdermal (topical), transmucosal, and rectal administration. Solutions or suspensions used for parenteral, intradermal, or subcutaneous application can include the following components: a sterile diluent such as water for injection, saline solution, fixed oils, polyethylene glycols, glycerine, propylene glycol or other synthetic solvents; antibacterial agents such as benzyl alcohol or methyl parabens; antioxidants such as ascorbic acid or sodium bisulfite; chelating agents such as ethylenediaminetetraacetic acid; buffers such as acetates, citrates or phosphates and

agents for the adjustment of tonicity such as sodium chloride or dextrose. pH can be adjusted with acids or bases, such as hydrochloric acid or sodium hydroxide. The parenteral preparation can be enclosed in ampoules, disposable syringes or multiple dose vials made of glass or plastic.

[0245] Pharmaceutical compositions suitable for injectable use include sterile aqueous solutions (where water soluble) or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersions. For intravenous administration, suitable carriers include physiological saline, bacteriostatic water, Cremophor EM™ (BASF, Parsippany, N.J.) or phosphate buffered saline (PBS). In all cases, the composition must be sterile and should be fluid to the extent that easy syringability exists. It must be stable under the conditions of manufacture and storage and must be preserved against the contaminating action of microorganisms such as bacteria and fungi. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, a pharmaceutically acceptable polyol like glycerol, propylene glycol, liquid poly-ethylene glycol, and suitable mixtures thereof. The proper fluidity can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of dispersion and by the use of surfactants. Prevention of the action of microorganisms can be achieved by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, ascorbic acid, thimerosal, and the like. In many cases, it will be preferable to include isotonic agents, for example, sugars, polyalcohols such as mannitol, sorbitol, sodium chloride in the composition. Prolonged absorption of the injectable compositions can be brought about by including in the composition an agent which delays absorption, for example, aluminum monostearate and gelatin. Sterile injectable solutions can be prepared by incorporating the active compound (e.g., a polypeptide or antibody) in the required amount in an appropriate solvent with one or a combination of ingredients enumerated above, as required, followed by filtered sterilization. Generally, dispersions are prepared by incorporating the active compound into a sterile vehicle which contains a basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum drying and freeze-drying which yields a powder of the active ingredient plus any additional desired ingredient from a previously sterile-filtered solution thereof.

[0246] Oral compositions generally include an inert diluent or an edible carrier. They can be enclosed in gelatin capsules or compressed into tablets. For the purpose of oral therapeutic administration, the active compound can be incorporated with excipients and used in the form of tablets, troches, or capsules. Oral compositions can also be prepared using a fluid carrier for use as a mouthwash, wherein the compound in the fluid carrier is applied orally and swished and expectorated or swallowed.

[0247] Pharmaceutically compatible binding agents, and/or adjuvant materials can be included as part of the composition. The tablets, pills, capsules, troches and the like can contain any of the following ingredients, or compounds of a similar nature: a binder such as microcrystalline cellulose, gum tragacanth or gelatin; an excipient such as starch or lactose, a disintegrating agent such as alginic acid, Primogel, or corn starch; a lubricant such as magnesium stearate or steroles; a glidant such as colloidal silicon dioxide; a sweet-

ening agent such as sucrose or saccharin; or a flavoring agent such as peppermint, methyl salicylate, or orange flavoring.

[0248] For administration by inhalation, the compounds are delivered in the form of an aerosol spray from a pressurized container or dispenser which contains a suitable propellant, e.g., a gas such as carbon dioxide, or a nebulizer.

[0249] Systemic administration can also be by trans-mucosal or transdermal means. For trans-mucosal or transdermal administration, penetrants appropriate to the barrier to be permeated are used in the formulation. Such penetrants are generally known in the art, and include, for example, for trans-mucosal administration, detergents, bile salts, and fusidic acid derivatives. Trans-mucosal administration can be accomplished through the use of nasal sprays or suppositories. For trans-dermal administration, the active compounds are formulated into ointments, salves, gels, or creams as generally known in the art.

[0250] The compounds can also be prepared in the form of suppositories (e.g., with conventional suppository bases such as cocoa butter and other glycerides) or retention enemas for rectal delivery.

[0251] In one embodiment, the active compounds are prepared with carriers that will protect the compound against rapid elimination from the body, such as a controlled release formulation, including implants and microencapsulated delivery systems. Biodegradable, biocompatible polymers can be used, such as ethylene vinyl acetate, polyanhydrides, polyglycolic acid, collagen, polyorthoesters, and polylactic acid. Methods for preparation of such formulations will be apparent to those skilled in the art. The materials can also be obtained commercially from Alza Corporation and Nova Pharmaceuticals, Inc. Liposomal suspensions (including liposomes targeted to infected cells with monoclonal antibodies to viral antigens) can also be used as pharmaceutically acceptable carriers. These can be prepared according to methods known to those skilled in the art, for example, as described in U.S. Pat. No. 4,522,811.

[0252] It is especially advantageous to formulate oral or parenteral compositions in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form as used herein refers to physically discrete units suited as unitary dosages for the subject to be treated; each unit containing a predetermined quantity of active compound calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier. The specification for the dosage unit forms of the invention are dictated by and directly dependent on the unique characteristics of the active compound and the particular therapeutic effect to be achieved, and the limitations inherent in the art of compounding such an active compound for the treatment of individuals.

[0253] The pharmaceutical compositions can be included in a container, pack, or dispenser together with instructions for administration. For pharmaceutical compositions which include an antagonist of PRSS23 activity, a compound which reduces expression of PRSS23, or a compound which reduces expression or activity of a protein in the PRSS23 signaling pathway or any combination thereof, the instructions for administration will specify use of the composition for cardiovascular diseases, hematological diseases, neurological diseases, respiratory diseases, gastroenterological diseases and urological diseases. For pharmaceutical compositions which include an agonist of PRSS23 activity, a compound which increases expression of PRSS23, or a compound which increases expression or activity of a protein in the PRSS23

signaling pathway or any combination thereof, the instructions for administration will specify use of the composition for cardiovascular diseases, hematological diseases, neurological diseases, respiratory diseases, gastroenterological diseases and urological diseases.

Diagnostics

[0254] One embodiment of the invention describes PRSS23 as a biomarker for diagnostic use.

[0255] Use of PRSS23 as a biomarker in diagnostics is based by the comparison of PRSS23 level in a biological sample from a diseased mammal with the PRSS23 level in a control sample from a healthy or normal mammal. Does the PRSS23 level in the diseased mammal differs from the PRSS23 level in a normal or healthy mammal then the diseased mammal is diagnosed with a disease associated with an altered PRSS23 level. Furthermore, comparing PRSS23 levels of a biological sample from a diseased mammal with PRSS23 levels of control samples from mammals with a PRSS23-associated disease already diagnosed with different stages or severity of said disease, allows the diagnose of a PRSS23-associated disease of said first diseased mammal and specifying the severity of the PRSS23-associated disease. The biological sample is taken from the analogue tissue or body fluid than the control sample.

[0256] Normal or standard values for PRSS23 expression are established by using control samples from healthy or diseased mammalian subjects. A control sample can be obtained by collecting separate or combined body fluids or cell extracts taken from normal mammalian subjects, preferably human, achieving statistical relevant numbers. To obtain the normal or standard PRSS23 level of the control samples, the samples were subjected to suitable detection methods to detect PRSS23 polypeptide, polynucleotide or activity. The determination of PRSS23 level in a mammal subjected to diagnosis is performed analogously by collecting a biological sample from said mammal. Quantities of PRSS23 levels in biological samples from a mammal subjected to diagnosis are compared with the standard or normal values measured from a control sample. Deviation between standard value (determined from control sample) and subject value (determined from biological sample) establishes the parameters for diagnosing disease. Absolute quantification of PRSS23 levels measured from biological or control samples may be achieved by comparing those values with values obtained from an experiment in which a known amount of a substantially purified polypeptide is used.

[0257] Antibodies which specifically bind PRSS23 may be used for the diagnosis of disorders characterized by the expression of the biomarker PRSS23, or in diagnostic assays to monitor patients being treated achieving guidance for therapy for such a disease. Such a treatment includes medication suitable to treat such a disease, and treatment with PRSS23 polypeptides or polynucleotides, or agonists, antagonists, and inhibitors of PRSS23. Antibodies useful for diagnostic purposes may be prepared in the same manner as those described above for therapeutics. Diagnostic assays for PRSS23 include methods which utilize the antibody and a label to detect PRSS23 in human body fluids or in extracts of cells or tissues. The antibodies may be used with or without modification, and may be labeled by covalent or non-covalent joining with a reporter molecule. A wide variety of reporter molecules, several of which are described above, are known in the art and may be used.

[0258] A variety of protocols for measuring PRSS23, including ELISAs, RIAs, Planar Waveguide technology, and FACS, are known in the art and provide a basis for diagnosing altered or abnormal levels of PRSS23 expression. Planar Waveguide Technology bioassays are designed to perform multiplexed nucleic acid hybridization assays, immunoaffinity reactions and membrane receptor based assays with high sensitivity and selectivity. The recognition elements specific for the analytes of interest are bound onto the surface in small discrete spots; the transfer of the recognition elements onto the surface is performed using an adequate spotting technology, which requires only minute amounts of recognition elements. Such an arrangement of different recognition elements in an array format allows the simultaneous detection and quantification of hundreds to thousands of different analytes per sample including replicates.

[0259] Reactions on microarrays usually follow a typical scheme:

[0260] Recognition elements (e.g. oligonucleotides, cDNAs, or antibodies) are spotted onto the chemically modified planar waveguide surface with typical spot diameters of 100-200 μm . The remaining free binding sites on the surface subsequently are being blocked to reduce or eliminate non-specific binding. In a next step the sample (e.g. fluorescently labeled cDNA or pre-incubated analyte/fluorescently labeled antibody complex) is transferred onto the surface for incubation. The incubation time where a selective recognition and binding between recognition elements and corresponding target molecules (e.g. DNA—DNA hybridization or antigen—antibody interaction) occurs depends on the affinity between the analytes and the immobilized recognition elements. The resulting fluorescing spots can then be detected during read-out.

[0261] Due to the laterally resolved imaging of the fluorescence signals of the individual spots by a CCD-camera, a large variety of different analytes can be quantified simultaneously, requiring typically sample volumes in the range of 15 μl . Calibration and referencing spots allow for accurate quantification of analytes using just one chip and enable the establishment of dose response and time dependent activity profiles [Pawlak (2002), Duveneck (2002)].

[0262] Normal or standard values for PRSS23 expression are established by using control samples from healthy or diseased mammalian subjects. A control sample can be obtained by collecting separate or combined body fluids or cell extracts taken from normal mammalian subjects, preferably human, achieving statistical relevant numbers. To obtain normal or standard values the control samples are combined with an antibody to PRSS23 under conditions suitable for complex formation. The amount of standard complex formation may be quantified by various methods, preferably by photometric means. The determination of PRSS23 level in a mammal subjected to diagnosis is performed analogously by collecting a biological sample from said mammal, combining said sample with an antibody to PRSS23 and determination of complex formation. Quantities of PRSS23 expressed in biological samples from a mammal subjected to diagnosis are compared with the standard or normal values measured from a control sample. Deviation between standard value (determined from control sample) and subject value (determined from biological sample) establishes the parameters for diagnosing disease. Absolute quantification of PRSS23 levels measured from biological or control samples may be achieved by comparing those values with values obtained

from an experiment in which a known amount of a substantially purified polypeptide is used.

[0263] In another embodiment of the invention, the polynucleotides encoding PRSS23 may be used for diagnostic purposes. The polynucleotides which may be used include oligonucleotide sequences, complementary RNA and DNA molecules, and PNAs. The polynucleotides may be used to detect and quantified gene expression in control and biological samples in which expression of the bio-marker PRSS23 may be correlated with disease. The diagnostic assay may be used to distinguish between absence, presence, and excess expression of PRSS23, and to monitor regulation of PRSS23 levels during therapeutic intervention.

[0264] Polynucleotide sequences encoding PRSS23 may be used for the diagnosis of cardiovascular diseases, hematological diseases, neurological diseases, respiratory diseases, gastroenterological diseases and urological diseases associated with expression of PRSS23. The polynucleotide sequences encoding PRSS23 may be used in Southern, Northern, or dot-blot analysis, or other membrane-based technologies; in PCR technologies; in dipstick, pin, and ELISA assays; bDNA (branched DNA technology) and Planar Waveguide Technology; and in microarrays utilizing a biological sample from diseased mammals to detect altered PRSS23 expression. Such qualitative or quantitative methods are well known in the art.

[0265] In a particular aspect, the nucleotide sequences encoding PRSS23 may be useful in assays that detect the presence of associated disorders, particularly those mentioned above. The nucleotide sequences encoding PRSS23 may be labeled by standard methods and added to a biological sample from diseased mammals under conditions suitable for the formation of hybridization complexes. After a suitable incubation period, the sample is washed and the signal is quantified and compared with a standard value. If the amount of signal in the patient sample is altered from that of a comparable control sample, the nucleotide sequences have hybridized with nucleotide sequences in the sample, and the presence of altered levels of nucleotide sequences encoding PRSS23 in the sample indicates the presence of the associated disorder. Such assays may also be used to evaluate the efficacy of a particular therapeutic treatment regimen in animal studies, in clinical trials, or in monitoring the treatment of an individual patient.

[0266] In another embodiment of the invention PRSS23 activity can be used for diagnostic purposes. The activity of PRSS23 molecules of the present invention can be measured using a variety of assays that measure PRSS23 activity. For example, PRSS23 enzyme activity can be assessed by a standard protease assay (see, for example, [U.S. Pat. No. 5,057, 414]). Those of skill in the art are aware of a variety of substrates suitable for in vitro assays, such as SucAla-Ala-Pro-Phe-pNA, fluorescein mono-p-guanidinobenzoate hydrochloride, benzyloxycarbonyl-L-Arginyl-5-benzylester, Nalpha-Benzoyl-L-arginine ethyl ester hydrochloride, and the like. In addition, protease assay kits available from commercial sources, such as Calbiochem™ (San Diego, Calif.). For general references, see Barrett (Ed.), *Methods in Enzymology, Proteolytic Enzymes: Serine and Cysteine Peptidases* (Academic Press Inc. 1994), and Barrett et al., (Eds.), *Handbook of Proteolytic Enzymes* (Academic Press Inc. 1998).

[0267] In order to provide a basis for the diagnosis of cardiovascular diseases, hematological diseases, neurological

diseases, respiratory diseases, gastroenterological diseases and urological diseases associated with expression of PRSS23, a normal or standard profile for expression is established. This may be accomplished by combining body fluids or cell extracts taken from normal subjects, either animal or human, with a sequence, or a fragment thereof, encoding PRSS23, under conditions suitable for hybridization or amplification. Quantification of PRSS23 levels measured from biological or control samples may be achieved by comparing those values with values obtained from an experiment in which a known amount of a substantially purified polynucleotide is used. Standard values obtained from normal samples may be compared with values obtained from samples from patients who are symptomatic for a disorder. Deviation from standard values is used to establish the presence of a disorder.

Biomarker

Use of PRSS23 as a Biomarker

[0268] One of ordinary skill in the art knows several methods and devices for the detection and analysis of the markers of the instant invention. With regard to polypeptides or proteins in patient test samples, immunoassay devices and methods are often used. These devices and methods can utilize labelled molecules in various sandwich, competitive, or non-competitive assay formats, to generate a signal that is related to the presence or amount of an analyte of interest. Additionally, certain methods and devices, such as biosensors and optical immunoassays, may be employed to determine the presence or amount of analytes without the need for a labelled molecule.

[0269] Preferably the markers are analyzed using an immunoassay, although other methods are well known to those skilled in the art (for example, the measurement of marker RNA levels). The presence or amount of a marker is generally determined using antibodies specific for each marker and detecting specific binding. Any suitable immunoassay may be utilized, for example, enzyme-linked immunoassays (ELISA), radioimmunoassay (RIAs), competitive binding assays, planar waveguide technology, and the like. Specific immunological binding of the antibody to the marker can be detected directly or indirectly. Direct labels include fluorescent or luminescent tags, metals, dyes, radionuclides, and the like, attached to the antibody. Indirect labels include various enzymes well known in the art, such as alkaline phosphatase, horseradish peroxidase and the like. For an example of how this procedure is carried out on a machine, one can use the RAMP Biomedical device, called the Clinical Reader sup™, which uses the fluorescent tag method, though the skilled artisan will know of many different machines and manual protocols to perform the same assay. Diluted whole blood is applied to the sample well. The red blood cells are retained in the sample pad, and the separated plasma migrates along the strip. Fluorescent dyed latex particles bind to the analyte and are immobilized at the detection zone. Additional particles are immobilized at the internal control zone. The fluorescence of the detection and internal control zones are measured on the RAMP Clinical Reader sup™, and the ratio between these values is calculated. This ratio is used to determine the analyte concentration by interpolation from a lot-specific standard curve supplied by the manufacturer in each test kit for each assay.

[0270] The use of immobilized antibodies specific for the markers is also contemplated by the present invention and is well known by one of ordinary skill in the art. The antibodies could be immobilized onto a variety of solid supports, such as magnetic or chromatographic matrix particles, the surface of an assay plate (such as microtiter wells), pieces of a solid substrate material (such as plastic, nylon, paper), and the like. An assay strip could be prepared by coating the antibody or a plurality of antibodies in an array on solid support. This strip could then be dipped into the test sample and then processed quickly through washes and detection steps to generate a measurable signal, such as a coloured spot.

[0271] The analysis of a plurality of markers may be carried out separately or simultaneously with one test sample. Several markers may be combined into one test for efficient processing of a multiple of samples. In addition, one skilled in the art would recognize the value of testing multiple samples (for example, at successive time points) from the same individual. Such testing of serial samples will allow the identification of changes in marker levels over time. Increases or decreases in marker levels, as well as the absence of change in marker levels, would provide useful information about the disease status that includes, but is not limited to identifying the approximate time from onset of the event, the presence and amount of salvagable tissue, the appropriateness of drug therapies, the effectiveness of various therapies, identification of the severity of the event, identification of the disease severity, and identification of the patient's outcome, including risk of future events.

[0272] An assay consisting of a combination of the markers referenced in the instant invention may be constructed to provide relevant information related to differential diagnosis. Such a panel may be constructed using 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, or more or individual markers. The analysis of a single marker or subsets of markers comprising a larger panel of markers could be carried out methods described within the instant invention to optimize clinical sensitivity or specificity in various clinical settings.

[0273] The analysis of markers could be carried out in a variety of physical formats as well. For example, the use of microtiter plates or automation could be used to facilitate the processing of large numbers of test samples. Alternatively, single sample formats could be developed to facilitate immediate treatment and diagnosis in a timely fashion, for example, in ambulatory transport or emergency room settings. Particularly useful physical formats comprise surfaces having a plurality of discrete, addressable locations for the detection of a plurality of different analytes. Such formats include protein microarrays, or "protein chips" and capillary devices.

[0274] Cardiac markers serve an important role in the early detection and monitoring of cardiovascular disease. Markers of disease are typically substances found in a bodily sample that can be easily measured. The measured amount can correlate to underlying disease pathophysiology, presence or absence of a current or imminent cardiac event, probability of a cardiac event in the future. In patients receiving treatment for their condition the measured amount will also correlate with responsiveness to therapy. Markers can include elevated levels of blood pressure, cholesterol, blood sugar, homocysteine and C-reactive protein (CRP). However, current markers, even in combination with other measurements or risk factors, do not adequately identify patients at risk, accurately detect events (i.e., heart attacks), or correlate with therapy.

For example, half of patients do not have elevated serum cholesterol or other traditional risk factors.

[0275] Use of markers in diagnosis of cardiac conditions is described in, for example, Alpert et al. (2000); Newby et al. (2001); de Lemos et al. (2002); Boersma et al. (2002); Christenson et al. (2001), each of which is incorporated by reference in its entirety.

Cardiovascular Biomarker

BNP (as an Example for Cardiovascular Biomarkers)

[0276] B-type natriuretic peptide (BNP), also called brain-type natriuretic peptide is a 32 amino acid, 4 kDa peptide that is involved in the natriuresis system to regulate blood pressure and fluid balance. The precursor to BNP is synthesized as a 108-amino acid molecule, referred to as "pre pro BNP," that is proteolytically processed into a 76-amino acid N-terminal peptide (amino acids 1-76), referred to as "NT pro BNP" and the 32-amino acid mature hormone, referred to as BNP or BNP 32 (amino acids 77-108). It has been suggested that each of these species—NT pro-BNP, BNP-32, and the pre pro BNP—can circulate in human plasma. The 2 forms, pre pro BNP and NT pro BNP, and peptides which are derived from BNP, pre pro BNP and NT pro BNP and which are present in the blood as a result of proteolyses of BNP, NT pro BNP and pre pro BNP, are collectively described as markers related to or associated with BNP. Proteolytic degradation of BNP and of peptides related to BNP have also been described in the literature and these proteolytic fragments are also encompassed by the term "BNP related peptides". BNP and BNP-related peptides are predominantly found in the secretory granules of the cardiac ventricles, and are released from the heart in response to both ventricular volume expansion and pressure overload. Elevations of BNP are associated with raised atrial and pulmonary wedge pressures, reduced ventricular systolic and diastolic function, left ventricular hypertrophy, and myocardial infarction [Sagnella, (1998)]. Furthermore, there are numerous reports of elevated BNP concentration associated with congestive heart failure and renal failure. While BNP and BNP-related peptides are likely not specific for ACS, they may be sensitive markers of ACS because they may indicate not only cellular damage due to ischemia, but also a perturbation of the natriuretic system associated with ACS. The term "BNP" as used herein refers to the mature 32-amino acid BNP molecule itself. As the skilled artisan will recognize, however, other markers related to BNP may also serve as diagnostic or prognostic indicators in patients with ACS. For example, BNP is synthesized as a 108-amino acid pre pro-BNP molecule that is proteolytically processed into a 76-amino acid "NT pro BNP" and the 32-amino acid BNP molecule. Because of its relationship to BNP, the concentration of NT pro-BNP molecule can also provide diagnostic or prognostic information in patients. The phrase "marker related to BNP or BNP related peptide" refers to any polypeptide that originates from the pre pro-BNP molecule, other than the 32-amino acid BNP molecule itself. Thus, a marker related to or associated with BNP includes the NT pro-BNP molecule, the pro domain, a fragment of BNP that is smaller than the entire 32-amino acid sequence, a fragment of pre pro-BNP other than BNP, and a fragment of the pro domain.

Biomarker Classes

[0277] PRSS23 could be used as a biomarker for cardiovascular diseases, hematological diseases, neurological

diseases, respiratory diseases, gastroenterological diseases and urological diseases in different classes:

- [0278]** Disease Biomarker: a biomarker that relates to a clinical outcome or measure of disease.
- [0279]** Efficacy Biomarker: a biomarker that reflects beneficial effect of a given treatment.
- [0280]** Staging Biomarker: a biomarker that distinguishes between different stages of a chronic disorder.
- [0281]** Surrogate Biomarker: a biomarker that is regarded as a valid substitute for a clinical outcomes measure.
- [0282]** Toxicity Biomarker: a biomarker that reports a toxicological effect of a drug on an in vitro or in vivo system.
- [0283]** Mechanism Biomarker: a biomarker that reports a downstream effect of a drug.
- [0284]** Target Biomarker: a biomarker that reports interaction of the drug with its target.
- [0285]** One embodiment of the invention is a method of use of PRSS23 as a biomarker for a disease comprising:
- [0286]** (a) obtaining a biological sample from a mammal,
- [0287]** (b) measuring the level of PRSS23 in the biological sample,
- [0288]** (c) obtaining a control sample from a mammal,
- [0289]** (d) measuring the level of PRSS23 in the control sample,
- [0290]** (e) comparing the level of PRSS23 in the biological sample with the level of PRSS23 in a control sample, and
- [0291]** (f) diagnosing a disease based upon the PRSS23 level of the biological sample in comparison to the control sample.
- [0292]** The biological sample in step (a) of the methods is in a preferred embodiment a biological sample comprised in a group of samples consisting of a blood sample, a plasma sample, a serum sample, a tissue sample, a oral mucosa sample, a saliva sample, an interstitial fluid sample or an urine sample. The blood sample is for example a whole blood sample, a fractionated blood sample, a platelet sample, a neutrophil sample, a leukocyte sample, a white blood cell sample, a monocyte sample, a red blood cell sample, a granulocyte sample, and a erythrocyte sample. A tissue sample is for example a sample collected from muscle, adipose, heart or skin.
- [0293]** In a preferred embodiment PRSS23 is used as a biomarker diagnosing a disease which is associated with altered PRSS23 levels. Another preferred embodiment PRSS23 is used as a biomarker for identifying an individual risk for developing a disease, or for predicting an adverse outcome in a patient diagnosed with a disease,
- [0294]** Use of PRSS23 as a disease biomarker in diagnostics is based by the comparison of PRSS23 level in a biological sample from a diseased mammal with the PRSS23 level in a control sample from a healthy or normal mammal or a group of healthy or normal mammals. Does the PRSS23 level in the diseased mammal differs from the PRSS23 level in a normal or healthy mammal then the diseased mammal is diagnosed with a disease associated with altered PRSS23 level.
- [0295]** Furthermore, using PRSS23 as a staging biomarker, the PRSS23 levels of a diseased mammal are compared with PRSS23 levels of a mammal with a PRSS23-associated disease already diagnosed with different stages or severity of

said disease, allows the diagnose of said first diseased mammal specifying the severity of the PRSS23-associated disease.

[0296] A control sample can be a sample taken from a mammal. A control sample can be a previously taken sample from a mammal, as a PRSS23 level in a control sample can be a predetermined level of PRSS23 measured in a previously taken sample. The level of PRSS23 in a control sample or in a biological sample can be determined for example as a relative value and as an absolute value. A previously measured PRSS23 level from a control sample can be for example stored in a database, in an interne publication, in an electronically accessible form, in a publication. Comparing the level of PRSS23 of a biological sample to a control sample may be comparing relative values or absolute quantified values.

[0297] Another embodiment is a method of use of PRSS23 as a biomarker for guiding a therapy of a disease comprising:

- [0298]** (a) obtaining a baseline level of PRSS23 in biological sample from a diseased mammal,
- [0299]** (b) administering to the diseased mammal a treatment for the disease,
- [0300]** (c) obtaining one or more subsequent biological samples from the diseased mammal
- [0301]** (d) measuring the level of PRSS23 in the one or more subsequent biological samples,
- [0302]** (e) comparing the level of PRSS23 in the one or more subsequent biological samples with the baseline sample, and
- [0303]** (f) determining whether increased dosages, additional or alternative treatments are necessary based on PRSS23 levels obtained from one or more subsequent biological samples compared to the baseline PRSS23 level.

[0304] In a preferred embodiment PRSS23 is used as a biomarker for guiding a therapy in a disease which is associated with altered PRSS23 levels.

[0305] Use of PRSS23 as a disease, efficacy or surrogate endpoint biomarker in diagnostics is based by the comparison of PRSS23 level in a biological sample from a diseased mammal before treatment (the baseline sample level) with the PRSS23 level in subsequent samples from said mammal receiving a treatment for the disease. Does the PRSS23 level in the baseline sample differs from the PRSS23 level in the subsequent samples then the therapy can be considered as successful. Does the PRSS23 level in the baseline sample does not differ or differs only slightly from the PRSS23 level in the subsequent samples then the therapy can be considered as not successful. If the therapy is considered not successful increased dosages of the same therapy, repeat of the same therapy or an alternative treatment which is different from the first therapy can be considered.

[0306] The biological sample in step (a) of the methods is in a preferred embodiment a biological sample comprised in a group of samples consisting of a blood sample, a plasma sample, a serum sample, a tissue sample, a oral mucosa sample, a saliva sample, an interstitial fluid sample or an urine sample. The blood sample is for example a whole blood sample, a fractionated blood sample, a platelet sample, a neutrophil sample, a leukocyte sample, a white blood cell sample, a monocyte sample, a red blood cell sample, a granulocyte sample, and a erythrocyte sample. A tissue sample is for example a sample collected from muscle, adipose, heart, skin or a biopsy.

[0307] In a preferred embodiment the level of PRSS23 is determined by determining the level of PRSS23 polynucleotide.

[0308] In another preferred embodiment the level of PRSS23 is determined by determining the level of PRSS23 polypeptide.

[0309] In a further preferred embodiment the level of PRSS23 is determined by determining the level of PRSS23 activity.

[0310] In a preferred embodiment the disease associated with PRSS23 is comprised in a group of diseases consisting of cardiovascular diseases, hematological diseases, neurological diseases, respiratory diseases, gastroenterological diseases, and urological diseases. In a more preferred embodiment the cardiovascular disease associated with PRSS23 is comprised in a group of diseases consisting of congestive heart failure, pulmonary hypertension, left ventricular dysfunction, and right ventricular dysfunction, myocardial infarction, coronary occlusion, disease, ischemic heart disease, cardiac hypertrophy disorder, cardiac fibrosis disorders.

[0311] In a preferred embodiment of the invention the mammal is a human.

[0312] In a preferred embodiment of the invention the level of PRSS23 of the biological sample is elevated compared to the control sample.

[0313] Another embodiment of the present invention prefers the use of PRSS23 in combination with the use of one or more biomarkers, more preferably with biomarkers used in diagnosing PRSS23-associated diseases.

[0314] In a preferred embodiment of the invention the use of PRSS23 is combined with the use of one or more biomarkers which are comprised in a group of biomarkers consisting of CRTAC, FN1, NPR3, LTBP2, TGFB2, or CTGF, BNP, ANP, Troponin, CRP, Myoglobin, CK-MB and metabolites.

[0315] In a further preferred embodiment the use of PRSS23 is combined with the use of one or more clinical biomarkers which are comprised in a group of biomarkers consisting of blood pressure, heart rate, pulmonary artery pressure, or systemic vascular resistance.

[0316] In a further preferred embodiment the use of PRSS23 is combined with the use of one or more diagnostic imaging methods which are comprised in a group of methods consisting of PET (Positron Emission Tomography), CT (Computed Tomography), ultrasonic, SPECT (Single Photon Emission Computed Tomography), Echocardiography, or Impedance Cardiography.

[0317] In a further preferred embodiment the use of PRSS23 is combined with the use of one or more diagnostic imaging methods which are comprised in a group of methods consisting of PET (Positron Emission Tomography), CT (Computed Tomography), ultrasonic, SPECT (Single Photon Emission Computed Tomography), Echocardiography, Impedance Cardiography, blood pressure, heart rate, pulmonary artery pressure, systemic vascular resistance, CRTAC, FN1, NPR3, LTBP2, TGFB2, or CTGF, BNP, ANP, Troponin, CRP, Myoglobin, CK-MB, and metabolites.

[0318] In a further preferred embodiment is a kit for identifying an individual risk for developing a disease, for predicting a disease or an adverse outcome in a patient diagnosed with a disease, or for guiding a therapy in a patient with a disease, the kit comprising one or more antibodies which specifically binds PRSS23, detection means, one or more containers for collecting and or holding the biological sample, and an instruction for its use.

[0319] Another preferred embodiment is a kit for identifying an individual risk for developing a disease, for predicting a disease or an adverse outcome in a patient diagnosed with a disease, or for guiding a therapy in a patient with a disease, the kit comprising one or more probes or primers for detecting PRSS23 mRNA, detection means, one or more containers for collecting and or holding the biological sample, and an instruction for its use.

[0320] Another preferred embodiment is a kit for identifying an individual risk for developing a disease, for predicting a disease or an adverse outcome in a patient diagnosed with a disease, or for guiding a therapy in a patient with a disease, the kit comprising one or more substrates for detecting PRSS23 activity, detection means, one or more containers for collecting and or holding the biological sample, and an instruction for its use.

Determination of a Therapeutically Effective Dose

[0321] The determination of a therapeutically effective dose is well within the capability of those skilled in the art. A therapeutically effective dose refers to that amount of active ingredient which increases or decreases PRSS23 activity relative to PRSS23 activity which occurs in the absence of the therapeutically effective dose. For any compound, the therapeutically effective dose can be estimated initially either in cell culture assays or in animal models, usually mice, rabbits, dogs, or pigs. The animal model also can be used to determine the appropriate concentration range and route of administration. Such information can then be used to determine useful doses and routes for administration in humans.

[0322] Therapeutic efficacy and toxicity, e.g., ED_{50} (the dose therapeutically effective in 50% of the population) and LD_{50} (the dose lethal to 50% of the population), can be determined by standard pharmaceutical procedures in cell cultures or experimental animals. The dose ratio of toxic to therapeutic effects is the therapeutic index, and it can be expressed as the ratio, LD_{50}/ED_{50} . Pharmaceutical compositions which exhibit large therapeutic indices are preferred. The data obtained from cell culture assays and animal studies is used in formulating a range of dosage for human use. The dosage contained in such compositions is preferably within a range of circulating concentrations that include the ED_{50} with little or no toxicity. The dosage varies within this range depending upon the dosage form employed, sensitivity of the patient, and the route of administration. The exact dosage will be determined by the practitioner, in light of factors related to the subject that requires treatment. Dosage and administration are adjusted to provide sufficient levels of the active ingredient or to maintain the desired effect. Factors which can be taken into account include the severity of the disease state, general health of the subject, age, weight, and gender of the subject, diet, time and frequency of administration, drug combination(s), reaction sensitivities, and tolerance/response to therapy. Long-acting pharmaceutical compositions can be administered every 3 to 4 days, every week, or once every two weeks depending on the half-life and clearance rate of the particular formulation.

[0323] Normal dosage amounts can vary from 0.1 micrograms to 100,000 micrograms, up to a total dose of about 1 g, depending upon the route of administration. Guidance as to particular dosages and methods of delivery is provided in the literature and generally available to practitioners in the art. Those skilled in the art will employ different formulations for nucleotides than for proteins or their inhibitors. Similarly, delivery of polynucleotides or polypeptides will be specific to

particular cells, conditions, locations, etc. If the reagent is a single-chain antibody, polynucleotides encoding the antibody can be constructed and introduced into a cell either ex vivo or in vivo using well-established techniques including, but not limited to, transferrin-polycation-mediated DNA transfer, transfection with naked or encapsulated nucleic acids, liposome-mediated cellular fusion, intracellular transportation of DNA-coated latex beads, protoplast fusion, viral infection, electroporation, "gene gun", and DEAE- or calcium phosphate-mediated transfection.

[0324] If the expression product is mRNA, the reagent is preferably an antisense oligonucleotide or a ribozyme. Polynucleotides which express antisense oligonucleotides or ribozymes can be introduced into cells by a variety of methods, as described above. Preferably, a reagent reduces expression of PRSS23 gene or the activity of PRSS23 by at least about 10, preferably about 50, more preferably about 75, 90, or 100% relative to the absence of the reagent. The effectiveness of the mechanism chosen to decrease the level of expression of PRSS23 gene or the activity of PRSS23 can be assessed using methods well known in the art, such as hybridization of nucleotide probes to PRSS23-specific mRNA, quantitative RT-PCR, immunologic detection of PRSS23, or measurement of PRSS23 activity.

[0325] In any of the embodiments described above, any of the pharmaceutical compositions of the invention can be administered in combination with other appropriate therapeutic agents. Selection of the appropriate agents for use in combination therapy can be made by one of ordinary skill in the art, according to conventional pharmaceutical principles. The combination of therapeutic agents can act synergistically to effect the treatment or prevention of the various disorders described above. Using this approach, one may be able to achieve therapeutic efficacy with lower dosages of each agent, thus reducing the potential for adverse side effects. Any of the therapeutic methods described above can be applied to any subject in need of such therapy, including, for example, mammals such as dogs, cats, cows, horses, rabbits, monkeys, and most preferably, humans.

[0326] Nucleic acid molecules of the invention are those nucleic acid molecules which are contained in a group of nucleic acid molecules consisting of (i) nucleic acid molecules encoding a polypeptide comprising the amino acid sequence of SEQ ID NO: 3 or SEQ ID NO: 4, (ii) nucleic acid molecules comprising the sequence of SEQ ID NO: 1 or SEQ ID NO: 2, (iii) nucleic acid molecules having the sequence of SEQ ID NO: 1 or SEQ ID NO: 2, (iv) nucleic acid molecules the complementary strand of which hybridizes under stringent conditions to a nucleic acid molecule of (i), (ii), or (iii), (v) nucleic acid molecules the sequence of which differs from the sequence of a nucleic acid molecule of (iii) due to the degeneracy of the genetic code, (vi) nucleic acid molecules which have a sequence identity of at least 80%, 85%, 90%, 95%, 98% or 99%; and (vii) wherein the polypeptide encoded by said nucleic acid molecules of (i)-(vi) have PRSS23 activity.

[0327] Polypeptides of the invention are those polypeptides which are contained in a group of polypeptides consisting of (i) polypeptides having the sequence of SEQ ID NO: 3 or 4, (ii) polypeptides comprising the sequence of SEQ ID NO: 3 or 4, (iii) polypeptides encoded by nucleic acid molecules of the invention and (iv) polypeptides which show at least 99%, 98%, 95%, 90%, or 80% identity with a polypeptide of (i), (ii), or (iii).

[0328] An object of the invention is a method of screening for therapeutic agents useful in the treatment of a disease comprised in a group of diseases consisting of cardiovascular diseases, hematological diseases, neurological diseases, respiratory diseases, gastroenterological diseases and urological diseases in a mammal comprising the steps of (i) contacting a test compound with a PRSS23 polypeptide, (ii) detect binding of said test compound to said PRSS23 polypeptide. E.g., compounds that bind to the PRSS23 polypeptide are identified potential therapeutic agents for such a disease.

[0329] Another object of the invention is a method of screening for therapeutic agents useful in the treatment of a disease comprised in a group of diseases consisting of cardiovascular diseases, hematological diseases, neurological diseases, respiratory diseases, gastroenterological diseases and urological diseases in a mammal comprising the steps of (i) determining the activity of a PRSS23 polypeptide at a certain concentration of a test compound or in the absence of said test compound, (ii) determining the activity of said polypeptide at a different concentration of said test compound. E.g., compounds that lead to a difference in the activity of the PRSS23 polypeptide in (i) and (ii) are identified potential therapeutic agents for such a disease.

[0330] Another object of the invention is a method of screening for therapeutic agents useful in the treatment of a disease comprised in a group of diseases consisting of cardiovascular diseases, hematological diseases, neurological diseases, respiratory diseases, gastroenterological diseases and urological diseases in a mammal comprising the steps of (i) determining the activity of a PRSS23 polypeptide at a certain concentration of a test compound, (ii) determining the activity of a PRSS23 polypeptide at the presence of a compound known to be a regulator of a PRSS23 polypeptide. E.g., compounds that show similar effects on the activity of the PRSS23 polypeptide in (i) as compared to compounds used in (ii) are identified potential therapeutic agents for such a disease.

[0331] Other objects of the invention are methods of the above, wherein the step of contacting is in or at the surface of a cell.

[0332] Other objects of the invention are methods of the above, wherein the cell is in vitro.

[0333] Other objects of the invention are methods of the above, wherein the step of contacting is in a cell-free system.

[0334] Other objects of the invention are methods of the above, wherein the polypeptide is coupled to a detectable label.

[0335] Other objects of the invention are methods of the above, wherein the compound is coupled to a detectable label.

[0336] Other objects of the invention are methods of the above, wherein the test compound displaces a ligand which is first bound to the polypeptide.

[0337] Other objects of the invention are methods of the above, wherein the polypeptide is attached to a solid support.

[0338] Other objects of the invention are methods of the above, wherein the compound is attached to a solid support.

[0339] Another object of the invention is a method of screening for therapeutic agents useful in the treatment of a disease comprised in a group of diseases consisting of cardiovascular diseases, hematological diseases, neurological diseases, respiratory diseases, gastroenterological diseases and urological diseases in a mammal comprising the steps of (i) contacting a test compound with a PRSS23 polynucleotide, (ii) detect binding of said test compound to said

PRSS23 polynucleotide. Compounds that, e.g., bind to the PRSS23 polynucleotide are potential therapeutic agents for the treatment of such diseases.

[0340] Another object of the invention is the method of the above, wherein the nucleic acid molecule is RNA.

[0341] Another object of the invention is a method of the above, wherein the contacting step is in or at the surface of a cell.

[0342] Another object of the invention is a method of the above, wherein the contacting step is in a cell-free system.

[0343] Another object of the invention is a method of the above, wherein the polynucleotide is coupled to a detectable label.

[0344] Another object of the invention is a method of the above, wherein the test compound is coupled to a detectable label.

[0345] Another object of the invention is a method of diagnosing a disease comprised in a group of diseases consisting of cardiovascular diseases, hematological diseases, neurological diseases, respiratory diseases, gastroenterological diseases and urological diseases in a mammal comprising the steps of (i) determining the amount of a PRSS23 polynucleotide in a sample taken from said mammal, (ii) determining the amount of PRSS23 polynucleotide in healthy and/or diseased mammal. A disease is diagnosed, e.g., if there is a substantial similarity in the amount of PRSS23 polynucleotide in said test mammal as compared to a diseased mammal.

[0346] Another object of the invention is a pharmaceutical composition for the treatment of a disease comprised in a group of diseases consisting of cardiovascular diseases, hematological diseases, neurological diseases, respiratory diseases, gastroenterological diseases and urological diseases in a mammal comprising a therapeutic agent which binds to a PRSS23 polypeptide.

[0347] Another object of the invention is a pharmaceutical composition for the treatment of a disease comprised in a group of diseases consisting of cardiovascular diseases, hematological diseases, neurological diseases, respiratory diseases, gastroenterological diseases and urological diseases in a mammal comprising a therapeutic agent which regulates the activity of a PRSS23 polypeptide.

[0348] Another object of the invention is a pharmaceutical composition for the treatment of a disease comprised in a group of diseases consisting of cardiovascular diseases, hematological diseases, neurological diseases, respiratory diseases, gastroenterological diseases and urological diseases in a mammal comprising a therapeutic agent which regulates the activity of a PRSS23 polypeptide, wherein said therapeutic agent is (i) a small molecule, (ii) an RNA molecule, (iii) an antisense oligonucleotide, (iv) a polypeptide, (v) an antibody, or (vi) a ribozyme.

[0349] Another object of the invention is a pharmaceutical composition for the treatment of a disease comprised in a group of diseases consisting of cardiovascular diseases, hematological diseases, neurological diseases, respiratory diseases, gastroenterological diseases and urological diseases in a mammal comprising a PRSS23 polynucleotide.

[0350] Another object of the invention is a pharmaceutical composition for the treatment of a disease comprised in a group of diseases consisting of cardiovascular diseases, hematological diseases, neurological diseases, respiratory diseases, gastroenterological diseases and urological diseases in a mammal comprising a PRSS23 polypeptide.

[0351] Another object of the invention is the use of regulators of a PRSS23 for the preparation of a pharmaceutical composition for the treatment of a disease comprised in a group of diseases consisting of cardiovascular diseases, hematological diseases, neurological diseases, respiratory diseases, gastroenterological diseases and urological diseases in a mammal.

[0352] Another object of the invention is a method for the preparation of a pharmaceutical composition useful for the treatment of a disease comprised in a group of diseases consisting of cardiovascular diseases, hematological diseases, neurological diseases, respiratory diseases, gastroenterological diseases and urological diseases in a mammal comprising the steps of (i) identifying a regulator of PRSS23, (ii) determining whether said regulator ameliorates the symptoms of a disease comprised in a group of diseases consisting of cardiovascular diseases, hematological diseases, neurological diseases, respiratory diseases, gastroenterological diseases and urological diseases in a mammal; and (iii) combining of said regulator with an acceptable pharmaceutical carrier.

[0353] Another object of the invention is the use of a regulator of PRSS23 for the regulation of PRSS23 activity in a mammal having a disease comprised in a group of diseases consisting of cardiovascular diseases, hematological diseases, neurological diseases, respiratory diseases, gastroenterological diseases and urological diseases.

[0354] The uses, methods or compositions of the invention are useful for each single disease comprised in a group of diseases consisting of cardiovascular diseases, hematological diseases, neurological diseases, respiratory diseases, gastroenterological diseases and urological diseases.

[0355] The examples below are provided to illustrate the subject invention. These examples are provided by way of illustration and are not included for the purpose of limiting the invention.

EXAMPLES

Example 1

Search for Homologous Sequences in Public Sequence Data Bases

[0356] The degree of homology can readily be calculated by known methods. Preferred methods to determine homology are designed to give the largest match between the sequences tested. Methods to determine homology are codified in publicly available computer programs such as BestFit, BLASTP, BLASTN, and FASTA. The BLAST programs are publicly available from NCBI and other sources in the internet.

[0357] For PRSS23 the following hits to known sequences were identified by using the BLAST algorithm [Altschul S F, Madden T L, Schaffer A A, Zhang J, Zhang Z, Miller W, Lipman D J; Nucleic Acids Res 1997 Sep. 1; 25(17): 3389-402] and the following set of parameters: matrix=BLOSUM62 and low complexity filter. The following databases were searched: NCBI (non-redundant database) and DERWENT patent database (Geneseq).

[0358] The following hits were found:

>dbj|DD200139.1| Methods and compositions for treating cardiovascular disease, Length=1647, Score=3265 bits (1647), Expect=0.0, Identities=1647/1647 (100%), Strand=Plus/Plus

>emb|AX578034.1| Sequence 156 from Patent WO02081745, Length=1647, Score=3265 bits (1647), Expect=0.0, Identities=1647/1647 (100%), Strand=Plus/Plus
>emb|CS247591.1| Sequence 43 from Patent EP1612281, Length=1647, Score=3265 bits (1647), Expect=0.0, Identities=1647/1647 (100%), Strand=Plus/Plus
>gb|AF015287.1|AF015287 *Homo sapiens* serine protease mRNA, complete cds, Length=1647, Score=3265 bits (1647), Expect=0.0

Identities=1647/1647 (100%), Strand=Plus/Plus

[0359] >gb|BC001278.1| *Homo sapiens* protease, serine, 23, mRNA (cDNA clone MGC:5107, IMAGE:3460054), complete cds, Length=1662, Score=3263 bits (1646), Expect=0.0, Identities=1646/1646 (100%), Strand=Plus/Plus
>emb|AL136914.1|HSM801880 *Homo sapiens* mRNA; cDNA DKFZp586B0719 (from clone DKFZp586B0719), Length=1649, Score=3255 bits (1642), Expect=0.0, Identities=1645/1646 (99%), Strand=Plus/Plus
>dbj|BD261715.1|12 human secreted proteins, Length=1654, Score=3251 bits (1640), Expect=0.0, Identities=1643/1644 (99%), Strand=Plus/Plus

Example 2

Expression Profiling

[0360] Total cellular RNA was isolated from cells by one of two standard methods: 1) guanidine isothiocyanate/Cesium chloride density gradient centrifugation [Kellogg, (1990)]; or with the Tri-Reagent protocol according to the manufacturer's specifications (Molecular Research Center, Inc., Cincinnati, Ohio). Total RNA prepared by the Tri-reagent protocol was treated with DNase I to remove genomic DNA contamination.

[0361] For relative quantitation of the mRNA distribution of PRSS23, total RNA from each cell or tissue source was first reverse transcribed. 85 µg of total RNA was reverse transcribed using 1 µmole random hexamer primers, 0.5 mM each of dATP, dCTP, dGTP and dTTP (Qiagen, Hilden, Germany), 3000 U RnaseQut (Invitrogen, Groningen, Netherlands) in a final volume of 680 µl. The first strand synthesis buffer and Omniscript reverse transcriptase (2 u/µl) were from (Qiagen, Hilden, Germany). The reaction was incubated at 37° C. for 90 minutes and cooled on ice. The volume was adjusted to 6800 µl with water, yielding a final concentration of 12.5 ng/µl of starting RNA.

[0362] For relative quantitation of the distribution of PRSS23 mRNA in cells and tissues the Applied Bioscience 7900HT Sequence Detection system was used according to the manufacturer's specifications and protocols. PCR reactions were set up to quantitate PRSS23 and the housekeeping genes HPRT (hypoxanthine phosphoribosyltransferase), GAPDH (glyceraldehyde-3-phosphate dehydrogenase), β-actin, and others. Forward and reverse primers and probes for PRSS23 were designed using the Applied Bioscience ABI Primer Express™ software and were synthesized by Eurogentec (Belgium). The PRSS23 forward primer sequence was: Primed (SEQ ID NO: 8). The PRSS23 reverse primer sequence was Primer2 (SEQ ID NO: 10). Probe1 (SEQ ID NO: 9), labelled with FAM (carboxyfluorescein succinimidyl ester) as the reporter dye and TAMRA (carboxytetramethylrhodamine) as the quencher, is used as a probe for PRSS23. The following reagents were prepared in a total of 20 µl: 1×qPCR-MasterMix (Eurogentec; Belgium) and Probe1

(SEQ ID NO: 9), PRSS23 forward and reverse primers each at 200 nM, 200 nM PRSS23 FAM/TAMRA-labelled probe, and 5 µl of template cDNA. Thermal cycling parameters were 2 min at 50° C., followed by 10 min at 95° C., followed by 40 cycles of melting at 95° C. for 15 sec and annealing/extending at 60° C. for 1 min.

Calculation of Corrected CT Values

[0363] The CT (threshold cycle) value is calculated as described in the "Quantitative determination of nucleic acids" section. The CF-value (factor for threshold cycle correction) is calculated as follows:

[0364] 1. PCR reactions were set up to quantitate the housekeeping genes (HKG) for each cDNA sample.

[0365] 2. CT_{HKG} -values (threshold cycle for housekeeping gene) were calculated as described in the "Quantitative determination of nucleic acids" section.

[0366] 3. CT_{HKG} -mean values (CT mean value of all HKG tested on one cDNAs) of all HKG for each cDNA are calculated (n =number of HKG):

$$[0367] \quad CT_{HKG-n} \text{-mean value} = (CT_{HKG1} \text{-value} + CT_{HKG2} \text{-value} + \dots + CT_{HKG-n} \text{-value})/n$$

[0368] 4. CT_{panel} mean value (CT mean value of all HKG in all tested cDNAs)=

$$[0369] \quad (CT_{HKG1} \text{-mean value} + CT_{HKG2} \text{-mean value} + \dots + CT_{HKG-y} \text{-mean value})/y$$

[0370] (y =number of cDNAs)

[0371] 5. CF_{cDNA-n} (correction factor for cDNA n)= CT_{panel} - CT_{HKG-n} -mean value

[0372] 6. CT_{cDNA-n} (CT value of the tested gene for the cDNA n)+ CF_{cDNA-n} (correction factor for cDNA n)= $CT_{cor-cDNA-n}$ (corrected CT value for a gene on cDNA n)

Calculation of Relative Expression

[0373] Definition: highest $CT_{cor-cDNA-n} \neq 40$ is defined as $CT_{cor-cDNA} [high]$

$$\text{Relative Expression} = 2^{(CT_{cor-cDNA} [high] - CT_{cor-cDNA-n})}$$

Tissues

[0374] The expression of PRSS23 was investigated in the tissues in table 1.

Expression Profile

[0375] The results of the mRNA-quantification (expression profiling) is shown in Table 1.

TABLE 1

Relative expression of PRSS23 in various human tissues.	
Tissue	Relative Expression
T-cells peripheral blood CD4+	18
T-cells peripheral blood CD4+	69
T-cells peripheral blood CD4+ D117 II virus infected	34
T-cells peripheral blood CD4+ D34 virus infected	151
monocytes	239
monocytes HIV-1 infected	282
fetal heart	49
heart	399
heart	2539
heart	1201

TABLE 1-continued

Relative expression of PRSS23 in various human tissues.	
Tissue	Relative Expression
heart myocardial infarction	3281
heart myocardial infarction	2665
heart myocardial infarction	776
pericardium	976
heart atrium (right)	6747
heart atrium (right)	5518
heart atrium (left)	1808
heart atrium (left)	5955
heart ventricle (left)	224
heart ventricle (left)	2647
heart ventricle (right)	1017
heart apex	2106
Purkinje fibers	1783
interventricular septum	855
fetal aorta	352
aorta	2721
aorta	630
arcus aorta	508
aorta valve	3929
artery	495
coronary artery	1409
coronary artery	428
pulmonary artery	446
carotid artery	929
mesenteric artery	714
arteria radialis	1152
vein	1722
pulmonic valve	891
vein (saphena magna)	205
(caval) vein	936
coronary artery endothel cells	23010
coronary artery smooth muscle primary cells	2628
aortic smooth muscle cells	9742
pulmonary artery smooth muscle cells	16845
aortic endothel cells	33225
HUVEC cells	45703
pulmonary artery endothel cells	11747
iliac artery endothel cells	44453
skin	3641
adrenal gland	714
thyroid	4360
thyroid tumor	10735
pancreas	1287
esophagus	820
esophagus tumor	3956
stomach	6165
stomach tumor	2896
colon	3717
colon tumor	1428
small intestine	3616
ileum	1783
ileum tumor	1468
ileum chronic inflammation	0
rectum	11268
rectum tumor	8780
fetal liver	158
liver	413
liver	107
liver	290
liver liver cirrhosis	3616
liver tumor	6472
HuH-7 cells	4576
leukocytes (peripheral blood)	739
Jurkat (T-cells)	0
Raji (B-cells)	19
bone marrow	69
spleen	474
erythrocytes	671
lymphnode	1596
thymus	942

TABLE 1-continued

Relative expression of PRSS23 in various human tissues.	
Tissue	Relative Expression
thrombocytes	296
bone marrow stromal cells	6383
bone marrow CD71+ cells	57
bone marrow CD33+ cells	4
bone marrow CD34+ cells	43
bone marrow CD15+ cells	5
cord blood CD71+ cells	296
cord blood CD34+ cells	27
neutrophils cord blood	0
T-cells peripheral blood CD8+	288
monocytes peripheral blood CD14+	21
B-cells peripheral blood CD19+	147
neutrophils peripheral blood	9
skeletal muscle	1563
cartilage	2684
adipose	7591
adipose	9476
fetal adipose	4040
adipose (subcutaneous) BMI 21.74	13
adipose (subcutaneous) BMI 35.04	25
brain	292
cerebellum	676
cerebral cortex	709
frontal lobe	1758
occipital lobe	734
parietal lobe	1652
temporal lobe	734
substantia nigra	231
caudatum	495
hippocampus	338
thalamus	553
posteroventral thalamus	2469
dorsalmedial thalamus	704
hypothalamus	1389
dorsal root ganglia	2106
spinal cord	929
spinal cord (ventral horn)	776
spinal cord (dorsal horn)	626
glial tumor H4 cells	6937
retina	580
fetal lung	5557
fetal lung fibroblast IMR-90 cells	16271
fetal lung fibroblast MRC-5 cells	11666
lung	2957
lung	3350
lung tumor	1872
lung COPD	4973
trachea	8540
primary bronchia	3126
secondary bronchia	7332
bronchial epithelial cells	21321
bronchial smooth muscle cells	28329
small airway epithelial cells	1938
testis	2304
HeLa cells (cervix tumor)	8719
placenta	3848
uterus	5367
uterus tumor	11666
ovary	2402
ovary tumor	2210
breast	1468
breast tumor	14067
mammary gland	2077
prostate	22227
prostate	9810
prostate BPH	8841
prostate tumor	14263
bladder	19893
bladder	13034
bladder	15936
ureter	6165

TABLE 1-continued

Relative expression of PRSS23 in various human tissues.	
Tissue	Relative Expression
penis	8192
corpus cavernosum	16047
fetal kidney	12246
kidney	4905
kidney tumor	843
renal epithelial cells	12766
HEK 293 cells	3191

Example 3

Antisense Analysis

[0376] Knowledge of the correct, complete cDNA sequence coding for PRSS23 enables its use as a tool for antisense technology in the investigation of gene function. Oligonucleotides, cDNA or genomic fragments comprising the antisense strand of a polynucleotide coding for PRSS23 are used either in vitro or in vivo to inhibit translation of the mRNA. Such technology is now well known in the art, and antisense molecules can be designed at various locations along the nucleotide sequences. By treatment of cells or whole test animals with such antisense sequences, the gene of interest is effectively turned off. Frequently, the function of the gene is ascertained by observing behavior at the intracellular, cellular, tissue or organismal level (e.g., lethality, loss of differentiated function, changes in morphology, etc.).

[0377] In addition to using sequences constructed to interrupt transcription of a particular open reading frame, modifications of gene expression is obtained by designing antisense sequences to intron regions, promoter/enhancer elements, or even to trans-acting regulatory genes.

Example 4

Expression of PRSS23

[0378] Expression of PRSS23 is accomplished by subcloning the cDNAs into appropriate expression vectors and transfecting the vectors into expression hosts such as, e.g., *E. coli*. In a particular case, the vector is engineered such that it contains a promoter for β -galactosidase, upstream of the cloning site, followed by sequence containing the amino-terminal Methionine and the subsequent seven residues of β -galactosidase. Immediately following these eight residues is an engineered bacteriophage promoter useful for artificial priming and transcription and for providing a number of unique endonuclease restriction sites for cloning.

[0379] Induction of the isolated, transfected bacterial strain with Isopropyl- β -D-thiogalactopyranoside (IPTG) using standard methods produces a fusion protein corresponding to the first seven residues of β -galactosidase, about 15 residues of "linker", and the peptide encoded within the cDNA. Since cDNA clone inserts are generated by an essentially random process, there is probability of 33% that the included cDNA will lie in the correct reading frame for proper translation. If the cDNA is not in the proper reading frame, it is obtained by deletion or insertion of the appropriate number of bases using well known methods including in vitro mutagenesis, digestion with exonuclease III or mung bean nuclease, or the inclusion of an oligonucleotide linker of appropriate length.

[0380] The PRSS23 cDNA is shuttled into other vectors known to be useful for expression of proteins in specific hosts. Oligonucleotide primers containing cloning sites as well as a segment of DNA (about 25 bases) sufficient to hybridize to stretches at both ends of the target cDNA is synthesized chemically by standard methods. These primers are then used to amplify the desired gene segment by PCR. The resulting gene segment is digested with appropriate restriction enzymes under standard conditions and isolated by gel electrophoresis. Alternately, similar gene segments are produced by digestion of the cDNA with appropriate restriction enzymes. Using appropriate primers, segments of coding sequence from more than one gene are ligated together and cloned in appropriate vectors. It is possible to optimize expression by construction of such chimeric sequences.

[0381] Suitable expression hosts for such chimeric molecules include, but are not limited to, mammalian cells such as Chinese Hamster Ovary (CHO) and human 293 cells., insect cells such as Sf9 cells, yeast cells such as *Saccharomyces cerevisiae* and bacterial cells such as *E. coli*. For each of these cell systems, a useful expression vector also includes an origin of replication to allow propagation in bacteria, and a selectable marker such as the β -lactamase antibiotic resistance gene to allow plasmid selection in bacteria. In addition, the vector may include a second selectable marker such as the neomycin phosphotransferase gene to allow selection in transfected eukaryotic host cells. Vectors for use in eukaryotic expression hosts require RNA processing elements such as 3' polyadenylation sequences if such are not part of the cDNA of interest.

[0382] Additionally, the vector contains promoters or enhancers which increase gene expression. Such promoters are host specific and include MMTV, SV40, and metallothionein promoters for CHO cells; trp, lac, tac and T7 promoters for bacterial hosts; and alpha factor, alcohol oxidase and PGH promoters for yeast. Transcription enhancers, such as the rous sarcoma virus enhancer, are used in mammalian host cells. Once homogeneous cultures of recombinant cells are obtained through standard culture methods, large quantities of recombinantly produced PRSS23 are recovered from the conditioned medium and analyzed using chromatographic methods known in the art. For example, PRSS23 can be cloned into the expression vector pcDNA3, as exemplified herein. This product can be used to transform, for example, HEK293 or COS by methodology standard in the art. Specifically, for example, using Lipofectamine (Gibco BRL catalog no. 18324-020) mediated gene transfer.

Example 5

Isolation of Recombinant PRSS23

[0383] PRSS23 is expressed as a chimeric protein with one or more additional polypeptide domains added to facilitate protein purification. Such purification facilitating domains include, but are not limited to, metal chelating peptides such as histidine-tryptophan modules that allow purification on immobilized metals [Appa Rao, 1997] and the domain utilized in the FLAGS extension/affinity purification system (Immunex Corp., Seattle, Wash.). The inclusion of a cleavable linker sequence such as Factor Xa or enterokinase (Invitrogen, Groningen, The Netherlands) between the purification domain and the PRSS23 sequence is useful to facilitate expression of PRSS23.

[0384] The following example provides a method for purifying PRSS23.

[0385] PRSS23 is generated using the baculovirus expression system BAC-TO-BAC (GIBCO BRL) based on *Autographa californica* nuclear polyhedrosis virus (AcNPV) infection of *Spodoptera frugiperda* insect cells (SF9 cells).

[0386] cDNA encoding proteases cloned into either the donor plasmid pFASTBAC1 or pFASTBAC-HT which contain a mini-Tn7 transposition element. The recombinant plasmid is transformed into DH10BAC competent cells which contain the parent bacmid bMON14272 (AcNPV infectious DNA) and a helper plasmid. The mini-Tn7 element on the pFASTBAC donor can transpose to the attTn7 attachment site on the bacmid thus introducing the protease gene into the viral genome. Colonies containing recombinant bacmids are identified by disruption of the lacZ gene. The protease/bacmid construct can then be isolated and infected into insect cells (SD cells) resulting in the production of infectious recombinant baculovirus particles and expression of either unfused recombinant enzyme (pFastbac1) or PRSS23-His fusion protein (pFastbacHT).

[0387] Cells are harvested and extracts prepared 24, 48 and 72 hours after transfection. Expression of PRSS23 is confirmed by coomassie staining after sodium dodecyl sulphate-polyacrylamide gel electrophoresis (SDS-PAGE) and western blotting onto a PVDF membrane of an unstained SDS-PAGE. The protease-His fusion protein is detected due to the interaction between the Ni-NTA HRP conjugate and the His-tag which is fused to PRSS23.

Example 6

Production of PRSS23 Specific Antibodies

[0388] Two approaches are utilized to raise antibodies to PRSS23, and each approach is useful for generating either polyclonal or monoclonal antibodies. In one approach, denatured protein from reverse phase HPLC separation is obtained in quantities up to 75 mg. This denatured protein is used to immunize mice or rabbits using standard protocols; about 100 µg are adequate for immunization of a mouse, while up to 1 mg might be used to immunize a rabbit. For identifying mouse hybridomas, the denatured protein is radioiodinated and used to screen potential murine B-cell hybridomas for those which produce antibody. This procedure requires only small quantities of protein, such that 20 mg is sufficient for labeling and screening of several thousand clones.

[0389] In the second approach, the amino acid sequence of an appropriate PRSS23 domain, as deduced from translation of the cDNA, is analyzed to determine regions of high antigenicity. Oligopeptides comprising appropriate hydrophilic regions are synthesized and used in suitable immunization protocols to raise antibodies. The optimal amino acid sequences for immunization are usually at the C-terminus, the N-terminus and those intervening, hydrophilic regions of the polypeptide which are likely to be exposed to the external environment when the protein is in its natural conformation.

[0390] Typically, selected peptides, about 15 residues in length, are synthesized using an Applied Biosystems Peptide Synthesizer Model 431A using fmoc-chemistry and coupled to keyhole limpet hemocyanin (KLH; Sigma, St. Louis, Mo.) by reaction with M-maleimidobenzoyl-N-hydroxy-succinimide ester, MBS. If necessary, a cysteine is introduced at the N-terminus of the peptide to permit coupling to KLH. Rabbits are immunized with the peptide-KLH complex in complete

Freund's adjuvant. The resulting antisera are tested for anti-peptide activity by binding the peptide to plastic, blocking with 1% bovine serum albumin, reacting with antisera, washing and reacting with labeled (radioactive or fluorescent), affinity purified, specific goat anti-rabbit IgG.

[0391] Hybridomas are prepared and screened using standard techniques. Hybridomas of interest are detected by screening with labeled PRSS23 to identify those fusions producing the monoclonal antibody with the desired specificity. In a typical protocol, wells of plates (FAST; Becton-Dickinson, Palo Alto, Calif.) are coated during incubation with affinity purified, specific rabbit anti-mouse (or suitable antispecies 1 g) antibodies at 10 mg/ml. The coated wells are blocked with 1% bovine serum albumin, (BSA), washed and incubated with supernatants from hybridomas. After washing the wells are incubated with labeled PRSS23 at 1 mg/ml. Supernatants with specific antibodies bind more labeled PRSS23 than is detectable in the background. Then clones producing specific antibodies are expanded and subjected to two cycles of cloning at limiting dilution. Cloned hybridomas are injected into pristane-treated mice to produce ascites, and monoclonal antibody is purified from mouse ascitic fluid by affinity chromatography on Protein A. Monoclonal antibodies with affinities of at least 10^8M^{-1} , preferably 10^9 to 10^{10}M^{-1} or stronger, are typically made by standard procedures.

Example 7

Diagnostic Test Using PRSS23 Specific Antibodies

[0392] Particular PRSS23 antibodies are useful for investigating signal transduction and the diagnosis of infectious or hereditary conditions which are characterized by differences in the amount or distribution of PRSS23 or downstream products of an active signaling cascade.

[0393] Diagnostic tests for PRSS23 include methods utilizing antibody and a label to detect PRSS23 in human body fluids, membranes, cells, tissues or extracts of such. The polypeptides and antibodies of the present invention are used with or without modification. Frequently, the polypeptides and antibodies are labeled by joining them, either covalently or noncovalently, with a substance which provides for a detectable signal. A wide variety of labels and conjugation techniques are known and have been reported extensively in both the scientific and patent literature. Suitable labels include radionuclides, enzymes, substrates, cofactors, inhibitors, fluorescent agents, chemiluminescent agents, chromogenic agents, magnetic particles and the like.

[0394] A variety of protocols for measuring soluble or membrane-bound PRSS23, using either polyclonal or monoclonal antibodies specific for the protein, are known in the art. Examples include enzyme-linked immunosorbent assay (ELISA), radioimmunoassay (RIA) and fluorescent activated cell sorting (FACS). A two-site monoclonal-based immunoassay utilizing monoclonal antibodies reactive to two non-interfering epitopes on PRSS23 is preferred, but a competitive binding assay may be employed.

Example 8

Purification of Native PRSS23 Using Specific Antibodies

[0395] Native or recombinant PRSS23 is purified by immunoaffinity chromatography using antibodies specific for

PRSS23. In general, an immunoaffinity column is constructed by covalently coupling the anti-TRH antibody to an activated chromatographic resin.

[0396] Polyclonal immunoglobulins are prepared from immune sera either by precipitation with ammonium sulfate or by purification on immobilized Protein A (Pharmacia LKB Biotechnology, Piscataway N.J.). Likewise, monoclonal antibodies are prepared from mouse ascites fluid by ammonium sulfate precipitation or chromatography on immobilized Protein A. Partially purified immunoglobulin is covalently attached to a chromatographic resin such as CnBr-activated Sepharose (Pharmacia LKB Biotechnology). The antibody is coupled to the resin, the resin is blocked, and the derivative resin is washed according to the manufacturer's instructions.

[0397] Such immunoaffinity columns are utilized in the purification of PRSS23 by preparing a fraction from cells containing PRSS23 in a soluble form. This preparation is derived by solubilization of whole cells or of a subcellular fraction obtained via differential centrifugation (with or without addition of detergent) or by other methods well known in the art. Alternatively, soluble PRSS23 containing a signal sequence is secreted in useful quantity into the medium in which the cells are grown.

[0398] A soluble PRSS23-containing preparation is passed over the immunoaffinity column, and the column is washed under conditions that allow the preferential absorbance of PRSS23 (e.g., high ionic strength buffers in the presence of detergent). Then, the column is eluted under conditions that disrupt antibody/protein binding (e.g., a buffer of pH 2-3 or a high concentration of a chaotrope such as urea or thiocyanate ion), and PRSS23 is collected.

Example 9

Drug Screening

[0399] This invention is particularly useful for screening therapeutic compounds by using PRSS23 or fragments thereof in any of a variety of drug screening techniques.

[0400] The following example provides a system for drug screening measuring the protease activity.

[0401] The recombinant protease-His fusion protein can be purified from the crude lysate by metal-affinity chromatography using Ni-NTA agarose. This allows the specific retention of the recombinant material (since this is fused to the His-tag) whilst the endogenous insect proteins are washed off. The recombinant material is then eluted by competition with imidazol.

[0402] The activity of PRSS23 molecules of the present invention can be measured using a variety of assays that measure PRSS23 activity. For example, PRSS23 enzyme activity can be assessed by a standard *in vitro* serine/metallo/. . . protease assay (see, for example, [U.S. Pat. No. 5,057, 414]). Those of skill in the art are aware of a variety of substrates suitable for *in vitro* assays, such as SucAla-Ala-Pro-Phe-pNA, fluorescein mono-p-guanidinobenzoate hydrochloride, benzyloxycarbonyl-L-Arginyl-5-benzylester, Nalpha-Benzoyl-L-arginine ethyl ester hydrochloride, and the like. In addition, protease assay kits available from commercial sources, such as Calbiochem™ (San Diego, Calif.). For general references, see Barrett (Ed.), *Methods in Enzymology, Proteolytic Enzymes: Serine and Cysteine Pepti-*

dases (Academic Press Inc. 1994), and Barrett et al., (Eds.), *Handbook of Proteolytic Enzymes* (Academic Press Inc. 1998).

Example 10

Rational Drug Design

[0403] The goal of rational drug design is to produce structural analogs of biologically active polypeptides of interest or of small molecules with which they interact, agonists, antagonists, or inhibitors. Any of these examples are used to fashion drugs which are more active or stable forms of the polypeptide or which enhance or interfere with the function of a polypeptide *in vivo*.

[0404] In one approach, the three-dimensional structure of a protein of interest, or of a protein-inhibitor complex, is determined by x-ray crystallography, by computer modeling or, most typically, by a combination of the two approaches. Both the shape and charges of the polypeptide must be ascertained to elucidate the structure and to determine active site(s) of the molecule. Less often, useful information regarding the structure of a polypeptide is gained by modeling based on the structure of homologous proteins. In both cases, relevant structural information is used to design efficient inhibitors. Useful examples of rational drug design include molecules which have improved activity or stability or which act as inhibitors, agonists, or antagonists of native peptides.

[0405] It is also possible to isolate a target-specific antibody, selected by functional assay, as described above, and then to solve its crystal structure. This approach, in principle, yields a pharmacore upon which subsequent drug design is based. It is possible to bypass protein crystallography altogether by generating anti-idiotypic antibodies (anti-ids) to a functional, pharmacologically active antibody. As a mirror image of a mirror image, the binding site of the anti-ids is expected to be an analog of the original receptor. The anti-id is then used to identify and isolate peptides from banks of chemically or biologically produced peptides. The isolated peptides then act as the pharmacore.

[0406] By virtue of the present invention, sufficient amount of polypeptide are made available to perform such analytical studies as X-ray crystallography. In addition, knowledge of the PRSS23 amino acid sequence provided herein provides guidance to those employing computer modeling techniques in place of or in addition to x-ray crystallography.

Example 11

Identification of Other Members of the Signal Transduction Complex

[0407] Labeled PRSS23 is useful as a reagent for the purification of molecules with which it interacts. In one embodiment of affinity purification, PRSS23 is covalently coupled to a chromatography column. Cell-free extract derived from synovial cells or putative target cells is passed over the column, and molecules with appropriate affinity bind to PRSS23. PRSS23-complex is recovered from the column, and the PRSS23-binding ligand disassociated and subjected to N-terminal protein sequencing. The amino acid sequence information is then used to identify the captured molecule or to design degenerate oligonucleotide probes for cloning the relevant gene from an appropriate cDNA library.

[0408] In an alternate method, antibodies are raised against PRSS23, specifically monoclonal antibodies. The mono-

clonal antibodies are screened to identify those which inhibit the binding of labeled PRSS23. These monoclonal antibodies are then used therapeutically.

Example 12

Use and Administration of Antibodies, Inhibitors, or Antagonists

[0409] Antibodies, inhibitors, or antagonists of PRSS23 or other treatments and compounds that are limiters of signal transduction (LSTs), provide different effects when administered therapeutically. LSTs are formulated in a nontoxic, inert, pharmaceutically acceptable aqueous carrier medium preferably at a pH of about 5 to 8, more preferably 6 to 8, although pH may vary according to the characteristics of the antibody, inhibitor, or antagonist being formulated and the condition to be treated. Characteristics of LSTs include solubility of the molecule, its half-life and antigenicity/immunogenicity. These and other characteristics aid in defining an effective carrier. Native human proteins are preferred as LSTs, but organic or synthetic molecules resulting from drug screens are equally effective in particular situations.

[0410] LSTs are delivered by known routes of administration including but not limited to topical creams and gels; transmucosal spray and aerosol; transdermal patch and bandage; injectable, intravenous and lavage formulations; and orally administered liquids and pills particularly formulated to resist stomach acid and enzymes. The particular formulation, exact dosage, and route of administration is determined by the attending physician and varies according to each specific situation.

[0411] Such determinations are made by considering multiple variables such as the condition to be treated, the LST to be administered, and the pharmacokinetic profile of a particular LST. Additional factors which are taken into account include severity of the disease state, patient's age, weight, gender and diet, time and frequency of LST administration, possible combination with other drugs, reaction sensitivities, and tolerance/response to therapy. Long acting LST formulations might be administered every 3 to 4 days, every week, or once every two weeks depending on half-life and clearance rate of the particular LST.

[0412] Normal dosage amounts vary from 0.1 to 10^5 μg , up to a total dose of about 1 g, depending upon the route of administration. Guidance as to particular dosages and methods of delivery is provided in the literature; see U.S. Pat. Nos. 4,657,760; 5,206,344; or 5,225,212. Those skilled in the art employ different formulations for different LSTs. Administration to cells such as nerve cells necessitates delivery in a manner different from that to other cells such as vascular endothelial cells.

[0413] It is contemplated that abnormal signal transduction, trauma, or diseases which trigger PRSS23 activity are treatable with LSTs. These conditions or diseases are specifically diagnosed by the tests discussed above, and such testing should be performed in suspected cases of viral, bacterial or fungal infections, allergic responses, mechanical injury associated with trauma, hereditary diseases, lymphoma or carcinoma, or other conditions which activate the genes of lymphoid or neuronal tissues.

Example 13

Production of Non-Human Transgenic Animals

[0414] Animal model systems which elucidate the physiological and behavioral roles of the PRSS23 are produced by

creating nonhuman transgenic animals in which the activity of the PRSS23 is either increased or decreased, or the amino acid sequence of the expressed PRSS23 is altered, by a variety of techniques. Examples of these techniques include, but are not limited to: 1) Insertion of normal or mutant versions of DNA encoding a PRSS23, by microinjection, electroporation, retroviral transfection or other means well known to those skilled in the art, into appropriately fertilized embryos in order to produce a transgenic animal or 2) homologous recombination of mutant or normal, human or animal versions of these genes with the native gene locus in transgenic animals to alter the regulation of expression or the structure of these PRSS23 sequences. The technique of homologous recombination is well known in the art. It replaces the native gene with the inserted gene and hence is useful for producing an animal that cannot express native PRSS23s but does express, for example, an inserted mutant PRSS23, which has replaced the native PRSS23 in the animal's genome by recombination, resulting in underexpression of the transporter. Microinjection adds genes to the genome, but does not remove them, and the technique is useful for producing an animal which expresses its own and added PRSS23, resulting in overexpression of the PRSS23.

[0415] One means available for producing a transgenic animal, with a mouse as an example, is as follows: Female mice are mated, and the resulting fertilized eggs are dissected out of their oviducts. The eggs are stored in an appropriate medium such as cesiumchloride M2 medium. DNA or cDNA encoding PRSS23 is purified from a vector by methods well known to the one skilled in the art. Inducible promoters may be fused with the coding region of the DNA to provide an experimental means to regulate expression of the transgene. Alternatively or in addition, tissue specific regulatory elements may be fused with the coding region to permit tissue-specific expression of the transgene. The DNA, in an appropriately buffered solution, is put into a microinjection needle (which may be made from capillary tubing using a piper puller) and the egg to be injected is put in a depression slide. The needle is inserted into the pronucleus of the egg, and the DNA solution is injected. The injected egg is then transferred into the oviduct of a pseudopregnant mouse which is a mouse stimulated by the appropriate hormones in order to maintain false pregnancy, where it proceeds to the uterus, implants, and develops to term. As noted above, microinjection is not the only method for inserting DNA into the egg but is used here only for exemplary purposes.

Example 14

Use of PRSS23 as a Biomarker, Therapeutic and Diagnostic Target in Cardiovascular Disease (DOCA)

[0416] The DOCA-salt hypertensive rat model is a well established model of left ventricular hypertrophy.

[0417] Uninephrectomized male Sprague-Dawley rats weighing 300-350 g were given 1% NaCl in drinking water and subcutaneous injections of deoxycorticosterone acetate (DOCA, 30 mg/kg once weekly) for four weeks. Untreated rats without uninephrectomy served as control rats.

[0418] After four weeks DOCA-salt rats showed a significant increase in the tibia length-corrected left ventricular mass (DOCA-salt: 25.87 ± 0.84 mg/mm vs. control: 21.03 ± 0.60 mg/mm). At this time point heart and Li-Heparin plasma samples were taken for expression analysis.

[0419] Total cellular RNA was isolated with the Trizol-Reagent protocol according to the manufacturer's specifications (Invitrogen; USA). Total RNA prepared by the Trizol-reagent protocol was treated with DNase I to remove genomic DNA contamination.

[0420] For relative quantitation of the mRNA distribution of PRSS23, total RNA from each sample was first reverse transcribed. 1 µg of total RNA was reverse transcribed using ImProm-II Reverse Transcription System (Promega, USA) according to the manufacturer's protocol. The final volume was adjusted to 200 µl with water.

[0421] For relative quantitation of the distribution of PRSS23 mRNA the Applied Bioscience ABI 7900HT Sequence Detection system was used according to the manufacturer's specifications and protocols. PCR reactions were set up to quantitate PRSS23 and the housekeeping gene L32. Forward and reverse primers and probes for PRSS23 were designed using the Applied Bioscience ABI Primer Express™ software and were synthesized by Eurogentec (Belgium). The PRSS23 forward primer sequence was: Primer1 (SEQ ID NO: 5). The PRSS23 reverse primer sequence was Primer2 (SEQ ID NO: 7). Probe1 (SEQ ID NO: 6), labelled with FAM (carboxyfluorescein succinimidyl ester) as the reporter dye and TAMRA (carboxytetramethylrhodamine) as the quencher, is used as a probe for PRSS23. The following reagents were prepared in a total of 20 µl: 1× qPCR-MasterMix (Eurogentec; Belgium) and Probe1 (SEQ ID NO: 6), PRSS23 forward and reverse primers each at 200 nM, 200 nM PRSS23 FAM/TAMRA-labelled probe, and 5 µl of template cDNA. Thermal cycling parameters were 2 min at 50° C., followed by 10 min at 95° C., followed by 40 cycles of melting at 95° C. for 15 sec and annealing/extending at 60° C. for 1 min.

Calculation of Relative Expression

[0422] The CT (threshold cycle) value is calculated as described in the "Quantitative determination of nucleic acids" section.

$$\text{deltaCT} = \text{CT}_{\text{prss23}} - \text{CT}_{\text{L32}}$$

$$\text{relative expression} = 2^{-(15 - \text{deltaCT})}$$

[0423] The results of the mRNA-quantification (expression profiling) is shown in FIG. 11.

Example 14

Use of PRSS23 as a Biomarker, therapeutic and diagnostic Target in cardio-vascular disease (Occlusion)

[0424] In the chronic myocardial infarction model in rat [Pfeffer et al, (1979)] left coronary artery ligation is performed under isoflurane anaesthesia. Following a left thoracotomy at the fourth intercostal space, the pericardium is opened and the heart briefly exteriorized. The left coronary artery (LAD) is chronically ligated. In sham operated animals the LAD stays open. The chest is closed and animals are weaned from the ventilator and placed in cages with free access to food and water. One week after LAD occlusion application of test compounds is started. Heart tissue and plasma samples are analyzed 9 weeks after induction of the infarct towards plasma markers and expression profiles.

[0425] Total cellular RNA was isolated with the Trizol-Reagent protocol according to the manufacturer's specifica-

tions (Invitrogen; USA). Total RNA prepared by the Trizol-reagent protocol was treated with DNase I to remove genomic DNA contamination.

[0426] For relative quantitation of the mRNA distribution of PRSS23, total RNA from each sample was first reverse transcribed. 1 µg of total RNA was reverse transcribed using ImProm-II Reverse Transcription System (Promega, USA) according to the manufacturer's protocol. The final volume was adjusted to 200 µl with water.

[0427] For relative quantitation of the distribution of PRSS23 mRNA the Applied Bioscience ABI 7900HT Sequence Detection system was used according to the manufacturer's specifications and protocols. PCR reactions were set up to quantitate PRSS23 and the housekeeping gene L32. Forward and reverse primers and probes for PRSS23 were designed using the Applied Bioscience ABI Primer Express™ software and were synthesized by Eurogentec (Belgium). The PRSS23 forward primer sequence was: Primer1 (SEQ ID NO: 5). The PRSS23 reverse primer sequence was Primer2 (SEQ ID NO: 7). Probe1 (SEQ ID NO: 6), labelled with FAM (carboxyfluorescein succinimidyl ester) as the reporter dye and TAMRA (carboxytetramethylrhodamine) as the quencher, is used as a probe for PRSS23. The following reagents were prepared in a total of 20 µl: 1× qPCR-MasterMix (Eurogentec; Belgium) and Probe1 (SEQ ID NO: 6), PRSS23 forward and reverse primers each at 200 nM, 200 nM PRSS23 FAM/TAMRA-labelled probe, and 5 µl of template cDNA. Thermal cycling parameters were 2 min at 50° C., followed by 10 min at 95° C., followed by 40 cycles of melting at 95° C. for 15 sec and annealing/extending at 60° C. for 1 min.

Calculation of Relative Expression

[0428] The CT (threshold cycle) value is calculated as described in the "Quantitative determination of nucleic acids" section.

$$\text{deltaCT} = \text{CT}_{\text{prss23}} - \text{CT}_{\text{L32}}$$

$$\text{relative expression} = 2^{-(15 - \text{deltaCT})}$$

[0429] The results of the mRNA-quantification (expression profiling) is shown in FIG. 12.

Example 15

Use of PRSS23 as a Biomarker, Therapeutic and Diagnostic Target in Cardio-Vascular Disease (Monocrotalin)

[0430] Adult male Sprague-Dawley rats weighing 250 to 300 g were given a single subcutaneous injection of either 60 mg/kg Monocrotaline or vehicle.

[0431] The Monocrotaline (MCT)-treated rat is a widely used animal model for pulmonary arterial hypertension. After subcutaneous injection the pyrrolizidine alkaloid MCT is activated by the liver to the toxic MCT pyrrole, which causes endothelial injury in the pulmonary vasculature within few days with subsequent remodeling of small pulmonary arteries (de novo muscularization and medial hypertrophy). In the present study, MCT induced severe, progressive pulmonary hypertension in all animals.

[0432] Four weeks after a single MCT injection, the rats displayed threefold elevated right ventricular systolic pressure (placebo MCT: 77.62±4.17 mmHg vs. control: 26.4±1.12 mmHg; mean±sem), accompanied by a reduction of sys-

temic arterial pressure, cardiac index, arterial oxygenation and central venous oxygen saturation. In accordance with these results, an impressive right heart hypertrophy was observed (right ventricle/left ventricle+septum ratio placebo MCT: 0.62 ± 0.03 vs. control: 0.26 ± 0.01).

[0433] Heart and Li-Heparin plasma samples were taken for expression analysis four weeks after the MCT injection.

[0434] Total cellular RNA was isolated with the Trizol-Reagent protocol according to the manufacturer's specifications (Invitrogen; USA). Total RNA prepared by the Trizol-reagent protocol was treated with DNase I to remove genomic DNA contamination.

[0435] For relative quantitation of the mRNA distribution of PRSS23, total RNA from each sample was first reverse transcribed. 1 μ g of total RNA was reverse transcribed using ImProm-II Reverse Transcription System (Promega, USA) according to the manufacturer's protocol. The final volume was adjusted to 200 μ l with water.

[0436] For relative quantitation of the distribution of PRSS23 mRNA the Applied Bioscience ABI 7900HT Sequence Detection system was used according to the manufacturer's specifications and protocols. PCR reactions were set up to quantitate PRSS23 and the housekeeping gene L32. Forward and reverse primers and probes for PRSS23 were designed using the Applied Bioscience ABI Primer Express™ software and were synthesized by Eurogentec (Belgium). The PRSS23 forward primer sequence was: Primer1 (SEQ ID NO: 5). The PRSS23 reverse primer sequence was Primer2 (SEQ ID NO: 7). Probe1 (SEQ ID NO: 6), labelled with FAM (carboxyfluorescein succinimidyl ester) as the reporter dye and TAMRA (carboxytetramethylrhodamine) as the quencher, is used as a probe for PRSS23. The following reagents were prepared in a total of 20 μ l: 1 \times qPCR-MasterMix (Eurogentec; Belgium) and Probe1 (SEQ ID NO: 6), PRSS23 forward and reverse primers each at 200 nM, 200 nM PRSS23 FAM/TAMRA-labelled probe, and 5 p. 1 of template cDNA. Thermal cycling parameters were 2 min at 50° C., followed by 10 min at 95° C., followed by 40 cycles of melting at 95° C. for 15 sec and annealing/extending at 60° C. for 1 min.

Calculation of Relative Expression

[0437] The CT (threshold cycle) value is calculated as described in the "Quantitative determination of nucleic acids" section.

$$\text{deltaCT} = \text{CT}_{\text{prss23}} - \text{CT}_{\text{L32}}$$

$$\text{relative expression} = 2^{-(15 - \text{deltaCT})}$$

[0438] The results of the mRNA-quantification (expression profiling) is shown in FIG. 13.

Example 16

Microarray Experiments

[0439] Total RNA extracted from cardiac tissue and was purified using an affinity resin column (RNeasy; Qiagen, Hilden, Germany), quantified by spectrophotometry (absorbance 260 nm), and the quality of RNA was assessed by microfluidics electrophoretic separation with a Bioanalyzer (Agilent Technologies, Palo Alto, USA). Purified total RNA (1 mg) was converted to cDNA using the Superscript Choice cDNA synthesis kit (Invitrogen, Carlsbad, Calif., USA), incorporating a T7-(dT)₂₄ primer. Double-stranded cDNA was then purified by affinity resin column (Clean up Kit, Qiagen, Hilden, Germany) with ethanol extraction. Purified cDNA was used as a template for in vitro transcription reac-

tion for the synthesis of biotinylated cRNA using an Enzo BioArray HighYield RNA transcription labeling kit (Affymetrix, Santa Clara, Calif.), and further purified using an affinity resin column (Clean up Kit, Qiagen, Hilden, Germany). After purification, in vitro cRNA was fragmented in buffer containing magnesium at 95° C. for 35 min. Fragmented cRNA was hybridized onto the Affymetrix GeneChip Human Genome U133 Plus 2.0 Array. Briefly, 15 mg fragmented cRNA was added along with control cRNA (BioB, BioC, and BioD), herring sperm DNA (10 mg/ml), 10% DMSO, and acetylated BSA (50 mg/ml) to the hybridization buffer. The hybridization mixture was heated at 99° C. for 5 min, incubated at 45° C. for 5 min, centrifuged for 5 min at 13,000 rpm, and injected into the microarray. After hybridization at 45° C. for 16 h rotating at 60 rpm, the array was washed and stained with the Affymetrix Fluidics Protocols-antibody amplification for Eukaryotic Targets, and scanned using an Affymetrix microarray scanner (GeneChip Scanner 3000 7G system) at 570 nm.

Example 17

Microarray Expression Data from Human Heart of CHF Patients with Left Ventricular Assist Devices

[0440] Implantation of left ventricular assist devices (LVAD) often is the only possible means of supporting patients with end-stage heart failure in the form of bridging to transplantation (see [Clegg et al. (2005)] for a review). Like the heart, the LVAD is a pump. One end hooks up to the left ventricle—that's the chamber of the heart that pumps blood out of the lungs and into the body. The other end hooks up to the aorta, the body's main artery. A tube passes from the device through the skin. The outside of the tube is covered with a special material to aid in healing and allow the skin to regrow. The LVAD is implanted during open-heart surgery. Recent reports demonstrate that LVAD support may be associated with adaptive remodeling of the ventricular myocardium, including reduced LV mass, wall thickness and myocyte diameter, changes in LV pressure-volume relationships and reversal of LV chamber dilation [Li et al. (2001)].

[0441] Myocardial samples of the left ventricle were collected during cardiac surgery from 32 heart failure patients at the time of cardiac transplantation or insertion of a mechanical assist device. Corresponding myocardial specimen are designed as pre- and post-LVAD samples. All procedures involving human tissue use were approved by the institutional review boards of the "Heart- and Diabetes-Center North Rhine Westphalia, Bad Oeynhausen, Germany. Consent was obtained from patients before tissue harvest. Samples were immediately frozen in liquid nitrogen and pulverized using pestle and mortar. Total RNA was isolated according to standard procedures.

Example 18

Data Analysis from Microarray Experiments

[0442] Raw data analysis and scaling were performed in Microarray Suite 5.0 software (Affymetrix), and normalization and further analysis in expressionist Pro 3.0 (Genedata). Results for HG-U133 Plus 2.0 arrays were subjected to global scaling with a target intensity of 100.

[0443] Base-2 logarithms were calculated for all expression values and taken for subsequent statistical analysis. To analyze the differential expression between the two groups, non-failing hearts (N) and pre-operation hearts (P), a two-tailed Student's test was applied to the expression values under the assumption of equal variances. A resultant p-value of less or equal than 0.05 was taken as indicator for significant differential expression.

REFERENCES

- [0444] EP 1 069 188
 [0445] EP 1 275 733
 [0446] EP 1 308 459
 [0447] EP 1 560 025
 [0448] EP 1 612 281
 [0449] U.S. Pat. No. 4,522,811
 [0450] U.S. Pat. No. 5,057,414
 [0451] U.S. Pat. No. 5,283,317
 [0452] U.S. Pat. No. 5,565,332
 [0453] U.S. Pat. No. 5,723,323.
 [0454] U.S. Pat. No. 5,747,334
 [0455] U.S. Pat. No. 5,783,384
 [0456] U.S. Pat. No. 5,885,814
 [0457] U.S. Pat. No. 5,985,629
 [0458] WO 84/03564
 [0459] WO 93/03151
 [0460] WO 94/13804
 [0461] WO 00/47750
 [0462] WO 02/06492
 [0463] WO 02/26958
 [0464] WO 02/47670
 [0465] WO 03/051370
 [0466] WO 02/081745
 [0467] Altschul S F, Madden T L, Schaffer A A, Zhang J, Zhang Z, Miller W, Lipman D J; *Nucleic Acids Res* 1997 Sep. 1; 25(17): 3389-402
 [0468] Appa Rao et al., 1997, *Protein Expr Purif* Nov, 11(2): 201-8
 [0469] Alpert, J. S., et al. *J. Am. Coll. Cardiol.* 2000; 36:959-69
 [0470] Avalle et al., *Ann. NY Acad. Sci.* 864:118 (1998).
 [0471] Barnes, 2000, *Chest*, 117:10 S14S
 [0472] Barrett et al., (Eds.), *Handbook of Proteolytic Enzymes* (Academic Press Inc. 1998).
 [0473] Barrett (Ed.), *Methods in Enzymology, Proteolytic Enzymes: Serine and Cysteine Peptidases* (Academic Press Inc. 1994)
 [0474] Boersma, E., et al. *Lancet* 2002; 359:189-98
 [0475] Botstein et al., 1980, *Am J Hum Genet.* 32: 314-31
 [0476] Clegg et al. 2005, *Health Technol Assess.* November; 9(45):1-148
 [0477] Colbere-Garapin et al., 1981, *J. Mol. Biol.* 150, 1-14
 [0478] Christenson, R. H., et al., *Clin. Chem.* 2001; 47:464-470
 [0479] Cunningham and Wells, *J. Mol. Biol.* 234:554 (1993).
 [0480] DesGroseillers et al. (2001), *DNA Cell Biol.* August; 20(8):493-8.
 [0481] de Lemos, J. A., et al. *J. Am. Coll. Cardiol.* 2002; 40:238-44
 [0482] Duveneck G. L. et al, *Analytica Chimica Acta* 469 (2002), 49-61.
 [0483] Engelhard et al., 1994, *Proc. Nat. Acad. Sci.* 91, 3224-3227
 [0484] Friboulet et al., *Appl. Biochem. Biotechnol.* 47:229 (1994)
 [0485] Gergen and Weiss, 1992, *Am Rev Respir Dis* 146: 823-824
 [0486] Gibson et al., 1996, *Genome Research* 6: 995-1001
 [0487] Haseloff et al., 1988, *Nature* 334, 585-591
 [0488] Heid et al., 1996, *Genome Research* 6: 986-994
 [0489] Holland et al., 1991, *PNAS* 88: 7276-7280
 [0490] Ifon et al. 2005, *Cancer Cell Int.* June 22; 5:19.
 [0491] Jeffreys et al., 1985, *Nature* 316: 76-9
 [0492] Johnson et al., 1989, *Endoc. Rev.* 10, 317-331
 [0493] Joron et al., *Ann. NY Acad. Sci.* 672:216 (1992)
 [0494] Karlsson, *Immunol. Methods* 145:229 (1991)
 [0495] Kellogg et al., 1990, *Anal. Biochem.* 189:202-208
 [0496] Lam, 1997, *Anticancer Drug Res.* 12(3):145-67
 [0497] Li et al. 2001, *Circulation.* September 4; 104(10): 1147-52
 [0498] Livak et al., 1995, *PCR Methods and Applications* 357-362
 [0499] Logan, Shenk, 1984, *Proc. Natl. Acad. Sci.* 81, 3655-3659
 [0500] Lowy et al., 1980, *Cell* 22, 817-23
 [0501] Maddox et al., 1983, *J. Exp. Med.* 158, 1211-1216
 [0502] Monfardini et al., *Proc. Assoc. Am. Physicians* 108: 420 (1996)
 [0503] McConnell et al., 1992, *Science* 257, 1906-1912
 [0504] Nicholls et al., 1993, *J. Immunol. Meth.* 165, 81-91
 [0505] Newby, L. K., et al. *Circulation* 2001:103; 1832-7
 [0506] Pawlak M. et al, *Proteomics* 2(4) (2002), 383-393
 [0507] Pentecost et al. 2005, *Mol Cell Endocrinol.* June 30; 238(1-2):9-25.
 [0508] Piatak et al., 1993, *BioTechniques* 14:70-81
 [0509] Piatak et al., 1993, *Science* 259:1749-1754
 [0510] Porath et al., 1992, *Prot. Exp. Purif.* 3, 263-281
 [0511] Pfeffer et al., *Circ Res.* 1979 April; 44(4):503-12.
 [0512] Roberge et al., 1995, *Science* 269, 202-204
 [0513] Sagnella, G. A., *Clinical Science* 95:519-529, 1998
 [0514] Scott and Smith (1990) *Science* 249:386-390
 [0515] Sjolander, Urbaniczky, 1991, *Anal. Chem.* 63, 2338-2345
 [0516] Szabo et al., 1995, *Curr. Opin. Struct. Biol.* 5, 699-705
 [0517] Thomas, 1980, *Proc. Nat. Acad. Sci.*, 77:5201-5205
 [0518] Uhlmann et al., 1987, *Tetrahedron. Lett.* 215, 3539-3542
 [0519] Weber et al., 1990, *Genomics* 7: 524-30
 [0520] Wigler et al., 1977, *Cell* 11, 223-32
 [0521] Wigler et al., 1980, *Proc. Natl. Acad. Sci.* 77, 3567-70

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 10

<210> SEQ ID NO 1
 <211> LENGTH: 1647
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

-continued

<400> SEQUENCE: 1

```

ggcaggcatg ggagccgcgc gctctctccc ggcgcccaca cctgtctgag cggcgcagcg      60
agccgcggcc cgggggggct gctcggcgcg gaacagtgct cggcatggca gggattccag      120
ggctectctt ccttctcttc tttctgctct gtgctgttgg gcaagtgagc ccttacagtg      180
ccccctggaa acccacttgg cctgcatacc gcctcctctg cgtcttgcce cagtctaccc      240
tcaatttagc caagccagac tttggagccg aagccaaatt agaagtatct tcttcatgtg      300
gaccccagtg tcataaggga actcacttgc ccacttacga agaggccaag caatatctgt      360
cttatgaaac gctctatgcc aatggcagcc gcacagagac gcagggtggc atctacatcc      420
tcagcagtag tggagatggg gcccaacacc gagactcagg gtcttcagga aagtctcgaa      480
ggaagcggca gatttatggc tatgacagca ggttcagcat ttttgggaag gacttctctc      540
tcaactaccc ttttcaaca tcagtgaagt tatccacggg ctgcaccggc accttgggtg      600
cagagaagca tgtcctcaca gctgccact gcatacacga tggaaaaacc tatgtgaaag      660
gaaccagaa gcttcagtg ggcttcttaa agcccaagtt taaagatggt ggtcgagggg      720
ccaacgactc cacttcagcc atgcccagc agatgaaatt tcagtggatc cgggtgaaac      780
gcacccatgt gcccaagggt tggatcaagg gcaatgcaa tgacatcggc atggattatg      840
attatgcctt cctggaactc aaaaagcccc acaagagaaa atttatgaag attgggggtga      900
gccctctctc taagcagctg ccagggggca gaattcaact ctctggttat gacaatgacc      960
gaccaggcaa tttggtgat cgcttctgtg acgtcaaaga cgagacctat gacttctctt     1020
accagcaatg cgatgccccg ccaggggcca gcgggtcttg ggtctatgtg aggatgtgga     1080
agagacagca gcagaagtgg gagcgaaaaa ttattggcat ttttccaggg caccagtggg     1140
tggacatgaa tggttcccca caggatttca acgtggctgt cagaatcact cctctcaaat     1200
atgccagat ttgctatttg attaaaggaa actacctgga ttgtaggag gggtgacaca     1260
gtgttccctc ctggcagcaa ttaagggtct tcatgttctt atttaggag aggccaaatt     1320
gtttttgtc attggcgtgc acacgtgtgt gtgtgtgtgt gtgtgtgtgt aagggtctct     1380
ataatctttt acctatttct tacaattgca agatgactgg ctttactatt tgaaaactgg     1440
tttgtgtatc atatcatata tcatttaagc agtttgaagg catacttttg catagaata     1500
aaaaaaaaac tgatttgggg caatgaggaa tatttgacaa ttaagttaat cttcacgttt     1560
ttgcaaacct tgatttttat ttcatctgaa cttgtttcaa agatttatat taaatatttg     1620
gcatacaaga gaaaaaaaaa aaaaaaa                                     1647
    
```

<210> SEQ ID NO 2

<211> LENGTH: 1877

<212> TYPE: PRT

<213> ORGANISM: Rattus norvegicus

<400> SEQUENCE: 2

```

Cys Gly Cys Gly Cys Gly Cys Gly Thr Cys Cys Thr Cys Gly Gly Gly
 1           5           10          15
Gly Cys Gly Cys Cys Cys Ala Cys Ala Cys Cys Thr Gly Thr Cys Thr
          20          25          30
Gly Ala Gly Cys Gly Gly Cys Gly Cys Ala Cys Gly Gly Cys Cys Gly
          35          40          45
Cys Gly Gly Cys Cys Cys Cys Gly Gly Cys Gly Gly Gly Cys Thr Gly
    
```

-continued

50	55	60																	
Cys	Thr	Cys	Cys	Ala	Cys	Gly	Cys	Gly	Gly	Thr	Ala	Gly	Cys	Ala	Cys				
65					70					75					80				
Thr	Cys	Ala	Gly	Cys	Ala	Thr	Gly	Gly	Cys	Thr	Gly	Gly	Ala	Ala	Thr				
			85						90						95				
Cys	Cys	Cys	Ala	Gly	Gly	Ala	Cys	Thr	Cys	Cys	Thr	Cys	Ala	Thr	Cys				
			100					105						110					
Cys	Thr	Thr	Cys	Thr	Cys	Cys	Thr	Thr	Gly	Thr	Cys	Cys	Thr	Gly	Cys				
		115					120						125						
Thr	Cys	Thr	Gly	Thr	Gly	Thr	Gly	Thr	Thr	Cys	Ala	Thr	Gly	Cys	Ala				
		130					135					140							
Gly	Gly	Thr	Gly	Ala	Gly	Thr	Cys	Cys	Cys	Thr	Ala	Cys	Ala	Ala	Cys				
		145			150					155					160				
Gly	Thr	Thr	Cys	Cys	Gly	Thr	Gly	Gly	Ala	Ala	Ala	Cys	Cys	Ala	Ala				
			165						170					175					
Cys	Ala	Thr	Gly	Gly	Cys	Cys	Gly	Gly	Cys	Thr	Thr	Ala	Cys	Cys	Gly				
			180					185						190					
Cys	Cys	Thr	Cys	Cys	Cys	Thr	Ala	Thr	Ala	Gly	Thr	Cys	Thr	Thr	Gly				
		195					200						205						
Cys	Cys	Thr	Cys	Ala	Gly	Thr	Cys	Thr	Ala	Cys	Cys	Cys	Thr	Cys	Ala				
		210					215						220						
Ala	Gly	Thr	Thr	Ala	Gly	Cys	Cys	Ala	Ala	Ala	Cys	Cys	Ala	Gly	Ala				
		225			230					235					240				
Cys	Thr	Thr	Cys	Gly	Ala	Thr	Gly	Cys	Cys	Ala	Ala	Ala	Gly	Cys	Ala				
			245						250					255					
Ala	Ala	Ala	Thr	Thr	Gly	Gly	Ala	Gly	Gly	Thr	Gly	Thr	Cys	Cys	Thr				
			260					265						270					
Cys	Cys	Thr	Cys	Ala	Thr	Gly	Thr	Gly	Gly	Ala	Cys	Cys	Cys	Cys	Ala				
		275					280						285						
Gly	Thr	Gly	Thr	Cys	Ala	Cys	Ala	Ala	Gly	Gly	Gly	Ala	Ala	Cys	Ala				
		290				295					300								
Cys	Cys	Gly	Cys	Thr	Gly	Cys	Cys	Cys	Ala	Cys	Cys	Thr	Ala	Cys	Gly				
		305			310					315					320				
Ala	Ala	Gly	Ala	Gly	Gly	Cys	Cys	Ala	Ala	Gly	Cys	Ala	Gly	Thr	Ala				
			325						330					335					
Cys	Cys	Thr	Thr	Thr	Cys	Cys	Thr	Ala	Thr	Gly	Ala	Ala	Ala	Cys	Cys				
		340						345						350					
Cys	Thr	Thr	Thr	Ala	Thr	Gly	Cys	Cys	Ala	Ala	Cys	Gly	Gly	Cys	Ala				
		355					360						365						
Gly	Cys	Cys	Gly	Cys	Ala	Cys	Ala	Gly	Ala	Gly	Ala	Cys	Thr	Cys	Ala				
		370				375						380							
Gly	Gly	Thr	Gly	Gly	Gly	Cys	Ala	Thr	Cys	Thr	Ala	Cys	Ala	Thr	Cys				
		385			390					395					400				
Cys	Thr	Cys	Ala	Gly	Cys	Ala	Ala	Cys	Gly	Gly	Thr	Gly	Ala	Ala	Gly				
			405						410					415					
Gly	Cys	Ala	Gly	Gly	Gly	Cys	Ala	Cys	Gly	Cys	Ala	Gly	Cys	Ala	Gly				
			420						425					430					
Ala	Gly	Ala	Cys	Thr	Cys	Gly	Gly	Ala	Gly	Gly	Cys	Cys	Gly	Cys	Ala				
		435					440						445						
Gly	Gly	Gly	Ala	Ala	Ala	Thr	Cys	Thr	Cys	Gly	Cys	Ala	Gly	Gly	Ala				
		450				455						460							

-continued

Ala Gly Ala Gly Gly Cys Ala Gly Ala Thr Thr Thr Ala Thr Gly Gly
465 470 475 480

Cys Thr Ala Thr Gly Ala Thr Gly Gly Cys Ala Gly Ala Thr Thr Thr
485 490 495

Ala Gly Cys Ala Thr Thr Thr Thr Cys Gly Gly Gly Ala Ala Gly Gly
500 505 510

Ala Cys Thr Thr Cys Cys Thr Gly Cys Thr Cys Ala Ala Cys Thr Ala
515 520 525

Thr Cys Cys Cys Thr Thr Cys Thr Cys Ala Ala Cys Ala Thr Cys Gly
530 535 540

Gly Thr Gly Ala Ala Gly Thr Thr Gly Thr Cys Thr Ala Cys Thr Gly
545 550 555 560

Gly Cys Thr Gly Cys Ala Cys Cys Gly Gly Cys Ala Cys Cys Cys Thr
565 570 575

Gly Gly Thr Gly Gly Cys Ala Gly Ala Gly Ala Ala Gly Cys Ala Cys
580 585 590

Gly Thr Cys Cys Thr Cys Ala Cys Thr Gly Cys Thr Gly Cys Cys Cys
595 600 605

Ala Cys Thr Gly Cys Ala Thr Ala Cys Ala Cys Gly Ala Thr Gly Gly
610 615 620

Cys Ala Ala Ala Ala Cys Cys Thr Ala Thr Gly Thr Gly Ala Ala Ala
625 630 635 640

Gly Gly Ala Ala Cys Ala Cys Ala Gly Ala Ala Ala Cys Thr Cys Cys
645 650 655

Gly Ala Gly Thr Gly Gly Gly Cys Thr Thr Cys Cys Thr Gly Ala Ala
660 665 670

Gly Cys Cys Cys Ala Ala Gly Thr Ala Thr Ala Ala Ala Gly Ala Thr
675 680 685

Gly Gly Thr Gly Cys Cys Gly Gly Ala Gly Gly Gly Gly Ala Cys Ala
690 695 700

Ala Cys Ala Gly Thr Thr Cys Ala Ala Gly Cys Thr Cys Ala Gly Cys
705 710 715 720

Thr Cys Thr Gly Gly Thr Ala Gly Ala Gly Ala Ala Gly Ala Thr Gly
725 730 735

Ala Ala Ala Thr Thr Thr Cys Ala Gly Thr Gly Gly Ala Thr Cys Cys
740 745 750

Gly Thr Gly Thr Gly Ala Ala Ala Cys Gly Cys Ala Cys Cys Cys Ala
755 760 765

Thr Gly Thr Gly Cys Cys Cys Ala Ala Gly Gly Gly Gly Thr Gly Gly
770 775 780

Ala Thr Cys Ala Ala Gly Gly Gly Cys Ala Ala Thr Gly Cys Cys Ala
785 790 795 800

Ala Cys Gly Ala Cys Ala Thr Thr Gly Gly Cys Ala Thr Gly Gly Ala
805 810 815

Thr Thr Ala Thr Gly Ala Cys Thr Ala Cys Gly Cys Cys Cys Thr Thr
820 825 830

Cys Thr Gly Gly Ala Ala Cys Thr Cys Ala Ala Gly Ala Ala Ala Cys
835 840 845

Cys Cys Cys Ala Cys Ala Ala Ala Ala Gly Ala Ala Ala Gly Thr Thr
850 855 860

-continued

Cys Ala Thr Gly Ala Ala Gly Ala Thr Thr Gly Gly Thr Gly Thr Gly
 865 870 875 880
 Ala Gly Thr Cys Cys Thr Cys Cys Ala Gly Cys Gly Ala Ala Gly Cys
 885 890 895
 Ala Gly Cys Thr Cys Cys Cys Gly Gly Gly Gly Gly Cys Ala Gly
 900 905 910
 Gly Ala Thr Cys Cys Ala Cys Thr Thr Cys Thr Cys Ala Gly Gly Cys
 915 920 925
 Thr Ala Cys Gly Ala Cys Ala Ala Thr Gly Ala Cys Cys Gly Gly Cys
 930 935 940
 Cys Cys Gly Gly Cys Ala Ala Thr Thr Thr Gly Gly Thr Gly Thr Ala
 945 950 955 960
 Cys Cys Gly Cys Thr Thr Cys Thr Gly Thr Gly Ala Thr Gly Thr Cys
 965 970 975
 Ala Ala Ala Gly Ala Thr Gly Ala Gly Ala Cys Ala Thr Ala Cys Gly
 980 985 990
 Ala Cys Cys Thr Thr Cys Thr Thr Thr Ala Thr Cys Ala Gly Cys Ala
 995 1000 1005
 Gly Thr Gly Thr Gly Ala Thr Gly Cys Cys Cys Ala Gly Cys Cys
 1010 1015 1020
 Cys Gly Gly Gly Gly Cys Cys Ala Gly Thr Gly Gly Thr Thr Cys
 1025 1030 1035
 Ala Gly Gly Gly Thr Cys Thr Ala Thr Gly Thr Gly Ala Gly
 1040 1045 1050
 Gly Ala Thr Gly Thr Gly Gly Ala Ala Gly Ala Gly Ala Cys Cys
 1055 1060 1065
 Ala Cys Ala Gly Cys Ala Gly Ala Ala Ala Thr Gly Gly Gly Ala
 1070 1075 1080
 Ala Ala Gly Gly Ala Ala Ala Thr Thr Ala Thr Cys Gly Gly
 1085 1090 1095
 Cys Ala Thr Cys Thr Thr Thr Thr Cys Ala Gly Gly Gly Cys Ala
 1100 1105 1110
 Cys Cys Ala Gly Thr Gly Gly Gly Thr Gly Gly Ala Cys Ala Thr
 1115 1120 1125
 Gly Ala Ala Thr Gly Gly Cys Thr Cys Thr Cys Cys Gly Cys Ala
 1130 1135 1140
 Gly Gly Ala Thr Thr Thr Cys Ala Ala Cys Gly Thr Gly Gly Cys
 1145 1150 1155
 Ala Gly Thr Thr Ala Gly Ala Ala Thr Cys Ala Cys Ala Cys Cys
 1160 1165 1170
 Thr Cys Thr Thr Ala Ala Ala Thr Ala Thr Gly Cys Cys Cys Ala
 1175 1180 1185
 Gly Ala Thr Thr Thr Gly Cys Thr Ala Thr Thr Gly Gly Ala Thr
 1190 1195 1200
 Thr Ala Ala Ala Gly Gly Ala Ala Ala Cys Thr Ala Cys Cys Thr
 1205 1210 1215
 Gly Gly Ala Thr Thr Gly Cys Ala Gly Gly Gly Ala Gly Gly Gly
 1220 1225 1230
 Gly Thr Gly Ala Cys Ala Thr Gly Cys Gly Thr Cys Thr Thr Cys
 1235 1240 1245
 Thr Thr Gly Cys Cys Ala Gly Cys Ala Cys Cys Gly Ala Thr Gly

-continued

1250	1255	1260
Gly Thr Cys Gly Thr Thr Thr Thr Gly Cys Ala Cys Thr Cys Ala		
1265	1270	1275
Thr Thr Gly Thr Ala Gly Gly Ala Gly Ala Gly Ala Gly Gly Cys		
1280	1285	1290
Cys Ala Gly Cys Thr Thr Thr Ala Thr Ala Thr Cys Ala Thr Thr		
1295	1300	1305
Gly Ala Cys Thr Cys Thr Thr Gly Thr Gly Gly Thr Gly Thr Gly		
1310	1315	1320
Thr Gly Thr Cys Ala Cys Ala Thr Ala Gly Thr Ala Thr Cys Thr		
1325	1330	1335
Thr Thr Thr Ala Cys Cys Thr Ala Ala Thr Ala Thr Thr Cys Thr		
1340	1345	1350
Thr Cys Ala Ala Ala Thr Ala Gly Cys Ala Ala Ala Ala Cys Thr		
1355	1360	1365
Ala Thr Thr Gly Gly Thr Thr Cys Thr Ala Thr Thr Ala Thr Thr		
1370	1375	1380
Thr Gly Ala Ala Ala Ala Thr Thr Gly Thr Thr Gly Thr Gly Thr		
1385	1390	1395
Gly Cys Gly Thr Thr Ala Thr Ala Gly Cys Ala Thr Thr Thr Ala		
1400	1405	1410
Ala Gly Cys Ala Gly Thr Cys Thr Gly Ala Ala Ala Gly Cys Ala		
1415	1420	1425
Thr Ala Cys Thr Thr Thr Thr Thr Ala Cys Ala Thr Ala Gly Ala		
1430	1435	1440
Gly Ala Cys Thr Thr Thr Ala Ala Ala Ala Thr Ala Thr Thr Gly		
1445	1450	1455
Ala Thr Gly Thr Thr Cys Gly Gly Gly Thr Thr Ala Ala Thr Ala		
1460	1465	1470
Gly Gly Gly Gly Ala Thr Ala Thr Thr Thr Gly Gly Cys Ala Ala		
1475	1480	1485
Gly Thr Ala Ala Ala Thr Thr Ala Ala Ala Cys Thr Thr Thr Cys		
1490	1495	1500
Ala Gly Cys Thr Thr Thr Thr Gly Gly Ala Gly Ala Gly Cys Thr		
1505	1510	1515
Cys Thr Gly Ala Thr Thr Thr Thr Thr Thr Gly Cys Cys Thr Thr Ala		
1520	1525	1530
Thr Cys Cys Ala Ala Ala Cys Ala Thr Gly Cys Thr Thr Cys Ala		
1535	1540	1545
Gly Ala Gly Gly Thr Thr Thr Ala Thr Ala Thr Cys Ala Ala Ala		
1550	1555	1560
Thr Ala Thr Gly Thr Gly Ala Cys Ala Thr Ala Cys Cys Gly Gly		
1565	1570	1575
Gly Ala Ala Thr Ala Thr Gly Gly Ala Thr Thr Cys Thr Thr Ala		
1580	1585	1590
Thr Gly Thr Thr Thr Gly Thr Ala Thr Ala Thr Gly Thr Ala Thr		
1595	1600	1605
Ala Thr Gly Thr Cys Thr Thr Cys Thr Gly Ala Gly Ala Gly Thr		
1610	1615	1620
Cys Ala Thr Ala Cys Ala Thr Ala Thr Ala Thr Thr Gly Ala Thr		
1625	1630	1635

-continued

Ala Thr Thr Thr Thr Thr Thr Thr Ala Ala Thr Gly Thr Gly Thr
 1640 1645 1650

Gly Gly Thr Thr Ala Thr Ala Gly Thr Gly Thr Thr Thr Cys Cys
 1655 1660 1665

Ala Ala Ala Thr Gly Ala Cys Ala Ala Thr Ala Gly Cys Ala Ala
 1670 1675 1680

Ala Gly Cys Thr Thr Gly Cys Ala Ala Thr Gly Gly Gly Cys Ala
 1685 1690 1695

Ala Thr Cys Ala Thr Ala Ala Thr Gly Thr Thr Thr Cys Cys Gly
 1700 1705 1710

Ala Thr Thr Cys Ala Ala Ala Cys Ala Thr Thr Thr Ala Cys Ala
 1715 1720 1725

Thr Ala Gly Thr Ala Gly Thr Cys Cys Thr Thr Gly Ala Ala Gly
 1730 1735 1740

Ala Gly Ala Ala Ala Cys Ala Gly Thr Ala Ala Thr Thr Thr Ala
 1745 1750 1755

Thr Thr Gly Gly Cys Thr Cys Thr Ala Thr Thr Gly Ala Thr Ala
 1760 1765 1770

Cys Ala Cys Ala Thr Gly Thr Ala Ala Ala Gly Cys Thr Gly Thr
 1775 1780 1785

Ala Thr Cys Thr Thr Ala Cys Ala Gly Thr Gly Cys Ala Gly Ala
 1790 1795 1800

Gly Ala Ala Thr Thr Cys Cys Cys Ala Gly Gly Cys Thr Gly Cys
 1805 1810 1815

Thr Thr Thr Thr Ala Gly Thr Thr Thr Thr Gly Ala Ala Ala Ala
 1820 1825 1830

Thr Ala Ala Ala Ala Cys Thr Cys Thr Gly Thr Cys Thr Thr Cys
 1835 1840 1845

Thr Ala Ala Ala Ala Ala Ala Ala Ala Ala Ala Ala Ala Ala Ala
 1850 1855 1860

Ala Ala Ala Ala Ala Ala Ala Ala Ala Ala Ala Ala Ala Ala
 1865 1870 1875

<210> SEQ ID NO 3
 <211> LENGTH: 382
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 3

Met Ala Gly Ile Pro Gly Leu Leu Phe Leu Leu Phe Phe Leu Leu Cys
 1 5 10 15

Ala Val Gly Gln Val Ser Pro Tyr Ser Ala Pro Trp Lys Pro Thr Trp
 20 25 30

Pro Ala Tyr Arg Leu Pro Val Val Leu Pro Gln Ser Thr Leu Asn Leu
 35 40 45

Ala Lys Pro Asp Phe Gly Ala Glu Ala Lys Leu Glu Val Ser Ser Ser
 50 55 60

Cys Gly Pro Gln Cys His Lys Gly Thr Pro Leu Pro Thr Tyr Glu Glu
 65 70 75 80

Ala Lys Gln Tyr Leu Ser Tyr Glu Thr Leu Tyr Ala Asn Gly Ser Arg
 85 90 95

Thr Glu Thr Gln Val Gly Ile Tyr Ile Leu Ser Ser Ser Gly Asp Gly

-continued

100			105			110									
Ala	Gln	His	Arg	Asp	Ser	Gly	Ser	Ser	Gly	Lys	Ser	Arg	Arg	Lys	Arg
	115						120					125			
Gln	Ile	Tyr	Gly	Tyr	Asp	Ser	Arg	Phe	Ser	Ile	Phe	Gly	Lys	Asp	Phe
	130						135					140			
Leu	Leu	Asn	Tyr	Pro	Phe	Ser	Thr	Ser	Val	Lys	Leu	Ser	Gly	Cys	Thr
	145				150					155					160
Gly	Thr	Leu	Val	Ala	Glu	Lys	His	Val	Leu	Thr	Ala	Ala	His	Cys	Ile
			165						170						175
His	Asp	Gly	Lys	Thr	Tyr	Val	Lys	Gly	Thr	Gln	Lys	Leu	Arg	Val	Gly
			180						185				190		
Phe	Leu	Lys	Pro	Lys	Phe	Lys	Asp	Gly	Gly	Arg	Gly	Ala	Asn	Asp	Ser
			195					200				205			
Thr	Ser	Ala	Met	Pro	Glu	Gln	Met	Lys	Phe	Gln	Trp	Ile	Arg	Val	Lys
			210				215					220			
Arg	Thr	His	Val	Pro	Lys	Gly	Trp	Ile	Lys	Gly	Asn	Ala	Asn	Asp	Ile
					230					235					240
Gly	Met	Asp	Tyr	Asp	Tyr	Ala	Leu	Leu	Glu	Leu	Lys	Lys	Pro	His	Lys
				245					250						255
Arg	Lys	Phe	Met	Lys	Ile	Gly	Val	Ser	Pro	Pro	Ala	Lys	Gln	Leu	Pro
			260						265				270		
Gly	Gly	Arg	Ile	His	Phe	Ser	Gly	Tyr	Asp	Asn	Asp	Arg	Pro	Gly	Asn
			275						280			285			
Leu	Val	Tyr	Arg	Phe	Cys	Asp	Val	Lys	Asp	Glu	Thr	Tyr	Asp	Leu	Leu
			290				295				300				
Tyr	Gln	Gln	Cys	Asp	Ala	Gln	Pro	Gly	Ala	Ser	Gly	Ser	Gly	Val	Tyr
					310				315						320
Val	Arg	Met	Trp	Lys	Arg	Gln	Gln	Gln	Lys	Trp	Glu	Arg	Lys	Ile	Ile
				325					330						335
Gly	Ile	Phe	Ser	Gly	His	Gln	Trp	Val	Asp	Met	Asn	Gly	Ser	Pro	Gln
			340						345				350		
Asp	Phe	Asn	Val	Ala	Val	Arg	Ile	Thr	Pro	Leu	Lys	Tyr	Ala	Gln	Ile
			355				360					365			
Cys	Tyr	Trp	Ile	Lys	Gly	Asn	Tyr	Leu	Asp	Cys	Arg	Glu	Gly		
			370				375					380			

<210> SEQ ID NO 4
 <211> LENGTH: 383
 <212> TYPE: PRT
 <213> ORGANISM: Rattus norvegicus

<400> SEQUENCE: 4

Met	Ala	Gly	Ile	Pro	Gly	Leu	Leu	Ile	Leu	Leu	Leu	Val	Leu	Leu	Cys
1				5					10						15
Val	Phe	Met	Gln	Val	Ser	Pro	Tyr	Asn	Val	Pro	Trp	Lys	Pro	Thr	Trp
			20					25					30		
Pro	Ala	Tyr	Arg	Leu	Pro	Ile	Val	Leu	Pro	Gln	Ser	Thr	Leu	Lys	Leu
			35				40					45			
Ala	Lys	Pro	Asp	Phe	Asp	Ala	Lys	Ala	Lys	Leu	Glu	Val	Ser	Ser	Ser
			50			55					60				
Cys	Gly	Pro	Gln	Cys	His	Lys	Gly	Thr	Pro	Leu	Pro	Thr	Tyr	Glu	Glu
					70					75					80

-continued

Ala Lys Gln Tyr Leu Ser Tyr Glu Thr Leu Tyr Ala Asn Gly Ser Arg
85 90 95

Thr Glu Thr Gln Val Gly Ile Tyr Ile Leu Ser Asn Gly Glu Gly Arg
100 105 110

Ala Arg Ser Arg Asp Ser Glu Ala Ala Gly Lys Ser Arg Arg Lys Arg
115 120 125

Gln Ile Tyr Gly Tyr Asp Gly Arg Phe Ser Ile Phe Gly Lys Asp Phe
130 135 140

Leu Leu Asn Tyr Pro Phe Ser Thr Ser Val Lys Leu Ser Thr Gly Cys
145 150 155 160

Thr Gly Thr Leu Val Ala Glu Lys His Val Leu Thr Ala Ala His Cys
165 170 175

Ile His Asp Gly Lys Thr Tyr Val Lys Gly Thr Gln Lys Leu Arg Val
180 185 190

Gly Phe Leu Lys Pro Lys Tyr Lys Asp Gly Ala Gly Gly Asp Asn Ser
195 200 205

Ser Ser Ser Ala Leu Val Glu Lys Met Lys Phe Gln Trp Ile Arg Val
210 215 220

Lys Arg Thr His Val Pro Lys Gly Trp Ile Lys Gly Asn Ala Asn Asp
225 230 235 240

Ile Gly Met Asp Tyr Asp Tyr Ala Leu Leu Glu Leu Lys Lys Pro His
245 250 255

Lys Arg Lys Phe Met Lys Ile Gly Val Ser Pro Pro Ala Lys Gln Leu
260 265 270

Pro Gly Gly Arg Ile His Phe Ser Gly Tyr Asp Asn Asp Arg Pro Gly
275 280 285

Asn Leu Val Tyr Arg Phe Cys Asp Val Lys Asp Glu Thr Tyr Asp Leu
290 295 300

Leu Tyr Gln Gln Cys Asp Ala Gln Pro Gly Ala Ser Gly Ser Gly Val
305 310 315 320

Tyr Val Arg Met Trp Lys Arg Pro Gln Gln Lys Trp Glu Arg Lys Ile
325 330 335

Ile Gly Ile Phe Ser Gly His Gln Trp Val Asp Met Asn Gly Ser Pro
340 345 350

Gln Asp Phe Asn Val Ala Val Arg Ile Thr Pro Leu Lys Tyr Ala Gln
355 360 365

Ile Cys Tyr Trp Ile Lys Gly Asn Tyr Leu Asp Cys Arg Glu Gly
370 375 380

<210> SEQ ID NO 5
<211> LENGTH: 17
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: primer 1

<400> SEQUENCE: 5

cgccacacacc tgtctga

17

<210> SEQ ID NO 6
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: primer 2

-continued

```

<400> SEQUENCE: 6
ctcggcgcgagg aacagtgtctc                                     20

<210> SEQ ID NO 7
<211> LENGTH: 17
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: probe 1

<400> SEQUENCE: 7
tggaatccct gccatgc                                         17

<210> SEQ ID NO 8
<211> LENGTH: 17
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: primer 3

<400> SEQUENCE: 8
cgcccacacc tgtctga                                         17

<210> SEQ ID NO 9
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: primer 4

<400> SEQUENCE: 9
ctgctccacg cggtagcact ca                                   22

<210> SEQ ID NO 10
<211> LENGTH: 17
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: probe 2

<400> SEQUENCE: 10
tgggattcca gccatgc                                         17

```

1. A method of using PRSS23 as a biomarker for a disease, comprising:

- i) obtaining a biological sample from a mammal,
- ii) measuring the level of PRSS23 in the biological sample,
- iii) comparing the level of PRSS23 in the biological sample with the level of PRSS23 in a control sample, and
- iv) diagnosing a disease based upon the PRSS23 level of the biological sample in comparison to the control sample.

2. A method of using PRSS23 as a biomarker for guiding a therapy of a disease, comprising:

- i) obtaining a baseline level of PRSS23 in biological sample from a diseased mammal,
- ii) administering to the diseased mammal a treatment for the disease,
- iii) obtaining one or more subsequent biological samples from the diseased mammal

iv) measuring the level of PRSS23 in the one or more subsequent biological samples,

v) comparing the level of PRSS23 in the one or more subsequent biological samples with the baseline sample, and

vi) determining whether increased dosages, additional or alternative treatments are necessary based on PRSS23 levels obtained from one or more subsequent biological samples compared to the baseline PRSS23 level.

3. The method of claim 1 wherein the biological sample is selected from the group consisting of a blood sample, a plasma sample, a serum sample, a tissue sample, a oral mucosa sample, a saliva sample, an interstitial fluid sample, and a urine sample.

4. The method of claim 1 wherein the level of PRSS23 is determined by determining the level of PRSS23 polynucleotide.

5. The method of claim 1 wherein the level of PRSS23 is determined by determining the level of PRSS23 polypeptide.

6. The method of claim 1 wherein the level of PRSS23 is determined by determining the level of PRSS23 activity.

7. The method of claim 1 wherein the disease is selected from the group consisting of cardiovascular diseases, hematological diseases, neurological diseases, respiratory diseases, gastroenterological diseases, and urological diseases.

8. The method of claim 1 wherein the mammal is a human.

9. The method of claim 1 wherein the level of PRSS23 of the biological sample is elevated compared to the control sample.

10. The method of claim 1 wherein the use of PRSS23 is combined with the use of one or more biomarkers.

11. The method of claim 1 wherein the use of PRSS23 is combined with the use of one or more biomarkers which are selected from the group consisting of CRTAC, FN1, NPR3, LTBP2, TGFB2, and CTGF.

12. The method of claim 1 wherein the use of PRSS23 is combined with the use of one or more biomarkers which are selected from the group consisting of BNP, ANP, Troponin, CRP, Myoglobin, CK-MB, and metabolites thereof.

13. The method of claim 1 wherein the use of PRSS23 is combined with the use of one or more clinical biomarkers which are selected from the group consisting of blood pressure, heart rate, pulmonary artery pressure, and systemic vascular resistance.

14. The method of claim 1 wherein the use of PRSS23 is combined with the use of one or more diagnostic imaging methods which are selected from the group consisting of PET (Positron Emission Tomography), CT (Computed Tomography), ultrasonic, SPECT (Single Photon Emission Computed Tomography), Echocardiography, and Impedance Cardiography.

15. A kit for identifying an individual risk for developing a disease, for predicting a disease or an adverse outcome in a patient diagnosed with a disease, or for guiding a therapy in a patient with a disease, the kit comprising one or more antibodies which specifically bind PRSS23, one or more containers for collecting and or holding a biological sample, and an instruction for its use, wherein the disease is associated with an altered PRSS23 level.

16. A kit for identifying an individual risk for developing a disease, for predicting a disease or an adverse outcome in a patient diagnosed with a disease, or for guiding a therapy in a patient with a disease, the kit comprising one or more probes or primers for detecting PRSS23 mRNA, one or more containers for collecting and or holding a biological sample, and an instruction for its use, wherein the disease is associated with an altered PRSS23 level.

* * * * *

专利名称(译)	PRSS23作为生物标志物，治疗和诊断目标		
公开(公告)号	US20100104501A1	公开(公告)日	2010-04-29
申请号	US12/445494	申请日	2007-10-02
[标]申请(专利权)人(译)	GOLZ STEFAN SUMMER HOLGER GEERTS ANDREAS BRUEGGEMEIER ULF ALBRECHT BARBARA KLEIN MARTINA STEPPAN SONJA ELLINGHAUS PETER ðURSO DONATELLA SEEWALD MICHAEL MILTING HENDRIK		
申请(专利权)人(译)	GOLZ STEFAN SUMMER HOLGER GEERTS ANDREAS BRUEGGEMEIER ULF ALBRECHT BARBARA KLEIN MARTINA STEPPAN SONJA ELLINGHAUS PETER ðURSO DONATELLA SEEWALD MICHAEL MILTING HENDRIK		
当前申请(专利权)人(译)	拜耳知识产权GMBH		
[标]发明人	GOLZ STEFAN SUMMER HOLGER GEERTS ANDREAS BRUGGEMEIER ULF ALBRECHT BARBARA KLEIN MARTINA STEPPAN SONJA ELLINGHAUS PETER DURSO DONATELLA SEEWALD MICHAEL MILTING HENDRIK		
发明人	GOLZ, STEFAN SUMMER, HOLGER GEERTS, ANDREAS BRUGGEMEIER, ULF ALBRECHT, BARBARA KLEIN, MARTINA STEPPAN, SONJA ELLINGHAUS, PETER D'URSO, DONATELLA SEEWALD, MICHAEL MILTING, HENDRIK		

