



US 20110003318A1

(19) **United States**(12) **Patent Application Publication**
Ervin, JR.(10) **Pub. No.: US 2011/0003318 A1**(43) **Pub. Date: Jan. 6, 2011**(54) **NUCLEOTIDE AND PROTEIN SEQUENCE OF
MAMMASTATIN AND METHODS OF USE**(75) Inventor: **Paul R. Ervin, JR.**, Ann Arbor, MI
(US)


Correspondence Address:

Casimir Jones, S.C.**2275 DEMING WAY, SUITE 310****MIDDLETON, WI 53562 (US)**(73) Assignee: **THE REGENTS OF THE
UNIVERSITY OF MICHIGAN,**
ANN ARBOR, MI (US)(21) Appl. No.: **12/846,386**(22) Filed: **Jul. 29, 2010****Related U.S. Application Data**(63) Continuation of application No. 11/838,654, filed on
Aug. 14, 2007, now Pat. No. 7,816,097, which is a
continuation of application No. 10/459,218, filed on
Jun. 11, 2003, now Pat. No. 7,256,277, which is a
continuation of application No. 09/643,476, filed on
Aug. 22, 2000, now Pat. No. 6,599,495, which is acontinuation of application No. 08/994,076, filed on
Dec. 19, 1997, now Pat. No. 6,500,937, which is a
continuation-in-part of application No. 08/943,828,
filed on Oct. 3, 1997, now abandoned.(60) Provisional application No. 60/027,315, filed on Oct.
3, 1996.**Publication Classification**(51) **Int. Cl.****G01N 33/53** (2006.01)**C07K 16/18** (2006.01)(52) **U.S. Cl. 435/7.21; 436/501; 435/7.92; 530/388.1**

(57)

ABSTRACT

A nucleic acid sequence encoding Mammastatin, a specific mammary cell growth inhibitor. Mammastatin is encoded by a single nucleic acid sequence and has an approximate molecular weight of 44 kDa in its inactive, non-phosphorylated form. Normal mammary cells express functional phosphorylated forms having approximate molecular weights of 53 kDa and 49 kDa. Metastatic mammary cells either do not express Mammastatin at all, or do not express the 53 kDa or 49 kDa, phosphorylated forms. Mammary cancer cells are inhibited in their growth by the administration of phosphorylated mammastatin.

Expression of Recombinant Mammastatin in Cos-7 Cells

Expression of Recombinant Mammastatin in Cos-7 Cells




FIG. 1




FIG. 2

Figure 3

Expression of Recombinant Mammastatin



MCF-7 Cell Growth with Mammastatin Transcription and Translation Products





Figure 4**Growth Inhibitory Activity in Cos-7 Cell Conditioned Media.**




FIG. 5




FIG. 6


Figure 7

FIG. 8




FIG. 9

Mammastatin Standard Curve






Figure 10**Variation of Mammastatin
in Normal Female Serum**

Figure 11**Variation of Mammastatin in FO Patient Serum During Treatment**




FIG. 12




FIG. 13A




FIG. 13B




FIG. 13C




FIG. 14A




FIG. 14B





FIG. 14C

Figure 15

Serum Mammastatin Dot Blot Assay



2




FIG. 16

NUCLEOTIDE AND PROTEIN SEQUENCE OF MAMMASTATIN AND METHODS OF USE

[0001] This is a Continuation-in-Part of U.S. patent application Ser. No. 08/943,828 filed on Oct. 3, 1997, which is hereby incorporated by reference for all purposes.

FIELD OF THE INVENTION

[0002] This invention relates to compositions and methods for the diagnosis and treatment of breast cancer. More particularly, the invention is related to nucleic acid sequences encoding Mammastatin, a protein useful in the diagnosis and treatment of breast cancer,

BACKGROUND OF THE INVENTION

[0003] Breast cancer is a disease that kills over 45,000 women each year in the United States alone. Over 180,000 new cases of breast cancer are diagnosed annually, and it is estimated that one in eight women will develop breast cancer. These numbers indicate that breast cancer is one of the most dangerous diseases facing women today. Cancer research has been unable to determine the cause of breast cancer, and has not found a suitable method of therapy or prevention.

[0004] A woman diagnosed with breast cancer may be treated with surgery, hormone therapy, chemotherapy, and radiation. If the patient develops metastatic disease, radiation and high dose chemotherapy are required to ablate the cancer in remote areas such as the brain, bone, and liver.

[0005] The current therapies available for the treatment of breast cancer are toxic, dangerous, costly, and many are ineffective, especially in the treatment of metastatic disease. The table below was extracted from Churchill Livingston, Clinical Oncology, 1995, and summarizes data available on the current methods of treatment and expected survival rates.

[0006] Currently, there are no therapies that are effective for long term treatment of breast cancer that has metastasized to lymph nodes or distal sites. Local disease can be effectively treated by surgery, if all of the cancer can be removed. A new therapy for the effective treatment of breast cancer that could stop the growth of breast cancer and of cells derived from metastatic cancer is urgently needed. Such a therapy would be useful in the treatment of localized breast cancer, in long term treatment of metastatic disease, and as a follow-up treatment after surgical removal of tumors. Other applications include a growth inhibitor as a primary therapy and for preventative use.

[0007] Detection methods for breast cancer, such as mammogram, physical exam, CAT-scan, and ultrasound, have significantly improved early detection of breast cancer. However, with these methods, a suspected tumor must still be surgically removed for pathological examination to determine if the tumor is benign or malignant, and to attempt to determine the tissue type and grade of the malignancy. This pathological diagnosis helps to determine what subsequent treatment protocols may be used.

[0008] For breast cancer, these methods are generally inconclusive, as adequate breast cancer tumor markers are not available. Available markers such as CA 15-3 and CA 27-29 are used as indicators of metastases, however, they are not specific. There is a great need for diagnostic tools and methods that can effectively and reliably diagnose breast cancer, e.g., using new and specific breast cancer markers. In addition, a reliable and simple method for the early detection and diagnosis of breast cancer is greatly needed. Preferably, such an early detection method would identify breast cancer in its early stages, track progression of breast cancer through advanced metastatic disease, and diagnose the propensity of a patient to develop breast cancer or to develop advanced dis-

Treatment	Method	Effect	Toxicity	Result	Survival
adriamycin	bolus	kill cancer cells	high	can induce remission	+14 months
cyclophosphate	bolus	kill cancer cells	high	can induce remission	+16 months
methotrexate	infusion	kill cancer cells	high	can induce remission	+16 months
5F uracil	infusion	kill cancer cells	high	can induce remission	+18 months
mix of above	mixed	kill cancer cells	high	can induce remission	+22 months
taxol	bolus	kill cancer cells	high	can induce remission	+12 months
estrogen	oral	may stop growth	low	can induce remission	+6 months
tamoxifen	oral	may stop growth	low	may stop progression	+12 months
mastectomy	surgery	remove tumor	low	may eliminate cancer	+5 years*
lumpectomy	surgery	remove tumor	low	may eliminate cancer	+5 years*
surgery and tamoxifen	combination	combination	low	may eliminate cancer	+7-10 years*
radiation	mechanical	kill cancer cells	high	can induce remission	+14 months

*assumes there are no micrometastases

ease. Most preferably, the diagnostic method could be used without tissue biopsy, e.g., by analysis of a body fluid such as blood.

[0009] Human mammary tissues undergo a burst of proliferative activities at the onset of menarche and during each menstrual cycle. Studies on the effects of estrogen on mammary tissues and tumors indicate that estrogen is a primary growth-initiating factor for mammary tissues. Estradiol-sensitive growth factors have been characterized. In addition, mammary cell growth factors which are not hormonal in nature have also been described.

[0010] Specific growth factors which have been shown to have a stimulating effect on mammary tissue growth include platelet-derived growth factor (PDGF), insulin-like growth factor (IGF-1) and transforming growth factor (TGF) alpha. TGF-beta, on the other hand, has been shown to suppress mammary tissue growth.

[0011] The regulation of mammary cell growth is of great importance in the diagnosis and treatment of breast cancer. Neoplastic growth of mammary tissues, if unchecked, can develop into uncontrollably-proliferating malignant tumors, which are the cause of death of thousands of women yearly. A growth inhibition factor capable of specifically suppressing mammary cell growth would provide a dynamic tool for use in the diagnosis and treatment of breast cancer.

[0012] Thus, it would be of great utility to isolate and characterize a specific mammary cell growth inhibitor, to identify its nucleic acid sequence and amino acid sequence, and to recombinantly express the inhibitor as a purified protein. Diagnostic and therapeutic methods using the nucleic acid sequence and/or recombinantly produced inhibitor would be of great utility in the diagnosis and treatment of breast cancer.

SUMMARY OF THE INVENTION

[0013] A specific mammary cell growth inhibitor, Mammastatin, has been isolated from normal human mammary cells and characterized. It has now been found that Mammastatin is produced by normal mammary cells, but not by breast cancer cells. Furthermore, it has now been found that the reduction or absence of Mammastatin in the blood correlate with the presence of breast cancer. Administration of active Mammastatin prevents growth of breast cancer cells.

[0014] The nucleic acid sequence encoding Mammastatin has now been cloned, sequenced, and expressed recombinantly in host cells as an active inhibitor of mammary cell growth. The isolated and characterized nucleic acid sequence (Sequence ID No: 1) and its deduced amino acid sequence (Sequence ID No: 2) provide unique and specific tools for use in the diagnosis and treatment of breast cancer.

[0015] The present invention provides an isolated and purified nucleic acid sequence encoding Mammastatin, a specific protein inhibitor of mammary cell growth, and particularly of mammary cancer cell growth. The invention also includes plasmids and vectors containing the Mammastatin nucleic acid sequence, amino acid sequence of Mammastatin, and methods, kits, and compositions utilizing the Mammastatin nucleic acid or amino acid sequences to produce purified mammary cell growth inhibitor and in the diagnosis and treatment of breast cancer. The inventive compositions include probes and primers that specifically hybridize to the Mammastatin nucleic acid sequence and its RNA products.

[0016] The invention further includes a method for treating breast cancer by administering Mammastatin.

BRIEF DESCRIPTION OF THE DRAWINGS

[0017] FIG. 1 is a Western Blot showing expression of recombinant Mammastatin in Eucaryotic Cos-7 cells.

[0018] FIG. 2 is an immunoblot showing expression of Mammastatin in insect cells.

[0019] FIG. 3 is a graph showing inhibition of mammary cell growth by recombinant Mammastatin produced by in vitro transcription and translation.

[0020] FIG. 4 is a graph showing growth inhibition in human mammary cancer cell growth by treatment with conditioned medium of Cos-7 cells transfected with Mammastatin cDNA.

[0021] FIG. 5 is a Western Blot showing relative amounts of 53, 49 and 44 kD Mammastatin in normal and cancerous human mammary cells.

[0022] FIG. 6 is an immunoblot showing phosphatase digestion of Mammastatin.

[0023] FIG. 7 is a graph showing the effect of phosphatase on the activity of Mammastatin.

[0024] FIG. 8 is a Western Blot showing Mammastatin from normal and cancerous human mammary cells, as well as in mixed cultures of normal and cancerous cells.

[0025] FIG. 9 is a graph showing Mammastatin in normal human serum as analyzed by ELISA.

[0026] FIG. 10 is a graph showing a Mammastatin ELISA standard curve.

[0027] FIG. 11 is a graph showing Mammastatin levels in breast cancer patients over the course of treatment.

[0028] FIG. 12 is a Western blot showing expression of Mammastatin induced by retrovirus.

[0029] FIGS. 13A, 13B and 13C are graphs showing the effect of Mammastatin treatment on MCF7 tumor cells in nude mice.

[0030] FIGS. 14A, 14B and 14C are graphs showing the effect of Mammastatin treatment on tumor cells in nude mice.

[0031] FIG. 15 is a dot blot assay showing Mammastatin in blood from normal females versus the absence of Mammastatin in blood from breast cancer patients.

[0032] FIG. 16 is an immunoblot of mammalian cells expressing recombinant mammastatin.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

Mammastatin

[0033] Mammastatin is a protein growth inhibitor produced and secreted by normal human mammary epithelial cells. A mammary cell growth inhibitor was first described as an inhibitory protein activity present in media conditioned by the growth of normal human mammary cells. The inhibitory activity was identified in conditioned medium from normal human mammary cells, but not in media conditioned by the growth of human mammary cancer cells. The inhibitory activity was determined by bioassay and antibody development to reside in three proteins, having the approximate molecular weights of 53, 49 and 44 kDa (Ervin, Paul R., Doctoral Dissertation University of Michigan, 1995).

[0034] It has now been determined that a specific mammary cell growth inhibitor, Mammastatin, is expressed as a 44 kD protein which is phosphorylatedly increasing the molecular weight to 49 kD and 53 kD. The non-ph 44 kD form is not an

active inhibitor, whereas the phosphorylate 49 kD and 53 kD forms inhibit growth of breast cancer cells. The active 53 and/or 49 kD phosphoprotein is expressed by normal human mammary cells, but is not generally produced by human mammary carcinoma cells. Some carcinoma cells make the 4410 protein that lacks phosphorylation and is inactive.

[0035] The table below summarizes data showing expression and activity of Mammastatin in normal and cancerous cells and tissues.

Cell Type*	Number	44 kDa**	53/49 kDa**	produce inhibitor	are inhibited
Normal primary cultures	42	+/-	++++	42/42	2/2
Normal breast tissue	5	+	++		
Mammary Cell Line	16			0/16	12/12
Type A	11	+	-	0/11	8/8
Type B	5	-	-	0/5	4/4
Breast tumor lysate	25				
Type A	17	+	-		
Type B	8	-	-		
Non-mammary cell lines	8			0/2	0/8
Type A	3	+	-***	0/1	0/3
Type B	5	-	-***	0/1	0/5

*Carcinoma cells in which 44 kDa Mammastatin was detected (Type A) or not (Type B)

**(-) No expression (+++++) intense expression

***Two cell lines, BxPc3 and A253 expressed proteins identified as 53/49 kD, but neither cell line produced inhibitory activity.

[0036] Dose response studies with human mammary carcinoma cells indicates that carcinoma cell growth is 50-70% inhibited with 10 ng/ml of Mammastatin and blocked completely with 25-50 ng/ml. Highly metastatic cells such as MDA-MB-435 and MDA-MB-231 required 50 ng/ml to stop growth. In vitro and in vivo clinical data experiments indicate the effect is reversible, and that repeated administration of the inhibitor is required to arrest carcinoma cell growth at the lower concentrations. At doses above 50 ng/ml, however, Mammastatin appears to induce apoptosis, as indicated by histology, e.g. cell necrosis.

[0037] Since Mammastatin is a natural growth inhibitor that blocks mammary carcinoma cell growth, and since no tumors make active Mammastatin, Mammastatin replacement therapy is ideal for therapeutic treatment of breast cancer. The clinical data provided in the examples below demonstrate the effectiveness of Mammastatin replacement therapy.

[0038] The nucleic acid sequence encoding Mammastatin protein has now been isolated, characterized, sequenced (Sequence ID NO:1), determined to encode all three (53, 49, and 44 kD) molecular weight proteins, and given the name "Mammastatin". Differences in the molecular weight of the three forms has been determined to be caused by the extent of the protein's phosphorylation. Mammastatin produced by normal human mammary cells (NHMC) in culture and recombinantly expressed Mammastatin inhibit the growth of human mammary carcinoma cells, and is useful as a therapeutic agent in the treatment of breast cancer.

[0039] Analysis of human sera from normal women and from breast cancer patients indicates that decreased blood levels of Mammastatin correlate with advancing breast cancer. Screening and monitoring blood serum for the presence

of this active inhibitor as described in the examples below provides a specific and effective diagnostic tool.

Nucleic Acid Sequence

[0040] The nucleic acid sequence encoding Mammastatin DNA (SEQ ID NO:1) and its deduced amino acid sequence (SEQ ID NO:2) are shown in Table 1. The sequence was identified by cloning and sequencing of Mammastatin cDNA from a normal human mammary cell cDNA library, as described more fully in the Examples below. Chromatographically purified inhibitor had not previously been sufficiently isolated to permit its amino acid analysis, and early attempts to sequence the protein inhibitor by standard techniques failed. Attempts to screen a cDNA library using antibodies raised against chromatographically purified inhibitor protein failed to generate an active clone. To overcome these problems, the gene encoding Mammastatin was identified by peptide sequencing and degenerate oligonucleotide screening of a normal human mammary cell cDNA library.

[0041] Concentrated protein produced by normal human mammary cells was affinity purified using an anti-Mammastatin antibody raised against chromatographically purified inhibitor. Purified protein fractions were supplemented with a small amount (10^5 cpm) of 32 P labeled as tracers. The labeled tracer protein was purified from conditioned media of cells grown in the presence of 35 P, as described more fully in the examples below. The protein was cleaved with cyanogen bromide, and cleaved fragments were identified as Mammastatin by autoradiographic analysis of 32 P-labeled protein. The most abundant labeled peptides generated by the cleavage were sequenced.

[0042] Two peptides, selected as having unique amino acid sequences (seq. ID Nos. 3 and 4), were used to produce degenerate oligonucleotides. The degenerate oligonucleotides were then used to screen a normal human mammary cell cDNA library.

[0043] One clone, labeled pMammA, hybridized to oligonucleotides from both selected peptides. This clone was further characterized, and was shown to express protein recognized by anti-Mammastatin antibodies. The clone has been verified as encoding Mammastatin by Northern blot analysis, in vitro transcription and translation assays, and growth inhibition assays. A pcDNA3 clone containing the Mammastatin cDNA insert (pMammB) was deposited with the American Type Culture Collection and given Accession Number 97451. The recombinant protein expressed from pMammB (Sequence ID No: 2) has been detected by immunoblot of transfected mammalian cell lines and has been demonstrated to possess growth inhibitory activities against mammary cancer cells. The cDNA clone has been completely sequenced (see Example 3) and found to be unique to the BLAST DNA database.

[0044] The nucleic acid sequence of the invention (Sequence ID No: 1) encodes human Mammastatin, which functions to inhibit the growth of human mammary cells, normal and cancerous. The term "human" is not intended to limit the source of the protein nor to limit its inhibitory effects only to human cells and tissues. It is understood that the nucleic acid sequence and amino acid sequence of Mammastatin in individuals may vary somewhat, without altering the structure or function of the protein. Further, one skilled in biochemistry will appreciate that modifications of the nucleic acid or amino acid sequence may be made without altering the structure and/or function of the molecule. For example, the nucleic acid

sequence may be modified to permit optimal expression of the desired amino acid sequence using known optimal codons for a particular cellular host.

[0045] The nucleic acid sequence of the invention is useful in producing large quantities of highly purified Mammastatin protein for use in therapeutic and diagnostic methods in the treatment of breast cancer.

Anti-Mammastatin Antibodies:

[0046] Several anti-Mammastatin antibodies have been produced and characterized. See, for example, PCT application WO 89/11491 published 30 Nov. 1989. These antibodies were raised against chromatographically purified inhibitor protein, and have been demonstrated to block the inhibitory effect of Mammastatin protein on mammary cell growth.

[0047] Available anti-Mammastatin antibodies include 7G6 and 3C6, commercially available from Neomarkers (Freemont, Calif.) and 6B8. Hybridoma cells producing 6B8 antibody are available from the American Type Culture Collection (ATCC No. HB 10152). Each of these antibodies binds to all three molecular weight forms of Mammastatin and are useful in immunological assays, including dot blots and Western blots. The 7G6 antibody is preferred for Western blot analysis or for ELISA analysis of denatured protein samples. The antibodies 3G6 and 6B8 may be used in ELISA assays, e.g., under conditions specified in the examples.

[0048] Additional antibodies can be produced using standard methods known for producing monoclonal or polyclonal antibodies. The antigen used to produce antibodies may be derived from culture of NHMC or from recombinantly expressed Mammastatin.

Diagnostic Method

[0049] The invention further provides an in vitro assay for detecting active, inhibitory Mammastatin in patient samples, including tissues, cells, and fluids. Breast cancer disease and advancing metastatic disease is diagnosed by correlating the presence and type of Mammastatin protein in a patient's sample with that of normal or cancerous human mammary cells. A patient's blood or tissue sample is analyzed for Mammastatin protein, e.g., for the abundance of Mammastatin protein and/or for the molecular weight forms of Mammastatin. As discussed below, the absence or loss of Mammastatin, particularly of the higher molecular weight, phosphorylated forms of Mammastatin, is correlated with breast cancer and indicative of advancing metastatic disease.

[0050] Analysis of Mammastatin is preferably by immunoassay, including ELISA or Western Blot analysis of a patient's blood samples, using anti-Mammastatin antibodies. Preferably, recombinant Mammastatin standards are used to provide a standard curve for reliable quantitation of inhibitor levels. Such immunoassays are exemplified by the dot-blot assays and Western blot assays shown in the examples below. In an alternative preferred embodiment of the invention, tissue samples, such as tumor biopsies, are analyzed by immunohistochemistry, or by culturing a patient's tumor cells and examining the cultures for expression of Mammastatin.

[0051] In a particularly preferred embodiment, an assay for the diagnosis of breast cancer includes at least two specific antibodies: an antibody to identify the sampled breast tissue as epithelial tissue, such as an anti-cytokeratin antibody, and an anti-Mammastatin antibody. For example, using an immunoblot format, tissue suspected of containing breast cancer

cells is homogenized, separated on an SDS/PAGE gel, transferred to membrane, and probed with both anti-keratin and anti-Mammastatin antibodies. Isotype specific second antibodies that are conjugated to a suitable marker system such as peroxidase or alkaline phosphatase are used to detect bound antibodies. Membranes containing bound first and second antibodies are then developed using known colorimetric or fluorometric techniques and quantitated by known methods.

[0052] In the most preferred embodiment, the sample is analyzed for the phosphorylated forms of Mammastatin, such as by Western Blot, using anti-Mammastatin antibodies. A decline or absence of the high molecular weight (53/49 kD) Mammastatin correlates with advancing breast cancer.

Recombinant Expression Vectors and Transformed Cells

[0053] Recombinant expression vectors of the invention are useful for production and amplification of purified Mammastatin protein and portions thereof, and for easy isolation of Mammastatin protein and portions thereof to be used in diagnostic and therapeutic methods.

[0054] A target sequence, such as all or a portion of the 2A34 kb Mammastatin cDNA (SEQ ID NO: 1), is cloned into a suitable nucleic acid sequence expression vector such as pUC18, pKC30, pBR322, pKK177-3, pET-3, pcDNA3 (In Vitrogen) for COS and CHO cells, and pAcG3X baculovirus expression vector (PharMingen, San Diego, Calif.) for expression in insect cells, and like, known expression systems by standard methods. Commercially available expression vectors provide for cloning of a target sequence into a site of the vector such that the target sequence is operably linked to transcriptional and translational control regions.

[0055] The expression vector is then introduced into suitable host cells using known methods such as calcium phosphate precipitation, liposome mediated transformation, protoplast transformation, electroporation, and the like. Suitable host cells include COS and CHO cells, High 5 and SF9 insect cells, baculovirus, and yeast cells. Other host cells include *E. coli* strains such as *E. coli* DH5a, and avirulent isogenic *Salmonella* spp. such as *S. typhimurium* deletion mutants lacking adenylate cyclase and cAMP receptor protein, *Salmonella* mutants in *aro* genes, and other *Salmonella* vaccine strains as described in *Bio/Tech*, 6:693 (1988).

[0056] Preferably, the cellular host is a Eukaryotic cell, capable of expressing the protein with proper folding and kinase activity to produce a phosphorylated, active inhibitor. Host cells may be screened by transfection with cDNA encoding Mammastatin. Analysis of the protein produced by the transformed cells, e.g. by immunoblot, and the ability of the protein to inhibit mammary cell growth, for example MCF7 cell growth, as described in the examples recited below, can be used to screen potential host cell systems.

[0057] Host cells transformed with the target nucleic acid sequence are screened by a variety of methods including colony hybridization or reactivity with antibodies specific for Mammastatin protein. A transformed cell is a suitable host cell carrying a pcDNA3 or other plasmid or vector containing a nucleic acid sequence encoding Mammastatin. One such plasmid is the pcDNA plasmid (pMammB) carrying the 2.4 kb BamHI-XhoI insert from pMammA, deposited with the American Type Culture Collection in Rockville, Md. on Feb. 22, 1996, and was given Accession No. 97451. (See Example 5.)

[0058] An expression vector containing the specific target DNA sequence is used to generate all or a portion of Mam-

mastatin protein, by in vitro transcription and translation by insertion into cellular hosts for protein production. Proteins produced from the expression vector system inhibit the growth of mammary cells, normal and cancerous. (See Example 7.) Eucaryotic cells, e.g., Cos7 host cells, transfected with the vector express and secrete Mammastatin into the conditioned medium. Conditioned medium inhibited the growth of normal and cancerous mammary cells. (See Example 8.)

Amino Acid Sequence

[0059] The Mammastatin protein (Sequence ID No: 2) is a polypeptide of about 2400 amino acid residues having the sequence deduced from the nucleic acid sequence (SEQ. ID NO: 1) and shown in Table 1. Protein expressed from the cloned Mammastatin nucleic acid sequence (Seq. ID No. 1) inhibits growth of breast cancer cells (MCF-7).

[0060] Recombinant Mammastatin protein can be efficiently produced in purified form and in large quantities. Purified recombinant Mammastatin is useful as a reliable standard for diagnostic assays of the inhibitor in patient samples. Recombinant Mammastatin protein is also useful as a purified therapeutic agent to inhibit or prevent the growth of breast cancer cells.

Therapeutic Use

[0061] Mammastatin protein for therapeutic use is produced from NHMC cultures under serum free conditions or by recombinant means. Mammastatin phosphoprotein is used therapeutically to inhibit mammary cell growth, e.g., in the treatment of breast cancer. Preferably, Mammastatin is produced in higher eucaryotic cells to achieve phosphorylation of the protein. Recombinant Mammastatin protein is produced in host cells or by synthetic means.

[0062] Functional Mammastatin is administered to patients by known methods, for the administration of phosphoprotein, preferably by injection, to increase inhibitor levels in the bloodstream and increase the inhibitor's interactions with mammary cells.

[0063] The protein may be delivered to the patient by methods known in the field for delivery of phosphorylated protein therapeutic agents. In general, the inhibitor is mixed with a delivery vehicle and administered by injection.

[0064] The dosage of inhibitor to be administered may be determined by one skilled in the art, and will vary with the type of treatment modality and extent of disease. Since Mammastatin inhibits approximately 50% of mammary cancer cell growth at a concentration of 10 ng/ml and stops growth at about 20-25 ng/ml in vitro, a useful therapeutic dosage range is about 2.5 ug to about 250 ug administered daily dose. Preferred is approximately 125 ug daily administered dose. The aim of the administration is to result in a final body dose that is in the physiological or slightly higher range (50-75 ng/ml). Higher doses of inhibitor (>50 ng/ml) appear to induce apoptosis, as seen in histology of treated cells. For clinical use, the preferred dosage range is about 500 ng/ml for initial treatment of metastatic disease, followed by a maintenance dosage of about 50 ng/ml. Initial clinical studies, reported in the examples below, indicate an administered daily dose of about 50 ng/ml to about 750 ng/ml is sufficient to induce remission in Stage IV breast cancer patients.

[0065] Since active Mammastatin is a phosphorylated protein, it is anticipated that multiple doses of the inhibitor will

be required to maintain growth inhibiting levels of Mammastatin in the patient's blood. Also, since Mammastatin generally acts as a cytostatic agent rather than a cytotoxic agent, it is expected that a maximum effect of the inhibitor will require regular maintenance of inhibitor levels in breast cancer patients.

[0066] In its preferred use, Mammastatin is administered in high dosages (>50 ng/ml, preferably about 50-500 ng/ml) to induce tumor regression. Lower, maintenance doses (<50 ng/ml, preferably 20-50 ng/ml) are used to prevent cancer cell growth.

[0067] Clinical experience with administered Mammastatin in Stage IV breast cancer patients indicates a useful dose is that which maintains physiological levels of Mammastatin in the blood. Administration is preferably daily, but, may be, for example, by continuous infusion, by slow release depot, or by injection once every 2-3 days. Anecdotal evidence suggests continuous administration may induce feedback inhibition, thus, a preferred administration scheme is to administer daily dose of Mammastatin for approximately 25-28 days, followed by 2-5 days without administration.

Diagnostic Use

[0068] Assays of the present invention for detecting the presence of the functional inhibitor in human tissue and serum are useful in screening patients for breast cancer, for screening the population for those at high risk of developing breast cancer, for detecting early onset of breast cancer, and for monitoring patient levels of inhibitor during treatment. For example, analysis of a patient's blood Mammastatin may indicate a reduced amount of high molecular weight, phosphorylated Mammastatin, as compared with a normal control or with the patient's prior Mammastatin profile. Such a change is correlated with increased risk of breast cancer, with early onset of breast cancer, and with advancing metastatic breast cancer. Diagnostic assay for phosphorylated, active, 49/53 kD Mammastatin preferably is by Western blot immunoassay, e.g. ELISA, or using specific anti-Mammastatin antibodies. Screening, for example, in serum, is preferably by immunoassay, e.g., dot blot assay.

[0069] For best results, the patient samples should be assayed within a short time of sampling (within one week), stored at 4° C. (less than one year), or frozen for long term storage. Most preferably, samples are frozen until time of assay.

Assay Kit

[0070] In a specific embodiment of the invention, an assay kit for the detection of Mammastatin in a patient's fluid and/or breast tissue is provided. The preferred screening assay is an immunoassay such as a dot blot assay to detect or quantitate Mammastatin in blood serum. Such a screening kit includes anti-Mammastatin antibodies and optionally a control antibody and/or Mammastatin controls or standards. A second screening assay analyzes Mammastatin in breast tissue. Preferably, the assay kit contains necessary reagents and tools for reacting the tissue with an antibody to specifically determine that the tissue is breast epithelium, e.g., an anti-cytokeratin antibody, and a specific anti-Mammastatin antibody. The commercially available antibody mixture, pan-keratin (Sigma) is a preferred anti-cytokeratin antibody.

[0071] A negative assay for Mammastatin could be caused by either the presence of a breast cancer tumor, or by non-

epithelial breast tissue. Use of the anti-cytokeratin antibody guards against false positive assays. Epithelial cells of the breast that do not stain with the anti-Mammastatin antibody or which only express the 44 kD Mammastatin are transformed cells. Thus, by first identifying the tissue as breast epithelium, e.g., isolated from breast tissue and positive with the anti-cytokeratin antibody, and then identifying a second positive reaction with anti-Mammastatin antibody, false positives are avoided.

[0072] Because about 30% of the breast cancer cells studied to date express non-phosphorylated inactive, 44 kD Mammastatin, the preferred method of analysis is to differentiate between the 53/49 kD and 44 kD forms, e.g. by Western blot analysis.

[0073] The invention is further defined by reference to the following examples:

Example 1

Human Mammary Cell cDNA Library

[0074] A cDNA library was prepared from human mammary cells obtained from reduction mammoplasties (UM Hospital). Total RNA was isolated from the mammary cells by cesium chloride gradient. From the total RNA preparation, mRNA was isolated. The methods used were those described in Garner I., "Isolation of total and poly A+ RNA from animal cells", *Methods Mol. Biol.* (1994) 28:41-7.

[0075] Reverse transcriptase in the presence of the isolated mRNA produced cDNA that was then ligated to EcoRI linkers. The cDNA was inserted into EcoRI cut T4 DNA ligase-treated Lambda Zap, and amplified in XL1-blue *E. coli*, following the method described in Short J M., et al. (1988) *Nucleic Acids Research* 16: 7583.

Example 2

Preparation of Mammastatin Oligonucleotides

[0076] The normal human mammary cell cDNA library prepared in Example 1 was screened for the presence of nucleic acids encoding Mammastatin using degenerate oligonucleotides. The degenerate oligonucleotides were derived as follows:

[0077] Normal human mammary cells were obtained from the Plastic Surgery Department of the University of Michigan Hospital or from the Cooperative Human Tissue Network. The tissue was reduced by collagenase treatment generally following the procedure described in Soule, et al., *In Vitro* 22:6 (1986).

[0078] Mammary cells were grown to confluence in 175 cm² flasks in DMEM/F12 low calcium media formulated with 40 μ M CaCl₂ and supplemented with 5% CHELEX treated equine serum (Sigma), 0.1 μ g/ml cholera toxin (Sigma), 0.5 μ g/ml hydrocortisone (Sigma), 10 ng/ml epidermal growth factor (EGF, Collaborative Research, Bedford Mass.), 10 pg/ml insulin, and 1 μ g/ml penicillin/streptomycin following the method described in Soule, et al., *In vitro* 22:6 (1986). Equine serum was treated with CHELEX resin for three hours at room temperature to remove serum calcium.

[0079] Cell lysates were prepared by rinsing cells with TBS and scraping from the flask with a Teflon scraper. Cells were collected by centrifugation and lysed with 8M Urea, 50 mM TRIS pH 7.5, 0.5% Beta-mercaptoethanol, 0.5% TRITON X-100 (lysis buffer) and three minutes of sonication on ice.

[0080] The cell lysates were fractionated on DEAE-Sephacel anion exchange resin (Sigma) equilibrated with lysis buffer. Lysates were loaded onto the resin filled columns (50 ml disposable, Bio Rad) and washed with ten column volumes of the lysis buffer. Material flowed through the columns with only gravity feed. Fractions were eluted with a salt gradient produced by continuous gravity feed of elution buffer containing 5M NaCl into a closed mixing chamber initially containing elution buffer (250 ml of 8M urea and 50 mM TRIS pH 7.5) in the absence of salt.

[0081] Elution fractions (2 ml) were collected with a Gibson fraction collector, and were analyzed for the presence of mammary cell growth inhibitor by dot blot with the anti-Mammastatin antibody, 7G6, described above.

[0082] Positive fractions were pooled and dialyzed into lysis buffer with 50 mM NaCl, and were again separated on an identical ion exchange column and eluted with a continuous decreasing pH gradient (pH 8 to pH3) in elution buffer with 50 mM NaCl. (To produce the pH gradient, pH3 buffered urea was continuously mixed with the initial pH8 buffer.) Fractions (2 ml) were collected and analyzed with the 7G6 antibody as described above.

[0083] Positive fractions were again pooled and concentrated to 1/10 the original volume by filtered centrifugation (Amicon Centriprep, 10 kD cutoff). The concentrated pool was size fractionated by preparative SDS polyacrylamide gel electrophoresis (PAGE) along with prestained molecular weight standards (Sigma).

[0084] Protein contained in the molecular weight range between 40 and 60 kD was excised from the gel in 0.5 cm strips or fractions. Electroelution of the protein from each gel strip was carried out by placing the gel strip in 1 ml of running buffer (192 mM glycine, 25 mM TRIS pH 8.3, 0.1% SDS) in dialysis tubing. The tubing was placed in a submarine electrophoresis apparatus and electroeluted overnight at 25 volts. Current was reversed for 2 minutes and running buffer, now containing the electroeluted protein, was removed. Purity of the eluted protein was checked by analytic SDS PAGE with silver-staining, and also by immunoblot with the 7G6 antibody, following the procedure described in Towbin et al., *J. Clin. Chem. Clin. Biochem.* 27:495-501 (1989). Fractions that were at least 70% pure as determined by silver-stained PAGE were pooled, concentrated, and lyophilized to powder form.

[0085] The pooled protein was cleaved with cyanogen bromide by resuspending lyophilized powder in 500 μ l of 70% formic acid and incubating overnight at room temperature (about 20 hours) with 20 mg/ml of cyanogen bromide (Sigma). The methods used are described in Freemont, et al., *Arch. Biochem. Biophys.* 228:342-352 (1986). Cyanogen bromide-cleaved protein samples were dialyzed into double distilled, deionized water and again concentrated and lyophilized to powder.

[0086] Cyanogen bromide cleavage generated multiple peptides from the original protein sample, which were separated by preparative 15% SDS PAGE and transferred onto PVDF membrane by electroelution.

[0087] In addition to the protein obtained from mammary cell lysates, protein was also isolated from normal human mammary cell conditioned medium. Normal cells were incubated with 8 ml DMEM lacking phosphates and supplemented with 200 μ Ci/ml ³²P-ortho-phosphate and 1% dia-

lyzed fetal bovine sera. Cells were allowed to grow for 24 hours in the presence of the ^{32}P before conditioned media was collected.

[0088] The collected conditioned media was concentrated 5× by Amicon filtration with 10 kD exclusion limit. Concentrated media was rinsed once with PBS on filtration membranes to remove excess unincorporated phosphate and was further fractionated by S-200 SEPHACRYL (Pharmacia, Upsala, Sweden) molecular sieve chromatography (100 cm×0.75 cm column) eluted with PBS. Both the filter and the column permit removal of unincorporated ^{32}P from the sample. One ml fractions were collected from the column, and labeled fractions identified by scintillation counting. Radioactive fractions were pooled and analyzed by SDS PAGE with silver staining and autoradiography. The pooled protein was concentrated, lyophilized to powder, and combined with the larger mass of unlabeled protein purified as described above, before cyanogen bromide cleavage. The addition of labeled protein provided a convenient means of tracing cyanogen bromide cleavage fragments containing phosphorylated Mammastatin peptides. Cleaved peptides were separated on preparative PAGE as described above.

[0089] After radioactive proteins were cyanogen bromide cleaved, separated, transferred to PVDF membrane, and exposed to Xray film, two labeled bands of approximately 20 and 22 kD were seen. These two peptides were excised from membranes and sequenced by Edman degradation methods at the University of Michigan Biomedical Research Core Facility using methods described in Ullah Alt et. al., *Biochem. Biophys. Res. Comm.* 203:182-189 (1994). The amino acid sequences of each of the two peptides was compared with known database sequences using the NIH "BLAST" server. The two peptides appeared to be unique.

[0090] A particularly unique portion of each sequence was used to produce degenerate oligonucleotides, using the standard third position degeneracy according to the method described in Jerala, *Biotechniques* 13:564-567 (1992). From the 20 kD peptide, the sequence "gly-gln-leu-glu-tyr-gln-asp-leu-arg" (Seq ID No. 3) was used; from the 22 kD peptide, the sequence "tyr-glu-arg-asp-leu-lys-gly-arg-asp-pro-val-ala-ala" (Seq ID No. 4) was used to generate multiple species of oligonucleotides. The degenerate oligonucleotides were purified by high pressure liquid chromatography.

SEQ. ID NO.	Peptide
3	gly gln leu glu tyr gln asp leu arg
4	tyr glu arg asp leu lys gly arg asp pro val ala ala

[0091] The degenerate oligonucleotides were end-labeled with ^{32}P -gamma ATP and T4 DNA polynucleotide kinase (BRL, Bethesda, Md.) and resuspended in T4 DNA kinase buffer (60 mM TRIS pH 7.8, 10 mM MgCl_2 , 15 mM beta-mercaptoethanol) at 1.5 mg/ml. Oligonucleotides (250 μM) were then incubated with 0.33 μM ATP, 5 units kinase in 25 μl kinase buffer, for two hours at 37° C. Incorporation of ^{32}P -phosphate was determined by TCA precipitation (15% TCA, 4° C., 15 minutes). Typical incorporation was 10⁹ cpm/ μg DNA.

Example 3

Screening Mammary Cell cDNA Library with Degenerate Oligonucleotides

[0092] Bacteria infected with phage prepared for Example 1, containing a normal mammary cell cDNA insert, were plated on 15 cm NZCYM (10 g, NZ amine (Bohringer Mannheim), 5 g NaCl, 5 g yeast extract, 2 g MgSO_4 , 1 g casamino acids) plates in top agar (1/10 dilution of infected bacterial cultures to 6 ml of 7% NZYM top agar) and allowed to incubate eight hours at 37° C. Plates containing plaques were overlaid with nitrocellulose for 15 minutes before denaturation of phage. Phage was denatured by blotting filters (DNA side up) on Whatman paper saturated with 0.5 M NaOH, 1.5 M NaCl for 5 minutes. Filters were rinsed with H_2O before incubating for 5 minutes in 1 M TRIS pH 7.0, 1.5 M NaCl followed by 20×SSC and 2×SSC, each for minutes. Filters were dried and baked for 1 hour at 80° C. or placed under ultraviolet light to immobilize DNA. Baked filters were washed for 30 minutes in 2×SSC with 1% SDS and then prehybridized with 50% deionized formamide, 5×Denhart's solution, 1% SDS, 5×SSC and 100 $\mu\text{g}/\text{ml}$ sheared salmon sperm DNA overnight at 37° C.

[0093] Filters were hybridized with the labeled degenerate oligonucleotide prepared as described for Example 2 in pre-hybridization buffer to which 10⁷ cpm/ml of heat-denatured (95° C., 5 minutes) labeled degenerate oligonucleotide had been added. Hybridizations were performed at 37° C. for 24 hours. Filters were washed with 2×SSC for thirty minutes at 37° C. followed by 3 washes in 2×SSC plus 1% SDS at 50° C. for thirty minutes. Filters were rinsed with 2×SSC briefly, dried and exposed to Kodak AR-5 film for 24-48 hours to identify positive plaques.

[0094] Positive plaques were isolated from agar plugs excised using a reversed 200 μl sterile pipette tip, and resuspended in SM buffer overnight at 4° C. Secondary and tertiary plates (10 cm) were made using XL1-B infected with 1/10,000 dilution of phage containing SM buffer, to bacteria, in NZCYM (with 1 mM MgSO_4). Plaques were produced by incubating infected bacteria for 8 hours as described above, and were then transferred to nitrocellulose before screening with labeled degenerate oligonucleotides. Screening was performed essentially as described in Krocze R A., *J Chromatogr* 618:133-45 (1993), using 10⁷ cpm/ml of labeled DNA for hybridizations and a final wash stringency of 2×SSC at 50° C. for thirty minutes.

[0095] The clone selected for further analysis was one recognized by both of the degenerate oligonucleotides. This clone was given the name "pMamma".

Example 4

Sequencing of Mammastatin cDNA

[0096] The positive clone obtained in Example 3, pMamma, was sequenced by an automated sequencer at the Biomedical Research Core Facility at the University of Michigan and also by dideoxy DNA sequencing using 15% DNA sequencing gels and radiolabeling the DNA fragments with ^{35}S nucleotides. The methods used are described in Lasken R S., et al. *Proc Natl Acad Sci USA* 82:1301-5 (1985).

[0097] The recognized error rate of automatic sequences is about 5%. Therefore, the clone deposited is resequenced for confirmation of the nucleotide sequence, particularly mindful of areas suspected of potential errors, as noted.

[0098] The sequence information originally obtained included numerous stop codons in all reading frames and apparent errors, particularly in the GC rich areas. The original clone (pMammA) was carefully resequenced. The nucleic acid sequence obtained was free of internal stop codons and is shown in Table 1 (Seq ID No: 1).

Example 5

Subcloning the Mammastatin cDNA into an Expression Vector

[0099] The Mammastatin cDNA insert, pMammA, was subcloned into the expression vector, pcDNA 3 (Invitrogen). The Mammastatin cDNA was isolated by digesting the pMammA plasmid obtained as described for Example 4 with BamHI and XhoI restriction endonucleases. The restriction enzymes cut the plasmid at the ends of the Mammastatin clone insert, creating a linear plasmid fragment and a linear insert fragment. The digested sample was placed in the wells of a 1.2% agarose gel submerged in an electrophoresis apparatus, a 50V current was applied for two hours. Electrophoresis separates DNA fragments on the basis of size with the larger plasmid DNA fragment having the slower migration rate on the gel. The portion of the agarose gel containing the 2.4 kb was visualized by ethidium bromide staining and observing the gel over an ultra-violet light box. The 2.4 kb Mammastatin fragment was cut from the gel and placed into dialysis tubing and the DNA was electroeluted into tris-borate buffer, TBE: (0.089M Tris-borate, 0.089M boric acid, 0.002M EDTA) that was collected and precipitated with ethanol.

[0100] The pcDNA3 plasmid DNA was modified to accept the Mammastatin cDNA fragment during ligation. pcDNA3 plasmid was digested with BamHI and XhoI restriction endonucleases and after digestion was complete, the DNA was incubated for one hour in the presence of calf intestinal phosphatase to remove 5' phosphates. The pcDNA3 sample was then phenol extracted and ethanol precipitated.

[0101] The pcDNA3 and the Mammastatin 2.4 kb cDNA fragment were ligated together. The 2.4 kb Mammastatin fragment and the linear pcDNA3 plasmid were mixed in a 3:1 ratio in the presence of T4 DNA ligase. The ligation reaction was allowed to incubate for one hour and then stored at 4° C. overnight. After the ligation reaction was completed the DNA was used to transform *E. coli* competent cells. Subcloning was verified by purifying plasmid DNA from ampicillin selected colonies. The plasmids were digested with the restriction endonucleases BamHI and XhoI. The digested DNA samples were placed in an agarose gel and separated by electrophoresis. A plasmid containing the correct size Mammastatin DNA fragment was designated pMammB, and was deposited with the American Type Culture Collection (ATCC) on Feb. 22, 1996, and given accession number: ATCC 97451.

Example 6

Transfection and Protein Expression from the Mammastatin cDNA Sequence

[0102] Cos-7 cells do not express immunoreactive proteins that co-migrate with the Mammastatin proteins. pMammB and PCDNA3 were used to transfect Cos-7 monkey fibroblast cells using LIPOFECTIN® (BRL, Life Technologies, Bethesda, Md.) using the manufacturers suggested protocol.

The transfected cells were grown for two days prior to harvest. Transfected cells were removed from plates by trypsinization of cells using standard protocols. (2.5 ml's of Trypsin (0.25% SIGMA) was incubated in flasks of cells at 37° C. for 5 minutes. A 7.5 ml aliquot of RPMI media with 10% FBS (fetal bovine serum) was added and cells were collected by centrifugation.) Cells were counted by hemocytometer and lysed in SDS PAGE sample loading buffer at 10⁷ cells/ml. Cell lysates were separated on 8-15% SDS-PAGE gradient gels (Biorad) and transferred to a nylon membrane using methods described in Towbin H., et al., *J. Clin Chem Clin Biochem* (1989 August) 27(8):495-501. The membrane was probed with anti-Mammastatin monoclonal antibody 7G6. Bound antibody was detected with peroxidase conjugated GAM-IgM and developed by ECL (Amersham).

[0103] As shown in FIG. 1, Cos-7 cells transfected with pMammB (lanes C,D) expressed immunoreactive proteins that co-migrated with Mammastatin protein (lane A). Cos-7 cells transfected with the empty vector PCDNA3 alone did not express immunoreactive proteins when immunoblot experiments were performed (lane B).

DNA/AA Sequence

[0104]

Lane A	NHMC (25 µg) - control
Lane B	Cos pcDNA3 cell lysate (25 µg) - control
Lane C	Cos-pMammB cell lysate (10 µg)
Lane D	Cos-pMammB cell lysate (20 µg)

[0105] The immunoblot experiments illustrate the pMammB clone contains a cDNA insert capable of synthesizing a protein with the size and immunologic characteristics of Mammastatin. In addition, immunoreactive proteins of 44, 49 and 53 kD were expressed in Cos-7 cells transfected with pMammB. These proteins migrated at the same molecular weight as the Mammastatin proteins previously identified in normal human mammary cells. This group of immunoreactive proteins was not identified in Cos-7 cells transfected with the empty vector, pcDNA3.

[0106] In the particular assay shown in FIG. 1, the NHMC control shows an unusually high amount of 44 kD Mammastatin. This is an artifact produced by long term (>1 yr) storage of the NHMC standard at 4° C., causing degradation of the higher molecular weight forms, over time. When fresher NHMC samples (<1 yr old) or frozen samples are used, the 44 kD protein is always less abundant than the higher molecular weight forms.

Example 7

GST Fusion

[0107] The Mammastatin clone can be similarly subcloned into a baculovirus expression system. The pMammA insert has been subcloned into a pAcG3X vector obtained commercially from Pharmgen (San Diego, Calif.). This vector allows production of Mammastatin as a fusion protein with glutathione S-transferase (GST), having a portion of the GST gene upstream of the coding site.

[0108] The pMammA insert was subcloned by preparing sets of PCR primers that contained Bam HI (5') and Sma I (3') restriction enzyme recognition sites, a small, non-specific

region, and a portion of the Mammastatin sequence. Three sets of primers, each shifted in reading frame, were prepared. The primers hybridized to the pMammA clones and in a typical PCR reaction with pMammA template DNA, amplified a pMammA PCR product capable of insertion into the reading frame of the GST gene in pAcG3X. The vector was then used to transfect High 5 (InVitrogen) host insect cells, and express a GST-Mammastatin fusion protein that was easily purified from host insect cells using glutathione resin (glutathione agarose, Qiagen, Chatsworth, Calif.).

[0109] To prepare DNA for insertion into the BamH1, Sma 1 restriction site of pAcG3X (PharMingen, San Diego, Calif.), primer sets were prepared in three reading frames to include, for the 5' primer, the BamH1 recognition site (GGATCC), a portion of the pMammA sequence, and some 5' sequence from the pBluescript vector. The 3' primers were identical, and included the Sma1 recognition sequence (GGG CCC), a portion of the pMammA sequence, and some pBluescript sequence.

[0110] The primer sets used are shown in the following table:

Seq. ID No:	
5	5' Primers (in three reading frames)* 5'-TGG GAT CCC TTC GCC ACG AGC ACG GTG-3'
6	5'-TGG GAT CCT TCG CCA CGA GCA CGG-3'
7	5'-TGG GAT CCC CTT CGC CAC GAG CAC-3'
8	3' Primer 5'- TTT TTT TTT TTT GGG CCC TTA AGT-3'**

*Bam H1 site underlined **Sma 1 site underlined

[0111] Only one primer set (Seq. ID NOS. 6 and 8) produced clones capable of coding for active inhibitory Mammastatin. The active clones, when used to transform High 5 cells, produced Mammastatin that was immunologically reactive in the transformed cells (see FIG. 2).

[0112] Other known eukaryotic expression systems may similarly be used to produce Mammastatin protein.

Example 8

Inhibition Assay with Proteins Produced by In Vitro Transcription and Translation

[0113] In vitro transcription of pMammB, Mammastatin cDNA was performed using a Stratagene, Express RNA transcription kit to produce Mammastatin RNA. The RNA produced was translated into protein using the Stratagene In Vitro Express translation kit. Mammastatin protein produced from translation of the Mammastatin RNA was shown to inhibit mammary cell growth in culture.

[0114] Cultures of MCF-7 cells were treated with protein products produced in the translation assays described above. Protein products (5% by volume, culture medium) were added to cells in 12-well plates containing 1 ml medium per well. Parallel cultures were treated with both the translation product and the anti-Mammastatin antibody 3C6, at 30 µg/ml final concentration.

[0115] As a negative control, cultures were treated with protein products translated with the Stragene In vitro Express Translation kit incubated in the absence of Mammastatin

cDNA (i.e., employ vector). These lysates do not have the proper machinery to produce the Mammastatin protein.

[0116] All cultures were allowed to grow for six days after being treated with the protein products and the cell number of each sample was calculated using a Coulter counter. There were triplicate samples of each culture condition so that the cell number of each sample was averaged and percent inhibition was determined by comparison to the reticulocyte lysate treated control cells.

[0117] As shown in FIG. 3, the protein translation product of pMammB inhibited MCF-7 cell growth. This inhibition was greatly reduced or blocked in the presence of anti-Mammastatin antibody, 3C6.

Example 9

Inhibition of Mammary Cells with Proteins Present Within Conditioned Media Obtained From Growing Cos-7 Cells Transfected with pMammB

[0118] Mammary cell growth inhibition experiments were performed using conditioned media obtained from Cos-7 cells transfected with pMammB as described for Example 6. Mammastatin is a secreted protein and is found in conditioned media of cells expressing the protein. The growth inhibition caused by conditioned media was blocked by the addition of anti-Mammastatin antibody.

[0119] MCF-7 cells were plated at 10⁴ cells/ml in MEM supplemented with 10% non-essential amino acids and FBS (SIGMA). Cells were allowed to attach overnight and were then supplemented with 10% by volume of conditioned media (3 day culture) from either: (1) Cos-7 cells transfected with the empty vector pcDNA (Negative control), (2) Cos-7 cells transfected with pMammB (pMammB-Cos), (3) NHMC-conditioned media, or (4) non-conditioned media. Parallel MCF-7 cultures were supplemented with 30 µg/ml of 3C6 blocking antibody. Treated MCF-7 cells were allowed to grow for six days and were then counted by hemocytometer.

[0120] Inhibition of cell growth was determined by comparing the growth of MCF-7 cells incubated in conditioned media with the growth of MCF-7 cells incubated in control, non-conditioned media. Data are shown in FIG. 4, and demonstrate that conditioned media from pMammB-transformed cells inhibited mammary cancer cell growth as efficiently as did normal human mammary cell conditioned media. This inhibition was blocked in the presence of anti-Mammastatin antibody.

Example 10

Three Immunologically Reactive Anti-Mammastatin Proteins

[0121] Whole normal human mammary cells (NHMC) and mammary carcinoma cells in tissue culture cells were lysed, and cell lysate proteins were separated by SDS/PAGE as described above and in Ervin, Paul, 1995, Doctoral dissertation, University of Michigan, Chapter 2. Lysed cell samples were separated on 10% SDS-PAGE in a Mini-Protein II apparatus (25 µg/sample). Proteins were transferred to nitrocellulose and probed with the anti-Mammastatin monoclonal antibody 7G6 or IgM control antibody, alkaline phosphatase conjugated second antibody, goat anti-mouse IgM was utilized with an NBT/BCIP substrate system to detect positive antibody reactions colorimetrically. The data are shown in FIG. 5.

CARCINOMA CELLS	
LANE 1	ZR-75-1
LANE 2	MDA MB 435
LANE 3	4MCF-7
LANE 4	T47D
LANE 5	NHMC-14 positive control
LANE 6	NHMC-14 positive control with the 38C13 antibody
NORMAL CELLS	
LANE 7	NHMC-17
LANE 8	NHMC-16
LANE 9	NHMC-15
LANE 10	NHMC-14
LANE 11	NHMC-6
LANE 12	NHMC-14 positive control

[0122] As shown in FIG. 5, normal human mammary cells expressed a doublet of proteins migrating at 49 and 53 kD that were strongly recognized by the anti-Mammastatin monoclonal antibody and a third weakly immuno-reactive 44 kD protein. The four tumor cell lines tested expressed either a 4410 immuno-reactive protein alone (lanes 1,4) or no immuno-reactive protein at all (lanes 2, 3).

[0123] The above data is representative of experiments performed on normal cells from 42 different reduction mammo-plasty patients over a period of several years. Expression of the 44 kD protein in normal cells and cancer cell lines varied in intensity with each preparation.

Example 11

Mammastatin is a Phosphoprotein

[0124] Cellular phosphorylated proteins of mammary cells were labeled with ^{32}P by supplementing normal mammary cell cultures with ^{32}P -orthophosphate (200 $\mu\text{Ci}/\text{ml}$) for 24 hours. Conditioned media was concentrated 5 $\times$ by Amicom Centrifugation with a 30 kD molecular weight restriction. Concentrated media was rinsed once with PBS on filtration membranes to remove excess unincorporated phosphate and fractionated by S-200 SEPHACRYL (Pharmacia, Upsala, Sweden) molecular sieve chromatography (100 $\text{cm} \times 0.75$ cm column) with PBS elution buffer. Immunoblots were prepared as described above and probed with the 7G6 antibody.

[0125] A radiolabeled 53 kD Mammastatin protein was identified in conditioned media by immunoprecipitation. This analysis indicated Mammastatin is a secreted phosphoprotein. Since secreted phosphoproteins are uncommon, Brefeldin A treatment of cells was utilized to determine whether Mammastatin was present in conditioned media due to secretion or to cell breakage or leaking. Brefeldin A is a fungal compound that blocks the secretion of proteins from eukaryotic cells. Brefeldin A inhibits normal endoplasmic reticulum and golgi function and blocks vesicle formation (Ervin, Paul, 1995, Dissertation, Page 25). Since most secreted proteins are liberated from the cell by a process of exocytosis from membrane bound vesicles, blocking vesicle formation blocks secretion of many proteins. When NHMC are grown in the presence of Brefeldin A, phosphorylated Mammastatin is not identified in conditioned media.

[0126] To determine the amino acid residues that are phosphorylated in Mammastatin protein, radiolabeled 53 kD protein was subjected to phospho-amino acid analysis. NHMC

cells were incubated with ^{32}P -orthophosphate for 24 hours. Cell lysates were then immunoprecipitated with the anti-Mammastatin antibody 7G6 and purified as follows. The 53 kD protein was digested with trypsin and hydrolyzed with acid. Two dimensional thin layer chromatography was used to analyze the phosphorylated amino acids of Mammastatin ^{32}P -amino acids were mixed with phospho-ser/thr/tyr controls and loaded at the origin (0) of a 2D TLC plate (20 cm). The samples were separated into two dimensions: 1st dimension—pH 1.9 Buffer (50 ml formic acid, 156 ml glacial acetic acid/2000 ml (1794 H_2O), 20 minutes @ 1.5 K volts; rotate clockwise; 2nd dimension—pH 3.5 Buffer (10 ml pyridine, 100 ml's glacial acetic acid:1890 ml H_2O) for 16 minutes @ 1.3 K volts.

[0127] The TLC plates were stained with ninhydrin and exposed to film. Phospho-amino acid analysis demonstrated the 53 kD Mammastatin protein contained three types of phosphorylated amino acid residues by comparing autoradiographs to ninhydrin stained phospho-amino acid standards.

[0128] Threonine (Th) was the most abundant phosphorylated amino acid followed by serine (S) and Tyrosine (Ty), the least abundant phosphorylated species. However, the relative abundance of phosphoamino acid residues may not be representative of that in the native protein, since acid hydrolysis can free phosphate from phosphotyrosyl residues.

Example 12

One Mammastatin Protein with Varied Phosphorylation

[0129] Cellular phosphorylation of proteins can be modulated by phosphatases and kinases. Mammastatin is differentially phosphorylated in normal and tumor cell lysates due to differential activities of Mammastatin phosphatases. The effect of phosphatase on Mammastatin in NHMC lysates was examined.

[0130] NHMC were grown to confluence in low calcium media and collected by scraping into TBS. Cells were washed with TBS and resuspended at 2 mg/ml in acetate buffer pH 6.6 with 0.5% Triton X-100. 5 $\mu\text{g}/\text{ml}$ of either *Yersinia* phosphatase (YOP) (Stuckey, et al., *Nature* 370:571-5 (1994)) or *Yersinia* phosphatase mutant (MYOP) containing an active site mutation was used to digest cell lysates for six hours at 37° C. (YOP and MYOP were gifts from Dr. S. Jack Dixon, University of Michigan, Biochemistry Department). As shown in FIG. 6, digestion of normal human mammary cell lysates with *Yersinia* phosphatase (YOP) resulted in a reduced amount of 53 kD Mammastatin protein identified by anti-Mammastatin immunoblot (lane A). In contrast, digestion with the *Yersinia* phosphatase mutant (MYOP, lane B), did not alter identification of the 53 kD Mammastatin protein. These results indicate identification of the 53 kD Mammastatin protein by immunoblot is a convenient measure of the state of phosphorylation of the Mammastatin protein.

[0131] Conditioned medium incubated in the presence of *Yersinia* phosphatase (YOP), as described above, was used to treat MCF-7 cells. As previously observed, NHMC conditioned medium inhibits the growth of MCF-7 cells, and this inhibition is blocked by anti-Mammastatin antibodies. As shown in FIG. 7, treatment of NHMC conditioned medium with YOP abrogates this inhibitory activity. As a control, treatment of NHMC conditioned media with a YOP mutant lacking phosphatase activity (MYOP) was tested. This mutant had no effect on the inhibitory activity of NHMC

conditioned media. Immunoprecipitation of the conditioned media with the anti-Mammastatin antibody 7G6 removed the inhibitory activity.

[0132] TCA precipitation indicated that incubation of conditioned media with YOP removed about 50% of incorporated phosphate. As shown above, YOP also removed the 53 kD species from NHMC lysates (FIG. 6).

Example 13

Phosphorylated Mammastatin Produced by Normal But Not Cancerous Mammary Cells

[0133] Normal and transformed mammary cells were labeled with ^{32}P orthophosphate. Carcinoma cell lines were grown in the media as suggested by the ATCC, with the exception of MCF-7 cells which were grown in MEM (Celox) supplemented with 10% FBS, non-essential amino acids, and insulin (10 mg/l). ^{32}P -orthophosphate labeling of cellular proteins was performed in phosphate-free DMEM (ICN) containing 2% dialysed FBS. Cells were incubated 24 hours at 37° C. with 200 $\mu\text{Ci}/\text{ml}$ of ^{32}P -phosphate. After 48 hours, conditioned media was collected from cell cultures and concentrated 5 $\times$. Conditioned media was washed with TBS and concentrated on Amicon filters with a 10 kD mw cut-off. The cell layer was scraped (using a Teflon cell scraper) into lysis buffer, 1.5 ml/flask (0.5% TritonX-100, 2.01% SDS at deoxycholate) from cell lysates and conditioned media.

[0134] Mammastatin proteins were immunoprecipitated by adding 5 μg 7G6 anti-Mammastatin antibody per 500 μl of 5 $\times$ concentrated media or cell lysate and incubating at room temperature for 1.5 hours. Goat anti-mouse IgM second antibody (5 $\mu\text{g}/0.5\text{ ml}$) was added and the mixture incubated an additional hour. Protein G PLUS/A Agarose® slurry (Oncogene Science) was added and the mixture incubated 1.5 hours at room temperature to immobilize antibody complexes.

[0135] The complexes were washed 6 $\times$ with lysis buffer, each wash followed by centrifugation at 3000 $\times$ g. SDS-PAGE loading buffer (50 μl) was added before the sample was heated to 100° C. for 3 minutes. Supernatants were resolved by SDS-PAGE, transferred to nitrocellulose, and exposed to Kodak X-AR film.

[0136] Phosphate labeling of NHMC proteins and subsequent immunoprecipitation identified 49 and 53 kD phosphoproteins in NHMC. The 49 and 53 kD phosphoproteins were not recognized in carcinoma cell lines. Carcinoma cell lines MCF-7, T47D, ZR-75-1 and MDA-MB-435 expressed a 44 kD immunoreactive protein, but this protein did not label with ^{32}P -orthophosphate.

[0137] This study indicates more incorporated phosphate with increasing molecular weight of Mammastatin. Lack of phosphorylation of Mammastatin in transformed cell lines correlates with lack of higher molecular weight forms of the protein and lack of Mammastatin inhibitory activity.

Example 14

Mammastatin Kinase & Phosphatase

[0138] Flasks of normal or carcinoma cells were grown to 75% confluence. Cell cultures were washed three times with TBS and then scraped into TBS with a Teflon scraper. Cell suspensions were pelleted at 1000 g by centrifugation and then resuspended in a small volume of TBS. An aliquot of each type of cell was removed for protein quantitation. Protein concentrations were then equalized at 2 mg/ml in lysis

buffer (FBS with 0.5% Triton X-100 and 5 $\mu\text{g}/\text{ml}$ each of aprotinin, leupeptin, and PMSF). Equal masses of normal and tumor cell proteins were mixed and incubated at 37° C. for three hours. Parallel mixtures of normal and carcinoma cell lysates were performed in the presence of 10 nM Orthovanadate (NaVO_4), a phosphatase inhibitor. The mixture was then separated by SDS/PAGE and analyzed by Western Blot using the 7G6 antibody. The data are shown in FIG. 8.

LANE A	ZR-75-1 Lysate (30 μg)
LANE B	NHMC Lysate (30 μg)
LANE C	NHMC (30 μg) + ZR 75 (30 μg) + 10 nM NaVO_4
LANE D	NHMC (30 μg) + ZR 75 (30 μg)

[0139] As shown in FIG. 8, cancer cells (ZR-75-1) (lane A) did not produce 53/49 kD Mammastatin, as compared with NHMC (lane B). Mixing of normal and cancer cell proteins, in the presence of proteinases, reduces the amount of active, 5310 inhibitor (lane D). However, in the presence of the tyrosine-phosphatase inhibitor NaVO_4 , the 53 species is retained in the mix (lane C). These results indicate that carcinoma cells express phosphatase activity capable of eliminating phosphorylated forms of Mammastatin.

[0140] Expression of Mammastatin in normal and transformed cell lines can be measured quantitatively by Western blot analysis. Using anti-Mammastatin monoclonal antibodies, it has been demonstrated that there is a consistent difference in expression of this protein between mammary carcinoma cells and cells derived from normal mammary epithelium. Mammastatin was recognized in normal human female mammary tissue as 44, 49, and 53 kD species by Western blot analysis with anti-Mammastatin monoclonal antibody 7G6. In mammary carcinoma cells, there was inconsistent recognition of a 44 kD species, but never 49 or 53 kD immunoreactive forms. When the 49 and 53 kD forms are identified in normal cells they are phosphorylated. The 44 kD species is not phosphorylated. It is therefore possible to use immunoblot analysis to determine if Mammastatin is phosphorylated by observing the expression of the 44 and 49, and 53 kD species of Mammastatin.

Example 15

Identification of Mammastatin in Human Sera

[0141] An enzyme-linked immunosorbent assay (ELISA) was established to detect Mammastatin, using the purified anti-Mammastatin monoclonal antibodies 6B8 and 3C6.

[0142] The antibody 6B8 was used to coat Immulon 1 96-well microtiter plates (Immulon Corp.) at a concentration of 10 $\mu\text{g}/\text{ml}$ or 100 $\mu\text{l}/\text{well}$, for three (3) hours at room temperature, or overnight at 4° C. Plates were blocked with 2% BSA (Sigma) in TBS (150 mM NaCl, 100 mM Tris pH 7.4) for 30 minutes and were then incubated with either purified Mammastatin or sample sera diluted 50% in 2% BSA solution for 1.5 hours at 37° C. Microtiter plates were washed for 5 minutes, three times with 300 $\mu\text{l}/\text{well}$ of TBS plus 0.1% Triton X-100 before addition of second antibody.

[0143] Second antibody was biotinylated 3C6. Antibody was biotinylated by incubation with biotin, N-hydroxy succinimide ester (Sigma) in 0.1 M NaHCO_3 for two hours at room temperature and 16 hours at 4° C. Antibody was dialyzed into 1M NaCl, 50 mM Tris pH 7.4, 0.02% Azide (NaN_3 , Sigma) before use or storage.

[0144] Biotinylated anti-Mammastatin antibody was added at 1 µg/ml and 100 µl/well, in a 2% BSA/TBS solution and incubated for 1.5 hours at 37° C. Microliter plates were washed 5 times for 5 minutes with TBS plus 0.1% Triton X-100 as described above. Second antibody was identified with alkaline phosphatase conjugated streptavidin (Southern Biotechnology) and incubated for one hour at a dilution of 1/1000 in 2% BSA/TBS, 100 µl/well for all samples.

[0145] ELISA assays were developed colorimetrically with PNPP (para-nitrophenyl phosphate Sigma), 1 mg/ml in alkaline phosphatase buffer (10 mM diethanolamine pH 9.5 (Sigma), 0.50 mM MgCl₂ (Sigma)). Microtitre plates were read on an ELISA reader at 405 nm at fifteen minute and thirty minute intervals.

[0146] Using chromatographically purified Mammastatin isolated from cell lysates or conditioned media, a standard curve was established for the ELISA indicating sensitivity of the assay for Mammastatin in the low nanogram range. (See FIG. 9). Quantitation of Mammastatin levels in normal human volunteer sera was performed in serum samples collected at two day intervals for one month, from a volunteer. Mammastatin levels in normal human female sera were detectable by this assay and varied between about 10 and 50 ng/ml (FIG. 10).

[0147] Mammastatin levels were also measured in sera collected from breast cancer patients. Patients diagnosed at the University of Michigan Breast Care Center with node negative breast cancer were tested for Mammastatin expression in sera throughout the course of their treatment. The data are shown in FIG. 11 and summarized below. Serum samples were collected from breast cancer patients during the entire course of their treatment on a hormonal cycling, combined modality protocol with Cytosin, Adriamycin, Methotrexate, and 5Fu. Serum was separated from whole blood, after clotting, by centrifugation and stored at -20° C. until use. ELISA assay using 150 µl serum at 50% in 0.5% NFDm in duplicate were performed, using the 6B8 and 3C6 anti-Mammastatin antibodies in an enzyme linked—"sandwich" assay. The standard curve was generated with chromatographically purified Mammastatin and was comparable to that shown in FIG. 9.

[0148] Expression of Mammastatin varied among patients and fluctuated during the course of their treatment. It was consistently observed that Mammastatin levels became undetectable with progression to metastatic disease.

[0149] Patients diagnosed with breast cancer had low levels of Mammastatin in serum at the time of diagnosis as compared with levels in normal patient serum. Mammastatin levels generally rose on the hormonal cycling, adjuvant chemotherapy protocol. Levels of Mammastatin fluctuated on this protocol. Mammastatin levels were undetectable in patients with advanced disease, before death. The patient data sorted into four groups, as shown in the table below.

[0150] I. Group of patients whose serum Mammastatin levels continued to raise during therapy.

[0151] II. Group of patients whose serum Mammastatin levels increased initially during therapy, but then became undetectable.

[0152] III. Group of patients whose serum Mammastatin levels rose during therapy, but then fluctuated widely.

[0153] IV. Group of patients who had low serum Mammastatin levels which became undetectable with therapy.

Summary of Mammastatin Levels in Patient Sera

[0154]

Group	Number	Days Followed	Outcome
I.	4 p	280 +/- 100	Remission
II.	14 p	500 +/- 220	Deceased
III.	10 p	380 +/- 280	Variable
IV.	5 p	290 +/- 150	Deceased

Example 16

In Vivo Efficacy of Mammastatin

[0155] CD-1 Nu/Nu homozygote, female, six week old mice (Charles River) were supplemented with Estrogen via slow release pellets, 0.72 mg/pellet, 60 day release of 17-beta estradiol (Innovative Research #SE-121). Estrogen supplemented mice were injected with 3×10⁶ MCF-7 cells 100 µl per injection in 60% matrigel. Two injections were administered, one per flank. After seven days of tumor cell growth, Mammastatin was administered. Test mice received 1, 2, or 5 µg of Mammastatin in production media at 2 day intervals for a period of six weeks. Control mice were injected with BSA, or were not injected with tumor, but with the inhibitor alone.

[0156] Tumor size was measured at the point of greatest diameter at weekly intervals and averaged for treatment group. The results are shown in FIGS. 13A-13C, with tumor size plotted as the mean diameter±standard deviation.

[0157] This animal study was repeated using MDA-231 tumor cells. Cells were injected at a concentration of 2×10⁶ cells per injection as described above for MCF-7 cells.

[0158] The results are shown in FIGS. 14A-14C.

[0159] The results shown were not as great as expected. The animals were injected by tail vein, resulting in less than the needed blood dose. Subsequent studies using intraperitoneal injection have resulted in more effective treatment. At doses of 5 ug/mouse and higher, tumor growth is abrogated.

Example 17

Retrovirus Expression of Mammastatin

[0160] The Mammastatin cDNA (2.4 kilobase (kb) insert) was subcloned into a retroviral expression vector. The vector was used to transfect 3T3 fibroblast cells. Transfected cells were harvested, lysed, and the cell lysate analyzed by Western Blot

[0161] As shown in FIG. 12, 3T3 cells transfected with the Mammastatin-carrying retrovirus, expressed phosphorylated Mammastatin.

Example 18

Expression of Mammastatin in Baculovirus and Cos 7 Cells

[0162] Monkey kidney cells, Cos7 cells, were transfected with nucleic acid sequences encoding Mammastatin. Tested groups (n=3) included the following:

GROUP	NUCLEIC ACID SEQUENCE
A	Control, normal human mammary cell lysate 25 µg
B	Control, Cos-pcDNA3 cell lysates - 25 µg
C	Cos-pMammB cell lysate - 10 µg
D	Cos-pMammB cell lysate - 10 µg

[0163] As shown in FIG. 16, induction of recombinant mammastatin expression in Cos-7 cells demonstrated that the mammastatin gene of pMammA and pMammB codes for authentic mammastatin. Furthermore, the observation that Cos-7 cells express the different forms of mammastatin associated with phosphorylation of the protein suggests that mammastatin will be phosphorylated and active when produced in eucaryotic cell lines other than human mammary cells. Stable transfectants have been selected to allow perpetual synthesis of recombinant mammastatin.

Example 19

Production of Mammastatin by Normal Human Breast Epithelial Cells in Culture

[0164] Healthy breast tissue was obtained from reduction mammoplasty, sterile, and direct from the operating room. The tissue was minced under sterile conditions in a laminar flow hood in a solution containing 4 units per gram of type III collagenase (Life Sciences, Bethesda, Md.). The minced tissue was incubated overnight in a shaking water bath at 37° C. to allow collagenase digestion.

[0165] Collagenase digested breast tissue, a viscous fluid containing a variety of cell types and lipid released from adipose cells, was centrifuged to separate lipid, aqueous solution, and other cell types. The collagenase-digested material was spun at 1000 rpm in a table top centrifuge at room temperature for 5 minutes. Adipose cells and free lipid partitioned to the top half of the centrifuge tube, and were withdrawn by aspiration and discarded. The aqueous supernatant positioned above the cell pellet was also withdrawn by aspiration and discarded. The remaining cell pellet was washed with sterile solutions of mammalian growth media, DMEM, pH 7.4. The washing was continued until the supernatant from the washes was no longer turbid (for example about 4 washes). The washed cells were resuspended in growth media and allowed to settle by gravity for 30 minutes at 40° C. Because red blood cells are enucleated and are less dense than nucleated epithelial cells, this procedure resulted in removal of the red blood cells from the sedimented epithelial cells, by withdrawing the supernatant containing the red blood cells. This sedimentation procedure was repeated until no red color remained in the cell pellet, e.g., about 2 times. The remaining cell pellet was resuspended in a nutrient rich DMEM/F12 growth media containing 5% equine serum, 10 µg/ml epidermal growth factor, 100 ng/ml of cholera toxin, 500 ng/ml hydrocortisone, 10 µg/ml insulin, 100 units/ml penicillin and streptomycin, and 1 mM concentration of calcium chloride. Physiological concentrations of calcium helped to promote cell attachment and outgrowth in cell culture. The cell suspensions were incubated in steril tissue culture flasks at 37° C. with a 5% CO₂ concentration.

[0166] Initial cultures of normal breast tissue contain a mixed cell population. The adipocytes, neurons, and vascular

tissue are significantly reduced by the differential centrifugation process described above. Connective tissue cells are present in significant amounts. To remove non-epithelial cells, a differential attachment method was used. Fibroblasts, neurons, and other cell types in breast tissue all attach to tissue culture plastic more rapidly than epithelial cells. In addition, all of these cell types are removed from tissue culture plastic by trypsin more rapidly than epithelial cells. To enrich the cultures for breast epithelial cells, cultures beginning to form a monolayer (5-7 days after initial plating) are treated with a trypsin:EDTA solution (250:1) molar ratio. The majority of cells were removed within 5 minutes of incubation at 37° C. The remaining attached cells were more than 90% epithelial breast cells. These cells were saved and returned to the growth medium described above with 40 µM calcium chloride. The fibroblast cells were removed from the trypsinized culture flasks, collected by centrifugation, resuspended in growth medium, and plated onto tissue culture plastic for 30 minutes at 37° C. The attached cells were predominantly fibroblasts. The cells that did not attach were significantly enriched for fibroblast cells (50-80%). These suspended cells were removed and allowed to settle in fresh tissue culture flasks. This process was repeated twice to obtain cell populations that were predominantly epithelial. Because cholera toxin promotes epithelial cell growth and inhibits fibroblast growth, and because fibroblasts do not grow well in reduced calcium, the cultures were approximately 100% epithelial within one week in the low calcium medium described above. These cultures of normal human mammary cells (NHMC) produced Mammastatin into the culture medium.

[0167] Nutrient medium used to grow the NHMC contains 5% equine serum, which is not acceptable for human injection. The equine serum proteins must either be purified away from the mammastatin protein, or the cells grown in medium devoid of the serum. Normal cells can only be maintained in the absence of serum for about seven to ten days. In order to produce a significant quantity of serum free mammastatin over a prolonged period of time, the cells were alternately grown in serum-free and serum-containing medium.

[0168] NHMC were grown to complete confluence in growth media as described above. The cells began to bud in solution as they grew, when cells covered the available surface of the flask. Budding cells were collected and transferred to new flasks. Confluent flasks were rinsed three times with sterile saline, with a five minute saline incubation between washes to remove serum protein. Cells were then provided with serum free "production medium" that was essentially the growth medium devoid of serum, cholera toxin, and hydrocortisone. Cells were maintained on the production medium for about 4 days (96 hours), with collection of the medium, and a return of the cells to growth medium for at least four days. The typical batch size for mammastatin produced in this way was 1-2 liters.

[0169] Mamastatin has also been produced in a Bioreactor, the Bioflow 3000, New Brunswick Scientific. In this perfusion reactor, cells were attached on tissue culture treated fibracell disks. The cell-attached disks were maintained in a basket in the reaction vessel and perfused with media. When NHMC were introduced to the reactor, they populate the fibracell disks and were fed by the perfusion of media. Conditioned media was harvested from the reactor and refrigerated.

Example 20

Dot Blot Serum Assay for Mammastatin

[0170] Serum from a 25 year old healthy female was obtained and compared with serum from a breast cancer

patient (Stage IV), an undiagnosed sibling of the patient, and from the patient's mother, whose family has a history of breast cancer. The serum samples were compared in an immunoassay for the presence of Mammastatin. Blood samples from multiple breast cancer patients taken on day of diagnosis were also analyzed in the immunoassay. Normal human mammary cell (NHMC) conditioned media was used as a standard control. Standard NHMC mammastatin contained approximately 50 ng/ml as determined in a dot blot assay with mammastatin protein standard chromatographically purified.

[0171] Individual blood samples were collected into vacutainer tubes, and the serum separated from whole blood. Serum samples (250 or 500 μ l volume) were applied without dilution to nitrocellulose by suction using a 96 well, S&S Dot-Blot manifold. Conditioned medium was prepared as described for Example 19. Samples on the nitrocellulose filters were washed with Triton X-100 in TBS, blocked with non-fat dry milk (5% in TBS) and incubated with 1 μ g/ml of first anti-mammastatin antibody (7G6 mouse IgM) in 5% non-fat dry milk for 1.5 hours at room temperature, followed by incubation with 1 μ g/ml of second antibody (goat anti-mouse IgM conjugated to alkaline phosphatase) in 5% non-fat dry milk for one hour and room temperature. The alkaline phosphatase color reaction was developed using nitroblue-tetrazolium and BCIP.

[0172] As shown in FIG. 13, the amount of Mammastatin in the sample was quantitated against the standard curve obtained from normal breast cell conditioned medium. Serum obtained from healthy females contained readily detectable amounts of Mammastatin, as indicated by darkly colored blots, whereas serum from diagnosed breast cancer patients, and from undiagnosed family members showed little or no Mammastatin.

[0173] Additional samples were obtained from breast cancer patients on day of diagnosis, from healthy members of a breast cancer patient's family, and from healthy females and males. The serum was processed as described above in order to analyze Mammastatin. The dot blots were evaluated as "negative or low" or "positive or high" to indicate the intensity of the developed color reaction. Data are shown in the following table.

Sample	Number	Negative or Low	Positive or High
Breast Cancer patient	89	83 (93%)	6 (7%)
Healthy female	11	2 (18%)	9 (82%)
Healthy member of high risk family	4	4 (100%)	0
Male	3	2	1

Example 21

Treatment of Human Breast Cancer Patients

[0174] Twenty-nine (29) Stage IV breast cancer patients with recurrent breast cancer, who had failed, or were failing on chemotherapeutic regimes were given access to Mammastatin protein. The protein was produced as described above for Example 19, and provided in production medium, with the required dose in a 3 ml injection volume. Patients administered the protein intravenously according to their prescribed regimen. In general, one daily dose was injected. The selected

dose was that which provided physiological amounts of Mammastatin in the patient's bloodstream, e.g., 5-50 ng/ml in healthy women. The dosage and frequency for each patient are indicated in the table below.

Patient Number	Dose (μ g)	Schedule	Result
1.	125	daily	complete remission followed by relapse*
2.	125	daily	complete remission; pain if therapy stopped
3.	75	daily	non-responder*
4.	75	daily	partial remission; possible immune reaction**
5.	75	daily	non-responder*
6.	125	daily	partial remission
7.	125	daily	partial remission*
8.	75	daily	non-responder*
9.	125	every third day	complete remission
10.	125	daily	non-responder
11.	125	daily	partial remission
12.	125	daily	non-responder#
13.	150	daily	non-responder*
14.	75	daily	non-responder#
15.	125	daily	partial remission
16.	125	daily	partial remission
17.	75	daily	non-responder, infection**
18.	125	daily	partial remission
19.	125	daily	partial remission
20.	125	daily	partial remission
21.	125	every other day	non-responder**; alternative therapy
22.	125	every other day	partial remission
23.	125	every other day	non-responder
24.	125	daily	non-responder
25.	125	daily	non-responder
26.	125	daily	partial remission
27.	125	daily	partial remission
28.	125	every other day	partial remission
29.	125	daily	
Total			% Responders % without liver involvement % without lung or liver involvement
29	17	59	81 89

*patient deceased

**Patient withdrawn from therapy

#evidence of improvement in jaundice prior to death

[0175] Of the group of 29 patients, six, having late stage liver disease, did not survive. These six patients displayed clinical evidence of liver failure before receiving Mammastatin, and were not helped by the treatments. One patient showed signs of decreased jaundice before her liver failed, but all six of these patients appeared to die of nitrogen toxicity common to patients with advanced liver cancer.

[0176] Of the remaining patients, two have died of their disease. One appeared to be disease-free after two months of Mammastatin therapy. This patient was removed from therapy, and relapsed within two months. The patient's disease was never brought back under control and she died of liver involvement. The second patient died after 4 months of treatment, having never shown any sign of response to the therapy. There was no sign of toxicity in any of these patients, although the dose of Mammastatin in these latter two patients was increased ten fold.

[0177] Of the 19 patients currently receiving Mammastatin therapy, the majority show signs of positive benefit and no signs of adverse reaction. It is unclear if three of these patients are receiving any benefit from Mammastatin. The other 16 patients show definite clinical signs of benefit including

decreased tumor markers (CA15-3 and CA27-29) to normal levels, decreased size of palpable tumor masses, decreased disease as evidenced on MRI scan, and decreased pain. Several of these patients show improvement to the point of being considered disease free. However, it has consistently been observed that denying these patients protein for periods of three to five days results in resumption of disease activity as evidenced by increased pain, even in patients that show no signs of disease. Resumption of protein treatments decreases or eliminates the symptoms of increased pain within 2-4 hours.

[0178] It has also been observed that Mammastatin levels in the blood decline after long term treatment, suggesting a negative feedback system. This decline in constant blood levels is successfully avoided by providing Mammastatin daily for a period of about 28 days, followed by 2-3 days without protein.

Example 22

Recombinant Mammastatin for Human Therapy

[0179] Recombinant Mammastatin has been produced in Cos-7 monkey kidney cells, chinese hamster ovary (CHO) cells, and Sf9 insect cells by transfecting the cells with a plasmid containing the Mammastatin cDNA sequence. The Mammastatin cDNA has been stably integrated into the genomes of these producing cell lines, and secrete protein immunoreactive with growth inhibitory activity.

[0180] To produce Mammastatin from these cells and isolate the protein for human use, the cell lines are grown in serum free medium for approximately 48 to 72 hours. The media is withdrawn and protein purified from conditioned medium, either by ion exchange chromatography in Tris buffer, pH7.5, using a sodium chloride gradient from about 0.1 M to about 0.5 M, collecting the Mammastatin fraction at about 0.2 M. The protein fraction is then dialyzed against normal saline, diluted if necessary, and filter sterilized.

[0181] In an alternative method, Mammastatin is produced as a fusion protein in Cos7 or Sf9 cells. The fusion protein contains a histidine tag (six histidine residues) and a Factor X proteinase cleavage site. The Mammastatin expressing cells are cultured, preferably in 1% serum-containing media, the conditioned media is collected and passed over a nickel chelating resin. The His-fusion protein adheres to the column, is washed with 50 mM TRIS, pH 7.5, 0.1 M NaCl, and is slowly eluted with TRIS-NaCl containing 10 Unit/ml Factor X proteinase. This liberates Mammastatin from the His fusion. Mammastatin is separated by molecular sieve chromatography, or by ion exchange chromatography as described above.

[0182] The above specification, examples and data provide a complete description of the manufacture and use of the composition of the invention. Since many embodiments of the invention can be made without departing from the spirit and scope of the invention, the invention resides in the claims hereinafter appended.

TABLE 1

MAMMASTATIN DNA SEQUENCE									
CGGCACGAGC	ACGGTGAAGA	GACATGAGAG	GTGTAGAATA	AGTGGGAGGC	CCCCGGCGCC	60			
CCCCCGGTGT	CCCCGCGAGG	GGCCCGCGGG	TCCGCCGGCC	CGCGGGCGCC	GGTGAAATAC	120			
CACTACTCTG	ATCGTTTTTT	CACTGACCCG	GTGAGGCGGG	GGGCGAGCCC	CGAGGGGCTC	180			
TCGCTTCTGG	CGCCAAGCGC	CCGGCCGCGC	GCCGGCCGGG	CGCGACCCGC	TCCGGGGACA	240			
GTGCCAGTGG	GGAGTTTGAC	TGGGGCGGTA	CACCTGTCAA	ACGGTAACGC	AGGTGTCTTA	300			
AGGCGAGCTC	AGGGAGGACA	AAACCTCCCG	TGGAGCAGAA	GGGCAAA	ATG ATC TTG	356			
					Met Ile Leu				
					1				
ATT TTC AGT	ACG AAT ACA	GAC CGT	GAA AGC	GGG GCC	TCA GAT	CTT CTG	404		
Ile Phe Ser	Thr Asn Thr	Asp Arg	Glu Ser	Gly Ala	Ser Asp	Leu Leu			
5		10		15					
ACC TTT TGG	GTT TTA AGC	AGG AGG	TGT CAG	AAA AGT	TAC CAC	AGG GAT	452		
Thr Phe Trp	Val Leu Ser	Arg Arg	Cys Gln	Lys Ser	Tyr His	Arg Asp			
20		25		30		35			
AAC TGG CTT	GTG GCG GCC	AAG CGT	TCA AAG	CGA CGT	CGC TTT	TTG ATC	500		
Asn Trp Leu	Val Ala Ala	Lys Arg	Ser Lys	Arg Arg	Arg Phe	Leu Ile			
	40		45			50			
CTT CGA TGT	CGG CTC TTC	CTA TCA	TTG GGA	AGC AGA	ATT CAC	CAA GCG	548		
Leu Arg Cys	Arg Leu Phe	Leu Ser	Leu Gly	Ser Arg	Ile His	Gln Ala			
	55		60		65				
TTG GAT TGT	TCA CCC ACT	AAT AGG	GAA CGT	GAG CTG	GGT TTA	GAC CGT	596		
Leu Asp Cys	Ser Pro Thr	Asn Arg	Glu Arg	Glu Leu	Gly Leu	Asp Arg			
	70		75		80				
CGT GAG ACA	GGT TTG TTT	ACC CTA	CTG ATG	ATG TGT	TGT TGC	CAT GGT	644		
Arg Glu Thr	Gly Leu Phe	Thr Leu	Leu Met	Met Cys	Cys Cys	Cys His			
	85		90		95				

TABLE 1-continued

MAMMASTATIN DNA SEQUENCE																
AAT Asn 100	CCT Pro	GCT Ala	CAG Gln	TAC Tyr	GAG Glu	AGG Arg	AAC Asn	CGC Arg	AGG Arg	TTC Phe	AGA Arg	CAT His	TTG Leu	GTG Val	TAT Tyr	692
					105					110					115	
GTG Val	CTT Leu	GGC Gly	TGG Trp	GGA Gly	GCC Ala	AAT Asn	GGG Gly	GCG Ala	AAG Lys	CTA Leu	CCA Pro	TCT Ser	GTG Val	GGA Gly	TTA Leu	740
					120					125					130	
TTA Leu	CTG Leu	AAC Asn	GCC Ala	TCT Ser	AAG Lys	TCA Ser	GAA Glu	TCC Ser	CGC Arg	CCA Pro	GGC Gly	GGA Gly	ACG Thr	ATA Ile	CGG Arg	788
					135					140					145	
CAG Gln	CGC Arg	CGC Arg	GGA Gly	GCC Ala	TCG Ser	GTT Val	GGC Gly	CTC Leu	GGA Gly	TGG Trp	CCG Pro	GTC Val	CCC Pro	CGC Arg	CTG Leu	836
					150					155					160	
TCC Ser	CCG Pro	CCG Pro	GCG Ala	GGC Gly	GCC Ala	CCC Pro	CCC Pro	CCT Pro	CCA Pro	CGC Arg	GCC Ala	CCG Pro	CGC Arg	GCG Ala	CGG Arg	884
					165					170					175	
GAG Glu	GGC Gly	GCG Ala	TGC Cys	CCC Pro	GCC Ala	GCG Ala	CGC Arg	CGG Arg	GAC Asp	CGG Arg	GGT Gly	CCG Pro	GTG Val	CGG Arg	AGT Ser	932
					180					185					190	
GCC Ala	CTT Leu	CGT Arg	CCT Pro	GGG Gly	AAA Lys	CGG Arg	GGC Gly	GCG Ala	GCC Ala	GGA Gly	AAG Lys	GCG Ala	GCC Ala	GCC Ala	CCC Pro	980
					200					205					210	
TCG Ser	CCC Pro	GTC Val	ACG Thr	CAC His	CGC Arg	ACG Thr	TTC Phe	GTG Val	CTC Leu	GTG Val	CCG Pro	AAT Asn	TCG Ser	GCA Ala	CGA Arg	1028
					215					220					225	
GTA Val	GCA Ala	CCA Pro	TTC Phe	ACA Thr	ATA Ile	GAC Asp	ATA Ile	CAA Gln	GTG Val	CAT His	GTA Val	TCT Ser	TTA Leu	TTA Leu	TAT Tyr	1076
					230					235					240	
AAT Asn	GAA Glu	TTC Phe	TTT Phe	TCC Ser	TTT Phe	GGG Gly	GAG Glu	ATA Ile	TCC Ser	AGT Ser	AGT Ser	GGG Gly	ATT Ile	GCT Ala	AGA Arg	1124
					245					250					255	
TCA Ser	CCT Pro	GGT Gly	AGT Ser	TCT Ser	ATT Ile	TCT Ser	GGT Gly	TTA Leu	TTG Leu	AGA Arg	AAT Asn	CTT Leu	CAT His	ACT Thr	GAT Asp	1172
					260					265					270	
TTC Phe	CAT His	AGA Arg	GGT Gly	TGT Cys	ACA Thr	AAT Asn	TTA Leu	CAT His	CCC Pro	TAC Tyr	CAA Gln	GTG Val	ATT Ile	TTT Phe	TTA Leu	1220
					280					285					290	
AAT Asn	ATG Met	AAA Lys	GAA Glu	TGG Trp	TCT Ser	GGA Gly	GAA Glu	ATG Met	CCC Pro	CTC Leu	ATT Ile	AGT Ser	ATC Ile	CCC Pro	CTT Leu	1268
					295					300					305	
TTA Leu	CCT Pro	CTC Leu	TAC Tyr	TGC Cys	AGA Arg	ATG Met	ACT Thr	TCA Ser	AGG Arg	GGT Gly	ACA Thr	GGT Gly	ATT Ile	TAC Tyr	AAG Lys	1316
					310					315					320	
TTT Phe	CAT His	TAT Tyr	ACA Thr	GAC Asp	AAA Lys	TTG Leu	AAT Asn	ATT Ile	GAA Glu	ATT Ile	TCT Ser	GCA Ala	TTA Leu	GAG Glu	GCA Ala	1364
					325					330					335	
CAG Gln	ATT Ile	TTA Leu	GGA Gly	TTC Phe	AAA Lys	GTT Val	GTA Val	AGA Arg	ACA Thr	AGG Arg	ACA Thr	AGT Ser	GCT Ala	CTA Leu	GGG Gly	1412
					340					345					350	
ACT Thr	TGC Cys	AAA Lys	GCT Ala	GGA Gly	ATT Ile	GGA Gly	AAT Asn	CTC Leu	AGA Arg	AGA Arg	AAT Asn	ACA Thr	TTT Phe	CTA Leu	GTA Val	1460
					360					365					370	
GTA Val	CCA Pro	CCA Pro	GCA Ala	TAT Tyr	ATT Ile	CTA Leu	CTG Leu	AAT Asn	TGG Trp	CTT Leu	TGT Cys	GAT Asp	CAT His	CAT His	TTA Leu	1508
					375					380					385	

TABLE 1-continued

MAMMASTATIN DNA SEQUENCE																					
TAC	CTA	CTT	ATT	AAA	ACT	AAT	GAA	AAG	GGT	TTA	TAT	CAA	ATA	TAC	TTT	1556					
Tyr	Leu	Leu	Ile	Lys	Thr	Asn	Glu	Lys	Gly	Leu	Tyr	Gln	Ile	Tyr	Phe						
		390					395					400									
AAG	GTA	AAA	AAA	TCA	AAT	TAT	AGG	AAA	AGC	TGT	TTT	CTT	TTG	CAT	TTT	1604					
Lys	Val	Lys	Lys	Ser	Asn	Tyr	Arg	Lys	Ser	Cys	Phe	Leu	Leu	His	Phe						
	405					410					415										
AAT	TTC	AAA	ACA	AAA	AAT	AGC	TAC	CGT	CTA	TTG	GGC	ATT	TAT	ACT	GTA	1652					
Asn	Phe	Lys	Thr	Lys	Asn	Ser	Tyr	Arg	Leu	Leu	Gly	Ile	Tyr	Thr	Val						
	420				425					430				435							
CCA	GAC	ACT	GTG	TTT	GTC	ACA	TTT	CAA	AAA	TGT	TCT	CAT	GGT	AAT	GTT	1700					
Pro	Asp	Thr	Val	Phe	Val	Thr	Phe	Gln	Lys	Cys	Ser	His	Gly	Asn	Val						
			440						445					450							
CAC	AAT	AAT	TCT	GTA	GGG	TGG	AGA	AAT	AGT	CTT	ACC	GTA	GTA	AGA	CTA	1748					
His	Asn	Asn	Ser	Val	Gly	Trp	Arg	Asn	Ser	Leu	Thr	Val	Val	Arg	Leu						
			455					460					465								
TTC	AGA	AAC	GAA	ACC	TCT	GAA	CCT	TGG	AGT	TCA	ACT	TGC	GCA	AAG	TTA	1796					
Phe	Arg	Asn	Glu	Thr	Ser	Glu	Pro	Trp	Ser	Ser	Thr	Cys	Ala	Lys	Leu						
		470					475					480									
GTA	ACA	GGA	CTA	GGA	CTT	GAA	CCT	GAA	CCA	TCA	CAC	TCC	AGA	TCT	CTC	1844					
Val	Thr	Gly	Leu	Gly	Leu	Glu	Pro	Glu	Pro	Ser	His	Ser	Arg	Ser	Leu						
	485					490					495										
CAT	ACC	ACA	CTG	CTA	GCA	CAT	GTG	CCT	GTC	ATC	TTA	TTC	CTG	GCT	CCC	1892					
His	Thr	Thr	Leu	Leu	Ala	His	Val	Pro	Val	Ile	Leu	Phe	Leu	Ala	Pro						
	500				505					510				515							
TKY	TTA	TTT	CCT	TTC	CCT	TCC	TCC	CAC	AAC	CCC	TTT	TTC	CCC	CCA	TTT	1940					
Xaa	Leu	Phe	Pro	Phe	Pro	Ser	Ser	His	Asn	Pro	Phe	Phe	Pro	Pro	Phe						
			520						525				530								
CTT	TTC	TTT	CTT	TTT	ATT	TGT	TAATTACATA				ACTAATACAT				GTTTATCAGA		ACAA	1995			
Leu	Phe	Phe	Leu	Phe	Ile	Cys															
			535																		
TTGATATAGC ACAAAAGGAT ATAAAGTACG GGTGAGTGAT AGCTCATCCC TGTAATCTAG 2055																					
CACTTTGGAA GGCCAAGGCA GGCAGATCAC TTGATCCAGA GTTCGAGACC AGCCTGGGCA 2115																					
ACATGGTGAA ACCCTGTCTC TACAAAAAAA TACAAAAATT TAGCCGGGCG TGCTGGCACA 2175																					
CACCTGTAGT CTCAGCTACT CTGAGGGCTG AGGTGGGAAG ATTGATTGAG CCCAGGAGGT 2235																					
GGAAGCTGCA GCAGTGCCT GAGATTGCGC CATTGCACTC CAGCCTGGGT GAGAGAGAGA 2295																					
GACCCTGTCT TCAAAAAAAA AAAAAAAAAA A																					

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 8

<210> SEQ ID NO 1

<211> LENGTH: 2326

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<220> FEATURE:

<221> NAME/KEY: CDS

<222> LOCATION: (348)..(1961)

<400> SEQUENCE: 1

-continued

cggcacgagc acggtgaaga gacatgagag gtgtagaata agtgggaggc ccccggcgcc	60
cccccggtgt ccccgcgagg ggcgcgagg tccgcccggc cgcgggcgcc ggtgaaatac	120
cactactctg atcgtttttt cactgaccgc gtgaggcggg gggcgagccc cgaggggctc	180
tcgcttctgg cgccaagcgc ccggccgcgc gccggccggg cgcgaccgc tccggggaca	240
gtgccagtgg ggagtttgac tggggcggtg cacctgtcaa acggtaacgc aggtgtccta	300
aggcgagctc agggaggaca aaacctcccg tggagcagaa gggcaaa atg atc ttg	356
Met Ile Leu	
1	
att ttc agt acg aat aca gac cgt gaa agc ggg gcc tca gat ctt ctg	404
Ile Phe Ser Thr Asn Thr Asp Arg Glu Ser Gly Ala Ser Asp Leu Leu	
5 10 15	
acc ttt tgg gtt tta agc agg agg tgt cag aaa agt tac cac agg gat	452
Thr Phe Trp Val Leu Ser Arg Arg Cys Gln Lys Ser Tyr His Arg Asp	
20 25 30 35	
aac tgg ctt gtg gcg gcc aag cgt tca aag cga cgt cgc ttt ttg atc	500
Asn Trp Leu Val Ala Ala Lys Arg Ser Lys Arg Arg Arg Phe Leu Ile	
40 45 50	
ctt cga tgt cgg ctc ttc cta tca ttg gga agc aga att cac caa cgg	548
Leu Arg Cys Arg Leu Phe Leu Ser Leu Gly Ser Arg Ile His Gln Ala	
55 60 65	
ttg gat tgt tca ccc act aat agg gaa cgt gag ctg ggt tta gac cgt	596
Leu Asp Cys Ser Pro Thr Asn Arg Glu Arg Glu Leu Gly Leu Asp Arg	
70 75 80	
cgt gag aca ggt ttg ttt acc cta ctg atg atg tgt tgt tgc cat ggt	644
Arg Glu Thr Gly Leu Phe Thr Leu Leu Met Met Cys Cys Cys His Gly	
85 90 95	
aat cct gct cag tac gag agg aac cgc agg ttc aga cat ttg gtg tat	692
Asn Pro Ala Gln Tyr Glu Arg Asn Arg Arg Phe Arg His Leu Val Tyr	
100 105 110 115	
gtg ctt ggc tgg gga gcc aat ggg gcg aag cta cca tct gtg gga tta	740
Val Leu Gly Trp Gly Ala Asn Gly Ala Lys Leu Pro Ser Val Gly Leu	
120 125 130	
tta ctg aac gcc tct aag tca gaa tcc cgc cca ggc gga acg ata cgg	788
Leu Leu Asn Ala Ser Lys Ser Glu Ser Arg Pro Gly Gly Thr Ile Arg	
135 140 145	
cag cgc cgc gga gcc tcg gtt ggc ctc gga tgg ccg gtc ccc cgc ctg	836
Gln Arg Arg Gly Ala Ser Val Gly Leu Gly Trp Pro Val Pro Arg Leu	
150 155 160	
tcc ccg ccg gcg ggc gcc ccc ccc cct cca cgc gcc ccg cgc gcg ccg	884
Ser Pro Pro Ala Gly Ala Pro Pro Pro Pro Arg Ala Pro Arg Ala Arg	
165 170 175	
gag ggc gcg tgc ccc gcc gcg cgc ccg gac ccg ggt ccg gtg ccg agt	932
Glu Gly Ala Cys Pro Ala Ala Arg Arg Asp Arg Gly Pro Val Arg Ser	
180 185 190 195	
gcc ctt cgt cct ggg aaa ccg ggc gcg gcc gga aag gcg gcc gcc ccc	980
Ala Leu Arg Pro Gly Lys Arg Gly Ala Ala Gly Lys Ala Ala Ala Pro	
200 205 210	
tcg ccc gtc acg cac cgc acg ttc gtg ctc gtg ccg aat tcg gca cga	1028
Ser Pro Val Thr His Arg Thr Phe Val Leu Val Pro Asn Ser Ala Arg	
215 220 225	
gta gca cca ttc aca ata gac ata caa gtg cat gta tct tta tta tat	1076
Val Ala Pro Phe Thr Ile Asp Ile Gln Val His Val Ser Leu Leu Tyr	
230 235 240	
aat gaa ttc ttt tcc ttt ggg gag ata tcc agt agt ggg att gct aga	1124

Asn 245	Glu Phe	Phe Ser	Phe Ser	Ser Ile	Phe 265	Gly Ser	Glu Gly	Ile Leu	Ser Leu	Ser Arg	Ser 270	Gly Asn	Ile Leu	Ala His	Arg Thr
tca Ser 260	cct Pro	ggg Gly	agt Ser	tct Ser	att Ile 265	tct Ser	ggg Gly	tta Leu	ttg Leu	aga Arg	aat Asn 270	ctt Leu	cat His	act Thr	gat Asp 275
1172															
ttc Phe	cat His	aga Arg	ggg Gly	tgt Cys 280	aca Thr	aat Asn	tta Leu	cat His	ccc Pro 285	tac Tyr	caa Gln	gtg Val	att Ile	ttt Phe 290	tta Leu
1220															
aat Asn	atg Met	aaa Lys 295	gaa Glu	tgg Trp	tct Ser	gga Gly	gaa Glu	atg Met	ccc Pro 300	ctc Leu	att Ile	agt Ser	atc Ile	ccc Pro	ctt Leu
1268															
tta Leu	cct Pro	ctc Leu 310	tac Tyr	tgc Cys	aga Arg	atg Met	act Thr	tca Ser	agg Arg	ggg Gly	aca Thr 320	ggg Gly	att Ile	tac Tyr	aag Lys
1316															
ttt Phe	cat His 325	tat Tyr	aca Thr	gac Asp	aaa Lys	ttg Leu 330	aat Asn	att Ile	gaa Glu	att Ile	tct Ser 335	gca Ala	tta Leu	gag Glu	gca Ala
1364															
cag Gln 340	att Ile	tta Leu	gga Gly	ttc Phe	aaa Lys 345	ggt Val	gta Val	aga Arg	aca Thr	agg Arg 350	aca Thr	agt Ser	gct Ala	cta Leu	ggg Gly 355
1412															
act Thr	tgc Cys	aaa Lys	gct Ala	gga Gly 360	att Ile	gga Gly	aat Asn	ctc Leu	aga Arg 365	aga Arg	aat Asn	aca Thr	ttt Phe	cta Leu 370	gta Val
1460															
gta Val	cca Pro	cca Pro	gca Ala 375	tat Tyr	att Ile	cta Leu	ctg Leu	aat Asn	tgg Trp 380	ctt Leu	tgt Cys	gat Asp 385	cat His	cat His	tta Leu
1508															
tac Tyr	cta Leu 390	ctt Leu	att Ile	aaa Lys	act Thr	aat Asn	gaa Glu 395	aag Lys	ggg Gly	tta Leu	tat Tyr	caa Gln 400	ata Ile	tac Tyr	ttt Phe
1556															
aag Lys	gta Val 405	aaa Lys	aaa Lys	tca Ser	aat Asn	tat Tyr 410	agg Arg	aaa Lys	agc Ser	tgt Cys	ttt Phe 415	ctt Leu	ttg Leu	cat His	ttt Phe
1604															
aat Asn 420	ttc Phe	aaa Lys	aca Thr	aaa Lys	aat Asn 425	agc Ser	tac Tyr	cgt Arg	cta Leu	ttg Leu 430	ggc Gly	att Ile	tat Tyr	act Thr	gta Val 435
1652															
cca Pro	gac Asp	act Thr	gtg Val	ttt Phe 440	gtc Val	aca Thr	ttt Phe	caa Gln	aaa Lys 445	tgt Cys	tct Ser	cat His	ggg Gly	aat Asn 450	gtt Val
1700															
cac His	aat Asn	aat Asn	tct Ser	gta Val 455	ggg Gly	tgg Trp	aga Arg	aat Asn 460	agt Ser	ctt Leu	acc Thr	gta Val 465	gta Val	aga Arg	cta Leu
1748															
ttc Phe	aga Arg 470	aac Asn	gaa Glu	acc Thr	tct Ser	gaa Glu	cct Pro	tgg Trp 475	agt Ser	tca Ser	act Thr	tgc Cys 480	gca Ala	aag Lys	tta Leu
1796															
gta Val 485	aca Thr	gga Gly	cta Leu	gga Gly	ctt Leu	gaa Glu 490	cct Pro	gaa Glu	cca Pro	tca Ser	cac His 495	tcc Ser	aga Arg	tct Ser	ctc Leu
1844															
cat His 500	acc Thr	aca Thr	ctg Leu	cta Leu	gca Ala 505	cat His	gtg Val	cct Pro	gtc Val	atc Ile 510	tta Leu	ttc Phe	ctg Leu	gct Ala	ccc Pro 515
1892															
tky Xaa	tta Leu	ttt Phe	cct Pro	ttc Phe 520	cct Pro	tcc Ser	tcc Ser	cac His	aac Asn 525	ccc Pro	ttt Phe	ttc Phe	ccc Pro	cca Pro 530	ttt Phe
1940															
ctt Leu	ttc Phe	ttt Phe	ctt Leu	ttt Phe 535	att Ile	tgt Cys	taattacata actaatacat				gtttatcaga				
1991															
acaattgata taqcacaaaa qqatataaaq tacqqgtqag tqataqctca tcacctqtaat															2051

-continued

```

ctagcacttt ggaaggccaa ggcaggcaga tcacttgatc cagagttcga gaccagcctg 2111
ggcaacatgg tgaaccctg tctctacaaa aaaatacaaa aatttagccg ggcgtgctgg 2171
cacacacctg tagtctcagc tactctgagg gctgaggtgg gaagattgat tgagcccagg 2231
aggtggaagc tgcagcagtg cgctgagatt gcgccattgc actccagcct gggtagagaga 2291
gagagaccct gtcttcaaaa aaaaaaaaaa aaaaa 2326

```

```

<210> SEQ ID NO 2
<211> LENGTH: 538
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (516)..(516)
<223> OTHER INFORMATION: The 'Xaa' at location 516 stands for Cys, or Phe.
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

```

```

<400> SEQUENCE: 2

```

```

Met Ile Leu Ile Phe Ser Thr Asn Thr Asp Arg Glu Ser Gly Ala Ser
1           5           10          15
Asp Leu Leu Thr Phe Trp Val Leu Ser Arg Arg Cys Gln Lys Ser Tyr
20          25          30
His Arg Asp Asn Trp Leu Val Ala Ala Lys Arg Ser Lys Arg Arg Arg
35          40          45
Phe Leu Ile Leu Arg Cys Arg Leu Phe Leu Ser Leu Gly Ser Arg Ile
50          55          60
His Gln Ala Leu Asp Cys Ser Pro Thr Asn Arg Glu Arg Glu Leu Gly
65          70          75          80
Leu Asp Arg Arg Glu Thr Gly Leu Phe Thr Leu Leu Met Met Cys Cys
85          90          95
Cys His Gly Asn Pro Ala Gln Tyr Glu Arg Asn Arg Arg Phe Arg His
100         105         110
Leu Val Tyr Val Leu Gly Trp Gly Ala Asn Gly Ala Lys Leu Pro Ser
115         120         125
Val Gly Leu Leu Leu Asn Ala Ser Lys Ser Glu Ser Arg Pro Gly Gly
130         135         140
Thr Ile Arg Gln Arg Arg Gly Ala Ser Val Gly Leu Gly Trp Pro Val
145         150         155         160
Pro Arg Leu Ser Pro Pro Ala Gly Ala Pro Pro Pro Arg Ala Pro
165         170         175
Arg Ala Arg Glu Gly Ala Cys Pro Ala Ala Arg Arg Asp Arg Gly Pro
180         185         190
Val Arg Ser Ala Leu Arg Pro Gly Lys Arg Gly Ala Ala Gly Lys Ala
195         200         205
Ala Ala Pro Ser Pro Val Thr His Arg Thr Phe Val Leu Val Pro Asn
210         215         220
Ser Ala Arg Val Ala Pro Phe Thr Ile Asp Ile Gln Val His Val Ser
225         230         235         240
Leu Leu Tyr Asn Glu Phe Phe Ser Phe Gly Glu Ile Ser Ser Ser Gly
245         250         255
Ile Ala Arg Ser Pro Gly Ser Ser Ile Ser Gly Leu Leu Arg Asn Leu
260         265         270

```

-continued

His	Thr	Asp	Phe	His	Arg	Gly	Cys	Thr	Asn	Leu	His	Pro	Tyr	Gln	Val
		275					280					285			
Ile	Phe	Leu	Asn	Met	Lys	Glu	Trp	Ser	Gly	Glu	Met	Pro	Leu	Ile	Ser
		290				295					300				
Ile	Pro	Leu	Leu	Pro	Leu	Tyr	Cys	Arg	Met	Thr	Ser	Arg	Gly	Thr	Gly
305					310					315					320
Ile	Tyr	Lys	Phe	His	Tyr	Thr	Asp	Lys	Leu	Asn	Ile	Glu	Ile	Ser	Ala
				325					330					335	
Leu	Glu	Ala	Gln	Ile	Leu	Gly	Phe	Lys	Val	Val	Arg	Thr	Arg	Thr	Ser
			340					345					350		
Ala	Leu	Gly	Thr	Cys	Lys	Ala	Gly	Ile	Gly	Asn	Leu	Arg	Arg	Asn	Thr
		355					360					365			
Phe	Leu	Val	Val	Pro	Pro	Ala	Tyr	Ile	Leu	Leu	Asn	Trp	Leu	Cys	Asp
		370				375					380				
His	His	Leu	Tyr	Leu	Leu	Ile	Lys	Thr	Asn	Glu	Lys	Gly	Leu	Tyr	Gln
385				390					395						400
Ile	Tyr	Phe	Lys	Val	Lys	Lys	Ser	Asn	Tyr	Arg	Lys	Ser	Cys	Phe	Leu
			405					410					415		
Leu	His	Phe	Asn	Phe	Lys	Thr	Lys	Asn	Ser	Tyr	Arg	Leu	Leu	Gly	Ile
		420					425					430			
Tyr	Thr	Val	Pro	Asp	Thr	Val	Phe	Val	Thr	Phe	Gln	Lys	Cys	Ser	His
		435				440					445				
Gly	Asn	Val	His	Asn	Asn	Ser	Val	Gly	Trp	Arg	Asn	Ser	Leu	Thr	Val
		450			455					460					
Val	Arg	Leu	Phe	Arg	Asn	Glu	Thr	Ser	Glu	Pro	Trp	Ser	Ser	Thr	Cys
465					470				475						480
Ala	Lys	Leu	Val	Thr	Gly	Leu	Gly	Leu	Glu	Pro	Glu	Pro	Ser	His	Ser
			485					490						495	
Arg	Ser	Leu	His	Thr	Thr	Leu	Leu	Ala	His	Val	Pro	Val	Ile	Leu	Phe
		500					505					510			
Leu	Ala	Pro	Xaa	Leu	Phe	Pro	Phe	Pro	Ser	Ser	His	Asn	Pro	Phe	Phe
		515				520					525				
Pro	Pro	Phe	Leu	Phe	Phe	Leu	Phe	Ile	Cys						
		530			535										

<210> SEQ ID NO 3
 <211> LENGTH: 9
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 3

Gly	Gln	Leu	Glu	Tyr	Gln	Asp	Leu	Arg
1					5			

<210> SEQ ID NO 4
 <211> LENGTH: 13
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 4

Tyr	Glu	Arg	Asp	Leu	Lys	Gly	Arg	Asp	Pro	Val	Ala	Ala
1				5					10			

-continued

```

<210> SEQ ID NO 5
<211> LENGTH: 27
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

```

```

<400> SEQUENCE: 5

```

```

tgggatacct tcgccacgag cacggtg

```

27

```

<210> SEQ ID NO 6
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

```

```

<400> SEQUENCE: 6

```

```

tgggatacctt cgcacagagc acgg

```

24

```

<210> SEQ ID NO 7
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

```

```

<400> SEQUENCE: 7

```

```

tgggataccc ttgccacga gcac

```

24

```

<210> SEQ ID NO 8
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

```

```

<400> SEQUENCE: 8

```

```

tttttttttt ttgggccctt aagt

```

24

1.-17. (canceled)

18. A method comprising the steps of:

- a) determining an amount of mammastatin in a patient's sample by contacting said mammastatin with an antibody that binds specifically to a mammastatin protein comprising SEQ ID NO: 1;
- b) comparing the amount of determined mammastatin to a control, said control comprising an amount of mammastatin from a normal, non-breast cancer female; and
- c) correlating an amount of mammastatin less than the control as a characteristic of breast cancer.

19. The method of claim 18, wherein said antibody is a monoclonal antibody.

20. The method of claim 18, wherein said antibody is selected from the group consisting of monoclonal antibody 7G6, monoclonal antibody 3C6, and monoclonal antibody 6B8.

21. The method of claim 18, wherein said sample comprises mammary tissue, mammary cells, or fluids from said patient.

22. The method of claim 18, wherein said patient's sample comprises blood or blood serum from said patient.

23. The method of claim 18, where said step of determining an amount of mammastatin comprises use of an immunoassay.

24. The method of claim 23, wherein the immunoassay is selected from the group consisting of a Western blot, a dot blot, and an enzyme-linked immunoabsorbent assay.

25. The method of claim 18, wherein said control comprises an amount of mammastatin of at least 10 ng/ml.

26. The method of claim 18, wherein said control comprises mammary cells, or mammary tissue, or fluid obtained from a normal, non-breast cancer female.

27. A monoclonal antibody specific for a mammastatin protein comprising SEQ ID NO: 1.

28. The monoclonal antibody of claim 27, wherein said mammastatin protein has an approximate molecular weight

selected from the group consisting of 44 kD, 49 kD and 53 kD.

29. The monoclonal antibody of claim **27**, wherein said antibody is selected from the group consisting of monoclonal

antibody 7G6, monoclonal antibody 3C6, and monoclonal antibody 6B8.

* * * * *

专利名称(译)	乳房抑制素的核苷酸和蛋白质序列及使用方法		
公开(公告)号	US20110003318A1	公开(公告)日	2011-01-06
申请号	US12/846386	申请日	2010-07-29
[标]申请(专利权)人(译)	密歇根大学		
申请(专利权)人(译)	密歇根大学董事会		
当前申请(专利权)人(译)	密歇根大学董事会		
[标]发明人	ERVIN JR PAUL R		
发明人	ERVIN, JR., PAUL R.		
IPC分类号	G01N33/53 C07K16/18 A61K38/00 A61K48/00 A61P35/00 C07K14/47 C12N15/09 C12N15/12 C12Q1/68 G01N33/574 G01N33/68		
CPC分类号	A61K38/00 A61K48/00 C07K14/4703 C07K16/18 G01N33/57488 C07K2319/00 C12Q1/6886 G01N33/57415 C07K2317/74 A61P35/00		
优先权	60/027315 1996-10-03 US		
外部链接	Espacenet USPTO		

摘要(译)

编码乳房抑制素(一种特定的乳腺细胞生长抑制剂)的核酸序列。乳房抑制素由单个核酸序列编码,并且其无活性的非磷酸化形式具有约44kDa的分子量。正常乳腺细胞表达具有约53kDa和49kDa的近似分子量的功能性磷酸化形式。转移性乳腺细胞或者根本不表达乳房抑制素,或者不表达53kDa或49kDa的磷酸化形式。通过施用磷酸化的乳腺抑素抑制乳腺癌细胞的生长。

Expression of Recombinant Mammatatin in Cos-7 Cells

