



US 20110123567A1

(19) **United States**

(12) **Patent Application Publication**
Bird et al.

(10) **Pub. No.: US 2011/0123567 A1**
(43) **Pub. Date: May 26, 2011**

(54) **RECOMBINANT RIFT VALLEY FEVER (RVF)
VIRUSES AND METHODS OF USE**

Related U.S. Application Data

(75) Inventors: **Brian H. Bird**, Woodland, CA
(US); **Cesar G. Albarino**, Atlanta,
GA (US); **Stuart T. Nichol**, Atlanta,
GA (US); **Thomas G. Ksiazek**,
Lilburn, GA (US)

(60) Provisional application No. 61/016,065, filed on Dec.
21, 2007, provisional application No. 61/042,987,
filed on Apr. 7, 2008.

Publication Classification

(73) Assignees: **THE GOVERNMENT OF THE
UNITED STATES OF AMERICA
AS REPRESENTED BY THE
SECRETARY OF THE
DEPARTMENT OF HEALTH
AND HUMAN SERVICES,
CENTERS FOR DISEASE
CONTROL AND PREVENTION,
ATLANTA, GA (US)**

(51) **Int. Cl.**
A61K 39/12 (2006.01)
C12N 7/01 (2006.01)
C12N 15/63 (2006.01)
C12N 5/00 (2006.01)
G01N 33/566 (2006.01)
G01N 33/53 (2006.01)

(52) **U.S. Cl.** **424/205.1**; 435/235.1; 435/320.1;
435/325; 435/7.92; 436/501

(57) **ABSTRACT**

Described herein are recombinant RVF viruses comprising deletions in one or more viral virulence genes, such as NSs and NSm. The recombinant RVF viruses, generated using a plasmid-based reverse genetics system, can be used as vaccines to prevent infection of RVF virus in livestock and humans. As described herein, the recombinant RVF viruses grow to high titers, provide protective immunity following a single injection and allow for the differentiation between vaccinated animals and animals infected with wild-type RVF virus.

(21) Appl. No.: **12/809,561**


(22) PCT Filed: **Dec. 16, 2008**

(86) PCT No.: **PCT/US2008/087023**


§ 371 (c)(1),
(2), (4) Date: **Jun. 18, 2010**

FIG. 1

A. Recombinant RVF viruses



B. Live infected cells



C. Fixed and gamma-irradiated cells




FIG. 2




FIG. 4

Pilot Study: Day 21 post-immunization

Inoculum	# animals	Vacc. Dose (PFU)	RVF IgG SUN ₁₀₀ mean ± (SEM)	PRNT ₅₀ mean ± (SEM)	RVF viremia (PFU/mL eq.)
					post-vaccination (day 1-4)
rRVF-ΔNSs:GFP	9	1x10 ⁷	2.14 ± 0.12*§	1:1480 ± (631)	0.0
rRVF-ΔNSs:GFP-ΔNSm	9	1x10 ⁷	1.24 ± 0.06*§	1:280 ± (120)	0.0
Neg. Control	3	0	-0.08 ± 0.06	1:10 ± (0.0)	0.0

Challenge Study: Day 26 post-immunization

Inoculum	# animals	Vacc. Dose (PFU)	RVF IgG SUN ₁₀₀ mean ± (SEM)	PRNT ₅₀ mean ± (SEM)	RVF viremia (PFU/mL eq.)		# Surviving Challenge
					post-vacc. (d1-7)	Peak post-challenge (day)	
rRVF-ΔNSs:GFP	10	1x10 ⁷	4.10 ± (0.12) *	1:640 (0.0)	0.0	10 animals = 0.0*	10/10*
rRVF-ΔNSs:GFP	10	1x10 ⁸	4.79 ± (0.11) *	1:7040 (3200)	0.0	9 = 0.0* 1 = 1.5x10 ¹ (d4)*	10/10*
rRVF-ΔNSs:GFP-ΔNSm	10	1x10 ⁷	3.94 ± (0.12) *	1:1120 (733)	0.0	9 = 0.0* 1 = 1.1x10 ² (d3)*	10/10*
rRVF-ΔNSs:GFP-ΔNSm	10	1x10 ⁸	4.54 ± (0.13) *	1:640 (0.0)	0.0	9 = 0.0* 1 = 7.0x10 ¹ (d3)*	10/10*
Neg. Control	5	0	-0.02 ± (0.09)	1:5 (2.9)	0.0	3 = 3.4x10 ¹ (d3) 2 = 1.6x10 ¹ (d4)	2/5

FIG. 5




FIG. 6




FIG. 7




FIG. 8





FIG. 9



RECOMBINANT RIFT VALLEY FEVER (RVF) VIRUSES AND METHODS OF USE

CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] This application claims the benefit of U.S. Provisional Application No. 61/016,065, filed Dec. 21, 2007, and U.S. Provisional Application No. 61/042,987, filed Apr. 7, 2008, which are herein incorporated by reference in their entirety.

FIELD

[0002] This disclosure concerns recombinant RVF viruses comprising deletions in viral virulence genes. These recombinant viruses can be used as vaccines to prevent RVF virus infection. This disclosure further relates to a reverse genetics system used to generate recombinant RVF viruses.

BACKGROUND

[0003] Rift Valley fever (RVF) virus (family Bunyaviridae, genus *Phlebovirus*) is a mosquito-borne pathogen of both livestock and humans found throughout Africa and more recently the Arabian peninsula. Historically, RVF virus has been the cause of either low-level endemic activity or large explosive epizootics/epidemics of severe disease throughout its range (Findlay et al., *Lancet ii*:1350-1351, 1931; Jouan et al., *Res. Virol.* 140:175-186, 1989; Ringot et al., *Emerg. Infect. Dis.* 10:945-947, 2004; Woods et al., *Emerg. Infect. Dis.* 8:138-144, 2002). RVF outbreaks are characterized by economically disastrous "abortion storms" with newborn animal mortality approaching 100% among livestock, especially sheep and cattle (Coetzer et al., *J. Vet. Res.* 49:11-17, 1982; Easterday et al., *Am. J. Vet. Res.* 23:470-479, 1962; Rippy et al., *Vet. Pathol.* 29:495-502, 1992).

[0004] Human infections typically occur either from an infected mosquito bite, percutaneous/aerosol exposure during the slaughter of infected animals, or via contact with aborted fetal materials. Human RVF disease is primarily a self-limiting febrile illness that in a small percentage (about 1-2%) of cases can progress to more serious and potentially lethal complications including hepatitis, delayed onset encephalitis, retinitis, blindness, or a hemorrhagic syndrome with a hospitalized case fatality of 10-20% (Madani et al., *Clin. Infect. Dis.* 37:1084-1092, 2003; McIntosh et al., *S. Afr. Med. J.* 58:83-806, 1980). Excessively heavy rainfall in semi-arid regions often precedes large periodic outbreaks of RVF virus activity, allowing for the abundant emergence of transovarially infected *Aedes* spp. mosquitoes and subsequent initiation of an outbreak by transmission of virus to livestock and humans via infected mosquito feeding (Linthicum et al., *Science* 285:397-400, 1999; Swanepoel et al., *Contributions to Epidemiology and Biostatistics* 3:83-91, 1981). The association with abnormally heavy rains provides some ability to predict periods and regions of high disease risk, which in turn provides a potential window of opportunity for targeted vaccination programs if a safe, inexpensive and highly efficacious single dose vaccine were available.

[0005] The ability of RVF virus to cross international and geographic boundaries and strain veterinary and public health infrastructures is well documented. In 1977, RVF virus was reported for the first time north of the Sahara desert where an extremely large outbreak affecting more than 200,000 people occurred along the Nile River basin in Egypt (Meegan et al.,

Contributions to Epidemiology and Biostatistics 3:100-113, 1981). Approximately ten years later in 1987, a large outbreak occurred in western Africa along the border of Mauritania and Senegal affecting an estimated 89,000 individuals (Jouan et al., *Res. Virol.* 140:175-186, 1989). Later, the virus was isolated for the first time outside of Africa (across the Red sea) in Saudi Arabia and Yemen and was found to be the cause of a large epizootic/epidemic in 2000 with an estimated 2000 human cases and 245 deaths (Anonymous, *Morb. Mortal. Wkly. Rep.* 49:982-5, 2000; Centers for Disease Control and Prevention, *Morb. Mortal. Wkly. Rep.* 49:1065-1066, 2000; Shoemaker et al., *Emerg. Infect. Dis.* 8:1415-1420, 2002).

[0006] Most recently, in late 2006 to early 2007, following heavy rainfall in eastern Africa, RVF virus emerged as the cause of a widespread outbreak that eventually resulted in 1062 reported human cases and 315 reported deaths. Associated with the outbreak were substantial economic losses among livestock in southern Somalia, Kenya, and northern Tanzania (Anonymous, *Morb. Mortal. Wkly. Rep.* 56:73-76, 2007). The ability of RVF virus to cause explosive outbreaks in previously unaffected regions accompanied by high morbidity and mortality during RVF epizootics/epidemics highlights the importance of developing high throughput screening tools for potential antiviral therapeutic agents and the development of safe and efficacious vaccines for this significant veterinary and public health threat.

SUMMARY

[0007] Disclosed are RVF viruses that are highly attenuated, immunogenic and contain precise molecular markers allowing for the differentiation of naturally infected and vaccinated animals (DIVA). Provided herein are recombinant RVF viruses, wherein the genome of the recombinant RVF viruses comprise (i) a full-length L segment; (ii) a full-length M segment or an M segment comprising a complete deletion of the NSm open reading frame (ORF); and (iii) an S segment comprising a complete deletion of the NSs ORF. In one embodiment, the NSs ORF of the recombinant RVF virus is replaced by the ORF of a reporter gene.

[0008] Also provided are immunogenic compositions comprising one or more of the recombinant RVF viruses described herein and a pharmaceutically acceptable carrier. Further provided is a method of immunizing a subject against RVF virus infection, comprising administering to the subject an immunogenic composition disclosed herein. The immunogenic compositions can be used for vaccination of livestock or humans.

[0009] Further provided is a collection of plasmids comprising (i) a plasmid encoding a full-length anti-genomic copy of the L segment of RVF virus; (ii) a plasmid encoding a full-length anti-genomic copy of the M segment of RVF virus, or an anti-genomic copy of the M segment of RVF virus comprising a complete deletion of the NSm ORF; and (iii) a plasmid encoding an anti-genomic copy of the S segment of RVF virus, wherein the S segment comprises a complete deletion of the NSs ORF. In some embodiments, the plasmids further comprise a T7 promoter and a hepatitis delta virus ribozyme. In one example, the NSs ORF of the S segment plasmid is replaced by the ORF of a reporter gene, such as a green fluorescent protein. Also provided are isolated host cells comprising a collection of plasmids provided herein.

[0010] Further provided is a method of preparing a recombinant RVF virus for immunization of a subject, comprising (i) transfecting cultured cells with the collection of plasmids

described herein; (ii) incubating the cells for 1 to 5 days; and (iii) collecting recombinant RVF virus from the cell supernatant.

[0011] Also provided are recombinant RVF viruses, wherein the genome of the recombinant RVF viruses comprise a full-length L segment, a full-length M segment and a full-length S segment, wherein the S segment further encodes the ORF of a reporter gene.

[0012] A reverse genetics system for producing recombinant RVF virus is also provided. The reverse genetics system consists of three plasmids, a plasmid that encodes an antigenomic copy of an S segment, a plasmid that encodes an M segment and a plasmid that encodes an L segment of RVF virus, wherein each plasmid comprises a T7 promoter and a hepatitis delta virus ribozyme. Further provided are recombinant RVF viruses produced using the reverse genetics system described herein.

[0013] The foregoing and other features and advantages will become more apparent from the following detailed description of several embodiments, which proceeds with reference to the accompanying figures.

BRIEF DESCRIPTION OF THE FIGURES

[0014] FIG. 1A is a schematic depiction of the rRVF cDNA plasmid constructions used to generate the rRVF virus stocks. FIG. 1B is a series of images showing direct live cell LTV imaging of Vero E6 cells 24 hours after infection with each recombinant rRVF virus (expressing eGFP). FIG. 1C is a series of images showing Vero E6 cells 24 hours after infection with rRVF-NSs-(Ala)₃GFP or rRVF-NSs-(Ala)₁₀GFP viruses at an MOI of 1. Cells were fixed and stained with monoclonal antibodies specific for eGFP, RVF NSs and counterstained with DAPI to confirm intranuclear co-localization of eGFP and NSs.

[0015] FIG. 2 is a graph showing the results of anti-RVF virus total IgG adjusted SUM_{OD} ELISA testing of all vaccinated (40) and sham-inoculated (5) control animals at day 26 post-immunization. A positive/negative cut-off value was established as the mean+3 standard deviations of the sham-inoculated SUM_{OD} values (open circles-dashed line). §=Significant differences between vaccinated and control groups (p-value <0.05).

[0016] FIGS. 3A-3D are a series of images showing representative results of indirect fluorescent testing of serum collected from (A) WF rats surviving challenge with RVF virus; (B) vaccinated WF rats (day 26 post-vaccination); (C) negative control sham inoculated rats (day 26 post-vaccination); and (D) naturally infected convalescent livestock (goat) sera obtained during the RVF virus outbreak in Saudi Arabia in 2000. The presence of anti-NP antibodies (left panels) and anti-NSs antibodies (right panels) is detected using Vero E6 cells expressing either NP or NSs, respectively. To confirm intranuclear accumulation of anti-NSs antibody, cells were counterstained with DAPI.

[0017] FIGS. 4A and 4B are tables showing data obtained from a vaccination pilot study (A) and challenge study (B).

[0018] FIG. 5 is a plasmid map of pRVS.

[0019] FIG. 6 is a plasmid map of pRVM.

[0020] FIG. 7 is a plasmid map of pRVL.

[0021] FIG. 8 is a plasmid map of pRVS-GFPANSs.

[0022] FIG. 9 is a plasmid map of pRVMANSm.

SEQUENCE LISTING

[0023] The nucleic and amino acid sequences listed in the accompanying sequence listing are shown using standard letter abbreviations for nucleotide bases, and three letter code for amino acids, as defined in 37 C.F.R. 1.822. Only one strand of each nucleic acid sequence is shown, but the complementary strand is understood as included by any reference to the displayed strand. In the accompanying sequence listing:

[0024] SEQ ID NO: 1 is the nucleotide sequence of the S segment of wild-type RVF virus strain ZH501, deposited under Genbank Accession No. DQ380149 on Jan. 31, 2007.

[0025] SEQ ID NO: 2 is the nucleotide sequence of the M segment of wild-type RVF virus strain ZH501, deposited under Genbank Accession No. DQ380200 on Jan. 31, 2007.

[0026] SEQ ID NO: 3 is the nucleotide sequence of the L segment of wild-type RVF virus strain ZH501, deposited under Genbank Accession No. DQ375406 on Jan. 31, 2007.

[0027] SEQ ID NO: 4 is the nucleotide sequence of primer RVS-35/KpnI.

[0028] SEQ ID NO: 5 is the nucleotide sequence of primer RVS+827/BglIII.

[0029] SEQ ID NO: 6 is the nucleotide sequence of primer eGFP+1/KpnI.

[0030] SEQ ID NO: 7 is the nucleotide sequence of primer eGFP-720/BgA

[0031] SEQ ID NO: 8 is the nucleotide sequence of primer RVS-829rev/KpnI.

[0032] SEQ ID NO: 9 is the nucleotide sequence of primer NSsGFP+10Ala/Fwd.

[0033] SEQ ID NO: 10 is the nucleotide sequence of primer NSsGFP+10Ala/Rev.

[0034] SEQ ID NO: 11 is the nucleotide sequence of plasmid pRVS.

[0035] SEQ ID NO: 12 is the nucleotide sequence of plasmid pRVM.

[0036] SEQ ID NO: 13 is the nucleotide sequence of plasmid pRVL.

[0037] SEQ ID NO: 14 is the nucleotide sequence of plasmid pRVS-GFPANSs.

[0038] SEQ ID NO: 15 is the nucleotide sequence of the plasmid pRVMANSm.

DETAILED DESCRIPTION

I. Abbreviations

- [0039]** BSL Bio-safety level
[0040] CPE Cytopathic effect
[0041] DIVA Differentiation of naturally infected and vaccinated animals
[0042] eGFP Enhanced green fluorescent protein
[0043] HRP Horseradish peroxidase
[0044] IFA Immunofluorescence assay
[0045] LD Lethal dose
[0046] MOI Multiplicity of infection
[0047] NP Nucleoprotein
[0048] NS Non-structural
[0049] OIE Office International des Epizooties
[0050] ORF Open reading frame
[0051] PFU Plaque-forming unit
[0052] PRNT Plaque reduction neutralization titers
[0053] RNA Ribonucleic acid
[0054] RT-PCR Reverse transcriptase polymerase chain reaction

- [0055] RVF Rift Valley fever
- [0056] SQ Subcutaneously
- [0057] USDA United States Department of Agriculture
- [0058] WF Wistar-furth
- [0059] WOA World Organization for Animal Health

II. Terms

[0060] Unless otherwise noted, technical terms are used according to conventional usage. Definitions of common terms in molecular biology may be found in Benjamin Lewin, *Genes V*, published by Oxford University Press, 1994 (ISBN 0-19-854287-9); Kendrew et al. (eds.), *The Encyclopedia of Molecular Biology*, published by Blackwell Science Ltd., 1994 (ISBN 0-632-02182-9); and Robert A. Meyers (ed.), *Molecular Biology and Biotechnology: a Comprehensive Desk Reference*, published by VCH Publishers, Inc., 1995 (ISBN 1-56081-569-8).

[0061] In order to facilitate review of the various embodiments of the disclosure, the following explanations of specific terms are provided:

[0062] Adjuvant: A substance or vehicle that non-specifically enhances the immune response to an antigen. Adjuvants can include a suspension of minerals (alum, aluminum hydroxide, or phosphate) on which antigen is adsorbed; or water-in-oil emulsion in which antigen solution is emulsified in mineral oil (for example, Freund's incomplete adjuvant), sometimes with the inclusion of killed mycobacteria (Freund's complete adjuvant) to further enhance antigenicity. Immunostimulatory oligonucleotides (such as those including a CpG motif) can also be used as adjuvants (for example, see U.S. Pat. Nos. 6,194,388; 6,207,646; 6,214,806; 6,218,371; 6,239,116; 6,339,068; 6,406,705; and 6,429,199). Adjuvants also include biological molecules, such as costimulatory molecules. Exemplary biological adjuvants include IL-2, RANTES, GM-CSF, TNF- α , INF- γ , G-CSF, LFA-3, CD72, B7-1, B7-2, OX-40L and 41 BBL.

[0063] Administer: As used herein, administering a composition to a subject means to give, apply or bring the composition into contact with the subject. Administration can be accomplished by any of a number of routes, such as, for example, topical, oral, subcutaneous, intramuscular, intraperitoneal, intravenous, intrathecal and intramuscular.

[0064] Ambisense: Refers to a genome or genomic segments having both positive sense and negative sense portions. For example, the S segment of a Phlebovirus, such as Rift Valley fever virus, is ambisense, encoding nucleoprotein (NP) in the negative sense and the non-structural protein (NSs) in the positive sense.

[0065] Animal: Living multi-cellular vertebrate organisms, a category that includes, for example, mammals and bird. Mammals include, but are not limited to, humans, non-human primates, dogs, cats, horses, sheep and cows. The term mammal includes both human and non-human mammals.

[0066] Antibody: An immunoglobulin molecule produced by B lymphoid cells with a specific amino acid sequence. Antibodies are evoked in humans or other animals by a specific antigen (immunogen). Antibodies are characterized by reacting specifically with the antigen in some demonstrable way, antibody and antigen each being defined in terms of the other. "Eliciting an antibody response" refers to the ability of an antigen or other molecule to induce the production of antibodies.

[0067] Antigen: A compound, composition, or substance that can stimulate the production of antibodies or a T-cell

response in an animal, including compositions that are injected or absorbed into an animal. An antigen reacts with the products of specific humoral or cellular immunity, including those induced by heterologous immunogens. In one embodiment, an antigen is a RVF virus antigen.

[0068] Anti-genomic: As used herein, "anti-genomic" refers to a genomic segment of a virus that encodes a protein in the orientation opposite to the viral genome. For example, Rift valley fever virus is a negative-sense RNA virus. However, the S segment is ambisense, encoding proteins in both the positive-sense and negative-sense orientations. "Anti-genomic" refers to the positive-sense orientation, while "genomic" refers to the negative-sense orientation.

[0069] Attenuated: In the context of a live virus, the virus is attenuated if its ability to infect a cell or subject and/or its ability to produce disease is reduced (for example, eliminated) compared to a wild-type virus. Typically, an attenuated virus retains at least some capacity to elicit an immune response following administration to an immunocompetent subject. In some cases, an attenuated virus is capable of eliciting a protective immune response without causing any signs or symptoms of infection. In some embodiments, the ability of an attenuated virus to cause disease in a subject is reduced at least about 10%, at least about 25%, at least about 50%, at least about 75% or at least about 90% relative to wild-type virus.

[0070] Fusion protein: A protein generated by expression of a nucleic acid sequence engineered from nucleic acid sequences encoding at least a portion of two different (heterologous) proteins. To create a fusion protein, the nucleic acid sequences must be in the same reading frame and contain internal stop codons.

[0071] Hepatitis delta virus ribozyme: A non-coding, catalytic RNA from the hepatitis delta virus. Ribozymes catalyze the hydrolysis of their own phosphodiester bonds or those of other RNA molecules.

[0072] Host cell: In the context of the present disclosure, a "host cell" is a cell of use with the RVF virus reverse genetics systems described herein. A suitable host cell is one that is capable of transfection with and expression of the plasmids of the RVF virus reverse genetics system. In one embodiment, the host cell is a cell expressing the T7 polymerase, such as, but not limited to BSR-T7/5 cells (Buchholz et al., *J. Virol.* 73(1):251-259, 1999).

[0073] Immune response: A response of a cell of the immune system, such as a B-cell, T-cell, macrophage or polymorphonucleocyte, to a stimulus such as an antigen. An immune response can include any cell of the body involved in a host defense response, including for example, an epithelial cell that secretes an interferon or a cytokine. An immune response includes, but is not limited to, an innate immune response or inflammation. As used herein, a protective immune response refers to an immune response that protects a subject from infection (prevents infection or prevents the development of disease associated with infection).

[0074] Immunogen: A compound, composition, or substance which is capable, under appropriate conditions, of stimulating an immune response, such as the production of antibodies or a T-cell response in an animal, including compositions that are injected or absorbed into an animal. As used herein, as "immunogenic composition" is a composition comprising an immunogen.

[0075] Immunize: To render a subject protected from an infectious disease, such as by vaccination.

[0076] Isolated: An “isolated” biological component (such as a nucleic acid, protein or virus) has been substantially separated or purified away from other biological components (such as cell debris, or other proteins or nucleic acids). Biological components that have been “isolated” include those components purified by standard purification methods. The term also embraces recombinant nucleic acids, proteins or viruses, as well as chemically synthesized nucleic acids or peptides.

[0077] Linker: One or more amino acids that serve as a spacer between two polypeptides of a fusion protein.

[0078] Livestock: Domesticated animals reared in an agricultural setting as a source of food or to provide labor. The term “livestock” includes, but is not limited to, cattle, deer, donkeys, goats, horses, mules, rabbits and sheep.

[0079] ORF (open reading frame): A series of nucleotide triplets (codons) coding for amino acids without any termination codons. These sequences are usually translatable into a peptide.

[0080] Operably linked: A first nucleic acid sequence is operably linked with a second nucleic acid sequence when the first nucleic acid sequence is placed in a functional relationship with the second nucleic acid sequence. For instance, a promoter is operably linked to a coding sequence if the promoter affects the transcription or expression of the coding sequence. Generally, operably linked DNA sequences are contiguous and, where necessary to join two protein-coding regions, in the same reading frame.

[0081] Pharmaceutically acceptable vehicles: The pharmaceutically acceptable carriers (vehicles) useful in this disclosure are conventional. *Remington's Pharmaceutical Sciences*, by E. W. Martin, Mack Publishing Co., Easton, Pa., 15th Edition (1975), describes compositions and formulations suitable for pharmaceutical delivery of one or more therapeutic compounds or molecules, such as one or more recombinant RVF viruses, and additional pharmaceutical agents.

[0082] In general, the nature of the carrier will depend on the particular mode of administration being employed. For instance, parenteral formulations usually comprise injectable fluids that include pharmaceutically and physiologically acceptable fluids such as water, physiological saline, balanced salt solutions, aqueous dextrose, glycerol or the like as a vehicle. For solid compositions (for example, powder, pill, tablet, or capsule forms), conventional non-toxic solid carriers can include, for example, pharmaceutical grades of mannitol, lactose, starch, or magnesium stearate. In addition to biologically-neutral carriers, pharmaceutical compositions to be administered can contain minor amounts of non-toxic auxiliary substances, such as wetting or emulsifying agents, preservatives, and pH buffering agents and the like, for example sodium acetate or sorbitan monolaurate.

[0083] Plasmid: A circular nucleic acid molecule capable of autonomous replication in a host cell.

[0084] Polypeptide: A polymer in which the monomers are amino acid residues which are joined together through amide bonds. When the amino acids are alpha-amino acids, either the L-optical isomer or the D-optical isomer can be used. The terms “polypeptide” or “protein” as used herein are intended to encompass any amino acid sequence and include modified sequences such as glycoproteins. The term “polypeptide” is specifically intended to cover naturally occurring proteins, as well as those which are recombinantly or synthetically pro-

duced. The term “residue” or “amino acid residue” includes reference to an amino acid that is incorporated into a protein, polypeptide, or peptide.

[0085] Conservative amino acid substitutions are those substitutions that, when made, least interfere with the properties of the original protein, that is, the structure and especially the function of the protein is conserved and not significantly changed by such substitutions. Examples of conservative substitutions are shown below.

Original Residue	Conservative Substitutions
Ala	Ser
Arg	Lys
Asn	Gln, His
Asp	Glu
Cys	Ser
Gln	Asn
Glu	Asp
His	Asn; Gln
Ile	Leu, Val
Leu	Ile; Val
Lys	Arg; Gln; Glu
Met	Leu; Ile
Phe	Met; Leu; Tyr
Ser	Thr
Thr	Ser
Trp	Tyr
Tyr	Trp; Phe
Val	Ile; Leu

[0086] Conservative substitutions generally maintain (a) the structure of the polypeptide backbone in the area of the substitution, for example, as a sheet or helical conformation, (b) the charge or hydrophobicity of the molecule at the target site, or (c) the bulk of the side chain.

[0087] The substitutions which in general are expected to produce the greatest changes in protein properties will be non-conservative, for instance changes in which (a) a hydrophilic residue, for example, seryl or threonyl, is substituted for (or by) a hydrophobic residue, for example, leucyl, isoleucyl, phenylalanyl, valyl or alanyl; (b) a cysteine or proline is substituted for (or by) any other residue; (c) a residue having an electropositive side chain, for example, lysyl, arginyl, or histadyl, is substituted for (or by) an electronegative residue, for example, glutamyl or aspartyl; or (d) a residue having a bulky side chain, for example, phenylalanine, is substituted for (or by) one not having a side chain, for example, glycine.

[0088] Preventing, treating or ameliorating a disease: “Preventing” a disease refers to inhibiting the full development of a disease. “Treating” refers to a therapeutic intervention that ameliorates a sign or symptom of a disease or pathological condition after it has begun to develop. “Ameliorating” refers to the reduction in the number or severity of signs or symptoms of a disease.

[0089] Promoter: A promoter is an array of nucleic acid control sequences which direct transcription of a nucleic acid. A promoter includes necessary nucleic acid sequences near the start site of transcription. A promoter also optionally includes distal enhancer or repressor elements. A “constitutive promoter” is a promoter that is continuously active and is not subject to regulation by external signals or molecules. In contrast, the activity of an “inducible promoter” is regulated by an external signal or molecule (for example, a transcrip-

tion factor). In one embodiment, the promoter is a T7 promoter (from bacteriophage T7).

[0090] Purified: The term “purified” does not require absolute purity; rather, it is intended as a relative term. Thus, for example, a purified peptide, protein, virus, or other active compound is one that is isolated in whole or in part from naturally associated proteins and other contaminants. In certain embodiments, the term “substantially purified” refers to a peptide, protein, virus or other active compound that has been isolated from a cell, cell culture medium, or other crude preparation and subjected to fractionation to remove various components of the initial preparation, such as proteins, cellular debris, and other components.

[0091] Recombinant: A recombinant nucleic acid, protein or virus is one that has a sequence that is not naturally occurring or has a sequence that is made by an artificial combination of two otherwise separated segments of sequence. This artificial combination is often accomplished by chemical synthesis or, more commonly, by the artificial manipulation of isolated segments of nucleic acids, for example, by genetic engineering techniques. In one embodiment, recombinant RVF virus is generated using the reverse genetics system described herein. In some examples, the recombinant RVF viruses comprise one or more deletions in a viral virulence factor, such as NSs and/or NSm. In other examples, the recombinant RVF viruses comprise a heterologous gene, such as a reporter gene.

[0092] Reporter gene: A reporter gene is a gene operably linked to another gene or nucleic acid sequence of interest (such as a promoter sequence). Reporter genes are used to determine whether the gene or nucleic acid of interest is expressed in a cell or has been activated in a cell.

[0093] Reporter genes typically have easily identifiable characteristics, such as fluorescence, or easily assayed products, such as an enzyme. Reporter genes can also confer antibiotic resistance to a host cell or tissue. Reporter genes include, for example, GFP (or eGFP) or other fluorescence genes, luciferase, β -galactosidase and alkaline phosphatase.

[0094] Reverse genetics: Refers to the process of introducing mutations (such as deletions, insertions or point mutations) into the genome of an organism or virus in order to determine the phenotypic effect of the mutation. For example, introduction of a mutation in a specific viral gene enables one to determine the function of the gene.

[0095] Rift Valley fever (RVF) virus: A virus belonging to the family Bunyaviridae and genus Phlebovirus. RVF virus has a single-stranded, negative-sense genome composed of three genome segments, S, M and L. The S segment is an ambisense genome segment, meaning it encodes proteins in both the positive-sense and negative-sense orientations. The RVF virus genome encodes both structural and non-structural proteins. A “structural” protein is a protein found in the virus particle, whereas a “non-structural” protein is only expressed in a virus-infected cell. RVF virus structural proteins include nucleoprotein (NP or N, used interchangeably), two glycoproteins (Gn and Ge) and the viral RNA-dependent RNA polymerase (L protein). Non-structural RVF virus proteins include NSs, NSm and the NSm+Gn fusion protein. As used herein, a “full-length” RVF virus genome segment is one containing no deletions. Full-length genome segments can contain mutations or substitutions, but retain the same length as the wild-type virus. A “complete deletion” of an ORF of a RVF virus genome segment means either every nucleotide encoding the ORF is deleted from genome segment, or nearly

every nucleotide encoding the ORF is deleted such that no proteins are translated from the ORF. Thus, a “complete deletion” includes genome segments retaining up to ten nucleotides encoding the ORF, such as one, two, three, four, five, six, seven, eight, nine or ten nucleotides. In other A number of RVF virus strains have been identified. In one embodiment described herein, the RVF virus strain is ZH501.

[0096] As used herein, plasmids encoding full-length RVF virus S, M and L genome segments are referred to as pRVS, pRVM, and pRVL, respectively. The S segment plasmid containing the eGFP ORF in place of the complete NSs ORF is referred to as pRVS-GFPANSSs. The M segment plasmid containing a complete deletion of the NSm ORF is referred to as pRVM- Δ NSm. The recombinant RVF viruses based upon the ZH501 genome that are generated using reverse genetics are referred to as either rRVF or rZH501. For example, recombinant RVF virus generated using the pRVS-GFPANSSs plasmid (and wild-type M and L plasmids), is referred to as rRVF- Δ NSs:GFP or rZH501- Δ NSs:GFP. Similarly, recombinant RVF virus generated using the pRVS-GFPANSSs plasmid and pRVM- Δ NSm plasmid (and wild-type L plasmid), referred to as rRVF- Δ NSs:GFP- Δ NSm or rZH501- Δ NSs:GFP- Δ NSm. Recombinant RVF virus comprising wild-type M and L segments, and an S segment encoding an NSs-eGFP fusion protein with a three alanine residue linker, is referred to as rRVF-NSs(Ala)₃GFP or rZH501-NSs(Ala)₃GFP. If the fusion protein comprises a ten alanine residue linker, the recombinant virus is referred to as rRVF-NSs(Ala)₁₀GFP or rZH501-NSs(Ala)₁₀GFP.

[0097] Sequence identity: The similarity between amino acid or nucleic acid sequences is expressed in terms of the similarity between the sequences, otherwise referred to as sequence identity. Sequence identity is frequently measured in terms of percentage identity (or similarity or homology); the higher the percentage, the more similar the two sequences are. Homologs or variants of a given gene or protein will possess a relatively high degree of sequence identity when aligned using standard methods.

[0098] Methods of alignment of sequences for comparison are well known in the art. Various programs and alignment algorithms are described in: Smith and Waterman, *Adv. Appl. Math.* 2:482, 1981; Needleman and Wunsch, *J. Mol. Biol.* 48:443, 1970; Pearson and Lipman, *Proc. Natl. Acad. Sci. U.S.A.* 85:2444, 1988; Higgins and Sharp, *Gene* 73:237-244, 1988; Higgins and Sharp, *CABIOS* 5:151-153, 1989; Comet et al., *Nucleic Acids Research* 16:10881-10890, 1988; and Pearson and Lipman, *Proc. Natl. Acad. Sci. U.S.A.* 85:2444, 1988. Altschul et al., *Nature Genet.* 6:119-129, 1994.

[0099] The NCBI Basic Local Alignment Search Tool (BLAST™) (Altschul et al., *J. Mol. Biol.* 215:403-410, 1990) is available from several sources, including the National Center for Biotechnology Information (NCBI, Bethesda, Md.) and on the Internet, for use in connection with the sequence analysis programs blastp, blastn, blastx, tblastn and tblastx.

[0100] Subject: Living multi-cellular vertebrate organisms, a category that includes both human and non-human mammals. Subjects include veterinary subjects, including livestock such as cows and sheep, and non-human primates.

[0101] Therapeutically effective amount: A quantity of a specified agent sufficient to achieve a desired effect in a subject being treated with that agent. For example, this may be the amount of a recombinant RVF virus useful for eliciting an immune response in a subject and/or for preventing infection by RVF virus. Ideally, in the context of the present dis-

closure, a therapeutically effective amount of a recombinant RVF virus is an amount sufficient to increase resistance to, prevent, ameliorate, and/or treat infection caused by RVF virus in a subject without causing a substantial cytotoxic effect in the subject. The effective amount of a recombinant RVF virus useful for increasing resistance to, preventing, ameliorating, and/or treating infection in a subject will be dependent on, for example, the subject being treated, the manner of administration of the therapeutic composition and other factors.

[0102] Transformed: A transformed cell is a cell into which has been introduced a nucleic acid molecule by molecular biology techniques. As used herein, the term transformation encompasses all techniques by which a nucleic acid molecule might be introduced into such a cell, including transfection with viral vectors, transformation with plasmid vectors, and introduction of naked DNA by electroporation, lipofection, and particle gun acceleration.

[0103] Vaccine: A preparation of immunogenic material capable of stimulating an immune response, administered for the prevention, amelioration, or treatment of infectious or other types of disease. The immunogenic material may include attenuated or killed microorganisms (such as attenuated viruses), or antigenic proteins, peptides or DNA derived from them. Vaccines may elicit both prophylactic (preventative) and therapeutic responses. Methods of administration vary according to the vaccine, but may include inoculation, ingestion, inhalation or other forms of administration. Inoculations can be delivered by any of a number of routes, including parenteral, such as intravenous, subcutaneous or intramuscular. Vaccines may be administered with an adjuvant to boost the immune response.

[0104] Unless otherwise explained, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this disclosure belongs. The singular terms "a," "an," and "the" include plural referents unless context clearly indicates otherwise. Similarly, the word "or" is intended to include "and" unless the context clearly indicates otherwise. Hence "comprising A or B" means including A, or B, or A and B. It is further to be understood that all base sizes or amino acid sizes, and all molecular weight or molecular mass values, given for nucleic acids or polypeptides are approximate, and are provided for description. Although methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present disclosure, suitable methods and materials are described below. All publications, patent applications, patents, and other references mentioned herein are incorporated by reference in their entirety. In case of conflict, the present specification, including explanations of terms, will control. In addition, the materials, methods, and examples are illustrative only and not intended to be limiting.

III. Overview of Several Embodiments

[0105] Disclosed herein are RVF virus vaccine candidates that are highly attenuated, immunogenic and contain precise molecular markers allowing for the differentiation of naturally infected and vaccinated animals (DIVA). The recombinant RVF viruses described herein, which in some embodiments contain complete deletions of one or both of the major virus virulence factors, NSs and NSm, are highly attenuated in vivo, induce robust anti-RVF antibody responses, provide protection from virulent virus challenge and allow for the assessment of vaccination status in animals on the basis of

NSs/NP serology. Further provided are recombinant RVF viruses comprising a reporter moiety, such as eGFP, which have a variety of applications, including as live virus research tools that can be useful for the rapid screening of antiviral therapeutic compounds.

[0106] Provided herein are recombinant RVF viruses, wherein the genome of the recombinant RVF viruses comprise (i) a full-length L segment; (ii) a full-length M segment or an M segment comprising a complete deletion of the NSm open reading frame (ORF); and (iii) an S segment comprising a complete deletion of the NSs ORF. In one embodiment, the NSs ORF of the recombinant RVF virus is replaced by the ORF of a reporter gene.

[0107] Reporter genes include, but are not limited to genes encoding fluorescent proteins, enzymes or antibiotic resistance. Any gene that produces a protein with a functional readout can be used as the reporter gene. In one example, the reporter gene is a green fluorescent protein (GFP), such as enhanced GFP (eGFP).

[0108] The genome of the recombinant RVF viruses provided herein can be derived from any strain or variant of RVF virus. In some embodiments, the genome is derived from ZH501, ZH548, SA75 or SPB 9800523. In a preferred embodiment, the genome is derived from RVF virus strain ZH501. The sequences of the S, M and L segments of the ZH501 strain are deposited under GenBank Accession Nos. DQ380149 (SEQ ID NO: 1), DQ380200 (SEQ ID NO: 2) and DQ375406 (SEQ ID NO: 3), respectively. The nucleotide sequences of the S, M and L segments need not be 100% identical to the sequences provided herein. In some examples, the S, M and L segments are at least 80%, at least 85%, at least 90%, at least 95%, or at least 99% identical to a known or disclosed RVF virus S, M or L segment, such as the S, M and L segments set forth as SEQ ID NO: 1, SEQ ID NO: 2 and SEQ ID NO: 3, respectively.

[0109] Also provided are immunogenic compositions comprising the recombinant RVF viruses described herein and a pharmaceutically acceptable carrier. Suitable pharmaceutical carriers are described herein and are well known in the art. The pharmaceutical carrier used depends on a variety of factors, including the route of administration. In one embodiment, the immunogenic composition further comprises an adjuvant. The adjuvant can be any substance that improves the immune response to the recombinant RVF virus.

[0110] Further provided is a method of immunizing a subject against RVF virus infection, comprising administering to the subject an immunogenic composition disclosed herein. In one embodiment, the subject is livestock. Livestock includes, but is not limited to sheep and cattle. In another embodiment, the subject is a human. In one example, the immunogenic composition is administered in a single dose. In another embodiment, the immunogenic composition is administered in multiple doses, such as two, three or four doses. When administered in multiple doses, the time period between doses can vary. In some cases, the time period is days, weeks or months. The immunogenic composition can be administered using any suitable route of administration. Generally, the recombinant RVF viruses are administered parenterally, such as intramuscularly, intravenously or subcutaneously.

[0111] Further provided is a collection of plasmids comprising (i) a plasmid encoding a full-length anti-genomic copy of the L segment of RVF virus; (ii) a plasmid encoding a full-length anti-genomic copy of the M segment of RVF virus, or an anti-genomic copy of the M segment of RVF virus

comprising a complete deletion of the NSm ORF; and (iii) a plasmid encoding an anti-genomic copy of the S segment of RVF virus, wherein the S segment comprises a complete deletion of the NSs ORF. In some embodiments, the plasmids further comprise a T7 promoter and a hepatitis delta virus ribozyme. In one example, the NSs ORF of the S segment plasmid is replaced by the ORF of a reporter gene. In one example, the reporter gene is green fluorescent protein. In some embodiments, the RVF virus is ZH501.

[0112] In one embodiment, the nucleotide sequence of the S segment plasmid is at least 95% identical to the nucleotide sequence of SEQ ID NO: 14. In particular examples, the nucleotide sequence of the S segment plasmid comprises the nucleotide sequence of SEQ ID NO: 14. In another embodiment, the nucleotide sequence of the M segment plasmid is at least 95% identical to the nucleotide sequence of SEQ ID NO: 12. In particular examples, the nucleotide sequence of the M segment plasmid comprises the nucleotide sequence of SEQ ID NO: 12. In another embodiment, the nucleotide sequence of the M segment plasmid is at least 95% identical to the nucleotide sequence of SEQ ID NO: 15. In particular examples, the nucleotide sequence of the M segment plasmid comprises the nucleotide sequence of SEQ ID NO: 15. In another embodiment, the nucleotide sequence of the L segment plasmid is at least 95% identical to the nucleotide sequence of SEQ ID NO: 13. In particular examples, the nucleotide sequence of the L segment plasmid comprises the nucleotide sequence of SEQ ID NO: 13.

[0113] Also provided are isolated host cells comprising the collection of plasmids provided herein. In one embodiment, the cells express T7 polymerase.

[0114] Further provided is a method of preparing a recombinant RVF virus for immunization of a subject, comprising (i) transfecting cultured cells with the collection of plasmids described herein; (ii) incubating the cells for 1 to 5 days; and (iii) collecting recombinant RVF virus from the cell supernatant. In one embodiment, the cells express T7 polymerase.

[0115] Further provided are recombinant RVF viruses, wherein the genome of the recombinant RVF viruses comprise a full-length L segment, a full-length M segment and a full-length S segment, wherein the S segment further encodes the ORF of a reporter gene. In one embodiment, the S segment encodes a NSs-reporter gene fusion protein. In one aspect, the reporter gene is fused to the C-terminus of NSs. In some examples, the fusion protein further comprises a linker. The linker can be any suitable combination of one or more amino acids. In one embodiment, the linker comprises 3 to 10 alanine residues. In another embodiment, the reporter gene is GFP.

[0116] Also provided herein is a reverse genetics system for producing recombinant RVF virus. The reverse genetics system consists of three plasmids, wherein a first plasmid encodes an anti-genomic copy of a S segment, a second plasmid encodes an anti-genomic copy of a M segment and a third plasmid encodes an anti-genomic copy of a L segment of RVF virus, and wherein each plasmid comprises a T7 promoter and a hepatitis delta virus ribozyme. In some embodiments, the RVF virus is ZH501. In some embodiments, the S, M and L segments are wild-type S, M and L segments. In another embodiment, the S segment comprises a deletion of the NSs ORF. In another embodiment, the M segment comprises a deletion of the NSm ORF. In another embodiment, the S segment comprises a deletion of the NSs ORF and the M segment comprises a deletion of the NSm ORF.

[0117] In one embodiment, the nucleotide sequence of the S segment plasmid is at least 95% identical to the nucleotide sequence of SEQ ID NO: 11. In particular examples, the nucleotide sequence of the S segment plasmid comprises the nucleotide sequence of SEQ ID NO: 11. In another embodiment, the nucleotide sequence of the S segment plasmid is at least 95% identical to the nucleotide sequence of SEQ ID NO: 14. In particular examples, the nucleotide sequence of the S segment plasmid comprises the nucleotide sequence of SEQ ID NO: 14. In another embodiment, the nucleotide sequence of the M segment plasmid is at least 95% identical to the nucleotide sequence of SEQ ID NO: 12. In particular examples, the nucleotide sequence of the M segment plasmid comprises the nucleotide sequence of SEQ ID NO: 12. In another embodiment, the nucleotide sequence of the M segment plasmid is at least 95% identical to the nucleotide sequence of SEQ ID NO: 15. In particular examples, the nucleotide sequence of the M segment plasmid comprises the nucleotide sequence of SEQ ID NO: 15. In another embodiment, the nucleotide sequence of the L segment plasmid is at least 95% identical to the nucleotide sequence of SEQ ID NO: 13. In particular examples, the nucleotide sequence of the L segment plasmid comprises the nucleotide sequence of SEQ ID NO: 13.

[0118] Typically, the three plasmids are transfected simultaneously into cells expressing T7 polymerase. After a sufficient period of time to allow for production of recombinant virus, such as about 1 to about 5 days, recombinant RVF virus is collected from the cell supernatant.

[0119] Further provided are recombinant RVF viruses produced using the reverse genetics system described herein. In some examples, the recombinant RVF viruses comprise a deletion in one or more viral virulence factors. In one embodiment, the recombinant viruses comprise a complete deletion of the NSs ORF. In another embodiment, the recombinant viruses comprise a complete deletion of the NSm ORF. In yet another embodiment, the recombinant viruses comprise a deletion in both the NSs and NSm ORFs. The recombinant **[0120]** RVF viruses can be used for both research and therapeutic purposes. For example, the recombinant RVF viruses described herein are suitable for use as vaccines to prevent, treat or ameliorate RVF virus infection in livestock and humans.

IV. Rift Valley Fever Virus Genome and Encoded Proteins

[0121] Like other members of the genus Phlebovirus, RVF virus has a negative-sense, single-stranded tripartite RNA genome composed of the S, M and L segments (Schmaljohn, and Hooper, *Bunyaviridae: The Viruses and their Replication In Fields' Virology*, Lippincott Williams & Wilkins, Philadelphia, Pa., 2001). The small (S) segment (about 1.6 kB) encodes, in an ambisense fashion, the virus nucleoprotein (NP) in the genomic (-) orientation, and the non-structural (NSs) protein in the anti-genomic (+) orientation (Albarino et al., *J. Virol.* 81:5246-5256, 2007). The medium (M) segment (about 3.8 kB) encodes a least four nested proteins in a single ORF, including two structural glycoproteins, Gn and Ge, and two nonstructural proteins, the 14 kD NSm and a 78 kD NSm+Gn fusion (Gerrard et al., *Virology* 359:459-465, 2007; Gerrard and Nichol, *Virology* 357:124-133, 2007). The M segment contains five conserved in-frame AUG-methionine start codons within the NSm protein coding region at anti-genomic sense positions 21, 135, 174, 411 and 426. Alternate

utilization of the AUGs at positions 21 or 135 results in expression of the 14 kDa NSm protein and the 78 kD NSm+Gn fusion protein (Suzich et al., *J. Virol.* 64:1549-1555, 1990; Gerrard and Nichol, *Virology* 357:124-133, 2007). The large (L) segment (about 6.4 kB) encodes the viral RNA-dependent RNA polymerase (L protein).

[0122] RVF virus NP and L proteins are required for viral RNA synthesis (Ikegami et al., *J. Virol.* 79:5606-5615, 2005; Lopez et al., *Virology* 69:3972-3979, 1995). Gn and Gc are believed to mediate binding to an as yet unidentified receptor. The 78kD NSm+Gn fusion protein has been reported to be dispensable for viral replication in cell culture (Won et al., *J. Virol.* 80:8274-8278, 2006).

[0123] Both nonstructural genes (NSs and NSm) have been reported to function as virus virulence factors and determinants of mammalian host pathogenesis (Bird et al., *Virology* 362:10-15, 2007; Vialat et al., *Virology* 74:1538-1543, 2000; Won et al., *J. Virol.* 24:13335-13345, 2007). NSs mediates the pan-downregulation of mRNA production by inhibition of RNA polymerase II activity (Billecocq et al., *J. Virol.* 78:9798-9806, 2004; Le May et al., *Cell* 116:541-550, 2004). Via this mechanism, the NSs protein performs a critical role in mammalian host pathogenesis by indirectly disrupting the host cell antiviral response (Bouloy et al., *J. Virol.*, 75:1371-1377, 2001; Muller et al., *Arm. J. Trop. Med. Hyg.* 53:405-411, 1995; Vialat et al., *Virology* 74:1538-1543, 2000).

[0124] Studies of the non-structural gene located on the RVF virus M segment (NSm) indicate it is dispensable for efficient RVF virus growth in both IFN-competent and IFN-deficient cell culture (Gerrard et al., *Virology* 359:459-465, 2007; Won et al., *J. Virol.* 80:8274-8278, 2006). However, further work utilizing a highly sensitive animal model revealed that recombinant RVF virus lacking the entire NSm coding region (rRVF-ΔNSm) was attenuated yet retained the ability to cause either acute-onset lethal hepatic necrosis or delayed-onset lethal neurologic disease in a minority (44%) of animals (Bird et al., *Virology* 362:10-15, 2007). Other studies have shown that NSm functions as a virus virulence factor by suppressing the host cell apoptotic pathway following infection (Won et al., *J. Virol.* 24:13335-13345, 2007).

V. Reverse Genetics System for RVF Virus

[0125] The ability to generate recombinant viruses containing selected mutations and/or deletions is a powerful tool for the development of virus vaccines. Reverse genetics systems have been described for a few viruses of the Bunyaviridae family, including Bunyamvera virus (Bridgen and Elliott, *Proc. Natl. Acad. Sci. USA* 93:15400-15404, 1996) and La Crosse virus (Blakqori and Weber, *J. Virol.* 79:10420-10428, 2005). Recently, a reverse genetics system for a vaccine strain (MP-12) of RVF virus was reported (U.S. Pre-Grant Publication No. 2007/0122431, herein incorporated by reference).

[0126] The recombinant RVF viruses described herein are generated using an optimized reverse genetics system capable of rapidly generating wild-type and mutant viruses (Bird et al., *Virology* 362:10-15, 2001; Gerrard et al., *Virology* 359:459-465, 2007, each of which is herein incorporated by reference). The RVF virus reverse genetics system is a T7 RNA polymerase-driven plasmid-based genetic system based on the genome of the virulent RVF virus Egyptian isolate ZH501. This system, described in detail in Bird et al. (*Virology* 362:10-15, 2001) and in the Examples below, includes three plasmids expressing anti-genomic copies of the S, M

and L segments of ZH501. As used herein, the plasmids are referred to as the pRVS, pRVM and pRVL plasmids, respectively.

[0127] Rescue of recombinant viruses is accomplished by simultaneous transfection of the three anti-genomic sense plasmids into cells stably expressing T7 polymerase (for example, BSR-T7/5 cells). The genome segments of each plasmid are flanked by a T7 promoter, which enables generation of the primary RNA transcript, and the hepatitis delta virus ribozyme, which removes extraneous nucleotides from the 3' end of the primary transcriptional products. The T7 RNA polymerase generated transcripts are identical copies of the RVF virus genome segments, with the exception of an extra G nucleotide on the 5' end derived from the T7 promoter. When expressed in transfected host cells, the pRVS, pRVM and pRVL plasmids generate anti-genomic sense copies of the S, M and L segments, respectively.

[0128] In one embodiment, the recombinant RVF viruses are generated using an S-segment plasmid that comprises a deletion in the NSs gene. In one example, the deletion is a deletion of the entire NSs ORF. In one aspect, the NSs ORF is replaced by the eGFP ORF. In another embodiment, the recombinant RVF viruses are generated using an S-segment plasmid comprising a deletion in the NSs gene and an M-segment plasmid comprising a deletion in the NSm ORF. In one example, the deletion is a deletion of the entire NSm ORF. In another embodiment, the recombinant RVF viruses are generated using full-length S, M and L plasmids, wherein one of the plasmids further encodes eGFP. In one example, the eGFP ORF is encoded by the S plasmid as a NSs-eGFP fusion protein.

VI. Use of Recombinant RVF Viruses

[0129] Recombinant RVF viruses generated using the reverse genetics system described herein can be used for both research and therapeutic purposes. Using this system, recombinant RVF viruses can be generated that contain precisely defined deletions of major virus virulence factors on one or more genome segments. For example, viruses can be produced that contain deletions of NSs and/or NSm. Accordingly, such recombinant RVF viruses can be used as vaccines to prevent infection of livestock and humans with wild-type RVF virus. The recombinant RVF viruses described herein can also be used as live-virus research tools, particularly those viruses that incorporate reporter genes, for instance a fluorescent protein such as GFP. For example, these viruses can be used for high-throughput screening of antiviral compounds in vitro.

[0130] Efforts to prevent RVF virus infection via vaccination began shortly after the first isolation of the virus in 1931 (Findlay and Daubney, *Lancet* ii:1350-1351, 1931). These earliest vaccines (MacKenzie, *J. Pathol. Bacteriol.* 40:65-73, 1935) and several that followed, including the currently available TSI-GSD-200 preparation, relied on formalin inactivation of live wild-type virus (Pittman et al., *Vaccine* 18:181-189, 1999; Randall et al., *J. Immunol.* 89:660-671, 1962). While capable of eliciting protective immune responses among livestock and humans, these inactivated vaccines typically require a series of 2 or 3 initial inoculations, followed by regular booster vaccinations to achieve and maintain protection (Pittman et al., *Vaccine* 18:181-189, 1999; Swanepoel et al., "Rift Valley fever" in *Infectious Diseases of livestock with special reference to South Africa*, pages 688-717, Oxford university Press, Cape Town). However, multiple dosing and

annual vaccination regimens are logistically difficult to implement and expensive to maintain, and thus are of limited practical value in resource-poor settings, especially in regard to control of RVF virus infection in livestock in enzootic settings. In addition, there have been problems in the past with quality control and “inactivated” vaccines causing disease.

[0131] In an effort to eliminate the necessity of booster inoculations, several live-attenuated vaccine candidates were developed for RVF virus with some, such as the Smithburn neurotropic strain, being employed in Africa. These vaccine candidates have relied upon the random introduction of attenuating mutations via serial passage in suckling mouse brain or tissue culture, in vitro passage in the presence of chemical mutagens, such as 5-fluorouracil, or as naturally occurring virus isolates (such as the Smithburn neurotropic strain, the Kenyan-IB8 strains, MP-12, or the Clone 13 isolate) (Caplen et al., *J. Gen. Virol.* 66:2271-2277, 1985; Coackley, *J. Pathol. Bacteriol.* 89:123-131, 1965; Moussa et al., *Am. J. Trop. Med. Hyg.* 35:660-662, 1986; Muller et al., *Am. J. Trop. Med. Hyg.* 53:405-411, 1995; Rossi and Turell, *J. Gen. Virol.* 69:817-823, 1988; Smithburn, *Br. J. Exp. Pathol.* 30:1-16, 1949).

[0132] Due to the technical limitations of these procedures, and the lack of complete genome sequence for many of the historically derived RVF virus vaccines, the exact underlying molecular mechanisms of attenuation for many of these live-attenuated RVF virus vaccines is either unknown (Smithburn neurotropic strain or Kenyan-IB8) or reliant on the combinatorial effects of multiple nucleotide or amino acid substitutions (MP-12) (Saluzzo and Smith, *Vaccine* 8:369-375, 1990; Takehara et al., *Virology* 169:452-457, 1989). Experimental and field experience with existing live-attenuated RVF virus vaccines demonstrated that in certain instances these vaccines retain the ability to cause teratogenic effects, abortion, and neural pathology in livestock or animal models. Thus, widespread use of these live-attenuated vaccines is problematic, especially in non-endemic areas, or during inter-epizootic/epidemic periods (Hunter et al., *Onderstepoort J. Vet. Res.* 69:95-98, 2002; Morrill et al., *Am. J. Vet. Res.* 58:1104-1109, 1997; Morrill et al., *Am. J. Vet. Res.* 58:1110-1114, 1997; Morrill and Peters, *Vaccine* 21:2994-3002, 2003).

[0133] While useful in many situations, several distinct disadvantages exist among live attenuated RNA virus vaccines prepared by the traditional techniques discussed above. Live-attenuated vaccines reliant on single or multiple nucleotide substitutions are at increased risk for reversion to virulent phenotypes due to the inherently high rate of viral RNA polymerase errors. The loss of attenuation via this mechanism among livestock and human live vaccines has been documented (Berkhout et al., *J. Virol.* 73:1138-1145, 1999; Catelli et al., *Vaccine* 24:6476-6482, 2006; Halstead et al., *Am. J. Trop. Med. Hyg.* 33:672-678, 1984; Hopkins and Yoder, *Avian Dis.* 30:221-223, 1986; Rahimi et al., *J. Clin. Virol.* 39:304-307, 2007).

[0134] The potential for a similar reversion event among live RVF virus vaccines dependent on attenuating nucleotide mutation was illustrated by recent genomic analyses of RVF virus that revealed an overall molecular evolution rate ($\sim 2.5 \times 10^{-4}$ nucleotide substitutions/site/year) similar to other single-stranded RNA viruses (Bird et al., *J. Virol.* 81:2805-2816, 2007). Due to error-prone polymerases, live-attenuated RNA virus vaccines prepared by multiple serial passage techniques involved in virus attenuation inherently consist of a

complex mixture of genomic micro-variants. In contrast, the origin of reverse genetics derived virus vaccine candidates is advantageous in that vaccine stocks can be generated directly or following limited amplification steps from precisely defined DNA plasmids. This approach allows for the simple production of virus vaccines following good manufacturing processes (GMP) with higher levels of genetic homogeneity.

[0135] Another significant drawback of all previously generated live-attenuated RVF virus vaccines is that they do not allow for differentiation of naturally infected from vaccinated animals (DIVA). This ability is important to augment efforts to contain an accidental or intentional release of wild-type RVF virus in previously unaffected areas (Henderson, *Biologicals* 33:203-209, 2005). As a high consequence pathogen, RVF virus has been classified as a category A Select Agent as defined by the United States Department of Health and Human Services and the United States Department of Agriculture (USDA), and is listed as a high consequence agent with potential for international spread (List A) by the Office International des Epizooties (OIE) (Le May et al., *Cell* 116: 541-550, 2004) of the World Organization for Animal Health (WOAH), thus greatly increasing the consequences for international livestock trade following the introduction of RVF virus into previously unaffected countries or epizootics in enzootic areas (USDA, 7 CFR Part 331 and 9 CFR Part 121, Federal Register RN 0579-AB47:13241-13292, 2005). Currently, OIE regulations require surveillance and absence of RVF virus activity for 2 years following an outbreak before resumption of disease free status and the subsequent easing of import/export trade restrictions (International Office of Epizootics, Terrestrial Animal Health Code, XI:2.2.14.1, 2007). The use of any current commercially available livestock vaccines does not permit the differentiation of vaccinated from naturally infected livestock, thus contraindicating the use of prophylactic vaccination in countries wishing to retain disease free status, or in those with ongoing/endemic RVF virus activity.

[0136] Thus, disclosed herein are infectious recombinant RVF viruses, generated using reverse genetics, containing either complete deletions of major virus virulence factors, NSs (rZH501- Δ NSs:GFP) or NSs and NSm (rZH501- Δ NSs:GFP- Δ NSm), which confer attenuated phenotypes in vivo, and which allow for the serologic differentiation of naturally infected and vaccinated animals by presence/absence of anti-RVF NP/anti-RVF NSs antibodies. As described herein, in vivo testing of these recombinant RVF (rRVF) viruses demonstrated that they were highly immunogenic and efficacious in the prevention of severe RVF virus disease and lethality (FIG. 2 and FIG. 4). In an initial pilot study, animals developed high end-point titers ($\geq 1:400$) of total anti-RVF virus IgG by day 21 post-vaccination that were significantly higher than sham-inoculated controls (p-value <0.05, Table 1). At no observed time point post-vaccination did any animal develop disease symptoms or vaccine virus induced viremia (FIG. 4).

[0137] Additional testing in a larger follow-up study confirmed these results with the majority of animals generating robust total anti-RVF IgG responses with typical titers $\geq 1:6400$ by day 26 post-vaccination. As with the pilot study, no vaccine virus induced viremia was detected. The immunologic response generated in the Δ NSs/ Δ NSm virus vaccinated animals was significantly higher than controls (p-value <0.05) and was sufficient to confer complete protection from

both clinical illness and lethality in 100% of vaccinated animals given a known lethal challenge dose of wild-type RVF virus (FIG. 4).

[0138] Direct comparisons of the level of protective immunity (PRNT₅₀ or total IgG) titers with previous studies utilizing other RVF virus vaccines are difficult due to vaccine used and species level differences in immunity. However, earlier studies utilizing the 3 dose regimen (day 0, 7 and 28) of inactivated TSI-GSD-200 vaccine in the WF rat model demonstrated protective efficacy against virulent virus challenge at PRNT₈₀ titers >1:40 (Anderson et al., *Vaccine* 9:710-714, 1991). Later, a large retrospective analysis of human volunteers (n=598), receiving the same recommended 3-dose regimen of this inactivated vaccine were found to develop mean PRNT₈₀ titers of 1:237 (Pittman et al., *Vaccine* 18:181-189, 1999). Additionally, a large study of the pathogenesis and neurovirulence of the live-attenuated MP-12 vaccine in rhesus macaques demonstrated PRNT₈₀ titers among vaccinated animals of \geq 1:640 (Morrill and Peters, *Vaccine* 21:2994-3002, 2003). As described herein, inoculation with a single dose of the recombinant RVF viruses resulted in mean PRNT₅₀ titers ranging from 1:640 to 1:7040, indicating that the level of neutralizing antibody is similar to or greater than that demonstrated in earlier RVF vaccine studies using multiple doses in animal model systems or among human volunteers.

[0139] Thus, the enhanced safety, attenuation, and reduced possibility of reversion to full virulence (via either RVF virus polymerase nucleotide substitution or gene segment reassortment with field-strains) afforded by the double genetic deletions of the entire RVF virus NSs and NSm genes, does not diminish overall vaccine efficacy. A high level of protective immunity was induced by a single dose of the rRVF viruses disclosed herein, with 37 of 40 total vaccinated animals developing a potentially sterilizing immunity as determined by the lack of any detectable post-challenge viremia (FIG. 4).

[0140] As described herein, animals immunized with either recombinant RVF virus do not have detectable anti-NSs antibodies. Thus, given the high-level anti-NSs antibody in survivor control animals, DIVA will be possible among animals immunized with these candidate vaccines based on the presence/absence of anti-NSs antibody (FIGS. 3A-C). Anti-NSs antibodies have also been detected in the serum of naturally infected convalescent livestock obtained during the outbreak in Saudi Arabia in 2000 (FIG. 3D), and in humans. Therefore, the use of these recombinant RVF viruses, combined with the further development of rapid ELISA or solid matrix-based differential detection assays for anti-NP/anti-NSs antibodies, can provide a robust DIVA field screening tool.

[0141] In addition, the recombinant RVF viruses described herein routinely grew to high titers in tissue culture and provided protective immunity following a single injection, thus likely reducing the overall economic cost of production, and potentially eliminating the need for resource intensive follow-up booster inoculations. Additionally, the precisely defined attenuating deletions and use of cDNA technology eliminates the potential risk of reversion to, or contamination from, virulent wild-type virus inherent in serially passaged or inactivated vaccine preparations, and may ease the federal/national regulatory approval process. While the recombinant RVF viruses can be targeted towards veterinary medical use, and thus indirectly the prevention of human RVF disease, the candidate vaccines can also provide effective prophylactic protection for humans, such as those in high risk occupational

settings, or in recognized risk groups following natural or intentional introduction of RVF virus into previously unaffected areas.

VII. Administration of Recombinant RVF Virus for Vaccination

[0142] Recombinant RVF viruses, or immunogenic compositions thereof, can be administered to a subject by any of the routes normally used for introducing recombinant virus into a subject. Methods of administration include, but are not limited to, intradermal, intramuscular, intraperitoneal, parenteral, intravenous, subcutaneous, vaginal, rectal, intranasal, inhalation or oral. Parenteral administration, such as subcutaneous, intravenous or intramuscular administration, is generally achieved by injection. Injectables can be prepared in conventional forms, either as liquid solutions or suspensions, solid forms suitable for solution or suspension in liquid prior to injection, or as emulsions. Injection solutions and suspensions can be prepared from sterile powders, granules, and tablets of the kind previously described. Administration can be systemic or local.

[0143] Immunogenic compositions are administered in any suitable manner, such as with pharmaceutically acceptable carriers. Pharmaceutically acceptable carriers are determined in part by the particular composition being administered, as well as by the particular method used to administer the composition. Accordingly, there is a wide variety of suitable formulations of pharmaceutical compositions of the present disclosure.

[0144] Preparations for parenteral administration include sterile aqueous or non-aqueous solutions, suspensions, and emulsions. Examples of non-aqueous solvents are propylene glycol, polyethylene glycol, vegetable oils such as olive oil, and injectable organic esters such as ethyl oleate. Aqueous carriers include water, alcoholic/aqueous solutions, emulsions or suspensions, including saline and buffered media. Parenteral vehicles include sodium chloride solution, Ringer's dextrose, dextrose and sodium chloride, lactated Ringer's, or fixed oils. Intravenous vehicles include fluid and nutrient replenishers, electrolyte replenishers (such as those based on Ringer's dextrose), and the like. Preservatives and other additives may also be present such as, for example, antimicrobials, anti-oxidants, chelating agents, and inert gases and the like.

[0145] Some of the compositions may potentially be administered as a pharmaceutically acceptable acid- or base-addition salt, formed by reaction with inorganic acids such as hydrochloric acid, hydrobromic acid, perchloric acid, nitric acid, thiocyanic acid, sulfuric acid, and phosphoric acid, and organic acids such as formic acid, acetic acid, propionic acid, glycolic acid, lactic acid, pyruvic acid, oxalic acid, malonic acid, succinic acid, maleic acid, and fumaric acid, or by reaction with an inorganic base such as sodium hydroxide, ammonium hydroxide, potassium hydroxide, and organic bases such as mono-, di-, trialkyl and aryl amines and substituted ethanolamines

[0146] Administration can be accomplished by single or multiple doses. The dose administered to a subject in the context of the present disclosure should be sufficient to induce a beneficial therapeutic response in a subject over time, or to inhibit or prevent RVF virus infection. The dose required will vary from subject to subject depending on the species, age, weight and general condition of the subject, the severity of the infection being treated, the particular immu-

nogenic composition being used and its mode of administration. An appropriate dose can be determined by one of ordinary skill in the art using only routine experimentation.

[0147] Provided herein are pharmaceutical compositions (also referred to as immunogenic compositions) which include a therapeutically effective amount of the recombinant RVF viruses alone or in combination with a pharmaceutically acceptable carrier. Pharmaceutically acceptable carriers include, but are not limited to, saline, buffered saline, dextrose, water, glycerol, ethanol, and combinations thereof. The carrier and composition can be sterile, and the formulation suits the mode of administration. The composition can also contain minor amounts of wetting or emulsifying agents, or pH buffering agents. The composition can be a liquid solution, suspension, emulsion, tablet, pill, capsule, sustained release formulation, or powder. The composition can be formulated as a suppository, with traditional binders and carriers such as triglycerides. Oral formulations can include standard carriers such as pharmaceutical grades of mannitol, lactose, starch, magnesium stearate, sodium saccharine, cellulose, and magnesium carbonate. Any of the common pharmaceutical carriers, such as sterile saline solution or sesame oil, can be used. The medium can also contain conventional pharmaceutical adjunct materials such as, for example, pharmaceutically acceptable salts to adjust the osmotic pressure, buffers, preservatives and the like. Other media that can be used with the compositions and methods provided herein are normal saline and sesame oil.

[0148] The recombinant RVF viruses described herein can be administered alone or in combination with other therapeutic agents to enhance antigenicity. For example, the recombinant viruses can be administered with an adjuvant, such as Freund incomplete adjuvant or Freund's complete adjuvant.

[0149] Optionally, one or more cytokines, such as IL-2, IL-6, IL-12, RANTES, GM-CSF, TNF- α , or IFN- γ , one or more growth factors, such as GM-CSF or G-CSF; one or more molecules such as OX-40L or 41 BBL, or combinations of these molecules, can be used as biological adjuvants (see, for example, Salgaller et al., 1998, *J. Surg. Oncol.* 68(2):122-38; Lotze et al., 2000, *Cancer J. Sci. Am.* 6(Suppl 1):S61-6; Cao et al., 1998, *Stem Cells* 16(Suppl 1):251-60; Kuiper et al., 2000, *Adv. Exp. Med. Biol.* 465:381-90). These molecules can be administered systemically (or locally) to the host.

[0150] A number of means for inducing cellular responses, both in vitro and in vivo, are known. Lipids have been identified as agents capable of assisting in priming CTL in vivo against various antigens. For example, as described in U.S. Pat. No. 5,662,907, palmitic acid residues can be attached to the alpha and epsilon amino groups of a lysine residue and then linked (for example, via one or more linking residues, such as glycine, glycine-glycine, serine, serine-serine, or the like) to an immunogenic peptide. The lipidated peptide can then be injected directly in a micellar form, incorporated in a liposome, or emulsified in an adjuvant. As another example, *E. coli* lipoproteins, such as tripalmitoyl-S-glycerylcysteinylseryl-serine can be used to prime tumor specific CTL when covalently attached to an appropriate peptide (see, Deres et al., *Nature* 342:561, 1989). Further, as the induction of neutralizing antibodies can also be primed with the same molecule conjugated to a peptide which displays an appropriate epitope, two compositions can be combined to elicit both humoral and cell-mediated responses where that is deemed desirable.

[0151] The following examples are provided to illustrate certain particular features and/or embodiments. These examples should not be construed to limit the disclosure to the particular features or embodiments described.

EXAMPLES

Example 1

Optimized Reverse Genetics System for Generation of Recombinant RVF Virus

[0152] This example describes the development of an improved reverse genetics system for generation of recombinant RVF viruses. The recombinant RVF viruses described herein are generated using an optimized reverse genetics system capable of rapidly generating wild-type and mutant viruses (Bird et al., *Virology* 362:10-15, 2001, herein incorporated by reference). The RVF virus reverse genetics system is a T7 RNA polymerase-driven plasmid-based genetic system that requires only three plasmids. Each plasmid individually expresses an anti-genomic copy of either the S, M or L segment. In this specific example, the S, M and L segments are derived from the genome of the virulent Egyptian RVF virus isolate ZH501. Nucleotide sequences of ZH501 S, M and L segments are provided herein as SEQ ID NOs: 1-3, respectively.

[0153] Rescue of recombinant viruses is accomplished by simultaneous transfection of the three anti-genomic sense plasmids into cells stably expressing T7 polymerase (for example, BSR-T7/5 cells). The genome segments of each plasmid are flanked by a T7 promoter, which enables generation of the primary RNA transcript, and the hepatitis delta virus ribozyme, which removes extraneous nucleotides from the 3' end of the primary transcriptional products. The T7 RNA polymerase generated transcripts are identical copies of the RVF-encoding plasmids, with the exception of an extra G nucleotide on the 5' end derived from the T7 promoter.

[0154] A previously reported method of producing recombinant RVF virus required the use of five expression plasmids (Gerrard et al., *Virology* 359:459-465, 2007). In addition to expression plasmids encoding RVF virus S, M and L segments, this system included plasmids encoding RVF virus N protein and RNA-dependent RNA polymerase (L protein). Northern blot and 3'RACE revealed that the S plasmid used in that system produced low levels of full length S-segment anti-genomic replication products. To produce a more efficient S segment plasmid, the full length S RNA from ZH501 was re-cloned into a version of the plasmid vector that was modified to remove the multiple cloning site. Within this backbone context, levels of N protein expression were significantly increased, allowing for subsequent rescue of recombinant virus without the need for a support plasmid encoding the N protein.

[0155] In addition, the optimized system described herein includes an M segment clone in which two non-synonymous mutations were corrected to match the wild-type sequence, allowing for exact in vivo comparisons between recombinant and wild-type viruses. These corrections were made in both the full-length M segment and the Δ NSm plasmid by site directed mutagenesis and restriction fragment exchange. These changes included removal of the XhoI restriction site.

After incorporating these changes into the reverse genetics system, virus rescue was accomplished by simultaneous transfection of only the three anti-genomic sense plasmids (either wild-type S, M and L segments, or deletion mutants thereof). The nucleotide sequences of exemplary wild-type S, M and L plasmids, and plasmids comprising deletions in the S and M segments, are provided herein as SEQ ID NOs: 11-15. Maps of each of these plasmids are shown in FIGS. 5-9.

[0156] After complete lysis of transfected cells, cell supernatants containing rescued viruses were clarified, diluted and virus passaged twice on Vero E6 cells. All recombinant

Construction of ΔNSs:GFP Deletion/Replacement Plasmid

[0160] To generate plasmids containing a complete deletion of the NSs ORF, replaced by the eGFP ORF (Townner et al., *Virology* 332(1):20-27, 2005, incorporate herein by reference), the eGFP ORF was amplified by PCR with forward and reverse primers containing the KpnI (GGTACC) and BglII (AGATCT) restriction sites, respectively. Utilizing the full-length RVF S-segment plasmid (pRVS; SEQ ID NO: 11), a strand specific PCR was conducted with primers designed to contain the KpnI and BglII restriction sites annealing immediately upstream to the NSs START codon, and immediately down-stream of the NSs STOP codon. Primer sequences are shown below in Table 1. Restriction site sequences are indicated by italics.

TABLE 1

Primer sequences for replacement of NSs with eGFP		
Primer	Sequence	SEQ ID NO:
RVS-35/KpnI	aaaaaaGGTACCGATATACTTGATAAGCACTAG	4
RVS + 827/BglII	aaaaaaAGATCTGATTAGAGGTTAAGGCTG	5
eGFP + KpnI	aaaaaaGGTACCATGGTGGAGCAAGGGCGAGGAG	6
eGFP-720/BglII	aaaaaaAGATCTTTACTTGTACAGCTCGTCCATG	7

viruses were found to grow similarly to wild-type ZH501 in Vero E6 cell culture. To confirm the exact molecular identity of all viruses used in this Example, complete genome sequence was obtained following previously described techniques (Bird et al., *J. Virol.* 81(6):2805-2816, 2007, herein incorporated by reference).

[0157] A total of 11 separate rescue attempts using a variety of plasmid concentrations ranging from 0.5 µg to 4 µg resulted in 100% recombinant virus rescue efficiency using only the three expression plasmids.

Example 2

Generation and Characterization of Recombinant RVF (rRVF) Viruses

[0158] This example describes the generation of recombinant RVF viruses comprising deletions in the ORF of NSs and/or NSm, which play a role in viral virulence. This example further describes a recombinant RVF virus comprising a reporter gene.

RVF Virus and Biosafety

[0159] All work with infectious RVF virus (wild-type or recombinant) was conducted within the Centers for Disease Control (CDC) bio-safety level 4 (BSL-4) laboratory. Low passage (FRhL+2, Vero E6+2) working stocks of wild-type strain ZH501, isolated originally from a fatal Egyptian human case in 1977, were used as challenge virus, and were prepared by passage on Vero E6 cell monolayers. The complete genome sequence of the S, M and L segments of the wild-type RVF virus strain ZH501 has been deposited under Genbank Accession Nos. DQ380149 (SEQ ID NO: 1), DQ380200 (SEQ ID NO: 2) and DQ375406 (SEQ ID NO: 3), respectively.

[0161] The resulting PCR fragment contained a complete deletion of the NSs ORF flanked by KpnI and BglII restriction sites. The eGFP and RVF-S plasmid amplicons were gel purified and ligated following standard molecular biology techniques. The resulting plasmid (pRVS-GFPΔNSs; SEQ ID NO: 14) contained a complete in-frame replacement of the NSs ORF by eGFP.

Construction of Plasmids Containing a NSs-eGFP Fusion Protein

[0162] In a second set of plasmid constructions, rRVF viruses were generated which contained the complete full-length genome and an insertion of the eGFP ORF. Two constructions were made containing in-frame fusions of the c-terminus of NSs with the eGFP ORF, separated by amino acid linker moieties of varying lengths. In the first construction, the full-length RVF-S segment plasmid backbone was modified using site directed mutagenesis PCR (QuickChange, Stratagene) with overlapping primers (shown in Table 2 below) containing a deletion/replacement of the NSs STOP codon with a linker section containing nucleotides encoding three alanine residues and the KpnI restriction site. Following this, both this amplicon and the pRVS-GFPΔNSs plasmid were digested with KpnI. The resulting restriction enzyme fragments were gel purified and ligated to generate the final construction pRVFS-NSs(A₃)eGFP. A second NSs-eGFP fusion peptide construction was created to increase the length of the eGFP linker moiety and eliminate the KpnI restriction site contained in the previous fusion peptide construction. To accomplish this, site directed mutagenesis (QuickChange, Stratagene) was employed starting with the pRVFS-NSs(A₃)eGFP backbone and overlapping primers containing the remote cutter restriction enzyme BsmBI and nucleotides encoding ten alanine residues (shown in Table 2 below).

TABLE 2

Primer sequences for construction of NSs-eGFP fusion proteins		
Primer	Sequence	SEQ ID NO:
RVS-829rev/KpnI	aaaaaaGGTACCTGCTGCTGCATCAACCTCAACAAA TCCATC	8
RVS + 827/BglII	aaaaaaAGATCTGATTAGAGGTTAAGGCTG	5
NSsGFP + 10Ala/Fwd	aaaaaaCGTCTCaGCAGCAGCAGCAGCAATGGTGAG CAAGGGCGAGGAG	9
NSsGFP + 10Ala/Rev	aaaaaaCGTCTCaCTGCTGCTGCTGCTGCATCAA CCTCAACAAATCCATC	10

[0163] Following PCR amplification, restriction enzyme digestion and re-ligation, the resulting construction contained a perfect in-frame fusion of the C-terminus of NSs and eGFP ORF in the context of the complete full-length RVF S segment genome, preserving the spacing and nucleotide sequence of both the NP and NSs transcription termination signals.

Generation and In Vitro Testing of Recombinant RVF Viruses

[0164] In all cases, rescue of recombinant viruses was accomplished using cDNA plasmids encoding the virulent RVF virus ZH501 strain. The basic design and construction of the full-length plasmids containing inserts of the complete S segment (pRVS; SEQ ID NO: 11), M segment (pRVM; SEQ ID NO: 12), and L segment (pRVL; SEQ ID NO: 13), and a plasmid containing a deletion of the NSm gene (pRVM-ΔNSm; SEQ ID NO: 15), have been described previously (Bird et al., *Virology* 362:10-15, 2007; Gerrard et al., *Virology* 359:459-465, 2007, each of which is herein incorporated by reference). Maps of the full-length S, M and L plasmids, and plasmids containing NSs and NSm deletions, are shown in FIGS. 5-9.

[0165] Anti-genomic sense plasmids representing the three genomic segments were transfected in 1 μg quantities with LT-1 (Mirus) at a ratio of 6:1 and transferred onto sub-confluent (approximately 60-70% confluent) monolayers of BSR-T7/5 cells stably expressing T7 polymerase (Buchholz et al., *J. Virol.* 73(1):251-259, 1999, herein incorporated by reference). Four or five days post transfection, the cell supernatant was clarified by low speed centrifugation and passed twice on confluent monolayers of Vero E6 cells. After passage and prior to use in subsequent experiments, the complete genome sequence of each rescued recombinant virus was confirmed by previously described techniques (Bird et al., *J. Virol.* 81:2805-2816, 2007, herein incorporated by reference).

Infected Live Cell (Direct) or Fixed Cell (Indirect) Fluorescent Antibody Detection of RVF NSs and eGFP Proteins

[0166] Vero E6 cells were seeded on glass coverslips and infected at a multiplicity of infection (MOI) of approximately 1.0 with either rZH501-ΔNSs:GFP, rZH501-ΔNSs:GFP-ΔNSm, rZH501-NSs(Ala)₃GFP or rZH501-NSs(Ala)₁₀GFP. At 24 hours post infection, cells were directly visualized by inverted ultraviolet microscopy (live cell) or were fixed in 10% formalin overnight. Following fixation, infected cells were gamma-irradiated (5.0×10^6 RAD) to inactivate any residual virus activity. After inactivation, cells were perme-

abilized (Triton X-100 0.01%) and incubated with monoclonal antibodies specific for either RVF NSs or eGFP protein following standard techniques.

Results

[0167] Rescue of all recombinant viruses used in this study was accomplished by transfection of three anti-genomic sense plasmids, each representing one of the three virus

[0168] RNA segments, without the requirement of supporting expression plasmids encoding virus structural proteins. Multiple rRVF viruses were generated containing an insertion of the reporter molecule eGFP into the virus S segment (FIG. 1A). Among these, two rRVF viruses were rescued containing an in-frame fusion of the C-terminus of the NSs protein with the N-terminus of eGFP, separated by a peptide linker of either three or ten alanine residues (rZH501-NSs(Ala)₃GFP and rZH501-NSs(Ala)₁₀GFP). Two rRVF viruses were created containing deletions of either the NSs alone or NSs/NSm genes in combination (rZH501-ΔNSs:GFP and rZH501-ΔNSs:GFP-ΔNSm). In both of these viruses, the NSs gene was replaced by the reporter molecule eGFP, preserving the native S segment ambisense RNA orientation. Both rRVF viruses were rescued upon the first attempt and grew to high titers routinely exceeding 1.0×10^6 PFU/mL in Vero E6 cell culture resulting in complete monolayer lysis.

[0169] Following passage on Vero E6 cells, cytoplasmic GFP signal was first observed approximately 10-12 hours post infection and appeared to spread rapidly throughout the cell monolayer prior to the first signs of extensive CPE and plaque formation (FIG. 1B).

[0170] Recombinant virus containing NSs-GFP fusion peptides (rZH501-NSs(Ala)₃GFP and rZH501-NSs(Ala)₁₀GFP) were also rescued on the first attempt but were found to grow to slightly lower titers (about 5.0×10^5 PFU/mL). NSs-GFP fusion protein was first localized in the cytoplasm of infected cells followed by perinuclear accumulation and eventual intranuclear migration followed by the formation of filamentous structures by 12-18 hours post-infection (FIGS. 1B and 1C) Stability of the eGFP reporter gene in all recombinant viruses reported here was followed for 15 serial passages (1:100 dilution between each passage) in Vero E6 cells during which time no decrease in the stability of eGFP expression was observed, with all infected cells expressing robust amounts of eGFP protein similar to that seen in early passages.

Example 3

In vivo Immunogenicity and Safety of rRVF Viruses

[0171] This example describes a study to determine the immunogenicity and safety of the recombinant RVF viruses described in the examples above.

Animal Immunization and Infection

[0172] A total of 66 female Wistar-furth (WF/NSd) (Harlan) rats 6-8 weeks of age (approximately 160 g) were used in this study. The animals were housed in micro-isolator pans and provided food and water ad libitum. All pans were kept in HEPA filtration racks following standard barrier care techniques within the BSL-4 laboratory for the duration of the experiment. All animals were inoculated subcutaneously (SQ) in the right hind flank with an inoculum (vaccine or challenge virus) prepared in 200 μ L of sterile physiologic saline. A total of eight animals that were administered sterile saline only served as sham inoculated animal controls (three in the pilot study, five in the vaccination/challenge study). All animals were examined daily post inoculation for signs of clinical illness, weight loss and respiratory distress. Animals found either in distress or moribund were immediately anesthetized using isoflurane and then euthanized using Beuthanasia solution (Schering-Plough) following standard techniques.

Pilot in vivo Immunogenicity and Safety Study

[0173] A total of 18 rats were administered 1.0×10^3 PFU SQ of either the rZH501- Δ NSs:GFP (nine animals) or rZH501- Δ NSs:GFP- Δ NSm (nine animals) with three animals serving as sham inoculated controls. Following vaccination, a small (approximately 25 μ L) sample of whole blood was obtained via the tail vein on days 1-4 to detect vaccine virus induced viremia. This whole blood sample was placed directly into 2 \times non-cellular lysis buffer (Applied Biosystems) for decontamination and transfer to a BSL-2 laboratory following standard protocols (Towner et al., *J. Infect. Dis.* S2002-S212, 2007, incorporated herein by reference) for subsequent RNA extraction and RVF specific q-RT-PCR. At day 21 post-vaccination, all animals were anesthetized using isoflurane, serum samples were collected for determination of total anti-RVF IgG titers, and the animals were euthanized using Beuthanasia solution (Schering-Plough).

Anti-RVF Virus Total IgG ELISA

[0174] Determination of anti-RVF IgG titers from vaccinated and control rats was performed essentially as described by Madani et al. (*Clin. Infect. Dis.* 37:1084-1092, 2003) with the following modifications necessary for rat specimens. Standard 96-well microtiter plates were coated overnight at 4 $^{\circ}$ C. with 100 μ L of gamma-irradiated RVF infected Vero E6 cell lysate diluted 1:2000 in (0.01M PBS, pH 7.2), or similarly diluted gamma-irradiated uninfected Vero E6 cell lysate, to serve as adsorption controls. Plates were washed 3 \times (PBS 0.01% Tween-20) and 100 μ L duplicate samples of rat sera were diluted 1:100-1:6400 in 4-fold dilutions in skim milk serum diluent and adsorbed for 1 hour at 37 $^{\circ}$ C. Plates were washed 3 \times (PBS 0.01% Tween-20) and 100 μ L of goat anti-Rat IgG HRP conjugate antibody (KPL, Gaithersburg, Md.) diluted 1:2500 was adsorbed for 1 hour at 37 $^{\circ}$ C. After a 3 \times wash, 100 μ L of ABTS substrate (KPL) was added and incubated for 30 minutes at 37 $^{\circ}$ C.

[0175] Plates were read at 410 nm with an absorbance correction of 490 nm for plate imperfections. The absorbance of the 1:100, 1:400, 1:1600, and 1:6400 dilutions were added and constituted a SUM_{OD} value for each specimen. The background adsorption of each animal serum to negative control Vero E6 cells was subtracted from the calculated SUM_{OD} value obtained from antigen positive Vero E6 cells and was recorded as an adjusted-SUM_{OD} value for each animal. Final end point dilution titers were determined as the reciprocal of the final serum dilution yielding an adjusted-SUM_{OD} of >0.20. A cut-off value for determining positive versus negative samples was established as the mean sample adjusted-SUM_{OD}+3 standard deviations obtained from the five sham-inoculated control animals.

Statistical Analyses

[0176] For all calculations, the analysis program XLSTAT (AddinSoft, USA) was utilized. Kaplan-Meier analyses were completed with log-rank and Wilcoxon tests of significance with an α -level setting of 0.05. Analyses of SUM_{OD} and viremia were completed utilizing a one-way ANOVA and a post-hoc Tukey's HSD test of significance with an α -level setting of 0.05.

Results

[0177] To gain a primary assessment of the relative in vivo characteristics of the rZH501- Δ NSs:GFP and rZH501- Δ NSs:GFP- Δ NSm recombinant viruses, groups of nine rats were inoculated with each rRVF virus at a dose of 1.0×10^3 PFU (FIG. 4A). Animals were monitored daily for signs of clinical illness and weight loss. At no time post-vaccination did any animal show signs of clinical illness, and all experienced average daily weight changes equivalent to sham-inoculated controls of approximately 0-5 g. All vaccinated rats were bled at days 1-4 post-inoculation to determine the titer of vaccine virus induced viremia. Using a highly sensitive q-RT-PCR assay, no animal at any time point analyzed post-vaccination developed a detectable viremia.

[0178] All animals were euthanized at day 21 post-vaccination and anti-RVF total IgG antibody titers were evaluated (FIG. 2). Testing revealed that the SUM_{OD} (mean \pm SEM) for all animals vaccinated with the rZH501- Δ NSs:GFP virus was 2.14 \pm 0.12, which corresponded to end point titers of 1:1600 in 66% (6/9) of animals, with the remaining 33% (3/9) having titers equal to 1:400. Among animals receiving the rZH501- Δ NSs:GFP- Δ NSm virus, SUM_{OD} (mean \pm SEM) was 1.24 \pm 0.06, with 89% (8/9) developing end-point dilution titers equal to 1:400. As expected, all sham-inoculated animals (N=3) were negative for detectable levels of anti-RVF total IgG antibody SUM_{OD} -0.08 \pm 0.06. All vaccinated animals in the rZH501- Δ NSs:GFP and rZH501- Δ NSs:GFP- Δ NSm virus groups developed statistically higher mean anti-RVF total IgG SUM_{OD} values compared with non-vaccinated controls (p-value <0.001 and p-value=0.003, respectively). Animals in the rZH501- Δ NSs:GFP virus group developed significantly higher mean SUM_{OD} values than animals given the rZH501- Δ NSs:GFP- Δ NSm vaccine (p-value=0.004). Plaque reduction neutralization titers (PRNT₅₀) testing was completed on a subset (four animals) chosen randomly from each vaccine virus group with two sham-inoculated animals serving as controls. The results showed mean PRNT₅₀ titers of 1:1480

(rZH501-ΔNSs:GFP) and 1:280 (rZH501-ΔNSs:GFP-ΔNSm), with sham-inoculated control animal titer of <1:10.

Example 4

Follow-Up Vaccination and Virulent Virus Challenge Study

[0179] This example describes the efficacy of recombinant RVF viruses comprising a deletion in the NSs ORF, or comprising a deletion in both the NSs and NSm ORFs, following challenge with wild-type virus.

Vaccination and Virus Challenge

[0180] A total of 20 rats were vaccinated in two dosage groups of ten animals each with either 1.0×10^3 or 1.0×10^4 PFU SQ of the rZH501-ΔNSs:GFP virus as described above. An additional 20 rats were inoculated in two dosage groups of ten animals each with either 1.0×10^3 or 1.0×10^4 PFU SQ of the rZH501-ΔNSs:GFP-ΔNSm virus as described above. A total of five animals served as sham-inoculated controls. On days 2, 4 and 7 post-vaccination, a small blood sample (approximately 25 μL) was collected from the tail vein and added directly to 2× non-cellular lysis buffer (Applied Biosystems) as described above for subsequent RNA extraction and q-RT-PCR.

[0181] At day 26 post-vaccination, all animals were briefly anesthetized using isoflurane vapor (3.0-3.5% atmosphere) (RC² Rodent Anesthesia system, VetEquip) and a 500 μL sample of whole blood was obtained. Serum was collected and stored at -70° C. for later determination of total anti-RVF IgG titers, PRNT₅₀ and anti-NSs/anti-NP specific antibody. All serum samples were surface decontaminated and inactivated by gamma-irradiation (5.0×10^6 RAD) following standard BSL-4 safety protocols prior to use in a BSL-2 laboratory.

[0182] At day 28 post-vaccination, all rats (vaccinated and sham controls) were challenged with 1.0×10^3 PFU SQ of virulent wild-type RVF virus strain ZH501, previously shown to result in lethal disease in Wistar-furth rats (Anderson et al., *Microb. Pathog.* 5:241-250, 1988; Anderson et al., *Am. J. Trop. Med. Hyg.* 44(5):475-80, 1991; Bird et al., *Virology* 362:10-15, 2007, each of which is herein incorporated by reference). On days 2, 3, 4 and 7 following challenge, a small blood sample was collected from the tail vein for subsequent RNA extraction and RVF specific q-RT-PCR to assess the level of viremia. At day 42 post-challenge, all animals surviving wild-type virus infection were bled via cardiac puncture under general anesthesia (isoflurane vapor, RC² Rodent Anesthesia system, VetEquip) followed by euthanasia (Beuthanasia solution, Schering-Plough).

Anti-RVF Virus Plaque Reduction and Neutralization Testing (PRNT₅₀)

[0183] The stock of RVF virus was diluted to 50 PFU in 300 μL of DMEM (1% Penicillin/Streptomycin) without FBS. Separately, aliquots of serum from vaccinated rats or from sham-inoculated controls corresponding to 21 days post-vaccination (pilot study) or 26 days (challenge study) were heat inactivated for 30 min at 56° C. After inactivation, serum dilutions of 1:10, 1:40, 1:160, 1:640, 1:2560 and 1:10240 were made in DMEM (1% Pen/Strep). Diluted rat serum (300 μL) was mixed with an equal volume of diluted virus and incubated overnight at 4° C. The following day, each mixture

of serum+RVF virus was used to infect confluent monolayers of Vero E6 cells in 12-well plates. After a 1 hr adsorption, the mixture was removed and a 1-2 ml nutrient agarose overlay (MEM 1×, 2% FBS, 1% Pen/Strep, 1% SeaPlaque agar) was added to the monolayers. After a five-day incubation, the cells were fixed with 10% formalin overnight. Following fixation, the agarose overlay was removed and the plates were surface decontaminated and gamma-irradiated (2.0×10^6 RAD) following standard BSL-4 safety procedures. After inactivation, the cell monolayer was stained with 1% crystal violet in PBS and plaques were enumerated. The calculated PRNT₅₀ titers correspond to the reciprocal titer of the last dilution resulting in a 50% reduction in the number of plaques when compared to controls.

RVF Virus Specific q-RT-PCR

[0184] Whole rat blood was assayed for the presence and quantity of RVF specific virus RNA as described previously (Bird et al., *J. Clin. Microbiol.* 45(11):3506-13, 2007, incorporated herein by reference). Quantification of total serum RVF RNA was calculated directly via interpolation from a standard curve generated from serial dilutions of stock RVF virus strain ZH501 of a known titer in whole rat blood extracted and processed in an identical manner with each experimental replicate q-RT-PCR run. Briefly, 25 μL of whole rat blood (either from vaccinated/challenged animals or stock virus serial dilutions) was added to 300 μL 2× non-cellular lysis buffer (Applied Biosystems) and total RNA was extracted (ABI 6100 nucleic acid workstation, Applied Biosystems.) After extraction, cDNA was generated by random hexamer priming (High Capacity cDNA kit, Applied Biosystems) followed by RVF specific q-PCR (Universal q-PCR master mix, Applied Biosystems). Results are reported as RVF PFU equivalents/mL of rat blood.

Results

Immunization Phase

[0185] Following the promising findings of robust immunogenicity and in vivo attenuation in the initial pilot study, a larger study was undertaken utilizing multiple doses of each recombinant RVF virus followed by virulent virus challenge. Groups of ten animals each were inoculated with either rZH501-ΔNSs:GFP or rZH501-ΔNSs:GFP-ΔNSm viruses at dosages of 1.0×10^3 or 1.0×10^4 PFU (FIG. 4B). Whole blood samples were assayed at day 2, 4, and 7 post-vaccination for the presence of detectable viremia. As was observed in the pilot study, at no time did any animal develop detectable vaccine viremia. Additionally, no clinical illness was observed among any vaccinated animals, and all experienced weight gains of approximately 1-5 g per day, similar to sham-inoculated controls.

[0186] At day 26 post-vaccination, serum samples were obtained to determine the level of total anti-RVF IgG, PRNT₅₀, and anti-NP/anti-NSs protein specific antibody production prior to subsequent challenge on day 28. All animals regardless of recombinant virus type or dose, developed high-titered total anti-RVF IgG antibody (FIG. 2). Among rats receiving the rZH501-ΔNSs:GFP virus, the mean SUM_{OD}±SEM was 4.10 ± 0.12 (1.0×10^3 dose group) and 4.79 ± 0.11 (1.0×10^4 dose group), which corresponded to 85% (17/20) of animals developing anti-RVF IgG end-point titers of 1:6400. The remaining three animals in the rZH501-ΔNSs:GFP virus group developed end-point titers equal to 1:1600. In animals receiving the rZH501-ΔNSs:GFP-ΔNSm virus,

the mean $SUM_{OD} \pm SEM$ was 3.94 ± 0.12 (1.0×10^3 dose group) and 4.54 ± 0.11 (1.0×10^4 dose group), which corresponded to 75% (15/20) developing anti-RVF IgG end-point titers of 1:6400, with the remaining 25% (5/20) attaining end-point titers of 1:1600. As was observed in the pilot study, all animals vaccinated with either rZH501- Δ NSs:GFP or rZH501- Δ NSs:GFP- Δ NSm viruses, regardless of dose, developed statistically significant higher mean SUM_{OD} values than sham-inoculated control animals (p-values all <0.05).

[0187] Similarly, PRNT₅₀ titers were found to be elevated above sham-inoculated controls among animals inoculated with the rZH501- Δ NSs:GFP virus with mean titers of 1:640 and 1:7040 in the 1.0×10^3 and 1.0×10^4 dose groups, respectively. Mean PRNT₅₀ titers among animals vaccinated with the rZH501- Δ NSs:GFP- Δ NSm were found to be similar with mean titers of 1:1120 and 1:640 in the 1.0×10^3 and 1.0×10^4 dose groups, respectively.

Challenge Phase

[0188] On day 28 post-vaccination, all rats were challenged with a known lethal dose (1.0×10^3 PFU) of virulent wild-type strain ZH501. All animals were monitored daily for signs of clinical illness and weight loss/gain for 42 days post-challenge. At no time post challenge (days 1-42) did any rat that received prior vaccination with either the rZH501- Δ NSs:GFP or rZH501- Δ NSs:GFP- Δ NSm viruses develop clinically detectable illness (ruffled fur, hunched posture or lethargy). At approximately day 2 post-challenge, a majority of animals suffered slight 1-5 g reductions in total body weight that was regained by day 5 post-challenge.

[0189] Rat whole blood was obtained on days 2, 3, 4 and 7 post-challenge and assayed for the presence of RVF virus RNA by a highly sensitive q-RT-PCR assay (Bird et al., *J. Clin. Microbiol.* 45(11):3506-13, 2007, incorporated herein by reference). Following challenge, low-level viremia was detected in a total of 3/40 vaccinated animals. Among the 20 animals that were vaccinated with rZH501- Δ NSs:GFP- Δ NSm, two animals developed a peak post-challenge viremia on day 3 of 1.1×10^2 and 7.0×10^1 PFU equivalents/mL of whole blood, respectively (FIG. 4B). In the 20 animals vaccinated with rZH501- Δ NSs:GFP, one animal was detected with a peak post-challenge viremia on day 4 of 1.5×10^1 PFU equivalents/mL of whole blood. In all three animals, the detectable viremia was transient and was resolved by day 7 post-challenge. No other vaccinated animals (37 total) had a detectable RVF virus viremia at any time point assayed post challenge. In sharp contrast, the five sham inoculated animals all suffered severe to lethal clinical illness with 3/5 animals succumbing to infection by day 3 post-challenge with peak viremia titers of 1.9×10^7 , 2.7×10^7 , and 3.1×10^7 PFU equivalents/mL. Two sham-inoculated animals did not succumb to infection but did develop severe clinical illness (ruffled fur, hunched back, lethargy) with a peak viremia post-challenge of 2.7×10^4 and 5.8×10^3 PFU equivalents/mL whole blood. As anticipated, both candidate vaccines (rZH501- Δ NSs:GFP or rZH501- Δ NSs:GFP- Δ NSm) significantly reduced post-challenge viremia (p-value <0.0001) regardless of dose. Additionally, Kaplan-Meier survivor analysis (log-rank and Wilcoxon tests) of survival post-challenge revealed a significant protective efficacy effect with both rRVF viruses regardless of dose when compared with sham-inoculated controls (p-value <0.001).

Example 5

Differentiation of Wild Type-Infected and Vaccinated Animals

[0190] This example describes how rRVF viruses lacking NSs can be used to differentiate animals that have been

infected with the rRVF virus and wild-type virus using methods to detect antibodies specific for NSs and NP.

Anti-RVF NSs and NP Differential Indirect Fluorescent Antibody (IFA) Assays

[0191] Plasmid constructions expressing only the RVF nucleoprotein (NP) or non-structural S(NSs) proteins were generated following techniques described previously (Niwa et al., *Gene* 108:193-199, incorporated herein by reference). Briefly, oligonucleotide primers were designed to anneal within the NP or NSs ORF with the addition of a SacI and BglII restriction site for cloning into a polymerase II-based expression plasmid, pCAGGS. The resulting PCR amplicons were agarose gel purified, digested with SacI and BglII, and ligated between the corresponding restriction sites of the pCAGGS vector. Prior to use, the resulting clones, pC-NP and pC-NSs, were sequenced using standard techniques.

[0192] Following confirmation of the molecular sequence, each plasmid was transfected separately on Vero E6 cells grown on glass coverslips in 1 μ g quantities at a 6:1 ratio with lipofectant solution (LT-1, Mirus). Following a 48-hour incubation, transfected cells expressing either the NP or NSs protein were fixed with 10% formalin. To assess the presence or absence of anti-NP or anti-NSs antibody, serum samples from all vaccinated and the two surviving sham-inoculated control animals were adsorbed separately for 1 hour on cells transfected with either NP or NSs. The presence or absence of anti-NP or anti-NSs adsorbed rat antibody was visualized by secondary adsorption of an AlexaFluor 594 nm (Giorgi et al., *Virology* 180:738-753, 1991) anti-rat total IgG (Molecular Probes/Invitrogen) antibody. Intra-nuclear localization of NSs protein and rat antibody was confirmed by DAPI counterstaining. To confirm the presence of anti-RVF virus NP and NSs antibodies among naturally infected animals, archived serum samples collected from goats in Saudi Arabia during the outbreak in 2000 were tested essentially as described above with anti-goat specific total IgG (AlexaFluor 488 nm, FITC, Molecular Probes/Invitrogen).

Results

[0193] Serum obtained from all vaccinated and sham vaccinated animals at day 26 post-vaccination was tested for the presence of anti-NP and anti-NSs specific antibodies utilizing Vero E6 cells transfected with plasmids expressing either NP or NSs proteins. As a positive control, the serum obtained from the two sham vaccinated animals that survived infection, and six additional control rat serum samples (taken from animals inoculated with a sub-lethal dose of RVF virus for validation purposes) were utilized. Control animals surviving infection developed high levels of both anti-NP and anti-NSs antibody with strong immunostaining of both in vitro expressed cytoplasmic NP and filamentous intra-nuclear accumulations of NSs protein (FIG. 3A). No anti-NS or anti-NP antibodies were detected in serum from control rats (FIG. 3C). In concordance with the anti-RVF total IgG data, all vaccinated animals demonstrated high anti-NP specific antibody levels, and as anticipated, no vaccinated animal, regardless of vaccine virus or dose, developed detectable anti-NSs specific antibody (FIG. 3B). As a further step toward the demonstration of the ability to differentiate naturally infected animals from those vaccinated with the vaccines provided herein, naturally infected livestock were also shown to pro-

duce anti-N Ss antibodies similar to those observed with the wild type virus infected rats (FIG. 3D).

Example 6

Safety and Efficacy of Recombinant RVF Virus Vaccines in Pregnant Ewes

[0194] The safety and efficacy of recombinant RVF virus vaccines can be evaluated in pregnant ewes according to procedures well known in the art (Morrill et al., *Am. J. Vet. Res.* 48(7):1042-1047, 1987; Baskerville et al., *Res. Vet. Sci.* 52:307-311, 1992, each of which is herein incorporated by reference). By way of example, rRVF viruses are evaluated in healthy, RVF virus sero-negative, crossbred ewes at approximately 12 weeks gestation. The ewes are housed in a biological containment facility and fed a daily ration of alfalfa hay and a commercial grain mix ration, with grass hay and water provided ad libitum. Pregnant ewes are administered recombinant RVF virus subcutaneously or intramuscularly at various doses. Mock-infected ewes serve as controls. Ewes are monitored daily for health. At various timepoints post-inoculation, blood, serum or other body fluid samples can be taken to assay RVF virus-induced viremia or anti-RVF virus antibody production as described above, or other desired biological endpoints. Lambs born to inoculated ewes are allowed to remain with their dams and suckle.

[0195] To test efficacy of the recombinant RVF viruses as vaccines, inoculated and sham-inoculated ewes are administered wild-type RVF virus at various doses. Lambs can also be challenged with live virus to determine whether maternal antibodies against recombinant RVF virus provide protection against natural infection. Animals are observed daily for signs of clinical illness, weight loss and respiratory distress. Animals that are in distress or moribund are immediately anesthetized and then euthanized. As described above, at various timepoints following inoculation, small blood samples can be taken to test for the presence of viral RNA. Serum samples can be collected to determine anti-RVF virus antibody titers.

Example 7

Safety and Efficacy of Recombinant RVF Virus Vaccines in a Rhesus Macaque Model for Human Disease

[0196] The safety and efficacy of recombinant RVF virus vaccines can be evaluated in non-human primates, such as rhesus macaques, according to procedures well known in the art (Morrill and Peters, *Vaccine* 21:2994-3002, 2003). By way of example, rRVF viruses are evaluated in adult (5-10 kg) and/or juvenile (1-3 kg) captive-bred rhesus monkeys (*Macaca mulatta*). Animals are housed in individual cages in a biosafety level 3 (BSL-3) biological containment facility maintained at constant room temperature with a 12 hour light/dark photoperiod. Inoculations are given, and rectal temperatures and venous blood samples are taken while the animals are under ketamine-xylazine anesthesia.

[0197] Sero-negative monkeys are inoculated intravenously or intramuscularly with various doses of recombinant RVF viruses. Mock-inoculated animals serve as controls. The animals are monitored daily for clinical signs of illness, including weakness, paralysis or any alteration of physical condition. At various timepoints post-inoculation, blood, serum or other body fluid samples can be taken to assay RVF virus-induced viremia, anti-RVF virus antibody production,

or other desired biological endpoints (for example, white blood cell count, red blood cell count, hematocrit, platelet count, AST and ALT). Moribund monkeys are euthanized and necropsied.

[0198] To test efficacy of the recombinant RVF viruses as vaccines, inoculated and sham-inoculated monkeys are administered wild-type RVF virus at various doses. Animals are observed daily for signs of clinical illness, weight loss and respiratory distress. Animals that are in distress or moribund are immediately anesthetized and then euthanized. As described above, at various timepoints following inoculation, small blood samples can be taken to test for the presence of viral RNA. Serum samples can be collected to determine anti-RVF virus antibody titers.

Example 8

Vaccination of Human Subjects with Recombinant RVF Virus

[0199] The safety and efficacy of recombinant RVF virus vaccines can be evaluated in human volunteers according to procedures well known in the art (Pittman et al., *Vaccine* 18:181-189, 2000, herein incorporated by reference). Typically, human volunteers are selected from those having occupations that put them at risk of infection with RVF virus, such as veterinarians in endemic areas. All volunteers are screened to ensure they are in good health. Informed consent is obtained from each volunteer prior to vaccination.

[0200] In this example, human volunteers are injected with candidate rRVF vaccine subcutaneously at an appropriate dose. The appropriate dose is the dose approved by the FDA, and can be determined from suitable animal studies conducted prior to human vaccination trials. Other routes of administration are possible, including intramuscular and intravenous. The vaccine can be administered as a single dose, or given in multiple doses, such as two, three or four doses. When administered in multiple doses, the booster doses can be administered at various time intervals, such as months to years. Serum samples can be obtained to determine neutralizing antibody titers and identify responder and non-responders to the vaccine.

[0201] Vaccinated volunteers are encouraged to return and report local or systemic reactions. Local reactions are assessed by grading pain and tenderness at the site of inoculation and/or axillary lymph nodes and measuring erythema and induration at the site. Systemic reaction parameters include fever, chills, headache, malaise, myalgia, arthralgia, sore throat, gastric upset, dizziness, photophobia and skin rash. Additional laboratory samples, including complete blood cell count, chemistry profile, β -HCG (in females), urinalysis, blood samples for viremia titrations, and oropharyngeal washing for viral isolated culture can be obtained. Using serum samples obtained from vaccinated individuals, plaque-reduction neutralization tests can be performed to determine how robust the immune response was to the particular rRVF virus. Vaccinated volunteers are also screened for the development of RVF virus infection.

[0202] This disclosure provides recombinant RVF viruses comprising deletions in virus virulence genes. The disclosure further provides methods of immunizing subjects at risk of infection with RVF virus with the recombinant viruses. It will be apparent that the precise details of the methods described may be varied or modified without departing from the spirit of

the described disclosure. We claim all such modifications and variations that fall within the scope and spirit of the claims below.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 15

<210> SEQ ID NO 1

<211> LENGTH: 1690

<212> TYPE: DNA

<213> ORGANISM: Rift Valley fever virus

<400> SEQUENCE: 1

```

acacaaagac ccctagtgac ttatcaagta tatcatggat tactttcctg tgatatctgt    60
tgatttcag agtggtcgac gtgttggtgc agtggagtac tttagaggag atggctctcc    120
caggatacct tattctatgg ttgggcccctg ttgtgtcttt ctcatgcacc atcgtcctag    180
tcacgaggtt cgcttgcgat tctctgattt ctacaatgac ggagaattcc cataccgagt    240
cggacttggg gactttgcat caaacgttgc acctccacca gcgaagcctt ttcagagact    300
tattgatcta ataggccata tgactcttag tgatttcaca aggttcccca atctgaaaga    360
agccatatcc tggcctcttg gagaaccctc actggcttcc tttgacctaa gctctactag    420
agtgcatagg aatgatgaca ttagaaggga tcagattgcc actctagcaa tgaggagtgt    480
caagatcacc aatgatctag aggactcctt tgttgctta cacaggatga tagcgactga    540
ggccatcctc agagggattg acctgtgact gttgccaggc tttgatctca tgtatgaggt    600
tgctcacgta cagtgcgttc ggcttctgca agcagcaaaa gaggacattt ctaatgctgt    660
agttccaaac tcagccctca ttgttcttat ggaggagagc ctgatgctgc gctcatcact    720
tcccagcatg atggggagaa acaactggat tccagttatt cctccaatcc cagatgttga    780
gatggaatca gaggaggaga gtgatgatga tggatttgtt gaggttgatt agaggttaag    840
gctgccccac cccccacccc ccaatccoga ccgtaacccc aactcccctt ccccccaacc    900
ccctgggcag ccacttaggc tgctgtcttg taagcctgag cggctgccat gacagcagct    960
gacggcttcc cattagaatc cacaagtoca aaggctttca agaattctct cctcttctca   1020
tggettataa agttgctatt cactgctgca ttcattggct gcgtgaaagt tgcagcaacc   1080
tcctcttttg ttctacctcg gaggtttggg ttgatgacct gggagaactg cagcagatac   1140
agagagtgag catctaatat tgcccttaga tagtctcctg gtagagaagg atccaccatg   1200
ccagcaaagc tggggtgcat catatgcctt gggtatgcag gggataggcc gtccatggtg   1260
gtcccagtgga caggaagcca ctactcaag acgaccaaaag cctggcatgt ccagccagcc   1320
agggcggcag caactcgtga tagagtcaac tcatcccggg aaggattccc ttccttttagc   1380
ttatacttgt tgatgagagc ctccacagtt gctttgcctt ctctcagcat tttcatcacc   1440
atcctcctgg gcttgttgcg acgagttaga gccagaacaa tcattttctt ggcacacctc   1500
tcccagtcag cccccacata ctgctttaag agttcgataa ctctacgggc atcaaacctc   1560
tgataagcaa actctcggac ccaactgttca atctcattgc ggteccactgc ttgagcagca   1620
aactggatcg caagctcttg atagtgttcc attattgtaa tagtgtttgt atctctaggg   1680
agcttttgtg                                     1690

```

-continued

```

<210> SEQ ID NO 2
<211> LENGTH: 3885
<212> TYPE: DNA
<213> ORGANISM: Rift Valley fever virus

<400> SEQUENCE: 2
acacaaagac ggtgcattaa atgtatgttt tattaacaat tctaatctcg gttctgggtg    60
gtgaagcggg tattagagtg tctctaagct ccacaagaga agagacctgc tttggtgact    120
ccactaacc agagatgatt gaaggagctt gggattcact cagagaggag gagatgccgg    180
aggagctctc ctgttctata tcaggcataa gagaggttaa gacctcaagc caggagttaa    240
acagggcatt aaaagccatc attgctgctg atggcttgaa caacatcacc tgccatggta    300
aggatcctga ggacaagatt tccctcataa agggctcctc tcacaaaaag cgggtgggga    360
tagttcgggtg tgagagacga agagatgcta agcaaatagg gagagaaacc atggcagggg    420
ttgcaatgac agtccttcca gccttagcag tttttgcttt ggcacctgtt gtttttgctg    480
aagaccccc tctcagaaac agaccaggga aggggcacaa ctacattgac gggatgactc    540
aggaggatgc cacatgcaaa cctgtgacat atgctggggc atgtagcagt tttgatgtct    600
tgcttgaaaa gggaaaattt ccccttttcc agtcgtatgc tcatcataga actctactag    660
aggcagttca cgacaccatc attgcaaagg ctgatccacc tagctgtgac cttcagagtg    720
ctcatgggaa cccctgcatg aaagagaaac tegtgatgaa gacacactgt ccaaagtact    780
accagtcagc tcattacctc aacaatgacg ggaaaatggc ttcagtcaag tgcctccta    840
agtatgagct cactgaggac tgcaactttt gtaggcagat gacaggtgct agcctgaaga    900
aggggtctta tcctctccaa gacttgtttt gtcagtcaag tgaggatgat ggatcaaaat    960
taaaaacaaa aatgaaaggg gtctgcgaag tgggggttca agcactcaaa aagtgtgatg   1020
gccaaactcag cactgcacat gaggttgctg cctttgcagt gtttaagaac tcaaagaagg   1080
tttatcttga taagcttgac cttaagactg aggagaatct gctaccagac tcatttgtct   1140
gtttcgagca taagggacag tacaaggaa caatggactc tggtcagact aagaggggagc   1200
tcaaaagctt tgatatctct cagtgcacca agattggagg acatggtagt aagaagtgca   1260
ctggggacgc agcattttgc tctgcttatg agtgcactgc tcagtagcc aatgcctatt   1320
gttcacatgc taatgggtca gggattgtgc agatacaagt atcaggggtc tggagaagc   1380
ctttatgtgt agggatagag agagtgggtg tgaagagaga actctctgcc aagcccatcc   1440
agagagttga gcottgcaca acttgataa ccaaatgtga gctcatgga ttggttgtcc   1500
gatcaacagg gttcaagata tcatcagcag ttgcttgtgc tagcggagtt tgcgtcacag   1560
gatcgagag tccttcacc gagattacac tcaagtatcc agggatatcc cagtcttctg   1620
ggggggacat aggggttcac atggcacacg atgatcagtc agttagctcc aaaatagtag   1680
ctcactgcc tcaccaggac ccgtgcttag tgcattggctg catagtgtgt gctcatggcc   1740
tgataaatta ccaggtcac actgctctca gtgcctttgt tgttgtgttt gtattcagtt   1800
ctattgcaat aatttgttta gctgttcttt atagggtgct taagtgcctg aagattgcc   1860
caaggaaagt tctgaatcca ctaatgtgga tcacagcctt catcagatgg atatataaga   1920
agatggttgc cagagtggca gacaacatta atcaagtgaa cagggaaata ggatggatgg   1980
aaggaggtca gttggttcta ggaaccctg cccctattcc togtcatgcc ccaatccac   2040
gttatagcac atacctgatg ttattattga ttgtctcata tgcacagca tgttcagaac   2100

```

-continued

```

tgattcaggc aagctccaga atcaccactt gctctacaga ggggtgtaac accaagtgt 2160
gactgtctgg cacagcattg atcagagcag ggtcagttgg ggcagaggct tgtttgatgt 2220
tgaagggggg caaggaagat caaaccaagt tcttaagat aaaaactgtc tcaagtgagc 2280
tatcatgcag ggagggccag agttattgga ctgggtcctt tagccctaaa tgtttgagct 2340
caaggagatg ccaccttgtc ggggaatgcc atgtgaatag gtgtctgtct tggagggaca 2400
atgaaacttc agcagagttt tcatattgtt gggaaagcac gaccatgcga gagaataagt 2460
gttttgagca atgtggagga tgggggtgtg ggtgtttcaa tgtgaacca tcttgcttat 2520
ttgtgcacac gtatctgcag tcagttagaa aagaggccct tagagttttt aactgtatcg 2580
actgggtgca taaactcact ctagagatca cagactttga tggctctgtt tcaacaatag 2640
acttgggagc atcatctagc cgtttcacaa actggggctc agttagctc tcaactggagc 2700
cagagggcat ctcaggctca aatagctttt ctttcattga gagcccaggc aaagggatg 2760
caattgttga tgagccatc tcagaaatc ctcggcaagg gttcttgggg gagatcaggt 2820
gcaattcaga gtccctcagtc ctgagtgctc atgaatcatg ccttagggca ccaaacctta 2880
tctcatacaa gcccatgata gatcaattgg agtgcacaac aaatctgatt gatccctttg 2940
ttgtcttga gaggggttct ctgccacaga caaggaatga taaaacctt gcagcttcaa 3000
aaggaaatag aggtgttcaa gctttctcta agggctctgt acaagctgat ctaactctga 3060
tgtttgacaa ttttgagggt gactttgtgg gagcagccgt atcttctgat gccgcttct 3120
taaatttgac aggttgctat tcttgcaatg caggggccag ggtctgctg tctatcacat 3180
ccacaggaac tggatctctc tctgccaca ataaggatgg gtctctgcat atagtccttc 3240
catcagagaa tggaaacaaa gaccagtgtc agatactaca cttcactgtg cctgaagtag 3300
aggaggagtt tatgtactct tgtgatggag atgagcggcc tctgttgggt aaggggacc 3360
tgatagccat tgatccattt gatgataggc gggaaagcagg gggggaatca acagttgtga 3420
atccaaaatc tggatcttgg aatttcttg actggtttc tggactcatg agttggtttg 3480
gagggcctct taaaactata ctctcattt gcctgtatgt tgcatatca attgggctct 3540
tttctcctt tatatatctt ggaagaacag gcctctctaa aatgtggctt gctgccacta 3600
agaaggctc atagatcagt acgtgtaaaa gcaatatgtt gaaataagta gacacaagca 3660
aacctaatta tgtaagtgtt gtacagatag gtcaaattat tggaaatcc aagcttagaa 3720
acttatgcaa taactctta gatgtaagct tagttgtaat ttgggtgggt ggggtgagc 3780
agcagcagtc tcaagtctt gtgaatattc tagttggcgt aatctgttt tgccagatta 3840
gctgggaatt aaactaactc tttgaagttg caccgtctt tgtgt 3885

```

<210> SEQ ID NO 3

<211> LENGTH: 6404

<212> TYPE: DNA

<213> ORGANISM: Rift Valley fever virus

<400> SEQUENCE: 3

```

acacaaaggc gcccaatcat ggattctata ttatcaaac agctggttga caagactggt 60
tttgtagag tgccaatcaa gcattttgac tgtacaatgc taactctggc acttccaaca 120
tttgatgttt ccaagatggg agatagaatt accatagact tcaatctgga tgatatacaa 180
ggagcatctg aaataggctc aactttgcta cctccatgt cgatagatgt ggaagatag 240

```

-continued

gccaatTTTg tTcacgattt cacettTggc cacttagctg acaagactga cagactgtta 300
atgcgtgagt tTcccatgat gaatgacggg tTtgatcatt tgagccctga catgatcatt 360
aaaactacat ctggcatgta caacatcgtt gagttcacca ccttagggg agatgaaaga 420
ggtgcattcc aggctgccat gactaaactc gctaagtatg aggttccttg tgagaacaga 480
tctcagggca ggactgttgt tctttatggt gttagtgtt atcggcatgg tgtatggtct 540
aatctggagc tagaggactc tgaagcagag gagatggttt ataggtaacag acttctctct 600
agtgtgatgg atgagctaag gacctgttcc ccagaactgt catccacaga tgaggaacta 660
gggaagactg agagagagtt gctagccatg gtctctctca tccaaataaa ttggtcagtc 720
acagaatctg tgtttccacc cttcagcaga gaaatgtttg acaggtttag atcctcccct 780
cccgatccag agtatatcac gaggatagtg agcagatgcc tcataaattc tcaagagaaa 840
ctcatcaata gttccttctt tgctgaaggg aatgataagg ctctgagatt tTcaaaaaac 900
gctgaagagt gttccttggc agtagagaga gccttaaatc agtatagagc agaagacaac 960
cttagggacc tcaatgacca caagtcaact attcagctgc ctccctggct gtcctatcat 1020
gatgtcagtg gcaaaagatct gtgcctctct cagggttag atgtgagagg ggaccatccc 1080
atgtgcaact tgtggaggga agtggtcacc tctgcaaac tagaggagat tgagaggatg 1140
cacgatgatg cagcagcaga acttgagttt gctcttctcg gagtaaagga caggccagat 1200
gagagaaaca gataccatag agtccaccta aatatgggct cagatgatag tgtctacata 1260
gctgctttag gagttaatgg aaagaagcat aaagcagaca ctttagtgca acaaatgaga 1320
gacaggagta aacagccttt ctccccagac cacgatgtgg atcacatata tgaatttctc 1380
tctgcatgct ctagtgaact gtgggcaaca gatgaggacc tgtacaacc tctctctgt 1440
gataaagagc ttagattggc agcccagagg attcatcagc catccttctc agaaaggggt 1500
tTcaatgaga tcataacaga gactacaaa tTcatgggaa gtaggatagg tTcatggtgc 1560
caaatggtca gcttgatagg agctgagcta tcagcttctg tTaaacaaca tgtcaagcct 1620
aactactttg tgattaaacg actactaggt tctgggattt tcttctaat caagcccact 1680
tccagcaaaa gccatatatt tgtgtctttt gcaattaagc gctcttctc ggcccttgat 1740
ctctccactt ccagggtttt caagccctac atagatgctg gggatctgtt agttactgac 1800
tttgttctt ataagctaag caagcttacc aaactctgca agtgcgttcc attaatggag 1860
tcctccttct cattctgggc agaagcattt ggaattccaa gctggaactt tgttggtgac 1920
ttgttcaggt cttcagactc tgcagcaatg gatgcctcat acatgggcaa actttcttta 1980
tTaaacctt tTgaagacaa agcagcaact gaagagttac agactattgc aagatatata 2040
atcatggagg gctttgtctc gccccagaa atccccaaac ctcaagat gacctctaag 2100
tttctaagg tTctcaggtc agagctgcag gtttacttat taaactgctt atgcagaact 2160
atccagagaa tagcagggta gcccttcatt cttaagaaga aggatgggtc tatatcctgg 2220
ggtggcatgt tcaatccttt tTcagggcgt cactgcttg atatgcaacc actcatcagc 2280
tgttgttaca atggttactt taaaaataaa gaagaagaga ctgagccttc gtccttctc 2340
gggatgtata agaaaaatc agaacttgag caccttagac cacagtcaga tgccttctg 2400
ggttcaaaag atccagaact tcccagaatg catgagtcca gtgttctca cttgaaggag 2460
gcttgcaatc atgctaagct agtcttgagg agcctctatg gacagaattt catggagcag 2520

-continued

atagacaacc agattattcg agagctcagt gggttgactc tagaaagggt ggccacactt	2580
aaggccacaa gcaactttaa tgagaattgg tatgtctata aggatgtagc agacaaaaac	2640
tacacaaggg ataaattatt agtgaagatg tcaaaatag cctctgaggg aaagagccta	2700
gctatccaga agtttgagga ttgtatgagg cagatagagt cacaaggatg catgcatatt	2760
tgtttgttta agaaacaaca gcatggaggt ctgagagaga tctatgtgat gggtgagag	2820
gaaagaattg ttaactcggg ggtggagaca atagccaggt ccatagggaa gttccttgct	2880
tctgataccc tctgtaaccc cccaataaa gtgaaaattc ctgagacaca tggcatcagg	2940
gcccgaagc aatgtaaggg gcctgtgtgg acttgtgcaa catcagatga tgcaaggaag	3000
tggaaccaag gccattttgt tacaagtgtt gccctcatgc tgtgtgagtt cacctctcct	3060
aatgggtggc cgctgatcat taggggatgc tcaatgttta ccaggaaaag gatgatgatg	3120
aatttgaatt atcttaagat cctggatggt catcgggagc ttgatattag agatgacttt	3180
gtgatggatc tcttcaaacg ttatcatggc gaggcagaag ttccatgggc ctttaagggc	3240
aaaacatatt tggaaccac aacagggatg atgcagggaa tactgcatta tacttcctca	3300
ctattacaca ccattcacca agaatacatc cggtccttgt ccttaagat attcaacctg	3360
aaggttgctc ctgagatgag caagggcctg gtttgtgaca tgatgcaagg atcagatgat	3420
agtagtatgc taatcagctt cccagctgat gatgagaagg ttcttaccag atgcaaagtg	3480
gccgcagcta tatgcttccg catgaagaag gagctgggag tgtacctgc catttacctc	3540
tcagagaagt ccacagcaaa cacagatttt gtgatggagt acaattctga attttattc	3600
cacaccagc atgttagacc aacgatcagg tggattgcag cttgttcag cctgccagaa	3660
gtgaaacac tagtagcccg ccaggaagag gcctctaacc taatgacttc agttactgag	3720
ggaggtgggt cattctcctt agctgcaatg attcagcaag ctcagtgcac tctccattac	3780
atgctgatgg gcatgggagt gtctgagcta ttcttagagt ataagaaggc agtgcggaag	3840
tggaatgacc ctggcctggg tttcttctctg cttgacaatc cttatgcgtg cggattggga	3900
ggtttcagat ttaatctctt caaagctatc accagaactg atttgcagaa gctatatgct	3960
ttcttcatga agaaggtaaa gggctcagct gctagggact gggcagatga agatgtcacc	4020
atcccagaaa cgtgtagcgt gagccaggt gccgctctaa ttcttagctc ctctctaaag	4080
tgggatccta ggaagaagtt tcagaaattg agagaccgtt tgaacatacc agagaactgg	4140
attgaactaa taaatgagaa tccagaggtg ctctatcggg ctcccagaac aggccagaa	4200
atattgttgc gcatcgaga gaaagtccat agcccaggtg ttgtgtcacc attgtcttct	4260
ggcaatgcag tttgtaaagt catggcctca gctgtatact tcttatcagc aacaattttt	4320
gaggacactg gacgtcctga gttcaacttc ttggaggatt ctaagtacag cttgctacaa	4380
aagatggctg catattctgg ctttcatggt tttaatgata tggagccaga agatatatta	4440
ttcttattcc cgaatattga ggaattagaa tcaactggatt ctatagttaa caacaagga	4500
gaaatagaca tcatcccaag agtcaacatc agggatgcaa cccaaaccag ggtcactatc	4560
tttaatgagc agaagacct cggacatct ccagagaagt tgggtgcaga caagtggttt	4620
gggactcaga agagtaggat agggcaaaaca accttctctg ctgaatggga aaagctaaag	4680
aaaattgtaa agtggttggg agacactcca gaagcaactc tagctcacac cccactgaat	4740
aaccatattc aagttaggaa tttcttctctg agaatggaaa gcaagcctag aacagtcaga	4800

-continued

```

ataacaggag ctccagtaaa gaagaggta ggggttagta agatagetat ggttatccgt 4860
gacaatttct cccggatggg ccatcttcga ggtgtagaag accttgctgg cttcactcgt 4920
agtgtgtcag ctgaaattct caagcacttt ctattctgta tactacaagg tccatacagt 4980
gagagctata agctacagct aatctacaga gtcttaagct cagtgtcaaa cgttgagata 5040
aaggaatcag atggtaagac aaaaaccaac ttgattgaa tccttcagag atttctagat 5100
ggatgatcag ttgtcccat aattgaagag atgggagccg gaacagtggtg tggattcatc 5160
aagagacaac aatctaaagt tgtgcagaac aaagtggctt attatggagt tgggatttgg 5220
agaggcttca tggatggata tcaggtccat ctagagatag aaaatgacat aggacagccc 5280
ccaaggctta ggaatgtcac aactaactgt cagagcagcc catgggacct gagtattcca 5340
ataaggcaat gggcagaaga catgggggtc acaacaacc aggattattc ctctaaatct 5400
agcagagggg ccagatattg gatgcattca ttcaggatgc aaggacctag caagccattt 5460
ggatgcccag tttatattat taagggtgat atgtcagatg tcatcagact gagaaaggag 5520
gaggtggaga tgaaagtacg gggctctact ctcaacttgt acaccaagca ccattctcat 5580
caggacctac acattctatc ttactactga tcagacaatg atctcagtc aggcatcttc 5640
aagtcaatat cagatgaggg ggtggctcaa gcctcgaat tatttgagag ggagccaagc 5700
aactgctggg tgagatgtga gtctgtagcc ccaaaattha tatcagccat ccttgagata 5760
tgtgagggga agagacagat aaggggaatt aacagaacca gactctcaga gattgtgaga 5820
atgtgtcttg aatcttcctt aagatcaaaa gtcggatcta tgttctcatt tgtcgccaat 5880
gtcagaggag cccatgatgt gattatgat gcgttaatgg atctaatgat agaggatgcc 5940
aagaacaatg cattcagca tgggtgtgac tgcatagagt tggatgttag tggcccttac 6000
gagatggagt cttttgatac atctgatgtc aatctctttg ggccagccca ttacaaggac 6060
atcagttcat tatctatgat tgcctatccc ttaatggata agtttgttga ttatgctatt 6120
tctaagatgg ggagagcctc agttaggaaa gttctagaaa caggtcggtg ctccagcaaa 6180
gactatgatt tatcaaaggt tctcttcaga actctacaga gaccagaaga aagcattagg 6240
atagatgatc tggaattata tgaggagaca gatgtggcgg atgacatgct aggctaagac 6300
caataagcaa agtcaggctt agatttaggg atactatgct agtattgaa tccatgtggg 6360
ttctgatact agcatagtgc tacaatattg ggcggtcttt gtgt 6404

```

```

<210> SEQ ID NO 4
<211> LENGTH: 33
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

```

```

<400> SEQUENCE: 4

```

```

aaaaaaggta ccgatatact tgataagcac tag 33

```

```

<210> SEQ ID NO 5
<211> LENGTH: 30
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

```

```

<400> SEQUENCE: 5

```

-continued

aaaaaaagat ctgattagag gttaaggctg 30

<210> SEQ ID NO 6
<211> LENGTH: 33
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 6

aaaaaaggta ccatggtgag caagggcgag gag 33

<210> SEQ ID NO 7
<211> LENGTH: 34
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 7

aaaaaaagat ctttacttgt acagctcgtc catg 34

<210> SEQ ID NO 8
<211> LENGTH: 42
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 8

aaaaaaggta cctgctgctg catcaacctc aacaaatcca tc 42

<210> SEQ ID NO 9
<211> LENGTH: 49
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 9

aaaaaacgtc tcagcagcag cagcagcaat ggtgagcaag ggcgaggag 49

<210> SEQ ID NO 10
<211> LENGTH: 53
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 10

aaaaaacgtc tcaactgctgc tgctgctgct gcatcaacct caacaaatcc atc 53

<210> SEQ ID NO 11
<211> LENGTH: 4594
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Plasmid

<400> SEQUENCE: 11

acacaaagct ccttagagat acaaacacta ttacaataat ggacaactat caagagcttg 60

gcatccagtt tgctgctcaa gcagtgacc gcaatgagat tgaacagtgg gtccgagagt 120

-continued

ttgcttatca agggtttgat gcccgtagag ttatcgaact cttaaagcag tatggtgggg	180
ctgactggga gaaggatgcc aagaaaatga ttgttctggc tctaactcgt ggcaacaagc	240
ccaggaggat gatgatgaaa atgtcgaaag aaggcaaagc aactgtggag gctctcatca	300
acaagtataa gctaaaggaa gggaaatcctt cccgggatga gttgactcta tcacgagttg	360
ctgccgcctt ggctggctgg acatgccagg ctttggctgt cttgagttag tggcttctctg	420
tactgggac taccatggac ggcatatccc ctgcataccc aaggcatatg atgcaccca	480
gctttgtctg catggtggat ccttctctac caggagacta tctaagggca atattagatg	540
ctcactctct gtatctctctg cagttctccc gggtcatcaa cccaaacctc cgaggtagaa	600
caaaagagga ggttgctgca acgttcacgc agccaatgaa tgcagcagtg aatagcaact	660
ttataagcca tgagaagagg agagaattct tgaaagcctt tggacttggt gatttcaatg	720
ggaagccgtc agctgctgtc atggcagccg ctccaggctta caagacagca gcctaagtgg	780
ctgccaggg ggttgggggg aaggggagtt ggggttacgg tcgggattgg ggggtggggg	840
gtggggcagc cttaacctct aatcaacctc aacaaatcca tcatcatcac tctcctctc	900
tgattccatc tcaacatctg ggattggagg aataactgga atccagttgt ttctcccat	960
catgctggga agtgatgagc gcagcatcag gctctctccc ataagaacaa tgagggtgga	1020
gtttgaact acagcattag aaatgtcctc ttttctctgt tgcagaagcc gaacgcactg	1080
tacgtgagca acotcataka tgagatcaaa gcctggcaac aggcacaggt caatccctct	1140
gaggatggcc tcagtcgcta tcatcctgtg taagccaaca aaggagtcct ctagatcatt	1200
ggtgatcttg caactcctca ttgctagagt ggcaatctga tcccttctaa tgtcatcatt	1260
cctatgcact ctagtagagc ttaggtcaaa gaaagccagt gagggttctc caagaggcca	1320
ggatattggt tcttcagat tggggaacct tgtgaaatca ctaagagtca tatggcctat	1380
tagatcaata agtctctgaa aaggctctgc tggtgagggt gcaacgtttg atgcaaagtc	1440
tccaagtccg actcggtagt ggaattctcc gacattgtag aaatcagaga atcgcaagcg	1500
aacctctgta ctaggacgat ggtgcatgag aaagacacaa cagggcccaa ccatagaata	1560
aggtatcctg ggaggacct ctctctaaa gtactccact gacacaacac gacgacct	1620
ctgcaaatca acagatatca caggaaagta atccatgata tacttgataa gcactagggg	1680
gtctttgtgt gggtcggcat ggcactctca cctcctcggc gtccgacctg ggcactcgaa	1740
ggaggacgtc gtccactcgg atggctaagg gagagctcgg atccggctgc taacaaagcc	1800
cgaaggaag ctgagttggc tgcctgccacc gctgagcaat aactagcata accccttggg	1860
gcctctaaac gggctctgag gggttttttg ctgaaaggag gaactatata cggatcgaga	1920
tcctctaggt acaagcctaa ttgtgtagca tctggcttac tgaagcagac cctatcatct	1980
ctctcgtaaa ctgccctcag agtcggtttg gttggacgaa cctctctgagt ttctggtaac	2040
gccgtcccgc acccggaaat ggtcagcgaa ccaatcagca gggctcatcgc tagccagatc	2100
ctctacgccg gacgcactgc ggccggcacc accggcgcca caggtgcggt tgetggcgcc	2160
tatatcgccg acatcaccga tggggaagat cgggctcgc acttcgggct catgagcct	2220
tgtttcggcg tgggtatggt ggcaggcccc gtggccgggg gactgttggg cgccatctcc	2280
ttgcaccatt ccttgcggcg cgggtgctca acggcctcaa cctactactg ggetgcttcc	2340
taatgcagga gtcgcataag ggagagcgtc gatatggtgc actctcagta caatctgctc	2400

-continued

tgatgcegca tagttaagcc agccccgaca ccgccaaca cccgctgacg cgecctgacg	2460
ggcttgtctg ctccccgcat ccgcttacag acaagctgtg acaaagggcc tcgtgatacg	2520
cctatnttta taggttaatg tcatgataat aatggtttct tagacgtcag gtggcacttt	2580
tcggggaat gtgcgcgaa cccctatttg tttatnttcc taaatacatt caaatatgta	2640
tccgctcatg agacaataac cctgataaat gcttcaataa tattgaaaa ggaagagtat	2700
gagtattcaa catttccgtg tcgcccctat tcctntttt gcggcatttt gecttccgtg	2760
ttttgctcac ccagaaacgc tgggtaaagt aaaagatgct gaagatcagt tgggtgcacg	2820
agtgggttac atcgaactgg atctcaacag cgtaagatc cttgagagtt ttcgccccga	2880
agaacgtttt ccaatgatga gcactnttaa agttctgcta tgtggcgcgg tattatcccc	2940
tattgacgcc gggcaagagc aactcggctg ccgcatacac tattctcaga atgacttggg	3000
tgagtactca ccagtcacag aaaagcatct tacggatggc atgacagtaa gagaattatg	3060
cagtgtctgc ataaccatga gtgataaac tcgcccaca ttacttctga caacgatcgg	3120
aggaccgaag gagctaaccg cttntttgca caacatgggg gatcatgtaa ctgccttga	3180
tcgttgggaa ccggagctga atgaagccat accaaacgac gagcgtgaca ccacgatgcc	3240
tgtagcaatg gcaacaacgt tcgcgaaact attaactggc gaactactta ctctagcttc	3300
ccggcaacaa ttaatagact ggatggaggc ggataaagtt gcaggaccac ttctgcgctc	3360
ggcccctccg gctggctgtg ttattgctga taaatctgga gccggtgagc gtgggtctcg	3420
cggtatcatt gcagcactgg ggcagatgg taagcccctc cgtatcgtag ttatctacac	3480
gacggggagt caggcaacta tggatgaac aaatagacag atcctgaga taggtgctc	3540
actgattaag cattggtaac tgctcagacca agtttactca tataacttt agattgattt	3600
aaaacttcat tnttaattta aaaggatcta ggtgaagatc cttnttgata atctcatgac	3660
caaaatccct taacgtgagt tntcgttcca ctgagcgtca gaccccgtag aaaagatcaa	3720
aggatcttct tgagatcctt tnttctgctg cgtaatctgc tgcttgcaaa caaaaaaac	3780
accgctacca gcggtggtt gtttgccgga tcaagagcta ccaactctt tccgaaggt	3840
aactggcttc agcagagcgc agataccaaa tactgtcctt ctagtgtagc cgtagttagg	3900
ccaccacttc aagaactctg tagcaccgcc tacatacctc gctctgctaa tcctgttacc	3960
agtggctgct gccagtgccg ataagctcgtg tcttaccggg ttggactcaa gacgatagtt	4020
accggataag gcgcagcggc cgggctgaac ggggggttcg tgcacacagc ccagcttgg	4080
gcgaacgacc tacaccgaac tgagatacct acagcgtgag ctatgagaaa gcgccacgct	4140
tcccgaaggg agaaaggcgg acaggtatcc ggtaaagcgc agggctcgaa caggagagcg	4200
cacgagggag cttccagggg gaaacgcctg gtatctttat agtcctgtcg ggttctcgca	4260
cctctgactt gacgctcgat tnttgtgatg ctctgtaggg gggcggagcc tatggaaaa	4320
cgccagcaac gcggcctttt taecggttcc ggccntttgc tggccttttg ctccatggt	4380
cttctctgct ttatccccctg attctgtgga taaccgtatt accgctttg agtgagctga	4440
taccgctcgc gcagccgaa cgaccgagcg cagcagatca gtgagcaggg aagcggaa	4500
gcgccaata gcgaaaccgc ctctccccgc gcggtggcgg attcattaat gcagggggat	4560
ctcgatcccc cgaaattaat acgactcact atag	4594

-continued

<210> SEQ ID NO 12
<211> LENGTH: 6789
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic plasmid

<400> SEQUENCE: 12

acacaaagac ggtgcattaa atgtatgttt tattaacaat tctaactctcg gttctgggtg 60
gtgaagcggg tattagagtg tctctaagct ccacaagaga agagacctgc tttgggtgact 120
ccactaacc agagatgatt gaaggagcct gggattcact cagagaggag gagatgccgg 180
aggagctctc ctgttctata tcaggcataa gagaggttaa gacctcaagc caggagttaa 240
acagggcatt aaaagccatc attgctgctg atggctttaa caacatcacc tgccatggta 300
aggatcctga ggacaagatt tccctcataa agggctcctc tcacaaaaag cgggtgggga 360
tagttcgggtg tgagagacga agagatgcta agcaaatagg gagagaaacc atggcagggga 420
ttgcaatgac agtccttcca gccttagcag tttttgcttt ggcacctgtt gtttttctg 480
aagaccccca tctcagaaac agaccagggg aggggcacaa ctacattgac gggatgactc 540
aggaggatgc cacatgcaaa cctgtgacat atgctggggc atgtagcagt tttgatgtct 600
tgcttgaaaa gggaaaatth ccccttttcc agtcgtatgc tcatcataga actctactag 660
aggcagttca cgacaccatc attgcaaagg ctgatccacc tagctgtgac cttcagagtg 720
ctcatgggaa cccctgcatg aaagagaaac tcgtgatgaa gacacactgt ccaaatgact 780
accagtgcgc tcattacctc aacaatgacg ggaaaatggc ttcagtcaag tgcctccta 840
agtatgagct cactgaggac tgcaactttt gtaggcagat gacaggtgct agcctgaaga 900
aggggtctta tctctccaa gacttgthtt gtcagtcaag tgaggatgat ggatcaaat 960
taaaaaaaa aatgaaaggg gtctgcgaag tgggggttca agcactcaaa aagtgtgatg 1020
gccaactcag cactgcacat gaggttgctc cctttgcagt gtttaagaac tcaagaagg 1080
tttatcttga taagcttgac ctttaagactg aggagaatct gctaccagac tcatttctct 1140
gtttcgagca taagggacag tacaaggaa caatggactc tggtcagact aagagggagc 1200
tcaaaagctt tgatatctct cagtgcacca agattggagg acatggtagt aagaagtgca 1260
ctggggacgc agcattttgc tctgcttatg agtgcactgc tcagtacgcc aatgcctatt 1320
gttccatgac taatgggtca gggattgtgc agatacaagt atcaggggtc tggagaagc 1380
ctttatgtgt agggatgag agagtggttg tgaagagaga actctctgcc aagcccatcc 1440
agagagttga gccttgaca acttgataa ccaaatgtga gcctcatgga ttggtgtctc 1500
gatcaacagg gttcaagata tcatcagcag ttgcttctgc tagcggagtt tgcgtcacag 1560
gatcgcagag tcttccacc gagattacac tcaagtatcc agggatatcc cagtcttctg 1620
ggggggacat aggggttcac atggcacacg atgatcagtc agttagctcc aaaatagtag 1680
ctcactgcc tccccaggac ccgtgtttag tgcattgctg catagtgtgt gctcatggcc 1740
tgataaatta ccagtgtcac actgcttca gtgcctttgt tgtgtgttt gtattcagtt 1800
ctattgcaat aattgttta gctgttcttt ataggtgtct taagtgcctg aagattgcc 1860
caaggaaagt tctgaatcca ctaatgtgga tcacagcctt catcagatgg atatataaga 1920
agatgggtgc cagagtggca gacaacatta atcaagttaa cagggaaata ggatggatgg 1980
aaggaggtca gttggttcta gggaaacctg cccctattcc tgcctatgcc ccaatccac 2040

-continued

gttatagcac atacctgatg ttattattga ttgtctcata tgcacagca tgttcagaac	2100
tgattcaggc aagctccaga atcaccactt gctctacaga ggggtttaac accaagtgtg	2160
gactgtctgg cacagcattg atcagagcag ggtcagttgg ggcagaggct tgtttgatgt	2220
tgaagggggt caaggaagat caaaccaagt tcttaagat aaaaactgtc tcaagtgagc	2280
tatcatgcag ggagggccag agttattgga ctgggtcctt tagccctaaa tgtttgagct	2340
caaggagatg ccaccttgtc ggggaatgcc atgtgaatag gtgtctgtct tggagggaca	2400
atgaaacttc agcagagttt tcatctgttg gggaaagcac gaccatgcga gagaataagt	2460
gttttgagca atgtggagga tgggggtgtg ggtgtttcaa tgtgaacca tcttgcttat	2520
ttgtgcacac gtatctgcag tcagttagaa aagaggccct tagagttttt aactgtatcg	2580
actgggtgca taaactcact cttagagatca cagactttga tggctctggt tcaacaatag	2640
acttgggagc atcatctagc cgtttcacaa actgggggtc agttagctc tcaactggagc	2700
cagagggcat ctcaggctca aatagctttt ctttcattga gagcccaggt aaagggtatg	2760
caattgttga tgaccattc tcagaaattc ctcggcaagg gttcttgggg gagatcaggt	2820
gcaattcaga gtctcagtc ctgagtgctc atgaatcatg ccttagggca ccaaacctta	2880
tctcatacaa gcccatgata gatcaattgg agtgcacaac aaatctgatt gatcccttg	2940
ttgtcttga gaggggttct ctgccacaga caaggaatga taaaaccttt gcagcttcaa	3000
aaggaaatag aggtgttcaa gctttctota agggtctgt acaagctgat ctaactctga	3060
tgtttgacaa ttttgaggtg gactttgtgg gagcagccgt atcttgtgat gccgccttct	3120
taaattgac aggttgctat tcttgcaatg caggggccag ggtctgctg tctatcacat	3180
ccacaggaac tggatctctc tctgccaca ataaggatgg gtctctgcat atagtccttc	3240
catcagagaa tggaaacaaa gaccagtgtc agatactaca ctctactgtg cctgaagtag	3300
aggaggagtt tatgtactct tgtgatggag atgagcggcc tctgttggtg aaggggacct	3360
tgatagccat tgatccattt gatgataggg gggaaagcagg gggggaatca acagttgtga	3420
atccaaaatc tggatcttgg aatttcttg actggttttc tggactcatg agttggtttg	3480
gagggcctct taaaactata ctctcattt gcctgtatgt tgcattatca attgggctct	3540
ttttcctcct tatatatctt ggaagaacag gcctctctaa aatgtggctt gctgccacta	3600
agaaggctc atagatcagt acgtgtaaaa gcaatatgtt gaaataagta gacacaagca	3660
aacctaatta tgtaagtgtt gtacagatag gtcaaattat tggaaatcc aagcttagaa	3720
acttatgcaa taatacttta gatgtaagct tagttgtaat ttgggtgggt ggggtgagc	3780
agcagcagtc tcaagtgtt gtgaatattc tagttggcgt aatcgtcttt tgcagatta	3840
gctgggaatt aaactaactc tttgaagttg caccgtctt tgtgtgggtc ggcattggat	3900
ctccacctcc tcgcggtccg acctgggcat ccgaaggagg acgtcgtcca ctcgatggc	3960
taagggagag ctcggtaccg gctgctaaca aagccgaaa ggaagctgag ttggctgctg	4020
ccaccgctga gcaataacta gcataacccc ttggggcctc taaacgggtc ttgaggggtt	4080
ttttgctgaa aggaggaact atatccggat cgagatcctc taggtacaag cctaatttg	4140
tagcatctgg cttactgaag cagaccctat catctctctc gtaaactgcc gtcagagtcg	4200
gtttggttg acgaacctc tgagttctg gtaacgccgt cccgcaccg gaaatggtea	4260
gcgaaccaat cagcagggtc atcgctagcc agatcctcta cgccggacgc atcggtggcg	4320

-continued

gcataccggc	cgccacaggt	gcggttgctg	gcgctatat	cgccgacatc	accgatgggg	4380
aagatcgggc	tcgccacttc	gggctcatga	gcgcttggtt	cggcgtgggt	atggtggcag	4440
gccccgtggc	cgggggactg	ttgggcgcca	tctccttgca	ccattccttg	cggcgcggt	4500
gctcaacggc	ctcaacctac	tactgggctg	cttcctaattg	caggagtgcg	ataagggaga	4560
gcgtcgatat	ggtgcactct	cagtacaatc	tgctctgatg	ccgcatagtt	aagccagccc	4620
cgacaccggc	caacaccggc	tgacgcgccc	tgacgggctt	gtctgctccc	ggcatccgct	4680
tacagacaag	ctgtgacaaa	gggcctcgtg	atagcctat	ttttataggt	taatgtcatg	4740
ataataatgg	tttcttagac	gtcaggtggc	acttttcggg	gaaatgtgcg	cggaaccctc	4800
atgtgtttat	ttttctaaat	acattcaaat	atgtatccgc	tcatgagaca	ataaccctga	4860
taaagtcttc	aataatattg	aaaagggaag	agtatgagta	ttcaacattt	ccgtgtcgcc	4920
cttattccct	tttttgccgg	attttgccctt	cctgtttttg	ctcaccagga	aacgctgggtg	4980
aaagtaaaag	atgtgaaga	tcagttgggt	gcacgagtgg	gttacatcga	actggatctc	5040
aacagcggta	agatccttga	gagttttcgc	cccgaagaac	gttttccaat	gatgagcact	5100
tttaaagttc	tgctatgtgg	gcggttatta	tcccgtattg	acgcgggcca	agagcaactc	5160
ggtcgcccga	tacactattc	tcagaatgac	ttggttgagt	actcaccagt	cacagaaaag	5220
catcttacgg	atggcatgac	agtaagagaa	ttatgcagtg	ctgccataac	catgagtgat	5280
aacactcggg	ccaacttact	tctgacaacg	atcggaggac	cgaaggagct	aaccgctttt	5340
ttgcacaaca	tgggggatca	tgtaactcgc	cttgatcgtt	gggaaccgga	gctgaatgaa	5400
gccataccaa	acgacgagcg	tgacaccacg	atgcctgtag	caatggcaac	aacgttgccg	5460
aaactattaa	ctggcgaaact	acttactcct	gcttcccggc	aacaattaat	agactggatg	5520
gaggcggata	aagttgcagg	accactctct	cgctcggccc	ttccggctgg	ctggtttatt	5580
gctgataaat	ctggagccgg	tgagcgtggg	tctcgcggta	tcattgcagc	actggggcca	5640
gatggtaagc	cctcccgtat	cgtagttatc	tacacgacgg	ggagtccagg	aactatggat	5700
gaacgaaata	gacagatcgc	tgagataggt	gcctcactga	ttaagcattg	gtaactgtca	5760
gaccaagttt	actcatatat	actttagatt	gattttaaac	ttcattttta	attttaaagg	5820
atctaggtga	agatcctttt	tgataatctc	atgaccaaaa	tccttaacg	tgagttttcg	5880
ttccactgag	cgctcagacc	cgtagaaaag	atcaaaggat	cttcttgaga	tccttttttt	5940
ctgcgcgtaa	tctgctgctt	gcaaacaaaa	aaaccaccgc	taccagcggg	ggtttgtttg	6000
ccggatcaag	agctaccaac	tctttttccg	aaggtaactg	gcttcagcag	agcgcagata	6060
ccaaatactg	tccttctagt	gtagccgtag	ttaggccacc	acttcaagaa	ctctgtagca	6120
ccgcctacat	acctcgctct	gctaactcctg	ttaccagtgg	ctgctgccag	tgccgataag	6180
tcgtgtctta	ccgggttgga	ctcaagacga	tagttaccgg	ataaggcgca	gcggtcgggc	6240
tgaacggggg	gttcgtgcac	acagcccagc	ttggagcgaa	cgacctacac	cgaactgaga	6300
tacctacagc	gtgagctatg	agaaagcgcc	acgcttcccc	aagggagaaa	ggcggacagg	6360
tatccggtaa	gcggcagggt	cggaacagga	gagcgcacga	gggagcttcc	agggggaaac	6420
gcctggatc	ttatagctc	tgctcgggtt	cgccacctct	gacttgagcg	tcgatttttg	6480
tgatgctcgt	cagggggggc	gagcctatgg	aaaaacgcca	gcaacgcggc	ctttttacgg	6540
ttcctggcct	tttctgctgc	ttttgctcac	atgttcttct	ctgcgttctc	ccctgattct	6600

-continued

```

gtggataacc gtattaccgc ctttgagtga gctgataccg ctcgccgcag ccgaacgacc 6660
gagcgcagcg agtcagtgag cgaggaagcg gaagagcgcc caatacgcga accgcctctc 6720
cccgcgcggt ggccgattca ttaatgcagg gggatctcga tcccgcgaaa ttaatacgcg 6780
tcactatag 6789

```

```

<210> SEQ ID NO 13
<211> LENGTH: 9575
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Plasmid

```

```

<400> SEQUENCE: 13

```

```

acacaaaggc gcccaatcat ggattctata ttatcaaac agctgggtga caagactggt 60
tttgtagag tgccaatcaa gcattttgac tgtacaatgc taactctggc acttccaaca 120
tttgatgttt ccaagatggt agatagaatt accatagact tcaatctgga tgatatacaa 180
ggagcatctg aaataggctc aactttgcta ccctccatgt cgatagatgt ggaagatatg 240
gccaattttg ttcacgattt cacctttggc cacttagctg acaagactga cagactgtta 300
atgcgtgagt tcccattgat gaatgacggg ttgatcatt tgagccctga catgatcatt 360
aaaactacat ctggcatgta caacatcgtt gagttcacca cctttagggg agatgaaaga 420
ggtgcatctc aggtgccat gactaaactc gctaagtatg aggttccttg tgagaacaga 480
tctcagggca ggactgttgt tctttatggt gttagtgtct atcggcatgg tgtatggtct 540
aatctggagc tagaggactc tgaagcagag gagatggttt ataggtacag acttgctctt 600
agtgtgatgg atgagctaag gacctgttcc ccagaactgt catccacaga tgaggaacta 660
gggaagactg agagagagtt gctagccatg gtctcctcca tccaaataaa ttggtcagtc 720
acagaatctg tgtttccacc cttcagcaga gaaatgttg acaggtttag atcctcccct 780
cccgatccag agtatatcac gaggatagtg agcagatgcc tcataaattc tcaagagaaa 840
ctcatcaata gttccttctt tgctgaaggg aatgataagg ctctgagatt ttcaaaaaac 900
gctgaagagt gttccttggc agtagagaga gccttaaact agtatagagc agaagacaac 960
cttagggacc tcaatgacca caagtcaact attcagctgc ctcctggct gtctatcat 1020
gatgtcgatg gcaaatgctc gtgcccctct cagggttag atgtgagagg ggaccatccc 1080
atgtgcaact tgtggaggga agtggtcacc tctgcaaacc tagaggagat tgagaggatg 1140
cacgatgatg cagcagcaga acttgagttt gctctttcgg gagtaaagga caggccagat 1200
gagagaaaaca gataccatag agtccacctc aatatgggct cagatgatag tgtctacata 1260
gctgctttag gagttaatgg aaagaagcat aaagcagaca ctttagtgca acaaatgaga 1320
gacaggagta aacagccttt ctcccagac cacgatgtgg atcacatctc tgaatttctc 1380
tctgcatgct ctagtactgt gtggcaaca gatgaggacc tgtacaacc tctctctgt 1440
gataaagagc ttagattggc agcccagagg attcatcagc catccttctc agaaaggggt 1500
ttcaatgaga tcataacaga gcactacaaa ttcattggaa gtaggatagg ttcattggtc 1560
caaatgttca gcttgatagg agctgagcta tcagcttctg ttaacaaca tgtcaagcct 1620
aactactttg tgattaaacg actactaggt tctgggattt tcttgctaat caagcccact 1680
tccagcaaaa gccatatatt tgtgtctttt gcaattaagc gctcttctg ggcccttgat 1740

```

-continued

ctctccactt ccagggtttt caagccctac atagatgctg gggatctggt agttactgac	1800
tttgtttctt ataagctaag caagcttacc aacctctgca agtgcgtttc attaatggag	1860
tcctccttct cattctgggc agaagcattt ggcattccaa gotggaactt tgttggtgac	1920
ttgttcaggt cttcagactc tgcagcaatg gatgcctcat acatgggcaa actttcttta	1980
ttaacccttt tggaagacaa agcagcaact gaagagttac agactattgc aagatatata	2040
atcatggagg gctttgtctc gccccagaa atcccaaac ctcaaatgat gacctctaag	2100
tttcttaagg ttctcaggtc agagctgcag gtttacttat taaactgctt atgcagaact	2160
atccagagaa tagcaggtga gcccttcatt cttaagaaga aggatgggtc tatatcctgg	2220
ggtggcatgt tcaatccttt ttcagggcgt cactgcttg atatgcaacc actcatcagc	2280
tgttggtaca atggttactt taaaaataa gaagaagaga ctgagccttc gtccctttct	2340
gggatgtata agaaaaatc agaacttgag caccttagac cacagtcaga tgccttcttg	2400
ggttacaag atccagaact tcccagaatg catgagttca gtgtttccta cttgaaggag	2460
gcttgcaatc atgctaagct agtcttgagg agcctctatg gacagaattt catggagcag	2520
atagacaacc agattattcg agagctcagt gggttgactc tagaaaggtt gcccacactt	2580
aaggcccaa gcaactttaa tgagaattgg tatgtctata aggatgtagc agacaaaaac	2640
tacacaaggg ataaattatt agtgaagatg tcaaaatag cctctgaggg aaagagccta	2700
gctatccaga agtttgagga ttgtatgagg cagatagagt cacaaggatg catgcatatt	2760
tgtttgttta agaaacaaca gcatggaggt ctgagagaga tctatgtgat gggtgacagag	2820
gaaagaattg tcaatcgtg ggtggagaca atagccaggt ccatagggaa gttctttgct	2880
tctgataccc tctgtaaccc ccccaataa gtgaaaatc ctgagacaca tggcatcagg	2940
gcccgaagc aatgtaaggc gcctgtgtgg acttgtgcaa catcagatga tgcaaggaag	3000
tggaaccaag gccattttgt tacaagttt gccctcatgc tgtgtgagtt cacctctcct	3060
aatgggtggc cgctgatcat taggggatgc tcaatgttta ccaggaaaag gatgatgatg	3120
aatttgaatt atcttaagat cctggatggt catcgggagc ttgatattag agatgacttt	3180
gtgatggatc tottcaaagc ttatcatggc gaggcagaag ttccatgggc ctttaaggc	3240
aaaacatatt tggaaccac aacagggatg atgcagggaa tactgcatta tacttctca	3300
ctattacaca ccattcacca agaatacatc cggctcttgt cctttaagat attcaacctg	3360
aaggttgctc ctgagatgag caagggcctg gtttgtgaca tgatgcaagg atcagatgat	3420
agtagtatgc taatcagctt cccagctgat gatgagaagg ttcttaccag atgcaaagtg	3480
gccgcagcta tatgcttccg catgaagaag gagctgggag tgaacctgc catttacctc	3540
tcagagaagt ccacagcaaa cacagatttt gtgatggagt acaattctga attttatctc	3600
cacaccagc atgttagacc aacgatcagg tggattgcag cttgttgag cctgccagaa	3660
gtgaaacac tagtagcccg ccaggaagag gcctctaacc taatgacttc agttactgag	3720
ggaggtgggt cattctcctt agctgcaatg attcagcaag ctcaagtgcac tctccattac	3780
atgctgatgg gcattggagt gtctgagcta ttcttagagt ataagaaggc agtgcgaag	3840
tggaatgacc ctggcctggg tttcttctg cttgacaatc cttatgcgtg cggattggga	3900
ggtttcagat ttaatctctt caaagctatc accagaactg atttgcagaa gctatatgct	3960
ttcttcatga agaaggtcaa gggctcagct gctagggact gggcagatga agatgtcacc	4020

-continued

atcccagaaa cgtgtagcgt gagcccaggt ggcgctctaa ttcttagctc ctctctaaaag 4080
tggggatcta ggaagaagtt tcagaaattg agagaccggt tgaacatacc agagaactgg 4140
attgaaactaa taaatgagaa tccagaggtg ctctatcggg ctcccagaac aggccagaaa 4200
atattgttgc gcattgcaga gaaagtcocat agcccaggtg ttgtgtcatc attgtcttct 4260
ggcaatgcag tttgtaaagt catggcctca gctgtatact tcttatcagc aacaattttt 4320
gaggacactg gacgtcctga gttcaacttc ttggaggatt ctaagtacag cttgctacaa 4380
aagatggctg catattctgg ctttcatggt tttaatgata tggagccaga agatatatta 4440
ttcttattcc cgaatattga ggaattagaa tcaactggatt ctatagtta caacaaggga 4500
gaaatagaca tcatcccaag agtcaacatc agggatgcaa cccaaaccag ggtcactatc 4560
tttaatgagc agaagaccct ccggacatct ccagagaagt tgggtgcaga caagtggttt 4620
gggactcaga agagttagt agggcaaaaca accttctctg ctgaatggga aaagctaaaag 4680
aaaattgtaa agtggttgga agacactcca gaagcaactc tagctcacac cccactgaat 4740
aaccatattc aagttaggaa tttctttgct agaatggaaa gcaagcctag aacagtcaga 4800
ataacaggag ctccagtaaa gaagaggtca ggggttagta agatagctat ggttatccgt 4860
gacaatttct ccggatggg ccactctoga ggtgtagaag acctgtctgg ctccactcgt 4920
agtgtgtcag ctgaaattct caagcacttt ctattctgta tactacaagg tccatacagt 4980
gagagctata agctacagct aatctacaga gtcttaagct cagtgtcaaa cgttgagata 5040
aaggaatcag atggaagac aaaaaccaac ttgattggaa tccttcagag atttctagat 5100
ggtgatcacg ttgtcccat aattgaagag atgggagccg gaacagtggg tggattcatc 5160
aagagacaac aatctaaagt tgtgcagaac aaagtggctc attatggagt tgggatttgg 5220
agaggcttca tggatggata tcaggtccat cttagatag aaaatgacat aggcagccc 5280
ccaaggctta ggaatgtcac aactaactgt cagagcagcc catgggacct gagtattcca 5340
ataaggcaat gggcagaaga catgggggtc acaacaacc aggattattc ctctaaatct 5400
agcagagggg ccagatattg gatgcattca ttcaggatgc aaggacctag caagccattt 5460
ggatgcccag tttatattat taagggtgat atgtcagatg tcatcagact gagaaaggag 5520
gaggtggaga tgaagtacg gggctctact ctcaacttgt acaccaagca ccattctcat 5580
caggacctac acattctatc ttacactgca tcagacaatg atctcagtcc aggcattttc 5640
aagtcaatat cagatgaggg ggtggctcaa gcctgcaat tatttgagag ggagccaagc 5700
aactgctggg tgagatgtga gtctgtagcc ccaaaattta taccagccat ccttgagata 5760
tgtgagggga agagacagat aaggggaatt aacagaacca gactctcaga gatttgagaga 5820
atgtgtctg aatcttccct aagatcaaaa gtcggatcta tgttctcatt tgtcgccaat 5880
gtcagaggag cccatgatgt tgattatgat gcgttaatgg atctaagtat agaggatgcc 5940
aagaacaatg cattcagtcg tgttgttgac tgcataagat tggatgttag tggcccttac 6000
gagatggagt cttttgatac atctgatgtc aatctctttg ggccagccca ttacaaggac 6060
atcagttcat tatctatgat tgcctatccc ttaatggata agtttgttga ttatgctatt 6120
tctaagatgg ggagagcctc agttaggaaa gttctagaaa caggtcggtg ctccagcaaa 6180
gactatgatt tatcaaaggt tctcttcaga actctacaga gaccagaaga aagcattag 6240
atagatgatc tgaattata tgaggagaca gatgtggcgg atgacatgct aggctaagac 6300

-continued

caataagcaa agtcaggctt agatttaggg atactatgct agtattggaa tccatgtggg	6360
ttctgatact agcatagtg tacaatattg ggcggtcttt gtgtgggtcg gcatggcatc	6420
tccacctcct cgcggtccga cctgggcac cgaaggagga cgtcgtccac tcggatggct	6480
aaggagcaa gcttagcggc caccgctgag caataactag cataaccctc tggggcctct	6540
aaacgggtct tgaggggttt ttaagccga attcgtaatc atgcatagc tgttctctgt	6600
gtgaaattgt tatccgctca caattccaca caacatacga gccggaagca taaagtgtaa	6660
agcctggggt gcctaagtg tgagctaact cacattaatt gcgttgctgc cactgcccgc	6720
ttccagtcg gaaaacctgt cgtgccagct gcattaatga atcggccaac gcgcggggag	6780
aggcggtttg cgtattgggc gctcttcgc ttctcgtc actgactgc tgcgctcggc	6840
cgttcggctg cggcagcgg tatcagctca ctcaaaggcg gtaatacggc tatccacaga	6900
atcaggggat aacgcaggaa agaacatgtg agcaaaaggc cagcaaaagg ccaggaaccg	6960
taaaaaggcc gcgttgctgg cgtttttcca taggtccgc cccctgacg agcatcacia	7020
aatcgacgc tcaagtcaga ggtggcgaaa cccgacagga ctataaagat accagcgtt	7080
tcccctgga agctccctgc tgcgctctcc tgttccgacc ctgcccgtta ccggatacct	7140
gtccgccttt ctcccttcgg gaagcgtggc gctttctcat agctcagct gtaggtatct	7200
cagttcggtg taggtcgttc gctccaagct gggctgtgtg cacgaacccc ccgttcagcc	7260
cgaccgctgc gccttatccg gtaactatcg tcttgagtc aaccggtaa gacacgactt	7320
atgccactg gcagcagcca ctggtaacag gattagcaga gcgaggtatg taggcgggtg	7380
tacagagttc ttgaagtgtg ggcctaacta cggctacact agaagaacag tatttggtat	7440
ctgcgctctg ctgaagccag ttacctcgg aaaaagagtt ggtagctctt gatccggcaa	7500
acaaaccacc gctgtagcgt gtggtttttt tgtttgcaag cagcagatta cgcgcagaaa	7560
aaaaggatct caagaagatc tttgatctt ttctacgggg tctgacgctc agtggaaagc	7620
aaactcacgt taagggattt tggctatgag attatcaaaa aggatcttca cctagatcct	7680
tttaaatata aatgaagtt ttaaatcaat ctaaagtata tatgagtaaa cttggtctga	7740
cagttaccaa tgcttaatac gtgaggcacc tatctcagcg atctgtctat ttcggtcatc	7800
catagttgcc tgactcccc tcgtgtagat aactacgata cgggagggct taccatctgg	7860
ccccagtgc gcaatgatac cgcgagaccc acgctcaccg gctccagatt tatcagcaat	7920
aaaccagcca gccggaaggc ccgagcgcag aagtgtcct gcaactttat ccgcctocat	7980
ccagtctatt aattgttgc gggagctag agtaagtagt tcgccagtta atagtttgcg	8040
caacgttgtt gccattgcta caggcatcgt ggtgtcacgc tcgtcgtttg gtatggcttc	8100
atcagctcc ggttcccaac gatcaaggcg agttacatga tccccatgt tgtgcaaaaa	8160
agcggtagc tcttcggtc ctccgatcgt tgcagaagt aagttggccg cagtgttatc	8220
actcatggtt atggcagcac tgcataatc tcttactgtc atgccatccg taagatgctt	8280
ttctgtgact ggtgagtact caaccaagtc attctgagaa tagtgatgc ggcgaccgag	8340
ttgctcttgc ccggcgtcaa tacgggataa taccgcgcca catagcagaa ctttaaaagt	8400
gctcatcatt gaaaaacgtt ctccggggcg aaaactctca aggatcttac cgtgttgag	8460
atccagttcg atgtaacca ctctgcacc caactgatct tcagcatctt ttactttcac	8520
cagcgtttct gggtagcaa aaacaggaag gcaaaatgcc gcaaaaaagg gaataagggc	8580

-continued

```

gacacggaaa tgttgaatac tcatactcct cctttttcaa tattattgaa gcatttatca 8640
gggttattgt ctcatgagcg gatacatatt tgaatgtatt tagaaaaata aacaaatagg 8700
ggttccgcgc acatttcccc gaaaagtgcc acctgacgct taagaaacca ttattatcat 8760
gacattaacc tataaaaata ggcgatcac gaggccttt cgtctcgcgc gtttcgggta 8820
tgacggtgaa aacctctgac acatgcagct cccggagacg gtcacagctt gtctgtaagc 8880
ggatgccggg agcagacaag cccgtcaggg cgcgtcagcg ggtgttggcg ggtgtcgggg 8940
ctggcttaac tatgcgcat cagagcagat tgtactgaga gtgcaccatt cgacgctctc 9000
ccttatgcga ctctgcatt aggaagcagc ccagtagtag gttgaggccg ttgagaccg 9060
ccgcccgaag gaatggtgca tgcaaggaga tggcgcccaa cagtccccg gccacggggc 9120
ctgccacat acccacgccc aaacaagcgc tcatgagccc gaagtggcga gcccgatctt 9180
ccccatcggg gatgctggcg atataggcgc cagcaaccgc acctgtggcg cgggtgatgc 9240
cggccacgat gcgtcggcg tagaggatct ggctagcgat gacctgctg attggttctc 9300
tgaccatttc cgggtcggcg acggcgttac cagaaactca gaaggttcgt ccaaccaaac 9360
cgactctgac ggcagtttc gagagagatg atagggtctg cttcagtaag ccagatgcta 9420
cacaattagg cttgtacata ttgtcgttag aacgcggcta caattaatac ataacctat 9480
gtatcataca catacgattt aggtgacact atagaataca agctagcttg gctgcaggt 9540
cgacttctag aggatcctaa tacgactcac tatag 9575

```

```

<210> SEQ ID NO 14
<211> LENGTH: 4534
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic plasmid

```

```

<400> SEQUENCE: 14

```

```

acacaaagct ccctagagat acaaacacta ttacaataat ggacaactat caagagcttg 60
cgatccagtt tgctctcaa gcagtgacc gcaatgagat tgaacagtgg gtccgagagt 120
ttgcttatca agggtttgat gcccgtagag ttatcgaact cttaaagcag tatggtgggg 180
ctgactggga gaaggatgcc aagaaaatga ttgttctggc tctaactcgt ggcaacaagc 240
ccaggaggat gatgatgaaa atgtcgaaag aaggcaaagc aactgtggag gctctcatca 300
acaagtataa gctaaaggaa gggaaatcct cccgggatga gttgactcta tcacgagttg 360
ctgccgcctt ggttggctgg acatgccagg ctttggctct cttgagttag tggcttctg 420
tacttgggac taocatggac ggcctatccc ctgcataccc aaggcatatg atgcaccca 480
gctttgctgg catggtggat cctctctac caggagacta tctaagggca atattagatg 540
ctcactctct gtatctgctg cagttctccc gggatcaca cccaaacctc cgaggtagaa 600
caaaagagga gtttctgca acgttcacgc agccaatgaa tgcagcagtg aatagcaact 660
ttataagcca tgagaagagg agagaattct tgaagcctt tggacttctg gatttcaatg 720
ggaagccgct agctgctgct atggcagccg ctccagctta caagacagca gcctaagtgg 780
ctgccaggg ggttgggggg aaggggagtt ggggttacgc tgggatttg ggggtggggg 840
gtggggcagc cttaacctct aatcagatct ttacttctac agctcgtcca tgccgagagt 900
gatcccgcg gcggtcacga actccagcag gacctgtga tcgcgcttct cgttggggtc 960

```

-continued

tttgctcagg gcggaactggg tgcctcaggta gtgggtgtcg ggcagcagca cggggccgctc	1020
gccgatgggg gtgttctgct ggtagtggtc gccgagctgc acgctgccgt cctcgatgtt	1080
gtggcggatc ttgaagttca ccttgatgcc gttcttctgc ttgtcggcca tgatatagac	1140
gttgtggctg ttgtagttgt actccagctt gtgccccagg atggtgccgt cctccttgaa	1200
gtcgatgccc ttcagctcga tgcggttcac cagggtgtcg ccctcgaact tcacctcggc	1260
gcgggtcttg tagttgccgt cgtccttgaa gaagatggtg cgctcctgga cgtagccttc	1320
gggcatggcg gacttgaaga agtcgtgctg cttcatgtgg tggggtagc ggctgaagca	1380
ctgcacgccg taggtcaggg tggtcacgag ggtgggccag ggcacgggca gcttgccggt	1440
ggtgcagatg aacttcaggg tcagcttgcc gtagggtgca tgcacctcgc cctcgccgga	1500
cacgctgaac ttgtggccgt ttacgtcgcg gtccagctcg accaggatgg gcaccacccc	1560
ggtgaacagc tctcgcctc tgcctcaccat ggtaccgata tacttgataa gcactagggg	1620
gtctttgtgt gggtcggcat ggcactcoca cctcctcgcg gtccgacctg ggcattcgaa	1680
ggaggacgct gtcactcgg atggctaagg gagagctcgg atccggctgc taacaaagcc	1740
cgaaggaag ctgagttggc tgcctgccacc gctgagcaat aactagcata accccttggg	1800
gcctctaaac gggctttgag gggttttttg ctgaaaggag gaactatc cggatcgaga	1860
tcctctaggt acaagcctaa ttgtgtagca tctggcttac tgaagcagac cctatcatct	1920
ctctcgtaaa ctgccctcag agtcggtttg gttggacgaa cctctgagt ttctggtaac	1980
gccgtcccgc acccggaaat ggtcagcgaa ccaatcagca gggctcctgc tagccagatc	2040
ctctacgccg gacgcatcgt ggcggcctc accggcgcca cagggtgccgt tgcctggccc	2100
tatatcgccg acatcaccga tggggaagat cgggctcgc acttcgggct catgagcgt	2160
tgtttcggcg tgggtatggt ggcaggcccc gtggccgggg gactgttggg cgccatctcc	2220
ttgaccatt ccttgcggcg gcggtgctca acggcctcaa cctactactg ggtgcttcc	2280
taatgcagga gtcgcataag ggagagcgtc gatatggtgc actctcagta caatctgctc	2340
tgatgccgca tagttaagcc agccccgaca cccgccaaca cccgctgacg cgcctgacg	2400
ggcttgtctg ctcccggcat ccgcttacag acaagctgtg acaaagggcc tctgatac	2460
cctattttta taggttaatg tcatgataat aatggtttct tagacgtcag gtggcacttt	2520
tccgggaaat gtgcgaggaa cccctatttg tttatttttc taaatacatt caaatatgta	2580
tccgctcatg agacaataac cctgataaat gcttcaataa tattgaaaaa ggaagagtat	2640
gagtatcaaa catttccgtg tgcctcttat tccctttttt gggcattttt gccttccgtg	2700
ttttgctcac ccagaaacgc tggtgaaagt aaaagatgct gaagatcagt tgggtgcacg	2760
agtggtttac atcgaaactg atctcaacag cggttaagatc cttgagagtt ttcgccccga	2820
agaacgtttt ccaatgatga gcacttttaa agttctgcta tgtggcgcgg tattatcccc	2880
tattgacgcc gggcaagagc aactcggctc ccgcatacac tattctcaga atgacttggt	2940
tgagtactca ccagtcacag aaaagcatct tacggatggc atgacagtaa gagaattatg	3000
cagtgtgccc ataaccatga gtgataaac tgcggccaac ttacttctga caacgatcgg	3060
aggaccgaag gagctaaccg cttttttgca caacatgggg gatcatgtaa ctccgcttga	3120
tcggtgggaa ccggagctga atgaagccat accaaacgac gagcgtgaca ccacgatgcc	3180
tgtagcaatg gcaacaacgt tgcgcaaac attaactggc gaactactta ctctagcttc	3240

-continued

```

ccggcaacaa ttaatagact ggatggaggg ggataaagt gcaggaccac ttctgcgctc 3300
ggcccttcgg gctggctggg ttattgctga taaatctgga gccggtgagc gtgggtctcg 3360
cggatcatt gcagcactgg gccagatgg taagccctcc cgtatcgtag ttatctacac 3420
gacggggagt caggcaacta tggatgaacg aaatagacag atcgtgaga taggtgcctc 3480
actgattaag cattggtaac tgcagacca agttactca tatatacttt agattgattt 3540
aaaacttcat ttttaattta aaaggatcta ggtgaagatc ctttttgata atctcatgac 3600
caaaatccct taacgtgagt tttcgttcca ctgagcgtca gaccccgtag aaaagatcaa 3660
aggatcttct tgagatcctt tttttctgcg cgtaatctgc tgcttgcaaa caaaaaaac 3720
accgctacca ggggtggttt gtttgccgga tcaagagcta ccaactcttt ttccgaaggt 3780
aactggcttc agcagagcgc agataccaaa tactgtcctt ctagtgtagc cgtagttagg 3840
ccaccacttc aagaactctg tagcaccgcc tacatacctc gctctgctaa tectgttacc 3900
agtggctgct gccagtgggc ataagtcgtg tcttaccggg ttggactcaa gacgatagtt 3960
accggataag gcgcagcggg cgggctgaac ggggggttcg tgcacacagc ccagcttggg 4020
gcgaacgacc tacaccgaac tgagatacct acagcgtgag ctatgagaaa gcgccacgct 4080
tcccgaaggg agaaaaggcg acaggtatcc ggtaagcggc agggctcgaa caggagagcg 4140
cacgagggag ctccagggg gaaacgcctg gtatctttat agtctctgctg ggtttcgcca 4200
cctctgactt gagcgtcgat tttgtgatg ctctgaggg gggcggagcc tatggaaaaa 4260
cgccagcaac gcggcctttt tacggttcct ggctttttgc tggccttttg ctcacatgtt 4320
ctttcctgcg ttatcccttg atctgtgga taaccgtatt accgctttg agtgagctga 4380
taccgctcgc cgcagccgaa cgaccgagcg cagcagctca gtgagcgagg aagcggaaaga 4440
gcgccaata cgcaaacccg ctctcccgc gcggtggcgg attcattaat gcagggggat 4500
ctcgatcccg cgaaattaat acgactcact atag 4534

```

```

<210> SEQ ID NO 15
<211> LENGTH: 6652
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic plasmid

```

```

<400> SEQUENCE: 15

```

```

acacaaagac ggtgcattaa accatggcag ggattgcaat gacagtcctt ccagccttag 60
cagtttttgc tttggcacct gttgtttttg ctgaagacc ccatctcaga aacagaccag 120
ggaaggggca caactacatt gacgggatga ctcaggagga tgccacatgc aaacctgtga 180
catatgctgg gccatgtagc agttttgatg tcttgcttga aaagggaana tttccccttt 240
tccagtcgta tgctcatcat agaactctac tagaggcagt tcacgacacc atcattgcaa 300
aggctgatcc acctagctgt gaccttcaga gtgctcatgg gaaccctgc atgaaagaga 360
aactcgtgat gaagacacac tgtccaaatg actaccagtc agtcattac ctcaacaatg 420
acgggaaaat ggcttcagtc aagtgcctc ctaagtatga gctcactgag gactgcaact 480
ttttaggca gatgacaggt gctagcctga agaaggggct ttatcctctc caagacttgt 540
tttgcagtc aagtgaggat gatggatcaa aattaaaaac aaaaatgaaa ggggtctgcg 600
aagtgggggt tcaagcactc aaaaagtgtg atggccaact cagcactgca catgaggttg 660

```

-continued

tgccctttgc agtgtttaag aactcaaaga aggtttatct tgataagctt gaccttaaga	720
ctgaggagaa tctgctacca gactcatttg tctgtttcga gcataaggga cagtacaaag	780
gaacaatgga ctctggtcag actaagaggg agctcaaaag ctttgatata tctcagtgcc	840
ccaagattgg aggacatggt agtaagaagt gcaactggga cgcagcattt tgctctgctt	900
atgagtgcac tgctcagtac gccaatgcct attgttcaca tgctaattggg tcagggattg	960
tgcaagatac agtatcaggg gtctggaaga agcctttatg tgtagggatg gagagagtg	1020
ttgtgaagag agaactctct gccaaagcca tccagagagt tgagccttgc acaacttga	1080
taaccaaagtg tgagcctcat ggattggttg tccgatcaac agggttcaag atatcatcag	1140
cagttgcttg tgctagcggg gtttgctca caggatcgca gagtccttcc accgagatta	1200
cactcaagta tccagggata tcccagctct ctggggggga catagggggt cacatggcac	1260
acgatgatca gtcagtttag tccaaaatag tagctcactg ccctcccag gacctgtgtt	1320
tagtgcattg ctgcatagtg tgtgctcatg gcctgataaa ttaccagtgt cacactgctc	1380
tcagtgccct tgtgtttgtg tttgtattca gttctattgc aataattgt ttagctgttc	1440
tttataggt gcttaagtgc ctgaagattg ccccaaggaa agttctgaat ccaactaatg	1500
ggatcacagc cttcatcaga tggatatata agaagatggt tgccagagtg gcagacaaca	1560
ttaatcaagt gaacagggaa ataggatgga tggaaaggagg tcagttggtt ctagggaaac	1620
ctgcccctat tctcgtcat gccccaatcc cactgtatag cacatactg atgttattat	1680
tgattgtctc atatgcatca gcatgttcag aactgattca ggcaagctcc agaactacca	1740
cttgctctac agaggggtgtt aacaccaagt gtagactgtc tggcacagca ttgatcagag	1800
cagggtcagt tggggcagag gcttgtttga tgttgaaggg ggtcaaggaa gatcaacca	1860
agttcttaaa gataaaaaat gtctcaagtg agctatcatg cagggagggc cagagttatt	1920
ggactgggtc ctttagcctt aaatgtttga gctcaaggag atgccacctt gtcggggaat	1980
gccatgtgaa taggtgtctg tcttgagggg acaatgaaac ttcagcagag ttttcatttg	2040
ttggggaaag cagcaccatg cgagagaata agtgttttga gcaatgtgga ggatgggggt	2100
gtgggtgttt caatgtgaac ccatcttgcct tattttgtgca cacgtatctg cagtcagtta	2160
gaaaagaggc ccttagagtt tttaactgta tcgactgggt gcataaaactc actctagaga	2220
tcacagactt tgatggctct gtttcaacaa tagacttggg agcatcatct agccgtttca	2280
caaaactggg ttcagtttag ctctcactgg acgcagaggg catctcaggc tcaaatagct	2340
tttctttcat tgagagccca ggtaaaagggt atgcaattgt tgatgagcca ttctcagaaa	2400
ttcctcggca agggttcttg ggggagatca ggtgcaattc agagtcctca gtcctgagtg	2460
ctcatgaatc atgccttagg gcaccaaacc ttatctcata caagcccatg atagatcaat	2520
tggagtgcac aacaaactct attgatccct ttgtgtctt tgagaggggt tctctgccac	2580
agacaaggaa tgataaaacc tttgcagctt caaaaggaaa tagaggtgtt caagctttct	2640
ctaagggctc tgtacaagct gatctaactc tgatgtttga caattttgag gtggactttg	2700
tgggagcagc cgtatcttgt gatgcgcctt tcttaaattt gacaggttgc tattcttgc	2760
atgcaggggc cagggctctg ctgtctatca catccacagg aactggatct ctctctgccc	2820
acaataagga tgggtctctg catatagtcc ttccatcaga gaatggaaca aaagaccagt	2880
gtcagatact acacttctct gtgectgaag tagaggagga gtttatgtac tcttgtgatg	2940

-continued

gagatgagcg gcctctgttg gtgaagggga cctgatagc cattgatcca tttgatgata	3000
ggcgggaagc agggggggaa tcaacagttg tgaatccaaa atctggatct tggaaattct	3060
ttgactgggt ttctggactc atgagttggt ttggagggcc tcttaaaact atactcctca	3120
tttgccctgta tgttgcatca tcaattgggc tctttttcct ccttatatat cttggaagaa	3180
caggcctctc taaaatgtgg cttgctgcca ctaagaaggc ctcatagatc agtacgtgta	3240
aaagcaatat gttgaaataa gtagacacaa gcaaacctaa ttatgtaagt gttgtacaga	3300
taggtcaaat tattggaata tccaagctta gaaacttatg caataaact ttagatgtaa	3360
gcttagttgt aatttggggt ggtggggtga ggcagcagca gtctcaagt cttgtgaata	3420
ttctagtgg cgtaatcgtc ttttgccaga ttagctggga attaaactaa ctctttgaag	3480
ttgcaccggt ctttgtgtgg gtcggcatgg catctccacc tctctcggt cggacctggg	3540
catccgaagc aggcagctgt ccaactcggat ggctaaggga gcaagcttag cggtcaccgc	3600
tgagcaataa ctgacataac cccttggggc ctctaaacgg gtcttgagg gttttttaag	3660
ccgaattcgt aatcatgtca tagctgttcc ctgtgtgaaa ttgttatccg ctcacaattc	3720
cacacaacat acgagccgga agcataaagt gtaaagcctg ggggtcctaa tgagtgagct	3780
aactcacatt aattgcgttg cgctcactgc ccgctttcca gtcgggaaac ctgtcgtgcc	3840
agctgcatta atgaatcggc caacgcgcgg ggagagggcg tttgcgtatt gggcgctctt	3900
ccgcttctc gctcactgac tcgctgcgct cggctgttcg gctcggcgca gcggtatcag	3960
ctcactcaaa ggcgtaata cggttatcca cagaatcagg ggataacgca ggaagaaca	4020
tgtgagcaaa aggccagcaa aagccagga accgtaaaaa ggcgcgcttg ctggcgttt	4080
tccataggct ccgccccct gacgagcacc acaaaaatcg acgctcaagt cagaggtggc	4140
gaaacccgac aggactataa agataaccagg cgtttcccc tggaagctcc ctctgctgct	4200
ctctgttcc gacctgccc cttaccggat acctgtccc ctttctcct tgggaagcg	4260
tggcgcttcc tcatagctca cgctgtaggt atctcagttc ggtgtaggtc gttcgtcca	4320
agctgggctg tgtgacgaa cccccgctc agcccagcc ctgcccctta tccgtaact	4380
atcgtcttga gtccaaccg gtaagacacg acttatcgc actggcagca gccactggt	4440
acaggattag cagagcgagg tatgtaggcg gtgctacaga gttcttgaag tggtagccta	4500
actacggcta cactagaaga acagtatttg gtatctgccc tctgctgaag ccagttacct	4560
tcggaaaaag agttggtagc tcttgatccc gcaaaaaac caccgctggt agcggtggtt	4620
ttttgtttg caagcagcag attacgcgca gaaaaaagg atctcaagaa gatcctttga	4680
tctttctac ggggtctgac gctcagtgga acgaaaactc acgtaagggt attttggtca	4740
tgagattatc aaaaaggatc ttcacctaga tctttttaa ttaaaatga agttttaa	4800
caatctaaag tatatatgag taaacttgggt ctgacagtta ccaatgctta atcagtgagg	4860
cacctatctc agcgatctgt ctatttctgt catccatagt tgcctgactc cccgtcgtgt	4920
agataactac gatacgggag ggcttaccat ctggccccag tgetgcaatg ataccgagag	4980
accacgctc accgctcca gattatcag caataaacca gccagccgga agggccgagc	5040
gcagaagtgg tctgcaact ttatccgct ccacccagtc tattaattgt tgcgggaag	5100
ctagagtaag tagttcgcca gttaatagtt tgcgcaactg tgttgccatt gctacaggca	5160
tcgtggtgct acgctcgtcg tttggtatgg cttcattcag ctccggttcc caacgatcaa	5220

-continued

```

ggcgagttac atgatccccc atgttgtgca aaaaagcggg tagctccttc ggtcctccga 5280
tcgttgtcag aagtaagtgt gccgcagtgt tatcactcat ggttatggca gcaactgcata 5340
attctcttac tgcctatgcca tccgtaagat gcttttctgt gactggtgag tactcaacca 5400
agtcattctg agaatagtgt atgcggcgac cgagttgctc ttgcccgcg tcaatacggg 5460
ataataccgc gccacatagc agaactttaa aagtgtctcat cattggaaaa cgttcttcgg 5520
ggcgaaaaact ctcaaggatc ttaccgctgt tgagatccag ttcgatgtaa cccactcgtg 5580
caccacaactg atcttcagca tcttttactt tcaccagcgt ttctgggtga gcaaaaaacag 5640
gaaggcaaaa tgccgcaaaa aaggaataa gggcgacacg gaaatgtga atactcatac 5700
tcttcctttt tcaatattat tgaagcattt atcagggtta ttgtctcatg agcggataca 5760
tatttgaatg tatttagaaa aataa caaaa taggggttcc ggcacattt ccccgaaaag 5820
tgccacctga cgtctaagaa accattatta tcatgacatt aacctataaa aataggcgtg 5880
tcacgaggcc ctttctctc gcgcgttctg gtgatgacgg tgaaaacctc tgacacatgc 5940
agctcccgga gacggtcaca gcttgtctgt aagcggatgc cgggagcaga caagcccgtc 6000
agggcgcgtc agcgggtgtt ggcgggtgtc ggggctggct taactatgcg gcatcagagc 6060
agattgtact gagagtgcac cattcgaagc tctcccttat gogactcctg cattaggaag 6120
cagcccagta gtaggttgag gccggtgagc accgcccggc caaggaatgg tgcattgcaag 6180
gagatggcgc ccaacagtc cccggccacg gggcctgcca ccataccac gccgaaacaa 6240
gcgctcatga gcccgaaagt ggcagcccga tcttccccat cggatgatgc ggcgatatag 6300
gcgcccagaa ccgcacctgt ggcgcccgtg atgcccggca cgatgcgtcc ggcgtagagg 6360
atctggctag cgatgacct gctgattggt tcgctgacca ttccgggtg cgggacggcg 6420
ttaccagaaa ctcagaaggt tcgtccaacc aaaccgactc tgacggcagt ttacgagaga 6480
gatgataggg tctgcttcag taagccagat gctacacaat taggcttga catattgtcg 6540
ttagaacgcg gctacaatta atacataacc ttatgtatca tacacatagc atttaggtga 6600
cactatagaa tacaagcttg ggctgcaggt cgactaatac gactcactat ag 6652

```

1. A recombinant Rift Valley fever (RVF) virus, wherein the genome of the recombinant RVF virus comprises (i) a full-length L segment; (ii) a full-length M segment or an M segment comprising a complete deletion of the NSm open reading frame (ORF); and (iii) an S segment comprising a complete deletion of the NSs ORF.

2. The recombinant RVF virus of claim 1, wherein the genome is derived from RVF virus strain ZH501.

3. The recombinant RVF virus of claim 1, wherein the nucleotide sequence of the L segment is at least 95% identical to the nucleotide sequence of Genbank Accession No. DQ375406 (SEQ ID NO: 3).

4. (canceled)

5. The recombinant RVF virus of claim 1, wherein the nucleotide sequence of the M segment is at least 95% identical to the nucleotide sequence of Genbank Accession No. DQ380200 (SEQ ID NO: 2).

6. (canceled)

7. The recombinant RVF virus of claim 1, wherein the NSs ORF is replaced by the ORF of a reporter gene.

8. The recombinant RVF virus of claim 7, wherein the reporter gene encodes a green fluorescent protein.

9. An immunogenic composition comprising the recombinant RVF virus of claim 1 and a pharmaceutically acceptable carrier.

10. A method of immunizing a subject against RVF virus infection, comprising administering to the subject the immunogenic composition of claim 9.

11. The method of claim 10, wherein the subject is livestock.

12. The method of claim 10, wherein the subject is a human.

13-14. (canceled)

15. A collection of plasmids comprising:

(i) a plasmid encoding a full-length anti-genomic copy of the L segment of RVF virus;

(ii) a plasmid encoding a full-length anti-genomic copy of the M segment of RVF virus, or an anti-genomic copy of the M segment of RVF virus comprising a complete deletion of the NSm ORF; and

- (iii) a plasmid encoding an anti-genomic copy of the S segment of RVF virus, wherein the S segment comprises a complete deletion of the NSs ORF.
16. The collection of claim 15, wherein the plasmids further comprise a T7 promoter and a hepatitis delta virus ribozyme.
17. The collection of claim 15, wherein the NSs ORF is replaced by the ORF of a reporter gene.
18. The collection of claim 17, wherein the reporter gene encodes a green fluorescent protein.
19. The collection of claim 15, wherein the RVF virus is ZH501.
20. The collection of claim 15, wherein the nucleotide sequence of the S segment plasmid is at least 95% identical to the nucleotide sequence of SEQ ID NO: 14.
21. The collection of claim 15, wherein the nucleotide sequence of the M segment plasmid is at least 95% identical to the nucleotide sequence of SEQ ID NO: 12 or SEQ ID NO: 15.
22. (canceled)
23. The collection of claim 15, wherein the nucleotide sequence of the L segment plasmid is at least 95% identical to the nucleotide sequence of SEQ ID NO: 13.
24. An isolated host cell comprising the collection of claim 15.
25. The isolated host cell of claim 24, wherein the cell expresses T7 polymerase.
26. A method of preparing a recombinant RVF virus for immunization of a subject, comprising:
- (i) transfecting cultured cells with the collection of plasmids of claim 15;
 - (ii) incubating the cells for 1 to 5 days; and
 - (iii) collecting recombinant RVF virus from the cell supernatant.
27. The method of claim 26, wherein the cells express T7 polymerase.
28. A recombinant RVF virus, wherein the genome of the recombinant RVF virus comprises a full-length L segment, a full-length M segment and a full-length S segment, wherein the S segment further encodes a reporter gene.
29. The recombinant RVF virus of claim 28, wherein the S segment encodes a NSs-reporter gene fusion protein.
30. The recombinant RVF virus of claim 29, wherein the reporter gene is fused to the C-terminus of NSs.
- 31-33. (canceled)
34. A reverse genetics system for producing recombinant RVF virus consisting of three plasmids, wherein a first plasmid encodes an anti-genomic copy of a S segment; a second plasmid encodes an anti-genomic copy of a M segment and a third plasmid encodes an anti-genomic copy of a L segment of RVF virus, and wherein each plasmid comprises a T7 promoter and a hepatitis delta virus ribozyme.
35. The reverse genetics system of claim 34, wherein the RVF virus is ZH501.
36. The reverse genetics system of claim 34, wherein the S, M and L segments are wild-type S, M and L segments.
37. The reverse genetics system of claim 34, wherein the S segment comprises a deletion of the NSs ORF.
38. The reverse genetics system of claim 34, wherein the M segment comprises a deletion of the NSm ORF.
39. The reverse genetics system of claim 34, wherein the S segment comprises a deletion of the NSs ORF and the M segment comprises a deletion of the NSm ORF.
40. The reverse genetics system of claim 34, wherein the nucleotide sequence of the S segment plasmid is at least 95% identical to the nucleotide sequence of SEQ ID NO: 11 or SEQ ID NO: 14.
41. (canceled)
42. The reverse genetics system of claim 34, wherein the nucleotide sequence of the M segment plasmid is at least 95% identical to the nucleotide sequence of SEQ ID NO: 12 or SEQ ID NO: 15.
43. (canceled)
44. The reverse genetics system of claim 34, wherein the nucleotide sequence of the L segment plasmid is at least 95% identical to the nucleotide sequence of SEQ ID NO: 13.
45. A recombinant RVF virus produced using the reverse genetics system of claim 34.
- 46-47. (canceled)
48. A method of differentiating a subject administered the recombinant RVF virus of claim 1 from a subject naturally infected with RVF virus, comprising:
- obtaining a body fluid sample from the subject; and
 - detecting the presence or absence of anti-NSs and anti-NP antibodies in the sample, wherein the presence of both anti-NP and anti-NSs antibodies in the sample indicates that the subject was naturally infected with RVF virus, and wherein the presence of anti-NP antibodies and absence of anti-NSs antibodies in the sample indicates that the subject was administered the recombinant RVF virus of claim 1.
49. A method of differentiating a subject administered the recombinant RVF virus of claim 1, wherein the recombinant RVF virus comprises an M segment with a complete deletion of the NSm ORF, from a subject naturally infected with RVF virus, comprising:
- obtaining a body fluid sample from the subject; and
 - detecting the presence or absence of anti-NSm and anti-NP antibodies in the sample, wherein the presence of both anti-NP and anti-NSm antibodies in the sample indicates that the subject was naturally infected with RVF virus, and wherein the presence of anti-NP antibodies and absence of anti-NSm antibodies indicates that the subject was administered the recombinant RVF virus of claim 1.
50. The method of claim 49, wherein the body fluid sample is blood or serum.
51. The method of claim 49, wherein detecting the presence or absence of anti-NP or anti-NSm antibodies comprises an indirect fluorescent antibody (IFA) assay, an ELISA assay or a solid matrix-based differential detection assay.
52. The method of claim 48, wherein detecting the presence or absence of anti-NP or anti-NSs antibodies comprises an IFA assay, an ELISA assay or a solid matrix-based differential detection assay.
53. (canceled)
54. The method of claim 49, wherein the subject is livestock or the subject is human.
55. The method of claim 48, wherein the subject is livestock or the subject is human.
56. The method of claim 48, wherein the body fluid sample is blood or serum.

专利名称(译)	重组裂谷热 (RVF) 病毒及其使用方法		
公开(公告)号	US20110123567A1	公开(公告)日	2011-05-26
申请号	US12/809561	申请日	2008-12-16
[标]申请(专利权)人(译)	美國政府 健康与人类服务美分，疾病控制		
申请(专利权)人(译)	美利坚合众国政府是代表的部秘书 卫生和人类服务部，疾病控制和预防中心		
当前申请(专利权)人(译)	美利坚合众国政府是代表的部秘书 卫生和人类服务部，疾病控制和预防中心		
[标]发明人	BIRD BRIAN H ALBARINO CESAR G NICHOL STUART T KSIAZEK THOMAS G		
发明人	BIRD, BRIAN H. ALBARINO, CESAR G. NICHOL, STUART T. KSIAZEK, THOMAS G.		
IPC分类号	A61K39/12 C12N7/01 C12N15/63 C12N5/00 G01N33/566 G01N33/53		
CPC分类号	A61K39/00 A61K39/12 A61K2039/525 C12N2760/12234 C12N2760/12221 A61K35/76 A61K2039/5254 C12N7/00 C12N2760/12211 Y02A50/397		
优先权	61/016065 2007-12-21 US 61/042987 2008-04-07 US		
其他公开文献	US8673629		
外部链接	Espacenet USPTO		

摘要(译)

本文描述了包含在一种或多种病毒毒力基因 (例如NS和NSm) 中缺失的重组RVF病毒。使用基于质粒的反向遗传学系统产生的重组RVF病毒可以用作疫苗以预防家畜和人中RVF病毒的感染。如本文所述，重组RVF病毒生长至高滴度，在单次注射后提供保护性免疫，并允许在接种的动物和用野生型RVF病毒感染的动物之间进行分化。

FIG. 1

