




US 20170157246A1

(19) **United States**(12) **Patent Application Publication****Visvanathan et al.**(10) **Pub. No.: US 2017/0157246 A1**(43) **Pub. Date: Jun. 8, 2017**(54) **BIOMARKERS USEFUL IN THE
TREATMENT OF IL-23A RELATED
DISEASES**(86) PCT No.: **PCT/US2015/041706**

§ 371 (c)(1),

(2) Date: **Jan. 18, 2017**(71) Applicant: **BOEHRINGER INGELHEIM
INTERNATIONAL GMBH**, Ingelheim
am Rhein (DE)**Related U.S. Application Data**(60) Provisional application No. 62/028,379, filed on Jul.
24, 2014.(72) Inventors: **Sudha Visvanathan**, Ridgefiels, CT
(US); **Patrick BAUM**, Ingelheim am
Rhein (DE); **Oliver KLEINER**,
Ingelheim am Rhein (DE); **Ulrich
KUNZ**, Ingelheim am Rhein (DE)**Publication Classification**(51) **Int. Cl.****A61K 39/395** (2006.01)**G01N 33/53** (2006.01)(52) **U.S. Cl.**CPC **A61K 39/395** (2013.01); **G01N 33/53**
(2013.01); **G01N 33/5308** (2013.01); **A61K**
2121/00 (2013.01)(73) Assignee: **BOEHRINGER INGELHEIM
INTERNATIONAL GMBH**, Ingelheim
am Rhein (DE)(57) **ABSTRACT**

This invention generally relates to biomarkers useful in the treatment of IL-23 related diseases, in particular inflammatory diseases such as psoriasis. The invention also relates to methods of using the biomarkers disclosed herein, for example in therapies utilizing IL-23 antagonists.

(21) Appl. No.: **15/327,144**(22) PCT Filed: **Jul. 23, 2015**





FIG. 1

FIG. 2

FIG. 2A
FIG. 2B

FIG. 2A






FIG. 2B




FIG. 3

BIOMARKERS USEFUL IN THE TREATMENT OF IL-23A RELATED DISEASES

SEQUENCE LISTING

[0001] The instant application contains a Sequence Listing which has been submitted electronically in ASCII format and is hereby incorporated by reference in its entirety. Said ASCII copy, created on Jul. 16, 2015, is named 09-0632-WO-1_SL.txt and is 25,626 bytes in size.

TECHNICAL FIELD OF THE INVENTION

[0002] This invention generally relates to biomarkers useful in the treatment of IL-23A related diseases, in particular inflammatory diseases such as psoriasis. The invention also relates to methods of using the biomarkers disclosed herein, for example in therapies utilizing IL-23A antagonists.

BACKGROUND OF THE INVENTION

[0003] Psoriasis is a chronic, immune-mediated, inflammatory skin disease, with a global incidence of approximately 2%, associated with significant morbidity and can have a substantial psychosocial impact on quality of life and well-being of patients. Plaque psoriasis is the most common form and affects approximately 80-90% of patients, manifesting as raised plaques on the skin; the disease usually begins in late adolescence and early adulthood and may persist through adult life. The extent of the affected body surface area (BSA) and the degree of skin manifestations, including erythema, induration, and scaling, defines the severity of psoriasis with approximately 20-30% of patients having moderate-to-severe disease.

[0004] Psoriasis is a multifactorial disease, of unknown aetiology, with autoimmune and autoinflammatory components. Multiple genome-wide association studies have linked variants in the genes for the IL-23 receptor to psoriasis susceptibility. Human IL-23 is primarily produced by antigen presenting cells and induces differentiation of T helper 17 (Th17) cells. This results in the production of IL-17 and IL-22, which may mediate the development of the epidermal hyperplasia and tissue inflammation observed in psoriasis.

[0005] There is therefore a need for improved means to follow the efficacy of treatment options against psoriasis, identify patients that will most benefit from these treatments in Psoriasis and other inflammatory diseases, and to determine and adjust the dosages of therapies for patients as may be needed.

SUMMARY OF THE INVENTION

[0006] The present invention addresses the above needs and provides biomarkers useful in the treatment of IL-23A related diseases.

[0007] In one embodiment, the present invention provides a method for detecting the presence or absence of a beneficial response in a patient after administration of an IL-23A antagonist, comprising: a) obtaining a biological sample from the patient; b) measuring in said sample the level of expression of one or more biomarkers; c) comparing the level to control value of the level of the biomarkers; and d) determining whether or not the difference in levels between the sample and the control reflects a beneficial response in the patient, wherein the one or more biomarkers are genes/

proteins associated with the IL-23/IL-17 axis (for example one or more of IL-23A, IL-23R, IL-22, IL-22RA1 RA2, IL-17A, IL-17F, IL-17RA, and IL-17RC), keratinocyte and epithelial cell differentiation (for example one or more of late cornified envelope protein, transglutaminase 1, and cornifelin), tissue inflammation (for example one or more of β -defensin 2, neutrophil gelatinase lipocalin, and S-100 proteins) or the IFN α pathway (for example one or more of IFIH1, ISG15, IRF7, IF144, MX1, MX2, STAT1, and TRIM22).

[0008] In one embodiment, the presence or absence of a beneficial response in the patient is detected prior to and after administration of the IL-23A antagonist.

[0009] In one embodiment, the control value of the value of a patient treated with a placebo. In one embodiment, the control value of the value of a patient treated with a placebo and the difference is the difference between the sample from a patient treated with the IL-23A antagonist and the placebo.

[0010] In one embodiment, the level of the gene or the protein of said one or more biomarker is measured. In one embodiment, the patient suffers from psoriasis.

[0011] In one embodiment, the control value is calculated using samples from subjects that do not suffer from psoriasis. In one embodiment, the control value, for example the placebo value, is determined using samples from known psoriasis patients, for example from placebo-treated psoriasis patients. In one embodiment, the control value is determined using at least one previous sample taken from the patient.

[0012] In one embodiment, the one or more biomarkers is S-100A7, neutrophil gelatinase lipocalin or β -defensin 2. In one embodiment, the biological sample is a skin biopsy, blood, plasma or serum sample. In one embodiment, the IL-23A antagonist is an anti-IL-23A antibody or an antigen binding fragment thereof. In one embodiment, the levels of biomarkers are determined by RNA sequencing or ELISA or another protein assay.

[0013] In one embodiment, the method further comprises continuing the administration of the IL-23A antagonist to the patient if the difference in levels between the sample and the control reflects a beneficial response in the patient. In one embodiment, the method further comprises continuing the administration of the IL-23A antagonist to the patient if the difference in levels from a patient treated with the antagonist versus the placebo reflects a beneficial response in the patient.

[0014] In a further embodiment, the present invention provides a method of determining whether a potential therapeutic agent is efficacious in the treatment of psoriasis comprising: a) obtaining a first biological sample from a psoriasis patient prior to being treated with the potential therapeutic agent; b) treating the psoriasis patient with the potential therapeutic agent; c) obtaining a second biological sample from the psoriasis patient after being treated with the potential therapeutic agent; d) measuring in said first and second sample the levels of expression of one or more biomarkers; and e) comparing the biomarker levels in the second sample to the levels in the first sample, wherein lower biomarker levels in the second sample than in the first sample indicate that the potential therapeutic agent is efficacious, and further wherein the one or more biomarkers are genes associated with the IL-23/IL-17 axis (for example one or more of IL-23A, IL-23R, IL-22, IL-22RA1 RA2, IL-17A, IL-17F, IL-17RA, and IL-17RC), keratinocyte and epithelial

cell differentiation (for example one or more of late cornified envelope protein, transglutaminase 1, and cornifelin), tissue inflammation (for example one or more of β -defensin 2, neutrophil gelatinase lipocalin, and S-100 proteins) and the IFN α pathway (for example one or more of IFIH1, ISG15, IRF7, IF144, MX1, MX2, STAT1, and TRIM22). In one embodiment, said step e) comprises comparing the biomarker levels in the second sample to the levels in the first sample, wherein lower biomarker levels in the second sample than in the first sample and correlation with improvement in a clinical efficacy measure ie PASI scores in case of Psoriasis indicates the potential therapeutic agent is efficacious.

[0015] In one embodiment, the method further comprises continuing the treatment of the patient if biomarker levels in the second sample is lower than the levels in the first sample.

[0016] In a further embodiment, the present invention provides a method of treating psoriasis in a subject comprising: a) determining whether to initiate treatment of the subject, modify the treatment dose, modify the dosing interval, or discontinue treatment, based on the method of any of the preceding claims; and b) modifying the treatment regimen based on the determination.

[0017] In a further embodiment, the present invention provides a method of monitoring patient response to a psoriasis treatment comprising:

[0018] a) obtaining a first biological sample from the patient;

[0019] b) measuring the level of one or more biomarkers in said first biological sample, wherein said one or more biomarkers are genes/proteins associated with the IL-23/IL-17 axis (for example one or more of IL-23A, IL-23R, IL-22, IL-22RA1 RA2, IL-17A, IL-17F, IL-17RA, and IL-17RC), keratinocyte and epithelial cell differentiation (for example one or more of late cornified envelope protein, transglutaminase 1, and cornifelin), tissue inflammation (for example one or more of β -defensin 2, neutrophil gelatinase lipocalin, and S-100 proteins) and the IFN α pathway (for example one or more of IFIH1, ISG15, IRF7, IF144, MX1, MX2, STAT1, and TRIM22);

[0020] c) administering a treatment compound to the patient;

[0021] d) obtaining a second biological sample from the patient;

[0022] e) measuring the level of said one or more biomarkers in said second biological sample; and

[0023] f) comparing the levels of the one or more biomarkers obtained from first and second biological samples;

[0024] wherein a decreased level of the one or more biomarkers in the second biological sample indicates an effective response. In one aspect, a decreased level of the one or more biomarkers in the second biological sample and correlation with improvement in a clinical efficacy measure ie PASI scores in case of Psoriasis indicates an effective response.

[0025] In a further embodiment, the present invention provides a method for monitoring patient compliance with a drug treatment protocol for psoriasis comprising:

[0026] a) obtaining a biological sample from said patient;

[0027] b) measuring the level of one or more biomarkers, wherein the one or more biomarkers are genes/

proteins associated with the IL-23/IL-17 axis (for example one or more of IL-23A, IL-23R, IL-22, IL-22RA1 RA2, IL-17A, IL-17F, IL-17RA, and IL-17RC), keratinocyte and epithelial cell differentiation (for example one or more of late cornified envelope protein, transglutaminase 1, and cornifelin), tissue inflammation (for example one or more of β -defensin 2, neutrophil gelatinase lipocalin, and S-100 proteins) and the IFN α pathway (for example one or more of IFIH1, ISG15, IRF7, IF144, MX1, MX2, STAT1, and TRIM22); and

[0028] c) determining if the level is decreased in the patient sample compared to the level in a control untreated sample;

[0029] wherein a decreased level indicates patient compliance with said drug treatment protocol.

[0030] In one embodiment, in any one of the methods above, the level of the one or more biomarkers in the second biological sample is decreased by at least about 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 65%, 70%, 75%, 80%, 85%, 90%, or 95% or more as compared to the level in the first biological sample.

[0031] In one embodiment, in any one of the methods above, the biological sample is a skin biopsy, blood, plasma or serum sample. In one embodiment, in any one of the methods above, the IL-23A antagonist is an anti-IL-23A antibody or an antigen binding fragment thereof. In one embodiment, in any one of the methods above, the levels of biomarkers are determined by RNA sequencing or ELISA or another protein assay.

[0032] In one embodiment, the present invention further provides a method of selecting a patient, for example using a method disclosed herein. In one embodiment, the present invention further provides a method of enriching a patient population for patients expected to have a beneficial response after treatment with an IL-23 antagonist, for example using a method of the present invention. In one embodiment, the present invention further provides a method of enriching a patient population for patients expected to have a beneficial response prior to or early after treatment with an IL-23 antagonist, for example using a method of the present invention.

[0033] In a further embodiment, the present invention further provides an ELISA kit comprising one or more antibodies or antigen binding fragments thereof that specifically bind to one or more biomarkers, wherein the one or more biomarkers is ie S-100 proteins, neutrophil gelatinase lipocalin or β -defensin 2. In one embodiment, the ELISA or other protein assay kit further comprises instructions for use of the kit prior to treatment or for monitoring psoriasis.

[0034] In one embodiment, in any one of the methods above, the anti-IL-23A antibody or an antigen binding fragment thereof as disclosed below.

[0035] In one embodiment, the IL-23A antagonist is an anti-IL-23p19 antibody or antigen-binding fragment thereof, wherein the antibody or antigen-binding fragment thereof comprises a light chain variable region comprising the amino acid sequence of SEQ ID NO:1 (CDR1-L); the amino acid sequence of SEQ ID NO:2 (CDR2-L); and the amino acid sequence of SEQ ID NO:3 (CDR3-L); and a heavy chain variable region comprising the amino acid sequence of SEQ ID NO: 4, 7, 8 or 9 (CDR1-H); the amino acid sequence of SEQ ID NO:5 (CDR2-H); and the amino acid sequence of SEQ ID NO:6 (CDR3-H).

[0036] In one embodiment, the IL-23A antagonist is an anti-IL-23p19 or antigen-binding fragment thereof antibody, wherein the antibody or antigen-binding fragment thereof comprises a light chain variable region comprising the amino acid sequence of SEQ ID NO:1 (CDR1-L); the amino acid sequence of SEQ ID NO:2 (CDR2-L); and the amino acid sequence of SEQ ID NO:3 (CDR3-L); and a heavy chain variable region comprising the amino acid sequence of SEQ ID NO:4 (CDR1-H); the amino acid sequence of SEQ ID NO:5 (CDR2-H); and the amino acid sequence of SEQ ID NO:6 (CDR3-H).

[0037] In one embodiment, the IL-23A antagonist is an anti-IL-23p19 or antigen-binding fragment thereof antibody, wherein the antibody or antigen-binding fragment thereof comprises a light chain variable region comprising the amino acid sequence of SEQ ID NO:1 (CDR1-L); the amino acid sequence of SEQ ID NO:2 (CDR2-L); and the amino acid sequence of SEQ ID NO:3 (CDR3-L); and a heavy chain variable region comprising the amino acid sequence of SEQ ID NO:7 (CDR1-H); the amino acid sequence of SEQ ID NO:5 (CDR2-H); and the amino acid sequence of SEQ ID NO:6 (CDR3-H).

[0038] In one embodiment, the IL-23A antagonist is an anti-IL-23p19 or antigen-binding fragment thereof antibody, wherein the antibody or antigen-binding fragment thereof comprises a light chain variable region comprising the amino acid sequence of SEQ ID NO:1 (CDR1-L); the amino acid sequence of SEQ ID NO:2 (CDR2-L); and the amino acid sequence of SEQ ID NO:3 (CDR3-L); and a heavy chain variable region comprising the amino acid sequence of SEQ ID NO:8 (CDR1-H); the amino acid sequence of SEQ ID NO:5 (CDR2-H); and the amino acid sequence of SEQ ID NO:6 (CDR3-H).

[0039] In one embodiment, the IL-23A antagonist is an anti-IL-23p19 or antigen-binding fragment thereof antibody, wherein the antibody or antigen-binding fragment thereof comprises a light chain variable region comprising the amino acid sequence of SEQ ID NO:1 (CDR1-L); the amino acid sequence of SEQ ID NO:2 (CDR2-L); and the amino acid sequence of SEQ ID NO:3 (CDR3-L); and a heavy chain variable region comprising the amino acid sequence of SEQ ID NO:9 (CDR1-H); the amino acid sequence of SEQ ID NO:5 (CDR2-H); and the amino acid sequence of SEQ ID NO:6 (CDR3-H).

[0040] In one embodiment, the IL-23A antagonist is an anti-IL-23p19 antibody or antigen-binding fragment thereof, wherein the antibody or antigen-binding fragment thereof comprises a light chain variable region comprising the amino acid sequence of any one of SEQ ID NO:10, 11, 12 or 13; and a heavy chain variable region comprising the amino acid sequence any one of SEQ ID NO:14, 15, 16 or 17.

[0041] In one embodiment, the IL-23A antagonist is an anti-IL-23p19 antibody or antigen-binding fragment thereof, wherein the antibody or antigen-binding fragment thereof comprises a light chain variable region comprising the amino acid sequence of SEQ ID NO:11 and a heavy chain variable region comprising the amino acid sequence SEQ ID NO:14.

[0042] In one embodiment, the IL-23A antagonist is an anti-IL-23p19 antibody or antigen-binding fragment thereof, wherein the antibody or antigen-binding fragment thereof comprises a light chain variable region comprising the

amino acid sequence of SEQ ID NO:11 and a heavy chain variable region comprising the amino acid sequence SEQ ID NO:15.

[0043] In one embodiment, the IL-23A antagonist is an anti-IL-23p19 antibody or antigen-binding fragment thereof, wherein the antibody or antigen-binding fragment thereof comprises a light chain variable region comprising the amino acid sequence of SEQ ID NO:10 and a heavy chain variable region comprising the amino acid sequence SEQ ID NO:14.

[0044] In one embodiment, the IL-23A antagonist is an anti-IL-23p19 antibody or antigen-binding fragment thereof, wherein the antibody or antigen-binding fragment thereof comprises a light chain variable region comprising the amino acid sequence of SEQ ID NO:10 and a heavy chain variable region comprising the amino acid sequence SEQ ID NO:15.

[0045] In one embodiment, the anti-IL-23p19 antibody is a humanized antibody. In one embodiment, the anti-IL-23p19 antibody is a monoclonal antibody. In one embodiment, the anti-IL-23p19 antibody is a full length antibody. In one embodiment, the anti-IL-23p19 antibody is a humanized monoclonal anti-IL-23p19 antibody, for example a full length humanized monoclonal antibody. In one embodiment, the antigen-binding fragment is a Fab, F(ab)₂, or single chain Fv fragment. In one embodiment, the antigen-binding fragment comprises a light chain variable region and a heavy chain variable region.

[0046] In one embodiment, the IL-23A antagonist is an antibody comprising the amino acid sequence SEQ ID NO:14 or 15 linked to a human IgG1, IgG2, IgG3, IgG4, IgM, IgA or IgE heavy chain constant region. An antibody comprising the amino acid sequence of SEQ ID NO: 14 or 15 linked to a human IgG1 heavy chain constant region. An antibody comprising the amino acid sequence of SEQ ID NO:10 or 11 linked to a human kappa or lambda light chain constant region. An antibody comprising the amino acid sequence of SEQ ID NO: 10 or 11 linked to a human kappa light chain constant region.

[0047] In one embodiment, the IL-23A antagonist is an antibody comprising the amino acid sequence of SEQ ID NO:14 or 15 linked to a human IgG1 heavy chain constant region; and the amino acid sequence of SEQ ID NO: 10 or 11 linked to a human kappa light chain constant region.

[0048] In one embodiment, the IL-23A antagonist is a humanized monoclonal anti-IL-23p19 antibody comprising a light chain variable region comprising the amino acid sequence selected from the group consisting of any one of SEQ ID NO:10, 11, 12 and 13 and a heavy chain variable region comprising the amino acid sequence selected from the group consisting of any one of SEQ ID NO:14, 15, 16 and 17.

[0049] In one embodiment, the IL-23A antagonist is a humanized monoclonal anti-IL-23p19 antibody comprising a light chain variable region comprising the amino acid sequence of SEQ ID NO:11 and a heavy chain variable region comprising the amino acid sequence SEQ ID NO:14.

[0050] In one embodiment, the IL-23A antagonist is a humanized monoclonal anti-IL-23p19 antibody comprising a light chain variable region comprising the amino acid sequence of SEQ ID NO:11 and a heavy chain variable region comprising the amino acid sequence SEQ ID NO:15.

[0051] In one embodiment, the IL-23A antagonist is a humanized monoclonal anti-IL-23p19 antibody comprising

a light chain variable region comprising the amino acid sequence of SEQ ID NO:10 and a heavy chain variable region comprising the amino acid sequence SEQ ID NO:14.

[0052] In one embodiment, the IL-23A antagonist is a humanized monoclonal anti-IL-23p19 antibody comprising a light chain variable region comprising the amino acid sequence of SEQ ID NO:10 and a heavy chain variable region comprising the amino acid sequence SEQ ID NO:15.

[0053] In one embodiment, the IL-23A antagonist is a humanized monoclonal anti-IL-23p19 antibody comprising a light chain comprising the amino acid sequence of SEQ ID NO:18 or 21 and a heavy chain comprising the amino acid sequence of SEQ ID NO:19 or 20.

[0054] In one embodiment, the IL-23A antagonist is a humanized monoclonal anti-IL-23p19 antibody comprising a light chain comprising the amino acid sequence of SEQ ID NO:18 and a heavy chain comprising the amino acid sequence of SEQ ID NO:19.

[0055] In one embodiment, the IL-23A antagonist is a humanized monoclonal anti-IL-23p19 antibody comprising a light chain comprising the amino acid sequence of SEQ ID NO:18 and a heavy chain comprising the amino acid sequence of SEQ ID NO:20.

[0056] In one embodiment, the IL-23A antagonist is a humanized monoclonal anti-IL-23p19 antibody comprising a light chain comprising the amino acid sequence of SEQ ID NO:21 and a heavy chain comprising the amino acid sequence of SEQ ID NO:19.

[0057] In one embodiment, the IL-23A antagonist is a humanized monoclonal anti-IL-23p19 antibody comprising a light chain comprising the amino acid sequence of SEQ ID NO: 21 and a heavy chain comprising the amino acid sequence of SEQ ID NO: 20.

[0058] In one embodiment, the IL-23A antagonist is Antibody A, Antibody B, Antibody C or Antibody D.

[0059] In one embodiment, the IL-23A antagonist is an antibody as disclosed in WO2007/005955, WO2007/024846, WO2007/027714, WO2007/076524, WO2008/103432 or WO2012/061448.

BRIEF DESCRIPTION OF THE FIGURES

[0060] FIG. 1: Mean improvements in PASI score for patients receiving IV or SC Antibody A. For Antibody A IV, N=24 to Week 12 and N=23 thereafter (patient, originally allocated to the Antibody A 5 mg/kg group, lost to follow up due to relocation and discontinued after Week 12). For Antibody A SC, N=15 throughout. IV, intravenous; PASI, psoriasis area severity index; SC, subcutaneous.

[0061] FIG. 2: Heat Map showing overall decreases in fold change expression (change from Baseline to Week 8) in a cluster of 79 genes from RNA-seq analysis post-treatment with Antibody A versus placebo with each column representing individual patients. The scale shows log 2-fold change in expression of each of the 79 genes with red representing fold-change increase and blue fold-change decrease (Antibody A versus placebo).

[0062] FIG. 3: Correlation plot of the cluster means of the 79 genes (from FIG. 2) for each patient versus PASI scores at Week 8, showing a significant correlation between reductions in expression of this gene cluster in Antibody A treated patients and lower PASI scores [Antibody A (dots) versus placebo (crosses)].

DETAILED DESCRIPTION

[0063] The p19 subunit of IL-23 (also referred to herein as “IL-23A”, “IL-23p19” and “p19 subunit”) is a 189 amino acid polypeptide containing a 21 aa leader sequence (Oppmann et al. *Immunity* 13:715 (2000), SEQ ID NO: 22). The biological activity of the molecule is only detected when it is partnered with the IL-12p40 subunit to form IL-23. IL-23 is predominantly expressed by activated dendritic cells (DCs) and phagocytic cells. The receptor for IL-23 was found to be composed of the IL-12Rβ1 subunit of IL-12 receptor partnered with a unique subunit called IL-23R (Parham et al. *J. Immunol.* 168:5699 (2002)). Expression of the receptor is detected primarily on memory T cells and NK cells. Thus, expression of this cytokine:receptor pair appears to be restricted to specific populations of immune cells. While it was first thought that IL-12 and IL-23 would share many functions, the data has shown the picture to be different. Whereas IL-12 has a predominant role in the production of Th1 cells, IL-23 was found to be critically involved in the production and maintenance of a recently recognized Th cell subset termed Th17 (Kikly et al. *Curr. Opin. Immunol.* 18:670 (2006), Kastelein et al. *Ann. Rev. Immunol.* 25:221 (2007)). These cells produce IL-17A, IL-17F, IL-22 and other pro-inflammatory cytokines such as IL-6 and TNF-α. As described below, animal model studies on the role of these Th17 cells show their importance as a driving force in chronic inflammation and autoimmunity.

```
SEQ ID NO: 22:
mlgsravmll lllpwtacgr avpggsspaw tqcqqslsqkl
ctlawisahpl vghmdlreeg deetndvph iqcgdgcdpqr
glrdnsqfcl qrihqglify eklldsfift gepslldpds
vgqlhasllg lsqllqpegh hwetqqipsl spsqpwqrll
lrfkilsrlq afvavaarvf ahgaatlsp
```

[0064] The present invention provides biomarkers useful in the treatment of IL-23A related diseases.

[0065] In one embodiment, the present invention provides a method for detecting the presence or absence of a beneficial response in a patient after administration of an IL-23A antagonist, comprising: a) obtaining a biological sample from the patient; b) measuring in said sample the level of expression of one or more biomarkers; c) comparing the level to control value of the level of the biomarkers; and d) determining whether or not the difference in levels between the sample and the control reflects a beneficial response in the patient, wherein the one or more biomarkers are genes/proteins associated with the IL-23/IL-17 axis (for example one or more of IL-23A, IL-23R, IL-22, IL-22RA1 RA2, IL-17A, IL-17F, IL-17RA, and IL-17RC), keratinocyte and epithelial cell differentiation (for example one or more of late cornified envelope protein, transglutaminase 1, and cornifelin), tissue inflammation (for example one or more of β-defensin 2, neutrophil gelatinase lipocalin, and S-100 proteins) or the IFNα pathway (for example one or more of IFIH1, ISG15, IRF7, IF144, MX1, MX2, STAT1, and TRIM22).

[0066] In one embodiment, the presence or absence of a beneficial response in the patient is detected prior to and after administration of the IL-23A antagonist.

[0067] In one embodiment, the control value of the value of a patient treated with a placebo. In one embodiment, the control value of the value of a patient treated with a placebo and the difference is the difference between the sample from a patient treated with the IL-23A antagonist and the placebo.

[0068] In one embodiment, the level of the gene or the protein of said one or more biomarker is measured. In one embodiment, the patient suffers from psoriasis.

[0069] In one embodiment, the control value is calculated using samples from subjects that do not suffer from psoriasis. In one embodiment, the control value, for example the placebo value, is determined using samples from known psoriasis patients, for example from placebo-treated psoriasis patients. In one embodiment, the control value is determined using at least one previous sample taken from the patient.

[0070] In one embodiment, the one or more biomarkers is S-100A7, neutrophil gelatinase lipocalin or β -defensin 2. In one embodiment, the biological sample is a skin biopsy, blood, plasma or serum sample. In one embodiment, the IL-23A antagonist is an anti-IL-23A antibody or an antigen binding fragment thereof. In one embodiment, the levels of biomarkers are determined by RNA sequencing or ELISA or another protein assay.

[0071] In one embodiment, the method further comprises continuing the administration of the IL-23A antagonist to the patient if the difference in levels between the sample and the control reflects a beneficial response in the patient. In one embodiment, the method further comprises continuing the administration of the IL-23A antagonist to the patient if the difference in levels from a patient treated with the antagonist versus the placebo reflects a beneficial response in the patient.

[0072] In a further embodiment, the present invention provides a method of determining whether a potential therapeutic agent is efficacious in the treatment of psoriasis comprising: a) obtaining a first biological sample from a psoriasis patient prior to being treated with the potential therapeutic agent; b) treating the psoriasis patient with the potential therapeutic agent; c) obtaining a second biological sample from the psoriasis patient after being treated with the potential therapeutic agent; d) measuring in said first and second sample the levels of expression of one or more biomarkers; and e) comparing the biomarker levels in the second sample to the levels in the first sample, wherein lower biomarker levels in the second sample than in the first sample indicate that the potential therapeutic agent is efficacious, and further wherein the one or more biomarkers are genes associated with the IL-23/IL-17 axis (for example one or more of IL-23A, IL-23R, IL-22, IL-22RA1 RA2, IL-17A, IL-17F, IL-17RA, and IL-17RC), keratinocyte and epithelial cell differentiation (for example one or more of late cornified envelope protein, transglutaminase 1, and cornifelin), tissue inflammation (for example one or more of β -defensin 2, neutrophil gelatinase lipocalin, and S-100 proteins) and the IFN α pathway (for example one or more of IFIH1, ISG15, IRF7, IF144, MX1, MX2, STAT1, and TRIM22). In one embodiment, said step e) comprises comparing the biomarker levels in the second sample to the levels in the first sample, wherein lower biomarker levels in the second sample than in the first sample and correlation with improvement in a clinical efficacy measure ie PASI scores in case of Psoriasis indicates the potential therapeutic agent is efficacious.

[0073] In one embodiment, the method further comprises continuing the treatment of the patient if biomarker levels in the second sample is lower than the levels in the first sample.

[0074] In a further embodiment, the present invention provides a method of treating psoriasis in a subject comprising: a) determining whether to initiate treatment of the subject, modify the treatment dose, modify the dosing interval, or discontinue treatment, based on the method of any of the preceding claims; and b) modifying the treatment regimen based on the determination.

[0075] In a further embodiment, the present invention provides a method of monitoring patient response to a psoriasis treatment comprising:

[0076] a) obtaining a first biological sample from the patient;

[0077] b) measuring the level of one or more biomarkers in said first biological sample, wherein said one or more biomarkers are genes/proteins associated with the IL-23/IL-17 axis (for example one or more of IL-23A, IL-23R, IL-22, IL-22RA1 RA2, IL-17A, IL-17F, IL-17RA, and IL-17RC), keratinocyte and epithelial cell differentiation (for example one or more of late cornified envelope protein, transglutaminase 1, and cornifelin), tissue inflammation (for example one or more of β -defensin 2, neutrophil gelatinase lipocalin, and S-100 proteins) and the IFN α pathway (for example one or more of IFIH1, ISG15, IRF7, IF144, MX1, MX2, STAT1, and TRIM22);

[0078] c) administering a treatment compound to the patient;

[0079] d) obtaining a second biological sample from the patient;

[0080] e) measuring the level of said one or more biomarkers in said second biological sample; and

[0081] f) comparing the levels of the one or more biomarkers obtained from first and second biological samples;

[0082] wherein a decreased level of the one or more biomarkers in the second biological sample indicates an effective response. In one aspect, a decreased level of the one or more biomarkers in the second biological sample and correlation with improvement in a clinical efficacy measure ie PASI scores in case of Psoriasis indicates an effective response.

[0083] In a further embodiment, the present invention provides a method for monitoring patient compliance with a drug treatment protocol for psoriasis comprising:

[0084] a) obtaining a biological sample from said patient;

[0085] b) measuring the level of one or more biomarkers, wherein the one or more biomarkers are genes/proteins associated with the IL-23/IL-17 axis (for example one or more of IL-23A, IL-23R, IL-22, IL-22RA1 RA2, IL-17A, IL-17F, IL-17RA, and IL-17RC), keratinocyte and epithelial cell differentiation (for example one or more of late cornified envelope protein, transglutaminase 1, and cornifelin), tissue inflammation (for example one or more of β -defensin 2, neutrophil gelatinase lipocalin, and S-100 proteins) and the IFN α pathway (for example one or more of IFIH1, ISG15, IRF7, IF144, MX1, MX2, STAT1, and TRIM22); and

[0086] c) determining if the level is decreased in the patient sample compared to the level in a control untreated sample;

[0087] wherein a decreased level indicates patient compliance with said drug treatment protocol.

[0088] In one embodiment, in any one of the methods above, the level of the one or more biomarkers in the second biological sample is decreased by at least about 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 65%, 70%, 75%, 80%, 85%, 90%, or 95% or more as compared to the level in the first biological sample.

[0089] In one embodiment, in any one of the methods above, the biological sample is a skin biopsy, blood, plasma or serum sample. In one embodiment, in any one of the methods above, the IL-23A antagonist is an anti-IL-23A antibody or an antigen binding fragment thereof. In one embodiment, in any one of the methods above, the levels of biomarkers are determined by RNA sequencing or ELISA or another protein assay.

[0090] In one embodiment, the present invention further provides a method of selecting a patient, for example using a method disclosed herein. In one embodiment, the present invention further provides a method of enriching a patient population for patients expected to have a beneficial response after treatment with an IL-23 antagonist, for example using a method of the present invention. In one embodiment, the present invention further provides a method of enriching a patient population for patients expected to have a beneficial response prior to or early after treatment with an IL-23 antagonist, for example using a method of the present invention.

[0091] In a further embodiment, the present invention further provides an ELISA kit comprising one or more antibodies or antigen binding fragments thereof that specifically bind to one or more biomarkers, wherein the one or more biomarkers is ieS-100 proteins, neutrophil gelatinase lipocalin or β -defensin 2. In one embodiment, the ELISA or other protein assay kit further comprises instructions for use of the kit prior to treatment or for monitoring psoriasis.

[0092] In one embodiment, in any one of the methods above, the anti-IL-23A antibody or an antigen binding fragment thereof as disclosed below.

[0093] In one aspect, the anti-IL-23p19 antibody is a humanized antibody. In one aspect, the anti-IL-23p19 antibody is a monoclonal antibody. In one aspect, the anti-IL-23p19 antibody is a full length antibody. In one aspect, the anti-IL-23p19 antibody is a humanized monoclonal antibody, for example a full length humanized monoclonal antibody.

[0094] An antibody or antigen-binding fragment thereof of the present invention recognizes specific "IL-23p19 antigen epitope" or "IL-23p19 epitope". As used herein these terms refer to a molecule (e.g., a peptide) or a fragment of a molecule capable of immunoreactivity with an anti-IL-23p19 antibody and, for example, include an IL-23p19 antigenic determinant recognized by the any of the antibodies having a light chain/heavy chain sequence combination of SEQ ID NO:11/14, 11/15, 10/14 or 10/15.

[0095] The generalized structure of antibodies or immunoglobulin is well known to those of skill in the art. These molecules are heterotetrameric glycoproteins, typically of about 150,000 daltons, composed of two identical light (L) chains and two identical heavy (H) chains and are typically referred to as full length antibodies. Each light chain is

covalently linked to a heavy chain by one disulfide bond to form a heterodimer, and the heterotrimeric molecule is formed through a covalent disulfide linkage between the two identical heavy chains of the heterodimers. Although the light and heavy chains are linked together by one disulfide bond, the number of disulfide linkages between the two heavy chains varies by immunoglobulin isotype. Each heavy and light chain also has regularly spaced intrachain disulfide bridges. Each heavy chain has at the amino-terminus a variable domain (V_H), followed by three or four constant domains (C_{H1} , C_{H2} , C_{H3} , and C_{H4}), as well as a hinge region between C_{H1} and C_{H2} . Each light chain has two domains, an amino-terminal variable domain (V_L) and a carboxy-terminal constant domain (C_L). The V_L domain associates non-covalently with the V_H domain, whereas the C_L domain is commonly covalently linked to the C_{H1} domain via a disulfide bond. Particular amino acid residues are believed to form an interface between the light and heavy chain variable domains (Chothia et al., 1985, J. Mol. Biol. 186:651-663). Variable domains are also referred herein as variable regions.

[0096] Certain domains within the variable domains differ extensively between different antibodies i.e., are "hypervariable." These hypervariable domains contain residues that are directly involved in the binding and specificity of each particular antibody for its specific antigenic determinant. Hypervariability, both in the light chain and the heavy chain variable domains, is concentrated in three segments known as complementarity determining regions (CDRs) or hypervariable loops (HVLs). CDRs are defined by sequence comparison in Kabat et al., 1991, In: Sequences of Proteins of Immunological Interest, 5th Ed. Public Health Service, National Institutes of Health, Bethesda, Md., whereas HVLs (also referred herein as CDRs) are structurally defined according to the three-dimensional structure of the variable domain, as described by Chothia and Lesk, 1987, J. Mol. Biol. 196: 901-917. These two methods result in slightly different identifications of a CDR. As defined by Kabat, CDR-L1 is positioned at about residues 24-34, CDR-L2, at about residues 50-56, and CDR-L3, at about residues 89-97 in the light chain variable domain; CDR-H1 is positioned at about residues 31-35, CDR-H2 at about residues 50-65, and CDR-H3 at about residues 95-102 in the heavy chain variable domain. The exact residue numbers that encompass a particular CDR will vary depending on the sequence and size of the CDR. Those skilled in the art can routinely determine which residues comprise a particular CDR given the variable region amino acid sequence of the antibody. The CDR1, CDR2, CDR3 of the heavy and light chains therefore define the unique and functional properties specific for a given antibody.

[0097] The three CDRs within each of the heavy and light chains are separated by framework regions (FR), which contain sequences that tend to be less variable. From the amino terminus to the carboxy terminus of the heavy and light chain variable domains, the FRs and CDRs are arranged in the order: FR1, CDR1, FR2, CDR2, FR3, CDR3, and FR4. The largely β -sheet configuration of the FRs brings the CDRs within each of the chains into close proximity to each other as well as to the CDRs from the other chain. The resulting conformation contributes to the antigen binding site (see Kabat et al., 1991, NIH Publ. No. 91-3242, Vol. I, pages 647-669), although not all CDR residues are necessarily directly involved in antigen binding.

[0098] FR residues and Ig constant domains are not directly involved in antigen binding, but contribute to antigen binding and/or mediate antibody effector function. Some FR residues are thought to have a significant effect on antigen binding in at least three ways: by noncovalently binding directly to an epitope, by interacting with one or more CDR residues, and by affecting the interface between the heavy and light chains. The constant domains are not directly involved in antigen binding but mediate various Ig effector functions, such as participation of the antibody in antibody dependent cellular cytotoxicity (ADCC), complement dependent cytotoxicity (CDC) and antibody dependent cellular phagocytosis (ADCP).

[0099] The light chains of vertebrate immunoglobulins are assigned to one of two clearly distinct classes, kappa (κ) and lambda (λ), based on the amino acid sequence of the constant domain. By comparison, the heavy chains of mammalian immunoglobulins are assigned to one of five major classes, according to the sequence of the constant domains: IgA, IgD, IgE, IgG, and IgM. IgG and IgA are further divided into subclasses (isotypes), e.g., IgG₁, IgG₂, IgG₃, IgG₄, IgA₁, and IgA₂. The heavy chain constant domains that correspond to the different classes of immunoglobulins are called α , β , ϵ , γ , and μ , respectively. The subunit structures and three-dimensional configurations of the classes of native immunoglobulins are well known.

[0100] The terms, “antibody”, “anti-IL-23p19 antibody”, “humanized anti-IL-23p19 antibody”, “humanized anti-IL-23p19 epitope antibody”, and “variant humanized anti-IL-23p19 epitope antibody” specifically encompass monoclonal antibodies (including full length monoclonal antibodies), polyclonal antibodies, multispecific antibodies (e.g., bispecific antibodies), and antibody fragments such as variable domains and other portions of antibodies that exhibit a desired biological activity, e.g., IL-23p19 binding. The term “monoclonal antibody” (mAb) refers to an antibody that is highly specific, being directed against a single antigenic determinant, an “epitope”. Therefore, the modifier “monoclonal” is indicative of antibodies directed to the identical epitope and is not to be construed as requiring production of the antibody by any particular method. It should be understood that monoclonal antibodies can be made by any technique or methodology known in the art; including e.g., the hybridoma method (Kohler et al., 1975, *Nature* 256: 495), or recombinant DNA methods known in the art (see, e.g., U.S. Pat. No. 4,816,567), or methods of isolation of monoclonal recombinantly produced using phage antibody libraries, using techniques described in Clackson et al., 1991, *Nature* 352: 624-628, and Marks et al., 1991, *J. Mol. Biol.* 222: 581-597.

[0101] The term “monomer” refers to a homogenous form of an antibody. For example, for a full-length antibody, monomer means a monomeric antibody having two identical heavy chains and two identical light chains.

[0102] Chimeric antibodies consist of the heavy and light chain variable regions of an antibody from one species (e.g., a non-human mammal such as a mouse) and the heavy and light chain constant regions of another species (e.g., human) antibody and can be obtained by linking the DNA sequences encoding the variable regions of the antibody from the first species (e.g., mouse) to the DNA sequences for the constant regions of the antibody from the second (e.g. human) species and transforming a host with an expression vector containing the linked sequences to allow it to produce a chimeric

antibody. Alternatively, the chimeric antibody also could be one in which one or more regions or domains of the heavy and/or light chain is identical with, homologous to, or a variant of the corresponding sequence in a monoclonal antibody from another immunoglobulin class or isotype, or from a consensus or germline sequence. Chimeric antibodies can include fragments of such antibodies, provided that the antibody fragment exhibits the desired biological activity of its parent antibody, for example binding to the same epitope (see, e.g., U.S. Pat. No. 4,816,567; and Morrison et al., 1984, *Proc. Natl. Acad. Sci. USA* 81: 6851-6855).

[0103] The terms, “antibody fragment”, “anti-IL-23p19 antibody fragment”, “anti-IL-23p19 epitope antibody fragment”, “humanized anti-IL-23p19 antibody fragment”, “humanized anti-IL-23p19 epitope antibody fragment”, “variant humanized anti-IL-23p19 epitope antibody fragment” refer to a portion of a full length anti-IL-23p19 antibody, in which a variable region or a functional capability is retained, for example, specific IL-23p19 epitope binding. Examples of antibody fragments include, but are not limited to, a Fab, Fab', F(ab')₂, Fd, Fv, scFv and scFv-Fc fragment, a diabody, a linear antibody, a single-chain antibody, a minibody, a diabody formed from antibody fragments, and multispecific antibodies formed from antibody fragments.

[0104] Full length antibodies can be treated with enzymes such as papain or pepsin to generate useful antibody fragments. Papain digestion is used to produce two identical antigen-binding antibody fragments called “Fab” fragments, each with a single antigen-binding site, and a residual “Fc” fragment. The Fab fragment also contains the constant domain of the light chain and the C_{H1} domain of the heavy chain. Pepsin treatment yields a F(ab')₂ fragment that has two antigen-binding sites and is still capable of cross-linking antigen.

[0105] Fab' fragments differ from Fab fragments by the presence of additional residues including one or more cysteines from the antibody hinge region at the C-terminus of the C_{H1} domain. F(ab')₂ antibody fragments are pairs of Fab' fragments linked by cysteine residues in the hinge region. Other chemical couplings of antibody fragments are also known.

[0106] “Fv” fragment contains a complete antigen-recognition and binding site consisting of a dimer of one heavy and one light chain variable domain in tight, non-covalent association. In this configuration, the three CDRs of each variable domain interact to define an antigen-binding site on the surface of the V_H-V_L dimer. Collectively, the six CDRs confer antigen-binding specificity to the antibody.

[0107] A “single-chain Fv” or “scFv” antibody fragment is a single chain Fv variant comprising the V_H and V_L domains of an antibody where the domains are present in a single polypeptide chain. The single chain Fv is capable of recognizing and binding antigen. The scFv polypeptide may optionally also contain a polypeptide linker positioned between the V_H and V_L domains in order to facilitate formation of a desired three-dimensional structure for antigen binding by the scFv (see, e.g., Pluckthun, 1994, In *The Pharmacology of monoclonal Antibodies*, Vol. 113, Rosenberg and Moore eds., Springer-Verlag, New York, pp. 269-315).

[0108] A “diabody” refers to small antibody fragments with two antigen-binding sites, which fragments comprise a heavy chain variable domain (V_{sub}.H) connected to a light

chain variable domain (V.sub.L) in the same polypeptide chain (V.sub.H-V.sub.L or V.sub.L-V.sub.H). Diabodies are described more fully in, e.g., Holliger et al. (1993) Proc. Natl. Acad. Sci. USA 90: 6444-6448.

[0109] Other recognized antibody fragments include those that comprise a pair of tandem Fd segments (V_H - C_{H1} - V_H - C_{H1}) to form a pair of antigen binding regions. These “linear antibodies” can be bispecific or monospecific as described in, for example, Zapata et al. 1995, Protein Eng. 8(10):1057-1062.

[0110] A “humanized antibody” or a “humanized antibody fragment” is a specific type of chimeric antibody which includes an immunoglobulin amino acid sequence variant, or fragment thereof, which is capable of binding to a predetermined antigen and which, comprises one or more FRs having substantially the amino acid sequence of a human immunoglobulin and one or more CDRs having substantially the amino acid sequence of a non-human immunoglobulin. This non-human amino acid sequence often referred to as an “import” sequence is typically taken from an “import” antibody domain, particularly a variable domain. In general, a humanized antibody includes at least the CDRs or HVLs of a non-human antibody, inserted between the FRs of a human heavy or light chain variable domain. The present invention describes specific humanized anti-IL-23p19 antibodies which contain CDRs derived from the mouse monoclonal antibodies or humanized CDRs shown in Tables 1 and 2 inserted between the FRs of human germline sequence heavy and light chain variable domains. It will be understood that certain mouse FR residues may be important to the function of the humanized antibodies and therefore certain of the human germline sequence heavy and light chain variable domains residues are modified to be the same as those of the corresponding mouse sequence.

[0111] In another aspect, a humanized anti-IL-23p19 antibody comprises substantially all of at least one, and typically two, variable domains (such as contained, for example, in Fab, Fab', F(ab')₂, Fabc, and Fv fragments) in which all, or substantially all, of the CDRs correspond to those of a non-human immunoglobulin, and specifically herein, all of the CDRs are mouse or humanized sequences as detailed in Tables 1 and 2 herein below and all, or substantially all, of the FRs are those of a human immunoglobulin consensus or germline sequence. In another aspect, a humanized anti-IL-23p19 antibody also includes at least a portion of an immunoglobulin Fc region, typically that of a human immunoglobulin. Ordinarily, the antibody will contain both the light chain as well as at least the variable domain of a heavy chain. The antibody also may include one or more of the C_{H1} , hinge, C_{H2} , C_{H3} , and/or C_{H4} regions of the heavy chain, as appropriate.

[0112] A humanized anti-IL-23p19 antibody can be selected from any class of immunoglobulins, including IgM, IgG, IgD, IgA and IgE, and any isotype, including IgG₁, IgG₂, IgG₃, IgG₄, IgA₁ and IgA₂. For example, the constant domain can be a complement fixing constant domain where it is desired that the humanized antibody exhibit cytotoxic activity, and the isotype is typically IgG₁. Where such cytotoxic activity is not desirable, the constant domain may be of another isotype, e.g., IgG₂. An alternative humanized anti-IL-23p19 antibody can comprise sequences from more than one immunoglobulin class or isotype, and selecting particular constant domains to optimize desired effector functions is within the ordinary skill in the art. In specific

embodiments, the present invention provides antibodies that are IgG1 antibodies and more particularly, are IgG1 antibodies in which there is a knock-out of effector functions.

[0113] The FRs and CDRs, or HVLs, of a humanized anti-IL-23p19 antibody need not correspond precisely to the parental sequences. For example, one or more residues in the import CDR, or HVL, or the consensus or germline FR sequence may be altered (e.g., mutagenized) by substitution, insertion or deletion such that the resulting amino acid residue is no longer identical to the original residue in the corresponding position in either parental sequence but the antibody nevertheless retains the function of binding to IL-23p19. Such alteration typically will not be extensive and will be conservative alterations. Usually, at least 75% of the humanized antibody residues will correspond to those of the parental consensus or germline FR and import CDR sequences, more often at least 90%, and most frequently greater than 95%, or greater than 98% or greater than 99%.

[0114] Immunoglobulin residues that affect the interface between heavy and light chain variable regions (“the V_L - V_H interface”) are those that affect the proximity or orientation of the two chains with respect to one another. Certain residues that may be involved in interchain interactions include V_L residues 34, 36, 38, 44, 46, 87, 89, 91, 96, and 98 and V_H residues 35, 37, 39, 45, 47, 91, 93, 95, 100, and 103 (utilizing the numbering system set forth in Kabat et al., Sequences of Proteins of Immunological Interest (National Institutes of Health, Bethesda, Md., 1987)). U.S. Pat. No. 6,407,213 also discusses that residues such as V_L residues 43 and 85, and V_H residues 43 and 60 also may be involved in this interaction. While these residues are indicated for human IgG only, they are applicable across species. Important antibody residues that are reasonably expected to be involved in interchain interactions are selected for substitution into the consensus sequence.

[0115] The terms “consensus sequence” and “consensus antibody” refer to an amino acid sequence which comprises the most frequently occurring amino acid residue at each location in all immunoglobulins of any particular class, isotype, or subunit structure, e.g., a human immunoglobulin variable domain. The consensus sequence may be based on immunoglobulins of a particular species or of many species. A “consensus” sequence, structure, or antibody is understood to encompass a consensus human sequence as described in certain embodiments, and to refer to an amino acid sequence which comprises the most frequently occurring amino acid residues at each location in all human immunoglobulins of any particular class, isotype, or subunit structure. Thus, the consensus sequence contains an amino acid sequence having at each position an amino acid that is present in one or more known immunoglobulins, but which may not exactly duplicate the entire amino acid sequence of any single immunoglobulin. The variable region consensus sequence is not obtained from any naturally produced antibody or immunoglobulin. Kabat et al., 1991, Sequences of Proteins of Immunological Interest, 5th Ed. Public Health Service, National Institutes of Health, Bethesda, Md., and variants thereof. The FRs of heavy and light chain consensus sequences, and variants thereof, provide useful sequences for the preparation of humanized anti-IL-23p19 antibodies. See, for example, U.S. Pat. Nos. 6,037,454 and 6,054,297.

[0116] Human germline sequences are found naturally in the human population. A combination of those germline genes generates antibody diversity. Germline antibody

sequences for the light chain of the antibody come from conserved human germline kappa or lambda v-genes and j-genes. Similarly the heavy chain sequences come from germline v-, d- and j-genes (LeFranc, M-P, and LeFranc, G, "The Immunoglobulin Facts Book" Academic Press, 2001).

[0117] As used herein, "variant", "anti-IL-23p19 variant", "humanized anti-IL-23p19 variant", or "variant humanized anti-IL-23p19" each refers to a humanized anti-IL-23p19 antibody having at least a light chain variable murine CDR from any of the sequences as shown in Table 1 or a heavy chain murine CDR sequence derived from the murine monoclonal antibody as shown in Table 2. Variants include those having one or more amino acid changes in one or both light chain or heavy chain variable domains, provided that the amino acid change does not substantially impair binding of the antibody to IL-23p19. Exemplary humanized antibodies produced herein include those designated as Antibody A, Antibody B, Antibody C and Antibody D, and the various light chains and heavy chains of the same are shown in SEQ ID Nos:18 and 21, and SEQ ID Nos:19 and 20, respectively.

[0118] An "isolated" antibody is one that has been identified and separated and/or recovered from a component of its natural environment. Contaminant components of the antibody's natural environment are those materials that may interfere with diagnostic or therapeutic uses of the antibody, and can be enzymes, hormones, or other proteinaceous or nonproteinaceous solutes. In one aspect, the antibody will be purified to at least greater than 95% isolation by weight of antibody.

[0119] An isolated antibody includes an antibody in situ within recombinant cells in which it is produced, since at least one component of the antibody's natural environment will not be present. Ordinarily however, an isolated antibody will be prepared by at least one purification step in which the recombinant cellular material is removed.

[0120] The term "antibody performance" refers to factors that contribute to antibody recognition of antigen or the effectiveness of an antibody in vivo. Changes in the amino acid sequence of an antibody can affect antibody properties such as folding, and can influence physical factors such as initial rate of antibody binding to antigen (k_a), dissociation constant of the antibody from antigen (k_d), affinity constant of the antibody for the antigen (Kd), conformation of the antibody, protein stability, and half life of the antibody.

[0121] The term "epitope tagged" when used herein, refers to an anti-IL-23p19 antibody fused to an "epitope tag". An "epitope tag" is a polypeptide having a sufficient number of amino acids to provide an epitope for antibody production, yet is designed such that it does not interfere with the desired activity of the humanized anti-IL-23p19 antibody. The epitope tag is usually sufficiently unique such that an antibody raised against the epitope tag does not substantially cross-react with other epitopes. Suitable tag polypeptides generally contain at least 6 amino acid residues and usually contain about 8 to 50 amino acid residues, or about 9 to 30 residues. Examples of epitope tags and the antibody that binds the epitope include the flu HA tag polypeptide and its antibody 12CA5 (Field et al., 1988 Mol. Cell. Biol. 8: 2159-2165; c-myc tag and 8F9, 3C7, 6E10, G4, B7 and 9E10 antibodies thereto (Evan et al., 1985, Mol. Cell. Biol. 5(12):3610-3616; and Herpes simplex virus glycoprotein D (gD) tag and its antibody (Paborsky et al. 1990, Protein Engineering 3(6): 547-553). In certain embodiments, the epitope tag is a "salvage receptor binding epitope". As used

herein, the term "salvage receptor binding epitope" refers to an epitope of the Fc region of an IgG molecule (such as IgG₁, IgG₂, IgG₃, or IgG₄) that is responsible for increasing the in vivo serum half-life of the IgG molecule.

[0122] For diagnostic as well as therapeutic monitoring purposes, the antibodies of the invention also may be conjugated to a label, either a label alone or a label and an additional second agent (prodrug, chemotherapeutic agent and the like). A label, as distinguished from the other second agents refers to an agent that is a detectable compound or composition and it may be conjugated directly or indirectly to a humanized antibody of the present invention. The label may itself be detectable (e.g., radioisotope labels or fluorescent labels) or, in the case of an enzymatic label, may catalyze chemical alteration of a substrate compound or composition that is detectable. Labeled humanized anti-IL-23p19 antibody can be prepared and used in various applications including in vitro and in vivo diagnostics.

[0123] In various aspects of the present invention one or more domains of the humanized antibodies will be recombinantly expressed. Such recombinant expression may employ one or more control sequences, i.e., polynucleotide sequences necessary for expression of an operably linked coding sequence in a particular host organism. The control sequences suitable for use in prokaryotic cells include, for example, promoter, operator, and ribosome binding site sequences. Eukaryotic control sequences include, but are not limited to, promoters, polyadenylation signals, and enhancers. These control sequences can be utilized for expression and production of humanized anti-IL-23p19 antibody in prokaryotic and eukaryotic host cells.

[0124] A nucleic acid sequence is "operably linked" when it is placed into a functional relationship with another nucleic acid sequence. For example, a nucleic acid presequence or secretory leader is operably linked to a nucleic acid encoding a polypeptide if it is expressed as a preprotein that participates in the secretion of the polypeptide; a promoter or enhancer is operably linked to a coding sequence if it affects the transcription of the sequence; or a ribosome binding site is operably linked to a coding sequence if it is positioned so as to facilitate translation. Generally, "operably linked" means that the DNA sequences being linked are contiguous, and, in the case of a secretory leader, contiguous and in reading frame. However, enhancers are optionally contiguous. Linking can be accomplished by ligation at convenient restriction sites. If such sites do not exist, synthetic oligonucleotide adaptors or linkers can be used.

[0125] As used herein, the expressions "cell", "cell line", and "cell culture" are used interchangeably and all such designations include the progeny thereof. Thus, "transformants" and "transformed cells" include the primary subject cell and cultures derived therefrom without regard for the number of transfers.

[0126] The term "mammal" for purposes of treatment refers to any animal classified as a mammal, including humans, domesticated and farm animals, and zoo, sports, or pet animals, such as dogs, horses, cats, cows, and the like. Preferably, the mammal is human.

[0127] A "disorder", as used herein, is any condition that would benefit from treatment with a humanized anti-IL-23p19 antibody described herein. This includes chronic and

acute disorders or diseases including those pathological conditions that predispose the mammal to the disorder in question.

[0128] As used herein, the term “IL-23-associated disorder” or “IL-23-associated disease” refers to a condition in which IL-23 activity contributes to the disease and typically where IL-23 is abnormally expressed. An IL-23-associated disorder includes diseases and disorders of the immune system, such as autoimmune disorders and inflammatory disorders. Such conditions include psoriasis, inflammatory bowel disease, for example ulcerative colitis or Crohn’s disease, and spondyloarthritis, for example ankylosing spondylitis, non-radiographic axial spondyloarthritis, peripheral spondyloarthritis or psoriatic arthritis.

[0129] The term “intravenous infusion” refers to introduction of an agent into the vein of an animal or human patient over a period of time greater than approximately 15 minutes, generally between approximately 30 to 90 minutes.

[0130] The term “intravenous bolus” or “intravenous push” refers to drug administration into a vein of an animal or human such that the body receives the drug in approximately 15 minutes or less, generally 5 minutes or less.

[0131] The term “subcutaneous administration” refers to introduction of an agent under the skin of an animal or human patient, preferable within a pocket between the skin and underlying tissue, by relatively slow, sustained delivery from a drug receptacle. Pinching or drawing the skin up and away from underlying tissue may create the pocket.

[0132] The term “subcutaneous infusion” refers to introduction of a drug under the skin of an animal or human patient, preferably within a pocket between the skin and underlying tissue, by relatively slow, sustained delivery from a drug receptacle for a period of time including, but not limited to, 30 minutes or less, or 90 minutes or less. Optionally, the infusion may be made by subcutaneous implantation of a drug delivery pump implanted under the skin of the animal or human patient, wherein the pump delivers a predetermined amount of drug for a predetermined period of time, such as 30 minutes, 90 minutes, or a time period spanning the length of the treatment regimen.

[0133] The term “subcutaneous bolus” refers to drug administration beneath the skin of an animal or human patient, where bolus drug delivery is less than approximately 15 minutes; in another aspect, less than 5 minutes, and in still another aspect, less than 60 seconds. In yet even another aspect, administration is within a pocket between the skin and underlying tissue, where the pocket may be created by pinching or drawing the skin up and away from underlying tissue.

[0134] The term “therapeutically effective amount” is used to refer to an amount of an active agent that relieves or ameliorates one or more of the symptoms of the disorder being treated. In another aspect, the therapeutically effective amount refers to a target serum concentration that has been shown to be effective in, for example, slowing disease progression. Efficacy can be measured in conventional ways, depending on the condition to be treated.

[0135] The terms “treatment” and “therapy” and the like, as used herein, are meant to include therapeutic as well as prophylactic, or suppressive measures for a disease or disorder leading to any clinically desirable or beneficial effect, including but not limited to alleviation or relief of one or more symptoms, regression, slowing or cessation of progression of the disease or disorder. Thus, for example, the

term treatment includes the administration of an agent prior to or following the onset of a symptom of a disease or disorder thereby preventing or removing one or more signs of the disease or disorder. As another example, the term includes the administration of an agent after clinical manifestation of the disease to combat the symptoms of the disease. Further, administration of an agent after onset and after clinical symptoms have developed where administration affects clinical parameters of the disease or disorder, such as the degree of tissue injury or the amount or extent of metastasis, whether or not the treatment leads to amelioration of the disease, comprises “treatment” or “therapy” as used herein. Moreover, as long as the compositions of the invention either alone or in combination with another therapeutic agent alleviate or ameliorate at least one symptom of a disorder being treated as compared to that symptom in the absence of use of the humanized anti-IL-23p19 antibody composition, the result should be considered an effective treatment of the underlying disorder regardless of whether all the symptoms of the disorder are alleviated or not.

[0136] The term “package insert” is used to refer to instructions customarily included in commercial packages of therapeutic products, that contain information about the indications, usage, administration, contraindications and/or warnings concerning the use of such therapeutic products.

Antibodies

[0137] The CDRs of selected antibodies used in the context of the present invention are shown in Table 1 and 2. The variable regions of selected antibodies used in the context of the present invention are shown in Table 3 and 4.

TABLE 1

LIGHT CHAIN CDR sequences		
L-CDR1	L-CDR2	L-CDR3
6B8 KASRDVAIAVA (SEQ ID NO: 1)	WASTRHT (SEQ ID NO: 2)	HQYSSYPPT (SEQ ID NO: 3)

TABLE 2

HEAVY CHAIN CDR sequences			
	H-CDR1	H-CDR2	H-CDR3
6B8	GNTFTDQTIH (SEQ ID NO: 4)	YIYPRDDSPKYNENFKG (SEQ ID NO: 5)	PDRSGYAWFIY (SEQ ID NO: 6)
Hu_6138-2	GYTFTDQTIH (SEQ ID NO: 7)	YIYPRDDSPKYNENFKG (SEQ ID NO: 5)	PDRSGYAWFIY (SEQ ID NO: 6)
Hu_6138-5	GFTFTDQTIH (SEQ ID NO: 8)	YIYPRDDSPKYNENFKG (SEQ ID NO: 5)	PDRSGYAWFIY (SEQ ID NO: 6)
Hu_6138-36/65	GGTFTDQTIH (SEQ ID NO: 9)	YIYPRDDSPKYNENFKG (SEQ ID NO: 5)	PDRSGYAWFIY (SEQ ID NO: 6)

TABLE 3

Humanized 6B8-VK Sequences	
6B8CVK-65	DIQMTQSPSSLSASVGDRTITCKASRDVAIAVAWY QQKPGKVPKLLFWASTRHTGVPDRFSGSGSGTDFT LTISLQPEDLADYYCHQYSSYPFTFGQGTKLEIK (SEQ ID NO: 10)
6B8CVK-66	DIQMTQSPSSLSASVGDRTITCKASRDVAIAVAWY QQKPGKVPKLLIYASTRHTGVPDRFSGSGSGTDFT LTISLQPEDVADYFCHQYSSYPFTFGSGTKLEIK (SEQ ID NO: 11)
6B8CVK-67	DIQMTQSPSSLSASVGDRTITCKASRDVAIAVAWY QQKPGKVPKLLIYASTRHTGVPDRFSGSGSGTDFT LTISLQPEDVATYYCHQYSSYPFTFGSGTKLEIK (SEQ ID NO: 12)
6B8CVK-78	DIQMTQSPSSLSASVGDRTITCKASRDVAIAVAWY QQKPGKVPKLLFWASTRHTGVPDRFSGSGSGTDFT LTISLQPEDLADYYCHQYSSYPFTFGSGTKLEIK (SEQ ID NO: 13)

TABLE 4

Humanized 6B8-VH Sequence	
6B8CVH-02	QVQLVQSGAEVKKPGSSVKVCKASGYTFTDQTIHW MRQAPGQGLEWIGYIYPRDPSPKYNENFKGKVTITA DKSTSTAYMELSSLRSEDTAVYYCAIPDRSGYAWFI YWGQGLTVTVSS (SEQ ID NO: 14)
6B8CVH-05	QVQLVQSGAEVKKPGSSVKVCKASGYTFTDQTIHW VRQAPGQGLEWIGYIYPRDPSPKYNENFKGKVTITA DKSTSTAYMELSSLRSEDTAVYYCAIPDRSGYAWFI YWGQGLTVTVSS (SEQ ID NO: 15)
6B8CVH-36	QVQLVQSGAEVKKPGSSVKVCKASGYTFTDQTIHW VRQAPGQGLEWIGYIYPRDPSPKYNENFKGKVTITA DKSTSTAYMELSSLRSEDTAVYYCAIPDRSGYAWFI YWGQGLTVTVSS (SEQ ID NO: 16)
6B8CVH-65	QVQLVQSGAEVKKPGSSVKVCKASGYTFTDQTIHW VRQAPGQGLEWIGYIYPRDPSPKYNENFKGKVTITA DKSTSTAYMELSSLRSEDTAVYYCARPDRSGYAWFI YWGQGLTVTVSS (SEQ ID NO: 17)

[0138] Selected combination of humanized light chain and heavy chain variable regions derived from mouse antibody 6B8 resulted in Antibodies A, B, C and D:

Antibody A: 6B8-IgG1 KO-2 with IgK-66 (heavy chain variable region 6B8CVH-02 and light chain variable region 6B8CVK-66);

Antibody B: 6B8-IgG1 KO-5 with IgK-66 (heavy chain variable region 6B8CVH-05 and light chain variable region 6B8CVK-66);

Antibody C: 6B8-IgG1 KO-2 with IgK-65 (heavy chain variable region 6B8CVH-02 and light chain variable region 6B8CVK-65);

Antibody D: 6B8-IgG1 KO-5 with IgK-65 (heavy chain variable region 6B8CVH-05 and light chain variable region 6B8CVK-65).

[0139] Antibodies A, B, C and D have the heavy and light chain sequences shown in Table 5.

TABLE 5

Heavy and Light Chain DNA and Amino Acid Sequences for Antibodies A, B, C, and D	
Anti-body A	IgK Chain #66: <u>DIQMTQSPSSLSASVGDRTITCKASRDVAIAVAWYQ</u> <u>QKPGKVPKLLIYASTRHTGVPDRFSGSGSGTDFTLT</u> <u>ISLQPEDVADYFCHQYSSYPFTFGSGTKLEIKRTVA</u> APSVFIFPPSDEQLKSGTASVCLNNFYPREAKVQW KVDNALQSGNSQESVTEQDSKDSSTYSLSSTLTLSKAD YEKHKVYACEVTHQGLSPVTKSFNRGEC (SEQ ID NO: 18) IgG1KO Heavy Chain #2: <u>QVQLVQSGAEVKKPGSSVKVCKASGYTFTDQTIHWM</u> <u>RQAPGQGLEWIGYIYPRDPSPKYNENFKGKVTITADK</u> <u>STSTAYMELSSLRSEDTAVYYCAIPDRSGYAWFIYWG</u> <u>QGLTVTVSSASTKGPSVFPLAPSSKSTSGGTAALGCL</u> VKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSL SSVVTVPSSSLGTQTYICNVNHKPSNTKVDKRVPEKS CDKTHCTCPPEAPEAAGGSPVFLPPPKDITLMISRT PEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPRE EQYNSYTRVSVLTIVLHQLDNLNGKEYCKVSNKALPAP PIEKTIKAKGQPREPQVYTLPPSREEMTKNQVSLTCL LVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDGSF FLYSKLTVDKSRWQQGNVFCSCVMHEALHNHYTQKSL SLSPG (SEQ ID NO: 19)
Anti-body B	IgK Chain #66: (SEQ ID NO: 18) IgG1KO Heavy Chain #5: <u>QVQLVQSGAEVKKPGSSVKVCKASGYTFTDQTIHWM</u> <u>RQAPGQGLEWIGYIYPRDPSPKYNENFKGKVTITADK</u> <u>STSTAYMELSSLRSEDTAVYYCAIPDRSGYAWFIYWG</u> <u>QGLTVTVSSASTKGPSVFPLAPSSKSTSGGTAALGCL</u> VKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSL SSVVTVPSSSLGTQTYICNVNHKPSNTKVDKRVPEKS CDKTHCTCPPEAPEAAGGSPVFLPPPKDITLMISRT PEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPRE EQYNSYTRVSVLTIVLHQLDNLNGKEYCKVSNKALPAP PIEKTIKAKGQPREPQVYTLPPSREEMTKNQVSLTCL LVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDGSF FLYSKLTVDKSRWQQGNVFCSCVMHEALHNHYTQKSL SLSPG (SEQ ID NO: 20)
Anti-body C	IgK Chain #65: <u>DIQMTQSPSSLSASVGDRTITCKASRDVAIAVAWYQ</u> <u>QKPGKVPKLLFWASTRHTGVPDRFSGSGSGTDFTLT</u> <u>ISLQPEDLADYYCHQYSSYPFTFGQGTKLEIKRTVA</u> APSVFIFPPSDEQLKSGTASVCLNNFYPREAKVQW KVDNALQSGNSQESVTEQDSKDSSTYSLSSTLTLSKAD YEKHKVYACEVTHQGLSPVTKSFNRGEC (SEQ ID NO: 21) IgG1KO Heavy Chain #5: (SEQ ID NO: 19)
Anti-body D	IgK Chain #65: (SEQ ID NO: 21) IgG1KO Heavy Chain #5: (SEQ ID NO: 20)

[0140] Light chains and heavy chain variable regions of Antibodies A, B, C, and D are underlined in Table 5 above.

[0141] In one aspect, a humanized antibody of the present invention is Antibody A, Antibody B, Antibody C or Antibody D. Accordingly, in one embodiment, a humanized antibody of the present invention comprises the light chain sequence of SEQ ID NO:18 and the heavy chain sequence of SEQ ID NO:19 (Antibody A). In another embodiment, a

humanized antibody of the present invention comprises the light chain sequence of SEQ ID NO:18 and the heavy chain sequence of SEQ ID NO:20 (Antibody B). In another embodiment, a humanized antibody of the present invention comprises the light chain sequence of SEQ ID NO:21 and the heavy chain sequence of SEQ ID NO:19 (Antibody C). In another embodiment, a humanized antibody of the present invention comprises the light chain sequence of SEQ ID NO:21 and the heavy chain sequence of SEQ ID NO:20 (Antibody D).

[0142] In a further embodiment, a humanized antibody of the present invention consists of the light chain sequence of SEQ ID NO:18 and the heavy chain sequence of SEQ ID NO:19 (Antibody A). In a further embodiment, a humanized antibody of the present invention consists of the light chain sequence of SEQ ID NO:18 and the heavy chain sequence of SEQ ID NO:20 (Antibody B). In a further embodiment, a humanized antibody of the present invention consists of the light chain sequence of SEQ ID NO:21 and the heavy chain sequence of SEQ ID NO:19 (Antibody C). In a further embodiment, a humanized antibody of the present invention consists of the light chain sequence of SEQ ID NO:21 and the heavy chain sequence of SEQ ID NO:20 (Antibody D).

[0143] In some embodiments, the humanized anti-IL-23p19 antibodies, including antigen-binding fragments thereof, such as heavy and light chain variable regions, comprise an amino acid sequence of the residues derived from Antibody A (light chain sequence=SEQ ID NO:18; heavy chain sequence=SEQ ID NO:19), Antibody B (light chain sequence=SEQ ID NO:18; heavy chain sequence=SEQ ID NO:20), Antibody C (light chain sequence=SEQ ID NO:21; heavy chain sequence=SEQ ID NO:19) or Antibody D (light chain sequence=SEQ ID NO:21; heavy chain sequence=SEQ ID NO:20).

[0144] In a further embodiment, the present invention provides an anti-IL-23p19 antibody or an antigen-binding fragment thereof that binds to human IL-23p19 at an epitope consisting of amino acid residues 108 to 126 and amino acid residues 137 to 151 of SEQ ID NO: 22.

[0145] In a further embodiment, the present invention provides an anti-IL-23p19 antibody or antigen-binding fragment thereof that competitively binds to human IL-23p19 with an antibody of the present invention, for example Antibody A, Antibody B, Antibody C or Antibody D described herein. The ability of an antibody or antigen-binding fragment to competitively bind to IL-23p19 can be measured using competitive binding assays known in the art.

[0146] In some embodiments, the present invention describes other humanized antibodies with light chain variable region sequences having the amino acid sequence set forth in of SEQ ID NO:10, 11, 12 or 13. In some embodiments, the present invention describes other humanized antibodies with heavy chain variable region sequences having the amino acid sequence set forth in of SEQ ID NO:14, 15, 16 or 17 (see Tables 3 and 4 above). The CDR sequences of these antibodies are shown in Tables 1 and 2. In particular, the present invention provides monoclonal antibodies with the combinations of light chain variable and heavy chain variable regions of SEQ ID NO: 11/14, 11/15, 10/14 or 10/15. Such variable regions can be combined with human constant regions.

[0147] In certain embodiments, the humanized anti-IL-23p19 antibody is an antibody fragment. Various antibody fragments have been generally discussed above and there are

techniques that have been developed for the production of antibody fragments. Fragments can be derived via proteolytic digestion of intact antibodies (see, e.g., Morimoto et al., 1992, Journal of Biochemical and Biophysical Methods 24:107-117; and Brennan et al., 1985, Science 229:81). Alternatively, the fragments can be produced directly in recombinant host cells. For example, Fab'-SH fragments can be directly recovered from *E. coli* and chemically coupled to form F(ab')₂ fragments (see, e.g., Carter et al., 1992, Bio/Technology 10:163-167). By another approach, F(ab')₂ fragments can be isolated directly from recombinant host cell culture. Other techniques for the production of antibody fragments will be apparent to the skilled practitioner. Accordingly, in one aspect, the present invention provides antibody fragments comprising the CDRs described herein, in particular one of the combinations of L-CDR1, L-CDR2, L-CDR3, H-CDR1, H-CDR2 and H-CDR3 described herein. In a further aspect, the present invention provides antibody fragments comprising the variable regions described herein, for example one of the combinations of light chain variable regions and heavy chain variable regions described herein.

[0148] Certain embodiments include an F(ab')₂ fragment of a humanized anti-IL-23p19 antibody comprise a light chain sequence of any of SEQ ID NO: 18 or 21 in combination with a heavy chain sequence of SEQ ID NO: 19 or 20. Such embodiments can include an intact antibody comprising such an F(ab')₂.

[0149] In some embodiments, the antibody or antibody fragment includes a constant region that mediates effector function. The constant region can provide antibody-dependent cellular cytotoxicity (ADCC), antibody-dependent cellular phagocytosis (ADCP) and/or complement-dependent cytotoxicity (CDC) responses against an IL-23 expressing target cell. The effector domain(s) can be, for example, an Fc region of an Ig molecule.

[0150] The effector domain of an antibody can be from any suitable vertebrate animal species and isotypes. The isotypes from different animal species differ in the abilities to mediate effector functions. For example, the ability of human immunoglobulin to mediate CDC and ADCC/ADCP is generally in the order of IgM≈IgG₁≈IgG₃>IgG₂>IgG₄ and IgG₁≈IgG₃>IgG₂/IgM/IgG₄, respectively. Murine immunoglobulins mediate CDC and ADCC/ADCP generally in the order of murine IgM≈IgG₃>>IgG_{2b}>IgG_{2a}>>IgG₁ and IgG_{2b}>IgG_{2a}>IgG₁>>IgG₃, respectively. In another example, murine IgG_{2a} mediates ADCC while both murine IgG_{2a} and IgM mediate CDC.

Polynucleotides, Vectors, Host Cells, and Recombinant Methods

[0151] Other embodiments encompass isolated polynucleotides that comprise a sequence encoding a humanized anti-IL-23p19 antibody, vectors, and host cells comprising the polynucleotides, and recombinant techniques for production of the humanized antibody. The isolated polynucleotides can encode any desired form of the anti-IL-23p19 antibody including, for example, full length monoclonal antibodies, Fab, Fab', F(ab')₂, and Fv fragments, diabodies, linear antibodies, single-chain antibody molecules, and multispecific antibodies formed from antibody fragments.

[0152] The polynucleotide(s) that comprise a sequence encoding a humanized anti-IL-23p19 antibody or a fragment or chain thereof can be fused to one or more regulatory or

control sequence, as known in the art, and can be contained in suitable expression vectors or host cell as known in the art. Each of the polynucleotide molecules encoding the heavy or light chain variable domains can be independently fused to a polynucleotide sequence encoding a constant domain, such as a human constant domain, enabling the production of intact antibodies. Alternatively, polynucleotides, or portions thereof, can be fused together, providing a template for production of a single chain antibody.

[0153] For recombinant production, a polynucleotide encoding the antibody is inserted into a replicable vector for cloning (amplification of the DNA) or for expression. Many suitable vectors for expressing the recombinant antibody are available. The vector components generally include, but are not limited to, one or more of the following: a signal sequence, an origin of replication, one or more marker genes, an enhancer element, a promoter, and a transcription termination sequence.

[0154] The humanized anti-IL-23p19 antibodies can also be produced as fusion polypeptides, in which the antibody is fused with a heterologous polypeptide, such as a signal sequence or other polypeptide having a specific cleavage site at the amino terminus of the mature protein or polypeptide. The heterologous signal sequence selected is typically one that is recognized and processed (i.e., cleaved by a signal peptidase) by the host cell. For prokaryotic host cells that do not recognize and process the humanized anti-IL-23p19 antibody signal sequence, the signal sequence can be substituted by a prokaryotic signal sequence. The signal sequence can be, for example, alkaline phosphatase, penicillinase, lipoprotein, heat-stable enterotoxin II leaders, and the like. For yeast secretion, the native signal sequence can be substituted, for example, with a leader sequence obtained from yeast invertase alpha-factor (including *Saccharomyces* and *Kluyveromyces* α -factor leaders), acid phosphatase, *C. albicans* glucoamylase, or the signal described in WO90/13646. In mammalian cells, mammalian signal sequences as well as viral secretory leaders, for example, the herpes simplex gD signal, can be used. The DNA for such precursor region is ligated in reading frame to DNA encoding the humanized anti-IL-23p19 antibody.

[0155] Expression and cloning vectors contain a nucleic acid sequence that enables the vector to replicate in one or more selected host cells. Generally, in cloning vectors this sequence is one that enables the vector to replicate independently of the host chromosomal DNA, and includes origins of replication or autonomously replicating sequences. Such sequences are well known for a variety of bacteria, yeast, and viruses. The origin of replication from the plasmid pBR322 is suitable for most Gram-negative bacteria, the 2-v. plasmid origin is suitable for yeast, and various viral origins (SV40, polyoma, adenovirus, VSV, and BPV) are useful for cloning vectors in mammalian cells. Generally, the origin of replication component is not needed for mammalian expression vectors (the SV40 origin may typically be used only because it contains the early promoter).

[0156] Expression and cloning vectors may contain a gene that encodes a selectable marker to facilitate identification of expression. Typical selectable marker genes encode proteins that confer resistance to antibiotics or other toxins, e.g., ampicillin, neomycin, methotrexate, or tetracycline, or alternatively, are complement auxotrophic deficiencies, or in

other alternatives supply specific nutrients that are not present in complex media, e.g., the gene encoding D-alanine racemase for *Bacilli*.

[0157] One example of a selection scheme utilizes a drug to arrest growth of a host cell. Those cells that are successfully transformed with a heterologous gene produce a protein conferring drug resistance and thus survive the selection regimen. Examples of such dominant selection use the drugs neomycin, mycophenolic acid, and hygromycin. Common selectable markers for mammalian cells are those that enable the identification of cells competent to take up a nucleic acid encoding a humanized anti-IL-23p19 antibody, such as DHFR (dihydrofolate reductase), thymidine kinase, metallothionein-I and -II (such as primate metallothionein genes), adenosine deaminase, ornithine decarboxylase, and the like. Cells transformed with the DHFR selection gene are first identified by culturing all of the transformants in a culture medium that contains methotrexate (MTx), a competitive antagonist of DHFR. An appropriate host cell when wild-type DHFR is employed is the Chinese hamster ovary (CHO) cell line deficient in DHFR activity (e.g., DG44).

[0158] Alternatively, host cells (particularly wild-type hosts that contain endogenous DHFR) transformed or co-transformed with DNA sequences encoding anti-IL-23p19 antibody, wild-type DHFR protein, and another selectable marker such as aminoglycoside 3'-phosphotransferase (APH), can be selected by cell growth in medium containing a selection agent for the selectable marker such as an aminoglycosidic antibiotic, e.g., kanamycin, neomycin, or G418. See, e.g., U.S. Pat. No. 4,965,199.

[0159] Where the recombinant production is performed in a yeast cell as a host cell, the TRP1 gene present in the yeast plasmid YRp7 (Stinchcomb et al., 1979, *Nature* 282: 39) can be used as a selectable marker. The TRP1 gene provides a selection marker for a mutant strain of yeast lacking the ability to grow in tryptophan, for example, ATCC No. 44076 or PEP4-1 (Jones, 1977, *Genetics* 85:12). The presence of the trp1 lesion in the yeast host cell genome then provides an effective environment for detecting transformation by growth in the absence of tryptophan. Similarly, Leu2p-deficient yeast strains such as ATCC 20,622 and 38,626 are complemented by known plasmids bearing the LEU2 gene.

[0160] In addition, vectors derived from the 1.6 μ m circular plasmid pKD1 can be used for transformation of *Kluyveromyces* yeasts. Alternatively, an expression system for large-scale production of recombinant calf chymosin was reported for *K. lactis* (Van den Berg, 1990, *Bio/Technology* 8:135). Stable multi-copy expression vectors for secretion of mature recombinant human serum albumin by industrial strains of *Kluyveromyces* have also been disclosed (Fleer et al., 1991, *Bio/Technology* 9:968-975).

[0161] Expression and cloning vectors usually contain a promoter that is recognized by the host organism and is operably linked to the nucleic acid molecule encoding an anti-IL-23p19 antibody or polypeptide chain thereof. Promoters suitable for use with prokaryotic hosts include phoA promoter, β -lactamase and lactose promoter systems, alkaline phosphatase, tryptophan (trp) promoter system, and hybrid promoters such as the tac promoter. Other known bacterial promoters are also suitable. Promoters for use in bacterial systems also will contain a Shine-Dalgarno (S.D.) sequence operably linked to the DNA encoding the humanized anti-IL-23p19 antibody.

[0162] Many eukaryotic promoter sequences are known. Virtually all eukaryotic genes have an AT-rich region located approximately 25 to 30 bases upstream from the site where transcription is initiated. Another sequence found 70 to 80 bases upstream from the start of transcription of many genes is a CNCAAT region where N may be any nucleotide. At the 3' end of most eukaryotic genes is an AATAAA sequence that may be the signal for addition of the poly A tail to the 3' end of the coding sequence. All of these sequences are suitably inserted into eukaryotic expression vectors.

[0163] Examples of suitable promoting sequences for use with yeast hosts include the promoters for 3-phosphoglycerate kinase or other glycolytic enzymes, such as enolase, glyceraldehyde-3-phosphate dehydrogenase, hexokinase, pyruvate decarboxylase, phosphofructokinase, glucose-6-phosphate isomerase, 3-phosphoglycerate mutase, pyruvate kinase, triosephosphate isomerase, phosphoglucose isomerase, and glucokinase.

[0164] Inducible promoters have the additional advantage of transcription controlled by growth conditions. These include yeast promoter regions for alcohol dehydrogenase 2, isocytocrome C, acid phosphatase, derivative enzymes associated with nitrogen metabolism, metallothionein, glyceraldehyde-3-phosphate dehydrogenase, and enzymes responsible for maltose and galactose utilization. Suitable vectors and promoters for use in yeast expression are further described in EP 73,657. Yeast enhancers also are advantageously used with yeast promoters.

[0165] Humanized anti-IL-23p19 antibody transcription from vectors in mammalian host cells is controlled, for example, by promoters obtained from the genomes of viruses such as polyoma virus, fowlpox virus, adenovirus (such as Adenovirus 2), bovine papilloma virus, avian sarcoma virus, cytomegalovirus, a retrovirus, hepatitis-B virus and Simian Virus 40 (SV40), from heterologous mammalian promoters, e.g., the actin promoter or an immunoglobulin promoter, or from heat-shock promoters, provided such promoters are compatible with the host cell systems.

[0166] The early and late promoters of the SV40 virus are conveniently obtained as an SV40 restriction fragment that also contains the SV40 viral origin of replication. The immediate early promoter of the human cytomegalovirus is conveniently obtained as a HindIII E restriction fragment. A system for expressing DNA in mammalian hosts using the bovine papilloma virus as a vector is disclosed in U.S. Pat. No. 4,419,446. A modification of this system is described in U.S. Pat. No. 4,601,978. See also Reyes et al., 1982, Nature 297:598-601, disclosing expression of human p-interferon cDNA in mouse cells under the control of a thymidine kinase promoter from herpes simplex virus. Alternatively, the Rous sarcoma virus long terminal repeat can be used as the promoter.

[0167] Another useful element that can be used in a recombinant expression vector is an enhancer sequence, which is used to increase the transcription of a DNA encoding a humanized anti-IL-23p19 antibody by higher eukaryotes. Many enhancer sequences are now known from mammalian genes (e.g., globin, elastase, albumin, α -feto-protein, and insulin). Typically, however, an enhancer from a eukaryotic cell virus is used. Examples include the SV40 enhancer on the late side of the replication origin (bp 100-270), the cytomegalovirus early promoter enhancer, the polyoma enhancer on the late side of the replication origin, and adenovirus enhancers. See also Yaniv, 1982, Nature

297:17-18 for a description of enhancing elements for activation of eukaryotic promoters. The enhancer may be spliced into the vector at a position 5' or 3' to the humanized anti-IL-23p19 antibody-encoding sequence, but is preferably located at a site 5' from the promoter.

[0168] Expression vectors used in eukaryotic host cells (yeast, fungi, insect, plant, animal, human, or nucleated cells from other multicellular organisms) can also contain sequences necessary for the termination of transcription and for stabilizing the mRNA. Such sequences are commonly available from the 5' and, occasionally 3', untranslated regions of eukaryotic or viral DNAs or cDNAs. These regions contain nucleotide segments transcribed as polyadenylated fragments in the untranslated portion of the mRNA encoding anti-IL-23p19 antibody. One useful transcription termination component is the bovine growth hormone polyadenylation region. See WO94/11026 and the expression vector disclosed therein. In some embodiments, humanized anti-IL-23p19 antibodies can be expressed using the CHEF system. (See, e.g., U.S. Pat. No. 5,888,809; the disclosure of which is incorporated by reference herein.)

[0169] Suitable host cells for cloning or expressing the DNA in the vectors herein are the prokaryote, yeast, or higher eukaryote cells described above. Suitable prokaryotes for this purpose include eubacteria, such as Gram-negative or Gram-positive organisms, for example, Enterobacteriaceae such as *Escherichia*, e.g., *E. coli*, *Enterobacter*, *Erwinia*, *Klebsiella*, *Proteus*, *Salmonella*, e.g., *Salmonella typhimurium*, *Serratia*, e.g., *Serratia marcescans*, and *Shigella*, as well as Bacilli such as *B. subtilis* and *B. licheniformis* (e.g., *B. licheniformis* 41 P disclosed in DD 266,710 published Apr. 12, 1989), *Pseudomonas* such as *P. aeruginosa*, and *Streptomyces*. One preferred *E. coli* cloning host is *E. coli* 294 (ATCC 31,446), although other strains such as *E. coli* B, *E. coli* X1776 (ATCC 31,537), and *E. coli* W3110 (ATCC 27,325) are suitable. These examples are illustrative rather than limiting.

[0170] In addition to prokaryotes, eukaryotic microbes such as filamentous fungi or yeast are suitable cloning or expression hosts for humanized anti-IL-23p19 antibody-encoding vectors. *Saccharomyces cerevisiae*, or common baker's yeast, is the most commonly used among lower eukaryotic host microorganisms. However, a number of other genera, species, and strains are commonly available and useful herein, such as *Schizosaccharomyces pombe*; *Kluyveromyces* hosts such as, e.g., *K. lactis*, *K. fragilis* (ATCC 12,424), *K. bulgaricus* (ATCC 16,045), *K. wickerhamii* (ATCC 24,178), *K. waltii* (ATCC 56,500), *B. drosophilae* (ATCC 36,906), *K. thermotolerans*, and *K. marxianus*; *yarrowia* (EP 402,226); *Pichia* pastors (EP 183,070); *Candida*; *Trichoderma reesia* (EP 244,234); *Neurospora crassa*; *Schwanniomyces* such as *Schwanniomyces occidentalis*; and filamentous fungi such as, e.g., *Neurospora*, *Penicillium*, *Tolypocladium*, and *Aspergillus* hosts such as *A. nidulans* and *A. niger*.

[0171] Suitable host cells for the expression of glycosylated humanized anti-IL-23p19 antibody are derived from multicellular organisms. Examples of invertebrate cells include plant and insect cells, including, e.g., numerous baculoviral strains and variants and corresponding permissive insect host cells from hosts such as *Spodoptera frugiperda* (caterpillar), *Aedes aegypti* (mosquito), *Aedes albopictus* (mosquito), *Drosophila melanogaster* (fruitfly), and *Bombyx mori* (silk worm). A variety of viral strains for

transfection are publicly available, e.g., the L-1 variant of *Autographa californica* NPV and the Bm-5 strain of *Bombyx mori* NPV, and such viruses may be used, particularly for transfection of *Spodoptera frugiperda* cells.

[0172] Plant cell cultures of cotton, corn, potato, soybean, petunia, tomato, and tobacco can also be utilized as hosts.

[0173] In another aspect, expression of humanized anti-IL-23p19 is carried out in vertebrate cells. The propagation of vertebrate cells in culture (tissue culture) has become routine procedure and techniques are widely available. Examples of useful mammalian host cell lines are monkey kidney CV1 line transformed by SV40 (COS-7, ATCC CRL 1651), human embryonic kidney line (293 or 293 cells subcloned for growth in suspension culture, (Graham et al., 1977, J. Gen. Virol. 36: 59), baby hamster kidney cells (BHK, ATCC CCL 10), Chinese hamster ovary cells/DHFR1 (CHO, Urlaub et al., 1980, Proc. Natl. Acad. Sci. USA 77: 4216; e.g., DG44), mouse sertoli cells (TM4, Mather, 1980, Biol. Reprod. 23:243-251), monkey kidney cells (CV1 ATCC CCL 70), African green monkey kidney cells (VERO-76, ATCC CRL-1587), human cervical carcinoma cells (HELA, ATCC CCL 2), canine kidney cells (MDCK, ATCC CCL 34), buffalo rat liver cells (BRL 3A, ATCC CRL 1442), human lung cells (W138, ATCC CCL 75), human liver cells (Hep G2, HB 8065), mouse mammary tumor (MMT 060562, ATCC CCL51), TR1 cells (Mather et al., 1982, Annals N.Y. Acad. Sci. 383: 44-68), MRC 5 cells, FS4 cells, and human hepatoma line (Hep G2).

[0174] Host cells are transformed with the above-described expression or cloning vectors for humanized anti-IL-23p19 antibody production and cultured in conventional nutrient media modified as appropriate for inducing promoters, selecting transformants, or amplifying the genes encoding the desired sequences.

[0175] The host cells used to produce a humanized anti-IL-23p19 antibody described herein may be cultured in a variety of media. Commercially available media such as Ham's F10 (Sigma-Aldrich Co., St. Louis, Mo.), Minimal Essential Medium ((MEM), (Sigma-Aldrich Co.), RPMI-1640 (Sigma-Aldrich Co.), and Dulbecco's Modified Eagle's Medium ((DMEM), Sigma-Aldrich Co.) are suitable for culturing the host cells. In addition, any of the media described in one or more of Ham et al., 1979, Meth. Enz. 58: 44, Barnes et al., 1980, Anal. Biochem. 102: 255, U.S. Pat. No. 4,767,704, U.S. Pat. No. 4,657,866, U.S. Pat. No. 4,927,762, U.S. Pat. No. 4,560,655, U.S. Pat. No. 5,122,469, WO 90/103430, and WO 87/00195 may be used as culture media for the host cells. Any of these media may be supplemented as necessary with hormones and/or other growth factors (such as insulin, transferrin, or epidermal growth factor), salts (such as sodium chloride, calcium, magnesium, and phosphate), buffers (such as HEPES), nucleotides (such as adenosine and thymidine), antibiotics (such as gentamicin), trace elements (defined as inorganic compounds usually present at final concentrations in the micromolar range), and glucose or an equivalent energy source. Other supplements may also be included at appropriate concentrations that would be known to those skilled in the art. The culture conditions, such as temperature, pH, and the like, are those previously used with the host cell selected for expression, and will be apparent to the ordinarily skilled artisan.

[0176] When using recombinant techniques, the antibody can be produced intracellularly, in the periplasmic space, or

directly secreted into the medium. If the antibody is produced intracellularly, the cells may be disrupted to release protein as a first step. Particulate debris, either host cells or lysed fragments, can be removed, for example, by centrifugation or ultrafiltration. Carter et al., 1992, Bio/Technology 10:163-167 describes a procedure for isolating antibodies that are secreted to the periplasmic space of *E. coli*. Briefly, cell paste is thawed in the presence of sodium acetate (pH 3.5), EDTA, and phenylmethylsulfonylfluoride (PMSF) over about 30 minutes. Cell debris can be removed by centrifugation. Where the antibody is secreted into the medium, supernatants from such expression systems are generally first concentrated using a commercially available protein concentration filter, for example, an Amicon or Millipore Pellicon ultrafiltration unit. A protease inhibitor such as PMSF may be included in any of the foregoing steps to inhibit proteolysis and antibiotics may be included to prevent the growth of adventitious contaminants. A variety of methods can be used to isolate the antibody from the host cell.

[0177] The antibody composition prepared from the cells can be purified using, for example, hydroxylapatite chromatography, gel electrophoresis, dialysis, and affinity chromatography, with affinity chromatography being a typical purification technique. The suitability of protein A as an affinity ligand depends on the species and isotype of any immunoglobulin Fc domain that is present in the antibody. Protein A can be used to purify antibodies that are based on human gamma1, gamma2, or gamma4 heavy chains (see, e.g., Lindmark et al., 1983 J. Immunol. Meth. 62:1-13). Protein G is recommended for all mouse isotypes and for human gamma3 (see, e.g., Guss et al., 1986 EMBO J. 5:1567-1575). A matrix to which an affinity ligand is attached is most often agarose, but other matrices are available. Mechanically stable matrices such as controlled pore glass or poly(styrenedivinyl)benzene allow for faster flow rates and shorter processing times than can be achieved with agarose. Where the antibody comprises a C_{H3} domain, the Bakerbond ABXTM resin (J. T. Baker, Phillipsburg, N.J.) is useful for purification. Other techniques for protein purification such as fractionation on an ion-exchange column, ethanol precipitation, reverse phase HPLC, chromatography on silica, chromatography on heparin SEPHAROSETM chromatography on an anion or cation exchange resin (such as a polyaspartic acid column), chromatofocusing, SDS-PAGE, and ammonium sulfate precipitation are also available depending on the antibody to be recovered.

[0178] Following any preliminary purification step(s), the mixture comprising the antibody of interest and contaminants may be subjected to low pH hydrophobic interaction chromatography using an elution buffer at a pH between about 2.5-4.5, typically performed at low salt concentrations (e.g., from about 0-0.25M salt).

Kits Using Biomarkers

[0179] One or more biomarkers disclosed herein (for example β -defensin 2, neutrophil gelatinase lipocalin, and S-100 proteins) can be developed into an analytical method, for example an assay kit, a test stick or another diagnostic device, for example for predicting response to an anti-IL-23p19 antibody either prior to treatment or early after initiation of treatment of patients. In one aspect, such analytical method includes a packaged combination of reagents in predetermined amounts, with or without a ana-

lytical device, with instructions for performing the diagnostic assay. Where specific antibodies are labeled with an enzyme or a dye or a radioactive isotope, the kit may include substrates and cofactors required by the enzyme such as a substrate precursor that provides the detectable chromophore or fluorophore. In addition, other additives may be included such as stabilizers, buffers (for example a block buffer or lysis buffer), and the like. The relative amounts of the various reagents may be varied widely to provide for concentrations in solution of the reagents that substantially optimize the sensitivity of the assay. The reagents may be provided as dry powders, usually lyophilized, including excipients that on dissolution will provide a reagent solution having the appropriate concentration or as solutions.

Therapeutic Uses

[0180] In another embodiment, a humanized anti-IL-23p19 antibody disclosed herein is useful in the treatment of various disorders associated with the expression of IL-23p19 as described herein. Methods for treating an IL-23 associated disorder comprise administering a therapeutically effective amount of a humanized anti-IL-23p19 antibody to a subject in need thereof.

[0181] The humanized anti-IL-23p19 antibody or agent is administered by any suitable means, including parenteral, subcutaneous, intraperitoneal, intrapulmonary, and intranasal, and, if desired for local immunosuppressive treatment, intralesional administration (including perfusing or otherwise contacting the graft with the antibody before transplantation). The humanized anti-IL-23p19 antibody or agent can be administered, for example, as an infusion or as a bolus. Parenteral infusions include intramuscular, intravenous, intraarterial, intraperitoneal, or subcutaneous administration. In addition, the humanized anti-IL-23p19 antibody is suitably administered by pulse infusion, particularly with declining doses of the antibody. In one aspect, the dosing is given by injections, most preferably intravenous or subcutaneous injections, depending in part on whether the administration is brief or chronic.

[0182] For the prevention or treatment of disease, the appropriate dosage of antibody will depend on a variety of factors such as the type of disease to be treated, as defined above, the severity and course of the disease, whether the antibody is administered for preventive or therapeutic purposes, previous therapy, the patient's clinical history and response to the antibody, and the discretion of the attending physician. The antibody is suitably administered to the patient at one time or over a series of treatments.

[0183] Depending on the type and severity of the disease, about 1 µg/kg to 20 mg/kg (e.g., 0.1-15 mg/kg) of antibody is an initial candidate dosage for administration to the patient, whether, for example, by one or more separate administrations, or by continuous infusion. A typical daily dosage might range from about 1 µg/kg to 100 mg/kg or more, depending on the factors mentioned above. For repeated administrations over several days or longer, depending on the condition, the treatment is sustained until a desired suppression of disease symptoms occurs. However, other dosage regimens may be useful. The progress of this therapy is easily monitored by conventional techniques and assays. An exemplary dosing regimen is that disclosed in WO 94/04188.

[0184] The term "suppression" is used herein in the same context as "amelioration" and "alleviation" to mean a lessening of one or more characteristics of the disease.

[0185] The antibody composition will be formulated, dosed, and administered in a fashion consistent with good medical practice. Factors for consideration in this context include the particular disorder being treated, the particular mammal being treated, the clinical condition of the individual patient, the cause of the disorder, the site of delivery of the agent, the method of administration, the scheduling of administration, and other factors known to medical practitioners. The "therapeutically effective amount" of the antibody to be administered will be governed by such considerations, and is the minimum amount necessary to prevent, ameliorate, or treat the disorder associated with IL-23 expression.

[0186] The antibody need not be, but is optionally, formulated with one or more agents currently used to prevent or treat the disorder in question. The effective amount of such other agents depends on the amount of humanized anti-IL-23p19 antibody present in the formulation, the type of disorder or treatment, and other factors discussed above. These are generally used in the same dosages and with administration routes as used hereinbefore or about from 1 to 99% of the heretofore employed dosages.

IL-23-Associated Disorders

[0187] The anti-IL-23p19 antibodies or agents are useful for treating or preventing an immunological disorder characterized by abnormal expression of IL-23, e.g., by inappropriate activation of immune cells (e.g., lymphocytes or dendritic cells). Such abnormal expression of IL-23 can be due to, for example, increased IL-23 protein levels.

[0188] Immunological diseases that are characterized by inappropriate activation of immune cells and that can be treated or prevented by the methods described herein can be classified, for example, by the type(s) of hypersensitivity reaction(s) that underlie the disorder. These reactions are typically classified into four types: anaphylactic reactions, cytotoxic (cytolytic) reactions, immune complex reactions, or cell-mediated immunity (CMI) reactions (also referred to as delayed-type hypersensitivity (DTH) reactions). (See, e.g., *Fundamental Immunology* (William E. Paul ed., Raven Press, N.Y., 3rd ed. 1993).) Immunological diseases include inflammatory diseases and autoimmune diseases.

[0189] Specific examples of immunological diseases include the following: psoriasis, inflammatory bowel disease, for example ulcerative colitis or Crohn's disease, and spondyloarthritis, for example ankylosing spondylitis, non-radiographic axial spondyloarthritis, peripheral spondyloarthritis or psoriatic arthritis.

[0190] In one aspect, the immunological disease is psoriasis. Psoriasis is a chronic inflammatory disease of the skin characterized by dysfunctional keratinocyte differentiation and hyperproliferation and marked accumulation of inflammatory T cells and dendritic cells. For example, the immunological disease includes plaque psoriasis, for example chronic plaque psoriasis, for example moderate to severe chronic plaque psoriasis, for example in patients who are candidates for systemic therapy or phototherapy. For example, the immunological disease includes palmar pustular psoriasis, guttate psoriasis, inverse psoriasis, pustular psoriasis or erythrodermic psoriasis.

Pharmaceutical Compositions and Administration Thereof

[0191] A composition comprising an IL-23p19 binding agent (e.g., an anti-IL-23p19 antibody) can be administered to a subject having or at risk of having an immunological disorder. The invention further provides for the use of a IL-23p19 binding agent (e.g., an anti-IL-23p19 antibody) in the manufacture of a medicament for prevention or treatment of an immunological disorder. The term "subject" as used herein means any mammalian patient to which an IL-23p19 binding agent can be administered, including, e.g., humans and non-human mammals, such as primates, rodents, and dogs. Subjects specifically intended for treatment using the methods described herein include humans. The antibodies or agents can be administered either alone or in combination with other compositions in the prevention or treatment of the immunological disorder.

[0192] Further examples of antibodies for use in such pharmaceutical compositions are also those that comprise a humanized antibody or antibody fragment having the light chain variable region amino acid sequence of any of SEQ ID NO:10, 11, 12 or 13. Preferred antibodies for use in such pharmaceutical compositions are also those that comprise a humanized antibody or antibody fragment having the heavy chain variable region amino acid sequence of any of SEQ ID NO:14, 15, 16 or 17.

[0193] Further examples of antibodies for use in such pharmaceutical compositions are also those that comprise a humanized antibody or antibody fragment having the light chain variable region and heavy chain variable region of any of SEQ ID NO: 11 and 14, SEQ ID NO: 11 and 15, SEQ ID NO: 10 and 14 or SEQ ID NO: 10 and 15.

[0194] Further examples of antibodies for use in such pharmaceutical compositions are also those that comprise a humanized antibody having the light chain region amino acid sequence of any of SEQ ID NO:18 or 21. Preferred antibodies for use in such pharmaceutical compositions are also those that comprise humanized antibody having the heavy chain variable region amino acid sequence of any of SEQ ID NO:19 or 20.

[0195] Further examples of antibodies for use in such pharmaceutical compositions are also those that comprise Antibody A, Antibody B, Antibody C or Antibody D.

[0196] Various delivery systems are known and can be used to administer the IL-23p19 binding agent. Methods of introduction include but are not limited to intradermal, intramuscular, intraperitoneal, intravenous, subcutaneous, intranasal, epidural, and oral routes. The IL-23p19 binding agent can be administered, for example by infusion, bolus or injection, and can be administered together with other biologically active agents such as chemotherapeutic agents. Administration can be systemic or local. In preferred embodiments, the administration is by subcutaneous injection. Formulations for such injections may be prepared in for example prefilled syringes that may be administered once every other week.

[0197] In specific embodiments, the IL-23p19 binding agent composition is administered by injection, by means of a catheter, by means of a suppository, or by means of an implant, the implant being of a porous, non-porous, or gelatinous material, including a membrane, such as a sialastic membrane, or a fiber. Typically, when administering the composition, materials to which the anti-IL-23p19 antibody or agent does not absorb are used.

[0198] In other embodiments, the anti-IL-23p19 antibody or agent is delivered in a controlled release system. In one embodiment, a pump may be used (see, e.g., Langer, 1990, *Science* 249:1527-1533; Sefton, 1989, *CRC Crit. Rev. Biomed. Eng.* 14:201; Buchwald et al., 1980, *Surgery* 88:507; Saudek et al., 1989, *N. Engl. J. Med.* 321:574). In another embodiment, polymeric materials can be used. (See, e.g., *Medical Applications of Controlled Release* (Langer and Wise eds., CRC Press, Boca Raton, Fla., 1974); *Controlled Drug Bioavailability, Drug Product Design and Performance* (Smolen and Ball eds., Wiley, New York, 1984); Ranger and Peppas, 1983, *Macromol. Sci. Rev. Macromol. Chem.* 23:61. See also Levy et al., 1985, *Science* 228:190; During et al., 1989, *Ann. Neurol.* 25:351; Howard et al., 1989, *J. Neurosurg.* 71:105.) Other controlled release systems are discussed, for example, in Langer, supra.

[0199] An IL-23p19 binding agent (e.g., an anti-IL-23p19 antibody) can be administered as pharmaceutical compositions comprising a therapeutically effective amount of the binding agent and one or more pharmaceutically compatible ingredients.

[0200] In typical embodiments, the pharmaceutical composition is formulated in accordance with routine procedures as a pharmaceutical composition adapted for intravenous or subcutaneous administration to human beings. Typically, compositions for administration by injection are solutions in sterile isotonic aqueous buffer. Where necessary, the pharmaceutical can also include a solubilizing agent and a local anesthetic such as lignocaine to ease pain at the site of the injection. Generally, the ingredients are supplied either separately or mixed together in unit dosage form, for example, as a dry lyophilized powder or water free concentrate in a hermetically sealed container such as an ampoule or sachette indicating the quantity of active agent. Where the pharmaceutical is to be administered by infusion, it can be dispensed with an infusion bottle containing sterile pharmaceutical grade water or saline. Where the pharmaceutical is administered by injection, an ampoule of sterile water for injection or saline can be provided so that the ingredients can be mixed prior to administration.

[0201] Further, the pharmaceutical composition can be provided as a pharmaceutical kit comprising (a) a container containing a IL-23p19 binding agent (e.g., an anti-IL-23p19 antibody) in lyophilized form and (b) a second container containing a pharmaceutically acceptable diluent (e.g., sterile water) for injection. The pharmaceutically acceptable diluent can be used for reconstitution or dilution of the lyophilized anti-IL-23p19 antibody or agent. Optionally associated with such container(s) can be a notice in the form prescribed by a governmental agency regulating the manufacture, use or sale of pharmaceuticals or biological products, which notice reflects approval by the agency of manufacture, use or sale for human administration.

[0202] Examples of pharmaceutical compositions used in the context of the present invention are disclosed in Example 2 hereinbelow.

[0203] The amount of the IL-23p19 binding agent (e.g., anti-IL-23p19 antibody) that is effective in the treatment or prevention of an immunological disorder can be determined by standard clinical techniques. In addition, *in vitro* assays may optionally be employed to help identify optimal dosage ranges. The precise dose to be employed in the formulation will also depend on the route of administration, and the stage of immunological disorder, and should be decided according

to the judgment of the practitioner and each patient's circumstances. Effective doses may be extrapolated from dose-response curves derived from in vitro or animal model test systems.

[0204] Generally, the dosage of an anti-IL-23p19 antibody or IL-23p19 binding agent administered to a patient with an immunological disorder is typically about 0.1 mg/kg to about 100 mg/kg of the subject's body weight. The dosage administered to a subject is about 0.1 mg/kg to about 50 mg/kg, about 1 mg/kg to about 30 mg/kg, about 1 mg/kg to about 20 mg/kg, about 1 mg/kg to about 15 mg/kg, or about 1 mg/kg to about 10 mg/kg of the subject's body weight.

[0205] Exemplary doses include, but are not limited to, from 1 ng/kg to 100 mg/kg. In some embodiments, a dose is about 0.5 mg/kg, about 1 mg/kg, about 2 mg/kg, about 3 mg/kg, about 4 mg/kg, about 5 mg/kg, about 6 mg/kg, about 7 mg/kg, about 8 mg/kg, about 9 mg/kg, about 10 mg/kg, about 11 mg/kg, about 12 mg/kg, about 13 mg/kg, about 14 mg/kg, about 15 mg/kg or about 16 mg/kg. The dose can be administered, for example, daily, once per week (weekly), twice per week, thrice per week, four times per week, five times per week, six times per week, biweekly or monthly, every two months, or every three months. In specific embodiments, the dose is about 0.5 mg/kg/week, about 1 mg/kg/week, about 2 mg/kg/week, about 3 mg/kg/week, about 4 mg/kg/week, about 5 mg/kg/week, about 6 mg/kg/week, about 7 mg/kg/week, about 8 mg/kg/week, about 9 mg/kg/week, about 10 mg/kg/week, about 11 mg/kg/week, about 12 mg/kg/week, about 13 mg/kg/week, about 14 mg/kg/week, about 15 mg/kg/week or about 16 mg/kg/week. In some embodiments, the dose ranges from about 1 mg/kg/week to about 15 mg/kg/week.

[0206] In some embodiments, the pharmaceutical compositions comprising the IL-23p19 binding agent can further comprise a therapeutic agent, either conjugated or unconjugated to the binding agent. The anti-IL-23p19 antibody or IL-23p19 binding agent can be co-administered in combination with one or more therapeutic agents for the treatment or prevention of immunological disorders.

[0207] Such combination therapy administration can have an additive or synergistic effect on disease parameters (e.g., severity of a symptom, the number of symptoms, or frequency of relapse).

[0208] With respect to therapeutic regimens for combinatorial administration, in a specific embodiment, an anti-IL-23p19 antibody or IL-23p19 binding agent is administered concurrently with a therapeutic agent. In another specific embodiment, the therapeutic agent is administered prior or subsequent to administration of the anti-IL-23p19 antibody or IL-23p19 binding agent, by at least an hour and up to several months, for example at least an hour, five hours, 12 hours, a day, a week, a month, or three months, prior or subsequent to administration of the anti-IL-23p19 antibody or IL-23p19 binding agent.

[0209] The invention is further described in the following examples, which are not intended to limit the scope of the invention.

EXAMPLES

Example 1: Study

Study Design

[0210] This is a multi-centre, randomised, parallel assignment, double-blind, single-rising-dose, placebo-controlled, within-dose cohort trial. Following a 6-week screening period, patients were assigned to one of the two experimen-

tal arms. In the first arm, patients were randomised and allocated to one of six sequential dose cohorts (0.01, 0.05, 0.25, 1, 3, and 5 mg/kg) and randomised to receive a single dose of Antibody A or placebo intravenously (IV); four patients were included in each cohort (3:1 ratio; Antibody A:placebo). In the second arm, patients were randomised and allocated to one of two parallel dose groups (0.25 and 1 mg/kg) and randomised to receive Antibody A or placebo subcutaneously (SC); seven patients were planned for each cohort (6:1 ratio; Antibody A to placebo). Sample size was based on commonly used values for single-rising-dose studies and was not calculated according to hypotheses testing.

[0211] Antibody A IV dose groups were enrolled sequentially and Antibody A SC dose groups were randomised in parallel. Active versus placebo allocation within each cohort was double-blinded, but, for both IV and SC arms, cohorts were not blinded and were known to investigators and patients.

[0212] Central randomisation and blinded assignment of medication was performed via an interactive response tool. The randomisation list was generated using a validated system incorporating a pseudo-random number generator and supplied seed number. Access to randomisation codes was controlled and documented.

[0213] All patients were followed up for 24 weeks. A protocol amendment permitted patients receiving Antibody A SC who had at least a 50% improvement in psoriasis area severity index ($\geq$ PASI 50) from baseline to Week 24 to be offered enrolment into an optional observational follow-up period.

Patients

[0214] Eligible patients were aged 18-75 years, with chronic moderate-to-severe plaque psoriasis, lasting $\geq$ 6 months, with BSA involvement of $\geq$ 10, PASI $\geq$ 12, and Static Physicians Global Assessment (sPGA) score of moderate and above. Female patients were not of childbearing potential and patients had a Body Mass Index of $\geq$ 18.5 but $<$ 40.

[0215] Key exclusion criteria included patients who had received prior treatment with: ustekinumab within 24 weeks, other biologic agents or psoralen and ultraviolet A (PUVA) within 12 weeks, UVB phototherapy and oral anti-psoriatic medications within 4 weeks, or topical anti-psoriasis medications (except emollients) within 2 weeks prior to study dosing. Patients with current or previous clinically significant psoriatic disease, medical condition other than psoriasis or medical examination finding were also excluded from this study on the basis of the investigator's clinical judgement. Patients with chronic or relevant acute infections including hepatitis and tuberculosis (or a positive interferon- γ release assay at screening) were also excluded from the study.

Study Medication and Dosage

[0216] For IV administration, Antibody A (10 mg/mL) or placebo solution was injected into a 100 mL IV infusion bag of normal saline solution. Antibody A (90 mg/mL) or placebo pre-filled syringes were administered SC without dilution; excess volume was discarded.

Endpoints

[0217] The primary endpoint was safety and tolerability of Antibody A and was assessed descriptively based on physical examination, vital signs, electrocardiogram (ECG), clinical laboratory tests, adverse event (AE) reporting, and assessment of local and global tolerability by the investigator. Secondary clinical endpoints included PASI percentage

change from baseline and sPGA at Weeks 4, 8, 12, and 24. Patient photographs were obtained as part of the study protocol. Secondary pharmacokinetic endpoints were also evaluated, along with pharmacodynamic endpoints.

Histopathology, Immunohistochemistry, and RNA-Sequencing

[0218] Skin 6 mm punch biopsy samples were obtained from all patients at Baseline from a representative psoriatic lesion and at Week 8 from the same area. Biopsies were processed for histopathology, immunohistochemistry, and RNA-sequencing analysis.

[0219] Tissue sections were stained. Histopathology was based on haematoxylin and eosin staining. Sections were also stained with specific antibodies and evaluated semi-quantitatively for the presence of keratinocyte-related markers (K16 and Ki67), markers associated with tissue inflammation (5-100A7, neutrophil gelatinase lipocalin, β -defensin 2), dendritic cell lysosome associated membrane glycoprotein (DC-LAMP), CD11c (dermal dendritic cell marker), and CD3 (activate T-cell marker). In addition, RNA was extracted from skin biopsy samples for subsequent global transcriptome-wide RNA sequencing using the Illumina Hi-Seq 2000 (Illumina Inc., San Diego, Calif.).

Statistical Analysis

[0220] Comparisons of safety, clinical efficacy and biomarkers of Antibody A were summarised between Antibody A dose groups and placebo descriptively. In patients receiving rescue treatments after experiencing an intolerable increase in disease activity, efficacy endpoints assessed after administration of restricted medications were censored or last pre-rescue observation was carried forward (continuous).

[0221] A generalised linear model on the RNA-sequence count data was fit using empirical Bayes estimates of the dispersion (variance). The model included different covariates, yielding estimates of the adjusted fold change and p-value for each transcript. The false discovery rate was computed for each transcript and a p-value threshold determined. Gene expression fold changes for each patient dataset were calculated according to the following equation: (Antibody A—treatment value at Week 8/value at Baseline)/(Placebo-treatment value at Week 8/value at Baseline).

[0222] The Gene Ontology (GO) pathway sub-analysis used an approach, which estimates the enrichment ratio and p-value for gene sets relative to the background (all human genes, Huang et al, Genome Biol. 2007; 8:R183. Published microarray data were downloaded from the Gene Expression Omnibus database (<http://www.ncbi.nlm.nih.gov/geo>) and a

robust multi-array average normalisation used. Empirical Bayes t-tests/ANOVA models computed fold changes and p-values were calculated for each gene.

Results

[0223] Of 73 patients screened, 39 were eligible; 24 were randomised to six Antibody A IV cohorts and 14 to two Antibody A SC cohorts. One patient originally randomised to receive 1 mg/kg SC inadvertently received the 0.25 mg/kg dose. An additional patient was, therefore, assigned to the 1 mg/kg cohort, so that seven and six patients were included in the 0.25 and 1 mg/kg SC cohort, respectively. All 39 patients received a single dose of Antibody A or placebo and completed the 24-week follow-up except one patient in the 5 mg/kg IV group who discontinued follow-up after Week 12 due to relocation.

[0224] Baseline demographics were similar across all treatment groups; patients had similar duration of disease and were comparable within the SC and IV cohorts. Mean PASI scores ranged from 157 to 22.8, with individual subject scores at randomisation ranging from 10.5 to 43.4.

[0225] Overall, in patients receiving IV or SC Antibody A, a 75, 90, and 100% decrease in PASI was achieved by 87% (27/31), 58% (18/31), and 16% (5/31) of patients at Week 12, respectively; no patients receiving placebo achieved these improvements in PASI score (Table 6). At Week 24, PASI 75 was achieved by 68% of Antibody A patients, overall (22/31); only one placebo patient (13%) achieved PASI 75 at Week 24. PASI 90 and 100 rates at Week 24 were achieved by 48% (15/31) and 29% (9/31) of Antibody A patients, respectively; no patients in the placebo cohort achieved these PASI improvements. Within 24 weeks of the single Antibody A dose, 96%, 84%, and 64% of patients in Antibody A dose groups ≥ 0.25 mg/kg (n=25) achieved PASI 75, PASI 90, and PASI 100, respectively.

[0226] Assessments of sPGA were consistent with PASI; all SC Antibody A patients achieving “clear” or “almost clear” at both Weeks 12 and 24 (Table 6) and clinical improvements were confirmed by lesion photographs.

[0227] Improvements in PASI scores in Antibody A cohorts were observed as early as Week 2. Further PASI score improvements were observed through to Week 8 and were maintained until Week 24 (FIG. 1). All SC Antibody A patients had PASI 75 at Week 24. A protocol amendment extended their follow-up beyond Week 24 to evaluate the durability of this response to Antibody A. Eight of 13 patients who received SC Antibody A were eligible and chose to enroll into this optional observational follow-up. Six of eight patients maintained PASI 100 at a range of 41 to 66 weeks after a single dose (data not shown).

TABLE 6

PASI score improvements and sPGA classification rates at Weeks 12 and 24									
Method of administration of Antibody A									
IV									
SC									
Antibody A dose	mg/kg	mg/kg	mg/kg	mg/kg	mg/kg	mg/kg	0.25 mg/kg	1 mg/kg	mg/kg
N	3	3	3	3	3	2	7		6
PASI 75									
Week 12	3	1	3	2	3	3	6		6
Week 24	1	0	3	2	2	1 ^a	7		6
PASI 90									
Week 12	3	0	2	2	2	2	4		3
Week 24	0	0	1	2	1	1 ^a	5		6

TABLE 6-continued

PASI score improvements and sPGA classification rates at Weeks 12 and 24									
PASI 100	Week 12	0	0	0	1	0	1	2	1
	Week 24	0	0	1	1	0	1 ^a	3	4
sPGA	Week 12	3	1	2	2	3	3	7	6
“clear” or “almost clear”	Week 24	1	0	1	2	2	1 ^a	7	6

Antibody A dose		Overall Antibody A		Placebo	
N		31		8	
PASI 75	Week 12	27		0	
	Week 24	22		1	
PASI 90	Week 12	18		0	
	Week 24	15		0	
PASI 100	Week 12	5		0	
	Week 24	9		0	
sPGA	Week 12	27		0	
“clear” or “almost clear”	Week 24	20		0	

[0228] Number of patients in each treatment group achieving PASI 75-100 and a favourable sPGA at Week 12 and Week 24. N values represent the number of patients per group at Baseline; all patients were analysed at Week 12 and Week 24, except where indicated. Overall Antibody A values are given as totals for the Antibody A groups.

[0229] ^aN=2 due to relocation of a patient in the 5 mg/kg group.

[0230] IV, intravenous; PASI, Psoriasis Area Severity Index; SC, subcutaneous; sPGA, Static Physicians Global Assessment.

Histopathology, Immunohistochemistry, and RNA-Sequencing

[0231] Single doses of Antibody A resulted in marked reductions from Baseline to Week 8 in hyperkeratosis with parakeratosis, epidermal acanthosis, and generalised inflammation in the dermis and epidermis. Treatment with Antibody A resulted in decreases in markers associated with thickening of the keratinocyte layer (K16) and hyperproliferation (Ki67), dermal infiltration by T-cells (CD3), neutro-

phils (neutrophil gelatinase lipocalin), dendritic cells (CD11c and DC-LAMP), and tissue inflammation (β -defensin 2, 5-100A8/A9) at Week 8. These changes in the skin biopsy markers were primarily observed in patients from IV and SC Antibody A dose cohorts ≥ 0.25 mg/kg compared with placebo, across the different markers.

[0232] Treatment with Antibody A also resulted in significant (>2 fold) reductions ($p < 0.005$) in the expression of 199 genes (IV) and 226 genes (SC) relative to placebo in patient skin biopsies (comparing Baseline to Week 8). The overlap between these two cohorts was 192 genes. This gene set included genes associated with the IL-23/IL-17 axis (IL-23A, IL-23R, IL-22, IL-22RA1 RA2, IL-17A, IL-17F, IL-17RA, and IL-17RC), keratinocyte and epithelial cell differentiation (late cornified envelope protein, transglutaminase 1, and cornifelin), tissue inflammation (β -defensin 2, neutrophil gelatinase lipocalin, and S-100A7/A8), and the IFN α pathway (IFIH1, ISG15, IRF7, 1F144, MX1, MX2, STAT1, and TRIM22). GO pathway analysis also showed significant modulation of the pathways represented by these genes (Table 7).

TABLE 7

Summary of GO: Pathway analysis of the 79 gene cluster post-treatment with Antibody A				
Term	Count	p p-value	Genes	Fold enrichment
GO: 0009615~response to virus	9	2.48E-08	IFIH1, ISG15, IRF7, RSAD2, IFI44, MX1, STAT1, MX2, TRIM22	18.31133
GO: 0031424~keratinization	4	9.02E-04	LCE3D, TGM1, CNFN, LCE3E	20.62981
GO: 0030216~keratinocyte differentiation	4	0.003113	LCE3D, TGM1, CNFN, LCE3E	13.44064
GO: 0030855~epithelial cell differentiation	5	0.003166	LCE3D, TGM1, CNFN, DHRS9, LCE3E	8.093814
GO: 0009913~epidermal cell differentiation	4	0.003981	LCE3D, TGM1, CNFN, LCE3E	12.32058
GO: 0009310~amine catabolic process	4	0.004984	ARG1, KYNU, HAL, SMOX	11.37285
REACT_13: Metabolism of amino acids	4	0.00735	ARG1, KYNU, HAL, SMOX	8.33865
GO: 0009063~cellular amino acid catabolic process	3	0.03655	ARG1, KYNU, HAL	9.783992

TABLE 7-continued

Summary of GO: Pathway analysis of the 79 gene cluster post-treatment with Antibody A				
Term	Count	p p-value	Genes	Fold enrichment
hsa04622: RIG-I-like receptor signaling pathway	3	0.043535	IFIH1, ISG15, IRF7	8.594366

[0233] Enriched RNAs following treatment with Antibody A were identified relative to background using a previously described methodology {Huang et al. Genome Biol 2007}. GO pathways were identified using published microarray data (<http://www.ncbi.nlm.nih.gov/geo>) and a robust multi-array average normalisation. Empirical Bayes t-tests or ANOVA models computed fold changes and p-values were calculated for each gene {Huang et al. Genome Biol 2007}. HGNC gene names are used.

[0234] GO, Gene ontology; RNA, ribonucleic acid; HGNC, HUGO Gene Nomenclature Committee.

[0235] Of the Antibody A-responsive gene set, expression of a subset of 79 genes was decreased following treatment with Antibody A and significantly correlated with PASI scores at Week 8 ($r=0.73$, $p=2 \times 10^{-6}$). Reductions in IL-23A and IL-23R expression were also significantly correlated with PASI scores at Week 8 ($r=0.57$ and $r=0.54$; $p<0.005$). In addition, Antibody A treatment led to normalisation of a gene set previously reported to distinguish lesional and non-lesional tissue from psoriasis patients. Importantly, this gene set had significant overlap (91%) with the Antibody A-responsive gene set that correlated with PASI scores.

[0236] The results are summarized in Table 8, which indicates the 79 gene transcripts, full names and fold changes (p values). Gene expression fold changes for each patient dataset were calculated according to the following equation: (Antibody A-treatment value at Week 8/value at Baseline)/(Placebo-treatment value at Week 8/value at Baseline).

TABLE 8

Transcript	Trt FC	p.value	Name
PRSS22	0.22	0.001309	protease, serine, 22
TYMP	0.187	0.001765	thymidine phosphorylase
MXD1	0.516	0.000498	MAX dimerization protein 1
DHRS9	0.333	0.003122	dehydrogenase/reductase (SDR family) member 9
ATP12A	0.131	0.002618	ATPase, H+/K+ transporting, nongastric, alpha polypeptide
HAL	0.348	0.006686	histidine ammonia-lyase
SMOX	0.368	0.000122	spermine oxidase
TGM1	0.29	5.52E-05	transglutaminase 1 (K polypeptide epidermal type I, protein-glutamine-gamma-glutamyltransferase)
VNN3	0.229	0.000265	vanin 3
PLA2G3	0.474	0.002278	phospholipase A2, group III
SLC5A1	0.4	0.001253	solute carrier family 5 (sodium/glucose cotransporter), member 1
APOL1	0.308	0.011199	apolipoprotein L, 1
GDPD3	0.429	0.000346	glycerophosphodiester phosphodiesterase domain containing 3

TABLE 8-continued

Transcript	Trt FC	p.value	Name
SMPD3	0.38	0.000115	sphingomyelin phosphodiesterase 3, neutral membrane (neutral sphingomyelinase II)
SQLE	0.503	0.000211	squalene epoxidase
CNFN	0.239	1.52E-05	cornifelin
HYAL4	0.218	4.60E-07	hyaluronoglucosaminidase 4
OAS3	0.29	0.001929	2'-5'-oligoadenylate synthetase 3, 100 kDa
OAS2	0.192	0.003705	2'-5'-oligoadenylate synthetase 2, 69/71 kDa
HRH2	0.373	0.002386	histamine receptor H2
IFIH1	0.425	0.008001	interferon induced with helicase C domain 1
STAT1	0.459	0.012805	signal transducer and activator of transcription 1, 91 kDa
KYNU	0.249	0.001811	kynureninase
FLVCR2	0.44	0.000454	feline leukemia virus subgroup C cellular receptor family, member 2
IFIT3	0.291	0.002822	interferon-induced protein with tetratricopeptide repeats 3
NT5C3	0.421	0.000717	5'-nucleotidase, cytosolic III
IFI6	0.103	0.000325	interferon, alpha-inducible protein 6
KLK10	0.336	0.000902	kallikrein-related peptidase 10
KLHDC7B	0.304	0.000148	kelch domain containing 7B
SCO2	0.339	0.004546	SCO2 cytochrome c oxidase assembly protein
HELZ2	0.365	0.000531	helicase with zinc finger 2, transcriptional coactivator
TRIM22	0.422	0.0375	tripartite motif containing 22
EPSTI1	0.26	0.003714	epithelial stromal interaction 1 (breast)
RSAD2	0.225	0.001466	radical S-adenosyl methionine domain containing 2
CMPK2	0.255	0.001387	cytidine monophosphate (UMP-CMP) kinase 2, mitochondrial
TCN1	0.083	0.000784	transcobalamin I (vitamin B12 binding protein, R binder family)
OASL	0.122	5.35E-05	2'-5'-oligoadenylate synthetase-like
DDX60	0.407	0.009311	DEAD (Asp-Glu-Ala-Asp) box polypeptide 60 ('DEAD (Asp-Glu-Ala-Asp)' disclosed as SEQ ID NO: 23)
IFI44L	0.211	0.005291	interferon-induced protein 44-like
IFI44	0.283	0.012865	interferon-induced protein 44
PARP9	0.391	0.010339	poly (ADP-ribose) polymerase family, member 9
HERC6	0.189	0.001701	HECT and RLD domain containing E3 ubiquitin protein ligase family member 6
EPHA2	0.438	5.27E-06	EPH receptor A2
UBE2L6	0.418	0.002472	ubiquitin-conjugating enzyme E2L6
MX1	0.185	0.000774	myxovirus (influenza virus) resistance 1, interferon-inducible protein p78 (mouse)
PLA2G2F	0.349	0.000271	phospholipase A2, group IIF
HK2	0.527	0.021317	hexokinase 2
GBAP1	0.366	0.000729	glucosidase, beta, acid pseudogene 1
PDZK1IP1	0.349	0.000932	PDZK1 interacting protein 1
LCE3D	0.144	0.004114	late cornified envelope 3D

TABLE 8-continued

Transcript	Ttt FC	p.value	Name
ZC3H12A	0.236	0.000427	zinc finger CCCH-type containing 12A
IFI27	0.289	0.006965	interferon, alpha-inducible protein 27
ANGPTL4	0.318	0.003141	angiopoietin-like 4
FUT3	0.342	0.001375	fucosyltransferase 3 (galactoside 3(4)-L-fucosyltransferase, Lewis blood group)
LRG1	0.237	0.001718	leucine-rich alpha-2-glycoprotein 1
PRSS27	0.192	0.00072	protease, serine 27
HPSE	0.211	0.000233	heparanase
FBXO45	0.507	0.002771	F-box protein 45
SLC26A9	0.282	0.000288	solute carrier family 26, member 9
TREX2	0.294	1.04E-05	three prime repair exonuclease 2
MX2	0.317	0.00491	myxovirus (influenza virus) resistance 2 (mouse)
PAPL	0.197	0.000138	iron/zinc purple acid phosphatase-like protein
USP18	0.364	0.000984	ubiquitin specific peptidase 18
FAM43A	0.382	4.01E-05	family with sequence similarity 43, member A
IRF7	0.272	3.53E-05	interferon regulatory factor 7
IFIT1	0.211	0.003096	interferon-induced protein with tetratricopeptide repeats 1
LCE3E	0.134	0.001434	late cornified envelope 3E
ISG15	0.124	0.000226	ISG15 ubiquitin-like modifier
C10orf99	0.279	0.000848	chromosome 10 open reading frame 99
RNF222	0.439	6.77E-05	ring finger protein 222
GM2A	0.398	0.000122	GM2 ganglioside activator
SERPINB13	0.331	0.00069	serpin peptidase inhibitor, clade B (ovalbumin), member 13
HIST1H4I	0.651	0.000158	histone cluster 1, H4i
FCHSD1	0.387	0.000306	FCH and double SH3 domains 1
KLK9	0.219	0.000436	kallikrein-related peptidase 9
CCDC19	0.5	6.02E-05	coiled-coil domain containing 19
LOC729041	0.337	0.002125	fatty acid amide hydrolase pseudogene
PCP4L1	0.449	0.015145	Purkinje cell protein 4 like 1
KLK9	0.201	0.008977	kallikrein-related peptidase 9

Example 2: Pharmaceutical Compositions

[0237] Examples of formulations suitable for an antibody of the present invention are shown below. Antibodies used in the formulations below are for example Antibody A, Antibody B, Antibody C or Antibody D.

Formulation 1:

[0238]

Components	Concentration [mmol/L]	Concentration [g/l]	Nominal Amount [mg/vial] V = 10.0 ml
Antibody		10.0	100.0
Succinic acid	0.7	0.083	0.8
Disodium succinate hexahydrate	24.3	6.564	65.6
Sodium chloride	125	7.305	73.1
Polysorbat 20	0.16	0.20	0.20
Water for Injection	—	Ad 1 L	Ad 1 mL

[0239] The pH of formulation 1 is typically in the range of pH 6.0 to 7.0, for example pH 6.5. This formulation is particularly suitable for intravenous administration.

[0240] Molecular weight (MW in g/mol) of used excipients: Disodium succinate hexahydrate=270.14 g/mol; Succinic acid=118.09 g/mol; Sodium chloride=58.44 g/mol.

[0241] The osmolality of the formulation is 300+/-30 mOsmol/kg, as determined using an Osmomat 030 (Gonotec GmbH, Berlin, Germany). The density at 20° C. of the formulation is approximately 1.0089 g/cm³, as determined using a measuring unit DMA 4500 (Anton Paar GmbH, Ostfildern-Scharnhausen, Germany).

Formulation 2:

[0242]

Components	Concentration [mmol/L]	Concentration [g/l]	Nominal Amount [mg/syringe] V = 1.0 ml
Antibody	0.6	90.0	90.0
Succinic acid	0.5	0.059	0.059
Disodium succinate hexahydrate	3.9	1.054	1.054
Sorbitol	225	41.00	41.00
Polysorbat 20	0.16	0.20	0.20
Water for Injection	—	Ad 1 L	Ad 1 mL

[0243] The pH of formulation 2 is typically in the range of pH 5.5 to 6.5, for example 5.5 to 6.1, for example the pH is 5.8. This formulation is particularly suitable for subcutaneous administration.

[0244] Molecular weight (MW in g/mol) of used excipients:

[0245] MW: Succinic acid (C₄H₆O₄)=118.09 g/mol

[0246] MW: Disodium succinate hexahydrate (C₄O₄Na₂H₄×6H₂O)=270.14 g/mol

[0247] MW: Sorbitol=182.17 g/mol

[0248] MW: Polysorbate 20=1227.72 g/mol

[0249] The osmolality of the formulation is 300+/-30 mOsmol/kg, as determined using an Osmomat 030 (Gonotec GmbH, Berlin, Germany). The density at 20° C. of the formulation is approximately 1.040 g/cm³, as determined using a measuring unit DMA 4500 (Anton Paar GmbH, Ostfildern-Scharnhausen, Germany).

Formulation 3:

[0250]

Components	Concentration [mmol/L]	Concentration [g/l]	Nominal Amount [mg/syringe] V = 1.0 ml
Antibody	0.6	90.0	90.0
Sorbitol	240	43.733	43.733
Polysorbat 20	0.16	0.20	0.20
Water for Injection	—	Ad 1 L	Ad 1 mL

[0251] The pH of formulation 3 is typically in the range of pH 5.5 to 6.5, for example 5.5 to 6.1, for example the pH is 5.8. This formulation is particularly suitable for subcutaneous administration.

[0252] Molecular weight (MW in g/mol) of used excipients:

[0253] MW: Sorbitol=182.17 g/mol

[0254] MW: Polysorbate 20=1227.72 g/mol.

[0255] The osmolality of the formulation is 300+/-30 mOsmol/kg, as determined using an Osmomat 030 (Gonotec GmbH, Berlin, Germany).

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 23

<210> SEQ ID NO 1

<211> LENGTH: 11

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<221> NAME/KEY: source

<223> OTHER INFORMATION: /note="Description of Artificial Sequence:
Synthetic peptide"

<400> SEQUENCE: 1

Lys Ala Ser Arg Asp Val Ala Ile Ala Val Ala
1 5 10

<210> SEQ ID NO 2

<211> LENGTH: 7

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<221> NAME/KEY: source

<223> OTHER INFORMATION: /note="Description of Artificial Sequence:
Synthetic peptide"

<400> SEQUENCE: 2

Trp Ala Ser Thr Arg His Thr
1 5

<210> SEQ ID NO 3

<211> LENGTH: 9

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<221> NAME/KEY: source

<223> OTHER INFORMATION: /note="Description of Artificial Sequence:
Synthetic peptide"

<400> SEQUENCE: 3

His Gln Tyr Ser Ser Tyr Pro Phe Thr
1 5

<210> SEQ ID NO 4

<211> LENGTH: 10

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<221> NAME/KEY: source

<223> OTHER INFORMATION: /note="Description of Artificial Sequence:
Synthetic peptide"

<400> SEQUENCE: 4

Gly Asn Thr Phe Thr Asp Gln Thr Ile His
1 5 10

<210> SEQ ID NO 5

<211> LENGTH: 17

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<221> NAME/KEY: source

<223> OTHER INFORMATION: /note="Description of Artificial Sequence:
Synthetic peptide"

<400> SEQUENCE: 5

Tyr Ile Tyr Pro Arg Asp Asp Ser Pro Lys Tyr Asn Glu Asn Phe Lys
1 5 10 15

-continued

Gly

<210> SEQ ID NO 6
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<221> NAME/KEY: source
<223> OTHER INFORMATION: /note="Description of Artificial Sequence:
Synthetic peptide"

<400> SEQUENCE: 6

Pro Asp Arg Ser Gly Tyr Ala Trp Phe Ile Tyr
1 5 10

<210> SEQ ID NO 7
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<221> NAME/KEY: source
<223> OTHER INFORMATION: /note="Description of Artificial Sequence:
Synthetic peptide"

<400> SEQUENCE: 7

Gly Tyr Thr Phe Thr Asp Gln Thr Ile His
1 5 10

<210> SEQ ID NO 8
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<221> NAME/KEY: source
<223> OTHER INFORMATION: /note="Description of Artificial Sequence:
Synthetic peptide"

<400> SEQUENCE: 8

Gly Phe Thr Phe Thr Asp Gln Thr Ile His
1 5 10

<210> SEQ ID NO 9
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<221> NAME/KEY: source
<223> OTHER INFORMATION: /note="Description of Artificial Sequence:
Synthetic peptide"

<400> SEQUENCE: 9

Gly Gly Thr Phe Thr Asp Gln Thr Ile His
1 5 10

<210> SEQ ID NO 10
<211> LENGTH: 107
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<221> NAME/KEY: source
<223> OTHER INFORMATION: /note="Description of Artificial Sequence:
Synthetic polypeptide"

<400> SEQUENCE: 10

Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly

-continued

```

1           5           10           15
Asp Arg Val Thr Ile Thr Cys Lys Ala Ser Arg Asp Val Ala Ile Ala
                20                25                30

Val Ala Trp Tyr Gln Gln Lys Pro Gly Lys Val Pro Lys Leu Leu Leu
                35                40                45

Phe Trp Ala Ser Thr Arg His Thr Gly Val Pro Asp Arg Phe Ser Gly
                50                55                60

Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro
        65                70                75                80

Glu Asp Leu Ala Asp Tyr Tyr Cys His Gln Tyr Ser Ser Tyr Pro Phe
                85                90                95

Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys
        100                105

<210> SEQ ID NO 11
<211> LENGTH: 107
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<221> NAME/KEY: source
<223> OTHER INFORMATION: /note="Description of Artificial Sequence:
        Synthetic polypeptide"

<400> SEQUENCE: 11
Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
1           5           10           15

Asp Arg Val Thr Ile Thr Cys Lys Ala Ser Arg Asp Val Ala Ile Ala
                20                25                30

Val Ala Trp Tyr Gln Gln Lys Pro Gly Lys Val Pro Lys Leu Leu Ile
                35                40                45

Tyr Trp Ala Ser Thr Arg His Thr Gly Val Pro Ser Arg Phe Ser Gly
                50                55                60

Ser Gly Ser Arg Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro
        65                70                75                80

Glu Asp Val Ala Asp Tyr Phe Cys His Gln Tyr Ser Ser Tyr Pro Phe
                85                90                95

Thr Phe Gly Ser Gly Thr Lys Leu Glu Ile Lys
        100                105

<210> SEQ ID NO 12
<211> LENGTH: 107
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<221> NAME/KEY: source
<223> OTHER INFORMATION: /note="Description of Artificial Sequence:
        Synthetic polypeptide"

<400> SEQUENCE: 12
Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
1           5           10           15

Asp Arg Val Thr Ile Thr Cys Lys Ala Ser Arg Asp Val Ala Ile Ala
                20                25                30

Val Ala Trp Tyr Gln Gln Lys Pro Gly Lys Val Pro Lys Leu Leu Leu
                35                40                45

Tyr Trp Ala Ser Thr Arg His Thr Gly Val Pro Ser Arg Phe Ser Gly
                50                55                60

```

-continued

Ser Gly Ser Arg Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro
 65 70 75 80

Glu Asp Val Ala Thr Tyr Tyr Cys His Gln Tyr Ser Ser Tyr Pro Phe
 85 90 95

Thr Phe Gly Ser Gly Thr Lys Leu Glu Ile Lys
 100 105

<210> SEQ ID NO 13
 <211> LENGTH: 107
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <221> NAME/KEY: source
 <223> OTHER INFORMATION: /note="Description of Artificial Sequence:
 Synthetic polypeptide"

<400> SEQUENCE: 13

Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
 1 5 10 15

Asp Arg Val Thr Ile Thr Cys Lys Ala Ser Arg Asp Val Ala Ile Ala
 20 25 30

Val Ala Trp Tyr Gln Gln Lys Pro Gly Lys Val Pro Lys Leu Leu Leu
 35 40 45

Phe Trp Ala Ser Thr Arg His Thr Gly Val Pro Asp Arg Phe Ser Gly
 50 55 60

Ser Gly Ser Arg Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro
 65 70 75 80

Glu Asp Leu Ala Asp Tyr Tyr Cys His Gln Tyr Ser Ser Tyr Pro Phe
 85 90 95

Thr Phe Gly Ser Gly Thr Lys Leu Glu Ile Lys
 100 105

<210> SEQ ID NO 14
 <211> LENGTH: 120
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <221> NAME/KEY: source
 <223> OTHER INFORMATION: /note="Description of Artificial Sequence:
 Synthetic polypeptide"

<400> SEQUENCE: 14

Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ser
 1 5 10 15

Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asp Gln
 20 25 30

Thr Ile His Trp Met Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Ile
 35 40 45

Gly Tyr Ile Tyr Pro Arg Asp Asp Ser Pro Lys Tyr Asn Glu Asn Phe
 50 55 60

Lys Gly Lys Val Thr Ile Thr Ala Asp Lys Ser Thr Ser Thr Ala Tyr
 65 70 75 80

Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95

Ala Ile Pro Asp Arg Ser Gly Tyr Ala Trp Phe Ile Tyr Trp Gly Gln
 100 105 110

-continued

Gly Thr Leu Val Thr Val Ser Ser
115 120

<210> SEQ ID NO 15
<211> LENGTH: 120
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<221> NAME/KEY: source
<223> OTHER INFORMATION: /note="Description of Artificial Sequence:
Synthetic polypeptide"

<400> SEQUENCE: 15

Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ser
1 5 10 15
Ser Val Lys Val Ser Cys Lys Ala Ser Gly Phe Thr Phe Thr Asp Gln
20 25 30
Thr Ile His Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met
35 40 45
Gly Tyr Ile Tyr Pro Arg Asp Asp Ser Pro Lys Tyr Asn Glu Asn Phe
50 55 60
Lys Gly Lys Val Thr Leu Thr Ala Asp Lys Ser Thr Ser Thr Ala Tyr
65 70 75 80
Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95
Ala Ile Pro Asp Arg Ser Gly Tyr Ala Trp Phe Ile Tyr Trp Gly Gln
100 105 110
Gly Thr Leu Val Thr Val Ser Ser
115 120

<210> SEQ ID NO 16
<211> LENGTH: 120
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<221> NAME/KEY: source
<223> OTHER INFORMATION: /note="Description of Artificial Sequence:
Synthetic polypeptide"

<400> SEQUENCE: 16

Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ser
1 5 10 15
Ser Val Lys Thr Ser Cys Lys Ala Ser Gly Gly Thr Phe Thr Asp Gln
20 25 30
Thr Ile His Trp Val Arg Gln Arg Pro Gly Gln Gly Leu Glu Trp Met
35 40 45
Gly Tyr Ile Tyr Pro Arg Asp Asp Ser Pro Lys Tyr Asn Glu Asn Phe
50 55 60
Lys Gly Arg Val Thr Ile Thr Ala Asp Lys Ser Thr Ser Thr Ala Tyr
65 70 75 80
Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95
Ala Ile Pro Asp Arg Ser Gly Tyr Ala Trp Phe Ile Tyr Trp Gly Gln
100 105 110
Gly Thr Leu Val Thr Val Ser Ser
115 120

-continued

<210> SEQ ID NO 17
<211> LENGTH: 120
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<221> NAME/KEY: source
<223> OTHER INFORMATION: /note="Description of Artificial Sequence:
Synthetic polypeptide"

<400> SEQUENCE: 17

Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ser
1 5 10 15
Ser Val Lys Val Ser Cys Lys Ala Ser Gly Gly Thr Phe Thr Asp Gln
20 25 30
Thr Ile His Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met
35 40 45
Gly Tyr Ile Tyr Pro Arg Asp Asp Ser Pro Lys Tyr Asn Glu Asn Phe
50 55 60
Lys Gly Arg Val Thr Leu Thr Ala Asp Lys Ser Thr Ser Thr Ala Tyr
65 70 75 80
Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Phe Cys
85 90 95
Ala Arg Pro Asp Arg Ser Gly Tyr Ala Trp Phe Ile Tyr Trp Gly Gln
100 105 110
Gly Thr Leu Val Thr Val Ser Ser
115 120

<210> SEQ ID NO 18
<211> LENGTH: 214
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<221> NAME/KEY: source
<223> OTHER INFORMATION: /note="Description of Artificial Sequence:
Synthetic polypeptide"

<400> SEQUENCE: 18

Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
1 5 10 15
Asp Arg Val Thr Ile Thr Cys Lys Ala Ser Arg Asp Val Ala Ile Ala
20 25 30
Val Ala Trp Tyr Gln Gln Lys Pro Gly Lys Val Pro Lys Leu Leu Ile
35 40 45
Tyr Trp Ala Ser Thr Arg His Thr Gly Val Pro Ser Arg Phe Ser Gly
50 55 60
Ser Gly Ser Arg Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro
65 70 75 80
Glu Asp Val Ala Asp Tyr Phe Cys His Gln Tyr Ser Ser Tyr Pro Phe
85 90 95
Thr Phe Gly Ser Gly Thr Lys Leu Glu Ile Lys Arg Thr Val Ala Ala
100 105 110
Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly
115 120 125
Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala
130 135 140
Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln
145 150 155 160

-continued

Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser
 165 170 175
 Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr
 180 185 190
 Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser
 195 200 205
 Phe Asn Arg Gly Glu Cys
 210

<210> SEQ ID NO 19
 <211> LENGTH: 449
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <221> NAME/KEY: source
 <223> OTHER INFORMATION: /note="Description of Artificial Sequence:
 Synthetic polypeptide"

<400> SEQUENCE: 19

Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ser
 1 5 10 15
 Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asp Gln
 20 25 30
 Thr Ile His Trp Met Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Ile
 35 40 45
 Gly Tyr Ile Tyr Pro Arg Asp Asp Ser Pro Lys Tyr Asn Glu Asn Phe
 50 55 60
 Lys Gly Lys Val Thr Ile Thr Ala Asp Lys Ser Thr Ser Thr Ala Tyr
 65 70 75 80
 Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95
 Ala Ile Pro Asp Arg Ser Gly Tyr Ala Trp Phe Ile Tyr Trp Gly Gln
 100 105 110
 Gly Thr Leu Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val
 115 120 125
 Phe Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala
 130 135 140
 Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser
 145 150 155 160
 Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val
 165 170 175
 Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro
 180 185 190
 Ser Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys
 195 200 205
 Pro Ser Asn Thr Lys Val Asp Lys Arg Val Glu Pro Lys Ser Cys Asp
 210 215 220
 Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Ala Ala Gly Gly
 225 230 235 240
 Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile
 245 250 255
 Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu
 260 265 270

-continued

Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His
 275 280 285
 Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg
 290 295 300
 Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys
 305 310 315 320
 Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu
 325 330 335
 Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr
 340 345 350
 Thr Leu Pro Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu
 355 360 365
 Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp
 370 375 380
 Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val
 385 390 395 400
 Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp
 405 410 415
 Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His
 420 425 430
 Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro
 435 440 445

Gly

<210> SEQ ID NO 20
 <211> LENGTH: 449
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <221> NAME/KEY: source
 <223> OTHER INFORMATION: /note="Description of Artificial Sequence:
 Synthetic polypeptide"

<400> SEQUENCE: 20

Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ser
 1 5 10 15
 Ser Val Lys Val Ser Cys Lys Ala Ser Gly Phe Thr Phe Thr Asp Gln
 20 25 30
 Thr Ile His Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met
 35 40 45
 Gly Tyr Ile Tyr Pro Arg Asp Asp Ser Pro Lys Tyr Asn Glu Asn Phe
 50 55 60
 Lys Gly Lys Val Thr Leu Thr Ala Asp Lys Ser Thr Ser Thr Ala Tyr
 65 70 75 80
 Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95
 Ala Ile Pro Asp Arg Ser Gly Tyr Ala Trp Phe Ile Tyr Trp Gly Gln
 100 105 110
 Gly Thr Leu Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val
 115 120 125
 Phe Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala
 130 135 140
 Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser
 145 150 155 160

-continued

Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val
 165 170 175
 Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro
 180 185 190
 Ser Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys
 195 200 205
 Pro Ser Asn Thr Lys Val Asp Lys Arg Val Glu Pro Lys Ser Cys Asp
 210 215 220
 Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Ala Ala Gly Gly
 225 230 235 240
 Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile
 245 250 255
 Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu
 260 265 270
 Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His
 275 280 285
 Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg
 290 295 300
 Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys
 305 310 315 320
 Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu
 325 330 335
 Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr
 340 345 350
 Thr Leu Pro Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu
 355 360 365
 Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp
 370 375 380
 Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val
 385 390 395 400
 Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp
 405 410 415
 Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His
 420 425 430
 Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro
 435 440 445

Gly

<210> SEQ ID NO 21
 <211> LENGTH: 214
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <221> NAME/KEY: source
 <223> OTHER INFORMATION: /note="Description of Artificial Sequence:
 Synthetic polypeptide"

<400> SEQUENCE: 21

Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
 1 5 10 15
 Asp Arg Val Thr Ile Thr Cys Lys Ala Ser Arg Asp Val Ala Ile Ala
 20 25 30
 Val Ala Trp Tyr Gln Gln Lys Pro Gly Lys Val Pro Lys Leu Leu Leu

-continued

35	40	45
Phe Trp Ala Ser Thr Arg His Thr Gly Val Pro Asp Arg Phe Ser Gly		
50	55	60
Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro		
65	70	75
Glu Asp Leu Ala Asp Tyr Tyr Cys His Gln Tyr Ser Ser Tyr Pro Phe		
85	90	95
Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys Arg Thr Val Ala Ala		
100	105	110
Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly		
115	120	125
Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala		
130	135	140
Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln		
145	150	155
Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser		
165	170	175
Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr		
180	185	190
Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser		
195	200	205
Phe Asn Arg Gly Glu Cys		
210		

<210> SEQ ID NO 22

<211> LENGTH: 189

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 22

Met Leu Gly Ser Arg Ala Val Met Leu Leu Leu Leu Leu Pro Trp Thr		
1	5	10
Ala Gln Gly Arg Ala Val Pro Gly Gly Ser Ser Pro Ala Trp Thr Gln		
20	25	30
Cys Gln Gln Leu Ser Gln Lys Leu Cys Thr Leu Ala Trp Ser Ala His		
35	40	45
Pro Leu Val Gly His Met Asp Leu Arg Glu Glu Gly Asp Glu Glu Thr		
50	55	60
Thr Asn Asp Val Pro His Ile Gln Cys Gly Asp Gly Cys Asp Pro Gln		
65	70	75
Gly Leu Arg Asp Asn Ser Gln Phe Cys Leu Gln Arg Ile His Gln Gly		
85	90	95
Leu Ile Phe Tyr Glu Lys Leu Leu Gly Ser Asp Ile Phe Thr Gly Glu		
100	105	110
Pro Ser Leu Leu Pro Asp Ser Pro Val Gly Gln Leu His Ala Ser Leu		
115	120	125
Leu Gly Leu Ser Gln Leu Leu Gln Pro Glu Gly His His Trp Glu Thr		
130	135	140
Gln Gln Ile Pro Ser Leu Ser Pro Ser Gln Pro Trp Gln Arg Leu Leu		
145	150	155
Leu Arg Phe Lys Ile Leu Arg Ser Leu Gln Ala Phe Val Ala Val Ala		
165	170	175

-continued

Ala Arg Val Phe Ala His Gly Ala Ala Thr Leu Ser Pro
180 185

<210> SEQ ID NO 23
<211> LENGTH: 4
<212> TYPE: PRT
<213> ORGANISM: Unknown
<220> FEATURE:
<221> NAME/KEY: source
<223> OTHER INFORMATION: /note="Description of Unknown: 'DEAD' box motif peptide"

<400> SEQUENCE: 23

Asp Glu Ala Asp
1

1. A method for detecting the presence or absence of a beneficial response in a patient after administration of an IL-23A antagonist, comprising:

- a) obtaining a biological sample from the patient;
- b) measuring in said sample the level of one or more biomarkers, or the level of expression of one or more biomarkers;
- c) comparing the level to control value of the level of the biomarkers; and
- d) determining whether or not the difference in levels between the sample and the control reflects a beneficial response in the patient,

wherein the one or more biomarkers are genes/proteins associated with the IL-23/IL-17 axis, keratinocyte and epithelial cell differentiation, tissue inflammation and the IFN α pathway.

2. The method of claim 1, wherein the level of the gene or the protein of said one or more biomarker is measured.

3. The method of claim 1, wherein the patient suffers from psoriasis.

4. The method of claim 1, wherein the control value is calculated using samples from subjects that do not suffer from psoriasis.

5. The method of claim 1, wherein the control value is determined using samples from known psoriasis patients.

6. The method of claim 1, wherein the control value is determined using at least one previous sample taken from the patient.

7. The method of claim 1, wherein the one or more biomarkers is S-100A7, neutrophil gelatinase lipocalin or β -defensin 2.

8. The method of claim 1, wherein the biological sample is a skin biopsy, blood, plasma or serum sample.

9. The method of claim 1, wherein the IL-23A antagonist is an anti-IL-23A antibody or an antigen binding fragment thereof.

10. The method claim 1, wherein the levels of biomarkers are determined by RNA sequencing or ELISA.

11. The method according to claim 1, further comprising continuing the administration of the IL-23A antagonist to the patient if the difference in levels between the sample and the control reflects a beneficial response in the patient.

12. A method of determining whether a potential therapeutic agent is efficacious in the treatment of psoriasis comprising:

- a) obtaining a first biological sample from a psoriasis patient prior to being treated with the potential therapeutic agent;
 - b) treating the psoriasis patient with the potential therapeutic agent;
 - c) obtaining a second biological sample from the psoriasis patient after being treated with the potential therapeutic agent;
 - d) measuring in said first and second sample the levels of expression of one or more biomarkers; and
 - e) comparing the biomarker levels in the second sample to the levels in the first sample,
- wherein lower biomarker levels in the second sample than in the first sample indicate that the potential therapeutic agent is efficacious, and further wherein the one or more biomarkers are genes associated with the IL-23/IL-17 axis, keratinocyte and epithelial cell differentiation, tissue inflammation and the IFN α pathway.

13. The method according to claim 12, wherein lower biomarker levels in the second sample than in the first sample correlate with improvement in clinical efficacy measures.

14. The method according to claim 12, further comprising continuing the treatment of the patient if biomarker levels in the second sample is lower than the levels in the first sample.

15. A method of treating psoriasis in a subject comprising:

- a) determining whether to initiate treatment of the subject, modify the treatment dose, modify the dosing interval, or discontinue treatment, based on the method of any of the preceding claims; and
- b) modifying the treatment regimen based on the determination.

16. A method of monitoring patient response to a psoriasis treatment comprising:

- a) obtaining a first biological sample from the patient;
- b) measuring the level of one or more biomarkers in said first biological sample, wherein said one or more biomarkers are genes/proteins associated with the IL-23/IL-17 axis, keratinocyte and epithelial cell differentiation, tissue inflammation and the IFN α pathway;
- c) administering a treatment compound to the patient;
- d) obtaining a second biological sample from the patient;
- e) measuring the level of said one or more biomarkers in said second biological sample; and
- f) comparing the levels of the one or more biomarkers obtained from first and second biological samples;

wherein a decreased level of the one or more biomarkers in the second biological sample indicates an effective response.

17. A method for monitoring patient compliance with a drug treatment protocol for psoriasis comprising:

- a) obtaining a biological sample from said patient;
- b) measuring the level of one or more biomarkers, wherein the one or more biomarkers are genes/proteins associated with the IL-23/IL-17 axis, keratinocyte and epithelial cell differentiation, tissue inflammation and the IFN α pathway; and
- c) determining if the level is decreased in the patient sample compared to the level in a control untreated sample;

wherein a decreased level indicates patient compliance with said drug treatment protocol.

18. The method of claim **15**, wherein the level of the one or more biomarkers in the second biological sample is

decreased by at least about 50%, 55%, 60%, 65%, 65%, 70%, 75%, 80%, 85%, 90%, or 95% or more as compared to the level in the first biological sample.

19. The method of claim **15**, wherein the biological sample is a skin biopsy, blood, plasma or serum sample.

20. The method of claim **15**, wherein the IL-23A antagonist is an anti-IL-23A antibody or an antigen binding fragment thereof.

21. The method of claim **15**, wherein the levels of biomarkers are determined by RNA sequencing or ELISA.

22. An ELISA kit comprising one or more antibodies or antigen binding fragments thereof that specifically bind to one or more biomarkers selected from the the group consisting of S-100 proteins, neutrophil gelatinase lipocalin and β -defensin 2.

23. The ELISA kit of claim **22**, further comprising instructions for use of the kit in monitoring psoriasis.

* * * * *

专利名称(译)	生物标志物可用于治疗il-23a相关疾病		
公开(公告)号	US20170157246A1	公开(公告)日	2017-06-08
申请号	US15/327144	申请日	2015-07-23
[标]申请(专利权)人(译)	百灵佳殷格翰国际股份有限公司		
申请(专利权)人(译)	勃林格殷格翰国际GMBH		
当前申请(专利权)人(译)	勃林格殷格翰国际GMBH		
[标]发明人	VISVANATHAN SUDHA BAUM PATRICK KLEINER OLIVER KUNZ ULRICH		
发明人	VISVANATHAN, SUDHA BAUM, PATRICK KLEINER, OLIVER KUNZ, ULRICH		
IPC分类号	A61K39/395 G01N33/53		
CPC分类号	A61K39/3955 A61K2121/00 G01N33/5308 G01N33/53 A61P1/04 A61P17/06 A61P19/02 A61P43/00 C12Q1/6883 C12Q2600/106 C12Q2600/158 G01N33/6863 G01N33/6869		
优先权	62/028379 2014-07-24 US		
其他公开文献	US10507241		
外部链接	Espacenet USPTO		

摘要(译)

本发明一般涉及用于治疗IL-23相关疾病，特别是炎性疾病如牛皮癣的生物标志物。本发明还涉及使用本文公开的生物标志物的方法，例如在利用IL-23拮抗剂的疗法中。

