



(19) **United States**

(12) **Patent Application Publication** (10) **Pub. No.: US 2002/0161191 A1**
Feder et al. (43) **Pub. Date: Oct. 31, 2002**

(54) **NOVEL IMIDAZOLINE RECEPTOR
HOMOLOGS**

(52) **U.S. Cl.** **530/350**

(76) **Inventors: John N. Feder**, Belle Mead, NJ (US);
Gabe Mintier, Hightstown, NJ (US);
Gene G. Kinney, Collegeville, PA
(US); **Chandra S. Ramanathan**,
Wallingford, CT (US)

(57) **ABSTRACT**

Correspondence Address:
Christopher A. Klein, Esq.
Bristol-Myers Squibb Company
Post Office Box 4000
Lawrenceville-Provinceline Road
Princeton, NJ 08543-4000 (US)

Novel imidazoline receptor homologs, designated imidazoline receptor related protein 1 (IMRRP1), imidazoline receptor related protein 1b (IMRRP1b) and derivatives thereof are described. Pharmaceutical compositions comprising at least one IMRRP1, IMRRP1b or a functional portion thereof are provided as are methods for producing IMRRP1, IMRRP1b or a functional portion thereof. In addition, nucleic acid sequences encoding polypeptides, oligonucleotides, fragments, portions or antisense molecules thereof, and expression vectors and host cells comprising polynucleotides that encode IMRRP1 or IMRRP1b are provided. The use of the nucleic acid sequences, polypeptide, peptide and antibodies for diagnosis and treatment of disorders or diseases associated with aberrant regulation of blood pressure, induction of feeding, stimulation of firing of locus coeruleus neurons, and stimulation of insulin release, as well as the aberrant induction of the expression of glial fibrillary acidic protein independent of the action of alpha-2 adrenoceptors, dysphoric premenstrual syndrome, neurodegenerative disorders such as Alzheimer's disease, opiate addiction, monoamine turnover and therefore nociception, aging, mood and stroke, salivary disorders and developmental disorders is also described.

(21) **Appl. No.:** **09/932,145**

(22) **Filed:** **Aug. 17, 2001**

Related U.S. Application Data

(60) **Provisional application No. 60/226,411**, filed on Aug. 18, 2000. **Provisional application No. 60/261,779**, filed on Jan. 16, 2001.

Publication Classification

(51) **Int. Cl.⁷** **C07K 1/00**; C07K 14/00;
C07K 17/00

ATGTTCCGGCTCCGCCCCCAGCGTCCCGTGGCCATGACGACCGCTCAGAGGGACTCCCTG
TTGTGGAAGCTCGCGGGTTGCTGCGGGAGTCCGGGATGTGGTCTGTCTGGCTGTAGC
ACCCTGAGCCTGCTGACTCCCACACTGCAACAGCTGAACCACGTATTTGAGCTGCACCTG
GGCCATGGGGCCCTGGCCAGACAGGCTTTGTGGCTCTGCCCTCCCATCCTGCCGACTCC
CCTGTTATTCTTCAGCTTCAGTTTCTCTTCGATGTGCTGCAGAAAACACTTTCACTCAAG
CTGGTCCATGTTGCTGGTCTGGCCCCACAGGGCCCATCAAGATTTTCCCCTTCAAATCC
CTTCGGCACCTGGAGCTCCGAGGTGTCCCCTCCACTGTCTGCATGGCCTCCGAGGCATC
TACTCCCAGCTGGAGACCCTGATTTGCAGCAGGAGCCTCCAGGCATTAGAGGAGCTCCTC
TCAGCCTGCGGCGCGACTTCTGCTCTGCCCTCCCTTGGCTGGCTCTGCTTTCTGCCAAC
TTCAGCTACAATGCACTGACCGCCTTAGACAGCTCCCTGCGCCTCTTGTFCAGCTCTGCGT
TTCTTGAACCTAAGCCACAATCAAGTCCAGGACTGTCAGGGATTCCTGATGGATTTGTGT
GAGCTCCACCATCTGGACATCTCCTATAATCGCCTGCATTTGGTGCCAAGAATGGGACCC
TCAGGGCTGCTCTGGGGTCTGATACTGCGAGGCAATGAGCTTCGGAGCCTGCATGGC
CTAGAGCAGCTGAGGAATCTGCGGCACCTGGATTTGGCATAACAACCTGCTGGAAGGACAC
CGGGAGCTGTCACTGTGGCTGCTGGCTGAGCTCCGCAAGCTCTACCTGGAGGGGAAC
CCTCTTTGGTTCACCCCTGAGCACCGAGCAGCCACTGCCAGTACTTGTACCCCGGGCC
AGGGATGCTGCTACTGGCTTCTTCTCGATGGCAAGGTCTTGTCACTGACAGATTTTCAG
ACTCACACATCCTTGGGGCTCAGCCCCATGGGCCACCFTTGGCCCTGGCCAGTGGGGAGT
ACTCCTGAAACCTCAGGTGGCCCTGACCTGAGTGACAGCCTCTCCTCAGGGGTGTGTG
ACCCAGCCCCTGCTTCATAAGGTTAAGAGCCGAGTCCGTGTGAGGCGGGCAAGCATCTCT
GAACCCAGTGATACGGACCCGGAGCCCCGAACTCTGAACCCCTCTCCGGCTGGATGGTTC
GTGCAGCAGCACCCGGAGCTGGAGCTCATGAGCAGCTTCCGGGAACGGTTCGGCCGCAAC
TGGCTGCAGTACAGGAGTCACCTGGAGCCCTCCGAAACCCTCTGCCGGCCACCCCACT

Figure 1A

ACTTCTGCACCCAGTGCACCTCCAGCCAGCTCCCAGGGCCCCGACACTGCACCCAGACCT
TCACCCCCGCAGGAGGAAGCCAGAGGCCCCAGGACTCACCACAGAAAATGTCAGAGGAG
GTCAGGGCGGAGCCACAGGAGGAGGAAGAGGAGAAGGAGGGGAAGGAGGAGAAGGAGGAG
GGGAGATGGTGGAAACAGGGAGAAGAGGAGGCAGGAGAGGAGGAAGAAGAGGAGCAGGAC
CAGAAGGAAGTGGAAAGCGGAACTCTGTGCCCCCTTGTGGTGTGTCCCCTGGAGGGCCT
GAGGGCATAACGGGGCAGGGAATGCTTTCTCAGGGTCACTTCTGCCACCTGTTTGAGGTG
GAACTCCAAGCAGCTCGCACCTTGGAGCGACTGGAGCTCCAGAGTCTGGAGGCAGCTGAG
ATAGAGCCGGAGGCCCAGGCCCAGAGGTCGCCCAGGCCACGGGCTCAGATCTGCTCCCT
GGAGCCCCATCCTCAGTCTGCGCTTCTCCTACATCTGCCCTGACCGGCAGTTGCGTCGC
TATTTGGTGTCTGGAGCCTGATGCCACGCAGCTGTCCAGGAGCTGCTTGCCGTGTTGACC
CCAGTCACCAATGTGGCTCGGGAACAGCTTGGGGAGGCCAGGGACCTCCTGCTGGGTAGA
TTCCAGTGTCTACGCTGTGGCCATGAGTTCAAGCCAGAGGAGCCCAGGATGGGATTAGAC
AGTGAGGAAGGCTGGAGGCCTCTGTTCCAAAAGACAGGGAGCGGAAACAGGGAGAGCAGT
CTCTGGCTCCTTCTCCGTTTGCCAGCCCTGTCTGCCACCCTCCTGGCCATGGTGACCACC
TTGACAGGGCCAAGAACAGCCCACCTCAGGCACCGAGCACCCGTGACCATGGTAGTTGGA
GCCTCAGTCCCCCCCCCTGAGCGCTGTGGCCTCCGCTCTGTGGACCACCGACTCCGGCTCT
TCCTGGATGTTGAGGTGFTCAGCGATGCCCAGGAGGAGTTCCAGTGCTGCCTCAAGGTGC
CAGTGGCATTGGCAGGCCACACTGGGGAGTTCATGTGCCTTGTGGTTGTGTCTGACCGCA
GGCTGTACCTGTTGA

Figure 1B.

ATGTTGGCTCCGCCCCCAGCGTCCCGTGGCCATGACGACCGCTCAGAGGGACTCCCTG
TTGTGGAAGCTCGCGGGGTTGCTGCGGGAGTCCGGGGATGTGGTCCTGTCTGGCTGTAGC
ACCCTGAGCCTGCTGACTCCCACACTGCAACAGCTGAACCACGTATTTGAGCTGCACCTG
GGCCATGGGGCCCTGGCCAGACAGGCTTTGTGGCTCTGCCCTCCCATCCTGCCGACTCC
CCTGTTATTCTTCAGCTTCAGTTTCTCTTCGATGTGCTGCAGAAAACACTTTCACTCAAG
CTGGTCCATGTTGCTGGTCCTGGCCCCACAGGGCCCATCAAGATTTCCCTTCAAATCC
CTTCGGCACCTGGAGCTCCGAGGTGTTCCCTCCACTGTCTGCATGGCCTCCGAGGCATC
TACTCCCAGCTGGAGACCCTGATTTGCAGCAGGAGCCTCCAGGCATTAGAGGAGCTCCTC
TCAGCCTGCGGCGGCGACTTCTGCTCTGCCCTCCCTTGGCTGGCTCTGCTTTCTGCCAAC
TTCAGCTACAATGCACTGACCGCCTTAGACAGCTCCCTGCGCCTCTTGTGAGCTCTGCGT
TTCTTGAACCTAAGCCACAATCAAGTCCAGGACTGTCAGGGATTCCCTGATGGATTTGTGT
GAGCTCCACCATCTGGACATCTCCTATAATCGCCTGCATTTGGTGCCAAGAATGGGACCC
TCAGGGGCTGCTCTGGGGTCTGATACTGCGAGGCAATGAGCTTCGGAGCCTGCATGGC
CTAGAGCAGCTGAGGAATCTGCGGCACCTGGATTTGGCATAACAACCTGCTGGAAGGACAC
CGGGAGCTGTCACCACTGTGGCTGCTGGCTGAGCTCCGCAAGCTCTACCTGGAGGGGAAC
CCTCTTTGGTTCCACCCTGAGCACCGAGCAGCCACTGCCAGTACTTGTCACCCCGGGCC
AGGGATGCTGCTACTGGCTTCTTCTCGATGGCAAGGTCTTGTCACTGACAGATTTTCAG
ACTCACACATCCTTGGGGCTCAGCCCCATGGGCCACCTTTGCCCTGGCCAGTGGGGAGT
ACTCCTGAAACCTCAGGTGGCCCTGACCTGAGTGACAGCCTCTCCTCAGGGGGTGTGTG
ACCCAGCCCCTGCTTCATAAGGTTAAGAGCCGAGTCCGTGTGAGGCGGGCAAGCATCTCT
GAACCCAGTGATACGGACCCGGAGCCCCGAACCTCTGAACCCCTCTCCGGCTGGATGGTTC
GTGCAGCAGCACCCGGAGCTGGAGCTCATGAGCAGCTTCCGGGAACGGTTCGGCCGCAAC

Figure 2A

TGGCTGCAGTACAGGAGTCACCTGGAGCCCTCCGGAAACCCTCTGCCGGCCACCCCCACT
ACTTCTGCACCCAGTGCACCTCCAGCCAGCTCCCAGGGCCCCGACACTGCACCCAGACCT
TCACCCCCGCAGGAGGAAGCCAGAGGCCCCCAGGAGTCACCACAGAAAATGTCAGAGGAG
GTCAGGGCGGAGCCACAGGAGGAGGAAGAGGAGAAGGAGGGGAAGGAGGAGAAGGAGGAG
GGGAGATGGTGGAAACAGGGAGAAGAGGAGGCAGGAGAGGAGGAAGAAGAGGAGCAGGAC
CAGAAGGAAGTGAAGCGGAACTCTGTGCCCCCTTGTGGTGTGTCCCCTGGAGGGGCCT
GAGGGCGTACGGGGCAGGGAATGCTTTCTCAGGGTCACTTCTGCCACCTGTTTGAGGTG
GAACTCCAAGCAGCTCGCACCTTGGAGCGACTGGAGCTCCAGAGTCTGGAGGCAGCTGAG
ATAGAGCCGGAGGCCAGGCCAGAGGTCGCCCAGGCCACGGGCTCAGATCTGCTCCCT
GGAGCCCCATCCTCAGTCTGCGCTTCTCCTACATCTGCCCTGACCGGCAGTTGCGTCGC
TATTTGGTGTCTGGAGCCTGATGCCACGCAGCTGTCCAGGAGCTGCTTGCCGTGTTGACC
CCAGTCACCAATGTGGCTCGGGAACAGCTTGGGGAGGCCAGGGACCTCCTGCTGGGTAGA
TTCCAGTGTCTACGCTGTGGCCATGAGTTCAAGCCAGAGGAGCCCAGGATGGGATTAGAC
AGTGAGGAAGGCTGGAGGCCTCTGTTCCAAAAGACAGAATCTCCTGCTGTGTGTCCTAAC
TGTGGTAGTGACCACGTGGTTCTCCTCGCTGTGTCTCGGGGAACCCCCAACAGGGAGCGG
AAACAGGGAGAGCAGTCTCTGGCTCCTTCTCCGTTTGCCAGCCCTGTCTGCCACCCTCCT
GGCCATGGTGACCACCTTGACAGGGCCAAGAACAGCCCACCTCAGGCACCGAGCACCCGT
GACCATGGTAGTTGGAGCCTCAGTCCCCCCCCTGAGCGCTGTGGCCTCCGCTCTGTGGAC
CACCGACTCCGGCTCTTCTGGATGTTGAGGTGTTTCAGCGATGCCAGGAGGAGTTCCAG
TGCTGCCTCAAGGTGCCAGTGGCATTGGCAGGCCACACTGGGGAGTTCATGTGCCTTGTG
GTTGTGTCTGACCGCAGGCTGTACCTGTTGAAGGTGACTGGGGAGATGCGTGAGCCTCCA
GCTAGCTGGCTGCAGCTGACCCTGGCTGTTCCCCTGCAGGATCTGAGTGGCATAGAGCTG

Figure 2B

GGCCTGGCAGGCCAGAGCCTGCCGGCTAGAGTGGGCAGCTGGGGCGGGCCGCTGTGTGCTG
CTGCCCCGAGATGCCAGGCATTGCCGGGCCTTCCTAGAGGAGCTCCTTGATGTCTTGCAG
TCTCTGCCCCCTGCCTGGAGGAAGTGTGTGAGTCCACAGAGGAGGAGGTCACCCCCCAG
CACCGGCTCTGGCCATTGCTGGAAAAAGACTCATCCTTGGAGGCTCGCCAGTTCTTCTAC
CTTCGGGCGTTTCCTGGTTGAAGGCCCTTCCACCTGCCTCGTATCCCTGTTGCTGACTCCG
TCCACCCTGTTTCCTGTTAGATGAGGATGCTGCAGGGTCCCCGGCAGAGCCCTCTCCTCCA
GCAGCATCTGGCGAAGCCTCTGAGAAGGTGCCTCCCTCGGGGCGGGCCCTGCTGTGCGT
GTCAGGGAGCAGCAGCCACTCAGCAGCCTGAGCTCCGTGCTGCTCTACCGCTCAGCCCCCT
GAGGACTTGCGGCTGCTCTTCTACGATGAGGTGTCCCGGCTGGAGAGCTTTTGGGCACTC
CGTGTGGTGTGTCAGGAGCAGCTGACAGCCCTGCTTGCCTGGATCCGGAACCATGGGAG
GAGCTGTTTTCCATCGGACTCCGGACAGTGATCCAAGAGGCGCTGGCCCTTGACCGATGA

Figure 2C.

MFGSAPQRPVAMTTAQRDSSLWKLGLLRESGDVVLSGCSTLSLLTPTLQQLNHVFEHLH
GPWGPGQTGFVALPSHPADSPVILQLQFLFDVLQKTLCLKLVHVAGPGPTGPIKIFPKS
LRHLELRGVPLHCLHGLRGIYSQLETLICSRSLQALEELLSACGGDFCSALPWLALLSAN
FSYNALTALDSSLRLLSALRFLNLSHNQVQDCQGFLMDLCELHHLDISYNRLHLVPRMGP
SGAALGVLIILRGNELRSLHGLEQLRNLRHLDLAYNLEGHRELSPLWLLAELRKLYLEGN
PLWFHPEHRAATAQYLSPRARDAATGFLLDGKVLSTDFQHTSLGLSPMGPPLPWPVGS
TPETSGGPDLSDSLSSGGVVTQPLLHKVKSRVRVRRASISEPSDTPPEPRTLNPSPAGWF
VQQHPELELMSSFRERFGRNWLYRSHLEPSGNPLPATPTTSAPSAPPASSQGPDTAPRP
SPPQEEARGPQESPQKMSEEVRAEPQEEEEKEGKKEEKEGEMVEQGEEOEAGEEEEEEOQ
QKEVEAELCRPLLVCPLGPEGIRGRECFLRVTS AHLFEVELQAARTLERLELQSLEAAE
IEPEAQAQRSPRPTGSDLLPGAPILSLRFSYICPDRQLRRYLVLLEPDAHA AVQELLAVLT
PVTNVAREQLGEARDLLLGRFQCLRCGHEFKPEEPRMGLDSEEGWRPLFQKTGSGNRESS
LWLLLRLPALSATLLAMVTTLTGPRTAHLRHRAPVTMVVGASVPPLSAVASALWTTDSGS
SWMLRCSAMPRRSSSAASRCQWHWQATLGSSCALWLCLTAGCTC

Figure 3.

MFGSAPQRPVAMTTAQRDSSLWKLAGLLRESGDVVLSGCSTLSLLTPTLQQLNHVFELHL
GPWGPQQTGFVALPSPHAPDSPVILQLQFLFDVLQKTLCLKLVHVAGPGPTGPIKIFPFKS
LRHLELRGVPLHCLHGLRGIYSQLETLICSRSLQALEELLSACGGDFCSALPWLALLSAN
FSYNALTALDSSLRLLSALRFLNLSHNQVQDCQGFMDLCELHHLDISYNRLHLVPRMGP
SGAALGVLIIRGNELRSLHGLEQLRNLRHLDLAYNLLLEGHRELSPLWLLAELRKLYLEGN
PLWFHPEHRAATAQYLSPRARDAATGFLLDGKVLSLTDFQTHTSLGLSPMGPPLPWPVGS
TPETSGGPDLSDSLSSGGVVTQPLLHKVKSRRVRRASISEPSDTPPEPRTLNPSPAGWF
VQQHPELELMSSFRERFGRNWLYRSHLEPSGNPLPATPTTSAPSAPPASSQGPDTAPRP
SPPQEEARGPQESPQKMSEEVRAEPQEEEEKEGKKEEKEEGEMVEQEEEEAGEEEEEEQD
QKEVEAELCRPLLVCPLGPEGVVRGREGFLRVTSALHFEVELQAARTLERLELQSLEAAE
IEPEAQAQRSPRPTGSDLLPGAPILSLRFSYICPDRQLRRYLVLEPDAHAHVQELLAVLT
PVTNVAREQLGEARDLLLGRFQCLRCGHEFKPEEPRMGLDSEEGWRPLFQKTESPAVCPN
CGSDHVLLAVSRGTPNRERKQGEQSLAPSPFASPVCHPPGHGDHLDRAKNSPPQAPSTR
DHGSWSLSPPERCGLRSVDHRLRLFLDVEVFSDAQEEFQCCLKVPVALAGHTGEFMCLV
VVSDRRLYLKVTGEMREPPASWLQLTLAVPLQDLSGIELGLAGQSLRLEWAAGAGRCVL
LPRDARHCRAFLEELLDVLQSLPPAWRNCVSATEEEVTPQHRLWPLLEKDSSEARQFFY
LRAFLVEGPSTCLVSLLLTPTLFLDDEDAAGSPAEPSPPAASGEASEKVPPSGPGPAVR
VREQQPLSSLSSVLLYRSAPEDLRLLFYDEVSRLESFWALRVVCQEQLTALLAWIREPWE
ELFSIGLRTVIQEALALDR

Figure 4.

>gi|6005788 imidazoline receptor candidate >gi|3462807|gb|AAC33104.1|
(AF082516) I-1 receptor candidate protein [Homo sapiens]
Length = 1504

Score = 68.3 bits (164), Expect = 4e-10
Identities = 69/256 (26%), Positives = 102/256 (38%), Gaps = 26/256 (10%)

Query: 107 VLQKTLCLKLVHVAGP-GPTG-----PIKIFPFKSLRHLRLRGVPLHCLHGLRGIY 156
+L T LK + V+G GP G P + FKSL +E+ + GL

Sbjct: 180 ILDFTCRLKYLKVSQTEGPPFGTSTNIQEQLLPFDLISIFKSLHQVEISHCDAKHIRGLVASK 239

Query: 157 SQLETLICSRSLQALEELL-----SACGGDFCSALP-WLALLSANFSYNXX 201
L TL S +++E+L + G + +P W AL + + S+N

Sbjct: 240 PTLATLSVRFSAATSMKEVLVPEASEFDEWEPEGTTLEGPVTAIVIPTWQALTTLDLSHNSI 299

Query: 202 XXXXXXXXXXXXXXXXXXXXXXXHNQVQDCQGFMDLCELHHLDISYNRLHLVPRMGPSGAALG 261
HN + L L L HLD+SYN+L + + +

Sbjct: 300 SEIDESVKLIPKIEFLDLSHNGLLVVDN-LQHLYNLVHLDLSYNKLSLEGLHTKLGNIK 358

Query: 262 VLILRGNELRSLHGLEQLRNLRHLDLAYNLLEGHRELSPLWLLAELRKLKLYLEGNPLWFHP 321
L L GN L SL GL +L +L +LDL N +E E+ + L L + L NPL P

Sbjct: 359 TLNLAGNLESLSGLHKLYSLVNLDLRDNRIEQMEEVRSIGSLPCLEHVSLNNPLSIIP 418

Query: 322 EHRAATAQYLSPRARD 337
++R RA +

Sbjct: 419 DYRTKVLQAQFGERASE 434

Figure 5.

```

101 TACGCTGTGGCCATGAGTTCAAGCCAGAGGAGCCCAGGATGGGATTAGAC 2150
    |||
1644 TACGCTGTGGCCATGAGTTCAAGCCAGAGGAGCCCAGGATGGGATTAGAC 1693
    .
2151 AGTGAGGAAGGCTGGAGGCCTCTGTTCCAAAAG..... 2183
    |||
1694 AGTGAGGAAGGCTGGAGGCCTCTGTTCCAAAAGACAGAATCTCCTGCTGT 1743
    .
    .
2184 .....ACAGGGAGCGGAAACAGGGAGAGCAGTCTCTGGCTCCTTCT 2224
    |||
1794 GAACCCCAACAGGGAGCGGAAACAGGGAGAGCAGTCTCTGGCTCCTTCT 1843
    .
2225 CCGTTTGCCAGCCCTGTCTGCCACCCTCCTGGCCATGGTGACCACCTTGA 2274
    |||
1844 CCGTTTGCCAGCCCTGTCTGCCACCCTCCTGGCCATGGTGACCACCTTGA 1893
    .
    .
2825 TCATCCTTGGAGGCTCGCCAGTTCTTCTACCTTCGGGCGTTCCTGGTTGA 2874
    |||
2444 TCATCCTTGGAGGCTCGCCAGTTCTTCTACCTTCGGGCGTTCCTGGTTGA 2493
    .
    .
3375 ATCTGGGCCCTCCATGACCTTCCACACTGGATGCCTCTTTCCCTGCAGG 3424
    |||
2494 .....AGG 2496
    .
3425 CCCTTCCACCTGCCTCGTATCCCTGTTGCTGACTCCGTCCACCCTGTTCC 3474
    |||
2497 CCCTTCCACCTGCCTCGTATCCCTGTTGCTGACTCCGTCCACCCTGTTCC 2546

```

Figure 6.

Query= sequence
(1114 letters)

Database: newnr
228,478 sequences; 162,186,938 total letters

Searching.....done

Sequences producing significant alignments:	Score	E
	(bits)	Value

gb AAF52305.1 (AE003611) CG9044 gene product [Drosophila melano...	127	5e-28
gi 6005788 imidazoline receptor candidate >gi 3462807 gb AAC3310...	68	4e-10
gb AAF57514.1 (AE003794) CG8595 gene product [Drosophila melano...	47	0.001

>gb|AAF52305.1| (AE003611) CG9044 gene product [Drosophila melanogaster]
Length = 1289

Score = 127 bits (317), Expect = 5e-28
Identities = 99/321 (30%), Positives = 149/321 (45%), Gaps = 11/321 (3%)

Query: 38 KLAGLLRESGDVVXXXXXXXXXXXXXXXXXNHVF-----ELHLGPWGPQTGFVALPSH 91
+LA LLR++GD + N F E+ G F +
Sbjct: 8 ELANLLRQNGDKILSSEFTLTLSGSLLRALNDSFTLIADTEIGTGAGYLQPQSFQVVKPI 67

Query: 92 PADSPVILQLQFLFDVLQKTLCLKLVHVAGPGP-TGPIKIFPFKSLRHLELRGVPLHCLH 150
A S V LQ + D +QKT LKL + G I I F++LR LE+ + + +
Sbjct: 68 NAKSSVFPDLQLVHDFVQKTTLLKLTYPSEHYFEGAIDIAKFRALRRLEVNKINIGQVV 127

Query: 151 GLRGIYSQLETLICSRSLQALELLSACGGDFCSALPWLALLSANFSYNXXXXXXXXXXXXX 210
G++ + QL+ LIC +SL ++++++ CGGD + W L +A+FSYN
Sbjct: 128 GIQPLRGQLQHLCVKSLTSVDDIITRCGGDNSNGFVWNLKLTADFSYNSLRSDTALEF 187

Query: 211 XXXXXXXXXXXXHNQVQDCQGFMDLCELHHLDISYNRLHLVPRMGPSGA-ALGVLILRGNE 269
HN++ + L L LD+SYN L +P+ L +L + N
Sbjct: 188 AQHLQHLNLRHNKLTVA-A-IKWLPHLKTLDLSYNCLTHLPQFHMEACKRLQLLNISNNY 246

Query: 270 LRSLHGLEQLRNLRLHLDLAYNLLEGHRELSPLWLLAELRKLYLEGNPLWFHPEHRAATAQ 329
+ L + +L L +LDL+ N L H +L PL L L L L+GNPL +P+HR ATAQ
Sbjct: 247 VEELLDVAKLDALYNLDLSDNCLLEHSQLLPLSALMSLIVLNQGNPLACNPKHRQATAQ 306

Query: 330 YLSPRARDAATGFLLDGKVL 350
YL A F+LD + L+
Sbjct: 307 YL--HKNSATVKFVLD FEPLT 325

Figure 7A

Score = 41.4 bits (95), Expect = 0.054
 Identities = 41/151 (27%), Positives = 62/151 (40%), Gaps = 20/151 (13%)

Query: 814 VDHRLRLFLDVEVFSDAQEEFQCCLKVPVALAGHTGEFMCLVVVSDRRLYLLKVTGEMRE 873
 +DHRL+L+ F + E F+ K + LVV+S+ + YL++ E +

Sbjct: 1018 IDHRLKLYFYQRKFKEDGEHFKWLAKGRIYNEQTQSLGEGLVVMSNCKCYLMEAFAPHD 1077

Query: 874 PPASWLQLTLAVPLQDLSGIELGLAGQSLRLEWAAGA-----GRCVLLPRDARHCRAF 926
 A WL+ ++V + L I+L L W G G VLL D

Sbjct: 1078 DVAKWLRQVSVAVNRLVAIDL-----LPWKLGLSF'TLKDWGGFVLLLDMLR---- 1125

Query: 927 LEELLDVLQSLPPAWRNCVSATEEEVTPQHR 957

E LL+ LQ +P C + VT H+

Sbjct: 1126 TESLLNYLQQIPLP-EQCKLNHQPSVTLSHQ 1155

>gi|6005788 imidazoline receptor candidate >gi|3462807|gb|AAC33104.1|
 (AF082516) I-1 receptor candidate protein [Homo sapiens]
 Length = 1504

Score = 68.3 bits (164), Expect = 4e-10
 Identities = 69/256 (26%), Positives = 102/256 (38%), Gaps = 26/256 (10%)

Query: 107 VLQKTLSLKLHVHAGP-GPTG-----PIKIFPFKSLRHLLELRGVPLHCLHGLRGIY 156
 +L T LK + V+G GP G P + FKSL +E+ + GL

Sbjct: 180 ILDFTCRLLKYLKVSGETEPPFGTNSIQEQLLPFDLSIFKSLHQVEISHCDAKHIRGLVASK 239

Query: 157 SQLETLICRSRLQALEELL-----SACGGDFCSALP-WLALLSANFSYNXX 201
 L TL S +++E+L + G + +P W AL + + S+N

Sbjct: 240 PTLATLSVRFSAATSMKEVLVPEASEFDEWEPEGTTLLEGPVTAVIPTWQALTTLDLSHNSI 299

Query: 202 XXXXXXXXXXXXXXXXXXXXHNQVQDCQGFMLDLCELHHLDISYNRLHLVPRMGPSGAALG 261
 HN + L L L HLD+SYN+L + + +

Sbjct: 300 SEIDESVKLIPKIEFLDLSHNGLLVVDN-LQHLYNLVHLDLSYNKLSLEGLHTKLGNIK 358

Query: 262 VLILRGNELRSLHGLEQLRNLRHLDLAYNLLLEGHRELSPLWLLAELRKLYLEGNPLWFHP 321
 L L GN L SL GL +L +L +LDL N +E E+ + L L + L NPL P

Sbjct: 359 TNLNLAGNLESLSGLHKLYSLVNLDLRDNRIEQMEEVRSIGSLPCLEHVSLNPLSIIP 418

Query: 322 EHRAATAQYLSRARD 337

++R RA +

Sbjct: 419 DYRTKVLAQFGERASE 434

Figure 7B

```

FL1-18_SPLICE_VARIANT      ---MFGSAPQRPVAMTTAQRDS---LLWKLAGLLRESGD-----
FL1-18                      ---MFGSAPQRPVAMTTAQRDS---LLWKLAGLLRESGD-----
Drosophila_melanogaster_CG9044  -----MDPQK-----ITELANLLRQNGD-----
imidazoline_receptor_candidate MATARTFGPEAREAPAKEARVVGSELVDTTYVYIIQVTDGSHEWTVKHRY
                               *::      :      :      :      :      *

FL1-18_SPLICE_VARIANT      -----VVLSGCSTLSLLTP-----
FL1-18                      -----VVLSGCSTLSLLTP-----
Drosophila_melanogaster_CG9044  -----KILSSBFTLTLSGS-----
imidazoline_receptor_candidate SDFHDLHEKLVLAERKIDKNLLPCKIIGKNSRSLVEKREKDEVLVYQKLL
                               :      .      *      .

FL1-18_SPLICE_VARIANT      -----TLQQLNHVFEHLHG-----PWG
FL1-18                      -----TLQQLNHVFEHLHG-----PWG
Drosophila_melanogaster_CG9044  -----LLRALNDSFTLIADT-----EIG
imidazoline_receptor_candidate AAFPQVTPRVLAHFLHFHFYEINGITAALAEELFEKGEQLLGAGEVFAIG
                               :      *      .      :      :      *

FL1-18_SPLICE_VARIANT      PG---Q-TGFVALPSHPADSPVILQQLFLFDVLQKTLSSLKLVHVAGPGP
FL1-18                      PG---Q-TGFVALPSHPADSPVILQQLFLFDVLQKTLSSLKLVHVAGPGP
Drosophila_melanogaster_CG9044  TGAGYVLPQSPQVVKPINAKSSVFPDLQVLVHDFVQKTTLLKLTYPFSEHY
imidazoline_receptor_candidate PLQLYAVTEQLQQGKPTCASGDAKTDLGHILDFTCRKLYLKVSGTEGPF
                               .      :      .      *      .      :      *      :      **      .

FL1-18_SPLICE_VARIANT      -----TGPIKIFPFKSLRHLELRGVPLHCLHGLRGIYSQLETLICRSR
FL1-18                      -----TGPIKIFPFKSLRHLELRGVPLHCLHGLRGIYSQLETLICRSR
Drosophila_melanogaster_CG9044  F-----EGAIDIAKFRALRRLEVNKINIGQVVGIQPLRGQLQHLCVKS
imidazoline_receptor_candidate TSNIQBQLLPDLSIFKSLHQVEISHCDAKHIRGLVASKPTLATLSVRFS
                               .      :      *      :      :      *      :      *      *      *

FL1-18_SPLICE_VARIANT      LQALELLSACGGDFCSALP-----WLALLSANFSYNALT
FL1-18                      LQALELLSACGGDFCSALP-----WLALLSANFSYNALT
Drosophila_melanogaster_CG9044  LTSVDDIITRCGGDNSNGFV-----WNELKTADFSYNSLR
imidazoline_receptor_candidate ATSMKEVLPPEASEFEWEPEGTTLEGPVTAIPTWQALTTLDLSHNSIS
                               :      :      :      .      .      .      *      *      :      :      *      :

FL1-18_SPLICE_VARIANT      ALDSSLRLLSALRFLNLSHNQVQDCQGFMDLCELHHLDISYNRLHLVPR
FL1-18                      ALDSSLRLLSALRFLNLSHNQVQDCQGFMDLCELHHLDISYNRLHLVPR
Drosophila_melanogaster_CG9044  SVDTALEFAQHLQHLNLRHNKLTVA-AIKWLPHLKTLTDLSDNCLTHLPQ
imidazoline_receptor_candidate EIDESVKLIPKIEFLDLSHNGLLVVD-NLQHLNVLVHLDLSYNKLSLEG
                               :*      :      :      :      .      :      *      *      **      :      *      :

FL1-18_SPLICE_VARIANT      -MGPSGAALGVLIIRGNELRSLHGLEQLRNLRHLDLAYNLEGHRELSPL
FL1-18                      -MGPSGAALGVLIIRGNELRSLHGLEQLRNLRHLDLAYNLEGHRELSPL
Drosophila_melanogaster_CG9044  FHMEACKRQLQNLNINSNYVEELLDVAKLDALYNLSDNCLLEHSQQLLPL
imidazoline_receptor_candidate -LHTKLGNIKTLNLAGNLLSLSGLHKLYSLVNLDLRDNRIEQMEEVRSI
                               :      *      :      .      *      .      :      *      *      ***      *      :      :      :

FL1-18_SPLICE_VARIANT      WLLAELRKLYLEGNPLWFHPEHRAATAQYLSPRARDAATGFLLDGKVLSL
FL1-18                      WLLAELRKLYLEGNPLWFHPEHRAATAQYLSPRARDAATGFLLDGKVLSL
Drosophila_melanogaster_CG9044  SALMSLIVLNLQGNPLACNPKHRQATAQYLHKNS--ATVKFVLDFEPLTK
imidazoline_receptor_candidate GSLPCLEHVSVLLNPLSIPDYRTKVLAQFGERA---SEVCLDDTVTTE
                               *      *      :      *      .      ***      *      :      .      :      :      .      **      :
    
```

Figure 8A

FL1-18_SPLICE_VARIANT
FL1-18
Drosophila_melanogaster_CG9044
imidazoline_receptor_candidate

TDFQTHTS-----LGLSPMGPP-LPWVPGSTPETSGGPDLSDLSLSSG--
TDFQTHTS-----LGLSPMGPP-LPWVPGSTPETSGGPDLSDLSLSSG--
AEKALTGSQKRWYISGLSHRSPRSTSMGINSSGASINTSDGSGFSSFGSQ
KELDTVEV-----LKAIQKAKE--VKSKLSNPEKKGED-SRLSAAPCI
: . . . * . . . * * :

FL1-18_SPLICE_VARIANT
FL1-18
Drosophila_melanogaster_CG9044
imidazoline_receptor_candidate

-----G-----VVTQPLLHKVKS-RVR--VRRASISEPS--DTD----
-----G-----VVTQPLLHKVKS-RVR--VRRASISEPS--DTD----
RSVSRGKKNYTLLEDNQSMDSQSSKRISCKIRTVDIESSEINTDAASV
R----PSSSPPTVAPASASLPQPILSNQGIMFVQEEALASSLSSTDS---
. . . . * **

FL1-18_SPLICE_VARIANT
FL1-18
Drosophila_melanogaster_CG9044
imidazoline_receptor_candidate

--PEPRT-----LNPSPAG-----WFVQOHPPEL
--PEPRT-----LNPSPAG-----WFVQOHPPEL
STPNRSE-----YEEEPDNHLETKKKIETLRLTYGNEWLKSNGNAEL
LTPBEHQPIAQGCSDSLESIPAG----QAASDDLDRDVPGAVGGASPEHAEF
* : . : * . : *

FL1-18_SPLICE_VARIANT
FL1-18
Drosophila_melanogaster_CG9044
imidazoline_receptor_candidate

EL-----MSSFREERFG-----RNWLQYRSHLE---PSGNP-----L
EL-----MSSFREERFG-----RNWLQYRSHLE---PSGNP-----L
MLGIE--TPQPTERERNES----RQLFNEYLGELSGFTEAKNDSEHNI
EVQVVPGSGQIIFLFTTCIGYTATNQDFIQLRSLTLIR--QAIERQLPAWI
: : : : * : : :

FL1-18_SPLICE_VARIANT
FL1-18
Drosophila_melanogaster_CG9044
imidazoline_receptor_candidate

PATPT-----TSAPSAPPASSQGPDTAPRP-----SPQE---EARG-
PATPT-----TSAPSAPPASSQGPDTAPRP-----SPQE---EARG-
SSTPTNNVLLASTFDATITPIKSEANDTSGQTLYETCTEGETNYESFGN
EAANQRE-EGQGEQGEEDDEEEEEEDVAENRYFEMGPPDVE---BEEGG
: : . . . * * * *

FL1-18_SPLICE_VARIANT
FL1-18
Drosophila_melanogaster_CG9044
imidazoline_receptor_candidate

-----PQESFQ-KMSEEVRAEP----QEED-----E
-----PQESFQ-KMSEEVRAEP----QEED-----E
NTTELSTEERPPDRHEELLRLYASSNAQDED-----P
GQGE-EEEEEEDEEAEEERLAEWALGADEDFLLEHIRILKVLWCFLIH
: * . * * : * :

FL1-18_SPLICE_VARIANT
FL1-18
Drosophila_melanogaster_CG9044
imidazoline_receptor_candidate

KEGKBEKEEGEMV--EQGEEAGEEEEEEQ-DQKEVEAEL----CRP---
KEGKBEKEEGEMV--EQGEEAGEEEEEEQ-DQKEVEAEL----CRP---
VSDAESDEETYIVYHEQKPSVFLFTISSN-FIREKDTLT----ERT---
VQGSTRQFAACLVLTDGFIADVFEIPHQESRGSSQHILSSLRVFCFPHGD
. . . : * : . . . : :

FL1-18_SPLICE_VARIANT
FL1-18
Drosophila_melanogaster_CG9044
imidazoline_receptor_candidate

LLVCPLEGPEG----VRGECFLRV TSA---HLFEVEL-----QAAR
LLVCPLEGPEG----IRGECFLRV TSA---HLFEVEL-----QAAR
KAKWSLKILESC---ERVRSNTLRINFD---TMRKDKQ-----ERIY
LTEFGFLMPELCLVLKVRHSENTLFIISDAANLHEFHADLRSCFAPQHMA
: * * * * : . . :

FL1-18_SPLICE_VARIANT
FL1-18
Drosophila_melanogaster_CG9044
imidazoline_receptor_candidate

TLERLELQSLEAAEIBPEAQORSRPTGS-----DLLPGAPIL-----
TLERLELQSLEAAEIBPEAQORSRPTGS-----DLLPGAPIL-----
CVENTLQCELEKLRDILSQRDLTEMNISIYRCV--NCLTQFTIEQK---
MLCSPILYGSHTSLQBFLRQLLTFYKVAGGCQERSQGCFFVYLVYSDKRM
: . . : * . . . : :

Figure 8B

```

FL1-18_SPLICE_VARIANT
FL1-18
Drosophila_melanogaster_CG9044
imidazoline_receptor_candidate
-----SLRFS-----YICPDRQ-----LRR-----YLVLEP-DA
-----SLRFS-----YICPDRQ-----LRR-----YLVLEP-DA
-----SKRYKA--KELRCPCRSVYVAEVELSSSLSKPS-GEVAAEPKLS
VQTAAGDYSGNIEWASCTLCSAVRRSCCAPSEAVKSAAPYWLLLTPOHL
: :. *. : *

FL1-18_SPLICE_VARIANT
FL1-18
Drosophila_melanogaster_CG9044
imidazoline_receptor_candidate
HAAVQELLAVLTPVTNVAREQ-----LGEARDLLLGRFQC-----L
HAAVQELLAVLTPVTNVAREQ-----LGEARDLLLGRFQC-----L
PAMIVEESPVEELAAANKEES----NSIGKSLASFLFYFDESSFDSNQS
NVIKADFNPMNPRGTHNCRNRNSFKLSRVPLSTVLLDPTRSCTQ-PRGAF
. :. : :. : : :

FL1-18_SPLICE_VARIANT
FL1-18
Drosophila_melanogaster_CG9044
imidazoline_receptor_candidate
RCG-----HEFKPBEPRMGLDSE-----EGWRP-LFQKTESPAVCP
RCG-----HEFKPBEPRMGLDSE-----EGWRP-LFQKTGS-----
VVGSSNTDRD-MEFRANESDVIDIISNPSQSSI EVLDPNYPVQSASRKTSEE
ADGHVLELLVGYRFVTAIFVLPHEKPFHFLRVYNQLRA-SLQDLKTVVIAK
* . * . : . : . * .

FL1-18_SPLICE_VARIANT
FL1-18
Drosophila_melanogaster_CG9044
imidazoline_receptor_candidate
NCGSDHVLLAVSRGTPNRRERKQGEQSLAPSPFASPVCHPPGHGDLHDLRA
--GNRESSLWLLLR-LP-----ALSATLLAMVTTLTGPRTAHL-RH
RRISQLPHLETIHDEVAK-SKSFIEREFQQLLAEQAQPTTPTAAPLAPA
TPGTGGSPQGSFADGQPAERRASNDQRPOEVPAAEALAPAVEVPAPAPAA
. . . . :.

FL1-18_SPLICE_VARIANT
FL1-18
Drosophila_melanogaster_CG9044
imidazoline_receptor_candidate
KNSP-P-QAPSTRDHGSW-----SLSPPPERCGLRSVDHRLRL
R-----APVTMVVG-----ASVPP-----
KSAV-PSHVPLTBSSSSGSVTDICTTYEQQATDAPQNLQNSLLTESSNS
ASASGPAKTAPABASTSALVP--EETPVEAPAPPPAEPAPAQYFSEHLIQ
. . * . : * .

FL1-18_SPLICE_VARIANT
FL1-18
Drosophila_melanogaster_CG9044
imidazoline_receptor_candidate
FLDVEV----FSDAQEEFQCCLKVVALAGHTGEFMCVVVSDRRLYLL
-----LSAVAS-----ALWTTDS-----
QVSGSD----AESNSRLKSAEDASLLPFASVPQSTNLLMSSSKKLIIESE
ATSEENQIPSHLPACPSLRHVASLRGSATIELFHSSTIAEVENEELRHLMW
. . :

FL1-18_SPLICE_VARIANT
FL1-18
Drosophila_melanogaster_CG9044
imidazoline_receptor_candidate
KVTGEMREPPASWLQTLTAVPLQDLSGIELGL--AGQSLRLEWAAGAG-
-----GSSWMLRCSAMPRRSS-----AASRCQWHWQATLGS
ATVFGTQPYKFNYSDFNDIDHRLKLYFYQRKF--KEDGEHFKWLAKGRI
SSVVFYQTPGLEVTACVLLSTKAVYFVLHDGLRRYFSEPLQDFWHQKNTD
. . : *

FL1-18_SPLICE_VARIANT
FL1-18
Drosophila_melanogaster_CG9044
imidazoline_receptor_candidate
-----RCVLLPRDARHCRAFLEELLVDVLSLPPAWRNCVS--ATEEEV
-----SCALWLCLTAGCTC-----
YN-EQTQSLGEGLVVMSNCKCYLMEAFAPHDVAKWLRQVSVAVNRLV
YNNSPFHISQCFVLKLSLDLQSVNVGLFDQHFRLTGSTPMQVVTCLTRDSY
.

FL1-18_SPLICE_VARIANT
FL1-18
Drosophila_melanogaster_CG9044
imidazoline_receptor_candidate
TPQHRLWPLLEKSSLEARQFFYLR-----AFLVEGPSTCLVS--
-----
AIDLPLPWKIGLSFTLKDWGGFVLLLDMLRTESSLNLYLQIPLPEQCK--
LTHCFLQHLMVVLSLERTPSEPEVDKDFYSEFGNKTTGKMENYELIHS

```

Figure 8C

Incyte-2499870 Imidazoline-related Receptor




Figure 9.

FL1 - Imidazoline Receptor




Figure 10.

NOVEL IMIDAZOLINE RECEPTOR HOMOLOGS

FIELD OF THE INVENTION

[0001] The invention relates to nucleic acid and amino acid sequences of novel imidazoline receptors and to the use of these sequences in the treatment of physical and neurological disorders.

BACKGROUND OF THE INVENTION

[0002] Imidazoline receptor (IMR) subtypes bind clonidine and imidazoline (Escriba et al., 1995). These compounds mediate the regulation of blood pressure, induction of feeding, stimulation of firing of locus coeruleus neurons, and stimulation of insulin release, as well as the induction of the expression of glial fibrillary acidic protein independent of the action of alpha-2 adrenoceptors. These receptors are pharmacologically important target for drugs that can mediate the aforementioned physiological conditions (Farsang and Kapocsi, 1999).

[0003] Non-adrenoceptor sites predominantly labeled by clonidine or para-amino clonidine are termed I₁-sites whereas those non-adrenoceptor sites predominantly labeled by idazoxan are termed I₂-sites. Imidazoline sites which are distinct from either I₁- or I₂ sites are termed I₃-sites. An example is an imidazoline receptor in the pancreas reported to enhance insulin secretion. Chan et al. (1993) *Eur. J. Pharmacol.* 230 375; Chan et al. (1994) *Br. J. Pharmacol.* 112 1065. The receptor is efaroxan sensitive and it a target for the treatment of type II diabetes. The site is also sensitive to agmatine, an insulin secretagogue, and to crude preparation of clonidine displacing substance (CDS).

[0004] Endogenous ligands of the imidazoline receptors are harmane, tryptamine and agmatine. There are also numerous compounds which are selective for either I₁-sites, e.g., clonidine, benazoline and rilmenidine, or I₂-sites, e.g. RS-45041-190, 2-BFI, BU 224, and BU 239. Many of these compounds are commercially available, for example, from Tocris Cookson, Inc., USA.

[0005] I₁-site selective drugs are promising for the treatment of hypertension, I₃-site selective drugs are promising for the treatment of diabetes, and I₂-site selective drugs affect monoamine turnover and therefore I₂ receptor ligands can affect a wide range of brain functions such as nociception, ageing, mood and stroke.

SUMMARY OF THE INVENTION

[0006] The present invention relates to a novel imidazoline receptor homologs, hereinafter designated imidazoline receptor related protein 1 (IMRRP1), imidazoline receptor related protein 1b (IMRRP1b) and derivatives thereof.

[0007] Accordingly, the invention relates to a substantially purified IMRRP1 having the amino acid sequence of **FIG. 3** (SEQ ID NO: 3), or functional portion thereof, and substantially purified IMRRP1b having the amino acid sequence of **FIG. 4** (SEQ ID NO: 4).

[0008] The present invention further provides a substantially purified soluble IMRRP1. In a particular aspect, the soluble IMRRP1 comprises the amino acid sequence of **FIG. 3** (SEQ ID NO: 3). The present invention further provides a substantially purified soluble IMRRP1b. In a

particular aspect, the soluble IMRRP1 comprises the amino acid sequence of **FIG. 4** (SEQ ID NO: 4).

[0009] The present invention provides pharmaceutical compositions comprising at least one IMRRP1, IMRRP1 or a functional portion thereof.

[0010] The present invention also provides methods for producing IMRRP1, IMRRP1b or a functional portion thereof.

[0011] One aspect of the invention relates to isolated and substantially purified polynucleotides that encode IMRRP1 or IMRRP1b. In a particular aspect, the polynucleotide comprises the nucleotide sequence of **FIG. 1** (SEQ ID NO: 1). In another aspect of the invention, the polynucleotide comprises the nucleotide sequence which encodes IMRRP1. In another aspect, the polynucleotide comprises the nucleotide sequence of **FIG. 2** (SEQ ID NO: 2). In another aspect of the invention, the polynucleotide comprises the nucleotide sequence which encodes IMRRP1b.

[0012] The invention also relates to a polynucleotide sequence comprising the complement of **FIGS. 1 or 2** (SEQ ID NO: 1 or 2) or variants thereof. In addition, the invention features polynucleotide sequences which hybridize under stringent conditions to a polynucleotide sequence of **FIGS. 1 or 2** (SEQ ID NO: 1 or 2).

[0013] The invention further relates to nucleic acid sequences encoding polypeptides, oligonucleotides, fragments, portions or antisense molecules thereof, and expression vectors and host cells comprising polynucleotides that encode IMRRP 1 or IMRRP1b.

[0014] It is another object of the present invention to provide methods for producing polynucleotide sequences encoding an imidazoline receptor.

[0015] Another aspect of the invention is antibodies which bind specifically to an imidazoline receptor or epitope thereof, for use as therapeutics and diagnostic agents.

[0016] Another aspect of the invention is an agonist, antagonist or inverse agonist of IMRRP1 or IMRRP1b.

[0017] The present invention provides methods for screening for agonists, antagonists and inverse agonists of the imidazoline receptors.

[0018] It is another object of the present invention to use the nucleic acid sequences, polypeptide, peptide and antibodies for diagnosis of disorders or diseases associated with aberrant regulation of blood pressure, induction of feeding, stimulation of firing of locus coeruleus neurons, and stimulation of insulin release, as well as the aberrant induction of the expression of glial fibrillary acidic protein independent of the action of alpha-2 adrenoceptors, dysphoric premenstrual syndrome, neurodegenerative disorders such as Alzheimer's disease, opiate addiction, monoamine turnover and therefore nociception, ageing, mood and stroke, salivary disorders and developmental disorders.

[0019] The present invention provides methods of preventing or treating disorders associated with aberrant regulation of blood pressure, induction of feeding, stimulation of firing of locus coeruleus neurons, and stimulation of insulin release, as well as methods of preventing or treating disorders associated with the aberrant induction of the expression of glial fibrillary acidic protein independent of the action of

alpha-2 adrenoceptors, dysphoric premenstrual syndrome, neurodegenerative disorders such as Alzheimer's disease, opiate addiction, monoamine turnover and therefore nociception, ageing, mood and stroke, salivary disorders and developmental disorders.

[0020] The present invention provides kits for screening and diagnosis of disorders associated with aberrant IMRRP1 or IMRRP1b.

BRIEF DESCRIPTION OF THE FIGURES

[0021] These and other objects, features and many of the attendant advantages of the invention will be better understood upon a reading of the detailed description of the invention when considered in connection with the accompanying drawings wherein:

[0022] FIGS. 1A and B show the polynucleotide sequence from Clone No. FL1-18 (SEQ ID NO: 1). Clone No. FL1-18 was deposited as ATCC Deposit No. PTA-2671 on Nov. 15, 2000 at the American Type Culture Collection, Patent Depository, 10801 University Boulevard, Manassas, Va. 20110-2209.

[0023] FIGS. 2A-C show the polynucleotide sequence from Clone No. FL1-18 splice variant (SEQ ID NO: 2).

[0024] FIG. 3 shows the polypeptide sequence from IMRRP1 (SEQ ID NO: 3).

[0025] FIG. 4 shows the polypeptide sequence from IMRRP1 (SEQ ID NO: 4).

[0026] FIG. 5 shows the comparison of IMRRP1 and human imidazoline receptor Accession Number NP_009115.

[0027] FIG. 6 shows the comparison of FL1-18 to Incyte 2499870. Top strand, FL1-18; bottom strand, Incyte 2499870.

[0028] FIGS. 7A and B show the comparison of FL1-18 splice variant to *Drosophila melanogaster* CG9044, and human imidazoline receptor Accession Number NP_009115.

[0029] FIGS. 8A-D show a comparison of FL1-18 splice variant, FL1-18, *Drosophila melanogaster* CG9044, and human imidazoline receptor Accession Number NP_009115.

[0030] FIG. 9 shows the expression profile of IMRRP1.

[0031] FIG. 10 shows the expression profile of IMRRP1.

DESCRIPTION OF THE INVENTION

[0032] "Nucleic acid sequence", as used herein, refers to an oligonucleotide, nucleotide, or polynucleotide, and fragments or portions thereof, and to DNA or RNA of genomic or synthetic origin which may be single- or double-stranded, and represent the sense or antisense strand. Similarly, "amino acid sequence" as used herein refers to an oligopeptide, peptide, polypeptide, or protein sequence, and fragments or portions thereof, and to naturally occurring or synthetic molecules.

[0033] Where "amino acid sequence" is recited herein to refer to an amino acid sequence of a naturally occurring protein molecule, "amino acid sequence" and like terms,

such as "polypeptide" or "protein" are not meant to limit the amino acid sequence to the complete, native amino acid sequence associated with the recited protein molecule.

[0034] "Peptide nucleic acid", as used herein, refers to a molecule which comprises an oligomer to which an amino acid residue, such as lysine, and an amino group have been added. These small molecules, also designated anti-gene agents, stop transcript elongation by binding to their complementary strand of nucleic acid (Nielsen, P. E. et al (1993) *Anticancer Drug Des.*, 8:53-63).

[0035] IMRRP1 and IMRRP1b, as used herein, refer to the amino acid sequences of substantially purified imidazoline receptor related proteins obtained from any species, particularly mammalian, including bovine, ovine, porcine, murine, equine, and preferably human, from any source whether natural, synthetic, semi-synthetic, or recombinant.

[0036] "Consensus", as used herein, refers to a nucleic acid sequence which has been resequenced to resolve uncalled bases, or which has been extended using XL-PCR (Perkin Elmer, Norwalk, Conn.) in the 5' and/or the 3' direction and resequenced, or which has been assembled from the overlapping sequences of more than one Incyte clone or publically available clone using the GELVIEW Fragment Assembly system (GCG, Madison, Wis.), or which has been both extended and assembled.

[0037] A "variant" of IMRRP1 or IMRRP1b, as used herein, refers to an amino acid sequence that is altered by one or more amino acids. The variant may have "conservative" changes, wherein a substituted amino acid has similar structural or chemical properties, e.g., replacement of leucine with isoleucine. More rarely, a variant may have "nonconservative" changes, e.g., replacement of a glycine with a tryptophan. Similar minor variations may also include amino acid deletions or insertions, or both. Guidance in determining which amino acid residues may be substituted, inserted, or deleted without abolishing biological or immunological activity may be found using computer programs well known in the art, for example, DNASTAR software.

[0038] A "deletion", as used herein, refers to a change in either amino acid or nucleotide sequence in which one or more amino acid or nucleotide residues, respectively, are absent.

[0039] An "insertion" or "addition", as used herein, refers to a change in an amino acid or nucleotide sequence resulting in the addition of one or more amino acid or nucleotide residues, respectively, as compared to the naturally occurring molecule.

[0040] A "substitution", as used herein, refers to the replacement of one or more amino acids or nucleotides by different amino acids or nucleotides, respectively.

[0041] The term "biologically active", as used herein, refers to a protein having structural, regulatory, or biochemical functions of a naturally occurring molecule. Likewise, "immunologically active" refers to the capability of the natural, recombinant, or synthetic imidazoline receptor, or any oligopeptide thereof, to induce a specific immune response in appropriate animals or cells and to bind with specific antibodies.

[0042] The term "agonist", as used herein, refers to a molecule which when bound to IMRRP1 or IMRRP1b,

increases the amount of, or prolongs the duration of, the activity of IMRRP1 or IMRRP1b. Agonists may include proteins, nucleic acids, carbohydrates, organic molecules or any other molecules which bind to IMRRP1 or IMRRP1b.

[0043] The term “antagonist”, as used herein, refers to a molecule which, when bound to IMRRP1 or IMRRP1b, decreases the biological or immunological activity of IMRRP1 or IMRRP1b. Antagonists and inhibitors may include proteins, nucleic acids, carbohydrates, organic molecules or any other molecules which bind to IMRRP1 or IMRRP1b.

[0044] The term “mimetic”, as used herein, refers to a molecule, the structure of which is developed from knowledge of the structure of IMRRP1 or IMRRP1b or portions thereof and, as such, is able to effect some or all of the actions of IMRRP1 or IMRRP1b.

[0045] The term “derivative”, as used herein, refers to the chemical modification of a nucleic acid encoding IMRRP1 or IMRRP1b or the encoded IMRRP1 or IMRRP1b. Illustrative of such modifications would be replacement of hydrogen by an alkyl, acyl, or amino group. A nucleic acid derivative would encode a polypeptide which retains essential biological characteristics of the natural molecule.

[0046] The term “substantially purified”, as used herein, refers to nucleic or amino acid sequences that are removed from their natural environment, isolated or separated, and are at least 60% free, preferably 75% free, and most preferably 90% or greater free from other components with which they are naturally associated.

[0047] “Amplification”, as used herein, refers to the production of additional copies of a nucleic acid sequence and is generally carried out using polymerase chain reaction (PCR) technologies well known in the art Dieffenbach, D. W. and G. S. Dveksler (1995), *PCR Primer, a Laboratory Manual*, Cold Spring Harbor Press, Plainview, N.Y).

[0048] The term “hybridization”, as used herein, refers to any process by which a strand of nucleic acid binds with a complementary strand through base pairing.

[0049] The term “hybridization complex”, as used herein, refers to a complex formed between two nucleic acid sequences by virtue of the formation of hydrogen bonds between complementary G and C bases and between complementary A and T bases; these hydrogen bonds may be further stabilized by base stacking interactions. The two complementary nucleic acid sequences hydrogen bond in an antiparallel configuration. A hybridization complex may be formed in solution (e.g., C_{0t} or R_{0t} analysis) or between one nucleic acid sequence present in solution and another nucleic acid sequence immobilized on a solid support (e.g., membranes, filters, chips, pins or glass slides to which cells have been fixed in situ hybridization).

[0050] The terms “complementary” or “complementarity”, as used herein, refer to the natural binding of polynucleotides under permissive salt and temperature conditions by base-pairing. For example, the sequence “A-G-T” binds to the complementary sequence “T-C-A”. Complementarity between two single-stranded molecules may be “partial”, in which only some of the nucleic acids bind, or it may be complete when total complementarity exists between single stranded molecules. The degree of comple-

mentarity between nucleic acid strands has significant effects on the efficiency and strength of hybridization between nucleic acid strands. This is of particular importance in amplification reactions, which depend upon binding between nucleic acids strands.

[0051] The term “homology”, as used herein, refers to a degree of complementarity. There may be partial homology or complete homology (i.e., identity). A partially complementary sequence is one that at least partially inhibits an identical sequence from hybridizing to a target nucleic acid; it is referred to using the functional term “substantially homologous.” The inhibition of hybridization of the completely complementary sequence to the target sequence may be examined using a hybridization assay (Southern or northern blot, solution hybridization and the like) under conditions of low stringency. A substantially homologous sequence or probe will compete for and inhibit the binding (i.e., the hybridization) of a completely homologous sequence or probe to the target sequence under conditions of low stringency. This is not to say that conditions of low stringency are such that non-specific binding is permitted; low stringency conditions require that the binding of two sequences to one another be a specific (i.e., selective) interaction. The absence of non-specific binding may be tested by the use of a second target sequence which lacks even a partial degree of complementarity (e.g., less than about 30% identity); in the absence of non-specific binding, the probe will not hybridize to the second non-complementary target sequence.

[0052] As known in the art, numerous equivalent conditions may be employed to comprise either low or high stringency conditions. Factors such as the length and nature (DNA, RNA, base composition) of the sequence, nature of the target (DNA, RNA, base composition, presence in solution or immobilization, etc.), and the concentration of the salts and other components (e.g., the presence or absence of formamide, dextran sulfate and/or polyethylene glycol) are considered and the hybridization solution may be varied to generate conditions of either low or high stringency different from, but equivalent to, the above listed conditions.

[0053] The term “stringent conditions”, as used herein, is the “stringency” which occurs within a range from about $T_m - 5^\circ \text{C}$. (5°C . below the melting temperature T_m of the probe) to about 20°C . to 25°C . below T_m . As will be understood by those of skill in the art, the stringency of hybridization may be altered in order to identify or detect identical or related polynucleotide sequences.

[0054] The term “antisense”, as used herein, refers to nucleotide sequences which are complementary to a specific DNA or RNA sequence. The term “antisense strand” is used in reference to a nucleic acid strand that is complementary to the “sense” strand. Antisense molecules may be produced by any method, including synthesis by ligating the gene(s) of interest in a reverse orientation to a viral promoter which permits the synthesis of a complementary strand. Once introduced into a cell, this transcribed strand combines with natural sequences produced by the cell to form duplexes. These duplexes then block either the further transcription or translation. In this manner, mutant phenotypes may be generated. The designation “negative” is sometimes used in reference to the antisense strand, and “positive” is sometimes used in reference to the sense strand.

[0055] The term “portion”, as used herein, with regard to a protein (as in “a portion of a given protein”) refers to fragments of that protein. The fragments may range in size from four amino acid residues to the entire amino acid sequence minus one amino acid. Thus, a protein “comprising at least a portion of the amino acid sequence of SEQ ID NO: 3 or 4” encompasses the full-length human IMRRP1 or IMRRP1 and fragments thereof.

[0056] “Transformation”, as defined herein, describes a process by which exogenous DNA enters and changes a recipient cell. It may occur under natural or artificial conditions using various methods well known in the art. Transformation may rely on any known method for the insertion of foreign nucleic acid sequences into a prokaryotic or eukaryotic host cell. The method is selected based on the host cell being transformed and may include, but is not limited to, viral infection, electroporation, lipofection, and partial bombardment. Such “transformed” cells include stably transformed cells in which the inserted DNA is capable of replication either as an autonomously replicating plasmid or as part of the host chromosome. They also include cells which transiently express the inserted DNA or RNA for limited periods of time.

[0057] The term “antigenic determinant”, as used herein, refers to that portion of a molecule that makes contact with a particular antibody (i.e., an epitope). When a protein or fragment of a protein is used to immunize a host animal, numerous regions of the protein may induce the production of antibodies which bind specifically to a given region or three-dimensional structure on the protein; these regions or structures are referred to as antigenic determinants. An antigenic determinant may compete with the intact antigen (i.e., the immunogen used to elicit the immune response) for binding to an antibody.

[0058] The terms “specific binding” or “specifically binding”, as used herein, in reference to the interaction of an antibody and a protein or peptide, mean that the interaction is dependent upon the presence of a particular structure (i.e., the antigenic determinant or epitope) on the protein; in other words, the antibody is recognizing and binding to a specific protein structure rather than to proteins in general. For example, if an antibody is specific for epitope “A”, the presence of a protein containing epitope A (or free, unlabeled A) in a reaction containing labeled “A” and the antibody will reduce the amount of labeled A bound to the antibody.

[0059] The term “sample”, as used herein, is used in its broadest sense. A biological sample suspected of containing nucleic acid encoding IMRRP1 or IMRRP1b or fragments thereof may comprise a cell, chromosomes isolated from a cell (e.g., a spread of metaphase chromosomes), genomic DNA (in solution or bound to a solid support such as for Southern analysis), RNA (in solution or bound to a solid support such as for northern analysis), cDNA (in solution or bound to a solid support), an extract from cells or a tissue, and the like.

[0060] The term “correlates with expression of a polynucleotide”, as used herein, indicates that the detection of the presence of ribonucleic acid that is similar to SEQ ID NOS: 1 or 2 by northern analysis is indicative of the presence of mRNA encoding IMRRP1 and IMRRP1b in a sample and thereby correlates with expression of the transcript from the polynucleotide encoding the protein.

[0061] “Alterations” in the polynucleotide of SEQ ID NOS: 1 and 2 as used herein, comprise any alteration in the sequence of polynucleotides encoding IMRRP1 and IMRRP1b including deletions, insertions, and point mutations that may be detected using hybridization assays. Included within this definition is the detection of alterations to the genomic DNA sequence which encodes IMRRP1 or IMRRP1b (e.g., by alterations in the pattern of restriction fragment length polymorphisms capable of hybridizing to SEQ ID NOS: 1 or 2), the inability of a selected fragment of SEQ ID NOS: 1 or 2 to hybridize to a sample of genomic DNA (e.g., using allele-specific oligonucleotide probes), and improper or unexpected hybridization, such as hybridization to a locus other than the normal chromosomal locus for the polynucleotide sequence encoding IMRRP1 or IMRRP1b (e.g., using fluorescent in situ hybridization (FISH) to metaphase chromosome spreads).

[0062] As used herein, the term “antibody” refers to intact molecules as well as fragments thereof, such as Fa, F(ab')₂, Fv, chimeric antibody, single chain antibody which are capable of binding the epitopic determinant. Antibodies that bind IMRRP1 or IMRRP1b polypeptides can be prepared using intact polypeptides or fragments containing small peptides of interest or prepared recombinantly for use as the immunizing antigen. The polypeptide or peptide used to immunize an animal can be derived from the transition of RNA or synthesized chemically, and can be conjugated to a carrier protein, if desired. Commonly used carriers that are chemically coupled to peptides include bovine serum albumin and thyroglobulin. The coupled peptide is then used to immunize the animal, e.g., a mouse, a rat, or a rabbit.

[0063] The term “humanized antibody”, as used herein, refers to antibody molecules in which amino acids have been replaced in the non-antigen binding regions in order to more closely resemble a human antibody, while still retaining the original binding ability.

[0064] The deposit(s) referred to herein will be maintained under the terms of the Budapest Treaty on the International Recognition of the Deposit of Microorganisms for purposes of Patent Procedure. These deposits are provided merely as convenience to those of skill in the art and are not an admission that a deposit is required under 35 U.S.C. §112. The sequence of the polynucleotides contained in the deposited materials, as well as the amino acid sequence of the polypeptides encoded thereby, are incorporated herein by reference and are controlling in the event of any conflict with any description of sequences herein. A license may be required to make, or sell the deposited materials, and no such license is hereby granted.

[0065] The invention is novel human imidazoline receptors referred to as IMRRP1 and IMRRP1b, polynucleotides encoding IMRRP1 and IMRRP1b, and the use of these compositions for the diagnosis, prevention, or treatment of disorders associated with aberrant cellular development, immune responses and inflammation, as well as organ and tissue transplantation rejection.

[0066] Human imidazoline receptor protein sequence was used as a probe to search the Incyte and public domain EST databases. The search program used was gapped BLAST (Altschul et al., 1997). The top EST hits from the BLAST results were searched back against the non-redundant protein and patent sequence databases. From this analysis, ESTs

encoding a potential novel imidazoline receptor was identified based on sequence homology. The Incyte EST (Clone ID: 2499870) was selected as a potential novel imidazoline receptor candidate for subsequent analysis.

[0067] A PCR primer pair, designed from the DNA sequence of Incyte clone-2499870 was used to amplify a piece of DNA from the clone in which the anti-sense strand of the amplified fragment was biotinylated on the 5' end. This biotinylated piece of double stranded DNA was denatured and incubated with a mixture of single-stranded covalently closed circular cDNA libraries which contain DNA corresponding to the sense strand. The cDNA libraries were total brain tissue libraries obtained from Gibco Life Technologies. Hybrids between the biotinylated DNA and the circular cDNA were captured on streptavidin magnetic beads. Upon thermal release of the cDNA from the biotinylated DNA, the single stranded cDNA was converted into double strands using a primer homologous to a sequence on the cDNA cloning vector. The double stranded cDNA was introduced into *E. coli* by electroporation and the resulting colonies were screen by PCR, using the original primer pair to identify the proper cDNA clones. One clone named FL1-18 was sequenced on both strands (FIG. 1). The deduced amino acid sequence corresponding to the nucleic acid sequence of clone FL1-18 is shown in FIG. 3.

[0068] A comparison of the FL1-18 cDNA to that of the partial clone found in the Incyte database (clone 2499870) revealed that at nucleotide position 1725 of the Incyte clone a small insertion of 25 bases occurs and at position 3375 of clone FL1-18 an insertion of 47 bases occurs. (See FIG. 6: Top strand, FL1-18; bottom strand, Incyte 2499870.) An alignment of the two DNA sequences, FL1-18 and Incyte 2499870 is shown in FIG. 8.

[0069] An alignment of the two DNA sequences, FL1-18 and Incyte 2499870 to the rough draft of the human genome, revealed that both insertions/deletion in the two sequences correspond to putative exons as determined by the conservation of splice donor and acceptor sequences on either side of the inserted DNA and hence represent different RNA splice forms of a transcript that originates from one genomic location (i.e., one gene).

[0070] The Incyte clone is missing approximately 450 bp of the 5'-end. Combining the 5'-end sequences of FL1-18 sequence with that of the Incyte clone creates a novel nucleotide sequence which is referred to the FL1-18 splice variant. Translation of this sequence produces a longer polypeptide chain than that of FL1-18 because of the elimination of an in frame stop caused by the lack of the small exon in FL1-18. The first 712 amino acid are identical, but after that the remaining 97 amino acids of FL1-18 differ. The second alternatively spliced exon found in the Incyte clone is a coding exon. Hence, these splice variants produce different length and possibly different functional proteins.

[0071] In one embodiment, the invention encompasses a polypeptide comprising the amino acid sequence of SEQ ID NO: 3 as shown in FIG. 3, or the amino acid sequence of SEQ ID NO: 4 as shown in FIG. 4. IMRRP1 and IMRRP1b share chemical and structural homology with the human imidazoline receptor, Accession number NP_009115. IMRRP1 and IMRRP1b also share chemical and structural homology with two *Drosophila* proteins identified as Accession number AAF52305 and Accession number AAF57514.

IMRRP1 shares 26% identity with the human imidazoline receptor, Accession number NP_009115, as illustrated in FIG. 5.

[0072] Expression profiling of imidazoline receptor homolog IMRRP1 showed expression in a variety of human tissue. The same PCR primer used in the cloning of imidazoline receptor IMRRP1 was used to measure the steady state levels of mRNA by quantitative PCR. Briefly, first strand cDNA was made from commercially available mRNA. The relative amount of cDNA used in each assay was determined by performing a parallel experiment using a primer pair for a gene expressed in equal amounts in all tissues, cyclophilin. The cyclophilin primer pair detected small variations in the amount of cDNA in each sample and these data were used for normalization of the data obtained with the primer pair for IMRRP1. The PCR data was converted into a relative assessment of the difference in transcript abundance amongst the tissues tested and the data is presented in FIG. 9.

[0073] The invention also encompasses IMRRP1 and IMRRP1b variants. Preferred IMRRP1 and IMRRP1b variants are those having at least 80%, and more preferably 90% or greater, amino acid identity to the IMRRP1 and IMRRP1b amino acid sequence of SEQ ID NOS: 3 and 4, respectively. Most preferred IMRRP1 and IMRRP1b variants are those having at least 95% amino acid sequence identity to SEQ ID NOS: 3 and 4, respectively.

[0074] The present invention provides isolated IMRRP1 and IMRRP1b and homologs thereof. Such proteins are substantially free of contaminating endogenous materials and, optionally, without associated nature-pattern glycosylation. Derivatives of the IMRRP1 and IMRRP1b receptors within the scope of the invention also include various structural forms of the primary protein which retain biological activity. Due to the presence of ionizable amino and carboxyl groups, for example, IMRRP1 and IMRRP1 proteins may be in the form of acidic or basic salts, or may be in neutral form. Individual amino acid residues may also be modified by oxidation or reduction.

[0075] The primary amino acid structure may be modified by forming covalent or aggregative conjugates with other chemical moieties, such as glycosyl groups, lipids, phosphate, acetyl groups and the like, or by creating amino acid sequence mutants. Covalent derivatives are prepared by linking particular functional groups to amino acid side chains or at the N- or C-termini.

[0076] The present invention farther encompassed fusion proteins comprising the amino acid sequence of IMRRP1 or IMRRP1b or portions thereof linked to an immunoglobulin Fc region. Depending on the portion of the Fc region used, a fusion protein may be expressed as a dimer, through formation of interchain disulfide bonds. If the fusion proteins are made with both heavy and light chains of an antibody, it is possible to form a protein oligomer with as many as four IMRRP1 and/or IMRRP1 regions.

[0077] The invention also encompasses polynucleotides which encode IMRRP1 and IMRRP1b. Accordingly, any nucleic acid sequence which encodes the amino acid sequence of IMRRP1 or IMRRP1b can be used to generate recombinant molecules which express IMRRP1 and IMRRP1b. In a particular embodiment, the invention

encompasses the polynucleotide comprising the nucleic acid sequence of SEQ ID NOS. 1 and 2 as shown in FIGS. 1 and 2.

[0078] It will be appreciated by those skilled in the art that as a result of the degeneracy of the genetic code, a multitude of nucleotide sequences encoding IMRRP1 and IMRRP1b, some bearing minimal homology to the nucleotide sequences of any known and naturally occurring gene, may be produced. Thus, the invention contemplates each and every possible variation of nucleotide sequence that could be made by selecting combinations based on possible codon choices. These combinations are made in accordance with the standard triplet genetic code as applied to the nucleotide sequence of naturally occurring IMRRP1 and IMRRP1b, and all such variations are to be considered as being specifically disclosed.

[0079] Although nucleotide sequences which encode IMRRP1 or IMRRP1b and their variants are preferably capable of hybridizing to the nucleotide sequence of the naturally occurring coding sequence for IMRRP1 or IMRRP1b under appropriately selected conditions of stringency, it may be advantageous to produce nucleotide sequences encoding IMRRP1 or IMRRP1b or their derivatives possessing a substantially different codon usage. Codons may be selected to increase the rate at which expression of the peptide occurs in a particular prokaryotic or eukaryotic host in accordance with the frequency with which particular codons are utilized by the host. Other reasons for substantially altering the nucleotide sequence encoding IMRRP1 or IMRRP1b and their derivatives without altering the encoded amino acid sequences include the production of RNA transcripts having more desirable properties, such as a greater half-life, than transcripts produced from the naturally occurring sequence.

[0080] The invention also encompasses production of DNA sequences, or portions thereof, which encode IMRRP1 or IMRRP1b and their derivatives, entirely by synthetic chemistry. After production, the synthetic sequence may be inserted into any of the many available expression vectors and cell systems using reagents that are well known in the art at the time of the filing of this application. Moreover, synthetic chemistry may be used to introduce mutations into a sequence encoding IMRRP1 or IMRRP1b or any portion thereof.

[0081] Also encompassed by the invention are polynucleotide sequences that are capable of hybridizing to the claimed nucleotide sequences, and in particular, those shown in SEQ ID NOS: 1 and 2, under various conditions of stringency. Hybridization conditions are based on the melting temperature (T_m) of the nucleic acid binding complex or probe, as taught in Wahl, G. M. and S. L. Berger (1987; *Methods Enzymol.* 152:399-407) and Kimmel, A. R. (1987; *Methods of Enzymol.* 152:507-511), and may be used at a defined stringency. In one embodiment, sequences include those capable of hybridizing under moderately stringent conditions (prewashing solution of 2×SSC, 0.5% SOS, 1.0 mM MEDTA, pH 8.0) and hybridization conditions of 50° C., 5×SSC, overnight, to the sequences encoding IMRRP1 or IMRRP1b and other sequences which are degenerate to those which encode IMRRP1 or IMRRP1b.

[0082] Altered nucleic acid sequences encoding IMRRP1 or IMRRP1b which are encompassed by the invention

include deletions, insertions, or substitutions of different nucleotides resulting in a polynucleotide that encodes the same or a functionally equivalent IMRRP1 or IMRRP1b. The encoded protein may also contain deletions, insertions, or substitutions of amino acid residues which produce a silent change and result in a functionally equivalent IMRRP1 or IMRRP1b. Deliberate amino acid substitutions may be made on the basis of similarity in polarity, charge, solubility, hydrophobicity, hydrophilicity, and/or the amphipathic nature of the residues as long as the biological activity of IMRRP1 and IMRRP1b is retained. For example, negatively charged amino acids may include aspartic acid and glutamic acid; positively charged amino acids may include lysine and arginine; and amino acids with uncharged polar head groups having similar hydrophilicity values may include leucine, isoleucine, and valine; glycine and alanine; asparagine and glutamine; serine and threonine; phenylalanine and tyrosine.

[0083] Also included within the scope of the present invention are alleles of the genes encoding IMRRP1 and IMRRP1b. As used herein, an "allele" or "allelic sequence" is an alternative form of the gene which may result from at least one mutation in the nucleic acid sequence. Alleles may result in altered mRNAs or polypeptides whose structure or function may or may not be altered. Any given gene may have none, one, or many allelic forms. Common mutational changes which give rise to alleles are generally ascribed to natural deletions, additions, or substitutions of nucleotides. Each of these types of changes may occur alone, or in combination with the others, one or more times in a given sequence.

[0084] Methods for DNA sequencing which are well known and generally available in the art may be used to practice any embodiments of the invention. The methods may employ such enzymes as the Klenow fragment of DNA polymerase I, SEQUENCE (US Biochemical Corp. Cleveland, Ohio), Taq polymerase (Perkin Elmer), thermostable T7 polymerase (Amersham, Chicago, Ill.), or combinations of recombinant polymerases and proofreading exonucleases such as the ELONGASE Amplification System marketed by Gibco BRL (Gaithersburg, Md.). Preferably, the process is automated with machines such as the Hamilton Micro Lab 2200 (Hamilton, Reno, Nev.), Peltier Thermal Cycler (PTC200; M J Research, Watertown, Mass.) and the ABI 377 DNA sequencers (Perkin Elmer).

[0085] The nucleic acid sequences encoding IMRRP1 or IMRRP1b may be extended utilizing a partial nucleotide sequence and employing various methods known in the art to detect upstream sequences such as promoters and regulatory elements. For example, one method which may be employed, "restriction-site" PCR, uses universal primers to retrieve unknown sequence adjacent to a known locus (Sarkar, G. (1993) *PCR Methods Applic.* 2:318-322). In particular, genomic DNA is first amplified in the presence of primer to linker sequence and a primer specific to the known region. The amplified sequences are then subjected to a second round of PCR with the same linker primer and another specific primer internal to the first one. Products of each round of PCR are transcribed with an appropriate RNA polymerase and sequenced using reverse transcriptase.

[0086] Inverse PCR may also be used to amplify or extend sequences using divergent primers based on a known region

(Triglia, T. et al. (1988) *Nucleic Acids Res.* 16:8186). The primers may be designed using OLIGO 4.06 Primer Analysis software (National Biosciences Inc., Plymouth, Minn.), or another appropriate program, to be 22-30 nucleotides in length, to have a GC content of 50% or more, and to anneal to the target sequence at temperatures about 68° C. to about 72° C. The method uses several restriction enzymes to generate a suitable fragment in the known region of a gene. The fragment is then circularized by intramolecular ligation and used as a PCR template.

[0087] Another method which may be used is capture PCR which involves PCR amplification of DNA fragments adjacent to a known sequence in human and yeast artificial chromosome DNA (Lagerstrom, M. et al. (1991) *PCR Methods Applic.* 1:111-119). In this method, multiple restriction enzyme digestions and ligations may also be used to place an engineered double-stranded sequence into an unknown portion of the DNA molecule before performing PCR.

[0088] Another method which may be used to retrieve unknown sequences is that of Parker, J. D. et al. (1991; *Nucleic Acids Res.* 19:3055-3060). Additionally, one may use PCR, nested primers, and PROMOTERFINDER libraries to walk in genomic DNA (Clontech, Palo Alto, Calif.). This process avoids the need to screen libraries and is useful in finding intron/exon junctions.

[0089] When screening for full-length cDNAs, it is preferable to use libraries that have been size-selected to include larger cDNAs. Also, random-primed libraries are preferable, in that they will contain more sequences which contain the 5' regions of genes. Use of a randomly primed library may be especially preferable for situations in which an oligo d(T) library does not yield a full-length cDNA. Genomic libraries may be useful for extension of sequence into the 5' and 3' non-transcribed regulatory regions.

[0090] Capillary electrophoresis systems which are commercially available may be used to analyze the size or confirm the nucleotide sequence of sequencing or PCR products. In particular, capillary sequencing may employ flowable polymers for electrophoretic separation, four different fluorescent dyes (one for each nucleotide) which are laser activated, and detection of the emitted wavelengths by a charge coupled device camera. Output/light intensity may be converted to electrical signal using appropriate software (e.g., GENOTYPER and SEQUENCE NAVIGATOR, Perkin Elmer) and the entire process from loading of samples to computer analysis and electronic data display may be computer controlled. Capillary electrophoresis is especially preferable for the sequencing of small pieces of DNA which might be present in limited amounts in a particular sample.

[0091] In another embodiment of the invention, polynucleotide sequences or fragments thereof which encode or fusion proteins or functional equivalents thereof, may be used in recombinant DNA molecules to direct expression of IMRRP1 or IMRRP1b in appropriate host cells. Due to the inherent degeneracy of the genetic code, other DNA sequences which encode substantially the same or a functionally equivalent amino acid sequence may be produced and these sequences may be used to clone and express IMRRP1 or IMRRP1b.

[0092] As will be understood by those of skill in the art, it may be advantageous to produce IMRRP1- or IMRRP1b-

encoding nucleotide sequences possessing non-naturally occurring codons. For example, codons preferred by a particular prokaryotic or eukaryotic host can be selected to increase the rate of protein expression or to produce a recombinant RNA transcript having desirable properties, such as a half-life which is longer than that of a transcript generated from the naturally occurring sequence.

[0093] The nucleotide sequences of the present invention can be engineered using methods generally known in the art in order to alter the IMRRP1 and IMRRP1b encoding sequences for a variety of reasons, including but not limited to, alterations which modify the cloning, processing, and/or expression of the gene product. DNA shuffling by random fragmentation and PCR reassembly of gene fragments and synthetic oligonucleotides may be used to engineer the nucleotide sequences. For example, site-directed mutagenesis may be used to insert new restriction sites, alter glycosylation patterns, change codon preference, produce splice variants, or introduce mutations, and so forth.

[0094] In another embodiment of the invention, natural, modified, or recombinant nucleic acid sequences encoding IMRRP1 or IMRRP1b may be ligated to a heterologous sequence to encode a fusion protein. For example, to screen peptide libraries for inhibitors of IMRRP1 or IMRRP1b activity, it may be useful to encode a chimeric IMRRP1 or IMRRP1b protein that can be recognized by a commercially available antibody. A fusion protein may also be engineered to contain a cleavage site located between the IMRRP1 or IMRRP1b encoding sequence and the heterologous protein sequence, so that IMRRP1 or IMRRP1b may be cleaved and purified away from the heterologous moiety.

[0095] In another embodiment, sequences encoding IMRRP1 or IMRRP1b may be synthesized, in whole or in part, using chemical methods well known in the art (see Caruthers, M. H. et al. (1980) *Nuci. Acids Res. Symp. Ser.* 215-223, Horn, T. et al. (1980) *Nucl. Acids Res. Symp. Ser.* 225-232). Alternatively, the protein itself may be produced using chemical methods to synthesize the amino acid sequence of IMRRP1 or IMRRP1b, or a portion thereof. For example, peptide synthesis can be performed using various solid-phase techniques (Roberge, J. Y. et al. (1995) *Science* 269:202-204) and automated synthesis may be achieved, for example, using the ABI 431A Peptide Synthesizer (Perkin Elmer).

[0096] The newly synthesized peptide may be substantially purified by preparative high performance liquid chromatography (e.g., Creighton, T. (1983) *Proteins, Structures and Molecular Principles*, W H Freeman and Co., New York, N.Y.), by reverse-phase high performance liquid chromatography, or other purification methods as are known in the art. The composition of the synthetic peptides may be confirmed by amino acid analysis or sequencing (e.g., the Edman degradation procedure; Creighton, supra). Additionally, the amino acid sequence of IMRRP1 or IMRRP1b, or any part thereof, may be altered during direct synthesis and/or combined using chemical methods with sequences from other proteins, or any part thereof, to produce a variant polypeptide.

[0097] In order to express a biologically active IMRRP1 or IMRRP1b the nucleotide sequences encoding IMRRP1 or IMRRP1b or functional equivalents, may be inserted into an appropriate expression vector, i.e., a vector which contains

the necessary elements for the transcription and translation of the inserted coding sequence.

[0098] Methods which are well known to those skilled in the art may be used to construct expression vectors containing sequences encoding and appropriate transcriptional and translational control elements. These methods include in vitro recombinant DNA techniques, synthetic techniques, and in vivo genetic recombination. Such techniques are described in Sambrook, J. et al. (1989) *Molecular Cloning, A Laboratory Manual*, Cold Spring Harbor Press, Plainview, N.Y., and Ausubel, F. M. et al. (1989) *Current Protocols in Molecular Biology*, John Wiley & Sons, New York, N.Y.

[0099] A variety of expression vector/host systems may be utilized to contain and express sequences encoding IMRRP1 or IMRRP1b. These include, but are not limited to, microorganisms such as bacteria transformed with recombinant bacteriophage, plasmid, or cosmid DNA expression vectors; yeast transformed with yeast expression vectors; insect cell systems infected with virus expression vectors (e.g., baculovirus); plant cell systems transformed with virus expression vectors (e.g., cauliflower mosaic virus, CaMV or tobacco mosaic virus, TMV) or with bacterial expression vectors (e.g., Ti or pBR322 plasmids); or animal cell systems.

[0100] The "control elements" or "regulatory sequences" are those non-translated regions of the vector-enhancers, promoters, 5' and 3' untranslated regions which interact with host cellular proteins to carry out transcription and translation. Such elements may vary in their strength and specificity. Depending on the vector system and host utilized, any number of suitable transcription and translation elements, including constitutive and inducible promoters, may be used. For example, when cloning in bacterial systems, inducible promoters such as the hybrid lacZ promoter of the BLUESCRIPT phagemid (Stratagene, LaJolla, Calif.) or PSPORT1 plasmid (Gibco BRL) and the like may be used. The baculovirus polyhedrin promoter may be used in insect cells. Promoters or enhancers derived from the genomes of plant cells (e.g., heat shock, RUBISCO; and storage protein genes) or from plant viruses (e.g., viral promoters or leader sequences) may be cloned into the vector. In mammalian cell systems, promoters from mammalian genes or from mammalian viruses are preferable. If it is necessary to generate a cell line that contains multiple copies of the sequence encoding IMRRP1 or IMRRP1b, vectors based on SV40 or EBV may be used with an appropriate selectable marker.

[0101] In bacterial systems, a number of expression vectors may be selected depending upon the use intended for IMRRP1 or IMRRP1b. For example, when large quantities are needed for the induction of antibodies, vectors which direct high level expression of fusion proteins that are readily purified may be used. Such vectors include, but are not limited to, the multifunctional *E. coli* cloning and expression vectors such as BLUESCRIPT (Stratagene), in which the sequence encoding IMRRP1 or IMRRP1b may be ligated into the vector in frame with sequences for the amino-terminal Met and the subsequent 7 residues of β -galactosidase so that a hybrid protein is produced, pIN vectors (Van Heeke, G. and S. M. Schuster (1989) *J. Biol. Chem.* 264:5503-5509); and the like. pGEX vectors (Promega, Madison, Wis.) may also be used to express foreign polypeptides, as fusion proteins with glutathione S-trans-

ferase (GST). In general, such fusion proteins are soluble and can easily be purified from lysed cells by adsorption to glutathione-agarose beads followed by elution in the presence of free glutathione. Proteins made in such systems may be designed to include heparin, thrombin, or factor XA protease cleavage sites so that the cloned polypeptide of interest can be released from the GST moiety at will.

[0102] In the yeast, *Saccharomyces cerevisiae*, a number of vectors containing constitutive or inducible promoters such as alpha factor, alcohol oxidase, and PGH may be used. For reviews, see Ausubel et al. (supra) and Grant et al. (1987) *Methods Enzymol.* 153:516-544.

[0103] In cases where plant expression vectors are used, the expression of sequences encoding IMRRP1 or IMRRP1b may be driven by any of a number of promoters. For example, viral promoters such as the 35S and 19S promoters of CaMV may be used alone or in combination with the omega leader sequence from TMV (Takamatsu, N. (1987) *EMBO J.* 6:307-311). Alternatively, plant promoters such as the small subunit of RUBISCO or heat shock promoters may be used (Coruzzi, G. et al. (1984) *EMBO J.* 3:1671-1680; Broglie, R. et al. (1984) *Science* 224:838-843; and Winter, J. et al. (1991) *Results Probl. Cell Differ.* 17:85-105). These constructs can be introduced into plant cells by direct DNA transformation or pathogen-mediated transfection. Such techniques are described in a number of generally available reviews (see, for example, Hobbs, S. or Murry, L. E. in McGraw Hill *Yearbook of Science and Technology* (1992) McGraw Hill, New York, N.Y.; pp. 191-196).

[0104] An insect system may also be used to express IMRRP1 or IMRRP1b. For example, in one such system, *Autographa californica* nuclear polyhedrosis virus (AcNPV) is used as a vector to express foreign genes in *Spodoptera frugiperda* cells or in *Trichoplusia larvae*. The sequences encoding IMRRP1 or IMRRP1b may be cloned into a non-essential region of the virus such as the polyhedrin gene, and placed under control of the polyhedrin promoter. Successful insertion of IMRRP1 or IMRRP1b will render the polyhedrin gene inactive and produce recombinant virus lacking coat protein. The recombinant viruses may then be used to infect, for example, *S. frugiperda* cells or *Trichoplusia larvae* in which IMRRP1 or IMRRP1b may be expressed (Engelhard, E. K. et al. (1994) *Proc. Nat. Acad. Sci.* 91:3224-3227).

[0105] In mammalian host cells, a number of viral-based expression systems may be utilized. In cases where an adenovirus is used as an expression vector, sequences encoding IMRRP1 or IMRRP1b may be ligated into an adenovirus transcription/translation complex consisting of the late promoter and tripartite leader sequence. Insertion in a non-essential E1 or E3 region of the viral genome may be used to obtain a viable virus which is capable of expressing IMRRP1 or IMRRP1b in infected host cells (Logan, J. and Shenk, T. (1984) *Proc. Natl. Acad. Sci.* 81:3655-3659). In addition, transcription enhancers, such as the Rous, sarcoma virus (RSV) enhancer, may be used to increase expression in mammalian host cells.

[0106] Specific initiation signals may also be used to achieve more efficient translation of sequences encoding IMRRP1 or IMRRP1b. Such signals include the ATG initiation codon and adjacent sequences. In cases where

sequences encoding IMRRP1 or IMRRP1b, their initiation codon, and upstream sequences are inserted into the appropriate expression vector, no additional transcriptional or translational control signals may be needed. However, in cases where only a coding sequence, or a portion thereof, is inserted, exogenous translational control signals including the ATG initiation codon should be provided. Furthermore, the initiation codon should be in the correct reading frame to ensure translation of the entire insert. Exogenous translational elements and initiation codons may be of various origins, both natural and synthetic. The efficiency of expression may be enhanced by the inclusion of enhancers which are appropriate for the particular cell system which is used, such as those described in the literature (Scharf, D. et al. (1994) *Results Probl. Cell Differ.* 20:125-162).

[0107] In addition, a host cell strain may be chosen for its ability to modulate the expression of the inserted sequences or to process the expressed protein in the desired fashion. Such modifications of the polypeptide include, but are not limited to, acetylation, carboxylation, glycosylation, phosphorylation, lipidation, and acylation. Post-translational processing which cleaves a "prepro" form of the protein may also be used to facilitate correct insertion, folding and/or function. Different host cells such as CHO, HeLa, MDCK, HEK293, and W138, which have specific cellular machinery and characteristic mechanisms for such post-translational activities, may be chosen to ensure the correct modification and processing of the foreign protein.

[0108] For long-term, high-yield production of recombinant proteins, stable expression is preferred. For example, cell lines which stably express IMRRP1 or IMRRP1b may be transformed using expression vectors which may contain viral origins of replication and/or endogenous expression elements and a selectable marker gene on the same or on a separate vector. Following the introduction of the vector, cells may be allowed to grow for 1-2 days in an enriched media before they are switched to selective media. The purpose of the selectable marker is to confer resistance to selection, and its presence allows growth and recovery of cells which successfully express the introduced sequences. Resistant clones of stably transformed cells may be proliferated using tissue culture techniques appropriate to the cell type.

[0109] Any number of selection systems may be used to recover transformed cell lines. These include, but are not limited to, the herpes simplex virus thymidine kinase (Wigler, M. et al. (1977) *Cell* 11:223-32) and adenine phosphoribosyltransferase (Lowy, I. et al. (1980) *Cell* 22:817-23) genes which can be employed in tk or apt cells, respectively. Also, antimetabolite, antibiotic or herbicide resistance can be used as the basis for selection; for example, dhfr which confers resistance to methotrexate (Wigler, M. et al. (1980) *Proc. Natl. Acad. Sci.* 77:3567-70); npt, which confers resistance to the aminoglycosides, neomycin and G-418 (Colbere-Garapin, F. et al (1981) *J. Mol. Biol.* 150:1-14); and als or pat, which confer resistance to chlorsulfuron and phosphinotricin acetyltransferase, respectively (Murry, supra). Additional selectable genes have been described, for example, trpB, which allows cells to utilize indole in place of tryptophan, or hisd, which allows cells to utilize histinol in place of histidine (Hartman, S. C. and R. C. Mulligan (1988) *Proc. Natl. Acad. Sci.* 85:8047-51). Recently, the use of visible markers has gained popularity with such markers

as anthocyanins, B glucuronidase and its substrate GUS, and luciferase and its substrate luciferin, being widely used not only to identify transformants, but also to quantify the amount of transient or stable protein expression attributable to a specific vector system (Rhodes, C. A. et al. (1995) *Methods Mol. Biol.* 55:121-131).

[0110] Although the presence/absence of marker gene expression suggests that the gene of interest is also present, its presence and expression may need to be confirmed. For example, if the sequence encoding IMRRP1 or IMRRP1b is inserted within a marker gene sequence, recombinant cells containing sequences encoding can be identified by the absence of marker gene function. Alternatively, a marker gene can be placed in tandem with a sequence encoding IMRRP1 or IMRRP1b under the control of a single promoter. Expression of the marker gene in response to induction or selection usually indicates expression of the tandem gene as well.

[0111] Alternatively, host cells which contain the nucleic acid sequence encoding IMRRP1 or IMRRP1b and express IMRRP1 or IMRRP1b may be identified by a variety of procedures known to those of skill in the art. These procedures include, but are not limited to, DNA-DNA or DNA-RNA hybridizations and protein bioassay or immunoassay techniques which include membrane, solution, or chip based technologies for the detection and/or quantification of nucleic acid or protein.

[0112] The presence of polynucleotide sequences encoding IMRRP1 or IMRRP1b can be detected by DNA-DNA or DNA-RNA hybridization or amplification using probes or portions or fragments of polynucleotides encoding IMRRP1 or IMRRP1b. Nucleic acid amplification based assays involve the use of oligonucleotides or oligomers based on the sequences encoding IMRRP1 or IMRRP1b to detect transformants containing DNA or RNA encoding IMRRP1 or IMRRP1b. As used herein "oligonucleotides" or "oligomers" refer to a nucleic acid sequence of at least about 10 nucleotides and as many as about 60 nucleotides, preferably about 15 to 30 nucleotides, and more preferably about 20-25 nucleotides, which can be used as a probe or amplicon.

[0113] A variety of protocols for detecting and measuring the expression of IMRRP1 or IMRRP1b, using either polyclonal or monoclonal antibodies specific for the proteins are known in the art. Examples include enzyme-linked immunosorbent assay (ELISA), radioimmunoassay (RIA), and fluorescence activated cell sorting (FACS). A two-site, monoclonal-based immunoassay utilizing monoclonal antibodies reactive to two non-interfering epitopes on IMRRP1 or IMRRP1b is preferred, but a competitive binding assay may be employed. These and other assays are described, among other places, in Hampton, R. et al. (1990; *Serological Methods, a Laboratory Manual*, APS Press, St Paul, Minn.) and Maddox, D. E. et al. (1983; *J. Exp. Med.* 158:1211-1216).

[0114] A wide variety of labels and conjugation techniques are known by those skilled in the art and may be used in various nucleic acid and amino acid assays. Means for producing labeled hybridization or PCR probes for detecting sequences related to polynucleotides encoding IMRRP1 or IMRR1b include oligolabeling, nick translation, end-labeling or PCR amplification using a labeled nucleotide. Alternatively, the sequences encoding IMRRP1 or IMRRP1b, or

any portions thereof may be cloned into a vector for the production of an mRNA probe. Such vectors are known in the art, are commercially available, and may be used to synthesize RNA probes in vitro by addition of an appropriate RNA polymerase such as T7, T3, or SP6 and labeled nucleotides. These procedures may be conducted using a variety of commercially available kits (Pharmacia & Upjohn, (Kalamazoo, Mich.); Promega (Madison Wis.); and U.S. Biochemical Corp., (Cleveland, Ohio)). Suitable reporter molecules or labels, which may be used, include radionuclides, enzymes, fluorescent, chemiluminescent, or chromogenic agents as well as substrates, cofactors, inhibitors, magnetic particles, and the like.

[0115] Host cells transformed with nucleotide sequences encoding IMRRP1 or IMRRP1b may be cultured under conditions suitable for the expression and recovery of the protein from cell culture. The protein produced by a recombinant cell may be secreted or contained intracellularly depending on the sequence and/or the vector used. As will be understood by those of skill in the art, expression vectors containing polynucleotides which encode IMRRP1 or IMRRP1b may be designed to contain signal sequences which direct secretion of IMRRP1 or IMRRP1b through a prokaryotic or eukaryotic cell membrane. Other recombinant constructions may be used to join sequences encoding IMRRP1 or IMRRP1b to nucleotide sequence encoding a polypeptide domain which will facilitate purification of soluble proteins. Such purification facilitating domains include, but are not limited to, metal chelating peptides such as histidine tryptophan modules that allow purification on immobilized metals, protein A domains that allow purification on immobilized immunoglobulin, and the domain utilized in the FLAGS extension/affinity purification system (Immunex Corp., Seattle, Wash.). The inclusion of cleavable linker sequences such as those specific for Factor XA or enterokinase (Invitrogen, San Diego, Calif.) between the purification domain and IMRRP1 or IMRRP1b may be used to facilitate purification. One such expression vector provides for expression of a fusion protein containing IMRRP1 or IMRRP1b and a nucleic acid encoding 6 histidine residues preceding a thioredoxin or an enterokinase cleavage site. The histidine residues facilitate purification on IMIAC (immobilized metal ion affinity chromatography) as described in Porath, J et al. (1992, *Prot. Exp. Purif.* 3:263-281) while the enterokinase cleavage site provides a means for purifying from the fusion protein. A discussion of vectors which contain fusion proteins is provided in Kroll, D. J. et al. 993; *DNA Cell Biol.* 12:441-453).

[0116] In addition to recombinant production, fragments of IMRRP1 or IMRRP1b may be produced by direct peptide synthesis using solid-phase techniques (Merrifield J. (1963) *J. Am. Chem. Soc.* 85:2149-2154). Protein synthesis may be performed using manual techniques or by automation. Automated synthesis may be achieved, for example, using Applied Biosystems 431A Peptide Synthesizer (Perkin Elmer). Various fragments of IMRRP1 or IMRRP1b can be chemically synthesized separately and combined using chemical methods to produce the full length molecule.

[0117] Chemical and structural homology exists among IMRRP1 or IMRRP1b and the human imidazoline receptor disclosed in *DNA Cell Biol.* 19 (6), 319-329 (2000). Furthermore, IMRRP1 and IMRRP1b are expressed in brain, bone marrow, heart, kidney, liver, lung, lymph node, pla-

centa, small intestine, spinal cord, spleen testis, and thymus tissues, many of which are associated with the regulation of blood pressure, induction of feeding, stimulation of firing of locus coeruleus neurons, and stimulation of insulin release, as well as the induction of the expression of glial fibrillary acidic protein independent of the action of alpha-2 adrenoceptors, dysphoric premenstrual syndrome, neurodegenerative disorders such as Alzheimer's disease, opiate addiction, monoamine turnover and therefore nociception, ageing, mood and stroke, salivary disorders and developmental disorders. IMRRP1 and IMRRP1b therefore play an important role in mammalian physiology.

[0118] In another embodiment a vector capable of expressing IMRRP1 or IMRRP1b, or a fragment or derivative thereof, may also be administered to a subject to treat or prevent a physical or psychological disorder, including those listed above.

[0119] In another embodiment, agonists or antagonists of IMRRP1 or IMRRP1b may be administered to a subject to treat or prevent a disorder associated with many neurological conditions and disorders including depression. In one aspect, antibodies which are specific for IMRRP1 or IMRRP1b may be used directly as an antagonist, or indirectly as a targeting or delivery mechanism for bringing a pharmaceutical agent to cells or tissue which express IMRRP1 or IMRRP1b.

[0120] In another embodiment, a vector expressing the complementary or antisense sequence of the polynucleotide encoding IMRRP1 or IMRRP1b may be administered to a subject to treat or prevent a disorder associated many neurological conditions and disorders including depression.

[0121] In another embodiment a vector expressing the complementary or antisense sequence of the polynucleotide encoding IMRRP1 or IMRRP1b may be administered to a subject to treat or many neurological conditions and disorders including depression associated with expression of IMRRP1 or IMRRP1b.

[0122] In other embodiments, any of the therapeutic proteins, antagonists, antibodies, agonists, antisense sequences or vectors described above may be administered in combination with other appropriate therapeutic agents. Selection of the appropriate agents for use in combination therapy may be made by one of ordinary skill in the art, according to conventional pharmaceutical principles. The combination of therapeutic agents may act synergistically to effect the treatment or prevention of the various disorders described above. Using this approach, one may be able to achieve therapeutic efficacy with lower dosages of each agent, thus reducing the potential for adverse side effects.

[0123] Agonists and antagonists or inhibitors of IMRRP1 or IMRRP1b may be produced using methods which are generally known in the art. For example, cloned receptors may be expressed in mammalian cells and compounds can be screened for activity. In addition, purified IMRRP1 or IMRRP1b may be used to produce antibodies or to screen libraries of pharmaceutical agents to identify those which specifically bind IMRRP1 or IMRRP1b.

[0124] Antibodies specific for IMRRP1 or IMRRP1b may be generated using methods that are well known in the art. Such antibodies may include, but are not limited to, polyclonal, monoclonal, chimeric, single chain, Fab fragments,

and fragments produced by a Fab expression library. Neutralizing antibodies, (i.e., those which inhibit dimer formation) are especially preferred for therapeutic use.

[0125] For the production of antibodies, various hosts including goats, rabbits, rats, mice, humans, and others, may be immunized by injection with or any fragment or oligopeptide of IMRRP1 or IMRRP1b which has immunogenic properties. Depending on the host species, various adjuvants may be used to increase immunological response. Such adjuvants include, but are not limited to, Ribi adjuvant R700 (Ribi, Hamilton, Mont.), incomplete Freund's adjuvant, mineral gels such as aluminum hydroxide, and surface active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, keyhole limpet hemocyanin, and dinitrophenol. Among adjuvants used in humans, BCG (bacillus Calmette Guérin) and *Corynebacterium parvum* are especially preferable.

[0126] It is preferred that the peptides, fragments, or oligopeptides used to induce antibodies to IMRRP1 or IMRRP1b have an amino acid sequence consisting of at least five amino acids, and more preferably at least 10 amino acids. It is also preferable that they are identical to a portion of the amino acid sequence of the natural protein, and they may contain the entire amino acid sequence of a small, naturally occurring molecule. The peptides, fragments or oligopeptides may comprise a single epitope or antigenic determinant or multiple epitopes. Short stretches of IMRRP1 or IMRRP1 amino acids may be fused with those of another protein such as keyhole limpet hemocyanin and antibody produced against the chimeric molecule.

[0127] Monoclonal antibodies to IMRRP1 or IMRRP1b may be prepared using any technique which provides for the production of antibody molecules by continuous cell lines in culture. These include, but are not limited to, the hybridoma technique, the human B-cell hybridoma technique, and the EBV-hybridoma technique (Kohler, G. et al. (1975) *Nature* 256:495-497; Kozbor, D. et al. (1985) *J. Immunol. Methods* 81:31-42, Cote, R. J. et al. (1983) *Proc. Natl. Acad. Sci.* 80:2026-2030; Cole, S. P. et al. (1984) *Mol. Cell Biol.* 62:109-120).

[0128] In addition, techniques developed for the production of "chimeric antibodies," the splicing of mouse antibody genes to human antibody genes to obtain a molecule with appropriate antigen specificity and biological activity can be used (Morrison, S. L. et al. (1984) *Proc. Natl. Acad. Sci.* 81:6851-6855; Neuberger, M. S. et al. (1984) *Nature* 312:604-608; Takeda, S. et al. (1985) *Nature* 314:452-454). Alternatively, techniques described for the production of single chain antibodies may be adapted, using methods known in the art, to produce IMRRP1- or IMRRP1b-specific single chain antibodies. Antibodies with related specificity, but of distinct idiotypic composition, may be generated by chain shuffling from random combinatorial immunoglobulin libraries (Burton D. R. (1991) *Proc. Natl. Acad. Sci.* 88:1120-3).

[0129] Antibodies may also be produced by inducing in vivo production in the lymphocyte population or by screening recombinant immunoglobulin libraries or panels of highly specific binding reagents as disclosed in the literature (Orlandi, R. et al. (1989) *Proc. Natl. Acad. Sci.* 86:3833-3837; Winter, G. et al. (1991) *Nature* 349:293-299).

[0130] Antibody fragments which contain specific binding sites for IMRRP1 or IMRRP1b may also be generated. For

example, such fragments include, but are not limited to, the F(ab')₂ fragments which can be produced by pepsin digestion of the antibody molecule and the Fab fragments which can be generated by reducing the disulfide bridges of the F(ab')₂ fragments. Alternatively, Fab expression libraries may be constructed to allow rapid and easy identification of monoclonal Fab fragments with the desired specificity (Huse, W. D. et al. (1989) *Science* 254:1275-1281).

[0131] Various immunoassays may be used for screening to identify antibodies having the desired specificity. Numerous protocols for competitive binding or immunoradiometric assays using either polyclonal or monoclonal antibodies with established specificities are well known in the art. Such immunoassays typically involve the measurement of complex formation between IMRRP1 or IMRRP1b and their specific antibody. A two-site, monoclonal-based immunoassay utilizing monoclonal antibodies reactive to two non-interfering IMRRP1 or IMRRP1b epitopes is preferred, but a competitive binding assay may also be employed (Maddox, supra).

[0132] In another embodiment of the invention, the polynucleotides encoding IMRRP1 or IMRRP1b or any fragment thereof or antisense molecules, may be used for therapeutic purposes. In one aspect, antisense to the polynucleotide encoding IMRRP1 or IMRRP1b may be used in situations in which it would be desirable to block the transcription of the mRNA. In particular, cells may be transformed with sequences complementary to polynucleotides encoding IMRRP1 or IMRRP1b. Thus, antisense molecules may be used to modulate IMRRP1 or IMRRP1b activity, or to achieve regulation of gene function. Such technology is well known in the art, and sense or antisense oligomers or larger fragments, can be designed from various locations along the coding or control regions of sequences encoding IMRRP1 or IMRRP1b.

[0133] Expression vectors derived from retroviruses, adenovirus, herpes or vaccinia viruses, or from various bacterial plasmids may be used for delivery of nucleotide sequences to the targeted organ, tissue or cell population. Methods which are well known to those skilled in the art can be used to construct recombinant vectors which will express antisense molecules complementary to the polynucleotides of the genes encoding IMRRP1 or IMRRP1b. These techniques are described both in Sambrook et al. (supra) and in Ausubel et al. (supra).

[0134] Genes encoding IMRRP1 or IMRRP1b can be turned off by transforming a cell or tissue with expression vectors which express high levels of a polynucleotide or fragment thereof which encodes IMRRP1 or IMRRP1b. Such constructs may be used to introduce untranslatable sense or antisense sequences into a cell. Even in the absence of integration into the DNA, such vectors may continue to transcribe RNA molecules until they are disabled by endogenous nucleases. Transient expression may last for a month or more with a non-replicating vector and even longer if appropriate replication elements are part of the vector system.

[0135] As mentioned above, modifications of gene expression can be obtained by designing antisense molecules, DNA, RNA, or PNA, to the control regions of the genes encoding IMRRP1 or IMRRP1b, i.e., the promoters, enhancers, and introns. Oligonucleotides derived from the

transcription initiation site, e.g., between positions -10 and +10 from the start site, are preferred. Similarly, inhibition can be achieved using "triple helix" base-pairing methodology. Triple helix pairing is useful because it causes inhibition of the ability of the double helix to open sufficiently for the binding of polymerases, transcription factors, or regulatory molecules. Recent therapeutic advances using triplex DNA have been described in the literature (Gee, J. E. et al. (1994) In: Huber, B. E. and B. L. Carr, *Molecular and Immunologic Approaches*, Futura Publishing Co., Mt. Kisco, N.Y). The antisense molecules may also be designed to block translation of mRNA by preventing the transcript from binding to ribosomes.

[0136] Ribozymes, enzymatic RNA molecules, may also be used to catalyze the specific cleavage of RNA. The mechanism of ribozyme action involves sequence-specific hybridization of the ribozyme molecule to complementary target RNA, followed by endonucleolytic cleavage. Examples which may be used include engineered hammerhead motif ribozyme molecules that can specifically and efficiently catalyze endonucleolytic cleavage of sequences encoding IMRRP1 or IMRRP1b.

[0137] Specific ribozyme cleavage sites within any potential RNA target are initially identified by scanning the target molecule for ribozyme cleavage sites which include the following sequences: GUA, GUU, and GUC. Once identified, short RNA sequences of between 15 and 20 ribonucleotides corresponding to the region of the target gene containing the cleavage site may be evaluated for secondary structural features which may render the oligonucleotide inoperable. The suitability of candidate targets may also be evaluated by testing accessibility to hybridization with complementary oligonucleotides using ribonuclease protection assays.

[0138] Antisense molecules and ribozymes of the invention may be prepared by any method known in the art for the synthesis of nucleic acid molecules. These include techniques for chemically synthesizing oligonucleotides such as solid phase phosphoramidite chemical synthesis. Alternatively, RNA molecules may be generated by in vitro and in vivo transcription of DNA sequences encoding IMRRP1 or IMRRP1b. Such DNA sequences may be incorporated into a wide variety of vectors with suitable RNA polymerase promoters such as T7 or SP6. Alternatively, these cDNA constructs that synthesize antisense RNA constitutively or inducibly can be introduced into cell lines, cells, or tissues.

[0139] RNA molecules may be modified to increase intracellular stability and half-life. Possible modifications include, but are not limited to, the addition of flanking sequences at the 5' and/or 3' ends of the molecule or the use of phosphorothioate or 2' O-methyl rather than by phosphodiesterase linkages within the backbone of the molecule. This concept is inherent in the production of PNAs and can be extended in all of these molecules by the inclusion of nontraditional bases such as inosine, queosine, and wybutosine, as well as acetyl-, methyl-, thio-, and similarly modified forms of adenine, cytidine, guanine, thymine, and uridine which are not as easily recognized by endogenous endonucleases.

[0140] Many methods for introducing vectors into cells or tissues are available and equally suitable for use in vivo, in vitro, and ex vivo. For ex vivo therapy, vectors may be

introduced into stem cells taken from the patient and clonally propagated for autologous transplant back into that same patient as disclosed in U.S. Pat. No. 5,399,493 and 5,437,994. Delivery by transfection and by liposome injections may be achieved using methods which are well known in the art.

[0141] Any of the therapeutic methods described above may be applied to any subject in need of such therapy, including, for example, mammals such as dogs, cats, cows, horses, rabbits, monkeys, and most preferably, humans.

[0142] An additional embodiment of the invention relates to the administration of a pharmaceutical composition, in conjunction with a pharmaceutically acceptable carrier, for any of the therapeutic effects discussed above. Such pharmaceutical compositions may consist of IMRRP1 or IMRRP1b, antibodies to IMRRP1 or IMRRP1b, mimetics, agonists, antagonists, or inhibitors of IMRRP1 or IMRRP1b. The compositions may be administered alone or in combination with at least one other agent, such as stabilizing compound, which may be administered in any sterile, biocompatible pharmaceutical carrier, including, but not limited to, saline, buffered saline, dextrose, and water. The compositions may be administered to a patient alone, or in combination with other agents, drugs, hormones, or biological response modifiers.

[0143] The pharmaceutical compositions utilized in this invention may be administered by any number of routes including, but not limited to, oral, intravenous, intramuscular, intraarterial, intramedullary, intrathecal, intraventricular, transdermal, subcutaneous, intraperitoneal, intranasal, enteral, topical, sublingual, or rectal means.

[0144] In addition to the active ingredients, these pharmaceutical compositions may contain suitable pharmaceutically acceptable carriers comprising excipients and auxiliaries which facilitate processing of the active compounds into preparations which can be used pharmaceutically. Further details on techniques for formulation and administration may be found in the latest edition of *Remington's Pharmaceutical Sciences* (Mack Publishing Co., Easton, Pa.).

[0145] Pharmaceutical compositions for oral administration can be formulated using pharmaceutically acceptable carriers well known in the art in dosages suitable for oral administration. Such carriers enable the pharmaceutical compositions to be formulated as tablets, pills, dragees, capsules, liquids, gels, syrups, slurries, suspensions, and the like, for ingestion by the patient.

[0146] Pharmaceutical preparations for oral use can be obtained through combination of active compounds with solid excipient, optionally grinding a resulting mixture, and processing the mixture of granules, after adding suitable auxiliaries, if desired, to obtain tablets or dragee cores. Suitable excipients are carbohydrate or protein fillers, such as sugars, including lactose, sucrose, mannitol, or sorbitol; starch from corn, wheat, rice, potato, or other plants; cellulose, such as methyl cellulose, hydroxypropylmethylcellulose, or sodium carboxymethylcellulose; gums including arabic and tragacanth, and proteins such as gelatin and collagen. If desired, disintegrating or solubilizing agents may be added, such as the cross-linked polyvinyl pyrrolidone, agar, alginic acid, or a salt thereof, such as sodium alginate.

[0147] Dragee cores may be used in conjunction with suitable coatings, such as concentrated sugar solutions, which may also contain gum arabic, talc, polyvinylpyrrolidone, carbopol gel, polyethylene glycol, and/or titanium dioxide, lacquer solutions, and suitable organic solvents or solvent mixtures. Dye-stuffs or pigments may be added to the tablets or dragee coatings for product identification or to characterize the quantity of active compound, i.e., dosage.

[0148] Pharmaceutical preparations which can be used orally include push-fit capsules made of gelatin, as well as soft, sealed capsules made of gelatin and a coating, such as glycerol or sorbitol. Push-fit capsules can contain active ingredients mixed with a filler or binders, such as lactose or starches, lubricants, such as talc or magnesium stearate, and, optionally, stabilizers. In soft capsules, the active compounds may be dissolved or suspended in suitable liquids, such as fatty oils, liquid, or liquid polyethylene glycol with or without stabilizers.

[0149] Pharmaceutical formulations suitable for parenteral administration may be formulated in aqueous solutions, preferably in physiologically compatible buffers such as Hanks' solution, Ringer's solution, or physiologically buffered saline. Aqueous injection suspensions may contain substances which increase the viscosity of the suspension, such as sodium carboxymethyl cellulose, sorbitol, or dextran. Additionally, suspensions of the active compounds may be prepared as appropriate oily injection suspensions. Suitable lipophilic solvents or vehicles include fatty oils such as sesame oil, or synthetic fatty acid esters, such as ethyl oleate or triglycerides, or liposomes. Optionally, the suspension may also contain suitable stabilizers or agents which increase the solubility of the compounds to allow for the preparation of highly concentrated solutions.

[0150] For topical or nasal administration, penetrants appropriate to the particular barrier to be permeated are used in the formulation. Such penetrants are generally known in the art.

[0151] The pharmaceutical compositions of the present invention may be manufactured in a manner that is known in the art, e.g., by means of conventional mixing, dissolving, granulating, dragee-making, levigating, emulsifying, encapsulating, entrapping, or lyophilizing processes.

[0152] The pharmaceutical composition may be provided as a salt and can be formed with many acids, including but not limited to, hydrochloric, sulfuric, acetic, lactic, tartaric, malic, succinic, etc. Salts tend to be more soluble in aqueous or other protonic solvents than are the corresponding free base forms. In other cases, the preferred preparation may be a lyophilized powder which may contain any or all of the following: 1-50 m-M histidine, 0.1%-2% sucrose, and 2-7% mannitol, at a pH range of 4.5 to 5.5, that is combined with buffer prior to use.

[0153] After pharmaceutical compositions have been prepared, they can be placed in an appropriate container and labeled for treatment of an indicated condition. For administration of IMRRP1 or IMRRP1b, such labeling would include amount, frequency, and method of administration.

[0154] Pharmaceutical compositions suitable for use in the invention include compositions wherein the active ingredients are contained in an effective amount to achieve the

intended purpose. The determination of an effective dose is well within the capability of those skilled in the art.

[0155] For any compound, the therapeutically effective dose can be estimated initially either in cell culture assays, e.g., of neoplastic cells, or in animal models, usually mice, rabbits, dogs, or pigs. The animal model may also be used to determine the appropriate concentration range and route of administration. Such information can then be used to determine useful doses and routes for administration in humans.

[0156] A therapeutically effective dose refers to that amount of active ingredient, for example IMRRP1 or IMRRP1b or fragments thereof antibodies of IMRRP1 or IMRRP1b, agonists, antagonists or inhibitors of IMRRP1 or IMRRP1b which ameliorates the symptoms or condition. Therapeutic efficacy and toxicity may be determined by standard pharmaceutical procedures in cell cultures or experimental animals, e.g., ED₅₀ (the dose therapeutically effective in 50% of the population) and LD₅₀ (the dose lethal to 50% of the population). The dose ratio of toxic to therapeutic effects is the therapeutic index, and it can be expressed as the ratio, LD₅₀/ED₅₀. Pharmaceutical compositions which exhibit large therapeutic indices are preferred. The data obtained from cell culture assays and animal studies is used in formulating a range of dosage for human use. The dosage contained in such compositions is preferably within a range of circulating concentrations that include the ED₅₀ with little or no toxicity. The dosage varies within this range depending upon the dosage form employed, sensitivity of the patient, and the route of administration.

[0157] The exact dosage will be determined by the practitioner, in light of factors related to the subject that requires treatment. Dosage and administration are adjusted to provide sufficient levels of the active moiety or to maintain the desired effect. Factors which may be taken into account include the severity of the disease state, general health of the subject, age, weight, and gender of the subject, diet, time and frequency of administration, drug combination(s), reaction sensitivities, and tolerance/response to therapy. Long-acting pharmaceutical compositions may be administered every 3 to 4 days, every week, or once every two weeks depending on half-life and clearance rate of the particular formulation.

[0158] Normal dosage amounts may vary from 0.1 to 100,000 microgram, up to a total dose of about 1 g, depending upon the route of administration. Guidance as to particular dosages and methods of delivery is provided in the literature and generally available to practitioners in the art. In one embodiment, dosages of IMRRP1 or IMRRP1b or fragment thereof from about 1 ng/kg/day to about 10 mg/kg/day, and preferably from about 500 ug/kg/day to about 5 mg/kg/day are expected to induce a biological effect. Those skilled in the art will employ different formulations for nucleotides than for proteins or their inhibitors. Similarly, delivery of polynucleotides or polypeptides will be specific to particular cells, conditions, locations, etc.

[0159] In another embodiment, antibodies which specifically bind IMRRP1 or IMRRP1b may be used for the diagnosis of conditions or diseases characterized by expression of IMRRP1 or IMRRP1b, or in assays to monitor patients being treated with IMRRP1 or IMRRP1b, agonists, antagonists or inhibitors. The antibodies useful for diagnostic purposes may be prepared in the same manner as those

described above for therapeutics. Diagnostic assays for IMRRP1 or IMRRP1b include methods which utilize the antibody and a label to detect it in human body fluids or extracts of cells or tissues. The antibodies may be used with or without modification, and may be labeled by joining them, either covalently or non-covalently, with a reporter molecule. A wide variety of reporter molecules which are known in the art may be used, several of which are described above.

[0160] A variety of protocols including ELISA, RIA, and FACS for measuring IMRRP1 or IMRRP1b are known in the art and provide a basis for diagnosing altered or abnormal levels of IMRRP1 or IMRRP1b expression. Normal or standard values for IMRRP1 or IMRRP1b expression are established by combining body fluids or cell extracts taken from normal mammalian subjects, preferably human, with antibody to IMRRP1 or IMRRP1b under conditions suitable for complex formation. The amount of standard complex formation may be quantified by various methods, but preferably by photometric means. Quantities of IMRRP1 or IMRRP1b expressed in subject samples, control and disease from biopsied tissues are compared with the standard values. Deviation between standard and subject values establishes the parameters for diagnosing disease.

[0161] In another embodiment of the invention, the polynucleotides encoding IMRRP1 or IMRRP1b may be used for diagnostic purposes. The polynucleotides which may be used include oligonucleotide sequences, antisense RNA and DNA molecules, and PNAs. The polynucleotides may be used to detect and quantitate gene expression in biopsied tissues in which expression of IMRRP1 or IMRRP1b may be correlated with disease. The diagnostic assay may be used to distinguish between absence, presence, and excess expression of IMRRP1 or IMRRP1b, and to monitor regulation of IMRRP1 or IMRRP1b levels during therapeutic intervention.

[0162] In one aspect, hybridization with PCR probes which are capable of detecting polynucleotide sequences, including genomic sequences, encoding IMRRP1 or IMRRP1b or closely related molecules, may be used to identify nucleic acid sequences which encode IMRRP1 or IMRRP1b. The specificity of the probe, whether it is made from a highly specific region, e.g., unique nucleotides in the 5' regulatory region, or a less specific region, e.g., especially in the 3' coding region, and the stringency of the hybridization or amplification (maximal, high, intermediate, or low) will determine whether the probe identifies only naturally occurring sequences encoding IMRRP1 or IMRRP1b, alleles, or related sequences.

[0163] Probes may also be used for the detection of related sequences, and should preferably contain at least 50% of the nucleotides from any of the IMRRP1 or IMRRP1b encoding sequences. The hybridization probes of the subject invention may be DNA or RNA and derived from the nucleotide sequence of SEQ ID NOS: 1 or 2 or from genomic sequence including promoter, enhancer elements, and introns of the naturally occurring IMRRP1 or IMRRP1b genes.

[0164] Means for producing specific hybridization probes for DNAs encoding IMRRP1 or IMRRP1b include the cloning of nucleic acid sequences encoding IMRRP1 or IMRRP1b or derivatives into vectors for the production of mRNA probes. Such vectors are known in the art, commer-

cially available, and may be used to synthesize RNA probes *in vitro* by means of the addition of the appropriate RNA polymerases and the appropriate labeled nucleotides. Hybridization probes may be labeled by a variety of reporter groups, for example, radionuclides such as ^{32}P or ^{35}S , or enzymatic labels, such as alkaline phosphatase coupled to the probe via avidin/biotin coupling systems, and the like.

[0165] Polynucleotide sequences encoding IMRRP1 or IMRRP1b may be used for the diagnosis of disorders associated with expression of IMRRP1 and IMRRP1b. Examples of such disorders or conditions include regulation of blood pressure, hypertension, induction of feeding, stimulation of firing of locus coeruleus neurons, and stimulation of insulin release, as well as the aberrant induction of the expression of glial fibrillary acidic protein independent of the action of alpha-2 adrenoceptors, dysphoric premenstrual syndrome, neurodegenerative disorders such as Alzheimer's disease, opiate addiction, monoamine turnover and therefore nociception, ageing, mood and stroke, salivary disorders and developmental disorders. The polynucleotide sequences encoding IMRRP1 or IMRRP1b may be used in Southern or northern analysis, dot blot, or other membrane-based technologies; in PCR technologies; or in dip stick, pin, ELISA or chip assays utilizing fluids or tissues from patient biopsies to detect altered IMRRP1 or IMRRP1b expression. Such qualitative or quantitative methods are well known in the art.

[0166] The nucleotide sequences encoding IMRRP1 or IMRRP1b may be labeled by standard methods, and added to a fluid or tissue sample from a patient under conditions suitable for the formation of hybridization complexes. After a suitable incubation period, the sample is washed and the signal is quantitated and compared with a standard value. If the amount of signal in the biopsied or extracted sample is significantly altered from that of a comparable control sample, the nucleotide sequences have hybridized with nucleotide sequences in the sample, and the presence of altered levels of nucleotide sequences encoding IMRRP1 or IMRRP1b in the sample indicates the presence of the associated disease. Such assays may also be used to evaluate the efficacy of a particular therapeutic treatment regimen in animal studies, in clinical trials, or in monitoring the treatment of an individual patient.

[0167] In order to provide a basis for the diagnosis of disease associated with expression of IMRRP1 or IMRRP1b, a normal or standard profile for expression is established. This may be accomplished by combining body fluids or cell extracts taken from normal subjects, either animal or human, with a sequence, or a fragment thereof, which encodes IMRRP1 or IMRRP1b, under conditions suitable for hybridization or amplification. Standard hybridization may be quantified by comparing the values obtained from normal subjects with those from an experiment where a known amount of a substantially purified polynucleotide is used. Standard values obtained from normal samples may be compared with values obtained from samples from patients who are symptomatic for disease. Deviation between standard and subject values is used to establish the presence of disease.

[0168] Once disease is established and a treatment protocol is initiated, hybridization assays may be repeated on a regular basis to evaluate whether the level of expression in the patient begins to approximate that which is observed in

the normal patient. The results obtained from successive assays may be used to show the efficacy of treatment over a period ranging from several days to months.

[0169] Additional diagnostic uses for oligonucleotides designed from the sequences encoding IMRRP1 or IMRRP1b may involve the use of PCR. Such oligomers may be chemically synthesized, generated enzymatically, or produced from a recombinant source. Oligomers will preferably consist of two nucleotide sequences, one with sense orientation ($5' \geq 3'$) and another with antisense ($3' \geq 5'$), employed under optimized conditions for identification of a specific gene or condition. The same two oligomers, nested sets of oligomers, or even a degenerate pool of oligomers may be employed under less stringent conditions for detection and/or quantitation of closely related DNA or RNA sequences.

[0170] Methods which may also be used to quantitate the expression of IMRRP1 or IMRRP1b include radiolabeling or biotinylating nucleotides, coamplification of a control nucleic acid, and standard curves onto which the experimental results are interpolated (Melby, P. C. et al. (1993) *J. Immunol. Methods*, 159:235-244; Duplaa, C. et al. (1993) *Anal. Biochem.* 229-236). The speed of quantitation of multiple samples may be accelerated by running the assay in an ELISA format where the oligomer of interest is presented in various dilutions and a spectrophotometric or calorimetric response gives rapid quantitation.

[0171] In another embodiment of the invention, the nucleic acid sequences which encode IMRRP1 or IMRRP1b may also be used to generate hybridization probes which are useful for mapping the naturally occurring genomic sequence. The sequences may be mapped to a particular chromosome or to a specific region of the chromosome using well known techniques. Such techniques include FISH, FACS, or artificial chromosome constructions, such as yeast artificial chromosomes, bacterial artificial chromosomes, bacterial PI constructions or single chromosome cDNA libraries as reviewed in Price, C. M. (1993) *Blood Rev.* 7:127-134, and Trask, B. J. (1991) *Trends Genet.* 7:149-154.

[0172] FISH (as described in Verma et al. (1988) *Human Chromosomes: A Manual of Basic Techniques* Pergamon Press, New York, N.Y.) may be correlated with other physical chromosome mapping techniques and genetic map data. Examples of genetic map data can be found in the 1994 Genome Issue of Science (265:1981f). Correlation between the location of the gene encoding IMRRP1 or IMRRP1b on a physical chromosomal map and a specific disease, or predisposition to a specific disease, may help delimit the region of DNA associated with that genetic disease. The nucleotide sequences of the subject invention may be used to detect differences in gene sequences between normal, carrier, or affected individuals.

[0173] In situ hybridization of chromosomal preparations and physical mapping techniques such as linkage analysis using established chromosomal markers may be used for extending genetic maps. Often the placement of a gene on the chromosome of another mammalian species, such as mouse, may reveal associated markers even if the number or arm of a particular human chromosome is not known. New sequences can be assigned to chromosomal arms, or parts thereof, by physical mapping. This provides valuable information to investigators searching for disease genes using

positional cloning or other gene discovery techniques. Once the disease or syndrome has been crudely localized by genetic linkage to a particular genomic region, for example, AT to 11q22-23 (Gatti, R. A. et al. (1988) *Nature* 336:577-580), any sequences mapping to that area may represent associated or regulatory genes for further investigation. The nucleotide sequence of the subject invention may also be used to detect differences in the chromosomal location due to translocation, inversion, etc. among normal, carrier, or affected individuals.

[0174] In another embodiment of the invention, IMRRP1 or IMRRP1b, their catalytic or immunogenic fragments or oligopeptides thereof can be used for screening libraries of compounds in any of a variety of drug screening techniques. The fragment employed in such screening may be free in solution, affixed to a solid support, borne on a cell surface, or located intracellularly. The formation of binding complexes, between IMRRP1 or IMRRP1b and the agent being tested, may be measured.

[0175] Another technique for drug screening which may be used provides for high throughput screening of compounds having suitable binding affinity to the protein of interest as described in published PCT application WO84/03564. In this method, as applied to IMRRP1 or IMRRP1b, large numbers of different small test compounds are synthesized on a solid substrate, such as plastic pins or some other surface. The test compounds are contacted with IMRRP1 or IMRRP1b or fragments thereof, and washed. Bound IMRRP1 or IMRRP1b are then detected by methods well known in the art. Purified IMRRP1 or IMRRP1b can also be coated directly onto plates for use in the aforementioned drug screening techniques. Alternatively, non-neutralizing antibodies can be used to capture the peptide and immobilize it on a solid support.

[0176] In another embodiment, one may use competitive drug screening assays in which neutralizing antibodies capable of binding IMRRP1 or IMRRP1 specifically compete with a test compound for binding IMRRP1 or IMRRP1b. In this manner, the antibodies can be used to detect the presence of any peptide which shares one or more antigenic determinants with IMRRP1 or IMRRP1b.

[0177] In additional embodiments, the nucleotide sequences which encode IMRRP1 or IMRRP1b may be used in any molecular biology techniques that have yet to be developed, provided the new techniques rely on properties of nucleotide sequences that are currently known, including, but not limited to, such properties as the triplet genetic code and specific base pair interactions.

[0178] The examples below are provided to illustrate the subject invention and are not included for the purpose of limiting the invention. All publications and patents mentioned in the specification are herein incorporated by reference. Various modifications and variations of the described method and system of the invention will be apparent to those skilled in the art without departing from the scope and spirit of the invention. Although the invention has been described in connection with specific preferred embodiments, it should be understood that the invention as claimed should not be unduly limited to such specific embodiments. Indeed, various modifications of the described modes for carrying out the invention which are obvious to those skilled in molecular biology or related fields are intended to be within the scope of the following claims.

EXAMPLES

Example I

[0179] Method of Isolation of cDNA Encoding IMRRP1 or IMRRP1b

[0180] Human imidazoline receptor protein sequence was used as a probe to search the Incyte and public domain EST databases. The search program used was gapped BLAST (Altschul et al., 1997). The top EST hits from the BLAST results were searched back against the non-redundant protein and patent sequence databases. From this analysis, ESTs encoding a potential novel imidazoline receptor was identified based on sequence homology. The Incyte EST (CloneID: 2499870) was selected as a potential novel imidazoline receptor candidate for subsequent analysis.

[0181] A PCR primer pair, designed from the DNA sequence of Incyte clone-2499870 was used to amplify a piece of DNA from the clone in which the anti-sense strand of the amplified fragment was biotinylated on the 5' end. This biotinylated piece of double stranded DNA was denatured and incubated with a mixture of single-stranded covalently closed circular cDNA libraries which contain DNA corresponding to the sense strand. The cDNA libraries were total brain tissue libraries obtained from Gibco Life Technologies. Hybrids between the biotinylated DNA and the circular cDNA were captured on streptavidin magnetic beads. Upon thermal release of the cDNA from the biotinylated DNA, the single stranded cDNA was converted into double strands using a primer homologous to a sequence on the cDNA cloning vector. The double stranded cDNA was introduced into *E. coli* by electroporation and the resulting colonies were screen by PCR, using the original primer pair, to identify the proper cDNA clones. One clone named FL1-18 was sequenced on both strands (FIG. 1).

Example II

[0182] Cellular and Tissue Distribution of IMRRP 1

[0183] The same PCR primer used in the cloning of imidazoline receptor IMRRP1 used to measure the steady state levels of mRNA by quantitative PCR. Briefly, first strand cDNA was made from commercially available mRNA. The relative amount of cDNA used in each assay was determined by performing a parallel experiment using a primer pair for a gene expressed in equal amounts in all tissues, cyclophilin. The cyclophilin primer pair detected small variations in the amount of cDNA in each sample and these data were used for normalization of the data obtained with the primer pair for IMRRP1. The PCR data was converted into a relative assessment of the difference in transcript abundance amongst the tissues tested and the data is presented in FIG. 9.

Example III

[0184] Labeling and Use of Hybridization Probes Hybridization probes derived from SEQ ID NOS: 1 or 2 are employed to screen cDNAs, genomic DNAs, or mRNAs. Although the labeling of oligonucleotides, consisting of about 20 base-pairs, is specifically described, essentially the same procedure is used with larger cDNA fragments. Oligonucleotides are designed using state-of-the-art software such as OLIGO 4.06 (National Biosciences), labeled by

combining 50 pmol of each oligomer and 250 μ Ci of [γ - 32 P] adenosine triphosphate (Amersham) and T4 polynucleotide kinase (DuPont NEN, Boston, Mass.). The labeled oligonucleotides are substantially purified with SEPHADEX G-25 superfine resin column (Pharmacia & Upjohn). A portion containing about 10^7 counts per minute of each of the sense and antisense oligonucleotides is used in a typical membrane based hybridization analysis of human genomic DNA digested with one of the following endonucleases (Ase I, Bgl II, Eco RI, Pst I, Xba I, or Pvu II: DuPont NEN).

[0185] The DNA from each digest is fractionated on a 0.7 percent agarose gel and transferred to nylon membranes (Nytran Plus, Schleicher & Schuell, Durham, N. H.). Hybridization is carried out for 16 hours at 40° C. To remove nonspecific signals, blots are sequentially washed at room temperature under increasingly stringent conditions up to 0.1x saline sodium citrate and 0.5% sodium dodecyl sulfate. After XOMATAR film (Kodak, Rochester, N.Y.) is exposed to the blots in a Phosphoimager cassette (Molecular Dynamics, Sunnyvale, Calif.) for several hours, hybridization patterns are compared visually.

Example IV

[0186] Antisense Molecules

[0187] Antisense molecules or nucleic acid sequence complementary to the IMRRP1 or IMRRP1b encoding sequences, or any part thereof, is used to inhibit in vivo or in vitro expression of naturally occurring IMRRP1 or IMRRP1b. Although use of antisense oligonucleotides, comprising about 20 base-pairs, is specifically described, essentially the same procedure is used with larger cDNA fragments. An oligonucleotide based on the coding sequences of IL-17R, as shown in FIGS. 1 and 2 is used to inhibit expression of naturally occurring IMRRP1 or IMRRP1b. The complementary oligonucleotide is designed from the unique 5' sequence as shown in FIGS. 1 or 2 and used either to inhibit transcription by preventing promoter binding to the upstream nontranslated sequence or translation of an IMRRP1 or IMRRP1b encoding transcript by preventing the ribosome from binding. Using an appropriate portion of the signal and 5' sequence of SEQ ID NOS: 1 or 2 an effective antisense oligonucleotide includes any 15-20 nucleotides spanning the region which translates into the signal or 5' coding sequence of the polypeptide as shown in FIGS. 1 and 2.

Example V

[0188] Production of IMRRP or IMRRP1b Specific Antibodies

[0189] IMRRP1 or IMRRP1b that is substantially purified using PAGE electrophoresis (Sambrook, supra), or other purification techniques, is used to immunize rabbits and to produce antibodies using standard protocols. The amino acid sequence from SEQ ID NOS: 3 or 4 is analyzed using DNASTAR software (DNASTAR Inc.) to determine regions of high immunogenicity and a corresponding oligopolypeptide is synthesized and used to raise antibodies by means known to those of skill in the art. Selection of appropriate epitopes, such as those near the C-terminus or in hydrophilic regions, is described by Ausubel et al. (supra) and others.

[0190] Typically, the oligopeptides are 15 residues in length, synthesized using an Applied Biosystems Peptide

Synthesizer Model 431A using fmoc-chemistry, and coupled to keyhole limpet hemacyanin (KLH, Sigma, St. Louis, Mo.) by reaction with N-maleimidobenzoyl-N-hydroxysuccinimide ester (MBS; Ausubel et al., supra). Rabbits are immunized with the oligopeptide-KLH complex in complete Freund's adjuvant. The resulting antisera are tested for antipeptide activity, for example, by binding the rabbit antisera, washing, and reacting with radioiodinated, goat and anti-rabbit IgG.

Example VI

[0191] Purification of Naturally Occurring IMRRP1 or IMRRP1b Using Specific Antibodies

[0192] Naturally occurring or recombinant IMRRP1 or IMRRP1b is substantially purified by immunoaffinity chromatography using antibodies specific for IMRRP1 or IMRRP1b. An immunoaffinity column is constructed by covalently coupling IMRRP1 or IMRRP1b specific antibody to an activated chromatographic resin, such as CNBr-activated SEPHAROSE (Pharmacia & Upjohn). After the coupling, the resin is blocked and washed according to the manufacturer's instructions.

[0193] Media containing IMRRP1 or IMRRP1b is passed over the immunoaffinity column, and the column is washed under conditions that allow the preferential absorbance of IMRRP1 or IMRRP1b (e.g., high ionic strength buffers in the presence of detergent). The column is eluted under conditions that disrupt antibody-IMRRP1 or IMRRP1b binding (e.g., buffer of pH 2-3 or a high concentration of a chaotrope, such as urea or thiocyanate ion), and IMRRP1 or IMRRP1b is collected.

Example VII

[0194] Identification of Molecules Which Interact with IMRRP 1 or IMRRP1b

[0195] IMRRP1 or IMRRP1b or biologically active fragments thereof are labeled with ^{125}I Bolton-Hunter reagent (Bolton et al. (1973) *Biochem. J.*, 133:529). Candidate molecules previously arrayed in the wells of a multi-well plate are incubated with the labeled IMRRP1 or IMRRP1b, washed and any wells with labeled IMRRP1 or IMRRP1b complex are assayed. Data obtained using different concentrations of IMRRP1 or IMRRP1b are used to calculate values for the number, affinity, and associate of IMRRP1 or IMRRP1b with the candidate molecules.

Example VIII

[0196] Expression Profiling of IMMRP1

[0197] Expression profiling in 12 tissue RNA samples was carried out to show the overall pattern of gene expression in the body. The same PCR primer pair, shown below as Incyte-2499870, that was used to identify IMMRP1 cDNA clones was used to measure the steady state levels of mRNA by quantitative PCR.

INCYTE-2499870-s
GCTGGAGACCCGTGATTTGCA (SEQ ID NO:5)

INCYTE-2499870-a
BTGGACTTGATTTGGCTTAGGTT (SEQ ID NO:6)

[0198] First strand cDNA was made from commercially available mRNA (Clontech, Stratagene, and Life Technologies) and subjected to real time quantitative PCR using a PE 5700 instrument (Applied Biosystems, Foster City, Calif.) which detects the amount of DNA amplified during each cycle by the fluorescent output of SYBR green, a DNA binding dye specific for double strands. The specificity of the primer pair for its target is verified by performing a thermal denaturation profile at the end of the run which gives an indication of the number of different DNA sequences present by determining melting T_m . In the case of the FGFR1ΔCP primer pair, only one DNA fragment was detected having a homogeneous melting point. Contributions of contaminating genomic DNA to the assessment of tissue abundance is controlled for by performing the PCR with first strand made with and without reverse transcriptase. In all cases, the contribution of material amplified in the no reverse transcriptase controls was negligible.

[0199] Small variations in the amount of cDNA used in each tube was determined by performing a parallel experiment using a primer pair for cyclophilin, a gene expressed in equal amounts in all tissues. These data were used to normalize the data obtained with the IMMRP1 primer pair. The PCR data was converted into a relative assessment of the difference in transcript abundance amongst the tissues tested and the data are presented in bar graph form in FIG. 10. Transcripts corresponding to IMMRP1 were found in all the additional RNAs tested with the highest amount present in the testis (like that of the first panel tested). Relatively high expression was also observed in the salivary gland and the fetal brain.

[0200] The quantitative PCT was performed by determining the number of reactions and amount of mix needed. All samples were run in triplicate, so each sample tube need 3.5 reactions worth of mix. This is determined by the following formula: $2 \times \# \text{ tissue samples} + 1 \text{ no template control} + 1 \text{ for pipetting error}$.

[0201] The reaction mixture was prepared as follows.

Components	vol/rxn
2 × SybrGreen Master Mix	25 microliters
water	23.5 microliters
primer mix (10 μM ea.)	0.5 microliters
cDNA (2.5 ng/μL)	1 microliter

[0202] An aliquot 171.5 μL of mix was added to each to sample tubes followed by the addition of 1 μL of cDNA. Each sample tube was mixed gently and spun down. An aliquot of 3×50 μL was added to an optical plate and analyzed.

References

- [0203] Altschul, S. F., Madden, T. L., Schaffer, A. A., Zhang, J., Zhang, Z., Miller, W., and Lipman, D. L. (1997). Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acid Res.* 25, 3389-3402.
- [0204] Escriba, P. V., Ozaita, A., Miralles, A., Reis, D. J., and Garcia-Sevilla, J. A. (1995). Molecular charac-

terization and isolation of a 45-kilodalton imidazoline receptor protein from the rat brain. *Molec. Brain. Res.* 32, 187-196.

[0205] Farsang, C., and Kapocsi, J. (1999). Imidazoline receptors: from discovery to antihypertensive therapy (facts and doubts). *Brain Res. Bull.* 49, 317-331.

[0206] Garcia-Sevilla, J. A., Escriba, P. V., and Guimon, J. (1999). Imidazoline receptors and human brain disorders. *Ann. N. Y. Acad. Sci.* 21, 392-409.

What is claimed is:

1. A substantially purified IMRRP1 polypeptide consisting of amino acid sequence SEQ ID NO: 3.

2. A substantially purified IMRRP1b polypeptide consisting of amino acid sequence SEQ ID NO: 4.

3. A substantially purified IMRRP1 polypeptide consisting of amino acid sequence SEQ ID NO: 3, wherein the amino acid sequence differs from SEQ ID NO: 3 by conservative substitutions.

4. A substantially purified IMRRP1b polypeptide consisting of amino acid sequence SEQ ID NO: 4, wherein the amino acid sequence differs from SEQ ID NO: 4 by conservative substitutions.

5. An IMRRP1 or IMRRP1b polypeptide according to claims 1 or 2 wherein the polypeptide is without native mammalian glycosylation.

6. A substantially purified fragment of the IMRRP1 polypeptide of claim 1.

7. A substantially purified fragment of the IMRRP1b polypeptide of claim 2.

8. A substantially purified IMRRP1 polypeptide encoded by a polynucleotide having nucleic acid sequence SEQ ID NO: 1.

9. A substantially purified IMRRP1b polypeptide encoded by a polynucleotide having nucleic acid sequence SEQ ID NO: 2.

10. A pharmaceutical composition comprising substantially purified IMRRP1 or substantially purified IMRRP1b or a fragment thereof and a pharmaceutically acceptable excipient.

11. A purified antibody which binds specifically to the polypeptide of any one of claims 1 or 2 or antigenic epitope thereof.

12. An isolated and purified polynucleotide encoding an IMRRP1 polypeptide or fragment thereof consisting of amino acid sequence SEQ ID NO: 3.

13. An isolated and purified polynucleotide encoding an IMRRP1b polypeptide or fragment thereof consisting of amino acid sequence SEQ ID NO: 4.

14. An isolated polynucleotide comprising a nucleic acid sequence having: (a) SEQ ID NO: 1, (b) a nucleic acid sequence degenerate from SEQ ID NO: 1 as a result of the genetic code, or a nucleic acid sequence complementary to either (a) or (b).

15. An isolated polynucleotide comprising a nucleic acid sequence having: (a) SEQ ID NO: 2, (b) a nucleic acid sequence degenerate from SEQ ID NO: 2 as a result of the genetic code, or a nucleic acid sequence complementary to either (a) or (b).

16. The isolated polynucleotide according to claims 14 or 15 wherein the complementary nucleic acid sequence hybridizes to either strand of a denatured, double-stranded polynucleotide comprising the nucleic acid under conditions

of moderate stringency in 50% formamide and 6×SSC, at 42° C. with washing conditions at 60° C., 0.5×SSC, 0.1% SDS.

17. An expression vector comprising the polynucleotide of any one of claims 12.

18. An expression vector according to claim 17 that expresses a soluble IMRRP1 or a soluble IMRRP1b polypeptide.

19. A host cell containing the expression vector of claim 18.

20. A method for producing an IMRRP1 or IMRRP1b polypeptide comprising the steps of:

(a) culturing the host cell of claim 19 under conditions suitable for the expression of the polypeptide; and

(b) recovering the polypeptide from the host cell culture.

21. A hybridization probe or primer comprising an oligonucleotide or polynucleotide of a sequence capable of hybridizing with a polynucleotide of SEQ ID NO: 1 or 2 under moderate to high stringency conditions characterized in that the sequence comprises 10 or more contiguous bases.

22. A hybridization probe or primer of claim 21 characterized in that it is capable of hybridizing with a polynucleotide of SEQ ID NO: 1 or 2 under high stringency conditions.

23. A method for detecting a polynucleotide encoding an IMRRP1 or IMRRP1b polypeptide or fragment thereof in a biological sample containing nucleic acid material, the method comprising the steps of:

(a) hybridizing the oligonucleotide of claim 21 or 22 to the nucleic acid material of the biological sample, thereby forming a hybridization complex; and

(b) detecting the hybridization complex, wherein the presence of the complex correlates with the presence of the polynucleotide encoding the IMRRP1 or IMRRP1b polypeptides or fragment thereof in the biological sample.

24. The method of claim 23, wherein the nucleic acid material of the biological sample is amplified by the polymerase chain reaction before the hybridizing step.

25. A method for detecting IMRRP1 or IMRRP1b polypeptides or antigenic fragments thereof in a sample, comprising:

(a) contacting the sample with an antibody specific for IMRRP1 or IMRRP1b polypeptides or antigenic fragment thereof under conditions in which an antigen-antibody complex can form between the antibody and the IMRRP1 or IMRRP1b polypeptides or antigenic fragment thereof in the sample; and

(b) detecting an antigen-antibody complex formed in step (a), wherein detection of the complex indicates the presence of the IMRRP1 or IMRRP1b polypeptides or antigenic fragments thereof in the sample.

26. A method of identifying candidate ligands which bind to an IMRRP1 or IMRRP1b polypeptide comprising:

(a) contacting a test compound with IMRRP1 or IMRRP1b polypeptide or ligand binding portion thereof,

(b) selecting as candidate ligands those test compounds which bind to IMRRP1 or IMRRP1b or ligand binding portion thereof.

27. The method according to claim 26, wherein IMRRP1 or IMRRP1 is soluble, bound to a substrate, or cell membrane associated.

28. The method according to claim 26, wherein the method is a competitive inhibition assay.

29. The method according to claim 26, wherein said binding is detected using an antibody.

* * * * *

专利名称(译)	新型咪唑啉受体同系物		
公开(公告)号	US20020161191A1	公开(公告)日	2002-10-31
申请号	US09/932145	申请日	2001-08-17
[标]申请(专利权)人(译)	FEDER JOHNñ MINTIER GABE KINNEY基因摹 RAMANATHAN CHANDRA小号		
申请(专利权)人(译)	FEDER约翰N. MINTIER GABE KINNEY基因G. RAMANATHAN CHANDRA S.		
当前申请(专利权)人(译)	FEDER约翰N. MINTIER GABE KINNEY基因G. RAMANATHAN CHANDRA S.		
[标]发明人	FEDER JOHN N MINTIER GABE KINNEY GENE G RAMANATHAN CHANDRA S		
发明人	FEDER, JOHN N. MINTIER, GABE KINNEY, GENE G. RAMANATHAN, CHANDRA S.		
IPC分类号	G01N33/50 A61K38/00 A61K39/395 A61P9/00 A61P25/04 A61P25/28 A61P25/36 A61P43/00 C07K14/705 C07K16/28 C12N1/15 C12N1/19 C12N1/21 C12N5/10 C12N15/09 C12N15/12 C12P21/02 G01N33/15 G01N33/53 G01N33/566 C07K1/00 C07K14/00 C07K17/00		
CPC分类号	C07K14/705 A61K38/00 A61P25/04 A61P25/28 A61P25/36		
优先权	60/226411 2000-08-18 US 60/261779 2001-01-16 US		
外部链接	Espacenet USPTO		

摘要(译)

描述了新的咪唑啉受体同系物，称为咪唑啉受体相关蛋白1 (IMRRP1)，咪唑啉受体相关蛋白1b (IMRRP1b) 及其衍生物。提供包含至少一种IMRRP1，IMRRP1b或其功能部分的药物组合物，以及产生IMRRP1，IMRRP1b或其功能部分的方法。另外，提供了编码其多肽，寡核苷酸，片段，部分或反义分子的核酸序列，以及包含编码IMRRP1或IMRRP1b的多核苷酸的表达载体和宿主细胞。核酸序列，多肽，肽和抗体用于诊断和治疗与血压异常调节，进食诱导，蓝斑神经元放电刺激和胰岛素释放刺激相关的疾病或疾病的用途，以及胶质纤维酸性蛋白表达的异常诱导，不依赖于α-2肾上腺素能受体，烦躁经前期综合征，阿尔茨海默病等神经退行性疾病，阿片成瘾，单胺转换，因此伤害感受，衰老，情绪和中风，唾液疾病和还描述了发育障碍。

```

ATGTTCCGCTCCGCCCCAGCGTCCCTGGCCATGACGACCGCTCAGAGGACTCCCTG
TTGTGGAGCTCCGCGGCTTCTGCGGGAGTCCGGGATGTGGTCTCCCTGCTGGCTGTAGC
ACCTGAGCTCTGACTCCGCACTGCAAGAGTGAACAGATATTGGCTGGACTG
GGCCATGGGCCCCCTGGCAGCAGGCTTTGTGGCTCCGCTCCATCTCCGAGCTCC
CCTGTATTCTTCAAGCTTCAAGTCTCTCTCGATGTCGCGAAGAACCTTCACTCAAG
CTGGTCCATGTGGCTGGTCTGGCCCCACAGGGCCGATCAAGATTTCCCTTCAAAATCC
CTTCGGACCTGGAGCTCCGAGTGTTCCTCCCTCCACTGTTGCATGGCTCCGAGGATC
TACTCCAGCTGGAGACCTGATTTGGAGGAGAGCTCCGAGGATTAGAGGAGTCTCTC
TCAGCTCGGGGCGACTCTGCTCTCCCTCCCTTGGCTGGCTCTGCTTCTGCGAAC
TTGAGTCAATGCACTGACCGCTTAGAGAGCTCCCTGGCGCTCTGTCAGCTCTGCT
TTCTTGAACCTAAGCCAGCAAGTCAAGTCAAGTCTGAGGATTCGATGATTTGTGT
GAGCTCCACATCGGACTCTCTTAATATCCCTGGATTTGGTGGCAAGATGGAGCC
TCAGGGCTGCTCTGGGGTCTGATACTCGAGCAATGAGCTTCGAGCTCGTAGGC
CTAGAGCTGAGGAATCTCGCGCACTGGATTTGGATACAACCTGCTGGAGGACAC
CGGGACTGTCAACTGTGGCTGGCTGAGCTCCGCAAGCTTACTAGGGGGAAC
CCTCTTGGTCCACCTGAGCACCGAGCAGCACTGCCAGTACTGTACCCCGGGCC
AGGATGCTGCTACTGGCTTCTCTCGATGCAAGTCTTGTCACTGACAGATTTTCAG
ACTCACACATCTTGGGCTCAGCCCATGGCCCACTTTGCCCTGGCAGTGGGAGT
ACTCTGAACTCAGGTGGCTTAAAGTCAAGTCAAGCTTCTCCAGGGGTTGTGTG
ACCAGCCCTGCTCATAGTTAAGGCTAAGCGGATCGTGTGAGGGGGAGTCTCTT
GAACCGATACAGGACCCGAGCCCGCACTTGAACCCCTCCCGCTGGATGGTTC
GTGAGCAGCACCCGAGCTGGAGCTATGAGAGCTTCCGGGAACTTCCGCGCAAC
TGGCTGAGTACAGGAGTCAAGTCCGAGCCCTCCGAAAGCTCTGCGGCGCCACCCCACT

```

Figure 1A