



(19) **United States**

(12) **Patent Application Publication** (10) **Pub. No.: US 2004/0062769 A1**
Scherf et al. (43) **Pub. Date: Apr. 1, 2004**

(54) **IDENTIFICATION OF THE DOMAIN OF PLASMODIUM FALCIPARUM ERYTHROCYTE MEMBRANE PROTEIN 1 (PFEMP1) THAT MEDIATES ADHESION TO CHONDROITIN SULFATE A**

(76) Inventors: **Arthur Scherf**, Paris (FR); **Louis H. Miller**, Bethesda, MD (US); **Benoit Gamain**, Washington, DC (US); **Dror I. Baruch**, Rockville, MD (US); **Pierre Buffet**, Paris (FR); **Christine Scheidig**, Savigny le Temple (FR); **Jurg Gysin**, St. Zacharie (FR); **Bruno Pouvelle**, Saint Maximin la Sainte Baume (FR); **Nobutaka Fujii**, Kyoto (JP); **Joseph Smith**, Fort Collins, CO (US)

Correspondence Address:
KNOBBE MARTENS OLSON & BEAR LLP
2040 MAIN STREET
FOURTEENTH FLOOR
IRVINE, CA 92614 (US)

(21) Appl. No.: **10/087,013**

(22) Filed: **Feb. 21, 2002**

Related U.S. Application Data

(63) Continuation of application No. PCT/US00/24195, filed on Sep. 1, 2000.
(60) Provisional application No. 60/152,023, filed on Sep. 1, 1999.

Publication Classification

(51) **Int. Cl.⁷** **A61K 39/395**; G01N 33/569;
G01N 33/53
(52) **U.S. Cl.** **424/151.1**; 435/7.22

(57) **ABSTRACT**

The present invention relates to the discovery of a var gene and corresponding protein that modulates adhesion of parasitized red blood cells to chondroitin sulfate A. Novel biological tools, prophylactics, therapeutics, diagnostics, and methods of use of the foregoing are also disclosed.




FIG. 1A





FIG. 1B



Surface expression in CHO cells

FIG. 1C




FIG. 2B




FIG. 2A

**IDENTIFICATION OF THE DOMAIN OF
PLASMODIUM FALCIPARUM ERYTHROCYTE
MEMBRANE PROTEIN 1 (PFEMP1) THAT
MEDIATES ADHESION TO CHONDROITIN
SULFATE A**

**CROSS-REFERENCE TO RELATED
APPLICATIONS**

[0001] This application is a continuation of international application number PCT/US00/24195, and claims the benefit of priority of international application number PCT/US00/24195 having international filing date of Sep. 1, 2000, designating the United States of America and published in English, which claims the benefit of priority of U.S. provisional patent application No. 60/152,023, filed Sep. 1, 1999; both of which are hereby expressly incorporated by reference in their entireties.

FIELD OF THE INVENTION

[0002] The present invention relates to the discovery of a var gene and corresponding protein that modulates adhesion of parasitized red blood cells to chondroitin sulfate A. Novel biological tools, prophylactics, therapeutics, diagnostics, and methods of use of the foregoing are also disclosed.

BACKGROUND OF THE INVENTION

[0003] *Plasmodium falciparum* malaria is more severe in pregnant women, especially during the first pregnancy (primigravida), and causes disease in the mother and fetal death even in those women who were previously immune. (Steketee, et al., *Am J Trop Med Hyg* 55, 2-7 (1996)). In the primigravida, massive numbers of parasitized red blood cells (PRBCs) sequester in the maternal circulation of the placenta, binding to chondroitin sulfate A (CSA). (Fried & Duffy, *Science* 272, 1502-1504 (1996)). Antibodies that develop after multiple pregnancies are associated with reduced PRBCs in the placenta and block CSA-binding of PRBCs. (Fried, et al., *Nature* 395, 851-2 (1998)).

[0004] Members of the recently described var gene family and their expressed proteins, *Plasmodium falciparum* Erythrocyte Membrane Protein-1 (PFEMP1), mediate PRBCs binding to several adhesion receptors such as CD36, intercellular adhesion molecule-1 (ICAM-1), and chondroitin sulfate A (CSA). (Baruch, et al., *Cell* 82, 77-87 (1995), (Smith, et al., *Cell* 82, 101-10 (1995), (Su, et al., *Cell* 82, 89-100 (1995), and (Scherf, et al., *Embo J* 17, 5418-5426 (1998)). Recent work on var gene switching has established that transcription of a particular var gene (termed "FCR3.varCSA") in parasites selected for binding to CSA but not in parasites selected for adhesion to CD36 or ICAM-1. (Scherf, et al., *Embo J* 17, 5418-5426 (1998)). Thus, var genes adhere dichotomously either to CD36 and other receptors on endothelium or to CSA in placenta and not to CD36.

[0005] Potential receptor domains in var genes include Duffy binding like (DBL) domains, named for their homology to the Duffy binding domain of *P. vivax* (Su, et al., *Cell* 82, 89-100 (1995)), and cysteine-rich interdomain regions (CIDR). The CIDR1 domain, located after the first DBL, was shown to mediate PRBCs adhesion to CD36. (Baruch, et al., *Cell* 82, 77-87 (1995) and (Baruch, et al., *Blood* 90, 3766-75 (1997)). DBL1 has been identified as a receptor for

binding PRBCs to uninfected RBCs in var genes from PRBCs that rosette normal RBCs. (Rowe, et al., *Nature* 388, 292-5 (1997) and (Chen, et al., *J Exp Med* 187, 15-23 (1998)). Although antibodies directed to two different domains of a var gene expressed in CSA-binding parasites reduced binding to CSA (Reeder, et al., *Proc Natl Acad Sci USA* 96, 5198-202 (1999)), the gene, protein and domains thereof that bind CSA have not been identified.

BRIEF SUMMARY OF THE INVENTION

[0006] The invention described herein concerns the discovery of molecules that are intimately involved in PRBC binding, sequestration, and the onset of maternal malaria. One such molecule is the product of the FCR3.varCSA gene, a 3,542-amino acid polypeptide called the FCR3.varCSA protein, which binds to CSA. Other molecules that mediate PRBC binding, sequestration, and the onset of maternal malaria include fragments of the FCR3.varCSA protein (e.g., polypeptides that comprise the CIDR1 and/or the DBL3 domains or portions thereof) and other varCSA proteins and fragments thereof including, but not limited to polypeptides having the A4 tres DBL3- γ and ItG2-CS2 DBL2- γ (SEQ. ID. Nos: 9 and 11, respectively) sequence. Furthermore, nucleic acids encoding these molecules can be used to modulate PRBC binding, sequestration, and the onset of maternal malaria.

[0007] The FCR3.varCSA gene was cloned and sequenced in its entirety and the FCR3.varCSA protein is predicted to have eight receptor-like domains. To further characterize the FCR3.varCSA-CSA complex, several adhesion assays (referred to as "varCSA characterization assays" or "FCR3.varCSA characterization assays") were performed. In some experiments, proteins encompassing various domains of FCR3.varCSA or other varCSA polypeptides were expressed on the surface of CHO cells and adhesion to various ligands was analyzed. From these characterization assays it was discovered that two Duffy-binding-like (DBL) domains (DBL3 and DBL7) of FCR3.varCSA were involved in adhesion to CSA. Further, it was found that DBL7, but not DBL3, bound chondroitin sulfate C (CSC), a negatively charged sugar that does not support PRBC adhesion. Competitive binding experiments employing exogenously added CSA prevented the interaction with DBL3, however, either competitor (i.e., exogenously added CSA or CSC) prevented adhesion to DBL7. Thus, evidence is provided herein that the DBL3 and/or CIDR1 domain of FCR3.varCSA are intimately involved in PRBC binding, sequestration, and the onset of maternal malaria.

[0008] Many different forms of var genes exist due to gene switching and it was believed that some of these gene products and fragments thereof also specifically bind CSA. To verify this hypothesis, several adhesion assays were conducted using CHO cells that cell-surface-express polypeptides having various types of varCSA domains. These experiments revealed that some domains of other varCSA molecules effectively bound CSA (e.g., A4 tres DBL3- γ (SEQ. ID. No.: 9) and ItG2-CS2 DBL2- γ (SEQ. ID. No.: 11)) while others (e.g., R29)BL2- γ (SEQ. ID. NO.: 7), A4 DBL4- γ (SEQ. ID. No.: 8), and FCR3 var3 DBL- γ (SEQ. ID. No.: 10) did not.

[0009] Several embodiments concern the interaction of FCR3.varCSA with CSA, the formation of a FCR3.varCSA-

CSA complex, PRBC binding, sequestration, and the onset of maternal malaria. For example, embodiments include the FCR3.varCSA-CSA complex, FCR3.varCSA protein, fragments of FCR3.varCSA protein (e.g., DBL3 and CIDR1), nucleic acids encoding these polypeptides, cells that have these nucleic acids, cells that express these polypeptides, antibodies that recognize these polypeptides, and software and hardware that have nucleotide or polypeptide information or protein modeling information corresponding to these sequences, as well as, data from FCR3.varCSA characterization assays and diagnostic profiles.

[0010] Other embodiments concern the interaction of other varCSA molecules including, but not limited to, varCSA molecules having the A4 tres DBL3- γ (SEQ. ID. No.: 9) and ItG2-CS2 DBL2- γ (SEQ. ID. No.: 11) sequences, with CSA and the formation of a varCSA-CSA complex. For example, embodiments include a varCSA-CSA complex, fragments of a varCSA protein (e.g., A4 tres DBL3- γ (SEQ. ID. No.: 9) and ItG2-CS2 DBL2- γ (SEQ. ID. No.: 11)), nucleic acids encoding these polypeptides, cells that have these nucleic acids, cells that express these polypeptides, antibodies that recognize these polypeptides, and software and hardware that have nucleotide or polypeptide information or protein modeling information corresponding to these sequences, as well as, data from varCSA characterization assays and diagnostic profiles.

[0011] Additionally, nucleic acids that complement nucleic acids encoding FCR3.varCSA or fragments of FCR3.varCSA or other varCSA molecules that bind CSA (e.g., A4 tres DBL3- γ (SEQ. ID. No.: 9) and ItG2-CS2 DBL2- γ (SEQ. ID. No.: 11) and cells that have these sequences are embodiments. Another aspect of the invention includes the use of therapeutic or prophylactic agents (e.g., FCR3.varCSA or fragments of FCR3.varCSA, A4 tres DBL3- γ (SEQ. ID. No.: 9) and ItG2-CS2 DBL2- γ (SEQ. ID. No.: 11) or nucleic acids encoding these compositions) to modulate adhesion to CSA and/or to generate an immune response in a patient. Further, methods of discovering such agents including approaches in rational drug design and combinatorial chemistry are also embodiments.

[0012] Other embodiments include biotechnological tools, diagnostic assays, diagnostic kits, and methods of use of the foregoing. For example, multimeric and multimerized FCR3.varCSA, fragments of FCR3.varCSA, A4 tres DBL3- γ , and ItG2-CS2 DBL2- γ and nucleic acids encoding these sequences or complementary sequences are used as biotechnological tools or diagnostic reagents. Diagnostic assays preferably measure the concentration or expression level of FCR3.varCSA or nucleic acid encoding FCR3.varCSA in tested subjects and compare these values to those obtained from healthy individuals or individuals that are infected with *Plasmodium falciparum* (FCR3.varCSA disease-state profiles). Additionally, some diagnostic assay embodiments measure the concentration or expression level of proteins or polypeptides comprising the A4 tres DBL3- γ and ItG2-CS2 DBL2- γ fragments or nucleic acids encoding these molecules in tested subjects and compare these values to those obtained from healthy individuals or individuals that are infected with *Plasmodium falciparum*. These varCSA disease-state profiles can be recorded on software and hardware and can be used to analyze disease-state profiles of tested subjects so as to identify the presence or prevalence of maternal malaria or progress of a treatment for maternal

malaria. Desirably, measurements of the concentration or expression level of the varCSA proteins or polypeptides or nucleic acids encoding these molecules are made from blood. These disease-state profiles are invaluable tools for the prognosis, diagnosis, and treatment of FCR3.varCSA-related diseases, including, but not limited to, maternal malaria.

[0013] Pharmaceuticals having FCR3.varCSA or fragments of FCR3.varCSA (e.g., DBL3 and/or CIDR1) or nucleic acids encoding these polypeptides or antibodies that recognize these molecules or agents that otherwise interact with FCR3.varCSA are also embodiments. The pharmaceutical embodiments may also comprise polypeptides having the A4 tres DBL3- γ and ItG2-CS2 DBL2- γ sequence or nucleic acids encoding these molecules. The pharmaceuticals described herein can also include carriers and other agents that promote delivery of the active ingredients.

[0014] Further, methods of treatment and prevention of malaria, specifically maternal malaria, are provided. Some methods of treatment and prevention of maternal malaria, involve identifying a subject in need of an agent that inhibits the association of a varCSA molecule (e.g., FCR3.varCSA) with CSA and administering to said subject a therapeutically effective dose of an agent that either inhibits adhesion of the varCSA molecule to CSA and/or promotes an immune response in a patient. Other methods involve identifying a patient in need of an agent that inhibits PRBC binding, sequestration, or the onset of maternal malaria and administering to said patient a composition comprising the CIDR1 domain or fragment thereof or an antibody that recognizes a CIDR1 domain. Preferably, this composition is derived from FCR3.varCSA in that it comprises a CIDR1 domain or antibody thereto or fragment thereof that is derived from FCR3.varCSA.

BRIEF DESCRIPTION OF THE DRAWINGS

[0015] **FIG. 1(A)** Overlapping clones of the FCR3.varCSA gene were isolated from genomic FCR3-CSA parasites and sequenced (gDNA). Regions amplified by RT-PCR (cDNA) from FCR3-CSA trophozoite mRNA confirm that the genomic FCR3.varCSA gene sequence is contiguous with the exception of the intron region. **(B)** Schematic domain organization of the FCR3.varCSA gene. An unusually small intron of 230 bp separates exon 1 and exon 2 of the FCR3.varCSA gene. The amino acid boundaries of the different DBL (Duffy Binding Like) and CIDR1 (Cysteine-rich Interdomain Region) domains are indicated. **(C)** Domain regions that were expressed on the surface of CHO-745 cells showing their amino acid boundaries.

[0016] **FIG. 2(A)** Binding of anti-biotin coated Dynabeads to CHO-745 cells expressing different domains of FCR3.varCSA incubated with CSA-biotin. The percentage of transfected cells that bound 4 or more beads are shown. **(B)** Inhibition of binding of CSA-biotin and CSC-biotin to DBL-3 and DBL-7 transfectants. Transfected cells were incubated with biotin-CSA or biotin-CSC without (control), or after preincubation with 200 μ g/ml CSA (+CSA) or CSC (+CSC). Binding is given as number of positive cells (A) or number of beads (B) per 100 cells. Error bars represent the standard deviation from three different experiments.

DETAILED DESCRIPTION OF THE
INVENTION

[0017] Several molecules that mediate PRBC binding, sequestration, and the onset of maternal malaria have been discovered. One such molecule is the product of the FCR3.varCSA gene, a 3,542-amino acid polypeptide called the FCR3.varCSA protein. Other molecules include fragments of the FCR3.varCSA protein (e.g., DBL3 and/or CIDR1) and other varCSA proteins and fragments thereof including, but not limited to, polypeptides having the A4 tres DBL3- γ and ItG2-CS2 DBL2- γ (SEQ. ID. Nos: 9 and 11, respectively) sequence. The adhesion of *Plasmodium falciparum* infected red blood cells, also referred to as "parasitized red blood cells (PRBCs), to chondroitin sulfate A (CSA) is intimately involved in sequestration of *P. falciparum*, and the manifestation of maternal malaria. Thus, many of the embodiments described herein can be used as biological tools, therapeutics, prophylactics, and diagnostics for the study, treatment and prevention of maternal malaria.

[0018] Embodiments of the invention include software and hardware comprising nucleic acid sequences encoding FCR3.varCSA or fragments thereof (e.g., nucleic acids encoding molecules that comprise DBL3 and/or CIDR1) or complements of these sequences and protein sequences corresponding to FCR3.varCSA and fragments of FCR3.varCSA (e.g., DBL3 and/or CIDR1). Preferred software and hardware have nucleic acid sequences that encode fragments of FCR3.varCSA that bind to chondroitin sulfate A (e.g., DBL3 and/or CIDR1) or amino acid sequences that correspond to regions of a var protein that bind CS A. Additionally, the software and hardware of the invention include embodiments that provide disease-state profiles that have information such as concentrations and expression levels of FCR3.varCSA (e.g., mRNA) or FCR3.varCSA detected in biological samples from healthy subjects, as well as, subjects suffering from malaria. The software and hardware embodiments of the invention are also used to further characterize FCR3.varCSA (e.g., to develop protein models of FCR3.varCSA, to identify homologous proteins, and to identify agents that interact with FCR3.varCSA) and to provide diagnostic and prognostic information that allows for the determination of the disease state of a tested individual.

[0019] Nucleic acids encoding full-length FCR3.varCSA or nucleic acids encoding fragments of FCR3.varCSA (e.g., DBL3 and/or CIDR1) are embodiments of the invention. Preferred nucleic acid embodiments include nucleic acids encoding fragments of FCR3.varCSA that bind to CSA (e.g., DBL3) or otherwise mediate PRBC binding, sequestration, and the onset of maternal malaria (e.g., CIDR1). Additionally, the nucleic acid embodiments of the invention include nucleic acids or derivatives thereof that are complementary to full-length FCR3.varCSA or fragments of FCR3.varCSA (e.g., antisense oligonucleotides and ribozymes). Preferred complementary nucleic acids of the invention include nucleic acids or derivatives thereof that are complementary to fragments of FCR3.varCSA that have a nucleotide sequence that encodes a polypeptide that binds to CSA (e.g., DBL3) or otherwise mediates PRBC binding, sequestration, and the onset of maternal malaria (e.g., CIDR1).

[0020] The nucleic acid embodiments can be manufactured as monomeric, multimeric, and multimerized agents.

The nucleic acid embodiments also include vectors, plasmids, and recombinant constructs having nucleic acids encoding full-length FCR3.varCSA or fragments of FCR3.varCSA. Additional embodiments are vectors, plasmids, and recombinant constructs having nucleic acids complementary to the full-length FCR3.varCSA or fragments of FCR3.varCSA. Cells having the nucleic acid embodiments described herein, including cells in animals having a nucleic acid embodiment created by genetic engineering (e.g., cells in a transgenic animal or an oocyte), are within the scope of aspects of the invention.

[0021] The nucleic acid embodiments also include nucleic acids encoding fragments of other varCSA proteins that bind CSA. For example, some embodiments concern nucleic acids that encode polypeptides comprising the A4 tres DBL3- γ and ItG2-CS2 DBL2- γ (SEQ. ID. Nos: 9 and 11, respectively) sequence or complements thereto. These nucleic acid embodiments can be manufactured as monomeric, multimeric, and multimerized agents and can be cloned into vectors, plasmids, and recombinant constructs. Furthermore, cells having nucleic acids that encode polypeptides comprising the A4 tres DBL3- γ and ItG2-CS2 DBL2- γ (SEQ. ID. Nos: 9 and 11, respectively) sequence or complements thereto, including cells in animals having a nucleic acid embodiment created by genetic engineering (e.g., cells in a transgenic animal or an oocyte), are embodiments.

[0022] Protein-based embodiments include full-length FCR3.varCSA or fragments of FCR3.varCSA. Preferred protein-based embodiments include fragments of FCR3.varCSA that have an amino acid sequence that encode a polypeptide that binds to CSA (e.g., DBL3) or otherwise mediates PRBC binding, sequestration, and the onset of maternal malaria (e.g., CIDR1). Additionally, the protein-based embodiments include protein derivatives or modifications of FCR3.varCSA or fragments of FCR3.varCSA including, but not limited to peptidomimetics. The protein-based embodiments can be manufactured as monomeric, multimeric, and multimerized agents. Cells having the protein-based embodiments, including cells in animals having a protein-based manufacture of the present invention (e.g., cells in a transgenic animal or an oocyte), are within the scope of aspects of the invention.

[0023] In some embodiments, the polypeptides described herein are used to generate antibodies. Preferred embodiments include polyclonal and monoclonal antibodies that recognize epitopes corresponding to regions of FCR3.varCSA (e.g., DBL3 and CIDR1). These antibodies have application in biological assays, therapeutics, and can be used to diagnose human disease by identifying the presence of FCR3.varCSA in a biological sample.

[0024] The protein-based embodiments also include other varCSA proteins and fragments thereof that bind CSA. For example, some embodiments concern polypeptides comprising the A4 tres DBL3- γ and ItG2-CS2 DBL2- γ (SEQ. ID. Nos: 9 and 11, respectively) sequence. As above, these embodiments can be manufactured as monomeric, multimeric, and multimerized agents. Embodiments also include cells having polypeptides comprising the A4 tres DBL3- γ and ItG2-CS2 DBL2- γ (SEQ. ID. Nos: 9 and 11, respectively) sequence, including cells in animals (e.g., cells in a transgenic animal or an oocyte). In some embodiments,

these polypeptides are used to generate monoclonal and/or polyclonal antibodies, which have diagnostic and therapeutic application.

[0025] Several types of assays that provide information about a particular varCSA molecule (e.g., FCR3.varCSA) or the formation of a particular varCSA-CSA complex (e.g., FCR3.varCSA-CSA complex) are embodiments. These assays are collectively referred to as "FCR3.varCSA characterization assays" or "varCSA characterization assays". One type of varCSA characterization assay concerns measuring the ability of FCR3.varCSA or fragments thereof to bind to CSA or fragments of CSA. For example, methods of performing characterization assays are provided, in which CSA or FCR3.varCSA is disposed on a support and is subsequently contacted with a ligand (e.g., FCR3.varCSA, or CSA, depending on the support-bound molecule) and FCR3.varCSA-mediated adhesion is determined. A similar binding assay can be employed in the presence of an inhibiting or enhancing molecule (a "modulator") such as a peptide or peptidomimetic (collectively referred to as a "peptide agent") or a chemical. The supports in these assays can be conventional resins, plastics, lipids, and cells. Thus, in some FCR3.varCSA characterization assays cells having FCR3.varCSA or a fragment thereof at the cell membrane (e.g., accomplished by transfection or liposome transfer) are used to identify agents that interfere with FCR3.varCSA mediated adhesion.

[0026] In some aspects, the modulation of FCR3.varCSA-mediated adhesion is accomplished by using a modulator that is a nucleic acid embodiment. For example, a construct encoding FCR3.varCSA is transfected into cells so as to raise the concentration of FCR3.varCSA and thereby promote FCR3.varCSA-mediated adhesion to CSA or, alternatively, a construct encoding a nucleic acid that is complementary to a nucleic acid encoding FCR3.varCSA (e.g., an antisense inhibitor or a ribozyme) is used to reduce the concentration of FCR3.varCSA and thereby inhibit FCR3.varCSA-mediated adhesion to CSA. Further, in some embodiments, nucleic acids encoding wild-type or mutant FCR3.varCSA or fragments of FCR3.varCSA or complements thereof are transfected and expressed in cells so as to modulate FCR3.varCSA-mediated adhesion or to induce an immune response or both.

[0027] According to other aspects, the modulation of FCR3.varCSA-mediated adhesion is achieved by using a modulator that is a protein-based embodiment. For example, FCR3.varCSA is delivered to cells by liposome-mediated transfer so as to raise the intracellular concentration of FCR3.varCSA and thereby promote FCR3.varCSA-mediated adhesion to CSA or, alternatively, wild-type or mutant FCR3.varCSA or fragments of FCR3.varCSA (e.g., DBL3 and/or CIDR1) are delivered to cells by liposome-mediated transfer so as to inhibit FCR3.varCSA-mediated adhesion to CSA or to induce an immune response or both. Peptidomimetics that resemble FCR3.varCSA or fragments thereof (e.g., DBL3 and/or CIDR1) are also modulators of the invention and can be used to effect FCR3.varCSA mediated adhesion or to induce an immune response or both. Many chemicals can also be modulators and can be identified by their ability to effect FCR3.varCSA mediated adhesion using the FCR3.varCSA characterization assays and teachings herein.

[0028] Approaches in rational drug design can be employed, for example, to identify novel agents that interact with FCR3.varCSA so as to modulate FCR3.varCSA-mediated adhesion or that can be used to induce an immune response in a patient. In these embodiments, protein models of FCR3.varCSA, fragments of FCR3.varCSA, and agents that interact with FCR3.varCSA or fragments of FCR3.varCSA are constructed and approaches in combinatorial chemistry are used to develop agents that modulate FCR3.varCSA-mediated adhesion to CSA or induce an immune response. Accordingly, novel agents that interact with FCR3.varCSA are developed, screened in a FCR3.varCSA characterization assay (e.g., a FCR3.varCSA adhesion assay), and the identity of each agent and its performance in a FCR3.varCSA characterization assay, its effect on the modulation FCR3.varCSA-mediated adhesion to CSA or its ability to induce an immune response is recorded on software or hardware. The recorded data can be used to create a library of FCR3.varCSA modulating agents. These libraries can be employed to identify more agents that modulate FCR3.varCSA-mediated adhesion to CSA and are valuable clinical tools for manufacturing and selecting an appropriate pharmaceutical to treat a particular type of Plasmodium.

[0029] The nucleic acid and protein-based embodiments described herein can also be used as biotechnological tools and probes in diagnostic assays. In some aspects, for example, the nucleic acid embodiments are employed as nucleic acid probes in hybridization assays, cloning, or as primers for Polymerase Chain Reaction (PCR). Similarly, the protein-based embodiments can be used, for example, to characterize FCR3.varCSA or other varCSA molecules that bind CSA, identify related proteins, and study varCSA-mediated adhesion to CSA.

[0030] In some diagnostic embodiments, nucleic acids complementary to full-length FCR3.varCSA or fragments of FCR3.varCSA are used to identify FCR3.varCSA nucleic acids (e.g., mRNA) present in a biological sample. In other diagnostic embodiments, nucleic acids complementary to nucleic acids that encode polypeptides comprising the A4 tres DBL3- γ and ItG2-CS2 DBL2- γ (SEQ. ID. Nos: 9 and 11, respectively) sequence are used to identify FCR3.varCSA nucleic acids (e.g., mRNA) present in a biological sample. In preferred diagnostic embodiments, however, nucleic acids complementary to fragments of FCR3.varCSA that comprise sequence not found in the nucleic acid encoding other var proteins are used to identify FCR3.varCSA nucleic acids (e.g., mRNA) present in a biological sample.

[0031] Depending on the type of Plasmodium present in the biological sample, the concentration or expression level of nucleic acid encoding FCR3.varCSA or other varCSA that binds CSA can differ. That is, an individual having one form of malaria can be infected with a type of parasite that produces a lower amount of a particular type of varCSA (e.g., FCR3.varCSA), or none at all. Additionally, healthy individuals will not express varCSA. Thus, malaria and, more specifically, a type of Plasmodium infection that leads to maternal malaria can be diagnosed by determining the concentration or expression level of a nucleic acid encoding a varCSA molecule that binds CSA (e.g., a mRNA encoding FCR3.varCSA).

[0032] For example, a FCR3.varCSA-disease state profile comprising a concentration range of a nucleic acid encoding FCR3.varCSA in a biological sample can be created for healthy and diseased individuals and these FCR3.varCSA disease state profiles can be compared to the concentrations or expression levels of a nucleic acid encoding FCR3.varCSA detected in a tested individual so as to predict or follow the disease state of that individual. Thus, in some embodiments, the term "FCR3.varCSA-disease state profile" refers to the concentration or expression level or concentration range or expression level range of a nucleic acid encoding FCR3.varCSA that is detected in a biological sample. Desirably, addressable arrays comprising nucleic acid probes complementary to the full-length FCR3.varCSA or fragments of FCR3.varCSA are used to create such FCR3.varCSA-disease state profiles. Such arrays or individual probes are also components of diagnostic kits.

[0033] In similar fashion to that discussed above, a FCR3.varCSA-disease state profile comprising concentration ranges or levels of FCR3.varCSA in healthy and diseased individuals can be created and can be used to predict or follow the disease state of an individual. In some embodiments, the term "FCR3.varCSA-disease state profile" refers to the concentration or expression level or concentration range or expression level range of a protein corresponding to FCR3.varCSA that is detected in a biological sample. Thus, by comparing a FCR3.varCSA-disease state profile from healthy individuals and subjects infected with *P. falciparum* from different regions of the world, with the FCR3.varCSA disease state profile from a tested subject, a clinician can rapidly diagnose whether the tested subject is infected with malaria and whether the type of Plasmodium will place the individual at risk for contracting forms of malaria that can lead to maternal malaria. Desirably, addressable arrays comprising antibodies that recognize epitopes of FCR3.varCSA are used to create such FCR3.varCSA-disease state profiles. Such arrays or antibodies are also components of diagnostic kits.

[0034] In some therapeutic and prophylactic embodiments, FCR3.varCSA, polypeptide fragments of FCR3.varCSA (e.g., DBL3 and/or CIDR1), nucleic acids encoding these molecules, and agents that interact with a FCR3.varCSA-CSA complex are incorporated into pharmaceuticals. In other therapeutic and prophylactic embodiments, polypeptides comprising the A4 tres DBL3- γ and ItG2-CS2 DBL2- γ (SEQ. ID. Nos: 9 and 11, respectively) sequence, nucleic acids encoding these molecules, and agents that interact with a varCSA-CSA complex are incorporated into pharmaceuticals. In still more embodiments, antibodies directed to the molecules above (preferably CIDR1) are provided to a subject to provide protection against PRBC binding, sequestration, and the onset of maternal malaria. These pharmaceuticals can be delivered by any conventional route including, but not limited to, transdermal, parenteral, gastrointestinal, transbronchial, and transalveolar. In addition to the active ingredients mentioned above, the pharmaceutical embodiments can comprise carriers, proteins, supports, adjuvants, or components that facilitate or enhance drug delivery. These pharmaceuticals can be employed in therapeutic protocols for the treatment and prevention of maternal malaria.

[0035] Because some aspects of the invention can be used to both inhibit adhesion of FCR3.varCSA to CSA and to

generate an immune response directed at FCR3.varCSA, embodiments that administer FCR3.varCSA or fragments thereof are therapeutically and prophylactically useful. By one approach, a subject at risk for contracting maternal malaria or a subject infected with *P. falciparum* is identified by conventional techniques or the diagnostic assays described herein and then is administered an effective amount of an agent that inhibits FCR3.varCSA-mediated adhesion to CSA and/or promotes an immune response in a patient. Other methods described herein concern the inhibition of the adhesion of other varCSA proteins to CSA including proteins comprising the A4 tres DBL3- γ and ItG2-CS2 DBL2- γ (SEQ. ID. Nos: 9 and 11, respectively) sequence. Similar to the approach above, this method is practiced by identifying a subject in need of an agent that disrupts the formation of a varCSA-CSA complex and administering said subject an effective amount of an agent that inhibits the formation of the varCSA-CSA complex. In still more methods of treatment and prevention of maternal malaria, a subject in need of an agent that mediates PRBC binding, sequestration, and the onset of maternal malaria is identified and then is provided a therapeutically sufficient amount of an agent that comprises a CIDR1 domain, fragment thereof, or antibody thereto. The discovery of the FCR3.varCSA gene and FCR3.varCSA protein and its characterization as a molecule that mediates adhesion of PRBCs to CSA is disclosed below.

[0036] Identification and Isolation of the Gene Encoding FCR3.varCSA and FCR3.varCSA Protein

[0037] The var gene of FCR3-CSA-PRBCs was cloned and sequenced to identify the domain of the parasite ligand that mediates adhesion to CSA. A specific sequence tag of the FCR3.varCSA gene corresponding to the DBL-1 (Scherf, et al., *Embo J* 17, 5418-5426 (1998)) was used to extend the gene sequence in the 5' and 3' directions. The Vectorette technique (Genosys Biotechnologies Inc.) was employed to perform the extension. (Scherf, et al., *Embo J* 17, 5418-5426 (1998)). A nucleic acid of 10,628 bp, which contains the entire extracellular region encoded by exon I (9,931 bp), the intracellular domain of FCR3.varCSA encoded by exon II (698 bp), and an unusually short intron of 230 bp was obtained. (See FIG. 1). The cloned DNA sequence predicts an open reading frame of 3,542 amino acids, and an overall structure that resembles published var sequences in that 7 DBL domains and a CIDR1 domain were found. (FIG. 1B).

[0038] The linear order of the DNA sequence obtained from genomic FCR3.varCSA was confirmed using overlapping PCR fragments from cDNA of FCR3.varCSA trophozoites (FIG. 1A) and a YAC clone (gift of Dr. M. Lanzer, University of Heidelberg) spanning most of the exon I of the FCR3.varCSA gene. Probes to FCR3.varCSA sequence corresponding to DBL-1, DBL3/4, and DBL6/7 were found to hybridize to an identical large transcript of about 13 kb in total RNA of FCR3-CSA trophozoites. In this experiment, an Hsp70 specific probe hybridizing to the *P. falciparum* heat shock gene transcript of approximately 3 kb was used as a control. RT-PCR of mRNA and PCR of the genomic DNA also proved that the sequence was contiguous (FIG. 1A).

[0039] The RT-PCR and Northern analysis were performed as described (Scherf, et al., *Embo J* 17, 5418-5426

(1998)), from total parasite RNA prepared using the TRI-ZOL (Life Technologies, Gaithersburg, Mich.) extraction method (Smith, et al., *Mol Biochem Parasitol* 97, 133-48 (1998)). More evidence that the FCR3.varCSA protein is involved in CSA adhesion is provided below.

[0040] FCR3.varCSA Codes for a Large Trypsin Sensitive Erythrocyte Surface Molecule that Binds to CSA

[0041] Surface iodination of FCR3-CSA trophozoite-infected RBCs identified a single molecule of about 400 kDa having the characteristics of a var gene product that binds CSA. (Baruch, et al., *Proc Natl Acad Sci USA* 93, 3497-502 (1996)). To perform this experiment, SDS extracts of surface iodinated FCR3-CSA and FCR3-CD36 trophozoites were separated on a gel. The labeled high molecular mass proteins of approx. 400 kDa and 250 kDa were observed to be sensitive to trypsinization. Thus, evidence supporting the conclusion that the cloned gene was a member of the var family included the fact that FCR3.varCSA of intact PRBCs were sensitive to trypsin digestion and efficient extraction of FCR3.varCSA could only be achieved in a denaturing detergent (2% SDS). Notably, the iodinated portion of FCR3.varCSA was sensitive to low concentrations of trypsin but the region of the molecule that binds to CSA was not sensitive under these conditions. The surface iodination and trypsin degradation experiments were performed on *P. falciparum* FCR3 parasites that were cultured and selected on the adhesion receptors CD36 and CSA, as described in Scherf, et al., *Embo J* 17, 5418-5426 (1998). Accordingly, mature intact PRBCs were selected by the receptor panning procedure, grown for 1-2 cycles and enriched to >75% by the plasmagel technique prior to iodination. (Pasvol, et al., *Annals of Tropical Medicine & Parasitology* 72, 87-8 (1978)).

[0042] Surface iodination was accomplished by sequential extraction with 1% Triton X-100 followed by 2% SDS and trypsinization (TPCK-treated trypsin, Sigma) of PRBC, as described in Baruch, et al., *Proc Natl Acad Sci USA* 93, 3497-502 (1996). Samples derived from iodination were separated on a 0.5% agarose/4% acrylamide composite gel, dried and exposed to Kodak X-Omat XAR-5 film (Wiesner, et al., *Parasitol. Today* 14, 38-40 (1998)). Prestained protein markers were used to verify the molecular size (Life Technologies, Gaithersburg, Md. and New England BioLabs Inc. Beverly, Mass.). Additionally, antibodies directed to the internal domain of MC.var1, a region conserved in var proteins, reacted with the FCR3.varCSA protein, thus, providing more evidence that the cloned molecule was a member of the var family.

[0043] Further proof that the cloned gene was a member of the var family was obtained from extensive adhesion assays (e.g., FCR3.varCSA characterization assays). In one set of experiments, iodinated FCR3.varCSA was captured by affinity purification. Accordingly, an affinity resin was made by incubating 30 µg of recombinant human thrombomodulin with 1×10^8 tosyl activated M-450 Dynabeads (4.5 µm diameter, Dynal A. S., Norway) in 1 ml of 0.1 M phosphate buffer pH 7.4 according to the protocol provided by the manufacturer. Next the affinity resin was incubated overnight at 4° C. with 15 µl of iodinated FCR3.varCSA extract (prepared by SDS extraction of FCR3-CSA PRBCs) diluted in 500 µl of BM pH 6.8 containing 1% BSA. Beads were subsequently

washed using a magnet (Dynal MPC) and processed. (Baruch, et al., *Proc Natl Acad Sci USA* 93, 3497-502 (1996)).

[0044] It was found that the cloned FCR3.varCSA molecule was variant in that it was absent in CD36-selected PRBCs (FCR3-CD36), which instead expressed an iodinated molecule of 250 kDa. Additionally, surface iodinated FCR3.varCSA, extracted from FCR3-CSA PRBCs, bound to human thrombomodulin-coated dynabeads, whereas, FCR3-CD36 PfEMP1 did not bind thrombomodulin. Thus, CSA-containing thrombomodulin affinity purified a red cell surface molecule having properties expected of a member of the var family. The purified molecule was found to be sensitive to trypsin treatment. To further understand the properties of FCR3.varCSA, more FCR3.varCSA characterization assays were performed, as provided below.

[0045] Adhesive Phenotype of Parasites Selected for CSA Binding

[0046] The binding characteristics of CSA-selected PRBCs resemble the adhesive phenotype observed in PRBCs isolated from placentas of malaria infected women, that is, binding to CSA but not to CD36. (Fried & Duffy, *Science* 272, 1502-1504 (1996)). (See TABLE 1). Furthermore, sera from multigravid women from Cameroon and Senegal block efficiently adhesion of FCR3-CSA-PRBCs to CSA. These adhesion properties differ from those of a CD 36-selected PRBCs that bind several receptors but not to CSA. In agreement with these clinical observations, the inventors have discovered that the CIDR1 domain of FCR3.varCSA does not bind to CD36, however, the CIDR1 domain of MC.var1, ITA4.var, and FVO.var efficiently bind to CD36.

[0047] The results presented in Table 1 are the product of several adhesion assays that were conducted as follows. A stable transfectant of CHO-745 cells (CSA negative) (Rogerson, et al., *J Exp Med* 182, 15-20 (1995)) permanently expressing cDNA's of CD36 (Berendt, et al., *Nature* 341, 57-9 (1989)), ICAM-1 (Simmons, et al., *Nature* 331, 624-7 (1988)), VCAM-1 (Osborn, et al., *Cell* 59, 1203-11 (1989)) and E-selectin (Bevilacqua, et al., *Science* 243, 1160-5 (1989)) was constructed using Fugene 6 transfection reagent (Roche Diagnostics GmbH, Germany). Additionally, a stably transformed HUVEC cell line was kindly provided by D. Paulin and P. Vicart (Vicart, et al., *J Cell Physiol* 157, 41-51 (1992)). Surface expression of PECAM-1, ICAM-1, E-selectin and VCAM-1 was analyzed using specific monoclonal antibodies (R&D Systems, Europe Ltd). The mAb anti CD36 was a gift of L. Edelman, Institut Pasteur.

TABLE 1

Binding characteristics of FCR3-CSA and FCR3-CD36 parasites to various host receptors		
Adhesion Receptor	FCR3-CSA	FCR3-CD36
^a human thrombomodulin ^{CSA}	8910 ± 352	34 ± 24
CSA	3545 ± 278	68 ± 26
BIOT-CSA	2866 ± 156	22 ± 15
BIOT-CSC	32 ± 12	nd
^b placenta	850 ± 230	58 ± 46
^c CHO ^{CSA}	3450 ± 234	23 ± 34
CHO ^{CD36}	45 ± 32	2035 ± 143
CHO ^{ICAM-1}	24 ± 21	679 ± 64

TABLE 1-continued

Binding characteristics of FCR3-CSA and FCR3-CD36 parasites to various host receptors		
Adhesion Receptor	FCR3-CSA	FCR3-CD36
CHO ^{VCAM-1}	46 ± 56	456 ± 69
CHO ^{E-selectin}	82 ± 34	235 ± 36
^d human thrombospondin (TSP-1)	45 ± 34	78 ± 53
^e HUVEC ^{PECAM-1, ICAM-1, VCAM-1}	124 ± 67	1879 ± 98

^aRecombinant human thrombomodulin carrying chondroitin sulfate A (hTM) produced in CHO cells. Binding of PRBCs to receptors bound to plastic is expressed as number of PRBCs per mm² ± SD of two independent experiments.

^bAdhesion of PRBCs to serial 7 μm cryosections of snap frozen placenta tissue was performed, as described elsewhere Gysin et al., *Mol. Bioch. Parasitol.* 88: 267 (1997). Only PRBCs adhesion on syncytiotrophoblasts and syncytial bridges were counted and expressed as the mean number of PRBCs ± SE per 20 high power microscopic fields (1000x Leitz Diaplan microscope).

^cCHO-745 cells (CSA negative) stably transfected with the human adhesion receptors CD36, ICAM-1, VCAM-1 and E-selectin are described below. Cytoadherence on confluent cells is expressed as number of PRBCs per mm² ± SD of two independent experiments.

^dCytoadhesion of PRECs was performed on purified human thrombospondin (TSP) (Sigma St. Louis) at 50 μg/ml in 20 mM Tris-HCl, pH 7.2, 150 mM NaCl, 2 mM CaCl spotted onto Petri dishes. Number of PRBCs per mm² ± SD of two independent experiments.

^eNonactivated human umbilical vein endothelial cells (HUVEC) express PCAM-1, ICAM-1 and VCAM-1 as detected on HUVEC by immunofluorescence using mAB9G11, 11C81 and 4B2 (R&D Systems, Europe Ltd.).

[0048] PRBC adhesion assays were performed on transfected CHO cells and fresh cryo-sections of human placenta as described in Gysin, J., et al., *Mol Biochem Parasitol* 88, 267-71 (1997). Accordingly, adhesion of plasmagel enriched PRBCs to various receptors coated on plastic was achieved by immobilizing 10 μl of receptor in PBS directly on Petri dishes (Falcon 1001) overnight at 4° C. The receptor concentration used included recombinant human thrombomodulin^{CSA} (hTM) (5 μg/ml), CSA (10 μg/ml, Sigma), CSC (10 g/ml, Sigma), Biot-CSA and Biot-CSC (100 μg/ml). The coated dots were blocked with 1% BSA and incubated with 10 μl of trophozoites (0.5% hematocrit) in binding medium (BM) (RPMI-1640 with 25 mM HEPES, pH 6.8) for 20 min at 37° C. Unaffixed cells were removed by washes in BM and the cells that remained joined to the plastic were fixed with 2% glutaraldehyde and stained in Giemsa for microscopic examination. Once it was understood that FCR3.varCSA mediated adhesion to CSA, domains of FCR3.varCSA were cloned into cells that do not express CSA so as to elucidate the regions of the molecule that mediate binding to CSA, as described in the next section.

[0049] The DBL-3 Domain of FCR3.varCSA Binds Specifically to CSA.

[0050] To identify domains of FCR3.varCSA that were involved in binding to CSA, PCR products spanning each single domain were cloned into the expression vector pSRA5 and were transfected into CHO-745 cells (a cell line that does not natively express CSA). (Smith, et al., *Mol Biochem Parasitol* 97, 133-48 (1998)). Stable transfectants that expressed these FCR3.varCSA regions on the surface of CHO-745 cells were then selected on a Fluorescent-activated cell sorter (FACS), expanded and were used for adhesion studies that employed techniques similar to those described in Smith, et al., *Mol Biochem Parasitol* 97: 133-48 (1998).

[0051] In preparation for these adhesion studies, biotinylated-CSA and biotinylated-CSC were developed as reagents to measure CSA binding to the FCR3.varCSA domains. The activity of the biotinylated compounds was identical to that of the non-biotinylated material (TABLE 1) in that PRBCs bound only to biotin-CSA. Conjugation of biotin to chondroitin sulfate A (Bovine Trachea, Sigma St. Louis) and chondroitin sulfate C (shark cartilage, Sigma St. Louis) was accomplished by an improvement of the method described by Shinohara, et al., *J Biochem (Tokyo)* 117: 1076-82 (1995). Briefly, Biotinyl-(aminocaproyl)₃-hydrazide was synthesized by Fmoc-based solid phase peptide synthesis. 0.71 mmole of P-alkoxybenzylalcohol-Wang resin (0.71 mmole/g 100-200 mesh, Watanabe Chemical Co., Japan) was treated with p-nitrophenylchloroformate (3.55 mmole, Seq) and pyridine (7.1 mmole, 10 eq) in CHCl₃ over night at room temperature. The resin was washed with CHCl₃ (6 times) and washed with dimethylformamide (DMF) (6 times). The resin in DMF was treated with NH₂NH₂.hydrate (7.1 mmole, 10eq) by shaking for 3 hours at room temperature and washed with DMF 6 times. The resulting hydrazinated resin in DMF was acylated with Fmoc-aminocaproic acid (3eq) 1-hydroxybenzotriazole (3 eq), and diisopropylcarbodiimide (3 eq) for coupling. 20% piperidine-DMF was used for deprotection and the reaction repeated twice more with anicaproic acid, followed by (+)-Biotin (Wako Pure Chemical Industry, Japan). The protected peptide resin was treated with TFA-m-cresol-ethanedithiol (9:0.5:0.5 cocktail) for deprotection and cleavage, purified by HPLC in 0.1% MeCNaq and characterized by mass-spectrometry. Conjugation between the biotinyl-(aminocaproyl)₃-hydrazide and chondroitin sulfates was carried out by reductive hydrazination of chondroitin sulfate via terminal aldehyde with NaBH₃(CN) in 1N AcOH at room temperature. This procedure gave a stable covalent conjugation. The conjugates were purified by gel filtration on Sephadex G-15 in 0.1N AcOH, analyzed by HPLC and the incorporation ratio was determined by combustive amino acid analysis as follows. Biotinyl-(aminocaproyl)₃-NHNH-chondroitin sulfate was hydrolyzed with 6 N HCL containing 0.2% phenol at 110° C. for 24 hours and an aliquot of the resulting hydrolysates was subjected to amino acid analyzer (Hitachi 835 A) equipped with Chromato-Integrator (Hitachi D-2500). An unknown peak corresponding to the same retention time of aminocaproic acid was observed with CSA. Thus, the molar ratio of conjugation was calculated from the difference between the biotin-CSA to CSA alone. The incorporation was only 15% to 25%.

[0052] Biot-CSA, Biot-CSC, or soluble CD36 were immobilized via mouse monoclonal antibodies (5 μg/ml) directed either against biotin (Sigma, St. Louis) or against an epitope tag incorporated into a recombinant CD36 molecule (mAb 179). (Smith, et al., *Mol Biochem Parasitol* 97, 133-48 (1998)). The adhesion assays employing biotinylated CSA and CSC were performed with approximately, 2x10⁶ sheep anti-mouse IgG M-450 Dynabeads were incubated overnight at 5° C. with 2 μg of mouse anti-biotin monoclonal antibody (Jackson. Immunoresearch Labs, West Grove, Pa.) in PBS with continuous agitation. The beads were washed 3 times with BM pH 7.2+1% BSA (BMB) and resuspended with 45 μl of BMB to 4x10⁷ beads/ml. Approximately, 100,000 transfected CHO-745 cells were grown for 48 h. on 4 glass cover slips in six wells plates. Cover slips were

transferred into a 12 wells plate containing 1 ml of BMB and 50 μ g of Biot-CSA or Biot-CSC (Sigma) and incubated 1 hour.

[0053] For inhibition assays, the cells were incubated for 1 hour with 200 μ g/ml of CSA or CSC (Sigma) before addition of the Biotin conjugated carbohydrates. The coverslips were washed 3 times in a basin containing BMB, transferred to a humidified chamber and incubated, 1 hour room temperature, with the coated Dynal beads (45 μ l of 4×10^6 beads 1 ml). The coverslips were then flipped cell-side down onto a stand and incubated for 3 minutes to allow unbound beads to settle by gravity. Coverslips were then washed 3 times with BMB, fixed with 2% paraformaldehyde (Polysciences) in PBS and the degree of bead associated with cells was examined. In some experiments, chondroitinase ABC (Fluka, Ronkonkoma N.Y.) at 1 U/ml was added to the cells, 1 h room temperature, prior to the addition of beads.

[0054] Of the eight receptor-like domains expressed on the surface of CHO-745 cells (a mutant cell line that does not express CSA, see **FIG. 1C**), DBL3 and DBL7 transfectants were found to bind biotin-CSA (**FIG. 2A**). No binding was observed when cells were incubated with biotin alone or when cells were treated with chondroitinase ABC after incubation with Biotin-CSA. Further, competition with CSA blocked the binding of biotin-CSA to both DBLs but competition with CSC, a molecule that does not block CSA-mediated PRBCs adhesion (Rogerson, et al., *J Exp Med* 182, 15-20 (1995) and (Robert, et al., *Res in Immunol* 146, 383-93 (1995)), had no effect on binding to DBL3 but did block adhesion of DBL7 (**FIG. 2B**). Still further, DBL7 expressed on CHO cells bound biotin-CSC, and this binding was inhibited by addition of CSA or CSC (**FIG. 2B**). Thus, the binding properties of DBL3, and not of DBL7, are compatible with the properties exhibited by CSA-adherent PRBCs (TABLE 1).

[0055] A previous study demonstrated that antibodies directed to two domains of a var gene (varCS2) from CSA-PRBCs reduced binding to CSA (Reeder, et al., *Proc Natl Acad Sci USA* 96, 5198-202 (1999)). However, the identity of a parasite CSA-binding ligand molecule was not known until this disclosure. It has been determined by direct binding studies that the FCR3.varCSA gene domain DBL-3 binds CSA. The DBL3 sequences described in Reeder, et al., *Proc Natl Acad Sci USA* 96, 5198-202 (1999) and the DBL3 sequences described herein share no specific homology other than the homology found among all DBL3 domains, and notably these regions of homology can be found in PRBCs that do not bind CSA. (Smith, et al., *Mol Biochem Parasitol* 97, 133-48 (1998)). The same is true for the CIDR1 domain. The CIDR1 domain of the FCR3-CSA var did not bind CD36, which is in full agreement with the failure of the PRBCs to bind CD36. This is in distinction from other CIDR1 domains from CD36-binding PRBCs that bind CD36. (Baruch, et al., *Blood* 90, 3766-75 (1997) and (Smith, et al., *Mol Biochem Parasitol* 97, 133-48 (1998)). Thus, the adhesion properties of a particular var protein or var domain cannot be predicted from its primary sequence.

[0056] Identification of the domain that binds CSA provides the molecular complement to the Fried and Duffy model of maternal malaria. (Fried & Duffy, *Science* 272, 1502-1504 (1996) and (Fried, et al., *Nature* 395, 851-2

(1998)). Accordingly, at the age of first pregnancy, most residents of endemic areas are clinically immune and develop a repertoire of anti-PfEMP1 antibodies against endothelial adherent PRBCs (CD36 binding PRBCs), but not to the CSA-binding placental adherent PRBCs. Primi-gravid women who do not yet display antibodies against the CSA binding ligand offer a new niche for sequestration and proliferation of those parasites. The findings disclosed herein establish that antigenic variation of PfEMP1, besides its role in immune evasion, contributes to drastic changes in parasite tropism. A switch to a PfEMP1 that mediates CSA adhesion is a significant molecular event involved in the disease process observed during the first pregnancy. The data published by Fried and Duffy (Fried & Duffy, *Science* 272, 1502-1504 (1996) and (Fried, et al., *Nature* 395, 851-2 (1998)) demonstrate that antibodies from multigravid women block binding of PRBCs from placenta to CSA. This blockade of adhesion is not specific for a particular clone, as sera from multigravid females block not only PRBCs from Africa but also PRBCs from other parts of the world. Although CSA-binding PfEMP1s vary in primary sequence, a conserved three-dimensional structure or conserved antigenic determinants among various CSA-adherent strains can exist.

[0057] To test this hypothesis, several adhesion assays were conducted using polypeptide fragments of other varCSA proteins. (See TABLE 2). As above, nucleic acids encoding each polypeptide were cloned into an expression vector and were transfected into CHO-745 cells. Stable transfectants that expressed the varCSA polypeptides on the surface of CHO-745 cells were then selected on a Fluorescent-activated cell sorter (FACS), expanded and were used for adhesion studies that employed techniques similar to those described in Smith, et al., *Mol Biochem Parasitol* 97: 133-48 (1998). The adhesion assays employed biotinylated CSA and CSC and were performed as described above. The results of these assays are provided in TABLE 2.

TABLE 2

varCSA polypeptide	Binding to CSA
R29 DBL2- γ (SEQ. ID. No.: 7)	-
A4 DBL4- γ (SEQ. ID. No.: 8)	-
A4 tres DBL3- γ (SEQ. ID. No.: 9)	++++
FCR3 var3 DBL- γ (SEQ. ID. No.: 10)	-
ItG2-CS2 DBL2- γ (SEQ. ID. No.: 11)	++++

[0058] These assays verified that some varCSA proteins and fragments thereof are able to bind while others do not. Further, these results provided evidence that varCSA proteins or fragments thereof that are able to bind CSA have similarity in structure despite differences in primary sequence. The section below provides several software and hardware embodiments of the invention, as well as, computational methods that can be used to further characterize a varCSA nucleic acid sequence and a varCSA polypeptide sequence, as well as, identify agents that inhibit varCSA-mediated adhesion to CSA.

[0059] Software and Hardware Embodiments

[0060] The FCR3.varCSA nucleic acid sequence and the FCR3.varCSA protein sequence were entered onto a computer readable medium for recording and manipulation. It

will be appreciated by those skilled in the art that a computer readable medium having the FCR3.varCSA nucleic acid sequence or the FCR3.varCSA protein sequence or both is useful for the determination of homologous sequences, structural and functional domains, and the construction of protein models for rational drug design. The functionality of a computer readable medium having the FCR3.varCSA nucleic acid sequence or the FCR3.varCSA protein sequence or both includes the ability to compare the sequence to others stored on databases, to ascertain structural and functional information, to develop protein models, and to conduct rational drug design.

[0061] The FCR3.varCSA nucleic acid sequence or the FCR3.varCSA protein sequence or both can be stored, recorded, and manipulated on any medium that can be read and accessed by a computer. As used herein, the words "recorded" and "stored" refer to a process for storing information on computer readable medium. A skilled artisan can readily adopt any of the presently known methods for recording information on computer readable medium to generate manufactures comprising the nucleotide or polypeptide sequence information of this embodiment.

[0062] A variety of data storage structures are available to a skilled artisan for creating a computer readable medium having recorded thereon a nucleotide or polypeptide sequence. The choice of the data storage structure will generally be based on the component chosen to access the stored information. Computer readable media include magnetically readable media, optically readable media, or electronically readable media. For example, the computer readable media can be a hard disc, a floppy disc, a magnetic tape, zip disk, CD-ROM, DVD-ROM, RAM, or ROM as well as other types of other media known to those skilled in the art. The computer readable media on which the sequence information is stored can be in a personal computer, a network, a server or other computer systems known to those skilled in the art.

[0063] Embodiments include systems, particularly computer-based systems that contain the sequence information described herein. The term "a computer-based system" refers to the hardware, software, and any database used to analyze the FCR3.varCSA nucleic acid sequence or the FCR3.varCSA protein sequence or both, or fragments of these biomolecules. The computer-based system preferably includes the storage media described above, and a processor for accessing and manipulating the sequence data. The hardware of the computer-based systems of this embodiment comprise a central processing unit (CPU) and a data database. A skilled artisan can readily appreciate that any one of the currently available computer-based systems are suitable.

[0064] In one particular embodiment, the computer system includes a processor connected to a bus that is connected to a main memory (preferably implemented as RAM) and a variety of secondary storage devices, such as a hard drive and removable medium storage device. The removable medium storage device may represent, for example, a floppy disk drive, a DVD drive, an optical disk drive, a compact disk drive, a magnetic tape drive, etc. A removable storage medium, such as a floppy disk, a compact disk, a magnetic tape, etc. containing control logic and/or data recorded therein (e.g., the FCR3.varCSA nucleic acid sequence or the FCR3.varCSA protein sequence or both or fragments

thereof) can be inserted into the removable storage device. The computer system includes appropriate software for reading the control logic and/or the data from the removable medium storage device once inserted in the removable medium storage device.

[0065] The FCR3.varCSA nucleic acid sequence or the FCR3.varCSA protein sequence or both can be stored in a well known manner in the main memory, any of the secondary storage devices, and/or a removable storage medium. Software for accessing and processing the FCR3.varCSA nucleic acid sequence or the FCR3.varCSA protein sequence or both (such as search tools, compare tools, and modeling tools etc.) reside in main memory during execution.

[0066] As used herein, "a database" refers to memory that can store nucleotide or polypeptide sequence information, protein model information, information on other peptides, chemicals, peptidomimetics, and other agents that interact with proteins, and values or results from varCSA characterization assays. Additionally, a "database" refers to a memory access component that can access manufactures having recorded thereon nucleotide or polypeptide sequence information, protein model information, information on other peptides, chemicals, peptidomimetics, and other agents that interact with proteins, and values or results from varCSA characterization assays. In other embodiments, a database stores a varCSA disease-state profile comprising concentrations or expression levels or concentration ranges or expression level ranges of FCR3.varCSA or FCR3.varCSA or both, for example, detected in biological samples from different subjects (e.g., subjects with and without a disease related to FCR3.varCSA). In more embodiments, a database stores a FCR3.varCSA disease-state profile comprising concentration ranges or levels of FCR3.varCSA detected in biological samples obtained from various tissue or fluid sources from diseased and healthy subjects. Many databases are known to those of skill in the art and several will be discussed below.

[0067] The sequence data on FCR3.varCSA or FCR3.varCSA or both can be stored and manipulated in a variety of data processor programs in a variety of formats. For example, the sequence data can be stored as text in a word processing file, such as MicrosoftWORD or WORDPERFECT, an ASCII file, a html file, or a pdf file in a variety of database programs familiar to those of skill in the art, such as DB2, SYBASE, or ORACLE.

[0068] A "search program" refers to one or more programs that are implemented on the computer-based system to compare a nucleotide or polypeptide sequence with other nucleotide or polypeptide sequences and agents including but not limited to peptides, peptidomimetics, and chemicals stored within a database. A search program also refers to one or more programs that compare one or more protein models to several protein models that exist in a database and one or more protein models to several peptides, peptidomimetics, and chemicals that exist in a database. A search program is used, for example, to compare regions of the FCR3.varCSA nucleic acid sequence or the FCR3.varCSA protein sequence or both that match sequences in nucleic acid and protein data bases so as to identify homologies and structural or functional motifs. Further, a search program is used to compare an unknown nucleic acid or protein sequence with the FCR3.varCSA nucleic acid sequence or the FCR3.varCSA protein sequence so as to identify homologies and related

structural or functional domains. Additionally, a search program is used to compare a FCR3.varCSA-disease state profile from a tested subject to FCR3.varCSA-disease state profiles from diseased and healthy subjects present in a database. Still further, a search program is used to compare values or results from FCR3.varCSA characterization assays.

[0069] A "retrieval program" refers to one or more programs that are implemented on the computer based system to identify a homologous nucleic acid sequence, a homologous protein sequence, or a homologous protein model. A retrieval program is also used to identify peptides, peptidomimetics, and chemicals that interact with a nucleic acid sequence, a protein sequence, or a protein model stored in a database. Further a retrieval program is used to identify a disease state of an individual by obtaining a FCR3.varCSA disease-state profile from the database that matches the FCR3.varCSA-disease state profile from the tested subject. Additionally, a retrieval program is used to obtain "a FCR3.varCSA-agent profile" that can be composed of a nucleic acid or polypeptide sequence or model thereof or one or more symbols that represent these sequences and/or models, an identifier that represents one or more FCR3.varCSA modulating agents, and a value or result from a FCR3.varCSA characterization assay. The discussion below describes embodiments of the invention having nucleic acids that encode FCR3.varCSA.

[0070] Use of Nucleic Acids Encoding FCR3. varCSA or Fragments of FCR3. varCSA

[0071] The cDNA sequence encoding FCR3.varCSA is provided in the sequence listing (SEQ. ID NO.: 1). Full-length FCR3.varCSA and fragments of FCR3.varCSA (e.g., nucleic acids encoding DBL3 and/or CIDR1) are embodiments of the invention. Further embodiments include nucleic acids that complement full-length FCR3.varCSA and nucleic acids that complement fragments of FCR3.varCSA (e.g., nucleic acids encoding DBL3 and/or CIDR1) and other nucleic acids that encode a polypeptide that binds to CSA. Desired embodiments include nucleic acids having at least 9 consecutive bases of FCR3.varCSA or a sequence complementary thereto. Preferred embodiments include a nucleic acid that encodes a polypeptide that binds to CSA or a nucleic acid that complements a nucleic acid that encodes a polypeptide that binds to CSA.

[0072] The nucleic acid embodiments of the invention can have from 9 to 10,628 consecutive nucleotides in length that encode a fragment of FCR3.varCSA or full-length FCR3.varCSA or a complementary nucleic acid, whose complement encodes a fragment of FCR3.varCSA or full-length FCR3.varCSA. However, one of skill in the art will appreciate that FCR3.varCSA nucleic acids can be joined to an exogenous nucleic acid so as to create a nucleic acid embodiment having virtually any length. Thus, a nucleic acid having a portion (9 to 10,627 consecutive nucleotides) or full-length FCR3.varCSA are embodiments of the invention. That is, a nucleic acid having less than or equal to 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 125, 150, 175, 200, 250,

300, 350, 400, 500, 600, 700, 800, 900, 1000, 1500, 2000, 3000, 4000, 5000, 6000, 7000, 8000, 9000, 10,000, 10,500, and 10,628 nucleotides are embodied. Preferably, the nucleic acid embodiments, however, comprise at least 12, 13, 14, 15, 16, 17, 18, or 19 consecutive nucleotides from FCR3.varCSA or a nucleic acid that complements FCR3.varCSA, as conditions dictate. Nucleic acid embodiments that comprise a fragment of FCR3.varCSA (e.g., nucleic acids encoding DBL3 and/or CIDR1) or a complement thereof can be determined by referring to the sequences provided in SEQ. ID. Nos.: 1 and 2 and FIG. 1.

[0073] More preferably, the nucleic acid embodiments comprise at least 20-30 consecutive nucleotides from FCR3.varCSA or a nucleic acid that complements FCR3.varCSA. In some cases, the nucleic acid embodiments comprise more than 30 nucleotides from the nucleic acids encoding FCR3.varCSA or a nucleic acid that complements FCR3.varCSA and in other cases, the nucleic acid embodiments comprise at least 40, at least 50, at least 75, at least 100, at least 150, or at least 200 consecutive nucleotides from the nucleic acids encoding FCR3.varCSA or a nucleic acid that complements FCR3.varCSA. These nucleic acid oligomers have biotechnological and diagnostic use, e.g., in nucleotide acid hybridization assays, Southern and Northern Blot analysis, etc. and the prognosis of FCR3.varCSA-related diseases.

[0074] Some embodiments comprise recombinant nucleic acids having all or part of the FCR3.varCSA gene or recombinant nucleic acids that complement all or part of FCR3.varCSA. Desirable embodiments comprise full-length FCR3.varCSA and fragments of FCR3.varCSA that encode a polypeptide that binds to CSA and nucleic acids that complement full-length FCR3.varCSA and fragments of FCR3.varCSA that encode a polypeptide that binds to CSA. A recombinant construct can be capable of replicating autonomously in a host cell. Alternatively, the recombinant construct can become integrated into the chromosomal DNA of the host cell. Such a recombinant polynucleotide comprises a polynucleotide of genomic or cDNA, of semi-synthetic or synthetic origin by virtue of human manipulation. Therefore, recombinant nucleic acids comprising sequences otherwise not naturally occurring are provided by embodiments of this invention. Although nucleic acids encoding FCR3.varCSA or nucleic acids having sequences that complement FCR3.varCSA as they appear in nature can be employed, they will often be altered, e.g., by deletion, substitution, or insertion and will be accompanied by sequence not present in humans.

[0075] The nucleic acid embodiments can be altered by mutation such as substitutions, additions, or deletions that provide for sequences encoding functionally equivalent molecules. Due to the degeneracy of nucleotide coding sequences, other DNA sequences that encode substantially the same FCR3.varCSA amino acid sequence as depicted in SEQ. ID NO.: 2 can be used in some embodiments of the present invention. These include, but are not limited to, nucleic acid sequences comprising all or portions of FCR3.varCSA or nucleic acids that complement all or part of FCR3.varCSA that have been altered by the substitution of different codons that encode a functionally equivalent amino acid residue within the sequence, thus producing a silent change.

[0076] In addition, recombinant FCR3.varCSA-encoding nucleic acid sequences and their complementary sequences can be engineered so as to modify processing or expression of FCR3.varCSA. For example, and not by way of limitation, the FCR3.varCSA gene can be combined with a promoter sequence and/or ribosome binding site, or a signal sequence can be inserted upstream of FCR3.varCSA-encoding sequences to permit secretion of FCR3.varCSA and thereby facilitate harvesting or bioavailability. Additionally, a given FCR3.varCSA nucleic acid can be mutated in vitro or in vivo, to create and/or destroy translation, initiation, and/or termination sequences, or to create variations in coding regions and/or form new restriction sites or destroy preexisting ones, or to facilitate further in vitro modification. Any technique for mutagenesis known in the art can be used, including but not limited to, in vitro site-directed mutagenesis. (Hutchinson et al., J. Biol. Chem. 253:6551 (1978)). Further, nucleic acids encoding other proteins or domains of other proteins can be joined to nucleic acids encoding FCR3.varCSA so as to create a fusion protein. The resulting fusion proteins are used as biotechnological tools or pharmaceuticals or both, as will be discussed below.

[0077] The nucleic acid embodiments can also be used as biotechnological tools for isolation procedures and diagnostic assays. By using the FCR3.varCSA nucleic acid sequence disclosed in the sequence listing (SEQ ID NO.: 1), probes that complement FCR3.varCSA can be designed and manufactured by oligonucleotide synthesis. These probes can be used to screen cDNA or genomic libraries so as to isolate natural sources of the nucleic acid embodiments of the present invention. Additionally, these probes can be used to isolate other nucleotide sequences capable of hybridizing to them. Further, sequences from nucleic acids complementing FCR3.varCSA, or portions thereof can be used to make oligonucleotide primers by conventional oligonucleotide synthesis for use in isolation and diagnostic procedures. These oligonucleotide primers can be used, for example, to isolate the nucleic acid embodiments of this invention by amplifying the sequences resident in genomic DNA or other natural sources by using the Polymerase Chain Reaction (PCR) or other nucleic acid amplification techniques. Further, the nucleic acid embodiments of the invention can be used to modulate FCR3.varCSA-mediated adhesion to CSA (e.g., by upregulating or downregulating the expression of FCR3.varCSA) and, therefore, have several uses in addition to biotechnological research including therapeutic and prophylactic applications, as will be discussed below. Alternatively, the nucleic acids encoding FCR3.varCSA or fragments thereof are manipulated using conventional techniques in molecular biology to create recombinant constructs that express FCR3.varCSA or fragments of FCR3.varCSA.

[0078] Embodiments also include nucleic acids encoding polypeptides that comprise A4 tres DBL3- γ (SEQ. ID. No.: 9) and ItG2-CS2 DBL2- γ (SEQ. ID. No.: 11) sequence or complements thereto or fragments thereof. These nucleic acid embodiments can be for example, less than or equal to 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 125, 150, 175, 200, 250, 300, 350, 400, 500, 600, 700, 800, 900, 1000, and

1050 nucleotides in length so long as the nucleic acid can bind CSA. As with the other nucleic acid embodiments described herein, the nucleic acids encoding polypeptides that comprise A4 tres DBL3- γ (SEQ. ID. No.: 9) and ItG2-CS2 DBL2- γ (SEQ. ID. No.: 11) sequence or complements thereto or fragments thereof can be incorporated into vectors, plasmids, expression constructs and organisms, including humans. The discussion that follows describes some of the expression constructs and protein embodiments of the invention.

[0079] FCR3.varCSA Polypeptides and Fragments of FCR3.varCSA

[0080] The FCR3.varCSA polypeptides or derivatives thereof, include but are not limited to, those containing as a primary amino acid sequence all of the amino acid sequence substantially as depicted in the sequence listing (SEQ. ID NO.: 2) and fragments of SEQ. ID. NO.: 2 at least three amino acids in length. Preferred polypeptide embodiments include domains of FCR3.varCSA (e.g., DBL3 and/or CIDR1), including altered sequences in which functionally equivalent amino acid residues are substituted for residues within the sequence resulting in a silent change. The sequence of these domains or fragments thereof can be determined by referring to SEQ. ID. No. 2 and FIG. 1.

[0081] Additionally, one or more amino acid residues within the FCR3.varCSA polypeptide of SEQ ID. NO.: 2 and fragments of SEQ. ID. NO.: 2 that comprise an amino acid sequence found in a peptide that binds CSA can be substituted by another amino acid of a similar polarity that acts as a functional equivalent, resulting in a silent alteration. Substitutes for an amino acid within the sequence can be selected from other members of the class to which the amino acid belongs. For example, the non-polar (hydrophobic) amino acids include alanine, leucine, isoleucine, valine, proline, phenylalanine, tryptophan, and methionine. The polar neutral amino acids include glycine, serine, threonine, cysteine, tyrosine, asparagine and glutamine. The positively charged (basic) amino acids include arginine, lysine, and histidine. The negatively charged (acidic) amino acids include aspartic acid and glutamic acid. The aromatic amino acids include phenylalanine, tryptophan, and tyrosine.

[0082] The FCR3.varCSA fragments can be less than or equal to 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 125, 150, 175, 200, 250, 300, 350, 400, 500, 600, 700, 800, 900, 1000, 1500, 2000, 2,500, 3000, 3,500, or 3,542 amino acids in length. One embodiment, for example, comprises a polypeptide fragment having the sequence EAEKELKEGKIPGFKRQMFYTFG DYRDILFG (SEQ. ID. NO.: 3). Desirable polypeptide embodiments comprise the sequence KELKEGKIPE (SEQ. ID. NO.: 4). Preferred polypeptide embodiments comprise the sequence KEGK (SEQ. ID. NO.: 5) and, more preferably, polypeptide embodiments comprise the sequence KX_1GX_2 (SEQ. ID. NO.: 6), wherein X_1 and X_2 are any amino acid. In other aspects of the invention, the FCR3.varCSA polypeptide of SEQ ID. NO.: 2 and fragments of SEQ. ID. NO.: 2 that comprise an amino acid sequence that binds to CSA, or

derivatives thereof are differentially modified during or after translation, e.g., by phosphorylation, glycosylation, cross-linking, acylation, proteolytic cleavage, linkage to an antibody molecule, membrane molecule, or other ligand. (Ferguson et al., *Ann. Rev. Biochem.* 57:285-320 (1988)).

[0083] Other embodiments include polypeptides that have homology to FCR3.varCSA and bind to CSA. By "homology to FCR3.varCSA" is meant either protein sequence homology or three-dimensional homology. As will be discussed below, several techniques exist to determine protein sequence homology and/or three-dimensional homology. These methods are routinely employed to discover related sequences and novel ligands, as well as, determine the extent of homology that one sequence, domain, or model has to a target sequence, domain, or model. Because the region of FCR3.varCSA (e.g., a region within a DBL3 domain) that mediates CSA adhesion can be quite small (e.g., 3, 4, 5, 6, 7, 8, 9, 10, 12, 15, 18, 20, 22, 25, 30 amino acids in length) embodiments of the invention can exhibit a vast degree of homology to full-length FCR3.varCSA. For example, a fusion protein having a small region of FCR3.varCSA can exhibit a low degree of overall homology to FCR3.varCSA yet retain the ability to bind CSA. Thus, embodiments of the invention can have from 1% homology to 100% homology to full-length FCR3.varCSA. That is, embodiments can have 1.0%, 2.0%, 3.0%, 4.0%, 5.0%, 6.0%, 7.0%, 8.0%, 9.0%, 10.0%, 11.0%, 12.0%, 13.0%, 14.0%, 15.0%, 16.0%, 17.0%, 18.0%, 19.0%, 20.0%, 21.0%, 22.0%, 23.0%, 24.0%, 25.0%, 26.0%, 27.0%, 28.0%, 29.0%, 30.0%, 31.0%, 32.0%, 33.0%, 34.0%, 35.0%, 36.0%, 37.0%, 38.0%, 39.0%, 40.0%, 41.0%, 42.0%, 43.0%, 44.0%, 45.0%, 46.0%, 47.0%, 48.0%, 49.0%, 50.0%, 51.0%, 52.0%, 53.0%, 54.0%, 55.0%, 56.0%, 57.0%, 58.0%, 59.0%, 60.0%, 61.0%, 62.0%, 63.0%, 64.0%, 65.0%, 66.0%, 67.0%, 68.0%, 69.0%, 70.0%, 71.0%, 72.0%, 73.0%, 74.0%, 75.0%, 76.0%, 77.0%, 78.0%, 79.0%, 80.0%, 81.0%, 82.0%, 83.0%, 84.0%, 85.0%, 86.0%, 87.0%, 88.0%, 89.0%, 90.0%, 91.0%, 92.0%, 93.0%, 94.0%, 95.0%, 96.0%, 97.0%, 98.0%, 99.0%, and 100.0% homology to FCR3.varCSA. Therefore, embodiments of the invention include polypeptides varying in size from 3 amino acids up to and including the full-length FCR3.varCSA protein that have 1%-100% homology to FCR3.varCSA and exhibit the ability to bind to CSA.

[0084] Preferred embodiments also include polypeptides that comprise A4 tres DBL3- γ (SEQ. ID. No.: 9) and ItG2-CS2 DBL2- γ (SEQ. ID. No.: 11) sequence. These embodiments can be for example, less than or equal to 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 125, 150, 175, 200, 250, 300, 348 amino acids in length so long as the peptide can bind CSA. As above, one or more amino acid residues within the polypeptide sequence of SEQ ID. NO.: 9 and/or 11 and fragments of these molecules that comprise an amino acid sequence found in a peptide that binds CSA can be substituted by another amino acid of a similar polarity that acts as a functional equivalent, resulting in a silent alteration.

[0085] In several embodiments, the FCR3.varCSA polypeptide (SEQ ID. NO.: 2) and polypeptides comprising

the A4 tres DBL3- γ (SEQ. ID. No.: 9) and the ItG2-CS2 DBL2- γ (SEQ. ID. No.: 11) sequence and fragments of these molecules are expressed in a cell line. The term "isolated" requires that the material be removed from its original environment (e.g., the natural environment if it is naturally occurring). For example, a naturally occurring nucleic acid or protein present in a living cell is not isolated, but the same nucleic acid or protein, separated from some or all of the coexisting materials in the natural system, is isolated. In accordance with this definition, FCR3.varCSA nucleic acid or FCR3.varCSA protein or nucleic acid or polypeptide fragments present in a cell lysate are "isolated". The term "purified" does not require absolute purity; rather it is intended as a relative definition. For example, recombinant nucleic acids and proteins are routinely purified to electrophoretic homogeneity, as detected by ethidium bromide staining or Coomassie staining, and are suitable in several assays despite having the presence of contaminants.

[0086] To express the protein embodiments described herein, nucleic acids containing the coding sequence for these molecules are obtained and cloned into a suitable expression vector such that the coding region is operably linked to a heterologous promoter. The nucleic acid encoding the protein or polypeptide to be expressed is operably linked to a promoter in an expression vector using conventional cloning technology. The expression vector can be in any of the mammalian, yeast, amphibian, insect, parasite, or bacterial expression systems known in the art. Commercially available vectors and expression systems are available from a variety of suppliers including Genetics Institute (Cambridge, Mass.), Stratagene (La Jolla, Calif.), Promega Madison, Wis.), and Invitrogen (San Diego, Calif.). If desired, to enhance expression and facilitate proper protein folding, the codon context and codon pairing of the sequence can be optimized for the particular expression organism in which the expression vector is introduced, as explained by Hatfield, et al., U.S. Pat. No. 5,082,767. Further, a secretory leader sequence can be incorporated so as to facilitate purification of the protein.

[0087] The following is provided as one exemplary method to express the proteins encoded by the nucleic acids described above. First, the methionine initiation codon for the gene and the poly A signal of the gene are identified. If the nucleic acid encoding the polypeptide to be expressed lacks a methionine to serve as the initiation site, an initiating methionine can be introduced next to the first codon of the nucleic acid using conventional techniques. Similarly, if the nucleic acid lacks a poly A signal, this sequence can be added to the construct by, for example, splicing out the Poly A signal from pSG5 (Stratagene) using BglI and Sall restriction endonuclease enzymes and incorporating it into the mammalian expression vector pXT1 (Stratagene). The vector pXT1 contains the LTRs and a portion of the gag gene from Moloney Murine Leukemia Virus. The position of the LTRs in the construct allow efficient stable transfection. The vector includes the Herpes Simplex Thymidine Kinase promoter and the selectable neomycin gene.

[0088] The nucleic acid encoding the polypeptide to be expressed can be obtained by PCR from the bacterial vector using oligonucleotide primers complementary to the nucleic acid and containing restriction endonuclease sequences for Pst I incorporated into the 5' primer and BglII at the 5' end of the corresponding cDNA 3' primer, taking care to ensure

that the nucleic acid is positioned in frame with the poly A signal. The purified fragment obtained from the resulting PCR reaction is digested with PstI, blunt ended with an exonuclease, digested with Bgl II, purified and ligated to pXT1, now containing a poly A signal and digested with BglIII. The ligated product is transfected into a suitable cell line, e.g., mouse NIH 3T3 cells, using Lipofectin (Life Technologies, Inc., Grand Island, N.Y.) under conditions outlined in the product specification. Positive transfectants are selected after growing the transfected cells in 600 ug/ml G418 (Sigma, St. Louis, Mo.). Preferably the expressed protein is released into the culture medium, thereby facilitating purification.

[0089] Another embodiment utilizes the "Xpress system for expression and purification" (Invitrogen, San Diego, Calif.). The Xpress system is designed for high-level production and purification of recombinant proteins from bacterial, mammalian, and insect cells. The Xpress vectors produce recombinant proteins fused to a short N-terminal leader peptide that has a high affinity for divalent cations. Using a nickel-chelating resin (Invitrogen), the recombinant protein can be purified in one step and the leader can be subsequently removed by cleavage with enterokinase.

[0090] One preferred vector for the expression of FCR3.varCSA and fragments of FCR3.varCSA is the pBlue-BacHis2 Xpress. The pBlueBacHis2 Xpress vector is a Baculovirus expression vector containing a multiple cloning site, an ampicillin resistance gene, and a lac z gene. By one approach, the FCR3.varCSA nucleic acid, or portion thereof is cloned into the pBlueBacHis2 Xpress vector and SF9 cells are infected. The expression protein is then isolated or purified according to the manufacturer's instructions. Several other cultured cell lines having recombinant constructs or vectors comprising FCR3.varCSA or portions thereof are embodiments of the present invention and their manufacture would be routine given the present disclosure.

[0091] Proteins in the culture medium can also be separated by gel electrophoresis. The separated proteins are then detected using techniques such as Coomassie or silver staining or by using antibodies against the protein. Coomassie, silver staining, and immunolabeling of proteins are techniques familiar to those skilled in the art. If desired, the proteins can also be ammonium sulfate precipitated or separated based on size or charge prior to electrophoresis.

[0092] The protein embodiments described herein can also be purified using standard immunochromatography techniques. In such procedures, a solution containing the protein, such as the culture medium or a cell extract, is applied to a column having antibodies against the protein attached to the chromatography matrix. The protein is allowed to bind the immunochromatography column. Thereafter, the column is washed to remove non-specifically bound proteins. The specifically bound protein is then released from the column and recovered using standard techniques.

[0093] Further, nucleic acids encoding a protein embodiment or portion thereof can be incorporated into expression vectors designed for use in purification schemes employing chimeric polypeptides. In one such strategy, for example, the coding sequence of FCR3.varCSA or portion thereof is inserted in frame with the gene encoding the other half of the chimera. The other half of the chimera may be β -globin or a nickel binding polypeptide encoding sequence. A chroma-

tography matrix having antibody to β -globin or nickel attached thereto is then used to purify the chimeric protein. Protease cleavage sites can be engineered between the β -globin gene or the nickel binding polypeptide and the FCR3.varCSA cDNA such as enterokinase. Thus, the two polypeptides of the chimera can be separated from one another by protease digestion.

[0094] One useful expression vector for generating β -globin chimerics is pSG5 (Stratagene), which encodes rabbit β -globin. Intron II of the rabbit β -globin gene facilitates splicing of the expressed transcript, and the polyadenylation signal incorporated into the construct increases the level of expression. These techniques as described are well known to those skilled in the art of molecular biology. Standard methods are published in methods texts such as Davis et al., (*Basic Methods in Molecular Biology*, L. G. Davis, M. D. Dibner, and J. F. Battey, ed., Elsevier Press, NY, 1986) and many of the methods are available from Stratagene, Life Technologies, Inc., or Promega. Polypeptide may additionally be produced from the construct using in vitro translation systems, such as the In vitro Express™ Translation Kit (Stratagene).

[0095] In addition to isolating or purifying the protein embodiments by using recombinant DNA techniques, these molecules can be prepared by chemical synthesis methods (such as solid phase peptide synthesis) using methods known in the art such as those set forth by Merrifield et al., *J. Am. Chem. Soc.* 85:2149 (1964), Houghten et al., *Proc. Natl. Acad. Sci. USA*, 82:51:32 (1985), and Stewart and Young (solid phase peptide synthesis, Pierce Chem Co., Rockford, Ill. (1984)). Such polypeptides can be synthesized with or without a methionine on the amino terminus. Chemically synthesized FCR3.varCSA and fragments of FCR3.varCSA can be oxidized using methods set forth in these references to form disulfide bridges. FCR3.varCSA and fragments of FCR3.varCSA can be employed as biologically active or immunological substitutes for natural, purified FCR3.varCSA and fragments of FCR3.varCSA, for example. Further, peptidomimetics that structurally and/or functionally resemble a peptide embodiment (e.g., FCR3.varCSA or fragments of FCR3.varCSA) can be made and evaluated for their ability to interact with CSA in a characterization assay or to induce an immune response in a subject. Several approaches to make peptidomimetics that resemble polypeptides have been described. A vast number of methods, for example, can be found in U.S. Pat. Nos. 5,288,707; 5,552,534; 5,811,515; 5,817,626; 5,817,879; 5,821,231; and 5,874,529.

[0096] Following synthesis or expression and isolation or purification of a protein embodiment, the isolated or purified molecules can be used to generate antibodies and w tools for identifying agents that interact with a varCSA and fragments of a varCSA. Antibodies that recognize FCR3.varCSA and fragments of FCR3.varCSA (e.g., CIDR1 and/or DBL3), as well as A4 tres DBL3- γ (SEQ. ID. No.: 9) and ItG2-CS2 DBL2- γ (SEQ. ID. No.: 11) or fragments thereof, for example, have many uses including, but not limited to, biotechnological applications, therapeutic/prophylactic applications, and diagnostic applications. Such antibodies include, but are not limited to, polyclonal, monoclonal, chimeric, single chain, Fab fragments and fragments produced by a Fab expression library. Neutralizing antibodies, e.g., those that inhibit FCR3.varCSA-mediated adhesion or

formation of a complex having varCSA and CSA, are especially preferred for diagnostics and therapeutics.

[0097] For the production of antibodies, various hosts including goats, rabbits, rats, mice, etc can be immunized by injection with a protein embodiment that has immunogenic properties. Depending on the host species, various adjuvants can be used to increase immunological response. Such adjuvants include but are not limited to Freund's, mineral gels such as aluminum hydroxide, and surface active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, keyhole limpet hemocyanin, and dinitrophenol. BCG (*Bacillus Calmette-Guerin*) and *Corynebacterium parvum* are potentially useful adjuvants.

[0098] Peptides used to induce specific antibodies can have an amino acid sequence consisting of at least three amino acids, preferably at least 10 or 15 amino acids. Desirably, short stretches of amino acids encoding fragments of a varCSA molecule (e.g., FCR3.varCSA, A4 tres DBL3-65, or ItG2-CS2 DBL2- γ) are fused with those of another protein such as keyhole limpet hemocyanin and antibody is produced against the chimeric molecule. While antibodies capable of specifically recognizing a varCSA molecule, for example, can be generated by injecting into mice synthetic 3-mer, 10-mer, and 15-mer peptides that correspond to the particular protein sequence, a more diverse set of antibodies can be generated by using a recombinant or purified protein embodiment.

[0099] For example, to generate antibodies to FCR3.varCSA and fragments of FCR3.varCSA, substantially pure FCR3.varCSA or a fragment of FCR3.varCSA (e.g., DBL3, CIDR1, A4 tres DBL3- γ , or ItG2-CS2 DBL2- γ) is isolated from a transfected or transformed cell. The concentration of the polypeptide in the final preparation is adjusted, for example, by concentration on an Amicon filter device, to the level of a few micrograms/ml. Monoclonal or polyclonal antibody to the polypeptide of interest can then be prepared as follows:

[0100] Monoclonal antibodies to a varCSA protein or a fragment thereof can be prepared using any technique that provides for the production of antibody molecules by continuous cell lines in culture. These include but are not limited to the hybridoma technique originally described by Koehler and Milstein (*Nature* 256:495-497 (1975), the human B-cell hybridoma technique (Kosbor et al. *Immunol Today* 4:72 (1983); Cote et al. *Proc Natl Acad Sci* 80:2026-2030 (1983), and the EBV-hybridoma technique Cole et al. *Monoclonal Antibodies and Cancer Therapy*, Alan R. Liss Inc, New York N.Y., pp 77-96 (1985). In addition, techniques developed for the production of "chimeric antibodies", the splicing of mouse antibody genes to human antibody genes to obtain a molecule with appropriate antigen specificity and biological activity can be used. (Morrison et al. *Proc Natl Acad Sci* 81:6851-6855 (1984); Neuberger et al. *Nature* 312:604-608 (1984); and Takeda et al. *Nature* 314:452-454 (1985). Alternatively, techniques described for the production of single chain antibodies (U.S. Pat. No. 4,946,778) can be adapted to produce specific single chain antibodies. Antibodies can also be produced by inducing in vivo production in the lymphocyte population or by screening recombinant immunoglobulin libraries or panels of highly specific binding reagents as disclosed in Orlandi et al., *Proc Natl Acad Sci* 86:3833-3837 (1989), and Winter G. and Milstein C; *Nature* 349:293-299 (1991).

[0101] Antibody fragments that contain specific binding sites for FCR3.varCSA (e.g., DBL3, CIDR1, A4 tres DBL3- γ , or ItG2-CS2 DBL2- γ) can also be generated. For example, such fragments include, but are not limited to, the F(ab')₂ fragments that can be produced by pepsin digestion of the antibody molecule and the Fab fragments that can be generated by reducing the disulfide bridges of the F(ab')₂ fragments. Alternatively, Fab expression libraries can be constructed to allow rapid and easy identification of monoclonal Fab fragments with the desired specificity. (Huse W. D. et al. *Science* 256:1275-1281 (1989)).

[0102] By one approach, monoclonal antibodies are made as follows. Briefly, a mouse is repetitively inoculated with a few micrograms of the selected protein or peptides derived therefrom over a period of a few weeks. The mouse is then sacrificed, and the antibody producing cells of the spleen isolated. The spleen cells are fused in the presence of polyethylene glycol with mouse myeloma cells, and the excess unfused cells destroyed by growth of the system on selective media comprising aminopterin (HAT media). The successfully fused cells are diluted and aliquots of the dilution placed in wells of a microtiter plate where growth of the culture is continued. Antibody-producing clones are identified by detection of antibody in the supernatant fluid of the wells by immunoassay procedures, such as ELISA, as originally described by Engvall, E., *Meth. Enzymol.* 70:419 (1980) and derivative methods thereof. Selected positive clones can be expanded and their monoclonal antibody product harvested for use. Detailed procedures for monoclonal antibody production are described in Davis, L. et al. *Basic Methods in Molecular Biology* Elsevier, N.Y. Section 21-2.

[0103] Polyclonal antiserum containing antibodies to heterogeneous epitopes of a single protein can be prepared by immunizing suitable animals with the expressed protein or peptides derived therefrom described above, which can be unmodified or modified to enhance immunogenicity. Effective polyclonal antibody production is affected by many factors related both to the antigen and the host species. For example, small molecules tend to be less immunogenic than others and may require the use of carriers and adjuvant. Also, host animals vary in response to site of inoculations and dose, with both inadequate or excessive doses of antigen resulting in low titer antisera. Small doses (ng level) of antigen administered at multiple intradermal sites appears to be most reliable. An effective immunization protocol for rabbits can be found in Vaitukaitis, J. et al. *J. Clin. Endocrinol. Metab.* 33:988-991(1971).

[0104] Booster injections can be given at regular intervals, and antiserum harvested when antibody titer thereof, as determined semi-quantitatively, for example, by double immunodiffusion in agar against known concentrations of the antigen, begins to fall. See, for example, Ouchterlony, O. et al., Chap. 19 in: *Handbook of Experimental Immunology* D. Wier (ed) Blackwell (1973). Plateau concentration of antibody is usually in the range of 0.1 to 0.2 mg/ml of serum (about 12 μ M). Affinity of the antisera for the antigen is determined by preparing competitive binding curves, as described, for example, by Fisher, D., Chap. 42 in: *Manual of Clinical Immunology*, 2d Ed. (Rose and Friedman, Eds.) Amer. Soc. For Microbiol., Washington, D.C. (1980). Antibody preparations prepared according to either protocol are useful in quantitative immunoassays that determine concen-

trations of antigen-bearing substances in biological samples; they are also used semi-quantitatively or qualitatively (e.g., in diagnostic embodiments that identify the presence of a varCSA molecule in biological samples).

[0105] Additionally, FCR3.varCSA, fragments of FCR3.varCSA, A4 tres DBL3- γ , or ItG2-CS2 DBL2- γ can be used to induce antibody production in humans. That is, these peptides whether made chemically or as detailed above, can be used as an antigen or vaccine so as to elicit an immune response in a patient. Accordingly, FCR3.varCSA, fragments of FCR3.varCSA, A4 tres DBL3- γ , or ItG2-CS2 DBL2- γ can be joined to or administered with another protein, carrier, support, or adjuvant so as to generate a pharmaceutical or vaccine that will induce potent immune response. Additionally, nucleic acids encoding FCR3.varCSA, fragments of FCR3.varCSA, A4 tres DBL3-7, or ItG2-CS2 DBL2- γ can be administered by themselves or with FCR3.varCSA, fragments of FCR3.varCSA, A4 tres DBL3- γ , or ItG2-CS2 DBL2- γ and, as above, can be joined to or administered with a protein, carrier, support, or adjuvant. These nucleic acids can be administered "naked" or can be incorporated into vectors. Vaccination protocols can include, for example, identifying a subject in need of a vaccine (e.g., pregnant women in regions populated with *P. falciparum*) and administering to said subject a therapeutically effective amount of either a protein or a nucleic acid-based vaccines or combinations of protein and nucleic acid vaccines. The next section describes the use of var characterization assays and methods to identify agents that modulate FCR3.varCSA-mediated adhesion.

[0106] Modulation of FCR3.varCSA-dependent Adhesion to CSA

[0107] The data above establishes that FCR3.varCSA, fragments of FCR3.varCSA, A4 tres DBL3- γ , or ItG2-CS2 DBL2- γ efficiently associate with CSA to form a varCSA-CSA complex. The formation of such a complex can be measured using many techniques common to immunology and receptor biology. By one approach, FCR3.varCSA dependent adhesion to CSA is analyzed by contacting a support having CSA or a representative fragment of CSA with FCR3.varCSA or a representative fragment of FCR3.varCSA. If the FCR3.varCSA or fragment thereof is detectably labeled (e.g., 125%, the association to immobilized CSA (or CSA fragment) can be directly determined by detecting the signal (e.g., scintillation counting). Alternatively, the association of FCR3.varCSA, fragments of FCR3.varCSA, A4 tres DBL3- γ , or ItG2-CS2 DBL2- γ with CSA can be determined indirectly by employing a detectably labeled antibody that has an epitope that corresponds to a region of FCR3.varCSA. In these assays, the support can be a resin, plastic, a membrane, a lipid, and a cell. Additionally, the varCSA or fragment thereof can be joined to a second support so as to more closely reproduce native conditions. Many varCSA characterization assays can be automated (e.g., high throughput screening employing a fluorescently labeled FCR3.varCSA or fragment thereof) so as to quickly identify regions of the molecule that are involved in binding to CSA. Values or results from these assays can be recorded on a computer readable media (e.g., in a database) and analyzed with a search program and retrieval program. Of course, embodiments of the invention include the converse of the assay described above. That is, immobilizing FCR3.varCSA, fragments of FCR3.varCSA, A4 tres DBL3-

γ , or ItG2-CS2 DBL2- γ on a support and detecting the adhesion of labeled CSA or fragments of CSA.

[0108] Additional embodiments include methods of identifying agents that modulate the formation of a varCSA-CSA complex. By one approach, an agent that modulates varCSA-CSA adhesion can be identified by contacting a support having CSA or a representative fragment thereof with FCR3.varCSA, fragments of FCR3.varCSA, A4 tres DBL3- γ , or ItG2-CS2 DBL2- γ in the presence of the agent. Detection of adhesion is accomplished, as described above, and successful agents are identified according to their ability to induce a desired modulation of the formation of the varCSA-CSA complex. As above, the support can be a resin, a membrane, plastic, a lipid, or a cell and the varCSA or fragment thereof can be joined to a second support so as to more nearly reproduce native binding conditions. In another approach, a support having a varCSA or a representative fragment thereof can be used to capture directly or indirectly labeled CSA or fragments of CSA. In some aspects, the fragments of FCR3.varCSA that are used have a polypeptide sequence that binds to CSA and is at least 80% homologous to FCR3.varCSA. As above, binding is conducted in the presence of the agent and FCR3.varCSA dependent adhesion to CSA is determined by the amount of labeled CSA bound to the immobilized FCR3.varCSA. In this method, the support can be a resin, a membrane, plastic, a lipid, and a cell and the CSA can also be joined to a second support to approximate native binding conditions.

[0109] In a preferred approach, an agent that modulates varCSA dependent adhesion to CSA is identified using a cell-based assay. Accordingly, cells are transfected with a construct comprising a nucleic acid sequence encoding a varCSA or a representative fragment thereof. Transfectants are brought in contact with a support having CSA (or CSA fragment) and, as above, binding is conducted in the presence of the agent. Adhesion to CSA is determined by the amount of labeled CSA bound to the varCSA (or fragment thereof) expressing cells. In this method, the support can be a resin, a membrane, plastic, a lipid, and another cell. The converse of this assay can also be performed. That is, CSA expressing cells can be bound to immobilized varCSA or fragments of varCSA in the presence of a modulator. Further, a two-cell adhesion assay employing a first cell that expresses CSA and a second cell that expresses a varCSA or fragment thereof can be performed. Accordingly, the inhibition of cell aggregation in the presence of a modulator indicates that the agent is effective at disrupting varCSA-mediated CSA adhesion.

[0110] In some aspects of the invention, nucleic acids encoding FCR3.varCSA, nucleic acids complementary to FCR3.varCSA, FCR3.varCSA protein, and polypeptide fragments of FCR3.varCSA are agents that modulate (e.g., inhibit or enhance) the formation of the FCR3.varCSA-CSA complex. Several embodiments are provided that inhibit the association of FCR3.varCSA in a FCR3.varCSA-CSA complex or otherwise inhibit PRBC binding, sequestration, and the onset of maternal malaria (collectively referred to as "FCR3.varCSA inhibitory agents"). One embodiment concerns an inhibitory agent that is an antisense oligonucleotide or ribozyme that hybridizes to nucleic acid encoding regions of a varCSA molecule (e.g., FCR3.varCSA, DBL3, CIDR1, A4 tres DBL3- γ , or ItG2-CS2 DBL2- γ). By "antisense oligonucleotide" is meant a nucleic acid or modified nucleic

acid including, but not limited to DNA, RNA, modified DNA or RNA (including branched chain nucleic acids and 2' O-methyl RNA) and PNA (polyamide nucleic acid).

[0111] Several ribozymes known to those of skill in the art can be easily designed to hybridize to nucleic acid sequence encoding a varCSA or fragment thereof and thereby inhibit the production of functional protein. Desirably, antisense oligonucleotides or ribozymes that hybridize to the start codon are used. In one embodiment, full length antisense FCR3.varCSA is used to significantly reduced FCR3.varCSA-dependent adhesion to CSA. Many other antisense oligonucleotides or ribozymes that interfere with the formation of a varCSA-CSA complex can be designed and screened by the methods detailed previously.

[0112] The antisense nucleic acids should have a length and melting temperature sufficient to permit formation of an intracellular duplex having sufficient stability to inhibit the expression of the mRNA in the duplex. Strategies for designing antisense nucleic acids suitable for use in gene therapy are disclosed in Green et al., *Ann. Rev. Biochem.*, 55:569-597 (1986) and Izant and Weintraub, *Cell*, 36:1007-1015 (1984). In some strategies, antisense molecules are obtained from a nucleotide sequence encoding FCR3.varCSA by reversing the orientation of the coding region with respect to a promoter so as to transcribe the opposite strand from that which is normally transcribed in the cell. Antisense molecules and ribozymes can be prepared by any method known in the art for the synthesis of RNA molecules. These include techniques for chemically synthesizing oligonucleotides such as solid phase phosphoramidite chemical synthesis.

[0113] Additionally, RNA molecules can be generated by in vitro and in vivo transcription of DNA sequences encoding FCR3.varCSA, fragments of FCR3.varCSA, A4 tres DBL3- γ , or ItG2-CS2 DBL2- γ . Such DNA sequences can be incorporated into a wide variety of vectors with suitable RNA polymerase promoters such as T7 or SP6. Further, antisense cDNA constructs that synthesize antisense RNA constitutively or inducibly can be introduced into cell lines, cells or tissues. Still further, oligonucleotides that are complementary to the mRNA encoding FCR3.varCSA, fragments of FCR3.varCSA, A4 tres DBL3- γ , or ItG2-CS2 DBL2- γ can be synthesized in vitro. Thus, antisense nucleic acids are capable of hybridizing to a varCSA mRNA to create a duplex. In some embodiments, the antisense sequences can contain modified sugar phosphate backbones to increase stability and make them less sensitive to RNase activity. Possible modifications include, but are not limited to, the addition of flanking sequences at the 5' and/or 3' ends of the molecule or the use of phosphorothioate or 2' O-methyl rather than phosphodiesterase linkages within the backbone of the molecule. This concept is inherent in the production of PNAs and can be extended in all of these molecules by the inclusion of nontraditional bases such as inosine, queosine and wybutosine as well as acetyl-, methyl-, thio- and similarly modified forms of adenine, cytidine, guanine, thymine, and uridine that are not as easily recognized by endogenous endonucleases. Further examples are described by Rossi et al., *Pharmacol. Ther.*, 50(2):245-254, (1991).

[0114] Various types of antisense oligonucleotides complementary to a nucleic acid encoding FCR3.varCSA,

fragments of FCR3.varCSA, A4 tres DBL3- γ , or ItG2-CS2 DBL2- γ can be used. In one preferred embodiment, stable and semi-stable antisense oligonucleotides described in International Application No. PCT WO94/23026 are used. In these molecules, the 3' end or both the 3' and 5' ends are engaged in intramolecular hydrogen bonding between complementary base pairs. These molecules are better able to withstand exonuclease attacks and exhibit increased stability compared to conventional antisense oligonucleotides. In another preferred embodiment, the antisense oligodeoxynucleotides described in International Application No. WO 95/04141 are used. In yet another preferred embodiment, the covalently cross-linked antisense oligonucleotides described in International Application No. WO 96/31523 are used. These double- or single-stranded oligonucleotides comprise one or more, respectively, inter- or intra-oligonucleotide covalent cross-linkages, wherein the linkage consists of an amide bond between a primary amine group of one strand and a carboxyl group of the other strand or of the same strand, respectively, the primary amine group being directly substituted in the 2' position of the strand nucleotide monosaccharide ring, and the carboxyl group being carried by an aliphatic spacer group substituted on a nucleotide or nucleotide analog of the other strand or the same strand, respectively.

[0115] The antisense oligodeoxynucleotides and oligonucleotides disclosed in International Application No. WO 92/18522 can also be used. These molecules are stable to degradation and contain at least one transcription control recognition sequence that binds to control proteins and are effective as decoys therefor. These molecules can contain "hairpin" structures, "dumbbell" structures, "modified dumbbell" structures, "cross-linked" decoy structures and "loop" structures. In another preferred embodiment, the cyclic double-stranded oligonucleotides described in European Patent Application No. 0 572 287 A2 are used. These ligated oligonucleotide "dumbbells" contain the binding site for a transcription factor and inhibit expression of the gene under control of the transcription factor by sequestering the factor. Use of the closed antisense oligonucleotides disclosed in International Application No. WO 92/19732 is also contemplated. Because these molecules have no free ends, they are more resistant to degradation by exonucleases than are conventional oligonucleotides. These oligonucleotides can be multifunctional, interacting with several regions that are not adjacent to the target mRNA.

[0116] The appropriate level of antisense nucleic acids required to inhibit formation of the varCSA-CSA complex can be determined using in vitro expression analysis and the varCSA characterization assays described herein. The antisense molecule can be introduced into the cells expressing FCR3.varCSA, fragments of FCR3.varCSA, A4 tres DBL3- γ , or ItG2-CS2 DBL2- γ by diffusion, injection, infection or transfection using procedures known in the art. For example, the antisense nucleic acids can be introduced into the body as a bare or naked oligonucleotide, oligonucleotide encapsulated in lipid, oligonucleotide sequence encapsulated by viral protein, or as an oligonucleotide operably linked to a promoter contained in an expression vector. The expression vector can be any of a variety of expression vectors known in the art, including retroviral or viral vectors, vectors capable of extrachromosomal replication, or integrating vectors. The vectors can be DNA or RNA.

[0117] The antisense molecules are introduced onto cell samples at a number of different concentrations preferably between $1 \times 10^{-10} \text{M}$ to $1 \times 10^{-4} \text{M}$. Once the minimum concentration that can adequately control gene expression is identified, the optimized dose is translated into a dosage suitable for use in vivo. For example, an inhibiting concentration in culture of 1×10^{-7} translates into a dose of approximately 0.6 mg/kg bodyweight. Levels of oligonucleotide approaching 100 mg/kg bodyweight or higher can be possible after testing the toxicity of the oligonucleotide in laboratory animals. It is additionally contemplated that cells from a vertebrate, such as a mammal or human, are removed, treated with the antisense oligonucleotide, and reintroduced into the vertebrate.

[0118] Ribozymes can also be used to reduce or eliminate expression of FCR3.varCSA, fragments of FCR3.varCSA, A4 tres DBL3- γ , or ItG2-CS2 DBL2- γ . Ribozymes are enzymatic RNA molecules capable of catalyzing the specific cleavage of RNA. The mechanism of ribozyme action involves sequence-specific hybridization of the ribozyme molecule to complementary target RNA, followed by endonucleolytic cleavage. Within the scope of aspects of the invention, are engineered hammerhead motif ribozyme molecules that can specifically and efficiently catalyze endonucleolytic cleavage of a sequence encoding FCR3.varCSA, for example. Specific ribozyme cleavage sites within any potential RNA target are initially identified by scanning the target molecule for ribozyme cleavage sites that include the following sequences, GUA, GUU and GUC. Once identified, short RNA sequences of between 15 and 20 ribonucleotides corresponding to the region of the target gene containing the cleavage site can be evaluated for secondary structural features that may render the oligonucleotide inoperable. The suitability of candidate targets can also be evaluated by testing accessibility to hybridization with complementary oligonucleotides using ribonuclease protection assays. Delivery of antisense and ribozyme agents by transfection and by liposome are quite well known in the art.

[0119] Another embodiment of an inhibitory agent is a polypeptide that interferes with the association of FCR3.varCSA, fragments of FCR3.varCSA, A4 tres DBL3- γ , or ItG2-CS2 DBL2- γ with CSA. Polypeptide fragments that inhibit the association of with CSA can be rapidly engineered and identified given the present disclosure and candidate polypeptides can contain regions of FCR3.varCSA, fragments of FCR3.varCSA, A4 tres DBL3- γ , ItG2-CS2 DBL2- γ or CSA. The screening of polypeptide fragments and mutant proteins that inhibit the association of a complex having these molecules would be routine given the present disclosure and assays described herein. For example, polypeptide FCR3.varCSA inhibitory agents can be identified by their ability to disrupt the formation of the FCR3.varCSA-CSA complex by employing conventional affinity chromatography techniques, sandwich assays, ELISA assays, or other binding assays known to those of skill in the art and described above. A screening method, for example, wherein the polypeptide FCR3.varCSA inhibitory agent is administered to cells expressing FCR3.varCSA in culture and cell lysates are analyzed by immunoprecipitation and Western blot can rapidly evaluate the polypeptide's ability to inhibit the association of a an FCR3.varCSA-CSA complex.

[0120] In another embodiment, concentrations of FCR3.varCSA, fragments of FCR3.varCSA, A4 tres DBL3- γ , or ItG2-CS2 DBL2- γ are raised in a cell so as to enhance adhesion to CSA. There may be many ways to raise the concentration of these molecules or CSA or both in a cell. Liposome-mediated transfer, is one approach to deliver FCR3.varCSA, fragments of FCR3.varCSA, A4 tres DBL3- γ , ItG2-CS2 DBL2- γ or CSA or both protein to a cell. Alternatively, the concentration of FCR3.varCSA, fragments of FCR3.varCSA, A4 tres DBL3- γ , ItG2-CS2 DBL2- γ or CSA or any combination thereof can be raised in a cell by transfecting constructs encoding these molecules. A construct for use in the transfection of FCR3.varCSA into cells in culture, for example, was described previously and many others can be developed by those of skill in the art.

[0121] Retroviral constructs for the delivery of nucleic acid encoding FCR3.varCSA, fragments of FCR3.varCSA, A4 tres DBL3- γ , or ItG2-CS2 DBL2- γ or complements thereof are also contemplated. Other embodiments of varCSA inhibitory or enhancing agents (collectively referred to as "modulators") include antibodies, peptidomimetics, and chemicals that inhibit or enhance adhesion to CSA. Several other methods for identifying agents that modulate the formation of a varCSA-CSA complex can be used. The discussion below describes methods of molecular modeling, combinatorial chemistry, and rational drug design, which can be used to identify molecules that interact with a varCSA molecule and thereby modulate the formation of a varCSA-CSA complex.

[0122] Methods of Rational Drug Design

[0123] Combinatorial chemistry is the science of synthesizing and testing compounds for bioactivity en masse, instead of one by one, the aim being to discover drugs and materials more quickly and inexpensively than was formerly possible. Although the following describes the use of FCR3.varCSA, fragments thereof and nucleic acids encoding these molecules to conduct rational drug design, it should be understood that these approaches can be used with other molecules that mediate PRBC binding, sequestration, and the onset of maternal malaria, including but not limited to fragments of FCR3.varCSA (e.g., DBL3 and CIDR1), A4 tres DBL3- γ , and ItG2-CS2 DBL2- γ . Thus, the description of FCR3.varCSA, fragments thereof and nucleic acids encoding these molecules are provided for the purposes of explanation and example.

[0124] In some embodiments, search programs are employed to compare regions of FCR3.varCSA that modulate the formation of a FCR3.varCSA-CSA complex with other molecules, such as peptides, peptidomimetics, and chemicals so that therapeutic interactions of the molecules can be predicted and new derivative molecules can be designed. In other embodiments, search programs are employed to compare regions of molecules that interact with FCR3.varCSA and, thereby modulate the formation of a FCR3.varCSA-CSA complex, with other molecules such as peptides, peptidomimetics, and chemicals, so that therapeutic interactions of the molecules can be predicted and new derivative FCR3.varCSA modulating agents can be designed. (Schneider, *Genetic Engineering News* December: page 20 (1998), Tempczyk et al., *Molecular Simulations Inc. Solutions* April (1997), and Butenhof, *Molecular Simulations Inc. Case Notes* (August 1998)). This process of

directed combinatorial chemistry is referred to as “rational drug design”. Libraries of molecules that resemble FCR3.varCSA or interact with FCR3.varCSA and, thereby inhibit or enhance the function of FCR3.varCSA (“modulate” FCR3.varCSA activity) can be created. In some contexts, the term “FCR3.varCSA modulating agent” or “modulators” includes FCR3.varCSA, polypeptide fragments corresponding to FCR3.varCSA, fusion proteins comprising FCR3.varCSA or polypeptide fragments of FCR3.varCSA, nucleic acids encoding these molecules, peptidomimetics, chemicals, and other molecules that mediate PRBC binding, sequestration, and the onset of maternal malaria.

[0125] One goal of rational drug design is to produce structural analogs of biologically active polypeptides of interest or of small molecules with which they interact (e.g., agonists, antagonists, null compounds) in order to fashion drugs that are, for example, more or less potent forms of the molecule. (See, e.g., Hodgson, *Bio. Technology* 9:19-21 (1991)). Rational drug design has been used to develop HIV protease inhibitors and agonists for five different somatostatin receptor subtypes. (Erickson et al., *Science* 249:527-533 (1990) and Berk et al., *Science* 282:737 (1998)).

[0126] By starting with the sequence or protein models of FCR3.varCSA, and/or fragments thereof, polypeptides having two-dimensional and/or three-dimensional homology can be rapidly identified. In a two-dimensional approach, a percent sequence identity can be determined by standard methods that are commonly used to compare the similarity and position of the amino acid of two polypeptides. Using a computer program such as BLAST or FASTA, two polypeptides are aligned for optimal matching of their respective amino acids (either along the full length of one or both sequences, or along a predetermined portion of one or both sequences). Such programs provide “default” opening penalty and a “default” gap penalty, and a scoring matrix such as PAM 250 (a standard scoring matrix; see Dayhoff et al., in: *Atlas of Protein Sequence and Structure*, Vol. 5, Supp. 3 (1978) can be used in conjunction with the computer program. The percent identity can then be calculated as:

$$\frac{\text{total number of identical matches}}{\text{[length of the longer sequence within the matched span + number of gaps introduced into the longer sequence in order to align the two sequences]}} \times 100$$

[0127] Accordingly, the protein sequence corresponding to FCR3.varCSA is compared to known sequences on a protein basis. Protein sequences corresponding to FCR3.varCSA are compared, for example, to known amino acid sequences found in Swissprot release 35, PIR release 53 and Genpept release 108 public databases using BLASTP with the parameter W=8 and allowing a maximum of 10 matches. In addition, the protein sequences encoding FCR3.varCSA are compared to publicly known amino acid sequences of Swissprot using BLASTX with the parameter E=0.001. Because the region involved in CSA binding can be as small as three amino acids, the embodied polypeptides can have the following degrees of homology to FCR3.varCSA: 1.0%, 2.0%, 3.0%, 4.0%, 5.0%, 6.0%, 7.0%, 8.0%, 9.0%, 10.0%, 11.0%, 12.0%, 13.0%, 14.0%, 15.0%, 16.0%, 17.0%, 18.0%, 19.0%, 20.0%, 21.0%, 22.0%,

23.0%, 24.0%, 25.0%, 26.0%, 27.0%, 28.0%, 29.0%, 30.0%, 31.0%, 32.0%, 33.0%, 34.0%, 35.0%, 36.0%, 37.0%, 38.0%, 39.0%, 40.0%, 41.0%, 42.0%, 43.0%, 44.0%, 45.0%, 46.0%, 47.0%, 48.0%, 49.0%, 50.0%, 51.0%, 52.0%, 53.0%, 54.0%, 55.0%, 56.0%, 57.0%, 58.0%, 59.0%, 60.0%, 61.0%, 62.0%, 63.0%, 64.0%, 65.0%, 66.0%, 67.0%, 68.0%, 69.0%, 70.0%, 71.0%, 72.0%, 73.0%, 74.0%, 75.0%, 76.0%, 77.0%, 78.0%, 79.0%, 80.0%, 81.0%, 82.0%, 83.0%, 84.0%, 85.0%, 86.0%, 87.0%, 88.0%, 89.0%, 90.0%, 91.0%, 92.0%, 93.0%, 94.0%, 95.0%, 96.0%, 97.0%, 98.0%, 99.0%, and 100.0% The candidate polypeptides are identified and are subsequently examined using the functional assays described herein. Candidate polypeptides that interact with FCR3.varCSA to modulate the formation of a FCR3.varCSA-CSA complex and thereby effect PRBC adhesion to CSA can be identified in this manner.

[0128] Additionally, a search program can be used to compare the three-dimensional structure of FCR3.varCSA or fragments of FCR3.varCSA with other known three-dimensional structures. Once candidate related structures are identified, these molecules can be made recombinantly or by peptide or chemical synthesis. The newly generated compounds are then screened in FCR3.varCSA characterization assays so as to identify modulators that interact with FCR3.varCSA and thereby effect the formation of a FCR3.varCSA-CSA complex.

[0129] In the past, the three-dimensional structure of proteins has been determined in a number of ways. Perhaps the best known way of determining protein structure involves the use of x-ray crystallography. A general review of this technique can be found in Van Holde, K. E. *Physical Biochemistry*, Prentice-Hall, N.J. pp. 221-239 (1971). Using this technique, it is possible to elucidate three-dimensional structure with good precision. Additionally, protein structure can be determined through the use of techniques of neutron diffraction, or by nuclear magnetic resonance (NMR). (See, e.g., Moore, W. J., *Physical Chemistry*, 4th Edition, Prentice-Hall, N.J. (1972)).

[0130] Alternatively, the protein model embodiments of the present invention can be constructed using computer-based protein modeling techniques. By one approach, the protein folding problem is solved by finding target sequences that are most compatible with profiles representing the structural environments of the residues in known three-dimensional protein structures. (See, e.g., Eisenberg et al., U.S. Pat. No. 5,436,850 issued Jul. 25, 1995). In another technique, the known three-dimensional structures of proteins in a given family are superimposed to define the structurally conserved regions in that family. This protein modeling technique also uses the known three-dimensional structure of a homologous protein to approximate the structure of a polypeptide of interest. (See e.g., Srinivasan, et al., U.S. Pat. No. 5,557,535 issued Sep. 17, 1996). Conventional homology modeling techniques have been used routinely to build models of proteases and antibodies. (Sowdhamini et al., *Protein Engineering* 10:207, 215 (1997)). Comparative approaches can also be used to develop three-dimensional protein models when the protein of interest has poor sequence identity to template proteins. In some cases, proteins fold into similar three-dimensional structures despite having very weak sequence identities. For example, the

three-dimensional structures of a number of helical cytokines fold in similar three-dimensional topology in spite of weak sequence homology.

[0131] The recent development of threading methods and “fuzzy” approaches now enables the identification of likely folding patterns and functional protein domains in a number of situations where the structural relatedness between target and template(s) is not detectable at the sequence level. By one method, fold recognition is performed using Multiple Sequence Threading (MST) and structural equivalences are deduced from the threading output using the distance geometry program DRAGON that constructs a low resolution model. A full-atom representation is then constructed using a molecular modeling package such as QUANTA.

[0132] According to this 3-step approach, candidate templates are first identified by using the novel fold recognition algorithm MST, which is capable of performing simultaneous threading of multiple aligned sequences onto one or more 3-D structures. In a second step, the structural equivalences obtained from the MST output are converted into interresidue distance restraints and fed into the distance geometry program DRAGON, together with auxiliary information obtained from secondary structure predictions. The program combines the restraints in an unbiased manner and rapidly generates a large number of low resolution model confirmations. In a third step, these low resolution model confirmations are converted into full-atom models and subjected to energy minimization using the molecular modeling package QUANTA. (See e.g., Aszódi et al., *Proteins: Structure, Function, and Genetics*, Supplement 1:38-42 (1997)).

[0133] In one approach, a three-dimensional structure of a polypeptide of interest (e.g., FCR3.varCSA, and/or fragments thereof or a FCR3.varCSA modulating agent) is determined by x-ray crystallography, NMR, or neutron diffraction and computer modeling, as described above. Useful protein models of the polypeptide of interest can also be gained by computer modeling alone. Combinatorial chemistry can then be employed to design derivatives of the polypeptide of interest based on the three-dimensional models. The candidate FCR3.varCSA modulating agents are then tested in functional assays. The assays, described herein and assays that evaluate the formation of a FCR3.varCSA-CSA complex in the presence of FCR3.varCSA or fragments thereof that will be apparent to one of skill in the art given the disclosure herein (referred to collectively as “FCR3.varCSA characterization assays”) are performed on the FCR3.varCSA modulating agents and groups of FCR3.varCSA modulating agents (wherein the grouping is based on the potency of modulation of the formation of a FCR3.varCSA-CSA complex) are identified and recorded on a computer readable media. Further cycles of modeling and FCR3.varCSA characterization assays can be employed to more narrowly define the parameters needed in an optimal FCR3.varCSA modulating agent.

[0134] By another approach, a FCR3.varCSA modulating agent that interacts with FCR3.varCSA can be manufactured and identified as follows. First, a molecular model of one or more FCR3.varCSA modulating agents or portions of FCR3.varCSA modulating agents that interact with FCR3.varCSA are created using one of the techniques discussed above or as known in the art. FCR3.varCSA modulating agents that are known to interact with

FCR3.varCSA include antibodies and fragments of CSA. Next, chemical libraries and databases are searched for molecules similar in structure to the known FCR3.varCSA modulating agents. Identified candidate FCR3.varCSA modulating agents are then screened in the FCR3.varCSA characterization assays, described above, and the agents that produce the desired response are used as templates for further library construction. Libraries of FCR3.varCSA modulating agents are synthesized on solid support beads by split-and-pool synthesis, a multistage process for producing very large numbers of compounds. The support-bound agents are then used in FCR3.varCSA characterization assays or “free mixtures” are created by cleaving the agent from the support and these free mixtures are screened in the FCR3.varCSA characterization assays. Compounds that produce desirable responses are identified, recorded on a computer readable media, and the process is repeated to select optimal FCR3.varCSA modulating agents.

[0135] Each FCR3.varCSA modulating agent and its response in a FCR3.varCSA characterization assay can be recorded on a computer readable media and a database or library of FCR3.varCSA modulating agents and respective responses in the FCR3.varCSA characterization assay can be generated. These databases or libraries can be used by researchers to identify important differences between active and inactive molecules so that compound libraries are enriched for FCR3.varCSA modulating agents that have favorable characteristics. Further, enrichment can be achieved by using approaches in dynamic combinatorial chemistry. (See e.g., Angnew, *Chem. Int. Ed.*, 37:2828 (1998)). For example, a target biomolecule, such as FCR3.varCSA, is joined to a support and is bound by the FCR3.varCSA modulating agents from the libraries generated above. The FCR3.varCSA resin bound with one or more candidate FCR3.varCSA modulating agents is removed from the binding reaction, the FCR3.varCSA modulating agents are eluted from the support, and are identified. Cycles of immobilized target binding assays are conducted, classes of FCR3.varCSA modulating agents that exhibit desired binding characteristics are identified, and this data is recorded on a computer readable media and is used to select more FCR3.varCSA modulating agents that produce a desired modulation of the formation of a FCR3.varCSA-CSA complex.

[0136] In addition, a peptide of interest (e.g., FCR3.varCSA, and/or fragments thereof or a FCR3.varCSA modulating agent) can be analyzed by an alanine scan (Wells, *Methods in Enzymol.* 202:390-411 (1991)) or other types of site-directed mutagenesis analysis. In alanine scan, for example, an amino acid residue is replaced by alanine, and its affect on the peptide's activity is measured by functional assays, such as the FCR3.varCSA characterization assays described herein. Each of the amino acid residues of the peptide is analyzed in this manner and the regions important for a specific modulation of the formation of a FCR3.varCSA-CSA complex are identified. Subsequently, these functionally important regions are recorded on a computer readable medium, stored in a first database in a computer system, and a search program is employed to generate protein models of the functionally important regions. Once protein models of the functionally important regions have been generated, a second database comprising one or more libraries having peptides, chemicals, peptidomimetics and other agents is accessed by a search program

and individual agents are compared to the protein models to identify agents that comprise homologous regions or domains that resemble the identified functionally important regions. Agents identified by the approach above are then tested in the FCR3.varCSA characterization assays and are used to construct multimeric agents and/or are incorporated into pharmaceuticals, as detailed below.

[0137] In another embodiment, computer modeling and the sequence-to-structure-to-function paradigm is exploited to identify more FCR3.varCSA modulating agents that modulate the formation of a FCR3.varCSA-CSA complex. By this approach, first the structure of a FCR3.varCSA modulating agent having a known response in a FCR3.varCSA characterization assay (e.g., FCR3.varCSA, and fragments thereof, and antibodies to FCR3.varCSA, is determined from its sequence using a threading algorithm, which aligns the sequence to the best matching structure in a structural database. Next, the protein's active site (i.e., the site important for a desired response in the FCR3.varCSA characterization assay) is identified and a "fuzzy functional form" (FFF)—a three-dimensional descriptor of the active site of a protein—is created. (See e.g., Fetrow et al., *J. Mol. Biol.* 282:703-711 (1998) and Fetrow and Skolnick, *J. Mol. Biol.* 281: 949-968 (1998)).

[0138] The FFFs are built by iteratively superimposing the protein geometries from a series of functionally related proteins with known structures. The FFFs are not overly specific, however, and the degree to which the descriptors can be relaxed is explored. In essence, conserved and functionally important residues for a desired response are identified and a set of geometric and conformational constraints for a specific function are defined in the form of a computer algorithm. The program then searches experimentally determined protein structures from a protein structural database for sets of residues that satisfy the specified constraints. In this manner, homologous three-dimensional structures can be compared and degrees (e.g., percentages of three-dimensional homology) can be ascertained.

[0139] By using this computational protocol, genome sequence data bases such as maintained by various organizations including: <http://www.tigr.org/tdb>; <http://www.genetics.wisc.edu>; <http://genome-www.stanford.edu/~ball>; <http://hiv-web.lanl.gov>; <http://www.ncbi.nlm.nih.gov>; <http://www.ebi.ac.uk>; <http://pasteur.fr/other/biology>; and <http://www-genome.wi.mit.edu>, can be rapidly screened for specific protein active sites and for identification of the residues at those active sites that resemble a desired molecule. Several other groups have developed databases of short sequence patterns or motifs designed to identify a given function or activity of a protein. These databases, notably Prosite (<http://expasy.hcuge.ch/spro/prosite.html>); Blocks (<http://www.blocks.fhrc.org>); and Prints (<http://www.biochem.ucl.ac.uk/lbms/dbbrowser/PRINTS/PRINTS.html>, use short stretches of sequence information to identify sequence patterns that are specific for a given function; thus they avoid the problems arising from the necessity of matching entire sequences. In this manner, new FCR3.varCSA modulating agents are rationally selected for further identification by FCR3.varCSA characterization assays, as described above. Rounds or cycles of functional assays on the molecules and derivatives thereof and further FFF refinement and database searching allows an investigator to more narrowly define

classes of FCR3.varCSA modulating agents that produce a desired modulation of the formation of a FCR3.varCSA-CSA complex.

[0140] Many computer programs and databases can be used with embodiments of the invention to identify agents that modulate FCR3.varCSA-mediated adhesion to CSA. The following list is intended not to limit the invention but to provide guidance to programs and databases that are useful with the approaches discussed above. The programs and databases that may be used include, but are not limited to: MacPattern (EMBL), DiscoveryBase (Molecular Applications Group), GeneMine (Molecular Applications Group), Look (Molecular Applications Group), MacLook (Molecular Applications Group), BLAST and BLAST2 (NCBI), BLASTN and BLASTX (Altschul et al., *J. Mol. Biol.* 215: 403 (1990)), FASTA (Pearson and Lipman, *Proc. Natl. Acad. Sci. USA*, 85: 2444 (1988)), Catalyst (Molecular Simulations Inc.), Catalyst/SHAPE (Molecular Simulations Inc.), Cerius².DBAccess (Molecular Simulations Inc.), HypoGen (Molecular Simulations Inc.), Insight II, (Molecular Simulations Inc.), Discover (Molecular Simulations Inc.), CHARM[™] (Molecular Simulations Inc.), Felix (Molecular Simulations Inc.), DelPhi, (Molecular Simulations Inc.), QuanteMM, (Molecular Simulations Inc.), Homology (Molecular Simulations Inc.), Modeler (Molecular Simulations Inc.), Modeller 4 (Sali and Blundell *J. Mol. Biol.* 234:217-241 (1997)), ISIS (Molecular Simulations Inc.), Quanta/Protein Design (Molecular Simulations Inc.), WebLab (Molecular Simulations Inc.), WebLab Diversity Explorer (Molecular Simulations Inc.), Gene Explorer (Molecular Simulations Inc.), SeqFold (Molecular Simulations Inc.), the EMBL/Swissprotein database, the MDL Available Chemicals Directory database, the MDL Drug Data Report data base, the Comprehensive Medicinal Chemistry database, Derwent's World Drug Index database, and the BioByteMasterFile database. Many other programs and data bases would be apparent to one of skill in the art given the present disclosure.

[0141] Libraries of information on FCR3.varCSA modulating agents with their corresponding response in FCR3.varCSA characterization assays can be generated by performing the rational drug design approaches above in conjunction with the FCR3.varCSA characterization assays. A record of the results for each FCR3.varCSA modulating agent is generated and groups of FCR3.varCSA modulating agents are identified and stored on a computer readable media. Databases of this information are valuable to investigators and clinicians for selecting the type of FCR3.varCSA modulating agent-based pharmaceutical to treat or elicit a particular response. Preferable libraries are created by performing the assays above on FCR3.varCSA and fragments thereof.

[0142] Many of the FCR3.varCSA modulating agents are provided in biotechnological tools, diagnostics, and pharmaceuticals as multimeric or multimerized agents or both that can be joined to a support. In the disclosure below, we discuss the preparation of multimeric supports and multimerized FCR3.varCSA modulating agents comprising FCR3.varCSA or fragments of FCR3.varCSA, complementary nucleic acids to FCR3.varCSA, FCR3.varCSA or fragments of FCR3.varCSA, antibodies or antibody fragments that recognize epitopes of FCR3.varCSA, and FCR3.varCSA fusion proteins.

[0143] Preparation of Multimeric Supports and Multimerized FCR3.varCSA Modulators

[0144] Biotechnological tools and components to prophylactic and therapeutic agents desirably provide FCR3.varCSA, fragments of FCR3.varCSA, A4 tres DBL3- γ , or ItG2-CS2 DBL2- γ antibodies or antibody fragments that recognize epitopes on these molecules, fusion proteins containing these molecules, nucleic acids encoding these molecules (and complementary nucleic acids thereof) in such a form or in such a way that a sufficient affinity, modulation of a varCSA-CSA complex formation or induction of an immune response is achieved. While a natural monomeric agent (that is, an agent that presents a discrete molecule, thus, carrying only one binding epitope or domain) can be sufficient to achieve a desired response, a synthetic agent or a multimeric agent (that is, an agent that presents multiple molecules, thus, having several binding epitopes or domains) often times can elicit a greater response. It should be noted that the term "multimeric" refers to the presence of more than one molecule on an agent, for example, several individual molecules of an antibody joined to a support, as distinguished from the term "multimerized" that refers to an agent that has more than one molecule joined as a single discrete compound molecule on a support, for example several antibody molecules joined to form a single compound molecule that is joined to a support.

[0145] A multimeric agent (synthetic or natural) that modulates the formation of a varCSA-CSA complex or induces an immune response is obtained by joining FCR3.varCSA, fragments of FCR3.varCSA, A4 tres DBL3- γ , or ItG2-CS2 DBL2- γ antibodies or antibody fragments that recognize epitopes on these molecules, fusion proteins containing these molecules, nucleic acids encoding these molecules (and complementary nucleic acids thereof), collectively referred to as "FCR3.varCSA modulating agents", "varCSA modulating agents" or "modulators", to a macromolecular support. These modulating agents including peptidomimetics and chemical molecules that resemble these ligands are also joined to supports so as to create the multimeric agents of the invention. A "support" can also be a carrier, a protein, a resin or any macromolecular structure used to join or immobilize a FCR3.varCSA modulating agent. Solid supports include, but are not limited to, the walls of wells of a reaction tray, test tubes, polystyrene beads, magnetic beads, nitrocellulose strips, membranes, microparticles such as latex particles, sheep (or other animal) red blood cells, Duracyte® artificial cells, and others. In several embodiments, the macromolecular support has a hydrophobic surface that interacts with a portion of the varCSA modulating agent by a hydrophobic non-covalent interaction. In some cases, the hydrophobic surface of the support is a polymer such as plastic or any other polymer in which hydrophobic groups have been linked such as polystyrene, polyethylene or polyvinyl. Additionally, the varCSA modulating agent is covalently bound to carriers including proteins and oligo/polysaccharides (e.g. cellulose, starch, glycogen, chitosane or aminated sepharose). In these later embodiments, a reactive group on a FCR3.varCSA modulating agent, such as a hydroxy or an amino group, is used to join to a reactive group on the carrier so as to create the covalent bond. Embodiments also comprise a support with a charged surface that interacts with the varCSA modulating agent. Additional embodiments comprise a support that has other reactive groups that are chemically activated so as to

attach a varCSA modulating agent, such as a peptide or chemical compound. For example, cyanogen bromide activated matrices, epoxy activated matrices, thio and thiopropyl gels, nitrophenyl chloroformate and N-hydroxy succinimide chloroformate linkages, or oxirane acrylic supports are used. (Sigma).

[0146] Inorganic carriers, such as silicon oxide material (e.g. silica gel, zeolite, diatomaceous earth or aminated glass) to which the varCSA modulating agent is covalently linked through a hydroxy, carboxy or amino group and a reactive group on the carrier are also embodiments. Carriers for use in the body, (i.e. for prophylactic or therapeutic applications) are desirably physiological, non-toxic and preferably, non-immunoresponsive. Contemplated carriers for use in the body include poly-L-lysine, poly-D, L-alanine and Chromosorb® (Johns-Manville Products, Denver Co.). Conjugated Chromosorb® (Synsorb-Pk) has been tested in humans for the prevention of hemolytic-uremic syndrome and was reported as not presenting adverse reactions. (Armstrong et al. *J. Infectious Diseases* 171:1042-1045 (1995)). For some embodiments, the administration of a "naked" carrier (i.e., lacking an attached varCSA modulating agent) that has the capacity to attach a varCSA modulating agent that modulates the formation of a varCSA-CSA complex inside the body of a subject is performed. By this approach, a "prodrug-type" therapy is administered in which the naked carrier is provided separately from the desired varCSA modulating agent and, once both are in the body, the carrier and the varCSA modulating agent assemble into a multimeric complex and modulate the formation of a varCSA-CSA complex.

[0147] In another embodiment, linkers, such as X linkers, of an appropriate length are inserted between the varCSA modulating agent and the support so as to encourage greater flexibility in the varCSA modulating agent and thereby overcome any steric hindrance that is presented by the support. The determination of an appropriate length of linker that allows for optimal binding and modulation of the formation of a varCSA-CSA complex, is made by screening the varCSA modulating agents with varying linkers in the varCSA characterization assays.

[0148] A composite support having more than one type of varCSA modulating agent is also an embodiment. A "composite support" is a carrier, a resin, or any macromolecular structure used to join or immobilize two or more different varCSA modulating agents that modulate the formation of a varCSA-CSA complex. The composite supports are also constructed by utilizing hydrophobic interactions and covalent linkages formed through reactive groups, as detailed above. Further, linkers, such as λ linkers, of an appropriate length between the varCSA modulating agents and the support are inserted in some embodiments so as to encourage greater flexibility in the molecule and overcome steric hindrance. The determination of an appropriate length of linker that allows for optimal binding and modulation of the formation of a varCSA-CSA complex, is made by screening the varCSA modulating agents with varying linkers in the varCSA characterization assays detailed in the present disclosure.

[0149] In other embodiments, the multimeric and composite supports discussed above have attached multimerized varCSA modulating agents so as to create a "multimerized-

multimeric support” and a “multimerized-composite support”, respectively. An embodiment of a multimerized varCSA modulating agent, for example, is obtained by creating an expression construct having two or more nucleotide sequences encoding the varCSA modulating agent joined together by using conventional techniques in molecular biology. The expressed fusion protein is one embodiment of a multimerized agent and is then joined to a support. A support having many such multimerized agents is termed a multimerized-multimeric support. The multimerized form of the varCSA modulating agent can be advantageous for many applications because of the ability to obtain an agent with a better ability to modulate the formation of a varCSA-CSA complex. The incorporation of linkers or spacers, such as flexible λ linkers, between the protein domains that make-up the multimerized agent can also be advantageous for some embodiments. The insertion of λ linkers of an appropriate length between protein binding domains, for example, encourages greater flexibility in the molecule and overcomes steric hindrance between the several proteins. Similarly, the insertion of linkers between the multimerized varCSA modulating agent and the support encourages greater flexibility and reduces steric hindrance presented by the support. The determination of an appropriate length of linker that allows for optimal binding and modulation of the formation of a varCSA-CSA complex can be accomplished by screening the varCSA modulating agents with varying linkers in the varCSA characterization assays detailed in this disclosure. In a similar fashion composite-multimerized-multimeric supports with and without linkers can be constructed by joining more than one different multimerized varCSA modulating agent to a support. The discussion that follows describes several diagnostic embodiments.

[0150] Diagnostic Embodiments

[0151] Several diagnostic and prognostic tools that detect the concentration and expression level of nucleic acids encoding a varCSA molecule (e.g., FCR3.varCSA) and the concentration and expression level of a varCSA molecule (e.g., FCR3.varCSA) in various tissues and fluids are used to determine whether an individual is suffering from maternal malaria or is likely to suffer from maternal malaria. Generally, the diagnostics and methods of use thereof can be classified according to whether the diagnostic detects the concentration or expression level of a nucleic acid or protein in a biological sample (e.g., blood). Accordingly, in some embodiments, the concentration and expression level of FCR3.varCSA in a biological sample can be determined by monitoring the amount of RNA in the sample. The detection of RNA encoding FCR3.varCSA in a sample indicates the existence or predilection to maternal malaria. Further, a detection of DNA encoding FCR3.varCSA in a biological sample indicates the existence or predilection to maternal malaria. Similarly, the concentration and expression level of FCR3.varCSA in a biological sample can be determined by monitoring the amount of FCR3.varCSA protein in the sample. The detection of FCR3.varCSA in a sample indicates the existence or predilection to maternal malaria. Other diagnostic approaches involve the detection of A4 tres DBL3- γ or ItG2-CS2 DBL2- γ and the detection of antibodies to FCR3.varCSA, fragments of FCR3.varCSA, A4 tres DBL3- γ , or ItG2-CS2 DBL2- γ .

[0152] For example, to determine the presence of FCR3.varCSA or FCR3.varCSA in a subject, first a biologi-

cal sample is obtained. Several methods known to those in the art can be employed to obtain a biological sample having red blood cells (e.g., phlebotomy). Once a biological sample from a subject in need of testing is obtained, many different techniques can be used to detect the concentration and expression level of FCR3.varCSA or FCR3.varCSA including, but not limited to, antibody-based detection techniques (e.g., ELISA, sandwich assays, immunoprecipitation, and immunoblots), bacteriophage display techniques, hybridization techniques (e.g., Southern and Northern), and enzymatic digestion (e.g., RNase protection) techniques. Some of these techniques can involve disposing the proteins and/or nucleic acids present in the biological sample on a support, and contacting the support with detection reagents such as antibodies to FCR3.varCSA or nucleic acid probes complementary to FCR3.varCSA mRNA. Desirably, the levels of expression or concentration of FCR3.varCSA or FCR3.varCSA or both from diseased and healthy individuals is compared to the level detected in the subject tested. Similar approaches can be used to detect A4 tres DBL3- γ , or ItG2-CS2 DBL2- γ or nucleic acids encoding these molecules.

[0153] In preferred embodiments, the nucleic acid embodiments are attached to a support in an ordered array wherein a plurality of nucleic acid probes are attached to distinct regions of the support that do not overlap with each other. Preferably, such an ordered array is designed to be “addressable” where the distinct locations of the probe are recorded and can be accessed as part of an assay procedure. In some embodiments, addressable nucleic acid arrays comprise a plurality of nucleic acid probes that complement a nucleic acid encoding FCR3.varCSA, fragments of FCR3.varCSA, A4 tres DBL3- γ , or ItG2-CS2 DBL2- γ . These probes are joined to a support in different known locations. The knowledge of the precise location of each nucleic acid probe makes these “addressable” arrays particularly useful in binding assays.

[0154] For example, an addressable array can comprise a support having several regions to which are joined a plurality nucleic acid probes that complement FCR3.varCSA. The nucleic acids from a preparation of several biological samples from a plurality of human subjects or a plurality of tissues or fluids from a single subject are labeled by conventional approaches (e.g., radioactivity or fluorescence) and the labeled samples are applied to the array under conditions that permit hybridization. If a nucleic acid in the sample hybridizes to a probe on the array, then a signal will be detected at a position on the support that corresponds to the location of the hybrid. Since the identity of each labeled sample is known and the region of the support on which the labeled sample was applied is known, an identification of the presence, concentration, and/or expression level can be rapidly determined. That is, by employing labeled standards of a known concentration of a nucleic acid encoding FCR3.varCSA, (e.g., RNA), an investigator can accurately determine the concentration of a nucleic acid encoding FCR3.varCSA in a sample and from this information can assess the expression level of FCR3.varCSA. Conventional methods in densitometry can also be used to more accurately determine the concentration or expression level of a nucleic acid encoding FCR3.varCSA. These approaches are easily automated using technology known to those of skill in the art of high throughput diagnostic analysis.

[0155] Additionally, an opposite approach to that presented above can be employed. Nucleic acids present in biological samples (e.g., tissues or fluids from one or more subjects or one or more sources in a subject's body) can be disposed on a support so as to create an addressable array. Preferably, the samples are disposed on the support at known positions that do not overlap. The presence of nucleic acids encoding FCR3.varCSA in each sample is determined by applying labeled nucleic acid probes that complement nucleic acids that encode FCR3.varCSA and detecting the presence of a signal at locations on the array that correspond to the positions at which the biological samples were disposed. Because the identity of the biological sample and its position on the array is known, an identification of the presence, concentration, and/or expression level of a nucleic acid encoding FCR3.varCSA is rapidly determined. That is, by employing labeled standards of a known concentration of a nucleic acid encoding FCR3.varCSA, (e.g., RNA), an investigator can accurately determine the concentration of a nucleic acid encoding FCR3.varCSA in a sample and from this information can assess the expression level of FCR3.varCSA. Conventional methods in densitometry can also be used to more accurately determine the concentration or expression level of a nucleic acid encoding FCR3.varCSA. These approaches are also easily automated using technology known to those of skill in the art of high throughput diagnostic analysis. Similar approaches can be used to determine the presence, absence, or amount of a nucleic acid encoding A4 tres DBL3- γ or ItG2-CS2 DBL2- γ .

[0156] Any addressable array technology known in the art can be employed with this aspect of the invention. One particular embodiment of polynucleotide arrays is known as Genechips™, and has been generally described in U.S. Pat. No. 5,143,854; PCT publications WO 90/15070 and 92/10092. These arrays are generally produced using mechanical synthesis methods or light directed synthesis methods, which incorporate a combination of photolithographic methods and solid phase oligonucleotide synthesis. (Fodor et al., *Science*, 251:767-777, (1991)). The immobilization of arrays of oligonucleotides on solid supports has been rendered possible by the development of a technology generally identified as "Very Large Scale Immobilized Polymer Synthesis" (VLSIPS™) in which, typically, probes are immobilized in a high density array on a solid surface of a chip. Examples of VLSIPS™ technologies are provided in U.S. Pat. Nos. 5,143,854 and 5,412,087 and in PCT Publications WO 90/15070, WO 92/10092 and WO 95/11995, which describe methods for forming oligonucleotide arrays through techniques such as light-directed synthesis techniques. In designing strategies aimed at providing arrays of nucleotides immobilized on solid supports, further presentation strategies were developed to order and display the oligonucleotide arrays on the chips in an attempt to maximize hybridization patterns and diagnostic information. Examples of such presentation strategies are disclosed in PCT Publications WO 94/12305, WO 94/11530, WO 97/29212 and WO 97/31256.

[0157] A wide variety of labels and conjugation techniques are known by those skilled in the art and can be used in various nucleic acid assays. There are several ways to produce labeled nucleic acids for hybridization or PCR (Polymerase Chain Reaction) including, but not limited to, oligolabeling, nick translation, end-labeling, or PCR amplification using a labeled nucleotide. Alternatively, a nucleic

acid encoding FCR3.varCSA, or any portion of it, can be cloned into a vector for the production of an mRNA probe. Such vectors are known in the art, are commercially available, and can be used to synthesize RNA probes in vitro by addition of an appropriate RNA polymerase such as T7, T3 or SP6 and labeled nucleotides. A number of companies such as Pharmacia Biotech (Piscataway N.J.), Promega (Madison Wis.), and U.S. Biochemical Corp (Cleveland Ohio) supply commercial kits and protocols for these procedures. Suitable reporter molecules or labels include those radionuclides, enzymes, fluorescent, chemiluminescent, or chromogenic agents, as well as, substrates, cofactors, inhibitors, magnetic particles and the like.

[0158] For diagnostic and prognostic purposes, nucleic acid probes having a sequence that complements a nucleic acid encoding FCR3.varCSA, fragments of FCR3.varCSA, A4 tres DBL3- γ , or ItG2-CS2 DBL2- γ or a portion thereof can be used to detect and quantitate gene expression in biological samples. Preferably, nucleic acid probes that are complementary to mRNA encoding FCR3.varCSA, fragments of FCR3.varCSA, A4 tres DBL3- γ , or ItG2-CS2 DBL2- γ are used to screen for polynucleotides present in blood. RNA-detection-based diagnostic assays, such as Northern hybridization, Northern dot blots, RNA in situ hybridization, and ELISA assays, are particularly useful to distinguish between the absence or presence of these mRNAs and/or to monitor mRNA levels during therapeutic intervention.

[0159] Embodiments also include the use of oligonucleotide sequences, antisense RNA and DNA molecules, and PNAs that complement a nucleic acid encoding FCR3.varCSA, fragments of FCR3.varCSA, A4 tres DBL3- γ , or ItG2-CS2 DBL2- γ for the determination of the concentration and expression level in the cells of a subject by RNA-based detection techniques. The form of such qualitative and/or quantitative methods can include Northern analysis, dot blot or other membrane-based technologies; PCR technologies; dip stick, pin, chip, and ELISA technologies. All of these techniques are well known in the art and are the basis of many commercially available diagnostic kits.

[0160] In one aspect, RNA probes complementary to FCR3.varCSA mRNA are used in assays that detect maternal malaria. Accordingly, the nucleotide sequence encoding FCR3.varCSA or a fragment thereof is used to design suitable RNA probes. The RNA probes are labeled by methods known in the art and are added to a DNase treated fluid or tissue sample from a subject under conditions suitable for the formation of hybridization complexes. Hybridization complexes are isolated or the sample is treated with an agent that removes unhybridized nucleic acids. After an incubation period, the sample is washed with a compatible fluid that optionally contains a dye (or other label requiring a developer) if the nucleotide has been labeled with an enzyme. After the compatible fluid is rinsed off, the dye is quantitated and compared with a standard. If the amount of dye in the sample is significantly elevated over that of a comparable control sample, the nucleotide sequence has hybridized with RNA in the sample, and the presence of elevated levels of RNA encoding FCR3.varCSA or a portion thereof in the sample indicates the presence of a FCR3.varCSA-related disease, such as cancer. A similar

approach can be used to determine the presence, absence or amount of an mRNA encoding A4 tres DBL3- γ , or ItG2-CS2 DBL2- γ .

[0161] Such assays can also be used to evaluate the efficacy of a particular therapeutic treatment regime in animal studies, in clinical trials, or in monitoring the treatment of an individual patient. In order to provide a basis for the diagnosis of disease, a normal or standard profile for FCR3.varCSA expression in isolated cells, extracts, or tissue can be established, for example. This is accomplished by combining body fluids or cell extracts taken from healthy subjects with RNA probes encoding FCR3.varCSA, or a portion thereof, under conditions suitable for hybridization. Standard hybridization can be quantified by comparing the values obtained for healthy and diseased subjects with a dilution series of FCR3.varCSA RNA run in the same experiment where a known amount of substantially purified FCR3.varCSA is used. Standard values obtained from samples from healthy and diseased subjects are then compared with values obtained from samples from the tested subjects. Deviation between standards and the values obtained for the subject tested establishes the presence or predilection for material malaria. A similar approach can be based on a profile constructed from the presence or amount of mRNA encoding A4 tres DBL3- γ , or ItG2-CS2 DBL2- γ from healthy and diseased individuals. w Additionally, PCR methods that can be used to quantitate the concentration and expression level of a particular molecule include radiolabeling (Melby P. C. et al. *J Immunol Methods* 159:235-44 (1993)) or biotinylating nucleotides (Duplaa C. et al. *Anal Biochem* 212:229-236 (1993)), coamplification of a control nucleic acid, and standard curves onto which the experimental results are interpolated. Quantitation of multiple samples can be processed more rapidly by running the assay in an ELISA format where the oligomer of interest is presented in various dilutions and a spectrophotometric or colorimetric response gives rapid quantitation. A definitive diagnosis of this type can allow health professionals to create a disease state profile for a patient, begin aggressive treatment for the malaria, and prevent further worsening of the condition. Similarly, further assays and reference to the changing disease state profile can help clinicians monitor the progress of a patient during treatment. That is, once a disease state is established, a therapeutic agent is administered and an initial disease state profile is generated. The assays above can be repeated on a regular basis to evaluate whether the values in the subject's disease state profile progresses toward or returns back to the initial disease state profile. Successive treatment profiles can be used to show the efficacy of treatment over a period of several days or several months.

[0162] As mentioned above, PCR technology can be used to identify and quantitate concentration and expression levels of a nucleic acid encoding FCR3.varCSA, fragments of FCR3.varCSA, A4 tres DBL3- γ , or ItG2-CS2 DBL2- γ . For amplification of mRNAs, it is within the scope of the present invention to reverse transcribe mRNA into cDNA followed by PCR (RT-PCR); or, to use a single enzyme for both steps as described in U.S. Pat. No. 5,322,770, or, to use Reverse Transcriptase Asymmetric Gap Ligase Chain Reaction (RT-AGLCR), as described by Marshall R. L. et al. (*PCR Methods and Applications* 4:80-84, 1994).

[0163] A variety of PCR techniques are familiar to those skilled in the art. For a review of PCR technology, see *Molecular Cloning to Genetic Engineering* White, B. A. Ed. in *Methods in Molecular Biology* 67: Humana Press, Totowa (1997) and the publication entitled "PCR Methods and Applications" (1991, Cold Spring Harbor Laboratory Press). In each of these PCR procedures, PCR primers on either side of the sequence to be amplified are added to a suitably prepared nucleic acid sample along with dNTPs and a thermostable polymerase such as Taq polymerase, Pfu polymerase, or Vent polymerase. The nucleic acid in the sample is denatured and the PCR primers are specifically hybridized to complementary nucleic acid sequences in the sample. The hybridized primers are extended. Thereafter, another cycle of denaturation, hybridization, and extension is initiated. The cycles are repeated multiple times to produce an amplified fragment containing the nucleic acid sequence between the primer sites. PCR has further been described in several patents including U.S. Pat. Nos. 4,683,195, 4,683,202 and 4,965,188.

[0164] The primers are selected to be substantially complementary to a portion of the sequence of a nucleic acid encoding FCR3.varCSA, fragments of FCR3.varCSA, A4 tres DBL3- γ , or ItG2-CS2 DBL2- γ and a portion of the sequence that complements the sequence of FCR3.varCSA, fragments of FCR3.varCSA, A4 tres DBL3- γ , or ItG2-CS2 DBL2- γ mRNA, thereby allowing the sequences between the primers to be amplified. The length of the primers for use with this aspect of the present invention be identical to most of the lengths of the nucleic acid embodiments provided previously. That is, primer length can be less than or equal to 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 125, 150, 175, 200, 250, 300, 350, 400, 500, 600, 700, 800, 900, 1000, 1500, 2000, 3000, 4000, 5000, 6000, 7000, 8000, 9000, 10,000, and 10,500 nucleotides. Preferably, however primers are 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29 and 30 nucleotides in length. Shorter primers tend to lack specificity for a target nucleic acid sequence and generally require cooler temperatures to form sufficiently stable hybrid complexes with the template. Longer primers are expensive to produce and can sometimes self-hybridize to form hairpin structures. The formation of stable hybrids depends on the melting temperature (T_m) of the DNA. The T_m depends on the length of the primer, the ionic strength of the solution and the G+C content. The higher the G+C content of the primer, the higher is the melting temperature because G:C pairs are held by three H bonds whereas A:T pairs have only two. The G+C content of the amplification primers of the present invention preferably ranges between 10 and 75%, more preferably between 35 and 60%, and most preferably between 40 and 55%. The appropriate length for primers under a particular set of assay conditions may be empirically determined by one of skill in the art.

[0165] The spacing of the primers determines the length of the segment to be amplified. In the context of the present invention amplified segments carrying nucleic acid sequence encoding fragments of FCR3.varCSA, fragments of FCR3.varCSA, A4 tres DBL3- γ , or ItG2-CS2 DBL2- γ can range in size from at least about 25 bp to 35 kbp. Amplifi-

cation fragments from 25-3000 bp are typical, fragments from 50-1000 bp are preferred and fragments from 100-600 bp are highly preferred. It will be appreciated that amplification primers for a nucleic acid encoding FCR3.varCSA, fragments of FCR3.varCSA, A4 tres DBL3- γ , or ItG2-CS2 DBL2- γ can be of any sequence that allows the specific amplification of any DNA fragment carrying nucleic acid sequence unique these molecules. Amplification primers can be labeled or immobilized on a solid support as described above.

[0166] The presence of FCR3.varCSA, A4 tres DBL3- γ , or ItG2-CS2 DBL2- γ protein can be detected by screening for the presence of the protein using conventional assays. For example, monoclonal antibodies immunoreactive with FCR3.varCSA can be used to screen biological samples for the presence, concentration, and expression level of FCR3.varCSA and, thereby, provide diagnostic information about FCR3.varCSA-related diseases. Such immunological assays can be done in many convenient formats.

[0167] In one embodiment, for example, antibodies are used to immunoprecipitate FCR3.varCSA, fragments of FCR3.varCSA, A4 tres DBL3- γ , or ItG2-CS2 DBL2- γ from solution and, in another embodiment, antibodies are used to react with FCR3.varCSA on Western or Immunoblots of a polyacrylamide gel. In desirable embodiments, antibodies are used to detect FCR3.varCSA, fragments of FCR3.varCSA, A4 tres DBL3- γ , or ItG2-CS2 DBL2- γ in paraffin or frozen sections, using immunocytochemical techniques. Favored embodiments for detecting FCR3.varCSA, fragments of FCR3.varCSA, A4 tres DBL3- γ , or ItG2-CS2 DBL2- γ include enzyme-linked immunosorbant assays (ELISA), radioimmunoassays (RIA), immunoradiometric assays (IRMA) and immunoenzymatic assays (IEMA), including sandwich assays using monoclonal and/or polyclonal antibodies. Exemplary sandwich assays are described by David et al., in U.S. Pat. Nos. 4,376,110 and 4,486,530.

[0168] In preferred protein-based diagnostic embodiments, antibodies of the present invention are attached to a support in an ordered array wherein a plurality of antibodies are attached to distinct regions of the support that do not overlap with each other. As with the nucleic acid-based arrays, the protein-based arrays are ordered arrays that are designed to be "addressable" such that the distinct locations are recorded and can be accessed as part of an assay procedure.

[0169] For example, in some embodiments, addressable antibody arrays comprise a plurality of antibodies that recognize FCR3.varCSA. These probes are joined to a support in different known locations. The knowledge of the precise location of each probe makes these "addressable" arrays particularly useful in binding assays. For example, an addressable array can comprise a support having several regions to which are joined a plurality antibody probes that recognize FCR3.varCSA. Proteins from a preparation of several biological samples from a plurality of human subjects or a plurality of tissues or fluids from a single subject are labeled by conventional approaches (e.g., radioactivity, calorimetrically, or fluorescently) and the labeled samples are applied to the array under conditions that permit binding. If a protein in the sample binds to an antibody probe on the array, then a signal will be detected at a position on the support that corresponds to the location of the antibody-

protein complex. Since the identity of each labeled sample is known and the region of the support on which the labeled sample was applied is known, an identification of the presence, concentration, and/or expression level is rapidly determined. That is, by employing labeled standards of a known concentration of FCR3.varCSA, an investigator can accurately determine the protein concentration of FCR3.varCSA in a sample and from this information can assess the expression level of FCR3.varCSA. Conventional methods in densitometry can also be used to more accurately determine the concentration or expression level of FCR3.varCSA. These approaches are easily automated using technology known to those of skill in the art of high throughput diagnostic analysis.

[0170] In another embodiment, an opposite approach to that presented above can be employed. Proteins present in biological samples (e.g., tissues or fluids from one or more subjects or one or more sources in a subject's body) can be disposed on a support so as to create an addressable array. Preferably, the protein samples are disposed on the support at known positions that do not overlap. The presence of a protein encoding FCR3.varCSA in each sample is then determined by applying labeled antibody probes that recognize epitopes of FCR3.varCSA and detecting a signal at locations on the array that correspond to the positions at which the biological samples were disposed. Because the identity of the biological sample and its position on the array is known, an identification of the presence, concentration, and/or expression level FCR3.varCSA is rapidly determined. That is, by employing labeled standards of a known concentration of FCR3.varCSA, an investigator can accurately determine the concentration of FCR3.varCSA in a sample and from this information can assess the expression level of FCR3.varCSA. Conventional methods in densitometry can also be used to more accurately determine the concentration or expression level of FCR3.varCSA. These approaches are also easily automated using technology known to those of skill in the art of high throughput diagnostic analysis. As detailed above, any addressable array technology known in the art can be employed with this aspect of the invention and display the protein arrays on the chips in an attempt to maximize antibody binding patterns and diagnostic information. Similar approaches can be used to detect the presence, absence, or amount of A4 tres DBL3- γ , or ItG2-CS2 DBL2- γ in a biological sample.

[0171] As discussed above, the presence or detection of FCR3.varCSA, fragments of FCR3.varCSA, A4 tres DBL3- γ , or ItG2-CS2 DBL2- γ can provide a diagnosis of a subject's disease state or predilection to disease and this information allows health professionals to create a disease state profile for a patient, begin aggressive treatment for the malaria, and prevent further worsening of the condition. Similarly, further assays and reference to the changing disease state profile can help clinicians monitor the progress of a patient during treatment. That is, once a disease state is established, a therapeutic agent is administered and an initial disease state profile is generated. The assays above can be repeated on a regular basis to evaluate whether the values in the subject's disease state profile progresses toward or returns back to the initial disease state profile. Successive treatment profiles can be used to show the efficacy of treatment over a period of several days or several months. Additional embodiments include the preparation of diagnostic kits comprising detection components such as antibodies

specific for FCR3.varCSA, fragments of FCR3.varCSA, A4 tres DBL3- γ , or ItG2-CS2 DBL2- γ or nucleic acid probes that detect the presence of these molecules. The detection component will typically be supplied in combination with one or more of the following reagents. A support capable of absorbing or otherwise binding RNA or protein will often be supplied. Available supports for this purpose include, but are not limited to, membranes of nitrocellulose, nylon or derivatized nylon that can be characterized by bearing an array of positively charged substituents, and Genechips™ or their equivalents. One or more enzymes, such as Reverse Transcriptase and/or Taq polymerase, can be furnished in the kit, as can dNTPs, buffers, or non-human polynucleotides like calf-thymus or salmon-sperm DNA. Results from the kit assays can be interpreted by a healthcare provider or a diagnostic laboratory. Alternatively, diagnostic kits are manufactured and sold to private individuals for self-diagnosis. The next section describes several embodiments that have therapeutic or prophylactic application or both.

[0172] Therapeutic and Prophylactic Applications

[0173] The varCSA modulating agents described herein are suitable for treatment of subjects either as a preventive measure to avoid maternal malaria, or as a therapeutic to treat subjects already afflicted with the disease. Although anyone could be treated with the agents of the invention as a prophylactic, the most suitable subjects are people at risk for maternal malaria. Such subjects include, but are not limited to, pregnant women living in regions of the world populated with *P. falciparum*.

[0174] The pharmacologically active compounds of this invention, including but not limited to, FCR3.varCSA, A4 tres DBL3- γ , ItG2-CS2 DBL2- γ , fragments of these molecules, nucleic acids encoding these molecules, and antibodies directed to these molecules, can be processed in accordance with conventional methods of galenic pharmacy to produce medicinal agents for administration to patients, e.g., mammals including humans. These FCR3.varCSA inhibitory agents can be incorporated into a pharmaceutical product with and without modification. Further, the manufacture of pharmaceuticals or therapeutic agents that deliver the inhibitory agent by several routes are aspects of the invention. For example, and not by way of limitation, DNA, RNA, and viral vectors having sequence encoding FCR3.varCSA or a polypeptide fragment of FCR3.varCSA are within the scope of aspects of the present invention. Nucleic acids encoding a desired FCR3.varCSA inhibitory agent can be administered alone or in combination with other varCSA inhibitory agents. Similarly nucleic acids encoding A4 tres DBL3- γ , or ItG2-CS2 DBL2- γ can be administered alone or in combination with other varCSA inhibitory agents.

[0175] The compounds described herein can be employed in admixture with conventional excipients, i.e., pharmaceutically acceptable organic or inorganic carrier substances suitable for parenteral, enteral (e.g., oral) or topical application that do not deleteriously react with the varCSA inhibitory agents. Suitable pharmaceutically acceptable carriers include, but are not limited to, water, salt solutions, alcohols, gum arabic, vegetable oils, benzyl alcohols, polyethylene glycols, gelatine, carbohydrates such as lactose, amylose or starch, magnesium stearate, talc, silicic acid, viscous paraffin, perfume oil, fatty acid monoglycerides and

diglycerides, pentaerythritol fatty acid esters, hydroxy methylcellulose, polyvinyl pyrrolidone, etc. The pharmaceutical preparations can be sterilized and if desired mixed with auxiliary agents, e.g., lubricants, preservatives, stabilizers, wetting agents, emulsifiers, salts for influencing osmotic pressure, buffers, coloring, flavoring and/or aromatic substances and the like that do not deleteriously react with the active compounds.

[0176] The effective dose and method of administration of a particular inhibitory agent formulation can vary based on the individual patient and the stage of the disease, as well as other factors known to those of skill in the art. Therapeutic efficacy and toxicity of such compounds can be determined by standard pharmaceutical procedures in cell cultures or experimental animals, e.g., ED50 (the dose therapeutically effective in 50% of the population) and LD50 (the dose lethal to 50% of the population). The dose ratio of toxic to therapeutic effects is the therapeutic index, and it can be expressed as the ratio, LD50/ED50. Pharmaceutical compositions that exhibit large therapeutic indices are preferred. The data obtained from cell culture assays and animal studies is used in formulating a range of dosage for human use. The dosage of such compounds lies preferably within a range of circulating concentrations that include the ED50 with little or no toxicity. The dosage varies within this range depending upon the dosage form employed, sensitivity of the patient, and the route of administration.

[0177] The exact dosage is chosen by the individual physician in view of the patient to be treated. Dosage and administration are adjusted to provide sufficient levels of the active moiety or to maintain the desired effect. Additional factors that may be taken into account include the prevalence of *P. falciparum* in the geographical vicinity of the patient, the severity of the disease state of the patient, age, and weight of the patient; diet, time and frequency of administration, drug combination(s), reaction sensitivities, and tolerance/response to therapy. Short acting pharmaceutical compositions are administered daily whereas long acting pharmaceutical compositions are administered every 2, 3 to 4 days, every week, or once every two weeks. Depending on half-life and clearance rate of the particular formulation, the pharmaceutical compositions of the invention are administered once, twice, three, four, five, six, seven, eight, nine, ten or more times per day.

[0178] Routes of administration of the varCSA inhibitory agents include, but are not limited to, transdermal, parenteral, gastrointestinal, transbronchial, and transalveolar. Transdermal administration is accomplished by application of a cream, rinse, gel, etc. capable of allowing the FCR3.varCSA inhibitory agent to penetrate the skin and enter the blood stream. Parenteral routes of administration include, but are not limited to, electrical or direct injection such as direct injection into a central venous line, intravenous, intramuscular, intraperitoneal or subcutaneous injection. Gastrointestinal routes of administration include, but are not limited to, ingestion and rectal. Transbronchial and transalveolar routes of administration include, but are not limited to, inhalation, either via the mouth or intranasally.

[0179] Compositions of the varCSA inhibitory agents suitable for transdermal administration include, but are not limited to, pharmaceutically acceptable suspensions, oils, creams, and ointments applied directly to the skin or incor-

porated into a protective carrier such as a transdermal device ("transdermal patch"). Examples of suitable creams, ointments, etc. can be found, for instance, in the Physician's Desk Reference. Examples of suitable transdermal devices are described, for instance, in U.S. Pat. No. 4,818,540 issued Apr. 4, 1989 to Chinen, et al.

[0180] Compositions of the varCSA inhibitory agents suitable for parenteral administration include, but are not limited to, pharmaceutically acceptable sterile isotonic solutions. Such solutions include, but are not limited to, saline and phosphate buffered saline for injection into a central venous line, intravenous, intramuscular, intraperitoneal, or subcutaneous injection of the FCR3.varCSA inhibitory agents.

[0181] Compositions of the varCSA inhibitory agents suitable for transbronchial and transalveolar administration include, but not limited to, various types of aerosols for inhalation. Devices suitable for transbronchial and transalveolar administration of the FCR3.varCSA inhibiting agents are also embodiments. Such devices include, but are not limited to, atomizers and vaporizers. Many forms of currently available atomizers and vaporizers can be readily adapted to deliver varCSA inhibitory agents.

[0182] Compositions of the varCSA inhibitory agents suitable for gastrointestinal administration include, but not limited to, pharmaceutically acceptable powders, pills or liquids for ingestion and suppositories for rectal administration. Due to the ease of use, gastrointestinal administration, particularly oral, is the preferred embodiment of the present invention.

[0183] Several methods of treatment and prevention of maternal malaria, which involve administration of the pharmaceutical embodiments of the invention are provided. In these aspects of the invention, FCR3.varCSA, fragments of FCR3.varCSA, A4 tres DBL3- γ , or ItG2-CS2 DBL2-7, nucleic acids encoding these molecules, and agents that interact with a varCSA-CSA complex are incorporated into pharmaceuticals and are administered to patients in need. Because aspects of the invention that incorporate a varCSA molecule or fragments thereof can both interrupt varCSA mediated adhesion and stimulate an immune response to these polypeptides, significant therapeutic and prophylactic benefit can be achieved by administration of these agents to patients in need. Thus, in some contexts, a therapeutic protocol can also be termed a method of vaccination. By one approach, a subject at risk for contracting maternal malaria or a subject infected with *P. falciparum* is identified by conventional techniques or the diagnostic assays described above and then a therapeutically or prophylactically beneficial amount of a varCSA molecule or fragment thereof is administered.

[0184] Although the invention has been described with reference to embodiments and examples, it should be understood that various modifications can be made without departing from the spirit of the invention. Accordingly, the invention is limited only by the following claims. All references cited herein are hereby expressly incorporated by reference.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 11

<210> SEQ ID NO 1

<211> LENGTH: 10628

<212> TYPE: DNA

<213> ORGANISM: Plasmodium falciparum

<400> SEQUENCE: 1

```

atgggggttct cctgcaaaata ttttattata aaaatgggga atgcagcatc atcattagag      60
ggagatgcta aaagccctat tataaaagaa agtcacaaaa gtgcaagaaa tgttttggaa      120
cgttatgcca aaaatataag acatccatca aaatatgcaa aagaacatgt ggattcgttg      180
aaaggggatt tgacgaaagc agaatttcgt ggtggtcctt ctacgccagt aaataagcat      240
aattattatt atccatattc atgtaattta gatcataagg aacatactaa tttacggtat      300
gatgatgtga atttgagaca tccttgccat ggtagagaac aaaaccgatt tgatgaagat      360
gaagaatctg aatgtggaaa taaaatacgt aattataaaa gaaaaaatga tgctatagcc      420
tgtgogccac ctagaagacg acatatgtgt gataaaaact tggaagctct aatgatata      480
aatacccaaa atattcatga tttattggga aatgtactag ttacagcaaa atacgaaggt      540
gaatcaattg ttaataatca tccacataaa ggaacttcag acgcttgtag tgctcttgca      600
cgaagttttg cagatatagg tgatattgta agaggaatag atatgtttaa accaaatgtc      660
catgacaaa tagaaacggg totccgagag gttttcaaga aaatacatga tggaatggaa      720

```

-continued

gatgaagtaa	aaaatgatta	caatcctgat	ggatctggaa	attattataa	attaagagaa	780
gcatggtgga	atgtgaatag	aaataaagta	tgggaagcta	taacatgtga	tgcatcatat	840
aaatctggat	atTTTTatgca	atcagaaagt	aatacaccat	tatTTTTcaaa	tcctaaatgc	900
ggccataaac	aaggaaaggt	tcctaccaat	ttagattatg	tcctcaata	tttacgttgg	960
ttcgacgaat	ggggagaaga	gttttgccga	aaaagaaata	ttaaattgaa	aaaggtcaag	1020
gactcctgtc	gtaatgacaa	agaacgctta	tattgtagtc	ataatggaca	tgattgtacg	1080
acaactatTT	ggaaaaaagg	tatTTTgcat	ttggataata	agtgtactga	ctgttcgact	1140
aaatgcaaag	TTTTTgaagt	ttggTTtaggg	aatcaacaag	aagcatttaa	aaaacaaaaa	1200
gaaaaaatatg	aaaaagaat	acaatcatat	ttatcgaacg	ataacaaatt	tgtcaataat	1260
attaatagtg	aatattataa	acaatTTTat	gaaaaactta	aggaaacgca	atatgcaact	1320
aatgacactt	TTTTaaatTT	actaaatgaa	ggaaagtatt	gtaaaggagg	attaccagga	1380
gaaaaggata	ttactTTTatc	taacagtgtc	gatgacaaag	ggatTTTTa	tcgttcagaa	1440
tattgccaag	tgTgtcccgga	ctgcggggtc	aaatgtgatg	gtataaaata	cacacacaaa	1500
tcagataatg	atcgtgaacg	tgtaaataat	gaagactata	aacctccatg	gggtgtgaag	1560
cctactaata	tcactgtcct	ttatagtTgt	aatgaacaag	gtgatattac	acaaaaatta	1620
gaaaatTTTT	gtaacagctc	aactaattac	aaagataaaa	ataatcaaaa	atgggaatgc	1680
tattataagg	atgaaaatat	aaatagatgt	aaactggaac	aaaactactga	aatcaataat	1740
gataatccta	agataaatatc	atTTcataat	TTTTTTgaaT	tatgggttac	atatttatta	1800
agggatacta	ttaagtggaa	tgacaaactt	aaaacttgta	taataatac	aaccacgcat	1860
tgtattgatg	aatgtaacag	aaattgctta	tgTTTTgaca	gatgggttaa	acaaaaagaa	1920
gaagaatgga	atagtataaa	gaaactgttc	acaaaaaaaa	agaatataca	gcaatcgtat	1980
tatagtaata	ttaataatct	TTTTgaaggt	tatTTTTTTa	aagttatgga	taaacttgac	2040
aaagatgaa	caaaatggaa	agaacttatg	gaaaatataa	aaagaaaaaa	aaatgagttt	2100
tcCaatTTTg	aaaataatag	ggactatTTa	gagaatgcaa	tagaactctt	gttagatcac	2160
ttaaaagaaa	ctgccacgat	atgTaaagac	aataatacaa	acgaagcatg	tgaaacatcc	2220
cataatgcaa	caacaaacc	gtgtgtTtaa	cctcgtggag	gcacgcaacc	cactaaaaat	2280
ataaaaagaaa	tagcacaata	cttTaaaagg	agTgcatacg	aggaaacacg	aaatcgtggT	2340
cttcataaat	Tgaaaggaaa	ggcacacgaa	ggtatatata	aacgtggggg	taggagaaaag	2400
gacttcaagg	acaatTTatg	tagaataatg	ataaaacatt	ctaatcgtaa	tcttggTTTT	2460
tcaaatggac	catgtgatgg	caaaggcaca	ggtgatggta	tacaaacaag	atTTgtcgta	2520
ggaaactgaat	gggaagtgga	tcgggaacac	atgcgTaaag	atcacgaaga	Tgttattatg	2580
cctcctagaa	gacgacatat	atgtacatcc	aatTTggaac	atTTacaaac	ggatgatcac	2640
ccactTaatg	gtaatattgt	Tgatgattta	gtTaaataat	cctTTTTTgg	ggatgttctt	2700
ctatcagcaa	aatatgaagc	aaacaagata	atacgaatgt	ataaagaaaa	gaataaccta	2760
aaggGCCca	aagaagTaac	TgaccCaaaa	caccagacaa	ctatctgtcg	agctatacgt	2820
tacagTTTTg	cagatatag	Tgatataat	cagggaagag	atctctggga	aagaaacggT	2880
gacatgTtaa	agctgcaagg	acatTTgTaa	actgTTTTTg	gTaatataca	taagTcactc	2940
aaaggCaag	gaaatgataa	atataatgat	gatgCCCCa	aatatTTaaa	atTgaggGaa	3000

-continued

aattggtggg aagctaatag agccaaagta tgggaagcca tgaatgtga tataaaatat	3060
ttgaaggata aatcgggaca ccaatcaaca caaagtagtt attgcgata tagtgatcat	3120
acaccattgg atgattatat cccacaaaa ttaagatgga tgaccgaatg ggcagaatgg	3180
tactgcaagg tgcagaaaa ggagtatgat aagttgaagg agaagtgtaa ggagtgtaa	3240
gataaggata atggccaagg ctgtacgaaa gagagtggta caggtgtac gaagtgcaca	3300
gaagcttgta atgaataata tgatataata ggattatgga aagaacaatg gaataata	3360
tcagataaat acaagaat acatgaacaa gcacaaatgt ctgttagtaa tagtggtatt	3420
gaagcttcca gtactgcaa aaatcatata gacaggaatg ttattgaatt tttgctgaa	3480
ttataccaac aaaatggtg caaaagtaat aaaagtgga ctagtatga aagtgtgtc	3540
attggtacta acaccacgta tgaatgtt ggagcatatc tccatgatac agaaaat	3600
gatgattgtc agtcacaaa tgagttttgt gatgaaaaa gtgatggtaa ggataacgaa	3660
aaatatgcct ttagagataa accacaggac catgatggtg cgtgtggtg taaaagtgga	3720
tcgaaaccga caaggtgaca gataaaacg aaaaaaaag cggagaaaa ggatacggaa	3780
tgtaaacag tgaatgatat acttaagaa aacgatgga agaacaagt agaagattgt	3840
catcaaaaa agaatagtaa tggatatcc gattggcaat gcgaaatat aaatttagt	3900
gaagaccctc gtgtgtgtat gcccctaga agacaaaagt tatgctaca tttcttggca	3960
aatgataatg aataaaaaa attacaatca caagtaatt taaaagaagc tttcatcaa	4020
tctgcagcag cagaacatt cttctcatg tattattata aaagtaagga tggtaagga	4080
aatgaactcg ataaagaat aaaaagaagc aaaattcctc ccgattttt gagatccatg	4140
ttctacacat ttggagatta tagagattt ttatttgaa cagatatatc aaaaggtcat	4200
ggtgagggaa gtaactaaa agagcaata gattctctt tcaaaaatg tgaccaaaaa	4260
tctcctaag gaaaacacg ccaagaatg tggacagaac atagtcatga gatatggaa	4320
gctatgctat gtgcactagt aaaaattgg gcaaaaaag atgattttac cgaaaactac	4380
ggttaacaaca acgtcaaat tagtgacaaa agcaccactt tggaggaatt tgcaaacga	4440
ccccagttt tacgatggct aaccgaatg tacgacgact attgctatac acgacaaaa	4500
tatttgaag atgtgcagga aaaaatgta tcaaatgacc aattgaagtg tgatacagaa	4560
tgtaataaga aatgcagga ctacgttaa tatatgaaa aaaaaaaga gtggattcca	4620
caagataaat attacaagga tgaacgcgac aaaaaagat tcgatagaca acacattggt	4680
gtaatggta cagactatac tgaacgaat gcaacagatt actgaacag gaaatttact	4740
gctagtgtg gtgataagcc tggaaagtgc tctgtgtac aaagaaatat acaattgta	4800
gaaaaacag cttactatga tgcgcacaaa cattgtgggt gcacaaaatt tattgaaat	4860
gacgacaaat atactaacat ttcgagtaa gataagtga aaggattagt aaaggagca	4920
aacacaggtg ctattaagt gcaaaacaaa ggtcctaata actacaataa cttgaaagaa	4980
ttgactgaag atgtgctttt tccttctcgt cgactacgta tatgtttca tgcattggat	5040
ggcaattata cagatccaga agttaagat gaaaatgggt tgcgaaaaag attgatgaa	5100
gtggcggcaa cggaaaggta caatttgggt caatactaca aagaaaaaa agaaaaagag	5160
aaaataaaaa cgtcggatgc gcacaaatat tcttatgagg tccgccttg tagtgctatg	5220
aaatatagtt tttatgatt aagagatata attctaggta ttgataattt ggaagatgaa	5280

-continued

aaacaaaaga	cogaggaaaa	tttgaagaaa	atatttaaca	aaaatggaac	atcagttggc	5340
aaaggaagtg	atagtactac	aggaaatccc	ggtagtactg	cgcgaaaatt	tttctggaac	5400
gaaaaataagg	aatgtgtgtg	gaacgcaatg	atatgcgggt	acaaacgtgg	tagggatgat	5460
ggaaatagtg	gaaatagtc	aagaagtgat	gaagatctaa	aaaaatgtgg	ttctgtacct	5520
tcagatgatg	attatcctat	gggaaaaaat	cgcgatgaag	gtactgcgta	tcagtttctt	5580
cgatggtttg	ccgaatgggg	tgaagatfff	tgcaaacata	aagaaaagga	attggagaaa	5640
ttggtagggg	cgtgtaatga	ttatacttgt	ggtgataatg	aagataaaag	aaagaaatgt	5700
acagatgcgt	gtacacaata	taaaaatff	attagtgagt	ggaaccaca	gtatgaaaa	5760
caaatcaaaa	aatatggta	gaataaagac	aaaatatatt	ccgagcatcc	tgtggcaaaa	5820
gatgcagagg	acgctcgcga	atatttagac	aaacaattaa	aaaaatffg	tgaaaataaa	5880
agtggagatt	gtgaatata	gtgtatgaaa	gatgtgtcca	cacagcgatt	aactgatggt	5940
aatagtcaaa	atatgcccg	atcattagac	gatgaaccaa	aagaagtga	aggaaagtgt	6000
aattgtcaag	tgccacgagg	tccaccacgt	gtacgaagg	aaacaccgtc	accacgggta	6060
tcactgatat	caaaagcgc	ggcatcgaaa	aaagaagcga	aaacagcgcc	gcctacaaaa	6120
cagccgaaaa	aagtggaaaa	tctaacaaca	gaaatgcgag	cacaaacacg	aaccgacga	6180
gcagcacaa	aaacacgaaa	acgaacatca	acagcaacaa	caacagaatc	tgacgtgggc	6240
acaatggtaa	aggccattct	ttcgaataaa	ccagatagca	gggggtggaat	agagggttgt	6300
aatccaaaa	cgtagggaca	atatacctaaa	tggggttgta	ttgtaggtaa	gtctaaagaa	6360
aatgaaaatg	gcataatgat	gcctcctag	agaaaaaat	tatgtataaa	taatatacaa	6420
tatttaaatt	atgaaactga	aaataagcgt	gacaatgata	taaaagaggc	ttttattaaa	6480
tgtgcagcaa	tagaaactca	atffttgtgg	ttaaaatata	taattgaaaa	tcctgcagca	6540
gaaaatgaat	tgcaaatgg	aacaattcca	gatgaattta	aaagaataat	gtattataca	6600
tatggtgatt	ataaagatat	gttttttgga	actgatattt	ctaatgataa	aaaaattata	6660
actgtaacaa	atagtgtaac	aaccattctc	aatgaaaata	ataagaaaa	acaggataaa	6720
aaaaaagatg	aagaattacg	taaaatattt	tgggagaaaa	ataaaaaatt	tatttgggaa	6780
ggaatgatat	atggattaac	ttatcatctc	acagacgaaa	acgaaaaaga	aaaaattaga	6840
gataattacc	agtacaatga	catgacacaa	ctgacgcctt	cccttgaaga	gtttgtaaaa	6900
aggccccaat	ttttgagatg	gttcacagaa	tgggcagaag	aatfftgtaa	taagaggaag	6960
gaacagttgt	taaaattgga	ggcgggctgt	aaggaatatg	agtgtaatgg	tagtaatgac	7020
ggtaagacac	aagaatgtgc	agaggcgtgt	gtaacatata	aaaatfftat	taagaagtgg	7080
aaaactgaat	atgaagaca	aagagaaaa	ttcaaaaagg	ataaagatgg	caaaaagtat	7140
aaggattata	cttctactga	aagagacata	gagaaggcaa	catgtgctca	tgaatattta	7200
aacatgaaat	taaaagaatt	atgtggcaat	aaggattggt	cttgtatgca	aaaaccttct	7260
tcacaactac	caaaaaacac	acaacaatca	caatcatccg	atgctaata	tatgccagaa	7320
tcgctggatt	atgttcctga	agaatffaac	aagtgtgagt	gtcctgaact	ttcaaaaaag	7380
ggatctatga	ttcatacaaa	aaaaattact	gaacctaaaa	tacctatgaa	ttgtgtagag	7440
aaagcagcat	attatffatc	taaagaagca	gaaaataata	tggatattac	cttgaaggaa	7500
aaatffatac	ctattgagtc	tacaaaggaa	aaggaaaagta	aaaatagttg	gactaataat	7560

-continued

aatccttgcg	atcctaagaa	accttatgca	cctgataaat	atataggaag	aagaaccct	7620
tgtgaaaata	gagaagaaa	tcgttttaag	gtagattatg	aatggaaatg	ttacaaaaat	7680
tcaaagtctt	atcaggagaa	aaaaagagta	tgtgtacctc	caagaagaga	acatatgtgc	7740
ttaaggaatt	tagatgaaat	taaaattgaa	agacttaagg	atagtaatta	tctcctaaaa	7800
atggttcgto	gaactgcacg	aaatgaagga	atagacataa	taaaaactt	caactcagag	7860
aacgggtgcg	caatgaatcc	aatatgtgat	actatgaaat	atagtttcgc	tgatctgggt	7920
gacatagtta	gaggaacaga	tatgttacga	attggtggtt	acttacctcc	cgtagaaata	7980
aaattatata	aggtttttga	atacatatat	ggaaaatgga	gaaataaaaa	taaaggtaga	8040
aataaataca	acgatgtaca	aacgtttcgt	tctgcttggt	gggatgctaa	tagaaaagat	8100
atttgaaag	caatgacgtg	caaagcacca	gaagatgcaa	aactttttag	aaaaggaaga	8160
atggatggat	ttgaacgcgt	aacattaata	caagataagt	gtggacataa	ggacgatcca	8220
cctggtgatg	attatatacc	tcaacgggtt	cgatggatga	ctgaatggtc	tgaatattat	8280
tgtaaagcac	tgatggaaga	attggaaaaa	tttaaaaaat	catgtgatca	ctgtaaaaca	8340
tctgacagat	gcaagaatga	ttatgatgaa	aataagtgtg	aacagtgtaa	aacgagatgt	8400
caagaatata	aaaattttgt	tcttaaatgg	aatctctat	tcgatataca	atcaaataaa	8460
tacaaagaat	tgtatgaaca	accaatatat	acaaaaatct	ctaactatga	tcatgttcaa	8520
aattttgtac	aaaagttgaa	aacttttaaa	agtgaatggt	ctgttgagag	cttttctgaa	8580
tatcttcatg	aaacaagtaa	gtgtttgaat	tataaattta	atgaaaatga	tggttcttcc	8640
aatatacgaa	catatgcttt	cgaagaaaca	ccaaaaagt	ataaagaagc	ttgcagttgt	8700
acactacctt	ctaagaatcc	attggataat	tgtcctaccg	atcaaaacaa	agatggatgt	8760
aaggaattac	aaacttttac	cttctgctcg	aagaatgatt	atgataataa	tcttgataat	8820
tggaacgcgt	accttgttct	taatagttca	gatgataaca	aagggtgatt	gattcctcca	8880
agaagaagac	atztatgtac	aagacctatc	actgcatata	attatagaaa	aggtgataaa	8940
gaaattttta	aaaaaaaaact	tcttacttct	gctttcagtc	aaggacaatt	gttaggtcaa	9000
aaatataaat	cggaagaaga	gttgtgcttt	gaggcaatga	aatatagtta	tgcagattat	9060
tccgatataa	ttaaaggaac	tgatatgatg	gacacttcat	tatctgaaaa	aattaaaaaa	9120
atatttgaaa	catcaaatga	agcaaccgaa	aatcgtaaaa	catggtggga	aaataataga	9180
cgtcagatag	ggcacgctat	gttatgtgga	tataaaattg	ctaactcaaa	agtaacatta	9240
gatgaaggat	ggtgtcaatt	accaaaggat	gaagaaacta	atcagtttct	tcgttggtta	9300
attgaatggg	caaagcaagc	atgtaaggaa	aagaaacatg	taagtgattc	attaaaaaca	9360
aaatgtcctc	gttcaaacga	agataatttt	gaagcgtcag	aattattaag	acaacctgga	9420
tgtcagaatg	atattagaaa	atatattagc	ttgaatatat	tgataaaaaa	tacaatggaa	9480
aatcctaata	taaaatataa	gcaattaaaa	gatcaatctt	caggtaatat	agacaataaa	9540
ccatctgaag	aaaatgttca	gtcatatata	aaatcaaaag	attctcaatg	cgctttggag	9600
ttaaagtata	taaatgaaat	agttacagga	acaaaaata	atgaaaataa	tgaattcaaa	9660
gaagtactaa	aaaaattata	tctgtgttta	tattttgttg	aagatgaaac	acacaaaaat	9720
catgtactag	atggaaatat	aaaagaagaa	gagcaaacag	ttcgtcctaa	agcactctat	9780
ttctttacac	ccatgtaga	ttctttctat	caagcacctt	tattctcaac	acatcgagta	9840

-continued

```

gcacaatatg atcctaaaaa tgatatattg aaaagtagta tctctgttgt tattgtatcg 9900
gcgttagggt tgatagcgct tcatttcgat aagaaaaaat tcaaatcgtc tgtggacttg 9960
ttgcgtatac tgaatatccc gcaaggagag tatggaatgc ctacgttga atccaaaaat 10020
aggtacatac catatagaag tgggtccatg aaaggcaaaa catatatata tatggaagga 10080
gatactagtg gagatgaaga taaatatatg tgggacttat cttctctga tattacttca 10140
tccgaaagtg agtgaaga attggatatt aatgatatat atgtaccagg tagtcctaaa 10200
tataaaacat tgatagaagt agtactagaa ccatcaaaaa gggatatacc aagtgatgat 10260
acaccaagta atgatacacc acgtacgaat agatttattg atgatgaatg gaatgaactg 10320
aaacatgatt ttgtatctca atatttacca aatacagaac caaataataa ttacaaaagt 10380
gcagatatto caatgaatac agaacctaactcttatt ctgataatcc tgaagaaaaa 10440
ccttttatta tatctattca tgatagggat ttatatactg gaaagaaat tagttataat 10500
attaatatga gtactaatac taataatgat attccaatga atgctagaaa tgattcttat 10560
agaggtatag atttaattaa tgattcacta gtggtgctaa acctattgat atatatgatg 10620
aagtattg 10628
    
```

```

<210> SEQ ID NO 2
<211> LENGTH: 3542
<212> TYPE: PRT
<213> ORGANISM: Plasmodium falciparum
    
```

<400> SEQUENCE: 2

```

Met Gly Phe Ser Cys Lys Tyr Phe Ile Ile Lys Met Gly Asn Ala Ala
 1          5          10          15
Ser Ser Leu Glu Gly Asp Ala Lys Ser Pro Ile Ile Lys Glu Ser His
          20          25          30
Lys Ser Ala Arg Asn Val Leu Glu Arg Tyr Ala Lys Asn Ile Arg His
          35          40          45
Pro Ser Lys Tyr Ala Lys Glu His Val Asp Ser Leu Lys Gly Asp Leu
          50          55          60
Thr Lys Ala Glu Phe Arg Gly Gly Pro Ser Thr Pro Val Asn Lys His
          65          70          75          80
Asn Tyr Tyr Tyr Pro Tyr Pro Cys Asn Leu Asp His Lys Glu His Thr
          85          90          95
Asn Leu Arg Tyr Asp Asp Val Asn Leu Arg His Pro Cys His Gly Arg
          100          105          110
Glu Gln Asn Arg Phe Asp Glu Asp Glu Glu Ser Glu Cys Gly Asn Lys
          115          120          125
Ile Arg Asn Tyr Lys Arg Lys Asn Asp Ala Ile Ala Cys Ala Pro Pro
          130          135          140
Arg Arg Arg His Met Cys Asp Lys Asn Leu Glu Ala Leu Asn Asp Ile
          145          150          155          160
Asn Thr Gln Asn Ile His Asp Leu Leu Gly Asn Val Leu Val Thr Ala
          165          170          175
Lys Tyr Glu Gly Glu Ser Ile Val Asn Asn His Pro His Lys Gly Thr
          180          185          190
Ser Asp Ala Cys Thr Ala Leu Ala Arg Ser Phe Ala Asp Ile Gly Asp
          195          200          205
Ile Val Arg Gly Ile Asp Met Phe Lys Pro Asn Val His Asp Lys Val
    
```

-continued

210		215		220		
Glu Thr Gly Leu Arg	Glu Val Phe Lys Lys	Ile His Asp Gly Met Glu	225	230	235	240
Asp Glu Val Lys Asn Asp Tyr Asn Pro Asp Gly Ser Gly Asn Tyr Tyr	245	250	255			
Lys Leu Arg Glu Ala Trp Trp Asn Val Asn Arg Asn Lys Val Trp Glu	260	265	270			
Ala Ile Thr Cys Asp Ala Ser Tyr Lys Ser Gly Tyr Phe Met Gln Ser	275	280	285			
Glu Ser Asn Thr Pro Leu Phe Ser Asn Pro Lys Cys Gly His Lys Gln	290	295	300			
Gly Lys Val Pro Thr Asn Leu Asp Tyr Val Pro Gln Tyr Leu Arg Trp	305	310	315			
Phe Asp Glu Trp Gly Glu Glu Phe Cys Arg Lys Arg Asn Ile Lys Leu	325	330	335			
Lys Lys Val Lys Asp Ser Cys Arg Asn Asp Lys Glu Arg Leu Tyr Cys	340	345	350			
Ser His Asn Gly His Asp Cys Thr Thr Thr Ile Trp Lys Lys Gly Ile	355	360	365			
Leu His Leu Asp Asn Lys Cys Thr Asp Cys Ser Thr Lys Cys Lys Val	370	375	380			
Phe Glu Val Trp Leu Gly Asn Gln Gln Glu Ala Phe Lys Lys Gln Lys	385	390	395			400
Glu Lys Tyr Glu Lys Glu Ile Gln Ser Tyr Leu Ser Asn Asp Asn Lys	405	410	415			
Phe Val Asn Asn Ile Asn Ser Glu Tyr Tyr Lys Gln Phe Tyr Glu Lys	420	425	430			
Leu Lys Glu Thr Gln Tyr Ala Thr Asn Asp Thr Phe Leu Asn Leu Leu	435	440	445			
Asn Glu Gly Lys Tyr Cys Lys Gly Gly Leu Pro Gly Glu Lys Asp Ile	450	455	460			
Thr Phe Thr Asn Ser Ala Asp Asp Lys Gly Ile Phe Tyr Arg Ser Glu	465	470	475			480
Tyr Cys Gln Val Cys Pro Asp Cys Gly Val Lys Cys Asp Gly Ile Lys	485	490	495			
Tyr Thr His Lys Ser Asp Asn Asp Arg Glu Arg Val Asn Asn Glu Asp	500	505	510			
Tyr Lys Pro Pro Trp Gly Val Lys Pro Thr Asn Ile Thr Val Leu Tyr	515	520	525			
Ser Gly Asn Glu Gln Gly Asp Ile Thr Gln Lys Leu Glu Asn Phe Cys	530	535	540			
Asn Ser Ser Thr Asn Tyr Lys Asp Lys Asn Asn Gln Lys Trp Glu Cys	545	550	555			560
Tyr Tyr Lys Asp Glu Asn Ile Asn Arg Cys Lys Leu Glu Gln Asn Thr	565	570	575			
Glu Ile Asn Asn Asp Asn Pro Lys Ile Ile Ser Phe His Asn Phe Phe	580	585	590			
Glu Leu Trp Val Thr Tyr Leu Leu Arg Asp Thr Ile Lys Trp Asn Asp	595	600	605			
Lys Leu Lys Thr Cys Ile Asn Asn Thr Thr Thr His Cys Ile Asp Glu	610	615	620			

-continued

Cys Asn Arg Asn Cys Leu Cys Phe Asp Arg Trp Val Lys Gln Lys Glu
 625 630 635 640
 Glu Glu Trp Asn Ser Ile Lys Lys Leu Phe Thr Lys Lys Lys Asn Ile
 645 650 655
 Gln Gln Ser Tyr Tyr Ser Asn Ile Asn Asn Leu Phe Glu Gly Tyr Phe
 660 665 670
 Phe Lys Val Met Asp Lys Leu Asp Lys Asp Glu Ala Lys Trp Lys Glu
 675 680 685
 Leu Met Glu Asn Ile Lys Arg Lys Lys Asn Glu Phe Ser Asn Leu Glu
 690 695 700
 Asn Asn Arg Asp Tyr Leu Glu Asn Ala Ile Glu Leu Leu Leu Asp His
 705 710 715 720
 Leu Lys Glu Thr Ala Thr Ile Cys Lys Asp Asn Asn Thr Asn Glu Ala
 725 730 735
 Cys Glu Thr Ser His Asn Ala Thr Thr Asn Pro Cys Val Lys Pro Arg
 740 745 750
 Gly Gly Thr Gln Pro Thr Lys Asn Ile Lys Glu Ile Ala Gln Tyr Phe
 755 760 765
 Lys Arg Ser Ala Tyr Glu Glu Ala Arg Asn Arg Gly Leu His Lys Leu
 770 775 780
 Lys Gly Lys Ala His Glu Gly Ile Tyr Lys Arg Gly Gly Arg Arg Lys
 785 790 795 800
 Asp Phe Lys Asp Asn Leu Cys Arg Ile Met Ile Lys His Ser Asn Arg
 805 810 815
 Asn Leu Gly Phe Ser Asn Gly Pro Cys Asp Gly Lys Gly Thr Gly Asp
 820 825 830
 Gly Ile Gln Thr Arg Phe Val Val Gly Thr Glu Trp Glu Val Asp Pro
 835 840 845
 Glu His Met Arg Lys Asp His Glu Asp Val Ile Met Pro Pro Arg Arg
 850 855 860
 Arg His Ile Cys Thr Ser Asn Leu Glu His Leu Gln Thr Asp Asp His
 865 870 875 880
 Pro Leu Asn Gly Asn Ile Val Asp Asp Leu Val Asn Asn Ser Phe Leu
 885 890 895
 Gly Asp Val Leu Leu Ser Ala Lys Tyr Glu Ala Asn Lys Ile Ile Arg
 900 905 910
 Met Tyr Lys Glu Lys Asn Asn Leu Lys Gly Pro Lys Glu Val Thr Asp
 915 920 925
 Pro Lys His Gln Thr Thr Ile Cys Arg Ala Ile Arg Tyr Ser Phe Ala
 930 935 940
 Asp Ile Gly Asp Ile Ile Arg Gly Arg Asp Leu Trp Glu Arg Asn Gly
 945 950 955 960
 Asp Met Val Lys Leu Gln Gly His Leu Glu Thr Val Phe Gly Asn Ile
 965 970 975
 His Lys Ser Leu Lys Gly Lys Gly Asn Asp Lys Tyr Asn Asp Asp Ala
 980 985 990
 Pro Lys Tyr Leu Lys Leu Arg Glu Asn Trp Trp Glu Ala Asn Arg Ala
 995 1000 1005
 Lys Val Trp Glu Ala Met Lys Cys Asp Ile Lys Tyr Leu Lys Asp Lys
 1010 1015 1020

-continued

Ser Gly His Gln Ser Thr Gln Ser Ser Tyr Cys Gly Tyr Ser Asp His
 1025 1030 1035 1040
 Thr Pro Leu Asp Asp Tyr Ile Pro Gln Lys Leu Arg Trp Met Thr Glu
 1045 1050 1055
 Trp Ala Glu Trp Tyr Cys Lys Val Gln Lys Lys Glu Tyr Asp Lys Leu
 1060 1065 1070
 Lys Glu Lys Cys Lys Glu Cys Lys Asp Lys Asp Asn Gly Gln Gly Cys
 1075 1080 1085
 Thr Lys Glu Ser Gly Thr Gly Cys Thr Lys Cys Thr Glu Ala Cys Asn
 1090 1095 1100
 Glu Tyr Asn Asp Ile Ile Gly Leu Trp Lys Glu Gln Trp Asn Ile Ile
 1105 1110 1115 1120
 Ser Asp Lys Tyr Lys Glu Leu His Glu Gln Ala Gln Met Ser Val Ser
 1125 1130 1135
 Asn Ser Gly Ile Glu Ala Ser Ser Thr Ala Lys Asn His Ile Asp Arg
 1140 1145 1150
 Asn Val Ile Glu Phe Leu Ser Glu Leu Tyr Gln Gln Asn Gly Gly Lys
 1155 1160 1165
 Ser Asn Lys Ser Gly Thr Ser Asp Glu Ser Ala Val Ile Gly Thr Asn
 1170 1175 1180
 Thr Thr Tyr Glu Asn Val Gly Ala Tyr Leu His Asp Thr Gly Asn Phe
 1185 1190 1195 1200
 Asp Asp Cys Gln Ser Gln Asn Glu Phe Cys Asp Glu Lys Ser Asp Gly
 1205 1210 1215
 Lys Asp Asn Glu Lys Tyr Ala Phe Arg Asp Lys Pro Gln Asp His Asp
 1220 1225 1230
 Gly Ala Cys Gly Cys Lys Ser Gly Ser Lys Pro Thr Arg Val Gln Ile
 1235 1240 1245
 Lys Thr Lys Lys Lys Ala Glu Glu Lys Asp Thr Glu Cys Lys Thr Val
 1250 1255 1260
 Asn Asp Ile Leu Lys Glu Asn Asp Gly Lys Lys Gln Val Glu Asp Cys
 1265 1270 1275 1280
 His Pro Lys Lys Asn Ser Asn Gly Tyr Pro Asp Trp Gln Cys Gly Asn
 1285 1290 1295
 Ile Asn Leu Val Glu Asp Pro Arg Val Cys Met Pro Pro Arg Arg Gln
 1300 1305 1310
 Lys Leu Cys Val His Phe Leu Ala Asn Asp Asn Glu Ile Lys Lys Leu
 1315 1320 1325
 Gln Ser Gln Val Asn Leu Lys Glu Ala Phe Ile Lys Ser Ala Ala Ala
 1330 1335 1340
 Glu Thr Phe Phe Ser Trp Tyr Tyr Tyr Lys Ser Lys Asp Gly Glu Gly
 1345 1350 1355 1360
 Asn Glu Leu Asp Lys Glu Leu Lys Glu Gly Lys Ile Pro Pro Ala Phe
 1365 1370 1375
 Leu Arg Ser Met Phe Tyr Thr Phe Gly Asp Tyr Arg Asp Phe Leu Phe
 1380 1385 1390
 Gly Thr Asp Ile Ser Lys Gly His Gly Glu Gly Ser Lys Leu Lys Glu
 1395 1400 1405
 Gln Ile Asp Ser Leu Phe Lys Asn Gly Asp Gln Lys Ser Pro Asn Gly
 1410 1415 1420
 Lys Thr Arg Gln Glu Trp Trp Thr Glu His Ser His Glu Ile Trp Glu

-continued

1425	1430	1435	1440
Ala Met Leu Cys Ala Leu Val Lys Ile Gly Ala Lys Lys Asp Asp Phe	1445	1450	1455
Thr Glu Asn Tyr Gly Tyr Asn Asn Val Lys Phe Ser Asp Lys Ser Thr	1460	1465	1470
Thr Leu Glu Glu Phe Ala Lys Arg Pro Gln Phe Leu Arg Trp Leu Thr	1475	1480	1485
Glu Trp Tyr Asp Asp Tyr Cys Tyr Thr Arg Gln Lys Tyr Leu Lys Asp	1490	1495	1500
Val Gln Glu Lys Cys Lys Ser Asn Asp Gln Leu Lys Cys Asp Thr Glu	1505	1510	1520
Cys Asn Lys Lys Cys Glu Asp Tyr Val Lys Tyr Met Lys Lys Lys Lys	1525	1530	1535
Glu Trp Ile Pro Gln Asp Lys Tyr Tyr Lys Asp Glu Arg Asp Lys Lys	1540	1545	1550
Arg Phe Asp Arg Gln His Ile Gly Val Met Val Thr Asp Tyr Thr Gly	1555	1560	1565
Thr Asn Ala Thr Asp Tyr Leu Asn Arg Lys Phe Thr Ala Ser Cys Gly	1570	1575	1580
Asp Lys Pro Gly Ser Ala Ser Val Val Gln Arg Asn Ile Gln Leu Leu	1585	1590	1600
Glu Lys Gln Ala Tyr Tyr Asp Ala Asp Lys His Cys Gly Cys Thr Lys	1605	1610	1615
Phe Ile Glu Asn Asp Asp Lys Tyr Thr Asn Ile Ser Ser Lys Asp Lys	1620	1625	1630
Cys Lys Gly Leu Val Lys Glu Ala Asn Thr Gly Ala Ile Lys Trp Gln	1635	1640	1645
Asn Lys Gly Pro Asn Asn Tyr Asn Asn Leu Lys Glu Leu Thr Glu Asp	1650	1655	1660
Val Leu Phe Pro Ser Arg Arg Leu Arg Ile Cys Phe His Ala Leu Asp	1665	1670	1680
Gly Asn Tyr Thr Asp Pro Glu Val Lys Asp Glu Asn Gly Leu Arg Lys	1685	1690	1695
Arg Leu Met Glu Val Ala Ala Thr Glu Gly Tyr Asn Leu Gly Gln Tyr	1700	1705	1710
Tyr Lys Glu Lys Lys Glu Lys Glu Lys Ile Lys Thr Ser Asp Ala His	1715	1720	1725
Lys Tyr Ser Tyr Glu Val Pro Pro Cys Ser Ala Met Lys Tyr Ser Phe	1730	1735	1740
Tyr Asp Leu Arg Asp Ile Ile Leu Gly Ile Asp Asn Leu Glu Asp Glu	1745	1750	1760
Lys Gln Lys Thr Glu Glu Asn Leu Lys Lys Ile Phe Asn Lys Asn Gly	1765	1770	1775
Thr Ser Val Gly Lys Gly Ser Asp Ser Thr Thr Gly Asn Pro Gly Ser	1780	1785	1790
Thr Ala Arg Lys Phe Phe Trp Asn Glu Asn Lys Glu Cys Val Trp Asn	1795	1800	1805
Ala Met Ile Cys Gly Tyr Lys Arg Gly Arg Asp Asp Gly Asn Ser Gly	1810	1815	1820
Asn Ser Ala Arg Ser Asp Glu Asp Leu Lys Lys Cys Gly Ser Val Pro	1825	1830	1840

-continued

Ser Asp Asp Asp Tyr Pro Met Gly Lys Asn Arg Asp Glu Gly Thr Ala
 1845 1850 1855

Tyr Gln Phe Leu Arg Trp Phe Ala Glu Trp Gly Glu Asp Phe Cys Lys
 1860 1865 1870

His Lys Glu Lys Glu Leu Glu Lys Leu Val Gly Ala Cys Asn Asp Tyr
 1875 1880 1885

Thr Cys Gly Asp Asn Glu Asp Lys Arg Lys Lys Cys Thr Asp Ala Cys
 1890 1895 1900

Thr Gln Tyr Lys Lys Phe Ile Ser Glu Trp Lys Pro Gln Tyr Glu Lys
 1905 1910 1915 1920

Gln Ile Lys Lys Tyr Gly Glu Asn Lys Asp Lys Ile Tyr Ser Glu His
 1925 1930 1935

Pro Val Ala Lys Asp Ala Glu Asp Ala Arg Glu Tyr Leu Asp Lys Gln
 1940 1945 1950

Leu Lys Lys Ile Cys Glu Asn Lys Ser Gly Asp Cys Glu Tyr Lys Cys
 1955 1960 1965

Met Lys Asp Val Ser Thr Gln Arg Leu Thr Asp Gly Asn Ser Gln Asn
 1970 1975 1980

Met Pro Ala Ser Leu Asp Asp Glu Pro Lys Glu Val Glu Gly Lys Cys
 1985 1990 1995 2000

Asn Cys Gln Val Pro Arg Gly Pro Pro Arg Val Arg Arg Glu Thr Pro
 2005 2010 2015

Ser Pro Arg Val Ser Leu Ile Ser Lys Ala Thr Ala Ser Lys Lys Glu
 2020 2025 2030

Ala Lys Thr Ala Pro Pro Thr Lys Gln Pro Lys Lys Val Glu Asn Leu
 2035 2040 2045

Thr Thr Glu Met Arg Ala Gln Thr Arg Thr Arg Arg Ala Ala Gln Gln
 2050 2055 2060

Thr Arg Lys Arg Thr Ser Thr Ala Thr Thr Thr Glu Ser Asp Val Gly
 2065 2070 2075 2080

Thr Met Val Lys Ala Ile Leu Ser Asn Lys Pro Asp Ser Arg Gly Gly
 2085 2090 2095

Ile Glu Gly Cys Asn Pro Lys Thr Tyr Gly Gln Tyr Pro Lys Trp Gly
 2100 2105 2110

Cys Ile Val Gly Lys Ser Lys Glu Asn Glu Asn Gly Ile Cys Met Pro
 2115 2120 2125

Pro Arg Arg Lys Lys Leu Cys Ile Asn Asn Ile Gln Tyr Leu Asn Tyr
 2130 2135 2140

Glu Thr Glu Asn Lys Arg Asp Asn Asp Ile Lys Glu Ala Phe Ile Lys
 2145 2150 2155 2160

Cys Ala Ala Ile Glu Thr Gln Phe Leu Trp Leu Lys Tyr Ile Ile Glu
 2165 2170 2175

Asn Pro Ala Ala Glu Asn Glu Leu Gln Asn Gly Thr Ile Pro Asp Glu
 2180 2185 2190

Phe Lys Arg Ile Met Tyr Tyr Thr Tyr Gly Asp Tyr Lys Asp Met Phe
 2195 2200 2205

Phe Gly Thr Asp Ile Ser Asn Asp Lys Lys Ile Ile Thr Val Thr Asn
 2210 2215 2220

Ser Val Thr Thr Ile Leu Asn Glu Asn Asn Lys Lys Lys Gln Asp Lys
 2225 2230 2235 2240

-continued

Lys Lys Asp Glu Glu Leu Arg Lys Ile Phe Trp Glu Lys Asn Lys Lys
 2245 2250 2255

Phe Ile Trp Glu Gly Met Ile Tyr Gly Leu Thr Tyr His Leu Thr Asp
 2260 2265 2270

Glu Asn Glu Lys Glu Lys Ile Arg Asp Asn Tyr Gln Tyr Asn Asp Met
 2275 2280 2285

Thr Lys Leu Thr Pro Ser Leu Glu Glu Phe Val Lys Arg Pro Gln Phe
 2290 2295 2300

Leu Arg Trp Phe Thr Glu Trp Ala Glu Glu Phe Cys Asn Lys Arg Lys
 2305 2310 2315 2320

Glu Gln Leu Leu Lys Leu Glu Ala Gly Cys Lys Glu Tyr Glu Cys Asn
 2325 2330 2335

Gly Ser Asn Asp Gly Lys Thr Gln Glu Cys Ala Glu Ala Cys Val Thr
 2340 2345 2350

Tyr Gln Asn Phe Ile Lys Lys Trp Lys Thr Glu Tyr Glu Arg Gln Arg
 2355 2360 2365

Glu Lys Phe Lys Lys Asp Lys Asp Gly Lys Lys Tyr Lys Asp Tyr Pro
 2370 2375 2380

Ser Thr Glu Arg Asp Ile Glu Lys Ala Thr Cys Ala His Glu Tyr Leu
 2385 2390 2395 2400

Asn Met Lys Leu Lys Glu Leu Cys Gly Asn Lys Asp Cys Ser Cys Met
 2405 2410 2415

Gln Lys Pro Ser Ser Gln Leu Pro Lys Thr Thr Gln Gln Ser Gln Ser
 2420 2425 2430

Ser Asp Ala Asn Asp Met Pro Glu Ser Leu Asp Tyr Val Pro Glu Glu
 2435 2440 2445

Phe Asn Lys Cys Glu Cys Pro Glu Leu Ser Lys Lys Gly Ser Met Ile
 2450 2455 2460

His Thr Lys Lys Ile Thr Glu Pro Lys Ile Pro Met Asn Cys Val Glu
 2465 2470 2475 2480

Lys Ala Ala Tyr Tyr Leu Ser Lys Glu Ala Glu Asn Asn Met Asp Ile
 2485 2490 2495

Thr Leu Lys Glu Lys Phe Ile Pro Ile Glu Ser Thr Lys Glu Lys Glu
 2500 2505 2510

Ser Lys Asn Ser Trp Thr Asn Asn Asn Pro Cys Asp Pro Lys Lys Pro
 2515 2520 2525

Tyr Ala Pro Asp Lys Tyr Ile Gly Arg Arg Asn Pro Cys Glu Asn Arg
 2530 2535 2540

Glu Glu Asn Arg Phe Lys Val Asp Tyr Glu Trp Lys Cys Tyr Lys Asn
 2545 2550 2555 2560

Ser Lys Phe Tyr Gln Glu Lys Lys Arg Val Cys Val Pro Pro Arg Arg
 2565 2570 2575

Glu His Met Cys Leu Arg Asn Leu Asp Glu Ile Lys Ile Glu Arg Leu
 2580 2585 2590

Lys Asp Ser Asn Tyr Leu Leu Lys Met Val Arg Arg Thr Ala Arg Asn
 2595 2600 2605

Glu Gly Ile Asp Ile Ile Lys Asn Phe Asn Ser Glu Asn Gly Cys Ala
 2610 2615 2620

Met Asn Pro Ile Cys Asp Thr Met Lys Tyr Ser Phe Ala Asp Leu Gly
 2625 2630 2635 2640

Asp Ile Val Arg Gly Thr Asp Met Leu Arg Ile Gly Gly Tyr Leu Pro

-continued

Asn Tyr Lys Ser Ala Asp Ile Pro Met Asn Thr Glu Pro Asn Thr Leu
 3460 3465 3470

Tyr Ser Asp Asn Pro Glu Glu Lys Pro Phe Ile Ile Ser Ile His Asp
 3475 3480 3485

Arg Asp Leu Tyr Thr Gly Lys Glu Ile Ser Tyr Asn Ile Asn Met Ser
 3490 3495 3500

Thr Asn Thr Asn Asn Asp Ile Pro Met Asn Ala Arg Asn Asp Ser Tyr
 3505 3510 3515 3520

Arg Gly Ile Asp Leu Ile Asn Asp Ser Leu Val Val Leu Asn Leu Leu
 3525 3530 3535

Ile Tyr Met Met Lys Tyr
 3540

<210> SEQ ID NO 3
 <211> LENGTH: 32
 <212> TYPE: PRT
 <213> ORGANISM: Plasmodium falciparum

<400> SEQUENCE: 3

Glu Ala Glu Lys Glu Leu Lys Glu Gly Lys Ile Pro Glu Gly Phe Lys
 1 5 10 15

Arg Gln Met Phe Tyr Thr Phe Gly Asp Tyr Arg Asp Ile Leu Phe Gly
 20 25 30

<210> SEQ ID NO 4
 <211> LENGTH: 10
 <212> TYPE: PRT
 <213> ORGANISM: Plasmodium falciparum

<400> SEQUENCE: 4

Lys Glu Leu Lys Glu Gly Lys Ile Pro Glu
 1 5 10

<210> SEQ ID NO 5
 <211> LENGTH: 4
 <212> TYPE: PRT
 <213> ORGANISM: Plasmodium falciparum

<400> SEQUENCE: 5

Lys Glu Gly Lys
 1

<210> SEQ ID NO 6
 <211> LENGTH: 6
 <212> TYPE: PRT
 <213> ORGANISM: Plasmodium falciparum
 <220> FEATURE:
 <221> NAME/KEY: VARIANT
 <222> LOCATION: (1)...(6)
 <223> OTHER INFORMATION: misc_difference
 <223> OTHER INFORMATION: Xaa = Any Amino Acid

<400> SEQUENCE: 6

Lys Xaa Asn Gly Xaa Asn
 1 5

<210> SEQ ID NO 7
 <211> LENGTH: 294
 <212> TYPE: PRT
 <213> ORGANISM: Plasmodium falciparum

-continued

<400> SEQUENCE: 7

Val Leu Gln Gln Lys Ser Asn Gly Ser Ile Asp Asn Cys Asn Ala Lys
 1 5 10 15
 Asn Arg Lys Lys Asn Glu Trp Gln Cys Asp Lys Asn Thr Phe Val Asp
 20 25 30
 Gly Asn Glu Gly Val Cys Met Pro Pro Arg Arg Lys Ser Ile Cys Ile
 35 40 45
 His Asn Leu Thr Leu Glu Glu Gln Thr Lys Asn Lys Tyr Gln Leu Arg
 50 55 60
 Glu Ala Phe Ile Lys Cys Ala Ala Lys Glu Thr Asn Leu Leu Trp Asp
 65 70 75 80
 Lys Tyr Lys Asn Asp Lys Asn Glu Ala Glu Glu Leu Leu Lys Lys Gly
 85 90 95
 Lys Ile Pro Glu Asp Phe Met Arg Ile Met Phe Tyr Thr Phe Gly Asp
 100 105 110
 Phe Arg Asp Phe Cys Leu Glu Asn Asp Met Gly Lys Asp Val Asp Lys
 115 120 125
 Val Lys Lys Asn Ile Asn Lys Val Phe Asn Asn Ser Ser Lys Arg Gly
 130 135 140
 Phe Lys Lys Ile Asp Pro Glu Asn Trp Trp Asn Glu Asn Gly Pro Gln
 145 150 155 160
 Ile Trp Asn Gly Met Leu Cys Ala Leu Ile His Ala Asp Thr Lys Asp
 165 170 175
 Ser Ile Lys Asn Lys Asp Asn Tyr Lys Tyr Glu Lys Val Thr Ile Leu
 180 185 190
 Ala Lys Arg Asp Gly Ser Asn Gly Met Thr Leu Ser Glu Phe Ala Lys
 195 200 205
 Lys Pro Lys Phe Leu Arg Trp Phe Val Glu Trp Tyr Asp Asp Tyr Cys
 210 215 220
 Lys Glu Arg Gln Lys Tyr Leu Thr Glu Val Ala Ser Thr Cys Lys Ser
 225 230 235 240
 Ile Asp Gly Gly Gln Leu Lys Cys Asp Arg Gly Cys Asn Asn Lys Cys
 245 250 255
 Asp Glu Tyr Lys Lys Tyr Met Arg Lys Lys Lys Glu Glu Trp Asn Leu
 260 265 270
 Gln Asp Lys Tyr Tyr Lys Asp Lys Arg Glu Asn Lys Gly Ile Asp Lys
 275 280 285
 Gly Pro Ile Gly Ile Ile
 290

<210> SEQ ID NO 8

<211> LENGTH: 407

<212> TYPE: PRT

<213> ORGANISM: Plasmodium falciparum

<400> SEQUENCE: 8

Asn Gly Gln Asp Gly Thr Lys Lys Ile Glu Glu Cys Asn Thr Lys Tyr
 1 5 10 15
 Tyr Pro Thr Lys Asn Asp Tyr Pro Gly Trp Asn Cys Thr Asp Lys Val
 20 25 30
 Ile Asn Arg Glu Glu Gly Ser Cys Met Pro Pro Arg Arg Gln Lys Leu
 35 40 45

-continued

Cys Ile His Asn Leu Glu His Leu Ser Glu Lys Ala Thr Glu Thr Glu
 50 55 60
 Leu Arg Lys Ala Phe Ile Glu Cys Ala Ala Ile Glu Thr Phe Trp Leu
 65 70 75 80
 Trp Asp Lys Tyr Lys Glu Asp Lys Lys Asp Glu Lys Lys Thr Glu Gly
 85 90 95
 Gly Gly Ile Ser Asp Asp Pro Asp Asp Pro Gln Lys Lys Leu Glu Gly
 100 105 110
 Gly Thr Ile Pro Glu Asp Phe Lys Arg Gln Met Phe Tyr Thr Tyr Gly
 115 120 125
 Asp Tyr Arg Asp Phe Leu Phe Gly Thr Asp Ile Ser Lys Gly His Gly
 130 135 140
 Lys Glu Ser Ala Leu Gly Lys Lys Ile Asp Ser Leu Phe Lys Asn Gly
 145 150 155 160
 Asp Gln Lys Ser Pro Ser Gly Lys Thr Pro Thr Glu Trp Trp Asn Asp
 165 170 175
 Tyr Gly Pro Asp Ile Trp Lys Gly Met Val Cys Gly Leu Ser His His
 180 185 190
 Ile Lys Asn Gly Asn Lys Glu Gln Leu Arg Lys Asn Leu Thr Asp Asn
 195 200 205
 Asn Lys Tyr Thr Lys Ile Ser Ser Lys Leu Glu Asp Phe Ala Ser Arg
 210 215 220
 Pro Gln Phe Leu Arg Trp Phe Ile Glu Trp Gly Asp Gln Phe Cys Arg
 225 230 235 240
 Glu Arg Val Val Lys Ile Asn Gln Leu Lys Thr Gly Cys Asn Glu Tyr
 245 250 255
 Glu Cys Gly Ser Gln Glu Asn Gly Lys Lys Glu Ala Cys Lys Asn Ala
 260 265 270
 Cys Glu Ala Tyr Lys Ser Trp Leu Lys Asp Trp Lys Asp Gln Tyr Glu
 275 280 285
 Gln Gln Thr Ala Lys Phe Asp Lys Asp Lys Lys Asp Lys Lys Phe Asp
 290 295 300
 Gly Thr Ser Ala Glu Val Asp Val Ala Ala Val Ser Ser Val His Glu
 305 310 315 320
 Tyr Leu Gln Glu Glu Leu Lys Asn Leu Cys Thr Lys Gly Asp Cys Ala
 325 330 335
 Cys Met Glu Lys Pro Ser Ala Gln Asp Glu Glu Thr Glu Leu Leu Gly
 340 345 350
 Gly Asn Tyr Phe Pro Glu Ala Met Asp Tyr Pro Pro Lys Glu Ile Gly
 355 360 365
 Glu Arg Cys Lys Cys Ala Ile Pro Ser Glu Pro Met Ser Cys Val Glu
 370 375 380
 Gln Ile Ala Lys His Leu Arg Glu Lys Ala Glu Lys Asn Val Lys Ile
 385 390 395 400
 Tyr Glu Ser Ser Leu Lys Gly
 405

<210> SEQ ID NO 9

<211> LENGTH: 351

<212> TYPE: PRT

<213> ORGANISM: Plasmodium falciparum

<400> SEQUENCE: 9

-continued

Pro Cys Lys Met Val Gln Lys Leu Ile Ser Glu Gln Ile Glu Lys Asn
 1 5 10 15
 Asn Ile His Asn Cys Lys Lys Thr Glu Asp Ala Lys Trp Lys Cys Glu
 20 25 30
 Asn Thr Lys Leu Gly Glu Asp Glu Gly Val Cys Met Pro Pro Arg Arg
 35 40 45
 Gln Asn Leu Cys Val His Tyr Leu Thr Lys Leu Asn Asp Asp Ser Lys
 50 55 60
 Glu Glu Asp Leu Arg Glu Ala Phe Ile Lys Ser Ala Ala Ala Glu Thr
 65 70 75 80
 Phe Leu Leu Arg Gln Tyr Tyr Asn Ser Lys Asn Val Glu Asp Asp Lys
 85 90 95
 Ile Leu His Arg Asp Met Ile Pro Pro Glu Phe Phe Arg Ser Met Phe
 100 105 110
 Tyr Thr Phe Gly Asp Tyr Arg Asp Ile Cys Leu Asp Thr Asp Ile Ser
 115 120 125
 Glu Lys Ile Ala Asp His Asp Val Thr Thr Ala Lys Lys Lys Ile Thr
 130 135 140
 Ala Val Phe Gln Lys Ile Gly Ser Lys Thr Thr Asn Gly Lys Lys Val
 145 150 155 160
 Leu Glu Arg Glu Gly Trp Trp Lys Glu Tyr Gly Leu Ser Ile Trp Lys
 165 170 175
 Gly Met Leu Cys Ala Leu Ser Tyr Asn Thr Glu Thr Lys Lys Met Asp
 180 185 190
 Glu Gly Val Arg Thr Tyr Leu Met Lys Tyr Ile Tyr Lys Asn Asn Asp
 195 200 205
 Ile Lys Glu Tyr Leu Glu Glu Phe Ala Ser Arg Pro Pro Phe Leu Arg
 210 215 220
 Trp Val Thr Glu Trp Gly Glu Asp Phe Val Lys Asn Arg Lys Lys Glu
 225 230 235 240
 Leu Val Ser Leu Lys Lys Lys Cys Asp Ser Cys Thr Leu Arg Asn Asn
 245 250 255
 Gly Thr Ser Asn Lys Thr Cys Asp Asp Asn Glu Asn Cys Gly Ala Cys
 260 265 270
 Lys Thr Gln Cys Glu Lys Tyr Lys Lys Trp Met Glu Arg Trp Lys Lys
 275 280 285
 His Tyr Ser Ser Gln Lys Lys Lys Phe Gln Leu Tyr Lys Asn Ser Ala
 290 295 300
 Thr Tyr Asn Asn Gly Leu Ala Val Lys Glu Ala Asn Ser Glu Thr Tyr
 305 310 315 320
 Lys Asn Asp Pro Glu Val Thr Glu Ala Asn Ser Ala Lys His Ala Arg
 325 330 335
 Asp Tyr Leu Lys Thr Gln Leu Glu Asn Met Ile Cys Thr Asn Gly
 340 345 350

<210> SEQ ID NO 10

<211> LENGTH: 311

<212> TYPE: PRT

<213> ORGANISM: Plasmodium falciparum

<400> SEQUENCE: 10

Glu Thr Asp Asp Ile Asp Gly Cys Asn Gln Lys Tyr Lys Ala Gly Lys

-continued

1	5	10	15																
Asp	Lys	Tyr	Pro	Gly	Trp	Asp	Cys	Asn	Ser	Gln	Ile	His	Thr	Thr	His				
			20					25					30						
Asn	Gly	Ala	Cys	Met	Pro	Pro	Arg	Arg	Gln	Lys	Leu	Cys	Val	Ser	Gly				
		35					40					45							
Leu	Thr	Lys	Thr	Asp	Arg	Ile	Lys	Ala	Ile	Glu	Tyr	Ile	Arg	Thr	Glu				
	50					55					60								
Phe	Ile	Lys	Ser	Ala	Ala	Ile	Glu	Thr	His	Phe	Ala	Trp	Asp	Arg	Tyr				
	65				70					75					80				
Lys	Glu	Asp	Asn	Gly	Glu	Ala	Glu	Ala	Glu	Leu	Lys	Asn	Gly	Asn	Ile				
				85					90					95					
Pro	Glu	Gly	Phe	Lys	Arg	Gln	Met	Tyr	Tyr	Thr	Phe	Gly	Asp	Tyr	Arg				
			100					105						110					
Asp	Ile	Phe	Phe	Gly	Arg	Asp	Ile	Ser	Thr	His	Ala	Tyr	Ile	Ser	Gly				
		115					120						125						
Val	Ser	Pro	Lys	Val	Ile	Thr	Ile	Leu	Glu	Lys	Glu	Asn	Asp	Ala	Lys				
		130				135						140							
Tyr	Ala	Ala	Lys	Gln	Asn	Ser	Asn	Asn	Glu	Leu	Leu	Asp	Asp	Trp	Trp				
	145				150						155				160				
Asp	Gln	His	Gly	Lys	Asp	Ile	Trp	Glu	Gly	Met	Leu	Cys	Ala	Leu	Thr				
				165					170						175				
His	Lys	Ile	Ser	Asp	Glu	Glu	Lys	Lys	Lys	Glu	Ile	Lys	Asn	Lys	Tyr				
			180					185						190					
Ser	Tyr	Lys	Lys	Leu	Asn	Glu	Ser	Pro	Lys	Gly	Ser	Asn	Lys	Val	Glu				
		195					200						205						
Asp	Phe	Ala	Lys	Lys	Pro	Gln	Phe	Leu	Arg	Trp	Phe	Ile	Glu	Trp	Gly				
	210					215						220							
Asp	Glu	Phe	Cys	Ala	Gln	Arg	Glu	Glu	Lys	Glu	Ala	Lys	Val	Lys	Val				
	225				230					235					240				
Ser	Cys	Ser	Asp	Ala	Lys	Asp	Tyr	Asp	Gly	Cys	Lys	Asn	Thr	Lys	Ser				
				245					250					255					
Asn	Ala	Ser	Cys	Val	Ser	Ala	Cys	Lys	Val	Tyr	Glu	Asp	Tyr	Ile	Thr				
			260				265							270					
Lys	Lys	Lys	Val	Glu	Tyr	Thr	Lys	Gln	Lys	Gly	Lys	Phe	Asp	Ala	Glu				
		275					280						285						
Lys	Ile	Thr	Asp	Lys	Glu	Gly	Tyr	Glu	Gly	Phe	Ser	Thr	Lys	Asp	Ala				
	290					295						300							
Ser	Glu	Tyr	Leu	Lys	Lys	Lys													
	305				310														

<210> SEQ ID NO 11
 <211> LENGTH: 308
 <212> TYPE: PRT
 <213> ORGANISM: Plasmodium falciparum

<400> SEQUENCE: 11

Gly	Asn	Asp	Gly	Ser	Asn	Glu	Ile	Ser	Gly	Cys	Asn	Pro	Lys	Glu	Ser				
1			5						10					15					
Tyr	Pro	Asp	Trp	Asp	Cys	Lys	Lys	Asn	Ile	Asp	Asn	Ser	His	Ser	Gly				
			20					25						30					

-continued

Ala Cys Met Pro Pro Arg Arg Gln Lys Leu Cys Val Arg Asp Leu Thr
35 40 45

Gln Gly Gly Glu Ile Arg Lys Pro Glu Asp Ile Leu Thr Lys Phe Ile
50 55 60

Asn Cys Ala Ala Lys Glu Thr His Phe Ala Trp His Lys Tyr Lys Lys
65 70 75 80

Asp Asn Val Asn Ala Glu Asn Glu Leu Lys Ser Gly Lys Ile Pro Glu
85 90 95

Gly Phe Arg Lys Gln Met Tyr Tyr Thr Phe Gly Asp Phe Arg Asp Ile
100 105 110

Phe Phe Gly Thr Asp Ile Ser Ser Cys Arg Tyr Ile Lys Asp Thr Ser
115 120 125

Gln Thr Ile Lys Ser Lys Leu Gly Asp Gln Ala Thr Thr Glu Lys Gly
130 135 140

Asp Thr His Ile Asp Asp Asn Lys Lys Leu Gln Glu Trp Trp Thr Ile
145 150 155 160

His Gly Pro Lys Ile Trp Glu Gly Met Leu Cys Ala Leu Thr Asn Gly
165 170 175

Leu Ser Glu Ser Glu Lys Lys Asn Ile Leu Gln Asp Tyr Ser Tyr Asn
180 185 190

Lys Leu Asn Asn Ala Glu Lys Asp Asp Cys Cys Leu Glu Lys Phe Ala
195 200 205

Ser Lys Pro Gln Phe Leu Arg Trp Tyr Val Glu Trp Ser Asp Glu Phe
210 215 220

Cys Arg Glu Arg Lys Lys Leu Glu Asp Lys Val Glu Asp Val Cys Ile
225 230 235 240

Lys Ala Lys Asp Tyr Glu Gly Cys Lys Asn Asn Lys Ser Asn Asn Ser
245 250 255

Cys Val Lys Val Cys Lys Glu Tyr Glu Asn Tyr Ile Thr Gly Lys Lys
260 265 270

Thr Gln Tyr Glu Ser Gln Glu Gly Lys Phe Asn Thr Glu Lys Arg Gln
275 280 285

Lys Lys Pro Glu Tyr Asn Ser Tyr Ser Lys Lys Asp Ala Ser Glu Tyr
290 295 300

Leu Lys Asp Lys
305

What is claimed is:

1. A method of detecting the concentration or expression level of FCR3.varCSA or FCR3.varCSA in a biological sample from a tested subject comprising the step of comparing the concentration or expression level of a first sequence selected from the group consisting of FCR3.varCSA gene, FCR3.varCSA RNA, FCR3.varCSA cDNA, and FCR3.varCSA polypeptide from the biological sample with the concentration or expression level of a second sequence selected from the group consisting of FCR3.varCSA gene, FCR3.varCSA RNA, FCR3.varCSA cDNA, and FCR3.varCSA polypeptide from a healthy subject or FCR3.varCSA gene, FCR3.varCSA RNA, FCR3.varCSA cDNA, and FCR3.varCSA polypeptide from a subject afflicted with malaria.

2. A method of making a FCR3.varCSA disease-state profile comprising:

providing a biological sample; and

detecting in the biological sample a concentration or expression level of FCR3.varCSA or FCR3.varCSA.

3. A purified or isolated nucleic acid comprising the sequence of SEQ ID NO: 1 or a sequence complementary thereto.

4. A purified or isolated nucleic acid comprising at least 9 consecutive bases of the sequence of SEQ ID NO.: 1 or a sequence complementary thereto.

5. The purified or isolated nucleic acid of claim 4, wherein said nucleic acid comprises a DBL3 domain or CIDR1 domain or fragment thereof.

6. A purified or isolated nucleic acid comprising at least 9 consecutive bases of the sequence of SEQ ID NO.: 1,

wherein the nucleic acid has a nucleotide sequence found in a var gene that encodes a protein that binds to chondroitin sulfate A that is at least 80% homologous to FCR3.varCSA or a sequence complementary to SEQ. ID. NO.: 1 or said var gene.

7. A purified or isolated nucleic acid encoding a polypeptide having the sequence of SEQ ID NO.: 2.

8. A recombinant construct comprising the coding region of SEQ ID NO.: 1 operably linked to a heterologous promoter.

9. A vector comprising the isolated DNA of claim 3.

10. A vector comprising the isolated DNA of claim 5.

11. An isolated nucleic acid molecule that hybridizes to SEQ. ID. NO. 1 at 37° C. in the presence of 0.5M NaPO₄ (pH 7) and 7% SDS and under wash conditions of 37° C., in 6×SSC and 0.2% SDS, wherein the nucleic acid molecule has a sequence complementary to a sequence found in a var gene that encodes a protein that binds to chondroitin sulfate A that is at least 80% homologous to FCR3.varCSA.

12. A purified or isolated protein comprising the sequence of SEQ ID NO.: 2.

13. The purified or isolated protein of claim 12, wherein at least one acidic amino acid contained therein is replaced with a different acidic amino acid.

14. The purified or isolated protein of claim 12, wherein at least one basic amino acid contained therein is replaced with a different basic amino acid.

15. The purified or isolated protein of claim 12, wherein at least one nonpolar amino acid contained therein is replaced with a different nonpolar amino acid.

16. The purified or isolated protein of claim 12, wherein at least one uncharged amino acid contained therein is replaced with a different uncharged amino acid.

17. The purified or isolated protein of claim 12, wherein at least one aromatic amino acid contained therein is replaced with a different aromatic amino acid.

18. A purified or isolated polypeptide comprising at least 3 consecutive amino acids of the sequence of SEQ ID NO.: 2, wherein the polypeptide binds to chondroitin sulfate A (CSA).

19. A purified or isolated polypeptide comprising at least 3 consecutive amino acids of the sequence of SEQ ID NO.: 2, wherein the amino acid sequence of the polypeptide is found in molecule that binds to chondroitin sulfate A (CSA) that is at least 80% homologous to FCR3.varCSA.

20. A purified or isolated polypeptide, wherein the polypeptide is at least 80% identical to the polypeptide having the amino acid sequence of SEQ. ID. NO. 1 as determined by FASTA or BLAST using default opening and gap penalties and a PAM 250 scoring matrix.

21. A purified or isolated polypeptide comprising a DBL3 or CIDR1 domain or fragment thereof consisting of at least 9 amino acids of SEQ. ID. No.:2.

22. A method of making a protein having the sequence of SEQ ID NO.: 2 comprising:

obtaining a cDNA comprising the sequence in SEQ ID NO.: 1;

inserting said cDNA in an expression vector such that said cDNA is operably linked to a promoter; and

introducing said expression vector into a host cell whereby said host cell produces the protein encoded by said cDNA.

23. The method of claim 22, further comprising isolating the protein.

24. An isolated FCR3.varCSA polypeptide that promotes adhesion to CSA wherein the polypeptide is selected from the group consisting of:

(a) a polypeptide having the amino acid sequence of SEQ. ID NO. 2

(b) a polypeptide encoded by the nucleic acid of claim 3, 4, 5, 6, or 7; and

(c) a polypeptide that is at least 70% identical to the polypeptide of (a) or (b) as determined by FASTA or BLAST using default opening and gap penalties and a PAM 250 scoring matrix.

25. A method for constructing a transformed host cell that expresses SEQ ID NO.: 3 comprising transforming the host cell with a recombinant DNA vector comprising the sequence of SEQ ID NO.: 1.

26. A cultured cell line comprising the vector of claim 9.

27. A cultured cell line comprising the vector of claim 10.

28. A purified or isolated antibody capable of specifically binding FCR3.varCSA, wherein the antibody recognizes an epitope found on a var protein that binds to chondroitin sulfate A that is unique to a class of var proteins having a structure that is at least 80% homologous to FCR3.varCSA.

29. The antibody of claim 28, wherein the antibody is a monoclonal antibody.

30. A purified or isolated antibody that binds DBL3 or CIDR1 of SEQ. ID. No. 2.

31. The antibody of claim 30, wherein the antibody is a monoclonal antibody.

32. An isolated or purified biological complex comprising FCR3.varCSA and a ligand for FCR3.varCSA.

33. An isolated or purified biological complex comprising a molecule selected from the group consisting of FCR3.varCSA, a fragment of FCR3.varCSA, A4 tres DBL3- γ , and ItG2-CS2 DBL2- γ joined to chondroitin sulfate A (CSA) or an analog thereof.

34. A method of identifying FCR3.varCSA dependent adhesion to chondroitin sulfate A (CSA) comprising:

providing a support having chondroitin sulfate A (CSA) or a representative fragment thereof;

contacting the support with FCR3.varCSA or a representative fragment thereof; and

detecting FCR3.varCSA dependent adhesion.

35. The method of claim 34, wherein the support is selected from the group consisting of a resin, a plastic reservoir, a lipid bilayer, and a cell.

36. The method of claim 34, wherein the FCR3.varCSA is joined to a second support selected from the group consisting of a resin, a plastic reservoir, a lipid bilayer, and a cell.

37. A method of identifying an agent that modulates FCR3.varCSA dependent adhesion to chondroitin sulfate A (CSA) comprising:

providing a support having chondroitin sulfate A (CSA) or a representative fragment thereof;

contacting the support with FCR3.varCSA or a representative fragment thereof;

contacting the support with the agent; and

detecting FCR3.varCSA dependent adhesion.

38. The method of claim 37, wherein the support is selected from the group consisting of a resin, a plastic reservoir, a lipid bilayer, and a cell.

39. The method of claim 37, wherein the FCR3.varCSA is joined to a second support selected from the group consisting of a resin, a plastic reservoir, a lipid bilayer, and a cell.

40. A method of identifying an agent that modulates FCR3.varCSA dependent adhesion to chondroitin sulfate A (CSA) comprising:

providing a support having FCR3.varCSA or a representative fragment thereof that has a polypeptide sequence found in a var protein that binds to chondroitin sulfate A and is at least 80% homologous to FCR3.varCSA.;

contacting the support with the agent; and

detecting FCR3.varCSA dependent adhesion to chondroitin sulfate A (CSA).

41. The method of claim 40, wherein the FCR3.varCSA is joined to a second support selected from the group consisting of a resin, a plastic reservoir, a lipid bilayer, and a cell.

42. The method of claim 40, wherein the support is selected from the group consisting of a resin, a plastic reservoir, a lipid bilayer, and a cell.

43. The method of claim 40, wherein the CSA is joined to a second support selected from the group consisting of a resin, a plastic reservoir, a lipid bilayer, and a cell.

44. A method of identifying an agent that modulates FCR3.varCSA dependent adhesion to chondroitin sulfate A (CSA) comprising:

providing a cell transfected with a construct comprising a nucleic acid sequence encoding FCR3.varCSA or a representative fragment thereof that has a nucleotide sequence found in a var gene that encodes a protein that binds to chondroitin sulfate A that is at least 80% homologous to FCR3.varCSA.;

contacting the cell with the agent; and

detecting FCR3.varCSA dependent adhesion to chondroitin sulfate A (CSA).

45. A method of identifying an agent that interacts with the sequence set forth in SEQ ID NO.: 2 or a representative fragment thereof that has an amino acid sequence not found in another var protein comprising:

transfecting a cell with a nucleic acid encoding the sequence set forth in SEQ ID NO.: 1 or 2 or a representative fragment thereof that has a nucleotide sequence encoding FCR3.varCSA or a representative fragment thereof that has a nucleotide sequence found in a var gene that encodes a protein that binds to chondroitin sulfate A that is at least 80% homologous to FCR3.varCSA.;

contacting the cell with the agent; and

detecting an interaction of the agent and a polypeptide encoded by the sequence set forth in SEQ ID NO.: 2 or a representative fragment thereof that has a nucleotide sequence found in a var gene that encodes a protein that binds to chondroitin sulfate A that is at least 80% homologous to FCR3.varCSA.

46. A method of preparing a therapeutic agent comprising: providing an agent that modulates FCR3.varCSA-dependent adhesion to CSA; and

mixing with the agent a pharmaceutically acceptable carrier.

47. A pharmaceutical comprising an antibody of claims 28, 29, 30, or 31 or an agent identified according to any one of claims 37-46 and a pharmaceutically acceptable carrier.

48. A method of treatment and prevention of maternal malaria comprising:

identifying a patient at risk for contracting maternal malaria or a patient afflicted with maternal malaria; and

administering a therapeutically effective amount of a nucleic acid complementary to at least 15 consecutive nucleotides of SEQ. ID. NO. 1, wherein the nucleic acid is complementary to a sequence encoding FCR3.varCSA or a representative fragment thereof that has a nucleotide sequence found in a var gene that encodes a protein that binds to chondroitin sulfate A that is at least 80% homologous to FCR3.varCSA.

49. The method of claim 48 further comprising administering a pharmaceutically acceptable carrier or adjuvant.

50. A method of treatment and prevention of maternal malaria comprising:

identifying a patient at risk for contracting maternal malaria or a patient afflicted with maternal malaria; and

administering a therapeutically effective amount of a polypeptide having at least 3 consecutive amino acids of SEQ. ID. NO. 2, wherein the amino acid sequence is found in a protein that binds chondroitin sulfate A that is at least 80% homologous to FCR3.varCSA.

51. The method of claim 50, further comprising administering a pharmaceutically acceptable carrier or adjuvant.

52. A method of treatment and prevention of maternal malaria comprising:

identifying a patient at risk for contracting maternal malaria or a patient afflicted with maternal malaria; and

administering a therapeutically effective amount of a peptide agent that corresponds to at least 3 consecutive amino acids of SEQ. ID. NO. 2, wherein the amino acid sequence is found in a protein that binds chondroitin sulfate A that is at least 80% homologous to FCR3.varCSA.

53. The method of claim 52, further comprising administering a pharmaceutically acceptable carrier or adjuvant.

54. A method of treatment and prevention of maternal malaria comprising:

identifying a patient at risk for contracting maternal malaria or a patient afflicted with maternal malaria; and

administering a therapeutically effective amount of a peptide agent that corresponds to at least 3 consecutive amino acids of SEQ. ID. NO. 9 or 11, wherein the amino acid sequence is found in a protein that binds chondroitin sulfate A.

55. The method of claim 54, further comprising administering a pharmaceutically acceptable carrier or adjuvant.

* * * * *

专利名称(译)	鉴定恶性疟原虫红细胞膜蛋白1 (PfEMP1) 的结构域介导与硫酸软骨素A的粘附		
公开(公告)号	US20040062769A1	公开(公告)日	2004-04-01
申请号	US10/087013	申请日	2002-02-21
[标]申请(专利权)人(译)	舍夫ARTHUR MILLER LOUIS ^ h GAMAIN BENOIT BARUCH DROR我 BUFFET PIERRE SCHEIDIG CHRISTINE GYSIN于尔格 POUVELLE BRUNO 藤井信孝 SMITH JOSEPH		
申请(专利权)人(译)	舍夫ARTHUR MILLER LOUIS H. GAMAIN BENOIT BARUCH DROR I. BUFFET PIERRE SCHEIDIG CHRISTINE GYSIN于尔格 POUVELLE BRUNO 藤井信孝 SMITH JOSEPH		
当前申请(专利权)人(译)	UNIVERSITE DE LA MEDITERRANEE 政府美利坚合众国，成为代表局局长，卫生和人类服务，THE系 巴斯德研究所		
[标]发明人	SCHERF ARTHUR MILLER LOUIS H GAMAIN BENOIT BARUCH DROR I BUFFET PIERRE SCHEIDIG CHRISTINE GYSIN JURG POUVELLE BRUNO FUJII NOBUTAKA SMITH JOSEPH		
发明人	SCHERF, ARTHUR MILLER, LOUIS H. GAMAIN, BENOIT BARUCH, DROR I. BUFFET, PIERRE SCHEIDIG, CHRISTINE GYSIN, JURG POUVELLE, BRUNO FUJII, NOBUTAKA SMITH, JOSEPH		
IPC分类号	A61K38/00 A61K39/00 A61K48/00 C07K14/445 C12N15/30 C12Q1/68 C12Q1/6883 G01N33/569 A61K39/395 G01N33/53		

CPC分类号	A61K38/00 A61K39/00 A61K48/00 A61K2039/505 G01N2333/445 C07K14/445 C12Q1/6883 C12Q2600/158 G01N33/56905 A61K2039/53 Y02A50/412
优先权	60/152023 1999-09-01 US
其他公开文献	US6855323
外部链接	Espacenet USPTO

摘要(译)

本发明涉及调节寄生的红细胞与硫酸软骨素的粘附的var基因和相应的蛋白质的发现。还公开了前述的新生物工具，预防剂，治疗剂，诊断剂和使用方法。



FIG. 1A




FIG. 1B