



US 20050142557A1

(19) **United States**

(12) **Patent Application Publication** (10) **Pub. No.: US 2005/0142557 A1**
Alleman et al. (43) **Pub. Date: Jun. 30, 2005**

(54) **POLYNUCLEOTIDES AND POLYPEPTIDES OF ANAPLASMA PHAGOCYTOPHILUM AND METHODS OF USING THE SAME**

(52) **U.S. Cl.** **435/6;** 435/69.1; 435/320.1; 435/252.3; 530/350; 536/23.7; 435/7.32; 530/388.4

(75) Inventors: **Arthur Rick Alleman**, Alachua, FL (US); **Anthony F. Barbet**, Archer, FL (US)

(57) **ABSTRACT**

Correspondence Address:
SALIWANCIK LLOYD & SALIWANCHIK
A PROFESSIONAL ASSOCIATION
PO BOX 142950
GAINESVILLE, FL 32614-2950 (US)

We have successfully sequenced and cloned the gene expressing the Major Surface Protein 5 (MSP5) of *A. phagocytophilum*. The recombinant MSP5 (rMSP5) protein has been tested using sera from humans and dogs infected with *A. phagocytophilum*. The polypeptide has been found to be immunogenic and useful as a diagnostic test antigen. The polypeptide antigen of the subject invention can provide the basis of a diagnostic assay that would allow the rapid, in-house, laboratory diagnosis of infection with *A. phagocytophilum* using a sample (e.g., serum, plasma, or whole blood) from an infected human or animal. Additionally, the subject invention provides methods of detecting the presence of *A. phagocytophilum* in biological or environmental samples utilizing antibodies provided by the subject invention. Furthermore, the use of the single antigen in the diagnosis of this important disease offers many advantages including enhanced test specificity, ease of testing and consistency of results using synthetically produced test antigens instead of cultured, whole organisms.

(73) Assignee: **University of Florida**, Gainesville, FL

(21) Appl. No.: **10/696,019**

(22) Filed: **Oct. 28, 2003**

Publication Classification

(51) **Int. Cl.⁷** **C12Q 1/68;** C07H 21/04; C07K 14/195; G01N 33/554; G01N 33/569




FIG 1




FIG 2

**POLYNUCLEOTIDES AND POLYPEPTIDES OF
ANAPLASMA PHAGOCYTOPHILUM AND
METHODS OF USING THE SAME**

[0001] *Anaplasma phagocytophilum* is a rickettsial agent that infects multiple species of mammals including humans, dogs, horses, cows and small ruminants. The current method of serological diagnosis relies on using whole, cultured organisms in an immunofluorescent antibody assay that is labor intensive, subjective in evaluation and nonspecific in reactivity. There remains a great need for an immunoassay system that simplifies the diagnosis and/or identification of *Anaplasma phagocytophilum* infections in individuals.

[0002] The currently existing methods for serological diagnosis of infection with *A. phagocytophilum* is the immunofluorescent antibody assay. This assay requires the use of whole, cultivated organisms to serve as diagnostic antigens. The production of these organisms is costly and labor intensive and there is no way to control antigen preparations of different lots to provide consistent immunoassay results.

[0003] The immunofluorescent assay itself is labor intensive and requires specialized equipment that is only available in commercial or research laboratories. Additionally, the assay requires highly skilled personnel to accurately interpret results. The rMSP5 could be synthetically produced as a peptide product, which would result in consistency in the test antigens. The synthetic peptide is adaptable to a format that would allow technicians in virtually any clinical setting to test subjects for antibodies to the organism. A positive result would be recognized by a visible color change on the test pad that could be interpreted by anyone with minimal training and skills.

BRIEF SUMMARY OF THE INVENTION

[0004] We have successfully sequenced and cloned the gene expressing the Major Surface Protein 5 (MSP5) of *A. phagocytophilum*. The recombinant MSP5 (rMSP5) protein has been tested using sera from humans and dogs infected with *A. phagocytophilum*. The polypeptide has been found to be immunogenic and useful as a diagnostic test antigen. The polypeptide antigen of the subject invention can provide the basis of a diagnostic assay that would allow the rapid, in-house, laboratory diagnosis of infection with *A. phagocytophilum* using a sample (e.g., serum, plasma, or whole blood) from an infected human or animal. Additionally, the subject invention provides methods of detecting the presence of *A. phagocytophilum* in biological or environmental samples utilizing antibodies provided by the subject invention. Furthermore, the use of the single antigen in the diagnosis of this important disease offers many advantages including enhanced test specificity, ease of testing and consistency of results using synthetically produced test antigens instead of cultured, whole organisms.

BRIEF DESCRIPTION OF THE DRAWINGS

[0005] **FIG. 1**—Coomassie Blue-stained gel of purified recombinant MSP5 (rMSP5) from *A. phagocytophilum*. Lane A contains molecular weight standards; Lanes B and C contain fractions of soluble rMSP5 in the supernatant of the eluted material; Lane D contains rMSP5 from a pelleted insoluble fraction.

[0006] **FIG. 2**—Western immunoblot using rMSP5 and peroxidase labeled anti-histidine antibody to identify the

recombinant protein with the histidine tag. Lanes B, and C contain rMSP5 from soluble fractions and Lane D contains rMSP5 from the insoluble pellet. Horseradish peroxidase labeled anti-His antibodies were as the expressed rMSP5 contained a polyhistidine tag.

BRIEF DESCRIPTION OF THE TABLES

[0007] Tables 1-3 provides exemplary polypeptide fragments of the polypeptides of the subject invention (e.g., the polypeptides of SEQ ID NO: 2, 3, and 4).

[0008] Tables 4-5 illustrate the results of enzyme linked immunosorbent (ELISA) assays using recombinant rMSP5 to detect antibodies in the serum of infected and non-infected humans (Table 4) and dogs (Table 5). Samples from non-infected humans were used to establish cutoff values for positive reactors. An optical density (OD) reading based upon the mean from the normal samples (at each dilution) plus three standard deviations (SD) was used as the upper limit of normal. Any samples with OD readings greater than the mean +3SD (of the normal samples at the same dilutions) were considered positive reactors. Human samples were tested at dilutions of 1/100, 1/300, and 1/1000. Canine samples were tested at dilutions of 1/100 and 1/300.

[0009] In Table 4, serum samples from non-infected humans (Normals 1-5) were compared to samples obtained from humans that were PCR positive for *A. phagocytophilum*. As illustrated in the table, all samples were serologically positive except for the sample obtained from 99-01291 (this sample was PCR positive but serologically negative when tested by a different assay).

[0010] In Table 5, ELISA tests using serum samples from non-infected dogs (Molly, CN76AC, CN115A, CN125C, CN35B, Dixie, CN54B, CN105C, Presley, and CN74H) and samples obtained from a commercial source (IDEXX Laboratories, Inc., Westbrook, Me.). Samples from IDEXX were blinded as 10 of the samples were positive for antibodies to *A. phagocytophilum* (samples 4, 6, 11, 12, 13, 15, 16, 17, 18, and 20). Samples with OD readings greater than the mean +3SD (of the normal samples at the same dilutions) were considered positive reactors.

BRIEF DESCRIPTION OF THE SEQUENCES

[0011] SEQ ID NO: 1 provides a polynucleotide sequence encoding a MSP5 polypeptide of the subject invention.

[0012] SEQ ID NO: 2 depicts a processed form (lacking a signal peptide) of a MSP5 polypeptide of the subject invention.

[0013] SEQ ID NO: 3 illustrates an unprocessed form of a MSP5 polypeptide of the subject invention (e.g., containing a signal peptide).

[0014] SEQ ID NO: 4 is a signal peptide of the subject invention.

**DETAILED DESCRIPTION OF THE
INVENTION**

[0015] The subject invention provides:

[0016] a) one or more:

[0017] 1) isolated, purified, and/or recombinant polypeptides comprising SEQ ID NO: 2, 3, or 4;

- [0018] 2) variant polypeptides having at least about 20% to 99.99% identity, preferably at least 60 to 99.99% identity to the polypeptide of SEQ ID NO: 2, 3, or 4 and which has at least one of the activities associated with the polypeptide of SEQ ID NO: 2, 3, or 4;
- [0019] 3) a fragment of the polypeptide of SEQ ID NO: 2, 3, or 4, or a variant polypeptide, wherein said polypeptide fragment or fragment of said variant polypeptide has substantially the same activity as the polypeptide of SEQ ID NO: 2, 3, or 4;
- [0020] 4) a multimeric polypeptide construct comprising a series of repeating elements that are, optionally, joined together by linker elements, wherein said repeating elements are selected from one, or more, of the following polypeptides: a) SEQ ID NO: 2; b) SEQ ID NO: 3; c) SEQ ID NO: 4; d) fragments of SEQ ID NO: 2; e) fragments of SEQ ID NO: 3; or f) fragments of SEQ ID NO: 4;
- [0021] 5) an epitope of a polypeptide selected from the group consisting of SEQ ID NO: 2, SEQ ID NO: 3, and SEQ ID NO: 4;
- [0022] 6) a multi-epitope construct comprising at least one epitope as set forth herein; or
- [0023] 7) a polypeptide according to embodiments a(1), a(2), a(3), a(4), a(5), or a(6) that further comprises a heterologous polypeptide sequence;
- [0024] b) a composition comprising a carrier and a polypeptide as set forth in a(1), a(2), a(3), a(4), a(5), a(6), or a(7), wherein said carrier is an adjuvant or a pharmaceutically acceptable excipient;
- [0025] c) methods of detecting the presence of antibodies in an individual infected with *A. phagocytophilum* comprising contacting a biological sample with a polypeptide or polypeptides as set forth in a(1), a(2), a(3), a(4), a(5), a(6), or a(7) and detecting the presence of an antigen/antibody complex;
- [0026] d) an improvement in methods of diagnosing or detecting an *A. phagocytophilum* infection in an individual, wherein the improvement comprises the use of an isolated, purified, and/or recombinant polypeptide as set forth in a(1), a(2), a(3), a(4), a(5), a(6), or a(7) in an immunoassay for the detection or diagnosis of an *A. phagocytophilum* infection;
- [0027] In the context of the instant invention, the terms “oligopeptide”, “polypeptide”, “peptide” and “protein” can be used interchangeably; however, it should be understood that the invention does not relate to the polypeptides in natural form, that is to say that they are not in their natural environment but that the polypeptides may have been isolated or obtained by purification from natural sources or obtained from host cells prepared by genetic manipulation (e.g., the polypeptides, or fragments thereof, are recombinantly produced by host cells, or by chemical synthesis). Polypeptides according to the instant invention may also contain non-natural amino acids, as will be described below. The terms “oligopeptide”, “polypeptide”, “peptide” and “protein” are also used, in the instant specification, to designate a series of residues, typically L-amino acids, connected one to the other, typically by peptide bonds between the α -amino and carboxyl groups of adjacent amino acids. Linker elements can be joined to the polypeptides of the subject invention through peptide, bonds or via chemical bonds (e.g., heterobifunctional chemical linker elements) as set forth below. Additionally, the terms “amino acid(s)” and “residue(s)” can be used interchangeably.
- [0028] Thus, the subject invention provides polypeptides comprising SEQ ID NOs: 2, 3 or 4 and/or polypeptide fragments of SEQ ID NOs: 2, 3 or 4. In some embodiments of the subject invention, polypeptide fragments of the subject invention are epitopes that are bound by antibodies or T-cell receptors are designated “epitopes”; in the context of the subject invention, “epitopes” are considered to be a subset of the invention designated as “fragments of SEQ ID NOs: 2, 3 or 4”. In yet another aspect of the invention, a “fragment” within the context of the subject invention comprises the signal peptide (signal sequence) of SEQ ID NO 4.
- [0029] Polypeptide fragments (and/or epitopes) according to the subject invention, usually comprise a contiguous span of or at least 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, or 73 amino acids of SEQ ID NO:2; in one embodiment, a fragment of SEQ ID No: 2 comprises, consists essentially of, or consists of amino acids 17-74 of SEQ ID NO:2. In other embodiments, the subject invention provides fragments of SEQ ID No: 3 that are, or that are at least, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, or 57 consecutive amino acids.
- [0030] Polypeptide fragments of the subject invention can be any integer in length from at least 3, preferably 4, and more preferably 5 consecutive amino acids to 1 amino acid less than a full length polypeptide of SEQ ID NO: 2, 3 or 4. Thus, for SEQ ID No: 2, a polypeptide fragment can be any integer of consecutive amino acids from 3 to 73 amino acids; for SEQ ID No: 3, a fragment can be any integer of consecutive amino acids from 5 to 57 consecutive amino acids, for example. The term “integer” is used herein in its mathematical sense and thus representative integers include: 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, and/or 74.
- [0031] Each polypeptide fragment of the subject invention can also be described in terms of its N-terminal and C-terminal positions. For example, combinations of N-terminal to C-terminal fragments of 6 contiguous amino acids to 1 amino acid less than the full length polypeptide of SEQ ID No: 2 are included in the present invention. Thus, a 6 consecutive amino acid fragment could occupy positions selected from the group consisting of 1-6, 2-7, 3-8, 4-9, 5-10, 6-11, 7-12, 8-13, 9-14, 10-15, 11-16, 12-17, 13-18, 14-19, 15-20, 16-21, 17-22, 18-23, 19-24, 20-25, 21-26, 22-27, 23-28, 24-29, 25-30, 26-31, 27-32, 28-33, 29-34, 30-35, 31-36, 32-37, 33-38, 34-39, 35-40, 36-41, 37-42, 38-43, 39-44, 40-45, 41-46, 42-47, 43-48, 44-49, 45-50,

46-51, 47-52, 48-53, 49-54, 50-55, 51-56, 52-57, 53-58, 54-59, 55-60, 56-61, 57-62, 58-63, 59-64, 60-65, 61-66, 62-67, 63-68, 64-69, 65-70, 66-71, 67-72, 68-73, and 69-74. A 70 consecutive amino acid fragment could occupy positions selected from the group consisting of 1-70, 2-71, 3-72, 4-73, and 5-74 of SEQ ID NO: 2.

[0032] Exemplary polypeptide fragments of SEQ ID NOs: 2, 3 or 4 are also set forth in Tables 1, 2, and 3 of the subject specification. As is indicated in the tables appended hereto, polypeptide fragments have an N-terminal amino acid residue that corresponds to an amino acid position of the SEQ ID NO: indicated in the table and a C-terminal amino acid residue that, likewise, corresponds to an amino acid position of the SEQ ID NO: indicated in the table. Any polypeptide fragment listed in the appended tables may be included or specifically excluded from the subject invention.

[0033] Fragments, as described herein, can be obtained by cleaving the polypeptides of the invention with a proteolytic enzyme (such as trypsin, chymotrypsin, or collagenase) or with a chemical reagent, such as cyanogen bromide (CNBr). Alternatively, polypeptide fragments can be generated in a highly acidic environment, for example at pH 2.5. Such polypeptide fragments may be equally well prepared by chemical synthesis or using hosts transformed with an expression vector according to the invention. The transformed host cells contain a nucleic acid, allowing the expression of these fragments, under the control of appropriate elements for regulation and/or expression of the polypeptide fragments. Various polypeptide fragments encompassed within the scope of the subject invention are provided in Table 3. The cleavage point indicated in the table is at the carboxy group of the amino acid indicated in the table; the numerical value indicated in Table 3 corresponds to the amino acid at the identical position in SEQ ID NO: 2.

[0034] In certain preferred embodiments, fragments of the polypeptides disclosed herein retain at least one property or activity of the full-length polypeptide from which the fragments are derived. Thus, fragments of the polypeptide of SEQ ID NOs: 2, 3 or 4 have one or more of the following properties or activities: a) the ability to: 1) specifically bind to antibodies specific for SEQ ID NO: 2, 3, 4; and/or 2) specifically bind antibodies found in an animal or human infected with *A. phagocytophilum*; b) the ability to bind to, and activate T-cell receptors (CTL (cytotoxic T-lymphocyte) and/or HTL (helper T-lymphocyte receptors)) in the context of MHC Class I or Class II antigen that are isolated or derived from an animal or human infected with *A. phagocytophilum*; 3) the ability to induce an immune response in an animal or human; 4) the ability to induce a protective immune response in an animal or human against *A. phagocytophilum*; and/or 5) the ability to direct the extracellular secretion of a polypeptide (e.g., a signal peptide such as SEQ ID NO: 4).

[0035] The polypeptides, and fragments thereof, may further comprise linker elements (L) that facilitate the attachment of the fragments to other molecules, amino acids, or polypeptide sequences. The linkers can also be used to attach the polypeptides, or fragments thereof, to solid support matrices for use in affinity purification protocols. Non-limiting examples of "linkers" suitable for the practice of the invention include chemical linkers (such as those sold by Pierce, Rockford, Ill.), or peptides that allow for the con-

nection combinations of polypeptides (see, for example, linkers such as those disclosed in U.S. Pat. Nos. 6,121,424, 5,843,464, 5,750,352, and 5,990,275, hereby incorporated by reference in their entirety).

[0036] In other embodiments, the linker element (L) can amino acid sequences. In other embodiments, the peptide linker has one or more of the following characteristics: a) it allows for the free rotation of the polypeptides that it links (relative to each other); b) it is resistant or susceptible to digestion (cleavage) by proteases; and c) it does not interact with the polypeptides it joins together. In various embodiments, a multimeric construct according to the subject invention includes a peptide linker and the peptide linker is 5 to 60 amino acids in length. More preferably, the peptide linker is 10 to 30, amino acids in length; even more preferably, the peptide linker is 10 to 20 amino acids in length. In some embodiments, the peptide linker is 17 amino acids in length.

[0037] Peptide linkers suitable for use in the subject invention are made up of amino acids selected from the group consisting of Gly, Ser, Asn, Thr and Ala. Preferably, the peptide linker includes a Gly-Ser element. In a preferred embodiment, the peptide linker comprises (Ser-Gly-Gly-Gly-Gly)_y, wherein y is 1, 2, 3, 4, 5, 6, 7, or 8. Other embodiments provide for a peptide linker comprising ((Ser-Gly-Gly-Gly-Gly)_y-Ser-Pro). In certain preferred embodiments, y is a value of 3, 4, or 5. In other preferred embodiment, the peptide linker comprises (Ser-Ser-Ser-Ser-Gly)_y, or ((Ser-Ser-Ser-Ser-Gly)_y-Ser-Pro), wherein y is 1, 2, 3, 4, 5, 6, 7, or 8. In certain preferred embodiments, y is a value of 3, 4, or 5. Where cleavable linker elements are desired, one or more cleavable linker sequences such as Factor Xa or enterokinase (Invitrogen, San Diego Calif.) can be used alone or in combination with the aforementioned linkers

[0038] Multimeric constructs of the subject invention typically comprise a series of repeating elements, optionally interspersed with other elements. As would be appreciated by one skilled in the art, the order in which the repeating elements occur in the multimeric polypeptide is not critical and any arrangement of the repeating elements as set forth herein can be provided by the subject invention. Thus, a "multimeric construct" according to the subject invention can provide a multimeric polypeptide comprising a series of polypeptides, polypeptide fragments, or epitopes that are, optionally, joined together by linker elements (either chemical linker elements or amino acid linker elements).

[0039] Non-limiting examples of multimeric polypeptide constructs according to the subject invention comprise: a) (SEQ ID NO: 2)_{2x}, (SEQ ID NO: 3)_{2x}, or (SEQ ID NO: 4)_{2x}; or b) [L_d-(SEQ ID NO:2)]_{2x}, [L_d-(SEQ ID NO:3)]_{2x}, [L_d-(SEQ ID NO:4)]_{2x}, [(SEQ ID NO: 2)_a-L_d-(SEQ ID NO:3)_b]_x, [(SEQ ID NO: 2)_a-L_d-(SEQ ID NO:4)_b]_x, [(SEQ ID NO: 3)_a-L_d-(SEQ ID NO:4)_b]_x, or [(SEQ ID NO: 2)_a-L_d-(SEQ ID NO:3)_b-L_e-(SEQ ID NO:4)_c]_x, wherein: 1) L is a linker element joined to the polypeptides of SEQ ID NO: 2, 3 or 4, or fragments of SEQ ID NO: 2 and/or 3; 2) x is an integer from 1 to 100; 3) a, b, and c can be the same, or different, and are an integer from 1 to 100; and 4) d and e can be the same, or different and are an integer from 0 to 100, preferably 0 to 10, and more preferably 0, 1, 2, 3, 4, 5, 6, 7, or 8, and even more preferably 3, 4, or 5. Some embodiments

provide for multimeric constructs of SEQ ID NO: 2 or SEQ ID NO: 3 wherein no linker elements are provided (e.g., a multimeric construct represented by the formula (SEQ ID NO: 2)_x or (SEQ ID NO: 3)_x, wherein x is an integer value from 2 to 200). The order and arrangement of SEQ ID NO: 2, 3, 4 and/or fragments thereof can be altered in any fashion and it is not necessary that the sequences alternate, and in the context of this aspect of the invention, "SEQ ID NO: 2", "SEQ ID NO: 3", and "SEQ ID NO: 4" can be interchanged with fragments of the given SEQ ID NO:.

[0040] A "variant polypeptide" (or polypeptide variant) is to be understood to designate polypeptides exhibiting, in relation to the natural polypeptide, certain modifications. These modifications can include a deletion, addition, or substitution of at least one amino acid, a truncation, an extension, a chimeric fusion, a mutation, or polypeptides exhibiting post-translational modifications. Among these homologous variant polypeptides, are those comprising amino acid sequences exhibiting between at least (or at least about) 20.00% to 99.99% (inclusive) identity to the full length, native, or naturally occurring polypeptide are another aspect of the invention. The aforementioned range of percent identity is to be taken as including, and providing written description and support for, any fractional percentage, in intervals of 0.01%, between 20.00% and, up to, including 99.99%. These percentages are purely statistical and differences between two polypeptide sequences can be distributed randomly and over the entire sequence length. Thus, variant polypeptides can have 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, or 99 percent identity with the polypeptide sequences of the instant invention. In a preferred embodiment, a variant or modified polypeptide exhibits at least 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, or 99 percent identity to SEQ ID Nos: 2, 3 or 4. Typically, the percent identity is calculated with reference to the full-length, native, and/or naturally occurring polypeptide (e.g. those polypeptides set forth in SEQ ID NO: 2, SEQ ID NO: 3 or SEQ ID NO: 4). In all instances, variant polypeptides retain at least one of the activities associated with the polypeptide set forth in SEQ ID Nos: 2, 3 or 4. In some embodiments, variant polypeptides retain at least 2, and preferably all of the activities associated with the polypeptide.

[0041] Variant polypeptides can also comprise one or more heterologous polypeptide sequences (e.g., tags that facilitate purification of the polypeptides of the invention (see, for example, U.S. Pat. No. 6,342,362, hereby incorporated by reference in its entirety; Altendorf et al [1999-WWW, 2000]"Structure and Function of the F₀ Complex of the ATP Synthase from *Escherichia Coli*," *J. of Experimental Biology* 203:19-28, The Co. of Biologists, Ltd., G. B.; Baneyx [1999]"Recombinant Protein Expression in *Escherichia coli*," *Biotechnology* 10:411-21, Elsevier Science Ltd.; Eihauer et al. [2001]"The FLAG™ Peptide, a Versatile Fusion Tag for the Purification of Recombinant Proteins," *J. Biochem Biophys Methods* 49:455-65; Jones et al. [1995]"*J. Chromatography* 707:3-22; Jones et al. [1995]"Current Trends in Molecular Recognition and Bioseparation," *J. of Chromatography A*. 707:3-22, Elsevier Science B. V.; Mar-

golin [2000]"Green Fluorescent Protein as a Reporter for Macromolecular Localization in Bacterial Cells," *Methods* 20:62-72, Academic Press; Puig et al. [2001]"The Tandem Affinity Purification (TAP) Method: A General Procedure of Protein Complex Purification," *Methods* 24:218-29, Academic Press; Sassenfeld [1990]"Engineering Proteins for Purification," *TibTech* 8:88-93; Sheibani [1999]"Prokaryotic Gene Fusion Expression Systems and Their Use in Structural and Functional Studies of Proteins," *Prep. Biochem. & Biotechnol.* 29(1):77-90, Marcel Dekker, Inc.; Skerra et al. [1999]"Applications of a Peptide Ligand for Streptavidin: the Strep-tag", *Biomolecular Engineering* 16:79-86, Elsevier Science, B. V.; Smith [1998]"Cookbook for Eukaryotic Protein Expression: Yeast, Insect, and Plant Expression Systems," *The Scientist* 12(22):20; Smyth et al. [2000]"Eukaryotic Expression and Purification of Recombinant Extracellular Matrix Proteins Carrying the Strep II Tag", *Methods in Molecular Biology*, 139:49-57; Unger [1997]"Show Me the Money: Prokaryotic Expression Vectors and Purification Systems," *The Scientist* 11(17):20, each of which is hereby incorporated by reference in their entireties), or commercially available tags from vendors such as such as STRATAGENE (La Jolla, Calif.), NOVAGEN (Madison, Wis.), QIAGEN, Inc., (Valencia, Calif.), or InVitrogen (San Diego, Calif.).

[0042] In other embodiments, polypeptides of the subject invention (e.g., SEQ ID NOS: 2, 3, and/or 4 or fragments thereof) can be fused to heterologous polypeptide sequences that have adjuvant activity (a polypeptide adjuvant). Non-limiting examples of such polypeptides include heat shock proteins (hsp) (see, for example, U.S. Pat. No. 6,524,825, the disclosure of which is hereby incorporated by reference in its entirety). As indicated supra, the signal peptide of SEQ ID NO: 4 can be used to direct the extracellular secretion of any protein to which they are operably linked. Signal peptides find application in simplifying protein purification techniques. In such applications, the extracellular secretion of the desired protein greatly facilitates purification by reducing the number of undesired proteins from which the desired protein must be selected. Thus, a signal peptide of the subject invention can be operably linked to a heterologous polypeptide to direct the secretion of said polypeptide from a transformed host cell.

[0043] Also included within the scope of the subject invention are at least one or more polypeptide fragments of SEQ ID NO: 2, 3, or 4 that are an "epitope". In the context of the subject invention, the term "epitope" is used to designate a series of residues, typically L-amino acids, connected one to the other, typically by peptide bonds between the α -amino and carboxyl groups of adjacent amino acids. The preferred CTL (or CD8⁺ T cell)-inducing peptides of the invention are 13 residues or less in length and usually consist of between about 8 and about 11 residues (e.g., 8, 9, 10 or 11 residues), preferably 9 or 10 residues. The preferred HTL (or CD4⁺ T cell)-inducing peptides are less than about 50 residues in length and usually consist of between about 6 and about 30 residues, more usually between about 12 and 25 (e.g., 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24 or 25), and often between about 15 and 20 residues (e.g., 15, 16, 17, 18, 19 or 20).

[0044] The nomenclature used to describe peptide compounds follows the conventional practice wherein the amino group is presented to the left (the N-terminus) and the

carboxyl group to the right (the C-terminus) of each amino acid residue. The L-form of an amino acid residue is represented by a capital single letter or a capital first letter of a three-letter symbol, and the D-form, for those amino acids having D-forms, is represented by a lower case single letter or a lower case three letter symbol. Glycine has no asymmetric carbon atom and is simply referred to as "Gly" or G. Symbols for the amino acids are as follows: (Single Letter Symbol; Three Letter Symbol Amino Acid) A; Ala; Alanine: C; Cys; Cysteine: D; Asp; Aspartic Acid: E; Glu; Glutamic Acid: F; Phe; Phenylalanine: G; Gly; Glycine: H; His; Histidine: I; Ile; Isoleucine: K; Lys; Lysine: L; Leu; Leucine: M; Met; Methionine: N; Asn; Asparagine: P; Pro; Proline: Q; Gln; Glutamine: R; Arg; Arginine: S; Ser; Serine: T; Thr; Threonine: V; Val; Valine: W; Trp; Tryptophan: Y; Tyr; Tyrosine. Amino acid "chemical characteristics" are defined as: Aromatic (F, W, Y); Aliphatic-hydrophobic (L, I, V, M); Small polar (S, T, C); Large polar (Q, N); Acidic (D, E); Basic (R, H, K); Non-polar: (P, A, G) Proline; Alanine; and Glycine. By way of example, amino acid substitutions can be carried out without resulting in a substantial modification of the associated activity (or activities) of the corresponding modified polypeptides; for example, the replacement of leucine with valine or isoleucine, of aspartic acid with glutamic acid, of glutamine with asparagine, of arginine with lysine, and the like, the reverse substitutions can be performed without substantial modification of the biological activity of the polypeptides.

[0045] In order to extend the life of the polypeptides according to the invention, it may be advantageous to use non-natural amino acids, for example in the D-form, or alternatively amino acid analogs, for example sulfur-containing forms of amino acids in the production of "variant polypeptides". Alternative means for increasing the life of polypeptides can also be used in the practice of the instant invention. For example, polypeptides of the invention, and fragments thereof, can be recombinantly modified to include elements that increase the plasma, or serum half-life of the polypeptides of the invention. These elements include, and are not limited to, antibody constant regions (see for example, U.S. Pat. No. 5,565,335, hereby incorporated by reference in its entirety, including all references cited therein), or other elements such as those disclosed in U.S. Pat. Nos. 6,319,691, 6,277,375, or 5,643,570, each of which is incorporated by reference in its entirety, including all references cited within each respective patent. Alternatively, the polynucleotides and genes of the instant invention can be recombinantly fused to elements, well known to the skilled artisan, that are useful in the preparation of immunogenic constructs for the purposes of vaccine formulation.

[0046] The subject invention also provides biologically active fragments (epitopes) of a polypeptide according to the invention and includes those peptides capable of eliciting an immune response directed against *A. phagocytophilum*, said immune response providing components (B-cells, antibodies, and/or components of the cellular immune response (e.g., helper, cytotoxic, and/or suppressor T-cells)) reactive with the fragment of said polypeptide; the intact, full length, unmodified polypeptide disclosed herein; or both a fragment of a polypeptide and the intact, full length, unmodified polypeptides disclosed herein.

[0047] The subject application also provides a composition comprising at least one isolated, recombinant, or puri-

fied polypeptide as set forth herein and at least one additional component. In various aspects of the invention, the additional component is a solid support (for example, microtiter wells, magnetic beads, non-magnetic beads, agarose beads, glass, cellulose, plastics, polyethylene, polypropylene, polyester, nitrocellulose, nylon, or polysulfone) and/or a pharmaceutically acceptable excipient or adjuvant known to those skilled in the art. In some aspects of the invention, the solid support provides an array of polypeptides of the subject invention or an array of polypeptides comprising combinations of various polypeptides of the subject invention. Compositions of the subject invention can also comprise additional antigens of interest (e.g., *Anaplasma marginale*, *Anaplasma centrale*, *Ehrlichia canis*, *Ehrlichia chaffeensis*, or *Cowdria ruminantium*, and/or *Borrelia burgdorferi*).

[0048] In one embodiment, the subject invention provides methods for eliciting an immune response in an individual comprising the administration of compositions comprising polypeptides according to the subject invention to an individual in amounts sufficient to induce an immune response in the individual. In some embodiments, a "protective" or "therapeutic immune response" is induced in the individual. A "protective immune response" or "therapeutic immune response" refers to a CTL (or CD8⁺ T cell) and/or an HTL (or CD4⁺ T cell), and/or an antibody response to an antigen derived from an infectious agent or a tumor antigen, which in some way prevents or at least partially arrests disease symptoms, side effects or progression. The protective immune response may also include an antibody response that has been facilitated by the stimulation of helper T cells (or CD4⁺ T cells). Additional methods of inducing an immune response in an individual are taught in U.S. Pat. No. 6,419,931, hereby incorporated by reference in its entirety. The term CTL can be used interchangeably with CD8⁺ T-cell(s) and the term HTL can be used interchangeably with CD4⁺ T-cell(s) throughout the subject application.

[0049] The term "individual" includes mammals which include, and are not limited to, apes, chimpanzees, orangutans, humans, monkeys or domesticated animals (pets) such as dogs, cats, guinea pigs, hamsters, Vietnamese pot-bellied pigs, rabbits, ferrets, cows, horses, goats and sheep. In a preferred embodiment, the methods of inducing an immune response contemplated herein are practiced on humans.

[0050] The composition administered to the individual may, optionally, contain an adjuvant and may be delivered in any manner known in the art for the delivery of immunogen to a subject. Compositions may also be formulated in any carriers, including for example, pharmaceutically acceptable carriers such as those described in E. W. Martin's *Remington's Pharmaceutical Science*, Mack Publishing Company, Easton, Pa. In preferred embodiments, compositions may be formulated in incomplete Freund's adjuvant, complete Freund's adjuvant, or alum.

[0051] In other embodiments, the subject invention provides for diagnostic assays based upon Western blot formats or standard immunoassays known to the skilled artisan. For example, antibody-based assays such as enzyme linked immunosorbent assays (ELISAs), radioimmunoassays (RIAs); lateral flow assays, reversible flow chromatographic binding assay (see, for example, U.S. Pat. No. 5,726,010, which is hereby incorporated by reference in its entirety),

immunochromatographic strip assays, automated flow assays, and assays utilizing peptide- or antibody-containing biosensors may be employed for the detection of: 1) the polypeptides, and fragments thereof, provided by the subject invention; or 2) antibodies that bind to the polypeptides or fragments thereof, provided by the subject invention; The assays and methods for conducting the assays are well-known in the art and the methods may test biological samples (e.g., serum, plasma, or blood) qualitatively (presence or absence of polypeptide) or quantitatively (comparison of a sample against a standard curve prepared using a polypeptide of the subject invention) for the presence of: a) one or more polypeptide of the subject invention, or 2) antibodies that bind to polypeptides of the subject invention.

[0052] Thus, the subject invention provides a method of detecting an *A. phagocytophilum* polypeptide, variant, or fragment of said polypeptide or variant, comprising contacting a sample with an antibody that specifically binds to: 1) a polypeptide, or fragment thereof, or 2) a variant, or a fragment thereof, and detecting the presence of an antibody-antigen complex. Alternatively, the subject invention provides a method of detecting antibodies to *A. phagocytophilum* comprising contacting a sample from an individual with: 1) a polypeptide of the subject invention, or fragment thereof, or 2) a variant of the subject invention, or a fragment thereof, and detecting the presence of an antibody-antigen complex. A sample can comprise a blood, serum, or tissue sample from an individual infected by *A. phagocytophilum*. Alternatively, a sample can comprise culture medium in which polypeptides of the subject invention (or fragments thereof) are expressed or transformed host cells (lysed or intact cells) expressing polypeptides (or fragments thereof) that are provided by the subject invention.

[0053] The antibody-based assays can be considered to be of four types: direct binding assays, sandwich assays, competition assays, and displacement assays. In a direct binding assay, either the antibody or antigen is labeled, and there is a means of measuring the number of complexes formed. In a sandwich assay, the formation of a complex of at least three components (e.g., antibody-antigen-antibody) is measured. In a competition assay, labeled antigen and unlabelled antigen compete for binding to the antibody, and either the bound or the free component is measured. In a displacement assay, the labeled antigen is pre-bound to the antibody, and a change in signal is measured as the unlabelled antigen displaces the bound, labeled antigen from the receptor.

[0054] Lateral flow assays can be conducted according to the teachings of U.S. Pat. No. 5,712,170 and the references cited therein. U.S. Pat. No. 5,712,170 and the references cited therein are hereby incorporated by reference in their entirety. Displacement assays and flow immunosensors useful for carrying out displacement assays are described in: (1) Kusterbeck et al., "Antibody-Based Biosensor for Continuous Monitoring", in *Biosensor Technology*, R. P. Buck et al., eds., Marcel Dekker, N.Y. pp. 345-350 (1990); Kusterbeck et al., "A Continuous Flow Immunoassay for Rapid and Sensitive Detection of Small Molecules", *Journal of Immunological Methods*, vol. 135, pp. 191-197 (1990); Ligler et al., "Drug Detection Using the Flow Immunosensor", in *Biosensor Design and Application*, J. Findley et al., eds., American Chemical Society Press, pp. 73-80 (1992); and Ogert et al., "Detection of Cocaine Using the Flow Immunosensor", *Analytical Letters*, vol. 25, pp. 1999-2019

(1992), all of which are incorporated herein by reference in their entirety. Displacement assays and flow immunosensors are also described in U.S. Pat. No. 5,183,740, which is also incorporated herein by reference in its entirety. The displacement immunoassay, unlike most of the competitive immunoassays used to detect small molecules, can generate a positive signal with increasing antigen concentration. One aspect of the invention allows for the exclusion of Western blots as a diagnostic assay, particularly where the Western blot is a screen of whole cell lysates of *A. phagocytophilum*, or related organisms, against immune serum of infected individuals. In another aspect of the invention, peptide, or polypeptide, based diagnostic assays utilize *A. phagocytophilum* peptides or polypeptides that have been produced either by chemical peptide synthesis or by recombinant methodologies.

[0055] The subject invention also provides methods of binding an antibody to a polypeptide of the subject invention comprising contacting a sample containing an antibody with a polypeptide under conditions that allow for the formation of an antibody-antigen complex. These methods can further comprise the step of detecting the formation of said antibody-antigen complex. In various aspects of this method, an immunoassay is conducted for the detection of *Anaplasma phagocytophilum*. Non-limiting examples of such immunoassays include enzyme linked immunosorbent assays (ELISAs), radioimmunoassays (RIAs), lateral flow assays, immunochromatographic strip assays, automated flow assays, Western blots, immunoprecipitation assays, reversible flow chromatographic binding assays, agglutination assays, and biosensors. Additional aspects of the invention provide for the use of an array of polypeptides when conducted the aforementioned methods of detection (the array can comprise polypeptides of the same or different sequence as well as polypeptides from one or more other organisms (e.g., *Anaplasma marginale*, *Anaplasma centrale*, *Ehrlichia canis*, *Ehrlichia chaffeensis*, or *Cowdria ruminantium*, and/or *Borrelia burgdorferi*).

[0056] The subject invention also concerns antibodies that bind to polypeptides of the invention. Antibodies that are immunospecific for the polypeptides as set forth herein are specifically contemplated. In various embodiments, antibodies that do not cross-react with other proteins (such as *A. marginale* MSP5) are also specifically contemplated. The antibodies of the subject invention can be prepared using standard materials and methods known in the art (see, for example, *Monoclonal Antibodies: Principles and Practice*, 1983; *Monoclonal Hybridoma Antibodies: Techniques and Applications*, 1982; *Selected Methods in Cellular Immunology*, 1980; *Immunological Methods, Vol. II*, 1981; *Practical Immunology*, and Kohler et al. [1975] *Nature* 256:495). These antibodies can further comprise one or more additional components, such as a solid support, a carrier or pharmaceutically acceptable excipient, or a label. In certain aspects of the invention, the antibodies of the invention does not cross-react with an antigen of *Anaplasma marginale*, *Anaplasma centrale*, *Ehrlichia canis*, *Ehrlichia chaffeensis*, or *Cowdria ruminantium*.

[0057] The term "antibody" is used in the broadest sense and specifically covers monoclonal antibodies (including full length monoclonal antibodies), polyclonal antibodies, multispecific antibodies (e.g., bispecific antibodies), and antibody fragments so long as they exhibit the desired

biological activity, particularly neutralizing activity. AAntibody fragments \equiv comprise a portion of a full length antibody, generally the antigen binding or variable region thereof. Examples of antibody fragments include Fab, Fab', F(ab')₂, and Fv fragments; diabodies; linear antibodies; single-chain antibody molecules; and multi-specific antibodies formed from antibody fragments.

[0058] The term Amonoclonal antibody \equiv as used herein refers to an antibody obtained from a population of substantially homogeneous antibodies, i.e., the individual antibodies comprising the population are identical except for possible naturally occurring mutations that may be present in minor amounts. Monoclonal antibodies are highly specific, being directed against a single antigenic site. Furthermore, in contrast to conventional (polyclonal) antibody preparations that typically include different antibodies directed against different determinants (epitopes), each monoclonal antibody is directed against a single determinant on the antigen. The modifier Amonoclonal \equiv indicates the character of the antibody as being obtained from a substantially homogeneous population of antibodies, and is not to be construed as requiring production of the antibody by any particular method. For example, the monoclonal antibodies to be used in accordance with the present invention may be made by the hybridoma method first described by Kohler et al. [1975] *Nature* 256: 495, or may be made by recombinant DNA methods (see, e.g., U.S. Pat. No. 4,816,567). The Amonoclonal antibodies \equiv may also be isolated from phage antibody libraries using the techniques described in Clackson et al. [1991] *Nature* 352: 624-628 and Marks et al. [1991] *J. Mol. Biol.* 222: 581-597, for example.

[0059] The monoclonal antibodies described herein specifically include "chimeric" antibodies (immunoglobulins) in which a portion of the heavy and/or light chain is identical with or homologous to corresponding sequences in antibodies derived from a particular species or belonging to a particular antibody class or subclass, while the remainder of the chain(s) is identical with or homologous to corresponding sequences in antibodies derived from another species or belonging to another antibody class or subclass, as well as fragments of such antibodies, so long as they exhibit the desired biological activity (U.S. Pat. No. 4,816,567; and Morrison et al. [1984] *Proc. Natl. Acad. Sci. USA* 81: 6851-6855). Also included are humanized antibodies, such as those taught in U.S. Pat. Nos. 6,407,213 or 6,417,337 which are hereby incorporated by reference in their entirety.

[0060] "Single-chain Fv" or "sFv" antibody fragments comprise the V_H and V_L domains of an antibody, wherein these domains are present in a single polypeptide chain. Generally, the Fv polypeptide further comprises a polypeptide linker between the V_H and V_L domains which enables the sFv to form the desired structure for antigen binding. For a review of sFv see, Pluckthun in *The Pharmacology of Monoclonal Antibodies* [1994] Vol. 113:269-315, Rosenberg and Moore eds. Springer-Verlag, New York.

[0061] The term Adiabodies \equiv refers to small antibody fragments with two antigen-binding sites, which fragments comprise a heavy chain variable domain (V_H) connected to a light chain variable domain (V_L) in the same polypeptide chain (V_H-V_L). Diabodies are described more fully in, for example, EP 404,097; WO 93/11161; and Hollinger et al. [1993] *Proc. Natl. Acad. Sci. USA* 90: 6444-6448. The term

Alinear antibodies \equiv refers to the antibodies described in Zapata et al. [1995] *Protein Eng.* 8(10):1057-1062.

[0062] An Aisolated \equiv antibody is one which has been identified and separated and/or recovered from a component of its natural environment. Contaminant components of its natural environment are materials which would interfere with diagnostic or therapeutic uses for the antibody, and may include enzymes, hormones, and other proteinaceous or nonproteinaceous solutes. In preferred embodiments the antibody will be purified (1) to greater than 95% by weight of antibody as determined by the Lowry method, and most preferably more than 99% by weight, (2) to a degree sufficient to obtain at least 15 residues of N-terminal or internal amino acid sequence by use of a spinning cup sequenator, or (3) to homogeneity by SDS-PAGE under reducing or nonreducing conditions using Coomassie blue or, preferably, silver stain. Isolated antibody includes the antibody in situ within recombinant cells since at least one component of the antibody's natural environment will not be present. Ordinarily, however, isolated antibody will be prepared by at least one purification step.

[0063] The terms "comprising", "consisting of" and "consisting essentially of" are defined according to their standard meaning. The terms may be substituted for one another throughout the instant application in order to attach the specific meaning associated with each term. The phrases "isolated" or "biologically pure" refer to material that is substantially or essentially free from components which normally accompany the material as it is found in its native state. Thus, isolated peptides in accordance with the invention preferably do not contain materials normally associated with the peptides in their in situ environment. "Link" or "join" refers to any method known in the art for functionally connecting peptides, including, without limitation, recombinant fusion, covalent bonding, disulfide bonding, ionic bonding, hydrogen bonding, and electrostatic bonding.

[0064] The subject invention also provides isolated, recombinant, and/or purified polynucleotide sequences comprising:

- [0065] a) a polynucleotide sequence encoding a polypeptide as set forth in paragraph 13(a);
- [0066] b) a polynucleotide sequence having at least about 20% to 99.99% identity to a polynucleotide sequence encoding a polypeptide SEQ ID NOs: 2, 3, or 4, wherein said polynucleotide encodes a polypeptide having at least one of the activities of SEQ ID NOs: 2, 3, or 4;
- [0067] c) a polynucleotide sequence comprising SEQ ID NO: 1;
- [0068] d) a polynucleotide sequence having at least about 20% to 99.99% identity to the polynucleotide sequence of SEQ ID NO: 1;
- [0069] e) a polynucleotide that is complementary to the polynucleotides set forth in (a), (b), (c), or (d);
- [0070] f) a genetic construct comprising a polynucleotide sequence as set forth in (a), (b), (c), (d), or (e);
- [0071] g) a vector comprising a polynucleotide or genetic construct as set forth in (a), (b), (c), (d), (e), or (f);

[0072] h) a host cell comprising a vector as set forth in (g);

[0073] m) a polynucleotide that hybridizes under low, intermediate or high stringency with a polynucleotide sequence as set forth in (a), (b), (c), (d), (e), (f), or (g); or

[0074] n) a probe comprising a polynucleotide according to (a), (b), (c), (d), (e), (f), or (g) and, optionally, a label or marker;

[0075] "Nucleotide sequence", "polynucleotide" or "nucleic acid" can be used interchangeably and are understood to mean, according to the present invention, either a double-stranded DNA, a single-stranded DNA or products of transcription of the said DNAs (e.g., RNA molecules). It should also be understood that the present invention does not relate to genomic polynucleotide sequences in their natural environment or natural state. The nucleic acid, polynucleotide, or nucleotide sequences of the invention can be isolated, purified (or partially purified), by separation methods including, but not limited to, ion-exchange chromatography, molecular size exclusion chromatography, or by genetic engineering methods such as amplification, subtractive hybridization, cloning, subcloning or chemical synthesis, or combinations of these genetic engineering methods.

[0076] A homologous polynucleotide or polypeptide sequence, for the purposes of the present invention, encompasses a sequence having a percentage identity with the polynucleotide or polypeptide sequences, set forth herein, of between at least (or at least about) 20.00% to 99.99% (inclusive). The aforementioned range of percent identity is to be taken as including, and providing written description and support for, any fractional percentage, in intervals of 0.01%, between 20.00% and, up to, including 99.99%. These percentages are purely statistical and differences between two nucleic acid sequences can be distributed randomly and over the entire sequence length. For example, homologous sequences can exhibit a percent identity of 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, or 99 percent with the sequences of the instant invention. Typically, the percent identity is calculated with reference to the full length, native, and/or naturally occurring polynucleotide. The terms "identical" or percent "identity", in the context of two or more polynucleotide or polypeptide sequences, refer to two or more sequences or subsequences that are the same or have a specified percentage of amino acid residues that are the same, when compared and aligned for maximum correspondence over a comparison window, as measured using a sequence comparison algorithm or by manual alignment and visual inspection.

[0077] Both protein and nucleic acid sequence homologies may be evaluated using any of the variety of sequence comparison algorithms and programs known in the art. Such algorithms and programs include, but are by no means limited to, TBLASTN, BLASTP, FASTA, TFASTA, and CLUSTALW (Pearson and Lipman, 1988, *Proc. Natl. Acad. Sci. USA* 85(8):2444-2448; Altschul et al., 1990, *J. Mol. Biol.* 215(3):403-410; Thompson et al., 1994, *Nucleic Acids Res.* 22(2):4673-4680; Higgins et al., 1996, *Methods Enzy-*

mol. 266:383-402; Altschul et al., 1990, *J. Mol. Biol.* 215(3):403-410; Altschul et al., 1993, *Nature Genetics* 3:266-272). Sequence comparisons are, typically, conducted using default parameters provided by the vendor or using those parameters set forth in the above-identified references, which are hereby incorporated by reference in their entireties.

[0078] A "complementary" polynucleotide sequence, as used herein, generally refers to a sequence arising from the hydrogen bonding between a particular purine and a particular pyrimidine in double-stranded nucleic acid molecules (DNA-DNA, DNA-RNA, or RNA-RNA). The major specific pairings are guanine with cytosine and adenine with thymine or uracil. A "complementary" polynucleotide sequence may also be referred to as an "antisense" polynucleotide sequence or an "antisense sequence".

[0079] Sequence homology and sequence identity can also be determined by hybridization studies under high stringency, intermediate stringency, and/or low stringency. Various degrees of stringency of hybridization can be employed. The more severe the conditions the greater the complementarity that is required for duplex formation. Severity of conditions can be controlled by temperature, probe concentration, probe length, ionic strength, time, and the like. Preferably, hybridization is conducted under low, intermediate, or high stringency conditions by techniques well known in the art, as described, for example, in Keller, G. H., M. M. Manak [1987] *DNA Probes*, Stockton Press, New York, N.Y., pp. 169-170.

[0080] For example, hybridization of immobilized DNA on Southern blots with ³²P-labeled gene-specific probes can be performed by standard methods (Maniatis et al. [1982] *Molecular Cloning. A Laboratory Manual*, Cold Spring Harbor Laboratory, New York). In general, hybridization and subsequent washes can be carried out under intermediate to high stringency conditions that allow for detection of target sequences with homology to the exemplified polynucleotide sequence. For double-stranded DNA gene probes, hybridization can be carried out overnight at 20-25° C. below the melting temperature (T_m) of the DNA hybrid in 6×SSPE, 5× Denhardt's solution, 0.1% SDS, 0.1 mg/ml denatured DNA. The melting temperature is described by the following formula (Beltz et al. [1983] *Methods of Enzymology*, R. Wu, L. Grossman and K. Moldave [eds.] Academic Press, New York 100:266-285).

[0081] T_m=81.5° C.+16.6 Log[Na⁺]+0.41(% G+C)-0.61(% formamide)-600/length of duplex in base pairs.

[0082] Washes are typically carried out as follows:

[0083] (1) twice at room temperature for 15 minutes in 1×SSPE, 0.1% SDS (low stringency wash);

[0084] (2) once at T_m-20° C. for 15 minutes in 0.2×SSPE, 0.1% SDS (intermediate stringency wash).

[0085] For oligonucleotide probes, hybridization can be carried out overnight at 10-20° C. below the melting temperature (T_m) of the hybrid in 6×SSPE, 5× Denhardt's solution, 0.1% SDS, 0.1 mg/ml denatured DNA. T_m for oligonucleotide probes can be determined by the following formula:

[0086] $T_m(^{\circ}\text{C.})=2(\text{number T/A base pairs})+4(\text{number G/C base pairs})$ (Suggs et al. [1981] *ICN-UCLA. Symp. Dev. Biol. Using Purified Genes*, D. D. Brown [ed.], Academic Press, New York, 23:683-693).

[0087] Washes can be carried out as follows:

[0088] (1) twice at room temperature for 15 minutes 1×SSPE, 0.1% SDS (low stringency wash);

[0089] 2) once at the hybridization temperature for 15 minutes in 1×SSPE, 0.1% SDS (intermediate stringency wash).

[0090] In general, salt, and/or temperature can be altered to change stringency. With a labeled DNA fragment >70 or so bases in length, the following conditions can be used:

Low:	1 or 2× SSPE, room temperature
Low:	1 or 2× SSPE, 42° C.
Intermediate:	0.2× or 1× SSPE, 65° C.
High:	0.1× SSPE, 65° C.

[0091] By way of another non-limiting example, procedures using conditions of high stringency can also be performed as follows: Pre-hybridization of filters containing DNA is carried out for 8 h to overnight at 65° C. in buffer composed of 6×SSC, 50 mM Tris-HCl (pH 7.5), 1 mM EDTA, 0.02% PVP, 0.02% Ficoll, 0.02% BSA, and 500 μg/ml denatured salmon sperm DNA. Filters are hybridized for 48 h at 65° C., the preferred hybridization temperature, in pre-hybridization mixture containing 100 μg/ml denatured salmon sperm DNA and 5-20×10⁶ cpm of ³²P-labeled probe. Alternatively, the hybridization step can be performed at 65° C. in the presence of SSC buffer, 1×SSC corresponding to 0.15M NaCl and 0.05 M Na citrate. Subsequently, filter washes can be done at 37° C. for 1 h in a solution containing 2×SSC, 0.01% PVP, 0.01% Ficoll, and 0.01% BSA, followed by a wash in 0.1×SSC at 50° C. for 45 min. Alternatively, filter washes can be performed in a solution containing 2×SSC and 0.1% SDS, or 0.5×SSC and 0.1% SDS, or 0.1×SSC and 0.1% SDS at 68° C. for 15 minute intervals. Following the wash steps, the hybridized probes are detectable by autoradiography. Other conditions of high stringency which may be used are well known in the art and as cited in Sambrook et al., 1989, *Molecular Cloning, A Laboratory Manual*, Second Edition, Cold Spring Harbor Press, N.Y., pp. 9.47-9.57; and Ausubel et al., 1989, *Current Protocols in Molecular Biology*, Green Publishing Associates and Wiley Interscience, N.Y. are incorporated herein in their entirety.

[0092] Another non-limiting example of procedures using conditions of intermediate stringency are as follows: Filters containing DNA are pre-hybridized, and then hybridized at a temperature of 60° C. in the presence of a 5×SSC buffer and labeled probe. Subsequently, filters washes are performed in a solution containing 2×SSC at 50° C. and the hybridized probes are detectable by autoradiography. Other conditions of intermediate stringency which may be used are well known in the art and as cited in Sambrook et al., 1989, *Molecular Cloning, A Laboratory Manual*, Second Edition, Cold Spring Harbor Press, N.Y., pp. 9.47-9.57; and Ausubel et al., 1989, *Current Protocols in Molecular Biology*, Green

Publishing Associates and Wiley Interscience, N.Y. are incorporated herein in their entirety.

[0093] Duplex formation and stability depend on substantial complementarity between the two strands of a hybrid and, as noted above, a certain degree of mismatch can be tolerated. Therefore, the probe sequences of the subject invention include mutations (both single and multiple), deletions, insertions of the described sequences, and combinations thereof, wherein said mutations, insertions and deletions permit formation of stable hybrids with the target polynucleotide of interest. Mutations, insertions and deletions can be produced in a given polynucleotide sequence in many ways, and these methods are known to an ordinarily skilled artisan. Other methods may become known in the future.

[0094] It is also well known in the art that restriction enzymes can be used to obtain functional fragments of the subject DNA sequences. For example, Bal31 exonuclease can be conveniently used for time-controlled limited digestion of DNA (commonly referred to as “erase-a-base” procedures). See, for example, Maniatis et al. [1982] *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory, New York; Wei et al. [1983] *J. Biol. Chem.* 258:13006-13512.

[0095] The present invention further comprises fragments of the polynucleotide sequences of the instant invention. Representative fragments of the polynucleotide sequences according to the invention will be understood to mean any nucleotide fragment having at least 5 successive nucleotides, preferably at least 12 successive nucleotides, and still more preferably at least 15, 18, or at least 20 successive nucleotides of the sequence from which it is derived. The upper limit for such fragments, is the total number of nucleotides found in the full-length sequence encoding a particular polypeptide (e.g., a polypeptide such as that of SEQ ID NO: 2). The term “successive” can be interchanged with the term “consecutive” or the phrase “contiguous span”. Thus, in some embodiments, a polynucleotide fragment may be referred to as “a contiguous span of at least X nucleotides, wherein X is any integer value beginning with 5; the upper limit for such fragments is one nucleotide less than the total number of nucleotides found in the full-length sequence encoding a particular polypeptide (e.g., a polypeptide comprising SEQ ID NO: 2).

[0096] In some embodiments, the subject invention includes those fragments capable of hybridizing under various conditions of stringency conditions (e.g., high or intermediate or low stringency) with a nucleotide sequence according to the invention; fragments that hybridize with a nucleotide sequence of the subject invention can be, optionally, labeled as set forth below.

[0097] The subject invention provides, in one embodiment, methods for the identification of the presence of nucleic acids according to the subject invention in transformed host cells or in cells isolated from an individual suspected of being infected by *A. phagocytophilum*. In these varied embodiments, the invention provides for the detection of nucleic acids in a sample (obtained from the individual or from a cell culture) comprising contacting a sample with a nucleic acid (polynucleotide) of the subject invention (such as an RNA, mRNA, DNA, cDNA, or other nucleic acid). In a preferred embodiment, the polynucleotide

is a probe that is, optionally, labeled and used in the detection system. Many methods for detection of nucleic acids exist and any suitable method for detection is encompassed by the instant invention. Typical assay formats utilizing nucleic acid hybridization includes, and are not limited to, 1) nuclear run-on assay, 2) slot blot assay, 3) northern blot assay (Alwine, et al., *Proc. Natl. Acad. Sci.* 74:5350), 4) magnetic particle separation, 5) nucleic acid or DNA chips, 6) reverse Northern blot assay, 7) dot blot assay, 8) in situ hybridization, 9) RNase protection assay (Melton, et al., *Nuc. Acids Res.* 12:7035 and as described in the 1998 catalog of Ambion, Inc., Austin, Tex.), 10) ligase chain reaction, 11) polymerase chain reaction (PCR), 12) reverse transcriptase (RT)-PCR (Berchtold, et al., *Nuc. Acids Res.* 17:453), 13) differential display RT-PCR (DDRT-PCR) or other suitable combinations of techniques and assays. Labels suitable for use in these detection methodologies include, and are not limited to 1) radioactive labels, 2) enzyme labels, 3) chemiluminescent labels, 4) fluorescent labels, 5) magnetic labels, or other suitable labels, including those set forth below. These methodologies and labels are well known in the art and widely available to the skilled artisan. Likewise, methods of incorporating labels into the nucleic acids are also well known to the skilled artisan.

[0098] Thus, the subject invention also provides detection probes (e.g., fragments of the disclosed polynucleotide sequences) for hybridization with a target sequence or the amplicon generated from the target sequence. Such a detection, probe will comprise a contiguous/consecutive span of at least 8, 9, 10, 11, 12, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, or 100 nucleotides. Labeled probes or primers are labeled with a radioactive compound or with another type of label as set forth above (e.g., 1) radioactive labels, 2) enzyme labels, 3) chemiluminescent labels, 4) fluorescent labels, or 5) magnetic labels). Alternatively, non-labeled nucleotide sequences may be used directly as probes or primers; however, the sequences are generally labeled with a radioactive element (^{32}P , ^{35}S , ^3H , ^{125}I) or with a molecule such as biotin, acetylaminofluorene, digoxigenin, 5-bromo-deoxyuridine, or fluorescein to provide probes that can be used in numerous applications.

[0099] Polynucleotides of the subject invention can also be used for the qualitative and quantitative analysis of gene expression using arrays or polynucleotides that are attached to a solid support. As used herein, the term array means a one-, two-, or multi-dimensional arrangement of full length polynucleotides or polynucleotides of sufficient length to permit specific detection of gene expression. Preferably, the fragments are at least 15 nucleotides in length. More preferably, the fragments are at least 100 nucleotides in length. More preferably, the fragments are more than 100 nucleotides in length. In some embodiments the fragments may be more than 500 nucleotides in length.

[0100] For example, quantitative analysis of gene expression may be performed with full-length polynucleotides of the subject invention, or fragments thereof, in a complementary DNA microarray as described by Schena et al. (*Science* 270:467-470, 1995; *Proc. Natl. Acad. Sci. U.S.A.* 93:10614-10619, 1996). Polynucleotides, or fragments thereof, are amplified by PCR and arrayed onto silylated microscope slides. Printed arrays are incubated in a humid chamber to allow rehydration of the array elements and

rinsed, once in 0.2% SDS for 1 min, twice in water for 1 min and once for 5 min in sodium borohydride solution. The arrays are submerged in water for 2 min at 95° C., transferred into 0.2% SDS for 1 min, rinsed twice with water, air dried and stored in the dark at 25° C.

[0101] mRNA is isolated from a biological sample and probes are prepared by a single round of reverse transcription. Probes are hybridized to 1 cm² microarrays under a 14×14 mm glass coverslip for 6-12 hours at 60° C. Arrays are washed for 5 min at 25° C. in low stringency wash buffer (1×SSC/0.2% SDS), then for 10 min at room temperature in high stringency wash buffer (0.1×SSC/0.2% SDS). Arrays are scanned in 0.1×SSC using a fluorescence laser scanning device fitted with a custom filter set. Accurate differential expression measurements are obtained by taking the average of the ratios of two independent hybridizations.

[0102] Quantitative analysis of the polynucleotides present in a biological sample can also be performed in complementary DNA arrays as described by Pietu et al. (*Genome Research* 6:492-503, 1996). The polynucleotides of the invention, or fragments thereof, are PCR amplified and spotted on membranes. Then, mRNAs originating from biological samples derived from various tissues or cells are labeled with radioactive nucleotides. After hybridization and washing in controlled conditions, the hybridized mRNAs are detected by phospho-imaging or autoradiography. Duplicate experiments are performed and a quantitative analysis of differentially expressed mRNAs is then performed.

[0103] Alternatively, the polynucleotide sequences of the invention may also be used in analytical systems, such as DNA chips. DNA chips and their uses are well known in the art and (see for example, U.S. Pat. Nos. 5,561,071; 5,753,439; 6,214,545; Schena et al., *BioEssays*, 1996, 18:427-431; Bianchi et al., *Clin. Diagn. Virol.*, 1997, 8:199-208; each of which is hereby incorporated by reference in their entirety) and/or are provided by commercial vendors such as Affymetrix, Inc. (Santa Clara, Calif.). In addition, the nucleic acid sequences of the subject invention can be used as molecular weight markers in nucleic acid analysis procedures.

[0104] The subject invention also provides for modified nucleotide sequences. Modified nucleic acid sequences will be understood to mean any nucleotide sequence that has been modified, according to techniques well known to persons skilled in the art, and exhibiting modifications in relation to the native, naturally occurring nucleotide sequences.

[0105] The subject invention also provides genetic constructs comprising: a) a polynucleotide sequence encoding a polypeptide comprising SEQ ID No: 2, 3, 4, or a fragment thereof; b) a polynucleotide sequence having at least about 20% to 99.99% identity to a polynucleotide sequence encoding a polypeptide comprising SEQ ID No: 2, 3, 4, or a fragment of SEQ ID NOs: 2, 3, or 4, wherein said polynucleotide encodes a polypeptide having at least one of the activities or a polypeptide comprising SEQ ID No: 2, 3, 4, or a fragment of SEQ ID NOs: 2, 3, or 4; c) a polynucleotide sequence encoding a fragment of a polypeptide comprising SEQ ID No: 2, 3, or 4, wherein said fragment has at least one of the activities of the polypeptide of SEQ ID No: 2, 3, or 4; d) a polynucleotide sequence comprising SEQ ID NO: 1; e) a polynucleotide sequence having at least about 20% to

99.99% identity to the polynucleotide sequence of SEQ ID NO: 1) a polynucleotide sequence encoding variant (e.g., a variant polypeptide) of the polypeptide of SEQ ID No: 2, 3, or 4, wherein said variant has at least one of the activities associated with the polypeptide of SEQ ID NO: 2, 3, or 4; f) a polynucleotide sequence encoding a fragment of a variant polypeptide as set forth in (e); g) a polynucleotide sequence encoding multimeric construct; or h) a polynucleotide that is complementary to the polynucleotides set forth in (a), (b), (c), (d), (e), (f), or (g). Genetic constructs of the subject invention can also contain additional regulatory elements such as promoters and enhancers and, optionally, selectable markers. In one aspect of the subject invention, the genetic construct comprises a promoter operably linked to a polynucleotide sequence encoding the signal peptide identified in SEQ ID NO: 5 which is operably linked to a polynucleotide sequence encoding a heterologous polypeptide.

[0106] Also within the scope of the subject instant invention are vectors or expression cassettes containing genetic constructs as set forth herein or polynucleotides encoding the polypeptides, set forth supra, operably linked to regulatory elements. The vectors and expression cassettes may contain additional transcriptional control sequences as well. The vectors and expression cassettes may further comprise selectable markers. The expression cassette may contain at least one additional gene, operably linked to control elements, to be co-transformed into the organism. Alternatively, the additional gene(s) and control element(s) can be provided on multiple expression cassettes. Such expression cassettes are provided with a plurality of restriction sites for insertion of the sequences of the invention to be under the transcriptional regulation of the regulatory regions. The expression cassette(s) may additionally contain selectable marker genes operably linked to control elements.

[0107] The expression cassette will include in the 5'-3' direction of transcription, a transcriptional and translational initiation region, a DNA sequence of the invention, and a transcriptional and translational termination regions. The transcriptional initiation region, the promoter, may be native or analogous, or foreign or heterologous, to the host cell. Additionally, the promoter may be the natural sequence or alternatively a synthetic sequence. By "foreign" is intended that the transcriptional initiation region is not found in the native plant into which the transcriptional initiation region is introduced. As used herein, a chimeric gene comprises a coding sequence operably linked to a transcriptional initiation region that is heterologous to the coding sequence.

[0108] Another aspect of the invention provides vectors for the cloning and/or the expression of a polynucleotide sequence taught herein. Vectors of this invention, including vaccine vectors, can also comprise elements necessary to allow the expression and/or the secretion of the said nucleotide sequences in a given host cell. The vector can contain a promoter, signals for initiation and for termination of translation, as well as appropriate regions for regulation of transcription. In certain embodiments, the vectors can be stably maintained in the host cell and can, optionally, contain signal sequences directing the secretion of translated protein. These different elements are chosen according to the host cell used. Vectors can integrate into the host genome or, optionally, be autonomously-replicating vectors.

[0109] The subject invention also provides for the expression of a polypeptide, peptide, fragment, or variant encoded by a polynucleotide sequence disclosed herein comprising the culture of a host cell transformed with a polynucleotide of the subject invention under conditions that allow for the expression of the polypeptide and, optionally, recovering the expressed polypeptide.

[0110] The disclosed polynucleotide sequences can also be regulated by a second nucleic acid sequence so that the protein or peptide is expressed in a host transformed with the recombinant DNA molecule. For example, expression of a protein or peptide may be controlled by any promoter/enhancer element known in the art. Promoters which may be used to control expression include, but are not limited to, the CMV-IE promoter, the SV40 early promoter region (Bernoist and Chambon, 1981, *Nature* 290:304-310), the promoter contained in the 3' long terminal repeat of Rous sarcoma virus (Yamamoto, et al., 1980, *Cell* 22:787-797), the herpes simplex thymidine kinase promoter (Wagner et al., 1981, *Proc. Natl. Acad. Sci. U.S.A.* 78:1441-1445), the regulatory sequences of the metallothionein gene (Brinster et al., 1982, *Nature* 296:39-42); prokaryotic vectors containing promoters such as the β -lactamase promoter (Villa-Kamaroff, et al., 1978, *Proc. Natl. Acad. Sci. U.S.A.* 75:3727-3731), or the tac promoter (DeBoer, et al., 1983, *Proc. Natl. Acad. Sci. U.S.A.* 80:21-25); see also "Useful proteins from recombinant bacteria" in *Scientific American*, 1980, 242:74-94; plant expression vectors comprising the nopaline synthetase promoter region (Herrera-Estrella et al., 1983, *Nature* 303:209-213) or the cauliflower mosaic virus 35S RNA promoter (Gardner, et al., 1981, *Nucl. Acids Res.* 9:2871), and the promoter of the photosynthetic enzyme ribulose biphosphate carboxylase (Herrera-Estrella et al., 1984, *Nature* 310:115-120); promoter elements from yeast or fungi such as the Gal 4 promoter, the ADC (alcohol dehydrogenase) promoter, PGK (phosphoglycerol kinase) promoter, and/or the alkaline phosphatase promoter.

[0111] The vectors according to the invention are, for example, vectors of plasmid or viral origin. In a specific embodiment, a vector is used that comprises a promoter operably linked to a protein or peptide-encoding nucleic acid sequence contained within the disclosed polynucleotide sequences, one or more origins of replication, and, optionally, one or more selectable markers (e.g., an antibiotic resistance gene). Expression vectors comprise regulatory sequences that control gene expression, including gene expression in a desired host cell. Exemplary vectors for the expression of the polypeptides of the invention include the pET-type plasmid vectors (Promega) or pBAD plasmid vectors (Invitrogen) or those provided in the examples below. Furthermore, the vectors according to the invention are useful for transforming host cells so as to clone or express the polynucleotide sequences of the invention.

[0112] The invention also encompasses the host cells transformed by a vector according to the invention. These cells may be obtained by introducing into host cells a nucleotide sequence inserted into a vector as defined above, and then culturing the said cells under conditions allowing the replication and/or the expression of the polynucleotide sequences of the subject invention.

[0113] The host cell may be chosen from eukaryotic or prokaryotic systems, such as for example bacterial cells,

(Gram negative or Gram positive), yeast cells (for example, *Saccharomyces cerevisiae* or *Pichia pastoris*), animal cells (such as Chinese hamster ovary (CHO) cells), plant cells, and/or insect cells using baculovirus vectors. In some embodiments, the host cells for expression of the polypeptides include, and are not limited to, those taught in U.S. Pat. Nos. 6,319,691, 6,277,375, 5,643,570, or 5,565,335, each of which is incorporated by reference in its entirety, including all references cited within each respective patent.

[0114] Furthermore, a host cell strain may be chosen which modulates the expression of the inserted sequences, or modifies and processes the gene product in the specific fashion desired. Expression from certain promoters can be elevated in the presence of certain inducers; thus, expression of the genetically engineered polypeptide may be controlled. Furthermore, different host cells have characteristic and specific mechanisms for the translational and post-translational processing and modification (e.g., glycosylation, phosphorylation) of proteins. Appropriate cell lines or host systems can be chosen to ensure the desired modification and processing of the foreign protein expressed. For example, expression in a bacterial system can be used to produce an unglycosylated core protein product. Expression in yeast will produce a glycosylated product. Expression in mammalian cells can be used to ensure "native" glycosylation of a heterologous protein. Furthermore, different vector/host expression systems may effect processing reactions to different extents.

[0115] The subject invention also concerns novel compositions that can be employed to elicit an immune response or a protective immune response. In this aspect of the invention, an amount of a composition comprising recombinant DNA or mRNA encoding a polynucleotide of the subject invention sufficient to elicit an immune response or protective immune response is administered to an individual. Signal sequences may be deleted from the nucleic acid encoding an antigen, of interest and the individual may be monitored for the induction of an immune response according to methods known in the art. A "protective immune response" or "therapeutic immune response" refers to a CTL (or CD8⁺ T cell) and/or an HTL (or CD4⁺ T cell) response to an antigen that, in some way, prevents or at least partially arrests disease symptoms, side effects or progression. The immune response may also include an antibody response that has been facilitated by the stimulation of helper T cells.

[0116] In another embodiment, the subject invention further comprises the administration of polynucleotide vaccines in conjunction with a polypeptide antigen, or composition thereof, of the invention. In a preferred embodiment, the antigen is the polypeptide that is encoded by the polynucleotide administered as the polynucleotide vaccine. As a particularly preferred embodiment, the polypeptide antigen is administered as a booster subsequent to the initial administration of the polynucleotide vaccine.

[0117] A further embodiment of the subject invention provides for the induction of an immune response to the novel *A. phagocytophilum* antigens disclosed herein (see, for example, the antigens and peptides set forth in the Tables and Sequence Listing attached hereto) using a "prime-boost" vaccination regimen known to those skilled in the art. In this aspect of the invention, a DNA vaccine or polypeptide antigen of the subject invention is administered to an indi-

vidual in an amount sufficient to "prime" the immune response of the individual. The immune response of the individual is then "boosted" via the administration of: 1) one or a combination of: a peptide, polypeptide, and/or full length polypeptide antigen of the subject invention (optionally in conjunction with an immunostimulatory molecule and/or an adjuvant); or 2) a viral vector that contains nucleic acid encoding one, or more, of the same or, optionally, different, antigens, multi-epitope constructs, and/or peptide antigens set forth herein. In some alternative embodiments of the invention, a gene encoding an immunostimulatory molecule may be incorporated into the viral vector used to "boost the immune response of the individual. Exemplary immunostimulatory molecules include, and are not limited to, IL-1, IL-2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-8, IL-9, IL-10, IL-11, IL-15, IL-16, IL-18, IL-23, IL-24, erythropoietin, G-CSF, M-CSF, platelet derived growth factor (PDGF), MSF, FLT-3 ligand, EGF, fibroblast growth factor (FGF; e.g., aFGF (FGF-1), bFGF (FGF-2), FGF-3, FGF-4, FGF-5, FGF-6, or FGF-7), insulin-like growth factors (e.g., IGF-1, IGF-2); vascular endothelial growth factor (VEGF); interferons (e.g., IFN- γ , IFN- α , IFN- β); leukemia inhibitory factor (LIF); ciliary neurotrophic factor (CNTF); oncostatin M; stem cell factor (SCF); transforming growth factors (e.g., TGF- α , TGF- β 1, TGF- β 2, TGF- β 3), or chemokines (such as, but not limited to, BCA-1/BLC-1, BRAK/Kcc, CXCL16, CXCR3, ENA-78/LIX, Eotaxin-1, Eotaxin-2/MIPF-2, Exodus-2/SLC, Fractalkine/Neurotactin, GRO α /MGSA, HCC-1, I-TAC, Lymphotactin/ATAC/SCM, MCP-1/MCAF, MCP-3, MCP-4, MDC/STCP-1, ABCD-1, MIP-1 α , MIP-1 β , MIP-2 α /GRO β , MIP-3 α /Exodus/LARC, MIP-3 β /Exodus-3/ELC, MIP-4/PARC/DC-CK1, PF-4, RANTES, SDF1 α , TARC, or TECK). Genes encoding these immunostimulatory molecules are known to those skilled in the art and coding sequences may be obtained from a variety of sources, including various patents databases, publicly available databases (such as the nucleic acid and protein databases found at the National Library of Medicine or the European Molecular Biology Laboratory), the scientific literature, or scientific literature cited in catalogs produced by companies such as Genzyme, Inc., R&D Systems, Inc, or InvivoGen, Inc. [see, for example, the 1995 Cytokine Research Products catalog, Genzyme Diagnostics, Genzyme Corporation, Cambridge Mass.; 2002 or 1995 Catalog of R&D Systems, Inc (Minneapolis, Minn.); or 2002 Catalog of InvivoGen, Inc (San Diego, Calif.) each of which is incorporated by reference in its entirety, including all references cited therein].

[0118] Methods of introducing DNA vaccines into individuals are well-known to the skilled artisan. For example, DNA can be injected into skeletal muscle or other somatic tissues (e.g., intramuscular injection). Cationic liposomes or biolistic devices, such as a gene gun, can be used to deliver DNA vaccines. Alternatively, iontophoresis and other means for transdermal transmission can be used for the introduction of DNA vaccines into an individual.

[0119] Viral vectors for use in the subject invention can have a portion of the viral genome is deleted to introduce new genes without destroying infectivity of the virus. The viral vector of the present invention is, typically, a non-pathogenic virus. At the option of the practitioner, the viral vector can be selected so as to infect a specific cell type, such as professional antigen presenting cells (e.g., macrophage or dendritic cells). Alternatively, a viral vector can be selected

that is able to infect any cell in the individual. Exemplary viral vectors suitable for use in the present invention include, but are not limited to poxvirus such as vaccinia virus, avipox virus, fowlpox virus, a highly attenuated vaccinia virus (such as Ankara or MVA [Modified Vaccinia Ankara]), retrovirus, adenovirus baculovirus and the like. In a preferred embodiment, the viral vector is Ankara or MVA.

[0120] General strategies for construction of vaccinia virus expression vectors are known in the art (see, for example, Smith and Moss Bio Techniques November/December, 306-312, 1984; U.S. Pat. No. 4,738,846 (hereby incorporated by reference in its entirety). Sutter and Moss (Proc. Nat'l. Acad. Sci U.S.A. 89:10847-10851, 1992) and Sutter et al. (Vaccine, 12(11):1032-40, 1994) disclose the construction and use as a vector, a non-replicating recombinant Ankara virus (MVA) which can be used as a viral vector in the present invention.

[0121] Compositions comprising the subject polynucleotides can include appropriate nucleic acid vaccine vectors (plasmids), which are commercially available (e.g., Vical, San Diego, Calif.) or other nucleic acid vectors (plasmids), which are also commercially available (e.g., Valenti, Burlingame, Calif.). Alternatively, compositions comprising viral vectors and polynucleotides according to the subject invention are provided by the subject invention. In addition, the compositions can include a pharmaceutically acceptable carrier, e.g., saline. The pharmaceutically acceptable carriers are well known in the art and also are commercially available. For example, such acceptable carriers are described in E. W. Martin's *Remington's Pharmaceutical Science*, Mack Publishing Company, Easton, Pa.

[0122] The subject invention also provides an assay that comprises the use of polynucleotides, as set forth herein, for the detection of *Anaplasma phagocytophilum*. In certain preferred embodiments, the polynucleotides used in the assay methods does not hybridize with a polynucleotide sequence from *Anaplasma marginale*, *Anaplasma centrale*, *Ehrlichia canis*, *Ehrlichia chaffeensis*, or *Cowdria ruminantium*. Some aspects of the invention provide for a method that comprises contacting a sample comprising a population of polynucleotides with a second population of polynucleotides under conditions that allow for the formation of an hybridization complex, wherein said second population of polynucleotides comprises polynucleotides that encode at least one polypeptide that is selected from the group consisting of: a) SEQ ID NO 2; b) SEQ ID NO: 3; c) SEQ ID NO: 4; d) fragments of SEQ ID NOs: 2, 3, or 4; e) a polypeptide as set forth in Tables 1 or 2 or 3; f) a variant polypeptide of SEQ ID NO: 2, 3, or 4, wherein said variant polypeptide specifically binds to an antibody that specifically binds to a polypeptide of SEQ ID NO: 2, 3, or 4; g) a variant polypeptide fragment of SEQ ID NO: 2, 3, or 4, wherein said variant polypeptide fragment specifically binds to an antibody that specifically binds to a polypeptide of SEQ ID NO: 2, 3, or 4 or a fragment of SEQ ID NO: 2, 3, or 4; h) a variant of a polypeptide as set forth in Tables 1-3, wherein said variant polypeptide specifically binds to an antibody that specifically binds to a polypeptide of SEQ ID NO: 2, 3, or 4 or a polypeptide as set forth in Table 1, 2 or 3; i) a heterologous polypeptide fused, in frame, to a polypeptide comprising: 1) SEQ ID NO: 2, 3, or 4; 2) fragments of SEQ ID NO: 2, 3, or 4; or 3) a polypeptide as set forth in Tables 1, 2 or 3; and j) mixtures of polypeptides

as set forth in a), b), c), d), e), f), g), h), or i). The method can further comprise the step of detecting the hybridization complex and the second population of polynucleotides can be an array of polynucleotides or the same or different sequence if desired. Some aspects of the invention provide for the use of polynucleotides that do not hybridize with a polynucleotide sequence from *Anaplasma marginale*, *Anaplasma centrale*, *Ehrlichia canis*, *Ehrlichia chaffeensis*, or *Cowdria ruminantium*.

TABLE 1

N-terminal amino acid position	C-terminal amino acid position
5-mer peptides of SEQ ID NO: 2	
1	5
2	6
3	7
4	8
5	9
6	10
7	11
8	12
9	13
10	14
11	15
12	16
13	17
14	18
15	19
16	20
17	21
18	22
19	23
20	24
21	25
22	26
23	27
24	28
25	29
26	30
27	31
28	32
29	33
30	34
31	35
32	36
33	37
34	38
35	39
36	40
37	41
38	42
39	43
40	44
41	45
42	46
43	47
44	48
45	49
46	50
47	51
48	52
49	53
50	54
51	55
52	56
53	57
54	58
55	59
56	60
57	61
58	62
59	63

TABLE 1-continued

N-terminal amino acid position	C-terminal amino acid position
60	64
61	65
62	66
63	67
64	68
65	69
66	70
67	71
68	72
69	73
70	74
6-mer peptides of SEQ ID NO: 2	
1	6
2	7
3	8
4	9
5	10
6	11
7	12
8	13
9	14
10	15
11	16
12	17
13	18
14	19
15	20
16	21
17	22
18	23
19	24
20	25
21	26
22	27
23	28
24	29
25	30
26	31
27	32
28	33
29	34
30	35
31	36
32	37
33	38
34	39
35	40
36	41
37	42
38	43
39	44
40	45
41	46
42	47
43	48
44	49
45	50
46	51
47	52
48	53
49	54
50	55
51	56
52	57
53	58
54	59
55	60
56	61
57	62
58	63
59	64
60	65

TABLE 1-continued

N-terminal amino acid position	C-terminal amino acid position
61	66
62	67
63	68
64	69
65	70
66	71
67	72
68	73
69	74
7-mer peptides of SEQ ID NO: 2	
1	7
2	8
3	9
4	10
5	11
6	12
7	13
8	14
9	15
10	16
11	17
12	18
13	19
14	20
15	21
16	22
17	23
18	24
19	25
20	26
21	27
22	28
23	29
24	30
25	31
26	32
27	33
28	34
29	35
30	36
31	37
32	38
33	39
34	40
35	41
36	42
37	43
38	44
39	45
40	46
41	47
42	48
43	49
44	50
45	51
46	52
47	53
48	54
49	55
50	56
51	57
52	58
53	59
54	60
55	61
56	62
57	63
58	64
59	65
60	66
61	67
62	68

TABLE 1-continued

N-terminal amino acid position	C-terminal amino acid position
63	69
64	70
65	71
66	72
67	73
68	74
8-mer peptides of SEQ ID NO: 2	
1	8
2	9
3	10
4	11
5	12
6	13
7	14
8	15
9	16
10	17
11	18
12	19
13	20
14	21
15	22
16	23
17	24
18	25
19	26
20	27
21	28
22	29
23	30
24	31
25	32
26	33
27	34
28	35
29	36
30	37
31	38
32	39
33	40
34	41
35	42
36	43
37	44
38	45
39	46
40	47
41	48
42	49
43	50
44	51
45	52
46	53
47	54
48	55
49	56
50	57
51	58
52	59
53	60
54	61
55	62
56	63
57	64
58	65
59	66
60	67
61	68
62	69
63	70
64	71
65	72

TABLE 1-continued

N-terminal amino acid position	C-terminal amino acid position
66	73
67	74
9-mer peptides of SEQ ID NO: 2	
1	9
2	10
3	11
4	12
5	13
6	14
7	15
8	16
9	17
10	18
11	19
12	20
13	21
14	22
15	23
16	24
17	25
18	26
19	27
20	28
21	29
22	30
23	31
24	32
25	33
26	34
27	35
28	36
29	37
30	38
31	39
32	40
33	41
34	42
35	43
36	44
37	45
38	46
39	47
40	48
41	49
42	50
43	51
44	52
45	53
46	54
47	55
48	56
49	57
50	58
51	59
52	60
53	61
54	62
55	63
56	64
57	65
58	66
59	67
60	68
61	69
62	70
63	71
64	72
65	73
66	74

TABLE 1-continued

N-terminal amino acid position	C-terminal amino acid position
<u>10-mer peptides of SEQ ID NO: 2</u>	
1	10
2	11
3	12
4	13
5	14
6	15
7	16
8	17
9	18
10	19
11	20
12	21
13	22
14	23
15	24
16	25
17	26
18	27
19	28
20	29
21	30
22	31
23	32
24	33
25	34
26	35
27	36
28	37
29	38
30	39
31	40
32	41
33	42
34	43
35	44
36	45
37	46
38	47
39	48
40	49
41	50
42	51
43	52
44	53
45	54
46	55
47	56
48	57
49	58
50	59
51	60
52	61
53	62
54	63
55	64
56	65
57	66
58	67
59	68
60	69
61	70
62	71
63	72
64	73
65	74
<u>11-mer peptides of SEQ ID NO: 2</u>	
1	11
2	12
3	13
4	14

TABLE 1-continued

N-terminal amino acid position	C-terminal amino acid position
5	15
6	16
7	17
8	18
9	19
10	20
11	21
12	22
13	23
14	24
15	25
16	26
17	27
18	28
19	29
20	30
21	31
22	32
23	33
24	34
25	35
26	36
27	37
28	38
29	39
30	40
31	41
32	42
33	43
34	44
35	45
36	46
37	47
38	48
39	49
40	50
41	51
42	52
43	53
44	54
45	55
46	56
47	57
48	58
49	59
50	60
51	61
52	62
53	63
54	64
55	65
56	66
57	67
58	68
59	69
60	70
61	71
62	72
63	73
64	74
<u>12-mer peptides of SEQ ID NO: 2</u>	
1	12
2	13
3	14
4	15
5	16
6	17
7	18
8	19
9	20
10	21
11	22

TABLE 1-continued

N-terminal amino acid position	C-terminal amino acid position
12	23
13	24
14	25
15	26
16	27
17	28
18	29
19	30
20	31
21	32
22	33
23	34
24	35
25	36
26	37
27	38
28	39
29	40
30	41
31	42
32	43
33	44
34	45
35	46
36	47
37	48
38	49
39	50
40	51
41	52
42	53
43	54
44	55
45	56
46	57
47	58
48	59
49	60
50	61
51	62
52	63
53	64
54	65
55	66
56	67
57	68
58	69
59	70
60	71
61	72
62	73
63	74
<u>13-mer peptides of SEQ ID NO: 2</u>	
1	13
2	14
3	15
4	16
5	17
6	18
7	19
8	20
9	21
10	22
11	23
12	24
13	25
14	26
15	27
16	28
17	29
18	30
19	31

TABLE 1-continued

N-terminal amino acid position	C-terminal amino acid position
20	32
21	33
22	34
23	35
24	36
25	37
26	38
27	39
28	40
29	41
30	42
31	43
32	44
33	45
34	46
35	47
36	48
37	49
38	50
39	51
40	52
41	53
42	54
43	55
44	56
45	57
46	58
47	59
48	60
49	61
50	62
51	63
52	64
53	65
54	66
55	67
56	68
57	69
58	70
59	71
60	72
61	73
62	74
<u>14-mer peptides of SEQ ID NO: 2</u>	
1	14
2	15
3	16
4	17
5	18
6	19
7	20
8	21
9	22
10	23
11	24
12	25
13	26
14	27
15	28
16	29
17	30
18	31
19	32
20	33
21	34
22	35
23	36
24	37
25	38
26	39
27	40
28	41

TABLE 1-continued

N-terminal amino acid position	C-terminal amino acid position
29	42
30	43
31	44
32	45
33	46
34	47
35	48
36	49
37	50
38	51
39	52
40	53
41	54
42	55
43	56
44	57
45	58
46	59
47	60
48	61
49	62
50	63
51	64
52	65
53	66
54	67
55	68
56	69
57	70
58	71
59	72
60	73
61	74
<u>15-mer peptides of SEQ ID NO: 2</u>	
1	15
2	16
3	17
4	18
5	19
6	20
7	21
8	22
9	23
10	24
11	25
12	26
13	27
14	28
15	29
16	30
17	31
18	32
19	33
20	34
21	35
22	36
23	37
24	38
25	39
26	40
27	41
28	42
29	43
30	44
31	45
32	46
33	47
34	48
35	49
36	50
37	51
38	52

TABLE 1-continued

N-terminal amino acid position	C-terminal amino acid position
39	53
40	54
41	55
42	56
43	57
44	58
45	59
46	60
47	61
48	62
49	63
50	64
51	65
52	66
53	67
54	68
55	69
56	70
57	71
58	72
59	73
60	74
<u>16-mer peptides of SEQ ID NO: 2</u>	
1	16
2	17
3	18
4	19
5	20
6	21
7	22
8	23
9	24
10	25
11	26
12	27
13	28
14	29
15	30
16	31
17	32
18	33
19	34
20	35
21	36
22	37
23	38
24	39
25	40
26	41
27	42
28	43
29	44
30	45
31	46
32	47
33	48
34	49
35	50
36	51
37	52
38	53
39	54
40	55
41	56
42	57
43	58
44	59
45	60
46	61
47	62
48	63
49	64

TABLE 1-continued

N-terminal amino acid position	C-terminal amino acid position
50	65
51	66
52	67
53	68
54	69
55	70
56	71
57	72
58	73
59	74
<u>17-mer peptides of SEQ ID NO: 2</u>	
1	17
2	18
3	19
4	20
5	21
6	22
7	23
8	24
9	25
10	26
11	27
12	28
13	29
14	30
15	31
16	32
17	33
18	34
19	35
20	36
21	37
22	38
23	39
24	40
25	41
26	42
27	43
28	44
29	45
30	46
31	47
32	48
33	49
34	50
35	51
36	52
37	53
38	54
39	55
40	56
41	57
42	58
43	59
44	60
45	61
46	62
47	63
48	64
49	65
50	66
51	67
52	68
53	69
54	70
55	71
56	72
57	73
58	74

TABLE 1-continued

N-terminal amino acid position	C-terminal amino acid position
<u>18-mer peptides of SEQ ID NO: 2</u>	
1	18
2	19
3	20
4	21
5	22
6	23
7	24
8	25
9	26
10	27
11	28
12	29
13	30
14	31
15	32
16	33
17	34
18	35
19	36
20	37
21	38
22	39
23	40
24	41
25	42
26	43
27	44
28	45
29	46
30	47
31	48
32	49
33	50
34	51
35	52
36	53
37	54
38	55
39	56
40	57
41	58
42	59
43	60
44	61
45	62
46	63
47	64
48	65
49	66
50	67
51	68
52	69
53	70
54	71
55	72
56	73
57	74
<u>19-mer peptides of SEQ ID NO: 2</u>	
1	19
2	20
3	21
4	22
5	23
6	24
7	25
8	26
9	27
10	28
11	29
12	30

TABLE 1-continued

N-terminal amino acid position	C-terminal amino acid position
13	31
14	32
15	33
16	34
17	35
18	36
19	37
20	38
21	39
22	40
23	41
24	42
25	43
26	44
27	45
28	46
29	47
30	48
31	49
32	50
33	51
34	52
35	53
36	54
37	55
38	56
39	57
40	58
41	59
42	60
43	61
44	62
45	63
46	64
47	65
48	66
49	67
50	68
51	69
52	70
53	71
54	72
55	73
56	74
20-mer peptides of SEQ ID NO: 2	
1	20
2	21
3	22
4	23
5	24
6	25
7	26
8	27
9	28
10	29
11	30
12	31
13	32
14	33
15	34
16	35
17	36
18	37
19	38
20	39
21	40
22	41
23	42
24	43
25	44
26	45
27	46

TABLE 1-continued

N-terminal amino acid position	C-terminal amino acid position
28	47
29	48
30	49
31	50
32	51
33	52
34	53
35	54
36	55
37	56
38	57
39	58
40	59
41	60
42	61
43	62
44	63
45	64
46	65
47	66
48	67
49	68
50	69
51	70
52	71
53	72
54	73
55	74
21-mer peptides of SEQ ID NO: 2	
1	21
2	22
3	23
4	24
5	25
6	26
7	27
8	28
9	29
10	30
11	31
12	32
13	33
14	34
15	35
16	36
17	37
18	38
19	39
20	40
21	41
22	42
23	43
24	44
25	45
26	46
27	47
28	48
29	49
30	50
31	51
32	52
33	53
34	54
35	55
36	56
37	57
38	58
39	59
40	60
41	61
42	62
43	63

TABLE 1-continued

N-terminal amino acid position	C-terminal amino acid position
44	64
45	65
46	66
47	67
48	68
49	69
50	70
51	71
52	72
53	73
54	74
<u>22-mer peptides of SEQ ID NO: 2</u>	
1	22
2	23
3	24
4	25
5	26
6	27
7	28
8	29
9	30
10	31
11	32
12	33
13	34
14	35
15	36
16	37
17	38
18	39
19	40
20	41
21	42
22	43
23	44
24	45
25	46
26	47
27	48
28	49
29	50
30	51
31	52
32	53
33	54
34	55
35	56
36	57
37	58
38	59
39	60
40	61
41	62
42	63
43	64
44	65
45	66
46	67
47	68
48	69
49	70
50	71
51	72
52	73
53	74
<u>23-mer peptides of SEQ ID NO: 2</u>	
1	23
2	24
3	25
4	26
5	27

TABLE 1-continued

N-terminal amino acid position	C-terminal amino acid position
6	28
7	29
8	30
9	31
10	32
11	33
12	34
13	35
14	36
15	37
16	38
17	39
18	40
19	41
20	42
21	43
22	44
23	45
24	46
25	47
26	48
27	49
28	50
29	51
30	52
31	53
32	54
33	55
34	56
35	57
36	58
37	59
38	60
39	61
40	62
41	63
42	64
43	65
44	66
45	67
46	68
47	69
48	70
49	71
50	72
51	73
52	74
<u>24-mer peptides of SEQ ID NO: 2</u>	
1	24
2	25
3	26
4	27
5	28
6	29
7	30
8	31
9	32
10	33
11	34
12	35
13	36
14	37
15	38
16	39
17	40
18	41
19	42
20	43
21	44
22	45
23	46
24	47

TABLE 1-continued

N-terminal amino acid position	C-terminal amino acid position
25	48
26	49
27	50
28	51
29	52
30	53
31	54
32	55
33	56
34	57
35	58
36	59
37	60
38	61
39	62
40	63
41	64
42	65
43	66
44	67
45	68
46	69
47	70
48	71
49	72
50	73
51	74
<u>25-mer peptides of SEQ ID NO: 2</u>	
1	25
2	26
3	27
4	28
5	29
6	30
7	31
8	32
9	33
10	34
11	35
12	36
13	37
14	38
15	39
16	40
17	41
18	42
19	43
20	44
21	45
22	46
23	47
24	48
25	49
26	50
27	51
28	52
29	53
30	54
31	55
32	56
33	57
34	58
35	59
36	60
37	61
38	62
39	63
40	64
41	65
42	66
43	67
44	68

TABLE 1-continued

N-terminal amino acid position	C-terminal amino acid position
45	69
46	70
47	71
48	72
49	73
50	74
<u>26-mer peptides of SEQ ID NO: 2</u>	
1	26
2	27
3	28
4	29
5	30
6	31
7	32
8	33
9	34
10	35
11	36
12	37
13	38
14	39
15	40
16	41
17	42
18	43
19	44
20	45
21	46
22	47
23	48
24	49
25	50
26	51
27	52
28	53
29	54
30	55
31	56
32	57
33	58
34	59
35	60
36	61
37	62
38	63
39	64
40	65
41	66
42	67
43	68
44	69
45	70
46	71
47	72
48	73
49	74
<u>27-mer peptides of SEQ ID NO: 2</u>	
1	27
2	28
3	29
4	30
5	31
6	32
7	33
8	34
9	35
10	36
11	37
12	38
13	39
14	40

TABLE 1-continued

N-terminal amino acid position	C-terminal amino acid position
15	41
16	42
17	43
18	44
19	45
20	46
21	47
22	48
23	49
24	50
25	51
26	52
27	53
28	54
29	55
30	56
31	57
32	58
33	59
34	60
35	61
36	62
37	63
38	64
39	65
40	66
41	67
42	68
43	69
44	70
45	71
46	72
47	73
48	74
<u>28-mer peptides of SEQ ID NO: 2</u>	
1	28
2	29
3	30
4	31
5	32
6	33
7	34
8	35
9	36
10	37
11	38
12	39
13	40
14	41
15	42
16	43
17	44
18	45
19	46
20	47
21	48
22	49
23	50
24	51
25	52
26	53
27	54
28	55
29	56
30	57
31	58
32	59
33	60
34	61
35	62
36	63
37	64

TABLE 1-continued

N-terminal amino acid position	C-terminal amino acid position
38	65
39	66
40	67
41	68
42	69
43	70
44	71
45	72
46	73
47	74
<u>29-mer peptides of SEQ ID NO: 2</u>	
1	29
2	30
3	31
4	32
5	33
6	34
7	35
8	36
9	37
10	38
11	39
12	40
13	41
14	42
15	43
16	44
17	45
18	46
19	47
20	48
21	49
22	50
23	51
24	52
25	53
26	54
27	55
28	56
29	57
30	58
31	59
32	60
33	61
34	62
35	63
36	64
37	65
38	66
39	67
40	68
41	69
42	70
43	71
44	72
45	73
46	74
<u>30-mer peptides of SEQ ID NO: 2</u>	
1	30
2	31
3	32
4	33
5	34
6	35
7	36
8	37
9	38
10	39
11	40
12	41
13	42

TABLE 1-continued

N-terminal amino acid position	C-terminal amino acid position
14	43
15	44
16	45
17	46
18	47
19	48
20	49
21	50
22	51
23	52
24	53
25	54
26	55
27	56
28	57
29	58
30	59
31	60
32	61
33	62
34	63
35	64
36	65
37	66
38	67
39	68
40	69
41	70
42	71
43	72
44	73
45	74
<u>31-mer peptides of SEQ ID NO: 2</u>	
1	31
2	32
3	33
4	34
5	35
6	36
7	37
8	38
9	39
10	40
11	41
12	42
13	43
14	44
15	45
16	46
17	47
18	48
19	49
20	50
21	51
22	52
23	53
24	54
25	55
26	56
27	57
28	58
29	59
30	60
31	61
32	62
33	63
34	64
35	65
36	66
37	67
38	68
39	69

TABLE 1-continued

N-terminal amino acid position	C-terminal amino acid position
40	70
41	71
42	72
43	73
44	74
<u>32-mer peptides of SEQ ID NO: 2</u>	
1	32
2	33
3	34
4	35
5	36
6	37
7	38
8	39
9	40
10	41
11	42
12	43
13	44
14	45
15	46
16	47
17	48
18	49
19	50
20	51
21	52
22	53
23	54
24	55
25	56
26	57
27	58
28	59
29	60
30	61
31	62
32	63
33	64
34	65
35	66
36	67
37	68
38	69
39	70
40	71
41	72
42	73
43	74
<u>33-mer peptides of SEQ ID NO: 2</u>	
1	33
2	34
3	35
4	36
5	37
6	38
7	39
8	40
9	41
10	42
11	43
12	44
13	45
14	46
15	47
16	48
17	49
18	50
19	51
20	52
21	53

TABLE 1-continued

N-terminal amino acid position	C-terminal amino acid position
22	54
23	55
24	56
25	57
26	58
27	59
28	60
29	61
30	62
31	63
32	64
33	65
34	66
35	67
36	68
37	69
38	70
39	71
40	72
41	73
42	74
34-mer peptides of SEQ ID NO: 2	
1	34
2	35
3	36
4	37
5	38
6	39
7	40
8	41
9	42
10	43
11	44
12	45
13	46
14	47
15	48
16	49
17	50
18	51
19	52
20	53
21	54
22	55
23	56
24	57
25	58
26	59
27	60
28	61
29	62
30	63
31	64
32	65
33	66
34	67
35	68
36	69
37	70
38	71
39	72
40	73
41	74
35-mer peptides of SEQ ID NO: 2	
1	35
2	36
3	37
4	38
5	39
6	40
7	41

TABLE 1-continued

N-terminal amino acid position	C-terminal amino acid position
8	42
9	43
10	44
11	45
12	46
13	47
14	48
15	49
16	50
17	51
18	52
19	53
20	54
21	55
22	56
23	57
24	58
25	59
26	60
27	61
28	62
29	63
30	64
31	65
32	66
33	67
34	68
35	69
36	70
37	71
38	72
39	73
40	74
36-mer peptides of SEQ ID NO: 2	
1	36
2	37
3	38
4	39
5	40
6	41
7	42
8	43
9	44
10	45
11	46
12	47
13	48
14	49
15	50
16	51
17	52
18	53
19	54
20	55
21	56
22	57
23	58
24	59
25	60
26	61
27	62
28	63
29	64
30	65
31	66
32	67
33	68
34	69
35	70
36	71
37	72
38	73

TABLE 1-continued

N-terminal amino acid position	C-terminal amino acid position
39	74
<u>37-mer peptides of SEQ ID NO: 2</u>	
1	37
2	38
3	39
4	40
5	41
6	42
7	43
8	44
9	45
10	46
11	47
12	48
13	49
14	50
15	51
16	52
17	53
18	54
19	55
20	56
21	57
22	58
23	59
24	60
25	61
26	62
27	63
28	64
29	65
30	66
31	67
32	68
33	69
34	70
35	71
36	72
37	73
38	74
<u>38-mer peptides of SEQ ID NO: 2</u>	
1	38
2	39
3	40
4	41
5	42
6	43
7	44
8	45
9	46
10	47
11	48
12	49
13	50
14	51
15	52
16	53
17	54
18	55
19	56
20	57
21	58
22	59
23	60
24	61
25	62
26	63
27	64
28	65
29	66
30	67

TABLE 1-continued

N-terminal amino acid position	C-terminal amino acid position
31	68
32	69
33	70
34	71
35	72
36	73
37	74
<u>39-mer peptides of SEQ ID NO: 2</u>	
1	39
2	40
3	41
4	42
5	43
6	44
7	45
8	46
9	47
10	48
11	49
12	50
13	51
14	52
15	53
16	54
17	55
18	56
19	57
20	58
21	59
22	60
23	61
24	62
25	63
26	64
27	65
28	66
29	67
30	68
31	69
32	70
33	71
34	72
35	73
36	74
<u>40-mer peptides of SEQ ID NO: 2</u>	
1	40
2	41
3	42
4	43
5	44
6	45
7	46
8	47
9	48
10	49
11	50
12	51
13	52
14	53
15	54
16	55
17	56
18	57
19	58
20	59
21	60
22	61
23	62
24	63
25	64
26	65

TABLE 1-continued

N-terminal amino acid position	C-terminal amino acid position
27	66
28	67
29	68
30	69
31	70
32	71
33	72
34	73
35	74
41-mer peptides of SEQ ID NO: 2	
1	41
2	42
3	43
4	44
5	45
6	46
7	47
8	48
9	49
10	50
11	51
12	52
13	53
14	54
15	55
16	56
17	57
18	58
19	59
20	60
21	61
22	62
23	63
24	64
25	65
26	66
27	67
28	68
29	69
30	70
31	71
32	72
33	73
34	74
42-mer peptides of SEQ ID NO: 2	
1	42
2	43
3	44
4	45
5	46
6	47
7	48
8	49
9	50
10	51
11	52
12	53
13	54
14	55
15	56
16	57
17	58
18	59
19	60
20	61
21	62
22	63
23	64
24	65
25	66
26	67

TABLE 1-continued

N-terminal amino acid position	C-terminal amino acid position
27	68
28	69
29	70
30	71
31	72
32	73
33	74
43-mer peptides of SEQ ID NO: 2	
1	43
2	44
3	45
4	46
5	47
6	48
7	49
8	50
9	51
10	52
11	53
12	54
13	55
14	56
15	57
16	58
17	59
18	60
19	61
20	62
21	63
22	64
23	65
24	66
25	67
26	68
27	69
28	70
29	71
30	72
31	73
32	74
44-mer peptides of SEQ ID NO: 2	
1	44
2	45
3	46
4	47
5	48
6	49
7	50
8	51
9	52
10	53
11	54
12	55
13	56
14	57
15	58
16	59
17	60
18	61
19	62
20	63
21	64
22	65
23	66
24	67
25	68
26	69
27	70
28	71
29	72
30	73

TABLE 1-continued

N-terminal amino acid position	C-terminal amino acid position
31	74
<u>45-mer peptides of SEQ ID NO: 2</u>	
1	45
2	46
3	47
4	48
5	49
6	50
7	51
8	52
9	53
10	54
11	55
12	56
13	57
14	58
15	59
16	60
17	61
18	62
19	63
20	64
21	65
22	66
23	67
24	68
25	69
26	70
27	71
28	72
29	73
30	74
<u>46-mer peptides of SEQ ID NO: 2</u>	
1	46
2	47
3	48
4	49
5	50
6	51
7	52
8	53
9	54
10	55
11	56
12	57
13	58
14	59
15	60
16	61
17	62
18	63
19	64
20	65
21	66
22	67
23	68
24	69
25	70
26	71
27	72
28	73
29	74
<u>47-mer peptides of SEQ ID NO: 2</u>	
1	47
2	48
3	49
4	50
5	51
6	52
7	53

TABLE 1-continued

N-terminal amino acid position	C-terminal amino acid position
8	54
9	55
10	56
11	57
12	58
13	59
14	60
15	61
16	62
17	63
18	64
19	65
20	66
21	67
22	68
23	69
24	70
25	71
26	72
27	73
28	74
<u>48-mer peptides of SEQ ID NO: 2</u>	
1	48
2	49
3	50
4	51
5	52
6	53
7	54
8	55
9	56
10	57
11	58
12	59
13	60
14	61
15	62
16	63
17	64
18	65
19	66
20	67
21	68
22	69
23	70
24	71
25	72
26	73
27	74
<u>49-mer peptides of SEQ ID NO: 2</u>	
1	49
2	50
3	51
4	52
5	53
6	54
7	55
8	56
9	57
10	58
11	59
12	60
13	61
14	62
15	63
16	64
17	65
18	66
19	67
20	68
21	69

TABLE 1-continued

N-terminal amino acid position	C-terminal amino acid position
22	70
23	71
24	72
25	73
26	74
<u>50-mer peptides of SEQ ID NO: 2</u>	
1	50
2	51
3	52
4	53
5	54
6	55
7	56
8	57
9	58
10	59
11	60
12	61
13	62
14	63
15	64
16	65
17	66
18	67
19	68
20	69
21	70
22	71
23	72
24	73
25	74
<u>51-mer peptides of SEQ ID NO: 2</u>	
1	51
2	52
3	53
4	54
5	55
6	56
7	57
8	58
9	59
10	60
11	61
12	62
13	63
14	64
15	65
16	66
17	67
18	68
19	69
20	70
21	71
22	72
23	73
24	74
<u>52-mer peptides of SEQ ID NO: 2</u>	
1	52
2	53
3	54
4	55
5	56
6	57
7	58
8	59
9	60
10	61
11	62
12	63
13	64

TABLE 1-continued

N-terminal amino acid position	C-terminal amino acid position
14	65
15	66
16	67
17	68
18	69
19	70
20	71
21	72
22	73
23	74
<u>53-mer peptides of SEQ ID NO: 2</u>	
1	53
2	54
3	55
4	56
5	57
6	58
7	59
8	60
9	61
10	62
11	63
12	64
13	65
14	66
15	67
16	68
17	69
18	70
19	71
20	72
21	73
22	74
<u>54-mer peptides of SEQ ID NO: 2</u>	
1	54
2	55
3	56
4	57
5	58
6	59
7	60
8	61
9	62
10	63
11	64
12	65
13	66
14	67
15	68
16	69
17	70
18	71
19	72
20	73
21	74
<u>55-mer peptides of SEQ ID NO: 2</u>	
1	55
2	56
3	57
4	58
5	59
6	60
7	61
8	62
9	63
10	64
11	65
12	66
13	67
14	68

TABLE 1-continued

N-terminal amino acid position	C-terminal amino acid position
15	69
16	70
17	71
18	72
19	73
20	74
56-mer peptides of SEQ ID NO: 2	
1	56
2	57
3	58
4	59
5	60
6	61
7	62
8	63
9	64
10	65
11	66
12	67
13	68
14	69
15	70
16	71
17	72
18	73
19	74
57-mer peptides of SEQ ID NO: 2	
1	57
2	58
3	59
4	60
5	61
6	62
7	63
8	64
9	65
10	66
11	67
12	68
13	69
14	70
15	71
16	72
17	73
18	74
58-mer peptides of SEQ ID NO: 2	
1	58
2	59
3	60
4	61
5	62
6	63
7	64
8	65
9	66
10	67
11	68
12	69
13	70
14	71
15	72
16	73
17	74
59-mer peptides of SEQ ID NO: 2	
1	59
2	60
3	61
4	62
5	63

TABLE 1-continued

N-terminal amino acid position	C-terminal amino acid position
6	64
7	65
8	66
9	67
10	68
11	69
12	70
13	71
14	72
15	73
16	74
60-mer peptides of SEQ ID NO: 2	
1	60
2	61
3	62
4	63
5	64
6	65
7	66
8	67
9	68
10	69
11	70
12	71
13	72
14	73
15	74
61-mer peptides of SEQ ID NO: 2	
1	61
2	62
3	63
4	64
5	65
6	66
7	67
8	68
9	69
10	70
11	71
12	72
13	73
14	74
62-mer peptides of SEQ ID NO: 2	
1	62
2	63
3	64
4	65
5	66
6	67
7	68
8	69
9	70
10	71
11	72
12	73
13	74
63-mer peptides of SEQ ID NO: 2	
1	63
2	64
3	65
4	66
5	67
6	68
7	69
8	70
9	71
10	72
11	73
12	74

TABLE 1-continued

N-terminal amino acid position	C-terminal amino acid position
<u>64-mer peptides of SEQ ID NO: 2</u>	
1	64
2	65
3	66
4	67
5	68
6	69
7	70
8	71
9	72
10	73
11	74
<u>65-mer peptides of SEQ ID NO: 2</u>	
1	65
2	66
3	67
4	68
5	69
6	70
7	71
8	72
9	73
10	74
<u>66-mer peptides of SEQ ID NO: 2</u>	
1	66
2	67
3	68
4	69
5	70
6	71
7	72
8	73
9	74
<u>67-mer peptides of SEQ ID NO: 2</u>	
1	67
2	68
3	69
4	70
5	71
6	72
7	73
8	74
<u>68-mer peptides of SEQ ID NO: 2</u>	
1	68
2	69
3	70
4	71
5	72
6	73
7	74
<u>69-mer peptides of SEQ ID NO: 2</u>	
1	69
2	70
3	71
4	72
5	73
6	74
<u>70-mer peptides of SEQ ID NO: 2</u>	
1	70
2	71
3	72
4	73
5	74

TABLE 1-continued

N-terminal amino acid position	C-terminal amino acid position
<u>71-mer peptides of SEQ ID NO: 2</u>	
1	71
2	72
3	73
4	74
<u>72-mer peptides of SEQ ID NO: 2</u>	
1	72
2	73
3	74
<u>73-mer peptides of SEQ ID NO: 2</u>	
1	73
2	74

[0123]

TABLE 2

N-terminal amino acid position	C-terminal amino acid position
<u>5-mer peptides of SEQ ID NO: 3</u>	
1	5
2	6
3	7
4	8
5	9
6	10
7	11
8	12
9	13
10	14
11	15
12	16
13	17
14	18
15	19
16	20
17	21
18	22
19	23
20	24
21	25
22	26
23	27
24	28
25	29
26	30
27	31
28	32
29	33
30	34
31	35
32	36
33	37
34	38
35	39
36	40
37	41
38	42
39	43
40	44
41	45
42	46
43	47
44	48
45	49

TABLE 2-continued

N-terminal amino acid position	C-terminal amino acid position
46	50
47	51
48	52
49	53
50	54
51	55
52	56
53	57
54	58
55	59
56	60
57	61
58	62
59	63
60	64
61	65
62	66
63	67
64	68
65	69
66	70
67	71
68	72
69	73
70	74
71	75
72	76
73	77
74	78
75	79
76	80
77	81
78	82
79	83
80	84
81	85
82	86
83	87
84	88
85	89
6-mer peptides of SEQ ID NO: 3	
1	6
2	7
3	8
4	9
5	10
6	11
7	12
8	13
9	14
10	15
11	16
12	17
13	18
14	19
15	20
16	21
17	22
18	23
19	24
20	25
21	26
22	27
23	28
24	29
25	30
26	31
27	32
28	33
29	34
30	35
31	36

TABLE 2-continued

N-terminal amino acid position	C-terminal amino acid position
32	37
33	38
34	39
35	40
36	41
37	42
38	43
39	44
40	45
41	46
42	47
43	48
44	49
45	50
46	51
47	52
48	53
49	54
50	55
51	56
52	57
53	58
54	59
55	60
56	61
57	62
58	63
59	64
60	65
61	66
62	67
63	68
64	69
65	70
66	71
67	72
68	73
69	74
70	75
71	76
72	77
73	78
74	79
75	80
76	81
77	82
78	83
79	84
80	85
81	86
82	87
83	88
84	89
7-mer peptides of SEQ ID NO: 3	
1	7
2	8
3	9
4	10
5	11
6	12
7	13
8	14
9	15
10	16
11	17
12	18
13	19
14	20
15	21
16	22
17	23
18	24

TABLE 2-continued

N-terminal amino acid position	C-terminal amino acid position
19	25
20	26
21	27
22	28
23	29
24	30
25	31
26	32
27	33
28	34
29	35
30	36
31	37
32	38
33	39
34	40
35	41
36	42
37	43
38	44
39	45
40	46
41	47
42	48
43	49
44	50
45	51
46	52
47	53
48	54
49	55
50	56
51	57
52	58
53	59
54	60
55	61
56	62
57	63
58	64
59	65
60	66
61	67
62	68
63	69
64	70
65	71
66	72
67	73
68	74
69	75
70	76
71	77
72	78
73	79
74	80
75	81
76	82
77	83
78	84
79	85
80	86
81	87
82	88
83	89
8-mer peptides of SEQ ID NO: 3	
1	8
2	9
3	10
4	11
5	12
6	13

TABLE 2-continued

N-terminal amino acid position	C-terminal amino acid position
7	14
8	15
9	16
10	17
11	18
12	19
13	20
14	21
15	22
16	23
17	24
18	25
19	26
20	27
21	28
22	29
23	30
24	31
25	32
26	33
27	34
28	35
29	36
30	37
31	38
32	39
33	40
34	41
35	42
36	43
37	44
38	45
39	46
40	47
41	48
42	49
43	50
44	51
45	52
46	53
47	54
48	55
49	56
50	57
51	58
52	59
53	60
54	61
55	62
56	63
57	64
58	65
59	66
60	67
61	68
62	69
63	70
64	71
65	72
66	73
67	74
68	75
69	76
70	77
71	78
72	79
73	80
74	81
75	82
76	83
77	84
78	85
79	86

TABLE 2-continued

N-terminal amino acid position	C-terminal amino acid position
80	87
81	88
82	89
9-mer peptides of SEQ ID NO: 3	
1	9
2	10
3	11
4	12
5	13
6	14
7	15
8	16
9	17
10	18
11	19
12	20
13	21
14	22
15	23
16	24
17	25
18	26
19	27
20	28
21	29
22	30
23	31
24	32
25	33
26	34
27	35
28	36
29	37
30	38
31	39
32	40
33	41
34	42
35	43
36	44
37	45
38	46
39	47
40	48
41	49
42	50
43	51
44	52
45	53
46	54
47	55
48	56
49	57
50	58
51	59
52	60
53	61
54	62
55	63
56	64
57	65
58	66
59	67
60	68
61	69
62	70
63	71
64	72
65	73
66	74
67	75
68	76

TABLE 2-continued

N-terminal amino acid position	C-terminal amino acid position
69	77
70	78
71	79
72	80
73	81
74	82
75	83
76	84
77	85
78	86
79	87
80	88
81	89
10-mer peptides of SEQ ID NO: 3	
1	10
2	11
3	12
4	13
5	14
6	15
7	16
8	17
9	18
10	19
11	20
12	21
13	22
14	23
15	24
16	25
17	26
18	27
19	28
20	29
21	30
22	31
23	32
24	33
25	34
26	35
27	36
28	37
29	38
30	39
31	40
32	41
33	42
34	43
35	44
36	45
37	46
38	47
39	48
40	49
41	50
42	51
43	52
44	53
45	54
46	55
47	56
48	57
49	58
50	59
51	60
52	61
53	62
54	63
55	64
56	65
57	66
58	67

TABLE 2-continued

N-terminal amino acid position	C-terminal amino acid position
59	68
60	69
61	70
62	71
63	72
64	73
65	74
66	75
67	76
68	77
69	78
70	79
71	80
72	81
73	82
74	83
75	84
76	85
77	86
78	87
79	88
80	89
<u>11-mer peptides of SEQ ID NO: 3</u>	
1	11
2	12
3	13
4	14
5	15
6	16
7	17
8	18
9	19
10	20
11	21
12	22
13	23
14	24
15	25
16	26
17	27
18	28
19	29
20	30
21	31
22	32
23	33
24	34
25	35
26	36
27	37
28	38
29	39
30	40
31	41
32	42
33	43
34	44
35	45
36	46
37	47
38	48
39	49
40	50
41	51
42	52
43	53
44	54
45	55
46	56
47	57
48	58
49	59

TABLE 2-continued

N-terminal amino acid position	C-terminal amino acid position
50	60
51	61
52	62
53	63
54	64
55	65
56	66
57	67
58	68
59	69
60	70
61	71
62	72
63	73
64	74
65	75
66	76
67	77
68	78
69	79
70	80
71	81
72	82
73	83
74	84
75	85
76	86
77	87
78	88
79	89
<u>12-mer peptides of SEQ ID NO: 3</u>	
1	12
2	13
3	14
4	15
5	16
6	17
7	18
8	19
9	20
10	21
11	22
12	23
13	24
14	25
15	26
16	27
17	28
18	29
19	30
20	31
21	32
22	33
23	34
24	35
25	36
26	37
27	38
28	39
29	40
30	41
31	42
32	43
33	44
34	45
35	46
36	47
37	48
38	49
39	50
40	51
41	52

TABLE 2-continued

N-terminal amino acid position	C-terminal amino acid position
42	53
43	54
44	55
45	56
46	57
47	58
48	59
49	60
50	61
51	62
52	63
53	64
54	65
55	66
56	67
57	68
58	69
59	70
60	71
61	72
62	73
63	74
64	75
65	76
66	77
67	78
68	79
69	80
70	81
71	82
72	83
73	84
74	85
75	86
76	87
77	88
78	89
13-mer peptides of SEQ ID NO: 3	
1	13
2	14
3	15
4	16
5	17
6	18
7	19
8	20
9	21
10	22
11	23
12	24
13	25
14	26
15	27
16	28
17	29
18	30
19	31
20	32
21	33
22	34
23	35
24	36
25	37
26	38
27	39
28	40
29	41
30	42
31	43
32	44
33	45
34	46

TABLE 2-continued

N-terminal amino acid position	C-terminal amino acid position
35	47
36	48
37	49
38	50
39	51
40	52
41	53
42	54
43	55
44	56
45	57
46	58
47	59
48	60
49	61
50	62
51	63
52	64
53	65
54	66
55	67
56	68
57	69
58	70
59	71
60	72
61	73
62	74
63	75
64	76
65	77
66	78
67	79
68	80
69	81
70	82
71	83
72	84
73	85
74	86
75	87
76	88
77	89
14-mer peptides of SEQ ID NO: 3	
1	14
2	15
3	16
4	17
5	18
6	19
7	20
8	21
9	22
10	23
11	24
12	25
13	26
14	27
15	28
16	29
17	30
18	31
19	32
20	33
21	34
22	35
23	36
24	37
25	38
26	39
27	40
28	41

TABLE 2-continued

N-terminal amino acid position	C-terminal amino acid position
29	42
30	43
31	44
32	45
33	46
34	47
35	48
36	49
37	50
38	51
39	52
40	53
41	54
42	55
43	56
44	57
45	58
46	59
47	60
48	61
49	62
50	63
51	64
52	65
53	66
54	67
55	68
56	69
57	70
58	71
59	72
60	73
61	74
62	75
63	76
64	77
65	78
66	79
67	80
68	81
69	82
70	83
71	84
72	85
73	86
74	87
75	88
76	89
15-mer peptides of SEQ ID NO: 3	
1	15
2	16
3	17
4	18
5	19
6	20
7	21
8	22
9	23
10	24
11	25
12	26
13	27
14	28
15	29
16	30
17	31
18	32
19	33
20	34
21	35
22	36
23	37

TABLE 2-continued

N-terminal amino acid position	C-terminal amino acid position
24	38
25	39
26	40
27	41
28	42
29	43
30	44
31	45
32	46
33	47
34	48
35	49
36	50
37	51
38	52
39	53
40	54
41	55
42	56
43	57
44	58
45	59
46	60
47	61
48	62
49	63
50	64
51	65
52	66
53	67
54	68
55	69
56	70
57	71
58	72
59	73
60	74
61	75
62	76
63	77
64	78
65	79
66	80
67	81
68	82
69	83
70	84
71	85
72	86
73	87
74	88
75	89
16-mer peptides of SEQ ID NO: 3	
1	16
2	17
3	18
4	19
5	20
6	21
7	22
8	23
9	24
10	25
11	26
12	27
13	28
14	29
15	30
16	31
17	32
18	33
19	34

TABLE 2-continued

N-terminal amino acid position	C-terminal amino acid position
20	35
21	36
22	37
23	38
24	39
25	40
26	41
27	42
28	43
29	44
30	45
31	46
32	47
33	48
34	49
35	50
36	51
37	52
38	53
39	54
40	55
41	56
42	57
43	58
44	59
45	60
46	61
47	62
48	63
49	64
50	65
51	66
52	67
53	68
54	69
55	70
56	71
57	72
58	73
59	74
60	75
61	76
62	77
63	78
64	79
65	80
66	81
67	82
68	83
69	84
70	85
71	86
72	87
73	88
74	89
17-mer peptides of SEQ ID NO: 3	
1	17
2	18
3	19
4	20
5	21
6	22
7	23
8	24
9	25
10	26
11	27
12	28
13	29
14	30
15	31
16	32

TABLE 2-continued

N-terminal amino acid position	C-terminal amino acid position
17	33
18	34
19	35
20	36
21	37
22	38
23	39
24	40
25	41
26	42
27	43
28	44
29	45
30	46
31	47
32	48
33	49
34	50
35	51
36	52
37	53
38	54
39	55
40	56
41	57
42	58
43	59
44	60
45	61
46	62
47	63
48	64
49	65
50	66
51	67
52	68
53	69
54	70
55	71
56	72
57	73
58	74
59	75
60	76
61	77
62	78
63	79
64	80
65	81
66	82
67	83
68	84
69	85
70	86
71	87
72	88
73	89
18-mer peptides of SEQ ID NO: 3	
1	18
2	19
3	20
4	21
5	22
6	23
7	24
8	25
9	26
10	27
11	28
12	29
13	30
14	31

TABLE 2-continued

N-terminal amino acid position	C-terminal amino acid position
15	32
16	33
17	34
18	35
19	36
20	37
21	38
22	39
23	40
24	41
25	42
26	43
27	44
28	45
29	46
30	47
31	48
32	49
33	50
34	51
35	52
36	53
37	54
38	55
39	56
40	57
41	58
42	59
43	60
44	61
45	62
46	63
47	64
48	65
49	66
50	67
51	68
52	69
53	70
54	71
55	72
56	73
57	74
58	75
59	76
60	77
61	78
62	79
63	80
64	81
65	82
66	83
67	84
68	85
69	86
70	87
71	88
72	89
19-mer peptides of SEQ ID NO: 3	
1	19
2	20
3	21
4	22
5	23
6	24
7	25
8	26
9	27
10	28
11	29
12	30
13	31

TABLE 2-continued

N-terminal amino acid position	C-terminal amino acid position
14	32
15	33
16	34
17	35
18	36
19	37
20	38
21	39
22	40
23	41
24	42
25	43
26	44
27	45
28	46
29	47
30	48
31	49
32	50
33	51
34	52
35	53
36	54
37	55
38	56
39	57
40	58
41	59
42	60
43	61
44	62
45	63
46	64
47	65
48	66
49	67
50	68
51	69
52	70
53	71
54	72
55	73
56	74
57	75
58	76
59	77
60	78
61	79
62	80
63	81
64	82
65	83
66	84
67	85
68	86
69	87
70	88
71	89
20-mer peptides of SEQ ID NO: 3	
1	20
2	21
3	22
4	23
5	24
6	25
7	26
8	27
9	28
10	29
11	30
12	31
13	32

TABLE 2-continued

N-terminal amino acid position	C-terminal amino acid position
14	33
15	34
16	35
17	36
18	37
19	38
20	39
21	40
22	41
23	42
24	43
25	44
26	45
27	46
28	47
29	48
30	49
31	50
32	51
33	52
34	53
35	54
36	55
37	56
38	57
39	58
40	59
41	60
42	61
43	62
44	63
45	64
46	65
47	66
48	67
49	68
50	69
51	70
52	71
53	72
54	73
55	74
56	75
57	76
58	77
59	78
60	79
61	80
62	81
63	82
64	83
65	84
66	85
67	86
68	87
69	88
70	89
21-mer peptides of SEQ ID NO: 3	
1	21
2	22
3	23
4	24
5	25
6	26
7	27
8	28
9	29
10	30
11	31
12	32
13	33
14	34

TABLE 2-continued

N-terminal amino acid position	C-terminal amino acid position
15	35
16	36
17	37
18	38
19	39
20	40
21	41
22	42
23	43
24	44
25	45
26	46
27	47
28	48
29	49
30	50
31	51
32	52
33	53
34	54
35	55
36	56
37	57
38	58
39	59
40	60
41	61
42	62
43	63
44	64
45	65
46	66
47	67
48	68
49	69
50	70
51	71
52	72
53	73
54	74
55	75
56	76
57	77
58	78
59	79
60	80
61	81
62	82
63	83
64	84
65	85
66	86
67	87
68	88
69	89
22-mer peptides of SEQ ID NO: 3	
1	22
2	23
3	24
4	25
5	26
6	27
7	28
8	29
9	30
10	31
11	32
12	33
13	34
14	35
15	36
16	37

TABLE 2-continued

N-terminal amino acid position	C-terminal amino acid position
17	38
18	39
19	40
20	41
21	42
22	43
23	44
24	45
25	46
26	47
27	48
28	49
29	50
30	51
31	52
32	53
33	54
34	55
35	56
36	57
37	58
38	59
39	60
40	61
41	62
42	63
43	64
44	65
45	66
46	67
47	68
48	69
49	70
50	71
51	72
52	73
53	74
54	75
55	76
56	77
57	78
58	79
59	80
60	81
61	82
62	83
63	84
64	85
65	86
66	87
67	88
68	89
<u>23-mer peptides of SEQ ID NO: 3</u>	
1	23
2	24
3	25
4	26
5	27
6	28
7	29
8	30
9	31
10	32
11	33
12	34
13	35
14	36
15	37
16	38
17	39
18	40
19	41

TABLE 2-continued

N-terminal amino acid position	C-terminal amino acid position
20	42
21	43
22	44
23	45
24	46
25	47
26	48
27	49
28	50
29	51
30	52
31	53
32	54
33	55
34	56
35	57
36	58
37	59
38	60
39	61
40	62
41	63
42	64
43	65
44	66
45	67
46	68
47	69
48	70
49	71
50	72
51	73
52	74
53	75
54	76
55	77
56	78
57	79
58	80
59	81
60	82
61	83
62	84
63	85
64	86
65	87
66	88
67	89
<u>24-mer peptides of SEQ ID NO: 3</u>	
1	24
2	25
3	26
4	27
5	28
6	29
7	30
8	31
9	32
10	33
11	34
12	35
13	36
14	37
15	38
16	39
17	40
18	41
19	42
20	43
21	44
22	45
23	46

TABLE 2-continued

N-terminal amino acid position	C-terminal amino acid position
24	47
25	48
26	49
27	50
28	51
29	52
30	53
31	54
32	55
33	56
34	57
35	58
36	59
37	60
38	61
39	62
40	63
41	64
42	65
43	66
44	67
45	68
46	69
47	70
48	71
49	72
50	73
51	74
52	75
53	76
54	77
55	78
56	79
57	80
58	81
59	82
60	83
61	84
62	85
63	86
64	87
65	88
66	89
<u>25-mer peptides of SEQ ID NO: 3</u>	
1	25
2	26
3	27
4	28
5	29
6	30
7	31
8	32
9	33
10	34
11	35
12	36
13	37
14	38
15	39
16	40
17	41
18	42
19	43
20	44
21	45
22	46
23	47
24	48
25	49
26	50
27	51
28	52

TABLE 2-continued

N-terminal amino acid position	C-terminal amino acid position
29	53
30	54
31	55
32	56
33	57
34	58
35	59
36	60
37	61
38	62
39	63
40	64
41	65
42	66
43	67
44	68
45	69
46	70
47	71
48	72
49	73
50	74
51	75
52	76
53	77
54	78
55	79
56	80
57	81
58	82
59	83
60	84
61	85
62	86
63	87
64	88
65	89
<u>26-mer peptides of SEQ ID NO: 3</u>	
1	26
2	27
3	28
4	29
5	30
6	31
7	32
8	33
9	34
10	35
11	36
12	37
13	38
14	39
15	40
16	41
17	42
18	43
19	44
20	45
21	46
22	47
23	48
24	49
25	50
26	51
27	52
28	53
29	54
30	55
31	56
32	57
33	58
34	59

TABLE 2-continued

N-terminal amino acid position	C-terminal amino acid position
35	60
36	61
37	62
38	63
39	64
40	65
41	66
42	67
43	68
44	69
45	70
46	71
47	72
48	73
49	74
50	75
51	76
52	77
53	78
54	79
55	80
56	81
57	82
58	83
59	84
60	85
61	86
62	87
63	88
64	89
<u>27-mer peptides of SEQ ID NO: 3</u>	
1	27
2	28
3	29
4	30
5	31
6	32
7	33
8	34
9	35
10	36
11	37
12	38
13	39
14	40
15	41
16	42
17	43
18	44
19	45
20	46
21	47
22	48
23	49
24	50
25	51
26	52
27	53
28	54
29	55
30	56
31	57
32	58
33	59
34	60
35	61
36	62
37	63
38	64
39	65
40	66
41	67

TABLE 2-continued

N-terminal amino acid position	C-terminal amino acid position
42	68
43	69
44	70
45	71
46	72
47	73
48	74
49	75
50	76
51	77
52	78
53	79
54	80
55	81
56	82
57	83
58	84
59	85
60	86
61	87
62	88
63	89
<u>28-mer peptides of SEQ ID NO: 3</u>	
1	28
2	29
3	30
4	31
5	32
6	33
7	34
8	35
9	36
10	37
11	38
12	39
13	40
14	41
15	42
16	43
17	44
18	45
19	46
20	47
21	48
22	49
23	50
24	51
25	52
26	53
27	54
28	55
29	56
30	57
31	58
32	59
33	60
34	61
35	62
36	63
37	64
38	65
39	66
40	67
41	68
42	69
43	70
44	71
45	72
46	73
47	74
48	75
49	76

TABLE 2-continued

N-terminal amino acid position	C-terminal amino acid position
50	77
51	78
52	79
53	80
54	81
55	82
56	83
57	84
58	85
59	86
60	87
61	88
62	89
29-mer peptides of SEQ ID NO: 3	
1	29
2	30
3	31
4	32
5	33
6	34
7	35
8	36
9	37
10	38
11	39
12	40
13	41
14	42
15	43
16	44
17	45
18	46
19	47
20	48
21	49
22	50
23	51
24	52
25	53
26	54
27	55
28	56
29	57
30	58
31	59
32	60
33	61
34	62
35	63
36	64
37	65
38	66
39	67
40	68
41	69
42	70
43	71
44	72
45	73
46	74
47	75
48	76
49	77
50	78
51	79
52	80
53	81
54	82
55	83
56	84
57	85
58	86

TABLE 2-continued

N-terminal amino acid position	C-terminal amino acid position
59	87
60	88
61	89
30-mer peptides of SEQ ID NO: 3	
1	30
2	31
3	32
4	33
5	34
6	35
7	36
8	37
9	38
10	39
11	40
12	41
13	42
14	43
15	44
16	45
17	46
18	47
19	48
20	49
21	50
22	51
23	52
24	53
25	54
26	55
27	56
28	57
29	58
30	59
31	60
32	61
33	62
34	63
35	64
36	65
37	66
38	67
39	68
40	69
41	70
42	71
43	72
44	73
45	74
46	75
47	76
48	77
49	78
50	79
51	80
52	81
53	82
54	83
55	84
56	85
57	86
58	87
59	88
60	89
31-mer peptides of SEQ ID NO: 3	
1	31
2	32
3	33
4	34
5	35
6	36

TABLE 2-continued

N-terminal amino acid position	C-terminal amino acid position
7	37
8	38
9	39
10	40
11	41
12	42
13	43
14	44
15	45
16	46
17	47
18	48
19	49
20	50
21	51
22	52
23	53
24	54
25	55
26	56
27	57
28	58
29	59
30	60
31	61
32	62
33	63
34	64
35	65
36	66
37	67
38	68
39	69
40	70
41	71
42	72
43	73
44	74
45	75
46	76
47	77
48	78
49	79
50	80
51	81
52	82
53	83
54	84
55	85
56	86
57	87
58	88
59	89
32-mer peptides of SEQ ID NO: 3	
1	32
2	33
3	34
4	35
5	36
6	37
7	38
8	39
9	40
10	41
11	42
12	43
13	44
14	45
15	46
16	47
17	48
18	49

TABLE 2-continued

N-terminal amino acid position	C-terminal amino acid position
19	50
20	51
21	52
22	53
23	54
24	55
25	56
26	57
27	58
28	59
29	60
30	61
31	62
32	63
33	64
34	65
35	66
36	67
37	68
38	69
39	70
40	71
41	72
42	73
43	74
44	75
45	76
46	77
47	78
48	79
49	80
50	81
51	82
52	83
53	84
54	85
55	86
56	87
57	88
58	89
33-mer peptides of SEQ ID NO: 3	
1	33
2	34
3	35
4	36
5	37
6	38
7	39
8	40
9	41
10	42
11	43
12	44
13	45
14	46
15	47
16	48
17	49
18	50
19	51
20	52
21	53
22	54
23	55
24	56
25	57
26	58
27	59
28	60
29	61
30	62
31	63

TABLE 2-continued

N-terminal amino acid position	C-terminal amino acid position
32	64
33	65
34	66
35	67
36	68
37	69
38	70
39	71
40	72
41	73
42	74
43	75
44	76
45	77
46	78
47	79
48	80
49	81
50	82
51	83
52	84
53	85
54	86
55	87
56	88
57	89
<u>34-mer peptides of SEQ ID NO: 3</u>	
1	34
2	35
3	36
4	37
5	38
6	39
7	40
8	41
9	42
10	43
11	44
12	45
13	46
14	47
15	48
16	49
17	50
18	51
19	52
20	53
21	54
22	55
23	56
24	57
25	58
26	59
27	60
28	61
29	62
30	63
31	64
32	65
33	66
34	67
35	68
36	69
37	70
38	71
39	72
40	73
41	74
42	75
43	76
44	77
45	78

TABLE 2-continued

N-terminal amino acid position	C-terminal amino acid position
46	79
47	80
48	81
49	82
50	83
51	84
52	85
53	86
54	87
55	88
56	89
<u>35-mer peptides of SEQ ID NO: 3</u>	
1	35
2	36
3	37
4	38
5	39
6	40
7	41
8	42
9	43
10	44
11	45
12	46
13	47
14	48
15	49
16	50
17	51
18	52
19	53
20	54
21	55
22	56
23	57
24	58
25	59
26	60
27	61
28	62
29	63
30	64
31	65
32	66
33	67
34	68
35	69
36	70
37	71
38	72
39	73
40	74
41	75
42	76
43	77
44	78
45	79
46	80
47	81
48	82
49	83
50	84
51	85
52	86
53	87
54	88
55	89
<u>36-mer peptides of SEQ ID NO: 3</u>	
1	36
2	37
3	38

TABLE 2-continued

N-terminal amino acid position	C-terminal amino acid position
4	39
5	40
6	41
7	42
8	43
9	44
10	45
11	46
12	47
13	48
14	49
15	50
16	51
17	52
18	53
19	54
20	55
21	56
22	57
23	58
24	59
25	60
26	61
27	62
28	63
29	64
30	65
31	66
32	67
33	68
34	69
35	70
36	71
37	72
38	73
39	74
40	75
41	76
42	77
43	78
44	79
45	80
46	81
47	82
48	83
49	84
50	85
51	86
52	87
53	88
54	89
<u>37-mer peptides of SEQ ID NO: 3</u>	
1	37
2	38
3	39
4	40
5	41
6	42
7	43
8	44
9	45
10	46
11	47
12	48
13	49
14	50
15	51
16	52
17	53
18	54
19	55
20	56

TABLE 2-continued

N-terminal amino acid position	C-terminal amino acid position
21	57
22	58
23	59
24	60
25	61
26	62
27	63
28	64
29	65
30	66
31	67
32	68
33	69
34	70
35	71
36	72
37	73
38	74
39	75
40	76
41	77
42	78
43	79
44	80
45	81
46	82
47	83
48	84
49	85
50	86
51	87
52	88
53	89
<u>38-mer peptides of SEQ ID NO: 3</u>	
1	38
2	39
3	40
4	41
5	42
6	43
7	44
8	45
9	46
10	47
11	48
12	49
13	50
14	51
15	52
16	53
17	54
18	55
19	56
20	57
21	58
22	59
23	60
24	61
25	62
26	63
27	64
28	65
29	66
30	67
31	68
32	69
33	70
34	71
35	72
36	73
37	74
38	75

TABLE 2-continued

N-terminal amino acid position	C-terminal amino acid position
39	76
40	77
41	78
42	79
43	80
44	81
45	82
46	83
47	84
48	85
49	86
50	87
51	88
52	89
<u>39-mer peptides of SEQ ID NO: 3</u>	
1	39
2	40
3	41
4	42
5	43
6	44
7	45
8	46
9	47
10	48
11	49
12	50
13	51
14	52
15	53
16	54
17	55
18	56
19	57
20	58
21	59
22	60
23	61
24	62
25	63
26	64
27	65
28	66
29	67
30	68
31	69
32	70
33	71
34	72
35	73
36	74
37	75
38	76
39	77
40	78
41	79
42	80
43	81
44	82
45	83
46	84
47	85
48	86
49	87
50	88
51	89
<u>40-mer peptides of SEQ ID NO: 3</u>	
1	40
2	41
3	42
4	43

TABLE 2-continued

N-terminal amino acid position	C-terminal amino acid position
5	44
6	45
7	46
8	47
9	48
10	49
11	50
12	51
13	52
14	53
15	54
16	55
17	56
18	57
19	58
20	59
21	60
22	61
23	62
24	63
25	64
26	65
27	66
28	67
29	68
30	69
31	70
32	71
33	72
34	73
35	74
36	75
37	76
38	77
39	78
40	79
41	80
42	81
43	82
44	83
45	84
46	85
47	86
48	87
49	88
50	89
<u>41-mer peptides of SEQ ID NO: 3</u>	
1	41
2	42
3	43
4	44
5	45
6	46
7	47
8	48
9	49
10	50
11	51
12	52
13	53
14	54
15	55
16	56
17	57
18	58
19	59
20	60
21	61
22	62
23	63
24	64
25	65

TABLE 2-continued

N-terminal amino acid position	C-terminal amino acid position
26	66
27	67
28	68
29	69
30	70
31	71
32	72
33	73
34	74
35	75
36	76
37	77
38	78
39	79
40	80
41	81
42	82
43	83
44	84
45	85
46	86
47	87
48	88
49	89
<u>42-mer peptides of SEQ ID NO: 3</u>	
1	42
2	43
3	44
4	45
5	46
6	47
7	48
8	49
9	50
10	51
11	52
12	53
13	54
14	55
15	56
16	57
17	58
18	59
19	60
20	61
21	62
22	63
23	64
24	65
25	66
26	67
27	68
28	69
29	70
30	71
31	72
32	73
33	74
34	75
35	76
36	77
37	78
38	79
39	80
40	81
41	82
42	83
43	84
44	85
45	86
46	87

TABLE 2-continued

N-terminal amino acid position	C-terminal amino acid position
47	88
48	89
<u>43-mer peptides of SEQ ID NO: 3</u>	
1	43
2	44
3	45
4	46
5	47
6	48
7	49
8	50
9	51
10	52
11	53
12	54
13	55
14	56
15	57
16	58
17	59
18	60
19	61
20	62
21	63
22	64
23	65
24	66
25	67
26	68
27	69
28	70
29	71
30	72
31	73
32	74
33	75
34	76
35	77
36	78
37	79
38	80
39	81
40	82
41	83
42	84
43	85
44	86
45	87
46	88
47	89
<u>44-mer peptides of SEQ ID NO: 3</u>	
1	44
2	45
3	46
4	47
5	48
6	49
7	50
8	51
9	52
10	53
11	54
12	55
13	56
14	57
15	58
16	59
17	60
18	61
19	62
20	63

TABLE 2-continued

N-terminal amino acid position	C-terminal amino acid position
21	64
22	65
23	66
24	67
25	68
26	69
27	70
28	71
29	72
30	73
31	74
32	75
33	76
34	77
35	78
36	79
37	80
38	81
39	82
40	83
41	84
42	85
43	86
44	87
45	88
46	89
45-mer peptides of SEQ ID NO: 3	
1	45
2	46
3	47
4	48
5	49
6	50
7	51
8	52
9	53
10	54
11	55
12	56
13	57
14	58
15	59
16	60
17	61
18	62
19	63
20	64
21	65
22	66
23	67
24	68
25	69
26	70
27	71
28	72
29	73
30	74
31	75
32	76
33	77
34	78
35	79
36	80
37	81
38	82
39	83
40	84
41	85
42	86
43	87
44	88
45	89

TABLE 2-continued

N-terminal amino acid position	C-terminal amino acid position
46-mer peptides of SEQ ID NO: 3	
1	46
2	47
3	48
4	49
5	50
6	51
7	52
8	53
9	54
10	55
11	56
12	57
13	58
14	59
15	60
16	61
17	62
18	63
19	64
20	65
21	66
22	67
23	68
24	69
25	70
26	71
27	72
28	73
29	74
30	75
31	76
32	77
33	78
34	79
35	80
36	81
37	82
38	83
39	84
40	85
41	86
42	87
43	88
44	89
47-mer peptides of SEQ ID NO: 3	
1	47
2	48
3	49
4	50
5	51
6	52
7	53
8	54
9	55
10	56
11	57
12	58
13	59
14	60
15	61
16	62
17	63
18	64
19	65
20	66
21	67
22	68
23	69
24	70
25	71

TABLE 2-continued

N-terminal amino acid position	C-terminal amino acid position
26	72
27	73
28	74
29	75
30	76
31	77
32	78
33	79
34	80
35	81
36	82
37	83
38	84
39	85
40	86
41	87
42	88
43	89
48-mer peptides of SEQ ID NO: 3	
1	48
2	49
3	50
4	51
5	52
6	53
7	54
8	55
9	56
10	57
11	58
12	59
13	60
14	61
15	62
16	63
17	64
18	65
19	66
20	67
21	68
22	69
23	70
24	71
25	72
26	73
27	74
28	75
29	76
30	77
31	78
32	79
33	80
34	81
35	82
36	83
37	84
38	85
39	86
40	87
41	88
42	89
49-mer peptides of SEQ ID NO: 3	
1	49
2	50
3	51
4	52
5	53
6	54
7	55
8	56
9	57

TABLE 2-continued

N-terminal amino acid position	C-terminal amino acid position
10	58
11	59
12	60
13	61
14	62
15	63
16	64
17	65
18	66
19	67
20	68
21	69
22	70
23	71
24	72
25	73
26	74
27	75
28	76
29	77
30	78
31	79
32	80
33	81
34	82
35	83
36	84
37	85
38	86
39	87
40	88
41	89
50-mer peptides of SEQ ID NO: 3	
1	50
2	51
3	52
4	53
5	54
6	55
7	56
8	57
9	58
10	59
11	60
12	61
13	62
14	63
15	64
16	65
17	66
18	67
19	68
20	69
21	70
22	71
23	72
24	73
25	74
26	75
27	76
28	77
29	78
30	79
31	80
32	81
33	82
34	83
35	84
36	85
37	86
38	87

TABLE 2-continued

N-terminal amino acid position	C-terminal amino acid position
39	88
40	89
<u>51-mer peptides of SEQ ID NO: 3</u>	
1	51
2	52
3	53
4	54
5	55
6	56
7	57
8	58
9	59
10	60
11	61
12	62
13	63
14	64
15	65
16	66
17	67
18	68
19	69
20	70
21	71
22	72
23	73
24	74
25	75
26	76
27	77
28	78
29	79
30	80
31	81
32	82
33	83
34	84
35	85
36	86
37	87
38	88
39	89
<u>52-mer peptides of SEQ ID NO: 3</u>	
1	52
2	53
3	54
4	55
5	56
6	57
7	58
8	59
9	60
10	61
11	62
12	63
13	64
14	65
15	66
16	67
17	68
18	69
19	70
20	71
21	72
22	73
23	74
24	75
25	76
26	77
27	78
28	79

TABLE 2-continued

N-terminal amino acid position	C-terminal amino acid position
29	80
30	81
31	82
32	83
33	84
34	85
35	86
36	87
37	88
38	89
<u>53-mer peptides of SEQ ID NO: 3</u>	
1	53
2	54
3	55
4	56
5	57
6	58
7	59
8	60
9	61
10	62
11	63
12	64
13	65
14	66
15	67
16	68
17	69
18	70
19	71
20	72
21	73
22	74
23	75
24	76
25	77
26	78
27	79
28	80
29	81
30	82
31	83
32	84
33	85
34	86
35	87
36	88
37	89
<u>54-mer peptides of SEQ ID NO: 3</u>	
1	54
2	55
3	56
4	57
5	58
6	59
7	60
8	61
9	62
10	63
11	64
12	65
13	66
14	67
15	68
16	69
17	70
18	71
19	72
20	73
21	74
22	75

TABLE 2-continued

N-terminal amino acid position	C-terminal amino acid position
23	76
24	77
25	78
26	79
27	80
28	81
29	82
30	83
31	84
32	85
33	86
34	87
35	88
36	89

55-mer peptides of SEQ ID NO: 3

1	55
2	56
3	57
4	58
5	59
6	60
7	61
8	62
9	63
10	64
11	65
12	66
13	67
14	68
15	69
16	70
17	71
18	72
19	73
20	74
21	75
22	76
23	77
24	78
25	79
26	80
27	81
28	82
29	83
30	84
31	85
32	86
33	87
34	88
35	89

56-mer peptides of SEQ ID NO: 3

1	56
2	57
3	58
4	59
5	60
6	61
7	62
8	63
9	64
10	65
11	66
12	67
13	68
14	69
15	70
16	71
17	72
18	73
19	74
20	75

TABLE 2-continued

N-terminal amino acid position	C-terminal amino acid position
21	76
22	77
23	78
24	79
25	80
26	81
27	82
28	83
29	84
30	85
31	86
32	87
33	88
34	89

57-mer peptides of SEQ ID NO: 3

1	57
2	58
3	59
4	60
5	61
6	62
7	63
8	64
9	65
10	66
11	67
12	68
13	69
14	70
15	71
16	72
17	73
18	74
19	75
20	76
21	77
22	78
23	79
24	80
25	81
26	82
27	83
28	84
29	85
30	86
31	87
32	88
33	89

58-mer peptides of SEQ ID NO: 3

1	58
2	59
3	60
4	61
5	62
6	63
7	64
8	65
9	66
10	67
11	68
12	69
13	70
14	71
15	72
16	73
17	74
18	75
19	76
20	77
21	78
22	79

TABLE 2-continued

N-terminal amino acid position	C-terminal amino acid position
23	80
24	81
25	82
26	83
27	84
28	85
29	86
30	87
31	88
32	89
59-mer peptides of SEQ ID NO: 3	
1	59
2	60
3	61
4	62
5	63
6	64
7	65
8	66
9	67
10	68
11	69
12	70
13	71
14	72
15	73
16	74
17	75
18	76
19	77
20	78
21	79
22	80
23	81
24	82
25	83
26	84
27	85
28	86
29	87
30	88
31	89
60-mer peptides of SEQ ID NO: 3	
1	60
2	61
3	62
4	63
5	64
6	65
7	66
8	67
9	68
10	69
11	70
12	71
13	72
14	73
15	74
16	75
17	76
18	77
19	78
20	79
21	80
22	81
23	82
24	83
25	84
26	85
27	86
28	87

TABLE 2-continued

N-terminal amino acid position	C-terminal amino acid position
29	88
30	89
5-mer peptides of SEQ ID NO: 4	
1	5
2	6
3	7
4	8
5	9
6	10
7	11
8	12
9	13
10	14
6-mer peptides of SEQ ID NO: 4	
1	6
2	7
3	8
4	9
5	10
6	11
7	12
8	13
9	14
7-mer peptides of SEQ ID NO: 4	
1	7
2	8
3	9
4	10
5	11
6	12
7	13
8	14
8-mer peptides of SEQ ID NO: 4	
1	8
2	9
3	10
4	11
5	12
6	13
7	14
61-mer peptides of SEQ ID NO: 3	
1	61
2	62
3	63
4	64
5	65
6	66
7	67
8	68
9	69
10	70
11	71
12	72
13	73
14	74
15	75
16	76
17	77
18	78
19	79
20	80
21	81
22	82
23	83
24	84
25	85
26	86
27	87

TABLE 2-continued

N-terminal amino acid position	C-terminal amino acid position
28	88
29	89
<u>62-mer peptides of SEQ ID NO: 3</u>	
1	62
2	63
3	64
4	65
5	66
6	67
7	68
8	69
9	70
10	71
11	72
12	73
13	74
14	75
15	76
16	77
17	78
18	79
19	80
20	81
21	82
22	83
23	84
24	85
25	86
26	87
27	88
28	89
<u>63-mer peptides of SEQ ID NO: 3</u>	
1	63
2	64
3	65
4	66
5	67
6	68
7	69
8	70
9	71
10	72
11	73
12	74
13	75
14	76
15	77
16	78
17	79
18	80
19	81
20	82
21	83
22	84
23	85
24	86
25	87
26	88
27	89
<u>64-mer peptides of SEQ ID NO: 3</u>	
1	64
2	65
3	66
4	67
5	68
6	69
7	70
8	71
9	72
10	73

TABLE 2-continued

N-terminal amino acid position	C-terminal amino acid position
11	74
12	75
13	76
14	77
15	78
16	79
17	80
18	81
19	82
20	83
21	84
22	85
23	86
24	87
25	88
26	89
<u>9-mer peptides of SEQ ID NO: 4</u>	
1	9
2	10
3	11
4	12
5	13
6	14
<u>10-mer peptides of SEQ ID NO: 4</u>	
1	10
2	11
3	12
4	13
5	14
<u>11-mer peptides of SEQ ID NO: 4</u>	
1	11
2	12
3	13
4	14
<u>12-mer peptides of SEQ ID NO: 4</u>	
1	12
2	13
3	14
<u>13-mer peptides of SEQ ID NO: 4</u>	
1	13
2	14
<u>65-mer peptides of SEQ ID NO: 3</u>	
1	65
2	66
3	67
4	68
5	69
6	70
7	71
8	72
9	73
10	74
11	75
12	76
13	77
14	78
15	79
16	80
17	81
18	82
19	83
20	84
21	85
22	86
23	87
24	88
25	89

TABLE 2-continued

N-terminal amino acid position	C-terminal amino acid position
<u>66-mer peptides of SEQ ID NO: 3</u>	
1	66
2	67
3	68
4	69
5	70
6	71
7	72
8	73
9	74
10	75
11	76
12	77
13	78
14	79
15	80
16	81
17	82
18	83
19	84
20	85
21	86
22	87
23	88
24	89
<u>67-mer peptides of SEQ ID NO: 3</u>	
1	67
2	68
3	69
4	70
5	71
6	72
7	73
8	74
9	75
10	76
11	77
12	78
13	79
14	80
15	81
16	82
17	83
18	84
19	85
20	86
21	87
22	88
23	89
<u>68-mer peptides of SEQ ID NO: 3</u>	
1	68
2	69
3	70
4	71
5	72
6	73
7	74
8	75
9	76
10	77
11	78
12	79
13	80
14	81
15	82
16	83
17	84
18	85
19	86
20	87

TABLE 2-continued

N-terminal amino acid position	C-terminal amino acid position
21	88
22	89
<u>69-mer peptides of SEQ ID NO: 3</u>	
1	69
2	70
3	71
4	72
5	73
6	74
7	75
8	76
9	77
10	78
11	79
12	80
13	81
14	82
15	83
16	84
17	85
18	86
19	87
20	88
21	89
<u>70-mer peptides of SEQ ID NO: 3</u>	
1	70
2	71
3	72
4	73
5	74
6	75
7	76
8	77
9	78
10	79
11	80
12	81
13	82
14	83
15	84
16	85
17	86
18	87
19	88
20	89
<u>71-mer peptides of SEQ ID NO: 3</u>	
1	71
2	72
3	73
4	74
5	75
6	76
7	77
8	78
9	79
10	80
11	81
12	82
13	83
14	84
15	85
16	86
17	87
18	88
19	89
<u>72-mer peptides of SEQ ID NO: 3</u>	
1	72
2	73
3	74

TABLE 2-continued

N-terminal amino acid position	C-terminal amino acid position
4	75
5	76
6	77
7	78
8	79
9	80
10	81
11	82
12	83
13	84
14	85
15	86
16	87
17	88
18	89
<u>73-mer peptides of SEQ ID NO: 3</u>	
1	73
2	74
3	75
4	76
5	77
6	78
7	79
8	80
9	81
10	82
11	83
12	84
13	85
14	86
15	87
16	88
17	89
<u>74-mer peptides of SEQ ID NO: 3</u>	
1	74
2	75
3	76
4	77
5	78
6	79
7	80
8	81
9	82
10	83
11	84
12	85
13	86
14	87
15	88
16	89
<u>75-mer peptides of SEQ ID NO: 3</u>	
1	75
2	76
3	77
4	78
5	79
6	80
7	81
8	82
9	83
10	84
11	85
12	86
13	87
14	88
15	89
<u>76-mer peptides of SEQ ID NO: 3</u>	
1	76
2	77

TABLE 2-continued

N-terminal amino acid position	C-terminal amino acid position
3	78
4	79
5	80
6	81
7	82
8	83
9	84
10	85
11	86
12	87
13	88
14	89
<u>77-mer peptides of SEQ ID NO: 3</u>	
1	77
2	70
3	71
4	72
5	73
6	74
7	75
8	76
9	77
10	78
11	79
12	80
13	81
14	82
15	83
16	84
17	85
18	86
19	87
20	88
21	89
<u>78-mer peptides of SEQ ID NO: 3</u>	
1	78
2	71
3	72
4	73
5	74
6	75
7	76
8	77
9	78
10	79
11	80
12	81
13	82
14	83
15	84
16	85
17	86
18	87
19	88
20	89
<u>79-mer peptides of SEQ ID NO: 3</u>	
1	79
2	72
3	73
4	74
5	75
6	76
7	77
8	78
9	79
10	80
11	81
12	82
13	83
14	84

TABLE 2-continued

N-terminal amino acid position	C-terminal amino acid position
15	85
16	86
17	87
18	88
19	89
80-mer peptides of SEQ ID NO: 3	
1	80
2	73
3	74
4	75
5	76
6	77
7	78
8	79
9	80
10	81
11	82
12	83
13	84
14	85
15	86
16	87
17	88
18	89
81-mer peptides of SEQ ID NO: 3	
1	81
2	82
3	83
4	84
5	85
6	86
7	87
8	88
9	89
82-mer peptides of SEQ ID NO: 3	
1	82
2	83
3	84
4	85
5	86
6	87
7	88
8	89
83-mer peptides of SEQ ID NO: 3	
1	83
2	84
3	85
4	86
5	87
6	88
7	89
84-mer peptides of SEQ ID NO: 3	
1	84
2	85
3	86
4	87
5	88
6	89
85-mer peptides of SEQ ID NO: 3	
1	85
2	86
3	87
4	88
5	89

TABLE 2-continued

N-terminal amino acid position	C-terminal amino acid position
86-mer peptides of SEQ ID NO: 3	
1	86
2	87
3	88
4	89
87-mer peptides of SEQ ID NO: 3	
1	87
2	88
3	89
88-mer peptides of SEQ ID NO: 3	
1	88
2	89

[0124]

TABLE 3

Name of enzyme	No. of cleavages	Positions of cleavage sites within SEQ ID No: 2 (cleavage site is C terminal to amino acid number)
Asparaginyl endopeptidase	4	16 25 39 52
Cyanogen bromide	3	18 59 62
Chymotrypsin-high specificity (C-term to [FYW], not before P)	8	3 6 8 15 38 54 63 65
Chymotrypsin-low specificity (C-term to [FYWML], not before P)	18	3 4 6 8 15 18 22 35 38 41 54 59 60 61 62 63 65 70
Glutamyl endopeptidase	2	30 45
Lysyl endopeptidase	5	11 12 21 57 69
(2-nitro-5-thiocyanobenzoic acid)	2	67 71
Pepsin (pH = 1.3)	23	2 3 3 4 5 6 7 8 15 21 22 35 37 38 40 41 53 54 61 62 63 64 65
Pepsin (pH > 2)	19	2 3 3 4 15 21 22 35 37 38 40 41 53 54 61 62 63 64 65
Proteinase K	34	1 2 3 4 6 8 9 14 15 20 22 25 27 28 29 32 34 35 36 38 39 41 42 48 49 52 54 55 60 61 63 65 71 74
Staphylococcal peptidase I	2	30 45
Thermolysin	26	1 2 3 8 13 14 19 21 27 28 33 34 35 37 41 47 48 54 58 59 60 61 62 64 70 73
Trypsin	5	11 12 21 57

[0125]

TABLE 4

Supe. 2/3 Negative Controls					
1 Ab	Normal 1	Normal 2	Normal 3	Normal 4	Normal 5
1/100	0.576	0.26	0.394	0.386	0.482
1/300	0.277	0.22	0.219	0.267	0.37
1/1000	0.143	0.147	0.142	0.132	0.134
1/100		1/300		1/1000	
*N = 5		*N = 5		*N = 5	
Mean: 0.4196		Mean: 0.2706		Mean: 0.1396	
*Highest: 0.576		*Highest: 0.37		*Highest: 0.147	
*Lowest: 0.26		*Lowest: 0.219		*Lowest: 0.132	
Variance = 0.316		Variance = 0.151		Variance = 0.015	
Standard Dev. = 0.079		S.D. = 0.03775		S.D. = 0.00375	
2 Stand. Dev. = 0.158		2 S.D. = 0.0755		2 S.D. = 0.0075	
3 Stand. Dev. = 0.237		3 S.D. = 0.11325		3 S.D. = 0.01125	
2x Mean: 0.8392		2x Mean: 0.5412		2x Mean: 0.2792	
2 S.D. + Mean: 0.5776		2 S.D. + Mean: 0.3461		2 S.D. + Mean: 0.1471	
3 S.D. + Mean: 0.6566		3 S.D. + Mean: 0.38385		3 S.D. + Mean: 0.15085	
Supe 2/3					
1 Ab	#179	#178	#80439	#80255	#99-01291
1/100	0.731	1.162	1.37	1.252	0.468
1/300	0.438	0.675	0.995	0.63	0.34
1/1000	0.22	0.382	0.396	0.282	0.147

[0126]

TABLE 5

Negative Controls					
1 Ab	Mollie	CN76AC	CN115A Jun. 11, 1998	CN125C	CN35B
1/100	1.738	2.074	2.096	2.494	1.815
1/300	1.091	1.499	1.284	1.63	1.055
1 Ab	Dixie	CN54B	CN105C	Presley	CN74H
1/100	1.823	2.429	2.127	1.532	1.703
1/300	1.336	1.698	1.496	0.947	1.114
1/100		1/300			
*N = 10		*N = 10			
Mean: 1.9831		Mean: 1.315			
*Highest: 2.494		*Highest: 1.698			
*Lowest: 1.532		*Lowest: 0.947			
Variance = 0.962		Variance = 0.751			
Standard Dev. = 0.2405		S.D. = 0.18775			
2 Stand. Dev. = 0.481		2 S.D. = 0.3755			
3 Stand. Dev. = 0.7215		3 S.D. = 0.56325			
2x Mean: 3.9662		2x Mean: 2.63			

TABLE 5-continued

Negative Controls					
2 S.D. + Mean: 2.4641			2 S.D. + Mean: 1.6905		
3 S.D. + Mean: 2.7046			3 S.D. + Mean: 1.87825		
Supe 2/3					
1 Ab	Indexx 1	Indexx 2	Indexx 3	Indexx 4	Indexx 5
1/100	2.409	2.029	2.061	2.821	2.11
1/300	1.618	1.565	1.278	2.422	1.24
1 Ab	Indexx 6	Indexx 7	Indexx 8	Indexx 9	Indexx 10
1/100	2.51	2.012	1.883	1.743	2.268
1/300	1.985	1.24	1.11	1.041	1.596
1 Ab	Indexx 11	Indexx 12	Indexx 13	Indexx 14	Indexx 15
1/100	2.685	2.669	2.788	1.999	3.18
1/300	1.79	2.052	1.992	1.385	2.919
1 Ab	Indexx 16	Indexx 17	Indexx 18	Indexx 19	Indexx 20
1/100	3.023	2.471	2.443	2.466	3.184
1/300	2.544	1.907	2.155	1.788	2.821

[0127]

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 4

<210> SEQ ID NO 1
 <211> LENGTH: 742
 <212> TYPE: DNA
 <213> ORGANISM: Anaplasma phagocytophilum
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (29)..(250)
 <223> OTHER INFORMATION: Coding sequence of processed MSP5 polypeptide
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (29)..(610)
 <223> OTHER INFORMATION: 5' and 3' ends of cloned MSP5 insert

<400> SEQUENCE: 1

```

aaataaggag gaataaacca tggcccttac tgtgtttctg gggatttcgt atgttaacaa      60
aaaggaata  ttacagtaca tgggtgctaa attaagtagt actgacactg tggttgaaca      120
agcgccgata ttgcttctt ttactgatct gataaatcaa gaaggtcagg ttgttagtag      180
cactgatttt gctgggaagc acatgttgct tatgttcggc ttttcttcat gcaagcacat      240
atgtcctgcg gaacttggtg tggatcgcga actttttaaac aagttgggtg agactgotga      300
taaacttcaa gctgttttca taactattga tcccaaaaat gacactgttg aaaggttaaa      360
tgagtatcac aaagcctttg atagtcgaat tcagatgctg actggggatg aggaagttat      420
tcgcaatgta gtgaataatt ataaggtata cgtaggtgag tctgatagtg aaggagatat      480
taatcactca tcgtttttgt acttggttga tgccgacgga agatatgtag ggcattttgc      540
tcctgatttt gatgaatacg aaagtcaagt aggtagactt tttgatttcg ttaataagta      600
ccctacttct aagggcgaat tcgaagctta cgtagaacaa aaactcatct cagaagagga      660
tctgaatagc gcgctcgacc atcatcatca tcatcattga gtttaaacgg tctccagctt      720
ggctgttttg gcggatgaga ga                                          742

```

<210> SEQ ID NO 2
 <211> LENGTH: 74
 <212> TYPE: PRT
 <213> ORGANISM: Anaplasma phagocytophilum

<400> SEQUENCE: 2

```

Thr Val Phe Leu Gly Tyr Ser Tyr Val Asn Lys Lys Gly Ile Phe Ser
1           5           10           15

Asp Met Gly Ala Lys Leu Ser Ser Thr Asp Thr Val Val Glu Gln Ala
          20           25           30

Pro Ile Leu Ala Ser Phe Thr Asp Leu Ile Asn Gln Glu Gly Gln Val
          35           40           45

Val Ser Ser Thr Asp Phe Ala Gly Lys His Met Leu Leu Met Phe Gly
          50           55           60

Phe Ser Ser Cys Lys His Ile Cys Pro Ala
65           70

```

<210> SEQ ID NO 3
 <211> LENGTH: 89
 <212> TYPE: PRT
 <213> ORGANISM: Anaplasma phagocytophilum

-continued

<400> SEQUENCE: 3

```

Met Lys Thr Phe Lys Thr Val Ser Asn Leu Leu Leu Val Ala Thr
1           5           10          15
Val Phe Leu Gly Tyr Ser Tyr Val Asn Lys Lys Gly Ile Phe Ser Asp
          20           25           30
Met Gly Ala Lys Leu Ser Ser Thr Asp Thr Val Val Glu Gln Ala Pro
          35           40           45
Ile Leu Ala Ser Phe Thr Asp Leu Ile Asn Gln Glu Gly Gln Val Val
          50           55           60
Ser Ser Thr Asp Phe Ala Gly Lys His Met Leu Leu Met Phe Gly Phe
65           70           75           80
Ser Ser Cys Lys His Ile Cys Pro Ala
          85

```

<210> SEQ ID NO 4

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Anaplasma phagocytophilum

<220> FEATURE:

<221> NAME/KEY: MISC_FEATURE

<222> LOCATION: (1)..(15)

<223> OTHER INFORMATION: MSP5 signal peptide

<400> SEQUENCE: 4

```

Met Lys Thr Phe Lys Thr Val Ser Asn Leu Leu Leu Val Ala
1           5           10          15

```

We claim:

1. An isolated, recombinant, or purified polypeptide comprising:

- a) SEQ ID NO 2;
- b) SEQ ID NO: 3;
- c) SEQ ID NO: 4;
- d) fragments of SEQ ID NOs: 2, 3, or 4;
- e) a polypeptide as set forth in Tables 1, 2, or 3;
- f) a variant polypeptide of SEQ ID NO: 2, 3, or 4, wherein said variant polypeptide specifically binds to an antibody that specifically binds to a polypeptide of SEQ ID NO: 2, 3, or 4;
- g) a variant polypeptide fragment of SEQ ID NO: 2, 3, or 4, wherein said variant polypeptide fragment specifically binds to an antibody that specifically binds to a polypeptide of SEQ ID NO: 2, 3, or 4 or a fragment of SEQ ID NO: 2, 3, or 4;
- h) a variant of a polypeptide as set forth in Table 1, 2 or 3, wherein said variant polypeptide specifically binds to an antibody that specifically binds to a polypeptide of SEQ ID NO: 2, 3, or 4 or a polypeptide as set forth in Table 1, 2 or 3;
- i) a heterologous polypeptide fused, in frame, to a polypeptide comprising: 1) SEQ ID NO: 2, 3, or 4; 2) fragments of SEQ ID NO: 2, 3, or 4; or 3) a polypeptide as set forth in Tables 1, 2 or 3;

j) a multimeric construct comprising: a) SEQ ID NO: 2, 3 or 4; b) fragments of SEQ ID NO: 2, 3 or 4; c) combinations of SEQ ID NO: 2, 3, 4, or fragments or variants thereof.

2. A composition comprising at least one isolated, recombinant, or purified polypeptide according to claim 1 and an additional component.

3. The composition according to claim 2, wherein said additional component is a solid support.

4. The composition according to claim 3, wherein said solid support is selected from the group consisting of microtiter wells, magnetic beads, non-magnetic beads, agarose beads, glass, cellulose, plastics, polyethylene, polypropylene, polyester, nitrocellulose, nylon, and polysulfone.

5. The composition according to claim 2, wherein said additional component is a pharmaceutically acceptable excipient.

6. The composition according to claim 3, wherein said solid support provides an array of polypeptides and said array of polypeptides is selected from the group consisting of:

- a) SEQ ID NO 2;
- b) SEQ ID NO: 3;
- c) SEQ ID NO: 4;
- d) fragments of SEQ ID NOs: 2, 3, or 4;
- e) a polypeptide as set forth in Tables 1, 2 or 3;
- f) a variant polypeptide of SEQ ID NO: 2, 3, or 4, wherein said variant polypeptide specifically binds to an antibody that specifically binds to a polypeptide of SEQ ID NO: 2, 3, or 4;

- g) a variant polypeptide fragment of SEQ ID NO: 2, 3, or 4, wherein said variant polypeptide fragment specifically binds to an antibody that specifically binds to a polypeptide of SEQ ID NO: 2, 3, or 4 or a fragment of SEQ ID NO: 2, 3, or 4;
- h) a variant of a polypeptide as set forth in Table 1 or 2, wherein said variant polypeptide specifically binds to an antibody that specifically binds to a polypeptide of SEQ ID NO: 2, 3, or 4 or a polypeptide as set forth in Table 1 or 2;
- i) a heterologous polypeptide fused, in frame, to a polypeptide comprising: 1) SEQ ID NO: 2, 3, or 4; 2) fragments of SEQ ID NO: 2, 3, or 4; or 3) a polypeptide as set forth in Tables 1 or 2; and
- j) combinations of said polypeptides and variants.
7. The composition of claim 6, further comprising an additional antigen of interest.
8. A method of binding an antibody to a polypeptide comprising contacting a sample containing an antibody with a polypeptide under conditions that allow for the formation of an antibody-antigen complex, wherein said polypeptide is selected from the group consisting of:
- SEQ ID NO 2;
 - SEQ ID NO: 3;
 - SEQ ID NO: 4;
 - fragments of SEQ ID NOs: 2, 3 or 4;
 - a polypeptide as set forth in Tables 1, 2 or 3;
 - a variant polypeptide of SEQ ID NO: 2, 3, or 4, wherein said variant polypeptide specifically binds to an antibody that specifically binds to a polypeptide of SEQ ID NO: 2, 3, or 4;
 - a variant polypeptide fragment of SEQ ID NO: 2, 3, or 4, wherein said variant polypeptide fragment specifically binds to an antibody that specifically binds to a polypeptide of SEQ ID NO: 2, 3, or 4 or a fragment of SEQ ID NO: 2, 3, or 4;
 - a variant of a polypeptide as set forth in Table 1, 2 or 3, wherein said variant polypeptide specifically binds to an antibody that specifically binds to a polypeptide of SEQ ID NO: 2, 3, or 4 or a polypeptide as set forth in Table 1, 2 or 3;
 - a heterologous polypeptide fused, in frame, to a polypeptide comprising: 1) SEQ ID NO: 2, 3, or 4; 2) fragments of SEQ ID NO: 2, 3, or 4; or 3) a polypeptide as set forth in Tables 1, 2 or 3; and
 - mixtures of polypeptides as set forth in a), b), c), d), e), f), g), h), or i)
9. The method according to claim 8, further comprising the step of detecting the formation of said antibody-antigen complex.
10. The method according to claim 8, wherein said method is an immunoassay.
11. The method according to claim 10, wherein said immunoassay is selected from the group consisting of enzyme linked immunosorbent assays (ELISAs), radioimmunoassays (RIAs), lateral flow assays, immunochromatographic strip assays, automated flow assays, Western blots, immunoprecipitation assays, reversible flow chromatographic binding assays, agglutination assays, and biosensors.
12. The method according to claim 8, wherein said method is performed using an array of polypeptides.
13. The method according to claim 12, wherein said array of polypeptides comprises the same polypeptide.
14. The method of claim 12, wherein said array of polypeptides comprises the same or different epitopes of the polypeptide of SEQ ID NOs: 2, 3, or 4.
15. The method of claim 12, wherein said array of polypeptides further comprises isolated polypeptides from other organisms of interest.
16. The method of claim 15, wherein said other organisms of interest are selected from the group consisting of *Borrelia burgdorferi*, *Ehrlichia canis*, *Ehrlichia chaffeensis*, *Ehrlichia ruminantium*, *Anaplasma marginale*, and combinations of said organisms of interest.
17. In a method of detecting the presence of antibodies that specifically bind to *Anaplasma phagocytophilum* or antigens thereof, the improvement comprising the use of at least one isolated, recombinant, purified polypeptide comprising:
- SEQ ID NO 2;
 - SEQ ID NO: 3;
 - SEQ ID NO: 4;
 - fragments of SEQ ID NOs: 2, 3, or 4;
 - a polypeptide as set forth in Tables 1, 2 or 3;
 - a variant polypeptide of SEQ ID NO: 2, 3, or 4, wherein said variant polypeptide specifically binds to an antibody that specifically binds to a polypeptide of SEQ ID NO: 2, 3, or 4;
 - a variant of a polypeptide fragment of a polypeptide fragment of SEQ ID NO: 2, 3, or 4, wherein said variant polypeptide fragment specifically binds to an antibody that specifically binds to a polypeptide of SEQ ID NO: 2, 3, or 4, a fragment of SEQ ID NO: 2, 3, or 4, or a polypeptide as set forth in Table 1, 2 or 3;
 - a variant of a polypeptide as set forth in Table 1, 2 or 3, wherein said variant polypeptide specifically binds to an antibody that specifically binds to a polypeptide of SEQ ID NO: 2, 3, or 4 or a polypeptide as set forth in Table 1, 2 or 3;
 - a heterologous polypeptide fused, in frame, to a polypeptide comprising: 1) SEQ ID NO: 2, 3, or 4; 2) fragments of SEQ ID NO: 2, 3, or 4; or 3) a polypeptide as set forth in Table 1, 2 or 3; or
 - a combination of polypeptides as set forth in a), b), c), d), e), f), g), h), or i).
18. An isolated or purified polynucleotide comprising:
- a polynucleotide sequence encoding a polypeptide selected from the group consisting of:
 - SEQ ID NO 2;
 - SEQ ID NO: 3;
 - SEQ ID NO: 4;
 - fragments of SEQ ID NOs: 2, 3, or 4;

- 5) a polypeptide as set forth in Tables 1, 2 or 3;
- 6) a variant polypeptide of SEQ ID NO: 2, 3, or 4, wherein said variant polypeptide specifically binds to an antibody that specifically binds to a polypeptide of SEQ ID NO: 2, 3, or 4;
- 7) a variant polypeptide fragment of SEQ ID NO: 2, 3, or 4, wherein said variant polypeptide fragment specifically binds to an antibody that specifically binds to a polypeptide of SEQ ID NO: 2, 3, or 4 or a fragment of SEQ ID NO: 2, 3, or 4;
- 8) a variant of a polypeptide as set forth in Table 1, 2 or 3, wherein said variant polypeptide specifically binds to an antibody that specifically binds to a polypeptide of SEQ ID NO: 2, 3, or 4 or a polypeptide as set forth in Table 1, 2 or 3;
- 9) a heterologous polypeptide fused, in frame, to a polypeptide comprising: 1) SEQ ID NO: 2, 3, or 4; 2) fragments of SEQ ID NO: 2, 3, or 4; or 3) a polypeptide as set forth in Tables 1, 2 or 3;
- 10) a multimeric construct comprising: a) SEQ ID NO: 2, 3 or 4; b) fragments of SEQ ID NO: 2, 3 or 4; c) combinations of SEQ ID NO: 2, 3, 4, or fragments or variants thereof.
- b) a polynucleotide sequence having at least about 60% to 99.99% identity to a polynucleotide sequence encoding a polypeptide according to (a)(1) through (a)(10);
- c) a polynucleotide sequence comprising SEQ ID NO: 1;
- d) a polynucleotide sequence having at least about 60% to 99.99% identity to the polynucleotide sequence of SEQ ID NO: 1;
- e) a polynucleotide that is complementary to the polynucleotides set forth in (a), (b), (c), or (d);
- f) a genetic construct comprising a polynucleotide sequence as set forth in (a), (b), (c), (d), or (e);
- g) a vector comprising a polynucleotide or genetic construct as set forth in (a), (b), (c), (d), (e), or (f);
- h) a polynucleotide that hybridizes under low, intermediate or high stringency with a polynucleotide sequence as set forth in (a), (b), (c), (d), (e), (f), or (g); or
- i) a probe comprising a polynucleotide according to (a), (b), (c), (d), (e), (f), (g) (h) and, optionally, a label or marker.
19. An antibody that specifically binds to a polypeptide selected from the group consisting of:
- 1) SEQ ID NO 2;
 - 2) SEQ ID NO: 3;
 - 3) SEQ ID NO: 4;
 - 4) fragments of SEQ ID NOs: 2, 3, or 4;
 - 5) a polypeptide as set forth in Table 1, 2 or 3;
 - 6) a variant polypeptide of SEQ ID NO: 2, 3, or 4, wherein said variant polypeptide specifically binds to an antibody that specifically binds to a polypeptide of SEQ ID NO: 2, 3, or 4;
 - 7) a variant polypeptide fragment of SEQ ID NO: 2, 3, or 4, wherein said variant polypeptide fragment specifically binds to an antibody that specifically binds to a polypeptide of SEQ ID NO: 2, 3, or 4 or a fragment of SEQ ID NO: 2, 3, or 4;
 - 8) a variant of a polypeptide as set forth in Table 1, 2 or 3, wherein said variant polypeptide specifically binds to an antibody that specifically binds to a polypeptide of SEQ ID NO: 2, 3, or 4 or a polypeptide as set forth in Table 1, 2 or 3;
 - 9) a heterologous polypeptide fused, in frame, to a polypeptide comprising 1) SEQ ID NO: 2, 3, or 4; 2) fragments of SEQ ID NO: 2, 3, or 4; or 3) a polypeptide as set forth in Table 1, 2 or 3;
- 7) a variant polypeptide fragment of SEQ ID NO: 2, 3, or 4, wherein said variant polypeptide fragment specifically binds to an antibody that specifically binds to a polypeptide of SEQ ID NO: 2, 3, or 4 or a fragment of SEQ ID NO: 2, 3, or 4;
- 8) a variant of a polypeptide as set forth in Table 1, 2 or 3, wherein said variant polypeptide specifically binds to an antibody that specifically binds to a polypeptide of SEQ ID NO: 2, 3, or 4 or a polypeptide as set forth in Table 1, 2 or 3;
- 9) a heterologous polypeptide fused, in frame, to a polypeptide comprising: 1) SEQ ID NO: 2, 3, or 4; 2) fragments of SEQ ID NO: 2, 3, or 4; or 3) a polypeptide as set forth in Tables 1 or 2;
20. The antibody according to claim 19, further comprising an additional component.
21. The antibody according to claim 20, wherein said additional component is a solid support.
22. The antibody according to claim 20, wherein said additional component is a carrier.
23. The antibody according to claim 22, wherein said carrier is a pharmaceutically acceptable excipient.
24. The antibody according to claim 20, wherein said additional component is a label.
25. The antibody according to claim 19, wherein said antibody does not cross-react with an antigen of *Anaplasma marginale*, *Anaplasma centrale*, *Ehrlichia canis*, *Ehrlichia chaffeensis*, or *Cowdria ruminantium*.
26. A host cell comprising a polynucleotide according to claim 18.
27. A method of making a recombinant host cell comprising the introduction of a polynucleotide into a host cell, wherein said polynucleotide comprises:
- a) a polynucleotide sequence encoding a polypeptide selected from the group consisting of:
 - 1) SEQ ID NO 2;
 - 2) SEQ ID NO: 3;
 - 3) SEQ ID NO: 4;
 - 4) fragments of SEQ ID NOs: 2, 3, or 4;
 - 5) a polypeptide as set forth in Tables 1, 2 or 3;
 - 6) a variant polypeptide of SEQ ID NO: 2, 3, or 4, wherein said variant polypeptide specifically binds to an antibody that specifically binds to a polypeptide of SEQ ID NO: 2, 3, or 4;
 - 7) a variant polypeptide fragment of SEQ ID NO: 2, 3, or 4, wherein said variant polypeptide fragment specifically binds to an antibody that specifically binds to a polypeptide of SEQ ID NO: 2, 3, or 4 or a fragment of SEQ ID NO: 2, 3, or 4;
 - 8) a variant of a polypeptide as set forth in Table 1, 2 or 3, wherein said variant polypeptide specifically binds to an antibody that specifically binds to a polypeptide of SEQ ID NO: 2, 3, or 4 or a polypeptide as set forth in Table 1, 2 or 3;
 - 9) a heterologous polypeptide fused, in frame, to a polypeptide comprising 1) SEQ ID NO: 2, 3, or 4; 2) fragments of SEQ ID NO: 2, 3, or 4; or 3) a polypeptide as set forth in Table 1, 2 or 3;

- 10) a multimeric construct comprising: a) SEQ ID NO: 2, 3 or 4; b) fragments of SEQ ID NO: 2, 3 or 4; c) combinations of SEQ ID NO: 2, 3, 4, or fragments or variants thereof.
- b) a polynucleotide sequence having at least about 60% to 99.99% identity to a polynucleotide sequence encoding a polypeptide according to (a)(1) through (a)(10);
- c) a polynucleotide sequence comprising SEQ ID NO: 1;
- d) a polynucleotide sequence having at least about 20% to 99.99% identity to the polynucleotide sequence of SEQ ID NO: 1;
- e) a polynucleotide that is complementary to the polynucleotides set forth in (a), (b), (c), or (d);
- f) a genetic construct comprising a polynucleotide sequence as set forth in (a), (b), (c), (d), or (e);
- g) a vector comprising a polynucleotide or genetic construct as set forth in (a), (b), (c), (d), (e), or (f);
- h) a polynucleotide that hybridizes under low, intermediate or high stringency with a polynucleotide sequence as set forth in (a), (b), (c), (d), (e), (f), or (g).
- 28.** The method according to claim 27, further comprising the step of culturing said recombinant host cell.
- 29.** The method according to claim 27, further comprising the step of culturing said transformed host cell under conditions that allow for the expression of a polypeptide encoded by said polynucleotide.
- 30.** The method according to claim 29, further comprising the step of recovering said expressed polypeptide.
- 31.** In a nucleic acid based assay for the detection of *Anaplasma phagocytophilum*, the improvement comprising the use of a polynucleotide according to claim 18.
- 32.** The method according to claim 31, wherein said polynucleotide does not hybridize with a polynucleotide sequence from *Anaplasma marginale*, *Anaplasma centrale*, *Ehrlichia canis*, *Ehrlichia chaffeensis*, or *Cowdria ruminantium*.
- 33.** A method of hybridizing polynucleotides comprising contacting a sample comprising a population of polynucleotides with a second population of polynucleotides under conditions that allow for the formation of an hybridization complex, wherein said second population of polynucleotides comprises polynucleotides that encode at least one polypeptide that is selected from the group consisting of:
- SEQ ID NO 2;
 - SEQ ID NO: 3;
 - SEQ ID NO: 4;
 - fragments of SEQ ID NOs: 2, 3, or 4;
 - a polypeptide as set forth in Tables 1 or 2;
 - a variant polypeptide of SEQ ID NO: 2, 3, or 4, wherein said variant polypeptide specifically binds to an antibody that specifically binds to a polypeptide of SEQ ID NO: 2, 3, or 4;
 - a variant polypeptide fragment of SEQ ID NO: 2, 3, or 4, wherein said variant polypeptide fragment specifically binds to an antibody that specifically binds to a polypeptide of SEQ ID NO: 2, 3, or 4 or a fragment of SEQ ID NO: 2, 3, or 4;
 - a variant of a polypeptide as set forth in Table 1 or 2, wherein said variant polypeptide specifically binds to an antibody that specifically binds to a polypeptide of SEQ ID NO: 2, 3, or 4 or a polypeptide as set forth in Table 1 or 2;
 - a heterologous polypeptide fused, in frame, to a polypeptide comprising: 1) SEQ ID NO: 2, 3, or 4; 2) fragments of SEQ ID NO: 2, 3, or 4; or 3) a polypeptide as set forth in Tables 1 or 2; and
 - mixtures of polypeptides as set forth in a), b), c), d), e), f), g), h), or i)
- 34.** The method according to claim 32, further comprising the step of detecting the hybridization complex.
- 35.** The method according to claim 32, wherein said second population of polynucleotides is an array of polynucleotides or the same or different sequence.
- 36.** The method according to claim 31, wherein said polynucleotide does not hybridize with a polynucleotide sequence from *Anaplasma marginale*, *Anaplasma centrale*, *Ehrlichia canis*, *Ehrlichia chaffeensis*, or *Cowdria ruminantium*.
- 37.** A method of inducing an immune response comprising the administration of:
- a polypeptide antigen comprising:
 - SEQ ID NO 2;
 - SEQ ID NO: 3
 - SEQ ID NO: 4;
 - fragments of SEQ ID NOs: 2, 3, or 4;
 - a polypeptide as set forth in Table 1, 2 or 3;
 - a variant polypeptide of SEQ ID NO: 2, 3, or 4, wherein said variant polypeptide specifically binds to an antibody that specifically binds to a polypeptide of SEQ ID NO: 2, 3, or 4;
 - a variant polypeptide fragment of SEQ ID NO: 2, 3, or 4, wherein said variant polypeptide fragment specifically binds to an antibody that specifically binds to a polypeptide of SEQ ID NO: 2, 3, or 4 or a fragment of SEQ ID NO: 2, 3, or 4;
 - a variant of a polypeptide as set forth in Table 1, 2 or 3, wherein said variant polypeptide specifically binds to an antibody that specifically binds to a polypeptide of SEQ ID NO: 2, 3, or 4 or a polypeptide as set forth in Table 1, 2 or 3;
 - a heterologous polypeptide fused, in frame, to a polypeptide comprising: 1) SEQ ID NO: 2, 3, or 4; 2) fragments of SEQ ID NO: 2, 3, or 4; or 3) a polypeptide as set forth in Table 1, 2 or 3; and
 - mixtures of polypeptides as set forth in a), b), c), d), e), f), g), h), or i);
 - a polynucleotide encoding at least one polypeptide antigen that is selected from the group consisting of:
 - SEQ ID NO 2;
 - SEQ ID NO: 3;
 - SEQ ID NO: 4;
 - fragments of SEQ ID NOs: 2, 3, or 4;

- e) a polypeptide as set forth in Table 1, 2 or 3;
 - f) a variant polypeptide of SEQ ID NO: 2, 3, or 4, wherein said variant polypeptide specifically binds to an antibody that specifically binds to a polypeptide of SEQ ID NO: 2, 3, or 4;
 - g) a variant polypeptide fragment of SEQ ID NO: 2, 3, or 4, wherein said variant polypeptide fragment specifically binds to an antibody that specifically binds to a polypeptide of SEQ ID NO: 2, 3, or 4 or a fragment of SEQ ID NO: 2, 3, or 4;
 - h) a variant of a polypeptide as set forth in Table 1, 2 or 3, wherein said variant polypeptide specifically binds to an antibody that specifically binds to a polypeptide of SEQ ID NO: 2, 3, or 4 or a polypeptide as set forth in Table 1, 2 or 3;
 - i) a heterologous polypeptide fused, in frame, to a polypeptide comprising: 1) SEQ ID NO: 2, 3, or 4; 2) fragments of SEQ ID NO: 2, 3, or 4; or 3) a polypeptide as set forth in Table 1, 2 or 3; and
 - j) mixtures of polypeptides asset forth in a), b), c), d), e), f), g), h), or i); or
- 3) the administration of at least one of said polypeptide antigens and at least one of said polynucleotides encoding an antigen.

* * * * *

专利名称(译)	无形体嗜吞噬细胞的多核苷酸和多肽及其使用方法		
公开(公告)号	US20050142557A1	公开(公告)日	2005-06-30
申请号	US10/696019	申请日	2003-10-28
[标]申请(专利权)人(译)	佛罗里达大学		
申请(专利权)人(译)	佛罗里达大学		
当前申请(专利权)人(译)	佛罗里达州研究基金会大学		
[标]发明人	ALLEMAN ARTHUR RICK BARBET ANTHONY F		
发明人	ALLEMAN, ARTHUR RICK BARBET, ANTHONY F.		
IPC分类号	A61K38/00 A61K39/00 A61K39/02 C07H21/04 C07K1/00 C07K14/00 C07K14/195 C07K14/29 C07K16/00 C07K16/12 C12N C12N1/20 C12Q1/68 G01N33/53 G01N33/554 G01N33/569		
CPC分类号	A61K39/00 C07K14/29 G01N2469/20 G01N2333/29 C07K16/1246 Y02A50/57		
其他公开文献	US7304139		
外部链接	Espacenet USPTO		

摘要(译)

我们已成功测序并克隆了表达A. phagocytophilum的主要表面蛋白5 (MSP5) 的基因。已经使用来自感染A. phagocytophilum的人和狗的血清测试了重组MSP5 (rMSP5) 蛋白。已发现该多肽具有免疫原性, 可用作诊断试验抗原。本发明的多肽抗原可以提供诊断测定的基础, 该测定允许使用来自感染的样品 (例如, 血清, 血浆或全血) 对A. phagocytophilum感染进行快速的, 内部的实验室诊断。人或动物。另外, 本发明提供了利用本发明提供的抗体检测生物或环境样品中A. phagocytophilum的存在的方法。此外, 在诊断这种重要疾病中使用单一抗原提供了许多优点, 包括增强的测试特异性, 易于测试和使用合成产生的测试抗原而不是培养的整个生物体的结果的一致性。




FIG 1