



US 20060104974A1

(19) **United States**(12) **Patent Application Publication****Davis et al.**(10) **Pub. No.: US 2006/0104974 A1**(43) **Pub. Date: May 18, 2006**(54) **CD147 BINDING MOLECULES AS
THERAPEUTICS**

now abandoned, which is a continuation-in-part of
application No. 09/034,607, filed on Mar. 3, 1998,
now abandoned.

(76) Inventors: **C. Geoffrey Davis**, Burlingame, CA
(US); **Russell W. Blacher**, Castro
Valley, CA (US); **Jose R. Corvalan**,
Foster City, CA (US); **Alan R. Culwell**,
Carlsbad, CA (US); **Larry L. Green**,
San Francisco, CA (US); **Joanne Hales**,
Fremont, CA (US); **Nancy Havrilla**,
Oakland, CA (US); **Vladimir E.**
Ivanov, Fremont, CA (US); **John A.**
Lipani, Livermore, CA (US); **Qiang**
Liu, Foster City, CA (US); **Richard F.**
Weber, San Francisco, CA (US);
Xiao-Dong Yang, Palo Alto, CA (US)

Correspondence Address:
FISH & NEAVE IP GROUP
ROPES & GRAY LLP
1251 AVENUE OF THE AMERICAS FL C3
NEW YORK, NY 10020-1105 (US)

(21) Appl. No.: **09/784,950**(22) Filed: **Feb. 15, 2001****Related U.S. Application Data**

(63) Continuation of application No. PCT/US99/04583,
filed on Mar. 3, 1999, which is a continuation-in-part
of application No. 09/244,253, filed on Feb. 3, 1999,

Publication Classification(51) **Int. Cl.****A61K 39/395** (2006.01)**C07K 16/28** (2006.01)(52) **U.S. Cl.** **424/144.1; 530/388.22**

(57)

ABSTRACT

In accordance with the present invention, we have discov-
ered that the molecule CD147 as expressed on certain cells,
such as T-cells, B-cells, and/or monocytes, can be utilized
for the treatment of a variety of diseases. In particular, we
have demonstrated that antibodies that bind to CD147 and
that result in the killing of such cells, for example, through
the binding of complement, is efficacious in the treatment of
diseases. Diseases in which such treatment appears effica-
cious include, without limitation: graft versus host disease
(GVHD), organ transplant rejection diseases (including,
without limitation, renal transplant, ocular transplant, and
others), cancers (including, without limitation, cancers of
the blood (i.e., leukemias and lymphomas), pancreatic, and
others), autoimmune diseases, inflammatory diseases, and
others.




FIG.1





FIG.2B




FIG.3




FIG.4




FIG.5




FIG. 6B








FIG. 6A







FIG. 8C

FIG. 8B

FIG. 8A






FIG. 9B




FIG. 9D




FIG. 9A




FIG. 9C




FIG. 10A




FIG. 10C




FIG. 10E




FIG. 10B




FIG. 10D




FIG. 10F




FIG. 11A




FIG. 11C




FIG. 11E




FIG. 11B




FIG. 11D




FIG. 11F

CELL TYPE	SURFACE MARKERS	CDC DEPLETION
RESTING T CELLS ACTIVATED T CELLS	CD3 ⁺ CD25 ⁻ CD3 ⁺ (CD4 ⁺ /CD8 ⁺)CD25 ⁺	NO YES
RESTING B CELLS ACTIVATED B CELLS	CD20 ⁺ CD25 ⁻ CD20 ⁺ CD25 ⁺	NO YES
RESTING MONOCYTES ACTIVATED MONOCYTES	CD14 ⁺ CD25 ⁻ CD14 ⁺ CD25 ⁺	YES YES

FIG.12

CELL	CELL TYPE	CBL Ag EXPRESSION	CBL CDC
CEM	T CELL	++	+
JURKAT	T CELL	++	-
U937	MONOCYTE	++	+
A431	EPIDERMAL	++	-
SW948	COLON	+++	-
MDA468	BREAST	+	-

FIG.13

CELL	CELL TYPE	CBL EXPRESSION	CBL CDC	CD55	CD59
CEM	T CELL	++	+	-	+
JURKAT	T CELL	++	-	+	+
U937	MONOCYTE	++	+	+	-
A431	EPIDERMAL	++	-	+	+
SW948	COLON	+++	-	+	+
MDA468	BREAST	+	-	+	+

FIG.14




FIG. 15A




FIG. 15B




FIG. 15C




FIG. 16A




FIG. 16B




FIG. 16C




FIG. 17




FIG. 18




FIG. 19A




FIG. 19B




FIG. 19C




FIG. 20A




FIG. 20B




FIG. 20C




FIG.21




FIG.22




FIG.23A

E. Coli




FIG. 23B

IgM Antibody Sequences

CEM 10.1 C3 Heavy cDNA

GGACTGTTGA	AGCCTTCGGA	GACCCTGTCC	CTCACCTGCG	CTGTCTATGG	50
TGGGTCCTTC	AGTGGTTACT	ACTGGAGCTG	GATCCGCCAG	CCCCCAGGGA	100
AGGGGCTGGA	GTGGATTGGG	GAAATCAATC	ATAGTGGAAG	CACCAACTAC	150
AACCCGTCCC	TCAAGAGTCG	AGTCACCATA	TCAGTAGACA	CGTCCAAGAA	200
CCAGTTCTCC	CTGAAGCTGA	GCTCTGTGAC	CGCNGCGGAC	ACGGCTGTGT	250
ATTACTGTGC	GAGAGGCACT	ACGGAATATT	ACTACTACTA	CTACGGTATG	300
GACGTCTGGG	GCCAAGGGAC	CACGGTCACC	GTCTCCTCAG	GGAGTGCATC	350
CGCCCCAACC	CTTTTCCCCC	TCGTCTCCTG	TGAGAATTCC	CCGTCGGATA	400
CGAGCAGCGT	GGCCGTTGGC	TGCCTCGCAC	AGGACTTCCT	TCCCGACTYC	450
ATCACTTTCT	CCTGGAAATA	CAAGAACAAC	TCTGACATCA	GCAGCACCCG	500
GGGCTTCCCA	TCAGTCCTGA	GAGGGGGCAA	GTACGCAGCC	ACCTCACAGG	550
TGCTGCTGCC	TTCCAAGGAC	GTCATGCAGG	GCACAGACGA	ACACGTGGTG	600
ACGGGATCCA	AAGAGTA				617

(SEQ ID NO:62)

CEM 10.1 C3 Heavy Protein

GLLKPSETLS	LTCVYGGSF	SGYYWSWIRQ	PPGKGLEWIG	EINHSGSTNY	50
NPSLKSRTVI	SVDTSKNQFS	LKLSSVTAAD	TAVYYCARGT	TEYYYYYYGM	100
DVWQGTTVT	VSSGSASAPT	LFPLVSCENS	PSDTSSVAVG	CLAQDFLPDX	150
ITFSWKYKNN	SDISSTRGFP	SVLRGGKYAA	TSQVLLPSKD	VMQGTDEHVV	200
TGSKE					205

(SEQ ID NO:23)

CEM 10.1 C3 Kappa cDNA

CTCTCCCTGC	CCGTCACCCC	TGGAGAGCCG	GCCTCCATCT	CCTGCAGGTC	50
TAGTCAGAGC	CTCCTGCATA	GTAATGGATA	CAACTATTTG	GATTGGTACC	100
TGCAGAAGCC	AGGGCAGTCT	CCACAGCTCC	TGATCTATTT	GGGTTCTAAT	150
CGGGCCTCCG	GGGTCCCTGA	CAGGTTCACT	GGCAGTGGAT	CAGGCACAGA	200
TTTTACACTG	AAAATCAGCA	GAGTGGAGGC	TGAGGATGTT	GGGATTTATT	250
ACTGCATGCA	GACTCGACAA	ACTCCTCGGA	CGTTCGGCCA	AGGGACCAAG	300
GTGGAAATCA	AACGAACTGT	GGCTGCACCA	TCTGTCTTCA	TCTTCCCGCC	350
ATCTGATGAG	CAGTTGAAAT	CTGGAAGTGC	CTCTGTTGTG	TGCCTGCTGA	400
ATAACTTCTA	TCCCAGAGAG	GCCAAAGAGC	ATCAAAAGAG	TCCA	444

(SEQ ID NO:63)

CEM 10.1 C3 Kappa Protein

LSPVTPGEP	ASISCRSSQS	LLHSNGYNYL	DWYLQKPGQS	PQLLIYLGSN	50
RASGVPDFRS	GSGSGDFTL	KISRVEADV	GIYYCMQTRQ	TPRTFGQGTK	100
VEIKRTVAAP	SVFIFPPSDE	QLKSGTASVV	CLLNNFYPRE	AKEHQKSP	148

(SEQ ID NO:24)

FIG. 24

IgM Antibody Sequences

CEM 10.1 G10 Heavy cDNA

CTGGTGAAGC	CTTCGGAGAC	CCTGTCCCTC	ACCTGCACTG	TCTCTGGTGG	50
CTCCATCAGT	AGTTACTACT	GGAAGTGGAT	CCGGCAGCCC	CCAGGGAAGG	100
GACTGGAGTG	GATTGGGTAT	ATCTATTACA	GTGGGAGCAC	CAACTACAAC	150
CCCTCCCTCA	AGAGTCGAGT	CACCATATCA	GTAGACACGT	CCAAGAACCA	200
GTTCTCCCTG	AAGCTGAGCT	CTGTGACCGC	TGCGGACACG	GCCGTGTATT	250
ACTGTGCGAG	AGATAGGGGA	GTGGGAGCTA	CTGGTTTTGA	CTACTGGGGC	300
CAGGGAACCC	TGGTCACCGT	CTCCTCAGGG	AGTGCATCCG	CCCCAACCCCT	350
TTTCCCCCTC	GTCTCCTGTG	AGAATTCCCC	GTCCGATACG	AGCAGCGTGG	400
CCGTTGGCTG	CCTCGCACAG	GACTTCCTTC	CCGACTCCAT	CACTTTCTCC	450
TGGAAATACA	AGAACAACCTC	TGACATCAGC	AGCACCCGGG	GCTTCCCATC	500
AGTCCTGAGA	GGGGGCAAGT	ACGCAGCCAC	CTCACAGGTG	CTGCTGCCTT	550
CCAAGGACGT	CATGCAGGGC	ACAGACGAAC	ACAAGGTGTG	CGA	593

(SEQ ID NO:64)

CEM 10.1 G10 Heavy Protein

LVKPSSETLSL	TCTVSGGSIS	SYYNWIRQP	PGKLEWIGY	IYYSGSTNYN	50
PSLKSRVTIS	VDTSKNQFSL	KLSSVTAADT	AVYYCARDRG	VGATGFDYWG	100
QGTLLVTVSSG	SASAPTLFPL	VSCENSPSDT	SSVAVGCLAQ	DFLPDSITFS	150
WKYKNNSDIS	STRGFPSVLR	GGKYAATSQV	LLPSKDVMQG	TDEHKVC	197

(SEQ ID NO:25)

CEM 10.1 G10 Kappa cDNA

AGCCAGTCTC	CATCCTCCCT	GTCTGCATCT	GTAGGAGAGA	GAGTCACCAT	50
CACTTGCCGG	GCAAGTCAGG	GCATTAGAGA	TGAATTAGGC	TGGTATCAGC	100
AGAAACCAGG	GAAAGCCCCCT	AAGCGCCTGA	TCTATGTTGC	ATCCAGTTTG	150
CAAAGTGGGG	TCCCATCAAG	GTTTCAGCGGC	AGTGGATCTG	GGACAGAATT	200
CACTCTCACA	ATCAGCAGCC	TGCAGCCTGA	AGATTTTGCA	ACTTATTACT	250
GTCTACAGCA	TAATGGTTAC	CCTCGGACGT	TCGGCCAAGG	GACCAAGGTG	300
GAAATCAAAC	GAAGTGTGGC	TGCACCATCT	GTCTTCATCT	TCCCGCCATC	350
TGATGAGCAG	TTGAAATCTG	GAAGTGCCTC	TGTTGTGTGC	CTGCTGAATA	400
ACTTCTATCC	CAGAGAGGCC	AAAGAGCATC	AAAAGAGTCC	A	441

(SEQ ID NO:65)

CEM 10.1 G10 Kappa Protein

SQSPSSLSAS	VGERVTITCR	ASQGIRDELG	WYQQKPGKAP	KRLIYVASSL	50
QSGVPSRFSG	SGSGTEFTLT	ISSLPEDFA	TYYCLQHNGY	PRTEGQGTKV	100
EIKRTVAAPS	VFIFPPSDEQ	LKSGTASVVC	LLNNFYPREA	KEHQKSP	147

(SEQ ID NO:26)

FIG. 25

IgM Antibody Sequences

CEM 10.12 F3 Heavy cDNA

AAGAAGCCTG	GGGCCTCAGT	GAAGGTCTCC	TGCAAGGCTT	CTGGATACAC	50
CTTCACCACT	TATGATATCA	ACTGGGTGCG	ACAGGCCACT	GGACAAGGGC	100
TTGAGTGGAT	GGGATGGATG	AACCCTAACA	GTGGTAACAC	AGGCTATGCA	150
CAGAAGTTCC	AGGGCAGAGT	CACCATGAAC	AGGAACACCT	CCATAAGCAC	200
AGCCTACATG	GAGCTGAGCA	GCCTGAGATC	TGAGGACACG	GCCGTGTATT	250
ACTGTGCGAG	AGGGGGTCAT	GGTGGGAGCT	ACTTCTACTC	CTAYTACGGT	300
ATGGACGTCT	GGGGCCAGGG	GACCACGGTC	ACCGTCTCCT	CAGGGAGTGC	350
ATCCGCCCCA	ACCCTTTTCC	CCCTCGTCTC	CTGTGAGAAAT	TCCCCGTCGG	400
ATACGAGCAG	CGTGGCCGTT	GGCTGCCTCG	CACAGGACTT	CCTTCCCGAC	450
TCCATCACTT	TCTCCTGGAA	ATACAAGAAC	AACTCTGACA	TCAGCAGCAC	500
CCGGGGCTTC	CCATCAGTCC	TGAGAGGGGG	CAAGTACGCA	GCCACCTCAC	550
AGGTGCTGCT	GCCTTCCAAG	GACGTCATGC	AGGGCACAGA	CGAACACGTG	600
GTGTGCAAAC					610

(SEQ ID NO:66)

CEM 10.12 F3 Heavy Protein

KKPGASVKVS	CKASGYTFTS	YDINWVRQAT	GQGLEWMGWM	NPNSGNTGYA	50
QKFGQGRVTM	RNTSISTAYM	ELSSLRSED	AVYYCARGGH	GGSYFYFYYG	100
MDVWQGQTTV	TVSSGSASAP	TLFPLVSCEN	SPSDTSSVAV	GCLAQDFLPD	150
SITFSWKYKN	NSDISSTRGF	PSVLRGGKYA	ATSQVLLPSK	DVMQGTDEHV	200
VCK					203

(SEQ ID NO:27)

CEM 10.12 F3 Kappa cDNA

CACTCCCTGG	CTGTGTCTCT	GGGCGAGAGG	GCCACCATCA	ACTGCAAGTC	50
CAGCCAGAGT	GTTTTATACA	GTTTTAACA	TAAGAACTAC	TTAGCTTGGT	100
ACCAGCAGAA	ACCAGGACAG	CCTCCTAAGC	TGCTCATTTA	CTGGGCATCT	150
ACCCGGGAAT	CCGGGGTCCC	TGACCGATT	GGTGGCAGCG	GGTCTGGGAC	200
AGATTTCACT	CTCACCATCA	GCAGCCTGCA	GGCTGAAGAT	GTGGCAGTTT	250
ATTACTGTCA	GCAATATTAT	AGTACTCCTM	GGACGTTCCG	CCAAGGGACC	300
AAGGTGGAAA	TCAAACGAAC	TGTGGCTGCA	CCATCTGTCT	TCATCTTCCC	350
GCCATCTGAT	GAGCAGTTGA	AATCTGGAAC	TGCCTCTGTT	GTGTGCCTGC	400
TGAATAACTT	CTATCCCAGA	GAGGCCAAAG	AGCATCAAAA	GAGTCCA	447

(SEQ ID NO:67)

CEM 10.12 F3 Kappa Protein

HSLAVSLGER	ATINCKSSQS	VLYSFNNKNY	LAWYQQKPGQ	PPKLLIYWAS	50
TRESGVPDRF	GGSGSGTDFT	LTISLQAED	VAVYYCQQYY	STPRTFGQGT	100
KVEIKRTVAA	PSVFIFPPSD	EQLKSGTASV	VCLLNIFYPR	EAKEHQKSP	149

(SEQ ID NO:28)

FIG. 26

IgM Antibody Sequences

CEM 10.12 G5 Heavy

GAGGTGAAGA	AGCCTGGGGC	CTCAGTGAAG	GTCTCCTGCA	AGGCTTCTGG	50
ATACACCTTC	ACCAGTTATG	ATATCAACTG	GGTGCGACAG	GCCACTGGAC	100
AAGGGCTTGA	GTGGATGGGA	TGGATGAACC	CTAACAGTGG	TAACACAGGC	150
TATGCACAGA	AGTTCCAGGG	CAGAGTCACC	ATGACCAGGA	ACACCTCCAT	200
AAGCACAGCC	TACATGGAGC	TGAGCAGCCT	GAGATCTGAG	GACACGGCCG	250
TGTATTACTG	TGCGAGAGAG	GAGTGGCTGG	TACGTTACTA	CGGTATGGAC	300
GTCTGGGGCC	AAGGGACCAC	GGTCACCGTC	TCCTCAGGGA	GTGCATCCGC	350
CCCAACCCCTT	TTCCCCCTCG	TCTCCTGTGA	GAATTCCCCG	TCGGATACGA	400
GCAGCGTGGC	CGTTGGCTGC	CTCGCACAGG	ACTTCCTTCC	CGACTCCATC	450
ACTTTCTCCT	GGAAATACAA	GAACAACTCT	GACATCAGCA	GCACCCGGGG	500
CTTCCCATCA	GTCCTGAGAG	GGGGCAAGTA	CGCAGCCACC	TCACAGGTGC	550
TGCTGCCTTC	CAAGGACGTC	ATGCAGGGCA	CAGACGAACA	CAAGGTGTG	599

(SEQ ID NO:68)

CEM 10.12 G5 Heavy Protein

EVKKPGASVK	VSKASGYTF	TSYDINWVRQ	ATQGQLEWMG	WMNPNSGNTG	50
YAQKFQGRVT	MTRNTSISTA	YMELSSLRSE	DTAVYYCARE	EWLVRYYGMD	100
VWQGQTTVTV	SSGSASAPTL	FPLVSCENSP	SDTSSVAVGC	LAQDFLPDSI	150
TFSWKYKNNS	DISSTRGFPS	VLRGGKYAAT	SQVLLPSKDV	MQGTDEHKV	199

(SEQ ID NO:29)

CEM 10.12 G5 Kappa cDna

GGCCAGTCTC	CATCCTCCCT	GTCTGCATCT	GTAGGAGACA	GAGTCACCAT	50
CACTTGCCCG	GCAAGTCAGG	ACATTAGAGA	TAATTTAGGC	TGGTATCAGC	100
AGAAACCAGG	GAAAGCCCCT	AAGCGCCTGA	TCTATGCTGC	ATCCAATTTG	150
CAAAGTGGGG	TCCCATCAAG	GTTTCAGCGG	AGTGGATCTG	GGACAGAATT	200
CACTCTCACA	ATCAGCAGCC	TGCAGCCTGA	AGATTTTGCA	ACTTATTACT	250
GTCTACAGTA	TAAAACTTAC	CCGTGGACGT	TCGGCCAAGG	GACCAAGGTG	300
GAAATCAAAC	GAAGTGTGGC	TGCACCATCT	GTCTTCATCT	TCCCGCCATC	350
TGATGAGCAG	TTGAAATCTG	GAAGTGCCTC	TGTTGTGTGC	CTGCTGAATA	400
ACTTCTATCC	CAGAGAGGMC	AAAGAGCATC	AAAAGAGTCC	A	441

(SEQ ID NO:69)

CEM 10.12 G5 Kappa Protein

GQSPSSLSAS	VGDRVITICR	ASQDIRDNLG	WYQKPGKAP	KRLIYAASNL	50
QSGVPSRFSG	SGSGTEFTLT	ISSLPEDFA	TYYCLQYKTY	PWTFGQGTKV	100
EIKRTVAAPS	VFIFPPSDEQ	LKSGTASVVC	LLNNFYPREX	KEHQKSP	147

(SEQ ID NO:30)

FIG. 27

IgM Antibody Sequences

CEM 13.12 Heavy cDNA

AAGCTTCCGG	AGACCTGTC	CCTCACCTGC	GCTGTCTATG	GTGGGTCCTT	50
CAGTGGTTAC	TACTGGAGCT	GGATCCGCCA	GCCCCAGGG	AAGGGGCTGG	100
AGTGGATTGG	GGAAATCAAT	CATAGTGGAA	GCACCAACTA	CAACCCGTCC	150
CTCAAGAGTC	GAGTCACCAT	ATCAGTAGAC	ACGTCCAAGA	ACCAGTTCTC	200
CCTGAAGCTG	AGCTCTGTGA	CGCCGCGGA	CACGGCTGTG	TATTACTGTG	250
CGAGAGGGGC	AGCTGAATAT	TACTACTACT	ACTACGGTAT	GGACGTCTGG	300
GGCCAAGGGA	CCACGGTCAC	CGTCTCCTCA	GGGAGTGCAT	CCGCCCCAAC	350
CCTTTTCCCC	CTCGTCTCCT	GTGAGAATTC	CCCGTCGGAT	ACGAGCAGCG	400
TGGCCGTTGG	CTGCCTCGCA	CAGGACTTCC	TTCCCGACTY	CATCACTTTC	450
TYCTGGAAT	ACAAGAACAA	CTCTGACATC	AGCAGCACCC	GGGGCTTCCC	500
ATCAGTCCTG	AGAGGGGGCA	AGTACGCAGC	CACCTCACAG	GTGCTGCTGC	550
CTTCCAAGGA	CGTCATGCAG	GGCACAGACG	AACACGTGGT	GACGGGATCC	600
AAAGAGT					607

(SEQ ID NO:70)

CEM 13.12 Heavy Protein

KLPETLSLTC	AVYGGSFSGY	YWSWIRQPPG	KGLEWIGEIN	HSGSTNYNPS	50
LKSRVTISVD	TSKNQFSLKL	SSVTAADTAV	YYCARGAAEY	YYYYYGMDVW	100
GQGTTVTVSS	GSASAPTLFP	LVSCENSPSD	TSSVAVGCLA	QDFLPDXITF	150
XWKYKNNSDI	SSTRGFPSVL	RGGKYAATSQ	VLLPSKDVMQ	GTDEHVVTS	200
KE					202

(SEQ ID NO:31)

CEM 13.12 Kappa cDNA

ATGCCCCGTCA	CCCCTGGAGA	GCCGGCCTCC	ATCTCCTGCA	GGTCTAGTCA	50
GAGCCTCCTG	CATAGTAATG	GATACAATA	TTTGGACTGG	TACCTGCAGA	100
AGCCAGGGCA	GTCTCCACAG	CTCCTGATCT	ATTTGGGTTC	TAATCGGGCC	150
TCCGGGGTCC	CTGACAGGTT	CAGTGGCAGT	GGATCAGGCA	CAGATTTTAC	200
ACTGAAAATC	AGCAGAGTGG	AGGCTGAGGA	TGTTGGGATT	TATTACTGCA	250
TGCAAAGTCT	ACAAATTCCC	CGGCTTTTTCG	GCCCTGGGAC	CAAAGTGGAT	300
ATCAAACGAA	CTGTGGCTGC	ACCATCTGTC	TTCATCTTCC	CGCCATCTGA	350
TGAGCAGTTG	AAATCTGGAA	CTGCCTCTGT	TGTGTGCCTG	CTGAGTAACT	400
TCTATCCAG	AGAGGCCAAA	GTACAGTGGA	A		431

(SEQ ID NO:71)

CEM 13.12 Kappa Protein

MPVTPGEPAS	ISCRSSQSLL	HSNGYNYLDW	YLQKPGQSPQ	LLIYLGSNRA	50
SGVPDRFSGS	SGGTDFTLKI	SRVEAEDVGI	YYCMQSLQIP	RLFGPGTKVD	100
IKRTVAAPSV	FIFPPSDEQL	KSGTASVVCL	LSNFYPREAK	VQW	143

(SEQ ID NO:32)

FIG. 28

IgM Antibody Sequences

CEM 13.5 Heavy cDNA

TCGGAGACCC	TGTCCCTCAC	CTGCGCTGTC	TATGGTGGGT	CCTTCAGTGG	50
TTACTACTGG	AGCTGGATCC	GCCAGCCCCC	AGGGAAGGGG	CTGGAGTGGA	100
TTGGGGAAAT	CAATCATAGT	GGAAGCACCA	ACTACAACCC	GTCCCTCAAG	150
AGTCGAGTCA	CCATATCAGT	AGACACGTCC	AAGAACCAGT	TCTCCCTGAA	200
GCTGAGTTCT	GTGACCGCCG	CGGACACGGC	TGTGTATTAC	TGTGCGAGAG	250
GCGGGACTAC	AGTAACTTTT	GATGCTTTTG	ATATCTGGGG	CCAAGGGACA	300
ATGGTCACCG	TCTCTTCAGG	GAGTGCATCC	GCCCCAACCC	TTTTCCCCCT	350
CGTCTCCTGT	GAGAATTCCC	CGTCGGATAC	GAGCAGCGTG	GCCGTTGGCT	400
GCCTCGCACA	GGACTTCCTT	CCCGACTCCA	TCACTTTCTC	CTGGAAATAC	450
AAGAACAAC	CTGACATCAG	CAGCACCCGG	GGCTTCCCAT	CAGTCCTGAG	500
AGGGGGCAAG	TACGCAGCCA	CCTCACAGGT	GCTGCTGCCT	TCCAAGGACG	550
TCATGCAGGG	CACAGACGAA				570

(SEQ ID NO:72)

CEM 13.5 Heavy Protein

SETLSLTCAV	YGGSFSGYYW	SWIRQPPGKG	LEWIGEINHS	GSTNYPNPSLK	50
SRVTISVDTS	KNQFSLKLSS	VTAADTAVYY	CARGGTTVTF	DAFDIWQOGT	100
MVTVSSGSAS	APTLFPLVSC	ENSPSDTSSV	AVGCLAQDFL	PDSITFSWKY	150
KNNSDISSTR	GFPSVLRGGK	YAATSQVLLP	SKDVMQGTDE		190

(SEQ ID NO:33)

CEM [13.5] Kappa cDNA

CTGGCTGTGT	CTCTGGGCGA	GAGGGCCACC	ATCAACTGCA	AGTCCAGCCA	50
GAGTGTTTTA	TACAGTTTTA	ACAATAAGAA	CTACTTAGCT	TGGTACCAGC	100
AGAAACCAGG	ACAGCCTCCT	AAGCTGCTCA	TTTACTGGGC	ATCTACCCGG	150
GAATCCGGGG	TCCCTGACCG	ATTCACTGGC	AGCGGGTCTG	GGACAGATTT	200
CACTCTCACC	ATCAGCAGCC	TGCAGGCTGA	AGATGTGGCA	GTTTATTACT	250
GTCAGCAATA	TTATAGTACT	CCTCGGACGT	TCGGCCAAGG	GACCAAGGTG	300
GAAATCAAAC	GAAGTGTGGC	TGCACCATCT	GTCTTCATCT	TCCCGCCATC	350
TGATGAGCAG	TTGAAATCTG	GAAGTGCCTC	TGTTGTGTGC	CTGCTGAATA	400
ACTTCTATCC	CAGAGAGGCC	AAAGTACAGT	GGAAGGTGAT	C	441

(SEQ ID NO:73)

CEM 13.5 Kappa Protein

LAVSLGERAT	INCKSSQSVL	YSFNNKNYLA	WYQQKPGQPP	KLLIYWASTR	50
ESGVPDRFSG	SGSGTDFTLT	ISSLQAEDVA	VYYCQYYST	PRTFGQGTKV	100
EIKRTVAAPS	VFIFPPSDEQ	LKSGTASVVC	LLNNFYPREA	KVQWKVI	147

(SEQ ID NO:34)

FIG. 29

IgG Antibody Sequences

2.4.4 Heavy cDNA

AACCCACAGA	CGACCCTCAC	GCTGACCTGC	ACCTTCTCTG	GGTTCTCACT	50
CATTACCCGT	GGAGTGGGTG	TGGATTGGAT	CCGTCAGCCC	CCAGGAAAGG	100
CCCTGCAGTG	GCTCGCACTC	ATTTATTGGA	ATGATGATAA	GCGCTACAGT	150
CCATCTCTGA	AGAGCAGGCT	CACCATCACC	AAGGACACCT	CCAAAAACCA	200
GGTGGTCCTC	ACAATGACCA	ACATGGACCC	TGTGGACACA	GCCACATATT	250
ACTGTGCACA	CCATTTCTTT	GATAGTAGTG	GTTATTACCC	TTTTGACTCC	300
TGGGGCCAGG	GAACCCTGGT	CTCCGTCTCC	TCAGCCTCCA	CCAAGGGCCC	350
ATCGGTCTTC	CCCCTGGCGC	CCTGCTCCAG	GAGCACCTCC	GAGAGCACAG	400
CGGCCCTGGG	CTGCCTGGTC	AAGGACTACT	TCCCCGAACC	GGTGACG	447

(SEQ ID NO:74)

2.4.4 Heavy Protein

NPQTLTLTC	TFSGFSLITR	GVGVDWIRQP	PGKALQWLAL	IYWNDDKRY	50
PSLKSRLTIT	KDTSKNQVVL	TMTNMDPVD	ATYYCAHFF	DSSGYYPFDS	100
WGQGLVSVS	SASTKGPSVF	PLAPCSRSTS	ESTAALGCLV	KDYFPEPVT	149

(SEQ ID NO:35)

2.4.4 Kappa cDNA

GTGACTCAGT	CTCCACTCTC	TCTGTCCGTC	ACCCCTGGAC	AGCCGGCCTC	50
CATCTCCTGC	AAGTCTAGTC	AGAGCCTCCT	GCATAGTGAT	GGAAAGACCT	100
ATTTGTATTG	GTACCTGCAG	AAGCCAGGCC	AGCCTCCACA	GCTCCTGATC	150
TATGAAGCTT	TCAACCGGTT	CTCTGGAGTG	CCAGATAGGT	TCAGTGGCAG	200
CGGGTCAGGG	ACAGATTTC	CACTGAAAAT	CAGCCGGGTG	GAGGCTGAGG	250
ATGTTGGACT	TTATTATTGC	ATGCAAAGTA	TAGAGCTTCC	GTTCACTTTC	300
GGCGGAGGGA	CCAAGGTGGA	GATCAAACGA	ACTGTGGCTG	CACCATCTGT	350
CTTCATCTTC	CCGCCATCTG	ATGAGCAGTT	GAAATCTGGA	ACTGCCTCTG	400
TTGTGTGCCT	GCTGAATAAC	TTCTATCCCA	GAAAAGAAAG	AGTCR	445

(SEQ ID NO:75)

2.4.4 Kappa Protein

VTQSPLSLSV	TPGQPASISC	KSSQSLHSD	GKTYLYWYLQ	KPGQPPQLLI	50
YEA FNRFSGV	PDRFSGSGSG	TDFTLKISR	EAEDVGLYYC	MQSIELPF	100
GGG TKVEIKR	TVAAPSVFIF	PPSDEQLKSG	TASVVCLLNN	FYPRKERV	148

(SEQ ID NO:36)

FIG. 30

IgG Antibody Sequences

2.1.1 Heavy cDNA

GGGGAAGGCC	TGGTCAAGCC	TGGGGGGTCC	CTGAGACTCT	CCTGTGCAGC	50
CTCTGGATTG	ACCTTCAGTA	GCTATAGCAT	GAAGTGGGTC	CGCCAGGCTC	100
CAGGGAAGGG	GCTGGAGTGG	GTCTCATCCA	TTAGTAGTAG	TAGTAGTTAC	150
ATATACTACG	CAGACTCAGT	GAAGGGCCGA	TTCAACCATCT	CCAGAGACAA	200
CGCCAAGAAC	TCACTGTATC	TGCAAATGAA	CAGCCTGAGA	GCCGAGGACA	250
CGGCTGTGTA	TTACTGTGCG	AGGGATAGCA	GTGGCTGGTA	TGAGGACTAC	300
TTTGACTACT	GGGGCCAGGG	AACCTGGTTC	ACCGTCTCCT	CAGCCTCCAC	350
CAAGGGCCCCA	TCGGTCTTCC	CCCTGGCGCC	CTGCTCCAGG	AGCACCTCCG	400
AGAGCACAGC	GGCCCTGGGC	TGCCTGGTCA	AGGACTACTT	CCCCGAACCG	450
GTGACGGTGT	CGTGGAAGTC	AGGCGCTCTG	ACCAGCGGCG	TGCACACCTT	500
CCCAGCTGTC	CTACAGTCA				519

(SEQ ID NO:76)

2.1.1 Heavy Protein

GEGLVKPGGS	LRLSCAASGF	TFSSYSMNWV	RQAPGKGLEW	VSSISSSSSY	50
IYYADSVKGR	FTISRDNKN	SLYLQMNLSR	AEDTAVYYCA	RDSSGWYEDY	100
FDYWGGQTLV	TVSSASTKGP	SVFPLAPCSR	STSESTAALG	CLVKDYFPEP	150
VTVSWNSGAL	TSGVHTFPAV	LQS			173

(SEQ ID NO:37)

2.1.1 Kappa cDNA

CTTGACATCC	AGCTGACCCA	GTCTCCGTCC	TCACTGTCTG	CATCTGTAGG	50
AGACAGAGTC	ACCATCACTT	GTGCGGCGAG	TCAGGACATT	AGCATTATT	100
TAGCCTGGTT	TCAGCAGAGA	CCAGGGAAAG	CCCCTAAGTC	CCTGATCTAT	150
GCTGCATCCA	GTTTGCAAAG	TGGGGTCCCA	TCAAAGTTCA	GCGGCAGTGG	200
ATCTGGGACA	GATTTCACTC	TCACCATCAG	CAGCCTGCAG	CCTGAAGATT	250
TTGCAACTTA	TTACTGCCAA	CAATATAATA	GTTATCCATT	CACCTTTCGGG	300
CCC					303

(SEQ ID NO:77)

2.1.1 Kappa Protein

LDIQLTQSPS	SLSASVGDRV	TITCRASQDI	SIYLAWFQQR	PGKAPKSLIY	50
AASSLQSGVP	SKFSGSGSGT	DFTLTISLQ	PEDFATYYCQ	QYNSYPFTFG	100
P					101

(SEQ ID NO:38)

FIG. 31

IgG Antibody Sequences

2.3.2 Heavy cDNA

CTGACCTGCA	CCTTCTCTGG	GTTCTCACTC	ATTACCCGTG	GAGTGGGTGT	50
GGATTGGATC	CGTCAGCCCC	CAGGAAAGGC	CCTGCAGTGG	CTCGCACTCA	100
TTTATTGGAA	TGATGATAAG	CGCTACAGTC	CATCTCTGAA	GAGCAGGCTC	150
ACCATCACCA	AGGACACCTC	CAAAAACCAG	GTGGTCCTCA	CAATGACCAA	200
CATGGACCCT	GTGGACACAG	CCACATATTA	CTGTGCACAC	CATTTCTTTG	250
ATAGTAGTGG	TTATTACCCT	TTTGACTCCT	GGGGCCAGGG	AACCCTGGTC	300
TCCGTCTCCT	CAGCCTCCAC	CAAGGGCCCA	TCGGTCTTCC	CCCTGGCGCC	350
CTGCTCCAGG	AGCACCTCCG	AGAGCACAGC	GGCCCTGGGC	TGCCTGGTCA	400
AGGACTACTT	CCCCGAACCG	GTGACGGTGT	CGTGGAACCT	AGGCGCTCTG	450
ACCAGCGGCG	TGCACACCTT	CCAGCTG			477

(SEQ ID NO:78)

2.3.2 Heavy Protein

LTCTFSGFSL	ITRGVGVDWI	RQPPGKALQW	LALIYWDDK	RYSPSLKSRL	50
TITKDTSKNQ	VVLTMNMDP	VDATYYCAH	HFFDSSGYYP	FDSWGQGLV	100
SVSSASTKGP	SVFPLAPCSR	STSESTAALG	CLVKDYFPEP	VTVSWNSGAL	150
TSGVHTFQL					159

(SEQ ID NO:39)

FIG. 32

IgG Antibody Sequences

2.6.1 Heavy cDNA

GGGGGAGGCT	TGGTACAGCC	TGGGGGGTCC	CTGAGACTCT	CCTGTGCAGC	50
CTCTGGATTC	ACTTTTAGCA	GCTATGCCAT	GAGCTGGGTC	CGCCAGGCTC	100
CAGGGAAGGG	GCTGGAGTGG	GTCTCAACTA	TTAGTGTTAG	TGGTATTACC	150
ACATACTACG	TAGACTCCGT	GAAGGGCCGG	TTCACCATCT	CCAGAGACAA	200
TTCCAAGAAC	ATTCTGTATC	TGCAAATGAA	CAGCCTGAGA	GCCGAGGACA	250
CGGCCGTATA	TTACTGTGCG	AAACGGATT	TTGGAGTGGT	CTGGGGCCAG	300
GGAACCCTGG	TCACCGTCTC	CTCAGCCTCC	ACCAAGGGCC	CATCGGTCTT	350
CCCCCTGGCG	CCCTGCTCCA	GGAGCACCTC	CGAGAGCACA	GCGGCCCTGG	400
GCTGCCTGGT	CAAGGACTAC	TTCCCCGAAC	CGGTGACGGT	GTCGTGGAAC	450
TTAGGCGCTC	TGACCAGCGG	CGTGCACACC	TTCCCAGCTG	TCCTACAGTC	500
CTA					503

(SEQ ID NO:79)

2.6.1 Heavy Protein

GGGLVQPGGS	LRLSCAASGF	TFSSYAMSWV	RQAPGKGLEW	VSTISVSGIT	50
TYYVDSVKGR	FTISRDN SKN	ILYLQMN SLR	AEDTAVYYCA	KRIFGVVWGO	100
GT LVTVSSAS	TKGPSVFPLA	PCSRSTSEST	AALGCLVKDY	FPEPVTVSWN	150
LGALTSGVHT	FPAVLQS				167

(SEQ ID NO:40)

2.6.1 Kappa cDNA

GGAATTCGGC	TTGATATTCA	GCTGACTCAG	TCTCCATCCT	CACTGTCTGC	50
ATCTGTAGGA	GACAGAGTCA	CCATCACTTG	TCGGGCGAGT	CAGGGCATT	100
GCATTTATTT	AGCCTGGTTT	CAGCAGAGAC	CAGGGAAAGC	CCCTAAGTCC	150
CTGATCTATG	CTGCATCCAG	TTTGCAAAGT	GGGGTCCCAT	CAAAGTTCAG	200
CGGCAGTGGA	TCTGGGACAG	ATTTCACTCT	CACCATCAGC	AGCCTGCAGC	250
CTGAAGATTT	TGCAACTTAT	TACTGCCAAC	AATATAATAG	TTACCCATT	300
ACTTTCGGCC	CTGGGACCAA	AGTGGATATC	AAACGAACTG	TGGCTGCACC	350
ATCTGTCTTC	ATCTTCCCGC	CATCTGATGA	GCAGTTGAAA	TCTGGAAGT	400
CCTCTGTTGT	GTGCCTGCTG	AATAACTTCT	ATCCCAGAGA	GGCCAAAGTA	450
CAGTGGAAGG	TGGATAACGC	CCTCCAATCG	GGTAAGCCGA	ATTC	494

(SEQ ID NO:80)

2.6.1 Kappa Protein

GIRLDIQLTQ	SPSSLSASVG	DRVITICRAS	QGISIYLAWF	QQRPGKAPKS	50
LIYAASSLQS	GVPSKFSGSG	SGTDFTLTIS	SLQPEDFATY	YCQYNSYPF	100
TFGPGTKVDI	KRTVAAPSVF	IFPPSDEQLK	SGTASVVCLL	NNFYPPREAKV	150
QWKVDNALQS	GKPN				164

(SEQ ID NO:41)

FIG. 33

IgM Antibody Sequences

CEM 10.1 C3 Heavy Protein

GLLKPSETLSLTCAVYGGSGYYSWI RQPPGKGLEWIGEINHSGSTNYPNPSLKSRVTISVDTSKNQFS
 CDR1 CDR2
 LKLSVTAADTAVYYCARGTEYYYYYGGMDVWGQGTTVTVSSGSASAPTLFPLVSCENSPSDTSSVAVG
 CDR3
 CLAQDELDPXITFSWKYKNNSDISSTRGFPSVLRGGKYAATSQVLLPSKDVMQGTDEHVVTGSKE

CEM 10.1 C3 Kappa Protein

LSLPVTPGEPASISCRSSQSLHNSGNYLDWYLOKPGQSPQLLIYLGSNRASGVPDRFSGSGSGTDFTL
 CDR1 CDR2
 KISRVEAEDVGIIYYCMQTRQTPRTFGQGTKVEIKRTVAAPSVEIFPPSDEQLKSGTASVVCLLNNEYPRE
 CDR3
 AKEHQKSP

FIG. 34

IgM Antibody Sequences

CEM 10.1 G10 Heavy Protein

LVKPSETLSLTCTVSGGSISSYYWNWIRQPPGKGLEWIGIYIYSGSTNYPNPSLKSRVTISVDTSKNQFSL
CDR1 CDR2
KLSVTAADTAVYYCARDRGVGATGEDYWGQGT~~LVTVSSG~~SASAP~~TLFPLV~~SCENSPSDTSSVAVGCLAQ
CDR3
DFLPDSITFSWKYKNNSDISSTRGFPSVL~~RGGKYA~~TSQVLLPSKDV~~MQGTDE~~HKVC

CEM 10.1 G10 Kappa Protein

SQSPSSLSASVGERVTITCRASQGIRDELGWYQQKPGKAPKRLIYWASSLQSGVPSRFSGSGSGTEFTLT
CDR2

ISSLPEDFATYYCLQHNGYPRTFGQGTKVEIKRTVAAPSVEIFFPSDEQLKSGTASVVCLLNNFYPREA
CDR3

KEHQKSE

FIG. 35

IgM Antibody Sequences

CEM 10.12 F3 Heavy Protein

KKPGASVKVSCKASGYTFTSYDINWVRQATGQGLEWMGMPNSGNTGYAQKFQGRVTYNRNTSISTAYM
 CDR1 CDR2
 ELSSLRSEDTAVYYCARGGHGGSFYFYYGMDVWGQGTITVTVSSGSAAPTLEPLVSCENSPDTSVAV
 CDR3
 GCCLAQDFLPDSITFSWKYKNNSDISSTRGFPSVLRGGKYAATSQVLLPSKDV MQGTDEHVVCK

CEM 10.12 F3 Kappa Protein

HSLAVSLGERATINCKSSQSVLYSENNKNYLAWYQQKPGQPCDR2KLIIYWASTRESGVDPDRFGGSGSGTDFT
 LTISSLQAEDVAVYYCQQYYSTPRTFGQGTKVEIKRTVAAPSVEIFPPSDEQLKSGTASVVCLLNNEYPRCDR3
 EAKEHQKSP

*IgM Antibody Sequences**CEM 10.12 G5 Heavy Protein*

EVKKPGASVKVSCKASGYTETSYDINWVRQATGQGLEWMGMNPNSGNTGYAQKFQGRVTMTNRNTSISTA
CDR1
YMESSLRSEDVAVYYCAREEWLVRYYGMDVWGQGT^{CDR3}TVTVSSGSASAPTLFPLVSCENSPSDTSSVAVGC
LAQDFLPDSITFSWKYKNNSDISSSTRGFPSVLRGGKYAATSQVLLPSKDV MQGTDEHKV

CEM 10.12 G5 Kappa Protein

GQSPSSLSASVGDRTITCRASQDIRDNLGWYQQKPGKAPKRLIYAA^{CDR2}SNLQSGVPSRFSGSGSGTEFTLT
CDR1
ISSLQPEDEATYYCLQYKTYPWTFGQGT^{CDR3}KVEIKRTVAAPS^{CDR2}VFI^{CDR1}FPSPDEQLKSGTASVVCLLNNFYPREX
KEHQKSP

FIG. 37

IgM Antibody SequencesCEM 13.12 Heavy Protein

KLPE^{CDR1}TLSCAVYGGSE^{CDR2}FGYYWSWIRQPPGKGLEWIGEINHSGSTNYP^{CDR2}SLKSRVTISVDTSKNQFSLKL
SSVTAADTAVY^{CDR3}YCARGAAEY^{CDR3}YYYYGMDVWGQGT^{CDR3}TVTVSSGASAP^{CDR3}TLFPLVSCENS^{CDR3}PSDTSSVAVGCLA
QDFLPDXITFXWKYKNNSDISSTRGFPSVLRGGKYAATSQVLLPSKDV^{CDR3}MQGTDEHVVTGSKE

CEM 13.12 Kappa Protein

MPVTPGEPASISCRSSQ^{CDR1}LLHSNGYNYLDWYLQKPGQSP^{CDR2}QLLIYLGSNRASGV^{CDR2}PDREFSGSGGTDTLKI
SRVEAEDVGIYYCMQSLQI^{CDR3}PRLEGP^{CDR3}GTKVDIKRTVAAPSVEIFPPSDEQLKSGTASV^{CDR3}VCLLSN^{CDR3}FYPREAK

VQW

FIG. 38

IgM Antibody Sequences

CEM 13.5 Heavy Protein

SETLSLTCAVYGGSESGYYWSWIRQPPGKGLEWIGEINHGSTNYPNPSLKSRVTISVDTSKNQFSLKLSS
CDR1
 VTAADTAVYYCARGGTTVTEDAFDIWGQGTMTVSSSGSASAPTLFPLVSCENSPSDTSVAVGCLAQDFL
CDR3
 PDSITFSWKYKNNSDISSTRGFPSVLRGGKYAATSQVLLPSKDVMMQGTDE

CEM 13.5 Kappa Protein

LAVSLGERATINCKSSQSVLYSFNNKNYLAWYQQKPGQPPKLLIYWASTRESGVPDRESGSGSGTDFTLT
CDR1
 ISSLQAEDVAVYYCQQYYSTPRTEFGQGTKVEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREA
CDR3
 KVQWKVI

FIG. 39

IgG Antibody Sequences

2.4.4 Heavy Protein

NPQTTLTCTFSGFSLITRGVVDWIRQPPGKALQWLALIYW^{CDR1}NDDKRYSPSLKSR^{CDR2}LTITKDTSKNQVVL
 TMTNMDPVDTATYYCAHHFFEDSSGYYPEDSWGQGTLSVSSASTKGPSVFPLAPCSRSTSESTAALGCLV
 KDYFPEPVT

2.4.4 Kappa Protein

VTQSPLSLVTPGQPASISCKSSQSL^{CDR1}LHSDGKTYLYWLQKPGQPPQLLIYEAFNRFSGVPDRFSGSGG
 TDETLKISRVEAEDVGLYYCMQSIELPFTFEGGKVEIKRTVAAPSVEIFPPSDEQLKSGTASVVCLLNN
 FYPRKERV

FIG. 40

IgG Antibody Sequences

2.1.1 Heavy Protein

GGLVKPGGSLRLSCAASGETFESSYSMNWVRQAPGKGLEWVSSISSSSSYIYYADSVKGRFTISRDNAKN
 CDR1 CDR2
 SLYLQMNSLRAEDTAVYICARDSSGWYEDYFDYWGGTLVTVSSASTKGPSVFPLAPCSRSTSESTAALG
 CDR3
 CLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQS

2.1.1 Kappa Protein

LDIQLTQSPSSLASVGDRVTITCRASQDISIYLAWFQQRP GKAPKSLIYAASSLQSGVPSKFSGSGSGT
CDR1 CDR2

DFTLTISLQPEDFATYYCQQYNSYPETFPG
CDR3

FIG. 41

IgG Antibody Sequences

2.3.2 Heavy Protein

LTCTESGFSLITRGVGVDWIRQPPGKALQWLALIYWNDKRYSPSLKSRLTITKDTSKNQVVLTMTNMDP
 CDR1 CDR2
VDATATYCAHHEFDSSGYYPEDSWGQGTLSVSSASTKGPSVFPLAPCSRSTSESTAALGCLVKDYFPEP
 CDR3
 VTVSWNSGALTSGVHTFQL

2.3.2 Kappa Protein

FIG. 42

IgG Antibody Sequences**2.6.1 Heavy Protein**

GGGLVQPGGSLRLSCAASGETFSSYAMSWVRQAPGKGLEWVSTISVSGITTYYYVDSVKGRFTISRDN SKN
CDR2
ILYLQMNSLRAEDTAVYYCAKRIEGVVWGQGLTVTVSSASTKGPSVFPLAPCSRSTSESTAALGCLVKDY
CDR3
FPEPVTVSWNLGALTSGVHTFPAVLQS

2.6.1 Kappa Protein

GIRLDIQLTQSPSSLASVGDRTITCRASQGISIYLAWFQQRP GKAPKSLIYAASSLQSGVP SKESGSG
CDR1
SGTDFTLTISSLQPEDFATYYCQQYN SYPETFGPGTKVDIKRTVAAPSVEIFPPSDEQLKSGTASVVCLL
CDR3
NNFYPREAKVQWKVDNALQSGKEN

FIG. 43




FIG. 44A




FIG. 44B




FIG. 45A




FIG. 45B




FIG. 45C




FIG. 46




FIG. 47




FIG.48




FIG.49




FIG.50A




FIG. 50B




FIG. 50C




FIG. 50D




FIG.50E




FIG.50F

CD147 BINDING MOLECULES AS THERAPEUTICS

BACKGROUND OF THE INVENTION

[0001] 1. Summary of the Invention

[0002] In accordance with the present invention, we have discovered that the molecule CD147 as expressed on certain cells, such as T-cells, B-cells, and/or monocytes, can be utilized as a target for the treatment of a variety of diseases. In particular, we have demonstrated that an antibody that binds to CD147 and that results in the killing of such cells, for example, through the binding of complement, is efficacious in the treatment of diseases. Diseases in which such treatment appears efficacious include, without limitation, graft versus host disease (GVHD), organ transplant rejection diseases (including, without limitation, renal transplant, ocular transplant, and others), cancers (including, without limitation, cancers of the blood (i.e., leukemias and lymphomas) and pancreatic), autoimmune diseases (including, without limitation, lupus), inflammatory diseases (including, without limitation, arthritis), and others.

[0003] 2. Background of the Technology

[0004] In about 1982, a group from UCLA reported the generation of antibodies cytotoxic to human leukemia cells in mice through immunization with acute leukemia cells followed by formation of hybridomas and screening of the hybridomas in a microcytotoxicity assay in which toxicity of the antibody against the immunizing cells and normal lymphocytes was assayed. See U.S. Pat. Nos. 5,330,896 and 5,643,740, the disclosures of which are hereby incorporated by reference in their entirety. One hybridoma was recovered that was cytotoxic to tumor cells but non-toxic to normal cells (except activated T-cells, activated B-cells, and monocytes were also killed). Such hybridoma was cloned and isolated and deposited with the ATCC as HB 8214. The monoclonal antibody expressed by this hybridoma was designated CBL1, and is a murine IgM. The group further demonstrated that the antibody was reactive with an antigenic determinant that appeared to be present in the cytoplasm of both activated and nonactivated cells. However, the antigenic determinant appeared to be present on the extracellular membrane of only certain circulating cells, including, activated T-cells, activated B-cells, and resting and activated monocytes, but not present extracellularly on other circulating nonactivated cells.

[0005] The group also endeavored to isolate the antigen responsible for the observations. The patents characterize the antigenic determinant recognized by the CBL-1 antibody as being a molecule that:

[0006] (i) is present on the cell membrane and within the cytoplasm of tumor cells and activated lymphocytes;

[0007] (ii) is present in the cytoplasm of unstimulated normal peripheral blood lymphocytes but when these cells are stimulated by antigens or by mitogens, said antigen appears also on the cell membrane;

[0008] (iii) is present on lymphocytes activated in vitro by mitogens;

[0009] (iv) is capable of binding to CBL1 monoclonal antibody which is produced by the hybridoma cell line having the ATCC number HB8214;

[0010] (v) functions as an autocrine growth factor produced by tumor cells and activated lymphocytes;

[0011] (vi) binds to the surface membrane of tumor cells and stimulates the growth of these cells and cells of the lymphoid series;

[0012] (vii) is present in the medium from growing cancer cells and in the serum of patients with cancer and diseases in which activated lymphocytes are present; and

[0013] (viii) has a molecular weight of approximately 15,000 daltons.

[0014] No improved identification of the antigen to which the CBL1 antibody binds has been accomplished with respect to the UCLA group's papers and patents. Nevertheless, the CBL1 antibody has been effective in patients in the treatment of a variety of diseases including: graft versus host disease (GVHD) and kidney transplant rejection. See e.g., Heslop et al. *The Lancet* 346:805-806 (1995) (GVHD); Benamin *Clinical Trial Monitor* Abstract No. 13385 (1995); Takahashi et al. *The Lancet* 2:1155-1158 (1983) (kidney allograft rejection); Takahashi *Transplantation Proceedings* 17:10-12 (1985) (kidney allograft rejection); Oei et al. *Transplantation Proceedings* 17:13-16 (1985) (kidney allograft rejection). In connection with such studies, there has been no evidence of safety concerns or cross-reactivity. The following papers relate to additional characterization of the CBL1 antibody: Billing et al. *Hybridoma* 1:303-311 (1982); Billing et al. *Clin. Exp. Immunol.* 49:142-148 (1982); Chatterjee et al. *Hybridoma* 1:369-377 (1982); Billing R. and Chatterjee S. *Transplantation Proceedings* 15:649-650 (1983); Kinukawa T. and Terasaki P. I. *Transplantation Proceedings* 1:993-998 (1985); Billing in *Monoclonal Antibodies: Diagnostic and Therapeutic Use in Tumor and Transplantation* Ch. 9, 85-90 (Chatterjee ed., PSG Publ. Co., Inc. (1985)); Billing et al. in *Monoclonal Antibodies: Diagnostic and Therapeutic Use in Tumor and Transplantation* Ch. 2, 11-19 (Chatterjee ed., PSG Publ. Co., Inc. (1985)).

[0015] Human Graft Versus Host Disease (GVHD) was first described by Mathe et al. in 1960 (Mathe et al. "Nouveaux essais de greffe de moelle osseuse homologue apres irradiation totale chez des enfants atteints de leucemie aigue en remission. Le probleme du syndrome secondaire chez l'homme" *Rev Fr Etud Clin Biol* 15:115-161 (1960)). Essentially GVHD is the clinical manifestation of an immunological reaction between donor cells and host tissue. The clinical syndrome consists of skin rash, gastro-intestinal symptoms, and hepatic dysfunction seen usually within two weeks of allogeneic bone marrow transplant. The immunopathogenesis requires recognition of host antigens by immunocompetent donor cells; immunosuppressed host (recipient); and alloantigenic differences to exist between donor and recipient. The immunocompetent donor cells are mature T-cells (Ferrara J L and Deeg H J "Graft versus Host Disease" *NEJM* 324:667 (1991) and the clinical severity of the disease correlates with the number of T-cells transferred to the patient (Ferrara J L and Deeg H J "Graft versus Host Disease" *NEJM* 324:667 (1991)).

[0016] The clinical features of acute GvHD include dermatitis, jaundice and gastro-intestinal involvement. These symptoms may occur alone or in any combination and can

range from mild to life-threatening. Skin involvement is the most common manifestation. The most severe manifestation of skin involvement includes bullous lesions similar to third degree burns. Jaundice is brought about from an elevated bilirubin with and without alteration of other liver enzymes. Gastro-intestinal involvement includes watery diarrhea. This diarrhea can be voluminous and bloody, causing life-threatening fluid and electrolyte losses as well as a portal of entry for infections. Other patients may experience severe ileus. Upper GI involvement is less common. This presents as anorexia, dyspepsia, food intolerance and nausea/vomiting. Most patients with GI involvement require total parenteral nutrition (TPN) support.

[0017] Strategies for prevention and possibly treatment should be and sometimes are, directed towards removal of T-cells from the donor marrow or toward blocking their activation. However, the T-depleted marrow results in a higher rate of graft failure that is usually fatal. An additional concern associated with T-depleted marrow is the increased relapse rate in marrow recipients with a primary diagnosis of leukemia. A graft versus leukemia effect, mediated by donor T-cells, also mitigates against using a T-depleted marrow in allogeneic bone marrow transplantation.

[0018] Clinically significant acute GVHD (Grades II-IV) occurs in up to 50% of patients who receive a marrow from a HLA genotypically identical sibling. If unrelated matched donors are used, the incident increases to 80% in some studies. The greater the HLA incompatibility, the greater the incidence and severity of GVHD.

[0019] The primary treatment for acute GvHD is prevention. Prevention regimens include the use of immunosuppression therapy and T-cell depletion of the donor cells. "Standard" first-line therapy consists of glucocorticoids. Approximately 20-25% of patients achieve a complete response and patients who do not respond have a poor outcome. Those patients who continue to require treatment with steroids are susceptible to all of the untoward effects of steroid use. These untoward effects include increased susceptibility to infections, GI bleed, altered metabolic states, hypertension, etc.

[0020] Glucocorticoids, cyclosporine, methotrexate, cyclophosphamide have all been used in prevention as well as treatment of GVHD. Anti-thymocyte globulin (ATG) has been used for many years. All of these agents are potentially quite toxic. Monoclonal antibodies such as anti-Interleukin-2 and immunotoxins like anti-CD5-ricin have been used and found to be of limited success. A humanized anti-TAC was used for prophylaxis of GVHD but failed in the treatment protocols.

[0021] Because of the indication that CBL1 was effective in treating GVHD, we undertook additional investigations of the CBL1 antibody. In connection with such additional work, we have now demonstrated that the CBL1 antibody, in fact, appears to bind to and be efficacious with respect to the CD147 antigen as expressed on certain cells, such as T-cells, B-cells, and/or monocytes through the process of complement dependent cytotoxicity (killing).

[0022] CD147 is a member of the immunoglobulin (Ig) superfamily that is expressed on a large number of different cells in a variety of tissues. It was originally named human Basigin (for basic immunoglobulin superfamily) and was first

cloned in about 1991. (Miyauchi et al. *J Biochem (Tokyo)* 110:770-774 (1991); Kanekura et al. *Cell Struct Funct* 16:23-30 (1991); Miyauchi et al. *J Biochem (Tokyo)* 110:770-774 (1991)). The molecule is composed of approximately 269 amino acids (Miyauchi et al. *J Biochem (Tokyo)* 110:770-774 (1991)) and is a glycoprotein with about 40% of its molecular weight made up of carbohydrate, having a predicted deglycosylated molecular weight of approximately 27 KD and a fully glycosylated molecular weight of between 43-66 KD (Kanekura et al. *Cell Struct Funct* 16:23-30 (1991)). The Basigin gene was mapped to Chromosome 19p13.3 (Kaname et al. *Cytogenet Cell Genet* 64:195-197 (1993)).

[0023] The molecule has been identified to possess homology with, or identity to, a number of other molecules, including:

[0024] Mouse Basigin (Miyauchi et al. *J Biochem (Tokyo)* 107:316-323 (1990); Joseph et al. *Adv Exp Med Biol* 342:389-391(1993); Kaname et al. *J Biochem (Tokyo)* 118:717-724 (1995));

[0025] Rabbit Basigin (Schuster et al. *Biochim Biophys Acta* 1311:13-19 (1996));

[0026] Mouse gp42 (Altruda et al. *Gene* 85:445-451 (1989); Imboden et al. *J Immunol* 143:3100-3103 (1989); Cheng et al. *Biochim Biophys Acta* 1217:307-311 (1994));

[0027] Chicken HT7 or 5A11 (Albrecht et al. *Brain Res* 535:49-61 (1990); Seulberger et al. *EMBO J* 9:2151-2158 (1990); Miyauchi et al. *J Biochem (Tokyo)* 110:770-774 (1991); Janzer et al. *Adv Exp Med Biol* 331:217-221 (1993); Lobrinus et al. *Brain Res Dev Brain Res* 70:207-211 (1992); Seulberger et al. *Neurosci Lett* 140:93-97 (1992); Fadool J M & Linser P J *J Neurochem* 60:1354-136 (1993); Fadool J M & Linser P J *Dev Dyn* 196:252-262 (1993); Unger et al. *Adv Exp Med Biol* 331:211-215 (1993); Rizzolo L J & Zhou S J *Cell Sci* 108:3623-3633 (1995); Ikeda et al. *Neurosci Lett* 209:149-152 (1996); Fadool J M & Linser P J *Biochem Biophys Res Commun* 229:280-286 (1996));

[0028] Neurothelin (Schlosshauer B & Herzog K H *J Cell Biol* 110:1261-1274 (1990); Schlosshauer B *Development* 113:129-140 (1991); Schlosshauer B *BioEssays* 15:341-346 (1993); Schlosshauer et al. *Eur J Cell Biol* 68:159-166 (1995));

[0029] M6 leukocyte activation antigen (Felzmann et al. *J Clin Immunol* 11:205-212 (1991); Gadd et al. *Rheumatol Int* 12:153-157 (1992); Kasinrerker et al. *J Immunol* 149:847-854 (1992));

[0030] OX-47 (Fossum et al. *Eur J Immunol* 21:671-679 (1991); Fossum et al. *Eur J Immunol* 21:671-679 (1991); Casselia et al. *J Anat* 189:407-415 (1996));

[0031] Mo3 (Mizukami et al. *J Immunol* 147:1331-1337 (1991));

[0032] CE9 (Petruszak et al. *J Cell Biol* 114:917-927 (1991); Scott L J & Hubbard A L *J Biol Chem* 267:6099-6106 (1992); Nehme et al. *J Cell Biol* 120:687-694 (1993); Cesario M M & Bartles J R *J Cell Sci* 107:561-570 (1994); Cesario et al. *Dev Biol* 169:473-486 (1995); Nehme et al. *Biochem J* 310:693-698 (1995));

[0033] EMMPRIN (Biswas et al. *Cancer Res* 55:434 (1995); DeCastro et al. *J Invest Dermatol* 106:1260-1265 (1996));

[0034] RET-PE2 (Finnemann et al. *Invest Ophthalmol Vis Sci* 38:2366-2374 (1997));

[0035] Ok^a Blood Group Antigen (Spring et al. *Eur J Immunol* 27:891-897 (1997)); and

[0036] 1W5 (Seulberger et al. *EMBO J* 9:2151-2158 (1990)).

[0037] Indeed, Seulberger et al. *Neurosci Lett* 140:93-97 (1992) demonstrated that HT7, Neurothelin, Basigin, gp42 and OX-47 were each names for one molecule which is a developmentally regulated immunoglobulin-like surface glycoprotein which is present on blood-brain barrier endothelium, epithelial tissue barriers, and neurons. Further, Kasinrerker et al. *J Immunol* 149:847-854 (1992) demonstrated that the human leukocyte activation antigen M6 is a member of the Ig superfamily and is the species homologue of rat OX47, mouse Basigin, and chicken HT7 antigens. EMMPRIN was demonstrated to be identical to the M6 antigen and human Basigin (Biswas et al. *Cancer Res* 55:434 (1995)). See also Guo et al. "Characterization of the gene for human EMMPRIN, a tumor cell surface inducer of matrix metalloproteinases" *Gene* 220:99-108 (1998) conducted additional characterization of the gene for human EMMPRIN;

[0038] Through its homology with the related molecules, CD147 has been shown or postulated to have a role in a number of physiological processes, diseases, and/or conditions. For example, an early role postulated for the molecule was activity in the blood-brain barrier. Such relationship was first demonstrated with respect to the chick HT7 antigen (Risau et al. *EMBO J* 5:3179-3183 (1986); Albrecht et al. *Brain Res* 535:49-61 (1990); Seulberger et al. *EMBO J* 9:2151-2158 (1990); Janzer et al. *Adv Exp Med Biol* 331:217-221 (1993); Lobrinus et al. *Brain Res Dev* 70:207-211 (1992); Unger et al. *Adv Exp Med Biol* 331:211-215 (1993)). A similar relationship was observed in connection with Neurothelin (Schlosshauer B & Herzog K H *J Cell Biol* 110:1261-1274 (1990); Schlosshauer B *Development* 113:129-140 (1991); Schlosshauer B *BioEssays* 15:341-346 (1993); Schlosshauer et al. *Eur J Cell Biol* 68:159-166 (1995)). The molecule has also been postulated to be involved in development and activation of various cells, for example: lymphocyte activated killer (LAK) cell activation (Imboden et al. *J Immunol* 143:3100-3103 (1989)), T-cell activation (Paterson et al. *Mol Immunol* 24:1281-1290 (1987); Kirsch et al. *Tissue Antigens* 50:147-152 (1997)), leukocyte activation (Fossum et al. *Eur J Immunol* 21:671-679 (1991); Fossum et al. *Eur J Immunol* 21:671-679 (1991)), and mononuclear phagocyte activation (Mizukami et al. *J Immunol* 147:1331-1337 (1991)). Other regulatory, signaling, and recognition functions have also been postulated, for instance: MHC function (Miyachi et al. *J Biochem (Tokyo)* 107:316-323 (1990)), signal transduction and membrane transport (Kasinrerker et al. *J Immunol* 149:847-854 (1992); Berditchevski et al. *J Biol Chem* 272:29174-29180 (1997)), cellular recognition (Fadool J M & Linser P *J Dev Dyn* 196:252-262 (1993); Kaname et al. *Cytogenet Cell Genet* 64:195-197 (1993)), cellular adhesion (Miyachi et al. *J Biochem (Tokyo)* 110:770-774 (1991); Seulberger et al. *Neurosci Lett* 140:93-97 (1992); Joseph et al. *Adv Exp Med Biol* 342:389-391 (1993); Sudou et al. *J Biochem (Tokyo)* 117:271-275 (1995)), intercellular stimulation and matrix metalloproteinase synthesis (Biswas et al. *Cancer*

Res 55:434 (1995)), tissue remodeling (Guo et al. *J Biol Chem* 272:24-27 (1997)), metabolism, and sperm development and maturation (Petruszak et al. *J Cell Biol* 114:917-927 (1991); Nehme et al. *J Cell Biol* 120:687-694 (1993); Cesario M M & Bartles J R *J Cell Sci* 107:561-570 (1994); Cesario et al. *Dev Biol* 169:473-486 (1995)). CD147 also appears to have a role in retinal development and disease, see Marmorstein et al. "Morphogenesis of the retinal pigment epithelium: toward understanding retinal degenerative diseases" *Ann N Y Acad Sci* 857:1-12 (1998) (suggested that N-CAM and EMMPRIN are potentially important molecules in other RPE functions necessary for photoreceptor survival). See also Marmorstein et al. "Apical polarity of N-CAM and EMMPRIN in retinal pigment epithelium resulting from suppression of basolateral signal recognition" *J Cell Biol* 142:697-710 (1998).

[0039] The molecule has also been investigated for a potential association in both rheumatoid and reactive arthritis (Felzmann et al. *J Clin Immunol* 11:205-212 (1991); Gadd et al. *Rheumatol Int* 12:153-157 (1992)) and renal disease (Schuster et al. *Biochim Biophys Acta* 1311:13-19 (1996)). Moreover, certain clear associations between the molecule and cancer have also been indicated (Biswas *Biochem Biophys Res Commun* 109:1026 (1982); Miyachi et al. *J Biochem (Tokyo)* 110:770-774 (1991); Biswas et al. *Cancer Res* 55:434 (1995); Guo et al. *J Biol Chem* 272:24-27 (1997); Guo et al. *J Biol Chem* 272:24-27 (1997)). See also Lim et al. "Tumor-derived EMMPRIN (extracellular matrix metalloproteinase inducer) stimulates collagenase transcription through MAPK p38" *FEBS Lett* 441:88-92 (1998); van den Oord et al. "Expression of gelatinase B and the extracellular matrix metalloproteinase inducer EMMPRIN in benign and malignant pigment cell lesions of the skin" *Am J Pathol* 151:665-70 (1997); Polette et al. "Tumor collagenase stimulatory factor (TCSF) expression and localization in human lung and breast cancers" *J Histochem Cytochem* 45:703-9 (1997).

[0040] A mouse model in which the Basigin gene was knocked-out has been examined (Igakura et al. *Biochem Biophys Res Commun* 224:33-36 (1996)). The work indicated that the molecule was not necessarily active in the blood-brain barrier. However, the work indicated that there was enhanced interaction in connection with lymphocyte activation as well as an abnormal response to irritating odors. Later work indicated certain abnormalities in sensory and memory functions in such model (Naruhashi et al. *Biochem Biophys Res Commun* 236:733-737 (1997)).

[0041] In connection with the expression of CD147, see Woodhead et al. "From sentinel to messenger: an extended phenotypic analysis of the monocyte to dendritic cell transition" *Immunology* 94:552-9 (1998) demonstrated that CD147 was expressed on dendritic cells, Ghannadan et al. "Phenotypic characterization of human skin mast cells by combined staining with toluidine blue and CD antibodies" *J Invest Dermatol* 111:689-95 (1998) demonstrated that clustered CD antigens (including CD147) were detectable on skin mast cells, Mutin et al. "Immunologic phenotype of cultured endothelial cells: quantitative analysis of cell surface molecules" *Tissue Antigens* 50:449-58 (1997) discussed quantitative analysis of cell surface molecules on cultured endothelial cells (HUVEC).

[0042] In view of the foregoing, CD147 has been implicated as a potentially useful target for the treatment of

diseases. However, at the same time, CD147 is expressed in and on many cells that are widely distributed amongst many tissues. For example, the Ok^a blood group antigen is expressed on virtually all cells (Williams et al. *Immunogenetics* 27:322-329 (1988)). OX-47 has been disclosed to be on most immature cells, endothelial cells, and cells with excitable membranes (Fossum et al. *Eur J Immunol* 21:671-679 (1991)). Similarly, Basigin was demonstrated to be expressed not only in endothelial cells but was also found in a variety of tissues, including, the spleen, small intestine, kidney, and liver in relatively high levels and in small quantities in the testes (Kanekura et al. *Cell Struct Funct* 16:23-30 (1991)). CE9 was disclosed to be widely expressed on rat hepatocytes (Scott L J & Hubbard A L *J Biol Chem* 267:6099-6106 (1992)). Seulberger et al. *Neurosci Lett* 140:93-97 (1992) demonstrated that the HT7 molecule (which is identical to Neurothelin, Basigin, gp42, and OX-47) was expressed on the blood-brain barrier, choroid plexus (blood-CNS fluid barrier), retinal epithelium (blood-eye barrier), neurons, kidney tubules, some endothelium, epithelium, and epithelial tissue barriers. The CE9 antigen (which was demonstrated to possess identity to the OX-47 antigen) is expressed, to some extent, in virtually all rat tissues (Nehme et al. *Biochem J* 310:693-698 (1995)). Because of the broad tissue distribution, there would be a number of concerns related to the safety of any therapy that inhibited or killed cells expressing it.

[0043] There is some evidence that there may be different forms of CD147, stemming from, for example, differential glycosylation or alternative splicing of the molecule (Kanekura et al. *Cell Struct Funct* 16:23-30 (1991) (Basigin); Schlosshauer B *Development* 113:129-140 (1991) (Neurothelin); Fadool J M & Linser P J *J Neurochem* 60:1354-136 (1993) (5A11/HT7); Nehme et al. *J Cell Biol* 120:687-694 (1993) (CE9); DeCastro et al. *J Invest Dermatol* 106:1260-1265 (1996) (EMMPRIN); Spring et al. *Eur J Immunol* 27:891-897 (1997) (Ok^a)).

BRIEF DESCRIPTION OF THE DRAWING FIGURES

[0044] **FIG. 1** is a 12% SDS-PAGE/Western Blot showing the binding of particular antibodies to CEM cell membrane extracts lysates. Lane A: rabbit-anti-mouse-hn-RNP-K protein antibody; Lane B: ABX-CBL antibody; Lane C: 2.6.1 antibody (also referred to herein as cem2.6 and ABX-Rb2); Lane D: anti-CD147 antibody (Pharminogen); and Lane E: anti-CD147 antibody (RDI). Sample: 5 microliters CEM Cell Extract.

[0045] **FIGS. 2A-2B** is an analysis of the components obtained from the CBL1 antibody produced by the hybridoma cell line having ATCC Deposit No. BB 8214. The data demonstrate that the CBL1 IgM antibody produced by the HB 8214 hybridoma is the active component that inhibits MLR in the presence of complement.

[0046] **FIG. 3** is a graph comparing the inhibition of MLR using antibodies from various CBL1 subclones in comparison to CBL1.

[0047] **FIG. 4** is a graph comparing MLR inhibition utilizing ABX-CBL in the presence of rabbit and human complement.

[0048] **FIG. 5** is a graph comparing the activity of the ABX-CBL antibody and the 2.6.1 antibody (also referred to

as cem 2.6) in inhibiting the MLR assay. The data demonstrate that the 2.6.1 antibody is not an effective inhibitor.

[0049] **FIGS. 6A-6B:** FACS analyses of activated lymphocytes demonstrating co-expression of CD147 and CD25.

[0050] **FIGS. 7A-7D:** FACS analyses of PBMC demonstrating the selective upregulation of CD25 upon stimulation, and the specific depletion of the same cells after treatment with ABX-CBL and complement. **FIG. 7A:** untreated PBMC. **FIGS. 7B and 7D:** PBMC stimulated with ConA. **FIG. 7C:** PBMC stimulated with ConA, then treated with ABX-CBL plus complement.

[0051] **FIGS. 8A-8D** compare FACS analyses of PBMC demonstrating the selective upregulation of CD25 upon stimulation, and the specific depletion of the same cells after treatment with ABX-CBL and complement. **FIG. 8A:** PBMC+ConA; **FIG. 8B:** CBL-1 only/Medium; **FIG. 8C:** Complement only/Medium; **FIG. 8D:** CBL-1+complement/Medium. M1: CD25 high (depleted); M2: CD25 low (undepleted); M3: CD25 null (undepleted).

[0052] **FIGS. 9A-9D** show another series of FACS analyses of PBMC demonstrating the selective upregulation of CD25 and CD147 upon stimulation.

[0053] **FIGS. 10A-10F** show a comparison of activated T-cells (**FIGS. 10A-10B**), activated monocytes, (**FIGS. 10C-10D**) and activated B-cells (**FIGS. 10E-10F**) before and after treatment with ABX-CBL and complement and demonstrating the specific depletion of the same cells upon treatment with ABX-CBL and complement.

[0054] **FIGS. 11A-11F** shows a similar comparison of subpopulations of activated T-cells (**FIGS. 11A-11B**), activated B-cells (**FIGS. 11C-11D**), and activated monocytes (**FIGS. 11E-11F**) before and after treatment with ABX-CBL and complement. The data demonstrate the specific depletion of the same cells upon treatment with ABX-CBL and complement.

[0055] **FIG. 12** illustrates that the mode of action of ABX-CBL is by depleting leukocyte subpopulations. The table compares cell type, surface markers, and Complement-Dependent Cytotoxicity (CDC) depletion of leukocyte subpopulations.

[0056] **FIG. 13** is a table comparing cell, cell type, CD147 expression, and CDC after treatment of the cells with ABX-CBL and complement. The data demonstrate that not all cells that express CD147 are killed upon such treatment.

[0057] **FIG. 14** is a table summarizing the expression of CDC resistant molecules on CBL-1⁺ cells. The chart compares cell, cell type, CD147 expression, CDC after treatment of the cells with ABX-CBL and complement, and expression of the complement inhibitory molecules CD55 and CD59. The data demonstrate that of these cells, only cells that do not express both CD55 and CD59 are killed upon such treatment.

[0058] **FIGS. 15A-15C** present FACS analyses showing the expression of CD147 on the human endothelial cell line ECV-304.

[0059] **FIGS. 16A-16C** present FACS analyses showing the expression of CD147 on the human endothelial cell line HUVEC-C.

[0060] **FIG. 17** is a graph showing the effects of ABX-CBL and complement on the human endothelial cell line ECV-304 in comparison to the effects of the same on CEM cells.

[0061] **FIG. 18** is a graph showing the effect of ABX-CBL on human endothelial cell line HUVEC-C in comparison to the effects of the same on CEM cells.

[0062] **FIGS. 19A-19C** present FACS analyses showing the expression of the complement inhibitory molecules CD46, CD55, and CD59 on the human endothelial cell line ECV-304.

[0063] **FIGS. 20A-20C** present FACS analyses showing the expression of the complement inhibitory molecules CD46, CD55, and CD59 on the human endothelial cell line HUVEC-C.

[0064] **FIG. 21** is a schematic diagram of the vector utilized for cloning and expression of CD147 cDNA in COS cells.

[0065] **FIG. 22** is a schematic diagram of the pBK-CMN phagemid vector utilized for cloning and expression of CD147 cDNA in COS and *E. coli* cells.

[0066] **FIG. 23** is a SDS-PAGE/Western Blot of CD147 expressed in COS cells (**FIG. 23A**) and *E. coli* (**FIG. 23B**). **FIGS. 23A-23B**: Antibodies: Pharmingen (panel A), 2.6.1 (panel B), and ABX-CBL (panel C). **FIG. 23A**: 5 μ L CEM cell membrane extract (Lane 1); 7.5 μ L control vector transfected COS cell extract (Lane 2); 7.5 μ L CD147 transfected COS cell extract (Lane 3). **FIG. 23B**: Clone 1: CD147-Transfected, uninduced (Lane 1); Clone 1: CD147-Transfected, induced (Lane 2); Clone 5: Control Vector Transfected, uninduced (Lane 3); Clone 5: Control Vector Transfected, induced (Lane 4).

[0067] **FIGS. 24-33** are heavy chain and kappa chain cDNA and protein sequences of or for the antibodies: CEM 10.1 C3 (**FIG. 24**), CEM 10.1 G10 (**FIG. 25**), CEM 10.12 F3 (**FIG. 26**), CEM 10.12 G5 (**FIG. 27**), CEM 13.12 (**FIG. 28**), CEM 13.5 (**FIG. 29**), 2.4.4 (**FIG. 30**), 2.1.1 (**FIG. 31**), 2.3.2 (**FIG. 32**), and 2.6.1 (**FIG. 33**).

[0068] **FIGS. 34-43** are heavy chain and kappa chain protein sequences of or for the antibodies: CEM 10.1 C3 (**FIG. 34**), CEM 10.1 G10 (**FIG. 35**), CEM 10.12 F3 (**FIG. 36**), CEM 10.12 G5 (**FIG. 37**), CEM 13.12 (**FIG. 38**), CEM 13.5 (**FIG. 39**), 2.4.4 (**FIG. 40**), 2.1.1 (**FIG. 41**), 2.3.2 (**FIG. 42**), and 2.6.1 (**FIG. 43**) showing CDR positions.

[0069] **FIGS. 44A-44B** show the amino acid sequences and structure of human heavy chains derived from CBL-1 specific hybridomas showing alignment against the germline V-segment genes.

[0070] **FIGS. 45A-45C** and **FIG. 46** show amino acid sequences and structure of human kappa chains derived from CBL-1 specific hybridomas, showing alignment against the germline V-segment genes.

[0071] **FIG. 47** is a restriction map of the vector pWBFNP MCS that was utilized for the construction and cloning of certain constructs in accordance with the invention.

[0072] **FIG. 48** is a schematic restriction map of the vector pIK6.1+Puro that was utilized for the construction and cloning of certain constructs in accordance with the invention.

[0073] **FIG. 49** shows a comparison of the activity of the ABX-CBL antibody and the 2.6.1 multimeric IgM antibody (also known as ABX-Rb2) in inhibiting the MLR assay, demonstrating that the 2.6.1 multimeric IgM antibody is effective in inhibition of MLR. C: Rabbit complement.

[0074] **FIGS. 50A-50F** provide additional detail of the cloning strategy utilized in connection with the generation of CD147-IgG2 and gp42-IgG2 fusion proteins for use in connection with the generation of surrogate antibodies for use in animal models.

SUMMARY OF THE INVENTION

[0075] In accordance with a first aspect of the present invention, there is provided an isolated monoclonal antibody having an isotype that fixes complement and a variable region that binds to the epitope on CD147 bound by the IgM monoclonal antibody ABX-CBL, with the proviso that the antibody is not CBL1. In a preferred embodiment, the antibody in the presence of complement acts to selectively kill cells selected from the group consisting of activated T-cells, activated B-cells, and monocytes but is substantially non-toxic to resting T-cells and resting B-cells. In another preferred embodiment, the antibody is a human antibody. In another preferred embodiment, the antibody has an isotype selected from the group consisting of murine IgM, murine IgG2a, murine IgG2b, murine IgG3, human IgM, human IgG1, and human IgG3.

[0076] In accordance with a second aspect of the present invention, there is provided an isolated monoclonal antibody having an isotype that fixes complement and a variable region that binds to CD147 on populations of activated T-cells, activated B-cells, and resting or activated monocytes, that, in the presence of complement, selectively depletes such populations through complement mediated killing while being substantially nontoxic to other cells, with the proviso that the antibody is not CBL1. In a preferred embodiment, the antibody is a human antibody. In another preferred embodiment, the antibody has an isotype selected from the group consisting of murine IgM, murine IgG2a, murine IgG2b, murine IgG3, human IgM, human IgG1, and human IgG3.

[0077] In accordance with a third aspect of the present invention, there is provided an isolated monoclonal antibody having the following characteristics: binds to CD147; shows a binding against CEM cell lysates on Western blot similar to that provided in **FIG. 1**; an isotype selected from the group consisting of murine IgM, murine IgG2a, murine IgG2b, murine IgG3, human IgM, human IgG1, and human IgG3, competes with ABX-CBL for binding to CD147; cross reacts with hn-RNP-k protein; binds to a consensus sequence on CD147 comprising RVRSS; selectively kills activated T-cells, activated B-cells, and monocytes in a MLR assay only in the presence of complement; and is substantially non-toxic to cells expressing CD55 and CD59, with and without the presence of complement, with the proviso that the antibody is not CBL1.

[0078] In accordance with a fourth aspect of the present invention, there is provided a method to select an anti-CD147 antibodies for the treatment of disease, comprising: generating antibodies that bind to CD147 and that are capable of binding complement; assaying the antibodies for one or more of the following properties: competition with

ABX-CBL for binding to CD147; capability to selectively kill activated T-cells, activated B-cells, and monocytes in a MLR assay only in the presence of complement; and being substantially non-toxic to cells expressing CD55 and CD59, with and without the presence of complement, with the proviso that the antibody is not CBL1. In a preferred embodiment, the method comprises assaying the antibodies for binding to CEM cell lysates on Western blot in a manner similar to that provided in FIG. 1. In another preferred embodiment, the method comprises assaying the antibodies for binding to a consensus sequence in a peptide of RXRS. In another preferred embodiment, the method comprises assaying the antibodies for cross reaction with hn-RNP-k protein. In another preferred embodiment, the method comprises assaying the antibodies for binding to a form of CD147 expressed by COS cells and *E. coli* cells.

[0079] In accordance with a fifth aspect of the present invention, there is provided a method for preventing or lessening the severity of disease, comprising providing to a subject in need of such treatment an antibody that has an isotype that fixes complement and a variable region that binds to CD147 on populations of activated T-cells, activated B-cells, and resting or activated monocytes, that, in the presence of complement, selectively depletes such populations through complement mediated killing while being substantially nontoxic to other cells, with the proviso that the antibody is not CBL1. In a preferred embodiment, the antibody is a human antibody. In another preferred embodiment, the antibody has an isotype is selected from the group consisting of murine IgM, murine IgG2a, murine IgG2b, murine IgG3, human IgM, human IgG1, and human IgG3.

[0080] In accordance with a sixth aspect of the present invention, there is provided a method to prevent or lessen the severity of GVHD, comprising providing to a subject in need of such treatment an antibody that has an isotype that fixes complement and a variable region that binds to CD147 on populations of activated T-cells, activated B-cells, and resting or activated monocytes, that, in the presence of complement, selectively depletes such populations through complement mediated killing while being substantially non-toxic to other cells, with the proviso that the antibody is not CBL1. In a preferred embodiment, the antibody is a human antibody. In another preferred embodiment, the antibody has an isotype is selected from the group consisting of murine IgM, murine IgG2a, murine IgG2b, murine IgG3, human IgM, human IgG1, and human IgG3.

[0081] In accordance with a seventh aspect of the present invention, there is provided a monoclonal antibody that binds to an epitope on CD147 comprising the consensus sequence RVRSH, wherein the antibody is not CBL1. In a preferred embodiment, the antibody is a human antibody.

[0082] In accordance with an eighth aspect of the present invention, there is provided an isolated peptide comprising the sequence selected from the group consisting of RXRS, RXRSH, RVRSH, and RVRSH. In a preferred embodiment, the peptide is used for the generation of antibodies.

[0083] In accordance with a ninth aspect of the present invention, there is provided a human monoclonal antibody that binds to CD147.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

Discussion of the Present Invention

[0084] The pharmaceutical agent ABX-CBL was derived from the hybridoma cell line expressing the CBL1 antibody. CBL1 is a murine IgM, anti-human lymphoblastoid monoclonal antibody that was raised in Balb/c mice immunized with the T cell acute lymphoblastic leukemia cell line (T-ALL) CEM (Billing et al. "Monoclonal and heteroantibody reacting with different common antigens common to human blast cells and monocytes" *Hybridoma* 1:303-311 (1982)). Following fusion of the splenocytes and selection in HAT medium, supernatants from hybridoma-containing wells were screened by microcytotoxicity assay for reactivity with CEM cells. Hybridomas that tested positive in this assay were further screened for their ability to discriminate between resting lymphocytes and blast cells. CBL1 was selected for further study because it showed selectivity for blast cells (Billing et al. "Monoclonal and heteroantibody reacting with different common antigens common to human blast cells and monocytes" *Hybridoma* 1:303-311 (1982)). The CBL1 antibody was deposited with the ATCC as HB 8214.

[0085] The assignee of the present application, Abgenix, Inc., Fremont, Calif., acquired CBL1 in 1997 and determined that the hybridoma line deposited with the ATCC as HB 8214 was not entirely pure. Rather, it was actually a mix of two distinct hybridoma lines, one producing an IgG and one producing an IgM. Following subcloning, a pure IgM producer as well as a pure IgG producer were derived. Through a series of in vitro experiments described herein, it was demonstrated that the IgM antibody mediated the activities previously attributed to the CBL1 hybridoma. Only the IgM is biologically active in inhibition of complement mediated lysis of cells in a mixed lymphocyte reaction assay (MLR). The mechanism of inhibition is via antibody mediated complement-dependent cytotoxicity (CDC) because the inhibition is specific and complement-dependent, as discussed herein. Therefore, in connection with our work described herein, using conventional techniques, we subcloned the line to produce a cell line producing solely the IgM. Further, the HB 8214 cell line expressing the CBL1 antibody possessed a second kappa light chain (MOPC-21) which appears to have been derived from the myeloma fusion partner, a P3 myeloma cell line, that was used to prepared the original hybridoma cell line. Our subcloned hybridoma cell line possesses and expresses both light chains and the ABX-CBL antibody appears to contain both light chains. IgM antibodies generally possess a pentameric structure, where five heavy and light chain dimers are associated. With the two light chains in the ABX-CBL antibody, we expect that the IgM pentameric structure of the ABX-CBL antibody contains both light chains in various ratios of light chains to form pentamers with homodimeric, heterodimeric, and homo- and heterodimeric combinations.

[0086] In order to manufacture the ABX-CBL antibody for use in preclinical and clinical development, we utilized hollow fiber cell culture technology through contract manufacturing with Goodwin Biotechnology, Plantation, Fla. The growth medium is a serum free formulation HYBRIDOMA-SFM supplied by Gibco Life Technologies.

[0087] The stability of the Master Cell Bank (MCB) of ABX-CBL was determined by single cell subcloning. Cells

were subcloned showing >95% stability for the single cell colony producers. The ABX-CBL MCB also showed stable antibody production for more than 130 generations in culture. The manufacturing process in hollow fiber bioreactors is an approximately 40 day growth process that is equivalent to approximately 130 generations.

[0088] Primary purification of the monoclonal antibody from the cell culture supernatant is performed using Protein A affinity chromatography. Incubation at low pH following elution is performed as a viral inactivation step. The material is further purified by anion exchange chromatography. This provides for residual protein A and DNA removal. The final step in the purification process is a filtration of the material to provide additional viral removal.

[0089] The formulated bulk drug substance is stored at 2-8° C. prior to vialing. Using aseptic techniques, the antibody is filled in liquid form from the bulk containers into 5 mL glass vials. The vials are stored and shipped at 2-8° C. ABX-CBL is a murine IgM, anti-human lymphoblastoid monoclonal antibody raised to a T-ALL (Acute Lymphoblastic Leukemia) cell line (CEM). ABX-CBL is formulated in 20 mM sodium citrate and 120 mM sodium chloride at a pH of 6.0.

[0090] As used herein, the term "ABX-CBL" is used to refer to the purified and reactive IgM antibody derived from the original cell line deposited with the ATCC as HB 8214. The sequence of the ABX-CBL heavy and light chains are discussed above and presented as SEQ ID NO.: 18 and SEQ ID NO.: 19, respectively.

[0091] We have now demonstrated that the active agent of the CBL1 antibody and ABX-CBL binds to the CD147 antigen as expressed on certain cells, such as T-cells, B-cells, and/or monocytes. Accordingly, it is expected that the CD147 antigen, can be utilized as a target for the treatment of a variety of diseases. Since the CBL1 antibody has been effective in patients in the treatment of the diseases mentioned above, and based upon the results discussed herein, it is expected that additional CD147 based therapeutics will be similarly effective. Thus, in accordance with the present invention, we have discovered that the molecule CD147 as expressed on certain cells, such as T-cells, B-cells, and/or monocytes, can be utilized for the treatment of a variety of diseases. In particular, we have demonstrated that antibodies that bind to CD147 and that result in the killing of such cells, for example, through the activation of complement, is efficacious in the treatment of diseases. Diseases in which such treatment appears efficacious include, without limitation: graft versus host disease (GVHD), organ transplant rejection diseases (including, without limitation, renal transplant, corneal transplant, and others), cancers (including, without limitation, cancers of the blood (i.e., leukemias and lymphomas), and pancreatic), autoimmune diseases, inflammatory diseases, and others.

[0092] As was mentioned above, CBL1 had not previously been indicated to bind to CD147. Further, the particular epitope or antigen to which the CBL1 antibody bound was unknown or at least relatively uncharacterized. Thus, because of the apparent safety and therapeutic efficacy of the CBL1 antibody, we were interested in determining the precise antigen or epitope to which the CBL1 and our ABX-CBL antibody bound. Further, we were interested in

further understanding the manner in which the CBL1 antibody was efficacious, particularly in connection with the treatment of GVHD.

[0093] By way of reference, the hybridoma line deposited with the ATCC as HB 8214 was not entirely pure. The line produced an IgG antibody and an IgM antibody. Only the IgM is biologically active in inhibition of complement mediated lysis of cells in a mixed lymphocyte reaction assay (MLR). The mechanism of inhibition is via antibody mediated complement-dependent cytotoxicity (CDC) because the inhibition is specific and complement-dependent, as discussed herein. Therefore, in connection with our work described herein, we subcloned the line to produce a cell line producing solely the IgM. Further, the HB 8214 cell line expressing the CBL1 antibody possessed a second kappa light chain (MOPC-21) which appears to have been derived from the myeloma fusion partner, a P3 myeloma cell line, that was used to prepare the original hybridoma cell line. Our subcloned hybridoma cell line possesses and expresses both light chains and the ABX-CBL antibody appears to contain both light chains. IgM antibodies generally possess a pentameric structure, where five heavy and light chain dimers are associated. With the two light chains in the ABX-CBL antibody, we expect that the IgM pentameric structure of the ABX-CBL antibody contains both light chains in various ratios of light chains to form pentamers with homodimeric, heterodimeric, and homo- and heterodimeric combinations.

[0094] The role of the MOPC-21 light chain in CBL1 and ABX-CBL binding was unknown. In connection with our work, we endeavored to clarify the role of the MOPC-21 light chain through, for example, preparation of hybridoma subclones that express only the ABX-CBL light chain or the MOPC-21 light chain. One approach that we utilized was to fuse the ABX-CBL hybridoma with a mouse myeloma cell line to achieve light chain shuffling. Upon generation of hybridomas expressing only the MOPC-21 light chain or the ABX-CBL light chain, we were able to conduct certain characterizations to distinguish the role of the two light chains in ABX-CBL binding.

Definitions

[0095] Unless otherwise defined, scientific and technical terms used in connection with the present invention shall have the meanings that are commonly understood by those of ordinary skill in the art. Further, unless otherwise required by context, singular terms shall include pluralities and plural terms shall include the singular. Generally, nomenclatures utilized in connection with, and techniques of, cell and tissue culture, molecular biology, and protein and oligo- or polynucleotide chemistry and hybridization described herein are those well known and commonly used in the art. Standard techniques are used for recombinant DNA, oligonucleotide synthesis, and tissue culture and transformation (e.g., electroporation, lipofection, etc.). Enzymatic reactions and purification techniques are performed according to manufacturer's specifications or as commonly accomplished in the art or as described herein. The foregoing techniques and procedures are generally performed according to conventional methods well known in the art and as described in various general and more specific references that are cited and discussed throughout the present specification. See e.g., Sambrook et al. *Molecular Cloning. A Laboratory Manual*

(2d ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1989)), which is incorporated herein by reference. The nomenclatures utilized in connection with, and the laboratory procedures and techniques of, analytical chemistry, synthetic organic chemistry, and medicinal and pharmaceutical chemistry described herein are those well known and commonly used in the art. Standard techniques are used for chemical syntheses, chemical analyses, pharmaceutical preparation, formulation, and delivery, and treatment of patients.

[0096] As utilized in accordance with the present disclosure, the following terms, unless otherwise indicated, shall be understood to have the following meanings:

[0097] The term “isolated polynucleotide” as used herein shall mean a polynucleotide of genomic, cDNA, or synthetic origin or some combination thereof, which by virtue of its origin the “isolated polynucleotide” (1) is not associated with all or a portion of a polynucleotide in which the “isolated polynucleotide” is found in nature, (2) is operably linked to a polynucleotide which it is not linked to in nature, or (3) does not occur in nature as part of a larger sequence.

[0098] The term “isolated protein” referred to herein means a protein of cDNA, recombinant RNA, or synthetic origin or some combination thereof, which by virtue of its origin, or source of derivation, the “isolated protein” (1) is not associated with proteins found in nature, (2) is free of other proteins from the same source, e.g. free of murine proteins, (3) is expressed by a cell from a different species, or (4) does not occur in nature.

[0099] The term “polypeptide” is used herein as a generic term to refer to native protein, fragments, or analogs of a polypeptide sequence. Hence, native protein, fragments, and analogs are species of the polypeptide genus.

[0100] The term “naturally-occurring” as used herein as applied to an object refers to the fact that an object can be found in nature. For example, a polypeptide or polynucleotide sequence that is present in an organism (including viruses) that can be isolated from a source in nature and which has not been intentionally modified by man in the laboratory or otherwise is naturally-occurring.

[0101] The term “operably linked” as used herein refers to positions of components so described are in a relationship permitting them to function in their intended manner. A control sequence “operably linked” to a coding sequence is ligated in such a way that expression of the coding sequence is achieved under conditions compatible with the control sequences.

[0102] The term “control sequence” as used herein refers to polynucleotide sequences that are necessary to effect the expression and processing of coding sequences to which they are ligated. The nature of such control sequences differs depending upon the host organism; in prokaryotes, such control sequences generally include promoter, ribosomal binding site, and transcription termination sequence; in eukaryotes, generally, such control sequences include promoters and transcription termination sequence. The term “control sequences” is intended to include, at a minimum, all components whose presence is essential for expression and processing, and can also include additional components whose presence is advantageous, for example, leader sequences and fusion partner sequences.

[0103] The term “polynucleotide” as referred to herein means a polymeric form of nucleotides of at least 10 bases in length, either ribonucleotides or deoxynucleotides or a modified form of either type of nucleotide. The term includes single and double stranded forms of DNA.

[0104] The term “oligonucleotide” referred to herein includes naturally occurring, and modified nucleotides linked together by naturally occurring, and non-naturally occurring oligonucleotide linkages. Oligonucleotides are a polynucleotide subset generally comprising a length of 200 bases or fewer. Preferably oligonucleotides are 10 to 60 bases in length and most preferably 12, 13, 14, 15, 16, 17, 18, 19, or 20 to 40 bases in length. Oligonucleotides are usually single stranded, e.g. for probes; although oligonucleotides may be double stranded, e.g. for use in the construction of a gene mutant. Oligonucleotides of the invention can be either sense or antisense oligonucleotides.

[0105] The term “naturally occurring nucleotides” referred to herein includes deoxyribonucleotides and ribonucleotides. The term “modified nucleotides” referred to herein includes nucleotides with modified or substituted sugar groups and the like. The term “oligonucleotide linkages” referred to herein includes oligonucleotide linkages such as phosphorothioate, phosphorodithioate, phosphoroselenoate, phosphorodiselenoate, phosphoroanilothioate, phosphoraniladate, phosphoramidate, and the like. See e.g., LaPlanche et al. *Nucl. Acids Res.* 14:9081 (1986); Stec et al. *J. Am. Chem. Soc.* 106:6077 (1984); Stein et al. *Nucl. Acids Res.* 16:3209 (1988); Zon et al. *Anti-Cancer Drug Design* 6:539 (1991); Zon et al. *Oligonucleotides and Analogues: A Practical Approach*, pp. 87-108 (F. Eckstein, Ed., Oxford University Press, Oxford England (1991)); Stec et al. U.S. Pat. No. 5,151,510; Uhlmann and Peyman *Chemical Reviews* 90:543 (1990), the disclosures of which are hereby incorporated by reference. An oligonucleotide can include a label for detection, if desired.

[0106] The term “selectively hybridize” referred to herein means to detectably and specifically bind. Polynucleotides, oligonucleotides and fragments thereof in accordance with the invention selectively hybridize to nucleic acid strands under hybridization and wash conditions that minimize appreciable amounts of detectable binding to nonspecific nucleic acids. High stringency conditions can be used to achieve selective hybridization conditions as known in the art and discussed herein. Generally, the nucleic acid sequence homology between the polynucleotides, oligonucleotides, and fragments of the invention and a nucleic acid sequence of interest will be at least 80%, and more typically with preferably increasing homologies of at least 85%, 90%, 95%, 99%, and 100%. Two amino acid sequences are homologous if there is a partial or complete identity between their sequences. For example, 85% homology means that 85% of the amino acids are identical when the two sequences are aligned for maximum matching. Gaps (in either of the two sequences being matched) are allowed in maximizing matching; gap lengths of 5 or less are preferred with 2 or less being more preferred. Alternatively and preferably, two protein sequences (or polypeptide sequences derived from them of at least 30 amino acids in length) are homologous, as this term is used herein, if they have an alignment score of at more than 5 (in standard deviation units) using the program ALIGN with the mutation data matrix and a gap penalty of 6 or greater. See Dayhoff,

M. O., in *Atlas of Protein Sequence and Structure*, pp. 101-110 (Volume 5, National Biomedical Research Foundation (1972)) and Supplement 2 to this volume, pp. 1-10. The two sequences or parts thereof are more preferably homologous if their amino acids are greater than or equal to 50% identical when optimally aligned using the ALIGN program. The term "corresponds to" is used herein to mean that a polynucleotide sequence is homologous (i.e., is identical, not strictly evolutionarily related) to all or a portion of a reference polynucleotide sequence, or that a polypeptide sequence is identical to a reference polypeptide sequence. In contradistinction, the term "complementary to" is used herein to mean that the complementary sequence is homologous to all or a portion of a reference polynucleotide sequence. For illustration, the nucleotide sequence "TATAC" corresponds to a reference sequence "TATAC" and is complementary to a reference sequence "GTATA".

[0107] The following terms are used to describe the sequence relationships between two or more polynucleotide or amino acid sequences: "reference sequence", "comparison window", "sequence identity", "percentage of sequence identity", and "substantial identity". A "reference sequence" is a defined sequence used as a basis for a sequence comparison; a reference sequence may be a subset of a larger sequence, for example, as a segment of a full-length cDNA or gene sequence given in a sequence listing or may comprise a complete cDNA or gene sequence. Generally, a reference sequence is at least 18 nucleotides or 6 amino acids in length, frequently at least 24 nucleotides or 8 amino acids in length, and often at least 48 nucleotides or 16 amino acids in length. Since two polynucleotides or amino acid sequences may each (1) comprise a sequence (i.e., a portion of the complete polynucleotide or amino acid sequence) that is similar between the two molecules, and (2) may further comprise a sequence that is divergent between the two polynucleotides or amino acid sequences, sequence comparisons between two (or more) molecules are typically performed by comparing sequences of the two molecules over a "comparison window" to identify and compare local regions of sequence similarity. A "comparison window", as used herein, refers to a conceptual segment of at least 18 contiguous nucleotide positions or 6 amino acids wherein a polynucleotide sequence or amino acid sequence may be compared to a reference sequence of at least 18 contiguous nucleotides or 6 amino acid sequences and wherein the portion of the polynucleotide sequence in the comparison window may comprise additions, deletions, substitutions, and the like (i.e., gaps) of 20 percent or less as compared to the reference sequence (which does not comprise additions or deletions) for optimal alignment of the two sequences. Optimal alignment of sequences for aligning a comparison window may be conducted by the local homology algorithm of Smith and Waterman *Adv. Appl. Math.* 2:482 (1981), by the homology alignment algorithm of Needleman and Wunsch *J. Mol. Biol.* 48:443 (1970), by the search for similarity method of Pearson and Lipman *Proc. Natl. Acad. Sci. (U.S.A.)* 85:2444 (1988), by computerized implementations of these algorithms (GAP, BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics Software Package Release 7.0, (Genetics Computer Group, 575 Science Dr., Madison, Wis.), Geneworks, or MacVector software packages), or by inspection, and the best alignment (i.e., resulting in the highest percentage of homology over the comparison window) generated by the various methods is selected.

[0108] The term "sequence identity" means that two polynucleotide or amino acid sequences are identical (i.e., on a nucleotide-by-nucleotide or residue-by-residue basis) over the comparison window. The term "percentage of sequence identity" is calculated by comparing two optimally aligned sequences over the window of comparison, determining the number of positions at which the identical nucleic acid base (e.g., A, T, C, G, U, or I) or residue occurs in both sequences to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the comparison window (i.e., the window size), and multiplying the result by 100 to yield the percentage of sequence identity. The terms "substantial identity" as used herein denotes a characteristic of a polynucleotide or amino acid sequence, wherein the polynucleotide or amino acid comprises a sequence that has at least 85 percent sequence identity, preferably at least 90 to 95 percent sequence identity, more usually at least 99 percent sequence identity as compared to a reference sequence over a comparison window of at least 18 nucleotide (6 amino acid) positions, frequently over a window of at least 24-48 nucleotide (8-16 amino acid) positions, wherein the percentage of sequence identity is calculated by comparing the reference sequence to the sequence which may include deletions or additions which total 20 percent or less of the reference sequence over the comparison window. The reference sequence may be a subset of a larger sequence.

[0109] As used herein, the twenty conventional amino acids and their abbreviations follow conventional usage. See *Immunology—A Synthesis* (2nd Edition, E. S. Golub and D. R. Gren, Eds., Sinauer Associates, Sunderland, Mass. (1991)), which is incorporated herein by reference. Stereoisomers (e.g., D-amino acids) of the twenty conventional amino acids, unnatural amino acids such as α -, α -disubstituted amino acids, N-alkyl amino acids, lactic acid, and other unconventional amino acids may also be suitable components for polypeptides of the present invention. Examples of unconventional amino acids include: 4-hydroxyproline, γ -carboxyglutamate, ϵ -N,N,N-trimethyllysine, ϵ -N-acetyllysine, O-phosphoserine, N-acetyls erine, N-formylmethionine, 3-methylhistidine, 5-hydroxylysine, σ -N-methylarginine, and other similar amino acids and imino acids (e.g., 4-hydroxyproline). In the polypeptide notation used herein, the lefthand direction is the amino terminal direction and the righthand direction is the carboxy-terminal direction, in accordance with standard usage and convention.

[0110] Similarly, unless specified otherwise, the lefthand end of single-stranded polynucleotide sequences is the 5' end; the lefthand direction of double-stranded polynucleotide sequences is referred to as the 5' direction. The direction of 5' to 3' addition of nascent RNA transcripts is referred to as the transcription direction; sequence regions on the DNA strand having the same sequence as the RNA and which are 5' to the 5' end of the RNA transcript are referred to as "upstream sequences"; sequence regions on the DNA strand having the same sequence as the RNA and which are 3' to the 3' end of the RNA transcript are referred to as "downstream sequences".

[0111] As applied to polypeptides, the term "substantial identity" means that two peptide sequences, when optimally aligned, such as by the programs GAP or BESTFIT using default gap weights, share at least 80 percent sequence

identity, preferably at least 90 percent sequence identity, more preferably at least 95 percent sequence identity, and most preferably at least 99 percent sequence identity. Preferably, residue positions which are not identical differ by conservative amino acid substitutions. Conservative amino acid substitutions refer to the interchangeability of residues having similar side chains. For example, a group of amino acids having aliphatic side chains is glycine, alanine, valine, leucine, and isoleucine; a group of amino acids having aliphatic-hydroxyl side chains is serine and threonine; a group of amino acids having amide-containing side chains is asparagine and glutamine; a group of amino acids having aromatic side chains is phenylalanine, tyrosine, and tryptophan; a group of amino acids having basic side chains is lysine, arginine, and histidine; and a group of amino acids having sulfur-containing side chains is cysteine and methionine. Preferred conservative amino acids substitution groups are: valine-leucine-isoleucine, phenylalanine-tyrosine, lysine-arginine, alanine-valine, glutamic-aspartic, and asparagine-glutamine.

[0112] As discussed herein, minor variations in the amino acid sequences of antibodies or immunoglobulin molecules are contemplated as being encompassed by the present invention, providing that the variations in the amino acid sequence maintain at least 75%, more preferably at least 80%, 90%, 95%, and most preferably 99%. In particular, conservative amino acid replacements are contemplated. Conservative replacements are those that take place within a family of amino acids that are related in their side chains. Genetically encoded amino acids are generally divided into families: (1) acidic=aspartate, glutamate; (2) basic=lysine, arginine, histidine; (3) non-polar=alanine, valine, leucine, isoleucine, proline, phenylalanine, methionine, tryptophan; and (4) uncharged polar=glycine, asparagine, glutamine, cysteine, serine, threonine, tyrosine. More preferred families are: serine and threonine are aliphatic-hydroxy family; asparagine and glutamine are an amide-containing family; alanine, valine, leucine and isoleucine are an aliphatic family; and phenylalanine, tryptophan, and tyrosine are an aromatic family. For example, it is reasonable to expect that an isolated replacement of a leucine with an isoleucine or valine, an aspartate with a glutamate, a threonine with a serine, or a similar replacement of an amino acid with a structurally related amino acid will not have a major effect on the binding or properties of the resulting molecule, especially if the replacement does not involve an amino acid within a framework site. Whether an amino acid change results in a functional peptide can readily be determined by assaying the specific activity of the polypeptide derivative. Assays are described in detail herein. Fragments or analogs of antibodies or immunoglobulin molecules can be readily prepared by those of ordinary skill in the art. Preferred amino- and carboxy-termini of fragments or analogs occur near boundaries of functional domains. Structural and functional domains can be identified by comparison of the nucleotide and/or amino acid sequence data to public or proprietary sequence databases. Preferably, computerized comparison methods are used to identify sequence motifs or predicted protein conformation domains that occur in other proteins of known structure and/or function. Methods to identify protein sequences that fold into a known three-dimensional structure are known. Bowie et al. *Science* 253:164 (1991). Thus, the foregoing examples demonstrate that those of skill in the art can recognize sequence motifs

and structural conformations that may be used to define structural and functional domains in accordance with the invention.

[0113] Preferred amino acid substitutions are those which: (1) reduce susceptibility to proteolysis, (2) reduce susceptibility to oxidation, (3) alter binding affinity for forming protein complexes, (4) alter binding affinities, and (5) confer or modify other physicochemical or functional properties of such analogs. Analogs can include various muteins of a sequence other than the naturally-occurring peptide sequence. For example, single or multiple amino acid substitutions (preferably conservative amino acid substitutions) may be made in the naturally-occurring sequence (preferably in the portion of the polypeptide outside the domain(s) forming intermolecular contacts. A conservative amino acid substitution should not substantially change the structural characteristics of the parent sequence (e.g., a replacement amino acid should not tend to break a helix that occurs in the parent sequence, or disrupt other types of secondary structure that characterizes the parent sequence). Examples of art-recognized polypeptide secondary and tertiary structures are described in *Proteins, Structures and Molecular Principles* (Creighton, Ed., W. H. Freeman and Company, New York (1984)); *Introduction to Protein Structure* (C. Branden and J. Tooze, eds., Garland Publishing, New York, N.Y. (1991)); and Thornton et al. *Nature* 354:105 (1991), which are each incorporated herein by reference.

[0114] The term "polypeptide fragment" as used herein refers to a polypeptide that has an amino-terminal and/or carboxy-terminal deletion, but where the remaining amino acid sequence is identical to the corresponding positions in the naturally-occurring sequence deduced, for example, from a full-length cDNA sequence. Fragments typically are at least 5, 6, 8 or 10 amino acids long, preferably at least 14 amino acids long, more preferably at least 20 amino acids long, usually at least 50 amino acids long, and even more preferably at least 70 amino acids long. The term "analog" as used herein refers to polypeptides which are comprised of a segment of at least 25 amino acids that has substantial identity to a portion of a deduced amino acid sequence and which has at least one of the following properties: (1) specific binding to a CD147, under suitable binding conditions, (2) ability to modify CD147's binding to its ligand or receptor, or (3) ability to kill or inhibit growth of CD147 expressing cells in vitro or in vivo. Typically, polypeptide analogs comprise a conservative amino acid substitution (or addition or deletion) with respect to the naturally-occurring sequence. Analogs typically are at least 20 amino acids long, preferably at least 50 amino acids long or longer, and can often be as long as a full-length naturally-occurring polypeptide.

[0115] Peptide analogs are commonly used in the pharmaceutical industry as non-peptide drugs with properties analogous to those of the template peptide. These types of non-peptide compound are termed "peptide mimetics" or "peptidomimetics". Fauchere, *J. Adv. Drug Res.* 15:29 (1986); Veber and Freidinger *TINS* p.392 (1985); and Evans et al. *J. Med. Chem.* 30:1229 (1987), which are incorporated herein by reference. Such compounds are often developed with the aid of computerized molecular modeling. Peptide mimetics that are structurally similar to therapeutically useful peptides may be used to produce an equivalent therapeutic or prophylactic effect. Generally, peptidomimet-

ics are structurally similar to a paradigm polypeptide (i.e., a polypeptide that has a biochemical property or pharmacological activity), such as human antibody, but have one or more peptide linkages optionally replaced by a linkage selected from the group consisting of: $-\text{CH}_2\text{NH}-$, $-\text{CH}_2\text{S}-$, $-\text{CH}_2-\text{CH}_2-$, $-\text{CH}=\text{CH}-$ (cis and trans), $-\text{COCH}_2-$, $-\text{CH}(\text{OH})\text{CH}_2-$, and $-\text{CH}_2\text{SO}-$, by methods well known in the art. Systematic substitution of one or more amino acids of a consensus sequence with a D-amino acid of the same type (e.g., D-lysine in place of L-lysine) may be used to generate more stable peptides. In addition, constrained peptides comprising a consensus sequence or a substantially identical consensus sequence variation may be generated by methods known in the art (Rizo and Gierasch *Ann. Rev. Biochem.* 61:387 (1992), incorporated herein by reference); for example, by adding internal cysteine residues capable of forming intramolecular disulfide bridges which cyclize the peptide.

[0116] "Antibody" or "antibody peptide(s)" refer to an intact antibody, or a binding fragment thereof that competes with the intact antibody for specific binding. Binding fragments are produced by recombinant DNA techniques, or by enzymatic or chemical cleavage of intact antibodies. Binding fragments include Fab, Fab', F(ab')₂, Fv, and single-chain antibodies. An antibody other than a "bispecific" or "bifunctional" antibody is understood to have each of its binding sites identical. An antibody substantially inhibits adhesion of a receptor to a counterreceptor when an excess of antibody reduces the quantity of receptor bound to counterreceptor by at least about 20%, 40%, 60% or 80%, and more usually greater than about 85% (as measured in an *in vitro* competitive binding assay).

[0117] The term "epitope" includes any protein determinant capable of specific binding to an immunoglobulin or T-cell receptor. Epitopic determinants usually consist of chemically active surface groupings of molecules such as amino acids, sugar, or other carbohydrate side chains and usually have specific three dimensional structural characteristics, as well as specific charge characteristics. An antibody is said to specifically bind an antigen when the dissociation constant is $\leq 1 \mu\text{M}$, preferably $\leq 100 \text{ nM}$ and most preferably $\leq 10 \text{ nM}$.

[0118] The term "agent" is used herein to denote a chemical compound, a mixture of chemical compounds, a biological macromolecule, or an extract made from biological materials.

[0119] As used herein, the terms "label" or "labeled" refers to incorporation of a detectable marker, e.g., by incorporation of a radiolabeled amino acid or attachment to a polypeptide of biotinylated moieties that can be detected by marked avidin (e.g., streptavidin containing a fluorescent marker or enzymatic activity that can be detected by optical or calorimetric methods). In certain situations, the label or marker can also be therapeutic. Various methods of labeling polypeptides and glycoproteins are known in the art and may be used. Examples of labels for polypeptides include, but are not limited to, the following: radioisotopes or radionuclides (e.g., ³H, ¹⁴C, ¹⁵N, ³⁵S, ⁹⁰Y, ⁹⁹Tc, ¹¹¹In, ¹²⁵I, ¹³¹I), fluorescent labels (e.g., FITC, rhodamine, lanthanide phosphors), enzymatic labels (e.g., horseradish peroxidase, β -galactosidase, luciferase, alkaline phosphatase), chemiluminescent, biotinyl groups, predetermined polypeptide epitopes recog-

nized by a secondary reporter (e.g., leucine zipper pair sequences, binding sites for secondary antibodies, metal binding domains, epitope tags). In some embodiments, labels are attached by spacer arms of various lengths to reduce potential steric hindrance.

[0120] The term "pharmaceutical agent or drug" as used herein refers to a chemical compound or composition capable of inducing a desired therapeutic effect when properly administered to a patient. Other chemistry terms herein are used according to conventional usage in the art, as exemplified by *The McGraw-Hill Dictionary of Chemical Terms* (Parker, S., Ed., McGraw-Hill, San Francisco (1985)), incorporated herein by reference).

[0121] The term "substantially non-toxic to resting T-cells and resting B-cells" as used herein means, preferably, that the antibody in the presence of complement at at least a 2-fold lower level of depletion of resting cells occurs than the level of depletion of activated T- and B-cells. More preferably, there is at least a 5-fold lower level of cell depletion of resting cells compared to the level of depletion of activated cells. And, most preferably, there would be no detectable depletion of resting cells.

ABX-CBL Antigen Identification and Characterization

[0122] We undertook two primary approaches to the identification and characterization of the antigen to which the ABX-CBL antibody bound (i) an immunoaffinity purification approach and (ii) a classical protein purification approach.

[0123] Immunoaffinity Purification

[0124] We investigated immunoaffinity purification of the antigen to which the CBL1 antibody bound. The antigen to which the CBL1 antibody bound appeared to be highly expressed on CEM cells which is a T lymphoblastoid cell line derived by Foley et al. *Cancer* 18:522-529 (1965) and available from the ATCC, Rockville, Md. (ATCC No. CCL-119). Immunoaffinity purification using the native ABX-CBL antibody was frustrated by the fact that the ABX-CBL antibody is an IgM antibody having a pentameric structure and prone to nonspecific interactions *in vitro*. Therefore, we prepared human IgG2 antibodies against CEM cells and tested for competition with the ABX-CBL antibody in binding assays with CEM cells. Such human antibodies were prepared in accordance with Mendez et al. *Nature Genetics* 15:146-156 (1997) and U.S. patent application Ser. No. 08/759,620, filed Dec. 3, 1996, the disclosures of which are hereby incorporated by reference herein in their entirety, through the immunization of XenoMouseTM animals with CEM cells, followed by fusions, and screening of the resulting hybridoma supernatants against CEM cells and in FACS competition assays with the ABX-CBL antibody. In the FACS competition assays, inhibition of the binding of ABX-CBL antibodies, labeled with FITC, to CEM cells was analyzed, both alone and in the presence of hybridoma supernatants containing human antibodies reactive with the CEM cells.

[0125] Four hybridoma clones were isolated and determined, in this manner, to be that were highly competitive with the ABX-CBL antibody in binding to the CEM cells. One hybridoma clone, designated 2.6.1, was selected for further analysis. We generated ascites to each of the hybridomas, including the 2.6.1 hybridoma, in SCID mice and

purified the 2.6.1 antibody using a Protein A affinity purification process using standard conditions. From the purified 2.6.1 antibody, we prepared an immunoaffinity column. To prepare the column, the purified 2.6.1 antibody was conjugated to CNBr activated Sepharose-4B, according to the manufacturer's specifications. Approximately 8.4 mg of the antibody was conjugated to about 2.0 g of the activated Sepharose. We passed cell lysates of CEM cells through the column and eluted the components that bound. The elution product was analyzed by Western blotting and probing with both the ABX-CBL antibody and the 2.6.1 antibody. Based upon preliminary data, the 2.6.1 antibody bound most intensely to a molecule or molecules contained within a diffuse band from about 45-55 KD, while the ABX-CBL antibody showed binding with a low intensity to a similar diffuse band from about 45-55 KD. Through use of preparative gel electrophoresis and electroblotting techniques, we isolated a portion of the 45-55 KD band and obtained a partial amino acid sequence of the molecule (35/40 residues). The resulting sequence information was analyzed through a protein database search (Protein Identification Resource (PIR) R47.0, December 1995) and the sequence comparison data indicated that the molecule was CD147.

[0126] Protein Purification and Sequencing

[0127] In connection with our work related to the characterization of the antigen to which the ABX-CBL antibody bound, we saw significant ABX-CBL binding on Western blots to molecules localized in relatively sharp bands at 35 KD and 62 KD. The intensity of this 35 KD band appeared to vary from prep to prep, depending on culture age and other conditions not completely understood. Therefore, we initially purified the 62 KD material. Because the N-terminus was blocked, we cleaved the protein with CNBr and sequenced two of the peptides that resulted from the cleavage. The resulting sequence information was analyzed through a protein database search (Protein Identification Resource (PIR) R47.0, December 1995) and the sequence comparison data indicated that the molecule was heterogeneous ribonuclear protein k (hnRNP-k). Such molecule is an intracellular component, and, accordingly, does not conform to the observations that the ABX-CBL antibody appeared to recognize an extracellular component. Nevertheless, the identification of this molecule may be useful in connection with further understanding of the binding of ABX-CBL to CD147, for example in connection with epitope elucidation.

[0128] Characterization of the 35 KD band can also be undertaken for similar reasons. In such an approach, the 35 KD molecule can be purified in a similar manner to that utilized in connection with the 62 KD band mentioned above. The purified material from the 35 KD band can be characterized to further understand any potential structural differences between material contained in the 45-55 KD CD147 band. The material contained in the 35 KD band can be sequenced to either demonstrate that the material is CD147 or to determine epitopic information related to ABX-CBL's binding to CD147.

[0129] Further Elucidation of CD147 Binding and Epitopic Analysis of ABX-CBL

[0130] As was discussed above, another area of exploration is in connection with the elucidation of the binding of the ABX-CBL antibody to the CD147 molecule. Because of the safety and efficacy of the ABX-CBL antibody, we expect

that molecules, particularly antibodies, that mimic the binding of the ABX-CBL antibody to CD147 should possess a similar safety profile. Thus, in order to further understand the binding of the ABX-CBL antibody to CD147, we have undertaken, or designed, experiments in order to elucidate the same. Our experiments include (i) cloning of CD147 and expression in eukaryotic (COS) cells, (ii) expression in prokaryotic (*E. coli*) cells, and (iii) screening of random peptide libraries utilizing phage display techniques.

[0131] Cloning of CD147 and Expression in COS Cells

[0132] We cloned CD147 cDNA from a Jurkat library (Stratagene), prepared constructs for transfection, and transfected COS cells with the CD147 cDNA. Transfected cells were analyzed for expression of CD147 utilizing FACS analysis and Western blotting in connection with the ABX-CBL antibody, the 2.6.1 antibody, and the Pharmingen antibody mentioned above. COS cells transfected with CD147 cDNA showed binding to each of the antibodies in each of the FACS and Western blot analyses. In contrast, COS cells transfected with control vectors were negative for binding with each of the 2.6.1 and ABX-CBL antibodies. With respect to the Pharmingen antibody, certain background staining was observed in cells transfected with control vectors on FACS and no binding on Western blot analysis. The transfected cells showed significant binding over background on FACS and were positive on Western blot analysis. Our results confirm that the ABX-CBL and the 2.6.1 antibodies bind to CD147.

[0133] Expression of CD147 in *E. Coli* Cells

[0134] Utilizing a slightly modified vector, we also transfected *E. coli* cells with the CD147 cDNA. The *E. coli* cells so transfected were capable of expression of the CD147 molecule as evidenced by Western blotting analysis of each of the ABX-CBL, 2.6.1, and Pharmingen antibodies. Since the prokaryotic *E. coli* cells should not glycosylate the expressed CD147, it was expected that the molecular weight of the CD147 expressed by the *E. coli* should closely approximate the predicted, unglycosylated molecular weight of CD147 of about 27 KD. Indeed, in each case, binding of the three antibodies on Western blot analysis was observed to a band between about 27 and 30 KD.

[0135] This data further confirms that the ABX-CBL and the 2.6.1 antibodies bind to CD147. Further, the evidence indicates that ABX-CBL binding to CD147 is not directly based on carbohydrate binding, i.e., that ABX-CBL does not bind directly to a carbohydrate epitope on CD147. Such data, however, does not eliminate the possibility that binding to CD147 is influenced by the presence of carbohydrate or glycosylation.

[0136] Screening Utilizing Phage Display

[0137] In order to further elucidate the binding of the ABX-CBL antibody to CD147, we undertook phage display experiment. Such experiments were conducted through panning a phage library expressing random peptides for binding with the ABX-CBL and 2.6.1 antibodies to determine if we could isolate peptides that bound. If successful, certain epitope information can be gleaned from the peptides that bind.

[0138] In general, the phage libraries expressing random peptides were purchased from New England Biolabs (7-mer

and 12-mer libraries, Ph.D.-7 Peptide 7-mer Library Kit and Ph.D.-12 Peptide 12-mer Library Kit, respectively) based on a bacteriophage M13 system. The 7-mer library represents a diversity of approximately 2.0×10^9 independent clones, which represents most, if not all, of the $20^7 = 1.28 \times 10^9$ possible 7-mer sequences. The 12-mer library contains approximately 1.9×10^{10} independent clones and represents only a very small sampling of the potential sequence space of $20^{12} = 4.1 \times 10^{15}$ 12-mer sequences. Each of 7-mer and 12-mer libraries were panned or screened in accordance with the manufacturer's recommendations in which plates were coated with an antibody to capture the appropriate antibody (goat anti-human IgG Fc for the 2.6.1 antibody and goat anti-mouse κ chain for the ABX-CBL antibody) followed by washing. Bound phage were eluted with 0.2 M glycine-HCl, pH 2.2. After 3 rounds of selection/amplification at constant stringency (0.5% Tween), through use of DNA sequencing, we characterized a total of 5 clones from the 7-mer library and 6 clones from the 12-mer library reactive with the ABX-CBL antibody and a total of 6 clones from each of the 7-mer and 12-mer libraries reactive with the 2.6.1 antibody. Reactivity of the peptides was determined by ELISA. For an additional discussion of epitope analysis of peptides see also Scott, J. K. and Smith, G. P. *Science* 249:386-390 (1990); Cwirla et al. *PNAS USA* 87:6378-6382 (1990); Felici et al. *J. Mol. Biol.* 222:301-310 (1991), and Kuwabara et al. *Nature Biotechnology* 15:74-78 (1997).

[0139] No consensus sequence was readily apparent for reactivity of the 2.6.1 antibody with CD147. However, sequence alignment of the characterized 7-mer and 12-mer sequences against the amino acid sequence of CD147 yielded a number of matches for a single sequence within CD147 from residue number 177 through residue number 188 (ITLRVRSH (SEQ ID NO: 1)). In particular, each of the 7-mers contained sequence matches (represented by *) to 3 or more residues within this sequence of CD147:

[0140] Further, 4 of the 12-mers contained sequence matches (represented by *) to 3 or more residues within this sequence of CD147, with 4 matches for 12-mer peptide number 1 and for 6 matches of 12-mer peptide number 2:

7-mer sequences				
1.	EE	RLR S	Y	(SEQ ID NO:2)
2.	YE	RVR W	Y	(SEQ ID NO:3)
3.	EE	RLR S	Y	(SEQ ID NO:4)
4.	AE	RIR S	I	(SEQ ID NO:5)
5.	EE	RLR S	Y	(SEQ ID NO:6)

```

12-mer sequences
      * * * *
1.   TVHGDL  RLR S   LP   (SEQ ID NO:7)

      * * * * *
2.   TNDIGL  ROR S   HS   (SEQ ID NO:8)

```

-continued

* * *

3. SPLLDGQ RER S Y (SEQ ID NO:9)

* * *

4. YDLPM RSR S YPG (SEQ ID NO:10)

[0141] These results indicate a consensus sequence of RXRS (SEQ ID NO:11) that is present in 10 of the sequenced clones. Accordingly, we had a synthetic peptide prepared (AnaSpec Incorporated, San Jose, Calif.) which spanned residues 169-183 of CD147 with the following sequence (with —OH representing carboxy terminus):

KGSDQAIITLVRSH-OH (SEQ ID NO:12)
| |
169 184

[0142] Below, the amino acid sequence of CD147 is provided with the 15-mer peptide's sequence indicated by double underlining and the RXRSH (SEQ ID NO:13) consensus sequence indicated in bold. In addition, putative N-linked glycosylation sites of CD147 are shown as underlined and italics:

CD147 Sequence (SEQ ID NO:14)

MAAALFVLLGFALLGTHGASGAAGTVFTTVEDLGSKILLTCSLNDSATEV

TGHRWLKGGVVLKEDALPGQKTEFKVDSDDQWGEYSYCVFLPEPMGTANIQ

LHGPPRVKAVKSSSEHINEGETAMLVCKSESVPPVTDWANYKITDSEDKAL

MNGSESRSFFVSSSQGRSELHIENLNMEADPGQYRCNGTSSKGSDQAIITLR

VRSLLAALWPLFLGIVAEVLVLVTIIFYEKRRKPEDVLDDDDAGSAPLKS

SGOHONDKGKNVRORNSS

[0143] The 15-mer peptide was assayed using ELISA and it was determined that the ABX-CBL antibody specifically bound to the peptide. Further, neither the 2.6.1 antibody nor a control murine IgM antibody bound to the peptide. However, based on a competition study between the CD147 antigen and the 15-mer peptide, the ABX-CBL antibody's binding to the 15-mer peptide can only be measured when the 15-mer peptide is coated on plates and not when the peptide is in solution. Indeed, in competition experiments in which the ABX-CBL antibody is bound to either the peptide or the CD147 antigen coated to plates, the ABX-CBL antibody is not removed or replaced by the peptide in solution even at high concentrations. Nevertheless, the binding of the ABX-CBL antibody to the 15-mer peptide can be specifically competed by the CD147 antigen and positive phage preparations mentioned above but not with non-specific antigen (i.e., L-Selectin isolated from cell membrane or human plasma) or the negative phage preparations mentioned above. Similarly, the binding of the ABX-CBL antibody to the CD147 antigen can be specifically competed by positive phage preparations as compared to negative phage preparation in competition assays using preincubation.

[0144] These results indicate that while the sequence within CD147 that contains the consensus sequence RXRSH is important to the binding of the ABX-CBL antibody to

CD147, it does not fully explain ABX-CBL's binding to CD147. Indeed, the data also suggests that the consensus sequence contained either in the 15-mer peptide when bound to the plate or the reactive phage materials when tethered to the phage coat protein binds more tightly to the ABX-CBL antibody than does the free peptide in solution. Taken together, while not wishing to bound to any particular theory or mode of operation, it is possible that CD147 possesses certain conformations that are not well mimicked in the 15-mer peptide in solution. Nevertheless, the above epitopic information is important to understanding the manner in which the ABX-CBL antibody binds to CD147 and to producing other candidate molecules against CD147 as a therapeutic target.

[0145] It is interesting to note that in addition to the results above in connection with the presence of the RXRSH consensus sequence within CD147, we also looked for the presence of the consensus sequence within the hn-RNP-k protein to which ABX-CBL also appears to bind. Such analyses were conducted by sequence alignment against the phage derived peptides discussed above. Two sequences were found which possessed statistically interesting matches:

[0146] First, there was a match (indicated by *) of 5 amino acids with the 7-mer peptide number 4:

```

      *  **  **
PE  RIL  SI          (SEQ ID NO:15)
|
84

```

[0147] Second, there was a match (indicated by *) of 5 amino acids with the 12-mer peptide number 1:

```

      *  *  *  *
GGS  RAR  NLP        (SEQ ID NO:16)
|          |
300        306

```

[0148] The amino acid sequence of the hn-RNP-k protein is provided below with such sequences indicated by double underlining. In addition, a number of RXR sequence motifs are present in the hn-RNP-k protein's sequence which are also indicated by underlining:

```

hn-RNP-k Protein Sequence
                                     (SEQ ID NO:17)
METEQPEETFPNTETNGEFGKRPADMEEEQAFKRSRNTDEMVELRILLQ
SKNAGAVIGKGGKNIKALRTDYNASVSPDSSGPERILSISADIETIGEI
LKKIIPTELEGLQLPSPTATSQLPLESDAVECLNYQHYKGSDFDCLELL
IHQSLAGGIIGVKGAKIKELRENTQTITKLFQECPPHSTDRVVLIGGKPD
RVVECIKIILDLISESPIKGRAQPYDPNFYDETYDYGFTMMFDDRRGRP
VGFPMPGRGGGFDMPPPGRGGRPMPSPRRDYDDMSPRGGPPPPPPGRGGRC
GSRARNLPLPPPPPPRGDLMAYDRGRPGDRYDGMVGFSADETWDSDAID
TWSPEWQMayEQGGSGYDYSAGGRGSYDGLGGPIITTQVTIPKDLAG
SIIGKGGQRIKQIRHESGASIKIDEPLGSEDRIITITGTQDQIQNAQYL
LQNSVKQYSGKIFF

```

[0149] Without wishing to be bound to any particular theory or mode of operation, it is possible that the binding of the ABX-CBL antibody to the hn-RNP-k protein is partially explained by the presence of these motifs within the protein.

[0150] Discussion of Results of Antigen Identification and Analysis

[0151] It is interesting to note that the ABX-CBL antibody appears to bind to the 45-55 KD band with less intensity than it does the 35 KD band in CEM cell lysates. However, without wishing to be bound to any particular theory or mode of operation of the ABX-CBL antibody, the 35 KD band could either represent another epitope or could be an alternative form of CD147. Indeed, as discussed above, there is evidence in the literature for alternative splicings of CD147 or differential glycosylation. See e.g., Kanekura et al. *Cell Struct Funct* 16:23-30 (1991) (Basigin); Schlo-sshauer B *Development* 113:129-140 (1991) (Neurothelin); Fadool J M & Linser P J *J Neurochem* 60:1354-136 (1993) (5A11/HT7); Nehme et al. *J Cell Biol* 120:687-694 (1993) (CE9); DeCastro et al. *J Invest Dermatol* 106:1260-1265 (1996) (EMMPRIN); Spring et al. *Eur J Immunol* 27:891-897 (1997) (Ok^a). Anecdotal evidence indicates that a 35 KD band could correspond to a singly-glycosylated form of CD147. See Kanekura et al. *Cell Struct Funct* 16:23-30 (1991). Further, it is also interesting to note that in comparisons of Western blots produced by two commercially available anti-CD147 antibodies (RDI-CBL535 (an anti-CD147 IgG2 antibody), available from RDI, Flanders, N.J., and 36901A (an anti-CD147 IgG1 antibody), available from Pharmingen, San Diego, Calif.) to the ABX-CBL and 2.6.1 antibodies indicates that each of the commercially available antibodies recognize a molecule that has a molecular weight around 35 KD and appearing similar to the 35 KD band recognized by the ABX-CBL antibody. However, the 45-55 KD diffuse band is more intense. See FIG. 1.

[0152] Based upon preliminary data, another interesting observation is that in the immunoaffinity purification mentioned above, when the effluent product from the 2.6.1 antibody was probed with the ABX-CBL antibody, the 35 KD band was no longer visible by Western blot. Rather, the ABX-CBL antibody appeared to bind to the diffuse band from 45-55 KD with relatively low intensity.

[0153] Further, our results in phage display experiments indicates that the ABX-CBL antibody and the 2.6.1 antibody bind to different epitopes. However, from our work related to the expression of CD147 in *E. coli* cells and based on the phage display work, the ABX-CBL antibody appears to recognize a protein epitope of CD147 and glycosylation, alone, does not appear responsible for ABX-CBL binding to CD147.

[0154] Nevertheless, in light of all of the foregoing, taken together, our results and data indicate that the ABX-CBL antibody does bind to the CD147 antigen. However, the ABX-CBL antibody appears to preferentially recognize a different epitope than recognized by the 2.6.1 or commercially available antibodies. Our finding that the ABX-CBL antibody binds to the CD147 antigen is indicative that a form of CD147 as expressed on particular cells is a viable therapeutic target for the treatment of disease.

Functional Understanding of the Mode of CD147 Therapy

[0155] As mentioned above, the CBL1 antibody has been used extensively in the treatment of GVHD in patients. Indeed, about a number of GVHD patients have been treated using the CBL1 antibody with a high percent success rate. Corneal and renal transplant studies have shown similar efficacy. Further, no signs of safety concerns or adverse effects have been observed. This is striking, given that, as discussed above, CD147 is so widely expressed in various tissues and cells of man. One would be concerned that an antibody to CD147 could cause a variety of adverse effects. Accordingly, we also endeavored to study the mechanism through which the CBL1 antibody operated to result in the treatment of disease, focused on models relevant to the reversal of GVHD. Understanding the mechanism could assist in elucidating why the CBL1 antibody is efficacious in patients and could also provide an understanding of how to use the antigen to which the CBL1 antibody binds, CD147, in the treatment of disease.

[0156] There are several possible explanations related to the safety and specificity of the CBL1 antibody in the treatment of disease. Without limitation, these include (i) that there is a unique role of complement mediated cell killing (complement dependent cytotoxicity, CDC), (ii) that certain cells in becoming activated become sensitive to CBL1 binding and cell killing, (iii) that there are particular protective elements in certain cellular populations that render the cells resistant to CBL1 induced CDC, (iv) that CD147 expression levels are higher in given populations of cells (which could also be relevant to CDC), and (v) that the CBL1 antibody binds to a particular form of CD147 expressed on certain cellular populations (as discussed above). Each of these roles will be discussed in additional detail below.

[0157] Complement Mediated Killing of Cells

[0158] The role of complement mediated cell killing (complement dependent cytotoxicity, CDC) in connection with the CBL1 antibody has been studied previously and we have additionally studied its role extensively.

[0159] Past Work with CBL1

[0160] The UCLA group mentioned above (see e.g., U.S. Pat. Nos. 5,330,896 and 5,643,740) provided certain evidence that the CBL1 antibody operated through killing of certain activated cell populations while the antibody did not react with non-activated cells. For example, in microcytotoxicity assays, the CBL1 antibody was disclosed to kill activated lymphocytic cells but not non-activated lymphocytic or other normal cells. Further, the patents disclose that the cell killing operated through complement mediated killing of the cells.

[0161] Further Demonstration of the Role of CDC

[0162] Indeed, in our work, we have further demonstrated that CBL1 and ABX-CBL operates through complement mediated cell killing. We have utilized a mixed lymphocyte reaction (MLR) assay or a modified MLR assay in our work. The MLR assay provides an in vitro system for assaying proliferation of alloreactive T-lymphocytes. In this manner, the MLR assay is an excellent model of GVHD in patients receiving bone marrow transplant (BMT). In the MLR assay, MHC mismatch lymphocytes from two individuals are co-

cultured. Typically the assays are set up so that the lymphocytes from one patient are inactivated by, for example, radiation (the "stimulators") and the lymphocytes from the other patient are able to act as "Responders" and proliferate and undergo extensive blast transformation. After a suitable period of co-culture, the extent of proliferation of the cells can be quantified by adding tritium-labeled thymidine ($[^3\text{H}]$ thymidine) to the culture medium and monitoring uptake of the label into the DNA of the Responder lymphocytes.

[0163] In our work, use of the CBL1 antibody by itself, the isotype-matched control mouse IgM antibody by itself (**FIG. 2**), or complement (either human or rabbit) by itself in an MLR or ConA induced lymphocyte proliferation assay is ineffective in inhibiting T-cell proliferation. See **FIGS. 2-5**. However, when both complement and the CBL1 and/or ABX-CBL antibody are present, T-cell proliferation is inhibited in a dose dependent manner. See **FIGS. 2-5**. The human IgG2 antibody 2.6.1 is ineffective in inhibiting T-cell proliferation in the same assay, either by itself, or in combination with complement. See **FIG. 5**. This is expected, since the 2.6.1 antibody as a gamma-2 isotype is notoriously less efficient in complement mediated lysis than is an IgM antibody, such as the CBL1 or ABX-CBL antibody.

[0164] Role of Cellular Activation Levels

[0165] We have also studied whether certain cells in becoming activated become sensitive to ABX-CBL binding and cell killing.

[0166] Indeed, we have demonstrated in our work that the T-cell activation marker, CD25 (the alpha-2 subunit of the IL-2 receptor), appears to be expressed in high levels in the same cellular populations as those expressing the antigen to which the ABX-CBL antibody binds. See **FIG. 6**. This finding provided a useful marker to detect whether activated cells were depleted in connection with the MLR assay. Where the MLR assay is conducted utilizing ABX-CBL alone, complement alone, or ABX-CBL and complement in combination, it is only in those experiments where ABX-CBL and complement are used in combination that CD25 expressing cell populations are depleted. See **FIGS. 7-11**. In particular, **FIG. 8** shows cells expressing low levels of CD25. The selective killing of different cell populations are shown in **FIGS. 10-12**.

[0167] Role of Density or Expression Levels of CD147 in CDC

[0168] We have also considered whether CD147 expression levels are higher in given populations of cells (which could also be relevant to CDC).

[0169] In flow cytometry studies with peripheral blood mononuclear cells (PBMC) with the ABX-CBL antibody, we have noticed that, prior to the addition of complement, there are populations of cells that appear to express high and low levels of CD147. After complement is added, there are populations of cells that appear to correspond to the low level expressers mentioned above. It appears that these results could be indicative of density of CD147 expression levels on the cell surface. Density can play a role in CDC through providing additional antigen binding sites to allow for distortion of the antibody which is the first step in triggering the complement cascade. Upon distortion of the antibody, the factor C1q binds first and the cascade proceeds.

[0170] Whether the expression level (or, density) of CD147 in cellular populations plays a role in the therapeutic efficacy of the ABX-CBL antibody can be assayed through analyzing the expression levels of the CD147 molecule in various cellular populations. Generally, the experiments are conducted where beads having various known quantities of the CD147 antigen on their surface are prepared and analyzed on FACS (i.e., utilizing a FITC-labeled anti-CD147 IgG antibody) in order to generate approximately 10-20 data points of different quantities of antigen on the beads. A linear regression curve is prepared from such data. Thereafter, cells expressing the CD147 antigen can be run through FACS and the relative quantities of antigen on the surface of the cells can be calculated from the linear regression curve.

[0171] Presence and Role of Protective Elements in Cellular Populations

[0172] We have also studied whether there is a correlation between certain cellular protective elements in particular cellular populations that inhibit CDC induced by ABX-CBL binding and fixing of complement.

[0173] In connection with this work, we have investigated various cells to which the ABX-CBL antibody binds and considered whether such cells were (i) killed and (ii) if so, was the mechanism similar to complement mediated lysis. In the experiment, we looked for ABX-CBL antibody binding to a number of cells (and, thus, the antigen to which the ABX-CBL antibody binds is expressed upon such cells). Those cells to which ABX-CBL would bind were then tested for complement mediated lysis through treatment with the ABX-CBL antibody and complement. Two T-cell lines (CEM and Jurkat cells), a monocyte line (U937 cells), and three tumor cell lines (A431 (epidermal), SW948 (colon), and MDA468 (breast)), each of which bound the ABX-CBL antibody were examined. Despite the expression on such cells lines, the ABX-CBL antibody is very specific about which cells are killed, being restricted to the CEM T-cell line and U937 monocyte line. See FIG. 13. We also analyzed two endothelial cell lines (i) ECV-304 (ATCC CRL-1998) is a spontaneously transformed immortal EC established from the vein of an apparently normal human umbilical cord and carrying EC characteristics and (ii) HUVEC-C (ATCC CRL-1730) is an EC line derived from the vein of a normal human umbilical cord. Using FACS, we found that the ECV-304 and HUVEC-C lines each stained positive against the 2.6.1, Pharmingen, and ABX-CBL antibodies suggesting that these ECs do express CD147 on the surface. FIGS. 15 and 16, respectively. We then carried out in vitro Alamar-blue based CDC assay and demonstrated that both EC lines were resistant to ABX-CBL mediated CDC in the presence of human complement. See FIGS. 17 and 18, respectively.

[0174] In order to further understand why cells that all appear to express CD147 would not be killed by the ABX-CBL antibody in the presence of complement, we looked into CD46, CD55, and CD59 expression in such cells. Each of CD46 (membrane cofactor protein, MCP), CD55 (decay accelerating factor, DAF), and CD59 (membrane attack complex inhibitor, MACI) have been implicated as complement inhibitory molecules. See e.g., Liszewski et al. *Annu. Rev. Immunol.* 9:431 (1991) and Loveland et al. "Coordinate functions of multiple complement regulating molecules, CD46, CD55 and CD59" *Transpl. Proc.* 26:1070 (1994) related to CD46, Kinoshita et al. "Distribution of decay-

accelerating factor in the peripheral blood of normal individuals and patients with paroxysmal nocturnal hemoglobinuria" *J. Exp. Med.* 162:75 (1985) and Loveland et al. "Coordinate functions of multiple complement regulating molecules, CD46, CD55 and CD59" *Transpl. Proc.* 26:1070 (1994) related to CD55, and Whitlow et al. "H19, a surface membrane molecule involved in T-cell activation, inhibits channel formation by human complement" *Cell. Immunol.* 126: 176 (1990), Loveland et al. "Coordinate functions of multiple complement regulating molecules, CD46, CD55 and CD59" *Transpl. Proc.* 26:1070 (1994), and Davies, A. and Lachmann, P. J. "Membrane defense against complement lysis: the structure and biological properties of CD59" *Immunol. Res.* 12: 258 (1993) related to CD59. Accordingly, we considered whether there was differential expression of either, or both, of these molecules on the cell lines tested above. Indeed, all of the cells, except the CEM line and the U937 line, expressed both of the molecules. And, indeed, the endothelial cell line ECV-304 expressed all three, CD46, CD55, and CD59. FIGS. 19 and 20, respectively. In contrast, the CEM line expressed only CD59 and the U937 line expressed only CD55. See FIG. 14. This data is useful in connection with the prediction of cells that could be selectively eradicated by ABX-CBL and consequently targeted in connection with anti-CD147 in accordance with the present invention.

[0175] Discussion of Function of ABX-CBL/CD147 Based Therapy

[0176] From the foregoing, it is clear that CBL1 and ABX-CBL operates to kill cells through the activation of complement. The combination of ABX-CBL and complement only kill activated T-cells (both CD4⁺ and CD8⁺), activated B-cells, and monocytes, but does not effect resting T-cells and B-cells because such cells do not appear to express CD147 at the same level as the activated cells. It is important, to note that monocytes are also killed by ABX-CBL and complement. This data provides an explanation for the operation of ABX-CBL therapy in diseases, such as GVHD, because, ABX-CBL selectively depletes those effector cells (activated T- and B-cells) and the antigen presenting cells (monocytes and B-cells) which ordinarily would lead to further T-cell activation.

[0177] The mode of operation of the ABX-CBL antibody, and future therapeutic molecules directed against CD147, in this regard appears to be at least partially related to, or dependent upon, each of the above-discussed functional characteristics: (i) complement mediated lysis, (ii) cellular activation, (iii) expression levels of CD147 and/or density of CD147 on the cell surface, and (iv) the absence of expression of one or more of the complement inhibitory molecules on the cell surface. Accordingly, through use of this information, it is possible to design functional assays for the prediction of efficacy of a CD147 based therapeutic.

[0178] Indeed, the desirability of mimicking ABX-CBL binding and efficacy is highlighted based upon a preliminary tissue distribution study of the ABX-CBL antibody. In the study, ABX-CBL is widely distributed throughout a variety of tissues. However, the majority of the distribution is likely to be due to nonspecific binding. Nevertheless, there appears to be specific binding in endothelial cells (venules, arterioles, but not capillary beds), smooth muscle, and some mesothelium. Also, the lymphoreticular tissues appear to be

bound, although, the staining seems to be restricted to large lymphocytes, presumably activated blasts. From the study conducted, it was difficult to distinguish intracellular from extracellular staining. A certain amount of cytoplasmic staining was clearly evident and could have been related to hn-RNP-k binding.

Discussion of Results; Utilization of the ABX-CBL Antibody for the Design of Therapeutics

[0179] The above in vitro work with the ABX-CBL antibody, in combination with the association of the ABX-CBL antibody with the CD147 antigen herein, provide the first evidence that antibodies to CD147 that are capable of complement mediated killing could provide an efficacious approach to the treatment of disease. Moreover, because of CD147's wide distribution and expression in the body and the tissue binding information that indicates that the CBL1 and ABX-CBL antibody associates with many tissues, the excellent prior clinical experience with the CBL1 antibody was difficult to reconcile unless CBL1 and ABX-CBL are, for example, specific to forms of CD147 expressed on certain cells or that other factors associated with complement mediated cell killing limit the CBL1 and ABX-CBL antibody's effects to particular tissues or perhaps a combination thereof.

[0180] Criteria for Generation of CD147 Based Therapeutics

[0181] From the foregoing, it is clear that the ABX-CBL antibody provides a powerful tool for the development of other CD147 based therapeutics. First, because of the extreme safety demonstrated to date with the CBL1 and ABX-CBL antibody, it is desirable to mimic the binding of the ABX-CBL antibody as closely as possible. Second, because of the apparent efficacy of the CBL1 antibody it is desirable, at least initially, that any new therapeutic mediate complement fixation and lysis. Accordingly, in connection with the design of other CD147 based therapeutics, it is expected that through simulating the binding (or structural aspects) and mode of operation (or functional aspects) of ABX-CBL in the therapeutic candidates, safety and efficacy can be expected.

[0182] Structural Considerations

[0183] In connection with simulating or mimicking the structural aspects of ABX-CBL binding, we expect to be able readily generate antibodies that bind to CD147 in a similar manner as ABX-CBL. With the information discussed above, we know at least three levels of detail related to ABX-CBL's binding to CD147: (i) ABX-CBL appears to bind, if not preferentially, to a form of CD147 expressed on the population of cells selected from the group consisting of activated T-cells, activated B-cells, and monocytes, (ii) ABX-CBL shows clear and specific binding to 62 KD and 35 KD molecular species on Western blot analysis, and (iii) ABX-CBL appears very specific to an epitope on CD147 (and potentially a similar epitope on hn-RNP-k protein) defined by the consensus sequence RXRSH. In addition, ABX-CBL can be utilized to "structurally" compare, screen, or act as a functional assay for additional antibody candidates to CD147 through competition studies.

[0184] As will be appreciated, the above information provides highly useful information to the generation of additional antibody candidates. Put another way, antibody

candidates that are generated that possess one or more of the above-characteristics are more likely to possess similar activity to the ABX-CBL antibody. An antibody candidate that possesses greater numbers of similar characteristics is likely to be a very close mimic to the ABX-CBL antibody and, accordingly, would likely exhibit similar safety and efficacy data as the ABX-CBL antibody.

[0185] In addition, as was discussed above, we expect to be able to generate additional information related to the binding of the ABX-CBL antibody to CD147 through certain experiments designed to elucidate ABX-CBL binding, for example, through:

[0186] Additional mapping experiments related to the binding of CD147 to the ABX-CBL antibody. One such set of experiments relate to depletion experiments in which the ABX-CBL antibody bound to CD147 is cleaved with protease and the resulting products scanned with mass spectroscopy and the process repeated as necessary. Another such set of experiments relate to the isolation, purification, and understanding of the 35 KD species recognized by the ABX-CBL antibody. One method of accomplishing this is through the classical purification of the 35 KD molecule as discussed above in connection with the 62 KD species (hn-RNP-k protein). Another approach is the immunoaffinity purification of the 35 KD band through the generation of, for example, Fab fragments of the ABX-CBL antibody and binding the same to a column as discussed above in connection with the immunoaffinity purification conducted with the 2.6.1 antibody.

[0187] Experiments directed to understanding CD147 cellular development. For example, the development of CD147 on the cell surface can be gleaned through conducting "pulse-chase" experiments. In such experiments, cells (such as CEM cells) growing in culture (Met⁻ media) are "pulsed" with S³⁵-Met for a sufficient time periods (and varied time periods) for the label to be enrolled into the cellular protein synthesis. Thereafter, cells are washed with "cold" medium and CD147 on the cell surface can be immunoprecipitated and subjected to autoradiography. Information can be gained related to potential alternative splicings, glycosylation levels, and other developmental differences of the expressed CD147 molecules.

[0188] Experiments related to the role of glycosylation levels to ABX-CBL binding to CD147 can also be queried through reaction of CD147 with various glycosidases (see e.g., Mizukami et al. *J. Immunol.* 147:1331-1337 (1991), Schlosshauer *Development* 113:129-140 (1991), Fadool and Linser *J. Neurochemistry* 60:1354-1364 (1993)) and considering ABX-CBL binding to the various forms.

[0189] Functional Considerations

[0190] Once, or, concurrently with determining whether, one is satisfied with the "structure" of an antibody candidate (i.e., in connection with the antibody's binding to CD147), in accordance with the present invention, we have provided detailed functional criteria that appear important to the ABX-CBL antibody's in vivo efficacy that can be utilized to determine whether an antibody candidate is likely to operate in a similar manner to the ABX-CBL antibody. Such features include (i) cell killing through CDC, (ii) apparent effect of

density or expression of the CD147 molecule on cellular populations, and (iii) the role of protective factors (for example, CD46, CD55, and CD59) on cellular populations.

[0191] As will be appreciated, the above information provides highly useful information to the generation of additional antibody candidates. Put another way, antibody candidates that are generated that possess one or more of the above-characteristics are more likely to possess similar activity to the ABX-CBL antibody. An antibody candidate that possesses greater numbers of similar characteristics is likely to be a very close mimic to the ABX-CBL antibody and, accordingly, would likely exhibit similar safety and efficacy data as the ABX-CBL antibody.

[0192] In Vivo Models

[0193] Each of the foregoing features, whether structural or functional, can essentially be carried out in vitro. Of course, however, prior to proceeding into man with therapeutic candidates it is desirable to generate in vivo data to ensure that operation of the antibody candidate will be safe and efficacious in vivo. In connection with GVHD, there are several animal models that have been shown to be highly predictive of the operation of therapeutic candidates in man. Such models include:

[0194] Murine model (Hakim F T & Mackall C L "The Immune System: Effector and Target of Graft-Versus-Host Disease" in *Graft-vs. Host Disease* (Ferrara et al. eds, 2d edition, Marcel Dekker, Inc., NY (1997)).

[0195] Canine Model (Storb et al. *Blood* 89:3048-3054 (1997); Yu et al. *Bone Marrow Transplantation* 17:649-653 (1996), Raff et al. *Transplantation* 54:813-820 (1992); and Deeg et al. *Transplantation* 37:62-65 (1984))

[0196] Primate Skin Graft Model (Chatterjee et al. *Hybridoma* 1:369-377 (1982) and Billing R. and Chatterjee S. *Transplantation Proceedings* 15:649-650 (1983))

[0197] As will be appreciated, in order such models to be predictive, it is necessary that the antibody candidate is reactive with the endogenous form of CD147 in the animal.

Construction of Antibodies

[0198] An excellent model in which to generate therapeutic molecules targeting CD147 is in connection with the generation of antibodies. Antibodies can be generated with relative ease and are also capable of ready screening. In recent years, it has become possible to generate different "types" of antibodies; from conventional murine antibodies through human antibodies generated from transgenic animals. Within that spectrum, antibodies can also be generated through display techniques (i.e., phage), murine or other antibodies can be humanized, and the like. Some of these techniques are discussed below.

[0199] In connection with the generation of antibodies through immunization techniques, both classical and advanced immunization techniques can be used. By classical, we mean that animals can simply be immunized with the antigen, lymphocytic cells fused with myeloma cells, and hybridomas screened therefrom. By advanced, we mean that either immunization schemes can be biased or, instead of simply forming hybridomas, lymphocytic cells can be used

directly to form display libraries and screened using, for example, phage or other display technologies. Such techniques are conventional in the art and are discussed in additional detail below. In connection with biasing immunizations, one can immunize with CD147, followed by immunization with peptides, such as the 15-mer peptide mentioned above. In this manner, there is a higher probability of generating antibodies that possess specificity and affinity for selected epitopes for example. Thus, it is expected that antibodies having specificity for the RXRSH consensus sequence in CD147, as discussed above, can be more readily generated. It will be appreciated that such immunization techniques can be utilized in connection with standard fusions and screening procedures or advanced screening procedures. Another set of advanced immunization techniques are related to techniques of antigen presentation (i.e., DEC systems) and techniques to augment the immune response (i.e., CD140 systems) in the animal in which the immunization is being undertaken.

[0200] Generation of Human Antibodies from Transgenic Animals

[0201] The generation of fully human antibodies, for example, from transgenic animals, is very attractive. Fully human antibodies are expected to minimize the immunogenic and allergic responses intrinsic to mouse or mouse-derived Mabs and thus to increase the efficacy and safety of the administered antibodies. The use of fully human antibodies can be expected to provide a substantial advantage in the treatment of chronic and recurring human diseases, such as inflammation, autoimmunity, and cancer, which often require repeated antibody administrations.

[0202] One approach that has been utilized in connection with the generation of human antibodies is the construction of mouse strains that are deficient in mouse antibody production but that possess large fragments of the human Ig loci so that such mice would produce a large repertoire of human antibodies in the absence of mouse antibodies. Large human Ig fragments preserve the large variable gene diversity as well as the proper regulation of antibody production and expression. By exploiting the mouse machinery for antibody diversification and selection and the lack of immunological tolerance to human proteins, the reproduced human antibody repertoire in these mouse strains yields high affinity antibodies against any antigen of interest, including human antigens. Using hybridoma technology, antigen-specific human Mabs with the desired specificity can be readily produced and selected.

[0203] This general strategy was demonstrated in connection with the generation of the first XenoMouse strains as published in 1994. See Green et al. *Nature Genetics* 7:13-21 (1994). The XenoMouse strains were engineered with 245 kb and 190 kb-sized germline configuration fragments of the human heavy chain loci and kappa light chain loci, respectively, which contained core variable and constant region sequences. The human Ig containing yeast artificial chromosomes (YACs) proved to be compatible with the mouse system for both rearrangement and expression of antibodies, and were capable of substituting for the inactivated mouse Ig genes. This was demonstrated by their ability to induce B-cell development and to produce an adult-like human repertoire of fully human antibodies and to generate antigen-specific human Mabs. These results also

suggested that introduction of larger portions of the human Ig loci containing greater numbers of V genes, additional regulatory elements, and human Ig constant regions might recapitulate substantially the full repertoire that is characteristic of the human humoral response to infection and immunization.

[0204] Such approach is further discussed and delineated in U.S. patent application Ser. No. 07/466,008, filed Jan. 12, 1990, Ser. No. 07/610,515, filed Nov. 8, 1990, Ser. No. 07/919,297, filed Jul. 24, 1992, Ser. No. 07/922,649, filed Jul. 30, 1992, filed Ser. No. 08/031,801, filed Mar. 15, 1993, Ser. No. 08/112,848, filed Aug. 27, 1993, Ser. No. 08/234,145, filed Apr. 28, 1994, Ser. No. 08/376,279, filed Jan. 20, 1995, Ser. No. 08/430,938, Apr. 27, 1995, Ser. No. 08/464,584, filed Jun. 5, 1995, Ser. No. 08/464,582, filed Jun. 5, 1995, Ser. No. 08/463,191, filed Jun. 5, 1995, Ser. No. 08/462,837, filed Jun. 5, 1995, Ser. No. 08/486,853, filed Jun. 5, 1995, Ser. No. 08/486,857, filed Jun. 5, 1995, Ser. No. 08/486,859, filed Jun. 5, 1995, Ser. No. 08/462,513, filed Jun. 5, 1995, Ser. No. 08/724,752, filed Oct. 2, 1996, and Ser. No. 08/759,620, filed Dec. 3, 1996. See also European Patent No., EP 0 463 151 B1, grant published Jun. 12, 1996, International Patent Application No., WO 94/02602, published Feb. 3, 1994, International Patent Application No., WO 96/34096, published Oct. 31, 1996, and PCT Application No. PCT/US96/05928, filed Apr. 29, 1996. The disclosures of each of the above-cited patents and applications are hereby incorporated by reference in their entirety.

[0205] In an alternative approach, others, including GenPharm International, Inc., have utilized a "minilocus" strategy. In the minilocus strategy, an exogenous Ig locus is mimicked through the inclusion of pieces (individual genes) from the Ig locus. Thus, one or more V_H genes, one or more D_H genes, one or more J_H genes, a mu constant region, and a second constant region (preferably a gamma constant region) are formed into a construct for insertion into an animal. This approach is described in U.S. Pat. No. 5,545,807 to Surani et al., U.S. Pat. Nos. 5,545,806, 5,625,825, 5,661,016, 5,633,425, and 5,625,126, each to Lonberg and Kay, U.S. Pat. No. 5,643,763 to Dunn and Choi, U.S. Pat. No. 5,612,205 to Kay et al., U.S. Pat. No. 5,591,669 to Krimpenfort and Berns, and GenPharm International U.S. patent application Ser. No. 07/574,748, filed Aug. 29, 1990, Ser. No. 07/575,962, filed Aug. 31, 1990, Ser. No. 07/810,279, filed Dec. 17, 1991, Ser. No. 07/853,408, filed Mar. 18, 1992, Ser. No. 07/904,068, filed Jun. 23, 1992, Ser. No. 07/990,860, filed Dec. 16, 1992, Ser. No. 08/053,131, filed Apr. 26, 1993, Ser. No. 08/096,762, filed Jul. 22, 1993, Ser. No. 08/155,301, filed Nov. 18, 1993, Ser. No. 08/161,739, filed Dec. 3, 1993, Ser. No. 08/165,699, filed Dec. 10, 1993, Ser. No. 08/209,741, filed Mar. 9, 1994, Ser. No. 08/544,404, filed Oct. 10, 1995, the disclosures of which are hereby incorporated by reference. See also International Patent Application Nos. WO 97/13852, published Apr. 17, 1997, WO 94/25585, published Nov. 10, 1994, WO 93/12227, published Jun. 24, 1993, WO 92/22645, published Dec. 23, 1992, WO 92/03918, published Mar. 19, 1992, the disclosures of which are hereby incorporated by reference in their entirety. See further Taylor et al., 1992, Chen et al., 1993, Tuailon et al., 1993, Choi et al., 1993, Lonberg et al., (1994), Taylor et al., (1994), and Tuailon et al., (1995), the disclosures of which are hereby incorporated by reference in their entirety.

[0206] The inventors of Surani et al., cited above, and assigned to the Medical Research Counsel (the "MRC"), produced a transgenic mouse possessing an Ig locus through use of the minilocus approach. The inventors on the GenPharm International work, cited above, Lonberg and Kay, following the lead of the present inventors, proposed inactivation of the endogenous mouse Ig locus coupled with substantial duplication of the Surani et al. work.

[0207] An advantage of the minilocus approach is the rapidity with which constructs including portions of the Ig locus can be generated and introduced into animals. Commensurately, however, a significant disadvantage of the minilocus approach is that, in theory, insufficient diversity is introduced through the inclusion of small numbers of V, D, and J genes. Indeed, the published work appears to support this concern. B-cell development and antibody production of animals produced through use of the minilocus approach appear stunted. Therefore, the present inventors have consistently urged introduction of large portions of the Ig locus in order to achieve greater diversity and in an effort to reconstitute the immune repertoire of the animals.

[0208] It will be appreciated that through use of the above-technology, human antibodies can be generated to, for example, CD147 expressing cells, CD147 itself, forms of CD147, epitopes or peptides thereof, and expression libraries thereto (see e.g. U.S. Pat. No. 5,703,057) through immunization of a transgenic mouse therewith, forming hybridomas, and screening the resulting hybridomas as described above for the activities described above.

[0209] Indeed, through use of the above-discussed technology, we prepared a panel of human monoclonal antibodies that bind CD147 through immunization of XenoMouse™ strains of transgenic mice (see Mendez et al., (1997), supra. and U.S. patent application Ser. No. 08/759,620, filed Dec. 3, 1996), available from Abgenix, Inc., Fremont, Calif. Such antibodies were further screened for their ability to compete with ABX-CBL for binding with CD147. In such panel, both human IgG2 and human IgM antibodies were detected that bound to CD147 and were capable of competition with ABX-CBL for binding to CD147. The hybridomas expressing such antibodies were designated as follows:

[0210] IgMs: CEM 10.1 C3, CEM 10.1 G10, CEM 10.12 F3, CEM 10.12 G5 CEM 13.12, CEM 13.5; and

[0211] IgG2s: 2.4.4, 2.1.1, 2.3.2, 2.6.1.

[0212] Each of the above antibodies were sequenced through isolating cDNAs encoding them from the corresponding hybridomas through RT-PCR. Germline gene identifications were made and the sequences of the antibodies compared to the germline sequences. Germline gene identifications are provided in the following Table:

TABLE 1

Antibody	Heavy/ Light	V_H or V_K	D	J_H or J_K
CEM 10-1 C3	Heavy	V4-34	D2/D2-15	JH6b
	Light	A3/A19/DPK15		JK1
CEM 10.1 G10	Heavy	DP71 (V4-59)	D1-26	JH6b
	Light	A30		JK1
				(not identical seq)

TABLE 1-continued

Antibody	Heavy/ Light	V _H or V _K	D	J _H or J _K
CEM 10.12 F3	Heavy	DP15 (V1-8)	D1-26	JH6b
	Light	B3/DPK24		JK1
CEM 10.12 G5	Heavy	DP15 (V1-8)	D6-19	JH6b
	Light	A30		JK1
CEM 13.12	Heavy	V4-34	D2-2/D4	JH6b
	Light	A3/A19/DPK15		JK3
CEM 13.5	Heavy	DP77-WH16 (3-21)	D6-19	JH4b
	Light	B3/DPK24		JK1 (not identical seq)
2.4.4	Heavy	VII-5	D21-9/D3-22	JH4b
	Light	A2 DPK12		JK4
2.1.1	Heavy	DP77	D6-19	JH4b
	Light	LFVK431		JK3
2.3.2	Heavy	VII-5	D21-9/D3-22	JH4b
	Light	A2 DPK12		JK4
2.6.1	Heavy	DP47	DXP4	JH4b
	Light	LFVK431		JK3

[0213] Germline sequences of the V_H, D, J_H, V_K, and J_K genes are available on GenBank. The sequences of certain of the antibodies were compared to transcripts of the germline V-gene segments to observe somatic mutations in the amino acid sequences. Such sequence comparisons are shown in **FIGS. 44 through 46**. cDNA sequences and protein transcripts of and for each of the antibodies are shown in **FIGS. 24 through 33**. In addition, CDRs, according to Kabat numbering scheme, of the heavy chains and kappa light chains of the antibodies are shown in **FIGS. 34 through 43**.

[0214] It will be appreciated that CDRs of the above antibodies are generally very important in connection with antibody binding to an antigen. Accordingly, it will be understood that a variety of FR and other modifications can be made in and to antibodies that do not modify an antibody binding the epitope on an antigen. Thus, an important factor in an antibody's activity is the epitope on an antigen to which an antibody binds. So long as the epitope binding is conserved, in many ways it may matter little if the primary sequence of the antibody is modified. Therefore, where sequences are discussed herein, it is submitted that the sequence of an antibody may initially define an efficacious epitope on the antigen, however, once the epitope is identified by the antigen, any antibody that binds to the same epitope on the is contemplated herein.

[0215] In view of a number of tests that were conducted, the 2.6.1 IgM antibody was chosen for additional development. As will be appreciated, all of the IgMs that were generated were monovalent. Accordingly, in order to prepare a fully human multimeric IgM antibody, we cloned the human J-chain gene from human buffy coat cells, prepared a first expression vector containing the 2.6.1 kappa light chain cDNA and the J-chain cDNA and a second expression

vector containing the 2.6.1 heavy chain cDNA, cotransfected DHFR⁻ Chinese hamster ovary cells with the two vectors, and selected clones expressing the multimeric IgM.

[0216] The 2.6-1 IgM+J-Chain antibody was capable of acting in ADCC as shown in **FIG. 50**.

[0217] Humanization and Display Technologies

[0218] As was discussed above in connection with human antibody generation, there are advantages to producing antibodies with reduced immunogenicity. To a degree, this can be accomplished in connection with techniques of humanization and display techniques using appropriate libraries. It will be appreciated that murine antibodies or antibodies from other species can be humanized or primatized using techniques well known in the art. See e.g., Winter and Harris *Immunol Today* 14:43-46 (1993) and Wright et al. *Crit. Reviews in Immunol.* 12:125-168 (1992). Further, human antibodies or antibodies from other species can be generated through display-type technologies, including, without limitation, phage display, retroviral display, ribosomal display, and other techniques, using techniques well known in the art and the resulting molecules can be subjected to additional maturation, such as affinity maturation, as such techniques are well known in the art. Wright and Harris, supra., Hanes and Pluchthau *PNAS USA* 94:4937-4942 (1997) (ribosomal display), Parmley and Smith *Gene* 73:305-318 (1988) (phage display), Scott *TIBS* 17:241-245 (1992), Cwirla et al. *PNAS USA* 87:6378-6382 (1990), Russel et al. *Nucl. Acids Research* 21:1081-1085 (1993), Hoganboom et al. *Immunol. Reviews* 130:43-68 (1992), and Chiswell and McCafferty *TIBTECH* 10:80-84 (1992). If display technologies are utilized to produce antibodies that are not human, such antibodies can be humanized as described above.

[0219] Using these techniques, antibodies can be generated to CD147 expressing cells, CD147 itself, forms of CD147, epitopes or peptides thereof, and expression libraries thereto (see e.g. U.S. Pat. No. 5,703,057) which can thereafter be screened as described above for the activities described above.

[0220] Further, the sequence for the active antibody from the deposited hybridoma cell line expressing the ABX-CBL antibody was previously unknown. In view of our findings discussed above that the IgM antibody was the entity responsible for the activity of the CBL1 antibody and the fact that neither the presence nor the absence of the MOPC21 light chain appeared to be advantageous nor detrimental to the activity of the antibody, we cloned the heavy chain and the kappa light chains from the IgM (ABX-CBL) producing hybridoma through RT-PCR and sequenced the cDNAs. The results of such sequencing studies, including the cDNA sequences of the heavy chain and kappa light chain and the protein transcripts thereof are shown below:

ABX-CBL Heavy Chain Nucleotide Sequence

(SEQ ID NO: 81)

```

ATGTACTTGG GACTGAACCTA TGTATTCATA GTTTTCTCT TAAATGGTGT 50
CCAGAGTGAA GTGAAGCTTG AGGAGTCTGG AGGAGGCTTG GTGCAACCTG 100
GAGGATCCAT GAAACTCTCC TGTGTTGCCT CTGGATTAC TTTCAAGTAAC 150
TACTGGATGA ACTGGGTCCG CCAGTCTCCA GAGAAGGGGC TTGAGTGGGT 200
TGCTGAAATT AGATTGAANF CTAATAATTA TGCAACACAT TATGCGGAGT 250

```

-continued

CTGTGAAAGG	GAGGTTACCC	ATCTCAAGAG	ATGATTCCAA	AAGTAGTGTG	300
TACCTGCAAA	TGAACAACTT	AAGAGCTGAA	GACACTGGCA	TTTATTACTG	350
TACGGATTAC	GATGCTTACT	GGGGCCAAGG	GACTCTGGTC	ACTGTCTCTG	400
CAGAGAGTCA	GTCCTTCCCA	AATGTCTTCC	CCCTCGTCTC	CTGCGAGAGC	450
CCCTGTGCTG	ATAAGAATCT	GGTGGCCATG	GGCTGCCTGG	CCCGGGACTT	500
CCTGCCAGC	ACCATTTCTT	TCACCTGGAA	CTACCAGAAC	AACACTGAAG	550
TCATCCAGGG	TATCAGAACC	TTCCCAACAC	TGAGGACAGG	GGGCAAGTAC	600
CTAGCCACCT	CGCAGGTGTT	GCTGTCTCCC	AAGAGCATCC	TTGAAGGTTT	650
AGATGAATAC	CTGGTATGCA	AAATCCACTA	CGGAGGCAAA	AACAGAGATC	700
TGCATGTGCC	CATTCCAGCT	GTCGCAGAGA	TGAACCCCAA	TGTAAATGTG	750
TTCTGTCCAC	CACGGGATGG	CTTCTCTGGC	CCTGCACCAC	GCAAGTCTAA	800
ACTCATCTGC	GAGGCCACGA	ACTTCACTCC	AAAACCGATC	ACAGTATCCT	850
GGCTAAAGGA	TGGGAAGCTC	GTGGAATCTG	GCTTCACCAC	AGATCCGGTG	900
ACCATCGAGA	ACAAAGGATC	CACACCCCAA	ACCTACAAGG	TCATAAGCAC	950
ACTTACCATC	TCTGAAATCG	ACTGGCTGAA	CCTGAATGTG	TACACCTGCC	1000
GTGTGGATCA	CAGGGGTCTC	ACCTTCTTGA	AGAAGCTGTC	CTCCACATGT	1050
GCTGCCAGTC	CCTCCACAGA	CATCCTAACC	TTACCATACC	CCCCCTCCTT	1100
TGCCGACATC	TTCTCAGCA	AGTCCGCTAA	CCTGACCTGT	CTGGTCTCAA	1150
ACCTGGCAAC	CTATGAAACC	CTGAATATCT	CCTGGGCTTC	TCAAAGTGGT	1200
GAACCACTGG	AAACCAAAAT	TAAATCATG	GAAAGCCATC	CCAATGGCAC	1250
CTTCAGTGCT	AAGGGTGTGG	CTAGTGTTTG	TGTGGAAGAC	TGGAATAACA	1300
GGAAGGAATT	TGTGTGTACT	GTGACTCACA	GGGATCTGCC	TTCACCACAG	1350
AAGAAATTC	TCTCAAAACC	CAATGAGGTG	CACAAACATC	CACCTGTGTT	1400
GTACCTGCTG	CCACCAGCTC	GTGAGCAACT	GAACCTGAGG	GAGTCAGCCA	1450
CAGTCACCTG	CCTGGTGAAG	GGCTTCTCTC	CTGCAGACAT	CAGTGTGCAG	1500
TGGCTTCAGA	GAGGGCAACT	CTTGCCCCAA	GAGAAGTATG	TGACCAGTGC	1550
CCCAGTGCCA	GAGCCTGGGG	CCCCAGGCTT	CTACTTTACC	CACAGCATCC	1600
TGACTGTGAC	AGAGGAGGAA	TGGAATCCG	GAGAGACCTA	TACCTGTGTT	1650
GTAGGCCACG	AGGCCCTGCC	ACACCTGGTG	ACCGAGAGGA	CCGTGGACAA	1700
GTCCACTGGT	AAACCCACAC	TGTACAATGT	CTCCCTGATC	ATGTCTGACA	1750
CAGGCGGCAC	CTGCTATTGA	CCAT			1774

ABX-CBL Heavy Chain Protein Sequence

(SEQ ID NO:18)

EVKLEESGGG	LVQPGGSMKL	SCVASGFTFS	NYWMNWVRQS	PEKLEWVAE	50
IRLKSNNYAT	HYAESVKGRF	TISRDDSKSS	VYLQMNLLRA	EDTGIYYCTD	100
YDAYWGGQTL	VTVSAESQSF	PNVFPLVSC	SPLSDKNLVA	MGCLARDFLP	150
STISFTWNYQ	NNTFVIQIR	TFPTLRGTGG	YLATSQVLLS	PKSILEGSDE	200
YLVCKIHYGG	KNRDLHVP	IPVAVEMNPVN	VFVPPRDGFS	GPAPRKSCLI	250
CEATNFTPKP	ITVSWLKD	KGKLVESGFTTDP	VTIENKGSTP	QTYKVISTLT	300
ISEIDWLNLN	VYTCRVDHRG	LTFLLKNVSS	CAASPSTDIL	TFTIPPSFAD	350
IFLSKSANLT	CLVSNLATYE	TLNISWASQS	GEPLTKIKI	MESHPPNGTFS	400
AKGVASVCVE	DWNNRKEFVC	TVTHRDLPSP	QKKFISKPNE	VHKHPPAVYL	450
LPPAREQLNL	RESATVTCLV	KGFSPADISV	QWLQRGQLLP	QEKYVTSAPM	500
PEPGAPGFYF	THSILTVTEE	EWNSGETYTC	VVGHEALPHL	VTERTVDKST	550
GKPTLYNVSL	IMSDTGGTCY				570

ABX-CBL Light Chain Protein Sequence

(SEQ ID NO:19)

KFLLVSAHDR	VTITCKASQS	VSNDVAWYQQ	KPGQSPKLLI	YYASNRYTGV	50
PDRFTGSGYG	TDFTFTISTV	QAEDLAVYFC	QDYSSPYTF	GGGTYKLEIKR	100
ADAAPTVSIF	PPSSQLTSG	GASVVCFLNN	FYPKDINVKW	KIDGSEQRNG	150
VLNSWTDQDS	KDSTYSMSST	LTLTKDEYER	HNSYTCEATH	KTSTSPIVKS	200
FNRNEC					206

[0221] As will be appreciated, through utilization of the sequence, it is possible to prepare a humanized version of the ABX-CBL antibody. In general, the nucleotide sequences encoding the CDRs are grafted into human framework (FR) sequences using conventional techniques. Alternatively, amino acid residues in the framework regions surrounding the CDRs (i.e., residues in FR1 and FR2, surrounding CDR1, FR2 and FR3, surrounding CDR2, and/or FR3 and FR4, surrounding CDR3) are modified through mutagenesis of cDNAs encoding the same also using conventional techniques. In either case, the modified cDNAs encoding the humanized kappa light chain and the heavy chain are generally then introduced into a cell line for expression (i.e., NSO, CHO, or the like) either directly,

through cotransfection, or through use of the cell-cell fusion techniques described in U.S. patent application Ser. No. 08/730,639, filed Oct. 11, 1996 or International Patent Application No. WO 98/16654, published Apr. 23, 1998. Thereafter, the humanized antibodies are expressed and assayed for binding and other functional attributes. The molecules can be iteratively modified at the DNA level as desired or necessary to achieve improved binding or other functional attributes of the antibodies. For example, in certain cases, it is necessary to reintroduce murine sequences within the human FRs to improve binding. A good step-by-step introduction to humanization and demonstrating how conventional humanization has become in the art is provided on the internet <http://www.cryst.bbk.ac.uk/~ubcg07s/>.

[0222] In general, at the same time, or during the process, the constant region would be switched from the murine IgM to another human constant region (such as a human IgM constant region, without or without the J-chain, as discussed above) to prepare a humanized chimeric antibody.

[0223] Additional Criteria for Antibody Therapeutics

[0224] As discussed herein, the function of the ABX-CBL antibody appears important to at least a portion of its mode of operation. By function, we mean, by way of example, the activity of the ABX-CBL antibody is CDC. Accordingly, it is desirable in connection with the generation of antibodies as therapeutic candidates against CD147 that the antibodies be capable of fixing complement and participating in CDC. There are a number of isotypes of antibodies that are capable of the same, including, without limitation, the following: murine IgM, murine IgG2a, murine IgG2b, murine IgG3, human IgM, human IgG1, and human IgG3. It will be appreciated that antibodies that are generated need not initially possess such an isotype but, rather, the antibody as generated can possess any isotype and the antibody can be isotype switched thereafter using conventional techniques that are well known in the art. Such techniques include the use of direct recombinant techniques (see e.g., U.S. Pat. No. 4,816,397), cell-cell fusion techniques (see e.g., U.S. patent application Ser. No. 08/730,639, filed Oct. 11, 1996), among others.

[0225] In the cell-cell fusion technique, a myeloma or other cell line is prepared that possesses a heavy chain with any desired isotype and another myeloma or other cell line is prepared that possesses the light chain. Such cells can, thereafter, be fused and a cell line expressing an intact antibody can be isolated.

[0226] By way of example, the 2.6.1 antibody discussed herein is a human anti-CD147 IgG2 antibody. If such antibody possessed desired binding to the CD147 molecule, it could be readily isotype switched to generate an human IgM, human IgG1, or human IgG3 isotype, while still possessing the same variable region (which defines the antibody's specificity and some of its affinity). Such molecule would then be capable of fixing complement and participating in CDC, in a similar manner to the ABX-CBL antibody.

[0227] Accordingly, as antibody candidates are generated that meet desired "structural" attributes as discussed above, they can generally be provided with at least certain of the desired "functional" attributes through isotype switching.

Design and Generation of Other Therapeutics

[0228] In accordance with the present invention and based on the activity of the ABX-CBL antibody with respect to CD147, it is now also possible to design other therapeutic modalities beyond ordinary antibody moieties, including, without limitation, advanced antibody therapeutics, such as bispecific antibodies, immunotoxins, and radiolabeled therapeutics, generation of peptide therapeutics, gene therapies, particularly intrabodies, antisense therapeutics, and small molecules.

[0229] In connection with the generation of advanced antibody therapeutics, it may be possible to sidestep the dependence on complement for cell killing that we have

demonstrated is necessary for the function of the ABX-CBL antibody through the use of bispecifics, immunotoxins, or radiolabels, for example.

[0230] For example, in connection with bispecific antibodies, bispecific antibodies can be generated that comprise (i) two antibodies one with a specificity to CD147 and another to a second molecule that are conjugated together, (ii) a single antibody that has one chain specific to CD147 and a second chain specific to a second molecule, or (iii) a single chain antibody that has specificity to CD147 and the other molecule. Such bispecific antibodies can be generated using techniques that are well known for example, in connection with (i) and (ii) see e.g., Fanger et al. *Immunol Methods* 4:72-81 (1994) and Wright and Harris, *supra.* and in connection with (iii) see e.g., Traunecker et al. *Int. J. Cancer (Suppl.)* 7:51-52 (1992). In each case, the second specificity can be made to the heavy chain activation receptors, including, without limitation, CD16 or CD64 (see e.g., Deo et al. 18:127 (1997)) or CD89 (see e.g., Valerius et al. *Blood* 90:4485-4492 (1997)). Bispecific antibodies prepared in accordance with the foregoing would be likely to kill cells expressing CD147, and particularly those cells in which the ABX-CBL antibody is effective.

[0231] In connection with immunotoxins, antibodies can be modified to act as immunotoxins utilizing techniques that are well known in the art. See e.g., Vitetta *Immunol Today* 14:252 (1993). See also U.S. Pat. No. 5,194,594. In connection with the preparation of radiolabeled antibodies, such modified antibodies can also be readily prepared utilizing techniques that are well known in the art. See e.g. Junghans et al. in *Cancer Chemotherapy and Biotherapy* 655-686 (2d edition, Chafner and Longo, eds., Lippincott Raven (1996)). See also U.S. Pat. Nos. 4,681,581, 4,735,210, 5,101,827, 5,102,990 (RE 35,500), 5,648,471, and 5,697,902. Each of immunotoxins and radiolabeled molecules would be likely to kill cells expressing CD147, and particularly those cells in which the ABX-CBL antibody is effective.

[0232] In connection with the generation of therapeutic peptides, through the utilization of structural information related to CD147 and antibodies thereto, such as the ABX-CBL antibody (as discussed below in connection with small molecules) or screening of peptide libraries, therapeutic peptides can be generated that are directed against CD147. Design and screening of peptide therapeutics is discussed in connection with Houghten et al. *Biotechniques* 13:412-421 (1992), Houghten *PNAS USA* 82:5131-5135 (1985), Pinalla et al. *Biotechniques* 13:901-905 (1992), Blake and Litz-Davis *BioConjugate Chem.* 3:510-513 (1992). Immunotoxins and radiolabeled molecules can also be prepared, and in a similar manner, in connection with peptidic moieties as discussed above in connection with antibodies.

[0233] Assuming that the CD147 molecule (or a form, such as a splice variant or alternate form) is functionally active in a disease process, it will also be possible to design gene and antisense therapeutics thereto through conventional techniques. Such modalities can be utilized for modulating the function of CD147. In connection therewith the discovery of the present invention allows design and use of functional assays related thereto. A design and strategy for antisense therapeutics is discussed in detail in International Patent Application No. WO 94/29444. Design and strategies for gene therapy are well known. However, in particular, the

use of gene therapeutic techniques involving intrabodies could prove to be particularly advantageous. See e.g., Chen et al. *Human Gene Therapy* 5:595-601 (1994) and Marasco *Gene Therapy* 4:11-15 (1997). General design of and considerations related to gene therapeutics is also discussed in International Patent Application No- WO 97/38137.

[0234] Small molecule therapeutics can also be envisioned in accordance with the present invention. Drugs can be designed to modulate the activity of CD147 based upon the present invention. Knowledge gleaned from the structure of the CD147 molecule and its interactions with other molecules in accordance with the present invention, such as the ABX-CBL antibody, CD46, CD55, CD59, and others can be utilized to rationally design additional therapeutic modalities. In this regard, rational drug design techniques such as X-ray crystallography, computer-aided (or assisted) molecular modeling (CAMM), quantitative or qualitative structure-activity relationship (QSAR), and similar technologies can be utilized to focus drug discovery efforts. Rational design allows prediction of protein or synthetic structures which can interact with the molecule or specific forms thereof which can be used to modify or modulate the activity of CD147. Such structures can be synthesized chemically or expressed in biological systems. This approach has been reviewed in Capsey et al. *Genetically Engineered Human Therapeutic Drugs* (Stockton Press, NY (1988)). Further, combinatorial libraries can be designed and synthesized and used in screening programs, such as high throughput screening efforts.

Therapeutic Administration and Formulations

[0235] It will be appreciated that administration of therapeutic entities in accordance with the invention will be administered with suitable carriers, excipients, and other agents that are incorporated into formulations to provide improved transfer, delivery, tolerance, and the like. A multitude of appropriate formulations can be found in the formulary known to all pharmaceutical chemists: Remington's Pharmaceutical Sciences (15th ed, Mack Publishing Company, Easton, Pa. (1975)), particularly Chapter 87 by Blaug, Seymour, therein. These formulations include, for example, powders, pastes, ointments, jellies, waxes, oils, lipids, lipid (cationic or anionic) containing vesicles (such as Lipofectin™), DNA conjugates, anhydrous absorption pastes, oil-in-water and water-in-oil emulsions, emulsions carbowax (polyethylene glycols of various molecular weights), semi-solid gels, and semi-solid mixtures containing carbowax. Any of the foregoing mixtures may be appropriate in treatments and therapies in accordance with the present invention, provided that the active ingredient in the formulation is not inactivated by the formulation and the formulation is physiologically compatible and tolerable with the route of administration. See also Powell et al. "Compendium of excipients for parenteral formulations" *PDA J Pharm Sci Technol.* 52:238-311 (1998) and the citations therein for additional information related to excipients and carriers well known to pharmaceutical chemists.

EXAMPLES

[0236] The following examples, including the experiments conducted and results achieved, are provided for illustrative purposes only and are not to be construed as limiting upon the present invention.

Experiment 1 Generation of Human Antibodies

[0237] Human antibodies were prepared in accordance with Mendez et al. *Nature Genetics* 15:146-156 (1997) and U.S. patent application Ser. No. 08/759,620, filed Dec. 3, 1996, the disclosures of which are hereby incorporated by reference herein in their entirety, through the immunization of XenoMouse™ animals with CEM cells, followed by fusions, and screening of the resulting hybridoma supernatants against CEM cells and in competition assays with the ABX-CBL antibody (Example 2).

Experiment 2 Immuno-affinity Purification of ABX-CBL Antigen

[0238] We undertook immunoaffinity purification of the antigen to which the ABX-CBL antibody bound. The antigen to which the CBL1 and ABX-CBL antibody bound appeared to be highly expressed on CEM cells. Immunoaffinity purification using the native ABX-CBL antibody was frustrated by the fact that the ABX-CBL antibody is an IgM antibody having a pentameric structure. Therefore, we prepared human IgG2 antibodies (Example 1), followed by fusions, and screening of the resulting hybridoma supernatants against CEM cells and tested for competition with the ABX-CBL antibody in binding assays with CEM cells using FACS. In the FACS competition assays, inhibition of the binding of ABX-CBL antibodies, labeled with FITC, to CEM cells was analyzed, both alone and in the presence of the anti-CEM human antibodies.

[0239] We obtained four hybridoma clones from the fusions that produced monoclonal antibodies that bound to the CEM cells and that were highly competitive with the ABX-CBL antibody in binding to the CEM cells. One hybridoma clone, designated 2.6.1 appeared most competitive.

[0240] We generated ascites to each of the four hybridoma clones, including the 2.6.1 hybridoma, in SCID mice and purified the 2.6.1 antibody using a Protein A affinity purification process using standard conditions. From the purified 2.6.1 antibody, we prepared an immunoaffinity column. To prepare the column, the purified 2.6.1 antibody was conjugated to CNBr activated Sepharose-4B, according to the manufacturer's specifications. Approximately 8.4 mg of the antibody was conjugated to about 2.0 g of the activated Sepharose. We passed cell lysates of CEM cells through the column and eluted the components that bound. The elution product was analyzed by SDS-PAGE electrophoresis, Western blotting, ELISAs, and BiaCore reactivity against CEM cell lysates.

[0241] The elution product that was purified from CEM cell lysates was demonstrated to be CD147 upon our sequencing of the diffuse band corresponding to 45-55 KD that we observed on Western Blot analysis after reaction with each of the 2.6.1 antibody and the ABX-CBL antibody. As will be observed from FIG. 1, the 2.6.1 antibody bound most intensely to a molecule or molecules contained within a diffuse band from about 45-55 KD, while the ABX-CBL antibody showed binding with a lower intensity to a similar band from about 45-55 KD.

[0242] Sequencing was accomplished upon a portion of the 45-55 KD band that was isolated through use of preparative gel electrophoresis and electroblotting techniques using a Perkin Elmer sequencer. We obtained a partial amino

acid sequence of the molecule (between 35 through 40 residues). The resulting sequence information was analyzed through a protein database search (Protein Identification Resource (PIR) R47.0, December 1995) and the sequence comparison data indicated that the molecule was CD147.

[0243] Western blots on CEM lysates were generally accomplished as follows:

[0244] CEM cells were homogenized in 10 mM Tris pH 7.5, 150 mM NaCl, 1% Triton X-100, and protease inhibitors to generate CEM extracts at 5×10^8 cells/ml. The extract (5 μ l) were electrophoresed on 12% SDS-PAGE gels and then blotted onto PVDF. The blot was cut into 5 strips in preparation for antibody staining. All first antibody staining was done at 14 g/ml in 1% gelatin/PBST buffer. All AP labeled second antibody was done at a dilution of 1:1000 in the same buffer. The rabbit-anti-mouse-hnRNP-k Protein antibody was supplied to us by Dr. Karol Bomstzyk at the University of Washington. Each of the ABX-CBL, Pharmingen, and 2.6.1 antibodies are described further herein.

Experiment 3 Purification of 62 KD Band

[0245] In order to purify the material contained in the 62 KD band, CEM whole cell lysates were prepared from approximately 3×10^{10} cells. The lysates were extracted and concentrated to provide about 3.8 mg of protein. A portion of the recovered protein was subjected to a series of chromatography steps: size exclusion, anion exchange, hydrophobic interaction, reversed phase, and microbore reversed phase. In each step, the fraction showing binding to the ABX-CBL antibody on Western blot was carried on to the next step. Following microbore reversed phase chromatography, approximately 5×10^{-6} grams of protein was recovered and a portion of the protein subjected to gel electrophoresis and electroblotting to generate approximately 90% pure 62 KD protein.

[0246] A direct N-terminal sequence was attempted, however, the molecule possessed a blocked N-terminus. Thus, the material was digested with CNBr and preparative gel electrophoresis and electroblotting were conducted, yielding bands at approximately 12 KD and 32.5 KD. The blotted fragments were sequenced and the resulting sequence results were analyzed through protein database searches (Protein Identification Resource (PIR) R47.0, December 1995). The sequence comparison data indicated that the molecule was heterogeneous ribonuclear protein k (hnRNP-k), with the 12 KD band having residues 360 and up (after Methionine; 359) and the 32.5 KD band having residues 43 and up (after Methionine; 42).

Experiment 4 CD147 ELISA Assay

[0247] We have utilized the enriched purified antigen obtained from CEM cell lysates to develop a specific ELISA assay for the detection of the expression of CD147 in a secreted or membrane bound form. In the assay, we immobilize the CD147 antigen (for example, the CD147 antigen that is affinity purified from CEM cell lysates) in the wells of plates. Binding of the antigen can be accomplished using conventional techniques. Thereafter, the plates containing the antigen can be used for the detection of antibodies that are reactive with it using conventional techniques. We have demonstrated that each of the commercially available anti-CD147 antibodies (RDI-CBL535 (a murine anti-CD147 IgG2b antibody), available from RDI, Flanders, N.J., and

36901A (a murine anti-CD147 IgG1 antibody), available from Pharmingen, San Diego, Calif.), the ABX-CBL antibody, and the human antibodies that we have generated in Example 2 react specifically in this assay.

[0248] The present ELISA assay is useful as a screening system for detecting antibodies that bind to the CD147 antigen.

Experiment 5 Evidence Related to Role of 35 KD Band

[0249] As mentioned above, anecdotal evidence indicates that a 35 KD band could correspond to a singly-glycosylated form of CD147. See Kanekura et al. *Cell Struct Funct* 16.23-30 (1991). Further, it is also interesting to note that in comparisons of Western blots produced by two commercially available anti-CD147 antibodies (RDI-CBL535 (a murine anti-CD147 IgG2b antibody), available from RDI, Flanders, N.J., and 36901A (a murine anti-CD147 IgG1 antibody), available from Pharmingen, San Diego, Calif.) to the ABX-CBL and 2.6.1 antibodies indicates that each of the commercially available antibodies recognize a molecule that has a molecular weight around 35 KD and appearing similar to the 35 KD band recognized by the ABX-CBL antibody. See FIG. 1. Another interesting observation is that in the immunoaffinity purification mentioned above, when the effluent product from the 2.6.1 antibody was probed with the ABX-CBL antibody, the 35 KD band was no longer visible by Western blot. Rather, the ABX-CBL antibody appeared to bind to the diffuse band from 45-55 KD with relatively low intensity (similar to that shown in FIG. 1). This evidence indicates that the ABX-CBL antibody could bind preferentially to a different epitope on, or a different form of, CD147 than the 2.6.1 antibody and the commercially available antibodies.

Experiment 6 Complement Mediated Cell Killing

[0250] The UCLA group mentioned above (see e.g., U.S. Pat. Nos. 5,330,896 and 5,643,740) provided certain evidence that the CBL1 antibody operated through killing of certain activated cell populations while the antibody did not react with non-activated cells. For example, in a microcytotoxicity assay, the CBL1 antibody was disclosed to kill activated lymphocytic cells but not other normal cells.

[0251] In connection with this experiment, the following materials and procedures were utilized:

Mixed Lymphocyte Reaction

[0252] Mixed lymphocyte reaction (MLR) is an in vitro system for assaying T lymphocyte proliferation in cell-mediated responses. A cell-mediated response is an in vitro assay of effector cytotoxic function, which can also be assayed in vivo by graft-versus-host reaction in experimental animals. When co-culturing allogeneic lymphocytes in MLR the cells undergo extensive blast transformation and cell proliferation. Thus, MLR can be quantified by adding tritium-labeled thymidine ($[^3\text{H}]$ thymidine) to the culture medium and monitoring uptake of label into DNA of the dividing lymphocytes.

[0253] To determine the function and quality CBL-1 and ABX-CBL antibody we used MLR to test the ability of CBL-1 and ABX-CBL to inhibit lymphocyte proliferative responses. Peripheral blood mononuclear cells were isolated from two HLA mismatched individuals by Ficoll-Paque gradient centrifugation. Allogeneic lymphocytes were

mixed (1:1) and co-cultured (total of 5×10^5 cells/well in 96-well plate) in vitro for six days. Lymphocytes from one individual were irradiated with 3000 rads prior to the culture. CBL-1 and ABX-CBL antibody plus either 10% rabbit or 25% human complement were added to the culture 24 h prior to the end of the culture. The culture was pulsed with [3 H]methyl-thymidine (Amersham) overnight and harvested on day 6. Lymphocyte proliferative response was determined by measuring [3 H]-thymidine incorporation. Percentage inhibition was calculated as the cpm in the absence of antibody minus the cpm in the presence of antibody divided by the cpm in the absence of antibody.

ConA Stimulated Lymphocyte Proliferation

[0254] Human PBMC were isolated as described above and stimulated by the mitogen Concanavalin A (ConA) at 5 μ g/ml for 48 h. Antibodies with or without 10% complement were added to the culture 24 h prior to the end of the culture. The culture was pulsed with [3 H]-methyl-thymidine overnight and harvested next day. Lymphocyte proliferative response was determined by measuring [3 H]-thymidine incorporation. Percentage inhibition was calculated as the cpm in the absence of antibody minus the cpm in the presence of antibody divided by the cpm in the absence of antibody.

FACS Analysis of Cell Surface Molecules

[0255] For cell surface expression of different surface molecules, immunofluorescent staining and analysis on a FACSvantage (Becton Dickinson, San Jose, Calif.) have been described (FACSscan Manual. Becton Dickinson, San Jose, Calif.). Monoclonal antibodies anti-CD3-PE, anti-CD4-PE, anti-CD8-PE, anti-CD14-PE, anti-CD20-PE, anti-CD25-FITC and anti-CD25-PE were obtained from Becton Dickinson. Anti-CD55-FITC and anti-CD59-FITC were purchased from Pharmingen (San Diego, Calif.). ABX-CBL and cem2.6.1 were conjugated with FITC and PE, respectively, at Abgenix.

Complement-dependent Cytotoxicity Assay Using Alamar Blue

[0256] Complement-dependent cytotoxicity (CDC) assay was performed as described (Gazzano-Santoro et al. "A non-radioactive complement-dependent cytotoxicity assay for anti-CD20 monoclonal antibody" *J. Immunol. Methods* 202:163-171 (1997)). Fifty microliters of a cell suspension of 10^6 cells/ml, 50 μ l of various concentrations of antibodies and 50 μ l of a 10% rabbit or human complement were added to flat-bottomed 96-well tissue culture plate and incubated for 2 hours at 37° C. and 5% CO₂. Fifty microliters of Alamar blue (Accumed International) were then added (final 10%) and the incubation continued for another 5 hours. The plates were allowed to cool to room temperature for 10 minutes on a shaker and the fluorescence was read using a 96-well fluorometer with excitation at 530 nm and emission at 590 nm. Results were expressed in relative fluorescence units (RFU).

[0257] In our work, we have demonstrated that CBL1 and ABX-CBL operate through complement mediated cell killing. Use of the CBL1 antibody by itself, the isotype-matched control mouse IgM antibody by itself (FIG. 2), or complement (either human or rabbit) by itself in the MLR or modified MLR assay (ConA induced lymphocyte proliferation assay) is ineffective in inhibiting T-cell proliferation.

See FIGS. 2-5. However, when both complement and the CBL1 or ABX-CBL antibody are present, T-cell proliferation is inhibited in a dose dependent manner. See FIGS. 2-5. The human IgG2 antibody 2.6.1 is ineffective in inhibiting T-cell proliferation in the same assay, either by itself, or in combination with complement. See FIG. 5. This is expected, since the 2.6.1 antibody as a gamma-2 is notoriously less efficient in complement mediated lysis than is an IgM antibody, such as the ABX-CBL antibody.

[0258] The combination of CBL1 or ABX-CBL and complement only kill activated T-cells (both CD4⁺ and CD8⁺), activated B-cells, and monocytes, but does not effect resting T-cells and B-cells because such cells do not express CD147. It is important, to note that monocytes are also killed by ABX-CBL and complement. This data provides an explanation for the operation of ABX-CBL therapy in diseases, such as GVHD, because, ABX-CBL selectively depletes those effector cells (activated T- and B-cells) and the antigen presenting cells (monocytes and B-cells) which ordinarily would lead to further T-cell activation.

Experiment 7 Evidence Related to Cellular Activation

[0259] Using techniques described in Experiment 6, we also demonstrated that the CD25 marker appears to be expressed in high levels in the same cellular populations as those expressing the antigen to which the ABX-CBL antibody binds. See FIG. 6. This finding provided a useful marker to detect whether the cells expressing CD25 were depleted in connection with the MLR assay. Where the MLR assay is conducted utilizing a variety of activated cell populations, CD25 expressing cell populations are depleted only in those treated with the ABX-CBL antibody plus complement. See FIGS. 7-11. The selective killing of different cell populations are shown in FIGS. 10-12.

Experiment 8 Evidence Related to the Role of Expression Levels of CD147

[0260] We have also considered whether CD147 expression levels are higher in given populations of cells (which could also be relevant to CDC).

[0261] In flow cytometry studies with peripheral blood mononuclear cells (PBMC) with the ABX-CBL antibody, we have noticed that, prior to the addition of complement, there are populations of cells that appear to express high and low levels of CD147. After complement is added, there are populations of cells that appear to correspond to the low level expressers mentioned above. It appears that these results could be indicative of density of CD147 expression levels on the cell surface. Density can play a role in CDC through providing additional antigen binding sites to allow for distortion of the antibody which is the first step in triggering the complement cascade. Upon distortion of the antibody, the factor clq binds first and the cascade proceeds.

[0262] Whether the expression level (or, density) of CD147 in cellular populations plays a role in the therapeutic efficacy of the ABX-CBL antibody can be assayed through analyzing the expression levels of the CD147 molecule in various cellular populations. Generally, the experiments are conducted where beads having various known quantities of the CD147 antigen on their surface are prepared and analyzed on FACS (i.e., utilizing a FITC-labeled anti-CD147 IgG antibody) in order to generate approximately 10-20 data points of different quantities of antigen on the beads. A linear

regression curve is prepared from such data. Thereafter, cells expressing the CD147 antigen can be run through FACS and the relative quantities of antigen on the surface of the cells can be calculated from the linear regression curve.

Experiment 9 Evidence Related to the Role of Complement Inhibitory Molecules

[0263] Further, in order to consider the cellular specificity of the mode of operation of the ABX-CBL antibody, we investigated various cells to which the ABX-CBL antibody binds and considered whether such cells were killed in a manner similar to complement mediated lysis. In connection with this work, we have investigated various cells to which the ABX-CBL antibody binds and considered whether such cells were (i) killed and (ii) if so, was the mechanism similar to complement mediated lysis. In the experiment, we looked for ABX-CBL antibody binding to a number of cells (and, thus, the antigen to which the ABX-CBL antibody binds is expressed upon such cells). Those cells to which ABX-CBL would bind were then tested for complement mediated lysis through treatment with the ABX-CBL antibody and complement. Two T-cell lines (CEM and Jurkat cells), a monocyte line (U937 cells), and three tumor cell lines (A431 (epidermal), SW948 (colon), and MDA468 (breast)), each of which bound the ABX-CBL antibody were examined. Despite the expression on such cells lines, the ABX-CBL antibody is very specific about which cells are killed, being restricted to the CEM T-cell line and U937 monocyte line. See FIG. 13. We also analyzed two endothelial cell lines (i) ECV-304 (ATCC CRL-1998) is a spontaneously transformed immortal EC established from the vein of an apparently normal human umbilical cord and carrying EC characteristics and (ii) HUV-EC-C (ATCC CRL-1730) is an EC line derived from the vein of a normal human umbilical cord. Using FACS, we found that each of the ECV-304 and HUVEC-C lines stained positive against the 2.6.1, Pharmingen, and ABX-CBL antibodies suggesting that these ECs do express CD147 on the surface. FIGS. 15 and 16, respectively. We then carried out in vitro Alamar-blue based CDC assay and demonstrated that both EC lines were resistant to ABX-CBL mediated CDC in the presence of human complement. See FIGS. 17 and 18, respectively.

[0264] In order to farther understand why cells that all appear to express CD147 would not be killed by the ABX-CBL antibody in the presence of complement, we looked into CD46, CD55, and CD59 expression in such cells. Each of CD46 (membrane cofactor protein, MCP), CD55 (decay accelerating factor, DAF), and CD59 (membrane attack complex inhibitor, MACI) have been implicated as complement inhibitory molecules. See e.g., Liszewski et al. *Annu. Rev. Immunol.* 9:431 (1991) and Loveland et al. "Coordinate functions of multiple complement regulating molecules, CD46, CD55 and CD59" *Transpl. Proc.* 26:1070 (1994) related to CD46, Kinoshita et al. "Distribution of decay-accelerating factor in the peripheral blood of normal individuals and patients with paroxysmal nocturnal hemoglobinuria" *J. Exp. Med.* 162:75 (1985) and Loveland et al. "Coordinate functions of multiple complement regulating molecules, CD46, CD55 and CD59" *Transpl. Proc.* 26:1070 (1994) related to CD55, and Whitlow et al. "H19, a surface membrane molecule involved in T-cell activation, inhibits channel formation by human complement" *Cell. Immunol.* 126: 176 (1990), Loveland et al. "Coordinate functions of multiple complement regulating molecules, CD46, CD55

and CD59" *Transpl. Proc.* 26:1070 (1994), and Davies, A. and Lachmann, P. J. "Membrane defense against complement lysis: the structure and biological properties of CD59" *Immunol. Res.* 12: 258 (1993) related to CD59. Accordingly, we considered whether there was differential expression of either, or both, of these molecules on the cell lines tested above. Indeed, all of the cells, except the CEM line and the U937 line, expressed both of the molecules. And, indeed, the endothelial cell lines HUVEC-C and ECV-304 expressed all three, CD46, CD55, and CD59. FIGS. 19 and 20, respectively. In contrast, the CEM line expressed only CD59 and the U937 line expressed only CD55. See FIG. 14. This data is useful in connection with the prediction of cells that could be selectively eradicated by ABX-CBL and consequently targeted in connection with anti-CD147 in accordance with the present invention.

Experiment 10 Cloning and Expression of CD147 in Eukaryotic Cells and Binding of Antibodies

[0265] In the present experiment, we cloned full length CD147 cDNA through use of PCR in connection with the Jurkat Zapp Express phagemid DNA (Stratagene).

[0266] The following PCR primers were utilized, based on the CD147 sequence reported by Miyauchi et al. *J. Biochem.* 110:770-774 (1991) (Gene Bank Accession No. D45131):

5' : 5' -GACTACGAATTCTTGTAGGACCGCGAGG (SEQ ID NO:42)
AATAGG-3'

3' : 5' GACTACGGGCCCGGTGAGAACTTGAATCT (SEQ ID NO:43)
TGCAAGC-3'

[0267] A 949 base pair PCR product was isolated whose open reading frame encoded the 269 amino acid CD147 protein. The PCR product was digested with EcoRI and ApaI and ligated into the EcoRI and ApaI sites of mammalian expression vectors pWBFNP (FIG. 21) and pBKCMV (Stratagene) (FIG. 22) (digested with NheI/SpeI to remove the lac promoter and the lacZ ATG between positions 1300 and 1098) to create the vectors CD147/pWBFNP and CD147/pBKCMV(delta-NheI/SpeI) respectively. In the constructs, eukaryotic expression of CD147 is driven from the cytomegalovirus (CMV) immediate early promoter. CD147/pWBFNP, CD147/pBKCMV(delta-NheI/SpeI) and control vectors pWBFNP and pBKCMV were transiently transfected into monkey kidney (COS-7) cells by the CAPO₄ method. Cells were harvested 60 hours later, washed in PBS and stained with anti-ABX-CBL-FITC, anti-CEM2.6.1/anti-HulG-FITC, or anti-CD147-FITC (Pharmingen) and analyzed by FACS analysis and Western blot analysis (see FIG. 23A). The blot was accomplished using procedures described in Example 3.

[0268] FACS analysis revealed an increase in specific cell surface staining with all three antibodies only on COS cells transfected with vectors expressing CD147 cDNA (CD147/pWBFNP and CD147/pBKCMV (delta-NheI/SpeI)). COS cells transfected with CD147 cDNA showed binding to each of the antibodies in each of the FACS and Western blot analyses. In contrast, COS cells transfected with control vectors were negative for binding with each of the 2.6.1 and ABX-CBL antibodies. With respect to the Pharmingen antibody, certain background staining was observed in cells transfected with control vectors on FACS and no binding on

Western blot analysis. The transfected cells showed significant binding over background on FACS and were positive on Western blot analysis. Our results confirm that the ABX-CBL and the 2.6.1 antibodies bind to CD147.

Experiment 11 Cloning and Expression of CD147 in Eukaryotic Cells and Binding of Antibodies

[0269] Utilizing a slightly modified vector, we also transfected *E. coli* cells with the CD147 cDNA. In the experiment, CD147 cDNA generated as above was subcloned into pBKCMV (Stratagene) (FIG. 22). CD147/pBKCMV plasmid DNA was transformed into *E. coli* strain XL1-Blue MRF⁺ (Stratagene). Cultures were grown in LB media supplemented with kanamycin at 50 µg/ml to OD₆₀₀ of 0.7 then for an additional 3 hours in the presence of 1 mM isopropyl-B-D-thio-galactopyranoside (IPTG). Cells were harvested by centrifugation and stored frozen at -20° C. The *E. coli* cells so transfected were capable of expression of the CD147 molecule as evidenced by Western blotting analysis of each of the ABX-CBL, 2.6.1, and Pharmingen antibodies. Since the prokaryotic *E. coli* cells should not glycosylate the expressed CD147, it was expected that the molecular weight of the CD147 expressed by the *E. coli* should closely approximate the predicted, unglycosylated molecular weight of CD147 of about 27 KD. Indeed, in each case, binding of the three antibodies on Western blot analysis was observed to a band between about 27 and 30 KD. FIG. 23B. The blot was accomplished using procedures described in Example 3.

[0270] This data further confirms that the ABX-CBL and the 2.6.1 antibodies bind to CD147. Further, the evidence indicates that ABX-CBL binding to CD147 is not directly based on carbohydrate binding, i.e., that ABX-CBL does not bind directly to a carbohydrate epitope on CD147. Such data, however, does not eliminate the possibility that binding to CD147 is influenced by the presence of carbohydrate or glycosylation.

Experiment 12 Epitope Analysis

[0271] In order to further elucidate the binding of the ABX-CBL antibody to CD147, we undertook phage display experiment. Such experiments were conducted through panning a phage library expressing random peptides for binding with the ABX-CBL and 2.6.1 antibodies to determine if we could isolate peptides that bound. If successful, certain epitope information can be gleaned from the peptides that bind.

[0272] In general, the phage libraries expressing random peptides were purchased from New England Biolabs (7-mer and 12-mer libraries, Ph.D.-7 Peptide 7-mer Library Kit and Ph.D.-12 Peptide 12-mer Library Kit, respectively) based on a bacteriophage M13 system. The 7-mer library represents a diversity of approximately 2.0×10^9 independent clones, which represents most, if not all, of the $20^7 = 1.28 \times 10^9$ possible 7-mer sequences. The 12-mer library contains approximately 1.9×10^9 independent clones and represents only a very small sampling of the potential sequence space of $20^{12} = 4.1 \times 10^{15}$ 12-mer sequences. Each of 7-mer and 12-mer libraries were panned or screened in accordance with the manufacturer's recommendations in which plates were coated with an antibody to capture the appropriate antibody (goat anti-human IgG Fc for the 2.6.1 antibody and goat anti-mouse µ chain for the ABX-CBL antibody) followed by washing. Bound phage were eluted with 0.2 M glycine-HCl,

pH 2.2. After 3 rounds of selection/amplification at constant stringency (0.5% Tween), through use of DNA sequencing, we characterized a total of 5 clones from the 7-mer library and 6 clones from the 12-mer library reactive with the ABX-CBL antibody and a total of 6 clones from each of the 7-mer and 12-mer libraries reactive with the 2.6.1 antibody. Reactivity of the peptides was determined by ELISA. For an additional discussion of epitope analysis of peptides see also Scott, J. K. and Smith, G. P. *Science* 249:386-390 (1990); Cwirla et al. *PNAS USA* 87:6378-6382 (1990); Felici et al. *J. Mol. Biol.* 222:301-310 (1991), and Kuwabara et al. *Nature Biotechnology* 15:74-78 (1997).

[0273] No consensus sequence was readily apparent for reactivity of the 2.6.1 antibody with CD147. However, sequence alignment of the characterized 7-mer and 12-mer sequences against the amino acid sequence of CD147 yielded a number of matches for a single sequence within CD147 from residue number 177 through residue number 188 (ITLRVRSH (SEQ ID NO:1)). In particular, each of the 7-mers contained sequence matches (represented by *) to 3 or more residues within this sequence of CD147:

7-mer sequences						
			*	*	*	
1.	EE	RLR	S	Y		(SEQ ID NO:2)

2.	YE	RVR	W	Y		(SEQ ID NO:3)
			*	*	*	
3.	EE	RLR	S	Y		(SEQ ID NO:4)
			*	*	*	
4.	AE	RIR	S	I		(SEQ ID NO:5)
			*	*	*	
5.	EE	RLR	S	Y		(SEQ ID NO:6)

[0274] Further, 4 of the 12-mers contained sequence matches (represented by *) to 3 or more residues within this sequence of CD147, with 4 matches for 12-mer peptide number 1 and for 6 matches of 12-mer peptide number 2:

12-mer sequences						
			*	*	*	*
1.	TVHGD	RLR	S	LP		(SEQ ID NO:7)
			*	*	*	*
2.	TNDIGL	RQR	S	HS		(SEQ ID NO:8)
			*	*	*	*
3.	SPLLDGQ	RER	S	Y		(SEQ ID NO:9)
			*	*	*	*
4.	YDLPM	RSR	S	YPG		(SEQ ID NO:10)
			*			
5.	SLAPLWY	YSR	H	G		(SEQ ID NO:20)
6.	HTPETAPLPATV					(SEQ ID NO:21)
						(no binding)

[0275] These results indicate a consensus sequence of RXRS (SEQ ID NO:11) that is present in 10 of the sequenced clones. Accordingly, we had a synthetic peptide prepared (AnaSpec Incorporated, San Jose, Calif.) which spanned residues 169-183 of CD147 with the following sequence (with —OH representing carboxy terminus):

KGSDQAIITLRVRSRSH-OH (SEQ ID NO:12)
 | |
 170 184

[0276] Below, the amino acid sequence of CD147 is provided with the 15-mer peptide's sequence indicated by double underlining and the RXRSRSH (SEQ ID NO:13) consensus sequence indicated in bold. In addition, putative N-linked glycosylation sites of CD147 are shown as underlined and italics:

CD147 Sequence (SEQ ID NO:14)
 MAAALFVLLGFGALLGTHGASGAAGTVFTTVEDLGSKILLTCSLNDSTAEV
 TGHRLWLKGGVVLKEDALPGQKTEFKVDSDDQWGEYSCVFLPEPMGTANIQ
 LHGPPIRVKAVKSSSEHINEGETAMLVCKSESVPVTDWAWYKITDSEDKAL
 MNGSESRRFFVSSSQGRSELHIENLNMEADPGQYRCNGTSKSGSDQAIITL
TCAGAAGAAG TGAAGTCAAGLAALWPFLLGIVAEVLVLVTII-
 FIYEKRRKPEDVLDLDDAGSAPLK
 SSGQHQNDRKGNVRQRNSS

[0277] The 15-mer peptide was assayed using ELISA and it was determined that the ABX-CBL antibody specifically bound to the peptide. Further, neither the 2.6.1 antibody nor a control murine IgM antibody bound to the peptide. However, based on a competition study between the CD147 antigen and the 15-mer peptide, the ABX-CBL antibody's binding to the 15-mer peptide can only be measured when the 15-mer peptide is coated on plates and not when the peptide is in solution. Indeed, in competition experiments in which the ABX-CBL antibody is bound to either the peptide or the CD147 antigen coated to plates, the ABX-CBL antibody is not removed or replaced by the peptide in solution even at high concentrations. Nevertheless, the binding of the ABX-CBL antibody to the 15-mer peptide can be specifically competed by the CD147 antigen and positive phage preparations mentioned above but not with non-specific antigen (i.e., L-Selectin isolated from cell membrane or human plasma) or the negative phage preparations mentioned above. Similarly, the binding of the ABX-CBL antibody to the CD147 antigen can be specifically competed by positive phage preparations as compared to negative phage preparation in competition assays using preincubation.

[0278] These results indicate that while the sequence within CD147 that contains the consensus sequence RXRSRSH is important to the binding of the ABX-CBL antibody to CD147, it does not fully explain ABX-CBL's binding to CD147. Indeed, the data also suggests that the consensus sequence contained either in the 15-mer peptide when bound to the plate or the reactive phage materials when tethered to the phage coat protein binds more tightly to the ABX-CBL antibody than does the free peptide in solution. Taken together, while not wishing to bound to any particular theory or mode of operation, it is possible that CD147 possesses certain conformations that are not well mimicked in the 5-mer peptide in solution. Nevertheless, the above epitopic information is important to understanding the manner in

which the ABX-CBL antibody binds to CD147 and to producing other candidate molecules against CD147 as a therapeutic target.

[0279] It is interesting to note that in addition to the results above in connection with the presence of the RXRSRSH consensus sequence within CD147, we also looked for the presence of the consensus sequence within the hn-RNP-k protein to which ABX-CBL also appears to bind. Such analyses were conducted by sequence alignment against the phage derived peptides discussed above. Two sequences were found which possessed statistically interesting matches:

[0280] First, there was a match (indicated by *) of 5 amino acids with the 7-mer peptide number 4:

* * * (SEQ ID NO:15)
 PE RIL SI
 |
 84

[0281] Second, there was a match (indicated by *) of 5 amino acids with the 12-mer peptide number 1:

* * * * (SEQ ID NO:16)
 GGS RAR NLP
 |
 300 306

[0282] The amino acid sequence of the hn-RNP-k protein is provided below with such sequences indicated by double underlining. In addition, a number of RXR sequence motifs are present in the hn-RNP-k protein's sequence which are also indicated by underlining:

hn-RNP-k Protein Sequence (SEQ ID NO:17)
 METEQPEETFPNTETNGEFGKRPADMEEEQAFKSRNNTDEMVELRILLQ
 SKNAGAVIGKGGKNIKALRTDYNASVSPDSSGPERILSISADIETIGEI
 LKKIIPITLLEGLQLPSPTATSQLPLESDAVECLNYQHYKSGDFDCELRL
 IHQSLAGGIIGVKAKIKELRENTQTITKLFQECPPHSTDRVVLIGGKPD
 RVVECICKIILDLISESPIKGRAQPYDPNFDYDYGFTMMFDDRGRGP
 VGFPMPGRGGFDRMPGRGGRPMPPSRDYDDMSPRRGPPIPPPPGRGGRG
GSARNLPLPPPPPPRGDLMAYDRRGRPGDRYDGMVGF
 SADETWDSAITWSPSEWQMAPYEPQGGSGYDYSYAGGRGSYGLGGPIITT
 QVTIPKDLAGSIIGKGGRIKQIRHESGASIKIDEPLEGSEDRIITITGT
 QDQIQNAQYLLQNSVKQYSGKFF

[0283] Without wishing to be bound to any particular theory or mode of operation, it is possible that the binding of the ABX-CBL antibody to the hn-RNP-k protein is partially explained by the presence of these motifs within the protein.

Experiment 13 Expression of CD147 and Binding of Antibodies

[0284] Indeed, the desirability of mimicking ABX-CBL binding and efficacy is highlighted based upon a preliminary

tissue distribution study of the ABX-CBL antibody. In the study, ABX-CBL is widely distributed throughout a variety of tissues. However, the majority of the distribution is likely to be due to nonspecific binding. Nevertheless, there appears to be specific binding in endothelial cells (venules, arterioles, but not capillary beds), smooth muscle, and some mesothelium. Also, the lymphoreticular tissues appear to be bound, although, the staining seems to be restricted to large lymphocytes, presumably activated blasts. From the study conducted, it was difficult to distinguish intracellular from extracellular staining. A certain amount of cytoplasmic staining was clearly evident and could have been related to hn-RNP-k binding.

Experiment 14 Analysis of Activity of MOPC21 Light Chain Activity in ABX-CBL Antibody

[0285] Two different techniques were utilized to endeavor to study the role of the MOPC21 light in ABX-CBL activity. In each technique, efforts were made to segregate the MOPC21 light chain from the cell line producing the IgM antibody. In the first technique, segregation was effected by fusion of the ABX-CBL IgM producing cell line with another cell line (NSO). In the second technique, segregation by spontaneous loss variants was endeavored. The fusion technique was successful and work was stopped on the second technique.

[0286] In the fusion technique, in general, NSO cells were transfected with a puromycin containing vector to create a puromycin⁻ NSO cell line. The ABX-CBL IgM producing cell line was grown in HAT medium was fused with the puromycin⁺ NSO cell line.

[0287] In general, fusions are accomplished in accordance with the following techniques and procedures:

Preparation of Cells

[0288] Prior to fusion, parental cell lines for use in the fusion are grown up and maintained in medium containing DMEM high, 10% FBS, 1% non-essential amino acids, 1% pen-strep, and 1% L-glutamine.

[0289] On the day prior to fusion, each of the parental cell lines are prepared and split to provide a cell density of approximately 10^5 cells/ml. On the day of the fusion, cells are counted and the fusion is commenced when, and assuming, that cell count for each of the parental cell lines are within the range of about $1.5-2.5 \times 10^5$ cells/ml. Sufficient quantities of each of the parental cell lines to make up 5×10^6 cells each are withdrawn from the cultures and added to a 50 ml centrifugation tube and the cells are pelleted at 1200 rpm for approximately 5 minutes. Concurrently with the preparation of the cells, incomplete DMEM, PEG, and double selection media are prewarmed in an incubator bath. Following pelleting, cells are resuspended in 20 ml incomplete DMEM and pelleted again. Thereafter, the cells are resuspended in 5 ml incomplete DMEM and the two parental cell lines are pooled in a single tube and pelleted again to form a co-pellet containing both of the parental cell lines. The co-pellet is resuspended in 10 ml incomplete DMEM and again pelleted. All of the supernatant is then removed from the co-pellet and the cells are ready for fusion.

Fusion

[0290] Following removal of all of the supernatant, 1 ml PEG- 1500 is added over the course of 1 minute to the

co-pellet while stirring. After addition of the PEG is completed, either gentle stirring with a pipet is continued for 1 minute or the suspended co-pellet can be allowed to stand for 1 minute. Thereafter, 10 ml of incomplete DMEM is added to the co-pellet over the course of 5 minutes with slow stirring. The mixture is then centrifuged at about 1200 rpm for 5 minutes and following centrifugation, the supernatant is aspirated off, and 10 ml of complete double selection medium is added and gently stirred into the cells. The cells are then plated at 100 μ l/well into 10 96-well microtiter plates and placed into an incubator (37° C. with 10% CO₂) where they are not disturbed for 1 week. After the passage of a week, plates are fed by adding 100 μ l of complete double selection medium to each well.

[0291] Double selection medium is prepared depending upon the marker gene utilized in connection with the parental cell lines. In the majority of our experiments, the selectable markers conferring puromycin, hygromycin, of hypoxanthine and thymidine resistance are utilized. Concentrations required to obtain complete cell killing of NS/0 cells were determined through use of kill curves and resulted in our use of 6 micrograms/ml of puromycin and 350 micrograms/ml of hygromycin. In connection with HPRT resistance, we used HAT media supplement from Sigma using standard conditions.

[0292] In the present case, cells were selected for puromycin⁺/HAT resistance. Individual clones were picked based on selection and clones were expanded in 96-well plates. Plates were split ($\frac{1}{2}$ for freezer stock, $\frac{1}{2}$ for growth). Total RNA was isolated from the growth plates using the Qiagen 96-well RNA isolation kit according to the manufacturer's instructions. Primers were designed based on conserved sites on the MOPC21 and the ABX-CBL kappa chains that would amplify fragments of the chains which contained unique restriction sites in the respective chains, as follows:

Restriction site	Chain	Position
AgeI (BsrFI)	MOPC21	135
BstYI	MOPC21	173
KpnI	ABX-CBL	85
NsiI	ABX-CBL	130
XcmI	MOPC21	58

[0293]

5 prime: 5'-GCA GTC TCC TAA ACT GCT (SEQ ID NO:44)
positions 99-116 allows
analysis of BstYI
restriction site;

or

5 prime: 5'-ACC TGC AAG GCC AGT (SEQ ID NO:45)
positions 40-54 allows
analysis of NsiI or KpnI
restriction sites,

3 prime: 5'-CAC TCA TTC CTG TTG AAG. (SEQ ID NO:46)

[0294] Accordingly, through amplification with the above primers, followed by digestion with the appropriate restriction enzymes, presence or absence of MOPC21 or ABX-

CBL could be readily detected on agarose gel electrophoresis. Through use of the above techniques, at least 6 variants were obtained that lost the MOPC21 light chain expression but retained the ABX-CBL kappa. No variants were directly obtained that lost ABX-CBL kappa chain expression and retained the MOPC21 chain expression. However, we isolated a cell line that appeared to be a minimal producer of ABX-CBL light chain and subcloned the line. It turned out to be a mixed cell line of a heterogeneous MOPC21/ABX-CBL light chain producer and a MOPC21 light chain only producer. Accordingly, we isolated the MOPC21 only producer after subcloning.

[0295] MOPC21 only light chain containing and ABX-CBL only light chain containing antibodies were compared and supported the conclusion that the presence or absence of the MOPC21 light chain did not appear to substantially impact antibody binding or properties of the antibodies. Although, it did appear that the MOPC21 only light chain containing antibody did not bind as intensely on Western blotting to CEM cells or CD147.

Experiment 15 Generation and Characterization of Human Antibodies to CD147

[0296] In accordance with Experiment 1, we generated a panel of fully human anti-CD147 antibodies. Antibodies were screened by ELISA for binding with CD147 and FACs for ability to compete with ABX-CBL. Certain of such

through 33. In addition, CDRs, according to Kabat numbering scheme, of the heavy chains and kappa light chains of the antibodies are shown in **FIGS. 34 through 43**.

[0297] In view of a number of tests that were conducted, particularly, competition studies between ABX-CBL and the certain of the antibodies, the 2.6.1 IgM antibody was chosen for additional development.

Experiment 16 Generation of 2.6.1 Expression Vectors for the Generation of IgG1, IgM, and Multimeric IgM Antibodies

[0298] In order to investigate the ability of the 2.6.1 antibody to operate in ADCC, similar to the CBL1 and ABX-CBL antibodies, we were interested in preparing IgM and IgG1 isotypes of the 2.6.1 antibody. The isotype switching of the 2.6.1 antibody from an IgG2 to an IgG1 was relatively simple. Whereas, the switching of the 2.6.1 antibody to a multimeric IgM required certain additional steps.

[0299] As will be appreciated, all of the IgMs that were generated from XenoMouse animals were monovalent. Accordingly, in order to prepare a fully human multimeric IgM antibody, we first were required to clone the human J-chain gene from human buffy coat cells. The sequence of the human J-chain cDNA is shown below with the 5'-untranslated portion shown in bold, italics and underlining:

```

                                     (SEQ ID NO:47)
TCAGAAGAAG TGAAGTCAAGATGAAGAACC ATTGCTTTT CTGGGGAGTC      50
CTGGCGGTTT TTATTAAGGC TGTTCATGTG AAAGCCAAG AAGATGAAAG      100
GATTGTCTCT GTTGACAACA AATGTAAGTG TGCCCGGATT ACTTCCAGGA      150
TCATCCCTTC TTCCGAAGAT CCTAATGAGG ACATTGTGGA GAGAAACATC      200
CGAATTATTG TTCCTCTGAA CAACAGGGAG AATATCTCTG ATCCCACCTC      250
ACCATTTGAGA ACCAGATTTG TGTACCATTG GTCTGACCTC TGTAPAAAAT      300
GTGATCTTAC AGAAGTGGAG CTGGATAATC AGATAGTTAC TGCTACCCAG      350
AGCAATATCT GTGATGAAGA CAGTGCTACA GAGACCTGCT ACACTTATGA      400
CAGAAACAAG TGCTACACAG CTGTGGTCCC ACTCGTATAT GGTGGTGAGA      450
CCAAATGGT  GGAAACAGCC TTAACCCAG ATGCTGCTA TCCTGACTAA      500

```

antibodies were sequenced. The sequences of certain of the antibodies were compared to transcripts of the germline

[0300] The J-chain gene encodes the human J-chain with the following sequence.

```

                                     (SEQ ID NO:22)
MKNHLLFWGV LAVFIKAVHV KAQEDERIVL VDNKCKCARI TSRIIRSSSED      50
PNEDIVERNI RIIVPLNRE NISDPTSPLR TRFVYHLSDL CKKCDPTEVE      100
LDNQIVTATQ SNICDEDSAT ETCYTYDRNK CYTAVVPLVY GGETKMOVETA      150
LTPDACYPD                                     159

```

V-gene segments to somatic mutations in the amino acid sequences. Such sequence comparisons are shown in **FIGS. 44 through 46**. cDNA sequences and protein transcripts of and for each of the antibodies are shown in **FIGS. 24**

[0301] The following primers, retrofitted with the indicated restriction sites for further cloning, were designed for amplifying the human J-chain cDNA out of RT-PCR prepared materials from human Buffy coat cells:

5'- GAATTC AGA AGA AGT GAA GTC (SEQ ID NO:48)
EcoRI

3'- GTCGAC TAT GCA GTC AGC AAT GAC (SEQ ID NO:49)
Sali

[0302] The J-chain cDNA and the 2.6.1 kappa gene isolated through RT-PCR were amplified using the above primers and a 500 base pair PCR product was isolated whose open reading frame encoded the 159 amino acid J-chain protein. The PCR product was cloned into the TA cloning kit (Invitrogen) and had an EcoRI restriction site on each end. This vector was digested with EcoRI and the digest cloned into pWBFNP MCS (**FIG. 47**) that was cut with EcoRI and treated with CIP. Orientation of the insert was determined through digestion with PvuII which created differently sized fragments based on orientation (PvuII sites were present in the pWBFNP MCS vector as shown in **FIG. 47** and at position 421 in the J-chain insert. This vector was called pWBj1

[0303] The 2.6.1 kappa chain was amplified by RT-PCR using the following primers:

5 prime:
5' TGC AGG AAT CAG ACC CAG TC (SEQ ID NO:50)

3 prime:
5' GTC AGG CTG GAA CTG AGG AGC A (SEQ ID NO:51)

[0304] using the TA cloning kit providing EcoRI sites on each end of the VJCK insert. The kappa chain was sequenced. The kappa cDNA was EcoRI digested and cloned into the EcoRI site in pWBFNP MCS. Orientation was determined based on fragment size by NotI and PstI digestion of the NotI site in pWBFNP MCS and the PstI site contained at position 243 of the kappa insert shown in **FIG. 33**. This vector was called pWBK1.

[0305] In order to allow insertion of the J-chain expression cassette into pWBK1 from pWBj1, pWBK1 was cut with PacI and blunted and recut with AvrII and pWBj1 was cut with SpeI and blunted and recut with AvrII and the blunted SpeI/AvrII fragment was cloned into pWBK1 blunt PacI/AvrII to yield pWBK1(J). pWBK1(J) contained expression cassettes for both the 2.6.1 kappa chain and the J-chain.

[0306] pWBK1(J) was further modified to contain DHFR resistance through cloning DEFR through NotI digestion from a vector pWB DHFR (containing DHFR at NotI) into pWBK1(J) at the NotI site. This vector was called pWBK1(J) DHFR.

[0307] In order to make an IgG1 expression vector, the 2.6.1 heavy chain was amplified through RT-PCR using the TA cloning vector (Invitrogen) using the following primers:

5 prime:
5' TCA TTT GGT GAT CAG CAC T (SEQ ID NO:52)

3 prime:
5' GCT AGC TGA GGA GAC GGT GAC CAG G (SEQ ID NO:53)
3' gamma 1 NheI (introduces a
NheI restriction site)

The resulting product contained only the VDJ cDNA sequences and not the constant region. The sequence was confirmed by sequencing. This vector was utilized to prepare an IgG1 expression vector as described below.

[0308] pWBFNP MCS was digested with EcoRI and treated with CIP and the EcoRI digest from the TA vector, above, was cloned into the vector. Orientation was determined by size through digestion with NheI, which confirmed the insertion, followed by digestion with NotI. This vector was called pWBVDJ261NheI. pWBVDJ261NheI was cut with XhoI and blunted and recut with NheI. A human gammal construct was cloned in from a pWBFNP vector containing the gammal constant region between NheI and EcoRI sites was cut with EcoRI and blunted and recut with NheI. This vector was called pWBVDJ261G1 (or pWBjIgG1). A puromycin cassette was cloned in from a pIK6.1+puro vector (**FIG. 48**) which was cut with HindIII and blunted and recut with AvrII. The pWBjIgG1 was cut with PacI and blunted and recut with AvrII and the puro cassette was cloned therein. This vector was called pWBjIgG1 Puro.

[0309] In order to make an IgM expression vector, the 2.6.1 heavy chain was amplified through RT-PCR using the TA cloning vector (Invitrogen) using the following primers:

5 prime:
5' TCA TTT GGT GAT CAG CAC T (SEQ ID NO:54)

3 prime:
5' GGA TCC TGA GGA GAC GGT GAC G (SEQ ID NO:55)
3' Mu BamHI (introduces BamHI
restriction site)

The resulting product contained only the VDJ cDNA sequences and not the constant region. The sequence was confirmed by sequencing. This vector was utilized to prepare an IgM expression vector as described below.

[0310] pWBFNP MCS was digested with EcoRI and treated with CIP and the EcoRI digest from the TA vector, above, was cloned into the vector. Orientation was determined by size through digestion with BamHI, which confirmed the insertion, followed by digestion with NotI. This vector was called pWBVDJ261BamHI.

[0311] A human Mu construct was PCR amplified from a yeast artificial chromosome construct, YAC 2CM, described in Mendez et al., (1997), supra. and U.S. patent application Ser. No. 08/759,620, filed Dec. 3, 1996, through RT-PCR using the TA cloning vector (Invitrogen) using the following primers:

5 prime:
5' GGA TTA GCA TCC GCC CCA ACC CTT (SEQ ID NO:56)
(which introduced a BamHI
restriction site on the
5' end)

3' prime:
5' GTC GAC GCA CAC ACA GAG CGG CCA (SEQ ID NO:57)

[0312] The vector pWBVDJ261BamI was cut with BamHI and recut with XhoI. The TA cloning vector containing the Mu insert was cut with BamHI and XhoI (which is another site in the TA vector) and was cloned into the BamHI/XhoI sites of pWBVDJ261BamI. The resulting vector was called pWBVDJ261IgM (or pWBIGM). The vector was further equipped with a puromycin cassette in the same manner as described above in connection with the construction of pWBIGG1 Puro. The resulting vector was called pWBIGM Puro.

Experiment 17 Generation of Cell Line Expressing 2.6.1 IgG1 Antibodies

[0313] In order to generate a cell line expressing the 2.6.1 IgG1 antibody, we cotransfected DHFR⁻ CHO cells with The pWBIGG1 Puro vector and the pWBK1 DHFR vector through electroporation. This was accomplished by taking a stock of approximately 2×10^7 DHFR CHO cells and electroporating at 290 V, 960 μ FD, 200 μ g of linearized plasmid DNA plus 200 μ g of carrier DNA. Cells were seeded in α^+ medium and allowed to grow for two days. 8×10^5 cells were seeded in 10 cm dish in α^- medium with 4 μ g/ml puromycin selection medium. Cells were incubated for 4-5 days and then transferred to α^- medium with 0.5 μ M MTX at 5×10^5

Experiment 19 Characterization of the 2.6.1 IgG1 and Multimeric IgM Antibodies

[0317] In order to assess the function of the 2.6.1 IgG1 and multimeric IgM antibodies, we assayed the antibodies in several assays. Each of the 2.6.1 IgG1 and multimeric IgM bound to CEM cells and bound to CD25⁺ activated human peripheral blood cells in a similar manner to the CBL1 and ABX-CBL. The antibodies were assayed in a potency and a lysis assay, in the same manner described above. In connection with these experiments, the 2.6.1 multimeric IgM antibody appeared approximately as active as CBL1 and ABX-CBL. Further, the 2.6.1 multimeric IgM antibody was capable of acting in ADCC as shown in FIG. 50.

Experiment 20 Affinity Measurement of the 2.6.1 Multimeric IgM Antibodies

[0318] We also examined the affinity of the 2.6.1 multimeric IgM antibody in comparison to ABX-CBL and certain other forms of the 2.6.1 antibody. Affinity measurements were conducted as described in Mendez et al., (1997), supra. and U.S. patent application Ser. No. 08/759,620, filed Dec. 3, 1996. The results are shown in the following Table:

TABLE 2

Antibody	Ig class	On-rates ka ($M^{-1}s^{-1}$)	Off-rates kd (s^{-1})	KA kd/ka (M^{-1})	KD ka/kd (M)	BLAcore surface Hu rCD147- IgG [RU]
ABX-CBL	M IgM	7.25×10^5	3.76×10^{-4}	1.39×10^9	5.18×10^{-10}	791
ABX-CBL	M IgM	6.34×10^4	4.94×10^{-3}	1.28×10^7	7.84×10^{-8}	791
	monomer					
CEM	Hu IgG2	8.20×10^5	3.75×10^{-4}	2.19×10^9	4.57×10^{-10}	791
2.6.1						
CEM2.6.1	Hu IgG2	7.17×10^5	4.03×10^{-4}	1.78×10^9	5.61×10^{-10}	242
CEM2.6.1	Hu IgM	6.52×10^5	2.03×10^{-4}	3.21×10^9	3.12×10^{-10}	242
CEM2.6.1	Hu IgM	2.63×10^5	1.67×10^{-3}	1.57×10^8	6.39×10^{-9}	242
	monomer					
CEM2.6.1	Hu IgG1	3.13×10^5	2.01×10^{-4}	1.55×10^9	6.43×10^{-10}	242

cells per 10 cm dish. Cells were incubated for approximately 14 days for selection and, thereafter, clones were picked and expanded and assayed for ability to bind to CD147 and the presence of IgG1.

[0314] We recovered a number of clones expressing a 2.6.1 antibody with a gamma-1 isotype that bound specifically to CD147.

Experiment 18 Generation of Cell Line Expressing 2.6.1 Multimeric IgM Antibodies

[0315] In order to generate a cell line expressing the 2.6.1 multimeric IgM antibody, we cotransfected DHFR⁻ CHO cells with The pWBIGM Puro vector and the pWBK1(J) DHFR vector through electroporation. The same techniques described in Experiment 18 were utilized.

[0316] We recovered a number of clones expressing a 2.6.1 antibody with a multimeric Mu isotype that bound specifically to CD147.

Experiment 21 Human Clinical Trial with ABX-CBL Antibody

Phase II Clinical Trial of ABX-CBL

[0319] A. Background

[0320] As we mentioned above, in view of the positive results observed with respect to the CBL1 antibody, we undertook clinical trials utilizing ABX-CBL. The first such trial was a Phase II, multicenter, open label, dose escalation clinical trial examining multiple intravenous infusions of four doses of ABX-CBL in patients with steroid resistant GVHD. The trial enrolled patients with acute GVHD who were unresponsive to at least three days of treatment with corticosteroids and who had a severity index of at least B according to a modified IBMTR Severity Index (Rowlings et al. "IBMTR severity index for grading acute graft-versus-host disease: retrospective comparison with glucksberg grade" *British Journal of Haematology* 97: 855-864 (1997)).

In the trial, four different doses were administered intravenously in a dose escalation design using an induction regimen of seven days followed by a maintenance dose of twice weekly for two weeks. Patients were followed for 8 weeks after completion of the treatment course. Long-term safety follow-up has been instituted.

[0321] The study was designed with three primary objectives and four secondary objectives under review, as follows:

[0322] Primary Objectives (i) to assess the safety of multiple doses of ABX-CBL in patients with steroid resistant acute GVHD; (ii) to determine the maximum tolerated IV dose of ABX-CBL in patients with steroid resistant acute GVHD; and (iii) to determine the pharmacokinetics of multiple doses of ABX-CBL in patients with steroid resistant acute GVHD.

[0323] Secondary Objectives (i) to assess the clinical efficacy of four different doses of ABX-CBL in patients with steroid resistant acute GVHD; (ii) to assess a dose response of ABX-CBL; (iii) to assess long-term safety in patients with acute GVHD who have received multiple doses of ABX-CBL; and (iv) to assess the long-term survival in patients with acute GVHD who have received multiple doses of ABX-CBL.

[0324] Determination of the dosing for ABX-CBL was considered essential. As discussed above, the initial clinical trials conducted with the CBL1 antibody utilized ascites fluid that was not purified. Thus, the concentration of the antibody within the materials given to patients was not known. Further, because CBL1 was generated in a cell line that was not producing solely the IgM, but also an IgG, the concentration of the IgM antibody given to patients was even less clear.

[0325] In order to assess the above objectives, a four cohort trial plan was established with the following dose cohorts of patients:

[0326] Cohort 1: 0.01 mg/kg

[0327] Cohort 2: 0.1 mg/kg

[0328] Cohort 3: 0.3 mg/kg

[0329] Cohort 4: 1.0 mg/kg

[0330] Patients in all cohorts were to receive, and received, up to 11 intravenous infusions of ABX-CBL. ABX-CBL was infused over 2 hours via a syringe pump. The dosing schedule was as follows: daily times 7 days, followed by twice a week for two weeks. Safety evaluations were conducted prior to advancing to the next dose cohort. 27 patients were enrolled across all 4 of the dose cohorts.

[0331] During the conduct of the study, adverse events were observed in patients in the third cohort, receiving 0.3 mg/kg of ABX-CBL. There, several of the patients experienced myalgia or myalgia-like symptoms. As a result, the third dose cohort (0.3 mg/kg) was determined as the maximum tolerated dose. Thus, the fourth dose cohort was reduced to a dosage of 0.2 mg/kg so that the actual dosing utilized in the study was as follows:

[0332] Cohort 1: 0.01 mg/kg

[0333] Cohort 2: 0.1 mg/kg

[0334] Cohort 3: 0.3 mg/kg

[0335] Cohort 4: 0.2 mg/kg

[0336] The eligibility requirements for patients to enter the study were as follows:

[0337] One year old or older

[0338] Stem Cell transplant within 100 days

[0339] Steroid resistant acute GVHD with a severity index of B, C, or D

[0340] No experimental drugs or devices within 30 days of enrollment unless mutually agreed upon by the investigator, the sponsor's medical monitor, and the FDA

[0341] ANC>500/mm³ with or without GCSF or GMCSF

[0342] Patients were screened and assigned to a treatment cohort once the patient met the eligibility criteria. Standard post stem cell transplant treatment was continued.

[0343] Once dosing was initiated, patients were infused with ABX-CBL the applicable dose for their dose cohort daily for 7 days (referred to as an induction regimen) followed by infusions 2 times per week for two weeks (referred to as a maintenance regimen). Patients were followed for 8 weeks following their infusions (visits are weekly for 4 weeks followed by a visit 4 weeks later) for safety and clinical effect. Further, patients who received at least one infusion of ABX-CBL were scheduled to participate in a long term follow up program to evaluate the long term safety of ABX-CBL and long term survival.

[0344] Safety was assessed by monitoring adverse events while on study as well as vital signs during the infusion of ABX-CBL. Further, patients received frequent physical exams and underwent extensive laboratory studies. Laboratory studies included complete blood counts, T-cell subsets, serum chemistries, and urinalyses at regular intervals as outlined below. Baseline CPK with isoenzymes were obtained on all patients and patients who experienced any infusion related adverse experiences were reanalyzed with CPK and isoenzymes. In addition, patients were monitored for Human Anti Mouse Antibody (HAMA) response by ELISA. Further, five patients in each cohort were assigned to have pharmacokinetic blood samples for pK profile.

[0345] Clinical effect of ABX-CBL was assessed by evaluating changes to the overall score of acute GVHD based upon the modified IBMTR Severity Index (Rowlings et al. "IBMTR severity index for grading acute graft-versus-host disease: retrospective comparison with glucksberg grade" *British Journal of Haematology* 97: 855-864 (1997)), time to response, duration of response, time and incidence of flare of acute GVHD, and length of hospitalization.

[0346] B. Protocol Procedures

[0347] In connection with the trial, the following tests, observation schedules, preparations of the study medication were utilized:

TABLE 3

TESTS AND OBSERVATION SCHEDULE												
		ON STUDY										
PERIOD	SCREEN	TREATMENT						TREATMENT FOLLOW UP			LT. FOLLOW UP	
WEEK		0	1		2		3, 4, 5, 6	10		Q 6 mos		
DAY		0	1-6	9	13	16	20	23, 30, 37, 44	72	100^		
Visit #	1	2	3-8	9	10	11	12	13-16	17	0th	19+	
Procedures:												
Informed Consent	X											
Elig for enrollment	X											
Medical History	X											
Serum pregnancy	X ⁴											
Hgt/Wgt ¹	X		X ⁷	X ⁷	X ⁷	X ⁷	X ⁷	X	X	X		
Phys. Exam(C/A)	C		A ⁷	A ⁷	A ⁷	A ⁷	A ⁷	C	C	A		
Vital Signs	X	X ⁶	X ⁶	X ⁶	X ⁶	X ⁶	X ⁶	X	X	X		
Mod IBMTR score	X	X ⁵	X ⁷	X ⁷	X ⁷	X ⁷	X ⁷	X	X	X		
KPS/Lansky	X		X ⁷	X ⁷	X ⁷	X ⁷	X ⁷	X	X	X		
CBC/DIFF/PLAT	X ²	X ⁵	X ⁷	X ⁷	X ⁷	X ⁷	X ⁷	X	X	X		
Serum Chem	X ²	X ⁵	X ⁷	X ⁷	X ⁷	X ⁷	X ⁷	X	X	X		
CPK-III (mm) ³	X ²		X ³	X ³	X ³	X ³	X ³	X ³	X ³			
CD 3, 4, 8, 19		X ⁷	X ⁷	X ⁷		X ⁷		X	X			
Blood for HAMA	X ¹³	X ¹³		X ⁷				X ⁹	X ¹⁰			
Blood for pK		X ⁸	X ⁸	X ⁸			X ⁸					
UA	X ²			X ⁷		X ⁷		X ⁹	X			
Study med adm		X	X	X	X	X	X					
Intercurrent Illness	X									➔	X ¹¹	
AE'S		X								➔	X ¹²	
Concomitant meds	X									➔	X ¹¹	
ECG/CXR	X*	Data will be collected as tests completed per routine patient care										
Biopsy assessment		Data will be collected on biopsies as they occur based on pt status										

[^]This visit occurs 100 days post allogeneic stem cell transplant not post infusion of study medication

*Obtain if not completed Within 7 days of randomization (ECG if patient is ≥ 16 years)

C/A C = Complete PE, A = Abbreviated PE

¹Obtain Height at first visit only

²Obtain within 48 hours of randomization request

³Obtain at baseline, obtain if patient experiences any infusion related AE's (refer to protocol)

⁴Serum pregnancy will be obtained on females of child bearing potential (refer to protocol)

⁵Obtain if results are >8 hours from the start of the infusion of the study medication

⁶Obtain vital signs (T, P, R, BP) prior to the start of the infusion (maximum of 10 mins), q 15 mins during the first hour of the infusion, followed by 90, 120, 180, 240, 300, and 360 minutes after the start of the infusion

⁷Obtain prior to the start of the infusion (maximum of 12 hours)

⁸Obtain just prior to the start of the first infusion and the following timepoints after the completion of the first infusion; 15 and 30 minutes, 1, 2, 4, 8, 12, 18, and 24 hours (before the 2nd infusion) and at 4 hours after the completion of the Days 9 and 20 infusions (assigned patients only)

⁹Obtain at weeks 4 and 6 only

TABLE 3-continued

TESTS AND OBSERVATION SCHEDULE

¹⁰Obtain during long term follow up if + at end of study¹¹GVHD status and current treatment(s)¹²Resolve any ongoing AEs¹³Obtain HAMA during screen if the patient previously received a murine derived product. This needs to be negative in order for the patient to qualify. If the patient has never received a murine product, this is to be obtained on Day 0 prior to the start of the infusion

[0348] In connection with the trial, it was preferred that the modified IBMTR Severity Index scoring was completed by the same physician.

[0349] The following labs were to be completed by laboratory at each clinical site (local lab):

HEMATOLOGY

CBC w/ differential
White blood cells count (WBC)
WBC differential (diff)
bands/stabs
neutrophils
EOS
basophils
lymphocytes
monocytes
Red blood cell count (RBC)
Hemoglobin (Hgb)
Hematocrit (Hct)
Platelet count (Plt)

URINALYSIS

Specific gravity
PH
Protein
Glucose
Ketones

SERUM CHEMISTRY

Sodium (Na)
Potassium (K)
Chloride (Cl)
Bicarbonate (HCO₃)
Glucose
Blood Urea Nitrogen (BUN)
Creatinine (Cr)
Uric acid
Albumin
Total protein
Total bilirubin (bili)
Alkaline Phosphatase (alk phos)
Alanine aminotransferase (ALT, SGPT)
Aspartate aminotransferase (AST, SGOT)
Calcium (Ca)
Phosphate (PO₄)
CPK-III isoenzyme* (mm) (skeletal muscle)

Females only:

serum Pregnancy
(if applicable)

*Obtain CPK with isoenzymes at baseline and post infusion on any patients with infusion related AE's.

[0350] In addition, the following lab assessments were completed by a central testing laboratory:

[0351] T cell subset (CD3, 4, 8) and CD19: lymphocyte count, %, and CD4:CD8 ratio)

[0352] Further, the following lab assessment were completed by Abgenix, Inc.:

[0353] ELISA for HAMA

[0354] pK (a minimum of 5 patients/cohort to include those who previously received a murine product)

[0355] The Study Medication (ABX-CBL) was prepared and administered as follows:

[0356] ABX-CBL is a protein so it requires gentle handling to avoid foaming. The avoidance of foaming during product handling, preparation, and administration is important because foaming can lead to denaturation of the protein product. The pharmacist prepared each dose of study medication. The dose was based upon the patient's weight prior to randomization and the patient's cohort assignment, therefore the patient will receive the same dose for all 11 infusions. The pharmacist prepared the syringe and filter (filter supplied by Abgenix) and sent this to the patient unit for patient dosing.

[0357] Infusion setup: The infusion syringe was prepared using aseptic techniques. The appropriate volume of study medication was drawn up into the syringe(s), followed by the calculated volume of the pyrogen-free 0.9% sodium chloride solution, USP (saline solution). A 0.22 micron low-protein binding filter was attached and the tubing was primed to minimize fluid loss and according to the manufacturer's instructions.

[0358] Infusion volume: The total infusion volume (study medication+saline solution) to be infused for each infusion (0.01, 0.1, 0.3, 0.2 mg/kg) is equal to the patient's weight in kg. Below are examples:

TABLE 4

Pt's Weight (kg)	Cohort Assignment	Total Infusion Volume (mL)
70 kg	0.01	70 mL
70 kg	0.1	70 mL
70 kg	0.3	70 mL
70 kg	0.2	70 mL

[0359] The formula below was used to determine the volume of study medication and saline solution for each dose.

[0360] a. Dose required=patient's weight×mg/kg (mg/kg is based upon cohort assignment)

[0361] b. ABX-CBL Volume required=dose/study medication concentration (1 mg/mL)

[0362] c. Number of vials required=volume of ABX-CBL required (b above)/5 mL (each vial contains 5 mL of ABX-CBL)

[0363] d. Total volume to be administered: For all treatment cohorts the patients received a total volume which was equal to their weight in kg (a 15 kg patient will receive a total of 15 mL, a 70 kg will receive 70 mL, etc.)

TABLE 5

Example:	
Patient weighs 70 kg and is assigned to receive 0.3 mg/kg	
a.	70 kg $\times$ 0.3 mg/kg = 21 mg
b.	21 mg = 21 mL
c.	21 mL/5 mL per vial = 4.2 vials, therefore 5 vials are required
d.	70 mL (total volume) - 21 mL (study med volume) = 49 mL (saline solution)

[0364] The labeled, filled infusion syringe was sent to the patient unit for infusion, making sure that all clamps on the infusion set were closed to prevent leakage of the study medication and/or normal saline. All caps were secured in place to maintain a closed system. The sponsor provided the label for the infusion syringe and this label will contain the following:

[0365] space to record the patient study ID and initial

[0366] space to record the date and time the study medication was prepared along with the expiration date and time

[0367] space to record the initials of the person who prepared the study medication and the infusion set

[0368] Infusion instructions:

[0369] "Caution: New Drug-Limited by Federal Law To Investigational Use"

[0370] Administer infusion over 2 hours via syringe pump

[0371] Do not mix with any other medication.*

[0372] Space to specify the infusion rate based upon the total volume. For a 70 mL volume, the infusion rate would be 35 mL/hour.

[0373] The person preparing the study medication was responsible for completing the above information on the label.

[0374] For the infusion, most patients had an indwelling central line, therefore a new catheter was not be required as long as there is a dedicated line for the infusion of ABX-CBL. During the administration of ABX-CBL no other medications were to be infused via the specific port or IV line. If a central line was not available, ABX-CBL could also be infused in a peripheral intravenous line. Because this was a trial, ABX-CBL was not mixed with other medications. If another medication was previously infused in the port, the lumen was flushed with 3-5 cc of normal saline (depending on the size catheter, lumen used, and patient's size) to clear any pre-existing medications from the line and the new infusion setup from the pharmacist was attached to the port or 3-way stopcock (not piggy backed onto another line) for infusion.

[0375] The protocol was composed of four study periods: screen, treatment, treatment follow up, and long term follow up.

[0376] 1. Screen Period

[0377] The screen period began the day the patient or the patient's legal guardian signs the informed consent and ends

at treatment assignment notification. Patients could be screened for enrollment into this study up to 100 days after stem cell transplant. Patients who failed to develop steroid-resistant acute GVHD were not enrolled into the study.

[0378] Each patient must understand and have signed an IRB approved informed consent form. If the patient was a minor, the patient's legal guardian was to sign the informed consent form.

[0379] The following procedures were to be completed after the informed consent form is signed but prior to requesting treatment assignment. The results of these procedures were not more than 8 hours old unless otherwise indicated. These procedures include:

[0380] a. Complete medical history

[0381] b. Complete physical examination, which includes weight (this is the weight to be used to determine the required dose of study medication throughout the study) and height

[0382] c. Vital signs (oral temperature, resting pulse, respiration, and blood pressure)

[0383] d. Medication history and stem cell transplant treatment history from 30 days prior to requesting treatment assignment

[0384] e. Modified IBMTR Severity Index for acute GVHD

[0385] f. Assessment of intercurrent illness(es)

[0386] g. Karnofsky Performance Scale (KPS) (age ≥ 16 years) or Lansky Scale (age <16 years)

[0387] h. The following lab results were obtained if not obtained within 48 hours prior to randomization:

[0388] CBC with diff and platelets

[0389] Serum Chemistry (refer to Appendix V)

[0390] Baseline CPK-UIL isoenzyme (mm)

[0391] Serum Pregnancy test. This may be waived for women who are not of child bearing potential or who, in the opinion of the investigator, are sterile due to the pre conditioning for the stem cell transplant

[0392] Urinalysis

[0393] i. CXR if not completed within the previous 7 days

[0394] j. ECG if not completed within the previous 7 days for all patients 16 years of age or older.

[0395] k. Obtain serum specimen to be assayed by Abgenix for the determination of a positive HACA/HAMA for any patient who previously received a murine chimeric or fully murine product. This sample was to be shipped on dry ice overnight to Abgenix and results were generally available within 24 hours of Abgenix's receipt of the sample.

[0396] After the above were completed and the investigator determined that the patient was eligible for treatment, the clinical center requested (via fax) the cohort assignment

from the sponsor. The clinical center generally received notification of the treatment assignment by fax within 3 hours of the request.

[0397] 2. Treatment Period

[0398] The treatment period began when the clinical site received the patient's treatment assignment and ended when the patient completed the infusion regimen (11 doses). This period generally lasted a maximum of three weeks. The patient was considered "on study" once the patient was dosed and was considered "off study" after the completion of the week 10 visit procedures or when the patient withdrew from the study.

[0399] 3. Week 0, day 0

[0400] Pre-Infusion Procedures:

[0401] The pharmacist would prepare the study medication for infusion while the following visit procedures are being completed:

[0402] a. Update any changes in concomitant medications or intercurrent illnesses

[0403] b. Modified IBMTR Severity Index if the previous score was obtained greater than 8 hours prior to the start of the infusion

[0404] c. Blood draw for the following:

[0405] CBC with diff and platelets (if previous results are >8 hours from the start of the infusion of study medication)

[0406] Serum chemistry (if previous results are >8 hours from the start of the infusion of study medication)

[0407] CD3,4, 8, & 19

[0408] Baseline HAMA (patients who had blood drawn for HAMA as part of their eligibility screen procedure do not need to have this sample obtained)

[0409] Baseline pK sample up to 10 minutes prior to the start of the infusion (for assigned patients only)

[0410] Study Medication Infusion Procedures:

[0411] The pharmacist prepared the study medication such that the maximum total volume to be infused is dependent upon the patient's weight and cohort assignment (total volume of study medication and normal saline). The study medication was generally infused over 2 hours and the patient was closely monitored during the infusion and for the following 4 hours for any untoward reactions to the infusion. As of the start of the infusion of the study medication, the patient was monitored for adverse events on an ongoing basis. The sponsor was notified immediately of any suspected infusion related adverse experiences (cytokine release syndrome: fever, chills, rigors/shakes, hypotension, and rash or hypersensitivity reaction: fever, chills, bradycardia/cardiac arrest, respiratory arrest, acute respiratory distress syndrome, rash/urticaria, pancytopenia, increased liver transaminases, and arthralgias/myalgias). If an infusion reaction is suspected and the patient experiences myalgias or any muscular problems, CPK-III isoenzyme (mm) were obtained.

[0412] Infusion Vital Signs

[0413] During the infusion, vital signs (T, P, R, BP) were obtained just prior to the start of the infusion (a maximum of 10 minutes prior to the start of the infusion), every 15 minutes during the first hour of the infusion (4 sets), followed by 90 minutes after the start of the infusion, and at the completion of the infusion (120 minutes after the start of the infusion). Vital signs were generally obtained hourly for the next 4 hours (4 sets at 180, 240, 300, and 360 minutes after the start of the infusion). After the infusion vital signs have been completed, vital signs were monitored according to the established guidelines used by the clinical center.

[0414] Pharmacokinetic Blood Samples:

[0415] Blood for pK analysis was obtained from at least 5 patients in each cohort. All patients enrolled in study who previously received a murine product had pK assessments completed. Blood samples were generally obtained at the following times after the completion of the first infusion; 15 and 30 minutes, 1, 2, 4, 8, 12, 18, and 24 hours. The 24 hour post infusion sample was obtained prior to the start of the second infusion of ABX-CBL.

[0416] 4. Week 0, days 1-6

[0417] The patient received a daily infusion of the study medication for 7 consecutive days (induction regimen). Each subsequent infusion generally began at the same time as the first infusion (± 60 minutes). The dose was based upon the pre-enrollment weight, therefore, the patient will receive the same dose throughout the treatment period. Data was collected on any patients having an ECG or CXR completed at any time during the treatment period, otherwise routine ECGs and CXRs were not required. The same will hold true for any biopsies completed during this period.

[0418] The following procedures were generally completed within 12 hours prior to the start of each infusion unless otherwise noted:

[0419] a. Abbreviated physical exam (refer to Appendix II)

[0420] b. Weight

[0421] c. KPS or Lansky Scale

[0422] d Modified IBMTR Severity Index

[0423] e. Update any changes in concomitant medications or intercurrent illnesses

[0424] f. Adverse experience assessment

[0425] g. Blood draw for the following (refer to Appendix V for test to be processed by the local labs and those to be processed by the central lab):

[0426] CBC with diff and platelets

[0427] Serum chemistry

[0428] CD3,4, 8, & 19

[0429] Pharmacokinetic sample for assigned patients only and obtain prior to the start of the Day 1 infusion only (this is the 24 hour post infusion 1 sample).

[0430] Study Medication Infusion:

[0431] The study medication was infused over 2 hours following the above procedures. If an infusion reaction was

suspected and the patient experiences myalgias or any muscular problems, CPK-III isoenzyme (mm) were obtained.

[0432] Infusion Vital Signs:

[0433] Vital signs (T, P, R, BP) were obtained according to the schedule described for the first infusion.

[0434] 5. Week 1 (study days 9 and 13)

[0435] At the completion of the induction regimen, the patients were infused with the study medication twice a week for two weeks (maintenance regimen). The start time of each infusion in the maintenance regimen was generally ± 60 minutes from the start time of the first infusion (Day 0). The following procedures were generally completed within 12 hours prior to the start of each infusion unless otherwise noted.

[0436] a. Abbreviated physical exam

[0437] b. Weight

[0438] c. KPS or Lansky Scale

[0439] d. Modified IBMTR Severity Index

[0440] e. Update any changes in concomitant medications or intercurrent illnesses

[0441] f. Adverse experience assessment

[0442] g. Urinalysis (study day 9 only)

[0443] h. Blood draw for the following(refer to Appendix V for test to be processed by the local labs and those to be processed by the central lab):

[0444] CBC with diff and platelets

[0445] Serum chemistry

[0446] CD3, 4, 8, & 19 (day 9 only)

[0447] HAMA (day 9 only)

[0448] Study Medication Infusion:

[0449] The study medication was infused over 2 hours and the procedures described above were again followed. If an infusion reaction was suspected and the patient experiences myalgias or any muscular problems, CPK-III isoenzyme (mm) were obtained.

[0450] Infusion Vital Signs:

[0451] Vital sign regimen described above was utilized.

[0452] Pharmacokinetic Sample:

[0453] A blood sample for pK analysis was obtained about 4 hours after the completion of the Day 9 infusion.

[0454] 6. Week 2 (study days 16 and 20)

[0455] This was the second week of the maintenance regimen (dosing is twice a week for two consecutive weeks). The start time of each infusion was generally ± 60 minutes from the start time of the Day 0 infusion. The following procedures were generally completed within 12 hours prior to the start of each infusion unless otherwise noted:

[0456] a. Abbreviated physical exam

[0457] b. Weight

[0458] c. KPS or Lansky

[0459] d. Modified IBMTR Severity Index

[0460] e. Update any changes in concomitant medications or intercurrent illnesses

[0461] f. Adverse experience assessment

[0462] g. Urinalysis (day 16 only)

[0463] h. Blood draw for the following(refer to Appendix V for test to be processed by the local labs and those to be processed by the central lab):

[0464] CBC with diff and platelets

[0465] Serum chemistry

[0466] CD3, 4, 8, & 19 (day 16 only)

[0467] Study Medication Infusion:

[0468] The study medication was infused over 2 hours and the same procedures described above were followed. If an infusion reaction was suspected and the patient experiences myalgias or any muscular problems, CPK-III isoenzyme (mm) were obtained.

[0469] Infusion Vital Signs:

[0470] Vital sign regimen described above was utilized.

[0471] Pharmacokinetic Sample:

[0472] A blood sample for pK analysis was obtained about 4 hours after the completion of the Day 20 infusion.

[0473] 7. Treatment Follow up Period (Weeks 3-10)

[0474] The treatment follow up period began after the completion of the Day 20 visit and ended at the completion of the week 10 visit. There were five visits during this period. When the patient completed the week 10 visit the patient was considered "off study". If a patient is discharged from the clinical center during this study period, every attempt was made to complete a telephone assessment in place of an office visit. Weeks 3, 4, 5, 6, and 10 were treatment follow up visits. Safety, efficacy or signs of relapse was assessed at these visits. Patients who were partial or complete responders and have a flare of their GVHD were allowed to withdraw from the study and enroll into a separate open label, compassionate treatment protocol. Any biopsies, ECGs, and/or CXRs completed during the treatment follow up period were completed per routine patient care as specified at each clinical center, however, the data from these procedures was collected. Any patients who experienced a suspected infusion related adverse experience with myalgias or any muscular problems and who had elevated mm (isoenzyme which becomes elevated when there is muscular necrosis or inflammation) levels generally had a routine CPK-III (mm) sample obtained throughout the remainder of the study

[0475] 8. Week 3 (Study Day 23)

[0476] The following procedures were completed at this visit:

[0477] a. Complete physical exam, vital signs, and weight

[0478] b. KPS or Lansky

[0479] c. Modified IBMTR Severity Index

- [0480] d. Update any changes in concomitant medications or intercurrent illnesses
- [0481] e. Adverse experience assessment
- [0482] f. Hospitalization status (in patient or discharge)
- [0483] g. Blood draw for the following:
 - [0484] CBC with diff and platelets
 - [0485] Serum chemistry (refer to Appendix V)
 - [0486] CD 3, 4, 8, & 19
- [0487] 9. Week 4 (Study Day 30±1)
- [0488] The following procedures were completed at this visit:
 - [0489] a. Complete physical exam, vital signs, and weight
 - [0490] b. KPS or Lansky
 - [0491] c. Modified IBMTR Severity Index
 - [0492] d. Update any changes in concomitant medications or intercurrent illnesses
 - [0493] e. Adverse experience assessment
 - [0494] f. Hospitalization status (in/outpatient or discharge from clinical center)
 - [0495] g. Urinalysis
 - [0496] h. Blood draw for the following:
 - [0497] CBC with diff and platelets
 - [0498] Serum chemistry (refer to Appendix V)
 - [0499] CD 3, 4, 8, & 19
 - [0500] HAMA
- [0501] 10. Week 5 (Study Day 37±1)
- [0502] The following procedures were completed at this visit:
 - [0503] a. Complete physical exam, vital signs, and weight
 - [0504] b. KPS or Lansky
 - [0505] c. Modified IBMTR Severity Index
 - [0506] d. Update any changes in concomitant medications or intercurrent illnesses
 - [0507] e. Adverse experience assessment
 - [0508] f. Hospitalization status (in/outpatient or discharge from clinical center)
 - [0509] g. Blood draw for the following:
 - [0510] CBC with diff and platelets
 - [0511] Serum chemistry (refer to Appendix V)
 - [0512] CD 3, 4, 8, & 19
- [0513] 11. Week 6 (Study Day 44±1)
- [0514] The following procedures were completed at this visit:
 - [0515] a. Complete physical exam, vital signs, and weight
 - [0516] b. KPS or Lansky
 - [0517] c. Modified IBMTR Severity Index
 - [0518] d. Update any changes in concomitant medications or intercurrent illnesses
 - [0519] e. Adverse experience assessment
 - [0520] f. Hospitalization status (in/outpatient or discharge from clinical center)
 - [0521] g. Urinalysis
 - [0522] h. Blood draw for the following:
 - [0523] CBC with diff and platelets
 - [0524] Serum chemistry (refer to Appendix V)
 - [0525] CD 3, 4, 8, & 19
 - [0526] HAMA
- [0527] 12. Week 10 (Study Day 72±2)
- [0528] At the completion of this visit the patient was considered "off study".
- [0529] The following procedures were completed at this visit:
 - [0530] a. Complete physical exam, vital signs, and weight
 - [0531] b. KPS or Lansky
 - [0532] c. Modified IBMTR Severity Index
 - [0533] d. Update any changes in concomitant medications or intercurrent illnesses
 - [0534] e. Adverse experience assessment
 - [0535] f. Hospitalization status (in/outpatient or discharge from clinical center)
 - [0536] g. Urinalysis
 - [0537] h. Blood draw for the following:
 - [0538] CBC with diff and platelets
 - [0539] Serum chemistry (refer to Appendix V)
 - [0540] CD 3, 4, 8, & 19
 - [0541] HAMA (if any patient has a positive HAMA, blood draws for HAMA will be requested during the Long Term Follow up Period)
- [0542] 13. Additional Visit Timepoint (Day 100 post stem cell transplant)
- [0543] Most patients were assessed 100 days post stem cell transplant. The order in which this visit occurs in relationship to the protocol visits varied on a patient by patient basis depending on when acute GVHD develops post stem cell transplant. Regardless of when day 100 occurs, the following procedures were completed at this visit (if the patient had been discharged from the clinical center every effort was made to obtain this information through a phone call to the patient and the patient's private physician):
 - [0544] a. Abbreviated physical exam, vital signs, and weight
 - [0545] b. KPS or Lansky

- [0546] c. Modified IBMTR Severity Index
- [0547] d. Update any changes in concomitant medications or intercurrent illnesses
- [0548] e. Adverse experience assessment
- [0549] f. Hospitalization status (in/outpatient or discharge from clinical center)
- [0550] g. Blood draw for the following:
- [0551] CBC with diff and platelets
 - [0552] Serum chemistry (refer to Appendix V)
- [0553] 14. Long Term Follow up Period
- [0554] The long term follow up period begins the day after the completion of the week 10 visit and is planned to continue for 10 years or until the patient withdraws consent to be followed. The primary purpose of the long term follow up period is to determine long term safety of ABX-CBL and to determine the long term survival. The patient will be assessed every 6 months from their week 10 visit. These assessments will occur either by telephone interview or by office visit. Long term follow up data may be obtained by the sponsor, Abgenix, Inc., from the primary physician provided that the patient/legal guardian has provided written consent. All data will be entered into the database using the patient's unique study ID. The following information should be obtained during these phone calls or visits:
- [0555] a. Determine the patient's assessment of their health status, this includes the closeout any AE's that were ongoing at the last "on study" visit
 - [0556] b. Determine the onset of any of the following:
 - [0557] Death
 - [0558] Opportunistic Infections
 - [0559] Other immune impairments
 - [0560] Other cancer(s)
 - [0561] Congenital abnormality
 - [0562] If female, if pregnant, status of baby (after pregnancy)
 - [0563] c. Determine if the patient is active in any other research (investigative products and/or devices) since the previous visit/call.
- [0564] If the long term follow up visit data is obtained by the transplant team at the clinical center, a copy of each visit assessment will be faxed to the sponsor within 10 working days of the phone call/visit.
- [0565] C. Determination of HAMA
- [0566] This assay was designed to study the immunogenicity of ABX-CBL in human subjects to detect human antibodies against ABX-CBL (human anti-mCBL antibody) in human serum (human anti-murine antibody, HAMA, response).
- [0567] Materials:
- [0568] Negative Control, pool of HAMA negative sera (from Blood Centers of the Pacific, Irwin Blood Center, SF, Calif.) tested and pooled, stored at -20°C .
 - [0569] Positive Control, pool of HAMA positive sera (from immunizing XenoMouse mice (Abgenix, Inc.) with ABX-CBL and removal and pooling of serum), stored at -20°C .
 - [0570] ABX-CBL, 5 $\mu\text{g}/50\ \mu\text{L}$ (100 $\mu\text{g}/\text{mL}$), Abgenix, Lot No. 097-104-1, stored at -20°C . or equivalent
 - [0571] Biotinylated ABX-CBL (ABX-CBL-biotin), Abgenix, Lot No. J090-112 or equivalent
 - [0572] Streptavidin-HRP, Southern Biotechnology, Cat. No. 7100-05 or equivalent
 - [0573] O-phenylenediamine dihydrochloride (OPD) Substrate Tablets, 20 mg, Sigma, Cat. No. P-7288 or equivalent
 - [0574] O-phenylenediamine dihydrochloride (OPD) Substrate Tablets, 10 mg, Sigma, Cat. No. P-8287 or equivalent
 - [0575] Hydrogen Peroxide, 30%, Sigma, Cat. No. H-1009 or equivalent
 - [0576] Deionized, reverse osmosis purified water (DiH_2O) or equivalent
 - [0577] Coating ELISA Plate: Thaw a vial of ABX-CBL 5 $\mu\text{g}/50\ \mu\text{L}$ (100 $\mu\text{g}/\text{mL}$) at room temperature for 2-5 minutes. Vortex on low speed for 3-5 seconds. Add 48 μL of ABX-CBL 5 $\mu\text{g}/50\ \mu\text{L}$ (100 $\mu\text{g}/\text{mL}$) to 12 mL of Coating Buffer (NaHCO_3 at 16.8 gms/1.8 L DiWater to pH 9.6 w/5N NaOH) in a 15 mL conical tube. Vortex the coating solution on low speed for 3-5 seconds. Pour the coating solution into a reagent reservoir. Using a multi-channel pipettor, add 100 μL of coating solution to each well. Cover plate with plastic plate sealer. Incubate plate at $2-8^{\circ}\text{C}$. for 16-24 hours. Wash the plates with 1 $\times$ Wash Buffer (50 nL Tween 20 in 10 L 10 $\times$ PBS diluted by 10) using a plate washer. Using the multi-channel pipettor, add 100 μL of Blocking Buffer (20 gms BSA in 400 mL 10 $\times$ PBS, 0.4 gms. Thimerosal, 4 mL Tween 20, diluted to 4 L DiWater) to each well. Cover plate with plate sealer and incubate for 1 hour at room temperature.
 - [0578] Preparation of Positive Control: Thaw 1 vial of positive control (HAMA positive serum) at room temperature for 10-20 minutes. Vortex positive control for 3-5 seconds on low speed. Avoid air bubbles. Add 20 μL of positive control to 180 μL of Blocking Buffer in a micro-centrifuge tube. In well A1 and A2 of a low binding 96-well plate, add 20 μL of diluted positive control above to 180 μL of Blocking Buffer. Mix. Mix well by aspirating and dispensing the solution 5 times. Avoid air bubbles. Prepare 2 fold serial dilutions of the positive control. Note: Each plate should include the positive control in duplicate in columns 1 and 2. The following procedure is for one plate. Add 100 μL of Blocking Buffer to wells B3, B4 through H3, H4 on the plate as above. Using a multi-channel pipettor, transfer 100 μL of the solution in wells A1 and A2 to B1 and B2, respectively. Mix well by aspirating and dispensing 100 μL of the solution 5 times. Avoid bubbles. Transfer 100 μL of the solution from wells B1 and B2 to wells C1 and C2, respectively. Mix well by aspirating and dispensing 100 μL of the solution 5 times. Avoid bubbles. Continue dilutions down the plate from row to row with the last dilution in Row G (wells G1 and G2). Leave the Blocking Buffer in Row H as blank controls.

[0579] Preparation of Negative Control: Thaw negative control at room temperature for 20-30 minutes. Vortex the negative control for 3-5 seconds on low speed before transferring to the ELISA plate. Dilute negative control by adding 20 μ L to 980 μ L of blocking Buffer.

[0580] Preparation of Sample: Note 1: Serum samples should be prepared in a designated area. Note 2 Wear gloves when handling serum and follow Universal Precautions. Thaw serum samples at room temperature for 20-30 minutes. Vortex serum samples for 3-5 seconds on low speed. Dilute serum samples 1:50 by adding 20 μ L of a serum sample to 980 μ L Blocking Buffer in a titer tube. Mix the diluted samples by aspirating and dispensing 50 μ L of the solution 5 times. Avoid bubbles. Wash the coated ELISA plate from Step 7.3.2 using a plate washer. Transfer 50 μ L of positive control, negative control, samples and blank to the ELISA plate as above. Cover the ELISA plate with plastic plate sealer and incubate for two hours at room temperature. Shake the plate on low speed.

[0581] Preparation of ABX-CBL-biotin: Note: Minimum of 10 mL of diluted ABX-CBL-biotin is needed for each ELISA plate. Final dilution may be adjusted according to the potency of the reagent. Vortex ABX-CBL-biotin for 3-5 seconds on low speed. Dilute 15 μ L of ABX-CBL-biotin into 1.4-85 mL of Blocking Buffer in a microcentrifuge tube. Total dilution is 1:100. Dilute 1200 μ L of 1:100 diluted ABX-CBL-biotin into 10.80 mL of Blocking Buffer. Total dilution is 1:1000. Vortex for 3-5 seconds on low speed. Wash the coated ELISA plate using a plate washer. Using a multi-channel pipettor, add 100 μ L of 1:1000 diluted ABX-CBL-biotin to each well of the ELISA plate. Cover the plate with plastic plate sealer and incubate for 1 hour at room temperature.

[0582] Preparation of Streptavidin-BPP: Note: Minimum of 10 mL of diluted Streptavidin-HRP is needed for each ELISA plate. Final dilution may be adjusted according to the potency of the reagent. Vortex Streptavidin-BRP for 3-5 seconds on low speed. Dilute 10 μ L of Streptavidin-HRP into 990 μ L of Blocking Buffer in a microcentrifuge tube. Total dilution is 1:100. Dilute 250 μ L of 1:100 diluted Streptavidin-HRP into 12.25 mL of Blocking Buffer. Total dilution is 1:5000. Vortex for 3-5 seconds on low speed. Wash the ELISA plate from above using a plate washer. Using multi-channel pipettor, add 100 μ L of 1:5,000 diluted Streptavidin-HRP to each well of the ELISA plate. Incubate the plate for 15 min at room temperature.

[0583] Preparation of Substrate Solution: Note 1: Minimum of 10 mL of Substrate Solution is needed for each ELISA plate. Note 2: Prepare Substrate Solution fresh prior to use. To make 12 mL of Substrate Solution, add one 10 mg OPD tablet, and 12 μ L of 30% H_2O_2 into 12 mL of Substrate Buffer in a conical tube. Dissolve the tablet by leaving the tube at room temperature for 3-5 minutes. Vortex the solution for 3-5 seconds prior to adding to the plate. Wash the ELISA plate from above using a plate washer. Using a multi-channel pipettor, add 100 μ L of Substrate Solution into each well and incubate for 15 minutes. Using a multi-channel pipettor, add 50 μ L of Stop Solution (2 M H_2SO_4) to each well.

[0584] Reading ELISA plate(s): Set wavelength at, 492 nm and check automix function to premix plate for 5 seconds before reading plate. Use reduction function (Check

L1) to subtract the calculated blank for the assay. Samples and controls are blanked against the buffer blank. Read plate using the SPECTRAmax 250 spectrophotometer within 30 minutes of stopping the assay.

[0585] As discussed above, the present assay was utilized for patient samples in connection with the present clinical trials and no patients tested positive for a HAMA response.

[0586] D. Determination of pK

[0587] The present assay was utilized in connection with pharmacokinetic (pK) studies to measure the presence of ABX-IL8 in human serum.

[0588] Materials:

[0589] ABX-CBL, anti-mouse CBL antibody, 5 μ g/50 μ L (100 μ g/mL), Abgenix, Lot No 69-21-4 or equivalent

[0590] High, Medium and Low Positive Controls, ABX-CBL: 69-21-3, 69-21-2, 69-21-1 or equivalent

[0591] Goat anti-mouse IgM, Caltag, Cat. No. M31500, Lot No. 3501 or equivalent

[0592] Goat anti-mouse IGM-HRP, Caltag, Cat. No. M31507, Lot No. 2301 or equivalent

[0593] Normal human serum

[0594] O-phenylenediamine dihydrochloride (OPD) Substrate Tablets, 20 mg, Sigma, Cat. No. P-7288 or equivalent

[0595] O-phenylenediamine dihydrochloride (OPD) Substrate Tablets, 10 mg, Sigma, Cat. No. P-8287 or equivalent

[0596] Hydrogen Peroxide, 30%, Sigma, Cat. No. H-1009 or equivalent

[0597] Deionized, reverse osmosis purified water (DiH_2O) or equivalent

[0598] Buffers and solutions that are used herein are the same as the buffers and solutions described in connection with the HAMA assay unless described otherwise

[0599] Coating ELISA Plate: Note: Minimum of 10 mL of coating solution is needed for each ELISA plate. Pull vial of goat anti-mouse IgM (1 mg/mL) from the 2-8° C. refrigerator. Let stand for 2-5 minutes at room temperature. Vortex on low speed for 3-5 seconds. Add 3 μ L goat anti-mouse IgM (1 mg/mL) to 15 mL of Coating Buffer in a 15 mL conical tube. Vortex the coating solution on low speed for 3-5 seconds. Pour the coating solution into a reagent reservoir. Using a multi-channel pipettor, add 100 μ L of coating solution to each well. Cover the plate with a plastic plate sealer. Incubate at 2-8° C. for 16-24 hours. Wash the plate with 1 $\times$ Wash Buffer using a plate washer.

[0600] Blocking ELISA Plate: Using the multi-channel pipettor, add 200 μ L of Blocking Buffer to each well. Cover plate with plastic plate sealer and incubate for 1 hour at room temperature.

[0601] Preparation of Standard: Note 1: Blocking Buffer used in Sections 8.4 and 8.6 (except 8.4.4.1 and 8.6.3) contains 1% serum from untreated human subjects. Minimum of 9 mL of Blocking Buffer is needed for each plate. To make 10 mL of Blocking Buffer containing 1% serum,

add 100 μ L serum to 9.9 nL of Blocking Buffer in a conical tube. Vortex on low speed for 3-5 seconds. Thaw 1 vial of ABX-CBL standard (100 μ g/mL) at room temperature for 10-20 minutes. Vortex 100 μ g/mL ABX-CBL on low speed for 3-5 seconds. Avoid bubbles.

[0602] Initial Dilution of Standard: Using a single channel pipette, add 40 μ L of 100 μ g/mL stock to 360 μ L of Blocking Buffer in a 1.7 mL microcentrifuge tube. Mix well. This is a 1:10 dilution equal to 10 μ g/mL. Using a single channel pipette, add 40 μ L of the previous 1:10 dilution (10 μ g/mL) into 460 μ L of Blocking Buffer in a 1.7 mL microcentrifuge tube. Mix well. This dilution is equal to a concentration of 800 ng/mL. Mix the diluted standard by vortexing on low speed for 3-5 seconds. Avoid bubbles. Prepare 2 fold serial dilutions of the standard. Note: Each blank low binding ELISA plate should include the standard in duplicate in columns 1 and 2. The following procedure is for one plate. Add 100 μ L of Blocking Buffer to Wells B1, B2 through H1, H2. Transfer 200 μ L of 800 ng/mL standard to Wells A1 and A2. Using a multi-channel pipette, transfer 100 μ L of the solution in Wells A1 and A2 to Wells B1 and B2, respectively. Mix well by aspirating and dispensing 100 μ L of the solution 5 times. Avoid bubbles. Transfer 100 μ L of the solution from Wells B1 and B2 to Wells C1 and C2, respectively. Mix well by aspirating and dispensing 100 μ L of the solution 5 times. Avoid bubbles. Continue dilutions down the plate from row to row with the last dilution in Wells H1 and H2.

[0603] Preparation of Positive Controls: Note: One vial of high, medium and low control is needed for each assay plate. Thaw 1 vial of high, medium and low controls at room temperature for 10-20 minutes. Vortex the controls for 3-5 seconds on low speed before transferring to the ELISA plate.

[0604] Preparation of Sample: Thaw serum samples at room temperature for 30 minutes. Vortex serum samples on low speed for 3-5 seconds prior to dilutions. Dilute serum samples 1:10 by adding 20 μ L of a serum sample to 180 μ L Blocking Buffer (without 1% serum) in Row A of a blank plate. Mix well by aspirating and dispensing 100 μ L of the solution 5 times. Avoid bubbles. Prepare two fold serial dilutions of the sample. Using a multi-channel pipette, add 100 μ L of Blocking Buffer to Row B through Row H. Transfer 100 μ L of the diluted samples from Step 8.6.3 to Row B. Mix as above. Continue to transfer 100 μ L of the samples from Row B to Row C, from Row C to Row D, and so on to Row H. Mix samples after each transfer by aspirating and dispensing 100 μ L of the solution 5 times. Avoid bubbles. Wash the plate with 1 $\times$ Wash Buffer using a plate washer. Transfer 50 μ L diluted standard, controls and samples from blank plate to the ELISA plate. Start from Row H, then go to Row G and so on up to Row A. Check plate template to add additional wells of buffer blank. Cover the plate with a plastic plate sealer and incubate for two hours at room temperature.

[0605] Prepare RP-conjugated detection antibody: Note: Minimum of 10 mL of diluted HRP-conjugated antibody is needed for each plate. Mix goat anti-mouse IgM-HRP by vortexing on low speed for 3-5 seconds. Dilute goat anti-mouse IgM-HRP to 1:1500 by adding 8 μ L of goat anti-mouse IgM-HRP to 12 mL of Blocking Buffer in a 15 mL conical tube. Vortex. Wash the plate with 1 $\times$ Wash Buffer using a plate washer. Using a multi-channel pipette, add 100

μ L of diluted goat anti-mouse IgM-HRP (from Step 8.10.2) to each well of the plate. Cover the plate with a plastic plate sealer and incubate for 1 hour at room temperature.

[0606] Prepare Substrate Solution: Note 1: Minimum of 10 mL of Substrate Solution is needed for each plate. Prepare Substrate Solution fresh prior to use. To make 12 mL of Substrate Solution, add one 10 mg OPD) tablet and 12 μ L of 30% H₂O₂ to 12 mL of Substrate Buffer in a conical tube. Dissolve the tablet by leaving the tube at room temperature for 3-5 minutes. Vortex the solution for 3-5 seconds prior to adding to the plate. Wash the plate with 1 $\times$ Wash Buffer using a plate washer. Using a multi-channel pipettor, add 100 μ L of Substrate Solution into each well and incubate for 15 minutes.

[0607] Stopping ELISA reaction: Using a multi-channel pipette, add 50 μ L of Stop Solution to each well.

[0608] Reading ELISA plate(s): Set wavelength at 492 nm and check automix function to premix plate for 5 seconds before reading plate. Use reduction function (check LI) to subtract the calculated blank for the assay. Standard, controls and samples are blanked against the buffer blank. Read plate(s) using the SPECTRAmax 250 or equivalent spectrophotometer within 30 minutes of stopping the assay, Operation and Maintenance of the Molecular Devices SPECTRAmax 250 Microplate Spectrophotometer.

[0609] Data Analysis: The OD for the standard is used to calculate the standard curve. Use "4-parameter fit" to curve fit the standard. Sample and control concentrations are calculated automatically by the software from the standard curve. The following criteria must be met in order for the assay to be valid: Only use OD's <4.0 for standard, controls and samples. Compare the results for the assay controls (High, Medium and Low). The values for the controls must fall within 20% of expected concentration and with coefficient of variation (CV) $\leq$ 20%. The CV of the standards between ST03 and ST06 must be $\leq$ 20%. The correlation coefficient of the standard curve of the assay must be $\geq$ 0.990.

[0610] The present assay was utilized for determining the pharmacokinetics of the ABX-CBL antibody in the present clinical trials. The results from our preliminary determinations of pKs in patients utilizing the above-assay are shown in **FIG. 1**.

[0611] E. Results

[0612] Herein, we describe the results that were observed in the treatment of patients with acute GVHD with ABX-CBL.

[0613] In the trial, twenty-seven patients were enrolled across the four dose levels. The lower doses were completed prior to enrolling in the higher dose cohorts. Patients who were treated at the higher dose in the original third cohort (0.3 mg/kg) experienced myalgia or myalgia-like symptoms. Abgenix determined this dose to be the Maximum Tolerated Dose (MTD) and revised the last dose from 1.0 mg/kg to 0.2 mg/kg (mid dose between the MTD and the dose prior to the MTD).

[0614] Once the 4 dose cohorts were filled, additional patients were enrolled at a dose level of 0.15 mg/kg to 0.2 mg/kg. As of Jan. 13, 1999, a total of 44 patients (17 additional patients) have been enrolled. Data continues to be

collected on these additional 17 patients. This data will be presented as it becomes available.

[0615] All data presented herein are based upon the initial 27 patients except for the Serious Adverse Event (SAE) Summaries. The SAE Summaries relate to all patients as of Jan. 13, 1999.

[0616] Patients had to receive a minimum of 4 infusions of ABX-CBL to be evaluated for efficacy. Of the twenty-seven patients enrolled, 23 met this criteria. Excluding the patients in cohort 1 (the no-effect dose). There was an overall response rate of 73% with a mean duration of 32 days.

[0617] Other than the incidence of myalgia, ABX-CBL was well tolerated. All patients were, and remain, negative for HAMA, and no reports of hypersensitivity to ABX-CBL have been received.

[0618] 1. Demographics:

[0619] Of the twenty-seven patients enrolled, 21 were adults (age 16 or older) and 6 were pediatric (Table 4). Twenty-four patients were recipients of an allogeneic bone marrow transplant, and the other three received peripheral stem cells. The mean duration from the date of transplant to enrollment into this study was 48 days. Seven patients were entered into the study with an IBMTR grade of B, 10 with a grade of C and 10 with D. (Table 5). Table 6 lists the baseline score for the 23 patients evaluated for efficacy.

TABLE 6

	GENDER/AGE CATEGORY		
	MALE	FEMALE	TOTAL
ADULT	13	8	21
PEDIATRIC (<16 YRS)	4	2	6
TOTAL	17	10	27

[0620]

TABLE 7

BASELINE IBMTR SEVERITY SCORE-ALL PATIENTS				
COHORT	B n(%)	C n(%)	D n(%)	TOTAL N
1 (0.01 mg/kg)	2 (22%)	3 (33%)	4 (44%)	9
2 (0.1 mg/kg)	2 (29%)	3 (42%)	2 (29%)	7
3 (0.3 mg/kg)	1 (50%)	1 (50%)	0	2
4 (0.2 mg/kg)	2 (22%)	3 (33%)	4 (44%)	9
TOTAL	7	10	10	27

[0621]

TABLE 8

BASELINE IBMTR SEVERITY SCORE FOR EVALUABLE PATIENTS				
COHORT	B n (%)	C n (%)	D n (%)	TOTAL N
1 (0.01 mg/kg)	2 (25%)	3 (38%)	3 (38%)	8
2 (0.1 mg/kg)	1 (17%)	3 (50%)	2 (33%)	6

TABLE 8-continued

BASELINE IBMTR SEVERITY SCORE FOR EVALUABLE PATIENTS				
COHORT	B n (%)	C n (%)	D n (%)	TOTAL N
3 (0.3 mg/kg)	1 (50%)	1 (50%)	0	2
4 (0.2 mg/kg)	2 (29%)	2 (29%)	3 (43%)	7
TOTAL	6	9	8	23

[0622] 2. Efficacy:

[0623] Patients eligible for enrollment into this study required a minimum IBMTR score of B. Patients who demonstrated at least a 2 index decrease in overall IBMTR score were considered responders. Those who decreased to no score, meaning there was no acute GvHD present, were considered to be complete responders. Only patients who received 4 or more infusions of ABX-CBL are included in the efficacy analyses. (Table 7)

TABLE 9

COHORT	EVALUATED FOR EFFICACY		MEAN DURATION OF RESPONSE (n)
	(n)	n (%)	
1 (0.01 mg/kg)	8	3 (38%)	24 days (3)
2 (0.1 mg/kg)	6	4 (67%)	11 days (3)
3 (0.3 mg/kg)	2	2 (100%)	69 days (1)
4 (0.2 mg/kg)	7	4 (57%)*	41 days (3)
TOTAL	23	13 (57%)	36 days

*One patient responded to additional therapy with ABX-CBL in the ABX-CB-9702 protocol and is not included in the above table.

[0624] Overall, thirteen (57%) of the twenty-three patients demonstrated a response to ABX-CBL in ABX-CB-9701. The mean duration was 36 days. One additional patient who rolled over into protocol described below responded to additional therapy. This brings the overall response rate to 61%. The assumption going into the study was that the dose of 0.01 mg/kg would be the no effect dose. Assuming this dose to have no effect, the response rate was 73% (11 of 15 patients). With this assumption, the mean duration of response was 32 days.

[0625] The duration of response seems to increase as the dose is increased. One patient, [0108], was an outlier for duration in the first cohort. This patient's duration lasted at least 59 days. The duration may be longer, but the study ended at Day 72.

[0626] [Patient 0816] experienced severe myalgia at the 0.3 mg/kg dose level during the first infusion. This patient was continued at a decreased dose of 0.2 mg/kg for all subsequent infusions. Because of the change in dose, this patient is evaluated in the 0.2 mg/kg cohort for efficacy and in the 0.3 mg/kg for safety.

[0627] Only one patient in the lowest dose cohort and both patients in the highest dose level completed the study through Day 72. Four of the six patients in the 0.1 mg/kg dose group completed the study, and 4 of the 7 in the 0.2

mg/kg dose group completed. All patients who demonstrated a complete response also completed this study through Day 72.

[0628] 3. Safety:

[0629] All patients who received any amount of ABX-CBL were evaluated for safety. ABX-CBL was well tolerated with the exception of myalgia, which became the Dose Limiting Toxicity (DLT). The incidence of myalgia increased in relationship to an increase in the dose administered. This led to the Maximum Tolerated Dose (MTD) at 0.3 mg/kg. The onset of the myalgia ranged from 20-60 minutes into the infusion and usually resolved within 1-2 hours after the completion of the infusion. Of the 14 patients who experienced any grade of myalgia, two required being withdrawn from this study due to the myalgia. All myalgias resolved without sequelae except for one patient in whom myalgia persisted. This last incidence is under further evaluation and clarification. Table 6 summarizes the incidence of myalgia by severity and dose. Patients with adverse events listed as myalgia graded as “not related” or “unlikely” and with a baseline disease of myalgia are not included in the this table.

TABLE 10

INCIDENCE OF MYALGIA AND OUTCOME									
		0.01 mg/kg (n = 9)		0.1 mg/kg (n = 7)		0.3 mg/kg (n = 3)		0.2 mg/kg (n = 8)	
SEVERITY	n (%)	Study status	n (%)	Study status	n (%)	Study status	n (%)	Study status	n (%)
SEVERE			1	W/D	1	con't dec. dose	3	con't W/D	
MODERATE	2	Con't	1	Con't			1	Con't	
MILD	1	Con't	1	Con't			1	Con't	

W/D = withdrew from the study related to the myalgia

[0630] Abgenix continues to investigate the causality of myalgia and any possible inter-relationships. The following causes have been ruled out as a predisposing factor to those who do develop myalgia:

[0631] alteration in electrolytes

[0632] responders vs non responders

[0633] type of transplant

[0634] type of donor

[0635] steroid dose

[0636] Eleven Serious Adverse Experiences in eleven patients have been reported with ABX-CBL. Five “severe” events, all myalgia related, are listed as “probable” for the relationship to ABX-CBL. One event, “hepatic failure of unknown etiology” is listed as “suspected”. The remaining SAEs are listed as “unlikely” or “not related”.

[0637] Twenty-three of these events were evaluated as probably related to ABX-CBL and 7 as suspected. All other events were reported as “unlikely” or “not related”.

[0638] Of the 23 “probable” adverse events, all except 2 were myalgia related. One patient experienced moderate “fatigue” which resolved without sequelae. The other expe-

rienced moderate “hemolysis” which resolved with a sequelae of increased Liver Function Tests (LFT).

[0639] Of the seven events evaluated as “suspected” to be related to ABX-CBL, 1 event was severe, 4 were moderate, and 2 were mild in severity. All of these events resolved without sequelae. The severe event was “edema”. The four moderate events occurred in 4 patients and consisted of “moderate decrease in uric acid”, “fever/chills”, “hypotension”, and “fever”. The two mild events occurred in two patients and consisted of “low grade fever following study drug” and “chills”.

[0640] HAMA testing on all 27 patients has been negative through the patients’ last study visit.

[0641] Lymphocyte counts were drawn from all patients just prior to the first infusion and at regular intervals throughout the study. Of the patients who enrolled into ABX-CB-9701, approximately 50% could not be evaluated on the basis of the immunocompromised state secondary to both BMT and their ongoing GvHD. Patients who are post stem cell transplant are immunodeficient secondary to their conditioning regimen as well as an exacerbation of their immunodeficient state from acute GvHD. To date, ABX-CBL does not appear to have an untoward effect on the T-cell counts.

Phase II Clinical Trial of ABX-CBL—Rescue Protocol

[0642] As patients completed the above-described Phase II trial, we also initiated a second Phase II continuation trial for such patients to continue to receive ABX-CBL for any flares of GVHD experienced. The continuation trial was designed as an open label clinical trial for patients with acute GVHD who have previous exposure to ABX-CBL. Those patients who had acute GVHD of grades II/III/IV severity, as discussed above, were eligible.

[0643] In the trial, all patients are receiving, or will receive, up to 7 intravenous doses (1st treatment course) of ABX-CBL. The medication will be infused over 2 hours via a syringe pump for 7 consecutive days. The dose will be 0.2 mg/kg (approximate dose used effectively in clinical trial described above). If the first treatment course produced a therapeutic effect (complete or partial response), patients may receive a second treatment course prior to the onset of chronic GVHD, or day 200 post primary transplant whichever is reached first. The second treatment course with ABX-CBL will be handled on a case by case basis through a discussion with the medical monitor and the investigator.

[0644] The objectives of this trial were as follows:

[0645] To assess the safety of continued dosing with ABX-CBL in patients with acute GVHD.

[0646] To determine the clinical effect of repeat treatments of ABX-CBL in patients with flare of acute GVHD or patients who were previous treatment failures with ABX-CBL.

[0647] To allow treatment for patients who failed to demonstrate a clinical effect at a lower dose of ABX-CBL and/or to provide treatment for previous responders to ABX-CBL who are experiencing a flare of their acute GVHD.

[0648] To assess flare rates after initial treatment with AdBX-CBL.

[0649] All of the procedures described above in connection with the initial clinical trial were utilized in connection with this study, with only minor modifications.

Dosing, Dose Regimen, and Treatment with ABX-CBL

[0650] In view of the foregoing discussion and results, ABX-CBL provides a profound treatment for GVHD and likely other disease etiologies wherein lymphatic cells are deleteriously or undesirably activated. The results presented herein demonstrate that through administration of a dose of ABX-CBL greater than about 0.1 mg/kg and less than about 0.4 mg/kg of the antibody is efficacious in connection with the treatment of such disease etiologies. Preferably, the dose is from about 0.1 mg/kg to about 0.3 mg/kg and more preferably from about 0.15 mg/kg to about 0.2 mg/kg. Further, the dosing regimen disclosed herein of an induction regimen (plural daily infusions, herein daily for 7 days) followed by a maintenance regimen (periodic infusions, herein twice weekly for two weeks) appears to assist in remission of GVBD and certainly lessens the severity of patients' GVHD between flares of the disease.

[0651] As will be appreciated, both the purified ABX-C, BL, discussed in detail in the present invention and other anti-CD147 antibodies, such as those discussed herein, will be similarly efficacious.

[0652] In addition to GVHD, therapeutics in accordance with the present invention will likely be efficacious with respect to diseases having an etiology characterized by a harmful presence of activated T cells, B cells, or monocytes. As an example, GVHD is one such disease. However, many inflammatory diseases and autoimmune diseases can be characterized as sharing such an etiology. Further the therapies of the invention will likely be efficacious in the following disease etiologies, including, without limitation: graft versus host disease (GVHD), organ transplant rejection diseases (including, without limitation, renal transplant, ocular transplant, and others), cancers (including, without limitation, cancers of the blood (i.e., leukemias and lymphomas), pancreatic, and others), autoimmune diseases, inflammatory diseases (including without limitations arthritis, rheumatoid arthritis), and others.

Experiment 22 Surrogate Antibodies that Bind to Murine gp42 for Animal Models

[0653] As discussed above, certain animal models are contemplated in connection with the present invention. One of the simplest animal models is the mouse. The 2.6.1 antibody did not bind to mouse gp42 (basigin or mouse CD147). Accordingly, we undertook the generation of anti-mouse gp42 antibodies from rats that could be utilized as a surrogate antibody to ABX-CBL and/or the 2.6.1 antibodies for use in such models. Described below is cloning strategy utilized to prepare fusion proteins for immunization of rats and the preliminary characterization of antibodies generated therefrom. The cloning strategy described below is further detailed in **FIGS. 51 and 52.**

Cloning of Hu-CD147IgG2 Fusion Protein

[0654] The following PCR primers were utilized, based on the CD147 sequence reported by Miyauchi et al. *J. Biochem.* 110:770-774 (1991) (Gene Bank Accession # D45131)

5 prime:

(SEQ ID NO:58)

5'-GACTACGAATTCGGACCGCGGAGGAATAGGAATCATG-3'
and

3 prime:

(SEQ ID NO:59)

5'-GGATGGTGTGGTAGCTAGCACGCGGAGCGTGATGATGGCCTG-3'

[0655] A 626bp PCR product was amplified from CD147/pBCKMV plasmid DNA template that encoded the amino terminal 202 amino acid residues of the extracellular domain of CD147. The PCR product was digested with EcoR1 and Nhe1 and ligated into pIK1.1Hu-CD4IgG2 expression vector digested with EcoR1 and Nhe1. The resulting construct, pIKHu-CD147IgG2 encodes a fusion protein consisting of the N-terminal 202 amino acids of CD147 the last four C-terminal residues of the extracellular domain of CD4 in frame with the hinge CH2 and CH3 domains of Hu IgG2.

Cloning of Mu-GP42IgG2 Fusion Protein

[0656] The following PCR primers were utilized, based on the GP42 sequence reported by Kanekura et al. *Cell Struct. Funct.* 16:23-30 (1991) (Gene Bank Accession # Y16256):

5 prime:

(SEQ ID NO:60)

5'-GACTACGAATTCACGAGGCGACATGGCGGCGGC-3'
and

3 prime:

(SEQ ID NO:61)

5'-GGATGGTGTGGTAGCTAGCACGCGAGTGAGATGGTTTCCCG-3'

[0657] A 659bp PCR product was amplified from mouse lymph node cDNA and encodes the amino terminal 206 amino acid residues of the extracellular domain of GP42. The PCR product was digested with EcoR1 and Nhe1 and ligated into pIK1.1Hu-CD4IgG2 expression vector digested with EcoR1 and Nhe1 to create pIKMu-GP42 IgG2.

Stable CHO Cell Line Engineering

[0658] The EcoR1/Bgl2 fragments from pIK Hu-CD147IgG2 and pIKMu-GP42IgG2 were cloned into the expression vector pWBFNP DHBR digested with EcoR1/Bgl2. PWBFPN DBFR is a derivative of pWBFNP into which a DHFR cDNA under the transcriptional control of SV40 promoter/enhancer and SV40 poly A is cloned at the NotI site. The resulting constructs, Hu-CD147IgG2 DHFR and Mu-GP42IgG2 DHFR were introduced into DHFR deficient CHO cell lines by CaPO₄ mediated transfection. Stable lines were selected for their ability to grow in the absence of exogenous thymidine, glycine and purines. Clones secreting elevated levels of fusion proteins as judged by SDS-PAGE were suspension adapted to spinner flasks in serum-free media. Mu-GP42IgG2 and Hu-CD147IgG2 fusion proteins were purified from culture media by protein A chromatography.

[0659] Following generation of the fusion proteins, rats were immunized using conventional techniques and hybridomas generated also using conventional techniques. Anti-

bodies secreted by such hybridonias could then be utilized as surrogate antibodies in certain animal models, particularly, murine models.

Incorporation by Reference

[0660] All references cited herein, including patents, patent applications, papers, text books, and the like, and the references cited therein, to the extent that they are not already, are hereby incorporated herein by reference in their entirety.

Equivalents

[0661] The foregoing description, Figures, and Examples detail certain preferred embodiments of the invention and describes the best mode contemplated by the inventors. It will be appreciated, however, that no matter how detailed the foregoing may appear in text, the invention may be practiced in many ways and the invention should be construed in accordance with the appended claims and any equivalents thereof.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 105

<210> SEQ ID NO 1

<211> LENGTH: 8

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 1

Ile Thr Leu Arg Val Arg Ser His
1 5

<210> SEQ ID NO 2

<211> LENGTH: 7

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 2

Glu Glu Arg Leu Arg Ser Tyr
1 5

<210> SEQ ID NO 3

<211> LENGTH: 7

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 3

Tyr Glu Arg Val Arg Trp Tyr
1 5

<210> SEQ ID NO 4

<211> LENGTH: 7

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 4

Glu Glu Arg Leu Arg Ser Tyr
1 5

<210> SEQ ID NO 5

<211> LENGTH: 7

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 5

Ala Glu Arg Ile Arg Ser Ile
1 5

<210> SEQ ID NO 6

<211> LENGTH: 7

-continued

<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 6

Glu Glu Arg Leu Arg Ser Tyr
1 5

<210> SEQ ID NO 7
<211> LENGTH: 12
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 7

Thr Val His Gly Asp Leu Arg Leu Arg Ser Leu Pro
1 5 10

<210> SEQ ID NO 8
<211> LENGTH: 12
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 8

Thr Asn Asp Ile Gly Leu Arg Gln Arg Ser His Ser
1 5 10

<210> SEQ ID NO 9
<211> LENGTH: 12
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 9

Ser Pro Leu Leu Asp Gly Gln Arg Glu Arg Ser Tyr
1 5 10

<210> SEQ ID NO 10
<211> LENGTH: 12
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 10

Tyr Asp Leu Pro Met Arg Ser Arg Ser Tyr Pro Gly
1 5 10

<210> SEQ ID NO 11
<211> LENGTH: 4
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<223> OTHER INFORMATION: Xaa may be any amino acid

<400> SEQUENCE: 11

Arg Xaa Arg Ser
1

<210> SEQ ID NO 12
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 12

Lys Gly Ser Asp Gln Ala Ile Ile Thr Leu Arg Val Arg Ser His
1 5 10 15

-continued

<210> SEQ ID NO 13
 <211> LENGTH: 5
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens
 <220> FEATURE:
 <223> OTHER INFORMATION: Xaa may be any amino acid

<400> SEQUENCE: 13

Arg Xaa Arg Ser His
 1 5

<210> SEQ ID NO 14
 <211> LENGTH: 269
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 14

Met Ala Ala Ala Leu Phe Val Leu Leu Gly Phe Ala Leu Leu Gly Thr
 1 5 10 15
 His Gly Ala Ser Gly Ala Ala Gly Thr Val Phe Thr Thr Val Glu Asp
 20 25 30
 Leu Gly Ser Lys Ile Leu Leu Thr Cys Ser Leu Asn Asp Ser Ala Thr
 35 40 45
 Glu Val Thr Gly His Arg Trp Leu Lys Gly Gly Val Val Leu Lys Glu
 50 55 60
 Asp Ala Leu Pro Gly Gln Lys Thr Glu Phe Lys Val Asp Ser Asp Asp
 65 70 75 80
 Gln Trp Gly Glu Tyr Ser Cys Val Phe Leu Pro Glu Pro Met Gly Thr
 85 90 95
 Ala Asn Ile Gln Leu His Gly Pro Pro Arg Val Lys Ala Val Lys Ser
 100 105 110
 Ser Glu His Ile Asn Glu Gly Glu Thr Ala Met Leu Val Cys Lys Ser
 115 120 125
 Glu Ser Val Pro Pro Val Thr Asp Trp Ala Trp Tyr Lys Ile Thr Asp
 130 135 140
 Ser Glu Asp Lys Ala Leu Met Asn Gly Ser Glu Ser Arg Phe Phe Val
 145 150 155 160
 Ser Ser Ser Gln Gly Arg Ser Glu Leu His Ile Glu Asn Leu Asn Met
 165 170 175
 Glu Ala Asp Pro Gly Gln Tyr Arg Cys Asn Gly Thr Ser Ser Lys Gly
 180 185 190
 Ser Asp Gln Ala Ile Ile Thr Leu Arg Val Arg Ser His Leu Ala Ala
 195 200 205
 Leu Trp Pro Phe Leu Gly Ile Val Ala Glu Val Leu Val Leu Val Thr
 210 215 220
 Ile Ile Phe Ile Tyr Glu Lys Arg Arg Lys Pro Glu Asp Val Leu Asp
 225 230 235 240
 Asp Asp Asp Ala Gly Ser Ala Pro Leu Lys Ser Ser Gly Gln His Gln
 245 250 255
 Asn Asp Lys Gly Lys Asn Val Arg Gln Arg Asn Ser Ser
 260 265

<210> SEQ ID NO 15
 <211> LENGTH: 7
 <212> TYPE: PRT

-continued

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 15

Pro Glu Arg Ile Leu Ser Ile
1 5

<210> SEQ ID NO 16

<211> LENGTH: 9

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 16

Gly Gly Ser Arg Ala Arg Asn Leu Pro
1 5

<210> SEQ ID NO 17

<211> LENGTH: 463

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 17

Met Glu Thr Glu Gln Pro Glu Glu Thr Phe Pro Asn Thr Glu Thr Asn
1 5 10 15Gly Glu Phe Gly Lys Arg Pro Ala Glu Asp Met Glu Glu Glu Gln Ala
20 25 30Phe Lys Arg Ser Arg Asn Thr Asp Glu Met Val Glu Leu Arg Ile Leu
35 40 45Leu Gln Ser Lys Asn Ala Gly Ala Val Ile Gly Lys Gly Gly Lys Asn
50 55 60Ile Lys Ala Leu Arg Thr Asp Tyr Asn Ala Ser Val Ser Val Pro Asp
65 70 75 80Ser Ser Gly Pro Glu Arg Ile Leu Ser Ile Ser Ala Asp Ile Glu Thr
85 90 95Ile Gly Glu Ile Leu Lys Lys Ile Ile Pro Thr Leu Glu Glu Gly Leu
100 105 110Gln Leu Pro Ser Pro Thr Ala Thr Ser Gln Leu Pro Leu Glu Ser Asp
115 120 125Ala Val Glu Cys Leu Asn Tyr Gln His Tyr Lys Gly Ser Asp Phe Asp
130 135 140Cys Glu Leu Arg Leu Leu Ile His Gln Ser Leu Ala Gly Gly Ile Ile
145 150 155 160Gly Val Lys Gly Ala Lys Ile Lys Glu Leu Arg Glu Asn Thr Gln Thr
165 170 175Thr Ile Lys Leu Phe Gln Glu Cys Cys Pro His Ser Thr Asp Arg Val
180 185 190Val Leu Ile Gly Gly Lys Pro Asp Arg Val Val Glu Cys Ile Lys Ile
195 200 205Ile Leu Asp Leu Ile Ser Glu Ser Pro Ile Lys Gly Arg Ala Gln Pro
210 215 220Tyr Asp Pro Asn Phe Tyr Asp Glu Thr Tyr Asp Tyr Gly Gly Phe Thr
225 230 235 240Met Met Phe Asp Asp Arg Arg Gly Arg Pro Val Gly Phe Pro Met Arg
245 250 255Gly Arg Gly Gly Phe Asp Arg Met Pro Pro Gly Arg Gly Gly Arg Pro
260 265 270

-continued

Met Pro Pro Ser Arg Arg Asp Tyr Asp Asp Met Ser Pro Arg Arg Gly
 275 280 285

Pro Pro Pro Pro Pro Gly Arg Gly Gly Arg Gly Gly Ser Arg Ala
 290 295 300

Arg Asn Leu Pro Leu Pro Pro Pro Pro Pro Arg Gly Gly Asp Leu
 305 310 315 320

Met Ala Tyr Asp Arg Arg Gly Arg Pro Gly Asp Arg Tyr Asp Gly Met
 325 330 335

Val Gly Phe Ser Ala Asp Glu Thr Trp Asp Ser Ala Ile Asp Thr Trp
 340 345 350

Ser Pro Ser Glu Trp Gln Met Ala Tyr Glu Pro Gln Gly Gly Ser Gly
 355 360 365

Tyr Asp Tyr Ser Tyr Ala Gly Gly Arg Gly Ser Tyr Gly Asp Leu Gly
 370 375 380

Gly Pro Ile Ile Thr Thr Gln Val Thr Ile Pro Lys Asp Leu Ala Gly
 385 390 395 400

Ser Ile Ile Gly Lys Gly Gly Gln Arg Ile Lys Gln Ile Arg His Glu
 405 410 415

Ser Gly Ala Ser Ile Lys Ile Asp Glu Pro Leu Glu Gly Ser Glu Asp
 420 425 430

Arg Ile Ile Thr Ile Thr Gly Thr Gln Asp Gln Ile Gln Asn Ala Gln
 435 440 445

Tyr Leu Leu Gln Asn Ser Val Lys Gln Tyr Ser Gly Lys Phe Phe
 450 455 460

<210> SEQ ID NO 18
 <211> LENGTH: 570
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 18

Glu Val Lys Leu Glu Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
 1 5 10 15

Ser Met Lys Leu Ser Cys Val Ala Ser Gly Phe Thr Phe Ser Asn Tyr
 20 25 30

Trp Met Asn Trp Val Arg Gln Ser Pro Glu Lys Gly Leu Glu Trp Val
 35 40 45

Ala Glu Ile Arg Leu Lys Ser Asn Asn Tyr Ala Thr His Tyr Ala Glu
 50 55 60

Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asp Ser Lys Ser Ser
 65 70 75 80

Val Tyr Leu Gln Met Asn Asn Leu Arg Ala Glu Asp Thr Gly Ile Tyr
 85 90 95

Tyr Cys Thr Asp Tyr Asp Ala Tyr Trp Gly Gln Gly Thr Leu Val Thr
 100 105 110

Val Ser Ala Glu Ser Gln Ser Phe Pro Asn Val Phe Pro Leu Val Ser
 115 120 125

Cys Glu Ser Pro Leu Ser Asp Lys Asn Leu Val Ala Met Gly Cys Leu
 130 135 140

Ala Arg Asp Phe Leu Pro Ser Thr Ile Ser Phe Thr Trp Asn Tyr Gln
 145 150 155 160

Asn Asn Thr Glu Val Ile Gln Gly Ile Arg Thr Phe Pro Thr Leu Arg

-continued

165					170					175						
Thr	Gly	Gly	Lys	Tyr	Leu	Ala	Thr	Ser	Gln	Val	Leu	Leu	Ser	Pro	Lys	
			180				185						190			
Ser	Ile	Leu	Glu	Gly	Ser	Asp	Glu	Tyr	Leu	Val	Cys	Lys	Ile	His	Tyr	
			195				200						205			
Gly	Gly	Lys	Asn	Arg	Asp	Leu	His	Val	Pro	Ile	Pro	Ala	Val	Ala	Glu	
			210				215						220			
Met	Asn	Pro	Asn	Val	Asn	Val	Phe	Val	Pro	Pro	Arg	Asp	Gly	Phe	Ser	
225						230						235			240	
Gly	Pro	Ala	Pro	Arg	Lys	Ser	Lys	Leu	Ile	Cys	Glu	Ala	Thr	Asn	Phe	
						245			250						255	
Thr	Pro	Lys	Pro	Ile	Thr	Val	Ser	Trp	Leu	Lys	Asp	Gly	Lys	Leu	Val	
			260						265			270				
Glu	Ser	Gly	Phe	Thr	Thr	Asp	Pro	Val	Thr	Ile	Glu	Asn	Lys	Gly	Ser	
			275			280						285				
Thr	Pro	Gln	Thr	Tyr	Lys	Val	Ile	Ser	Thr	Leu	Thr	Ile	Ser	Glu	Ile	
			290			295						300				
Asp	Trp	Leu	Asn	Leu	Asn	Val	Tyr	Thr	Cys	Arg	Val	Asp	His	Arg	Gly	
305						310						315			320	
Leu	Thr	Phe	Leu	Lys	Asn	Val	Ser	Ser	Thr	Cys	Ala	Ala	Ser	Pro	Ser	
			325						330						335	
Thr	Asp	Ile	Leu	Thr	Phe	Thr	Ile	Pro	Pro	Ser	Phe	Ala	Asp	Ile	Phe	
			340						345			350				
Leu	Ser	Lys	Ser	Ala	Asn	Leu	Thr	Cys	Leu	Val	Ser	Asn	Leu	Ala	Thr	
			355			360						365				
Tyr	Glu	Thr	Leu	Asn	Ile	Ser	Trp	Ala	Ser	Gln	Ser	Gly	Glu	Pro	Leu	
			370			375						380				
Glu	Thr	Lys	Ile	Lys	Ile	Met	Glu	Ser	His	Pro	Asn	Gly	Thr	Phe	Ser	
385						390						395			400	
Ala	Lys	Gly	Val	Ala	Ser	Val	Cys	Val	Glu	Asp	Trp	Asn	Asn	Arg	Lys	
			405						410						415	
Glu	Phe	Val	Cys	Thr	Val	Thr	His	Arg	Asp	Leu	Pro	Ser	Pro	Gln	Lys	
			420						425			430				
Lys	Phe	Ile	Ser	Lys	Pro	Asn	Glu	Val	His	Lys	His	Pro	Pro	Ala	Val	
			435			440						445				
Tyr	Leu	Leu	Pro	Pro	Ala	Arg	Glu	Gln	Leu	Asn	Leu	Arg	Glu	Ser	Ala	
450						455						460				
Thr	Val	Thr	Cys	Leu	Val	Lys	Gly	Phe	Ser	Pro	Ala	Asp	Ile	Ser	Val	
465						470						475			480	
Gln	Trp	Leu	Gln	Arg	Gly	Gln	Leu	Leu	Pro	Gln	Glu	Lys	Tyr	Val	Thr	
			485						490						495	
Ser	Ala	Pro	Met	Pro	Glu	Pro	Gly	Ala	Pro	Gly	Phe	Tyr	Phe	Thr	His	
			500						505			510				
Ser	Ile	Leu	Thr	Val	Thr	Glu	Glu	Glu	Trp	Asn	Ser	Gly	Glu	Thr	Tyr	
			515			520						525				
Thr	Cys	Val	Val	Gly	His	Glu	Ala	Leu	Pro	His	Leu	Val	Thr	Glu	Arg	
530						535						540				
Thr	Val	Asp	Lys	Ser	Thr	Gly	Lys	Pro	Thr	Leu	Tyr	Asn	Val	Ser	Leu	
545						550						555			560	
Ile	Met	Ser	Asp	Thr	Gly	Gly	Thr	Cys	Tyr							
			565			570										

-continued

<210> SEQ ID NO 19

<211> LENGTH: 206

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 19

Lys Phe Leu Leu Val Ser Ala Gly Asp Arg Val Thr Ile Thr Cys Lys
1 5 10 15
Ala Ser Gln Ser Val Ser Asn Asp Val Ala Trp Tyr Gln Gln Lys Pro
20 25 30
Gly Gln Ser Pro Lys Leu Leu Ile Tyr Tyr Ala Ser Asn Arg Tyr Thr
35 40 45
Gly Val Pro Asp Arg Phe Thr Gly Ser Gly Tyr Gly Thr Asp Phe Thr
50 55 60
Phe Thr Ile Ser Thr Val Gln Ala Glu Asp Leu Ala Val Tyr Phe Cys
65 70 75 80
Gln Gln Asp Tyr Ser Ser Pro Tyr Thr Phe Gly Gly Gly Thr Lys Leu
85 90 95
Glu Ile Lys Arg Ala Asp Ala Ala Pro Thr Val Ser Ile Phe Pro Pro
100 105 110
Ser Ser Glu Gln Leu Thr Ser Gly Gly Ala Ser Val Val Cys Phe Leu
115 120 125
Asn Asn Phe Tyr Pro Lys Asp Ile Asn Val Lys Trp Lys Ile Asp Gly
130 135 140
Ser Glu Arg Gln Asn Gly Val Leu Asn Ser Trp Thr Asp Gln Asp Ser
145 150 155 160
Lys Asp Ser Thr Tyr Ser Met Ser Ser Thr Leu Thr Leu Thr Lys Asp
165 170 175
Glu Tyr Glu Arg His Asn Ser Tyr Thr Cys Glu Ala Thr His Lys Thr
180 185 190
Ser Thr Ser Pro Ile Val Lys Ser Phe Asn Arg Asn Glu Cys
195 200 205

<210> SEQ ID NO 20

<211> LENGTH: 12

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 20

Ser Leu Ala Pro Leu Trp Tyr Tyr Ser Arg His Gly
1 5 10

<210> SEQ ID NO 21

<211> LENGTH: 12

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 21

His Thr Pro Glu Thr Ala Pro Leu Pro Ala Thr Val
1 5 10

<210> SEQ ID NO 22

<211> LENGTH: 159

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

-continued

<400> SEQUENCE: 22

```

Met Lys Asn His Leu Leu Phe Trp Gly Val Leu Ala Val Phe Ile Lys
 1             5             10             15

Ala Val His Val Lys Ala Gln Glu Asp Glu Arg Ile Val Leu Val Asp
      20             25             30

Asn Lys Cys Lys Cys Ala Arg Ile Thr Ser Arg Ile Ile Arg Ser Ser
      35             40             45

Glu Asp Pro Asn Glu Asp Ile Val Glu Arg Asn Ile Arg Ile Ile Val
      50             55             60

Pro Leu Asn Asn Arg Glu Asn Ile Ser Asp Pro Thr Ser Pro Leu Arg
      65             70             75             80

Thr Arg Phe Val Tyr His Leu Ser Asp Leu Cys Lys Lys Cys Asp Pro
      85             90             95

Thr Glu Val Glu Leu Asp Asn Gln Ile Val Thr Ala Thr Gln Ser Asn
      100            105            110

Ile Cys Asp Glu Asp Ser Ala Thr Glu Thr Cys Tyr Thr Tyr Asp Arg
      115            120            125

Asn Lys Cys Tyr Thr Ala Val Val Pro Leu Val Tyr Gly Gly Glu Thr
      130            135            140

Lys Met Val Glu Thr Ala Leu Thr Pro Asp Ala Cys Tyr Pro Asp
      145            150            155

```

<210> SEQ ID NO 23

<211> LENGTH: 205

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<220> FEATURE:

<223> OTHER INFORMATION: Xaa may be any amino acid

<400> SEQUENCE: 23

```

Gly Leu Leu Lys Pro Ser Glu Thr Leu Ser Leu Thr Cys Ala Val Tyr
 1             5             10             15

Gly Gly Ser Phe Ser Gly Tyr Tyr Trp Ser Trp Ile Arg Gln Pro Pro
      20             25             30

Gly Lys Gly Leu Glu Trp Ile Gly Glu Ile Asn His Ser Gly Ser Thr
      35             40             45

Asn Tyr Asn Pro Ser Leu Lys Ser Arg Val Thr Ile Ser Val Asp Thr
      50             55             60

Ser Lys Asn Gln Phe Ser Leu Lys Leu Ser Ser Val Thr Ala Ala Asp
      65             70             75             80

Thr Ala Val Tyr Tyr Cys Ala Arg Gly Thr Thr Glu Tyr Tyr Tyr Tyr
      85             90             95

Tyr Tyr Gly Met Asp Val Trp Gly Gln Gly Thr Thr Val Thr Val Ser
      100            105            110

Ser Gly Ser Ala Ser Ala Pro Thr Leu Phe Pro Leu Val Ser Cys Glu
      115            120            125

Asn Ser Pro Ser Asp Thr Ser Ser Val Ala Val Gly Cys Leu Ala Gln
      130            135            140

Asp Phe Leu Pro Asp Xaa Ile Thr Phe Ser Trp Lys Tyr Lys Asn Asn
      145            150            155            160

Ser Asp Ile Ser Ser Thr Arg Gly Phe Pro Ser Val Leu Arg Gly Gly
      165            170            175

Lys Tyr Ala Ala Thr Ser Gln Val Leu Leu Pro Ser Lys Asp Val Met

```

```

<210> SEQ ID NO 25
<211> LENGTH: 197
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 25

Leu Val Lys Pro Ser Glu Thr Leu Ser Leu Thr Cys Thr Val Ser Gly
  1             5             10             15
Gly Ser Ile Ser Ser Tyr Tyr Trp Asn Trp Ile Arg Gln Pro Pro Gly
             20             25             30
Lys Gly Leu Glu Trp Ile Gly Tyr Ile Tyr Tyr Ser Gly Ser Thr Asn
             35             40             45
Tyr Asn Pro Ser Leu Lys Ser Arg Val Thr Ile Ser Val Asp Thr Ser
             50             55             60
Lys Asn Gln Phe Ser Leu Lys Leu Ser Ser Val Thr Ala Ala Asp Thr
             65             70             75             80
Ala Val Tyr Tyr Cys Ala Arg Asp Arg Gly Val Gly Ala Thr Gly Phe
             85             90             95
Asp Tyr Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser Gly Ser Ala
             100            105            110
Ser Ala Pro Thr Leu Phe Pro Leu Val Ser Cys Glu Asn Ser Pro Ser
             115            120            125
Asp Thr Ser Ser Val Ala Val Gly Cys Leu Ala Gln Asp Phe Leu Pro

```

-continued

130	135	140
Asp Ser Ile Thr Phe Ser Trp Lys Tyr Lys Asn Asn Ser Asp Ile Ser		
145	150	155 160
Ser Thr Arg Gly Phe Pro Ser Val Leu Arg Gly Gly Lys Tyr Ala Ala		
	165	170 175
Thr Ser Gln Val Leu Leu Pro Ser Lys Asp Val Met Gln Gly Thr Asp		
	180	185 190
Glu His Lys Val Cys		
195		

<210> SEQ ID NO 26

<211> LENGTH: 147

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 26

Ser Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly Glu Arg Val Thr		
1	5	10 15
Ile Thr Cys Arg Ala Ser Gln Gly Ile Arg Asp Glu Leu Gly Trp Tyr		
	20	25 30
Gln Gln Lys Pro Gly Lys Ala Pro Lys Arg Leu Ile Tyr Val Ala Ser		
	35	40 45
Ser Leu Gln Ser Gly Val Pro Ser Arg Phe Ser Gly Ser Gly Ser Gly		
	50	55 60
Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Glu Asp Phe Ala		
	65	70 75 80
Thr Tyr Tyr Cys Leu Gln His Asn Gly Tyr Pro Arg Thr Phe Gly Gln		
	85	90 95
Gly Thr Lys Val Glu Ile Lys Arg Thr Val Ala Ala Pro Ser Val Phe		
	100	105 110
Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly Thr Ala Ser Val		
	115	120 125
Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala Lys Glu His Gln		
	130	135 140
Lys Ser Pro		
145		

<210> SEQ ID NO 27

<211> LENGTH: 203

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 27

Lys Lys Pro Gly Ala Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr		
1	5	10 15
Thr Phe Thr Ser Tyr Asp Ile Asn Trp Val Arg Gln Ala Thr Gly Gln		
	20	25 30
Gly Leu Glu Trp Met Gly Trp Met Asn Pro Asn Ser Gly Asn Thr Gly		
	35	40 45
Tyr Ala Gln Lys Phe Gln Gly Arg Val Thr Met Asn Arg Asn Thr Ser		
	50	55 60
Ile Ser Thr Ala Tyr Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr		
	65	70 75 80
Ala Val Tyr Tyr Cys Ala Arg Gly Gly His Gly Gly Ser Tyr Phe Tyr		

-continued

85										90										95									
Ser	Tyr	Tyr	Gly	Met	Asp	Val	Trp	Gly	Gln	Gly	Thr	Thr	Val	Thr	Val														
			100					105					110																
Ser	Ser	Gly	Ser	Ala	Ser	Ala	Pro	Thr	Leu	Phe	Pro	Leu	Val	Ser	Cys														
		115					120					125																	
Glu	Asn	Ser	Pro	Ser	Asp	Thr	Ser	Ser	Val	Ala	Val	Gly	Cys	Leu	Ala														
		130				135					140																		
Gln	Asp	Phe	Leu	Pro	Asp	Ser	Ile	Thr	Phe	Ser	Trp	Lys	Tyr	Lys	Asn														
145					150					155					160														
Asn	Ser	Asp	Ile	Ser	Ser	Thr	Arg	Gly	Phe	Pro	Ser	Val	Leu	Arg	Gly														
			165					170					175																
Gly	Lys	Tyr	Ala	Ala	Thr	Ser	Gln	Val	Leu	Leu	Pro	Ser	Lys	Asp	Val														
			180				185						190																
Met	Gln	Gly	Thr	Asp	Glu	His	Val	Val	Cys	Lys																			
		195					200																						

<210> SEQ ID NO 28

<211> LENGTH: 149

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 28

His	Ser	Leu	Ala	Val	Ser	Leu	Gly	Glu	Arg	Ala	Thr	Ile	Asn	Cys	Lys			
1				5					10					15				
Ser	Ser	Gln	Ser	Val	Leu	Tyr	Ser	Phe	Asn	Asn	Lys	Asn	Tyr	Leu	Ala			
		20					25						30					
Trp	Tyr	Gln	Gln	Lys	Pro	Gly	Gln	Pro	Pro	Lys	Leu	Leu	Ile	Tyr	Trp			
		35				40						45						
Ala	Ser	Thr	Arg	Glu	Ser	Gly	Val	Pro	Asp	Arg	Phe	Gly	Gly	Ser	Gly			
		50				55					60							
Ser	Gly	Thr	Asp	Phe	Thr	Leu	Thr	Ile	Ser	Ser	Leu	Gln	Ala	Glu	Asp			
65				70					75					80				
Val	Ala	Val	Tyr	Tyr	Cys	Gln	Gln	Tyr	Tyr	Ser	Thr	Pro	Arg	Thr	Phe			
			85					90						95				
Gly	Gln	Gly	Thr	Lys	Val	Glu	Ile	Lys	Arg	Thr	Val	Ala	Ala	Pro	Ser			
		100						105					110					
Val	Phe	Ile	Phe	Pro	Pro	Ser	Asp	Glu	Gln	Leu	Lys	Ser	Gly	Thr	Ala			
		115					120					125						
Ser	Val	Val	Cys	Leu	Leu	Asn	Asn	Phe	Tyr	Pro	Arg	Glu	Ala	Lys	Glu			
		130				135					140							
His	Gln	Lys	Ser	Pro														
145																		

<210> SEQ ID NO 29

<211> LENGTH: 199

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 29

Glu	Val	Lys	Lys	Pro	Gly	Ala	Ser	Val	Lys	Val	Ser	Cys	Lys	Ala	Ser			
1				5					10					15				
Gly	Tyr	Thr	Phe	Thr	Ser	Tyr	Asp	Ile	Asn	Trp	Val	Arg	Gln	Ala	Thr			
		20					25						30					
Gly	Gln	Gly	Leu	Glu	Trp	Met	Gly	Trp	Met	Asn	Pro	Asn	Ser	Gly	Asn			

-continued

35					40					45					
Thr	Gly	Tyr	Ala	Gln	Lys	Phe	Gln	Gly	Arg	Val	Thr	Met	Thr	Arg	Asn
50						55					60				
Thr	Ser	Ile	Ser	Thr	Ala	Tyr	Met	Glu	Leu	Ser	Ser	Leu	Arg	Ser	Glu
65					70					75					80
Asp	Thr	Ala	Val	Tyr	Tyr	Cys	Ala	Arg	Glu	Glu	Trp	Leu	Val	Arg	Tyr
				85					90					95	
Tyr	Gly	Met	Asp	Val	Trp	Gly	Gln	Gly	Thr	Thr	Val	Thr	Val	Ser	Ser
			100					105					110		
Gly	Ser	Ala	Ser	Ala	Pro	Thr	Leu	Phe	Pro	Leu	Val	Ser	Cys	Glu	Asn
			115				120					125			
Ser	Pro	Ser	Asp	Thr	Ser	Ser	Val	Ala	Val	Gly	Cys	Leu	Ala	Gln	Asp
	130					135					140				
Phe	Leu	Pro	Asp	Ser	Ile	Thr	Phe	Ser	Trp	Lys	Tyr	Lys	Asn	Asn	Ser
145					150					155					160
Asp	Ile	Ser	Ser	Thr	Arg	Gly	Phe	Pro	Ser	Val	Leu	Arg	Gly	Gly	Lys
				165					170					175	
Tyr	Ala	Ala	Thr	Ser	Gln	Val	Leu	Leu	Pro	Ser	Lys	Asp	Val	Met	Gln
			180					185					190		
Gly	Thr	Asp	Glu	His	Lys	Val									
	195														

<210> SEQ ID NO 30

<211> LENGTH: 147

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<220> FEATURE:

<223> OTHER INFORMATION: Xaa may be any amino acid

<400> SEQUENCE: 30

Gly	Gln	Ser	Pro	Ser	Ser	Leu	Ser	Ala	Ser	Val	Gly	Asp	Arg	Val	Thr
1				5					10					15	
Ile	Thr	Cys	Arg	Ala	Ser	Gln	Asp	Ile	Arg	Asp	Asn	Leu	Gly	Trp	Tyr
		20					25					30			
Gln	Gln	Lys	Pro	Gly	Lys	Ala	Pro	Lys	Arg	Leu	Ile	Tyr	Ala	Ala	Ser
		35				40						45			
Asn	Leu	Gln	Ser	Gly	Val	Pro	Ser	Arg	Phe	Ser	Gly	Ser	Gly	Ser	Gly
	50				55						60				
Thr	Glu	Phe	Thr	Leu	Thr	Ile	Ser	Ser	Leu	Gln	Pro	Glu	Asp	Phe	Ala
65				70						75					80
Thr	Tyr	Tyr	Cys	Leu	Gln	Tyr	Lys	Thr	Tyr	Pro	Trp	Thr	Phe	Gly	Gln
			85						90					95	
Gly	Thr	Lys	Val	Glu	Ile	Lys	Arg	Thr	Val	Ala	Ala	Pro	Ser	Val	Phe
			100					105					110		
Ile	Phe	Pro	Pro	Ser	Asp	Glu	Gln	Leu	Lys	Ser	Gly	Thr	Ala	Ser	Val
		115					120					125			
Val	Cys	Leu	Leu	Asn	Asn	Phe	Tyr	Pro	Arg	Glu	Xaa	Lys	Glu	His	Gln
	130					135					140				
Lys	Ser	Pro													
145															

<210> SEQ ID NO 31

<211> LENGTH: 202

<212> TYPE: PRT

-continued

<213> ORGANISM: Homo sapiens

<220> FEATURE:

<223> OTHER INFORMATION: Xaa may be any amino acid

<400> SEQUENCE: 31

Lys Leu Pro Glu Thr Leu Ser Leu Thr Cys Ala Val Tyr Gly Gly Ser
 1 5 10 15
 Phe Ser Gly Tyr Tyr Trp Ser Trp Ile Arg Gln Pro Pro Gly Lys Gly
 20 25 30
 Leu Glu Trp Ile Gly Glu Ile Asn His Ser Gly Ser Thr Asn Tyr Asn
 35 40 45
 Pro Ser Leu Lys Ser Arg Val Thr Ile Ser Val Asp Thr Ser Lys Asn
 50 55 60
 Gln Phe Ser Leu Lys Leu Ser Ser Val Thr Ala Ala Asp Thr Ala Val
 65 70 75 80
 Tyr Tyr Cys Ala Arg Gly Ala Ala Glu Tyr Tyr Tyr Tyr Tyr Gly
 85 90 95
 Met Asp Val Trp Gly Gln Gly Thr Thr Val Thr Val Ser Ser Gly Ser
 100 105 110
 Ala Ser Ala Pro Thr Leu Phe Pro Leu Val Ser Cys Glu Asn Ser Pro
 115 120 125
 Ser Asp Thr Ser Ser Val Ala Val Gly Cys Leu Ala Gln Asp Phe Leu
 130 135 140
 Pro Asp Xaa Ile Thr Phe Xaa Trp Lys Tyr Lys Asn Asn Ser Asp Ile
 145 150 155 160
 Ser Ser Thr Arg Gly Phe Pro Ser Val Leu Arg Gly Gly Lys Tyr Ala
 165 170 175
 Ala Thr Ser Gln Val Leu Leu Pro Ser Lys Asp Val Met Gln Gly Thr
 180 185 190
 Asp Glu His Val Val Thr Gly Ser Lys Glu
 195 200

<210> SEQ ID NO 32

<211> LENGTH: 143

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 32

Met Pro Val Thr Pro Gly Glu Pro Ala Ser Ile Ser Cys Arg Ser Ser
 1 5 10 15
 Gln Ser Leu Leu His Ser Asn Gly Tyr Asn Tyr Leu Asp Trp Tyr Leu
 20 25 30
 Gln Lys Pro Gly Gln Ser Pro Gln Leu Leu Ile Tyr Leu Gly Ser Asn
 35 40 45
 Arg Ala Ser Gly Val Pro Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr
 50 55 60
 Asp Phe Thr Leu Lys Ile Ser Arg Val Glu Ala Glu Asp Val Gly Ile
 65 70 75 80
 Tyr Tyr Cys Met Gln Ser Leu Gln Ile Pro Arg Leu Phe Gly Pro Gly
 85 90 95
 Thr Lys Val Asp Ile Lys Arg Thr Val Ala Ala Pro Ser Val Phe Ile
 100 105 110
 Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly Thr Ala Ser Val Val
 115 120 125

-continued

Cys Leu Leu Ser Asn Phe Tyr Pro Arg Glu Ala Lys Val Gln Trp
130 135 140

<210> SEQ ID NO 33
<211> LENGTH: 190
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 33

Ser Glu Thr Leu Ser Leu Thr Cys Ala Val Tyr Gly Gly Ser Phe Ser
1 5 10 15
Gly Tyr Tyr Trp Ser Trp Ile Arg Gln Pro Pro Gly Lys Gly Leu Glu
20 25 30
Trp Ile Gly Glu Ile Asn His Ser Gly Ser Thr Asn Tyr Asn Pro Ser
35 40 45
Leu Lys Ser Arg Val Thr Ile Ser Val Asp Thr Ser Lys Asn Gln Phe
50 55 60
Ser Leu Lys Leu Ser Ser Val Thr Ala Ala Asp Thr Ala Val Tyr Tyr
65 70 75 80
Cys Ala Arg Gly Gly Thr Thr Val Thr Phe Asp Ala Phe Asp Ile Trp
85 90 95
Gly Gln Gly Thr Met Val Thr Val Ser Ser Gly Ser Ala Ser Ala Pro
100 105 110
Thr Leu Phe Pro Leu Val Ser Cys Glu Asn Ser Pro Ser Asp Thr Ser
115 120 125
Ser Val Ala Val Gly Cys Leu Ala Gln Asp Phe Leu Pro Asp Ser Ile
130 135 140
Thr Phe Ser Trp Lys Tyr Lys Asn Asn Ser Asp Ile Ser Ser Thr Arg
145 150 155 160
Gly Phe Pro Ser Val Leu Arg Gly Gly Lys Tyr Ala Ala Thr Ser Gln
165 170 175
Val Leu Leu Pro Ser Lys Asp Val Met Gln Gly Thr Asp Glu
180 185 190

<210> SEQ ID NO 34
<211> LENGTH: 147
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 34

Leu Ala Val Ser Leu Gly Glu Arg Ala Thr Ile Asn Cys Lys Ser Ser
1 5 10 15
Gln Ser Val Leu Tyr Ser Phe Asn Asn Lys Asn Tyr Leu Ala Trp Tyr
20 25 30
Gln Gln Lys Pro Gly Gln Pro Pro Lys Leu Leu Ile Tyr Trp Ala Ser
35 40 45
Thr Arg Glu Ser Gly Val Pro Asp Arg Phe Ser Gly Ser Gly Ser Gly
50 55 60
Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Ala Glu Asp Val Ala
65 70 75 80
Val Tyr Tyr Cys Gln Gln Tyr Tyr Ser Thr Pro Arg Thr Phe Gly Gln
85 90 95
Gly Thr Lys Val Glu Ile Lys Arg Thr Val Ala Ala Pro Ser Val Phe
100 105 110

-continued

Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly Thr Ala Ser Val
 115 120 125

Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala Lys Val Gln Trp
 130 135 140

Lys Val Ile
 145

<210> SEQ ID NO 35
 <211> LENGTH: 149
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 35

Asn Pro Gln Thr Thr Leu Thr Leu Thr Cys Thr Phe Ser Gly Phe Ser
 1 5 10 15

Leu Ile Thr Arg Gly Val Gly Val Asp Trp Ile Arg Gln Pro Pro Gly
 20 25 30

Lys Ala Leu Gln Trp Leu Ala Leu Ile Tyr Trp Asn Asp Asp Lys Arg
 35 40 45

Tyr Ser Pro Ser Leu Lys Ser Arg Leu Thr Ile Thr Lys Asp Thr Ser
 50 55 60

Lys Asn Gln Val Val Leu Thr Met Thr Asn Met Asp Pro Val Asp Thr
 65 70 75 80

Ala Thr Tyr Tyr Cys Ala His His Phe Phe Asp Ser Ser Gly Tyr Tyr
 85 90 95

Pro Phe Asp Ser Trp Gly Gln Gly Thr Leu Val Ser Val Ser Ser Ala
 100 105 110

Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Cys Ser Arg Ser
 115 120 125

Thr Ser Glu Ser Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr Phe
 130 135 140

Pro Glu Pro Val Thr
 145

<210> SEQ ID NO 36
 <211> LENGTH: 148
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 36

Val Thr Gln Ser Pro Leu Ser Leu Ser Val Thr Pro Gly Gln Pro Ala
 1 5 10 15

Ser Ile Ser Cys Lys Ser Ser Gln Ser Leu Leu His Ser Asp Gly Lys
 20 25 30

Thr Tyr Leu Tyr Trp Tyr Leu Gln Lys Pro Gly Gln Pro Pro Gln Leu
 35 40 45

Leu Ile Tyr Glu Ala Phe Asn Arg Phe Ser Gly Val Pro Asp Arg Phe
 50 55 60

Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile Ser Arg Val
 65 70 75 80

Glu Ala Glu Asp Val Gly Leu Tyr Tyr Cys Met Gln Ser Ile Glu Leu
 85 90 95

Pro Phe Thr Phe Gly Gly Gly Thr Lys Val Glu Ile Lys Arg Thr Val
 100 105 110

-continued

Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys
 115 120 125

Ser Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg
 130 135 140

Lys Glu Arg Val
 145

<210> SEQ ID NO 37
 <211> LENGTH: 173
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 37

Gly Glu Gly Leu Val Lys Pro Gly Gly Ser Leu Arg Leu Ser Cys Ala
 1 5 10 15

Ala Ser Gly Phe Thr Phe Ser Ser Tyr Ser Met Asn Trp Val Arg Gln
 20 25 30

Ala Pro Gly Lys Gly Leu Glu Trp Val Ser Ser Ile Ser Ser Ser Ser
 35 40 45

Ser Tyr Ile Tyr Tyr Ala Asp Ser Val Lys Gly Arg Phe Thr Ile Ser
 50 55 60

Arg Asp Asn Ala Lys Asn Ser Leu Tyr Leu Gln Met Asn Ser Leu Arg
 65 70 75 80

Ala Glu Asp Thr Ala Val Tyr Tyr Cys Ala Arg Asp Ser Ser Gly Trp
 85 90 95

Tyr Glu Asp Tyr Phe Asp Tyr Trp Gly Gln Gly Thr Leu Val Thr Val
 100 105 110

Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Cys
 115 120 125

Ser Arg Ser Thr Ser Glu Ser Thr Ala Ala Leu Gly Cys Leu Val Lys
 130 135 140

Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu
 145 150 155 160

Thr Ser Gly Val His Thr Phe Pro Ala Val Leu Gln Ser
 165 170

<210> SEQ ID NO 38
 <211> LENGTH: 101
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 38

Leu Asp Ile Gln Leu Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val
 1 5 10 15

Gly Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Asp Ile Ser Ile
 20 25 30

Tyr Leu Ala Trp Phe Gln Gln Arg Pro Gly Lys Ala Pro Lys Ser Leu
 35 40 45

Ile Tyr Ala Ala Ser Ser Leu Gln Ser Gly Val Pro Ser Lys Phe Ser
 50 55 60

Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln
 65 70 75 80

Pro Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln Tyr Asn Ser Tyr Pro
 85 90 95

-continued

Phe Thr Phe Gly Pro
100

<210> SEQ ID NO 39

<211> LENGTH: 159

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 39

Leu Thr Cys Thr Phe Ser Gly Phe Ser Leu Ile Thr Arg Gly Val Gly
1 5 10 15
Val Asp Trp Ile Arg Gln Pro Pro Gly Lys Ala Leu Gln Trp Leu Ala
20 25 30
Leu Ile Tyr Trp Asn Asp Asp Lys Arg Tyr Ser Pro Ser Leu Lys Ser
35 40 45
Arg Leu Thr Ile Thr Lys Asp Thr Ser Lys Asn Gln Val Val Leu Thr
50 55 60
Met Thr Asn Met Asp Pro Val Asp Thr Ala Thr Tyr Tyr Cys Ala His
65 70 75 80
His Phe Phe Asp Ser Ser Gly Tyr Tyr Pro Phe Asp Ser Trp Gly Gln
85 90 95
Gly Thr Leu Val Ser Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val
100 105 110
Phe Pro Leu Ala Pro Cys Ser Arg Ser Thr Ser Glu Ser Thr Ala Ala
115 120 125
Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser
130 135 140
Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Gln Leu
145 150 155

<210> SEQ ID NO 40

<211> LENGTH: 167

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 40

Gly Gly Gly Leu Val Gln Pro Gly Gly Ser Leu Arg Leu Ser Cys Ala
1 5 10 15
Ala Ser Gly Phe Thr Phe Ser Ser Tyr Ala Met Ser Trp Val Arg Gln
20 25 30
Ala Pro Gly Lys Gly Leu Glu Trp Val Ser Thr Ile Ser Val Ser Gly
35 40 45
Ile Thr Thr Tyr Tyr Val Asp Ser Val Lys Gly Arg Phe Thr Ile Ser
50 55 60
Arg Asp Asn Ser Lys Asn Ile Leu Tyr Leu Gln Met Asn Ser Leu Arg
65 70 75 80
Ala Glu Asp Thr Ala Val Tyr Tyr Cys Ala Lys Arg Ile Phe Gly Val
85 90 95
Val Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser Ala Ser Thr Lys
100 105 110
Gly Pro Ser Val Phe Pro Leu Ala Pro Cys Ser Arg Ser Thr Ser Glu
115 120 125
Ser Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro
130 135 140

-continued

Val Thr Val Ser Trp Asn Leu Gly Ala Leu Thr Ser Gly Val His Thr
145 150 155 160

Phe Pro Ala Val Leu Gln Ser
165

<210> SEQ ID NO 41
<211> LENGTH: 164
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 41

Gly Ile Arg Leu Asp Ile Gln Leu Thr Gln Ser Pro Ser Ser Leu Ser
1 5 10 15

Ala Ser Val Gly Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Gly
20 25 30

Ile Ser Ile Tyr Leu Ala Trp Phe Gln Gln Arg Pro Gly Lys Ala Pro
35 40 45

Lys Ser Leu Ile Tyr Ala Ala Ser Ser Leu Gln Ser Gly Val Pro Ser
50 55 60

Lys Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser
65 70 75 80

Ser Leu Gln Pro Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln Tyr Asn
85 90 95

Ser Tyr Pro Phe Thr Phe Gly Pro Gly Thr Lys Val Asp Ile Lys Arg
100 105 110

Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln
115 120 125

Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr
130 135 140

Pro Arg Glu Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser
145 150 155 160

Gly Lys Pro Asn

<210> SEQ ID NO 42
<211> LENGTH: 35
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 42

gactacgaat tcttgtagga ccggcgagga atagg 35

<210> SEQ ID NO 43
<211> LENGTH: 37
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 43

gactacgggc ccggtgagaa cttggaatct tgcaagc 37

<210> SEQ ID NO 44
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: primer

<400> SEQUENCE: 44

-continued

gcagttctcct aaactgct 18

<210> SEQ ID NO 45
<211> LENGTH: 15
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: primer

<400> SEQUENCE: 45

acctgcaagg ccagt 15

<210> SEQ ID NO 46
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: primer

<400> SEQUENCE: 46

cactcattcc tgttgaag 18

<210> SEQ ID NO 47
<211> LENGTH: 500
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 47

tcagaagaag tgaagtcaag atgaagaacc atttgctttt ctggggagtc ctggcggttt 60

ttattaaggc tgtttcatgtg aaagcccaag aagatgaaag gattgttctt gttgacaaca 120

aatgtaagtg tgcccggatt acttccagga tcatccgttc ttccgaagat cctaattgagg 180

acattgtgga gagaacatc cgaattattg ttccctctgaa caacaggagg aatatctctg 240

atcccacctc accattgaga accagatttg tgtaccattt gtctgacctc tgtaaaaaat 300

gtgatccctac agaagtgagg ctggataatc agatagttac tgctaccag agcaatatct 360

gtgatgaaga cagtgctaca gagacctgct acacttatga cagaacaag tgctacacag 420

ctgtggtccc actcgtatat ggtggtgaga ccaaatggt ggaaacagcc ttaacccag 480

atgcctgcta tcctgactaa 500

<210> SEQ ID NO 48
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: primer

<400> SEQUENCE: 48

gaattcagaa gaagtgaagt c 21

<210> SEQ ID NO 49
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: primer

<400> SEQUENCE: 49

gtcgactatg cagtcagcaa tgac 24

-continued

<210> SEQ ID NO 50
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: primer

<400> SEQUENCE: 50

tgcaggaatc agaccagtc 20

<210> SEQ ID NO 51
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: primer

<400> SEQUENCE: 51

gtcaggctgg aactgaggag ca 22

<210> SEQ ID NO 52
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: primer

<400> SEQUENCE: 52

tcatttggtg atcagcact 19

<210> SEQ ID NO 53
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: primer

<400> SEQUENCE: 53

gctagctgag gagacggtga ccagg 25

<210> SEQ ID NO 54
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: primer

<400> SEQUENCE: 54

tcatttggtg atcagcact 19

<210> SEQ ID NO 55
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: primer

<400> SEQUENCE: 55

ggatcctgag gagacggtga cg 22

<210> SEQ ID NO 56
<211> LENGTH: 24

-continued

<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: primer

<400> SEQUENCE: 56
ggattagcat ccgccccaac cctt 24

<210> SEQ ID NO 57
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: primer

<400> SEQUENCE: 57
gtcgacgcac acacagagcg gccca 24

<210> SEQ ID NO 58
<211> LENGTH: 37
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 58
gactacgaat tcggaccggc gaggaatagg aatcatg 37

<210> SEQ ID NO 59
<211> LENGTH: 43
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 59
ggatggtggtt ggtagctagc acgcggagcg tgatgatggc ctg 43

<210> SEQ ID NO 60
<211> LENGTH: 33
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 60
gactacgaat tcacgagcg acatggcggc ggc 33

<210> SEQ ID NO 61
<211> LENGTH: 43
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 61
ggatggtggtt ggtagctagc acacgcagtg agatggtttc ccg 43

<210> SEQ ID NO 62
<211> LENGTH: 617
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<223> OTHER INFORMATION: "n" may be A, T, C, G, unknown or other

<400> SEQUENCE: 62
ggactgttga agccttcgga gaccctgtcc ctcacctgcg ctgtctatgg tgggtccttc 60
agtggttact actggagctg gatccgccag cccccaggga aggggctgga gtggattggg 120
gaaatcaatc atagtgggaag caccaactac aaccctgcc tcaagagtcg agtcaccata 180

-continued

tcagtagaca cgtccaagaa ccagttctcc ctgaagctga gctctgtgac cgengcggac	240
acggctgtgt attactgtgc gagaggcact acggaatatt actactacta ctacggtatg	300
gacgtctggg gccaaaggac cacggtcacc gtctcctcag ggagtgcac cgccccaacc	360
cttttcccc tcgtctcctg tgagaattcc ccgtcggata cgagcagcgt ggccgttggc	420
tgccctgcac aggacttcct tcccgactyc atcactttct cctggaaata caagaacaac	480
tctgacatca cgagcaccgc gggcttccca tcagtcctga gagggggcaa gtacgcagcc	540
acctcacagg tgctgctgcc ttccaaggac gtcctgcagg gcacagacga acacgtggtg	600
acgggatcca aagagta	617

<210> SEQ ID NO 63
 <211> LENGTH: 444
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 63

ctctccctgc ccgtcaccgc tggagagccg gcctccatct cctgcaggtc tagtcagagc	60
ctcctgcata gtaatggata caactatttg gattgggtacc tgcagaagcc agggcagctc	120
ccacagctcc tgatctatct ggggttcta atcgccctccg gggctccctga cagggttcagt	180
ggcagtggtat caggcacaga ttttacctg aaaatcagca gagtggaggc tgaggatgtt	240
gggatttatt actgcctgca gactcgacaa actcctcgga cgttcggcca agggaccaag	300
gtggaaatca aacgaactgt ggctgcacca tctgtcttca tcttcccgc atctgatgag	360
cagttgaaat ctggaactgc ctctgtgtg tgctgctga ataacttcta tcccagagag	420
gccaagagc atcaaaagag tcca	444

<210> SEQ ID NO 64
 <211> LENGTH: 593
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 64

ctggtgaagc ctccggagac cctgtccctc acctgcactg tctctggtgg ctccatcagt	60
agttactact ggaactggat ccggcagccc ccagggaagg gactggagtg gattgggtat	120
atctattaca gtgggagcac caactacaac ccctccctca agagtcgagt caccatatca	180
gtagacacgt ccaagaacca gttctccctg aagctgagct ctgtgaccgc tgcggacacg	240
gccgtgtatt actgtgcgag agatagggga gtgggagcta ctggttttga ctactggggc	300
cagggaaccc tggtcaccgt ctctcaggg agtgcacccg cccaaccct tttcccctc	360
gtctcctgtg agaattcccc gtccgatacg agcagcgtgg ccgttggtg cctcgacacg	420
gacttccttc ccgactccat cactttctcc tggaaatata agaacaactc tgacatcagc	480
agcaccggg gcttcccatc agtcctgaga gggggcaagt acgcagccac ctacaggtg	540
ctgctgcctt ccaaggacgt catgcagggc acagacgaac acaaggtgtg cga	593

<210> SEQ ID NO 65
 <211> LENGTH: 441
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 65

-continued

```

agccagtctc catcctccct gtctgcatct gtaggagaga gagtoacccat cacttgccgg      60
gcaagtcagg gcattagaga tgaattaggc tggatcagc agaaaccagg gaaagcccct      120
aagcgcctga tctatgttgc atccagtttg caaagtgggg tcccatcaag gttcagcggc      180
agtggatctg ggacagaatt cactctcaca atcagcagcc tgcagcctga agattttgca      240
acttattact gtctacagca taatggttac cctcggacgt tcggccaagg gaccaagggtg      300
gaaatcaaac gaactgtggc tgcaccatct gtcttcatct tcccgccatc tgatgagcag      360
ttgaaatctg gaactgcctc tgttgtgtgc ctgctgaata acttctatcc cagagaggcc      420
aaagagcatc aaaagagtcc a                                                  441

```

```

<210> SEQ ID NO 66
<211> LENGTH: 610
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 66

```

```

aagaagcctg gggcctcagt gaaggctctc tgcaaggctt ctggatacac cttcaccagt      60
tatgatatca actgggtgcg acagggccact ggacaagggc ttgagtggat gggatggatg      120
aaccctaaca gtggtaaacac aggcctatgca cagaagttcc agggcagagt caccatgaac      180
aggaacacct ccataagcac agcctacatg gagctgagca gcctgagatc tgaggacacg      240
gccgtgtatt actgtgcgag aggggggtcat ggtgggagct acttotactc ctaytacggt      300
atggacgtct ggggccaggg gaccacggtc accgtctcct cagggagtgc atccgccccca      360
acccttttcc ccctcgtctc ctgtgagaat tccccgtcgg atacgagcag cgtggccggt      420
ggctgcctcg cacaggactt ccttcccgcac tccatcactt tctcctggaa atacaagaac      480
aactctgaca tcagcagcac ccggggcttc ccatcagtcc tgagaggggg caagtacgca      540
gccacctcac aggtgctgct gccttccaag gacgtcatgc agggcacaga cgaacacgtg      600
gtgtgcaaac                                                                610

```

```

<210> SEQ ID NO 67
<211> LENGTH: 447
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 67

```

```

cactccctgg ctgtgtctct gggcgagagg gccaccatca actgcaagtc cagccagagt      60
gttttataca gttttaacaa taagaactac ttagcttggt accagcagaa accaggacag      120
cctcctaagc tgetcattta ctgggcatct acccggaat ccgggggtccc tgaccgattc      180
ggtggcagcg ggtctgggac agatttcact ctcaccatca gcagcctgca ggctgaagat      240
gtggcagttt attactgtca gcaatattat agtactcctm ggacgttcgg ccaaggggacc      300
aaggtggaat tcaaacgaac tgtggctgca ccatctgtct tcattctccc gccatctgat      360
gagcagttga aatctggaac tgcctctggt gtgtgcctgc tgaataactt ctatcccaga      420
gaggccaaag agcatcaaaa gagtcca                                          447

```

```

<210> SEQ ID NO 68
<211> LENGTH: 599
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

```

-continued

<400> SEQUENCE: 68

gaggtgaaga agcctggggc ctcagtgaag gtctcctgca aggtctcttg atacaccttc	60
accagttatg atatcaactg ggtgcgacag gccactggac aagggcttga gtggatggga	120
tggatgaacc ctaacagtgg taacacaggc tatgcacaga agttccaggg cagagtcacc	180
atgaccagga acacctccat aagcacagcc tacatggagc tgagcagcct gagatctgag	240
gacacggccg tgtattactg tgcgagagag gagtggctgg tacgttacta cggtatggac	300
gtctggggcc aagggaccac ggtcacgctc tcctcaggga gtgcacccgc cccaacctt	360
ttccccctcg tctcctgtga gaattccccg tcggatacga gcagcgtggc cgttggtgc	420
ctcgcacagg acttccttcc cgactccatc actttctcct ggaaatacaa gaacaactct	480
gacatcagca gcaccgggg cttcccatca gtctgagag ggggcaagta cgcagccacc	540
tcacaggtgc tgctgccttc caaggacgct atgcagggca cagacgaaca caaggtgtg	599

<210> SEQ ID NO 69

<211> LENGTH: 441

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 69

ggccagtctc catcctccct gtctgcatct gtaggagaca gagtcacat cacttgccgg	60
gcaagtcagg acattagaga taatttaggc tggatcagc agaaaccagg gaaagcccct	120
aagcgcctga tctatgtgc atccaatttg caaagtggg tcccatcaag gttcagcggc	180
agtggatctg ggacagaatt cactctcaca atcagcagcc tgcagcctga agattttgca	240
acttattact gtctacagta taaaacttac ccgtggacgt tcggccaagg gaccaagggtg	300
gaaatcaaac gaactgtggc tgcacatct gtcttcatct tcccgccatc tgatgagcag	360
ttgaaatctg gaactgcctc tgttgtgtgc ctgctgaata acttctatcc cagagaggmc	420
aaagagcatc aaaagagtcc a	441

<210> SEQ ID NO 70

<211> LENGTH: 607

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 70

aagcttcagg agaccctgtc cctcacctgc gctgtctatg gtgggtcctt cagtggttac	60
tactggagct ggatccgcca gccccaggg aaggggctgg agtggattgg ggaaatcaat	120
catagtggaa gcaccaacta caaccgtcc ctcaagagtc gagtcacat atcagtagac	180
acgtccaaga accagttctc cctgaagctg agctctgtga ccgcccggga cagggtgtg	240
tattactgtg cgagaggggc agctgaatat tactactact actacggtat ggacgtctgg	300
ggccaaggga ccacggtcac cgtctcctca gggagtgcac ccgcccac cttttcccc	360
ctcgtctcct gtgagaattc ccgctcggat acgagcagcg tggccgttgg ctgcctcgca	420
caggacttcc ttcccacty catcactttc tyctggaaat acaagaacaa ctctgacatc	480
agcagcaccg ggggcttccc atcagtcctg agagggggca agtacgcagc cacctcacag	540
gtgctgtgtc cttccaagga cgtcatgcag ggcacagacg aacacgtggg gacgggatcc	600
aaagagt	607

-continued

<210> SEQ ID NO 71
<211> LENGTH: 431
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 71

atgcccgtca cccctggaga gccggcctcc atctcctgca ggtctagtca gagcctcctg	60
catagtaatg gatacaacta tttggactgg tacctgcaga agccagggca gtctccacag	120
ctcctgatct atttgggttc taatcgggcc tccgggggtcc ctgacagggt cagtggcagt	180
ggatcaggca cagattttac actgaaaatc agcagagtgg aggctgagga tgttgggatt	240
tattactgca tgcaaatgtc acaaatcccc cggcttttcg gccctgggac caaagtggat	300
atcaaacgaa ctgtggctgc accatctgtc ttcattctcc cgccatctga tgagcagttg	360
aaatctggaa ctgcctctgt tgtgtgcctg ctgagtaact tctatcccag agaggccaaa	420
gtacagtgga a	431

<210> SEQ ID NO 72
<211> LENGTH: 570
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 72

tccggagacc tgctccctcac ctgcgctgtc tatgggtggg ccttcagtgg ttactactgg	60
agctggatcc gccagccccc aggggaaggg ctggagtgga ttggggaaat caatcatagt	120
ggaagcacca actacaaccc gtccctcaag agtcgagtca ccatatcagt agacacgtcc	180
aagaaccagt tctccctgaa gctgagttct gtgaccgccg cggacacggc tgtgtattac	240
tgtgcgagag gcgggactac agtaactttt gatgcttttg atatctgggg ccaagggaca	300
atggtcaccg tctcttcagg gagtgcaccc gcccacaccc ttttccccct cgtctcctgt	360
gagaattccc cgtcggatac gagcagcgtg gccgttggct gcctcgacac ggacttcctt	420
cccgaactcca tcaattttct ctggaaatac aagaacaact ctgacatcag cagcaccggg	480
ggcttcccat cagtccctgag agggggcaag tacgcagcca cctcacaggt gctgctgcct	540
tccaaggacg tcatgcaggg cacagacgaa	570

<210> SEQ ID NO 73
<211> LENGTH: 441
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 73

ctggctgtgt ctctgggcca gagggccacc atcaactgca agtccagcca gagtgtttta	60
tacagtttta acaataagaa ctacttagct tggtagcagc agaaaccagg acagcctcct	120
aagctgctca tttactgggc atctaccogg gaatccgggg tccctgaccg attcagtggc	180
agcgggtctg ggacagattt cactctcacc atcagcagcc tgcaggctga agatgtggca	240
gtttattact gtcagcaata ttatagtact cctcggacgt tcggccaagg gaccaagggtg	300
gaaatcaaac gaactgtggc tgcacatctt gtcttcatct tcccgccatc tgatgagcag	360
ttgaaatctg gaactgcctc tgttgtgtgc ctgctgaata acttctatcc cagagaggcc	420
aaagtacagt ggaaggatgat c	441

-continued

<210> SEQ ID NO 74

<211> LENGTH: 447

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 74

```
aaccacaga cgacctcac gctgacctgc accttctctg ggttctcact cattaccgt    60
ggagtgggtg tggattggt cgcacagccc ccaggaaagg cctgcagtg gctcgactc    120
atttattgga atgatgataa gcgtacagtc ccatctctga agagcaggct caccatcacc    180
aaggacacct ccaaaaacca ggtggtcctc acaatgacca acatggaccc tgtggacaca    240
gccacatatt actgtgcaca ccatttcttt gatagtagtg gttattaccc tttgactcc    300
tggggccagg gaacctgggt ctccgtctcc tcagcctcca ccaagggccc atcgggtctc    360
cccttgccgc cctgtccag gagcacctcc gagagcacag cggccctggg ctgcctggtc    420
aaggactact tccccgaacc ggtgacg                                     447
```

<210> SEQ ID NO 75

<211> LENGTH: 445

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 75

```
gtgactcagt ctccactctc tctgtccgtc acccctggac agccggcctc catctcctgc    60
aagtctagtc agagcctcct gcatagtgat ggaaagacct atttgattg gtacctgcag    120
aagccaggcc agcctccaca gctcctgac tatgaagctt tcaaccggtt ctctggagtg    180
ccagataggt tcagtggcag cgggtcaggg acagatttca cactgaaaat cagccgggtg    240
gaggctgagg atgttggtact ttattattgc atgcaaagta tagagcttcc gttcactttc    300
ggcggaggga ccaagggtga gatcaaagca actgtggctg caccatctgt cttcatcttc    360
ccgccatctg atgagcagtt gaaatctgga actgcctctg ttgtgtgcct gctgaataac    420
ttctatccca gaaaagaaa agtcr                                     445
```

<210> SEQ ID NO 76

<211> LENGTH: 519

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 76

```
ggggaaggcc tggtaagcc tgggggggtc ctgagactct cctgtgcagc ctctggattc    60
accttcagta gctatagcat gaactgggtc cgccaggctc cagggaaggg gctggagtgg    120
gtctcatcca ttagtagtag tagtagttac atatactacg cagactcagt gaagggccga    180
ttcacctct ccagagacaa gcccaagaac tcactgtatc tgcaaatgaa cagcctgaga    240
gccgaggaca cggtgtgta ttactgtgcg agggatagca gtggctggtg tgaggactac    300
tttgactact ggggccaggg aacctgggtc accgtctcct cagcctccac caagggccca    360
tcggtcttcc ccctggcgcc ctgctccagg agcacctccg agagcacagc ggccctgggc    420
tgcttggtca aggactactt ccccgaaacc gtgacggtgt cgtggaactc aggcgctctg    480
accagcggcg tgcacacctt cccagctgtc ctacagtca                                     519
```

-continued

<210> SEQ ID NO 77

<211> LENGTH: 303

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 77

cttgacatcc agctgaccca gtctccgtcc tcactgtctg catctgtagg agacagagtc	60
accatcactt gtcgggagag tcaggacatt agcatttatt tagcctggtt tcagcagaga	120
ccagggaag ccctaagtc cctgatctat gctgcatcca gtttgcaaag tggggcca	180
tcaaagtca gcggcagtgg atctgggaca gatttcactc tcaccatcag cagcctgcag	240
cctgaagatt ttgcaactta ttactgcca caatataata gttatccatt cactttcggg	300
ccc	303

<210> SEQ ID NO 78

<211> LENGTH: 477

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 78

ctgacctgca cttctctctg gttctcactc attaccctg gagtgggtgt ggattggatc	60
cgtcagcccc caggaaagc cctgcagtgg ctgcactca tttattggaa tgatgataag	120
cgctacagtc catctctgaa gagcaggctc accatcacca aggacacctc caaaaaccag	180
gtggtctca caatgaccaa catggacct gtggacacag ccacatatta ctgtgcacac	240
catttctttg atagtagtgg ttattacct tttgactcct ggggccaggg aacctggtc	300
tccgtctcct cagcctccac caagggccca tcggtcttcc cctggcgcc ctgctccagg	360
agcacctccg agagcacagc ggcctgggc tgctgtgtca aggactactt ccccgaaaccg	420
gtgacggtgt cgtggaactc aggcgctctg accagcgcg tgacacacctt ccagctg	477

<210> SEQ ID NO 79

<211> LENGTH: 503

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 79

gggggaggct tggtacagcc tggggggtcc ctgagactct cctgtgcagc ctctggatcc	60
acttttagca gctatgccat gagctgggtc cgccaggctc cagggaagg gctggagtgg	120
gtctcaacta ttagtgtagg tggattacc acatactacg tagactcctg gaagggccg	180
ttcacatct ccagagacaa ttccaagaac attctgtatc tgcaaatgaa cagcctgaga	240
gccgaggaca cggccgtata ttactgtgcg aaacggattt ttggagtggc ctggggccag	300
ggaacctggt tcacctctc ctacgcctcc accaagggcc catcggtctt cccctggcg	360
ccctgctcca ggagcacctc cgagagcaca gcggccctgg gctgcctggt caaggactac	420
ttccccgaac cggtgacggt gtcgtggaac ttaggcgctc tgaccagcgg cgtgcacacc	480
ttcccagctg tcctacagtc cta	503

<210> SEQ ID NO 80

<211> LENGTH: 494

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 80

-continued

```

ggaattcggc ttgatattca gctgactcag tctccatcct cactgtctgc atctgtagga    60
gacagagtca ccatcacttg tcgggagagt cagggcatta gcatttattt agcctgggtt    120
cagcagagac cagggaaagc ccctaagtcc ctgatctatg ctgcattccag ttgcaaagt    180
gggggtcccat caaagttcag cggcagtgga tctgggacag atttcactct caccatcagc    240
agcctgcagc ctgaagattt tgcaacttat tactgccaac aatataatag ttaccatttc    300
actttcggcc ctgggaccaa agtggatata aaacgaactg tggctgcacc atctgtcttc    360
atcttcccg cactctgatga gcagttgaaa tctggaactg cctctgttgt gtgcctgctg    420
aataacttct atcccagaga ggccaaagta cagtgggaag tggataacgc cctccaatcg    480
ggtaagccga attc                                         494

```

```

<210> SEQ ID NO 81
<211> LENGTH: 1774
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 81

```

```

atgtacttgg gactgaacta tgtattcata gtttttctct taaatggtgt ccagagtga    60
gtgaagcttg aggagtctgg aggaggcttg gtgcaacctg gaggatccat gaaactctcc    120
tgtgttgctt ctggattcac ttccagtaac tactggatga actgggtccg ccagtctcca    180
gagaaggggc ttgagtgggt tgctgaaatt agattgaaat ctaataatta tgcaacacat    240
tatgctggagt ctgtgaaagg gaggttcacc atctcaagag atgattccaa aagtagtgtc    300
tacctgcaaa tgaacaactt aagagctgaa gacactggca ttattactg tacggattac    360
gatgcttact ggggccaagg gactctggtc actgtctctg cagagagtca gtccttccca    420
aatgtcttcc ccctcgtctc ctgcgagagc ccctgtctg ataagaatct ggtggccatg    480
ggctgccttg cccgggactt cctgccagc accatttcct tcacctgga ctaccagaac    540
aacactgaag tcattccagg tatcagaacc ttcccaacac tgaggacagg gggcaagtac    600
ctagccacct cgcagggtgt gctgtctccc aagagcatcc ttgaaggttc agatgaatac    660
ctggtatgca aaatccacta cggaggcaaa aacagagatc tgcattgtgc cattccagct    720
gtcgagagaa tgaaccccaa tgtaaatgtg ttctgtccac caggggatgg cttctctggc    780
cctgcaccac gcaagtctaa actcatctgc gagggcagca acttcactcc aaaaccgac    840
acagtatcct ggctaaagga tgggaagctc gtggaatctg gcttcaccac agatccggtg    900
accatcgaga acaaaggatc cacaccccaa acctacaagg tcataagcac acttaccatc    960
tctgaaatcg actggctgaa cctgaatgtg tacacctgcc gtgtggatca caggggtctc    1020
accttcttga agaactgtgc ctccacatgt gctgccagtc cctccacaga catcctaacc    1080
ttcaccatcc cccctcctt tgccgacatc ttctcagca agtccgctaa cctgacctgt    1140
ctggtctcaa acctggcaac ctatgaaacc ctgaatatct cctgggcttc tcaaagtgg    1200
gaaccacttg aaacaaaaa taaatcatg gaaagccatc ccaatggcac cttcagtgt    1260
aaggggtgtg ctagtgtttg tgtggaagac tggaataaca ggaaggaatt tgtgtgtact    1320
gtgactcaca gggatctgcc ttaccacag aagaaattca tctcaaaacc caatgaggtg    1380
cacaacatc cacctgctgt gtacctgctg ccaccagctc gtgagcaact gaacctgagg    1440
gagtcagcca cagtcacctg cctgggtgaag ggcttctctc ctgcagacat cagtgtgcag    1500

```

-continued

```

tggttcaga gagggcaact cttgccccaa gagaagtatg tgaccagtgc cccgatgcca 1560
gagcctgggg ccccaggctt ctactttacc cacagcatcc tgactgtgac agaggaggaa 1620
tggaactccg gagagaccta tacctgtgtt gtaggccacg aggccttgcc acacctggtg 1680
accgagagga ccgtggacaa gtccactggg aaacccacac tgtacaatgt ctccctgac 1740
atgtctgaca caggcggcac ctgctattga ccat 1774

```

```

<210> SEQ ID NO 82
<211> LENGTH: 81
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 82

```

```

Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Ser Tyr Asp Ile
 1             5             10            15
Asn Trp Val Arg Gln Ala Thr Gly Gln Gly Leu Glu Trp Met Gly Trp
          20            25            30
Met Asn Pro Asn Ser Gly Asn Thr Gly Tyr Ala Gln Lys Phe Gln Gly
          35            40            45
Arg Val Thr Met Thr Arg Asn Thr Ser Ile Ser Thr Ala Tyr Met Glu
          50            55            60
Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Tyr Cys Ala Arg
          65            70            75            80
Gly

```

```

<210> SEQ ID NO 83
<211> LENGTH: 92
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 83

```

```

Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Ser Tyr Asp Ile
 1             5             10            15
Asn Trp Val Arg Gln Ala Thr Gly Gln Gly Leu Glu Trp Met Gly Trp
          20            25            30
Met Asn Pro Asn Ser Gly Asn Thr Gly Tyr Ala Gln Lys Phe Gln Gly
          35            40            45
Arg Val Thr Met Thr Arg Asn Thr Ser Ile Ser Thr Ala Tyr Met Glu
          50            55            60
Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Tyr Cys Ala Arg
          65            70            75            80
Glu Glu Trp Leu Val Arg Tyr Tyr Gly Met Asp Val
          85            90

```

```

<210> SEQ ID NO 84
<211> LENGTH: 96
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 84

```

```

Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Ser Tyr Asp Ile
 1             5             10            15
Asn Trp Val Arg Gln Ala Thr Gly Gln Gly Leu Glu Trp Met Gly Trp
          20            25            30

```

-continued

```

Met Asn Pro Asn Ser Gly Asn Thr Gly Tyr Ala Gln Lys Phe Gln Gly
    35                      40                      45

Arg Val Thr Met Asn Arg Asn Thr Ser Ile Ser Thr Ala Tyr Met Glu
    50                      55                      60

Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Tyr Cys Ala Arg
    65                      70                      75                      80

Gly Gly His Gly Gly Ser Tyr Phe Tyr Ser Tyr Tyr Gly Met Asp Val
    85                      90                      95

```

```

<210> SEQ ID NO 85
<211> LENGTH: 78
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 85

```

```

Ser Leu Thr Cys Ala Val Tyr Gly Gly Phe Ser Gly Tyr Tyr Trp Ser
  1              5              10              15

Trp Ile Arg Gln Pro Pro Gly Lys Gly Leu Glu Trp Ile Gly Glu Ile
    20              25              30

Asn His Ser Gly Ser Thr Asn Tyr Asn Pro Ser Leu Lys Ser Arg Val
    35              40              45

Thr Ile Ser Val Asp Thr Ser Lys Asn Gln Phe Ser Leu Lys Leu Ser
    50              55              60

Ser Val Thr Ala Ala Asp Thr Ala Val Tyr Tyr Cys Ala Arg
    65              70              75

```

```

<210> SEQ ID NO 86
<211> LENGTH: 92
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 86

```

```

Ser Leu Thr Cys Ala Val Tyr Gly Gly Phe Ser Gly Tyr Tyr Trp Ser
  1              5              10              15

Trp Ile Arg Gln Pro Pro Gly Lys Gly Leu Glu Trp Ile Gly Glu Ile
    20              25              30

Asn His Ser Gly Ser Thr Asn Tyr Asn Pro Ser Leu Lys Ser Arg Val
    35              40              45

Thr Ile Ser Val Asp Thr Ser Lys Asn Gln Phe Ser Leu Lys Leu Ser
    50              55              60

Ser Val Thr Ala Ala Asp Thr Ala Val Tyr Tyr Cys Ala Arg Gly Ala
    65              70              75              80

Ala Glu Tyr Tyr Tyr Tyr Tyr Tyr Gly Met Asp Val
    85              90

```

```

<210> SEQ ID NO 87
<211> LENGTH: 92
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 87

```

```

Ser Leu Thr Cys Ala Val Tyr Gly Gly Phe Ser Gly Tyr Tyr Trp Ser
  1              5              10              15

Trp Ile Arg Gln Pro Pro Gly Lys Gly Leu Glu Trp Ile Gly Glu Ile
    20              25              30

Asn His Ser Gly Ser Thr Asn Tyr Asn Pro Ser Leu Lys Ser Arg Val

```

-continued

35	40	45
Thr Ile Ser Val Asp Thr Ser Lys Asn Gln Phe Ser Leu Lys Leu Ser		
50	55	60
Ser Val Thr Ala Ala Asp Thr Ala Val Tyr Tyr Cys Ala Arg Gly Thr		
65	70	75 80
Thr Glu Tyr Tyr Tyr Tyr Tyr Tyr Gly Met Asp Val		
85	90	

<210> SEQ ID NO 88
 <211> LENGTH: 79
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 88

Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Gly Ile Arg Asn Asp		
1	5	10 15
Leu Gly Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Arg Leu Ile		
20	25	30
Tyr Ala Ala Ser Ser Leu Gln Ser Gly Val Pro Ser Arg Phe Ser Gly		
35	40	45
Ser Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro		
50	55	60
Glu Asp Phe Ala Thr Tyr Tyr Cys Leu Gln His Asn Ser Tyr Pro		
65	70	75

<210> SEQ ID NO 89
 <211> LENGTH: 79
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 89

Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Asp Ile Arg Asp Asn		
1	5	10 15
Leu Gly Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Arg Leu Ile		
20	25	30
Tyr Ala Ala Ser Asn Leu Gln Ser Gly Val Pro Ser Arg Phe Ser Gly		
35	40	45
Ser Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro		
50	55	60
Glu Asp Phe Ala Thr Tyr Tyr Cys Leu Gln Tyr Lys Thr Tyr Pro		
65	70	75

<210> SEQ ID NO 90
 <211> LENGTH: 79
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 90

Glu Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Gly Ile Arg Asp Glu		
1	5	10 15
Leu Gly Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Arg Leu Ile		
20	25	30
Tyr Val Ala Ser Ser Leu Gln Ser Gly Val Pro Ser Arg Phe Ser Gly		
35	40	45
Ser Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro		
50	55	60

-continued

Glu Asp Phe Ala Thr Tyr Tyr Cys Leu Gln His Asn Gly Tyr Pro
65 70 75

<210> SEQ ID NO 91
<211> LENGTH: 85
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 91

Glu Arg Ala Thr Ile Asn Cys Lys Ser Ser Gln Ser Val Leu Tyr Ser
1 5 10 15
Ser Asn Asn Lys Asn Tyr Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln
20 25 30
Pro Pro Lys Leu Leu Ile Tyr Trp Ala Ser Thr Arg Glu Ser Gly Val
35 40 45
Pro Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr
50 55 60
Ile Ser Ser Leu Gln Ala Glu Asp Val Ala Val Tyr Tyr Cys Gln Gln
65 70 75 80
Tyr Tyr Ser Thr Pro
85

<210> SEQ ID NO 92
<211> LENGTH: 86
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 92

Glu Arg Ala Thr Ile Asn Cys Lys Ser Ser Gln Ser Val Leu Tyr Ser
1 5 10 15
Phe Asn Asn Lys Asn Tyr Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln
20 25 30
Pro Pro Lys Leu Leu Ile Tyr Trp Ala Ser Thr Arg Glu Ser Gly Val
35 40 45
Pro Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr
50 55 60
Ile Ser Ser Leu Gln Ala Glu Asp Val Ala Val Tyr Tyr Cys Gln Gln
65 70 75 80
Tyr Tyr Ser Thr Arg Thr
85

<210> SEQ ID NO 93
<211> LENGTH: 86
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 93

Glu Arg Ala Thr Ile Asn Cys Lys Ser Ser Gln Ser Val Leu Tyr Ser
1 5 10 15
Phe Asn Asn Lys Asn Tyr Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln
20 25 30
Pro Pro Lys Leu Leu Ile Tyr Trp Ala Ser Thr Arg Glu Ser Gly Val
35 40 45
Pro Asp Arg Phe Gly Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr
50 55 60

-continued

```
Ile Ser Ser Leu Gln Ala Glu Asp Val Ala Val Tyr Tyr Cys Gln Gln
65              70              75              80
```

```
Tyr Tyr Ser Thr Arg Thr
85
```

```
<210> SEQ ID NO 94
<211> LENGTH: 84
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
```

```
<400> SEQUENCE: 94
```

```
Glu Pro Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Leu Leu His Ser
1              5              10              15
```

```
Asn Gly Tyr Asn Tyr Leu Asp Trp Tyr Leu Gln Lys Pro Gly Gln Ser
20              25              30
```

```
Pro Gln Leu Leu Ile Tyr Leu Gly Ser Asn Arg Ala Ser Gly Val Pro
35              40              45
```

```
Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile
50              55              60
```

```
Ser Arg Val Glu Ala Glu Asp Val Gly Val Tyr Tyr Cys Met Gln Ala
65              70              75              80
```

```
Leu Gln Thr Pro
```

```
<210> SEQ ID NO 95
<211> LENGTH: 86
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
```

```
<400> SEQUENCE: 95
```

```
Glu Pro Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Leu Leu His Ser
1              5              10              15
```

```
Asn Gly Tyr Asn Tyr Leu Asp Trp Tyr Leu Gln Lys Pro Gly Gln Ser
20              25              30
```

```
Pro Gln Leu Leu Ile Tyr Leu Gly Ser Asn Arg Ala Ser Gly Val Pro
35              40              45
```

```
Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile
50              55              60
```

```
Ser Arg Val Glu Ala Glu Asp Val Gly Ile Tyr Tyr Cys Met Gln Thr
65              70              75              80
```

```
Arg Gln Thr Pro Arg Thr
85
```

```
<210> SEQ ID NO 96
<211> LENGTH: 86
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
```

```
<400> SEQUENCE: 96
```

```
Glu Pro Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Leu Leu His Ser
1              5              10              15
```

```
Asn Gly Tyr Asn Tyr Leu Asp Trp Tyr Leu Gln Lys Pro Gly Gln Ser
20              25              30
```

```
Pro Gln Leu Leu Ile Tyr Leu Gly Ser Asn Arg Ala Ser Gly Val Pro
35              40              45
```

```
Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile
50              55              60
```

-continued

Ser Arg Val Glu Ala Glu Asp Val Gly Ile Tyr Tyr Cys Met Gln Ser
65 70 75 80

Leu Gln Ile Pro Arg Leu
85

<210> SEQ ID NO 97

<211> LENGTH: 80

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 97

Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Gly Ile Ser Asn Tyr
1 5 10 15

Leu Ala Trp Phe Gln Gln Lys Pro Gly Lys Ala Pro Lys Ser Leu Ile
20 25 30

Tyr Ala Ala Ser Ser Leu Gln Ser Gly Val Pro Ser Lys Phe Ser Gly
35 40 45

Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro
50 55 60

Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln Tyr Asn Ser Tyr Pro Pro
65 70 75 80

<210> SEQ ID NO 98

<211> LENGTH: 83

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 98

Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Asp Ile Ser Ile Tyr
1 5 10 15

Leu Ala Trp Phe Gln Gln Arg Pro Gly Lys Ala Pro Lys Ser Leu Ile
20 25 30

Tyr Ala Ala Ser Ser Leu Gln Ser Gly Val Pro Ser Lys Phe Ser Gly
35 40 45

Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro
50 55 60

Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln Tyr Asn Ser Tyr Phe Thr
65 70 75 80

Phe Gly Pro

<210> SEQ ID NO 99

<211> LENGTH: 83

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 99

Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Gly Ile Ser Ile Tyr
1 5 10 15

Leu Ala Trp Phe Gln Gln Arg Pro Gly Lys Ala Pro Lys Ser Leu Ile
20 25 30

Tyr Ala Ala Ser Ser Leu Gln Ser Gly Val Pro Ser Lys Phe Ser Gly
35 40 45

Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro
50 55 60

Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln Tyr Asn Ser Tyr Phe Thr

-continued

65	70	75	80
Phe Gly Pro			
<210> SEQ ID NO 100			
<211> LENGTH: 18			
<212> TYPE: DNA			
<213> ORGANISM: Homo sapiens			
<220> FEATURE:			
<221> NAME/KEY: CDS			
<222> LOCATION: (1)..(18)			
<400> SEQUENCE: 100			
gtg cta gct ttc cag gag			18
Val Leu Ala Phe Gln Glu			
1	5		
<210> SEQ ID NO 101			
<211> LENGTH: 6			
<212> TYPE: PRT			
<213> ORGANISM: Homo sapiens			
<400> SEQUENCE: 101			
Val Leu Ala Phe Gln Glu			
1	5		
<210> SEQ ID NO 102			
<211> LENGTH: 21			
<212> TYPE: DNA			
<213> ORGANISM: Homo sapiens			
<220> FEATURE:			
<221> NAME/KEY: CDS			
<222> LOCATION: (1)..(21)			
<400> SEQUENCE: 102			
cgc gtg cta gct ttc cag gag			21
Arg Val Leu Ala Phe Gln Glu			
1	5		
<210> SEQ ID NO 103			
<211> LENGTH: 7			
<212> TYPE: PRT			
<213> ORGANISM: Homo sapiens			
<400> SEQUENCE: 103			
Arg Val Leu Ala Phe Gln Glu			
1	5		
<210> SEQ ID NO 104			
<211> LENGTH: 21			
<212> TYPE: DNA			
<213> ORGANISM: Homo sapiens			
<220> FEATURE:			
<221> NAME/KEY: CDS			
<222> LOCATION: (4)..(21)			
<400> SEQUENCE: 104			
cgt gtg cta gct ttc cag gag			21
Val Leu Ala Phe Gln Glu			
1	5		
<210> SEQ ID NO 105			
<211> LENGTH: 6			
<212> TYPE: PRT			
<213> ORGANISM: Homo sapiens			

-continued

<400> SEQUENCE: 105

Val Leu Ala Phe Gln Glu
1 5

1. An isolated human monoclonal antibody having an isotype that fixes complement and a variable region that binds to the epitope on CD147 bound by the IgM monoclonal antibody ABX-CBL, with the proviso that the antibody is not CBL1 (ATCC HB 8214).

2. The antibody of claim 1, wherein the antibody in the presence of complement acts to selectively kill cells selected from the group consisting of activated T-cells, activated B-cells, and monocytes but is substantially non-toxic to resting T-cells and resting B-cells.

3. canceled

4. The antibody of claim 1, wherein the isotype is selected from the group consisting of murine IgM, murine IgG2a, murine IgG2b, murine IgG3, human IgM, human IgG1, and human IgG3.

5. canceled

6. The antibody of claim 2, wherein the isotype is selected from the group consisting of murine IgM, murine IgG2a, murine IgG2b, murine IgG3, human IgM, human IgG1, and human IgG3.

7. An isolated human monoclonal antibody having an isotype that fixes complement and a variable region that binds to CD147 on populations of activated T-cells, activated B-cells, and resting or activated monocytes, that, in the presence of complement, selectively depletes such populations through complement mediated killing while being substantially nontoxic to other cells, with the proviso that the antibody is not CBL1 (ATCC HB 8214).

8. canceled

9. The antibody of claim 7, wherein the isotype is selected from the group consisting of murine IgM, murine IgG2a, murine IgG2b, murine IgG3, human IgM, human IgG1, and human IgG3.

10. An isolated human monoclonal antibody having the following characteristics:

- (a) binds to CD147;
- (b) shows a binding against CEM cell lysates on Western blot similar to that provided in **FIG. 1**;
- (c) an isotype selected from the group consisting of murine IgM, murine IgG2a, murine IgG2b, murine IgG3, human IgM, human IgG1, and human IgG3;
- (d) competes with ABX-CBL for binding to CD147;
- (e) cross reacts with hn-RNP-k protein;
- (f) binds to a consensus sequence on CD147 comprising RVRS (SEQ ID NO:106);
- (g) selectively kills activated T-cells, activated B-cells, and monocytes in an MLR assay only in the presence of complement; and
- (h) is substantially non-toxic to cells expressing CD55 and CD59, with and without the presence of complement,

with the proviso that the antibody is not CBL1 (ATCC HB 8214).

11. A method to select an anti-CD147 antibodies for the treatment of disease, comprising:

generating antibodies that bind to CD147 and that are capable of binding complement;

assaying the antibodies for one or more of the following properties:

- (a) competition with ABX-CBL for binding to CD147;
- (b) capability to selectively kill activated T-cells, activated B-cells, and monocytes in a MLR assay only in the presence of complement; and
- (c) being substantially non-toxic to cells expressing CD55 and CD59, with and without the presence of complement,

with the proviso that the antibody is not CBL1.

12. The method of claim 11, further comprising the following property:

- (d) binding to CEM cell lysates on Western blot in a manner similar to that provided in **FIG. 1**.

13. The method of claim 11, further comprising the following property:

- (e) binding to a consensus sequence in a peptide of RXRS (SEQ ID NO:11).

14. The method of claim 11, further comprising the following property:

- (f) cross reacts with hn-RNP-k protein.

15. The method of claim 11, further comprising the following property:

- (g) binding to a form of CD147 expressed by COS cells and *E. coli* cells.

16. A method to treat disease, comprising providing an antibody that has an isotype that fixes complement and a variable region that binds to CD147 on populations of activated T-cells, activated B-cells, and resting or activated monocytes, that, in the presence of complement, selectively depletes such populations through complement mediated killing while being substantially nontoxic to other cells, with the proviso that the antibody is not CBL1.

17. The method of claim 16, wherein the antibody is a human antibody.

18. The method of claim 16, wherein the isotype is selected from the group consisting of murine IgM, murine IgG2a, murine IgG2b, murine IgG3, human IgM, human IgG1, and human IgG3.

19. A method to treat GVHD, comprising providing an antibody that has an isotype that fixes complement and a variable region that binds to CD147 on populations of activated T-cells, activated B-cells, and resting or activated monocytes, that, in the presence of complement, selectively

depletes such populations through complement mediated killing while being substantially nontoxic to other cells, with the proviso that the antibody is not CBL1.

20. The method of claim 19, wherein the antibody is a human antibody.

21. The method of claim 19, wherein the isotype is selected from the group consisting of murine IgM, murine IgG2a, murine IgG2b, murine IgG3, human IgM, human IgG1, and human IgG3.

22. A human monoclonal antibody that binds to an epitope on CD147 comprising the consensus sequence RVRSH (SEQ ID NO:107), wherein the antibody is not CBL1 (ATCC HB 8214).

23. canceled

24. An isolated peptide comprising the sequence selected from the group consisting of RXRS (SEQ ID NO:11, RXRSH (SEQ ID NO:13, RVRSH (SEQ ID NO:106), and RVRSH (SEQ ID NO:107).

25. A method of producing an antibody that binds to the peptide of claim 24 comprising the steps of immunizing an animal with the peptide and collecting the antibody.

26. A human monoclonal antibody that binds to CD147.

27. A kit for the treatment of diseases having an etiology characterized by a harmful presence of activated T cells, B cells, or monocytes, comprising:

(a) a liquid preparation comprising an amount of an anti-CD147 antibody in a pharmaceutically acceptable carrier and

(b) instructions on administering said preparation to a patient suffering from a disease having the etiology characterized by a harmful presence of activated T cells, B cells, or monocytes to provide a dosage in the range of from about 0.1 mg/kg to about 0.3 mg/kg of the antibody.

28. The kit of claim 27, wherein the antibody comprises ABX-CBL.

29. The kit of claim 27, wherein the instructions further include instructions for the administration of the antibody in a series of administrations to provide a dosage in the range of from about 0.1 mg/kg to about 0.3 mg/kg of the antibody in each administration.

30. The kit of claim 27, wherein the disease comprises GVHD.

31. An article of manufacture for use in the treatment of diseases having an etiology characterized by a harmful presence of activated T cells, B cells, or monocytes, comprising:

(a) a sterile vial;

(b) an anti-CD147 monoclonal antibody in a pharmaceutically acceptable carrier contained within the vial; and

(c) instructions for administration of the antibody to a patient suffering from such a disease in a manner to provide a dosage in the range of from about 0.1 mg/kg to about 0.3 mg/kg of the antibody in each administration.

32. The article of claim 31, wherein the antibody comprises ABX-CBL.

33. The article of claim 31, wherein the instructions further include instructions for the administration of the antibody in a series of administrations to provide a dosage in the range of from about 0.1 mg/kg to about 0.3 mg/kg of the antibody in each administration.

34. The kit of claim 31, wherein the disease comprises GVHD.

35. A kit for the treatment of diseases having an etiology characterized by a harmful presence of activated T cells, B cells, or monocytes, comprising:

(a) a liquid preparation comprising an amount of an anti-CD147 antibody designated ABX-CBL in a pharmaceutically acceptable carrier and

(b) instructions on administering said preparation to a patient suffering from a disease having the etiology characterized by a harmful presence of activated T cells, B cells, or monocytes in a series of administrations to provide a dosage in the range of from about 0.1 mg/kg to about 0.3 mg/kg of the antibody in each administration.

36. The kit of claim 35, wherein the instructions further include instructions for the administration of the antibody in a series of administrations to provide a dosage in the range of from about 0.1 mg/kg to about 0.3 mg/kg of the antibody in each administration.

37. The kit of claim 35, wherein the disease comprises GVHD.

38. An article of manufacture for use in the treatment of diseases having an etiology characterized by a harmful presence of activated T cells, B cells, or monocytes, comprising:

(a) a sterile vial;

(b) an anti-CD147 monoclonal antibody designated ABX-CBL in a pharmaceutically acceptable carrier contained within the vial; and

(c) instructions for administration of the antibody to a patient suffering from such a disease in a manner to provide a dosage in the range of from about 0.1 mg/kg to about 0.3 mg/kg of the antibody in each administration.

39. The article of claim 38, wherein the instructions further include instructions for the administration of the antibody in a series of administrations to provide a dosage in the range of from about 0.1 mg/kg to about 0.3 mg/kg of the antibody in each administration.

40. The article of claim 38, wherein the disease comprises GVHD.

41. A kit for the treatment GVHD, comprising:

(a) a liquid preparation comprising an amount of an anti-CD147 antibody in a pharmaceutically acceptable carrier and

(b) instructions on administering said preparation to a patient suffering from GVHD in a series of administrations to provide a dosage in the range of from about 0.1 mg/kg to about 0.3 mg/kg of the antibody in each administration.

42. The kit of claim 41, wherein the antibody comprises ABX-CBL.

43. The kit of claim 41, wherein the instructions further include instructions for the administration of the antibody in a series of administrations to provide a dosage in the range of from about 0.1 mg/kg to about 0.3 mg/kg of the antibody in each administration.

44. An article of manufacture for use in the treatment of GVHD, comprising:

- (a) a sterile vial;
- (b) an anti-CD147 monoclonal antibody in a pharmaceutically acceptable carrier contained within the vial; and
- (c) instructions for administration of the antibody to a patient suffering from GVHD in a manner to provide a dosage in the range of from about 0.1 mg/kg to about 0.3 mg/kg of the antibody in each administration.

45. The article of claim 44, wherein the antibody comprises ABX-CBL.

46. The article of claim 44, wherein the instructions further include instructions for the administration of the antibody in a series of administrations to provide a dosage in the range of from about 0.1 mg/kg to about 0.3 mg/kg of the antibody in each administration.

47. A kit for the treatment of GVHD, comprising:

- (a) a liquid preparation comprising an amount of an anti-CD147 antibody designated ABX-CBL in a pharmaceutically acceptable carrier and
- (b) instructions on administering said preparation to a patient suffering from GVHD in a series of administrations to provide a dosage in the range of from about 0.1 mg/kg to about 0.3 mg/kg of the antibody in each administration.

48. The kit of claim 47, wherein the instructions further include instructions for the administration of the antibody in a series of administrations to provide a dosage in the range of from about 0.1 mg/kg to about 0.3 mg/kg of the antibody in each administration.

49. An article of manufacture for use in the treatment of GVHD, comprising:

- (a) a sterile vial;
- (b) an anti-CD147 monoclonal antibody designated ABX-CBL in a pharmaceutically acceptable carrier contained within the vial; and
- (c) instructions for administration of the antibody to a patient suffering from GVHD in a manner to provide a dosage in the range of from about 0.1 mg/kg to about 0.3 mg/kg of the antibody in each administration.

50. The article of claim 49, wherein the instructions further include instructions for the administration of the antibody in a series of administrations to provide a dosage in the range of from about 0.1 mg/kg to about 0.3 mg/kg of the antibody in each administration.

51. A pharmaceutical composition, comprising an anti-CD147 monoclonal antibody designated ABX-CBL in a pharmaceutically acceptable diluent, buffer, or excipient.

52. The pharmaceutical composition of claim 51, wherein the antibody is provided in a dosage from about 0.1 mg/kg and about 0.2 mg/kg.

53. A method for the treatment of diseases having an etiology characterized by a harmful presence of activated T cells, B cells, or monocytes, comprising administering a liquid preparation comprising an amount of an anti-CD147 antibody in a pharmaceutically acceptable carrier to a patient suffering from such disease.

54. The method of claim 53, wherein the antibody comprises ABX-CBL.

55. The method of claim 53, wherein the administration is conducted to provide a dosage in the range of from about 0.1 mg/kg to about 0.3 mg/kg of the antibody in each administration.

56. The method of claim 53, wherein the disease comprises GVHD.

57. A method for the treatment of GVHD comprising administering a liquid preparation comprising an amount of an anti-CD147 antibody in a pharmaceutically acceptable carrier to a patient suffering from GVHD.

58. The method of claim 57, wherein the antibody comprises ABX-CBL.

59. The method of claim 57, wherein the administration is conducted to provide a dosage in the range of from about 0.1 mg/kg to about 0.3 mg/kg of the antibody in each administration.

60. The human antibody according to claim 26, wherein the heavy chain has an amino acid sequence selected from the group consisting of: SEQ ID NO:23; SEQ ID NO:25; SEQ ID NO:27; SEQ ID NO:29; SEQ ID NO:31; SEQ ID NO:33; SEQ ID NO:35; SEQ ID NO:37; SEQ ID NO:39; SEQ ID NO:40.

61. The human antibody according to claim 26, wherein the light chain has an amino acid sequence selected from the group consisting of: SEQ ID NO:24; SEQ ID NO:26; SEQ ID NO:28; SEQ ID NO:30; SEQ ID NO:32; SEQ ID NO:34; SEQ ID NO:36; SEQ ID NO:38; and SEQ ID NO:41.

* * * * *

专利名称(译)	CD147结合分子作为治疗剂		
公开(公告)号	US20060104974A1	公开(公告)日	2006-05-18
申请号	US09/784950	申请日	2001-02-15
[标]申请(专利权)人(译)	DAVIS C G 布拉谢罗素W 科尔巴兰JOSEř CULWELL ALAN - [R 绿色LARRY大号 HALES JOANNE 豪夫里洛NANCY 伊万诺夫VLADIMIRê LIPANI约翰 刘强 WEBER RICHARD°F 杨肖冬		
申请(专利权)人(译)	DAVIS C G 布拉谢罗素W 科尔巴兰JOSEř CULWELL ALAN - [R 绿色LARRY大号 HALES JOANNE 豪夫里洛NANCY 伊万诺夫VLADIMIRê LIPANI约翰 LIU QIANG WEBER RICHARD°F 杨晓东		
当前申请(专利权)人(译)	DAVIS C G 布拉谢罗素W 科尔巴兰JOSEř CULWELL ALAN - [R 绿色LARRY大号 HALES JOANNE 豪夫里洛NANCY 伊万诺夫VLADIMIRê LIPANI约翰 LIU QIANG WEBER RICHARD°F 杨晓东		
[标]发明人	DAVIS C GEOFFREY BLACHER RUSSELL W CORVALAN JOSE R CULWELL ALAN R GREEN LARRY L HALES JOANNE HAVRILLA NANCY IVANOV VLADIMIR E LIPANI JOHN A LIU QIANG		

	WEBER RICHARD F YANG XIAO DONG
发明人	DAVIS, C. GEOFFREY BLACHER, RUSSELL W. CORVALAN, JOSE R. CULWELL, ALAN R. GREEN, LARRY L. HALES, JOANNE HAVRILLA, NANCY IVANOV, VLADIMIR E. LIPANI, JOHN A. LIU, QIANG WEBER, RICHARD F. YANG, XIAO-DONG
IPC分类号	A61K39/395 C07K16/28 G01N33/53 A61P19/02 A61P29/00 A61P35/00 A61P35/02 A61P37/02 A61P37/06 A61P37/08 C07K14/705 C12N15/09 C12P21/08 G01N33/577
CPC分类号	A61K2039/505 C07K14/70596 C07K16/2803 C07K16/2896 C07K2317/21 C07K2317/732 C07K2317/734 C07K2317/92 C07K2319/00 C07K2319/30 C07K2317/34 A61P19/02 A61P29/00
优先权	PCT/US1999/004583 1999-03-03 WO 09/244253 1999-02-03 US 09/034607 1998-03-03 US
外部链接	Espacenet USPTO

根据本发明，我们发现某些细胞如T细胞，B细胞和/或单核细胞上表达分子CD147可用于治疗多种疾病。特别地，我们已经证明，与CD147结合并导致例如通过补体结合杀死这些细胞的抗体在疾病的治疗中是有效的。这种治疗似乎有效的疾病包括但不限于：移植物抗宿主病（GVHD），器官移植排斥疾病（包括但不限于肾移植，眼部移植等），癌症（包括但不限于癌症）。血液（即白血病和淋巴瘤），胰腺和其他），自身免疫疾病，炎症性疾病等。

[illegible]