



US 20020127568A1

(19) **United States**
 (12) **Patent Application Publication** (10) **Pub. No.: US 2002/0127568 A1**
Meyers (43) **Pub. Date: Sep. 12, 2002**

(54) **47324, A NOVEL HUMAN G-PROTEIN AND USES THEREFOR**

Publication Classification

(51) **Int. Cl.⁷** **C12Q 1/68**; G01N 33/53;
 C07H 21/04; C12P 21/02;
 C12N 5/06; C07K 14/705;
 C07K 16/28
 (52) **U.S. Cl.** **435/6**; 435/7.1; 435/69.1;
 435/325; 435/320.1; 530/350;
 530/388.22; 536/23.5

(76) Inventor: **Rachel Meyers**, Newton, MA (US)

Correspondence Address:
Carolyn A. Favorito
Morrison & Foerster LLP
Suite 500
3811 Valley Centre Drive
San Diego, CA 92130-2332 (US)

(57) **ABSTRACT**
 The invention provides isolated nucleic acids molecules, designated 47324 nucleic acid molecules, which encode novel G-protein family members, preferably Ras family members. The invention also provides antisense nucleic acid molecules, recombinant expression vectors containing 47324 nucleic acid molecules, host cells into which the expression vectors have been introduced, and nonhuman transgenic animals in which a 47324 gene has been introduced or disrupted. The invention still further provides isolated 47324 proteins, fusion proteins, antigenic peptides and anti-47324 antibodies. Diagnostic methods utilizing compositions of the invention are also provided.

(21) Appl. No.: **09/945,173**

(22) Filed: **Aug. 31, 2001**

Related U.S. Application Data

(60) Provisional application No. 60/229,293, filed on Sep. 1, 2000.

M A S L D R V K V L V L G D S

AGGGNAGGCAGTGGCAAG ATG GCG TCC CTG GAT CCG GTG AAG GTA CTG GTG TTG GGA GAC TCA SEQ ID NO:2
 ↑SEQ ID NO:3→ SEQ ID NO:1

G V G K S S L V H L L C Q N Q V L G N P 35
 GGT GTT GGG AAA TCT TCG TTA GTC CAT CTC CTA TGC CAA AAT CAA GTG CTG GGA AAT CCA 105

S W T V G C S V D V R V H D Y K E G T P 55
 TCA TGG ACT GTG GGC TGC TCA GTG GAT GTC AGA GTT CAT GAT TAC AAA GAA GGA ACC CCA 165

E E K T Y Y I E L W D V G G S V G S A S 75
 GAA GAG AAG ACC TAC TAC ACA GAA TTA TGG GAT GTT GGA GGC TCT GTG GGC AGT GCC AGC 225

S V K S T R A V F Y N S V N G I I F V H 95
 AGC GTG AAA AGC ACA AGA GCA GTA TTC TAC AAC TCC GTA AAT GGT ATT ATT TTC GTA CAC 285

D L T N K K S S Q N L R R W S L E A L N 115
 GAC TTA ACA AAT AAG AAG TCC TCC CAA AAC TTG CGT CGT TGG TCA TTG GAA GCT CTC AAC 345

R D L V P T G V L V T N G D Y D Q E Q F 135
 AGG GAT TTG GTG CCA ACT GGA GTC TTG GTG ACA AAT GGG GAT TAT GAT CAA GAA CAG TTT 405

A D N Q I P L L V I G T K L D Q I H E T 155
 GCT GAT AAC CAA ATA CCA CTG TTG GTA ATA GGG ACT AAA CTG GAC CAG ATT CAT GAA ACA 465

K R H E V L T R T A F L A E D F N P E E 175
 AAG CGC CAT GAA GTT TTA ACT AGG ACT GCT TTC CTG GCT GAG GAT TTC AAT CCA GAA GAA 525

I N L D C T N P R Y L A A G S S N A V K 195
 ATT AAT TTG GAC TGC ACA AAT CCA CGG TAC TTA GCT GCA GGT TCT TCC AAT GCT GTC AAG 585

L S R F F D K V I E K R Y F L R E G N Q 215
 CTC AGT AGG TTT TTT GAT AAG GTC ATA GAG AAG AGA TAC TTT TTA AGA GAA GGT AAT CAG 645

I P G F P D R K R F G A G T L K S L H Y 235
 ATT CCA GGC TTT CCT GAT CCG AAA AGA TTT GGG GCA GGA ACA TTA AAG AGC CTT CAT TAT 705

D * 237
 GAC TGA 711
 ←SEQ ID NO:3↑

ATTACATCATCCTTTTGAAGAGTGAGCAAGCAGTGGCAGTCTTTTCACAGCTCATCTTGCTGTGTTC AATTATTACCAT
 CACAGCCTTTTAAACAAAATCATCTTAAATGCTACCCTTCAGCCTTACCCTTTAATGGAAAAATGAAAGGAAGTGACAA
 TACGGGAGGTCCAAACTTTGTCCCTGTTCTCTGTGTCTTACCTTTCTGTCCCTGTGATAGATTATGTAAGGCCTT
 GTGTAATATGAGATGTTGTCAAAATGATGCAGTAAATGAGCAATGACAGTGTACTGCAGAGAAAATTTACTCTTGCCT
 AGAACTGGAGGGTTTTTATGGGTCTGTAATTTTCCCACTCATTGCTGAAAGCTTAATTAAGTACTTCAAAAACGTAT
 CTCCATTGTTTTACCTTCTTGAGGGGAACGGTCTTGTAAACAGCCCTGAGTTGTCTACCCCAACAACTCTCTGTCAAT
 TTCAAAGATGCAAAATGGTGTATTTAATTGTCTCCACCATTGTACACACAGGAATGCCTAATAATAGCARCCCTTGT
 CTCCTCTCTCTCCTTTGCAAAATGGCTCAGTGACTGGAAGAGCCGACTAATAGCCAGAGTTAATATAAATACAAAT

FIG. 1A

TAATAATACATAGAGAACAGCAATACCAGAAAAAGAATTCTGGTAAAATGATGTGAAAAATTGACAGCTCCCTCACT
CTTAAGGTTGCTGCTATATACAGTCTAGGTTTTCTGTTGGAAATAGGTAGGGTAAAATCTAAGACCTGCACAAGGGCA
GTGAGAGACATTTACAGCCTCCTCTCTATTTGTTTTTTAAGGAAAAGTCAACTCCTGAAATGTCCCTTAGCTATAATC
AGAAACTAAGAATATTATTCTGTGTCAACAATGTATTTATGGAGAGAAGTAAAAATAAGTTCCACAGCAACACAAAA
CATGAATTATTGAACTA

FIG. 1B




FIG. 2

```

*->GlgflkiflklglwnKEMRILiLGLDnAGKTTILykLklge.ivtt SEQ ID NO:4
      ++l +   +++L+LG + GK+ +++ L +++ + +
47324  2  -----ASLDR----VKVLVLGDSGVGKSSLVHLLCQNQvLGNP 35

      iPTiGFNVetveykN.....ikFtvWDvGGQ.....eslRPlW
      + T+G v v + + ++++++++ + WDvGG ++ ++ +s R +
47324  36 SWTVGCSV-DVRVHDykegtpeektYYIELWDVGGsvgsassvKSTRAV- 83

      rnYfpnTdavIfvVDSaDreRieeakeELhaLLneeeL.....
      + +++++IfV D ++ + + L + e +++ +++ ++
47324  84 --FYNSVNGIIFVHDLTNKK----SSQLRRWSLEALNrdlvptgvlvtn 127

      .....adAplLifANKQDlpgAmseaEi...reaLgLhelkgskg
      ++ +++++ + plL+ + K D + +E+ +++ L +
47324  128 gdydqeqfadNQIPLLVIgTKLDQIHETKRHEVltrTAFLAEDFNP-E-- 174

      kVtl.e.gdRpWeiiggcsAvkGeGLyEGldWLSnnikkr<-*
      ++ l+ ++ R + ++Avk l + + +++++
47324  175 EINLdCtNPRYLAAGSSNAVK-----LSRFFDKVIEK 206
    
```

FIG. 3

```

*->KlVliGDSGVGKSsLliRfTdnkFveeYipTIGvDFyktvevD... SEQ ID NO:5
      K++++GDSGVGKSsL++ +n+ T+G+ ++ ++ +++
47324  8  KVLVLGDSGVGKSSLVHLLCQNQVVLGNPSWTVGCSVDVRVHDYKegt 54

      ..GktvkLqIWDTAGQERFralrpa.....YYRgAqGfLLVYDITs
      ++ kt+ ++WD G ++ +a++ +++ +Y +G++ V D T+
47324  55 peEKTYIELWDVGG----SVGSAssvkstravFYNSVNGIIFVHDLTN 99

      rdSFenvkkWleeilr..had.....kdenvpivLVGNKc
      ++S +n+++W e l+++ +++ +++++ +++++ d+ +p + +G+K
47324  100 KKSSQNLRRWSLEALNrdLVptgvlvtnngdydqeqfADNQIPLLVIgTKL 149

      DLeddedlelte.gqkRvVsteeGealAkelgA.....lpFmETSA
      D + e++ +V t ++ lA++++ ++ + + ++ +++ +
47324  150 DQIH-----EtKR-HEVLTRTAF-LAEDFNPeeinldctnPRYLAAGS 190

      KtntNVeeaFeelareilkkvsevnvnlDqpakkkkSkCcil<-*
      +++ F+ +++++ + +++++++ + +k+ +
47324  191 SNAVKLSRFFDKVIEKRYFLREGNQIPGFPPD-RKRFGAGTLK 231
    
```

FIG. 4

Query: 35 PSWTV-GCSVDVRVHDYKEGTPEEKT--YYIELWDXXXXXXXXXXXXXTRAVFYNSVNGI 91
 P+WT S D + Y TP Y++E +D R FY +++GI
 Sbjct: 69 PTWTTSPSSSESENYPYMRSTPTTTNLLYFVEFYDLNSDWRMCRQQ---RESFYKNIDGI 125 SEQ ID
 NO:6

Query: 92 IFVHDLTNKSSQNLRWSLEALNRDLVPTGVLVTVNGDYDQEQFADNQIPLLVIKTKLDQ 151
 + V+++ S +L W + L + + + + P+LV+GT LD+
 Sbjct: 126 VLVYNMLELSSQDSLHDWLYDPLRQICKHRHLRI-----RSILKNHNAPILVVGTVNLDK 179

Query: 152 IHETKRHEVLTRTAFLAEDFNPEEINLDCTNPRYLAAGSSNAVKLSRFFDKVIE-KRYF 209
 + L R +A N EE+ ++C +P+ S N KL F ++VIE K F
 Sbjct: 180 L----MRRPLRRRGSIAHQNLNVEMLVNCCLDPQSFVDKSRNQKLYGFLNRVIEFKEQF 234

FIG. 5a

Query: 7 VKVLVLGDSGVGKSSLVHLLCQNVLGNP-SWTVG---CSVDVRVHDYKE 52
 V++L+LGD GVGK+SL +L+ ++ P S TVG V VR+H+Y +
 Sbjct: 12 VRILMLGDRGVGKTSLTNLMATTEITPTPDSRTVGEESWHVQVRLHEYSK 61 SEQ ID NO:7

FIG. 5b

Query: 121 TGVLVTVNGD--YDQEQFAD-NQIPLLVIKTKLDQIHETKRHEVL--TRTAFLAEDFNPEE 175
 T L T+G +D E+F Q P+LV+GTKLD + E KRH + + +A+ EE
 Sbjct: 19 TDNLCTDGHILPDMEEFLGATQTPILVMGTKLDLLE-KRHPKMGVKKPGGIADKCGAEE 77 SEQ ID
 NO:8

Query: 176 INLDCTNPRYLAAGSSNAVKLSRFFDKVIEKRYFLR-----EGNQIPGFPDRKRFGAG 228
 I L+C N R LAAG+++AVKLSRFFD+VIE R LR + PDR+RFG
 Sbjct: 78 IWLNCRNSRSLAAGTTDAVKLSRFFDRVIENRKALRAALAFGVSSNAVSPDRRRFGPT 137

Query: 229 TLK 231
 + K
 Sbjct: 138 SAK 140

FIG. 6

Query: 6 RVKVLVLGDSGVGKSSLVHLLCQNQVVLGNPSWTVGCSVDVVRVHDYKEGTPEEKTYTIE 63
 +V+VLV+GDSGVGK+SLVHL+ + + P T+GC+V V+ Y G+P + I+
 Sbjct: 80 QVRVLVVGDSGVGKTSLVHLINKGSSIVRPPQTIGCTVGVKHITY--GSPASSSSSIQ 135 SEQ ID NO:9

FIG. 7

Query: 8 KVLVLGDSGVGKSSLVHLL--CQNQ-----VLGNPSWTVGCSVDVVRVHDYKEGT 54
 K+++GDSGVGK+SL++ L +N V+ + T+G +V+ T
 Sbjct: 7 KIVIIIGDSGVGKTSLLNKLRFTEENSFTEEDPTTRTVVDSYKSTIGVDFNVKTTIEVVDT 66 SEQ ID
 NO:10

Query: 55 PEEKTYTIELWDXXXXXXXXXXXXXXXXXTRAVFYNSVNGIIFVHDLTNKKSS-QNLRRSLEA 113
 + K +++WD A +Y I V+D+ + +SS +N +W E
 Sbjct: 67 -DGKNIKLIWDTAGQERYRSMLTSM EA-YYRGAEADIIVYDVDSSESSFENQTKWLKEI 124

Query: 114 LNRDLVPTGVLVTNGDYDQEQFADNQPILLVIGTKLD-QI---HETKRHEVLTRTAFLAE 169
 L +N E+ ++N +P++++G K D ++ E + + T A+
 Sbjct: 125 LRH-----ASN-----EEASEN-VPIILVGNKADLEVPNPEEVVEEKEEASTEEBAQ 170

Query: 170 DFNPEEINLDCTNPRYLAAGSSNAVKLSRFFDKVIEKRYFLREGNQIPGFPDRKRF 225
 F EE L P + + + F +++ R L++ +I D++++
 Sbjct: 171 SF-AEEKGLGVV-PFIETSAKTTGTNVVEEVFQELV--REILKKKKEIQEKADQEKY 222

FIG. 8

47324, A NOVEL HUMAN G-PROTEIN AND USES THEREFOR

[0001] This application claims benefit of priority from U.S. application Ser. No. 60/229,293 filed Sep. 1, 2000, which is hereby incorporated by reference in its entirety.

BACKGROUND OF THE INVENTION

[0002] The G protein superfamily (e.g., heterotrimeric and small G proteins) encompasses a diverse array of proteins which regulate a complex range of biological processes, including the regulation of protein synthesis, cellular trafficking (e.g., vesicular and nuclear transport), regulation of the cell cycle, growth, differentiation, apoptosis, and cytoskeletal rearrangements (Cerione et al. (1996) *Curr. Op. Cell Biol.* 8:216-222; Cherfils et al. (1999) *Trends Biochem. Sci.* 24:306-311). The common motif among this important family of proteins is the presence of a GTP-binding domain (Alberts et al. (1994) *Molecular Biology of the Cell*, Garland Publishing, Inc., New York, N.Y. pp. 206-207, 641). These proteins act as molecular switches that can cycle between active (GTP-bound) and inactive (GDP-bound) states (Bourne et al. (1990) *Nature*, 348:125-132). In the active state, G proteins are able to interact with a broad range of effector molecules. These effector molecules constitute components of a variety of signaling cascades. Upon hydrolysis of bound GTP, the G protein switches to the inactive state, a step that is facilitated by GTPase activating proteins (GAPs) (Scheffzek et al. (1998) *Trends Biochem. Sci.* 23:257-262; Gambelin and Smerdon (1998) *Curr. Opinion in Struct. Biol.* 8:195-201).

[0003] Activation of G proteins is mediated by the exchange of GDP for GTP. Dissociation of GDP from the inactive small G protein is facilitated by a class of proteins known as guanine nucleotide exchange factors (GEFs) or GTP-releasing factors. The small G protein is then able to bind GTP and undergo conformational changes which allow it to interact with effector molecules.

[0004] GEFs consist of four families based on sequence similarity among family members and on selectivity of small G protein activated by the GEF, including GEFs of Ran, ARF, Ras, and Rho (also known as the Dbl homology (DH) domain-containing GEFs). GEF family members all contain a GEF homology domain amino terminal to a pleckstrin homology (PH) domain, and most contain other functional domains commonly found in signaling molecules (Cerione et al. and Cherfils et al., supra). For example, the GEF family members Dbs and Vav both have Src homology (SH3) domains at their carboxyl termini (Whitehead et al. (1995) *Oncogene* 10:713-721).

[0005] Many GEF family members have been identified to date including Dbl, Ost, Tiam-1, Ect-2, Vav, Lbc, FGD1, Dbs, Lfc, Tim, Brc, Abr, Sos, and Ras GEF. These proteins are found in various tissues including adrenal gland, brain, gonad, heart, keratinocyte, kidney, liver, lung, mammary epithelial, myeloid, pancreas, placenta, spleen, skeletal muscle, testis, and fetal brain and heart, and in diffuse B-cell lymphomas, osteosarcomas, T-lymphoma cells, and myeloid leukemias. The Sos protein is ubiquitous (Cerione et al., supra).

[0006] It is the regulated cycling between active and inactive states of G proteins that allows for proper trans-

duction of many vital cellular signals. Indeed, the regulation of GTP/GDP levels in the cell by small G proteins and their accessory GEF molecules, has been implicated in a number of diseases, including oncogenesis and metastasis, facio-genital dysplasia and chronic myelogenous leukemia (Cerione et al., supra).

SUMMARY OF THE INVENTION

[0007] The present invention is based, in part, on the discovery of a novel human G-protein, referred to herein as "47324". The nucleotide sequence of a cDNA encoding 47324 is shown in SEQ ID NO:1, and the amino acid sequence of a 47324 polypeptide is shown in SEQ ID NO:2. In addition, the nucleotide sequence of the coding region is depicted in SEQ ID NO:3.

[0008] Accordingly, in one aspect, the invention features a nucleic acid molecule which encodes a 47324 protein or polypeptide, e.g., a biologically active portion of the 47324 protein. In a preferred embodiment, the isolated nucleic acid molecule encodes a polypeptide having the amino acid sequence of SEQ ID NO:2. In other embodiments, the invention provides an isolated 47324 nucleic acid molecule having the nucleotide sequence shown in SEQ ID NO:1, SEQ ID NO:3, or the sequence of the DNA insert of the plasmid deposited with ATCC Accession Number _____. In still other embodiments, the invention provides nucleic acid molecules that are substantially identical (e.g., naturally occurring allelic variants) to the nucleotide sequence shown in SEQ ID NO:1, SEQ ID NO:3, or the sequence of the DNA insert of the plasmid deposited with ATCC Accession Number _____. In other embodiments, the invention provides a nucleic acid molecule which hybridizes under stringent hybridization conditions to a nucleic acid molecule comprising the nucleotide sequence of SEQ ID NO:1, SEQ ID NO:3, or the sequence of the DNA insert of the plasmid deposited with ATCC Accession Number _____, wherein the nucleic acid encodes a full length 47324 protein or an active fragment thereof.

[0009] In a related aspect, the invention further provides nucleic acid constructs which include a 47324 nucleic acid molecule described herein. In certain embodiments, the nucleic acid molecules of the invention are operatively linked to native or heterologous regulatory sequences. Also included, are vectors and host cells containing the 47324 nucleic acid molecules of the invention e.g., vectors and host cells suitable for producing 47324 nucleic acid molecules and polypeptides.

[0010] In another related aspect, the invention provides nucleic acid fragments suitable as primers or hybridization probes for the detection of 47324-encoding nucleic acids.

[0011] In still another related aspect, isolated nucleic acid molecules that are antisense to a 47324 encoding nucleic acid molecule are provided.

[0012] In another aspect, the invention features, 47324 polypeptides, and biologically active or antigenic fragments thereof that are useful, e.g., as reagents or targets in assays applicable to treatment and diagnosis of 47324-mediated or -related disorders. In another embodiment, the invention provides 47324 polypeptides having a 47324 activity. Preferred polypeptides are 47324 proteins including at least one Ras family domain, and, preferably, having a 47324 activity, e.g., a 47324 activity as described herein.

[0013] In other embodiments, the invention provides 47324 polypeptides, e.g., a 47324 polypeptide having the amino acid sequence shown in SEQ ID NO:2; the amino acid sequence encoded by the cDNA insert of the plasmid deposited with ATCC Accession Number _____; an amino acid sequence that is substantially identical to the amino acid sequence shown in SEQ ID NO:2; or an amino acid sequence encoded by a nucleic acid molecule having a nucleotide sequence which hybridizes under stringent hybridization conditions to a nucleic acid molecule comprising the nucleotide sequence of SEQ ID NO:1, SEQ ID NO:3, or the sequence of the DNA insert of the plasmid deposited with ATCC Accession Number _____, wherein the nucleic acid encodes a full length 47324 protein or an active fragment thereof.

[0014] In a related aspect, the invention further provides nucleic acid constructs which include a 47324 nucleic acid molecule described herein.

[0015] In a related aspect, the invention provides 47324 polypeptides or fragments operatively linked to non-47324 polypeptides to form fusion proteins.

[0016] In another aspect, the invention features antibodies and antigen-binding fragments thereof, that react with, or more preferably specifically bind 47324 polypeptides.

[0017] In another aspect, the invention provides methods of screening for compounds that modulate the expression or activity of the 47324 polypeptides or nucleic acids.

[0018] In still another aspect, the invention provides a process for modulating 47324 polypeptide or nucleic acid expression or activity, e.g. using the screened compounds. In certain embodiments, the methods involve treatment of conditions related to aberrant activity or expression of the 47324 polypeptides or nucleic acids, such as conditions involving aberrant or deficient cellular proliferation or differentiation.

[0019] The invention also provides assays for determining the activity of or the presence or absence of 47324 polypeptides or nucleic acid molecules in a biological sample, including for disease diagnosis.

[0020] In further aspect the invention provides assays for determining the presence or absence of a genetic alteration in a 47324 polypeptide or nucleic acid molecule, including for disease diagnosis.

[0021] In another aspect, the invention features a two dimensional array having a plurality of addresses, each address of the plurality being positionally distinguishable from each other address of the plurality, and each address of the plurality having a unique capture probe, e.g., a nucleic acid or peptide sequence. At least one address of the plurality has a capture probe that recognizes a 47324 molecule. In one embodiment, the capture probe is a nucleic acid, e.g., a probe complementary to a 47324 nucleic acid sequence. In another embodiment, the capture probe is a polypeptide, e.g., an antibody specific for 47324 polypeptides. Also featured is a method of analyzing a sample by contacting the sample to the aforementioned array and detecting binding of the sample to the array.

[0022] Other features and advantages of the invention will be apparent from the following detailed description, and from the claims.

BRIEF DESCRIPTION OF THE DRAWINGS

[0023] FIGS. 1A-B depicts a cDNA sequence (SEQ ID NO:1) and predicted amino acid sequence (SEQ ID NO:2) of human 47324. The methionine-initiated open reading frame of human 47324 (without the 5' and 3' untranslated regions) extends from nucleotide position 1 to position 711 of SEQ ID NO:3, including the terminal codon.

[0024] FIG. 2 depicts a hydropathy plot of human 47324. Relatively hydrophobic residues are shown above the dashed horizontal line, and relatively hydrophilic residues are below the dashed horizontal line. The cysteine residues (cys) are indicated by short vertical lines just below the hydropathy trace. The numbers corresponding to the amino acid sequence of human 47324 are indicated. Polypeptides of the invention include fragments which include: all or part of a hydrophobic sequence, e.g., a sequence above the dashed line, e.g., the sequence from about amino acid 6 to 14, and from about 63 to 73 of SEQ ID NO:2; all or part of a hydrophilic sequence, e.g., a sequence below the dashed line, e.g., the sequence from about amino acid 48 to 58, from about 98 to 108, and from about 205 to 220 of SEQ ID NO:2; or a sequence which includes a Cys.

[0025] FIG. 3 depicts an alignment of the ADP-ribosylation factor family domain of human 47324 with a consensus amino acid sequence derived from a hidden Markov model (HMM) from PFAM. The upper sequences are the consensus amino acid sequence (SEQ ID NO:4), while the lower amino acid sequences correspond to amino acids 2 to 206 of SEQ ID NO:2.

[0026] FIG. 4 depicts an alignment of the Ras family domain of human 47324 with a consensus amino acid sequence derived from a hidden Markov model (HMM) from PFAM. The upper sequences are the consensus amino acid sequence (SEQ ID NO:5), while the lower amino acid sequences correspond to amino acids 8 to 231 of SEQ ID NO:2.

[0027] FIGS. 5a-b depicts a BLAST alignment of human 47324 with a consensus amino acid sequence derived from a ProDomain No. PD314684, "CG15399" (Release 2001.1; <http://www.toulouse.inra.fr/prodom.html>). The lower sequence is amino acid residues 69 to 234 and 12 to 61 of the 242 amino acid consensus sequence (SEQ ID NOs:6 and 7), while the upper amino acid sequence corresponds to the "CG15399" domain of human 47324, amino acid residues 35 to 209 and 7 to 52 of SEQ ID NO:2.

[0028] FIG. 6 depicts a BLAST alignment of human 47324 with a consensus amino acid sequence derived from a ProDomain No. PD302329, "CG4789" (Release 2001.1; <http://www.toulouse.inra.fr/prodom.html>). The lower sequence is amino acid residues 19 to 140 of the 155 amino acid consensus sequence (SEQ ID NO:8), while the upper amino acid sequence corresponds to the "CG4789" domain of human 47324, amino acid residues 121 to 231 of SEQ ID NO:2.

[0029] FIG. 7 depicts a BLAST alignment of human 47324 with a consensus amino acid sequence derived from a ProDomain No. PD301653, "GB|AAF03448.1" (Release 2001.1; <http://www.toulouse.inra.fr/prodom.html>). The lower sequence is amino acid residues 80 to 135 of the 152 amino acid consensus sequence (SEQ ID NO:9), while the upper amino acid sequence corresponds to the

“GB|AAF03448.1” domain of human 47324, amino acid residues 6 to 63 of SEQ ID NO:2.

[0030] FIG. 8 depicts a BLAST alignment of human 47324 with a consensus amino acid sequence derived from a ProDomain No. PD000015, “GTP-binding lipoprotein prenylation transport Ras-related factor initiation ADP-ribosylation small family” (Release 2001.1; <http://www.toulouse.inra.fr/prodom.html>). The lower sequence is amino acid residues 7 to 222 of the 229 amino acid consensus sequence (SEQ ID NO:10), while the upper amino acid sequence corresponds to the “GTP-binding lipoprotein prenylation transport Ras-related factor initiation ADP-ribosylation small family” domain of human 47324, amino acid residues 8 to 225 of SEQ ID NO:2.

[0031] Other features and advantages of the invention will be apparent from the following detailed description, and from the claims.

DETAILED DESCRIPTION

[0032] Human 47324

[0033] The human 47324 sequence (FIGS. 1A-B; SEQ ID NO:1), which is approximately 1694 nucleotides long including untranslated regions, contains a predicted methionine-initiated coding sequence of about 711 nucleotides (nucleotides 19-729 of SEQ ID NO:1; SEQ ID NO:3), including the terminal codon. The coding sequence encodes a 236 amino acid protein (SEQ ID NO:2).

[0034] This mature protein form is approximately 236 amino acid residues in length (from about amino acid 1 to amino acid 236 of SEQ ID NO:2). Human 47324 contains the following regions or other structural features:

[0035] an ADP-ribosylation factor family domain located at about amino acid residues 2-206 of SEQ ID NO:2;

[0036] a Ras family domain located at about amino acid residues 8-231;

[0037] one predicted glycosaminoglycan attachment site (PS00002) located at about amino acids 15-18 of SEQ ID NO:2;

[0038] two predicted cAMP- and cGMP-dependent protein kinase phosphorylation sites (PS00004) located at about amino acid residues 100-103 and 107-110 of SEQ ID NO:2;

[0039] five predicted protein kinase C phosphorylation sites (PS00005) located at about amino acid residues 76-78, 79-81, 98-100, 155-157, and 229-231 of SEQ ID NO:2;

[0040] three predicted casein kinase II phosphorylation sites (PS00006) located at about amino acid residues 54-57, 126-129, and 147-150 of SEQ ID NO:2;

[0041] one predicted located at about amino acid residues 202-208 of SEQ ID NO:2;

[0042] five predicted N-myristoylation sites (PS00008) located at about amino acid residues 16-21, 68-73, 122-127, 189-194, and 228-233 of SEQ ID NO:2;

[0043] one predicted prokaryotic membrane lipoprotein lipid attachment site (PS00013) located at about amino acid residues 31-41 of SEQ ID NO:2;

[0044] one predicted ATP/GTP-binding site motif A (P-loop) site (PS00017) located at about amino acid residues 13-20 of SEQ ID NO:2; and

[0045] one predicted Sigma-54 interaction domain STP-binding region A signature site (PS00675) located at about amino acid residues 9-22 of SEQ ID NO:2.

[0046] For general information regarding PFAM identifiers, PS prefix and PF prefix domain identification numbers, refer to Sonnhammer et al. (1997) *Protein* 28:405-420 and <http://www.psc.edu/general/software/packages/pfam/pfam.html>.

[0047] A plasmid containing the nucleotide sequence encoding human 47324 was deposited with American Type Culture Collection (ATCC), 10801 University Boulevard, Manassas, Va. 20110-2209, on _____ and assigned Accession Number _____. This deposit will be maintained under the terms of the Budapest Treaty on the International Recognition of the Deposit of Microorganisms for the Purposes of Patent Procedure. This deposit was made merely as a convenience for those of skill in the art and is not an admission that a deposit is required under 35 U.S.C. §112.

[0048] The 47324 protein contains a significant number of structural characteristics in common with members of the Ras family. The term “family” when referring to the protein and nucleic acid molecules of the invention means two or more proteins or nucleic acid molecules having a common structural domain or motif and having sufficient amino acid or nucleotide sequence homology as defined herein. Such family members can be naturally or non-naturally occurring and can be from either the same or different species. For example, a family can contain a first protein of human origin as well as other distinct proteins of human origin, or alternatively, can contain homologues of non-human origin, e.g., rat or mouse proteins. Members of a family can also have common functional characteristics.

[0049] As used herein, the term “Ras family” includes a molecule which is involved in the signal transduction pathway associated with a large number of extracellular signals. The Ras family can interact with a large number of molecules following activation by interaction with GTP. Ras family molecules are involved in the growth, development, and differentiation of cells, in the regulation of cellular homeostasis, in the metabolism and catabolism of biochemical molecules necessary for energy production or storage, in intra- or intercellular signaling, in metabolism or catabolism of metabolically important biomolecules, and in the removal of potentially harmful compounds from the interior of the cell. Examples of Ras family members include Ras, Rap1, Ral and Rhex. The Ras family molecules of the present invention provide novel diagnostic targets and therapeutic agents to control Ras family-associated disorders.

[0050] As used herein, a “47324 activity”, “biological activity of 47324” or “functional activity of 47324”, refers to an activity exerted by a 47324 protein, polypeptide or nucleic acid molecule on e.g., a 47324-responsive cell or on a 47324 substrate, e.g., a lipid or protein substrate, as determined in vivo or in vitro. In one embodiment, a 47324

activity is a direct activity, such as an association with a 47324 target molecule. A “target molecule” or “binding partner” is a molecule with which a 47324 protein binds or interacts in nature, e.g., a molecule in which the 47324 protein activates a kinase activity. A 47324 activity can also be an indirect activity, e.g., a cellular signaling activity mediated by interaction of the 47324 protein with a 47324 ligand. For example, the 47324 proteins of the present invention can have one or more of the following activities: 1) activation by association with a substrate or binding partner (e.g., a Guanine nucleotide exchange factor); 2) association with a lipid bilayer due to post-translational modifications (e.g., methylation and farnesylation); 3) modulation of signal transduction (e.g., signal transduction cascades involving small GTP-binding proteins); 4) control of cell morphology and differentiation; 5) mediation of cellular proliferation; 6) mediation of apoptosis; 7) modulation of tumor inhibition; and 8) regulation of cellular trafficking (e.g. vesicular transport); 9) regulation of active cytoskeleton organization; 10) the ability to antagonize or inhibit, competitively or non-competitively, any of 1-7. Thus, the 47324 molecules can act as novel diagnostic targets and therapeutic agents for controlling G protein-related disorders, for example, such as those diseases associated with the activities described above. As the 47324 molecules have homology to known G proteins, in particular, ras-family proteins, they are expected to be involved in controlling similar disorders.

[0051] To identify the presence of a “Ras family” domain in a 47324 protein sequence, and make the determination that a polypeptide or protein of interest has a particular profile, the amino acid sequence of the protein can be searched against a database of HMMs (e.g., the Pfam database, release 2.1) using the default parameters (http://www.sanger.ac.uk/Software/Pfam/HMM_search). For example, the hnmmsf program, which is available as part of the HMMER package of search programs, is a family specific default program for MILPAT0063 and a score of 15 is the default threshold score for determining a hit. Alternatively, the threshold score for determining a hit can be lowered (e.g., to 8 bits). A description of the Pfam database can be found in Sonhammer et al., (1997) *Proteins* 28(3):405-420 and a detailed description of HMMs can be found, for example, in Gribskov et al., (1990) *Meth. Enzymol.* 183:146-159; Gribskov et al., (1987) *Proc. Natl. Acad. Sci. USA* 84:4355-4358; Krogh et al., (1994) *J. Mol. Biol.* 235:1501-1531; and Stultz et al., (1993) *Protein Sci.* 2:305-314, the contents of which are incorporated herein by reference.

[0052] A 47324 polypeptide can include an “ADP-ribosylation factor family domain” or regions homologous with a “ADP-ribosylation factor family domain”. As used herein, the term “ADP-ribosylation factor family domain” includes an amino acid sequence of about 100-250 amino acid residues in length and having a bit score for the alignment of the sequence to the ADP-ribosylation factor family domain (HMM) of at least 8. Preferably, an ADP-ribosylation factor family domain includes at least about 150-250 amino acids, more preferably about 175-250 amino acid residues, or about 200-225 amino acids and has a bit score for the alignment of the sequence to the ADP-ribosylation factor family domain (HMM) of at least 16 or greater. The ADP-ribosylation factor family domain (HMM) has been assigned the PFAM Accession PF00025 (<http://pfam.wustl.edu/>).

An alignment of the ADP-ribosylation factor family domain (amino acids 2 to 206 of SEQ ID NO:2) of human 47324 with a consensus amino acid sequence derived from a hidden Markov model is depicted in **FIG. 3**.

[0053] Preferably, the ADP-ribosylation factor family domain includes the following amino acid consensus sequence having Prosite signatures as PS01316, or sequences homologous thereto: E-x(5)-G-x-[SAG]-x(2)-[IV]-x-D-[LIV]-x(2)-[ST]-G-x-T-[LM] (SEQ ID NO:11). In the above conserved motif, and other motifs described herein, the standard IUPAC one-letter code for the amino acids is used. Each element in the pattern is separated by a dash (-); square brackets ([]) indicate the particular residues that are accepted at that position; x indicates that any residue is accepted at that position; and numbers in parentheses (()) indicate the number of residues represented by the accompanying amino acid. The ADP-ribosylation factor family domain corresponds to about amino acids 2-206 of SEQ ID NO:2. The ADP-ribosylation factor family domain (HMM) has been assigned the PFAM Accession Number PF00025 (<http://genome.wustl.edu/Pfam/html>).

[0054] In a preferred embodiment 47324 polypeptide or protein has an “ADP-ribosylation factor family domain” or a region which includes at least about 100-250 more preferably about 150-250 or 200-225 amino acid residues and has at least about 60%, 70%, 80%, 90%, 95%, 99%, or 100% homology with a “Ras family domain,” e.g., the ADP-ribosylation factor family domain of human 47324 (e.g., amino acid residues 2-206 of SEQ ID NO:2).

[0055] A 47324 polypeptide can also include a “Ras family domain” or regions homologous with a “Ras family domain”. As used herein, the term “Ras family domain” includes an amino acid sequence of about 100-250 amino acid residues in length and having a bit score for the alignment of the sequence to the Ras family domain (HMM) of at least 8. Preferably, a Ras family domain includes at least about 100-250 amino acids, more preferably about 150-250 amino acid residues, or about 200-225 amino acids and has a bit score for the alignment of the sequence to the Ras family domain (HMM) of at least 16 or greater. The Ras family domain (HMM) has been assigned the PFAM Accession PF00071 (<http://pfam.wustl.edu/>). An alignment of the Ras family domain (amino acids 8 to 231 of SEQ ID NO:2) of human 47324 with a consensus amino acid sequence derived from a hidden Markov model is depicted in **FIG. 4**.

[0056] In a preferred embodiment 47324 polypeptide or protein has a “Ras family domain” or a region which includes at least about 100-250 more preferably about 150-250 or 200-225 amino acid residues and has at least about 60%, 70%, 80%, 90%, 95%, 99%, or 100% homology with a “Ras family domain,” e.g., the Ras family domain of human 47324 (e.g., amino acid residues 8-231 of SEQ ID NO:2).

[0057] An additional method to identify domains in a 47324 protein sequence, and make the determination that a polypeptide or protein of interest has a particular profile, the amino acid sequence of the protein can be searched against a SMART database (Simple Modular Architecture Research Tool, <http://smart.embl-heidelberg.de/>) of HMMs as described in Schultz et al. (1998), *Proc. Natl. Acad. Sci. USA* 95:5857 and Schultz et al. (2000) *Nucl. Acids Res* 28:231. The database contains domains identified by profiling with

the hidden Markov models of the HMMer2 search program (R. Durbin et al. (1998) *Biological sequence analysis: probabilistic models of proteins and nucleic acids*. Cambridge University Press.; <http://hmmerr.wustl.edu/>). The database also is extensively annotated and monitored by experts to enhance accuracy. A search was performed against the HMM database resulting in the identification of an "AAA_5" domain in the amino acid sequence of human 47324 at about residues 29 to 241 of SEQ ID NO:2; a "rab_sub_5" domain in the amino acid sequence of human 47324 at about residues 7-187 of SEQ ID NO:2; a "ras_sub_4" domain in the amino acid sequence of human 47324 at about residues 4-209 of SEQ ID NO:2; a "rho_sub_3" domain in the amino acid sequence of human 47324 at about residues 9-183 of SEQ ID NO:2 (see FIG. 1).

[0058] For further identification of domains in a 47324 protein sequence, and make the determination that a polypeptide or protein of interest has a particular profile, the amino acid sequence of the protein can be searched against a database of domains, e.g., the ProDom database (Corpet et al. (1999), *Nucl. Acids Res.* 27:263-267). The ProDom protein domain database consists of an automatic compilation of homologous domains. Current versions of ProDom are built using recursive PSI-BLAST searches (Altschul S F et al. (1997) *Nucleic Acids Res.* 25:3389-3402; Gouzy et al. (1999) 23:333-340) of the SWISS-PROT 38 and TREMBL protein databases. The database automatically generates a consensus sequence for each domain.

[0059] A BLAST search was performed against the HMM database resulting in the identification of regions homologous to ProDom family PD314684 ("CG15399" SEQ ID NO:6 and 7, ProDomain Release 2001.1; <http://www.toulouse.inra.fr/prodom.html>). An alignment of the "CG15399" domain (amino acids 35-209 and 7-52 of SEQ ID NO:2) of human 47324 with consensus amino acid sequences (SEQ ID NOs:6 and 7) derived from a hidden Markov model is depicted in FIGS. 5a and 5b. The consensus sequence for SEQ ID NO:6 is 25% identical over amino acids 35 to 209 and for SEQ ID NO:7 is 44% identical over amino acids 7 to 52 of SEQ ID NO:2 as shown in FIGS. 5a and 5b.

[0060] A BLAST search was performed against the HMM database resulting in the identification of regions homologous to ProDom family PD302329 ("CG4789" SEQ ID NO:8, ProDomain Release 2001.1; <http://www.toulouse.inra.fr/prodom.html>). An alignment of the "CG4789" domain (amino acids 121-231 of SEQ ID NO:2) of human 47324 with consensus amino acid sequences (SEQ ID NO:8) derived from a hidden Markov model is depicted in FIG. 6. The consensus sequence for SEQ ID NO:8 is 43% identical over amino acids 121 to 131 of SEQ ID NO:2 as shown in FIG. 6.

[0061] A BLAST search was performed against the HMM database resulting in the identification of regions homologous to ProDom family PD301653 ("GB|AAF03448.1" SEQ ID NO:9, ProDomain Release 2001.1; <http://www.toulouse.inra.fr/prodom.html>). An alignment of the "GB|AAF03448.1" domain (amino acids 6-63 of SEQ ID NO:2) of human 47324 with consensus amino acid sequences (SEQ ID NO:9) derived from a hidden Markov model is depicted in FIG. 7. The consensus sequence for SEQ ID NO:9 is 44% identical over amino acids 6 to 63 of SEQ ID NO:2 as shown in FIG. 7.

[0062] A BLAST search was performed against the HMM database resulting in the identification of regions homologous to ProDom family PD000015 ("GTP-binding lipoprotein prenylation transport Ras-related factor initiation ADP-ribosylation small family" SEQ ID NO:10, ProDomain Release 2001.1; <http://www.toulouse.inra.fr/prodom.html>). An alignment of the "GTP-binding lipoprotein prenylation transport Ras-related factor initiation ADP-ribosylation small family" domain (amino acids 8-225 of SEQ ID NO:2) of human 47324 with consensus amino acid sequences (SEQ ID NO:10) derived from a hidden Markov model is depicted in FIG. 8. The consensus sequence for SEQ ID NO:10 is 21% identical over amino acids 8 to 225 of SEQ ID NO:2 as shown in FIG. 8.

[0063] In a preferred embodiment, a 47324 family member can include at least one ADP-ribosylation factor family domain (PFAM Accession Number PF00025) or at least one Ras family domain (PFAM Accession Number PF00071). Furthermore, a 47324 family member can include at least one glycosaminoglycan attachment site (PS00002); at least one, and preferably two cAMP- and cGMP-dependent protein kinase phosphorylation sites (PS00004); at least one, two, three, four, and preferably five protein kinase C phosphorylation sites (PS00005); at least one, two, and preferably three casein kinase II phosphorylation sites (PS00006); at least one tyrosine kinase phosphorylation site (PS00007); at least one, two, three, four, and preferably five N-myristoylation sites (PS00008); at least one prokaryotic membrane lipoprotein lipid attachment site (PS00013); at least one ATP/GTP-binding site motif A (P-loop) site (PS00017); and at least one Sigma-54 interaction domain STP-binding region A signature site (PS00675).

[0064] As used herein, a "Ras family-associated disorder" includes a disorder, disease or condition which is caused or characterized by a misregulation (e.g., downregulation or upregulation) of a Ras family-mediated activity. Ras family-associated disorders can detrimentally affect cellular functions such as cellular proliferation, growth, differentiation, or migration, cellular regulation of homeostasis, inter- or intra-cellular communication; tissue function, such as cardiovascular function, protection of cells from toxic compounds (e.g., carcinogens, toxins, mutagens, and toxic byproducts of metabolic activity (e.g., reactive oxygen species)), brain function, blood vessel development and function, hematopoiesis, and platelet production. Accordingly, 47324 protein may mediate various disorders, including cellular proliferative and/or differentiative disorders, cardiovascular disorders, liver disorders, pain and metabolic disorders, brain disorders, blood vessel disorders, hematopoietic disorders and platelet disorders. As the 47324 polypeptides of the invention may modulate 47324-mediated activities, they may be useful for developing novel diagnostic and therapeutic agents for 47324-mediated or related disorders, as described below.

[0065] Ras-family proteins are essential for signal transduction from receptor tyrosine kinases to serine/threonine kinases which are involved in cell growth and differentiation. Ras-family proteins that have undergone mutations may become permanently activated, resulting in continuous cell proliferation. An example of this is TC21, a Ras-like protein that has been found to be highly expressed in human teratocarcinoma cell lines. The 47324 polypeptides share a common domain with known Ras-family members and is

expected to have similar effects in cell proliferation. Accordingly, 47324 may play a role in cell proliferation and cancer, inflammation and apoptosis, and thus the 47324 compositions of the invention (e.g., nucleic acids, polypeptides, proteins, antibodies) can be used to modulate cell proliferation, e.g., in cancer, inflammation or apoptosis, and furthermore can be used in screening assays to identify agents for modulating cell proliferation, as well as in detection or diagnostic assays to identify conditions involving aberrant cell proliferation.

[0066] Examples of cellular proliferative and/or differentiative disorders include cancer, e.g., carcinoma, sarcoma, metastatic disorders or hematopoietic neoplastic disorders, e.g., leukemias. A metastatic tumor can arise from a multitude of primary tumor types, including but not limited to those of prostate, colon, lung, breast and liver origin.

[0067] As used herein, the terms “cancer”, “hyperproliferative” and “neoplastic” refer to cells having the capacity for autonomous growth, i.e., an abnormal state or condition characterized by rapidly proliferating cell growth. Hyperproliferative and neoplastic disease states may be categorized as pathologic, i.e., characterizing or constituting a disease state, or may be categorized as non-pathologic, i.e., a deviation from normal but not associated with a disease state. The term is meant to include all types of cancerous growths or oncogenic processes, metastatic tissues or malignantly transformed cells, tissues, or organs, irrespective of histopathologic type or stage of invasiveness. “Pathologic hyperproliferative” cells occur in disease states characterized by malignant tumor growth. Examples of non-pathologic hyperproliferative cells include proliferation of cells associated with wound repair.

[0068] The terms “cancer” or “neoplasms” include malignancies of the various organ systems, such as affecting lung, breast, thyroid, lymphoid, gastrointestinal, and genito-urinary tract, as well as adenocarcinomas which include malignancies such as most colon cancers, renal-cell carcinoma, prostate cancer and/or testicular tumors, non-small cell carcinoma of the lung, cancer of the small intestine and cancer of the esophagus.

[0069] The term “carcinoma” is art recognized and refers to malignancies of epithelial or endocrine tissues including respiratory system carcinomas, gastrointestinal system carcinomas, genitourinary system carcinomas, testicular carcinomas, breast carcinomas, prostatic carcinomas, endocrine system carcinomas, and melanomas. Exemplary carcinomas include those forming from tissue of the cervix, lung, prostate, breast, head and neck, colon and ovary. The term also includes carcinosarcomas, e.g., which include malignant tumors composed of carcinomatous and sarcomatous tissues. An “adenocarcinoma” refers to a carcinoma derived from glandular tissue or in which the tumor cells form recognizable glandular structures.

[0070] The term “sarcoma” is art recognized and refers to malignant tumors of mesenchymal derivation.

[0071] The 47324 nucleic acid and protein of the invention can be used to monitor, treat and/or diagnose a variety of proliferative disorders. E.g., such disorders include hematopoietic neoplastic disorders. As used herein, the term “hematopoietic neoplastic disorders” includes diseases involving hyperplastic/neoplastic cells of hematopoietic ori-

gin, e.g., arising from myeloid, lymphoid or erythroid lineages, or precursor cells thereof. Preferably, the diseases arise from poorly differentiated acute leukemias, e.g., erythroblastic leukemia and acute megakaryoblastic leukemia. Additional exemplary myeloid disorders include, but are not limited to, acute promyeloid leukemia (APML), acute myelogenous leukemia (AML) and chronic myelogenous leukemia (CML) (reviewed in Vaickus, L., (1991) *Crit. Rev. in Oncol./Hematol.* 11:267-97); lymphoid malignancies include, but are not limited to acute lymphoblastic leukemia (ALL) which includes B-lineage ALL and T-lineage ALL, chronic lymphocytic leukemia (CLL), prolymphocytic leukemia (PLL), hairy cell leukemia (HLL) and Waldenstrom's macroglobulinemia (WM). Additional forms of malignant lymphomas include, but are not limited to non-Hodgkin lymphoma and variants thereof, peripheral T cell lymphoma, adult T cell leukemia/lymphoma (ATL), cutaneous T-cell lymphoma (CTCL), large granular lymphocytic leukemia (LGL), Hodgkin's disease and Reed-Stemberg disease.

[0072] Examples of disorders involving the heart or “cardiovascular disorder” include, but are not limited to, a disease, disorder, or state involving the cardiovascular system, e.g., the heart, the blood vessels, and/or the blood. A cardiovascular disorder can be caused by an imbalance in arterial pressure, a malfunction of the heart, or an occlusion of a blood vessel, e.g., by a thrombus. Examples of cardiovascular disorders include but are not limited to, hypertension, atherosclerosis, coronary artery spasm, coronary artery disease, arrhythmias, heart failure, including but not limited to, cardiac hypertrophy, left-sided heart failure, and right-sided heart failure; ischemic heart disease, including but not limited to angina pectoris, myocardial infarction, chronic ischemic heart disease, and sudden cardiac death; hypertensive heart disease, including but not limited to, systemic (left-sided) hypertensive heart disease and pulmonary (right-sided) hypertensive heart disease; valvular heart disease, including but not limited to, valvular degeneration caused by calcification, such as calcification of a congenitally bicuspid aortic valve, and mitral annular calcification, and myxomatous degeneration of the mitral valve (mitral valve prolapse), rheumatic fever and rheumatic heart disease, infective endocarditis, and noninfected vegetations, such as nonbacterial thrombotic endocarditis and endocarditis of systemic lupus erythematosus (Libman-Sacks disease), carcinoid heart disease, and complications of artificial valves; myocardial disease, including but not limited to dilated cardiomyopathy, hypertrophic cardiomyopathy, restrictive cardiomyopathy, and myocarditis; pericardial disease, including but not limited to, pericardial effusion and hemopericardium and pericarditis, including acute pericarditis and healed pericarditis, and rheumatoid heart disease; neoplastic heart disease, including but not limited to, primary cardiac tumors, such as myxoma, lipoma, papillary fibroelastoma, rhabdomyoma, and sarcoma, and cardiac effects of noncardiac neoplasms; congenital heart disease, including but not limited to, left-to-right shunts—late cyanosis, such as atrial septal defect, ventricular septal defect, patent ductus arteriosus, and atrioventricular septal defect, right-to-left shunts—early cyanosis, such as tetralogy of fallot, transposition of great arteries, truncus arteriosus, tricuspid atresia, and total anomalous pulmonary venous connection, obstructive congenital anomalies, such as coarctation of aorta,

pulmonary stenosis and atresia, and aortic stenosis and atresia, disorders involving cardiac transplantation, and congestive heart failure.

[0073] A cardiovascular disease or disorder also includes an endothelial cell disorder. As used herein, an "endothelial cell disorder" includes a disorder characterized by aberrant, unregulated, or unwanted endothelial cell activity, e.g., proliferation, migration, angiogenesis, or vascularization; or aberrant expression of cell surface adhesion molecules or genes associated with angiogenesis, e.g., TIE-2, FLT and FLK. Endothelial cell disorders include tumorigenesis, tumor metastasis, psoriasis, diabetic retinopathy, endometriosis, Grave's disease, ischemic disease (e.g., atherosclerosis), and chronic inflammatory diseases (e.g., rheumatoid arthritis).

[0074] Disorders which may be treated or diagnosed by methods described herein include, but are not limited to, disorders associated with an accumulation in the liver of fibrous tissue, such as that resulting from an imbalance between production and degradation of the extracellular matrix accompanied by the collapse and condensation of preexisting fibers. The methods described herein can be used to diagnose or treat hepatocellular necrosis or injury induced by a wide variety of agents including processes which disturb homeostasis, such as an inflammatory process, tissue damage resulting from toxic injury or altered hepatic blood flow, and infections (e.g., bacterial, viral and parasitic). For example, the methods can be used for the early detection of hepatic injury, such as portal hypertension or hepatic fibrosis. In addition, the methods can be employed to detect liver fibrosis attributed to inborn errors of metabolism, for example, fibrosis resulting from a storage disorder such as Gaucher's disease (lipid abnormalities) or a glycogen storage disease, A1-antitrypsin deficiency; a disorder mediating the accumulation (e.g., storage) of an exogenous substance, for example, hemochromatosis (iron-overload syndrome) and copper storage diseases (Wilson's disease), disorders resulting in the accumulation of a toxic metabolite (e.g., tyrosinemia, fructosemia and galactosemia) and peroxisomal disorders (e.g., Zellweger syndrome). Additionally, the methods described herein may be useful for the early detection and treatment of liver injury associated with the administration of various chemicals or drugs, such as for example, methotrexate, isoniazid, oxyphenisatin, methyl-dopa, chlorpromazine, tolbutamide or alcohol, or which represents a hepatic manifestation of a vascular disorder such as obstruction of either the intrahepatic or extrahepatic bile flow or an alteration in hepatic circulation resulting, for example, from chronic heart failure, veno-occlusive disease, portal vein thrombosis or Budd-Chiari syndrome.

[0075] Additionally, 47324 may play an important role in the regulation of metabolism or pain disorders. Diseases of metabolic imbalance include, but are not limited to, obesity, anorexia nervosa, cachexia, lipid disorders, and diabetes. Examples of pain disorders include, but are not limited to, pain response elicited during various forms of tissue injury, e.g., inflammation, infection, and ischemia, usually referred to as hyperalgesia (described in, for example, Fields, H. L., (1987) *Pain*, New York: McGraw-Hill); pain associated with musculoskeletal disorders, e.g., joint pain; tooth pain; headaches; pain associated with surgery; pain related to irritable bowel syndrome; or chest pain.

[0076] Disorders involving the brain include, but are not limited to, disorders involving neurons, and disorders involving glia, such as astrocytes, oligodendrocytes, ependymal cells, and microglia; cerebral edema, raised intracranial pressure and herniation, and hydrocephalus; malformations and developmental diseases, such as neural tube defects, forebrain anomalies, posterior fossa anomalies, and syringomyelia and hydromyelia; perinatal brain injury; cerebrovascular diseases, such as those related to hypoxia, ischemia, and infarction, including hypotension, hypoperfusion, and low-flow states—global cerebral ischemia and focal cerebral ischemia—infarction from obstruction of local blood supply, intracranial hemorrhage, including intracerebral (intraparenchymal) hemorrhage, subarachnoid hemorrhage and ruptured berry aneurysms, and vascular malformations, hypertensive cerebrovascular disease, including lacunar infarcts, slit hemorrhages, and hypertensive encephalopathy; infections, such as acute meningitis, including acute pyogenic (bacterial) meningitis and acute aseptic (viral) meningitis, acute focal suppurative infections, including brain abscess, subdural empyema, and extradural abscess, chronic bacterial meningoencephalitis, including tuberculosis and mycobacterioses, neurosyphilis, and neuroborreliosis (Lyme disease), viral meningoencephalitis, including arthropod-borne (Arbo) viral encephalitis, *Herpes simplex* virus Type 1, *Herpes simplex* virus Type 2, *Vari-cella-zoster* virus (*Herpes zoster*), cytomegalovirus, poliomyelitis, rabies, and human immunodeficiency virus 1, including HIV-1 meningoencephalitis (subacute encephalitis), vacuolar myelopathy, AIDS-associated myopathy, peripheral neuropathy, and AIDS in children, progressive multifocal leukoencephalopathy, subacute sclerosing panencephalitis, fungal meningoencephalitis, other infectious diseases of the nervous system; transmissible spongiform encephalopathies (prion diseases); demyelinating diseases, including multiple sclerosis, multiple sclerosis variants, acute disseminated encephalomyelitis and acute necrotizing hemorrhagic encephalomyelitis, and other diseases with demyelination; degenerative diseases, such as degenerative diseases affecting the cerebral cortex, including Alzheimer disease and Pick disease, degenerative diseases of basal ganglia and brain stem, including Parkinsonism, idiopathic Parkinson disease (paralysis agitans), progressive supranuclear palsy, corticobasal degeneration, multiple system atrophy, including striatonigral degeneration, Shy-Drager syndrome, and olivopontocerebellar atrophy, and Huntington disease; spinocerebellar degenerations, including spinocerebellar ataxias, including Friedreich ataxia, and ataxia-telanglectasia, degenerative diseases affecting motor neurons, including amyotrophic lateral sclerosis (motor neuron disease), bulbospinal atrophy (Kennedy syndrome), and spinal muscular atrophy; inborn errors of metabolism, such as leukodystrophies, including Krabbe disease, metachromatic leukodystrophy, adrenoleukodystrophy, Pelizaeus-Merzbacher disease, and Canavan disease, mitochondrial encephalomyopathies, including Leigh disease and other mitochondrial encephalomyopathies; toxic and acquired metabolic diseases, including vitamin deficiencies such as thiamine (vitamin B₁) deficiency and vitamin B₁₂ deficiency, neurologic sequelae of metabolic disturbances, including hypoglycemia, hyperglycemia, and hepatic encephalopathy, toxic disorders, including carbon monoxide, methanol, ethanol, and radiation, including combined methotrexate and radiation-induced injury; tumors, such as gliomas, including

astrocytoma, including fibrillary (diffuse) astrocytoma and glioblastoma multiforme, pilocytic astrocytoma, pleomorphic xanthoastrocytoma, and brain stem glioma, oligodendroglioma, and ependymoma and related paraventricular mass lesions, neuronal tumors, poorly differentiated neoplasms, including medulloblastoma, other parenchymal tumors, including primary brain lymphoma, germ cell tumors, and pineal parenchymal tumors, meningiomas, metastatic tumors, paraneoplastic syndromes, peripheral nerve sheath tumors, including schwannoma, neurofibroma, and malignant peripheral nerve sheath tumor (malignant schwannoma), and neurocutaneous syndromes (phakomatoses), including neurofibromatosis, including Type 1 neurofibromatosis (NF1) and TYPE 2 neurofibromatosis (NF2), tuberous sclerosis, and Von Hippel-Lindau disease.

[0077] Disorders involving blood vessels include, but are not limited to, responses of vascular cell walls to injury, such as endothelial dysfunction and endothelial activation and intimal thickening; vascular diseases including, but not limited to, congenital anomalies, such as arteriovenous fistula, atherosclerosis, and hypertensive vascular disease, such as hypertension; inflammatory disease—the vasculitides, such as giant cell (temporal) arteritis, Takayasu arteritis, polyarteritis nodosa (classic), Kawasaki syndrome (mucocutaneous lymph node syndrome), microscopic polyangiitis (microscopic polyarteritis, hypersensitivity or leukocytoclastic angitis), Wegener granulomatosis, thromboangitis obliterans (Buerger disease), vasculitis associated with other disorders, and infectious arteritis; Raynaud disease; aneurysms and dissection, such as abdominal aortic aneurysms, syphilitic (luetic) aneurysms, and aortic dissection (dissecting hematoma); disorders of veins and lymphatics, such as varicose veins, thrombophlebitis and phlebothrombosis, obstruction of superior vena cava (superior vena cava syndrome), obstruction of inferior vena cava (inferior vena cava syndrome), and lymphangitis and lymphedema; tumors, including benign tumors and tumor-like conditions, such as hemangioma, lymphangioma, glomus tumor (glomangioma), vascular ectasias, and bacillary angiomatosis, and intermediate-grade (borderline low-grade malignant) tumors, such as Kaposi sarcoma and hemangioendothelioma, and malignant tumors, such as angiosarcoma and hemangiopericytoma; and pathology of therapeutic interventions in vascular disease, such as balloon angioplasty and related techniques and vascular replacement, such as coronary artery bypass graft surgery.

[0078] Examples of hematopoietic disorders include, but are not limited to, autoimmune diseases (including, for example, diabetes mellitus, arthritis (including rheumatoid arthritis, juvenile rheumatoid arthritis, osteoarthritis, psoriatic arthritis), multiple sclerosis, encephalomyelitis, myasthenia gravis, systemic lupus erythematosus, autoimmune thyroiditis, dermatitis (including atopic dermatitis and eczematous dermatitis), psoriasis, Sjögren's Syndrome, Crohn's disease, aphthous ulcer, iritis, conjunctivitis, keratoconjunctivitis, ulcerative colitis, asthma, allergic asthma, cutaneous lupus erythematosus, scleroderma, vaginitis, proctitis, drug eruptions, leprosy reversal reactions, erythema nodosum leprosum, autoimmune uveitis, allergic encephalomyelitis, acute necrotizing hemorrhagic encephalopathy, idiopathic bilateral progressive sensorineural hearing loss, aplastic anemia, pure red cell anemia, idiopathic thrombocytopenia, polychondritis, Wegener's granulomatosis, chronic active hepatitis, Stevens-Johnson syndrome,

idiopathic sprue, lichen planus, Graves' disease, sarcoidosis, primary biliary cirrhosis, uveitis posterior, and interstitial lung fibrosis), graft-versus-host disease, cases of transplantation, and allergy such as, atopic allergy.

[0079] The 47324 protein, fragments thereof, and derivatives and other variants of the sequence in SEQ ID NO:2 are collectively referred to as "polypeptides or proteins of the invention" or "47324 polypeptides or proteins". Nucleic acid molecules encoding such polypeptides or proteins are collectively referred to as "nucleic acids of the invention" or "47324 nucleic acids." 47324 molecules refer to 47324 nucleic acids, polypeptides, and antibodies.

[0080] As used herein, the term "nucleic acid molecule" includes DNA molecules (e.g., a cDNA or genomic DNA) and RNA molecules (e.g., an mRNA) and analogs of the DNA or RNA generated, e.g., by the use of nucleotide analogs. The nucleic acid molecule can be single-stranded or double-stranded, but preferably is double-stranded DNA.

[0081] The term "isolated or purified nucleic acid molecule" includes nucleic acid molecules which are separated from other nucleic acid molecules which are present in the natural source of the nucleic acid. For example, with regards to genomic DNA, the term "isolated" includes nucleic acid molecules which are separated from the chromosome with which the genomic DNA is naturally associated. Preferably, an "isolated" nucleic acid is free of sequences which naturally flank the nucleic acid (i.e., sequences located at the 5' and/or 3' ends of the nucleic acid) in the genomic DNA of the organism from which the nucleic acid is derived. For example, in various embodiments, the isolated nucleic acid molecule can contain less than about 5 kb, 4kb, 3kb, 2kb, 1 kb, 0.5 kb or 0.1 kb of 5' and/or 3' nucleotide sequences which naturally flank the nucleic acid molecule in genomic DNA of the cell from which the nucleic acid is derived. Moreover, an "isolated" nucleic acid molecule, such as a cDNA molecule, can be substantially free of other cellular material, or culture medium when produced by recombinant techniques, or substantially free of chemical precursors or other chemicals when chemically synthesized.

[0082] As used herein, the term "hybridizes under stringent conditions" describes conditions for hybridization and washing. Stringent conditions are known to those skilled in the art and can be found in *Current Protocols in Molecular Biology*, John Wiley & Sons, N.Y. (1989), 6.3.1-6.3.6. Aqueous and nonaqueous methods are described in that reference and either can be used. A preferred, example of stringent hybridization conditions are hybridization in 6x sodium chloride/sodium citrate (SSC) at about 45° C., followed by one or more washes in 0.2xSSC, 0.1% SDS at 50° C. Another example of stringent hybridization conditions are hybridization in 6x sodium chloride/sodium citrate (SSC) at about 45° C., followed by one or more washes in 0.2xSSC, 0.1% SDS at 55° C. A further example of stringent hybridization conditions are hybridization in 6x sodium chloride/sodium citrate (SSC) at about 45° C., followed by one or more washes in 0.2xSSC, 0.1% SDS at 60° C. Preferably, stringent hybridization conditions are hybridization in 6x sodium chloride/sodium citrate (SSC) at about 45° C., followed by one or more washes in 0.2xSSC, 0.1% SDS at 65° C. Particularly preferred stringency conditions (and the conditions that should be used if the practitioner is uncertain about what conditions should be applied to determine if a

molecule is within a hybridization limitation of the invention) are 0.5M Sodium Phosphate, 7% SDS at 65° C., followed by one or more washes at 0.2×SSC, 1% SDS at 65° C. Preferably, an isolated nucleic acid molecule of the invention that hybridizes under stringent conditions to the sequence of SEQ ID NO:1, or SEQ ID NO:3, corresponds to a naturally-occurring nucleic acid molecule.

[0083] As used herein, a “naturally-occurring” nucleic acid molecule refers to an RNA or DNA molecule having a nucleotide sequence that occurs in nature (e.g., encodes a natural protein).

[0084] As used herein, the terms “gene” and “recombinant gene” refer to nucleic acid molecules which include an open reading frame encoding a 47324 protein, preferably a mammalian 47324 protein, and can further include non-coding regulatory sequences, and introns.

[0085] An “isolated” or “purified” polypeptide or protein is substantially free of cellular material or other contaminating proteins from the cell or tissue source from which the protein is derived, or substantially free from chemical precursors or other chemicals when chemically synthesized. In one embodiment, the language “substantially free” means preparation of 47324 protein having less than about 30%, 20%, 10% and more preferably 5% (by dry weight), of non-47324 protein (also referred to herein as a “contaminating protein”), or of chemical precursors or non-47324 chemicals. When the 47324 protein or biologically active portion thereof is recombinantly produced, it is also preferably substantially free of culture medium, i.e., culture medium represents less than about 20%, more preferably less than about 10%, and most preferably less than about 5% of the volume of the protein preparation. The invention includes isolated or purified preparations of at least 0.01, 0.1, 1.0, and 10 milligrams in dry weight.

[0086] A “non-essential” amino acid residue is a residue that can be altered from the wild-type sequence of 47324 (e.g., the sequence of SEQ ID NO:1, SEQ ID NO:3, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____) without abolishing or more preferably, without substantially altering a biological activity, whereas an “essential” amino acid residue results in such a change. For example, amino acid residues that are conserved among the polypeptides of the present invention, e.g., those present in the Ras family domain, are predicted to be particularly unamenable to alteration.

[0087] A “conservative amino acid substitution” is one in which the amino acid residue is replaced with an amino acid residue having a similar side chain. Families of amino acid residues having similar side chains have been defined in the art. These families include amino acids with basic side chains (e.g., lysine, arginine, histidine), acidic side chains (e.g., aspartic acid, glutamic acid), uncharged polar side chains (e.g., glycine, asparagine, glutamine, serine, threonine, tyrosine, cysteine), nonpolar side chains (e.g., alanine, valine, leucine, isoleucine, proline, phenylalanine, methionine, tryptophan), beta-branched side chains (e.g., threonine, valine, isoleucine) and aromatic side chains (e.g., tyrosine, phenylalanine, tryptophan, histidine). Thus, a predicted non-essential amino acid residue in a 47324 protein is preferably replaced with another amino acid residue from the same side chain family. Alternatively, in another embodiment, muta-

tions can be introduced randomly along all or part of a 47324 coding sequence, such as by saturation mutagenesis, and the resultant mutants can be screened for 47324 biological activity to identify mutants that retain activity. Following mutagenesis of SEQ ID NO:1, SEQ ID NO:3, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____, the encoded protein can be expressed recombinantly and the activity of the protein can be determined.

[0088] As used herein, a “biologically active portion” of a 47324 protein includes a fragment of a 47324 protein which participates in an interaction between a 47324 molecule and a non-47324 molecule. Biologically active portions of a 47324 protein include peptides comprising amino acid sequences sufficiently homologous to or derived from the amino acid sequence of the 47324 protein, e.g., the amino acid sequence shown in SEQ ID NO:2, which include less amino acids than the full length 47324 proteins, and exhibit at least one activity of a 47324 protein. Typically, biologically active portions comprise a domain or motif with at least one activity of the 47324 protein, e.g., Ras family activity. A biologically active portion of a 47324 protein can be a polypeptide which is, for example, 10, 25, 50, 100, 200 or more amino acids in length. Biologically active portions of a 47324 protein can be used as targets for developing agents which modulate a 47324 mediated activity, e.g., Ras family activity.

[0089] Calculations of homology or sequence identity between sequences (the terms are used interchangeably herein) are performed as follows.

[0090] To determine the percent identity of two amino acid sequences, or of two nucleic acid sequences, the sequences are aligned for optimal comparison purposes (e.g., gaps can be introduced in one or both of a first and a second amino acid or nucleic acid sequence for optimal alignment and non-homologous sequences can be disregarded for comparison purposes). In a preferred embodiment, the length of a reference sequence aligned for comparison purposes is at least 30%, preferably at least 40%, more preferably at least 50%, even more preferably at least 60%, and even more preferably at least 70%, 80%, 90%, 100% of the length of the reference sequence (e.g., when aligning a second sequence to the 47324 amino acid sequence of SEQ ID NO:2 having 236 amino acid residues, at least 71, preferably at least 94, more preferably at least 118, even more preferably at least 142, and even more preferably at least 165, 189, 212 or 236 amino acid residues are aligned). The amino acid residues or nucleotides at corresponding amino acid positions or nucleotide positions are then compared. When a position in the first sequence is occupied by the same amino acid residue or nucleotide as the corresponding position in the second sequence, then the molecules are identical at that position (as used herein amino acid or nucleic acid “identity” is equivalent to amino acid or nucleic acid “homology”). The percent identity between the two sequences is a function of the number of identical positions shared by the sequences, taking into account the number of gaps, and the length of each gap, which need to be introduced for optimal alignment of the two sequences.

[0091] The comparison of sequences and determination of percent identity between two sequences can be accomplished using a mathematical algorithm. In a preferred

embodiment, the percent identity between two amino acid sequences is determined using the Needleman and Wunsch (*J. Mol. Biol.* (48):444-453 (1970)) algorithm which has been incorporated into the GAP program in the GCG software package (available at <http://www.gcg.com>), using either a Blossum 62 matrix or a PAM250 matrix, and a gap weight of 16, 14, 12, 10, 8, 6, or 4 and a length weight of 1, 2, 3, 4, 5, or 6. In yet another preferred embodiment, the percent identity between two nucleotide sequences is determined using the GAP program in the GCG software package (available at <http://www.gcg.com>), using a NWSgapdna.CMP matrix and a gap weight of 40, 50, 60, 70, or 80 and a length weight of 1, 2, 3, 4, 5, or 6. A particularly preferred set of parameters (and the one that should be used if the practitioner is uncertain about what parameters should be applied to determine if a molecule is within a sequence identity or homology limitation of the invention) is using a Blossum 62 scoring matrix with a gap open penalty of 12, a gap extend penalty of 4, and a frameshift gap penalty of 5.

[0092] The percent identity between two amino acid or nucleotide sequences can be determined using the algorithm of E. Meyers and W. Miller (CABIOS, 4:11-17 (1989)) which has been incorporated into the ALIGN program (version 2.0), using a PAM120 weight residue table, a gap length penalty of 12 and a gap penalty of 4.

[0093] The nucleic acid and protein sequences described herein can be used as a "query sequence" to perform a search against public databases to, for example, identify other family members or related sequences. Such searches can be performed using the NBLAST and XBLAST programs (version 2.0) of Altschul, et al., (1990) *J. Mol. Biol.* 215:403-10. BLAST nucleotide searches can be performed with the NBLAST program, score=100, wordlength=12 to obtain nucleotide sequences homologous to 47324 nucleic acid molecules of the invention. BLAST protein searches can be performed with the XBLAST program, score=50, wordlength=3 to obtain amino acid sequences homologous to 47324 protein molecules of the invention. To obtain gapped alignments for comparison purposes, Gapped BLAST can be utilized as described in Altschul et al., (1997) *Nucleic Acids Res.* 25(17):3389-3402. When utilizing BLAST and Gapped BLAST programs, the default parameters of the respective programs (e.g., XBLAST and NBLAST) can be used. See <http://www.ncbi.nlm.nih.gov>.

[0094] Particular 47324 polypeptides of the present invention have an amino acid sequence substantially identical to the amino acid sequence of SEQ ID NO:2. In the context of an amino acid sequence, the term "substantially identical" is used herein to refer to a first amino acid that contains a sufficient or minimum number of amino acid residues that are i) identical to, or ii) conservative substitutions of aligned amino acid residues in a second amino acid sequence such that the first and second amino acid sequences can have a common structural domain and/or common functional activity. For example, amino acid sequences that contain a common structural domain having at least about 60%, or 65% identity, likely 75% identity, more likely 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identity ID NO:2 are termed substantially identical.

[0095] In the context of nucleotide sequence, the term "substantially identical" is used herein to refer to a first nucleic acid sequence that contains a sufficient or minimum

number of nucleotides that are identical to aligned nucleotides in a second nucleic acid sequence such that the first and second nucleotide sequences encode a polypeptide having common functional activity, or encode a common structural polypeptide domain or a common functional polypeptide activity. For example, nucleotide sequences having at least about 60%, or 65% identity, likely 75% identity, more likely 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identity to SEQ ID NO:1 or 3 are termed substantially identical.

[0096] "Misexpression or aberrant expression", as used herein, refers to a non-wild type pattern of gene expression, at the RNA or protein level. It includes: expression at non-wild type levels, i.e., over or under expression; a pattern of expression that differs from wild type in terms of the time or stage at which the gene is expressed, e.g., increased or decreased expression (as compared with wild type) at a predetermined developmental period or stage; a pattern of expression that differs from wild type in terms of decreased expression (as compared with wild type) in a predetermined cell type or tissue type; a pattern of expression that differs from wild type in terms of the splicing size, amino acid sequence, post-translational modification, or biological activity of the expressed polypeptide; a pattern of expression that differs from wild type in terms of the effect of an environmental stimulus or extracellular stimulus on expression of the gene, e.g., a pattern of increased or decreased expression (as compared with wild type) in the presence of an increase or decrease in the strength of the stimulus.

[0097] "Subject", as used herein, can refer to a mammal, e.g., a human, or to an experimental or animal or disease model. The subject can also be a non-human animal, e.g., a horse, cow, goat, or other domestic animal.

[0098] A "purified preparation of cells", as used herein, refers to, in the case of plant or animal cells, an in vitro preparation of cells and not an entire intact plant or animal. In the case of cultured cells or microbial cells, it consists of a preparation of at least 10% and more preferably 50% of the subject cells.

[0099] Various aspects of the invention are described in further detail below.

[0100] Isolated Nucleic Acid Molecules

[0101] In one aspect, the invention provides, an isolated or purified, nucleic acid molecule that encodes a 47324 polypeptide described herein, e.g., a full length 47324 protein or a fragment thereof, e.g., a biologically active portion of 47324 protein. Also included is a nucleic acid fragment suitable for use as a hybridization probe, which can be used, e.g., to identify nucleic acid molecule encoding a polypeptide of the invention, 47324 mRNA, and fragments suitable for use as primers, e.g., PCR primers for the amplification or mutation of nucleic acid molecules.

[0102] In one embodiment, an isolated nucleic acid molecule of the invention includes the nucleotide sequence shown in SEQ ID NO:1, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____, or a portion of any of these nucleotide sequences. In one embodiment, the nucleic acid molecule includes sequences encoding the human 47324 protein (i.e., "the coding region", from nucleotides 19-729 of SEQ ID NO:1, including the terminal codon), as well as 5' untrans-

lated sequences (nucleotides 1-18 of SEQ ID NO:1). Alternatively, the nucleic acid molecule can include only the coding region of SEQ ID NO:1 (e.g., nucleotides 19-729 of SEQ ID NO:1, corresponding to SEQ ID NO:3) and, e.g., no flanking sequences which normally accompany the subject sequence. In another embodiment, the nucleic acid molecule encodes a sequence corresponding to the mature protein of SEQ ID NO:2.

[0103] In another embodiment, an isolated nucleic acid molecule of the invention includes a nucleic acid molecule which is a complement of the nucleotide sequence shown in SEQ ID NO:1, SEQ ID NO:3, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____, or a portion of any of these nucleotide sequences. In other embodiments, the nucleic acid molecule of the invention is sufficiently complementary to the nucleotide sequence shown in SEQ ID NO:1, SEQ ID NO:3, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____ such that it can hybridize to the nucleotide sequence shown in SEQ ID NO:1, SEQ ID NO:3, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____, thereby forming a stable duplex.

[0104] In one embodiment, an isolated nucleic acid molecule of the present invention includes a nucleotide sequence which is at least about 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or more homologous to the nucleotide sequence shown in SEQ ID NO:1, SEQ ID NO:3, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____. In the case of an isolated nucleic acid molecule which is longer than or equivalent in length to the reference sequence, e.g., SEQ ID NO:1, or SEQ ID NO:3, the comparison is made with the full length of the reference sequence. Where the isolated nucleic acid molecule is shorter than the reference sequence, e.g., shorter than SEQ ID NO:1, or SEQ ID NO:3, the comparison is made to a segment of the reference sequence of the same length (excluding any loop required by the homology calculation).

[0105] 47324 Nucleic Acid Fragments

[0106] A nucleic acid molecule of the invention can include only a portion of the nucleic acid sequence of SEQ ID NO:1, SEQ ID NO:3, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____. For example, such a nucleic acid molecule can include a fragment which can be used as a probe or primer or a fragment encoding a portion of a 47324 protein, e.g., an immunogenic or biologically active portion of a 47324 protein. A fragment can comprise: nucleotides 40-711 of SEQ ID NO:1, which encodes a Ras family domain of human 47324. The nucleotide sequence determined from the cloning of the 47324 gene allows for the generation of probes and primers designed for use in identifying and/or cloning other 47324 family members, or fragments thereof, as well as 47324 homologues, or fragments thereof, from other species.

[0107] In another embodiment, a nucleic acid includes a nucleotide sequence that includes part, or all, of the coding region and extends into either (or both) the 5' or 3' noncoding region. Other embodiments include a fragment which includes a nucleotide sequence encoding an amino acid

fragment described herein. Nucleic acid fragments can encode a specific domain or site described herein or fragments thereof, particularly fragments thereof which are at least 150 amino acids in length. Fragments also include nucleic acid sequences corresponding to specific amino acid sequences described above or fragments thereof. Nucleic acid fragments should not be construed as encompassing those fragments that may have been disclosed prior to the invention.

[0108] A nucleic acid fragment can include a sequence corresponding to a domain, region, or functional site described herein. A nucleic acid fragment can also include one or more domain, region, or functional site described herein. Thus, for example, the nucleic acid fragment can include a Ras family domain. In a preferred embodiment the fragment is at least, 50, 100, 200, 300, 400, 500, 600, 700, or 900 base pairs in length.

[0109] 47324 probes and primers are provided. Typically a probe/primer is an isolated or purified oligonucleotide. The oligonucleotide typically includes a region of nucleotide sequence that hybridizes under stringent conditions to at least about 7, 12 or 15, preferably about 20 or 25, more preferably about 30, 35, 40, 45, 50, 55, 60, 65, or 75 consecutive nucleotides of a sense or antisense sequence of SEQ ID NO:1, SEQ ID NO:3, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____, or of a naturally occurring allelic variant or mutant of SEQ ID NO:1, SEQ ID NO:3, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____.

[0110] In a preferred embodiment the nucleic acid is a probe which is at least 5 or 10, and less than 200, more preferably less than 100, or less than 50, base pairs in length. It should be identical, or differ by 1, or less than in 5 or 10 bases, from a sequence disclosed herein. If alignment is needed for this comparison the sequences should be aligned for maximum homology. "Looped" out sequences from deletions or insertions, or mismatches, are considered differences.

[0111] A probe or primer can be derived from the sense or anti-sense strand of a nucleic acid which encodes a Ras family domain (e.g., about nucleotides 40-711 of SEQ ID NO:1).

[0112] In another embodiment a set of primers is provided, e.g., primers suitable for use in a PCR, which can be used to amplify a selected region of a 47324 sequence, e.g., a region described herein. The primers should be at least 5, 10, or 50 base pairs in length and less than 100, or less than 200, base pairs in length. The primers should be identical, or differs by one base from a sequence disclosed herein or from a naturally occurring variant. E.g., primers suitable for amplifying all or a portion of any of the following regions are provided: a Ras family domain (e.g., about nucleotides 40-711 of SEQ ID NO:1).

[0113] A nucleic acid fragment can encode an epitope bearing region of a polypeptide described herein.

[0114] A nucleic acid fragment encoding a "biologically active portion of a 47324 polypeptide" can be prepared by isolating a portion of the nucleotide sequence of SEQ ID NO:1, SEQ ID NO:3, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession

Number _____, which encodes a polypeptide having a 47324 biological activity (e.g., the biological activities of the 47324 proteins as described herein), expressing the encoded portion of the 47324 protein (e.g., by recombinant expression in vitro) and assessing the activity of the encoded portion of the 47324 protein. For example, a nucleic acid fragment encoding a biologically active portion of 47324 includes a Ras family domain (e.g., about nucleotides 40-711 of SEQ ID NO:1). A nucleic acid fragment encoding a biologically active portion of a 47324 polypeptide, may comprise a nucleotide sequence which is greater than 300-1200 or more nucleotides in length.

[0115] In preferred embodiments, nucleic acids include a nucleotide sequence which is about 300, 400, 500, 600, 700, 800, 900, 1000, 1100, 1200, 1300, 1400 nucleotides in length and hybridizes under stringent hybridization conditions to a nucleic acid molecule of SEQ ID NO:1, or SEQ ID NO:3, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____.

[0116] 47324 Nucleic Acid Variants

[0117] The invention further encompasses nucleic acid molecules that differ from the nucleotide sequence shown in SEQ ID NO:1, SEQ ID NO:3, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____. Such differences can be due to degeneracy of the genetic code (and result in a nucleic acid which encodes the same 47324 proteins as those encoded by the nucleotide sequence disclosed herein. In another embodiment, an isolated nucleic acid molecule of the invention has a nucleotide sequence encoding a protein having an amino acid sequence which differs, by at least 1, but less than 5, 10, 20, 50, or 100 amino acid residues that shown in SEQ ID NO:2. If alignment is needed for this comparison the sequences should be aligned for maximum homology. "Looped" out sequences from deletions or insertions, or mismatches, are considered differences.

[0118] Nucleic acids of the inventor can be chosen for having codons, which are preferred, or non preferred, for a particular expression system. E.g., the nucleic acid can be one in which at least one codon, at preferably at least 10%, or 20% of the codons has been altered such that the sequence is optimized for expression in *E. coli*, yeast, human, insect, or CHO cells.

[0119] Nucleic acid variants can be naturally occurring, such as allelic variants (same locus), homologs (different locus), and orthologs (different organism) or can be non-naturally occurring. Non-naturally occurring variants can be made by mutagenesis techniques, including those applied to polynucleotides, cells, or organisms. The variants can contain nucleotide substitutions, deletions, inversions and insertions. Variation can occur in either or both the coding and non-coding regions. The variations can produce both conservative and non-conservative amino acid substitutions (as compared in the encoded product).

[0120] In a preferred embodiment, the nucleic acid differs from that of SEQ ID NO:1, SEQ ID NO:3, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____, e.g., as follows: by at least one but less than 10, 20, 30, or 40 nucleotides; at least one but less than 1%, 5%, 10% or 20% of the in the subject

nucleic acid. If necessary for this analysis the sequences should be aligned for maximum homology. "Looped" out sequences from deletions or insertions, or mismatches, are considered differences.

[0121] Orthologs, homologs, and allelic variants can be identified using methods known in the art. These variants comprise a nucleotide sequence encoding a polypeptide that is 50%, at least about 55%, typically at least about 70-75%, more typically at least about 80-85%, and most typically at least about 90-95% or more identical to the amino acid sequence shown in SEQ ID NO:2 or a fragment of this sequence. Such nucleic acid molecules can readily be obtained as being able to hybridize under stringent conditions, to the nucleotide sequence shown in SEQ ID NO:3 or a fragment of this sequence. Nucleic acid molecules corresponding to orthologs, homologs, and allelic variants of the 47324 cDNAs of the invention can further be isolated by mapping to the same chromosome or locus as the 47324 gene. Preferred variants include those that are correlated with Ras family activity.

[0122] Allelic variants of 47324, e.g., human 47324, include both functional and non-functional proteins. Functional allelic variants are naturally occurring amino acid sequence variants of the 47324 protein within a population that maintain the ability to modulate the phosphorylation state of itself or another protein or polypeptide. Functional allelic variants will typically contain only conservative substitution of one or more amino acids of SEQ ID NO:2, or substitution, deletion or insertion of non-critical residues in non-critical regions of the protein. Non-functional allelic variants are naturally-occurring amino acid sequence variants of the 47324, e.g., human 47324, protein within a population that do not have the ability to activate signal transduction. Non-functional allelic variants will typically contain a non-conservative substitution, a deletion, or insertion, or premature truncation of the amino acid sequence of SEQ ID NO:2, or a substitution, insertion, or deletion in critical residues or critical regions of the protein.

[0123] Moreover, nucleic acid molecules encoding other 47324 family members and, thus, which have a nucleotide sequence which differs from the 47324 sequences of SEQ ID NO:1, SEQ ID NO:3, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____ are intended to be within the scope of the invention.

[0124] Antisense Nucleic Acid Molecules, Ribozymes and Modified 47324 Nucleic Acid Molecules

[0125] In another aspect, the invention features, an isolated nucleic acid molecule which is antisense to 47324. An "antisense" nucleic acid can include a nucleotide sequence which is complementary to a "sense" nucleic acid encoding a protein, e.g., complementary to the coding strand of a double-stranded cDNA molecule or complementary to an mRNA sequence. The antisense nucleic acid can be complementary to an entire 47324 coding strand, or to only a portion thereof (e.g., the coding region of human 47324 corresponding to SEQ ID NO:3). In another embodiment, the antisense nucleic acid molecule is antisense to a "non-coding region" of the coding strand of the nucleotide sequence encoding 47324 (e.g., the 5' and 3' untranslated regions).

[0126] An antisense nucleic acid can be designed such that it is complementary to the entire coding region of 47324

mRNA, but more preferably is an oligonucleotide which is antisense to only a portion of the coding or noncoding region of 47324 mRNA. For example, the antisense oligonucleotide can be complementary to the region surrounding the translation start site of 47324 mRNA, e.g., between the -10 and +10 regions of the target gene nucleotide sequence of interest. An antisense oligonucleotide can be, for example, about 7, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, or more nucleotides in length.

[0127] An antisense nucleic acid of the invention can be constructed using chemical synthesis and enzymatic ligation reactions using procedures known in the art. For example, an antisense nucleic acid (e.g., an antisense oligonucleotide) can be chemically synthesized using naturally occurring nucleotides or variously modified nucleotides designed to increase the biological stability of the molecules or to increase the physical stability of the duplex formed between the antisense and sense nucleic acids, e.g., phosphorothioate derivatives and acridine substituted nucleotides can be used. The antisense nucleic acid also can be produced biologically using an expression vector into which a nucleic acid has been subcloned in an antisense orientation (i.e., RNA transcribed from the inserted nucleic acid will be of an antisense orientation to a target nucleic acid of interest, described further in the following subsection).

[0128] The antisense nucleic acid molecules of the invention are typically administered to a subject (e.g., by direct injection at a tissue site), or generated in situ such that they hybridize with or bind to cellular mRNA and/or genomic DNA encoding a 47324 protein to thereby inhibit expression of the protein, e.g., by inhibiting transcription and/or translation. Alternatively, antisense nucleic acid molecules can be modified to target selected cells and then administered systemically. For systemic administration, antisense molecules can be modified such that they specifically bind to receptors or antigens expressed on a selected cell surface, e.g., by linking the antisense nucleic acid molecules to peptides or antibodies which bind to cell surface receptors or antigens. The antisense nucleic acid molecules can also be delivered to cells using the vectors described herein. To achieve sufficient intracellular concentrations of the antisense molecules, vector constructs in which the antisense nucleic acid molecule is placed under the control of a strong pol II or pol III promoter are preferred.

[0129] In yet another embodiment, the antisense nucleic acid molecule of the invention is an α -anomeric nucleic acid molecule. An α -anomeric nucleic acid molecule forms specific double-stranded hybrids with complementary RNA in which, contrary to the usual, β -units, the strands run parallel to each other (Gaultier et al., (1987) *Nucleic Acids Res.* 15:6625-6641). The antisense nucleic acid molecule can also comprise a 2'-o-methylribonucleotide (Inoue et al., (1987) *Nucleic Acids Res.* 15:6131-6148) or a chimeric RNA-DNA analogue (Inoue et al., (1987) *FEBS Lett.* 215:327-330).

[0130] In still another embodiment, an antisense nucleic acid of the invention is a ribozyme. A ribozyme having specificity for a 47324-encoding nucleic acid can include one or more sequences complementary to the nucleotide sequence of a 47324 cDNA disclosed herein (i.e., SEQ ID NO:1, or SEQ ID NO:3), and a sequence having known catalytic sequence responsible for mRNA cleavage (see U.S.

Pat. No. 5,093,246 or Haselhoff and Gerlach, (1988) *Nature* 334:585-591). For example, a derivative of a *Tetrahymena* L-19 IVS RNA can be constructed in which the nucleotide sequence of the active site is complementary to the nucleotide sequence to be cleaved in a 47324-encoding mRNA. See, e.g., Cech et al. U.S. Pat. No. 4,987,071; and Cech et al. U.S. Pat. No. 5,116,742. Alternatively, 47324 mRNA can be used to select a catalytic RNA having a specific ribonuclease activity from a pool of RNA molecules. See, e.g., Bartel, D. and Szostak, J. W. (1993) *Science* 261:1411-1418.

[0131] 47324 gene expression can be inhibited by targeting nucleotide sequences complementary to the regulatory region of the 47324 (e.g., the 47324 promoter and/or enhancers) to form triple helical structures that prevent transcription of the 47324 gene in target cells. See generally, Helene, C., (1991) *Anticancer Drug Des.* 6(6):569-84; Helene, C. et al., (1992) *Ann. N.Y. Acad. Sci.* 660:27-36; and Maher, L. J., (1992) *Bioassays* 14(12):807-15. The potential sequences that can be targeted for triple helix formation can be increased by creating a so-called "switchback" nucleic acid molecule. Switchback molecules are synthesized in an alternating 5'-3', 3'-5' manner, such that they base pair with first one strand of a duplex and then the other, eliminating the necessity for a sizeable stretch of either purines or pyrimidines to be present on one strand of a duplex.

[0132] The invention also provides detectably labeled oligonucleotide primer and probe molecules. Typically, such labels are chemiluminescent, fluorescent, radioactive, or colorimetric.

[0133] A 47324 nucleic acid molecule can be modified at the base moiety, sugar moiety or phosphate backbone to improve, e.g., the stability, hybridization, or solubility of the molecule. For example, the deoxyribose phosphate backbone of the nucleic acid molecules can be modified to generate peptide nucleic acids (see Hyrup B. et al., (1996) *Bioorganic & Medicinal Chemistry* 4 (1): 5-23). As used herein, the terms "peptide nucleic acid" or "PNA" refers to a nucleic acid mimic, e.g., a DNA mimic, in which the deoxyribose phosphate backbone is replaced by a pseudopeptide backbone and only the four natural nucleobases are retained. The neutral backbone of a PNA can allow for specific hybridization to DNA and RNA under conditions of low ionic strength. The synthesis of PNA oligomers can be performed using standard solid phase peptide synthesis protocols as described in Hyrup B. et al., (1996) supra; Perry-O'Keefe et al., *Proc. Natl. Acad. Sci.* 93: 14670-675.

[0134] PNAs of 47324 nucleic acid molecules can be used in therapeutic and diagnostic applications. For example, PNAs can be used as antisense or antigene agents for sequence-specific modulation of gene expression by, for example, inducing transcription or translation arrest or inhibiting replication. PNAs of 47324 nucleic acid molecules can also be used in the analysis of single base pair mutations in a gene, (e.g., by PNA-directed PCR clamping); as 'artificial restriction enzymes' when used in combination with other enzymes, (e.g., S1 nucleases (Hyrup B., (1996) supra)); or as probes or primers for DNA sequencing or hybridization (Hyrup B. et al., (1996) supra; Perry-O'Keefe supra).

[0135] In other embodiments, the oligonucleotide may include other appended groups such as peptides (e.g., for targeting host cell receptors in vivo), or agents facilitating

transport across the cell membrane (see, e.g., Letsinger et al., (1989) *Proc. Natl. Acad. Sci. USA* 86:6553-6556; Lemaitre et al., (1987) *Proc. Natl. Acad. Sci. USA* 84:648-652; PCT Publication No. W088/09810) or the blood-brain barrier (see, e.g., PCT Publication No. W089/10134). In addition, oligonucleotides can be modified with hybridization-triggered cleavage agents (See, e.g., Krol et al., (1988) *Bio-Techniques* 6:958-976) or intercalating agents. (See, e.g., Zon, (1988) *Pharm. Res.* 5:539-549). To this end, the oligonucleotide may be conjugated to another molecule, (e.g., a peptide, hybridization triggered cross-linking agent, transport agent, or hybridization-triggered cleavage agent).

[0136] The invention also includes molecular beacon oligonucleotide primer and probe molecules having at least one region which is complementary to a 47324 nucleic acid of the invention, two complementary regions one having a fluorophore and one a quencher such that the molecular beacon is useful for quantitating the presence of the 47324 nucleic acid of the invention in a sample. Molecular beacon nucleic acids are described, for example, in Lizardi et al., U.S. Pat. No. 5,854,033; Nazarenko et al., U.S. Pat. No. 5,866,336, and Livak et al., U.S. Pat. No. 5,876,930.

[0137] Isolated 47324 Polypeptides

[0138] In another aspect, the invention features, an isolated 47324 protein, or fragment, e.g., a biologically active portion, for use as immunogens or antigens to raise or test (or more generally to bind) anti-47324 antibodies. 47324 protein can be isolated from cells or tissue sources using standard protein purification techniques. 47324 protein or fragments thereof can be produced by recombinant DNA techniques or synthesized chemically.

[0139] Polypeptides of the invention include those which arise as a result of the existence of multiple genes, alternative transcription events, alternative RNA splicing events, and alternative translational and postranslational events. The polypeptide can be expressed in systems, e.g., cultured cells, which result in substantially the same postranslational modifications present when expressed the polypeptide is expressed in a native cell, or in systems which result in the alteration or omission of postranslational modifications, e.g., glycosylation or cleavage, present when expressed in a native cell.

[0140] In a preferred embodiment, a 47324 polypeptide has one or more of the following characteristics:

[0141] (i) activation of a signal transduction pathway;

[0142] (ii) it has a molecular weight, e.g., a deduced molecular weight, amino acid composition or other physical characteristic of the polypeptide of SEQ ID NO:2;

[0143] (iii) it has an overall sequence similarity of at least 50%, preferably at least 60%, more preferably at least 70, 80, 90, or 95%, with a polypeptide of SEQ ID NO:2;

[0144] (iv) it has a Ras family domain which preferably has an overall sequence similarity of about 70%, 80%, 90% or 95% with amino acid residues 8-231 of SEQ ID NO:2;

[0145] (v) it has at least 70%, preferably 80%, and most preferably 95% of the cysteines found in the amino acid sequence of the native protein.

[0146] In a preferred embodiment the 47324 protein, or fragment thereof, differs from the corresponding sequence in SEQ ID NO:2. In one embodiment it differs by at least one but by less than 15, 10 or 5 amino acid residues. In another it differs from the corresponding sequence in SEQ ID NO:2 by at least one residue but less than 20%, 15%, 10% or 5% of the residues in it differ from the corresponding sequence in SEQ ID NO:2. (If this comparison requires alignment the sequences should be aligned for maximum homology. "Looped" out sequences from deletions or insertions, or mismatches, are considered differences.) The differences are, preferably, differences or changes at a non-essential residue or a conservative substitution. In a preferred embodiment the differences are not in the Ras family domain. In another embodiment one or more differences are in the Ras family domain.

[0147] Other embodiments include a protein that contain one or more changes in amino acid sequence, e.g., a change in an amino acid residue which is not essential for activity. Such 47324 proteins differ in amino acid sequence from SEQ ID NO:2, yet retain biological activity.

[0148] In one embodiment, a biologically active portion of a 47324 protein includes a Ras family domain. Moreover, other biologically active portions, in which other regions of the protein are deleted, can be prepared by recombinant techniques and evaluated for one or more of the functional activities of a native 47324 protein.

[0149] In a preferred embodiment, the 47324 protein has an amino acid sequence shown in SEQ ID NO:2. In other embodiments, the 47324 protein is substantially identical to SEQ ID NO:2. In yet another embodiment, the 47324 protein is substantially identical to SEQ ID NO:2 and retains the functional activity of the protein of SEQ ID NO:2, as described in detail above. Accordingly, in another embodiment, the 47324 protein is a protein which includes an amino acid sequence at least about 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 98% or more identical to SEQ ID NO:2.

[0150] 47324 Chimeric or Fusion Proteins

[0151] In another aspect, the invention provides 47324 chimeric or fusion proteins. As used herein, a 47324 "chimeric protein" or "fusion protein" includes a 47324 polypeptide linked to a non-47324 polypeptide. A "non-47324 polypeptide" refers to a polypeptide having an amino acid sequence corresponding to a protein which is not substantially homologous to the 47324 protein, e.g., a protein which is different from the 47324 protein and which is derived from the same or a different organism. The 47324 polypeptide of the fusion protein can correspond to all or a portion e.g., a fragment described herein of a 47324 amino acid sequence. In a preferred embodiment, a 47324 fusion protein includes at least one (or two) biologically active portion of a 47324 protein. The non-47324 polypeptide can be fused to the N-terminus or C-terminus of the 47324 polypeptide.

[0152] The fusion protein can include a moiety which has a high affinity for a ligand. For example, the fusion protein can be a GST-47324 fusion protein in which the 47324 sequences are fused to the C-terminus of the GST sequences. Such fusion proteins can facilitate the purification of recombinant 47324. Alternatively, the fusion protein can be a 47324 protein containing a heterologous signal sequence at

its N-terminus. In certain host cells (e.g., mammalian host cells), expression and/or secretion of 47324 can be increased through use of a heterologous signal sequence.

[0153] Fusion proteins can include all or a part of a serum protein, e.g., an IgG constant region, or human serum albumin.

[0154] The 47324 fusion proteins of the invention can be incorporated into pharmaceutical compositions and administered to a subject in vivo. The 47324 fusion proteins can be used to affect the bioavailability of a 47324 substrate. 47324 fusion proteins may be useful therapeutically for the treatment of disorders caused by, for example, (i) aberrant modification or mutation of a gene encoding a 47324 protein; (ii) mis-regulation of the 47324 gene; and (iii) aberrant post-translational modification of a 47324 protein.

[0155] Moreover, the 47324-fusion proteins of the invention can be used as immunogens to produce anti-47324 antibodies in a subject, to purify 47324 ligands and in screening assays to identify molecules which inhibit the interaction of 47324 with a 47324 substrate.

[0156] Expression vectors are commercially available that already encode a fusion moiety (e.g., a GST polypeptide). A 47324-encoding nucleic acid can be cloned into such an expression vector such that the fusion moiety is linked in-frame to the 47324 protein.

[0157] Variants of 47324 Proteins

[0158] In another aspect, the invention also features a variant of a 47324 polypeptide, e.g., which functions as an agonist (mimetics) or as an antagonist. Variants of the 47324 proteins can be generated by mutagenesis, e.g., discrete point mutation, the insertion or deletion of sequences or the truncation of a 47324 protein. An agonist of the 47324 proteins can retain substantially the same, or a subset, of the biological activities of the naturally occurring form of a 47324 protein. An antagonist of a 47324 protein can inhibit one or more of the activities of the naturally occurring form of the 47324 protein by, for example, competitively modulating a 47324-mediated activity of a 47324 protein. Thus, specific biological effects can be elicited by treatment with a variant of limited function. Preferably, treatment of a subject with a variant having a subset of the biological activities of the naturally occurring form of the protein has fewer side effects in a subject relative to treatment with the naturally occurring form of the 47324 protein.

[0159] Variants of a 47324 protein can be identified by screening combinatorial libraries of mutants, e.g., truncation mutants, of a 47324 protein for agonist or antagonist activity.

[0160] Libraries of fragments e.g., N terminal, C terminal, or internal fragments, of a 47324 protein coding sequence can be used to generate a variegated population of fragments for screening and subsequent selection of variants of a 47324 protein.

[0161] Variants in which a cysteine residues is added or deleted or in which a residue which is glycosylated is added or deleted are particularly preferred.

[0162] Methods for screening gene products of combinatorial libraries made by point mutations or truncation, and for screening cDNA libraries for gene products having a

selected property. Recursive ensemble mutagenesis (REM), a new technique which enhances the frequency of functional mutants in the libraries, can be used in combination with the screening assays to identify 47324 variants (Arkin and Yourvan, (1992) *Proc. Natl. Acad. Sci. USA* 89:7811-7815; Delgrave et al., (1993) *Protein Engineering* 6(3):327-331).

[0163] Cell based assays can be exploited to analyze a variegated 47324 library. For example, a library of expression vectors can be transfected into a cell line, e.g., a cell line, which ordinarily responds to 47324 in a substrate-dependent manner. The transfected cells are then contacted with 47324 and the effect of the expression of the mutant on signaling by the 47324 substrate can be detected, e.g., by measuring Ras family activity. Plasmid DNA can then be recovered from the cells which score for inhibition, or alternatively, potentiation of signaling by the 47324 substrate, and the individual clones further characterized.

[0164] In another aspect, the invention features a method of making a 47324 polypeptide, e.g., a peptide having a non-wild type activity, e.g., an antagonist, agonist, or super agonist of a naturally occurring 47324 polypeptide, e.g., a naturally occurring 47324 polypeptide. The method includes: altering the sequence of a 47324 polypeptide, e.g., altering the sequence, e.g., by substitution or deletion of one or more residues of a non-conserved region, a domain or residue disclosed herein, and testing the altered polypeptide for the desired activity.

[0165] In another aspect, the invention features a method of making a fragment or analog of a 47324 polypeptide a biological activity of a naturally occurring 47324 polypeptide. The method includes: altering the sequence, e.g., by substitution or deletion of one or more residues, of a 47324 polypeptide, e.g., altering the sequence of a non-conserved region, or a domain or residue described herein, and testing the altered polypeptide for the desired activity.

[0166] Anti-47324 Antibodies

[0167] In another aspect, the invention provides an anti-47324 antibody. The term "antibody" as used herein refers to an immunoglobulin molecule or immunologically active portion thereof, i.e., an antigen-binding portion. Examples of immunologically active portions of immunoglobulin molecules include F(ab) and F(ab)₂ fragments which can be generated by treating the antibody with an enzyme such as pepsin.

[0168] The antibody can be a polyclonal, monoclonal, recombinant, e.g., a chimeric or humanized, fully human, non-human, e.g., murine, or single chain antibody. In a preferred embodiment it has effector function and can fix complement. The antibody can be coupled to a toxin or imaging agent.

[0169] A full-length 47324 protein or, antigenic peptide fragment of 47324 can be used as an immunogen or can be used to identify anti-47324 antibodies made with other immunogens, e.g., cells, membrane preparations, and the like. The antigenic peptide of 47324 should include at least 8 amino acid residues of the amino acid sequence shown in SEQ ID NO:2 and encompasses an epitope of 47324. Preferably, the antigenic peptide includes at least 10 amino acid residues, more preferably at least 15 amino acid residues, even more preferably at least 20 amino acid residues, and most preferably at least 30 amino acid residues.

[0170] Fragments of 47324 which include, e.g., residues 205-220 of SEQ ID NO:2 can be, e.g., used as immunogens, or used to be hydrophilic regions of the 47324 protein. Similarly, a fragment of 47324 which includes, e.g., residues 63-73 of SEQ ID NO:2 can be used to make an antibody against what is believed to be a hydrophobic region of the 47324 protein; a fragment of 47324 which includes, e.g., residues 8-231 of SEQ ID NO:2 can be used to make an antibody against what is believed to be the Ras family region of the 47324 protein.

[0171] Antibodies reactive with, or specific for, any of these regions, or other regions or domains described herein are provided.

[0172] In a preferred embodiment the antibody fails to bind an Fc receptor, e.g. it is a type which does not support Fc receptor binding or has been modified, e.g., by deletion or other mutation, such that it does not have a functional Fc receptor binding region.

[0173] Preferred epitopes encompassed by the antigenic peptide are regions of 47324 are located on the surface of the protein, e.g., hydrophilic regions, as well as regions with high antigenicity. For example, an Emini surface probability analysis of the human 47324 protein sequence can be used to indicate the regions that have a particularly high probability of being localized to the surface of the 47324 protein and are thus likely to constitute surface residues useful for targeting antibody production.

[0174] In a preferred embodiment the antibody binds an epitope on any domain or region on 47324 proteins described herein.

[0175] Chimeric, humanized, but most preferably, completely human antibodies are desirable for applications which include repeated administration, e.g., therapeutic treatment (and some diagnostic applications) of human patients.

[0176] Chimeric and humanized monoclonal antibodies, comprising both human and non-human portions, can be made using standard recombinant DNA techniques. Such chimeric and humanized monoclonal antibodies can be produced by recombinant DNA techniques known in the art, for example using methods described in Robinson et al. International Application No. PCT/US86/02269; Akira, et al. European Patent Application 184,187; Taniguchi, M., European Patent Application 171,496; Morrison et al. European Patent Application 173,494; Neuberger et al. PCT International Publication No. WO 86/01533; Cabilly et al. U.S. Pat. No. 4,816,567; Cabilly et al. European Patent Application 125,023; Better et al. (1988) *Science* 240:1041-1043; Liu et al. (1987) *Proc. Natl. Acad. Sci. USA* 84:3439-3443; Liu et al. (1987) *J. Immunol.* 139:3521-3526; Sun et al. (1987) *Proc. Natl. Acad. Sci. USA* 84:214-218; Nishimura et al. (1987) *Canc. Res.* 47:999-1005; Wood et al. (1985) *Nature* 314:446-449; and Shaw et al. (1988) *J. Natl. Cancer Inst.* 80:1553-1559; Morrison, S. L. (1985) *Science* 229:1202-1207; Oi et al. (1986) *BioTechniques* 4:214; Winter U.S. Pat. No. 5,225,539; Jones et al. (1986) *Nature* 321:552-525; Verhoeyan et al. (1988) *Science* 239:1534; and Beidler et al. (1988) *J. Immunol.* 141:4053-4060.

[0177] Completely human antibodies are particularly desirable for therapeutic treatment of human patients. Such antibodies can be produced using transgenic mice that are

incapable of expressing endogenous immunoglobulin heavy and light chains genes, but which can express human heavy and light chain genes. See, for example, Lonberg and Huszar (1995) *Int. Rev. Immunol.* 13:65-93; and U.S. Pat. Nos. 5,625,126; 5,633,425; 5,569,825; 5,661,016; and 5,545,806. In addition, companies such as Abgenix, Inc. (Fremont, Calif.) and Medarex, Inc. (Princeton, N.J.), can be engaged to provide human antibodies directed against a selected antigen using technology similar to that described above.

[0178] Completely human antibodies that recognize a selected epitope can be generated using a technique referred to as "guided selection." In this approach a selected non-human monoclonal antibody, e.g., a murine antibody, is used to guide the selection of a completely human antibody recognizing the same epitope. This technology is described by Jespers et al. (1994) *Bio/Technology* 12:899-903).

[0179] The anti-47324 antibody can be a single chain antibody. A single-chain antibody (scFV) can be engineered (see, for example, Colcher, D. et al. (1999) *Ann. NY Acad. Sci.* 880:263-80; and Reiter, Y. (1996) *Clin. Cancer Res.* 2:245-52). The single chain antibody can be dimerized or multimerized to generate multivalent antibodies having specificities for different epitopes of the same target 47324 protein.

[0180] In a preferred embodiment, the antibody has reduced or no ability to bind an Fc receptor. For example, it is an isotype or subtype, fragment or other mutant, which does not support binding to an Fc receptor, e.g., it has a mutagenized or deleted Fc receptor binding region. An antibody (or fragment thereof) may be conjugated to a therapeutic moiety such as a cytotoxin, a therapeutic agent or a radioactive metal ion. A cytotoxin or cytotoxic agent includes any agent that is detrimental to cells. Examples include taxol, cytochalasin B, gramicidin D, ethidium bromide, emetine, mitomycin, etoposide, tenoposide, vincristine, vinblastine, colchicin, doxorubicin, daunorubicin, dihydroxy anthracin dione, mitoxantrone, mithramycin, actinomycin D, 1-dehydrotestosterone, glucocorticoids, procaine, tetracaine, lidocaine, propranolol, and puromycin and analogs or homologs thereof. Therapeutic agents include, but are not limited to, antimetabolites (e.g., methotrexate, 6-mercaptopurine, 6-thioguanine, cytarabine, 5-fluorouracil decarbazine), alkylating agents (e.g., mechlorethamine, thioepa chlorambucil, melphalan, carmustine (BSNU) and lomustine (CCNU), cyclophosphamide, busulfan, dibromomannitol, streptozotocin, mitomycin C, and cis-dichlorodiamine platinum (II) (DDP) cisplatin), anthracyclines (e.g., daunorubicin (formerly daunomycin) and doxorubicin), antibiotics (e.g., dactinomycin (formerly actinomycin), bleomycin, mithramycin, and anthramycin (AMC)), and anti-mitotic agents (e.g., vincristine and vinblastine).

[0181] The conjugates of the invention can be used for modifying a given biological response, the drug moiety is not to be construed as limited to classical chemical therapeutic agents. For example, the drug moiety may be a protein or polypeptide possessing a desired biological activity. Such proteins may include, for example, a toxin such as abrin, ricin A, pseudomonas exotoxin, or diphtheria toxin; a protein such as tumor necrosis factor, α -interferon, β -interferon, nerve growth factor, platelet derived growth factor, tissue plasminogen activator; or, biological response modi-

fiers such as, for example, lymphokines, interleukin-1 ("IL-1"), interleukin-2 ("IL-2"), interleukin-6 ("IL-6"), granulocyte macrophage colony stimulating factor ("GM-CSF"), granulocyte colony stimulating factor ("G-CSF"), or other growth factors.

[0182] Alternatively, an antibody can be conjugated to a second antibody to form an antibody heteroconjugate as described by Segal in U.S. Pat. No. 4,676,980.

[0183] An anti-47324 antibody (e.g., monoclonal antibody) can be used to isolate 47324 by standard techniques, such as affinity chromatography or immunoprecipitation. Moreover, an anti-47324 antibody can be used to detect 47324 protein (e.g., in a cellular lysate or cell supernatant) in order to evaluate the abundance and pattern of expression of the protein. Anti-47324 antibodies can be used diagnostically to monitor protein levels in tissue as part of a clinical testing procedure, e.g., to determine the efficacy of a given treatment regimen. Detection can be facilitated by coupling (i.e., physically linking) the antibody to a detectable substance (i.e., antibody labelling). Examples of detectable substances include various enzymes, prosthetic groups, fluorescent materials, luminescent materials, bioluminescent materials, and radioactive materials. Examples of suitable enzymes include horseradish peroxidase, alkaline phosphatase, β -galactosidase, or acetylcholinesterase; examples of suitable prosthetic group complexes include streptavidin-biotin and avidin/biotin; examples of suitable fluorescent materials include umbelliferone, fluorescein, fluorescein isothiocyanate, rhodamine, dichlorotriazinylamine fluorescein, dansyl chloride or phycoerythrin; an example of a luminescent material includes luminol; examples of bioluminescent materials include luciferase, luciferin, and aequorin, and examples of suitable radioactive material include ^{125}I , ^{131}I , ^{35}S or ^3H .

[0184] In preferred embodiments, an antibody can be made by immunizing with a purified 47324 antigen, or a fragment thereof, e.g., a fragment described herein, a membrane associated antigen, tissues, e.g., crude tissue preparations, whole cells, preferably living cells, lysed cells, or cell fractions.

[0185] Antibodies which bind only a native 47324 protein, only denatured or otherwise non-native 47324 protein, or which bind both, are within the invention. Antibodies with linear or conformational epitopes are within the invention. Conformational epitopes sometimes can be identified by identifying antibodies which bind to native but not denatured 47324 protein.

[0186] Recombinant Expression Vectors, Host Cells and Genetically Engineered Cells

[0187] In another aspect, the invention includes, vectors, preferably expression vectors, containing a nucleic acid encoding a polypeptide described herein. As used herein, the term "vector" refers to a nucleic acid molecule capable of transporting another nucleic acid to which it has been linked and can include a plasmid, cosmid or viral vector. The vector can be capable of autonomous replication or it can integrate into a host DNA. Viral vectors include, e.g., replication defective retroviruses, adenoviruses and adeno-associated viruses.

[0188] A vector can include a 47324 nucleic acid in a form suitable for expression of the nucleic acid in a host cell.

Preferably the recombinant expression vector includes one or more regulatory sequences operatively linked to the nucleic acid sequence to be expressed. The term "regulatory sequence" includes promoters, enhancers and other expression control elements (e.g., polyadenylation signals). Regulatory sequences include those which direct constitutive expression of a nucleotide sequence, as well as tissue-specific regulatory and/or inducible sequences. The design of the expression vector can depend on such factors as the choice of the host cell to be transformed, the level of expression of protein desired, and the like. The expression vectors of the invention can be introduced into host cells to thereby produce proteins or polypeptides, including fusion proteins or polypeptides, encoded by nucleic acids as described herein (e.g., 47324 proteins, mutant forms of 47324 proteins, fusion proteins, and the like).

[0189] The recombinant expression vectors of the invention can be designed for expression of 47324 proteins in prokaryotic or eukaryotic cells. For example, polypeptides of the invention can be expressed in *E. coli*, insect cells (e.g., using baculovirus expression vectors), yeast cells or mammalian cells. Suitable host cells are discussed further in Goeddel, *Gene Expression Technology: Methods in Enzymology* 185, Academic Press, San Diego, Calif. (1990). Alternatively, the recombinant expression vector can be transcribed and translated in vitro, for example using T7 promoter regulatory sequences and T7 polymerase.

[0190] Expression of proteins in prokaryotes is most often carried out in *E. coli* with vectors containing constitutive or inducible promoters directing the expression of either fusion or non-fusion proteins. Fusion vectors add a number of amino acids to a protein encoded therein, usually to the amino terminus of the recombinant protein. Such fusion vectors typically serve three purposes: 1) to increase expression of recombinant protein; 2) to increase the solubility of the recombinant protein; and 3) to aid in the purification of the recombinant protein by acting as a ligand in affinity purification. Often, a proteolytic cleavage site is introduced at the junction of the fusion moiety and the recombinant protein to enable separation of the recombinant protein from the fusion moiety subsequent to purification of the fusion protein. Such enzymes, and their cognate recognition sequences, include Factor Xa, thrombin and enterokinase. Typical fusion expression vectors include pGEX (Pharmacia Biotech Inc; Smith, D. B. and Johnson, K. S., (1988) *Gene* 67:31-40), pMAL (New England Biolabs, Beverly, Mass.) and pRIT5 (Pharmacia, Piscataway, N.J.) which fuse glutathione S-transferase (GST), maltose E binding protein, or protein A, respectively, to the target recombinant protein.

[0191] Purified fusion proteins can be used in 47324 activity assays, (e.g., direct assays or competitive assays described in detail below), or to generate antibodies specific for 47324 proteins. In a preferred embodiment, a fusion protein expressed in a retroviral expression vector of the present invention can be used to infect bone marrow cells which are subsequently transplanted into irradiated recipients. The pathology of the subject recipient is then examined after sufficient time has passed (e.g., six (6) weeks).

[0192] To maximize recombinant protein expression in *E. coli* is to express the protein in host bacteria with an impaired capacity to proteolytically cleave the recombinant protein (Gottesman, S., *Gene Expression Technology: Meth-*

ods in Enzymology 185, Academic Press, San Diego, Calif. (1990) 119-128). Another strategy is to alter the nucleic acid sequence of the nucleic acid to be inserted into an expression vector so that the individual codons for each amino acid are those preferentially utilized in *E. coli* (Wada et al., (1992) *Nucleic Acids Res.* 20:2111-2118). Such alteration of nucleic acid sequences of the invention can be carried out by standard DNA synthesis techniques.

[0193] The 47324 expression vector can be a yeast expression vector, a vector for expression in insect cells, e.g., a baculovirus expression vector or a vector suitable for expression in mammalian cells.

[0194] When used in mammalian cells, the expression vector's control functions are often provided by viral regulatory elements. For example, commonly used promoters are derived from polyoma, Adenovirus 2, cytomegalovirus and Simian Virus 40.

[0195] In another embodiment, the recombinant mammalian expression vector is capable of directing expression of the nucleic acid preferentially in a particular cell type (e.g., tissue-specific regulatory elements are used to express the nucleic acid). Non-limiting examples of suitable tissue-specific promoters include the albumin promoter (liver-specific; Pinkert et al., (1987) *Genes Dev.* 1:268-277), lymphoid-specific promoters (Calame and Eaton, (1988) *Adv. Immunol.* 43:235-275), in particular promoters of T cell receptors (Winoto and Baltimore, (1989) *EMBO J.* 8:729-733) and immunoglobulins (Baneji et al., (1983) *Cell* 33:729-740; Queen and Baltimore, (1983) *Cell* 33:741-748), neuron-specific promoters (e.g., the neurofilament promoter; Byrne and Ruddle, (1989) *Proc. Natl. Acad. Sci. USA* 86:5473-5477), pancreas-specific promoters (Edlund et al., (1985) *Science* 230:912-916), and mammary gland-specific promoters (e.g., milk whey promoter; U.S. Pat. No. 4,873, 316 and European Application Publication No. 264,166). Developmentally-regulated promoters are also encompassed, for example, the murine hox promoters (Kessel and Gruss, (1990) *Science* 249:374-379) and the α -fetoprotein promoter (Campes and Tilghman, (1989) *Genes Dev.* 3:537-546).

[0196] The invention further provides a recombinant expression vector comprising a DNA molecule of the invention cloned into the expression vector in an antisense orientation. Regulatory sequences (e.g., viral promoters and/or enhancers) operatively linked to a nucleic acid cloned in the antisense orientation can be chosen which direct the constitutive, tissue specific or cell type specific expression of antisense RNA in a variety of cell types. The antisense expression vector can be in the form of a recombinant plasmid, phagemid or attenuated virus. For a discussion of the regulation of gene expression using antisense genes see Weintraub, H. et al., *Antisense RNA as a molecular tool for genetic analysis, Reviews—Trends in Genetics*, Vol. 1(1) 1986.

[0197] Another aspect the invention provides a host cell which includes a nucleic acid molecule described herein, e.g., a 47324 nucleic acid molecule within a recombinant expression vector or a 47324 nucleic acid molecule containing sequences which allow it to homologously recombine into a specific site of the host cell's genome. The terms "host cell" and "recombinant host cell" are used interchangeably herein. Such terms refer not only to the particular subject

cell but rather also to the progeny or potential progeny of such a cell. Because certain modifications may occur in succeeding generations due to either mutation or environmental influences, such progeny may not, in fact, be identical to the parent cell, but are still included within the scope of the term as used herein.

[0198] A host cell can be any prokaryotic or eukaryotic cell. For example, a 47324 protein can be expressed in bacterial cells such as *E. coli*, insect cells, yeast or mammalian cells (such as Chinese hamster ovary cells (CHO) or COS cells). Other suitable host cells are known to those skilled in the art.

[0199] Vector DNA can be introduced into host cells via conventional transformation or transfection techniques. As used herein, the terms "transformation" and "transfection" are intended to refer to a variety of art-recognized techniques for introducing foreign nucleic acid (e.g., DNA) into a host cell, including calcium phosphate or calcium chloride co-precipitation, DEAE-dextran-mediated transfection, lipofection, or electroporation

[0200] A host cell of the invention can be used to produce (i.e., express) a 47324 protein. Accordingly, the invention further provides methods for producing a 47324 protein using the host cells of the invention. In one embodiment, the method includes culturing the host cell of the invention (into which a recombinant expression vector encoding a 47324 protein has been introduced) in a suitable medium such that a 47324 protein is produced. In another embodiment, the method further includes isolating a 47324 protein from the medium or the host cell.

[0201] In another aspect, the invention features, a cell or purified preparation of cells which include a 47324 transgene, or which otherwise misexpress 47324. The cell preparation can consist of human or non-human cells, e.g., rodent cells, e.g., mouse or rat cells, rabbit cells, or pig cells. In preferred embodiments, the cell or cells include a 47324 transgene, e.g., a heterologous form of a 47324, e.g., a gene derived from humans (in the case of a non-human cell). The 47324 transgene can be misexpressed, e.g., overexpressed or underexpressed. In other preferred embodiments, the cell or cells include a gene which misexpress an endogenous 47324, e.g., a gene the expression of which is disrupted, e.g., a knockout. Such cells can serve as a model for studying disorders which are related to mutated or mis-expressed 47324 alleles or for use in drug screening.

[0202] In another aspect, the invention features, a human cell, e.g., a hematopoietic stem cell, transformed with nucleic acid which encodes a subject 47324 polypeptide.

[0203] Also provided are cells or a purified preparation thereof, e.g., human cells, in which an endogenous 47324 is under the control of a regulatory sequence that does not normally control the expression of the endogenous 47324 gene. The expression characteristics of an endogenous gene within a cell, e.g., a cell line or microorganism, can be modified by inserting a heterologous DNA regulatory element into the genome of the cell such that the inserted regulatory element is operably linked to the endogenous 47324 gene. For example, an endogenous 47324 gene, e.g., a gene which is "transcriptionally silent," e.g., not normally expressed, or expressed only at very low levels, may be activated by inserting a regulatory element which is capable

of promoting the expression of a normally expressed gene product in that cell. Techniques such as targeted homologous recombinations, can be used to insert the heterologous DNA as described in, e.g., Chappel, U.S. Pat. No. 5,272,071; WO 91/06667, published on May 16, 1991.

[0204] Transgenic Animals

[0205] The invention provides non-human transgenic animals. Such animals are useful for studying the function and/or activity of a 47324 protein and for identifying and/or evaluating modulators of 47324 activity. As used herein, a "transgenic animal" is a non-human animal, preferably a mammal, more preferably a rodent such as a rat or mouse, in which one or more of the cells of the animal includes a transgene. Other examples of transgenic animals include non-human primates, sheep, dogs, cows, goats, chickens, amphibians, and the like. A transgene is exogenous DNA or a rearrangement, e.g., a deletion of endogenous chromosomal DNA, which preferably is integrated into or occurs in the genome of the cells of a transgenic animal. A transgene can direct the expression of an encoded gene product in one or more cell types or tissues of the transgenic animal, other transgenes, e.g., a knockout, reduce expression. Thus, a transgenic animal can be one in which an endogenous 47324 gene has been altered by, e.g., by homologous recombination between the endogenous gene and an exogenous DNA molecule introduced into a cell of the animal, e.g., an embryonic cell of the animal, prior to development of the animal.

[0206] Intronic sequences and polyadenylation signals can also be included in the transgene to increase the efficiency of expression of the transgene. A tissue-specific regulatory sequence(s) can be operably linked to a transgene of the invention to direct expression of a 47324 protein to particular cells. A transgenic founder animal can be identified based upon the presence of a 47324 transgene in its genome and/or expression of 47324 mRNA in tissues or cells of the animals. A transgenic founder animal can then be used to breed additional animals carrying the transgene. Moreover, transgenic animals carrying a transgene encoding a 47324 protein can further be bred to other transgenic animals carrying other transgenes. 47324 proteins or polypeptides can be expressed in transgenic animals or plants, e.g., a nucleic acid encoding the protein or polypeptide can be introduced into the genome of an animal. In preferred embodiments the nucleic acid is placed under the control of a tissue specific promoter, e.g., a milk or egg specific promoter, and recovered from the milk or eggs produced by the animal. Suitable animals are mice, pigs, cows, goats, and sheep.

[0207] The invention also includes a population of cells from a transgenic animal, as discussed herein.

[0208] Uses

[0209] The nucleic acid molecules, proteins, protein homologues, and antibodies described herein can be used in one or more of the following methods: a) screening assays; b) predictive medicine (e.g., diagnostic assays, prognostic assays, monitoring clinical trials, and pharmacogenetics); and c) methods of treatment (e.g., therapeutic and prophylactic).

[0210] The isolated nucleic acid molecules of the invention can be used, for example, to express a 47324 protein (e.g., via a recombinant expression vector in a host cell in

gene therapy applications), to detect a 47324 mRNA (e.g., in a biological sample) or a genetic alteration in a 47324 gene, and to modulate 47324 activity, as described further below. The 47324 proteins can be used to treat disorders characterized by insufficient or excessive production of a 47324 substrate or production of 47324 inhibitors. In addition, the 47324 proteins can be used to screen for naturally occurring 47324 substrates, to screen for drugs or compounds which modulate 47324 activity, as well as to treat disorders characterized by insufficient or excessive production of 47324 protein or production of 47324 protein forms which have decreased, aberrant or unwanted activity compared to 47324 wild-type protein. Such disorders include those characterized by aberrant signaling or aberrant, e.g., Ras family activity. Moreover, the anti-47324 antibodies of the invention can be used to detect and isolate 47324 proteins, regulate the bioavailability of 47324 proteins, and modulate 47324 activity.

[0211] A method of evaluating a compound for the ability to interact with, e.g., bind, a subject 47324 polypeptide is provided. The method includes: contacting the compound with the subject 47324 polypeptide; and evaluating ability of the compound to interact with, e.g., to bind or form a complex with the subject 47324 polypeptide. This method can be performed in vitro, e.g., in a cell free system, or in vivo, e.g., in a two-hybrid interaction trap assay. This method can be used to identify naturally occurring molecules which interact with subject 47324 polypeptide. It can also be used to find natural or synthetic inhibitors of subject 47324 polypeptide. Screening methods are discussed in more detail below.

[0212] Screening Assays:

[0213] The invention provides methods (also referred to herein as "screening assays") for identifying modulators, i.e., candidate or test compounds or agents (e.g., proteins, peptides, peptidomimetics, peptoids, small molecules or other drugs) which bind to 47324 proteins, have a stimulatory or inhibitory effect on, for example, 47324 expression or 47324 activity, or have a stimulatory or inhibitory effect on, for example, the expression or activity of a 47324 substrate. Compounds thus identified can be used to modulate the activity of target gene products (e.g., 47324 genes) in a therapeutic protocol, to elaborate the biological function of the target gene product, or to identify compounds that disrupt normal target gene interactions.

[0214] In one embodiment, the invention provides assays for screening candidate or test compounds which are substrates of a 47324 protein or polypeptide or a biologically active portion thereof. In another embodiment, the invention provides assays for screening candidate or test compounds which bind to or modulate the activity of a 47324 protein or polypeptide or a biologically active portion thereof.

[0215] The test compounds of the present invention can be obtained using any of the numerous approaches in combinatorial library methods known in the art, including: biological libraries; peptoid libraries [libraries of molecules having the functionalities of peptides, but with a novel, non-peptide backbone which are resistant to enzymatic degradation but which nevertheless remain bioactive] (see, e.g., Zuckermann, R. N. et al., *J. Med. Chem.* 1994, 37: 2678-85); spatially addressable parallel solid phase or solution phase libraries; synthetic library methods requiring

deconvolution; the 'one-bead one-compound' library method; and synthetic library methods using affinity chromatography selection. The biological library and peptoid library approaches are limited to peptide libraries, while the other four approaches are applicable to peptide, non-peptide oligomer or small molecule libraries of compounds (Lam, K. S. (1997) *Anticancer Drug Des.* 12:145).

[0216] Examples of methods for the synthesis of molecular libraries can be found in the art, for example in: DeWitt et al. (1993) *Proc. Natl. Acad. Sci. U.S.A.* 90:6909; Erb et al., (1994) *Proc. Natl. Acad. Sci. USA* 91:11422; Zuckermann et al., (1994). *J. Med. Chem.* 37:2678; Cho et al., (1993) *Science* 261:1303; Carrell et al., (1994) *Angew. Chem. Int. Ed. Engl.* 33:2059; Carell et al., (1994) *Angew. Chem. Int. Ed. Engl.* 33:2061; and in Gallop et al., (1994) *J. Med. Chem.* 37:1233.

[0217] Libraries of compounds may be presented in solution (e.g., Houghten, (1992) *Biotechniques* 13:412-421), or on beads (Lam, (1991) *Nature* 354:82-84), chips (Fodor, (1993) *Nature* 364:555-556), bacteria or spores (Ladner, U.S. Pat. No. 5,223,409), plasmids (Cull et al., (1992) *Proc. Natl. Acad. Sci. USA* 89:1865-1869) or on phage (Scott and Smith, (1990) *Science* 249:386-390); (Devlin, (1990) *Science* 249:404-406); (Cwirla et al., (1990) *Proc. Natl. Acad. Sci.* 87:6378-6382); (Felici, (1991) *J. Mol. Biol.* 222:301-310); (Ladner supra.).

[0218] In one embodiment, an assay is a cell-based assay in which a cell which expresses a 47324 protein or biologically active portion thereof is contacted with a test compound, and the ability of the test compound to modulate 47324 activity is determined. Determining the ability of the test compound to modulate 47324 activity can be accomplished by monitoring, for example, Ras family activity. The cell, for example, can be of mammalian origin, e.g., human. Cell homogenates, or fractions, preferably membrane containing fractions, can also be tested.

[0219] The ability of the test compound to modulate 47324 binding to a compound, e.g., a 47324 substrate, or to bind to 47324 can also be evaluated. This can be accomplished, for example, by coupling the compound, e.g., the substrate, with a radioisotope or enzymatic label such that binding of the compound, e.g., the substrate, to 47324 can be determined by detecting the labeled compound, e.g., substrate, in a complex. Alternatively, 47324 could be coupled with a radioisotope or enzymatic label to monitor the ability of a test compound to modulate 47324 binding to a 47324 substrate in a complex. For example, compounds (e.g., 47324 substrates) can be labeled with ^{125}I , ^{35}S , ^{14}C , or ^3H , either directly or indirectly, and the radioisotope detected by direct counting of radioemission or by scintillation counting. Alternatively, compounds can be enzymatically labeled with, for example, horseradish peroxidase, alkaline phosphatase, or luciferase, and the enzymatic label detected by determination of conversion of an appropriate substrate to product.

[0220] The ability of a compound (e.g., a 47324 substrate) to interact with 47324 with or without the labeling of any of the interactants can be evaluated. For example, a microphysiometer can be used to detect the interaction of a compound with 47324 without the labeling of either the compound or the 47324. McConnell, H. M. et al., (1992) *Science* 257:1906-1912. As used herein, a "microphysiom-

eter" (e.g., Cytosensor) is an analytical instrument that measures the rate at which a cell acidifies its environment using a light-addressable potentiometric sensor (LAPS). Changes in this acidification rate can be used as an indicator of the interaction between a compound and 47324.

[0221] In yet another embodiment, a cell-free assay is provided in which a 47324 protein or biologically active portion thereof is contacted with a test compound and the ability of the test compound to bind to the 47324 protein or biologically active portion thereof is evaluated. Preferred biologically active portions of the 47324 proteins to be used in assays of the present invention include fragments which participate in interactions with non-47324 molecules, e.g., fragments with high surface probability scores.

[0222] Soluble and/or membrane-bound forms of isolated proteins (e.g., 47324 proteins or biologically active portions thereof) can be used in the cell-free assays of the invention. When membrane-bound forms of the protein are used, it may be desirable to utilize a solubilizing agent. Examples of such solubilizing agents include non-ionic detergents such as n-octylglucoside, n-dodecylglucoside, n-dodecylmalto-side, octanoyl-N-methylglucamide, decanoyl-N-methylglu-camide, Triton® X-100, Triton® X-114, Thesit®, Isotride-cypoly(ethylene glycol ether)_n, 3-[(3-cholamidopropyl)dimethylamino]-1-propane sulfonate (CHAPS), 3-[(3-cholamidopropyl)dimethylamminio]-2-hydroxy-1-propane sulfonate (CHAPSO), or N-dodecyl-N,N-dimethyl-3-ammonio-1-propane sulfonate.

[0223] Cell-free assays involve preparing a reaction mixture of the target gene protein and the test compound under conditions and for a time sufficient to allow the two components to interact and bind, thus forming a complex that can be removed and/or detected.

[0224] In one embodiment, assays are performed where the ability of an agent to block Ras family activity within a cell is evaluated.

[0225] The interaction between two molecules can also be detected, e.g., using fluorescence energy transfer (FET) (see, for example, Lakowicz et al., U.S. Pat. No. 5,631,169; Stavrianopoulos, et al., U.S. Pat. No. 4,868,103). A fluorophore label on the first, 'donor' molecule is selected such that its emitted fluorescent energy will be absorbed by a fluorescent label on a second, 'acceptor' molecule, which in turn is able to fluoresce due to the absorbed energy. Alternately, the 'donor' protein molecule may simply utilize the natural fluorescent energy of tryptophan residues. Labels are chosen that emit different wavelengths of light, such that the 'acceptor' molecule label may be differentiated from that of the 'donor'. Since the efficiency of energy transfer between the labels is related to the distance separating the molecules, the spatial relationship between the molecules can be assessed. In a situation in which binding occurs between the molecules, the fluorescent emission of the 'acceptor' molecule label in the assay should be maximal. An FET binding event can be conveniently measured through standard fluorometric detection means well known in the art (e.g., using a fluorimeter).

[0226] In another embodiment, determining the ability of the 47324 protein to bind to a target molecule can be accomplished using real-time Biomolecular Interaction Analysis (BIA) (see, e.g., Sjolander, S. and Urbaniczky, C.,

(1991) *Anal. Chem.* 63:2338-2345 and Szabo et al., (1995) *Curr. Opin. Struct. Biol.* 5:699-705). "Surface plasmon resonance" or "BIA" detects biospecific interactions in real time, without labeling any of the interactants (e.g., BIAcore). Changes in the mass at the binding surface (indicative of a binding event) result in alterations of the refractive index of light near the surface (the optical phenomenon of surface plasmon resonance (SPR)), resulting in a detectable signal which can be used as an indication of real-time reactions between biological molecules.

[0227] In one embodiment, the target gene product or the test substance is anchored onto a solid phase. The target gene product/test compound complexes anchored on the solid phase can be detected at the end of the reaction. Preferably, the target gene product can be anchored onto a solid surface, and the test compound, (which is not anchored), can be labeled, either directly or indirectly, with detectable labels discussed herein.

[0228] It may be desirable to immobilize either 47324, an anti-47324 antibody or its target molecule to facilitate separation of complexed from uncomplexed forms of one or both of the proteins, as well as to accommodate automation of the assay. Binding of a test compound to a 47324 protein, or interaction of a 47324 protein with a target molecule in the presence and absence of a candidate compound, can be accomplished in any vessel suitable for containing the reactants. Examples of such vessels include microtiter plates, test tubes, and micro-centrifuge tubes. In one embodiment, a fusion protein can be provided which adds a domain that allows one or both of the proteins to be bound to a matrix. For example, glutathione-S-transferase/47324 fusion proteins or glutathione-S-transferase/target fusion proteins can be adsorbed onto glutathione sepharose beads (Sigma Chemical, St. Louis, Mo.) or glutathione derivatized microtiter plates, which are then combined with the test compound or the test compound and either the non-adsorbed target protein or 47324 protein, and the mixture incubated under conditions conducive to complex formation (e.g., at physiological conditions for salt and pH). Following incubation, the beads or microtiter plate wells are washed to remove any unbound components, the matrix immobilized in the case of beads, complex determined either directly or indirectly, for example, as described above. Alternatively, the complexes can be dissociated from the matrix, and the level of 47324 binding or activity determined using standard techniques.

[0229] Other techniques for immobilizing either a 47324 protein or a target molecule on matrices include using conjugation of biotin and streptavidin. Biotinylated 47324 protein or target molecules can be prepared from biotin-NHS (N-hydroxy-succinimide) using techniques known in the art (e.g., biotinylation kit, Pierce Chemicals, Rockford, Ill.), and immobilized in the wells of streptavidin-coated 96 well plates (Pierce Chemical).

[0230] In order to conduct the assay, the non-immobilized component is added to the coated surface containing the anchored component. After the reaction is complete, unreacted components are removed (e.g., by washing) under conditions such that any complexes formed will remain immobilized on the solid surface. The detection of complexes anchored on the solid surface can be accomplished in a number of ways. Where the previously non-immobilized

component is pre-labeled, the detection of label immobilized on the surface indicates that complexes were formed. Where the previously non-immobilized component is not pre-labeled, an indirect label can be used to detect complexes anchored on the surface; e.g., using a labeled antibody specific for the immobilized component (the antibody, in turn, can be directly labeled or indirectly labeled with, e.g., a labeled anti-Ig antibody).

[0231] In one embodiment, this assay is performed utilizing antibodies reactive with 47324 protein or target molecules but which do not interfere with binding of the 47324 protein to its target molecule. Such antibodies can be derivatized to the wells of the plate, and unbound target or 47324 protein trapped in the wells by antibody conjugation. Methods for detecting such complexes, in addition to those described above for the GST-immobilized complexes, include immunodetection of complexes using antibodies reactive with the 47324 protein or target molecule, as well as enzyme-linked assays which rely on detecting an enzymatic activity associated with the 47324 protein or target molecule.

[0232] Alternatively, cell free assays can be conducted in a liquid phase. In such an assay, the reaction products are separated from unreacted components, by any of a number of standard techniques, including but not limited to: differential centrifugation (see, for example, Rivas, G., and Minton, A. P., *Trends Biochem Sci* Aug.18, 1993 (8):284-7); chromatography (gel filtration chromatography, ion-exchange chromatography); electrophoresis (see, e.g., Ausubel, F. et al., eds. *Current Protocols in Molecular Biology* 1999, J. Wiley: New York.); and immunoprecipitation (see, for example, Ausubel, F. et al., eds. *Current Protocols in Molecular Biology* 1999, J. Wiley: New York). Such resins and chromatographic techniques are known to one skilled in the art (see, e.g., Heegaard, N. H., *J. Mol. Recognit.* 1998 Winter;11(1-6):141-8; Hage, D. S., and Tweed, S. A., *J. Chromatogr. B Biomed. Sci. Appl.* Oct. 10, 1997;699(1-2):499-525). Further, fluorescence energy transfer may also be conveniently utilized, as described herein, to detect binding without further purification of the complex from solution.

[0233] In a preferred embodiment, the assay includes contacting the 47324 protein or biologically active portion thereof with a known compound which binds 47324 to form an assay mixture, contacting the assay mixture with a test compound, and determining the ability of the test compound to interact with a 47324 protein, wherein determining the ability of the test compound to interact with a 47324 protein includes determining the ability of the test compound to preferentially bind to 47324 or biologically active portion thereof, or to modulate the activity of a target molecule, as compared to the known compound.

[0234] The target gene products of the invention can, in vivo, interact with one or more cellular or extracellular macromolecules, such as proteins. For the purposes of this discussion, such cellular and extracellular macromolecules are referred to herein as "binding partners." Compounds that disrupt such interactions can be useful in regulating the activity of the target gene product. Such compounds can include, but are not limited to molecules such as antibodies, peptides, and small molecules. The preferred target genes/products for use in this embodiment are the 47324 genes

herein identified. In an alternative embodiment, the invention provides methods for determining the ability of the test compound to modulate the activity of a 47324 protein through modulation of the activity of a downstream effector of a 47324 target molecule. For example, the activity of the effector molecule on an appropriate target can be determined, or the binding of the effector to an appropriate target can be determined, as previously described.

[0235] To identify compounds that interfere with the interaction between the target gene product and its cellular or extracellular binding partner(s), e.g., a substrate, a reaction mixture containing the target gene product and the binding partner is prepared, under conditions and for a time sufficient, to allow the two products to form complex. In order to test an inhibitory agent, the reaction mixture is provided in the presence and absence of the test compound. The test compound can be initially included in the reaction mixture, or can be added at a time subsequent to the addition of the target gene and its cellular or extracellular binding partner. Control reaction mixtures are incubated without the test compound or with a placebo. The formation of any complexes between the target gene product and the cellular or extracellular binding partner is then detected. The formation of a complex in the control reaction, but not in the reaction mixture containing the test compound, indicates that the compound interferes with the interaction of the target gene product and the interactive binding partner. Additionally, complex formation within reaction mixtures containing the test compound and normal target gene product can also be compared to complex formation within reaction mixtures containing the test compound and mutant target gene product. This comparison can be important in those cases wherein it is desirable to identify compounds that disrupt interactions of mutant but not normal target gene products.

[0236] These assays can be conducted in a heterogeneous or homogeneous format. Heterogeneous assays involve anchoring either the target gene product or the binding partner onto a solid phase, and detecting complexes anchored on the solid phase at the end of the reaction. In homogeneous assays, the entire reaction is carried out in a liquid phase. In either approach, the order of addition of reactants can be varied to obtain different information about the compounds being tested. For example, test compounds that interfere with the interaction between the target gene products and the binding partners, e.g., by competition, can be identified by conducting the reaction in the presence of the test substance. Alternatively, test compounds that disrupt preformed complexes, e.g., compounds with higher binding constants that displace one of the components from the complex, can be tested by adding the test compound to the reaction mixture after complexes have been formed. The various formats are briefly described below.

[0237] In a heterogeneous assay system, either the target gene product or the interactive cellular or extracellular binding partner, is anchored onto a solid surface (e.g., a microtiter plate), while the non-anchored species is labeled, either directly or indirectly. The anchored species can be immobilized by non-covalent or covalent attachments. Alternatively, an immobilized antibody specific for the species to be anchored can be used to anchor the species to the solid surface.

[0238] In order to conduct the assay, the partner of the immobilized species is exposed to the coated surface with or

without the test compound. After the reaction is complete, unreacted components are removed (e.g., by washing) and any complexes formed will remain immobilized on the solid surface. Where the non-immobilized species is pre-labeled, the detection of label immobilized on the surface indicates that complexes were formed. Where the non-immobilized species is not pre-labeled, an indirect label can be used to detect complexes anchored on the surface; e.g., using a labeled antibody specific for the initially non-immobilized species (the antibody, in turn, can be directly labeled or indirectly labeled with, e.g., a labeled anti-Ig antibody). Depending upon the order of addition of reaction components, test compounds that inhibit complex formation or that disrupt preformed complexes can be detected.

[0239] Alternatively, the reaction can be conducted in a liquid phase in the presence or absence of the test compound, the reaction products separated from unreacted components, and complexes detected; e.g., using an immobilized antibody specific for one of the binding components to anchor any complexes formed in solution, and a labeled antibody specific for the other partner to detect anchored complexes. Again, depending upon the order of addition of reactants to the liquid phase, test compounds that inhibit complex or that disrupt preformed complexes can be identified.

[0240] In an alternate embodiment of the invention, a homogeneous assay can be used. For example, a preformed complex of the target gene product and the interactive cellular or extracellular binding partner product is prepared in that either the target gene products or their binding partners are labeled, but the signal generated by the label is quenched due to complex formation (see, e.g., U.S. Pat. No. 4,109,496 that utilizes this approach for immunoassays). The addition of a test substance that competes with and displaces one of the species from the preformed complex will result in the generation of a signal above background. In this way, test substances that disrupt target gene product-binding partner interaction can be identified.

[0241] In yet another aspect, the 47324 proteins can be used as "bait proteins" in a two-hybrid assay or three-hybrid assay (see, e.g., U.S. Pat. No. 5,283,317; Zervos et al., (1993) *Cell* 72:223-232; Madura et al., (1993) *J. Biol. Chem.* 268:12046-12054; Bartel et al., (1993) *Biotechniques* 14:920-924; Iwabuchi et al., (1993) *Oncogene* 8:1693-1696; and Brent WO94/10300), to identify other proteins, which bind to or interact with 47324 ("47324-binding proteins" or "47324-bp") and are involved in 47324 activity. Such 47324-bps can be activators or inhibitors of signals by the 47324 proteins or 47324 targets as, for example, downstream elements of a 47324-mediated signaling pathway.

[0242] The two-hybrid system is based on the modular nature of most transcription factors, which consist of separable DNA-binding and activation domains. Briefly, the assay utilizes two different DNA constructs. In one construct, the gene that codes for a 47324 protein is fused to a gene encoding the DNA binding domain of a known transcription factor (e.g., GAL-4). In the other construct, a DNA sequence, from a library of DNA sequences, that encodes an unidentified protein ("prey" or "sample") is fused to a gene that codes for the activation domain of the known transcription factor. (Alternatively the: 47324 protein can be the fused to the activator domain.) If the "bait" and the "prey"

proteins are able to interact, *in vivo*, forming a 47324-dependent complex, the DNA-binding and activation domains of the transcription factor are brought into close proximity. This proximity allows transcription of a reporter gene (e.g., LacZ) which is operably linked to a transcriptional regulatory site responsive to the transcription factor. Expression of the reporter gene can be detected and cell colonies containing the functional transcription factor can be isolated and used to obtain the cloned gene which encodes the protein which interacts with the 47324 protein.

[0243] In another embodiment, modulators of 47324 expression are identified. For example, a cell or cell free mixture is contacted with a candidate compound and the expression of 47324 mRNA or protein evaluated relative to the level of expression of 47324 mRNA or protein in the absence of the candidate compound. When expression of 47324 mRNA or protein is greater in the presence of the candidate compound than in its absence, the candidate compound is identified as a stimulator of 47324 mRNA or protein expression. Alternatively, when expression of 47324 mRNA or protein is less (statistically significantly less) in the presence of the candidate compound than in its absence, the candidate compound is identified as an inhibitor of 47324 mRNA or protein expression. The level of 47324 mRNA or protein expression can be determined by methods described herein for detecting 47324 mRNA or protein.

[0244] In another aspect, the invention pertains to a combination of two or more of the assays described herein. For example, a modulating agent can be identified using a cell-based or a cell free assay, and the ability of the agent to modulate the activity of a 47324 protein can be confirmed *in vivo*, e.g., in an animal.

[0245] This invention further pertains to novel agents identified by the above-described screening assays. Accordingly, it is within the scope of this invention to further use an agent identified as described herein (e.g., a 47324 modulating agent, an antisense 47324 nucleic acid molecule, a 47324-specific antibody, or a 47324-binding partner) in an appropriate animal model to determine the efficacy, toxicity, side effects, or mechanism of action, of treatment with such an agent. Furthermore, novel agents identified by the above-described screening assays can be used for treatments as described herein.

[0246] Detection Assays

[0247] Portions or fragments of the nucleic acid sequences identified herein can be used as polynucleotide reagents. For example, these sequences can be used to: (i) map their respective genes on a chromosome e.g., to locate gene regions associated with genetic disease or to associate 47324 with a disease; (ii) identify an individual from a minute biological sample (tissue typing); and (iii) aid in forensic identification of a biological sample. These applications are described in the subsections below.

[0248] Chromosome Mapping

[0249] The 47324 nucleotide sequences or portions thereof can be used to map the location of the 47324 genes on a chromosome. This process is called chromosome mapping. Chromosome mapping is useful in correlating the 47324 sequences with genes associated with disease.

[0250] Briefly, 47324 genes can be mapped to chromosomes by preparing PCR primers (preferably 15-25 bp in

length) from the 47324 nucleotide sequences. These primers can then be used for PCR screening of somatic cell hybrids containing individual human chromosomes. Only those hybrids containing the human gene corresponding to the 47324 sequences will yield an amplified fragment.

[0251] A panel of somatic cell hybrids in which each cell line contains either a single human chromosome or a small number of human chromosomes, and a full set of mouse chromosomes, can allow easy mapping of individual genes to specific human chromosomes. (D'Eustachio P. et al., (1983) *Science* 220:919-924).

[0252] Other mapping strategies e.g., *in situ* hybridization (described in Fan, Y. et al., (1990) *Proc. Natl. Acad. Sci. USA*, 87:6223-27), pre-screening with labeled flow-sorted chromosomes, and pre-selection by hybridization to chromosome specific cDNA libraries can be used to map 47324 to a chromosomal location.

[0253] Fluorescence *in situ* hybridization (FISH) of a DNA sequence to a metaphase chromosomal spread can further be used to provide a precise chromosomal location in one step. The FISH technique can be used with a DNA sequence as short as 500 or 600 bases. However, clones larger than 1,000 bases have a higher likelihood of binding to a unique chromosomal location with sufficient signal intensity for simple detection. Preferably 1,000 bases, and more preferably 2,000 bases will suffice to get good results at a reasonable amount of time. For a review of this technique, see Verma et al., *Human Chromosomes: A Manual of Basic Techniques* (Pergamon Press, New York 1988).

[0254] Reagents for chromosome mapping can be used individually to mark a single chromosome or a single site on that chromosome, or panels of reagents can be used for marking multiple sites and/or multiple chromosomes. Reagents corresponding to noncoding regions of the genes actually are preferred for mapping purposes. Coding sequences are more likely to be conserved within gene families, thus increasing the chance of cross hybridizations during chromosomal mapping.

[0255] Once a sequence has been mapped to a precise chromosomal location, the physical position of the sequence on the chromosome can be correlated with genetic map data. (Such data are found, for example, in V. McKusick, *Mendelian Inheritance in Man*, available on-line through Johns Hopkins University Welch Medical Library). The relationship between a gene and a disease, mapped to the same chromosomal region, can then be identified through linkage analysis (co-inheritance of physically adjacent genes), described in, for example, Egeland, J. et al., (1987) *Nature*, 325:783-787.

[0256] Moreover, differences in the DNA sequences between individuals affected and unaffected with a disease associated with the 47324 gene, can be determined. If a mutation is observed in some or all of the affected individuals but not in any unaffected individuals, then the mutation is likely to be the causative agent of the particular disease. Comparison of affected and unaffected individuals generally involves first looking for structural alterations in the chromosomes, such as deletions or translocations that are visible from chromosome spreads or detectable using PCR based on that DNA sequence. Ultimately, complete sequencing of

genes from several individuals can be performed to confirm the presence of a mutation and to distinguish mutations from polymorphisms.

[0257] Tissue Typing

[0258] 47324 sequences can be used to identify individuals from biological samples using, e.g., restriction fragment length polymorphism (RFLP). In this technique, an individual's genomic DNA is digested with one or more restriction enzymes, the fragments separated, e.g., in a Southern blot, and probed to yield bands for identification. The sequences of the present invention are useful as additional DNA markers for RFLP (described in U.S. Pat. No. 5,272,057).

[0259] Furthermore, the sequences of the present invention can also be used to determine the actual base-by-base DNA sequence of selected portions of an individual's genome. Thus, the 47324 nucleotide sequences described herein can be used to prepare two PCR primers from the 5' and 3' ends of the sequences. These primers can then be used to amplify an individual's DNA and subsequently sequence it. Panels of corresponding DNA sequences from individuals, prepared in this manner, can provide unique individual identifications, as each individual will have a unique set of such DNA sequences due to allelic differences.

[0260] Allelic variation occurs to some degree in the coding regions of these sequences, and to a greater degree in the noncoding regions. Each of the sequences described herein can, to some degree, be used as a standard against which DNA from an individual can be compared for identification purposes. Because greater numbers of polymorphisms occur in the noncoding regions, fewer sequences are necessary to differentiate individuals. The noncoding sequences of SEQ ID NO:1 can provide positive individual identification with a panel of perhaps 10 to 1,000 primers which each yield a noncoding amplified sequence of 100 bases. If predicted coding sequences, such as those in SEQ ID NO:3 are used, a more appropriate number of primers for positive individual identification would be 500-2,000.

[0261] If a panel of reagents from 47324 nucleotide sequences described herein is used to generate a unique identification database for an individual, those same reagents can later be used to identify tissue from that individual. Using the unique identification database, positive identification of the individual, living or dead, can be made from extremely small tissue samples.

[0262] Use of Partial 47324 Sequences in Forensic Biology

[0263] DNA-based identification techniques can also be used in forensic biology. To make such an identification, PCR technology can be used to amplify DNA sequences taken from very small biological samples such as tissues, e.g., hair or skin, or body fluids, e.g., blood, saliva, or semen found at a crime scene. The amplified sequence can then be compared to a standard, thereby allowing identification of the origin of the biological sample.

[0264] The sequences of the present invention can be used to provide polynucleotide reagents, e.g., PCR primers, targeted to specific loci in the human genome, which can enhance the reliability of DNA-based forensic identifications by, for example, providing another "identification

marker" (i.e. another DNA sequence that is unique to a particular individual). As mentioned above, actual base sequence information can be used for identification as an accurate alternative to patterns formed by restriction enzyme generated fragments. Sequences targeted to noncoding regions of SEQ ID NO:1 (e.g., fragments derived from the noncoding regions of SEQ ID NO:1 having a length of at least 20 bases, preferably at least 30 bases) are particularly appropriate for this use.

[0265] The 47324 nucleotide sequences described herein can further be used to provide polynucleotide reagents, e.g., labeled or labelable probes which can be used in, for example, an in situ hybridization technique, to identify a specific tissue, e.g., a tissue containing Ras family activity. This can be very useful in cases where a forensic pathologist is presented with a tissue of unknown origin. Panels of such 47324 probes can be used to identify tissue by species and/or by organ type.

[0266] In a similar fashion, these reagents, e.g., 47324 primers or probes can be used to screen tissue culture for contamination (i.e. screen for the presence of a mixture of different types of cells in a culture).

[0267] Predictive Medicine

[0268] The present invention also pertains to the field of predictive medicine in which diagnostic assays, prognostic assays, and monitoring clinical trials are used for prognostic (predictive) purposes to thereby treat an individual.

[0269] Generally, the invention provides, a method of determining if a subject is at risk for a disorder related to a lesion in or the misexpression of a gene which encodes 47324.

[0270] Such disorders include, e.g., a disorder associated with the misexpression of 47324, or Ras family related disorder.

[0271] The method includes one or more of the following:

[0272] detecting, in a tissue of the subject, the presence or absence of a mutation which affects the expression of the 47324 gene, or detecting the presence or absence of a mutation in a region which controls the expression of the gene, e.g., a mutation in the 5' control region;

[0273] detecting, in a tissue of the subject, the presence or absence of a mutation which alters the structure of the 47324 gene;

[0274] detecting, in a tissue of the subject, the misexpression of the 47324 gene, at the mRNA level, e.g., detecting a non-wild type level of a mRNA;

[0275] detecting, in a tissue of the subject, the misexpression of the gene, at the protein level, e.g., detecting a non-wild type level of a 47324 polypeptide.

[0276] In preferred embodiments the method includes: ascertaining the existence of at least one of: a deletion of one or more nucleotides from the 47324 gene; an insertion of one or more nucleotides into the gene, a point mutation, e.g., a substitution of one or more nucleotides of the gene, a gross chromosomal rearrangement of the gene, e.g., a translocation, inversion, or deletion.

[0277] For example, detecting the genetic lesion can include: (i) providing a probe/primer including an oligonucleotide containing a region of nucleotide sequence which hybridizes to a sense or antisense sequence from SEQ ID NO:1 naturally occurring mutants thereof or 5' or 3' flanking sequences naturally associated with the 47324 gene; (ii) exposing the probe/primer to nucleic acid of the tissue; and detecting, by hybridization, e.g., in situ hybridization, of the probe/primer to the nucleic acid, the presence or absence of the genetic lesion.

[0278] In preferred embodiments detecting the misexpression includes ascertaining the existence of at least one of: an alteration in the level of a messenger RNA transcript of the 47324 gene; the presence of a non-wild type splicing pattern of a messenger RNA transcript of the gene; or a non-wild type level of 47324.

[0279] Methods of the invention can be used prenatally or to determine if a subject's offspring will be at risk for a disorder.

[0280] In preferred embodiments the method includes determining the structure of a 47324 gene, an abnormal structure being indicative of risk for the disorder.

[0281] In preferred embodiments the method includes contacting a sample from the subject with an antibody to the 47324 protein or a nucleic acid, which hybridizes specifically with the gene. These and other embodiments are discussed below.

[0282] Diagnostic and Prognostic Assays

[0283] The presence, level, or absence of 47324 protein or nucleic acid in a biological sample can be evaluated by obtaining a biological sample from a test subject and contacting the biological sample with a compound or an agent capable of detecting 47324 protein or nucleic acid (e.g., mRNA, genomic DNA) that encodes 47324 protein such that the presence of 47324 protein or nucleic acid is detected in the biological sample. The term "biological sample" includes tissues, cells and biological fluids isolated from a subject, as well as tissues, cells and fluids present within a subject. A preferred biological sample is serum. The level of expression of the 47324 gene can be measured in a number of ways, including, but not limited to: measuring the mRNA encoded by the 47324 genes; measuring the amount of protein encoded by the 47324 genes; or measuring the activity of the protein encoded by the 47324 genes.

[0284] The level of mRNA corresponding to the 47324 gene in a cell can be determined both by in situ and by in vitro formats.

[0285] The isolated mRNA can be used in hybridization or amplification assays that include, but are not limited to, Southern or Northern analyses, polymerase chain reaction analyses and probe arrays. One preferred diagnostic method for the detection of mRNA levels involves contacting the isolated mRNA with a nucleic acid molecule (probe) that can hybridize to the mRNA encoded by the gene being detected. The nucleic acid probe can be, for example, a full-length 47324 nucleic acid, such as the nucleic acid of SEQ ID NO:1, or the DNA insert of the plasmid deposited with ATCC as Accession Number _____, or a portion thereof, such as an oligonucleotide of at least 7, 15, 30, 50, 100, 250 or 500 nucleotides in length and sufficient to

specifically hybridize under stringent conditions to 47324 mRNA or genomic DNA. Other suitable probes for use in the diagnostic assays are described herein.

[0286] In one format, mRNA (or cDNA) is immobilized on a surface and contacted with the probes, for example by running the isolated mRNA on an agarose gel and transferring the mRNA from the gel to a membrane, such as nitrocellulose. In an alternative format, the probes are immobilized on a surface and the mRNA (or cDNA) is contacted with the probes, for example, in a two-dimensional gene chip array. A skilled artisan can adapt known mRNA detection methods for use in detecting the level of mRNA encoded by the 47324 genes.

[0287] The level of mRNA in a sample that is encoded by one of 47324 can be evaluated with nucleic acid amplification, e.g., by rtPCR (Mullis, 1987, U.S. Pat. No. 4,683,202), ligase chain reaction (Barany, 1991, *Proc. Natl. Acad. Sci. USA* 88:189-193), self sustained sequence replication (Guatelli et al., 1990, *Proc. Natl. Acad. Sci. USA* 87:1874-1878), transcriptional amplification system (Kwoh et al., 1989, *Proc. Natl. Acad. Sci. USA* 86:1173-1177), Q-Beta Replicase (Lizardi et al., 1988, *Bio/Technology* 6:1197), rolling circle replication (Lizardi et al., U.S. Pat. No. 5,854,033) or any other nucleic acid amplification method, followed by the detection of the amplified molecules using techniques known in the art. As used herein, amplification primers are defined as being a pair of nucleic acid molecules that can anneal to 5' or 3' regions of a gene (plus and minus strands, respectively, or vice-versa) and contain a short region in between. In general, amplification primers are from about 10 to 30 nucleotides in length and flank a region from about 50 to 200 nucleotides in length. Under appropriate conditions and with appropriate reagents, such primers permit the amplification of a nucleic acid molecule comprising the nucleotide sequence flanked by the primers.

[0288] For in situ methods, a cell or tissue sample can be prepared/processed and immobilized on a support, typically a glass slide, and then contacted with a probe that can hybridize to mRNA that encodes the 47324 gene being analyzed.

[0289] In another embodiment, the methods further contacting a control sample with a compound or agent capable of detecting 47324 mRNA, or genomic DNA, and comparing the presence of 47324 mRNA or genomic DNA in the control sample with the presence of 47324 mRNA or genomic DNA in the test sample.

[0290] A variety of methods can be used to determine the level of protein encoded by 47324. In general, these methods include contacting an agent that selectively binds to the protein, such as an antibody with a sample, to evaluate the level of protein in the sample. In a preferred embodiment, the antibody bears a detectable label. Antibodies can be polyclonal, or more preferably, monoclonal. An intact antibody, or a fragment thereof (e.g., Fab or F(ab')₂) can be used. The term "labeled", with regard to the probe or antibody, is intended to encompass direct labeling of the probe or antibody by coupling (i.e., physically linking) a detectable substance to the probe or antibody, as well as indirect labeling of the probe or antibody by reactivity with a detectable substance. Examples of detectable substances are provided herein.

[0291] The detection methods can be used to detect 47324 protein in a biological sample in vitro as well as in vivo. In

vitro techniques for detection of 47324 protein include enzyme linked immunosorbent assays (ELISAs), immunoprecipitations, immunofluorescence, enzyme immunoassay (EIA), radioimmunoassay (RIA), and Western blot analysis. In vivo techniques for detection of 47324 protein include introducing into a subject a labeled anti-47324 antibody. For example, the antibody can be labeled with a radioactive marker whose presence and location in a subject can be detected by standard imaging techniques.

[0292] In another embodiment, the methods further include contacting the control sample with a compound or agent capable of detecting 47324 protein, and comparing the presence of 47324 protein in the control sample with the presence of 47324 protein in the test sample.

[0293] The invention also includes kits for detecting the presence of 47324 in a biological sample. For example, the kit can include a compound or agent capable of detecting 47324 protein or mRNA in a biological sample; and a standard. The compound or agent can be packaged in a suitable container. The kit can further comprise instructions for using the kit to detect 47324 protein or nucleic acid.

[0294] For antibody-based kits, the kit can include: (1) a first antibody (e.g., attached to a solid support) which binds to a polypeptide corresponding to a marker of the invention; and, optionally, (2) a second, different antibody which binds to either the polypeptide or the first antibody and is conjugated to a detectable agent.

[0295] For oligonucleotide-based kits, the kit can include: (1) an oligonucleotide, e.g., a detectably labeled oligonucleotide, which hybridizes to a nucleic acid sequence encoding a polypeptide corresponding to a marker of the invention or (2) a pair of primers useful for amplifying a nucleic acid molecule corresponding to a marker of the invention. The kit can also include a buffering agent, a preservative, or a protein-stabilizing agent. The kit can also include components necessary for detecting the detectable agent (e.g., an enzyme or a substrate). The kit can also contain a control sample or a series of control samples which can be assayed and compared to the test sample contained. Each component of the kit can be enclosed within an individual container and all of the various containers can be within a single package, along with instructions for interpreting the results of the assays performed using the kit.

[0296] The diagnostic methods described herein can identify subjects having, or at risk of developing, a disease or disorder associated with misexpressed or aberrant or unwanted 47324 expression or activity. As used herein, the term "unwanted" includes an unwanted phenomenon involved in a biological response such as pain or deregulated cell proliferation.

[0297] In one embodiment, a disease or disorder associated with aberrant or unwanted 47324 expression or activity is identified. A test sample is obtained from a subject and 47324 protein or nucleic acid (e.g., mRNA or genomic DNA) is evaluated, wherein the level, e.g., the presence or absence, of 47324 protein or nucleic acid is diagnostic for a subject having or at risk of developing a disease or disorder associated with aberrant or unwanted 47324 expression or activity. As used herein, a "test sample" refers to a biological sample obtained from a subject of interest, including a biological fluid (e.g., serum), cell sample, or tissue.

[0298] The prognostic assays described herein can be used to determine whether a subject can be administered an agent (e.g., an agonist, antagonist, peptidomimetic, protein, peptide, nucleic acid, small molecule, or other drug candidate) to treat a disease or disorder associated with aberrant or unwanted 47324 expression or activity. For example, such methods can be used to determine whether a subject can be effectively treated with an agent for a Ras family related disorder.

[0299] The methods of the invention can also be used to detect genetic alterations in a 47324 gene, thereby determining if a subject with the altered gene is at risk for a disorder characterized by misregulation in 47324 protein activity or nucleic acid expression, such as a Ras family related disorder. In preferred embodiments, the methods include detecting, in a sample from the subject, the presence or absence of a genetic alteration characterized by at least one of an alteration affecting the integrity of a gene encoding a 47324-protein, or the mis-expression of the 47324 gene. For example, such genetic alterations can be detected by ascertaining the existence of at least one of 1) a deletion of one or more nucleotides from a 47324 gene; 2) an addition of one or more nucleotides to a 47324 gene; 3) a substitution of one or more nucleotides of a 47324 gene, 4) a chromosomal rearrangement of a 47324 gene; 5) an alteration in the level of a messenger RNA transcript of a 47324 gene, 6) aberrant modification of a 47324 gene, such as of the methylation pattern of the genomic DNA, 7) the presence of a non-wild type splicing pattern of a messenger RNA transcript of a 47324 gene, 8) a non-wild type level of a 47324-protein, 9) allelic loss of a 47324 gene, and 10) inappropriate post-translational modification of a 47324-protein.

[0300] An alteration can be detected without a probe/primer in a polymerase chain reaction, such as anchor PCR or RACE PCR, or, alternatively, in a ligation chain reaction (LCR), the latter of which can be particularly useful for detecting point mutations in the 47324-gene. This method can include the steps of collecting a sample of cells from a subject, isolating nucleic acid (e.g., genomic, mRNA or both) from the sample, contacting the nucleic acid sample with one or more primers which specifically hybridize to a 47324 gene under conditions such that hybridization and amplification of the 47324-gene (if present) occurs, and detecting the presence or absence of an amplification product, or detecting the size of the amplification product and comparing the length to a control sample. It is anticipated that PCR and/or LCR may be desirable to use as a preliminary amplification step in conjunction with any of the techniques used for detecting mutations described herein.

[0301] Alternative amplification methods include: self sustained sequence replication (Guatelli, J. C. et al., (1990) *Proc. Natl. Acad. Sci. USA* 87:1874-1878), transcriptional amplification system (Kwoh, D. Y. et al., (1989) *Proc. Natl. Acad. Sci. USA* 86:1173-1177), Q-Beta Replicase (Lizardi, P. M. et al., (1988) *Bio-Technology* 6:1197), or other nucleic acid amplification methods, followed by the detection of the amplified molecules using techniques known to those of skill in the art.

[0302] In another embodiment, mutations in a 47324 gene from a sample cell can be identified by detecting alterations in restriction enzyme cleavage patterns. For example,

sample and control DNA is isolated, amplified (optionally), digested with one or more restriction endonucleases, and fragment length sizes are determined, e.g., by gel electrophoresis and compared. Differences in fragment length sizes between sample and control DNA indicates mutations in the sample DNA. Moreover, the use of sequence specific ribozymes (see, for example, U.S. Pat. No. 5,498,531) can be used to score for the presence of specific mutations by development or loss of a ribozyme cleavage site.

[0303] In other embodiments, genetic mutations in 47324 can be identified by hybridizing a sample and control nucleic acids, e.g., DNA or RNA, two-dimensional arrays, e.g., chip based arrays. Such arrays include a plurality of addresses, each of which is positionally distinguishable from the other. A different probe is located at each address of the plurality. The arrays can have a high density of addresses, e.g., can contain hundreds or thousands of oligonucleotide probes (Cronin, M. T. et al., (1996) *Human Mutation* 7: 244-255; Kozal, M. J. et al., (1996) *Nature Medicine* 2:753-759). For example, genetic mutations in 47324 can be identified in two dimensional arrays containing light-generated DNA probes as described in Cronin, M. T. et al., supra. Briefly, a first hybridization array of probes can be used to scan through long stretches of DNA in a sample and control to identify base changes between the sequences by making linear arrays of sequential overlapping probes. This step allows the identification of point mutations. This step is followed by a second hybridization array that allows the characterization of specific mutations by using smaller, specialized probe arrays complementary to all variants or mutations detected. Each mutation array is composed of parallel probe sets, one complementary to the wild-type gene and the other complementary to the mutant gene.

[0304] In yet another embodiment, any of a variety of sequencing reactions known in the art can be used to directly sequence the 47324 gene and detect mutations by comparing the sequence of the sample 47324 with the corresponding wild-type (control) sequence. Automated sequencing procedures can be utilized when performing the diagnostic assays ((1995) *Biotechniques* 19:448), including sequencing by mass spectrometry.

[0305] Other methods for detecting mutations in the 47324 gene include methods in which protection from cleavage agents is used to detect mismatched bases in RNA/RNA or RNA/DNA heteroduplexes (Myers et al., (1985) *Science* 230:1242; Cotton et al., (1988) *Proc. Natl. Acad. Sci. USA* 85:4397; Saleeba et al., (1992) *Methods Enzymol.* 217:286-295).

[0306] In still another embodiment, the mismatch cleavage reaction employs one or more proteins that recognize mismatched base pairs in double-stranded DNA (so called "DNA mismatch repair" enzymes) in defined systems for detecting and mapping point mutations in 47324 cDNAs obtained from samples of cells. For example, the mutY enzyme of *E. coli* cleaves A at G/A mismatches and the thymidine DNA glycosylase from HeLa cells cleaves T at G/T mismatches (Hsu et al., (1994) *Carcinogenesis* 15:1657-1662; U.S. Pat. No. 5,459,039).

[0307] In other embodiments, alterations in electrophoretic mobility will be used to identify mutations in 47324 genes. For example, single strand conformation polymorphism (SSCP) may be used to detect differences in electro-

phoretic mobility between mutant and wild type nucleic acids (Orita et al., (1989) *Proc. Natl. Acad. Sci. USA*: 86:2766, see also Cotton, (1993) *Mutat. Res.* 285:125-144; and Hayashi, (1992) *Genet. Anal. Tech. Appl.* 9:73-79). Single-stranded DNA fragments of sample and control 47324 nucleic acids will be denatured and allowed to renature. The secondary structure of single-stranded nucleic acids varies according to sequence, the resulting alteration in electrophoretic mobility enables the detection of even a single base change. The DNA fragments may be labeled or detected with labeled probes. The sensitivity of the assay may be enhanced by using RNA (rather than DNA), in which the secondary structure is more sensitive to a change in sequence. In a preferred embodiment, the subject method utilizes heteroduplex analysis to separate double stranded heteroduplex molecules on the basis of changes in electrophoretic mobility (Keen et al., (1991) *Trends Genet.* 7:5).

[0308] In yet another embodiment, the movement of mutant or wild-type fragments in polyacrylamide gels containing a gradient of denaturant is assayed using denaturing gradient gel electrophoresis (DGGE) (Myers et al., (1985) *Nature* 313:495). When DGGE is used as the method of analysis, DNA will be modified to insure that it does not completely denature, for example by adding a GC clamp of approximately 40 bp of high-melting GC-rich DNA by PCR. In a further embodiment, a temperature gradient is used in place of a denaturing gradient to identify differences in the mobility of control and sample DNA (Rosenbaum and Reissner, (1987) *Biophys. Chem.* 265:12753).

[0309] Examples of other techniques for detecting point mutations include, but are not limited to, selective oligonucleotide hybridization, selective amplification, or selective primer extension (Saiki et al., (1986) *Nature* 324:163; Saiki et al., (1989) *Proc. Natl. Acad. Sci. USA* 86:6230).

[0310] Alternatively, allele specific amplification technology which depends on selective PCR amplification may be used in conjunction with the instant invention. Oligonucleotides used as primers for specific amplification may carry the mutation of interest in the center of the molecule (so that amplification depends on differential hybridization) (Gibbs et al., (1989) *Nucleic Acids Res.* 17:2437-2448) or at the extreme 3' end of one primer where, under appropriate conditions, mismatch can prevent, or reduce polymerase extension (Prossner, (1993) *Tibtech* 11:238). In addition it maybe desirable to introduce a novel restriction site in the region of the mutation to create cleavage-based detection (Gasparini et al., (1992) *Mol. Cell Probes* 6:1). It is anticipated that in certain embodiments amplification may also be performed using Taq ligase for amplification (Barany, (1991) *Proc. Natl. Acad. Sci. USA* 88:189). In such cases, ligation will occur only if there is a perfect match at the 3' end of the 5' sequence making it possible to detect the presence of a known mutation at a specific site by looking for the presence or absence of amplification.

[0311] The methods described herein may be performed, for example, by utilizing pre-packaged diagnostic kits comprising at least one probe nucleic acid or antibody reagent described herein, which may be conveniently used, e.g., in-clinical settings to diagnose patients exhibiting symptoms or family history of a disease or illness involving a 47324 gene.

[0312] Use of 47324 Molecules as Surrogate Markers

[0313] The 47324 molecules of the invention are also useful as markers of disorders or disease states, as markers for precursors of disease states, as markers for predisposition of disease states, as markers of drug activity, or as markers of the pharmacogenomic profile of a subject. Using the methods described herein, the presence, absence and/or quantity of the 47324 molecules of the invention may be detected, and may be correlated with one or more biological states in vivo. For example, the 47324 molecules of the invention may serve as surrogate markers for one or more disorders or disease states or for conditions leading up to disease states. As used herein, a "surrogate marker" is an objective biochemical marker which correlates with the absence or presence of a disease or disorder, or with the progression of a disease or disorder (e.g., with the presence or absence of a tumor). The presence or quantity of such markers is independent of the disease. Therefore, these markers may serve to indicate whether a particular course of treatment is effective in lessening a disease state or disorder. Surrogate markers are of particular use when the presence or extent of a disease state or disorder is difficult to assess through standard methodologies (e.g., early stage tumors), or when an assessment of disease progression is desired before a potentially dangerous clinical endpoint is reached (e.g., an assessment of cardiovascular disease may be made using cholesterol levels as a surrogate marker, and an analysis of HIV infection may be made using HIV RNA levels as a surrogate marker, well in advance of the undesirable clinical outcomes of myocardial infarction or fully-developed AIDS). Examples of the use of surrogate markers in the art include: Koomen et al. (2000) *J. Mass. Spectrom.* 35: 258-264; and James (1994) *AIDS Treatment News Archive* 209.

[0314] The 47324 molecules of the invention are also useful as pharmacodynamic markers. As used herein, a "pharmacodynamic marker" is an objective biochemical marker which correlates specifically with drug effects. The presence or quantity of a pharmacodynamic marker is not related to the disease state or disorder for which the drug is being administered; therefore, the presence or quantity of the marker is indicative of the presence or activity of the drug in a subject. For example, a pharmacodynamic marker may be indicative of the concentration of the drug in a biological tissue, in that the marker is either expressed or transcribed or not expressed or transcribed in that tissue in relationship to the level of the drug. In this fashion, the distribution or uptake of the drug may be monitored by the pharmacodynamic marker. Similarly, the presence or quantity of the pharmacodynamic marker may be related to the presence or quantity of the metabolic product of a drug, such that the presence or quantity of the marker is indicative of the relative breakdown rate of the drug in vivo. Pharmacodynamic markers are of particular use in increasing the sensitivity of detection of drug effects, particularly when the drug is administered in low doses. Since even a small amount of a drug may be sufficient to activate multiple rounds of marker (e.g., a 47324 marker) transcription or expression, the amplified marker may be in a quantity which is more readily detectable than the drug itself. Also, the marker may be more easily detected due to the nature of the marker itself; for example, using the methods described herein, anti-47324 antibodies may be employed in an immune-based detection system for a 47324 protein marker, or 47324-specific radio-

labeled probes may be used to detect a 47324 mRNA marker. Furthermore, the use of a pharmacodynamic marker may offer mechanism-based prediction of risk due to drug treatment beyond the range of possible direct observations. Examples of the use of pharmacodynamic markers in the art include: Matsuda et al. U.S. Pat. No. 6,033,862; Hattis et al. (1991) *Env. Health Perspect.* 90: 229-238; Schentag (1999) *Am. J. Health-Syst. Pharm.* 56 Suppl. 3: S21-S24; and Nicolau (1999) *Am. J. Health-Syst. Pharm.* 56 Suppl. 3: S16-S20.

[0315] The 47324 molecules of the invention are also useful as pharmacogenomic markers. As used herein, a "pharmacogenomic marker" is an objective biochemical marker which correlates with a specific clinical drug response or susceptibility in a subject (see, e.g., McLeod et al. (1999) *Eur. J. Cancer* 35(12): 1650-1652). The presence or quantity of the pharmacogenomic marker is related to the predicted response of the subject to a specific drug or class of drugs prior to administration of the drug. By assessing the presence or quantity of one or more pharmacogenomic markers in a subject, a drug therapy which is most appropriate for the subject, or which is predicted to have a greater degree of success, may be selected. For example, based on the presence or quantity of RNA, or protein (e.g., 47324 protein or RNA) for specific tumor markers in a subject, a drug or course of treatment may be selected that is optimized for the treatment of the specific tumor likely to be present in the subject. Similarly, the presence or absence of a specific sequence mutation in 47324 DNA may correlate 47324 drug response. The use of pharmacogenomic markers therefore permits the application of the most appropriate treatment for each subject without having to administer the therapy.

[0316] Pharmaceutical Compositions

[0317] The nucleic acid and polypeptides, fragments thereof, as well as anti-47324 antibodies (also referred to herein as "active compounds") of the invention can be incorporated into pharmaceutical compositions. Such compositions typically include the nucleic acid molecule, protein, or antibody and a pharmaceutically acceptable carrier. As used herein the language "pharmaceutically acceptable carrier" includes solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents, and the like, compatible with pharmaceutical administration. Supplementary active compounds can also be incorporated into the compositions.

[0318] A pharmaceutical composition is formulated to be compatible with its intended route of administration. Examples of routes of administration include parenteral, e.g., intravenous, intradermal, subcutaneous, oral (e.g., inhalation), transdermal (topical), transmucosal, and rectal administration. Solutions or suspensions used for parenteral, intradermal, or subcutaneous application can include the following components: a sterile diluent such as water for injection, saline solution, fixed oils, polyethylene glycols, glycerine, propylene glycol or other synthetic solvents; antibacterial agents such as benzyl alcohol or methyl parabens; antioxidants such as ascorbic acid or sodium bisulfite; chelating agents such as ethylenediaminetetraacetic acid; buffers such as acetates, citrates or phosphates and agents for the adjustment of tonicity such as sodium chloride or dextrose. pH can be adjusted with acids or bases, such as hydrochloric acid or sodium hydroxide. The parenteral

preparation can be enclosed in ampoules, disposable syringes or multiple dose vials made of glass or plastic.

[0319] Pharmaceutical compositions suitable for injectable use include sterile aqueous solutions (where water soluble) or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersion. For intravenous administration, suitable carriers include physiological saline, bacteriostatic water, Cremophor EL™ (BASF, Parsippany, N.J.) or phosphate buffered saline (PBS). In all cases, the composition must be sterile and should be fluid to the extent that easy syringability exists. It should be stable under the conditions of manufacture and storage and must be preserved against the contaminating action of microorganisms such as bacteria and fungi. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (for example, glycerol, propylene glycol, and liquid polyethylene glycol, and the like), and suitable mixtures thereof. The proper fluidity can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of dispersion and by the use of surfactants. Prevention of the action of microorganisms can be achieved by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, ascorbic acid, thimerosal, and the like. In many cases, it will be preferable to include isotonic agents, for example, sugars, polyalcohols such as manitol, sorbitol, sodium chloride in the composition. Prolonged absorption of the injectable compositions can be brought about by including in the composition an agent which delays absorption, for example, aluminum monostearate and gelatin.

[0320] Sterile injectable solutions can be prepared by incorporating the active compound in the required amount in an appropriate solvent with one or a combination of ingredients enumerated above, as required, followed by filtered sterilization. Generally, dispersions are prepared by incorporating the active compound into a sterile vehicle which contains a basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum drying and freeze-drying which yields a powder of the active ingredient plus any additional desired ingredient from a previously sterile-filtered solution thereof.

[0321] Oral compositions generally include an inert diluent or an edible carrier. For the purpose of oral therapeutic administration, the active compound can be incorporated with excipients and used in the form of tablets, troches, or capsules, e.g., gelatin capsules. Oral compositions can also be prepared using a fluid carrier for use as a mouthwash. Pharmaceutically compatible binding agents, and/or adjuvant materials can be included as part of the composition. The tablets, pills, capsules, troches and the like can contain any of the following ingredients, or compounds of a similar nature: a binder such as microcrystalline cellulose, gum tragacanth or gelatin; an excipient such as starch or lactose, a disintegrating agent such as alginic acid, Primogel, or corn starch; a lubricant such as magnesium stearate or Sterotes; a glidant such as colloidal silicon dioxide; a sweetening agent such as sucrose or saccharin; or a flavoring agent such as peppermint, methyl salicylate, or orange flavoring.

[0322] For administration by inhalation, the compounds are delivered in the form of an aerosol spray from pressured

container or dispenser which contains a suitable propellant, e.g., a gas such as carbon dioxide, or a nebulizer.

[0323] Systemic administration can also be by transmucosal or transdermal means. For transmucosal or transdermal administration, penetrants appropriate to the barrier to be permeated are used in the formulation. Such penetrants are generally known in the art, and include, for example, for transmucosal administration, detergents, bile salts, and fusidic acid derivatives. Transmucosal administration can be accomplished through the use of nasal sprays or suppositories. For transdermal administration, the active compounds are formulated into ointments, salves, gels, or creams as generally known in the art.

[0324] The compounds can also be prepared in the form of suppositories (e.g., with conventional suppository bases such as cocoa butter and other glycerides) or retention enemas for rectal delivery.

[0325] In one embodiment, the active compounds are prepared with carriers that will protect the compound against rapid elimination from the body, such as a controlled release formulation, including implants and microencapsulated delivery systems. Biodegradable, biocompatible polymers can be used, such as ethylene vinyl acetate, polyanhydrides, polyglycolic acid, collagen, polyorthoesters, and polylactic acid. Methods for preparation of such formulations will be apparent to those skilled in the art. The materials can also be obtained commercially from Alza Corporation and Nova Pharmaceuticals, Inc. Liposomal suspensions (including liposomes targeted to infected cells with monoclonal antibodies to viral antigens) can also be used as pharmaceutically acceptable carriers. These can be prepared according to methods known to those skilled in the art, for example, as described in U.S. Pat. No. 4,522,811.

[0326] It is advantageous to formulate oral or parenteral compositions in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form as used herein refers to physically discrete units suited as unitary dosages for the subject to be treated; each unit containing a predetermined quantity of active compound calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier.

[0327] Toxicity and therapeutic efficacy of such compounds can be determined by standard pharmaceutical procedures in cell cultures or experimental animals, e.g., for determining the LD₅₀ (the dose lethal to 50% of the population) and the ED₅₀ (the dose therapeutically effective in 50% of the population). The dose ratio between toxic and therapeutic effects is the therapeutic index and it can be expressed as the ratio LD₅₀/ED₅₀. Compounds which exhibit high therapeutic indices are preferred. While compounds that exhibit toxic side effects may be used, care should be taken to design a delivery system that targets such compounds to the site of affected tissue in order to minimize potential damage to uninfected cells and, thereby, reduce side effects.

[0328] The data obtained from the cell culture assays and animal studies can be used in formulating a range of dosage for use in humans. The dosage of such compounds lies preferably within a range of circulating concentrations that include the ED₅₀ with little or no toxicity. The dosage may vary within this range depending upon the dosage form

employed and the route of administration utilized. For any compound used in the method of the invention, the therapeutically effective dose can be estimated initially from cell culture assays. A dose may be formulated in animal models to achieve a circulating plasma concentration range that includes the IC_{50} (i.e., the concentration of the test compound which achieves a half-maximal inhibition of symptoms) as determined in cell culture. Such information can be used to more accurately determine useful doses in humans. Levels in plasma may be measured, for example, by high performance liquid chromatography.

[0329] As defined herein, a therapeutically effective amount of protein or polypeptide (i.e., an effective dosage) ranges from about 0.001 to 30 mg/kg body weight, preferably about 0.01 to 25 mg/kg body weight, more preferably about 0.1 to 20 mg/kg body weight, and even more preferably about 1 to 10 mg/kg, 2 to 9 mg/kg, 3 to 8 mg/kg, 4 to 7 mg/kg, or 5 to 6 mg/kg body weight. The protein or polypeptide can be administered one time per week for between about 1 to 10 weeks, preferably between 2 to 8 weeks, more preferably between about 3 to 7 weeks, and even more preferably for about 4, 5, or 6 weeks. The skilled artisan will appreciate that certain factors may influence the dosage and timing required to effectively treat a subject, including but not limited to the severity of the disease or disorder, previous treatments, the general health and/or age of the subject, and other diseases present. Moreover, treatment of a subject with a therapeutically effective amount of a protein, polypeptide, or antibody can include a single treatment or, preferably, can include a series of treatments.

[0330] For antibodies, the preferred dosage is 0.1 mg/kg of body weight (generally 10 mg/kg to 20 mg/kg). If the antibody is to act in the brain, a dosage of 50 mg/kg to 100 mg/kg is usually appropriate. Generally, partially human antibodies and fully human antibodies have a longer half-life within the human body than other antibodies. Accordingly, lower dosages and less frequent administration is often possible. Modifications such as lipidation can be used to stabilize antibodies and to enhance uptake and tissue penetration (e.g., into the brain). A method for lipidation of antibodies is described by Cruikshank et al., ((1997) *J. Acquired Immune Deficiency Syndromes and Human Retrovirology* 14:193).

[0331] The present invention encompasses agents which modulate expression or activity. An agent may, for example, be a small molecule. For example, such small molecules include, but are not limited to, peptides, peptidomimetics (e.g., peptoids), amino acids, amino acid analogs, polynucleotides, polynucleotide analogs, nucleotides, nucleotide analogs, organic or inorganic compounds (i.e., including heteroorganic and organometallic compounds) having a molecular weight less than about 10,000 grams per mole, organic or inorganic compounds having a molecular weight less than about 5,000 grams per mole, organic or inorganic compounds having a molecular weight less than about 1,000 grams per mole, organic or inorganic compounds having a molecular weight less than about 500 grams per mole, and salts, esters, and other pharmaceutically acceptable forms of such compounds.

[0332] Exemplary doses include milligram or microgram amounts of the small molecule per kilogram of subject or sample weight (e.g., about 1 microgram per kilogram to

about 500 milligrams per kilogram, about 100 micrograms per kilogram to about 5 milligrams per kilogram, or about 1 microgram per kilogram to about 50 micrograms per kilogram. It is furthermore understood that appropriate doses of a small molecule depend upon the potency of the small molecule with respect to the expression or activity to be modulated. When one or more of these small molecules is to be administered to an animal (e.g., a human) in order to modulate expression or activity of a polypeptide or nucleic acid of the invention, a physician, veterinarian, or researcher may, for example, prescribe a relatively low dose at first, subsequently increasing the dose until an appropriate response is obtained. In addition, it is understood that the specific dose level for any particular animal subject will depend upon a variety of factors including the activity of the specific compound employed, the age, body weight, general health, gender, and diet of the subject, the time of administration, the route of administration, the rate of excretion, any drug combination, and the degree of expression or activity to be modulated.

[0333] An antibody (or fragment thereof) may be conjugated to a therapeutic moiety such as a cytotoxin, a therapeutic agent or a radioactive metal ion. A cytotoxin or cytotoxic agent includes any agent that is detrimental to cells. Examples include taxol, cytochalasin B, gramicidin D, ethidium bromide, emetine, mitomycin, etoposide, tenoposide, vincristine, vinblastine, colchicin, doxorubicin, daunorubicin, dihydroxy anthracin dione, mitoxantrone, mithramycin, actinomycin D, 1-dehydrotestosterone, glucocorticoids, procaine, tetracaine, lidocaine, propranolol, and puromycin and analogs or homologs thereof. Therapeutic agents include, but are not limited to, antimetabolites (e.g., methotrexate, 6-mercaptopurine, 6-thioguanine, cytarabine, 5-fluorouracil decarbazine), alkylating agents (e.g., mechlorethamine, thiopepa chlorambucil, melphalan, carmustine (BSNU) and lomustine (CCNU), cyclophosphamide, busulfan, dibromomannitol, streptozotocin, mitomycin C, and cis-dichlorodiamine platinum (II) (DDP) cisplatin), anthracyclines (e.g., daunorubicin (formerly daunomycin) and doxorubicin), antibiotics (e.g., dactinomycin (formerly actinomycin), bleomycin, mithramycin, and anthramycin (AMC)), and anti-mitotic agents (e.g., vincristine and vinblastine).

[0334] The conjugates of the invention can be used for modifying a given biological response, the drug moiety is not to be construed as limited to classical chemical therapeutic agents. For example, the drug moiety may be a protein or polypeptide possessing a desired biological activity. Such proteins may include, for example, a toxin such as abrin, ricin A, pseudomonas exotoxin, or diphtheria toxin; a protein such as tumor necrosis factor, .alpha.-interferon, .beta.-interferon, nerve growth factor, platelet derived growth factor, tissue plasminogen activator; or, biological response modifiers such as, for example, lymphokines, interleukin-1 ("IL-1"), interleukin-2 ("IL-2"), interleukin-6 ("IL-6"), granulocyte macrophage colony stimulating factor ("GM-CSF"), granulocyte colony stimulating factor ("G-CSF"), or other growth factors.

[0335] Alternatively, an antibody can be conjugated to a second antibody to form an antibody heteroconjugate as described by Segal in U.S. Pat. No. 4,676,980.

[0336] The nucleic acid molecules of the invention can be inserted into vectors and used as gene therapy vectors. Gene

therapy vectors can be delivered to a subject by, for example, intravenous injection, local administration (see U.S. Pat. No. 5,328,470) or by stereotactic injection (see e.g., Chen et al., (1994) *Proc. Natl. Acad. Sci. USA* 91:3054-3057). The pharmaceutical preparation of the gene therapy vector can include the gene therapy vector in an acceptable diluent, or can comprise a slow release matrix in which the gene delivery vehicle is imbedded. Alternatively, where the complete gene delivery vector can be produced intact from recombinant cells, e.g., retroviral vectors, the pharmaceutical preparation can include one or more cells which produce the gene delivery system.

[0337] The pharmaceutical compositions can be included in a container, pack, or dispenser together with instructions for administration.

[0338] Methods of Treatment:

[0339] The present invention provides for both prophylactic and therapeutic methods of treating a subject at risk of (or susceptible to) a disorder or having a disorder associated with aberrant or unwanted 47324 expression or activity. With regards to both prophylactic and therapeutic methods of treatment, such treatments may be specifically tailored or modified, based on knowledge obtained from the field of pharmacogenomics. As used herein, the term "treatment" is defined as the application or administration of a therapeutic agent to a patient, or application or administration of a therapeutic agent to an isolated tissue or cell line from a patient, who has a disease, a symptom of disease or a predisposition toward a disease, with the purpose to cure, heal, alleviate, relieve, alter, remedy, ameliorate, improve or affect the disease, the symptoms of disease or the predisposition toward disease. A therapeutic agent includes, but is not limited to, small molecules, peptides, antibodies, ribozymes and antisense oligonucleotides. "Pharmacogenomics", as used herein, refers to the application of genomics technologies such as gene sequencing, statistical genetics, and gene expression analysis to drugs in clinical development and on the market. More specifically, the term refers the study of how a patient's genes determine his or her response to a drug (e.g., a patient's "drug response phenotype", or "drug response genotype".) Thus, another aspect of the invention provides methods for tailoring an individual's prophylactic or therapeutic treatment with either the 47324 molecules of the present invention or 47324 modulators according to that individual's drug response genotype. Pharmacogenomics allows a clinician or physician to target prophylactic or therapeutic treatments to patients who will most benefit from the treatment and to avoid treatment of patients who will experience toxic drug-related side effects.

[0340] In one aspect, the invention provides a method for preventing in a subject, a disease or condition associated with an aberrant or unwanted 47324 expression or activity, by administering to the subject a 47324 or an agent which modulates 47324 expression or at least one 47324 activity. Subjects at risk for a disease which is caused or contributed to by aberrant or unwanted 47324 expression or activity can be identified by, for example, any or a combination of diagnostic or prognostic assays as described herein. Administration of a prophylactic agent can occur prior to the manifestation of symptoms characteristic of the 47324 aberrance, such that a disease or disorder is prevented or, alternatively, delayed in its progression. Depending on the

type of 47324 aberrance, for example, a 47324, 47324 agonist or 47324 antagonist agent can be used for treating the subject. The appropriate agent can be determined based on screening assays described herein.

[0341] It is possible that some 47324 disorders can be caused, at least in part, by an abnormal level of gene product, or by the presence of a gene product exhibiting abnormal activity. As such, the reduction in the level and/or activity of such gene products would bring about the amelioration of disorder symptoms.

[0342] As discussed, successful treatment of 47324 disorders can be brought about by techniques that serve to inhibit the expression or activity of target gene products. For example, compounds, e.g., an agent identified using an assays described above, that proves to exhibit negative modulatory activity, can be used in accordance with the invention to prevent and/or ameliorate symptoms of 47324 disorders. Such molecules can include, but are not limited to peptides, phosphopeptides, small organic or inorganic molecules, or antibodies (including, for example, polyclonal, monoclonal, humanized, anti-idiotypic, chimeric or single chain antibodies, and FAb, F(ab')₂ and FAb expression library fragments, scFV molecules, and epitope-binding fragments thereof).

[0343] Further, antisense and ribozyme molecules that inhibit expression of the target gene can also be used in accordance with the invention to reduce the level of target gene expression, thus effectively reducing the level of target gene activity. Still further, triple helix molecules can be utilized in reducing the level of target gene activity. Antisense, ribozyme and triple helix molecules are discussed above.

[0344] It is possible that the use of antisense, ribozyme, and/or triple helix molecules to reduce or inhibit mutant gene expression can also reduce or inhibit the transcription (triple helix) and/or translation (antisense, ribozyme) of mRNA produced by normal target gene alleles, such that the concentration of normal target gene product present can be lower than is necessary for a normal phenotype. In such cases, nucleic acid molecules that encode and express target gene polypeptides exhibiting normal target gene activity can be introduced into cells via gene therapy method. Alternatively, in instances in that the target gene encodes an extracellular protein, it can be preferable to co-administer normal target gene protein into the cell or tissue in order to maintain the requisite level of cellular or tissue target gene activity.

[0345] Another method by which nucleic acid molecules may be utilized in treating or preventing a disease characterized by 47324 expression is through the use of aptamer molecules specific for 47324 protein. Aptamers are nucleic acid molecules having a tertiary structure which permits them to specifically bind to protein ligands (see, e.g., Osborne, et al., *Curr. Opin. Chem. Biol.* 1997, 1(1): 5-9; and Patel, D. J., *Curr. Opin. Chem. Biol.* Jun. 1, 1997 (1):32-46). Since nucleic acid molecules may in many cases be more conveniently introduced into target cells than therapeutic protein molecules may be, aptamers offer a method by which 47324 protein activity may be specifically decreased without the introduction of drugs or other molecules which may have pluripotent effects.

[0346] Antibodies can be generated that are both specific for target gene product and that reduce target gene product

activity. Such antibodies may, therefore, be administered in instances whereby negative modulatory techniques are appropriate for the treatment of 47324 disorders. For a description of antibodies, see the Antibody section above.

[0347] In circumstances wherein injection of an animal or a human subject with a 47324 protein or epitope for stimulating antibody production is harmful to the subject, it is possible to generate an immune response against 47324 through the use of anti-idiotypic antibodies (see, for example, Herlyn, D., *Ann. Med.* 1999;31(1):66-78; and Bhattacharya-Chatterjee, M., and Foon, K. A., *Cancer Treat. Res.* 1998;94:51-68). If an anti-idiotypic antibody is introduced into a mammal or human subject, it should stimulate the production of anti-anti-idiotypic antibodies, which should be specific to the 47324 protein. Vaccines directed to a disease characterized by 47324 expression may also be generated in this fashion.

[0348] In instances where the target antigen is intracellular and whole antibodies are used, internalizing antibodies may be preferred. Lipofectin or liposomes can be used to deliver the antibody or a fragment of the Fab region that binds to the target antigen into cells. Where fragments of the antibody are used, the smallest inhibitory fragment that binds to the target antigen is preferred. For example, peptides having an amino acid sequence corresponding to the Fv region of the antibody can be used. Alternatively, single chain neutralizing antibodies that bind to intracellular target antigens can also be administered. Such single chain antibodies can be administered, for example, by expressing nucleotide sequences encoding single-chain antibodies within the target cell population (see e.g., Marasco et al., (1993, *Proc. Natl. Acad. Sci. USA* 90:7889-7893).

[0349] The identified compounds that inhibit target gene expression, synthesis and/or activity can be administered to a patient at therapeutically effective doses to prevent, treat or ameliorate 47324 disorders. A therapeutically effective dose refers to that amount of the compound sufficient to result in amelioration of symptoms of the disorders.

[0350] Toxicity and therapeutic efficacy of such compounds can be determined by standard pharmaceutical procedures in cell cultures or experimental animals, e.g., for determining the LD₅₀ (the dose lethal to 50% of the population) and the ED₅₀ (the dose therapeutically effective in 50% of the population). The dose ratio between toxic and therapeutic effects is the therapeutic index and it can be expressed as the ratio LD₅₀/ED₅₀. Compounds that exhibit large therapeutic indices are preferred. While compounds that exhibit toxic side effects can be used, care should be taken to design a delivery system that targets such compounds to the site of affected tissue in order to minimize potential damage to uninfected cells and, thereby, reduce side effects.

[0351] The data obtained from the cell culture assays and animal studies can be used in formulating a range of dosage for use in humans. The dosage of such compounds lies preferably within a range of circulating concentrations that include the ED₅₀ with little or no toxicity. The dosage can vary within this range depending upon the dosage form employed and the route of administration utilized. For any compound used in the method of the invention, the therapeutically effective dose can be estimated initially from cell culture assays. A dose can be formulated in animal models

to achieve a circulating plasma concentration range that includes the IC₅₀ (i.e., the concentration of the test compound that achieves a half-maximal inhibition of symptoms) as determined in cell culture. Such information can be used to more accurately determine useful doses in humans. Levels in plasma can be measured, for example, by high performance liquid chromatography.

[0352] Another example of determination of effective dose for an individual is the ability to directly assay levels of "free" and "bound" compound in the serum of the test subject. Such assays may utilize antibody mimics and/or "biosensors" that have been created through molecular imprinting techniques. The compound which is able to modulate 47324 activity is used as a template, or "imprinting molecule", to spatially organize polymerizable monomers prior to their polymerization with catalytic reagents. The subsequent removal of the imprinted molecule leaves a polymer matrix which contains a repeated "negative image" of the compound and is able to selectively rebind the molecule under biological assay conditions. A detailed review of this technique can be seen in Ansell, R. J. et al., (1996) *Current Opinion in Biotechnology* 7:89-94 and in Shea, K. J., (1994) *Trends in Polymer Science* 2:166-173. Such "imprinted" affinity matrixes are amenable to ligand-binding assays, whereby the immobilized monoclonal antibody component is replaced by an appropriately imprinted matrix. An example of the use of such matrixes in this way can be seen in Vlatakis, G. et al., (1993) *Nature* 361:645-647. Through the use of isotope-labeling, the "free" concentration of compound which modulates the expression or activity of 47324 can be readily monitored and used in calculations of IC₅₀.

[0353] Such "imprinted" affinity matrixes can also be designed to include fluorescent groups whose photon-emitting properties measurably change upon local and selective binding of target compound. These changes can be readily assayed in real time using appropriate fiberoptic devices, in turn allowing the dose in a test subject to be quickly optimized based on its individual IC₅₀. A rudimentary example of such a "biosensor" is discussed in Kriz, D. et al., (1995) *Analytical Chemistry* 67:2142-2144.

[0354] Another aspect of the invention pertains to methods of modulating 47324 expression or activity for therapeutic purposes. Accordingly, in an exemplary embodiment, the modulatory method of the invention involves contacting a cell with a 47324 or agent that modulates one or more of the activities of 47324 protein activity associated with the cell. An agent that modulates 47324 protein activity can be an agent as described herein, such as a nucleic acid or a protein, a naturally-occurring target molecule of a 47324 protein (e.g., a 47324 substrate or receptor), a 47324 antibody, a 47324 agonist or antagonist, a peptidomimetic of a 47324 agonist or antagonist, or other small molecule.

[0355] In one embodiment, the agent stimulates one or more 47324 activities. Examples of such stimulatory agents include active 47324 protein and a nucleic acid molecule encoding 47324. In another embodiment, the agent inhibits one or more 47324 activities. Examples of such inhibitory agents include antisense 47324 nucleic acid molecules, anti-47324 antibodies, and 47324 inhibitors. These modulatory methods can be performed in vitro (e.g., by culturing the cell with the agent) or, alternatively, in vivo (e.g., by

administering the agent to a subject). As such, the present invention provides methods of treating an individual afflicted with a disease or disorder characterized by aberrant or unwanted expression or activity of a 47324 protein or nucleic acid molecule. In one embodiment, the method involves administering an agent (e.g., an agent identified by a screening assay described herein), or combination of agents that modulates (e.g., upregulates or downregulates) 47324 expression or activity. In another embodiment, the method involves administering a 47324 protein or nucleic acid molecule as therapy to compensate for reduced, aberrant, or unwanted 47324 expression or activity.

[0356] Stimulation of 47324 activity is desirable in situations in which 47324 is abnormally downregulated and/or in which increased 47324 activity is likely to have a beneficial effect. For example, stimulation of 47324 activity is desirable in situations in which a 47324 is downregulated and/or in which increased 47324 activity is likely to have a beneficial effect. Likewise, inhibition of 47324 activity is desirable in situations in which 47324 is abnormally upregulated and/or in which decreased 47324 activity is likely to have a beneficial effect.

[0357] The 47324 molecules can act as novel diagnostic targets and therapeutic agents for controlling one or more of cellular proliferative and/or differentiative disorders, cardiovascular disorders, liver disorders, brain disorders, blood vessel disorders, hematopoietic disorders and platelet disorders, as described above, as well as disorders associated with bone metabolism, hormonal disorders, immune disorders, heart disorders, and viral diseases.

[0358] Aberrant expression and/or activity of 47324 molecules may mediate disorders associated with bone metabolism. "Bone metabolism" refers to direct or indirect effects in the formation or degeneration of bone structures, e.g., bone formation, bone resorption, etc., which may ultimately affect the concentrations in serum of calcium and phosphate. This term also includes activities mediated by 47324 molecules effects in bone cells, e.g. osteoclasts and osteoblasts, that may in turn result in bone formation and degeneration. For example, 47324 molecules may support different activities of bone resorbing osteoclasts such as the stimulation of differentiation of monocytes and mononuclear phagocytes into osteoclasts. Accordingly, 47324 molecules that modulate the production of bone cells can influence bone formation and degeneration, and thus may be used to treat bone disorders. Examples of such disorders include, but are not limited to, osteoporosis, osteodystrophy, osteomalacia, rickets, osteitis fibrosa cystica, renal osteodystrophy, osteosclerosis, anti-convulsant treatment, osteopenia, fibrogenesis-imperfecta ossium, secondary hyperparathyroidism, hypoparathyroidism, hyperparathyroidism, cirrhosis, obstructive jaundice, drug induced metabolism, medullary carcinoma, chronic renal disease, rickets, sarcoidosis, glucocorticoid antagonism, malabsorption syndrome, steatorrhea, tropical sprue, idiopathic hypercalcemia and milk fever.

[0359] Ras family-associated or related disorders also include hormonal disorders, such as conditions or diseases in which the production and/or regulation of hormones in an organism is aberrant. Examples of such disorders and diseases include type I and type II diabetes mellitus, pituitary disorders (e.g., growth disorders), thyroid disorders (e.g.,

hypothyroidism or hyperthyroidism), and reproductive or fertility disorders (e.g., disorders which affect the organs of the reproductive system, e.g., the prostate gland, the uterus, or the vagina; disorders which involve an imbalance in the levels of a reproductive hormone in a subject; disorders affecting the ability of a subject to reproduce; and disorders affecting secondary sex characteristic development, e.g., adrenal hyperplasia).

[0360] Ras family-associated or related disorders also include immune disorders, such as autoimmune disorders or immune deficiency disorders, e.g., congenital X-linked infantile hypogammaglobulinemia, transient hypogammaglobulinemia, common variable immunodeficiency, selective IgA deficiency, chronic mucocutaneous candidiasis, or severe combined immunodeficiency.

[0361] Disorders involving the heart, include but are not limited to, heart failure, including but not limited to, cardiac hypertrophy, left-sided heart failure, and right-sided heart failure; ischemic heart disease, including but not limited to angina pectoris, myocardial infarction, chronic ischemic heart disease, and sudden cardiac death; hypertensive heart disease, including but not limited to, systemic (left-sided) hypertensive heart disease and pulmonary (right-sided) hypertensive heart disease; valvular heart disease, including but not limited to, valvular degeneration caused by calcification, such as calcific aortic stenosis, calcification of a congenitally bicuspid aortic valve, and mitral annular calcification, and myxomatous degeneration of the mitral valve (mitral valve prolapse), rheumatic fever and rheumatic heart disease, infective endocarditis, and noninfected vegetations, such as nonbacterial thrombotic endocarditis and endocarditis of systemic lupus erythematosus (Libman-Sacks disease), carcinoid heart disease, and complications of artificial valves; myocardial disease, including but not limited to dilated cardiomyopathy, hypertrophic cardiomyopathy, restrictive cardiomyopathy, and myocarditis; pericardial disease, including but not limited to, pericardial effusion and hemopericardium and pericarditis, including acute pericarditis and healed pericarditis, and rheumatoid heart disease; neoplastic heart disease, including but not limited to, primary cardiac tumors, such as myxoma, lipoma, papillary fibroelastoma, rhabdomyoma, and sarcoma, and cardiac effects of noncardiac neoplasms; congenital heart disease, including but not limited to, left-to-right shunts—late cyanosis, such as atrial septal defect, ventricular septal defect, patent ductus arteriosus, and atrioventricular septal defect, right-to-left shunt—early cyanosis, such as tetralogy of fallot, transposition of great arteries, truncus arteriosus, tricuspid atresia, and total anomalous pulmonary venous connection, obstructive congenital anomalies, such as coarctation of aorta, pulmonary stenosis and atresia, and aortic stenosis and atresia, and disorders involving cardiac transplantation.

[0362] Additionally, 47324 molecules may play an important role in the etiology of certain viral diseases, including but not limited to, Hepatitis B, Hepatitis C and Herpes Simplex Virus (HSV). Modulators of 47324 activity could be used to control viral diseases. The modulators can be used in the treatment and/or diagnosis of viral infected tissue or virus-associated tissue fibrosis, especially liver and liver fibrosis. Also, 47324 modulators can be used in the treatment and/or diagnosis of virus-associated carcinoma, especially hepatocellular cancer.

[0363] Pharmacogenomics

[0364] The 47324 molecules of the present invention, as well as agents, or modulators which have a stimulatory or inhibitory effect on 47324 activity (e.g., 47324 gene expression) as identified by a screening assay described herein can be administered to individuals to treat (prophylactically or therapeutically) 47324 associated disorders (e.g., Ras family related disorders) associated with aberrant or unwanted 47324 activity. In conjunction with such treatment, pharmacogenomics (i.e., the study of the relationship between an individual's genotype and that individual's response to a foreign compound or drug) may be considered. Differences in metabolism of therapeutics can lead to severe toxicity or therapeutic failure by altering the relation between dose and blood concentration of the pharmacologically active drug. Thus, a physician or clinician may consider applying knowledge obtained in relevant pharmacogenomics studies in determining whether to administer a 47324 molecule or 47324 modulator as well as tailoring the dosage and/or therapeutic regimen of treatment with a 47324 molecule or 47324 modulator.

[0365] Pharmacogenomics deals with clinically significant hereditary variations in the response to drugs due to altered drug disposition and abnormal action in affected persons. See, for example, Eichelbaum, M. et al. (1996) *Clin. Exp. Pharmacol. Physiol.* 23(10-11):983-985 and Linder, M. W. et al. (1997) *Clin. Chem.* 43(2):254-266. In general, two types of pharmacogenetic conditions can be differentiated. Genetic conditions transmitted as a single factor altering the way drugs act on the body (altered drug action) or genetic conditions transmitted as single factors altering the way the body acts on drugs (altered drug metabolism). These pharmacogenetic conditions can occur either as rare genetic defects or as naturally-occurring polymorphisms. For example, glucose-6-phosphate dehydrogenase deficiency (G6PD) is a common inherited enzymopathy in which the main clinical complication is haemolysis after ingestion of oxidant drugs (anti-malarials, sulfonamides, analgesics, nitrofurans) and consumption of fava beans.

[0366] One pharmacogenomics approach to identifying genes that predict drug response, known as "a genome-wide association", relies primarily on a high-resolution map of the human genome consisting of already known gene-related markers (e.g., a "bi-allelic" gene marker map which consists of 60,000-100,000 polymorphic or variable sites on the human genome, each of which has two variants.) Such a high-resolution genetic map can be compared to a map of the genome of each of a statistically significant number of patients taking part in a Phase II/III drug trial to identify markers associated with a particular observed drug response or side effect. Alternatively, such a high-resolution map can be generated from a combination of some ten million known single nucleotide polymorphisms (SNPs) in the human genome. As used herein, a "SNP" is a common alteration that occurs in a single nucleotide base in a stretch of DNA. For example, a SNP may occur once per every 1000 bases of DNA. A SNP may be involved in a disease process, however, the vast majority may not be disease-associated. Given a genetic map based on the occurrence of such SNPs, individuals can be grouped into genetic categories depending on a particular pattern of SNPs in their individual genome. In such a manner, treatment regimens can be

tailored to groups of genetically similar individuals, taking into account traits that may be common among such genetically similar individuals.

[0367] Alternatively, a method termed the "candidate gene approach", can be utilized to identify genes that predict drug response. According to this method, if a gene that encodes a drug's target is known (e.g., a 47324 protein of the present invention), all common variants of that gene can be fairly easily identified in the population and it can be determined if having one version of the gene versus another is associated with a particular drug response.

[0368] Alternatively, a method termed the "gene expression profiling", can be utilized to identify genes that predict drug response. For example, the gene expression of an animal dosed with a drug (e.g., a 47324 molecule or 47324 modulator of the present invention) can give an indication whether gene pathways related to toxicity have been turned on.

[0369] Information generated from more than one of the above pharmacogenomics approaches can be used to determine appropriate dosage and treatment regimens for prophylactic or therapeutic treatment of an individual. This knowledge, when applied to dosing or drug selection, can avoid adverse reactions or therapeutic failure and thus enhance therapeutic or prophylactic efficiency when treating a subject with a 47324 molecule or 47324 modulator, such as a modulator identified by one of the exemplary screening assays described herein.

[0370] The present invention further provides methods for identifying new agents, or combinations, that are based on identifying agents that modulate the activity of one or more of the gene products encoded by one or more of the 47324 genes of the present invention, wherein these products may be associated with resistance of the cells to a therapeutic agent. Specifically, the activity of the proteins encoded by the 47324 genes of the present invention can be used as a basis for identifying agents for overcoming agent resistance. By blocking the activity of one or more of the resistance proteins, target cells, e.g., cancer cells, will become sensitive to treatment with an agent that the unmodified target cells were resistant to.

[0371] Monitoring the influence of agents (e.g., drugs) on the expression or activity of a 47324 protein can be applied in clinical trials. For example, the effectiveness of an agent determined by a screening assay as described herein to increase 47324 gene expression, protein levels, or upregulate 47324 activity, can be monitored in clinical trials of subjects exhibiting decreased 47324 gene expression, protein levels, or downregulated 47324 activity. Alternatively, the effectiveness of an agent determined by a screening assay to decrease 47324 gene expression, protein levels, or downregulate 47324 activity, can be monitored in clinical trials of subjects exhibiting increased 47324 gene expression, protein levels, or upregulated 47324 activity. In such clinical trials, the expression or activity of a 47324 gene, and preferably, other genes that have been implicated in, for example, a 47324-associated disorder can be used as a "read out" or markers of the phenotype of a particular cell.

[0372] Other Embodiments

[0373] In another aspect, the invention features, a method of analyzing a plurality of probes. The method is useful, e.g.,

for analyzing gene expression. The method includes: providing a two dimensional array having a plurality of addresses, each address of the plurality being positionally distinguishable from each other address of the plurality having a unique capture probe, e.g., wherein the capture probes are from a cell or subject which express or mis express 47324 or from a cell or subject in which a 47324 mediated response has been elicited, e.g., by contact of the cell with 47324 nucleic acid or protein, or administration to the cell or subject 47324 nucleic acid or protein; contacting the array with one or more inquiry probe, wherein an inquiry probe can be a nucleic acid, polypeptide, or antibody (which is preferably other than 47324 nucleic acid, polypeptide, or antibody); providing a two dimensional array having a plurality of addresses, each address of the plurality being positionally distinguishable from each other address of the plurality, and each address of the plurality having a unique capture probe, e.g., wherein the capture probes are from a cell or subject which does not express 47324 (or does not express as highly as in the case of the 47324 positive plurality of capture probes) or from a cell or subject which in which a 47324 mediated response has not been elicited (or has been elicited to a lesser extent than in the first sample); contacting the array with one or more inquiry probes (which is preferably other than a 47324 nucleic acid, polypeptide, or antibody), and thereby evaluating the plurality of capture probes. Binding, e.g., in the case of a nucleic acid, hybridization with a capture probe at an address of the plurality, is detected, e.g., by signal generated from a label attached to the nucleic acid, polypeptide, or antibody.

[0374] The capture probes can be a set of nucleic acids from a selected sample, e.g., a sample of nucleic acids derived from a control or non-stimulated tissue or cell.

[0375] The method can include contacting the 47324 nucleic acid, polypeptide, or antibody with a first array having a plurality of capture probes and a second array having a different plurality of capture probes. The results of each hybridization can be compared, e.g., to analyze differences in expression between a first and second sample. The first plurality of capture probes can be from a control sample, e.g., a wild type, normal, or non-diseased, non-stimulated, sample, e.g., a biological fluid, tissue, or cell sample. The second plurality of capture probes can be from an experimental sample, e.g., a mutant type, at risk, disease-state or disorder-state, or stimulated, sample, e.g., a biological fluid, tissue, or cell sample.

[0376] The plurality of capture probes can be a plurality of nucleic acid probes each of which specifically hybridizes, with an allele of 47324. Such methods can be used to diagnose a subject, e.g., to evaluate risk for a disease or disorder, to evaluate suitability of a selected treatment for a subject, to evaluate whether a subject has a disease or disorder. 47324 is associated with Ras family activity, thus it is useful for disorders associated with abnormal Ras family activity.

[0377] The method can be used to detect SNPs, as described above.

[0378] In another aspect, the invention features, a method of analyzing 47324, e.g., analyzing structure, function, or relatedness to other nucleic acid or amino acid sequences. The method includes: providing a 47324 nucleic acid or amino acid sequence; comparing the 47324 sequence with

one or more preferably a plurality of sequences from a collection of sequences, e.g., a nucleic acid or protein sequence database; to thereby analyze 47324.

[0379] The method can include evaluating the sequence identity between a 47324 sequence and a database sequence. The method can be performed by accessing the database at a second site, e.g., over the internet. Preferred databases include GenBank™ and SwissProt.

[0380] In another aspect, the invention features, a set of oligonucleotides, useful, e.g., for identifying SNP's, or identifying specific alleles of 47324. The set includes a plurality of oligonucleotides, each of which has a different nucleotide at an interrogation position, e.g., an SNP or the site of a mutation. In a preferred embodiment, the oligonucleotides of the plurality identical in sequence with one another (except for differences in length). The oligonucleotides can be provided with differential labels, such that an oligonucleotide which hybridizes to one allele provides a signal that is distinguishable from an oligonucleotides which hybridizes to a second allele.

[0381] The sequences of 47324 molecules are provided in a variety of mediums to facilitate use thereof. A sequence can be provided as a manufacture, other than an isolated nucleic acid or amino acid molecule, which contains a 47324 molecule. Such a manufacture can provide a nucleotide or amino acid sequence, e.g., an open reading frame, in a form which allows examination of the manufacture using means not directly applicable to examining the nucleotide or amino acid sequences, or a subset thereof, as they exist in nature or in purified form.

[0382] A 47324 nucleotide or amino acid sequence can be recorded on computer readable media. As used herein, "computer readable media" refers to any medium that can be read and accessed directly by a computer. Such media include, but are not limited to: magnetic storage media, such as floppy discs, hard disc storage medium, and magnetic tape; optical storage media such as compact disc and CD-ROM; electrical storage media such as RAM, ROM, EPROM, EEPROM, and the like; and general hard disks and hybrids of these categories such as magnetic/optical storage media. The medium is adapted or configured for having thereon 47324 sequence information of the present invention.

[0383] As used herein, the term "electronic apparatus" is intended to include any suitable computing or processing apparatus of other device configured or adapted for storing data or information. Examples of electronic apparatus suitable for use with the present invention include stand-alone computing apparatus; networks, including a local area network (LAN), a wide area network (WAN) Internet, Intranet, and Extranet; electronic appliances such as personal digital assistants (PDAs), cellular phones, pagers, and the like; and local and distributed processing systems.

[0384] As used herein, "recorded" refers to a process for storing or encoding information on the electronic apparatus readable medium. Those skilled in the art can readily adopt any of the presently known methods for recording information on known media to generate manufactures comprising the 47324 sequence information.

[0385] A variety of data storage structures are available to a skilled artisan for creating a computer readable medium

having recorded thereon a 47324 nucleotide or amino acid sequence of the present invention. The choice of the data storage structure will generally be based on the means chosen to access the stored information. In addition, a variety of data processor programs and formats can be used to store the nucleotide sequence information of the present invention on computer readable medium. The sequence information can be represented in a word processing text file, formatted in commercially-available software such as WordPerfect and Microsoft Word, or represented in the form of an ASCII file, stored in a database application, such as DB2, Sybase, Oracle, or the like. The skilled artisan can readily adapt any number of data processor structuring formats (e.g. text file or database) in order to obtain computer readable medium having recorded thereon the nucleotide sequence information of the present invention.

[0386] By providing the 47324 nucleotide or amino acid sequences of the invention in computer readable form, the skilled artisan can routinely access the sequence information for a variety of purposes. For example, one skilled in the art can use the nucleotide or amino acid sequences of the invention in computer readable form to compare a target sequence or target structural motif with the sequence information stored within the data storage means. A search is used to identify fragments or regions of the sequences of the invention which match a particular target sequence or target motif.

[0387] The present invention therefore provides a medium for holding instructions for performing a method for determining whether a subject has a Ras family-associated or another 47324-associated disease or disorder or a pre-disposition to a Ras family-associated or another 47324-associated disease or disorder, wherein the method comprises the steps of determining 47324 sequence information associated with the subject and based on the 47324 sequence information, determining whether the subject has a Ras family-associated or another 47324-associated disease or disorder and/or recommending a particular treatment for the disease, disorder, or pre-disease condition.

[0388] The present invention further provides in an electronic system and/or in a network, a method for determining whether a subject has a Ras family-associated or another 47324-associated disease or disorder or a pre-disposition to a disease associated with 47324, wherein the method comprises the steps of determining 47324 sequence information associated with the subject, and based on the 47324 sequence information, determining whether the subject has a Ras family-associated or another 47324-associated disease or disorder or a pre-disposition to a Ras family-associated or another 47324-associated disease or disorder, and/or recommending a particular treatment for the disease, disorder, or pre-disease condition. The method may further comprise the step of receiving phenotypic information associated with the subject and/or acquiring from a network phenotypic information associated with the subject.

[0389] The present invention also provides in a network, a method for determining whether a subject has a Ras family-associated or another 47324-associated disease or disorder or a pre-disposition to a Ras family-associated or another 47324-associated disease or disorder, said method comprising the steps of receiving 47324 sequence information from the subject and/or information related thereto,

receiving phenotypic information associated with the subject, acquiring information from the network corresponding to 47324 and/or corresponding to a Ras family-associated or another 47324-associated disease or disorder, and based on one or more of the phenotypic information, the 47324 information (e.g., sequence information and/or information related thereto), and the acquired information, determining whether the subject has a Ras family-associated or another 47324-associated disease or disorder or a pre-disposition to a Ras family-associated or another 47324-associated disease or disorder. The method may further comprise the step of recommending a particular treatment for the disease, disorder, or pre-disease condition.

[0390] The present invention also provides a business method for determining whether a subject has a Ras family-associated or another 47324-associated disease or disorder or a pre-disposition to a Ras family-associated or another 47324-associated disease or disorder, said method comprising the steps of receiving information related to 47324 (e.g., sequence information and/or information related thereto), receiving phenotypic information associated with the subject, acquiring information from the network related to 47324 and/or related to a Ras family-associated or another 47324-associated disease or disorder, and based on one or more of the phenotypic information, the 47324 information, and the acquired information, determining whether the subject has a Ras family-associated or another 47324-associated disease or disorder or a pre-disposition to a Ras family-associated or another 47324-associated disease or disorder. The method may further comprise the step of recommending a particular treatment for the disease, disorder, or pre-disease condition.

[0391] The invention also includes an array comprising a 47324 sequence of the present invention. The array can be used to assay expression of one or more genes in the array. In one embodiment, the array can be used to assay gene expression in a tissue to ascertain tissue specificity of genes in the array. In this manner, up to about 7600 genes can be simultaneously assayed for expression, one of which can be 47324. This allows a profile to be developed showing a battery of genes specifically expressed in one or more tissues.

[0392] In addition to such qualitative information, the invention allows the quantitation of gene expression. Thus, not only tissue specificity, but also the level of expression of a battery of genes in the tissue if ascertainable. Thus, genes can be grouped on the basis of their tissue expression per se and level of expression in that tissue. This is useful, for example, in ascertaining the relationship of gene expression in that tissue. Thus, one tissue can be perturbed and the effect on gene expression in a second tissue can be determined. In this context, the effect of one cell type on another cell type in response to a biological stimulus can be determined. In this context, the effect of one cell type on another cell type in response to a biological stimulus can be determined. Such a determination is useful, for example, to know the effect of cell-cell interaction at the level of gene expression. If an agent is administered therapeutically to treat one cell type but has an undesirable effect on another cell type, the invention provides an assay to determine the molecular basis of the undesirable effect and thus provides the opportunity to co-administer a counteracting agent or otherwise treat the undesired effect. Similarly, even within a single cell type,

undesirable biological effects can be determined at the molecular level. Thus, the effects of an agent on expression of other than the target gene can be ascertained and counteracted.

[0393] In another embodiment, the array can be used to monitor the time course of expression of one or more genes in the array. This can occur in various biological contexts, as disclosed herein, for example development of a Ras family-associated or another 47324-associated disease or disorder, progression of Ras family-associated or another 47324-associated disease or disorder, and processes, such a cellular transformation associated with the Ras family-associated or another 47324-associated disease or disorder.

[0394] The array is also useful for ascertaining the effect of the expression of a gene on the expression of other genes in the same cell or in different cells (e.g., ascertaining the effect of 47324 expression on the expression of other genes). This provides, for example, for a selection of alternate molecular targets for therapeutic intervention if the ultimate or downstream target cannot be regulated.

[0395] The array is also useful for ascertaining differential expression patterns of one or more genes in normal and abnormal cells. This provides a battery of genes (e.g., including 47324) that could serve as a molecular target for diagnosis or therapeutic intervention.

[0396] As used herein, a "target sequence" can be any DNA or amino acid sequence of six or more nucleotides or two or more amino acids. A skilled artisan can readily recognize that the longer a target sequence is, the less likely a target sequence will be present as a random occurrence in the database. Typical sequence lengths of a target sequence are from about 10 to 100 amino acids or from about 30 to 300 nucleotide residues. However, it is well recognized that commercially important fragments, such as sequence fragments involved in gene expression and protein processing, may be of shorter length.

[0397] Computer software is publicly available which allows a skilled artisan to access sequence information provided in a computer readable medium for analysis and comparison to other sequences. A variety of known algorithms are disclosed publicly and a variety of commercially available software for conducting search means are and can be used in the computer-based systems of the present invention. Examples of such software include, but are not limited to, MacPattern (EMBL), BLASTN and BLASTX (NCBI).

[0398] Thus, the invention features a method of making a computer readable record of a sequence of a 47324 sequence which includes recording the sequence on a computer readable matrix. In a preferred embodiment the record includes one or more of the following: identification of an ORF; identification of a domain, region, or site; identification of the start of transcription; identification of the transcription terminator; the full length amino acid sequence of the protein, or a mature form thereof; the 5' end of the translated region.

[0399] In another aspect, the invention features a method of analyzing a sequence. The method includes: providing a 47324 sequence, or record, in computer readable form; comparing a second sequence to the 47324 sequence; thereby analyzing a sequence. Comparison can include

comparing to sequences for sequence identity or determining if one sequence is included within the other, e.g., determining if the 47324 sequence includes a sequence being compared. In a preferred embodiment the 47324 or second sequence is stored on a first computer, e.g., at a first site and the comparison is performed, read, or recorded on a second computer, e.g., at a second site. E.g., the 47324 or second sequence can be stored in a public or proprietary database in one computer, and the results of the comparison performed, read, or recorded on a second computer. In a preferred embodiment the record includes one or more of the following: identification of an ORF; identification of a domain, region, or site; identification of the start of transcription; identification of the transcription terminator; the full length amino acid sequence of the protein, or a mature form thereof; the 5' end of the translated region. This invention is further illustrated by the following examples which should not be construed as limiting. The contents of all references, patents and published patent applications cited throughout this application are incorporated herein by reference.

EXAMPLES

Example 1

Identification and Characterization of Human 47324 cDNAs

[0400] The human 47324 sequence (FIGS. 1A-B; SEQ ID NO:1), which is approximately 1694 nucleotides long including untranslated regions, contains a predicted methionine-initiated coding sequence of about 711 nucleotides (nucleotides 19-729 of SEQ ID NO:1; SEQ ID NO:3). The coding sequence encodes a 236 amino acid protein (SEQ ID NO:2).

Example 2

Tissue Distribution of 47324 mRNA

[0401] Northern blot hybridizations with various RNA samples can be performed under standard conditions and washed under stringent conditions, i.e., 0.2×SSC at 65° C. A DNA probe corresponding to all or a portion of the 47324 cDNA (SEQ ID NO:1) can be used. The DNA was radioactively labeled with ³²P-dCTP using the Prime-It Kit (Stratagene, La Jolla, Calif.) according to the instructions of the supplier. Filters containing mRNA from mouse hematopoietic and endocrine tissues, and cancer cell lines (Clontech, Palo Alto, Calif.) can be probed in ExpressHyb hybridization solution (Clontech) and washed at high stringency according to manufacturer's recommendations.

Example 3

Gene Expression Analysis

[0402] Total RNA was prepared from various human tissues by a single step extraction method using RNA STAT-60 according to the manufacturer's instructions (TelTest, Inc). Each RNA preparation was treated with DNase I (Ambion) at 37° C. for 1 hour. DNase I treatment was determined to be complete if the sample required at least 38 PCR amplification cycles to reach a threshold level of fluorescence using β-2 microglobulin as an internal ampli-

con reference. The integrity of the RNA samples following DNase I treatment was confirmed by agarose gel electrophoresis and ethidium bromide staining. After phenol extraction cDNA was prepared from the sample using the SUPERSRIPT™ Choice System following the manufacturer's instructions (GibcoBRL). A negative control of RNA without reverse transcriptase was mock reverse transcribed for each RNA sample.

[0403] Human 47324 expression was measured by TaqMan® quantitative PCR (Perkin Elmer Applied Biosystems) in cDNA prepared from a variety of normal and diseased (e.g., cancerous) human tissues or cell lines.

[0404] Probes were designed by PrimerExpress software (PE Biosystems) based on the sequence of the human 47324 gene. Each human 47324 gene probe was labeled using FAM (6-carboxyfluorescein), and the β -2-microglobulin reference probe was labeled with a different fluorescent dye, VIC. The differential labeling of the target gene and internal reference gene thus enabled measurement in same well. Forward and reverse primers and the probes for both β -2-microglobulin and target gene were added to the TaqMan® Universal PCR Master Mix (PE Applied Biosystems). Although the final concentration of primer and probe could vary, each was internally consistent within a given experiment. A typical experiment contained 200 nM of forward and reverse primers plus 100 nM probe for β -2 microglobulin and 600 nM forward and reverse primers plus 200 nM probe for the target gene. TaqMan matrix experiments were carried out on an ABI PRISM 7700 Sequence Detection System (PE Applied Biosystems). The thermal cycler conditions were as follows: hold for 2 min at 50° C. and 10 min at 95° C., followed by two-step PCR for 40 cycles of 95° C. for 15 sec followed by 60° C. for 1 min.

[0405] The following method was used to quantitatively calculate human 47324 gene expression in the various tissues relative to β -2 microglobulin expression in the same tissue. The threshold cycle (Ct) value is defined as the cycle at which a statistically significant increase in fluorescence is detected. A lower Ct value is indicative of a higher mRNA concentration. The Ct value of the human 47324 gene is normalized by subtracting the Ct value of the β -2 microglobulin gene to obtain a Δ Ct value using the following formula: Δ Ct=Ct_{human 47324} - Ct _{β -2 microglobulin}. Expression is then calibrated against a cDNA sample showing a comparatively low level of expression of the human 47324 gene. The Δ Ct value for the calibrator sample is then subtracted from Δ Ct for each tissue sample according to the following formula: $\Delta\Delta$ Ct= Δ Ct_{sample} - Δ Ct_{calibrator}. Relative expression is then calculated using the arithmetic formula given by 2^{- $\Delta\Delta$ Ct}. Expression of the target human 47324 gene in each of the tissues tested is then graphically represented as discussed in more detail below.}}

[0406] TaqMan real-time quantitative RT-PCR is used to detect the presence of RNA transcript corresponding to human 47324 relative to a no template control in a panel of human tissues or cells. It is found that the highest expression of 47324 orthologs are expressed in normal brain cortex tissue, erythroid cell lines, and HUVEC (human umbilical vein endothelial cells), as shown in table 1. Relatively high expression is also seen in skeletal muscle, coronary smooth muscle cells, lung tumor tissue, normal artery and megakaryocytes. It is also of note that there is increased expres-

sion of 47324 in prostate tumor, colon tumor, lung tumor and fibrotic liver compared to normal prostate, colon, lung and liver tissue. In addition, there is decreased expression of 47324 in diseased aorta tissue, breast tumor and ovary tumor tissue compared to normal artery, breast, and ovary tissue.

TABLE 1

Tissue Type	Mean	β 2 Mean	$\Delta\Delta$ Ct	Expression
Artery normal	29.4	22.68	6.71	9.5188
Aorta diseased	32.02	22.97	9.06	1.8736
Vein normal	30.41	20.8	9.62	1.2753
Coronary SMC	27.34	21.14	6.2	13.6024
HUVEC	27.23	21.61	5.62	20.3335
Hemangioma	29.31	20.23	9.09	1.8414
Heart normal	28.73	21.21	7.53	5.4294
Heart CHF	28.72	21.32	7.39	5.9413
Kidney	27.75	20.5	7.25	6.5923
Skeletal Muscle	28.36	22.4	5.96	16.0643
Adipose normal	30.3	21.11	9.19	1.7181
Pancreas	30.16	22.54	7.62	5.0834
primary osteoblasts	30.86	21.07	9.79	1.1296
Osteoclasts (diff)	31.83	18	13.82	0.0689
Skin normal	31.8	22.54	9.26	1.631
Spinal cord normal	30.95	21.49	9.46	1.4199
Brain Cortex normal	27.57	22.41	5.17	27.7763
Brain Hypothalamus normal	30.13	22.73	7.4	5.9208
Nerve	31.59	22.41	9.17	1.736
DRG (Dorsal Root Ganglion)	29.61	21.89	7.73	4.7102
Breast normal	30.57	21.4	9.17	1.736
Breast tumor	31.25	21.16	10.08	0.9239
Ovary normal	29	20.25	8.76	2.3146
Ovary Tumor	28.66	18.97	9.7	1.2065
Prostate Normal	31.11	20.04	11.07	0.4668
Prostate Tumor	28.29	20.65	7.64	5.0134
Salivary glands	31.34	20.11	11.24	0.4134
Colon normal	31.06	19.08	11.98	0.2484
Colon Tumor	30.05	21.93	8.13	3.582
Lung normal	30.31	18.69	11.62	0.3188
Lung tumor	27.45	20.9	6.55	10.6722
Lung COPD	30.84	19.09	11.76	0.2893
Colon IBD	30.47	18.11	12.36	0.1909
Liver normal	30.79	20.39	10.4	0.7375
Liver fibrosis	29.61	21.2	8.41	2.9399
Spleen normal	30.29	18.92	11.37	0.3778
Tonsil normal	29.34	18.84	10.49	0.6929
Lymph node normal	30.79	20.54	10.25	0.8212
Small intestine normal	31.56	20.52	11.04	0.4766
Macrophages	29.09	17.41	11.68	0.3048
Synovium	32.44	20.32	12.12	0.2254
BM-MNC	31.66	19.34	12.33	0.1942
Activated PBMC	29.55	18.21	11.34	0.3858
Neutrophils	27.74	18.34	9.4	1.4802
Megakaryocytes	26.15	19.32	6.83	8.7895
Erythroid	26.59	21.07	5.51	21.9445
positive control	26.03	20.79	5.24	26.4608

Example 4

Recombinant Expression of 47324 in Bacterial Cells

[0407] In this example, 47324 is expressed as a recombinant glutathione-S-transferase (GST) fusion polypeptide in *E. coli* and the fusion polypeptide is isolated and characterized. Specifically, 47324 is fused to GST and this fusion polypeptide is expressed in *E. coli*, e.g., strain PEB199. Expression of the GST-47324 fusion protein in PEB199 is induced with IPTG. The recombinant fusion polypeptide is purified from crude bacterial lysates of the induced PEB199 strain by affinity chromatography on glutathione beads. Using polyacrylamide gel electrophoretic analysis of the polypeptide purified from the bacterial lysates, the molecular weight of the resultant fusion polypeptide is determined.

Example 5

Expression of Recombinant 47324 Protein in COS Cells

[0408] To express the 47324 gene in COS cells, the pcDNA/Amp vector by Invitrogen Corporation (San Diego, Calif.) is used. This vector contains an SV40 origin of replication, an ampicillin resistance gene, an *E. coli* replication origin, a CMV promoter followed by a polylinker region, and an SV40 intron and polyadenylation site. A DNA fragment encoding the entire 47324 protein and an HA tag (Wilson et al. (1984) *Cell* 37:767) or a FLAG tag fused in-frame to its 3' end of the fragment is cloned into the polylinker region of the vector, thereby placing the expression of the recombinant protein under the control of the CMV promoter.

[0409] To construct the plasmid, the 47324 DNA sequence is amplified by PCR using two primers. The 5' primer contains the restriction site of interest followed by approximately twenty nucleotides of the 47324 coding sequence starting from the initiation codon; the 3' end sequence contains complementary sequences to the other restriction site of interest, a translation stop codon, the HA tag or FLAG tag and the last 20 nucleotides of the 47324 coding sequence. The PCR amplified fragment and the pcDNA/Amp vector are digested with the appropriate restriction enzymes and the vector is dephosphorylated using the CIAP enzyme (New England Biolabs, Beverly, Mass.). Preferably the two restriction sites chosen are different so that the 47324 gene is inserted in the correct orientation. The ligation mixture is transformed into *E. coli* cells (strains HB101, DH5 α , SURE, available from Stratagene Cloning Systems, La Jolla, Calif., can be used), the transformed culture is plated on ampicillin media plates, and resistant colonies are selected. Plasmid DNA is isolated from transformants and examined by restriction analysis for the presence of the correct fragment.

[0410] COS cells are subsequently transfected with the 47324-pcDNA/Amp plasmid DNA using the calcium phosphate or calcium chloride co-precipitation methods, DEAE-dextran-mediated transfection, lipofection, or electroporation. Other suitable methods for transfecting host cells can be found in Sambrook, J., Fritsh, E. F., and Maniatis, T. *Molecular Cloning: A Laboratory Manual*. 2nd, ed., Cold Spring Harbor Laboratory, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 1989. The expression of the 47324 polypeptide is detected by radiolabelling (³⁵S-methionine or ³⁵S-cysteine available from NEN, Boston, Mass., can be used) and immunoprecipitation (Harlow, E. and Lane, D. *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 1988) using an HA specific monoclonal antibody. Briefly, the cells are labeled for 8 hours with ³⁵S-methionine (or ³⁵S-cysteine). The culture media are then collected and the cells are lysed using detergents (RIPA buffer, 150 mM NaCl, 1% NP-40, 0.1% SDS, 0.5% DOC, 50 mM Tris, pH 7.5). Both the cell lysate and the culture media are precipitated with an HA specific monoclonal antibody. Precipitated polypeptides are then analyzed by SDS-PAGE.

[0411] Alternatively, DNA containing the 47324 coding sequence is cloned directly into the polylinker of the pcDNA/Amp vector using the appropriate restriction sites.

The resulting plasmid is transfected into COS cells in the manner described above, and the expression of the 47324 polypeptide is detected by radiolabelling and immunoprecipitation using a 47324 specific monoclonal antibody.

Equivalents

[0412] Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents to the specific embodiments of the invention described herein. Such equivalents are intended to be encompassed by the following claims.

What is claimed is:

1. An isolated 47324 nucleic acid molecule selected from the group consisting of:

- a) a nucleic acid molecule comprising a nucleotide sequence which is at least 60% identical to the nucleotide sequence of SEQ ID NO:1, SEQ ID NO:3, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____;
- b) a nucleic acid molecule comprising a fragment of at least 15 nucleotides of the nucleotide sequence of SEQ ID NO:1, SEQ ID NO:3, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____;
- c) a nucleic acid molecule which encodes a polypeptide comprising the amino acid sequence of SEQ ID NO:2, or the amino acid sequence encoded by the cDNA insert of the plasmid deposited with the ATCC as Accession Number _____;
- d) a nucleic acid molecule which encodes a fragment of a polypeptide comprising the amino acid sequence of SEQ ID NO:2, or the amino acid sequence encoded by the cDNA insert of the plasmid deposited with the ATCC as Accession Number _____, wherein the fragment comprises at least 15 contiguous amino acids of SEQ ID NO:2, or the amino acid sequence encoded by the cDNA insert of the plasmid deposited with the ATCC as Accession Number _____;
- e) a nucleic acid molecule which encodes a naturally occurring allelic variant of a polypeptide comprising the amino acid sequence of SEQ ID NO:2, or the amino acid sequence encoded by the cDNA insert of the plasmid deposited with the ATCC as Accession Number _____, wherein the nucleic acid molecule hybridizes to a nucleic acid molecule comprising SEQ ID NO:1, SEQ ID NO:3, or a complement thereof, under stringent conditions;
- f) a nucleic acid molecule comprising the nucleotide sequence of SEQ ID NO:1, SEQ ID NO:3, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____; and
- g) a nucleic acid molecule which encodes a polypeptide comprising the amino acid sequence of SEQ ID NO:2, or the amino acid sequence encoded by the cDNA insert of the plasmid deposited with the ATCC as Accession Number _____.

2. The isolated nucleic acid molecule of claim 1, which is the nucleotide sequence SEQ ID NO:1.

3. A host cell which contains the nucleic acid molecule of claim 1.

4. An isolated 47324 polypeptide selected from the group consisting of:

- a) a polypeptide which is encoded by a nucleic acid molecule comprising a nucleotide sequence which is at least 60% identical to a nucleic acid comprising the nucleotide sequence of SEQ ID NO:1, SEQ ID NO:3, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____, or a complement thereof;
- b) a naturally occurring allelic variant of a polypeptide comprising the amino acid sequence of SEQ ID NO:2, or the amino acid sequence encoded by the cDNA insert of the plasmid deposited with the ATCC as Accession Number _____, wherein the polypeptide is encoded by a nucleic acid molecule which hybridizes to a nucleic acid molecule comprising SEQ ID NO:1, SEQ ID NO:3, or a complement thereof under stringent conditions;
- c) a fragment of a polypeptide comprising the amino acid sequence of SEQ ID NO:2, or the amino acid sequence encoded by the cDNA insert of the plasmid deposited with the ATCC as Accession Number _____, wherein the fragment comprises at least 15 contiguous amino acids of SEQ ID NO:2; and
- d) the amino acid sequence of SEQ ID NO:2.

5. An antibody which selectively binds to a polypeptide of claim 4.

6. A method for producing a polypeptide selected from the group consisting of:

- a) a polypeptide comprising the amino acid sequence of SEQ ID NO:2, or the amino acid sequence encoded by the cDNA insert of the plasmid deposited with the ATCC as Accession Number _____;
- b) a polypeptide comprising a fragment of the amino acid sequence of SEQ ID NO:2, or the amino acid sequence encoded by the cDNA insert of the plasmid deposited with the ATCC as Accession Number _____, wherein the fragment comprises at least 15 contiguous amino acids of SEQ ID NO:2, or the amino acid sequence encoded by the cDNA insert of the plasmid deposited with the ATCC as Accession Number _____;
- c) a naturally occurring allelic variant of a polypeptide comprising the amino acid sequence of SEQ ID NO:2, or the amino acid sequence encoded by the cDNA insert of the plasmid deposited with the ATCC as Accession Number _____, wherein the polypeptide is encoded by a nucleic acid molecule which hybridizes to a nucleic acid molecule comprising SEQ ID NO:1 or SEQ ID NO:3; and
- d) the amino acid sequence of SEQ ID NO:2;

comprising culturing the host cell of claim 3 under conditions in which the nucleic acid molecule is expressed.

7. A method for detecting the presence of a nucleic acid molecule of claim 1 or a polypeptide encoded by the nucleic acid molecule in a sample, comprising:

a) contacting the sample with a compound which selectively hybridizes to the nucleic acid molecule of claim 1 or binds to the polypeptide encoded by the nucleic acid molecule; and

b) determining whether the compound hybridizes to the nucleic acid or binds to the polypeptide in the sample.

8. A kit comprising a compound which selectively hybridizes to a nucleic acid molecule of claim 1 or binds to a polypeptide encoded by the nucleic acid molecule and instructions for use.

9. A method for identifying a compound which binds to a polypeptide or modulates the activity of the polypeptide of claim 4 comprising the steps of:

a) contacting a polypeptide, or a cell expressing a polypeptide of claim 4 with a test compound; and

b) determining whether the polypeptide binds to the test compound or determining the effect of the test compound on the activity of the polypeptide.

10. A method for modulating the activity of a polypeptide of claim 4 comprising contacting the polypeptide or a cell expressing the polypeptide with a compound which binds to the polypeptide in a sufficient concentration to modulate the activity of the polypeptide.

11. A method of identifying a nucleic acid molecule associated with a disorder comprising:

a) contacting a sample from a subject with or at risk of developing a disorder comprising nucleic acid molecules with a hybridization probe comprising at least 25 contiguous nucleotides of SEQ ID NO:1 defined in claim 2; and

b) detecting the presence of a nucleic acid molecule in the sample that hybridizes to the probe, thereby identifying a nucleic acid molecule associated with a disorder.

12. A method of identifying a nucleic acid associated with a disorder comprising:

a) contacting a sample from a subject having a disorder or at risk of developing a disorder comprising nucleic acid molecules with a first and a second amplification primer, the first primer comprising at least 25 contiguous nucleotides of SEQ ID NO:1 defined in claim 2 and the second primer comprising at least 25 contiguous nucleotides from the complement of SEQ ID NO:1;

b) incubating the sample under conditions that allow nucleic acid amplification; and

c) detecting the presence of a nucleic acid molecule in the sample that is amplified, thereby identifying the nucleic acid molecule associated with a disorder.

13. A method of identifying a polypeptide associated with a disorder comprising:

a) contacting a sample comprising polypeptides with a 47324 binding partner of the 47324 polypeptide defined in claim 4; and

b) detecting the presence of a polypeptide in the sample that binds to the 47324 binding partner, thereby identifying the polypeptide associated with a disorder.

14. A method of identifying a subject having a disorder or at risk for developing a disorder comprising:

a) contacting a sample obtained from the subject comprising nucleic acid molecules with a hybridization

probe comprising at least 25 contiguous nucleotides of SEQ ID NO:1 defined in claim 2; and

- b) detecting the presence of a nucleic acid molecule in the sample that hybridizes to the probe, thereby identifying a subject having a disorder or at risk for developing a disorder.

15. A method of identifying a subject having a disorder or at risk for developing a disorder comprising:

- a) contacting a sample obtained from the subject comprising nucleic acid molecules with a first and a second amplification primer, the first primer comprising at least 25 contiguous nucleotides of SEQ ID NO:1 defined in claim 2 and the second primer comprising at least 25 contiguous nucleotides from the complement of SEQ ID NO:1;
- b) incubating the sample under conditions that allow nucleic acid amplification; and
- c) detecting the presence of a nucleic acid molecule in the sample that is amplified, thereby identifying a subject having a disorder or at risk for developing a disorder.

16. A method of identifying a subject having a disorder or at risk for developing a disorder comprising:

- a) contacting a sample obtained from the subject comprising polypeptides with a 47324 binding partner of the 47324 polypeptide defined in claim 4; and
- b) detecting the presence of a polypeptide in the sample that binds to the 47324 binding partner, thereby identifying a subject having a disorder or at risk for developing a disorder.

17. A method for identifying a compound capable of treating a disorder characterized by aberrant 47324 nucleic acid expression or 47324 polypeptide activity comprising assaying the ability of the compound to modulate 47324 nucleic acid expression or 47324 polypeptide activity, thereby identifying a compound capable of treating a disorder characterized by aberrant 47324 nucleic acid expression or 47324 polypeptide activity.

18. A method for treating a subject having a disorder or at risk of developing a disorder comprising administering to the subject a 47324 modulator of the nucleic acid molecule defined in claim 1 or the polypeptide encoded by the nucleic acid molecule or contacting a cell with a 47324 modulator.

19. The method of claim 18, wherein the 47324 modulator is

- a) a small molecule;
- b) peptide;
- c) phosphopeptide;
- d) anti-47324 antibody;
- e) a 47324 polypeptide comprising the amino acid sequence of SEQ ID NO:2, or a fragment thereof;
- f) a 47324 polypeptide comprising an amino acid sequence which is at least 90 percent identical to the amino acid sequence of SEQ ID NO:2, wherein the percent identity is calculated using the ALIGN program for comparing amino acid sequences, a PAM120 weight residue table, a gap length penalty of 12, and a gap penalty of 4; or

- g) an isolated naturally occurring allelic variant of a polypeptide consisting of the amino acid sequence of SEQ ID NO:2, wherein the polypeptide is encoded by a nucleic acid molecule which hybridizes to a complement of a nucleic acid molecule consisting of SEQ ID NO:1 at 6×SSC at 45° C., followed by one or more washes in 0.2×SSC, 0.1% SDS at 65° C.

20. The method of claim 18, wherein the 47324 modulator is

- a) an antisense 47324 nucleic acid molecule;
- b) is a ribozyme;
- c) the nucleotide sequence of SEQ ID NO:1, or a fragment thereof;
- d) a nucleic acid molecule encoding a polypeptide comprising an amino acid sequence which is at least 90 percent identical to the amino acid sequence of SEQ ID NO:2, wherein the percent identity is calculated using the ALIGN program for comparing amino acid sequences, a PAM120 weight residue table, a gap length penalty of 12, and a gap penalty of 4;
- e) a nucleic acid molecule encoding a naturally occurring allelic variant of a polypeptide comprising the amino acid sequence of SEQ ID NO:2, wherein the nucleic acid molecule which hybridizes to a complement of a nucleic acid molecule consisting of SEQ ID NO:1 at 6×SSC at 45° C., followed by one or more washes in 0.2×SSC, 0.1% SDS at 65° C.; or
- f) a gene therapy vector.

21. A method for evaluating the efficacy of a treatment of a disorder, in a subject, comprising:

- treating a subject with a protocol under evaluation;
- assessing the expression level of a 47324 nucleic acid molecule defined in claim 1 or 47324 polypeptide encoded by the 47324 nucleic acid molecule,

wherein a change in the expression level of 47324 nucleic acid or 47324 polypeptide after the treatment, relative to the level before the treatment, is indicative of the efficacy of the treatment of a disorder.

22. A method of diagnosing a disorder in a subject, comprising:

- evaluating the expression or activity of a 47324 nucleic acid molecule defined in claim 1 or a 47324 polypeptide encoded by the 47324 nucleic acid molecule, such that a difference in the level of 47324 nucleic acid or 47324 polypeptide relative to a normal subject or a cohort of normal subjects is indicative of a disorder.

23. The method defined in claim 18, wherein the disorder is cancer or aberrant cellular proliferation and/or differentiation, cardiovascular disorders, liver disorders, pain and metabolic disorders, brain disorders, blood vessel disorders, hematopoietic disorders, and platelet disorders.

24. The method defined in claim 23, wherein the cancer or aberrant cellular proliferation and/or differentiation is breast, ovarian, prostate, colon, or lung cancer.

专利名称(译)	47324, 一种新型人类g蛋白, 并用于此		
公开(公告)号	US20020127568A1	公开(公告)日	2002-09-12
申请号	US09/945173	申请日	2001-08-31
[标]申请(专利权)人(译)	MEYERS RACHEL		
申请(专利权)人(译)	MEYERS RACHEL		
当前申请(专利权)人(译)	MEYERS RACHEL		
[标]发明人	MEYERS RACHEL		
发明人	MEYERS, RACHEL		
IPC分类号	C07K14/47 A61K38/00 C12N15/12 C12Q1/68 G01N33/53 C07H21/04 C12P21/02 C12N5/06 C07K14/705 C07K16/28		
CPC分类号	A61K38/00 C07K14/4722		
优先权	60/229293 2000-09-01 US		
外部链接	Espacenet USPTO		

摘要(译)

本发明提供了分离的核酸分子, 命名为47324核酸分子, 其编码新的G蛋白家族成员, 优选Ras家族成员。本发明还提供了反义核酸分子, 含有47324个核酸分子的重组表达载体, 已导入表达载体的宿主细胞, 以及其中已引入或破坏47324基因的非人转基因动物。本发明还进一步提供了分离的47324蛋白, 融合蛋白, 抗原肽和抗47324抗体。还提供了利用本发明组合物的诊断方法。

```

AGGAAAGCGGCTGGCAAG      H A S L D R V K V L V L G D S      SEQ ID NO:1
ATG GGG TCC CTC GAT CGG GTG AAG GTA CTG GTG GGA GAC TCA      SEQ ID NO:2
      S L W N L L C Q L Q V E G H F
G V G K S S T TCG TTA GTC CAT CTC CTA TGC CAA AAT CAA CTG GGA AAT CCA      35
GGT GTT GGG AAA TCT TCG TTA GTC CAT CTC CTA TGC CAA AAT CAA CTG GGA AAT CCA      105
      S M T V G C S V D V R V H D Y K E G T F E
TCA TGG ACT GTG GGC TCC TCA GGG GAT GGG AGA GTT GAT GAT TAC AAA GAA GGA ACC CCA      165
      E E K T Y TAC TAC TCA GAA TTA TGG GAT V G G S V G G S A S A S
GAA GAG AAG ACC TAC TAC TCA GAA TTA TGG GAT V G G S V G G S A S A S      225
      S V K S T R A V E V F S V R G L T F V H
AGC GTG AAA AGC ACA AGA GCA GTA TCC TAC AAG TCC GTA ANT GGT ATT ATT TTC GTA GAC      285
      D L T N K K S S Q N L R R W S L E A L N
GAC TTA ACA AAT AAG AAG TCC TCC GAA AAG TGG GGT CTT TGG TCA TGG GAA GCT CTC AAC      345
      R D L V R C G V L V T N G D Y D Q E Q F
AGG GAT TCG CTG CCA ACT GGA GTC TTG GTG ACA AAT GGG GAT TAT GAT CAA GAA CAG TTT      405
      A D N Q I F L L V T S T R K L D Q T R E T
GCT GAT AAG CAA ATA CCA CTG TTG GTA ATA GGG ACT ABA CTG GAC CGG AAT GAT GAA AGA      465
      K R H E V L T R T A F L A E D F N R E E
AAG GCG CAT GAA GTT TTA ACT AAG ACT GCT TTC CTG GCT GAG GAT TTC AAT CCA GAA GAA      525
      I N E D C E N R V D A A G S S H A V K
ATT AAT TTG GAC TGC ACA AAT CCA CGG TAC TTA GCT GCA GGT CCG TCC AAT GCT GTC AAG      585
      L S R R F F D R V T E K R V F L R E G H Q
CTC AGT AAG TTT TTT GAT AAG GTC ATA GAG AAG AGA TGC TTT TTA AGA GAA GGT AAT GAG      645
      I P G F F D R K R F G A G T L K S L K Y
ATT CCA GGC TTT CCT GAT CGG AAA AGA TTT GGG GCA GGA ACA TTA AAG AGC CTT CAT TAT      705
      D
      GAC TGA
      SEQ ID NO:3
ATTACACTCATCTCTTGGAAAGATGAGCAAGCGAGTGGCAGTCTTTCACAGCTCATCTTGCCTGTGTTCAATTAATACCAAT
CACAGGCTTTTAAACAANAATCATCTTAAAAATGCTACCCCTTACGCTTACCCCTTAAATGGAANAATGAAAGGAAGTACAA
TACGGGAGGTCCAAACTTTGCTCCCTCTCTGTCTCTACCTTCTCTGCTCCCTGTATACAGACTATGTAAGAAAGCCCT
GGTAAATATAGAGCTGTGCAAAAATGATGCAAAATGAGCAATGACAGTGTACTGCAGAGAAATTTACTCTTGCCT
AGAACTGGAGGGTTTTATAGGCTCTGCAATTTCCCACTCATTTGCTGAAAGCTTAAATTAAGTACTCCAAAAGCAT
CTCATTTTACTCTCTCTGAGGGGAGGGCTCTTACAGAGCCCTGATTTGTCACCCCAACAATCTCTGTGAT
TCAAAATGCAAAAATGCTGATTTAAATTTGCTCCAGCATGTCACACACAGGAATGCTTAATAATAGCAACCCCTG
TCCCTCTCTCTCTTTCGAAATGGCTCAGTGTGAGAGGCGGCAATATAGCCAGCTAAATATAAATACAAAT

```

FIG. 1A