



US 20030176647A1

(19) **United States**

(12) **Patent Application Publication** (10) **Pub. No.: US 2003/0176647 A1**  
(43) **Pub. Date:** **Sep. 18, 2003**

---

(54) **ANTIBODIES TO OCIF-BINDING MOLECULES**

(75) Inventors: **Kyoji Yamaguchi**, Saitama (JP); **Hisataka Yasuda**, Tochigi (JP); **Nobuaki Nakagawa**, Tochigi (JP); **Nobuyuki Shima**, Tochigi (JP); **Masahiko Kinosaki**, Tochigi (JP); **Eisuke Tsuda**, Tochigi (JP); **Masaaki Goto**, Tochigi (JP); **Kazuki Yano**, Tochigi (JP); **Akihiro Tomoyasu**, Tochigi (JP); **Fumie Kobayashi**, Tochigi (JP); **Naohiro Washida**, Tochigi (JP); **Ken Takahashi**, Tochigi (JP); **Tomonori Morinaga**, Tochigi (JP); **Kanji Higashio**, Saitama (JP)

Correspondence Address:

**ANDREW S. BRENC**  
**ARNOLD AND PORTER**  
**555 TWELFTH STREET, NW**  
**IP DOCKETING DEPARTMENT, ROOM 1126**  
**WASHINGTON, DC 20004-1206 (US)**

(73) Assignee: **Sankyo Co., Ltd.**, Tokyo (JP)

(21) Appl. No.: **10/167,182**

(22) Filed: **Jun. 11, 2002**

#### Related U.S. Application Data

(62) Division of application No. 09/202,455, filed on Dec. 15, 1998.

#### (30) Foreign Application Priority Data

Apr. 15, 1998	(WO) .....	PCT/JP98/01728
Apr. 15, 1997	(JP) .....	97808
Jun. 9, 1997	(JP) .....	151434
Aug. 12, 1997	(JP) .....	217897
Aug. 21, 1997	(JP) .....	224803
Dec. 2, 1997	(JP) .....	332241

#### Publication Classification

(51) **Int. Cl.<sup>7</sup>** ..... **C07K 14/725**; C07H 21/04;  
C12P 21/02; C12N 5/06  
(52) **U.S. Cl.** ..... **530/350**; 530/388.25; 435/69.1;  
435/320.1; 435/325; 536/23.5

#### (57) ABSTRACT

A novel protein which binds to Osteoclastogenesis Inhibitory Factor (OCIF-binding molecule; OBM), a process for preparing the same, DNA encoding said protein, a protein having an amino acid sequence encoded by this DNA, a method for producing said protein by genetic engineering technique, and a pharmaceutical composition containing said protein.

Screening methods for a substance for controlling expression of said protein using said protein and the DNA, a substance which inhibits or modulates the biological activity of said protein, or a receptor which transmits the action of said protein through binding to said protein, the substance obtained by the screening methods, and a pharmaceutical composition which contains this substance.

An antibody for said protein, a process for preparing the same, a measuring method of said protein using the antibody, and a medicine comprising this antibody.

FIG.1 (A) (B)

kDa	1	2	3	1	2
-----	---	---	---	---	---





FIG.3




FIG.4




FIG.5




FIG.6




FIG. 7

kDa 1 2 3

220 -  
97.4 -  
66 -  
46 -  
30 -  
21.5 -  
14.5 -

FIG. 8




FIG.9




FIG.10




FIG.11



FIG.12




FIG.13




FIG.14




FIG.15




FIG.16




FIG.17




FIG.18




FIG.19




FIG.20




FIG.21




FIG.22




FIG. 23




FIG. 24




FIG. 25




FIG. 26




FIG. 27




FIG. 28




FIG. 29




FIG. 30




FIG. 31




FIG. 32





FIG. 33



**ANTIBODIES TO OCIF-BINDING MOLECULES****FIELD OF TECHNOLOGY**

**[0001]** The present invention relates to a novel protein (OCIF-binding molecule, the protein may be hereinafter called OBM) which binds to osteoclastogenesis inhibitory factor (hereinafter it may be called OCIF) and a method to produce this protein.

**[0002]** The present invention also relates to DNA encoding this protein, proteins containing the amino acid sequence encoded by this DNA, a method for the preparation of this protein utilizing genetic engineering techniques, and pharmaceutical compositions comprising this protein.

**[0003]** The present invention further relates to methods for screening, using this protein and the DNA, substances to control the expression of this protein, substances inhibiting or regulating the biological activity of this protein, or receptors transducing the signal of the protein by interacting with this protein, to substances obtained by the screening, and to pharmaceutical compositions which comprise the resulting substances.

**[0004]** The present invention further relates to antibodies against this protein, methods for preparing the antibodies, and pharmaceutical compositions comprising these antibodies. Background Art: U.S. Pat. No. 5,440,839, 5

**[0005]** Bone metabolism is dependent on the overall activity of osteoblasts which control bone formation and osteoclasts which control bone resorption. Abnormality of bone metabolism is considered to be caused by an imbalance of the bone formation and the bone resorption. Osteoporosis, hypercalcemia, Paget's disease, renal osteodystrophy, chronic rheumatitis, osteoarthritis, and the like are known as diseases accompanying abnormality of bone metabolism. Osteoporosis is a typical disease caused by such abnormality of bone metabolism. This disease is generated when bone resorption by osteoclasts exceeds bone formation by osteoblasts. The disease is characterized by a decrease in both the bone calcified material and the bone matrix. Although the mechanism of this disease is not completely elucidated, the disease causes aches in bones, makes them fragile, and may result in fracturing. This disease is becoming a social problem because it increases the number of bedridden aged persons as the aged population becomes larger. Development of therapeutic agent for this disease is urgently desired. Disease due to a decrease in bone mass is expected to be cured by suppressing bone resorption, accelerating bone formation, or improving the balance between bone resorption and formation. Bone formation is expected to increase by accelerating proliferation, differentiation, or activation of osteoblasts which form bone, or by suppressing proliferation, differentiation, or activation of osteoclasts which resorb bone. In recent years, strong interest has been directed to hormones, low molecular weight substances, or physiologically active proteins exhibiting such activities, and energetic basic research and development is underway on these subjects.

**[0006]** Drugs such as a calcitonin agents, active-form vitamin D<sub>3</sub> agents, hormone agents containing estradiol, ipriflavon, vitamin K<sub>2</sub>, and bisphosphonate compounds have already been known as drugs to treat and shorten the treatment period of diseases related to bone. Clinical tests

are in progress on active-form vitamin D<sub>3</sub> derivatives, estradiol derivatives, and bisphosphonate compounds of the second and the third generation to develop therapeutic agents with excellent efficacy and minimal side effects.

**[0007]** However, therapies using these agents were found not necessarily satisfactory in terms of efficacy and therapeutic results. Development of novel therapeutic agents which are safer and with higher efficacy is urgently desired. Some agents used for the treatment of diseases related to bone metabolism are used only limitedly due to their side effects. Furthermore, treatments using two or more agents in combination are currently the mainstream in the treatment of diseases related to bone metabolism such as osteoporosis. From such a point of view, development of drugs having action mechanisms different from those of conventional drugs, and exhibiting a higher efficacy and minimal side effects is desired.

**[0008]** As mentioned above, the cells controlling bone metabolism are osteoblasts and osteoclasts. These cells are known to have close mutual interactions called "coupling". Specifically, cytokines such as Interleukins 1(IL-1), 3(IL-3), 6(IL-6), and 11(IL-11), granulocytic macrophage-colony stimulating factor (GM-CSF), macrophage-colony stimulating factor (M-CSF), Interferon- $\gamma$  (IFN- $\gamma$ ), tumor necrosis factor  $\alpha$  (TNF- $\alpha$ ), and transforming growth factor- $\beta$  (TGF- $\beta$ ), secreted by osteoblastic stromal cells are known to accelerate or suppress differentiation or maturation of osteoclasts (Raisz: Disorders of Bone and Mineral Metabolism, 287-311, 1992; Suda et al.: Principles of Bone Biology, 87-102, 1996; Suda et al.: Endocrine Reviews, 4, 266-270, 1995, Lacey et al.: Endocrinology, 136, 2369-2376, 1995). It has been reported that osteoblastic stromal cells play an important role in the differentiation and maturation of osteoclasts, as well as in osteoclast functions such as bone resorption by mature osteoclasts, through cell-to-cell contact with immature osteoclast precursors or mature osteoclasts.

**[0009]** A factor called osteoclast differentiation factor (ODF, Suda et al.: Endocrine Rev. 13, 66-80, 1992; Suda et al.: Bone 17, 87S-91S, 1995) is thought to be expressed on the membrane of osteoblastic stromal cells and involved in the formation of osteoclasts through cell-to-cell contact. According to this hypothesis, an ODF receptor is present in the precursor cells of osteoclasts. However, so far neither the ODF nor the receptor has been purified or identified. There are also no reports relating to their characteristics, action mechanism, or structure. Thus, the mechanism involved in differentiation and maturation of osteoclasts has not yet been sufficiently elucidated. Clarification of this mechanism will greatly contribute not only to the basic medicine, but also to the development of novel drugs for the treatment of diseases associated with abnormality of bone metabolism.

**[0010]** The present inventors have conducted extensive studies in view of this situation and discovered an osteoclastogenesis inhibitory factor (OCIF) in a culture broth of human embryonic lung fibroblast, IMR-90 (ATCC Deposition No. CCL186) (WO 96/26217).

**[0011]** The present inventors have been successful in cloning DNA encoding OCIF, production of recombinant OCIF in animal cells, and confirmation of in vivo pharmaceutical effects (improving effect on bone metabolism, etc.) of the recombinant OCIF. OCIF is expected to be used as an agent for the prevention or treatment of diseases related to

abnormality of bone metabolism, with higher efficacy than conventional drugs and less side effects.

#### DISCLOSURE OF THE INVENTION

[0012] The present inventors have searched for a protein which binds to osteoclastogenesis inhibitory factor (OCIF) and discovered that an OCIF-binding protein is specifically expressed on the osteoblastic stromal cells cultured in the presence of a bone resorption factor such as active-form vitamin D<sub>3</sub> and parathyroid hormone (PTH). In addition, the present inventors have investigated the characteristics and physiological functions of this OCIF-binding protein and found that the protein exhibits biological activity of a factor which supports or promotes the osteoclast differentiation and maturation from immature precursors of osteoclasts. These findings have led to the completion of the present invention. Further investigation into the protein of the present invention has proven that this is an important protein controlling the differentiation and maturation of osteoclasts from immature precursors of osteoclasts in a co-culture system of the osteoblastic stromal cells and spleen cells. The success in identification and isolation of the protein which functions as a factor supporting or promoting differentiation and maturation of osteoclasts in the present invention has enabled screening for a novel medicine useful for abnormality of bone metabolism based on mechanism of bone metabolism utilizing the protein of the present invention.

[0013] Accordingly, an object of the present invention is to provide a novel protein (OCIF-binding molecule or OBM) which binds to osteoclastogenesis inhibitory factor (OCIF), and a method to produce this protein.

[0014] Another object of the present invention is to provide DNA encoding this protein, proteins containing an amino acid sequence encoded by this DNA, a method for producing this protein utilizing genetic engineering techniques, and pharmaceutical compositions comprising this protein.

[0015] A further object of the present invention is to provide methods for screening substances which control expression of this protein using this protein and the DNA, substances inhibiting or regulating the biological activity of this protein, receptors transducing the action of the protein by binding to the protein, substances obtained by the screening, and pharmaceutical compositions which comprises these substances.

[0016] A still further object of the present invention is to provide antibodies against this protein, methods for preparing the antibodies, and pharmaceutical compositions comprising these antibodies.

[0017] The protein of the present invention has the following physicochemical properties and biological activity.

[0018] (a) Affinity: specifically binds to the osteoclastogenesis inhibitory factor (OCIF) and exhibits high affinity to OCIF (dissociation constant on cell membrane:  $K_d=10^{-9}$  M or less);

[0019] (b) Molecular weight: has a molecular weight of approximately 30,000-40,000 when determined by SDS-polyacrylamide gel electrophoresis (SDS-PAGE) under non-reducing conditions and an appar-

ent molecular weight of approximately 90,000-110,000 when cross-linked to a monomer form OCIF; and

[0020] (c) Biological activity: exhibits activity supporting or promoting osteoclast differentiation and maturation in a co-culture system of the mouse osteoblastic stromal cells and mouse spleen cells in the presence of bone resorption factors such as active-form vitamin D<sub>3</sub> and parathyroid hormone (PTH).

[0021] A co-culture system of ST2, a mouse osteoblastic stromal cell line, and mouse spleen cells in the presence of active-form vitamin D<sub>3</sub> or PTH is well known as a typical in vitro culture system for osteoclast formation. The cells expressing the protein of the present invention can be determined by testing the binding of OCIF to mouse osteoblastic stromal cells or mouse spleen cells cultured in the presence or absence of active-form vitamin D<sub>3</sub>. The protein of the present invention is specified as the protein which is induced specifically on the osteoblastic stromal cells cultured in the presence of an osteotropic factor such as active-form vitamin D<sub>3</sub> or PTH. In addition, the protein of the invention can be specified as a protein exhibiting biological activity supporting or promoting differentiation and maturation of osteoclasts from the following results. That is, the osteoclast formation is inhibited dose dependently by the addition of 1 to 40 ng/ml of OCIF to the above-mentioned co-culture system in the presence of the active-form vitamin D<sub>3</sub>, the time course of expression of the protein of the present invention on ST2 cells in the presence of active-form vitamin D<sub>3</sub> well correlates with the time course of osteoclast formation in the co-culture. In addition, the amount of protein of the present invention expressed on ST2 cells correlates with the capability of the cells to support the osteoclast formation, and the binding of OCIF to the protein of the present invention on the ST2 cells completely suppresses osteoclasts formation.

[0022] The affinity of the protein of the present invention to OCIF can be evaluated by labeling OCIF and examining the binding of the labeled OCIF to the surface of animal cell membrane. OCIF can be labeled by a conventional protein-labeling method such as radioisotope or fluorescent labeling. Labeling of tyrosine residues with <sup>125</sup>I can be given as a specific example of labeling of the OCIF with an radioisotope. Labeling methods such as iodogen method, chloramine T method, and enzymatic method can be utilized. The binding of the labeled OCIF to the surface membrane of animal cell can be tested by a conventional method. The addition of unlabeled OCIF to the medium used for the binding assay to a concentration, 100 to 400 times the concentration of labeled OCIF, ensures measurement of non-specific binding. The amount of specific binding of OCIF can be calculated by subtracting the amount of non-specific binding from the total amount of binding of the labeled OCIF. The affinity of the protein of the present invention expressed on the cell membrane to OCIF can be evaluated by changing the amount of labeled OCIF and analyzing the specific binding by Scatchard plot.

[0023] The determined affinity of the protein of the present invention to OCIF is approximately 100-500 pM. The protein of the present invention is specified by a high affinity (dissociation constant on cell membrane:  $K_d=10^{-9}$  M or

less) to osteoclastogenesis inhibitory factor (OCIF). The molecular weight of OBM can be accessed by gel filtration chromatography, SDS-PAGE, or the like. SDS-PAGE is preferred in order to accurately determine the molecular weight. The OBM is specified as a protein having a molecular weight of approximately 40,000 ( $40,000\pm 4,000$ ) under reducing conditions.

[0024] The protein of the present invention can be obtained from mouse osteoblastic stromal cell line, ST2, mouse preadipocyte cell line, PA6, human osteoblastic cell lines, or other osteoblastic cells selected from mammals such as humans, mice, or rats. As the substances to induce expression of the protein of the present invention, osteotropic factors such as active-form vitamin D<sub>3</sub> (calcitriol), parathyroid hormone (PTH), interleukin (IL)-1, IL-6, IL-11, Oncostatin M, and leukemia inhibitory factor (LIF) can be given. These substances can be added in the concentration of  $10^{-8}$  M (active-form vitamin D<sub>3</sub> and PTH), 10 ng/ml (IL-11), or 1 ng/ml (Oncostatin M). IL-6 is preferably used at a concentration of 20 ng/ml with 500 ng/ml soluble IL-6 receptor. Preferably, confluent cells of mouse osteoblastic stromal cell line, ST2, cultured in  $\alpha$ -MEM medium to which  $10^{-8}$  M of active-form vitamin D<sub>3</sub>,  $10^{-7}$  M of dexamethasone, and 10% fetal bovine serum were added can be used. The cultured cells may be collected by scraping with a cell scraper. The collected cells may be stored at -80° C. until use.

[0025] The protein of the present invention can be purified efficiently from the membrane fractions of the collected cells. The membrane fractions can be prepared by a conventional method which is used to prepare intracellular organella. Various types of protease inhibitors may be added to the buffer solution used for the preparation of the membrane fractions. Examples of the protease inhibitors include serine protease inhibitors, thiol protease inhibitors, and metaprotease inhibitors. PMSF, APMSF, EDTA, o-phenanthroline, leupeptine, pepstatin A, aprotinin, soybean trypsin inhibitor are given as specific examples. A Dounce homogenizer, a polytron homogenizer, or a ultrasonic processor can be used to homogenize the cells. The cell homogenate is suspended in a buffer solution containing 0.5 M sucrose and centrifuged for 10 minutes at 600×g, to separate the nucleus and undisrupted cells as precipitate. The supernatant is centrifuged for 90 minutes at 150,000×g to obtain a membrane fractions as precipitate. The obtained membrane fraction is treated by various types of detergents to efficiently solubilize and extract the protein of the present invention from the cell membrane. Detergents which are commonly used to solubilize cell membrane proteins, such as CHAPS (3-[3-cholamidopropyl]-dimethylammonio]-1-propanesulfonate), Triton X-100, Nikkol, and n-octylglycoside, can be used. Preferably, 0.5% CHAPS is added to the membrane fraction and the mixture is stirred for 2 hours at 4° C. to solubilize the protein of the present invention. The sample thus prepared is centrifuged at 150,000×g for 60 minutes to obtain the solubilized membrane fraction as a supernatant.

[0026] The protein of the present invention can be purified from the solubilized membrane fraction with a column, gel, or resin coupled with OCIF. The immobilized OCIF may be that isolated from a culture broth of human embryonic lung fibroblasts, IMR-90, described in WO 96/26217 or rOCIF prepared using genetic engineering techniques. rOCIF can

be prepared by introducing human cDNA, mouse cDNA, or rat cDNA into an expression vector according to a conventional method, transducing the constructed vector in animal cells such as CHO cells, BHK cells, or Namalwa cells, or in insect cells to produce rOCIF, and purifying rOCIF. Obtained OCIF has a molecular weight of approximately 60 kDa (monomer-form) or 120 kDa (dimer-form). The dimer-form OCIF is preferable for immobilization. Given as examples of the gels and resins to which OCIF is immobilized are ECH Sepharose 4B, EAH Sepharose 4B, Thiopropyl Sepharose 6B, CNBr-activated Sepharose 4B, activated CH Sepharose 4B, Epoxy activated Sepharose 6B, activated thiol Sepharose 4B (these are manufactured by Pharmacia Co.), TSKgel AF-Epoxy Toyopal 650, TSKgel AF-Amino Toyopal 650, TSKgel AF-Formyl Toyopal 650, TSKgel AF-Carboxy Toyopal 650, TSKgel AF-Tresyl Toyopal 650 (these are manufactured by Tosoh Corporation), Amino-Cellulofine, Carboxy-Cellulofine, FMP activated Cellulofine, Formyl-Cellulofine (these are manufactured by Seikagaku Kogyo Co.), Affigel 10, Affigel 15, and Affiprep 10 (these are manufactured by BioRad Co.). As columns to which OCIF is immobilized, HiTrap NHS-activated column (Pharmacia Co.), TSKgel Tresyl-5PW (Tosoh Corporation), etc. can be given. As a specific example of the method for immobilizing OCIF to a HiTrap NHS-activated column (1 ml, Pharmacia Co.), the following method can be given. Specifically, 1 ml of 0.2M NaHCO<sub>3</sub>/0.5 M NaCl solution (pH 8.3) containing 13.0 mg of OCIF is injected to the column to perform coupling reaction at room temperature for 30 minutes. 0.5 M ethanolamine/0.5 M NaCl (pH 8.3) and 0.1 M acetic acid/0.5 M NaCl (pH 4.0) are sequentially applied to the column. Then, the column is again washed with 0.5 M ethanolamine/0.5 M NaCl (pH 8.3) and the column is allowed to stand for one hour at room temperature to block excess active groups. The column is sequentially washed twice with 0.5 M ethanolamine/0.5 M NaCl (pH 8.3) and 0.1 M acetic acid/0.5 M NaCl (pH 4.0), and then washed with 50 mM Tris/1M NaCl/0.1% CHAPS solution (pH 7.5), thereby obtaining a OCIF-immobilized column. The protein of the present invention can be efficiently purified by a OCIF-immobilized column prepared in this manner, or an OCIF-immobilized gel or resin.

[0027] It is desirable to add the various above-mentioned protease inhibitors to the buffer solutions used for the purification of the protein to suppress degradation of the protein of the present invention. The protein of the present invention can be purified by loading the above-mentioned solubilized membrane fraction on the OCIF-immobilized column or by mixing with the OCIF-immobilized gel or resin, and eluting the protein from the column, gel, or resin with acid, various protein denaturing agents, cacodylate buffer, and the like. It is desirable to use an acid for elution and to neutralize immediately after elution to minimize denaturation of the protein of the present invention. As the acidic solution used for elution, 0.1 M glycine-hydrochloric acid solution (pH 3.0), 0.1 M glycine-hydrochloric acid solution (pH 2.0), 0.1 M sodium citrate solution (pH 2.0), and the like can be given.

[0028] The protein of the present invention can be further purified by conventional purification methods used for purification of various proteins from biological materials and by various purification methods utilizing the physicochemical properties of this protein. To concentrate solutions containing the protein of the present invention, conventional tech-

niques used in the purification process for proteins such as ultra filtration, freeze drying, and salting-out, can be used. Ultra filtration with Centricon-10 (BioRad Co.), for example, is preferably used. As a means for the purification, various techniques conventionally utilized for the purification of proteins, such as ion exchange chromatography, gel filtration chromatography, hydrophobic chromatography, reverse phase chromatography, and preparative electrophoresis, are used in combination. More specifically, it is possible to purify the protein of the present invention by a combination of gel filtration chromatography with Superose 12 column (Pharmacia Co.) and reverse phase chromatography. To detect the protein of the present invention in the purification process, the binding activity of the protein of the present invention to the immobilized OCIF is examined or the material bound to the immobilized OCIF is immuno precipitated with an anti-OCIF antibody and analyzed by SDS-polyacrylamide gel electrophoresis (SDS-PAGE).

[0029] The obtained protein of the present invention is useful as an agent for treating diseases caused by abnormality of bone metabolism such as osteopetrosis or as a reagent for research and diagnosis of these diseases.

[0030] The present invention further provides DNA encoding a novel protein (OCIF-binding molecule or OBM) which binds to osteoclastogenesis inhibitory factor, proteins containing the amino acid sequence encoded by this DNA, a method for the preparation of this protein by the genetic engineering technique, and pharmaceutical compositions comprising this protein. Furthermore, the present invention provides methods for screening substances to regulate expression of OBM, a method for screening substances inhibiting or modifying the biological activity of OBM, or a method for screening receptors transducing the action of OBM by binding to OBM, and pharmaceutical compositions which comprises substances obtained as a result of the screening.

[0031] The novel protein OBM which is encoded by the DNA of the present invention has the following physico-chemical properties and biological activity.

[0032] (a) binds specifically to osteoclastogenesis inhibitory factor (OCIF),

[0033] (b) has a molecular weight of approximately 40,000 ( $\pm 4,000$ ) when determined by SDS-PAGE under reducing conditions and an apparent molecular weight of approximately 90,000-110,000 when crosslinked to monomer-form OCIF, and

[0034] (c) exhibits activity supporting or promoting differentiation and maturation of osteoclasts.

[0035] Human osteoclastogenesis inhibitory factor (OCIF) which is used as a probe to identify the DNA encoding OBM, the OCIF-binding molecule of the present invention, and to evaluate properties of OBM can be isolated from a culture broth of a human embryonic lung fibroblast cell line, IMR-90, according to WO No. 96/26217. Recombinant human OCIF, recombinant mouse OCIF, recombinant rat OCIF, and the like can also be used for the isolation and identification of the DNA coding OBM. These recombinant OCIF proteins can be produced by inserting DNA fragments encoding these proteins into expression vectors according to conventional methods, expressing in animal cells such as CHO cells, BHK cells, or Namalwa cells, or in insect cells, and purifying them.

[0036] As a method for isolating cDNA encoding a target protein (cDNA cloning), the method comprising determination of a partial amino acid sequence of the protein and isolation of the target cDNA by hybridization utilizing the nucleotide sequence corresponding to the amino acid sequence can be employed. Another available method, even in the case where the amino acid sequence of the protein is not known, comprises constructing a cDNA library in a expression vector, introducing the cDNA into cells, and screening for the expression of the target protein to isolate the objective cDNA (expression cloning method, D'Andrea et al.: Cell 57, 277-285, 1989; Fukunaga et al.: Cell 61, 341-350, 1990). In the expression cloning method, suitable host cells such as bacteria, yeast, animal cells, and the like are selected depending on the objective. In many cases, animal cells are selected as the host cells for cloning cDNA encoding a protein such as the protein of the present invention which is considered to be present in animal cell membrane surface. Normally, host cells showing high efficiency for DNA transfection and achieving expression of the introduced DNA at high levels are selected. One of such animal cells is the monkey kidney cells, COS-7, used in the present invention. Because SV40 large T antigen is expressed in the COS-7 cell, a plasmid which has a replicator of SV40 can be present as an episome of multiple copies in the cell, so that a high level of expression is expected. In addition, because expression of a target protein by COS-7 cells reaches a maximum within a few days after introduction of DNA, the cell is suitable for rapid screening. A combination of this host cell with a plasmid capable of high expression ensures gene expression of an extremely high level. The factor exhibiting the greatest influence on the expression of a gene on a plasmid is a promoter. Promoters such as SRα promoter and cytomegalovirus-derived promoters are used as high expression promoters. To screen for the cDNA encoding a membrane protein by the expression cloning strategy, screening procedures such as binding method, panning method, or film emulsion method are used.

[0037] The present invention relates to DNA encoding the protein (OBM) which specifically binds to OCIF, isolated by the combination of the expression cloning strategy and the screening by the binding method, to the expressed protein, and to screening of physiologically active substances using the DNA or the expressed protein. OBM encoded by the DNA of the present invention can be detected by labeling OCIF and testing the binding of the labeled OCIF to membrane surface of an animal cell. OCIF can be labeled by a conventional labeling method such as radioisotope labeling method or fluorescent labeling method which is used for labeling common proteins. Labeling tyrosine residues by  $^{125}\text{I}$  can be given as a specific example of labeling OCIF with a radioisotope. Labeling methods such as the iodogene method, chloramine T method, and enzymatic method can be utilized. The binding of labeled OCIF to the animal cell membrane surface can be tested by conventional methods. The addition of unlabeled OCIF to the medium used for the test to a concentration, 100 to 400 times the concentration of labeled OCIF, enables quantification of the amount of non-specific binding. The amount of specific binding of OCIF can be calculated by subtracting the amount of non-specific binding from the total amount of binding of the labeled OCIF.

[0038] The present inventors assumed that there is interaction between the factor involved in differentiation of

osteoclasts and OCIF. Based on this assumption, to isolate the protein to which recombinant OCIF binds, the inventors screened the expression library prepared from mRNA of mouse osteoblastic stromal cell line, ST2, according to the following method. Specifically, DNA synthesized using ST2 mRNA was inserted into an expression vector for animal cells and the vector with the insert was introduced into monkey kidney COS-7 cells. The objective protein expressed on the COS-7 cells was screened using OCIF labeled with <sup>125</sup>I as a probe. As a result, DNA encoding the protein which binds specifically to OCIF was isolated. The nucleotide sequence of the DNA encoding this OCIF-binding molecule (OCIF-binding molecule; OBM) was then determined. Moreover, OBM encoded by this DNA was found to bind specifically and strongly to OCIF, on the cell membrane.

[0039] Comparatively mild conditions for hybridization of DNA in the present invention are the conditions, for example, wherein DNA is transferred to a nylon membrane and immobilized thereto according to conventional methods and hybridized in a buffer solution for hybridization with a probe DNA labeled with an isotope at a temperature of 40-70° C. for about 2 hours to overnight, followed by washing in 0.5×SSC (0.075 M sodium chloride and 0.0075 M sodium citrate) at 45° C. for 10 minutes. Specifically, Highbond N (Amersham Co.) is used as the nylon membrane to transfer and immobilize DNA thereon. DNA is then hybridized with a probe DNA labeled with <sup>32</sup>P in a rapid hybridization buffer (Amersham Co.) at 65° C. for 2 hours, followed by washing with 0.5×SSC (0.075 M sodium chloride and 0.0075 M sodium citrate) at 45° C. for 10 minutes.

[0040] A co-culture system of mouse osteoblastic stromal cells, ST2, and mouse spleen cells in the presence of active-form vitamin D<sub>3</sub> or PTH is well known as a typical in vitro culture system for osteoclast-formation. The protein of the present invention is specified as the protein which is induced specifically on the osteoblastic stromal cells cultured in the presence of an agent which accelerates bone resorption such as active-form vitamin D<sub>3</sub> or PTH. In addition, because of the fact that formation of osteoclasts is stimulated by the addition of the protein encoded by the DNA of the present invention to mouse spleen cells cultured even in the absence of active-form vitamin D<sub>3</sub> or PTH, OBM which is encoded by the DNA of the present invention is considered to be involved in the differentiation and maturation of osteoclasts.

[0041] Recombinant OBM can be produced by inserting the DNA of the present invention into an expression vector to construct a plasmid and introducing the plasmid into various cells or microorganisms to express recombinant OBM. As a host in which recombinant OBM is expressed, mammalian cells such as COS-7, CHO, Namalwa, or bacteria such as *Escherichia coli* can be used. OBM may be expressed as a membrane-bound-form protein using the full length DNA or as a secretion-form or a soluble-form protein by removing the portion encoding the transmembrane domain. The produced recombinant OBM can be efficiently purified using a suitable combination of conventional purification methods used for common proteins, such as affinity chromatography using OCIF-immobilized columns, ion exchange chromatography, and gel filtration chromatography. The obtained protein of the present invention is useful as an agent for treating diseases caused by abnormality of

bone metabolism such as osteopetrosis or as a reagent for research and diagnosis of such diseases.

[0042] The following screening operations can be carried out using the protein OBM encoded by the DNA of the present invention: (1) screening of substances which regulate expression of OBM, (2) screening of substances which specifically bind to OBM and inhibit the biological activity of OBM, and (3) screening of proteins which are present in osteoclast precursor cells and transduce the biological activity of OBM (OBM receptor). It is also possible to develop antagonists and agonists using this OBM receptor. In the combinatorial chemistry using the above-mentioned OBM or OBM receptor, a peptide library used for the screening of the antagonists or agonists can be prepared by the following method. Specifically, one of the methods is split method (Lam et al.; *Nature* 354, 82-84, 1991). According to this method, synthetic carriers (beads) each comprising a specific amino acid (unit) bound thereto are prepared separately for all units. The synthesized carriers are mixed altogether and divided into portions equal to the number of the units. Then, the next units are bound. This procedure is repeated "n" times to produce a library containing carriers to which "n" units are bound. According to this synthetic method, each carrier pool has one type of sequence. Therefore, it is possible to identify a peptide specifically binding to the protein of the present invention by selecting the pool which gives a signal positive in this screening method using the protein of the present invention, and determining the amino acid sequence of the peptide bound on the pool. Another method is phage display method which utilizes phage carrying synthetic DNA which encode peptides with random amino acid sequences. The method has the advantage of increasing the number of molecules in the library as compared with the above-mentioned synthetic peptide library method, but has the disadvantage of less variety for a given number of molecules because there can be particular sequences which are missing in the library if the phages are unable to express those sequences. In the phage display method, the screening system using the protein of the present invention can also be applied to determine the nucleotide sequence encoding the peptide. That is, the phage specifically binding to the protein of the present invention is concentrated by panning, the selected phage is amplified in *Escherichia coli*, and the nucleotide sequence encoding the peptide is determined. In addition, a peptide exhibiting high specificity and high affinity to OBM or OBM receptor can be screened from a peptide library using the screening systems mentioned above in (2) and (3) by screening in the presence of OBM or OCIF while increasing the concentration of OBM or OCIF. Only positive carrier pools or phages are selected in this manner. For example, low molecular weight peptide agonists exhibiting an EPO (erythropoietin)-like activity were screened from a peptide library using a receptor of erythropoietin (EPO) which is a hematopoietic hormone, the tertiary structure of this substance was analyzed, and based on this tertiary structure, low molecular weight substances (antagonist) exhibiting the EPO-like activity were synthesized (Nicholas et al.; *Science*, 273, 458-463, 1996).

[0043] The present inventors have previously discovered using the osteoclastogenesis inhibitory factor, OCIF, that an OCIF-binding protein is specifically expressed on osteoblastic stromal cell line, ST2, cultured in the presence of a osteotropic factor such as active-form vitamin D<sub>3</sub> or par-

thyroid hormone (PTH). The inventors further found that this protein exhibits a biological activity to support or stimulate differentiation or maturation of osteoclasts from immature osteoclast precursor cells, and clarified various physicochemical properties and the biological activity of this protein by purification thereof. In order to compare the recombinant OBM expressed by the DNA of the present invention and the above-mentioned purified natural type protein which specifically binds to OCIF, the present inventors investigated the physicochemical properties and biological activities of the two proteins. As a result, the two proteins were confirmed ① to be both membrane-bound proteins which specifically bind to OCIF, ② to have molecular weights of approximately 40,000 determined by SDS-PAGE, and ③ to have apparent molecular weights of about 90,000-110,000 when cross-linked to a monomer form OCIF. Not only are these physicochemical properties identical, but both proteins exhibit a biological activity to support or stimulate differentiation or maturation of osteoclasts, suggesting the possibility that these are the same protein. In addition, a rabbit anti-OBM polyclonal antibody produced using the purified protein prepared by expressing the DNA of the present invention by a genetic engineering technique (recombinant OBM) was confirmed to cross react with the above-described purified natural type protein, to inhibit specific binding of this purified natural type protein and OCIF in the same manner as the antibody inhibits specific binding of OBM and OCIF. Based on these results, it is clear that the recombinant OBM expressed by the DNA of the present invention is identical to the natural type protein which specifically binds to OCIF.

[0044] To isolate a gene (cDNA) encoding human OCIF-binding protein (hereinafter called human OBM) which specifically binds to OCIF and exhibits the activity to support and stimulate differentiation and maturation of osteoclasts from mouse spleen cells in the same manner as the natural type or recombinant mouse OBM dose, a cDNA library prepared from mRNA derived from human lymph nodes was screened using a human OBM cDNA fragment as a probe. The human OBM cDNA fragment was obtained by polymerase chain reaction (PCR) in accordance with the method mentioned above using both cDNA prepared from human lymph node as a template and the primer which was prepared from mouse OBM cDNA. As a result, cDNA encoding the human protein which specifically binds to OCIF was isolated and the nucleotide sequence of the cDNA encoding this human OCIF-binding protein molecule (i.e. the cDNA encoding human OBM) was determined. Similar to mouse OBM, this human OBM encoded by the cDNA has characteristics to bind to OCIF strongly and specifically on the cell membrane and exhibits the activity to support and promote differentiation and maturation of osteoclasts from mouse spleen cells. Specifically, the present invention provides DNA encoding novel human OBM protein which binds to osteoclastogenesis inhibitory factor (OCIF), a protein which possesses the amino acid sequence encoded by the DNA, a method for producing the protein exhibiting characteristics of specifically binding to OCIF and the activity to support and promote differentiation and maturation of osteoclasts from mouse spleen cells by genetic engineering techniques, pharmaceutical compositions comprising this protein for the treatment of diseases caused by abnormality of bone metabolism, a method for screening substances regulating expression of human OBM, a method

for screening substances which inhibit or modulate the activity of human OBM by binding to it, a method for screening receptors which bind to human OBM and transmit the action of OBM, and a pharmaceutical compositions comprising the substances obtained by these screenings.

[0045] The present invention further provides DNA encoding novel human OBM protein which specifically binds to OCIF and exhibits the biological activity to support and promote differentiation and maturation of osteoclasts, a protein which possesses the amino acid sequence encoded by the DNA, a method for producing the protein exhibiting characteristics of specifically binding to OCIF and the activity to support and promote differentiation and maturation of osteoclasts by genetic engineering techniques, and pharmaceutical compositions comprising this protein for the treatment of diseases causing abnormality of bone metabolism. Furthermore, the present invention provides a method for screening substances regulating expression of human OBM, a method for screening substances which inhibit or modulate the activity of human OBM by binding to it, a method for screening receptors binding to human OBM and transmitting the action of OBM, antibodies against human OCIF binding protein, and pharmaceutical compositions comprising these antibodies for the prevention or treatment of diseases causing abnormality of bone metabolism.

[0046] The novel, human OCIF-binding protein molecule (OBM) which is encoded by the DNA of the present invention has the following physicochemical properties and biological activity.

[0047] (a) binds specifically to osteoclastogenesis inhibitory factor (OCIF) (WO 96/26217),

[0048] (b) has a molecular weight of approximately 40,000 ( $\pm$ 5,000) when determined by SDS-PAGE under reducing conditions and an apparent molecular weight of approximately 90,000-110,000 when crosslinked with a monomer form OCIF, and

[0049] (c) exhibits activity to support and stimulate differentiation and maturation of osteoclasts.

[0050] Mouse OBM cDNA which encodes mouse OCIF-binding protein and used as a probe to isolate and identify the cDNA encoding human OBM of the present invention, can be isolated according to the above-mentioned method from a cDNA library of mouse osteoblastic stromal cell line, ST2. Human osteoclastogenesis inhibitory factor (OCIF) which is necessary to evaluate the properties and the biological activity of the protein obtained by expression of human OBM cDNA, can be prepared according to the method described in WO 96/26217 by isolating from a culture broth of human fibroblast cell line, IMP-90, or by genetic engineering techniques using the DNA encoding OCIF. Recombinant human OCIF, recombinant mouse OCIF, recombinant rat OCIF, or the like can be used for the assessment of the properties and biological activity of human OBM. These recombinant OCIF can be obtained according to conventional methods by inserting cDNA into an expression vector, expressing the cDNA in animal cells such as CHO cells, BHK cells, or Namalwa cells, or in insect cells, and purifying the expressed proteins.

[0051] The following methods can be used to isolate human cDNA encoding the target protein (cDNA cloning).  
① A method comprising purifying the protein, determining

the partial amino acid sequence of the protein, and isolating the target cDNA by hybridization using the DNA fragment comprising nucleotide sequence corresponding to the amino acid sequence as a probe, ② a method applied even in the case where the amino acid sequence of the protein is not known, which comprises constructing a cDNA library in a expression vector, introducing the cDNA library into cells, and screening for the expression of the target protein to isolate the objective cDNA (expression cloning method), and ③ a method of isolating cDNA encoding the target human protein from the cDNA library constructed using human cells or tissues by hybridization or by the use of polymerase chain reaction (PCR) using the cDNA encoding the protein of mammalian origin (other than human) which possesses the same characteristics and biological activity as the target protein of human origin as a probe, assuming that the cDNA probe has high homology with the human-origin cDNA which to be cloned. Based on the assumption that human OBM cDNA has a high homology with mouse OBM cDNA, it is possible to determine which cells or tissues produce human OBM by Northern hybridization method using the mouse OBM cDNA as a probe. Human OBM cDNA can be obtained by the following method using the mouse OBM primer prepared from the mouse OBM cDNA. Human OBM cDNA fragments can be prepared by the PCR method using cDNA prepared from human OBM-producing tissues such as human lymph nodes as a template. These human OBM cDNA fragments are used as probes for screening the cDNA library of human OBM-producing cells or tissues which were identified according to the method mentioned above. The present invention relates to the DNA encoding human OBM which has characteristics of specific binding to OCIF and exhibits activity to support and promote differentiation and maturation of osteoclasts. Because the OBM which is encoded by the DNA of the present invention is a membrane-bound type protein which comprises a transmembrane domain, this protein can be detected by labeling OCIF and by examining the binding of the labeled OCIF to the surface of animal cells in which the cDNA of the present invention was expressed. The above-described labeling method using radioisotope or fluoresceine conventionally applied to labeling proteins can be used for labeling OCIF.

[0052] The molecular weight of the protein expressed by the human OBM cDNA of the present invention can be accessed by gel filtration chromatography, SDS-PAGE, or the like. In order to accurately determine the molecular weight, it is desirable to use the SDS-PAGE method, by which human OBM was specified as a protein having a molecular weight of approximately 40,000 ( $40,000 \pm 5,000$ ) under reducing conditions.

[0053] Comparatively mild conditions for hybridization of DNA in the present invention are the conditions, for example, wherein DNA is transferred to a nylon membrane and immobilized thereto according to a conventional method and hybridized with a probe DNA labeled with an isotope in a buffer solution for hybridization at a temperature of 40-70° C. for about 2 hours to overnight, followed by washing in 0.5×SSC (0.075 M sodium chloride and 0.0075 M sodium citrate) at 45° C. for 10 minutes. Specifically, Highbond N (Amersham Co.) is used as the nylon membrane to transfer and immobilize DNA thereon. The DNA is then hybridized with a probe DNA labeled with  $^{32}\text{P}$  in a rapid hybridization

buffer (Amersham Co.) at 65° C. for 2 hours, followed by washing with 0.5×SSC at 45° C. for 10 minutes.

[0054] A co-culture system of mouse osteoblastic stromal cells, ST2, and mouse spleen cells in the presence of active-form vitamin D<sub>3</sub> or PTH is well known as a typical in vitro culture system for osteoclast-formation. Interaction by adhesion of osteoblastic stromal cells and spleen cells and presence of an osteotropic factor such as active-form vitamin D<sub>3</sub> or PTH are indispensable for the osteoclasts formation in this in vitro culture system. In this in vitro culture system, COS cells, monkey kidney cells having no osteoclast formation-supporting capability, acquire capability to support osteoclasts formation from spleen cells in the absence of an osteotropic factor when the cDNA of the present invention was expressed as osteoblastic stromal cell line ST2 did. Based on the fact that the cDNA of the present invention encodes a protein comprising a transmembrane domain form, this cDNA can be expressed as a secretion form or soluble-form by removing the part which encodes this transmembrane domain. It was confirmed that osteoclasts can be formed by the addition of the secretion form human OBM to the above-mentioned in vitro culture system in the absence of osteotropic factors. Based on these results, the human OBM which is encoded by the cDNA of the present invention is specified as the factor involved in the differentiation and maturation of osteoclasts.

[0055] A recombinant human OBM can be prepared by inserting the cDNA of the present invention into an expression vector, preparing a human OBM expression plasmid, introducing the plasmid into various cell strains and expressing OBM in the cells. Mammalian cells such as COS-7, CHO, Namalwa cells, or bacteria such as *Escherichia coli* can be used as a host for expressing OBM. In this case, OBM may be expressed as a membrane-bound-form protein, using the full length DNA, or as a secretion-form or soluble-form protein by removing the part encoding the transmembrane domain. The recombinant OBM thus produced can be efficiently purified using a suitable combination of conventional purification methods used for common proteins such as affinity chromatography using OCIF immobilized columns, ion exchange chromatography, and gel filtration chromatography. Human OBM of the present invention thus obtained is useful as an agent for treating diseases caused by abnormality of bone metabolism such as osteopetrosis or as a reagent for research and diagnosis of such diseases.

[0056] The following screening operations can be carried out using the protein OBM encoded by the DNA of the present invention: (1) screening of substances which can regulate expression of human OBM, (2) screening of substances which specifically bind to human OBM and inhibit or modify the biological activity of OBM, and (3) screening of human proteins which are present in osteoclast precursor cells and transmit the biological activity of human OBM (human OBM receptor). It is also possible to develop antagonists and agonists using this human OBM receptor. In the combinatorial chemistry using the human OBM or human OBM receptor, peptide libraries required for the screening of antagonists or agonists can be produced by the same method as used for the screening using mouse OBM. A peptide with extremely high specificity and affinity can be obtained by screening peptide libraries using human OBM instead of mouse OBM.

[0057] Although this OBM is very useful as mentioned above and antibodies specifically recognizing OBM and enzyme immunoassay using these antibodies are indispensable in determination of OBM concentration, no antibodies useful for the access of OBM concentration have been so far available. In addition, an anti-OBM antibody or anti-sOBM antibody which neutralizes the biological activity of OBM or sOBM is supposed to suppress the activity of OBM or sOBM, specifically the activity to induce osteoclasts formation. These are expected to be useful as therapeutic agents to treat abnormality of bone metabolism. However, no such antibodies have so far been available.

[0058] In view of this situation, the present inventors have conducted extensive studies. As a result, the present inventors have found antibodies (anti-OBM/sOBM antibodies) which recognize both OBM, a membrane-bound protein which specifically binds to osteoclastogenesis inhibitory factor (OCIF), and soluble OBM(sOBM) which lack a transmembrane domain. Accordingly, the present invention provides antibodies (anti-OBM/sOBM antibodies) which recognizes both OBM, a membrane-bound protein which specifically binds to osteoclastogenesis inhibitory factor (OCIF), and sOBM which lack a transmembrane domain; a method for the preparation thereof; a method for determination of OBM and sOBM concentrations using these antibodies; and agents for the prevention or treatment of diseases resulting from abnormality of bone metabolism.

[0059] The present invention relates to antibodies (anti-OBM/sOBM antibodies) which recognize both the OBM, a membrane-bound protein which specifically binds to osteoclastogenesis inhibitory factor (OCIF), and soluble OBM (sOBM) which lack a transmembrane domain; a method for the preparation thereof; a method for quantifying OBM and sOBM using these antibodies; and agents for the prevention or treatment of diseases resulting from abnormality of bone metabolism.

[0060] The antibodies of the present invention exhibit activity of neutralizing the osteoclastogenesis accelerating activity which is the biological activity of OBM and sOBM and comprises the antibodies having the following characteristics:

[0061] (a) polyclonal antibody which recognizes both mouse OBM and mouse sOBM (anti-mouse OBM/sOBM polyclonal antibody),

[0062] (b) polyclonal antibody which recognizes both human OBM and human sOBM (anti-human OBM/sOBM polyclonal antibody),

[0063] (c) monoclonal antibodies which recognizes both mouse OBM and mouse sOBM (anti-mouse OBM/sOBM monoclonal antibodies),

[0064] (d) monoclonal antibodies which recognize both human OBM and human sOBM (anti-human OBM/sOBM monoclonal antibodies), and

[0065] (e) anti-human OBM/sOBM monoclonal antibodies which crossreact to both mouse OBM and mouse sOBM.

[0066] The polyclonal antibody which recognizes both mouse OBM and mouse sOBM (hereinafter called anti-mouse OBM/sOBM polyclonal antibody) and the polyclonal antibody which recognizes both human OBM and human

sOBM (hereinafter called anti-human OBM/sOBM polyclonal antibody) were produced by the following method. The purified mouse OBM used as an antigen for immunization can be obtained according to the above-mentioned method. Especially, mouse osteoblastic stromal cell line, ST2, was treated with active-form vitamin D<sub>3</sub>, and OBM on the cell membrane was purified using an OCIF-immobilized column and gel filtration chromatography, thereby obtaining natural mouse OBM (native OBM). The above-mentioned mouse OBM cDNA (Sequence Table, Sequence No. 15) or human OBM cDNA (Sequence Table, Sequence ID No. 12) was inserted into an expression vector according to conventional methods. Recombinant mouse OBM (Sequence Table, Sequence ID No. 1) and recombinant human OBM (Sequence Table, Sequence ID No. 11) can be obtained by expressing cDNA in animal cells such as CHO cells, BHK cells, Namalwa, or COS-7 cells, insect cells or *Escherichia coli*, and purifying them using the same purification methods as mentioned above. These may be used as antigens for immunization. In this instance, purifying a large amount and a high level of mouse OBM or human OBM which are membrane-bound proteins is a task requiring a great deal of labor. On the other hand, as mentioned above, OBM which is a membrane-bound protein and a soluble OBM (sOBM) which is obtained by deleting transmembrane domain of OBM are known to be almost the same in their osteoclast differentiation and maturation activities. It is possible to use mouse sOBM and human sOBM which are relatively easily expressed and purified to a high level, as antigens for immunization.

[0067] Mouse sOBM (Sequence Table, Sequence ID No. 16) and human sOBM (Sequence Table, Sequence ID No. 17) can be obtained by adding a nucleotide sequence encoding a known signal sequence originating from the other secretion protein in the upstream side of the 5' end of, respectively, mouse sOBM cDNA (Sequence Table, Sequence ID No. 18) and human sOBM cDNA (Sequence Table, Sequence ID No. 19), inserting these into an expression vector by the use of genetic engineering techniques, causing these proteins to be expressed in host cells such as various animal cells, insect cells, or *Escherichia coli*, and purifying the resultant products. The antigens for immunization thus obtained are dissolved in phosphate buffered saline (PBS), mixed with the same volume of Freund's complete adjuvant to emulsify the solution if required, and subcutaneously administered to animals about once a week to immunize these animals several times. A booster injection is given when the antibody titer reaches a maximum. Exsanguination is performed 10 days after the booster administration. The resulting antiserum is treated with ammonium sulfate precipitation. IgG fraction is purified using an anion exchange chromatography or purified by protein A- or protein G-Sepharose column chromatography after diluting the antiserum two-fold with Binding Buffer<sup>TM</sup> (BioRad Co.), to obtain the anti-mouse or anti-human OBM/sOBM polyclonal antibody.

[0068] The monoclonal antibodies of the present invention can be obtained according to the following method. In the same manner as in the case of the polyclonal antibodies, natural mouse OBM (native OBM), recombinant mouse or human OBM, or recombinant mouse or human sOBM can be used as immunogens to prepare monoclonal antibodies. Hybridomas are produced according to conventional methods by immunizing mammals with these antigens or by

immunizing lymphocytes in vitro and fusing the immunized cells with myeloma cells. By analyzing the hybridoma culture supernatant thus obtained by a solid phase ELISA method, antibody-producing hybridomas recognizing the highly purified antigen are selected. The resulting hybridomas are cloned and established as stable antibody-producing hybridoma clones. These hybridomas are cultured to obtain the antibodies. Small mammals such as mice or rats are commonly used to produce hybridomas. Animals are immunized by intravenously or intraperitoneally injecting the antigen diluted to a suitable concentration using a suitable solvent such as physiological salt solution. Optionally, Freund's complete adjuvant maybe used together with antigen. These are usually injected 3-4 times, once a week or every two weeks. The immunized animals are dissected three days after final immunization. Splenocytes from the removed spleen are used as immunized cells. As mouse myeloma to be fused with immunized cells, p3/x63-Ag8, p3-U1, NS-1, MPC-11, SP-2/0, FO, P3x63 Ag8.653, and S194 can be given. A cell line such as R-210 is given as the cell of rat origin. Human antibodies are produced by immunizing human B lymphocytes in vitro and fusing the immunized cells with human myeloma cells or a cell line transformed with EB virus. The fusion of the immunized cells and myeloma cells can be carried out according to a conventional method such as the method of Koehler and Milstein (Koehler et al.: *Nature* 256, 495-497 (1975)). A method using electric pulse is also applicable. Immunized lymphocytes and myeloma cells are mixed at a conventionally accepted ratio and fused in an FCS-free (fetal bovine serum-free) culture medium with an addition of polyethylene glycol, and cultured in an FCS-containing HAT selection medium to select fused cells (hybridomas). Next, the hybridomas which produce antibodies were selected by using a conventional antibody detection method such as an ELISA, a plaque technique, Ouchterlony method, or aggregation method, to establish stable hybridomas. The hybridomas established in this way can be subcultured by a conventional culture method or can be stored by freezing as required. A hybridoma can be cultured by a conventional method to collect the culture supernatant or implanted in the abdominal cavity of mammals to obtain the antibody from the ascitic fluid. The antibody in the culture supernatant or ascitic fluid can be purified by a conventional method such as salting out, ion exchange and gel filtration chromatography, or protein A or protein G affinity chromatography. Almost all monoclonal antibodies obtained using sOBM as an antigen can specifically recognize not only sOBM but also OBM (such antibodies are hereinafter called anti-OBM/sOBM monoclonal antibodies). These antibodies can be used for the quantification of OBM or sOBM. The amounts of OBM and sOBM can be quantified by labeling these antibodies with a radioisotope or an enzyme and by applying the labeled antibodies to a quantification system known as a radioimmunoassay (RIA) or enzyme immunoassay (EIA). Using these quantification systems, the amount of sOBM in a biological sample such as blood or urine can be determined with ease at high sensitivity. In addition, the amount of OBM binding to a tissue or surface of cells can be measured with ease at high sensitivity utilizing a binding assay using these antibodies.

[0069] When an antibody is used as a medication for humans, it is desirable to use a human-type anti-human OBM/sOBM antibody in view of antigenicity. The human-type anti-human OBM/sOBM antibody can be prepared

according to the following methods ①, ②, or ③. In the method ①, human lymphocytes collected from human peripheral blood or spleen are immunized with an antigen human OBM or human sOBM in vitro in the presence of IL-4. The resulting immunized human lymphocytes are fused with K6H6/B5 (ATCC CRL1823) which is a hetero hybridoma of mouse and human, and screened to obtain the objective antibody producing hybridoma. The antibodies produced by the resulting antibody producing hybridomas are human type anti-human OBM/sOBM monoclonal antibodies. The antibodies neutralizing the activity of human OBM/sOBM are selected from these antibodies. However, in general, it is difficult to produce an antibody exhibiting high affinity to an antigen by the method of immunizing human lymphocytes in vitro. Therefore, in order to obtain monoclonal antibodies with high affinity to human OBM and sOBM, it is necessary to increase the affinity of the human-type anti-human OBM/sOBM monoclonal antibodies obtained by the above method. This can be done according to the following method. First, a random mutation is introduced into CDR region (particularly CDR3 region) of a human-type anti-human OBM/sOBM monoclonal antibody which neutralize OBM but have a low affinity, and make the phage to express protein. Phages which can strongly bind to human OBM/sOBM which are selected by a phage display method using plates on which human OBM/sOBM antigens are immobilized. The selected phages are grown in *Escherichia coli*. The amino acid sequence of the CDR which exhibits high affinity is determined from the nucleotide sequence of the DNA cloned in the phage. The thus-obtained DNA encoding the human type anti-human OBM/sOBM monoclonal antibodies is introduced into a commonly used expression vector for mammalian cells to produce the human type anti-human OBM/sOBM monoclonal antibodies. The target human type anti-human OBM/sOBM monoclonal antibodies exhibiting high affinity and capable of neutralizing the biological activity of human OBM/sOBM can be selected from these monoclonal antibodies. In the method ②, mouse type anti-human OBM/sOBM monoclonal antibodies are produced according to the same method as in the present invention using BALB/cmouse (Koehler et al.: *Nature* 256, 495-49, 1975), and monoclonal antibodies which can neutralize the biological activity of human OBM/sOBM and exhibiting high affinity are selected. These high affinity mouse anti-human OBM/sOBM monoclonal antibodies can be converted into human-type using the CDR-grafting technique (Winter and Milstein: *Nature* 349, 293-299, 1991) by implanting its CDR regions (CDR-1, 2 and 3) into the CDR regions of human IgG. In the method ③, human peripheral blood lymphocytes are implanted into a severe combined immune deficiency (SCID) mouse. Because the implanted SCID mouse can produce human antibodies (Mosier D. E. et al.: *Nature* 335, 256-259, 1988; Duchosal M. A. et al.: *Nature* 355, 258-262, 1992), lymphocytes which can produce the human monoclonal antibodies having specificity to human OBM/sOBM can be collected by screening SCID mouse immunized with human OBM or sOBM. The resulting lymphocytes are fused with K6H6/B5 (ATCC CRL1823) which is a heterohybridoma of mouse and human, according to the procedure described above for the human antibodies in the method ①. The resulting hybridomas are screened to obtain hybridomas which can produce the objective human monoclonal antibodies. The thus-obtained hybridomas are cul-

tured to produce large amounts of the objective human monoclonal antibodies. The antibodies can be purified by the above-mentioned purification method. In addition, it is possible to produce recombinant human monoclonal antibodies in large amounts by constructing a cDNA library from the hybridoma which can produce the objective human monoclonal antibodies to obtain a gene (cDNA) encoding the objective human-type monoclonal antibodies by cloning, inserting this gene into a suitable expression vector by using genetic engineering techniques, and expressing the monoclonal antibodies in host cells such as various animal cells, insect cells, or *Escherichia coli*. A large amounts of purified human monoclonal antibodies can be obtained by purifying from the resulting culture supernatant by the purification methods mentioned above.

[0070] The antibodies which can neutralize the biological activity of OBM/sOBM can be obtained from the anti-OBM/sOBM monoclonal antibodies produced according to this method. The antibodies which neutralize the biological activity of OBM/sOBM are expected to be useful as agents for the treatment or prevention of bone metabolism abnormality because of their capability of blocking in vivo biological activity of OBM/sOBM, specifically the capability of preventing the induction osteoclast formation. The activity of anti-OBM/sOBM antibodies to neutralize the biological activity of OBM or sOBM can be measured by determining the activity to suppress osteoclast formation in the in vitro system. Specifically, the following in vitro osteoclastogenesis culture system can be given: ① a co-culture system of mouse osteoblastic stromal cell strain, ST2 cells, and mouse spleen cells in the presence of active-form vitamin D<sub>3</sub> and dexamethasone, ② a co-culture system comprising OBM expressing monkey kidney cell strain, COS-7, immobilizing the OBM-expressing cells with formaldehyde, and culturing mouse spleen cells on those cells in the presence of M-CSF, and ③ a culture system of mouse spleen cells in the presence of recombinant sOBM and M-CSF. The osteoclastogenesis-inhibitory activity of the anti-OBM/sOBM antibodies can be measured by adding the anti-OBM/sOBM antibodies at various concentrations to these culture systems and investigating their effects on osteoclast formation. The osteoclastogenesis-inhibitory activity of the anti-OBM/sOBM antibodies can also be evaluated by measuring their bone resorption-inhibitory activity utilizing experimental animals in vivo. Especially, ovariectomized animal model is given as an animal model with progressive osteoclast formation. The osteoclastogenesis-inhibitory activity of the anti-OBM/sOBM antibodies can be determined by administering the anti-OBM/sOBM antibodies to such experimental animals and evaluating the suppression of bone resorption (a bone density increasing activity).

[0071] The thus-obtained antibodies capable of neutralizing the OBM/sOBM biological activity are useful in pharmaceutical compositions, particularly pharmaceutical compositions to prevent or treat bone metabolism abnormality or as antibodies for an immunological diagnosis of such diseases. The preparations comprising the antibodies of the present invention can be administered either orally or non-orally. Such preparations can be safely administered to humans or animals as pharmaceutical compositions which contain the antibodies recognizing OBM and/or sOBM as an active component. As the forms of pharmaceutical composition, injection agents including intravenous drip, suspen-

tory agents, sublingual agents, percutaneous absorption agents, and the like are given. Because monoclonal antibodies are macromolecule proteins, they not only readily adhere to a glass container such as a vial or a syringe, but also are easily denatured by physicochemical factors such as heat, pH, or humidity. Therefore, the preparations should be stabilized by the addition of stabilizers, pH adjusters, buffering agents, solubilizing agents, or detergents. As the stabilizers, amino acids such as glycine and alanine, saccharides such as dextran 40 and mannose, and sugar alcohols such as sorbitol, mannitol, and xylitol can be given. These stabilizers may be used either individually or in combinations of two or more. The amount of stabilizers to be added is preferably from 0.01 to 100 times, particularly preferably from 0.1 to 10 times, the amount of the antibody. The addition of these stabilizers increases storage stability of liquid preparations or lyophilized products thereof. Phosphate buffers and citrate buffers are given as examples of the buffering agents. The buffering agents not only adjust the pH of the liquid preparations or aqueous solutions obtained by re-dissolving the lyophilized products thereof, but also increase stability and solubility of the antibody. It is desirable to add the buffering agent in an amount to make from 1 mM to 10 mM concentration of the liquid preparation or of the aqueous solution prepared from the lyophilized product. Polysolbate 20, Pulluronic F-68, and polyethylene glycol are given as examples of the detergent. A particularly preferred example is Polysolbate 80. These detergents may be used either individually or in combinations of two or more. Macromolecule proteins such as an antibody is easily adhere to glass containers. Adherence to containers of the antibody in a liquid preparation or in an aqueous solution prepared by re-dissolving a lyophilized product can be prevented by adding such detergents at a concentration from 0.001 to 1.0%. The preparations comprising the antibodies of the present invention can be obtained by adding stabilizers, buffering agents, or agents which prevent adsorption. When the preparations are used as injection agents for medication or for animals, such injection agents should preferably have an osmotic pressure ratio of 1 to 2. The osmotic pressure ratio can be adjusted by increasing or decreasing the amount of sodium chloride when making the preparations. The amount of an antibody in a preparation can be suitably adjusted depending on the disease, route of administration, and the like. A dose of a human antibody to humans may be changed depending on the affinity of the antibody to human OBM/sOBM, especially, on the dissociation constant (Kd value) to human OBM/sOBM. The higher the affinity (or the smaller the Kd value), the less the dose to be administered to humans to obtain a certain medicinal effect. Because a human-type antibody has a long half-life in blood of about 20 days, it is sufficient to administer it to humans at a dose of about 0.1-100 mg/kg once or more in a 1-30 day period.

#### BRIEF DESCRIPTION OF THE DRAWINGS

[0072] FIG. 1 shows the result of SDS-PAGE of mouse OBM protein of the present invention obtained in Example 3.

[0073] <Explanation of Symbols>

[0074] (A): Lane 1: Molecular weight markers

[0075] Lane 2: A partially purified sample (Gly-HCl (pH 2.0) elution fraction) obtained from ST2 cells cultured in the presence of active-form vitamin D<sub>3</sub> and dexamethasone.

[0076] Lane 3: A partially purified sample (Gly-HCl (pH 2.0) elution fraction) obtained from ST2 cells cultured in the absence of active-form vitamin D<sub>3</sub> and dexamethasone.

[0077] (B): Lane 1: Molecular weight markers

[0078] Lane 2: Mouse OBM protein of the present invention after purification by reverse phase high performance liquid chromatography (Example 3)

[0079] FIG. 2 shows the result of the binding assay of <sup>125</sup>I-labeled OCIF to osteoblastic stromal cells, ST2, in Example 4.

[0080] FIG. 3 shows the osteoclast formation capability of osteoblastic stromal cells ST2 from different generations in Example 5(1).

[0081] <Explanation of Symbols>

[0082] 1: Ability of ST2 cells from about a 10<sup>th</sup> subculture to support osteoclast formation.

[0083] 2: Ability of ST2 cells from about a 40<sup>th</sup> subculture to support osteoclast formation.

[0084] FIG. 4 shows change with the passage of time in expression of the protein of the present invention on the cell membrane of osteoblastic stromal cells cultured in the presence of active-form vitamin D<sub>3</sub> and dexamethasone in Example 5(2).

[0085] FIG. 5 shows change with the passage of time in osteoclast formation in the co-culture system of Example 5(2).

[0086] FIG. 6 shows the inhibitory effect on osteoclast formation when treated with OCIF for different culture periods during the co-culture period in Example 5(3).

[0087] FIG. 7 shows the results of a crosslinking test of <sup>125</sup>I-labeled OCIF with the protein of the present invention in Example 6.

[0088] <Explanation of Symbols>

[0089] Lane 1: <sup>125</sup>I-labeled OCIF-CDD1

[0090] Lane 2: <sup>125</sup>I-labeled OCIF-CDD1 crosslinked with ST2 cells

[0091] Lane 3: <sup>125</sup>I-labeled OCIF-CDD1 crosslinked in the presence of 400-fold excess of unlabeled OCIF

[0092] FIG. 8 shows the result of SDS-PAGE in Example 9.

[0093] <Explanation of Symbols>

[0094] Lane 1: Proteins of pOBM291-transfected COS-7 cells immunoprecipitated in the absence of OCIF

[0095] Lane 2: Proteins of pOBM291-transfected COS-7 cells immunoprecipitated in the presence of OCIF

[0096] FIG. 9 shows the results of analysis of binding capability of <sup>125</sup>I-labeled OCIF to COS-7 cells transfected with pOBM291 in Example 10.

[0097] <Explanation of Symbols>

[0098] Lanes 1 and 2: The amount of the <sup>125</sup>I-labeled OCIF binding to COS-7 cells transfected with pOBM291

[0099] Lanes 3 and 4: The amount of the <sup>125</sup>I-labeled OCIF binding to COS-7 cells transfected with pOBM291 in the presence of 400-fold excess of unlabeled OCIF

[0100] FIG. 10 shows the result of a crosslinking test using OCIF labeled with <sup>125</sup>I in Example 11.

[0101] <Explanation of Symbols>

[0102] Lane 1: <sup>125</sup>I-labeled OCIF

[0103] Lane 2: <sup>125</sup>I-labeled OCIF crosslinked with COS-7 cells transfected with pOBM291

[0104] Lane 3: <sup>125</sup>I-labeled OCIF crosslinked with COS-7 cells transfected with pOBM291 in the presence of 400-fold excess of unlabeled OCIF

[0105] FIG. 11 shows the result of a Northern Blot in Example 12.

[0106] <Explanation of Symbols>

[0107] Lane 1: RNA originating from ST2 cells cultured without addition of Vitamin D and dexamethasone

[0108] Lane 2: RNA originating from ST2 cells cultured with the addition of Vitamin D and dexamethasone

[0109] FIG. 12 shows the OCIF-binding capability of the proteins in the conditioned medium at various OCIF concentrations in Example 13(2).

[0110] <Explanation of Symbols>

[0111] ○: pCEP4

[0112] ●: pCEP sOBM

[0113] FIG. 13 shows the OCIF-binding capability of the protein in the conditioned medium at various proportions of the conditioned medium in Example 13(2).

[0114] <Explanation of Symbols>

[0115] ○: pCEP4

[0116] ●: pCEP sOBM

[0117] FIG. 14 shows the result of SDS-PAGE of a fusion protein consisting of thioredoxin and mouse OBM expressed in *Escherichia coli* in Example 14(2).

[0118] <Explanation of Symbols>

[0119] Lane 1: Molecular weight markers

[0120] Lane 2: Soluble protein fractions originating from GI724/pTrxFus

[0121] Lane 3: Soluble protein fractions originating from GI724/pTrxOBM25

[0122] FIG. 15 shows the OCIF-binding capability at various proportions of soluble protein fractions in Example 14(3).

[0123] <Explanation of Symbols>

[0124] □: GI724/pTrxFus

[0125] ○: GI724/pTrxOBM25

[0126] FIG. 16 shows the OCIF-binding capability of soluble protein fractions (1%) at various concentrations of OCIF in Example 14(3).

[0127] <Explanation of Symbols>

[0128] □: GI724/pTrxFus

[0129] ○: GI724/pTrxOBM25

[0130] FIG. 17 shows the results of inhibition of specific binding to OCIF of mouse OBM obtained by expression of mouse OBM cDNA of the present invention and purification or natural OCIF-binding protein by a rabbit anti-mouse OBM antibody.

[0131] <Explanation of Symbols>

[0132] 1: Purified OBM prepared by expression of the cDNA in the presence of the antibody, OBM+<sup>125</sup>I-OCIF

[0133] 2: The natural protein in the presence of the antibody+<sup>125</sup>I-OCIF

[0134] 3: Mouse OBM prepared by expression of the cDNA in the absence of the antibody, mouse OBM+<sup>125</sup>I-OCIF

[0135] 4: The natural protein in the absence of the antibody+<sup>125</sup>I-OCIF

[0136] 5: 3+unlabeled OCIF (400-fold more than <sup>125</sup>I-OCIF)

[0137] 6: 4+unlabeled OCIF (400-fold more than <sup>125</sup>I-OCIF)

[0138] FIG. 18 shows the result of SDS-PAGE of human OBM protein expressed by the cDNA of the present invention.

[0139] <Explanation of Symbols>

[0140] Lane 1: Molecular weight markers

[0141] Lane 2: Proteins of COS-7 cells transfected with phOBM (an expression vector containing a cDNA of the present invention), immunoprecipitated with a rabbit anti-OCIF polyclonal antibody in the absence of OCIF

[0142] Lane 3: Proteins of COS-7 cells transfected with phOBM (an expression vector containing a cDNA of the present invention), immunoprecipitated with a rabbit anti-OCIF polyclonal antibody in the presence of OCIF

[0143] FIG. 19 shows the result of analysis of binding of OCIF to COS-7 cells transfected with phOBM, an expression vector containing a cDNA of the present invention.

[0144] <Explanation of Symbols>

[0145] Lane 1: COS-7 cells transfected with phOBM and the addition of <sup>125</sup>I-OCIF.

[0146] Lane 2: COS-7 cells transfected with phOBM and the addition of <sup>125</sup>I-OCIF, in the presence of a 400-fold more unlabeled OCIF

[0147] FIG. 20 shows the result of crosslinking of human OBM, which is a protein encoded by a cDNA of the present invention, with <sup>125</sup>I-OCIF (monomer-type).

[0148] <Explanation of Symbols>

[0149] Lane 1: <sup>125</sup>I-OCIF

[0150] Lane 2: The crosslinked products of <sup>125</sup>I-OCIF with the proteins on the membrane of COS-7 cells transfected with phOBM.

[0151] Lane 3: The crosslinked products of <sup>125</sup>I-OCIF with the proteins on the membrane of COS-7 cells transfected with phOBM, in the presence of a 400-fold more unlabeled OCIF.

[0152] FIG. 21 shows the OCIF-binding capability of the protein (secreted-form hOBM) in the conditioned medium at various OCIF concentrations in Example 23(2).

[0153] <Explanation of Symbols>

[0154] ○: Conditioned medium of 293-EBNA cells transfected with pCEP4, which does not contain cDNA encoding secreted-form human OBM

[0155] ●: Conditioned medium of 293-EBNA cells transfected with pCEPshOBM, which contains cDNA encoding secreted-form human OBM

[0156] FIG. 22 shows the OCIF-binding capability of the protein (secreted-form human OBM) in the conditioned medium at a specific OCIF concentration while changing the amount of conditioned medium added in Example 23(2).

[0157] <Explanation of Symbols>

[0158] ○: Conditioned medium of 293-EBNA cells transfected with pCEP4, which does not contain cDNA encoding secreted-form human OBM

[0159] ●: Conditioned medium of 293-EBNA cells transfected with pCEPshOBM, which contains cDNA encoding secreted-form human OBM

[0160] FIG. 23 shows the result of SDS-PAGE of a fusion protein consisting of thioredoxin and human OBM expressed in *Escherichia coli*.

[0161] <Explanation of Symbols>

[0162] Lane 1: Molecular weight markers

[0163] Lane 2: Soluble protein fractions originating from *Escherichia coli* GI724/pTrxFus

[0164] Lane 3: Soluble protein fractions originating from *Escherichia coli* GI724/pTrxhOBM

[0165] FIG. 24 shows the OCIF-binding capability of the fusion protein consisting of thioredoxin and human OBM to OCIF, when the amount of the soluble protein fraction originating from *Escherichia coli* including the fusion protein added was varied in Example 24(3).

[0166] <Explanation of Symbols>

[0167] ○: Soluble protein fractions originating from *Escherichia coli* GI724/pTrxFus

[0168] ●: Soluble protein fractions originating from *Escherichia coli* GI724/pTrxshOBM

[0169] FIG. 25 shows the OCIF-binding capability of the fusion protein of thioredoxin and human OBM in soluble protein fractions originating from *Escherichia coli* to OCIF in various concentrations in Example 24(3).

[0170] <Explanation of Symbols>

[0171] ○: Soluble protein fractions originating from *Escherichia coli* GI724/pTrxFus

[0172] ●: Soluble protein fractions originating from *Escherichia coli* GI724/pTrxshOBM

[0173] FIG. 26 shows the result of quantifying human OBM and human sOBM by the sandwich ELISA method using the rabbit anti-human OBM/sOBM polyclonal antibody of the present invention.

[0174] <Explanation of Symbols>

[0175] □: Human OBM

[0176] ●: Human sOBM

[0177] FIG. 27 shows the result of quantifying human OBM and human sOBM by the sandwich ELISA method using the anti-human OBM/sOBM monoclonal antibodies of the present invention.

[0178] <Explanation of Symbols>

[0179] □: Human OBM

[0180] ●: Human sOBM

[0181] FIG. 28 shows the result of quantifying mouse OBM and sOBM by the sandwich ELISA method using the anti-human OBM/sOBM monoclonal antibodies of the present invention which cross react mouse OBM and sOBM.

[0182] <Explanation of Symbols>

[0183] □: Mouse OBM

[0184] ●: Mouse sOBM

[0185] FIG. 29 shows the activity of the fusion protein consisting of thioredoxin and mouse OBM to stimulate human osteoclast-like cell formation.

[0186] FIG. 30 shows the suppression of the anti-OBM/sOBM antibody of the bone resorption activity stimulated by vitamin D<sub>3</sub>.

[0187] FIG. 31 shows the suppression of the anti-OBM/sOBM antibody of the bone resorption activity stimulated by prostaglandin E<sub>2</sub> (PGE<sub>2</sub>).

[0188] FIG. 32 shows the suppression by the anti-OBM/sOBM antibody of the bone-resorbing activity stimulated by parathyroid hormone (PTH).

[0189] FIG. 33 shows the suppression by the anti-OBM/sOBM antibody of the bone-resorbing activity stimulated by interleukin 1  $\alpha$  (IL-1)

#### BEST MODE FOR CARRYING OUT THE INVENTION

##### Examples

[0190] The present invention will be described in more detail by way of examples which are given for the purpose

of illustration of the invention and are not limiting thereof in any way of the remainder of the disclosure.

##### Example 1

[0191] Preparation of the Protein of the Present Invention

[0192] (1) Large-Scale Cultivation of ST2 Cells

[0193] Mouse osteoblastic stromal cell line ST2 (RIKEN CELL BANK RCB0224) was cultured using a-MEM containing 10% fetal bovine serum. ST2 cells cultured to confluence in a 225 cm<sup>2</sup> T flask for adherent-cell culture were treated with trypsin and harvested from the T flask. After washing, the cells were transferred to five 225 cm<sup>2</sup> T flasks. After the addition of 60 ml of a-MEM containing 10<sup>-8</sup> M active-form vitamin D<sub>3</sub> (Calcitriol), 10<sup>-7</sup> M dexamethasone, and 10% fetal bovine serum, cells in each flask were cultured for 7-10 days in a CO<sub>2</sub> incubator. The cultured ST2 cells were harvested using a cell scraper and stored at -80° C. until use.

[0194] (2) Preparation of Membrane Fraction and Solubilization of Membrane-Bound Proteins

[0195] To the ST2 cells (volume, about 12 ml) described in Example 1(1), cultured using eighty 225 cm<sup>2</sup> T flasks, was added three times the volume (36 ml) of 10 mM Tris-HCl buffer (pH 7.2) containing protease inhibitors (2 mM APMSFP, 2 mM EDTA, 2 mM o-phenanthroline, 1 mM leupeptin, 1  $\mu$ g/ml pepstatin A, and 100 unit/ml aprotinin). After vigorously agitating for 30 seconds using a vortex mixer, the cells were allowed to stand for 10 minutes on ice. The cells were homogenized using a homogenizer (DOUNCE TISSUE GRINDER, A syringe, WHEATON SCIENTIFIC Co.). The same volume (48 ml) of 10 mM Tris-HCl buffer (pH 7.2) containing the above-mentioned protease inhibitors, 0.5 M sucrose, 0.1 M potassium chloride, 10 mM magnesium chloride, and 2 mM calcium chloride was added to the homogenized cells. After stirring, the mixture was centrifuged at 600xg for 10 minutes at 4° C., thereby separating nuclei and non-homogenized cells as precipitate. The supernatant obtained by the centrifuge was centrifuged at 150,000xg for 90 minutes at 4° C., to obtain membrane fraction of the ST2 cells as precipitate. Eight ml of 10 mM Tris-HCl buffer (pH 7.2) containing the above-mentioned protease inhibitors, 150 mM sodium chloride, and 0.1 M sucrose was added to this membrane fraction. After the addition of 200  $\mu$ l of 20% CHAPS (3-[3-cholamidopropyl]-dimethylammonio]-1-propanesulfonate, Sigma Co.), the mixture was stirred for 2 hours at 4° C. The mixture was then centrifuged at 150,000xg for 60 minutes at 4° C., to obtain supernatant as solubilized membrane fraction.

##### Example 2

[0196] Purification of the Protein of the Present Invention

[0197] (1) Preparation of OCIF-Immobilized Affinity Column

[0198] After replacing iso-propanol in a HiTrap NHS-activated column (1 ml, manufactured by Pharmacia Co.) with 1 mM hydrochloric acid, 1 ml of 0.2 M NaHCO<sub>3</sub>/0.5 M NaCl solution (pH 8.3) containing 13.0 mg of recombinant OCIF prepared by the method of WO 96/26217 was added to the column using a syringe (5 ml, manufactured by Terumo Corp.), to effect coupling reaction at room tempera-

ture for 30 minutes. The column was fed with 3 ml of 0.5 M ethanolamine/0.5 M NaCl (pH 8.3) and 3 ml of 0.1 M acetic acid/0.5 M NaCl (pH 4.0) three times in turn to inactivate excess active groups, then the solution was replaced with 0.5 M ethanolamine/0.5 M NaCl (pH 8.3). After allowing to stand at room temperature for 1 hour, the column was washed twice alternately with 0.5 M ethanolamine/0.5 M NaCl (pH 8.3) and 0.1 M acetic acid/0.5 M NaCl (pH 4.0), and the solution was then replaced with 50 mM Tris/1 M NaCl/0.1% CHAPS buffer (pH 7.5).

**[0199]** (2) Purification of the Protein of the Present Invention Using OCIF-Immobilized Affinity Column

**[0200]** The purification of the OCIF-binding protein was carried out at 4°C, unless otherwise indicated. The above-mentioned OCIF-immobilized affinity column was equilibrated with 10 mM Tris-hydrochloride buffer (pH 7.2) to which the protease inhibitors described in Example 1(2), 0.15 M sodium chloride, and 0.5% CHAPS were added. About 8 ml of the solubilized membrane fraction described in Example 1(2) was applied to the column at a flow rate of 0.01 ml/minute. Then, the column was washed with 10 mM Tris-hydrochloride buffer (pH 7.2) to which the above-mentioned protease inhibitors, 0.15 M sodium chloride, and 0.5% CHAPS was added, for 100 minutes at a flow rate of 0.5 ml/minute. Next, the proteins adsorbed to the column was eluted with 0.1 M glycine-hydrochloride buffer (pH 3.3) containing the protease inhibitors, 0.2 M sodium chloride, and 0.5% CHAPS for 50 minutes at a flow rate of 0.1 ml/minute. In the same manner, the proteins adsorbed to the column was eluted with 0.1 M sodium citrate buffer (pH 2.0) containing the protease inhibitors, 0.2 M sodium chloride, and 0.5% CHAPS for 50 minutes at a flow rate of 0.1 ml/minute. The eluate was collected in 0.5 ml fractions. Each fraction was immediately neutralized by the addition of 2M Tris solution. The fractions derived from the elution with these buffers (each fraction consisting of 1.0-5.0 ml of eluate) were concentrated to 50-100 µl using Centricon-10 (manufactured by Amicon of U.S.A.). OCIF was added to a portion of each concentrated fraction and immunoprecipitated with anti-OCIF polyclonal antibody. The precipitated fractions were treated with SDS and subjected to SDS-PAGE. Fractions (Fr. No. 3-10) in which the band of the protein with specific binding ability to OCIF appeared were regarded as the protein fractions of the present invention.

**[0201]** (3) Purification of the Protein of the Present Invention by Gel Filtration

**[0202]** The concentrated OCIF-binding protein (the fractions obtain by the elution with 0.1 M glycine-hydrochloride buffer (pH 3.3) and 0.1 M sodium citrate buffer (pH 2.0)) prepared in Example 2(2) was applied to a Superose 12 HR10/30 column (1.0×30 cm, manufactured by Pharmacia Co.) which was equilibrated with 10 mM Tris-HCl, 0.5 M NaCl, 0.5% CHAPS (pH 7.0) and developed with the equilibration buffer as a mobile phase at a flow rate of 0.5 ml/min, and each 0.5 ml fraction was collected. The fractions containing the protein of the present invention (Fr. Nos. 27-32) were identified according to the same method as described above. Each of the fractions was concentrated using Centricon-10 (a product of Amicon).

**[0203]** (4) Purification by Reverse Phase High Performance Liquid Chromatography

**[0204]** The above-mentioned OCIF-binding protein purified by the gel filtration was applied to a C<sub>4</sub> column (2.1×250

mm, Vydac, USA) which was equilibrated with 0.1% trifluoroacetic acid (TFA) and 30% acetonitrile. The proteins bound to the column were eluted with linear gradients of acetonitrile from 30% to 55% for the first 50 minutes and from 55% to 80% during the next 10 minutes at a flow rate of 0.2 ml/min. Peaks of eluted proteins were detected by measuring optical density at 215 nm. Proteins in the different peaks were analyzed to identify the fractions containing the protein of the present invention, and a highly purified protein of the present invention was obtained.

### Example 3

**[0205]** SDS-PAGE of the Purified Protein of the Present Invention

**[0206]** The solubilized membrane fraction prepared from ST2 cells which were cultured in the presence or absence of active-form vitamin D<sub>3</sub> was subjected to purification with the OCIF-immobilized affinity column. The purified preparations were subjected to SDS-PAGE. As shown in FIG. 1(A), a major protein band with MW of about 30,000-40,000 was detected only in the purified preparation from ST2 cells which was cultured in the presence of the active-form vitamin D<sub>3</sub>, proving that the protein which specifically binds to OCIF (i.e. the protein of the present invention) can be selectively purified by the OCIF-immobilized affinity column. However, bands of several proteins (other than the protein of the present invention) which non-specifically bind to carriers or spacers of the OCIF-immobilized column were detected in both of the purified preparations. The proteins other than the protein of the present invention were removed according to the above-described method by gel filtration and C4 reverse phase chromatography. SDS-PAGE of the obtained highly purified protein of the present invention is shown in FIG. 1(B). The highly purified protein of the present invention was found to be electrophoretically homogeneous and had a molecular weight of about 30,000-40,000.

### Example 4

**[0207]** Binding Test of OCIF to Osteoblasts

**[0208]** (1) Preparation of <sup>125</sup>I-Labeled OCIF

**[0209]** OCIF was labeled with <sup>125</sup>I by the Iodogen method. Specifically, 20 µl of 2.5 mg/ml Iodogen-chloroform solution was transferred to a 1.5 ml Eppendorf tube and chloroform was evaporated off at 40°C, to obtain a tube coated with Iodogen. The tube was washed three times with 400 µl of 0.5 M sodium phosphate buffer (Na-Pi, pH 7.0). Five µl of 0.5 M Na-Pi (pH 7.0) was added to the tube. Immediately after the addition of 1.3 µl (18.5 MBq) of Na-<sup>125</sup>I solution (NEZ-033H2O, manufactured by Amersham Co.), 10 µl of 1 mg/ml rOCIF solution (monomer type or dimer type) was added to the tube. After mixing with a vortex mixer, the mixture was allowed to stand at room temperature for 30 seconds. The solution was transferred to a tube containing 80 µl of a solution of 10 mg/ml potassium iodide in 0.5 M Na-Pi (pH 7.0) and 5 µl of a phosphate buffered saline containing 5% bovine serum albumin, and stirred. The mixture was applied to a spin column (1 ml, G-25 fine, manufactured by Pharmacia Co.) which was equilibrated with phosphate buffered saline containing 0.25% bovine serum albumin and the column was centrifuged for 5 minutes at 2,000 rpm. Four hundred µl of a

phosphate buffered saline containing 0.25% bovine serum albumin was added to the fraction eluted from the column and the mixture was stirred. A 2  $\mu$ l of the aliquot was removed to measure the radioactivity using a gamma counter. The radiochemical purity of the  $^{125}$ I-labeled OCIF was determined by measuring the radioactivity precipitated with 10% TCA. The biological activity of the  $^{125}$ I-labeled OCIF was measured according to the method described in WO 96/26217. The concentration of the  $^{125}$ I-labeled OCIF was measured by the ELISA according to the following procedure.

[0210] (2) Measurement of the Concentration of  $^{125}$ I-Labeled OCIF by ELISA

[0211] One hundred  $\mu$ l of 50 mM NaHCO<sub>3</sub> (pH 9.6) in which the anti-OCIF rabbit polyclonal antibody described in WO 96/26217 was dissolved to a concentration of 2  $\mu$ g/ml was added to each well of a 96-well immuno-plate (MaxiSorp<sup>TM</sup>, a product of Nunc Co.). The plate was allowed to stand overnight at 4° C. After removing the solution by suction, 300  $\mu$ l of Block Ace<sup>TM</sup> (Snow Brand Milk Products Co., Ltd.)/phosphate buffered saline (25/75) solution was added to each well. The plate was then allowed to stand for two hours at room temperature. After removing the solution by suction, the wells were washed three times with phosphate buffered saline containing 0.01% Polysorbate 80 (P-PBS). Next, 300  $\mu$ l of Block Ace<sup>TM</sup>/phosphate buffered saline (25/75) solution to which  $^{125}$ I-labeled OCIF or the standard OCIF preparation was mixed, was added to each well. The plate was then allowed to stand for two hours at room temperature. After removing the solution by suction, each well was washed six times with 200  $\mu$ l of P-PBS.

[0212] One hundred  $\mu$ l of Block Ace<sup>TM</sup> (Snow Brand Milk Products Co., Ltd.)/phosphate buffered saline (25/75) solution containing peroxidase labeled rabbit anti-OCIF polyclonal antibody was added to each well. The plate was allowed to stand for two hours at room temperature. After removing the solution by suction, the wells were washed six times with 200  $\mu$ l P-PBS. Then, 100  $\mu$ l of TMB solution (TMB Soluble Reagent, High Sensitivity, Scytek Co.) was added to each well. After incubating at room temperature for 2-3 minutes, 100  $\mu$ l of stopping solution (Stopping Reagent, Scytek Co.) was added to each well. Absorbance of each well was measured at 490 nm using a microplate reader. The concentration of  $^{125}$ I-labeled OCIF was determined from a calibration curve prepared using the standard preparation of OCIF.

[0213] (3) Binding Test of OCIF to Osteoblasts or Spleen Cells

[0214] Mouse osteoblastic stromal cell line ST2 or spleen cells were suspended in  $\alpha$ -MEM containing 10% fetal bovine serum (FBS), either with or without 10<sup>-8</sup> M active-form vitamin D<sub>3</sub> (Calcitriol) and 10<sup>-7</sup> M dexamethasone, to a concentration of 4 $\times$ 10<sup>4</sup> cells/ml (ST2 cells) or 2 $\times$ 10<sup>6</sup> cells/ml (spleen cells), respectively. Each cell suspension was innoculated into a 24-well micro plate. The cells were cultured for 4 days in a CO<sub>2</sub> incubator. After washing the cells with  $\alpha$ -MEM, 200  $\mu$ l of medium for the binding test ( $\alpha$ -MEM to which 0.2% bovine serum albumin, 20 mM Hepes buffer, and 0.2% Na<sub>3</sub> were supplemented), containing 20 ng/ml of above-described  $^{125}$ I-labeled OCIF (monomer form or dimer form), was added to each well. To the wells for the measurement of non-specific binding, 200  $\mu$ l of

the medium for the binding test containing 8  $\mu$ g/ml of rOCIF (400 times concentration) in addition to 20 ng/ml of  $^{125}$ I-labeled OCIF was added. The cells were cultured for one hour in a CO<sub>2</sub> incubator and washed 3 times with 1 ml of a phosphate buffered saline. In this procedure, spleen cells were washed by centrifuging the 24-well plate in each washing step, because the spleen cells were non-adherent. After washing, 500  $\mu$ l of 0.1 N NaOH solution was added to each well and the plate was allowed to stand for 10 minutes at room temperature to dissolve the cells. The amount of RI in each well was measured by a gamma counter.

[0215] As shown in FIG. 2,  $^{125}$ I-labeled OCIF did not bind to the cultured spleen cells, but specifically bound only to the osteoblastic stromal cells which were cultured in the presence of active-form vitamin D<sub>3</sub>. The results indicated that the protein of the present invention is a membrane bound protein induced by active-form vitamin D<sub>3</sub> and dexamethasone on osteoblastic stromal cells.

#### Example 5

[0216] Biological Activity of the Protein of the Present Invention

[0217] (1) Osteoclasts-Formation Supported by Osteoblastic Stromal Cells

[0218] The osteoclasts formation-supporting capability of osteoblastic stromal cells was evaluated by measuring tartric acid resistant acid phosphatase activity (TRAP activity) of the formed osteoclasts. Specifically, spleen cells (2 $\times$ 10<sup>5</sup> cells/100  $\mu$ l/well) from a ddY mouse (8-12 weeks old) and mouse osteoblastic stromal cells ST2 (5 $\times$ 10<sup>3</sup> cells/100  $\mu$ l/well) were suspended in  $\alpha$ -MEM to which 10<sup>-8</sup> M active-form vitamin D<sub>3</sub>, 10<sup>-7</sup> M dexamethasone, and 10% fetal bovine serum were added. The cells were innoculated into 96-well plates and cultured for one week in a CO<sub>2</sub> incubator. After washing each well with phosphate buffered saline, 100  $\mu$ l of ethanol/acetone (1:1) was added to each well, and the cells were immobilized at room temperature for one minute. After immobilization, 100  $\mu$ l of 50 mM citrate buffer (pH 4.5) containing 5.5 mM p-nitrophenol phosphate and 10 mM sodium tartarate was added to each well. After 15 minutes of reaction at room temperature, 0.1 N NaOH solution was added to each well and absorbance at 405 nm was measured using a microplate reader. The results of osteoclasts-formation by ST2 cells with a passage number of about 10 or 40 after purchasing the cells from RIKEN CELL BANK are shown in FIG. 3. The results indicate that the ST2 cells with a higher passage number exhibit more potent ability to support osteoclasts-formation.

[0219] (2) Time Course Change of Expression of the Protein of the Present Invention on Membrane of Osteoblastic Stromal Cells in a Culture System Which Include Active-Form Vitamin D<sub>3</sub> and Dexamethasone and Time Course Change of Osteoclasts-Formation in the Co-Culture System

[0220] In the same manner as in Example 4(3), osteoblastic stromal cell ST2 was cultured for 7 days in the presence of active-form vitamin D<sub>3</sub> and dexamethasone. The OCIF-binding test was carried out using  $^{125}$ I-labeled OCIF (monomer type) as described in the experiment in Example 4(1). Non-specific binding was measured by competing  $^{125}$ I-OCIF binding to ST2 cells with 400-fold concentration of

unlabeled OCIF. As a result, it was confirmed that the amount of specific binding of  $^{125}\text{I}$ -labeled OCIF increase in accordance with increase in culture period in the presence of active-form vitamin D<sub>3</sub> and dexamethasone. Specifically, as shown in **FIGS. 4 and 5**, the protein of the present invention was expressed on the surface of ST2 cells by active-form vitamin D<sub>3</sub> in accordance with increase in culture period and the expression reached a maximum on the fourth day of culture. On the other hand, osteoclast-like cells are formed by coculturing mouse spleen cells and ST2 cells in the presence of active-form vitamin D<sub>3</sub>. TRAP (a marker enzyme of osteoclasts)-positive mononuclear pre-osteoclast-like cells are formed on the third or fourth day of the culture. More differentiated and mature TRAP-positive multinuclear cells are formed on the fifth to sixth day of the culture. A good correlation between time-course of the expression of the protein of the present invention and osteoclasts-formation was thus demonstrated.

**[0221]** (3) Inhibition of Osteoclasts Formation by OCIF Treatment for Different Period During the Co-Culture

**[0222]** To make it clear that the protein of the present invention is a factor involved in the osteoclasts-formation, the cells were treated with 100 mg/ml OCIF for different culture periods during the six day co-culture period described in the above-mentioned Example 5(2) (two consecutive days in the six-day period, except for the 5th day for which a one-day period was applied). As a result, as shown in **FIG. 6**, OCIF treatment at 48-96 hours after start of the culture at which expression of the protein of the present invention on ST2 cells is maximal was found to be most effective for inhibiting formation of osteoclasts. Specifically, it was confirmed that OCIF controls osteoclast formation by binding to ST2 cells via the protein of the present invention.

**[0223]** Based on the results of the above experiments, the protein of the present invention was confirmed to be induced on cell membrane of osteoblastic stromal cells by active-form vitamin D<sub>3</sub> and dexamethasone and to exhibit a biological activity to support or accelerate differentiation or maturation of osteoclasts.

#### Example 6

**[0224]** Crosslinking Test for  $^{125}\text{I}$ -Labeled OCIF and the Protein of the Present Invention

**[0225]** To identify the protein of the present invention more clearly, the protein of the present invention was crosslinked with  $^{125}\text{I}$ -labeled OCIF. Mouse osteoblastic stromal cell line ST2 was cultured for four days in the presence or absence of active-form vitamin D<sub>3</sub> and dexamethasone in the same manner as described in Example 4(3). After washing the cells with 1 ml of phosphate buffered saline, 200  $\mu\text{l}$  of medium for binding test ( $\alpha$ -MEM to which 0.2% bovine serum albumin, 20 mM Hepes buffer, and 0.2% Na<sub>3</sub>N) and the culture medium was serially diluted at a rate to 1/2 with the culture medium not containing  $^{125}\text{I}$ -labeled OCIF. Solutions for measuring non-specific binding were prepared by further adding 400-fold concentration of monomer-form OCIF to these solutions. Two hundred  $\mu\text{l}$  of the prepared solutions were added to the above-mentioned-wells with ST2 cells cultured for 4 days (passage number, about 10) in the presence of  $10^{-8}$  M active-form vitamin D<sub>3</sub> (Calcitriol) and  $10^{-7}$  M dexamethasone, to assess binding of  $^{125}\text{I}$ -labeled OCIF in the same method as described in Example 4(3). The results were subjected to Scatchard Plot analysis to determine the dissociation constant of OCIF and OCIF-binding protein and the number (site) of OCIF-binding protein per a ST2 cell. As a result, the dissociation constant of OCIF and the protein of the present invention was found to be 280 pM, and the number of the site of OCIF-binding protein per a ST2 cell was approximately 33,000/cell. Based on the finding in Example 5(1) that osteoclasts-formation supported by the ST2 cells with passage number about 40 was more extensive than that with passage number about 10, the number (the site) of the protein of the present invention expressed on the ST2 cell with a passage number about 40 was assessed. The number (site) was 58,000/cell and was clearly larger than the ST2 cells with passage number about 10, indicating that the amount of the protein of the present invention expressed on ST2 cells is related to their potency to support osteoclasts-formation. The results indicated that the protein of the present invention is a factor that supports or induces differentiation or maturation of osteoclasts.

arin. Five hundred  $\mu\text{l}$  of phosphate buffered saline containing 100  $\mu\text{g}/\text{ml}$  crosslinking agent, DSS (Disuccinimidyl suberate, Pierce Co.), was added to each well and the plate was kept for 10 minutes at 0° C. The wells were washed twice with 1 ml of phosphate buffered saline at 0° C. One hundred  $\mu\text{l}$  of 20 mM Hepes buffer containing 1% Triton X-100, 10  $\mu\text{M}$  pepstatin, 10  $\mu\text{M}$  leupeptin, 2 mM PMSF (phenylmethylsulfonyl fluoride), 10  $\mu\text{M}$  antipain, and 2 mM EDTA, was then added to each well. The plate was allowed to stand for 30 minutes at room temperature to dissolve the cells. Fifteen  $\mu\text{l}$  of these samples were treated with SDS under non-reducing conditions according to conventional method and subjected to SDS-polyacrylamide gel electrophoresis(4-20% polyacrylamide gradient, manufactured by Daiichi Chemical Co., Ltd.). After electrophoresis, the gels were dried and exposed to BioMax MS film (manufactured by Kodak) for 24 hours at -80° C. using BioMax MS intensifying screens (manufactured by Kodak). After exposure, the film was developed by conventional method. A band of crosslinking product with a molecular weight of 90,000-110,000 was detected when the  $^{125}\text{I}$ -labeled OCIF (monomer type, 60 kDa) was used. When the  $^{125}\text{I}$ -labeled OCIF-CDD1 (31 kDa) was used, a band of crosslinking product of about 70-80 kDa (average, 78 kDa) was detected as shown in **FIG. 7**.

#### Example 7

**[0226]** Analysis of the Protein of the Present Invention Expressed on ST Cells by Scatchard Plot

**[0227]** The above-mentioned  $^{125}\text{I}$ -labeled OCIF (monomer type) was added to a concentration of 1,000 pM to the culture medium for binding test ( $\alpha$ -MEM containing 0.2% bovine serum albumin, 20 mM Hepes buffer, and 0.2% Na<sub>3</sub>N) and the culture medium was serially diluted at a rate to 1/2 with the culture medium not containing  $^{125}\text{I}$ -labeled OCIF. Solutions for measuring non-specific binding were prepared by further adding 400-fold concentration of monomer-form OCIF to these solutions. Two hundred  $\mu\text{l}$  of the prepared solutions were added to the above-mentioned-wells with ST2 cells cultured for 4 days (passage number, about 10) in the presence of  $10^{-8}$  M active-form vitamin D<sub>3</sub> (Calcitriol) and  $10^{-7}$  M dexamethasone, to assess binding of  $^{125}\text{I}$ -labeled OCIF in the same method as described in Example 4(3). The results were subjected to Scatchard Plot analysis to determine the dissociation constant of OCIF and OCIF-binding protein and the number (site) of OCIF-binding protein per a ST2 cell. As a result, the dissociation constant of OCIF and the protein of the present invention was found to be 280 pM, and the number of the site of OCIF-binding protein per a ST2 cell was approximately 33,000/cell. Based on the finding in Example 5(1) that osteoclasts-formation supported by the ST2 cells with passage number about 40 was more extensive than that with passage number about 10, the number (the site) of the protein of the present invention expressed on the ST2 cell with a passage number about 40 was assessed. The number (site) was 58,000/cell and was clearly larger than the ST2 cells with passage number about 10, indicating that the amount of the protein of the present invention expressed on ST2 cells is related to their potency to support osteoclasts-formation. The results indicated that the protein of the present invention is a factor that supports or induces differentiation or maturation of osteoclasts.

## Example 8

[0228] Cloning of OBMcDNA

[0229] (1) Extraction of RNA from Mouse ST2 Cells

[0230] Mouse osteoblastic stromal cell line ST2 (RIKEN CELL BANK, RCB0224) was cultured in  $\alpha$ -MEM (Gibco BRL Co.) containing 10% fetal bovine serum. ST2 cells cultured to confluent in a 225 cm<sup>2</sup> T-flask for adherent cells were treated with trypsin to harvest the cells from the T-flask. The cells were washed and transferred to five 225 cm<sup>2</sup> T-flasks. Sixty ml of  $\alpha$ -MEM containing 10<sup>-8</sup> M active-form vitamin D<sub>3</sub> (Calcitriol, Wako Pure Chemicals Co., Ltd.), 10<sup>-7</sup> M dexamethasone, and 10% fetal bovine serum was added to each flask and the cells were cultured for 5 days in a CO<sub>2</sub> incubator. Total RNA was extracted from the cultured ST2 cells using ISOGEN (Wako Pure Chemicals Co., Ltd.) Poly A<sup>+</sup>RNA was prepared from about 600  $\mu$ g of the total RNA using an Oligo(dT)-cellulose column (5'-3' Prime Co.). About 8  $\mu$ g of Poly A<sup>+</sup>RNA was obtained.

[0231] (2) Construction of Expression Library

[0232] Double-stranded cDNA was synthesized from 2  $\mu$ g of polyA<sup>+</sup>RNA obtained in Example 8(1) using a Great Lengths cDNA Synthesis kit (Clontech Co.) according to the instruction in the manual. Specifically, 2  $\mu$ g of polyA<sup>+</sup>RNA and Oligo(dT)<sub>25</sub>(dN) primer were mixed and distilled water was added to the mixture to make the final volume to 6.25  $\mu$ l. After incubation for about 3 minutes at 70° C., the mixture was cooled on ice for 2 minutes. To the mixture were added 2.2  $\mu$ l of distilled water, 2.5  $\mu$ l of 5x First-strand buffer, 0.25  $\mu$ l of 100 mM DTT (dithiothreitol), 0.5  $\mu$ l of PRIME RNase inhibitor (1 U/ml) (5'-3' Prime Co.), 0.5  $\mu$ l of [ $\alpha$ -<sup>32</sup>P]dCTP (Amersham Co., 3000 Ci/mmol) diluted 5-fold with distilled water to make 2  $\mu$ Ci/ $\mu$ l, 0.65  $\mu$ l of dNTP (20 mM each), and 1.25  $\mu$ l (250 unit) of MMLV(RNaseH<sup>-</sup>) reverse transcriptase. The mixture was incubated for 90 minutes at 42° C., followed by the further addition of 62.25  $\mu$ l of distilled water, 20  $\mu$ l of 5x second-strand buffer, 0.75  $\mu$ l of dNTP (20 mM each), and 5  $\mu$ l of Second-strand enzyme cocktail. The resulting mixture was maintained at 16° C. for two hours. Then, 7.5 units of T4DNA polymerase was added to this reaction mixture. After incubation at 16° C. for 30 minutes, the reaction was terminated by the addition of 5  $\mu$ l of 0.2M EDTA. After a phenol-chloroform treatment, the product was precipitated with ethanol. An EcoRI-SalI-NotI linker (Clontech Co.) was attached to the ends of the resultant double-stranded cDNA. Then, the ends were phosphorylated and the product was applied on a size fractionation column to obtain cDNA with a length more than 500 bp. DNA was precipitated with ethanol, dissolved in water and ligated to pcDL-SR  $\alpha$ 296 (Molecular and Cellular Biology, Vol. 8, pp 466-472, 1988) which had been cut with a restriction enzyme EcoRI (Takara Shuzo Co.) and treated with CIAP (calf intestine alkaline phosphatase, Takara Shuzo Co.).

[0233] (3) Screening of Expression Library by Means of Binding to OCIF

[0234] An *escherichia coli* strain, XL2 Blue MRF' (Toyobo Co., Ltd.), was transformed using the DNA produced in Example 8(2) and cultured on L-Carbenisilin agar (1% tryptan, 0.5% yeast extract, 1% NaCl, 60  $\mu$ g/ml carbenisilin, 1.5% agar) prepared in a 24-well plastic plates, to produce about 100 colonies per well. Transformants in each

well were suspended in 3 ml of Terrific Broth ampicillin culture medium (1.2% tryptan, 2.4% yeast extract, 0.4% glycerol, 0.017 M KH<sub>2</sub>PO<sub>4</sub>, 0.072 M K<sub>2</sub>HPO<sub>4</sub>, 100  $\mu$ g/ml ampicillin) and cultured at 37° C. overnight with shaking. Cells were collected by centrifugation to prepare plasmid DNA using a QIAwell kit (QIAGEN Co.). DNA concentration was determined by measuring absorbance at 260 nm. DNA was concentrated by precipitating with ethanol and dissolved in distilled water to a concentration of 200 ng/ $\mu$ l. Five hundred DNA pools, each of which was obtained from about 100 colonies were prepared and were used for transfection into COS-7 cells (RIKEN CELL BANK, RCB0539). COS-7 cells were seeded into DMEM containing 10% fetal bovine serum in each well of 24-well plates at a cell density of 8 $\times$ 10<sup>4</sup> cells/well and cultured overnight at 37° C. in a CO<sub>2</sub> incubator. Next day, the culture medium was removed and the cells were washed with serum-free DMEM culture medium. The above-described plasmid DNA which was previously diluted with an OPTI-MEM culture medium (Gibco BRL Co.) and mixed with Lipofectamine (a transfection reagent, manufactured by Gibco BRL Co.) according to the protocol supplied with Lipofectamine. After 15 minutes, the mixture was added to the cells in each well. The amount of Lipofectamine and DNA used were, respectively, 1  $\mu$ g and 4  $\mu$ l per well. After 5 hours, the culture medium was removed and 1 ml of DMEM culture medium (Gibco BRL Co.) containing 10% fetal bovine serum was added to each well. The plates were incubated for 2-3 days at 37° C. in a CO<sub>2</sub> incubator(5% CO<sub>2</sub>). The COS-7 cells transfected and cultured for 2-3 days in this manner were washed with a serum-free DMEM culture medium. Then, 200  $\mu$ l of a culture medium for the binding assay (serum-free DMEM culture medium containing 0.2% calf serum albumin, 20 mM Hepes buffer, 0.1 mg/ml heparin, and 0.02% NaN<sub>3</sub>) with 20 ng/ml of <sup>125</sup>I-labeled OCIF added thereto was added to each well. After culturing for one hour at 37° C. in a CO<sub>2</sub> incubator (5% CO<sub>2</sub>), the cells were washed twice with 500  $\mu$ l of a phosphate buffered saline containing 0.1 mg/ml heparin. After washing, 500  $\mu$ l of 0.1 N NaOH solution was added to each well. The plates were allowed to stand for 10 minutes at room temperature to lyse the cells. The amount of <sup>125</sup>I in each well was measured using a gamma counter (Packard Co.). One DNA pool containing cDNA encoding the protein which specifically binds to OCIF was found by screening a total of 500 pools. The DNA pool containing the cDNA was further divided, and the above-described transfection and screening operations were repeated to isolate the cDNA which encodes the protein which binds to OCIF. The plasmid containing this cDNA was named pOBM291. The *Escherichia coli* containing this plasmid was deposited with The National Institute of Bioscience and Human Technology, Agency of Industrial Science and Technology, Biotechnology Laboratory, as pOBM291 on May 23, 1997 under the deposition No. FERM BP-5953.

[0235] The methods of labeling OCIF with <sup>125</sup>I and quantitative analysis of the <sup>125</sup>I-labeled OCIF by ELISA are shown below. Labeling of OCIF with <sup>125</sup>I was carried out according to the Iodogen method. Twenty  $\mu$ l of 25 mg/ml Iodogen-chloroform solution was added to a 1.5 ml Eppendorf tube and chloroform was evaporated by heating at 40° C., to prepare an Iodogen-coated tube. The tube was washed three times with 400  $\mu$ l of 0.5 M sodium phosphate buffer (Na-Pi, pH 7.0), and 5  $\mu$ l of 0.5 M Na-Pi (pH 7.0) was added. Immediately after the addition of 1.3  $\mu$ l (18.5 MBq) of

Na-<sup>125</sup>I solution (NEZ-033H20, Amersham Co.), 10  $\mu$ l of 1 mg/ml rOCIF solution (monomer type or dimer type) was added to the tube. After mixing the contents with a vortex mixer, the tube was allowed to stand at room temperature for 30 seconds. The solution in the tube was transferred to a tube to which 80  $\mu$ l of 10 mg/ml potassium iodide, 0.5 M Na-Pi (pH 7.0) and 5  $\mu$ l of a phosphate buffered saline containing 5% bovine serum albumin (BSA-PBS) were previously added. After stirring, the mixture was applied to a spin column (1 ml, G-25 fine, manufactured by Pharmacia Co.) equilibrated with BSA-PBS, and the column was centrifuging for 5 minutes at 2000 rpm. Four hundred  $\mu$ l of BSA-PBS was added to the fraction eluted from the column. After stirring, 2  $\mu$ l of an aliquot of this solution was sampled to measure the radioactivity by a gamma counter. The radiochemical purity of the <sup>125</sup>I-labeled OCIF solution thus prepared was determined by measuring radioactivity precipitated by 10% TCA. The biological activity of the <sup>125</sup>I-labeled OCIF was measured according to the method of WO 96/26217. The concentration of the <sup>125</sup>I-labeled OCIF was determined by the ELISA as follows. Specifically, 100  $\mu$ l of 50 mM NaHCO<sub>3</sub> (pH 9.6) in which the anti-OCIF rabbit polyclonal antibody described in WO 96/26217 was dissolved to a concentration of 2  $\mu$ g/ml was added to each well of a 96-well immuno-plate (MaxiSorp, a product of Nunc Co.). The plate was allowed to stand over night at 4° C. After removing the solution by suction, 300  $\mu$ l of Block Ace™ (Snow Brand Milk Products Co., Ltd.)/phosphate buffered saline (25/75) (B-PBS) was added to each well. The plate was then allowed to stand for two hours at room temperature. After removing the solution by suction, the wells were washed three times with phosphate buffered saline containing 0.01% Polysorbate 80(P-PBS). Next, 100  $\mu$ l of B-PBS containing <sup>125</sup>I-labeled OCIF or standard OCIF was added to each well. The plate was then allowed to stand for two hours at room temperature. After removing the solution by suction, each well was washed six times with 200  $\mu$ l of P-PBS. One hundred  $\mu$ l of peroxidase-labeled rabbit anti-OCIF polyclonal antibody diluted with B-PBS was added to each well. The plate was allowed to stand for two hours at room temperature. After removing the solution by suction, the wells were washed six times with 200  $\mu$ l of P-PBS. Then, 100  $\mu$ l of TMB solution (TMB Soluble Reagent, High Sensitivity, Scytek Co.) was added to each well. After incubating the plate at room temperature for 2-3 minutes, 100  $\mu$ l of stopping solution (Stopping Reagent, Scytek Co.) was added to each well. Absorbance at 450 nm of each well was measured using a microplate reader. The concentration of <sup>125</sup>I-labeled OCIF was determined based on a calibration curve drawn using the standard preparation of OCIF.

[0236] (4) Determination of the Nucleotide Sequence of the cDNA Encoding the Entire Amino Acid Sequence of OBM

[0237] The nucleotide sequence of the OBM cDNA obtained in the Example 8(3) was determined using a Taq DyeDeoxy Terminator Cycle Sequencing kit (a product of Perkin Elmer Co.). Specifically, the nucleotide sequence of the insert fragment was directly determined using pOBM291 as a template. Fragments with a length of about 1.0 kb and 0.7 kb which were obtained by digesting pOBM291 with a restriction enzyme EcoRI were inserted into the EcoRI site of plasmid pUC19 (Takara Shuzo Co.) The nucleotide sequences of these fragments were also determined. The following primers were used: primer SRR2 which was used

to determine nucleotide sequences of DNA fragments inserted into pcDL-SR α296, M13PrimerM3 and M13PrimerRV (both manufactured by Takara Shuzo Co.) which were used to determine the nucleotide sequences of DNA fragments inserted into plasmid pUC19, and synthesized primer OBM#8 designed based on the nucleotide sequence of OBMcDNA. Sequences of these primers are shown as the Sequence ID No. 3 to No 6 in the sequence table.

[0238] In addition, the nucleotide sequence of OBMcDNA determined is shown as Sequence ID No. 2 and the amino acid sequence determined therefrom is shown as the Sequence ID No. 1.

#### Example 9

[0239] Expression of the Protein Encoded by the cDNA of the Present Invention

[0240] Plasmid pOBM291 was transfected into COS-7 cells in each well of a 6-well plate using Lipofectamine and the transfected COS-7 cells were cultured for two days in DMEM containing 10% fetal bovine serum. The medium was replaced with a cysteine-methionine-free DMEM (Dainippon Seiyaku Co. Ltd.) (800  $\mu$ l/well) containing 5% dialyzed fetal bovine serum. The cells were cultured for 15 minutes, followed by the addition of 14  $\mu$ l of Express Protein Labeling Mix (10 mCi/ml, manufactured by NEN Co.). After culturing for four hours, 200  $\mu$ l of DMEM including 10% fetal bovine serum was added. After one hour culturing, the cells were washed twice with PBS. Then, 0.5 ml of a TSA buffer (10 mM Tris-HCl (pH 8.0) containing 0.14 M NaCl, 0.025% NaN<sub>3</sub>), containing 1% TritonX-100, 1% bovine hemoglobin, 10  $\mu$ g/ml leupeptin, 0.2 TIU/ml aprotinin, 1 mM PMSF, was added and the mixture was allowed to stand for one hour on ice. Cells were disrupted by pipetting and centrifuged at 3000 $\times$ g for 10 minutes at 4° C. to obtain supernatant. 200  $\mu$ l of dilution buffer (TSA buffer containing 0.1% TritonX-100, 0.1% bovine hemoglobin, 10  $\mu$ g/ml leupeptin, 0.2 TIU/ml aprotinin, 1 mM PMSF) was added to 100  $\mu$ l of this supernatant. The mixture was shaken for one hour at 4° C. with protein A Sepharose (50  $\mu$ l). The resultant mixture was centrifuged at 1500 $\times$ g for one minute at 4° C. to collect supernatant, and thereby fraction(s) which is non-specifically adsorbed to Protein A Sepharose was removed. OCIF (1  $\mu$ g) was added to this supernatant and the mixture was shaken at 4° C. for one hour to achieve the binding of OCIF to OBM. Anti-OCIF polyclonal antibody (50  $\mu$ g) was added and the mixture was shaken for one hour at 4° C. Then, Protein A Sepharose (10  $\mu$ l) was added and the mixture was shaken for an additional hour at 40° C., followed by centrifuge at 1500 $\times$ g for 1 minute at 40° C. to collect precipitate. The precipitate was washed twice with dilution buffer, twice with a bovine hemoglobin-free dilution buffer, once with TSA buffer, and once with 50 mM Tris-HCl (pH 6.5). After washing, SDS buffer (0.125 M Tris-HCl, 4% sodium dodecylsulfate, 20% glycerol, 0.002% Bromophenol Blue, pH 6.8) containing 10%  $\beta$ -mercaptoethanol was added to the precipitate. The mixture was heated for 5 minutes at 100° C. and subjected to SDS-PAGE (12.5% polyacrylamide gel, Daiichi Chemical Co., Ltd.). The gel was fixed according to a conventional method. Isotope signals were amplified using Amplify™ (Amersham Co.) and the sample was exposed to Bio Max MR film (KODAK Co.) at -80° C. The results are shown in FIG. 8, which indicates that the

protein encoded by the cDNA of the present invention has a molecular weight of about 40,000.

#### Example 10

[0241] Binding of the Protein Encoded by the cDNA of the Present Invention to OCIF

[0242] Plasmid pOBM291 was transfected into COS cells in each well of a 24-well plate using Lipofectamine. After culturing for 2-3 days, the cells were washed with serum-free DMEM culture medium. 200  $\mu$ l of culture medium for the binding assay (serum-free DMEM culture medium containing 0.2% calf serum, albumin, 20 mM Hepes, 0.1 mg/ml heparin, and 0.2% NaN<sub>3</sub>), supplemented with 20 ng/ml <sup>125</sup>I-labeled OCIF, was added to the wells. To the other wells, 200  $\mu$ l of culture medium for the binding assay to which 8  $\mu$ g/ml of unlabelled OCIF had been added, in addition to 20 ng/ml <sup>125</sup>I-labeled OCIF, was added. After culturing for one hour at 37° C. in a CO<sub>2</sub> incubator (5% CO<sub>2</sub>), the cells were washed twice with 500  $\mu$ l of phosphate buffered saline containing 0.1 mg/ml of heparin. Then, 500  $\mu$ l of 0.1 N NaOH solution was added to each well and the plate was allowed to stand for 10 minutes at room temperature to dissolve the cells. The amount of <sup>125</sup>I in each well was measured by a gamma counter. As a result, as shown in FIG. 9, the <sup>125</sup>I-labeled OCIF was found to bind only to the cells in which plasmid pOBM291 was transfected. In addition, the binding was confirmed to be conspicuously inhibited by the addition of (unlabeled) OCIF at a 400-fold concentration. These results have demonstrated that the protein OBM encoded by the cDNA on plasmid pOBM291 specifically binds to OCIF on the surface of the transfected COS-7 cells.

#### Example 11

[0243] Crosslinking of <sup>125</sup>I-Labeled OCIF and the Protein Encoded by the cDNA of the Present Invention

[0244] Crosslinking of <sup>125</sup>I-labeled monomer type OCIF and the protein encoded by the cDNA of the present invention was carried out to investigate the characteristics of the protein encoded by the cDNA of the present invention in further detail. After transfection of plasmid pOBM291 into COS-7 cells according to the method used in the Example 8(3), 200  $\mu$ l of the culture medium for the binding assay, as described above, containing <sup>125</sup>I-labeled OCIF (25 ng/ml) was added to the wells. The culture medium for the binding assay to which unlabeled OCIF at a 400-fold concentration was added in addition to <sup>125</sup>I-labeled OCIF was added to the other wells. After culturing for one hour at 37° C. in a CO<sub>2</sub> incubator (5% CO<sub>2</sub>), the cells were washed twice with 500  $\mu$ l of phosphate buffered saline containing 0.1 mg/ml heparin. Five hundred  $\mu$ l of phosphate buffered saline containing 100  $\mu$ g/ml of a crosslinking agent, DSS (disuccinimidyl suberate, manufactured by Pierce Co.) was added to the cells, followed by a reaction for 10 minutes at 0° C. The cells in these wells were washed twice with 1 ml of cold phosphate buffered saline (0° C.). After the addition of 100  $\mu$ l of 20 mM Hepes buffer containing 1% Triton X-100 (Wako Pure Chemicals Co., Ltd.), 2 mM PMSF (Phenylmethylsulfonyl fluoride, Sigma Co.), 10  $\mu$ M Pepstatin (Wako Pure Chemicals Co., Ltd.), 10  $\mu$ M Leupeptin (Wako Pure Chemicals Co., Ltd.), 10  $\mu$ M antipain (Wako Pure Chemicals Co., Ltd.) and 2 mM EDTA (Wako Pure Chemicals Co., Ltd.), the

wells were allowed to stand for 30 minutes at room temperature to dissolve the cells. Fifteen  $\mu$ l aliquots of these samples were heated in the presence of SDS under reducing conditions according to a conventional method and subjected to SDS-electrophoresis using 4-20% polyacrylamide gradient gel (Daichi Pure Chemical Co., Ltd.). After the electrophoresis, the gel was dried and exposed for 24 hours at -80° C. to a BioMax MS film (Kodak Co.) using a BioMax MS sensitization screen (Kodak Co.). The exposed film was developed according to a conventional method. As a result, a band with a molecular weight of a range of 90,000-110,000, shown in FIG. 10, was detected by crosslinking the <sup>125</sup>I-labeled monomer type OCIF and the protein encoded by the cDNA of the present invention.

#### Example 12

[0245] Northern Blotting Analysis

[0246] ST2 cells cultured to become confluent in a 25 cm<sup>2</sup>T flask for attached-cell cultures were treated with trypsin and stripped from the T flask. After washing, the cells were seeded into a 225 cm<sup>2</sup> T flask and cultured for 4 days in a CO<sub>2</sub> incubator with 60 ml of an  $\alpha$ -MEM culture medium containing 10<sup>-8</sup> M active-form vitamin D<sub>3</sub>, 10<sup>-7</sup> M dexamethasone, and 10% fetal bovine serum. Total RNA was extracted from the cultured ST2 cells using ISOGEN (Wako Pure Chemicals Co., Ltd.). The total RNA was also extracted in the same manner from ST2 cells which were cultured in the absence of the active-form vitamin D<sub>3</sub> and dexamethasone. After the addition of 2.0  $\mu$ l of 5 $\times$  gel electrophoresis buffer solution (0.2 M morpholino propane sulfonic acid, pH 7.0, 50 mM sodium acetate, 5 mM EDTA) and 3.5  $\mu$ l of formaldehyde, and 10.0  $\mu$ l of formamide to 20  $\mu$ g (4.5  $\mu$ l) of each of the total RNAs, the mixtures were incubated for 15 minutes at 55° C. and subjected to electrophoresis. The gel for electrophoresis was prepared according to the formulation of 1.0% agarose, 2.2 M deionized formaldehyde, 40 mM morpholinopropane sulfonic acid (pH 7.0), 10 mM sodium acetate, and 1 mM EDTA. The electrophoresis was carried out in a buffer solution of 40 mM morpholinopropane sulfonic acid, pH 7.0, 10 mM sodium acetate, and 1 mM EDTA. After the electrophoresis, RNA was transferred onto nylon membrane. About 1.0 kb DNA fragment was obtained by digesting pOBM291 with a restriction enzyme, EcoRI. Hybridization was carried out using this DNA fragment, labeled with a Megaprime DNA labeling kit (Amersham Co.) and  $\alpha$ -<sup>32</sup>P-dCTP (Amersham Co.), as a probe. As a result, as shown in FIG. 11, it was confirmed that when ST2 cells were cultured in the presence of active-form vitamin D<sub>3</sub> and dexamethasone, gene expression of the protein encoded by the cDNA of the present invention (OBM) is induced strongly.

#### Example 13

[0247] Osteoclasts Formation Supporting Capability of the Protein Encoded by the cDNA of the Present Invention

[0248] pOBM291 was transfected into COS cells according to the same method described in the Example 8(3). After three days, trypsinized cells were washed once with phosphate buffered saline solution by centrifugation, then fixed with PBS containing 1% paraformaldehyde at room temperature for 5 minutes, followed by washing with PBS six times by centrifugation. 700  $\mu$ l of 1 $\times$ 10<sup>6</sup>/ml mouse spleen

cells and 350  $\mu$ l of 4 $\times$ 10<sup>4</sup>/ml ST2 cells which were suspended in  $\alpha$ -MEM culture medium containing 10<sup>-8</sup> M active-form vitamin D<sub>3</sub>, 10<sup>-7</sup> M dexamethasone, and 10% fetal bovine serum, were added to a 24-well plate. TC insert (Nunc Co.) was set in each well. The above-described fixed COS cells (350  $\mu$ l) diluted to various concentrations with the above-mentioned culture medium and OCIF solution (50  $\mu$ l) were added to the TC insert and cultured for 6 days at 37° C. As a result, it was confirmed that the osteoclasts formation inhibitive activity of OCIF can be inhibited by the protein encoded by the cDNA of the present invention.

#### Example 14

[0249] Expression of Secreted-Form OBM

[0250] (1) Construction of a Plasmid for the Expression of Secreted-Form OBM

[0251] A PCR reaction was carried out using OBM HF (Sequence Table, Sequence ID No. 7) and OBM XR (Sequence Table, Sequence ID No. 8) as primers and pOBM291 as a template. After purification by agarose gel electrophoresis, the product was digested with restriction enzymes HindIII and EcoRI, and further purified by agarose gel electrophoresis. The purified fragment (0.6 kb), Hind III/EcoRI fragment (5.2 kb) of pSec TagA (Invitrogen Co.), and EcoRI/Pmacl fragment (0.32 kb) of OBM cDNA were ligated using a ligation kit ver. 2 (Takara Shuzo Co.) *Escherichia coli* DH5 $\alpha$  was transformed using the reaction product. Plasmids were purified by means of alkali SDS method from the resulting ampicillin resistant strains and digested with restriction enzymes to select a plasmid with fragments of a length of 0.6 Kb and 0.32 kb being inserted into pSec TagA. Selected plasmid was identified as having a sequence encoding the secreted-form OBM (nucleotide sequence: 338-1355 in Sequence ID No. 2, amino acid sequence: 72-316 in the Sequence ID No. 1) by sequencing using a dye terminator cycle sequencing FS kit (Perkin Elmer Co.). This plasmid was digested with restriction enzymes NheI and Xhol to isolate a fragment (1.0 kb) containing the secreted-form OBM cDNA by agarose gel electrophoresis. This fragment was inserted into the NheI/XhoI fragment (10.4 kb) of an expression vector, pCEP4 (Invitrogen Co.), using a ligation kit and *Escherichia coli* DH5  $\alpha$  was transformed using the reaction product thereof. Plasmids were purified by alkali SDS method from the resulting ampicillin resistant strains and digested with restriction enzymes to select an *Escherichia coli* strain having the secreted-form OBM expression plasmid (pCEP sOBM) with the correct structure. The *Escherichia coli* strain containing the pCEP sOBM was cultured and pCEP sOBM was purified using QIA filter plasmid midi kit (QIAGEN Co.).

[0252] (2) Expression of Secreted-Form OBM

[0253] 293-EBNA cells were suspended in IMDM containing 10% FCS (IMDM-10% FCS) and seeded into a 24-well plate coated with collagen (manufactured by Sumitomo Bakelite Co., Ltd.) in a cell density of 2 $\times$ 10<sup>5</sup>/2 ml/well and cultured overnight. The cells were transfected with 1  $\mu$ g of pCEP sOBM or pCEP4 using 4  $\mu$ l of Lipofectamine (Gibco Co.). After culturing for two days in 0.5 ml of a serum-free IMDM or IMDM-10% FCS, the conditioned medium was collected. Expression of the secreted-form OBM in the conditioned medium was confirmed as follows.

Sodium hydrogen carbonate was added to the conditioned medium to a final concentration of 0.1 M and the solution was added to a 96-well plate. The plate was allowed to stand overnight at 4° C., thereby immobilizing OBM in the conditioned medium on the 96-well plate. The plate was filled with a Block Ace™ (Snow Brand Milk Products Co., Ltd.) solution diluted four-fold with PBS (B-PBS) and allowed to stand for two hours at room temperature to block residual binding sites of the plate. After the addition to each well of 100  $\mu$ l of 3-100 ng/ml of OCIF which was diluted with B-PBS, the plate was allowed to stand for two hours at 37° C., followed by washing with PBS containing 0.05% Tween 20(PBS-T). Then, 100  $\mu$ l of a peroxidase-labeled rabbit anti-OCIF polyclonal antibody described in WO 96/26217 which was diluted with B-PBS was added to each well. After allowing to stand for two hours at 37° C., the wells were washed six times with PBS-T. Then, a TMB solution (TMB Soluble Reagent, High Sensitivity, Scytek Co.) was added in the amount of 100  $\mu$ l per well and allowed to stand at room temperature for about 10 minutes, whereupon the reaction was terminated by the addition of 100  $\mu$ l of a termination solution (Stopping Reagent, Scytek Co.) to each well. Absorbance at 450 nm of each well was measured by a microplate reader. The results are shown in FIG. 12 which indicates that the absorbance at 450 nm increased according to the concentration of the added OCIF in the plate in which the conditioned medium of the cells transfected with pCEP sOBM was immobilized. On the other hand, no increase in absorbance was seen in the plate in which the conditioned medium of the cells transfected with vector pCEP4 was immobilized. FIG. 13 shows the results of an experiment wherein the proportion of the conditioned medium which is used for immobilization was changed within a range of 5-90% and a specific concentration of OCIF (50 ng/ml) was added. It can be seen that the absorbance at 450 nm increased according to the increase in the proportion of the conditioned medium in the plate wherein the conditioned medium of the cells transfected with pCEP-sOBM was immobilized, whereas no such increase in absorbance was seen in the plate wherein the conditioned medium of the cells transfected with vector pCEP4 was immobilized. From these results, it was confirmed that secreted-form OBM is produced into the conditioned medium of the cells transfected with pCBP sOBM.

#### Example 15

[0254] Expression of Thioredoxin-OBM Fusion Protein (Trx-OBM)

[0255] (1) Construction of a Thioredoxin-OBM Fusion Protein (Trx-OBM) Expression Vector

[0256] Ten  $\mu$ l of 10 $\times$  ExTaq buffer (Takara Shuzo Co.), 8  $\mu$ l of 10 mM dNTP (Takara Shuzo Co.), 77.5  $\mu$ l of sterilized distilled water, 2  $\mu$ l of an aqueous solution of pOBM291 (10 ng/ $\mu$ l), 1  $\mu$ l of primer OBM3 (100 pmol/ $\mu$ l, Sequence Table, Sequence ID No. 9), 1  $\mu$ l of primer OBMSalR2 (100 pmol/ $\mu$ l, Sequence Table, Sequence ID No. 10), and 0.5  $\mu$ l of ExTaq (5  $\mu$ l) (Takara Shuzo Co.) were mixed and reacted (PCR reaction) in a micro centrifuge tube. After reacting at 95° C. for 5 minutes, at 50° C. for one second, at 55° C. for one minute, at 74° C. for one second, and at 72° C. for 5 minutes, a cycle reaction consisting of a reaction at 96° C. for one minute, at 50° C. for one second, at 55° C. for one minute, at 74° C. for one second, and at 72° C. for 3

minutes, was repeated 25 times. From the total reaction liquid DNA fragment of about 750 bp was purified by 1% agarose gel electrophoresis using QIAEX II gel extraction kit (QIAGEN Co.). The whole amount of purified DNA fragment was digested with restriction enzymes SaII and EcoRI (Takara Shuzo Co.) and subjected to an 1.5% agarose gel electrophoresis to purify a DNA fragment of about 160 bp (Fragment 1), which was dissolved in 20  $\mu$ l of sterilized distilled water. In the same manner, a DNA fragment of about 580 bp (Fragment 2) obtained by digesting 4  $\mu$ g of pOBM291 with restriction enzymes BamH1 and EcoRI (Takara Shuzo Co.) and a DNA fragment of about 3.6 kb (Fragment 3) obtained by digesting 2  $\mu$ g of pTrxFus (Invitrogen Co.) with restriction enzymes BamH1 and SaII (Takara Shuzo Co.) were respectively purified and dissolved in 20  $\mu$ l of sterilized distilled water. The QIAEXII gel extraction kit was used for the purification of DNA fragments. Fragments 1-3 were ligated by incubating at 16° C. for 2.5 hours using DNA ligation kit ver.2 (Takara Shuzo Co.). Using the ligation reaction liquid, *Escherichia coli* strain GI724 (Invitrogen Co.) was transformed according to the method described in the Instruction Manual of ThioFusion Expression System (Invitrogen Co.). A microorganism strain with plasmid in which the OBM cDNA fragment (nucleotide sequence: 350-1111 in the Sequence ID-No. 2, amino acid sequence: 76-316 in the Sequence ID No. 1) is fused in frame to a thioredoxin gene was selected from the resulting ampicillin resistant transformants by the analysis of restriction maps obtained by digestion with restriction enzymes and DNA sequence determination. The microorganism strain thus obtained was named GI724/pTrxOBM25.

[0257] (2) Expression of OBM in *Escherichia coli*

[0258] GI724/pTrxOBM25 and GI724 containing pTrxFus (GI724/pTrxFus) were respectively cultured six hours with shaking at 30° C. in 2 ml of RMG-Amp culture medium (0.6% Na<sub>2</sub>HPO<sub>4</sub>, 0.3% KH<sub>2</sub>PO<sub>4</sub>, 0.05% NaCl, 0.1% NH<sub>4</sub>Cl, 1.2% casamino acid (Difco Co.), 1% glycerol, 1 mM MgCl<sub>2</sub>, and 100  $\mu$ g/ml ampicillin (Sigma Co.), pH 7.4). The broth 0.5 ml of the broth was added to 50 ml of Induction culture medium (0.6% Na<sub>2</sub>HPO<sub>4</sub>, 0.3% KH<sub>2</sub>PO<sub>4</sub>, 0.05% NaCl, 0.1% NH<sub>4</sub>Cl, 0.2% casamino acid, 0.5% glucose, 1 mM MgCl<sub>2</sub>, 100  $\mu$ g/ml ampicillin, pH 7.4) and cultured with shaking at 30° C. When OD<sub>600 nm</sub> reached about 0.5, L-tryptophan was added to a final concentration of 0.1 mg/ml, followed by shaking the culture at 30° C. for an additional 6 hours. The culture broth was centrifuged at 3000xg to collect the cells, which were suspended in 12.5 ml of PBS (10 mM phosphate buffer, 0.15 M NaCl, pH 7.4). The suspension was subjected to an ultrasonic generator (Ultrasonics Co.) to disrupt the cells. The disrupted cells were centrifuged at 7000xg for 30 minutes to obtain a supernatant liquid as a soluble protein fraction. Ten  $\mu$ l of this soluble protein fraction was subjected to SDS polyacrylamide (10%) electrophoresis under reducing conditions. As a result, a band with a molecular weight of 40 kDa which was not detected in the soluble protein fraction of GI724/pTrxFus was found in the soluble protein fraction of GI724/pTrxOBM25 (FIG. 14). Accordingly, it was confirmed that a fusion protein (Trx-OBM) of thioredoxin and OBM was expressed in *Escherichia coli*.

[0259] (3) Binding Capability of Trx-OBM to OCIF

[0260] Binding of the expressed Trx-OBM to OCIF was confirmed according to the following experiment. Anti-

thioredoxin antibody (Invirogen Co.) which was diluted to 5000-fold with 10 mM sodium hydrogen carbonate solution was added to a 96-well immunoplate (Nunc Co.) in the amount of 100  $\mu$ l per well. After being allowed to stand overnight at 4° C., the liquid in the wells was discarded. Two hundred  $\mu$ l of a solution prepared by diluting Block Ace™ (Snow Brand Milk Products Co., Ltd.) two-fold with PBS (BA-PBS) was added to each well. After being allowed to stand for one hour at room temperature, the solution was discarded and soluble protein fractions originating from the above-described GI724/pTrxOBM25 or GI724/pTrxFus, each diluted with BA-PBS in various concentrations were added to each well in the amount of 100  $\mu$ l. After being allowed to stand for two hours at room temperature, each well was washed three times with PBS-T and charged with 100  $\mu$ l of OCIF (100 ng/ml) which was diluted with BA-PBS. After being allowed to stand for two hours at room temperature, each well was washed three times with PBS-T and charged with 100  $\mu$ l of peroxidase-labeled rabbit anti-OCIF polyclonal antibody (described in WO 96/26217) which was diluted 2,000-fold with BA-PBS. After being allowed to stand for two hours at room temperature, each well was washed six times with PBS-T and charged with 100  $\mu$ l of TMB solution (TMB Soluble Reagent, High Sensitivity, Scytek Co.). After being allowed to stand for about 10 minutes at room temperature, each well was charged with 100  $\mu$ l of termination solution (Stopping Reagent, Scytek Co.). Absorbance of each well at 450 nm was measured by a microplate reader. The results are shown in FIG. 15. There was no difference in absorbance between the sample with the soluble protein fraction originating from GI724/pTrxFus added thereto and the sample without the addition of this soluble protein fraction. On the other hand, the absorbance increased in the samples to which the soluble protein fraction originating from GI724/pTrxOBM25 was added in proportion to the concentration of the soluble protein fraction. The results of the other experiment wherein the dilution rate of the soluble protein fraction was maintained constant (1%) while adding OCIF diluted with BA-PBS in different concentrations (0-100 ng/ml) are shown in FIG. 16. It can be seen that the absorbance remained low at any concentrations of OCIF in samples using a soluble protein fraction originating from GI724/pTrxFus, whereas the absorbance increased in proportion to the OCIF concentration in the samples to which the soluble protein fraction originating from GI724/pTrxOBM25 was added. Based on these results, it was confirmed that Trx-OBM which is produced from GI724/pTrxOBM25 has a capability of binding to OCIF.

[0261] (4) Large-Scale Cultivation of *Escherichia coli* Which Produces Trx-OBM

[0262] GI724/pTrxOBM25 cells were spread on RMG-Amp agar (0.6% Na<sub>2</sub>PO<sub>4</sub>, 0.3% KH<sub>2</sub>PO<sub>4</sub>, 0.05% NaCl, 0.1% NH<sub>4</sub>Cl, 2% casamino acid, 1% glycerol, 1 mM MgCl<sub>2</sub>, 100  $\mu$ g/ml ampicillin, 1.5% agar, pH 7.4) using a platinum transfer loop. The cells were cultured overnight at 30° C. The cultured cells were suspended in 10 ml of Induction medium. The suspension was added 5 ml for each to two 2 l Erlenmeyer flasks containing 500 ml of Induction medium and cultured at 30° C. with shaking. When the OD<sub>600 nm</sub> reached about 0.5, L-tryptophan was added to a final concentration of 0.1 mg/ml. Culturing with shaking was continued for six hours at 30° C. The culture broth was centrifuged for 20 minutes at 3000xg to collect the cells, which were suspended in 160 ml of PBS. The suspension

was subjected to an ultrasonic generator (Ultrasonics Co.) to disrupt the cells. The supernatant liquid was centrifuged for 30 minutes at 7000×g to obtain a soluble protein fraction.

[0263] (5) Preparation of OCIF-Immobilized Affinity Column

[0264] Two g of TSKgel AF-Tolesyl Toyopal 650 (Tosoh Corp.) and 40 ml of 1.0 M potassium phosphate buffer (pH 7.5) containing 35.0 mg of recombinant OCIF, which was prepared according to the method described in WO 96/26217, were mixed. The mixture was gently shaken overnight at 4°C to effect a coupling reaction. The reaction mixture was centrifuged to remove the supernatant. To inactivate excess active residues, 40 ml of 0.1 M Tris-HCl buffer (pH 7.5) was added to the precipitated carrier and the mixture was gently shaken at room temperature for one hour. The carrier in a column was washed with 0.1 M glycine-HCl buffer (pH 3.3) containing 0.01% Polysorbate 80 and 0.2 M NaCl and 0.1 M sodium citrate buffer (pH 2.0) containing 0.01% Polysorbate 80 and 0.2 M NaCl. The carrier in the column was equilibrated by charging twice with 10 mM sodium phosphate buffer (pH 7.4) containing 0.01% Polysorbate 80.

[0265] (6) Purification of Trx-OBM Using OCIF-Immobilized Affinity Column

[0266] Unless otherwise indicated, purification of Trx-OBM was carried out at 4°C. The above-mentioned OCIF-immobilized affinity carrier (10 ml) and the soluble protein fraction (120 ml) prepared in Example 15(4) were mixed. The mixture was gently shaken overnight at 4°C in four 50-ml centrifuge tubes using a rotor. An Econo-column™ (internal diameter: 1.5 cm, length: 15 cm, manufactured by BioRad Co.) was filled with the carrier in the mixture. The column was charged with 300 ml of PBS containing 0.01% Polysorbate 80, 100 ml of 10 mM sodium phosphate buffer (pH 7.0) containing 0.01% Polysorbate 80 and 2 M NaCl, and 100 ml of 0.1 M glycine-HCl buffer (pH 3.3) containing 0.01% Polysorbate 80 and 0.2 M NaCl, in that order. Next, proteins adsorbed in the column were eluted with 0.1 M sodium citrate buffer (pH 2.0) containing 0.01% Polysorbate 80 and 0.2 M NaCl. The eluate was collected in 5 ml portions. Each fraction thus collected was immediately neutralized with addition 10% volume of 2 M Tris buffer (pH 8.0). Presence or absence of Trx-OBM in the eluted fractions was determined according to the method previously described in Example 15(3) (the binding capability to OCIF). The fractions containing Trx-OBM were collected and purified further.

[0267] (7) Purification of Trx-OBM by Gel Filtration

[0268] About 25 ml of Trx-OBM fractions obtained in Example 15(6) was concentrated to about 0.5 ml by centrifuge using Centriplus 10 and Centricon 10 (Amicon Co.). This sample was applied to a Superose 12 HR 10/30 column (1.0×30 cm, Pharmacia Co.) previously equilibrated with PBS containing 0.01% Polysorbate 80. For the separation, PBS containing 0.01% Polysorbate 80 was used as a mobile phase at a flow rate of 0.25 ml/min. The eluate from the column was collected in 0.25 ml portions. The Trx-OBM in the thus-collected fractions was detected by the same method as previously described in Example 15(3) and by SDS-polyacrylamide electrophoresis (10-15% polyacrylamide gel, Pharmacia Co.) using Phast System (Pharmacia

Co.) and silver staining. Fractions (Fr. 20-23) containing purified Trx-OBM were collected and the protein concentration of Trx-OBM was determined. The measurement of the protein concentration was carried out using bovine serum albumin as a standard substance using DC-Protein assay kit (BioRad Co.).

Example 16

[0269] Osteoclast Formation-Inducing Activity of OBM

[0270] pOBM291 and pcDL-SR α296 were respectively transfected into COS-7 cells using Lipofectamine (Gibco Co.). The cells were cultured in DMEM containing 10% FCS for one day, trypsinized, plated on cover slips (15 mm round shape, manufactured by Matsunami Co.) in 24-well plates at 5×10<sup>4</sup> cells per well, and cultured for 2 days. The culture plate was washed once with PBS. The cells were fixed with PBS containing 1% paraformaldehyde at room temperature for 8 minutes. The plate on which the fixed cells were attached was washed 6 times with PBS, then 700 μl of mouse spleen cells suspended at 1×10<sup>6</sup>/ml in α-MEM containing 10<sup>-8</sup> M active-form vitamin D<sub>3</sub>, 10<sup>-7</sup> M dexamethasone, and 10% fetal bovine serum were added to each well. Millicell PCF (Millipore Co.) was set in each well and a suspension of ST2 cells in the above-mentioned culture medium (4×10<sup>4</sup>/ml) were added, 700 μl per well, into the Millicell PCF followed by incubation at 37°C for 6 days. After the culture, the Millicell PCF was removed, the plate was washed once with PBS, and the cells were fixed with acetone-ethanol solution (50:50) for one minute. Then, cells exhibiting tartaric acid-resistant acid phosphatase activity (TRAP), which is a specific marker for osteoclast, were selectively stained using LEUKOCYTE ACID PHOSPHATASE kit (Sigma Co.). As a result of microscopic observation, TRAP-positive cells were not detected in the wells in which COS-7 cells transfected with pcDL-SR α296 were fixed. In contrast, 45±18 (average±standard deviation, n=3) TRAP positive cells were observed in the wells in which COS-7 cells transfected with pOBM291 were fixed. Moreover, it was also confirmed that calcitonin bound to these TRAP positive cells. Based on these findings, it has been proven that OBM has osteoclast formation-inducing activity.

Example 17

[0271] Osteoclast Formation-Inducing Activity of Trx-OBM and Secreted-Form OBM

[0272] Mouse spleen cells were suspended in α-MEM containing 10<sup>-8</sup> M active-form vitamin D<sub>3</sub>, 10<sup>-7</sup> M dexamethasone, and 10% fetal bovine serum at a concentration of 2×10<sup>6</sup>/ml. The suspension was added to a 24 well plate in the amount of 350 μl per well. Each well was then charged with 350 μl of a solution prepared by diluting purified Trx-OBM with the above-mentioned culture medium (40 ng/ml), 350 μl of solution prepared by 10-fold diluting conditioned medium which was produced by culturing 293-EBNA cells, in which pCEP sOBM or pCEP4 were transfected, in IMDM-10% FCS, with the above-mentioned culture medium, or 350 μl only of the above-mentioned culture medium. The Millicell PCF (Millipore Co.) was set on each well, to which 600 μl of ST2 cells which was suspended in the above-mentioned culture medium (4×10<sup>4</sup>/ml) were added. After cultured for six days, Millicell PCF was

removed. The plate was washed once with PBS and the cells were fixed with acetone-ethanol solution (50:50) for one minute. Then, the cells exhibiting the tartaric acid resistant acidic phosphatase activity (TRAP activity) were selectively stained using LEUKOCYTE ACID PHOSPHATASE kit (Sigma Co.). The result of microscopic observation revealed that no cells exhibiting the TRAP activity were detected in the wells to which no Trx-OBM was added, whereas  $106 \pm 21$  (average  $\pm$  standard deviation, n=3) TRAP-positive cells were observed in the wells to which Trx-OBM was added. Similarly, while no cells exhibiting TRAP activity were detected in the wells to which conditioned medium of 293-EBNA transfected with pCEP4 had been added,  $120 \pm 31$  (average  $\pm$  standard deviation, n=3) TRAP positive cells were observed in the wells to which conditioned medium of 293-EBNA transfected with pCEPsOBM had been added. Moreover, it was also confirmed that calcitonin binds to these TRAP positive cells. These results have proven that Trx-OBM and secreted-form OBM exhibit osteoclast formation-inducing activity.

#### Example 18

##### Identity of the Protein OBM Expressed by the cDNA of the Present Invention and the Natural Type OCIF-Binding Protein of the Present Invention

[0273] (1) Preparation of Rabbit Anti-OBM Polyclonal Antibody

[0274] Three male Japanese white rabbits (weight: 2.5-3.0 kg, supplied by Kitayama Labes Co.) were immunized with the purified OBM (thioredoxin-OBM fusion protein) produced according to the method in Examples 14(6) and 14(7) by subcutaneously injecting 1 ml/dose of emulsion prepared by mixing 200  $\mu$ g/ml of the purified OBM with equal volume of Freund's complete adjuvant (DIFCO Co.), six times, once a week. Ten days after the last immunization, the rabbits were exsanguinated. Antibody was purified from the serum as follows. Ammonium sulfate was added to the antiserum which was diluted two-fold with PBS to a final concentration of 40% (w/v %). After being allowed to stand for one hour at 4° C., the mixture was centrifuged for 20 minutes at 8000 $\times$ g to obtain a precipitate. The precipitate was dissolved in a small amount of PBS, dialyzed against PBS at 4° C., and loaded to a Protein G-Sepharose column (manufactured by Pharmacia Co.) After washed with PBS, the adsorbed immunoglobulin G was eluted with 0.1 M glycine-HCl buffer solution (pH 3.0). The eluate was immediately neutralized with 1.5 M Tris-HCl buffer (pH 8.7). After dialyzing the eluted protein fractions against PBS, the absorbance at 280 nm was measured to determine the protein concentration (E<sup>1%</sup> 13.5). Anti-OBM antibody labeled with horseradish peroxidase was prepared using a maleimide-activated paroxidase kit (Pierce Co.) as follows. 80  $\mu$ g of N-succinimide-S-acetyl thioacetic acid was added to 1 mg of the purified antibody and reacted at room temperature for 30 minutes. Five mg of hydroxylamine was added to the resulting mixture to deacetylate the antibody. The modified antibody was fractionated using a polyacrylamide desalting column. The protein fractions were mixed with 1 mg of maleimide-activated peroxidase and reacted for one hour at room temperature to obtain enzyme-labeled antibody.

[0275] (2) Capability of Rabbit Anti-OBM Polyclonal Antibody to Inhibit Specific Binding of the Protein (OBM) Expressed by the cDNA of the Present Invention or the Natural Type Protein of the Present Invention With OCIF

[0276] Purified OBM (thioredoxin-OBM fused protein) obtained according to the method described in the Examples 15(6) and 15(7) and the natural type purified OCIF-binding protein of the Example 2(4) were dissolved respectively in 0.1 M sodium carbonate buffer to a concentration of 2  $\mu$ g/ml. An aliquot of each solution was added 100  $\mu$ l per well respectively to a 96-well immunoplate (manufactured by Nunc Co.). The plate was allowed to stand overnight at 4° C. 200  $\mu$ l of 50% Block Ace was added to each well and the plate was allowed to stand at room temperature for one hour. After washing each well three times with PBS containing 0.1% Polysorbate 20 (P20-PBS), 100  $\mu$ l of rabbit anti-OBM antibody solution which was dissolved in 25% Block Ace prepared with P20-PBS to a concentration of 200  $\mu$ g/ml or 100  $\mu$ l of 25% Block Ace (containing no antibody) was added to each well, followed by incubation at 37° C. for one hour. Each well was washed three times with P20-PBS and charged with 100  $\mu$ l/well of a binding test solution (P20-PBS containing 0.2% calf serum albumin, 20 mM Hepes, and 0.1 mg/ml heparin) to which 20 ng/ml of <sup>125</sup>I-labeled OCIF described in the Example 8(3) was added. Alternatively, each well was charged with 100  $\mu$ l/well of another binding test solution containing 8  $\mu$ g/ml of unlabeled OCIF in addition to 20 ng/ml <sup>125</sup>I-labeled OCIF. After incubating these immunoplates at 37° C. for one hour, the wells were washed with P20-PBS six times. The amount of <sup>125</sup>I in each well was measured by a gamma counter. The results are shown in FIG. 17. As shown in the figure, both the purified OBM expressed using the cDNA of the present invention and the protein that specifically bind the natural type OCIF-specifically binding protein of the present invention do not bind to the <sup>125</sup>I-labeled OCIF at all, when they were treated with the rabbit anti-OBM polyclonal antibody, whereas both proteins bound <sup>125</sup>I-labeled OCIF when untreated with the antibody. The binding of both proteins to <sup>125</sup>I-labeled OCIF was confirmed to be clearly specific, because those bindings are significantly inhibited by the addition of 400-fold concentration of unlabelled OCIF (8  $\mu$ g/ml). Based on the results described above, the rabbit anti-OBM polyclonal antibody recognizes both the OBM which is the protein expressed using the cDNA of the present invention and the natural-type OCIF-binding protein of the present invention, and it inhibits the specific binding of these proteins with OCIF.

#### Example 19

[0277] Cloning of Human OBM cDNA

[0278] (1) Preparation of Mouse OBM Primer

[0279] The mouse OBM primers prepared according to the method of the Examples (OBM#3 and OBM#8) described above, were used for screening of human OBM cDNA. The sequences are shown in the Sequence Table, Sequence ID No. 9 and No. 6, respectively.

[0280] (2) Isolation of Human OBM cDNA Fragment by PCR

[0281] Human OBM cDNA fragments were obtained by PCR using the mouse OBM cDNA primers prepared in (1)

above and Human Lymph Node Marathon ready cDNA (Clontech Co.) as a template. The conditions for PCR were shown as follows:

10 × EX Taq buffer (Takara Shuzo Co.)	2 $\mu$ l
2.5 mM dNTP	1.6 $\mu$ l
cDNA solution	1 $\mu$ l
EX Taq (Takara Shuzo Co.)	0.2 $\mu$ l
Distilled water	14.8 $\mu$ l
40 $\mu$ M primer OBM#3	0.2 $\mu$ l
40 $\mu$ M primer OBM#8	0.2 $\mu$ l

[0282] These solutions were mixed in a microfuge tube and pre-incubated at 95° C. for 2 minutes, followed by 40 cycles of a three-stage reaction consisted of reactions at 95° C. for 30 seconds, at 57° C. for 30 seconds, and at 72° C. for 2.5 minutes. After the reaction, the solution was incubated for 5 minutes at 72° C. and a portion of the solution was subjected to electrophoresis on an agarose gel. A DNA fragment (about 690 bp) amplified by the mouse OBM cDNA primers described above was detected.

[0283] (3) Purification of the Human OBM cDNA Fragment Amplified by PCR and Determination of the Nucleotide Sequence

[0284] The human OBM cDNA fragment obtained in the Example 19(2) was separated by-electrophoresis on an agarose gel and further purified using a QIAEX gel extraction kit (Qiagen Co.). PCR was again performed using the purified human OBM cDNA fragment as a template and the mouse OBM cDNA primers described above, to produce a large quantity of the human OBM cDNA fragment. The DNA fragment was purified by a QIAEX gel extraction kit in the same manner as above. The nucleotide sequence of the purified human OBM cDNA fragment was determined using Taq Dye Deoxy Terminator Cycle Sequencing FS kit (Perkin Elmer Co.) using OBM#3 or OBM#8 (Sequence ID No. 9 and No. 6 respectively) as a primer. When compared with the sequence of corresponding area of the mouse OBM cDNA, the nucleotide sequence of the human OBM cDNA fragment showed 80.7% homology with that of the mouse OBM cDNA.

[0285] (4) Screening of a Full-Length Human OBM cDNA by Hybridization Using the Human OBM cDNA Fragment (about 690bp) as a Probe

[0286] A full-length human OBM cDNA was screened using the human OBM cDNA fragment (about 690 bp) that was purified in the Example 19(3) and labeled with [ $\alpha$ <sup>32</sup>P] dCTP using a Megaprime DNA Labeling kit (Amersham Co.). Human Lymph Node 5'-STRETCH PLUS cDNA library (Clontech Co., the U.S.A) was screened using the DNA probe. According to the manufacturer's protocol, *Escherichia coli* C600 Hfl was infected with the recombinant phage for 15 minutes at 37° C. The infected *Escherichia coli* was added to an LB agar (1% trypton, 0.5% yeast extract, 1% NaCl, 0.7% agar) which was heated at 45° C. The LB agar was poured onto an LB agar plate containing 1.5% agar. After overnight incubation at 37° C., HyBond-N<sup>TM</sup> (Amersham Co.) was placed to the plate on which plaques were produced and stored for about 3 minutes. According to a conventional method, the filter was treated with alkaline solution, neutralized, and dipped in 2 $\times$ SSC

solution. DNA was then immobilized onto the filter using the UV CROSSLINKER (Stratagene Co.). The resulting filter was dipped into Rapid-hyb buffer (Amersham Co.). After pretreatment for 15 minutes at 65° C., the filter was placed in Rapid-hyb buffer containing the heat-denatured human OBM cDNA fragment (about 690 bp, 5 $\times$ 10<sup>5</sup> cpm/ml) described above. After overnight hybridization at 65° C., the filter was washed with 2 $\times$ SSC, 1 $\times$ SSC, and 0.1 $\times$ SSC, each containing 0.1% SDS, in this order respectively for 15 minutes at 65° C. Several positive clones obtained were further purified by repeating the screening twice. A clone possessing an insert (about 2.2 kb) was selected from the purified clones and was used in the following experiments. This purified phage was named  $\lambda$ hOBM. About 10  $\mu$ g of DNA was obtained from the purified  $\lambda$ hOBM using a QIAGEN Lambda kit (Qiagen Co.) according to the manufacturer's protocol. The DNA was digested with restriction enzyme Sall and subjected to electrophoresis on an agarose gel to separate the hOBM insert cDNA (about 2.2 kb). This DNA fragment purified using the QIAEX gel extraction kit (Qiagen Co.) was digested with restriction enzyme Sall and inserted into plasmid pUC19 (MBI Co.) which was previously digested with a restriction enzyme Sall and dephosphorylated, using a DNA ligation kit ver. 2 (Takara Shuzo Co.). *Escherichia coli* DH 5 $\alpha$  (Gibco BRL Co.) was transformed with the pUC19 containing the resulting DNA fragment. The resulting transformant was named pUC19hOBM. The transformant was grown and pUC19hOBM in which the human OBM cDNA (about 2.2 kb) was inserted and purified by a conventional method.

[0287] (5) Determination of Nucleotide Sequence of cDNA Encoding the Entire Amino Acid Sequence of Human OBM

[0288] The nucleotide sequence of the resulting human OBM cDNA obtained in Example 19(4) was determined using the Taq Dye Deoxy Terminator Cycle Sequencing FS kit (Perkin Elmer Co.). Specifically, the nucleotide sequence of the inserted fragment was determined using pUC19hOBM as a template. As primers, primers for the determination of the nucleotide sequence of the inserted fragment DNA in pUC19hOBM, M13 Primer M3 and M13 Primer RV (manufactured by Takara Shuzo Co.), and a synthetic primer, human OBM#8, designed based on the nucleotide sequence of the human OBM cDNA fragment (about 690 bp) were used.

[0289] The nucleotide sequence of the primers used, M13 Primer M3 and M13 Primer RV, are respectively shown as the Sequence ID No. 4 and No. 5. The amino acid sequence of human OBM deduced from the nucleotide sequence of human OBM cDNA is shown in the Sequence Table as Sequence ID No. 11. The nucleotide sequence of human OBM cDNA is shown as Sequence ID No. 12.

[0290] The *Escherichia coli* which was transformed by the pUC19hOBM, which is the plasmid containing the resulting human OBM cDNA, was deposited in National Institute of Bioscience and Human Technology, Agency of Industrial Science and Technology, on Aug. 13, 1997 as deposition No. FERM BP-6058.

## Example 20

[0291] Radioiodination of OCIF with  $^{125}\text{I}$  and Quantitative Analysis of  $^{125}\text{I}$ -Labeled OCIF by ELISA

[0292] OCIF was labeled with  $^{125}\text{I}$  using the IODO-GEN method. Twenty  $\mu\text{l}$  of 2.5 mg/ml IODO-GEN-chloroform solution was transferred to a 1.5 ml Eppendorf tube and the chloroform was evaporated at 40° C., thereby providing a tube coated with IODO-GEN. The tube was washed three times with 400  $\mu\text{l}$  of 0.5 M sodium phosphate buffer solution (Na-Pi, pH 7.0), followed by the addition of 5  $\mu\text{l}$  of 0.5 M Na-Pi (pH 7.0). To this tube was added 1.3  $\mu\text{l}$  (18.5 MBq) of Na- $^{125}\text{I}$  solution (Amersham Co., NEZ-033H), immediately followed by the addition of 10  $\mu\text{l}$  of 1 mg/ml OCIF solution (monomer type or dimer type). The mixture was mixed in a vortex mixer and allowed to stand at room temperature for 30 seconds. This solution was transferred to a tube to which 80  $\mu\text{l}$  of 0.5 M Na-Pi (pH 7.0) solution containing 10 mg/ml potassium iodine and 5  $\mu\text{l}$  of a phosphate buffered saline solution containing 5% bovine serum albumin (BSA-PBS) were previously added. The solution was mixed, applied to a spin column (1 ml, G-25 Sephadex fine, manufactured by Pharmacia Co.) which was equilibrated with BSA-PBS in advance, and centrifuged for 5 minutes at 2,000 rpm. Four hundred  $\mu\text{l}$  of BSA-PBS was added to the fractions eluted from the column. After mixing, 2  $\mu\text{l}$  of the solution was used to measure the radioactivity by a gamma counter. The radiochemical purity of the  $^{125}\text{I}$ -labeled OCIF solution obtained above was measured by counting the radioactivity of fractions precipitated by 10% trichloroacetic acid (TCA).

[0293] The biological activity of the  $^{125}\text{I}$ -labeled OCIF was measured according to the method described in WO 96/26217. The concentration of the  $^{125}\text{I}$ -labeled OCIF was measured using the ELISA method as follows. Specifically, 50 mM NaHCO<sub>3</sub> (pH 9.6) in which rabbit anti-OCIF polyclonal antibody described in the WO 96/26217 was dissolved to a concentration of 2  $\mu\text{g}/\text{ml}$  was added to each well of a 96-well immunoplate (MaxiSorp™, manufactured by Nunc Co.) in the amount of 100  $\mu\text{l}$  per well. After these wells were allowed to stand overnight at 4° C., solution was removed. Then the wells were charged with a mixed aqueous solution of Block Ace™ (Snow Brand Milk Products Co., Ltd.) and a phosphate buffered saline solution (25:75) (B-PBS) in the amount of 200  $\mu\text{l}/\text{well}$ . The plate was then allowed to stand for two hours at room temperature. After the solution was removed, the wells were washed three times with a phosphate buffered saline solution containing 0.01% Polysolvate 80 (P-PBS). Next, B-PBS containing  $^{125}\text{I}$ -labeled OCIF sample or the standard OCIF was added in the amount of 100  $\mu\text{l}/\text{well}$ . The plate was then allowed to stand for two hours at room temperature. After the solution was removed, each well was washed six times with 200  $\mu\text{l}$  of P-PBS. A solution prepared by diluting peroxidase-labeled rabbit anti-OCIF polyclonal antibody with B-PBS was added in the amount of 100  $\mu\text{l}/\text{well}$ . The plate was allowed to stand for two hours at room temperature. After the solution was removed, the wells were washed six times with 200  $\mu\text{l}$  of P-PBS. Then, a TMB solution (TMB Soluble Reagent, High Sensitivity, Scytek Co.) was added in the amount of 100  $\mu\text{l}/\text{well}$ . After being allowed to stand at room temperature for 2-3 minutes, 100  $\mu\text{l}$  of a termination solution (Stopping Reagent, Scytek Co.) was added to each well. Absorbance of each well was measured at 450 nm using a microplate reader. The concentration of  $^{125}\text{I}$ -labeled OCIF was determined with a calibration curve prepared using the standard OCIF.

## Example 21

[0294] Expression of the Protein Encoded by cDNA of the Present Invention

[0295] (1) Construction of hOBM Expression Vector for Animal Cells

[0296] pUChOBM was digested with restriction enzyme SalI and a DNA fragment (about 2.2 kb) were purified by electrophoresis on an 1% agarose gel. The ends of the DNA fragments were blunted using a DNA blunting kit (Takara Shuzo Co.) (blunted hOBMcDNA fragment). Expression plasmid pcDL-SRα296 (Molecular and Cellar Biology, Vol. 8, pp 466-472 (1988)) was digested with restriction enzyme EcoRI, blunted with blunting kit and ligated with the blunted hOBM cDNA fragment using DNA ligation kit ver. 2. *Escherichia coli* DH $\alpha$  was transformed with the ligation reaction. A plasmid in the resulting ampicillin resistant transformant was subjected to digestion with restriction enzyme to analyze the DNA restriction map and determine the DNA sequence. As a result, a strain having a plasmid in which hOBM cDNA is inserted in the same direction of transcription as that of SRα promotor was selected. The microorganism strain was named DH5  $\alpha$ /phOBM.

[0297] (2) Expression of Human OBM in COS-7 Cells

[0298] *Escherichia coli* DH5  $\alpha$ /phOBM was cultured and plasmid phOBM was purified using Qiafilter Plasmid Midi kit (Qiagen Co.) phOBM was transfected using Lipofectamine into COS-7 cells in the wells of a 6-well plate and cultured for two days in DMEM containing 10% fetal bovine serum. The culture medium was replaced with cysteine-methionine-free DMEM (manufactured by Dainippon Seiyaku Co., Ltd.) to which 5% dialysed fetal bovine serum was added (88  $\mu\text{l}/\text{well}$ ). The cells were incubated for 15 minutes, followed by addition of 14  $\mu\text{l}$  of Express Protein LabelingMix (NEN Co., 10 mCi/ml). After four hours incubation, 200  $\mu\text{l}$  of DMEM containing 10% fetal bovine serum was added to each well. The cells were cultured for one hour and washed twice with PBS. Then, 0.5 ml of a TSA buffer (10 mM Tris-HCl containing 0.14 M NaCl and 0.025% Na<sub>3</sub>N, pH 8.0) containing 1% Triton X-100, 1% bovine hemoglobin, 10  $\mu\text{g}/\text{ml}$  leupeptin, 0.2 TIU/ml aprotinin, and 1 mM PMSF was added to each well and the mixtures were allowed to stand for one hour on ice. The cells were mixed by pipetting and centrifuged at 3,000×g, for 10 minutes at 4° C., to obtain supernatants. Two hundred  $\mu\text{l}$  of a dilution buffer (TSA buffer containing 0.1% Triton X-100, 0.1% bovine hemoglobin, 10  $\mu\text{g}/\text{ml}$  leupeptin, 0.2 TIU/ml aprotinin, and 1 mM PMSF) was added to 100  $\mu\text{l}$  of the supernatant from each well. The resulting mixtures were agitated at 4° C. for one hour together with Protein A Sepharose (50  $\mu\text{l}$ ) and centrifuged at 1,500×g for one minute at 4° C., to collect supernatants, thereby removing the protein which non-specifically adsorbed Protein A Sepharose. OCIF (1  $\mu\text{g}$ ) was added to the supernatants and the mixtures were agitated for one hour at 4° C. to bind human OBM and OCIF. Then, rabbit anti-OCIF polyclonal antibody (50  $\mu\text{g}$ ) was added, followed by agitation at 4° C. for one hour. Protein A Sepharose (10  $\mu\text{l}$ ) was added to the resulting solution, followed by agitation at 4° C. for an additional hour. The mixtures thus obtained were centrifuged for 1 minute at 1,500×g at 4° C. to collect precipitates. The precipitates were washed twice with a dilution buffer, twice with bovine hemoglobin-free dilution buffer, once

with TCA buffer, and once with 50 mM Tris-HCl (pH 6.5). After addition of SDS buffer containing 10%  $\beta$ -mercaptoethanol (0.125 M Tris-HCl, 4% sodium dodecylsulfate, 20% glycerol, 0.002% Bromophenol Blue, pH 6.8), the mixture was heated for 5 minutes at 100° C. and subjected to SDS-PAGE (12.5% polyacrylamide gel, Daiichi Pure Chemical Co.). The gel was fixed and dried according to a conventional method. After isotope signals were enhanced using Amplify™ (Amersham Co.), the dried gel was subjected to autoradiography at -80° C. using Bio Max MR film (Kodak Co.). The results are shown in FIG. 18, which shows that the molecular weight of the protein encoded by the cDNA of the present invention is about 40,000.

#### Example 22

[0299] Binding of the Protein Encoded by cDNA of the Present Invention and OCIF

[0300] PhOBM, which was purified in the same manner as in the Example 21(2), was transfected into COS-7 cells in each well of a 24-well plate using Lipofectamine. After cultured for 2 to 3 days, the cells were washed with serum-free DMEM. Two hundred  $\mu$ l of a culture medium for a binding test medium (serum-free DMEM to which 0.2% bovine serum albumin, 20 mM Hepes buffer solution, 0.1 mg/ml heparin, and 0.2% NaN<sub>3</sub> were added) containing 20 ng/ml of <sup>125</sup>I-labeled OCIF was added to the wells. To the other wells, 200  $\mu$ l of culture medium for the binding test medium containing 8  $\mu$ g/ml of unlabeled OCIF in addition to 20 ng/ml of <sup>125</sup>I-labeled OCIF, was added. After incubation for one hour at 37° C. in a CO<sub>2</sub> incubator (5% CO<sub>2</sub>), the cells were washed twice with 500  $\mu$ l of a phosphate buffered saline solution containing 0.1 mg/ml of heparin. Then, 500  $\mu$ l of 0.1 N NaOH solution was added to each well and the plate was allowed to stand for 10 minutes at room temperature to dissolve the cells. The radioactivity of <sup>125</sup>I in the wells was measured by a gamma counter. As a result, as shown in FIG. 19, it was confirmed that the <sup>125</sup>I-labeled OCIF binds only to the cells transfected with phOBM. Moreover, the binding was significantly inhibited by adding 400-fold excess unlabeled OCIF (8  $\mu$ g/ml). Based on the results described above, the protein, human OBM encoded by the cDNA in the phOBM was confirmed to specifically bind to OCIF on the surface of COS-7 cells.

#### Example 23

[0301] Crosslinking of <sup>125</sup>I-Labeled OCIF and the Protein Encoded by the cDNA of the Present Invention

[0302] Crosslinking of <sup>125</sup>I-labeled monomer type OCIF and the protein encoded by the cDNA of the present invention was carried out to further investigate the characteristics of the protein encoded by the cDNA of the present invention. After constructing expression vector phOBM and transfecting into COS-7 cells according to the method used in the Examples 21(1) and 21(2), 200  $\mu$ l of binding test medium containing <sup>125</sup>I-labeled OCIF (25 ng/ml) described above was added. The binding test medium to which unlabeled OCIF was added at a 400-fold concentration in addition to <sup>125</sup>I-labeled OCIF was used for the other wells. After cultured for one hour at 37° C. in a CO<sub>2</sub> incubator (5% CO<sub>2</sub>), the cells were washed twice with 500  $\mu$ l of phosphate buffered saline containing 0.1 mg/ml heparin. Five hundred  $\mu$ l of phosphate buffered saline in which 100  $\mu$ g/ml of a

crosslinking agent (DSS: disuccinimidyl suberate, manufactured by Pierce Co.) was dissolved and added to the cells, followed by incubation for 10 minutes at 0° C. The cells in these wells were washed twice with 1 ml of ice-cold phosphate buffered saline. After an addition of 100  $\mu$ l of 20 mM Hepes buffer solution containing 1% Triton X-100 (Wako Pure Chemicals Co., Ltd.), 2 mM PMSF (Phenylmethylsulfonyl fluoride, Sigma Co.), 10  $\mu$ M Pepstatin (Wako Pure Chemicals Co., Ltd.), 10  $\mu$ M leupeptin (Wako Pure Chemicals Co., Ltd.), 10  $\mu$ M antipain (Wako Pure Chemicals Co., Ltd.) and 2 mM EDTA (Wako Pure Chemicals Co., Ltd.), the wells were allowed to stand for 30 minutes at room temperature to dissolve the cells. These samples (15  $\mu$ l aliquots) were treated with SDS under reducing conditions according to a conventional method and subjected to SDS-electrophoresis using 4-20% polyacrylamide gradient gel (Daiichi Pure Chemical Co., Ltd.). After electrophoresis, the gel was dried and subjected to autoradiography for 24 hours at -80° C. using BioMax MS film (Kodak Co.) and BioMax MS sensitization screen (Kodak Co.). The film subjected to autoradiography was developed according to a conventional method. As a result, a band of a molecular weight in the range of 90,000-110,000, shown in FIG. 20, was detected by crosslinking of <sup>125</sup>I-labeled monomer type OCIF and the protein encoded by the cDNA of the present invention.

#### Example 24

[0303] Expression of Secreted-Form Human OBM

[0304] (1) Construction of Secreted-Form Human OBM Expression Plasmid

[0305] A PCR was carried out using human OBM SF (Sequence Table, Sequence ID No. 13) and mouse OBM #8 (Sequence Table, Sequence ID No. 6) as primers and pUC19hOBM as a template. After purified by electrophoresis on an agarose gel, the product was digested with restriction enzymes SpII and HindIII, and further purified by electrophoresis on an agarose gel to obtain a purified fragment (0.27 kb). Human OBM cDNA was partially digested with restriction enzyme DraI and DNA fragments digested with DraI at one site were purified by electrophoresis on an agarose gel. The purified fragment was further digested with restriction enzyme HindIII. The 0.53 kb DraI/HindII fragment was purified by electrophoresis on an agarose gel. The purified fragment was ligated with the 0.27 kb SpII/HindIII fragment derived from the PCR described above using ligation kit ver. 2 (Takara Shuzo Co.) together with HindIII/EcoRI fragment (5.2 kb) of pSec TagA (Invirogen Co.). *Escherichia coli* DH5 $\alpha$  was transformed using the reaction product. Plasmids were purified by alkali SDS method from the resulting ampicillin resistant transformants and digested with restriction enzymes to select a plasmid containing 0.27 kb and 0.53 kb fragments as inserts in pSec TagA. This plasmid was confirmed to have a sequence encoding the secreted human OBM by sequencing using Tag dye-deoxy-terminator cycle sequencing FS kit (Perkin Elmer Co.). The plasmid was digested with restriction enzymes NheI and Xhol to prepare a fragment (0.8 kb) corresponding to the secreted human OBM cDNA by electrophoresis on an agarose gel. This fragment was inserted into the NheI and Xhol fragment (10.4 kb) of an expression vector pCEP4 (Invirogen Co.) using ligation kit and *Escherichia coli* DH5 $\alpha$  was transformed using the reaction product. Plasmids were purified by alkali-SDS method from the resulting ampicillin resistant transformants and digested with restriction enzymes to select an *Escherichia coli* having the

expression plasmid for secreted-form human OBM (pCEP-shOBM). The *Escherichia coli* containing the pCEPshOBM was cultured and pCEPshOBM was purified using Qiafilter plasmid midi kit (Qiagen Co.).

[0306] (2) Expression of Secreted-Form OBM

[0307] 293-EBNA cells were suspended in IMDM containing 10% FCS (IMDM-10% FCS), added into a 24-well plate coated with collagen (manufactured by Sumitomo Bakelite Co., Ltd.) in a cell density of  $2 \times 10^5$ /2 ml/well and cultured overnight. The cells were transfected with 1  $\mu$ g of pCEPshOBM or pCEP4 using 4  $\mu$ l of Lipofectamine (Gibco Co.). After cultured for two days in 0.5 ml of a serum-free IMDM or IMDM-10% FCS, the culture supernatants were collected. Expression of the secreted human OBM in the culture supernatant was detected as follows. Sodium bicarbonate was added to the culture supernatants to a final concentration of 0.1 M and the mixtures were added to a 96-well plate. The plate was allowed to stand overnight at 4°C, thereby immobilizing human OBM in the culture supernatants on the 96-well plate. The plate was blocked using Block Ace™ (Snow Brand Milk Products Co., Ltd.) solution four-fold diluted with PBS (B-PBS) and allowed to stand for two hours at room temperature. After adding 3-100 ng/ml of OCIF which was diluted with B-PBS to each well, the plate was allowed to stand for two hours at 37°C, followed by wash with PBS containing 0.05% Polysolvate 20(P-PBS). Then, 100  $\mu$ l of a peroxidase-labeled rabbit anti-OCIF polyclonal antibody described in WO 96/26217 which was diluted with B-PBS was added to each well. After allowing to stand for two hours at 37°C, the wells were washed six times with P-PBS. Then, TMB solution (TMB Soluble Reagent, High Sensitivity, Scytek Co.) was added in the amount of 100  $\mu$ l per well and the mixture was allowed to stand at room temperature for about 10 minutes. The reaction was terminated by the addition of 100  $\mu$ l of termination solution (Stopping Reagent, Scytek Co.) to each well. Absorbance at 450 nm for each well was measured by a microplate reader. The results are shown in FIG. 21, which indicates that the absorbance at 450 nm increased according to the concentration of the added OCIF in the plate in which the conditioned medium of the cells transfected with pCEP-shOBM was immobilized. On the other hand, no increase in absorbance was seen in the wells in which the conditioned medium of the cells transfected with vector pCEP4 was immobilized. FIG. 22 shows the results of an experiment wherein the proportion of the conditioned medium used for immobilization was changed within a range of 5-90% in the presence of a constant concentration of OCIF (50 ng/ml). The absorbance at 450 nm increased according to the increase in the proportion of the conditioned medium in the plate wherein the conditioned medium of the cells transfected with pCEPshOBM was immobilized, whereas no such increase in absorbance was seen in the plate wherein the conditioned medium of the cells transfected with vector pCEP4 was immobilized. From these results, it was confirmed that secreted-form human OBM is produced in the conditioned medium of the cells transfected with pCBP-shOBM.

Example 25

[0308] Expression of Thioredoxin-Human OBM Fusion Protein (Trx-hOBM)

[0309] (1) Construction of a Thioredoxin-Human OBM Fusion Protein (Trx-hOBM) Expression Vector

[0310] Ten  $\mu$ l of 10 ExTaq buffer (Takara Shuzo Co.), 8  $\mu$ l of 10 mM dNTP (Takara Shuzo Co.), 77.5  $\mu$ l of sterilized

distilled water, 2  $\mu$ l of an aqueous solution of pUC19hOBM(10 ng/ $\mu$ l), 1  $\mu$ l of primer, mouse OBM #3 (100 pmol/ $\mu$ l, Sequence Table, Sequence ID No. 9), 1  $\mu$ l of primer, hOBM SalR2 (100 pmol/ $\mu$ l, Sequence Table, Sequence ID No. 14), and 0.5  $\mu$ l of ExTaq (5 u/ $\mu$ l) (Takara Shuzo Co.) were mixed and reacted (PCR) in a micro centrifugel tube. After the reaction at 95°C. for 5 minutes, at 50°C. for one second, at 55°C. for one minute, at 74°C. for one second, and at 72°C. for 5 minutes, a cycle reaction consisting of a reaction at 96°C. for one minute, at 50°C. for one second, at 55°C. for one minute, at 74°C. for one second, and at 72°C. for 3 minutes, was repeated 25 times. From the total reaction mixture DNA fragment (750 bp) was purified. The whole amount of purified DNA fragment was digested with restriction enzymes SalII (Takara Shuzo Co.) and BspHI (New England Biolabs Co.), and subjected to electrophoresis on a 1% agarose gel to obtain purified DNA fragment (Fragment 1, about 320 bp). The fragment was dissolved in 20  $\mu$ l of sterilized distilled water. In the same manner, DNA fragment (Fragment 2, about 450 bp) obtained by digesting 4  $\mu$ g of pUC19hOBM with restriction enzymes BamHI, and BspHI (Takara Shuzo Co.) and DNA fragment (Fragment 3, about 3.6 kb), obtained by digesting 2  $\mu$ g of pTrxFus (Invitrogen Co.) with restriction enzymes BamHI, and SalII (Takara Shuzo Co.) were respectively purified and dissolved in 20  $\mu$ l of sterilized distilled water. The QIAEXII gel extraction kit was used for purification of the DNA fragments. Fragments 1-3 were ligated by incubating at 16°C. for 2.5 hours using DNA ligation kit ver.2 (Takara Shuzo Co.). Using the ligation reaction, *Escherichia coli* GI724 (Invirogen Co.) was transformed according to the method described in the Instruction Manual of ThioFusion Expression System (Invirogen Co.). A microorganism strain with plasmid in which the hOBM cDNA fragment is fused in frame to thioredoxin gene was selected from the resulting ampicillin resistant transformants by analysis of DNA restriction map obtained by digestion with restriction enzyme and by determination of DNA sequence. The microorganism strain thus obtained was named GI724/pTrx-hOBM.

[0311] (2) Expression of Trx-hOBM in *Escherichia coli*

[0312] GI724/pTrxhOBM and GI724 containing pTrxFus (GI724/pTrxFus) were respectively cultured six hours with shaking at 30°C. in 2 ml of RMG-Amp medium (0.6%  $\text{Na}_2\text{HPO}_4$ , 0.3%  $\text{KH}_2\text{PO}_4$ , 0.05% NaCl, 0.1%  $\text{NH}_4\text{Cl}$ , 2% casamino acid, 1% glycerol, 1 mM  $\text{MgCl}_2$ , 100  $\mu\text{g}/\text{ml}$  ampicillin, pH 7.4). The broth(0.5 ml) was added to 50 ml of Induction medium (0.6%  $\text{Na}_2\text{HPO}_4$ , 0.3%  $\text{KH}_2\text{PO}_4$ , 0.05% NaCl, 0.1%  $\text{NH}_4\text{Cl}$ , 0.2% casamino acid, 0.5% glucose, 1 mM  $\text{MgCl}_2$ , 100  $\mu\text{g}/\text{ml}$  ampicillin, pH 7.4) and cultured with shaking at 30°C. When  $\text{OD}_{600\text{ nm}}$  reached about 0.5, L-tryptophan was added to a final concentration of 0.1 mg/ml, followed by culturing with shaking at 30°C. for an additional 6 hours. The culture broth was centrifuged at 3000xg to collect the cells, which were then suspended in 12.5 ml of PBS. The suspension was subjected to an ultrasonic generator (Ultrasonics Co.) to disrupt the cells. The disrupt cells were centrifuged at 7000xg for 30 minutes to obtain a supernatant liquid as a soluble protein fraction. Ten  $\mu$ l of this soluble protein fraction was subjected to SDS polyacrylamide (10%) electrophoresis under reducing conditions. As a result, as shown in FIG. 23, a band with a molecular weight of 40,000 which was not detected in the soluble protein fraction of GI724/pTrxFus was found in the soluble protein fraction of GI724/pTrxhOBM. Accordingly,

it was confirmed that a fusion protein (Trx-hOBM) of thioredoxin and human OBM was expressed in *Escherichia coli*.

[0313] (3) Binding Capability of Trx-hOBM to OCIF

[0314] Binding of the expressed Trx-hOBM to OCIF was confirmed according to the following experiment. Anti-thioredoxin antibody (Invirogen Co.) which was diluted 5000-fold with 10 mM sodium hydrogen carbonate solution was added to a 96-well immunoplate (Nunc Co.) in the amount of 100  $\mu$ l per well. After being allowed to stand overnight at 4° C., the liquid in the wells was discarded. Two hundred  $\mu$ l of a solution prepared by diluting Block Ace™ (Snow Brand Milk Products Co., Ltd.) two-fold with PBS (BA-PBS) was added to each well. After being allowed to stand for one hour at room temperature, the wells were washed three times with P-PBS. The soluble protein fractions originating from the above-described GI724/pTrxhOBM or GI724/pTrxFus, each diluted with BA-PBS in various concentrations were added to each well in the amount of 100  $\mu$ l. After being allowed to stand for two hours at room temperature, each well was washed three times with P-PBS and charged with 100  $\mu$ l of OCIF (100 ng/ml) which was diluted with BA-PBS. After being allowed to stand for two hours at room temperature, each well was washed three times with P-PBS and charged with 100  $\mu$ l of peroxidase-labeled anti-OCIF antibody (described in WO 96/26217) which was diluted 2,000-fold with BA-PBS. After being allowed to stand for two hours at room temperature, each well was washed six times with P-PBS and charged with 100  $\mu$ l of TMB solution. After being allowed to stand for about 10 minutes at room temperature, each well was charged with 100  $\mu$ l of termination solution (Stopping Reagent). Absorbance of each well at 450 nm was measured by a microplate reader. The results are shown in FIG. 24. There was no difference in the absorbance between the sample with the soluble protein fraction originating from GI724/pTrxFus added thereto and the sample without the addition of this soluble protein fraction. On the other hand, the absorbance increased in the samples to which the soluble protein fraction originating from GI724/pTrxhOBM was added in proportion to the concentration of the soluble protein fraction. The results of the other experiment wherein the dilution rate of the soluble protein fraction was maintained constant (1%) while adding OCIF diluted with BA-PBS in different concentrations (0-100 ng/ml) are shown in FIG. 25. It can be seen that the absorbance remained low at any concentrations of OCIF in samples using a soluble protein fraction originating from GI724/pTrxFus, whereas the absorbance increased in proportion to the OCIF concentration in the samples to which the soluble protein fraction originating from GI724/pTrxhOBM was added. Based on these results, it was confirmed that Trx-hOBM which is produced from GI724/pTrxhOBM has a capability of binding to OCIF.

[0315] (4) Large-Scale Cultivation of *Escherichia coli* Which Produces Trx-hOBM

[0316] GI724/pTrxhOBM cells were spread on RMG-Amp agar (0.6% Na<sub>2</sub>HPO<sub>4</sub>, 0.3% KH<sub>2</sub>PO<sub>4</sub>, 0.05% NaCl, 0.1% NH<sub>4</sub>Cl, 2% casamino acid, 1.5% agar, pH 7.4) using a platinum transfer 100 p. The cells were cultured overnight at 30° C. The cultured cells were suspended in 10 ml of Induction medium. The suspension was added (5 ml for each) to two 2 1 Erlenmeyer flasks containing 500 ml of

Induction medium and cultured at 30° C. with shaking. When the OD<sub>600 nm</sub> reached about 0.5, L-tryptophan was added to a final concentration of 0.1 mg/ml. Culturing with shaking was continued for six hours at 30° C. The culture broth was centrifuged for 20 minutes at 3000×g to collect the cells, which were suspended in 160 ml of PBS. The suspension was subjected to an ultrasonic generator (Ultrasonics Co.) to disrupt the cells. The supernatant liquid was centrifuged for 30 minutes at 7000×g to obtain a soluble protein fraction.

[0317] (5) Preparation of OCIF-Immobilized Affinity Column

[0318] Two g of TSKgel AF-Tolesyl Toyopal 650 (Tosoh Corp.) and 40 ml of 1.0 M potassium phosphate buffer (pH 7.5) containing 35.0 mg of recombinant OCIF, which was prepared according to the method described in WO 96/26217, were mixed. The mixture was gently shaken overnight at 4° C. to effect a coupling reaction. The reaction mixture was centrifuged to remove the supernatant. To inactivate excess active residues, 40 ml of 0.1 M Tris-HCl buffer (pH 7.5) was added to the precipitated carrier and the mixture was gently shaken at room temperature for one hour. The carrier in a column was washed with 0.1 M glycine-HCl buffer (pH 3.3) containing 0.01% Polysorbate 80 and 0.2 M NaCl and 0.1 M sodium citrate buffer (pH 2.0) containing 0.01% Polysorbate 80 and 0.2 M NaCl. The carrier in the column was equilibrated by charging twice with 10 mM sodium phosphate buffer (pH 7.4) containing 0.01% Polysorbate 80.

[0319] (6) Purification of Trx-hOBM Using OCIF-Immobilized Affinity Column

[0320] Unless otherwise indicated, purification of Trx-hOBM was carried out at 4° C. The above-mentioned OCIF-immobilized affinity carrier (10 ml) and the soluble protein fraction (120 ml) prepared in Example 25(4) were mixed. The mixture was gently shaken overnight at 4° C. in four 50 ml centrifugel tubes using a rotor. An Econocolumn™ (internal diameter: 1.5 cm, length: 15 cm, manufactured by BioRad Co.) was filled with the carrier in the mixture. The column was charged with 300 ml of PBS containing 0.01% Polysorbate 80, 100 ml of 10 mM sodium phosphate buffer (pH 7.0) containing 0.01% Polysorbate 80 and 2.0 M NaCl, and 100 ml of 0.1 M glycine-HCl buffer (pH 3.3) containing 0.01% Polysorbate 80 and 0.2 M NaCl, in that order. Next, proteins adsorbed in the column were eluted with 0.1 M sodium citrate buffer (pH 2.0) containing 0.01% Polysorbate 80 and 0.2 M NaCl. The eluate was collected in 5 ml portions. Each fraction thus collected was immediately neutralized with addition of 10% volume of 2 M Tris buffer solution (pH 8.0). Presence or absence of Trx-hOBM in the eluted fractions was determined according to the method previously described in Example 25(3) (the binding capability to OCIF). The fractions containing Trx-hOBM were collected and purified further.

[0321] (7) Purification of Trx-hOBM by Gel Filtration

[0322] About 25 ml of Trx-hOBM fractions obtained in Example 25(6) was concentrated to about 0.5 ml by centri-

fuging using Centriplus 10 and Centricon 10 (Amicon Co.). This sample was applied to a Superose 12 HR 10/30 column (1.0×30 cm, Pharmacia Co.) previously equilibrated with PBS containing 0.01% Polysorbate 80. For the separation, PBS containing 0.01% Polysorbate 80 was used as a mobile phase at a flow rate of 0.25 ml/min. The eluate from the column was collected in 0.25 ml portions. The Trx-hOBM in the thus-collected fractions was detected by the same method as previously described in the Example 25(3) and SDS-PAGE. Fractions containing purified Trx-hOBM were collected and the protein concentration of Trx-hOBM was determined. The measurement of the protein concentration was carried out using bovine serum albumin as a standard substance using DC-Protein assay kit (BioRad Co.).

#### Example 26

[0323] Osteoclast Formation-Inducing Activity of hOBM  
 [0324] phOBM and pcDL-SR $\alpha$ 296 were respectively transfected into COS-7 cells using Lipofectamine (Gibco Co.). The cells were cultured for one day in DMEM containing 10% FCS, trypsinized, plated on cover slips (15 mm round shape, manufactured by Matsunami Co.) in 24-well plates at 5×10<sup>4</sup> cells per well, and cultured for 2 days. The culture plate was washed once with PBS. The cells were fixed with PBS containing 1% paraformaldehyde at room temperature for 8 minutes. The plate on which the fixed cells were attached was washed 6 times with PBS, then 700  $\mu$ l of mouse spleen cells suspended at 1×10<sup>6</sup>/ml in  $\alpha$ -MEM containing 10<sup>-8</sup> M active-form vitamin D<sub>3</sub>, 10<sup>-7</sup> M dexamethasone, and 10% fetal bovine serum were added to each well. Millicell PCF (Millipore Co.) was set in each well and a suspension of ST2 cells in the above-mentioned culture medium (4×10<sup>4</sup>/ml) were added, 700  $\mu$ l per well, into the Millicell PCF followed by incubation at 37° C. for 6 days. After the culture, the Millicell PCF was removed, the plate was washed once with PBS, and the cells were fixed with acetone-ethanol solution (50:50) for one minute. Then, the cells exhibiting tartaric acid-resistant acid phosphatase activity (TRAP), which is a specific marker for osteoclast, were selectively stained using LEUKOCYTE ACID PHOSPHATASE kit (Sigma Co.). As a result of microscopic observation, TRAP-positive cells were not detected in the wells in which COS-7 cells transfected with pcDL-SR  $\alpha$ 296 were fixed. In contrast, 65±18 (average±standard deviation, n=3) TRAP positive cells were observed in the wells in which COS-7 cells transfected with phOBM were fixed. Moreover, expression of calcitonin receptor was confirmed by the fact that <sup>125</sup>I-labeled salmon calcitonin (Amersham Co.) specifically bound to these TRAP positive cells. Based on these findings, it has been proven that human OBM, which is the protein encoded by cDNA of the present invention, has osteoclast formation-inducing activity.

#### Example 27

[0325] Osteoclast Formation-Inducing Activity of Trx-hOBM and Secreted-Form Human OBM

[0326] Mouse spleen cells were suspended in  $\alpha$ -MEM containing 10<sup>-8</sup>M active-form vitamin D<sub>3</sub>, 10<sup>-7</sup>M dexamethasone, and 10% fetal bovine serum at a concentration of 2×10<sup>6</sup>/ml. The suspension was added to a 24 well plate in the amount of 350  $\mu$ l per well. Each well was then charged with 350  $\mu$ l of a solution prepared by diluting purified Trx-hOBM

with the above-mentioned culture medium (40 ng/ml), 350  $\mu$ l of solution prepared by 10-fold diluting a conditioned medium which was produced by culturing 293-EBNA cells, onto which pCEPshOBM or pCEP4 were transfected, in a culture medium IMDM-10% FCS, with above-mentioned culture medium, or 350  $\mu$ l only of the above-mentioned culture medium. The Millicell PCF (Millipore Co.) was placed on each well, to which 600  $\mu$ l of ST2 cells which were suspended in the above-mentioned culture medium (4×10<sup>4</sup>/ml) were added. After cultured for six days, the Millicell PCF was removed. The plate was washed once with PBS and the cells were fixed by acetone-ethanol solution (50:50) for one minute. Then, the cells exhibiting the activity of tartaric acid resistant acidic phosphatase (TRAP activity) were selectively stained using LEUKOCYTE ACID PHOSPHATASE kit (Sigma Co.). The results of microscopic observation revealed that no cells exhibiting the TRAP activity were detected in the wells to which no Trx-hOBM was added, whereas 115±19 (average±standard deviation, n=3) TRAP-positive cells were observed in the wells to which Trx-hOBM was added. Similarly, while no cells exhibiting TRAP activity were detected in the wells to which conditioned medium of 293-EBNA cells transfected with pCEP4 had been added, 125±23 (average±standard deviation, n=3) TRAP positive cells were observed in the wells to which conditioned medium of 293-EBNA cells transfected with pCEPshOBM had been added. Moreover, expression of calcitonin receptor was confirmed by the fact that <sup>125</sup>I-labeled salmon calcitonin (Amersham Co.) specifically binds to these TRAP positive cells. These results have proven that Trx-hOBM and secreted-form hOBM exhibit osteoclast formation-inducing activity.

#### Example 28

[0327] Preparation of Polyclonal Antibody

[0328] Mouse SOBM or human SOBM used as an immunogen was prepared according to the method described above. Especially, mouse sOBM cDNA (cDNA (Sequence ID No. 18) encoding mouse sOBM (Sequence ID No. 16) which does not have the membrane binding region of the mouse OBM due to absence of the amino acids from the N-terminal down to the 72nd amino acid) or human sOBM cDNA (cDNA (Sequence ID No. 19) encoding human sOBM (Sequence ID No. 17) which does not have the membrane binding region of human OBM due to absence of the amino acids from the N-terminal down to the 71st amino acid) was ligated with a Hind III/EcoRV fragment (5.2 kb) of the expression vector psec TagA (InVitrogen Co.) including the nucleotide sequence encoding a signal peptide of  $\kappa$ -chain of immunoglobulin, together with an EcoRI/PmaCl fragment (0.32 kb) of OBM cDNA, using a ligation kit ver. 2 (Takara Shuzo Co.). *Escherichia coli* DH5 $\alpha$  was transformed with the reaction product. The plasmids obtained from the resulting ampicillin resistant strains were purified by the alkali SDS and digested with an restriction enzyme to select a plasmid with 0.6 Kb and 0.32 kb fragments inserted into pSec TagA. The sequence of this plasmid was identified using the Dyedeoxyterminator Cycle Sequencing FS kit (product of Perkin Elmer Co.). As a result, it was confirmed that this plasmid has a sequence encoding mouse or human sOBM. After plasmid was digested with restriction enzymes NheI/XhoI, a fragment (1.0 kb) corresponding to secretion form OBM cDNA was recovered by agarose gel electrophoresis. This fragment was inserted into an NheI/XhoI

fragment (10.4 kb) of the expression vector pCEP4 (Invitrogen Co.) using a ligation kit. *Escherichia coli* DH5 $\alpha$  was transformed using the reaction product. Plasmids were purified by the alkali SDS from the resulting ampicillin resistant strains. Analyzing these plasmids by digesting with a restriction enzyme, *Escherichia coli* possessing a secretion type OBM expression plasmid (pCEP sOBM) having the objective structure was selected. The *Escherichia coli* strain having the pCEP sOBM was cultured and pCEP sOBM was purified using a Qiafilter plasmid midi kit (Qiagen Co.). Next, 293-EBNA cells were suspended in IMDM (IMDM-10% FCS) containing 10% FCS and plated onto a 24-well plate coated with collagen (product of Sumitomo Bakelite Co., Ltd.) at a cell density of  $2 \times 10^5$  cells/2 ml/well. After culturing overnight, the cells were transformed with 1  $\mu$ g of pCEP sOBM/pCEP4 using 4  $\mu$ l of Lipofectamine (Gibco Co.) and further cultured for two days in 0.5 ml of serum-free IMDM or IMDM-10% FCS. The culture supernatant was recovered. A cell line with high productivity of recombinant mouse soluble OBM (msOBM) or human soluble OBM (hsOBM) was screened as follows. Sodium bicarbonate was added to the culture supernatant which is assumed to contain msOBM or hsOBM to a final concentration of 0.1 M. One hundred  $\mu$ l of the culture supernatant was added to each well in 96-well immunoplates (Nunc Co.) and allowed to stand overnight at 4° C., thereby msOBM or hsOBM in the culture supernatant was immobilized on each well. To each well, 200  $\mu$ l of Block Ace<sup>TM</sup> (Snow Brand Milk Products Co., Ltd.) solution diluted four-fold with PBS (B-PBS) was added and the plates were allowed to stand for two hours at room temperature. After washing each well in the plates three times with PBS (P-PBS) containing 0.1% Polysorbate 20, 100  $\mu$ l of each recombinant OCIF (rOCIF) solution (3-100 ng/ml) diluted serially with P-PBS was added to each well in the plates. The plates were allowed to stand for two hours at 37° C. After washing the plates three times with PBS, 100  $\mu$ l of a peroxidase-labeled anti-OCIF polyclonal antibody (WO 96/26217) diluted with B-PBS was added to each well. After allowing to stand for two hours at 37° C., the wells were washed six times with P-PBS. Then, 100  $\mu$ l of TMB solution (TMB Soluble Reagent, High Sensitivity, ScyTek Co.) was added to each well in the plates and the plates were allowed to stand at room temperature for about 10 minutes, subsequently the reaction was terminated by adding 100  $\mu$ l of a stopping solution (Stopping Reagent, ScyTek Co.) to each well. Absorbance at 450 nm of each well was measured using a microplate reader. It was confirmed that the absorbance increased remarkably in proportion to concentration of the added OCIF in the plates in which msOBM or hsOBM in the culture supernatant of the cell line producing msOBM or hsOBM was immobilized therein.

[0329] The cell line that exhibited a high rate of increase in absorbance was selected as a strain with high productivity. Thus-related 293-EBNA cells with high productivity of msOBM or hsOBM were cultured on a large scale in an IMDM medium containing 5% FCS, using 25 T-flasks (T-225). After the cell reached to confluent, a fresh culture medium was added to each T-225 flask in the amount of 100 ml per flask and cells were cultured for 3-4 days, to collect the culture supernatant. These procedures were repeated four times to obtain 10 L of the culture supernatant containing msOBM or hsOBM. Purified msOBM (10 mg) or hsOBM (12 mg), which shows homogeneous band (molecu-

lar weight: 32 kDa) on SDS-polyacrylamide gel electrophoresis, were obtained from the culture supernatant by means of affinity chromatography on an OCIF-immobilized column and gel filtration chromatography according to the method described in examples 25(6) and 25(7). Each thus-obtained purified preparation was used as an antigen for immunization. Each protein antigen obtained was dissolved in phosphate buffered saline (PBS) to a concentration of 200  $\mu$ g/ml and emulsified with an equivalent volume of Freund's complete adjuvant. One ml of the emulsion was subcutaneously immunized to each of three Japanese white rabbits about once every week. A booster injection was given when the antibody titer reached a peak. Whole blood was collected 10 days thereafter. The serum was diluted two-fold with a binding buffer for protein A sepharose chromatography (BioRad Co.) and applied to a protein A column equilibrated with the same buffer. After washing the column extensively with the same buffer, the anti-sOBM antibody adsorbed to the column was eluted with an elution buffer (BioRad Co.) or 0.1 M glycine-HCl buffer, pH 3.0. To neutralize the eluate immediately, the eluate was fractionated using test tubes containing a small amount of 1.0 M Tris-HCl (pH 8.0). The eluate was dialyzed against PBS overnight at 4° C. The antibody content in the antibody solution was measured by the Lowry method using bovine IgG as a standard protein. Thus, about 10 mg of purified immunoglobulin (IgG) containing the polyclonal antibody of the present invention per 1 ml of rabbit antiserum was obtained.

#### Example 29

[0330] Measurement of OBM and sOBM by ELISA Using Polyclonal Antibody

[0331] A sandwich ELISA was constructed using the rabbit anti-human sOBM polyclonal antibody obtained in Example 28 as the solid phase antibody and enzyme-labeled antibody. Peroxidase (POD)-labeled antibody was prepared according to the method of Ishikawa (Ishikawa et al., J. Imunoassay, Vol. 4, 209-327, 1983).

[0332] The anti-human sOBM polyclonal antibody obtained in the Example 28 was dissolved in a 0.1 M NaHCO<sub>3</sub> to a concentration of 2  $\mu$ g/ml. One hundred  $\mu$ l of the resulting solution was added to each well in 96-well immunoplates (Nunc Co.), which was then allowed to stand at room temperature overnight. Next, 200  $\mu$ l of 50% Block Ace<sup>TM</sup> (Snow Brand Milk Co., Ltd.) was added to each well and the plates were allowed to stand for one hour at room temperature. The wells were washed three times with PBS containing 0.1% Polysorbate 20 (washing buffer).

[0333] Human OBM was expressed according to the method of Example 26 and purified according to the method of Example 2. The purified human OBM and the purified human SOBM prepared in example 28 were serially diluted with the first reaction buffer (0.2 M Tris-HCl buffer, pH 7.2, containing 40% Block Ace and 0.1% Polysorbate 20), respectively, and 100 ml of the diluted solution was added to each well in the plates. The plates were allowed to stand at room temperature for two hours, and washed three times with the above-mentioned washing buffer. Subsequently, 100  $\mu$ l of POD-labeled anti-human sOBM polyclonal antibody diluted 1000-fold with the second reaction buffer (0.1 M Tris-HCl buffer, pH 7.2, containing 25% Block Ace and 0.1% Polysorbate 20) was added to each well in the plates.

After the plates were allowed to stand at room temperature for two hours, each well was washed three times with the washing buffer. Next, 100  $\mu$ l of enzyme substrate solution (TMB, ScyTek Co.) was added to each well in the plates, and the plates were allowed to stand for 10 minutes, followed by the addition of 100  $\mu$ l of a reaction termination solution (Stopping reagent, ScyTek Co.) to stop the enzyme reaction. The absorbance at 450 nm of each well was measured using a microplate reader. The results are shown in FIG. 26. The sandwich ELISA using a rabbit anti-human sOBM polyclonal antibody recognized almost equally human sOBM (molecular weight, about 32 kDa) and human OBM (molecular weight, about 40 kDa), with a measurement sensitivity of about  $12.5 \times 10^{-3}$  pmol/ml (human OBM: about 500 pg/ml, human sOBM: about 400  $\mu$ g/ml). The measurement of mouse sOBM and mouse OBM by ELISA using the rabbit anti-mouse sOBM polyclonal antibody obtained in the Example 28 was able to be carried out in the same manner. It was confirmed that an extremely small amount of mouse sOBM or mouse OBM can be measured with almost the same sensitivity as described above.

[0334] As mentioned above, the anti-human sOBM polyclonal antibody of the present invention prepared in the Example 28 can equally recognize both the human sOBM and human OBM antigens. Therefore, the antibody was named anti-human OBM/sOBM polyclonal antibody. Similarly, the anti-mouse sOBM polyclonal antibody prepared in the Example 28 can equally recognize both the mouse sOBM and mouse OBM antigens. This antibody was therefore named anti-mouse OBM/sOBM polyclonal antibody.

#### Example 30

[0335] Preparation of Monoclonal Antibody

[0336] The purified human sOBM prepared in the Example 28 was used as the antigen for immunization. The purified human sOBM was dissolved in physiological saline solution to a concentration of 10  $\mu$ g/ml and emulsified by mixing with an equivalent volume of Freund's complete adjuvant. The emulsion was intraperitoneally administered to BALB/c mice at a dose of 200  $\mu$ l three times, once a week, to immunize mice. Next, the equivalent volume of the Freund's complete adjuvant was added to a physiological saline solution containing 5  $\mu$ g/ml of human sOBM and the mixture was sufficiently emulsified. This emulsion was injected intraperitoneally to BALB/c mice at a dose of 200  $\mu$ l, once a week for four weeks for immunization. One week after the fourth immunization, 100  $\mu$ l of a physiological saline solution containing 10  $\mu$ g/ml of human sOBM was intravenously administered to the BALB/c mice as a booster. After three days, the spleen was extracted and spleen cells were separated. The spleen cells were fused with mouse myeloma cells, P3x63-Ag8.653 according to a conventional method (Koehler, G. and Milstein, C., *Nature*, 256, 495 (1975)). The suspended fused cells were cultured for 10 days in an HAT medium containing hypoxanthine, aminopterin, and thymidine. After the myeloma cells were dead and hybridomas appeared, the HAT medium was replaced with an aminopterin-free HAT medium, and the cell culture was continued.

#### Example 31

[0337] Selection of Hybridoma and Cloning

[0338] Appearance of hybridomas was recognized 10 days after cell fusion in Example 30. Monoclonal antibodies

recognizing the human sOBM with high affinity and hybridomas producing these antibodies were selected according to the following procedure using the improved solid phase ELISA which is described below. In addition, to select the anti-OBM monoclonal antibody which recognizes both human sOBM and mouse sOBM, mouse sOBM prepared in the Example 27 was used in addition to human sOBM as the antigen for the solid phase ELISA. The human sOBM and mouse sOBM were respectively dissolved in a 0.1 M sodium bicarbonate solution at a concentration of 5  $\mu$ g/ml. Fifty ml of each antigen solution was added to each well in 96-well immunoplates (Nunc Co.). The plates were allowed to stand at 4°C overnight to immobilize the antigens. The antigen solution in each well was discarded. Each well was then filled with 200  $\mu$ l of 50% Block Ace™ (Snow Brand Milk Products Co., Ltd.) and allowed to stand at room temperature for one hour. After each well was washed with phosphate buffered saline solution (PBS-P) containing 0.1% Polysorbate 20, 40  $\mu$ l of calf serum (Hiclone Inc.) was added to each well. Subsequently, 10  $\mu$ l of each hybridoma culture supernatant was added to each well and each well was incubated at room temperature for two hours in the presence of 80% calf serum. The object of the solid phase ELISA in the presence of 80% calf serum is to select a hybridoma which produce an antibody which can detect a very small amount of human sOBM or mouse sOBM even in a solution containing high concentration of protein and in the presence of an immunoreaction interfering substance derived from serum, i.e. a hybridoma which can produce an antibody with a high affinity for human sOBM or mouse sOBM. After the reaction at room temperature for two hours, the plates were washed with PBS-P and subsequently, 50  $\mu$ l of peroxidase-labeled anti-mouse IgG (KPL Co.) diluted 5000-fold with physiological saline solution containing 25% Block Ace was added to each well. After the reaction at room temperature for two hours, the plate was washed three times with PBS-P. After the addition of 50  $\mu$ l of an enzyme substrate solution (TMB, ScyTek Co.) to each well, the reaction was continued at room temperature for five minutes. The enzymatic reaction was stopped by the addition of 50  $\mu$ l of a termination solution (stopping reagent, ScyTek Co.). Hybridomas which produce antibodies recognizing human sOBM or mouse sOBM were selected by measuring absorbance at 450 nm of each well using a microplate reader (Immune Reader NJ2000™, Nippon InterMed Co.). Hybridomas producing antibodies exhibiting particularly high absorbance ( $OD_{450\text{ nm}}$ ) were selected. Cloning of these hybridomas by a limiting dilution method was repeated 3 to 5 times to establish stable hybridomas. Hybridomas exhibiting particularly high antibody productivity were selected among the established antibody-producing hybridoma clones.

#### Example 32

[0339] Production and Purification of Monoclonal Antibody

[0340] The antibody-producing hybridomas obtained in the Example 31, i.e. high affinity antibody-producing hybridoma which recognizes human sOBM and hybridoma which produces an antibody showing cross-reactivity to the mouse sOBM were cultured, respectively. Each hybridoma was implanted intraperitoneally to BALB/c mice ( $1 \times 10^6$  cells per mouse) to which pristan (Aldrich Co.) was administered one week previously. After about 2-3 weeks, accumulated ascites were collected. The monoclonal antibody, which recognizes human sOBM of the present invention or both the human sOBM and mouse sOBM in the ascites, was purified according to the purification method of anti-OBM/

sOBM polyclonal antibodies using a Protein A column described in the Example 28. The purified monoclonal antibody was thus obtained from the ascites by Protein A column chromatography (Pharmacia Co.).

### Example 33

#### [0341] Antigen Specificity of Monoclonal Antibody

[0342] The antigen specificity of a monoclonal antibody which specifically recognizes human sOBM and the monoclonal antibody exhibiting cross-reactivity to both the human sOBM and mouse sOBM was investigated using human sOBM, human intact OBM having a membrane binding region, mouse SOBM, and mouse intact sOBM having a membrane binding region. More than thirty kinds of monoclonal antibody were obtained. The results on several representative antibodies are shown in Table 1. As a result, it was found that most anti-human sOBM monoclonal antibodies which specifically recognize human sOBM also recognize the human intact OBM having a membrane binding region, but not the mouse OBM and the mouse intact OBM which has a membrane binding region. On the other hand, it was found that only a few monoclonal antibodies recognizing both the human sOBM and mouse sOBM were obtained and that these antibodies exhibit cross-reactivity to both the human OBM and mouse OBM. These results show that there is a common antigen-recognizing sites, namely a common epitopes, in both the human OBM and mouse OBM. Based on the fact that the anti-human sOBM monoclonal antibody prepared using the human sOBM as an immune antigen also equally recognizes human OBM having a membrane binding region. Anti-human sOBM monoclonal antibody was named the anti-human OBM/sOBM monoclonal antibody.

TABLE 1

Antibody	Antigen			
	hsOBM	hOBM	MsOBM	mOBM
H-OBM 1	+	+	-	-
H-OBM 2	+	+	-	-
H-OBM 3	+	+	-	-
H-OBM 4	+	+	-	-
H-OBM 5	+	+	-	-
H-OBM 6	+	+	-	-
H-OBM 7	+	+	-	-
H-OBM 8	+	+	-	-
H-OBM 9	+	+	+	+
H-OBM 10	+	+	-	-
H-OBM 11	+	+	-	-
H-OBM 12	+	+	-	-
H-OBM 13	+	+	+	+
H-OBM 14	+	+	-	-

hsOBM: human soluble OBM,

hOBM: human membrane bonding type OBM,

msOBM: mouse soluble OBM,

mOBM: human membrane bonding type OBM

### Example 34

#### [0343] Determination of Class and Subclass of Monoclonal Antibody

[0344] The class and subclass of the monoclonal antibody of the present invention were determined by the immunoglobulin class and subclass analysis kit (Amersham Co.) according to the protocol indicated. The results on repre-

sentative monoclonal antibodies are shown in Table 2. As shown in Table 2, the majority of anti-human OBM/sOBM monoclonal antibodies were IgG<sub>1</sub>, the others were IgG<sub>2a</sub> and IgG<sub>2b</sub>. Light chains for all antibodies were κ chains.

TABLE 2

Antibody	IgG <sub>1</sub>	IgG <sub>2a</sub>	IgG <sub>2b</sub>	IgG <sub>3</sub>	IgA	κ
H-OBM 8	-	+	-	-	-	+
H-OBM 9	+	-	-	-	-	+
H-OBM 10	+	-	-	-	-	+
H-OBM 11	+	-	-	-	-	+
H-OBM 12	-	-	+	-	-	+
H-OBM 13	+	-	-	-	-	+
H-OBM 14	+	-	-	-	-	+

### Example 35

#### [0345] Measurement of the Dissociation Constant (K<sub>d</sub> value) of Monoclonal Antibody

[0346] The dissociation constant of the monoclonal antibody was measured according to a known method (Betrand Friguet et al.: Journal of Immunological Methods, 77, 305-319, 1986). That is, the purified antibody obtained in the Example 32 was diluted with 0.4 M Tris-HCl buffer (a primary buffer, pH 7.4) containing 40% Block Ace and 0.1% Polysorbate 20 to give a concentration of 5 ng/ml. The solution was mixed with an equivalent volume of a diluted solution of purified human soluble OBM (hsOBM) obtained in Example 28 in the primary buffer with a concentration range of 6.25 to 10 μg/ml. The mixture was allowed to stand for 15 hours at 4° C. to bind the hsOBM to the monoclonal antibody. After 15 hours, the antibody not bound to the hsOBM (10 μg/ml, 100 μl/well) was measured using an immobilized solid phase ELISA to calculate the dissociation constant of the monoclonal antibody to the hsOBM. In addition, affinity to msOBM of an antibody, which is a monoclonal antibody for the hsOBM and also exhibits the cross-reactivity to mouse soluble OBM (msOBM), was measured according to the same method except for using msOBM instead of the hsOBM. Dissociation constant of antibodies, which exhibit high affinity to each antigen and are useful for enzymatic immunoassay and binding assay, are shown in Table 3.

TABLE 3

Antibody	Subclass	Antigen	Dissociation constant Kd(M)
H-OBM 1	IgG <sub>1</sub> (κ)	hsOBM	$1 \times 10^{-11} < k_d < 1 \times 10^{-10}$
H-OBM 4	IgG <sub>1</sub> (κ)	hsOBM	$1 \times 10^{-11} < k_d < 1 \times 10^{-10}$
H-OBM 9	IgG <sub>1</sub> (κ)	hsOBM	$1 \times 10^{-9} < k_d < 1 \times 10^{-8}$
H-OBM 9	IgG <sub>1</sub> (κ)	msOBM	$1 \times 10^{-8} < k_d < 1 \times 10^{-7}$

[0347] As a result, the dissociation constants (Kd) of H-OBM 1 and H-OBM 4 which are the antibodies specific to human soluble OBM (hsOBM) were in the order of  $10^{-11}$  M, indicating the high affinity to hsOBM. On the other hand, the Kd value of the antibody H-OBM 9 which recognizes both the hsOBM and mouse soluble OBM (msOBM) was in the order of  $10^{-8}$  M to msOBM and in the order of  $10^{-9}$  M to hsOBM. In addition, the dissociation constant of the other antibody which recognizes both antigens in the Table 1, i.e. the dissociation constant of H-OBM 13 for each antigen,

was the same as that of H-OBM 9, and these two antibodies belong to the same subclass. These findings suggest the possibility that they are the identical antibodies which recognize the same epitope of each antigen.

#### Example 36

**[0348]** Measuring Method of Human OBM and sOBM by Sandwich ELISA Using Anti-Human OBM/sOBM Monoclonal Antibodies

**[0349]** A sandwich ELISA was constructed using the two high affinity monoclonal antibodies obtained in Example 35, H-OBM 1 and H-OBM 4, respectively as a solid phase antibody and an enzyme-labeled antibody. Labeling of the antibody was carried out using a maleimide activated-peroxidase kit (Piers Co.). The antibody, H-OBM 1, was dissolved in a 0.1 M sodium bicarbonate solution to a concentration of 10  $\mu$ g/ml, and 100  $\mu$ l of the solution was added to each well in 96-well immunoplates (Nunc company). After being allowed to stand overnight at 4° C. to immobilize the antibody, the solution was discarded and 300  $\mu$ l of 50% Block Ace™ solution was added to each well in the plates. Each well in the plates was blocked by allowing to stand at room temperature for two hours. After blocking, the plates were washed with phosphate buffered saline containing 0.1% Polysorbate 20 (PBS-P). Human OBM (hOBM) and human soluble OBM (hsOBM) were respectively diluted with 0.4 M Tris-HCl buffer, pH 7.4, containing 40% Block Ace™ (Snow Brand Milk Products Co., Ltd.) and 0.1% Polysorbate 20 (Wako Pure Chemicals Co., Ltd.) (the first reaction buffer) to prepare test samples with various concentrations. These test samples with different concentrations were added to each well in the amount of 100  $\mu$ l per well and reacted to the antibody, H-OBM 1 immobilized on each well by incubating at room temperature for two hours. After two hours, the plates were washed with PBS-P. Next, 100  $\mu$ l of a solution of POD-labeled H-OBM 4 antibody in 0.2 M Tris-HCl buffer, pH 7.4, containing 25% Block Ace™ and 0.1% Polysorbate 20 (the second reaction buffer) was added to each well, followed by further incubating at room temperature for two hours. The plates were then washed with PBS-P and 100  $\mu$ l of an enzyme substrate solution (TMB, ScyTek Co.) was added to each well to start enzyme reaction. The enzyme reaction was terminated by the addition of 100  $\mu$ l of a reaction termination solution (stopping reagent, ScyTek Co.) to each well. The absorbance of each well at 450 nm was measured using a microplate reader. The results are shown in **FIG. 27**.

**[0350]** As a result, it was confirmed that the sandwich ELISA constructed using the two anti-human OBM/sOBM monoclonal antibodies, H-OBM 1 and H-OBM 4 with high affinity for human OBM/sOBM prepared in the Example 35, equally recognizes human OBM and human sOBM, and is able to measure a very small amount of human OBM or human sOBM at a quantitative limit of about  $1.25 \times 10^{-3}$  to  $2.5 \times 10^{-3}$  pmol/ml (about 50-100  $\mu$ g/ml for human OBM with a molecular weight of 40 kDa, about 40-80  $\mu$ g/ml for human sOBM with a molecular weight of 32 kDa). The hybridomas which produce these two anti-human OBM/sOBM monoclonal antibodies, H-OBM 1 and H-OBM 4 were named H-OBM1 and H-OBM4, respectively. The hybridoma producing anti-human OBM/sOBM monoclonal antibody (H-OBM 9) which recognizes mouse OBM and mouse sOBM and also has an osteoclast formation-inhibi-

tory activity was named H-OBM9. These hybridomas were deposited with the National Institute of Bioscience and Human Technology, the Agency of Industrial Science and Technology, on Nov. 5, 1993 with Deposition Nos. FERM BP-6264 (H-OBM 1), FERM BP-6265 (H-OBM 4), and FERM BP-6266 (H-OBM 9).

#### Example 37

**[0351]** Measurement of Mouse OBM and Mouse sOBM Using Anti-Human OBM/sOBM Monoclonal Antibody Which Recognizes Mouse OBM and Mouse sOBM

**[0352]** A sandwich ELISA was constructed using the anti-human OBM/sOBM monoclonal antibody, H-OBM 9, which recognizes mouse OBM and mouse sOBM obtained as an solid phase antibody in the Examples 33 and 35, and the anti-mouse OBM/sOBM polyclonal antibody as an enzyme-labeled antibody obtained in the example 28. The mouse OBM and mouse sOBM were respectively diluted with the first reaction buffer to give various concentrations in the same manner as in the Example 35 and then measured sOBM according to the method described in the Example 36. The results are shown in **FIG. 28**. As a result, it was found that mouse OBM and mouse sOBM can be similarly measured using H-OBM 9 which is the anti-human OBM/sOBM monoclonal antibody recognizing the mouse OBM and mouse sOBM of the present invention. As shown by the result of Example 35, this anti-human OBM/sOBM monoclonal antibody H-OBM 9 has a high dissociation constant relative to the mouse sOBM, namely it has a comparatively low affinity to mouse sOBM. The sensitivity in the measurement of mouse OBM (molecular weight, about 40 kDa) and mouse sOBM (molecular weight, about 32 kDa) by this ELISA assay was about  $25 \times 10^{-3}$  pmol/ml (about 1 ng/ml for mouse OBM and about 0.8 ng/ml for mouse sOBM).

#### Example 38

**[0353]** Osteoclastogenesis-Inhibitory Activity of Anti-OBM/sOBM Antibody

**[0354]** It is known that osteoclast-like cells (OCL) are induced by co-culture of mouse spleen cells and ST2 cells (mouse bone marrow-derived stromal cells; Endocrinology, 125, 1805-1813 (1989)). Capability of the anti-OBM/sOBM antibody to inhibit the OCL formation when added to the co-culture system was studied. Because the mouse OBM is expressed in this co-culture system, a rabbit anti-mouse OBM/sOBM polyclonal antibody which recognizes mouse OBM and an anti-human OBM/sOBM monoclonal antibody (H-OBM 9) which recognizes both human OBM and mouse OBM antigens were used as the antibodies in this example. Seven hundred microliters per well of each anti-OBM antibody diluted serially with  $\alpha$ -MEM containing 10% FCS and 350  $\mu$ l/well of male mouse splenocytes ( $2 \times 10^6$ /ml)suspended in the same medium described above were added to each well in a 24-well plate (Nunc). Next, ST2 cells trypsinized and suspended in the above-mentioned culture medium containing  $4 \times 10^{-8}$  M Vitamin D<sub>3</sub> and  $4 \times 10^{-7}$  M Dexamethazone ( $8 \times 10^4$  cells/ml) were added to each well in the amount of 350  $\mu$ l/well, followed by culturing for six days

at 37° C. After the plates were washed once with PBS, cells in each well were fixed with a mixture of ethanol and acetone (50:50) for one hour at room temperature. The plates were dried in air, and 500  $\mu$ l of substrate solution was added to each well according to the protocol of the LEUKOCYTE ACID PHOSPHATASE kit (Sigma Co.), followed by incubating for 55 minutes at 37° C. Only the cells exhibiting the tartaric acid-resistant acid phosphatase activity (TRAP activity), which is a specific marker for osteoclasts, were stained by this reaction. The plates were washed once with distilled water and dried in air, and the number of TRAP-positive cells was counted. The results are shown in Table 4. As shown in Table 4, both the rabbit anti-mouse OBM/sOBM polyclonal antibody and the anti-human OBM/sOBM monoclonal antibody, H-OBM 9, which recognizes mouse OBM inhibited OCL formation in a dose-dependent manner. These antibodies were found to possess osteoclastogenesis-inhibitory activity like osteoclastogenesis-inhibitory factor, OCIF/OPG, and thus are promising as a therapeutic agent for treating bone metabolism abnormality symptoms.

TABLE 4

Amount of antibody ( $\mu$ g/ml)	Number of TRAP-positive multinucleates	
	Rabbit anti-mouse OBM/sOBM polyclonal antibody	Mouse anti-human OBM/sOBM monoclonal antibody (H-OBM 9)
0	1155 $\pm$ 53	1050 $\pm$ 45
10	510 $\pm$ 24	650 $\pm$ 25
100	10 $\pm$ 3	15 $\pm$ 4

(Average  $\pm$  standard deviation, n = 3)

## Example 39

[0355] Human Osteoclast Formation-Inducing Activity of Trx-OBM

[0356] Mononuclear cells were prepared from whole blood collected from the vein of a healthy adult by density gradient using Histopaque (Sigma Co.) according to the protocol attached thereto. The mononuclear cells were suspended at a cell density of  $1.3 \times 10^6$ /ml in  $\alpha$ -MEM containing  $10^{-7}$  M Dexamethasone, 200 ng/ml macrophage colony stimulating factor (The Green Cross Corp.), 10% fetal bovine serum, and purified Trx-OBM (0-100 ng/ml) obtained in Example 15. The cell suspension was added to each well in 48-well plates in the amount of 300  $\mu$ l per well, and the cells were cultured at 37° C. for three days. After the culture broth was replaced with the above-mentioned culture medium, the cells were cultured at 37° C. for four days. The cultured cells having tartaric acid resistant acid phosphatase activity (TRAP activity) were selectively stained according to the method described in Example 5. The number of stained multinucleates was measured by microscope observation. The results are shown in FIG. 29. It was confirmed that TRAP-positive multinucleates were induced in a dose dependent manner by addition of Trx-OBM, while no TRAP-positive cells were detected in the wells to which Trx-OBM was not added. Moreover, these TRAP-positive multinucleates were found positive to vitronectin receptor which is a marker for osteoclasts. Furthermore, when similar cell culture was carried out on ivory slices placed on each well in a 48-well plate, pit formation was observed on the ivory slices only in the presence of Trx-OBM. Based on

these findings, Trx-OBM was formed to have the activity of inducing human osteoclast formation.

## Example 40

[0357] Inhibition of Bone Resorbing Activity by Anti-OBM/sOBM Antibody

[0358] [ $^{45}$ Ca]-CaCl<sub>2</sub> solution (Amersham Co.) was subcutaneously injected to ddY mouse (Japan SLC Co.) in the 15th day of pregnancy at a dose of 25  $\mu$ Ci per mouse to label the bone of the fetus with  $^{45}$ Ca. Next day, the mouse was sacrificed to obtain the fetus. The forefoot of the fetus was drawn and the skin and muscle were removed to obtain the long bones. The cartilage was removed to obtain the shafts of long bones. The shafts of long bones were floated one by one in 0.5 ml of culture medium (BGJb medium (GIBCO BRL company) containing a 0.2% bovine serum albumin (Sigma Co.)) in each well in 24-well plates, and cultured for 24 hours at 37° C. in 5% CO<sub>2</sub>. After the pre-cultivation, the bones were transferred into various fresh culture media (0.5

ml), each containing one of different bone resorbing factors (vitamin D<sub>3</sub>, prostaglandins E<sub>2</sub>, parathyroid hormone, interleukin 1 $\alpha$ ), and normal rabbit IgG (100  $\mu$ g/ml; as a control), or the rabbit anti-OBM/sOBM polyclonal antibody prepared in Example 28, followed by further cultivation for 72 hours. After the cultivation, the long bones were placed in 0.5 ml of an aqueous solution of 5% trichloroacetic acid (Wako Pure Chemicals Co., Ltd.), and allowed to stand at room temperature for more than 3 hours to decalcify. Five ml of a scintillator (AQUASOL-2, PACKARD Co.) was added to the culture broth and the extract of the trichloroacetic acid solution (each 0.5 ml) to measure the radioactivity of  $^{45}$ Ca, whereby the ratio of the  $^{45}$ Ca which was liberated into the culture broth by bone resorption was calculated. The results are shown in FIGS. 30 to 33. As a result, vitamin D<sub>3</sub> ( $10^{-8}$  M) was found to increase the bone resorbing activity, but the rabbit anti-OBM/sOBM polyclonal antibody suppressed the bone resorption stimulated by vitamin D<sub>3</sub> in a concentration-dependent manner, completely inhibiting the increased bone resorption at a concentration of 100  $\mu$ g/ml (FIG. 30). Prostaglandins E<sub>2</sub> ( $10^{-6}$  M) and parathyroid hormone (100 ng/ml) also increased the bone resorbing activity. However, addition of 100  $\mu$ g/ml of the rabbit anti-OBM/sOBM polyclonal antibody almost completely inhibited the bone resorption stimulated by prostaglandins E<sub>2</sub> and parathyroid hormone (FIGS. 31 and 32). On the other hand, normal rabbit IgG (100  $\mu$ g/ml), which was used as a positive control, did not affect the bone resorbing activity induced by prostaglandins E<sub>2</sub> and parathyroid hormone. Bone resorption was also increased by interleukin 1 $\alpha$  (10 ng/ml), but significantly inhibited by the addition of rabbit anti-OBM/sOBM poly-

clonal antibody (100  $\mu$ g/ml) (FIG. 23). Based on these results, it is clear that the antibody of the present invention is a superior substance as a bone resorption inhibitor. The results obtained by similar experiment using H-OBM 9 which is a mouse anti-human OBM/sOBM antibody, confirmed that this antibody exhibits an almost equivalent bone resorption-inhibitory effect as the rabbit anti-OBM/sOBM polyclonal antibody.

[0359] Industrial Applicability

[0360] The present invention provides a novel protein that specifically binds to osteoclastogenesis-inhibitory factor (OCIF), a process for preparing the protein, a screening method for a substance which controls expression of this protein using this protein, a screening method for a substance which inhibits or modulates the activity of this protein, a screening method for the receptor which transmits the activity of this protein by binding thereto, a pharmaceutical composition which contains the substance obtained by these screening methods, an antibody for the said protein, and an agent for treating bone metabolism abnormality using the antibody.

[0361] Moreover, the present invention provides a DNA encoding a novel protein (OCIF-binding molecule) which binds to osteoclastogenesis-inhibitory factor (OCIF), a protein which possesses an amino acid sequence encoded by the DNA, a method for preparing the protein specifically binding to OCIF using said DNA by a genetic engineering technique, and an agent comprising said protein for treating bone metabolism acatostasia. Furthermore, the present invention provides a screening method for a substance which controls expression of the OCIF-binding molecule, a screening method for a substance which inhibits or modulates the activity of the OCIF-binding molecule by binding thereto, a screening method for the receptor which transmits the activity of the OCIF-binding molecule by binding thereto, and a pharmaceutical composition which contains the substance obtained by these screening methods.

[0362] Still further, the present invention provides a DNA encoding a novel human protein capable of binding to osteoclastogenesis-inhibitory factor, OCIF (human OCIF-binding molecule, human OBM), a protein containing an amino acid sequence encoded by the DNA, a process for preparing a protein having characteristics of specifically binding to OCIF and exhibiting a biological activity to support and promote the osteoclast differentiation and maturation by means of genetic engineering technique, and an agent for treating bone metabolism abnormality using the protein. Furthermore, the present invention provides a screening method for a substance which controls expression of the OCIF-binding molecule, a screening method for a substance which inhibits or modulates the activity of the OCIF-binding molecule by binding thereto, a screening method for the receptor which transmits the biological activity of the OCIF-binding molecule by binding thereto, a pharmaceutical composition which contains the substance obtained by these screening methods, an antibody to human OCIF-binding protein, and an agent for preventing and/or treating bone metabolism abnormality symptoms using the antibody.

[0363] In addition, the present invention provides antibodies which recognize both antigens (anti-OBM/sOBM antibodies), one is a membrane-bound protein which specifically binds to OCIF (OCIF binding molecule; OBM) and the other a soluble OBM (sOBM) which does not have a membrane binding region, a process for preparing the antigen, a method for measuring the OBM and SOBM using these antibodies, and an agent for preventing and/or treating bone metabolism abnormality symptoms using the antibody as an effective component.

[0364] The protein and antibody prepared by the process of the present invention are useful as medicines and/or reagents for research and test purposes.

[0365] Description of Deposited Microorganisms

[0366] (1) Name and address of the depository organization to which microorganism was deposited

[0367] Agency of Industrial Science and Technology  
1-3, Higashi 1-Chome, Tsukuba-shi, Ibaraki-ken,  
Japan (postal code 305)

[0368] Date of deposition to the depository organization  
May 23, 1997

[0369] The deposition number FERM BP-5953

[0370] (2) Name and address of the depository organization to which microorganism was deposited

[0371] Agency of Industrial Science and Technology  
1-3, Higashi 1-Chome, Tsukuba-shi, Ibaraki-ken,  
Japan (postal code 305)

[0372] Date of deposition to the depository organization  
Aug. 13, 1997

[0373] The deposition number FERM BP-6058

[0374] (3) Name and address of the depository organization to which microorganism was deposited

[0375] Agency of Industrial Science and Technology  
1-3, Higashi 1-Chome, Tsukuba-shi, Ibaraki-ken,  
Japan (postal code 305)

[0376] Date of deposition to the depository organization  
Nov. 5, 1997 (Original deposition date)

[0377] The deposition number FERM BP-6264

[0378] (4) Name and address of the depository organization to which microorganism was deposited

[0379] Agency of Industrial Science and Technology  
1-3, Higashi 1-Chome, Tsukuba-shi, Ibaraki-ken,  
Japan (postal code 305)

[0380] Date of deposition to the depository organization  
Nov. 5, 1997 (Original deposition date)

[0381] The deposition number FEPM BP-6265

[0382] (5) Name and address of the depository organization to which microorganism was deposited

[0383] Agency of Industrial Science and Technology  
1-3, Higashi 1-Chome, Tsukuba-shi, Ibaraki-ken,  
Japan (postal code 305)

[0384] Date of deposition to the depository organization Nov. 5, 1997 (Original deposition date)

[0385] The deposition number FERM BP-6266

[0386] Sequence ID No. 1

[0387] Sequence Length: 316

[0388] Sequence Type: amino acid

[0389] Strandedness: 1

[0390] Topology: linear

[0391] Molecule Type: protein

[0392] Sequence:

```

Met Arg Arg Ala Ser Arg Asp Tyr Gly Lys Tyr Leu Arg Ser Ser
 1           5           10          15

Glu Glu Met Gly Ser Gly Pro Gly Val Pro His Glu Gly Pro Leu
20          25          30

His Pro Ala Pro Ser Ala Pro Ala Pro Pro Pro Ala Ala
35          40          45

Ser Arg Ser Met Phe Leu Ala Leu Leu Gly Leu Gly Leu Gly Gln
50          55          60

Val Val Cys Ser Ile Ala Leu Phe Leu Tyr Phe Arg Ala Gln Met
65          70          75

Asp Pro Asn Arg Ile Ser Glu Asp Ser Thr His Cys Phe Tyr Arg
80          85          90

Ile Leu Arg Leu His Glu Asn Ala Gly Leu Gln Asp Ser Thr Leu
95          100         105

Glu Ser Glu Asp Thr Leu Pro Asp Ser Cys Arg Arg Met Lys Gln
110         115         120

Ala Phe Gln Gly Ala Val Gln Lys GIu Leu Gln His Ile Val Gly
125         130         135

Pro Gln Arg Phe Ser Gly Ala Pro Ala Met Met Glu Gly Ser Trp
140         145         150

Leu Asp Val Ala Gln Arg Gly Lys Pro Glu Ala Gln Pro Phe Ala
155         160         165

His Leu Thr Ile Asn Ala Ala Ser Ile Pro Ser Gly Ser His Lys
170         175         180

Val Thr Leu Ser Ser Trp Tyr His Asp Arg GIy Trp Ala Lys Ile
185         190         195

Ser Asn Met Thr Leu Ser Asn Gly Lys Leu Arg Val Asn Gln Asp
200         205         210

Gly Phe Tyr Tyr Leu Tyr Ala Asn Ile Cys Phe Arg His His Glu
215         220         225

Thr Ser Gly Ser Val Pro Thr Asp Tyr Leu Gln Leu Met Val Tyr
230         235         240

Val Val Lys Thr Ser Ile Lys Ile Pro Ser Ser His Asn Leu Met
245         250         255

Lys Gly Gly Ser Thr Lys Asn Trp Ser Gly Asn Ser Glu Phe His
260         265         270

Phe Tyr Ser Ile Asn Val Gly Gly Phe Phe Lys Leu Arg Ala Gly
275         280         285

Glu Glu Ile Ser Ile Gln Val Ser Asn Pro Ser Leu Leu Asp Pro
290         295         300

Asp Gln Asp Ala Thr Tyr Phe Gly Ala Phe Lys Val Gln Asp Ile
305         310         315

Asp
316

```

[0393] Sequence ID No. 2	[0406] Sequence:
[0394] Sequence Length: 1538	[0407] AAACGAAAAA AACCAGAAAG G 21
[0395] Sequence Type: nucleic acid	[0408] Sequence ID No. 4
[0396] Strandedness: 1	[0409] Sequence Length: 17
[0397] Topology: linear	[0410] Sequence Type: nucleic acid
[0398] Molecule Type: cDNA to mRNA	[0411] Strandedness: 1
[0399] Sequence:	[0412] Topology: linear

GCCAGGACCT CTGTGAACCG GTCGGCGCGG GGGCCGCCTG GCCGGGAGTC TGCTCGCGG 60  
 TGGGTGGCGG AGGAAGGGAG AGAACGATCG CGGAGCACCG CGCCCGAACT CGGGCGCGG 120  
 CGCCATGCGC CGGGCCAGCC GAGACTACGG CAAGTACCTG CGCAGCTCGG AGCAGATGCG 180  
 CAGCGGCCCC GGCGTCCCAC ACGAGGGTCC GCTGCACCCC GCGCCTCTG CACCGGCTCC 240  
 GGCGCCGCCA CCCGCCGCCT CCCGCTCCAT GTTCCTGGCC CTCCCTGGGC TGGGACTGGG 300  
 CCAGGTGGTC TGCAGCATCG CTCTGTTCTC GTACTTTCGA GCGCAGATGG ATCCTAACAG 360  
 AATATCAGAA GACAGCACTC ACTGCTTTA TAGAATCCTG AGACTCCATG AAAACGCAGG 420  
 TTTGCAGGAC TCGACTCTGG AGAGTGAAGA CACACTACCT GACTCCTGCA GGAGGATGAA 480  
 ACAAGCCTTT CAGGCGGCCG TGCAGAAGGA ACTGCAACAC ATTGTGGGGC CACAGCGCTT 540  
 CTCAGGAGCT CCAGCTATGA TGGAAGGCTC ATGGTTGGAT GTGGCCAGC GAGGCAAGCC 600  
 TGAGGCCAG CCATTTGCAC ACCTCACCAT CAATGCTGCC AGCATCCCAT CGCGTTCCCA 660  
 TAAAGTCACT CTGTCCTCTT GGTACCACGA TCGAGGCTGT GCCAAGATCT CTAACATGAC 720  
 GTTAAGCAAC GGAAAATAA CGCTTAACCA AGATGCCTTC TATTACCTGT ACGCCAACAT 780  
 TTGCTTCGG CATCATGAAA CATCGGAAAG CGTACCTACA GACTATCTTC AGGTGATGGT 840  
 GTATGTCGTT AAAACCAGCA TCAAAATCCC AAGTTCTCAT AACCTGATCA AAGGAGGGAG 900  
 CACGAAAAC TGGTCGGGC ATTCTGAATT CCACCTTTAT TCCATAAATG TTGGGGCATT 960  
 TTTCAAGCTC CGAGCTGGTG AAGAAATTAG CATTCAAGGTG TCCAACCTT CCCTGCTGGA 1020  
 TCCCGATCAA GATGCGACGT ACTTTGGGC TTTCAAAGTT CAGGACATAG ACTGAGACTC 1080  
 ATTCGTGGA ACATTAGCAT GGATGTCTTA GATGTTGGA AACTTCTAA AAAATGGATG 1140  
 ATGTCTATAC ATGTGTAAGA CTACTAAGAG ACATGGCCA CGGTGTATGA AACTCACAGC 1200  
 CCTCTCTCTT GAGCCTGTAC AGGTTGTGA TATGTAAGT CCATAGGTGA TGTTAGATTG 1260  
 ATCGTGATTA CACAACGTT TTACAATTGTT GTAATGATT CCTAGAATTG AACCAGATTG 1320  
 GGAGAGGTAT TCCGATGCTT ATGAAAACT TACACGTGAG CTATGGAAGG GGGTCACAGT 1380  
 CTCTGGGTCT AACCCCTGGA CATGTGCCAC TGAGAACCTT GAAATTAAGA GGATGCCATG 1440  
 TCATTGAAA GAAATGATAG TGTGAAGGGT TAAGTTCTTT TGAATTGTTA CATTGCGCTG 1500  
 GGACCTGCAA ATAAGTTCTT TTTTCTAAT GAGGAGAG 1538

[0400] Sequence ID No. 3	[0413] Molecule Type: other nucleic acid (synthetic DNA)
[0401] Sequence Length: 21	[0414] Sequence:
[0402] Sequence Type: nucleic acid	[0415] GTAAAACGAC GGCCAGT 17
[0403] Strandedness: 1	[0416] Sequence ID No. 5
[0404] Topology: linear	[0417] Sequence Length: 17
[0405] Molecule Type: other nucleic acid (synthetic DNA)	[0418] Sequence Type: nucleic acid

[0419] Strandedness: 1  
[0420] Topology: linear  
[0421] Molecule Type: other nucleic acid (synthetic DNA)  
[0422] Sequence:  
[0423] CAGGAAACAG CTATGAC 17  
[0424] Sequence ID No. 6  
[0425] Sequence Length: 22  
[0426] Sequence Type: nucleic acid  
[0427] Strandedness: 1  
[0428] Topology: linear  
[0429] Molecule Type: other nucleic acid (synthetic DNA)  
[0430] Sequence:  
[0431] AAGCCCCAAA GTACGTCGCA TC 22  
[0432] Sequence ID No. 7  
[0433] Sequence Length: 26  
[0434] Sequence Type: nucleic acid  
[0435] Strandedness: 1  
[0436] Topology: linear  
[0437] Molecule Type: other nucleic acid (synthetic DNA)  
[0438] Sequence:  
[0439] CGAAGCTTTC GAGCGCAGAT GGATCC  
26  
[0440] Sequence ID No. 8  
[0441] Sequence Length: 27  
[0442] Sequence Type: nucleic acid  
[0443] Strandedness: 1  
[0444] Topology: linear  
[0445] Molecule Type: other nucleic acid (synthetic DNA)  
[0446] Sequence:  
[0447] CCTCTAGAGT CTATGTCCTG AAGTTTG  
27  
[0448] Sequence ID No. 9  
[0449] Sequence Length: 20  
[0450] Sequence Type: nucleic acid  
[0451] Strandedness: 1  
[0452] Topology: linear  
[0453] Molecule Type: other nucleic acid (synthetic DNA)  
[0454] Sequence:  
[0455] ATCAGAAGAC AGCACTCACT 20  
[0456] Sequence ID No. 10  
[0457] Sequence Length: 33  
[0458] Sequence Type: nucleic acid  
[0459] Strandedness: 1  
[0460] Topology: linear  
[0461] Molecule Type: other nucleic acid (synthetic DNA)  
[0462] Sequence:  
[0463] GGGGTCGACC TAGGACATCC  
ATGCTAATGT TCC 33  
[0464] Sequence ID No. 11  
[0465] Sequence Length: 317  
[0466] Sequence Type: amino acid  
[0467] Strandedness: 1  
[0468] Topology: linear  
[0469] Molecule Type: protein  
[0470] Sequence:

Met Arg Arg Ala Ser Arg Asp Tyr Thr Lys Tyr Leu Arg Gly Ser			
Glu	5	10	15
Glu Met Gly Gly Pro Gly Ala Pro His Glu Gly Pro Leu His			
Ala	20	25	30
Pro Pro Pro Pro Ala Pro His Gln Pro Pro Ala Ala Ser Arg Ser			
Met	35	40	45
Phe Val Ala Leu Leu Gly Leu Gly Leu Gly Gln Val Val Cys Ser			
Val	50	55	60
Ala Leu Phe Phe Tyr Phe Arg Ala Gln Met Asp Pro Asn Arg Ile			
Ser	65	70	75
Glu Asp Gly Thr His Cys Ile Tyr Arg Ile Leu Arg Leu His Glu			
Asn	85	90	95

**-continued**

Ala Asp Phe Gln Asp Thr Thr Leu Glu Ser Gln Asp Thr Lys Leu  
 Ile 100 105 110  
 Pro Asp Ser Cys Arg Arg Ile Lys Gln Ala Phe Gln Gly Ala Val  
 Gln 115 120 125  
 Lys Glu Leu Gln His Ile Val Gly Ser Gln His Ile Arg Ala Glu  
 Lys 130 135 140  
 Ala Met Val Asp Gly Ser Trp Leu Asp Leu Ala Lys Arg Ser Lys  
 Leu 145 150 155 160  
 Glu Ala Gln Pro Phe Ala His Leu Thr Ile Asn Ala Thr Asp Ile  
 Pro 165 170 175  
 Ser Gly Ser His Lys Val Ser Leu Ser Ser Trp Tyr His Asp Arg  
 Gly 180 185 190  
 Trp Ala Lys Ile Ser Asn Met Thr Phe Ser Asn Gly Lys Leu Ile  
 Val 195 200 205  
 Asn Gln Asp Gly Phe Tyr Tyr Leu Tyr Ala Asn Ile Cys Phe Arg  
 His 210 215 220  
 His Glu Thr Ser Gly Asp Leu Ala Thr Glu Tyr Leu Gln Leu Met  
 Val 225 230 235 240  
 Tyr Val Thr Lys Thr Ser Ile Lys Ile Pro Ser Ser His Thr Leu  
 Met 245 250 255  
 Lys Gly Gly Ser Thr Lys Tyr Trp Ser Gly Asn Ser Glu Phe His  
 Phe 260 265 270  
 Tyr Ser Ile Asn Val Gly Phe Phe Lys Leu Arg Ser Gly Glu  
 Glu 275 280 285  
 Ile Ser Ile GIu Val Ser Asn Pro Ser Leu Leu Asp Pro Asp Gln  
 Asp 290 295 300  
 Ala Thr Tyr Phe Gly Ala Phe Lys Val Arg Asp Ile Asp  
 305 310 315

[0471] Sequence ID No. 12

[0472] Sequence Length: 954

[0473] Sequence Type: nucleic acid

[0474] Strandedness: 1

[0475] Topology: linear

[0476] Molecule Type: cDNA to mRNA

[0477] Sequence:

ATGCGCCGCG CCAGCAGAGA CTACACCAAG TACCTGCGTG GCTCGGAGGA GATGGGGGCC 60  
 GGCCCCGGAG CCCCCGACGA GGGCCCCCTG CACGCCCGC CGCCGCCTGC GCGCACCGAG 120  
 CCCCTGCCC CCTCCCGCTC CATGTTCGTG GCCCTCCTGG GGCTGGGGCT GGCCCAGGTT 180

-continued

GTCTGCAGCG TCGCCCTGTT CTTCTATTTC AGAGCGCAGA TGCATCTAA TAGAATATCA 240  
 GAAGATGGCA CTCACTGCAT TTATAGAATT TTGAGACTCC ATGAAAATGC AGATTTCAA 300  
 GACACAACTC TGGAGAGTCAGA AGATACAAAAA TTAATACCTG ATTCACTGTAG GAGAATTAAA 360  
 CAGCCCTTTC AAGGAGCTGT GCAAAACGAA TTACAACATA TCGTTGGATC ACAGCACATC 420  
 AGAGCAGAGA AAGCGATGGT GGATGGCTCA TGGTTAGATC TGGCCAAGAG GAGCAAGCTT 480  
 GAAGCTCAGC CTTTGCTCA TCTCACTATT AATGCCACCG ACATCCCATC TGGTTCCCAT 540  
 AAAGTGAGTC TGTCCCTTTG GTACCATGAT CGCGTTGGG CCAAGATCTC CAACATGACT 600  
 TTTAGCAATG CAAAACATAAGTAAATCAG GATGGCTTTT ATTACCTGTA TGCCAAACATT 660  
 TGCTTTCGAC ATCATGAAAC TTCAGGAGAC CTAGCTACAG AGTATCTTCA ACTAATGGTG 720  
 TACGTCACTA AAACCCAGCAT CAAAATCCCA AGTTCTCATA CCCTGATGAA AGGAGGAAGC 780  
 ACCAAGTATT GGTCAAGGGAA TTCTGAATTC CATTTCATT CCATAAACGT TGGTGGATTT 840  
 TTTAAGTTAC GGTCTGGAGA GGAAATCAGC ATCGAGGTCT CCAACCCCTC CTTACTGGAT 900  
 CGGGATCAGG ATGCAACATA CTTTGGGCT TTTAAAGTTG GAGATATAGA TTGA 954

[0478] Sequence ID No. 13

[0479] Sequence Length: 27

[0480] Sequence Type: nucleic acid

[0481] Strandedness: 1

[0482] Topology: linear

[0483] Molecule Type: other nucleic acid (synthetic DNA)

[0484] Sequence:

[0485] GGC GTACGCA GAGCGCAGAT  
 GGATCCT 27

[0486] Sequence ID No. 14

[0487] Sequence Length: 34

[0488] Sequence Type: nucleic acid

[0489] Strandedness: 1

[0490] Topology: linear

[0491] Molecule Type: other nucleic acid (synthetic DNA)

[0492] Sequence:

[0493] GGGGTCGACC ATCCAGGAAA TATCAT-AACA CTCC 34

[0494] Sequence ID No. 15

[0495] Sequence Length: 951

[0496] Sequence Type: nucleic acid

[0497] Strandedness: 1

[0498] Topology: linear

[0499] Molecule Type: cDNA to mRNA

[0500] Sequence:

ATGCGCCGGG CCAGCCGAGA CTACGGCAAG TACCTGCGCA GCTCGGAGGA GATGGGCAGC 60  
 GGCCCCCGCG TCCACACGA GGGTCCGCTG CACCCCGCGC CTTCTGCACC GGCTCCGGCG 120  
 CCGCCACCCCG CGCCCTCCCG CTCCATGTTC CTGGCCCTCC TGGGGCTGCG ACTGGGCCAG 180  
 GTGGTCTGCA GCATCGCTCT GTTCCGTAC TTTGAGCGC AGATGGATCC TAACAGAATA 240  
 TCAGAAGACA GCACACTCGT CTTTATAGA ATCCTGAGAC TCCATGAAA CGCAGGTTTG 300  
 CAGGACTCGA CTCTGGAGAG TGAAGACACA CTACCTGACT CCTGCAGGAG GATGAAACAA 360  
 GCCTTCAGG GGGCCGTGCA GAAGGAACTG CAACACATTG TGGGGCCACA GCGCTTCTCA 420  
 GGAGCTCCAG CTATGATGGA AGGCTCATGG TTGCATGTGG CCCAGCGAGG CAAGCCTGAG 480  
 GCCCAGCCAT TTGCACACCT CACCATCAAT GCTGCCAGCA TCCCACGGG TTCCCATAAA 540  
 GTCACCTGTG CCTCTTGGTA CCACGATCGA GGCTGGGCCA AGATCTCTAA CATGACGTTA 600

**-continued**

ACCAACGGAA AACTAAGGGT TAACCAAGAT GGCTTCTATT ACCTGTACGC CAACATTGC 660  
 TTTCGGCATC ATGAAACATC GGGAAAGCGTA CCTACAGACT ATCITCAGCT GATGGTGTAT 720  
 GTCGTTAAAA CCAGCATCAA AATCCAAGT TCTCATAACC TGATGAAAGG AGGGAGCACG 780  
 AAAAACTAAA CGGGCAATTG TGAATTCCAC TTTTATTCCA TAAATGTTGG GGGATTTTC 840  
 AAGCTCCGAG CTGGTGAAGA AATTAGCATT CAGGTGTCCA ACCCTCCCT GCTGCATCCG 900  
 GATCAAGATG CGACGTACTT TGGGGCTTTC AAAGTTCAAG ACATAGACTG A 951

[0501] Sequence ID No. 16

[0508] Sequence ID No. 17

[0502] Sequence Length: 244

[0509] Sequence Length: 246

[0503] Sequence Type: amino acid

[0510] Sequence Type: amino acid

[0504] Strandedness: 1

[0511] Strandedness: 1

[0505] Topology: linear

[0512] Topology: linear

[0506] Molecule Type: protein

[0513] Molecule Type: protein

[0507] Sequence:

Ala Gln Met Asp Pro Asn Arg Ile Ser Glu Asp Ser Thr His Cys Phe  
 1 5 10 15

Tyr Arg Ile Leu Arg Leu His Glu Asn Ala Gly Leu Gln Asp Ser Thr  
 20 25 30

Leu Glu Ser Glu Asp Thr Leu Pro Asp Ser Cys Arg Arg Met Lys Gln  
 35 40 45

Ala Phe Gln Gly Ala Val Gln Lys Glu Leu Gln His Ile Val Gly Pro  
 50 55 60

Gln Arg Phe Ser Gly Ala Pro Ala Met Met Glu Gly Ser Trp Leu Asp  
 65 70 75 80

Val Ala Gln Arg Gly Lys Pro Glu Ala Gln Pro Phe Ala His Leu Thr  
 85 90 95

Ile Asn Ala Ala Ser Ile Pro Ser Gly Ser His Lys Val Thr Leu Ser  
 100 105 110

Ser Trp Tyr His Asp Arg Gly Trp Ala Lys Ile Ser Asn Met Thr Leu  
 115 120 125

Ser Asn Gly Lys Leu Arg Val Asn Gln Asp Gly Phe Tyr Tyr Leu Tyr  
 130 135 140

Ala Asn Ile Cys Phe Arg His His Glu Thr Ser Gly Ser Val Pro Thr  
 145 150 155 160

Asp Tyr Leu Gln Leu Met Val Tyr Val Val Lys Thr Ser Ile Lys Ile  
 165 170 175

Pro Ser Ser His Asn Leu Met Lys Gly Gly Ser Thr Lys Asn Trp Ser  
 180 185 190

Gly Asn Ser Glu Phe His Phe Tyr Ser Ile Asn Val Gly Gly Phe Phe  
 195 200 205

Lys Leu Arg Ala Gly Glu Ile Ser Ile Gln Val Ser Asn Pro Ser  
 210 215 220

Leu Leu Asp Pro Asp Gln Asp Ala Thr Tyr Phe Gly Ala Phe Lys Val  
 225 230 235 240

Gln Asp Ile Asp

[0514] Sequence:

Ala Gln Met Asp Pro Asn Arg Ile Ser Glu Asp Gly Thr His Cys Ile  
 1 5 10 15  
 Tyr Arg Ile Leu Arg Leu His Glu Asn Ala Asp Phe Gln Asp Thr Thr  
 20 25 30  
 Leu Glu Ser Gln Asp Thr Lys Leu Ile Pro Asp Ser Cys Arg Arg Ile  
 35 40 45  
 Lys Gln Ala Phe Gln Gly Ala Val Gln Lys Glu Leu Gln His Ile Val  
 50 55 60  
 Gly Ser Gln His Ile Arg Ala Glu Lys Ala Met Val Asp Gly Ser Trp  
 65 70 75 80  
 Leu Asp Leu Ala Lys Arg Ser Lys Leu Glu Ala Gln Pro Phe Ala His  
 85 90 95  
 Leu Thr Ile Asn Ala Thr Asp Ile Pro Ser Gly Ser His Lys Val Ser  
 100 105 110  
 Leu Ser Ser Trp Tyr His Asp Arg Gly Trp Ala Lys Ile Ser Asn Met  
 115 120 125  
 Thr Phe Ser Asn Gly Lys Leu Ile Val Asn Gln Asp Gly Phe Tyr Tyr  
 130 135 140  
 Leu Tyr Ala Asn Ile Cys Phe Arg His His Glu Thr Ser Gly Asp Leu  
 145 150 155 160  
 Ala Thr Glu Tyr Leu Gln Leu Met Val Tyr Val Thr Lys Thr Ser Ile  
 165 170 175  
 Lys Ile Pro Ser Ser His Thr Leu Met Lys Gly Gly Ser Thr Lys Tyr  
 180 185 190  
 Trp Ser Gly Asn Ser Glu Phe His Phe Tyr Ser Ile Asn Val Gly Gly  
 195 200 205  
 Phe Phe Lys Leu Arg Ser Gly Glu Ile Ser Ile Glu Val Ser Asn  
 210 215 220  
 Pro Ser Leu Leu Asp Pro Asp Gln Asp Ala Thr Tyr Phe Gly Ala Phe  
 225 230 235 240  
 Lys Val Arg Asp Ile Asp  
 245

[0515] Sequence ID No. 18

[0516] Sequence Length: 735

[0517] Sequence Type: nucleic acid

[0518] Strandedness: 1

[0519] Topology: linear

[0520] Molecule Type: cDNA to mRNA

[0521] Sequence:

GCGCAGATGG ATCCTAACAG AATATCAGAA GACAGCACTC ACTGCTTTA TAGAATCCTG 60  
 AGACTCCATG AAAACGCAGG TTTGCAGGAC TCGACTCTGG AGAGTGAAGA CACACTACCT 120  
 GACTCCTGCA GGAGGATGAA ACAAGCCTT CAGGGCCCCG TCCAGAAGGA ACTGCAACAC 180  
 ATTGTGGTGC CACAGCGCTT CTCAGGAGCT CCAGCTATGA TGGAAGGCTC ATGGTTGGAT 240  
 GTGCCCCAGC GAGCCAAGCC TGAGGCCAG CCATTTGCAC ACCTCACCAT CAATGCTGCC 300  
 AGCATCCCAT CGGGTTCCCA TAAAGTCACT CTGTCCTCTT GGTACCACGA TCGAGGCTGG 360

**-continued**

GCCAAGATCT CTAACATGAC GTTAAGCAAC GGAAAACATAA GGTTAACCA AGATGGCTTC 420  
 TATTACCTGT ACGCCAACAT TTGCTTCGG CATCATGAAA CATCGGGAAAG CGTACCTACA 480  
 GACTATCTTC AGCTGATGGT GTATGTCGTT AAAACCAGCA TCAAAATCCC AAGTTCTCAT 540  
 AACCTGATGA AAGGAGGGAG CACGAAAAAC TGGTCCGGCA ATTCTGAATT CCACTTTAT 600  
 TCCATAAATG TTGGGGGATT TTTCAAGCTC CGAGCTGGTG AAGAAATTAG CATTACGTG 660  
 TCCAACCCTT CCCTGCTGGA TCCGGATCAA GATGCGACGT ACTTTGGGC TTTCAAAGTT 720  
 CAGGACATAG ACTGA 735

- [0522] Sequence ID No. 19
- [0523] Sequence Length: 741
- [0524] Sequence Type: nucleic acid
- [0525] Strandedness: 1
- [0526] Topology: linear
- [0527] Molecule Type: cDNA to mRNA
- [0528] Sequence:

GCGCAGATGG ATCCTAATAG AATATCAGAA GATGGCACTC ACTGCATTAA TAGAATTTG 60  
 AGACTCCATG AAAATGCAGA TTTTCAAGAC ACAACTCTGG AGAGTCAGA TACAAAATTA 120  
 ATACCTGATT CATGTAGGAG AATTAAACAG GCCTTCAAG GAGCTGTGCA AAAGGAATTA 180  
 CAACATATCG TTGGATCACA GCACATCAGA GCAGAGAAAG CGATGGTGGA TGGCTCATGG 240  
 TTAGATCTGG CCAAGACGAG CAAGCTTGAA GCTCAGCCTT TTGCTCATCT CACTATTAAAT 300  
 GCCACCGACA TCCCCTGTTCCATAAA GTGAGTCTGT CCTCTGGTA CCATGATCGG 360  
 GGTTGGGCCA AGATCTCAA CATGACTTTT AGCAATGGAA AACTAATAGT TAATCAGGAT 420  
 GGCTTTATT ACCTGTATGC CAACATTGTC TTTCGACATC ATGAAACTTC AGGAGACCTA 480  
 GCTACAGAGT ATCTTCAACT AATGGTGTAC GTCACTAAAA CCAGCATCAA AATCCCAAGT 540  
 TCTCATAACCC TGATGAAACG AGGAAGCACC AAGTATTGGT CAGGGAAATTC TGAATTCCAT 600  
 TTTTATTCCA TAAACGTTGG TGGATTTF AAGTTACGGT CTGGAGAGGA AATCAGCCTC 660  
 GAGGTCTCCA ACCCTCCTT ACTCGATCCG GATCAGGATG CAACATACTT TGGGGCTTT 720  
 AAAGTTCGAG ATATAGATTG A 741

- [0529]

---

 SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 19

<210> SEQ ID NO 1  
 <211> LENGTH: 316  
 <212> TYPE: PRT  
 <213> ORGANISM: *Mus musculus*  
 <400> SEQUENCE: 1

Met Arg Arg Ala Ser Arg Asp Tyr Gly Lys Tyr Leu Arg Ser Ser Glu  
 1 5 10 15

---

-continued

Glu Met Gly Ser Gly Pro Gly Val Pro His Glu Gly Pro Leu His Pro  
 20 25 30

Ala Pro Ser Ala Pro Ala Pro Ala Pro Pro Pro Ala Ala Ser Arg Ser  
 35 40 45

Met Phe Leu Ala Leu Leu Gly Leu Gly Leu Gly Gln Val Val Cys Ser  
 50 55 60

Ile Ala Leu Phe Leu Tyr Phe Arg Ala Gln Met Asp Pro Asn Arg Ile  
 65 70 75 80

Ser Glu Asp Ser Thr His Cys Phe Tyr Arg Ile Leu Arg Leu His Glu  
 85 90 95

Asn Ala Gly Leu Gln Asp Ser Thr Leu Glu Ser Glu Asp Thr Leu Pro  
 100 105 110

Asp Ser Cys Arg Arg Met Lys Gln Ala Phe Gln Gly Ala Val Gln Lys  
 115 120 125

Glu Leu Gln His Ile Val Gly Pro Gln Arg Phe Ser Gly Ala Pro Ala  
 130 135 140

Met Met Glu Gly Ser Trp Leu Asp Val Ala Gln Arg Gly Lys Pro Glu  
 145 150 155 160

Ala Gln Pro Phe Ala His Leu Thr Ile Asn Ala Ala Ser Ile Pro Ser  
 165 170 175

Gly Ser His Lys Val Thr Leu Ser Ser Trp Tyr His Asp Arg Gly Trp  
 180 185 190

Ala Lys Ile Ser Asn Met Thr Leu Ser Asn Gly Lys Leu Arg Val Asn  
 195 200 205

Gln Asp Gly Phe Tyr Tyr Leu Tyr Ala Asn Ile Cys Phe Arg His His  
 210 215 220

Glu Thr Ser Gly Ser Val Pro Thr Asp Tyr Leu Gln Leu Met Val Tyr  
 225 230 235 240

Val Val Lys Thr Ser Ile Lys Ile Pro Ser Ser His Asn Leu Met Lys  
 245 250 255

Gly Gly Ser Thr Lys Asn Trp Ser Gly Asn Ser Glu Phe His Phe Tyr  
 260 265 270

Ser Ile Asn Val Gly Gly Phe Phe Lys Leu Arg Ala Gly Glu Glu Ile  
 275 280 285

Ser Ile Gln Val Ser Asn Pro Ser Leu Leu Asp Pro Asp Gln Asp Ala  
 290 295 300

Thr Tyr Phe Gly Ala Phe Lys Val Gln Asp Ile Asp  
 305 310 315

<210> SEQ ID NO 2  
 <211> LENGTH: 1538  
 <212> TYPE: DNA  
 <213> ORGANISM: Mus musculus

<400> SEQUENCE: 2

gccaggacct ctgtgaacctc gtcggggcgg gggccgcctg gcccggagtc tgctcggcgg 60  
 tgggtggccg aggaagggag agaacgtatcg cggagcaggg cggccgaaact cccggcgcgg 120  
 cgccatgcgc cggccgcagcc gagactacgg caagtacctg cgcagctcgg aggagatggg 180  
 cagcggccccc ggcgtccac acgagggtcc gctgcacccc ggccttctg caccggctcc 240  
 ggcgcgcaca cccgcgcctc cccgctccat gttcctggcc ctccctgggc tgggactggg 300  
 ccaggtggtc tgcagcatcg ctctgttccct gtactttcga ggcgcagatgg atccctaacag 360

**-continued**


---

aatatcgaa gacagcactc actgcttta tagaatcctg agactccatg aaaacgcagg	420
tttgcaggac tcgactctgg agagtgaaga cacactacct gactcctgca ggaggatgaa	480
acaaggcctt cagggggccg tgcagaagga actgcaacac attgtggggc cacagcgctt	540
ctcaggagct ccagctatga tggaggcctc atggttggat gtggcccaagc gaggcaagcc	600
tgaggccag ccatttgac acctcaccat caatgctgcc agcatccat cgggttccca	660
taaagtcact ctgtccctt ggtaccacga tcgaggctgg gccaagatct ctaacatgac	720
gttaagcaac ggaaaactaa gggtaacca agatggcttc tattacctgt acgccaacat	780
ttgcttcgg catcatgaaa catcgaaag cgtacctaca gactatcttc agctgatgg	840
gtatgtcggtt aaaaccagca tcaaataccc aagttctcat aacctgtatga aaggaggag	900
cacaaaaaac tggtcggca attctgaatt ccactttat tccataaatg ttggggatt	960
tttcaagctc cgagctgggtt aagaaattag cattcagggt tccaaaccctt ccctgctgga	1020
tccggatcaa gatgcgacgt actttggggc tttcaaagtt caggacatag actgagactc	1080
atttcgtgga acattagcat ggatgtccata gatgtttgga aacttcttaa aaaatggatg	1140
atgtctatac atgtgtttaa ctactaagag acatggccca cggtgtatga aactcacagc	1200
cctctctt gggctgtac aggttgttata tatgtaaagt ccatagggtatgtttagattc	1260
atggtgattt cacaacgggtt ttacaatttt gtaatgattt cctagaattt aaccagattt	1320
ggagaggtat tccgatgtttt atgaaaaact tacacgttacatgttacatgttacatgtt	1380
ctctgggtctt aacccttggaa catgtgccac tgagaacctt gaaattaaga ggatgcccattt	1440
tcattgtttaa gaaatgtatgtt gttttttttt tttttttttt tttttttttt tttttttttt	1500
ggacctgcaaa ataagttctt tttttctaat gaggagag	1538

<210> SEQ ID NO 3  
 <211> LENGTH: 21  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence:primer  
 <400> SEQUENCE: 3

aaacgcaaaa aaccagaaag g 21

<210> SEQ ID NO 4  
 <211> LENGTH: 17  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence:primer  
 <400> SEQUENCE: 4

gtaaaacgac ggccagt 17

<210> SEQ ID NO 5  
 <211> LENGTH: 17  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence:primer  
 <400> SEQUENCE: 5

caggaaacag ctatgac 17

---

-continued

---

<210> SEQ ID NO 6  
<211> LENGTH: 22  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence:primer OBM #8

<400> SEQUENCE: 6

aagccccaaa gtacgtcgca tc 22

<210> SEQ ID NO 7  
<211> LENGTH: 26  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence:OBM HF

<400> SEQUENCE: 7

cgaagcttgc gacgcgacat ggcattcc 26

<210> SEQ ID NO 8  
<211> LENGTH: 27  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence:OBM XR

<400> SEQUENCE: 8

cctctagagt ctatgtcctg aagtttg 27

<210> SEQ ID NO 9  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence:OBM3

<400> SEQUENCE: 9

atcagaagac agcactcact 20

<210> SEQ ID NO 10  
<211> LENGTH: 33  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence:OBMSalR2

<400> SEQUENCE: 10

ggggtcgacc taggacatcc atgctaatgt tcc 33

<210> SEQ ID NO 11  
<211> LENGTH: 317  
<212> TYPE: PRT  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 11

Met Arg Arg Ala Ser Arg Asp Tyr Thr Lys Tyr Leu Arg Gly Ser Glu  
1 5 10 15

Glu Met Gly Gly Pro Gly Ala Pro His Glu Gly Pro Leu His Ala  
20 25 30

---

-continued

---

Pro Pro Pro Pro Ala Pro His Gln Pro Pro Ala Ala Ser Arg Ser Met  
 35 40 45  
 Phe Val Ala Leu Leu Gly Leu Gly Leu Gly Gln Val Val Cys Ser Val  
 50 55 60  
 Ala Leu Phe Phe Tyr Phe Arg Ala Gln Met Asp Pro Asn Arg Ile Ser  
 65 70 75 80  
 Glu Asp Gly Thr His Cys Ile Tyr Arg Ile Leu Arg Leu His Glu Asn  
 85 90 95  
 Ala Asp Phe Gln Asp Thr Thr Leu Glu Ser Gln Asp Thr Lys Leu Ile  
 100 105 110  
 Pro Asp Ser Cys Arg Arg Ile Lys Gln Ala Phe Gln Gly Ala Val Gln  
 115 120 125  
 Lys Glu Leu Gln His Ile Val Gly Ser Gln His Ile Arg Ala Glu Lys  
 130 135 140  
 Ala Met Val Asp Gly Ser Trp Leu Asp Leu Ala Lys Arg Ser Lys Leu  
 145 150 155 160  
 Glu Ala Gln Pro Phe Ala His Leu Thr Ile Asn Ala Thr Asp Ile Pro  
 165 170 175  
 Ser Gly Ser His Lys Val Ser Leu Ser Ser Trp Tyr His Asp Arg Gly  
 180 185 190  
 Trp Ala Lys Ile Ser Asn Met Thr Phe Ser Asn Gly Lys Leu Ile Val  
 195 200 205  
 Asn Gln Asp Gly Phe Tyr Tyr Leu Tyr Ala Asn Ile Cys Phe Arg His  
 210 215 220  
 His Glu Thr Ser Gly Asp Leu Ala Thr Glu Tyr Leu Gln Leu Met Val  
 225 230 235 240  
 Tyr Val Thr Lys Thr Ser Ile Lys Ile Pro Ser Ser His Thr Leu Met  
 245 250 255  
 Lys Gly Gly Ser Thr Lys Tyr Trp Ser Gly Asn Ser Glu Phe His Phe  
 260 265 270  
 Tyr Ser Ile Asn Val Gly Phe Phe Lys Leu Arg Ser Gly Glu Glu  
 275 280 285  
 Ile Ser Ile Glu Val Ser Asn Pro Ser Leu Leu Asp Pro Asp Gln Asp  
 290 295 300  
 Ala Thr Tyr Phe Gly Ala Phe Lys Val Arg Asp Ile Asp  
 305 310 315

<210> SEQ ID NO 12  
 <211> LENGTH: 954  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 12

```

atgcgcccgccagccagccatcacaccaag tacctgcgtg gctcgaggaa gatggccggc 60
ggccccggag ccccgacgaa gggcccccctg cacgccccgc cgccgcctgc gccgcacca 120
ccccctgcgg cctcccgctc catgttcgtg gcccctctgg ggctggggct gggccaggtt 180
gtctgcagcg tgcgcctgtt cttcttatttc agagcgcaga tggatcccaa tagaatatca 240
gaagatggca ctcactgcat ttatagaatt ttgagactcc atgaaaatgc agatttcaa 300
gacacaactc tggagagtca agataaaaaa ttaataccctg attcatgttag gagaattaaa 360
caggcctttc aaggagctgt gcaaaaaggaa ttacaacata tcgttggatc acagcacatc 420

```

**-continued**


---

agagcagaga aagcgatggt ggatggctca tggtagatc tggccaagag gagcaagctt	480
gaagctcagc cttttgtca tctcaattt aatgccacgg acatccatc tggttccat	540
aaagtggatc tgcctcttg gtaccatgtat cggggttggg ccaagatctc caacatgact	600
tttagcaatg gaaaactaat agttaatcag gatggctttt attaccgtta tgccaacatt	660
tgctttcgac atcatgaaac ttcaaggagac ctatctacag agtatacttca actaatggtg	720
tacgtcacta aaaccagcat caaaatccca agttctata ccctgtatgaa aggaggaagc	780
accaagtatt ggtcaggggaa ttctgaattt catttttatt ccataaacgt tggggattt	840
tttaagttac ggtctggaga ggaaatcagc atcgagggtct ccaacccttc cttactggat	900
ccggatcagg atgcaacata ctttgggct tttaaagttc gagatataaga ttga	954
<210> SEQ ID NO 13	
<211> LENGTH: 27	
<212> TYPE: DNA	
<213> ORGANISM: Artificial Sequence	
<220> FEATURE:	
<223> OTHER INFORMATION: Description of Artificial Sequence:human OBM SF	
<400> SEQUENCE: 13	
ggcgtacgca gagcgcagat ggatcct	27
<210> SEQ ID NO 14	
<211> LENGTH: 34	
<212> TYPE: DNA	
<213> ORGANISM: Artificial Sequence	
<220> FEATURE:	
<223> OTHER INFORMATION: Description of Artificial Sequence:synthetic DNA	
<400> SEQUENCE: 14	
gggttcgacc atccaggaaa tatcataaca ctcc	34
<210> SEQ ID NO 15	
<211> LENGTH: 951	
<212> TYPE: DNA	
<213> ORGANISM: Mus musculus	
<400> SEQUENCE: 15	
atgcgcggg ccagccgaga ctacggcaag tacctgcgca gtcggagga gatggcagc	60
ggcccccggcg tcccacacgaa gggccggctg caccggcgcc tttctgcacc ggctccggcg	120
ccggccaccccg ccgcctcccg ctccatgttc ctggccctcc tggggctggg actggggccag	180
gtggtctgca gcatcgatct gttcgttac ttccgagcgc agatggatcc taacagaata	240
tcagaagaca gcactcactg cttttataga atccgtgagac tccatgaaaa cgcaggtttg	300
caggactcga ctctggagag tgaagacaca ctacctgact cctgcaggag gatgaaacaa	360
gcctttcagg gggccgtgca gaaggaactg caacacatgg tggggccaca ggcgttctca	420
ggagctccag ctatgtatgg aggctcatgg ttggatgtgg cccagcggagg caagctgag	480
gcccgccat ttgcacacccat caccatcaat gctgccagca tcccatcggtt ttccataaa	540
gtcactctgt cctcttggta ccacgtcga ggctggccca agatcttaa catgacgttta	600
agcaacggaa aactaagggt taaccaagat ggcttctatt acctgtacgc caacatttg	660
tttcggcattc atgaaacatc gggaaagcgta cctacagact atcttcagct gatggtgat	720

**-continued**


---

```

gtcgtaaaa ccagcatcaa aatcccaagt tctcataacc tcatgaaagg agggagcacg    780
aaaaactggt cgggcaattc tgaattccac ttttattcca taaatgttgg gggattttc    840
aagctccgag ctggtaaga aatttagcatt caggtgtcca acccttcct gctggatccg    900
gatcaagatg cgacgtactt tggggcttc aaagttcagg acatagactg a           951

```

<210> SEQ ID NO 16

<211> LENGTH: 244

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 16

```

Ala Gln Met Asp Pro Asn Arg Ile Ser Glu Asp Ser Thr His Cys Phe
 1           5           10           15

```

```

Tyr Arg Ile Leu Arg Leu His Glu Asn Ala Gly Leu Gln Asp Ser Thr
 20          25          30

```

```

Leu Glu Ser Glu Asp Thr Leu Pro Asp Ser Cys Arg Arg Met Lys Gln
 35          40          45

```

```

Ala Phe Gln Gly Ala Val Gln Lys Glu Leu Gln His Ile Val Gly Pro
 50          55          60

```

```

Gln Arg Phe Ser Gly Ala Pro Ala Met Met Glu Gly Ser Trp Leu Asp
 65          70          75          80

```

```

Val Ala Gln Arg Gly Lys Pro Glu Ala Gln Pro Phe Ala His Leu Thr
 85          90          95

```

```

Ile Asn Ala Ala Ser Ile Pro Ser Gly Ser His Lys Val Thr Leu Ser
100          105          110

```

```

Ser Trp Tyr His Asp Arg Gly Trp Ala Lys Ile Ser Asn Met Thr Leu
115          120          125

```

```

Ser Asn Gly Lys Leu Arg Val Asn Gln Asp Gly Phe Tyr Tyr Leu Tyr
130          135          140

```

```

Ala Asn Ile Cys Phe Arg His His Glu Thr Ser Gly Ser Val Pro Thr
145          150          155          160

```

```

Asp Tyr Leu Gln Leu Met Val Tyr Val Val Lys Thr Ser Ile Lys Ile
165          170          175

```

```

Pro Ser Ser His Asn Leu Met Lys Gly Gly Ser Thr Lys Asn Trp Ser
180          185          190

```

```

Gly Asn Ser Glu Phe His Phe Tyr Ser Ile Asn Val Gly Gly Phe Phe
195          200          205

```

```

Lys Leu Arg Ala Gly Glu Glu Ile Ser Ile Gln Val Ser Asn Pro Ser
210          215          220

```

```

Leu Leu Asp Pro Asp Gln Asp Ala Thr Tyr Phe Gly Ala Phe Lys Val
225          230          235          240

```

Gln Asp Ile Asp

<210> SEQ ID NO 17

<211> LENGTH: 246

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 17

```

Ala Gln Met Asp Pro Asn Arg Ile Ser Glu Asp Gly Thr His Cys Ile
 1           5           10           15

```

```

Tyr Arg Ile Leu Arg Leu His Glu Asn Ala Asp Phe Gln Asp Thr Thr
 20          25          30

```

---

-continued

---

Leu Glu Ser Gln Asp Thr Lys Leu Ile Pro Asp Ser Cys Arg Arg Ile  
 35 40 45  
 Lys Gln Ala Phe Gln Gly Ala Val Gln Lys Glu Leu Gln His Ile Val  
 50 55 60  
 Gly Ser Gln His Ile Arg Ala Glu Lys Ala Met Val Asp Gly Ser Trp  
 65 70 75 80  
 Leu Asp Leu Ala Lys Arg Ser Lys Leu Glu Ala Gln Pro Phe Ala His  
 85 90 95  
 Leu Thr Ile Asn Ala Thr Asp Ile Pro Ser Gly Ser His Lys Val Ser  
 100 105 110  
 Leu Ser Ser Trp Tyr His Asp Arg Gly Trp Ala Lys Ile Ser Asn Met  
 115 120 125  
 Thr Phe Ser Asn Gly Lys Leu Ile Val Asn Gln Asp Gly Phe Tyr Tyr  
 130 135 140  
 Leu Tyr Ala Asn Ile Cys Phe Arg His His Glu Thr Ser Gly Asp Leu  
 145 150 155 160  
 Ala Thr Glu Tyr Leu Gln Leu Met Val Tyr Val Thr Lys Thr Ser Ile  
 165 170 175  
 Lys Ile Pro Ser Ser His Thr Leu Met Lys Gly Gly Ser Thr Lys Tyr  
 180 185 190  
 Trp Ser Gly Asn Ser Glu Phe His Phe Tyr Ser Ile Asn Val Gly Gly  
 195 200 205  
 Phe Phe Lys Leu Arg Ser Gly Glu Glu Ile Ser Ile Glu Val Ser Asn  
 210 215 220  
 Pro Ser Leu Leu Asp Pro Asp Gln Asp Ala Thr Tyr Phe Gly Ala Phe  
 225 230 235 240  
 Lys Val Arg Asp Ile Asp  
 245

<210> SEQ ID NO 18  
 <211> LENGTH: 735  
 <212> TYPE: DNA  
 <213> ORGANISM: Mus musculus  
 <400> SEQUENCE: 18

```

gccccatgg atcctaacag aatatcagaa gacagcactc actgccttta tagaattcctg 60
agactccatg aaaacgcagg tttgcaggac tcgactctgg agagtgaaga cacactacct 120
gactcctgca ggaggatgaa acaagcctt cagggggccg tgcagaagga actgcaacac 180
atgtgtgggc cacagcgctt ctcaggagct ccagctatga tggaggctc atggttggat 240
gtggcccgac gaggcaagcc tgaggccag ccatttgac acctcacat caatgctgcc 300
agcatcccat cgggttccca taaagtcact ctgtcctt ggtaccacga tcgaggctgg 360
gccaagatct ctaacatgac gttaagcaac ggaaaactaa gggtaacca agatggcttc 420
tattacctgt acgccaacat ttgccttcgg catcatgaaa catcggaaag cgtacctaca 480
gactatcttc agctgatggt gtatgtcggt aaaaccagca tcaaaatccc aagttctcat 540
aacctgatga aaggaggaggac cacgaaaaac tggtcggca attctgaatt ccacttttat 600
tccataaaatg ttgggggatt tttcaagctc cgagctggtg aagaaattag cattcaggtg 660
tccaaaccctt ccctgctgga tccggatcaa gatgcgacgt actttggggc tttcaaaatg 720
caggacatag actga 735
  
```

---

-continued

---

<210> SEQ ID NO 19  
 <211> LENGTH: 741  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 19

gcgcagatgg atccataatag aatatcagaa gatggcactc actgcattta tagaattttg	60
agactccatg aaaatgcaga ttttcaagac acaactctgg agagtcaaga tacaaaatta	120
atacctgatt catgttaggag aattaaacag gccttcaag gagctgtgca aaaggaatta	180
caacatatcg ttggatcaca gcacatcaga gcagagaaag cgatggtgg tggctcatgg	240
ttagatctgg ccaagaggag caagcttcaa gctcagccctt ttgcctatct cactattaaat	300
gccaccgaca tcccatctgg ttcccataaa gtgagtcgt cctcttgta ccatgatcg	360
ggttgggcca agatctccaa catgactttt agcaatggaa aactaatagt taatcaggat	420
ggcttttattt acctgtatgc caacatggc ttgcacatc atgaaacttc aggagaccta	480
gctacagagt atcttcaact aatgggtgtac gtcactaaaa ccagcatcaa aatcccaagt	540
tctcataccctt tcatgaaagg aggaaggcacc aagtatttggt cagggaaattc tgaattccat	600
ttttatttccaa taaacgttgg tggatttttt aagttacggt ctggagagga aatcagcatc	660
gaggctctccaa acccctccctt actggatccg gatcaggatg caacatactt tggggctttt	720
aaagttcgag atatagattt a	741

---

What is claimed is:

1. A novel protein having the following physicochemical characteristics and biological activity:
  - (a) affinity: specifically binds to osteoclastogenesis inhibitory factor (OCIF) and exhibits high affinity to OCIF (dissociation constant on cell membrane:  $K_d=10^{-9}$  M or less);
  - (b) molecular weight: has a molecular weight of about 30,000-40,000 or about  $40,000\pm4,000$  when measured using SDS-polyacrylamide electrophoresis under non-reducing conditions and an apparent molecular weight of about 90,000-110,000 when cross-linked to a monomer-type OCIF; and
  - (c) biological activity: exhibits activity supporting or promoting osteoclasts differentiation and maturation in a co-culture system of the osteoblastic stromal cells and spleen cells in the presence of such stimulating factors of bone resorption as active-form vitamin D<sub>3</sub> and parathyroid hormone (PTH).
2. A process for the preparation of the protein of claim 1, comprising culturing cells of an osteoblast-like cell line or stromal cells originating from bone marrow, in the presence of stimulating factors of bone resorption, preparing a membrane fraction of the cells, solubilizing the membrane protein using a detergent, and purifying the protein using an OCIF-immobilized affinity column.
3. A screening method using the protein of claim 1 for a substance which specifically binds to the protein of claim 1 and inhibits or modulates the biological activity of this protein.
4. A screening method using the protein of claim 1 for a substance which specifically binds to the protein of claim 1 and transmits the biological activity of this protein.
5. A pharmaceutical composition containing the substance obtained by the screening method described in claim 3 or 4.
6. An antibody to the protein of claim 1.
7. An agent for treating bone metabolism abnormality comprising the antibody of claim 6 as an active component.
8. A DNA encoding the amino acid sequence of Sequence ID No. 1 in the Sequence Table.
9. A DNA which consists of the nucleotide sequence of Sequence ID No. 2 in the Sequence Table.
10. A DNA with one or more nucleotides deleted, substituted, added, or inserted in the nucleotide sequence of Sequence ID No. 2 in the Sequence Table, and encoding a protein capable of specifically binding to Osteoclastogenesis Inhibitory Factor and suppressing its activity.
11. A DNA which hybridizes either the DNA of claim 8 or the DNA of claim 10 under relatively mild conditions.
12. A DNA having 70% or more homology to the DNA of any one of the claims 8-11.
13. A protein obtained by the expression of the DNA of any one of the claims 8-12 by genetic engineering technique.
14. A protein of claim 13 which exhibits the activity to support and promote osteoclast differentiation or maturation.
15. The protein having the amino acid sequence of Sequence ID No. 1 in the Sequence Table.
16. A process for producing a protein by genetic engineering technique using the DNA of any one of the claims 8-12.

17. A screening method for a substance which controls expression of the protein encoded by the DNA of any one of the claims **8-12**.

18. A screening method for a substance which specifically binds to the protein encoded by the DNA of any one of the claims **8-12** and inhibits or modulates its biological activity.

19. A screening method for a substance which specifically binds to the protein encoded by the DNA of any one of the claims **8-12** and transmits the biological activity of this protein through the binding.

20. A pharmaceutical composition comprising the substance obtained by the screening method described in claims **17-19**.

21. A DNA encoding the protein including the amino acid sequence consisting of the amino acid residues from No. 76 to No. 316 of Sequence ID No. 1 in the Sequence Table, or according to a fragment, analogue, or variant of the protein.

22. A DNA encoding the protein including the amino acid sequence consisting of the amino acid residues from No. 72 to No. 316 of Sequence ID No. 1 in the Sequence Table, or encoding a fragment, analogue, or variant of the protein.

23. A DNA encoding the protein including the amino acid sequence of Sequence ID No. 1 in the Sequence Table, or a fragment, analogue, or variant of the protein.

24. The DNA of any one of the claims **21-23**, which encodes a protein specifically binding to Osteoclastogenesis Inhibitory Factor, OCIF, and exhibiting the activity of suppressing its biological activity.

25. A DNA of any one of the claims **21-24** which has a nucleotide sequence of Sequence ID No. 2 in the Sequence Table.

26. A DNA which hybridizes to the DNA of any one of claims **21-25** under relatively mild conditions.

27. A DNA having 70% or more homology to the DNA of any one of claims **21-26**.

28. A protein including a polypeptide which is obtained by expression of the DNA of any one of the claims **21-27** by genetic engineering technique.

29. The protein of claim 28 exhibiting a biological activity to support and promote the osteoclast differentiation and maturation.

30. A protein of claim 28 or **29** including the amino acid sequence of Sequence ID No. 1 in the Sequence Table, or a fragment, analogue, or variant of the protein.

31. A process for producing a protein by genetic engineering technique using the DNA of any one of the claims **21-27**.

32. A screening method for a substance which controls expression of the protein encoded by the DNA of any one of the claims **21-27**.

33. A screening method for a substance which specifically binds to the protein encoded by DNA of any one of the claims **21-27** and inhibits or modulates its biological activity.

34. A screening method for a substance which specifically binds to a protein encoded by the DNA of any one of claims **21-27** and transmits the biological activity of this protein through the binding.

35. A pharmaceutical composition comprising the substance obtained by the screening method of any one of claims **32-34**.

36. A DNA encoding the protein including the amino acid sequence of Sequence ID No. 11 in the Sequence Table, or a fragment, analogue, or variant of the protein.

37. The DNA of claim 36, which encodes a protein specifically binding to Osteoclastogenesis Inhibitory Factor, OCIF, and exhibiting the activity of suppressing its biological activity.

38. A DNA of any one of claims **36-37** which has a nucleotide sequence of Sequence ID No. 12 in the Sequence Table.

39. A DNA which hybridizes to the DNA of any one of claims **36-38** under relatively mild conditions.

40. A DNA having 70% or more homology to the DNA of any one of claims **36-39**.

41. A protein including a polypeptide which is obtained by expression of the DNA of any one of the claims **36-40** by genetic engineering technique.

42. A protein of claim 41 which exhibits the activity to support and promote the differentiation into and the maturation of osteoclasts.

43. A protein of claim 41 or **42** including the amino acid sequence of Sequence ID No. 11 in the Sequence Table, or a fragment, analogue, or variant of the protein.

44. A process for producing a protein by a genetic engineering technique using the DNA of any one of the claims **36-40**.

45. A screening method for a substance which controls expression of the protein encoded by the DNA of any one of the claims **36-40**.

46. A screening method for a substance which specifically binds to the protein encoded by the DNA of any one of claims **36-40** and inhibits or modulates its biological activity.

47. A screening method for a substance which specifically binds to the protein encoded by the DNA of any one of claims **36-40** and transmits the biological activity of this protein through the binding.

48. A pharmaceutical composition comprising the substance obtained by the screening method any one of claims **45-47**.

49. An antibody which recognizes a membrane-bound protein molecule (OBM) which binds specifically to Osteoclastogenesis Inhibitory Factor (OCIF) and/or a soluble OBM (sOBM), which is deficient in the transmembrane region.

50. The antibody of claim 49 which recognizes mouse OBM and/or mouse sOBM.

51. The antibody of claim 49 which recognizes human OBM and/or human sOBM.

52. The antibody of claim 49 which is a polyclonal antibody.

53. The antibody of claim 49 which is a monoclonal antibody.

54. A monoclonal antibody of claim 53 which has cross-reactivity to human OBM, human sOBM, mouse OBM, and mouse sOBM.

55. The monoclonal antibody of claim 53 which possesses following characteristics:

- molecular weight: about 150,000,
- subclass: IgG<sub>1</sub> or IgG<sub>2</sub>, and
- light chain: κ chain

56. A process for preparing a polyclonal antibody comprising immunizing an animal with mouse or human OBM or sOBM and purifying from blood of the animal.

57. A hybridoma which produces the monoclonal antibody of any one of claims **53-55**.

**58.** The hybridomas with deposition numbers FERM BP-6264, FERM BP-6265, or FERM BP-6266.

**59.** A process for preparing a monoclonal antibody comprising cultivating the hybridoma of claim 57 or **58** to produce a monoclonal antibody in the culture broth and purifying and recovering the monoclonal antibody from the culture broth.

**60.** A method for measuring OBM, comprising using the antibody of any one of the claims **49-55**.

**61.** A method for measuring sOBM, comprising using the antibody of any one of claims **49-55**.

**62.** A method for measuring OBM or sOBM, comprising using the antibody of any one of claims **49-55** as an solid phase antibody or an enzyme-labeled antibody.

**63.** A pharmaceutical composition containing the antibody of any one of claims **49-55** as an active component.

**64.** An agent for preventing or treating bone metabolism diseases comprising the antibody of any one of claims **49-57** as an active component.

\* \* \* \* \*

专利名称(译)	针对OCIF结合分子的抗体		
公开(公告)号	<a href="#">US20030176647A1</a>	公开(公告)日	2003-09-18
申请号	US10/167182	申请日	2002-06-11
[标]申请(专利权)人(译)	株式会社三共		
申请(专利权)人(译)	SANKYO CO. , LTD.		
当前申请(专利权)人(译)	SANKYO CO. , LTD.		
[标]发明人	YAMAGUCHI KYOJI YASUDA HISATAKA NAKAGAWA NOBUAKI SHIMA NOBUYUKI KINOSAKI MASAHIKO TSUDA EISUKE GOTO MASAAKI YANO KAZUKI TOMOYASU AKIHIRO KOBAYASHI FUMIE WASHIDA NAOHIRO TAKAHASHI KEN MORINAGA TOMONORI HIGASHIO KANJI		
发明人	YAMAGUCHI, KYOJI YASUDA, HISATAKA NAKAGAWA, NOBUAKI SHIMA, NOBUYUKI KINOSAKI, MASAHIKO TSUDA, EISUKE GOTO, MASAAKI YANO, KAZUKI TOMOYASU, AKIHIRO KOBAYASHI, FUMIE WASHIDA, NAOHIRO TAKAHASHI, KEN MORINAGA, TOMONORI HIGASHIO, KANJI		
IPC分类号	C07K14/715 A61K A61K38/00 A61K39/395 A61K45/00 A61P3/14 A61P19/00 A61P19/02 A61P19/08 A61P19/10 A61P29/00 B01J20/281 C07K C07K1/22 C07K14/47 C07K14/525 C07K14/705 C07K16/18 C07K16/28 C12N C12N5/10 C12N5/20 C12N15/02 C12N15/09 C12N15/12 C12P C12P21/02 C12P21 /08 C12Q G01N G01N30/88 G01N33/15 G01N33/50 G01N33/53 G01N33/543 G01N33/566 G01N33 /577 C07K14/725 C07H21/04 C12N5/06		
CPC分类号	A61K38/00 G01N2800/108 C07K14/525 C07K14/705 C07K14/70578 C07K16/28 C07K16/2875 C07K16 /2878 C07K2316/96 C07K2317/92 C07K2319/00 G01N33/566 G01N2333/525 G01N2333/70578 A61K2039/505 A61P3/14 A61P19/00 A61P19/02 A61P19/04 A61P19/08 A61P19/10 A61P29/00 A61P43/00 C07K2317/76		
优先权	PCT/JP1998/001728 1998-04-15 WO 2008000978 1997-04-15 JP 1989051434 1997-06-09 JP		

1990017897 1997-08-12 JP

1990024803 1997-08-21 JP

1991032241 1997-12-02 JP

其他公开文献

US7527790

外部链接

[Espacenet](#) [USPTO](#)

### 摘要(译)

一种结合破骨细胞生成抑制因子 (OCIF结合分子; OBM) 的新蛋白质，其制备方法，编码所述蛋白质的DNA，具有该DNA编码的氨基酸序列的蛋白质，通过遗传产生所述蛋白质的方法工程技术和含有所述蛋白质的药物组合物。使用所述蛋白质和DNA控制所述蛋白质表达的物质的筛选方法，抑制或调节所述蛋白质的生物活性的物质，或通过与所述蛋白质结合而传递所述蛋白质的作用的受体，所获得的物质通过筛选方法和含有该物质的药物组合物。用于所述蛋白质的抗体，其制备方法，使用所述抗体测量所述蛋白质的方法，以及包含该抗体的药物。

