




US 20100136581A1

(19) **United States**(12) **Patent Application Publication**
Wands et al.(10) **Pub. No.: US 2010/0136581 A1**(43) **Pub. Date: Jun. 3, 2010**(54) **DIAGNOSIS AND TREATMENT OF
MALIGNANT NEOPLASMS**(75) Inventors: **Jack R. Wands**, Waban, MA (US);
Suzanne M. de la Monte, East
Greenwich, RI (US); **Nedim Ince**,
Boston, MA (US); **Rolf I. Carlson**,
Boston, MA (US)

Correspondence Address:

**MINTZ, LEVIN, COHN, FERRIS, GLOVSKY
AND POPEO, P.C**
ONE FINANCIAL CENTER
BOSTON, MA 02111 (US)(73) Assignee: **Rhode Island Hospital**,
Providence, RI (US)(21) Appl. No.: **12/625,729**(22) Filed: **Nov. 25, 2009****Related U.S. Application Data**(60) Continuation of application No. 11/376,941, filed on
Mar. 15, 2006, now abandoned, which is a continua-
tion of application No. 09/903,023, filed on Jul. 11,
2001, now Pat. No. 7,094,556, which is a division of
application No. 09/436,184, filed on Nov. 8, 1999, now
abandoned.**Publication Classification**(51) **Int. Cl.**
G01N 33/53

(2006.01)

(52) **U.S. Cl.** **435/7.9**(57) **ABSTRACT**The invention features a method for diagnosing a malignant
neoplasm in a mammal by contacting a bodily fluid from the
mammal with an antibody which binds to an human aspartyl
(asparaginy) beta-hydroxylase (HAAH) polypeptide and
methods of treating malignant neoplasms by inhibiting
HAAH.




Fig. 1




Fig. 2




Fig. 3A




Fig. 3B




Fig. 4




Fig. 5A




Fig. 5B




Fig. 5C





Fig. 5D

**Fig. 6**




Fig. 7




Fig. 8




Fig. 9

DIAGNOSIS AND TREATMENT OF MALIGNANT NEOPLASMS

CROSS-REFERENCES TO RELATED APPLICATIONS

[0001] This application is a continuation of U.S. application Ser. No. 11/376,941 filed Mar. 15, 2006, which is a continuation of U.S. patent Ser. No. 09/903,023, filed Jul. 11, 2001, which is a division of U.S. patent application Ser. No. 09/436,184, filed Nov. 8, 1999, each of which is herein incorporated by reference in its entirety.

STATEMENT AS TO FEDERALLY SPONSORED RESEARCH

[0002] This invention was made with U.S. Government support under National Institutes of Health grants CA-35711, AA-02666, AA-02169, and AA11431. The government has certain rights in the invention.

BACKGROUND OF THE INVENTION

[0003] Primary malignant central nervous system (CNS) neoplasms, particularly glioblastomas, are highly fatal due to their aggressive and widespread infiltration of the brain and resistance to anti-cancer treatments. Although progress has been made in unraveling the pathological mechanisms underlying CNS cancers as well as other cancer types, tumor specific therapeutic approaches and methods of diagnosis have been largely elusive.

SUMMARY OF THE INVENTION

[0004] The invention features a method for diagnosing a malignant neoplasm in a mammal by contacting a bodily fluid from the mammal with an antibody which binds to a human aspartyl (asparaginy) beta-hydroxylase (HAAH) polypeptide under conditions sufficient to form an antigen-antibody complex and detecting the antigen-antibody complex. Malignant neoplasms detected in this manner include those derived from endodermal tissue, e.g., colon cancer, breast cancer, pancreatic cancer, liver cancer, and cancer of the bile ducts. Neoplasms of the central nervous system (CNS) such as primary malignant CNS neoplasms of both neuronal and glial cell origin and metastatic CNS neoplasms are also detected. Patient derived tissue samples, e.g., biopsies of solid tumors, as well as bodily fluids such as a CNS-derived bodily fluid, blood, serum, urine, saliva, sputum, lung effusion, and ascites fluid, are contacted with an HAAH-specific antibody.

[0005] The assay format is also useful to generate temporal data used for prognosis of malignant disease. A method for prognosis of a malignant neoplasm of a mammal is carried out by (a) contacting a bodily fluid from the mammal with an antibody which binds to an HAAH polypeptide under conditions sufficient to form an antigen-antibody complex and detecting the antigen-antibody complex; (b) quantitating the amount of complex to determine the level of HAAH in the fluid; and (c) comparing the level of HAAH in the fluid with a normal control level of HAAH. An increasing level of HAAH over time indicates a progressive worsening of the disease, and therefore, an adverse prognosis.

[0006] The invention also includes an antibody which binds to HAAH. The antibody preferably binds to a site in the carboxyterminal catalytic domain of HAAH. Alternatively, the antibody binds to an epitope that is exposed on the surface of the cell. The antibody is a polyclonal antisera or mono-

clonal antibody. The invention encompasses not only an intact monoclonal antibody, but also an immunologically-active antibody fragment, e.g., a Fab or (Fab)₂ fragment; an engineered single chain Fv molecule; or a chimeric molecule, e.g., an antibody which contains the binding specificity of one antibody, e.g., of murine origin, and the remaining portions of another antibody, e.g., of human origin. Preferably the antibody is a monoclonal antibody such as FB50, 5C7, 5E9, 19B, 48A, 74A, 78A, 86A, HA238A, HA221, HA 239, HA241, HA329, or HA355. Antibodies which bind to the same epitopes as those monoclonal antibodies are also within the invention.

[0007] An HAAH-specific intrabody is a recombinant single chain HAAH-specific antibody that is expressed inside a target cell, e.g., tumor cell. Such an intrabody binds to endogenous intracellular HAAH and inhibits HAAH enzymatic activity or prevents HAAH from binding to an intracellular ligand. HAAH-specific intrabodies inhibit intracellular signal transduction, and as a result, inhibit growth of tumors which overexpress HAAH.

[0008] A kit for diagnosis of a tumor in a mammal contains an HAAH-specific antibody. The diagnostic assay kit is preferentially formulated in a standard two-antibody binding format in which one HAAH-specific antibody captures HAAH in a patient sample and another HAAH-specific antibody is used to detect captured HAAH. For example, the capture antibody is immobilized on a solid phase, e.g., an assay plate, an assay well, a nitrocellulose membrane, a bead, a dipstick, or a component of an elution column. The second antibody, i.e., the detection antibody, is typically tagged with a detectable label such as a colorimetric agent or radioisotope.

[0009] Also within the invention is a method of inhibiting tumor growth in a mammal, which is carried out by administering to the mammal a compound which inhibits expression or enzymatic activity of HAAH. Preferably, the compound is substantially pure nucleic acid molecule such as an HAAH antisense DNA, the sequence of which is complementary to a coding sequence of HAAH. Expression of HAAH is inhibited by contacting mammalian cells, e.g., tumor cells, with HAAH antisense DNA or RNA, e.g., a synthetic HAAH antisense oligonucleotide. For example, HAAH antisense nucleic acid is introduced into glioblastoma cells or other tumor cells which overexpress HAAH. Binding of the antisense nucleic acid to an HAAH transcript in the target cell results in a reduction in HAAH production by the cell. By the term "antisense nucleic acid" is meant a nucleic acid (RNA or DNA) which is complementary to a portion of an mRNA, and which hybridizes to and prevents translation of the mRNA. Preferably, the antisense DNA is complementary to the 5' regulatory sequence or the 5' portion of the coding sequence of HAAH mRNA (e.g., a sequence encoding a signal peptide or a sequence within exon 1 of the HAAH gene). Standard techniques of introducing antisense DNA into the cell may be used, including those in which antisense DNA is a template from which an antisense RNA is transcribed. The method is to treat tumors in which expression of HAAH is upregulated, e.g., as a result of malignant transformation of the cells. The length of the oligonucleotide is at least 10 nucleotides and may be as long as the naturally-occurring HAAH transcript. Preferably, the length is between 10 and 50 nucleotides, inclusive. More preferably, the length is between 10 and 20 nucleotides, inclusive.

[0010] By "substantially pure DNA or RNA" is meant that the nucleic acid is free of the genes which, in the naturally-

occurring genome of the organism from which the DNA of the invention is derived, flank a HAAH gene. The term therefore includes, for example, a recombinant nucleic acid which is incorporated into a vector, into an autonomously replicating plasmid or virus, or into the genomic DNA of a procaryote or eucaryote at a site other than its natural site; or which exists

specific intrabody is also useful to bind to HAAH and inhibit intracellular HAAH enzymatic activity, e.g., by binding to an epitope in the catalytic domain of HAAH. Other compounds such as L-mimosine or hydroxypyridone are administered directly into a tumor site or systemically to inhibit HAAH hydroxylase activity.

TABLE 1

Amino acid sequence of HAAH	
MAQRKNAKSS GNSSSSSGSGS GSTSAGSSSP GARRETKHGG HKNGRKGGLS GTSFFTWFMV	61
IALLGWVTSV AVVWFDLVGY EEVLGKGLGIY DADGDGDFDV DDAKVLLGLK ERSTSEPAVP	121
PEEAEPHTEP EEQVPVEAEP QNIEDEAKEQ IQSLLHEMVH AEHVEGEDLQ QEDGPTGEPQ	181
QEDDEFLMAT DVDDRPFETLE PEVSHEETEH SYHVEETVSQ DCNQDMEEMM SEQENPDSSSE	241
PVVEDERLHH DTDDVTYQVY EEQAVYEPL ENEGIEITEVT APPEDNPVED SQVIVEEVSI	301
FPVEEQQVVP PETNRKTDPP EQKAKVKKKK PKLLNKFDKT IKAELDAAEK LRKRKGKIEEA	361
VNAFKELVRK YPQSPRARYG KAQCEDDLAE KRSNEVLRG AIETYQEVAS LPDVPADLLK	421
LSLKRRSDRQ QFLGHMRGSL LTLQRLVQLF PNDTSLKNDL GVGYLLIGDN DNAKKVYEEV	481
LSVTPNMGFA KVHYGFILKA QNKIAESIPY LKEGIESGDP GTDDGRFYFH LGDAMQVRGN	541
KEAYKWYELG HKRGHFASVW QRSLYNVNGL KAQPWWTPKE TGYTELKLSL ERNWKILRDE	601
GLAVMDKAKG LFLPEDENLR EKGDSQFTL WQQGRRNENA CKGAPKTCTL LEKFPTTGC	661
RRGQIKYSIM HPGTHVWPHT GPTNCRLRMH LGLVIPKEGC KIRCANETRT WEEGKVLIFF	721
DSFEHEVWQD ASSFRLTFIV DVWHPELTPQ QRRSLPAI	
(SEQ ID NO: 2; GENBANK Accession No. S83325; His motif is underlined; conserved sequences within the catalytic domain are designated by bold type)	

as a separate molecule (e.g., a cDNA or a genomic or cDNA fragment produced by PCR or restriction endonuclease digestion) independent of other sequences. It also includes a recombinant nucleic acid which is part of a hybrid gene encoding additional polypeptide sequence such as a nucleic acid encoding an chimeric polypeptide, e.g., one encoding an antibody fragment linked to a cytotoxic polypeptide. Alternatively, HAAH expression is inhibited by administering a ribozyme or a compound which inhibits binding of Fos or Jun to an HAAH promoter sequence.

[0011] Compounds, which inhibit an enzymatic activity of HAAH, are useful to inhibit tumor growth in a mammal. By enzymatic activity of HAAH is meant hydroxylation of an epidermal growth factor (EGF)-like domain of a polypeptide. For example an EGF-like domain has the consensus sequence CX₂CX₄CX₁₀CXCX₈C (SEQ ID NO:1). HAAH hydroxylase activity is inhibited intracellularly. For example, a dominant negative mutant of HAAH (or a nucleic acid encoding such a mutant) is administered. The dominant negative HAAH mutant contains a mutation which changes a ferrous iron binding site from histidine of a naturally-occurring HAAH sequence to a non-iron-binding amino acid, thereby abolishing the hydroxylase activity of HAAH. The histidine to be mutated, e.g., deleted or substituted, is located in the carboxyterminal catalytic domain of HAAH. For example, the mutation is located between amino acids 650-700 (such as the His motif, underlined sequence of SEQ ID NO:2) the native HAAH sequence. For example, the mutation is at residues 671, 675, 679, or 690 of SEQ ID NO:2. An HAAH-

[0012] For example, a compound which inhibits HAAH hydroxylation is a polypeptide that binds a HAAH ligand but does not transduce an intracellular signal or a polypeptide which contains a mutation in the catalytic site of HAAH. Such a polypeptide contains an amino acid sequence that is at least 50% identical to a naturally-occurring HAAH amino acid sequence or a fragment thereof and which has the ability to inhibit HAAH hydroxylation of substrates containing an EGF-like repeat sequence. More preferably, the polypeptide contains an amino acid sequence that is at least 75%, more preferably at least 85%, more preferably at least 95% identical to SEQ ID NO: .

[0013] A substantially pure HAAH polypeptide or HAAH-derived polypeptide such as a mutated HAAH polypeptide is preferably obtained by expression of a recombinant nucleic acid encoding the polypeptide or by chemically synthesizing the protein. A polypeptide or protein is substantially pure when it is separated from those contaminants which accompany it in its natural state (proteins and other naturally-occurring organic molecules). Typically, the polypeptide is substantially pure when it constitutes at least 60%, by weight, of the protein in the preparation. Preferably, the protein in the preparation is at least 75%, more preferably at least 90%, and most preferably at least 99%, by weight, HAAH. Purity is measured by any appropriate method, e.g., column chromatography, polyacrylamide gel electrophoresis, or HPLC analysis. Accordingly, substantially pure polypeptides include recombinant polypeptides derived from a eucaryote but produced in *E. coli* or another procaryote, or in a eucaryote other than that from which the polypeptide was originally derived.

[0014] Nucleic acid molecules which encode such HAAH or HAAH-derived polypeptides are also within the invention.

TABLE 2

HAAH cDNA sequence	
cggaccgtgc aatggcccag cgtaagaatg ccaagagcag cggcaacagc agcagcagcg	61
gctccggcag cggtagcacg agtgcgggca gcagcagccc cggggcccg agagagacaa	121
agcatggagg acacaagaat gggaggaaag gcggactctc gggaacttca ttcttcacgt	181
ggtttatggt gattgcattg ctgggcgtct ggacatctgt agctgtcgtt tggtttgatc	241
ttgttgacta tgaggaaagt ctaggaaaac taggaatcta tgatgctgat ggtgatggag	301
atthttgatgt ggatgatgcc aaagtthttat taggacttaa agagagatct acttcagagc	361
cagcagtcgc gccagaagag gctgagccac aactgagcc cgaggagcag gttcctgtgg	421
aggcagaacc ccagaatata gaagatgaag caaaagaaca aattcagtcc cttctccatg	481
aatgggtaca cgcagaacat gttgagggag aagacttgca acaagaagat ggaccacacag	541
gagaaccaca acaagaggat gatgagtttc ttatggcgac tgatgtagat gatagatttg	601
agaccctgga acctgaagta tctcatgaag aaaccgagca tagttaccac gtggaagaga	661
cagtttcaca agactgtaat caggatatgg aagagatgat gtctgagcag gaaaatccag	721
attccagtga accagtagta gaagatgaaa gattgcacca tgatacagat gatgtaacat	781
accaagtcta tgaggaaaca gcagtatatg aacctctaga aaatgaaggg atagaaatca	841
cagaagtaac tgctccccc gaggataatc ctgtagaaga ttcacaggta attgtagaag	901
aagtaagcat tttctctgtg gaagaacagc aggaagtacc accagaaaca aatagaaaaa	961
cagatgatcc agaacaaaa gcaaaagtta agaaaagaa gcctaaactt ttaaataaat	1021
ttgataagac tattaaagct gaacttgatg ctgcagaaaa actccgtaaa aggggaaaaa	1081
ttgaggaagc agtgaatgca tttaaagaac tagtacgcaa ataccctcag agtccacgag	1141
caagatatgg gaaggcgcag tgtgaggatg atttggtgga gaagaggaga agtaatgagg	1201
tgctacgtgg agccatcgag acctaccaag aggtggccag cctacctgat gtccctgcag	1261
acctgctgaa gctgagtttg aagcgtcgtc cagacaggca acaatttcta ggtcatatga	1321
gaggttccct gcttaccctg cagagattag ttcaactatt tccaatgat acttccttaa	1381
aaaatgacct tggcgtggga tacctcttga taggagataa tgacaatgca aagaaagttt	1441
atgaagaggt gctgagtgtg acacctaatg atggctttgc taaagtccat tatggcttca	1501
tcctgaaggc acagaacaaa attgctgaga gcacccata tttaaaggaa ggaatagaat	1561
ccggagatcc tggcactgat gatgggagat tttatttcca cctgggggat gccatgcaga	1621
gggttgggaa caaagaggca tataagtggg atgagcttgg gcacaagaga ggacactttg	1681
catctgtctg gcaacgctca ctctacaatg tgaatggact gaaagcacag ccttggtgga	1741
ccccaaaaga aacgggctac acagagttag taaagtcttt agaagaaaac tggaaagtaa	1801
tccgagatga aggccttgca gtgatggata aagccaaagg tctcttcctg cctgaggatg	1861
aaaacctgag ggaagaaagg gactggagcc agttcacgct gtggcagcaa ggaagaagaa	1921
atgaaaatgc ctgcaaaagg gctcctaaaa cctgtacctt actagaaaag tccccgaga	1981
caacaggatg cagaagagga cagatcaaat attccatcat gcacccggg actcacgtgt	2041
ggccgcacac agggcccaca aactgcaggc tccgaatgca cctgggcttg gtgattccca	2101
aggaaggtcg caagattcga tgtgccaacg agaccaggac ctgggaggaa ggcaaggtgc	2161

TABLE 2-continued

HAAH cDNA sequence
tcacatcttga tgactccttt gagcacgagg tatggcagga tgcctcatct ttccggctga 2221
tattcatcgt ggaagtgtgg catccggaac tgacaccaca gcagagacgc agccttccag 2281
caatttagca tgaattcatg caagcttggg aaactctgga gaga
(SEQ ID NO:3; GENBANK Accession No. S83325; codon encoding
initiating methionine is underlined).

[0015] Methods of inhibiting tumor growth also include administering a compound which inhibits HAAH hydroxylation of a NOTCH polypeptide. For example, the compound inhibits hydroxylation of an EGF-like cysteine-rich repeat sequence in a NOTCH polypeptide, e.g., one containing the consensus sequence CDXXXCXXKXGNGXCDXXCN-NAACXXDGXDC (SEQ ID NO:4). Polypeptides containing an EGF-like cysteine-rich repeat sequence are administered to block hydroxylation of endogenous NOTCH.

[0016] Growth of a tumor which overexpresses HAAH is also inhibited by administering a compound which inhibits signal transduction through the insulin receptor substrate (IRS) signal transduction pathway. Preferably the compound inhibits IRS phosphorylation. For example, the compound is a peptide or non-peptide compound which binds to and inhibits phosphorylation at residues 46, 465, 551, 612, 632, 662, 732, 941, 989, or 1012 of SEQ ID NO:5. Compounds include polypeptides such as those which block an IRS phosphorylation site such as a Glu/Tyr site. Antibodies such as those which bind to a carboxyterminal domain of IRS containing a phosphorylation site block IRS phosphorylation, and as a consequence, signal transduction along the pathway. Inhibition of IRS phosphorylation in turn leads to inhibition of cell proliferation. Other compounds which inhibit IRS phosphorylation include vitamin D analogue EB1089 and Wortmannin.

[0017] HAAH-overproducing tumor cells were shown to express HAAH both intracellularly and on the surface of the tumor cell. Accordingly, a method of killing a tumor cell is carried out by contacting such a tumor cell with a cytotoxic agent linked to an HAAH-specific antibody. The HAAH-specific antibody (antibody fragment, or ligand which binds to extracellular HAAH) directs the chimeric polypeptide to the surface of the tumor cell allowing the cytotoxic agent to damage or kill the tumor cell to which the antibody is bound. The monoclonal antibody binds to an epitope of HAAH such as an epitope exposed on the surface of the cell or in the catalytic site of HAAH. The cytotoxic composition preferentially kills tumor cells compared to non-tumor cell.

[0018] Screening methods to identify anti-tumor agents which inhibit the growth of tumors which overexpress HAAH are also within the invention. A screening method used to determine whether a candidate compound inhibits HAAH enzymatic activity includes the following steps: (a) providing a HAAH polypeptide, e.g., a polypeptide which contains the carboxyterminal catalytic site of HAAH; (b) providing a polypeptide comprising an EGF-like domain; (c) contacting the HAAH polypeptide or the EGF-like polypeptide with the candidate compound; and (d) determining hydroxylation of the EGF-like polypeptide of step (b). A decrease in hydroxylation in the presence of the candidate compound compared to

that in the absence of said compound indicates that the compound inhibits HAAH hydroxylation of EGF-like domains in proteins such as NOTCH.

[0019] Anti-tumor agents which inhibit HAAH activation of NOTCH are identified by (a) providing a cell expressing HAAH; (b) contacting the cell with a candidate compound; and (c) measuring translocation of activated NOTCH to the nucleus of said cell. Translocation is measured by using a reagent such as an antibody which binds to a 110 kDa activation fragment of NOTCH. A decrease in translocation in the presence of the candidate compound compared to that in the absence of the compound indicates that the compound inhibits HAAH activation of NOTCH, thereby inhibiting NOTCH-mediated signal transduction and proliferation of HAAH-overexpressing tumor cells.

[0020] Nucleotide and amino acid comparisons described herein were carried out using the Lasergene software package (DNASTAR, Inc., Madison, Wis.). The MegAlign module used was the Clustal V method (Higgins et al., 1989, CABIOS 5(2):151-153). The parameter used were gap penalty 10, gap length penalty 10.

[0021] Hybridization is carried out using standard techniques, such as those described in Ausubel et al. (*Current Protocols in Molecular Biology*, John Wiley & Sons, 1989). "High stringency" refers to nucleic acid hybridization and wash conditions characterized by high temperature and low salt concentration, e.g., wash conditions of 65° C. at a salt concentration of 0.1×SSC. "Low" to "moderate" stringency refers to DNA hybridization and wash conditions characterized by low temperature and high salt concentration, e.g., wash conditions of less than 60° C. at a salt concentration of at least 1.0×SSC. For example, high stringency conditions include hybridization at 42° C. in the presence of 50% formamide; a first wash at 65° C. in the presence of 2×SSC and 1% SDS; followed by a second wash at 65° C. in the presence of 0.1×SSC. Lower stringency conditions suitable for detecting DNA sequences having about 50% sequence identity to an HAAH gene sequence are detected by, for example, hybridization at about 42° C. in the absence of formamide; a first wash at 42° C., 6×SSC, and 1% SDS; and a second wash at 50° C., 6×SSC, and 1% SDS.

[0022] Other features and advantages of the invention will be apparent from the following description of the preferred embodiments thereof, and from the claims.

BRIEF DESCRIPTION OF THE DRAWINGS

[0023] FIG. 1 is a bar graph showing colony formation induced by transient transfection of NIH-3T3 cells with various HAAH cDNAs. Colony formation was induced by transient transfection with 10 µg DNA. In contrast, the mutant murine

AAH construct without enzymatic activity has no transforming activity. The data is presented as mean number of transformed foci \pm SEM.

[0024] FIG. 2 is a bar graph showing the results of a densitometric analysis of a Western blot assay of proteins produced by various murine AAH stably transfected cell clones. In clones 7 and 18, there was a modest increase in HAAH gene expression, while the overexpression was to a lesser degree in clone 16.

[0025] FIGS. 3A-B are bar graphs showing colony formation in soft agar exhibited by HAAH stably transfected clones compared to HAAH enzymatic activity. FIG. 3A shows a measurement of murine AAH enzymatic activity in clones 7, 16 and 18, and FIG. 3B shows colony formation exhibited by clones 7, 16 and 18. Data is presented as mean number of colonies 10 days after plating \pm SEM. All three clones with modest increases in HAAH enzymatic activity, that correlated with protein expression, exhibited anchorage independent growth.

[0026] FIG. 4 is a bar graph showing tumor formation in nude mice injected with transfected clones overexpressing murine AAH. Tumor growth was assessed after 30 days. Mean tumor weight observed in mice injected with clones 7, 16 and 18 as compared to mock DNA transfected clone. All animals injected with clones overexpressing HAAH developed tumors.

[0027] FIGS. 5A-D are bar graphs showing increased AAH expression in PNET2 (FIGS. 5A, 5C) and SH-Sy5y (FIG. 5B) cells treated with retinoic acid (FIGS. 5A, 5B) or phorbol ester myristate (PMA; FIG. 5C) to induce neurite outgrowth as occurs during tumor cell invasion. The cells were treated with 10 μ M retinoic acid or 100 nM PMA for 0, 1, 2, 3, 4, or 7 days. Cell lysates were analyzed by Western blot analysis using an HAAH-specific monoclonal antibody to detect the 85 kDa AAH protein. The levels of immunoreactivity were measured by volume densitometry (arbitrary units). The graphs indicate the mean \pm S.D. of results obtained from three separate experiments. In FIG. 5D, PNET2 cells were treated for 24 hours with sub-lethal concentrations of H₂O₂ to induce neurite retraction. Viability of greater than 90% of the cells was demonstrated by Trypan blue dye exclusion. Similar results were obtained for SH-Sy5y cells.

[0028] FIG. 6 is a bar graph showing the effects of AAH over-expression on the levels of anti-apoptosis (Bcl-2), cell cycle-mitotic inhibitor (p16 and p21/Waf1), and proliferation (proliferating cell nuclear antigen; PCNA) molecules. PNET2 neuronal cells were stably transfected with the full-length human cDNA encoding AAH (pHAAH) or empty vector (pcDNA). AAH gene expression was under control of a CMV promoter. Western blot analysis was performed with cell lysates prepared from cultures that were 70 to 80 percent confluent. Protein loading was equivalent in each lane. Replicate blots were probed with the different antibodies. Bar graphs depict the mean S.D.'s of protein expression levels measured in three experiments. All differences are statistically significant by Student T-test analysis ($P < 0.01$ – $P < 0.001$).

[0029] FIG. 7 is a diagram of showing the components of the IRS-1 signal transduction pathway.

[0030] FIG. 8 is a line graph showing growth curves generated in cells expressing the antisense HAAH compared to controls expressing GFP.

[0031] FIG. 9 is a diagram of the functional domains of the hIRS-1 protein and structural organization of the point

mutants. All mutant and "wild type" hIRS-1 proteins construct contain a FLAG (F) epitope (DYKDDDDK; SEQ ID NO:7) at the C-terminus. PH and PTB indicate pleckstrin homology and phosphotyrosine binding, regions, respectively.

DETAILED DESCRIPTION

[0032] HAAH is a protein belonging to the (α -ketoglutarate dependent dioxygenase family of prolyl and lysyl hydroxylases which play a key role in collagen biosynthesis. This molecule hydroxylates aspartic acid or asparagine residues in EGF-like domains of several proteins in the presence of ferrous iron. These EGF-like domains contain conserved motifs, that form repetitive sequences in proteins such as clotting factors, extracellular matrix proteins, LDL receptor, NOTCH homologues or NOTCH ligand homologues.

[0033] The alpha-ketoglutarate-dependent dioxygenase aspartyl (asparaginyl) beta-hydroxylase (AAH) specifically hydroxylates one aspartic or asparagine residue in EGF-like domains of various proteins. The 4.3-kb cDNA encoding the human AspH (hAspH) hybridizes with 2.6 kb and 4.3 kb transcripts in transformed cells, and the deduced amino acid sequence of the larger transcript encodes a protein of about kDa. Both in vitro transcription and translation and Western blot analysis also demonstrate a 56-kDa protein that may result from posttranslational cleavage of the catalytic C terminus.

[0034] An physiological function of AAH is the post-translational beta-hydroxylation of aspartic acid in vitamin K-dependent coagulation proteins. However, the abundant expression of AAH in several malignant neoplasms, and low levels of AAH in many normal cells indicate a role for this enzyme in malignancy. The AAH gene is also highly expressed in cytotrophoblasts, but not syncytiotrophoblasts of the placenta. Cytotrophoblasts are invasive cells that mediate placental implantation. The increased levels of AAH expression in human cholangiocarcinomas, hepatocellular carcinomas, colon cancers, and breast carcinomas were primarily associated with invasive or metastatic lesions. Moreover, overexpression of AAH does not strictly reflect increased DNA synthesis and cellular proliferation since high levels of AAH immunoreactivity were observed in 100 percent of cholangiocarcinomas, but not in human or experimental disease processes associated with regeneration or nonneoplastic proliferation of bile ducts. AAH overexpression and attendant high levels of beta hydroxylase activity lead to invasive growth of transformed neoplastic cells. Detection of an increase in HAAH expression is useful for early and reliable diagnosis of the cancer types which have now been characterized as over-expressing this gene product.

Diagnosis of Malignant Tumors

[0035] HAAH is overexpressed in many tumors of endodermal origin and in at least 95% of CNS tumors compared to normal noncancerous cells. An increase in HAAH gene product in a patient-derived tissue sample (e.g., solid tissue or bodily fluid) is carried out using standard methods, e.g., by Western blot assays or a quantitative assay such as ELISA. For example, a standard competitive ELISA format using an HAAH-specific antibody is used to quantify patient HAAH levels. Alternatively, a sandwich ELISA using a first antibody as the capture antibody and a second HAAH-specific antibody as a detection antibody is used.

[0036] Methods of detecting HAAH include contacting a component of a bodily fluid with an HAAH-specific antibody bound to solid matrix, e.g., microtiter plate, bead, dipstick. For example, the solid matrix is dipped into a patient-derived sample of a bodily fluid, washed, and the solid matrix is contacted with a reagent to detect the presence of immune complexes present on the solid matrix.

[0037] Proteins in a test sample are immobilized on (bound to) a solid matrix. Methods and means for covalently or noncovalently binding proteins to solid matrices are known in the art. The nature of the solid surface may vary depending upon the assay format. For assays carried out in microtiter wells, the solid surface is the wall of the well or cup. For assays using beads, the solid surface is the surface of the bead. In assays using a dipstick (i.e., a solid body made from a porous or fibrous material such as fabric or paper) the surface is the surface of the material from which the dipstick is made. Examples of useful solid supports include nitrocellulose (e.g., in membrane or microtiter well form), polyvinyl chloride (e.g., in sheets or microtiter wells), polystyrene latex (e.g., in beads or microtiter plates, polyvinylidene fluoride (known as IMMULON™), diazotized paper, nylon membranes, activated beads, and Protein A beads. The solid support containing the antibody is typically washed after contacting it with the test sample, and prior to detection of bound immune complexes. Incubation of the antibody with the test sample is followed by detection of immune complexes by a detectable label. For example, the label is enzymatic, fluorescent, chemiluminescent, radioactive, or a dye. Assays which amplify the signals from the immune complex are also known in the art, e.g., assays which utilize biotin and avidin.

[0038] An HAAH-detection reagent, e.g., an antibody, is packaged in the form of a kit, which contains one or more HAAH-specific antibodies, control formulations (positive and/or negative), and/or a detectable label. The assay may be in the form of a standard two-antibody sandwich assay format known in the art.

Production of HAAH-Specific Antibodies

[0039] Anti-HAAH antibodies were obtained by techniques well known in the art. Such antibodies are polyclonal or monoclonal. Polyclonal antibodies were obtained, for example, by the methods described in Ghose et al., *Methods in Enzymology*, Vol. 93, 326-327, 1983. An HAAH polypeptide, or an antigenic fragment thereof, was used as the immunogen to stimulate the production of polyclonal antibodies in the antisera of rabbits, goats, sheep, or rodents. Antigenic polypeptides for production of both polyclonal and monoclonal antibodies useful as immunogens include polypeptides which contain an HAAH catalytic domain. For example, the immunogenic polypeptide is the full-length mature HAAH protein or an HAAH fragment containing the carboxyterminal catalytic domain e.g., an HAAH polypeptide containing the His motif of SEQ ID NO:2.

[0040] Antibodies which bind to the same epitopes as those antibodies disclosed herein as identified using standard methods, e.g., competitive binding assays, known in the art.

[0041] Monoclonal antibodies were obtained by standard techniques. Ten µg of purified recombinant HAAH polypeptide was administered to mice intraperitoneally in complete Freund's adjuvant, followed by a single boost intravenously (into the tail vein) 3-5 months after the initial inoculation. Antibody-producing hybridomas were made using standard methods. To identify those hybridomas producing antibodies

that are highly specific for an HAAH polypeptide, hybridomas were screened using the same polypeptide immunogen used to immunize. Those antibodies which were identified as having HAAH-binding activity are also screened for the ability to inhibit HAAH catalytic activity using the enzymatic assays described below. Preferably, the antibody has a binding affinity of at least about 10^8 liters/mole and more preferably, an affinity of at least about 10^9 liters/mole.

[0042] Monoclonal antibodies are humanized by methods known in the art, e.g., MAbs with a desired binding specificity can be commercially humanized (Scotgene, Scotland; Oxford Molecular, Palo Alto, Calif.).

[0043] HAAH-specific intrabodies are produced as follows. Following identification of a hybridoma producing a suitable monoclonal antibody, DNA encoding the antibody is cloned. DNA encoding a single chain HAAH-specific antibody in which heavy and light chain variable domains are separated by a flexible linker peptide is cloned into an expression vector using known methods (e.g., Marasco et al., 1993, *Proc. Natl. Acad. Sci. USA* 90:7889-7893 and Marasco et al., 1997, *Gene Therapy* 4:11-15). Such constructs are introduced into cells, e.g., using standard gene delivery techniques for intracellular production of the antibodies. Intracellular antibodies, i.e., intrabodies, are used to inhibit signal transduction by HAAH. Intrabodies which bind to a carboxyterminal catalytic domain of HAAH inhibit the ability of HAAH to hydroxylate EGF-like target sequences.

[0044] Methods of linking HAAH-specific antibodies (or fragments thereof) which bind to cell surface exposed epitopes of HAAH on the surface of a tumor cell are linked to known cytotoxic agents, e.g., ricin or diphtheria toxin, using known methods.

[0045] Deposit of Biological Materials

[0046] Under the terms of the Budapest Treaty on the International Recognition of the Deposit of Microorganisms for the Purpose of Patent Procedure, hybridoma FB501 (which produces monoclonal antibody FB50; designated ATCC accession no. PTA 3386), hybridoma HA386A (which produces monoclonal antibody 86A; designated ATCC accession no. 3385), hybridoma HA15C7A (which produces monoclonal antibody 5C7; designated ATCC accession no. 3383), and hybridoma HA219B (which produces monoclonal antibody 19B; designated ATCC accession no. 3384) were deposited on May 17, 2001, with the American Type Culture Collection (ATCC) of 10801 University Boulevard, Manassas, Va. 20110-2209 USA.

[0047] Applicants' assignee represents that the ATCC is a depository affording permanence of the deposit and ready accessibility thereto by the public if a patent is granted. All restrictions on the availability to the public of the material so deposited will be irrevocably removed upon the granting of a patent. The material will be available during the pendency of the patent application to one determined by the Commissioner to be entitled thereto under 37 CFR 1.14 and 35 U.S.C. 122. The deposited material will be maintained with all the care necessary to keep it viable and uncontaminated for a period of at least five years after the most recent request for the furnishing of a sample of the deposited plasmid, and in any case, for a period of at least thirty (30) years after the date of deposit or for the enforceable life of the patent, whichever period is longer.

[0048] Applicant's assignee acknowledges its duty to replace the deposit should the depository be unable to furnish a sample when requested due to the condition of the deposit.

Methods of Treating Malignant Tumors

[0049] Patients with tumors characterized as overexpressing HAAH as such tumors of endodermal origin or CNS tumors are treated by administering HAAH antisense nucleic acids.

[0050] Antisense therapy is used to inhibit expression of HAAH in patients suffering from hepatocellular carcinomas, cholangiocarcinomas, glioblastomas and neuroblastomas. For example, an HAAH antisense strand (either RNA or DNA) is directly introduced into the cells in a form that is capable of binding to the mRNA transcripts. Alternatively, a vector containing a sequence which, which once within the target cells, is transcribed into the appropriate antisense mRNA, may be administered. Antisense nucleic acids which hybridize to target mRNA decrease or inhibit production of the polypeptide product encoded by a gene by associating with the normally single-stranded mRNA transcript, thereby interfering with translation and thus, expression of the protein. For example, DNA containing a promoter, e.g., a tissue-specific or tumor specific promoter, is operably linked to a DNA sequence (an antisense template), which is transcribed into an antisense RNA. By "operably linked" is meant that a coding sequence and a regulatory sequence(s) (i.e., a promoter) are connected in such a way as to permit gene expression when the appropriate molecules (e.g., transcriptional activator proteins) are bound to the regulatory sequence(s).

[0051] Oligonucleotides complementary to various portions of HAAH mRNA are tested in vitro for their ability to decrease production of HAAH in tumor cells (e.g., using the FOCUS hepatocellular carcinoma (HCC) cell line) according to standard methods. A reduction in HAAH gene product in cells contacted with the candidate antisense composition compared to cells cultured in the absence of the candidate composition is detected using HAAH-specific antibodies or other detection strategies. Sequences which decrease production of HAAH in in vitro cell-based or cell-free assays are then be tested in vivo in rats or mice to confirm decreased HAAH production in animals with malignant neoplasms.

[0052] Antisense therapy is carried out by administering to a patient an antisense nucleic acid by standard vectors and/or gene delivery systems. Suitable gene delivery systems may include liposomes, receptor-mediated delivery systems, naked DNA, and viral vectors such as herpes viruses, retroviruses, adenoviruses and adeno-associated viruses, among others. A reduction in HAAH production results in a decrease in signal transduction via the IRS signal transduction pathway. A therapeutic nucleic acid composition is formulated in a pharmaceutically acceptable carrier. The therapeutic composition may also include a gene delivery system as described above. Pharmaceutically acceptable carriers are biologically compatible vehicles which are suitable for administration to an animal: e.g., physiological saline. A therapeutically effective amount of a compound is an amount which is capable of producing a medically desirable result such as reduced production of an HAAH gene product or a reduction in tumor growth in a treated animal.

[0053] Parenteral administration, such as intravenous, subcutaneous, intramuscular, and intraperitoneal delivery routes, may be used to deliver nucleic acids or HAAH-inhibitory peptides or non-peptide compounds. For treatment of CNS

tumors, direct infusion into cerebrospinal fluid is useful. The blood-brain barrier may be compromised in cancer patients, allowing systemically administered drugs to pass through the barrier into the CNS. Liposome formulations of therapeutic compounds may also facilitate passage across the blood-brain barrier.

[0054] Dosages for any one patient depends upon many factors, including the patient's size, body surface area, age, the particular nucleic acid to be administered, sex, time and route of administration, general health, and other drugs being administered concurrently. Dosage for intravenous administration of nucleic acids is from approximately 10^6 to 10^{22} copies of the nucleic acid molecule.

[0055] Ribozyme therapy is also be used to inhibit HAAH gene expression in cancer patients. Ribozymes bind to specific mRNA and then cut it at a predetermined cleavage point, thereby destroying the transcript. These RNA molecules are used to inhibit expression of the HAAH gene according to methods known in the art (Sullivan et al., 1994, J. Invest. Derm. 103:85 S-89S; Czubayko et al., 1994, J. Biol. Chem. 269:21358-21363; Mahieu et al., 1994, Blood 84:3758-65; Kobayashi et al. 1994, Cancer Res. 54:1271-1275).

Methods of Identifying Compounds that Inhibit HAAH Enzymatic Activity

[0056] Aspartyl (asparaginyl) beta-hydroxylase (HAAH) activity is measured in vitro or in vivo. For example, HAAH catalyzes posttranslational modification of β carbon of aspartyl and asparaginyl residues of EGF-like polypeptide domains. An assay to identify compounds which inhibit hydroxylase activity is carried out by comparing the level of hydroxylation in an enzymatic reaction in which the candidate compound is present compared to a parallel reaction in the absence of the compound (or a predetermined control value). Standard hydroxylase assays carried out in a test-tube are known in the art, e.g., Lavaissiere et al., 1996, J. Clin. Invest. 98:1313-1323; Jia et al., 1992, J. Biol. Chem. 267:14322-14327; Wang et al., 1991, J. Biol. Chem. 266:14004-14010; or Gronke et al., 1990, J. Biol. Chem. 265:8558-8565. Hydroxylase activity is also measured using carbon dioxide ($^{14}\text{CO}_2$, capture assay) in a 96-well microtiter plate format (Zhang et al., 1999, Anal. Biochem. 271:137-142). These assays are readily automated and suitable for high throughput screening of candidate compounds to identify those with hydroxylase inhibitory activity.

[0057] Candidate compound which inhibit HAAH activation of NOTCH are identified by detecting a reduction in activated NOTCH in a cell which expresses or overexpresses HAAH, e.g., FOCUS HCC cells. The cells are cultured in the presence of a candidate compound. Parallel cultures are incubated in the absence of the candidate compound. To evaluate whether the compound inhibits HAAH activation of NOTCH, translocation of activated NOTCH to the nucleus of the cell is measured. Translocation is measured by detecting a 110 kDa activation fragment of NOTCH in the nucleus of the cell. The activation fragment is cleaved from the large (approximately 300 kDa) transmembrane NOTCH protein upon activation. Methods of measuring NOTCH translocation are known, e.g., those described by Song et al., 1999, Proc. Natl. Acad. Sci. U.S.A. 96:6959-6963 or Capobianco et al., 1997, Mol. Cell. Biol. 17:6265-6273. A decrease in translocation in the presence of the candidate compound compared to that in the absence of the compound indicates that the compound inhibits

its HAAH activation of NOTCH, thereby inhibiting NOTCH-mediated signal transduction and proliferation of HAAH-overexpressing tumor cells.

[0058] Methods of screening for compounds which inhibit phosphorylation of IRS are carried out by incubating IRS-expressing cells in the presence and absence of a candidate compound and evaluating the level of IRS phosphorylation in the cells. A decrease in phosphorylation in cells cultured in the presence of the compound compared to in the absence of the compound indicates that the compound inhibits IRS-1 phosphorylation, and as a result, growth of HAAH-overexpressing tumors. Alternatively, such compounds are identified in an *in vitro* phosphorylation assay known in the art, e.g., one which measured phosphorylation of a synthetic substrate such as poly(Glu/Tyr).

Example 1

Increased Expression of HAAH is Associated with Malignant Transformation

[0059] HAAH is a highly conserved enzyme that hydroxylates EGF-like domains in transformation associated proteins. The HAAH gene is overexpressed in human hepatocellular carcinomas and cholangiocarcinomas. HAAH gene expression was found to be undetectable during bile duct proliferation in both human disease and rat models compared to cholangiocarcinoma. Overexpression of HAAH in NIH-3T3 cells was associated with generation of a malignant phenotype, and enzymatic activity was found to be required for cellular transformation. The data described below indicate that overexpression of HAAH is linked to cellular transformation of biliary epithelial cells.

[0060] To identify molecules that are specifically overexpressed in transformed malignant cells of human hepatocyte origin, the FOCUS hepatocellular carcinoma (HCC) cell line was used as an immunogen to generate monoclonal antibodies (mAb) that specifically or preferentially recognize proteins associated with the malignant phenotype. A lambda GT11 cDNA expression library derived from HepG2 HCC cells was screened, and HAAH-specific mAb produced against the FOCUS cell line was found to recognize an epitope on a protein encoded by an HAAH cDNA. The HAAH enzyme was found to be upregulated in several different human transformed cell lines and tumor tissues compared to adjacent human tissue counterparts. The overexpressed HAAH enzyme in different human malignant tissues was found to be catalytically active.

[0061] HAAH gene expression was examined in proliferating bile ducts and in NIH 3T3 cells. Its role in the generation of the malignant phenotype was measured by the formation of transformed foci, growth in soft agar as an index of anchorage independent growth and tumor formation in nude mice. The role of enzymatic activity in the induction of transformed phenotype was measured by using a cDNA construct with a mutation in the catalytic site that abolished hydroxylase activity. The results indicated that an increase in expression of HAAH gene is associated with malignant transformation of bile ducts.

[0062] The following materials and methods were used to generate the data described below.

[0063] Antibodies

[0064] The FB50 monoclonal antibody was generated by cellular immunization of Balb/C mice with FOCUS HCC cells. A monoclonal anti-Dengue virus antibody was used as

a non-relevant control. The HBOH2 monoclonal antibody was generated against a 52 kDa recombinant HAAH polypeptide and recognizes the catalytic domain of beta-hydroxylase from mouse and human proteins. Polyclonal anti-HAAH antibodies cross-react with rat hydroxylase protein. Control antibody anti-Erk-1 was purchased from Santa Cruz Biotechnology, Inc, CA. Sheep anti-mouse and donkey anti-rabbit antisera labeled with horseradish peroxidase were obtained from Amersham, Arlington Heights, Ill.

[0065] Constructs

[0066] The murine full length AAH construct (pNH376) and the site-directed mutation construct (pNH376-H660) with abolished catalytic activity were cloned into the eukaryotic expression vector pcDNA3 (Invitrogen Corp., San Diego, Calif.). The full length human AAH was cloned into prokaryotic expression vector pBC-SK+ (Stratagene, La Jolla, Calif.). The full length human AAH (GENBANK Accession No. 583325) was subcloned into the EcoRI site of the pcDNA3 vector.

[0067] Animal Model of Bile Duct Proliferation

[0068] Rats were divided into 9 separate groups of 3 animals each except for group 9 which contained 5 rats. Group 1 was the non-surgical control group, and group 2 was the sham-operated surgical control. The remaining groups underwent common bile duct ligation to induce intrahepatic bile duct proliferation and were evaluated at 6, 12, 24, 48 hours and 4, 8 and 16 days as shown in Table 3. Animals were asphyxiated with CO₂, and liver samples were taken from left lateral and median lobes, fixed in 2% paraformaldehyde and embedded in paraffin. Liver samples (5 µm) were cut and stained with hematoxylin and eosin to evaluate intrahepatic bile duct proliferation. Immunohistochemistry was performed with polyclonal anti-HAAH antibodies that cross-react with the rat protein to determine levels of protein expression.

[0069] Bile Duct Proliferation Associated with Primary Sclerosing Cholangitis (PSC)

[0070] Liver biopsy samples were obtained from 7 individuals with PSC and associated bile duct proliferation. These individuals were evaluated according to standard gastroenterohepatological protocols. Patients were 22-46 years of age and consisted of 4 males and 3 females. Four had associated inflammatory bowel disease (3 ulcerative colitis and 1 Crohn's colitis). All patients underwent a radiological evaluation including abdominal ultrasonography and endoscopic retrograde cholangiopancreatography to exclude the diagnosis of extrahepatic biliary obstruction. Tissue sections were prepared from paraffin embedded blocks and were evaluated by hematoxylin and eosin staining for bile duct proliferation. Expression of HAAH was determined by immunohistochemistry using an HAAH-specific monoclonal antibody such as FB50.

[0071] Immunohistochemistry

[0072] Liver tissue sections (5 µm) were deparaffinized in xylene and rehydrated in graded alcohol. Endogenous peroxidase activity was quenched by a 30-minute treatment with 0.6% H₂O₂ in 60% methanol. Endogenous biotin was masked by incubation with avidin-biotin blocking solutions (Vector Laboratories, Burlingame, Calif.). The FB50 mAb (for PSC samples) and polyclonal anti-HAAH-hydroxylase antibodies (for rat liver samples) were added to slides in a humidified chamber at 4° C. overnight. Immunohistochemical staining was performed using a standard avidin-biotin horseradish peroxidase complex (ABC) method using Vectastain Kits

with diaminobenzidine (DAB) as the chromogen according to manufacturer's instructions (Vector Laboratories, Inc., Burlingame, Calif.). Tissue sections were counterstained with hematoxylin, followed by dehydration in ethanol. Sections were examined by a light microscopy for bile duct proliferation and HAAH protein expression. Paraffin sections of cholangiocarcinoma and placenta were used as positive controls, and hepatosteatosis samples were used as a negative controls. To control for antibody binding specificity, adjacent sections were immunostained in the absence of a primary antibody, or using non-relevant antibody to Dengue virus. As a positive control for tissue immunoreactivity, adjacent sections of all specimens were immunostained with monoclonal antibody to glyceraldehyde 3-phosphate dehydrogenase.

[0073] Western Blot Analysis

[0074] Cell lysates were prepared in a standard radioimmunoprecipitation assay (RIPA) buffer containing protease inhibitors. The total amount of protein in the lysates was determined by Bio-Rad colorimetric assay (Bio Rad, Hercules, Calif.) followed by 10% sodium dodecyl sulphate-polyacrylamide gel electrophoresis (SDS-PAGE), transferred to PVDF membranes, and subjected to Western blot analysis using FB50, HBOH2, anti-Erk-1 (used as an internal control for protein loading) as primary, sheep anti-mouse and donkey anti-rabbit antisera labeled with horseradish peroxidase as secondary antibodies. Antibody binding was detected with enhanced chemiluminescence reagents (SuperSignal, Pierce Chemical Company, Rockford, Ill.) and film autoradiography. The levels of immunoreactivity were measured by volume densitometry using NIH Image software.

[0075] Enzymatic Activity Assay

[0076] AAH activity was measured in cell lysates using the first EGF-like domain of bovine protein S as substrate where ^{14}C -labeled α -ketoglutarate hydroxylates the domain releasing ^{14}C containing CO_2 according to standard methods, e.g., those described by Jia et al., 1992, J. Biol. Chem. 267:14322-14327; Wang et al., 1991, J. Biol. Chem. 266:14004-14010; or Gronke et al., 1990, J. Biol. Chem. 265:8558-8565. Incubations were carried out at 37° C. for 30 min in a final volume of 40 μl containing 48 μg of crude cell extract protein and 75 μM EGF substrate.

[0077] Cell Transfection Studies

[0078] The NIH-3T3 cells were cultured in Dulbecco's modified Eagle's medium (DMEM; Mediatech, Washington, D.C.) supplemented with 10% heat-inactivated fetal calf serum (FCS; Sigma Chemical Co., St. Louis, Mo.), 1% L-glutamine, 1% non-essential amino acids and 1% penicillin-streptomycin (GIBCO BRL, Life Technologies, Inc., Grand Island, N.Y.). Subconfluent NIH-3T3 cells (3×10^5 cells/60-mm dish) were transfected with 10 μg of one of the following plasmids: 1) non-recombinant pcDNA3 vector (Invitrogen Corp., San Diego, Calif.) as a negative control; 2) pNH376-H660, the murine AAH cDNA that was mutated in the catalytic domain and cloned into the pcDNA3 vector driven by a CMV promoter; 3) pNH376, the wild type murine AAH cDNA cloned into the pcDNA3 vector; 4) pCDHH, wild type human AAH cDNA cloned into the pcDNA3 vector; or 5) pLNCX-UP1, a cDNA that encodes v-Src oncogene (positive control). Cells were transfected using the calcium phosphate transfection kit according to manufacturer's instructions (5 Prime-3 Prime, Inc., Boulder, Colo.). Comparison of cellular transfection efficiency was assessed with the various constructs. For this procedure, confluent plates obtained 48 hours after transfection were split and reseeded

into 12 separate 6-cm dishes, and 6 of them were made to grow in the presence of 400 $\mu\text{g}/\text{ml}$ G-418 (GIBCO BRL, Life Technologies, Inc., Grand Island, N.Y.) containing medium. The number of G-418 resistant foci was determined at 14 days after transfection and used to correct for any variability in transfection efficiency.

[0079] Transformation Assay

[0080] The NIH-3T3 cells were transfected with the various constructs and allowed to reach confluence after 48 hours as described above. Each 6 cm dish was split and seeded into 12 different 6 cm dishes. While 6 of them were made to grow in the presence of G-418 to detect transfection efficiency, the other six were grown in complete medium without G-418 and with a medium change every 4th day. The number of transformed foci were counted in these plates without G-418 and expressed as transformed foci per μg transfected DNA.

[0081] Anchorage-Independent Cell Growth Assay

[0082] A limiting dilution technique (0.15 cell/well of a flat bottom 96-well-plate) was performed on transfectants grown in G-418 in order to isolate cell clones with different levels of HAAH activity as measured by Western blot analysis and enzymatic assay of hydroxylase activity. Cloned cell lines (1.0×10^4 cells) were suspended in complete medium containing 0.4% low-melting agarose (SeaPlaque GTG Agarose; FMC Bioproducts, Rockland, Me.) and laid over a bottom agar mixture consisting of complete medium with 0.53% low-melting agarose. Each clone was assayed in triplicate. The clones were seeded under these conditions and 10 days later the size (positive growth >0.1 mm in diameter) and number of foci were determined.

[0083] Tumorigenicity in Nude Mice

[0084] The same clones as assessed in the anchorage independent growth assay were injected into nude mice and observed for tumor formation. Tumorigenicity was evaluated using 10 animals in each of 4 groups (Charles River Labs., Wilmington, Mass.). Group 1 received 1×10^7 cells stably transfected with mock DNA, Group 2-4 received 1×10^7 cells of clones stably transfected with pNH376 and expressing various levels of murine HAAH protein. Nude mice were kept under pathogen-free conditions in a standard animal facility. Thirty days after tumor cell inoculation, the animals were sacrificed using isoflurane (Aerrane, Anaquest, N.J.) containing chambers and the tumors were carefully removed and weight determined.

[0085] Animal Model of Bile Duct Proliferation

[0086] Following ligation of the common bile duct, intrahepatic bile duct proliferation was evident at 48 hours. Tissue samples obtained 8 and 16 days following common bile duct ligation revealed extensive bile duct proliferation as shown in Table 3.

TABLE 3

Bile duct proliferation and HAAH expression at different intervals after common bile duct ligation			
Group	Surgical Procedure	Microscopy*	Immunohistochemistry
1	no surgery	normal	negative
2	sham surgery	normal	negative
3	6 hours post ligation	normal	negative
4	12 hours post ligation	normal	negative

TABLE 3-continued

Bile duct proliferation and HAAH expression at different intervals after common bile duct ligation			
Group	Surgical Procedure	Microscopy*	Immunohistochemistry
5	24 hours post ligation	normal	negative
6	48 hours post ligation	minimal bile duct prolif.	negative
7	4 days post ligation	moderate bile duct prolif.	negative
8	8 days post ligation	extensive bile duct prolif.	negative
9	16 days post ligation	extensive bile duct prolif.	negative

*Investigation was performed under light microscopy following a hematoxylin and eosin staining.

[0087] Immunohistochemical staining failed to detect presence of HAAH in proliferating bile ducts at any time. Analysis of HAAH expression in bile ducts derived from sham surgical controls was also negative, while all samples exhibited positive immunoreactivity with control antibodies to glyceraldehyde 3-phosphate dehydrogenase. Thus, bile duct proliferation was not associated with increased HAAH expression in this standard animal model system.

[0088] HAAH Expression in PSC

[0089] The liver biopsy specimens from patients with PSC exhibited bile duct proliferation accompanied by periductal fibrosis and a mononuclear inflammatory cell infiltrate without evidence of dysplasia. Adjacent sections immunostained with the an HAAH-specific monoclonal antibody had no detectable HAAH immunoreactivity in proliferating bile ducts. In contrast, sections of cholangiocarcinoma that were immunostained simultaneously using the same antibody and detection reagents manifested intense levels of HAAH immunoreactivity in nearly all tumor cells, whereas adjacent sections of the cholangiocarcinomas exhibited a negative immunostaining reaction with monoclonal antibody to Dengue virus. These findings indicate that HAAH expression was associated with malignant transformation rather than non-cancerous cellular proliferation of intrahepatic bile ducts.

[0090] HAAH Associated Transformation of NIH-3T3 Cells

[0091] The transforming capability of the murine and human AAH genes, as well as the murine AAH mutant construct without enzymatic activity were compared to mock DNA (negative control) and v-Src transfected NIH-3T3 cells (positive control). The transforming capability of murine AAH was found to be 2-3 times that of vector DNA control as shown in FIG. 1. The transforming capacity of the human gene was greater than that observed with the murine AAH (32 ± 1.5 versus 13 ± 2.6 transformed foci, respectively). The murine and human AAH transfected cells formed large foci, resembling those of v-Src transfected fibroblasts, compared to the occasional much smaller foci observed in cells transfected with vector DNA that displayed the contact inhibition of fibroblast cell lines. Parallel experiments performed using the mutant pNH376-H660 construct without enzymatic activity revealed no transforming activity. This finding indicates that the enzymatic activity of HAAH is required for the transforming activity exhibited by the HAAH gene.

[0092] Anchorage-Independent Cell Growth Assay

[0093] After transient transfection with the murine AAH construct, several different transformed foci were isolated for dilutional cloning experiments to establish stable transfected cell clones with different levels of HAAH gene expression. Nine different cloned cell lines were selected for further study. The expression level of the HAAH protein was determined by Western blot analysis. Clones 7 and 18 had a modest increase in HAAH protein expression, yet formed large colonies in soft agar (FIG. 2). Protein loading was equivalent in all lanes as shown by immunoblotting of the same membranes with an anti-Erk-1 monoclonal antibody. The increased protein expression was associated with increased enzymatic activity as shown in FIG. 3. The capability of these clones to exhibit anchorage independent cell growth in soft agar is presented in FIG. 3. All 3 clones with increased HAAH gene expression demonstrated anchorage independent cell growth compared to the mock DNA transfected clone.

[0094] Tumor Formation in Nude Mice

[0095] The 3 clones with increased HAAH gene expression were evaluated for the ability to form tumors in nude mice. Tumor size in the mouse given clone 18 was compared to a mock DNA transfected clone. Clones 7, 16 and 18 were highly transformed in this assay and produced large tumors with a mean weight of 2.5, 0.9 and 1.5 grams, respectively (FIG. 4). These data indicate that overexpression of HAAH contributes to induction and maintenance of the malignant phenotype in vivo.

[0096] High Level HAAH Expression is Indicative of Malignancy

[0097] In order to determine if HAAH expression was associated with malignancy rather than increased cell turnover, two models of bile duct proliferation were studied. In the animal model, ligation of the common bile duct induced extensive intrahepatic bile duct proliferation, yet there was no evidence of HAAH gene expression under these experimental conditions as shown in Table 3. Similarly, HAAH gene expression was assessed in a human disease model associated with bile duct proliferation since PSC is an autoimmune liver disease associated with destruction as well as proliferation of the intra and extrahepatic bile ducts. PSC is premalignant disease, and a significant proportion of affected individuals will eventually develop cholangiocarcinoma. However, no evidence for increased HAAH gene expression in the presence of extensive bile duct proliferation.

[0098] Having established that HAAH protein levels were elevated in cholangiocarcinoma and not in normal or proliferating bile ducts, the role of HAAH in the generation of a malignant phenotype was studied. The HAAH gene was transfected into NIH-3T3 cells and cellular changes, e.g., increased formation of transformed foci, colony growth in soft agar and tumor formation in nude mice associated with malignant transformation, were evaluated. The full-length murine and human AAH genes were cloned into expression constructs and transiently transfected into NIH-3T3 cells. An increased number of transformed foci was detected in cells transfected both with the murine and human AAH genes as compared to mock DNA transfected controls. The increased number of transformed foci, after controlling for transfection efficiency, was not as high compared to v-Src gene transfected cells used as a positive control. The enzymatic activity of the HAAH gene was required for a malignant phenotype because a mutant construct which abolished the catalytic site had no transforming properties. Several stable transfectants

and cloned NIH-3T3 cell lines with a modest increase in HAAH protein levels and enzymatic activity were established. Such cell lines were placed in soft agar to examine anchorage independent cell growth as another property of the malignant phenotype. All cell lines grew in soft agar compared to mock DNA transfected control, and there was a positive correlation between the cellular level of HAAH gene expression and the number and size of colonies formed. Three of these cloned cell lines formed tumors in nude mice. All three cell lines with increased HAAH expression were oncogenic as shown by the development of large tumors as another well-known characteristic of the transformed phenotype.

[0099] To determine whether cellular changes induced by overexpression of HAAH were related to the enzymatic function, a site-directed mutation was introduced into the gene that changed the ferrous iron binding site from histidine to lysine at 660th position of mouse HAAH thereby abolishing hydroxylase activity of the murine HAAH. A corresponding mutation in HAAH is used as a dominant negative mutant to inhibit HAAH hydroxylase activity. The pNH376-H660 construct had no transformation activity indicating cellular changes of the malignant phenotype induced by overexpression depends on the enzymatic activity of the protein.

[0100] Notch receptors and their ligands have several EGF-like domains in the N-terminal region that contain the putative consensus sequence for beta-hydroxylation. Notch ligands are important elements of the Notch signal transduction pathway and interaction of Notch with its ligands occurs by means of EGF-like domains of both molecules. Point mutations affecting aspartic acid or asparagine residues in EGF-like domains that are the targets for beta-hydroxylation by HAAH reduce calcium binding and protein-protein interactions involved in the activation of downstream signal transduction pathways. Overexpression of HAAH and Notch protein hydroxylation by HAAH contributes to malignancy. Tumor growth is inhibited by decreasing Notch protein hydroxylation by HAAH

[0101] The data presented herein is evidence that high-level HAAH expression is linked to malignant transformation. An increase in expression of the HAAH cDNA in NIH-3T3 cells induced a transformed phenotype manifested by increased numbers of transformed foci, anchorage-independent growth, and tumorigenesis in nude mice. In addition, intact HAAH-enzyme was found to be required for HAAH-associated transformation. Accordingly, inhibition of as little as 20% of endogenous HAAH enzymatic activity or expression confers a therapeutic benefit. For example, clinical benefit is achieved by 50%-70% inhibition of HAAH expression or activity after administration of an HAAH inhibitory compound compared to the level associated with untreated cancer cell or a normal noncancerous cell.

[0102] HAAH is regulated at the level of transcription. Only modest increases in HAAH expression and enzyme activity were required for cellular transformation. These results indicate that increased HAAH gene expression and enzyme activity contribute to the generation or maintenance of the transformed phenotype and that decreasing transcription of the HAAH gene or decreasing enzymatic activity of the HAAH gene product leads to a decrease in malignancy. Accordingly, HAAH transcription is inhibited by administering compounds which decrease binding of Fos and/or Jun (elements which regulate HAAH transcription) to HAAH promoter sequences.

[0103] Since HAAH is up-regulated with malignant transformation of bile duct epithelium, and HAAH immunoreactivity is detectable on tumor cell surface membranes, HAAH is also a molecule to which to target a cytotoxic agent, e.g., by linking the cytotoxic agent to a compound that binds to HAAH expressed on the surface of a tumor cell. Assay of HAAH protein levels in either biological fluids such as bile, or cells obtained by fine needle aspiration is a diagnostic marker of human cholangiocarcinoma.

Example 2

Expression of AAH and Growth and Invasiveness of Malignant CNS Neoplasms

[0104] AAH is abundantly expressed in carcinomas and trophoblastic cells, but not in most normal cells, including those of CNS origin. High levels of AAH expression were observed in 15 of 16 glioblastomas, 8 of 9 anaplastic oligodendrogliomas, and 12 of 12 primitive neuroectodermal tumors (PNETs). High levels of AAH immunoreactivity were primarily localized at the infiltrating edges rather than in the central portions of tumors. Double-label immunohistochemical staining demonstrated a reciprocal relationship between AAH and tenascin, a substrate for AAH enzyme activity. PNET2 neuronal cell lines treated with phorbol ester myristate or retinoic acid to stimulate neuritic extension and invasive growth exhibited high levels of AAH expression, whereas H₂O₂-induced neurite retraction resulted in down-regulation of AAH. PNET2 neuronal cells that stably over-expressed the human AAH cDNA had increased levels of PCNA and Bcl-2, and reduced levels of p21/Waf1 and p16, suggesting that AAH overexpression results in enhanced pathological cell proliferation, cell cycle progression, and resistance to apoptosis. In addition, the reduced levels of p16 observed in AAH-transfectants indicate that AAH over-expression confers enhanced invasive growth of neoplastic cells since deletion or down-regulation of the p16 gene correlates with more aggressive and invasive in vivo growth of glioblastomas. Increased AAH immunoreactivity was detected at the infiltrating margins of primary malignant CNS neoplasms, further indicating a role of HAAH in tumor invasiveness.

[0105] The following materials and methods were used to generate the data described below.

[0106] Analysis of AAH Immunoreactivity in Primary Human Malignant CNS Neoplasms:

[0107] AAH immunoreactivity was examined in surgical resection specimens of glioblastoma (N=16), anaplastic oligodendroglioma (N=9), and primitive neuroectodermal tumor (PNET; supratentorial neuroblastomas (N=3) and medulloblastomas (N=9)). The histopathological sections were reviewed to confirm the diagnoses using standard criteria. Paraffin sections from blocks that contained representative samples of viable solid tumor, or tumor with adjacent intact tissue were studied. Sections from normal adult post-mortem brains (N=4) were included as negative controls. AAH immunoreactivity was detected using qn HAAH-specific monoclonal antibody. Immunoreactivity was revealed by the avidin-biotin horseradish peroxidase complex method (Vector ABC Elite Kit; Vector Laboratories, Burlingame, Calif.) using 3-3' diaminobenzidine (DAB) as the chromogen (24) and hematoxylin as a counterstain.

[0108] Tenascin and laminin are likely substrates for AAH due to the presence of EGF-like repeats within the molecules. Double-immunostaining studies were performed to co-local-

ize AAH with tenascin or laminin. The AAH immunoreactivity was detected by the ABC method with DAB as the chromogen, and tenascin or laminin immunoreactivity was detected by the avidin-biotin alkaline phosphatase complex method (Vector Laboratories, Burlingame, Calif.) with BCIP/NBT as the substrate. As positive and negative controls, adjacent sections were immunostained with monoclonal antibody to glial fibrillary acidic protein (GFAP) and Hepatitis B surface antigen. All specimens were batch immunostained using the same antibody dilutions and immunodetection reagents.

[0109] Cell Lines and Culture Conditions

[0110] Studies were conducted to determine whether AAH expression was modulated with neurite (filopodia) extension (sprouting) as occurs with invasive growth of malignant neoplasms. Human PNET2 CNS-derived and SH-Sy5y neuroblastoma cells were cultured and stimulated for 0, 1, 2, 3, 5, or 7 days with 100 nM phorbol 12-ester 13-acetate or 10 μ M retinoic acid to induce sprouting. In addition, to examine the effects of neurite retraction on AAH expression, subconfluent cultures were treated for 24 hours with low concentrations (10-40 μ M) of H₂O₂. For both studies, AAH expression was evaluated by Western blot analysis using the an HAAH-specific antibody.

[0111] Generation of PNET2 AAH-Transfected Clones

[0112] The full-length human AAH cDNA (SEQ ID NO:3) was ligated into the pcDNA3.1 mammalian expression vector in which gene expression was under the control of a CMV promoter (Invitrogen Corp., San Diego, Calif.). PNET2 cells were transfected with either pHAAH or pcDNA3 (negative control) using Cellfectin reagent (Gibco BRL, Grand Island, N.Y.). Neomycin-resistant clones were selected for study if the constitutive levels of AAH protein expression were increased by at least two-fold relative to control (pcDNA3) as detected by Western blot analysis. To determine how AAH overexpression altered the expression of genes that modulate the transformed phenotype, the levels of proliferating cell nuclear antigen (PCNA), p53, p21/Waf1, Bcl-2, and p16 were measured in cell lysates prepared from subconfluent cultures of AAH (N=5) and pcDNA3 (N=5) stably transfected clones. PCNA was used as marker of cell proliferation. p53, p21/Waf1, and Bcl-2 levels were examined to determine whether cells that over-expressed AAH were more prone to cell cycle progression and more resistant to apoptosis. The levels of p16 were assessed to determine whether AAH over-expression has a role in tumor invasiveness.

[0113] Western Blot Analysis

[0114] Cells grown in 10 cm² dishes were lysed and homogenized in a standard radioimmunoprecipitation assay RIPA buffer containing protease and phosphatase inhibitors. The supernatants collected after centrifuging the samples at 12,000 \times g for 10 minutes to remove insoluble debris were used for Western blot analysis. Protein concentration was measured using the BCA assay (Pierce Chemical Co, Rockford, Ill.). Samples containing 60 μ g of protein were electrophoresed in sodium dodecyl sulfate polyacrylamide gels (SDS-PAGE) and subjected to Western blot analysis. Replicate blots were probed with the individual antibodies. Immunoreactivity was detected with horseradish peroxidase conjugated IgG (Pierce Chemical Co, Rockford, Ill.) and enhanced chemiluminescence reagents. To quantify the levels of protein expression, non-saturated autoradiographs were subjected to volume densitometry using NIH Image software, version 1.6. Statistical comparisons between pHAAH and pcDNA3 transfected cells were made using Student T tests.

[0115] Antibodies

[0116] HAAH-specific monoclonal antibody generated against the FOCUS hepatocellular carcinoma cells were used to detect AAH immunoreactivity. Monoclonal antibodies to tenascin, and glial fibrillary acidic protein, and rabbit polyclonal antibody to laminin were purchased from Sigma Co (St. Louis, Mo.). Rabbit polyclonal antibody to human p16 was purchased from Santa Cruz Biotechnology Inc. (Santa Cruz, Calif.). The 5C3 negative control monoclonal antibody to Hepatitis B surface antigen was generated using recombinant protein and used as a negative control.

[0117] AAH Immunoreactivity in Primary Malignant Brains Tumors

[0118] AAH immunoreactivity was detected in 15 of 16 glioblastomas, 8 of 9 anaplastic oligodendrogliomas, and all PNETs. AAH immunoreactivity was localized in the cytoplasm, nucleus, and cell processes. The tissue distribution of AAH immunoreactivity was notable for the intense labeling localized at the interfaces between tumor and intact brain, and the conspicuously lower levels of immunoreactivity within the central portions of the tumors. High levels of AAH immunoreactivity were also observed in neoplastic cells distributed in the subpial zones, leptomeninges, Virchow-Robin perivascular spaces, and in individual or small clusters of neoplastic cells that infiltrated the parenchyma. In contrast, AAH immunoreactivity was not detectable in normal brain. The distribution of AAH immunoreactivity appeared not to be strictly correlated with DNA synthesis since the density of nuclei in mitosis (1-5%) was similar in the central and peripheral portions of the tumors.

[0119] Relationship Between AAH and Tenascin Immunoreactivity in Glioblastomas

[0120] Tenascin is an extracellular matrix-associated antigen expressed in malignant gliomas. Tenascin contains EGF-like domains within the molecule, a substrate for HAAH hydroxylation. To localize AAH in relation to tenascin immunoreactivity in malignant brain tumors, double-label immunohistochemical staining was performed in which AAH was detected using a brown chromogen (DAB), and tenascin, a blue chromogen (BCIP/NBT). Adjacent sections were similarly double-labeled to co-localize AAH with laminin, another EGF domain containing extracellular matrix molecule expressed in the CNS. Intense levels of tenascin immunoreactivity were observed in perivascular connective tissue and in association with glomeruloid proliferation of endothelial cells. The double-labeling studies demonstrated a reciprocal relationship between AAH and tenascin immunoreactivity such that high levels of AAH were associated with low or undetectable tenascin, and low levels of AAH were associated with abundant tenascin immunoreactivity. Although laminins are also likely substrates for AAH enzyme activity due to the EGF repeats within the molecules, double labeling studies revealed only low levels of laminin immunoreactivity throughout the tumors and at interfaces between tumor and intact tissue.

[0121] Analysis of AAH Expression in Neuronal Cell Lines Treated with PMA or RA

[0122] Neuritic sprouting/filopodia extension marks invasive growth of neoplastic neuronal cells. PMA activates protein kinase C signal transduction pathways that are involved in neuritic sprouting. Retinoic acid binds to its own receptor and the ligand-receptor complex translocates to the nucleus where it binds to specific consensus sequences present in the promoter/enhancer regions of target genes involved in neu-

ritic growth. Both PNET2 and SH-Sy5y cells can be induced to sprout by treatment with PMA (60-120 nM) or retinoic acid (5-10 μ M). FIGS. 5A-D depict data from representative Western blot autoradiographs; the bar graphs correspond to the mean \pm S.D. of results obtained from three experiments. Western blot analysis with the FB50 antibody detected doublet bands corresponding to protein with an molecular mass of approximately 85 kDa. Untreated PNET2 cells had relatively low levels of AAH immunoreactivity (FIG. 5A), whereas untreated SH-Sy5y cells had readily detected AAH expression (FIG. 5B). Untreated PNET2 cells exhibited polygonal morphology with coarse, short radial cell processes, whereas SH-Sy5y cells were slightly elongated and spontaneously extend fine tapered processes. Both cell lines manifested time-dependent increases in the levels of AAH immunoreactivity following either RA (FIGS. 5A and 5B) or PMA (FIG. 5C) stimulation and neurite extension. In PNET2 cells, the levels of AAH protein increased by at least two-fold 24 hours after exposure to RA or PMA, and high levels of AAH were sustained throughout the 7 days of study. In SH-Sy5y cells, the RA- or PMA-stimulated increases in AAH expression occurred more gradually and were highest after 7 days of treatment (FIG. 5B).

[0123] To examine the effect of AAH expression on neurite retraction, PNET2 and SH-Sy5y cells were treated with low concentrations (8-40 μ M) of H₂O₂. After 24 hours exposure to up to 40 μ M H₂O₂, although most cells remained viable (Trypan blue dye exclusion), they exhibited neurite retraction and rounding. Western blot analysis using the FB50 antibody demonstrated H₂O₂ dose-dependent reductions in the levels of AAH protein (FIG. 5D).

[0124] Effects of AAH Over-Expression in PNET2 Cells

[0125] To directly assess the role of AAH overexpression in relation to the malignant phenotype, PNET2 cells were stably transfected with the human full-length cDNA with gene expression under control of a CMV promoter (pHAAH). Neomycin-resistant clones that had at least two-fold higher levels of AAH immunoreactivity relative to neomycin-resistant pcDNA3 (mock) clones were studied. Since aggressive behavior of malignant neoplasms is associated with increased DNA synthesis, cell cycle progression, resistance to apoptosis, and invasive growth, the changes in phenotype associated with constitutive over-expression of AAH were characterized in relation to PCNA, p21/Waf1, p53, Bcl-2, and p16. PCNA was used as an index of DNA synthesis and cell proliferation. p21/Waf1 is a cell cycle inhibitor. Expression of the p53 tumor-suppressor gene increases prior to apoptosis, whereas bcl-2 inhibits apoptosis and enhances survival of neuronal cells. p16 is an oncosuppressor gene that is often either down-regulated or mutated in infiltrating malignant neoplasms.

[0126] Five pHAAH and 5 pcDNA3 clones were studied. Increased levels of AAH expression in the pHAAH transfected clones was confirmed by Western (FIG. 6) and Northern blot analyses. Western blot analysis using cell lysates from cultures that were 70 to 80 percent confluent demonstrated that constitutively increased levels of AAH expression (approximately 85 kDa; $P<0.05$) in pHAAH-transfected cells were associated with significantly increased levels of PCNA (approximately 35 kDa; $P<0.01$) and Bcl-2 (approximately 25 kDa; $P<0.05$), and reduced levels of p21/Waf1 (approximately 21 kDa; $P<0.001$) and p16 (approximately 16 kDa; $P<0.001$) (FIG. 6). However, the pHAAH stable transfectants also exhibited higher levels of wild-type p53 (approximately 53-55 kDa). Although AAH expression (85 kDa protein) in

the stable transfectants was increased by only 75 to 100 percent, the levels of p16 and p21/Waf1 were sharply reduced, and PCNA increased by nearly two-fold (FIG. 6).

[0127] Increased AAH Expression is Indicative of Growth and Invasiveness of Malignant CNS Neoplasms

[0128] The data described herein demonstrates that AAH overexpression is a diagnostic tool by which to identify primary malignant CNS neoplasms of both neuronal and glial cell origin. Immunohistochemical staining studies demonstrated that AAH overexpression was detectable mainly at the interfaces between solid tumor and normal tissue, and in infiltrating neoplastic cells distributed in the subpial zones, leptomeninges, perivascular spaces, and parenchyma. In vitro experiments demonstrated that AAH gene expression was modulated with neurite (filopodium) extension and invasiveness and down-regulated with neurite retraction. In addition, PNET2 cells stably transfected with the AAH cDNA exhibited increased PCNA and bcl-2, and reduced Waf1/p21 and p16 expression. Therefore, AAH overexpression contributes to the transformed phenotype of CNS cells by modulating the expression of other genes that promote cellular proliferation and cell cycle progression, inhibit apoptosis, or enhance tumor cell invasiveness.

[0129] The data demonstrated readily detectable AAH mRNA transcripts (4.3 kDa and 2.6 kDa) and proteins (85 kDa and 50-56 kDa) in PNET2 and SH-Sy5y cells, but not in normal brain. Correspondingly, high levels of AAH immunoreactivity were observed in 35 of the 37 in malignant primary CNS-derived neoplasms studied, whereas the 4 normal control brains had no detectable AAH immunoreactivity. The presence of high-level AAH immunoreactivity at the infiltrating margins and generally not in the central portions of the tumors indicates that AAH overexpression is involved in the invasive growth of CNS neoplasms. Administration of compounds which decrease AAH expression or enzymatic activity inhibits proliferation of CNS tumors which overexpress AAH, as well as metastases of CNS tumors to other tissue types.

[0130] The AAH enzyme hydroxylates EGF domains of a number of proteins. Tenascin, an extracellular matrix molecule that is abundantly expressed in malignant gliomas, contains EGF-like domains. Since tenascin promotes tumor cell invasion, its abundant expression in glioblastomas represents an autocrine mechanism of enhanced tumor cell growth vis-à-vis the frequent overexpression of EGF or EGF-like receptors in malignant glial cell neoplasms. Analysis of the functional domains of tenascins indicated that the mitogenic effects of this family of molecules are largely mediated by the fibronectin domains, and that the EGF-like domains inhibit growth, cell process elongation, and matrix invasion. Therefore, hydroxylation of the EGF-like domains by AAH represents an important regulatory factor in tumor cell invasiveness.

[0131] Double-label immunohistochemical staining studies demonstrated a reciprocal relationship between AAH and tenascin immunoreactivity such that high levels AAH immunoreactivity present at the margins of tumors were associated with low levels of tenascin, and low levels of AAH were often associated with high levels of tenascin. These observations indicated that AAH hydroxylation of EGF-like domains of tenascin alters the immunoreactivity of tenascin protein, and in so doing, facilitates the invasive growth of malignant CNS neoplasms into adjacent normal tissue and perivascular spaces.

[0132] AAH immunoreactivity was examined in PNET2 and SH-Sy5y neuronal cells induced to undergo neurite extension with PMA or retinoic acid, or neurite retraction by exposure to low doses of H₂O₂. AAH expression was sharply increased by PMA- or retinoic acid-induced neurite (filopodium) extension, and inhibited by H₂O₂-induced neurite retraction and cell rounding. Neurite or filopodium extension and attachment to extracellular matrix are required for tumor cell invasion in the CNS. The EGF-like domains of tenascin inhibit neuritic and glial cell growth into the matrix during development.

[0133] To directly examine the role of AAH overexpression in relation to the transformed phenotype, genes modulated with DNA synthesis, cell cycle progression, apoptosis, and tumor invasiveness were examined in neuronal cell clones that stably over-expressed the human AAH cDNA. The findings of increased PCNA and reduced Waf1/p21 immunoreactivity indicated that AAH overexpression enhances cellular proliferation and cell cycle progression. In addition, the finding of increased Bcl-2 expression indicated that AAH overexpression contributes to the transformed phenotype by increasing cellular resistance to apoptosis. The apparently contradictory finding of higher levels of p53 in the cells that overexpressed AAH is explained by the observation that high levels of wildtype p53 in immature neuronal cells were associated with neuritic growth (invasiveness) rather than apoptosis. Levels of p16 were reduced (compared to normal cells) or virtually undetectable in cells that constitutively overexpressed AAH; a deletion mutation of the p16 gene has been correlated with invasive growth and more rapid progression of malignant neoplasms, including those of CNS origin. These data indicate that p16 expression is modulated by AAH.

Example 3

Increased HAAH Production and IRS-Mediated Signal Transduction

[0134] IRS-1 mediated signal transduction pathway is activated in 95% of human HCC tumors compared to the adjacent uninvolved liver tissue. HAAH is a downstream effector gene involved in this signal transduction pathway. HAAH gene upregulation is closely associated with overexpression of IRS-1 in HCC tumors as revealed by immunohistochemical staining and Western blot analysis. A high level of HAAH protein is expressed in HCC and cholangiocarcinoma compared to normal hepatocytes and bile ducts. Both of these tumors also exhibit high level expression of IRS-1 by immunohistochemical staining. FOCUS HCC cell clones stably transfected with a C-terminal truncated dominant negative mutant of IRS-1, which blocks insulin and IGF-1 stimulated signal transduction, was associated with a striking reduction in HAAH gene expression in liver. In contrast, transgenic mice overexpressing IRS-1 demonstrate an increase in HAAH gene expression by Western blot analysis. Insulin stimulation of FOCUS HCC cells (20 and 40 U) in serum free medium and after 16 hr of serum starvation demonstrated upregulation of HAAH gene expression. These data indicate that HAAH gene expression is a downstream effector of the IRS-1 signal transduction pathway.

Example 4

Effects of HAAH Expression Levels on the Characteristics of the Malignant Phenotype

[0135] Overexpression of IRS-1 in NIH 3T3 cells induces transformation. The full-length murine HAAH construct was

cloned into the pcDNA3 eukaryotic expression vector. A second murine construct encoded HAAH with abolished catalytic activity due to a site directed mutation. The full-length human HAAH cDNA was cloned into the pcDNA3 expression vector as well as a plasmid that encodes v-src which was used as a positive control for transformation activity. Standard methods were used for transfection of NIH 3T3 cells, control for transfection efficiency, assays of HAAH enzymatic activity, transformation by analysis of foci formation, anchorage-independent cell growth assays and analysis of tumorigenicity in nude mice. The data indicate that HAAH overexpression is associated with generation of a malignant phenotype.

TABLE 4

Overexpression of enzymatically active HAAH indicates malignancy			
cDNA	# of foci ± S.D. ^b	NIH 3T3 clone	# of colonies ^c
pcDNA3 (mock)	6.0 ± 3.3	pcDNA (mock)	0.4 ± 0.5
murine HAAH	14.0 ± 2.9	clone 18 ^d	6.2 ± 2.9
mutant murine HAAH ^a	1.6 ± 1.0	clone 16 ^e	4.7 ± 6.5
human HAAH	32.0 ± 5.4		
v-src	98.0 ± 7.1		

^aenzymatically inactive HAAH

^bP < 0.01 compared to mock and mutant murine HAAH

^cP < 0.001 compared to mock

^dClone 18 is a stable cloned NIH 3T3 cell line that overexpresses human HAAH by approximately two fold.

^eClone 16 is a stable cloned NIH 3T3 cell line that overexpresses human HAAH by about 50%.

[0136] These data indicate that overexpression of HAAH is associated with formation of transformed foci. Enzymatic activity is required for cellular transformation to occur. Cloned NIH 3T3 cell lines with increased human HAAH gene expression grew as solid tumors in nude mice. HAAH is a downstream effector gene of the IRS-1 signal transduction pathway.

Example 5

Inhibition of HAAH Gene Expression

[0137] The FOCUS HCC cell line from which the human HAAH gene was initially cloned has a level of HAAH expression that is approximately 3-4 fold higher than that found in normal liver. To make an HAAH antisense construct, the full length human HAAH cDNA was inserted in the opposite orientation into a retroviral vector containing a G418 resistant gene, and antisense RNA was produced in the cells. Shorter HAAH antisense nucleic acids, e.g., those corresponding to exon 1 of the HAAH gene are also used to inhibit HAAH expression.

[0138] FOCUS cells were infected with this vector and the level of HAAH was determined by Western blot analysis. A reduction in HAAH gene expression was observed. Growth rate and morphologic appearance of cells infected with a retrovirus containing a nonrelevant Green Fluorescent Protein (GFP) also inserted in the opposite orientation as a control (FIG. 8). Cells (harboring the HAAH antisense construct) exhibited a substantial change in morphology characterized by an increase in the cytoplasm to nuclear ratio as well as assuming cell shape changes that were reminiscent of normal adult hepatocytes in culture. Cells with reduced HAAH levels

grew at a substantially slower rate than retroviral infected cells expressing antisense (GFP) (control) as shown in FIG. 8. A reduction in HAAH gene expression was associated with a more differentiated noncancerous “hepatocyte like” phenotype. Expression of HAAH antisense sequences are used to inhibit tumor growth rate. Reduction of HAAH cellular levels results in a phenotype characterized by reduced formation of transformed foci, low level or absent anchorage independent growth in soft agar, morphologic features of differentiated hepatocytes as determined by light and phase contrast microscopy, and no tumor formation (as tested by inoculating the cells into nude mice).

Example 6

Human IRS-1 Mutants

[0139] Insulin/IGF-1 stimulated expression of HAAH in HCC cell lines. Dominant-negative IRS-1 cDNAs mutated in the plextrin and phosphotyrosine (PTB) domains, and Grb2, Syp and PI3K binding motifs located in the C-terminus of the

molecule were constructed. Human IRS-1 mutant constructs were generated to evaluate how HAAH gene expression is upregulated by activation of the IRS-1 growth factor signal transduction cascade. Specific mutations in the C terminus of the hIRS-1 molecule abolished the various domains which bind to SH2-effector proteins such as Grb2, Syp and PI3K. The human IRS-1 protein contains the same Grb2 and Syp binding motifs of 897YVNI (underlined in Table 5, below and 1180YIDL (underlined in Table 5, below), respectively, as the rat IRS-1 protein. Mutants of hIRS-1 were constructed by substitution of a TAT codon (tyrosine) with a TTT codon (phenylalanine), in these motifs by use of oligonucleotide-directed mutagenesis using the following primers: (5'-GGGGGAATTTGTCAATA-3' (SEQ ID NO:8) and 5'-GAATTTGTTAATATTG-3' (SEQ ID NO:9), respectively). The cDNAs of hIRS-1 (wild-type) and mutants (tyrosine 897-to-phenylalanine and tyrosine 1180-to-phenylalanine) were subcloned into the pBK-CMV expression vector and designated as hIRS-1-wt, 897F, Δ-Grb2), 1180F, and ΔSyp.

TABLE 5

Human IRS-1 amino acid sequence		
MASPPESDGF	SDVRKVGYLK KPKSMHKRFF VLRAASEAGG PARLEYEYENE KWRHKSSAP	61
KRSIPLESCF	NINKRADSKN KHLVALYTRD EHFAIADSE AEQDSWYQAL LQLHNRAKGH	121
HDGAAALGAG	GGGGSCSGSS GLGEAGEDLS YGDVPPGPAP KEVWQVILKP KGLGQTKNLI	181
GIYRLCLTSK	TISFVKLNSE AAAVLQLMN IRRCGHSENF FFIEVGRSAV TGPGEFWMQV	241
DDSVVAQNMH	ETILEAMRAM SDEFPRRSKS QSSSNCSNPI SVPLRRHHLN NPPPSQVGLT	301
RRSRTESITA	TSPASMVGK PGFVRVRASS DGEGTMSRPA SVDGSPVSPS TNRTHAHRHR	361
GSARLHPPLN	HSRSIPMPAS RCSPSATSPV SLSSSSSTSGH GSTSDCLFPR RSSASVSGSP	421
SDGGFISSDE	YGSSPCDFRS SFRSVTPDSL GHTPPARGEE ELSNYICMGK KGPSTLTAPN	481
GHYILSRGGN	GHRCTPGTGL GTSPALAGDE AASAADLDNR FRKRTHSAGT SPTITHQKTP	541
SQSSVASIEE	YTEMPPAYPP GGGSGGRLPG HRHSFVPTTR SYPEEGLEMH PLERRGGHHR	601
PDSSTLHTDD	GYMPMSPGVA PVPGRKGSG DYMPMSPKSV SAPQIINPI RRHPQRVDPN	661
GYMMSPSGG	CSPDIGGGS SSSSSSNAV SGTSGYGLWT NGVGGHSHV LPHPKPPVES	721
SGGKLLPCTG	DYMNMSVPGD SNTSSPSDCY YGPEDPQHKP VLSYYSLPRS FKHTQRPGE	781
EEGARHQHLR	LSTSSGRLLY AATADDSSSS TSSDSLGGY CGARLEPSLP HPHHQVLQPH	841
LPRKVDTAQ	TNSRLARPTR LSLGDPKAST LPRAREQQQQ QPLLHPPEP KSPGEYVNI	901
FGSDQSGYLS	GPVAFHSSPS VRCPSQLQPA PREEETGTEE YMKMDLGPGR RAAWQESTGV	961
EMGRLGPAPP	GAASICRPTR AVPSRRGDYM TMQMSPCRQS YVDTSPAAPV SYADMRTGIA	1021
AEVSLPRAT	MAAASSSSAA SASPTGPQGA AELAHHSSLL GGPQGPQGM S AFTRVNLSPN	1081
RNQSAAKIRA	DPQGCRRRHS SETFSSTPSA TRVGNTVPFG AGAAVGGGGG SSSSESDVKR	1141
HSSASFENVW	LRPGELGGAP KEPAKLCGAA GGLENLNYI <u>DL</u> DLVKDFKQ CPQECTPEPQ	1201
PPTPPPHQPP LGSGESSSTR RSSEDL SAYS SISFQKQPED RQ (SEQ ID NO: 5; GENBANK Accession No. J0670; pleckstrin domain spans residues 11-113, inclusive; Phosphate-binding residues include 46, 465, 551, 612, 632, 662, 732, 941, 989, or 1012 of SEQ ID NO: 5)		

TABLE 6

Human IRS-1 cDNA		
cggcggcgcg gtcggagggg gccggcgcg agagccagac gccgccgctt gttttggttg	61	
gggctctcgg caactctcgg aggaggagga ggaggaggga ggaggggaga agtaactgca	121	
gcggcagcgc cctcccgagg aacaggcgctc ttccccgaac ccttcccaaa cctcccccat	181	
ccccctctgc ccttgctccc tcccctcctc cccagccgccc tggagcgagg ggcagggatg	241	
agttctgtccc tccggccgggt ccccgagctgc agtggctgcc cgggtatcggt tcgcatggaa	301	
aagccacttt ctccaccgccc cgagatgggg ccggatgggg ctgcagagga cgcgcccgcg	361	
ggcggcgcca gcagcagcag cagcagcagc agcaacagca acagccgcag cgcgcgggtc	421	
tctgcgactg agctggtatt tgggcggctg gtggcggtg ggagcgttgg ggggtgggag	481	
gaggcgaagg aggaggagga accccgtgca acgttgggac ttggcaaccc gcctccccct	541	
gcccgaaggat atttaatttg cctcgggaat cgctgcttcc agaggggaac tcaggaggga	601	
aggcgcgcgc gcgcgcgcgc tccctggaggg gcaccgcagg gacccccgac tgtcgcttcc	661	
ctgtgccgga ctccagccgg ggccagcaga gatgcatctt cgctccttcc tgggtggcggc	721	
ggcggctgag aggagacttg gctctcggag gatcggggct gccctcacc cggacgcact	781	
gcctccccgc cggcgtaag gcgccgaaaa ctccggctcgg gctctctcct gggctcagca	841	
gctgcgtcct ccttcagctg cccctccccg gcgcgggggg cggcggtgat ttcagagtcg	901	
gggtttctgc tgcctccagc cctgtttgca tgtgccgggc cgcggcgagg agcctccgccc	961	
ccccaccggt ttgtttttcg gagcctccct ctgctcagcg ttggtggtgg cggtggcagc	1021	
atggcgagcc ctccggagag cgatggcttc tcggacgtgc gcaagggtgg ctacctgcgc	1081	
aaacccaaga gcatgcacaa acgcttcttc gtactcgcgc cggccagcga ggctgggggc	1141	
ccggcgcgcc tcgagtaacta cgagaacgag aagaagtggc ggcacaagtc gagcgcccc	1201	
aaacgctoga tcccccttga gagctgcttc aacatcaaca agcgggctga ctccaagaac	1261	
aagcacctgg tggctctcta caccggggac gagcactttg ccacgcgggc ggacagcgag	1321	
gccgagcaag acagctggta ccaggctctc ctacagctgc acaaccgtgc taagggccac	1381	
cacgacggag ctgcggccct cggggcgagg ggtggtgggg gcagctgcag cggcagctcc	1441	
ggccttggtg aggtggggga ggaattgagc tacgggtgacg tgccccagg acccgcatc	1501	
aaagaggtct ggcaagtgat cctgaagccc aagggcctgg gtcagacaaa gaacctgatt	1561	
ggtatctacc gcctttgcct gaccagcaag accatcagct tcgtgaagct gaactcggag	1621	
gcagcggccg tgggtctgca gctgatgaac atcaggcgct gtggccactc ggaaaacttc	1681	
ttcttcatcg aggtggggccg ttctgccgtg acggggccccg gggagttctg gatgcagggtg	1741	
gatgactctg tgggtggcca gaacatgcac gagaccatcc tggaggccat gcggggccatg	1801	
agtgatgagt tccgccctcg cagcaagagc cagtctctgt ccaactgctc taaccccatc	1861	
agcgtccccc tgcgcgggga ccactctaac aatcccccg cagccaggt ggggctgacc	1921	
cgcgatacgc gactgagag catcacgcgc acctccccg ccagcatggt gggcggaag	1981	
ccaggctcct tccgtgtccg gcctccagt gacggcgaag gcaccatgct ccgcccagcc	2041	
tcggtggacg gcagccctgt gagtccagc accaacagaa cccacgccca ccggcatcgg	2101	
ggcagcgcgc ggtgcaccc cccgctcaac cacagccgct ccaccccat gccggcttcc	2161	
cgctgctcgc ctccggccac cagcccggtc agtctgtcgt ccagtagcac cagtggccat	2221	

TABLE 6-continued

Human IRS-1 cDNA					
ggctccacct	cggattgtct	cttcccacgg	cgatctagt	cttcggtgtc	tggttcccc 2281
agcgatggcg	gtttcatctc	ctcggatgag	tatggctcca	gtccctgcga	tttccggagt 2341
tccttccgca	gtgtcactcc	ggattccctg	ggccacaccc	caccagcccg	cggtagggag 2401
gagctaagca	actatatctg	catgggtggc	aaggggccct	ccacctgac	cgcctccaac 2461
ggtcactaca	ttttgtctcg	gggtggcaat	ggccaccgct	gcacccagg	aacaggcttg 2521
ggcacgagtc	cagccttggc	tggggatgaa	gcagccagt	ctgcagatct	ggataatcgg 2581
ttccgaaaga	gaactcactc	ggcaggcaca	tcccctacca	ttaccacca	gaagaccccg 2641
tcccagtcct	cagtggcttc	cattgaggag	tacacagaga	tgatgcctgc	ctaccacca 2701
ggaggtggca	gtggaggccg	actgccggga	cacaggcact	cgccttcgt	gcccacccgc 2761
tcctaccag	aggagggtct	ggaaatgcac	cccttgagc	gtcggggggg	gcaccaccgc 2821
ccagacagct	ccacctcca	cacggatgat	ggctacatgc	ccatgtcccc	aggggtggcc 2881
ccagtgccta	gtggccgaaa	gggcagtgga	gactatatgc	ccatgagccc	caagagcgta 2941
tctgccccac	agcagatcat	caatcccac	agacgccac	cccagagagt	ggaccccaat 3001
ggctacatga	tgatgtcccc	cagcgggtgc	tgtctcctg	acattggagg	tggccccagc 3061
agcagcagca	gcagcagcaa	gcgcgtccct	tccgggacca	gctatggaaa	gctgtggaca 3121
aacggggtag	ggggccacca	ctctcatgtc	ttgcctcacc	ccaaaccccc	agtggagagc 3181
agcgggtgta	agctcttacc	ttgcacaggt	gactacatga	acatgtcacc	agtgggggac 3241
tccaacacca	gcagccctc	cgactgtac	tacggccctg	aggaccccca	gcacaagcca 3301
gtctctctct	actactcatt	gccaaagatc	tttaagcaca	cccagcgcgc	cggggagccg 3361
gaggaggggtg	cccggcatca	gcacctccgc	ctttccacta	gctctggtcg	ccttctctat 3421
gctgcaacag	cagatgatcc	ttcctcttcc	accagcagcg	acagcctggg	tgggggatac 3481
tgcggggcta	ggctggagcc	cagccttcca	catccccacc	atcaggttct	gcagcccat 3541
ctgcctcgaa	aggtggacac	agctgtcag	accaatagcc	gcctggcccg	gcccacgagg 3601
ctgtccctgg	gggateccaa	ggccagcacc	ttacctcggg	cccagagcga	gcagcagcag 3661
cagcagccct	tgtgcacccc	tccagagccc	aagagcccg	gggaatatgt	caatattgaa 3721
tttgggagtg	atcagctctg	ctacttgtct	ggcccggtgg	ctttccacag	ctcaccttct 3781
gtcaggtgtc	catcccagct	ccagccagct	cccagagagg	aagagactgg	cactgaggag 3841
tacatgaaga	tggacctggg	gccggggccg	agggcagcct	ggcaggagag	cactggggtc 3901
gagatgggca	gactggggcc	tgcacctccc	ggggctgcta	gcatttgcag	gcctaccccg 3961
gcagtgccta	gcagccgggg	tgactacatg	accatgcaga	tgagttgtcc	ccgtcagagc 4021
tacgtggaca	cctcgccagc	tgcctctgta	agctatgtg	acatgcgaac	aggcattgct 4081
gcagaggagg	tgagcctgcc	cagggccacc	atggctgtg	cctcctcacc	ctcagcagcc 4141

TABLE 6-continued

Human IRS-1 cDNA	
tctgcttccc cgactggggc tcaaggggca gcagagctgg ctgcccactc gtcctctgctg	4201
ggggggccac aaggacctgg gggcatgagc gccttcaccc ggggtgaacct cagtcctaac	4261
cgcaaccaga gtgccaaagt gatccgtgca gaccacaag ggtgccggcg gaggcatagc	4321
tccgagactt tctcctcaac acccagtgcc acccggtggt gcaacacagt gccctttgga	4381
gcggggcgag cagtaggggg cggtggcggt agcagcagca gcagcgagga tgtgaaacgc	4441
cacagctctg ctctctttga gaatgtgtgg ctgaggcctg gggagcttgg gggagccccc	4501
aaggaggccag ccaaaactgtg tggggctgct gggggtttgg agaattgtct taactacata	4561
gacctggatt tggtaacaga cttcaaacag tgccttcagg agtgcacccc tgaaccgcag	4621
cctccccac cccaccccc tcatcaaccc ctgggcagcg gtgagagcag ctccacccgc	4681
cgctcaagtg aggatttaag cgcctatgcc agcatcagtt tccagaagca gccagaggac	4741
cgtcagtagc tcaactggac atcacagcag aatgaagacc taaatgacct cagcaaatcc	4801
tcttctaact catgggtacc cagactctaa atatttcatg attcacaact aggacctcat	4861
atcttctca tcagtagatg gtacgatgca tccatttcag tttgtttact ttatccaatc	4921
ctcaggattt cattgactga actgcacggt ctatattgtg ccaagcgaaa aaaaaaatg	4981
cactgtgaca ccagaataat gagtctgcat aaacttcac ttcaacctta aggacttagc	5041
tggccacagt gagctgatgt ccccaccacc gtgtcatgag agaattgggt tactctcaat	5101
gcattttcaa gatacatctt atctgctgct gaaactgtgt acgacaaagc atcattgtaa	5161
attatttcat acaaaactgt tcacgttggg tggagagagt attaaatatt taacataggt	5221
tttgatttat atgtgtaatt ttttaaatga aaatgtaact tttcttacag cacatctttt	5281
ttttggatgt gggatggagg tatacaatgt tctgttgtaa agagtggagc aaatgcttaa	5341
aacaaggctt aaaagagtag aatagggtat gatccttggt ttaagattgt aattcagaaa	5401
acataatata agaatacatg tgccatagat ggttctcaat tgtatagtta tatttgctga	5461
tactatctct tgtcatataa acctgatggt gagctgagtt ccttataaga attaacttta	5521
attttgattt ttttctgta agacaatagg ccatgttaat taaactgaag aaggatatat	5581
ttggctgggt gttttcaaat gtcagcttaa aattggtaat tgaatggaag caaaattata	5641
agaagaggaa attaaagtct tccattgcat gtattgtaaa cagaaggaga tgggtgatcc	5701
cttcaattca aaagctctct ttggaatgaa caatgtgggc gtttgtaaat tctggaaatg	5761
tctttctatt cataataaac tagatactgt tgatctttta aaaaaaaaaa aaaaaaaaaa	5821

(SEQ ID NO: 6; GENBANK Accession No. NM 005544)

[0140] The double mutation of tyrosine 897 and 1180 was constructed by replacement of 3'-sequences coding 897F by the same region of 1180F using restriction enzymes NheI and EcoRI, and this construct was called 897F1180F or ΔGrb2 ΔSyp. The expression plasmids were under control of a CMV promoter (hIRS-1-wt, ΔGrb2, ΔSyp, ΔGrb2, ΔSyp and pBK-CMV (mock) and linearized at the 3'-end of poly A signal sequences by MluI restriction enzymes followed by purification. A similar approach was used to change the tyrosine residue to phenylalanine at positions 613 and 942 to create the double PI3K mutant construct (ΔPI3K). The hIRS-1 mutants have a FLAG epitope (DYKDDDDK (SEQ ID NO:6)+stop

codon) added to the C-terminus by PCR. This strategy allows to distinguish the mutant protein from "wild type" hIRS-1 in stable transfected cell lines. The mutants are used to define the link between the IRS signal transduction pathway and activation of HAAH as a downstream effector gene and identify compounds to inhibit transduction along the pathway to inhibit growth of tumors characterized by HAAH overexpression. Antibodies or other compounds which bind to phosphorylation sites or inhibit phosphorylation at those sites are used to inhibit signal transduction and thus proliferation of HAA-overexpressing tumors.

[0141] Other embodiments are within the following claims.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 9

<210> SEQ ID NO 1
<211> LENGTH: 36
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Consensus
EGF-like domain
<220> FEATURE:
<221> NAME/KEY: VARIANT
<222> LOCATION: (2)..(8)
<223> OTHER INFORMATION: Wherein Xaa is any amino acid
<220> FEATURE:
<221> NAME/KEY: VARIANT
<222> LOCATION: (10)..(13)
<223> OTHER INFORMATION: Wherein Xaa is any amino acid.
<220> FEATURE:
<221> NAME/KEY: VARIANT
<222> LOCATION: (15)..(24)
<223> OTHER INFORMATION: Wherein Xaa is any amino acid.
<220> FEATURE:
<221> NAME/KEY: VARIANT
<222> LOCATION: (26)
<223> OTHER INFORMATION: Wherein Xaa is any amino acid.
<220> FEATURE:
<221> NAME/KEY: VARIANT
<222> LOCATION: (28)..(35)
<223> OTHER INFORMATION: Wherein Xaa is any amino acid.

<400> SEQUENCE: 1

Cys Xaa Xaa Xaa Xaa Xaa Xaa Xaa Cys Xaa Xaa Xaa Xaa Cys Xaa Xaa
1 5 10 15
Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Cys Xaa Cys Xaa Xaa Xaa Xaa Xaa
20 25 30
Xaa Xaa Xaa Cys
35

<210> SEQ ID NO 2
<211> LENGTH: 758
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 2

Met Ala Gln Arg Lys Asn Ala Lys Ser Ser Gly Asn Ser Ser Ser Ser
1 5 10 15
Gly Ser Gly Ser Gly Ser Thr Ser Ala Gly Ser Ser Ser Pro Gly Ala
20 25 30
Arg Arg Glu Thr Lys His Gly Gly His Lys Asn Gly Arg Lys Gly Gly
35 40 45
Leu Ser Gly Thr Ser Phe Phe Thr Trp Phe Met Val Ile Ala Leu Leu
50 55 60
Gly Val Trp Thr Ser Val Ala Val Val Trp Phe Asp Leu Val Asp Tyr
65 70 75 80
Glu Glu Val Leu Gly Lys Leu Gly Ile Tyr Asp Ala Asp Gly Asp Gly
85 90 95
Asp Phe Asp Val Asp Asp Ala Lys Val Leu Leu Gly Leu Lys Glu Arg
100 105 110
Ser Thr Ser Glu Pro Ala Val Pro Pro Glu Glu Ala Glu Pro His Thr
115 120 125

-continued

Glu	Pro	Glu	Glu	Gln	Val	Pro	Val	Glu	Ala	Glu	Pro	Gln	Asn	Ile	Glu
130						135					140				
Asp	Glu	Ala	Lys	Glu	Gln	Ile	Gln	Ser	Leu	Leu	His	Glu	Met	Val	His
145				150					155					160	
Ala	Glu	His	Val	Glu	Gly	Glu	Asp	Leu	Gln	Gln	Glu	Asp	Gly	Pro	Thr
			165					170						175	
Gly	Glu	Pro	Gln	Gln	Glu	Asp	Asp	Glu	Phe	Leu	Met	Ala	Thr	Asp	Val
			180					185						190	
Asp	Asp	Arg	Phe	Glu	Thr	Leu	Glu	Pro	Glu	Val	Ser	His	Glu	Glu	Thr
		195					200					205			
Glu	His	Ser	Tyr	His	Val	Glu	Glu	Thr	Val	Ser	Gln	Asp	Cys	Asn	Gln
	210					215					220				
Asp	Met	Glu	Glu	Met	Met	Ser	Glu	Gln	Glu	Asn	Pro	Asp	Ser	Ser	Glu
225					230					235					240
Pro	Val	Val	Glu	Asp	Glu	Arg	Leu	His	His	Asp	Thr	Asp	Asp	Val	Thr
			245						250					255	
Tyr	Gln	Val	Tyr	Glu	Glu	Gln	Ala	Val	Tyr	Glu	Pro	Leu	Glu	Asn	Glu
		260					265							270	
Gly	Ile	Glu	Ile	Thr	Glu	Val	Thr	Ala	Pro	Pro	Glu	Asp	Asn	Pro	Val
	275						280					285			
Glu	Asp	Ser	Gln	Val	Ile	Val	Glu	Glu	Val	Ser	Ile	Phe	Pro	Val	Glu
	290					295					300				
Glu	Gln	Gln	Glu	Val	Pro	Pro	Glu	Thr	Asn	Arg	Lys	Thr	Asp	Asp	Pro
305					310					315					320
Glu	Gln	Lys	Ala	Lys	Val	Lys	Lys	Lys	Lys	Pro	Lys	Leu	Leu	Asn	Lys
		325							330					335	
Phe	Asp	Lys	Thr	Ile	Lys	Ala	Glu	Leu	Asp	Ala	Ala	Glu	Lys	Leu	Arg
		340						345					350		
Lys	Arg	Gly	Lys	Ile	Glu	Glu	Ala	Val	Asn	Ala	Phe	Lys	Glu	Leu	Val
	355						360					365			
Arg	Lys	Tyr	Pro	Gln	Ser	Pro	Arg	Ala	Arg	Tyr	Gly	Lys	Ala	Gln	Cys
	370					375					380				
Glu	Asp	Asp	Leu	Ala	Glu	Lys	Arg	Arg	Ser	Asn	Glu	Val	Leu	Arg	Gly
385					390					395					400
Ala	Ile	Glu	Thr	Tyr	Gln	Glu	Val	Ala	Ser	Leu	Pro	Asp	Val	Pro	Ala
			405						410					415	
Asp	Leu	Leu	Lys	Leu	Ser	Leu	Lys	Arg	Arg	Ser	Asp	Arg	Gln	Gln	Phe
		420						425					430		
Leu	Gly	His	Met	Arg	Gly	Ser	Leu	Leu	Thr	Leu	Gln	Arg	Leu	Val	Gln
	435						440					445			
Leu	Phe	Pro	Asn	Asp	Thr	Ser	Leu	Lys	Asn	Asp	Leu	Gly	Val	Gly	Tyr
	450					455					460				
Leu	Leu	Ile	Gly	Asp	Asn	Asp	Asn	Ala	Lys	Lys	Val	Tyr	Glu	Glu	Val
465					470					475					480
Leu	Ser	Val	Thr	Pro	Asn	Asp	Gly	Phe	Ala	Lys	Val	His	Tyr	Gly	Phe
			485						490					495	
Ile	Leu	Lys	Ala	Gln	Asn	Lys	Ile	Ala	Glu	Ser	Ile	Pro	Tyr	Leu	Lys
		500						505					510		
Glu	Gly	Ile	Glu	Ser	Gly	Asp	Pro	Gly	Thr	Asp	Asp	Gly	Arg	Phe	Tyr
	515						520					525			
Phe	His	Leu	Gly	Asp	Ala	Met	Gln	Arg	Val	Gly	Asn	Lys	Glu	Ala	Tyr

-continued

530					535					540					
Lys	Trp	Tyr	Glu	Leu	Gly	His	Lys	Arg	Gly	His	Phe	Ala	Ser	Val	Trp
545					550					555					560
Gln	Arg	Ser	Leu	Tyr	Asn	Val	Asn	Gly	Leu	Lys	Ala	Gln	Pro	Trp	Trp
			565						570					575	
Thr	Pro	Lys	Glu	Thr	Gly	Tyr	Thr	Glu	Leu	Val	Lys	Ser	Leu	Glu	Arg
			580					585					590		
Asn	Trp	Lys	Leu	Ile	Arg	Asp	Glu	Gly	Leu	Ala	Val	Met	Asp	Lys	Ala
		595					600					605			
Lys	Gly	Leu	Phe	Leu	Pro	Glu	Asp	Glu	Asn	Leu	Arg	Glu	Lys	Gly	Asp
	610					615					620				
Trp	Ser	Gln	Phe	Thr	Leu	Trp	Gln	Gln	Gly	Arg	Arg	Asn	Glu	Asn	Ala
	625					630					635				640
Cys	Lys	Gly	Ala	Pro	Lys	Thr	Cys	Thr	Leu	Leu	Glu	Lys	Phe	Pro	Glu
			645						650					655	
Thr	Thr	Gly	Cys	Arg	Arg	Gly	Gln	Ile	Lys	Tyr	Ser	Ile	Met	His	Pro
			660					665						670	
Gly	Thr	His	Val	Trp	Pro	His	Thr	Gly	Pro	Thr	Asn	Cys	Arg	Leu	Arg
		675						680				685			
Met	His	Leu	Gly	Leu	Val	Ile	Pro	Lys	Glu	Gly	Cys	Lys	Ile	Arg	Cys
	690					695					700				
Ala	Asn	Glu	Thr	Arg	Thr	Trp	Glu	Glu	Gly	Lys	Val	Leu	Ile	Phe	Asp
	705					710					715				720
Asp	Ser	Phe	Glu	His	Glu	Val	Trp	Gln	Asp	Ala	Ser	Ser	Phe	Arg	Leu
			725						730					735	
Ile	Phe	Ile	Val	Asp	Val	Trp	His	Pro	Glu	Leu	Thr	Pro	Gln	Gln	Arg
			740					745						750	
Arg	Ser	Leu	Pro	Ala	Ile										
			755												

<210> SEQ ID NO 3

<211> LENGTH: 2324

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 3

```

cggaccgtgc aatggcccag cgtaagaatg ccaagagcag cggcaacagc agcagcagcg      60
gctccggcag cggtagcacg agtgcgggca gcagcagccc cggggcccgg agagagacaa      120
agcatggagg acacaagaat gggaggaaag gcggaacttc ttcttcacgt      180
ggtttatggt gattgcattg ctgggcgtct ggacatctgt agctgtcgtt tggtttgatc      240
ttgttgacta tgaggaagtt ctaggaaaac taggaatcta tgatgctgat ggtgatggag      300
atthtgatgt ggatgatgcc aaagttttat taggacttaa agagagatct acttcagagc      360
cagcagtcgc gccagaagag gctgagccac aactgagcc cgaggagcag gttcctgtgg      420
aggcagaacc ccagaatatc gaagatgaag caaaagaaca aattcagtc cttctccatg      480
aaatggtaca cgcagaacat gttgaggagg aagacttgca acaagaagat ggaccacag      540
gagaaccaca acaagaggat gatgagtttc ttatggcgac tgatgtagat gatagatttg      600
agaccctgga acctgaagta tctcatgaag aaaccgagca tagttaccac gtggaagaga      660
cagtttcaca agactgtaat caggatatgg aagagatgat gtctgagcag gaaaatccag      720

```

-continued

attccagtga accagtagta gaagatgaaa gattgcacca tgatacagat gatgtaacat	780
accaagtcta tgaggaacaa gcagtatatg aacctctaga aaatgaaggg atagaaatca	840
cagaagtaac tgctccccct gaggataatc ctgtagaaga ttcacaggta attgtagaag	900
aagtaagcat ttttctgtg gaagaacagc aggaagtacc accagaaaca aatagaaaaa	960
cagatgatcc agaacaaaaa gcaaaagtta agaaaaagaa gcctaaactt ttaaataaat	1020
ttgataagac tattaaagct gaacttgatg ctgcagaaaa actccgtaaa aggggaaaaa	1080
ttgaggaagc agtgaatgca tttaaagaac tagtacgcaa ataccctcag agtccacgag	1140
caagatatgg gaaggcgagc tgtgaggatg atttggtgga gaagaggaga agtaatgagg	1200
tgctacgtgg agccatcgag acctaccaag aggtggccag cctacctgat gtccctgcag	1260
acctgctgaa gctgagtgtt aagcgtcgct cagacaggca acaatttcta ggtcatatga	1320
gaggttccct gcttaccctg cagagattag ttcaactatt tccaatgat acttccttaa	1380
aaaatgacct tggcgtggga tacctcttga taggagataa tgacaatgca aagaaagtgt	1440
atgaagaggt gctgagtgtg acacctaatg atggctttgc taaagtccat tatggcttca	1500
tcctgaaggc acagaacaaa attgctgaga gcacccata tttaaaggaa ggaatagaat	1560
ccggagatcc tggcactgat gatgggagat ttattttcca cctgggggat gccatgcaga	1620
gggttgggaa caaagaggca tataagtgtg atgagcttgg gcacaagaga ggacactttg	1680
catctgtctg gcaacgctca ctctacaatg tgaatggact gaaagcacag ccttggtgga	1740
ccccaaaaga aacgggctac acagagttag taaagtcttt agaaagaaac tggaaagttaa	1800
tccgagatga aggccttgca gtgatggata aagccaaagg tctcttctg cctgaggatg	1860
aaaacctgag gaaaaaaggg gactggagcc agttcacgct gtggcagcaa ggaagaagaa	1920
atgaaaatgc ctgcaaagga gctcctaaaa cctgtacctt actagaaaag tccccgaga	1980
caacaggatg cagaagagga cagatcaaat attccatcat gcacccccgg actcacgtgt	2040
ggccgcacac agggcccaca aactgcaggc tccgaatgca cctgggcttg gtgattccca	2100
aggaaggctg caagattcga tgtgccaacg agaccaggac ctgggaggaa ggcaagggtg	2160
tcactcttga tgactccttt gagcacgagg tatggcagga tgctcatct ttcgggtga	2220
tattcatcgt ggaatgtgtg catccggaac tgacaccaca gcagagacgc agccttcag	2280
caatttagca tgaattcatg caagcttggg aaactctgga gaga	2324

<210> SEQ ID NO 4

<211> LENGTH: 31

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: EGF-like cysteine-rich repeat

<220> FEATURE:

<221> NAME/KEY: VARIANT

<222> LOCATION: (3)..(5)

<223> OTHER INFORMATION: Wherein any Xaa may be any amino acid

<220> FEATURE:

<221> NAME/KEY: VARIANT

<222> LOCATION: (6)..(7)

<223> OTHER INFORMATION: Wherein Xaa is any amino acid.

<220> FEATURE:

<221> NAME/KEY: VARIANT

<222> LOCATION: (10)

<223> OTHER INFORMATION: Wherein Xaa is any amino acid.

<220> FEATURE:

<221> NAME/KEY: VARIANT

-continued

<222> LOCATION: (14)
<223> OTHER INFORMATION: Wherein Xaa is any amino acid.
<220> FEATURE:
<221> NAME/KEY: VARIANT
<222> LOCATION: (17)..(18)
<220> FEATURE:
<221> NAME/KEY: VARIANT
<222> LOCATION: (25)..(26)
<223> OTHER INFORMATION: Wherein Xaa is any amino acid.
<220> FEATURE:
<221> NAME/KEY: VARIANT
<222> LOCATION: (29)
<223> OTHER INFORMATION: Wherein Xaa is any amino acid.

<400> SEQUENCE: 4

Cys Asp Xaa Xaa Xaa Cys Xaa Xaa Lys Xaa Gly Asn Gly Xaa Cys Asp
1 5 10 15

Xaa Xaa Cys Asn Asn Ala Ala Cys Xaa Xaa Asp Gly Xaa Asp Cys
20 25 30

<210> SEQ ID NO 5
<211> LENGTH: 1242
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 5

Met Ala Ser Pro Pro Glu Ser Asp Gly Phe Ser Asp Val Arg Lys Val
1 5 10 15

Gly Tyr Leu Arg Lys Pro Lys Ser Met His Lys Arg Phe Phe Val Leu
20 25 30

Arg Ala Ala Ser Glu Ala Gly Gly Pro Ala Arg Leu Glu Tyr Tyr Glu
35 40 45

Asn Glu Lys Lys Trp Arg His Lys Ser Ser Ala Pro Lys Arg Ser Ile
50 55 60

Pro Leu Glu Ser Cys Phe Asn Ile Asn Lys Arg Ala Asp Ser Lys Asn
65 70 75 80

Lys His Leu Val Ala Leu Tyr Thr Arg Asp Glu His Phe Ala Ile Ala
85 90 95

Ala Asp Ser Glu Ala Glu Gln Asp Ser Trp Tyr Gln Ala Leu Leu Gln
100 105 110

Leu His Asn Arg Ala Lys Gly His His Asp Gly Ala Ala Ala Leu Gly
115 120 125

Ala Gly Gly Gly Gly Gly Ser Cys Ser Gly Ser Ser Gly Leu Gly Glu
130 135 140

Ala Gly Glu Asp Leu Ser Tyr Gly Asp Val Pro Pro Gly Pro Ala Phe
145 150 155 160

Lys Glu Val Trp Gln Val Ile Leu Lys Pro Lys Gly Leu Gly Gln Thr
165 170 175

Lys Asn Leu Ile Gly Ile Tyr Arg Leu Cys Leu Thr Ser Lys Thr Ile
180 185 190

Ser Phe Val Lys Leu Asn Ser Glu Ala Ala Ala Val Val Leu Gln Leu
195 200 205

Met Asn Ile Arg Arg Cys Gly His Ser Glu Asn Phe Phe Phe Ile Glu
210 215 220

Val Gly Arg Ser Ala Val Thr Gly Pro Gly Glu Phe Trp Met Gln Val
225 230 235 240

Asp Asp Ser Val Val Ala Gln Asn Met His Glu Thr Ile Leu Glu Ala

-continued

245								250					255				
Met	Arg	Ala	Met	Ser	Asp	Glu	Phe	Arg	Pro	Arg	Ser	Lys	Ser	Gln	Ser		
			260					265					270				
Ser	Ser	Asn	Cys	Ser	Asn	Pro	Ile	Ser	Val	Pro	Leu	Arg	Arg	His	His		
		275					280					285					
Leu	Asn	Asn	Pro	Pro	Pro	Ser	Gln	Val	Gly	Leu	Thr	Arg	Arg	Ser	Arg		
	290					295					300						
Thr	Glu	Ser	Ile	Thr	Ala	Thr	Ser	Pro	Ala	Ser	Met	Val	Gly	Gly	Lys		
305					310						315				320		
Pro	Gly	Ser	Phe	Arg	Val	Arg	Ala	Ser	Ser	Asp	Gly	Glu	Gly	Thr	Met		
				325					330					335			
Ser	Arg	Pro	Ala	Ser	Val	Asp	Gly	Ser	Pro	Val	Ser	Pro	Ser	Thr	Asn		
			340					345					350				
Arg	Thr	His	Ala	His	Arg	His	Arg	Gly	Ser	Ala	Arg	Leu	His	Pro	Pro		
		355					360					365					
Leu	Asn	His	Ser	Arg	Ser	Ile	Pro	Met	Pro	Ala	Ser	Arg	Cys	Ser	Pro		
	370					375					380						
Ser	Ala	Thr	Ser	Pro	Val	Ser	Leu	Ser	Ser	Ser	Ser	Thr	Ser	Gly	His		
385					390					395					400		
Gly	Ser	Thr	Ser	Asp	Cys	Leu	Phe	Pro	Arg	Arg	Ser	Ser	Ala	Ser	Val		
				405					410					415			
Ser	Gly	Ser	Pro	Ser	Asp	Gly	Gly	Phe	Ile	Ser	Ser	Asp	Glu	Tyr	Gly		
			420					425					430				
Ser	Ser	Pro	Cys	Asp	Phe	Arg	Ser	Ser	Phe	Arg	Ser	Val	Thr	Pro	Asp		
		435					440					445					
Ser	Leu	Gly	His	Thr	Pro	Pro	Ala	Arg	Gly	Glu	Glu	Glu	Leu	Ser	Asn		
	450					455					460						
Tyr	Ile	Cys	Met	Gly	Gly	Lys	Gly	Pro	Ser	Thr	Leu	Thr	Ala	Pro	Asn		
465				470						475					480		
Gly	His	Tyr	Ile	Leu	Ser	Arg	Gly	Gly	Asn	Gly	His	Arg	Cys	Thr	Pro		
			485						490					495			
Gly	Thr	Gly	Leu	Gly	Thr	Ser	Pro	Ala	Leu	Ala	Gly	Asp	Glu	Ala	Ala		
			500					505					510				
Ser	Ala	Ala	Asp	Leu	Asp	Asn	Arg	Phe	Arg	Lys	Arg	Thr	His	Ser	Ala		
		515					520					525					
Gly	Thr	Ser	Pro	Thr	Ile	Thr	His	Gln	Lys	Thr	Pro	Ser	Gln	Ser	Ser		
	530					535						540					
Val	Ala	Ser	Ile	Glu	Glu	Tyr	Thr	Glu	Met	Met	Pro	Ala	Tyr	Pro	Pro		
545					550					555					560		
Gly	Gly	Gly	Ser	Gly	Gly	Arg	Leu	Pro	Gly	His	Arg	His	Ser	Ala	Phe		
			565						570					575			
Val	Pro	Thr	Arg	Ser	Tyr	Pro	Glu	Glu	Gly	Leu	Glu	Met	His	Pro	Leu		
			580					585					590				
Glu	Arg	Arg	Gly	Gly	His	His	Arg	Pro	Asp	Ser	Ser	Thr	Leu	His	Thr		
		595					600					605					
Asp	Asp	Gly	Tyr	Met	Pro	Met	Ser	Pro	Gly	Val	Ala	Pro	Val	Pro	Ser		
	610					615					620						
Gly	Arg	Lys	Gly	Ser	Gly	Asp	Tyr	Met	Pro	Met	Ser	Pro	Lys	Ser	Val		
625					630					635					640		
Ser	Ala	Pro	Gln	Gln	Ile	Ile	Asn	Pro	Ile	Arg	Arg	His	Pro	Gln	Arg		
			645						650					655			

-continued

Val	Asp	Pro	Asn	Gly	Tyr	Met	Met	Met	Ser	Pro	Ser	Gly	Gly	Cys	Ser	660	665	670	
Pro	Asp	Ile	Gly	Gly	Gly	Pro	Ser	Ser	Ser	Ser	Ser	Ser	Ser	Asn	Ala	675	680	685	
Val	Pro	Ser	Gly	Thr	Ser	Tyr	Gly	Lys	Leu	Trp	Thr	Asn	Gly	Val	Gly	690	695	700	
Gly	His	His	Ser	His	Val	Leu	Pro	His	Pro	Lys	Pro	Pro	Val	Glu	Ser	705	710	715	720
Ser	Gly	Gly	Lys	Leu	Leu	Pro	Cys	Thr	Gly	Asp	Tyr	Met	Asn	Met	Ser	725	730	735	
Pro	Val	Gly	Asp	Ser	Asn	Thr	Ser	Ser	Pro	Ser	Asp	Cys	Tyr	Tyr	Gly	740	745	750	
Pro	Glu	Asp	Pro	Gln	His	Lys	Pro	Val	Leu	Ser	Tyr	Tyr	Ser	Leu	Pro	755	760	765	
Arg	Ser	Phe	Lys	His	Thr	Gln	Arg	Pro	Gly	Glu	Pro	Glu	Glu	Gly	Ala	770	775	780	
Arg	His	Gln	His	Leu	Arg	Leu	Ser	Thr	Ser	Ser	Gly	Arg	Leu	Leu	Tyr	785	790	795	800
Ala	Ala	Thr	Ala	Asp	Ser	Ser	Ser	Ser	Ser	Thr	Ser	Ser	Asp	Ser	Leu	805	810	815	
Gly	Gly	Gly	Tyr	Cys	Gly	Ala	Arg	Leu	Glu	Pro	Ser	Leu	Pro	His	Pro	820	825	830	
His	His	Gln	Val	Leu	Gln	Pro	His	Leu	Pro	Arg	Lys	Val	Asp	Thr	Ala	835	840	845	
Ala	Gln	Thr	Asn	Ser	Arg	Leu	Ala	Arg	Pro	Thr	Arg	Leu	Ser	Leu	Gly	850	855	860	
Asp	Pro	Lys	Ala	Ser	Thr	Leu	Pro	Arg	Ala	Arg	Glu	Gln	Gln	Gln	Gln	865	870	875	880
Gln	Gln	Pro	Leu	Leu	His	Pro	Pro	Glu	Pro	Lys	Ser	Pro	Gly	Glu	Tyr	885	890	895	
Val	Asn	Ile	Glu	Phe	Gly	Ser	Asp	Gln	Ser	Gly	Tyr	Leu	Ser	Gly	Pro	900	905	910	
Val	Ala	Phe	His	Ser	Ser	Pro	Ser	Val	Arg	Cys	Pro	Ser	Gln	Leu	Gln	915	920	925	
Pro	Ala	Pro	Arg	Glu	Glu	Glu	Thr	Gly	Thr	Glu	Glu	Tyr	Met	Lys	Met	930	935	940	
Asp	Leu	Gly	Pro	Gly	Arg	Arg	Ala	Ala	Trp	Gln	Glu	Ser	Thr	Gly	Val	945	950	955	960
Glu	Met	Gly	Arg	Leu	Gly	Pro	Ala	Pro	Pro	Gly	Ala	Ala	Ser	Ile	Cys	965	970	975	
Arg	Pro	Thr	Arg	Ala	Val	Pro	Ser	Ser	Arg	Gly	Asp	Tyr	Met	Thr	Met	980	985	990	
Gln	Met	Ser	Cys	Pro	Arg	Gln	Ser	Tyr	Val	Asp	Thr	Ser	Pro	Ala	Ala	995	1000	1005	
Pro	Val	Ser	Tyr	Ala	Asp	Met	Arg	Thr	Gly	Ile	Ala	Ala	Glu	Glu	Val	1010	1015	1020	
Ser	Leu	Pro	Arg	Ala	Thr	Met	Ala	Ala	Ala	Ser	Ser	Ser	Ser	Ala	Ala	1025	1030	1035	1040
Ser	Ala	Ser	Pro	Thr	Gly	Pro	Gln	Gly	Ala	Ala	Glu	Leu	Ala	Ala	His	1045	1050	1055	

-continued

Ser Ser Leu Leu Gly Gly Pro Gln Gly Pro Gly Gly Met Ser Ala Phe
1060 1065 1070

Thr Arg Val Asn Leu Ser Pro Asn Arg Asn Gln Ser Ala Lys Val Ile
1075 1080 1085

Arg Ala Asp Pro Gln Gly Cys Arg Arg Arg His Ser Ser Glu Thr Phe
1090 1095 1100

Ser Ser Thr Pro Ser Ala Thr Arg Val Gly Asn Thr Val Pro Phe Gly
1105 1110 1115 1120

Ala Gly Ala Ala Val Gly Gly Gly Gly Gly Ser Ser Ser Ser Ser Glu
1125 1130 1135

Asp Val Lys Arg His Ser Ser Ala Ser Phe Glu Asn Val Trp Leu Arg
1140 1145 1150

Pro Gly Glu Leu Gly Gly Ala Pro Lys Glu Pro Ala Lys Leu Cys Gly
1155 1160 1165

Ala Ala Gly Gly Leu Glu Asn Gly Leu Asn Tyr Ile Asp Leu Asp Leu
1170 1175 1180

Val Lys Asp Phe Lys Gln Cys Pro Gln Glu Cys Thr Pro Glu Pro Gln
1185 1190 1195 1200

Pro Pro Pro Pro Pro Pro Pro His Gln Pro Leu Gly Ser Gly Glu Ser
1205 1210 1215

Ser Ser Thr Arg Arg Ser Ser Glu Asp Leu Ser Ala Tyr Ala Ser Ile
1220 1225 1230

Ser Phe Gln Lys Gln Pro Glu Asp Arg Gln
1235 1240

<210> SEQ ID NO 6

<211> LENGTH: 5828

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 6

```

cgggcggcgcg gtcggagggg gccggcgcg agagccagac gccgccgctt gttttggttg      60
gggctctcgg caactctccg aggaggagga ggaggaggga ggaggggaga agtaactgca      120
gcggcagcgc cctcccgagg aacaggcgctc ttccccgaac ccttcccaaa cctcccccat      180
ccccctctgc ccttgtcccc tccctctctc ccagccgcgc tggagcgagg ggcagggatg      240
agtctgtccc tccggccggt cccagctgc agtggtgcc cggtatcggt tcgcatggaa      300
aagccacttt ctccaccgc cgagatgggc ccgatgggg ctgcagagga cgcgcccgcg      360
ggcggcggca gcagcagcag cagcagcagc agcaacagca acagccgcag cgcgcgggtc      420
tctgcgactg agctggtatt tgggcggctg gtggcggtg ggacggttg ggggtgggag      480
gaggcgaagg aggagggaga accccgtgca acgttgggac ttggcaaccc gcctccccct      540
gccaaggat atttaatttg cctcgggaat cgctgcttcc agaggggaac tcaggaggga      600
aggcgcgcg gcgcgcgcgc tccctggagg gcaccgcagg gacccccgac tgtcgctcc      660
ctgtgccgga ctccagccgg ggcgacgaga gatgcattt cgctccttcc tgggtggcggc      720
ggcggtgag aggagacttg gctctcggag gatcggggct gccctcacc cggaacgaact      780
gcctccccgc cggcgtgaag gcgccgaaaa ctccggtcgg gctctctcct gggctcagca      840
gctgcgtcct ccttcagctg cccctccccg gcgcgggggg cggcgtggat ttcagagtcg      900
gggtttctgc tgctccagc cctgtttgca tgtgccgggc cgcggcgagg agcctccgcc      960

```

-continued

ccccaccg	ttgttttctg	gagcctccct	ctgctcagcg	ttggtggtgg	cgtggtgcagc	1020
atggcgagcc	ctccgagag	cgatggcttc	tcggacgtgc	gcaaggtggg	ctacctgcgc	1080
aaaccaaga	gcatacaca	acgtttcttc	gtactgcgcg	cggccagcga	ggctgggggc	1140
cggcgcgcc	tcgagtacta	cgagaacgag	aagaagtggc	ggcacaagtc	gagcgccccc	1200
aaacgctcga	tcccccttga	gagctgcttc	aacatcaaca	agcgggctga	ctccaagaac	1260
aagcacctgg	tggtctctta	caccggggac	gagcactttg	ccatcgcggc	ggacagcgag	1320
gccgagcaag	acagctggta	ccaggctctc	ctacagctgc	acaaccgtgc	taagggccac	1380
cacgacggag	ctcgccctcc	cggggcggga	ggtggtgggg	gcagctgcag	cggcagctcc	1440
ggccttggtg	aggctgggga	ggacttgagc	tacggtgacg	tgcccccagg	accgcatttc	1500
aaagaggtct	ggcaagtgat	cctgaagccc	aagggcctgg	gtcagacaaa	gaacctgatt	1560
ggtatctacc	gcctttgcct	gaccagcaag	accatcagct	tcgtgaagct	gaactcggag	1620
gcagcgcccg	tggtgtctga	gctgatgaac	atcaggcgct	gtggccactc	ggaaaacttc	1680
ttcttcacgc	aggtagggcg	ttctgccgtg	acggggcccg	gggagttctg	gatgcagggtg	1740
gatgactctg	tggtggccca	gaacatgcac	gagaccatcc	tggaggccat	gcggggccatg	1800
agtgatgagt	tccgcctctg	cagcaagagc	cagtctctct	ccaactgtct	taaccccatc	1860
agcgtccccc	tgcgcgggca	ccatctcaac	aatccccgcg	ccagccaggt	ggggctgacc	1920
cgcgatcac	gcactgagag	catcacccgc	acctcccccg	ccagcatggt	ggcggggaag	1980
ccaggctcct	tccgtgtccg	cgcctccagt	gacggcgaag	gcaccatgtc	ccgccagacc	2040
tcggtggacg	gcagccctgt	gagtcaccag	accaacagaa	cccacgcca	ccggcatcgg	2100
ggcagcgccc	gggtgcaccc	cccgtcaaac	cacagccgct	ccatcccat	gccggttccc	2160
cgtctctcgc	cttcggccac	cagcccggtc	agtctgtcgt	ccagtagcac	cagtggccat	2220
ggctccacct	cggattgtct	cttccacagg	cgatctagt	cttcgggtgc	tggttccccc	2280
agcgatggcg	gtttcatctc	ctcggatgag	tatggctcca	gtccctcgca	tttccggagt	2340
tccttccgca	gtgtcactcc	ggattccctg	ggccacaccc	caccagcccg	cggtagggag	2400
gagctaagca	actatatctg	catgggtggc	aaggggcctc	ccacctgac	cgccecaaac	2460
ggtcactaca	ttttgtctcg	gggtggcaat	ggccaccgct	gcacccagc	aacaggcttg	2520
ggcacgagtc	cagccttggc	tggggatgaa	gcagccagtg	ctgcagatct	ggataatcgg	2580
ttccgaaaga	gaactcactc	ggcaggcaca	tcccctacca	ttacccacca	gaagaccccg	2640
tcccagtcct	cagtggcttc	cattgaggag	tacacagaga	tgatgcctgc	ctaccacca	2700
ggaggtggca	gtggaggccg	actgccggga	cacaggcact	cgcctctcgt	gcccaccgcg	2760
tcctaccacg	aggagggtct	ggaaatgcac	cccttgagc	gtcggggggg	gcaccaccgc	2820
ccagacagct	ccacctcca	cacggatgat	ggctacatgc	ccatgtcccc	aggggtggcc	2880
ccagtgcaca	gtggccgaaa	gggcagtgga	gactatatgc	ccatgagccc	caagagcgta	2940
tctgccccac	agcagatcat	caatcccatc	agacgccatc	cccagagagt	ggaccccaat	3000
ggctacatga	tgatgtcccc	cagcgggtggc	tgtctctctg	acattggagg	tggccccagc	3060
agcagcagca	gcagcagcaa	gcgcgtccct	tccgggacca	gctatggaaa	gctgtggaca	3120
aacggggtag	ggggccacca	ctctcatgtc	ttgcctcacc	ccaaaccccc	agtggagagc	3180
agcgggtgga	agctcttacc	ttgcacaggt	gactacatga	acatgtcacc	agtgggggac	3240

-continued

tccaacacca gcagcccctc cgactgctac tacggccctg aggacccccca gcacaagcca	3300
gtcctctcct actactcatt gccaaagatcc tttaagcaca cccagcgccc cggggagccg	3360
gaggaggggtg cccggcatca gcacctccgc ctttccacta gctctggteg ccttctctat	3420
gctgcaacag cagatgattc ttctcttcc accagcagcg acagcctggg tgggggatac	3480
tgcggggcta ggttgagcc cagccttcca catccccacc atcaggttct gcagccccat	3540
ctgcctcgaa aggtggacac agctgctcag accaatagcc gcctggcccc gcccaagagg	3600
ctgtccctgg gggatcccaa ggcagcacc ttacctcggg cccgagagca gcagcagcag	3660
cagcagccct tctgcaccc tccagagccc aagagcccg gggaatatgt caatattgaa	3720
tttgggagtg atcagtctgg ctacttgtct ggcccggtag ctttccacag ctcaccttct	3780
gtcaggtgtc catccagct ccagccagct cccagagagg aagagactgg cactgaggag	3840
tacatgaaga tggacctggg gccggggccgg agggcagcct ggcaggagag cactggggtc	3900
gagatgggca gactgggcc tgcacctccc ggggctgcta gcatttgag gcctaccgg	3960
gcagtggcca gcagccgggg tgactacatg accatgcaga tgagttgtcc ccgtcagagc	4020
tacgtggaca cctcgccagc tgcccctgta agctatgtg acatgcgaac aggcattgct	4080
gcagaggagg tgagctgcc cagggccacc atggtctgtg cctcctcacc ctcagcagcc	4140
tctgtctccc cgactggggc tcaaggggca gcagagctgg ctgcccactc gtccctgctg	4200
ggggggccac aaggacctgg gggcatgagc gccttcaccc gggtgaaact cagtccaaac	4260
cgcaaccaga gtgccaaagt gatccgtgca gaccacaaag ggtgccggcg gaggcatagc	4320
tccgagactt tctcctcaac acccagtgcc acccggttg gcaacacagt gccctttgga	4380
gcggggggcag cagtaggggg cgggtggcgt agcagcagca gcagcgagga tgtgaaacgc	4440
cacagctctg cttcctttga gaatgtgtgg ctgaggcctg gggagcttg gggagcccc	4500
aaggagccag ccaaaactgt tggggctgct gggggtttg agaatggtct taactacata	4560
gacctggatt tggtaagga cttcaaacag tgccctcagg agtgcacccc tgaaccgcag	4620
cctccccac ccccccccc tcatcaaccc ctgggcagcg gtgagagcag ctcacccgc	4680
cgctcaagtg aggatttaag cgctatgcc agcatcagtt tccagaagca gccagaggac	4740
cgtcagtagc tcaactggac atcacagcag aatgaagacc taaatgacct cagcaaatcc	4800
tcttctaact catgggtacc cagactctaa atatttcag attcacaact aggacctcat	4860
atcttctca tcagtagatg gtacgatgca tccatttcag tttgtttact ttatccaac	4920
ctcaggattt cattgactga actgcacgtt ctatatttg ccaagcgaaa aaaaaaatg	4980
cactgtgaca ccagaataat gagtctgcat aaacttcac ttcaacctta aggacttagc	5040
tggccacagt gagctgatg gcccaaccac gtgtcatgag agaatgggtt tactctcaat	5100
gcattttcaa gatacatctc atctgtgct gaaactgtg acgacaaagc atcattgtaa	5160
attatttcac acaaaactgt tcacgttggg tggagagagt attaaatatt taacataggt	5220
tttgatttat atgtgtaatt ttttaaatga aaatgtaact tttcttacag cacatcttt	5280
ttttggatgt gggatggagg tatacaatgt tctgttgtaa agagtggagc aaatgcttaa	5340
aacaaggctt aaaagagtag aatagggtat gatccttggt ttaagattgt aattcagaaa	5400
acataatata agaatcatag tgccatagat ggttctcaat tgtatagtta tatttgctga	5460
tactatctct tgtcatataa acctgatggt gagctgagtt ccttataaga attaatctta	5520

-continued

```

attttgtatt ttttctgta agacaatagg ccatgttaat taaactgaag aaggatatat 5580
ttggctgggt gttttcaaat gtcagcttaa aattggtaat tgaatggaag caaaattata 5640
agaagaggaa attaaagtct tccattgcat gtattgtaaa cagaaggaga tgggtgattc 5700
cttcaattca aaagctctct ttggaatgaa caatgtgggc gtttgtaaat tctggaaatg 5760
tctttctatt cataataaac tagatactgt tgatctttta aaaaaaaaaa aaaaaaaaaa 5820
aaaaaaaaa 5828

```

```

<210> SEQ ID NO 7
<211> LENGTH: 8
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: FLAG
        epitope

```

```

<400> SEQUENCE: 7

```

```

Asp Tyr Lys Asp Asp Asp Asp Lys
 1           5

```

```

<210> SEQ ID NO 8
<211> LENGTH: 17
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:Primer

```

```

<400> SEQUENCE: 8

```

```

gggggaattt gtcaata 17

```

```

<210> SEQ ID NO 9
<211> LENGTH: 16
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:Primer

```

```

<400> SEQUENCE: 9

```

```

gaatttggtta atattg 16

```

What is claimed is:

1. A method for diagnosing a malignant neoplasm in a mammal, comprising contacting a bodily fluid from said mammal with an antibody or fragment thereof which binds to an human aspartyl (asparaginy) beta-hydroxylase (HAAH) polypeptide under conditions sufficient to form an antigen-antibody complex and detecting the antigen-antibody complex, wherein an increase in antigen-antibody complex indicates an increase in HAAH level compared to a normal noncancerous control, said increase being indicative of a malignant neoplasm and wherein said antibody is a monoclonal antibody produced by hybridoma ATCC designation 3386.

2. The method of claim 1, wherein said neoplasm is derived from endodermal tissue.

3. The method of claim 1, wherein said neoplasm is selected from the group consisting of colon cancer, breast cancer, pancreatic cancer, liver cancer, and cancer of the bile ducts.

4. The method of claim 1, wherein said neoplasm is a cancer of the central nervous system (CNS).

5. The method of claim 1, wherein said bodily fluid is selected from the group consisting of a CNS-derived bodily fluid, blood, serum, urine, saliva, sputum, lung effusion, and ascites fluid.

6. The method of claim 5, wherein said bodily fluid is serum.

7. The method of claim 1, wherein the antigen-antibody complex is detected by a label selected from consisting of an enzymatic label, a fluorescent label, a chemiluminescent label, a radioactive label, and a dye label.

8. The method of claim 1, wherein said fragment of said monoclonal antibody is a single chain Fv molecule.

9. The method of claim 1, wherein said neoplasm is a hepatocellular carcinoma.

10. The method of claim 1, wherein said neoplasm is a cholangiocarcinoma.

11. The method of claim 1, wherein said neoplasm is a glioblastoma.

12. The method of claim 1, wherein said neoplasm is a neuroblastoma.

13. The method of claim 1, wherein said neoplasm is a pancreatic cancer.

14. The method of claim 1, wherein said tumor is a neuro-ectodermal tumor.

15. The method of claim 1, wherein said antibody comprises a first HAAH-specific antibody and a second HAAH-specific antibody.

16. A method for prognosis of a malignant neoplasm of a mammal, comprising:

contacting a bodily fluid from said mammal with an antibody or fragment thereof which binds to an HAAH polypeptide under conditions sufficient to form an antigen-antibody complex and detecting the antigen-antibody complex;

quantitating the amount of complex to determine the level of HAAH in said fluid; and

comparing the level of HAAH in said fluid with a normal noncancerous control level of HAAH, wherein increasing levels of HAAH over time indicates an adverse prognosis, and wherein said antibody is a monoclonal antibody produced by hybridoma ATCC designation 3386.

17. The method of claim 11, wherein said fragment of said monoclonal antibody is a single chain Fv molecule.

18. The method of claim 11, wherein the antigen-antibody complex is detected by a label selected from consisting of an enzymatic label, a fluorescent label, a chemiluminescent label, a radioactive label, and a dye label.

19. The method of claim 1, wherein said neoplasm is a cancer of the central nervous system (CNS).

20. A method for diagnosing a malignant neoplasm in a mammal, comprising contacting a bodily tissue from said mammal with an antibody or fragment thereof which binds to an human aspartyl (asparaginy) beta-hydroxylase (HAAH) polypeptide under conditions sufficient to form an antigen-antibody complex and detecting the antigen-antibody complex, wherein an increase in antigen-antibody complex indicates an increase in HAAH level compared to a normal noncancerous control, said increase being indicative of a malignant neoplasm, wherein said antibody is a monoclonal antibody produced by hybridoma ATCC designation 3386, and wherein said fragment of said monoclonal antibody is a single chain Fv molecule.

* * * * *

专利名称(译)	恶性肿瘤的诊断和治疗		
公开(公告)号	US20100136581A1	公开(公告)日	2010-06-03
申请号	US12/625729	申请日	2009-11-25
[标]申请(专利权)人(译)	罗德岛医院		
申请(专利权)人(译)	罗德岛医院		
当前申请(专利权)人(译)	罗德岛医院		
[标]发明人	WANDS JACK R DE LA MONTE SUZANNE M INCE NEDIM CARLSON ROLF I		
发明人	WANDS, JACK R. DE LA MONTE, SUZANNE M. INCE, NEDIM CARLSON, ROLF I.		
IPC分类号	G01N33/53 A61K31/00 A61K31/4412 A61K31/45 A61K31/7088 A61K38/43 A61K39/395 A61K45/00 A61K48/00 A61P35/00 C07K14/47 C07K16/40 C12N9/02 C12N15/09 C12N15/113 C12P21/08 C12Q1 /02 C12Q1/26 G01N33/573 G01N33/574 G01N33/577		
CPC分类号	A61K31/00 G01N2500/10 A61K2039/505 C07K14/47 C07K16/40 C12N9/0071 C12N15/1137 C12N2310/111 C12N2310/315 C12N2799/027 C12Q1/26 C12Y114/11016 G01N33/574 G01N33 /57484 G01N2500/00 A61K31/4412		
外部链接	Espacenet USPTO		

摘要(译)

本发明的特征在于一种通过使来自哺乳动物的体液与结合人天冬氨酰(天冬酰胺酰)β-羟化酶(HAAH)多肽的抗体接触以及通过抑制HAAH治疗恶性肿瘤的方法来诊断哺乳动物的恶性肿瘤的方法。

