



(19) **United States**

(12) **Patent Application Publication**
Frost et al.

(10) **Pub. No.: US 2010/0203063 A1**
(43) **Pub. Date: Aug. 12, 2010**

(54) **PESTIVIRUS SPECIES**

Publication Classification

(75) Inventors: **Melinda Jane Frost**, New South
Wales (AU); **Peter Daniel**
Kirkland, New South Wales (AU);
Deborah Susan Finlaison, New
South Wales (AU)

Correspondence Address:
KNOBBE MARTENS OLSON & BEAR LLP
2040 MAIN STREET, FOURTEENTH FLOOR
IRVINE, CA 92614 (US)

(73) Assignee: **Minister for Primary Industries**
For And On Behalf Of The State
Of New South Wales, Sydney (AU)

(21) Appl. No.: **12/298,026**

(22) PCT Filed: **Apr. 20, 2007**

(86) PCT No.: **PCT/AU2007/000521**

§ 371 (c)(1),
(2), (4) Date: **Jan. 15, 2009**

(30) **Foreign Application Priority Data**

Apr. 21, 2006 (AU) 2006902089

(51) Int. Cl.	
<i>A61K 39/42</i>	(2006.01)
<i>C07H 21/02</i>	(2006.01)
<i>C07H 21/04</i>	(2006.01)
<i>C07K 14/005</i>	(2006.01)
<i>C12Q 1/70</i>	(2006.01)
<i>G01N 33/53</i>	(2006.01)
<i>C12Q 1/68</i>	(2006.01)
<i>C12P 21/04</i>	(2006.01)
<i>A61K 39/12</i>	(2006.01)
<i>A61K 31/7088</i>	(2006.01)
<i>A61K 38/16</i>	(2006.01)
<i>C12N 15/63</i>	(2006.01)
<i>C12N 5/10</i>	(2006.01)
<i>C12P 21/06</i>	(2006.01)
<i>A61P 31/12</i>	(2006.01)
(52) U.S. Cl.	424/159.1; 536/23.1; 530/350;
	435/5; 435/7.1; 435/6; 435/69.6; 424/204.1;
	424/186.1; 514/44 R; 514/12; 435/320.1;
	435/325; 435/69.1

(57) **ABSTRACT**

The application relates to a pestivirus, designated PMC virus, that is associated with porcine myocarditis syndrome, and the gene and protein sequences derived therefrom. The application further relates to detection methods, vaccine therapeutics, and diagnostic methods using the PMC virus or gene/protein sequences derived therefrom.

Figure 1

gtataacgac	agtagttcaa	gtgtcgttat	gcatcattgg	ccataacaaa	ttatctaatt	60
tggaataggg	acctgcgacc	tgtacgaagg	ccgagcgtcg	gtagccattc	cgactagtag	120
gactagtaca	aataggtcaa	ctgggtgagc	aggtgagtg	gctgcagcgg	ctaagcgggtg	180
agtacaccgt	attcgtcaac	agggtgctact	ggaaggatc	acccactagc	gatgcctgtg	240
tggaacgagga	catgtccaag	ccaatgttat	cagtagcggg	ggtcggttact	gagaaagctg	300
cccagaatgg	gtagttgcac	atacagctcg	ataggatgcc	ggcggatgcc	ctgtattttg	360
accagtataa	atattatccg	ttgtaaagca	tatgaatact	tttactttta	atacatatgg	420
agggagtgg	gaaggaaaca	tgttcttttag	aactgcaccc	acgccgccac	caggggtgcca	480
agaaccgggt	tacacaagca	caatgagacc	aatttttggc	gaaccccatc	cacctctaca	540
caaacacagc	acgttaaaat	tgccacattg	gagggggatc	aaaacaatta	gagttaagaa	600
gagagaattg	ccaaagaagg	gcgattgtag	caactcaaca	acagctcca	cttcgggggt	660
gtacgttgaa	tttagggctg	tgttctataa	agattcacag	ggcacgggat	accatcgtgt	720
accgctagaa	ctttgtacaa	accaagagag	gtgcgagggg	tccaagtgtg	tagggagaat	780
gacagggctc	gatggcaggt	tgtacaacgt	tttagtatgt	ccggacgatt	gtatcctctt	840
tgagagacac	tgtagaggtc	aaacagtcgt	cctgaaatgg	atltccaacc	ccttgacatc	900
accactttgg	gtccagagtt	gttctgacga	caaaggagca	aaacccaagg	tgaaccataa	960
agacgacagg	atgaagcaag	gaaaaatagt	gacaaagcct	aaagagactg	aagcagatca	1020
aaaaactaga	ccaccagatg	ccacgatagt	ggttgacggg	cagaagtatc	aggtgaggaa	1080
gaaggggaaa	gcgaaaccca	agactcaaga	cggcttatac	cacaacaaga	acaaaccaga	1140
agcgtccagg	aagaagcttg	agaaggcctt	gctagcatgg	gcaatattag	cctgcctatt	1200
ggtggtaccg	gtagggcca	ccaacgtgac	acaatgggaa	ttatgggaca	ataaaagtac	1260
tacagacata	catagcgtca	tgttttctag	agggattaaa	aggagtctgc	atggaatttg	1320
gcccaacaaa	atctgcaaa	ggatccctac	acatctagca	gcagactatg	aactgaaaag	1380
gattcacggg	atgggtggatg	caagcccat	gaccaacttc	acatggtgta	ggctacagag	1440
acatgagtg	aacaagcatg	ggtggtgcaa	ctggtacaat	atagagccgt	ggatcaactc	1500
catgaataat	acccaaggac	tattaaacac	tggagacaat	ttcactgagt	gcgcagtcac	1560
atgcaggtat	gatgcagact	taggggtgaa	tatagtgact	caagccagga	ctactccaac	1620
tatcctgact	ggctgtaaga	aagggcacia	cttctctttc	tcaggggagg	tcagggcctc	1680
accctgcaac	tttgagttaa	ctgctgaaga	cttgctcagg	atcatggatc	acaccaaactg	1740
cgagggattt	acctacttcg	gggaaggaa	cgttgacggg	tacaccgagg	tagtagagaa	1800
ggccagggtca	agtggtttca	gggctctcac	atggttgtcg	agtaagattg	aaaacaccaa	1860
gaaaaaaata	ttcggagctg	aagccagtc	ttactgcccc	gtggctaa	gggtcttcaa	1920
cattatttat	accaacaatt	gcaccccgct	tggactgcca	gataagtcaa	aaattatag	1980
accaggaacc	tttgacatca	gtggcaggga	tgaattcata	tttccaaaac	tcccctacca	2040
cgtagatgac	ttcattctac	tgagcttaat	tgcaatgtct	gattttgtct	cagagacatc	2100
aagtataatc	taactggctt	tgactacact	aatgccaagt	aatgacaaca	gggactctcg	2160
gatggacctg	gacccaaata	aactaaacct	tactgcaact	aaatccgtgg	caagtgtggt	2220
ccctacatcg	gtgaatgtgt	taggtgaatg	gggtgtgcgt	aaaccaagtt	ggtggcctta	2280
ttccgocgaa	atcactaatc	tgataggagg	tgatcatcacc	gtggcagact	tagttataca	2340
gaccattgaa	gaattgctaa	atltgtggac	cgaagcaaca	gctgtggcat	ttctggctgc	2400
tctaataaaa	atlttttagag	gccagccgat	ccaagcggta	gcatggttaa	tcatcatagg	2460
gggagcaca	gccaaaacct	gcaaccctga	atltcatgtac	gcattagcga	aaaataccag	2520
cataggttca	ttaggaccag	aatcactgac	gacaagggtg	taccaactaa	ccagcggttt	2580
caaactcact	gacagcacga	ttgaagtcac	ctgtgtgggt	gctaacaatga	ggattcatgt	2640
agtgtgcccc	cttgtaagtg	acagatattt	ggccataaac	caccctagag	cactgccaac	2700
aacggcgtgg	ttcaggaaaa	tacacactca	gcatgaggta	ccaagagaaa	gaatcatgag	2760
tgagtcacaaa	aggaggtaca	ctgtccttg	tggttctaaa	ccagtgggtg	ggtcaacaac	2820
acaattcaac	ccaatatcta	tatctacccc	aagctttgaa	cttgaatgcc	ctaggggttg	2880
gactggggct	gtagagtgtg	cactagtctc	ccatcaact	ctgacaacag	agactatatt	2940
cacatacagg	aagcccaaac	cattcggact	tgaaaactgg	tgcaagtata	cagtgggtgga	3000
gaaagggatc	ctgtattctt	gtaaatlttg	gggcaattca	acatgcatca	aagggcttat	3060
agttaaaagg	caacgggaag	acaaagtaag	gtactgtgaa	tggtgtggtt	ataagttcag	3120
ttcaccacaa	ggactgcctc	agtatccact	gggattgtgt	gagaaaagac	aatcagaagg	3180
actcagggat	tatgggtgact	tcccatgctg	caacaacggc	acttgtattg	acaagaagg	3240
tagtgtgcaa	tgtctacatag	gggataagaa	agttaccgtg	aagctgtata	atgcctcact	3300
atltggcccc	tgtccctgca	aacctatagt	gtataactcc	caggggcccc	cagcgcctaa	3360
gacctgcact	tataggtggg	cctcaacatt	agaaaaataa	tattatgaa	ccagggacag	3420
ctactaccag	caatacatta	taaagtcagg	gtaacaaat	tggtttgatc	tacagcaaaa	3480

ggatcatgtg	gcagactgga	tcacaaaata	ctttccaata	ataatagtgg	ccttggttagg	3540
gggcagaggg	accttgtggg	tgttgatagc	ttatgagttg	ctaactcagt	atgaggtagt	3600
aggagacgag	aacatagtgg	otcaagctga	agccttggtg	atcggaaaaca	tcttgatgag	3660
tttagactta	gagataatta	gctgcttcct	tctggtgttg	atcgtggtga	aaaaacaagc	3720
tgtcaggaga	acgttggctt	tactgtttca	ttggataact	atgaacccat	tccagtcagt	3780
aatgatcaca	gtggtctact	tcgtcggttt	ggtgagggcc	gaagagggaa	ctaaagaggg	3840
tagtacaagc	ggggcaacca	tccatgtagt	tgcaatactg	ttattcctct	tgtaccacac	3900
agtgaagtat	aaggacttta	acatagcaat	gatcttactt	ataacattgt	ccctgaaaag	3960
ctcatcctac	atacatacca	gcttgtatga	aattccattg	cttgtggctg	taataagctt	4020
cacatgctcc	atatacattt	ttgacttgca	ggtaaagagc	aagctagtgg	ccccaactat	4080
aggataaatt	ggagttacc	tagcaatgag	agttttgtgg	ctggtaaggg	aaatgactat	4140
accaaccccc	tctgtgtcca	ttagtctgat	agatccaaaag	atggtcataa	tactctactt	4200
gataccctca	actattacag	tcaatcacaa	cctagaccta	gcaagttatt	gcttgaaact	4260
gggacctttt	atcctatcat	tcctaacaat	gtgggtggat	gttgtcatcc	tccgtctcat	4320
gctgccttgg	tacgaactag	taaaagtcta	ctacctaana	aagaagaaaag	aggatgtgga	4380
aacatggttc	caaaattcag	gaatatccac	ccaagaaact	tcccatacag	gatttgattt	4440
ttctagcccc	gggaggggag	tgcacacact	accaatgcaa	aataaaaacca	aattttgtag	4500
gactgcttac	atgactgtac	taagggcttt	ggtgataaca	gccatcagca	gtgtctggaa	4560
accaataaatt	ttagcagaac	tcctaataga	ggcagtgtat	tggacacaca	ttaaaatagc	4620
caaaagaattg	gcgggggtcaa	gcaggttcgt	tgctaggttc	attgcatcta	ttatagagtt	4680
gaattggggc	atggacgaaa	aagaagcatt	tcggtacaaa	agattttacc	tattatcatc	4740
caaaaataaca	gatctaattg	ttaagcacia	aatccaaaat	gagacagtaa	aatcctgggt	4800
tgaagaaact	gaaatatttg	gaatacaaaa	agtggcaatg	gtgataaggg	ctcattctct	4860
gagtttggag	ccaaatgcca	tcctttgctc	cgtttgtgaa	gaaaaacaaa	atcaaaaagc	4920
caaaaggccc	tgccttaagt	gtggtagtag	aggcactcaa	ataaagtgtg	ggctgacact	4980
ggccagagtt	gaggaagaac	attacaanaa	aatatacatc	ctcgaaggcc	aagatgaaac	5040
tcccattgagg	aaagaagaaa	gacagcaagt	aaacttatgt	tctaggggtg	ctctgttcc	5100
taggaatctt	cctatcttag	cttcaaaaaa	caaatacctt	cttgtaggca	atctgggtat	5160
ggaattgcaa	gatttggaaa	gtatgggatg	gatcattcga	gggccagccg	tctgcaagaa	5220
gataatacac	catgagaaat	gcaggccttc	aataccagac	aaactcatgg	cattctcgg	5280
gattatgcct	aggggagtta	caccaagagc	ccctacacgg	ttccctgtgt	ccttctgtaa	5340
gataaagcgg	ggttttgaga	ccggctgggc	ctacacacac	cctggagggg	taagtagtgt	5400
gatgcatgtc	accgctgggt	cggatatata	tgtcaatgac	tcaataggga	ggacaaaaat	5460
ccagtgccaa	gacaaaaaca	ctacaacaga	tgagtgtgaa	tatggtgtga	aaacagactc	5520
aggggtgctct	gatggagctc	ggtgctatgt	catcaaccc	gaagcaacca	acatagcagg	5580
gaccaagggg	gccatggtac	acctgaggaa	agctggagga	gagttcaact	gcgtgactgc	5640
ccagggtacc	cccgccttct	ataatctaaa	gaacttaaaa	ggatggtcag	gcctgcctat	5700
ctttgaaagct	gccacaggaa	gagtggtagg	aagggtaaaa	gcaggaaaaa	acactgacaa	5760
tgctccaaca	accattatgt	cagggaagca	agtggcaaaa	ccatcagagt	gtgacctaga	5820
atcagtggtg	aggaaactag	agacaatgaa	cagaggggaa	ttcaaacaaag	tgactctggc	5880
tacagcgcca	ggaaagacaa	ccatgctacc	aaagctgtta	atagaatcca	taggcaggca	5940
taagagagtg	ttagtactga	tcccgttgag	agctgcagcg	gagggggtgt	accagtacat	6000
gagaaccaaa	cacccaagca	tatctttcaa	cttgaggata	ggggatctga	aagaaggtga	6060
catggcaact	gggatcacct	atgcctctta	tgggtacttc	tgccaaatgg	acatgcctag	6120
actggagaat	gcaatgaagg	aataccacta	tattttcttg	gatgaatata	actgtgccc	6180
accgaaacag	ttggcagtga	tgtcaaaaa	acataggttc	ggtgaatcag	ttagggtaat	6240
agccatgacc	gccacgcca	ccgggactgt	gagcacaaaca	gggcagaaat	tcacaattga	6300
ggaggtggta	gtacctgaag	tgatgaaggg	ggaggacctt	gctgatgatt	acatcgaaat	6360
agcaggggtg	aaggtgcca	agaaagagtt	agagggtaac	gtactgactt	ttgtgctac	6420
aaaggagatg	cagcogaaa	atcaaccaca	atcaaccaca	cagggataca	atgctggata	6480
ctacttcagt	ggagaagatc	catcatccct	goggacaact	acttctaagt	caccatata	6540
agtagttgca	accaatgcca	ttgaaatccg	ggtaacctta	ccggaccttg	atacagtaat	6600
agatacaggc	atgaagtgtg	aaaagagact	aagaatcgaa	aacaaagctc	cctacatcgt	6660
aacaggactg	aaaagaatgg	ctatacaaac	gggggagcaa	gctcaaagaa	aaggtaggg	6720
aggcagggtt	aaacctggga	ggtacttgag	aggacctgaa	aacactgcag	gtgaaaagga	6780
ctatcactat	gaccttttac	aggcacagag	gtacggcatc	caagactcaa	taaacatcac	6840
caagtctttc	aggagatga	actatgattg	gcatattat	gaggaagacc	cgtaaagat	6900
tgcccaatta	gagttgctaa	acacactcct	gatctcaagg	gatctgccag	tagtaacaaa	6960
aaatctgatg	cccgcacaaa	cacatcccga	acctatacaa	ttggcttaca	atagtttaga	7020
aacctctgta	ccggtggcat	tcccaaaagt	gaaaaatgga	gaagtcactg	acgcacatga	7080
aacttacgag	ttgatgacct	gtaggaagct	tgagaaagac	ccccctatat	acctgtatgc	7140
aacagaagaa	gaagatctcg	tagtggacat	actgggattg	aaatggccag	acgccacaga	7200
gagggctgtc	ttggaagtgc	aagacgacct	gggcccagatc	acaggtttat	ctgcagggga	7260

gacagcttta	ctcatagccc	tattagggtg	ggtgggctac	gaagccttgg	tgaagaggca	7320
cgtgcctatg	gtgacagaca	tatacacccct	agaagatgaa	aaattggaag	acactacaca	7380
cctacaattt	gccccagatg	atctgaacaa	ttcagatacc	attgagctoc	aagacttatc	7440
gaatcaccaa	atccaacaaa	ttctagaagg	tgggaaggaa	tatgtcggcc	aagcctacca	7500
attcctcagg	ttgcaagctg	agagggctgc	caactcagac	aaaggcaaga	aagcaatggc	7560
agcggcccca	ttactagccc	acaagttcct	ggaatacttg	caagagcatg	caggtgacat	7620
aaagaagtat	ggtctatggg	gggtccacac	agcattgtat	aacagcataa	aagaagact	7680
gggtcacgaa	actgcattcg	catctctggg	tataaaatgg	attgcctttt	cctcagatgg	7740
agtccccggg	atgattaagc	aagcagcagt	agacttgggtg	gtatactata	taatcaacag	7800
gcctgagtat	caaggggata	aggagacaca	gaatgcaggt	agacaatttg	ttggctccct	7860
ttttgtttca	tgtctagcag	agtacacatt	caaaaacttc	aataaatcag	cattagaagg	7920
attgatcgag	cctgccttaa	gctatctacc	ctacgcttca	agcgcactaa	agttattcct	7980
accgactaga	ttgaaagtg	tagtgatact	gtccactact	atatacagaa	catacttatc	8040
aatcaggaaa	ggatctagtc	agggtttagc	cgggctggca	gttagctcag	cgatggagat	8100
catgaaccag	aacccaatca	gcgtggctat	tgcactggca	ctaggagtcc	gagcaatagc	8160
ggcacataat	gccattgaga	gcagtgaggc	aaaaaggact	ctcctgatga	aggtctttgt	8220
taagaacttt	ttggaccaag	cagccactga	tgagcttgtg	aaagagaacc	ctgagaagat	8280
cataatggca	gtgtttgagg	gcattcaaac	agctggaat	ccattgagac	ttgtatacca	8340
tctatatgca	atgtttctaca	aaggggtggac	tgcccgggaa	atagctgaaa	aaaccogtgg	8400
taggaacatt	tttgtgttaa	caatatttga	aggattggaa	atgttaggcc	tggatgccga	8460
ctcaaaatgg	agaaatctga	gctctaatta	tcttattgat	gcagtgaaga	aaatcattga	8520
aaaaatgact	aaaacagcaa	caagcttcac	ctacagcttt	ttgaaatctt	tgcttctctg	8580
ccccttctcg	tgtactaaat	cagaaagaga	tccaagaata	gggtggcccc	aaaaagacta	8640
cgactaccctc	gaggtccgat	gcgcttgtgg	gtataacagg	agagctataa	aaagagactc	8700
aggacctgtg	ttatgggaga	ccttagagga	gacgggtcca	gagtactgcc	acaacagagg	8760
tgaagggggg	ctcagcaatg	tgaagactac	tagatgcttt	gtccaaggag	aggaaatccc	8820
tccaatttga	ctgaggaaa	gagtaggtga	gatgttggtc	aaggggtgtt	cattcagaat	8880
agattttgat	aaagcaaga	tactttcaac	agacaagtgg	aaggtaccac	atagggcagt	8940
tacatcaatc	tttgaggatt	ggcaggggat	tggttacaga	gaggcttacc	tagggacca	9000
accagactat	gggggtctgg	tgcccagatc	ttgtgtaact	gtaacaaaac	aagggttaac	9060
attcttgaaa	actgccagag	gcattggctt	cacgactgac	ctgaccatcc	agaacatcaa	9120
aatgctgata	gctacatgct	tcaagaacaa	ggtgaaggaa	ggggagatac	cagctacgat	9180
tgaaggggaa	acatggatca	acataccact	agtgaatgag	gacaccggga	ccattaaacc	9240
aagcttcggg	gaaagagtga	ttcccgaacc	atatgaggag	gaccactctg	aaggcccaag	9300
tgtaatcggt	gaaacaggag	gcatagccat	caaccaaata	gggtcaatc	cacaatccag	9360
tacatgtgga	acagttttta	cagcagtgaa	ggatctgtgc	caaacagtta	gtaataaagc	9420
caagaatata	aaaattgggt	tttcgggaagg	ccaataccca	ggtccagggg	ttgcaaagaa	9480
gcactgaac	cagctcatac	aagatgaaga	cccaaaacca	ttcatatttg	tttgtggtcc	9540
tgacaagtca	atgtctaato	gggcaaaaac	tgcgaggaac	atcaagagaa	tcaccaccac	9600
aaacactgag	aaattcagag	acttggcaaa	aaacaagaaa	ttgataattg	tgctgttagg	9660
tgatagatac	catgaagata	tagaaaagta	tgcagacttc	aagggcacct	tcttgaccag	9720
acaaaccttg	gaagcactag	caagtgccaa	agctgtagag	aaggacatga	ccaagaaaga	9780
agcagcaaga	gtattggcaa	tggaaagaaa	ggatgaaacta	gaactoccag	ggtggctgca	9840
tacagatgca	cccgaattcc	tagacattac	taaggacaac	atcacacatc	acctaataag	9900
ggacatgcag	agtctgagag	aaagagcagg	ggagatagga	gcaaaggcca	ccactcaaat	9960
cactaagaaa	gggagtgtat	acacaatcaa	tctgagtagc	tggtgggagt	cagagaggtt	10020
ggcatctttg	gaacctttgt	tccgggaact	actatctaaa	tgcaaggccag	tggacagggg	10080
gacatataag	aaattgtcatt	ttgcaacagc	agcccactt	gcccggaggaa	actgggtacc	10140
ggtagcacca	gttgtacatc	ttggggaaat	tccggtaaag	aagaaaaaga	ctctccccta	10200
cgaggcatac	aagctcctaa	aagagatggg	tgactcggag	aaggaaattcc	ataaacccagt	10260
gagcagggaa	aaacaccaat	ggatactgaa	caaagtgaaa	actggtgggtg	acctcggctt	10320
aaaaaatcta	gtatgtccag	gtagggttgg	agaaccaatc	ctaagagaga	agaagaaatt	10380
caacatttac	aacaagagga	ttaccagtac	tatgttatca	gtagggataa	ggccagaaaa	10440
attgccagtg	gtaagagccc	agaccagtac	caaagaattt	catgaagcaa	taagggacaa	10500
aatagacaaa	aaagcaaaaa	cacagacccc	aggcctacac	aaagaattgt	tggagatatt	10560
caactcaata	tgtgccatcc	cgaacttag	aaatacctac	aaagaggttg	attgggacgt	10620
tctaacctca	ggcataaata	ggaaaagggtg	agccgggtac	ttcgaaaaaa	tgaacatag	10680
ggagatcata	gatagtgaca	aaaaatcagt	ggaacaactc	ataaagagaa	tgaatcagg	10740
gctagaattc	aactactatg	agactgcaat	accaaaaaat	gagaagaggg	cagtggtaga	10800
tgattggatg	gaaggtgact	atgtagaaga	aaaaagacca	agagtatac	agtatcctga	10860
ggcaaaagtg	agatttagcta	taaccaaaagt	aatgtataac	tgggtcaagc	agaagcctat	10920
agtaatccct	ggatacgaag	gtaagactcc	tttgtttcat	gttttcgaca	aggtccacaa	10980
agaatggaaa	aatttcaaca	gtccagttgc	agtcagtttt	gacactaaag	cctgggacac	11040


acaagtaaca	cccaaagacc	ttctcctcat	atcagaaatc	caaaagtatt	attacaagaa	11100
agaataccat	agattcatag	ataatttgac	cgagaaaatg	gtggaggtag	cagtggtttg	11160
tgaagacgga	aacgtctaca	taagagaagg	tcagagggga	agtggccaac	cagacactag	11220
cgcaggtaat	agtatgttga	atgtactgac	tatgatatat	gccttctgca	aagctaactc	11280
catcccttac	tcagccttcc	acagggtagc	aaagatacat	gtgtgtggag	atgatggttt	11340
cttgataact	gagaaaagt	ttggtgaggc	ctttgcatc	aaggggcctc	aaattttgat	11400
ggaagcagga	aaaccacaaa	aacttatagg	tgaatttggg	ctgaaattgg	catataaatt	11460
tgatgacatt	gaattttgct	cgcatacacc	aataaaggtc	aggtgggctg	acaacaacac	11520
atcatacatg	ccggaagag	acacagctac	cattctagct	aaaatggcaa	cccgccttga	11580
ctctagtggg	gagaggggga	ccgagggata	cgagctggcc	gtggccttca	gtttcttact	11640
aatgtattct	tggaaccccc	tggtagaag	aatatgcctg	cttgtcatgt	ctacaattga	11700
cacaaaagaa	gctagcctaaa	ataacactat	atatacattt	aggggggatc	ccataggtgc	11760
ctacacagag	gtaattgggt	ataggctgga	ccaactaaaa	cagacagagt	tctctaaatt	11820
ggctcagctg	aatttgtcaa	tggcaatact	tcaaataatac	aataaaaaaca	caaccaagag	11880
actcatcgaa	gattgtgtga	aacttggcaa	ccaaaataag	caaatattgg	tgaatgcaga	11940
ccgtttgatc	agcaagaaaa	cgggctacac	atatgagcca	acagctggcc	acactaagat	12000
aggcaagcac	tatgaagaaa	tcaacctgct	gaaagataca	ccacaaaaaa	ctgtctacca	12060
aggaaactgaa	aggtata					12077

Figure 2

GGSEEGNMFF	RTAPTPPPGC	QEPVYTSTMR	PIFGEPHPL	HKHSTLKLPH	WRGIKTIRVK	60
KRELPKKGDC	SNSTTAPTSG	VYVELGAVFY	KDYTGTVYHR	VPLELCTNQE	RCEGSKCVGR	120
MTGSDGRLYN	VLVCPDDCIL	FERHCRGQTV	VLKWSNPLT	SPLWVQSCSD	DKGAKPKVKP	180
KDDRMKQGKI	VTKPKETead	QKTRPPDATI	VVDGQKYQVR	KKGKAKPKTQ	DGLYHNKNKP	240
EASRKKLEKA	LLAWAILACL	LVVPGVSTNV	TQWNLWDNKS	TTDIHSVMFS	RGIKRSLHGI	300
WPTQICKGIP	THLAADYELK	RIHGMDVdASP	MTNFTCCRLQ	RHEWNKHGWC	NWYNIePWIN	360
LMNNTQGLLN	TGDNFTECAV	TCRYDADLGV	NIVTQARTTP	TILTGCKKGH	NFSFSGEVRA	420
SPCNFELTAE	DLLRIMDHTN	CEGFYFEGEG	IVDGYTEVVE	KARSSGFRAL	TWLSSKIENf	480
KKKIFGAEAS	PYCPVAKRVF	NIIYTNCTP	LGLPDKSKI	GPGTFDISGR	DEFIFPKLPY	540
HVDDFILLSL	IAMSDFAPET	SSIIYLALHY	LMPSNDNRDF	VMDLDPNKLN	LTATKSVASV	600
VPTSVNVLGE	WVCVKPSWwP	YSAEITNLIG	GVITVADLVI	KTIBELLNLW	TEATAVAFLA	660
ALIKIFRQGP	IQAVAWLIII	GGAQAOTCNP	EFMYALAKNT	SIGSLGPESL	TTRWYQLTSG	720
FKLTDSTIEV	TCVGANMRIH	VVCPLVSDRY	LAINHPRALP	TTAWFRKIHT	QHEVPRERIM	780
SESKRRYTCP	CGSKPVVRST	TQFNPIsIST	PSFELECPRG	WTGAVECTLV	SPSTLTtTETI	840
FTYRKPKPFG	LENWCKYTVV	BKGILYSCKF	GGNSTCIKGL	IVKGQREDKV	RYCEWCGYKF	900
SSPNGLPQYP	LGLCEKEQSE	GLRDYGDfPC	CNNGTCIDKE	GSVQCYIGDK	KVTVKLYNAS	960
LLAPMPCKPI	VYNSQGPPAP	KTCTYRWAST	LENKYIEPRD	SYQQYIIKS	GYQYWFDLTA	1020
KDHVADWITK	YFPIIIVALL	GGRGTLWVLI	AYELLTQYEV	VGDENIVAQA	EALVIGNILM	1080
SLDLEIISCF	LLLLLIVVKQ	AVRRTLALLF	HWITMNPfQS	VMITVVYFVG	LVRAEEGTKE	1140
GSTSGPPIHV	VAILLFLLYH	TVKYKDFNIA	MILLITLSLK	SSSYHTSly	EIPLLVAVIS	1200
LTCsIYIFDL	QVKSKLIVAPT	IGIIGVTLAM	RVLWLVRQMT	IPTPSVSISL	IDPKMVILLY	1260
LISLTITVNH	NLDLASYCLK	LGPFILSFLT	MWVDVVILLL	MLPWYELVKV	YYLKKKKEDV	1320
ETWFQNSGIS	TQETSPYGFd	FSSPGEgVHT	LPMQNKTfKC	RTAYMTVLRA	LVTTAISSVW	1380
KPIILAELLI	EAVYWTHIKI	AKELAGSSRF	VARFIASIE	LNWAMDEKEA	SRYKRFYLLS	1440
SKITDLMVKH	KIQNETVKSW	FEETEIfGfIQ	KVAMVIRAHs	LSLEPNAILC	SVCEBKQNOk	1500
AKRPPCKCGS	RGTQIKCGLT	LAEFEEHYK	KIYILEGQDE	TPMRKEERQQ	VTYVSRGALF	1560
LRNLPIlASK	NKYLLVGNLg	MELQDLESgM	WIIRGPAVCK	KIHHEKCRP	SIPDKLMAFF	1620
GIMPRGVTPR	APTRFFVSLl	KIRRGfETGw	AYTHPGGVSS	VMHVtAGSDI	YVNDsIGRTK	1680
IQCDKNtTfT	DECBYGVKTD	SGCSDGARCy	VINPEATNIA	GTKGAMVHLR	KAGGEFNCVT	1740
AQTFPAFYNL	KNLKGWSGLP	IFEAATGRVv	GRVKAGKNTD	NAPTTIMSGT	QVAKPSECdL	1800
ESVVRKLETM	NRGEFKQVTL	ATGAGKtTML	PKLLIESIGR	HKRVLVLIPL	RAAABGVYQY	1860
MRTKHPSISF	NLRIGDLKEG	DMATGITYAS	YGYFCQMDMP	RLENAMKEYH	YIFLDEYHCA	1920
TPEQLAVMSK	IHRFGESVRV	IAMTATPSGT	VSTTGQKfTI	BEVVVPEVMK	GEDLADDYIE	1980
IAGLKVPKKE	LEGNVLTfVP	TRKMASETAK	KLTTQGYNAG	YYFSGEDPSS	LRTTTSKSPY	2040
IVVATNAIES	GVTLPDLDTV	IDTGMKCEKR	LRIENKAPYI	VTGLKRMAIT	TGEQAQRKGR	2100
VGRVKPGRYL	RGPENTAGEK	DYHYDLLQAQ	RYGIQDSINI	TKSFREMNyD	WALYEEDPLK	2160
IAQLELLNtL	LISRDLPVVT	KNLMARTTHP	EPIQLAYNSL	ETPVPVAFPK	VKNGEVtDAH	2220
ETYBLMTCRK	LEGDPIYLY	ATEEEDLVVD	ILGLKWPdAT	ERAVLEVQDA	LGQITGLSAG	2280
ETALLIALLG	WVGYEALVKR	HVPMVTDIYT	LEDEKLEDTT	HLQFAPDDLN	NSDTTELQDL	2340
SNHQIQQILE	GGKEYVGQAY	QFLRLQAERA	ANSDKGKKAM	AAAPLLAHKF	LEYLQEBHAGD	2400
IKKYGLWGVH	TALYNSIKER	LGHETAFASL	VIKWIAFSSD	GVPGMIKQAA	VDLVVYYIIN	2460
RPEYQGDKET	QNAGRQFVGS	LfVSClAEYt	FKNFNKSALE	GLIEPALSyL	PyASSALKLf	2520
LPTRLESVVI	LSTTYRtTYL	SIRKGSsQCL	AGLAVSSAME	IMNQNPISVA	IAlALGVGAI	2580
AAHNAIESSE	AKRTLLMKVF	VKNFLDQAAT	DELVKENPEK	IIMAVFEGIQ	TAGNPLRLVY	2640
HLYAMFYKGW	TAABTAEKTA	GRNIFVLTIF	EGLEMLGLDA	DSKWRNLSSN	YLIDAVKKII	2700
EKMtKTATSF	TYSFLKSLLP	APfSCTKSER	DPRIGWPQKD	YDYLEVRCAC	GYNRRAIKRD	2760
SGPVLWETLE	ETGPEYCHNR	GERGLSNVKT	TRCFVQGEBI	PPIALRKGVG	EMLVKGVsFR	2820
IDFDKDKILS	TDKWKVPHRA	VTSIFEDWQG	IGYREAYLGT	KPDYGGLVPR	SCVTVTKQGL	2880
TFLKTARGMA	FTTDLTIQNI	KMLIATCFKN	KVKEGEIPAT	IEGETWINIP	LVNEDTGTIK	2940
PSFGERVlPE	PyEBDPLEGP	SVIVETGGIA	INQIGVNPQS	STCGTVFTAV	KDLcQTvsNk	3000
AKNIKIGFSE	GQYPGPgVAK	KTLNQLIQDE	DPKPFIFVCG	SDKsMSNRAK	TARNIKRITf	3060
TTPEKFRDLA	KNKKLlIVLL	GDRYHEDIeK	YADfKGtFLT	RQTLBALASA	KAVEKDMTKK	3120
EAARVLAMEE	KDELELPGWL	HTDAPKFLDI	TKDNITHHLI	GDMQSLRERA	GEIGAKATTQ	3180
ITKKGsvYTI	NLSTWwESER	LASLEPLFRE	LLSKCRPVDR	ETyKNCHfAT	AAQLAGGNwV	3240
PVAPVHLGE	IPVKKKKTLP	YEAYKLLKEM	VDSEKEFHkP	VSREKHQWIL	NKVKTGGDLG	3300
LKNLVCpGRV	GEPILREKKK	FNIYNKRITS	TMSVSGIRPE	KLPVVRAQTS	TKEFHBAIRD	3360
KIDKANTQT	PGLHKELLEI	FNSICALPEL	RNTYKEVDWD	VLTSGINRKG	AAGYfEKMNl	3420
GEIIDSdKKS	VEQLIKRMKS	GLEFNYYETA	IPKNEKRAVV	DDWMEGDYVE	EKRPRVlOYP	3480

EAKMRLAITK	VMYNWVKQKP	IVIPGYEGKT	PLFHVFDKVV	KWKNFNSPV	AVSFDTKAWD	3540
TQVTPKDLLL	ISEIQKYYYK	KEYHRFIDNL	TEKMVEVPVV	CEDGNVYIRE	GQRGSGQPD	3600
SAGNSMLNVL	TMIYAFCKAN	SIPYSAFHRV	AKIHVCGDDG	FLITEKSFGE	AFAIKGPQIL	3660
MEAGKPQKLI	GEFGLKLAYK	FDDIEFCSHT	PIKVRWADNN	TSYMPGRDTA	TILAKMATRL	3720
DSSGERGTEG	YELAVAFSFL	LMYSWNPLVR	RICLLVMSTI	DTKEASQNTT	IYTFRGDPIG	3780
AYTEVIGYRL	DQLKQTEFSK	LAQLNLSMAI	LQIYNKNTTK	RLIEDCVKLG	NQNKQILVNA	3840
DRLISKKTGY	TYEPTAGHTK	IGKHYYEINL	LKDTPQKTVY	QGTERY		3886

Figure 3



Keylist :

- PCR Product
- NADL Reference Sequence gi 8625549
-{...}F or R Primers: [...] Indicates bp locations based on NADL sequence
R Indicates reverse primer
F Indicates forward primer

Figure 4




Figure 5A





Figure 5B

41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60



61 62 63 64 65 66 67 68 69 70 71 72 73 74 75 76 77 78 79 80



81 82 83 84 85 86 87 88 89 90 91 92 93 94 95 96 97 98 99




Figure 6A




Figure 6B

41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60 61 62 63 64 65 66 67 68 69 70 71 72 73 74 75 76 77 78 79 80



81 82 83 84 85 86 87



Figure 7A




Figure 7B




Figure 8A

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20



21 22 23 24 25 26 27 28 29 30 31 32 33




Figure 8B

34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53




Figure 9





Figure 10



PESTIVIRUS SPECIES

FIELD OF THE INVENTION

[0001] The present invention relates to a novel pestivirus, and gene sequences derived from the same. The invention further relates to detection methods, vaccines, therapeutics, and diagnostic methods using the sequences of the present invention.

BACKGROUND ART

[0002] Pestiviruses cause highly contagious and often fatal diseases of pigs, cattle and sheep, which are characterised by damage to the respiratory and gastrointestinal tracts and immune system and can run an acute or chronic course. Infection of the reproductive system may cause embryonic and foetal death, congenital defects and the birth of persistently infected animals. Outbreaks of the diseases associated with pestivirus infections occur in many countries and can cause large economic losses.

[0003] The Pestivirus genus of the Flaviviridae comprises three structurally, antigenically and genetically closely related member species: Classical swine fever (CSF) or hog cholera (Francki et al. 1991. *Flaviviridae*, In the Fifth report of the International Committee on Taxonomy of Viruses, *Archiv. Virol. Suppl.* 2, Springer Verlag, Vienna p. 223-233); Bovine viral diarrhoea virus (BVDV) which mainly affects cattle, and Border disease virus (BDV) which mainly affects sheep (Moennig and Plagemann (1992) *Adv. Virus Res.* 41: 53-98; Moormann et al., (1990) *Virology* 177: 184-198; Becher et al. (1994) *Virology* 198: 542-551). Recent studies indicate that there may be several less well recognised viruses that warrant separate taxonomic classification, perhaps as separate species (Avalos-Ramirez et al (2001) *Virology* 286: 456-465)

[0004] The genomes of pestiviruses consist of a positive strand RNA molecule of about 12.5 kb (Renard et al. (1985) *DNA* 4: 429-438; Moormann and Hulst (1988) *Virus Res.* 11: 281-291; Becher et al. (1994) *Virology* 198: 542-551). However, the positive strand RNA genomes of several cytopathogenic BVDV strains may be considerably larger (Meyers et al. (1991) *Virology* 180: 602-616; Meyers et al. (1992) *Virology* 191: 368-386; Qi et al. (1992) *Virology* 189: 285-292).

[0005] An inherent property of viruses with a positive strand RNA genome is that their genomic RNA is infectious, i.e. after transfection of this RNA in cells that support viral replication, infectious virus is produced. As expected, the genomic (viral) RNA of pestiviruses is also infectious (Moennig and Plagemann, (1992) *Adv. Virus Res.* 41: 53-98).

[0006] In 2003 an outbreak of stillbirths and pre-weaning deaths of piglets occurred on two farms in New South Wales, Australia (McOrist et al, (2004) *Aust Vet J.* 82: 509-511). Key features of the clinical presentation and pathology findings suggested that this disease outbreak was novel and probably due to a virus. Extensive testing for known viruses and some bacteria failed to identify an aetiological agent. To avoid confusion with other important diseases in pigs, the term "porcine myocarditis syndrome" (abbreviated as "PMC") was ascribed to the disease, and the term "PMC virus" given to presumptive agent. Subsequently, the causative agent was identified as a novel pestivirus. The name Bungowanah is proposed for this new virus.

[0007] The present invention addresses a need in the art for methods of detecting and/or treating infections caused by the novel PMC virus.

SUMMARY OF THE INVENTION

[0008] The invention provides an isolated RNA nucleotide sequence corresponding to the PMC virus nucleotide sequence depicted in SEQ ID NO:1, or sequences substantially homologous to SEQ ID NO:1, or fragments thereof.

[0009] The invention also provides the isolated DNA nucleotide sequence of the PMC virus of SEQ ID NO:1, or sequences substantially homologous to SEQ ID NO:1, or fragments thereof.

[0010] The invention further provides polypeptides encoded by the above RNA and DNA nucleotide sequences and fragments thereof, and/or an isolated PMC virus amino acid sequence as shown in SEQ ID NO: 2 and fragments thereof.

[0011] In another aspect, the invention provides methods for detecting the presence of a PMC virus amino acid sequence in a sample, comprising the steps of:

[0012] a) contacting a sample suspected of containing a PMC virus amino acid sequence with an antibody that specifically binds to the PMC virus amino acid sequence under conditions which allow for the formation of reaction complexes comprising the antibody and the PMC virus amino acid sequence; and

[0013] b) detecting the formation of reaction complexes comprising the antibody and PMC virus amino acid sequence in the sample, wherein detection of the formation of reaction complexes indicates the presence of PMC virus amino acid sequence in the sample.

[0014] The invention also provides methods for detecting the presence of a PMC virus antibody in a sample, comprising the steps of:

[0015] a) contacting a sample suspected of containing a PMC virus antibody with an amino acid sequence under conditions which allow for the formation of reaction complexes comprising the PMC virus antibody and the amino acid sequence; and

[0016] b) detecting the formation of reaction complexes comprising the antibody and amino acid sequence in the sample, wherein detection of the formation of reaction complexes indicates the presence of PMC virus antibody in the sample.

[0017] Additionally, the invention provides an in vitro method for evaluating the level of PMC virus antibodies in a biological sample comprising the steps of:

[0018] a) detecting the formation of reaction complexes in a biological sample according to the method noted above; and

[0019] b) evaluating the amount of reaction complexes formed, which amount of reaction complexes corresponds to the level of PMC virus antibodies in the biological sample.

[0020] The invention also provides an in vitro method for evaluating the level of PMC virus polypeptides in a biological sample comprising the steps of:

[0021] a) detecting the formation of reaction complexes in a biological sample according to the method noted above; and

- [0022] b) evaluating the amount of reaction complexes formed, which amount of reaction complexes corresponds to the level of PMC virus polypeptide in the biological sample.
- [0023] The present invention further provides methods for detecting the presence or absence of PMC virus in a biological sample, which comprise the steps of:
- [0024] a) bringing the biological sample into contact with a polynucleotide probe or primer comprising a PMC virus polynucleotide of the invention under suitable hybridising conditions; and
- [0025] b) detecting any duplex formed between the probe or primer and nucleic acid in the sample.
- [0026] The present invention also relates to a method for the detection of PMC virus nucleic acids present in a biological sample, comprising:
- [0027] a) amplifying the nucleic acid with at least one primer as defined above,
- [0028] b) detecting the amplified nucleic acids.
- [0029] The present invention also relates to a method for the detection of PMC virus nucleic acids present in a biological sample, comprising:
- [0030] a) hybridizing the nucleic acids of the biological sample at appropriate conditions with one or more probes as defined above,
- [0031] b) washing under appropriate conditions, and
- [0032] c) detecting the hybrids formed.
- [0033] In a further aspect, the present invention provides a method for the generation of antibodies comprising the steps of:
- [0034] a) providing a PMC virus polypeptide sequence to a subject; and
- [0035] b) collecting the antibodies generated in the subject against the polypeptide.
- [0036] In another aspect of the invention, there is provided a vaccine composition comprising a PMC virus polypeptide or fragment thereof. The invention also provides a vaccine composition comprising a PMC virus nucleotide or fragment thereof that encodes for a PMC virus polypeptide.
- [0037] Pharmaceutical compositions comprising a PMC virus polypeptide that enhances the immunocompetence of the host individual and elicits specific immunity against the PMC virus are further provided by the invention.
- [0038] The present invention also provides therapeutic compositions comprising polynucleotide sequences and/or antibodies prepared against the polypeptides of the invention. The present invention further provides therapeutic compositions comprising PMC virus nucleic acid sequences as well as antisense and ribozyme polynucleotide sequences hybridisable to a polynucleotide sequence encoding a PMC virus amino acid sequence according to the invention.
- [0039] The present invention provides for the use of PMC virus amino acid sequences and/or antibodies according to the invention, for manufacture of a medicament for modulation of a disease associated with PMC virus. The present invention additionally provides for the use of polynucleotide sequences of the invention, as well as antisense and ribozyme polynucleotide sequences hybridisable to a polynucleotide sequence encoding a PMC virus amino acid sequence according to the invention, for manufacture of a medicament for modulation of a disease associated with PMC virus.
- [0040] The present invention further provides a method of inducing a protective immune response in an animal or human against PMC virus comprising the steps of:
- [0041] a) administering to said animal or human an effective amount of a composition of the invention.
- [0042] The present invention also provides methods for enhancing an animal's immunocompetence and the activity of its immune effector cells against a PMC virus comprising the step of:
- [0043] a) administering a composition comprising a therapeutically effective amount of a PMC virus peptide or polypeptide.
- [0044] In addition, the present invention provides a live vector comprising the PMC virus and a heterologous polynucleotide.
- [0045] In another aspect of the invention, there is provided a method of screening for drugs comprising the steps of:
- [0046] a) contacting an agent with a PMC virus amino acid sequence or fragment thereof and
- [0047] b) assaying for the presence of a complex between the agent and the PMC virus amino acid sequence or fragment.
- [0048] The present invention also provides a method of screening for ligands of the proteins of the PMC virus comprising the steps of:
- [0049] a) contacting a ligand with a PMC virus amino acid sequence or fragment thereof and
- [0050] b) assaying for the presence of a complex between the PMC virus amino acid sequence or fragment and a ligand.
- [0051] In a further aspect of the invention, a test kit may be prepared for the demonstration of the presence of PMC virus comprising:
- [0052] (a) a predetermined amount of at least one labelled immunochemically reactive component obtained by the direct or indirect attachment of the present PMC virus amino acid sequence or a specific binding partner thereto, to a detectable label;
- [0053] (b) other reagents; and
- [0054] (c) directions for use of said kit.
- [0055] Additionally, the invention provides a test kit for the demonstration of the presence of PMC virus comprising:
- [0056] (a) a predetermined amount of at least one labelled antibody to the PMC virus;
- [0057] (b) other reagents; and
- [0058] (c) directions for use of said kit.
- [0059] The invention also provides a test kit for the demonstration of the presence of PMC virus comprising:
- [0060] (a) a predetermined amount of at least one labelled polypeptide derived from the PMC virus;
- [0061] (b) other reagents; and
- [0062] (c) directions for use of said kit.
- [0063] Additionally the present invention provides a test kit prepared for the demonstration of the presence of PMC virus comprising:
- [0064] (a) a predetermined amount of at least one labelled nucleic acid sequence derived from the PMC virus;
- [0065] (b) other reagents; and
- [0066] (c) directions for use of said kit.
- [0067] The present invention also provides a recombinant expression vector comprising a PMC virus nucleic acid sequence or a part thereof as defined above, operably linked to prokaryotic, eukaryotic or viral transcription and translation control elements.
- [0068] The invention further relates to the hosts (prokaryotic or eukaryotic cells) which are transformed by the above

mentioned vectors and recombinants and which are capable of expressing said RNA and/or DNA fragments.

[0069] The present invention also relates to a method for the production of a recombinant PMC virus polypeptide, comprising the steps of:

[0070] a) transforming an appropriate cellular host with a recombinant vector, in which a PMC virus polynucleotide sequence or a part thereof has been inserted under the control of appropriate regulatory elements,

[0071] b) culturing said transformed cellular host under conditions enabling the expression of said insert, and,

[0072] c) harvesting said polypeptide.

[0073] According to another embodiment the present invention provides methods for preparing a PMC virus amino acid sequence, comprising the steps of:

[0074] (a) culturing a cell containing a vector as described above under conditions that provide for expression of the PMC virus amino acid sequence; and

[0075] (b) recovering the expressed PMC virus sequence.

BRIEF DESCRIPTION OF THE DRAWINGS

[0076] FIG. 1 shows the DNA sequence of the PMC virus of the present invention;

[0077] FIG. 2 shows the protein sequence of the PMC virus of the present invention;

[0078] FIG. 3 shows a map of the location of primers used to sequence the whole virus, the dotted lines underneath are the length of the PCR products produced and sequenced;

[0079] FIG. 4 shows an ethidium bromide stained 0.8% gel of SISPA applied to DNA and RNA of adaptor PCR (run on Corbett and Eppendorf cyclers). Arrows indicate where gel was cut to collect bands for purification and cloning (e.g. ER1=Eppendorf PCR machine, RNA preparation, gel position 1). Lane 1 Eppendorf machine RNA SISPA 10 ul of PCR product; Lane 2 Eppendorf machine DNA SISPA 10 ul of PCR product; Lane 3 Eppendorf machine RNA SISPA 40 ul of PCR product; Lane 4 Eppendorf machine DNA SISPA 40 ul of PCR product; Lane 5 Eppendorf machine Blank 40 ul of PCR control; Lane 6 Corbett machine RNA SISPA 40 ul of PCR product; Lane 7 Corbett machine DNA SISPA 40 ul of PCR product; Lane 8 Corbett machine blank 40 ul of PCR product; Lane 9 100 bp marker.

[0080] FIG. 5 shows an ethidium bromide stained 1% gel of SISPA applied to DNA and RNA simultaneously to screen colonies for inserts (e.g. ER3 1=Eppendorf PCR machine, RNA sample position 3 colony 1). Lane 1 ER3 1; Lane 2 ER3 2; Lane 3 ER3 3; Lane 4 ER3 4; Lane 5 ER3 5; Lane 6 ER3 6; Lane 7 ER3 7; Lane 8 ER3 8; Lane 9 ER3 9; Lane 10 ER3 10; Lane 11 ER3 11; Lane 12 ER3 12; Lane 13 Marker 100 bp; Lane 14 ER4 1; Lane 15 ER4 2; Lane 16 ER4 3; Lane 17 ER4 4; Lane 18 ER4 5; Lane 19 ER4 6; Lane 20 ER4 7; Lane 21 ER4 8; Lane 22 ER4 9; Lane 23 ER4 10; Lane 24 ER4 11; Lane 25 ER4 12; Lane 26 ER5 1; Lane 27 ER5 2; Lane 28 ER5 3; Lane 29 ER5 4; Lane 30 ER5 5; Lane 31 ER5 6; Lane 32 ER5 7; Lane 33 Marker 100 bp; Lane 34 ER5 8; Lane 35 ER5 9; Lane 36 ER5 10; Lane 37 ER5 11; Lane 38 ER5 12; Lane 39 ER6 1; Lane 40 ER6 2; Lane 41 ER6 3; Lane 42 ER6 4; Lane 43 ER6 5; Lane 44 ER6 6; Lane 45 ER6 7; Lane 46 ER6 8; Lane 47 ER6 9; Lane 48 ER6 10; Lane 49 ER6 11; Lane 50 ER6 12; Lane 51 ER7 1; Lane 52 ER7 2; Lane 53 Marker 100 bp; Lane 54 ER7 3; Lane 55 ER7 4; Lane 56 ER7 5; Lane 57 ER7 6; Lane 58 ER7 7; Lane 59 ER7 8; Lane 60 ER7 10; Lane 61 ER7 11; Lane 62 ER7 12; Lane 63 ER8 1;

Lane 64 ER8 2; Lane 65 ER8 3; Lane 66 ER8 4; Lane 67 ER8 5; Lane 68 ER8 6; Lane 69 ER8 7; Lane 70 ER8 8; Lane 71 ER8 9; Lane 72 ER8 10; Lane 73 Marker 100 bp; Lane 74 ER8 11; Lane 75 ER8 12; Lane 76 ER9 1; Lane 77 ER9 2; Lane 78 ER9 3; Lane 79 ER9 4; Lane 80 ER9 5; Lane 81 Marker 100 bp; Lane 82 ER9 6; Lane 83 ER9 7; Lane 84 ER9 8; Lane 85 ER9 9; Lane 86 ER9 10; Lane 87 ER9 11; Lane 88; Lane 89 ER10 2; Lane 90 ER10 3; Lane 91 ER10 4; Lane 92 ER10 5; Lane 93 ER10 6; Lane 94 ER10 7; Lane 95 ER10 8; Lane 96 ER10 9; Lane 97 ER10 10; Lane 98 ER10 11; Lane 99 ER10 12.

[0081] FIG. 6 shows an ethidium bromide stained 1% gel of PCR carried out to screen of colonies for DNA (Eppendorf cycler). Lane 1 ED2 1=Eppendorf machine, DNA gel cut out 2, colony 1; Lane 2 ED2 2; Lane 3 ED2 3; Lane 4 ED2 4; Lane 5 ED2 5; Lane 6 ED2 6; Lane 7 ED2 7; Lane 8 ED2 8; Lane 9 ED2 9; Lane 10 ED2 10; Lane 11 ED2 11; Lane 12 ED2 12; Lane 13 Marker 100 bp; Lane 14 ED3 1; Lane 15 ED3 2; Lane 16 ED3 3; Lane 17 ED3 4; Lane 18 ED3 5; Lane 19 ED3 6; Lane 20 ED3 7; Lane 21 ED3 8; Lane 22 ED3 9; Lane 23 ED3 10; Lane 24 ED3 11; Lane 25 ED3 12; Lane 26 ED4 1; Lane 27 ED4 2; Lane 28 ED4 3; Lane 29 ED4 4; Lane 30 ED4 5; Lane 31 ED4 6; Lane 32 ED4 7; Lane 33 Marker 100 bp; Lane 34 ED4 8; Lane 35 ED4 9; Lane 36 ED4 10; Lane 37 ED4 11; Lane 38 ED4 12; Lane 39 ED5 1; Lane 40 ED5 2; Lane 41 ED5 3; Lane 42 ED5 4; Lane 43 ED5 5; Lane 44 ED5 6; Lane 45 ED5 7; Lane 46 ED5 8; Lane 47 ED5 9; Lane 48 ED5 10; Lane 49 ED5 11; Lane 50 ED5 12; Lane 51 ED6 1; Lane 52 ED6 2; Lane 53 ED6 3; Lane 54 ED6 4; Lane 55 ED6 5; Lane 56 ED6 6; Lane 57 ED6 7; Lane 58 ED6 8; Lane 59 ED6 9; Lane 60 Marker 100 bp; Lane 61 ED6 10; Lane 62 ED6 11; Lane 63 ED6 12; Lane 64 ED7 1; Lane 65 ED7 2; Lane 66 ED7 3; Lane 67 ED7 4; Lane 68 ED7 5; Lane 69 ED7 6; Lane 70 ED7 7; Lane 71 ED7 8; Lane 72 ED7 9; Lane 73 ED7 10; Lane 74 ED7 11; Lane 75 ED7 12; Lane 76 ED8 1; Lane 77 ED8 2; Lane 78 ED8 3; Lane 79 ED8 4; Lane 80 ED8 5; Lane 81 ED8 6; Lane 82 ED8 7; Lane 83 ED8 8; Lane 84 ED8 9; Lane 85 ED8 10; Lane 86 ED8 11; Lane 87 ED8 12.

[0082] FIG. 7 shows an ethidium bromide stained 1% gel of PCR carried out to screen of colonies for RNA inserts (Corbett cycler). Lane 1 CR2 1=Corbett machine, RNA gel position 2, colony 1; Lane 2 CR2 2; Lane 3 CR2 3; Lane 4 CR2 4; Lane 5 CR2 5; Lane 6 CR2 6; Lane 7 Marker 100 bp; Lane 8 Marker 100 bp; Lane 9 CR2 7; Lane 10 CR2 8; Lane 11 CR2 9; Lane 12 CR2 10; Lane 13 CR2 11; Lane 14 CR2 12; Lane 15 CR3 1; Lane 16 CR3 2; Lane 17 CR3 3; Lane 18 CR3 4; Lane 19 CR3 5; Lane 20 CR3 6; Lane 21 CR3 7; Lane 22 CR3 8; Lane 23 CR3 9; Lane 24 CR3 10; Lane 25 CR3 11; Lane 26 CR3 12; Lane 27 Marker 100 bp; Lane 28 Marker 100 bp; Lane 29 CR4 1; Lane 30 CR4 2; Lane 31 CR4 3; Lane 32 CR4 4; Lane 33 CR4 5; Lane 34 CR4 6; Lane 35 CR4 7; Lane 36 CR4 8; Lane 37; CR4 9; Lane 38 CR4 10; Lane 39 CR4 11; Lane 40 CR4 12; Lane 41 marker 100 bp; Lane 42 marker 100 bp; Lane 43 CR5 1; Lane 44 CR5 2; Lane 45 CR5 3; Lane 46 CR5 4; Lane 47 CR5 5; Lane 48 CR5 6; Lane 49 CR5 7; Lane 50 CR5 8; Lane 51 CR5 9; Lane 52 CR5 10; Lane 53 CR5 11; Lane 54 PCR Blank control; Lane 55 marker 100 bp.

[0083] FIG. 8 shows an ethidium bromide stained 1% gel of PCR carried out to screen colonies for DNA (Corbett cycler). Lane 1 marker 100 bp; Lane 2 CD3 1=Corbett machine, DNA gel cut out 3, colony 1; Lane 3 CD3 2; Lane 4 CD3 3; Lane 5 CD3 4; Lane 6 CD3 5; Lane 7 CD3 6; Lane 8 CD3 7; Lane 9 CD3 8; Lane 10 CD3 9; Lane 11 CD3 10; Lane 12 CD3 11; Lane 13 CD3 12; Lane 14 CD4 1; Lane 15 CD4 2; Lane 16

CD4 3; Lane 17 CD4 4; Lane 18 CD4 5; Lane 19 CD4 6; Lane 20 marker 100 bp; Lane 21 marker 100 bp; Lane 22 CD4 7; Lane 23 CD4 8; Lane 24 CD4 9; Lane 25 CD4 10; Lane 26 CD4 11; Lane 27 CD4 12; Lane 28 CD5 1; Lane 29 CD5 2; Lane 30 CD5 3; Lane 31 CD5 4; Lane 32 CD5 5; Lane 33 CD5 6; Lane 34 CD5 7; Lane 35 CD5 8; Lane 36 CD5 9; Lane 37 CD5 10; Lane 38 CD5 11; Lane 39 CD5 12; Lane 40 marker 100 bp; Lane 41 marker 100 bp; Lane 42 CD6 1; Lane 43 CD6 2; Lane 44 CD6 3; Lane 45 CD6 4; Lane 46 CD6 5; Lane 47 CD6 6; Lane 48 CD6 7; Lane 49 CD6 8; Lane 50 CD6 9; Lane 51 CD6 10; Lane 52 CD6 11; Lane 53 CD6 12.

[0084] FIG. 9 shows an ethidium bromide stained 1.5% gel of PCR carried out to confirm authenticity of viral sequence for virus confirmation by nRT-PCR. PCR results confirmed the presence of Pestivirus in clinical specimens (lanes 3, 8 and 23) while EMCV was not present (lane 28) (lanes marked + are PCR positive). Lane 1 Marker 100 bp; Lane 2 Blank CR3 9 primers; Lane 3 SISPA sera CR39 primers; Lane 4 NADL +ve control CR39 primers; Lane 5 EMCV -ve control CR39 primers; Lane 6; Lane 7 Blank ER510 primers; Lane 8 SISPA sera ER510 primers; Lane 9 NADL +ve control ER510 primers; Lane 10 EMCV -ve control ER510 primers; Lane 11; Lane 12 Blank ER55 primers; Lane 13 SISPA sera ER55 primers; Lane 14 NADL +ve control ER55 primers; Lane 15 EMCV -ve control ER55 primers; Lane 16; Lane 17; Lane 18; Lane 19; Lane 20 Marker 100 bp; Lane 21 Marker 100 bp; Lane 22 Blank ER62 primers; Lane 23 SISPA sera ER62 primers; Lane 24 NADL +ve control ER62 primers; Lane 25 EMCV -ve control ER62 primers; Lane 26; Lane 27 Blank ER41 primers; Lane 28 SISPA sera ER41 primers; Lane 29 NADL +ve control ER41 primers; Lane 30 EMCV -ve control ER41 primers; Lane 31; Lane 32; Lane 33; Lane 34; Lane 35; Lane 36; Lane 37; Lane 38; Lane 39; Lane 40 marker 100 bp.

[0085] FIG. 10 shows a hydrophobicity plot of the PMC virus protein sequence.

DETAILED DESCRIPTION OF THE INVENTION

New Pestivirus

[0086] In accordance with this invention, a new pestivirus has been discovered that differs genetically from known pestiviruses. The new virus is characterised by the RNA sequence corresponding to that shown in SEQ ID NO: 1. The sequence has been deposited as Genbank reference EF100713.

[0087] The new virus is hereinafter generally referred to as PMC virus and the condition caused by infection with the PMC virus is PMC.

[0088] The PMC virus genome comprises a single open reading frame (ORF), encoding a number of genes. The genes encoded by the ORF of PMC correspond to those of other pestiviruses, being the Npro, capsid, E0, E1, E2, P7, NS2, NS3, NS4A, NS4B, NS5A and NS5B genes.

[0089] The PMC virus is approximately 40% similar to other pestiviruses on a nucleic acid sequence level. At the protein level, PMC virus has 46-71% identity and 63-83% similarity with other pestiviruses. A comparative analysis of both the nucleic acid and deduced amino acid sequences would suggest that PMC virus is sufficiently unique to warrant consideration for classification as a new species within the pestivirus genus.

Open Reading Frames, Encoded Genes, Features of Genome

[0090] The nucleotide sequence of SEQ ID NO:1 encodes a single ORF encoding a number of different genes. The

genes encoded by SEQ ID NO:1 correspond to the Npro, capsid, E0, E1, E2, P7, NS2, NS3, NS4A, NS4B, NS5A and NS5B genes of other pestiviruses.

[0091] The approximate location of the genes of PMC, based on sequence comparison with gi12657941, is indicated in Table 1.

TABLE 1

<u>Location of proteins within PMC nucleic acid open-reading frame.</u>	
PROTEIN	APPROXIMATE DNA POSITION
NPro	419-922
Capsid	923-1219
E0	1220-1885
E1	1886-2473
E2	2474-3604
P7	3605-3820
NS2	3821-5224
NS3	5225-7252
NS4A	7253-7441
NS4B	7442-8482
NS5A	8483-9997
NS5B	9998-12077

TABLE 2

<u>Location of proteins within PMC protein open-reading frame.</u>	
PROTEIN	APPROXIMATE AMINO ACID POSITION
NPro	1-167
Capsid	168-267
E0	268-489
E1	490-685
E2	686-1062
P7	1063-1134
NS2	1135-1602
NS3	1603-2278
NS4A	2279-2341
NS4B	2342-2688
NS5A	2689-3193
NS5B	3194-3886

Nucleic Acid Sequences

RNA

[0092] The invention provides an isolated RNA nucleotide sequence corresponding to the PMC virus nucleotide sequence depicted in SEQ ID NO:1, or sequences substantially homologous to SEQ ID NO:1, or fragments thereof. The invention further provides an RNA sequence comprising the complement of the PMC virus RNA genome, or fragments thereof.

[0093] The RNA sequence may also correspond to a fragment of SEQ ID NO:1. Preferably, the fragment is selected from the following locations of SEQ ID NO:1: position 419-922, 923-1219, 1220-1885, 1886-2473, 2474-3604, 3605-3820, 3821-5224, 5225-7252, 7253-7441, 7442-8482, 8483-9997, 9998-12077. Alternatively, the fragment may be selected from any one of SEQ ID NOs:3-15.

[0094] Substantial homology or identity exists when a PMC virus polynucleotide sequence or fragment thereof will hybridise to another PMC virus polynucleotide (or a complementary strand thereof) under selective hybridisation conditions.

[0095] Selective hybridisation may be under low, moderate or high stringency conditions, but is preferably under high stringency.

[0096] Typically, selective hybridisation will occur when there is at least about 55% identity over a stretch of at least about 14 nucleotides, preferably at least about 65%, more preferably at least about 75% and most preferably at least about 90%. The length of homology comparison, as described, may be over longer stretches and in certain embodiments will often be over a stretch of at least about nine nucleotides, usually at least about 20 nucleotides, more usually at least about 24 nucleotides, typically at least about 28 nucleotides, more typically at least about 32 nucleotides and preferably at least about 36 or more nucleotides.

[0097] Thus, the polynucleotide sequences of the invention preferably have at least 75%, more preferably at least 85%, more preferably at least 90% homology to the sequences shown in the sequence listings herein. More preferably there is at least 95%, more preferably at least 98%, homology. Nucleotide homology comparisons may be conducted as described below for polypeptides. A preferred sequence comparison program is the GCG Wisconsin Bestfit program.

[0098] In the context of the present invention, a homologous sequence is taken to include a nucleotide sequence which is at least 60, 70, 80 or 90% identical, preferably at least 95 or 98% identical at the nucleic acid level over at least 20, 50, 100, 200, 300, 500 or 819 nucleotides with the corresponding nucleotide sequences set out in SEQ ID NO:1. In particular, homology should typically be considered with respect to those regions of the sequence that encode contiguous amino acid sequences known to be essential for the function of one or more of PMC virus proteins, rather than non-essential neighbouring sequences.

[0099] PMC virus polynucleotide sequence fragments of the invention will preferably be at least 15 nucleotides in length, more preferably at least 20, 30, 40, 50, 100 or 200 nucleotides in length. Generally, the shorter the length of the polynucleotide sequence, the greater the homology required to obtain selective hybridisation. Consequently, where a polynucleotide sequence of the invention consists of less than about 30 nucleotides, it is preferred that the percentage identity is greater than 75%, preferably greater than 90% or 95% compared with the polynucleotide sequences set out in the sequence listings herein. Conversely, where a polynucleotide sequence of the invention consists of, for example, greater than 50 or 100 nucleotides, the percentage identity compared with the polynucleotide sequences set out in the sequence listings herein may be lower, for example greater than 50%, preferably greater than 60 or 75%.

[0100] Nucleic acid sequences according to the present invention which are homologous to the sequences as represented by a SEQ ID NO: 1 can be characterized and isolated according to any of the techniques known in the art, such as amplification by means of sequence-specific primers, hybridization with sequence-specific probes under more or less stringent conditions, serological screening methods or via the LiPA typing system.

DNA

[0101] The DNA of the new PMC virus also is provided. The DNA sequence is preferably derived from the RNA sequences described above. Most preferably, the DNA sequence is that shown in SEQ ID NO: 1 or fragments thereof.

[0102] The invention also provides DNA fragments hybridisable with the genomic RNA of PMC. The DNA or DNA fragment sequence may be derived from the cDNA sequence of the PMC virus or fragments thereof. The DNA, cDNA or fragments thereof may be in the form of recombinant DNAs.

[0103] The DNA sequence may also be a fragment of SEQ ID NO:1. Preferably, the fragment is selected from the following locations of SEQ ID NO:1: position 419-922, 923-1219, 1220-1885, 1886-2473, 2474-3604, 3605-3820, 3821-5224, 5225-7252, 7253-7441, 7442-8482, 8483-9997, 9998-12077.

Variant Nucleic Acids

[0104] Nucleic acid sequences and fragments, which would include some deletions or mutations which would not substantially alter their ability to hybridizing with the genome of PMC virus, are also provided by the present invention. Such variants are to be considered as forming obvious equivalents of the RNA, DNA or fragments referred to above.

[0105] Other preferred variant nucleic acid sequences of the present invention include sequences which are redundant as a result of the degeneracy of the genetic code compared to any of the above-given nucleic acid sequences of the present invention. These variant nucleic acid sequences will thus encode the same amino acid sequences as the nucleic acid sequences they are derived from. Preferably, the RNAs of these variants, and the related cDNAs derived from said RNAs, are hybridisable to corresponding parts of the RNA and cDNA of PMC virus.

[0106] Also included within the present invention are sequence variants of the DNA sequence of SEQ ID NO: 1 or corresponding RNA sequence or fragments thereof, containing either deletions and/or insertions of one or more nucleotides, especially insertions or deletions of 1 or more codons.

[0107] Also included are substitutions of some non-essential nucleotides by others (including modified nucleotides and/or inosine).

[0108] Particularly preferred variant polynucleotides of the present invention also include sequences which hybridise under stringent conditions with any of the nucleic acid sequences of the present invention. Thus, sequences which show a high degree of homology (similarity) to any of the nucleic acid sequences of the invention as described above are preferred. Particularly preferred are sequences which are at least 80%, 85%, 90%, 95% or more homologous to said nucleic acid sequences of the invention. Preferably, said sequences will have less than 20%, 15%, 10%, or 5% variation of the original nucleotides of said nucleic acid sequences.

Probes and Primers

[0109] Primer and probes are further provided, which can be made starting from any RNA or DNA sequence or sequence fragment according to the invention. Preferably, such probes or primers are between about 5 to 50 nucleotides long, more preferably from about 10 to 25 nucleotides. Probes and primers of the present invention may be used in PCR, sequencing reactions, hybridisation reactions and other applications known to the skilled person.

[0110] The present invention also relates to an oligonucleotide primer comprising part of SEQ ID NO: 1, said primer being able to act as a primer for specifically amplifying the nucleic acid of the PMC virus. Preferably, the primer is a single stranded DNA oligonucleotide sequence capable of

acting as a point of initiation for synthesis of a primer extension product which is complementary to the nucleic acid strand to be copied. The specific length and sequence of the primer used will depend on the complexity of the required DNA or RNA targets, as well as on the conditions of primer use, such as temperature and ionic strength. The fact that amplification primers do not have to match exactly with corresponding template sequence to warrant proper amplification is amply documented in the literature (Kwok et al., 1990).

[0111] The amplification method used can be either polymerase chain reaction (PCR; Saiki et al., 1988), ligase chain reaction (LCR; Landgren et al., 1988; Wu & Wallace, 1989; Barany, 1991), nucleic acid sequence-based amplification (NASBA; Guatelli et al., 1990; Compton, 1991), transcription-based amplification system (TAS; Kwok et al., 1989), strand displacement amplification (SDA; Duck, 1990; Walker et al., 1992) or amplification by means of Q β replicase (Lizardi et al., 1988; Lomeli et al., 1989) or any other suitable method to amplify nucleic acid molecules using primer extension. During amplification, the amplified products can be conveniently labelled either using labelled primers or by incorporating labelled nucleotides. Labels may be isotopic (³²P, ³⁵S, etc.) or non-isotopic (biotin, digoxigenin, etc.). The amplification reaction is repeated between 20 and 70 times, advantageously between 25 and 45 times.

[0112] The present invention also relates to an oligonucleotide probe comprising part of SEQ ID NO:1, with said probe being able to act as a hybridisation probe for the PMC virus. Preferably, the probe can be used for specific detection and/or classification into types and/or subtypes of PMC virus. Preferably, the probe is a single stranded sequence-specific oligonucleotide sequence which has a sequence that is complementary to the target sequence of the PMC virus to be detected.

[0113] Those skilled in the art will recognise that the stringency of hybridisation will be affected by such conditions as salt concentration, temperature, or organic solvents, in addition to the base composition, length of the complementary strands and the number of nucleotide base mismatches between the hybridising nucleic acids. Stringent temperature conditions will generally include temperatures in excess of 30° C., typically in excess of 37° C., and preferably in excess of 45° C. Stringent salt conditions will ordinarily be less than 1000 mM, typically less than 500 mM, and preferably less than 200 mM. However, the combination of parameters is much more important than the measure of any single parameter. An example of stringent hybridisation conditions is 65° C. and 0.1 \times SSC (1 \times SSC=0.15 M NaCl, 0.015 M sodium citrate pH 7.0).

[0114] Optionally, the probe of the invention is labelled and/or attached to a solid substrate. The solid substrate can refer to any substrate to which an oligonucleotide probe can be coupled, provided that it retains its hybridization characteristics and provided that the background level of hybridization remains low. Usually the solid substrate will be a microtiter plate, a membrane (e.g. nylon or nitrocellulose) or a microsphere (bead). Prior to application to the membrane or fixation it may be convenient to modify the nucleic acid probe in order to facilitate fixation or improve the hybridization efficiency. Such modifications may encompass homopolymer tailing, coupling with different reactive groups such as aliphatic groups, NH₂ groups, SH groups, carboxylic groups, or coupling with biotin or haptens.

[0115] The probes of the invention may include also an isolated polynucleotide attached to a label or reporter molecule and may be used to isolate other polynucleotide sequences, having sequence similarity by standard methods. For techniques for preparing and labelling probes see, e.g. Sambrook et al., (1989) or Ausubel et al., (2001).

[0116] Oligonucleotides according to the present invention and used as primers or probes may also contain or consist of nucleotide analogues such as phosphorothioates (Matsukura et al., 1987), alkylphosphorates (Miller et al., 1979) or peptide nucleic acids (Nielsen et al., 1991; Nielsen et al., 1993) or may contain intercalating agents (Asseline et al., 1984). The introduction of these modifications may be advantageous in order to positively influence characteristics such as hybridization kinetics, reversibility of the hybrid-formation, biological stability of the oligonucleotide molecules, etc.

[0117] Recombinant DNAs containing fragments of the DNA sequence of PMC virus are also provided by the present invention, and may be used as, for example, probes. Preferably, the plasmid used to generate the recombinant DNA is a plasmid amplifiable in prokaryotic or eukaryotic cells and carrying said fragments. For example, using cloned DNA containing a DNA fragment of PMC virus as a molecular hybridization probe, either by marking with radionucleotides or with fluorescent reagents, PMC virus RNA may be detected directly, for example, in blood, body fluids and blood products.

Nucleic Acid Arrays

[0118] PMC virus polynucleotide sequences (preferably in the form of probes) may also be immobilised to a solid phase support for the detection of PMC virus. Alternatively the PMC virus polynucleotide sequences will form part of a library of DNA molecules that may be used to detect simultaneously a number of different genes from PMC virus. In a further alternate form of the invention, PMC virus polynucleotide sequences together with other polynucleotide sequences (such as from other bacteria or viruses) may be immobilised on a solid support in such a manner permitting identification of the presence of PMC virus and/or any of the other polynucleotide sequences bound onto the solid support.

[0119] Techniques for producing immobilised libraries of DNA molecules have been described in the art. Generally, most prior art methods describe the synthesis of single-stranded nucleic acid molecule libraries, using for example masking techniques to build up various permutations of sequences at the various discrete positions on the solid substrate. U.S. Pat. No. 5,837,832 describes an improved method for producing DNA arrays immobilised to silicon substrates based on very large scale integration technology. In particular, U.S. Pat. No. 5,837,832 describes a strategy called "tiling" to synthesize specific sets of probes at spatially defined locations on a substrate that may be used to produce the immobilised DNA libraries of the present invention. U.S. Pat. No. 5,837,832 also provides references for earlier techniques that may also be used. Thus polynucleotide sequence probes may be synthesised in situ on the surface of the substrate.

[0120] Alternatively, single-stranded molecules may be synthesised off the solid substrate and each pre-formed sequence applied to a discrete position on the solid substrate. For example, polynucleotide sequences may be printed directly onto the substrate using robotic devices equipped with either pins or piezo electric devices.

[0121] The library sequences are typically immobilised onto or in discrete regions of a solid substrate. The substrate may be porous to allow immobilisation within the substrate or substantially non-porous, in which case the library sequences are typically immobilised on the surface of the substrate. The solid substrate may be made of any material to which polypeptides can bind, either directly or indirectly. Examples of suitable solid substrates include flat glass, silicon wafers, mica, ceramics and organic polymers such as plastics, including polystyrene and polymethacrylate. It may also be possible to use semi-permeable membranes such as nitrocellulose or nylon membranes, which are widely available. The semi-permeable membranes may be mounted on a more robust solid surface such as glass. The surfaces may optionally be coated with a layer of metal, such as gold, platinum or other transition metal. A particular example of a suitable solid substrate is the commercially available BiaCore™ chip (Pharmacia Biosensors).

[0122] Preferably, the solid substrate is generally a material having a rigid or semi-rigid surface. In preferred embodiments, at least one surface of the substrate will be substantially flat, although in some embodiments it may be desirable to physically separate regions for different polymers with, for example, raised regions or etched trenches. It is also preferred that the solid substrate is suitable for the high density application of DNA sequences in discrete areas of typically from 50 to 100 μm, giving a density of 10000 to 40000 dots/cm⁻².

[0123] The solid substrate is conveniently divided up into sections. This may be achieved by techniques such as photo-etching, or by the application of hydrophobic inks, for example teflon-based inks (Cel-line, USA).

[0124] Discrete positions, in which each different member of the library is located may have any convenient shape, e.g., circular, rectangular, elliptical, wedge-shaped, etc.

[0125] Attachment of the polynucleotide sequences to the substrate may be by covalent or non-covalent means. The polynucleotide sequences may be attached to the substrate via a layer of molecules to which the library sequences bind. For example, the polynucleotide sequences may be labelled with biotin and the substrate coated with avidin and/or streptavidin. A convenient feature of using biotinylated polynucleotide sequences is that the efficiency of coupling to the solid substrate can be determined easily. Since the polynucleotide sequences may bind only poorly to some solid substrates, it is often necessary to provide a chemical interface between the solid substrate (such as in the case of glass) and the nucleic acid sequences. Examples of suitable chemical interfaces include hexaethylene glycol. Another example is the use of polylysine coated glass, the polylysine then being chemically modified using standard procedures to introduce an affinity ligand. Other methods for attaching molecules to the surfaces of solid substrate by the use of coupling agents are known in the art, see for example WO98/49557.

[0126] Binding of complementary polynucleotide sequences to the immobilised nucleic acid library, may be determined by a variety of means such as changes in the optical characteristics of the bound polynucleotide sequence (i.e. by the use of ethidium bromide) or by the use of labelled nucleic acids, such as polypeptides labelled with fluorophores. Other detection techniques that do not require the use of labels include optical techniques such as optoacoustics, reflectometry, ellipsometry and surface plasmon resonance (see WO97/49989).

[0127] Thus, the present invention provides a solid substrate having immobilized thereon at least one polynucleotide of the present invention, preferably two or more different polynucleotide sequences of the present invention. In a preferred embodiment the solid substrate further comprises polynucleotide sequences derived from genes other than the PMC virus polynucleotide sequence.

Antisense Nucleic Acids and Ribozymes

[0128] The present invention also extends to the preparation of antisense nucleotides and ribozymes that may be used to interfere with the expression of PMC virus amino acid sequences at the translational level. This approach utilises antisense nucleic acid and ribozymes to block translation of a specific mRNA, either by masking that mRNA with an antisense nucleic acid or cleaving it with a ribozyme.

[0129] Antisense nucleic acids are DNA or RNA molecules that are complementary to at least a portion of a specific mRNA molecule [See: Weintraub, (1990) *Sci. Am.*, 262:40-46; Marcus-Sekura, (1988) *Anal. Biochem.*, 172:289-295]. In the cell, they hybridise to that mRNA, forming a double-stranded molecule. The cell does not translate an mRNA complexed in this double-stranded form. Therefore, antisense nucleic acids interfere with the expression of mRNA into protein. Oligomers of about fifteen nucleotides and molecules that hybridise to the AUG initiation codon will be particularly efficient, since they are easy to synthesize and are likely to pose fewer problems than larger molecules when introducing them into infected cells. Antisense methods have been used to inhibit the expression of many genes in vitro [Hambor et al., (1988) *J. Exp. Med.*, 168:1237-1245].

[0130] Ribozymes are RNA molecules possessing the ability to specifically cleave other single-stranded RNA molecules in a manner somewhat analogous to DNA restriction endonucleases. Ribozymes were discovered from the observation that certain mRNAs have the ability to excise their own introns. By modifying the nucleotide sequence of these RNAs, researchers have been able to engineer molecules that recognise specific nucleotide sequences in an RNA molecule and cleave it [Cech, (1988) *J. Am. Med. Assoc.*, 260:3030-3034]. Because they are sequence-specific, only mRNAs with particular sequences are inactivated. Investigators have identified two types of ribozymes, *Tetrahymena*-type and "hammerhead"-type. *Tetrahymena*-type ribozymes recognize four-base sequences, while "hammerhead"-type recognize eleven- to eighteen-base sequences. The longer the recognition sequence, the more likely it is to occur exclusively in the target mRNA species. Therefore, hammerhead-type ribozymes are preferable to *Tetrahymena*-type ribozymes for inactivating a specific mRNA species and eighteen base recognition sequences are preferable to shorter recognition sequences.

[0131] The PMC polynucleotide sequences described herein may thus be used to prepare antisense molecules against, and ribozymes that cleave, mRNAs for PMC virus amino acid sequences, thus inhibiting expression of the PMC virus polynucleotide sequences.

Polypeptide Sequences

Polypeptides

[0132] The invention also covers polypeptides encoded by the above RNA and DNA nucleotide sequences and fragments thereof. The invention further provides an isolated

PMC virus amino acid sequence as shown in SEQ ID NO: 2 and fragments thereof. More desirably, the PMC virus amino acid sequence is provided in substantially purified form. Further provided are polypeptide fragments having lower molecular weights and having peptide sequences or fragments in common with those shown in SEQ ID NO:2.

[0133] The term "isolated" is used to describe a PMC virus amino acid sequence that has been separated from components that accompany it in its natural state. Further, a PMC virus amino acid sequence is "substantially purified" when at least about 60 to 75% of a sample exhibits a single PMC virus amino acid sequence. A substantially purified PMC virus amino acid sequence will typically comprise about 60 to 90% W/W of a PMC virus amino acid sequence sample, more usually about 95%, and preferably will be over about 99% pure. Protein purity or homogeneity may be indicated by a number of means well known in the art, such as polyacrylamide gel electrophoresis of a protein sample, followed by visualizing a single PMC virus amino acid sequence band upon staining the gel. For certain purposes, higher resolution may be provided by using HPLC or other means well known in the art which are utilised for application.

[0134] The invention further contemplates fragments of the PMC virus amino acid sequence. A PMC virus amino acid sequence fragment is a stretch of amino acid residues of at least about five to seven contiguous amino acids, often at least about seven to nine contiguous amino acids, typically at least about nine to 13 contiguous amino acids and, most preferably, at least about 20 to 30 or more contiguous amino acids.

[0135] In a highly preferred form of the invention the fragments exhibit ligand-binding, immunological activity and/or other biological activities characteristic of PMC virus amino acid sequences. More preferably, the fragments possess immunological epitopes consistent with those present on native PMC virus amino acid sequences.

[0136] As used herein, "epitope" refers to an antigenic determinant of a polypeptide. An epitope could comprise three amino acids in a spatial conformation that is unique to the epitope. Generally, an epitope consists of at least five amino acids, and more usually consists of at least 8-10 amino acids. Methods of determining the spatial conformation of such amino acids are known in the art.

[0137] Preferred PMC virus amino acid sequences of the invention will have one or more biological properties (eg in vivo, in vitro or immunological properties) of the native full-length PMC virus amino acid sequence. Alternatively, fragments of the full-length PMC virus amino acid sequence may have one or more biological properties of one or more of the genes which the full length amino acid sequence encodes.

[0138] Preferably, the fragments of the full length PMC virus amino acid sequence SEQ ID NO:2 are chosen from the following locations in SEQ ID NO:2: 1-167, 168-267, 268-489, 490-685, 686-1062, 1063-1134, 1135-1602, 1603-2278, 2279-2341, 2342-2688, 2689-3193, 3194-3886. Alternatively, the fragment may be selected from any one of SEQ ID NOs:16-27.

[0139] Non-functional PMC virus amino acid sequences are also included within the scope of the invention since they may be useful, for example, as antagonists of PMC virus genes. The biological properties of analogues, fragments, or derivatives relative to wild type may be determined, for example, by means of biological assays.

[0140] PMC virus amino acid sequences, including analogues, fragments and derivatives, can be prepared syntheti-

cally (e.g., using the well known techniques of solid phase or solution phase peptide synthesis). Preferably, solid phase synthetic techniques are employed. Alternatively, PMC virus amino acid sequences of the invention can be prepared using well known genetic engineering techniques, as described infra.

[0141] In yet another embodiment, PMC virus amino acid sequences can be purified (e.g., by immunoaffinity purification) from a biological fluid, such as but not limited to whole blood, plasma, faeces, serum, or urine from animals, including pigs, cattle, sheep, chickens, human beings, dogs, horses, and fish.

Variant Polypeptides

[0142] PMC virus amino acid sequence analogues preferably include those having an amino acid sequence wherein one or more of the amino acids is substituted with another amino acid, which substitutions do not substantially alter the biological activity of the molecule.

[0143] In the context of the invention, an analogous sequence is taken to include a PMC virus amino acid sequence which is at least 60, 70, 80 or 90% homologous, preferably at least 95 or 98% homologous at the amino acid level over at least 20, 50, 100 or 200 amino acids, with the amino acid sequence set out in SEQ ID NO:1. In particular, homology should typically be considered with respect to those regions of the sequence known to be essential for the function of the protein or proteins encoded by the PMC virus RNA, rather than non-essential neighbouring sequences.

[0144] Although homology can be considered in terms of similarity (i.e. amino acid residues having similar chemical properties/functions), in the context of the present invention it is preferred to express homology in terms of sequence identity. The terms "substantial homology" or "substantial identity", when referring to PMC virus amino acid sequences, indicate that the PMC virus amino acid sequence in question exhibits at least about 70% identity with an entire naturally-occurring PMC amino acid sequence or portion thereof, usually at least about 80% identity and preferably at least about 90 or 95% identity.

[0145] In a highly preferred form of the invention, a PMC virus amino acid sequence analogue will have 80% or greater amino acid sequence identity to the PMC virus amino acid sequence set out in SEQ ID NO:2. Examples of PMC virus amino acid sequence analogues within the scope of the invention include the amino acid sequence of SEQ ID NO:2 wherein: (a) one or more aspartic acid residues is substituted with glutamic acid; (b) one or more isoleucine residues is substituted with leucine; (c) one or more glycine or valine residues is substituted with alanine; (d) one or more arginine residues is substituted with histidine; or (e) one or more tyrosine or phenylalanine residues is substituted with tryptophan.

[0146] PMC virus amino acid sequence derivatives are also provided by the invention and include PMC virus amino acid sequences, analogues or fragments thereof which are substantially homologous in primary structure but which include chemical and/or biochemical modifications or unusual amino acids. Such modifications include, for example, acetylation, carboxylation, phosphorylation, glycosylation, ubiquitination, labelling, (e.g., with radionucleotides), and various enzymatic modifications, as will be readily appreciated by those well skilled in the art.

[0147] In one form of the invention the chemical moieties suitable for derivatisation are selected from among water soluble polymers. The polymer selected should be water soluble so that the protein to which it is attached does not precipitate in an aqueous environment, such as a physiological environment. Preferably, for therapeutic use of the end-product preparation, the polymer will be pharmaceutically acceptable. One skilled in the art will be able to select the desired polymer based on considerations such as whether the polymer/protein conjugate will be used therapeutically, and if so, the desired dosage, circulation time, resistance to proteolysis and other considerations. For the present proteins and peptides, these may be ascertained using the assays provided herein.

[0148] The water soluble polymer may be selected from the group consisting of, for example, polyethylene glycol, copolymers of ethylene glycol/propylene glycol, carboxymethylcellulose, dextran, polyvinyl alcohol, polyvinyl pyrrolidone, poly-1,3-dioxolane, poly-1,3,6-trioxane, ethylene/maleic anhydride copolymer, polyaminoacids (either homopolymers or random copolymers), and dextran or poly (n-vinyl pyrrolidone)polyethylene glycol, propylene glycol homopolymers, polypropylene oxide/ethylene oxide copolymers, polyoxyethylated polyols and polyvinyl alcohol. Polyethylene glycol propionaldehyde may provide advantages in manufacturing due to its stability in water.

[0149] In another form of the invention the amino acid sequences may be modified to produce a longer half life in an animal host, for example, by fusing one or more antibody fragments (such as an Fc fragment) to the amino or carboxyl end of a PMC virus amino acid sequence.

[0150] Where the PMC virus amino acid sequence is to be provided in a labelled form, a variety of methods for labelling amino acid sequences are well known in the art and include radioactive isotopes such as ³²P, ligands which bind to labelled antiligands (eg, antibodies), fluorophores, chemiluminescent agents, enzymes and antiligands which can serve as specific binding pair members for a labelled ligand. The choice of label depends on the sensitivity required, stability requirements, and available instrumentation. Methods of labelling amino acid sequences are well known in the art [See, e.g., Sambrook et al., *Molecular Cloning: A Laboratory Manual*, Second Edition, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1989); and Ausubel, F., Brent, R., Kingston, R. E., Moore, D. D., Seidman, J. G., Smith, J. A., Struhl, K. *Current protocols in molecular biology*. Greene Publishing Associates/Wiley Intersciences, New York (2001)].

[0151] The PMC virus amino acid sequences of the invention, if soluble, may be coupled to a solid-phase support, e.g., nitrocellulose, nylon, column packing materials (e.g., Sepharose beads), magnetic beads, glass wool, plastic, metal, polymer gels, cells, or other substrates. Such supports may take the form, for example, of beads, wells, dipsticks, or membranes.

[0152] The invention also provides for fusion polypeptides, comprising PMC virus amino acid sequences and fragments. Thus PMC virus amino acid sequences may be fusions between two or more PMC virus amino acid sequences or between a PMC virus amino acid sequence and a related protein. Likewise, heterologous fusions may be constructed which would exhibit a combination of properties or activities of the derivative proteins. For example, ligand-binding or other domains may be "swapped" between different fusion

polypeptides or fragments. Such homologous or heterologous fusion polypeptides may display, for example, altered strength or specificity of binding. Fusion partners include immunoglobulins, bacterial beta-galactosidase, trpE, protein A, beta-lactamase, alpha amylase, alcohol dehydrogenase and yeast alpha mating factor.

[0153] Modified PMC virus amino acid sequences may be synthesised using conventional techniques, or may be encoded by a modified polynucleotide sequence and produced using recombinant nucleic acid methods. The modified polynucleotide sequence may also be prepared by conventional techniques. Fusion proteins will typically be made by either recombinant nucleic acid methods or may be chemically synthesised.

Diagnostics

[0154] In accordance with another embodiment the invention provides diagnostic and prognostic methods to detect the presence of PMC virus using PMC virus glycoproteins, proteins and other peptides and polypeptides (whether obtained in a purified state from PMC virus preparations, or by chemical synthesis) and/or antibodies derived there from and/or PMC virus polynucleotide sequences.

[0155] Diagnostic and prognostic methods will generally be conducted using a biological sample obtained from an animal, such as a pig. A "sample" refers to a sample of tissue or fluid suspected of containing a PMC polynucleotide or polypeptide from an animal, but not limited to, e.g., whole blood, blood cells, plasma, serum, milk, faecal samples, tissue and samples of in vitro cell culture constituents.

Polypeptide/Antibody-Based Diagnostics

[0156] Means are provided for the detection of proteins of PMC virus, particularly for the diagnosis of PMC or for the detection of antibodies against PMC virus or its proteins, particularly in subjects afflicted with PMC or more generally in asymptomatic carriers and in animal derived products such as meat. Such methods are also referred to as immunoassays.

[0157] The invention thus provides a method for detecting the presence of a PMC virus amino acid sequence in a sample, comprising the steps of:

[0158] a) contacting a sample suspected of containing a PMC virus amino acid sequence with an antibody that specifically binds to the PMC virus amino acid sequence under conditions which allow for the formation of reaction complexes comprising the antibody and the PMC virus amino acid sequence; and

[0159] b) detecting the formation of reaction complexes comprising the antibody and PMC virus amino acid sequence in the sample, wherein detection of the formation of reaction complexes indicates the presence of PMC virus amino acid sequence in the sample.

[0160] Particularly the invention relates to an in vitro process of diagnosis making use of an amino acid sequence encoding an envelope glycoprotein or of a polypeptide bearing an epitope of a glycoprotein from PMC virus or any other viral protein (structural or non-structural) for the detection of anti-PMC virus antibodies in serum, milk or body fluids. Preferably, the antibody used in the above methods binds to the E0, E1, E2, NS2, NS3, NS4A, NS4B and/or NS5A, NS5B proteins of PMC virus.

[0161] The invention also provides a method for detecting the presence of a PMC virus antibody in a sample, comprising the steps of:

[0162] a) contacting a sample suspected of containing a PMC virus antibody with an amino acid sequence under conditions which allow for the formation of reaction complexes comprising the PMC virus antibody and the amino acid sequence; and

[0163] b) detecting the formation of reaction complexes comprising the antibody and amino acid sequence in the sample, wherein detection of the formation of reaction complexes indicates the presence of PMC virus antibody in the sample.

[0164] A method is also provided for the detection of anti-PMC virus antibodies, comprising the steps of:

[0165] a) depositing a predetermined amount of one or several PMC virus antigens onto a solid support such as a microplate;

[0166] b) introducing increasing dilutions of a biological fluid (e.g., blood serum or plasma, milk, cerebrospinal fluid, lymphatic fluid or other body fluids) onto the antigens and incubating;

[0167] c) washing the solid support with an appropriate buffer;

[0168] d) adding specific labelled antibodies directed against the antibodies of the subject; and

[0169] e) detecting the antigen-antibody-antibody complex formed, which is then indicative of the presence of PMC virus antibodies in the biological fluid.

[0170] Preferably, the antibody used in these methods is derived from an affinity-purified polyclonal antibody, and more preferably a mAb. In addition, it is preferable for the antibody molecules used herein be in the form of Fab, Fab', F(ab')₂ or F(v) portions or whole antibody molecules.

[0171] Particularly preferred methods for detecting PMC virus based on the above methods include enzyme linked immunosorbent assays, radioimmunoassays, immunoradiometric assays and immunoenzymatic assays, including sandwich assays using monoclonal and/or polyclonal antibodies.

[0172] Three such procedures that are especially useful utilise either PMC virus amino acid sequences (or fragments thereof) labelled with a detectable label, antibody Ab₁ labelled with a detectable label, or antibody Ab₂ labelled with a detectable label. The procedures may be summarized by the following equations wherein the asterisk indicates that the particle is labelled and "AA" stands for the PMC virus amino acid sequence:



[0173] The procedures and their application are all familiar to those skilled in the art and accordingly may be utilised within the scope of the present invention. The "competitive" or "blocking" procedure, Procedure A, is described in U.S. Pat. Nos. 3,654,090 and 3,850,752. Procedure B is representative of well-known competitive assay techniques. Procedure C, the "sandwich" procedure, is described in U.S. Pat. Nos. RE 31,006 and 4,016,043. Still other procedures are known, such as the "double antibody" or "DASP" procedure.

[0174] In each instance, the PMC virus amino acid sequences form complexes with one or more antibody(ies) or

binding partners and one member of the complex is labelled with a detectable label. The fact that a complex has formed and, if desired, the amount thereof, can be determined by known methods applicable to the detection of labels.

[0175] It will be seen from the above that a characteristic property of Ab₂ is that it will react with Ab₁. This is because Ab₁, raised in one mammalian species, has been used in another species as an antigen to raise the antibody, Ab₂. For example, Ab₂ may be raised in goats using rabbit antibodies as antigens. Ab₂ therefore would be anti-rabbit antibody raised in goats. For purposes of this description and claims, Ab₁ will be referred to as a primary antibody, and Ab₂ will be referred to as a secondary or anti-Ab₁ antibody.

[0176] The labels most commonly employed for these studies are radioactive elements, enzymes, chemicals that fluoresce when exposed to ultraviolet light, and others.

[0177] A number of fluorescent materials are known and can be utilised as labels. These include, for example, fluorescein, rhodamine and auramine. A particular detecting material is anti-rabbit antibody prepared in goats and conjugated with fluorescein through an isothiocyanate.

[0178] The PMC virus amino acid sequences or their binding partners can also be labelled with a radioactive element or with an enzyme. The radioactive label can be detected by any of the currently available counting procedures. The preferred isotope may be selected from ³H, ¹⁴C, ³²P, ³⁵S, ³⁶Cl, ⁵¹Cr, ⁵⁷Co, ⁵⁸Co, ⁵⁹Fe, ⁹⁰Y, ¹²⁵I, ¹³¹I, and ¹⁸⁶Re.

[0179] Enzyme labels are likewise useful, and can be detected by any of the presently utilized colorimetric, spectrophotometric, fluorospectrophotometric, amperometric or gasometric techniques. The enzyme is conjugated to the selected particle by reaction with bridging molecules such as carbodiimides, diisocyanates, glutaraldehyde and the like. Many enzymes, which can be used in these procedures, are known and can be utilized. The preferred enzymes are peroxidase, β-glucuronidase, β-D-glucosidase, β-D-galactosidase, urease, glucose oxidase plus peroxidase and alkaline phosphatase. U.S. Pat. Nos. 3,654,090, 3,850,752 and 4,016,043 are referred to by way of example for their disclosure of alternate labelling material and methods.

[0180] In another embodiment of the invention there are provided in vitro methods for evaluating the level of PMC virus antibodies in a biological sample comprising the steps of:

[0181] a) detecting the formation of reaction complexes in a biological sample according to the method noted above; and

[0182] b) evaluating the amount of reaction complexes formed, which amount of reaction complexes corresponds to the level of PMC virus antibodies in the biological sample.

[0183] Preferably, the antibody used in the above methods binds to the E0, E1, E2, NS2, NS3, NS4A, NS4B and/or NS5A, NS5B proteins of PMC virus.

[0184] In another embodiment of the invention there are provided in vitro methods for evaluating the level of PMC virus polypeptides in a biological sample comprising the steps of:

[0185] a) detecting the formation of reaction complexes in a biological sample according to the method noted above; and

- [0186]** b) evaluating the amount of reaction complexes formed, which amount of reaction complexes corresponds to the level of PMC virus polypeptide in the biological sample.
- [0187]** Preferably, the polypeptide used in the above methods encodes the E0, E1, E2, NS2, NS3, NS4A, NS4B and/or NS5A, NS5B proteins of PMC virus.
- [0188]** Further there are provided in vitro methods for monitoring therapeutic treatment of a disease associated with PMC virus in an animal host comprising evaluating, as describe above, the levels of PMC virus antibodies in a series of biological samples obtained at different time points from an animal host undergoing such therapeutic treatment.
- [0189]** The methods for detecting polypeptides using antibodies, or immunoassays, according to the present invention may utilize antigens from the different domains of the new and unique polypeptide sequences of the present invention that maintain linear (in case of peptides) and conformational epitopes (in case of polypeptides) recognized by antibodies in the sera from subjects infected with PMC virus.
- [0190]** It is within the scope of the invention to use, for instance, single or specific oligomeric antigens, dimeric antigens, as well as combinations of single or specific oligomeric antigens.
- [0191]** The PMC virus antigens of the present invention may be employed in virtually any assay format that employs a known antigen to detect antibodies. Of course, a format that denatures the PMC virus conformational epitope should be avoided or adapted.
- [0192]** A common feature of all of these detection methods is that the antigen is contacted with the test specimen suspected of containing PMC virus antibodies under conditions that permit the antigen to bind to any such antibody present in the component. Such conditions will typically be physiologic temperature, pH and ionic strength, using an appropriate predetermined quantity of antigen. The incubation of the antigen with the specimen is followed by detection of immune complexes comprised of the antigen and antibodies derived from the specimen typically by using a labelled second antibody that is directed against the immunoglobulins of the test animal species.
- [0193]** Design of the immunoassays is subject to a great deal of variation, and many formats are known in the art. Protocols may, for example, use solid supports, or immunoprecipitation. Assays which amplify the signals from the immune complex are also known; examples of which are assays which utilize biotin and avidin or streptavidin, and enzyme-labelled and mediated immunoassays, such as ELISA assays. Furthermore, the immunoassay may be, without limitation, in a heterogeneous or in a homogeneous format, and of a standard or competitive type.
- [0194]** In a heterogeneous format, the polypeptide is typically bound to a solid matrix or support to facilitate separation of the sample from the polypeptide after incubation. Examples of solid supports that can be used are nitrocellulose (e.g., in membrane or microtiter well form), polyvinyl chloride (e.g., in sheets or microtiter wells), polystyrene latex (e.g., in beads or microtiter plates, polyvinylidene fluoride (known as Immunolon™), diazotized paper, nylon membranes, activated beads, and Protein A beads. For example, Dynatech Immunolon™ 1 or Immunolon™ 2 microtiter plates or 0.25 inch polystyrene beads (Precision Plastic Ball) can be used in the heterogeneous format. The solid support containing the antigenic polypeptides is typically washed after separating it from the test sample, and prior to detection of bound antibodies. Both standard and competitive formats are known in the art.
- [0195]** In a homogeneous format, the test sample is incubated with the combination of antigens in solution. For example, it may be under conditions that will precipitate any antigen-antibody complexes which are formed. Both standard and competitive formats for these assays are known in the art.
- [0196]** In a standard format, the amount of PMC virus antibodies in the antibody-antigen complexes is directly monitored. This may be accomplished by determining whether labelled anti-xenogeneic (e.g. anti-swine) antibodies which recognize an epitope on anti-PMC virus antibodies will bind due to complex formation. In a competitive format, the amount of PMC virus antibodies in the sample is deduced by monitoring the competitive effect on the binding of a known amount of labelled antibody (or other competing ligand) in the complex.
- [0197]** Complexes formed comprising anti-PMC virus antibody (or in the case of competitive assays, the amount of competing antibody) are detected by any of a number of known techniques, depending on the format. For example, unlabelled PMC virus antibodies in the complex may be detected using a conjugate of anti-xenogeneic Ig complexed with a label (e.g. an enzyme label).
- [0198]** In an immunoprecipitation or agglutination assay format, the reaction between the PMC virus antigens and the antibody forms a network that precipitates from the solution or suspension and forms a visible layer or film of precipitate. If no anti-PMC antibody is present in the test specimen, no visible precipitate is formed.
- [0199]** There currently exist three specific types of particle agglutination (PA) assays. These assays are used for the detection of antibodies to various antigens when coated to a support. One type of this assay is the haemagglutination assay using red blood cells (RBCs) that are sensitized by passively adsorbing antigen (or antibody) to the RBC. The addition of specific antigen antibodies present in the body component, if any, causes the RBCs coated with the purified antigen to agglutinate.
- [0200]** To eliminate potential non-specific reactions in the hemagglutination assay, two artificial carriers may be used instead of RBC in the PA. The most common of these are latex particles. However, gelatin particles may also be used. The assays utilizing either of these carriers are based on passive agglutination of the particles coated with purified antigens.
- #### Nucleic Acid-Based Diagnostics
- [0201]** The present invention further provides methods for detecting the presence or absence of PMC virus in a biological sample, which comprise the steps of:
- [0202]** c) bringing the biological sample into contact with a polynucleotide probe or primer comprising a PMC virus polynucleotide of the invention under suitable hybridising conditions; and
- [0203]** d) detecting any duplex formed between the probe or primer and nucleic acid sequences in the sample.
- [0204]** According to one embodiment of the invention, detection of PMC virus may be accomplished by directly amplifying PMC virus polynucleotide sequences from biological sample, using known techniques and then detecting the presence of PMC virus polynucleotide sequences.

[0205] The present invention thus also relates to a method for the detection of PMC virus nucleic acids present in a biological sample, comprising:

[0206] c) amplifying the nucleic acid with at least one primer as defined above,

[0207] d) detecting the amplified nucleic acids.

[0208] Preferably, the nucleic acid is extracted and/or purified (eg from a from a tissue sample) prior to amplification.

[0209] The present invention also relates to a method for the detection of PMC virus nucleic acids present in a biological sample, comprising:

[0210] d) hybridizing the nucleic acids of the biological sample at appropriate conditions with one or more probes as defined above,

[0211] e) washing under appropriate conditions, and

[0212] f) detecting the hybrids formed.

[0213] Preferably, the hybridizing conditions are denatured conditions.

[0214] Preferably, the nucleic acid is extracted and/or purified (eg from a from a tissue sample) prior to hybridisation. More preferably, the nucleic acid sample is amplified with at least one primer as defined above, after extraction or at least prior to hybridisation. Preferably, said probes are attached to a solid substrate or detected in a liquid phase by photometric or fluorogenic detection or by other methods of visualisation such as by agarose gel electrophoresis.

[0215] The present invention also relates to a method as defined above, wherein said nucleic acids are labelled during or after amplification.

[0216] Suitable assay methods for purposes of the present invention to detect hybrids formed between the oligonucleotide probes and the nucleic acid sequences in a sample may comprise any of the assay formats known in the art, such as the conventional dot-blot format, sandwich hybridization or reverse hybridization. For example, the detection can be accomplished using a dot blot format, the unlabelled amplified sample being bound to a membrane, the membrane being incorporated with at least one labelled probe under suitable hybridization and wash conditions, and the presence of bound probe being monitored.

[0217] An alternative and preferred method is a "reverse" dot-blot format, in which the amplified sequence contains a label. In this format, the unlabelled oligonucleotide probes are bound to a solid support and exposed to the labelled sample under appropriate stringent hybridization and subsequent washing conditions. It is to be understood that also any other assay method which relies on the formation of a hybrid between the nucleic acids of the sample and the oligonucleotide probes according to the present invention may be used.

[0218] In one form of the invention, the target nucleic acid sequence is amplified by PCR and then detected using any of the specific methods mentioned above. Other useful diagnostic techniques for detecting the presence of PMC virus polynucleotide sequences include, but are not limited to: 1) allele-specific PCR; 2) single stranded conformation analysis; 3) denaturing gradient gel electrophoresis; 4) RNase protection assays; 5) the use of proteins which recognize nucleotide mismatches, such as the *E. coli* mutS protein; 6) allele-specific oligonucleotides; and 7) fluorescent in situ hybridisation.

[0219] In addition to the above methods, PMC virus polynucleotide sequences may be detected using conventional probe technology. When probes are used to detect the presence of the PMC virus polynucleotide sequences, the biological

sample to be analysed, such as blood or serum, may be treated, if desired, to extract the nucleic acids. The sample polynucleotide sequences may be prepared in various ways to facilitate detection of the target sequence; e.g. denaturation, restriction digestion, electrophoresis or dot blotting. The targeted region of the sample polynucleotide sequence usually must be at least partially single-stranded to form hybrids with the targeting sequence of the probe. If the sequence is naturally single-stranded, denaturation will not be required. However, if the sequence is double-stranded, the sequence will probably need to be denatured. Denaturation can be carried out by various techniques known in the art.

[0220] Sample polynucleotide sequences and probes are incubated under conditions that promote stable hybrid formation of the target sequence in the probe with the putative PMC virus polynucleotide sequence in the sample. Preferably, high stringency conditions are used in order to prevent false positives.

[0221] Detection, if any, of the resulting hybrid is usually accomplished by the use of labelled probes. Alternatively, the probe may be unlabelled, but may be detectable by specific binding with a ligand that is labelled, either directly or indirectly. Suitable labels and methods for labelling probes and ligands are known in the art, and include, for example, radioactive labels which may be incorporated by known methods (e.g., nick translation, random priming or kinasings), biotin, fluorescent groups, chemiluminescent groups (e.g., dioxetanes, particularly triggered dioxetanes), enzymes, antibodies and the like. Variations of this basic scheme are known in the art, and include those variations that facilitate separation of the hybrids to be detected from extraneous materials and/or that amplify the signal from the labelled moiety.

[0222] It is also contemplated within the scope of this invention that the nucleic acid probe assays of this invention may employ a cocktail of nucleic acid probes and/or primers capable of detecting PMC virus polynucleotide sequences. Thus, in one example to detect the presence of PMC virus polynucleotide sequences in a cell sample, more than one probe complementary to PMC virus polynucleotide sequences is employed and in particular the number of different probes is alternatively 2, 3, or 5 different nucleic acid probe sequences.

[0223] Additionally, the present invention provides a method for detecting viral RNA or DNA comprising the steps of:

[0224] a) immobilizing PMC virus on a support (e.g., a nitrocellulose filter);

[0225] b) disrupting the virion; and

[0226] c) hybridizing with a probe.

[0227] Preferably, the probe is labelled. More preferably, the probe is radiolabelled or fluorescent- or enzyme-labelled. Such an approach to detection of virus has already been developed for Hepatitis B virus in peripheral blood (Scotto J. et al. Hepatology (1983), 3, 379-384).

[0228] The present invention also provides a method for rapid screening of genomic DNA derived from the tissue of subjects with PMC virus related symptoms to detect proviral PMC virus related DNA or RNA present in the tissues. Thus, the present invention also provides a method for screening the tissue of subjects comprising the steps of:

[0229] a) extracting DNA from tissue;

[0230] b) restriction enzyme cleavage of said DNA;

[0231] c) electrophoresis of the fragments; and

[0232] d) Southern blotting of genomic DNA from tissues and subsequent hybridization with labelled cloned PMC virus DNA.

[0233] Hybridization in situ can also be used.

Antigenic Polypeptide Production

[0234] Viral RNA and DNA according to the invention can be used for expressing PMC viral antigens for diagnostic purposes, as well as for the production of a vaccine against PMC virus. The methods which can be used to achieve expression of antigenic polypeptides are multifold:

a) DNA can be transfected into mammalian cells with appropriate selection markers by a variety of techniques, such as calcium phosphate precipitation, polyethylene glycol, protoplast-fusion, etc and the resultant proteins purified.

b) DNA fragments corresponding to genes can be cloned into expression vectors for *E. coli*, yeast or mammalian cells and the resultant proteins purified.

c) The proviral RNA or DNA can be "shot-gunned" (fragmented) into prokaryotic expression vectors to generate fusion polypeptides. Recombinants, producing antigenically competent fusion proteins, can be identified by simply screening the recombinants with antibodies against PMC virus antigens.

[0235] Particular reference in this respect is made to those portions of the genome of PMC virus which, in the figures, are shown to belong to open reading frames and which encode the products having the polypeptide sequences shown. Preferably, the nucleic acid sequences used in the above methods encode the E0, E1, E2, NS2, NS3, NS4A, NS4B and/or NS5A, NS5B proteins of PMC. Preferably, polypeptides are provided containing sequences in common with polypeptides comprising antigenic determinants included in the proteins encoded and expressed by the PMC virus genome.

Antibodies

Antibodies to PMC Proteins

[0236] The different peptides according to this invention can also be used themselves for the production of antibodies, preferably monoclonal antibodies specific for the respective different peptides. Thus, according to the invention, PMC virus amino acid sequences produced recombinantly or by chemical synthesis and fragments or other derivatives or analogues thereof, including fusion proteins, may be used as an immunogen to generate antibodies that recognize the PMC virus amino acid sequence. Such antibodies include but are not limited to polyclonal, monoclonal, chimeric, single chain, Fab fragments and a Fab expression library.

[0237] Thus, the present invention provides a method for the generation of antibodies comprising the steps of:

[0238] a) providing a PMC virus polypeptide sequence to a subject; and

[0239] b) collecting the antibodies generated in the subject against the polypeptide.

[0240] Preferably, the polypeptide used to generate the antibody is antigenic. More preferably, the polypeptide is chosen from the list comprising the E0, E1, E2, NS2, NS3, NS4A, NS4B and/or NS5A or NS5B proteins of PMC virus. More preferably, the protein used to generate the antibody is the E0, E2, NS2 and/or NS3 proteins or a fragment or derivative thereof. For example, in a highly preferred embodiment, a composition of the invention comprises both a PMC virus E0/E2 complex and an PMC virus NS2/NS3 complex.

[0241] A molecule is "antigenic" when it is capable of specifically interacting with an antigen recognition molecule of the immune system, such as an immunoglobulin (antibody) or T cell antigen receptor. An antigenic amino acid sequence contains at least about 5, and preferably at least about 10, amino acids. An antigenic portion of a molecule can be that portion that is immunodominant for antibody or T cell receptor recognition, or it can be a portion used to generate an antibody to the molecule by conjugating the antigenic portion to a carrier molecule for immunization. A molecule that is antigenic need not be itself immunogenic, i.e., capable of eliciting an immune response without a carrier.

[0242] An "antibody" is any immunoglobulin, including antibodies and fragments thereof, that binds a specific epitope. The term encompasses polyclonal, monoclonal, and chimeric antibodies, the last mentioned described in further detail in U.S. Pat. Nos. 4,816,397 and 4,816,567, as well as antigen binding portions of antibodies, including Fab, F(ab')₂ and F(v) (including single chain antibodies). Accordingly, the phrase "antibody molecule" in its various grammatical forms as used herein contemplates both an intact immunoglobulin molecule and an immunologically active portion of an immunoglobulin molecule containing the antibody combining site. An "antibody combining site" is that structural portion of an antibody molecule comprised of heavy and light chain variable and hypervariable regions that specifically binds antigen.

[0243] Exemplary antibody molecules are intact immunoglobulin molecules, substantially intact immunoglobulin molecules and those portions of an immunoglobulin molecule that contain the paratope, including those portions known in the art as Fab, Fab', F(ab')₂ and F(v), which portions are preferred for use in the therapeutic methods described herein.

[0244] Fab and F(ab')₂ portions of antibody molecules are prepared by the proteolytic reaction of papain and pepsin, respectively, on substantially intact antibody molecules by methods that are well-known. See for example, U.S. Pat. No. 4,342,566 to Theofilopolous et al. Fab' antibody molecule portions are also well-known and are produced from F(ab')₂ portions followed by reduction with mercaptoethanol of the disulfide bonds linking the two heavy chain portions, and followed by alkylation of the resulting protein mercaptan with a reagent such as iodoacetamide. An antibody containing intact antibody molecules is preferred herein.

[0245] The phrase "monoclonal antibody" in its various grammatical forms refers to an antibody having only one species of antibody combining site capable of immunoreacting with a particular antigen. A monoclonal antibody thus typically displays a single binding affinity for any antigen with which it immunoreacts.

[0246] For the production of hybridomas secreting said monoclonal antibodies, conventional production and screening methods can be used. These monoclonal antibodies, which themselves are part of the invention, provide very useful tools for the identification and even determination of relative proportions of the different polypeptides or proteins in biological samples, particularly animals samples containing PMC virus or related viruses.

[0247] Adjuvants include, but are not limited to, complete Freund's adjuvant, incomplete Freund's adjuvant, saponin, mineral gels such as aluminium hydroxide, surface active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil or hydrocarbon emulsions, keyhole limpet hemocyanins, dinitrophenol, and potentially useful human

adjuvants such as BCG (bacille Calmette-Guerin) and *Corynebacterium parvum*. Preferably, the adjuvant is pharmaceutically acceptable.

[0248] Further examples of adjuvants which may be effective include but are not limited to: N-acetyl-muramyl-L-threonyl-D-isoglutamine (thr-MDP), N-acetyl-nor-muramyl-L-alanyl-D-isoglutamine (CGP 11637, referred to as nor-MDP), N-acetylmuramyl-L-alanyl-D-isoglutaminyl-L-alanine-2-(1'-2'-dipalmitoyl-sn-glycero-3-hydroxyphosphoryloxy)-ethylamine (CGP 19835A, referred to as MTP-PE), and RIBI, which contains three components extracted from bacteria, monophosphoryl lipid A, trehalose dimycolate and cell wall skeleton (MPL+TDM+CWS) in a 2% squalene/Tween 80 emulsion.

[0249] Additional examples of adjuvants and other agents include aluminium hydroxide, aluminium phosphate, aluminium potassium sulfate (alum), beryllium sulfate, silica, kaolin, carbon, water-in-oil emulsions, oil-in-water emulsions, muramyl dipeptide, bacterial endotoxin, lipid X, *Corynebacterium parvum* (*Propionibacterium acnes*), *Bordetella pertussis*, polyribonucleotides, sodium alginate, lanolin, lysolecithin, vitamin A, saponin, immuno stimulating complexes (ISCOMs), liposomes, levamisole, DEAE-dextran, blocked copolymers or other synthetic adjuvants. Such adjuvants are available commercially from various sources, for example, Merck Adjuvant 65 (Merck and Company, Inc., Rahway, N.J.) or Freund's Incomplete Adjuvant and Complete Adjuvant (Difco Laboratories, Detroit, Mich.).

[0250] Typically, adjuvants such as Amphigen (oil-in-water), Alhydrogel (aluminium hydroxide), or a mixture of Amphigen and Alhydrogel are used. Only aluminium hydroxide is approved for human use.

[0251] The proportion of immunogenic polypeptide and adjuvant can be varied over a broad range so long as both are present in effective amounts. For example, aluminium hydroxide can be present in an amount of about 0.5% of the vaccine mixture (Al₂O₃ basis). Conveniently, the vaccines are formulated to contain a final concentration of immunogen in the range of from 0.2 to 200 µg/ml, preferably 5 to 50 µg/ml, most preferably 15 µg/ml.

[0252] After formulation, the vaccine may be incorporated into a sterile container which is then sealed and stored at a low temperature, for example 4° C., or it may be freeze-dried. Lyophilisation permits long-term storage in a stabilised form.

[0253] The vaccines are conventionally administered parenterally, by injection, for example, either subcutaneously or intramuscularly. Additional formulations which are suitable for other modes of administration include suppositories and, in some cases, oral formulations. For suppositories, traditional binders and carriers may include, for example, polyalkylene glycols or triglycerides; such suppositories may be formed from mixtures containing the active ingredient in the range of 0.5% to 10%, preferably 1% to 2%. Oral formulations include such normally employed excipients as, for example, pharmaceutical grades of mannitol, lactose, starch, magnesium stearate, sodium saccharine, cellulose, magnesium carbonate, and the like. These compositions take the form of solutions, suspensions, tablets, pills, capsules, sustained release formulations or powders and contain 10% to 95% of active ingredient, preferably 25% to 70%. Where the vaccine composition is lyophilised, the lyophilised material may be reconstituted prior to administration, e.g. as a suspension. Reconstitution is preferably effected in buffer

[0254] Capsules, tablets and pills for oral administration to a patient may be provided with an enteric coating comprising, for example, Eudragit "S", Eudragit "L", cellulose acetate, cellulose acetate phthalate or hydroxypropylmethyl cellulose.

[0255] The PMC virus polypeptides of the invention may be formulated into the vaccine as neutral or salt forms. Pharmaceutically acceptable salts include the acid addition salts (formed with free amino groups of the peptide) and which are formed with inorganic acids such as, for example, hydrochloric or phosphoric acids, or such organic acids such as acetic, oxalic, tartaric and maleic. Salts formed with the free carboxyl groups may also be derived from inorganic bases such as, for example, sodium, potassium, ammonium, calcium, or ferric hydroxides, and such organic bases as isopropylamine, trimethylamine, 2-ethylamino ethanol, histidine and procaine.

[0256] Compositions of the present invention may further comprise antigenic polypeptides that are not coupled to PMC virus polypeptides and/or biologically active molecules whose primary purpose is not to serve as an antigen but to modulate the immune response in some other aspect. Examples of biologically active molecules that modulate the immune system of an animal or human subject include cytokines.

[0257] The term "cytokine" refers to any secreted polypeptide that influences the function of other cells mediating an immune response. Some examples of cytokines include, but are not limited to, interleukin-1 α (IL-1 α), interleukin-1 β (IL-1 β), interleukin-2 (IL-2), interleukin-3 (IL-3), interleukin-4 (IL-4), interleukin-5 (IL-5), interleukin-6 (IL-6), interleukin-7 (IL-7), interleukin-8 (IL-8), interleukin-9 (IL-9), interleukin-10 (IL-10), interleukin-11 (IL-11), interleukin-12 (IL-12), interferon- α (IFN- α), interferon- β (IFN- β), interferon- γ (IFN- γ), tumour necrosis factor- α (TNF- α), tumour necrosis factor- β (TNF- β), granulocyte colony stimulating factor (G-CSF), granulocyte/macrophage colony stimulating factor (GM-CSF), and transforming growth factor- β (TGF- β).

[0258] Various procedures known in the art may be used for the production of polyclonal antibodies to PMC virus amino acid sequences, or fragment, derivative or analogues thereof.

[0259] For the production of antibody, various host animals can be immunised by injection with the PMC virus amino acid sequence, or a derivative (e.g., fragment or fusion protein) thereof, including but not limited to rabbits, mice, rats, sheep, goats, etc.

[0260] In one embodiment, the PMC virus amino acid sequences or fragment thereof can be conjugated to an immunogenic carrier, e.g., bovine serum albumin (BSA) or keyhole limpet hemocyanin (KLH).

[0261] Various adjuvants may be used to increase the immunological response, depending on the host species, including but not limited to Freund's (complete and incomplete), mineral gels such as aluminum hydroxide, surface active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, keyhole limpet hemocyanins, dinitrophenol, and potentially useful human adjuvants such as BCG (bacille Calmette-Guerin) and *Corynebacterium parvum*.

[0262] For preparation of monoclonal antibodies directed toward the PMC virus amino acid sequences, or fragments, analogues, or derivatives thereof, any technique that provides for the production of antibody molecules by continuous cell lines in culture may be used. These include but are not limited

to the hybridoma technique originally developed by Kohler et al., (1975) *Nature*, 256:495-497, the trioma technique, the human B-cell hybridoma technique [Kozbor et al., (1983) *Immunology Today*, 4:72], and the EBV-hybridoma technique to produce human monoclonal antibodies [Cole et al., (1985) in *Monoclonal Antibodies and Cancer Therapy*, pp. 77-96, Alan R. Liss, Inc.]. Immortal, antibody-producing cell lines can be created by techniques other than fusion, such as direct transformation of B lymphocytes with oncogenic DNA, or transfection with Epstein-Barr virus. See, e.g., U.S. Pat. Nos. 4,341,761; 4,399,121; 4,427,783; 4,444,887; 4,451,570; 4,466,917; 4,472,500; 4,491,632; and 4,493,890.

[0263] In an additional embodiment of the invention, monoclonal antibodies can be produced in germ-free animals. According to the invention, swine antibodies may be used and can be obtained by using swine hybridomas or by transfecting B cells with PMC virus in vitro. In fact, according to the invention, techniques developed for the production of "chimeric antibodies" [Morrison et al., (1984) *J. Bacteriol.*, 159:870; Neuberger et al., (1984) *Nature*, 312:604-608; Takeda et al., (1985) *Nature*, 314:452-454] by splicing the genes from a mouse antibody molecule specific for a PMC amino acid sequence together with genes from an antibody molecule of appropriate biological activity can be used; such antibodies are within the scope of this invention. Such chimeric antibodies are preferred for use in therapy of intestinal diseases or disorders, since the antibodies are much less likely than xenogenic antibodies to induce an immune response, in particular an allergic response, themselves.

[0264] According to the invention, techniques described for the production of single chain antibodies (U.S. Pat. No. 4,946,778) can be adapted to produce PMC virus amino acid sequence-specific single chain antibodies. An additional embodiment of the invention utilises the techniques described for the construction of Fab expression libraries [Huse et al., (1989) *Science*, 246:1275-1281] to allow rapid and easy identification of monoclonal Fab fragments with the desired specificity for a PMC virus amino acid sequence, or its derivatives, or analogues.

[0265] Antibody fragments, which contain the idiotype of the antibody molecule, can be generated by known techniques. For example, such fragments include but are not limited to: the F(ab')₂ fragment which can be produced by pepsin digestion of the antibody molecule; the Fab' fragments which can be generated by reducing the disulfide bridges of the F(ab')₂ fragment, and the Fab fragments which can be generated by treating the antibody molecule with papain and a reducing agent.

Screening for Antibodies

[0266] In the production of antibodies, screening for the desired antibody can be accomplished by techniques known in the art, e.g., radioimmunoassay, ELISA, "sandwich" immunoassays, immunoradiometric assays, gel diffusion precipitin reactions, immunodiffusion assays, in situ immunoassays (using colloidal gold, enzyme or radioisotope labels, for example), Western blots, precipitation reactions, agglutination assays (e.g., gel agglutination assays, hemagglutination assays), complement fixation assays, immunofluorescence assays, protein A assays, and immunoelectrophoresis assays, etc.

[0267] In one embodiment, antibody binding is detected by detecting a label on the primary antibody. In another embodiment, the primary antibody is detected by detecting binding

of a secondary antibody or reagent to the primary antibody. In a further embodiment, the secondary antibody is labelled. Many means are known in the art for detecting binding in an immunoassay and are within the scope of the present invention. For example, to select antibodies that recognise a specific epitope of a PMC virus amino acid sequence, one may assay generated hybridomas for a product that binds to a PMC virus amino acid sequence fragment containing such epitope. For selection of an antibody specific to a PMC virus amino acid sequence from a particular species of animal, one can select on the basis of positive binding with PMC virus amino acid sequence expressed by or isolated from cells of that species of animal.

Labelling Antibodies

[0268] Advantageously, the labelling of the anti-immunoglobulin antibodies is achieved by an enzyme selected from among those which are capable of hydrolysing a substrate, which substrate undergoes a modification of its radiation-absorption, at least within a predetermined band of wavelengths. The detection of the substrate, preferably comparatively with respect to a control, then provides a measurement of the likelihood of exposure of an animal to the virus, or of the effective presence, of the disease.

[0269] Thus, preferred methods of immunoenzymatic and also immunofluorescent detections, in particular according to the ELISA technique, are provided. Titrations may be determinations by immunofluorescence or direct or indirect immunoenzymatic determinations. Quantitative titrations of antibodies on the serums studied can be made.

Epitopes Bearing Fragments

[0270] Antibodies according to the present invention may be generated using polypeptide fragments (or molecules, particularly glycoproteins having the same polypeptidic backbone as the polypeptides mentioned hereinabove) bearing an epitope characteristic of a protein or glycoprotein of PMC virus. The polypeptide or molecule may further have N-terminal and C-terminal extremities respectively either free or, independently from each other, covalently bonded to amino acids other than those which are normally associated with them in the larger polypeptides or glycoproteins of the PMC virus, which last mentioned amino acids are then free or belong to another polypeptidic sequence.

Conjugation to Increases Immunogenicity

[0271] Peptide sequences of small size bearing an epitope or immunogenic determinant, (eg those which are readily generated by chemical synthesis), may require coupling or covalent conjugation to a physiologically acceptable and non-toxic carrier molecule in order to increase their in vivo immunogenic character and thus enhance the production of antibodies.

[0272] Particularly, the invention relates to antibodies generated using hybrid polypeptides containing any of the epitope bearing-polypeptides which have been defined more specifically hereinabove, recombined with other polypeptides fragments normally foreign to the PMC virus proteins, having sizes sufficient to provide increased immunogenicity to the epitope-bearing-polypeptide. The foreign polypeptide fragments are preferably immunogenically inert and/or do not interfere with the immunogenic properties of the epitope-bearing-polypeptide.

[0273] Such hybrid polypeptides, which may contain from 5 up to 150, even 250 amino acids, usually consist of the expression products of a vector which contains a nucleic acid sequence encoding said epitope-bearing-polypeptide expressible under the control of a suitable promoter or replicon in a suitable host.

[0274] Said epitope-bearing-polypeptides, particularly those whose N-terminal and C-terminal amino acids are free, may also be generated by chemical synthesis according to techniques well known in the chemistry of proteins.

[0275] Examples of carrier molecules or macromolecular supports which can be used for making the conjugates according to the invention are natural proteins, such as tetanic toxoid, ovalbumin, serum-albumins, hemocyanins, etc. Synthetic macromolecular carriers, for example polysines or poly (D-L-alanine)-poly(L-lysine), can also be used. Other types of macromolecular carriers that can be used, which generally have molecular weights higher than 20,000, are known from the literature.

[0276] The conjugates can be synthesized by known processes such as are described by Frantz and Robertson [*Infection & Immunity*, 33, 193-198 (1981)] and by P. E. Kauffman [*Applied and Environmental Microbiology*], October 1981 Vol. 42, No. 4, pp. 611-614]. For instance, the following coupling agents can be used: glutaric aldehyde, ethyl chloroformate, water-soluble carbodiimides such as (N-ethyl-N'(3-dimethylamino-propyl) carbodiimide, HCl), diisocyanates, bis-diazobenzidine, di- and trichloro-s-triazines, cyanogen bromides and benzaquinone, as well as the coupling agents mentioned in *Scand. J. Immunol.*, 1978, vol. 8, pp. 7-23 (Avrameas, Ternynck, Guesdon).

[0277] Any coupling process can be used for bonding one or several reactive groups of the peptide, on the one hand, and one or several reactive groups of the carrier, on the other hand. Coupling is advantageously achieved between the carboxyl and amine groups carried by the peptide and the carrier in the presence of a coupling agent of the type used in protein synthesis, e.g., 1-ethyl-3-(3-dimethylaminopropyl)-carbodiimide, N-hydroxybenzotriazole, etc. Coupling between amine groups respectively borne by the peptide and the carrier can also be made with glutaraldehyde, for instance, according to the method described by Boquet et al. (1982) *Molec. Immunol.*, 19, 1441-1549, when the carrier is haemocyanin.

[0278] The immunogenicity of epitope-bearing-peptides can also be increased by oligomerisation thereof, for example in the presence of glutaraldehyde or any other suitable coupling agent. In particular, the invention relates to the water soluble immunogenic oligomers thus obtained, comprising particularly from 2 to 10 monomer units.

Vaccines

[0279] The invention also relates to vaccine compositions whose active principle is a polypeptide or fragment thereof of the present invention i.e. the hereinabove disclosed polypeptides of PMC virus, fusion polypeptides or oligopeptides, in association with a suitable pharmaceutically or physiologically acceptable carrier. The present invention further provides immunogenic polypeptides, and more particularly protective polypeptides, for use in the preparation of vaccine compositions against PMC or related syndromes.

[0280] Thus, the present invention provides a vaccine composition comprising a PMC virus polypeptide or fragment thereof.

[0281] Preferably, the polypeptide is an antigenic polypeptide. More preferably, the vaccine further comprises a pharmaceutically acceptable carrier or diluent.

[0282] The invention also provides a vaccine composition comprising a PMC virus nucleotide or fragment thereof that encodes for a PMC virus polypeptide.

[0283] The term "vaccine" as used herein, refers to mean any composition of the invention containing PMC virus peptide or polypeptide or nucleotide sequences coding for PMC virus polypeptides having at least one antigenic determinant which, when administered to a animal, is capable of stimulating an immune response against the antigenic determinant. It will be understood that the term vaccine does not necessarily imply that the composition will provide a complete protective response. Rather a therapeutic effect will be sufficient.

[0284] The phrase "immune response" refers to any cellular process that is produced in the animal following stimulation with an antigen and is directed toward the elimination of the antigen from the animal. The immune response typically is mediated by one or more populations of cells characterized as being lymphocytic and/or phagocytic in nature.

[0285] A vaccine may generate an immune response that blocks the infectivity, either partially or fully, of an infectious agent. The administration of the vaccine of the present invention may be for either a prophylactic or therapeutic purpose. When provided prophylactically, the vaccine is provided in advance of any exposure to PMC virus or in advance of any symptom of any symptoms due to PMC virus infection. The prophylactic administration of the immunogen serves to prevent or attenuate any subsequent infection by PMC virus in a mammal or reduce the severity of infection and/or symptoms. When provided therapeutically, the vaccine is provided at (or shortly after) the onset of the infection or at the onset of any symptom of infection or disease caused by PMC virus. The therapeutic administration of the vaccine serves to attenuate the infection or disease.

[0286] The immune response generated against an introduced PMC virus peptide or polypeptide will be dictated by the amino acid constitution of the antigenic peptide or polypeptide. Such determinants may define either humoral or cell mediated antigenic regions. Without being limited to any particular mode of action, it is contemplated that the immune response generated by the PMC virus peptide or polypeptide will preferably include both humoral and cell mediated immune responses. Where a cell mediated immune response is effected it preferably leads to a T cell cascade, and more specifically by means of a cytotoxic T cell cascade.

[0287] The term "cytotoxic T cell", as used herein, refers to any T lymphocyte expressing the cell surface glycoprotein marker CD8+ that is capable of targeting and lysing a target cell which bears a major histocompatibility class I (MHC Class I) complex on its cell surface and is infected with an intracellular pathogen.

[0288] Preferably, the vaccine composition is developed to generate antibodies against the E0 and E2 envelope glycoproteins and the NS2 and NS3 non-structural proteins.

[0289] The vaccine compositions of the present invention may be used to vaccinate animals and humans against infectious diseases, preferably against PMC. The term "animal" includes: mammals such as farm animals including sheep, goats, pigs, cows, horses, llamas, household pets such as dogs and cats, and primates; birds, such as chickens, geese and ducks; fish; and reptiles such as crocodiles and alligators.

[0290] The vaccine composition according to the invention preferably contains a nucleotide sequence as described above, either as such or as a vaccine strain or in a vector or host organism, or a polypeptide as described above, in an amount effective for producing protection against a pestivirus infection. The vaccine can also be a multipurpose vaccine comprising other immunogens or nucleotides encoding these. The vaccines can furthermore contain conventional carriers, adjuvants, solubilizers, emulsifiers, preservatives etc. The vaccines according to the invention can be prepared by conventional methods.

[0291] Preferably, the active principle is a peptide containing less than 250 amino acid units, preferably less than 150, particularly from 5 to 150 amino acid residues, as deducible from the complete genome of PMC virus.

[0292] The term 'effective amount' refers to an amount of epitope-bearing polypeptide sufficient to induce an immunogenic response in the subject to which it is administered either in a single dose or as part of a series of doses. Preferably, the effective amount is sufficient to effect prophylaxis or treatment, as defined above. The exact amount necessary will vary according to the application. For vaccine applications or for the generation of polyclonal antiserum/antibodies, for example, the effective amount may vary depending on the taxonomic group or species of subject to be treated (e.g. nonhuman primate, primate, etc.), the age and general health and physical condition of the subject, the severity of the condition being treated, the capacity of the subject's immune system to synthesize antibodies, the degree of protection desired, the formulation of the vaccine, the strain of infecting PMC virus, the particular polypeptide selected and its mode of administration, and other relevant factors. It is also believed that effective amounts will be found within a relatively large, non-critical range. An appropriate effective amount can be readily determined using only routine experimentation.

[0293] By way of example, suitable dosages of the vaccine compositions are those which are effective to elicit antibodies *in vivo*, in the host, particularly a porcine host. Suitable doses range from 10 to 500 μg of polypeptide, protein or glycoprotein, for instance 50 to 100 μg . Other preferred ranges of proteins for prophylaxis of PMC are 0.01 to 1000 $\mu\text{g}/\text{dose}$, preferably 0.1 to 100 $\mu\text{g}/\text{dose}$. Several doses may be needed per subject in order to achieve a sufficient immune response and subsequent protection against PMC.

[0294] The immunogenic compositions are conventionally administered using standard procedures, for example, intravenously, subcutaneously, intramuscularly, intraorbitally, ophthalmically, intraventricularly, intracranially, intracapsularly, intraspinaly, intracisternally, intraperitoneally, buccal, rectally, vaginally, intranasally, orally or by aerosol administration.

[0295] Preferably, the immunogenic composition is administered parenterally, typically by injection, for example, subcutaneously or intramuscularly. However, additional formulations suitable for other methods of administration include oral formulations and suppositories or prepared for pulmonary, nasal or other forms of administration. Dosage treatment may be a single dose schedule or a multiple dose schedule.

[0296] The mode of administration of the immunogenic vaccine compositions prepared in accordance with the invention will necessarily depend upon such factors as the stability

of the immunogenic compositions under physiological conditions, the intensity of the immune response required etc.

[0297] The vaccine compositions of the invention may be co-administered with additional immune response enhancers or biological response modifiers including, but not limited to, the cytokines IFN- α , IFN- γ , IL-2, IL-4, IL-6, TNF, or other cytokine-affecting immune cells. In accordance with this aspect of the invention, the PMC virus peptide or polypeptide is administered in combination therapy with a therapeutically active amount of one or more of these cytokines. In addition, conventional antibiotics may be coadministered with the PMC virus peptide or polypeptide. The choice of suitable antibiotics will however be dependent upon the disease in question.

[0298] Parenteral Delivery

[0299] The compounds provided herein can be administered by any parenteral techniques such as subcutaneous, intravenous and intraperitoneal injections. Typically, such vaccines are prepared either as liquid solutions or suspensions; solid forms suitable for solution in, or suspension in, liquid prior to injection may also be prepared. The preparation may also be emulsified, or the protein encapsulated in liposomes. The active immunogenic ingredients are often mixed with excipients and carriers, which are pharmaceutically acceptable and compatible with the active ingredient. Suitable excipients are, for example, water, saline, dextrose, glycerol, ethanol, or the like and combinations thereof.

[0300] In addition, if desired, the vaccine may contain minor amounts of auxiliary substances such as wetting or emulsifying agents, pH buffering agents, and/or adjuvants which enhance the effectiveness of the vaccine.

[0301] Oral Delivery

[0302] Contemplated for use herein are oral solid dosage forms, which are described generally in Martin, Remington's Pharmaceutical Sciences, 18th Ed. (1990 Mack Publishing Co. Easton Pa. 18042) at Chapter 89, which is herein incorporated by reference. Solid dosage forms include tablets, capsules, pills, troches or lozenges, cachets or pellets. Also, liposomal or proteinoid encapsulation may be used to formulate the present compositions (as, for example, proteinoid microspheres reported in U.S. Pat. No. 4,925,673). Liposomal encapsulation may be used and the liposomes may be derivatised with various polymers (E.g., U.S. Pat. No. 5,013, 556). A description of possible solid dosage forms for the therapeutic is given by Marshall, in Modern Pharmaceutics, Chapter 10, Banker and Rhodes ed., (1979), herein incorporated by reference. In general, the formulation will include a PMC virus polypeptide or polynucleotide, and inert ingredients which allow for protection against the stomach environment, and release of the biologically active material in the intestine.

[0303] Also specifically contemplated are oral dosage forms of PMC virus polypeptides or polynucleotides. In this respect the PMC virus polypeptides or polynucleotides may be chemically modified so that oral delivery is efficacious. Generally, the chemical modification contemplated is the attachment of at least one moiety to the protein (or peptide) molecule itself, where said moiety permits (a) inhibition of proteolysis; and (b) uptake into the blood stream from the stomach or intestine. Also desired is the increase in overall stability of the protein and increase in circulation time in the body. Examples of such moieties include: polyethylene glycol, copolymers of ethylene glycol and propylene glycol, carboxymethyl cellulose, dextran, polyvinyl alcohol, polyvi-

nyl pyrrolidone and polyproline. Abuchowski et al., 1981, supra; Newmark et al., *J. Appl. Biochem.*, 4:185-189 (1982). Other polymers that could be used are poly-1,3-dioxolane and poly-1,3,6-tioxocane. Preferred for pharmaceutical usage, as indicated above, are polyethylene glycol moieties.

[0304] For PMC virus polypeptides or polynucleotides the location of release may be the stomach, the small intestine (the duodenum, the jejunum, or the ileum), or the large intestine. One skilled in the art has available formulations that will not dissolve in the stomach, yet will release the material in the duodenum or elsewhere in the intestine. Preferably, the release will avoid the deleterious effects of the stomach environment, either by protection of the complex or by release of the biologically active material beyond the stomach environment, such as in the intestine.

[0305] To ensure full gastric resistance, a coating impermeable to at least pH 5.0 is essential. Examples of the more common inert ingredients that are used as enteric coatings are cellulose acetate trimellitate (CAT), hydroxypropylmethylcellulose phthalate (HPMCP), HPMCP 50, HPMCP 55, polyvinyl acetate phthalate (PVAP), Eudragit L30D, Aquateric, cellulose acetate phthalate (CAP), Eudragit L, Eudragit S, and Shellac. These coatings may be used as mixed films.

[0306] A coating or mixture of coatings can also be used on tablets, which are not intended for protection against the stomach. This can include sugar coatings, or coatings which make the tablet easier to swallow. Capsules may consist of a hard shell (such as gelatin) for delivery of dry therapeutic i.e. powder; for liquid forms, a soft gelatin shell may be used. The shell material of cachets could be thick starch or other edible paper. For pills, lozenges, molded tablets or tablet triturates, moist massing techniques can be used.

[0307] The therapeutic can be included in the formulation as fine multiparticulates in the form of granules or pellets of particle size about 1 mm. The formulation of the material for capsule administration could also be as a powder, lightly compressed plugs or even as tablets. The therapeutic could be prepared by compression.

[0308] Colorants and flavoring agents may all be included. For example, PMC virus polypeptides or polynucleotides may be formulated (such as by liposome or microsphere encapsulation) and then further contained within an edible product, such as a refrigerated beverage containing colorants and flavoring agents.

[0309] One may dilute or increase the volume of the therapeutic with an inert material. These diluents could include carbohydrates, especially mannitol, alpha-lactose, anhydrous lactose, cellulose, sucrose, modified dextrans and starch. Certain inorganic salts may be also be used as fillers including calcium triphosphate, magnesium carbonate and sodium chloride. Some commercially available diluents are Fast-Flo, Emdex, STA-Rx 1500, Emcompress and Avicell.

[0310] Disintegrants may be included in the formulation of the therapeutic into a solid dosage form. Materials used as disintegrants include but are not limited to starch including the commercial disintegrant based on starch, Explotab. Sodium starch glycolate, Amberlite, sodium carboxymethylcellulose, ultramylopectin, sodium alginate, gelatin, orange peel, acid carboxymethyl cellulose, natural sponge and bentonite may all be used. Another form of the disintegrants are the insoluble cationic exchange resins. Powdered gums may be used as disintegrants and as binders and these can include powdered gums such as agar, Karaya or tragacanth. Alginic acid and its sodium salt are also useful as disintegrants.

[0311] Binders may be used to hold the therapeutic agent together to form a hard tablet and include materials from natural products such as acacia, tragacanth, starch and gelatin. Others include methyl cellulose (MC), ethyl cellulose (EC) and carboxymethyl cellulose (CMC). Polyvinyl pyrrolidone (PVP) and hydroxypropylmethyl cellulose (HPMC) could both be used in alcoholic solutions to granulate the therapeutic.

[0312] An antifrictional agent may be included in the formulation of the therapeutic to prevent sticking during the formulation process. Lubricants may be used as a layer between the therapeutic and the die wall and these can include but are not limited to: stearic acid including its magnesium and calcium salts, polytetrafluoroethylene (PTFE), liquid paraffin, vegetable oils and waxes. Soluble lubricants may also be used such as sodium lauryl sulphate, magnesium lauryl sulphate, polyethylene glycol of various molecular weights, and Carbowax 4000 and 6000.

[0313] Glidants that might improve the flow properties of the complex during formulation and to aid rearrangement during compression might be added. The glidants may include starch, talc, pyrogenic silica and hydrated silicoaluminate.

[0314] To aid dissolution of the therapeutic into the aqueous environment, a surfactant might be added as a wetting agent. Surfactants may include anionic detergents such as sodium lauryl sulphate, dioctyl sodium sulphosuccinate and dioctyl sodium sulfonate. Cationic detergents might be used and could include benzalkonium chloride or benzethonium chloride. The list of potential nonionic detergents that could be included in the formulation as surfactants are lauromacrogol 400, polyoxyl 40 stearate, polyoxyethylene hydrogenated castor oil 10, 50 and 60, glycerol monostearate, polysorbate 40, 60, 65 and 80, sucrose fatty acid ester, methyl cellulose and carboxymethyl cellulose. These surfactants could be present in the formulation of the complex either alone or as a mixture in different ratios.

[0315] Additives which potentially enhance uptake of the complex are for instance the fatty acids oleic acid, linoleic acid and linolenic acid.

[0316] Controlled release formulation may be desirable. The complex could be incorporated into an inert matrix which permits release by either diffusion or leaching mechanisms i.e., gums. Slowly degenerating matrices may also be incorporated into the formulation. Another form of a controlled release of this therapeutic is by a method based on the Oros therapeutic system (Alza Corp.), i.e. the drug is enclosed in a semipermeable membrane which allows water to enter and push drug out through a single small opening due to osmotic effects. Some enteric coatings also have a delayed release effect.

[0317] Other coatings may be used for the formulation. These include a variety of sugars which could be applied in a coating pan. The therapeutic agent could also be given in a film-coated tablet; the materials used in this instance are divided into 2 groups. The first are the nonenteric materials and include methyl cellulose, ethyl cellulose, hydroxyethyl cellulose, methylhydroxy-ethyl cellulose, hydroxypropyl cellulose, hydroxypropyl-methyl cellulose, sodium carboxymethyl cellulose, providone and the polyethylene glycols. The second group consists of the enteric materials that are commonly esters of phthalic acid.

[0318] A mix of materials might be used to provide the optimum film coating. Film coating may be carried out in a pan coater or in a fluidized bed or by compression coating.

[0319] Pulmonary Delivery

[0320] Also contemplated herein is pulmonary delivery of vaccine composition. The PMC virus polypeptides or polynucleotides may be delivered to the lungs of an animal while inhaling and traverses across the lung epithelial lining to the blood-stream.

[0321] Contemplated for use in the practice of this invention are a wide range of mechanical devices designed for pulmonary delivery of therapeutic products, including but not limited to nebulizers, metered-dose inhalers, and powder inhalers, all of which are familiar to those skilled in the art.

[0322] Some specific examples of commercially available devices suitable for the practice of this invention are the Ultravent nebulizer, manufactured by Mallinckrodt, Inc., St. Louis, Mo.; the Acorn II nebulizer, manufactured by Marquest Medical Products, Englewood, Colo.; the Ventolin metered dose inhaler, manufactured by Glaxo Inc., Research Triangle Park, N.C.; and the Spinhaler powder inhaler, manufactured by Fisons Corp., Bedford, Mass.

[0323] All such devices require the use of formulations suitable for the dispensing of the complex. Typically, each formulation is specific to the type of device employed and may involve the use of an appropriate propellant material, in addition to the usual diluents, adjuvants and/or carriers useful in therapy. Also, the use of liposomes, microcapsules or microspheres, inclusion complexes, or other types of carriers is contemplated. Chemically modified proteins may also be prepared in different formulations depending on the type of chemical modification or the type of device employed.

[0324] Formulations suitable for use with a nebulizer, either jet or ultrasonic, will typically comprise the complex suspended in water at a concentration of about 0.1 to 25 mg of biologically active protein per ml of solution. The formulation may also include a buffer and a simple sugar (e.g., for protein stabilization and regulation of osmotic pressure). The nebulizer formulation may also contain a surfactant, to reduce or prevent surface induced aggregation of the protein caused by atomization of the solution in forming the aerosol.

[0325] Formulations for use with a metered-dose inhaler device will generally comprise a finely divided powder containing the complex suspended in a propellant with the aid of a surfactant. The propellant may be any conventional material employed for this purpose, such as a chlorofluorocarbon, a hydrochlorofluorocarbon, a hydrofluorocarbon, or a hydrocarbon, including trichlorofluoromethane, dichlorodifluoromethane, dichlorotetrafluoroethanol, and 1,1,1,2-tetrafluoroethane, or combinations thereof. Suitable surfactants include sorbitan trioleate and soya lecithin. Oleic acid may also be useful as a surfactant.

[0326] Formulations for dispensing from a powder inhaler device will comprise a finely divided dry powder containing the complex and may also include a bulking agent, such as lactose, sorbitol, sucrose, or mannitol in amounts which facilitate dispersal of the powder from the device, e.g., 50 to 90% by weight of the formulation. The protein (or derivative) should most advantageously be prepared in particulate form with an average particle size of less than 10 microns, most preferably 0.5 to 5 microns, for most effective delivery to the distal lung.

Nasal Delivery

[0327] Nasal delivery of the vaccine comprising PMC virus polypeptides or polynucleotides is also contemplated. Nasal

delivery allows the passage of the protein to the blood stream directly after administering the therapeutic product to the nose, without the necessity for deposition of the product in the lung. Formulations for nasal delivery include those with dextran or cyclodextran.

Therapeutic Compositions

Polypeptide Based Therapies

[0328] The PMC virus polypeptides according to present invention also can be used as a prophylactic or therapeutic, which may be utilised for the purpose of stimulating humoral and cell mediated responses in animals, such as swine, thereby providing protection against infection with PMC virus. Natural infection with PMC virus induces circulating antibody titres against PMC virus. Therefore, PMC virus amino acid sequence or parts thereof, have the potential to form the basis of a systemically or orally administered prophylactic or therapeutic to provide protection against PMC.

[0329] Thus, the invention provides pharmaceutical compositions comprising a PMC virus polypeptide that enhances the immunocompetence of the host individual and elicits specific immunity against pathogens, preferably PMC virus.

[0330] The therapeutic regimens and pharmaceutical compositions of the invention are described elsewhere in the specification. These compositions are believed to have the capacity to prevent the onset and progression of infectious disease such as PMC.

[0331] Preferably the compositions are combined with a pharmaceutically acceptable carrier or diluent to produce a pharmaceutical composition (which may be for human or animal use). Compositions of the invention comprising PMC virus polypeptides may also be combined with suitable components to obtain vaccine compositions. Accordingly, in one embodiment the present invention provides a PMC virus amino acid sequence or fragments thereof described herein in a therapeutically effective amount admixed with a pharmaceutically acceptable carrier, diluent, or excipient.

[0332] The phrase "therapeutically effective amount" is used herein to mean an amount sufficient to reduce by at least about 15%, preferably by at least 50%, more preferably by at least 90%, and most preferably prevent, a clinically significant deficit in the activity, function and response of the animal host. Alternatively, a therapeutically effective amount is sufficient to cause an improvement in a clinically significant condition in the animal host or to stimulate by at least about 15%, preferably by at least 50%, more preferably by at least 90%, and most preferably completely, a animal's immune system, causing it to generate an immunological memory against the antigenic determinant.

[0333] The phrase "pharmaceutically acceptable" refers to molecular entities and compositions that are physiologically tolerable and do not typically produce an allergic or similarly untoward reaction, such as gastric upset and the like, when administered to an animal. The term "carrier" refers to a diluent, adjuvant, excipient, or vehicle with which the compound is administered. Such pharmaceutical carriers can be sterile liquids, such as water and oils, including those of petroleum, animal, vegetable or synthetic origin, such as peanut oil, soybean oil, mineral oil, sesame oil and the like. Water or saline solutions and aqueous dextrose and glycerol solutions are preferably employed as carriers, particularly for injectable solutions. Suitable pharmaceutical carriers are

described in Martin, *Remington's Pharmaceutical Sciences*, 18th Ed., Mack Publishing Co., Easton, Pa., (1990).

[0334] In a more specific form of the invention there are provided pharmaceutical compositions comprising therapeutically effective amounts of PMC virus amino acid sequence or an analogue, fragment or derivative product thereof together with pharmaceutically acceptable diluents, preservatives, solubilizers, emulsifiers, adjuvants and/or carriers. Such compositions include diluents of various buffer content (e.g., Tris-HCl, acetate, phosphate), pH and ionic strength and additives such as detergents and solubilizing agents (e.g., Tween 80, Polysorbate 80), anti-oxidants (e.g., ascorbic acid, sodium metabisulfite), preservatives (e.g., Thimersol, benzyl alcohol) and bulking substances (e.g., lactose, mannitol). The material may be incorporated into particulate preparations of polymeric compounds such as polylactic acid, polyglycolic acid, etc. or into liposomes. Hylauronic acid may also be used. Such compositions may influence the physical state, stability, rate of in vivo release, and rate of in vivo clearance of the present proteins and derivatives. See, e.g., Martin, *Remington's Pharmaceutical Sciences*, 18th Ed. 1990, Mack Publishing Co., Easton, Pa., pp 1435-1712 that are herein incorporated by reference. The compositions may be prepared in liquid form, or may be in dried powder, such as lyophilised form.

[0335] The present invention also provides for the use of PMC virus amino acid sequences according to the invention, for manufacture of a medicament for modulation of a disease associated with PMC virus.

Antibody Based Therapeutics

[0336] The present invention also provides therapeutic compositions comprising antibodies prepared against the polypeptides of the invention.

[0337] The antibodies can be used directly as antiviral agents. To prepare antibodies, a host animal is immunized using one or more PMC virus proteins bound to a carrier as described above for vaccines. The host serum or plasma is collected following an appropriate time interval to provide a composition comprising antibodies reactive with the protein (s) of the virus particle. The gamma globulin fraction or the IgG antibodies can be obtained, for example, by use of saturated ammonium sulfate or DEAE Sephadex, or other techniques known to those skilled in the art. The antibodies are substantially free of many of the adverse side effects which may be associated with other anti-viral agents such as drugs.

[0338] Such therapeutic antibody compositions may additionally contain one or more of the additional agents described above in relation to polypeptide therapeutics.

[0339] The present invention provides for the use of antibodies against the PMC virus according to the invention, for manufacture of a medicament for modulation of a disease associated with PMC virus.

Polynucleotide Base Therapy

[0340] The present invention further provides therapeutic compositions comprising PMC virus nucleic acid sequences as well as antisense and ribozyme polynucleotide sequences hybridisable to a polynucleotide sequence encoding a PMC virus amino acid sequence according to the invention.

[0341] Polynucleotide sequences encoding antisense constructs or ribozymes for use in therapeutic methods are desirably administered directly as a naked nucleic acid construct.

Uptake of naked nucleic acid constructs is enhanced by several known transfection techniques, for example those including the use of transfection agents. Example of these agents include cationic agents (for example calcium phosphate and DEAE-dextran) and lipofectants (for example Lipofectam™ and Transfectam™). Typically, nucleic acid constructs are mixed with the transfection agent to produce a composition.

[0342] Alternatively the antisense construct or ribozymes may be combined with a pharmaceutically acceptable carrier or diluent to produce a pharmaceutical composition. Suitable carriers and diluents include isotonic saline solutions, for example phosphate-buffered saline. The composition may be formulated for parenteral, intramuscular, intravenous, subcutaneous, intraocular, oral or transdermal administration.

[0343] Also addressed by the present invention is the use of polynucleotide sequences of the invention, as well as antisense and ribozyme polynucleotide sequences hybridisable to a polynucleotide sequence encoding a PMC virus amino acid sequence according to the invention, for manufacture of a medicament for modulation of a disease associated with PMC virus.

Administration of Therapeutic Compositions

[0344] It will be appreciated that therapeutic compositions provided accordingly to the invention may be administered by any means known in the art. Therapeutic compositions may be for administration by injection, or prepared for oral, pulmonary, nasal or other forms of administration. The mode of administration of the therapeutic compositions prepared in accordance with the invention will necessarily depend upon such factors as the stability of the complex under physiological conditions, the intensity of the immune response required etc.

[0345] Preferably, the pharmaceutical compositions for administration are administered by injection, orally, or by the pulmonary, or nasal route.

[0346] Preferably, the therapeutic compositions are administered using standard procedures, for example, intravenously, subcutaneously, intramuscularly, intraorbitally, ophthalmically, intraventricularly, intracranially, intracapsularly, intraspinally, intracisternally, intraperitoneally, buccal, rectally, vaginally, intranasally, orally or by aerosol administration.

[0347] The PMC virus amino acid sequence or antibodies derived there from, or polynucleotide sequences are more preferably delivered by intravenous, intra-arterial, intraperitoneal, intramuscular, or subcutaneous routes of administration. Alternatively, the PMC virus amino acid sequence or antibodies derived there from, properly formulated, can be administered by nasal or oral administration. The routes of administration described are intended only as a guide since a skilled practitioner will be able to determine readily the optimum route of administration and any dosage for any particular animal and condition.

[0348] The present invention further provides a method of inducing a protective immune response in an animal or human against a PMC virus comprising the steps of:

[0349] a) administering to said animal or human an effective amount of a composition of the invention.

[0350] The present invention also provides methods for enhancing an animal's immunocompetence and the activity of its immune effector cells against a PMC virus comprising the step of:

[0351] a) administering a composition comprising a therapeutically effective amount of a PMC virus peptide or polypeptide.

Live Vector Delivery Agent

[0352] In another aspect of the invention, the PMC virus may be used as a live vector for delivery of recombinant antigens.

[0353] Thus, the present invention provides a live vector comprising the PMC virus and a heterologous polynucleotide.

[0354] Preferably, the heterologous polynucleotide is operably linked to the polynucleotide sequence of the PMC virus, such that expression of the polynucleotide sequence of the PMC virus also leads to expression of the heterologous polynucleotide sequence.

[0355] Furthermore, the PMC virus may have one or more sections of autologous polynucleotide sequence removed. Removal of such sequence may preferably render the live virus attenuated in pathogenicity in a host subject.

[0356] For example, the PMC virus may be used as a delivery vector to deliver gene sequences that encode a protein from a second infective agent into a subject to be vaccinated against the second infective agent. The second infective agent may be a virus (such as classical swine fever virus), a bacteria, a parasite etc.

[0357] Alternatively, the PMC virus may be used as a delivery vector to deliver antigens from some other source. For example, a PMC virus vector may be used to deliver antigenic proteins to a subject to stimulate the subject to make antibodies against the antigenic proteins that may be collected for purposes such as use in diagnostic kits etc.

Drug Screening Assays

[0358] The present invention also provides assays that are suitable for identifying substances such as drugs, agents or ligands that bind to PMC virus amino acid sequences. In addition, assays are provided that are suitable for identifying substances that interfere with PMC virus amino acid sequences. Assays are also provided that test the effects of candidate substances identified in preliminary *in vitro* assays on intact cells in whole cell assays.

[0359] Thus, the present invention provides a method of screening for drugs comprising the steps of:

[0360] a) contacting an agent with a PMC virus amino acid sequence or fragment thereof and

[0361] b) assaying for the presence of a complex between the agent and the PMC virus amino acid sequence or fragment.

[0362] The present invention also provides a method of screening for ligands of the proteins of the PMC virus comprising the steps of:

[0363] a) contacting a ligand with a PMC virus amino acid sequence or fragment thereof and

[0364] b) assaying for the presence of a complex between the PMC virus amino acid sequence or fragment and a ligand.

[0365] One type of assay for identifying substances such as drugs, agents or ligands that bind to PMC virus amino acid sequences involves contacting a PMC virus amino acid sequence, which is immobilised on a solid support, with a non-immobilised candidate substance and determining whether and/or to what extent the PMC virus amino acid

sequences and candidate substance bind to each other. Alternatively, the candidate substance may be immobilised and the PMC virus amino acid sequence non-immobilised.

[0366] In a preferred assay method, the PMC virus amino acid sequence is immobilised on beads such as agarose beads. Typically this is achieved by expressing the component as a GST-fusion protein in bacteria, yeast or higher eukaryotic cell lines and purifying the GST-fusion protein from crude cell extracts using glutathione-agarose beads. The binding of the candidate substance to the immobilised PMC virus amino acid sequence is then determined. This type of assay is known in the art as a GST pull-down assay. Again, the candidate substance may be immobilised and the PMC virus amino acid sequence non-immobilised.

[0367] It is also possible to perform this type of assay using different affinity purification systems for immobilising one of the components, for example Ni-NTA agarose and hexahistidine-tagged components.

[0368] Binding of the PMC virus amino acid sequence to the candidate substance may be determined by a variety of methods well known in the art. For example, the non-immobilised component may be labelled (with for example, a radioactive label, an epitope tag or an enzyme-antibody conjugate). Alternatively, binding may be determined by immunological detection techniques. For example, the reaction mixture can be Western blotted and the blot probed with an antibody that detects the non-immobilised component. ELISA techniques may also be used.

[0369] Candidate substances are typically added to a final concentration of from 1 to 1000 nmol/ml, more preferably from 1 to 100 nmol/ml. In the case of antibodies, the final concentration used is typically from 100 to 500 µg/ml, more preferably from 200 to 300 µg/ml.

[0370] In a competitive binding assay the PMC virus amino acid sequence or fragment is typically labelled. Free PMC virus amino acid sequence or fragment is separated from that present in a protein:protein complex, and the amount of free (i.e., uncomplexed) label is a measure of the binding of the agent being tested to the PMC virus amino acid sequence or its interference with PMC virus amino acid sequence:ligand binding, respectively.

[0371] Another technique for drug screening provides high throughput screening for compounds having suitable binding affinity to the PMC virus amino acid sequence and is described in detail in PCT Application WO 84/03564, published on Sep. 13, 1984. Briefly stated, large numbers of different small peptide test compounds are synthesised on a solid substrate, such as plastic pins or some other surface. The peptide test compounds are reacted with PMC virus amino acid sequence and washed. Bound PMC virus amino acid sequence is then detected by methods well known in the art.

[0372] This invention also contemplates the use of competitive drug screening assays in which antibodies capable of specifically binding the PMC virus amino acid sequence compete with a test compound for binding to the PMC virus amino acid sequence or fragments thereof. In this manner, the antibodies can be used to detect the presence of any peptide that shares one or more antigenic determinants of the PMC virus amino acid sequence.

Kits

[0373] In a further embodiment of this invention, kits may be prepared to determine the presence or absence of PMC virus in suspected infected animals and/or to quantitatively

measure PMC infection. In accordance with the testing techniques discussed above, one class of such kits will contain at least the labelled PMC virus amino acid sequence or its binding partner, for instance an antibody specific thereto, and directions depending upon the method selected, e.g., "competitive," "sandwich," "DASP" and the like. The kits may also contain peripheral reagents such as buffers, stabilizers, etc.

[0374] Thus, kits for PMC virus serum immunoassay may be either (a) a sandwich type immunoassay, employing a first anti-PMC virus antibody as capture or detector antibody and a second anti-PMC virus antibody as a detector or capture antibody to complement the first anti-PMC virus antibody, or (b) a competitive type immunoassay, employing a anti-PMC virus antibody with a labelled PMC virus antigen or a PMC virus antigen attached to a solid phase.

[0375] Accordingly, a test kit may be prepared for the demonstration of the presence of PMC virus comprising:

[0376] (a) a predetermined amount of at least one labelled immunochemically reactive component obtained by the direct or indirect attachment of the present PMC virus amino acid sequence or a specific binding partner thereto, to a detectable label;

[0377] (b) other reagents; and

[0378] (c) directions for use of said kit.

[0379] More specifically, the diagnostic test kit may comprise:

[0380] (a) a known amount of the PMC virus amino acid sequence as described above (or a binding partner) generally bound to a solid phase to form an immunosorbent, or in the alternative, bound to a suitable tag, or there are a plural of such end products, etc;

[0381] (b) if necessary, other reagents; and

[0382] (c) directions for use of said test kit.

[0383] In a further variation, the test kit may be prepared and used for the purposes stated above, which operates according to a predetermined protocol (e.g. "competitive," "sandwich," "double antibody," etc.), and comprises:

[0384] (a) a labelled component which has been obtained by coupling the PMC virus amino acid sequence to a detectable label;

[0385] (b) one or more additional immunochemical reagents of which at least one reagent is a ligand or an immobilized ligand, which ligand is selected from the group consisting of:

[0386] (i) a ligand capable of binding with the labelled component (a);

[0387] (ii) a ligand capable of binding with a binding partner of the labelled component (a);

[0388] (iii) a ligand capable of binding with at least one of the component(s) to be determined; or

[0389] (iv) a ligand capable of binding with at least one of the binding partners of at least one of the component(s) to be determined; and

[0390] (c) directions for the performance of a protocol for the detection and/or determination of one or more components of an immunochemical reaction between the PMC virus amino acid sequence and a specific binding partner thereto.

Kits to Detect Antibodies

[0391] The invention also provides diagnostic kits for the in vitro detection of antibodies against the PMC virus, which kits comprise any of the polypeptides identified herein and all

the biological and chemical reagents, as well as equipment, necessary for performing diagnostic assays.

[0392] Accordingly, the invention provides a kit for demonstrating the presence of PMC virus comprising:

[0393] (a) a predetermined amount of at least one labelled antibody to the PMC virus;

[0394] (b) other reagents; and

[0395] (c) directions for use of said kit.

[0396] Preferably, the polypeptide used in the kit is an antigenic or epitope bearing polypeptide. Most preferably, the polypeptide is a polypeptide encoding, but not exclusively limited to, the E0, E2, NS2 or NS3 protein.

[0397] Preferred kits comprise all reagents required for carrying out ELISA assays. Thus preferred kits will include, in addition to any of said polypeptides, suitable buffers and anti-species immunoglobulins, which anti-species immunoglobulins are labelled either by an immunofluorescent molecule or by an enzyme. In the last instance, preferred kits also comprise a substrate hydrolysable by the enzyme and providing a signal, particularly modified absorption of a radiation, at least in a determined wavelength, which signal is then indicative of the presence of antibody in the biological fluid to be assayed with said kit. Kits may also include labelled monoclonal or polyclonal antibodies that are directed against PMC virus epitopes and these labelled antibodies may be used to block or compete with antibodies from the test specimen. If the activity of the labelled antibody is blocked, no or a reduced reaction will occur and it can be deduced that the test specimen contains antibodies to PMC virus.

[0398] The present invention also relates to a diagnostic kit for use in detecting the presence of PMC virus antibodies, said kit comprising at least one peptide as defined above, with said peptide being preferably bound to a solid support.

[0399] The peptide, for example, can be attached to a variety of different solid supports to enable the washing away of unreacted reagents during the course of using the kit. These include: microwells, coated test tubes, coated magnetic particles, wands or sticks, and membranes (nitrocellulose and others).

[0400] Preferably, the peptides are attached to specific locations on the solid support. More preferably, the solid support is a membrane strip and said peptides are coupled to the membrane in the form of parallel lines. Preferably, the peptide used in the kit is an antigenic or epitope bearing peptide.

[0401] The PMC virus antigens of the present invention will typically be packaged in the form of a kit for use in these immunoassays. The kit will normally contain, in separate containers, the PMC virus antigen, control antibody formulations (positive and/or negative), labelled antibody when the assay format requires the same and signal generating reagents (e.g. enzyme substrate) if the label does not generate a signal directly. The PMC virus antigen may be already bound to a solid support or may be provided separately, with reagents for binding it to the solid support. Instructions (e.g. written, tape, CD-ROM, etc.) for carrying out the assay usually will be included in the kit.

[0402] Immunoassays that utilize PMC virus antigens are useful in screening samples (such as blood, serum, plasma, milk, body fluids) to detect if the subject from which the tissue was derived has been exposed to or infected with PMC virus.

[0403] The solid support used in the kits of the present invention can include polymeric or glass beads, nitrocellulose, microparticles, microwells of a reaction tray, test tubes and magnetic beads.

[0404] The signal generating compound can include an enzyme, a luminescent compound, a fluorophore such as fluorescein, a time-resolved fluorescent probe such as a europium chelate, a chromogen, a radioactive element, a chemiluminescent compound such as an acridinium ester or particles such as colloidal gold, plain latex, or dyed latex. Examples of enzymes include alkaline phosphatase, horseradish peroxidase and beta-galactosidase.

Kits to Detect Polypeptides and Antigens

[0405] The present invention further provides a diagnostic kit for use in detecting the presence of PMC virus proteins.

[0406] Accordingly, the invention provides a kit for demonstrating the presence of PMC virus comprising:

[0407] (a) a predetermined amount of at least one labelled polypeptide derived from the PMC virus;

[0408] (b) other reagents; and

[0409] (c) directions for use of said kit.

[0410] Preferably, said antibody is bound to a solid support. The antibody can be attached to a variety of different solid supports to enable the washing away of unreacted reagents during the course of using the kit. These include: microwells, coated test tubes, coated magnetic particles, wands or sticks, and membranes (nitrocellulose and others). Preferably, the antibodies are attached to specific locations on a solid substrate.

[0411] The anti-PMC virus antibody can be attached to the solid support by a variety of means such as passive adsorption, covalent coupling, or by using a solid phase pre-coated with a secondary binder such as protein A, protein G, a secondary antibody specific for the primary antibody, avidin, or an antibody specific for a particular ligand (i.e.: biotin, dinitrophenol, fluorescein, and others). In the case of avidin or any of the ligand specific antibodies, it is necessary to covalently attach the ligand to the anti-PMC virus antibody.

[0412] For example, ELISA kits may be used to detect the presence of antigens to PMC virus in a sample to demonstrate that an animal is suffering from PMC or is, for example, a non-symptomatic carrier of the virus.

[0413] Preferably, the protein to be detected using the present kit is an antigen or an epitope bearing region of a PMC virus protein. Most preferably, the antibody binds to the E0, E2, NS2 or NS3 protein of PMC.

Kits to Detect Nucleic Acid Sequences

[0414] The invention also provides kits for screening animals suspected of being infected with PMC virus, or to confirm that an animal is infected with PMC virus, by detecting PMC virus nucleic acid sequences.

[0415] Accordingly, the invention provides a kit for demonstrating the presence of PMC virus comprising:

[0416] (a) a predetermined amount of at least one labelled nucleic acid sequence derived from the PMC virus;

[0417] (b) other reagents; and

[0418] (c) directions for use of said kit.

[0419] For example, the polynucleotide sequence may be one or more primers, such as those exemplified above, and the

instructions for use may be instructions to perform PCR on RNA or DNA extracted from a tissue sample from a subject.

Vectors, Host Cells Etc

Vectors

[0420] The present invention also provides a recombinant expression vector comprising a PMC virus nucleic acid sequence or a part thereof as defined above, operably linked to prokaryotic, eukaryotic or viral transcription and translation control elements.

[0421] The invention further relates to the hosts (prokaryotic or eukaryotic cells) which are transformed by the above mentioned vectors and recombinants and which are capable of expressing said RNA and/or DNA fragments.

[0422] According to another embodiment the present invention provides methods for preparing a PMC virus amino acid sequence, comprising the steps of:

[0423] (a) culturing a host cell containing a vector as described above under conditions that provide for expression of the PMC virus amino acid sequence; and

[0424] (b) recovering the expressed PMC virus sequence.

[0425] This procedure can also be accompanied by the step of:

[0426] (c) subjecting the amino acid sequence to protein purification.

[0427] The present invention also relates to a method for the production of a recombinant PMC virus polypeptide, comprising the steps of:

[0428] a) transforming an appropriate cellular host with a recombinant vector, in which a PMC virus polynucleotide sequence or a part thereof has been inserted under the control of appropriate regulatory elements,

[0429] b) culturing said transformed cellular host under conditions enabling the expression of said insert, and,

[0430] c) harvesting said polypeptide.

[0431] Vectors provided by the present invention will typically comprise a PMC virus polynucleotide sequence encoding the desired amino acid sequence and preferably transcription and translational regulatory sequences operably linked to the amino acid encoding sequence so as to allow for the expression of the antigenic polypeptide in the cell. Preferably, the vector will include appropriate prokaryotic, eukaryotic or viral promoter sequence followed by the PMC virus nucleotide sequences as defined above. The recombinant vector of the present invention may preferably allow the expression of any one of the PMC virus polypeptides as defined above in a prokaryotic, or eukaryotic host or in living mammals when injected as naked RNA or DNA.

[0432] The vector may comprise a plasmid, a cosmid, a phage, or a virus or a transgenic animal. Particularly useful for vaccine development may be BCG or adenoviral vectors, as well as avipox recombinant viruses. Examples of such expression vectors are described in Sambrook et al., (1989) supra or Ausubel et al., (2001) supra. Many useful vectors are known in the art and may be obtained from such vendors as Stratagene, New England Biolabs, Promega Biotech, and others.

[0433] It may be desirable to use regulatory control sequences that allow for inducible expression of the antigenic polypeptide, for example in response to the administration of an exogenous molecule. Alternatively, temporal control of

expression of the antigenic polypeptide may occur by only introducing the polynucleotide into the cell when it is desired to express the polypeptide.

[0434] It may also be convenient to include an N-terminal secretion signal so that the antigenic polypeptide is secreted into the cell medium.

[0435] Expression vectors may also include, for example, an origin of replication or autonomously replicating sequence and expression control sequences, a promoter, an enhancer and necessary processing information sites, such as ribosome-binding sites, RNA splice sites, polyadenylation sites, transcriptional terminator sequences, and mRNA stabilising sequences. Secretion signals may also be included where appropriate, from secreted polypeptides of the same or related species, which allow the protein to cross and/or lodge in cell membranes, and thus attain its functional topology, or to be secreted from the cell. Such vectors may be prepared by means of standard recombinant techniques well known in the art and discussed, for example, in Sambrook et al., (1989) or Ausubel et al., (2001).

[0436] An appropriate promoter and other necessary vector sequences will be selected so as to be functional in the host, and may include, when appropriate, those naturally associated with outer membrane lipoprotein genes.

[0437] Promoters such as the *trp*, *lac* and phage promoters, tRNA promoters and glycolytic enzyme promoters may be used in prokaryotic hosts. Useful yeast promoters include promoter regions for metallothionein, 3-phosphoglycerate kinase or other glycolytic enzymes such as enolase or glyceraldehyde-3-phosphate dehydrogenase, enzymes responsible for maltose and galactose utilization, and others. Vectors and promoters suitable for use in yeast expression are further described in Hitzeman et al., EP 73,675A. Appropriate non-native mammalian promoters might include the early and late promoters from SV40 or promoters derived from murine Moloney leukaemia virus, avian sarcoma viruses, adenovirus II, bovine papilloma virus or polyoma. In addition, the construct may be joined to an amplifiable gene (e.g., DHFR) so that multiple copies of the gene may be made.

[0438] While such expression vectors may replicate autonomously, they may also replicate by being inserted into the genome of the host cell, by methods well known in the art.

[0439] Expression and cloning vectors will likely contain a selectable marker, a gene encoding a protein necessary for survival or growth of a host cell transformed with the vector. The presence of this gene ensures growth of only those host cells that express the inserts. Typical selection genes encode proteins that a) confer resistance to antibiotics or other toxic substances, e.g. ampicillin, neomycin, methotrexate, etc.; b) complement auxotrophic deficiencies, or c) supply critical nutrients not available from complex media, e.g., the gene encoding D-alanine racemase for *Bacilli*. The choice of the proper selectable marker will depend on the host cell, and appropriate markers for different hosts are well known in the art.

[0440] Vectors containing PMC virus polynucleotide sequences can be transcribed *in vitro* and the resulting RNA introduced into the host cell by well-known methods, e.g., by injection, or the vectors can be introduced directly into host cells by methods well known in the art, which vary depending on the type of cellular host, including electroporation; transfection employing calcium chloride, rubidium chloride, calcium phosphate, DEAE-dextran, or other substances; microprojectile bombardment; lipofection; infection (where the

vector is an infectious agent, such as a retroviral genome); and other methods. The introduction of PMC virus polynucleotide sequences into the host cell may be achieved by any method known in the art, including, *inter alia*, those described above.

[0441] In a preferred embodiment, the PMC virus polynucleotide is part of a viral vector, such as a baculovirus vector, or infectious virus, such as a baculovirus. This provides a convenient system since not only can recombinant viral stocks can be maintained and stored until ready for use. Desirably, the nucleotide sequence encoding the antigenic peptide or polypeptides is inserted into a recombinant baculovirus that has been genetically engineered to produce antigenic peptide or polypeptides, for instance, by following the methods of Smith et al (1983) *Mol Cell Biol* 12: 2156-2165. A number of viral transfer vectors allow more than one polynucleotide sequence encoding a polypeptide to be inserted into the same vector so that they can be co-expressed by the same recombinant virus.

Host Cells

[0442] To produce a cell capable of expressing PMC virus amino acid sequences, preferably polynucleotide sequences of the invention are incorporated into a recombinant vector, which is then introduced into a host prokaryotic or eukaryotic cell.

[0443] The invention also provides host cells transformed or transfected with a PMC virus polynucleotide sequence. Preferred host cells include yeast, filamentous fungi, plant cells, insect, amphibian, avian species, bacteria, mammalian cells, and human cells in tissue culture. Illustratively, such host cells are selected from the group consisting of *E. coli*, *Pseudomonas*, *Bacillus*, *Streptomyces*, yeast, CHO, R1.1, B-W, L-M, COS1, COS 7, BSC1, BSC40, BMT10, and Sf9 cells.

[0444] Large quantities of PMC virus polynucleotide sequence of the invention may be prepared by expressing PMC virus polynucleotide sequences or portions thereof in vectors or other expression vehicles in compatible prokaryotic or eukaryotic host cells. The most commonly used prokaryotic hosts are strains of *Escherichia coli*, although other prokaryotes, such as *Bacillus subtilis* or *Pseudomonas* may also be used. Examples of commonly used mammalian host cell lines are VERO and HeLa cells, Chinese hamster ovary (CHO) cells, and WI38, BHK, and COS cell lines, although it will be appreciated by the skilled practitioner that other cell lines may be appropriate.

[0445] Also provided are mammalian cells containing a PMC virus polynucleotide sequences modified *in vitro* to permit higher expression of PMC virus amino acid sequence by means of a homologous recombinational event consisting of inserting an expression regulatory sequence in functional proximity to the PMC virus amino acid sequence encoding sequence.

[0446] The invention is not limited to the production of one antigenic polypeptide at a time in the host cell. Multiple polynucleotides encoding different antigenic polypeptides of interest may be introduced into the same host cell. The polynucleotides may be part of the same nucleic acid molecule or separate nucleic acid molecules.

General

[0447] Those skilled in the art will appreciate that the invention described herein is susceptible to variations and

modifications other than those specifically described. It is to be understood that the invention includes all such variations and modifications. The invention also includes all of the steps, features, compositions and compounds referred to or indicated in the specification, individually or collectively and any and all combinations or any two or more of the steps or features.

[0448] The present invention is not to be limited in scope by the specific embodiments described herein, which are intended for the purpose of exemplification only. Functionally equivalent products, compositions and methods are clearly within the scope of the invention as described herein.

[0449] The entire disclosures of all publications (including patents, patent applications, journal articles, laboratory manuals, books, or other documents) cited herein are hereby incorporated by reference. No admission is made that any of the references constitute prior art or are part of the common general knowledge of those working in the field to which this invention relates.

[0450] As used herein the term “derived” and “derived from” shall be taken to indicate that a specific integer may be obtained from a particular source albeit not necessarily directly from that source.

[0451] Throughout this specification, unless the context requires otherwise, the word “comprise”, or variations such as “comprises” or “comprising”, will be understood to imply the inclusion of a stated integer or group of integers but not the exclusion of any other integer or group of integers.

[0452] Other definitions for selected terms used herein may be found within the detailed description of the invention and apply throughout. Unless otherwise defined, all other scientific and technical terms used herein have the same meaning as commonly understood to one of ordinary skill in the art to which the invention belongs.

EXAMPLES

[0453] The following examples serve to more fully describe the manner of using the above-described invention, as well as to set forth the best modes contemplated for carrying out various aspects of the invention. It is understood that these methods in no way serve to limit the true scope of this invention, but rather are presented for illustrative purposes.

Example 1

Sample Preparation

[0454] Tissue samples were extracted and prepared using a method whose main basis was derived from Allander et al (2001) “A virus discovery method incorporating DNase treatment and its application to the identification of two bovine parvovirus species.” Proc Natl Acad Sci USA. 98(20): 11609-14, with some modifications to improve the efficiency from Baugh et al (2001) “Quantitative analysis of mRNA amplification by in vitro transcription.” Nucleic Acids Res. 29(5): E29. However, the methods were modified to improve efficiency.

1. Preparation of Serum Samples:

[0455] a) Obtain at least 240 μ L of supernatant from a tissue homogenate or serum and divide into 2 \times 120 μ L

[0456] b) To each 120 μ L of sample add 240 μ L of PBS or H₂O (or take 50 μ L sera+100 μ L PBS)

[0457] c) Filter diluted sample through two separate 0.2 μ m filters by centrifuging at 2000 \times g (wash top of filter and keep at -20° C.)

[0458] d) Add 25 μ L DNase I (250 U) to each tube of filtered sample and incubate at 37° C. for 2 hr

[0459] e) Add 1 μ L of RNase Cocktail (500 U RNase A, 20000 U RNase T1) to each tube and incubate at RT for 1 hr.

[0460] f) Take 1 tube of treated sample (360 μ L) for RNA extraction and one tube for DNA extraction (add 500 μ L DNAeasy AL+50 μ L proteinase K etc and elute in 50 μ L water).

2. RNA Extraction:

[0461] a) Divide sample into 90 μ L lots and add 600 μ L RLT, ie 4 \times 690 μ L

[0462] b) Homogenize by passing through 21 G syringe at least 5 \times

[0463] c) Add 690 μ L of 70% ethanol to each tube of sample and mix by pipetting

[0464] d) Apply 700 μ L of sample to column at a time and centrifuge for 15 sec at 10,000 rpm. Place flow through waste in a 5 ml container and keep at -80° C.

[0465] e) Add 700 μ L of buffer RW1 to the column and centrifuge for 15 sec at 10,000 rpm. Discard flow through material and collection tube.

[0466] f) Transfer column to a new tube and add 500 μ L of RPE centrifuge for 15 sec at 10,000 rpm, discard flow through material

[0467] g) Repeat step (f) using same tube but centrifuge for 2 min at 10,000 rpm.

[0468] h) Transfer column to a new tube and centrifuge for 1 min at 10,000 rpm.

[0469] i) Elute the RNA in 20 μ L of RNase free water, let the water sit on the column for 1 minute before centrifuging. Reuse the eluate and centrifuge for 1 min at 10,000 rpm to collect any left over RNA on column.

[0470] j) Store RNA at -80° C. until needed.

3. DNA Extraction:

[0471] a) To 360 μ L of sample add 36 μ L of proteinase K and 360 μ L of buffer AL, mix by vortex, incubate at 70° C. for 10 minutes.

[0472] b) Add 360 μ L of 100% ethanol, mix by vortexing

[0473] c) Pipette mixture from step (b) into DNAeasy column and centrifuge at 8,000 rpm for 1 minute. Place flow-through into a tube and store at -80° C.

[0474] d) Place column in a new tube and add 500 μ L of AW1 spin at 8,000 rpm for 1 minute. Discard flow through and tube.

[0475] e) Place column in a new tube and add 500 μ L of AW2 and spin at 13,000 rpm for 3 minutes. Discard flow through and spin for another 1 minute and discard flow through and tube.

[0476] f) Place column in a new tube, add 50 μ L of water and let sit for 1 minute. Spin at 8,000 rpm for 1 min and collect eluate. Reapply the 50 μ L eluate and spin again.

[0477] g) Store DNA at -80° C. until needed.

RNA Sequence-independent Single Primer Amplification (SISPA) for Double Stranded RNA Viruses

[0478] The SISPA method employed was developed from that of Baugh et al and Allander et al, to maximise yield and product length while minimising template-independent side

reactions. However, the present method is applied to low yield viral RNA, not total mRNA and a melting step has been added.

4. First Strand cDNA Synthesis

[0479] a) Mix Together the Following:

[0480] 1 ul random hexamers (10 μmol)

[0481] 8 ul-9 ul RNA (in H₂O)

[0482] b) Mix, heat 90° C. 3 minutes, spin and put on ice

[0483] c) On ice add:

1 st strand buffer	4 ul
0.1M dTT	2 ul
5 mM dNTP	2 ul
SSIII (400 U)	1 ul
T4gene32	1 ul

1st strand buffer: 50 mM Tris-HCl (pH 8.3), 75 mM KCl, 3 mM MgCl₂

[0484] d) Mix, spin and heat at 50° C. for 30 minutes

[0485] e) Add another 1 ul of SSIII and leave for another 30 min at 50° C.

[0486] f) Heat inactivate at 70° C. for 10 minutes and then place on ice

5. Second Strand cDNA Synthesis

[0487] a) On ice mix:

H ₂ O	87 ul
5X 2 nd strand buffer	30 ul
5 mM dNTPs	6 ul
DNA polymerase (40 U)	4 ul
<i>E. coli</i> DNA ligase (10 U)	1 ul
RNase H (2 U)	2 ul
1 st strand DNA mix (step 1)	20 ul

2nd Strand Buffer: 20 mM Tris-HCl (pH 6.9), 4.6 mM MgCl₂, 90 mM KCl, 0.15 mM b-NAD⁺, 10 mM (NH₄)₂SO₄

[0488] b) Mix, spin and incubate at 16° C. for 2 hrs.
NOTE: can start DNA SISPA whilst this incubation is underway.

[0489] c) Add 10 ul (10 U) T4 DNA polymerase (1 u/ul) and incubate at 16° C. for 15 min.

[0490] d) Heat 2nd strand synthesis at 72° C. 10 minutes, let cool to 37° C.

6. Clean up DNA

[0491] a) Spin phase lock at 13,000 rpm for 30 sec at 4° C.

[0492] b) Add 150 ul of step 2 reaction

[0493] c) Add equal volume phenol/chloroform 160 ul

[0494] d) Shake lightly

[0495] e) Spin at 13000 rpm 5 minutes, 4° C.

[0496] f) Transfer upper phase to new tube ~160 ul

[0497] g) Precipitate DNA add 100% ethanol 2.5V i.e 375 ul and 1 ul glycogen (20 mg/ml) Leave at -20° C. for 2 hrs or O/N

[0498] h) Spin at 13000 rpm 20 minutes, remove S/N off pellet

[0499] i) Wash pellet 1x70% ethanol 13000 rpm 5 min at 4° C.

[0500] j) Take pellet up in 35 ul of water *NOTE: can stop here and freeze at -80° C. until the DNA SISPA sample is also ready.*

DNA SISPA

7. Second DNA Strand Synthesis

[0501] a) Mix together the following:

DNA	50 ul
10 pmol random hexamers (10 pmol/ul)	1 ul
5 U 3'-5' exo Klenow fragment DNA polymerase	1 ul
Buffer (supplied with Klenow fragment DNA polymerase)	1 ul
5 mM dNTP	1 ul
T4gene32	1 ul

[0502] b) Leave at 37° C. for 1 hr

8. Clean up DNA

[0503] a) Spin phase lock at 13,000 rpm for 30 sec at 4° C.

[0504] b) Add 60 ul of step 1 reaction

[0505] c) Add equal volume phenol/chloroform 60 ul

[0506] d) Shake lightly

[0507] e) Spin at 13,000 rpm 5 minutes, 4° C.

[0508] f) Transfer upper phase to new tube ~60 ul

[0509] g) Precipitate DNA add 100% ethanol 2.5V i.e 150 ul and 1 ul glycogen (20 mg/ml) Leave at -20° C. for 2 hrs or overnight

[0510] h) Spin at 13,000 rpm for 20 minutes, remove supernatant off pellet

[0511] i) Wash pellet 1x70% ethanol 13,000 rpm 5 min at 4° C.

[0512] j) Take pellet up in 44 ul of water *NOTE: can stop here and freeze at -80° C. until the RNA SISPA sample is also ready.*

Generation of Recombinant Nucleic Acid Sequences

9. Restriction Digest

[0513] a) Add 10 U Csp 6.1 (i.e 1 ul of 10 U/ul stock) to 35 ul of sample, add 4 ul of Buffer B and 5 ul of Csp6I

[0514] b) Incubate at 37° C. for 2 hr

[0515] c) Inactivate at 65° C. for 20 minutes

10. Dephosphorylate digested DNA

[0516] a) To inactivated restriction digest (50 ul) add:

[0517] 6 ul of 10x CIP dephosphorylation buffer

[0518] 0.3 ul of CIP 18 U/ul

[0519] 3.7 ul water

CIP Dephosphorylase buffer 1x: 0.05M Tris-HCl, 0.1 mM EDTA, pH8.5

[0520] b) Incubate at 37° C. for 30 minutes

[0521] c) Add another 0.3 ul of CIP 18 U/ul and incubate at 37° C. for 30 minutes

11. Clean Up DNA

[0522] a) Spin phase lock at 13,000 rpm for 30 sec at 4° C.

[0523] b) Add 60 ul dephosphorylated DNA

[0524] c) Add equal volume (60 ul) phenol/chloroform

[0525] d) Shake lightly

[0526] e) Spin at 13,000 rpm 5 minutes, 4° C.

[0527] f) Transfer upper phase to new tube ~50 ul

- [0528] g) Precipitate DNA add 2.5 volumes 100% ethanol (150 μ l) and 1 μ l glycogen (20 mg/ml) Leave at -20° C. for 2 hrs or overnight
- [0529] h) Spin at 13,000 rpm 20 minutes, remove supernatant off pellet
- [0530] i) Wash pellet 1 \times 70% ethanol, spin 13,000 rpm 5 min at 4° C.
- [0531] j) Dessicate for 2-3 minutes or air dry for 15 minutes
- [0532] k) Reconstitute in 5.8 μ l H_2O .

12. Adaptor Ligation

- [0533] a) Mix together:

T4 DNA ligase (5 U/ μ l)	1.2 μ l
5x Ligase Buffer	2 μ l
50 pmol adaptor (phosphorylated ends)	1 μ l
DNA from Step 3.	5.8 μ l

Ligase buffer 5X: 330 mM Tris-HCl, 25 mM $MgCl_2$, 25 mM DTT, 5 mM ATP, pH 7.5

- [0534] b) Incubate 4° C. for 1 hr and 16° C. overnight

13. PCR Reaction (Results FIG. 2)

- [0535] a) Set up the following mix:

Ligated DNA (step 4)	2 μ l
50 pmol NBam24	1 μ l
5 mM dNTP	2 μ l
2 mM $MgCl_2$	2 μ l
10X PCR Buffer	5 μ l
H_2O	38 μ l

10X PCR buffer: 100 mM Tris-HCl, 500 mM KCl (pH 8.3)

- [0536] b) Heat at 72° C. for 3 minutes
- [0537] c) Add 0.5 μ l Taq DNA polymerase (5 U/ μ l)
- [0538] d) Run cycle:
- [0539] 72° C. for 5 minutes
- [0540] (94° C. for 1 minute, 72° C. for 3 minutes) \times 40 hold at 4° C.
- [0541] e) Run 10 μ l and 40 μ l of product on 1.0% EtBr gel (leave a well between them to make purification easier)

14. Cloning PCR Product

- [0542] a) Cut out sections of smeared region from gel as a lot of the dominant bands can be contaminating sequence from the products used in the methods, rather than the actual sample. Bands can also be hard to see if they are in the smeared regions.
- [0543] b) Clean up DNA from agarose using the Minielute Gel Extraction Kit (Qiagen)
- [0544] 1. Excise the DNA fragment from the agarose gel with a clean, sharp scalpel.
- [0545] 2. Weigh the gel slice in a colourless tube. Add 3 volumes of Buffer QG to 1 volume of gel (100 mg \sim 100 μ l).
- [0546] 3. Incubate at 50° C. for 10 min (or until the gel slice has completely dissolved). To help dissolve gel, mix by vortexing the tube every 2-3 min during the incubation.
- [0547] 4. After the gel slice has dissolved completely, check that the colour of the mixture is yellow (similar to Buffer QG without dissolved agarose). Note: If the

colour of the mixture is orange or violet, add 10 μ l of 3 M sodium acetate, pH 5.0, and mix. The colour of the mixture will turn to yellow.

- [0548] 5. Add 1 gel volume of isopropanol to the sample and mix by inverting the tube several times.
- [0549] 6. Place a MinElute column in a provided 2 ml collection tube in a suitable rack.
- [0550] 7. To bind DNA, apply the sample to the MinElute column, and centrifuge for 1 min.
- [0551] 8. Discard the flow-through and place the MinElute column back in the same collection tube.
- [0552] 9. Add 500 μ l of Buffer QG to the spin column and centrifuge for 1 min.
- [0553] 10. Discard the flow-through and place the MinElute column back in the same collection tube.
- [0554] 11. To wash, add 750 μ l of Buffer PE to the MinElute column and centrifuge for 1 min.
- [0555] 12. Discard the flow-through and centrifuge the MinElute column for an additional 1 min at $\geq 10,000\times g$ ($\sim 13,000$ rpm).
- [0556] 13. Place the MinElute column into a clean 1.5 ml microcentrifuge tube.
- [0557] 14. To elute DNA, add 10 μ l of Buffer EB (10 mM Tris.Cl, pH 8.5) or H_2O to the centre of the membrane, let the column stand for 1 min, and then centrifuge for 1 min.
- [0558] c) For ligations and cloning use Invitrogen TA Cloning[®] Kit Version V 7. Set up the 10 μ l ligation reaction as follows:

Fresh PCR product	6 μ l
10X Ligation Buffer	1 μ l
pCR Φ 2.1 vector (25 ng/ μ l)	2 μ l
T4 DNA Ligase (4.0 Weiss units)	1 μ l

Incubate the ligation reaction at 14° C. overnight, or at -20° C. until you are ready for transformation.

- [0559] d) Transform One Shot[®] Competent Cells.
- [0560] 1. Centrifuge vials containing the ligation reactions briefly and place them on ice.
- [0561] 2. Thaw on ice one 50 μ l vial of frozen One Shot[®] Competent Cells (enough for 2 ligations).
- [0562] 3. Pipette 2 μ l of each ligation reaction into 25 μ l of competent cells and mix by stirring gently with the pipette tip.
- [0563] 4. Incubate the vials on ice for 30 minutes. Store the remaining ligation mixtures at -20° C.
- [0564] 5. Heat shock the cells for 30 seconds at 42° C. without shaking. Immediately transfer the vials to ice.
- [0565] 6. Add 125 μ l of room temperature SOC medium to each vial.
- [0566] 7. Shake the vials horizontally at 37° C. for 1 hour at 225 rpm in a shaking incubator.
- [0567] 8. Spread 50 μ l to 100 μ l from each transformation vial on LB agar plates containing ~ 80 mg/ml X-Gal and 100 μ g/ml ampicillin.
- [0568] 9. Incubate plates overnight at 37° C. Place plates at 4° C. for 2-3 hours to allow for proper colour development.

15. Screening Colonies for Inserts and Sequencing (Results FIG. 3)

[0569] a) Use HotStarTaqMaster Mix (50 μ l/well of plate):

1X	110X (sufficient for one plate)
25 μ l HotStarTaqMaster Mix (vortex)	2750 μ l
12.5 μ l M13-20f (50 pmol)	1375 μ l
12.5 μ l M13-20f (50 pmol)	1375 μ l
add 50 μ l per well of the plate	

To make the M13-20f (50 pmol) and M13r (50 pmol) stocks: mix 0.5 μ l of 100 μ M primer with 12 μ l of water i.e. 500 μ l of 100 μ M stock primer + 1200 μ l water (from HotStarTaq Kit).

[0570] b) Place sterile aluminium foil over the plate containing the HotStar TaqMaster Mix. Stab through the foil to make a hole, and then stab a bacterial colony into each well of the plate.

[0571] c) Take off aluminium foil and add strip caps to seal plate.

[0572] d) Run PCR protocol:

[0573] 95° C. for 15 min

[0574] (94° C. for 30 s, 50° C. for 30 s, 72° C. for 1 min) \times 30

[0575] 72° C. for 1 min

[0576] 4° C. hold.

[0577] e) Run 5-10 μ l of PCR on gel

[0578] f) Use Qiagen Mini elute to clean up the remaining PCR product to sequence.

Example 2

Enzyme Linked Immunosorbent Assay to Detect Antibodies to PMC Virus

[0579] 1. Clone and express the PMC virus protein of interest (eg E2, NS3) in baculovirus and purify the expressed protein. This purified protein can be used as an antigen to detect specific antibodies to the PMC virus proteins of interest.

2. Coat 'medium binding' 96 well microplates (50 μ l per well) with antigen diluted in carbonate buffer (0.05M Carbonate buffer 1 \times (pH 9.6): Na₂CO₃ (1.59 gm); NaHCO₃ (2.93 gm) water to 1 L). Hold overnight at room temperature (18-25° C.).

3. Dilute samples and controls (Negative, High and Low Positive) 1/100 in sample diluent (phosphate buffered saline (pH 7.3) solution containing 1% skim milk powder and 0.05% Tween 20).

4. Wash plates 5 times with PBS-Tween and tap to dry.

5. Transfer diluted samples and controls to the ELISA plate in duplicate: 50 μ l to each well.

6. Incubate at 37° C. for 1 hr in a humidified container.

7. Wash plates 5 times with PBS-Tween, rotate and wash 5 more times, then tap to dry.

8. Dilute conjugate (antiporcine IgG, horseradish peroxidase conjugated) in sample diluent and add 50 μ l to each well.

9. Incubate at 37° C. for 1 hr in a humidified container.

10. Wash plates 10 times with PBS-Tween, then 5 times with purified water.

11. Develop by adding 100 μ l of TMB substrate to each well. Incubate at 37° C. in the dark for about 10 min until target OD is achieved for controls. A commercially available TMB substrate can be used (eg. Boehringer Mannheim Corp., Pierce Chemical Co., and Kirkegaard & Perry Laboratories).

12. Stop by adding 100 μ l of 1M sulphuric acid.

13. Read OD values at 450 nm.

14. Calculate results.

Example 3

Enzyme Linked Immunosorbent Assay to Detect Antigens of PMC Virus

[0580] It should be noted that working solutions of the detector reagent and enzyme conjugate reagents should be made within approximately 1 hour of anticipated use and then stored at 4° C.

[0581] Materials

[0582] ELISA Wash Buffer—10 \times concentrate: 1 M Tris; HCl (6.25 Normal) for pH adjustment; 0.01% Thimerosal; and 5% Tween 20.

[0583] Detector Reagent—10 \times concentrate: 25% Ethylene Glycol, 0.01% Thimerosal, approximately 5% biotinylated goat anti-PMC virus antibody, and 0.06% yellow food colouring in PBS (pH 7.4). The working Detector Reagent is prepared by mixing 1 part of the Detector Reagent—10 \times concentrate, 1 part of NSB Reagent 10 \times concentrate, and 8 parts of Reagent Diluent Buffer. This working agent should be prepared within approximately 1 hour of anticipated use.

[0584] NSB Reagent—10 \times concentrate: 25% Ethylene Glycol, 0.01% Thimerosal, 0.2% Mouse IgG, 0.06% red food colouring in PBS (pH 7.4).

[0585] Reagent Diluent Buffer: 2.5% Bovine Serum Albumin, 0.01% Thimerosal, and 1.0% bovine gamma globulin in PBS (pH 7.4).

[0586] Enzyme Conjugate Reagent—10 \times concentrate: 25% Ethylene Glycol, 0.01% Thimerosal, streptavidin-biotinylated horseradish peroxidase complex (dilution approximately 1 to 700), 0.1% rabbit albumin, and 0.02% rabbit gamma globulin in PBS (pH7.4). Working Enzyme Conjugate Reagent should be prepared by mixing 1 part of Enzyme Conjugate Reagent—10 \times concentrate, 1 part of NSB Reagent—10 \times concentrate, and 8 parts of Reagent Diluent buffer. This working reagent should be prepared within approximately 1 hour of anticipated use.

[0587] Negative Control: 1% Igepal, and 0.01% Thimerosal in PBS (pH 7.4).

[0588] Positive Control: 1% Igepal, 0.01% Thimerosal, 1% Bovine Serum Albumin, PMC virus culture (dilution approximately 1:20) and 50 μ M phenyl methyl sulfonyl fluoride in PBS (pH7.4).

[0589] Method

1. Prepare specimens by standard methods. For samples containing cells (tissues, white blood cells) homogenise the tissue and add sample lysis buffer (1% NP40). Allow at least 1 hour for antigen extraction and mix continually.

2. Clarify specimens by centrifuging for 15 minutes at approximately 2000 g;

3. Coat 96 well microplates with purified polyclonal antiserum raised against PMC virus antigens (100 μ l/well). Alternatively, a mixture of anti-PMC virus monoclonal antibodies may be used. Each 96-well tray is coated overnight at room temperature with 0.1 ml per well of a solution containing purified antibody at 5 μ g/ml and bovine serum albumin at 10 μ g/ml in carbonate buffer (pH9.6). Following the coating, each tray is washed three times with ELISA wash buffer and allowed to dry overnight at 4° C. A foil pouch is used to encase each tray after drying, and a desiccant is included inside each pouch to remove moisture.

4. Wash ELISA plates 3 times by pipetting 0.2 ml of ELISA Wash Buffer into each well and tap or pipette dry prior to the addition of sample.
5. Block ELISA plates with Blocking solution 1 (200 uL/well) for 30 min at 37° C. in a humidified container.
6. Transfer 100 uL of each specimen (including controls) to the ELISA plate;
7. Incubate plates for 60 min at 37° C. in a humidified container;
8. Wash ELISA plates 5 times with ELISA Wash Solution;
9. Block ELISA plates with Blocking Solution 2 (150 uL) for 30 min at 37° C. in a humidified container;
- 10 Wash ELISA plates 5 times;
11. Add Detector Reagent containing biotinylated anti-PMC virus monoclonal antibody (100 uL) to all wells;
- 12 Incubate plates for 60 min at 37° C. in a humidified container;
- 13 Wash plates 5 times;
14. Add Enzyme Conjugate Reagent containing streptavidin-biotinylated horseradish peroxidase complex and add 100 uL to all wells;
15. Incubate plates for 30 min at 37° C. in a humidified container;
16. Wash plates 10 times;
17. Prepare and add 100 uL of TMB substrate solution to all wells. A commercially available TMB substrate may be used (eg. Boehringer Mannheim Corp, Pierce Chemical Co, and Kirkegaard & Perry Laboratories).
18. Incubate plates for approx 10 min at room temperature in the dark;
19. Stop reaction with 1M sulphuric acid (100 uL per well);
20. Read ODs on ELISA plate reader at 450 nm;
21. Calculate results.

Example 4

Detection of PMC Virus RNA by Reverse Transcriptase (RT) Polymerase Chain Reaction (PCR)

- [0590]** a) Extract RNA from the test specimen as described in Example 1. Include in all steps of the reactions known positive and negative controls and a 'blank'.
- [0591]** b) Reverse transcribe (RT) the RNA as follows:
- [0592]** 1. Mix together the following:

random hexamers (50 pmol)	1 ul
RNA (in H ₂ O)	9 ul

- [0593]** 2. Heat at 90° C. for 3 minutes, spin and put on ice
- [0594]** 3. On ice add:

1st strand buffer	4 ul
0.1M dTT	2 ul
5 mM dNTP	2 ul
SSIII (200 U)	1 ul

- [0595]** 4. Mix, spin heat at 45° C. for 60 minutes.
- [0596]** 5. Heat inactivate at 70° C. for 10 minutes
- [0597]** 6. Place on ice.
- [0598]** c) Set up 1st round PCR
- [0599]** 1. Mix together the following PCR reagents

RT	5 ul
Forward primer 4 uM	1 ul
Reverse primer 4 uM	1 ul
Hotstart PCR mix (Qiagen)	12.5 ul
Water	5.5 ul

(see Table 3 for 1st reaction PCR primers)

- [0600]** 2. Cycle the PCR machine at:
- [0601]** 95° C. for 15 minutes
- [0602]** (94° C. for 30 sec, 50° C. for 30 sec, 72° C. for 1 min)×40
- [0603]** 72° C. for 1 min
- [0604]** 4° C. hold
- [0605]** d) Set up Nested PCR
- [0606]** 1. Mix together the following PCR reagents:

1st PCR product	1 ul
Forward nested primer 20 uM	1 ul
Reverse nested primer 20 uM	1 ul
Hotstart PCR mix (Qiagen)	12.5 ul
Water	9.5 ul

(see Table 3 for nested PCR primers. If no nested primer is listed, use 1st PCR primer)

- [0607]** 2. Cycle the PCR machine at
- [0608]** 95° C. for 15 minutes
- [0609]** (94° C. for 30 sec, 50° C. for 30 sec, 72° C. for 1 min)×25
- [0610]** 72° C. for 1 min
- [0611]** 4° C. hold
- [0612]** e) Run 5 ul of nested PCR product on a 1.5% ethidium bromide gel for 1 hour. Depending on the primers used, the expected size of the product is as listed in Table 1.

TABLE 3

<u>Primers for PCR detection of PMC virus</u>				
Clone	Virus	*Primer name	Primer Sequence (5' to 3')	Nested Product size
CR3 9	Pestivirus	CR39F (63)	CACATCTAGCAGCAGACTATGA	103 bp
		CR39R (190)	GTACCAGTTGCACCACCC	
		CR39FN (87)	TGAAAAGGATTTCACGG	
ER5 10	Pestivirus	ER510F (7)	AAACCGACGAAGTAGACC	114 bp
		ER510R (213)	AGACGAGAACATAGTGCC	
		ER510FN (68)	GAAACAGTAAAGCCAACG	
		ER510RN (182)	CTGGTAATCGAAACATC	

TABLE 3-continued

Primers for PCR detection of PMC virus				
Clone	Virus	*Primer name	Primer Sequence (5' to 3')	Nested Product size
ER6 2	Pestivirus	ER62F (203)	GGGACCGAGGGATACGA	98 bp
		ER62FN (373)	AGAGGTAATTGGGTAT	
		ER62R (637)	CAGCAGGTTGATTCTTCAT	
		ER62RN (516)	TTGCCAAGTTTCAC	
ER5 5	Pestivirus	ER55F (31)	AAACCGCCGAAGTAAACC	143 bp
		ER55R (214)	CTGGAGCCCTGGTAATGG	
		ER55FN (64)	GACGGGAATGGGTTCA	
		ER55RN (162)	TAGGTGCTTCTTATTGGTAT	

*F = forward primer, R = reverse primer, FN = forward nested primer, RN = reverse nested primer

Example 5

Determination of Full length viral sequence

[0613] Once the authenticity of the presence of PMC virus sequence has been confirmed in a sample by PCR, the entire viral sequence can be acquired by designing PCR primers to span the gaps between the clones (refer to Table 4). RT-PCR was carried out as either a two step (RT then PCR) or one step RT-PCR reaction.

1. RT Reaction:

[0614] a) Mix together the following:

[0615] 1 ul random hexamers (50 pmol)

[0616] 4 ul RNA

[0617] 4 ul Rnase free water

[0618] b) Heat 70° C. 10 minutes, spin and put on ice

[0619] c) On ice add

[0620] 4 ul 1st strand buffer

[0621] 2 ul 0.1M dTT

[0622] 2 ul 5 mM dNTP

[0623] 2 ul SSIII (400 U)

[0624] d) Mix, spin heat at 42° C. for 60 minutes.

[0625] e) Heat inactivate at 70° C. for 10 minutes

[0626] f) Place on ice

2. PCR Reaction:

[0627] a) Mix together the following PCR reagents

RT	1 ul
Forward primer 20 uM	1 ul
Reverse primer 20 uM	1 ul
Hotstart PCR mix (Qiagen)	12.5 ul
Water	13.5 ul

(see Table 4 for PCR primers)

[0628] b) Cycle the PCR machine at:

[0629] 95° C. for 15 minutes

[0630] (94° C. for 30 sec, 47° C. for 30 sec, 72° C. for 2 min)×40

[0631] 72° C. for 1 min

[0632] 4° C.—hold

3. One Step RT-PCR Method

[0633] a) Mix together the following reagents from the SSIII RT-PCR Kit

2x reaction mix	25 ul
Forward primer 30 uM	1 ul
Reverse primer 30 uM	1 ul
SSIII RT/Platinum mix	2 ul for products 2.5 kb or less 4 ul for products 2.5 kb or more
Water	15.8 ul for products 2.5 kb or less 13.8 ul for products 2.5 kb or more

(see Table 4 for PCR primers)

[0634] b) Cycle the PCR machine at:

[0635] 50° C. for 50 minutes

[0636] 94° C. for 2 min

[0637] (94° C. for 15 sec, 50° C. for 30 sec, 68° C. for 1 min/kb)×40

[0638] 68° C. for 5 min

[0639] 4° C.—hold

[0640] RT-PCR product of interest was PCR spin cleaned and cloned into the Invitrogen TA cloning vector PCR2.1 (see Example 1). Positive clones were then identified and sent for sequencing, as described in Example 1.

[0641] The primers used for sequencing were M13r, m13-20, primers in Table 4 and primers designed specifically for sequencing (see Table 5).

[0642] Plasmid sequence, PCR primers and poor sequence reads were removed from the sequence before being used in the program Bioedit (Hall, T. A. (1999) BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. Nucl. Acids. Symp. Ser. 41:95-98). Bioedit allowed the construction of contigs and the production of the full length consensus sequence for the virus.

TABLE 4

Primers designed to PCR the gaps between the SISPA clones sequences			
Region to	Primer Sequence		Product size
PCR	*Primer names	(5' to 3')	
5'UTR-Erns	JFP1F	CATGCCCATAGTAGGAC	1338 bp
	JFRR3R	ACCGAGTTRCACCAMCCAT	
Erns-P7	CR39-Er55PCR	AGGGCTCTCACATGGTTGTC	1810 bp
	ER55-510-512 R	CCATTACCAGGGCTCCAG	
Erns-NS5A	CR39F(63)	CACATCTAGCAGCAGAC-TATGA	2349 bp
	ER55RN(162)	TAGGTGCTTCTTATTGGTAT	
P7-NS5A	ER55-510-512 F	CGTTGGCCTTTACT-GTTTCATTG	5560 bp
	CR316-CR24R	TCCCCGAAGCTTGGTTTAAT	
NS3-NS5A	NS3F	GTCAGGCCTGCCTATCTTTG	4431 bp
	CR316-CR24R	TCCCCGAAGCTTGGTTTAAT	
NS5A-NS5B	CR316-CR24F	CGGGACCATTAAACCAAGC	2440 bp
	ER62-ER63R	CAGGGGGTTCCAAGAATACA	

*F = forward primer, R = reverse primer

TABLE 5

Primers designed for sequencing			
Protein location of primer	*Primer names	Primer Sequence (5' to 3')	
5 UTR	5utr(140)R	GGTGTACTCACCCTTAGCC	
NPRO	NPRO(630)RS	TTGCTACAATCGCCCTTCTT	
NPRO	NPRO(779)FS	AGGGAGAATGACAGGGTCTG	
Capsid	capsid(927)FS	ACA AAGGAGCAAAACCCAAG	
ERNS	EO(1365)RS	GTCACGTTGGTGGACCCTAC	
E1	E1(2402)RS	AGCCAGAAATGCCACAGC	
E1	E1(2606)FS	ACCTGTGTGGGTGCTAACAT	
E2	E2(3086)RS	TTACTTTGTCTTCCCGTTGC	
NS2	NS2(4409)FS	CCAAGAACTTCCCCATACG	
NS2	ns2(4460)RS	TTCCACATCCTCTTCTTCTTTT	
NS3	NS3(5170)RS	GCTGGCCCTCGAATGATCCA	
NS3	NS3(5468)FS	GTTCCCTGTGCTTGTGCTGA	
NS3	NS3(5670)RS	TGTTTTTGTCTTGGCACTGG	
NS3	NS3(6296)FS	GAGCACAACAGGGCAGAAAT	
NS3	NS3(6479)RS	CCATCTTCTTGTAGGCACA	
NS3	NS3F(6525)F	GTCAGGCCTGCCTATCTTTG	
NS3	NS3(7153)FS	GGAGAAGTCACTGACGCACA	

TABLE 5-continued

Primers designed for sequencing			
Protein location of primer	*Primer names	Primer Sequence (5' to 3')	
NS3	NS3(7241)RS	GCCATTTCAATCCCAGTATG	
NS4B	NS4B(7715)FS	GGGGTCCACACAGCATTGTA	
NS4B	NS4B(7893)RS	CCCTTGATACTCACGCCTGT	
NS4B	NS4B(8532)FS	GCCGACTCAAATGGAGAAA	
NS5A	NS5A(8810)RS	GCCACCTTATCTTGGATCTC	
NS5B	NS5B(10889)FS	AAATGAGAAGAGGGCAGTGG	
NS5B	NS5BF-10936	AAGGCCACCACTCAAATCAC	
NS5B	NS5BR-12039	AGGCTTCTGCTTGACCCAGT	

*FS = forward primer, RS = reverse primer

NOTE:

Numbers in brackets are estimated locations on Reference pestivirus strain NADL.

Example 6

UTR Sequences

[0643] 5'RACE and 3'Race were used to acquire the 5'UTR and 3'UTR sequences.

1. 5' RACE Method

[0644] Sequence data from the complete 5' untranslated region (UTR) was generated using rapid amplification of cDNA ends (RACE, BD), as described by BD Biosciences Clontech with the following modifications. PMC virus-specific primer CR24R (5'TCCCCGGAAGCTTGGTTTAAT 3') was used to generate the cDNA. Hotstart PCR (Qiagen) was carried out with primers CR39R (Table 3) and BD Universal primer A mix (5'CTAATACGACTCACTATAGGGCAAG-CAGTGGTATCAACGCAGAGT3' and 5'CTAATACGACTCACTATAGGGC3') with an annealing temperature of 67° C. and extension time of 2 minutes. The PMC virus specific primer N^{Fro}(630)RS (Table 5) and BD nested Universal Primer A (5'AAGCAGTGGTATCAACGCAGAT3') were used for the Hotstart Nested PCR, with an annealing temperature of 55° C. and an extension time 2 minutes. Nested PCR products were cleaned, cloned and sequenced.

2. 3' RACE Method

[0645] Sequence data from the complete 3' untranslated region was generated by first adding a poly (A) tail to the viral RNA, using Epicentre's A-Plus Ploy(A) polymerase tailing Kit for 8 minutes. This was followed by rapid amplification of cDNA ends (RACE, BD), as described by BD Biosciences Clontech with the following modifications. Hotstart PCR (Qiagen) was carried out with primers ER62F (Table 3) and BD Universal primer A mix (5'CTAATACGACTCACTATAGGGCAAGCAGTGGTATCAACGCAGAGT3' and 5'CTAATACGACTCACTATAGGGC3') with an annealing temperature of 65° C. and extension time of 2 minutes. The PMC virus specific primer NS5B(12100)F (Table 5) and BD nested Universal Primer A (5'AAGCAGTGGTATCAACG-

CAGAT3') were used for the Hotstart Nested PCR, with an annealing temperature of 65° C. and an extension time 2 minutes. Nested PCR products were cleaned, cloned and sequenced.

Example 7

Real Time PCR

[0646] The following primers and a matching probe based on Taqman® technology were developed:

Forward primer: CAGTTGGTGTGATCCATGATCCT
 Reverse primer: GGCTCACCTGCAACTTT
 Probe: 6FAMAAGTCTTCAGCAGTTAACTMGENFQ

[0647] Similar primer/probe combinations may be developed for other segments of the PMC genome.

[0648] A Real Time PCR assay was carried out using the following steps:

- a) Extract RNA from the test specimen. Include in all steps of the reactions known positive and negative controls and a 'blank'.
- b) Prepare reaction mixture (volumes per sample) as follows:

2x Mastermix (Roche)	12.5 uL
40x Multiscribe	0.625 uL
Forward primer	1 uL
Reverse Primer	1 uL
Taqman Probe	1 uL
Template (sample)	2 uL
Water	6.875 uL

- c) Set up cycling conditions for the PCR cycler available (the cycles below are appropriate for a Cepheid Smartcycler)

[0649] Cycle the PCR machine at:

[0650] Stage 1: Repeat 1x

[0651] 48° C. for 30 min

[0652] 95° C. for 10 min

[0653] Stage 2: Repeat 45x

[0654] 95° C. for 15 secs

[0655] 58° C. for 30 secs each

- d) Determine results using the Smartcycler software using cycle-threshold (CT) values. A CT value of <35 is considered to be positive. Values between 35-40 are suspicious and values >40 are negative.

Example 8

Production of Recombinant Baculoviruses and Expression of Recombinant PMC Virus Proteins

1. Cloning of PCR Fragments

[0656] PCR products are purified with PCR SPIN-CLEAN™ columns (Progen Industries, Limited), according to the manufacturer's instructions. If the PCR reaction produces non-specific bands in addition to the required product, or subcloning from another plasmid was necessary, the DNA can be further purified by elution from a 0.8% agarose gel, using a modification of the method described by Heery (1990).

[0657] Purified PCR fragments are digested and ligated into pBlueBacHis A, B or C baculovirus transfer vectors

(MaxBac Baculovirus Expression System, Invitrogen Corporation) containing compatible cohesive overhangs, using standard cloning protocols (Sambrook et al., 1989; Current Protocols in Molecular Biology, 1991). A, B or C vectors provide three different reading frames to achieve protein expression in the baculovirus expression system.

2. Transformation of Baculovirus Plasmids with the PCR Fragments

[0658] The ligations are transformed into competent *E. coli* strain Top 10 (Invitrogen Corporation), Genotype: FmcrA D(mrr-hsdRMS-mcrBC) f80lacZDM15 DlacX74 deoR recA1 araD139 D(ara-leu)7697 galU galK rpsL endA1 nupG, and/or Sure® *E. coli* (Stratagene), Genotype: e14⁻(McrA⁻)D (mcrCB-hsdSMR-mrr) 171 endA1 supE44 thi-1 gyrA96 rel A1 lac recB recJ sbcc umuc::Tn5 (kan^r) uurC[F⁺ proAB lac^Z D m15 Tn10(Tet^r)]^c. Protocols for the preparation of competent cells and transformation of the bacteria are taken from the Invitrogen MaxBac Baculovirus Expression System Manual Version 1.8.

[0659] Screening bacterial clones for plasmid containing PCR fragment and plasmid purification for transfection

[0660] Bacterial clones containing pBlueBacHis+PCR fragment are identified by growing colonies, extracting the plasmids using the boiling miniprep method described in Sambrook, et al. (1989), and then undertaking restriction digests of the plasmids to verify those containing the correct-sized insert. Recombinant plasmids are purified to a level suitable for transfection reactions using plasmid purification kits (QIAGEN Pty Ltd., tip-20 or tip-100 columns), according to the manufacturer's instructions.

3. Production of Purified Recombinant Baculoviruses by Cationic Liposome Transfection of Sf9 Cells to Produce Recombinant Baculoviruses

[0661] Recombinant baculoviruses are produced by co-transfecting linearised wild-type *Autographa californica* nuclear polyhedrosis virus (AcMNPV) DNA and baculovirus transfer vector containing PCR fragment into Sf9 cells, by the technique of cationic liposome mediated transfection. This is carried out according to the Invitrogen MaxBac Baculovirus Expression System Manual Version 1.8.

4. Plaque Purifying Recombinant Baculoviruses

[0662] Recombinant virus is plaque purified three times before virus master stocks are prepared, ensuring the virus is cloned from a single particle and no wild-type virus is present. Plaque assays are set up according the Invitrogen MaxBac Baculovirus Expression System Manual Version 1.8.

[0663] After each round of plaque purification, the recombinant viruses are screened using a modified Pestivirus antigen-capture ELISA (PACE) (Shannon et al., 1991). The modified method involves supernatant + cells (50 µl/well) being added directly to a blocked, washed ELISA plate, and the plate incubated for 1 hr at 37° C. Antibody solution (50 µl/well) is then added. The antibody used is either biotinylated goat anti-pestivirus antiserum or individual anti-PMC virus monoclonal antibodies (mAbs). The plate is incubated overnight at 22° C., then developed as described by Shannon et al. (1991), omitting the incubation with biotinylated anti-mouse IgG for samples that are reacted with the biotinylated goat antiserum.

5. Recombinant Baculovirus Master, Seed and Working Stocks

[0664] The master virus stock for each of the recombinant baculoviruses constructed are made according the Invitrogen

MaxBac Baculovirus Expression System Manual Version 1.8. The titre of the stock is determined by a plaque assay, as described above, except that the cells are overlaid with 1.5% carboxymethylcellulose (CMC, BDH; 6% CMC in deionised water, diluted 1 in 4 with complete TC100+X-gal [125 µg/ml, Boehringer Mannheim]). After 7 days, the blue plaques are counted to give the virus titre.

[0665] The seed and working stock are made from the master and seed stock, respectively using a low MOI of 0.1 to 0.5 pfu/ml. All virus stocks are stored at 4° C. for use in vaccine production. For long term storage of Master, Seed and Working stocks, each recombinant virus is ampouled and frozen at -80° C.

6. Optimisation of Recombinant Protein Production

[0666] Sf9 insect-cell suspensions, adapted to Sf-900 II Serum Free Media according to the protocol described by Gibco BRL (1995), are used to optimise recombinant protein expression. Two conical flasks, containing 50 ml cells (1.5×10^6 cells per ml), are infected with recombinant baculovirus at a high and low MOI, between 0.1 and 5.0. A third flask acts

as an uninfected control culture. The 3 flasks are incubated with shaking at 28° C., and 5 ml aliquots removed at 24 hr intervals for up to 7 days.

[0667] The samples are centrifuged at room temperature (RT) for 10 min at 900×g, and the supernatants carefully removed. The pellets and supernatants are stored at -20° C. until daily sampling is completed. The amount of specific, recombinant pestivirus protein in the samples is then determined using the modified PACE described above. The cell pellets are reconstituted in 200 µl or 250 µl NP-40 (1% [v/v] in PBS), vortexed and centrifuged at RT for 10 min at 900×g. Serial dilutions of the pellet extract (in 1% [v/v] NP40) are assayed. The culture supernatants are assayed undiluted, as well as serially diluted (in 1% [v/v] NP40). If cell viability is reduced at a higher rate of infection, then an MOI of 0.1 to 2 is more appropriate.

[0668] Modifications of the above-described modes of carrying out the various embodiments of this invention will be apparent to those skilled in the art based on the above teachings related to the disclosed invention. The above embodiments of the invention are merely exemplary and should not be construed to be in any way limiting.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 27

<210> SEQ ID NO 1

<211> LENGTH: 12077

<212> TYPE: DNA

<213> ORGANISM: PMC Virus

<400> SEQUENCE: 1

```

gtataacgac agtagttcaa gtgtcgttat gcatcattgg ccataacaaa ttatctaatt      60
tggaataggg acctgcgacc tgtacgaagg ccgagcgtcg gtagccattc cgactagtag      120
gactagtaca aataggtcaa ctggttgagc aggtgagtgt gctgcagcgg ctaagcggtg      180
agtacaccgt attcgtcaac aggtgctact ggaaaggatc acccactagc gatgcctgtg      240
tggacgagga catgtccaag ccaatgttat cagtagcggg ggtcgttact gagaaagctg      300
cccagaatgg gtagttgcac atacagtctg ataggatgcc ggcggatgcc ctgtattttg      360
accagtataa atattatccg ttgtaaaaga tatgaatact tttactttta atacatatgg      420
agggagtgag gaaggaaaca tgttctttag aactgcaacc acgccgccac cagggtgcca      480
agaaccgggt tacacaagca caatgagacc aatttttggc gaaccaccatc cacctctaca      540
caaacacagc acgttaaaat tgccacattg gagggggatc aaaacaatta gagttaagaa      600
gagagaattg ccaaagaagg gcgattgtag caactcaaca acagctccca cttcgggggt      660
gtacgtttaa ttaggggctg tgttctataa agattacacg ggcacggtat accatcgtgt      720
accgctagaa ctttgtacaa accaagagag gtgcgagggg tccaagtgtg tagggagaat      780
gacagggctc gatggcaggt tgtacaacct tttagtatgt ccggacgatt gtatcctctt      840
tgagagacac tgtagaggtc aaacagtcgt cctgaaatgg atttccaacc ccttgacatc      900
accactttgg gtccagagtt gttctgacga caaaggagca aaaccaagg tgaaaccaaa      960
agacgacagc atgaagcaag gaaaaatagt gacaaagcct aaagagactg aagcagatca     1020
aaaaactaga ccaccagatg ccacgatagt ggttgacggg cagaagtatc aggtgaggaa     1080

```

-continued

gaaggggaaa	gcgaaaccca	agactcaaga	cggcttatac	cacaacaaga	acaaaccaga	1140
agcgtccagg	aagaagcttg	agaaggcctt	gctagcatgg	gcaatattag	cctgcctatt	1200
ggtggtaccg	gtagggtcca	ccaacgtgac	acaatggaac	ttatgggaca	ataaaagtac	1260
tacagacata	catagcgtca	tgttttctag	agggattaaa	aggagtctgc	atggaatttg	1320
gcccacacaa	atctgcaaag	ggatccctac	acatctagca	gcagactatg	aactgaaaag	1380
gattcacggg	atggtggatg	caagcccat	gaccaacttc	acatgttgta	ggctacagag	1440
acatgagtgg	aacaagcatg	ggtggtgcaa	ctggtacaat	atagagccgt	ggatcaatct	1500
catgaataat	acccaaggac	tattaaacac	tggagacaat	ttcactgagt	gcgcagtcac	1560
atgcaggtat	gatgcagact	taggggtgaa	tatagtact	caagccagga	ctactccaac	1620
tatcctgact	gctgtgaaga	aagggcacia	cttctcttcc	tcaggggagg	tcagggcctc	1680
accctgcaac	tttgagttaa	ctgctgaaga	cttgctcagg	atcatggatc	acaccaactg	1740
cgagggattt	acctacttcg	gggaaggaat	cgttgacggt	tacaccgagg	tagtagagaa	1800
ggccagggtca	agtgggttca	gggctctcac	atggttctgc	agtaagattg	aaaacaccaa	1860
gaaaaaata	ttcggagctg	aagccagtc	ttactgcca	gtggctaaga	gggtcttcaa	1920
cattatttat	accaacaatt	gcaccccgt	tggactgcca	gataagtcaa	aaattatagg	1980
accaggaacc	ttgacatca	gtggcaggga	tgaattcata	tttccaaaac	tcccctacca	2040
cgtagatgac	ttcattctac	tgagcttaat	tgcaatgtct	gattttgctc	cagagacatc	2100
aagtataatc	tacctggctt	tgactacct	aatgccaagt	aatgacaaca	gggacttcgt	2160
gatggacctg	gacccaata	aactaaacct	tactgcaact	aaatccgtgg	caagtgtggt	2220
ccctacatcg	gtgaatgtgt	taggtgaatg	ggtgtgcgtc	aaaccaagtt	ggtggcctta	2280
ttccgccgaa	atcactaatc	tgataggagg	tgatcatcacc	gtggcagact	tagttatcaa	2340
gaccattgaa	gaattgctaa	atgtgtggac	cgaagcaaca	gctgtggcat	ttctggctgc	2400
tctaataaaa	attttttagag	gccagccgat	ccaagcggta	gcattggttaa	tcacatagag	2460
gggagcacia	gcccacacct	gcaaccctga	attcatgtac	gcattagcga	aaaataccag	2520
cataggttca	ttaggaccag	aatcactgac	gacaaggtgg	taccaactaa	ccagcggttt	2580
caaaactact	gacagcacga	ttgaagtcac	ctgtgtgggt	gctaacaatga	ggattcatgt	2640
agtgtgcccc	cttghtaagt	acagatattt	ggccataaac	caccctagag	caactgccaac	2700
aacggcgtgg	ttcaggaaaa	tacacactca	gcatgaggta	ccaagagaaa	gaatcatgag	2760
tgagtcaaaa	aggaggtaca	cttgtccttg	tggttctaaa	ccagtgggtga	ggtcaacaac	2820
acaattcaac	ccaatatcta	tatctacccc	aagctttgaa	cttgaatgcc	ctagggggtg	2880
gactggggct	gtagagtgtg	cactagtctc	cccatcaact	ctgacaacag	agactatatt	2940
cacatacagg	aagcccaaac	cattcggact	tgaaaactgg	tgcaagtata	cagtgtgga	3000
gaaagggatc	ctgtattctt	gtaaatttgg	gggcaattca	acatgcatca	aagggcttat	3060
agttaaaagg	caacgggaag	acaaagtaag	gtactgtgaa	tggtgtggtt	ataagttcag	3120
ttcaccacaa	ggactgcctc	agtatccact	gggattgtgt	gagaaagaac	aatcagaagg	3180
actcagggat	tatggtgact	tcccatgctg	caacaacggc	acttgatttg	acaaagaagg	3240
tagtgtgcaa	tgctacatag	gggataagaa	agttaccgtg	aagctgtata	atgcctcact	3300
attggcccc	atgcctgca	aaccatagt	gtataactcc	caggggcccc	cagcgcctaa	3360

-continued

gacctgact tatagtgagg cctcaacatt agaaaataaa tattatgaac ccaggacag	3420
ctactaccag caatacatta taaagtcagg gtatcaatat tggtttgatc tcacagcaaa	3480
ggatcatgtg gcagactgga tcacaaaata ctttccaata ataatagtg ccttgtagg	3540
gggcagaggc accttgtagg tgttgatagc ttatgagttg ctaactcagt atgaggtagt	3600
aggagacgag aacatagtg ctaagctga agccctgga atcggaaca tcttgatgag	3660
tttagactta gagataatta gctgcttct tctgttggtg atcgtggtga aaaaacaagc	3720
tgtcaggaga acgttggtt tactgtttca ttggataact atgaacccat tccagtcagt	3780
aatgatcaca gtgggtact tctgtggtt ggtgagggcc gaagagggaa ctaaagaggg	3840
tagtacaagc gggccaccaa tccatgtagt tgcaatactg ttattcctct tgtaccacac	3900
agtgaagtat aaggacttta acatagcaat gatcttactt ataacattgt cctgaaaag	3960
ctcatcctac atacatacca gcttgatga aattccattg cttgtggctg taataagtct	4020
cacatgctcc atatacattt ttgacttgca ggtaaagagc aagctagtgg cccaactat	4080
aggataaatt ggagttacc tagcaatgag agttttgtgg ctggaaggc aaatgactat	4140
accaacccc tctgtgtcca ttagtctgat agatccaaag atggcataa tactctactt	4200
gatatcccta actattacag tcaatcacia cctagaccta gcaagttatt gcttgaact	4260
gggacctttt atcctatcat tccatacaat gtgggtggat gttgtcatcc tctgtctcat	4320
gctgccttg tacgaactga taaaagtcta ctacctaaaa aagaagaaag aggatgtgga	4380
aacatggttc caaaattcag gaatatccac ccaagaaact tcccatacg gatttgattt	4440
ttctagcccc ggggagggg tgcacacact accaatgcaa aataaaacca aattttgtag	4500
gactgcttac atgactgtac taagggcttt ggtgataaca gccatcagca gtgtctggaa	4560
accaataatt ttagcagaac tccataaga gccagtgat tggacacaca ttaaaatagc	4620
caaagaattg gcgggtgcaa gcaggttctg tctaggttc attgcatcta ttatagagtt	4680
gaattgggcc atggacgaaa aagaagcatc tccgtacaaa agattttacc tattatcatc	4740
caaaaataca gatctaattg ttaagcacia aatccaaaat gagacagtaa aatcctggtt	4800
tgaagaaact gaaatatttg gaatacaaaa agtggcaatg gtgataaggc ctcatctct	4860
gagtttgag ccaaatgcca tcccttgctc cgtttgtgaa gaaaaacaaa atcaaaaagc	4920
caaaaaggcc tgccttaagt gtggtagtag aggcactcaa ataaagtgtg ggtgacact	4980
ggccgagttt gaggaagaac attacaaaa aatatacatc ctggaaggcc aagatgaaac	5040
tccatgaggg aaagaagaaa gacagcaagt aacttatgtc tctaggggtg ctctgttct	5100
taggaatctt cctatcttag cttcaaaaa caaatccta cttgtaggca atctgggtat	5160
ggaattgcaa gatttggaag gtatgggatg gatcattcga gggccagccg tctgcaagaa	5220
gataatacac catgagaaat gcaggccttc aataccagac aaactcatgg cattctcgg	5280
gattatgctt aggggagtta caccaagagc ccctacacgg ttccctgtgt ccttgctgaa	5340
gataagacgg ggttttgaga ccggctgggc ctacacacac cctggagggg taagtagtgt	5400
gatgatgtc accgctgggt cggatatata tgtcaatgac tcaataggga ggacaaaaat	5460
ccagtgcaa gacaaaaaca ctacaacaga tgagtgtgaa tatggtgtga aaacagactc	5520
agggtgctct gatggagctc ggtgctatgt catcaaccct gaagcaacca acatagcagg	5580
gaccaagggg gccatggtac acctgaggaa agctggagga gagttcaact gcgtgactgc	5640

-continued

ccagggtacc cccgccttct ataactctaaa gaacttaaaa ggatggtcag gctgcctat	5700
ctttgaagct gccacaggaa gagtggtagg aagggtaaaa gcaggaaaa acactgacaa	5760
tgctccaaca accattatgt cagggacgca agtggcaaaa ccatcagagt gtgacctaga	5820
atcagtggtg aggaaactag agacaatgaa cagaggggaa ttcaacaag tgactctggc	5880
tacaggcgca ggaagacaaa ccatgctacc aaagctgta atagaatcca taggcaggca	5940
taagagagtg ttagtactga tcccgttgag agctgcagcg gagggggtgt accagtacat	6000
gagaacaaa cacccaagca tatctttcaa cttgaggata ggggatctga aagaagggtga	6060
catggcaact gggatcacct atgcctctta tgggtacttc tgccaaatgg acatgcctag	6120
actggagaat gcaatgaagg aataccacta tattttcttg gatgaatc actgtgccac	6180
accagaacag ttggcagtg tgctcaaaa acataggctc ggtgaatcag ttagggtaat	6240
agccatgacc gccacgcat cccggactgt gagcacaaca gggcagaaat tcacaattga	6300
ggaggtggta gtacctgaag tgatgaaggg ggaggacctt gctgatgatt acatcgaaat	6360
agcagggttg aagggtccaa agaagagatt agagggtaac gtactgactt ttgtgcctac	6420
aaggaagatg gcatcgaaa cagcaaaaa attaaccaca cagggataca atgctggata	6480
ctacttcagt ggagaagatc catcatccct gcggcaact acttctaagt caccatata	6540
agtagttgca accaatgcc ttagaatccgg ggtaacctta cgggaccttg atacagtaat	6600
agatacagc atgaagtgtg aaaagagact aagaatcgaa aacaaagctc cctacatcgt	6660
aacaggactg aaaagaatgg ctataacaac gggggagcaa gctcaaagaa aaggtagggt	6720
aggcagggtt aaacctggga ggtacttgag aggacctgaa aacctgcag gtgaaaagga	6780
ctatcactat gaccttttac aggcacagag gtacggcatc caagactcaa taaacatcac	6840
caagtcttc agggagatga actatgattg ggcattatat gaggaagacc cgttaagat	6900
tgccaatta gaggttgctaa acacactcct gatctcaagg gatctgcag tagtaacaaa	6960
aaatctgatg gccccacaaa cacatccga acctatacaa ttggcttaca atagtttaga	7020
aaccctgta cgggtggcat tcccaaaagt gaaaaatgga gaagtactg acgcacatga	7080
aacttacgag ttgatgacct gtaggaaget tgagaaagac cccctatat acctgtatgc	7140
aacagaagaa gaagatctcg tagtggacat actgggattg aaatggccag acgccacaga	7200
gagggctgct ttggaagtgc aagacgccct gggccagatc acaggtttat ctgcagggga	7260
gacagcttta ctcatagccc tattagggtg ggtgggctac gaagccttg tgaagaggca	7320
cgtgcctatg gtgacagaca tatacaccct agaagatgaa aaattggaag aactacaca	7380
cctacaattt gcccagatg atctgaacaa ttcagatacc attgagctcc aagacttatc	7440
gaatcaccaa atccaacaaa ttctagaagg tgggaaggaa tatgtcggcc aagcctacca	7500
attcctcagg ttgcaagctg agagggctgc caactcagac aaaggcaaga aagcaatggc	7560
agcggcccca ttactagccc acaagttcct ggaatacttg caagagcatg caggtgacat	7620
aaagaagtat ggtctatggg ggttocacac agcattgtat aacagcataa aagaaagact	7680
gggtcacgaa actgcattcg catctctggt tataaaatgg attgcctttt cctcagatgg	7740
agtcccgagg atgattaagc aagcagcagt agacttgggt gtatactata taatcaacag	7800
gctgagat caaggggata agggagacaca gaatgcaggt agacaatttg ttggctccct	7860
ttttgtttca tgtctagcag agtacacatt caaaaacttc aataaatcag cattagaagg	7920

-continued

attgatcgag cctgccttaa gctatctacc ctacgcttca agcgactaa agttatctct	7980
accgactaga cttgaaagtg tagtgatact gtccactact atatacagaa catacttacc	8040
aatcaggaaa ggatctagtc agggtttagc cgggctggca gttagctcag cgatggagat	8100
catgaaccag aaccaatca gcgtggctat tgcactggca ctaggagtcg gagcaatagc	8160
ggcacataat gccattgaga gcagtgagcc aaaaaggact ctctgatga aggtctttgt	8220
taagaacttt ttggaccaag cagccactga tgagcttggt aaagagaacc ctgagaagat	8280
cataatggca gtgtttgagg gcattcaaac agctggaaat ccattgagac ttgtatacca	8340
tctatatgca atgtttctaa aagggtggac tgcgctggaa atagctgaaa aaaccgctgg	8400
taggaacatt tttgtgttaa caatattga aggattggaa atgttaggcc tggatgccga	8460
ctcaaatgg agaactctga gctctaatta tcttattgat gcagtgaaga aaatcattga	8520
aaaaatgact aaaacagcaa caagcttcac ctacagcttt ttgaaatctt tgcttcctgc	8580
ccccttctcg tgtactaaat cagaagaga tccaagaata ggggtggccc aaaaagacta	8640
cgactacctc gaggtccgat gcgcttggtg gtataacagg agagctataa aaagagactc	8700
aggacctgtg ttatgggaga ccttagagga gacgggtcca gagtactgcc acaacagagg	8760
tgaagggggc ctacagcaat tgaagactac tagatgcttt gtccaaggag aggaaatccc	8820
tccaattgca ctgaggaaaag gagtaggatga gatgttggtc aagggtgttt cattcagaat	8880
agattttgat aaagacaaga tactttcaac agacaagtgg aaggtaccac atagggcagt	8940
tacatcaatc tttgaggatt ggcaggggat tggttacaga gaggcttacc tagggaccaa	9000
accagactat gggggtctgg tgcccagatc ttgtgtaact gtaacaaaac aagggttaac	9060
attcttgaat actgccagag gcatggcttt cactgactgac ctgaccatcc agaactcaa	9120
aatgctgata gctacatgct tcaagaacaa ggtgaaggaa ggggagatc cagctacgat	9180
tgaaggggaa acatggatca acataccact agtgaatgag gacaccggga ccattaacc	9240
aagcttcggg gaaagagtga ttcccgaacc atatgaggag gaccacttg aaggccaag	9300
tgtaatcggt gaaacaggag gcatagccat caaccaata ggggtcaatc cacaatccag	9360
tacatgtgga acagttttta cagcagtga ggatctgtgc caaacagtta gtaataaagc	9420
caagaatc aaatgggtg tttcggaagg ccaataccca ggtccagggg ttgcaaagaa	9480
gacactgaac cagctcatac aagatgaaga cccaaaacca ttcatattg tttgtggctc	9540
tgacaagtca atgtctaatc gggcaaaaac tgcgaggaac atcaagagaa tcaccaccac	9600
aacacctgag aaattcagag acttgcaaaa aaacaagaaa ttgataattg tctgttagg	9660
tgatagatc catgaagata tagaaaagta tgcagacttc aaggacact tcttgaccag	9720
acaaaccttg gaagcactag caagtgccaa agctgtagag aaggacatga ccaagaaaga	9780
agcagcaaga gtattggcaa tggaaagaaa ggatgaacta gaactcccag ggtggctgca	9840
tacagatgca cccaaattcc tagacattac taaggacaac atcacacatc acctaatagg	9900
ggacatgcag agtctgagag aaagagcagg ggagatagga gcaaaggcca cactcaaat	9960
cactaagaaa gggagtgtat acacaatcaa tctgagtacg tgggtgggagt cagagaggtt	10020
ggcatctttg gaacctttgt tccgggaact actatctaaa tgcaggccag tggacagggg	10080
gacatataag aattgtcatt ttgcaacagc agcccaactt gccggaggaa actgggtacc	10140
ggtagacca gttgtacatc ttggggaaat tccggtaaaag aagaaaaaga ctctccccta	10200

-continued

```

cgaggcatac aagctcctaa aagagatggt tgactcggag aaggaattcc ataaaccagt 10260
gagcagggaa aaacaccaat ggatactgaa caaagtgaag actggtggtg acctcggctt 10320
aaaaaatcta gtatgtccag gtaggggttg agaaccaatc ctaagagaga agaagaaatt 10380
caacatttac aacaagagga ttaccagtac tatgttatca gtagggataa ggccagaaaa 10440
attgccagtg gtaagagccc agaccagtac caaagaattt catgaagcaa taagggacaa 10500
aatagacaaa aaagcaaaaca cacagacccc aggcctacac aaagaattgt tggagatatt 10560
caactcaata tgtgccatcc ccgaacttag aaatacctac aaagaggttg attgggacgt 10620
tctaacctca gccataaata ggaaaggtgc agccgggtac ttcgaaaaaa tgaacatagg 10680
ggagatcata gatagtgaca aaaaatcagt ggaacaactc ataaagagaa tgaatcagg 10740
gctagaatc aactactatg agactgcaat accaaaaaat gagaagaggg cagtggtaga 10800
tgattggatg gaaggtgact atgtagaaga aaaaagacca agagtcatac agtatcctga 10860
ggcaaagatg agattagcta taaccaaagt aatgtataac tgggtcaagc agaagcctat 10920
agtaatccct ggatacgaag gtaagactcc tttgtttcat gttttcgaca aggtccacaa 10980
agaatggaaa aatttcaaca gtccagtgc agtcagtttt gacctaaag cctgggacac 11040
acaagtaaca cccaaagacc ttctcctcat atcagaaatc caaagattt attacaagaa 11100
agaataccat agattcatag ataatttgac cgagaaaatg gtggaggtac cagtggtttg 11160
tgaagacgga aacgtctaca taagagaagg tcagagggga agtggcaac cagacactag 11220
cgcaggtaat agtatgttga atgtactgac tatgatata gcttctgca aagctaactc 11280
catcccttac tcagccttc acagggtagc aaagatacat gtgtgtggag atgatggttt 11340
cttgataact gagaaaagt ttggtgagc ctttgcgac aaggggcctc aaattttgat 11400
ggaagcagga aaaccacaaa aacttatagg tgaatttga ctgaaattgg catataaatt 11460
tgatgacatt gaattttgct cgcatacacc aataaaggtc aggtgggctg acaacaacac 11520
atcatacatg cccggaagag acacagctac cattctagct aaaatggcaa cccgccttga 11580
ctctagtggg gagaggggga ccgagggata cgagctggcc gtggcctca gtttcttact 11640
aatgtattct tggaaacccc tggtagaag aatatgcctg cttgtcatgt ctacaattga 11700
cacaaaagaa gctagccaaa ataactat atatacatt aggggggac ccataggtgc 11760
ctacacagag gtaattgggt ataggtgga ccaactaaaa cagacagagt tctctaatt 11820
ggctcagctg aatttgtcaa tggcaact tcaatatac aataaaaaca caaccaagag 11880
actcatcga gattgtgtga aacttgcaa ccaaaataag caaatattgg tgaatgcaga 11940
ccgtttgatc agcaagaaaa cgggctacac atatgagcca acagctggcc actaagat 12000
aggcaagcac tatgaagaaa tcaacctgct gaaagataca ccacaaaaa ctgtctacca 12060
aggaactgaa aggtata 12077

```

```

<210> SEQ ID NO 2
<211> LENGTH: 3886
<212> TYPE: PRT
<213> ORGANISM: PMC Virus

```

```

<400> SEQUENCE: 2

```

```

Gly Gly Ser Glu Glu Gly Asn Met Phe Phe Arg Thr Ala Pro Thr Pro
1          5          10          15
Pro Pro Gly Cys Gln Glu Pro Val Tyr Thr Ser Thr Met Arg Pro Ile

```

-continued

20					25					30					
Phe	Gly	Glu	Pro	His	Pro	Pro	Leu	His	Lys	His	Ser	Thr	Leu	Lys	Leu
	35						40					45			
Pro	His	Trp	Arg	Gly	Ile	Lys	Thr	Ile	Arg	Val	Lys	Lys	Arg	Glu	Leu
	50					55					60				
Pro	Lys	Lys	Gly	Asp	Cys	Ser	Asn	Ser	Thr	Thr	Ala	Pro	Thr	Ser	Gly
65					70						75				80
Val	Tyr	Val	Glu	Leu	Gly	Ala	Val	Phe	Tyr	Lys	Asp	Tyr	Thr	Gly	Thr
				85					90					95	
Val	Tyr	His	Arg	Val	Pro	Leu	Glu	Leu	Cys	Thr	Asn	Gln	Glu	Arg	Cys
			100						105					110	
Glu	Gly	Ser	Lys	Cys	Val	Gly	Arg	Met	Thr	Gly	Ser	Asp	Gly	Arg	Leu
		115					120					125			
Tyr	Asn	Val	Leu	Val	Cys	Pro	Asp	Asp	Cys	Ile	Leu	Phe	Glu	Arg	His
	130					135						140			
Cys	Arg	Gly	Gln	Thr	Val	Val	Leu	Lys	Trp	Ile	Ser	Asn	Pro	Leu	Thr
145					150					155					160
Ser	Pro	Leu	Trp	Val	Gln	Ser	Cys	Ser	Asp	Asp	Lys	Gly	Ala	Lys	Pro
				165					170					175	
Lys	Val	Lys	Pro	Lys	Asp	Asp	Arg	Met	Lys	Gln	Gly	Lys	Ile	Val	Thr
			180					185						190	
Lys	Pro	Lys	Glu	Thr	Glu	Ala	Asp	Gln	Lys	Thr	Arg	Pro	Pro	Asp	Ala
		195					200					205			
Thr	Ile	Val	Val	Asp	Gly	Gln	Lys	Tyr	Gln	Val	Arg	Lys	Lys	Gly	Lys
	210					215					220				
Ala	Lys	Pro	Lys	Thr	Gln	Asp	Gly	Leu	Tyr	His	Asn	Lys	Asn	Lys	Pro
225						230					235				240
Glu	Ala	Ser	Arg	Lys	Lys	Leu	Glu	Lys	Ala	Leu	Leu	Ala	Trp	Ala	Ile
				245					250					255	
Leu	Ala	Cys	Leu	Leu	Val	Val	Pro	Val	Gly	Ser	Thr	Asn	Val	Thr	Gln
			260					265					270		
Trp	Asn	Leu	Trp	Asp	Asn	Lys	Ser	Thr	Thr	Asp	Ile	His	Ser	Val	Met
	275						280					285			
Phe	Ser	Arg	Gly	Ile	Lys	Arg	Ser	Leu	His	Gly	Ile	Trp	Pro	Thr	Gln
	290					295					300				
Ile	Cys	Lys	Gly	Ile	Pro	Thr	His	Leu	Ala	Ala	Asp	Tyr	Glu	Leu	Lys
305						310					315				320
Arg	Ile	His	Gly	Met	Val	Asp	Ala	Ser	Pro	Met	Thr	Asn	Phe	Thr	Cys
				325					330					335	
Cys	Arg	Leu	Gln	Arg	His	Glu	Trp	Asn	Lys	His	Gly	Trp	Cys	Asn	Trp
			340						345					350	
Tyr	Asn	Ile	Glu	Pro	Trp	Ile	Asn	Leu	Met	Asn	Asn	Thr	Gln	Gly	Leu
	355						360					365			
Leu	Asn	Thr	Gly	Asp	Asn	Phe	Thr	Glu	Cys	Ala	Val	Thr	Cys	Arg	Tyr
	370					375					380				
Asp	Ala	Asp	Leu	Gly	Val	Asn	Ile	Val	Thr	Gln	Ala	Arg	Thr	Thr	Pro
385						390					395				400
Thr	Ile	Leu	Thr	Gly	Cys	Lys	Lys	Gly	His	Asn	Phe	Ser	Phe	Ser	Gly
				405					410					415	
Glu	Val	Arg	Ala	Ser	Pro	Cys	Asn	Phe	Glu	Leu	Thr	Ala	Glu	Asp	Leu
			420						425					430	

-continued

Leu Arg Ile Met Asp His Thr Asn Cys Glu Gly Phe Thr Tyr Phe Gly
 435 440 445
 Glu Gly Ile Val Asp Gly Tyr Thr Glu Val Val Glu Lys Ala Arg Ser
 450 455 460
 Ser Gly Phe Arg Ala Leu Thr Trp Leu Ser Ser Lys Ile Glu Asn Thr
 465 470 475 480
 Lys Lys Lys Ile Phe Gly Ala Glu Ala Ser Pro Tyr Cys Pro Val Ala
 485 490 495
 Lys Arg Val Phe Asn Ile Ile Tyr Thr Asn Asn Cys Thr Pro Leu Gly
 500 505 510
 Leu Pro Asp Lys Ser Lys Ile Ile Gly Pro Gly Thr Phe Asp Ile Ser
 515 520 525
 Gly Arg Asp Glu Phe Ile Phe Pro Lys Leu Pro Tyr His Val Asp Asp
 530 535 540
 Phe Ile Leu Leu Ser Leu Ile Ala Met Ser Asp Phe Ala Pro Glu Thr
 545 550 555 560
 Ser Ser Ile Ile Tyr Leu Ala Leu His Tyr Leu Met Pro Ser Asn Asp
 565 570 575
 Asn Arg Asp Phe Val Met Asp Leu Asp Pro Asn Lys Leu Asn Leu Thr
 580 585 590
 Ala Thr Lys Ser Val Ala Ser Val Val Pro Thr Ser Val Asn Val Leu
 595 600 605
 Gly Glu Trp Val Cys Val Lys Pro Ser Trp Trp Pro Tyr Ser Ala Glu
 610 615 620
 Ile Thr Asn Leu Ile Gly Gly Val Ile Thr Val Ala Asp Leu Val Ile
 625 630 635 640
 Lys Thr Ile Glu Glu Leu Leu Asn Leu Trp Thr Glu Ala Thr Ala Val
 645 650 655
 Ala Phe Leu Ala Ala Leu Ile Lys Ile Phe Arg Gly Gln Pro Ile Gln
 660 665 670
 Ala Val Ala Trp Leu Ile Ile Ile Gly Gly Ala Gln Ala Gln Thr Cys
 675 680 685
 Asn Pro Glu Phe Met Tyr Ala Leu Ala Lys Asn Thr Ser Ile Gly Ser
 690 695 700
 Leu Gly Pro Glu Ser Leu Thr Thr Arg Trp Tyr Gln Leu Thr Ser Gly
 705 710 715 720
 Phe Lys Leu Thr Asp Ser Thr Ile Glu Val Thr Cys Val Gly Ala Asn
 725 730 735
 Met Arg Ile His Val Val Cys Pro Leu Val Ser Asp Arg Tyr Leu Ala
 740 745 750
 Ile Asn His Pro Arg Ala Leu Pro Thr Thr Ala Trp Phe Arg Lys Ile
 755 760 765
 His Thr Gln His Glu Val Pro Arg Glu Arg Ile Met Ser Glu Ser Lys
 770 775 780
 Arg Arg Tyr Thr Cys Pro Cys Gly Ser Lys Pro Val Val Arg Ser Thr
 785 790 795 800
 Thr Gln Phe Asn Pro Ile Ser Ile Ser Thr Pro Ser Phe Glu Leu Glu
 805 810 815
 Cys Pro Arg Gly Trp Thr Gly Ala Val Glu Cys Thr Leu Val Ser Pro
 820 825 830

-continued

Ser Thr Leu Thr Thr Glu Thr Ile Phe Thr Tyr Arg Lys Pro Lys Pro
 835 840 845

Phe Gly Leu Glu Asn Trp Cys Lys Tyr Thr Val Val Glu Lys Gly Ile
 850 855 860

Leu Tyr Ser Cys Lys Phe Gly Gly Asn Ser Thr Cys Ile Lys Gly Leu
 865 870 875 880

Ile Val Lys Gly Gln Arg Glu Asp Lys Val Arg Tyr Cys Glu Trp Cys
 885 890 895

Gly Tyr Lys Phe Ser Ser Pro Asn Gly Leu Pro Gln Tyr Pro Leu Gly
 900 905 910

Leu Cys Glu Lys Glu Gln Ser Glu Gly Leu Arg Asp Tyr Gly Asp Phe
 915 920 925

Pro Cys Cys Asn Asn Gly Thr Cys Ile Asp Lys Glu Gly Ser Val Gln
 930 935 940

Cys Tyr Ile Gly Asp Lys Lys Val Thr Val Lys Leu Tyr Asn Ala Ser
 945 950 955 960

Leu Leu Ala Pro Met Pro Cys Lys Pro Ile Val Tyr Asn Ser Gln Gly
 965 970 975

Pro Pro Ala Pro Lys Thr Cys Thr Tyr Arg Trp Ala Ser Thr Leu Glu
 980 985 990

Asn Lys Tyr Tyr Glu Pro Arg Asp Ser Tyr Tyr Gln Gln Tyr Ile Ile
 995 1000 1005

Lys Ser Gly Tyr Gln Tyr Trp Phe Asp Leu Thr Ala Lys Asp His
 1010 1015 1020

Val Ala Asp Trp Ile Thr Lys Tyr Phe Pro Ile Ile Ile Val Ala
 1025 1030 1035

Leu Leu Gly Gly Arg Gly Thr Leu Trp Val Leu Ile Ala Tyr Glu
 1040 1045 1050

Leu Leu Thr Gln Tyr Glu Val Val Gly Asp Glu Asn Ile Val Ala
 1055 1060 1065

Gln Ala Glu Ala Leu Val Ile Gly Asn Ile Leu Met Ser Leu Asp
 1070 1075 1080

Leu Glu Ile Ile Ser Cys Phe Leu Leu Leu Leu Ile Val Val Lys
 1085 1090 1095

Lys Gln Ala Val Arg Arg Thr Leu Ala Leu Leu Phe His Trp Ile
 1100 1105 1110

Thr Met Asn Pro Phe Gln Ser Val Met Ile Thr Val Val Tyr Phe
 1115 1120 1125

Val Gly Leu Val Arg Ala Glu Glu Gly Thr Lys Glu Gly Ser Thr
 1130 1135 1140

Ser Gly Pro Pro Ile His Val Val Ala Ile Leu Leu Phe Leu Leu
 1145 1150 1155

Tyr His Thr Val Lys Tyr Lys Asp Phe Asn Ile Ala Met Ile Leu
 1160 1165 1170

Leu Ile Thr Leu Ser Leu Lys Ser Ser Ser Tyr Ile His Thr Ser
 1175 1180 1185

Leu Tyr Glu Ile Pro Leu Leu Val Ala Val Ile Ser Leu Thr Cys
 1190 1195 1200

Ser Ile Tyr Ile Phe Asp Leu Gln Val Lys Ser Lys Leu Val Ala
 1205 1210 1215

Pro Thr Ile Gly Ile Ile Gly Val Thr Leu Ala Met Arg Val Leu

-continued

1220	1225	1230
Trp Leu Val Arg Gln Met	Thr Ile Pro Thr Pro	Ser Val Ser Ile
1235	1240	1245
Ser Leu Ile Asp Pro Lys	Met Val Ile Ile Leu	Tyr Leu Ile Ser
1250	1255	1260
Leu Thr Ile Thr Val Asn	His Asn Leu Asp Leu	Ala Ser Tyr Cys
1265	1270	1275
Leu Lys Leu Gly Pro Phe	Ile Leu Ser Phe Leu	Thr Met Trp Val
1280	1285	1290
Asp Val Val Ile Leu Leu	Leu Met Leu Pro Trp	Tyr Glu Leu Val
1295	1300	1305
Lys Val Tyr Tyr Leu Lys	Lys Lys Glu Asp	Val Glu Thr Trp
1310	1315	1320
Phe Gln Asn Ser Gly Ile	Ser Thr Gln Glu Thr	Ser Pro Tyr Gly
1325	1330	1335
Phe Asp Phe Ser Ser Pro	Gly Glu Gly Val His	Thr Leu Pro Met
1340	1345	1350
Gln Asn Lys Thr Lys Phe	Cys Arg Thr Ala Tyr	Met Thr Val Leu
1355	1360	1365
Arg Ala Leu Val Ile Thr	Ala Ile Ser Ser Val	Trp Lys Pro Ile
1370	1375	1380
Ile Leu Ala Glu Leu Leu	Ile Glu Ala Val Tyr	Trp Thr His Ile
1385	1390	1395
Lys Ile Ala Lys Glu Leu	Ala Gly Ser Ser Arg	Phe Val Ala Arg
1400	1405	1410
Phe Ile Ala Ser Ile Ile	Glu Leu Asn Trp Ala	Met Asp Glu Lys
1415	1420	1425
Glu Ala Ser Arg Tyr Lys	Arg Phe Tyr Leu Leu	Ser Ser Lys Ile
1430	1435	1440
Thr Asp Leu Met Val Lys	His Lys Ile Gln Asn	Glu Thr Val Lys
1445	1450	1455
Ser Trp Phe Glu Glu Thr	Glu Ile Phe Gly Ile	Gln Lys Val Ala
1460	1465	1470
Met Val Ile Arg Ala His	Ser Leu Ser Leu Glu	Pro Asn Ala Ile
1475	1480	1485
Leu Cys Ser Val Cys Glu	Glu Lys Gln Asn Gln	Lys Ala Lys Arg
1490	1495	1500
Pro Cys Pro Lys Cys Gly	Ser Arg Gly Thr Gln	Ile Lys Cys Gly
1505	1510	1515
Leu Thr Leu Ala Glu Phe	Glu Glu Glu His Tyr	Lys Lys Ile Tyr
1520	1525	1530
Ile Leu Glu Gly Gln Asp	Glu Thr Pro Met Arg	Lys Glu Glu Arg
1535	1540	1545
Gln Gln Val Thr Tyr Val	Ser Arg Gly Ala Leu	Phe Leu Arg Asn
1550	1555	1560
Leu Pro Ile Leu Ala Ser	Lys Asn Lys Tyr Leu	Leu Val Gly Asn
1565	1570	1575
Leu Gly Met Glu Leu Gln	Asp Leu Glu Ser Met	Gly Trp Ile Ile
1580	1585	1590
Arg Gly Pro Ala Val Cys	Lys Lys Ile Ile His	His Glu Lys Cys
1595	1600	1605

-continued

Arg	Pro	Ser	Ile	Pro	Asp	Lys	Leu	Met	Ala	Phe	Phe	Gly	Ile	Met
1610						1615						1620		
Pro	Arg	Gly	Val	Thr	Pro	Arg	Ala	Pro	Thr	Arg	Phe	Pro	Val	Ser
1625						1630						1635		
Leu	Leu	Lys	Ile	Arg	Arg	Gly	Phe	Glu	Thr	Gly	Trp	Ala	Tyr	Thr
1640						1645						1650		
His	Pro	Gly	Gly	Val	Ser	Ser	Val	Met	His	Val	Thr	Ala	Gly	Ser
1655						1660						1665		
Asp	Ile	Tyr	Val	Asn	Asp	Ser	Ile	Gly	Arg	Thr	Lys	Ile	Gln	Cys
1670						1675						1680		
Gln	Asp	Lys	Asn	Thr	Thr	Thr	Asp	Glu	Cys	Glu	Tyr	Gly	Val	Lys
1685						1690						1695		
Thr	Asp	Ser	Gly	Cys	Ser	Asp	Gly	Ala	Arg	Cys	Tyr	Val	Ile	Asn
1700						1705						1710		
Pro	Glu	Ala	Thr	Asn	Ile	Ala	Gly	Thr	Lys	Gly	Ala	Met	Val	His
1715						1720						1725		
Leu	Arg	Lys	Ala	Gly	Gly	Glu	Phe	Asn	Cys	Val	Thr	Ala	Gln	Gly
1730						1735						1740		
Thr	Pro	Ala	Phe	Tyr	Asn	Leu	Lys	Asn	Leu	Lys	Gly	Trp	Ser	Gly
1745						1750						1755		
Leu	Pro	Ile	Phe	Glu	Ala	Ala	Thr	Gly	Arg	Val	Val	Gly	Arg	Val
1760						1765						1770		
Lys	Ala	Gly	Lys	Asn	Thr	Asp	Asn	Ala	Pro	Thr	Thr	Ile	Met	Ser
1775						1780						1785		
Gly	Thr	Gln	Val	Ala	Lys	Pro	Ser	Glu	Cys	Asp	Leu	Glu	Ser	Val
1790						1795						1800		
Val	Arg	Lys	Leu	Glu	Thr	Met	Asn	Arg	Gly	Glu	Phe	Lys	Gln	Val
1805						1810						1815		
Thr	Leu	Ala	Thr	Gly	Ala	Gly	Lys	Thr	Thr	Met	Leu	Pro	Lys	Leu
1820						1825						1830		
Leu	Ile	Glu	Ser	Ile	Gly	Arg	His	Lys	Arg	Val	Leu	Val	Leu	Ile
1835						1840						1845		
Pro	Leu	Arg	Ala	Ala	Ala	Glu	Gly	Val	Tyr	Gln	Tyr	Met	Arg	Thr
1850						1855						1860		
Lys	His	Pro	Ser	Ile	Ser	Phe	Asn	Leu	Arg	Ile	Gly	Asp	Leu	Lys
1865						1870						1875		
Glu	Gly	Asp	Met	Ala	Thr	Gly	Ile	Thr	Tyr	Ala	Ser	Tyr	Gly	Tyr
1880						1885						1890		
Phe	Cys	Gln	Met	Asp	Met	Pro	Arg	Leu	Glu	Asn	Ala	Met	Lys	Glu
1895						1900						1905		
Tyr	His	Tyr	Ile	Phe	Leu	Asp	Glu	Tyr	His	Cys	Ala	Thr	Pro	Glu
1910						1915						1920		
Gln	Leu	Ala	Val	Met	Ser	Lys	Ile	His	Arg	Phe	Gly	Glu	Ser	Val
1925						1930						1935		
Arg	Val	Ile	Ala	Met	Thr	Ala	Thr	Pro	Ser	Gly	Thr	Val	Ser	Thr
1940						1945						1950		
Thr	Gly	Gln	Lys	Phe	Thr	Ile	Glu	Glu	Val	Val	Val	Pro	Glu	Val
1955						1960						1965		
Met	Lys	Gly	Glu	Asp	Leu	Ala	Asp	Asp	Tyr	Ile	Glu	Ile	Ala	Gly
1970						1975						1980		

-continued

Leu	Lys	Val	Pro	Lys	Lys	Glu	Leu	Glu	Gly	Asn	Val	Leu	Thr	Phe
1985						1990					1995			
Val	Pro	Thr	Arg	Lys	Met	Ala	Ser	Glu	Thr	Ala	Lys	Lys	Leu	Thr
2000						2005					2010			
Thr	Gln	Gly	Tyr	Asn	Ala	Gly	Tyr	Tyr	Phe	Ser	Gly	Glu	Asp	Pro
2015						2020					2025			
Ser	Ser	Leu	Arg	Thr	Thr	Thr	Ser	Lys	Ser	Pro	Tyr	Ile	Val	Val
2030						2035					2040			
Ala	Thr	Asn	Ala	Ile	Glu	Ser	Gly	Val	Thr	Leu	Pro	Asp	Leu	Asp
2045						2050					2055			
Thr	Val	Ile	Asp	Thr	Gly	Met	Lys	Cys	Glu	Lys	Arg	Leu	Arg	Ile
2060						2065					2070			
Glu	Asn	Lys	Ala	Pro	Tyr	Ile	Val	Thr	Gly	Leu	Lys	Arg	Met	Ala
2075						2080					2085			
Ile	Thr	Thr	Gly	Glu	Gln	Ala	Gln	Arg	Lys	Gly	Arg	Val	Gly	Arg
2090						2095					2100			
Val	Lys	Pro	Gly	Arg	Tyr	Leu	Arg	Gly	Pro	Glu	Asn	Thr	Ala	Gly
2105						2110					2115			
Glu	Lys	Asp	Tyr	His	Tyr	Asp	Leu	Leu	Gln	Ala	Gln	Arg	Tyr	Gly
2120						2125					2130			
Ile	Gln	Asp	Ser	Ile	Asn	Ile	Thr	Lys	Ser	Phe	Arg	Glu	Met	Asn
2135						2140					2145			
Tyr	Asp	Trp	Ala	Leu	Tyr	Glu	Glu	Asp	Pro	Leu	Lys	Ile	Ala	Gln
2150						2155					2160			
Leu	Glu	Leu	Leu	Asn	Thr	Leu	Leu	Ile	Ser	Arg	Asp	Leu	Pro	Val
2165						2170					2175			
Val	Thr	Lys	Asn	Leu	Met	Ala	Arg	Thr	Thr	His	Pro	Glu	Pro	Ile
2180						2185					2190			
Gln	Leu	Ala	Tyr	Asn	Ser	Leu	Glu	Thr	Pro	Val	Pro	Val	Ala	Phe
2195						2200					2205			
Pro	Lys	Val	Lys	Asn	Gly	Glu	Val	Thr	Asp	Ala	His	Glu	Thr	Tyr
2210						2215					2220			
Glu	Leu	Met	Thr	Cys	Arg	Lys	Leu	Glu	Lys	Asp	Pro	Pro	Ile	Tyr
2225						2230					2235			
Leu	Tyr	Ala	Thr	Glu	Glu	Glu	Asp	Leu	Val	Val	Asp	Ile	Leu	Gly
2240						2245					2250			
Leu	Lys	Trp	Pro	Asp	Ala	Thr	Glu	Arg	Ala	Val	Leu	Glu	Val	Gln
2255						2260					2265			
Asp	Ala	Leu	Gly	Gln	Ile	Thr	Gly	Leu	Ser	Ala	Gly	Glu	Thr	Ala
2270						2275					2280			
Leu	Leu	Ile	Ala	Leu	Leu	Gly	Trp	Val	Gly	Tyr	Glu	Ala	Leu	Val
2285						2290					2295			
Lys	Arg	His	Val	Pro	Met	Val	Thr	Asp	Ile	Tyr	Thr	Leu	Glu	Asp
2300						2305					2310			
Glu	Lys	Leu	Glu	Asp	Thr	Thr	His	Leu	Gln	Phe	Ala	Pro	Asp	Asp
2315						2320					2325			
Leu	Asn	Asn	Ser	Asp	Thr	Ile	Glu	Leu	Gln	Asp	Leu	Ser	Asn	His
2330						2335					2340			
Gln	Ile	Gln	Gln	Ile	Leu	Glu	Gly	Gly	Lys	Glu	Tyr	Val	Gly	Gln
2345						2350					2355			
Ala	Tyr	Gln	Phe	Leu	Arg	Leu	Gln	Ala	Glu	Arg	Ala	Ala	Asn	Ser

-continued

2360	2365	2370
Asp Lys Gly Lys Lys Ala Met Ala Ala Ala Pro Leu Leu Ala His 2375 2380 2385		
Lys Phe Leu Glu Tyr Leu Gln Glu His Ala Gly Asp Ile Lys Lys 2390 2395 2400		
Tyr Gly Leu Trp Gly Val His Thr Ala Leu Tyr Asn Ser Ile Lys 2405 2410 2415		
Glu Arg Leu Gly His Glu Thr Ala Phe Ala Ser Leu Val Ile Lys 2420 2425 2430		
Trp Ile Ala Phe Ser Ser Asp Gly Val Pro Gly Met Ile Lys Gln 2435 2440 2445		
Ala Ala Val Asp Leu Val Val Tyr Tyr Ile Ile Asn Arg Pro Glu 2450 2455 2460		
Tyr Gln Gly Asp Lys Glu Thr Gln Asn Ala Gly Arg Gln Phe Val 2465 2470 2475		
Gly Ser Leu Phe Val Ser Cys Leu Ala Glu Tyr Thr Phe Lys Asn 2480 2485 2490		
Phe Asn Lys Ser Ala Leu Glu Gly Leu Ile Glu Pro Ala Leu Ser 2495 2500 2505		
Tyr Leu Pro Tyr Ala Ser Ser Ala Leu Lys Leu Phe Leu Pro Thr 2510 2515 2520		
Arg Leu Glu Ser Val Val Ile Leu Ser Thr Thr Ile Tyr Arg Thr 2525 2530 2535		
Tyr Leu Ser Ile Arg Lys Gly Ser Ser Gln Gly Leu Ala Gly Leu 2540 2545 2550		
Ala Val Ser Ser Ala Met Glu Ile Met Asn Gln Asn Pro Ile Ser 2555 2560 2565		
Val Ala Ile Ala Leu Ala Leu Gly Val Gly Ala Ile Ala Ala His 2570 2575 2580		
Asn Ala Ile Glu Ser Ser Glu Ala Lys Arg Thr Leu Leu Met Lys 2585 2590 2595		
Val Phe Val Lys Asn Phe Leu Asp Gln Ala Ala Thr Asp Glu Leu 2600 2605 2610		
Val Lys Glu Asn Pro Glu Lys Ile Ile Met Ala Val Phe Glu Gly 2615 2620 2625		
Ile Gln Thr Ala Gly Asn Pro Leu Arg Leu Val Tyr His Leu Tyr 2630 2635 2640		
Ala Met Phe Tyr Lys Gly Trp Thr Ala Ala Glu Ile Ala Glu Lys 2645 2650 2655		
Thr Ala Gly Arg Asn Ile Phe Val Leu Thr Ile Phe Glu Gly Leu 2660 2665 2670		
Glu Met Leu Gly Leu Asp Ala Asp Ser Lys Trp Arg Asn Leu Ser 2675 2680 2685		
Ser Asn Tyr Leu Ile Asp Ala Val Lys Lys Ile Ile Glu Lys Met 2690 2695 2700		
Thr Lys Thr Ala Thr Ser Phe Thr Tyr Ser Phe Leu Lys Ser Leu 2705 2710 2715		
Leu Pro Ala Pro Phe Ser Cys Thr Lys Ser Glu Arg Asp Pro Arg 2720 2725 2730		
Ile Gly Trp Pro Gln Lys Asp Tyr Asp Tyr Leu Glu Val Arg Cys 2735 2740 2745		

-continued

Ala Cys Gly Tyr Asn Arg Arg Ala Ile Lys Arg Asp Ser Gly Pro 2750 2755 2760
Val Leu Trp Glu Thr Leu Glu Glu Thr Gly Pro Glu Tyr Cys His 2765 2770 2775
Asn Arg Gly Glu Arg Gly Leu Ser Asn Val Lys Thr Thr Arg Cys 2780 2785 2790
Phe Val Gln Gly Glu Glu Ile Pro Pro Ile Ala Leu Arg Lys Gly 2795 2800 2805
Val Gly Glu Met Leu Val Lys Gly Val Ser Phe Arg Ile Asp Phe 2810 2815 2820
Asp Lys Asp Lys Ile Leu Ser Thr Asp Lys Trp Lys Val Pro His 2825 2830 2835
Arg Ala Val Thr Ser Ile Phe Glu Asp Trp Gln Gly Ile Gly Tyr 2840 2845 2850
Arg Glu Ala Tyr Leu Gly Thr Lys Pro Asp Tyr Gly Gly Leu Val 2855 2860 2865
Pro Arg Ser Cys Val Thr Val Thr Lys Gln Gly Leu Thr Phe Leu 2870 2875 2880
Lys Thr Ala Arg Gly Met Ala Phe Thr Thr Asp Leu Thr Ile Gln 2885 2890 2895
Asn Ile Lys Met Leu Ile Ala Thr Cys Phe Lys Asn Lys Val Lys 2900 2905 2910
Glu Gly Glu Ile Pro Ala Thr Ile Glu Gly Glu Thr Trp Ile Asn 2915 2920 2925
Ile Pro Leu Val Asn Glu Asp Thr Gly Thr Ile Lys Pro Ser Phe 2930 2935 2940
Gly Glu Arg Val Ile Pro Glu Pro Tyr Glu Glu Asp Pro Leu Glu 2945 2950 2955
Gly Pro Ser Val Ile Val Glu Thr Gly Gly Ile Ala Ile Asn Gln 2960 2965 2970
Ile Gly Val Asn Pro Gln Ser Ser Thr Cys Gly Thr Val Phe Thr 2975 2980 2985
Ala Val Lys Asp Leu Cys Gln Thr Val Ser Asn Lys Ala Lys Asn 2990 2995 3000
Ile Lys Ile Gly Phe Ser Glu Gly Gln Tyr Pro Gly Pro Gly Val 3005 3010 3015
Ala Lys Lys Thr Leu Asn Gln Leu Ile Gln Asp Glu Asp Pro Lys 3020 3025 3030
Pro Phe Ile Phe Val Cys Gly Ser Asp Lys Ser Met Ser Asn Arg 3035 3040 3045
Ala Lys Thr Ala Arg Asn Ile Lys Arg Ile Thr Thr Thr Thr Pro 3050 3055 3060
Glu Lys Phe Arg Asp Leu Ala Lys Asn Lys Lys Leu Ile Ile Val 3065 3070 3075
Leu Leu Gly Asp Arg Tyr His Glu Asp Ile Glu Lys Tyr Ala Asp 3080 3085 3090
Phe Lys Gly Thr Phe Leu Thr Arg Gln Thr Leu Glu Ala Leu Ala 3095 3100 3105
Ser Ala Lys Ala Val Glu Lys Asp Met Thr Lys Lys Glu Ala Ala 3110 3115 3120

-continued

Arg Val 3125	Leu Ala Met Glu 3130	Glu Lys Asp Glu Leu 3135	Leu Pro Gly
Trp Leu 3140	His Thr Asp Ala Pro 3145	Lys Phe Leu Asp Ile 3150	Thr Lys Asp
Asn Ile 3155	Thr His His Leu Ile 3160	Gly Asp Met Gln Ser 3165	Leu Arg Glu
Arg Ala 3170	Gly Glu Ile Gly Ala 3175	Lys Ala Thr Thr Gln 3180	Ile Thr Lys
Lys Gly 3185	Ser Val Tyr Thr Ile 3190	Asn Leu Ser Thr Trp 3195	Trp Glu Ser
Glu Arg 3200	Leu Ala Ser Leu Glu 3205	Pro Leu Phe Arg Glu 3210	Leu Leu Ser
Lys Cys 3215	Arg Pro Val Asp Arg 3220	Glu Thr Tyr Lys Asn 3225	Cys His Phe
Ala Thr 3230	Ala Ala Gln Leu Ala 3235	Gly Gly Asn Trp Val 3240	Pro Val Ala
Pro Val 3245	Val His Leu Gly Glu 3250	Ile Pro Val Lys Lys 3255	Lys Lys Thr
Leu Pro 3260	Tyr Glu Ala Tyr Lys 3265	Leu Leu Lys Glu Met 3270	Val Asp Ser
Glu Lys 3275	Glu Phe His Lys Pro 3280	Val Ser Arg Glu Lys 3285	His Gln Trp
Ile Leu 3290	Asn Lys Val Lys Thr 3295	Gly Gly Asp Leu Gly 3300	Leu Lys Asn
Leu Val 3305	Cys Pro Gly Arg Val 3310	Gly Glu Pro Ile Leu 3315	Arg Glu Lys
Lys Lys 3320	Phe Asn Ile Tyr Asn 3325	Lys Arg Ile Thr Ser 3330	Thr Met Leu
Ser Val 3335	Gly Ile Arg Pro Glu 3340	Lys Leu Pro Val Val 3345	Arg Ala Gln
Thr Ser 3350	Thr Lys Glu Phe His 3355	Glu Ala Ile Arg Asp 3360	Lys Ile Asp
Lys Lys 3365	Ala Asn Thr Gln Thr 3370	Pro Gly Leu His Lys 3375	Glu Leu Leu
Glu Ile 3380	Phe Asn Ser Ile Cys 3385	Ala Ile Pro Glu Leu 3390	Arg Asn Thr
Tyr Lys 3395	Glu Val Asp Trp Asp 3400	Val Leu Thr Ser Gly 3405	Ile Asn Arg
Lys Gly 3410	Ala Ala Gly Tyr Phe 3415	Glu Lys Met Asn Ile 3420	Gly Glu Ile
Ile Asp 3425	Ser Asp Lys Lys Ser 3430	Val Glu Gln Leu Ile 3435	Lys Arg Met
Lys Ser 3440	Gly Leu Glu Phe Asn 3445	Tyr Tyr Glu Thr Ala 3450	Ile Pro Lys
Asn Glu 3455	Lys Arg Ala Val Val 3460	Asp Asp Trp Met Glu 3465	Gly Asp Tyr
Val Glu 3470	Glu Lys Arg Pro Arg 3475	Val Ile Gln Tyr Pro 3480	Glu Ala Lys
Met Arg 3485	Leu Ala Ile Thr Lys 3490	Val Met Tyr Asn Trp 3495	Val Lys Gln
Lys Pro	Ile Val Ile Pro Gly	Tyr Glu Gly Lys Thr	Pro Leu Phe

-continued

3500		3505		3510
His Val Phe Asp Lys Val	His Lys Glu Trp Lys	Asn Phe Asn Ser		
3515	3520	3525		
Pro Val Ala Val Ser Phe	Asp Thr Lys Ala Trp	Asp Thr Gln Val		
3530	3535	3540		
Thr Pro Lys Asp Leu Leu	Leu Ile Ser Glu Ile	Gln Lys Tyr Tyr		
3545	3550	3555		
Tyr Lys Lys Glu Tyr His	Arg Phe Ile Asp Asn	Leu Thr Glu Lys		
3560	3565	3570		
Met Val Glu Val Pro Val	Val Cys Glu Asp Gly	Asn Val Tyr Ile		
3575	3580	3585		
Arg Glu Gly Gln Arg Gly	Ser Gly Gln Pro Asp	Thr Ser Ala Gly		
3590	3595	3600		
Asn Ser Met Leu Asn Val	Leu Thr Met Ile Tyr	Ala Phe Cys Lys		
3605	3610	3615		
Ala Asn Ser Ile Pro Tyr	Ser Ala Phe His Arg	Val Ala Lys Ile		
3620	3625	3630		
His Val Cys Gly Asp Asp	Gly Phe Leu Ile Thr	Glu Lys Ser Phe		
3635	3640	3645		
Gly Glu Ala Phe Ala Ile	Lys Gly Pro Gln Ile	Leu Met Glu Ala		
3650	3655	3660		
Gly Lys Pro Gln Lys Leu	Ile Gly Glu Phe Gly	Leu Lys Leu Ala		
3665	3670	3675		
Tyr Lys Phe Asp Asp Ile	Glu Phe Cys Ser His	Thr Pro Ile Lys		
3680	3685	3690		
Val Arg Trp Ala Asp Asn	Asn Thr Ser Tyr Met	Pro Gly Arg Asp		
3695	3700	3705		
Thr Ala Thr Ile Leu Ala	Lys Met Ala Thr Arg	Leu Asp Ser Ser		
3710	3715	3720		
Gly Glu Arg Gly Thr Glu	Gly Tyr Glu Leu Ala	Val Ala Phe Ser		
3725	3730	3735		
Phe Leu Leu Met Tyr Ser	Trp Asn Pro Leu Val	Arg Arg Ile Cys		
3740	3745	3750		
Leu Leu Val Met Ser Thr	Ile Asp Thr Lys Glu	Ala Ser Gln Asn		
3755	3760	3765		
Asn Thr Ile Tyr Thr Phe	Arg Gly Asp Pro Ile	Gly Ala Tyr Thr		
3770	3775	3780		
Glu Val Ile Gly Tyr Arg	Leu Asp Gln Leu Lys	Gln Thr Glu Phe		
3785	3790	3795		
Ser Lys Leu Ala Gln Leu	Asn Leu Ser Met Ala	Ile Leu Gln Ile		
3800	3805	3810		
Tyr Asn Lys Asn Thr Thr	Lys Arg Leu Ile Glu	Asp Cys Val Lys		
3815	3820	3825		
Leu Gly Asn Gln Asn Lys	Gln Ile Leu Val Asn	Ala Asp Arg Leu		
3830	3835	3840		
Ile Ser Lys Lys Thr Gly	Tyr Thr Tyr Glu Pro	Thr Ala Gly His		
3845	3850	3855		
Thr Lys Ile Gly Lys His	Tyr Glu Glu Ile Asn	Leu Leu Lys Asp		
3860	3865	3870		
Thr Pro Gln Lys Thr Val	Tyr Gln Gly Thr Glu	Arg Tyr		
3875	3880	3885		

-continued

```

<210> SEQ ID NO 3
<211> LENGTH: 418
<212> TYPE: DNA
<213> ORGANISM: PMC Virus

<400> SEQUENCE: 3
gtataacgac agtagttaa gtgtcgttat gcatcattgg ccataacaaa ttatctaatt    60
tggaataggg acctgcgacc tgtacgaagg ccgagcgtcg gtagccattc cgactagtag    120
gactagtaca aataggtcaa ctggttgagc aggtgagtgt gctgcagcgg ctaagcggtg    180
agtacaccgt attcgtcaac aggtgctact ggaaggatc acccactagc gatgcctgtg    240
tggacgagga catgtccaag ccaatgttat cagtagcggg ggtagttact gagaaagctg    300
cccagaatgg gtatgtgac atacagctcg ataggatgcc ggcggatgcc ctgtattttg    360
accagtataa atattatccg ttgtaaagca tatgaatact tttactttta atacatat    418

<210> SEQ ID NO 4
<211> LENGTH: 504
<212> TYPE: DNA
<213> ORGANISM: PMC Virus

<400> SEQUENCE: 4
ggagggagtg aggaagaaa catgttcttt agaactgcac ccacgccgcc accaggggtg    60
caagaaccgg ttacacaaag cacaatgaga ccaatttttg gcgaacccca tccaccteta    120
cacaacaca gcacgttaaa attgccacat tggaggggga tcaaaacaat tagagttaag    180
aagagagaat tgccaagaa gggcgattgt agcaactcaa caacagctcc cacttcgggg    240
gtgtacgttg aattaggggc tgtgttctat aaagattaca cgggcacggg ataccatcgt    300
gtaccgctag aactttgtac aaaccaagag aggtgcgagg gatccaagtg tgtagggaga    360
atgacagggg ctgatggcag gttgtacaac gttttagtat gtccggacga ttgtatcctc    420
tttgagagac actgtagagg tcaaacagtc gtccctgaaat ggatttccaa ccccttgaca    480
tcaccacttt gggtcagag ttgt                    504

<210> SEQ ID NO 5
<211> LENGTH: 297
<212> TYPE: DNA
<213> ORGANISM: PMC Virus

<400> SEQUENCE: 5
tctgacgaca aaggagcaaa acccaagggt aaaccaaag acgacaggat gaagcaagga    60
aaaatagtga caaagcctaa agagactgaa gcagatcaaa aaactagacc accagatgcc    120
acgatagtgg ttgacgggca gaagtatcag gtgaggaaga aggggaaagc gaaacccaag    180
actcaagacg gcttatacca caacaagaac aaaccagaag cgtccaggaa gaagcttgag    240
aaggccttgc tagcatgggc aatattagcc tgcctattgg tggtaccggt aggggtcc    297

<210> SEQ ID NO 6
<211> LENGTH: 666
<212> TYPE: DNA
<213> ORGANISM: PMC Virus

<400> SEQUENCE: 6
accaacgtga cacaatggaa cttatgggac aataaaagta ctacagacat acatagcgtc    60

```

-continued

```

atgttttcta gagggattaa aaggagtctg catggaattt ggcccacaca aatctgcaaa 120
gggatcccta cacatctagc agcagactat gaactgaaaa ggattcacgg gatggtggat 180
gcaagcccca tgaccaactt cacatgttgt aggctacaga gacatgagtg gaacaagcat 240
gggtggtgca actggtacaa tatagagccg tggatcaatc tcatgaataa taccaagga 300
ctattaaaca ctggagacaa ttctactgag tgcgcagtca catgcaggta tgatgcagac 360
ttaggggtga atatagtgac tcaagccagg actactccaa ctatcctgac tggctgtaag 420
aaagggcaca acttctcttt ctcaggggag gtcagggcct caccctgcaa ctttgagtta 480
actgctgaag acttgctcag gatcatggat cacaccaact gcgagggatt tacctacttc 540
ggggaaggaa tcgttgacgg ttacaccgag gtagtagaga aggccaggtc aagtggtttc 600
agggtctca catggttgtc gagtaagatt gaaaacacca agaaaaaat attcggagct 660
gaagcc 666

```

```

<210> SEQ ID NO 7
<211> LENGTH: 588
<212> TYPE: DNA
<213> ORGANISM: PMC Virus

```

```

<400> SEQUENCE: 7

```

```

agtccttact gccagtggc taagagggtc ttcaacatta tttatacaa caattgcacc 60
ccgcttgac tgccagataa gtcaaaaatt ataggaccag gaaccttga catcagtggc 120
agggatgaat tcatatttcc aaaactcccc taccacgtag atgacttcat tctactgagc 180
ttaattgcaa tgtctgattt tgetccagag acatcaagta taatctacct ggctttgac 240
tacctaatgc caagtaatga caacagggac ttctgtgatgg acctggaccc aaataaacta 300
aaccttactg caactaaatc cgtggcaagt gtggtcccta catcgggtgaa tgtgttaggt 360
gaatgggtgt gcgtcaaac aagtgtgtgg ccttattccg cggaaatcac taatctgata 420
ggagggtgca tcacctgggc agacttagtt atcaagacca ttgaagaatt gctaaatttg 480
tggaccgaag caacagctgt ggcatttctg gctgctctaa taaaaattt tagaggccag 540
ccgatccaag cggtagcatg gttaatcadc atagggggag cacaagcc 588

```

```

<210> SEQ ID NO 8
<211> LENGTH: 1131
<212> TYPE: DNA
<213> ORGANISM: PMC Virus

```

```

<400> SEQUENCE: 8

```

```

caaacctgca accctgaatt catgtacgca ttagcgaaaa ataccagcat aggttcatta 60
ggaccagaat cactgacgac aaggtggtac caactaacca gcggtttcaa actcactgac 120
agcacgattg aagtcacctg tgtgggtgct aacatgagga ttcatgtagt gtgccactt 180
gtaagtgaca gatatttggc cataaacccac cctagagcac tgccaacaac ggcgtggttc 240
aggaaaatac acactcagca tgaggtacca agagaaagaa tcatgagtga gtcaaaaagg 300
aggtacactt gtccttgtgg ttctaaacca gtggtgaggt caacaacaca attcaacca 360
atatctatat ctaccceaag ctttgaactt gaatgccta ggggttgac tggggtgta 420
gagtgtacac tagtctccc atcaactctg acaacagaga ctatattcac atacaggaag 480
cccaaaccat tcgacttga aaactggtgc aagtatacag tggtgagaa agggatcctg 540

```

-continued

```

tattcttgta aatttggggg caattcaaca tgcacaaaag ggcttatagt taaagggcaa    600
cggaagaca aagtaaggta ctgtgaatgg tgtgggtata agttcagttc accaaatgga    660
ctgcctcagt atccactggg attgtgtgag aaagaacaat cagaaggact cagggattat    720
ggtgacttcc catgctgcaa caacggcact tgtattgaca aagaaggtag tgtgcaatgc    780
tacatagggg ataagaaagt taccgtgaag ctgtataatg cctcactatt ggcccccatg    840
ccctgcaaac ccatagtgta taactcccag gggccccag cgcctaagac ctgcacttat    900
aggtgggct caacattaga aaataaatat tatgaacca gggacagcta ctaccagcaa    960
tacattataa agtcagggtg tcaatattgg tttgatctca cagcaaagga tcatgtggca  1020
gactggatca caaaactact tccaataata atagtggcct tgtagggggg cagaggcacc  1080
ttgtgggtgt tgatagctta tgagttgcta actcagtatg aggtagtagg a          1131

```

```

<210> SEQ ID NO 9
<211> LENGTH: 216
<212> TYPE: DNA
<213> ORGANISM: PMC Virus

```

```

<400> SEQUENCE: 9

```

```

gacgagaaca tagtggctca agctgaagcc ctggtaatcg gaaacatctt gatgagtta    60
gacttagaga taattagctg cttccttctg ttgttgatcg tggtgaaaaa acaagctgtc    120
aggagaacgt tggctttact gtttcattgg ataactatga acccattcca gtcagtaatg    180
atcacagtgg tctacttctg cggtttggg agggcc          216

```

```

<210> SEQ ID NO 10
<211> LENGTH: 1404
<212> TYPE: DNA
<213> ORGANISM: PMC Virus

```

```

<400> SEQUENCE: 10

```

```

gaagagggaa ctaaaggagg tagtacaagc gggccaccaa tccatgtagt tgcaatactg    60
ttattcctct tgtaccacac agtgaagtat aaggacttta acatagcaat gatccttact    120
ataacattgt cctgaaaag ctcacctac atacatacca gcttgatga aattccattg    180
cttgtggctg taataagtct cacatgctcc atatacatt ttgacttga ggtaaagagc    240
aagctagtgg cccaactat aggtataatt ggagttacc tagcaatgag agttttgtgg    300
ctggtaaagg aaatgactat accaaccctc tctgtgtcca ttagtctgat agatccaaag    360
atggtcataa tactctactt gatatcccta actattacag tcaatcaca cctagaccta    420
gcaagttatt gcttgaact gggacctttt atcctatcat tccatacaat gtgggtggat    480
gttgcacatc tctgtctcat gctgccttgg tacgaactag taaaagtcta ctacctaaaa    540
aagaagaaag aggatgtgga aacatggctc caaaattcag gaatatccac ccaagaaact    600
tcccatacag gatttgattt ttctagcccc ggggaggagg tgcacacact accaatgcaa    660
aataaaacca aattttgtag gactgcttac atgactgtac taagggtttt ggtgataaca    720
gccatcagca gtgtctggaa accaataatt ttagcagaac tcctaataga ggcagtgtat    780
tggacacaca ttaaaatagc caaagaattg gcgggttcaa gcaggttcgt tgctaggttc    840
attgcatcta ttatagagtt gaattgggccc atggacgaaa aagaagcacc tcggtacaaa    900
agattttacc tattatcatc caaaataaca gatctaattg ttaagcaca aatccaaaat    960

```

-continued

```

gagacagtaa aatcctgggt tgaagaaact gaaatatttg gaatacaaaa agtggcaatg 1020
gtgataaggg ctcaattctct gagtttggag ccaaatgcca tcctttgctc cgttttgtaa 1080
gaaaaacaaa atcaaaaagc caaaaggccc tgcctaagt gtggtagtag aggcactcaa 1140
ataaagtgtg ggctgacact ggccgagttt gaggaagaac attacaaaaa aatatacatc 1200
ctcgaaggcc aagatgaaac tcccatgagg aaagaagaaa gacagcaagt aacttatgtc 1260
tctaggggtg ctctgttctc taggaatctt cctatcttag cttcaaaaaa caaataccta 1320
ctttaggca atctgggat ggaattgcaa gatttggaaa gtatgggatg gatcattcga 1380
gggccagccg tctgcaagaa gata 1404

```

```

<210> SEQ ID NO 11
<211> LENGTH: 2028
<212> TYPE: DNA
<213> ORGANISM: PMC virus

```

```

<400> SEQUENCE: 11

```

```

ataccatg agaaatgacg gccttcaata ccagacaaac tcatggcatt ctctgggatt 60
atgcctaggg gagttacacc aagagccctt acacggttcc ctgtgtcctt gctgaagata 120
agacgggggt ttgagaccgg ctgggcctac acacaccctg gaggggtaag tagtgtgatg 180
catgtcaccg ctgggtcggg tatatatgtc aatgactcaa tagggaggac aaaaatccag 240
tgccaagaca aaaacactac aacagatgag tgtgaatatg gtgtgaaaac agactcaggg 300
tgctctgatg gagctcgggt ctatgtcctc aaccctgaag caaccaacat agcaggggacc 360
aagggggcca tggtagacct gaggaaagct ggaggagagt tcaactgcgt gactgcccag 420
ggtagccccc ccttctataa tctaagaaac ttaaaaggat ggtagggcct gcctatcttt 480
gaagctgcca caggaagagt ggtaggaagg gtaaaagcag gaaaaaacac tgacaatgct 540
ccaacaacca ttatgtcagg gacgcaagtg gcaaaacct cagagtgtga cctagaatca 600
gtggtgagga aactagagac aatgaacaga ggggaattca aacaagtgac tctggctaca 660
ggcgcaggaa agacaacct gctaccaaaag ctgttaatat aatccatagg caggcataag 720
agagtgttag tactgatccc gttgagagct gcagcggagg ggggtgacca gtacatgaga 780
accaaaccac caagcatatc tttcaacttg aggatagggg atctgaaaga aggtgacatg 840
gcaactggga tcacctatgc ctcttatggg tacttctgcc aaatggacat gcctagactg 900
gagaatgcaa tgaaggaata ccactatatt ttcttggatg aatatcactg tgccacacca 960
gaacagttag cagtgtatgc aaaaatacat aggttcgggtg aatcagttag ggtaatagcc 1020
atgaccgcca cgccatccgg gactgtgagc acaacagggc agaaattcac aattgaggag 1080
gtggtagtag ctgaagtgat gaagggggag gaccttctgt atgattacat cgaaatagca 1140
gggttgaagg tgccaagaa agagtttagg ggtaacgtac tgacttttgt gcctacaagg 1200
aagatggcat cgaaaacagc aaaaaatta accacacagg gatacaatgc tggatactac 1260
ttcagtggag aagatccatc atccctgcgg acaactactt ctaagtcacc atatatagta 1320
gttgcaacca atgccattga atccggggta accttaccgg accttgatac agtaatagat 1380
acaggcatga agtgtgaaaa gagactaaga atcgaaaaca aagctcccta catcgtaaca 1440
ggactgaaaa gaatggctat aacaacgggg gagcaagctc aaagaaaaag tagggtaggc 1500
agggttaaac ctgggaggta cttgagagga cctgaaaaca ctgcagggtg aaaggactat 1560

```

-continued

```

cactatgacc tttacaggc acagaggtag ggcattccaag actcaataaa catcaccaag 1620
tctttcaggg agatgaacta tgattgggca ttatatgagg aagaccggtt aaagattgcc 1680
caattagagt tgtaaacac actcctgac tcaagggatc tgccagtagt acaaaaaaat 1740
ctgatggccc gcacaacaca tcccgaacct atacaattgg cttacaatag tttagaacc 1800
cctgtaccgg tggcattccc aaaagtgaaa aatggagaag tcaactgacgc acatgaaact 1860
tacgagttga tgacctgtag gaagcttgag aaagaccccc ctatatacct gtatgcaaca 1920
gaagaagaag atctcgtagt ggacatactg ggattgaaat ggccagacgc cacagagagg 1980
gctgtcttgg aagtgcgaaga cgccctgggc cagatcacag gtttatct 2028

```

```

<210> SEQ ID NO 12
<211> LENGTH: 189
<212> TYPE: DNA
<213> ORGANISM: PMC Virus

```

```

<400> SEQUENCE: 12

```

```

gcaggggaga cagctttact catagcccta ttagggtggg tgggctacga agccttggtg 60
aagagggcagc tgcctatggt gacagacata tacaccctag aagatgaaaa attggaagac 120
actacacacc tacaatttgc cccagatgat ctgaacaatt cagataccat tgagctccaa 180
gacttatcg 189

```

```

<210> SEQ ID NO 13
<211> LENGTH: 1041
<212> TYPE: DNA
<213> ORGANISM: PMC Virus

```

```

<400> SEQUENCE: 13

```

```

aatcaccaaa tccaacaaat tctagaaggt ggaaggaat atgtcgcca agcctaccaa 60
ttcctcaggt tgcaagctga gagggtgcc aactcagaca aaggcaagaa agcaatggca 120
gcgccccat tactagccca caagttcctg gaatacttgc aagagcatgc aggtgacata 180
aagaagtatg gtctatgggg ggtccacaca gcattgtata acagcataaa agaaagactg 240
ggtcacgaaa ctgcattcgc atctctgggt ataaatgga ttgccttttc ctcaagatgga 300
gtcccgggga tgattaagca agcagcagta gacttggtgg tatactatat aatcaacagg 360
cctgagtatc aaggggataa ggagacacag aatgcaggta gacaatttgt tggctccctt 420
tttgtttcat gtctagcaga gtacacattc aaaaacttca ataaatcagc attagaagga 480
ttgatcgagc ctgccttaag ctatctaccc tacgcttcaa gcgcactaaa gttattccta 540
ccgactagac ttgaaagtgt agtgatactg tccactacta tatacagaac atacttatca 600
atcaggaaag gatctagtca gggtttagcc gggctggcag ttagctcagc gatggagatc 660
atgaaccaga acccaatcag cgtggctatt gcactggcac taggagtcgg agcaatagcg 720
gcacataatg ccattgagag cagtgaggca aaaaggactc tcctgatgaa ggtcctttgtt 780
aagaactttt tggaccaagc agccactgat gagcttgtga aagagaaccc tgagaagatc 840
ataatggcag tgtttgaggg cattcaaaca gctggaaatc cattgagact tgtataccat 900
ctatatgcaa tgttctacaa aggggtgact gccgcggaaa tagctgaaaa aaccgctggt 960
aggaacattt ttgtgttaac aatattttaa ggattggaaa tgttaggcct ggatgccgac 1020
tcaaatgga gaaatctgag c 1041

```

-continued

<210> SEQ ID NO 14
 <211> LENGTH: 1515
 <212> TYPE: DNA
 <213> ORGANISM: PMC Virus

<400> SEQUENCE: 14

```
tctaattatc ttattgatgc agtgaagaaa atcattgaaa aaatgactaa aacagcaaca    60
agcttcacct acagcttttt gaaatctttg cttcctgccc ccttctcgtg tactaaatca    120
gaaagagatc caagaatagg gtggcccaaa aaagactacg actacctcga ggtccgatgc    180
gcttgtgggt ataacaggag agctataaaa agagactcag gacctgtgtt atgggagacc    240
ttagaggaga cgggtccaga gtactgccac aacagagggt aaagggggct cagcaatgtg    300
aagactacta gatgctttgt ccaaggagag gaaatccctc caattgcact gaggaagga    360
gtaggtgaga tgttggtcaa ggggtgttca ttcagaatag attttgataa agacaagata    420
ctttcaacag acaagtggaa ggtaccacat agggcagtta catcaatctt tgaggattgg    480
cagggatttg gttacagaga ggcttaccta gggaccaaac cagactatgg gggctcgtgtg    540
cccagatctt gtgtaactgt aacaaaaaaa gggttaacat tcttgaaaac tgccagaggc    600
atggctttca cgactgacct gaccatccag aacatcaaaa tgctgatagc tacatgcttc    660
aagaacaagg tgaaggaagg ggagatacca gctacgattg aaggggaaac atggatcaac    720
ataccactag tgaatgagga caccgggacc attaaaccaa gcttcgggga aagagtgatt    780
cccgaacct atgaggagga cccacttgaa ggcccaagtg taatcgttga aacaggaggc    840
atagccatca accaaaatagg ggtcaatcca caatccagta catgtggaac agtttttaca    900
gcagtgaagg atctgtgcca aacagttagt aataaagcca agaatatcaa aattgggttt    960
tcggaaggcc aataccagg tccaggggtt gcaaagaaga cactgaacca gctcatacaa   1020
gatgaagacc caaaaccatt catatttgtt tgtggctctg acaagtcaat gtctaacg    1080
gcaaaaaactg cgaggaacat caagagaatc accaccacaa cacctgagaa attcagagac   1140
ttggcaaaaa acaagaattt gataattgtg ctgttaggtg atagatacca tgaagatata   1200
gaaaagtatg cagacttcaa gggcaccttc ttgaccagac aaaccttggga agcactagca   1260
agtgccaaag ctgtagagaa ggacatgacc aagaaagaag cagcaagagt attggcaatg   1320
gaagaaaagg atgaactaga actcccaggg tggctgcata cagatgcacc caaatccta   1380
gacattacta aggacaacat cacacatcac ctaatagggg acatgcagag tctgagagaa   1440
agagcagggg agataggagc aaaggccacc actcaaatca ctaagaaagg gagtgtatac   1500
acaatcaatc tgagt                                     1515
```

<210> SEQ ID NO 15
 <211> LENGTH: 2080
 <212> TYPE: DNA
 <213> ORGANISM: PMC Virus

<400> SEQUENCE: 15

```
acgtggtggg agtcagagag gttggcatct ttggaacctt tgttccggga actactatct    60
aaatgcaggc cagtggacag ggagacatat aagaattgtc attttgcaac agcagcccaa   120
cttgccggag gaaactgggt accggtagca ccagttgtac atcttgggga aattccggta   180
aagaagaaaa agactctccc ctacagggca tacaagctcc taaagagat ggttgactcg   240
```

-continued

```

gagaaggaat tccataaacc agtgagcagg gaaaaacacc aatggatact gaacaaagtg 300
aaaactgggtg gtgacctcgg cttaaaaaat ctagtatgtc caggtagggt tggagaacca 360
atcctaagag agaagaagaa attcaacatt tacaacaaga ggattaccag tactatgtta 420
tcagtaggga taaggccaga aaaattgcca gtggtaagag cccagaccag taccaagaa 480
tttcatgaag caataagga caaaatagac aaaaaagcaa acacacagac cccaggccta 540
cacaagaat tgttggagat attcaactca atatgtgcca tccccgaact tagaaatacc 600
tacaagagg ttgattggga cgttctaacc tcaggcataa ataggaaagg tgcagccggg 660
tacttcgaaa aaatgaactc aggggagatc atagatagtg acaaaaaatc agtggaaaca 720
ctcataaaga gaatgaaatc agggctagaa ttcaactact atgagactgc aataccaaaa 780
aatgagaaga gggcagtggg agatgattgg atggaagggt actatgtaga agaaaaaga 840
ccaagagtca tacagtatcc tgaggcaaaag atgagattag ctataacca agtaatgtat 900
aactgggtca agcagaagcc tatagtaatc cctggatagc aaggtaaagc tcctttgttt 960
catgttttcg acaaggtcca caaagaatgg aaaaatttca acagtccagt tgcagtcagt 1020
tttgacacta aagcctggga cacacaagta acaccaaaag accttctcct catatcagaa 1080
atccaaaagt attattacaa gaaagaatac catagattca tagataatth gaccgagaaa 1140
atggtggagg taccagtggg ttgtgaagac ggaaacgtct acataagaga aggtcagagg 1200
ggaagtggtc aaccagacac tagcgcaggt aatagtatgt tgaatgtact gactatgata 1260
tatgccttct gcaaagctaa tcctcatcct tactcagcct tccacagggt agcaaagata 1320
catgtgtgtg gagatgatgg tttcttgata actgagaaaa gttttgtga ggcctttgcg 1380
atcaaggggc ctcaaattht gatggaagca ggaaaaccac aaaaacttat aggtgaatth 1440
ggactgaaat tggcatataa atttgatgac attgaattht gctcgcatac accaataaag 1500
gtcaggtggg ctgacaacaa cacatcatac atgcccggaa gagacacagc taccattcta 1560
gctaaaatgg caaccgcct tgactctagt ggggagaggg ggaccgaggg atacgagctg 1620
gccgtggcct tcagtttctt actaatgtat tcttggaaac ccctggtaag aagaatatgc 1680
ctgcttgc tgtctacaat tgacacaaaa gaagctagcc aaaataacac tatatataca 1740
tttagggggg atcccatagc tgcctacaca gaggtaattg ggtataggct ggaccaacta 1800
aaacagacag agttctctaa atggctcag ctgaattht caatggcaat acttcaataa 1860
tacaataaaa acacaaccaa gagactcacc gaagattgtg tgaacttgg caacaaaat 1920
aagcaaatat tggatgaatc agaccgtttg atcagcaaga aaacgggcta cacatatgag 1980
ccaacagctg gccacactaa gataggcaag cactatgaag aaatcaacct gctgaaagat 2040
acaccacaaa aaactgtcta ccaaggaact gaaaggtata 2080

```

<210> SEQ ID NO 16
 <211> LENGTH: 167
 <212> TYPE: PRT
 <213> ORGANISM: PMC Virus

<400> SEQUENCE: 16

```

Gly Gly Ser Glu Glu Gly Asn Met Phe Phe Arg Thr Ala Pro Thr Pro
1           5           10           15
Pro Pro Gly Cys Gln Glu Pro Val Tyr Thr Ser Thr Met Arg Pro Ile
20           25           30

```

-continued

Phe Gly Glu Pro His Pro Pro Leu His Lys His Ser Thr Leu Lys Leu
 35 40 45
 Pro His Trp Arg Gly Ile Lys Thr Ile Arg Val Lys Lys Arg Glu Leu
 50 55 60
 Pro Lys Lys Gly Asp Cys Ser Asn Ser Thr Thr Ala Pro Thr Ser Gly
 65 70 75 80
 Val Tyr Val Glu Leu Gly Ala Val Phe Tyr Lys Asp Tyr Thr Gly Thr
 85 90 95
 Val Tyr His Arg Val Pro Leu Glu Leu Cys Thr Asn Gln Glu Arg Cys
 100 105 110
 Glu Gly Ser Lys Cys Val Gly Arg Met Thr Gly Ser Asp Gly Arg Leu
 115 120 125
 Tyr Asn Val Leu Val Cys Pro Asp Asp Cys Ile Leu Phe Glu Arg His
 130 135 140
 Cys Arg Gly Gln Thr Val Val Leu Lys Trp Ile Ser Asn Pro Leu Thr
 145 150 155 160
 Ser Pro Leu Trp Val Gln Ser
 165

<210> SEQ ID NO 17
 <211> LENGTH: 99
 <212> TYPE: PRT
 <213> ORGANISM: PMC Virus

<400> SEQUENCE: 17

Ser Asp Asp Lys Gly Ala Lys Pro Lys Val Lys Pro Lys Asp Asp Arg
 1 5 10 15
 Met Lys Gln Gly Lys Ile Val Thr Lys Pro Lys Glu Thr Glu Ala Asp
 20 25 30
 Gln Lys Thr Arg Pro Pro Asp Ala Thr Ile Val Val Asp Gly Gln Lys
 35 40 45
 Tyr Gln Val Arg Lys Lys Gly Lys Ala Lys Pro Lys Thr Gln Asp Gly
 50 55 60
 Leu Tyr His Asn Lys Asn Lys Pro Glu Ala Ser Arg Lys Lys Leu Glu
 65 70 75 80
 Lys Ala Leu Leu Ala Trp Ala Ile Leu Ala Cys Leu Leu Val Val Pro
 85 90 95
 Val Gly Ser

<210> SEQ ID NO 18
 <211> LENGTH: 222
 <212> TYPE: PRT
 <213> ORGANISM: PMC Virus

<400> SEQUENCE: 18

Thr Asn Val Thr Gln Trp Asn Leu Trp Asp Asn Lys Ser Thr Thr Asp
 1 5 10 15
 Ile His Ser Val Met Phe Ser Arg Gly Ile Lys Arg Ser Leu His Gly
 20 25 30
 Ile Trp Pro Thr Gln Ile Cys Lys Gly Ile Pro Thr His Leu Ala Ala
 35 40 45
 Asp Tyr Glu Leu Lys Arg Ile His Gly Met Val Asp Ala Ser Pro Met
 50 55 60

-continued

```

Thr Asn Phe Thr Cys Cys Arg Leu Gln Arg His Glu Trp Asn Lys His
65          70          75          80

Gly Trp Cys Asn Trp Tyr Asn Ile Glu Pro Trp Ile Asn Leu Met Asn
85          90          95

Asn Thr Gln Gly Leu Leu Asn Thr Gly Asp Asn Phe Thr Glu Cys Ala
100         105         110

Val Thr Cys Arg Tyr Asp Ala Asp Leu Gly Val Asn Ile Val Thr Gln
115         120         125

Ala Arg Thr Thr Pro Thr Ile Leu Thr Gly Cys Lys Lys Gly His Asn
130         135         140

Phe Ser Phe Ser Gly Glu Val Arg Ala Ser Pro Cys Asn Phe Glu Leu
145         150         155         160

Thr Ala Glu Asp Leu Leu Arg Ile Met Asp His Thr Asn Cys Glu Gly
165         170         175

Phe Thr Tyr Phe Gly Glu Gly Ile Val Asp Gly Tyr Thr Glu Val Val
180         185         190

Glu Lys Ala Arg Ser Ser Gly Phe Arg Ala Leu Thr Trp Leu Ser Ser
195         200         205

Lys Ile Glu Asn Thr Lys Lys Lys Ile Phe Gly Ala Glu Ala
210         215         220

```

<210> SEQ ID NO 19

<211> LENGTH: 196

<212> TYPE: PRT

<213> ORGANISM: PMC Virus

<400> SEQUENCE: 19

```

Ser Pro Tyr Cys Pro Val Ala Lys Arg Val Phe Asn Ile Ile Tyr Thr
1          5          10          15

Asn Asn Cys Thr Pro Leu Gly Leu Pro Asp Lys Ser Lys Ile Ile Gly
20         25         30

Pro Gly Thr Phe Asp Ile Ser Gly Arg Asp Glu Phe Ile Phe Pro Lys
35         40         45

Leu Pro Tyr His Val Asp Asp Phe Ile Leu Leu Ser Leu Ile Ala Met
50         55         60

Ser Asp Phe Ala Pro Glu Thr Ser Ser Ile Ile Tyr Leu Ala Leu His
65         70         75         80

Tyr Leu Met Pro Ser Asn Asp Asn Arg Asp Phe Val Met Asp Leu Asp
85         90         95

Pro Asn Lys Leu Asn Leu Thr Ala Thr Lys Ser Val Ala Ser Val Val
100        105        110

Pro Thr Ser Val Asn Val Leu Gly Glu Trp Val Cys Val Lys Pro Ser
115        120        125

Trp Trp Pro Tyr Ser Ala Glu Ile Thr Asn Leu Ile Gly Gly Val Ile
130        135        140

Thr Val Ala Asp Leu Val Ile Lys Thr Ile Glu Glu Leu Leu Asn Leu
145        150        155        160

Trp Thr Glu Ala Thr Ala Val Ala Phe Leu Ala Ala Leu Ile Lys Ile
165        170        175

Phe Arg Gly Gln Pro Ile Gln Ala Val Ala Trp Leu Ile Ile Ile Gly
180        185        190

Gly Ala Gln Ala
195

```

-continued

```

<210> SEQ ID NO 20
<211> LENGTH: 377
<212> TYPE: PRT
<213> ORGANISM: PMC Virus

<400> SEQUENCE: 20

Gln Thr Cys Asn Pro Glu Phe Met Tyr Ala Leu Ala Lys Asn Thr Ser
1          5          10          15

Ile Gly Ser Leu Gly Pro Glu Ser Leu Thr Thr Arg Trp Tyr Gln Leu
20          25          30

Thr Ser Gly Phe Lys Leu Thr Asp Ser Thr Ile Glu Val Thr Cys Val
35          40          45

Gly Ala Asn Met Arg Ile His Val Val Cys Pro Leu Val Ser Asp Arg
50          55          60

Tyr Leu Ala Ile Asn His Pro Arg Ala Leu Pro Thr Thr Ala Trp Phe
65          70          75          80

Arg Lys Ile His Thr Gln His Glu Val Pro Arg Glu Arg Ile Met Ser
85          90          95

Glu Ser Lys Arg Arg Tyr Thr Cys Pro Cys Gly Ser Lys Pro Val Val
100         105         110

Arg Ser Thr Thr Gln Phe Asn Pro Ile Ser Ile Ser Thr Pro Ser Phe
115        120        125

Glu Leu Glu Cys Pro Arg Gly Trp Thr Gly Ala Val Glu Cys Thr Leu
130        135        140

Val Ser Pro Ser Thr Leu Thr Thr Glu Thr Ile Phe Thr Tyr Arg Lys
145        150        155        160

Pro Lys Pro Phe Gly Leu Glu Asn Trp Cys Lys Tyr Thr Val Val Glu
165        170        175

Lys Gly Ile Leu Tyr Ser Cys Lys Phe Gly Gly Asn Ser Thr Cys Ile
180        185        190

Lys Gly Leu Ile Val Lys Gly Gln Arg Glu Asp Lys Val Arg Tyr Cys
195        200        205

Glu Trp Cys Gly Tyr Lys Phe Ser Ser Pro Asn Gly Leu Pro Gln Tyr
210        215        220

Pro Leu Gly Leu Cys Glu Lys Glu Gln Ser Glu Gly Leu Arg Asp Tyr
225        230        235        240

Gly Asp Phe Pro Cys Cys Asn Asn Gly Thr Cys Ile Asp Lys Glu Gly
245        250        255

Ser Val Gln Cys Tyr Ile Gly Asp Lys Lys Val Thr Val Lys Leu Tyr
260        265        270

Asn Ala Ser Leu Leu Ala Pro Met Pro Cys Lys Pro Ile Val Tyr Asn
275        280        285

Ser Gln Gly Pro Pro Ala Pro Lys Thr Cys Thr Tyr Arg Trp Ala Ser
290        295        300

Thr Leu Glu Asn Lys Tyr Tyr Glu Pro Arg Asp Ser Tyr Tyr Gln Gln
305        310        315        320

Tyr Ile Ile Lys Ser Gly Tyr Gln Tyr Trp Phe Asp Leu Thr Ala Lys
325        330        335

Asp His Val Ala Asp Trp Ile Thr Lys Tyr Phe Pro Ile Ile Ile Val
340        345        350

Ala Leu Leu Gly Gly Arg Gly Thr Leu Trp Val Leu Ile Ala Tyr Glu

```


-continued

Thr	Asp	Ser	Gly	Cys	Ser	Asp	Gly	Ala	Arg	Cys	Tyr	Val	Ile	Asn	Pro
			100					105					110		
Glu	Ala	Thr	Asn	Ile	Ala	Gly	Thr	Lys	Gly	Ala	Met	Val	His	Leu	Arg
		115					120					125			
Lys	Ala	Gly	Gly	Glu	Phe	Asn	Cys	Val	Thr	Ala	Gln	Gly	Thr	Pro	Ala
	130					135					140				
Phe	Tyr	Asn	Leu	Lys	Asn	Leu	Lys	Gly	Trp	Ser	Gly	Leu	Pro	Ile	Phe
145					150				155						160
Glu	Ala	Ala	Thr	Gly	Arg	Val	Val	Gly	Arg	Val	Lys	Ala	Gly	Lys	Asn
				165					170					175	
Thr	Asp	Asn	Ala	Pro	Thr	Thr	Ile	Met	Ser	Gly	Thr	Gln	Val	Ala	Lys
			180					185					190		
Pro	Ser	Glu	Cys	Asp	Leu	Glu	Ser	Val	Val	Arg	Lys	Leu	Glu	Thr	Met
		195					200					205			
Asn	Arg	Gly	Glu	Phe	Lys	Gln	Val	Thr	Leu	Ala	Thr	Gly	Ala	Gly	Lys
	210					215					220				
Thr	Thr	Met	Leu	Pro	Lys	Leu	Leu	Ile	Glu	Ser	Ile	Gly	Arg	His	Lys
225					230					235					240
Arg	Val	Leu	Val	Leu	Ile	Pro	Leu	Arg	Ala	Ala	Ala	Glu	Gly	Val	Tyr
			245						250					255	
Gln	Tyr	Met	Arg	Thr	Lys	His	Pro	Ser	Ile	Ser	Phe	Asn	Leu	Arg	Ile
			260					265					270		
Gly	Asp	Leu	Lys	Glu	Gly	Asp	Met	Ala	Thr	Gly	Ile	Thr	Tyr	Ala	Ser
	275						280					285			
Tyr	Gly	Tyr	Phe	Cys	Gln	Met	Asp	Met	Pro	Arg	Leu	Glu	Asn	Ala	Met
	290					295					300				
Lys	Glu	Tyr	His	Tyr	Ile	Phe	Leu	Asp	Glu	Tyr	His	Cys	Ala	Thr	Pro
305				310						315					320
Glu	Gln	Leu	Ala	Val	Met	Ser	Lys	Ile	His	Arg	Phe	Gly	Glu	Ser	Val
				325					330					335	
Arg	Val	Ile	Ala	Met	Thr	Ala	Thr	Pro	Ser	Gly	Thr	Val	Ser	Thr	Thr
			340					345						350	
Gly	Gln	Lys	Phe	Thr	Ile	Glu	Glu	Val	Val	Val	Pro	Glu	Val	Met	Lys
		355					360					365			
Gly	Glu	Asp	Leu	Ala	Asp	Asp	Tyr	Ile	Glu	Ile	Ala	Gly	Leu	Lys	Val
	370					375					380				
Pro	Lys	Lys	Glu	Leu	Glu	Gly	Asn	Val	Leu	Thr	Phe	Val	Pro	Thr	Arg
385					390					395					400
Lys	Met	Ala	Ser	Glu	Thr	Ala	Lys	Lys	Leu	Thr	Thr	Gln	Gly	Tyr	Asn
			405						410					415	
Ala	Gly	Tyr	Tyr	Phe	Ser	Gly	Glu	Asp	Pro	Ser	Ser	Leu	Arg	Thr	Thr
			420					425					430		
Thr	Ser	Lys	Ser	Pro	Tyr	Ile	Val	Val	Ala	Thr	Asn	Ala	Ile	Glu	Ser
		435					440					445			
Gly	Val	Thr	Leu	Pro	Asp	Leu	Asp	Thr	Val	Ile	Asp	Thr	Gly	Met	Lys
	450					455					460				
Cys	Glu	Lys	Arg	Leu	Arg	Ile	Glu	Asn	Lys	Ala	Pro	Tyr	Ile	Val	Thr
465					470					475					480
Gly	Leu	Lys	Arg	Met	Ala	Ile	Thr	Thr	Gly	Glu	Gln	Ala	Gln	Arg	Lys
			485						490					495	
Gly	Arg	Val	Gly	Arg	Val	Lys	Pro	Gly	Arg	Tyr	Leu	Arg	Gly	Pro	Glu

-continued

65		70		75		80									
Gly	His	Glu	Thr	Ala	Phe	Ala	Ser	Leu	Val	Ile	Lys	Trp	Ile	Ala	Phe
				85					90					95	
Ser	Ser	Asp	Gly	Val	Pro	Gly	Met	Ile	Lys	Gln	Ala	Ala	Val	Asp	Leu
			100					105						110	
Val	Val	Tyr	Tyr	Ile	Ile	Asn	Arg	Pro	Glu	Tyr	Gln	Gly	Asp	Lys	Glu
			115					120					125		
Thr	Gln	Asn	Ala	Gly	Arg	Gln	Phe	Val	Gly	Ser	Leu	Phe	Val	Ser	Cys
							135							140	
Leu	Ala	Glu	Tyr	Thr	Phe	Lys	Asn	Phe	Asn	Lys	Ser	Ala	Leu	Glu	Gly
							150							155	160
Leu	Ile	Glu	Pro	Ala	Leu	Ser	Tyr	Leu	Pro	Tyr	Ala	Ser	Ser	Ala	Leu
							165								175
Lys	Leu	Phe	Leu	Pro	Thr	Arg	Leu	Glu	Ser	Val	Val	Ile	Leu	Ser	Thr
							180								190
Thr	Ile	Tyr	Arg	Thr	Tyr	Leu	Ser	Ile	Arg	Lys	Gly	Ser	Ser	Gln	Gly
							200							205	
Leu	Ala	Gly	Leu	Ala	Val	Ser	Ser	Ala	Met	Glu	Ile	Met	Asn	Gln	Asn
							210							215	220
Pro	Ile	Ser	Val	Ala	Ile	Ala	Leu	Ala	Leu	Gly	Val	Gly	Ala	Ile	Ala
							225							230	240
Ala	His	Asn	Ala	Ile	Glu	Ser	Ser	Glu	Ala	Lys	Arg	Thr	Leu	Leu	Met
							245							250	255
Lys	Val	Phe	Val	Lys	Asn	Phe	Leu	Asp	Gln	Ala	Ala	Thr	Asp	Glu	Leu
							260							265	270
Val	Lys	Glu	Asn	Pro	Glu	Lys	Ile	Ile	Met	Ala	Val	Phe	Glu	Gly	Ile
							275							280	285
Gln	Thr	Ala	Gly	Asn	Pro	Leu	Arg	Leu	Val	Tyr	His	Leu	Tyr	Ala	Met
							290							295	300
Phe	Tyr	Lys	Gly	Trp	Thr	Ala	Ala	Glu	Ile	Ala	Glu	Lys	Thr	Ala	Gly
							305							310	315
Arg	Asn	Ile	Phe	Val	Leu	Thr	Ile	Phe	Glu	Gly	Leu	Glu	Met	Leu	Gly
							320							325	330
Leu	Asp	Ala	Asp	Ser	Lys	Trp	Arg	Asn	Leu	Ser					
							340							345	

<210> SEQ ID NO 26

<211> LENGTH: 505

<212> TYPE: PRT

<213> ORGANISM: PMC Virus

<400> SEQUENCE: 26

Ser	Asn	Tyr	Leu	Ile	Asp	Ala	Val	Lys	Lys	Ile	Ile	Glu	Lys	Met	Thr
1			5						10					15	
Lys	Thr	Ala	Thr	Ser	Phe	Thr	Tyr	Ser	Phe	Leu	Lys	Ser	Leu	Leu	Pro
			20						25					30	
Ala	Pro	Phe	Ser	Cys	Thr	Lys	Ser	Glu	Arg	Asp	Pro	Arg	Ile	Gly	Trp
			35						40					45	
Pro	Gln	Lys	Asp	Tyr	Asp	Tyr	Leu	Glu	Val	Arg	Cys	Ala	Cys	Gly	Tyr
			50						55					60	
Asn	Arg	Arg	Ala	Ile	Lys	Arg	Asp	Ser	Gly	Pro	Val	Leu	Trp	Glu	Thr
65					70					75					80

-continued

Leu Glu Glu Thr Gly Pro Glu Tyr Cys His Asn Arg Gly Glu Arg Gly
 85 90 95
 Leu Ser Asn Val Lys Thr Thr Arg Cys Phe Val Gln Gly Glu Glu Ile
 100 105 110
 Pro Pro Ile Ala Leu Arg Lys Gly Val Gly Glu Met Leu Val Lys Gly
 115 120 125
 Val Ser Phe Arg Ile Asp Phe Asp Lys Asp Lys Ile Leu Ser Thr Asp
 130 135 140
 Lys Trp Lys Val Pro His Arg Ala Val Thr Ser Ile Phe Glu Asp Trp
 145 150 155 160
 Gln Gly Ile Gly Tyr Arg Glu Ala Tyr Leu Gly Thr Lys Pro Asp Tyr
 165 170 175
 Gly Gly Leu Val Pro Arg Ser Cys Val Thr Val Thr Lys Gln Gly Leu
 180 185 190
 Thr Phe Leu Lys Thr Ala Arg Gly Met Ala Phe Thr Thr Asp Leu Thr
 195 200 205
 Ile Gln Asn Ile Lys Met Leu Ile Ala Thr Cys Phe Lys Asn Lys Val
 210 215 220
 Lys Glu Gly Glu Ile Pro Ala Thr Ile Glu Gly Glu Thr Trp Ile Asn
 225 230 235 240
 Ile Pro Leu Val Asn Glu Asp Thr Gly Thr Ile Lys Pro Ser Phe Gly
 245 250 255
 Glu Arg Val Ile Pro Glu Pro Tyr Glu Glu Asp Pro Leu Glu Gly Pro
 260 265 270
 Ser Val Ile Val Glu Thr Gly Gly Ile Ala Ile Asn Gln Ile Gly Val
 275 280 285
 Asn Pro Gln Ser Ser Thr Cys Gly Thr Val Phe Thr Ala Val Lys Asp
 290 295 300
 Leu Cys Gln Thr Val Ser Asn Lys Ala Lys Asn Ile Lys Ile Gly Phe
 305 310 315 320
 Ser Glu Gly Gln Tyr Pro Gly Pro Gly Val Ala Lys Lys Thr Leu Asn
 325 330 335
 Gln Leu Ile Gln Asp Glu Asp Pro Lys Pro Phe Ile Phe Val Cys Gly
 340 345 350
 Ser Asp Lys Ser Met Ser Asn Arg Ala Lys Thr Ala Arg Asn Ile Lys
 355 360 365
 Arg Ile Thr Thr Thr Thr Pro Glu Lys Phe Arg Asp Leu Ala Lys Asn
 370 375 380
 Lys Lys Leu Ile Ile Val Leu Leu Gly Asp Arg Tyr His Glu Asp Ile
 385 390 395 400
 Glu Lys Tyr Ala Asp Phe Lys Gly Thr Phe Leu Thr Arg Gln Thr Leu
 405 410 415
 Glu Ala Leu Ala Ser Ala Lys Ala Val Glu Lys Asp Met Thr Lys Lys
 420 425 430
 Glu Ala Ala Arg Val Leu Ala Met Glu Glu Lys Asp Glu Leu Glu Leu
 435 440 445
 Pro Gly Trp Leu His Thr Asp Ala Pro Lys Phe Leu Asp Ile Thr Lys
 450 455 460
 Asp Asn Ile Thr His His Leu Ile Gly Asp Met Gln Ser Leu Arg Glu
 465 470 475 480
 Arg Ala Gly Glu Ile Gly Ala Lys Ala Thr Thr Gln Ile Thr Lys Lys

-continued

Val Ala Val Ser Phe Asp Thr Lys Ala Trp Asp Thr Gln Val Thr Pro
 340 345 350
 Lys Asp Leu Leu Leu Ile Ser Glu Ile Gln Lys Tyr Tyr Tyr Lys Lys
 355 360 365
 Glu Tyr His Arg Phe Ile Asp Asn Leu Thr Glu Lys Met Val Glu Val
 370 375 380
 Pro Val Val Cys Glu Asp Gly Asn Val Tyr Ile Arg Glu Gly Gln Arg
 385 390 395 400
 Gly Ser Gly Gln Pro Asp Thr Ser Ala Gly Asn Ser Met Leu Asn Val
 405 410 415
 Leu Thr Met Ile Tyr Ala Phe Cys Lys Ala Asn Ser Ile Pro Tyr Ser
 420 425 430
 Ala Phe His Arg Val Ala Lys Ile His Val Cys Gly Asp Asp Gly Phe
 435 440 445
 Leu Ile Thr Glu Lys Ser Phe Gly Glu Ala Phe Ala Ile Lys Gly Pro
 450 455 460
 Gln Ile Leu Met Glu Ala Gly Lys Pro Gln Lys Leu Ile Gly Glu Phe
 465 470 475 480
 Gly Leu Lys Leu Ala Tyr Lys Phe Asp Asp Ile Glu Phe Cys Ser His
 485 490 495
 Thr Pro Ile Lys Val Arg Trp Ala Asp Asn Asn Thr Ser Tyr Met Pro
 500 505 510
 Gly Arg Asp Thr Ala Thr Ile Leu Ala Lys Met Ala Thr Arg Leu Asp
 515 520 525
 Ser Ser Gly Glu Arg Gly Thr Glu Gly Tyr Glu Leu Ala Val Ala Phe
 530 535 540
 Ser Phe Leu Leu Met Tyr Ser Trp Asn Pro Leu Val Arg Arg Ile Cys
 545 550 555 560
 Leu Leu Val Met Ser Thr Ile Asp Thr Lys Glu Ala Ser Gln Asn Asn
 565 570 575
 Thr Ile Tyr Thr Phe Arg Gly Asp Pro Ile Gly Ala Tyr Thr Glu Val
 580 585 590
 Ile Gly Tyr Arg Leu Asp Gln Leu Lys Gln Thr Glu Phe Ser Lys Leu
 595 600 605
 Ala Gln Leu Asn Leu Ser Met Ala Ile Leu Gln Ile Tyr Asn Lys Asn
 610 615 620
 Thr Thr Lys Arg Leu Ile Glu Asp Cys Val Lys Leu Gly Asn Gln Asn
 625 630 635 640
 Lys Gln Ile Leu Val Asn Ala Asp Arg Leu Ile Ser Lys Lys Thr Gly
 645 650 655
 Tyr Thr Tyr Glu Pro Thr Ala Gly His Thr Lys Ile Gly Lys His Tyr
 660 665 670
 Glu Glu Ile Asn Leu Leu Lys Asp Thr Pro Gln Lys Thr Val Tyr Gln
 675 680 685
 Gly Thr Glu Arg Tyr
 690

1. An isolated RNA nucleotide sequence comprising:
 - a) SEQ ID NO:1, wherein thymidine (t) nucleotides are substituted with uridine (u) nucleotides;
 - b) an RNA sequence substantially homologous to the nucleotide sequence of (a);
 - c) an RNA sequence comprising the complement of the nucleotide sequence of (a);
 - d) fragments of any one of (a), (b) or (c);
 - e) any one of SEQ ID NOs:3-15; or
 - f) variants of any one of (a)-(e).
2. The sequence fragment of claim 1 selected from the group consisting of the following locations of SEQ ID NO: 1: position 419-922, 923-1219, 1220-1885, 1886-2473, 2474-3604, 3605-3820, 3821-5224, 5225-7252, 7253-7441, 7442-8482, 8483-9997 and 9998-12077.
3. An isolated DNA sequence comprising:
 - a) SEQ ID NO: 1;
 - b) a DNA sequence substantially homologous to SEQ ID NO:1;
 - c) a DNA sequence comprising the complement of SEQ ID NO:1;
 - d) fragments of any one of (a), (b) or (c);
 - e) any one of SEQ ID NOs:3-15; or
 - f) variants of any one of (a)-(e).
4. The sequence of claim 3 selected from the group consisting of the following locations of SEQ ID NO:1: position 419-922, 923-1219, 1220-1885, 1886-2473, 2474-3604, 3605-3820, 3821-5224, 5225-7252, 7253-7441, 7442-8482, 8483-9997 and 9998-12077.
5. An isolated amino acid sequence comprising:
 - a) SEQ ID NO:2;
 - b) polypeptide encoded by the nucleotide of claim 1;
 - c) any one of SEQ ID NOs:16-27;
 - d) fragments of (a) or (b);
 - e) variants of any one of (a)-(d).
6. The sequence of claim 5 selected from the group consisting of the following locations in SEQ ID NO:2: 1-167, 168-267, 268-489, 490-685, 686-1062, 1063-1134, 1135-1602, 1603-2278, 2279-2341, 2342-2688, 2689-3193 and 3194-3886.
7. A method for detecting the presence of a PMC virus amino acid sequence according to claim 5 in a sample, comprising the steps of:
 - a) contacting a sample suspected of containing a PMC virus amino acid sequence with an antibody that specifically binds to the PMC virus amino acid sequence under conditions which allow for the formation of reaction complexes comprising the antibody and the PMC virus amino acid sequence; and
 - b) detecting the formation of reaction complexes comprising the antibody and PMC virus amino acid sequence in the sample, wherein detection of the formation of reaction complexes indicates the presence of PMC virus amino acid sequence in the sample.
8. A method for detecting the presence of an anti-PMC virus antibody in a sample, comprising the steps of:
 - a) contacting a sample suspected of containing an anti-PMC virus antibody with an amino acid sequence according to claim 5 under conditions which allow for the formation of reaction complexes comprising the PMC virus antibody and the amino acid sequence; and
 - b) detecting the formation of reaction complexes comprising the antibody and amino acid sequence in the sample, wherein detection of the formation of reaction complexes indicates the presence of PMC virus antibody in the sample.
9. A method for the detection and evaluating the levels of anti-PMC virus antibodies in a subject, comprising the steps of:
 - a) depositing a predetermined amount of one or several PMC virus antigens comprising an amino acid sequence according to claim 5 onto a solid support such as a microplate;
 - b) introducing increasing dilutions of a biological fluid (e.g., blood serum or plasma, milk, cerebrospinal fluid, lymphatic fluid or other body fluids) onto the antigens and incubating;
 - c) washing the solid support with an appropriate buffer;
 - d) adding specific labelled antibodies directed against the antibodies of the subject;
 - e) detecting the antigen-antibody-antibody complex formed, which is then indicative of the presence of anti-PMC virus antibodies in the biological fluid; and
 - f) evaluating the amount of reaction complexes formed, which amount of reaction complexes corresponds to the level of PMC virus antibodies in the biological sample.
10. The method of claim 9 wherein the levels of PMC virus antibodies are determined in a series of biological samples obtained at different time points from an animal host undergoing therapeutic treatment.
11. A method for detecting the presence or absence of PMC virus in a biological sample, comprising the steps of:
 - a) bringing the biological sample into contact with a polynucleotide probe or primer comprising:
 - i. an isolated RNA nucleotide sequence according to claim 1, or
 - ii. an isolated DNA sequence comprising:
 - 1) SEQ ID NO: 1;
 - 2) a DNA sequence substantially homologous to SEQ ID NO:1;
 - 3) a DNA sequence comprising the complement of SEQ ID NO:1;
 - 4) fragments of any one of (1), (2) or (3);
 - 5) any one of SEQ ID NOs:3-15; or
 - 6) variants of any one of (1)-(5),
 under suitable hybridizing conditions; and
 - b) detecting any duplex formed between the probe or primer and nucleic acid sequences in the sample.
12. A method for the detection of PMC virus nucleic acids present in a biological sample, comprising:
 - a) amplifying the DNA sequence of claim 3 with at least one primer,
 - b) detecting the amplified nucleic acids.
13. A method for the detection of PMC virus nucleic acids present in a biological sample, comprising:
 - a) hybridizing the nucleic acids of the biological sample at appropriate conditions with one or more probes comprising:
 - i. an isolated RNA nucleotide sequence according to claim 1, or
 - ii. an isolated DNA sequence comprising:
 - 1) SEQ ID NO: 1;
 - 2) a DNA sequence substantially homologous to SEQ ID NO:1;
 - 3) a DNA sequence comprising the complement of SEQ ID NO:1;
 - 4) fragments of any one of (1), (2) or (3);

- 5) any one of SEQ ID NOs:3-15; or
6) variants of any one of (1)-(5),
b) washing under appropriate conditions, and
c) detecting the hybrids formed.
- 14.** A method for detecting viral RNA or DNA comprising the steps of:
a) immobilizing PMC virus on a support;
b) disrupting the virion; and
c) hybridizing the nucleic acids of the virion with a probe comprising:
i. an isolated RNA nucleotide sequence according to claim 1, or
ii. an isolated DNA sequence comprising:
1) SEQ ID NO: 1;
2) a DNA sequence substantially homologous to SEQ ID NO:1;
3) a DNA sequence comprising the complement of SEQ ID NO:1;
4) fragments of any one of (1), (2) or (3);
5) any one of SEQ ID NOs:3-15; or
6) variants of any one of (1)-(5).
- 15.** A method for screening the tissue of subjects for PMC virus comprising the steps of:
a) extracting DNA from tissue;
b) restriction enzyme cleavage of said DNA;
c) electrophoresis of the fragments; and
d) Southern blotting of genomic DNA from tissues and subsequent hybridization with a labelled cloned PMC virus DNA sequence according to claim 3.
- 16.** A method for the generation of antibodies comprising the steps of:
a) providing a PMC virus polypeptide sequence comprising an amino acid sequence of claim 5 to a subject; and
b) collecting the antibodies generated in the subject against the polypeptide.
- 17.** The method of claim 16 wherein the polypeptide is further coupled or covalently conjugated to a physiologically acceptable carrier molecule.
- 18.** A vaccine composition comprising a PMC virus polypeptide that comprises an amino acid sequence according to claim 5.
- 19.** A vaccine composition comprising:
i. an isolated RNA nucleotide sequence according to claim 1, or
ii. an isolated DNA sequence comprising:
1) SEQ ID NO: 1;
2) a DNA sequence substantially homologous to SEQ ID NO:1;
3) a DNA sequence comprising the complement of SEQ ID NO:1;
4) fragments of any one of (1), (2) or (3);
5) any one of SEQ ID NOs:3-15; or
6) variants of any one of (1)-(5).
- 20.** The composition of claim 18 wherein the composition further comprises a pharmaceutically acceptable carrier or diluent and/or an adjuvant.
- 21.** A method to vaccinate an animal against PMC comprising the steps of: providing a vaccine according to claim 18 to the animal in an amount effective for producing protection against a pestivirus infection.
- 22.** The method of claim 21 wherein the immunogenic composition is administered by a method selected from the list comprising: intravenously, subcutaneously, intramuscularly, intraorbitally, ophthalmically, intraventricularly, intrac-

ranially, intracapsularly, intraspinaly, intracisternally, intraperitoneally, buccal, rectally, vaginally, intranasally, orally or aerosol administration.

23. A pharmaceutical composition comprising a PMC virus amino acid sequence according to claim 5 in a therapeutically effective amount admixed with a pharmaceutically acceptable carrier, diluent, or excipient.

24. A pharmaceutical composition comprising antibodies prepared against PMC polypeptides comprising an amino acid sequence of claim 5.

25. A therapeutic composition comprising a nucleic acid sequence selected from the group consisting of:

- i) an isolated RNA nucleotide sequence comprising:
1) SEQ ID NO:1, wherein thymidine (t) nucleotides are substituted with uridine (u) nucleotides;
2) an RNA sequence substantially homologous to the nucleotide sequence of (1);
3) an RNA sequence comprising the complement of the nucleotide sequence of (1);
4) fragments of any one of (1), (2) or (3);
5) any one of SEQ ID NOs:3-15; or
6) variants of any one of (1)-(5);
ii. an isolated DNA sequence comprising:
1) SEQ ID NO: 1;
2) a DNA sequence substantially homologous to SEQ ID NO:1;
3) a DNA sequence comprising the complement of SEQ ID NO:1;
4) fragments of any one of (1), (2) or (3);
5) any one of SEQ ID NOs:3-15; or
6) variants of any one of (1)-(5); and
iii. an antisense or ribozyme polynucleotide sequence hybridizable to a polynucleotide sequence encoding a PMC virus amino acid sequence according to claim 5.

26. A method of ameliorating a disease associated with PMC virus comprising administering to a subject a PMC virus amino acid sequence according to claim 5, wherein said PMC virus amino acid sequence ameliorates said disease associated with PMC virus.

27. A method of ameliorating a disease associated with PMC virus comprising administering to a subject antibodies against the PMC virus, wherein said antibodies against the PMC virus ameliorate said disease associated with PMC virus.

28. A method of ameliorating a disease associated with PMC virus comprising administering to a subject a PMC virus nucleic acid sequence, or an antisense or ribozyme polynucleotide sequence hybridizable to a polynucleotide sequence encoding a PMC virus amino acid sequence according to claim 5, wherein said a PMC virus nucleic acid sequence, or an antisense or ribozyme polynucleotide sequence hybridizable to a polynucleotide sequence encoding a PMC virus amino acid sequence ameliorates said disease associated with PMC virus.

29. A method of inducing a protective immune response in an animal or human against a PMC virus comprising the step of:

- a) administering to said animal or human an effective amount of a composition comprising anti-PMC virus antibodies that are specific to an amino acid sequence according to claim 5.

30. A method for enhancing an animal's immunocompetence and the activity of its immune effector cells against a PMC virus comprising the step of:

- a) administering a composition comprising a therapeutically effective amount of a PMC virus peptide or polypeptide that comprise an amino acid according to claim 5.
- 31.** A live vector comprising:
- i. an isolated RNA nucleotide sequence according to claim 1, or
 - ii. an isolated DNA sequence comprising:
 - 1) SEQ ID NO: 1;
 - 2) a DNA sequence substantially homologous to SEQ ID NO:1;
 - 3) a DNA sequence comprising the complement of SEQ ID NO:1;
 - 4) fragments of any one of (1), (2) or (3);
 - 5) any one of SEQ ID NOs:3-15; or
 - 6) variants of any one of (1)-(5)
- and a heterologous polynucleotide.
- 32.** The live vector of claim 31 wherein the heterologous polynucleotide is operably linked to the polynucleotide sequence of the PMC virus, such that expression of the polynucleotide sequence of the PMC virus also leads to expression of the heterologous polynucleotide sequence.
- 33.** The vector of claim 31 wherein one or more sections of autologous polynucleotide sequence are removed such that the live virus is rendered attenuated in pathogenicity in a host subject.
- 34.** A method of screening for drugs comprising the steps of:
- a) contacting an agent with a PMC virus amino acid sequence according to claim 5 and
 - b) assaying for the presence of a complex between the agent and the PMC virus amino acid sequence.
- 35.** A method of screening for ligands of the proteins of the PMC virus comprising the steps of:
- a) contacting a ligand with a PMC virus amino acid sequence according to claim 5; and
 - b) assaying for the presence of a complex between the PMC virus amino acid sequence or fragment and a ligand.
- 36.** A kit for the demonstration of the presence of PMC virus comprising:
- (a) a predetermined amount of at least one labelled immunochemically reactive component obtained by the direct or indirect attachment of a PMC virus amino acid sequence according to claim 5 or a specific binding partner thereto, to a detectable label;
 - (b) other reagents; and
 - (c) directions for use of said kit.
- 37.** A kit for demonstrating the presence of PMC virus comprising:
- (a) a predetermined amount of at least one labelled antibody specific for an amino acid sequence of claim 5;
 - (b) other reagents; and
 - (c) directions for use of said kit.
- 38.** A diagnostic kit for demonstrating the presence of PMC virus comprising:
- (a) a predetermined amount of at least one labelled polypeptide comprising an amino acid sequence of claim 5 derived from the PMC virus;
 - (b) other reagents; and
 - (c) directions for use of said kit.
- 39.** A kit for demonstrating the presence of PMC virus comprising:
- (a) a predetermined amount of at least one labelled nucleic acid sequence derived from the PMC virus, wherein said nucleic acid sequence comprises:
 - i. an isolated RNA nucleotide sequence according to claim 1, or
 - ii. an isolated DNA sequence comprising:
 - 1) SEQ ID NO: 1;
 - 2) a DNA sequence substantially homologous to SEQ ID NO:1;
 - 3) a DNA sequence comprising the complement of SEQ ID NO:1;
 - 4) fragments of any one of (1), (2) or (3);
 - 5) any one of SEQ ID NOs:3-15; or
 - 6) variants of any one of (1)-(5);
 - (b) other reagents; and
 - (c) directions for use of said kit.
- 40.** A recombinant expression vector comprising a PMC virus nucleic acid sequence or a part thereof of claim 1, operably linked to prokaryotic, eukaryotic or viral transcription and translation control elements.
- 41.** A host cell transformed by the vector of claim 40.
- 42.** A method for preparing a PMC virus amino acid sequence according to claim 5, comprising the steps of:
- (a) culturing a host cell containing a vector according to claim 40 under conditions that provide for expression of the PMC virus amino acid sequence; and
 - (b) recovering the expressed PMC virus sequence.
- 43.** The composition of claim 19 wherein the composition further comprises a pharmaceutically acceptable carrier or diluent and/or an adjuvant.
- 44.** A method to vaccinate an animal against PMC comprising the steps of: providing a vaccine according to claim 19 to the animal in an amount effective for producing protection against a pestivirus infection.
- 45.** The method of claim 44 wherein the immunogenic composition is administered by a method selected from the list comprising: intravenously, subcutaneously, intramuscularly, intraorbitally, ophthalmically, intraventricularly, intracranially, intracapsularly, intraspinaly, intracisternally, intraperitoneally, buccal, rectally, vaginally, intranasally, orally or aerosol administration.
- 46.** A recombinant expression vector comprising a PMC virus nucleic acid sequence or a part thereof of claim 3, operably linked to prokaryotic, eukaryotic or viral transcription and translation control elements.
- 47.** A host cell transformed by the vector of claim 46.
- 48.** A method for preparing a PMC virus amino acid sequence, comprising the steps of:
- (a) culturing a host cell containing a vector according to claim 46 under conditions that provide for expression of the PMC virus amino acid sequence; and
 - (b) recovering the expressed PMC virus sequence

* * * * *

专利名称(译)	瘟病毒种		
公开(公告)号	US20100203063A1	公开(公告)日	2010-08-12
申请号	US12/298026	申请日	2007-04-20
[标]申请(专利权)人(译)	MINIST初级IND FOR & ON新南威尔士州代表国家		
申请(专利权)人(译)	代表新南威尔士州的初级工业部长		
当前申请(专利权)人(译)	代表新南威尔士州的初级工业部长		
[标]发明人	FROST MELINDA JANE KIRKLAND PETER DANIEL FINLAISON DEBORAH SUSAN		
发明人	FROST, MELINDA JANE KIRKLAND, PETER DANIEL FINLAISON, DEBORAH SUSAN		
IPC分类号	A61K39/42 C07H21/02 C07H21/04 C07K14/005 C12Q1/70 G01N33/53 C12Q1/68 C12P21/04 A61K39/12 A61K31/7088 A61K38/16 C12N15/63 C12N5/10 C12P21/06 A61P31/12		
CPC分类号	A61K39/12 C07K14/005 C12N7/00 C12N2770/24321 G01N2469/20 C12Q1/701 G01N33/56983 G01N2333/18 C12N2770/24322 A61K39/00 A61P31/12 A61P31/14 A61K2039/53 C12N2770/24334 C12Q2600/136		
优先权	2006902089 2006-04-21 AU		
其他公开文献	US8202977		
外部链接	Espacenet USPTO		

摘要(译)

本申请涉及与猪心肌炎综合征相关的瘟病毒，称为PMC病毒，以及由其衍生的基因和蛋白质序列。本申请还涉及使用PMC病毒或其衍生的基因/蛋白质序列的检测方法，疫苗治疗剂和诊断方法。

Figure 1

```

gatacaacgac agtagtctcga gctgctgctat gcaatcattgg ccaatacaaaa tcatctcaatt 60
tggaaaatggg acctgcaaac tglcaagaggg scggagcctgg gtaacacatcc cgaactcgtag 120
gactcagatca atcaggtcaaa ctgggtgagc aggttagctg gctcagaggg cttaagcggctg 180
gtacacacctt attctctcaac aggtctactt ggaagagctt acccaactgac ggaactcctg 240
ctcagagatgg gtagttgcaac acaaacctctt ataggtatctt gctcagatccc cgtcattcttg 300
accggtataa atattatccg tctgtaaaaga ta tpaatactt tttaacttta acaaatatctg 360
aggggtggag gaagaaaaa caatgagaca caatgagacc aaatgacccc agcagccccc cagggtatcca 420
agaacccggtc tcccaagca caatgagacc aaatctctgg gaaccccctc caactcaca 480
caaacagatc agtcaaaat tgcacacttt gagggtgctt gaaccccctc caactcaca 540
ggagatcttg ccaagagagj gtagt tptat caactcaaca aagctcoca ctctgggggt 600
tagctgtgaa tcaagggctg tgcctatata agatcaaacj ggcacgggat accctcctgt 720
atcgttagaa ctctgcaaa tgcagagatc tttagatagt ccggagatc gtaactcctt 780
gcaagggctc gatggaggt tglcaaaagt cctgaagtag atcccaaac ccttgaacac 840
tcaagatcac tctagaggtc taacaagcgt cctgaagtag atcccaaac ccttgaacac 900
accacttgg gtcagaggtt gctctgagca caaagggcca aaacccaagg tgaacaacaaa 960
agaagccagg atpagaagj gaaaaatgag gctcagagctt caaagagctg aagagatcca 1020
aaaacctaga ccccaagctg ccaagatagc ggttgaaggg caaagatctc aggtgagaaa 1080
gaagggcaaa gcaaaaacca agactcaaga agctctcaac ctctggggac caaaaacctc 1140
aggtccagg agagagctc agaaagctct gtagatagg gcaaatctag cctgcaatct 1200
ggtatgacag gctgagcaca ccaagctgca aggtctaaa agggatcgg atggaaattg 1260
gccaacaaaa atctgcaagj gbatccctca acactgaaa gctcagatcg aactgaaa 1320
gatccaggj atgtggatg cagagccact gaacacttc acatgctga ggtcacaagj 1380
caatgaatac acccaagggc tatcaaac caatgagacaa tggagacaaa ctactgggtt gctcagatc 1440
atggagatc gatcagact taggttcaaa taactgact caagccaggj caactcaac 1500
tatctgact ggtctgaagj aggggcaaa atctctctt tcaagggatg tcaagggctc 1560
accctgcaac tctgattaa ctgctgaaga ctctgctagg atcattggct caacacactg 1620
ggagatctt actctctagj gggagagatc gttgagctcg agtaagatcg aaaaacccaa 1680
ggtagatca agtggctca gggctctcac atggctgctg agtaagatcg aaaaacccaa 1740
cgaagatctt cctgagctg aggcagatc ttactgcaaa gctcagagg tagagatcaa 1800
catattatc accaacatt gcaaccctct tggactgcca gtaagatcaa aattctagj 1860
accaggaaca tctgactca gttggagagg tagattctat tctccaacac tccctcaatca 2040
cgttagtgc tccattctca tggatctaat tgaactgctc gaattctgct cagagacatc 2100
agcaatact cactcctgct tgcctacact aaacccagct aatcaacaa gggatctctg 2160
gttggactcg gcccaaatc aactaaacct taactcaact aaaccctggj caagctctgct 2220
cctcaacagj gcaaatgctc taagttgggtt gttcaaacct gttgctgctt aaacacagct 2280
tccagcaaaa accactaac tptatggggj tgcatacaac gttgagact tegtatcaa 2340
gcaacatgaa gaactgctc atctatgac cgaagcaaca gctctggatc tctgctgctg 2400
tcaataaaa atctctagj gcccagctt ccaagggatc gaatggatcaa tcaactatgg 2460
ggagatcaca gccaacagj gcaacccaga atctctgca gactctgaga caaagctg 2520
caaatcact gcaagcaga tggagctcc ctctgctgctt gcaacatgga caagctgct 2580
caatgctca tcaagccagj aatcactgaa gaaagggjg taacacaaa caagctgct 2640
atgtctcca ctctctagj aagatctc taataaaa gttcaaaa gttcaaaa gttcaaaa 2700
aacggctggj tcaagaaaa tcaaacatca gcttaggata caaagagaaa gaactctagj 2760
tjctctaaa agggatgca ctctgactg tggcttcaa cctctggjg ggtcaaac 2820
caatctaac caaatctca tctcaacccc aggtctgaa ctctgactgc ctaggggtg 2880
gactggggct gtaggtgata accctcaact cctcaactca actgagcaac agatctatg 2940
caatctaac agcccaaac caatggact tgaaacctg tggagatcaa caaggttga 3000
gaagggatct cgtactctct gctactctgj gggatctca aactgcaaca caaggtat 3060
agctaaaggj caaggggagj caaagcaagj gctactgtga tgggtggctt aaggttga 3120
tcaacaaact ggtactctct ggtactctgj ggaagagaa ggtactgaa aaggttga 3180
actcagggat tctgtgact tccctctgct caaacagggj actctgactg caaagatgg 3240
tgggtcaca tgcactagj gggcaagaa ggtcaccctg aagctatca aaggttga 3300
actggccccc atggcctga acccaactg gataactcc aggggccc cagggctaa 3360
gcaactcact tctaggtggj cctcaactt ggaataaaa tatctgagc caaggtgag 3420
cactacccg caatactta taagctagg gataactat tggcttgact tcaagcaaa 3480

```