



US 20120329052A1

(19) **United States**(12) **Patent Application Publication**
Peltonen et al.(10) **Pub. No.: US 2012/0329052 A1**(43) **Pub. Date: Dec. 27, 2012**(54) **IDENTIFICATION OF A DNA VARIANT
ASSOCIATED WITH ADULT TYPE
HYPOLACTASIA**(75) Inventors: **Leena Peltonen**, Helsinki (FI); **Nabil
Enattah**, Helsinki (FI); **Irma Järvelä**,
Helsinki (FI); **Timo Sahi**, Helsinki (FI);
Erkki Savilahti, Hus (FI); **Joseph
Terwilliger**, New York, NY (US)(73) Assignee: **National Public Health Institute**,
Helsinki (FI)(21) Appl. No.: **13/594,731**(22) Filed: **Aug. 24, 2012****Related U.S. Application Data**(60) Division of application No. 10/775,501, filed on Feb.
9, 2004, now Pat. No. 8,252,537, which is a continua-
tion of application No. PCT/EP02/08963, filed on
Aug. 9, 2002.(60) Provisional application No. 60/315,955, filed on Aug.
31, 2001.(30) **Foreign Application Priority Data**

Aug. 10, 2001 (EP) 01119377.8

Aug. 14, 2001 (EP) 01119528.6

Publication Classification(51) **Int. Cl.****C12Q 1/68** (2006.01)**C07H 21/04** (2006.01)(52) **U.S. Cl.** **435/6.11**; 536/23.2; 536/24.33;
536/24.31(57) **ABSTRACT**

The present invention relates to a nucleic acid molecule comprising a 5' portion of an intestinal lactase-phlorizine hydrolase (LPH) gene contributing to or indicative of the adult-type hypolactasia. The present invention further relates to methods for testing for the presence of or predisposition to adult-type hypolactasia that are based on the analysis of an SNP contained in the above recited nucleic acid molecule. Additionally, the present invention relates to diagnostic composition and kit useful in the detection of the presence of or predisposition to adult-type hypolactasia.




Fig. 1




Fig.2

Fig. 3

Marker	Haplotype										Total
LPH13	6	6	6	6	6	6	6	6	6	6	6
LPH2	4	4	4	4	4	4	4	4	4	4	4
LPH1	3	2	2	2	1	1	4	4	4	4	4
AC7	5	5	5	5	5	5	5	5	5	5	5
AC3	2	2	2	4	2	2	2	2	2	2	2
AC4	5	5	5	5	5	5	5	5	5	5	5
AC5	6	6	5	6	6	3	6	5	5	3	3
Lactase persistent alleles	20	4	1	2	2	1	1	1	1	1	33
Lactase non-persistent alleles	3	2	0	0	0	0	2	0	0	2	54

SEQ ID NO: 1

ACCTTTCATTTCAGGAAAAATGTACTTAGACCCTACAATGTACTAGTAGGC
CTCTGCGCTGGCAATACAGATAAGATAATGTAGcCCCTGGCCTCAAAGGA
ACTCTCCTCCTTAGGTTGCATTTGTATAATGTTTGATTTTATAGATTGTTCTT
TGAGCCCTGCATTCCACGAGGATAGGTC

FIG. 4**SEQ ID NO: 2**

TAAGAACATTTTACACTCTTCAGTATAAAGAAGTCAGAATACCCCTACCCT
ATCAGTAAAGGCCTATAAGTTACCATTAAAAAGATGTCCTTAAAAACAGC
ATTCTCAGCTGGGCgCGGTGGCTCACACCTTTGTCCCAGTACTTTGGGAAG
CCGAGGTGGGTGGATCACCTGAGGTCAG

FIG. 5

SEQ ID NO: 3

ATCAGAGTCACTTTGATATGATGAGAGCAGAGATAAACAGATTTGTTGCA
TGTTTTTAATCTTTGGTATGGGACATACTAGAATTCAGTGCAAATACATTT
TTATGTAAGTGTGAATGCTCATACGACCATGGAATTCCTCCCTTTAAAGA
GCTTGGTAAGCATTGAGTGTAGTTGTTAGACGGAGACGATCACGTCATA
GTTTATAGAGTGCATAAAGACGTAAGTTACCATTTAATACCTTTTCATTGAG
GAAAAATGTACTTAGACCCTACAATGTACTAGTAGGCCTCTGCGCTGGCA
ATACAGATAAGATAATGTAGTCCCTGGCCTCAAAGGAACTCTCCTCCTTAG
GTTGCATTTGTATAATGTTTGATTTTTAGATTGTTCTTTGAGCCCTGCATTC
CACGAGGATAGGTCAGTGGGTATTAACGAGGTAAAAGGGGAGTAGTACG
AAAGGGCATTCAAGCGTCCCCTCTTCGCTTCAACCAAAGCAGCCCTGCGT
TTTCCTAGTTTTATTAATAGGTTTGATGTAAGGTCGCTTTTAAAAGGGGG
TTTGGCTTTTTTTTACAGTGTGACTGAGGTATAATTTATAAAAAGGGAAAT
GTATGGCATGGTGAGTTTTTTTACATACATCCTTGTGAATACCCAGCTCAA
GATCCAAAACATTTCCATAATTTTCAGAAAGTTCCAAACCCCTGCCTCTTTT
CAGTCTTAGCCCTCTTCCCCTGAAGTAACCACTGTTCCGACTTCAATCACT
ACTTTTATCCACAGGTTAATTTTTTTGGCTTTTTTTCCACTAAATTTTCAAAT
TCTTTGATATGGTACTTTACTATTGACGAAGTACTTTTACACTAGGTTATTT
AATATTCTTTGATTCACCCAATATTTAGGGAACACCTGTAGGGGACAAAA
AATGAATGAGAGCCCCCTGCCTTCCATTGCTGCTAATCTGGTGGAACGAG
ACATGTATTTAATTAAGCATGTAAAAAATAGAGTGGGTGATGAAATAATC
TATATACTAAATCCCCATGACACACAGTTTACCTATGTAACAAACCTGCAT
GTGTACCCCGAACCTAAATATAAGTTGGAAATTAACAAAAAACGAGA
GGGAGAATAGAGCATCACAACCAGAGTGCTGAGATGAATTAATTTATTAC
CAAAGAAGGAGGAGGACTCAGGGAGGTGCCGACGTTTAAACCCAGTCAC
TGAAGGGTGTGCAGAATTTGGATAGGCAAGATACCCTGGGACAAGGTCAT
TCTAAAACCATGCTAACATTTGTACTTTTTTTTTTTCATTGTGATAGTTCTGA
AATGAGTTGCATAAACTGGTACATGTCTTAGGGCAGTCTCTAATTGATTT
TTATTTTGTCTATTTTTTAAAAATTAGTCTTCAAATAGCAGATTCACATGAT
ATTAAAAATATATGCACATAAATTATATACACAAATATATTTTCTGAATGAA
ATTTAGTATCTGCATATATTTAAGAGCTATTTCTGTCTCATATGTTTATAAT
CTTCATCCATTAAAAAACTTTTGTAGGCCTTCTCACTCTAAGATTATA
AAAAATTCTCCCATTTTACCTAGCTAGTTTTCTAGTTGTTCCAAAACCA
TTTATTGAACAATCCATCTTTTTGACACTGGTTTGGCATGCCTTAATTATAT
ATTCTTGTGTGTGTTAGGATCTCCTTTTGGACTTTCCATTCTGTTTATTGAG
TCTTATCAGCTCCTCTTACATTGGTACCATGATGTTTTAATCTATGGGGCTT
TGTAGTTTTAAATGTAGGGCTAGTTCCAGCGCATTGTTCTCTATCAGCTGTT
AGGAACTTAGAAATCAGCTTGCTCTGTTTTAAAGAAAAACCTGGTATTTTT
TTATCAGTATAACATTCTATTTATATTAACCTGAAGAATTGAAAACATCTA
TGATTTTTCTTATTCAGTAACGTATCACTTAGAATAGGTTAGGTTGTACTA
CTATAAAATCTCAGCTGCATAAAACAATTTTTTTTTTGTGCTGTGCTACACAT
CCATTAGGTCATCAAGGGACTCACCTTGTCAGTTACTCAGAGATTCAGG
CTGATATAAAGGTTTGATCTTGACATACGCTTTCATGATGACAGAAAGCA

FIG. 6 (SHEET 1 OF 2)

GGGAAGAGAAGGTGGTGAGCCATGTGCTTTCTCCCCCTTCTATCCAGAAA
TGACACATACTCACATTTTCATTCGCCAGAGAAATTAACATGGCCCCCTCCTA
AGTTCAAATGGATAGAGAAATGCCTTCCTACCAGGTGCCCAGAAATTAGAA
GAGCAAACATTTGTGAACAGTTCTGAGTACCACAAATACCGTTATCTTTCC
ACTTAAGTCTTCTGTTTCACTCAGTAGTGCTTTAAACTTTTCTTCATATGTT
TTTCAGTGTTTCTTGTTGAATTTCTTGATATTTTATCATGTTTGTTTCGTA
GGAGTAGCCTTTTTTCCATTTTCATTTTCTGGCTGGTTTCATTGCTGGTTGT
TTTTTGTGTTTGTGTTTGTGTTTGTGAGATGGAGTCTCACTCTGTCGCCCAGGCT
GGAGTGCAGTGTCACAATCTCGGCTCACTGCAACCTCTGCCTCCCAGGTTC
AAGCGATTCTTCTTTCTCAGCCTCCTGAGTAGCTGGGATTACAGGCATGTG
CCACCATGCCCAGCTAATTTTTTATATTTTATAGTAGAGATGGGGTTTCTCC
ATGTTGGTCAGGCTGGTCTCAAACCTCCAATCTCAGGTGATCCGCCTGCCT
CTGCCTTCCAAAGTGCTGGGATTATAGACATGAGCCACCGTGCCTGGCCT
AGTTCTTATGGGATGTATATGTCTTTGGATTCATATGATATGTATATATGTT
TATATTTCTACAAGTACATACCTAGGAGTGGAATTGTTGGGTCATAGGTTA
ATGCATGTTTTTCTGCCAAACAGTTGTGTCAATTTCTGTTTTACCGCTGTG
AATGAGAGTTGTTCTACCTTCTTGACAACACTTGATATTGTCAGTCATTTT
AGCCATTCTGGTGAATTTATAGTGCTATTTCTGTGTGTGTAAGAGAGAGAA
TGAGAGAGGGTGTGTTGTGAGAAAACCAAAGCAACACTGTGAGAGTGTGTG
TGTTTGTGAGAAAACCAAATACTACTGTGATTTTCATTGGGAGAGAAA
ATCTGTTTGGTATATCAAAAAAAGTAGCTTAATTACTTCATCATTATTGGT
TTAGGT

FIG. 6 (SHEET 2 OF 2)

SEQ ID NO: 4

TAAGAACATTTTACACTCTTCAGTATAAAGAAGTCAGAATACCCCTACCCCT
ATCAGTAAAGGCCTATAAGTTACCATTAAAAAGATGTCCTTAAAAACAGC
ATTCTCAGCTGGGCaCGGTGGCTCACACCTTTGTCCCAGTACTTTGGGAAG
CCGAGGTGGGTGGATCACCTGAGGTCAGGAGTTCGAGACCAGCCTGGCCA
ACATGGCGAAAACCCATTTTCTCTACTAAAAATACAAAAATTAGCCGGGC
ATGGTGGCGGGTGCTTGTGGTCCCAGCTACTCAAGAGGCTGAGGTGGGAG
GATCACTGAGCCCAGGAGGTGGAGGCTGCATTGAGCCAAGATTGTGCCAC
TGCACTCCAGCCTGGGTGACAGAGCGAGACTCTGTCTCAAAAAAACCAAA
ACAAAAAAAACCCAGCATTCTTTAGTAAATAATTCATAGTTTTCTTCATCT
AGAATTTAAAATTGTGATAGTTGATCAGCATGTCCTGAGCACGTGTGTTTG
CTGTTACTAGTTTAGATCGGTAGATGTGTATATAAGTTATAGGTATAAAAT
CAATCCTGAGTTGACACAAGGTTTTGATGTTGAGTACAAGTACAGTAAGT
GTATATTTTTAGTTATGCTCTTAGTTTTAAGTCAATTGTGTGGTTCTTTCTA
GCTTTAGGATCTGTTGAATTATCTTCCTTAGAAAAGGGAGTTAAGAATCTT
CACTTACCTATCTTCTACTTGTTTGGAGAATAGAAGAGTCCCTGTGGTAGC
AGACTTTGTGAGTTTACTTGTAATTTTCCATCTGAAAGACTGTTCTTGTTTT
TCGTGATGAAGTCTTGCTCTGTCGCCCAGGCTGGAGTGCAGTGGTGCAAC
CTTGGCTCACTGCAACCTCTGCCTCCCGGGTTCAAGCAATTCTCCTGCCTC
AGCCTCCCGAGTATCTGGGATTACAGGTGCACACCACCACACCTGGCTAA
TTTTTGATTTTCAGTAGAGACGGGGTTTCACCATGTTGGCCAGGCTGGTC
TCGAACTCTTGACCTCATGATCAGCCACCTCAGCCTTCCAAAGTGCTGGG
ATTACAGGTGTGAGCCCCACACTCGGCCGTTGTTGTTTTTAAGAGACAG
GGTCTCACTCTGTACCTAACCTGGAGTACAGTGGCAATCATGGCTCACTG
TAACCTCAAATGCCCCGGCCTTAGTGAAGCGTTCTTCCTGCCTTGGCCTCCC
AAAGTGCTGGGATTACAAGTGTGAGCCATGCATCCAGCTTGAAAGACAGC
TTCTTAGGCTTGATTTGTTTGGTTACAGG

FIG. 7

SEQ ID NO: 5

ATCAGAGTCACTTTGATATGATGAGAGCAGAGATAAACAGATTTGTTGCA
TGTTTTTAATCTTTGGTATGGGACATACTAGAATTCAGTCAAATACATTT
TTATGTAAGTGTGAAATGCTCATAACGACCATGGAATTCCTCCCTTTAAAGA
GCTTGGTAAGCATTTGAGTGTAGTTGTTAGACGGAGACGATCACGTCATA
GTTTATAGAGTGCATAAAGACGTAAGTTACCATTTAATACCTTTTCATTTCAG
GAAAAATGTACTTAGACCCTACAATGTACTAGTAGGCCTCTGCGCTGGCA
ATACAGATAAGATAATGTAGCCCTGGCCTCAAAGGAACTCTCCTCCTTAG
GTTGCATTTGTATAATGTTTGATTTTTAGATTGTTCTTTGAGCCCTGCATTC
CACGAGGATAGGTGAGTGGGTATTAACGAGGTAAAAGGGGAGTAGTACG
AAAGGGCATTCAAGCGTCCCATCTTCGCTTCAACCAAAGCAGCCCTGCGT
TTTCCTAGTTTTATTAATAGGTTTGATGTAAGGTCGTCTTTGAAAAGGGGG
TTTGGCTTTTTTTTACAGTGTGACTGAGGTATAATTTATAAAAAGGGAAAT
GTATGGCATGGTGAGTTTTTTTACATACATCCTTGTGAATACCCAGCTCAA
GATCCAAAACATTTCCATAATTTTCAGAAAGTTCCAAACCCCTGCCTCTTTT
CAGTCTTAGCCCTCTTCCCCTGAAGTAACCACTGTTCCGACTTCAATCACT
ACTTTTATCCACAGGTTAATTTTTTTGGCTTTTTTCCACTAAATTTTCAAAT
TCTTTGATATGGTACTTTACTATTGACGAAGTACTTTCACACTAGGTTATTT
AATATTCTTTGATTCACCCAATATTTAGGGAACACCTGTAGGGGACAAAA
AATGAATGAGAGCCCCTGCCTTCCATTGCTGCTAATCTGGTGGGAACGAG
ACATGTATTTAATTAAGCATGTAAAAAATAGAGTGGGTGATGAAATAATC
TATATACTAAATCCCCATGACACACAGTTTACCTATGTAACAAACCTGCAT
GTGTACCCCCGAACCTAAAATATAAGTTGGAAATTAAAAAAAACGAGA
GGGAGAATAGAGCATCACAACCAGAGTGCTGAGATGAATTACTTTATTAC
CAAAGAAGGAGGAGGACTCAGGGAGGTGCCGACGTTTAAACCCAGTCAC
TGAAGGGTGTGCAGAATTTGGATAGGCAAGATACCCTGGGACAAGGTCAT
TCTAAAACCATGCTAACATTTGTACTTTTTTTTTTCATTGTGATAGTTCCTGA
AATGAGTTGCATAAACTGGTACATGTCTTAGGGCAGTCTCTAATTGATTT
TTATTTTGTCTATTTTTTAAAAATTAGTCTTCAAATAGCAGATTCACATGAT
ATTAAAATATATGCACATAAATTATATACACAAATATATTTTCTGAATGAA
ATTTAGTATCTGCATATATTTAAGAGCTATTTCTGTCTCATATGTTTCATAAT
CTTCATCCATTAAAAAACTTTTTGTTAGGCCTTTCTCACTCTAAGATTATA
AAAAATTCTCCCATTTATTTACCTAGCTAGTTTTCTAGTTGTTCCAAAACCA
TTTATTGAACAATCCATCTTTTTGACACTGGTTTGGCATGCCTTAATTATAT
ATTCTTGTGTGTGTAGGATCTCCTTTTGGACTTTCCATTCTGTTTCATTGAG
TCTTATCAGCTCCTCTTACATTGGTACCATGATGTTTTAATCTATGGGGCTT
TGTAGTTTTAAATGTAGGGCTAGTTCAGCGCATTGTTCTCTATCAGCTGTT
AGGAACTTAGAAATCAGCTTGCTCTGTTTTAAAGAAAAACCTGGTATTTTT
TTATCAGTATAACATTCTATTTATATTAACCTGAAGAATTGAAAACATCTA
TGATTTTTCTTATTCAGTAACGTATCACTTAGAATAGGTTAGGTTGTACTA
CTATAAAATCTCAGCTGCATAAAACAATTTTTTTTTTGCTTGTGCTACACAT
CCATTAGGTCATCAAGGGACTCACCTTGTCAAGTTACTCAGAGATTCAGG

FIG. 8 (SHEET 1 OF 2)

CTGATATAAAGGTTTGATCTTGACATACGCTTTCATGATGACAGAAAGCA
GGGAAGAGAAGGTGGTGAGCCATGTGCTTCTCCCCCTTCTATCCAGAAA
TGACACATACTCACATTTCAATTCGCCAGAGAAATTAACATGGCCCCCTCCTA
AGTTCAAATGGATAGAGAAAATGCCTTCCTACCAGGTGCCCAGAATTAGAA
GAGCAAACATTTGTGAACAGTTCTGAGTACCACAAATACCGTTATCTTTCC
ACTTAAGTCTTCTGTTTCACTCAGTAGTGCTTTAACTTTTCTTCATATGTT
TTTCAGTGTTTCTTGTTGAATTTCTTGATATTTTATCATGTTTGTTTCGTA
GGAGTAGCCTTTTTTCCATTTCAATTTCTGGCTGGTTTCATTGCTGGTTGT
TTTTTGTGTTTGTGTTTGTGTTTGTGAGATGGAGTCTCACTCTGTCGCCCAGGCT
GGAGTGCAGTGTCACAATCTCGGCTCACTGCAACCTCTGCCTCCCAGGTTC
AAGCGATTCTTCTTCTCAGCCTCCTGAGTAGCTGGGATTACAGGCATGTG
CCACCATGCCCAGCTAATTTTTTATATTTTAGTAGAGATGGGGTTTCTCC
ATGTTGGTCAGGCTGGTCTCAAACCTCCAATCTCAGGTGATCCGCCTGCCT
CTGCCTTCCAAAGTGCTGGGATTATAGACATGAGCCACCGTGCCTGGCCT
AGTTCTTATGGGATGTATATGTCTTTGGATTTCATATGATATGTATATATGTT
TATATTTCTACAAGTACATACCTAGGAGTGGAATTGTTGGGTCATAGGTTA
ATGCATGTTTTTCTGCCAAACAGTTGTGTCAATTTCTGTTTTACCGCTGTG
AATGAGAGTTGTTCTACCTTCTTGACAACACTTGATATTGTCAGTCATTTT
AGCCATTCTGGTGAATTTATAGTGCTATTTCTGTGTGTGTAAGAGAGAGAA
TGAGAGAGGGTGTTTGTGAGAAAACCAAAGCAACACTGTGAGAGTGTGTG
TGTTTGTGAGAAAACCAAATACTACTGTGATTTTCATTGGGAGAAA
ATCTGTTTGGTATATCAAAAAAGTAGCTTAATTACTTCATCATTATTGGT
TTAGGT

Fig. 8 (SHEET 2 OF 2)

SEQ ID NO: 6

TAAGAACATTTTACACTCTTCAGTATAAAGAAGTCAGAATACCCCTACCCT
ATCAGTAAAGGCCTATAAGTTACCATTAAAAAGATGTCCTTAAAAACAGC
ATTCTCAGCTGGGCgCGGTGGCTCACACCTTTGTCCCAGTACTTTGGGAAG
CCGAGGTGGGTGGATCACCTGAGGTCAGGAGITCGAGACCAGCCTGGCCA
ACATGGCGAAAACCCATTTTCTCTACTAAAAATACAAAAATTAGCCGGGC
ATGGTGGCGGGTGCTTGTGGTCCCAGCTACTCAAGAGGCTGAGGTGGGAG
GATCACTGAGCCCAGGAGGTGGAGGCTGCATTGAGCCAAGATTGTGCCAC
TGCACTCCAGCCTGGGTGACAGAGCGAGACTCTGTCTCAAAAAAACAAA
ACAAAAAAAACCCAGCATTCTTTAGTAAATAATTCATAGTTTTCTTCATCT
AGAATTTAAAAATTGTGATAGTTGATCAGCATGTCCTGAGCACGTGTGTTTG
CTGTTACTAGTTTATAGATCGGTAGATGTGTATATAAGTTATAGGTATAAAAT
CAATCCTGAGTTGACACAAGGTTTTTGATGTTGAGTACAAGTACAGTAAGT
GTATATTTTTAGTTATGCTCTTAGTTTTTAAGTCAATTGTGTGGTTCTTTCTA
GCTTTAGGATCTGTTGAATTATCTTCCTTAGAAAAGGGAGTTAAGAATCTT
CACTTACCTATCTTCTACTTGTTTGGAGAATAGAAGAGTCCCTGTGGTAGC
AGACTTTGTGAGTTTACTTGTAATTTCCATCTGAAAGACTGTTCTTGTTTT
TCGTGATGAAGTCTTGCTCTGTGCGCCAGGCTGGAGTGCAGTGGTGCAAC
CTTGGCTCACTGCAACCTCTGCCTCCCGGGTTCAAGCAATTCTCCTGCCTC
AGCCTCCCGAGTATCTGGGATTACAGGTGCACACCACCACACCTGGCTAA
TTTTTGATTTTTCAGTAGAGACGGGGTTTCACCATGTTGGCCAGGCTGGTC
TCGAACTCTTGACCTCATGATCAGCCCACCTCAGCCTTCCAAAGTGCTGGG
ATTACAGGTGTGAGCCCCCAGCTCGGCCGTTGTTGTTTTTTAAGAGACAG
GGTCTCACTCTGTACCTAACCTGGAGTACAGTGGCAATCATGGCTCACTG
TAACCTCAAATGCCCGGCCTTAGTGAAGCGTTCTTCCTGCCTTGGCCTCCC
AAAGTGCTGGGATTACAAGTGTGAGCCATGCATCCAGCTTGAAAGACAGC
TTCTTAGGCTTGATTTGTTTGTTACAGG

FIG. 9

IDENTIFICATION OF A DNA VARIANT ASSOCIATED WITH ADULT TYPE HYPOLACTASIA

[0001] The present application is a divisional of U.S. patent application Ser. No. 10/775,501, filed Feb. 9, 2004, which application is a continuation application of PCT/EP02/08963, filed Aug. 9, 2002, which claims priority to European Patent Application No. 01119377.8, filed Aug. 10, 2001, European Patent Application No. 01119528.6, filed Aug. 14, 2001, and U.S. Provisional Application No. 60/315,955, filed Aug. 31, 2001, the teachings of which are hereby incorporated by reference in their entirety.

[0002] The present invention relates to a nucleic acid molecule comprising a 5' portion of an intestinal lactase-phlorizin hydrolase (LPH) gene contributing to or indicative of the adult-type hypolactasia wherein said nucleic acid molecule is selected from the group consisting of (a) a nucleic acid molecule having or comprising the nucleic acid sequence of SEQ ID NO:1, the sequence of SEQ ID NO:1 is also depicted in FIG. 4 and comprised in the sequence as depicted in FIG. 8; (b) a nucleic acid molecule having or comprising the nucleic acid sequence of SEQ ID NO:2, the sequence of SEQ ID NO:2 is also depicted in FIG. 5 and comprised in the sequence as depicted in FIG. 9; (c) a nucleic acid molecule of at least 20 nucleotides in the complementary strand of which hybridizes under stringent conditions to the nucleic acid molecule of (a) or (b), wherein said polynucleotide/nucleic acid molecule has at a position corresponding to position -13910 5' from the LPH gene a cytosine residue; and (d) a nucleic acid molecule of at least 20 nucleotides the complementary strand of which hybridizes under stringent conditions to the nucleic acid molecule of (a) or (b), wherein said polynucleotide/nucleic acid molecule has at a position corresponding to position -22018 5' from the LPH gene a guanine residue. The present invention further relates to methods for testing for the presence of or predisposition to adult-type hypolactasia that are based on the analysis of an SNP contained in the above recited nucleic acid molecule. Additionally, the present invention relates to diagnostic composition and kit useful in the detection of the presence of or predisposition to adult-type hypolactasia.

[0003] A variety of documents are cited throughout this specification. The disclosure content of these documents, including manufacturer's manuals and catalogues, is hereby incorporated by reference.

[0004] Lactase-phlorizin hydrolase enzyme (LPH), which is exclusively expressed by intestinal epithelial cells, hydrolyses lactose, sugar of milk, into glucose and galactose¹. The expression of the LPH enzyme dramatically declines to very low levels at the weaning period in mammals when lactose is no longer an essential part of the diet. In humans, the condition known as adult-type hypolactasia or lactase non-persistence, affects most populations and severely limits the use of fresh milk among adults due to lactose intolerance. The age of onset of lactase non-persistence status varies between populations, ranging from 1-2 years of age among the Thais to 10-20 years of age among the Finns²⁻³. However, in Northern European and a few other ethnic groups, LPH activity persists throughout life in the majority of adults, a condition known as lactase persistence. The phenotype lactase persistence/non-persistence has been shown to be genetically determined, the persistent status being dominant over the non-persistent status⁴⁻⁶.

[0005] The state of the art diagnosis of adult-type hypolactasia is based on the lactose tolerance test (LTT). After over-

night fasting (10 hours), 1 g/kg of lactose is given as a 12.5% solution, the maximum dose being 50 g. Capillary blood samples are taken before and 20 and 30 min after lactose ingestion. The glucose concentration is determined by the glucose oxidase method (Hjelm and de Verdier 1963). Abdominal symptoms on the day of LTT are noted. A maximum rise in blood glucose concentration of 1.1 mmol/l or more was taken as a sign of lactose malabsorption (Gudman-Hoyer and Harnum 1968, Jussila 1970, Sahi 1972). LTT contains a 10% risk for false positive and negative diagnoses, i.e. the sensitivity and specificity of

[0006] LTT is about 90% (Isokoski et al. 1972, Newcomer et al. 1975, Sahi 1983). The accuracy of LTT can be improved by giving 0.3 g/kg ethanol that inhibits the metabolism of galactose in the liver (Tygstrup and Lundqvist 1962) and 15 min later 1 g/kg lactose as 12.5% solution.

[0007] Children with maximum rises of less than 0.2 mg/100 ml in the first or repeated LTT have been sent for small-intestinal biopsy that is taken through gastroscopy. This is an invasive procedure that needs expertise and is usually performed at university hospitals by specialists in gastroenterology only. Biopsy samples are examined with a dissection microscope and histologically, and the mucosal maltase, sucrase and lactase activities are determined (Lau-niala et al. 1964). The diagnosis of hypolactasia in children is justified if the histology of the intestinal biopsy is normal and lactase activity is less than 20 U/g protein and lactase/sucrase ratio less than 0.30, or in the LTT with ethanol administration a maximum rise in blood glucose concentration of less than 20 mg/100 ml and in galactose concentration of 5 mg/100 ml or less (Sahi et al, 1972) is demonstrated. As described above, the current methods to diagnose adult-type hypolactasia are laborious. LTT is inexact and therefore, an invasive procedure, gastroscopy is needed before the diagnosis can be ascertained. Since adult-type hypolactasia is very common and the major cause of nonspecific abdominal symptoms (in one third of patients complaining stomach pain), there is a clear need to improve the diagnostics of this common health problem.

[0008] Yet, so far no biochemical test that is easy to handle and, at the same time, provides quick and accurate results has been developed. Elucidation of the cause of the disease on the genomic DNA/expression level has equally been unsuccessful. Thus, the sequencing of the coding and promoter regions of the LPH gene in adults has revealed no DNA-variations which correlate with lactase persistence/non-persistence, nor has evidence emerged of splice variants or mRNA editing variants associated with this trait⁷⁻⁸. Previous studies have shown that the lactase persistence/non-persistence trait is possibly controlled by cis-acting element(s) residing within or adjacent to the lactase gene, and strong linkage disequilibrium (LD) has been observed across the 70 kb haplotype spanning the lactase gene^{9,10}. Several studies report evidence that the main control of the LPH gene expression operates at the level of transcription regulation¹¹⁻¹³. However, it has been suggested that variation influencing both transcriptional and posttranscriptional control of expression of the LPH gene may be involved in the etiology of adult-type hypolactasia¹⁴⁻¹⁵.

[0009] In view of the above, the technical problem underlying the present invention was to provide means and methods that allow for an accurate and convenient diagnosis of adult-type hypolactasia or of a predisposition to this disease.

[0010] The solution to said technical problem is achieved by the embodiments characterized in the claims.

[0011] Thus, the present invention relates to a nucleic acid molecule comprising a 5' portion of an intestinal lactase-phlorizine hydrolase (LPH) gene contributing to or indicative of adult-type hypolactasia wherein said nucleic acid molecule is selected from the group consisting of (a) a nucleic acid molecule having or comprising the nucleic acid sequence of SEQ ID NO: 1, the sequence of SEQ ID NO:1 is also depicted in FIG. 4 and comprised in the sequence as depicted in FIG. 8; (b) a nucleic acid molecule having or comprising the nucleic acid sequence of SEQ ID NO: 2, the sequence of SEQ ID NO:2 is also as depicted in FIG. 5 and comprised in the sequence as depicted in FIG. 9; (c) a nucleic acid molecule of at least 20 nucleotides the complementary strand of which hybridizes under stringent conditions to the nucleic acid molecule of (a) or (b), wherein said polynucleotide/nucleic acid molecule has at a position corresponding to position -13910 5' from the LPH gene a cytosine residue; and (d) a nucleic acid molecule of at least 20 nucleotides the complementary strand of which hybridizes under stringent conditions to the nucleic acid molecule of (a) or (b), wherein said polynucleotide/nucleic acid molecule has at a position corresponding to position -22018 5' from the LPH gene a guanine residue.

[0012] In accordance with the invention, the term "intestinal lactase-phlorizine hydrolase (LPH) gene" denotes a gene that encodes an enzyme having the activity of hydrolyzing lactose into its components glucose and galactose. The enzyme is characterized by E.C. 3.2.1.23.62.

[0013] The term "adult-type hypolactasia" refers to a condition also known as lactose intolerance, which is an autosomal recessive condition resulting from the "physiological" decline of the lactase-phlorizin hydrolase (LPH) enzyme activity in intestinal cells in a significant proportion of the global population.

[0014] The term "contributing to or indicative of adult-type hypolactasia", refers to the fact that the SNPs and thus the corresponding nucleic acid molecules found are indicative of the condition and possibly also causative therefore. Accordingly, this term necessarily requires that the recited 5' position is indicative of the condition. Said term, on the other hand, does not necessarily require that the 5' portion is causative or contributes to the condition. Yet, said term does not exclude a causative or contributory role of either or both SNPs.

[0015] The term "which hybridizes under stringent conditions" refers to hybridization conditions that are well known to or can be established by the person skilled in the art according to conventional protocols. The term most advantageously refers to highly stringent conditions. Appropriate stringent conditions for each sequence may be established on the basis of well-known parameters such as temperature, composition of the nucleic acid molecules, salt conditions etc.: see, for example, Sambrook et al., "Molecular Cloning, A Laboratory Manual"; CSH Press, Cold Spring Harbor, 1989 or Higgins and Hames (eds.), "Nucleic acid hybridization, a practical approach", IRL Press, Oxford 1985 (reference 54), see in particular the chapter "Hybridization Strategy" by Britten & Davidson, 3 to 15. Typical (highly stringent) conditions comprise hybridization at 65° C. in 0.5×SSC and 0.1% SDS or hybridization at 42° C. in 50% formamide, 4×SSC and 0.1% SDS. Hybridization is usually followed by washing to remove unspecific signal. Washing conditions include conditions such as 65° C., 0.2×SSC and 0.1% SDS or 2×SSC and 0.1% SDS or 0.3×SSC and 0.1% SDS at 25° C.-65° C.

[0016] As disclosed herein above, the present invention also relates to a hybridizing nucleic acid molecules of at least

20 nucleotides; see (c) and (d) herein above. Yet, the present invention also relates to a nucleic acid molecule of at least 50, at least 100, at least 150, or at least 200 nucleotides. Preferably, said hybridizing fragments comprise at least 25, at least 50, or at least 75 nucleotides, at least 100 nucleotides, 5' and 3' of the position -13910 as defined in (c) or of position -22018 as defined in (d) herein above.

[0017] The term "nucleic acid molecule" refers both to naturally and non-naturally occurring nucleic acid molecules. Non-naturally occurring nucleic acid molecules include cDNA as well as derivatives such as PNA.

[0018] The term "nucleic acid molecule [. . .] comprising the nucleic acid sequence of SEQ ID NO:" throughout this specification refers to nucleic acid molecules that are at least 1 nucleotide longer than the nucleic acid molecule specified by the SEQ ID NO. At the same time, these nucleic acid molecules extend, at a maximum, 30000 nucleotides over the 5' and/or 3' end of the nucleic acid molecule of the invention specified e.g. by the SEQ ID NO: 2 or 1, 3 or 4.

[0019] Surprisingly, it was found in accordance with the present invention that the two hypolactasia-associated variants locate at a considerable distance from the LPH gene, positioned in different introns of the MCM6 gene. MCM6 is a member of a gene family (MCM 2-7), required for the initiation of DNA replication ensuring that it takes place only once during the cell cycle³¹. MCM6, unlike LPH, is not restricted in its tissue distribution and there is no correlation in the levels of MCM6 and LPH transcripts¹⁸. These findings would suggest that these two genes do not share any functionally significant cis-acting elements providing tissue specificity or developmental regulation¹⁸. Most probably the identified variants have different functional significance for the expression of the LPH and MCM6 genes. Further surprisingly, based on complete association to hypolactasia they (or one of them) are associated to age-dependent down regulation of the transcript level of the LPH gene in the intestinal epithelium but have little or no effect on the transcription of the MCM6.

[0020] Experimentally, using linkage, allelic association and extended haplotype analysis carried out in nine extended Finnish families the adult-type hypolactasia locus was restricted to a 47 kb interval on 2q21. The sequence analysis of the region revealed a single nucleotide polymorphism (SNP), C/T-13910 that completely cosegregated with adult-type hypolactasia in all Finnish families and in a sample set of 236 individuals from four different populations. Another SNP G/A-22018 residing 8 kb telomeric from C/T -13910 was associated with the trait in all but 7 cases. The prevalence of C/T -13910 SNP in 1047 DNA samples reflected the reported prevalence of adult-type hypolactasia in three different populations providing additional evidence for its importance for the trait.

[0021] The surprising finding referred to above for the first time allows the establishment of test systems that are based on the molecular analysis of the recited single nucleotide polymorphisms upstream of the LPH gene. Whereas both SNPs provide for a solid basis for the diagnosis of or the diagnosis of a predisposition to adult-type hypolactasia, it is preferred that the nucleotide position -13910 is analyzed, either alone or in combination with nucleotide position -22018. This is because the SNP at position -13910 was associated in 100% of the analysed cases with the disease whereas the SNP at position -22018 was associated in only 98% of all cases with adult-type hypolactasia. Nevertheless, analyses of nucleotide

position -22018 alone will usually also provide a sound basis for a diagnosis of a predisposition to adult-type hypolactasia.

[0022] Due to the abundance of established methods for assessing for the presence of SNPs, it is now possible to conveniently, in a short amount of time, at low cost, with high accuracy and without significant trouble for the person under investigation, diagnose a genetic predisposition to adult-type hypolactasia.

[0023] The invention further relates to a nucleic acid molecule comprising a 5' portion of an intestinal lactase-phlorizin hydrolase (LPH) gene wherein said nucleic acid molecule is selected from the group consisting of (a) a nucleic acid molecule having or comprising the nucleic acid sequence of SEQ ID NO:3, the sequence of SEQ ID NO:3 is also depicted in FIG. 6; (b) a nucleic acid molecule having or comprising the nucleic acid sequence of SEQ ID NO:4, the sequence of SEQ ID NO:4 is also depicted in FIG. 7; (c) a nucleic acid molecule the complementary strand of which hybridizes under stringent conditions to the nucleic acid molecule of (a) or (b), wherein said polynucleotide/nucleic acid molecule has at a position corresponding to position -13910 of the LPH gene a thymidine residue; and (d) a nucleic acid molecule the complementary strand of which hybridizes under stringent conditions to the nucleic acid molecule of (a) or (b), wherein said polynucleotide/nucleic acid molecule has at a position corresponding to position -22018 of the LPH gene a adenosine residue.

[0024] This embodiment of the present invention may conveniently be used to demonstrate that a person does not suffer from adult-type hypolactasia and has no predisposition therefor. Further, this nucleic acid molecule reflecting the "wild-type" situation of the position -13910 or -22018 upstream of the LPH gene may be used as a control means in experiments where a predisposition to adult-type hypolactasia is tested for. For testing, methods as described throughout this specification may be used.

[0025] In a preferred embodiment of the invention the nucleic acid molecule is genomic DNA.

[0026] This preferred embodiment of the invention reflects the fact that usually the analysis would be carried out on the basis of genomic DNA from body fluid, cells or tissue isolated from the person under investigation.

[0027] In a further preferred embodiment of the invention said genomic DNA is part of a gene.

[0028] In accordance with the invention, it is preferred that at least one of the introns of the MCM6 gene harboring position -13910 or position -22018 relative to the LPH gene is analyzed.

[0029] In addition, the invention relates to a fragment of the nucleic acid molecule as described herein above having at least 14 nucleotides wherein said fragment comprises nucleotide position -13910 or nucleotide position -22018 (upstream) of the LPH gene.

[0030] The fragment of the invention may be of natural as well as of (semi)synthetic origin. Thus, the fragment may, for example, be a nucleic acid molecule that has been synthesized according to conventional protocols of organic chemistry. Importantly, the nucleic acid fragment of the invention comprises nucleotide position -13910 or nucleotide position -22018 upstream of the LPH gene. In these positions, the fragment may have either the wild-type nucleotide or the nucleotide contributing to or indicative of adult-type hypolactasia (also referred to as the "mutant" sequence). Conse-

quently, the fragment of the invention may be used, for example, in assays differentiating between the wild-type and the mutant sequence.

[0031] It is further preferred that the fragment of the invention consists of at least 17 nucleotides, more preferred at least 21 nucleotides, and most preferred at least 25 nucleotides such as 30 nucleotides.

[0032] Furthermore, the invention relates to a nucleic acid molecule which is complementary to the nucleic acid molecule as described herein above.

[0033] This embodiment of the invention comprising at least 14 nucleotides and covering at least position -13910 or position -22018 of the sequence upstream of the LPH gene is particularly useful in the analysis of the genetic setup in the recited positions in hybridization assays. Thus, for example, a 15mer exactly complementary either to the wild-type sequence (i.e. a T in position -13910 or an A in position -22018) or to the variants contributing to or indicative of adult-type hypolactasia (i.e. a C in position -13910 or a G in position -22018) may be used to differentiate between the polymorphic variants. This is because a nucleic acid molecule labeled with a detectable label not exactly complementary to the DNA in the analyzed sample will not give rise to a detectable signal, if appropriate hybridization and washing conditions are chosen.

[0034] In this regard, it is important to note that the nucleic acid molecule of the invention, the fragment thereof as well as the complementary nucleic acid molecule may be detectably labeled. Detectable labels include radioactive labels such as ³H, or ³²P or fluorescent labels. Labeling of nucleic acids is well understood in the art and described, for example, in Sambrook et al., loc. cit.

[0035] In addition, the invention relates to a vector comprising the nucleic acid molecule as described herein above. The vector of the invention may either contain a nucleic acid molecule comprising the wild-type sequence(s) or it may contain a nucleic acid molecule comprising the mutant sequence(s).

[0036] The vectors may particularly be plasmids, cosmids, viruses or bacteriophages used conventionally in genetic engineering that comprise the nucleic acid molecule of the invention. Preferably, said vector is an expression vector and/or a gene transfer or targeting vector. Expression vectors derived from viruses such as retroviruses, vaccinia virus, adeno-associated virus, herpes viruses, or bovine papilloma virus, may be used for delivery of the nucleic acid molecule of the invention into targeted cell population. Methods which are well known to those skilled in the art can be used to construct recombinant viral vectors; see, for example, the techniques described in Sambrook et al., loc. cit. and Ausubel et al., *Current Protocols in Molecular Biology*, Green Publishing Associates and Wiley Interscience, N.Y. (1989). Alternatively, the nucleic acid molecules and vectors of the invention can be reconstituted into liposomes for delivery to target cells. The vectors containing the nucleic acid molecules of the invention can be transferred into the host cell by well-known methods, which vary depending on the type of cellular host. For example, calcium chloride transfection is commonly utilized for prokaryotic cells, whereas, e.g., calcium phosphate or DEAE-Dextran mediated transfection or electroporation may be used for other cellular hosts; see Sambrook, supra.

[0037] Such vectors may comprise further genes such as marker genes which allow for the selection of said vector in a suitable host cell and under suitable conditions. Preferably,

the nucleic acid molecule of the invention is operatively linked to expression control sequences allowing expression in prokaryotic or eukaryotic cells. Expression of said polynucleotide comprises transcription of the polynucleotide into a translatable mRNA. Regulatory elements ensuring expression in eukaryotic cells, preferably mammalian cells, are well known to those skilled in the art. They usually comprise regulatory sequences ensuring initiation of transcription and, optionally, a poly-A signal ensuring termination of transcription and stabilization of the transcript, and/or an intron further enhancing expression of said polynucleotide. Additional regulatory elements may include transcriptional as well as translational enhancers, and/or naturally-associated or heterologous promoter regions. Possible regulatory elements permitting expression in prokaryotic host cells comprise, e.g., the PL, lac, trp or tac promoter in *E. coli*, and examples for regulatory elements permitting expression in eukaryotic host cells are the AOX1 or GAL1 promoter in yeast or the CMV-, SV40-, RSV-promoter (Rous sarcoma virus), CMV-enhancer, SV40-enhancer or a globin intron in mammalian and other animal cells. Beside elements which are responsible for the initiation of transcription such regulatory elements may also comprise transcription termination signals, such as the SV40-poly-A site or the tk-poly-A site, downstream of the polynucleotide. Optionally, the heterologous sequence can encode a fusion protein including an C- or N-terminal identification peptide imparting desired characteristics, e.g., stabilization or simplified purification of expressed recombinant product. In this context, suitable expression vectors are known in the art such as Okayama-Berg cDNA expression vector pcDV1 (Pharmacia), pCDM8, pRc/CMV, pcDNA1, pcDNA3, the Echo™ Cloning System (Invitrogen), pSPORT1 (GIBCO BRL) or pRevTet-On/pRevTet-Off or pCI (Promega). Preferably, the expression control sequences will be eukaryotic promoter systems in vectors capable of transforming or transfecting eukaryotic host cells, but control sequences for prokaryotic hosts may also be used.

[0038] As mentioned above, the vector of the present invention may also be a gene transfer or targeting vector. Gene therapy, which is based on introducing therapeutic genes into cells by ex-vivo or in-vivo techniques is one of the most important applications of gene transfer. Suitable vectors and methods for in-vitro or in-vivo gene therapy are described in the literature and are known to the person skilled in the art; see, e.g., Giordano, *Nature Medicine* 2 (1996), 534-539; Schaper, *Circ. Res.* 79 (1996), 911-919; Anderson, *Science* 256 (1992), 808-813; Isner, *Lancet* 348 (1996), 370-374; Muhlhauser, *Circ. Res.* 77 (1995), 1077-1086; Wang, *Nature Medicine* 2 (1996), 714-716; WO94/29469; WO 97/00957, Schaper, *Current Opinion in Biotechnology* 7 (1996), 635-640, or Kay et al. (2001) *Nature Medicine*, 7, 33-40 and references cited therein. The polynucleotides and vectors of the invention may be designed for direct introduction or for introduction via liposomes, or viral vectors (e.g. adenoviral, retroviral) into the cell. Preferably, said cell is a germ line cell, embryonic cell, or egg cell or derived therefrom, most preferably said cell is a stem cell. Gene therapy is envisaged with the wild-type nucleic acid molecule only.

[0039] The invention as well relates to a primer or primer pair, wherein the primer or primer pair hybridizes under (highly) stringent conditions to the nucleic acid as described herein above comprising nucleotide position -13910 or -22018 of the LPH gene or to the complementary strand thereof.

[0040] Preferably, the primers of the invention have a length of at least 14 nucleotides such as 17 or 21 nucleotides. It is further preferred that the primers have a maximum length of 24 nucleotides. Hybridization or lack of hybridization of a primer under appropriate conditions to a genome sequence comprising either position -13910 or position -22018 coupled with an appropriate detection method such as an elongation reaction or an amplification reaction may be used to differentiate between the polymorphic variants and then draw conclusions with regard to, e.g., the predisposition of the person under investigation for adult-type hypolactasia. The present invention envisages two types of primers/primer pairs. One type hybridizes to a sequence comprising the mutant sequence. In other words, the primer is exactly complementary to a sequence that contains the C in position -13910 or the G in position -22018 or to the complementary strand thereof. The other type of primer is exactly complementary to a sequence having a T in position -13910 or an A in position -22018 or to the complementary strand thereof. Since hybridization conditions would preferably be chosen to be stringent enough, contacting of e.g. a primer exactly complementary to the mutant sequence with a wild-type allele would not result in efficient hybridization due to the mismatch formation. After washing, no signal would be detected due to the removal of the primer.

[0041] Additionally, the invention relates to a non-human host transformed with the vector of the invention as described herein above. The host may either carry the mutant or the wild-type sequence. Upon breeding etc. the host may be heterozygous or homozygous for one or both SNPs.

[0042] The host of the invention may carry the vector of the invention either transiently or stably integrated into the genome. Methods for generating the non-human host of the invention are well known in the art. For example, conventional transfection protocols described in Sambrook et al., loc. cit., may be employed to generate transformed bacteria (such as *E. coli*) or transformed yeasts. The non-human host of the invention may be used, for example, to elucidate the onset of adult-type hypolactasia.

[0043] In a preferred embodiment of the invention the non-human host is a bacterium, a yeast cell, an insect cell, a fungal cell, a mammalian cell, a plant cell, a transgenic animal or a transgenic plant.

[0044] Whereas *E. coli* is a preferred bacterium, preferred yeast cells are *S. cerevisiae* or *Pichia pastoris* cells. Preferred fungal cells are *Aspergillus* cells and preferred insect cells include *Spodoptera frugiperda* cells. Preferred mammalian cells are colon carcinoma cell lines showing expression of the LPH enzyme and include CaCo2-cells.

[0045] A method for the production of a transgenic non-human animal, for example transgenic mouse, comprises introduction of the aforementioned polynucleotide or targeting vector into a germ cell, an embryonic cell, stem cell or an egg or a cell derived therefrom. The non-human animal can be used in accordance with a screening method of the invention described herein. Production of transgenic embryos and screening of those can be performed, e.g., as described by A. L. Joyner Ed., *Gene Targeting, A Practical Approach* (1993), Oxford University Press. The DNA of the embryonal membranes of embryos can be analyzed using, e.g., Southern blots with an appropriate complementary nucleic acid molecule; see supra. A general method for making transgenic non-human animals is described in the art, see for example WO 94/24274. For making transgenic non-human organisms

(which include homologously targeted non-human animals), embryonal stem cells (ES cells) are preferred. Murine ES cells, such as AB-1 line grown on mitotically inactive SNL76/7 cell feeder layers (McMahon and Bradley, *Cell* 62:1073-1085 (1990)) essentially as described (Robertson, E. J. (1987) in *Teratocarcinomas and Embryonic Stem Cells: A Practical Approach*, E. J. Robertson, ed. (Oxford: IRL Press), p. 71-112) may be used for homologous gene targeting. Other suitable ES lines include, but are not limited to, the E14 line (Hooper et al., *Nature* 326:292-295 (1987)), the D3 line (Doetschman et al., *J. Embryol. Exp. Morph.* 87:27-45 (1985)), the CCE line (Robertson et al., *Nature* 323:445-448 (1986)), the AK-7 line (Zhuang et al., *Cell* 77:875-884 (1994)). The success of generating a mouse line from ES cells bearing a specific targeted mutation depends on the pluripotency of the ES cells (i. e., their ability, once injected into a host developing embryo, such as a blastocyst or morula, to participate in embryogenesis and contribute to the germ cells of the resulting animal). The blastocysts containing the injected ES cells are allowed to develop in the uteri of pseudopregnant nonhuman females and are born as chimeric mice. The resultant transgenic mice are chimeric for cells having the desired nucleic acid molecule are backcrossed and screened for the presence of the correctly targeted transgene(s) by PCR or Southern blot analysis on tail biopsy DNA of offspring so as to identify transgenic mice heterozygous for the nucleic acid molecule of the invention.

[0046] The transgenic non-human animals may, for example, be transgenic mice, rats, hamsters, dogs, monkeys (apes), rabbits, pigs, or cows. Preferably, said transgenic non-human animal is a mouse. The transgenic animals of the invention are, inter alia, useful to study the phenotypic expression/outcome of the nucleic acids and vectors of the present invention. Furthermore, the transgenic animals of the present invention are useful to study the developmental expression of the LPH enzyme, for example in the rodent intestine. It is furthermore envisaged, that the non-human transgenic animals of the invention can be employed to test for therapeutic agents/compositions or other possible therapies which are useful to ameliorate adult-type hypolactasia.

[0047] In addition, the invention relates to an antibody or aptamer or phage that specifically binds to the mutant nucleic acid molecule of the invention but not to the corresponding wild type nucleic acid molecule.

[0048] The antibody may be tested for binding and used in any serologic technique well known in the art, such as agglutination techniques in tubes, gels, solid phase and capture techniques with or without secondary antibodies, or in flow cytometry with or without immunofluorescence enhancement (see, for example, techniques described in Harlow and Lane "Antibodies, A Laboratory Manual", CSH Press, Cold Spring Harbor, USA, 1988 (see reference 53)).

[0049] In line with the invention, the antibody specifically recognizes an epitope comprising position -13910 (wherein the nucleotide is C) or position -22018 (wherein the nucleotide is G). It does not or essentially does not cross-react with an epitope comprising position -13910 with a T in this position nor with the epitope comprising position -22018 with a G in this position. Specificity of an antibody which may be generated according to standard protocols, may be tested by contacting with DNA molecules carrying the wild-type and the mutant sequence such as in an ELISA assay. Only those

antibodies will be selected that produce a signal over background with the mutant sequence but not with the wild-type sequence.

[0050] The antibody of the invention may be a monoclonal antibody or an antibody derived from or comprised in a polyclonal antiserum. The term "antibody", as used in accordance with the present invention, further comprises fragments of said antibody such as Fab, F(ab')₂, Fv or scFv fragments; see, for example, Harlow and Lane⁵³, loc. cit. The antibody or the fragment thereof may be of natural origin or may be (semi) synthetically produced. Such synthetic products also comprise non-proteinaceous as semi-proteinaceous material that has the same or essentially the same binding specificity as the antibody of the invention. Such products may, for example, be obtained by peptidomimetics.

[0051] The term "aptamer" is well known in the art and defined, e.g., in Osborne et al., *Curr. Opin. Chem. Biol.* 1 (1997), 5-9 (see reference 51) or in Stall and Szoka, *Pharm. Res.* 12 (1995), 465-483 (see reference 52).

[0052] Moreover, the invention relates to an antibody or aptamer or phage that specifically binds to the wild-type nucleic acid molecule as described herein above but not to the corresponding mutant sequence contributing to or indicative of adult-type hypolactasia. The statements with respect to specificity etc. made for the antibody which is specific for the mutant sequence apply mutatis mutandis here.

[0053] Furthermore, the invention relates to a pharmaceutical composition comprising the wild-type nucleic acid molecule as described herein above.

[0054] The pharmaceutical composition of the invention may be used in gene therapy approaches, particularly in somatic gene therapy.

[0055] The wild-type nucleic acid molecule referred to above and contained in the pharmaceutical composition of the invention may be combined with a pharmaceutically acceptable carrier and/or diluent.

[0056] Examples of suitable pharmaceutical carriers are well known in the art and include phosphate buffered saline solutions, water, emulsions, such as oil/water emulsions, various types of wetting agents, sterile solutions etc. Compositions comprising such carriers can be formulated by well known conventional methods. These pharmaceutical compositions can be administered to the subject at a suitable dose. Administration of the suitable compositions may be effected by different ways, e.g., by intravenous, intraperitoneal, subcutaneous, intramuscular, topical, intradermal, intranasal or intrabronchial administration. The dosage regimen will be determined by the attending physician and clinical factors. As is well known in the medical arts, dosages for any one patient depends upon many factors, including the patient's size, body surface area, age, the particular compound to be administered, sex, time and route of administration, general health, and other drugs being administered concurrently. A typical dose can be, for example, in the range of 0.001 to 1000 pg of nucleic acid for expression or for inhibition of expression; however, doses below or above this exemplary range are envisioned, especially considering the aforementioned factors. Dosages will vary but a preferred dosage for intravenous administration of DNA is from approximately 10⁶ to 10¹² copies of the DNA molecule. Progress can be monitored by periodic assessment. The compositions of the invention may be administered locally or systemically. Administration will generally be parenterally, e.g., intravenously; DNA may also be administered directly to the target site, e.g., by biolistic

delivery to an internal or external target site or by catheter to a site in an artery. Preparations for parenteral administration include sterile aqueous or non-aqueous solutions, suspensions, and emulsions. Examples of non-aqueous solvents are propylene glycol, polyethylene glycol, vegetable oils such as olive oil, and injectable organic esters such as ethyl oleate. Aqueous carriers include water, alcoholic/aqueous solutions, emulsions or suspensions, including saline and buffered media. Parenteral vehicles include sodium chloride solution, Ringer's dextrose, dextrose and sodium chloride, lactated Ringer's, or fixed oils. Intravenous vehicles include fluid and nutrient replenishers, electrolyte replenishers (such as those based on Ringer's dextrose), and the like. Preservatives and other additives may also be present such as, for example, antimicrobials, anti-oxidants, chelating agents, and inert gases and the like.

[0057] Additionally, the invention relates to a diagnostic composition comprising the nucleic acid molecule as described herein above, the vector as described herein above, the primer or primer pair as described herein above, and/or the antibody aptamer and/or phage as described herein above.

[0058] The diagnostic composition is useful for assessing the genetic status of a person with respect to his or her predisposition to develop adult-type hypolactasia or with regard to the diagnosis of the acute condition. The various possible components of the diagnostic composition may be packaged in one or more vials, in a solvent or otherwise such as in lyophilized form. If dissolved in a solvent, the diagnostic composition is preferably cooled to at least +8° C. to +4° C. Freezing may be preferred in other instances.

[0059] The invention also relates to a method for testing for the presence or predisposition of adult-type hypolactasia or associated trait comprising testing a sample obtained from a prospective patient or from a person suspected of carrying such a predisposition to the presence of the nucleic acid molecule as described herein above in a homozygous or heterozygous state. In varying embodiments, it may be tested either for the presence of the wild-type sequence(s) or of the mutant sequence(s).

[0060] The method of the invention is useful for detecting the genetic set-up of said person/patient and drawing appropriate conclusions whether a condition from which said patient suffers is adult-type hypolactasia. Alternatively, it may be assessed whether a person not suffering from a condition carries a predisposition to adult-type hypolactasia. With regard to position -13910 upstream of the LPH gene, only if cytosine is found in a homozygous state, a condition would be diagnosed as adult-type hypolactasia or a corresponding predisposition would be manifest. On the other hand, if thymidine is found in a homozygous state or if the individual is heterozygous (C/T), then it may be concluded that a condition from which a patient suffers is not related to adult-type hypolactasia and further, that the patient does not carry a predisposition to develop this condition. It may, however, be concluded that children of persons carrying the heterozygous genotype may develop the condition if chromosome carrying the C residue is matched with a corresponding chromosome from the other parent.

[0061] The situation is similar and essentially the same conclusions apply for the analysis of the SNP in position -22018. A homozygously occurring G residue marks a predisposition to or the occurrence of acute adult-type hypolactasia. A heterozygous G/A state correlates with a high likelihood to not develop the condition. Individuals carrying A in a

homozygous state would not be expected to develop the condition. Similarly, patients suffering from a condition would be diagnosed not to suffer from adult-type hypolactasia.

[0062] In a preferred embodiment of the method of the invention said testing comprises hybridizing the complementary nucleic acid molecule as described herein above which is complementary to the nucleic acid molecule contributing to or indicative of adult-type hypolactasia or the nucleic acid molecule as described herein above which is complementary to the wild-type sequence as a probe under (highly) stringent conditions to nucleic acid molecules comprised in said sample and detecting said hybridization.

[0063] Again, depending on the nucleic acid probe used, either wild-type or mutant sequences (i.e. sequences contributing to or indicative of adult-type hypolactasia) would be detected. It is understood that hybridization conditions would be chosen such that a nucleic acid molecule complementary to wild-type sequences would not or essentially not hybridize to the mutant sequence. Similarly, a nucleic acid molecule complementary to the mutant sequence would not or would not essentially not hybridize to the wild-type sequence. In order to differentiate between results obtained from homozygous and heterozygous genotypes in the hybridization methods of the invention, one can for example monitor/detect the strength/intensity of the respective detection signal after the hybridization. To differentiate between wild-type homozygous, heterozygous and/or mutant homozygous alleles in the hybridization methods of the invention, internal control samples of the corresponding genotypes will be included in the analysis.

[0064] In a further preferred embodiment, the method of the invention further comprises digesting the product of said hybridization with a restriction endonuclease or subjecting the product of said hybridization to digestion with a restriction endonuclease and analyzing the product of said digestion.

[0065] This preferred embodiment of the invention allows by convenient means, the differentiation between an effective hybridization and a non-effective hybridization. For example, if the DNA sequence adjacent to position -13910 or position -22018 comprises an endonuclease restriction site, the hybridized product will be cleavable by an appropriate restriction enzyme upon an effective hybridization whereas a lack of hybridization will yield no double-stranded product or will not comprise the recognizable restriction site and, accordingly, will not be cleaved. In particular, the restriction enzymes specific for the sequence of the DNA-variant C/T₋₁₃₉₁₀ is CviI I, for the DNA-variant G/A₋₂₂₀₁₈ are HhaI and Aci I. Said restriction enzymes which cut rg/cy were found by the use of the program Webcutter. The analysis of the digestion product can be effected by conventional means, such as by gel electrophoresis which may be optionally combined by the staining of the nucleic acid with, for example, ethidium bromide. Combinations with further techniques such as Southern blotting are also envisaged.

[0066] Detection of said hybridization may be effected, for example, by an anti-DNA double-strand antibody or by employing a labeled oligonucleotide. Conveniently, the method of the invention is employed together with blotting techniques such as Southern or Northern blotting and related techniques. Labeling may be effected, for example, by standard protocols and includes labeling with radioactive markers, fluorescent, phosphorescent, chemiluminescent, enzymatic labels, etc. (see also above).

[0067] In accordance with the above, in another preferred embodiment of the method of the invention said probe is detectably labeled, e.g. by the methods and with the labels described herein above.

[0068] In yet another preferred embodiment of the method of the invention said testing comprises determining the nucleic acid sequence of at least a portion of the nucleic acid molecule as described herein above, said portion comprising nucleotide position -13910 and/or nucleotide position -22018 of the LPH gene.

[0069] Determination of the nucleic acid molecule may be effected in accordance with one of the conventional protocols such as the Sanger or Maxam/Gilbert protocols (see Sambrook et al., loc. cit., for further guidance).

[0070] In a further preferred embodiment of the method of the invention the determination of the nucleic acid sequence is effected by solid-phase minisequencing. Solid-phase minisequencing is based on quantitative analysis of the wild type and mutant nucleotide in a solution. First, the genomic region containing the mutation is amplified by PCR with one biotinylated and non-biotinylated primer where the biotinylated primer is attached to a streptavidin (SA) coated plate. The PCR-product is denatured to a single stranded form to allow a minisequencing primer to bind to this strand just before the site of the mutation. The tritium (H3) or fluorescence labeled mutated and wild type nucleotides together with nonlabeled dNTPs are added to the minisequencing reaction and sequenced using Taq-polymerase. The result is based on the amount of wild type and mutant nucleotides in the reaction measured by beta counter or fluorometer and expressed as an R-ratio. See also Syvänen AC, Sajantila A, Lukka M. *Am J Hum Genet* 1993; 52,46-59 and Suomalainen A and Syvanen A C. *Methods Mol Biol* 1996;65:73-79.

[0071] A preferred embodiment of the method of the invention further comprises, prior to determining said nucleic acid sequence, amplification of at least said portion of said nucleic acid molecule.

[0072] Preferably, amplification is effected by polymerase chain reaction (PCR). Other amplification methods such as ligase chain reaction may also be employed.

[0073] In a preferred embodiment of the method of the invention said testing comprises carrying out an amplification reaction wherein at least one of the primers employed in said amplification reaction is the primer as described herein above or belongs to the primer pair as described herein above, comprising assaying for an amplification product. In this embodiment and depending on the information the investigator/physician wishes to obtain, primers hybridizing either to the wild-type or mutant sequences may be employed.

[0074] The method of the invention will result in an amplification of only the target sequence, if said target sequence carries a sequence exactly complementary to the primer used for hybridization. This is because the oligonucleotide primer will under preferably (highly) stringent hybridization conditions not hybridize to the wild-type/mutant sequence—depending which type of primer is used—(with the consequence that no amplification product is obtained) but only to the exactly matching sequence. Naturally, combinations of primer pairs hybridizing to both SNPs may be used. In this case, the analysis of the amplification products expected (which may be no, one, two, three or four amplification product(s) if the second, non-differentiating primer is the same for each locus) will provide information on the genetic status of both positions -13910 and -22018.

[0075] In a preferred embodiment of the method of the invention said amplification is effected by or said amplification is the polymerase chain reaction (PCR).

[0076] The PCR is well established in the art. Typical conditions to be used in accordance with the present invention include for example a total of 35 cycles in a total of 50 μ l volume exemplified with a denaturation step at 93° C. for 3 minutes; an annealing step at 55° C. for 30 seconds; an extension step at 72° C. for 75 seconds and a final extension step at 72° C. for 10 minutes.

[0077] The invention furthermore relates to a method for testing for the presence or predisposition of adult-type hypolactasia comprising assaying a sample obtained from a human for specific binding to the antibody or aptamer or phage as described herein above. In this context a weaker staining for the presence of the antigen of the invention compared to homozygous wild type control samples (comprising two persistent alleles) is indicative for the heterozygous wild type (one persistent allele and one hypolactasic allele, whereas for the homozygous hypolactasic individual no staining is expected if the appropriate antibody is used. Preferably, the method of the invention is performed in the presence of control samples corresponding to all three possible allelic combinations as internal controls. Testing may be carried out with an antibody etc. specific for the wild-type or specific for the mutant sequence. Testing for binding may, again, involve the employment of standard techniques such as ELISAs; see, for example, Harlow and Lane⁵³, loc. cit.

[0078] In a preferred embodiment of the method of the invention said antibody or aptamer or phage is detectably labeled.

[0079] Whereas the aptamers are preferably radioactively labeled with ³H or ³²P or with a fluorescent marker as described above, the phage or antibody may either be labeled in a corresponding manner (with ¹²⁵I as the preferred radioactive label) or be labeled with a tag such as His-tag, FLAG-tag or myc-tag.

[0080] In a further preferred embodiment of the method of the invention the test is an immuno-assay.

[0081] In another preferred embodiment of the method of the invention said sample is blood, serum, plasma, fetal tissue, saliva, urine, mucosal tissue, mucus, vaginal tissue, fetal tissue obtained from the vagina, skin, hair, hair follicle or another human tissue.

[0082] In an additional preferred embodiment of the method of the invention said nucleic acid molecule from said sample is fixed to a solid support.

[0083] Fixation of the nucleic acid molecule to a solid support will allow an easy handling of the test assay and furthermore, at least some solid supports such as chips, silica wafers or microtiter plates allow for the simultaneous analysis of larger numbers of samples. Ideally, the solid support allows for an automated testing employing, for example, robotics devices.

[0084] In a particularly preferred embodiment of the method of the invention said solid support is a chip, a silica wafer, a bead or a microtiter plate.

[0085] Furthermore, the invention relates to the use of the nucleic acid molecule as described herein above for the analysis of the presence or predisposition of adult-type hypolactasia.

[0086] The nucleic acid molecule simultaneously allows for the analysis of the absence of the condition or the predisposition to the condition, as has been described in detail herein above.

[0087] In addition, the invention relates to a kit comprising the nucleic acid molecule as described herein above, the primer or primer pair as described herein above, the vector as described herein above, and/or the antibody aptamer and/or phage as described herein above in one or more containers.

[0088] The invention as well relates to the use of the nucleic acid molecule as described herein above or the vector as described herein above in gene therapy.

[0089] Gene therapy approaches have been discussed herein above in connection with the vector of the invention and equally apply here. It is of note that in accordance with this invention, also fragments of the nucleic acid molecules as defined herein above and as, in particular, depicted in SEQ ID NOs: 3 to 4 may be employed in gene therapy approaches. Said fragments comprise the nucleotide at position -13910 as defined in (c) herein above (and also shown in SEQ ID NO: 3) or position -22018 as defined in (d) herein above (and as shown in SEQ ID NO: 4). Preferably, said fragments comprise at least 200, at least 250, at least 300, at least 400 and most preferably at least 500 nucleotides.

[0090] In a preferred embodiment of the use of the invention said gene therapy treats or prevents adult-type hypolactasia.

BRIEF DESCRIPTION OF THE DRAWINGS

[0091] The figures show:

[0092] FIG. 1: The Finnish adult-type hypolactasia families studied. Blackened symbols indicate hypolactasic individuals, asterisk (*) indicate that no sample was available, question mark (?) indicates unknown affection status. ↑ indicates the individuals used for sequencing for SNP identification (Table 2).

[0093] FIG. 2: Physical map of adult-type hypolactasia locus. BAC clones are shown above the horizontal line. The three genes LPH, MCM6 and DARS are shown by thick black arrows with the tip pointed toward the 3' end of the gene above the black boxes. The position of ten polymorphic microsatellite markers used for fine mapping of the locus are shown. The backslash in the horizontal line denotes a gap in the sequence of the contig sequence. The position of marker D2S2169 was confirmed by bridging the gap with PAC 106020 isolated from the PAC library as described before⁴⁰. The organisation of the MCM6 gene is shown including the position of the lactase persistent phenotype-associated variants in introns 9 and 13 located 13.9 kb and 22 kb 5' of the first ATG of LPH.

[0094] FIG. 3: Extended haplotype analysis of the persistent chromosomes derived from Finnish adult-type hypolactasia families using seven closely linked microsatellite markers. The haplotypes representing the ancestral founder persistent chromosome are shaded. Only the haplotypes of non-persistent chromosomes that were also present in the persistent chromosomes are shown. On the basis of ancestral recombinations, the adult-type hypolactasia locus could be restricted to 47 kb interval between markers LPH1 and AC3.

[0095] FIG. 4: The sequence comprised in the sequence of intron 13 of the MCM6 gene (3220 bp) comprising the SNP at position -13910 in which the T, which is specific for the

lactase persistence, is substituted by a C. Said position is indicated by the use of a small letter. This sequence refers to SEQ ID NO:1.

[0096] FIG. 5: The sequence comprised in the sequence of intron 9 of the MCM6 gene (1295 bp) comprising the SNP at position -22018 in which the A, which is specific for the lactase persisting-type sequence is substituted by a G. Said position is indicated by the use of a small letter. This sequence refers to SEQ ID NO:2.

[0097] FIG. 6: The sequence of the lactase persisting-type intron 13 of the MCM6 gene (3220 bp) comprising at position -13910 a T. Said position is indicated by the use of a small letter. This sequence refers to SEQ ID NO:3.

[0098] FIG. 7: The sequence of the lactase persisting-type intron 9 of the MCM6 gene (1295 bp) comprising at position -22018 an A. Said position is indicated by the use of a small letter. This sequence refers to SEQ ID NO:4.

[0099] FIG. 8: The sequence of intron 13 of the MCM6 gene (3220 bp) comprising the SNP at position -13910 in which the T, which is specific for the lactase persisting-type sequence is substituted by a C. Said position is indicated by the use of a small letter. This sequence refers to SEQ ID NO:5.

[0100] FIG. 9: The sequence of intron 9 of the MCM6 gene (1295 bp) comprising the SNP at position -22018 in which the A, which is specific for the lactase persisting-type sequence is substituted by a G. Said position is indicated by the use of a small letter. This sequence refers to SEQ ID NO:6.

[0101] The examples illustrate the invention.

EXAMPLE 1

Linkage and Linkage Disequilibrium Analysis

[0102] Seven polymorphic microsatellite markers between D2S114 and D2S2385 flanking the LPH gene on 2q21 were analyzed in nine extended Finnish hypolactasia families (FIG. 1). Significant evidence for linkage was found with markers D2S314, D2S442, D2S2196 and D2S1334, with a maximum lod score of 7.67 at $\theta=0$ obtained with marker D2S2196 (Table 1). Obligatory recombination events were detected with marker D2S114 (family B, IV3), which defines the centromeric boundary for the lactase persistence/non-persistence locus, and with marker D2S2385 (family B, IV17) (FIG. 1, Table 1), which defines the telomeric boundary of the locus. To fine map the critical region, nine additional polymorphic markers were analyzed (Table 1). Linkage disequilibrium (LD) over the region was monitored conditional on the detected linkage treating the allele frequencies and the recombination fraction as nuisance parameters¹⁶⁻¹⁷. Six out of nine markers (LPH13, LPH2, LPH1, AC3, AC4, and AC10), spanning over ~200 kb interval showed highly significant evidence of LD ($p<10^{-4}$) whereas markers 3' from the LPH gene showed no evidence of LD (Table 1). Two markers, LPH2 and AC3, displayed the most significant linkage disequilibrium in the lactase persistence alleles ($p<10^{-7}$).

[0103] The family material consisted of nine extended Finnish pedigrees originally studied by Sahi⁵. All family material was tested for adult-type hypolactasia in the 1970s. The family material for this study was enlarged by collecting the DNA of the family members in the younger generations. The family material in this study consisted of 194 individuals in total (FIG. 1). The phenotypic status of all family members was confirmed by lactose tolerance tests with ethanol (LTTE)⁴⁻⁵ in all but 49 individuals. Gluten enteropathy has been excluded in all affected patients by measurement of the serum

IgA anti-tissue transglutaminase⁴⁵. DNA was extracted from blood samples taken from all participating family members in accordance with standard protocols⁴⁶, after obtaining informed consent. As a case-control study 196 random DNA samples isolated from jejunal biopsy specimens from which disaccharidase activities had been measured⁴⁷ at the Helsinki University Hospital were sequenced. DNA was isolated from intestinal biopsies according to the standard protocol⁴⁶. These series comprised 137 lactase persistent and 59 non-persistent samples. In addition DNA from nine Italian, kindly provided by M. Rossi, University of Naples, nine German DNA samples, kindly provided by M. Lentze, University of Bonn and twenty two South Korean, kindly provided by J. K. Seo, Seoul National University, intestinal biopsy sample specimens were analyzed (In the table: 23 Korean, 9 Italian and 7 Germans (One of the cases from Germany originated from South Korea)). The diagnosis was based on the measurement of disaccharidase activities. Finally, to determine the frequency of the C/T₋₁₃₉₁₀ variant in the Finnish population, the DNA of 938 anonymous Finnish blood donors from small parishes from Eastern and Western Finland and the DNA of 109 parents belonging to the CEPH families¹⁹ were analyzed. In addition, genomic DNA from a baboon (*Papio hemedryas* ussinus) isolated from liver biopsy using standard protocols⁴⁸ was analyzed. The study was approved by the Ethical Committees of the Helsinki University Hospital and the Finnish Red Cross Blood Transfusion Service.

EXAMPLE 2

Extended Haplotype Analysis

[0104] In the first stage ten highly polymorphic microsatellite markers flanking the LPH gene on 2q21 were analyzed as described elsewhere^{40,55}. Briefly, the ten highly polymorphic microsatellite markers on 2q in the vicinity of the lactase gene from The Genethon Resource Center⁵⁵ were analyzed with genetic distances as follows: cen—D2S114—1 cM—D2S1334—0 cM—D2S2196—0 cM—D2S442—2 cM—D2S314—2 cM—D2S2385—1 cM—D2S2288—1 cM—D2S397—1 cM—D2S150—1 cM—D2S132. The order of the markers has been mostly obtained from the physical YAC contig map of chromosome 2 (Chumakov et al. 1995⁵⁶) supplemented with the Génethon map. PCR was performed in a total volume of 15 µl containing 12 ng of template DNA, 5 pmol of primers, 0.2 mM of each nucleotide, 20 mM TrisHCl (pH 8.8), 15 mM (NH₄)₂SO₄, 1.5 mM MgCl₂, 0.1% Tween 20, 0.01% gelatin and 0.25 U Taq polymerase (Dynazyme, Finnzymes). One of the primers was radiolabeled at the 5' end with ³²P-γ-ATP. The reactions were performed in a multiwell microtitre plate for 35 cycles with denaturation at 94° C. for 30 s, annealing at various temperatures depending on the primers for 30 s and extension at 72° C. for 30 s; denaturation was set at 3 min and final extension at 5 min. The amplified fragments were separated on 6% polyacrylamide gel, and autoradiography was performed.

[0105] In the second stage, nine additional microsatellite markers within the contig constructed over the LPH gene were identified from the published genomic sequence of the BACs (NH034L23, NH0318L13, NH0218L22, and RP11-32911) using the Repeat Masker program (<http://ftp.genome.washington.edu/cgi-bin/RepeatMasker>). Primers flanking the repeats were synthesized. PCR conditions were as described elsewhere⁴⁰. The amplified fragments were separated on 6% polyacrylamide gel, and autoradiography was performed.

[0106] Pairwise lod scores were calculated by use of the MLINK option of the LINKAGE program package⁴⁹. Autosomal recessive inheritance for adult-type hypolactasia with complete penetrance, no sex difference in recombination fractions, and a disease allele frequency of 0.4 was assumed. Only individuals above 20 years of age were included in the study as the condition is manifested by that age in the Finnish population⁵⁻⁶. The affection status for individuals not confirmed by LTTE was regarded as unknown. Allele frequencies and heterozygosities for the markers were estimated from family material using the Downfreq program for purposes of the parametric linkage analysis⁴⁹. Additionally, pseudomarker linkage and linkage disequilibrium analyses were performed, assuming autosomal recessive mode of inheritance¹⁶. A test of LD was performed conditional on the detected linkage treating the allele frequencies and the recombination fraction as nuisance parameters^{16,49}. P-values from these analyses are shown in Table 1. Haplotypes were constructed manually for the microsatellite markers in this order: LPH1-LPH2-LPH13-AC7-AC3-AC4-AC5 (FIG. 3). A total of 54 non-persistent chromosomes and 33 persistent chromosomes in our family material were available for haplotype analysis.

[0107] The order of the closely linked markers was confirmed by assembling four BAC-clones NH0034L23, NH0218L22, NH0318L13 and 329110 in the critical region into one uninterrupted sequence segment. This contig extended from marker AC8 to the exon 10 of the aspartyl-tRNA synthetase (DARS) gene and covered a total of 222.5 kb (FIG. 2). Based on this physical map of the linked region, extended haplotypes with seven markers covering a 150 kb interval (cen-LPH13-LPH2-LPH1-AC7-AC3-AC4-AC5-tel) (FIG. 3) were constructed. One major haplotype was present in 20 persistence alleles (60%) versus 3 of the non-persistence alleles (5%), whereas a wide diversity of haplotypes was observed in non-persistence alleles. The remaining 40% of the haplotypes in the persistence alleles differed from the ancestral haplotype in a manner consistent with a breakdown of the haplotype by historical recombination events. Based on the conserved haplotype analysis, the locus for lactase persistence could be restricted to a 47 kb interval between markers LPH1 and AC3 (FIG. 3)

EXAMPLE 3

Sequence Analysis of the Adult-Type Hypolactasia Locus

[0108] The 47 kb region between the markers LPH1 and AC3 was amplified in overlapping PCR fragments from genomic DNA of several members of the nine hypolactase families and sequenced. The region contains the minichromosome maintenance (MCM6) gene¹⁸, which covers 36 kb of the critical 47 kb region (FIG. 2). No variations were detected in the coding region of the MCM6 gene but total of 52 variants; 43 SNPs and 9 deletion/insertion polymorphisms, were identified in the critical 47 kb region (Table 2). Only two of the variants (C/T₋₁₃₉₁₀, G/A₋₂₂₀₁₈) were associated with the lactase persistence/non-persistence trait in the Finnish families (Tables 2 and 3). The first associated variant, C/T₋₁₃₉₁₀, resides in intron 13 of the MCM6 gene at position -13910 by from the first ATG-codon of the LPH gene. The second associated variant, G/A₋₂₂₀₁₈, is located in intron 9 of the MCM6 gene at position -22018 from the first ATG-codon of the LPH gene (FIG. 2). These two variants, 8 kb apart from

each other, completely cosegregated with adult-type hypolactasia in nine extended Finnish families. All hypolactasic (non-persistent) family members were homozygous for both C₋₁₃₉₁₀ and G₋₂₂₀₁₈ (Table 3). Interestingly, both these variants reside in repeat elements, C/T₋₁₃₉₁₀ in an L2-derived element and G/A₋₂₂₀₁₈ in an Alu element.

[0109] Experimentally, three non-persistence, 2 homozygous persistence and 2 heterozygous persistence individuals sharing a similar haplotype across the critical region from our family material were used for sequencing in the first stage (FIG. 1). Using the published draft genomic sequence of the BACs: NH0034L23, NH0218L22, NH0318L23, and RP-329110 that covered the critical region of adult-type hypolactasia were assembled to one contig using Sequencher 4 software (Gene Codes Corporation). Oligonucleotide primers spanning the critical region between markers LPH1 and AC3 were designed (a list of oligonucleotide primers described herein below). PCR amplifications were carried out in a 50 pl volume with genomic DNA (100 ng), primers (20 ng each), dNTPs (200 μM), 0.5 U of Taq polymerase (Dynazyme, Finnzymes) in a standard buffer. Most PCR were amplified using the following PCR cycle conditions: an initial round of denaturation at 94° C. for 3 min, then 35 cycles at 94° C. at 30 s, 55° C. for 30 s, and 72° C. for 1.25 min and a final extension of 72° C. for 10 min, except that in cases where the size of the PCR products were more than 1 kb we used the Dynazyme extend kit (conditions are described herein below). Purified PCR products (15-40 ng) were cycle sequenced using BigDye terminator chemistry (PE Biosystems). Data were analyzed using ABI Sequencing Analysis 3.3 (PE Biosystems) and Sequencher 4.1 (Gene Codes).

Detection of the Lactase Variants by Sequencing:

[0110] PCR amplifications were carried out in a 50 pl volume with genomic DNA (100 ng), primers (20 ng each),

were analyzed using ABI Sequencing Analysis 3.3 (PE Biosystems) and Sequencher 4.1 (Gene Codes).

Screening of the Lactase Variants by Solid-Phase Minisequencing:

[0111] The DNA fragment spanning the C/T₋₁₃₉₁₀ variant was amplified using one biotinylated (5'-Bio-CCTCGT-TAATACCCACTGACCTA-3'; SEQ ID NO:9) primer and unbiotinylated (5'-GTCACCTTTGATATGATGAGAGCA-3'; SEQ ID NO:8) primer. For G/A₋₂₂₀₁₈ biotinylated (5'-Bio-TGCTCAGGACATGCTGATCAA-3'; SEQ ID NO:13) and one unbiotinylated 5'-CTACCCTATCAGTAAAGGCCTA-3'; SEQ ID NO:12) primer were used under conditions described above. 10 μl of the PCR product was captured in a streptavidin coated microtiter well (Lab systems, Finland). The wells were washed, and bound DNA was denaturated as described by Syvänen et al. (Am J Hum Genet. (1993), 52, 46-59) and Syvänen and Landegren (Hum Mutat. (1994), 3, 172-9). 50 μl of the minisequencing reaction mixture contained 10 pmoles of the minisequencing primers for C/T₋₁₃₉₁₀ (5'-GGCAATACAGATAAGATAATGTAG-3'; SEQ ID NO:10), G/A₋₂₂₀₁₈ (5'-AAAAACAGCATTCT-CAGCTGGGC-3'; SEQ ID NO:14), AND 0.1 μL of either H-dCTP, H-dGTP corresponding to the lactase non-persistence allele (115 Ci/mmol; Amersham, UK) or H-dTTP, H-sATP corresponding to the lactase persistence allele and 0.05 units of DNA polymerase (Dynazyme II, Finnzymes) in its buffer was added to each well. The microtiter plates were incubated for 20 min at 50° C., and the wells were washed. The detection was eluted and the eluted radioactivity was measured in a liquid scintillation counter (Rackbeta 1209, Wallac, Finland). Two parallel minisequencing reactions were carried out for each PCR product.

PCR primers and detection primer for the C/T ₋₁₃₉₁₀ variant:			
Forward PCR primer:	GTCACCTTTGATATGATGAGAGCA	Tm 58	SEQ ID NO: 8
Detection primer:	GGCAATACAGATAAGATAATGTAG	Tm 58	SEQ ID NO: 10
Bio-Reverse primer:	Bio-CCTCGTTAATACCCACTGACCTA	Tm 62	SEQ ID NO: 9
or			
	Bio-TAGGTCACTGGGTATTAACGAGGT		SEQ ID NO: 7
PCR primers and detection primer for the G/A ₋₂₂₀₁₈ variant:			
Forward PCR primer:	CTACCCTATCAGTAAAGGCCTA	Tm 58	SEQ ID NO: 12
Detection primer:	AAAAACAGCATTCTCAGCTGGGC	Tm 62	SEQ ID NO: 14
Bio-Reverse primer:	Bio-TGCTCAGGACATGCTGATCAA	Tm 62	SEQ ID NO: 13
or			
	Bio-TTGATCAGCATGTCCTGAGCA		SEQ ID NO: 11

dNTPs (200 μM), 0.5 U of Taq polymerase (Dynazyme, Finnzymes) in a standard buffer. Both PCRs were amplified using the following PCR cycle conditions: an initial round of denaturation at 94° C. for 3 min, then 35 cycles at 94° C. at 30 s, 55° C. for 30 s, and 72° C. for 1.25 min and a final extension of 72° C. for 10 min. PCR were purified by enzymatic reaction. Purified PCR products (15-40 ng) were cycle sequenced using BigDye terminator chemistry (PE Biosystems). Data

EXAMPLE 4

Monitoring the DNA-Variants in a Case/Control Study Sample

[0112] The frequency of the C/T₋₁₃₉₁₀ and G/A₋₂₂₀₁₈ variants was analyzed in DNA samples isolated from a total of 196 intestinal biopsy samples specimens which had been

analyzed for disaccharidase activity as a diagnostic test for hypolactasia. A total of 59 samples showed primary lactase deficiency. Six out of 59 cases (Table 3) were heterozygous GA for the G/A₋₂₂₀₁₈ variant, the remaining 53 being homozygous for the G allele. All 59 samples were homozygous for the C allele of the variant C/T₋₁₃₉₁₀.

[0113] Among the 173 cases showing lactase persistence, 74 were found to be homozygous for alleles T and A, 63 being heterozygous CT and GA and none being homozygous for alleles C and G at C/T₋₁₃₉₁₀ and G/A₋₂₂₀₁₈, respectively (Table 3).

[0114] To analyze these variants in other populations, DNA samples isolated from intestinal biopsy specimens from 40 non-Finnish cases with established disaccharidase deficiency were sequenced: 23 cases originated from South Korea, 9 from Italy and 8 from Germany. One Italian case was heterozygous GA for G/A₋₂₂₀₁₈ whereas all remaining 39 cases were homozygous CC and GG for C/T₋₁₃₉₁₀ variant and G/A₋₂₂₀₁₈ respectively (Table 3). An extended study gave rise to the data provided in Table 7 representing data of the complete association of C/T₋₁₃₉₁₀ variant with the biochemically verified hypolactasia (lactase non-persistence) in 400 individuals for 6 different populations. The G/A₋₂₂₀₁₈ variant was associated with the lactase non-persistence in 400 out of 401 cases.

EXAMPLE 5

Molecular Epidemiology of the Lactase Persistence Variant C/T₋₁₃₉₁₀

[0115] To monitor for the prevalence of the hypolactasia-associated variant in the Finnish population a solid-phase minisequencing method^{19,20} was used to screen DNA samples of 938 anonymous Finnish blood donors originating either from the Western early settlement region or the Eastern late settlement region of Finland (Table 4). Experimentally, the DNA fragment spanning the C/T₋₁₃₉₁₀ variant was amplified using one biotinylated (5'-CCTCGTTAATACCCCT-GACCTA-3'; SEQ ID NO:9) primer and unbiotinylated (5'-GTCACCTTGATATGATGAGAGCA-3'; SEQ ID NO:8) primer. For G/A₋₂₂₀₁₈ we used one biotinylated (5'-AGTCT-GTGGCATGTGTCTTCATG-3'; SEQ ID NO:15) and one unbiotinylated (5'-TGCTCAGGACATGCTGATCAACT-3'; SEQ ID NO:16) primer under conditions described above. 10 µl of the PCR product was captured in a streptavidin coated microtitre well (Lab system, Finland). The wells were washed, and the bound DNA was denatured as described previously^{19,20}, 50 µl of the minisequencing reaction mixture contain 10 pmoles of the minisequencing primers for G/A₋₂₂₀₀₅ (5'-GACAAAGGTGTGAGCCACCG-3'; SEQ ID NO:17), G/A₋₁₃₉₁₅ (5'-GGCAATACAGATAAGATAATG-TAG-3'; SEQ ID NO:10) and 0.1 µl of either-dCTP corresponding to the lactase non-persistence allele (115 Ci/mmol; Amersham, UK) or H-dTTP corresponding to the lactase persistence allele and 0.05 units of DNA polymerase (Dyna-zyme II, Finnzymes) in its buffer was added to each well. The microtiter plates were incubated for 20 min at 50° C., and the wells were washed. The detection primer was eluted, and the eluted radioactivity was measured in a liquid scintillation counter (Rackbeta 1209, Wallac, Finland). Two parallel minisequencing reactions were carried out for each PCR product. The overall prevalence of the putative hypolactasia genotype CC₋₁₃₉₁₀ (170 cases) was 18.1%, with higher prevalence (16.8% versus 18.9%) in the western than in the

eastern sample (Table 4). These values are in good agreement with the epidemiological study reporting the prevalence of 17% among Finnish speaking Finns with an increasing gradient from West to East². The same set of samples for the G/A₋₂₂₀₁₈ polymorphism was also genotyped, and the LD between these two SNPs monitored using the D' statistic²¹. They were found to be in almost complete LD (D'=0.98, p=7.62×10⁻¹¹, Table 5).

[0116] The prevalence of hypolactasia in different populations is known to vary greatly from less than 5% to almost 100%^{3,5}. To determine whether these changes in hypolactasia prevalence would correlate with the distribution of the genotype CC₋₁₃₉₁₀, the DNA of the parents of CEPH families²² was analyzed. CEPH families have been mainly collected from France, with reported prevalence of hypolactasia around 37%²³ and Utah, the Utah populations originating from Northern Europe with prevalence of hypolactasia less than 5%²⁴. Genotyping of the parents in CEPH families revealed that 41.2% (7 out of 7 samples) of French families have the genotype CC whereas only 7.6% (7 out of 92 samples of Utah families have the genotype CC (Table 4). Again, despite the small number of analyzed samples these figures agree with the values obtained in the epidemiological studies of hypolactasia in these populations^{23,24}.

[0117] Table 8 demonstrates that the observed prevalence of the variants well agrees with the described population frequencies of the lactose intolerance.

EXAMPLE 6

The Genealogy of the Lactase Persistence Variant C/T₋₁₃₉₁₀

[0118] Haplotype analysis in the Finnish families suggested that most if not all, lactase persistence alleles in Finland have descended from one common ancestor. Linkage disequilibrium was used to estimate the time of the introduction of the persistence allele into the Finnish population²⁵. Assuming 20 years generation time, this estimate would indicate that the founder mutation was introduced into the Finnish population some 9000-11400 years ago (Table 6). This is in good agreement with earliest signs of settlement in the Finnish mainland some 8000-9000 years ago²⁶ and would reasonably well coincide with the beginning of the dairy farming in 8000-10.000 BC²⁷. More importantly, the presence of the same DNA-variant in persistence alleles in different populations would suggest that this variant is even more ancient and the mutation has occurred before differentiation of the analyzed populations.

[0119] To get some insight into the phylogenetic origin of the lactase allele, intron 9 and part of intron 13 of the MCM6 gene of a Baboon (*Papio Hamadryas*) were sequenced. Genotype GG and CC was present in Baboons DNA at both G/A₋₂₂₀₁₈ and C/T₋₁₃₉₁₀. This could suggest that alleles G and C, respectively reflect the appearance of the ancestral allele, presenting the non-persistence type and a mutation has transformed this allele to create the persistence allele. This assumption is supported by the identification of the LD and shared haplotype in the persistence alleles versus a high diversity of alleles found in non-persistence alleles.

EXAMPLE 7

Pairwise LD of C/T and G/A Variants

[0120] Pairwise LD between C/T₋₁₃₉₁₀ and G/A₋₂₂₀₁₈ was estimated using the D' statistic²¹. Haplotype frequencies were

estimated by maximum likelihood using the EH program⁵⁰. D' is calculated as $\max(D/D_{\max}, D/D_{\min})$; where disequilibrium measure $D = h_{pq} - p \cdot q$, where h_{pq} is the frequency of the haplotype with rare allele at each locus, p and q are frequency of the rare alleles at loci 1 and 2, and $D_{\max} = \min p(1-p), q(1-q)$ if $D > 0$, and $D_{\min} = -\min pq, (1-p)(1-q)$ if $D < 0$. The significance of deviation of D' from 0 was determined using the statistic

$$D^2 \sqrt{\frac{N}{p(1-p)q(1-q)}}$$

which is distributed as χ^2 with 1 df²¹

[0121] Gene accessions numbers. For BACs NH0218L22, N0034L34, NH0318L13, and RP11-329I10 are AC012551, AC011893, AC011999 and AC016516 respectively. The accession numbers for human polymorphisms are GenBank AF395607-AF395615.

REFERENCES

- [0122] 1. Flatz, G. & Rothauwe, H. The human lactase polymorphism: physiology and genetics of lactose absorption and malabsorption. *Prog. Med. Genet.* 2, 205-249 (1977).
- [0123] 2. Sahi, T., Isokoski, M., Jussila, J. & Launiala, K. Lactose malabsorption in Finnish children of school age. *Acta Paediatr Scand.* 61, 11-16 (1972).
- [0124] 3. Wang, Y. et al. The genetically programmed down-regulation of lactase in children. *Gastroenterology*. 114:1230-1236 (1998).
- [0125] 4. Sahi, T., Isokoski, M., Jussila, J., Launiala, K. & Pyorala, K. Recessive inheritance of adult-type lactose malabsorption. *Lancet*. 823-826 (1973).
- [0126] 5. Sahi, T. The inheritance of selective adult-type lactose malabsorption. *Scand. J. Gastroenterol. suppl.* 30, 1-73 (1974).
- [0127] 6. Sahi, T. Genetics and epidemiology of adult-type hypolactasia. *Scand. J. Gastroenterol. Suppl.* 202, 7-20 (1994).
- [0128] 7. Boll, W., Wagner, P. & Mantei, N. Structure of the chromosomal gene and cDNAs coding for lactase-phlorizin hydrolase in human with adult-type hypolactasia or persistence of lactase. *Am. J. Hum. Genet.* 48, 889-902 (1991).
- [0129] 8. Mantei, N. et al. Complete primary structure of human and rabbit lactase-phlorizin hydrolase: implications for biosynthesis, membrane anchoring and evolution of the enzyme. *EMBO J.* 7, 2705-2713 (1988).
- [0130] 9. Wang, Y. et al. The lactase persistence/non-persistence polymorphism is controlled by a cis-acting element. *Hum. Mol. Genet.* 4, 657-662 (1995).
- [0131] 10. Harvey, C. B., Pratt, W. S., Islam, I., Whitehouse, D. B. & Swallow, D. M. DNA polymorphisms in the lactase gene: linkage disequilibrium across the 70 kb region. *Eur J. Hum. Genet.* 3, 27-41 (1995).
- [0132] 11. Escher, J. C. et al. Molecular basis of lactase levels in adult humans. *J. Clin. Invest.* 89, 480-483 (1992).
- [0133] 12. Lloyd, M. et al. Regulation of intestinal lactase in adult hypolactasia. *J. Clin. Invest.* 89, 524-529 (1992).
- [0134] 13. Fajardo, O., Naim, H. Y. & Lacey, S. W. The polymorphic expression of lactase in adults is regulated at the messenger RNA level. *Gastroenterology* 106, 1233-14.
- [0135] 14. Luigi, M. et al. Mosaic regulation of lactase in human adult-type *Gastroenterology* 112, 1506-1514 (1997).
- [0136] 15. Rossi, M. et al. Lactase persistence versus decline in human adults: Multifactorial events are involved in down-regulation after weaning. *Gastroenterology* 112, 1506-1514 (1997).
- [0137] 16. Göring, H. H. H. & Terwilliger, J. D. Linkage analysis in the presence of errors IV: Joint pseudomarker analysis of linkage and/or linkage disequilibrium on a mixture of pedigrees and singletons when mode of inheritance cannot be accurately specified. *Am. J. Hum. Genet.* 66, 1310-1327 (2000).
- [0138] 17. Terwilliger, J. D. & Göring, H. H. H. Gene mapping in the 20th and 21st centuries: Statistical methods, data analysis, and experimental design. *Hum. Biol.* 72, 63-132 (2000).
- [0139] 18. Harvey, C. B. et al. Regional localization of the lactase-phlorizin hydrolase, LCT, to chromosome 2q21. *Ann. Hum. Genet.* 57, 179-185 (1993).
- [0140] 19. Syvänen, A.-C., Sajantila, A., Lukka, M. Identification of individuals by analysis of biallelic DNA markers, using PCR and solid-phase minisequencing. *Am. J. Hum. Genet.* 52, 46-59 (1993).
- [0141] 20. Syvänen, A.-C. & Landegren, U. Detection of point mutations by solid-phase methods. *Hum. Mutat.* 3, 172-179 (1994).
- [0142] 21. Thompson, E. A., Deeb, S., Walker, D. & Motulsky, A. G. The detection of Linkage disequilibrium between closely linked markers: RFLPs at the AI-CIII Apolipoprotein genes. *Am. J. Hum. Genet.* 42, 113-124 (1998).
- [0143] 22. Dausset, J. et al. Centre d'étude du polymorphisme humain (CEPH): Collaborative genetic mapping of human genome. *Genomics* 6, 575-577 (1990).
- [0144] 23. Cuddenec, Y., Delbrück, H. & Flatz, G. Distribution of the adult lactase phenotypes—lactose absorber and malabsorber—in a group of 131 army recruit *Gastroenterol. Clin. Biol.* 6, 776-779 (1982).
- [0145] 24. McLellan, T., Jorde, L. B. & Skolnick, M. H. Genetic distance between the Utah Mormons and related populations. *Am. J. Hum. Genet.* 36, 836-857 (1984).
- [0146] 25. Terwilliger, J. D. A powerful likelihood method for the analysis of linkage disequilibrium between trait loci and one or more polymorphic marker loci. *Am. J. Hum. Genet.* 56, 777-787 (1995).
- [0147] 26. Nunez, M. G. A model of the early settlement of Finland. *Fennoscandia archaeologica* IV, 3-18 (1997).
- [0148] 27. Simoons, F. J. Primary adult lactose intolerance and the milking habit: a problem in biological and cultural interrelations. II. A cultural historical hypothesis. *Am. J. Dig. Dis.* 16, 695-710 (1970).
- [0149] 28. Varilo, T. et al. The age of human mutation: genealogical and linkage disequilibrium analysis of the CLN5 mutation in the Finnish population. *Am. J. Hum. Genet.* 58, 506-512 (1996).
- [0150] 29. Hästbacka, J. et al. Linkage disequilibrium mapping in isolated founder populations: diastrophic dysplasia in Finland. *Nature Genet.* 2: 204-211 (1992).
- [0151] 30. Harvey C. B. et al. Lactase haplotype frequencies in Caucasians: association with the lactase persistence/non persistence polymorphism. *Ann Hum Genet* 62, 215-223 (1998).
- [0152] 31. Ohtani, K. et al. Cell growth-regulated expression of mammalian MCM5 and MCM6 genes mediated by the transcription factor E2F. *Oncogene* 18, 2299-2309 (1999).

- [0153] 32. Smith, A. F. A. The origin of interspersed repeats in the human genome. *Curr. Opin. Genet. Dev.* 6, 743-748 (1996).
- [0154] 33. Kazazian, H. H. & Moran, J. V. The impact of L1 retrotransposons on the human genome. *Nature Genet.* 19, 19-24 (1998).
- [0155] 34. Moran, J. V., DeBerardinis, R. J. & Kazazian, H. H. Exon shuffling by L1 retrotransposition. *Science* 283, 1530-1534 (1999).
- [0156] 35. Wei, W. et al. Human L1 retrotransposition: cis preference versus trans complementation. *Mol. Cell. Biol.* 21, 1429-1439 (2001).
- [0157] 36. Donnelly, S. R., Hawkins, T. E. & Moss, S. E. A conserved nuclear element with a role in mammalian gene regulation. *Hum. Mol. Genet.* vol. 8, 9, 1723-1728 (1999).
- [0158] 37. Boeke, J. D. LINEs and Alus—the polyA connection. *Nature Genet.* 16, 6-7 (1997).
- [0159] 38. Jurka, J. Sequence patterns indicate an enzymatic involvement in integration of mammalian retrotransposons. *Proc. Natl. Acad. Sci. U.S.A.* 94, 1872-1877 (1997).
- [0160] 39. Savilahti E, Launiala K, Kuitunen P. Congenital lactase deficiency. *Arch. Dis. Child.* 58, 246-252 (1983).
- [0161] 40. Järvelä, I. et al. Assignment of the locus for congenital lactase deficiency to 2q21, in the vicinity of but separate from the lactase-phlorizin hydrolase gene. *Am. J. Hum. Genet.* 63, 1078-1085 (1998).
- [0162] 41. Simoons, F. J. The geographic hypothesis and lactose malabsorption. A weighing of the evidence. *Am. J. Dig. Dis.* 23, 963-980 (1978).
- [0163] 42. Flatz, G. & Rothauwe, H. W. The human lactase polymorphism: physiology and genetics of lactose absorption and malabsorption. *Prog. Med. Genet.* 2, 205-249 (1977).
- [0164] 43. McCracken, R. D. Lactase deficiency: an example of dietary evolution. *Curr. Anthropol.* 12, 479-517 (1971).
- [0165] 44. Arola, H. et al. Diagnosis of hypolactasia and lactose malabsorption. *Scand. J. Gastroenterol. Suppl.* 202, 26-35 (1994).
- [0166] 45. Sulkkanen, S. et al. Tissue transglutaminase autoantibody enzyme-linked immunosorbent assay in detecting celiac disease. *Gastroenterology* 115 (6), 1322-1328 (1998).
- [0167] 46. Sambrook, J., Fritsch, E. F. & Maniatis, T. *Molecular cloning: a laboratory manual*, (2nd ed). Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1989).
- [0168] 47. Messer, M. & Dahlqvist, A. A one-step ultramicro method for the assay of intestinal disaccharidases. *Anal. Biochem.* 14 (3), 376-92 (1966).
- [0169] 48. Cottingham, Jr. R. W., Idury, R. M. & Schaffer, A. A. Faster sequential genetic linkage computations. *Am. J. Hum. Genet.* 53, 252-263 (1993).
- [0170] 49. Goring, H. H. H. & Terwilliger, J. D. Linkage analysis in the presence of errors III:
- [0171] Marker loci and their map as nuisance parameters. *Am. J. Hum. Genet.* 66,1298-1309 (2000).
- [0172] 50. Terwilliger, J. D. & Ott, J. *Hand book of human genetic analysis.* Johns Hopkins University Press, Baltimore (1994).
- [0173] 51. Osborne et al., *Curr. Opin. Chem. Biol.* 1(1997), 5-9
- [0174] 52. Stall and Szoka, *Pharm. Res.* 12 (1995), 465-483
- [0175] 53. Harlow and Lane “Antibodies, A Laboratory Manual”, CSH Press, Cold Spring Harbor, USA, 1988
- [0176] 54. Higgins and Hames (eds.), “Nucleic acid hybridization, a practical approach”, IRL Press, Oxford 1985
- [0177] 55. Dib C, Faure S, Fizames C, Samson D, Drouot N, Vignal A, Millasseau P, Marc
- [0178] S, Hazan J, Seboun E, Lathrop M, Gyapay G, Morissette J, Weissenbach J. A comprehensive genetic map of the human genome based on 5,264 microsatellites. *Nature.* 1996 Mar. 14; 380 (6570):152-4.
- [0179] 56. Chumakov IM, Rigault P, Le Gall I, Bellanne-Chantelot C, Billault A, Guillou S,
- [0180] Soularue P, Guasconi G, Poullier E, Gros I, et al. A YAC contig map of the human genome. *Nature.* 1995 Sep. 28; 377 (6547 Suppl):175-297

TABLE 1

Linkage and Linkage Disequilibrium Analyses in adult-type hypolactasia families (fine mapping markers shown in bold)						
Marker	Lod score(Z) at θ					p-value ^a
	0.0	0.1	0.2	0.3	0.4	
D2S114	—∞	2.44	1.92	1.13	0.41	0.87195
P6112	2.76	2.20	1.45	0.75	0.22	0.66207
D2S1334	3.15	2.45	1.61	0.84	0.25	0.91039
AC8	2.26	1.99	1.36	0.71	0.21	0.53670
LPH13	3.67	2.94	1.96	1.03	0.31	4 × 10 ⁻⁶
LPH2	4.09	3.07	2.00	1.00	0.26	5.7 × 10 ⁻⁷
LPH1	5.91	4.52	2.96	1.53	0.46	5 × 10 ⁻⁶
AC7	3.63	2.60	1.66	0.83	0.23	0.03471
AC3	6.63	4.88	3.16	1.61	0.44	3.2 × 10 ⁻⁸
AC4	3.07	2.22	1.42	0.71	0.19	4 × 10 ⁻⁵
AC5	5.33	4.10	2.72	1.39	0.39	0.02166
AC10	6.60	4.99	3.25	1.65	0.46	1 × 10 ⁻⁵
D2S2196	7.67	5.62	3.62	1.85	0.54	0.00010
D2S442	3.81	3.08	2.08	1.03	0.27	0.22805
D2S314	4.22	3.61	2.50	1.37	0.45	0.27535
D2S2385	—∞	2.79	1.92	1.01	0.28	0.46457

^ap-values produced using linkage disequilibrium test given linkage^{16,49}

TABLE 2

The variations identified within adult-type hypolactasia locus in the Finnish Families								
Position ^a	Variant	Lactase persistence (Homozygous)		Lactase persistence (Heterozygous)		Lactase non-persistence		
		BIV4	AIV3	BIV8	CIV3	BIV9	DIV4	EIII2 ^b
-694	A→G	AA	AA	AG	AA	GG	N ^c	AA
-1640/50	T ₁₃ →T ₁₂	Tu _{13/13}	T _{13/13}	T _{13/13}	T _{13/13}	T _{13/13}	T _{12/12}	T _{12/12}
-2131	C→T	CC	CC	CT	CC	TT	CT*	TT
-3058/72	T ₁₅ →T ₁₆	T _{15/15}	T _{15/15}	T _{15/15}	T _{15/15}	T _{15/15}	T _{16/16}	T _{16/16}
-3075	G→T	GG	GG	GG	GG	GG	GG	TT

TABLE 2-continued

The variations identified within adult-type hypolactasia locus in the Finnish Families								
Position ^a	Variant	Lactase persistence (Homozygous)		Lactase persistence (Heterozygous)		Lactase non-persistence		
		BIV4	AIV3	BIV8	CIV3	BIV9	DIV4	EIII2 ^b
-4480	T→A	TT	TT	TA	TT	AA	TT	TT
-5440	C→T	CC	CC	CT	CC	TT	CC	CC
-5926	A→T	AA	AA	AA	AA	AA	TA	TT
-8540	G→A	GG	GG	GA	GA	AA	AG	AA
-8630	C→G	CC	CC	CG	CG	GG	GC	GG
-13495	T→C	TT	TT	TC	TT	CC	CT	CC
-13910	T→C	TT	TT	TC	TC	CC	CC	CC
-15239	G→A	GG	GG	GA	GG	AA	AG	AA
-15862	T→C	CC	CC	CT	CC	TT	TC	TT
-16568/79	T ₁₁ →T ₁₂	T _{11/11}	T _{11/11}	T _{11/12}	T _{11/11}	T _{12/12}	T _{11/11}	T _{12/12}
-16888	A→G	AA	AA	GA	AA	GG	GA	GG
-17300	C→T	CC	CC	CC	CC	CC	CT	TT
-19044	T→C	TT	TT	TC	TT	CC	CT	CC
-19519	T→C	TT	TT	TC	TT	CC	TT	TT
-20077	C→G	CC	CC	CG	CC	GG	GC	GG
-20486	G→A	GG	GG	GA	GG	AA	GG	GG
-21721/28	A ₇ →A ₆	A _{7/7}	A _{7/7}	A _{7/7}	A _{7/7}	A _{7/7}	A _{7/A6}	A _{7/7}
-21731	A→C	AA	AA	AA	AA	AA	CC	AA
-21736/43	A ₉ →A ₈	A _{9/9}	A _{9/9}	A _{9/A8}	A _{9/9}	A _{8/8}	A _{8/8}	A _{8/8}
-22018	G→A	AA	AA	AG	AG	GG	GG	GG
-22741	C→T	CC	CC	CC	CC	CC	N	TT
-22788	A→G	AA	AA	AG	AA	GG	N	GG
-23069	A→G	AA	AA	AG	AA	GG	N	GG
-23442	A→G	AA	AA	AA	AA	AA	N	GG
-23771	T→C	TT	TT	TT	TT	TT	N	CC
-25093/23	Δ30 bp	ΔΔ	ΔΔ	ΔΔ	ΔΔ	ΔΔ	N	II
-27310	A→G	AA	AA	AG	AA	GG	GA	GG
-27480	G→A	GG	GG	GA	GG	AA	AG	AA
-27807	A→C	AA	AA	AA	AA	AA	AC	CC
-30183	A→G	AA	AA	AG	AA	GG	AA	AA
-31268	A→G	AA	AA	AG	AA	GG	AA	AA
-31342	T→C	TT	TT	TT	TT	TT	CT	CC
-33645	C→T	CC	CC	CT	CC	TT	CC	CC
-35176	T→C	TT	TT	TC	TT	CC	CT	CC
-36254	C→T	CC	CC	CT	CC	TT	TC	TT
-36296	G→T	TT	TT	TG	TT	GG	TG	N
-36501	A→T	AA	AA	AT	AA	TT	AT	N
-36506/14	Δ9 bp	ΔΔ	ΔΔ	ΔI	ΔΔ	II	ΔI	N
-36671/77	T ₇ →T ₆	T _{7/7}	T _{7/7}	T _{7/6}	T _{7/7}	T _{6/6}	T _{7/7}	T _{7/7}
-37565	T→G	TT	TT	TG	TT	GG	GG	TG
-38276	G→C	GG	GG	GC	GG	CC	GG	GG
-39036	G→C	GG	N	GC	N	CC	N	N
-40608	G→C	GG	GG	GG	GG	GG	GC	CC
-41590	T→C	TT	TT	TC	TT	CC	CT	CC
-42081/82	ΔAG	AG	AG	AG/Δ	AG	ΔΔ	AG	AG
-42618	T→C	TT	TT	TC	TT	CC	TT	TT
-42893	G→A	GG	GG	GA	GG	AA	GG	GG

^aThe Number is from initiation translation codon (ATG) of the LPH gene using the compiled genomic sequence of the BACs NH034L23, NH0218L22, NH0318L13 and RP11-329I10,

^bthe individuals sequenced from the Finnish families studied and showed by arrow in FIG. 1,

^cnot determined

TABLE 3

Distribution of C/T ₁₃₉₁₀ & G/A ₂₂₀₁₈ genotypes in lactase persistent/non-persistent alleles								
	Genotype	C/T ₁₃₉₁₀			G/A ₂₂₀₁₈			Total
		CC	CT	TT	GG	GA	AA	
Family members	Lactase non-persistence	45	0	0	45	0	0	45
	Lactase persistence	0	32	13	0	32	13	45
Case-control samples								
Finnish	Lactase non-persistence	59	0	0	53	6	0	59
	Lactase persistence	0	63	74	0	63	74	137

TABLE 3-continued

Distribution of C/T ₁₃₉₁₀ & G/A ₂₂₀₁₈ genotypes in lactase persistent/non-persistent alleles								
Genotype		C/T ₁₃₉₁₀			G/A ₂₂₀₁₈			Total
		CC	CT	TT	GG	GA	AA	
Non-Finnish ^a	Lactase non-persistence	40	0	0	39	1	0	40
	Lactase persistence	0	5	0	0	5	0	5
Total	Lactase non-persistence						0	144
	Lactase persistence							187

^anon-Finnish samples consist of 23 South Korean, 9 Italian and 7 German individuals

TABLE 4

Prevalence of the C/T ₁₃₉₁₀ variant in population samples								
DNA samples	Genotype				Allele frequency (%)		% (CC)	
					C	T		
analysed	CC	CT	TT	Total	C	T	genotype	
I. Finnish population:								
1. Eastern regions	108	287	176	571	0.440	0.560	18.9%	
2. Western regions	62	159	146	367	0.385	0.615	16.8%	
Total	170	446	322	938	0.418	0.582	18.1%	
II. CEPH parents:								
1. Utah families	7	33	52	92	0.255	0.745	7.6%	
2. French families	7	9	1	17	0.676	0.324	41.2%	

A total of 938 DNA samples of anonymous Finnish blood donors from small parishes from Eastern and Western parts within Finland, and 109 DNA samples from CEPH parents. The prevalence of hypolactasia in the populations is reflected by the genotype frequencies of CC alleles.

TABLE 5

LD between C/T ₁₃₉₁₀ and G/A ₂₂₀₁₈ variants in random Finnish samples								
Genotype at	Genotype at C/T ₁₃₉₁₀				D'	χ^2 (1 df)	P-value	
G/A ₂₂₀₁₈	CC	CT	TT	Total				
GG	162	2	1	165				
GA	6	440	3	449				
AA	2	4	318	324				
Total	170	446	322	938	0.984	42.41	7.62×10^{-11}	

LD was calculated using D' statistic¹⁸, p value is the significance of D' from 0 as described in methods¹⁸.

TABLE 6

Estimation of the introduction of the C/T ₁₃₉₁₀ variant into Finnish population using DISLAMB program.								
Marker Allele	AC3		LPH2					
	Lactase persistence	Lactase non-persistence	Lactase persistence	Lactase non-persistence				
1	0	1	0	1				
2	31	10	0	20				
3	0	1	0	14				
4	2	9	32	15				
5	0	31	0	2				

TABLE 6-continued

Estimation of the introduction of the C/T ₁₃₉₁₀ variant into Finnish population using DISLAMB program.				
Marker Allele	AC3		LPH2	
	Lactase persistence	Lactase non-persistence	Lactase persistence	Lactase non-persistence
λ^a	0.838		0.999	
Θ^b	0.00031 (0.000038-0.00099)		0.0000 (0.00000-0.00052)	
n ^c	570		450	

^a λ is the proportion of increase of a certain allele in disease chromosomes (lactase persistence allele) relative to its population frequency (0.60).

^b Θ is the recombination fraction, reflected by the distance of the mutation from the closest marker, assuming 1cM = 1Mb

^cn is the number of generation since the introduction of the founder mutation into a population Applying $\lambda = \alpha (1 - \Theta)^n$ formula.

d: Hypothetical allele used in the calculations as Θ is zero and α is one.

TABLE 7

Prevalence of lactose intolerance variants in biochemically verified samples							
Population	Number	C/T ₁₃₉₁₀			G/A ₂₂₀₁₈		
		CC	CT	TT	GG	GA	AA
1. Finnish							
Lactase persistence	182	0	95	87	0	95	87
Lactase non-persistence	116	116	0	0	110	6	0
2. Italian							
Lactase persistence	7	0	7	0	0	7	0
Lactase non-persistence	23	23	0	0	22	1	0
3. German							
Lactase persistence	0	0	0	0	0	0	0
Lactase non-persistence	8	8	0	0	8	0	0
4. Somalian							
Lactase persistence	0	0	0	0	0	0	0
Lactase non-persistence	42	42	0	0	42	0	0
6. South Koreans							
Lactase persistence	0	0	0	0	0	0	0
Lactase non-persistence	23	23	0	0	23	0	0
Total	401	212	102	87	205	109	87

TABLE 8

Prevalence of lactose-intolerance variants in various population samples								
Population	Number	Genotype						% Prevalence
		C/T13910			G/A22018			of Lactase
		CC	CT	TT	GG	GA	AA	Persistence allele
South Koreans	23	23	0	0	23	0	0	0 *
France	17	7	9	1	6	10	1	59 *
Basques	85	7	44	34	13	35	37	92 *
Southern Italians	100	89	11	0	88	12	0	11 *
Somali	79	74	5	0	78	1	0	6
Utah	92	7	33	52	7	30	55	92 *
African Americans	96	76	15	5	78	12	5	21 *
Marrocans	90	62	25	3	65	22	3	31 *
Sarawhi (African)	57	29	26	2	28	26	3	49 *
Saami	30	20	10	0	21	9	0	33 *
Tibet	23	23	0	0	23	0	0	0
Eastern Finnish	571	108	287	176	107	288	176	81 *
Western Finnish	367	62	159	146	58	161	148	83 *
Finn-ugrian tribes								
Xan	20	19	1	0	19	1	0	5
Xm	20	19	1	0	19	1	0	5
Mansi	22	20	2	0	20	2	0	9
Lkomi	10	7	3	0	7	3	0	30
Erza	30	17	10	3	19	9	2	43
Moksa	30	13	17	0	14	16	0	57 *
Udmort	30	12	16	2	11	15	4	60 *
Pakistani tribes								
Kalash	30	30	0	0	28	2	0	0
Burusho	30	29	1	0	27	3	0	3
Hazara	14	13	1	0	11	3	0	7
Kashmiri	20	15	5	0	14	6	0	25
Makrani Baluch	29	19	10	0	19	8	1	34
Brahui	30	17	10	3	16	11	3	43
Makrani (Negroid)	29	16	10	3	16	10	3	45
Pathan	29	12	16	1	13	14	2	59 *
Indian	29	11	13	5	10	12	5	62 *
Total	2032							

* The prevalence of lactase persistence allele is correlated very well with the reported prevalences for the lactase persistence allele (Simoons Fj. The geographic hypothesis and lactose malabsorption Am J Dig Dis 1978 23 (11): 963-80)

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 17

<210> SEQ ID NO 1

<211> LENGTH: 180

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<220> FEATURE:

<223> OTHER INFORMATION: lactase persistence type intron 13 of the MCM6 gene with single nucleotide polymorphism (SNP) t substituted by c at position -13910 5' from the intestinal lactase-phlorizine hydrolase (LPH) gene

<400> SEQUENCE: 1

acctttcatt caggaaaaat gtacttagac cctacaatgt actagtaggc ctctgcgctg 60

gcaatacaga taagataatg tagccctgg cctcaaagga actctcctcc ttaggttgca 120

tttgataat gtttgatttt tagattgttc tttagccct gcattccacg aggataggtc 180

<210> SEQ ID NO 2

<211> LENGTH: 180

-continued

<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<223> OTHER INFORMATION: lactase persistence type intron 9 of the MCM6 gene with single nucleotide polymorphism (SNP) a substituted by g at position -22018 5' from the intestinal lactase-phlorizine hydrolase (LPH) gene

<400> SEQUENCE: 2

taagaacatt ttacactctt cagtataaag aagtcagaat acccctaccc tatcagtaaa 60
ggcctataag ttaccattaa aaagatgtcc ttaaaaacag cattctcagc tggggcggt 120
ggctcacacc ttgtcccag tactttggga agccgaggtg ggtggatcac ctgaggtcag 180

<210> SEQ ID NO 3
<211> LENGTH: 3213
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<223> OTHER INFORMATION: lactase persistence type intron 13 of the MCM6 gene with single nucleotide polymorphism (SNP) t at position -13910 5' from the intestinal lactase-phlorizine hydrolase (LPH) gene

<400> SEQUENCE: 3

atcagagtca ctttgatatg atgagagcag agataaacag atttggtgca tgtttttaat 60
ctttggtatg ggacatacta gaattcactg caaatacatt tttatgtaac tgttgaatgc 120
tcatacgacc atggaattct tccctttaa gagcttggtg agcatttgag tgtagttggt 180
agacggagac gatcacgtca tagtttatag agtgcataaa gacgtaagtt accatttaat 240
acctttcatt caggaaaaat gtacttagac cctacaatgt actagtaggc ctctgcgctg 300
gcaatacaga taagataatg tagtccttgg cctcaaagga actctcctcc ttaggttgca 360
tttgataat gtttgatttt tagattgttc ttgagccct gcattccacg aggataggtc 420
agtgggtatt aacgaggtaa aaggggagta gtacgaaagg gcattcaagc gtcccatctt 480
cgcttcaacc aaagcagccc tgcgttttcc tagttttatt aatagggttg atgtaaggtc 540
gtctttgaaa aggggggttg gctttttttt acagtgtgac tgagggtataa tttataaaaa 600
gggaaatgta tggcatggtg agttttttca catacatcct tgtgaatacc cagctcaaga 660
tccaaaacat ttccataatt tcagaaagtt ccaaaccctt gcctcttttc agtcttagcc 720
ctcttcccct gaagtaacca ctgttccgac ttcaatcact acttttatcc cacagggtta 780
ttttttggct tttttccact aaattttcaa attccttgat atggtacttt actattgacg 840
aagtactttc acactagggt atttaatat ctttgattca cccaatattt agggaacacc 900
tgtaggggac aaaaaatgaa tgagagcccc tgccttccat tgctgcta at ctggtgggaa 960
cgagacatgt atttaattaa gcatgtaaaa aatagagtgg gtgatgaaat aatctatata 1020
ctaaatcccc atgacacaca gtttacctat gtaacaaacc tgcattgtga cccccgaacc 1080
taaaatataa gttggaattt aaaaaaaa gagagggaga atagagcatc acaaccagag 1140
tgctgagatg aattacttta ttaccaaaagg aggaggagga ctcaggaggg tgccgacgtt 1200
taaaccagct cactgaaggg tgtgcagaat ttggataggc aagataccct gggacaaggt 1260
cattctaaaa ccatgctaac atttgtaact tttttttcat tgtgatagtt cctgaaatga 1320
gttgcataaa actggtacat gtcttagggc agtctcta at tgatttttat tttgttctat 1380
ttttaaaaat tagtcttcaa atagcagatt cacatgatat taaaatatat gcacataaat 1440

-continued

tatatacaca aatatat	ctgaatgaaa tttagtatct	gcataatatt aagagctatt	1500
tctgtctcat atgttcataa	tcttcaccca ttaaaaaaac	ttttgttagg cctttctcac	1560
tctaagatta taaaaaatc	tccattatt tacctagcta	gttttctagt tgttccaaaa	1620
ccatttattg aacaatccat	ctttttgaca ctggtttggc	atgccttaat tatatattct	1680
tgtgtgtgtt aggatctcct	tttggaacttt ccattctgtt	cattgagtct taccagctcc	1740
tcttacattg gtaccatgat	gttttaactct atggggcttt	gtagtttaaa ttaggggcta	1800
gttccagcgc attgttctct	atcagctgtt aggaacttag	aaatcagctt gctctgtttt	1860
aaagaaaaac ctggatattt	tttatcagta taacattcta	tttatattaa ctgaagaat	1920
tgaaaacatc tatgattttt	cctattcagt aacgtatcac	ttagaatagg ttaggttgta	1980
ctactataaa atctcagctg	cataaaacaa tttttttttg	cttgtgctac acatccatta	2040
ggtcacaaag ggactcacct	tgtcaagtta ctacagagatt	caggctgata taaaggtttg	2100
atcttgacat acgctttcat	gatgacagaa agcagggaag	agaagggtgt gagccatgtg	2160
ctttctcccc cttctatcca	gaaatgacac atactcacat	ttcattcgcc agagaaatta	2220
acatggcccc tcctaagttc	aaatggatag agaaatgcct	tcctaccagg tgcccagaat	2280
tagaagagca aacattttgtg	aacagttctg agtaccacaa	ataccgttat ctttccactt	2340
aagtctctctg tttcactcag	tagtgcttta aacttttctt	catatgtttt tcagtgttcc	2400
ttgttgaaatt tcttgatatt	ttatcatgtt tgttcgtact	gggagtagcc tttttttcca	2460
tttcattttc tggctggttt	cattgctggt tgtttttttg	ttttgttttg tttttgagat	2520
ggagtctcac tctgtcgccc	aggctggagt gcagtgtcac	aatctcggt cactgcaacc	2580
tctgcctccc aggttcaagc	gattcttctt tctcagcctc	ctgagtagct gggattacag	2640
gcattgtgcca ccatgcccag	ctaatttttt atatttttag	tagagatggg gtttctccat	2700
gttggtcagg ctggtctcaa	actcccaatc tcagggtgatc	cgctgcctc tgccttccaa	2760
agtgtctggga ttatagacat	gagccaccgt gcctggccta	gttcttatgg gatgtatatg	2820
tctttggatt catatgatat	gtatatatgt ttatatttct	acaagtacat acctaggagt	2880
ggaattgttg ggtcataggt	taatgcatgt ttttctgcca	aacagttgtg tcaatttctg	2940
ttttcaccgc tgtgaatgag	agttgttcta ccttcttgac	aacacttgat attgtcagtc	3000
attttagcca ttctggtgaa	tttatagtgc tatttctgtg	tgtgtaagag agagaatgag	3060
agagggtgtt tgtgagaaaa	ccaaagcaac actgtgagag	tgtgtgtgtt tgtgagaaaa	3120
ccaaaataca tactactgtg	atttcattgg gagaaaatct	gtttggtata tcaaaaaaag	3180
tagcttaatt acttcatcat	tattggttta ggt		3213

<210> SEQ ID NO 4

<211> LENGTH: 1296

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<220> FEATURE:

<223> OTHER INFORMATION: lactase persistence type intron 9 of the MCM6 gene with single nucleotide polymorphism (SNP) a at position -22018 5' from the intestinal lactase-phlorizine hydrolase (LPH) gene

<400> SEQUENCE: 4

taagaacatt ttacactctt	cagtataaag aagtcagaat	accctaccc tatcagtaaa	60
ggcctataag ttaccattaa	aaagatgtcc ttaaaaacag	cattctcagc tgggcacggt	120

-continued

ggctcacacc	tttgtccag	tactttggga	agccgagggtg	ggtggatcac	ctgaggtcag	180
gagttcgaga	ccagcctggc	caacatggcg	aaaacccatt	ttctctacta	aaaatacaaa	240
aattagccgg	gcacgtggc	gggtgcttgt	ggtcccagct	actcaagagg	ctgaggtggg	300
aggatcactg	agcccaggag	gtggaggctg	cattgagcca	agattgtgcc	actgcactcc	360
agcctgggtg	acagagcgag	actctgtctc	aaaaaaacca	aaacaaaaaa	aaccagcat	420
tctttagtaa	ataattcata	gttttcttca	tctagaattt	aaaattgtga	tagttgatca	480
gcacgtcctg	agcacgtgtg	tttgcctgta	ctagtttaga	tcggtagatg	tgtatataag	540
ttataggtat	aaaatcaatc	ctgagttgac	acaaggtttt	gatgttgagt	acaagtacag	600
taagtgtata	tttttagtta	tgctcttagt	tttaagtcaa	ttgtgtgggt	ctttctagct	660
ttaggatctg	ttgaattatc	ttccttagaa	aagggagtta	agaatcttca	cttacctatc	720
ttctacttgt	ttggagaata	gaagagtccc	tgtggtagca	gactttgtga	gtttacttgt	780
aattttccat	ctgaagact	gttcttgttt	ttcgtgatga	agtcttgctc	tgtcgcccag	840
gctggagtgc	agtgtgcaa	ccttggtcca	ctgcaacctc	tgctcccggg	gttcaagcaa	900
ttctcctgcc	tcagcctccc	gagtatctgg	gattacaggt	gcacaccacc	acacctggct	960
aatttttgta	ttttcagtag	agacgggggt	tcacatgtt	ggccaggctg	gtctcgaact	1020
cttgacctca	tgatcagccc	acctcagcct	tccaaagtgc	tgggattaca	ggtgtgagcc	1080
cccacactcg	gcggtgtgtg	ttttttaaga	gacaggttct	cactctgtca	cctaacctgg	1140
agtacagtgg	caatcatggc	tcaactgtaac	ctcaaatgcc	cggccttagt	gaagcgttct	1200
tcctgccttg	gcctcccaaa	gtgctgggat	tacaagtgtg	agccatgcat	ccagcttgaa	1260
agacagcttc	ttaggcttga	tttgtttggg	tacagg			1296

<210> SEQ ID NO 5

<211> LENGTH: 3213

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<220> FEATURE:

<223> OTHER INFORMATION: lactase persistence type intron 13 of the MCM6 gene with single nucleotide polymorphism (SNP) t substituted by c at position -13910 5' from the intestinal lactase-phlorizine hydrolase (LPH) gene

<400> SEQUENCE: 5

atcagagtca	ctttgatatg	atgagagcag	agataaacag	atttgttgca	tgtttttaat	60
ctttggtatg	ggacatacta	gaattcactg	caaatacatt	tttatgtaac	tgttgaatgc	120
tcatacgacc	atggaattct	tccttttaaa	gagcttggtg	agcatttgag	tgtagtgtgt	180
agacggagac	gatcacgtca	tagtttatag	agtgcataaa	gacgtaagtt	accatttaat	240
acctttcatt	caggaaaaat	gtacttagac	cctacaatgt	actagtaggc	ctctgcgctg	300
gcaatacaga	taagataatg	tagcccctgg	cctcaaagga	actctcctcc	ttaggttgca	360
tttgtataat	gtttgatttt	tagattgttc	tttgagccct	gcattccacg	aggataggtc	420
agtggttatt	aacgaggtaa	aaggggagta	gtacgaaagg	gcattcaagc	gtcccatctt	480
cgcttcaacc	aaagcagccc	tgctttttcc	tagttttatt	aatagggttg	atgtaaggtc	540
gtctttgaaa	aggggggttg	gctttttttt	acagtgtgac	tgaggataaa	tttataaaaa	600
gggaaatgta	tggcatggtg	agttttttca	catacatcct	tgtgaatacc	cagctcaaga	660
tccaaaacat	ttocataatt	tcagaaagtt	ccaaaccctc	gcctcttttc	agtccttagcc	720

-continued

ctcttcccct gaagtaacca ctgttccgac ttcaatcact acctttatcc cacaggttaa	780
ttttttggct tttttccact aaattttcaa attctttgat atggtacttt actattgacg	840
aagtactttc acactagggtt atttaatat ctttgattca cccaatat tt agggaacacc	900
tgtaggggac aaaaaatgaa tgagagcccc tgccttccat tgcgtgtaaat ctggtgggaa	960
cgagacatgt atttaattaa gcatgtaaaa aatagagtgg gtgatgaaat aatctatata	1020
ctaaatcccc atgacacaca gtttacctat gtaacaaacc tgcattgtga cccccaacc	1080
taaaatataa gttggaaatt aaaaaaaaa gagagggaga atagagcatc acaaccagag	1140
tgtcgagatg aattacttta ttaccaaaga aggaggagga ctgagggagg tgcgcagctt	1200
taaaccagct cactgaagggt tgtgcagaat ttggataggc aagataccct gggacaaggt	1260
cattctaaaa ccatgctaac atttgtactt tttttttcat tgtgatagtt cctgaaatga	1320
gttgcataaa actggtacat gtcttagggc agtctctaatt tgatttttat tttgttctat	1380
ttttaaaat tagtcttcaa atagcagatt cacatgatat taaaatatat gcacataaat	1440
tatatacaca aatatatttt ctgaatgaaa tttagtatct gcataatatt aagagctatt	1500
tctgtctcat atgttcataa tcttcatcca ttaaaaaaac ttttgttagg cctttctcac	1560
tctaagatta taaaaaatcc tccattattt tacctagcta gttttctagt tgttccaaaa	1620
ccattttattg aacaatccat ctttttgaca ctgggttggc atgccttaatt tatatattct	1680
tgtgtgtgtt aggatctcct tttggacttt ccattctgtt cattgagtct tatcagctcc	1740
tcttaccattg gtaccatgat gttttaatct atggggcttt gtagttaaaa tgtagggtta	1800
gttccagcgc attgttctct atcagctgtt aggaacttag aaatcagctt gctctgtttt	1860
aaagaaaaac ctggatattt tttatcagta taacattcta tttatattaa cttgaagaat	1920
tgaaaacatc tatgattttt cctattcagt aacgtatcac ttagaatagg ttaggttgta	1980
ctactataaa atctcagctg cataaaacaa tttttttttg cttgtgctac acatccatta	2040
gggtcatcaag ggactcacct tgtcaagtta ctgagagatt caggctgata taaagggttg	2100
atcttgacat acgctttcat gatgacagaa agcagggag agaggtgggt gagccatgtg	2160
ctttctcccc cttctatcca gaaatgacac atactcacat ttcattcgcc agagaaatta	2220
acatggcccc tcctaagttc aaatggatag agaaatgcct tcctaccagg tgcccagaat	2280
tagaagagca aacattttgtg aacagttctg agtaccacaa ataccgttat ctttccactt	2340
aagtctcttg tttcactcag tagtgcctta aacttttctt catatgtttt tcagtgtttc	2400
ttgttgaatt tcttgatatt ttatcatggt tgttcgtact gggagtagcc tttttttcca	2460
tttcattttc tggtgtgttt cattgtcgtt tgtttttttg ttttgttttg tttttgagat	2520
ggagtctcac tctgtcgccc aggctggagt gcagtgtcac aatctcggct cactgcaacc	2580
tctgcctccc aggttcaagc gattctctct tctcagcctc ctgagtagct gggattacag	2640
gcattgtgcca ccatgcccag ctaatttttt atatttttag tagagatggg gtttctccat	2700
gttggtcagg ctggtctcaa actcccaatc tcaggtgatc cgctgcctc tgccttccaa	2760
agtgtcggga ttatagacat gagccaccgt gcctggccta gttcttatgg gatgtatatg	2820
tctttggatt catatgatat gtatataagt ttatatttct acaagtacat acctaggagt	2880
ggaattgttg ggcataggt taatgcagt ttttctgcca aacagttgtg tcaatttctg	2940
ttttcaccgc tgtgaatgag agttgttcta ccttcttgac aacacttgat attgtcagtc	3000
attttagcca ttctggtgaa tttatagtgc tatttctgtg tgtgtaagag agagaatgag	3060

-continued

```

agaggggtgtt tgtgagaaaa ccaaagcaac actgtgagag tgtgtgtgtt tgtgagaaaa 3120
ccaaaataca tactactgtg atttcattgg gagaaaatct gtttgggtata tcaaaaaaag 3180
tagcttaatt acttcatcat tattggttta ggt 3213

```

```

<210> SEQ ID NO 6
<211> LENGTH: 1296
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<223> OTHER INFORMATION: lactase persistence type intron 9 of the MCM6
gene with single nucleotide polymorphism (SNP) a substituted by g
at position -22018 5' from the intestinal lactase-phlorizine
hydrolase (LPH) gene

```

```

<400> SEQUENCE: 6
taagaacatt ttacactctt cagtataaag aagtcagaat acccctaccc tatcagtaaa 60
ggcctataag ttaccattaa aaagatgtcc ttaaaaacag cattctcagc tgggcgcggt 120
ggctcacacc ttgtccccag tactttggga agccgagggt ggtggatcac ctgaggtcag 180
gagttcgaga ccagcctggc caacatggcg aaaaccatt ttcttacta aaaatacaaa 240
aattagccgg gcattgtggc ggggtgctgt ggtcccagct actcaagagg ctgaggtggg 300
aggatcactg agcccaggag gtggaggctg cattgagcca agattgtgcc actgcactcc 360
agcctgggtg acagagcgag actctgtctc aaaaaaacca aaacaaaaaa aaccagcat 420
tctttagtaa ataattcata gttttcttca tctagaattt aaaattgtga tagttgatca 480
gcattgtcctg agcacgtgtg tttgtgttta ctagttaga tcggttagatg tgtatataag 540
ttataggtat aaaatcaatc ctgagttgac acaaggtttt gatgttgagt acaagtacag 600
taagtgtata ttttagtta tgctcttagt ttttaagtcaa ttgtgtgggt ctttctagct 660
ttaggatctg ttgaattatc ttccttagaa aaggaggtta agaacttca cttacctatc 720
ttctacttgt ttggagaata gaagagtccc tgtggtagca gactttgtga gtttacttgt 780
aattttccat ctgaaagact gttcttgttt ttcgtgatga agtcttgctc tgcgcccag 840
gctggagtgc agtggtgcaa ccttggtcca ctgcaacctc tgcctcccgg gttcaagcaa 900
ttctcctgcc tcagcctccc gagtatctgg gattacaggt gcacaccacc acacctggct 960
aattttgtga ttttcagtag agacgggggt tcaccatgtt ggccaggctg gtctcgaact 1020
cttgacctca tgatcagccc acctcagcct tccaaagtgc tgggattaca ggtgtgagcc 1080
cccacactcg gccgttgttg ttttttaaga gacagggctc cactctgtca cctaacctgg 1140
agtacagtgg caatcatggc tcaactgaac ctcaaagcc cgcccttagt gaagcgttct 1200
tcctgccttg gcctcccaaa gtgctgggat tacaagtgtg agccatgcat ccagcttgaa 1260
agacagcttc ttaggcttga tttgtttggt tacagg 1296

```

```

<210> SEQ ID NO 7
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:biotinylated
PCR amplification primer (Bio-Reverse primer) for
C/T-13910 variant
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (1)
<223> OTHER INFORMATION: n = biotinylated t

```

-continued

<400> SEQUENCE: 7

naggtcagtg ggtattaacg aggt

24

<210> SEQ ID NO 8

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial
Sequence:unbiotinylated PCR amplification primer (Forward PCR
primer) for C/T-13910 variant

<400> SEQUENCE: 8

gtcactttga tatgatgaga gca

23

<210> SEQ ID NO 9

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence:biotinylated
PCR amplification primer (Bio-Reverse primer) for
C/T-13910 variant

<220> FEATURE:

<221> NAME/KEY: modified_base

<222> LOCATION: (1)

<223> OTHER INFORMATION: n = biotinylated c

<400> SEQUENCE: 9

nctcgттаат acccactgac cta

23

<210> SEQ ID NO 10

<211> LENGTH: 24

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial
Sequence:minisequencing primer for C/T-13910 variant, detection
primer for C/T-13910 variant

<400> SEQUENCE: 10

ggcaatacag ataagataat gtag

24

<210> SEQ ID NO 11

<211> LENGTH: 21

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence:biotinylated
PCR amplification primer (Bio-Reverse primer) for
G/A-22018 variant

<220> FEATURE:

<221> NAME/KEY: modified_base

<222> LOCATION: (1)

<223> OTHER INFORMATION: n = biotinylated t

<400> SEQUENCE: 11

ntgatcagca tgtcctgagc a

21

<210> SEQ ID NO 12

<211> LENGTH: 22

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial
Sequence:unbiotinylated PCR amplification primer (Forward PCR

-continued

primer) for G/A-22018 variant

<400> SEQUENCE: 12

ctaccctatc agtaaaggcc ta

22

<210> SEQ ID NO 13

<211> LENGTH: 21

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence:biotinylated
PCR amplification primer (Bio-Reverse primer) for
G/A-22018 variant

<220> FEATURE:

<221> NAME/KEY: modified_base

<222> LOCATION: (1)

<223> OTHER INFORMATION: n = biotinylated t

<400> SEQUENCE: 13

ngctcaggac atgctgatca a

21

<210> SEQ ID NO 14

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial
Sequence:minisequencing primer, detection primer for G/A-22018
variant

<400> SEQUENCE: 14

aaaaacagca ttctcagctg ggc

23

<210> SEQ ID NO 15

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence:biotinylated
PCR amplification primer for G/A-22018 variant

<220> FEATURE:

<221> NAME/KEY: modified_base

<222> LOCATION: (1)

<223> OTHER INFORMATION: n = biotinylated a

<400> SEQUENCE: 15

ngtctgtggc atgtgtcttc atg

23

<210> SEQ ID NO 16

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial
Sequence:unbiotinylated PCR amplification primer for G/A-22018
variant

<400> SEQUENCE: 16

tgctcaggac atgctgatca act

23

<210> SEQ ID NO 17

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial
Sequence:minisequencing primer for G/A-22018 variant

-continued

<400> SEQUENCE: 17

gacaaagggtg tgagccaccg

20

1. A nucleic acid molecule comprising a 5' portion of an intestinal lactase-phlorizine hydrolase (LPH) gene contributing to or indicative of adult-type hypolactasia wherein said nucleic acid molecule is selected from the group consisting of

- (a) a nucleic acid molecule having or comprising the nucleic acid sequence of SEQ ID NO: 2, the sequence of SEQ ID NO:2 is also depicted in FIG. 5 and comprised in the sequence as depicted in FIG. 9;
 - (b) a nucleic acid molecule of at least 20 nucleotides the complementary strand of which hybridizes under stringent conditions to the nucleic acid molecule of (a), wherein said polynucleotide/nucleic acid molecule has at a position corresponding to position -13910 5' from the LPH gene a cytosine residue;
 - (c) a nucleic acid molecule of at least 20 nucleotides the complementary strand of which hybridizes under stringent conditions to the nucleic acid molecule of (a), wherein said polynucleotide/nucleic acid molecule has at a position corresponding to position -22018 5' from the LPH gene a guanine residue;
 - (d) a nucleic acid molecule having or comprising the nucleic acid sequence of SEQ ID NO:4, the sequence of SEQ ID NO:4 is also depicted in FIG. 7;
 - (e) a nucleic acid molecule the complementary strand of which hybridizes under stringent conditions to the nucleic acid molecule of (d), wherein said polynucleotide/nucleic acid molecule has at a position corresponding to position -13910 of the LPH gene a thymidine residue; and
 - (f) a nucleic acid molecule the complementary strand of which hybridizes under stringent conditions to the nucleic acid molecule of (d), wherein said polynucleotide/nucleic acid molecule has at a position corresponding to position -22018 of the LPH gene a adenosine residue.
2. The nucleic acid molecule of claim 1 which is genomic DNA.
3. The nucleic acid molecule of claim 2 wherein said genomic DNA is part of a gene.
4. A fragment of the nucleic acid molecule of claim 1 having at least 14 nucleotides wherein said fragment comprises nucleotide position -13910 or nucleotide position -22018 of the LPH gene.
5. A nucleic acid molecule which is complementary to the nucleic acid molecule of claim 1.
6. A primer or primer pair, wherein the primer or primer pair hybridizes under stringent conditions to the nucleic acid molecule of claim 1 comprising nucleotide position -13910 or -22018 of the LPH gene or to the complementary strand thereof.
7. A method for testing for the presence or predisposition of adult-type hypolactasia or associated trait comprising testing a sample obtained from a prospective patient or from a person suspected of carrying such a predisposition for the presence of a nucleic acid molecule selected from the group consisting of

- (a) a nucleic acid molecule having or comprising the nucleic acid sequence of SEQ ID NO: 2, the sequence of SEQ ID NO:2 is also depicted in FIG. 5 and comprised in the sequence as depicted in FIG. 9;
 - (b) a nucleic acid molecule of at least 20 nucleotides the complementary strand of which hybridizes under stringent conditions to the nucleic acid molecule of (a), wherein said polynucleotide/nucleic acid molecule has at a position corresponding to position -13910 5' from the LPH gene a cytosine residue;
 - (c) a nucleic acid molecule of at least 20 nucleotides the complementary strand of which hybridizes under stringent conditions to the nucleic acid molecule of (a), wherein said polynucleotide/nucleic acid molecule has at a position corresponding to position -22018 5' from the LPH gene a guanine residue;
 - (d) a nucleic acid molecule having or comprising the nucleic acid sequence of SEQ ID NO:4, the sequence of SEQ ID NO:4 is also depicted in FIG. 7;
 - (e) a nucleic acid molecule the complementary strand of which hybridizes under stringent conditions to the nucleic acid molecule of (d), wherein said polynucleotide/nucleic acid molecule has at a position corresponding to position -13910 of the LPH gene a thymidine residue; and
 - (f) a nucleic acid molecule the complementary strand of which hybridizes under stringent conditions to the nucleic acid molecule of (d), wherein said polynucleotide/nucleic acid molecule has at a position corresponding to position -22018 of the LPH gene a adenosine residue.
8. The method of claim 7, wherein said testing comprises hybridizing the complementary nucleic acid molecule as a probe under stringent conditions to nucleic acid molecules comprised in said sample and detecting said hybridization.
9. The method of claim 8 further comprising digesting the product of said hybridization with a restriction endonuclease or subjecting the product of said hybridization to digestion with a restriction endonuclease and analyzing the product of said digestion.
10. The method of claim 8, wherein said probe is detectably labeled.
11. The method of claim 8, wherein said testing comprises determining the nucleic acid sequence of at least a portion of the nucleic acid molecule, said portion comprising nucleotide position -13910 and/or nucleotide position -22018 of the LPH gene.
12. The method of claim 11, wherein the determination of the nucleic acid sequence is effected by solid-phase minisequencing.
13. The method of claim 11 further comprising, prior to determining said nucleic acid sequence, amplification of at least said portion of said nucleic acid molecule.
14. The method of claim 13, wherein said amplification is effected by or said amplification is the polymerase chain reaction (PCR).

15. The method of claim **14**, wherein said sample is blood, serum, plasma, fetal tissue, saliva, urine, mucosal tissue, mucus, vaginal tissue, fetal tissue obtained from the vagina, skin, hair, hair follicle or another human tissue.

16. The method of claim **7**, wherein said nucleic acid molecule from said sample is fixed to a solid support.

17. The method of claim **16**, wherein said solid support is a chip, a silica wafer, a bead or a microtiter plate.

18. Kit comprising the nucleic acid molecule of claim **1**, a primer or primer pair and one or more containers.

* * * * *

专利名称(译)	鉴定与成人型低乳酸症相关的DNA变体		
公开(公告)号	US20120329052A1	公开(公告)日	2012-12-27
申请号	US13/594731	申请日	2012-08-24
申请(专利权)人(译)	国家公共卫生研究所		
当前申请(专利权)人(译)	国家公共卫生研究所		
[标]发明人	PELTONEN LEENA ENATTAH NABIL JARVELA IRMA SAHI TIMO SAVILAHTI ERKKI TERWILLIGER JOSEPH		
发明人	PELTONEN, LEENA ENATTAH, NABIL JARVELA, IRMA SAHI, TIMO SAVILAHTI, ERKKI TERWILLIGER, JOSEPH		
IPC分类号	C12Q1/68 C07H21/04 G01N33/53 A61K35/76 A61K38/00 A61K48/00 A61P3/00 A61P43/00 C07K16/18 C12N1/15 C12N1/19 C12N1/21 C12N5/10 C12N7/00 C12N9/24 C12N9/38 C12N15/09 C12Q1/34 C12Q1/70 G01N33/566		
CPC分类号	A01K2217/05 A61K38/00 A61K48/00 C12N9/2468 C12Q1/6883 C12N9/2471 C12Y302/01023 C12Y302/01062 C12N9/2402 C12Y302/01108 C12Q2600/156 A61P3/00 A61P43/00		
优先权	2001119377 2001-08-10 EP 2001119528 2001-08-14 EP 60/315955 2001-08-31 US		
其他公开文献	US10144979		
外部链接	Espacenet USPTO		

摘要(译)

本发明涉及一种核酸分子，其包含有助于或指示成人型低乳糜泻的58#39;部分肠乳糖酶 - 根赤碱水解酶 (LPH) 基因。本发明进一步涉及用于测试成人型低乳糜泻的存在或易感性的方法，所述方法基于对上述核酸分子中包含的SNP的分析。另外，本发明涉及用于检测成人型低乳糜泻的存在或易感性的诊断组合物和试剂盒。

