



US 20170184573A1

(19) **United States**(12) **Patent Application Publication**
Kaneko et al.(10) **Pub. No.: US 2017/0184573 A1**(43) **Pub. Date: Jun. 29, 2017**(54) **SURROGATE BIOMARKER FOR
EVALUATING INTRACEREBRAL AMYLOID
 β PEPTIDE ACCUMULATION AND METHOD
FOR ANALYSIS THEREOF****Publication Classification**(51) **Int. Cl.**
G01N 33/53 (2006.01)
(52) **U.S. Cl.**
CPC **G01N 33/5308** (2013.01); **G01N 33/53**
(2013.01); **G01N 2333/4709** (2013.01)(71) Applicants: **Shimadzu Corporation**, Kyoto-shi,
Kyoto (JP); **National Center for
Geriatrics and Gerontology**, Obu-shi,
Aichi (JP)(72) Inventors: **Naoki Kaneko**, Kyoto-shi, Kyoto (JP);
Katsuhiko Yanagisawa, Ryugasaki-shi,
Ibaraki (JP)(73) Assignees: **Shimadzu Corporation**, Kyoto-shi,
Kyoto (JP); **National Center for
Geriatrics and Gerontology**, Obu-shi,
Aichi (JP)(21) Appl. No.: **15/313,376**(22) PCT Filed: **May 19, 2015**(86) PCT No.: **PCT/JP2015/064386**

§ 371 (c)(1),

(2) Date: **Nov. 22, 2016**(30) **Foreign Application Priority Data**

May 22, 2014 (JP) 2014-106560

(57) **ABSTRACT**

Provided are a biomarker for evaluating a cerebral $A\beta$ accumulation state using an amyloid precursor protein (APP)-derived peptide in a living body-derived sample as an index, and a method for analysis thereof. A living body-derived sample derived from a test subject is measured, and at least one ratio selected from the group consisting of APP669-711/APP672-713 ($A\beta$ 1-42); APP672-709 ($A\beta$ 1-38)/APP672-713 ($A\beta$ 1-42); APP674-711 ($A\beta$ 3-40)/APP672-713 ($A\beta$ 1-42); APP672-710 ($A\beta$ 1-39)/APP672-713 ($A\beta$ 1-42); APP672-711 ($A\beta$ 1-40)/APP672-713 ($A\beta$ 1-42); and APP672-711 (Ox $A\beta$ 1-40)/APP672-713 ($A\beta$ 1-42) is calculated, and when each of the ratios of the test subject is higher than each of standard levels which are ratios of a person having normal cognitive function NC- who is negative for cerebral $A\beta$ accumulation, an accumulation amount of cerebral $A\beta$ of the test subject is determined as larger than an accumulation amount of cerebral $A\beta$ of the person having normal cognitive function NC-.


Formation Path of $A\beta$ Peptides and p3 Peptides from APP

Fig.1





Fig.2




Dunnett's test

* $0.05 > P \geq 0.01$, ** $0.01 > P \geq 0.001$, *** $P < 0.001$


Fig.3

(A) APP672-713 (A β 1-42) / SIL-A β 1-38



NC- vs NC+, AUC=0.789


(B) APP672-713 (A β 1-42) / SIL-A β 1-38



NC- vs MCI, AUC=0.746


Fig.4

(C) APP672-713 (A β 1-42) / SIL-A β 1-38




NC- vs AD, AUC=0.864

(D) APP672-713 (A β 1-42) / SIL-A β 1-38




NC- vs PIB+, AUC=0.808

Fig.5




Dunnett's test

* $0.05 > P \geq 0.01$, ** $0.01 > P \geq 0.001$, *** $P < 0.001$

Fig.6**(A)** APP672-709 (A β 1-38) / APP672-713 (A β 1-42)


NC- vs NC+, AUC=0.876

(B) APP672-709 (A β 1-38) / APP672-713 (A β 1-42)

NC- vs MCI, AUC=0.777


Fig.7

(C) APP672-709 (A β 1-38) / APP672-713 (A β 1-42)




NC- vs AD, AUC=0.757

(D) APP672-709 (A β 1-38) / APP672-713 (A β 1-42)



NC- vs PIB+, AUC=0.795

Fig.8




Dunnett's test

* $0.05 > P \geq 0.01$, ** $0.01 > P \geq 0.001$, *** $P < 0.001$


Fig.9

(A) APP674-711 (A β 3-40) / APP672-713 (A β 1-42)



NC- vs NC+, AUC=0.886


(B) APP674-711 (A β 3-40) / APP672-713 (A β 1-42)



NC- vs MCI, AUC=0.826


Fig.10

(C) APP674-711 ($A\beta$ 3-40) / APP672-713 ($A\beta$ 1-42)



NC- vs AD, AUC=0.973


(D) APP674-711 ($A\beta$ 3-40) / APP672-713 ($A\beta$ 1-42)



NC- vs PIB+, AUC=0.966

Fig.11

APP672-710(A β 1-39) / APP672-713(A β 1-42)




Dunnett's test

* $0.05 > P \geq 0.01$, ** $0.01 > P \geq 0.001$, *** $P < 0.001$


Fig.12

(A) APP672-710 (A β 1-39) / APP672-713 (A β 1-42)



NC- vs NC+, AUC=0.955


(B) APP672-710 (A β 1-39) / APP672-713 (A β 1-42)



NC- vs MCI, AUC=0.909


Fig.13

(C) APP672-710 (A β 1-39) / APP672-713 (A β 1-42)




NC- vs AD, AUC=0.914

(D) APP672-710 (A β 1-39) / APP672-713 (A β 1-42)



NC- vs PIB+, AUC=0.924

Fig.14




Dunnett's test

* $0.05 > P \geq 0.01$, ** $0.01 > P \geq 0.001$, *** $P < 0.001$


Fig.15

(A) APP672-711 ($A\beta$ 1-40) / APP672-713 ($A\beta$ 1-42)



NC- vs NC+, AUC=0.876


(B) APP672-711 ($A\beta$ 1-40) / APP672-713 ($A\beta$ 1-42)



NC- vs MCI, AUC=0.803


Fig.16

(C) APP672-711 (A β 1-40) / APP672-713 (A β 1-42)



NC- vs AD, AUC=0.743


(D) APP672-711 (A β 1-40) / APP672-713 (A β 1-42)



NC- vs PIB+, AUC=0.798

Fig.17

OxAPP672-711(OxA β 1-40) / APP672-713(A β 1-42)




Dunnett's test

* $0.05 > P \geq 0.01$, ** $0.01 > P \geq 0.001$, *** $P < 0.001$


Fig.18

(A) OxAPP672-711 (Ox $\text{A}\beta$ 1-40) / APP672-713 ($\text{A}\beta$ 1-42)



NC- vs NC+, AUC=0.897


(B) OxAPP672-711 (Ox $\text{A}\beta$ 1-40) / APP672-713 ($\text{A}\beta$ 1-42)



NC- vs MCI, AUC=0.843


Fig.19

(C) OxAPP672-711 (OxA β 1-40) / APP672-713 (A β 1-42)




NC- vs AD, AUC=0.985

(D) OxAPP672-711 (OxA β 1-40) / APP672-713 (A β 1-42)



NC- vs PIB+, AUC=0.876

Fig.20




Dunnett's test

* $0.05 > P \geq 0.01$, ** $0.01 > P \geq 0.001$, *** $P < 0.001$


Fig.21

(A) APP669-711 / APP672-713 ($A\beta$ 1-42)



NC- vs NC+, AUC=0.930


(B) APP669-711 / APP672-713 ($A\beta$ 1-42)



NC- vs MCI, AUC=0.966


Fig.22

(C) APP669-711 / APP672-713 ($A\beta$ 1-42)




NC- vs AD, AUC=0.997

(D) APP669-711 / APP672-713 ($A\beta$ 1-42)



NC- vs PIB+, AUC=0.969

Fig.23



SURROGATE BIOMARKER FOR EVALUATING INTRACEREBRAL AMYLOID β PEPTIDE ACCUMULATION AND METHOD FOR ANALYSIS THEREOF

TECHNICAL FIELD

[0001] The present invention pertains to the brain neuroscience field and the clinical medicine field, and relates to a surrogate biomarker for evaluating a cerebral amyloid β peptide (Aβ) accumulation state, and a method for analysis thereof. More specifically, the present invention relates to a surrogate biomarker for evaluating a cerebral Aβ accumulation state using, as an index, a level of Aβ and Aβ-like peptide generated by decomposition of amyloid precursor protein (APP) in a living body-derived sample, and a method for analysis thereof. The biomarker of the present invention is a marker to be used for, for example, presymptomatic diagnosis, screening for subjects of developing preventive intervention (pre-emptive therapeutic drug administration etc.) and evaluation of drug efficacy of therapeutic drugs and prophylactic drugs regarding Alzheimer's disease.

BACKGROUND ART

[0002] Alzheimer's disease (AD) is a principal cause of dementia, and occupies 50 to 60% of the entire dementia. The number of patients suffering from dementia was more than or equal to 24 million in the world in 2001, and is estimated to reach 81 million in 2040 (Non-Patent Document 1). It is considered that Aβ is deeply involved in development of Alzheimer's disease. Aβ is produced as a result of proteolysis of amyloid precursor protein (APP) which is a single-pass transmembrane protein and is composed of 770 amino acid residues, by β-secretase and γ-secretase (see FIG. 1). Appearance of senile plaques due to aggregation of Aβ accompanying fibrosis triggers aggregation and accumulation of tau protein inside neurocytes to cause nerve malfunction and neuronal cell death. It is considered that this results in extreme deterioration of the cognitive ability. It has long been known that Aβ mainly consists of 40 mer (Aβ1-40) and 42 mer (Aβ1-42) and migrates into cerebrospinal fluid (CSF) and blood. Further, in recent years, existence of Aβ-like peptides having lengths different from those of Aβ1-40 and Aβ1-42 in CSF has been reported (Non-Patent Document 2).

[0003] Alzheimer's disease develops latently and advances slowly. Diagnosis of Alzheimer's disease is made by performing ADAS-cog, MMSE, DemTect, SKT, or a test of cognitive function such as a clock drawing test for examining the clinical symptom, and examination of image findings of magnetic resonance imaging diagnosis (MRI), positron emission tomography (PET) and the like in combination. While MRI is an image diagnostic method capable of detecting cerebral degenerative atrophy, cerebral atrophy is not specific for Alzheimer's disease. Meanwhile, as an image diagnostic method that visualizes retention of a ligand molecule that specifically detects Aβ deposits (PiB: Pittsburgh compound-B), PiB-PET has been known. It has been found that thioflavin T-analogue (11C) PiB is retained in reflection of Aβ that gradually accumulates in a specific region of the brain of a patient suffering from mild cognitive impairment (MCI) or mild Alzheimer's disease, and hence PiB-PET is an optimum tool for detecting Aβ deposits. From the pathological findings of necropsy of Alzheimer's disease,

it is found that a large quantity of senile plaques has already accumulated even in the case of mild cognitive function decline. This leads the inference that aggregation and deposition of Aβ start quite long before exteriorization of dementia, and the result supporting this inference has been reported also in the findings of PiB-PET. However, PET examination requires massive equipment, and requires an expensive examination fee to perform one examination, and thus is not suited as a method that allows the citizen to widely have the examination.

[0004] A biomarker in blood or cerebrospinal fluid (CSF) is a useful index capable of detecting the development and progression of a disease on the molecular level. In Alzheimer's disease, a decrease in concentration of Aβ1-42 in CSF or concentration ratio of Aβ1-42/Aβ1-40, and an increase in total tau value or phosphorylation tau value are reported to be a useful diagnostic marker (Patent Document 1: JP-A-2010-19864, Non-Patent Document 3). However, collection of CSF is highly invasive, and is not suited as a method that allows the citizen to widely have the examination.

[0005] Under these circumstances, the potentiality of Aβ1-42 existing in blood as an Alzheimer's disease diagnostic marker is expected in a blood examination; however, it has been reported that the relationship between blood Aβ1-42 concentration and Alzheimer's disease development is low unlike the behavior of Aβ1-42 in CSF (Non-Patent Document 3). The reason for this has not been elucidated.

[0006] Also, Patent Document 2: JP-A-2013-63976 discloses a monoclonal antibody that does not recognize a soluble Aβ monomer, but specifically binds only to a soluble Aβ oligomer, and also discloses a diagnostic method of Alzheimer's disease using the antibody. The paragraph [0104] of the publication discloses a method in which when the ratio of Aβ oligomer to Aβ monomer in a sample of a subject is higher than that of a normal healthy person, the subject is determined as being a candidate for Alzheimer's disease.

[0007] Non-Patent Document 4 discloses existence of 22 kinds of APP-derived Aβ and Aβ-like peptides in human plasma by combination of immunoprecipitation and a mass spectrometer, and also discloses a method for quantifying these Aβ and Aβ-like peptides.

PRIOR ART DOCUMENTS

Patent Documents

- [0008]** Patent Document 1: JP-A-2010-19864
- [0009]** Patent Document 2: JP-A-2013-63976

Non-Patent Documents

- [0010]** Non-Patent Document 1: Blennow K, de Leon M J, Zetterberg H.: Alzheimer's disease. *Lancet*. 2006 Jul. 29; 368(9533): 387-403
- [0011]** Non-Patent Document 2: Portelius E, Westman-Brinkmalm A, Zetterberg H, Blennow K.: Determination of beta-amyloid peptide signatures in cerebrospinal fluid using immunoprecipitation-mass spectrometry. *J Proteome Res*. 2006 April; 5(4): 1010-6
- [0012]** Non-Patent Document 3: Hampel H, Shen Y, Walsh D M, Aisen P, Shaw L M, Zetterberg H, Trojanowski J Q, Blennow K.: Biological markers of amyloid beta-related mechanisms in Alzheimer's disease. *Exp Neurol*. 2010 June; 223(2): 334-46

[0013] Non-Patent Document 4: Kaneko N, Yamamoto R, Sato T A, Tanaka K.: Identification and quantification of amyloid beta-related peptides in human plasma using matrix-assisted laser desorption/ionization time-of-flight mass spectrometry. *Proc Jpn Acad Ser B Phys Biol Sci.* 2014; 90(3):104-17

SUMMARY OF THE INVENTION

Problems to be Solved by the Invention

[0014] It has been found that a large quantity of A β has been deposited before exteriorization of the cognitive function decline in an Alzheimer's disease (AD) patient. Although PiB-PET is effective for detecting A β accumulation, it requires high examination cost and long time for executing the examination, and thus is not a diagnostic method that allows for a majority of elderly people to easily undergo the examination. Therefore, a simplified analytical method capable of detecting increase in A β accumulation before exteriorization of clinical symptoms has been demanded.

[0015] As described above, generally, an examination method using a biomarker existing in blood or cerebrospinal fluid (CSF) as an index is an effective method capable of conveniently detecting the development and progression of a disease on the molecular level. Patent Document 1 and Non-Patent Document 3 described above have reported that in Alzheimer's disease, a decrease in concentration of A β 1-42 in CSF or concentration ratio of A β 1-42/A β 1-40, and an increase in total tau value or phosphorylation tau value are useful diagnostic markers. On the other hand, however, Non-Patent Document 3 has also reported that the relationship between blood A β 1-42 concentration and AD development is low unlike the case of CSF A β 1-42.

[0016] In previous reports regarding A β in blood, the correlatively with AD is examined only for concentrations of two kinds of A β 1-40 and A β 1-42 in blood. However, existence of shorter A β peptides that are cleaved on the N-terminal side or C-terminal side of A β 1-40 in CSF have been found besides A β 1-40 and A β 1-42 by a combinational method of immunoprecipitation and a mass spectrometer. In blood, it had been technically difficult to detect A β that is present in a smaller quantity than in CSF by immunoprecipitation and a mass spectrometer; however, as a result of improvement in the immunoprecipitation, existence of 22 kinds of APP-derived A β and A β -like peptides in human plasma has been elucidated, and a method for quantifying these peptides has been developed (Non-Patent Document 4).

[0017] An object of the present invention is to provide a biomarker for evaluating a cerebral A β accumulation state using an amyloid precursor protein (APP)-derived peptide in a living body-derived sample as an index, and a method for analysis thereof. In particular, an object of the present invention is to provide a biomarker for evaluating a cerebral A β accumulation state using amyloid precursor protein (APP)-derived A β and A β -like peptide in a blood sample as an index, and a method for analysis thereof. More specifically, an object of the present invention to provide a marker to be used for, for example, presymptomatic diagnosis, screening for subjects of developing preventive intervention (pre-emptive therapeutic drug administration etc.) and

evaluation of drug efficacy of therapeutic drugs and prophylactic drugs regarding Alzheimer's disease, and a method for analysis thereof.

Means for Solving the Problems

[0018] As a result of diligent efforts, the present inventors have found that specific APP-derived A β and A β -like peptide achieve the aforementioned object, and accomplished the present invention.

[0019] In the present description, "A β " is used as an abbreviation of an amyloid β peptide. That is, "A β " includes A β 1-40 and A β 1-42. A Peptide other than the A β generated by cleavage of amyloid precursor protein (APP) may be referred to as an A β -like peptide. A β and an A β -like peptide that are generated by cleavage of amyloid precursor protein (APP) may be referred to as "APP-derived peptide".

[0020] The present invention includes the following aspects.

(1) An analytical method for determining a cerebral A β accumulation state, the method comprising:

[0021] a measurement step of subjecting a living body-derived sample derived from a test subject to detection of a marker containing:

[0022] APP672-713 (A β 1-42) (SEQ ID NO.: 6), and

[0023] at least one selected from the group consisting of APP669-711 (SEQ ID NO.: 7), APP672-709 (A β 1-38) (SEQ ID NO.: 1), APP674-711 (A β 3-40) (SEQ ID NO.: 2), APP672-710 (A β 1-39) (SEQ ID NO.: 3), APP672-711 (A β 1-40) (SEQ ID NO.: 4), and OxAPP672-711 (OxA β 1-40) (SEQ ID NO.: 5),

to obtain measurement levels of:

[0024] APP672-713 (A β 1-42), and

[0025] the at least one selected from the group consisting of APP669-711, APP672-709 (A β 1-38), APP674-711 (A β 3-40), APP672-710 (A β 1-39), APP672-711 (A β 1-40), and OxAPP672-711 (OxA β 1-40) in the living body-derived sample;

[0026] a calculation step of calculating at least one ratio selected from the group consisting of:

[0027] a ratio of APP669-711 level to APP672-713 (A β 1-42) level: APP669-711/APP672-713 (A β 1-42);

[0028] a ratio of APP672-709 (A β 1-38) level to APP672-713 (A β 1-42) level: APP672-709 (A β 1-38)/APP672-713 (A β 1-42);

[0029] a ratio of APP674-711 (A β 1-40) level to APP672-713 (A β 1-42) level: APP674-711 (A β 3-40)/APP672-713 (A β 1-42);

[0030] a ratio of APP672-710 (A β 3-39) level to APP672-713 (A β 1-42) level: APP672-710 (A β 1-39)/APP672-713 (A β 1-42);

[0031] a ratio of APP672-711 (A β 1-40) level to APP672-713 (A β 1-42) level: APP672-711 (A β 1-40)/APP672-713 (A β 1-42); and

[0032] a ratio of OxAPP672-711 (OxA β 1-40) level to APP672-713 (A β 1-42) level: OxAPP672-711 (OxA β 1-40)/APP672-713 (A β 1-42); and

[0033] an evaluation step of determining that an amount of cerebral A β accumulation of the test subject is larger than an amount of cerebral A β accumulation of a person having normal cognitive function NC- who is negative for cerebral A β accumulation, when each of the ratios of the test subject is higher than each of standard levels which are ratios of the person having normal cognitive function NC-.

(2) The analytical method for determining a cerebral A β accumulation state according to the above (1), wherein the living body-derived sample is selected from the group consisting of blood, cerebrospinal fluid, urine, body secreting fluid, feces, saliva, and sputum.

(3) A marker for determining a cerebral A β accumulation state, the marker being selected from the group consisting of:

[0034] a combination of APP669-711 (SEQ ID NO.: 7) and APP672-713 (A β 1-42) (SEQ ID NO.: 6),

[0035] a combination of APP672-709 (A β 1-38) (SEQ ID NO.: 1) and APP672-713 (A β 1-42) (SEQ ID NO.: 6),

[0036] a combination of APP674-711 (A β 3-40) (SEQ ID NO.: 2) and APP672-713 (A β 1-42) (SEQ ID NO.: 6),

[0037] a combination of APP672-710 (A β 1-39) (SEQ ID NO.: 3) and APP672-713 (A β 1-42) (SEQ ID NO.: 6),

[0038] a combination of APP672-711 (A β 1-40) (SEQ ID NO.: 4) and APP672-713 (A β 1-42) (SEQ ID NO.: 6), and

[0039] a combination of OxAPP672-711 (OxAPP672-711) (SEQ ID NO.: 5) and APP672-713 (A β 1-42) (SEQ ID NO.: 6), in a living body-derived sample.

(4) An analytical method for determining a cerebral A β accumulation state, the method comprising:

[0040] a measurement step of subjecting a blood sample derived from a test subject to detection of a marker containing APP672-713 (A β 1-42) (SEQ ID NO.: 6), to obtain a measurement level of APP672-713 (A β 1-42) in the blood sample; and

[0041] an evaluation step of determining that an amount of cerebral A β accumulation of the test subject is larger than an amount of cerebral A β accumulation of a person having normal cognitive function NC— who is negative for cerebral A β accumulation, when the measurement level of APP672-713 (A β 1-42) of the test subject is lower than a standard level which is a level of APP672-713 (A β 1-42) of the person having normal cognitive function NC—.

(5) A method for determining efficacy of a medical intervention regarding a cerebral A β accumulation state, the method comprising, before and after a medical intervention performed for a test subject:

[0042] a measurement step of subjecting a living body-derived sample derived from the test subject to detection of a marker containing:

[0043] APP672-713 (A β 1-42) (SEQ ID NO.: 6), and

[0044] at least one selected from the group consisting of APP669-711 (SEQ ID NO.: 7), APP672-709 (A β 1-38) (SEQ ID NO.: 1), APP674-711 (A β 3-40) (SEQ ID NO.: 2), APP672-710 (A β 1-39) (SEQ ID NO.: 3), APP672-711 (A β 1-40) (SEQ ID NO.: 4), and OxAPP672-711 (OxAPP672-711) (SEQ ID NO.: 5),

to obtain measurement levels of:

[0045] APP672-713 (A β 1-42), and

[0046] the at least one selected from the group consisting of APP669-711, APP672-709 (A β 1-38), APP674-711 (A β 1-40), APP672-710 (A β 1-39), APP672-711 (A β 1-40), and OxAPP672-711 (OxAPP672-711) in the living body-derived sample; and

[0047] a calculation step of calculating at least one ratio selected from the group consisting of:

[0048] a ratio of APP669-711 level to APP672-713 (A β 1-42) level: APP669-711/APP672-713 (A β 1-42);

[0049] a ratio of APP672-709 (A β 1-38) level to APP672-713 (A β 1-42) level: APP672-709 (A β 1-38)/APP672-713 (A β 1-42);

[0050] a ratio of APP674-711 (A β 3-40) level to APP672-713 (A β 1-42) level: APP674-711 (A β 3-40)/APP672-713 (A β 1-42);

[0051] a ratio of APP672-710 (A β 1-39) level to APP672-713 (A β 1-42) level: APP672-710 (A β 1-39)/APP672-713 (A β 1-42);

[0052] a ratio of APP672-711 (A β 1-40) level to APP672-713 (A β 1-42) level: APP672-711 (A β 1-40)/APP672-713 (A β 1-42); and

[0053] a ratio of OxAPP672-711 (OxAPP672-711) level to APP672-713 (A β 1-42) level: OxAPP672-711 (OxAPP672-711)/APP672-713 (A β 1-42); and

[0054] comparing each of the ratios of the test subject before the medical intervention and each of the ratios of the test subject after the medical intervention to determine efficacy of the medical intervention regarding a cerebral A β accumulation state.

[0055] In the present invention, the term “level of marker” basically means a concentration, but may be other units applied correspondingly to concentration by a person skilled in the art. The term “test subject” includes human, and mammals other than human (rat, dog, cat etc.). In the present invention, the living body-derived sample is disposed of rather than being returned to the test subject (for example, subject) from which the biological sample is derived. The medical intervention includes administration of a therapeutic drug or a prophylactic drug, dietetic therapy, exercise therapy, learning therapy, surgical operation and the like.

Effects of the Invention

[0056] The present invention provides a marker for determining a cerebral A β accumulation state of a test subject, including a combination of APP672-713 (A β 1-42) (SEQ ID NO.: 6) and at least one selected from the group consisting of APP669-711 (SEQ ID NO.: 7), APP672-709 (A β 1-38) (SEQ ID NO.: 1), APP674-711 (A β 3-40) (SEQ ID NO.: 2), APP672-710 (A β 1-39) (SEQ ID NO.: 3), APP672-711 (A β 1-40) (SEQ ID NO.: 4), and OxAPP672-711 (OxAPP672-711) (SEQ ID NO.: 5) in a living body-derived sample. Also, the present invention provides a method for analysis of the marker.

[0057] By analyzing the marker in the living body-derived sample of the test subject, it is possible to discriminate whether cerebral A β accumulation in the test subject is less than or equal to the standard level or cerebral A β is excessively accumulated. The present invention is applicable to detection of not only the advanced stage of Alzheimer's disease in which cerebral A β is excessively accumulated and a cognitive functional disorder has appeared, but also the early stage of Alzheimer's disease in which cerebral A β is excessively accumulated but a cognitive functional disorder has not been appeared.

[0058] According to the present invention, as the living body-derived sample, not only blood, but also cerebrospinal fluid (CSF), urine, body secreting fluid, feces, saliva, and sputum can be used. Therefore, in the stage where the preventive method and the pre-emptive therapeutic method for Alzheimer's disease have established, analysis of a cerebral A β accumulation state for a person having normal cognitive function in a general medical examination, a complete physical examination and the like is effective for presymptomatic diagnosis of Alzheimer's disease.

[0059] By applying the present invention before and after a medical intervention performed for the test subject, it is

possible to evaluate the drug efficacy of a therapeutic drug or a prophylactic drug for Alzheimer's disease, or to evaluate the efficacy of other treatment. Also, the present invention is useful for follow-up of a patient suffering from Alzheimer's disease.

BRIEF DESCRIPTION OF THE DRAWINGS

[0060] FIG. 1 shows a diagram schematically showing the formation path of A β and p3 peptides by cleavages of amyloid precursor protein (APP).

[0061] FIG. 2 is a box-and-whisker plot showing the intensity ratio of APP672-713 (A β 1-42) to the internal standard SIL-A β 1-38 in each group (NC-, NC+, MCI, AD) for APP672-713 (A β 1-42) in Example 1.

[0062] FIG. 3(A) and FIG. 3(B) each show a receiver operating characteristic (ROC) curve of each group (NC+, MCI) versus NC- group for APP672-713 (A β 1-42)/SIL-A β 1-38.

[0063] FIG. 4(C) and FIG. 4(D) each show an ROC curve of each group (AD, PiB+) versus NC- group for APP672-713 (A β 1-42)/SIL-A β 1-38.

[0064] FIG. 5 is a box-and-whisker plot showing the intensity ratio of APP672-709 (A β 1-38) to APP672-713 (A β 1-42) in each group (NC-, NC+, MCI, AD) in Example 1.

[0065] FIG. 6(A) and FIG. 6(B) each show an ROC curve of each group (NC+, MCI) versus NC- group for APP672-709 (A β 1-38)/APP672-713 (A β 1-42).

[0066] FIG. 7(C) and FIG. 7(D) each show an ROC curve of each group (AD, PiB+) versus NC- group for APP672-709 (A β 1-38)/APP672-713 (A β 1-42).

[0067] FIG. 8 is a box-and-whisker plot showing the intensity ratio of APP674-711 (A β 3-40) to APP672-713 (A β 1-42) in each group (NC-, NC+, MCI, AD) in Example 1.

[0068] FIG. 9(A) and FIG. 9(B) each show an ROC curve of each group (NC+, MCI) versus NC- group for APP674-711 (A β 3-40)/APP672-713 (A β 1-42).

[0069] FIG. 10(C) and FIG. 10(D) each show an ROC curve of each group (AD, PiB+) versus NC- group for APP674-711 (A β 3-40)/APP672-713 (A β 1-42).

[0070] FIG. 11 is a box-and-whisker plot showing the intensity ratio of APP672-710 (A β 1-39) to APP672-713 (A β 1-42) in each group (NC-, NC+, MCI, AD) in Example 1.

[0071] FIG. 12(A) and FIG. 12(B) each show an ROC curve of each group (NC+, MCI) versus NC- group for APP672-710 (A β 1-39)/APP672-713 (A β 1-42).

[0072] FIG. 13(C) and FIG. 13(D) each show an ROC curve of each group (AD, PiB+) versus NC- group for APP672-710 (A β 1-39)/APP672-713 (A β 1-42).

[0073] FIG. 14 is a box-and-whisker plot showing the intensity ratio of APP672-711 (A β 1-40) to APP672-713 (A β 1-42) in each group (NC-, NC+, MCI, AD) in Example 1.

[0074] FIG. 15(A) and FIG. 15(B) each show an ROC curve of each group (NC+, MCI) versus NC- group for APP672-711 (A β 1-40)/APP672-713 (A β 1-42).

[0075] FIG. 16(C) and FIG. 16(D) each show an ROC curve of each group (AD, PiB+) versus NC- group for APP672-711 (A β 1-40)/APP672-713 (A β 1-42).

[0076] FIG. 17 is a box-and-whisker plot showing the intensity ratio of OxAPP672-711 (Ox A β 1-40) to APP672-713 (A β 1-42) in each group (NC-, NC+, MCI, AD) in Example 1.

[0077] FIG. 18(A) and FIG. 18(B) each show an ROC curve of each group (NC+, MCI) versus NC- group for OxAPP672-711 (Ox A β 1-40)/APP672-713 (A β 1-42).

[0078] FIG. 19(C) and FIG. 19(D) each show an ROC curve of each group (AD, PiB+) versus NC- group for OxAPP672-711 (Ox A β 1-40)/APP672-713 (A β 1-42).

[0079] FIG. 20 is a box-and-whisker plot showing the intensity ratio of APP669-711 to APP672-713 (A β 1-42) in each group (NC-, NC+, MCI, AD) in Example 1.

[0080] FIG. 21(A) and FIG. 21(B) each show an ROC curve of each group (NC+, MCI) versus NC- group for APP669-711/APP672-713 (A β 1-42).

[0081] FIG. 22(C) and FIG. 22(D) each show an ROC curve of each group (AD, PiB+) versus NC- group for APP669-711/APP672-713 (A β 1-42).

[0082] FIG. 23 is a scatter diagram of PiB accumulation mean value of cortical region (mcSUVR: mean cortical Standard Uptake Value Ratio) and APP669-711/APP672-713 (A β 1-42) in a total of 62 cases in Example 1. A probability ellipse ($p=0.95$) is indicated by a curve.

MODES FOR CARRYING OUT THE INVENTION

[1. Test Subject]

[0083] In the present invention, the test subject includes human, and mammals other than human (rat, dog, cat etc.). Hereinafter, the description will be made mainly for the case of human, but the same applies to mammals other than human.

[0084] In the method of the present invention, the subject may be any individuals including a person expected to be a normal healthy person regardless of past clinical history. For a person expected to be a normal healthy person, a cerebral A β accumulation state can be determined in a general medical examination, or a complete physical examination, preferably by a blood test, and the method is particularly effective for early detection/diagnosis of Alzheimer's disease. For a subject suspected to be a candidate for Alzheimer's disease as a result of ADAS-cog, MMSE, DemTect, SKT, or a test of cognitive function such as a clock drawing test for examining clinical symptom, and confirmation of image findings of magnetic resonance imaging diagnosis (MRI), positron emission tomography (PET) and the like, the method of the present invention can assist the diagnosis of Alzheimer's disease.

[2. Living Body-Derived Sample]

[0085] The marker of the present invention can be detected and analyzed in a living body-derived sample of a subject. Therefore, in the method of the present invention, a level of a marker in a living body-derived sample of a subject is analyzed.

[0086] The living body-derived sample can be selected from body fluids such as blood, cerebrospinal fluid (CSF), urine, body secreting fluid, saliva, and sputum; and feces. Among these, blood is preferred for diagnosis and presymp-

tomatic diagnosis of Alzheimer's disease in a general medical examination, a complete physical examination or the like.

[0087] The blood sample is a sample that is directly subjected to a measurement step of marker level, and includes whole blood, plasma, serum and the like. The blood sample can be prepared by appropriately treating whole blood collected from a test subject. The treatment performed in the case of preparing a blood sample from collected whole blood is not particularly limited, and any treatment that is clinically acceptable may be performed. For example, centrifugal separation or the like may be performed. The blood sample subjected to the measurement step may be appropriately stored at a low temperature such as freezing in the intermediate stage of the preparation step or in the post stage of the preparation step. In the present invention, the living body-derived sample such as a blood sample is disposed of rather than being returned to the subject from which it is derived.

[3. Marker]

[0088] The marker of the present invention includes a combination of APP672-713 (A β 1-42) (SEQ ID NO.: 6) and at least one selected from the group consisting of APP669-711 (SEQ ID NO.: 7), APP672-709 (A β 1-38) (SEQ ID NO.: 1), APP674-711 (A β 3-40) (SEQ ID NO.: 2), APP672-710 (A β 1-39) (SEQ ID NO.: 3), APP672-711 (A β 1-40) (SEQ ID NO.: 4), and OxAPP672-711 (Ox A β 1-40) (SEQ ID NO.: 5) in a living body-derived sample. For these markers, a significant difference has been observed between the level in the plasma sample from a person having normal cognitive function who is negative for cerebral A β accumulation and the level in the plasma sample from a subject having excessively accumulated cerebral A β .

APP672-713 (A β 1-42) (SEQ ID NO.: 6):
DAEFRHDSGYEVHHQKLVFFAEDVGSNKGAIIGLMVGGVVIA

APP669-711 (SEQ ID NO.: 7):
VKMDAEFRHDSGYEVHHQKLVFFAEDVGSNKGAIIGLMVGGVV

APP672-709 (A β 1-38) (SEQ ID NO.: 1):
DAEFRHDSGYEVHHQKLVFFAEDVGSNKGAIIGLMVGG

APP674-711 (A β 3-40) (SEQ ID NO.: 2):
EFRHDSGYEVHHQKLVFFAEDVGSNKGAIIGLMVGGVV

APP672-710 (A β 1-39) (SEQ ID NO.: 3):
DAEFRHDSGYEVHHQKLVFFAEDVGSNKGAIIGLMVGGV

APP672-711 (A β 1-40) (SEQ ID NO.: 4):
DAEFRHDSGYEVHHQKLVFFAEDVGSNKGAIIGLMVGGVV

OxAPP672-711 (Ox A β 1-40) (SEQ ID NO.: 5):
DAEFRHDSGYEVHHQKLVFFAEDVGSNKGAIIGLMVGGVV
(Met 706 is oxidized)

[0089] Amyloid precursor protein (APP) is a single-pass transmembrane protein and is composed of 770 amino acid residues. Amyloid precursor protein (APP) is proteolyzed by β secretase and γ secretase, and an amyloid β peptide (A β) is produced by the proteolysis. APP672-713 and A β 1-42 indicate the same peptide (SEQ ID NO.: 6). APP672-711 and A β 1-40 indicate the same peptide (SEQ ID NO.: 4).

[0090] Table 1 shows amino acid sequences of APP-derived peptides, and Table 2 shows theoretical average

masses of the same. For these peptides, analysis of a marker for determining a cerebral A β accumulation state was performed.

TABLE 1

SEQ ID NO.	APP-derived peptides	Sequence
		VKMDAEFRHDSGYEVHHQKLVFF 669 AEDVGSNKGAIIGLMVGGVVIA 713
1	APP672-709 (A β 1-38)	DAEFRHDSGYEVHHQKLVFFAED VGSNKGAIIGLMVGG
2	APP674-711 (A β 3-40)	EFRHDSGYEVHHQKLVFFAEDVG SNKGAIIGLMVGGVV
3	APP672-710 (A β 1-39)	DAEFRHDSGYEVHHQKLVFFAED VGSNKGAIIGLMVGGV
4	APP672-711 (A β 1-40)	DAEFRHDSGYEVHHQKLVFFAED VGSNKGAIIGLMVGGVV
5	OxAPP672-711 (Ox A β 1-40)	DAEFRHDSGYEVHHQKLVFFAED VGSNKGAIIGLMVGGVV
6	APP672-713 (A β 1-42)	DAEFRHDSGYEVHHQKLVFFAED VGSNKGAIIGLMVGGVVIA
7	APP669-711	VKMDAEFRHDSGYEVHHQKLVFF AEDVGSNKGAIIGLMVGGVV

TABLE 2

SEQ ID NO.	Theoretical average mass
1	4132.6
2	4144.7
3	4231.8
4	4330.9
5	4346.9
6	4515.1
7	4689.4

[0091] In Tables 1 to 2, "APP-derived peptides" indicates APP-derived peptides, and "Theoretical average mass" indicates theoretical average mass.

[0092] In Tables 1 to 2, OxAPP672-711 (Ox A β 1-40) of SEQ ID NO.: 17 indicates a peptide oxidized at Met 706 in APP672-711 (A β 1-40) of SEQ ID NO.: 16.

[0093] In Alzheimer's disease, there is familial Alzheimer's disease. The markers of the present invention may include mutation of familial Alzheimer's disease in the amino acid sequence that is generally known.

[0094] For familial Alzheimer's disease, for example, the following mutation sequences are known.

[0095] Swedish mutation: Amino acids KM in APP670-671 are mutated into NL.

[0096] Italian mutation: Amino acid A at APP673 is mutated into V.

[0097] Leuven mutation: Amino acid E at APP682 is mutated into K.

[0098] Icelandic mutation: Amino acid A at APP673 is mutated into T.

[0099] British mutation: Amino acid H at APP677 is mutated into R.

[0100] Tottori mutation: Amino acid D at APP678 is mutated into N.

[0101] Italian mutation: Amino acid A at APP673 is mutated into V.

[0102] Arctic mutation: Amino acid E at APP693 is mutated into G.

[0103] Iowa mutation: Amino acid D at APP694 is mutated into N.

[0104] Dutch mutation: Amino acid E at APP693 is mutated into Q.

[4. Analysis of Marker]

[0105] In the present description, the following classification is made from the view point of the degree of advance of Alzheimer's disease.

[0106] NC-: a person who is negative for cerebral A β accumulation, and does not have a cognitive impairment

[0107] NC+: a person who is positive for cerebral A β accumulation, but does not have a cognitive impairment

[0108] MCI: a person who is positive for cerebral A β accumulation, and has a mild cognitive impairment

[0109] AD: a person who is positive for cerebral A β accumulation, and has a cognitive impairment

[0110] PiB+: combined group of the group NC+, the group MCI and the group AD; that is, a person determined as positive for cerebral A β accumulation regardless of the presence or absence of a cognitive impairment

[0111] Normally, a cerebral A β accumulation state is determined by comparing PiB accumulation amount between the cerebral cortex and the white matter based on a PiB-PET image.

[0112] A method for analyzing the marker of the present invention includes:

[0113] a measurement step of subjecting a living body-derived sample derived from a test subject to detection of a marker containing APP672-713 (A β 1-42) (SEQ ID NO.: 6), and at least one selected from the group consisting of APP669-711 (SEQ ID NO.: 7), APP672-709 (A β 1-38) (SEQ ID NO.: 1), APP674-711 (A β 3-40) (SEQ ID NO.: 2), APP672-710 (A β 1-39) (SEQ ID NO.: 3), APP672-711 (A β 1-40) (SEQ ID NO.: 4), and OxAPP672-711 (Ox A β 1-40) (SEQ ID NO.: 5), to obtain measurement levels of APP672-713 (A β 3-42), and the at least one selected from the group consisting of APP669-711, APP672-709 (A β 1-38), APP674-711 (A β 3-40), APP672-710 (A β 1-39), APP672-711 (A β 1-40), and OxAPP672-711 (Ox A β 1-40) in the living body-derived sample;

[0114] a calculation step of calculating at least one ratio selected from the group consisting of:

[0115] a ratio of APP669-711 level to APP672-713 (A β 1-42) level: APP669-711/APP672-713 (A β 1-42);

[0116] a ratio of APP672-709 (A β 1-38) level to APP672-713 (A β 1-42) level: APP672-709 (A β 1-38)/APP672-713 (A β 1-42);

[0117] a ratio of APP674-711 (A β 3-40) level to APP672-713 (A β 1-42) level: APP674-711 (A β 3-40)/APP672-713 (A β 1-42);

[0118] a ratio of APP672-710 (A β 1-39) level to APP672-713 (A β 1-42) level: APP672-710 (A β 1-39)/APP672-713 (A β 1-42);

[0119] a ratio of APP672-711 (A β 1-40) level to APP672-713 (A β 1-42) level: APP672-711 (A β 1-40)/APP672-713 (A β 1-42); and

[0120] a ratio of OxAPP672-711 (Ox A β 1-40) level to APP672-713 (A β 1-42) level: OxAPP672-711 (Ox A β 1-40)/APP672-713 (A β 1-42); and

[0121] an evaluation step of determining that an amount of cerebral A β accumulation of the test subject is larger than an amount of cerebral A β accumulation of a person having normal cognitive function NC- who is negative for cerebral A β accumulation, when each of the ratios of the test subject is higher than each of standard levels which are ratios of the person having normal cognitive function NC-. This makes it possible to determine a cerebral A β accumulation state, or to assist the determination.

[0122] A method for analyzing the marker of the present invention includes:

[0123] a measurement step of subjecting a blood sample derived from a test subject to detection of a marker containing APP672-713 (A β 1-42) (SEQ ID NO.: 6) to obtain a measurement level of APP672-713 (A β 1-42) in the blood sample; and

[0124] an evaluation step of determining that an amount of cerebral A β accumulation of the test subject is larger than an amount of cerebral A β accumulation of a person having normal cognitive function NC- who is negative for cerebral A β accumulation, when the measurement level of APP672-713 (A β 1-42) of the test subject is lower than a standard level which is a level of APP672-713 (A β 1-42) of the person having normal cognitive function NC-. This makes it possible to determine a cerebral A β accumulation state, or to assist the determination.

[0125] The term "level of marker" basically means a concentration, but may be other units applied correspondingly to concentration by a person skilled in the art, for example, a detected ion intensity in the mass spectrometry. In the present invention, the concentration of the marker in the living body-derived sample is analyzed by comparing a measurement value and a standard value. For more accurate analysis, it is preferred that the measurement value and the standard value to be compared are values based on the living body-derived samples prepared in the same conditions (pre-treatment condition, storage condition and the like). As the standard level of the marker, a measurement value for a person having normal cognitive function who is negative for cerebral A β accumulation NC- can be used. Alternatively, a concentration standard value established for a person having normal cognitive function who is negative for intracerebral A β accumulation NC- may be used as the standard level of the marker.

[0126] A marker is measured, preferably, by test based on biological molecule specific affinity. The test based on biological molecule specific affinity is a method well known to a person skilled in the art and is not particularly limited, but is preferably an immunoassay. Specific examples of the immunoassay include competitive and non-competitive assays such as western blotting, radioimmunoassay, ELISA (Enzyme-Linked Immunosorbent Assay) (sandwich immunoassay, competitive assay, and direct binding assay are included), immunoprecipitation, precipitation reaction, immunodiffusion, immunoagglutination measurement, complement-binding reaction analysis, immunoradiometric assay, fluorescence immunoassay, and protein A immunoassay. In the immunoassay, an antibody that binds to the marker in a living body-derived sample is detected.

[0127] In the present invention, the measurement of the marker may be performed by using an immunoglobulin

having an antigen binding site capable of recognizing an amyloid precursor protein (APP)-derived peptide, or an antibody-immobilizing carrier prepared by using an immunoglobulin fragment having an antigen binding site capable of recognizing an amyloid precursor protein (APP)-derived peptide. By immunoprecipitation using the antibody-immobilizing carrier, a peptide in the sample can be detected by a mass spectrometer (Immunoprecipitation-Mass Spectrometry: IP-MS).

[0128] As shown in the section of EXAMPLES, a blood sample derived from a test subject is subjected to detection of a marker containing APP672-713 (A β 1-42) (SEQ ID NO.: 6) to obtain a measurement level of APP672-713 (A β 1-42) in the blood sample; and an amount of cerebral A β accumulation of the test subject is determined to be larger than an amount of cerebral AP accumulation of a person having normal cognitive function NC- who is negative for cerebral A β accumulation, when the measurement level of APP672-713 (A β 1-42) of the test subject is lower than a standard level which is a level of APP672-713 (A β 1-42) of the person having normal cognitive function NC-.

EXAMPLES

[0129] Hereinafter, the present invention will be described specifically with reference to examples, but is not limited to these examples. In the following, the amount of a matter indicated by % is based on weight when the matter is a subject, and based on volume when the matter is liquid unless otherwise indicated.

Experimental Example 1

(1) Evaluation of Performance of Alzheimer's Disease Diagnosis by Marker of Present Invention Using Clinical Sample

[0130] Plasma samples of cases classified into groups of NC-, NC+, MCI, and AD were prepared at National Center for Geriatrics and Gerontology.

[0131] NC-: a person who is negative for cerebral A β accumulation, and does not have a cognitive impairment

[0132] NC+: a person who is positive for cerebral A β accumulation, but does not have a cognitive impairment

[0133] MCI: a person who is positive for cerebral A β accumulation, and has a mild cognitive impairment

[0134] AD: a person who is positive for cerebral A β accumulation, and has a cognitive impairment

[0135] PiB+: combined group of the group NC+, the group MCI, and the group AD; that is, a person determined as positive for cerebral A β accumulation regardless of the presence or absence of a cognitive impairment

[0136] In order to determine the positivity or negativity of cerebral A β accumulation, PiB-PET images of the brains of the subjects were acquired. When the PiB accumulation amount of the cerebral cortex is larger than or equivalent to the non-specific PiB accumulation amount of the white matter, the subject was determined as positive. When only non-specific accumulation to the white matter was observed, and little accumulation was observed in the cortex, the subject was determined as negative. The cognitive impairment was determined in conformity with the NIA-AA criteria published in 2011.

[0137] Table 3 shows the feature and the number of cases of each group.

TABLE 3

Outline of Clinical Sample			
Group	n	PiB-PET	Dementia
NC-	22	-	-
NC+	11	+	-
MCI	12	+	+
AD	17	+	++

[0138] When the application to early diagnosis of Alzheimer's disease is considered, it is supposed that a blood marker showing variation in measurement level in the NC+ group, which is determined as positive for cerebral A β accumulation although does not have a cognitive impairment in comparison with the NC- group, is effective for diagnosis. In other words, it is important to find a blood marker having a diagnostic performance capable of discriminating between the positivity and the negativity of cerebral A β accumulation. From this viewpoint, criteria for evaluation were made in the marker analysis whether there is a difference in measurement level between each of other groups and the NC- group.

(2) Preparation of Antibody-Immobilizing Beads

[0139] 250 μ g of anti-A β antibody (6E10: Covance) recognizing the residues 3-8 of A β as an epitope was digested with 1250 μ L of Ficin agarose beads (Thermo) (33% slurry), and 100 μ g of anti-A β antibody (4G8: Covance) recognizing the residues 18-22 of A β as an epitope was digested with 500 ng of lysyl endopeptidase (LysC), and respective digests were separated and collected by size exclusion chromatography. The fractionated sample was examined by reducing and non-reducing SDS-PAGE, and a fraction corresponding to F(ab')₂ was pooled. These F(ab')₂ fractions of 6E10 and 4G8 were respectively reduced by 30 mM cysteamine to obtain F(ab'). Then 5 μ L (amount of beads 150 μ g) of amino magnetic beads (Dynabeads M-270 Amine: Invitrogen) was prepared, and PEG and the beads were covalently bound by causing NHS groups of SM(PEG)₂₄ to react with amino groups bound onto the surface of the beads at room temperature for 30 minutes. To SM(PEG)₂₄ bound to magnetic beads, each 0.25 μ g of 6E10 F(ab') and 4G8 F(ab') were added at the same time, and they were caused to react at room temperature for 2 hours to covalently bind a maleimide group and a thiol group. Finally, 0.4 mM L-cysteine was caused to react at room temperature for 30 minutes to block maleimide groups. The prepared antibody-immobilizing beads were stored at 4° C. before use.

(3) Preparation of Internal Standard Peptide

[0140] As an internal standard peptide, A β 1-38 (referred to as SIL-A β 1-38) that is a stable isotope-labeled AnaSpec (San Jose, Calif., USA) was used. In SIL-A β 1-38, carbon atoms in Phe and Ile are substituted by ¹³C. After dissolving a dry product of SIL-A β 1-38 in 50 mM NaOH, size exclusion chromatography (SEC) was performed by a Prominence HPLC System (Shimadzu Corp, Kyoto, Japan) installed with a COSMOSIL(R) 5Diol-120-II [7.5 mm I.D. $\times$ 600 mm] column (Nacalai Tesque, Kyoto). The settings were as follows: mobile phase: 40 mM Tris-HCl, pH 8.0, flow rate: 1 mL/min, column temperature: 25° C., detection wavelength: 214/280 nm. Part of the collected fractions was subjected to 15-20% Tricine-SDS-PAGE in non-reducing

condition, and bands were stained with Silver Staining kit (Invitrogen, Carlsbad, Calif., USA). The fraction identified as a monomer was diluted with 40 mM Tris-HCl, 150 mM NaCl, pH 8.0 containing 1 mg/mL bovine serum albumin, and then dispensed and stored at -80°C . Part of this was taken out, and the concentration of SIL-A β 1-38 was measured by using a Human Amyloid β (1-38) (FL) Assay Kit (Immuno-Biological Laboratories; Gunma, Japan).

(4) Measurement of Plasma A β and A β -Like Peptides

[0141] For a total of 62 cases, IP-MS was performed by using antibody-immobilizing beads with the use of SIL-A β 1-38 as the internal standard peptide.

[0142] Immunoprecipitation was performed in the following manner.

[0143] Into 250 μL of human plasma, 250 μL of a binding buffer (0.2% (w/v) n-Dodecyl- β -D-maltoside (DDM), 0.2% (w/v) n-Nonyl- β -D-thiomaltoside (NTM), 800 mM GlcNAc, 100 mM Tris-HCl, 300 mM NaCl, pH 7.4) containing 10 pM SIL-A β 1-38 was mixed. The precipitate contained in the plasma sample was centrifugally removed by using Ultrafree-MC, DV 0.65 μm , centrifugal filter devices. 500 μL of Protein G Plus Agarose (50% slurry; Pierce, Rockford, Ill.) was washed once with 400 μL of H_2O , and then washed three times with 400 μL of a washing buffer (0.1% (w/v) DDM, 0.1% (w/v) NTM, 50 mM Tris-HCl, 150 mM NaCl, pH 7.4). Into the resultant Protein G Plus Agarose, the previously obtained plasma sample was mixed, and incubated at 4°C . for 1 hour. After removing the Protein G Plus Agarose from the plasma sample, the plasma sample was mixed into 150 μg of antibody-immobilizing beads that was washed twice with an OTG-glycine buffer (1% n-Octyl- β -D-thiogluconide (OTG), 50 mM glycine, pH 2.8) and three times with 100 μL of a washing buffer, and the mixture was incubated at 4°C . for 1 hour to capture A β and A β -like peptides. Thereafter, the beads were washed once with 500 μL of a washing buffer, four times with 100 μL of a washing buffer, and twice with 20 μL of 50 mM ammonium acetate. Further, after washing once with 20 μL of H_2O , A β and A β -like peptides captured on the antibody-immobilizing beads were eluted with 2.5 μL of 70% acetonitrile containing 5 mM hydrochloric acid. Each 0.5 μL of the eluate was dropped into 4 wells on a μFocus MALDI PlateTM 900 μm . After mixing 0.5 μL of a matrix solution (0.5 mg/mL CHCA, 0.2% (w/v) MDPNA), measurement was performed by Linear TOF MS.

[0144] As a measurement value of A β and A β -like peptides in plasma, an averaged value of peak intensity ratios of each of A β and A β -like peptides to the internal standard peptide (SIL-A β 1-38) obtained in 4 wells measured by Linear TOF was used. For correcting a signal variation in MS peak intensity, the following criteria were provided. Four mass spectra are acquired in one time immunoprecipitation, and thus four peak intensity ratios are obtained for one peptide. A peak intensity ratio that is out of 0.7 to 1.3 times the median of the four peak intensity ratios in a certain peptide is regarded as an outlier, and is not used in the data processing for averaging. While the number of data of peak intensity ratio used for averaging of a certain peptide is four at maximum, the number of data decreases when the peak intensity is less than the detection limit ($S/N < 3$), or an outlier arises. When the number of data of peak intensity

ratio used in averaging is less than three, the intensity ratio at the peak is defined as not detectable (N/D) in the immunoprecipitation at that time.

[0145] FIG. 2 is a box-and-whisker plot showing the intensity ratio of APP672-713 (A β 1-42) to the internal standard SIL-A β 1-38 in each group for APP672-713 (A β 1-42). FIG. 3(A), FIG. 3(B), FIG. 4(C), and FIG. 4(D) each show an ROC curve of each group (NC+, MCI, AD, PiB+) versus NC- group for APP672-713 (A β 1-42)/SIL-A β 1-38.

[0146] In each box-and-whisker plot, the range indicated by the box in each group represents the intensity ratio contribution range (quartile range) of the samples whose concentration is rated between 25 to 75% of all specimens, and the horizontal lines shown above and below the box respectively indicate the maximum value and the minimum value of the samples within the range from the upper end and the lower end of the box to 1.5 times the quartile range, and the horizontal bar in the box indicates the median of the intensity ratio.

[0147] A statistical significant difference between each of other groups and NC- group was tested by using a Dunnett's test, and it was determined that there is a significant difference when $P < 0.05$. The detection limit or lower was set at 0. As a result, nine kinds of A β and A β -like peptides were detected in 60% or more cases, and among these, in APP672-713 (A β 1-42), there was a statistical significant difference in NC+, MCI, and AD in comparison with NC- (FIG. 2). For evaluating the diagnostic performance of APP672-713 (A β 1-42), ROC curves of NC+, MCI, AD, PiB+ groups versus NC- group were prepared. The areas under ROC curves (AUC) were NC- vs NC+=0.789, NC- vs MCI=0.746, NC- vs AD=0.864, NC- vs PiB+=0.808, and showed relatively high values (FIG. 3(A), FIG. 3(B), FIG. 4(C), and FIG. 4(D)).

[0148] As shown in FIG. 2, for APP672-713 (A β 1-42), the intensity ratio APP672-713 (A β 1-42)/SIL-A β 1-38 decreased in NC+, MCI, and AD with a statistical significant difference in comparison with NC-. As shown in FIG. 3(A), FIG. 3(B), FIG. 4(C), and FIG. 4(D), the AUC: NC- vs NC+=0.789, NC- vs MCI=0.746, NC- vs AD=0.864 revealed that APP672-713 (A β 1-42) had a relatively high ability capable of discrimination between NC- and NC+, NC- and MCI, and NC- and AD. The NC- vs PiB+=0.808 revealed that the performance capable of detecting the subject who is positive for cerebral A β accumulation is relatively high. Accordingly, this suggests that APP672-713 (A β 1-42) is a blood marker capable of estimating a cerebral A β accumulation state, and thus it has the possibility capable of being used for assisting diagnosis of Alzheimer's disease.

(5) More Specific Analysis

[0149] For observing a clearer difference between each group (NC+, MCI, AD) and NC-, the following investigation was further performed.

[0150] FIG. 5 is a box-and-whisker plot showing the intensity ratio of APP672-709 (A β 1-38) to APP672-713 (A β 1-42) in each group. FIG. 6(A), FIG. 6(B), FIG. 7(C), and FIG. 7(D) each show an ROC curve of each group (NC+, MCI, AD, PiB+) versus NC- group for APP672-709 (A β 1-38)/APP672-713 (A β 1-42).

[0151] FIG. 8 is a box-and-whisker plot showing the intensity ratio of APP674-711 (A β 3-40) to APP672-713 (A β 1-42) in each group. FIG. 9(A), FIG. 9(B), FIG. 10(C), and FIG. 10(D) each show an ROC curve of each group

(NC+, MCI, AD, PiB+) versus NC- group for APP674-711 (A β 3-40)/APP672-713 (A β 1-42).

[0152] FIG. 11 is a box-and-whisker plot showing the intensity ratio of APP672-710 (A β 1-39) to APP672-713 (A β 1-42) in each group. FIG. 12(A), FIG. 12(B), FIG. 13(C), and FIG. 13(D) each show an ROC curve of each group (NC+, MCI, AD, PiB+) versus NC- group for APP672-710 (A β 1-39)/APP672-713 (A β 1-42).

[0153] FIG. 14 is a box-and-whisker plot showing the intensity ratio of APP672-711 (A β 1-40) to APP672-713 (A β 1-42) in each group. FIG. 15(A), FIG. 15(B), FIG. 16(C), and FIG. 16(D) each show an ROC curve of each group (NC+, MCI, AD, PiB+) versus NC- group for APP672-711 (A β 1-40)/APP672-713 (A β 1-42).

[0154] FIG. 17 is a box-and-whisker plot showing the intensity ratio of OxAPP672-711 (OxA β 1-40) to APP672-713 (A β 1-42) in each group. FIG. 18(A), FIG. 18(B), FIG. 19(C), and FIG. 19(D) each show an ROC curve of each group (NC+, MCI, AD, PiB+) versus NC- group for OxAPP672-711 (OxA β 1-40)/APP672-713 (A β 1-42).

[0155] FIG. 20 is a box-and-whisker plot showing the intensity ratio of APP669-711 to APP672-713 (A β 1-42) in each group. FIG. 21(A), FIG. 21(B), FIG. 22(C), and FIG. 22(D) each show an ROC curve of each group (NC+, MCI, AD, PiB+) versus NC- group for APP669-711/APP672-713 (A β 1-42).

[0156] Among the nine kinds of A β and A β -like peptides detected in 60% or more cases, respective peak intensities of eight kinds other than APP672-713 (A β 1-42) were divided by the peak intensity of APP672-713 (A β 1-42), and the resultant values (ratios) were comparatively analyzed. As a result, in APP672-709 (A β 1-38)/APP672-713 (A β 1-42), APP674-711 (A β 3-40)/APP672-713 (A β 1-42), APP672-710 (A β 1-39)/APP672-713 (A β 1-42), APP672-711 (A β 1-40)/APP672-713 (A β 1-42), OxAPP672-711 (OxA β 1-40)/APP672-713 (A β 1-42), and APP669-711/APP672-713 (A β 1-42), a statistically significant increase was observed in NC+, MCI, and AD in comparison with NC- (FIGS. 5, 8, 11, 14, 17, and 20). It was demonstrated that APP669-711/APP672-713 (A β 1-42) in particular has a strong tendency to rise as Alzheimer's disease advances (FIG. 20).

[0157] For evaluating diagnostic performances of APP672-709 (A β 1-38)/APP672-713 (A β 1-42), APP674-711 (A β 3-40)/APP672-713 (A β 1-42), APP672-710 (A β 1-39)/APP672-713 (A β 1-42), APP672-711 (A β 1-40)/APP672-713 (A β 1-42), OxAPP672-711 (OxA β 1-40)/APP672-713 (A β 1-42), and APP669-711/APP672-713 (A β 1-42), ROC curves of NC+, MCI, AD, and PiB+ groups versus NC- group were prepared. As a result, in all these six kinds of ratios, high AUC was observed, revealing that the ability capable of discrimination between NC- and NC+, NC- and MCI, and NC- and AD is high, and the performance capable of detecting the subject who is positive for cerebral A β accumulation is high [FIG. 6(A), FIG. 6(B), FIG. 7(C), and FIG. 7(D); FIG. 9(A), FIG. 9(B), FIG. 10(C), and FIG. 10(D); FIG. 12(A), FIG. 12(B), FIG. 13(C), and FIG. 13(D); FIG. 15(A), FIG. 15(B), FIG. 16(C), and FIG. 16(D); FIG. 18(A), FIG. 18(B), FIG. 19(C), and FIG. 19(D); FIG. 21(A), FIG. 21(B), FIG. 22(C), and FIG. 22(D)].

[0158] Referring to FIG. 21(A), FIG. 21(B), FIG. 22(C), and FIG. 22(D), AUC of NC+, MCI, AD versus NC- group was 0.930 or more in the ROC curve of APP669-711/APP672-713 (A β 1-42) in particular, revealing that the ability capable of discrimination between NC- and NC+, NC-

and MCI, and NC- and AD is very high. NC- vs PiB+=0.969 also revealed that the performance capable of detecting the subject who is positive for cerebral A β accumulation is very high. Accordingly, this suggests that these six kinds of ratios are blood markers capable of estimating a cerebral A β accumulation state, and thus they have a possibility capable of being used for assisting diagnosis of Alzheimer's disease.

[0159] Next, for APP672-709 (A β 1-38)/APP672-713 (A β 1-42), APP674-711 (A β 3-40)/APP672-713 (A β 1-42), APP672-710 (A β 1-39)/APP672-713 (A β 1-42), APP672-711 (A β 1-40)/APP672-713 (A β 1-42), OxAPP672-711 (OxA β 1-40)/APP672-713 (A β 1-42), and APP669-711/APP672-713 (A β 1-42) in which an statistically significant increase was observed in NC+, MCI, AD in comparison with NC-, discrimination of Alzheimer's disease was evaluated by setting a cut-off point. In each ROC curve in FIGS. 6, 7, 9, 10, 12, 13, 15, 16, 18, 19, 21, and 22, "each peptide/APP672-713 (A β 1-42)" showing the highest value of "sensitivity-(1-specificity)" was set as a cut-off point. The set cut-off point, Specificity in that case, and Sensitivity, Positive Predictive Value (PPV), Negative Predictive Value (NPV), Accuracy of each group (NC+, MCI, AD, PiB+) versus NC- group are shown in Table 4.

TABLE 4

Evaluation of Diagnostic Performance				
	Sensitivity	Positive Predictive Value, PPV	Negative Predictive Value, NPV	Accuracy
APP672-709(A β 1-38)/APP672-713(A β 1-42) Cut-off point = 7.895, Specificity = 0.682				
NC- vs NC+	1.000	0.611	1.000	0.788
NC- vs MCI	0.833	0.588	0.882	0.735
NC- vs AD	0.765	0.650	0.789	0.718
NC- vs PiB+	0.850	0.829	0.714	0.790
APP674-711(A β 3-40)/APP672-713(A β 1-42) Cut-off point = 1.181, Specificity = 0.909				
NC- vs NC+	0.900	0.818	0.952	0.906
NC- vs MCI	0.833	0.833	0.909	0.882
NC- vs AD	1.000	0.895	1.000	0.949
NC- vs PiB+	0.923	0.947	0.870	0.918
APP672-710(A β 1-39)/APP672-713(A β 1-42) Cut-off point = 1.932, Specificity = 0.864				
NC- vs NC+	0.909	0.769	0.950	0.879
NC- vs MCI	0.833	0.769	0.905	0.853
NC- vs AD	0.941	0.842	0.950	0.897
NC- vs PiB+	0.900	0.923	0.826	0.887
APP672-711(A β 1-40)/APP672-713(A β 1-42) Cut-off point = 111.7, Specificity = 0.773				
NC- vs NC+	0.909	0.667	0.944	0.818
NC- vs MCI	0.750	0.643	0.850	0.765
NC- vs AD	0.647	0.688	0.739	0.718
NC- vs PiB+	0.750	0.857	0.630	0.758
OxAPP672-711(OxA β 1-40)/APP672-713(A β 1-42) Cut-off point = 3.431, Specificity = 0.864				
NC- vs NC+	0.818	0.750	0.905	0.848
NC- vs MCI	0.727	0.727	0.864	0.818
NC- vs AD	0.824	0.824	0.864	0.846
NC- vs PiB+	0.795	0.912	0.704	0.820
APP669-711/APP672-713(A β 1-42) Cut-off point = 0.914, Specificity = 0.955				
NC- vs NC+	0.818	0.900	0.913	0.909
NC- vs MCI	0.917	0.917	0.955	0.941

TABLE 4-continued

Evaluation of Diagnostic Performance				
	Sensitivity	Positive Predictive Value, PPV	Negative Predictive Value, NPV	Accuracy
NC- vs AD	1.000	0.944	1.000	0.974
NC- vs PiB+	0.925	0.974	0.875	0.935

[0160] In Table 4,

[0161] Cut-off point: cut-off point,

Positive Predictive Value (PPV)=(number of true positive)/(number of true positive+number of false positive),

Negative Predictive Value (NPV)=(number of true negative)/(number of true negative+number of false negative),

and

Accuracy=(number of true positive+number of true negative)/number of total cases.

[0162] APP669-711/APP672-713 ($A\beta$ 1-42) shows very high numerical values in all of Specificity, Sensitivity, Positive Predictive Value (PPV), Negative Predictive Value (NPV), and Accuracy, revealing that it has high ability capable of discriminating between NC- and NC+, MCI, AD, and PiB+, and in particular, it is effective for determination of positivity of cerebral $A\beta$ accumulation. The very high ability capable of discrimination between NC-, and NC+, MCI, and AD also indicates the applicability to assist diagnosis of Alzheimer's disease. Regarding other five kinds of ratios, they show high values in Specificity, Sensitivity, Positive Predictive Value (PPV), Negative Predictive Value (NPV), or Accuracy, revealing that they have high ability capable of discrimination between NC- and NC+, MCI, AD, and PiB+, and in particular, they show the possibility effective for determination of positivity of cerebral $A\beta$ accumulation. The high ability capable of discrimination between NC- and NC+, MCI, and AD also indicates the applicability to assist diagnosis of Alzheimer's disease.

(6) Correlation Analysis with PiB Measurement Value

[0163] Cerebral $A\beta$ accumulation is an important pathological index for Alzheimer's disease, and excessive accumulation of $A\beta$ is known to begin far in advance of exteriorization of dementia. PiB is a radioactive agent that specifically binds with $A\beta$ aggregate, and by measuring accumulation of PiB with the use of PET, it is possible to form an image of the cerebral $A\beta$ accumulation. In order to investigate whether APP669-711/APP672-713 ($A\beta$ 1-42) in plasma, which shows the ability capable of determining with high accuracy a subject who is positive for cerebral $A\beta$ accumulation, reflects a cerebral $A\beta$ accumulation state, correlation between APP669-711/APP672-713 ($A\beta$ 1-42) and PiB accumulation mean value of cortical region (mc-SUVR: mean cortical Standard Uptake Value Ratio) was analyzed.

[0164] FIG. 23 is a scatter diagram for a total of 62 cases, in which the horizontal axis represents the PiB accumulation mean value (mcSUVR) and the vertical axis represents APP669-711/APP672-713 ($A\beta$ 1-42) ratio.

[0165] The PiB accumulation mean value was measured by quantifying PiB accumulation of the cortex, and determining accumulation ratio of the cerebrum on the basis of the cerebellum. Correlation between APP669-711/APP672-713 ($A\beta$ 1-42) ratio obtained by IP-MS and PiB accumulation mean value (mcSUVR) obtained by PET was evaluated with Pearson product-moment correlation coefficient. As a result, the correlation coefficient (r) was 0.687, $p < 0.001$ (FIG. 23).

[0166] It was revealed that there is a significantly strong correlation between APP669-711/APP672-713 ($A\beta$ 1-42) ratio and PiB accumulation mean value (mcSUVR). This means that APP669-711/APP672-713 ($A\beta$ 1-42) ratio in plasma reflects a cerebral $A\beta$ accumulation state, and indicates that it has the possibility capable of being used as a blood marker for determining a cerebral $A\beta$ accumulation state.

[0167] The results demonstrated above indicate that the marker of the present invention is useful as a blood marker for determining a cerebral $A\beta$ accumulation state. This also has indicated the applicability to assist diagnosis of Alzheimer's disease and to presymptomatically diagnose Alzheimer's disease.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 7

<210> SEQ ID NO 1

<211> LENGTH: 38

<212> TYPE: PRT

<213> ORGANISM: Homo Sapiens

<400> SEQUENCE: 1

Asp Ala Glu Phe Arg His Asp Ser Gly Tyr Glu Val His His Gln Lys
1 5 10 15

Leu Val Phe Phe Ala Glu Asp Val Gly Ser Asn Lys Gly Ala Ile Ile
20 25 30

Gly Leu Met Val Gly Gly
35

<210> SEQ ID NO 2

-continued

<211> LENGTH: 38
<212> TYPE: PRT
<213> ORGANISM: Homo Sapiens

<400> SEQUENCE: 2

Glu Phe Arg His Asp Ser Gly Tyr Glu Val His His Gln Lys Leu Val
1 5 10 15

Phe Phe Ala Glu Asp Val Gly Ser Asn Lys Gly Ala Ile Ile Gly Leu
20 25 30

Met Val Gly Gly Val Val
35

<210> SEQ ID NO 3
<211> LENGTH: 39
<212> TYPE: PRT
<213> ORGANISM: Homo Sapiens

<400> SEQUENCE: 3

Asp Ala Glu Phe Arg His Asp Ser Gly Tyr Glu Val His His Gln Lys
1 5 10 15

Leu Val Phe Phe Ala Glu Asp Val Gly Ser Asn Lys Gly Ala Ile Ile
20 25 30

Gly Leu Met Val Gly Gly Val
35

<210> SEQ ID NO 4
<211> LENGTH: 40
<212> TYPE: PRT
<213> ORGANISM: Homo Sapiens

<400> SEQUENCE: 4

Asp Ala Glu Phe Arg His Asp Ser Gly Tyr Glu Val His His Gln Lys
1 5 10 15

Leu Val Phe Phe Ala Glu Asp Val Gly Ser Asn Lys Gly Ala Ile Ile
20 25 30

Gly Leu Met Val Gly Gly Val Val
35 40

<210> SEQ ID NO 5
<211> LENGTH: 40
<212> TYPE: PRT
<213> ORGANISM: Homo Sapiens

<400> SEQUENCE: 5

Asp Ala Glu Phe Arg His Asp Ser Gly Tyr Glu Val His His Gln Lys
1 5 10 15

Leu Val Phe Phe Ala Glu Asp Val Gly Ser Asn Lys Gly Ala Ile Ile
20 25 30

Gly Leu Met Val Gly Gly Val Val
35 40

<210> SEQ ID NO 6
<211> LENGTH: 42
<212> TYPE: PRT
<213> ORGANISM: Homo Sapiens

<400> SEQUENCE: 6

Asp Ala Glu Phe Arg His Asp Ser Gly Tyr Glu Val His His Gln Lys
1 5 10 15

-continued

Leu Val Phe Phe Ala Glu Asp Val Gly Ser Asn Lys Gly Ala Ile Ile
 20 25 30

Gly Leu Met Val Gly Gly Val Val Ile Ala
 35 40

<210> SEQ ID NO 7
 <211> LENGTH: 43
 <212> TYPE: PRT
 <213> ORGANISM: Homo Sapiens

<400> SEQUENCE: 7

Val Lys Met Asp Ala Glu Phe Arg His Asp Ser Gly Tyr Glu Val His
 1 5 10 15

His Gln Lys Leu Val Phe Phe Ala Glu Asp Val Gly Ser Asn Lys Gly
 20 25 30

Ala Ile Ile Gly Leu Met Val Gly Gly Val Val
 35 40

1. An analytical method for determining a cerebral A β accumulation state, the method comprising:

a measurement step of subjecting a living body-derived sample derived from a test subject to detection of a marker containing:

APP672-713 (A β 1-42) (SEQ ID NO.: 6), and

at least one selected from the group consisting of APP669-711 (SEQ ID NO.: 7), APP672-709 (A β 1-38) (SEQ ID NO.: 1), APP674-711 (A β 3-40) (SEQ ID NO.: 2), APP672-710 (A β 1-39) (SEQ ID NO.: 3), APP672-711 (A β 1-40) (SEQ ID NO.: 4), and OxAPP672-711 (OxAPP672-711) (SEQ ID NO.: 5),

to obtain measurement levels of:

APP672-713 (A β 1-42), and

the at least one selected from the group consisting of APP669-711, APP672-709 (A β 1-38), APP674-711 (A β 3-40), APP672-710 (A β 1-39), APP672-711 (A β 1-40), and OxAPP672-711 (OxAPP672-711) in the living body-derived sample;

a calculation step of calculating at least one ratio selected from the group consisting of:

a ratio of APP669-711 level to APP672-713 (A β 1-42) level: APP669-711/APP672-713 (A β 1-42);

a ratio of APP672-709 (A β 1-38) level to APP672-713 (A β 1-42) level: APP672-709 (A β 1-38)/APP672-713 (A β 1-42);

a ratio of APP674-711 (A β 3-40) level to APP672-713 (A β 1-42) level: APP674-711 (A β 3-40)/APP672-713 (A β 1-42);

a ratio of APP672-710 (A β 1-39) level to APP672-713 (A β 1-42) level: APP672-710 (A β 1-39)/APP672-713 (A β 1-42);

a ratio of APP672-711 (A β 1-40) level to APP672-713 (A β 1-42) level: APP672-711 (A β 1-40)/APP672-713 (A β 1-42); and

a ratio of OxAPP672-711 (OxAPP672-711) level to APP672-713 (A β 1-42) level: OxAPP672-711 (OxAPP672-711)/APP672-713 (A β 1-42); and

an evaluation step of determining that an amount of cerebral A β accumulation of the test subject is larger than an amount of cerebral A β accumulation of a

person having normal cognitive function NC- who is negative for cerebral A β accumulation, when each of the ratios of the test subject is higher than each of standard levels which are ratios of the person having normal cognitive function NC-.

2. The analytical method for determining a cerebral A β accumulation state according to claim 1, wherein the living body-derived sample is selected from the group consisting of blood, cerebrospinal fluid, urine, body secreting fluid, feces, saliva, and sputum.

3. A marker for determining a cerebral A β accumulation state, the marker being selected from the group consisting of:

a combination of APP669-711 (SEQ ID NO.: 7) and APP672-713 (A β 1-42) (SEQ ID NO.: 6),

a combination of APP672-709 (A β 1-38) (SEQ ID NO.: 1) and APP672-713 (A β 1-42) (SEQ ID NO.: 6),

a combination of APP674-711 (A β 3-40) (SEQ ID NO.: 2) and APP672-713 (A β 1-42) (SEQ ID NO.: 6),

a combination of APP672-710 (A β 1-39) (SEQ ID NO.: 3) and APP672-713 (A β 1-42) (SEQ ID NO.: 6),

a combination of APP672-711 (A β 1-40) (SEQ ID NO.: 4) and APP672-713 (A β 1-42) (SEQ ID NO.: 6), and

a combination of OxAPP672-711 (OxAPP672-711) (SEQ ID NO.: 5) and APP672-713 (A β 1-42) (SEQ ID NO.: 6), in a living body-derived sample.

4. An analytical method for determining a cerebral A β accumulation state, the method comprising:

a measurement step of subjecting a blood sample derived from a test subject to detection of a marker containing APP672-713 (A β 1-42) (SEQ ID NO.: 6), to obtain a measurement level of APP672-713 (A β 1-42) in the blood sample; and

an evaluation step of determining that an amount of cerebral A β accumulation of the test subject is larger than an amount of cerebral A β accumulation of a person having normal cognitive function NC- who is negative for cerebral A β accumulation, when the measurement level of APP672-713 (A β 1-42) of the test

subject is lower than a standard level which is a level of APP672-713 (A β 1-42) of the person having normal cognitive function NC-.

5. A method for determining efficacy of a medical intervention regarding a cerebral A β accumulation state, the method comprising, before and after a medical intervention performed for a test subject:

a measurement step of subjecting a living body-derived sample derived from the test subject to detection of a marker containing:

APP672-713 (A β 1-42) (SEQ ID NO.: 6), and

at least one selected from the group consisting of APP669-711 (SEQ ID NO.: 7), APP672-709 (A β 1-38) (SEQ ID NO.: 1), APP674-711 (A β 3-40) (SEQ ID NO.: 2), APP672-710 (A β 1-39) (SEQ ID NO.: 3), APP672-711 (A β 1-40) (SEQ ID NO.: 4), and OxAPP672-711 (OxA β 1-40) (SEQ ID NO.: 5),

to obtain measurement levels of:

APP672-713 (A β 1-42), and

the at least one selected from the group consisting of APP669-711, APP672-709 (A β 1-38), APP674-711 (A β 3-40), APP672-710 (A β 1-39), APP672-711 (A β 1-40), and OxAPP672-711 (OxA β 1-40) in the living body-derived sample; and

a calculation step of calculating at least one ratio selected from the group consisting of:

a ratio of APP669-711 level to APP672-713 (A β 1-42) level: APP669-711/APP672-713 (A β 1-42);

a ratio of APP672-709 (A β 1-38) level to APP672-713 (A β 1-42) level: APP672-709 (A β 1-38)/APP672-713 (A β 1-42);

a ratio of APP674-711 (A β 3-40) level to APP672-713 (A β 1-42) level: APP674-711 (A β 3-40)/APP672-713 (A β 1-42);

a ratio of APP672-710 (A β 1-39) level to APP672-713 (A β 1-42) level: APP672-710 (A β 1-39)/APP672-713 (A β 1-42);

a ratio of APP672-711 (A β 1-40) level to APP672-713 (A β 1-42) level: APP672-711 (A β 1-40)/APP672-713 (A β 1-42); and

a ratio of OxAPP672-711 (OxA β 1-40) level to APP672-713 (A β 1-42) level: OxAPP672-711 (OxA β 1-40)/APP672-713 (A β 1-42); and

comparing each of the ratios of the test subject before the medical intervention and each of the ratios of the test subject after the medical intervention to determine efficacy of the medical intervention regarding a cerebral A β accumulation state.

* * * * *

专利名称(译)	用于评估脑内淀粉样蛋白 β 肽积累的替代生物标志物及其分析方法		
公开(公告)号	US20170184573A1	公开(公告)日	2017-06-29
申请号	US15/313376	申请日	2015-05-19
[标]申请(专利权)人(译)	株式会社岛津制作所 NAT癌症CENT		
申请(专利权)人(译)	岛津制作所 国家中心的老年医学和老年病学		
当前申请(专利权)人(译)	岛津制作所 国家中心的老年医学和老年病学		
[标]发明人	KANEKO NAOKI YANAGISAWA KATSUHIKO		
发明人	KANEKO, NAOKI YANAGISAWA, KATSUHIKO		
IPC分类号	G01N33/53		
CPC分类号	G01N33/5308 G01N2333/4709 G01N33/53 G01N33/6896 G01N2800/52 G01N2800/56		
优先权	2014106560 2014-05-22 JP		
外部链接	Espacenet USPTO		

摘要(译)

本发明提供了使用来源于活体的样品中的淀粉样蛋白前体蛋白 (APP) 来源的肽作为指标来评价脑 $A\beta$ 积聚状态的生物标志物及其分析方法。测量衍生自测试受试者的活体衍生样品, 并且测量选自APP669-711 / APP672-713 ($A\beta$ 1-42) 的至少一种比例; APP672-709 ($A\beta$ 1-38) / APP672-713 ($A\beta$ 1-42); APP674-711 ($A\beta$ 3-40) / APP672-713 ($A\beta$ 1-42); APP672-710 ($A\beta$ 1-39) / APP672-713 ($A\beta$ 1-42); APP672-711 ($A\beta$ 1-40) / APP672-713 ($A\beta$ 1-42); 计算APP672-711 (Ox $A\beta$ 1-40) / APP672-713 ($A\beta$ 1-42), 并且当测试受试者的每个比率高于具有正常认知功能NC-的人的比率的每个标准水平时对于脑 $A\beta$ 积聚为阴性, 测试受试者的脑 $A\beta$ 的累积量被确定为大于具有正常认知功能NC-的人的脑 $A\beta$ 的累积量。

