



US 20040229804A1

(19) **United States**

(12) **Patent Application Publication** (10) **Pub. No.: US 2004/0229804 A1**
Jay (43) **Pub. Date: Nov. 18, 2004**

(54) **TRIBONECTINS**

Related U.S. Application Data

(75) Inventor: **Gregory D. Jay**, Norfolk, MA (US)

(63) Continuation of application No. 09/298,970, filed on Apr. 23, 1999, now Pat. No. 6,743,774.

Correspondence Address:

**MINTZ, LEVIN, COHN, FERRIS, GLOVSKY
AND POPEO, P.C.
ONE FINANCIAL CENTER
BOSTON, MA 02111 (US)**

Publication Classification

(51) **Int. Cl.⁷** **A61K 38/17; C07K 14/47**

(52) **U.S. Cl.** **514/12; 530/350**

(73) Assignee: **Rhode Island Hospital**, Providence, RI

(57) **ABSTRACT**

(21) Appl. No.: **10/858,595**

The invention features a tribonectin and a method of tribo-supplementation carried out by administering tribonectins directly to an injured or arthritic joint.

(22) Filed: **Jun. 1, 2004**

TRIBONECTINS

BACKGROUND OF THE INVENTION

[0001] The invention relates to lubrication of mammalian joints.

[0002] Osteoarthritis (OA) is the one of the most common form of joint disease. Factors which contribute to the development of OA include a family history of OA, previous damage to the joint through injury or surgery, and age of the joint, i.e., "wear and tear" of the articulating surfaces of the joint. OA is very common in older age groups, but can affect children as well.

[0003] Current treatment is directed to relieving pain and other symptoms of OA, e.g., by administering analgesics and anti-inflammatory drugs. Other therapeutic approaches include viscosupplementation by administering hyaluronic acid and derivatives thereof to joint tissue to increase the viscosity of synovial fluid.

SUMMARY OF THE INVENTION

[0004] The invention features a novel treatment for osteoarthritis and other degenerative joint diseases by tribosupplementation. Tribosupplementation is carried out by administering lubricating polypeptides directly to the injured or arthritic joint. Unlike viscosupplementation, tribosupplementation does not substantially increase the viscosity of the solution, e.g., synovial fluid, to which it is added. The viscosity of a solution to which a tribonectin is added increases no more than 10%, preferably no more than 5%, more preferably no more than 2%, more preferably no more than 1%. Most preferably, the viscosity of the solution to which a tribonectin is added is unaltered or decreases.

[0005] Accordingly, the invention provides a tribonectin. A tribonectin is an artificial boundary lubricant which contains at least one repeat of an amino acid sequence which is at least 50% identical to KEPAPTT (SEQ ID NO:3). A tribonectin is formulated for administration to a mammalian joint. Preferably, the tribonectin is a recombinant or chemically-synthesized lubricating polypeptide. For example, a tribonectin includes a substantially pure polypeptide the amino acid sequence of which includes at least one but less than 76 subunits. Each subunit contains at least 7 amino acids (and typically, 10 or fewer amino acids). The amino acid sequence of each subunit is at least 50% identical to SEQ ID NO:3, and a non-identical amino acid in the reference sequence is a conservative amino acid substitution. For example, one or both of the threonine residues are substituted with a serine residue. Preferably, the amino acid sequence of the subunit is identical to SEQ ID NO:3. The tribonectin may also contain one or more repeats of the amino acid sequence XXTTX (SEQ ID NO:4). Polypeptides or other compounds described herein are said to be "substantially pure" when they are within preparations that are at least 60% by weight (dry weight) the compound of interest. Preferably, the preparation is at least 75%, more preferably at least 90%, and most preferably at least 99%, by weight the compound of interest. Purity can be measured by any appropriate standard method, for example, by column chromatography, polyacrylamide gel electrophoresis, or HPLC analysis.

[0006] Where a particular polypeptide or nucleic acid molecule is said to have a specific percent identity to a reference polypeptide or nucleic acid molecule of a defined length, the percent identity is relative to the reference

polypeptide or nucleic acid molecule. Thus, a peptide that is 50% identical to a reference polypeptide that is 100 amino acids long can be a 50 amino acid polypeptide that is completely identical to a 50 amino acid long portion of the reference polypeptide. It can also be a 100 amino acid long polypeptide which is 50% identical to the reference polypeptide over its entire length.

[0007] A polypeptide or nucleic acid molecule which is "substantially identical" to a given reference polypeptide or nucleic acid molecule is a polypeptide or nucleic acid molecule having a sequence that has at least 85%, preferably 90%, and more preferably 95%, 98%, 99% or more identity to the sequence of the given reference polypeptide sequence or nucleic acid molecule. "Identity" has an art-recognized meaning and is calculated using well known published techniques, e.g., Computational Molecular Biology, 1988, Lesk A. M., ed., Oxford University Press, New York; Biocomputing: Informatics and Genome Projects, 1993, Smith, D. W., ed., Academic Press, New York; Computer Analysis of Sequence Data, Part I, 1994, Griffin, A. M. and Griffin, H. G., eds., Humana Press, New Jersey; Sequence Analysis in Molecular Biology, 1987, Heinje, G., Academic Press, New York; and Sequence Analysis Primer, 1991, Gribskov, M. and Devereux, J., eds., Stockton Press, New York). While there exist a number of methods to measure identity between two polynucleotide or polypeptide sequences, the term "identity" is well known to skilled artisans and has a definite meaning with respect to a given specified method. Sequence identity described herein is measured using the Sequence Analysis Software Package of the Genetics Computer Group (GCS), University of Wisconsin Biotechnology Center, 1710 University Avenue, Madison, Wis. 53705, with the default parameters as specified therein.

[0008] A tribonectin is characterized as reducing the coefficient of friction (μ) between bearing surfaces. For example, reduction of friction is measured in vitro by detecting a reduction in friction in a friction apparatus using latex:glass bearings. Reduction of friction is also measured in vivo, e.g., by measuring reduction of patient pain. Tribonectins of the invention are lubricating compositions. Polypeptides that have at least 50% (but less than 100%) amino acid sequence identity to a reference sequence are tested for lubricating function by measuring a reduction in the μ between bearing surfaces.

[0009] A tribonectin includes an O-linked oligosaccharide, e.g., an N-acetylgalactosamine and galactose in the form $\beta(1-3)\text{Gal-GalNAC}$. For example, KEPAPTT (SEQ ID NO:3) and XXTTTX (SEQ ID NO:4) repeat domains are glycosylated by $\beta(1-3)\text{Gal-GalNAC}$ (which may at times be capped with NeuAc in the form of $\beta(1-3)\text{Gal-GalNAC-NeuAc}$). The term "glycosylated" with respect to a polypeptide means that a carbohydrate moiety is present at one or more sites of the polypeptide molecule. For example, at least 10%, preferably at least 20%, more preferably at least 30%, and most preferably at least 40% of the tribonectin is glycosylated. Up to 50% or more of the tribonectin can be glycosylated. Percent glycosylation is determined by weight.

[0010] A tribonectin polypeptide can contain a substantially pure fragment of megakaryocyte stimulating factor (MSF). For example, the molecular weight of a substantially pure tribonectin having an amino acid sequence of a naturally-occurring tribonectin is in the range of 220-280 kDa. Preferably, the apparent molecular weight of a tribonectin is

less than 230 kDa, more preferably less than 250 kDa, and most preferably less than 280 kDa. A protein or polypeptide fragment is defined as a polypeptide which has an amino acid sequence that is identical to part, but not all, of the amino acid sequence of a naturally-occurring protein or polypeptide from which it is derived, e.g., MSF. The tribonectin may contain a polypeptide, the amino acid sequence of which is at least 50% identical to the sequence of residues 200-1140, inclusive, of SEQ ID NO:1, e.g., it contains the amino acid sequence of residues 200-1140, inclusive, of SEQ ID NO:1. In another example, the polypeptide contains an amino acid sequence that is at least 50% identical to the sequence of residues 200-1167, inclusive, of SEQ ID NO:1, e.g., one having the amino acid sequence identical to residues 200-1167, inclusive, of SEQ ID NO:1. The polypeptide contains an amino acid sequence that is at least 50% identical to the sequence of residues 200-1212, inclusive, of SEQ ID NO:1, e.g., the amino acid sequence of residues 200-1212, inclusive, of SEQ ID NO:1, or the polypeptide contains an amino acid sequence that is at least 50% identical to the sequence of residues 200-1263, inclusive, of SEQ ID NO:1, e.g., an amino acid sequence identical to residues 200-1263, inclusive, of SEQ ID NO:1. Preferably, the sequence of the polypeptide lacks the amino acid sequence of residues 1-24, inclusive, of SEQ ID NO:1 and/or the amino acid sequence of residues 67-104, inclusive of SEQ ID NO:1.

[0011] The invention also features an isolated nucleic acid molecule encoding a tribonectin. For example, the nucleic acid includes a sequence that is at least 50% identical to nucleotides 631-3453, inclusive, of SEQ ID NO:2. Preferably, the nucleic acid encodes a polypeptide with the amino acid sequence of residues 200-1140 of SEQ ID NO:1. For example, the nucleic acid has a nucleotide sequence identical to that of nucleotides 631-3453, inclusive, of SEQ ID NO:2, or a degenerate variant thereof. An isolated nucleic acid molecule is a nucleic acid molecule that is separated from the 5' and 3' coding sequences or non-coding sequences with which it is immediately contiguous in the naturally occurring genome of an organism. Isolated nucleic acid molecules include nucleic acid molecules which are not naturally occurring, e.g., nucleic acid molecules created by recombinant DNA techniques. For example, the nucleic acid of the invention includes nucleotides 631-3453, inclusive, of SEQ ID NO:2, but not nucleotides which are immediately contiguous to those sequences in the naturally-occurring genomic sequence or naturally-occurring cDNA.

[0012] Also within the invention is a method of lubricating a mammalian joint by contacting the joint with a tribonectin. The mammal is preferably a human, horse, dog, ox, donkey, mouse, rat, guinea pig, cow, sheep, pig, rabbit, monkey, or cat, and the joint is an articulating joint such as a knee, elbow, shoulder, hip, or any other weight-bearing joint. Tribonectins are administered intra-articularly. Therapeutic joint lubrication is also carried out by gene therapy, e.g., by contacting the joint or synovial fluid with a nucleic acid encoding a tribonectin. For example, nucleic acids are administered to a synovial cavity by intra-articular injection.

[0013] In addition to functioning as a boundary lubricant in a mammalian joint, a tribonectin is used as a boundary lubricant between soft mammalian tissues such as skin or internal organs or between a mammalian tissue and a medical device such as a prosthetic implant. Accordingly, the invention encompasses a biocompatible composition containing a tribonectin in a form suitable for the inhibition of tissue adhesion formation. For example, the tribonectin is in

the form of a film, membrane, foam, gel, or fiber. The term "film," as used herein, means a substance formed by compressing a foam or gel to a thin membrane, e.g., by casting into a flat mold and air drying to a thin membrane, or by compressing a gel or fibers, or by allowing or causing a gel or fibers to dehydrate. The term "foam," as used herein, means a substance with a porous structure formed, e.g., by introduction of air into a tribonectin solution, suspension, gels, or fiber. The term "bioabsorbable," as used herein, refers to the ability of a tissue-compatible material to degrade in the body after implantation, into nontoxic products which are eliminated from the body or metabolized. A "biocompatible" substance, as the term is used herein, is one that has no medically unacceptable toxic or injurious effects on biological function. Tribonectin compositions for the prevention of adhesions are also formulated as compositions suitable for extrusion, e.g., to form a mold upon which tissue can grow without adhering.

[0014] A method inhibiting adhesion formation between a first surface and a second surface in a mammal is carried out by placing a tribonectin between the first and second surfaces in an amount sufficient to prevent adhesion of the surfaces in the mammal. For example, one or both of the surfaces is a mammalian tissue, and a tribonectin placed between them prevents formation of adhesions during the healing process. Alternatively the first or the second surface (or both) is an artificial device such as an orthopedic implant. Tissues to be treated include those injured by surgical incision or trauma.

[0015] Also within the invention is a method for diagnosing osteoarthritis or a predisposition thereto by obtaining a biological sample from a mammal and measuring the amount of an MSF fragment in the biological sample. An increase in the amount compared to a control, e.g., a predetermined value associated with a negative diagnosis or a biological sample from a mammal known to be free of osteoarthritis, indicates that the mammal suffers from osteoarthritis or is predisposed to developing osteoarthritis. Any biological sample is suitable for testing in the diagnostic method; typically, the biological sample is synovial fluid, blood, serum, or urine. Preferably, the MSF fragment contains the amino acid sequence of SEQ ID NO:3. Alternatively, the MSF fragment contains the amino acid sequence of EPAPTT (SEQ ID NO:5; a product of trypsin cleavage of a tribonectin) or the amino acid sequence of PTTKEP (SEQ ID NO:6; a product of elastase cleavage of a tribonectin).

[0016] Other features and advantages of the invention will be apparent from the following description of the preferred embodiments thereof, and from the claims.

DETAILED DESCRIPTION

[0017] A human lubricating polypeptide was purified from synovial fluid and found to contain amino acids encoded by exons 2 and 4-12 of the MSF gene (but not exons 1 or 3). The gene encoding naturally-occurring full length MSF contains 12 exons, and the naturally-occurring MSF gene product contains 1404 amino acids with multiple polypeptide sequence homologies to vitronectin including hemopexin-like and somatomedin-like regions. Centrally-located exon 6 contains 940 residues. Exon 6 encodes a O-glycosylated mucin domain. A polypeptide encoded by nucleotides 631-3453 of SEQ ID NO:2 provides boundary lubrication of articular cartilage.

TABLE 1

MSF amino acid sequence
MAWKTLPIYLLLLLSVFVIQQVSSQDLSSCAGRCGEGYSRDATCNCNDYNCQHMECCPDF (SEQ ID NO:1)
KRVCTAELSCKGRCFESFERGRECDCAQCKKYDKCCPDYESFCAEVHNPTSPSSKKAP
PPSGASQTIKSTTKRSPKPNKKKKKVVIESEEITEHSVSENQESSSSSSSSSSSTIW
KIKSSKNSAANRELQKKLVKDNKKNRTKKKPTPKPPVDEAGSGLDNGDFKVTTPTDST
<u>TOHNKVSTSPKITTAKPINRPSLPPNSDTSKETSLTVNKETTIVETKETTITTNKOTSTDG</u>
<u>KEKTTSAKETOSIEKTSAKDLAPTSKVLAKPTPKAETTTKGPALTTPKEPTPTTPKEPAS</u>
<u>TTPKETPTTIKSAPTTPKEPAPTTTKSAPTTPKEPAPTTTKEPAPTTPEKEPAPTTTKEP</u>
<u>APTTTKSAPTTPKEPAPTTPKKPAPTTPKEPAPTTPEKEPTPTTPKEPAPTTKEPAPTTPK</u>
<u>EPAPTAPKKPAPTTPKEPAPTTPKEPAPTTTKEPSPTTPKEPAPTTTKSAPTTTKEPAPT</u>
<u>TTKSAPTTPKEPSPTTKEPAPTTPKEPAPTTPKKPAPTTPKEPAPTTPEKEPAPTTKKP</u>
<u>APTAPKEPAPTTPKETAPTTPKKLTPTTPEKLAPTTPEKPAPTTPEELAPTTPEEPTPTT</u>
<u>PEEPAPTTPKAAPNTPKEPAPTTPKEPAPTTPKEPAPTTPKETAPTTPKGTAPTTLKEP</u>
<u>APTTPKKPAPKELAPTTTKEPTSTTS DKPAPTTPKGTAPTTPKEPAPTTPEKEPAPTTPKG</u>
<u>TAPTTLKEPAPTTPKKPAPKELAPTTTKGPTSTTS DKPAPTTPKETAPTTPEKEPAPTTPK</u>
<u>KPAPTTPETPPPTTSEVSTPTTTKEPTTIHKSPDESTPELSAEPKALENSPKEPGVPT</u>
<u>TKTPAATKPEMTTAKDKTTERDLRTTPETTTAAPKMTKETATTEKTTESKITATTTTOV</u>
<u>TSTTTQDTPFKITTLKTTTLAPKVTTTKKTIITTEIMNKPEETA KPDRATNSKATTPK</u>
<u>POKPTKAPKKPTSTKKPKTMPRVKPKTTPTRKMTSTMPELNPTSRIA EAMLOTTTRPN</u>
<u>OTPN SKLVEVNPKSEDAGGAEGETPHMLLRPHVFMPEVTPDMDYLP RVPNQGI IINPMLS</u>
DETNICNGKPV DGLTTLRNGTLVAFRGHYFWMLSPFSPSPARRITEVWGIPSPIDTVFT
RCNCEGKTFFFKDSQYWRFTNDIKDAGYPKPIFKGFGGLTGQIVAALSTAKYKNWPESVY
FFKRGSIQQYIYKQEPVQKCPGRRPALNYPVYGEMTQVRRRRFERAIGPSQTHIRIQY
SPARLAYQDKGVLHNEVKVSILWRGLPNVVTSAISLPNIRKPDGYDYAFSKDQYYNIDV
PSRTARAITTRSGQTL SKVWYNCP

[0018]

TABLE 2

MSF cDNA
1 gcggccgcga ctattcggta cctgaaaaca acgatggcat ggaaaacact tcccatttac (SEQ ID NO:2)
61 ctgtgtgttg tgctgtctgt tttcgtgatt cagcaagttt catctcaaga tctatcaagc
121 tgtgcaggga gatgtgggga agggatttct agagatgcca cctgcaactg tgattataac
181 tgtcaacact acatggagtg tgcacctgat ttcaagagag tctgcactgc ggagctttcc
241 tgtaaaggcc gctgctttga gtccttcgag agaggaggag agtgtgactg cgacgcccaa
301 tgtaagaagt atgacaagtg tgtgtccgat tatgagagtt tctgtgcaga agtgcataat
361 cccacatcac caccatcttc aaagaaagca cctccacctt caggagcatc tcaaaccatc
421 aaatcaacaa ccaaacgttc acccaaacca ccaacaaga agaagactaa gaaagttata
481 gaatcagagg aaataacaga agaacattct gtttctgaaa atcaagagtc ctcctcctcc

TABLE 2-continued

MSF cDNA	
541	tcctcctctt cctcttcttc ttcaacaatt tggaaaatca agtcttccaa aaattcagct
	EXON 6
601	gctaatagag aattacagaa gaaactcaaa <u>gtaaaagata acaagaagaa cagaactaaa</u>
661	<u>aagaaaccta cccccaaacc accagttgta gatgaagctg gaagtggatt ggacaatggt</u>
721	<u>gacttcaagg tcacaactcc tgacacgtct accaccaac acaataaagt cagcacatct</u>
781	<u>cccaagatca caacagcaaa accaataaat ccagaccca gtcttcacc taattctgat</u>
841	<u>acatctaaag agacgtcttt gacagtgaat aaagagacaa cagttgaaac taaagaaact</u>
901	<u>actacaacaa ataacagac ttcaactgat ggaaaagaga agactacttc cgctaaagag</u>
961	<u>acacaaagta tagagaaac atctgctaaa gatttagcac ccacatctaa agtgctggct</u>
1021	<u>aaacctacac ccaaagctga aactacaacc aaaggccctg ctctcaccac tccaaggag</u>
1081	<u>cccacgcccc ccactcccaa ggagcctgca tctaccacac ccaaagagcc cacacctacc</u>
1141	<u>accatcaagt ctgcaccac ccccccaag gagcctgcac ccaccaccac caagtctgca</u>
1201	<u>cccaccactc ccaaggagcc tacaccacc accaccaagg agcctgcacc caccactccc</u>
1261	<u>aaggagcctg caccaccac caccaaggag cctgcaccca ccaccacaa gtctgcacc</u>
1321	<u>accactccca aggagcctgc acccaccacc ccaagaagc ctgcccaac taccccaag</u>
1381	<u>gagcctgcac ccaccactcc caaggagcct acaccacca ctccaagga gcctgcacc</u>
1441	<u>accaccaagg agcctgcacc caccactccc aaagagcctg caccactgc cccaagaag</u>
1501	<u>cctgccccaa ctacccccaa ggagcctgca cccaccactc ccaaggagcc tgcaccacc</u>
1561	<u>accaccaagg agccttcacc caccactccc aaggagcctg caccaccac caccaagtct</u>
1621	<u>gcaccacaca ctaccaagga gcctgcacc accactacca agtctgcacc caccactccc</u>
1681	<u>aaggagcctt caccaccac caccaaggaa cctgcaccca ccaactccaa ggagcctgca</u>
1741	<u>cccaccacc ccaagaagcc tgcccaact accccaagg agcctgcacc caccactccc</u>
1801	<u>aaggaacctg caccaccac caccaagaag cctgcaccca ccgctccaa agagcctgcc</u>
1861	<u>ccaactacc ccaaggagac tgcaccacc accccaaga agctcacgc caccacccc</u>
1921	<u>gagaagctcg caccaccac ccctcagaag ccgcaccca ccaccctga ggagctcgca</u>
1981	<u>cccaccacc ctgaggagcc cacaccacc accctgagg agcctgctcc caccactccc</u>
2041	<u>aaggcagcgg ctcccaacac ccctaaggag cctgctccaa ctaccctaa ggagcctgcc</u>
2101	<u>ccaactacc ctaaggagcc tgctccaact acccctaagg agactgctcc aactaccct</u>
2161	<u>aaagggactg ctccaactac cctcaaggaa cctgcaccca ctactccaa gaagcccgcc</u>
2221	<u>ccaaggagc ttgcaccac caccaccaag gagccacat ccaccactc tgacaagccc</u>
2281	<u>actccaacta cccctaagg gactgctcca actacccta aggaacctgc tccaactacc</u>
2341	<u>cctaaggagc ctgctccaac taccctaag gggactgctc caactacct caaggaacct</u>
2401	<u>gcaccaccta ctccaagaa gcctgcccc aaggagcttg caccaccac caccaagggg</u>
2461	<u>cccacatcca ccacctgta caaacctgct ccaactacac ctaaggagac tgctccaact</u>
2521	<u>accccaagg agcctgcacc cactacccc aagaagcctg ctccaactac tcctgagaca</u>
2581	<u>cctctccaa ccaattcaga ggtctctact ccaactacca ccaaggagcc taccactacc</u>
2641	<u>cacaaaagcc ctgatgaatc aactcctgag cttctctcag aaccacacc aaaagctctt</u>
2701	<u>gaaaacagtc ccaaggaaac tgggtgtacct acaactaaga ctctgcagc gactaaacct</u>

TABLE 2-continued

MSF cDNA	
2761	<u>gaaatgacta caacagctaa agacaagaca acagaaagag acttacgtac tacacctgaa</u>
2821	<u>actacaactg ctgcacctaa gatgacaaaa gagacagcaa ctacaacaga aaaaactacc</u>
2881	<u>gaatccaaaa taacagctac aaccacacaa gtaacatcta ccacaactca agataccaca</u>
2941	<u>ccattcaaaa ttactactct taaaacaact actcttgac ccaaagtaac tacaacaaaa</u>
3001	<u>aagacaatta ctaccactga gattatgaac aaacctgaag aaacagctaa accaaaagac</u>
3061	<u>agagctacta attctaaagc gacaactcct aaacctcaaa agccaaccaa agcaccctaaa</u>
3121	<u>aaaccactt ctaccacaaa gccaaaaaca atgcctagap tgagaaaacc aaagacgaca</u>
3181	<u>ccaactcccc gcaagatgac atcaacaatg ccagaattga accctacctc aagaatagca</u>
3241	<u>gaagccatgc tccaaaccac caccagacct aaccaaactc caaactccaa actagttgaa</u>
3301	<u>gtaaatccaa agagtgaaga tgcaggtggt gctgaaggag aaacacctca tatgctcctc</u>
3361	<u>aggcccatg tggtcatgcc tgaagttact cccgacatgg attacttacc gagagtaccc</u>
3421	<u>aatcaaggca ttatcatcaa tcccatgctt tccgatgaga ccaatatatg caatggtaag</u>
3481	<u>ccagtagatg gactgactac ttgcgcaat gggacattag ttgattccg aggtcattat</u>
3541	<u>ttctggatgc taagtccatt cagtccacca tctccagctc gcagaattac tgaagtttgg</u>
3601	<u>ggtattcctt ccccatgtga tactgttttt actaggtgca actgtgaagg aaaaactttc</u>
3661	<u>ttctttaagg attctcagta ctggcgtttt accaatgata taaaagatgc agggtacccc</u>
3721	<u>aaaccaattt tcaaaggatt tggaggacta actggacaaa tagcggcagc gctttcaaca</u>
3781	<u>gctaaatata agaactggcc tgaatctgtg tttttttca agagaggtgg cagcattcag</u>
3841	<u>cagtatatat ataaacagga acctgtacag aagtgcctg gaagaaggcc tgctctaaat</u>
3901	<u>tatccagtgt atggagaaat gacacaggtt aggagacgtc gctttgaacg tgctatagga</u>
3961	<u>ccttctcaaa cacacaccat cagaattcaa tattcacctg ccagactggc ttatcaagac</u>
4021	<u>aaaggtgtcc ttcataatga agttaaagt agtatactgt ggagaggact tccaaatgtg</u>
4081	<u>gttacctcag ctatatcact gcccaacatc agaaaacctg acggctatga ttactatgcc</u>
4141	<u>ttttctaaag atcaatacta taacattgat gtgcctagta gaacagcaag agcaattact</u>
4201	<u>actcgttctg ggcagacott atccaaagtc tgggtacaact gtccttagac tgatgagcaa</u>
4261	<u>aggaggagtc aactaatgaa gaaatgaata ataaattttg acactgaaaa acattttatt</u>
4321	<u>aataaagaat attgacatga gtataccagt ttatatataa aaatgtttt aaacttgaca</u>
4381	<u>atcattacac taaaacagat ttgataatct tattcacagt tgttattgtt tacagaccat</u>
4441	<u>ttaattaata ttctctctgt ttattcctcc tctccctccc attgcatggc tcacacctgt</u>
4501	<u>aaaagaaaa agaatacaat tgaatatatc ttttaagaat tcaaaactag tgtattcact</u>
4561	<u>taccctagtt cattataaaa aatatctagg cattgtggat ataaaactgt tgggtattct</u>
4621	<u>acaacttcaa tggaaattat tacaagcaga ttaatccctc tttttgtgac acaagtacaa</u>
4681	<u>tctaaaagtt atattggaaa acatggaaat attaaaattt tacactttta ctagctaaaa</u>
4741	<u>cataatcaca aagctttatc gtgtgtgata aaaaaattaa caatataatg gcaataggta</u>
4801	<u>gagatacaac aaatgaatat aacactataa cacttcatat tttccaaatc ttaatctgga</u>
4861	<u>tttaaggaag aaatcaataa atataaaata taagcacata tttatcatat atctaaggta</u>
4921	<u>tacaaatctg tctacatgaa gtttacagat tggtaaatat cacctgctca acatgtaatt</u>

TABLE 2-continued

MSF cDNA	
4981	atttaataaa actttggaac attaaaaaaa taaattggag gcttaaaaaa aaaaaaaaaa
5041	a

[0019]

TABLE 3

MSF Exon Boundaries		
Exon	Amino acid sequence in SEQ ID NO: 1	Nucleotide sequence in SEQ ID NO: 2
1	1-24, inclusive	34-105, inclusive
2	25-66, inclusive	106-231, inclusive
3	67-104, inclusive	232-345, inclusive
4	105-155, inclusive	346-498, inclusive
5	156-199, inclusive	499-630, inclusive
6	200-1140, inclusive	631-3453, inclusive
7	1141-1167, inclusive	3454-3534, inclusive
8	1168-1212, inclusive	3535-3670, inclusive
9	1213-1263, inclusive	3671-3822, inclusive
10	1264-1331, inclusive	3823-4026, inclusive

recombinant or chemically-produced lubricating polypeptide containing at least one repeat of the amino acid sequence KEPAPTT (SEQ ID NO:3) either identically or with conservative substitution is also administered to lubricate mammalian joints.

[0021] Production and Purification of Recombinant Lubricating Polypeptides

[0022] To produce recombinant polypeptides, DNA containing exon 6 of MSF (nucleotides 631-3453 of SEQ ID NO:2) in an appropriate expression vector is transfected into a cell. The DNA can also contain some or all of exon 7 (nucleotides 354-3534 of SEQ ID NO:2), exon 8 (nucleotides 3535-3670 of SEQ ID NO:2), or exon 9 (nucleotides 3671-3822 of SEQ ID NO:2) of the MSF gene. Primers for polymerase chain reaction (PCR) methods to generate DNA which spans various exons of MSF are shown below.

TABLE 4

PCR Primers		
MSF exons	Forward Primer	Reverse Primer
exon 2	5'AGATTATCAAGCTGT (SEQ ID NO:7) GCAGGGAG3'	5'TTTACAGGAAAGC (SEQ ID NO:8) TCCGCAGTG3'
exon 6	5'TCAAGGTCACAACTCC (SEQ ID NO:9) TGACACG3'	5'CTCTCGGTAAGTA (SEQ ID NO:10) ATCCATGTCGG3'
exons 2-12	5'TTGTTGCTGCTGTCTG (SEQ ID NO:11) TTTTCG3'	5'TGGATAAGGTCTG (SEQ ID NO:12) CCCAGAACGAG3'
exons 6-12	5'TCAAGGTCACAACTCC (SEQ ID NO:13) TGACACG3'	5'GATGGTGTGTGTT (SEQ ID NO:14) TGAGAAGGTCC3'

TABLE 3-continued

MSF Exon Boundaries		
Exon	Amino acid sequence in SEQ ID NO: 1	Nucleotide sequence in SEQ ID NO: 2
11	1332-1371, inclusive	4027-4146, inclusive
12	1372-1404, inclusive	4147-4245, inclusive

[0020] The boundary lubricant isolated from synovial fluid is an alternatively-spliced variant of MSF. This alternatively-spliced variant was found to be the composition present in synovial fluid that confers lubricating capabilities to the articular joint. The boundary lubricant isolated from human synovial fluid contains amino acids encoded by exons 2, and 4-12 of the MSF gene, i.e., the alternative splice variant lacks amino acids encoded by exons 1 and 3 of the MSF gene. A recombinant or chemically-produced polypeptide containing at least exon 6 (but not exons 1 or 3) of MSF is useful to prevent and/or treat osteoarthritic disease. A

[0023] Methods of designing forward and reverse primers used to make DNAs which encode tribonection polypeptides of varying lengths and which incorporate various exons of the MSF gene, e.g., to make polypeptide encoded by exons 2, 4-12; exons 6-9; and exons 2, 4-9, are well known in the art of molecular biology. Standard methods for transfecting cells with isolated nucleic acid are well known to those skilled in the art of molecular biology. For example, prokaryotic or eukaryotic cells in culture are transfected with the DNA of the invention operatively linked to expression control sequences appropriate for high-level expression in the cell. Such cells are useful for producing large amounts of the lubricating polypeptide, which are purified using standard methods. The lubricating polypeptides are used therapeutically for treatment or prevention of arthritic diseases. The polypeptides are also used to raise antibodies against a naturally-occurring or recombinantly-produced lubricating glycoproteins or glycopeptides.

[0024] For example, the recombinant gene product is expressed as a fusion protein and purified using a commer-

cially available expression and purification system, e.g., the pFLAG expression system (IBI). The expression systems that may be used for purposes of the invention include, but are not limited to, microorganisms such as bacteria (e.g., *E. coli* and *B. subtilis*) transformed with recombinant bacteriophage DNA, plasmid DNA, or cosmid DNA expression vectors containing the nucleic acid molecules described herein. For production of glycosylated polypeptides, eukaryotic expression systems are used. Yeast (for example, *Saccharomyces* and *Pichia*) transformed with recombinant yeast expression vectors containing the recombinant nucleic acid encoding a tribonectin polypeptide are used. Insect cell systems infected with recombinant virus expression vectors (for example, baculovirus) containing the nucleic acid molecules encoding a tribonectin and mammalian cell systems (e.g., COS, CHO, BHK, 293, VERO, HeLa, MDCK, WI38, and NIH 3T3 cells) harboring recombinant expression constructs containing promoters derived from the genome of mammalian cells (e.g., the metallothionein promoter) or from mammalian viruses (e.g., the adenovirus late promoter and the vaccinia virus 7.5K promoter) are also useful. In addition to clinical applications, recombinant polypeptides are injected into a rabbit or rodent to produce antibodies as described below.

[0025] The synovial fluid of an inflamed or injured joint contains proteolytic enzymes that degrade lubricating proteins or polypeptides. For example, infiltrating immune cells such as neutrophils secrete trypsin and/or elastase. Even a minor injury to an articulating joint or an inflammatory state can result in cellular infiltration and proteolytic enzyme secretion resulting in traumatic synovitis. Synovitis for a period of a few days or weeks can result in the loss of the cytoprotective layer of a joint, which in turn leads to the loss of cartilage. Non-lubricated cartilaginous bearings may experience premature wear which may initiate osteoarthritis. Individuals who clinically present with a traumatic effusion (e.g., "water on the knee") are predisposed to developing osteoarthritis; the elaboration of proteolytic enzymes degrades and depletes naturally-occurring lubricating compositions in the synovial fluid. Depletion of natural lubricating compositions occurs in other inflammatory joint diseases such as rheumatoid arthritis. Replacing or supplementing the synovial fluid of such injured joints with the lubricating compositions of the invention prevents the development of osteoarthritis in the long term (e.g., years, even decades later) and immediately lubricates the joint to minimize short term damage.

[0026] Analogs, homologs, or mimetics of lubricating peptides which are less susceptible to degradation in vivo are used to lubricate mammalian joints. Analogs can differ from the naturally-occurring peptides by amino acid sequence, or by modifications which do not affect the sequence, or both. Modifications (which do not normally alter primary sequence) include in vivo or in vitro chemical derivitization of polypeptides, e.g., acetylation or carboxylation. Also included are modifications of glycosylation, e.g., those made by modifying the glycosylation patterns of a polypeptide during its synthesis- and processing or in further processing steps, e.g., by exposing the polypeptide to enzymes which affect glycosylation, e.g., mammalian glycosylating or deglycosylating enzymes.

[0027] Where proteolytic degradation of the peptides following injection into the subject is a problem, replacement

of a particularly sensitive peptide bond with a noncleavable peptide mimetic bond renders the resulting peptide more stable, and thus more useful as a therapeutic. To render the therapeutic peptides less susceptible to cleavage by peptidases such as trypsin or elastase, the peptide bonds of a peptide may be replaced with an alternative type of covalent bond (a "peptide mimetic"). Trypsin, elastase, and other enzymes may be elaborated by infiltrating immune cells during joint inflammation. Trypsin cleaves a polypeptide bond on the carboxy-side of lysine and arginine; elastase cleaves on the carboxy-side of alanine, glycine. Thrombin, a serine protease which is present in hemorrhagic joints, cleaves a peptide bond on the carboxy-side of arginine. Collagenases are a family of enzymes produced by fibroblasts and chondrocytes when synovial metabolism is altered (e.g., during injury). These enzymes cut on the carboxy-side of glycine and proline. One or more peptidase-susceptible peptide bonds, e.g., those which appear in the KEPAPTT (SEQ ID NO:3) repeat sequence, are altered (e.g., replaced with a non-peptide bond) to make the site less susceptible to cleavage, thus increasing the clinical half-life of the therapeutic formulation.

[0028] Such mimetics, and methods of incorporating them into polypeptides, are well known in the art. Similarly, the replacement of an L-amino acid residue with a D-amino acid is a standard way of rendering the polypeptide less sensitive to proteolysis. Also useful are amino-terminal blocking groups such as t-butyloxycarbonyl, acetyl, theyl, succinyl, methoxysuccinyl, suberyl, adipyl, azelalyl, dansyl, benzylloxycarbonyl, fluorenylmethoxycarbonyl, methoxyazelalyl, methoxyadipyl, methoxysuberyl, and 2,4-dinitrophenyl.

[0029] Clinical formulations of tribonectins may also contain peptidase inhibitors such as N-methoxysuccinyl-Ala-Ala-Pro-Val chloromethylketone (an inhibitor of elastase). Other clinically acceptable protease inhibitors (e.g., as described in Berling et al., 1998, Int. J. Pancreatolgy 24:9-17) such as leupeptin, aprotinin, α 1-antitrypsin, α 2-macroglobulin, α 1-protease inhibitor, antichymotrypsin (ACHY), secretory leukocyte protease inhibitor (SLPI) are also co-administered with a tribonectin to reduce proteolytic cleavage and increase clinical halflife. A cocktail of two or more protease inhibitors can also be coadministered.

[0030] Compositions of tribonectin polypeptides or nucleic acids encoding the polypeptides are formulated in standard physiologically-compatible excipients known in the art, e.g., phosphate-buffered saline (PBS). Other formulations and methods for making such formulations are well known and can be found in, e.g., "Remington's Pharmaceutical Sciences". Tribonectins are also formulated with non-crosslinked hyaluronic acid preparations or viscosupplementation compositions, such as cross-linked hyaluronic acid preparations. When a tribonectin is added to a viscosupplement formulation, the interaction of the tribonectin with hyaluronic acid reduces the viscosity of the viscosupplement.

[0031] Methods of making a glycopeptide and determining % glycosylation are known in the art, e.g., as described in U.S. Pat. No. 5,767,254. The presence of N-acetylgalactosamine is indicative of the presence of O-linked oligosaccharides (or Ser/Thr-linked oligosaccharides) in which GalNAc is commonly found in O-glycosidic alpha-linkage directly to amino acid. The presence of O-linked oligosac-

charide is also detected by binding to Jacalin-Sepharose, an immobilized plant lectin that binds to the core disaccharide sequence Gal β (1-3)GalNAc linked to Ser/Thr in glycoproteins, or peanut agglutinin, which binds to β (1-3)Gal-GalNAc. O-linked oligosaccharides are distinguished from N-linked oligosaccharides using standard methods. For example, oligosaccharides in O-glycosidic linkage, but not in N-glycosidic linkage, are susceptible to release from peptide by treatment with mild base in the presence of sodium borohydride (50 mM NaOH, 1M NaBH₄, 16 hr at 45° C.) to cause a beta-elimination reaction.

[0032] Veterinary Applications

[0033] Canine osteoarthritis is a prevalent clinical disorder that is treated using the methods described herein. Osteoarthritis afflicts an estimated one in five adult dogs; an estimated 8 million dogs suffer from this degenerative, potentially debilitating disease. Yet, many owners do not recognize the signs of chronic canine pain. While any dog can develop osteoarthritis, those most at risk are large breeds, geriatric dogs, very active dogs (such as working or sporting animals), and those with inherited joint abnormalities such as hip or elbow dysplasia.

[0034] Equine degenerative joint disease such as osteoarthritis is a cause of lameness and impaired performance in horses. As with humans and other mammals, degenerative joint diseases which affect horses are progressive disorders of synovial joints characterized by articular cartilage degeneration and joint effusion. Acute or chronic trauma, overuse, developmental disease, joint instability and old age leads to synovitis, impaired chondrocyte metabolism, and the formation of fissures in the joint cartilage. Destructive enzymes such as trypsin, elastase, stromelysin and hyaluronidase are released into the joint where they degrade synovial fluid and cartilage components, resulting in decreased synovial fluid viscosity, poor lubrication, depressed cartilage metabolism and enhanced wear resulting in pain and cartilage erosion. Current therapeutic approaches include medications for pain relief and anti-inflammatory drugs. The compositions and methods described herein are useful to replenish the lubricating capabilities of the affected joint.

[0035] Administration of Therapeutic Polypeptides

[0036] Standard methods for delivery of peptides are used. Such methods are well known to those of ordinary skill in the art. For intra-articular administration, tribonection is delivered to the synovial cavity at a concentration in the range of 20-500 μ g/ml in a volume of approximately 0.1-2 ml per injection. For example, 1 ml of a tribonection at a concentration of 250 μ g/ml is injected into a knee joint using a fine (e.g., 14-22 gauge, preferably 18-22 gauge) needle. The compositions of the invention are also useful for parenteral administration, such as intravenous, subcutaneous, intramuscular, and intraperitoneal.

[0037] For prevention of surgical adhesions, the tribonections described herein are administered in the form of gel, foam, fiber or fabric. A tribonection formulated in such a manner is placed over and between damaged or exposed tissue interfaces in order to prevent adhesion formation between apposing surfaces. To be effective, the gel or film must remain in place and prevent tissue contact for a long enough time so that when the gel finally disperses and the tissues do come into contact, they will no longer have a

tendency to adhere. Tribonections formulated for inhibition or prevention of adhesion formation (e.g., in the form of a membrane, fabric, foam, or gel) are evaluated for prevention of post-surgical adhesions in a rat cecal abrasion model (Goldberg et al., In *Gynecologic Surgery and Adhesion Prevention*. Willey-Liss, pp. 191-204, 1993). Compositions are placed around surgically abraded rat ceca, and compared to non-treated controls (animals whose ceca were abraded but did not receive any treatment). A reduction in the amount of adhesion formation in the rat model in the presence of the tribonection formulation compared to the amount in the absence of the formulation indicates that the formulation is clinically effective to reduce tissue adhesion formation.

[0038] Tribonections are also used to coat artificial limbs and joints prior to implantation into a mammal. For example, such devices are dipped or bathed in a solution of a tribonection, e.g., as described in U.S. Pat. Nos. 5,709,020 or 5,702,456.

[0039] Lubricating polypeptides are at least about 10 amino acids (containing at least one KEPAPTT (SEQ ID NO:3)) or XXXTTX (SEQ ID NO:4) repeat), usually about 20 contiguous amino acids, preferably at least 40 contiguous amino acids, more preferably at least 50 contiguous amino acids, and most preferably at least about 60 to 80 contiguous amino acids in length. For example, the polypeptide is approximately 500 amino acids in length and contains 76 repeats of KEPAPTT (SEQ ID NO:3). The polypeptide is less than 1404 residues in length, e.g., it has the amino acid sequence of naturally-occurring MSF (SEQ ID NO:1) but lacks at least 5, 10, 15, 20, or 24 amino acids at the N-terminus of naturally-occurring MSF. Such peptides are generated by methods known to those skilled in the art, including proteolytic cleavage of a recombinant MSF protein, de novo synthesis, or genetic engineering, e.g., cloning and expression of at least exon 6, 7, 8, and/or 9 of the MSF gene.

[0040] Tribonection polypeptides are also biochemically purified. The enzyme chymotrypsin cleaves at sites which bracket amino acids encoded by exon 6 of the MSF gene. Thus, a polypeptide containing amino acids encoded by exon 6 of the MSF gene (but not any other MSF exons) is prepared from a naturally-occurring or recombinantly produced MSF gene product by enzymatic digestion with chymotrypsin. The polypeptide is then subjected to standard biochemical purification methods to yield a substantially pure polypeptide suitable for therapeutic administration, evaluation of lubricating activity, or antibody production.

[0041] Therapeutic compositions are administered in a pharmaceutically acceptable carrier (e.g., physiological saline). Carriers are selected on the basis of mode and route of administration and standard pharmaceutical practice. A therapeutically effective amount of a therapeutic composition (e.g., lubricating polypeptide) is an amount which is capable of producing a medically desirable result, e.g., boundary lubrication of a mammalian joint, in a treated animal. A medically desirable result is a reduction in pain (measured, e.g., using a visual analog pain scale described in Peyron et al., 1993, *J. Rheumatol.* 20 (suppl.39):10-15) or increased ability to move the joint (measured, e.g., using pedometry as described in Belcher et al., 1997, *J. Orthop. Trauma* 11:106-109). Another method to measure lubricity of synovial fluid after treatment is to reaspirate a small

volume of synovial fluid from the affected joint and test the lubricating properties in vitro using a friction apparatus as described herein.

[0042] As is well known in the medical arts, dosage for any one animal depends on many factors, including the animal's size, body surface area, age, the particular compound to be administered, sex, time and route of administration, general health, and other drugs being administered concurrently. Administration is generally local to an injured or inflamed joint. Alternatively, the polypeptides are administered via a timed-release implant placed in close proximity to a joint for slow release at the site of an injured or inflamed joint.

[0043] Gene Therapy

[0044] Gene therapy is carried out by administering to a mammal a nucleic acid encoding a therapeutic lubricating polypeptide, e.g., DNA encoding one or more repeats or the amino acid sequence KEPAPTT (SEQ ID NO:3) or DNA encoding a lubricating fragment of MSF, by standard vectors and/or gene delivery systems. Suitable gene delivery systems include liposomes, receptor-mediated delivery systems, naked DNA, and viral vectors such as herpes viruses, retroviruses, adenoviruses and adeno-associated viruses.

[0045] In addition to a gene delivery system as described above, the therapeutic composition may include a pharmaceutically acceptable carrier, e.g., a biologically compatible vehicle such as physiological saline, suitable for administration to an animal. A therapeutically effective amount of a nucleic acid or polypeptide composition is an amount which is capable of producing a medically desirable result in a treated animal, e.g., a reduction in pain associated with joint movement, an increase in lubricating function of synovial fluid.

[0046] Parenteral administration, such as intravenous, subcutaneous, intramuscular, and intraperitoneal delivery routes, may be used to deliver the compound. Preferably, therapeutic compositions such as nucleic acids or polypeptides are delivered intra-articularly. Dosage for any one patient depends upon many factors, including the patient's size, body surface area, age, the particular compound to be administered, sex, time and route of administration, general health, and other drugs (e.g., anti-inflammatory drugs, viscotherapeutic drugs) being administered concurrently. A preferred dosage for administration of nucleic acids is from approximately 10^6 to 10^{22} copies of the nucleic acid molecule.

[0047] DNA is introduced into target cells of the patient by standard vectors, e.g., a vector which contains DNA encoding a tribonectin operably linked to a promoter sequence. Suitable gene delivery systems may include liposomes, receptor-mediated delivery systems, naked DNA, and viral vectors such as herpes viruses, retroviruses, and adenoviruses, among others.

[0048] DNA may be administered locally using an adenovirus or adeno-associated virus delivery system using standard methods. For example, methods of delivering DNA intra-articularly to synovial fluid and methods of delivering DNA to cells from synovial fluid (e.g., synovial fibroblasts or chondrocytes) are described in U.S. Pat. No. 5,858,355. The only cis-acting sequences required for replication and packaging of recombinant adeno-associated virus (AAV) vector

are the AAV terminal repeats. Up to 4 kb of DNA is inserted between the terminal repeats without effecting viral replication or packaging. To package a recombinant AAV vector, a plasmid containing the terminal repeats and DNA encoding a therapeutic polypeptide is co-transfected into cells with a plasmid that expresses AAV rep and capsid proteins. The transfected cells are then infected with adeno-associated virus, and recombinant AAV virus containing the desired sequences is isolated from cells approximately 48-72 hours after transfection. Recombinant virus is then administered for gene therapy applications using known methods.

[0049] Electroporation is another method of introducing DNA into target cells, e.g., synovial fibroblasts or chondrocytes, ex vivo. Cells to be electroporated are placed into Hepes buffer saline (HBS) at a concentration of about 10^7 cells per ml. The DNA to be electroporated is added at a concentration of approximately 5-20 $\mu\text{g/ml}$ of HBS. The mixture is placed into an electroporation device and an electric field is applied according to standard protocols, e.g., in a range of between about 250 and 300 volts. Following introduction of DNA into synovial cells ex vivo, the genetically modified autologous synovial cells are transplanted back into the donor by intra-articular injection. Approximately 10^7 cells are injected intra-articularly into joints in a volume of approximately 1 ml.

[0050] Synovial cells into which DNA is introduced are obtained using routine methods, e.g., through an arthroscope. The arthroscope is a small, hollow rod inserted into the knee via a small puncture wound which allows access to a surgical instrument to recover synovial cells arthroscopically. In some cases, the synovial cells in arthroscopically excised tissue are aseptically recovered by enzymatic digestion of the connective tissue matrix. For example, the synovium is cut into pieces of approximately 1 mm diameter and digested sequentially with trypsin (0.2% w/v in Gey's Balanced Salt Solution) for 30 minutes at 37° C., and collagenase (0.2% w/v in Gey's Balanced Salt Solution) for 2 hours at 37° C. A suspension of genetically-modified cells is injected into a recipient mammalian joint. Intra-articular injections of this type are routine and carried out in the doctor's office without additional surgical intervention. Repeat injections are carried out as needed.

[0051] Alternatively, the DNA (naked or packaged in a virus) is formulated in a suitable pharmaceutical carrier and injected intra-articularly. Gene therapy is also administered as a prophylactic measure to prevent the development of osteoarthritis in those individuals determined to be highly susceptible of developing this disease, e.g., those who have suffered an acute joint injury. Direct intra-articular injection of a DNA encoding a therapeutic polypeptide into a joint results in transfection of the recipient synovial cells to allow expression of DNA.

[0052] Drugs which stimulate an endogenous tribonectin promoter, e.g., TGF β , may also be administered as described above to increase the level of synovial expression.

[0053] Production of Antibodies Specific for Synovial Lubricating Polypeptides

[0054] Antibodies specific for lubricating polypeptides are obtained by techniques well known in the art. Such antibodies can be polyclonal or monoclonal. Polyclonal antibodies can be obtained, for example, by the methods

described in Ghose et al., *Methods in Enzymology*, Vol. 93, 326-327, 1983. For example, a lubricating polypeptide encoded by nucleotides 632-3453 of SEQ ID NO:2 is used as an immunogen to stimulate the production of polyclonal antibodies in the antisera of a rabbit. Similar methods can be used to raise antisera in animals such as goats, sheep, and rodents.

[0055] Monoclonal antibodies are obtained by the well known process described by Milstein and Kohler in *Nature*, 256:495-497, 1975, or as modified by Gerhard, *Monoclonal Antibodies*, Plenum Press, 1980, pages 370-371. Hybridomas are screened to identify those producing antibodies that are highly specific for a synovial lubricating polypeptide. Preferably, the antibody has an affinity of at least about 10^8 liters/mole and more preferably, an affinity of at least about 10^9 liters/mole. Monoclonal or polyclonal antibodies provide a means of rapidly purifying large quantities of recombinant lubricating polypeptides.

[0056] In addition to antibodies which are directed to the peptide core of a tribonectin, an antibody directed to a sugar portion or to a glycopeptide complex of a tribonectin is desirable. To generate an antibody to the peptide core, a peptide spanning amino acids 200-350 of SEQ ID NO:1 is used. Shorter peptides, e.g. 8-15 amino acids in length, which are identical to an 8-15 amino acid portion of amino acids 200-350 of SEQ ID NO:1 are also used to generate such antibodies. Other peptides to be used as immunogens for antibodies specific for the peptide core of a tribonectin include those which are in the region of amino acids 24-66 of SEQ ID NO:1, amino acids 105-155 of SEQ ID NO:1, or amino acids 156-199 of SEQ ID NO:1. To generate antibodies which bind to a glycosylated tribonectin polypeptide (but not a deglycosylated or nonglycosylated form), the immunogen is preferably a glycopeptide, the amino acid sequence of which spans a highly glycosylated portion of a tribonectin, e.g. a peptide with an amino acid sequence of residues 200-1140 of SEQ ID NO:1. Shorter glycopeptides, e.g., 8-15 amino acids in length, within the same highly glycosylated region are also used as immunogens. Methods of generating antibodies to highly glycosylated biomolecules are known in the art, e.g., as described by Schneerson et al., 1980, *J. Exp. Med.* 152:361-376.

[0057] Methods of Diagnosis

[0058] Osteoarthritis is a disease that develops slowly and is difficult to diagnose until its late stages when joint pain often compels an individual to seek medical treatment. Early diagnosis of osteoarthritis or a predisposition to develop the disease allows early intervention to prevent or reduce the development of advanced osteoarthritis. The invention provides methods of early detection of this disease or a predisposition to develop it by testing bodily fluids such as serum or urine for the presence of fragments of naturally-occurring tribonectins or the presence of fragments of MSF. Detection and quantitation of such peptides in biological fluids is well known in the art. For example, a standard sandwich ELISA assay is carried out using two different antibodies (e.g., a first antibody which binds to an oligosaccharide portion of the glycopeptide and a second antibody which binds to the peptide core of the glycopeptide) to a naturally-occurring tribonectin. Alternatively, standard protein sequencing by liquid chromatography and mass spectroscopy, as is described below, is used to detect MSF fragments in bio-

logical samples. A control value is a predetermined value associated with a negative diagnosis; alternatively, a control sample is a biological sample from a mammal known to be free of osteoarthritis. An increase in the amount compared to a control value or sample indicates that the mammal suffers from osteoarthritis or is predisposed to developing osteoarthritis.

[0059] Characterization of a Tribonectin from Human Synovial Fluid

[0060] Aliquots of synovial fluid from patients undergoing diagnostic arthroscopy and total knee replacement were collected and assayed in the friction apparatus. In both cases, the synovial fluid was aspirated prior to initiation of any surgical procedure and immediately centrifuged at $10,000\times g$ at 4°C . for 2 hrs to remove cellular debris. Samples which were contaminated with blood were discarded. Aliquots with normal lubricating ability were pooled and stored at -20°C .

[0061] Purification and Isolation of a Tribonectin

[0062] Human synovial fluid (200 ml) was filtered through $0.22\text{ }\mu\text{m}$ sterile filter units (Nalgene) at 4°C . over two days. Retentate was scraped off filter membranes and resuspended with 50 mM NaAc buffer, pH 5.5, to the original synovial fluid volume containing proteolytic inhibitors: 1 mM phenylmethyl sulfonyl fluoride (PMSF), 1 mM parachloromercuribenzoic acid (PCMB), and 10 mM ethylenediamine tetraacetate (EDTA). Digestion of hyaluronic acid was carried out at 37°C . by streptomyces hyaluronidase at 1 U/ml of resuspended synovial fluid. The digest was loaded on a DEAE column (Whatman International, Maidstone, UK) settled volume of 300 ml, equilibrated with NaAc buffer, 50 mM and washed with 1.5 L of the same buffer. The material with lubricating activity was eluted off of the DEAE matrix with 1 M NaCl. A 1 L wash was collected and concentrated via a 500 ml Amicon flow cell with an XM-100 membrane (mw cutoff 100 kDa). The concentrated sample was dialyzed against 25 mM phosphate buffer, pH 7.4, containing 0.15 M NaCl and 0.5 mM CaCl_2 .

[0063] The DEAE-bound concentrate was loaded onto a peanut agglutinin (PNA)-agarose affinity column with a settled bed volume of 25 ml, equilibrated at room temperature with 25 mM phosphate and 0.15 M NaCl buffer, pH 7.4. Unbound protein was eluted with the same buffer until absorbance at 230 and 280 nm decreased to background. Material with lubricating activity was maximally eluted in the presence of a step-wise gradient of α -lactose at a concentration of 0.07 M in 25 mM Tris and 0.15 M NaCl at pH 7.4. This material was loaded onto an Actigel ALD agarose (Sterogene Bioseparations, Arcadia, Calif.) coupled via amine groups to a murine monoclonal antibody against human fibronectin (Zymed Laboratories Inc., San Francisco, Calif.) to remove fibronectin as a contaminant. Eluted material was assayed for purity on SDS-PAGE (5-15% acrylamide) stained with Coomassie blue and by HPLC.

[0064] Protein electrophoresis standards were from GibcoBRL (Grand Island, N.Y.), and DNA ladder standard was from FMC Bioproducts (Rockland, Me.).

[0065] High Pressure Liquid Chromatography

[0066] A μ Bondpak C18 $3.9\times 150\text{ mm}$ column (Waters, Milford, Mass.) was eluted in reverse phase with 45% (v/v) methanol (Sigma) and 5% (v/v) acetonitrile (Aldrich) HPLC

grade at 1 ml/min at 35° C. The eluate was assayed by a photo diode array detector PDA 996 (Waters), and material in peak fractions were analyzed by purity plots calculated using Millenium 32 software (Waters).

[0067] Friction Apparatus

[0068] A standard friction apparatus (e.g., an apparatus described by Jay et al., 1992, *Conn. Tiss. Res.* 28:71-88 or Jay et al., 1998, *J. Biomed. Mater. Res.* 40:414-418). Natural latex was oscillated against a ring of polished glass with a constant contact area of 1.59 cm². The bearing system was axially loaded within a gimbals system free to rotate around two perpendicular horizontal axes. Latex and glass as bearing materials were chosen because they offer a flat surface with small asperity heights on the order of 0.05 mm. Latex, like cartilage, is compliant. Within the gimbals system, these surfaces possess near perfect co-planarity. Accordingly, fluid wedges were not generated and only a thin layer of boundary fluid was present. The entraining velocity (i.e., sliding speed) was 0.37 mm/sec with a constant contact pressure of 0.35×10⁶ N/m².

[0069] The friction apparatus recorded displacements of the gimbals system around the vertical loading axis through a linear displacement voltage transducer, the output voltage of which was directly proportional to the magnitude of the frictional torque. The peak to peak amplitude of this signal was related to μ by a previous calibration with known frictional torque.

[0070] Test surfaces were cleaned extensively before use. A 3.8×3.8 cm piece of latex strapped onto the stainless steel stud was washed under running distilled deionized water (DDW) for 2 min. It was then placed in a shallow bath of 0.9% NaCl physiological saline (PS). The glass slide was scrubbed with a 1% (v/v) 7× detergent (Flow Laboratories, McLean, Va.) solution in DDW for 10 min and then allowed to soak in the same solution at 100° C. A 5 min. scrubbing was also performed with the hot 7× solution followed by rinsing for 2-4 min. under running DDW.

[0071] The μ was measured at 35° C. and was preceded by a baseline measurement of the μ with PS. Lubrication was manifested by a reduction of μ relative to the μ of PS. Negative delta μ values indicate lubrication, whereas positive values indicate friction. Addition of 200 μ l of PS and later 200 μ l of test lubricant was followed by bringing the bearing surfaces close enough so that the solution wet both surfaces. After 5 min for equilibration, the latex-coated bearing was brought to rest on the glass as it was oscillating. Peak to peak voltages were automatically recorded after 1, 3 and 5 mins. At this point, the surfaces were separated for 2 min. and then brought back together for another 5 min session. The 3 and 5 min. μ values of the last two 5 min. sessions typically stabilized and were recorded.

[0072] Human serum fibronectin was purchased from Sigma Chemical (F0895, St. Louis, Mo.) and dialyzed against PS before use in the friction apparatus.

[0073] Boundary lubricants exert their effect by changing the physico-chemical characteristics of a surface. Bearing surfaces must generate a mutual repulsion in order to be lubricated in the boundary mode. Typical room temperature examples of boundary lubricants are graphite, teflon and molybdenum sulphide. Such compositions reduce friction between bearing surfaces, and therefore, are used as positive

controls in assay to measure the lubricating properties of tribonectins. Tribonectins are boundary lubricants that can have an amphipathic character by coating non-biologic hydrophobic surfaces such as latex. The oligosaccharide component of a tribonectin networks with the surrounding aqueous environment. When the ultimate and penultimate sugars are removed from a naturally-occurring tribonectin purified from synovial fluid, the lubricating ability is eliminated.

[0074] The latex:glass arthrotripsometer offers an expedient way to test purified biological lubricating factors repetitively with reproducibility. Natural latex and polished glass represent bearing surfaces with little if any variation in physico-chemical characteristic from test to test. By contrast, resected cartilage apposed to either polished glass or cartilage itself will experience deformation that cannot be accurately controlled. The μ observed in a cartilage-cartilage bearing lubricated by synovial fluid was between 0.005 and 0.024. The values of μ in the latex:glass system were appreciably higher and typically 0.04 or less. Differences in μ between the bearing materials are attributed to the 80% (w/w) water content of cartilage.

[0075] Protein Sequencing by Liquid Chromatography and Mass Spectrometry (LCMS)

[0076] Standard LCMS was carried out on tryptic digests of the purified lubricating material described above. Excised bands from 2 mm thick 5-20% gradient SDS-PAGE gels (Bio-rad Laboratories, Hercules, Calif.) containing the lubricating material was analyzed. The material was deglycosylated by NaNase III and O-glycosidase DS (Glyko, Novato, Calif.). Deglycosylation was carried out with the above enzymes at activities of 0.17 U/ml and 0.10 U/ml, respectively, for 18 hrs in the presence of 0.5 mg/ml of a tribonectin purified from synovial fluid. In all cases, the gel slices were cut through the middle of the band and were 16 mm³ in size. All contact surfaces were carefully cleaned with 50% (v/v) acetonitrile. Sequence data was entered into the BLAST GENBANK® search algorithm and matches identified.

[0077] Isolation and Culture of Human Synovial Fibroblasts

[0078] Human synovium with a normal appearance was obtained from a 30 year old white male undergoing arthroscopy. Within 1 hr after surgery, the synovial tissue explant was washed three times with Dulbecco's calcium- and magnesium-free phosphate-buffered saline (GIBCO). Pieces 2 mm³ in size were placed in Dulbecco's modification of Eagle's medium (GIBCO), supplemented with 100 U of penicillin and 100 μ g of streptomycin per ml (GIBCO), containing 4 mg/ml of Clostridiopeptidase A (Worthington Biochemical CLS, 125-200 U/mg) sterilized through a 0.22 μ m filter (Nalge). The tissue fragments were further divided with scissors in a 100 mm plastic petri dish (Falcon) and incubated for 4 hrs in 20 ml of medium at 37° C. in a moist atmosphere of 5% carbon dioxide and 95% air.

[0079] The digest was well mixed many times by aspiration into and expulsion from a Pasteur pipette. An equal volume of 0.05% trypsin and 0.02% EDTA in modified Puck's Saline A (GIBCO) were added and incubation continued for a further hour under the same conditions. The suspension was centrifuged 10 min at 400×g at 23° C. and

washed three times each with 40 ml of calcium- and magnesium-free phosphate-buffered saline. The pellet was suspended in modified Eagle's medium (20 ml) supplemented with 10% fetal bovine serum (Flow Laboratories), 100 U of penicillin, and 100 mg of streptomycin per ml. Two milliliters of this final mixture were plated per 60 mm plastic petri dish (Falcon). Synovial fibroblasts were grown to confluence and cells harvested. Human skin fibroblasts (American Type Culture Collection (ATCC) Designation CCD-1099SK; ATCC, Manassas, Va.) which served as a control were also grown and harvested using the above procedure.

[0080] RNA Extraction and RT-PCR Analyses

[0081] RNA from synovial and skin fibroblasts was purified by RNeasy mini-columns and reagents (Qiagen, Crawley, Ltd., UK). Contaminating genomic DNA was removed by DNase (RNase free) (Qiagen). First strand cDNA was synthesized by reverse transcription and PCR amplification using the following oligonucleotide primers. MSF-exon 6 forward primer 5'-CCAAACCACAGTTGTAGATGAAGC-3' (SEQ ID NO:15) and MSF-exon 6 reverse primer 5'-GCGGAAGTAGTCTTCTCTTTCCATCAG-3' (SEQ ID NO:16). These primers correspond to nucleotide position numbers 674-698 and 953-926, respectively, of the human MSF gene (SEQ ID NO:2; GENBANK® accession number U70136). Thermal cycling conditions were 42° C. for 12 mins., 95° C. for 10 mins., followed by 43 cycles between 94° C.×20 secs and 55-65° C.×30-90 secs. A final extension for 7 mins was at 72° C. (Perkin Elmer Biosystems).

[0082] Alternative Splice Variant of MSF is a Tribonection

[0083] A lubricating polypeptide was purified from human synovial fluid using standard biochemical methods followed by affinity chromatography with peanut agglutinin. The final fraction, which solely possessed lubricating ability, contained a product with an apparent molecular weight of 280 kDa. Components with a molecular weight in excess of 280 kDa were not observed. LCMS performed on tryptic fragments from the 280 kDa excised band indicated the presence of two different proteins that matched in the BLAST search algorithm to fibronectin precursor and MSF (GenBank Accession No. U70136). Sequences of MSF were identified from both native and deglycosylated lubricating polypeptides. Accordingly, the purification scheme was terminated with an anti-fibronectin column resulting in the elimination of fibronectin as an impurity (as assayed by C18 analytical HPLC and purity plot analysis). In addition, lower molecular weight bands at 70 and 160 kDa on SDS-PAGE were absent from the purified tribonection preparation eluting from the anti-fibronectin column. The purified tribonection assayed in the friction apparatus was found to display boundary lubricating activity similar to that of whole synovial fluid (Table

5). By contrast, purified serum fibronectin raised friction indicating that synovial fluid lubricating ability was mediated by the purified tribonection.

TABLE 5

Friction coefficients for a tribonection purified from human synovial fluid and fibronectin (Mean ± SD; N = 3)			
LUBRICANT*	μ	μ (PS**)	$\Delta\mu$
Tribonection	0.047 ± .006	0.131 ± .007	-0.084 ± .004
HSF†	0.040 ± .005	0.135 ± .009	-0.095 ± .011
Fibronectin	0.181	0.136	+0.045 ± .005

*Tested at a concentration of 250 μ g/ml in PS.

**Physiological saline.

†Post-mortem human synovial fluid

[0084] Furthermore, LCMS of tryptic fragments identified portions of exons 6 through 9 of MSF, inclusively. Purified tribonection reacted to peanut agglutinin indicating the presence of β (1-3)Gal-GalNAC oligosaccharides by virtue of its purification. An increase in electrophoretic mobility was observed after digestion with NaNase III and O-glycosidase DS, indicating that the purified tribonection is highly glycosylated via O-linked oligosaccharides. The apparent molecular weight of deglycosylated tribonection purified from synovial fluid was 120 kDa.

[0085] RT-PCR analysis was completed using primers specific for nucleotide sequences encoding the N-terminal end of exon 6 of MSF. RT-PCR's using human synovial fibroblast RNA generated a 280 bp product, the predicted distance between the designed primers. Similar experiments without reverse transcriptase did not generate this product indicating that the RNA was free of genomic DNA. Purified RNA from skin fibroblasts did not produce any product using the same primers.

[0086] MSF was first isolated from human monocytes; a 25 kDa fragment of MSF was found to stimulate the development of megakaryocytes. MSF precursor protein is 1404 residues in size and constructed from 12 exons. Exon 6 appears to encode a centrally located mucin that is 940 residues in length. Exon 6 has homology to vitronectin, exons 2 and 3 appear homologous to somatomedin B-like regions, and exons 8, 9 are similar to hemopexin-like regions in vitronectin. Hemopexin is a serum heme scavenging protein that interacts with hyaluronate.

[0087] A tribonection purified from synovial fluid and an articular cartilage superficial zone protein (SZP) purified from articular cartilage share sequence identity with MSF but differ in their apparent molecular weights and amino acid sequences.

[0088] Other embodiments are within the following claims.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 16

<210> SEQ ID NO 1

<211> LENGTH: 1404

-continued

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 1

Met Ala Trp Lys Thr Leu Pro Ile Tyr Leu Leu Leu Leu Ser Val
1 5 10 15
Phe Val Ile Gln Gln Val Ser Ser Gln Asp Leu Ser Ser Cys Ala Gly
20 25 30
Arg Cys Gly Glu Gly Tyr Ser Arg Asp Ala Thr Cys Asn Cys Asp Tyr
35 40 45
Asn Cys Gln His Tyr Met Glu Cys Cys Pro Asp Phe Lys Arg Val Cys
50 55 60
Thr Ala Glu Leu Ser Cys Lys Gly Arg Cys Phe Glu Ser Phe Glu Arg
65 70 75 80
Gly Arg Glu Cys Asp Cys Asp Ala Gln Cys Lys Lys Tyr Asp Lys Cys
85 90 95
Cys Pro Asp Tyr Glu Ser Phe Cys Ala Glu Val His Asn Pro Thr Ser
100 105 110
Pro Pro Ser Ser Lys Lys Ala Pro Pro Pro Ser Gly Ala Ser Gln Thr
115 120 125
Ile Lys Ser Thr Thr Lys Arg Ser Pro Lys Pro Pro Asn Lys Lys Lys
130 135 140
Thr Lys Lys Val Ile Glu Ser Glu Glu Ile Thr Glu Glu His Ser Val
145 150 155 160
Ser Glu Asn Gln Glu Ser Ser Ser Ser Ser Ser Ser Ser Ser Ser
165 170 175
Ser Thr Ile Trp Lys Ile Lys Ser Ser Lys Asn Ser Ala Ala Asn Arg
180 185 190
Glu Leu Gln Lys Lys Leu Lys Val Lys Asp Asn Lys Lys Asn Arg Thr
195 200 205
Lys Lys Lys Pro Thr Pro Lys Pro Pro Val Val Asp Glu Ala Gly Ser
210 215 220
Gly Leu Asp Asn Gly Asp Phe Lys Val Thr Thr Pro Asp Thr Ser Thr
225 230 235 240
Thr Gln His Asn Lys Val Ser Thr Ser Pro Lys Ile Thr Thr Ala Lys
245 250 255
Pro Ile Asn Pro Arg Pro Ser Leu Pro Pro Asn Ser Asp Thr Ser Lys
260 265 270
Glu Thr Ser Leu Thr Val Asn Lys Glu Thr Thr Val Glu Thr Lys Glu
275 280 285
Thr Thr Thr Thr Asn Lys Gln Thr Ser Thr Asp Gly Lys Glu Lys Thr
290 295 300
Thr Ser Ala Lys Glu Thr Gln Ser Ile Glu Lys Thr Ser Ala Lys Asp
305 310 315 320
Leu Ala Pro Thr Ser Lys Val Leu Ala Lys Pro Thr Pro Lys Ala Glu
325 330 335
Thr Thr Thr Lys Gly Pro Ala Leu Thr Thr Pro Lys Glu Pro Thr Pro
340 345 350
Thr Thr Pro Lys Glu Pro Ala Ser Thr Thr Pro Lys Glu Pro Thr Pro
355 360 365
Thr Thr Ile Lys Ser Ala Pro Thr Thr Pro Lys Glu Pro Ala Pro Thr
370 375 380

Thr	Thr	Lys	Ser	Ala	Pro	Thr	Thr	Pro	Lys	Glu	Pro	Ala	Pro	Thr	Thr	385	400
																	390
Thr	Lys	Glu	Pro	Ala	Pro	Thr	Thr	Pro	Lys	Glu	Pro	Ala	Pro	Thr	Thr	405	410
																	415
Thr	Lys	Glu	Pro	Ala	Pro	Thr	Thr	Thr	Lys	Ser	Ala	Pro	Thr	Thr	Pro	420	425
																	430
Lys	Glu	Pro	Ala	Pro	Thr	Thr	Pro	Lys	Lys	Pro	Ala	Pro	Thr	Thr	Pro	435	440
																	445
Lys	Glu	Pro	Ala	Pro	Thr	Thr	Pro	Lys	Glu	Pro	Thr	Pro	Thr	Thr	Pro	450	455
																	460
Lys	Glu	Pro	Ala	Pro	Thr	Thr	Lys	Glu	Pro	Ala	Pro	Thr	Thr	Pro	Lys	465	480
																	470
Glu	Pro	Ala	Pro	Thr	Ala	Pro	Lys	Lys	Pro	Ala	Pro	Thr	Thr	Pro	Lys	485	490
																	495
Glu	Pro	Ala	Pro	Thr	Thr	Pro	Lys	Glu	Pro	Ala	Pro	Thr	Thr	Thr	Lys	500	505
																	510
Glu	Pro	Ser	Pro	Thr	Thr	Pro	Lys	Glu	Pro	Ala	Pro	Thr	Thr	Thr	Lys	515	520
																	525
Ser	Ala	Pro	Thr	Thr	Thr	Lys	Glu	Pro	Ala	Pro	Thr	Thr	Thr	Lys	Ser	530	535
																	540
Ala	Pro	Thr	Thr	Pro	Lys	Glu	Pro	Ser	Pro	Thr	Thr	Thr	Lys	Glu	Pro	545	550
																	555
Ala	Pro	Thr	Thr	Pro	Lys	Glu	Pro	Ala	Pro	Thr	Thr	Pro	Lys	Lys	Pro	565	570
																	575
Ala	Pro	Thr	Thr	Pro	Lys	Glu	Pro	Ala	Pro	Thr	Thr	Pro	Lys	Glu	Pro	580	585
																	590
Ala	Pro	Thr	Thr	Thr	Lys	Lys	Pro	Ala	Pro	Thr	Ala	Pro	Lys	Glu	Pro	595	600
																	605
Ala	Pro	Thr	Thr	Pro	Lys	Glu	Thr	Ala	Pro	Thr	Thr	Pro	Lys	Lys	Leu	610	615
																	620
Thr	Pro	Thr	Thr	Pro	Glu	Lys	Leu	Ala	Pro	Thr	Thr	Pro	Glu	Lys	Pro	625	630
																	635
Ala	Pro	Thr	Thr	Pro	Glu	Glu	Leu	Ala	Pro	Thr	Thr	Pro	Glu	Glu	Pro	645	650
																	655
Thr	Pro	Thr	Thr	Pro	Glu	Glu	Pro	Ala	Pro	Thr	Thr	Pro	Lys	Ala	Ala	660	665
																	670
Ala	Pro	Asn	Thr	Pro	Lys	Glu	Pro	Ala	Pro	Thr	Thr	Pro	Lys	Glu	Pro	675	680
																	685
Ala	Pro	Thr	Thr	Pro	Lys	Glu	Pro	Ala	Pro	Thr	Thr	Pro	Lys	Glu	Thr	690	695
																	700
Ala	Pro	Thr	Thr	Pro	Lys	Gly	Thr	Ala	Pro	Thr	Thr	Leu	Lys	Glu	Pro	705	710
																	715
Ala	Pro	Thr	Thr	Pro	Lys	Lys	Pro	Ala	Pro	Lys	Glu	Leu	Ala	Pro	Thr	725	730
																	735
Thr	Thr	Lys	Glu	Pro	Thr	Ser	Thr	Thr	Ser	Asp	Lys	Pro	Ala	Pro	Thr	740	745
																	750
Thr	Pro	Lys	Gly	Thr	Ala	Pro	Thr	Thr	Pro	Lys	Glu	Pro	Ala	Pro	Thr	755	760
																	765
Thr	Pro	Lys	Glu	Pro	Ala	Pro	Thr	Thr	Pro	Lys	Gly	Thr	Ala	Pro	Thr	770	775
																	780

-continued

Thr	Leu	Lys	Glu	Pro	Ala	Pro	Thr	Thr	Pro	Lys	Lys	Pro	Ala	Pro	Lys	
785					790					795					800	
Glu	Leu	Ala	Pro	Thr	Thr	Thr	Lys	Gly	Pro	Thr	Ser	Thr	Thr	Ser	Asp	
				805					810					815		
Lys	Pro	Ala	Pro	Thr	Thr	Pro	Lys	Glu	Thr	Ala	Pro	Thr	Thr	Pro	Lys	
				820				825						830		
Glu	Pro	Ala	Pro	Thr	Thr	Pro	Lys	Lys	Pro	Ala	Pro	Thr	Thr	Pro	Glu	
				835				840						845		
Thr	Pro	Pro	Pro	Thr	Thr	Ser	Glu	Val	Ser	Thr	Pro	Thr	Thr	Thr	Lys	
				850			855					860				
Glu	Pro	Thr	Thr	Ile	His	Lys	Ser	Pro	Asp	Glu	Ser	Thr	Pro	Glu	Leu	
865					870					875					880	
Ser	Ala	Glu	Pro	Thr	Pro	Lys	Ala	Leu	Glu	Asn	Ser	Pro	Lys	Glu	Pro	
				885					890					895		
Gly	Val	Pro	Thr	Thr	Lys	Thr	Pro	Ala	Ala	Thr	Lys	Pro	Glu	Met	Thr	
				900				905						910		
Thr	Thr	Ala	Lys	Asp	Lys	Thr	Thr	Glu	Arg	Asp	Leu	Arg	Thr	Thr	Pro	
				915				920					925			
Glu	Thr	Thr	Thr	Ala	Ala	Pro	Lys	Met	Thr	Lys	Glu	Thr	Ala	Thr	Thr	
				930			935					940				
Thr	Glu	Lys	Thr	Thr	Glu	Ser	Lys	Ile	Thr	Ala	Thr	Thr	Thr	Gln	Val	
945					950					955					960	
Thr	Ser	Thr	Thr	Thr	Gln	Asp	Thr	Thr	Pro	Phe	Lys	Ile	Thr	Thr	Leu	
				965					970						975	
Lys	Thr	Thr	Thr	Leu	Ala	Pro	Lys	Val	Thr	Thr	Thr	Lys	Lys	Thr	Ile	
				980					985					990		
Thr	Thr	Thr	Glu	Ile	Met	Asn	Lys	Pro	Glu	Glu	Thr	Ala	Lys	Pro	Lys	
			995				1000						1005			
Asp	Arg	Ala	Thr	Asn	Ser	Lys	Ala	Thr	Thr	Pro	Lys	Pro	Gln	Lys	Pro	
			1010				1015					1020				
Thr	Lys	Ala	Pro	Lys	Lys	Pro	Thr	Ser	Thr	Lys	Lys	Pro	Lys	Thr	Met	
1025					1030					1035					1040	
Pro	Arg	Val	Arg	Lys	Pro	Lys	Thr	Thr	Pro	Thr	Pro	Arg	Lys	Met	Thr	
				1045					1050					1055		
Ser	Thr	Met	Pro	Glu	Leu	Asn	Pro	Thr	Ser	Arg	Ile	Ala	Glu	Ala	Met	
				1060					1065					1070		
Leu	Gln	Thr	Thr	Thr	Arg	Pro	Asn	Gln	Thr	Pro	Asn	Ser	Lys	Leu	Val	
		1075					1080						1085			
Glu	Val	Asn	Pro	Lys	Ser	Glu	Asp	Ala	Gly	Gly	Ala	Glu	Gly	Glu	Thr	
		1090					1095						1100			
Pro	His	Met	Leu	Leu	Arg	Pro	His	Val	Phe	Met	Pro	Glu	Val	Thr	Pro	
1105					1110					1115					1120	
Asp	Met	Asp	Tyr	Leu	Pro	Arg	Val	Pro	Asn	Gln	Gly	Ile	Ile	Ile	Asn	
				1125					1130					1135		
Pro	Met	Leu	Ser	Asp	Glu	Thr	Asn	Ile	Cys	Asn	Gly	Lys	Pro	Val	Asp	
				1140					1145					1150		
Gly	Leu	Thr	Thr	Leu	Arg	Asn	Gly	Thr	Leu	Val	Ala	Phe	Arg	Gly	His	
		1155					1160						1165			
Tyr	Phe	Trp	Met	Leu	Ser	Pro	Phe	Ser	Pro	Pro	Ser	Pro	Ala	Arg	Arg	
		1170					1175						1180			
Ile	Thr	Glu	Val	Trp	Gly	Ile	Pro	Ser	Pro	Ile	Asp	Thr	Val	Phe	Thr	

-continued

1185	1190	1195	1200
Arg Cys Asn Cys Glu Gly Lys Thr Phe Phe Phe Lys Asp Ser Gln Tyr	1205	1210	1215
Trp Arg Phe Thr Asn Asp Ile Lys Asp Ala Gly Tyr Pro Lys Pro Ile	1220	1225	1230
Phe Lys Gly Phe Gly Gly Leu Thr Gly Gln Ile Val Ala Ala Leu Ser	1235	1240	1245
Thr Ala Lys Tyr Lys Asn Trp Pro Glu Ser Val Tyr Phe Phe Lys Arg	1250	1255	1260
Gly Gly Ser Ile Gln Gln Tyr Ile Tyr Lys Gln Glu Pro Val Gln Lys	1265	1270	1275
Cys Pro Gly Arg Arg Pro Ala Leu Asn Tyr Pro Val Tyr Gly Glu Met	1285	1290	1295
Thr Gln Val Arg Arg Arg Arg Phe Glu Arg Ala Ile Gly Pro Ser Gln	1300	1305	1310
Thr His Thr Ile Arg Ile Gln Tyr Ser Pro Ala Arg Leu Ala Tyr Gln	1315	1320	1325
Asp Lys Gly Val Leu His Asn Glu Val Lys Val Ser Ile Leu Trp Arg	1330	1335	1340
Gly Leu Pro Asn Val Val Thr Ser Ala Ile Ser Leu Pro Asn Ile Arg	1345	1350	1355
Lys Pro Asp Gly Tyr Asp Tyr Tyr Ala Phe Ser Lys Asp Gln Tyr Tyr	1365	1370	1375
Asn Ile Asp Val Pro Ser Arg Thr Ala Arg Ala Ile Thr Thr Arg Ser	1380	1385	1390
Gly Gln Thr Leu Ser Lys Val Trp Tyr Asn Cys Pro	1395	1400	

<210> SEQ ID NO 2

<211> LENGTH: 5041

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 2

gcggccgcga ctattcggta cctgaaaaca acgatggcat ggaaaacact tccattttac	60
ctgttgtgtg tgctgtctgt ttctgtgatt cagcaagttt catctcaaga tttatcaagc	120
tgtgcaggga gatgtgggga agggatttct agagatgcc cctgcaactg tgattataac	180
tgtcaacact acatggagtg ctgccctgat ttcaagagag tctgcaactg ggagctttcc	240
tgtaaaaggcc gctgctttga gtccttcgag agagggaggg agtgtgactg cgacgcccac	300
tgtaagaagt atgacaagtg ctgtcccgat tatgagagtt tctgtgcaga agtgcataat	360
cccacatcac caccatcttc aaagaaagca cctccacctt caggagcatc tcaaaccatc	420
aatcaacaa ccaaacgttc acccaaacca ccaacaaga agaagactaa gaaagttata	480
gaatcagagg aaataacaga agaacattct gtttctgaaa atcaagagtc ctctctctcc	540
tcctcctctt cctctcttct ttcaacaatt tggaatatca agtcttccaa aaattcagct	600
gctaatagag aattacagaa gaaactcaaa gtaaaagata acaagaagaa cagaactaaa	660
aagaaaccta ccccaaaccc accagttgta gatgaagctg gaagtggatt ggacaatggt	720
gacttcaagg tcacaactcc tgacacgtct accaccaac acaataaagt cagcacatct	780
cccaagatca caacagcaaa accaataaat ccagaccac gtcttccacc taattctgat	840

-continued

acatctaaag agacgtcttt gacagtgaat aaagagacaa cagttgaaac taaagaaact	900
actacaacaa ataaacagac ttcaactgat ggaaaagaga agactacttc cgctaaagag	960
acacaaagta tagagaaaac atctgctaaa gatttagcac ccacatctaa agtgctggct	1020
aaacctacac ccaaagctga aactacaacc aaaggccctg ctctcaccac tccaaggag	1080
cccacgccc cactcccaa ggagcctgca tctaccacac ccaaagagcc cacacctacc	1140
accatcaagt ctgcaccac caccccaag gagcctgcac ccaccaccac caagtctgca	1200
cccaccactc ccaaggagcc tgcaccacc accaccaagg agcctgcacc caccactccc	1260
aaggagcctg caccaccac caccaaggag cctgcacca ccaccacaa gtctgcacc	1320
accactcca aggagcctgc acccaccac ccaagaagc ctgcccacac taccccaag	1380
gagcctgcac ccaccactcc caaggagcct acaccacca ctccaagga gcctgcacc	1440
accaccaagg agcctgcacc caccactccc aaagagcctg caccactgc cccaagaag	1500
cctgcccac ctaccccac ggagcctgca cccaccactc ccaaggagcc tgcaccacc	1560
accaccaagg agccttcacc caccactccc aaggagcctg caccaccac caccaagtct	1620
gcaccacca ctaccaagga gcctgcacc accactacca agtctgcacc caccactccc	1680
aaggagcctt caccaccac caccaaggag cctgcacca cactcccaa ggagcctgca	1740
cccaccacc ccaagaagcc tgcccacact accccaagg agcctgcacc caccactccc	1800
aaggaacctg caccaccac caccaagaag cctgcacca ccgctccaa agagcctgcc	1860
ccaactacc ccaaggagac tgcaccacc accccaaga agctcagcc caccacccc	1920
gagaagctcg caccaccac cctgagaag ccgcaccca ccaccctga ggagctcgca	1980
cccaccacc ctgaggagcc cacaccacc acccctgagg agcctgctcc caccactccc	2040
aaggcagcgg ctccaacac ccctaaggag cctgctccaa ctaccctaa ggagcctgct	2100
ccaactacc ctaaggagcc tgctccaact acccctaagg agactgctcc aactaccct	2160
aaagggactg ctccaactac cctcaaggaa cctgcacca ctactccaa gaagcctgcc	2220
cccaaggagc ttgcaccac caccaccaag gagccacat ccaccctc tgacaagccc	2280
gctccaacta ccctaaggg gactgctcca actacccta aggagcctgc tccaactacc	2340
cctaaggagc ctgctccaac taccctaag gggactgctc caactaccct caaggaacct	2400
gcaccacta ctccaagaa gcctgcccc aaggagcttg caccaccac caccaagggg	2460
cccacatcca ccacctctga caagcctgct ccaactacac ctaaggagac tgctccaact	2520
acccccaagg agcctgcacc cactacccc aagaagcctg ctccaactac tcctgagaca	2580
cctcctcaa ccacttcaga ggtctctact ccaactacca ccaaggagcc taccactatc	2640
cacaaaagcc ctgatgaatc aactcctgag ctttctgcag aaccacacc aaaagctctt	2700
gaaaacagtc ccaaggaacc tgggtgtacct acaactaaga ctctgcagc gactaaacct	2760
gaaatgacta caacagctaa agacaagaca acagaagag acttacgtac tacacctgaa	2820
actacaactg ctgcacctaa gatgacaaaa gagacagcaa ctacaacaga aaaaactacc	2880
gaatccaaaa taacagctac aaccacacaa gtaacatcta ccacaactca agataccaca	2940
ccattcaaaa ttactactct taaaacaact actcttgac ccaagtaac tacaacaaaa	3000
aagacaatta ctaccactga gattatgaac aaacctgaag aaacagctaa accaaaagac	3060
agagctacta attctaagc gacaactcct aaacctcaaa agccaaccaa agcacccaaa	3120

-continued

aaacccactt ctaccaaaaa gccaaaaaca atgcctagag tgagaaaacc aaagacgaca	3180
ccaactcccc gcaagatgac atcaacaatg ccagaattga accctacctc aagaatagca	3240
gaagccatgc tccaaaccac caccagacct aaccaaactc caaactccaa actagttgaa	3300
gtaaatccaa agagtgaaga tgcagggtgt gctgaaggag aaacacctca tatgcttctc	3360
aggcccatg tgttcacgac tgaagttact cccgacatgg attacttacc gagagtaccc	3420
aatcaaggca ttatcatcaa tcccatgctt tccgatgaga ccaatatatg caatggtaag	3480
ccagtagatg gactgactac ttgcgcaat gggacattag ttgcattccg aggtcattat	3540
ttctggatgc taagtccatt cagtccacca tctccagctc gcagaattac tgaagtttg	3600
ggatttcctt ccccatgta tactgttttt actaggtgca actgtgaagg aaaaactttc	3660
ttctttaagg attctcagta ctggcgtttt accaatgata taaaagatgc agggtagccc	3720
aaaccaatth tcaaaggatt tggaggacta actggacaaa tagtggcagc gctttcaaca	3780
gctaaatata agaactggcc tgaatctgtg tatTTTTTca agagaggtgg cagcattcag	3840
cagtatatth ataaacagga acctgtacag aagtgccttg gaagaaggcc tgctctaaat	3900
tatccagtgt atggagaaat gacacaggtt aggagacgtc gctttgaacg tgctatagga	3960
ccttctcaaa cacacacat cagaattcaa tattcacctg ccagactggc ttatcaagac	4020
aaagggtgcc ttcataatga agttaagtgt agtatactgt ggagaggact tccaaatgtg	4080
gttacctcag ctatatcact gcccacatc agaaaacctg acggctatga ttactatgcc	4140
ttttctaaag atcaatacta taacattgat gtgcctagta gaacagcaag agcaattact	4200
actcgttctg ggcagacctt atccaaaagc tggtagaact gtccttagac tgatgagcaa	4260
aggaggagtc aactaatgaa gaaatgaata ataaattttg aactgaaaa acattttatt	4320
aataaagaat attgacatga gtataccagt ttatatataa aaatgttttt aaacttgaca	4380
atcattacac taaaacagat ttgataatct tattcacagt tgttattggt tacagaccat	4440
ttaattaata tttcctctgt ttattcctcc tctccctccc attgcatggc tcacacctgt	4500
aaaagaaaa agaatacaat tgaatatatc ttttaagaat tcaaaactag tgtattcact	4560
taccctagtt cattataaaa aatatctagg cattgtggat ataaaactgt tgggtattct	4620
acaacttcaa tggaaattat tacaagcaga ttaatccctc tttttgtgac acaagtacaa	4680
tctaaaagtt atattggaaa acatggaaat attaaaattt tacactttta ctagctaaaa	4740
cataatcaca aagctttatc gtgttgata aaaaaattaa caatataatg gcaataggta	4800
gagatacaac aaatgaatat aacactataa cacttcatat tttccaaatc ttaatttgga	4860
tttaagggaag aaatcaataa atataaaata taagcacata tttattatat atctaaggta	4920
tacaaatctg tctacatgaa gtttacagat tggtaaatat cacctgctca acatgtaatt	4980
atttaataaa actttggaac attaaaaaaa taaattggag gcttaaaaaa aaaaaaaaaa	5040
a	5041

<210> SEQ ID NO 3
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 3

Lys Glu Pro Ala Pro Thr Thr

-continued

1 5

<210> SEQ ID NO 4
<211> LENGTH: 6
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: VARIANT
<222> LOCATION: (1)...(6)
<223> OTHER INFORMATION: Xaa = Any Amino Acid

<400> SEQUENCE: 4

Xaa Xaa Thr Thr Thr Xaa
1 5

<210> SEQ ID NO 5
<211> LENGTH: 6
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 5

Glu Pro Ala Pro Thr Thr
1 5

<210> SEQ ID NO 6
<211> LENGTH: 6
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 6

Pro Thr Thr Lys Glu Pro
1 5

<210> SEQ ID NO 7
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 7

agatttatca agctgtgcag ggag 24

<210> SEQ ID NO 8
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 8

tttacaggaa agctccgcag tg 22

<210> SEQ ID NO 9
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 9

tcaaggtcac aactcctgac acg 23

<210> SEQ ID NO 10
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 10

-continued

ctctcggtaa gtaatccatg tcgg 24

<210> SEQ ID NO 11
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 11

ttgttgctgc tgtctgtttt cg 22

<210> SEQ ID NO 12
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 12

tggataaggt ctgccagaa cgag 24

<210> SEQ ID NO 13
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 13

tcaaggtcac aactcctgac acg 23

<210> SEQ ID NO 14
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 14

gatggtgtgt gtttgagaag gtcc 24

<210> SEQ ID NO 15
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 15

ccaaaccacc agttgtagat gaagc 25

<210> SEQ ID NO 16
<211> LENGTH: 28
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 16

gcggaagtag tcttctcttt tccatcag 28

What is claimed is:

1. A tribonectin comprising a polypeptide the amino acid sequence of which comprises at least one but less than 76 subunits, wherein

- (a) each subunit comprises at least 7 amino acids; and
- (b) the amino acid sequence of said subunit is at least 50% identical to SEQ ID NO:3, wherein a non-identical amino acid is a conservative amino acid substitution.
2. The tribonectin of claim 1, wherein the amino acid sequence of said subunit is SEQ ID NO:3.
3. The tribonectin of claim 1, wherein said tribonectin further comprises one or more repeats of the amino acid sequence of SEQ ID NO:4.
4. The tribonectin of claim 1, wherein said tribonectin is characterized as reducing the coefficient of friction between bearing surfaces.
5. The tribonectin of claim 1, wherein said tribonectin is characterized as reducing the coefficient of friction between bearing surfaces in vitro.
6. The tribonectin of claim 1, wherein said tribonectin is characterized as reducing the coefficient of friction between bearing surfaces in vivo.
7. The tribonectin of claim 1, wherein said tribonectin does not substantially increase the viscosity of a solution to which it is added.
8. The tribonectin of claim 1, wherein said tribonectin comprises an O-linked oligosaccharide.
9. The tribonectin of claim 1, wherein said oligosaccharide is an N-acetylgalactosamine-galactose.
10. The tribonectin of claim 1, wherein at least 10% of said tribonectin is glycosylated.
11. The tribonectin of claim 1, wherein at least 40% of said tribonectin is glycosylated.
12. The tribonectin of claim 1, wherein the molecular weight of said tribonectin is in the range of 200-280 kDa.
13. The tribonectin of claim 1, wherein said polypeptide comprises a fragment of megakaryocyte stimulating factor.
14. The tribonectin of claim 1, wherein said polypeptide comprises an amino acid sequence that is at least 50% identical to the sequence of residues 200-1140, inclusive, of SEQ ID NO:1.
15. The tribonectin of claim 14, wherein said polypeptide comprises the amino acid sequence of residues 200-1140, inclusive, of SEQ ID NO:1.
16. The tribonectin of claim 1, wherein said polypeptide comprises an amino acid sequence that is at least 50% identical to the sequence of residues 200-1167, inclusive, of SEQ ID NO:1.
17. The tribonectin of claim 16, wherein said polypeptide comprises the amino acid sequence of residues 200-1167, inclusive, of SEQ ID NO:1.
18. The tribonectin of claim 1, wherein said polypeptide comprises an amino acid sequence that is at least 50% identical to the sequence of residues 200-1212, inclusive, of SEQ ID NO:1.
19. The tribonectin of claim 18, wherein said polypeptide comprises the amino acid sequence of residues 200-1212, inclusive, of SEQ ID NO:1.
20. The tribonectin of claim 1, wherein said polypeptide comprises an amino acid sequence that is at least 50% identical to the sequence of residues 200-1263, inclusive, of SEQ ID NO:1.

21. The tribonectin of claim 20, wherein said polypeptide comprises the amino acid sequence of residues 200-1263, inclusive, of SEQ ID NO:1.

22. The tribonectin of claim 1, wherein said polypeptide lacks the amino acid sequence of residues 1-24, inclusive, of SEQ ID NO:1.

23. The tribonectin of claim 1, wherein said polypeptide lacks the amino acid sequence of residues 67-104, inclusive, of SEQ ID NO:1.

24. An isolated nucleic acid molecule encoding a tribonectin.

25. The nucleic acid of claim 24, wherein said nucleic acid comprises the sequence of nucleotides 631-3453, inclusive, of SEQ ID NO:2.

26. A method of lubricating a mammalian joint, comprising contacting said joint with the tribonectin of claim 1.

27. The method of claim 26, wherein said joint is an articulating joint of a human.

28. The method of claim 26, wherein said joint is an articulating joint of a dog.

29. The method of claim 26, wherein said joint is an articulating joint of a horse.

30. The method of claim 26, wherein said tribonectin is administered intra-articularly.

31. A method of lubricating a mammalian joint, comprising contacting said joint with the nucleic acid of claim 24.

32. A biocompatible composition comprising a tribonectin, wherein said composition is in a form suitable for the inhibition of tissue adhesion formation.

33. The composition of claim 32, wherein said tribonectin is in the form of a membrane, foam, gel, or fiber.

34. A method inhibiting adhesion formation between a first surface and a second surface in a mammal, said method comprising placing a tribonectin between said first and second surfaces in an amount sufficient to prevent adhesion of said surfaces in said mammal.

35. The method of claim 34, wherein said first surface and said second surface are both injured tissues of said mammal.

36. The method of claim 34, wherein said first or said second surface is an artificial device.

37. The method of claim 36, wherein said artificial device is an orthopedic implant.

38. The method of claim 34, wherein said tribonectin is in the form of a membrane, foam, gel, or fiber.

39. The method of claim 34, wherein said injury is due to a surgical incision.

40. The method of claim 34, wherein said injury is due to trauma.

41. A method for diagnosing an osteoarthritis or a pre-disposition thereto in a mammal, comprising measuring the amount of a fragment of megakaryocyte stimulating factor in a biological sample derived from said mammal, wherein an increase in said amount compared to a control indicates that said mammal suffers from osteoarthritis or is predisposed to developing osteoarthritis.

42. The method of claim 41, wherein said biological sample is synovial fluid, blood, serum, or urine.

43. The method of claim 41, wherein said MSF fragment comprises the amino acid sequence of SEQ ID NO:3.

44. The method of claim 41, wherein said MSF fragment comprises the amino acid sequence of SEQ ID NO:5.

45. The method of claim 41, wherein said MSF fragment comprises the amino acid sequence of SEQ ID NO:6.

专利名称(译)	Tribonectins		
公开(公告)号	US20040229804A1	公开(公告)日	2004-11-18
申请号	US10/858595	申请日	2004-06-01
[标]申请(专利权)人(译)	罗德岛医院		
申请(专利权)人(译)	罗德岛医院		
当前申请(专利权)人(译)	罗德岛医院		
[标]发明人	JAY GREGORY D		
发明人	JAY, GREGORY D.		
IPC分类号	G01N33/53 A61K38/00 A61K38/12 A61K48/00 A61L27/00 A61P19/02 A61P41/00 A61P43/00 C07K14/47 C07K14/475 C12N15/09 A61K38/17		
CPC分类号	A61K38/00 A61K48/00 C07K14/47 A61K38/1709 C07K14/475 A61P17/02 A61P19/02 A61P19/10		
其他公开文献	US7618941		
外部链接	Espacenet USPTO		

摘要(译)

本发明的特征在于三连蛋白和通过将三连蛋白直接给予受损或关节炎关节而进行的补骨脂补充方法。

TABLE 2	
MSF cDNA	
1	gcggccgcga ctattcggtg cctgaaaaca acgatggcat ggaaaacact tccatttac (SEQ ID NO:2)
61	ctgttgttgc tgcgtctgtg ttctgtgatt cagcaagttt catctcaaga tctatcaagc
121	tgtgcaggga gatgtgggga aggtatttot agagatgcoa octgcaactg tgattataac
181	tgtaaacact acatggagtg ctgocctgat ttcaagagag tctgcactgc ggagctttcc
241	tgtaagggcc gctgctttga gtccttcgag agagggaggg agtgtgactg cgacgcccac
301	tgtaagaagt atgacaaagt ctgtcccgat tatgagagtt tctgtgcaga agtgcataat
361	cccacatcac caccatotto aaagaaagca cctccacctt caggagcadc tcaaacatc
421	aaatcaacaa ccaaacgttc acccaacca ccaacaaga agaagactaa gaaagttata
481	gaatcagagc aaataacaga agaacattot gtttctgaaa atcaagagtc ctctctctcc