



US 20100075322A1

(19) **United States**

(12) **Patent Application Publication**
Hooper

(10) **Pub. No.: US 2010/0075322 A1**

(43) **Pub. Date: Mar. 25, 2010**

(54) **METHODS AND COMPOSITIONS FOR IDENTIFYING MYCOTOXINS AND FUNGAL SPECIES**

Publication Classification

(76) Inventor: **Dennis G. Hooper**, Lewisville, TX (US)

(51) **Int. Cl.**
C12Q 1/68 (2006.01)
G01N 33/566 (2006.01)
G01N 33/53 (2006.01)
G01N 33/48 (2006.01)

Correspondence Address:
BARNES & THORNBURG LLP
11 SOUTH MERIDIAN
INDIANAPOLIS, IN 46204 (US)

(52) **U.S. CL.** **435/6; 436/501; 435/7.8; 435/40.5**

(21) Appl. No.: **12/545,734**

(57) **ABSTRACT**


(22) Filed: **Aug. 21, 2009**

Related U.S. Application Data

(60) Provisional application No. 61/091,185, filed on Aug. 22, 2008.

The invention relates to a method of testing for a mycotoxin in patient tissue or body fluid and testing for a fungal species in the patient tissue or body fluid. The method can also be used to determine if a patient is at risk for or has developed a disease state related to a fungal infection, and to develop an effective treatment regimen for the patient.

URINE TRICOTHECENES



Tricothecenes Value (ppb)	Number of Tests
0 - 0.2	5
0.21 - 1.0	17
1.1 - 2.0	11
2.1 - 5.0	27
5.1 - 10	19
10.1 - 35	25
> 35*	5
Total:	109

URINE TRICOTHECENES





Fig. 1

Nasal Secretion TRICOTHECENES



Tricothecenes Testing - Nasal Secretions	
Tricothecenes Value (ppb)	Number of Tests
0 - 0.2	25
0.21 - 1.0	10
1.1 - 2.0	0
2.1 - 5.0	3
5.1 - 10	0
10.1 - 35	3
> 35	1
Total:	42

Fig. 2




Fig. 3




Fig. 4

METHODS AND COMPOSITIONS FOR IDENTIFYING MYCOTOXINS AND FUNGAL SPECIES

CROSS REFERENCE TO RELATED APPLICATIONS

[0001] This application claims priority under 35 U.S.C. §119(e) to U.S. Provisional Application Ser. No. 61/091,185, filed on Aug. 22, 2008, incorporated herein by reference in its entirety.

FIELD OF THE INVENTION

[0002] This invention relates to methods and compositions for detecting or identifying fungi and mycotoxins. More particularly, the invention relates to methods and compositions for detecting or identifying fungi and mycotoxins in the tissues or body fluid samples of patients.

BACKGROUND AND SUMMARY

[0003] Molds (i.e., toxigenic and other septate molds) are ubiquitous in the environment. Mold is the common name for various types of fungi. Molds are usually found in moist, warm environments. Because molds grow in wet or moist indoor environments, people are exposed to molds or their byproducts through either direct contact, or through the air, if molds or mold byproducts are aerosolized. Exposure to molds can cause a number of adverse effects including allergic reactions, asthma attacks, and infections, particularly in individuals with immune system deficiencies.

[0004] Adverse effects from molds may occur when individuals are exposed to large doses of chemicals, known as mycotoxins, which are fungal metabolites (Samson et al., 1985; Burge, 1990; Flannigan et al., 1991). Mycotoxins have toxic effects ranging from severe irritations, such as allergic reactions and asthma, to immuno-suppression and cancer. Most mycotoxins are cytotoxic and exert their effects by interfering with vital cellular processes such as protein, RNA, and DNA synthesis. As a result, mycotoxins may be damaging to the skin, the lungs, the gut, and the like. The combined outcome may increase the susceptibility of the exposed individual to infectious diseases and, possibly, to cancer. Almost all of the studies to date focus on disease induced by mycotoxins ingested in contaminated food (Baxter et al., 1981), but mycotoxins are secondary metabolites of fungal spores and can enter the body through the respiratory tract.

[0005] In heavily contaminated environments, neurotoxic symptoms related to airborne mycotoxin exposure have been reported (Croft et al., 1986). Skin is another potential route of exposure to the mycotoxins of several fungi which have caused cases of severe dermatosis (Vennewald and Wollina, 2005). These same molds may cause invasive mold infection among patients with diseases which render the patient immuno-suppressed such as leukemia, lymphoma, and many cancers (Kontoyiannis, D P et al, 2005). The mold infections in such patients are often fatal with a documented fatality rate of 92% (Paterson and Singh, 1999).

[0006] A definitive and early diagnosis of a fungal infection is crucial for patient treatment and management. A diagnosis of a fungal infection is often rendered late in the disease process, often even as late as autopsy (Kontoyiannis et al, 2000; Vogeser et al., 1997). The reasons for the late diagnosis of fungal infections include the lack of good clinical specimens, the difficulty in differentiating invasive mold infections

from other types of infections, the lack of identification of molds with special stains in pathological specimens (i.e., these assays have a high error rate, a low sensitivity, and low specificity), and the lack of an ability to obtain an antibody-based diagnosis in immuno-compromised patients.

[0007] Thus, a reliable, sensitive, specific, and rapid method for mold detection in patient body fluids and tissues is needed. Applicant's present invention is based on the idea that if mycotoxins can be identified in patient tissue or body fluids, the identification of mycotoxins may serve as a potential diagnostic method 1.) to identify patients at risk for developing disease states related to mold infections, or 2.) to rapidly determine the cause of diseases related to mold infections so that effective treatment regimens can be developed for patients exposed to molds and experiencing symptoms resulting from mold infection. Applicant's present invention is also based on the development of a reliable, sensitive, specific, and rapid method for detecting fungal DNA in patient body fluids and tissues, and the use of this assay in combination with mycotoxin detection.

[0008] The present invention provides methods for detecting and identifying, in patient tissue and body fluid specimens, 1.) mycotoxins produced by fungi, and 2.) fungal DNA from fungal spores. The present invention provides rapid, reliable, sensitive, and specific diagnostic tests for the presence of fungi and fungal toxins in patient tissue and body fluids. Applicant has developed mycotoxin and fungal DNA extraction procedures and has supplemented those methods by developing detection methods. The detection methods employ antibody-based identification for mycotoxins in combination with the amplification of DNA with primers that specifically and selectively amplify fungal DNA isolated from patient tissues and body fluids and DNA probes for detection.

[0009] In one embodiment, a method is provided of identifying a fungal species in a patient tissue or body fluid. The method comprises the steps of identifying a mycotoxin in a patient tissue or body fluid, and specifically identifying a fungal species in the mycotoxin positive patient tissue or body fluid. In another embodiment, the method can further comprise the steps of extracting and recovering the mycotoxin from the patient tissue or body fluid, and contacting the mycotoxin with an antibody directed against the mycotoxin. In another embodiment, the method can further comprise the steps of extracting and recovering DNA of the fungal species from the patient tissue or body fluid, amplifying the DNA, and hybridizing a probe to the DNA to specifically identify the fungal species.

[0010] In one embodiment, a method is provided of identifying a fungal species in a patient tissue or body fluid. The method comprises the steps of testing for the presence of a mycotoxin and not identifying a mycotoxin in a patient tissue or body fluid. The method further comprises the steps of extracting and recovering DNA of a fungal species from the patient tissue or body fluid, amplifying the DNA, and hybridizing a probe to the DNA to specifically identify the fungal species.

[0011] In one embodiment, a method is provided of identifying a mycotoxin in a patient tissue or body fluid. The method comprises the steps of extracting and recovering the mycotoxin from the patient tissue or body fluid, and contacting the mycotoxin with an antibody directed against the mycotoxin. The method further comprises the steps of extracting and recovering patient DNA from the patient tissue or body

fluid, amplifying the DNA, hybridizing a probe to the patient DNA, and not identifying a specific fungal species in the patient tissue or body fluid.

[0012] In another embodiment, a method is provided of determining if a patient is at risk for or has developed a disease state related to a fungal infection. The method comprises the steps of identifying a mycotoxin in a patient tissue or body fluid, specifically identifying a fungal species in the mycotoxin positive patient tissue or body fluid, and determining if the patient is at risk for or has developed the disease state related to the fungal infection. In another embodiment, the method further comprises the steps of extracting and recovering the mycotoxin from the patient tissue or body fluid, and contacting the mycotoxin with an antibody directed against the mycotoxin. In yet another embodiment, the method further comprises the steps of extracting and recovering DNA of the fungal species from the patient tissue or body fluid, amplifying the DNA, and hybridizing a probe to the DNA to specifically identify the fungal species.

[0013] In another illustrative embodiment the amplifying step is performed with primers that hybridize to the DNA. In another illustrative embodiment the DNA is amplified using PCR. In another illustrative embodiment, the PCR is real-time PCR. In another illustrative embodiment, the probe is fluorescently labeled. In another illustrative embodiment, the amplified sequence is internal transcribed spacer regions of nuclear ribosomal DNA. In another illustrative embodiment, the probe is bound to a bead dyed with a fluorochrome. In yet another illustrative embodiment, the method further comprises the step of quantifying the mycotoxin. In a further illustrative embodiment, the tissue is derived from a patient tissue biopsy and is in a 10% formalin solution or is in a paraffin block. In another illustrative embodiment, the antibody is bound to a bead dyed with a fluorochrome.

[0014] In various illustrative embodiments, the body fluids are selected from the group consisting of urine, nasal secretions, nasal washes, bronchial lavages, bronchial washes, spinal fluid, sputum, gastric secretions, seminal fluid, other reproductive tract secretions, lymph fluid, whole blood, serum, and plasma.

[0015] In various illustrative embodiments, the mycotoxin is selected from the group consisting of tricothecenes, Aflatoxin B1, Aflatoxin B2, Aflatoxin D1, Aflatoxin D2, and Ochratoxin A.

[0016] In various illustrative embodiments, the fungal species are selected from the group consisting of *S. chartarum*, *A. versicolor*, *A. niger*, *P. chrysogenum*, *P. verrucosum*, *G. candidum*, *A. flavus*, *A. fumigatus*, *A. nidulans*, *A. ochraceus*, *A. parviticus*, *A. sydowii*, *A. ustus*, *F. solani*, *P. aurantiogriseum*, *P. citrinum*, *P. corylophilum*, *P. crustosum*, *P. expansum*, *P. fellutanum*, *P. roquefortii*, *P. simplicissimum*, *S. echinata*, and *E. amstelodami*.

[0017] In yet another embodiment, a kit is provided. The kit comprises components for the extraction and recovery of a mycotoxin from body fluid or tissue of a patient, components for identification of the mycotoxin, and a purified nucleic acid probe and primer. In another embodiment, the components for identification of the mycotoxin include beads dyed with a fluorochrome and coupled to antibodies to the mycotoxin or to the mycotoxin or to a mycotoxin antigen. In another embodiment, the components for identification of the mycotoxin comprise an antibody directed against the mycotoxin.

BRIEF DESCRIPTION OF THE DRAWINGS

[0018] FIG. 1 shows Tricothecenes values for urine tests.

[0019] FIG. 2 shows Tricothecenes values for nasal secretions.

[0020] FIG. 3 shows fungal DNA coupling efficiency to microspheres.

[0021] FIG. 4 shows mycotoxin coupling efficiency to microspheres.

DESCRIPTION OF THE ILLUSTRATIVE EMBODIMENTS

[0022] The present invention relates to methods and compositions for identifying or detecting the presence of molds (i.e., fungi) in patient tissue and body fluids. The identification and detection methods are based on 1.) amplification of fungal DNA using a PCR-based method in combination with 2.) testing for mycotoxins in patient body fluids and tissues. The methods and compositions (e.g., primers and probes) for amplification of fungal DNA are highly specific and sensitive and avoid co-amplification of or do not co-amplify non-specific human or animal nucleic acids.

[0023] The methods and compositions for testing for mycotoxins to detect and quantify mycotoxins are also very specific and sensitive. These methods and compositions utilize antibody-based identification of mycotoxins. In illustrative embodiments, Enzyme Linked Immunosorbant Assay (ELISA), or affinity chromatography can be used to detect mycotoxins produced by toxic molds. Illustratively, the mycotoxins can be aflatoxins, ochratoxins, or tricothecenes (e.g., Verrucarins A, B and J, Roridin A, E, H, and L-2, Satratoxins F, G, and H, Verrucarol, isosatratoxin F, G, and H, and T-2). Illustrative of antibody-based assays that can be used to identify mycotoxins are the Tricothecene kit (Enviroligix, Inc., Portland, Me.), the AflaTest® (VICAM, Inc.), the OchraTest™ (VICAM Inc.), and Luminex®-based assays.

[0024] Aflatoxins are toxin metabolites produced by a variety of molds such as *Aspergillus flavus* and *Aspergillus parasiticus*. Aflatoxins are carcinogenic and can be present in grains, nuts, cottonseed and other commodities associated with human food or animal feeds. Crops may be contaminated by one or more of the four sub-types of aflatoxins, B1, B2, G1 and G2. Aflatoxin B1 is the most toxic and frequently detected aflatoxin. Aflatoxins have been implicated in human health disorders including hepatocellular carcinoma, Reye's syndrome, and chronic hepatitis.

[0025] Ochratoxin A is the most important and most commonly occurring of a structurally related group of compounds called ochratoxins. Ochratoxin A is produced by some species of *Aspergillus*, such as *A. ochraceus*, and by *Penicillium verrucosum*. Ochratoxin A is a potent toxin affecting mainly the kidneys, in which it can cause both acute and chronic lesions. A nephrotoxic effect has been demonstrated in all mammalian species, and both teratogenic and reproductive effects have been demonstrated. Ochratoxin A is known to affect the immune system in a number of mammalian species.

[0026] Some tricothecenes are macrocyclic mycotoxins. There are over one hundred tricothecenes that cause irritation and immunosuppressive effects (Ueno, Y. 1983 and Tuomi, et al. 2000). Most tricothecenes were originally isolated from species of *Fusarium*, but they also may be produced by other fungi, such as species of *Stachybotrys*, *Trichothecium*, and others (Kurata and Ueno, 1983 and Ueno, Y. 1983). *Stachybotrys chartarum* is known to produce a number of potent

mycotoxins in this family including Verrucarins B and J, Roridin E, Satratoxins F, G, and H, and isosatratoxin F, G, and H (Hinkley, et al. 2001 and Jarvis, et al, 1998). These mycotoxins are also known to be potent inhibitors of protein synthesis in eukaryotes.

[0027] In various illustrative embodiments, body fluids that can be tested for the presence of fungal DNA in combination with mycotoxins, include, but are not limited to, urine, nasal secretions, nasal washes, inner ear fluids, bronchial lavages, bronchial washes, alveolar lavages, spinal fluid, bone marrow aspirates, sputum, pleural fluids, synovial fluids, pericardial fluids, peritoneal fluids, saliva, tears, gastric secretions, stool, reproductive tract secretions, such as seminal fluid, lymph fluid, and whole blood, serum, or plasma. These samples can be prepared for testing as described herein. In various embodiments, tissue samples can include tissue biopsies of hospital patients or out-patients and autopsy specimens. As used herein, the term "tissue" includes, but is not limited to, biopsies, autopsy specimens, cell extracts, tissue sections, aspirates, tissue swabs, and fine needle aspirates.

[0028] In accordance with the invention the word "patient" means a human or an animal, such as a domestic animal (e.g., a dog or a cat). Accordingly, the methods and compositions disclosed herein can be used for both human clinical medicine and veterinary applications. Thus, the patient afflicted with a disease state related to a fungal infection can be a human, or in the case of veterinary applications, can be a laboratory, agricultural, domestic or wild animal. The present invention can be applied to patients including, but not limited to, humans, laboratory animals such rodents (e.g., mice, rats, hamsters, etc.), rabbits, monkeys, chimpanzees, domestic animals such as dogs, cats, and rabbits, agricultural animals such as cows, horses, pigs, sheep, goats, chickens, and wild animals in captivity such as bears, pandas, lions, tigers, leopards, elephants, zebras, giraffes, gorillas, dolphins, and whales.

[0029] The methods and compositions described herein can be used to test for microbial DNA (e.g., fungal DNA) in combination with microbial toxins (e.g., mycotoxins) in microbes selected from the group consisting of *Absidia coerulea*, *Absidia glauca*, *Absidia corymbifera*, *Acremonium strictum*, *Alternaria alternata*, *Apophysomyces elegans*, *Saksena vasiformis*, *Aspergillus flavus*, *Aspergillus oryzae*, *Aspergillus fumigatus*, *Neosartorya fischeri*, *Aspergillus niger*, *Aspergillus foetidus*, *Aspergillus phoenicis*, *Aspergillus nomius*, *Aspergillus ochraceus*, *Aspergillus ostianus*, *Aspergillus auricomus*, *Aspergillus parasiticus*, *Aspergillus sojae*, *Aspergillus restrictus*, *Aspergillus caesillus*, *Aspergillus conicus*, *Aspergillus sydowii*, *Aspergillus tamarii*, *Aspergillus terreus*, *Aspergillus ustus*, *Aspergillus versicolor*, *Aspergillus ustus*, *Aspergillus versicolor*, *Chaetomium globosum*, *Cladosporium cladosporioides*, *Cladosporium herbarum*, *Cladosporium sphaerospermum*, *Conidiobolus coronatus*, *Conidiobolus incongruus*, *Cunninghamella elegans*, *Emericella nidulans*, *Emericella rugulosa*, *Emericella quadrilineata*, *Apicoccum nigrum*, *Eurotium amstelodami*, *Eurotium chevalieri*, *Eurotium herbariorum*, *Eurotium rubrum*, *Eurotium repens*, *Geotrichum candidum*, *Geotrichum klebahnii*, *Memnoniella echinata*, *Mortierella polycephala*, *Mortierella wolfii*, *Mucor mucedo*, *Mucor amphibiorum*, *Mucor circinelloides*, *Mucor heimalis*, *Mucor indicus*, *Mucor racemosus*, *Mucor ramosissimus*, *Rhizopus azygosporus*, *Rhizopus homothalicus*, *Rhizopus microsporus*, *Rhizopus oligosporus*, *Rhizopus oryzae*,

Myrothecium verrucaria, *Myrothecium roridum*, *Paecilomyces lilacinus*, *Paecilomyces variotii*, *Penicillium freii*, *Penicillium verrucosum*, *Penicillium hirsutum*, *Penicillium alberechii*, *Penicillium aurantiogriseum*, *Penicillium polonicum*, *Penicillium viridicatum*, *Penicillium hirsutum*, *Penicillium brevicompactum*, *Penicillium chrysogenum*, *Penicillium griseofulvum*, *Penicillium glandicola*, *Penicillium coprophilum*, *Eupenicillium crustaceum*, *Eupenicillium egyptiacum*, *Penicillium crustosum*, *Penicillium citrinum*, *Penicillium sartoryi*, *Penicillium westlingi*, *Penicillium corylophilum*, *Penicillium decumbens*, *Penicillium echinulatum*, *Penicillium solitum*, *Penicillium camembertii*, *Penicillium commune*, *Penicillium echinulatum*, *Penicillium sclerotigenum*, *Penicillium italicum*, *Penicillium expansum*, *Penicillium fellutanum*, *Penicillium charlesii*, *Penicillium janthinellum*, *Penicillium raperi*, *Penicillium madriti*, *Penicillium gladioli*, *Penicillium oxalicum*, *Penicillium roquefortii*, *Penicillium simplicissimum*, *Penicillium ochrochloron*, *Penicillium spinulosum*, *Penicillium glabrum*, *Penicillium thomii*, *Penicillium pupure-scens*, *Eupenicillium lapidosum*, *Rhizomucor miehei*, *Rhizomucor pusillus*, *Rhizomucor variabilis*, *Rhizopus stolonifer*, *Scopulariopsis asperula*, *Scopulariopsis brevicaulis*, *Scopulariopsis fusca*, *Scopulariopsis brumptii*, *Scopulariopsis chartarum*, *Scopulariopsis sphaerospora*, *Trichoderma asperellum*, *Trichoderma hamatum*, *Trichoderma viride*, *Trichoderma harzianum*, *Trichoderma longibrachiatum*, *Trichoderma citroviride*, *Trichoderma atroviride*, *Trichoderma koningii*, *Ulocladium atrum*, *Ulocladium chartarum*, *Ulocladium botrytis*, *Walleimia sebi*, *Stachybotrys chartarum*, and the like.

[0030] In embodiments where the microbe is a fungal species, the microbe is typically selected from the group consisting of *S. chartarum*, *S. prolificans*, *A. versicolor*, *A. vesicularis*, *A. niger*, *P. chrysogenum*, *P. verrucosum*, *G. candidum*, *A. flavus*, *A. fumigatus*, *A. nidulans*, *A. ochraceus*, *A. paraciticus*, *A. sydowii*, *A. ustus*, *F. solani*, *F. chlamydosporum*, *P. aurantiogriseum*, *P. citrinum*, *P. corylophilum*, *P. crustosum*, *P. expansum*, *P. fellutanum*, *P. roquefortii*, *P. simplicissimum*, *S. echinata*, and *E. amstelodami*. In one embodiment, the molds (i.e., fungi) can be black, toxigenic molds.

[0031] In one illustrative embodiment, a method is provided of testing for mycotoxin and a specific fungal species in patient tissue or body fluid. The method comprises the steps of extracting and recovering DNA from the patient tissue or body fluid, amplifying the DNA, hybridizing a probe to the DNA to specifically identify a fungal species, and specifically identifying the fungal species. The method also comprises steps to identify a mycotoxin as described below.

[0032] In one embodiment, a method is provided of identifying a fungal species in a patient tissue or body fluid. The method comprises the steps of testing for the presence of a mycotoxin and not identifying a mycotoxin in a patient tissue or body fluid. The method further comprises the steps of extracting and recovering DNA of a fungal species from the patient tissue or body fluid, amplifying the DNA, and hybridizing a probe to the DNA to specifically identify the fungal species.

[0033] In one embodiment, a method is provided of identifying a mycotoxin in a patient tissue or body fluid. The method comprises the steps of extracting and recovering the mycotoxin from the patient tissue or body fluid, and contacting the mycotoxin with an antibody directed against the mycotoxin. The method further comprises the steps of extracting and recovering patient DNA from the patient tissue or body

fluid, amplifying the DNA, hybridizing a probe to the patient DNA, and not identifying a specific fungal species in the patient tissue or body fluid.

[0034] In some embodiments, real-time PCR-based methods can be used to amplify the fungal DNA and to detect and identify fungal DNA by hybridization of the probe to the fungal DNA. PCR is described in U.S. Pat. Nos. 4,683,202 and 4,800,159, incorporated herein by reference, and methods for PCR are well-known in the art. Real-time PCR combines amplification and simultaneous probe hybridization to

achieve sensitive and specific detection of infectious molds (i.e., fungi) in real-time thereby providing instant detection of molds. In this embodiment, the time to detect or identify the fungus and to obtain a diagnosis is greatly reduced. Real-time PCR is conducted according to methods well-known in the art. Exemplary probes and primers and their target DNAs, that can be used in accordance with the invention are shown below. "Primer F" refers to a forward primer and "Primer R" refers to a reverse primer which are well-known terms in the art.

Target 1 <i>S. chartarum</i>	
Probe 1 char: 5'-ttgcttcggcgggaacgccccg	(SEQ ID NO: 1)
Primer F2: 5'-gcgaggatcattaccgag	(SEQ ID NO: 2)
Primer R2: 5'-atcgatgccagagccaagag	(SEQ ID NO: 3)
Target 2 <i>A. versicolor</i>	
Probe 2 vers: 5'-cggggagccctctcgggggc	(SEQ ID NO: 4)
Primer F1: 5'-cgtaggtgaacctgcggaag	(SEQ ID NO: 5)
Primer R1: 5'-atcgatgccggaaccaagag	(SEQ ID NO: 6)
Target 3 <i>A. niger</i>	
Probe 3 niger: 5'-tgtctattgtacctgttcttc	(SEQ ID NO: 7)
Primer F14: 5'-cgtaggtgaacctgcggaag	(SEQ ID NO: 8)
Primer R1: 5'-atcgatgccggaaccaagag	(SEQ ID NO: 9)
Target 4 <i>P. chrysogenum</i>	
Probe 4 chry: 5'-ctctgtctgaagattgtagtctgagt	(SEQ ID NO: 10)
Primer F1: 5'-cgtaggtgaacctgcggaag	(SEQ ID NO: 11)
Primer R1: 5'-atcgatgccggaaccaagag	(SEQ ID NO: 12)
Target 5 <i>P. verrucosum</i>	
Probe 5 verru: 5'-cccgcctttgtggccgcc	(SEQ ID NO: 13)
Primer F1: 5'-cgtaggtgaacctgcggaag	(SEQ ID NO: 14)
Primer R1: 5'-atcgatgccggaaccaagag	(SEQ ID NO: 15)
Target 6 <i>G. candidum</i>	
For Geo F1H: 5'-ggatctcttgggtctcgtatc	(SEQ ID NO: 16)
Rev Geo R1H: 5'-cttgatctgaggtgaaatagtg	(SEQ ID NO: 17)
Probe 6 geo: 5'-aacgcacattgcactttgggtatc	(SEQ ID NO: 18)
Target 7 <i>A. flavus</i>	
Probe 7 flav: 5'-cccgccattcatggccgcccgg	(SEQ ID NO: 19)
Primer F1: 5'-cgtaggtgaacctgcggaag	(SEQ ID NO: 20)
Primer R1: 5'-atcgatgccggaaccaagag	(SEQ ID NO: 21)
Target 8 <i>A. fumigatus</i>	
Probe 8 fumi: 5'-aaagtatgcagtctgagttgattatc	(SEQ ID NO: 22)
Primer F1: 5'-cgtaggtgaacctgcggaag	(SEQ ID NO: 23)
Primer R1: 5'-atcgatgccggaaccaagag	(SEQ ID NO: 24)
Target 9 <i>A. nidulans</i>	
Probe 9 nid: 5'-cccagggggcgagccgccgg	(SEQ ID NO: 25)
Primer F1: 5'-cgtaggtgaacctgcggaag	(SEQ ID NO: 26)
Primer R1: 5'-atcgatgccggaaccaagag	(SEQ ID NO: 27)

-continued

Target 10 *A. ochraceus*
Probe 10 ochr: 5'-acaccaacgtgaacactgtctgaag (SEQ ID NO: 28)
Primer F1: 5'-cgtaggtgaacctgcggaag (SEQ ID NO: 29)
Primer R1: 5'-atcgatgccggaaccaagag (SEQ ID NO: 30)

Target 11 *A. paraciticus*
Probe 11 para: 5'-cgggcccgcgctcatggccg (SEQ ID NO: 31)
Primer F1: 5'-cgtaggtgaacctgcggaag (SEQ ID NO: 32)
Primer R1: 5'-atcgatgccggaaccaagag (SEQ ID NO: 33)

Target 12 *A. sydowii*
Probe 12 syd: 5'-ccctcggggcgagccgccc (SEQ ID NO: 34)
Primer F1: 5'-cgtaggtgaacctgcggaag (SEQ ID NO: 35)
Primer R1: 5'-atcgatgccggaaccaagag (SEQ ID NO: 36)

Target 13 *A. ustus*
Probe 13 ust: 5'-ccacaccgaacctctgttatagc (SEQ ID NO: 37)
Primer F1: 5'-cgtaggtgaacctgcggaag (SEQ ID NO: 38)
Primer R1: 5'-atcgatgccggaaccaagag (SEQ ID NO: 39)

Target 14 *F. solani*
Probe14salani: 5'-cgggaatagacggccccgtgaaac (SEQ ID NO: 40)
Primer F2: 5'-gcgaggatcattaccgag (SEQ ID NO: 41)
Primer R2: 5'-atcgatgccagagccaagag (SEQ ID NO: 42)

Target 15 *P. aurantiogriseum*
Probe 15 auran: 5'-cccgcctttactggccgccc (SEQ ID NO: 43)
Primer F1: 5'-cgtaggtgaacctgcggaag (SEQ ID NO: 44)
Primer R1: 5'-atcgatgccggaaccaagag (SEQ ID NO: 45)

Target 16 *P. citrinum*
Probe 16 citr: 5'-tgttgcctcgccggccccgc (SEQ ID NO: 46)
Primer F4: 5'-ggaaggatcattaccgagt (SEQ ID NO: 47)
Primer R1: 5'-atcgatgccggaaccaagag (SEQ ID NO: 48)

Target 17 *P. corylophilum*
Probe 17 corylo: 5'-ttattgtaccttgttcttcggcgg (SEQ ID NO: 49)
Primer F1: 5'-cgtaggtgaacctgcggaag (SEQ ID NO: 50)
Primer R1: 5'-atcgatgccggaaccaagag (SEQ ID NO: 51)

Target 18 *P. crustosum*
Probe 18 crust: 5'-cgatctccgggggacgggcc (SEQ ID NO: 52)
Primer F7: 5'-ctgtccgagcgtcattgctg (SEQ ID NO: 53)
Primer R5: 5'-cgaggaccggacgcggtg (SEQ ID NO: 54)

Target 19 *P. expansum*
Probe19expan: 5'-agacacccccgaactctgctgaa (SEQ ID NO: 55)
Primer F1: 5'-cgtaggtgaacctgcggaag (SEQ ID NO: 56)
Primer R1: 5'-atcgatgccggaaccaagag (SEQ ID NO: 57)

Target 20 *P. fellutanum*
Probe 20 fell: 5'-cccgcctgccaggccgccc (SEQ ID NO: 58)
Primer F1: 5'-cgtaggtgaacctgcggaag (SEQ ID NO: 59)
Primer R1: 5'-atcgatgccggaaccaagag (SEQ ID NO: 60)

-continued

Target 21 <i>P. roquefortii</i>	
Probe 21 roque: 5'-caccctgtttatttaccttattgc	(SEQ ID NO: 61)
Primer F1: 5'-cgtaggtgaacctgcggaag	(SEQ ID NO: 62)
Primer R1: 5'-atcgatgccggaaccaagag	(SEQ ID NO: 63)
Target 22 <i>P. simplicissimum</i>	
Probe 22 simpl: 5'-caccctgtttatcgctacctgttg	(SEQ ID NO: 64)
Primer F1: 5'-cgtaggtgaacctgcggaag	(SEQ ID NO: 65)
Primer R1: 5'-atcgatgccggaaccaagag	(SEQ ID NO: 66)
Target 23 <i>S. echinata</i>	
Probe 23 echin: 5'-ttgcttcggcgaggagccccg	(SEQ ID NO: 67)
Primer F2: 5'-gcgaggatcattaccgag	(SEQ ID NO: 68)
Primer R2: 5'-atcgatgccagagccaagag	(SEQ ID NO: 69)
Target 24 <i>E. amstelodami</i>	
Probe 24 amst: 5'-tgtctatctgtacctgttcttcg	(SEQ ID NO: 70)
Primer F1: 5'-cgtaggtgaacctgcggaag	(SEQ ID NO: 71)
Primer R1: 5'-atcgatgccggaaccaagag	(SEQ ID NO: 72)
Fungal Universal Group 1	
UP1: 5'-cctcggatcaggtaggatag	(SEQ ID NO: 73)
UF1: 5'-atgcctgtccgagcgtcatt	(SEQ ID NO: 74)
UR1: 5'-ttcctcgccttattgatag	(SEQ ID NO: 75)
Fungal Universal Group 2	
Up2: 5'-acggatctcttggctctggcatc	(SEQ ID NO: 76)
F2: 5'-gcgaggatcattaccgag	(SEQ ID NO: 77)
UR2: 5'-ttcactgaattctgcaattcac	(SEQ ID NO: 78)

[0035] Alternative illustrative embodiments for the Target 3 probe and primer F1 are 5'-cctctgccccgggcccgtg (SEQ ID NO: 79) and 5'-ggaaggatcattaccgagtg (SEQ ID NO: 80), respectively. An alternative illustrative embodiment for the Target 9 probe is 5'-ggagccccccaggggcgag (SEQ ID NO: 81). An alternative illustrative embodiment for the Target 12 probe is 5'-cggggaacccccctgggggc (SEQ ID NO: 82). An alternative illustrative embodiment for the Target 13 probe is 5'-tgcctccccccggggcgag (SEQ ID NO: 83). Alternative illustrative embodiments for the Target 18 probe, primer F7, and primer R5 are 5'-ggccccctccccgatctccg (SEQ ID NO: 84), 5'-agtgaatcatcgagctttgaac (SEQ ID NO: 85), and 5'-acctgatc-cgaggtcaacctg (SEQ ID NO: 86), respectively. An alternative illustrative embodiment for the Target 20 probe is 5'-cgggc-cgcctgccaggccg (SEQ ID NO: 87). An alternative illustrative embodiment for the Target 21 probe is 5'-ccggggggtttacac-ccccg (SEQ ID NO: 88). An alternative illustrative embodiment for the Target 22 probe is 5'-ccgggggcatctgcccccg (SEQ ID NO: 89).

[0036] In various embodiments, sample preparation (i.e., preparation of the target DNA) involves rupturing the cells (e.g., cells of the tissue or fungal spores in patient body fluid or tissue) and isolating the fungal DNA from the lysate. Techniques for rupturing cells and for isolation of DNA are well-known in the art. For example, cells may be ruptured by using a detergent or a solvent, such as phenol-chloroform. DNA may be separated from the lysate by physical methods including, but not limited to, centrifugation, pressure tech-

niques, or by using a substance with affinity for DNA, such as, for example, silica beads. After sufficient washing, the isolated DNA may be suspended in either water or a buffer. In other embodiments, commercial kits are available, such as Quiagen™, Nuclisens™, and Wizard™ (Promega), and Promega™. Methods for isolating DNA are described in Sambrook et al., "Molecular Cloning: A Laboratory Manual", 3rd Edition, Cold Spring Harbor Laboratory Press, (2001), incorporated herein by reference.

[0037] In embodiments where the fungal DNA is not detected, DNA in the lysate from the patient can be treated similarly.

[0038] In various embodiments described herein, the primers and probes used for amplification of the target DNA and for detection and identification of fungal DNA are oligonucleotides from about ten to about one hundred, more typically from about ten to about thirty or about six to about twenty-five base pairs long, but any suitable sequence length can be used. In illustrative embodiments, the primers and probes may be double-stranded or single-stranded, but the primers and probes are typically single-stranded. The primers and probes described herein are capable of specific hybridization, under appropriate hybridization conditions (e.g., appropriate buffer, ionic strength, temperature, formamide, and MgCl₂ concentrations), to a region of the target DNA. The primers and probes described herein are designed based on having a melting temperature within a certain range, and substantial complementarity to the target DNA. Methods for

the design of primers and probes are described in Sambrook et al., "Molecular Cloning: A Laboratory Manual", 3rd Edition, Cold Spring Harbor Laboratory Press, (2001), incorporated herein by reference.

[0039] The primers and probes described herein for use in PCR can be modified by substitution, deletion, truncation, and/or can be fused with other nucleic acid molecules wherein the resulting primers and probes hybridize specifically to the intended targets and are useful in the methods described herein for amplification of the target DNAs. Derivatives can also be made such as phosphorothioate, phosphotriester, phosphoramidate, and methylphosphonate derivatives, that specifically bind to single-stranded DNA or RNA (Goodchild, et al., *Proc. Natl. Acad. Sci.* 83:4143-4146 (1986)).

[0040] The invention encompasses isolated or substantially purified nucleic acids. An "isolated" or "purified" nucleic acid molecule is substantially free of other cellular material, or culture medium when produced by recombinant techniques, or substantially free of chemical precursors or other chemicals when chemically synthesized. Preferably, an "isolated" or "purified" nucleic acid is free of sequences that naturally flank the nucleic acid in the genomic DNA of the organism from which the nucleic acid is derived. For example, in various embodiments, the isolated or purified nucleic acid molecule can contain less than about 5 kb, 4 kb, 3 kb, 2 kb, 1 kb, 0.5 kb, or 0.1 kb of nucleotide sequences that naturally flank the nucleic acid molecule in genomic DNA of the cell from which the nucleic acid is derived.

[0041] Also within the scope of the invention are nucleic acids complementary to the probes and primers described herein, and those that hybridize to the nucleic acids described herein or those that hybridize to their complements under highly stringent conditions. In accordance with the invention "highly stringent conditions" means hybridization at 65° C. in 5×SSPE and 50% formamide, and washing at 65° C. in 0.5×SSPE. Conditions for low stringency and moderately stringent hybridization are described in Sambrook et al., "Molecular Cloning: A Laboratory Manual", 3rd Edition, Cold Spring Harbor Laboratory Press, (2001), incorporated herein by reference. In some illustrative aspects, hybridization occurs along the full-length of the nucleic acid.

[0042] Also included are nucleic acid molecules having about 60%, about 70%, about 75%, about 80%, about 85%, about 90%, about 95%, 96%, 97%, and 98% homology to the probes and primers described herein. Determination of percent identity or similarity between sequences can be done, for example, by using the GAP program (Genetics Computer Group, software; now available via Accelrys on <http://www.accelrys.com>), and alignments can be done using, for example, the ClustalW algorithm (VNTI software, InforMax Inc.). A sequence database can be searched using the nucleic acid sequence of interest. Algorithms for database searching are typically based on the BLAST software (Altschul et al., 1990). In some embodiments, the percent identity can be determined along the full-length of the nucleic acid.

[0043] As used herein, the term "complementary" refers to the ability of purine and pyrimidine nucleotide sequences to associate through hydrogen bonding to form double-stranded nucleic acid molecules. Guanine and cytosine, adenine and thymine, and adenine and uracil are complementary and can associate through hydrogen bonding resulting in the formation of double-stranded nucleic acid molecules when two nucleic acid molecules have "complementary" sequences.

The complementary sequences can be DNA or RNA sequences. The complementary DNA or RNA sequences are referred to as a "complement."

[0044] Techniques for synthesizing the probes and primers described herein are well-known in the art and include chemical syntheses and recombinant methods. Such techniques are described in Sambrook et al., "Molecular Cloning: A Laboratory Manual", 3rd Edition, Cold Spring Harbor Laboratory Press, (2001), incorporated herein by reference. Primers and probes can also be made commercially (e.g., CytoMol, Sunnyvale, Calif. or Integrated DNA Technologies, Skokie, Ill.). Techniques for purifying or isolating the probes and primers described herein are well-known in the art. Such techniques are described in Sambrook et al., "Molecular Cloning: A Laboratory Manual", 3rd Edition, Cold Spring Harbor Laboratory Press, (2001), incorporated herein by reference. The primers and probes described herein can be analyzed by techniques known in the art, such as restriction enzyme analysis or sequencing, to determine if the sequence of the primers and probes is correct.

[0045] In various embodiments of the methods and compositions described herein, the probes and primers can be labeled, such as with fluorescent compounds, radioactive isotopes, antigens, biotin-avidin, colorimetric compounds, or other labeling agents known to those of skill in the art, to allow detection and quantification of amplified DNA, such as by Real-Time PCR. In illustrative embodiments, the labels may include 6-carboxyfluorescein (FAMTM), TETTM (tetrachloro-6-carboxyfluorescein), JOETM (2,7-dimethoxy-4,5-dichloro-6-carboxyfluorescein), VICTM, HEX (hexachloro-6-carboxyfluorescein), TAMRATM (6-carboxy-N,N,N',N'-tetramethylrhodamine), BHQTM, SYBR[®] Green, Alexa 350, Alexa 430, AMCA, BODIPY 630/650, BODIPY 650/665, BODIPY-FL, BODIPY-R6G, BODIPY-TMR, BODIPY-TRX, Cascade Blue, Cy3, Cy5, 6-FAM, Fluorescein, Oregon Green 488, Oregon Green 500, Oregon Green 514, Pacific Blue, REG, Rhodamine Green, Rhodamine Red, ROX, and/or Texas Red.

[0046] Specificity of the probes and primers described herein was demonstrated by testing hybridization of the probe and primers sets against 23 different mold organisms (10 species of *Aspergillus*, 10 species of *Penicillium*, 2 species of *Stachybotrys*, and 1 species of *Fusarium*). There were no cross-over reactions or cross-over detection noted for any of the tested probe and primer sequences. Thus, the methods and compositions (e.g., primers and probes) for amplification of fungal DNA are highly specific and avoid co-amplification of or do not co-amplify non-specific nucleic acids.

[0047] Universal probes can be used to provide a method for determining the presence of fungal DNA before conducting target-specific assays. In one embodiment, universal probes and primers can be used to detect the presence of *Aspergillus* and *Penicillium* species (see probes and primers for Fungal Universal Group 1 below). In another embodiment, universal probes and primers can be used to detect the presence of *Stachybotrys* and *Fusarium* species (see probes and primers for Fungal Universal Group 2 below). In these embodiments, the probes and primers can be homologous for all targets of interest related to *Aspergillus*, *Penicillium*, *Stachybotrys*, and *Fusarium* species.

Fungal Universal Group 1
UP1: 5'-cctcggatcaggtaggatc (SEQ ID NO: 73)

-continued

UF1: 5'-atgcctgtccgagcgtcatt (SEQ ID NO: 74)

UR1: 5'-ttctcccgcttattgatatg (SEQ ID NO: 75)

Fungal Universal Group 2

Up2: 5'-acggatctcttggctctggcatc (SEQ ID NO: 76)

F2: 5'-gcgagggatcattaccgag (SEQ ID NO: 77)

UR2: 5'-ttcactgaattctgcaattcac (SEQ ID NO: 78)

[0048] In combination with the use of primers and probes to identify fungal species, a method is provided of identifying a mycotoxin in patient tissue or body fluid. The method comprises the steps of testing for the mycotoxin by extracting and recovering the mycotoxin from the patient tissue or body fluid, contacting the mycotoxin with an antibody directed against the mycotoxin, and identifying the mycotoxin. In some embodiments where a mycotoxin is not identified, patient samples are treated similarly although a mycotoxin is not detected.

[0049] Illustratively, patient (e.g., human or animal) tissue is received in 1.) a 10% formalin fluid or 2.) in a paraffin block in which the tissue has been fixed in formalin. In one embodiment for mycotoxin detection and quantitation, the tissue can then be processed by various dehydration steps and finally embedded in paraffin. In this embodiment, the tissue can then be cut in 3-5 micron samples. In an illustrative embodiment, approximately 25-35 mg of tissue can then be processed as described in Examples 2-6 for mycotoxin extraction. Illustratively, body fluids can be prepared as described in Examples 1 and 3-6 or by other methods known in the art. In another illustrative embodiment, patient body fluids can be tested for the presence of mycotoxins. Illustratively, any antigen associated with a fungus or with a mycotoxin can be detected, or a mycotoxin may not be detected.

[0050] The methods and compositions for detection and quantification of mycotoxins are very specific and sensitive (e.g., sensitivity of 1.0 ng/ml for aflatoxins, 0.2 ng/ml for Tricothecenes, and 2 ng/ml for ochratoxins). Specificity of mycotoxins was tested in each group (Tricothecenes, Aflatoxins, Ochratoxins) by testing known samples of mycotoxins (obtained from Trilogy Laboratories, Washington, Mo., and from Sigma, St. Louis, Mo.) in each mycotoxin test. There were no cross-over reactions or cross-over detection of mycotoxins between the groups. Thus, the methods and compositions for detection and quantification of mycotoxins are also very specific and sensitive. The methods and compositions described in Examples 2-6 utilize antibody-based identification of mycotoxins. In illustrative embodiments, Enzyme-Linked Immunosorbant Assay (ELISA), affinity chromatography, or a Luminex®-based assay can be used to detect mycotoxins produced by toxic molds. Illustratively, the mycotoxins can be aflatoxins, ochratoxins, or Tricothecenes (e.g., Verrucarins A, B and J, Roridin A, E, H, and L-2, Satratoxins F, G, and H, Verrucarol, isosatratoxin F, G, and H, and T-2).

[0051] Illustrative of antibody-based assays that can be used to identify and subsequently quantitate mycotoxins, or fungal or mycotoxin antigens, are the Tricothecene kit (Envirologix, Inc., Portland, Me.), the AflaTest® (VICAM, Inc.), and the OchraTest™ (VICAM Inc., Watertown, Mass.).

[0052] Another exemplary detection method for multiple mycotoxins in patient samples that have been exposed to

fungal targets belonging to, for example, *Aspergillus*, *Penicillium*, *Stachybotrys*, and *Fusarium* species is the Luminex® format (Luminex, Austin, Tex., see Examples 9 and 11-13). In one aspect of the invention, the Luminex® assay utilizes microspheres (beads) that are dyed with fluorochromes and that are coupled to antigens to detect antibodies, in patient body fluids or tissues, to mycotoxins, mycotoxin antigens, or other fungal antigens. In another embodiment, the microspheres are coupled to antibodies to detect, in patient body fluids or tissues, mycotoxins, mycotoxin antigens, or other fungal antigens. In this illustrative embodiment, the antibodies coupled to the microspheres can be polyclonal or monoclonal antibodies, but monoclonal antibodies are typically used. In another illustrative embodiment, the beads can be coupled to DNA probes to detect DNA specific to fungal species. Exemplary methods involving the Luminex® assay are described in more detail in Examples 9 and 11-13.

[0053] Mycotoxins are tested for or are identified and quantitated, for example, by using control samples of the body fluid or tissue to be analyzed where the control samples are obtained from patients with no documented history of exposure to molds or mycotoxins. For example, negative control samples can be obtained from autopsy specimens in which the patient had no exposure to mycotoxins or molds (e.g., victims of motor vehicle accidents, coronary artery disease, or myocardial infarction). For positive controls, for example, samples of negative tissue and/or body fluids can be spiked with known positive amounts of mycotoxins or spores prior to evaluation to generate a calibration curve (see Examples 4-6). Illustrative calibrators for the mycotoxins can be obtained from producers of the Tricothecene kit (Envirologix, Inc., Portland, Me.) and producers of the AflaTest® and OchraTest™ kits (VICAM Inc., Watertown, Mass.), or from Trilogy Laboratories (Washington, Mo.).

[0054] In another embodiment, a method is provided of determining if a patient is at risk for or has developed a disease state related to a fungal infection. The method comprises the steps of extracting and recovering a mycotoxin (i.e., a mycotoxin or a mycotoxin antigen) from a tissue or body fluid of the patient, contacting the mycotoxin (i.e., a mycotoxin or a mycotoxin antigen) with an antibody directed against the toxin, identifying the mycotoxin (i.e., a mycotoxin or a mycotoxin antigen), identifying the fungal species using the DNA-based methods described herein, and determining if the patient is at risk for or has developed the disease state related to the fungal infection. The method can further comprise the steps of extracting and recovering DNA of a specific fungal species from a tissue or body fluid of the patient, amplifying the DNA, hybridizing a probe to the DNA to specifically identify the fungal species, and specifically identifying the fungal species.

[0055] In one embodiment, a method is provided of identifying a fungal species in a patient tissue or body fluid. The method comprises the steps of testing for the presence of a mycotoxin and not identifying a mycotoxin in a patient tissue or body fluid. The method further comprises the steps of extracting and recovering DNA of a fungal species from the patient tissue or body fluid, amplifying the DNA, and hybridizing a probe to the DNA to specifically identify the fungal species.

[0056] In one embodiment, a method is provided of identifying a mycotoxin in a patient tissue or body fluid. The method comprises the steps of extracting and recovering the mycotoxin from the patient tissue or body fluid, and contact-

ing the mycotoxin with an antibody directed against the mycotoxin. The method further comprises the steps of extracting and recovering patient DNA from the patient tissue or body fluid, amplifying the DNA, hybridizing a probe to the patient DNA, and not identifying a specific fungal species in the patient tissue or body fluid.

[0057] In any embodiment involving “determining if the patient has developed the disease state related to the fungal infection,” this phrase means “diagnosing the patient with a fungal infection.”

[0058] The method embodiments described in the preceding paragraph provide methods of diagnosing fungal infections. Patients in need of diagnosis of a fungal infection can include cancer patients, post-operative patients, transplant patients, patients undergoing chemotherapy, immunosuppressed patients, and the like. These patients may experience symptoms of fungal infections including sinusitis, allergic reactions, headaches, and skin rashes. Patients in need of diagnosis may include humans or animals.

[0059] In one embodiment, for diagnosing fungal infections, kits are provided. The kits are useful for identifying, detecting, or quantitating fungal DNA and mycotoxins in a patient tissue or body fluid. The kit can contain the probes and/or primers described herein, components to extract and isolate fungal DNA, and components for DNA amplification, such as a heat stable DNA polymerase (e.g., Taq polymerase or Vent polymerase), buffers, MgCl₂, H₂O, and the like. The kit can also contain components to extract and isolate the mycotoxin (i.e., a mycotoxin or a mycotoxin antigen), antibody affinity matrices, ELISA plates, Luminex® beads, polyclonal or monoclonal antibodies, color development reagents, buffers, and the like. In one embodiment, the reagents can remain in liquid form. In another embodiment, the reagents can be lyophilized. In another illustrative embodiment, the kit can be used to detect other fungal antigens. The kits can also contain instructions for use.

[0060] In another embodiment, a kit comprising a purified nucleic acid with a sequence selected from the group consisting of SEQ ID NO: 1 to SEQ ID NO: 89 or a complement of a sequence selected from the group consisting of SEQ ID NO: 1 to SEQ ID NO: 89 is provided. The kit can comprise components for the extraction and recovery of fungal DNA and a mycotoxin from the body fluid or tissue of a patient. The kit can further comprise components for identification of the fungal DNA and the mycotoxin. The components for identification of the fungal DNA and the mycotoxin can include beads dyed with a fluorochrome and coupled to a probe for the fungal DNA or to antibodies to the mycotoxin or to the mycotoxin itself or to a mycotoxin antigen.

[0061] A calibration reagent (or multiple calibration reagents) can also be included in the kit and “calibration reagent” means any standard or reference material containing a known amount of the mycotoxin (i.e., a mycotoxin or a mycotoxin antigen). The sample suspected of containing the mycotoxin and the calibration reagent (or multiple calibration reagents) are assayed under similar conditions. The mycotoxin concentration is then calculated by comparing the results obtained for the unknown sample with the results obtained for the calibration reagent(s).

[0062] In accordance with one embodiment of the invention, a method of identifying a fungal species in a patient tissue or body fluid is described. A patient tissue or body fluid (e.g., blood, serum, spinal fluid, urine, sputum, or nasal washes) is first examined for the presence of mycotoxins. The

method comprises the steps of extracting and recovering the mycotoxin from the patient tissue or body fluid, contacting the mycotoxin with an antibody directed against the mycotoxin, and identifying the mycotoxin. The specimen is then tested for the presence of specific molds/fungi, the method further comprising the steps of extracting and recovering DNA of the fungal species from the patient tissue or body fluid, amplifying the DNA, hybridizing a probe to the DNA to specifically identify the fungal species, and specifically identifying the fungal species.

[0063] In one embodiment, a method is provided of identifying a fungal species in a patient tissue or body fluid. The method comprises the steps of testing for the presence of a mycotoxin and not identifying a mycotoxin in a patient tissue or body fluid. The method further comprises the steps of extracting and recovering DNA of a fungal species from the patient tissue or body fluid, amplifying the DNA, and hybridizing a probe to the DNA to specifically identify the fungal species.

[0064] In one embodiment, a method is provided of identifying a mycotoxin in a patient tissue or body fluid. The method comprises the steps of extracting and recovering the mycotoxin from the patient tissue or body fluid, and contacting the mycotoxin with an antibody directed against the mycotoxin. The method further comprises the steps of extracting and recovering patient DNA from the patient tissue or body fluid, amplifying the DNA, hybridizing a probe to the patient DNA, and not identifying a specific fungal species in the patient tissue or body fluid.

[0065] In various illustrative embodiments, the identification of a mycotoxin in a patient tissue or body fluid is followed by steps to specifically identify a corresponding mold or fungus in the patient tissue or body fluid. For example, if Tricothecenes are identified in the patient tissue or body fluid, steps for specifically identifying, e.g., *Stachybotrys chartarum*, *Stachybotrys echinata*, and/or *Fusarium solani* will follow. In another illustrative embodiment, if Aflatoxins are identified in the patient tissue or body fluid, steps for specifically identifying, e.g., *Aspergillus flavus*, *Aspergillus ochraceus*, *Aspergillus fumigatus*, *Aspergillus niger*, *Aspergillus ustus*, and/or *Aspergillus sydowii* will follow. In another illustrative embodiment, if Ochratoxin A is identified in the patient tissue or body fluid, steps for specifically identifying, e.g., *Penicillium chrysogenum*, *Penicillium fellutanum*, and/or *Penicillium verrucosum* will follow.

[0066] The following examples provide illustrative methods for carrying out the practice of the present invention. As such, these examples are provided for illustrative purposes only and are not intended to be limiting.

Example 1

Samples and Sample Preparation

[0067] Human urine was received in 5-10 ml quantities as first in the morning voided urines. Serums were received with the blood clot removed prior to receipt and a minimum of 1 ml of serum was frozen or used. Nasal secretions were obtained from hospital patients or out-patients. Fixed autopsy and surgical biopsy specimens were obtained from patients who had a history of exposure to mycotoxins or fungi. These samples were obtained from hospital pathology departments or coroners' offices. Tissue samples and body fluid samples were also obtained from patients who had no exposure to mycotoxins or

fungi and were sampled as a negative control group. Tissue specimens were cut using procedures described in Examples 2 and 7.

[0068] All specimens were placed into two groups. Group 1 comprised samples from individuals with no reported symptoms or known fungi or mycotoxin exposure. These samples served as negative controls and n values differed in each group of specimens sampled. Group 2 comprised samples from individuals with reported exposure to non-identified fungi or chemicals. Each test conducted had a different n value. Common symptoms of patients corresponding to group 2 samples included blurred vision, memory loss, fatigue, headache, nausea, loss of balance, cognitive deficits, rhinitis, sinusitis, rashes, and allergies. A detailed history and symptoms were provided to correspond to each patient sample.

[0069] Human urine was tested for the mycotoxins Aflatoxin B1, B2, G1, and G2, Ochratoxin A, and Tricothecenes. The studies described herein demonstrated that higher levels of mycotoxins were detected in early a.m. samples. Nasal secretions and washings were obtained by injection of 3-5 ml of sterile saline in each nostril of a patient. The patient was instructed to hold the saline in the nostrils for 30 seconds and then blow the saline into a sterile container held close to the nose. The specimen(s) were then collected and placed in containers.

[0070] Negative control samples of mycotoxins were made by dilution techniques for Aflatoxin (Sigma), Ochratoxin (Sigma) and Roridin A (for Tricothecenes). Samples of extracted and filtered human heart tissue, liver tissue, urine, and nasal secretions (including sputum) were spiked with various levels of the above named toxins. Each time a sample was evaluated, calibrators and negative and positive spiked tissues and fluids were also evaluated. Statistical analysis on all types of samples for mycotoxins were performed for sensitivity and specificity.

Example 2

Preparation of Tissues for Mycotoxin Extraction

[0071] Preparation of tissues for mycotoxin extraction from formalin fixed tissue and paraffin-embedded tissue from humans or animals was accomplished using the following procedure.

Specimens

[0072] Tissue was received as either tissue fixed in a 10% formalin solution or in a paraffin-embedded tissue block. Tissue can be stored indefinitely in either form. However, because of cross-linking of formalin and proteins which may give false negative readings for DNA, the tissue was not stored in formalin for greater than 6 months. A minimum of 25-35 mg of formalin-fixed tissue was required for mycotoxin extraction. A maximum of 3 grams of formalin-fixed tissue can be used.

Materials

[0073] Phosphate Buffered Saline (PBS; 0.9%), acid-washed silica beads (Cat #G1277; obtained from Sigma-

Aldrich), collection tubes (2 ml) screw cap, methanol (reagent grade, Sigma), and microcentrifuge tubes (2 ml) were used.

Procedure

[0074] For silica beads, 0.3 g±0.01 g of silica bead beating glass was added to a 2 ml screw cap tube making sure that there were no glass beads in the cap or around the rim. The tubes containing the beads were sterilized in an autoclave on the dry cycle for 10 minutes. If a large amount of tissue was evaluated, the tissue was placed in a blender and blended in PBS until well emulsified in the PBS. The sample was then filtered using simple gravity filtration through Whatman #9 filter paper.

[0075] The samples were recorded and assigned numbers in a sample log. 25-35 mg of paraffin-embedded tissue was then weighed and placed in a 2.0 ml screw cap tube. Methanol was added (1.0 ml reagent grade methanol) to the tube with the 0.3 g of silica beads and the sample was vortexed for 1 minute. The samples were bead beaten on the bead beater for 1 minute at the speed of 45. Then 500 µl of sample was removed and placed in 4.5 ml of PBS taking care not to remove the paraffin from the sample tube. The sample could then be used for extraction or could be frozen at -20 degrees centigrade to be used later in extraction and detection of aflatoxins, ochratoxins, and tricothecenes (see Examples 3-6).

Example 3

Preparation of Body Fluids for Mycotoxin Detection

[0076] Urine was received from a morning fresh first-voided specimen and stored at 1-6 degrees centigrade in a glass container. A urine analysis was conducted using a dipstick to measure pH, specific gravity, glucose, nitrates, ketones, and blood. The urine was examined for sediment and was centrifuged at 2500 rpm for 5 minutes if sediment was present. The supernatant was saved in a glass container for mycotoxin testing (storing in plastic was avoided to avoid a decrease in the detection level of tricothecenes). The urine was then used as indicated in Example 4 for aflatoxin determination, Example 5 for ochratoxin determination, and Example 6 for tricothecene determination.

[0077] Nasal secretions and mucous samples as well as washes were observed for mucous presence. If mucous was present, a solution of MUCOSOL™ (Alpha Tec Systems, Inc. Vancouver, Wash.) was prepared and added in equal amounts of body fluid to MUCOSOL™ in the secretions containing mucous. The specimen was then allowed to incubate 30 minutes at room temperature. The specimen was then centrifuged and the supernatant was removed. The sediment was then re-suspended in 10 ml of PBS. The specimen was then treated like any other body fluid and subjected to tests as described in Examples 4, 5, and 6. If testing for the presence of fungal DNA was desired, the specimen was then subjected to the tests described in Example 7.

[0078] Blood samples were obtained from the negative control group and exposed patients. Specimens were allowed to clot (no anticoagulant added) and then centrifuged for 10 minutes at 2000 rpm. Specimens were stored at 1-6 degrees centigrade for 48 hours or were frozen at -20 degrees centigrade for an indefinite period of time. Blood samples were extracted in a manner similar to that described by Garbis et al., *Anal. Chem.* 73:53589-64 (2001) and Hedman et al. *Arch.*

Tieremahr. 50:13-24 (1997). Serum samples were aliquoted in 200 μ l amounts into sterile 1.5 ml polystyrene microcentrifuge tubes. Immediately, 600 μ l of high performance HPLC grade acetonitrile (Fisher Scientific, Hampton, N.H.) was added. After 15 minutes, the samples were vortexed and centrifuged. The supernatants were transferred into clean 1.5 ml glass vials. Each sample was evaporated under a gentle stream of dry nitrogen and re-suspended in 100 μ l of pre-warmed sterile water. This was the final working solution for ELISA assays. Spinal fluid samples were analyzed as obtained from human patients. Samples were not processed before analysis.

Example 4

Detection of Aflatoxins in Body Fluids and Tissues

[0079] Aflatoxins are mycotoxins produced by the fungus *Aspergillus*. Samples from body fluids and tissues were prepared as described in Examples 2 and 3. Samples were then applied to an AflaTest® column (VICAM, L.P., Watertown, Mass.) which contains specific monoclonal antibodies directed against Aflatoxins B1, B2, G1, and G2. The aflatoxins bind to the antibodies on the column. The column was then washed to remove the immunoaffinity column of impurities. By passing an eluting solution through the column, the aflatoxin was removed from the antibody. The eluting solution was then read in a fluorometer. Aflatoxins B1, B2, G1, and G2 were detected using AflaTest® (VICAM, L.P., Watertown, Mass.) which is a well-known quantitative method for the detection of aflatoxins in grains and foodstuffs. This assay permits the measurement of all of the major aflatoxins (including B1, B2, G1, G2 and M1) without the use of toxic solvents (i.e., chloroform or methylene chloride).

[0080] In more detail, the procedure for body fluids and tissues was as follows. Tissue samples were prepared from fixed and paraffin-embedded tissues as described in Example 2 and body fluid samples were prepared as described in Example 3. One ml of the extracted tissue sample or body fluid was mixed with 10 ml of purified molecular grade water. The 10 ml of sample was then placed over an Aflatoxin® column (VICAM) and was filtered through the affinity column at a rate of 1-2 drops/second until air came through the column. Ten ml of an 80:20 solution of methanol:water was then passed through the column at a rate of 2 drops/second. Another 10 ml of purified water was passed through the affinity column at 2 drops/second. As an alternative procedure, ten ml of molecular grade water can be passed through the column at a rate of 2 drops/second. The affinity column was then eluted by passing 1.0 ml of HPLC grade methanol through the column at a rate of 1-2 drops/second and collecting all of the sample eluate (1 ml) in a glass cuvette. One ml of AflaTest® Developer (described below) containing bromine was added to the eluate in the cuvette. The specimen was then mixed well and the cuvette was placed in a calibrated fluorometer (Sequoia-Turner Model 450). The aflatoxin concentration was read after 60 seconds.

Preparation of AFLATEST® Developer Solution: Five ml of AflaTest® Developer concentrate solution (VICAM) was measured and placed in a 2 ounce amber glass bottle of a 50 ml bottle dispenser for the developer (VICAM Cat. no. 20600). Forty-five ml of purified water was added and mixed. The developer solution should be used within 8 hours after preparation. To ensure that the developer and water show no fluorescence, 1.0 ml of each was placed in the fluorometer and

read after 60 seconds. Results of the readings should be 0 ppb. Also, methanol blanks and purified water blanks should be read as 0 ppb. The results of the test were recorded in parts per billion (nanograms/milliliter).

Fluorometer Calibration: The Sequoia-Turner Fluorometer, Model 450, was calibrated as defined by the manufacturer's guidelines (VICAM). The machine was calibrated using standards supplied by VICAM (Green Standard=-2, and Red Standard=22). Known standards of Aflatoxin B1, B2, G1, and G2 were used (Trilogy Laboratories, Washington, Mo.). Standard Calibrators were made at 52 ppb, 26 ppb, and 2.6 ppb. **Pump and Set Up:** AflaTest® affinity chromatography was performed with the AflaTest® affinity column attached to a pump stand. The stand has a glass syringe barrel that serves as a reservoir for the column. A large plastic syringe with tubing and coupling provides air pressure to manually push liquids through the column. An adjustable air pump (VICAM Cat. no. 20650) can be attached to the pump tube instead of the large pump syringe barrel to operate without using hand pressure. Four and six position pump stands with aquarium pumps (VICAM cat. no. 21045) were used when testing multiple specimens.

Aflatoxin Panel Results:

[0081]

AFLATOXIN PANEL			
Specimen	0-0.5 ppb	>0.5 ppb	# of Tests
CSF	2	0	2
Nasal Secretions	19	4	23
Tissue Block	6	3	9
Urine	9	24	33
Other	3	0	3
Total:	39	31	70

Example 5

Detection of Ochratoxins in Body Fluids and Tissues

[0082] Ochratoxin A (OA) is detected using OchraTest® (VICAM, L.P., Watertown, Mass.) which is a well-known quantitative method for the detection of OA. This assay permits the measurement of the major OA without the use of toxic solvents (i.e., chloroform or methylene chloride). Ochratoxin is a mycotoxin produced by the fungus *Aspergillus ochraceus* and also by several species of *Penicillium* fungi. To measure ochratoxin levels, samples from body fluids and tissues are prepared by subjecting samples to the treatments described in Examples 2 and 3. The extracts are then applied to the OchraTest® column, which contains specific monoclonal antibodies for OA. The ochratoxin binds to the antibodies on the column. The column is then washed to remove impurities from the immunoaffinity column. By passing an eluting solution through the column, the ochratoxin is removed from the antibody. The eluting solution can then be measured in a fluorometer.

[0083] In more detail, the procedure for body fluids and tissues was as follows. Tissue samples were prepared from fixed and paraffin-embedded tissues as described in Example 2 and body fluid samples were prepared as described in Example 3. One ml of the extracted tissue sample, prepared sinus washes, or prepared urine was then mixed with 10 ml of PBS. The 10 ml of sample was then placed over an OchraTest® column (VICAM) and filtered through the affinity col-

umn at a rate of 1-2 drops/second until air came through the column. As an alternative procedure, 1.0 ml of sample can be mixed with 1.0 ml of 80:20 HPLC Grade Methanol:water solution. Then one (1) ml of the sample+80:20 HPLC Methanol:water solution can be added to 10 mls of PBS in each immunoaffinity column (OchraTest® column (VICAM)) and filtered through the affinity column at a rate of 1-2 drops/second until air came through the column. Ten ml of an 80:20 solution of methanol:water was then passed through the column at a rate of 2 drops/second. Ten ml of PBS was then passed through the affinity column at 2 drops/second. For the alternative procedure, these last two steps are not done. The affinity column was then washed by passing 10 ml of 1× Mycotoxin Wash Buffer (OchraTest® VICAM) through the column at a rate of 1-2 drops/second until air came through the column. The affinity column was then eluted by passing 1.5 ml of OchraTest® eluting solution through the column at a rate of 1 drop/second and collecting all of the sample eluate (1.5 ml) in a glass cuvette. The sample was mixed well and the cuvette was placed immediately into the calibrated fluorometer (Sequoia-Turner Model 450). The sample was read after 60 seconds. The results of the test are recorded in parts per billion (nanograms/milliliter).

Fluorometer Calibration: The Sequoia-Turner Fluorometer, Model 450, was calibrated as defined by the manufacturer's guidelines (VICAM). The machine was calibrated using standards supplied by VICAM (Green Standard=1.5 and Red Standard=23; Mycotoxin Calibration Standards (1 vial each of 3 levels) were used). Standards were prepared from controls purchased from Trilog Laboratories (Missouri). Standards were 50 ppb, 25 ppb, and 2.5 ppb. The lower limit of detection was determined to be 2.0 ppb using calibrators and known recovery rates of samples from the VICAM Immunoaffinity columns.

Pump and Set Up: OchraTest™ affinity chromatography was performed with the OchraTest™ affinity column attached to a pump stand. The stand has a glass syringe barrel that serves as a reservoir for the column. A large plastic syringe with tubing and coupling provides air pressure to manually push liquids through the column. An adjustable air pump (VICAM Cat. no. 20650) can be attached to the pump tube instead of the large pump syringe barrel to operate without using hand pressure. Four and six position pump stands with aquarium pumps (VICAM Cat. No. 21045) are used when testing multiple specimens.

Specimen	OCHRATOXIN PANEL		# of Tests
	0-2.0 ppb	>2.0 ppb	
CSF	1	0	1
Nasal Secretions	11	0	11
Tissue Block	6	1	7
Urine	18	4	22
Other	3	0	3
Total:	39	5	44

Example 6

Detection of Tricothecenes in Body Fluids and Tissues

[0084] This assay provides a procedure for the quantitative detection of Tricothecenes including Roridin A, E, H and L-2, Satratoxin G and H, Isosatratoxin F, Verrucarol A and J, and Verrucarol in human tissue and human body fluids treated as described in Examples 2 and 3.

[0085] Tricothecenes were measured using a kit provided by Envirologic Inc, (ENVIROLOGIX QuantiTox Kit (EP-100) for Tricothecenes; Envirologic, Inc. Portland, Me.).

The Envirologic kit is based on a competitive Enzyme-Linked Immunosorbent Assay (ELISA). In the test, Tricothecenes in a sample compete with enzyme (horseradish peroxidase)-labeled satratoxin for a limited number of antibody binding sites on the surface of the test wells. After a simple wash step, the outcome of the competition is visualized using a color development step. As with all competitive immunoassays, sample concentration is inversely proportional to color development (i.e., darker color=lower concentration of Tricothecenes in the sample and lighter color=higher concentration of Tricothecenes in the sample).

[0086] All plate kit components were stored at 4-8° C. when not in use. The kit components were never exposed to temperatures greater than 37° C. or less than 2° C. All reagents were allowed to reach ambient temperature before use. Reagents or test well strips from one plate kit were not used with reagents or test well strips from a different plate kit. Substrate was not exposed to sunlight during pipetting or while incubating samples in the test wells.

[0087] PBS Extraction/Dilution Buffer was prepared by making 1.0 liter of PBS, pH 7.4±0.05 in deionized water. PBS was stored refrigerated when not being used to prevent bacterial contamination. PBS was warmed to room temperature prior to use. Roridin A calibrators in PBS Extraction/Dilution Buffer were made daily in PBS using a positive-displacement pipet. At the end of the day, any unused calibrators were disposed of by adding to a 10% bleach solution. The calibrators were as follows:

[0088] 18.0 ppb Roridin A calibrator=20 µl of 900 ppb stock solution into 980 µl PBS in a glass tube.

[0089] 2.0 ppb Roridin A calibrator=100 µl of 18 ppb calibrator into 800 µl PBS in a glass tube.

[0090] 0.2 ppb Roridin A calibrator=10 µl of 18 ppb calibrator in 890 µl PBS in a glass tube.

[0091] Negative Calibrator: use an aliquot of the PBS stock.

[0092] Once all of the components had reached room temperature, the plate was removed from the pouch. All calibrators, sample extracts, and pipettes were organized so that the samples and enzyme conjugates could be added to the wells in 10 minutes or less. Fifty 50 µl of negative calibrator (NC), 50 µl of each Roridin A calibrator (C1-C3) and 50 µl of each sample extract (S1-S8) were added rapidly to their respective wells. Fifty µl of Enzyme Conjugate (Satratoxin Enzyme Conjugate), was rapidly added to each well.

[0093] The contents of the wells were mixed thoroughly by moving the plate frame in a rapid circular motion on the bench top for 30-45 seconds. The wells were covered with tape or parafilm and incubated at ambient temperature for 45 minutes with or without shaking on an orbital shaker at 200 rpm. After incubation, the covering was removed and the contents of the wells removed into a suitable container. The wells were flooded with water and then shaken to empty the wells a total of five times. The plate was then inverted several times on a paper towel to remove as much water as possible.

[0094] A 100 µl aliquot of substrate was then added to each well and the contents thoroughly mixed as described in the preceding paragraph. The wells were covered with new tape or parafilm and incubated for 15 minutes at ambient temperature with or without an orbital shaker. To stop the color development reaction, 100 µl of 1.0 N hydrochloric acid was added to each well and the samples were mixed thoroughly. This turned the well contents yellow. The plate was read within 30 minutes of the addition of the HCl using a Multiskan MCC 341 Microplate Reader at a wavelength of 450 nm.

[0095] New kit lots were tested in parallel with the kit lots currently in use. In order to test the new lot, one sample from the test batch was processed twice in the same protocol using both the new kit lot and the old kit lot. If the results from the new lot differed from the current lot, the lot test was repeated or the reagent was discarded.

[0096] A semi-log curve fit for the standard curve was used to plot the points of the calibrators. See below as example in the data reduction worksheet.

Calib. Conc.	rep1	rep 2	Avg. Value	% CV	B/Bo	Log (Conc.)	Conc.
Neg. Ctl. Abs. =	1.208	1.201	1.205	0.4%			
0.2 ppb Abs. =	1.023	1.084	1.054	4.1%	0.87	-0.70	0.20 $R^2 = -0.9980$
2.0 ppb Abs. =	0.657	0.671	0.664	1.5%	0.55	0.30	2.00 Slope = -0.3524
18 ppb Abs. =	0.359	0.372	0.366	2.5%	0.30	1.26	18.00 Intercept = 0.7950

Samples were run in duplicate and were plotted to give results in parts per billion (ppb) or nanograms/ml (see FIGS. 1 and 2).

TRICOTHECENES PANEL			
Specimen	0-0.2 ppb	>0.2 ppb	# of Tests
CSF	0	2	2
Nasal Secretions	26	16	42
Tissue Block	5	8	13
Urine	5	104	109
Other	4	2	6
Total:	40	132	172

Example 7
Detection of Fungal DNA

[0097] The target organisms for detection of fungal DNA using a PCR-based approach are shown in the following table. The table also indicates the primer-probe set (number in parentheses in the table) used for amplification and detection. The primers and probes used for amplification and detection of fungal DNA are also shown below.

Targets:				
<i>Aspergillus</i>	<i>Penicillium</i>	<i>Stachybotryous</i>	<i>Fusarium</i>	Control
(24) <i>amstelodami</i>	(15) <i>aurantiogriseum</i>	(1) <i>chartarum</i>	(14) <i>solani</i>	(6) GEO
(7) <i>flavus</i>	(4) <i>chrysogenum</i>	(23) <i>echinata</i>		
(8) <i>fumigatus</i>	(16) <i>citrinum</i>			
(9) <i>nidulans</i>	(17) <i>corylophilum</i>			
(3) <i>niger</i>	(18) <i>crustosum</i>			
(10) <i>ochraceus</i>	(20) <i>fellutanum</i>			
(11) <i>parasiticus</i>	(21) <i>roquefortii</i>			
(12) <i>sydowii</i>	(22) <i>simplisticum</i>			
(13) <i>ustus</i>	(5) <i>verrucosum</i>			
(2) <i>versicolor</i>	(19) <i>expansum</i>			

Primers and Probes:

[0098]

Target 1 <i>S. chartarum</i>	
Probe 1 char: 5'-ttgcttcgpcgggaacgccccg	(SEQ ID NO: 1)
Primer F2: 5'-gcccgggatcattaccgag	(SEQ ID NO: 2)
Primer R2: 5'-atcgatgccagaccaagag	(SEQ ID NO: 3)
Target 2 <i>A. versicolor</i>	
Probe 2 vers: 5'-cggggagccctctcgggggc	(SEQ ID NO: 4)
Primer F1: 5'-cgtaggtgaacctgcggaag	(SEQ ID NO: 5)
Primer R1: 5'-atcgatgccggaaccaagag	(SEQ ID NO: 6)
Target 3 <i>A. niger</i>	
Probe 3 niger: 5'-tgtctattgtacctgttgcttc	(SEQ ID NO: 7)
Primer F14: 5'-cgtaggtgaacctgcggaag	(SEQ ID NO: 8)
Primer R1: 5'-atcgatgccggaaccaagag	(SEQ ID NO: 9)
Target 4 <i>P. chrysogenum</i>	
Probe 4 chry: 5'-ctctgtctgaagattgtagtctgagt	(SEQ ID NO: 10)
Primer F1: 5'-cgtaggtgaacctgcggaag	(SEQ ID NO: 11)

-continued

Primer R1: 5'-atcgatgccggaaccaagag (SEQ ID NO: 12)

Target 5 *P. verrucosum*

Probe 5 verru: 5'-cccgcccttctgctggccgcc (SEQ ID NO: 13)

Primer F1: 5'-cgtaggtgaacctgcggaag (SEQ ID NO: 14)

Primer R1: 5'-atcgatgccggaaccaagag (SEQ ID NO: 15)

Target 6 *G. candidum*

For Geo F1H: 5'-ggatctcttggttctcgtatc (SEQ ID NO: 16)

Rev Geo R1H: 5'-cttgatctgaggttgaatagt (SEQ ID NO: 17)

Probe 6 geo: 5"-aacgcacattgcactttgggtatc (SEQ ID NO: 18)

Target 7 *A. flavus*

Probe 7 flav: 5'-cccgccattcatggccgcccgg (SEQ ID NO: 19)

Primer F1: 5'-cgtaggtgaacctgcggaag (SEQ ID NO: 20)

Primer R1: 5'-atcgatgccggaaccaagag (SEQ ID NO: 21)

Target 8 *A. fumigatus*

Probe 8 fumi: 5'-aaagtatgcagtctgagttgattatc (SEQ ID NO: 22)

Primer F1: 5'-cgtaggtgaacctgcggaag (SEQ ID NO: 23)

Primer R1: 5'-atcgatgccggaaccaagag (SEQ ID NO: 24)

Target 9 *A. nidulans*

Probe 9 nid: 5'-cccagggggcgagcccgg (SEQ ID NO: 25)

Primer F1: 5'-cgtaggtgaacctgcggaag (SEQ ID NO: 26)

Primer R1: 5'-atcgatgccggaaccaagag (SEQ ID NO: 27)

Target 10 *A. ochraceus*

Probe 10 ochr: 5'-acaccaacgtgaacctgtctgaag (SEQ ID NO: 28)

Primer F1: 5'-cgtaggtgaacctgcggaag (SEQ ID NO: 29)

Primer R1: 5'-atcgatgccggaaccaagag (SEQ ID NO: 30)

Target 11 *A. paraciticus*

Probe 11 para: 5'-cgggcccgccgtcatggccg (SEQ ID NO: 31)

Primer F1: 5'-cgtaggtgaacctgcggaag (SEQ ID NO: 32)

Primer R1: 5'-atcgatgccggaaccaagag (SEQ ID NO: 33)

Target 12 *A. sydowii*

Probe 12 syd: 5'-ccctcgggggcgagcccgg (SEQ ID NO: 34)

Primer F1: 5'-cgtaggtgaacctgcggaag (SEQ ID NO: 35)

Primer R1: 5'-atcgatgccggaaccaagag (SEQ ID NO: 36)

Target 13 *A. ustus*

Probe 13 ust: 5'-ccacaccgaacctcttgttatagc (SEQ ID NO: 37)

Primer F1: 5'-cgtaggtgaacctgcggaag (SEQ ID NO: 38)

Primer R1: 5'-atcgatgccggaaccaagag (SEQ ID NO: 39)

Target 14 *F. solani*

Probe14salani: 5'-cggaatagacggccccgtgaaac (SEQ ID NO: 40)

Primer F2: 5'-gcgaggatcattaccgag (SEQ ID NO: 41)

Primer R2: 5'-atcgatgccagagccaagag (SEQ ID NO: 42)

Target 15 *P. aurantiogriseum*

Probe 15 auran: 5'-cccgccttactggccgcccgg (SEQ ID NO: 43)

Primer F1: 5'-cgtaggtgaacctgcggaag (SEQ ID NO: 44)

-continued

Primer R1: 5'-atcgatgccggaaccaagag (SEQ ID NO: 45)

Target 16 *P. citrinum*
 Probe 16 citr: 5'-tggtgctcgggggccccgc (SEQ ID NO: 46)

Primer F4: 5'-ggaaggatcattaccgagtg (SEQ ID NO: 47)

Primer R1: 5'-atcgatgccggaaccaagag (SEQ ID NO: 48)

Target 17 *P. corylophilum*
 Probe 17 corylo: 5'-ttattgtaccttgcttcggcgg (SEQ ID NO: 49)

Primer F1: 5'-cgtaggtgaacctgcggaag (SEQ ID NO: 50)

Primer R1: 5'-atcgatgccggaaccaagag (SEQ ID NO: 51)

Target 18 *P. crustosum*
 Probe 18 crust: 5'-cgatctccggggacgggcc (SEQ ID NO: 52)

Primer F7: 5'-ctgtccgagcgtcattgctg (SEQ ID NO: 53)

Primer R5: 5'-cgaggaccggacgcggtg (SEQ ID NO: 54)

Target 19 *P. expansum*
 Probe19expan: 5'-agacacccccgaactctgctgaa (SEQ ID NO: 55)

Primer F1: 5'-cgtaggtgaacctgcggaag (SEQ ID NO: 56)

Primer R1: 5'-atcgatgccggaaccaagag (SEQ ID NO: 57)

Target 20 *P. fellutanum*
 Probe 20 fell: 5'-cccgcctgccaggcgcg (SEQ ID NO: 58)

Primer F1: 5'-cgtaggtgaacctgcggaag (SEQ ID NO: 59)

Primer R1: 5'-atcgatgccggaaccaagag (SEQ ID NO: 60)

Target 21 *P. roquefortii*
 Probe 21 roque: 5'-caccogtgtttatttaccttattgc (SEQ ID NO: 61)

Primer F1: 5'-cgtaggtgaacctgcggaag (SEQ ID NO: 62)

Primer R1: 5'-atcgatgccggaaccaagag (SEQ ID NO: 63)

Target 22 *P. simplicissimum*
 Probe 22 simpl: 5'-caccogtgtttatcgctacctgttg (SEQ ID NO: 64)

Primer F1: 5'-cgtaggtgaacctgcggaag (SEQ ID NO: 65)

Primer R1: 5'-atcgatgccggaaccaagag (SEQ ID NO: 66)

Target 23 *S. echinata*
 Probe 23 echin: 5'-ttgcttcggcgggagagccccg (SEQ ID NO: 67)

Primer F2: 5'-gcgaggatcattaccgag (SEQ ID NO: 68)

Primer R2: 5'-atcgatgccagagccaagag (SEQ ID NO: 69)

Target 24 *E. amstelodami*
 Probe 24 amst: 5'-tgtctatctgtaccctgttgcttcg (SEQ ID NO: 70)

Primer F1: 5'-cgtaggtgaacctgcggaag (SEQ ID NO: 71)

Primer R1: 5'-atcgatgccggaaccaagag (SEQ ID NO: 72)

Fungal Universal Group 1
 UP1: 5'-cctcggatcaggtaggatac (SEQ ID NO: 73)

UF1: 5'-atgcctgtccgagcgtcatt (SEQ ID NO: 74)

UR1: 5'-ttcctccgcttattgatatg (SEQ ID NO: 75)

-continued

Fungal Universal Group 2
 Up2: 5'-acggatctcttggctctggcatc (SEQ ID NO: 76)
 F2: 5'-gCGGAGGGATcattaccgag (SEQ ID NO: 77)
 UR2: 5'-ttcaactgaattctgcaattcac (SEQ ID NO: 78)

Extraction Methods:

[0099] Bead Beater Tube Preparation:

[0100] 1. 0.3 g±0.01 g of silica bead beating glass (Sigma-Aldrich Cat. no G1277) was added to 2 ml screw cap tube avoiding glass beads in the cap or around the rim.

[0101] 2. The tubes containing the beads were sterilized in an autoclave on the dry cycle for 10 minutes.

[0102] 3. The tubes were removed from the autoclave (proceed to the next step).

[0103] Solution Preparation:

[0104] 4. Buffers ATL (from DNeasy® Tissue Kit, Cat. no. 69506 (Quiagen, Stanford Valencia, Calif.)) and AL (from DNeasy® Tissue Kit, Cat. no. 69506) may form precipitates upon storage. If a precipitate formed in either buffer, the buffer was incubated at 55° C. until the precipitate fully dissolved.

[0105] 5. Buffers AW1 and AW2 (from DNeasy® Tissue Kit, Cat. no. 69506) were supplied as concentrates. Before using for the first time, the appropriate amounts of ethanol (96-100%) were added to Buffers AW1 and AW2 as indicated on the bottles.

[0106] 6. A 55° C. heat block and a 70° C. heat block were prepared for use in the assay.

[0107] Preparation of the Spore Solution or Tissue:

[0108] 7. If frozen material was used, it was equilibrated to room temperature.

[0109] 8. About 25.0 mg of paraffin-embedded tissue was weighed or 10.0 µl of spore solution was placed in a 2.0 ml screw cap tube.

[0110] 9. 180.0 µl of ATL Buffer and 20.0 µl of Proteinase K was added to each sample making sure that the lysate was not gelatinous.

[0111] 10. 10.0 µl of the Geo Spore reference DNA was added to each sample. (See Assay Specific Procedure for information regarding internal and external controls)

[0112] 11. All samples were bead beaten on the Bead Beater for 1 minute at the speed of 45.

[0113] 12. Samples were incubated at 55° C. on a pre-warmed heat block for 1 hour.

[0114] Extraction of Nucleic Acid:

[0115] 15. The samples were removed from the heat block and vortexed 15 seconds.

[0116] 16. 200 µl of Buffer AL was added and incubated at 70° C. for 10 minutes.

[0117] 17. The tubes were removed from the 70° C. heat block and add 200 µl of ethanol

[0118] 18. 200 µl of ethanol was added to each tube and vortexed.

[0119] 19. The mixture underneath the layer of paraffin was pipetted for each sample, making sure not to pipette the silica beads, into the corresponding DNeasy® Mini Spin Column 2 ml collection tube combo for that sample.

[0120] 20. The columns were centrifuged in a microcentrifuge at 8000 RPM for 1 minute. The collection tube containing the flow through was discarded.

[0121] 21. Each spin column was placed in a new 2.0 ml collection tube.

[0122] 22. 500.0 µl of Buffer AW1 was added to each column and centrifuged at 8000 RPM for 1 minute. The collection tube containing the flow through was discarded.

[0123] 23. Each spin column was placed in a new 2.0 ml collection tube.

[0124] 24. 500.0 µl of Buffer AW2 was added to each column and centrifuged at 13,000 RPM for 5 minute.

[0125] 25. The spin columns were removed carefully from the collection tubes so as not to splash nozzles. The collection tube containing the flow through was discarded.

[0126] 26. The spin columns were placed in their corresponding 1.5 ml elution tube.

[0127] 27. 100.0 µl of Buffer AE (from DNeasy® Tissue Kit, Cat. no. 69506) was placed into each spin column and incubated for 3 minutes at room temperature.

[0128] 28. The spin columns were centrifuged at 8000 RPM for 1 minute. The spin columns were discarded and capped and the extracted nucleic acid samples were stored at -20° C.

Real-Time PCR:

[0129] Preparation and Reaction Setup**[0130]** 1. Dilution of Probe Stocks

[0131] a. Resuspend the lyophilized probes in PCR grade water to a final concentration of 100 µM.

[0132] (Example: If the synthesis yields 15.03 nMoles, add 150.3 of PCR grade water to achieve 100 µM concentration)

[0133] 2. Dilution of Primer Stocks

[0134] a. Resuspend the lyophilized primers in PCR grade water to a final concentration of 100 µM.

[0135] (Example: If the synthesis yields 38.6 nMoles, add 386 µl of PCR grade water to achieve 100 µM concentration)

[0136] 3. Preparation of Primer/Probe Working Stock

[0137] a. See Appendix A for the working Stock setup for each target.

[0138] 4. Reaction Setup

[0139] a. The reaction setup for one reaction is shown below. In some cases the addition of MgCl₂ or varying concentrations of primer/probe mix is required for PCR. (See Appendix A)

DNA	5.0 µl
Primer/Probe Working Stock	3.5 µl (Final Concentration see appendix A)
OmniMix Beads	0.5 µl Beads (no volume contribution)
PCR Grade Water	16.5 µl
Total	25.0 µl

(Note that during reaction setup a master mix will be prepared for multiple reactions using the Smart Cycler PCR Worksheet (20.4007F). See following sections.)

[0140] 5. Smart Cycler Cycling Parameters (Omni Fungal I)

[0141] a. Omni Fungal I is the primary program used for the fungal real time assays and the run parameters for this program are outlined below. Cases may occur where changes to this program may be necessary for a specific target or specimen type. See SmartCycler Operation (20.2008S) for further instruction on programming and run optimization utilizing the Smart Cycler software.

[0142] Step 1 (1 Cycle)

[0143] Hot Start: 95° C. for 120 seconds

[0144] Step 2 (45 cycles)

[0145] Denature: 95° C. for 5 seconds

[0146] Anneal: 60° C. for 45 seconds

[0147] See Example PCR Worksheet: (Note: sheet has been truncated to show 3 target sets.)

II. Master Mix Setup

Reagent	Lot #	Volume (uL)	Reaction No.	Total Amount	Target
Set 1					
1 H2O	4532	16.5	6	99.0	(3) A niger
2 P/P Working Stock	040505	3.5	6	21.0	
3 Omni Mix (Bead)	2456	0.5	6	3.0	
4 MgCl2	NA	0.0	0	0.0	
Set 2					
1 H2O	4532	16.5	6	99.0	(2) A Versicolor
2 P/P Working Stock	020705	3.5	6	21.0	
3 Omni Mix (Bead)	2456	0.5	6	3.0	
4 MgCl2	NA	0.0	0	0.0	
Set 2					
1 H2O	4532	16.5	4	66.0	(6) Geo
2 P/P Working Stock	020705	3.5	4	14.0	
3 Omni Mix (Bead)	2456	0.5	4	2.0	
4 MgCl2	NA	0.0	0	0.0	

*Add MgCl₂ as needed per target subtract volume used from water added to maintain a 20 µl reaction. Add 20 µl of Master Mix to each tube and then add 5.0 µl of template for a total volume of 25.0 µl.

APPENDIX A

Target Working Stock Recipes

Target	P1 100uM	Final Conc.	P2 100uM	Final Conc.	Probe 100uM	Final Conc.	MgCl2 1 M	Final Conc.	Water	Final Volume
1	15.0 uL	.3 uM	15.0 uL	.3 uM	10.0 uL	.2 uM	NA	NA	660 uL	700 uL
2	30.0 uL	.6 uM	30.0 uL	.6 uM	20.0 uL	.4 uM	NA	NA	620 uL	700 uL
3	30.0 uL	.6 uM	30.0 uL	.6 uM	20.0 uL	.4 uM	30.0 uL	6 mM	590 uL	700 uL
4	15.0 uL	.3 uM	15.0 uL	.3 uM	10.0 uL	.2 uM	NA	NA	660 uL	700 uL
5	15.0 uL	.3 uM	15.0 uL	.3 uM	10.0 uL	.2 uM	NA	NA	660 uL	700 uL
6	15.0 uL	.3 uM	15.0 uL	.3 uM	10.0 uL	.2 uM	NA	NA	660 uL	700 uL
7	15.0 uL	.3 uM	15.0 uL	.3 uM	10.0 uL	.2 uM	NA	NA	660 uL	700 uL
8	15.0 uL	.3 uM	15.0 uL	.3 uM	10.0 uL	.2 uM	20.0	4 mM	640 uL	700 uL
9	15.0 uL	.3 uM	15.0 uL	.3 uM	10.0 uL	.2 uM	NA	NA	660 uL	700 uL
10	15.0 uL	.3 uM	15.0 uL	.3 uM	10.0 uL	.2 uM	NA	NA	660 uL	700 uL
11	15.0 uL	.3 uM	15.0 uL	.3 uM	10.0 uL	.2 uM	20.0	4 mM	640 uL	700 uL
12	15.0 uL	.3 uM	15.0 uL	.3 uM	10.0 uL	.2 uM	NA	NA	660 uL	700 uL
13	30.0 uL	.6 uM	30.0 uL	.6 uM	20.0 uL	.4 uM	NA	NA	620 uL	700 uL
14	45.0 uL	.9 uM	45.0 uL	.9 uM	30.0 uL	.6 uM	NA	NA	580 uL	700 uL
15	15.0 uL	.3 uM	15.0 uL	.3 uM	10.0 uL	.2 uM	NA	NA	660 uL	700 uL
16	15.0 uL	.3 uM	15.0 uL	.3 uM	10.0 uL	.2 uM	NA	NA	660 uL	700 uL
17	15.0 uL	.3 uM	15.0 uL	.3 uM	10.0 uL	.2 uM	NA	NA	660 uL	700 uL
18	30.0 uL	.6 uM	30.0 uL	.6 uM	20.0 uL	.4 uM	20.0 uL	4 mM	600 uL	700 uL
19	15.0 uL	.3 uM	15.0 uL	.3 uM	10.0 uL	.2 uM	20.0	4 mM	640 uL	700 uL
20	15.0 uL	.3 uM	15.0 uL	.3 uM	10.0 uL	.2 uM	NA	NA	660 uL	700 uL
21	15.0 uL	.3 uM	15.0 uL	.3 uM	10.0 uL	.2 uM	NA	NA	660 uL	700 uL

APPENDIX A-continued

Target Working Stock Recipes										
Target	P1 100uM	Final Conc.	P2 100uM	Final Conc.	Probe 100uM	Final Conc.	MgCl2 1 M	Final Conc.	Water	Final Volume
22	15.0 uL	.3 uM	15.0 uL	.3 uM	10.0 uL	.2 uM	20.0	4 mM	640 uL	700 uL
23	15.0 uL	.3 uM	15.0 uL	.3 uM	10.0 uL	.2 uM	NA	NA	660 uL	700 uL
24	15.0 uL	.3 uM	15.0 uL	.3 uM	10.0 uL	.2 uM	NA	NA	660 uL	700 uL

Mix target working stocks for each target as described in the table above by combining the amounts noted for Primer 1, Primer 2, Probe, MgCl₂ (if needed), and water. Use 3.5 µl of this working stock for each reaction performed.

Master Mix Preparation:

- [0148] 1. All steps were performed under sterile conditions.
- [0149] 2. After the water and beads had been pipetted into the individual tubes, the tubes were mixed until the beads (Cat no. Omni 1-100N-050; Cepheid, Sunnyvale, Calif.) were completely dissolved.
- [0150] 3. After the beads were dissolved, the appropriate primer/probe working stock was pipetted into each master mix tube as described in the PCR worksheet.
- [0151] 4. The solutions were mixed completely and the working stocks returned to the -20° C. freezer.
- [0152] 5. Controls—
- [0153] a. Internal Control—Every clinical sample processed was inoculated with spores from the internal control target Geometrica to show that a negative target result is a true negative result and not related to the extraction of the sample. The samples were processed through the extraction protocol and amplified and detected utilizing primer and probes specific for Geometrica.
- [0154] b. Positive Control—A positive control for each target of interest (Primer/Probe sets) was processed along with each clinical sample in each real-time PCR run. This positive control can be extracted from tissue or spore solutions but must be lot checked prior to use. The positive control shows that the primer/probe set for each target is not being inhibited and shows that a negative result is a true negative.
- [0155] c. Negative Control—A negative control for each target of interest (Primer/Probe sets) was processed along with each clinical sample in each real-time PCR run. This negative control can be extracted from tissue or water but must be lot checked prior to use. The negative control shows that the primer/probe set, water and extraction reagents for each target is not contaminated with the target and shows that a positive result is a true positive.
- [0156] Addition of Target Nucleic Acid:
- [0157] 1. 5.0 µl of the negative control, positive control and patient samples was pipetted into the appropriate reaction tubes.
- [0158] 2. The reaction tubes were centrifuged using the Smart Cycler® II modified centrifuge.
- [0159] 3. The tubes were returned to the cooling block and stored at 4° C. or the Smart Cycler Setup and Run was conducted.
- [0160] Smart Cycler Setup and Run:
- [0161] 1. The Omni Fungal I protocol or the appropriate protocol was selected for this real-time run.
- [0162] 2. For information regarding the operation of the Smart Cycler see SmartCycler Operation (20.2008S) (Smart Cycler® II Instrument; Cepheid, Sunnyvale, Calif.).
- [0163] Data Analysis:
- [0164] 1. After the run is completed the results were analyzed by reviewing
- [0165] Launch Internet Explorer Browser. Ink each site in the results table. If a specific sample tested was registered as positive by the software there was a positive in the results column for that sample. There was also a crossing point registered in the Ct column for that sample.
- [0166] 2. After reviewing the Results Table, the curves were reviewed for each sample by selecting the “FAM” or “Log FAM” of the “Views” menu.
- [0167] 3. With the graph selected, all samples that created a curve were present on the screen. Each sample was reviewed independently by clicking on the Site ID associated with the sample of interest located just to the right of the graph.
- [0168] 4. A sample was analyzed as positive by the software if the curve broke the baseline of 30 (default set in section above) before the end of the 45 cycles and negative if it did not break the baseline of 30 before the end of the 45 cycles.
- [0169] 5. Each sample was reviewed and then highlighted so that all sample curves were present on the graph.
- [0170] Results Interpretation:
- [0171] 1. Positive Result: A positive result is defined as any amplification observed crossing a baseline fluorescence of >30 between cycles 1 and 40 of the real-time PCR run.
- [0172] 2. Negative Result: A negative result is defined as no amplification observed crossing a baseline fluorescence of >30 between cycles 1 and 40 of the PCR run.
- [0173] 3. Equivocal Result: An equivocal result is defined as no amplification observed crossing a baseline fluorescence of >30 between cycles 1 and 40, a control out of range or questions regarding sample integrity.
- [0174] 4. Positive Control: A control that is positive for the target being tested and shows that the assay will show a positive in the presence of target spores and that there is not PCR inhibition.
- [0175] 5. Negative Control: A control that is negative for the target being tested and shows that the reagents or the sample were not contaminated with the target prior to the testing of the sample.
- [0176] 6. Internal Control: A control used to show that the extraction process is working fine for the purification

of nucleic acid from the clinical specimen and that a negative result is truly negative and not due to an issue associated with the extraction. (Note: the internal control must be positive for any sample to be reported as negative for a target.)

See Table Below:

[0177]

	Crossing Point	Positive Control	Negative Control	Internal Control
Reportable Result				
Positive Result	≧40	(+)	(-)	(+)
Positive Result	≧40	(-)	(-)	(+)
Positive Result	≧40	(+)	(-)	(-)
Positive Result	≧40	(-)	(-)	(-)
Negative Result	(-)	(+)	(-)	(+)
Negative Result	(-)	(+)	(+)	(+)
Negative Result	(-)	(-)	(+)	(+)
Un-reportable Result				
Positive Result	≧40	(+)	(+)	(+)
Positive Result	≧40	(-)	(+)	(+)
Positive Result	≧40	(+)	(+)	(-)
Positive Result	≧40	(-)	(+)	(-)
Negative Result	(-)	(-)	(-)	(+)
Negative Result	(-)	(+)	(-)	(-)
Negative Result	(-)	(+)	(+)	(-)
Equivocal Result				
Case by Case	Case by Case	Case by Case	Case by Case	Case by Case

In other illustrative embodiments, results can be determined based on a cycle range between cycles 1 and 45 of the PCR run or other useful ranges can be used.

DNA Results—

[0178]

Tissue	Pos Results/ tissue	Neg Results/ tissue	Isolates**
Lung	6	1	<i>Aspergillus fellutanum</i> (1) <i>Asp. niger</i> (2) <i>P. crysogenum</i> (3)
Liver	5	2	<i>Asp. flavus</i> (2) <i>Asp. Niger</i> (3)

-continued

Tissue	Pos Results/ tissue	Neg Results/ tissue	Isolates**
Brain	4	2	<i>Pen. versicolour</i> (2) <i>Asp. Fumigatus</i> (1) <i>Asp niger</i> (2) <i>Asp fumigatus</i> (2) <i>Asp. flavus</i> (2)
Skin	2	1	<i>Penicillium fellutanum</i> (1) <i>Pen. crysogenum</i> (1) <i>Asp. ustus</i> (1)
Respiratory###	3	6	<i>Asp. niger</i> (2) <i>Asp. flavus</i> (2) <i>Pen. versicolour</i> (1)

**some specimens revealed more than one isolate
###includes nasal secretions, inner ear fluids, and sputums

The test is 100% specific for the isolates tested for. 15 known negative (non-exposed) patients were tested in validation. No positive results with the probes in this patent were found. Of the 29 suspected exposed patients tested, 29 samples of tissue were tested. Of those 29 tissues/fluids, 20 specimens gave results of one or more organisms by DNA probing using RT-PCR.

Example 8

Detection of Fungal DNA Using Universal Primers

[0179] The purpose of the universal fungal assays is to check clinical samples for the presence of fungal DNA prior to running multiple fungal target specific assays. Two universal assays were designed for this purpose. Fungal assay UP1 was designed to detect the presence of fungal *Aspergillus* and *Penicillium* species. Fungal assay UP2 was designed to detect the presence of *Stachybotrys* and *Fusarium* species. Each assay is composed of two primers and one Taqman probe specific for fungal targets of the species described above.

[0180] To identify sequence present in 1.) all *Aspergillus* and *Penicillium* species, and in 2.) all *Stachybotrys* and *Fusarium* species a search was performed for public sequence and sequence was identified. A sequencing primer (oligonucleotide ~20 bases long) was designed specific for each of the targets of interest, and was ordered from a vender. The primers and target DNAs were sequenced, and the accuracy confirmed. Primers and probes were designed and were expected to be specific for all individual targets in each species group as described below. To check for specificity to the species groups described, the sequences of each assay were put through a “Blast” search against known sequences from fungal genomes. Initial results showed these sequences to be specific for the species as described above and not specific for clinically relevant targets outside the species. The following sequences were found to be homologous for all targets of interest related to *Aspergillus* and *Penicillium*:

Assay UP1

[0181]

(SEQ ID NO: 74)
UP1 atgctgtccgagcgtcatt (Forward Primer)

-continued

(SEQ ID NO: 75)
UR1 ttcctccgccttattgatatg (Reverse Primer)

(SEQ ID NO: 73)
UP1 cctcggatcaggtaggatac (Taqman probe)

The following sequences were found to be homologous for all targets of interest related to *Stachybotrys* and *Fusarium*:

Assay UP2

[0182]

(SEQ ID NO: 77)
F2 gggaggatcattaccgag (Forward Primer)

(SEQ ID NO: 78)
UR2 ttcactgaattctgcaattcac (Reverse Primer)

(SEQ ID NO: 76)
Up2 acggatctctggctctggcattc (Taqman probe)

Example 9

Detection of Mycotoxins Using Luminex®

[0183] The purpose of this assay is to utilize the Luminex® platform to detect approximately 3 toxin groups (trichothecenes, aflatoxins, and ochratoxins) in patient samples that have been exposed to fungal targets belonging to *Aspergillus*, *Penicillium*, *Stachybotrys*, and *Fusarium* species. The Luminex® assay utilizes microspheres (beads) that are coupled to antigens to detect antibodies against those specific antigens in a sample. Samples and coupled microspheres will be incubated in microtitration filter wells where antigen-antibody binding occurs. After incubation and washing, the appropriate detection antibody (e.g., biotinylated antibody) will be introduced and incubated during which antibody-antibody binding occurs. After incubation and washing, a reporter conjugate will be added and incubated where the biotin-binding reaction occurs.

[0184] In theory, each microsphere is color-coded into 100 different sets. Each bead set can be coated with a reagent to capture and detect a specific analyte from a sample. The Luminex® 100 has lasers that excite the internal dyes that identify the microsphere and any reporter dye captured during the assay. During the run on the Luminex®, several readings will be made on each of the bead sets. Potentially, this will create a multiplexing capability of up to 100 unique assays with one single sample.

Example 10

Detection of Fungal DNA and Mycotoxins in Human Patients

Findings for Patient #1 (Skin Biopsy)—

[0185] Positive for *P. chrysogenum* and *A. ustus*
Positive for Trichothecenes (38 ppb)

Findings for Patient #2 (Endometriosis DX in Ovary and Uterus)—

[0186] Positive for Trichothecenes (25 ppb in ovary pathology specimen)

Findings for Patient #3 (Autopsy Specimen)—

[0187] Positive for *A. fellutanum* (lung)
Positive for *A. flavus* and *A. niger* (liver)
Positive for *A. niger* (brain)

Findings for Patient #4 (Questionable Fungal Mass on Neck)—

[0188] Positive for *A. niger* (left and right inner ear)
Positive (left ear) for aflatoxin (9.5 ppb), ochratoxin (4.5 ppb), and Trichothecenes (0.62 ppb)
Positive (right ear) for aflatoxin (0.73 ppb) and Trichothecenes (0.62 ppb)

Findings for Patient #5 (Pilocystic Astrocytoma, Brain Tumor)

[0189] 6 year old female, Grade 1-2 Pilocystic astrocytoma, Brain
RT-PCR: Negative for 23 probes
Mycotoxins: aflatoxins (3.0 ppb), all other mycotoxins negative.

Findings for Patient #6 (Diagnosis: Pulmonary Fibrosis, Potential Lung Transplant Patient).

[0190] 38 year old male on oxygen. Previous lung biopsy: severe Interstitial fibrosis (UIP).
All bacterial and fungal cultures negative.
RT-PCR: negative for 23 probes.
Mycotoxins: aflatoxin (4 ppb), all other mycotoxin negative
Treated with antifungal agent, and, after six weeks, he improved and was taken off lung transplant emergent list. Oxygen use decreased significantly.

Findings for Patient #7: (Headache and Seizures)

[0191] 32 year old female with severe headaches and seizures. All diagnostic tests negative. All cultures negative.
RT-PCR: positive for *Stachybotrys chartarum*.

Findings for Patient #8 (Mouth Lesions for Two Years, No Resolution)

[0192] 84 year old female with three mouth lesions, chief complaint: pain and mouth feels like "cotton".
Biopsy: parakeratosis, ulceration, no malignant cells, chronic inflammation, increased vascularity. Diagnosis: consistent with history of aphthous stomatitis.
RT-PCR: positive for *Aspergillus sydowii*.

Findings for Patient # 9 (Chronic Headaches)

[0193] 40 year old male with chronic headaches
Pathology biopsy: Temporal dura and lobe: Perivascular inflammatory cell infiltrate. No neoplasm identified. No organisms noted by stain or culture or PCR for virus or bacteria.
RT-PCR: positive for *Aspergillus ustus* and *Emericella amstelodami*.

Mycotoxins: Negative

[0194] Findings for Patient #10 (Patient with Skin Lesions)
56 year old male exposed to *Stachybotrys*, *Aspergillus niger*, and *Penicillium* sp. (confirmed by Environmental evaluation).
RT-PCR: Skin Biopsy: Positive for *Stachybotrys chartarum*, *Aspergillus niger*.
Mycotoxins: Positive for Aflatoxin (4 ppb); Negative for trichothecenes.

Example 11

Detection of Mycotoxins

[0195] The purpose of this assay is to utilize the Luminex® platform to detect approximately 29 toxins in patient samples which include but are not limited to blood, spinal fluid, urine, nasal secretions, sputum, and tissue that have been exposed to fungal targets belonging to *Aspergillus*, *Penicillium*, *Stachybotrys*, and *Fusarium* species.

[0196] The Luminex® assay can, for example, utilize microspheres (beads) that are coupled to monoclonal antibodies to detect mycotoxin antigens. The monoclonal antibodies will be made or purchased for use in this procedure. The antigens are, for example, aflatoxin B1, B2, G1, and G2, Ochratoxin A, and a group of Trichothecenes. The group of Trichothecenes includes Roridin A, E, H and L-2, Satratoxin G and H, Isosratoxin F, Verrucaridin A and J, and Verrucarol, and T-2. Samples and coupled microspheres will be incubated in microtitration filter wells where antigen-antibody binding occurs. After incubation and washing, the appropriate detection antibody (e.g., biotinylated antibody) will be introduced and incubated during which antigen-antibody binding occurs. After incubation and washing, a reporter conjugate will be added and incubated where the biotin-binding reaction occurs.

[0197] Each microsphere is color-coded into, for example, 100 different sets. Each bead set can be coated with a reagent to capture and detect a specific analyte from a sample. The Luminex® 100 has lasers that excite the internal dyes that identify the microsphere and any reporter dye captured during the assay. During the run on the Luminex®, several readings will be made on each of the bead sets. Potentially, these assays will create a multiplexing capability of up to 100 unique assays with a single sample.

Example 12

Luminex Assay Using DNA Probes Bound to Beads

[0198] Microspheres (Luminex Corporation, Austin, Tex.) are 5.6 µm in diameter and are comprised of polystyrene, divinyl benzene, and methacrylic acid with surface carboxylate functionality for covalent attachment of biomolecules. The microspheres are internally dyed with red, infrared-emitting fluorochromes. Spectral addresses were created by adjusting the concentrations of each fluorochrome with each bead set. When the microsphere sets were analyzed with the Luminex 100 instrument (Luminex), each bead set was identified and classified by a distinct fluorescence signature pattern.

[0199] Coupling: Two microsphere sets were used for covalent coupling of *A. versicolor* and *F. solani* probes to carboxyl functional groups on the microsphere surfaces. A concentration of 5.0×10^6 microspheres were added to separate 1.5 ml microcentrifuge tubes and pelleted by microcentrifugation and the supernatant was aspirated. The microsphere pellet was resuspended and vortexed in 50 µl of 0.1M 2-(N-morpholino)-ethanesulfonic acid (MES), pH 4.5, 0.2 nanomole of each 5' amino-modified capture oligonucleotide or probe (*A. versicolor*: 5'-CGGGGAGC-CCTTCGGGGGC-3' (SEQ ID NO: 4) and *F. solani*: 5'-CGGGAATAGACGGCCCCGTGAAAC-3' (SEQ ID NO:

40)) and 25 µg of 1-ethyl-3-(3-dimethylaminopropyl)-carbodiimide hydrochloride (EDC) (10 mg/ml in dH₂O), and the mixture was incubated for 30 minutes at room temperature in the dark. The microsphere mixture was vortexed and 25 µg of EDC was added again and mixed, and the mixture was incubated for 30 min at room temperature in the dark. One milliliter of 0.02% Tween-20 (0.02% in dH₂O) was added to the coupled microspheres and microcentrifuged. The supernatant was removed and the pellet was resuspended in 100 µl of 1×Tris EDTA, pH 8.0 (TE) (diluted in dH₂O), by vortex and sonication (see FIG. 3).

[0200] Assay: A working mixture of the microspheres was prepared at a concentration of 150 microspheres per microliter in 1.5×TMAC (5M TMAC, 20% Sarkosyl, 1M Tris-HCL, 0.5M EDTA, H₂O), vortexed and sonicated. Thirty-three microliters of each microsphere mixture was added to the wells of a 96-well V-bottom PCR plate and 17 µl TE, pH 8.0, were added to the background wells. To each appropriate well 5' biotinylated complementary oligonucleotides (*A. versicolor*: 5'-GCCCCCGAGAGGGCTCCCCG-3' (SEQ ID NO: 90) and *F. solani*: 5'-GTTTCACGGGGCCGTCTATCCCG-3' (SEQ ID NO: 91)) (5 to 200 femtomoles) and TE, pH 8.0, to a total well volume of 50 µl were added. The reaction wells were mixed, covered and incubated at 96° C. for 3 min to denature any secondary structure. The reactions were incubated for 15 min at 52° C. and 25 µl of a 10 µg/ml dilution of streptavidin-R-phycoerythrin in 1×TMAC (1.5×TMAC diluted in dH₂O) was added, mixed and incubated for 5 min at 52° C. Fifty microliters from each well was analyzed at 52° C. with a minimum bead count set at 100 events.

DataType: Averaged Median		
Sample (fmol)	<i>A. versicolor</i>	Total Events
0	96	215
5	1143	213
10	1344	215
20	1420	223
50	2513	205
100	4467	224
200	5206	202

DataType: Averaged Median		
Sample (fmol)	<i>F. solani</i>	Total Events
0	66	224
5	1470	213
10	1682	215
20	1740	205
50	1884	221
100	1974	210
200	1978	223

-continued

copied from output Data Type: Median			
Location	Sample	<i>A. versicolor</i>	Total Events
1	0	73	220
2	5	1358	221
3	10	1461	227
4	20	1639	230
5	50	2512	201
6	100	3886	229
7	200	5279	201
8	0	56	200
9	0	118	209
10	5	928	205
11	10	1227	202
12	20	1201	215
13	50	2514	209
14	100	5047	218
15	200	5133	203
16	0	104	226

Data Type: Median			
Location	Sample	<i>F. solani</i>	Total Events
17	0	99	210
18	5	1582	214
19	10	1636	204
20	20	1800	200
21	50	1887	225
22	100	1932	212
23	200	1960	219
24	0	136	223
25	0	32	237
26	5	1359	211
27	10	1728	226
28	20	1681	210
29	50	1882	216
30	100	2017	207
31	200	1997	227
32	0	94	204

DNA Coupling Raw Data

[0201]

SN	LX10001089024
Session	022107couplingconfirmation.RTL
Operator	
Samples	32
Min Events	0

Results

[0202]

Location	Sample	Aver	Fsol	Total Events	Notes
Data Type: Median					
1	0	73		220	
2	5	1358		221	

Location	Sample	Aver	Fsol	Total Events	Notes
3	10	1461		227	
4	20	1639		230	
5	50	2512		201	
6	100	3886		229	
7	200	5279		201	
8	0	56		200	
9	0	118		209	
10	5	928		205	
11	10	1227		202	
12	20	1201		215	
13	50	2514		209	
14	100	5047		218	
15	200	5133		203	
16	0	104		226	
17	0		99	210	
18	5		1582	214	
19	10		1636	204	
20	20		1800	200	
21	50		1887	225	
22	100		1932	212	
23	200		1960	219	
24	0		136	223	
25	0		32	237	
26	5		1359	211	
27	10		1728	226	
28	20		1681	210	
29	50		1882	216	
30	100		2017	207	
31	200		1997	227	
32	0		94	204	
Data Type: Count					
1	0	100	120	220	
2	5	121	100	221	
3	10	100	127	227	
4	20	130	100	230	
5	50	100	101	201	
6	100	129	100	229	
7	200	100	101	201	
8	0	100	100	200	
9	0	109	100	209	
10	5	105	100	205	
11	10	102	100	202	
12	20	115	100	215	
13	50	109	100	209	
14	100	118	100	218	
15	200	103	100	203	
16	0	100	126	226	
17	0	110	100	210	
18	5	114	100	214	
19	10	104	100	204	
20	20	100	100	200	
21	50	125	100	225	
22	100	112	100	212	
23	200	119	100	219	
24	0	100	123	223	
25	0	100	137	237	
26	5	100	111	211	
27	10	126	100	226	
28	20	100	110	210	
29	50	116	100	216	
30	100	100	107	207	
31	200	100	127	227	
32	0	104	100	204	

The data indicate that DNA probes can be bound to beads and can be used to detect the presence of fungal DNA in samples.

Example 13

Luminex Indirect Assay Using Antigen Bound to Beads

[0203] Microspheres (Luminex Corporation, Austin, Tex.) are 5.6 µm in diameter and are comprised of polystyrene, divinyl benzene, and methacrylic acid with surface carboxylate functionality for covalent attachment of biomolecules. The microspheres are internally dyed with red, infrared-emitting fluorochromes. Spectral addresses were created by adjusting the concentrations each fluorochrome with each bead set. When the microsphere sets were analyzed with the Luminex 100 instrument (Luminex®), each bead set was identified and classified by a distinct fluorescence signature pattern.

[0204] Coupling: Two microsphere sets were used for covalent coupling of mycotoxin antigens. Aflatoxin (B1, B2, G1, G2) and Deoxynivalenol (DON) were used and required chemical modifications of carboxyl functional groups on the microsphere surfaces to amine groups due to the absence of an alpha-amino N-terminal group on either antigen. Adipic acid dihydrazide (ADH) was used to modify the carboxyl functional groups on the microsphere surfaces to provide an NH₂ group for coupling to each antigen. A concentration of 1.25×10⁷ microspheres were added to separate 1.5 ml microcentrifuge tubes and were washed by adding 500 µl of 100 mM 2-(N-morpholino)-ethanesulfonic acid (MES), pH 6.0, microcentrifuged at 10,000×g for 1 min at room temperature and the supernatant was aspirated. The microsphere pellet was resuspended and vortexed in 1 ml of ADH and 200 µl of 1-ethyl-3-(3-dimethylaminopropyl)-carbodiimide hydrochloride (EDC) (200 mg/ml in 100 mM MES, pH 6.0) and the tubes were rotated for 1 h at room temperature in the dark. The microspheres were then washed twice with 1 ml of 100 mM MES, pH 4.5, by microcentrifugation as described above, and the supernatants were discarded. Five hundred microliters of each antigen (500 ppb Aflatoxin and 125 ppb DON) were added to each pellet of ADH-modified microspheres and the mixture was incubated at room temperature for 4 h with rotation. To remove any noncovalently bound mycotoxin, the microspheres were washed twice by centrifugation with PBS-Tween and were blocked with PBS-Tween (see FIG. 4).

[0205] Immunoassay: In order to test the coupling efficiency of each antigen, Aflatoxin (B1, B2, G1, G2) and Deoxynivalenol (DON), to each microsphere set, an indirect immunoassay must be performed. For the competitive assay, each mycotoxin antibody will be titrated in a two-fold dilution and incubated in micotiter wells of a 96-well plate containing mycotoxin antigen-coupled microspheres. The MFI signal should increase as the antibody is diluted to reach optimal antibody dilution required for the competitive immunoassay. Two blanks will be identified with a "0" which indicates the absence of mycotoxin antibody and a signal of less than 50 MFI (Median Fluorescent Signal). The increasing signal indicating the labeled mycotoxin antibody binding to the antigen-coupled microspheres and the low signal in the blank wells, where no labeled mycotoxin antibody is present, prove that the mycotoxin antigens were effectively coupled to the microsphere sets and can be detected by the appropriate mycotoxin antibodies.

[0206] Fifty microliters of each microsphere mixture were added to the wells of a 1.2 µm-pore-size filter membrane microtiter plate. Fifty microliters of PBS-1% BSA were added to each background well. Sheep Anti-Aflatoxin and

Sheep Anti-DON antibodies were purified by ammonium sulfate precipitation and desalted utilizing desalting spin columns and were labeled with biotin using sulfosuccinimidyl-6-[biotin-amido]hexanoate. Each mycotoxin antibody was diluted in PBS-1% BSA in 2-fold dilutions and 50 µl of each dilution were added to the standard wells. The wells were mixed by pipetting and the samples were incubated, with shaking, at room temperature in the dark for 1 h. The wells were aspirated using a vacuum manifold filtration system and were washed twice with 100 µl of PBS-1% PBS-BSA. The microspheres were resuspended in 50 µl PBS-1% BSA and 50 µl of 4 µg/ml Streptavidin-R-phycoerythrin was added to each well, mixed and incubated, with shaking, at room temperature in the dark for 30 min. The wells were washed twice with 100 µl PBS-1% BSA and the samples were resuspended in 100 µl PBS-1% BSA. Seventy-five microliters of each sample were analyzed with a minimum bead count set at 100 events.

[0207] Data: Mycotoxin DON and Aflatoxin were each coupled to two sets of Luminex microspheres. Biotinylated Sheep Anti-DON and Sheep Anti-Aflatoxin were titrated in a two-fold dilution to show a dose response.

Aflatoxin Data Type: Median			
Location	Antibody Dose	Aflatoxin	Total Events
1	0	10	100
2	0	11	100
3	Neat	106	100
4	1:2	218	100
5	1:4	152	100
6	1:8	868	100
7	1:16	2442	100
8	1:32	4166	100

DON Data Type: Median			
Location	Antibody Dose	DON	Total Events
1	0	4	100
2	0	4	100
3	Neat	8	100
4	1:2	13	100
5	1:4	89	100
6	1:8	149	100
7	1:16	1451	100
8	1:32	2162	100

Protein Raw Data for Microsphere Binding for Detection of Mycotoxins.

[0208]

Program	Luminex 100 IS	
Build	2.2	
Date	3/3/2007	3:16:21 PM
SN	LX10001089024	

-continued

Session	030307Afla.modbead.biotinAb		
Operator			
Samples	8	Min Events	0

Results

[0209]

Location	Sample	Aflatoxin	Total Events
Data Type: Median			
1	Blank	10	100
2	Blank	10.5	100
3	Neat Alfa	105.5	100
4	1:02	217.5	100
5	1:02	152	100
6	1:02	868	100
7	1:02	2442	100
8	1:02	4165.5	100
Data Type: Count			
1	Blank	100	100
2	Blank	100	100
3	Neat Alfa	100	100
4	1:02	100	100
5	1:02	100	100
6	1:02	100	100
7	1:02	100	100
8	1:02	100	100

Program	Luminex 100 IS		
Build	2.2		
Date	3/3/2007 3:13:49 PM		
SN	LX10001089024		
Session	030307DONmodbead.antibiotin		
Operator			
Samples	8	Min Events	0

Location	Sample	DON	Total Events
Data Type: Median			
1	Blank	3.5	100
2	Blank	3.5	100
3	Neat Don	7.5	100
4	1:02	12.5	100
5	1:02	89	100
6	1:02	148.5	100
7	1:02	1450.5	100
8	1:02	2161.5	100
Data Type: Count			
1	Blank	100	100
2	Blank	100	100
3	Neat Don	100	100
4	1:02	100	100
5	1:02	100	100
6	1:02	100	100
7	1:02	100	100
8	1:02	100	100

The data indicate that mycotoxins or mycotoxin antigens can be bound to beads and can be used to detect in a competitive assay the presence of antibodies to mycotoxins in samples.

Example 14

Detection of Fungal DNA and Mycotoxins in Animals

[0210] Findings for Dog (Necropsy, High Exposure to Environmental Molds):

[0211] Liver and Lung:

[0212] Mycotoxins: tricothecenes 0.96 ppb

[0213] RT-PCR: *Penicillium chrysogenum* and *Aspergillus niger*

[0214] Findings for Horse (Urine):

[0215] Tricothecenes: 5.32 ppb

[0216] Findings for Cat (Necropsy):

[0217] Aflatoxin: 4 ppb

Example 15

Combination Method for Mycotoxins and Mold Detection in Specimens

[0218] Human or animal patients exhibiting symptoms of exposure to molds and/or mycotoxins were evaluated as follows. A patient tissue or body fluid was first examined for the presence of mycotoxins (e.g., Tricothecenes, Aflatoxins B1, B2, G1, G2, and Ochratoxin A, for example). If a mycotoxin was present, the specimen was then tested for the presence of molds/fungi as follows:

Mycotoxin Present	DNA Extraction Test
Tricothecenes	<i>Stachybotrys chartarum</i> <i>Stachybotrys echinata</i> <i>Fusarium solani</i>
Aflatoxins	<i>Aspergillus flavus</i> <i>Aspergillus ochraceus</i> <i>Aspergillus fumigatus</i> <i>Aspergillus niger</i> <i>Aspergillus ustus</i> <i>Aspergillus sydowii</i>
Ochratoxin A	<i>Penicillium chrysogenum</i> <i>Penicillium fellutanum</i> <i>Penicillium verrucosum</i>

Example 16

Combination Method for Mycotoxins and DNA Determination in Body Fluids and Tissue of Humans and Animals

[0219] Presence of *Stachybotrys chartarum* and *Aspergillus niger*—

[0220] The presence of mycotoxin and fungi was determined in a 57 year old Caucasian male with a history of indoor air exposure to *Stachybotrys*, *Aspergillus*, *Chaetomium*, *Penicillium*, and *Fusarium*. An initial skin biopsy showed Aflatoxin presence at 4 ppb. Real Time PCR was performed using the method described in Example 15 for

selection of probes. The results of RT-PCR of skin showed the presence of *Stachybotrys chartarum* and *Aspergillus niger*.

Presence of Mycotoxin—

[0221] The presence of mycotoxin was determined in a 60 year old Caucasian male with a history of exposure to molds in South America and who had developed severe headaches. An MRI of the brain showed a central lesion in or around the Pons. The Cerebral Spinal fluid (CSF) showed Aflatoxin levels present at 4 ppb. RT-PCR analysis was conducted on the CSF for the following organisms: *Aspergillus flavus*, *Aspergillus fumigatus*, *Aspergillus niger*, *Aspergillus ochraceus*, *Aspergillus parasiticus*, and *Aspergillus versicolor*. All results were negative.

[0222] The patient had the antifungal agent (voriconazole) administered until medical caregivers suggested the lesion seen on MRI could be an astrocytoma. The patient received chemotherapy, radiation, and steroids. The patient succumbed to the lesion within two weeks after discontinuing the voriconazole.

[0223] A brain autopsy showed a gliosis of the area near the pons of the patient. No fungal elements were seen on special stains. RT-PCR analysis of the brain biopsy using the method described in Example 15 for *Penicillium* and *Aspergillus* revealed the presence of *Penicillium verrucosum*.

Presence of *Stachybotrys echinata*, *Penicillium chrysogenum* and *Aspergillus ustus*—

[0224] The presence of mycotoxin and fungi was determined in a 47 year old Caucasian male with a long history of skin disorders diagnosed as psoriasis. A skin biopsy evaluation for mycotoxins showed Tricothecenes present at 1.36 ppb, and Aflatoxins, not present. Fungal culture on the skin lesion grew *Aspergillus fumigatus* and *Penicillium* sp. RT-PCR analysis of the skin lesion was conducted using the method described in Example 15. The results for the skin lesion showed the presence of *Stachybotrys echinata*, *Penicillium chrysogenum* and *Aspergillus ustus*.

Presence of *Aspergillus niger* and *Eurotium amstelodami*—

[0225] An autopsy was performed on a 36 year old female with a history of high exposure to environmental mold in the home. The mycotoxin evaluation of a mixture of tissues showed the presence of Aflatoxin at 1.0 ppb. RT-PCR products of the tissues were probed using the method described in Example 15. Results were positive for *Aspergillus niger* and *Eurotium amstelodami* in the brain and *Aspergillus niger* in the lung.

Presence of *Aspergillus niger*, *Stachybotryous chartarum*, and *Penicillium verrucosum*—

[0226] Studies were performed on a five year old Yorkshire Terrier that died of exposure to high levels of indoor mold. Necropsy results showed cardiopulmonary arrest. No molds were seen on special stains of tissues. Mycotoxin studies performed using liver, lung and brain tissues showed tricothecenes of 1.3 ppb and aflatoxins of 2.0 ppb. Results were positive for *Aspergillus niger*, *Stachybotryous chartarum*, and *Penicillium verrucosum*.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 91

<210> SEQ ID NO 1

<211> LENGTH: 22

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 1

ttgcttcggc gggaaacgcc cg

22

<210> SEQ ID NO 2

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 2

gcgaggggat cattaccgag

20

<210> SEQ ID NO 3

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

-continued

<400> SEQUENCE: 3
atcgatgcca gagccaagag 20

<210> SEQ ID NO 4
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
probe

<400> SEQUENCE: 4
cggggagccc tctcgggggc 20

<210> SEQ ID NO 5
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 5
cgtaggtgaa cctgcggaag 20

<210> SEQ ID NO 6
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 6
atcgatgccg gaaccaagag 20

<210> SEQ ID NO 7
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
probe

<400> SEQUENCE: 7
tgtctattgt acctgttgct tc 22

<210> SEQ ID NO 8
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 8
cgtaggtgaa cctgcggaag 20

<210> SEQ ID NO 9
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic

-continued

primer

<400> SEQUENCE: 9

atcgatgccg gaaccaagag 20

<210> SEQ ID NO 10
<211> LENGTH: 26
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 10

ctctgtctga agattgtagt ctgagt 26

<210> SEQ ID NO 11
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 11

cgtaggtgaa cctgcggaag 20

<210> SEQ ID NO 12
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 12

atcgatgccg gaaccaagag 20

<210> SEQ ID NO 13
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 13

cccgcctttg ctggcgcgc 19

<210> SEQ ID NO 14
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 14

cgtaggtgaa cctgcggaag 20

<210> SEQ ID NO 15
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence

-continued

<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 15

atcgatgccg gaaccaagag 20

<210> SEQ ID NO 16
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 16

ggatctcttg gttctcgtat c 21

<210> SEQ ID NO 17
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 17

cttgatctga ggttgaatag tg 22

<210> SEQ ID NO 18
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 18

aacgcacatt gcactttggg gtatc 25

<210> SEQ ID NO 19
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 19

cccgcattc atggccgccg gg 22

<210> SEQ ID NO 20
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 20

cgtaggtgaa cctgcggaag 20

<210> SEQ ID NO 21
<211> LENGTH: 20

-continued

<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 21
atcgatgccg gaaccaagag 20

<210> SEQ ID NO 22
<211> LENGTH: 26
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 22
aaagtatgca gtctgagttg attatc 26

<210> SEQ ID NO 23
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 23
cgtaggtgaa cctgcggaag 20

<210> SEQ ID NO 24
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 24
atcgatgccg gaaccaagag 20

<210> SEQ ID NO 25
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 25
cccagggggc gagccgcccg 20

<210> SEQ ID NO 26
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 26
cgtaggtgaa cctgcggaag 20

-continued

<210> SEQ ID NO 27
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 27

atcgatgccg gaaccaagag 20

<210> SEQ ID NO 28
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 28

acaccaacg tgaacactgt ctgaag 25

<210> SEQ ID NO 29
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 29

cgtaggtgaa cctgcggaag 20

<210> SEQ ID NO 30
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 30

atcgatgccg gaaccaagag 20

<210> SEQ ID NO 31
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 31

cgggcccgcc gtcatggcgg 20

<210> SEQ ID NO 32
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 32

cgtaggtgaa cctgcggaag 20

-continued

<210> SEQ ID NO 33
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 33

atc gat gcc g gaaccaagag 20

<210> SEQ ID NO 34
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
probe

<400> SEQUENCE: 34

ccctcggggg cgagccgccc 20

<210> SEQ ID NO 35
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 35

cgtaggtgaa cctgcggaag 20

<210> SEQ ID NO 36
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 36

atc gat gcc g gaaccaagag 20

<210> SEQ ID NO 37
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
probe

<400> SEQUENCE: 37

ccacaccgaa cctcttgta tagc 24

<210> SEQ ID NO 38
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 38

-continued

cgtaggtgaa cctgcggaag 20

<210> SEQ ID NO 39
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 39

atcgatgccg gaaccaagag 20

<210> SEQ ID NO 40
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 40

cgggaataga cggccccgtg aaac 24

<210> SEQ ID NO 41
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 41

gcggagggat cattaccgag 20

<210> SEQ ID NO 42
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 42

atcgatgcca gagccaagag 20

<210> SEQ ID NO 43
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 43

cccgccttta ctggccgccc g 21

<210> SEQ ID NO 44
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

-continued

<400> SEQUENCE: 44
cgtaggtgaa cctgcggaag 20

<210> SEQ ID NO 45
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 45
atcgatgccg gaaccaagag 20

<210> SEQ ID NO 46
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
probe

<400> SEQUENCE: 46
tgttgcctcg gggggccccg c 21

<210> SEQ ID NO 47
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 47
ggaaggatca ttaccgagtg 20

<210> SEQ ID NO 48
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 48
atcgatgccg gaaccaagag 20

<210> SEQ ID NO 49
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
probe

<400> SEQUENCE: 49
ttattgtacc ttgttgcttc ggcgg 25

<210> SEQ ID NO 50
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:

-continued

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 50

cgtaggtgaa cctgcggaag 20

<210> SEQ ID NO 51
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 51

atcgatgccg gaaccaagag 20

<210> SEQ ID NO 52
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 52

cgatctccgg gggacgggcc 20

<210> SEQ ID NO 53
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 53

ctgtccgagc gtcattgctg 20

<210> SEQ ID NO 54
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 54

cgaggaccgg acgcggtg 18

<210> SEQ ID NO 55
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 55

agacaccccc gaactctgcc tgaag 24

<210> SEQ ID NO 56
<211> LENGTH: 20
<212> TYPE: DNA

-continued

<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 56
cgtaggtgaa cctgcggaag 20

<210> SEQ ID NO 57
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 57
atcgatgccg gaaccaagag 20

<210> SEQ ID NO 58
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 58
cccgcctgcc aggccgccc 19

<210> SEQ ID NO 59
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 59
cgtaggtgaa cctgcggaag 20

<210> SEQ ID NO 60
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 60
atcgatgccg gaaccaagag 20

<210> SEQ ID NO 61
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 61
caccctgtt tatttacctt attgc 25

<210> SEQ ID NO 62

-continued

<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 62

cgtaggtgaa cctgcggaag 20

<210> SEQ ID NO 63
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 63

atcgatgccg gaaccaagag 20

<210> SEQ ID NO 64
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 64

caccctgtgt tatcgtaect tgttg 25

<210> SEQ ID NO 65
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 65

cgtaggtgaa cctgcggaag 20

<210> SEQ ID NO 66
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 66

atcgatgccg gaaccaagag 20

<210> SEQ ID NO 67
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 67

ttgcttcggc gggagagccc cg 22

-continued

<210> SEQ ID NO 68
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 68

gcggagggat cattaccgag 20

<210> SEQ ID NO 69
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 69

atcgatgcc a gagccaagag 20

<210> SEQ ID NO 70
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
probe

<400> SEQUENCE: 70

tgtctatctg taccctgttg ctctg 25

<210> SEQ ID NO 71
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 71

cgtaggtgaa cctgcggaag 20

<210> SEQ ID NO 72
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 72

atcgatgccg gaaccaagag 20

<210> SEQ ID NO 73
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
probe

<400> SEQUENCE: 73

-continued

cctcggatca ggtaggata c 21

<210> SEQ ID NO 74
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 74

atgcctgtcc gacgctcatt 20

<210> SEQ ID NO 75
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 75

ttcctccgct tattgatatg 20

<210> SEQ ID NO 76
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 76

acggatctct tggctctggc atc 23

<210> SEQ ID NO 77
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 77

gcggagggat cattaccgag 20

<210> SEQ ID NO 78
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 78

ttcactgaat tctgcaattc ac 22

<210> SEQ ID NO 79
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

-continued

<400> SEQUENCE: 79

cctctgcccc ccgggcccgt g 21

<210> SEQ ID NO 80

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 80

ggaaggatca ttaccgagtg 20

<210> SEQ ID NO 81

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 81

ggagcccccc agggggcgag 20

<210> SEQ ID NO 82

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 82

cggggaaccc cctcgggggc 20

<210> SEQ ID NO 83

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 83

tgcgctcccc ccgggggcag 20

<210> SEQ ID NO 84

<211> LENGTH: 21

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 84

ggccccgtcc ccgatctcc g 21

<210> SEQ ID NO 85

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic

-continued

primer

<400> SEQUENCE: 85

agtgaatcat cgagtctttg aac 23

<210> SEQ ID NO 86
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 86

acctgatccg aggtcaacct g 21

<210> SEQ ID NO 87
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 87

cgggcccgcc tgccaggccg 20

<210> SEQ ID NO 88
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 88

ccgggggggtt tacaccccg 20

<210> SEQ ID NO 89
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 89

ccggggggca tctgcccgcg g 21

<210> SEQ ID NO 90
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic oligonucleotide

<400> SEQUENCE: 90

gcccccgaga gggtccccg 20

<210> SEQ ID NO 91
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence

-continued

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic oligonucleotide

<400> SEQUENCE: 91

gtttcacggg gccgtctatt cccg

24

What is claimed is:

1. A method of identifying a fungal species in a patient tissue or body fluid, said method comprising the steps of: identifying a mycotoxin in a patient tissue or body fluid; and specifically identifying a fungal species in the mycotoxin positive patient tissue or body fluid.
2. The method of claim 1, further comprising the steps of: extracting and recovering the mycotoxin from the patient tissue or body fluid; and contacting the mycotoxin with an antibody directed against the mycotoxin.
3. The method of claim 1, further comprising the steps of: extracting and recovering DNA of the fungal species from the patient tissue or body fluid; amplifying the DNA; and hybridizing a probe to the DNA to specifically identify the fungal species.
4. The method of claim 3 wherein the amplifying step is performed with primers that hybridize to the DNA.
5. The method of claim 1 wherein the body fluids are selected from the group consisting of urine, nasal secretions, nasal washes, bronchial lavages, bronchial washes, spinal fluid, sputum, gastric secretions, seminal fluid, other reproductive tract secretions, lymph fluid, whole blood, serum, and plasma.
6. The method of claim 3 wherein the DNA is amplified using PCR.
7. The method of claim 6 wherein the PCR is real-time PCR.
8. The method of claim 3 wherein the probe is fluorescently labeled.
9. The method of claim 7 wherein the probe is fluorescently labeled.
10. The method of claim 4 wherein the amplified sequence is internal transcribed spacer regions of nuclear ribosomal DNA.
11. The method of claim 3 wherein the probe is bound to a bead dyed with a fluorochrome.

12. The method of claim 2 further comprising the step of quantifying the mycotoxin.

13. The method of claim 1 wherein the tissue is derived from a patient tissue biopsy and is in a 10% formalin solution or is in a paraffin block.

14. The method of claim 2 wherein the antibody is bound to a bead dyed with a fluorochrome.

15. A kit comprising components for the extraction and recovery of a mycotoxin from body fluid or tissue of a patient; components for identification of the mycotoxin; and a purified nucleic acid probe and primer.

16. The kit of claim 15 wherein the components for identification of the mycotoxin include beads dyed with a fluorochrome and coupled to antibodies to the mycotoxin or to the mycotoxin or to a mycotoxin antigen.

17. The kit of claim 15 wherein the components for identification of the mycotoxin comprise an antibody directed against the mycotoxin.

18. A method of identifying a fungal species in a patient tissue or body fluid, said method comprising the steps of:

testing for the presence of a mycotoxin in a patient tissue or body fluid;
extracting and recovering DNA of a fungal species from the patient tissue or body fluid;
amplifying the DNA; and
hybridizing a probe to the DNA to specifically identify the fungal species.

19. A method of identifying a mycotoxin in a patient tissue or body fluid, said method comprising the steps of:


extracting and recovering the mycotoxin from the patient tissue or body fluid;
contacting the mycotoxin with an antibody directed against the mycotoxin;
extracting and recovering patient DNA from the patient tissue or body fluid;
amplifying the DNA; and
hybridizing a probe to the patient DNA.

* * * * *

专利名称(译)	用于鉴定霉菌毒素和真菌物种的方法和组合物		
公开(公告)号	US20100075322A1	公开(公告)日	2010-03-25
申请号	US12/545734	申请日	2009-08-21
[标]申请(专利权)人(译)	HOOPER DENNIS 摹		
申请(专利权)人(译)	HOOPER DENNIS 摹		
[标]发明人	HOOPER DENNIS G		
发明人	HOOPER, DENNIS G.		
IPC分类号	C12Q1/68 G01N33/566 G01N33/53 G01N33/48		
CPC分类号	C12Q1/6883 G01N33/5308 C12Q1/6895		
优先权	61/091185 2008-08-22 US		
外部链接	Espacenet USPTO		

摘要(译)

本发明涉及一种在患者组织或体液中测试霉菌毒素并测试患者组织或体液中的真菌物种的方法。该方法还可用于确定患者是否处于与真菌感染相关的疾病状态或已经发展成与真菌感染相关的疾病状态，并为患者开发有效的治疗方案。



Tricothecenes Value (ppb)	Number of Tests
0 - 0.2	5
0.21 - 1.0	17
1.1 - 2.0	11
2.1 - 5.0	27
5.1 - 10	19
10.1 - 35	25
> 35*	5
Total:	109