



US00763557B2

(12) **United States Patent**
Arad

(10) **Patent No.:** **US 7,635,557 B2**
(45) **Date of Patent:** **Dec. 22, 2009**

(54) **ENZYMATIC DIAGNOSTIC TEST FOR SARS AND OTHER VIRAL DISEASES**

(75) Inventor: **Dorit Arad**, Tel Aviv (IL)

(73) Assignee: **MND Diagnostic Ltd.**, Nes Ziona (IL)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 242 days.

(21) Appl. No.: **10/875,133**

(22) Filed: **Jun. 23, 2004**

(65) **Prior Publication Data**

US 2005/0048473 A1 Mar. 3, 2005

Related U.S. Application Data

(60) Provisional application No. 60/480,605, filed on Jun. 23, 2003.

(51) **Int. Cl.**

C12Q 1/70 (2006.01)
C12Q 1/00 (2006.01)
C12Q 1/37 (2006.01)
G01N 33/53 (2006.01)
C12N 15/41 (2006.01)

(52) **U.S. Cl.** **435/5; 435/4; 435/7.72; 435/23; 435/24**

(58) **Field of Classification Search** None
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

4,748,116 A 5/1988 Simonsson

4,952,493 A * 8/1990 Kettner et al. 435/5
5,591,591 A 1/1997 Bronstein
6,037,137 A 3/2000 Komoriya
6,127,139 A 10/2000 Te Kokkele
6,243,980 B1 6/2001 Bronstein
6,287,767 B1 9/2001 Bronstein

FOREIGN PATENT DOCUMENTS

WO WO 2005/017191 2/2005

OTHER PUBLICATIONS

Chiu RW et al, "Automated extraction protocol for quantification of SARS-Coronavirus RNA in serum: an evaluation study" *BMC Infectious Diseases* 2006, 6:20-26.*

Loens K, et al. "Detection of Rhinoviruses by Tissue Culture and Two Independent Amplification Techniques, Nucleic Acid Sequence-Based Amplification and Reverse Transcription-PCR, in Children with Acute Respiratory Infections during a Winter Season" *J Clin Microbiol.* Jan. 2006;44(1):166-71.*

Cordingley, et al. Cleavage of Small Peptides in Vitro by Human Rhinovirus 14 3C Protease Expressed in *Escherichia coli*. *J. Virol.* 1889; 63(12): 5037-5045.*


(Continued)

Primary Examiner—Mary E Mosher
Assistant Examiner—Stuart W Snyder


(57) **ABSTRACT**

The present invention is directed towards methods, compositions and kits for testing for a virus in a sample. The methods determine the presence of a viral enzyme by contacting the sample with a peptidal compound capable of being cleaved by the viral enzyme to form peptidal compound fragments. Detection of a peptidal compound fragment confirms the presence of the virus.

12 Claims, 9 Drawing Sheets



Quenched tag




Signal Emitted

OTHER PUBLICATIONS


- Hata, et al. A simple purification and fluorescent assay method of the poliovirus 3C protease searching for specific inhibitors. *Journal of Virological Methods* 84 (2000) 117-126.*
- Manafi, et al. Fluorogenic and Chromogenic Substrates Used in Bacterial Diagnostics. *Microbiological Reviews*, Sep. 1991, vol. 55, No. 3: p. 335-348.*
- Wang et al. "Development of A Continuous Fluorescence Assay for Rhinovirus 14 3C Protease Using Synthetic Peptides", *Antiviral Chemistry & Chemotherapy*, 8(4): 303-310, 1997. p. 303, Abstract, p. 304, l-h Col., p. 306, r-h Col., Paragraph 2, p. 308, r-h Col., Paragraph 3.
- Wang et al. "Development of in Vitro Peptide Substrates for Human Rhinovirus-14 2A Protease", *Archives of Biochemistry and Biophysics*, 356(1): 12-18, 1998. p. 12, Abstract, p. 13, r-h Col., Paragraph 3, p. 15, r-h Col., Paragraph 2.
- Makinen, K. et al., *J. Gen. Virol.* vol. 81, pp. 2783-2789 (2000).
- Khan, A. R. et al. "Structural aspects of activation pathways of aspartic protease zymogens and viral 3C protease precursors", *Proc. Nat'l. Acad. Sci.*, vol. 96, No. 20, pp. 10968-10975 (1999).
- Rao, M. B. et al., "Molecular and Biotechnological Aspects of Microbial Proteases", *Microbiol. and Mol. Biology Rev.*, vol. 62, No. 3, p. 597-635 (1998).
- Corbalanya, A.E. and Snijder, E.J., "Viral cysteine proteinases, Perspectives in Drug Discovery and Design" 6: pp. 64-86 (1996).
- Dougherby, W.G. and Semler, B.L., "Expression of virus-encoded proteinases: functional and structural similarities with cellular enzymes", *Microbiol. Rev.* 57, pp. 781-822 (1993).
- John Ziebuhr et al. "Virus-encoded proteinases and proteolytic processing in the Nidovirales", *J. Gen. Vir.*, vol. 81, pp. 853-879 (2000).
- Anand K. et al., *Coronavirus Main Protease (3CL^{pro}) Structure: Basis of Design of Anti-SARS Drugs* Science, vol. 30, pp. 1763-1767 (2003).
- Sosnovtsev, S. V., et al., "Cleavage of the feline calicivirus capsid precursor is mediated by a virus-encoded proteinase", *J. Virol.*, vol. 72, pp. 3051-3059 (1998).
- Wang, W-K "Detection of SARS-associated Coronavirus in Throat Wash and Saliva in Early Diagnosis" *Emerg Infect Dis* [serial on the Internet] (Jun. 2004). Available from: <http://www.cdc.gov/ncidod/EID/vol10no7/03-1113.htm>.
- Wang Q.M. et al., "A continuous colorimetric assay for rhinovirus-14 3C protease using peptide p-nitroanilides as substrates" *Anal. Biochem.* vol. 252, pp. 238-245 (1997).
- Basak S. et al. "In vitro elucidation of substrate specificity and bioassay of proprotein convertase 4 using intramolecularly quenched fluorogenic peptides" *Biochem. J.* vol. 380, pp. 505-514 (2004).
- Orr D.C. et al. "Hydrolysis of a series of synthetic peptide substrates by the human rhinovirus 143C proteinase, cloned and expressed in *E. coli*", *J. Gen. Vir.* vol. 70, pp. 2931-2942 (1989).
- Rota, Paul A et al. "Characterization of a Novel Coronavirus Associated with Severe Acute Respiratory Syndrome" *Science*, vol. 300, pp. 1394-1399 (May 30, 2003).
- Marra, Marco A. et al., "The Genome Sequence of the SARS-Associated Coronavirus" *Science*, vol. 300, pp. 1399-1404 (May 30, 2003).
- Ksiazek, Thomas G. et al., "A Novel Coronavirus Associated with Severe Respiratory Syndrome" *New England Journal of Medicine* 348: 20, pp. 1953-1966 (Dec. 2, 2003).
- International Search Report for PCT/US2004/019897.

* cited by examiner

Figure 1



Quenched tag



Signal Emitted

	position Cleavage		Cleavage Position	Virus
	↓		Vp4/vp2	
SEQ ID NO:1	lmlkgapaln	spnveacgys	69	HRV14
SEQ ID NO:2	vlekgiptlq	spsveacgys	69	HRV16
SEQ ID NO:3	lmlkgapaln	spnveacgys	69	HRVb
SEQ ID NO:4	lmlktapaln	spnveacgys	69	Cxa21

Sequence Alignment:

```

HRVb          LMLKGAPALNSPNVEACGYS 20
Cxa21         LMLKTAPALNSPNVEACGYS 20
HRV14         LMLKGAPALNSPNVEACGYS 20
HRV16         VLEKGIPTLQSPSVEACGYS 20
              :: * *:*:**.*****
Profile       33-K-3P-L2SP2VEACGYS
    
```

			Vp2/vp3	
SEQ ID NO: 5	girksivpq	glptttlpgs	331	HRV14
SEQ ID NO: 6	sgaraktvvq	glpvvytpgs	330	HRV16
SEQ ID NO: 7	girksivpq	glptttlpgs	331	HRVb
SEQ ID NO: 8	lrnitvpvhq	glptmntpgs	341	Cxa21

```

HRV16         SGARAKTVV-QGLPVYVTPGS 20
Cxa21         LRNITVPVH-QGLPTMNTPGS 20
HRV14         -GIRSKSIVPQGLPTTTLPGS 20
HRVb          -GIRSKSIVPQGLPTTTLPGS 20
              : .: ****. ***
Profile       -----3-QGLP-----PGS
    
```

			Vp3/vp1	
SEQ ID NO: 9	fkrlmkdtq	tisqtvalte	557	HRV14
SEQ ID NO: 10	lhkqtgpitq	npveryvdev	568	HRV16
SEQ ID NO: 11	tisqtvalte	glgdeleevi	567	HRVb
SEQ ID NO: 12	sqskligrtq	giedlidtai	581	Cxa21

```

HRV14         FKLRLMKDTQTISQTVALTE----- 20
HRVb          -----TISQTVALTEGLGDELEEVI 20
Cxa21         -----SQSKLIGRTQGIEDLIDTAI 20
HRV16         -----LHKQTGPITQNP-VERYVDEV- 20
              . . . : : .
Profile       -----2--3--2-----
    
```

			Vp1/2A	
SEQ ID NO: 13	kkrkgdiksy	glgpryggii	856	HRV14
SEQ ID NO: 14	irprtnlttv	gpsdmyvhvg	853	HRV16
SEQ ID NO: 15	kkrkgdiksy	glgpryggii	856	HRVb
SEQ ID NO: 16	ltkvdsittf	gfghqnkavy	879	Cxa21

```

HRV14         KKRKGDIKSYGLGPRYGGIY 20
HRVb          KKRKGDIKSYGLGPRYGGIY 20
Cxa21         LTKVDSITTFGFGHQNKAVY 20
HRV16         IRPRTNLTTVGPSDMYVHVG 20
              .:: * . :
Profile       -----3-2-G-----3-
    
```

Figure 2 (A)

	position Cleavage		Cleavage Position	Virus
			2A/2B	
SEQ ID NO: 17	rqleciaeeq	glsdyitglg	1002	HRV14
SEQ ID NO: 18	lrhfhcaeeq	gitdyihmlg	995	HRV16
SEQ ID NO: 19	rqleciaeeq	glsdyitglg	1002	HRVb
SEQ ID NO: 20	wyeeeeameq	gitsyieslg	1028	Cxa21

Sequence Alignment:

```

HRV16      LRRHFCAEEQGITDYIHMLG 20
Cxa21      WVYEEEAMEQGITSYIESLG 20
HRV14      RQLECIAEEQGLSDYITGLG 20
HRVb      RQLECIAEEQGLSDYITGLG 20
           * **:::** ** **
Profile    -----A-EQG32-YI--LG
    
```

			2B/2C	
SEQ ID NO: 21	hfqvpwierq	andgwrkfn	1099	HRV14
SEQ ID NO: 22	wtqltyihke	sdswlkkfte	1090	HRV16
SEQ ID NO: 23	hfqvpwierq	andgwrkfn	1099	HRVb
SEQ ID NO: 24	lleipyvmrq	gdgwmkkfte	1125	Cxa21

```

HRV14      HFQVPYIERQANDGWFRKFN- 20
HRVb      HFQVPYIERQANDGWFRKFN- 20
Cxa21      LLEIPYVMRQG-DGWMKKFTE 20
HRV16      WTQLTYIHKES-DSWLKKFTE 20
           ::*: :: *.*:***.
Profile    ---3-Y3-5---D-W-5KF2-
    
```

			2C/3A	
SEQ ID NO: 25	itdsletlfq	gpvykdleid	1429	HRV14
SEQ ID NO: 26	vvdvmsaifq	gpismdkppp	1412	HRV16
SEQ ID NO: 27	itdsletlfq	gpvykdleid	1429	HRVb
SEQ ID NO: 28	igncmealfq	gplrykdlki	1453	Cxa21

```

HRV14      ITDSLETLFQGPV-YKDLEID 20
HRVb      ITDSLETLFQGPV-YKDLEID 20
Cxa21      IGNCMEALFQGPLYKDLKI- 20
HRV16      VVDVMSAIFQGPISMDKPPP- 20
           : : ::*:***: .. .
Profile    3---3--3FQGP3----3---
    
```

Figure 2(B)

	position Cleavage		Cleavage Position	Virus
SEQ ID NO: 29	viyklfaqtq	gpysgnpphn	1514	HRV14
SEQ ID NO: 30	iiyklfcsiq	gpysgepkpk	1489	HRV16
SEQ ID NO: 31	viyklfaqtq	gpysgnpphn	1514	HRVb
SEQ ID NO: 32	vmyklfagqq	gaytglpnkk	1541	Cxa21

Sequence Alignment:

```

HRV14      VIYKLFAQTQGPYSGNPPHN 20
HRVb      VIYKLFAQTQGPYSGNPPHN 20
HRV16     IIYKLFCSLQGPYSGEKPKK 20
Cxa21     VMYKLFAGQQGAYTGLPNKK 20
           ::****.  **.*:* *  :
Profile   33YKLF---QG3Y2G-P---
    
```

SEQ ID NO: 33	aptlrpvvvq	gpntefalsl	1537	HRV14
SEQ ID NO: 34	kvperrvvaq	gpeeefgmsi	1510	HRV16
SEQ ID NO: 35	aptlrpvvvq	gpntefalsl	1537	HRVb
SEQ ID NO: 36	vptirvakvq	gpgfdyavam	1563	Cxa21

```

HRV14      -APTLRPVVVQGPNTEFALS 20
HRVb      -APTLRPVVVQGPNTEFALS 20
HRV16     KVPERR-VVAQGPEEEFGMSI 20
Cxa21     -VPTIRVAKVQGPFGDYAVAM 20
           . *  *  . .***  :::::
Profile   -3P--RP3-3QGP--4--3-3
    
```

SEQ ID NO: 37	lkkqyfveq	gqviahkvr	1719	HRV14
SEQ ID NO: 38	llrsyfteq	gqiqiskhvk	1693	HRV16
SEQ ID NO: 39	lkkqyfveq	gqviahkvr	1719	HRVb
SEQ ID NO: 40	lkrsyftq	geiqwmrsk	1746	Cxa21

```

HRV16     LLRSYFTEQQGQIQISKHVK 20
Cxa21     LKRSYFTQNQGEIQWMRSSK 20
HRV14     LKKQYFVEKQGQVIARHKVR 20
HRVb     LKKQYFVEKQGQVIARHKVR 20
           *  :.*.::***:  :  :
Profile   L-52YF---QG-3-----5
    
```

profile key :

Group	Symbol	Amino acids
Aromatic	1	F Y W H
H-bonds	2	Q N S T C H
Hydrophobic	3	A L I V P C M
Acid	4	D E
alkaly	5	K R Q N H

Figure 2(C)

HRV14_2C_3A	-ITDSLETLFQ	GPV-YKDLEID-	20
HRVb_2C_3A	-ITDSLETLFQ	GPV-YKDLEID-	20
Cxa21_2C_3A	-IGNCMEALFQ	GPLRYKDLKI--	20
HRV16_2C_3A	-VVDVMSAIFQ	GPISMDKPPP--	20
HRV14_3A_3B	-VIYKLFAQTQ	GPYS-GNPPHN-	20
HRVb_3A_3B	-VIYKLFAQTQ	GPYS-GNPPHN-	20
Cxa21_3A_3B	-VMYKLFAGQQ	GAYT-GLPNKK-	20
HRV16_3A_3B	-IIYKLFCSLQ	GPYS-GEPKPK-	20
HRV14_3B_3C	-APTLRPVVVQ	GPN--TEFALSL	20
HRVb_3B_3C	-APTLRPVVVQ	GPN--TEFALSL	20
Cxa21_3B_3C	-VPTIRVAKVQ	GPG--FDYAVAM	20
HRV16_3B_3C	KVPERR-VVAQ	GPE--EEFGMSI	20
	*	*	

Group %	*	V	3			3		3		3	Q	G	P				4		
		50	75			75		75		75	100	100	92				67		
		I											A						
		30											8						
		A																	
		20																	
consensus		3				3		3		3	Q	G	P				4		

Figure 2 (D)

			Cleavage Position	Virus
SEQ ID NO: 41	reltrelnngg	avtryvdnnf	180	SARS
SEQ ID NO: 42	mskinkygle	vkpllyvdqy	246	BCov
SEQ ID NO: 43	dvvfgkrggg	nvtytdqylc	111	HCov

Sequence Alignment:

```

BCOV      MSKINKYGLEVKP-LLYVDQY-- 20
HCOV      --DV-VFGKRGGGNVTYTDQYLC 20
SARS      -REL-TRELNGGAVTRYVDNNF- 20
          .:      .      *.*:
Profile   ---3-----Y-D2---
    
```

SEQ ID NO: 44	tnnvfrlkgg	apikgvtfge	818	SARS
SEQ ID NO: 45	ldqawrvpca	grrvtfkeqp	851	BCov
SEQ ID NO: 46	lpvaftkaag	gkvsfsddve	897	HCov

```

BCOV      LDQAWRVPCAG--RRVTFKEQP- 20
SARS      TNNVFRKGGAPIKGVTFGE--- 20
HCOV      LPVAFTKAAGG---KVSFSDDVE 20
          .:      ...      *.*:
Profile   ---31-----V2F-4---
    
```

SEQ ID NO: 47	ittkislkkg	kivstcfklm	2740	SARS
SEQ ID NO: 48	lttpfslkkg	avfsyfvvvc	2750	BCov
SEQ ID NO: 49	atsivakqga	gdaghsltwl	2484	HCov

```

SARS      ITTKISLKGKIVSTCFKLM 20
BCOV      LTPFSLKGGAVFSYFVVC 20
HCOV      ATSIVAKQGAGDAGHSLTTL 20
          *: .: .*: . .
Profile   3T2----5G-----3
    
```

				4-5
SEQ ID NO: 50	qtsitsavlq	sgfrkmafps	3240	SARS
SEQ ID NO: 51	tasvstsflq	sgivkmvnpt	3246	BCov
SEQ ID NO: 52	ptvsygstlq	aglrkmaqps	2965	HCov

```

SARS      QTSITSAVLQSGFRKMAFPS 20
HCOV      PTVSYGSTLQAGLRKMAQPS 20
BCOV      TASVSTSFLQSGIVKMNPT 20
          :      : **:*: **. *:
Profile   -----LQ-G--KM3-P2
    
```

Figure 3(A)

			Cleavage Position 3CL	Virus
SEQ ID NO: 53	vrqcsqvtfq	gfkfkivkgt	3546	SARS
SEQ ID NO: 54	yqqlagiklq	skrtrlvkgi	3549	BCov
SEQ ID NO: 55	vkqmfqvnlg	sgkttsmfks	3267	HCov

Sequence Alignment:

SARS VRQCSGVTFQ-GKFKKIVKGT 20
 BCOV YQQLAGIKLQ-SKRTRLVKGI 20
 HCOV VKQMFQVNLQSGKTTSMFKS- 20
 :* *::* .* .:*.
 Profile -5Q3-G3---Q-K---3-K--

SEQ ID NO: 56	kpcikvatvq	skmsdvkcts	3836	SARS
SEQ ID NO: 57	vpievsqfq	skltdvkcan	3836	BCov
SEQ ID NO: 58	prtikvstvq	skltdlkctn	3546	HCov

BCOV VP-IIIEVSQFQSKLTDVKCAN 20
 HCOV -PRTIKVSTVQSKLTDLKCTN 20
 SARS KP-CIKVATVQSKMSDVKCTS 20
 * **:: .***::**::.
 Profile -P--I-V-2-QSK32D3KC-2

SEQ ID NO: 59	emldnratlq	aiasefsslp	3919	SARS
SEQ ID NO: 60	dyakdntvlq	alqsefvnma	3925	BCov
SEQ ID NO: 61	syfendsilq	svassfvgmp	3629	HCov

BCOV DYAKDNTVLQALQSEFVNMA 20
 HCOV SYFENDSILQSVASSFVGMP 20
 SARS EMLDNRATLQAIASEFSSLP 20
 . .: : **:: *.* .:..
 Profile -----LQ-3-S-F--33

SEQ ID NO: 62	lransavklq	nnelspvalr	4117	SARS
SEQ ID NO: 63	hnevsatvlq	nnelmpaklk	4122	BCov
SEQ ID NO: 64	lrcervvklq	nneimpqkkm	3824	HCov

BCOV HNEVSATVLQNNELMPAKLK 20
 HCOV LTCERVVKLQNNELMPAKLK 20
 SARS LRANSVAVKLQNNELSPVALR 20
 .. *****: * :..
 Profile -----3--LQNE3-P--35

Figure 3(B)

			Cleavage Position	
SEQ ID NO: 65	gslaatvrlq	agnatevpan	4230	SARS
SEQ ID NO: 66	gtisstvrlq	agtateyasn	4232	BCov
SEQ ID NO: 67	gyigatvrlq	agkqtefvsn	3933	HCov

Sequence Alignment:

BCOV GTISSTVRLQAGTATEYASN 20
 HCOV GYIGATVRLQAGKQTEFVSN 20
 SARS GSLAATVRLQAGNATEVPAN 20
 * :.:*****. ** :.*
 Profile G-3--TVRLQAG--TE-3-N

SEQ ID NO: 68	cdqlreplmq	sadastflnr	4369	SARS
SEQ ID NO: 69	scvstdttvq	skdtnflnr	4369	BCov
SEQ ID NO: 70	gctcdrtaiq	sfdnsylnr	4068	HCov

BCOV SC-VSTDTTVQSKDTN-FLNRV 20
 HCOV GC-TCDRTAIQSFDNS-YLNRV 20
 SARS -CDQLREPLMQSADASTFLNR- 20
 * . : ** * . : ***
 Profile -C-----3QS-D-2--LNR-

SEQ ID NO: 71	amyphtvlq	avgacvlcns	5301	SARS
SEQ ID NO: 72	nmylrsavmq	svgacvvcss	5297	BCov
SEQ ID NO: 73	smyekstvlq	aaglcvcgcs	4995	HCov

SARS AMYTPHTVLQAVGACVLCNS 20
 HCOV SMYEKSTVLQAAGLCVVCSS 20
 BCOV NMYLRSAVMQSVGACVVCSS 20
 ** :*:*:*. * **:*.*
 Profile -MY-2-V3Q-3G3CV3C-S

SEQ ID NO: 74	iprrnvatlq	aenvtqlfkd	5902	SARS
SEQ ID NO: 75	vpqavetrvg	cstnlfkdc	5900	BCov
SEQ ID NO: 76	ffeitmtdlq	sesscqlfkd	5592	HCov

SARS IPRRNVATLQAENVTGLFKD-- 20
 HCOV FFEITMTDLQSESSCGLFKD-- 20
 BCOV VPQAVETRVQCS--TNLFKDCS 20
 . . : :*.. .****
 Profile -----3Q-----2-LFKD--

Figure 3(C)

			Cleavage Position	Virus
SEQ ID NO: 77	nlwntftrlq	slenvaynvv	6429	SARS
SEQ ID NO: 78	nlwntftklq	slenvvynlv	6421	BCov
SEQ ID NO: 79	wqtftevnlg	gleniafnvv	6110	HCov

Sequence Alignment:

```

SARS      NLWNTFTR--LQSLENVAYNVV 20
BCOV     NLWNTFTK--LQSLENVVYNLV 20
HCov     --WQTFTEVNLQGLENI AFNVV 20
          . * : * * * .   * * . * * * : : * : *
Profile  --W2TFT---LQ-LEN33-N3V
    
```

SEQ ID NO: 80	hvetfypklq	asqawqpgva	6775	SARS
SEQ ID NO: 81	kvmtfyprlq	aasdwkpgys	6795	BCov
SEQ ID NO: 82	avatfypqlq	saewkcgysm	6458	HCov

```

BCOV     KVMTFYPR LQAASDWKPGYS- 20
HCov     AVATFY PQLQSA-EWKCGYSM 20
SARS     HVETFY PKLQASQAWQPGVA- 20
          * * * * : * * : : * : * :
Profile  -V-TFYP-LQ----W--G----
    
```

profile key :

Group	Symbol	Amino acids
Aromatic	1	F Y W H
H-bonds	2	Q N S T C H
Hydrophobic	3	A L I V P C M
Acid	4	D E
alkaly	5	K R Q N

```

QTSITSAVLQSGFRKMAFPS 20
VRQCSGVTFQ-GKFKKIVKG 20
KPCIKVATVQSKMSDVKCTS 20
EMLDNRATLQAIASEFSSLP 20
LRANSVAVKLQNNELSPVALR 20
GSLAATVRLQAGNATEVPAN 20
CDQLREPLMQSADASTFLNR-
AMYPHTVLQAVGACVLCNS 20
IPRRNVATLQAENV TGLFKD-20
WNTFTR--LQSLENVAYNVV 20
    
```

XXLQA(S)GXX

Figure 3(D)

ENZYMATIC DIAGNOSTIC TEST FOR SARS AND OTHER VIRAL DISEASES

RELATED APPLICATIONS

This application claims priority to U.S. Provisional Patent Application No. 60/480,605, filed Jun. 23, 2003, the entire content of which is fully incorporated herein by reference.

TECHNICAL FIELD OF THE INVENTION

The present invention provides a method, reagents and a kit for the detection of a virus in a test sample. The method involves the detection of an enzyme that is encoded by a viral nucleic acid during the replication of the virus. In one embodiment, the invention provides a method for detection of the Severe Acute Respiratory Syndrome (SARS) virus in a sample.

BACKGROUND

There is a continuing need for rapid sensitive methods for the detection of viral diseases. Rapid detection and identification of viral pathogens is essential to limit transfer and spread of viral disease and to monitor treatment methods. The need for such methods can be illustrated by the events surrounding the emergence of SARS.

SARS is a viral respiratory illness caused by the SARS-associated coronavirus (SARS-CoV). SARS was first reported in Asia in February 2003. Over the next few months, the illness spread to more than two dozen countries in North America, South America, Europe, and Asia before the SARS global outbreak of 2003 was contained. The symptoms of SARS included high fever, dry cough, and other flu like symptoms that can progress to pneumonia. Death occurred in approximately 15% of the infected patients. The contagious nature of SARS and its fast spread in several countries in Asia caused considerable concern throughout the world.

The SARS coronavirus belongs to a group of viruses similar to those causing the common cold. The SARS virus spreads by close person-to-person contact and is thought to be transmitted most readily by respiratory droplets produced when an infected person coughs or sneezes. Droplet spread can occur when droplets from an infected person are propelled a short distance (generally up to 3 feet) through the air and deposited on the mucus membranes of nearby persons. The virus also can spread when a person touches a surface contaminated with infectious droplets and then touches his or her mouth, nose, or eyes.

In most cases, symptoms of SARS first occur 2 to 5 days after exposure to the virus. However, the incubation period may be up to 2 weeks. Because of the extended incubation period, patients infected with the SARS virus may be placed in quarantine with those who are not infected. This may cause the uninfected patients and medical staff to become exposed to the SARS virus and to develop the symptoms of SARS.

Because of its ease of spread and long incubation period, it is critical to reliably determine the presence of the virus in a patient thought to be infected with SARS. Ideally, such a test should be sufficiently sensitive to detect the virus at an early stage of infection. The test should also be specific and have a low occurrence of false-positive and false-negative results.

Without a reliable test that can be used in the early stages of SARS infection, physicians and health care teams rely on a process of elimination, ruling out other known causes of the severe pneumonia before diagnosing SARS. However, a positive test result for another respiratory pathogen does not com-

pletely rule out infection with the SARS virus. Patients can be co-infected with both the SARS virus and other respiratory pathogens.

At present, two types of tests detect the presence of the SARS virus. The first of these is an enzyme immunoassay (EIA) test which detects serum antibody to SARS. The other test is a polymer chain reaction (PCR) test which detects the viral genetic material.

During the course of infection with the SARS virus, levels of specific anti-viral antibodies rise in the blood. Many of the tests currently available for the diagnosis of SARS are based on the detection of such antibodies. Such tests are typically either Enzyme Linked Immunoassays (ELISAs) or immunofluorescence assays (IFAs). With ELISAs, the antibodies cannot be detected until about 20 days after infection. IFAs can detect antibodies approximately 10 days after the initial infection. However, such assays are comparatively slow and require the growth of the virus in a cell culture.

The utility of antibody tests for detecting viruses, such as SARS, may be further limited due to the rapid mutation rate of some viruses. A Canadian study has detected the SARS virus in only 60% of those with SARS infection. Such results suggest that the virus is unstable and is mutating rapidly. This is not unexpected as coronaviruses are notorious for changing their outer surface antigens rapidly, a process termed antigenic drift.

Techniques such as the Polymer Chain Reaction (PCR) allow direct detection of the virus genetic material and, in theory, can detect infection at a very early stage. Many PCR tests use oligonucleotide microchip technology for detecting the virus with throat swabs, sputum or feces. Such tests typically take a few hours to perform and are relatively costly. In addition, currently available PCR methods give 40% of false positive and negative results, making the method ineffective.

Because of the deficiencies of presently available testing methods, there is a need for an improved test enabling the presence of viruses, such as the SARS virus, to be accurately detected at an early stage of infection. Such a test will benefit those showing symptoms of SARS by allowing for the monitoring of the course of their infection and subsequent recovery. In addition, a quick and effective test will benefit persons suspected of having the disease by allowing uninfected persons to be released from quarantine.

There is also the need for an automated test avoiding the need for manual intervention. Such a test will prevent spread of the disease due to infection during the testing process.

SUMMARY

One embodiment of the present invention provides a method for detecting a virus in a sample. The method involves contacting the sample with a peptidal compound capable of being cleaved at a cleavage point by an enzyme to form a first peptidal compound fragment and a second peptidal compound fragment. The enzyme is encoded by a nucleic acid of the virus during replication of the virus and cleaves a polyprotein encoded by the nucleic acid of the virus. A signaling moiety is linked to a portion of the peptidal compound that forms the first peptidal compound fragment. If the virus is present in the sample, the enzyme produced by the virus cleaves the peptidal compound. The virus is detected by observing the signal from the signaling moiety.

In another embodiment, a quenching moiety is linked to a portion of the peptidal compound present in the second peptidal compound fragment. The signaling moiety and the quenching moiety are linked to the peptidal compound at relative positions such that the quenching moiety quenches

the signal of the signaling moiety unless the peptidal compound is cleaved at the cleavage point.

In other embodiments, the virus detected is from either the Nidovirus or the Picornavirus virus families. In one embodiment the virus is the SARS virus. In another embodiment the virus is a rhinovirus.

Another aspect of the invention provides a kit for detecting the presence of a virus in a sample. The kit includes a reagent containing a peptidal compound capable of being cleaved at a cleavage point by an enzyme encoded by the viral nucleic acid to form a first peptidal compound fragment and a second peptidal compound fragment. A signaling moiety is linked to a portion of the peptidal compound that forms the first peptidal compound fragment.

Another aspect of the invention provides a composition for detecting a virus in a sample. The composition includes a peptidal compound capable of being cleaved at a cleavage point by an enzyme encoded by the virus to form a first peptidal compound fragment and a second peptidal compound fragment. A signaling moiety is linked to a portion of the peptidal compound that forms the first peptidal compound fragment and a quenching moiety is linked to a portion of the peptidal compound that forms the second peptidal compound fragment. The signaling moiety and the quenching moiety are linked to the peptidal compound at relative positions such that the quenching moiety quenches a signal of the signaling moiety unless the peptidal compound is cleaved at the cleavage point.

BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 is a schematic representation of a homogenous assay for the detection of a virus.

FIG. 2(A) is a Table showing polyprotein cleavage sites for selected species of Picornaviruses. HRV14: Human Rhinovirus 14; HRVb: Human Rhinovirus b; HRV16: Human Rhinovirus 16; Cxa21: Coxsackievirus A21.

FIG. 2(B) is a Table showing polyprotein cleavage sites for selected species of Picornaviruses. HRV14: Human Rhinovirus 14; HRVb: Human Rhinovirus b; HRV16: Human Rhinovirus 16; Cxa21: Coxsackievirus A21.

FIG. 2(C) is a Table showing polyprotein cleavage sites for selected species of Picornaviruses. HRV14: Human Rhinovirus 14; HRVb: Human Rhinovirus b; HRV16: Human Rhinovirus 16; Cxa21: Coxsackievirus A21.

FIG. 2(D) is a Table showing three polyprotein cleavage sites having high hydrolysis rates for selected species of picornaviruses. HRV14: Human Rhinovirus 14; HRVb: Human Rhinovirus b; HRV16: Human Rhinovirus 16; Cxa21: Coxsackievirus A21. A consensus peptide is also shown. SEQ ID NOS. of the cleavage sites are as follows:

Cleavage Site	SEQ ID NO
HRV 2C 3A	25
HRVb 2C 3A	27
Cxa21 2C 3a	28
HRV16 2C 3A	26
HRV14 3A 3B	29
HRVb 3A 3B	31
Cxa21 3A 3B	32
HRV16 3A 3B	30
HRV14 3B 3C	33
HRVb 3B 3C	35
Cxa21 3B 3C	36
HRV16 3B 3C	34

FIG. 3(A) is a Table showing polyprotein cleavage sites for selected species of Nidoviruses. SARS: Human Severe Acute Respiratory Syndrome Virus; BCov: Bovine Coronavirus; HCov: Human Coronavirus.

FIG. 3(B) is a Table showing polyprotein cleavage sites for selected species of Nidoviruses. SARS: Human Severe Acute Respiratory Syndrome Virus; BCov: Bovine Coronavirus; HCov: Human Coronavirus.

FIG. 3(C) is a Table showing polyprotein cleavage sites for selected species of Nidoviruses. SARS: Human Severe Acute Respiratory Syndrome Virus; BCov: Bovine Coronavirus; HCov: Human Coronavirus.

FIG. 3(D) is a Table showing polyprotein cleavage sites for selected species of nidoviruses. SARS: Human Severe Acute Respiratory Syndrome Virus; BCov: Bovine Coronavirus; HCov: Human Coronavirus. SEQ ID NOS. for the 20 residue sequences shown in alignment is as follows:

Cleavage Site	SEQ ID NO
QTSITSAVLQSGFRKMAFPS	50
VRQCSGVTFQ-GKFKKIVKG	53
KPCIKVATVQSKMSDVKCTS	56
EMLDNRATLQAIASEFSSLP	59
LRANSAVKLQNNELSPVALR	62
GSLAATVRLQAGNATEVPAN	65
CDQLREPLMQSADASTFNR	68
AMYPHTVTLQAVGACVLCNS	71
IPRRNVATLQAENVTLGLFKD	74
WNTFTR--LQSLENVAYNW	77

DESCRIPTION OF THE EMBODIMENTS

One embodiment of the present invention provides a method of detecting a virus. The method is based on a proteolytic enzyme assay that indicates the presence of the virus by detecting a viral enzyme present during replication of the virus.

Cleavage of the Viral Polyprotein

During the replication of many viruses, such as the SARS virus, human immunodeficiency virus, human papilloma virus, herpes virus, rhinovirus, picomavirus, coronavirus, hepatitis C virus, and others, the viral genetic material is transcribed to form a polyprotein, which is ultimately cleaved into two or more biologically active proteins.

The cleavage of the viral polyprotein into individual proteins is a critical part of the viral life cycle. Many viruses, including those of the adenovirus, baculovirus, comovirus, picomavirus, retrovirus, and togavirus families, encode proteases which cleave the viral polyprotein at these specific cleavage positions to form the active proteins required for viral replication. For example, polyprotein processing during replication of the Cocksfoot mottle virus (genus *Sobemovirus*) is described in Makinen, K. et al., J. Gen. Virol. vol. 81, pp 2783-89 (2000), the contents of which are incorporated by reference.

For the purposes of this invention, a "protease" is an enzyme that catalyses the hydrolysis of a peptide bond. Proteases are part of the hydrolase class of enzymes, which are selected from the enzymatic class having the general reaction equation:



Some virally-encoded proteases cleave only the polyprotein of a specific virus. Others cleave the polyprotein of more

than one type of virus. The specificity of protease action arises from the nature of the interaction of the protease at the cleavage region(s) of the polyprotein. A given protease will only cleave the polyprotein at a region having certain defined amino acid sequences. For example, the Rhinovirus HRV14 3C protease will cleave a polyprotein at a Glu-Gly, Glu-Ala or Glu-Gly dipeptide sequence. In addition, the rate of cleavage at these positions varies, depending on the peptide sequence of the polyprotein surrounding the cleavage position.

Thus, a virally-encoded protease can cleave the viral polyprotein at a number of different positions if the amino acid sequence at these positions is such that the cleavage reaction occurs. Similarly, if such sequences are conserved across a number of different viruses, a single protease may cleave the polyprotein of these viruses. The present method takes advantage of this specificity to provide detection methods that are specific for a single virus type or for more than one virus type.

It is believed, but not relied upon for that present invention, that during replication of certain viruses, viral-encoded proteases are autocleaved from the viral polyprotein at an early stage for the viral replication process. For example, the Picornavirus viral polyprotein is processed into the individual protein products by the viral 3C protease. This protease derives from a polypeptide sequence that is itself a part of the viral polyprotein. The mechanism by which this protease becomes cleaved from the viral polyprotein is described in Khan, A. R. et al. "Structural aspects of activation pathways of aspartic protease zymogens and viral 3C protease precursors", Proc. Nat'l. Acad. Sci., Vol. 96, No. 20, pp. 10968-75 (1999), the contents of which are incorporated by reference. For other viruses, the viral protease responsible for cleaving the polyprotein may be encoded separately from the polyprotein.

One embodiment of the present invention provides a method of detecting a virus in a sample obtained from a patient suspected of being infected with the virus. The method is applicable to the detection of the wide range of viruses producing a virally-encoded protease that cleaves the viral polyprotein. For the purposes of this invention, a "virally-encoded protease" or "viral protease" is defined as any protease encoded from the genetic material of a virus and which cleaves a peptide bond in the viral polyprotein during replication of the virus.

Virally-encoded proteases cleave the polyprotein of a wide range of virus families, including the Nidovirus, Herpesvirus, Adenovirus, Retrovirus, Picornavirus and Potyvirus families. Rao, M. B. et al., "Molecular and Biotechnological Aspects of Microbial Proteases", Microbiol. and Mol. Biology Rev., Vol. 62, No. 3, p. 597-635 (1998), the contents of which are incorporated by reference. Table 1 shows some examples of the proteases involved in polyprotein cleavage of viruses associated with various animal and plant diseases. Additional examples of such proteases are described in Corbalenya, A. E. and Snijder, E. J., "Viral cysteine proteinases, Perspectives in Drug Discovery and Design" 6: pp. 64-86 (1996) and Dougherby, W. G. and Semler, B. L., "Expression of virus-encoded proteinases: functional and structural similarities with cellular enzymes", Microbiol. Rev. 57, pp. 781-822 (1993), John Ziebuhr et al. "Virus-encoded proteinases and proteolytic processing in the Nidovirales", J. Gen. Vir., Vol. 81, pp. 853-79 (2000), the contents of which are incorporated by reference.

TABLE 1

EXAMPLES OF VIRAL-CODED PROTEASES INVOLVED IN POLYPROTEIN PROCESSING		
Virus	Protease	Associated Disease
<i>Picornaviridae</i>		
Enterovirus	3C (CSL), 2A(CSL)	Meningitis, gastro-intestinal infections
Coxsackievirus	3C	Common cold
Echovirus	3C	Summer flu, hand-foot-and-mouth
Poliovirus	3C	Poliomyelitis
Rhinovirus	3C (CSL), 2A(CSL)	Common cold, asthma exacerbation in allergies
Aphthovirus	3C (CSL), L (Cys)	Foot and mouth disease
Cardiovirus	3C (CSL)	Encephalitis, heart disease (mainly murine, affects other mammals)
Hepatovirus	3C (CSL)	Hepatitis A (chronic jaundice)
<i>Togoviridae</i>		
Alphaviruses	Cys and Ser	Equine encephalitis
Poxviruses	Ser	Smallpox
Rubiviruses	Cys	Rubella (German measles)
<i>Paramyxoviridae</i>		
Parainfluenza	Ser	Respiratory infection
RSV	Ser	Infant bronchiolitis, viral pneumonia
<i>Coronaviridae</i>	Cys, 3C-like CSL, Ser	Infant bronchiolitis, viral pneumonia
SARS	Cys, 3C like, PL1 PL2	Acute respiratory syndrome
Arterivirus	Cys	Pig disease
<i>Flaviviridae</i>		
Flavivirus	NS3 (Ser, unique), NS2B (Ser, unique)	
Yellow-fever virus	NS3 (Ser, unique)	Yellow-fever
HepC virus	NS3 (Ser, unique)	Hepatitis C
Pestivirus	Cys, Ser	Pigs, cattle and sheep disease
<i>Adenoviridae</i>		Acute upper respiratory, eye and intestinal tract, infant death
<i>Herpesviridae</i>	<i>Herpesviridae</i>	Herpes (systemic and topical)
<i>Retroviridae</i>		
HIV	Asp	AIDS
<i>Caliciviridae</i>	3C-like (CSL)	Rabbit hemorrhagic disease
<i>Potyviridae</i>		
Potyvirus	Nla (3C-like CSL), HC ^{pro} (Cys)	Potato disease
Bymovirus	3C-like (CSL), Cys	Plant disease
<i>Comoviridae</i>		
Comovirus	p24 (3C-like CSL)	Plant disease
Nepovirus	p23 (3C-like CSL)	Plant disease

Sources

- Gorbalenya, A. E. and Snijder, E. J., Viral cysteine proteinases, Perspectives in Drug Discovery and Design 6: pp. 64-86 (1996)
- Dougherby, W. G. and Semler, B. L., Expression of virus-encoded proteinases: functional and structural similarities with cellular enzymes, Microbiol. Rev., Vol. 57, pp. 781-822 (1993)

Proteases are subdivided into two major groups, depending on their site of action. Exopeptidases cleave the peptide bond proximal to the amino or carboxy termini of the substrate,

whereas endopeptidases cleave peptide bonds distant from the termini of the substrate. Based on the functional group present at the active site, proteases are further classified into four prominent sub-groups. These are serine proteases, aspartic proteases, cysteine proteases, and metalloproteases.

Viruses produce many different proteases, including members of the above classes. For example, the Hepatitis C virus genome encodes a specific serine protease and a metalloprotease (NS2 and NS3) and the Rhinovirus genome encodes cysteine proteases (3C and 2A). The SARS genome encodes a cysteine protease (3CL). Anand K. et al., "Coronavirus Main Protease (3CL^{pro}) Structure: Basis of Design of Anti-SARS Drugs" *Science*, Vol. 30, pp. 1763-67 (2003), the contents of which are incorporated by reference. In addition, the SARS genome encodes the PL1 and PL2 main proteases. Each member of the herpesvirus family encodes a unique serine protease. Adenoviruses code for a serine-centered, neutral protease specific for selected Gly-Ala bonds in several virus-encoded precursor proteins. Retroviruses encode an aspartyl protease, which is responsible for processing the gag and pol polyprotein precursors into the structural proteins of the mature virus.

Viral-encoded Enzymes

The cleavage of the polyprotein to produce functional viral proteins is a key step in the virus maturation process. The cleavage step occurs at an early stage of viral replication cycle. This requires that the viral-encoded proteases responsible for this process are themselves present at an early stage of replication. Because the proteases are present early in the cycle of infection, their detection allows viral infection to be confirmed at an early stage.

Tissue infected by a virus contains fully assembled virus particles as well as components of the viral particles. These components include the active proteases. See Sosnovtsev, S. V., et al., "Cleavage of the feline calicivirus capsid precursor is mediated by a virus-encoded proteinase", *J. Virol.*, Vol. 72, pp. 3051-59 (1998), the contents of which are incorporated by reference. It is also known that certain viruses, for example the Influenza types A and B viruses, produce surface glycoproteins with neuraminidase activity whose presence confirms infection with the virus. ZSTATFLU® Product Package Insert (ZymeTx, Inc., Oklahoma City, Okla. 73104)

Detection of Polyprotein Cleaving Proteases

In one embodiment of the present invention, the presence of a viral-encoded protease associated with a virus is detected by observing the cleavage of a peptidal compound having a specific cleavage site for the protease. The specificity of the cleavage reaction depends upon the amino acid sequence at the cleavage site. Specific cleavage sites for many viral-encoded proteases have been identified. See, for example, John Ziebuhr et al. "Virus-encoded proteinases and proteolytic processing in the Nidovirales", *J. Gen. Vir.*, 81, pp. 853-79 (2000), the contents of which is incorporated by reference. The viral cleavage sites are chosen such that their cleavage is specific for the type of virus to be detected. Table 2 shows examples of some viral proteases along with peptides that are specifically cleaved by them.

TABLE 2

EXAMPLES OF SPECIFIC CLEAVAGE PEPTIDES	
Protease	Specific Cleavage Peptides
Rhinovirus 3CL	Arg-Pro-Val-Val-Val-Gln-Gly-Pro-Asn SEQ ID NO: 83
Coronavirus TGEV 3CL	Ser-Thr-Leu-Gln-Ser-Gly-Leu-Arg-Lys SEQ ID NO: 84

TABLE 2-continued

EXAMPLES OF SPECIFIC CLEAVAGE PEPTIDES	
Protease	Specific Cleavage Peptides
SARS COV 3CL	Ala-Thr-Val-Arg-Leu-Gln-Ala-Gly-Phe SEQ ID NO: 85
	Val-Ser-Val-Asn-Ser-Thr-Leu-Gln-Ser- Gly-Leu-Arg-Lys-Met-Ala-Cys SEQ ID NO: 86

In the present invention, a region of the viral polyprotein containing a cleavage site for the protease to be detected is mimicked by a peptidal compound. For the purposes of this invention, the term "peptidal compound" is any compound containing a cleavage site for a protease encoded by the virus or viruses that are to be detected. Under appropriate conditions, the protease will cleave the peptidal compound to form a "first peptidal compound fragment" and a "second peptidal compound fragment". The conditions required for performing such enzyme cleavage reactions are well known for those skilled in the art.

A region of the peptidal compound that forms one of the peptidal compound fragments is tagged by directly or indirectly attaching a signaling moiety that allows that fragment to be detected. For the purposes of the invention, the "signaling moiety" can be any label that produces a detectable signal. For example, the signaling moiety can be a detectable label that produces a fluorescent, a chemiluminescent or a calorimetric signal.

In one embodiment, the cleavage site is specific for a single virus. This allows for a specific test that detects this virus and shows low cross-reactivity for other viruses. In another embodiment, the cleavage site that conserved across more than one virus. This allows for a single test that will detect more than one virus.

It is believed, but not relied upon for the present invention, that since polyprotein-cleaving proteases are not present on the viral surface, they are less subject to mutation than are viral-coat proteins. Also, since these proteases have an important role in the virus life cycle, significant mutation is unlikely.

The method of the present invention can be used to test for the presence of a virus in a sample contaminated with the virus. In one embodiment, the sample is taken from an organism. For example, the method can test for a virus in a sample taken from an animal or plant host. In one embodiment, the method tests for a virus in a sample taken from a human patient. The sample can be mucus, saliva, throat wash, blood, serum, plasma, urine, spinal fluid, sputum, tissue biopsy, bronchoalveolar fluid, vaginal fluid, tear fluid or another biological sample. For example, the SARS virus is known to be present in saliva. Wang, W-K "Detection of SARS-associated Coronavirus in Throat Wash and Saliva in Early Diagnosis" *Emerg Infect Dis* [serial on the Internet] (June 2004). Available from: <http://www.cdc.gov/ncidod/EID/vol11no7/03-1113.htm>. The invention includes methods where the sample is treated to disrupt cells to release viral components, including the protease, e.g. by using sonification. The presence of a virus is confirmed by detecting a viral protease produced by the virus. In one embodiment, the protease level is correlated with the level of viral replication or viral load.

In one embodiment of the invention, the peptidal compound includes a sequence of at least four amino acids residues from the sequence in the region of a polyprotein cleav-

age site, including the protease cleavage site. In another embodiment, the peptidal compound includes at least seven amino acids residues from the protease cleavage site. In another embodiment, the peptidal compound includes at least ten amino acids residues from the protease cleavage site. In yet another embodiment, the peptidal compound includes at least four amino acids from a sequence that is a consensus sequence of the sequences at the cleavage sites of two or more viruses. In other embodiments, the peptidal compound contains at least seven, or at least ten, amino acids from a consensus sequence of the cleavage sites of two or more viruses.

In another embodiment, the one or more amino acids present in the peptidal compound are modified or replaced by an analog. Analogs are amino acids that have a structure similar to the native compound but differ from it in respect to certain components or side chains.

Alternatively, one or more amino acids may be replaced by similar amino acids without altering the cleavage properties of the mimic peptidal compound. Useful conservative substitutions are shown in Table 3, "Preferred substitutions." Conservative substitutions whereby an amino acid of one class is replaced with another amino acid of the same type fall within the scope of the invention so long as the substitution does not materially alter the cleavage properties of the peptidal compound. Such substitutions can be tested using a method such as that described in Example 5 to determine whether the substitution alters the cleavage properties.

TABLE 3

Original residue	Preferred substitutions	
	Exemplary substitutions	Preferred substitutions
Ala (A)	Val, Leu, Ile	Val
Arg (R)	Lys, Gln, Asn	Lys
Asn (N)	Gln, His, Lys, Arg	Gln
Asp (D)	Glu	Glu
Cys (C)	Ser	Ser
Gln (Q)	Asn	Asn
Glu (E)	Asp	Asp
Gly (G)	Pro, Ala	Ala
His (H)	Asn, Gln, Lys, Arg	Arg
Ile (I)	Leu, Val, Met, Ala, Phe, Norleucine	Leu
Leu (L)	Norleucine, Ile, Val, Met, Ala, Phe	Ile
Lys (K)	Arg, Gln, Asn	Arg
Met (M)	Leu, Phe, Ile	Leu
Phe (F)	Leu, Val, Ile, Ala, Tyr	Leu
Pro (P)	Ala	Ala
Ser (S)	Thr	Thr
Thr (T)	Ser	Ser
Trp (W)	Tyr, Phe	Tyr
Tyr (Y)	Trp, Phe, Thr, Ser	Phe
Val (V)	Ile, Leu, Met, Phe, Ala, Norleucine	Leu

By selecting a peptidal compound to mimic a particular polyprotein cleavage site, the method of the present invention can be used to test for a wide range of viruses. In one embodiment, the virus detected is from the picornavirus family. This family of viruses includes the human rhinoviruses and the coxsackieviruses. FIGS. 2A-D show the sequences around known cleavage positions in the polyproteins encoded by some such viruses. The viruses shown are Human Rhinovirus 14 (HRV14), Human Rhinovirus b (HRVb), Human Rhinovirus 16 (HRV16) and Coxsackievirus A21 (Cxa21). The figures also show sequence alignment between the cleavage sites of the viruses and consensus peptide sequences. Sequence alignment was performed using the CLUSTAL W

(1.82) general purpose sequence analysis program. (European Bioinformatics Institute, available at www.ebi.ac.uk/clustalw).

In the test method for a particular virus, protease cleavage of a peptidal compound containing an amino acid sequence including one of the polypeptide cleavage regions is detected. By selecting a peptidal compound that shows low homology to sequences present in similar viruses, a specific test for the virus can be performed. The invention also includes methods where two or more peptidal compounds are cleaved, either by a single protease or by different proteases.

In another embodiment, the virus detected is a Nidovirus. This class of viruses includes the SARS virus, Bovine Coronavirus and Human Coronavirus. The SARS genome is available at: M. Marra et al., <http://www.bcgsc.ca/bioinfo/SARS>. The genome shows more than a 44% homology with the coronavirus genome and, like the coronavirus genome, encodes a 3CL protease (Sequence homology performed using the software packages available at EMBL (European Bioinformatics Institute, <http://www.ebi.ac.uk/emb1/index.html>)).

FIGS. 3A-D show the amino acid sequences around known cleavage positions in the polyproteins encoded by these viruses. Sequence alignments between the cleavage sites of the three viruses and consensus sequences are also shown. Sequence alignment was again performed using the CLUSTAL W (1.82) sequence analysis program. FIG. 3(A) shows that the cleavage regions centered at, for example, positions 180 or 818 of the SARS polyprotein shows little homology with the Human or Bovine Coronavirus polyprotein. By choosing peptide sequence mimicking the sequence of one of these cleavage positions, a specific test for the SARS virus can be performed. Similarly, by choosing a Human Coronavirus polyprotein sequence showing low homology to either the SARS or Bovine Coronavirus polyprotein, a specific test for Human Coronavirus can be performed.

Assays for Protease Cleaving Activity

One embodiment of the present invention provides a homogenous assay for the detection of a virus. The chosen peptidal compound is synthesized and linked to a signaling moiety at one side of the cleavage region and to a quencher moiety at the other side of the cleavage region. A "homogeneous assay" is an assay not requiring separation of signaling moiety from other assay components.

For the purposes of the invention, a "quencher moiety" is any substance that is capable of reducing or eliminating the signal emitted by the signaling moiety. For example, the quencher moiety may act by absorption of the signal emitted by the signaling moiety or by an energy transfer mechanism. The distance between the signaling moiety and the quencher moiety is such that presence of the quencher moiety substantially reduces or eliminates the signal emitted from the signaling moiety unless the peptidal compound is cleaved at a position resulting in separation of the signaling and quencher moieties.

In one embodiment, the signaling moiety and quencher moiety are separated by no more than 5 amino acid residues. In another embodiment, the signaling moiety and quencher moiety are separated by no more than 10 amino acid residues. In yet another embodiment, the signaling moiety and quencher moiety are separated by no more than 15 amino acid residues. In yet another embodiment, the signaling moiety and quencher moiety are separated by no more than 20 amino acid residues.

The peptidal compound is contacted with the sample being tested for the presence of a virus. If the virus is present in the

sample, the viral protease is also present. This protease cleaves the peptidal compound and a change in the signal from the signaling moiety can be observed. FIG. 1 shows a representation of the cleavage reaction. Such homogenous fluorescent and colorimetric assays are known to those skilled in the art. See, for example: Biochemistry, Allinger, Wang Q. M. et al., "A continuous calorimetric assay for rhinovirus-14 3C protease using peptide p-nitroanilides as substrates" Anal. Biochem. Vol. 252, pp. 238-45 (1997), and Basak S. et al. "In vitro elucidation of substrate specificity and bioassay of pro-

protein convertase 4 using intramolecularly quenched fluorogenic peptides" Biochem. J. Vol. 380, pp. 505-14 (2004) the contents of which are incorporated by reference. However, it is believed that such methods have not previously been utilized for the diagnosis of viral disease.

In another embodiment of the present invention, the signaling moiety is a chemiluminescent signaling moiety. The chemiluminescent signaling moiety is attached to one side of the cleavage region of the peptidal compound and a fluorescent accepting quencher moiety is attached at the other side of the cleavage region. U.S. Pat. No. 6,243,980, the contents of which are incorporated by reference, describes such a detection system, involving the use of a chemiluminescent 1,2-dioxetane compound as the signaling moiety. If the viral protease is not present in the sample, cleavage of the peptidal compound does not occur. The energy from the 1,2-dioxetane decomposition is transferred to the fluorescent accepting moiety and released at a wavelength distinct from the emission spectrum of the 1,2-dioxetane. If the peptidal compound is cleaved, the fluorescent accepting moiety is separated from the 1,2-dioxetane and a chemiluminescent emission from the dioxetane compound is observed.

In another embodiment, the signaling moiety is a fluorescent compound and the quencher moiety is a fluorescent compound having an excitation spectrum that overlaps the emission spectrum of the signaling moiety. Here, the two moieties are separated apart at a distance consistent with fluorescent resonance energy transfer so that the fluorescent moiety is capable of acting as a resonance energy donor.

In another embodiment, a quenching group, such as a non-fluorescent absorbing dye is used in place of the fluorescent accepting quenching moiety. Suitable quenching groups are described in U.S. Pat. No. 6,243,980, the contents of which are incorporated by reference.

In such a test method, the test sample is contacted with the peptidal compound under conditions that allow cleavage of the peptidal compound by the protease if the virus is present in the sample. In one embodiment, the temperature is controlled. For example, the temperature can be controlled at 37 deg C. to provide optimal conditions for the enzyme reaction. The signal from the cleaved peptidal compound fragment is then detected using a detection device appropriate to the label used.

Another embodiment of the present invention provides a heterogeneous assay for the detection of a virus. In one embodiment of the heterogeneous method, a first member of a first binding pair is linked to one side of the cleavage region of the peptidal compound. A first member of a second binding pair, or a signaling moiety, is linked to the other side of the cleavage region of the peptidal compound. A second member of the first binding pair is linked to a solid-phase. Alternatively, one side of the peptide can be linked directly to the solid-phase. A "heterogeneous assay" is an assay in which the solid-phase is separated from another assay component during the assay.

The peptidal compound is incubated with the solid-phase and a sample being tested for the presence of a virus. The

incubation conditions are such that binding occurs between the first and second members of the first binding pair. If a protease having the ability to cleave the peptidal compound at the cleavage site is present in the patient sample, the peptidal compound is cleaved causing the peptidal compound fragment linked to the first member of a second binding pair, or the signaling moiety, to become detached from the solid phase.

The solid-phase and liquid phase are then separated. If a signaling moiety is linked to the peptidal compound, the amount of signaling moiety in either the solid phase or the liquid phase is measured to determine the presence of the protease, and hence the virus, in the sample. If a first member of a second binding pair is linked to the peptidal compound, a second member of the second binding pair linked to a signaling moiety is incubated with the peptidal compound under conditions such that binding occurs between the first and second members of the second binding pair. The presence of the virus is then detected as above.

The above steps may be varied or combined without departing from the method of the present invention. For example, those skilled in the art will recognize that, in the heterogeneous assay method, the second member of the second binding pair linked to the signaling moiety can be contacted with the peptidal compound before or at the same time as the solid-phase is contacted with the peptidal compound.

Many different binding pairs may be used for the first and second binding pairs. However, the binding pairs must be chosen such that that the first binding pair does not interact with the second binding pair. Examples of such binding pairs are well known in the art and include biotin/avidin, biotin/streptavidin, antibody/antigen, antibody/hapten, binding protein/bound molecule and complementary nucleic acid sequences. The first member of each binding pair, or the signaling moiety, can be directly linked to the peptide by a covalent bond or indirectly via a spacer molecule having coupling functional groups at each end. Examples of such linkers include an alkyl, a glycol, an ether, a polyether, a polynucleotide and a polypeptide molecule.

Solid-phases suitable for use in the heterogeneous assay include, but are not limited to test tubes, microtiter plates, microtiter wells, beads, dipsticks, polymer microparticles, magnetic microparticles, nitrocellulose, chip arrays and other solid phases familiar to those skilled in the art. The signaling moiety used in the heterogeneous assay may be any label known to those skilled in the art. Such labels include radioactive, calorimetric, fluorescent and luminescent labels.

A heterogeneous chemiluminescent assay for the detection of proteases is described in U.S. Pat. No. 56,243,980, the contents of which are incorporated by reference. Here, the second member of the second binding pair is conjugated with an enzyme, such as alkaline phosphatase, which triggers 1,2-dioxetane to emit a chemiluminescent signal.

In one embodiment, the homogeneous or heterogeneous assay method of the present invention is automated so that a result can be obtained without the need for medical staff to be exposed to a patient thought to be infected by the viral disease under test. For example, the patient can be tested in a clean room (for example, but not limited to P3 type room). The patient can pick up, or get before entering the room, a diagnostic kit, which can include a solid phase coated with a labeled peptide of the type discussed above. For example, the solid phase can be a tissue which was previously immersed with peptide, or a test stick that can be from the type used to test pregnancy. The patient can supply a sample, such as a saliva sample, at a pre-prepared spot on the solid phase.

The solid phase containing the sample is then incubated to allow the enzymatic reaction to occur. In one embodiment, the reaction temperature is controlled at 37° C. to provide optimal conditions for the enzyme reaction. When the incubation is complete, the sample to be tested can be measured on a spectrophotometer, using a remote control, or a mechanical system operated manually from outside the room. The sample can be tested for a qualitative color or UV detection. After the test the sample can be discarded by an automated system, or a remote operated handle that trashes the sample.

Detection of New Emerging Viral Diseases

In another embodiment, the present invention provides a method for developing an assay for the detection of a newly emergent virus. Once the genome of the emergent virus is identified and its reproduction system known, viral proteases and those regions of the viral polyprotein that are cleaved by such proteases can be determined by examining the sequence homology between the sequence of the emergent virus and that of known viruses. Specific mimic peptide compounds bridging the regions cleaved by the protease are then prepared and tested to determine the activity of the polyprotein cleaving protease against each peptide. In one embodiment, the peptide showing the highest rate of cleavage is chosen and used in the preparation of the test for the virus. The cleavage rates of such compounds may be determined using a method such as that described in Orr D.C. et al. "Hydrolysis of a series of synthetic peptide substrates by the human rhinovirus 143C proteinase, cloned and expressed in *E. coli*", *J. Gen. Vir.* Vol. 70, pp. 2931-42 (1989), the contents of which are incorporated by reference.

The reason for the effectiveness of the method for new emerging viral diseases is that the basic function of the proteolytic activity is maintained in the overall mechanism of the virus maturation. Specific proteases can be rapidly analyzed once the genome of the new virus has been identified. Thus, a new protease from a new emerging virus is constructed based on the template of a protease from a similar known virus from the same family. For each virus family, a library of peptides can be synthesized and the rate of cleavage by the protease measured. Based on these results, a consensus peptide is chosen, as is shown in Example 5 for rhinovirus.

The proteolytic class of the enzyme can be classified using data in the literature. Several databases exist that include an extensive amount of information about enzymes as they relate to their designated class. For example, Swiss-Prot Enzyme nomenclature database at <http://kr.expasy.org/enzyme> or classification of enzymes at http://www-biol.paisley.ac.uk/marco/enzyme_electrode/Chapter1/page_1a.htm. The enzyme can be further classified upon sequencing the enzyme and comparing its homology to other enzymes. Often times, the sequence for the enzyme has already been determined and can thus be found in the literature or a database (e.g. Genbank) <http://www.ncbi.nlm.nih.gov/Genbank/> or UniProt/Swiss-Prot Protein Knowledgebase (European Bioinformatics Institute at <http://www.ebi.ac.uk/swissprot/> or GOR IV Secondary Structure Prediction Method (e.g. http://npsa-pbil.ibcp.fr/cgi-bin/npsa_automat.pl?page=NPSA/npsa_server.html). If the enzyme sequence has not been identified, the enzyme can be isolated according to a variety of techniques known to those skilled in the art. Additionally, the enzyme can be synthesized using recombinant DNA techniques that are commonly known to scientists practicing in the field of molecular biology (See Short Protocols in Molecular Biology, 2d. ed., John Wiley & Sons (1992), which is incorporated herein by reference.

Sequencing of the target protein can be performed at the amino acid level and/or the DNA level using techniques known to those skilled in such an art. To determine the DNA sequence of the target protein, the method discussed in Wilhelm Arrange, DNA sequencing strategies, Automated and Advanced Approaches, 15 BN 0971136832 (S. Zimmerman, ed.), which is also incorporated herein by reference, can be used. The determined sequence of the enzyme can be compared for homology to other known sequences of classified enzymes having an identified tertiary, or preferably quaternary structure. Information regarding such enzymes are located databases, such as the Brookhaven Crystallographic Data Bank (<http://pdb.pdb.bnl.gov>).

After categorizing the chemical reactions the enzymes undergo, the specific peptide sequence at the cleavage regions of the viral polyprotein is determined by comparing the genome sequence to a homologous genome. Mimic peptidal compounds are then prepared as described above. In one embodiment, the peptidal compounds contain peptides having 6-15 amino acids from the cleavage regions. A library of such compounds is prepared and the cleavage rate determined for each compound determined. Peptidal compounds having high cleavage rates are then selected for use in viral tests. Kits, Compositions and Reagents for the Detection of Viral Disease

The present invention also provides for kits for detecting a viral disease. In one embodiment, the kit contains at least a reagent comprising one the mimic peptidal compounds described above and a buffer in a package or container. In one embodiment, a signaling moiety is linked to a portion of the peptidal compound present in one peptidal compound fragment. In another embodiment, a quencher moiety linked to a portion of the peptidal compound present in the other peptidal compound fragment. The signaling moiety and the quencher moiety are linked to the peptidal compound at relative positions such that the quencher molecule quenches the signal of the reporter molecule unless the peptidal compound is cleaved at the cleavage point.

In another embodiment, the kit contains the following in one or more packages or containers: (a) a construct reagent comprising (i) a peptidal compound capable of being cleaved by the enzyme to form two peptidal compound fragments, (ii) a first member of a first binding pair linked to a portion of the peptidal compound present in one peptide fragment, and (iii) a first member of a second binding pair linked to a portion of the peptidal compound present in the other peptidal compound fragment, (b) a solid-phase reagent comprising a second member of the first binding pair is linked to a solid-phase, and (c) a signaling reagent comprising a second member of the second binding pair linked to a signaling moiety.

In yet another embodiment, the kit contains the following in one or more packages or containers: (a) a construct reagent comprising (i) a peptidal compound capable of being cleaved by the enzyme to form two peptidal compound fragments, (ii) a first member of a first binding pair linked to a portion of the peptidal compound present in one peptidal compound fragment, and (iii) a signaling moiety linked to a portion of the peptidal compound present in the other peptidal compound fragment and (b) a solid-phase reagent comprising a second member of the first binding pair is linked to a solid-phase.

When a kit is supplied, the different components may be packaged in separate containers and admixed immediately before use. Such packaging of the components separately may permit long-term storage without losing the active components' functions. Embodiments in which two or more of components (a)-(c) are found in the same container are also contemplated.

In another embodiment, the above kits further comprise one or more of the following reagents:

- (a) a wash buffer reagent for use using heterogeneous assays;
- (b) a negative control reagent free of a protease capable of cleaving the construct reagent;
- (c) a positive control containing a protease capable of cleaving the construct reagent;
- (d) a signal generation reagent for development of a detectable signal from the signaling moiety; and
- (d) a sample collection means such as a syringe, throat swab, or other sample collection device.

The reagents included in the kits can be supplied in containers of any sort such that the life of the different components are preserved and are not adsorbed or altered by the materials of the container. For example, sealed glass ampules may contain lyophilized reagents, or buffers that have been packaged under a neutral, non-reacting gas, such as nitrogen. Ampules may consist of any suitable material, such as glass, organic polymers, such as polycarbonate, polystyrene, etc.; ceramic, metal or any other material typically employed to hold similar reagents. Other examples of suitable containers include simple bottles that may be fabricated from similar substances as ampules, and envelopes, that may comprise foil-lined interiors, such as aluminum or an alloy. Other containers include test tubes, vials, flasks, bottles, syringes, or the like. Containers may have a sterile access port, such as a bottle having a stopper that can be pierced by a hypodermic injection needle. Other containers may have two compartments that are separated by a readily removable membrane that upon removal permits the components to be mixed. Removable membranes may be glass, plastic, rubber, etc.

Kits may also be supplied with instructional materials. Instructions may be printed on paper or other substrate, and/or may be supplied as an electronic-readable medium, such as a floppy disc, CD-ROM, DVD-ROM, Zip disc, videotape, audiotape, etc. Detailed instructions may not be physically associated with the kit; instead, a user may be directed to an internet web site specified by the manufacturer or distributor of the kit, or supplied as electronic mail.

In another embodiment, the invention provides a composition for detecting a virus in a sample. The composition includes a buffer and a mimic peptidal compound capable of being cleaved at a cleavage point by a protease produced by the virus to form a first peptidal compound fragment and a second peptidal compound fragment. A signaling moiety is linked to a portion of the peptidal compound present in the first peptidal compound fragment and a quenching moiety is linked to a portion of the peptidal compound present in the second peptidal compound fragment. The signaling moiety and the quenching moiety are linked to the peptidal compound at relative positions such that the quenching moiety quenches a signal of the signaling moiety.

In one embodiment, the virus is a Rhinovirus. In this embodiment, the composition can include a peptidal compound containing an amino acid sequence present one of the cleavage sites on the Rhinovirus polyprotein. For example, the amino acid sequence can be one of the Rhinovirus sequences listed in Table 2.

In another embodiment, the virus is the SARS virus. In this embodiment, can include a peptidal compound can containing an amino acid sequence present one of the cleavage sites on the SARS virus polyprotein. For example, the amino acid sequence can be one of the Rhinovirus sequences listed in Table 3.

A more complete understanding of the present invention can be obtained by reference to the following specific

Examples. The Examples are described solely for purposes of illustration and are not intended to limit the scope of the invention. Changes in form and substitution of equivalents are contemplated as circumstances may suggest or render expedient. Although specific terms have been employed herein, such terms are intended in a descriptive sense and not for purposes of limitations. Modifications and variations of the invention as hereinbefore set forth can be made without departing from the spirit and scope thereof, and, therefore, only such limitations should be imposed as are indicated by the appended claims.

EXAMPLES

Example 1

Detection of Protease Activity

Enzymatic activity for 3C protease can be detected using chromogenic substrates as described in: Wang, Q. M. et al. "A continuous colorimetric assay for rhinovirus-14 3C protease using peptide p-nitroanilides as substrates" Anal. Biochem. Vol. 252, pp. 238-45 (1997), the contents of which are incorporated by reference. Tagged substrates are used to determine the ability of the protease to cleave. The first peptide substrate used is tagged using p-nitroaniline. When p-nitroaniline is cleaved from the peptide, a signal is produced. The cleavage causes an aromatic pi-electron system to form, the presence of which absorbs in the 405 nm range of the electromagnetic spectrum. The nanomolar extinction coefficient of the cleaved p-nitroaniline is $10^4 \text{ mole}^{-1} \text{ cm}^{-1}$.

Alternatively, a substrate is constructed having a fluorescent tag attached to one end and a quencher attached to the other end. When the peptide is cleaved fluorescence is detected. Other tags use a similar principle using color reactions.

Example 2

Homogenous Fluorescence Resonance Energy Transfer Assay for 3C Protease

Human Rhinovirus serotype 1A (ATCC) is used to clone the 3C Protease into the expression vector pET16-b and transformed for production into the *E. coli* strain BL21-DE3-pLys-S. 3C Protease expression is induced with 1 mM IPTG at 25° C. and purified from the soluble protein extract by chromatography on a SourceQ (Pharmacia) followed by gel filtration. HRV 3CP activity was measured by fluorescence resonance energy transfer using a dimodified decapeptide substrate MOC-Arg-Ala-Glu-Leu-Gln-Gly-Pro-Tyr-Asp-Lys-DNP-NH₂ (SEQ ID NO: 113) (7-methoxy coumarin-4-acetic acid fluorochrome and dinitrophenol quencher) with a Km value of 16.8 μM. Inhibition was measured as a change in initial velocity (V₀) as a function of inhibitor (I) concentration and substrate (S) concentration.

Assays are performed in 100 uL volumes in a 96 well format at 30° C. containing 25 mM Tris HCl pH 8.0, 150 mM NaCl, 1 mM EDTA pH 8.0, 6 mM DTT, 2-6 uM substrate, 2% DMSO, 416 nM 3CP and inhibitor as needed. Fluorescence is monitored by excitation at 328 nm and emission at 393 nm with 10 nm cutoffs. Data are analyzed with the nonlinear regression analysis program EnzFitter (BioSoft) with the equation:

$$K_i = (I / ((V_{max} \times S) / V_0) / K_s) - I - S$$

Substrate concentrations used are lower than the K_m of the substrate (16.8 μ M) so no corrections for an S/K_m term were used.

Example 3

Using the CELLSCAN® Cytometer to Detect Protease Activity

The CELLSCAN® (Medis Technologies Ltd., New York, N.Y.) is a cytometer that can monitor the fluorescence intensity and polarization by using fluorescence probes within individual, non-adherent living cells and can also be used to detect protease activity. The heart of the CELLSCAN is a Cell carrier that contains up to 10,000 wells. A description of the CELLSCAN cytometer and its other uses for diagnosis of cancer and autoimmune diseases is available at: www.medisel.com.

A CELLSCAN probe is loaded with a specific peptidal compound. For example, the peptidal compound can contain a peptide that is specific for SARS. The peptidal compound is tagged with a fluorescent group on one side of the cleavage region and a quencher on the other. The sample under test, e.g. saliva or mucus, is loaded on the probe

If active protease exists, the CELLSCAN will detect fluorescence caused by the cleaved peptide. The presence of active protease and its concentration in the saliva is an indication of an active virus and serves as an indication for the contagious status of the patient.

Example 4

Paper-tissue-Based Automated System for the Detection of SARS

A solution of a color-tagged peptidal compound specific for 3Cl-SARS protease is prepared. The peptide is prepared in a pH 7.0 buffer solution in a millimolar concentration. A tissue (wet-wipes tissues) is immersed in the peptidal compound solution and is kept moist. The saliva or mucus specimen suspected of containing the SARS virus is put in contact with the tissue. If the SARS virus is present, the 3CL SARS protease cleaves the tagged peptide sequence and a color reaction occurs.

Several possibilities exist for detection of the colored reaction product. For a qualitative analysis, a color reaction may be detected visually. For a quantitative analysis, the tissue is transferred to spectrophotometric analyzer for either fluorescence or color detection. The process can be automated so as to protect those performing the assay from infection.

Example 5

Determining an Optimal Peptide Sequence for a Rhinovirus Assay

The cleavage sites of the Rhinovirus HRV14 3C protease are aligned as shown in Table 4. Table 4 also shows the QG dipeptides that appear in the HRV14 primary sequence, showing the six sites that are cleaved. The QA and EG dipeptide sites are also shown. One of each of these sites is known to be cleaved.

TABLE 4

QG DIPEPTIDES THAT APPEAR IN THE HRV14 PRIMARY SEQUENCE			
Location on polyprotein	Sequence	QG Site	Cleavage Site
331	SEQ ID NO: 87 KSIIV <u>QG</u> LPTTT		IB1C
1,002	SEQ ID NO: 88 CIAEB <u>QG</u> LSDYI		2A2B
1,429	SEQ ID NO: 89 LET <u>LFQGP</u> VYKD		2C3A
1,514	SEQ ID NO: 90 LFA <u>QTQGP</u> YSGN		3A3B
1,537	SEQ ID NO: 91 RPVV <u>VQGP</u> NTEF		3B3C
1,704	SEQ ID NO: 92 GGNGR <u>QGP</u> SAQL		
1,719	SEQ ID NO: 93 YFVEK <u>QGP</u> VIAR		3C3D
<u>QA Site</u>			
716	SEQ ID NO: 94 SNL <u>VVQA</u> MYVPH		
950	SEQ ID NO: 95 YPSR <u>FQA</u> GVMKG		
1,099	SEQ ID NO: 96 PYIER <u>QA</u> NDGWF		2B2C
1,139	SEQ ID NO: 97 NKVLP <u>QA</u> KEKLE		
1,485	SEQ ID NO: 98 ERAMN <u>QA</u> SMIIN		
<u>EG Site</u>			
205	SEQ ID NO: 99 QLASH <u>EG</u> GNVSV		
567	SEQ ID NO: 100 TVALT <u>EG</u> LGDLEL		IC1D
940	SEQ ID NO: 101 TNIWI <u>EG</u> SPYYP		
982	SEQ ID NO: 102 GLLT <u>AEG</u> SGYVC		
1,632	SEQ ID NO: 103 ISEDL <u>EG</u> VDTL		
2,001	SEQ ID NO: 104 EIYVV <u>EG</u> GMPSG		

The cleavage peptides that mimic the cleavage sites for rhinovirus HRV14 are constructed and presented in Table 5.

TABLE 5

CLEAVAGE PEPTIDES		
Cleavage Peptide	Cleavage site Mimicked	Peptide
SEQ ID NO: 105 DSLETLFQGPVYK	2C/3A	I
SEQ ID NO: 106 EAIAEEQGLSDYIT	2A/2B	II
SEQ ID NO: 107 VPYIERQANDGWFRK	2B/2C	III
SEQ ID NO: 108 RSKSIVPQGLPTTTY	1B/1C	IV
SEQ ID NO: 109 SQTVALTEGLGDELEEY	1C/1D	V
SEQ ID NO: 110 KLFAQTQGPYSGNP	3A/3B	VI
SEQ ID NO: 111 YRPVVVQGPNTFE	3B/3C	VII
SEQ ID NO: 112 KQYFVEKQGVVIAR	3C/3D	VIII

Next, the rate of hydrolysis for each of the peptidal compounds is determined. The assay used is a fluorometric, or HPLC based or UV based assay. Such assays are well known to those skilled in the art. One such method is described in Orr D. C. et al. "Hydrolysis of a series of synthetic peptide substrates by the human rhinovirus 14 3C proteinase, cloned and expressed in *E. coli*", J. Gen. Vir. Vol. 70, pp. 2931-42 (1989). Table 6 shows the relative rate of hydrolysis for the cleavage of synthetic peptides that mimic the HRV14 3C cleavage sites.

TABLE 6

RELATIVE RATE OF HYDROLYSIS FOR THE CLEAVAGE OF SYNTHETIC PEPTIDES THAT MIMIC THE HRV14 3C CLEAVAGE SITES (From Orr D. C. et al.)			
Relative Hydrolysis rate	Hydrolysis rate	Cleavage	Peptide
1.00	355	2C/3A	I
0.024	8	2A/2B	II
0.0005	0	2B/2C	III
0.0005	0	1C/1D	V
0.44	147	3A/3B	VI
0.11	37	3B/3C	VII
0.003	1	3C/3D	VIII

The best substrate is chosen based on the reaction rates. Based on the results, a library of different sized peptides is constructed and the analyzed by the same assay. For Rhinovirus 14 the best peptide is: Arg-Pro-Val-Val-Val-Gln-Gly-Pro-Asn.

Example 6

Choosing a Cleavage Peptide from Picornaviruses and Coxsaciviruses.

Table 7 shows an alignment of the amino acid sequences at the 2A.1C cleavage site for different Picornaviruses and Coxsaciviruses. The cleavage site is at the QG dipeptide. The sequences show a high degree of homogeneity. Such homogeneity can be used to identify a cleavage site in a particular virus if cleavage sites for other member of the virus family have been identified.

Once a cleavage site showing a high cleavage rate is identified for certain member of a virus family, a sequence at the cleavage site can be identified in a related virus on the basis of the homogeneity of the sequences.

TABLE 7

ALIGNMENT OF CLEAVAGE SITES FOR DIFFERENT CORNA VIRUSES AND COXSACIVIRUSES	
HUMAN COXSACIVIRUS COXA2 SED ID NO: 118	i g n c m e a l F Q G p l r y k d l k i
HUMAN COXSACIVIRUS COXA4 SED ID NO: 119	i g n c m e a l F Q G p i q y r d v m i
BOVINE ENTEROVIRUS BOVEV SED ID NO: 120	i g n v l e a l F Q G p v c y k p l r i
HUMAN COXSACIVIRUS COXA9 SED ID NO: 121	v g a t l e a l F Q G p p i y r e i k i
HUMAN COXSACIVIRUS COXB1 SED ID NO: 122	v g a t l e a l F Q G p p i y r e i k i
HUMAN COXSACIVIRUS COXB5 SED ID NO: 123	v g a t l e a l F Q G p p i y r e i k i
HUMAN ECHOVIRUS EC11G SED ID NO: 124	v g a t l e a l F Q G p p i y r e i k i

TABLE 7-continued

ALIGNMENT OF CLEAVAGE SITES FOR DIFFERENT CORNA VIRUSES AND COXACIVIRUSES	
HUMAN COXACIVIRUS COXB4 SED ID NO: 125	v g a t l e a l F Q G p p v y r e i k i
SWINE VESICULAR DISEASE VIRUS SVDVH SED ID NO: 126	v g a t l e a l F Q G p p v y r e i k i
SWINE VESICULAR DISEASE VIRUS SVDVU SED ID NO: 127	v g a t l e a l F Q G p p v y r e i k i
HUMAN COXACIVIRUS COXB3 SED ID NO: 128	v g t t l e a l F Q G p p v y r e i k i
HUMAN ENTEROVIRUS HUEV7 SED ID NO: 129	t q d k l e a l F Q G p p t f k e i k i
HUMAN RHINOVIRUS HRV1B SED ID NO: 130	v v d v m s a i F Q G p i s l d a p p p
HUMAN RHINOVIRUS HRV2 SED ID NO: 131	v v d v m t a i F Q G p i d m k n p p p
HUMAN RHINOVIRUS HRV89 SED ID NO: 132	a a q a m e a i F Q G i d l q s p p p p
HUMAN RHINOVIRUS HRV14 SED ID NO: 133	i t d s l e t l F Q G p v y k d l e i d

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 133

<210> SEQ ID NO 1

<211> LENGTH: 20

<212> TYPE: PRT

<213> ORGANISM: Human rhinovirus 14

<400> SEQUENCE: 1

Leu Met Leu Lys Gly Ala Pro Ala Leu Asn Ser Pro Asn Val Glu Ala
1 5 10 15

Cys Gly Tyr Ser
20

<210> SEQ ID NO 2

<211> LENGTH: 20

<212> TYPE: PRT

<213> ORGANISM: Human rhinovirus 16

<400> SEQUENCE: 2

Val Leu Glu Lys Gly Ile Pro Thr Leu Gln Ser Pro Ser Val Glu Ala
1 5 10 15

Cys Gly Tyr Ser
20

-continued

<210> SEQ ID NO 3
 <211> LENGTH: 20
 <212> TYPE: PRT
 <213> ORGANISM: Human Rhinovirus b

<400> SEQUENCE: 3

Leu Met Leu Lys Gly Ala Pro Ala Leu Asn Ser Pro Asn Val Glu Ala
 1 5 10 15

Cys Gly Tyr Ser
 20

<210> SEQ ID NO 4
 <211> LENGTH: 20
 <212> TYPE: PRT
 <213> ORGANISM: Coxsackievirus A21

<400> SEQUENCE: 4

Leu Met Leu Lys Thr Ala Pro Ala Leu Asn Ser Pro Asn Val Glu Ala
 1 5 10 15

Cys Gly Tyr Ser
 20

<210> SEQ ID NO 5
 <211> LENGTH: 20
 <212> TYPE: PRT
 <213> ORGANISM: Human rhinovirus 14

<400> SEQUENCE: 5

Gly Ile Arg Ser Lys Ser Ile Val Pro Gln Gly Leu Pro Thr Thr Thr
 1 5 10 15

Leu Pro Gly Ser
 20

<210> SEQ ID NO 6
 <211> LENGTH: 20
 <212> TYPE: PRT
 <213> ORGANISM: Human rhinovirus 16

<400> SEQUENCE: 6

Ser Gly Ala Arg Ala Lys Thr Val Val Gln Gly Leu Pro Val Tyr Val
 1 5 10 15

Thr Pro Gly Ser
 20

<210> SEQ ID NO 7
 <211> LENGTH: 20
 <212> TYPE: PRT
 <213> ORGANISM: Human Rhinovirus b

<400> SEQUENCE: 7

Gly Ile Arg Ser Lys Ser Ile Val Pro Gln Gly Leu Pro Thr Thr Thr
 1 5 10 15

Leu Pro Gly Ser
 20

<210> SEQ ID NO 8
 <211> LENGTH: 20
 <212> TYPE: PRT
 <213> ORGANISM: Coxsackievirus A21

<400> SEQUENCE: 8

Leu Arg Asn Ile Thr Val Pro Val His Gln Gly Leu Pro Thr Met Asn

-continued

1 5 10 15

Thr Pro Gly Ser
 20

<210> SEQ ID NO 9
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: Human rhinovirus 14

<400> SEQUENCE: 9

Phe Lys Leu Arg Leu Met Lys Asp Thr Gln Thr Ile Ser Gln Thr Val
1 5 10 15

Ala Leu Thr Glu
 20

<210> SEQ ID NO 10
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: Human rhinovirus 16

<400> SEQUENCE: 10

Leu His Lys Gln Thr Gly Pro Ile Thr Gln Asn Pro Val Glu Arg Tyr
1 5 10 15

Val Asp Glu Val
 20

<210> SEQ ID NO 11
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: Human Rhinovirus b

<400> SEQUENCE: 11

Thr Ile Ser Gln Thr Val Ala Leu Thr Gln Gly Leu Gly Asp Glu Leu
1 5 10 15

Glu Glu Val Ile
 20

<210> SEQ ID NO 12
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: Coxsackievirus A21

<400> SEQUENCE: 12

Ser Gln Ser Lys Leu Ile Gly Arg Thr Gln Gly Ile Glu Asp Leu Ile
1 5 10 15

Asp Thr Ala Ile
 20

<210> SEQ ID NO 13
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: Human rhinovirus 14

<400> SEQUENCE: 13

Lys Lys Arg Lys Gly Asp Ile Lys Ser Tyr Gly Leu Gly Pro Arg Tyr
1 5 10 15

Gly Gly Ile Tyr
 20

<210> SEQ ID NO 14
<211> LENGTH: 20
<212> TYPE: PRT

-continued

<213> ORGANISM: Human rhinovirus 16

<400> SEQUENCE: 14

Ile	Arg	Pro	Arg	Thr	Asn	Leu	Thr	Thr	Val	Gly	Pro	Ser	Asp	Met	Tyr
1				5					10					15	

Val	His	Val	Gly
			20

<210> SEQ ID NO 15

<211> LENGTH: 20

<212> TYPE: PRT

<213> ORGANISM: Human Rhinovirus b

<400> SEQUENCE: 15

Lys	Lys	Arg	Lys	Gly	Asp	Ile	Lys	Ser	Tyr	Gly	Leu	Gly	Pro	Arg	Tyr
1				5					10					15	

Gly	Gly	Ile	Tyr
			20

<210> SEQ ID NO 16

<211> LENGTH: 20

<212> TYPE: PRT

<213> ORGANISM: Coxsackievirus A21

<400> SEQUENCE: 16

Leu	Thr	Lys	Val	Asp	Ser	Ile	Thr	Thr	Phe	Gly	Phe	Gly	His	Gln	Asn
1				5					10					15	

Lys	Ala	Val	Tyr
			20

<210> SEQ ID NO 17

<211> LENGTH: 20

<212> TYPE: PRT

<213> ORGANISM: Human rhinovirus 14

<400> SEQUENCE: 17

Arg	Gln	Leu	Glu	Cys	Ile	Ala	Glu	Glu	Gln	Gly	Leu	Ser	Asp	Tyr	Ile
1				5					10					15	

Thr	Gly	Leu	Gly
			20

<210> SEQ ID NO 18

<211> LENGTH: 20

<212> TYPE: PRT

<213> ORGANISM: Human rhinovirus 16

<400> SEQUENCE: 18

Leu	Arg	His	Phe	His	Cys	Ala	Glu	Glu	Gln	Gly	Ile	Thr	Asp	Tyr	Ile
1				5					10					15	

His	Met	Leu	Gly
			20

<210> SEQ ID NO 19

<211> LENGTH: 20

<212> TYPE: PRT

<213> ORGANISM: Human Rhinovirus b

<400> SEQUENCE: 19

Arg	Gln	Leu	Glu	Cys	Ile	Ala	Glu	Glu	Gln	Gly	Leu	Ser	Asp	Tyr	Ile
1				5					10					15	

Thr	Gly	Leu	Gly
			20

-continued

<210> SEQ ID NO 20
 <211> LENGTH: 20
 <212> TYPE: PRT
 <213> ORGANISM: Coxsackievirus A21

<400> SEQUENCE: 20

Trp Val Tyr Glu Glu Glu Ala Met Glu Gln Gly Ile Thr Ser Tyr Ile
 1 5 10 15

Glu Ser Leu Gly
 20

<210> SEQ ID NO 21
 <211> LENGTH: 20
 <212> TYPE: PRT
 <213> ORGANISM: Human rhinovirus 14

<400> SEQUENCE: 21

His Phe Gln Val Pro Tyr Ile Glu Arg Gln Ala Asn Asp Gly Trp Phe
 1 5 10 15

Arg Lys Phe Asn
 20

<210> SEQ ID NO 22
 <211> LENGTH: 20
 <212> TYPE: PRT
 <213> ORGANISM: Human rhinovirus 16

<400> SEQUENCE: 22

Trp Thr Gln Leu Thr Tyr Ile His Lys Glu Ser Asp Ser Trp Leu Lys
 1 5 10 15

Lys Phe Thr Glu
 20

<210> SEQ ID NO 23
 <211> LENGTH: 20
 <212> TYPE: PRT
 <213> ORGANISM: Human Rhinovirus b

<400> SEQUENCE: 23

His Phe Gln Val Pro Tyr Ile Glu Arg Gln Ala Asn Asp Gly Trp Phe
 1 5 10 15

Arg Lys Phe Asn
 20

<210> SEQ ID NO 24
 <211> LENGTH: 20
 <212> TYPE: PRT
 <213> ORGANISM: Coxsackievirus A21

<400> SEQUENCE: 24

Leu Leu Glu Ile Pro Tyr Val Met Arg Gln Gly Asp Gly Trp Met Lys
 1 5 10 15

Lys Phe Thr Glu
 20

<210> SEQ ID NO 25
 <211> LENGTH: 20
 <212> TYPE: PRT
 <213> ORGANISM: Human rhinovirus 14

<400> SEQUENCE: 25

-continued

```
Ile Thr Asp Ser Leu Glu Thr Leu Phe Gln Gly Pro Val Tyr Lys Asp
1           5           10           15
```

```
Leu Glu Ile Asp
           20
```

```
<210> SEQ ID NO 26
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: Human rhinovirus 16

<400> SEQUENCE: 26
```

```
Val Val Asp Val Met Ser Ala Ile Phe Gln Gly Pro Ile Ser Met Asp
1           5           10           15
```

```
Lys Pro Pro Pro
           20
```

```
<210> SEQ ID NO 27
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: Human Rhinovirus b

<400> SEQUENCE: 27
```

```
Ile Thr Asp Ser Leu Glu Thr Leu Phe Gln Gly Pro Val Tyr Lys Asp
1           5           10           15
```

```
Leu Glu Ile Asp
           20
```

```
<210> SEQ ID NO 28
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: Cocksackievirus A21

<400> SEQUENCE: 28
```

```
Ile Gly Asn Cys Met Glu Ala Leu Phe Gln Gly Pro Leu Arg Tyr Lys
1           5           10           15
```

```
Asp Leu Lys Ile
           20
```

```
<210> SEQ ID NO 29
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: Human rhinovirus 14

<400> SEQUENCE: 29
```

```
Val Ile Tyr Lys Leu Phe Ala Gln Thr Gln Gly Pro Tyr Ser Gly Asn
1           5           10           15
```

```
Pro Pro His Asn
           20
```

```
<210> SEQ ID NO 30
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: Human rhinovirus 16

<400> SEQUENCE: 30
```

```
Ile Ile Tyr Lys Leu Phe Cys Ser Leu Gln Gly Pro Tyr Ser Gly Glu
1           5           10           15
```

```
Pro Lys Pro Lys
           20
```

```
<210> SEQ ID NO 31
<211> LENGTH: 20
```

-continued

<212> TYPE: PRT
 <213> ORGANISM: Human Rhinovirus b

<400> SEQUENCE: 31

Val Ile Tyr Lys Leu Phe Ala Gln Thr Gln Gly Pro Tyr Ser Gly Asn
 1 5 10 15

Pro Pro His Asn
 20

<210> SEQ ID NO 32
 <211> LENGTH: 20
 <212> TYPE: PRT
 <213> ORGANISM: Coxsackievirus A21

<400> SEQUENCE: 32

Val Met Tyr Lys Leu Phe Ala Gly Gln Gln Gly Ala Tyr Thr Gly Leu
 1 5 10 15

Pro Asn Lys Lys
 20

<210> SEQ ID NO 33
 <211> LENGTH: 20
 <212> TYPE: PRT
 <213> ORGANISM: Human rhinovirus 14

<400> SEQUENCE: 33

Ala Pro Thr Leu Arg Pro Val Val Val Gln Gly Pro Asn Thr Glu Phe
 1 5 10 15

Ala Leu Ser Leu
 20

<210> SEQ ID NO 34
 <211> LENGTH: 20
 <212> TYPE: PRT
 <213> ORGANISM: Human rhinovirus 16

<400> SEQUENCE: 34

Lys Val Pro Glu Arg Arg Val Val Ala Gln Gly Pro Glu Glu Glu Phe
 1 5 10 15

Gly Met Ser Ile
 20

<210> SEQ ID NO 35
 <211> LENGTH: 20
 <212> TYPE: PRT
 <213> ORGANISM: Human Rhinovirus b

<400> SEQUENCE: 35

Ala Pro Thr Leu Arg Pro Val Val Val Gln Gly Pro Asn Thr Glu Phe
 1 5 10 15

Ala Leu Ser Leu
 20

<210> SEQ ID NO 36
 <211> LENGTH: 20
 <212> TYPE: PRT
 <213> ORGANISM: Coxsackievirus A21

<400> SEQUENCE: 36

Val Pro Thr Ile Arg Val Ala Lys Val Gln Gly Pro Gly Phe Asp Tyr
 1 5 10 15

Ala Val Ala Met

-continued

20

<210> SEQ ID NO 37
 <211> LENGTH: 20
 <212> TYPE: PRT
 <213> ORGANISM: Human rhinovirus 14

<400> SEQUENCE: 37

Leu Lys Lys Gln Tyr Phe Val Glu Lys Gln Gly Gln Val Ile Ala Arg
 1 5 10 15

His Lys Val Arg
 20

<210> SEQ ID NO 38
 <211> LENGTH: 20
 <212> TYPE: PRT
 <213> ORGANISM: Human rhinovirus 16

<400> SEQUENCE: 38

Leu Leu Arg Ser Tyr Phe Thr Glu Gln Gln Gly Gln Ile Gln Ile Ser
 1 5 10 15

Lys His Val Lys
 20

<210> SEQ ID NO 39
 <211> LENGTH: 20
 <212> TYPE: PRT
 <213> ORGANISM: Human Rhinovirus b

<400> SEQUENCE: 39

Leu Lys Lys Gln Tyr Phe Val Glu Lys Gln Gly Gln Val Ile Ala Arg
 1 5 10 15

His Lys Val Arg
 20

<210> SEQ ID NO 40
 <211> LENGTH: 20
 <212> TYPE: PRT
 <213> ORGANISM: Coxsackievirus A21

<400> SEQUENCE: 40

Leu Lys Arg Ser Tyr Phe Thr Gln Asn Gln Gly Glu Ile Gln Trp Met
 1 5 10 15

Arg Ser Ser Lys
 20

<210> SEQ ID NO 41
 <211> LENGTH: 20
 <212> TYPE: PRT
 <213> ORGANISM: SARS: Human Severe Acute Respiratory Syndrome Virus

<400> SEQUENCE: 41

Arg Glu Leu Thr Arg Glu Leu Asn Gly Gly Ala Val Thr Arg Tyr Val
 1 5 10 15

Asp Asn Asn Phe
 20

<210> SEQ ID NO 42
 <211> LENGTH: 20
 <212> TYPE: PRT
 <213> ORGANISM: Bovine coronavirus

<400> SEQUENCE: 42

-continued

Met Ser Lys Ile Asn Lys Tyr Gly Leu Glu Val Lys Pro Leu Leu Tyr
 1 5 10 15

Val Asp Gln Tyr
 20

<210> SEQ ID NO 43
 <211> LENGTH: 20
 <212> TYPE: PRT
 <213> ORGANISM: Human coronavirus

<400> SEQUENCE: 43

Asp Val Val Phe Gly Lys Arg Gly Gly Gly Asn Val Thr Tyr Thr Asp
 1 5 10 15

Gln Tyr Leu Cys
 20

<210> SEQ ID NO 44
 <211> LENGTH: 20
 <212> TYPE: PRT
 <213> ORGANISM: SARS: Human Severe Acute Respiratory Syndrome Virus

<400> SEQUENCE: 44

Thr Asn Asn Val Phe Arg Leu Lys Gly Gly Ala Pro Ile Lys Gly Val
 1 5 10 15

Thr Phe Gly Glu
 20

<210> SEQ ID NO 45
 <211> LENGTH: 20
 <212> TYPE: PRT
 <213> ORGANISM: Bovine coronavirus

<400> SEQUENCE: 45

Leu Asp Gln Ala Trp Arg Val Pro Cys Ala Gly Arg Arg Val Thr Phe
 1 5 10 15

Lys Glu Gln Pro
 20

<210> SEQ ID NO 46
 <211> LENGTH: 20
 <212> TYPE: PRT
 <213> ORGANISM: Human coronavirus

<400> SEQUENCE: 46

Leu Pro Val Ala Phe Thr Lys Ala Ala Gly Gly Lys Val Ser Phe Ser
 1 5 10 15

Asp Asp Val Glu
 20

<210> SEQ ID NO 47
 <211> LENGTH: 20
 <212> TYPE: PRT
 <213> ORGANISM: SARS: Human Severe Acute Respiratory Syndrome Virus

<400> SEQUENCE: 47

Ile Thr Thr Lys Ile Ser Leu Lys Gly Gly Lys Ile Val Ser Thr Cys
 1 5 10 15

Phe Lys Leu Met
 20

<210> SEQ ID NO 48

-continued

<211> LENGTH: 20
 <212> TYPE: PRT
 <213> ORGANISM: Bovine coronavirus

<400> SEQUENCE: 48

Leu Thr Thr Pro Phe Ser Leu Lys Gly Gly Ala Val Phe Ser Tyr Phe
 1 5 10 15

Val Tyr Val Cys
 20

<210> SEQ ID NO 49
 <211> LENGTH: 20
 <212> TYPE: PRT
 <213> ORGANISM: Human coronavirus

<400> SEQUENCE: 49

Ala Thr Ser Ile Val Ala Lys Gln Gly Ala Gly Asp Ala Gly His Ser
 1 5 10 15

Leu Thr Trp Leu
 20

<210> SEQ ID NO 50
 <211> LENGTH: 20
 <212> TYPE: PRT
 <213> ORGANISM: SARS: Human Severe Acute Respiratory Syndrome Virus

<400> SEQUENCE: 50

Gln Thr Ser Ile Thr Ser Ala Val Leu Gln Ser Gly Phe Arg Lys Met
 1 5 10 15

Ala Phe Pro Ser
 20

<210> SEQ ID NO 51
 <211> LENGTH: 20
 <212> TYPE: PRT
 <213> ORGANISM: Bovine coronavirus

<400> SEQUENCE: 51

Thr Ala Ser Val Ser Thr Ser Phe Leu Gln Ser Gly Ile Val Lys Met
 1 5 10 15

Val Asn Pro Thr
 20

<210> SEQ ID NO 52
 <211> LENGTH: 20
 <212> TYPE: PRT
 <213> ORGANISM: Human coronavirus

<400> SEQUENCE: 52

Pro Thr Val Ser Tyr Gly Ser Thr Leu Gln Ala Gly Leu Arg Lys Met
 1 5 10 15

Ala Gln Pro Ser
 20

<210> SEQ ID NO 53
 <211> LENGTH: 20
 <212> TYPE: PRT
 <213> ORGANISM: SARS: Human Severe Acute Respiratory Syndrome Virus

<400> SEQUENCE: 53

Val Arg Gln Cys Ser Gly Val Thr Phe Gln Gly Lys Phe Lys Lys Ile
 1 5 10 15

-continued

Val Lys Gly Thr
20

<210> SEQ ID NO 54
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: Bovine coronavirus

<400> SEQUENCE: 54

Tyr Gln Gln Leu Ala Gly Ile Lys Leu Gln Ser Lys Arg Thr Arg Leu
1 5 10 15

Val Lys Gly Ile
20

<210> SEQ ID NO 55
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: Human coronavirus

<400> SEQUENCE: 55

Val Lys Gln Met Phe Gly Val Asn Leu Gln Ser Gly Lys Thr Thr Ser
1 5 10 15

Met Phe Lys Ser
20

<210> SEQ ID NO 56
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: SARS: Human Severe Acute Respiratory Syndrome Virus

<400> SEQUENCE: 56

Lys Pro Cys Ile Lys Val Ala Thr Val Gln Ser Lys Met Ser Asp Val
1 5 10 15

Lys Cys Thr Ser
20

<210> SEQ ID NO 57
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: Bovine coronavirus

<400> SEQUENCE: 57

Val Pro Ile Ile Glu Val Ser Gln Phe Gln Ser Lys Leu Thr Asp Val
1 5 10 15

Lys Cys Ala Asn
20

<210> SEQ ID NO 58
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: Human coronavirus

<400> SEQUENCE: 58

Pro Arg Thr Ile Lys Val Ser Thr Val Gln Ser Lys Leu Thr Asp Leu
1 5 10 15

Lys Cys Thr Asn
20

<210> SEQ ID NO 59
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: SARS: Human Severe Acute Respiratory Syndrome Virus

-continued

<400> SEQUENCE: 59

Glu Met Leu Asp Asn Arg Ala Thr Leu Gln Ala Ile Ala Ser Glu Phe
 1 5 10 15

Ser Ser Leu Pro
 20

<210> SEQ ID NO 60

<211> LENGTH: 20

<212> TYPE: PRT

<213> ORGANISM: Bovine coronavirus

<400> SEQUENCE: 60

Asp Tyr Ala Lys Asp Asn Thr Val Leu Gln Ala Leu Gln Ser Glu Phe
 1 5 10 15

Val Asn Met Ala
 20

<210> SEQ ID NO 61

<211> LENGTH: 20

<212> TYPE: PRT

<213> ORGANISM: Human coronavirus

<400> SEQUENCE: 61

Ser Tyr Phe Glu Asn Asp Ser Ile Leu Gln Ser Val Ala Ser Ser Phe
 1 5 10 15

Val Gly Met Pro
 20

<210> SEQ ID NO 62

<211> LENGTH: 20

<212> TYPE: PRT

<213> ORGANISM: SARS: Human Severe Acute Respiratory Syndrome Virus

<400> SEQUENCE: 62

Leu Arg Ala Asn Ser Ala Val Lys Leu Gln Asn Asn Glu Leu Ser Pro
 1 5 10 15

Val Ala Leu Arg
 20

<210> SEQ ID NO 63

<211> LENGTH: 20

<212> TYPE: PRT

<213> ORGANISM: Bovine coronavirus

<400> SEQUENCE: 63

His Asn Glu Val Ser Ala Thr Val Leu Gln Asn Asn Glu Leu Met Pro
 1 5 10 15

Ala Lys Leu Lys
 20

<210> SEQ ID NO 64

<211> LENGTH: 20

<212> TYPE: PRT

<213> ORGANISM: Human coronavirus

<400> SEQUENCE: 64

Leu Thr Cys Glu Arg Val Val Lys Leu Gln Asn Asn Glu Ile Met Pro
 1 5 10 15

Gly Lys Met Lys
 20

-continued

<210> SEQ ID NO 65
 <211> LENGTH: 20
 <212> TYPE: PRT
 <213> ORGANISM: SARS: Human Severe Acute Respiratory Syndrome Virus

<400> SEQUENCE: 65

Gly Ser Leu Ala Ala Thr Val Arg Leu Gln Ala Gly Asn Ala Thr Glu
 1 5 10 15
 Val Pro Ala Asn
 20

<210> SEQ ID NO 66
 <211> LENGTH: 20
 <212> TYPE: PRT
 <213> ORGANISM: Bovine coronavirus

<400> SEQUENCE: 66

Gly Thr Ile Ser Ser Thr Val Arg Leu Gln Ala Gly Thr Ala Thr Glu
 1 5 10 15
 Tyr Ala Ser Asn
 20

<210> SEQ ID NO 67
 <211> LENGTH: 20
 <212> TYPE: PRT
 <213> ORGANISM: Human coronavirus

<400> SEQUENCE: 67

Gly Tyr Ile Gly Ala Thr Val Arg Leu Gln Ala Gly Lys Gln Thr Glu
 1 5 10 15
 Phe Val Ser Asn
 20

<210> SEQ ID NO 68
 <211> LENGTH: 20
 <212> TYPE: PRT
 <213> ORGANISM: SARS: Human Severe Acute Respiratory Syndrome Virus

<400> SEQUENCE: 68

Cys Asp Gln Leu Arg Glu Pro Leu Met Gln Ser Ala Asp Ala Ser Thr
 1 5 10 15
 Phe Leu Asn Arg
 20

<210> SEQ ID NO 69
 <211> LENGTH: 20
 <212> TYPE: PRT
 <213> ORGANISM: Bovine coronavirus

<400> SEQUENCE: 69

Ser Cys Val Ser Thr Asp Thr Thr Val Gln Ser Lys Asp Thr Asn Phe
 1 5 10 15
 Leu Asn Arg Val
 20

<210> SEQ ID NO 70
 <211> LENGTH: 20
 <212> TYPE: PRT
 <213> ORGANISM: Human coronavirus

<400> SEQUENCE: 70

Gly Cys Thr Cys Asp Arg Thr Ala Ile Gln Ser Phe Asp Asn Ser Tyr
 1 5 10 15

-continued

Leu Asn Arg Val
20

<210> SEQ ID NO 71
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: SARS: Human Severe Acute Respiratory Syndrome Virus

<400> SEQUENCE: 71

Ala Met Tyr Thr Pro His Thr Val Leu Gln Ala Val Gly Ala Cys Val
1 5 10 15

Leu Cys Asn Ser
20

<210> SEQ ID NO 72
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: Bovine coronavirus

<400> SEQUENCE: 72

Asn Met Tyr Leu Arg Ser Ala Val Met Gln Ser Val Gly Ala Cys Val
1 5 10 15

Val Cys Ser Ser
20

<210> SEQ ID NO 73
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: Human coronavirus

<400> SEQUENCE: 73

Ser Met Tyr Glu Lys Ser Thr Val Leu Gln Ala Ala Gly Leu Cys Val
1 5 10 15

Val Cys Gly Ser
20

<210> SEQ ID NO 74
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: SARS: Human Severe Acute Respiratory Syndrome Virus

<400> SEQUENCE: 74

Ile Pro Arg Arg Asn Val Ala Thr Leu Gln Ala Glu Asn Val Thr Gly
1 5 10 15

Leu Phe Lys Asp
20

<210> SEQ ID NO 75
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: Bovine coronavirus

<400> SEQUENCE: 75

Val Pro Gln Ala Val Glu Thr Arg Val Gln Cys Ser Thr Asn Leu Phe
1 5 10 15

Lys Asp Cys Ser
20

<210> SEQ ID NO 76
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: Human coronavirus

-continued

<400> SEQUENCE: 76

Phe Phe Glu Ile Thr Met Thr Asp Leu Gln Ser Glu Ser Ser Cys Gly
 1 5 10 15

Leu Phe Lys Asp
 20

<210> SEQ ID NO 77

<211> LENGTH: 20

<212> TYPE: PRT

<213> ORGANISM: SARS: Human Severe Acute Respiratory Syndrome Virus

<400> SEQUENCE: 77

Asn Leu Trp Asn Thr Phe Thr Arg Leu Gln Ser Leu Glu Asn Val Ala
 1 5 10 15

Tyr Asn Val Val
 20

<210> SEQ ID NO 78

<211> LENGTH: 20

<212> TYPE: PRT

<213> ORGANISM: Bovine coronavirus

<400> SEQUENCE: 78

Asn Leu Trp Asn Thr Phe Thr Lys Leu Gln Ser Leu Glu Asn Val Val
 1 5 10 15

Tyr Asn Leu Val
 20

<210> SEQ ID NO 79

<211> LENGTH: 20

<212> TYPE: PRT

<213> ORGANISM: Human coronavirus

<400> SEQUENCE: 79

Trp Gln Thr Phe Thr Glu Val Asn Leu Gln Gly Leu Glu Asn Ile Ala
 1 5 10 15

Phe Asn Val Val
 20

<210> SEQ ID NO 80

<211> LENGTH: 20

<212> TYPE: PRT

<213> ORGANISM: SARS: Human Severe Acute Respiratory Syndrome Virus

<400> SEQUENCE: 80

His Val Glu Thr Phe Tyr Pro Lys Leu Gln Ala Ser Gln Ala Trp Gln
 1 5 10 15

Pro Gly Val Ala
 20

<210> SEQ ID NO 81

<211> LENGTH: 20

<212> TYPE: PRT

<213> ORGANISM: Bovine coronavirus

<400> SEQUENCE: 81

Lys Val Met Thr Phe Tyr Pro Arg Leu Gln Ala Ala Ser Asp Trp Lys
 1 5 10 15

Pro Gly Tyr Ser
 20

-continued

<210> SEQ ID NO 82
 <211> LENGTH: 20
 <212> TYPE: PRT
 <213> ORGANISM: Human coronavirus

<400> SEQUENCE: 82

Ala Val Ala Thr Phe Tyr Pro Gln Leu Gln Ser Ala Glu Trp Lys Cys
 1 5 10 15

Gly Tyr Ser Met
 20

<210> SEQ ID NO 83
 <211> LENGTH: 9
 <212> TYPE: PRT
 <213> ORGANISM: Human rhinovirus

<400> SEQUENCE: 83

Arg Pro Val Val Val Gln Gly Pro Asn
 1 5

<210> SEQ ID NO 84
 <211> LENGTH: 9
 <212> TYPE: PRT
 <213> ORGANISM: Human coronavirus

<400> SEQUENCE: 84

Ser Thr Leu Gln Ser Gly Leu Arg Lys
 1 5

<210> SEQ ID NO 85
 <211> LENGTH: 9
 <212> TYPE: PRT
 <213> ORGANISM: SARS: Human Severe Acute Respiratory Syndrome Virus

<400> SEQUENCE: 85

Ala Thr Val Arg Leu Gln Ala Gly Phe
 1 5

<210> SEQ ID NO 86
 <211> LENGTH: 16
 <212> TYPE: PRT
 <213> ORGANISM: SARS: Human Severe Acute Respiratory Syndrome Virus

<400> SEQUENCE: 86

Val Ser Val Asn Ser Thr Leu Gln Ser Gly Leu Arg Lys Met Ala Cys
 1 5 10 15

<210> SEQ ID NO 87
 <211> LENGTH: 12
 <212> TYPE: PRT
 <213> ORGANISM: Human rhinovirus 14

<400> SEQUENCE: 87

Lys Ser Ile Val Pro Gln Gly Leu Pro Thr Thr Thr
 1 5 10

<210> SEQ ID NO 88
 <211> LENGTH: 12
 <212> TYPE: PRT
 <213> ORGANISM: Human rhinovirus 14

<400> SEQUENCE: 88

Cys Ile Ala Glu Glu Gln Gly Leu Ser Asp Tyr Ile
 1 5 10

-continued

<210> SEQ ID NO 89
 <211> LENGTH: 12
 <212> TYPE: PRT
 <213> ORGANISM: Human rhinovirus 14

<400> SEQUENCE: 89

Leu Glu Thr Leu Phe Gln Gly Pro Val Tyr Lys Asp
 1 5 10

<210> SEQ ID NO 90
 <211> LENGTH: 12
 <212> TYPE: PRT
 <213> ORGANISM: Human rhinovirus 14

<400> SEQUENCE: 90

Leu Phe Ala Gln Thr Gln Gly Pro Tyr Ser Gly Asn
 1 5 10

<210> SEQ ID NO 91
 <211> LENGTH: 12
 <212> TYPE: PRT
 <213> ORGANISM: Human rhinovirus 14

<400> SEQUENCE: 91

Arg Pro Val Val Val Gln Gly Pro Asn Thr Glu Phe
 1 5 10

<210> SEQ ID NO 92
 <211> LENGTH: 12
 <212> TYPE: PRT
 <213> ORGANISM: Human rhinovirus 14

<400> SEQUENCE: 92

Gly Gly Asn Gly Arg Gln Gly Phe Ser Ala Gln Leu
 1 5 10

<210> SEQ ID NO 93
 <211> LENGTH: 12
 <212> TYPE: PRT
 <213> ORGANISM: Human rhinovirus 14

<400> SEQUENCE: 93

Tyr Phe Val Glu Lys Gln Gly Gln Val Ile Ala Arg
 1 5 10

<210> SEQ ID NO 94
 <211> LENGTH: 12
 <212> TYPE: PRT
 <213> ORGANISM: Human rhinovirus 14

<400> SEQUENCE: 94

Ser Asn Leu Val Val Gln Ala Met Tyr Val Pro His
 1 5 10

<210> SEQ ID NO 95
 <211> LENGTH: 12
 <212> TYPE: PRT
 <213> ORGANISM: Human rhinovirus 14

<400> SEQUENCE: 95

Tyr Pro Ser Arg Phe Gln Ala Gly Val Met Lys Gly
 1 5 10

-continued

<210> SEQ ID NO 96
 <211> LENGTH: 12
 <212> TYPE: PRT
 <213> ORGANISM: Human rhinovirus 14
 <400> SEQUENCE: 96
 Pro Tyr Ile Glu Arg Gln Ala Asn Asp Gly Trp Phe
 1 5 10

<210> SEQ ID NO 97
 <211> LENGTH: 12
 <212> TYPE: PRT
 <213> ORGANISM: Human rhinovirus 14
 <400> SEQUENCE: 97
 Asn Lys Val Leu Pro Gln Ala Lys Glu Lys Leu Glu
 1 5 10

<210> SEQ ID NO 98
 <211> LENGTH: 12
 <212> TYPE: PRT
 <213> ORGANISM: Human rhinovirus 14
 <400> SEQUENCE: 98
 Glu Arg Ala Met Asn Gln Ala Ser Met Ile Ile Asn
 1 5 10

<210> SEQ ID NO 99
 <211> LENGTH: 12
 <212> TYPE: PRT
 <213> ORGANISM: Human rhinovirus 14
 <400> SEQUENCE: 99
 Gln Leu Ala Ser His Glu Gly Gly Asn Val Ser Val
 1 5 10

<210> SEQ ID NO 100
 <211> LENGTH: 12
 <212> TYPE: PRT
 <213> ORGANISM: Human rhinovirus 14
 <400> SEQUENCE: 100
 Thr Val Ala Leu Thr Glu Gly Leu Gly Asp Glu Leu
 1 5 10

<210> SEQ ID NO 101
 <211> LENGTH: 12
 <212> TYPE: PRT
 <213> ORGANISM: Human rhinovirus 14
 <400> SEQUENCE: 101
 Thr Asn Ile Trp Ile Glu Gly Ser Pro Tyr Tyr Pro
 1 5 10

<210> SEQ ID NO 102
 <211> LENGTH: 12
 <212> TYPE: PRT
 <213> ORGANISM: Human rhinovirus 14
 <400> SEQUENCE: 102
 Gly Leu Leu Thr Ala Glu Gly Ser Gly Tyr Val Cys
 1 5 10

<210> SEQ ID NO 103
 <211> LENGTH: 12

-continued

<212> TYPE: PRT
<213> ORGANISM: Human rhinovirus 14

<400> SEQUENCE: 103

Ile Ser Glu Asp Leu Glu Gly Val Asp Ala Thr Leu
1 5 10

<210> SEQ ID NO 104
<211> LENGTH: 12
<212> TYPE: PRT
<213> ORGANISM: Human rhinovirus 14

<400> SEQUENCE: 104

Glu Ile Tyr Val Val Glu Gly Gly Met Pro Ser Gly
1 5 10

<210> SEQ ID NO 105
<211> LENGTH: 13
<212> TYPE: PRT
<213> ORGANISM: Human rhinovirus 14

<400> SEQUENCE: 105

Asp Ser Leu Glu Thr Leu Phe Gln Gly Pro Val Tyr Lys
1 5 10

<210> SEQ ID NO 106
<211> LENGTH: 14
<212> TYPE: PRT
<213> ORGANISM: Human rhinovirus 14

<400> SEQUENCE: 106

Glu Ala Ile Ala Glu Glu Gln Gly Leu Ser Asp Tyr Ile Thr
1 5 10

<210> SEQ ID NO 107
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Human rhinovirus 14

<400> SEQUENCE: 107

Val Pro Tyr Ile Glu Arg Gln Ala Asn Asp Gly Trp Phe Arg Lys
1 5 10 15

<210> SEQ ID NO 108
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Human rhinovirus 14

<400> SEQUENCE: 108

Arg Ser Lys Ser Ile Val Pro Gln Gly Leu Pro Thr Thr Thr Tyr
1 5 10 15

<210> SEQ ID NO 109
<211> LENGTH: 17
<212> TYPE: PRT
<213> ORGANISM: Human rhinovirus 14

<400> SEQUENCE: 109

Ser Gln Thr Val Ala Leu Thr Glu Gly Leu Gly Asp Glu Leu Glu Glu
1 5 10 15

Tyr

<210> SEQ ID NO 110
<211> LENGTH: 14

-continued

```

<212> TYPE: PRT
<213> ORGANISM: Human rhinovirus 14

<400> SEQUENCE: 110

Lys Leu Phe Ala Gln Thr Gln Gly Pro Tyr Ser Gly Asn Pro
1             5             10

<210> SEQ ID NO 111
<211> LENGTH: 13
<212> TYPE: PRT
<213> ORGANISM: Human rhinovirus 14

<400> SEQUENCE: 111

Tyr Arg Pro Val Val Val Gln Gly Pro Asn Thr Glu Phe
1             5             10

<210> SEQ ID NO 112
<211> LENGTH: 14
<212> TYPE: PRT
<213> ORGANISM: Human rhinovirus 14

<400> SEQUENCE: 112

Lys Gln Tyr Phe Val Glu Lys Gln Gly Gln Val Ile Ala Arg
1             5             10

<210> SEQ ID NO 113
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Human rhinovirus 1A
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<223> OTHER INFORMATION: MOC; fluorochrome
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (10)..()
<223> OTHER INFORMATION: DNP-NH2; Dinitrophenol quencher

<400> SEQUENCE: 113

Arg Ala Glu Leu Gln Gly Pro Tyr Asp Lys
1             5             10

<210> SEQ ID NO 114
<211> LENGTH: 19
<212> TYPE: PRT
<213> ORGANISM: Human Poliovirus- POLIM

<400> SEQUENCE: 114

Ile Gly Asn Cys Met Glu Ala Leu Phe Gln Gly Pro Gln Tyr Lys Asp
1             5             10             15

Leu Lys Ile

<210> SEQ ID NO 115
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: Human Poliovirus- POLIS

<400> SEQUENCE: 115

Ile Gly Asn Cys Met Glu Ala Leu Phe Gln Gly Pro Leu Gln Tyr Lys
1             5             10             15

Asp Leu Lys Ile
20

<210> SEQ ID NO 116
<211> LENGTH: 20
<212> TYPE: PRT

```

-continued

<213> ORGANISM: Human Poliovirus- POL32

<400> SEQUENCE: 116

Ile Gly Asn Cys Met Glu Ala Leu Phe Gln Gly Pro Leu Gln Tyr Lys
1 5 10 15

Asp Leu Lys Ile
 20

<210> SEQ ID NO 117

<211> LENGTH: 20

<212> TYPE: PRT

<213> ORGANISM: Human Poliovirus- POL3L

<400> SEQUENCE: 117

Ile Gly Asn Cys Met Glu Ala Leu Phe Gln Gly Pro Leu Gln Tyr Lys
1 5 10 15

Asp Leu Lys Ile
 20

<210> SEQ ID NO 118

<211> LENGTH: 20

<212> TYPE: PRT

<213> ORGANISM: Human Coxsacivirus -COXA2

<400> SEQUENCE: 118

Ile Gly Asn Cys Met Glu Ala Leu Phe Gln Gly Pro Leu Arg Tyr Lys
1 5 10 15

Asp Leu Lys Ile
 20

<210> SEQ ID NO 119

<211> LENGTH: 20

<212> TYPE: PRT

<213> ORGANISM: Human Coxsacivirus -COXA4

<400> SEQUENCE: 119

Ile Gly Asn Cys Met Glu Ala Leu Phe Gln Gly Pro Ile Gln Tyr Arg
1 5 10 15

Asp Val Met Ile
 20

<210> SEQ ID NO 120

<211> LENGTH: 20

<212> TYPE: PRT

<213> ORGANISM: Bovine enterovirus

<400> SEQUENCE: 120

Ile Gly Asn Val Leu Glu Ala Leu Phe Gln Gly Pro Val Cys Tyr Lys
1 5 10 15

Pro Leu Arg Ile
 20

<210> SEQ ID NO 121

<211> LENGTH: 20

<212> TYPE: PRT

<213> ORGANISM: Human Coxsacivirus -COXA9

<400> SEQUENCE: 121

Val Gly Ala Thr Leu Glu Ala Leu Phe Gln Gly Pro Pro Ile Tyr Arg
1 5 10 15

Glu Ile Lys Ile
 20

-continued

<210> SEQ ID NO 122
 <211> LENGTH: 20
 <212> TYPE: PRT
 <213> ORGANISM: Human Coxacivirus -COXB1

<400> SEQUENCE: 122

Val Gly Ala Thr Leu Glu Ala Leu Phe Gln Gly Pro Pro Ile Tyr Arg
 1 5 10 15

Glu Ile Lys Ile
 20

<210> SEQ ID NO 123
 <211> LENGTH: 20
 <212> TYPE: PRT
 <213> ORGANISM: Human Coxacivirus -COXB5

<400> SEQUENCE: 123

Val Gly Ala Thr Leu Glu Ala Leu Phe Gln Gly Pro Pro Ile Tyr Arg
 1 5 10 15

Glu Ile Lys Ile
 20

<210> SEQ ID NO 124
 <211> LENGTH: 20
 <212> TYPE: PRT
 <213> ORGANISM: Human Echovirus EC11G

<400> SEQUENCE: 124

Val Gly Ala Thr Leu Glu Ala Leu Phe Gln Gly Pro Pro Ile Tyr Arg
 1 5 10 15

Glu Ile Lys Ile
 20

<210> SEQ ID NO 125
 <211> LENGTH: 20
 <212> TYPE: PRT
 <213> ORGANISM: Human Coxacivirus -COXB4

<400> SEQUENCE: 125

Val Gly Ala Thr Leu Glu Ala Leu Phe Gln Gly Pro Pro Val Tyr Arg
 1 5 10 15

Glu Ile Lys Ile
 20

<210> SEQ ID NO 126
 <211> LENGTH: 20
 <212> TYPE: PRT
 <213> ORGANISM: Swine Vesicular Disease Virus SVDVH

<400> SEQUENCE: 126

Val Gly Ala Thr Leu Glu Ala Leu Phe Gln Gly Pro Pro Val Tyr Arg
 1 5 10 15

Glu Ile Lys Ile
 20

<210> SEQ ID NO 127
 <211> LENGTH: 20
 <212> TYPE: PRT
 <213> ORGANISM: Swine Vesicular Disease Virus SVDVU

<400> SEQUENCE: 127

-continued

Val	Gly	Ala	Thr	Leu	Glu	Ala	Leu	Phe	Gln	Gly	Pro	Pro	Val	Tyr	Arg
1				5					10					15	

Glu	Ile	Lys	Ile
			20

<210> SEQ ID NO 128
 <211> LENGTH: 20
 <212> TYPE: PRT
 <213> ORGANISM: Human Coxacivirus -COXB3

<400> SEQUENCE: 128

Val	Gly	Thr	Thr	Glu	Ala	Leu	Phe	Gln	Gly	Pro	Pro	Val	Tyr	Arg
1				5				10					15	

Glu	Ile	Lys	Ile
			20

<210> SEQ ID NO 129
 <211> LENGTH: 20
 <212> TYPE: PRT
 <213> ORGANISM: Human Enterovirus HUEV7

<400> SEQUENCE: 129

Thr	Gln	Asp	Lys	Leu	Glu	Ala	Leu	Phe	Gln	Gly	Pro	Pro	Thr	Phe	Lys
1				5					10					15	

Glu	Ile	Lys	Ile
			20

<210> SEQ ID NO 130
 <211> LENGTH: 20
 <212> TYPE: PRT
 <213> ORGANISM: Human rhinovirus 1B

<400> SEQUENCE: 130

Val	Val	Asp	Val	Met	Ser	Ala	Ile	Phe	Gln	Gly	Pro	Ile	Ser	Leu	Asp
1				5					10					15	

Ala	Pro	Pro	Pro
			20

<210> SEQ ID NO 131
 <211> LENGTH: 20
 <212> TYPE: PRT
 <213> ORGANISM: Human rhinovirus 2

<400> SEQUENCE: 131

Val	Val	Asp	Val	Met	Thr	Ala	Ile	Phe	Gln	Gly	Pro	Ile	Asp	Met	Lys
1				5					10					15	

Asn	Pro	Pro	Pro
			20

<210> SEQ ID NO 132
 <211> LENGTH: 20
 <212> TYPE: PRT
 <213> ORGANISM: Human rhinovirus 89

<400> SEQUENCE: 132

Ala	Ala	Gln	Ala	Met	Glu	Ala	Ile	Phe	Gln	Gly	Ile	Asp	Leu	Gln	Ser
1				5					10					15	

Pro	Pro	Pro	Pro
			20

<210> SEQ ID NO 133
 <211> LENGTH: 20

-continued

```

<212> TYPE: PRT
<213> ORGANISM: Human rhinovirus 14

<400> SEQUENCE: 133

Ile Thr Asp Ser Leu Glu Thr Leu Phe Gln Gly Pro Val Tyr Lys Asp
1           5           10           15
Leu Glu Ile Asp
                20
    
```

I claim:

1. A method for determining the presence of a Nidovirus or Picornavirus virus in a sample suspected of containing said virus, the method comprising:

Contacting a sample with a substrate peptide comprising at least four amino acid residues and capable of being cleaved at a cleavage point by an enzyme to form a cleaved first peptide fragment and a second peptide fragment; and

detecting the virus by observing the signal from a signaling moiety linked to the substrate peptide;

wherein the enzyme is encoded by nucleic acid of the virus and is present in the sample;

wherein said substrate peptide comprises an amino acid sequence identical to a sequence of a polyprotein cleavage region of said enzyme and including said cleavage point;

wherein said peptide cleavage fragments are peptide fragments of the substrate peptide;

wherein a signaling moiety is linked to a portion of the peptide present in the first peptide fragment; and wherein said method is a heterogeneous assay method.

2. The method of claim 1, wherein a quenching moiety is linked to a portion of the peptide present in the second peptide fragment and wherein the signaling moiety and the quenching moiety are linked to the peptide at relative positions such that the quenching moiety quenches the a signal of the signaling moiety unless the peptide is cleaved at the cleavage point.

3. The method of claim 1, further comprising contacting the sample with a second peptide comprising at least four

amino acid residues and capable of being cleaved at a cleavage point by a second enzyme, wherein the second enzyme is encoded by the viral nucleic acid, and wherein said second peptide comprises an amino acid sequence identical to a sequence of a polyprotein cleavage region of said second enzyme and including said cleavage point.

4. The method of claim 1, wherein the enzyme is a protease.

5. The method of claim 1, wherein the signal of the signaling moiety is selected from a group consisting of a moiety emitting a fluorescent signal, a colorimetric signal, and a chemiluminescent signal.

6. The method of claim 5, wherein the signal of the signaling moiety emits a fluorescent signal.

7. The method of claim 1, wherein the virus is detected in a sample taken from an animal thought to be infected by the virus.

8. The method of claim 7, wherein the animal is human.

9. The method of claim 8, wherein the sample is selected from a group consisting of mucus, saliva, blood, serum, plasma, urine, spinal fluid, sputum, tissue biopsy, bronchoalveolar fluid, and tears.

10. The method of claim 1, wherein the virus is a Rhinovirus.

11. The method of claim 1, wherein said peptide comprises at least seven amino acid residues.

12. The method of claim 1, wherein said peptide comprises at least ten amino acid residues.

* * * * *

专利名称(译)	SARS和其他病毒性疾病的酶学诊断试验		
公开(公告)号	US7635557	公开(公告)日	2009-12-22
申请号	US10/875133	申请日	2004-06-23
[标]申请(专利权)人(译)	阿拉德DORIT		
申请(专利权)人(译)	阿拉德DORIT		
当前申请(专利权)人(译)	MND诊断LTD.		
[标]发明人	ARAD DORIT		
发明人	ARAD, DORIT		
IPC分类号	C12Q1/70 C12N15/41 C12Q1/00 G01N33/53 C12Q1/37		
CPC分类号	C12Q1/37 G01N2333/165 G01N2333/095		
优先权	60/480605 2003-06-23 US		
其他公开文献	US20050048473A1		
外部链接	Espacenet USPTO		

摘要(译)

本发明涉及用于测试样品中病毒的方法，组合物和试剂盒。该方法通过使样品与能够被病毒酶切割的肽化合物接触以形成肽化合物片段来确定病毒酶的存在。肽化合物片段的检测证实了病毒的存在。

